

# Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.3-Tangent/99-4.3.10-c+d-x-<sup>^</sup>m-a+b-tan-<sup>^</sup>n

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 3:25am

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>detailed summary tables of results</b>	<b>20</b>
<b>3</b>	<b>Listing of integrals</b>	<b>43</b>
<b>4</b>	<b>Appendix</b>	<b>437</b>

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	3
1.2	Results . . . . .	4
1.3	Time and leaf size Performance . . . . .	7
1.4	Performance based on number of rules Rubi used . . . . .	9
1.5	Performance based on number of steps Rubi used . . . . .	10
1.6	Solved integrals histogram based on leaf size of result . . . . .	11
1.7	Solved integrals histogram based on CPU time used . . . . .	12
1.8	Leaf size vs. CPU time used . . . . .	13
1.9	list of integrals with no known antiderivative . . . . .	14
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	14
1.11	list of integrals solved by CAS but failed verification . . . . .	14
1.12	Timing . . . . .	15
1.13	Verification . . . . .	15
1.14	Important notes about some of the results . . . . .	15
1.15	Design of the test system . . . . .	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 63 ]. This is test number [ 99 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	100.00 ( 63 )	0.00 ( 0 )
Fricas	100.00 ( 63 )	0.00 ( 0 )
Rubi	98.41 ( 62 )	1.59 ( 1 )
Maple	92.06 ( 58 )	7.94 ( 5 )
Maxima	77.78 ( 49 )	22.22 ( 14 )
Giac	55.56 ( 35 )	44.44 ( 28 )
Mupad	50.79 ( 32 )	49.21 ( 31 )
Sympy	44.44 ( 28 )	55.56 ( 35 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	69.841	0.000	0.000	30.159
Mathematica	58.730	12.698	0.000	28.571
Maple	44.444	19.048	0.000	36.508
Fricas	39.683	31.746	0.000	28.571
Giac	19.048	7.937	0.000	73.016
Sympy	15.873	0.000	0.000	84.127
Maxima	12.698	36.508	0.000	50.794
Mupad	0.000	22.222	0.000	77.778

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Rubi	1	100.00	0.00	0.00
Maple	5	100.00	0.00	0.00
Maxima	14	35.71	0.00	64.29
Giac	28	100.00	0.00	0.00
Mupad	31	0.00	100.00	0.00
Sympy	35	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.



System	Mean time (sec)
Fricas	0.25
Rubi	0.63
Maple	0.64
Sympy	1.09
Maxima	1.89
Giac	3.52
Mupad	3.64
Mathematica	5.07

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	91.44	1.30	22.00	1.11
Sympy	143.93	1.25	19.00	1.00
Rubi	175.84	1.03	134.00	1.00
Fricas	275.24	1.77	130.00	1.58
Giac	295.46	1.58	23.00	1.10
Mathematica	300.63	1.46	171.00	1.17
Maple	433.60	1.73	118.00	1.00
Maxima	910.49	15.56	424.00	4.78

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

# 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{4, 5, 9, 10, 14, 15, 34, 35, 42, 43, 47, 48, 52, 53, 57, 58, 62, 63}

## 1.10 List of integrals solved by CAS but has no known antiderivative

**Rubi** {}

**Mathematica** {}

**Maple** {}

**Maxima** {}

**Fricas** {}

**Sympy** {}

**Giac** {}

**Mupad** {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

**Rubi** {}

**Mathematica** {44, 45, 49, 50, 59}

**Maple** {}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### 1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.



The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### 1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### 1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

### 1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

## 1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
June 27, 2013  
Design v0.01

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	21
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	24
2.3	Detailed conclusion table specific for Rubi results . . . . .	40

## 2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi . . . . .	21
2.1.2	Mma . . . . .	21
2.1.3	Maple . . . . .	22
2.1.4	Fricas . . . . .	22
2.1.5	Maxima . . . . .	22
2.1.6	Giac . . . . .	23
2.1.7	Mupad . . . . .	23
2.1.8	Sympy . . . . .	23

### 2.1.1 Rubi

**A grade** { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

**B grade** { }

**C grade** { }

**F normal fail** { 17 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.2 Mma

**A grade** { 1, 2, 3, 6, 8, 11, 12, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 40, 41, 46, 51, 54, 55, 56, 60 }

**B grade** { 7, 39, 44, 45, 49, 50, 59, 61 }

**C grade** { }

**F normal fail** { }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.3 Maple

**A grade** { 1, 2, 3, 6, 7, 8, 11, 12, 13, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 41, 46, 51 }

**B grade** { 39, 40, 44, 45, 49, 50, 54, 55, 56, 59, 60, 61 }

**C grade** { }

**F normal fail** { 16, 17, 36, 37, 38 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.4 Fricas

**A grade** { 8, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 46, 50, 51 }

**B grade** { 1, 2, 3, 6, 7, 11, 12, 13, 39, 40, 41, 44, 45, 49, 54, 55, 56, 59, 60, 61 }

**C grade** { }

**F normal fail** { }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.5 Maxima

**A grade** { 21, 22, 23, 27, 28, 32, 33, 41 }

**B grade** { 1, 2, 3, 6, 7, 8, 11, 12, 13, 39, 40, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

**C grade** { }

**F normal fail** { 16, 17, 36, 37, 38 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { 18, 19, 20, 24, 25, 26, 29, 30, 31 }

### 2.1.6 Giac

**A grade** { 18, 19, 20, 21, 24, 25, 26, 27, 29, 30, 31, 32 }

**B grade** { 8, 22, 23, 28, 33 }

**C grade** { }

**F normal fail** { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.7 Mupad

**A grade** { }

**B grade** { 3, 8, 16, 17, 18, 19, 20, 24, 25, 26, 29, 30, 31, 41 }

**C grade** { }

**F normal fail** { }

**F(-1) timeout fail** { 1, 2, 6, 7, 11, 12, 13, 21, 22, 23, 27, 28, 32, 33, 36, 37, 38, 39, 40, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

**F(-2) exception fail** { }

### 2.1.8 Sympy

**A grade** { 8, 18, 19, 20, 24, 25, 26, 29, 30, 31 }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 21, 22, 23, 27, 28, 32, 33, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }



## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column N.S. means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	133	106	125	243	286	0	0	0
N.S.	1	1.25	1.00	1.18	2.29	2.70	0.00	0.00	0.00
time (sec)	N/A	0.611	0.013	0.490	0.600	0.280	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	96	77	103	163	200	0	0	0
N.S.	1	1.25	1.00	1.34	2.12	2.60	0.00	0.00	0.00
time (sec)	N/A	0.465	0.009	0.429	0.682	0.259	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	61	54	78	92	122	0	0	129
N.S.	1	1.13	1.00	1.44	1.70	2.26	0.00	0.00	2.39
time (sec)	N/A	0.321	0.006	0.438	0.397	0.242	0.000	0.000	3.346

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	8	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	0.80	1.20	1.20
time (sec)	N/A	0.180	1.555	0.240	0.446	0.253	0.315	0.340	3.237

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.182	2.420	0.205	0.452	0.237	0.281	0.337	3.065

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	122	115	133	639	225	0	0	0
N.S.	1	1.24	1.17	1.36	6.52	2.30	0.00	0.00	0.00
time (sec)	N/A	0.587	0.686	0.632	0.397	0.256	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	73	87	160	108	257	144	0	0	0
N.S.	1	1.19	2.19	1.48	3.52	1.97	0.00	0.00	0.00
time (sec)	N/A	0.435	5.051	0.550	0.631	0.247	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	43	34	214	38	41	162	35
N.S.	1	1.00	1.43	1.13	7.13	1.27	1.37	5.40	1.17
time (sec)	N/A	0.245	0.146	0.459	0.524	0.251	0.216	0.604	0.163

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	243	14	10	14	14
N.S.	1	1.00	1.17	1.00	20.25	1.17	0.83	1.17	1.17
time (sec)	N/A	0.195	3.216	0.339	0.956	0.242	0.309	0.633	3.566

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	250	14	12	14	14
N.S.	1	1.00	1.17	1.00	20.83	1.17	1.00	1.17	1.17
time (sec)	N/A	0.197	2.442	0.311	0.447	0.237	0.287	0.671	2.928

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	245	370	251	1205	344	0	0	0
N.S.	1	1.20	1.80	1.22	5.88	1.68	0.00	0.00	0.00
time (sec)	N/A	1.300	6.536	0.355	0.471	0.274	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	149	172	180	736	240	0	0	0
N.S.	1	1.16	1.34	1.41	5.75	1.88	0.00	0.00	0.00
time (sec)	N/A	0.849	2.281	0.299	0.659	0.261	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	98	171	122	386	146	0	0	0
N.S.	1	1.09	1.90	1.36	4.29	1.62	0.00	0.00	0.00
time (sec)	N/A	0.480	4.577	0.271	0.783	0.257	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	534	14	10	14	14
N.S.	1	1.00	1.17	1.00	44.50	1.17	0.83	1.17	1.17
time (sec)	N/A	0.194	4.854	0.174	0.913	0.242	0.334	0.921	3.176

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	536	14	12	14	14
N.S.	1	1.00	1.17	1.00	44.67	1.17	1.00	1.17	1.17
time (sec)	N/A	0.194	2.584	0.188	0.713	0.242	0.284	0.962	2.826

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	18	0	0	16	0	0	50
N.S.	1	1.00	1.00	0.00	0.00	0.89	0.00	0.00	2.78
time (sec)	N/A	2.908	1.811	0.000	0.000	0.262	0.000	0.000	3.610

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	<b>F</b>	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	B
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	17	0	17	0	0	15	0	0	45
N.S.	1	0.00	1.00	0.00	0.00	0.88	0.00	0.00	2.65
time (sec)	N/A	0.000	0.916	0.000	0.000	0.250	0.000	0.000	5.453

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	189	206	278	170	0	164	258	188	423
N.S.	1	1.09	1.47	0.90	0.00	0.87	1.37	0.99	2.24
time (sec)	N/A	0.704	0.802	0.674	0.000	0.255	0.263	0.382	4.118

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	147	178	108	0	105	165	119	241
N.S.	1	1.07	1.30	0.79	0.00	0.77	1.20	0.87	1.76
time (sec)	N/A	0.481	0.493	0.520	0.000	0.251	0.190	0.359	3.706

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	90	96	50	0	55	92	62	105
N.S.	1	1.07	1.14	0.60	0.00	0.65	1.10	0.74	1.25
time (sec)	N/A	0.311	0.741	0.471	0.000	0.243	0.148	0.353	3.446

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	161	152	166	65	112	51	0	138	0
N.S.	1	0.94	1.03	0.40	0.70	0.32	0.00	0.86	0.00
time (sec)	N/A	0.731	0.610	0.601	0.282	0.252	0.000	0.371	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	167	224	96	119	83	0	1013	0
N.S.	1	0.99	1.33	0.57	0.71	0.49	0.00	6.03	0.00
time (sec)	N/A	0.745	0.767	0.508	0.585	0.259	0.000	2.299	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	227	225	285	216	158	130	0	532	0
N.S.	1	0.99	1.26	0.95	0.70	0.57	0.00	2.34	0.00
time (sec)	N/A	0.967	1.000	0.569	0.595	0.250	0.000	0.442	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	270	270	473	283	0	267	665	368	289
N.S.	1	1.00	1.75	1.05	0.00	0.99	2.46	1.36	1.07
time (sec)	N/A	0.532	2.541	0.746	0.000	0.251	0.415	0.434	4.301

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	202	202	282	173	0	163	418	217	183
N.S.	1	1.00	1.40	0.86	0.00	0.81	2.07	1.07	0.91
time (sec)	N/A	0.437	1.612	0.780	0.000	0.245	0.329	0.449	3.651

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	151	130	82	0	79	226	101	103
N.S.	1	1.00	0.86	0.54	0.00	0.52	1.50	0.67	0.68
time (sec)	N/A	0.388	1.070	0.801	0.000	0.248	0.258	0.416	3.130

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	305	305	211	114	194	84	0	404	0
N.S.	1	1.00	0.69	0.37	0.64	0.28	0.00	1.32	0.00
time (sec)	N/A	1.046	0.848	0.871	0.304	0.244	0.000	0.417	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	436	436	467	175	211	142	0	1967	0
N.S.	1	1.00	1.07	0.40	0.48	0.33	0.00	4.51	0.00
time (sec)	N/A	1.049	1.672	0.892	0.480	0.278	0.000	11.298	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	396	396	667	396	0	370	945	548	411
N.S.	1	1.00	1.68	1.00	0.00	0.93	2.39	1.38	1.04
time (sec)	N/A	0.670	3.764	1.209	0.000	0.245	0.538	0.580	4.249

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	294	294	405	238	0	221	588	315	263
N.S.	1	1.00	1.38	0.81	0.00	0.75	2.00	1.07	0.89
time (sec)	N/A	0.540	2.736	1.082	0.000	0.248	0.504	0.619	3.949

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	208	205	114	0	105	311	142	146
N.S.	1	1.00	0.98	0.55	0.00	0.50	1.49	0.68	0.70
time (sec)	N/A	0.493	1.516	0.884	0.000	0.246	0.343	0.559	3.940



Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	449	449	336	163	276	117	0	810	0
N.S.	1	1.00	0.75	0.36	0.61	0.26	0.00	1.80	0.00
time (sec)	N/A	1.980	1.137	0.934	0.315	0.255	0.000	0.584	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	712	712	833	254	297	197	0	2915	0
N.S.	1	1.00	1.17	0.36	0.42	0.28	0.00	4.09	0.00
time (sec)	N/A	2.063	2.853	1.063	0.431	0.247	0.000	30.478	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	520	115	51	23	24
N.S.	1	1.00	1.09	0.91	22.61	5.00	2.22	1.00	1.04
time (sec)	N/A	0.225	42.842	0.192	1.833	0.250	5.317	0.641	3.372

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	21	21	23	19	79	34	29	21	22
N.S.	1	1.00	1.10	0.90	3.76	1.62	1.38	1.00	1.05
time (sec)	N/A	0.211	9.865	0.163	0.500	0.242	2.400	0.357	3.337

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	98	98	133	0	0	85	0	0	0
N.S.	1	1.00	1.36	0.00	0.00	0.87	0.00	0.00	0.00
time (sec)	N/A	0.335	1.771	0.000	0.000	0.085	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	171	171	192	0	0	144	0	0	0
N.S.	1	1.00	1.12	0.00	0.00	0.84	0.00	0.00	0.00
time (sec)	N/A	0.432	4.489	0.000	0.000	0.087	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	251	251	269	0	0	198	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	0.79	0.00	0.00	0.00
time (sec)	N/A	0.507	13.029	0.000	0.000	0.087	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	152	342	500	672	500	0	0	0
N.S.	1	1.00	2.25	3.29	4.42	3.29	0.00	0.00	0.00
time (sec)	N/A	0.463	0.146	0.641	0.715	0.274	0.000	0.000	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	115	191	314	373	317	0	0	0
N.S.	1	1.00	1.66	2.73	3.24	2.76	0.00	0.00	0.00
time (sec)	N/A	0.413	0.080	0.583	0.520	0.261	0.000	0.000	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	87	143	130	160	0	0	161
N.S.	1	1.00	1.04	1.70	1.55	1.90	0.00	0.00	1.92
time (sec)	N/A	0.321	0.013	0.547	0.377	0.260	0.000	0.000	3.597

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	86	20	15	20	20
N.S.	1	1.00	1.11	1.00	4.78	1.11	0.83	1.11	1.11
time (sec)	N/A	0.208	1.914	0.217	0.440	0.243	0.663	0.389	3.534

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	142	31	17	20	20
N.S.	1	1.00	1.11	1.00	7.89	1.72	0.94	1.11	1.11
time (sec)	N/A	0.206	5.096	0.220	0.463	0.242	2.494	4.371	3.676

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	300	300	1337	952	2525	759	0	0	0
N.S.	1	1.00	4.46	3.17	8.42	2.53	0.00	0.00	0.00
time (sec)	N/A	0.764	7.326	1.603	1.726	0.276	0.000	0.000	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	229	229	649	575	1263	450	0	0	0
N.S.	1	1.00	2.83	2.51	5.52	1.97	0.00	0.00	0.00
time (sec)	N/A	0.601	6.833	1.389	0.912	0.264	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	135	200	238	527	219	0	0	0
N.S.	1	0.99	1.47	1.75	3.88	1.61	0.00	0.00	0.00
time (sec)	N/A	0.393	7.410	1.308	0.409	0.258	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	491	36	17	22	22
N.S.	1	1.00	1.10	1.00	24.55	1.80	0.85	1.10	1.10
time (sec)	N/A	0.224	19.689	0.366	0.827	0.252	1.091	0.565	4.172

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	609	47	19	22	22
N.S.	1	1.00	1.10	1.00	30.45	2.35	0.95	1.10	1.10
time (sec)	N/A	0.228	14.033	0.388	1.683	0.254	2.107	15.426	3.644

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	612	612	2594	1930	6861	1177	0	0	0
N.S.	1	1.00	4.24	3.15	11.21	1.92	0.00	0.00	0.00
time (sec)	N/A	1.285	7.568	1.210	10.883	0.268	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	436	427	1846	1138	3407	688	0	0	0
N.S.	1	0.98	4.23	2.61	7.81	1.58	0.00	0.00	0.00
time (sec)	N/A	0.912	7.164	1.058	1.972	0.290	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	277	275	277	493	1319	327	0	0	0
N.S.	1	0.99	1.00	1.78	4.76	1.18	0.00	0.00	0.00
time (sec)	N/A	0.594	8.323	0.915	0.670	0.266	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1935	52	17	22	22
N.S.	1	1.00	1.10	1.00	96.75	2.60	0.85	1.10	1.10
time (sec)	N/A	0.222	14.685	0.621	3.085	0.251	1.330	1.075	3.911

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	2194	63	19	22	22
N.S.	1	1.00	1.10	1.00	109.70	3.15	0.95	1.10	1.10
time (sec)	N/A	0.225	17.927	0.760	9.084	0.253	2.439	18.056	4.929

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	246	297	1468	983	1183	0	0	0
N.S.	1	1.01	1.22	6.04	4.05	4.87	0.00	0.00	0.00
time (sec)	N/A	0.972	1.676	0.780	0.568	0.282	0.000	0.000	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	188	236	940	715	834	0	0	0
N.S.	1	1.04	1.30	5.19	3.95	4.61	0.00	0.00	0.00
time (sec)	N/A	0.746	1.426	0.693	0.553	0.272	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	132	177	462	399	527	0	0	0
N.S.	1	1.06	1.42	3.70	3.19	4.22	0.00	0.00	0.00
time (sec)	N/A	0.478	1.560	0.637	0.603	0.272	0.000	0.000	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	279	27	17	22	22
N.S.	1	1.00	1.10	1.00	13.95	1.35	0.85	1.10	1.10
time (sec)	N/A	0.232	2.414	0.424	0.890	0.239	1.022	0.417	3.963

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	424	51	19	22	22
N.S.	1	1.00	1.10	1.00	21.20	2.55	0.95	1.10	1.10
time (sec)	N/A	0.227	4.558	0.317	1.726	0.252	1.759	4.307	4.091

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	848	848	1724	5959	4631	2537	0	0	0
N.S.	1	1.00	2.03	7.03	5.46	2.99	0.00	0.00	0.00
time (sec)	N/A	2.248	9.637	0.967	2.409	0.327	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	654	654	761	3901	2530	1576	0	0	0
N.S.	1	1.00	1.16	5.96	3.87	2.41	0.00	0.00	0.00
time (sec)	N/A	1.809	7.529	0.969	1.023	0.286	0.000	0.000	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	228	745	2001	1156	843	0	0	0
N.S.	1	1.07	3.48	9.35	5.40	3.94	0.00	0.00	0.00
time (sec)	N/A	0.771	7.889	0.862	0.705	0.268	0.000	0.000	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1498	55	19	22	22
N.S.	1	1.00	1.10	1.00	74.90	2.75	0.95	1.10	1.10
time (sec)	N/A	0.233	13.937	0.550	8.731	0.252	1.670	0.541	4.538

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1977	96	20	22	22
N.S.	1	1.00	1.10	1.00	98.85	4.80	1.00	1.10	1.10
time (sec)	N/A	0.234	15.452	0.652	27.223	0.248	3.181	22.558	3.988



## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [11] had the largest ratio of [1.3333299999999999]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	7	1.25	10	0.700
2	A	7	6	1.25	10	0.600
3	A	6	5	1.13	8	0.625
4	N/A	2	0	1.00	10	0.000
5	N/A	2	0	1.00	10	0.000
6	A	10	9	1.24	12	0.750
7	A	9	8	1.19	12	0.667
8	A	5	5	1.00	10	0.500
9	N/A	2	0	1.00	12	0.000
10	N/A	2	0	1.00	12	0.000
11	A	17	16	1.20	12	1.333
12	A	13	12	1.16	12	1.000
13	A	10	9	1.09	10	0.900
14	N/A	2	0	1.00	12	0.000
15	N/A	2	0	1.00	12	0.000
16	A	1	1	1.00	45	0.022
17	F	0	0	N/A	0.000	N/A
18	A	9	9	1.09	23	0.391
19	A	7	7	1.07	23	0.304
20	A	5	5	1.07	21	0.238
21	A	7	7	0.94	23	0.304

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	7	7	0.99	23	0.304
23	A	9	9	0.99	23	0.391
24	A	3	3	1.00	23	0.130
25	A	3	3	1.00	23	0.130
26	A	3	3	1.00	21	0.143
27	A	3	3	1.00	23	0.130
28	A	3	3	1.00	23	0.130
29	A	3	3	1.00	23	0.130
30	A	3	3	1.00	23	0.130
31	A	3	3	1.00	21	0.143
32	A	3	3	1.00	23	0.130
33	A	3	3	1.00	23	0.130
34	N/A	2	0	1.00	23	0.000
35	N/A	2	0	1.00	21	0.000
36	A	3	3	1.00	23	0.130
37	A	3	3	1.00	23	0.130
38	A	3	3	1.00	23	0.130
39	A	3	3	1.00	18	0.167
40	A	3	3	1.00	18	0.167
41	A	3	3	1.00	16	0.188
42	N/A	2	0	1.00	18	0.000
43	N/A	2	0	1.00	18	0.000
44	A	3	3	1.00	20	0.150
45	A	3	3	1.00	20	0.150
46	A	3	3	0.99	18	0.167
47	N/A	2	0	1.00	20	0.000
48	N/A	2	0	1.00	20	0.000
49	A	3	3	1.00	20	0.150
50	A	3	3	0.98	20	0.150
51	A	3	3	0.99	18	0.167
52	N/A	2	0	1.00	20	0.000
53	N/A	2	0	1.00	20	0.000
54	A	8	7	1.01	20	0.350

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
55	A	7	6	1.04	20	0.300
56	A	6	5	1.06	18	0.278
57	N/A	2	0	1.00	20	0.000
58	N/A	2	0	1.00	20	0.000
59	A	3	3	1.00	20	0.150
60	A	3	3	1.00	20	0.150
61	A	8	7	1.07	18	0.389
62	N/A	2	0	1.00	20	0.000
63	N/A	2	0	1.00	20	0.000

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int x^3 \tan(a + bx) dx$ . . . . .	45
3.2	$\int x^2 \tan(a + bx) dx$ . . . . .	52
3.3	$\int x \tan(a + bx) dx$ . . . . .	58
3.4	$\int \frac{\tan(a+bx)}{x} dx$ . . . . .	64
3.5	$\int \frac{\tan(a+bx)}{x^2} dx$ . . . . .	68
3.6	$\int x^3 \tan^2(a + bx) dx$ . . . . .	72
3.7	$\int x^2 \tan^2(a + bx) dx$ . . . . .	79
3.8	$\int x \tan^2(a + bx) dx$ . . . . .	85
3.9	$\int \frac{\tan^2(a+bx)}{x} dx$ . . . . .	90
3.10	$\int \frac{\tan^2(a+bx)}{x^2} dx$ . . . . .	95
3.11	$\int x^3 \tan^3(a + bx) dx$ . . . . .	100
3.12	$\int x^2 \tan^3(a + bx) dx$ . . . . .	111
3.13	$\int x \tan^3(a + bx) dx$ . . . . .	119
3.14	$\int \frac{\tan^3(a+bx)}{x} dx$ . . . . .	125
3.15	$\int \frac{\tan^3(a+bx)}{x^2} dx$ . . . . .	130
3.16	$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2 \sqrt{\tan(a + bx)} \right) dx$ . . . . .	135
3.17	$\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx$ . . . . .	139
3.18	$\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx$ . . . . .	143
3.19	$\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx$ . . . . .	150
3.20	$\int \frac{c+dx}{a+ia \tan(e+fx)} dx$ . . . . .	156
3.21	$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx$ . . . . .	161
3.22	$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$ . . . . .	167
3.23	$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$ . . . . .	174
3.24	$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^2} dx$ . . . . .	182
3.25	$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^2} dx$ . . . . .	189
3.26	$\int \frac{c+dx}{(a+ia \tan(e+fx))^2} dx$ . . . . .	195

3.27	$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$	200
3.28	$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx$	206
3.29	$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx$	213
3.30	$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^3} dx$	221
3.31	$\int \frac{c+dx}{(a+ia \tan(e+fx))^3} dx$	228
3.32	$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$	234
3.33	$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx$	241
3.34	$\int (c+dx)^m (a+ia \tan(e+fx))^2 dx$	250
3.35	$\int (c+dx)^m (a+ia \tan(e+fx)) dx$	255
3.36	$\int \frac{(c+dx)^m}{a+ia \tan(e+fx)} dx$	259
3.37	$\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^2} dx$	264
3.38	$\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^3} dx$	269
3.39	$\int (c+dx)^3 (a+b \tan(e+fx)) dx$	274
3.40	$\int (c+dx)^2 (a+b \tan(e+fx)) dx$	281
3.41	$\int (c+dx) (a+b \tan(e+fx)) dx$	287
3.42	$\int \frac{a+b \tan(e+fx)}{c+dx} dx$	292
3.43	$\int \frac{a+b \tan(e+fx)}{(c+dx)^2} dx$	296
3.44	$\int (c+dx)^3 (a+b \tan(e+fx))^2 dx$	300
3.45	$\int (c+dx)^2 (a+b \tan(e+fx))^2 dx$	308
3.46	$\int (c+dx) (a+b \tan(e+fx))^2 dx$	316
3.47	$\int \frac{(a+b \tan(e+fx))^2}{c+dx} dx$	321
3.48	$\int \frac{(a+b \tan(e+fx))^2}{(c+dx)^2} dx$	326
3.49	$\int (c+dx)^3 (a+b \tan(e+fx))^3 dx$	331
3.50	$\int (c+dx)^2 (a+b \tan(e+fx))^3 dx$	341
3.51	$\int (c+dx) (a+b \tan(e+fx))^3 dx$	350
3.52	$\int \frac{(a+b \tan(e+fx))^3}{c+dx} dx$	356
3.53	$\int \frac{(a+b \tan(e+fx))^3}{(c+dx)^2} dx$	361
3.54	$\int \frac{(c+dx)^3}{a+b \tan(e+fx)} dx$	366
3.55	$\int \frac{(c+dx)^2}{a+b \tan(e+fx)} dx$	375
3.56	$\int \frac{c+dx}{a+b \tan(e+fx)} dx$	383
3.57	$\int \frac{1}{(c+dx)(a+b \tan(e+fx))} dx$	389
3.58	$\int \frac{1}{(c+dx)^2(a+b \tan(e+fx))} dx$	394
3.59	$\int \frac{(c+dx)^3}{(a+b \tan(e+fx))^2} dx$	399
3.60	$\int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx$	408
3.61	$\int \frac{c+dx}{(a+b \tan(e+fx))^2} dx$	417
3.62	$\int \frac{1}{(c+dx)(a+b \tan(e+fx))^2} dx$	427
3.63	$\int \frac{1}{(c+dx)^2(a+b \tan(e+fx))^2} dx$	432

## 3.1 $\int x^3 \tan(a + bx) dx$

3.1.1	Optimal result . . . . .	45
3.1.2	Mathematica [A] (verified) . . . . .	45
3.1.3	Rubi [A] (verified) . . . . .	46
3.1.4	Maple [A] (verified) . . . . .	49
3.1.5	Fricas [B] (verification not implemented) . . . . .	49
3.1.6	Sympy [F] . . . . .	50
3.1.7	Maxima [B] (verification not implemented) . . . . .	50
3.1.8	Giac [F] . . . . .	51
3.1.9	Mupad [F(-1)] . . . . .	51

### 3.1.1 Optimal result

Integrand size = 10, antiderivative size = 106

$$\int x^3 \tan(a + bx) dx = \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{3x \text{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3i \text{PolyLog}(4, -e^{2i(a+bx)})}{4b^4}$$

output  $\frac{1}{4}ix^4 - x^3 \ln(1 + \exp(2i(bx+a))) / b + 3/2ix^2 \text{polylog}(2, -\exp(2i(bx+a))) / b^2 - 3/2x \text{polylog}(3, -\exp(2i(bx+a))) / b^3 - 3/4i \text{polylog}(4, -\exp(2i(bx+a))) / b^4$

### 3.1.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00

$$\int x^3 \tan(a + bx) dx = \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{3x \text{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3i \text{PolyLog}(4, -e^{2i(a+bx)})}{4b^4}$$

input `Integrate[x^3*Tan[a + b*x],x]`

output  $(I/4)x^4 - (x^3 \text{Log}[1 + E^{((2I)*(a + b*x)}))] / b + (((3I)/2)x^2 \text{PolyLog}[2, -E^{((2I)*(a + b*x)})]) / b^2 - (3*x \text{PolyLog}[3, -E^{((2I)*(a + b*x)})]) / (2*b^3) - (((3I)/4) \text{PolyLog}[4, -E^{((2I)*(a + b*x)})]) / b^4$

### 3.1.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.25, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$ , Rules used = {3042, 4202, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \tan(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^3 \tan(a + bx) dx \\
 & \quad \downarrow \text{4202} \\
 & \frac{ix^4}{4} - 2i \int \frac{e^{2i(a+bx)} x^3}{1 + e^{2i(a+bx)}} dx \\
 & \quad \downarrow \text{2620} \\
 & \frac{ix^4}{4} - 2i \left( \frac{3i \int x^2 \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{3011} \\
 & \frac{ix^4}{4} - 2i \left( \frac{3i \left( \frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{7163} \\
 & \frac{ix^4}{4} - 2i \left( \frac{3i \left( \frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left( \frac{i \int \text{PolyLog}(3, -e^{2i(a+bx)}) dx}{2b} - \frac{ix \text{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{2720}
 \end{aligned}$$

$$2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left( \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(3, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right)$$

↓ 7143

$$2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left( \frac{\operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right)$$

input `Int[x^3*Tan[a + b*x],x]`

output `(I/4)*x^4 - (2*I)*((-1/2*I)*x^3*Log[1 + E^((2*I)*(a + b*x))])/b + ((3*I)/2)*((I/2)*x^2*PolyLog[2, -E^((2*I)*(a + b*x))])/b - (I*((-1/2*I)*x*PolyLog[3, -E^((2*I)*(a + b*x))])/b + PolyLog[4, -E^((2*I)*(a + b*x))]/(4*b^2))/b)/b)`

### 3.1.3.1 Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`



rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4202 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x)))], x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_) + (f_)*(x_))^(m_)*PolyLog[n_, (d_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(p_)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### 3.1.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.18

method	result
risch	$\frac{ix^4}{4} - \frac{3x \operatorname{Li}_3(-e^{2i(bx+a)})}{2b^3} + \frac{2ia^3x}{b^3} + \frac{3ix^2 \operatorname{Li}_2(-e^{2i(bx+a)})}{2b^2} + \frac{3ia^4}{2b^4} - \frac{3i \operatorname{Li}_4(-e^{2i(bx+a)})}{4b^4} - \frac{2a^3 \ln(e^{i(bx+a)})}{b^4} - \frac{x^3 \ln(e^{2i(bx+a)})}{b^4}$

input `int(x^3*tan(b*x+a),x,method=_RETURNVERBOSE)`

output  $\frac{1}{4}Ix^4 - \frac{3}{2}x \operatorname{polylog}(3, -\exp(2I*(b*x+a)))/b^3 + 2I/b^3 * a^3 * x + \frac{3}{2}Ix^2 \operatorname{polylog}(2, -\exp(2I*(b*x+a)))/b^2 + 3/2I/b^4 * a^4 - 3/4I * \operatorname{polylog}(4, -\exp(2I*(b*x+a)))/b^4 - 2/b^4 * a^3 * \ln(\exp(I*(b*x+a))) - x^3 * \ln(\exp(2I*(b*x+a))+1)/b$

### 3.1.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 286 vs.  $2(83) = 166$ .

Time = 0.28 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.70

$$\int x^3 \tan(a + bx) dx = \frac{4b^3x^3 \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 4b^3x^3 \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 6ib^2x^2 \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - 6ib^2x^2 \operatorname{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right)}{b^4}$$

input `integrate(x^3*tan(b*x+a),x, algorithm="fricas")`

output  $-1/8*(4*b^3*x^3*\log(-2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 4*b^3*x^3*\log(-2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 6*I*b^2*x^2*dilog(2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1) - 6*I*b^2*x^2*dilog(2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1) + 6*b*x*polylog(3, (\tan(b*x + a)^2 + 2*I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 6*b*x*polylog(3, (\tan(b*x + a)^2 - 2*I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 3*I*polylog(4, (\tan(b*x + a)^2 + 2*I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 3*I*polylog(4, (\tan(b*x + a)^2 - 2*I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)))/b^4$

### 3.1.6 Sympy [F]

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(a + bx) dx$$

input `integrate(x**3*tan(b*x+a),x)`

output `Integral(x**3*tan(a + b*x), x)`

### 3.1.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 243 vs.  $2(83) = 166$ .

Time = 0.60 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.29

$$\int x^3 \tan(a + bx) dx = \frac{-3i (bx + a)^4 + 12i (bx + a)^3 a - 18i (bx + a)^2 a^2 + 12 a^3 \log(\sec(bx + a)) - 4(-4i (bx + a)^3 + 9i (bx + a)^2 a - 9i (bx + a) a^2) \arctan2(\sin(2bx + 2a), \cos(2bx + 2a) + 1) - 6(4i (bx + a)^2 - 6i (bx + a) a + 3i a^2) \operatorname{dilog}(-e^{(2I*bx + 2I*a)}) + 2(4(bx + a)^3 - 9(bx + a)^2 a + 9(bx + a) a^2) \log(\cos(2bx + 2a)^2 + \sin(2bx + 2a)^2 + 2\cos(2bx + 2a) + 1) + 6(4bx + a) \operatorname{polylog}(3, -e^{(2I*bx + 2I*a)}) + 12i \operatorname{polylog}(4, -e^{(2I*bx + 2I*a)})}{b^4}$$

input `integrate(x^3*tan(b*x+a),x, algorithm="maxima")`

output `-1/12*(-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b*x + a)^2*a^2 + 12*a^3*log(sec(b*x + a)) - 4*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a - 9*I*(b*x + a)*a^2)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) - 6*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilog(-e^(2*I*b*x + 2*I*a)) + 2*(4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(b*x + a)*a^2)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 6*(4*b*x + a)*polylog(3, -e^(2*I*b*x + 2*I*a)) + 12*I*polylog(4, -e^(2*I*b*x + 2*I*a)))/b^4`

### 3.1.8 Giac [F]

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(bx + a) dx$$

input `integrate(x^3*tan(b*x+a),x, algorithm="giac")`

output `integrate(x^3*tan(b*x + a), x)`

### 3.1.9 Mupad [F(-1)]

Timed out.

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(a + bx) dx$$

input `int(x^3*tan(a + b*x),x)`

output `int(x^3*tan(a + b*x), x)`

## 3.2 $\int x^2 \tan(a + bx) dx$

3.2.1	Optimal result . . . . .	52
3.2.2	Mathematica [A] (verified) . . . . .	52
3.2.3	Rubi [A] (verified) . . . . .	53
3.2.4	Maple [A] (verified) . . . . .	55
3.2.5	Fricas [B] (verification not implemented) . . . . .	55
3.2.6	Sympy [F] . . . . .	56
3.2.7	Maxima [B] (verification not implemented) . . . . .	56
3.2.8	Giac [F] . . . . .	56
3.2.9	Mupad [F(-1)] . . . . .	57

### 3.2.1 Optimal result

Integrand size = 10, antiderivative size = 77

$$\int x^2 \tan(a + bx) dx = \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3}$$

output  $1/3*I*x^3-x^2*\ln(1+\exp(2*I*(b*x+a)))/b+I*x*polylog(2,-\exp(2*I*(b*x+a)))/b^2-1/2*polylog(3,-\exp(2*I*(b*x+a)))/b^3$

### 3.2.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00

$$\int x^2 \tan(a + bx) dx = \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3}$$

input `Integrate[x^2*Tan[a + b*x],x]`

output  $(I/3)*x^3 - (x^2*\Log[1 + E^((2*I)*(a + b*x))])/b + (I*x*PolyLog[2, -E^((2*I)*(a + b*x))])/b^2 - PolyLog[3, -E^((2*I)*(a + b*x))]/(2*b^3)$

### 3.2.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.25, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4202, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \tan(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \tan(a + bx) dx \\
 & \quad \downarrow \text{4202} \\
 & \frac{ix^3}{3} - 2i \int \frac{e^{2i(a+bx)} x^2}{1 + e^{2i(a+bx)}} dx \\
 & \quad \downarrow \text{2620} \\
 & \frac{ix^3}{3} - 2i \left( \frac{i \int x \log(1 + e^{2i(a+bx)}) dx}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{3011} \\
 & \frac{ix^3}{3} - 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{2720} \\
 & \frac{ix^3}{3} - 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{7143} \\
 & \frac{ix^3}{3} - 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right)
 \end{aligned}$$

input `Int[x^2*Tan[a + b*x], x]`

output  $(I/3)*x^3 - (2*I)*((-1/2*I)*x^2*Log[1 + E^((2*I)*(a + b*x))]/b + (I*((I/2)*x*PolyLog[2, -E^((2*I)*(a + b*x))]/b - PolyLog[3, -E^((2*I)*(a + b*x))]/(4*b^2)))/b)$

### 3.2.3.1 Defintions of rubi rules used

rule 2620  $Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^{(n_)*((c_) + (d_)*(x_))^{(m_))}/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^{(n_)}), x\_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^{m-1}*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]$

rule 2720  $Int[u_, x\_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^{(n_)})^{(m)} /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]$

rule 3011  $Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^{(n_)}]*(f_) + (g_)*(x_))^{(m_)}, x\_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^{m-1}*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]$

rule 3042  $Int[u_, x\_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]$

rule 4202  $Int[(((c_) + (d_)*(x_))^{(m_)*tan[(e_) + (f_)*(x_)]}, x\_Symbol] := Simp[I*((c + d*x)^{m+1}/(d*(m+1))), x] - Simp[2*I Int[(c + d*x)^m*(E^{2*I*(e + f*x)})/(1 + E^{2*I*(e + f*x)})], x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]$

rule 7143  $Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^{(p_)}]/((d_) + (e_)*(x_)), x\_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]$

### 3.2.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.34

method	result	size
risch	$\frac{ix^3}{3} - \frac{2ia^2x}{b^2} - \frac{4ia^3}{3b^3} - \frac{x^2 \ln(e^{2i(bx+a)}+1)}{b} + \frac{ix \operatorname{Li}_2(-e^{2i(bx+a)})}{b^2} - \frac{\operatorname{Li}_3(-e^{2i(bx+a)})}{2b^3} + \frac{2a^2 \ln(e^{i(bx+a)})}{b^3}$	103

input `int(x^2*tan(b*x+a),x,method=_RETURNVERBOSE)`

output `1/3*I*x^3-2*I/b^2*a^2*x-4/3*I/b^3*a^3-x^2*ln(exp(2*I*(b*x+a))+1)/b+I*x*polylog(2,-exp(2*I*(b*x+a)))/b^2-1/2*polylog(3,-exp(2*I*(b*x+a)))/b^3+2/b^3*a^2*ln(exp(I*(b*x+a)))`

### 3.2.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 200 vs.  $2(62) = 124$ .

Time = 0.26 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.60

$$\int x^2 \tan(a + bx) dx = \frac{2b^2x^2 \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2b^2x^2 \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2i bx \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - 2i bx \operatorname{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right)}{4b^3}$$

input `integrate(x^2*tan(b*x+a),x, algorithm="fricas")`

output `-1/4*(2*b^2*x^2*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*b^2*x^2*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*I*b*x*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 2*I*b*x*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) + polylog(3, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + polylog(3, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)))/b^3`



### 3.2.6 Sympy [F]

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(a + bx) dx$$

input `integrate(x**2*tan(b*x+a),x)`

output `Integral(x**2*tan(a + b*x), x)`

### 3.2.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 163 vs.  $2(62) = 124$ .

Time = 0.68 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.12

$$\int x^2 \tan(a + bx) dx =$$

$$\frac{-2i (bx + a)^3 + 6i (bx + a)^2 a - 6i bx \operatorname{Li}_2(-e^{(2i bx + 2i a)}) - 6a^2 \log(\sec(bx + a)) - 6(-i(bx + a)^2 + 2i)}{-}$$

input `integrate(x^2*tan(b*x+a),x, algorithm="maxima")`

output `-1/6*(-2*I*(b*x + a)^3 + 6*I*(b*x + a)^2*a - 6*I*b*x*dilog(-e^(2*I*b*x + 2*I*a)) - 6*a^2*log(sec(b*x + a)) - 6*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 3*polylog(3, -e^(2*I*b*x + 2*I*a)))/b^3`

### 3.2.8 Giac [F]

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(bx + a) dx$$

input `integrate(x^2*tan(b*x+a),x, algorithm="giac")`

output `integrate(x^2*tan(b*x + a), x)`

**3.2.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(a + bx) dx$$

input `int(x^2*tan(a + b*x),x)`output `int(x^2*tan(a + b*x), x)`

### 3.3 $\int x \tan(a + bx) dx$

3.3.1	Optimal result . . . . .	58
3.3.2	Mathematica [A] (verified) . . . . .	58
3.3.3	Rubi [A] (verified) . . . . .	59
3.3.4	Maple [A] (verified) . . . . .	60
3.3.5	Fricas [B] (verification not implemented) . . . . .	61
3.3.6	Sympy [F] . . . . .	61
3.3.7	Maxima [B] (verification not implemented) . . . . .	62
3.3.8	Giac [F] . . . . .	62
3.3.9	Mupad [B] (verification not implemented) . . . . .	62

#### 3.3.1 Optimal result

Integrand size = 8, antiderivative size = 54

$$\int x \tan(a + bx) dx = \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2}$$

output  $\frac{1}{2}i x^2 - x \ln(1 + \exp(2i(bx+a))) / b + \frac{1}{2}i \operatorname{polylog}(2, -\exp(2i(bx+a))) / b^2$

#### 3.3.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int x \tan(a + bx) dx = \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2}$$

input `Integrate[x*Tan[a + b*x],x]`

output  $(\frac{I}{2})x^2 - (x \operatorname{Log}[1 + E^{((2I)(a + b*x))}]) / b + ((\frac{I}{2}) \operatorname{PolyLog}[2, -E^{((2I)(a + b*x))}]) / b^2$

### 3.3.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.13, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tan(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \tan(a + bx) dx \\
 & \quad \downarrow \text{4202} \\
 & \frac{ix^2}{2} - 2i \int \frac{e^{2i(a+bx)} x}{1 + e^{2i(a+bx)}} dx \\
 & \quad \downarrow \text{2620} \\
 & \frac{ix^2}{2} - 2i \left( \frac{i \int \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{2715} \\
 & \frac{ix^2}{2} - 2i \left( \frac{\int e^{-2i(a+bx)} \log(1 + e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \quad \downarrow \text{2838} \\
 & \frac{ix^2}{2} - 2i \left( -\frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right)
 \end{aligned}$$

input `Int[x*Tan[a + b*x],x]`

output `(I/2)*x^2 - (2*I)*((( -1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))])/b - PolyLog[2, -E^((2*I)*(a + b*x))]/(4*b^2))`

### 3.3.3.1 Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_)]], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4202 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

### 3.3.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

method	result
risch	$\frac{ix^2}{2} + \frac{2iax}{b} + \frac{ia^2}{b^2} - \frac{x \ln(e^{2i(bx+a)}+1)}{b} + \frac{i \operatorname{Li}_2(-e^{2i(bx+a)})}{2b^2} - \frac{2a \ln(e^{i(bx+a)})}{b^2}$
parts	$\frac{\ln(1+\tan^2(bx+a))x}{2b} - \frac{i \left( \ln(\tan(bx+a)-i) \ln(1+\tan^2(bx+a)) - \operatorname{dilog}\left(-\frac{i(\tan(bx+a)+i)}{2}\right) - \ln(\tan(bx+a)-i) \ln\left(-\frac{i(\tan(bx+a)+i)}{2}\right) - \ln(\tan(bx+a)+i) \ln(1+\tan^2(bx+a)) \right)}{2}$

input `int(x*tan(b*x+a), x, method=_RETURNVERBOSE)`

output  $1/2*I*x^2+2*I/b*a*x+I/b^2*a^2-x*\ln(\exp(2*I*(b*x+a))+1)/b+1/2*I*polylog(2,-\exp(2*I*(b*x+a)))/b^2-2/b^2*a*\ln(\exp(I*(b*x+a)))$

### 3.3.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 122 vs.  $2(41) = 82$ .

Time = 0.24 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.26

$$\int x \tan(a + bx) dx = \frac{2bx \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2bx \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + i \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - i \operatorname{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right)}{4b^2}$$

input `integrate(x*tan(b*x+a),x, algorithm="fricas")`

output  $-1/4*(2*b*x*\log(-2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 2*b*x*\log(-2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + I*dilog(2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1) - I*dilog(2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1))/b^2$

### 3.3.6 Sympy [F]

$$\int x \tan(a + bx) dx = \int x \tan(a + bx) dx$$

input `integrate(x*tan(b*x+a),x)`

output `Integral(x*tan(a + b*x), x)`

### 3.3.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 92 vs.  $2(41) = 82$ .

Time = 0.40 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.70

$$\int x \tan(a + bx) dx = \frac{-i b^2 x^2 + 2i bx \arctan(\sin(2bx + 2a), \cos(2bx + 2a) + 1) + bx \log(\cos(2bx + 2a)^2 + \sin(2bx + 2a) + 1)}{2b^2}$$

input `integrate(x*tan(b*x+a),x, algorithm="maxima")`

output `-1/2*(-I*b^2*x^2 + 2*I*b*x*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + b*x*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - I*dilog(-e^(2*I*b*x + 2*I*a)))/b^2`

### 3.3.8 Giac [F]

$$\int x \tan(a + bx) dx = \int x \tan(bx + a) dx$$

input `integrate(x*tan(b*x+a),x, algorithm="giac")`

output `integrate(x*tan(b*x + a), x)`

### 3.3.9 Mupad [B] (verification not implemented)

Time = 3.35 (sec) , antiderivative size = 129, normalized size of antiderivative = 2.39

$$\int x \tan(a + bx) dx = \frac{\pi \ln(\cos(bx)) + \text{polylog}(2, -e^{-a2i} e^{-bx2i}) \text{li} - \pi \ln(e^{-a2i} e^{-bx2i} + 1) + 2a \ln(e^{-a2i} e^{-bx2i} + 1) - \pi}{2b^2}$$

input `int(x*tan(a + b*x),x)`

output 
$$-(\text{polylog}(2, -\exp(-a*2i)*\exp(-b*x*2i))*1i - \pi*\log(\exp(b*x*2i) + 1) - \pi*\log(\exp(-a*2i)*\exp(-b*x*2i) + 1) + 2*a*\log(\exp(-a*2i)*\exp(-b*x*2i) + 1) + \pi*\log(\cos(b*x)) + b^2*x^2*1i - \log(\cos(a + b*x))*(2*a - \pi) + 2*b*x*\log(\exp(-a*2i)*\exp(-b*x*2i) + 1) + a*b*x*2i)/(2*b^2)$$



### 3.4 $\int \frac{\tan(a+bx)}{x} dx$

3.4.1	Optimal result . . . . .	64
3.4.2	Mathematica [N/A] . . . . .	64
3.4.3	Rubi [N/A] . . . . .	65
3.4.4	Maple [N/A] (verified) . . . . .	66
3.4.5	Fricas [N/A] . . . . .	66
3.4.6	Sympy [N/A] . . . . .	66
3.4.7	Maxima [N/A] . . . . .	67
3.4.8	Giac [N/A] . . . . .	67
3.4.9	Mupad [N/A] . . . . .	67

#### 3.4.1 Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\tan(a+bx)}{x} dx = \text{Int}\left(\frac{\tan(a+bx)}{x}, x\right)$$

output `Unintegrable(tan(b*x+a)/x,x)`

#### 3.4.2 Mathematica [N/A]

Not integrable

Time = 1.55 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a+bx)}{x} dx = \int \frac{\tan(a+bx)}{x} dx$$

input `Integrate[Tan[a + b*x]/x,x]`

output `Integrate[Tan[a + b*x]/x, x]`

### 3.4.3 Rubi [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(a + bx)}{x} dx$$

↓ 3042

$$\int \frac{\tan(a + bx)}{x} dx$$

↓ 4222

$$\int \frac{\tan(a + bx)}{x} dx$$

input `Int[Tan[a + b*x]/x,x]`

output `$Aborted`

#### 3.4.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4222 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], I^n*Unintegrable[(c + d*x)^m*Coth[(-I)*(e - Pi/2) - I*f*x]^n, x], I^n*Unintegrable[(c + d*x)^m*Tanh[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], (-1)^n*Unintegrable[(c + d*x)^m*Cot[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*Tan[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]`

**3.4.4 Maple [N/A] (verified)**

Not integrable

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan (bx + a)}{x} dx$$

input `int(tan(b*x+a)/x,x)`output `int(tan(b*x+a)/x,x)`**3.4.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan (bx + a)}{x} dx$$

input `integrate(tan(b*x+a)/x,x, algorithm="fricas")`output `integral(tan(b*x + a)/x, x)`**3.4.6 Sympy [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan (a + bx)}{x} dx$$

input `integrate(tan(b*x+a)/x,x)`output `Integral(tan(a + b*x)/x, x)`

**3.4.7 Maxima [N/A]**

Not integrable

Time = 0.45 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

input `integrate(tan(b*x+a)/x,x, algorithm="maxima")`output `integrate(tan(b*x + a)/x, x)`**3.4.8 Giac [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

input `integrate(tan(b*x+a)/x,x, algorithm="giac")`output `integrate(tan(b*x + a)/x, x)`**3.4.9 Mupad [N/A]**

Not integrable

Time = 3.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(a + bx)}{x} dx$$

input `int(tan(a + b*x)/x,x)`output `int(tan(a + b*x)/x, x)`

### 3.5 $\int \frac{\tan(a+bx)}{x^2} dx$

3.5.1	Optimal result	68
3.5.2	Mathematica [N/A]	68
3.5.3	Rubi [N/A]	69
3.5.4	Maple [N/A] (verified)	70
3.5.5	Fricas [N/A]	70
3.5.6	Sympy [N/A]	70
3.5.7	Maxima [N/A]	71
3.5.8	Giac [N/A]	71
3.5.9	Mupad [N/A]	71

#### 3.5.1 Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\tan(a+bx)}{x^2} dx = \text{Int}\left(\frac{\tan(a+bx)}{x^2}, x\right)$$

output `Unintegrable(tan(b*x+a)/x^2,x)`

#### 3.5.2 Mathematica [N/A]

Not integrable

Time = 2.42 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a+bx)}{x^2} dx = \int \frac{\tan(a+bx)}{x^2} dx$$

input `Integrate[Tan[a + b*x]/x^2,x]`

output `Integrate[Tan[a + b*x]/x^2, x]`

### 3.5.3 Rubi [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(a + bx)}{x^2} dx$$

↓ 3042

$$\int \frac{\tan(a + bx)}{x^2} dx$$

↓ 4222

$$\int \frac{\tan(a + bx)}{x^2} dx$$

input `Int[Tan[a + b*x]/x^2,x]`

output `$Aborted`

#### 3.5.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4222 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], I^n*Unintegrable[(c + d*x)^m*Coth[(-I)*(e - Pi/2) - I*f*x]^n, x], I^n*Unintegrable[(c + d*x)^m*Tanh[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], (-1)^n*Unintegrable[(c + d*x)^m*Cot[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*Tan[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]`

**3.5.4 Maple [N/A] (verified)**

Not integrable

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan (bx + a)}{x^2} dx$$

input `int(tan(b*x+a)/x^2,x)`output `int(tan(b*x+a)/x^2,x)`**3.5.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan (bx + a)}{x^2} dx$$

input `integrate(tan(b*x+a)/x^2,x, algorithm="fricas")`output `integral(tan(b*x + a)/x^2, x)`**3.5.6 Sympy [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan (a + bx)}{x^2} dx$$

input `integrate(tan(b*x+a)/x**2,x)`output `Integral(tan(a + b*x)/x**2, x)`

**3.5.7 Maxima [N/A]**

Not integrable

Time = 0.45 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

input `integrate(tan(b*x+a)/x^2,x, algorithm="maxima")`output `integrate(tan(b*x + a)/x^2, x)`**3.5.8 Giac [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

input `integrate(tan(b*x+a)/x^2,x, algorithm="giac")`output `integrate(tan(b*x + a)/x^2, x)`**3.5.9 Mupad [N/A]**

Not integrable

Time = 3.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)}{x^2} dx$$

input `int(tan(a + b*x)/x^2,x)`output `int(tan(a + b*x)/x^2, x)`



### 3.6 $\int x^3 \tan^2(a + bx) dx$

3.6.1	Optimal result . . . . .	72
3.6.2	Mathematica [A] (verified) . . . . .	72
3.6.3	Rubi [A] (verified) . . . . .	73
3.6.4	Maple [A] (verified) . . . . .	76
3.6.5	Fricas [B] (verification not implemented) . . . . .	76
3.6.6	Sympy [F] . . . . .	77
3.6.7	Maxima [B] (verification not implemented) . . . . .	77
3.6.8	Giac [F] . . . . .	78
3.6.9	Mupad [F(-1)] . . . . .	78

#### 3.6.1 Optimal result

Integrand size = 12, antiderivative size = 98

$$\int x^3 \tan^2(a + bx) dx = -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} - \frac{3ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} + \frac{3 \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^4} + \frac{x^3 \tan(a + bx)}{b}$$

output `-I*x^3/b-1/4*x^4+3*x^2*ln(1+exp(2*I*(b*x+a)))/b^2-3*I*x*polylog(2,-exp(2*I*(b*x+a)))/b^3+3/2*polylog(3,-exp(2*I*(b*x+a)))/b^4+x^3*tan(b*x+a)/b`

#### 3.6.2 Mathematica [A] (verified)

Time = 0.69 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.17

$$\int x^3 \tan^2(a + bx) dx = -\frac{x^4}{4} + \frac{2b^2 x^2 \left( \frac{2ibx}{1+e^{2ia}} + 3 \log(1 + e^{-2i(a+bx)}) \right) + 6ibx \operatorname{PolyLog}(2, -e^{-2i(a+bx)}) + 3 \operatorname{PolyLog}(3, -e^{-2i(a+bx)})}{2b^4} + \frac{x^3 \sec(a) \sec(a + bx) \sin(bx)}{b}$$

input `Integrate[x^3*Tan[a + b*x]^2,x]`

output 
$$-1/4*x^4 + (2*b^2*x^2*((2*I)*b*x)/(1 + E^((2*I)*a)) + 3*Log[1 + E^((-2*I)*(a + b*x))]) + (6*I)*b*x*PolyLog[2, -E^((-2*I)*(a + b*x))] + 3*PolyLog[3, -E^((-2*I)*(a + b*x))]/(2*b^4) + (x^3*Sec[a]*Sec[a + b*x]*Sin[b*x])/b$$

### 3.6.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.24, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 4203, 15, 3042, 4202, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \tan^2(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^3 \tan(a + bx)^2 dx \\
 & \quad \downarrow \text{4203} \\
 & -\frac{3 \int x^2 \tan(a + bx) dx}{b} - \int x^3 dx + \frac{x^3 \tan(a + bx)}{b} \\
 & \quad \downarrow \text{15} \\
 & -\frac{3 \int x^2 \tan(a + bx) dx}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3 \int x^2 \tan(a + bx) dx}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \\
 & \quad \downarrow \text{4202} \\
 & -\frac{3 \left( \frac{ix^3}{3} - 2i \int \frac{e^{2i(a+bx)} x^2}{1+e^{2i(a+bx)}} dx \right)}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \\
 & \quad \downarrow \text{2620} \\
 & -\frac{3 \left( \frac{ix^3}{3} - 2i \left( \frac{i \int x \log(1+e^{2i(a+bx)}) dx}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{3 \left( \frac{ix^3}{3} - 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int \operatorname{PolyLog}(2, -e^{2i(a+bx)} dx}{b} \right)}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{\frac{x^3 \tan(a+bx)}{b} - \frac{x^4}{4}} + \\
 & \quad \downarrow \text{2720} \\
 & \frac{3 \left( \frac{ix^3}{3} - 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)} de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{\frac{x^3 \tan(a+bx)}{b} - \frac{x^4}{4}} + \\
 & \quad \downarrow \text{7143} \\
 & \frac{3 \left( \frac{ix^3}{3} - 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{4b^2} \right)}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{\frac{x^3 \tan(a+bx)}{b} - \frac{x^4}{4}} +
 \end{aligned}$$

input `Int[x^3*Tan[a + b*x]^2,x]`

output `-1/4*x^4 - (3*((I/3)*x^3 - (2*I)*((( -1/2*I)*x^2*Log[1 + E^((2*I)*(a + b*x))])/b + (I*(((I/2)*x*PolyLog[2, -E^((2*I)*(a + b*x))])/b - PolyLog[3, -E^((2*I)*(a + b*x))]/(4*b^2)))/b))/b + (x^3*Tan[a + b*x])/b`

## 3.6.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_)^(m_.)))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4202 `Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`
- rule 4203 `Int[((c_.) + (d_.)*(x_)^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Simp[b*d*(m/(f*(n - 1))) Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Simp[b^2 Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]`

rule 7143 `Int [PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp [PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.6.4 Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.36

method	result
risch	$-\frac{x^4}{4} + \frac{2ix^3}{b(e^{2i(bx+a)}+1)} - \frac{2ix^3}{b} + \frac{6ia^2x}{b^3} + \frac{4ia^3}{b^4} + \frac{3x^2 \ln(e^{2i(bx+a)}+1)}{b^2} - \frac{3ix \operatorname{Li}_2(-e^{2i(bx+a)})}{b^3} + \frac{3 \operatorname{Li}_3(-e^{2i(bx+a)})}{2b^4} -$

input `int(x^3*tan(b*x+a)^2,x,method=_RETURNVERBOSE)`

output 
$$-1/4*x^4+2*I*x^3/b/(\exp(2*I*(b*x+a))+1)-2*I/b*x^3+6*I/b^3*a^2*x+4*I/b^4*a^3+3*x^2*\ln(\exp(2*I*(b*x+a))+1)/b^2-3*I*x*\operatorname{polylog}(2,-\exp(2*I*(b*x+a)))/b^3+3/2*\operatorname{polylog}(3,-\exp(2*I*(b*x+a)))/b^4-6/b^4*a^2*\ln(\exp(I*(b*x+a)))$$

### 3.6.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 225 vs.  $2(83) = 166$ .

Time = 0.26 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.30

$$\int x^3 \tan^2(a + bx) dx = \frac{b^4 x^4 - 4 b^3 x^3 \tan(bx + a) - 6 b^2 x^2 \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 6 b^2 x^2 \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 6i bx \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 6i bx \operatorname{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right)}{b^4}$$

input `integrate(x^3*tan(b*x+a)^2,x, algorithm="fricas")`

output 
$$-1/4*(b^4*x^4 - 4*b^3*x^3*\tan(b*x + a) - 6*b^2*x^2*\log(-2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 6*b^2*x^2*\log(-2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 6*I*b*x*dilog(2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1) + 6*I*b*x*dilog(2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1) - 3*\operatorname{polylog}(3, (\tan(b*x + a)^2 + 2*I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 3*\operatorname{polylog}(3, (\tan(b*x + a)^2 - 2*I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)))/b^4$$

### 3.6.6 Sympy [F]

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan^2(a + bx) dx$$

input `integrate(x**3*tan(b*x+a)**2,x)`

output `Integral(x**3*tan(a + b*x)**2, x)`

### 3.6.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 639 vs.  $2(83) = 166$ .

Time = 0.40 (sec) , antiderivative size = 639, normalized size of antiderivative = 6.52

$$\int x^3 \tan^2(a + bx) dx = \frac{2(bx + a - \tan(bx + a))a^3 - \frac{3((bx+a)^2 \cos(2bx+2a)^2 + (bx+a)^2 \sin(2bx+2a)^2 + 2(bx+a)^2 \cos(2bx+2a) + (bx+a)^2 - (\cos(2bx+2a) - \sin(2bx+2a))}{\cos(2bx+2a)}}{1}$$

input `integrate(x^3*tan(b*x+a)^2,x, algorithm="maxima")`

output `1/2*(2*(b*x + a - tan(b*x + a))*a^3 - 3*((b*x + a)^2*cos(2*b*x + 2*a)^2 + (b*x + a)^2*sin(2*b*x + 2*a)^2 + 2*(b*x + a)^2*cos(2*b*x + 2*a) + (b*x + a)^2 - (cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - 4*(b*x + a)*sin(2*b*x + 2*a))*a^2/(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 2*(I*(b*x + a)^4 - 4*I*(b*x + a)^3*a + 12*((b*x + a)^2 - 2*(b*x + a)*a + ((b*x + a)^2 - 2*(b*x + a)*a)*cos(2*b*x + 2*a) - (-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*sin(2*b*x + 2*a))*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + (I*(b*x + a)^4 - 4*(b*x + a)^3*(I*a + 2) + 24*(b*x + a)^2*a*cos(2*b*x + 2*a) - 12*(b*x*cos(2*b*x + 2*a) + I*b*x*sin(2*b*x + 2*a) + b*x)*dilog(-e^(2*I*b*x + 2*I*a)) - 6*(I*(b*x + a)^2 - 2*I*(b*x + a)*a + (I*(b*x + a)^2 - 2*I*(b*x + a)*a)*cos(2*b*x + 2*a) - ((b*x + a)^2 - 2*(b*x + a)*a)*sin(2*b*x + 2*a))*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - 6*(I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a) + I)*polylog(3, -e^(2*I*b*x + 2*I*a)) - ((b*x + a)^4 - 4*(b*x + a)^3*(a - 2*I) - 24*I*(b*x + a)^2*a*sin(2*b*x + 2*a))/(-4*I*cos(2*b*x + 2*a) + 4*sin(2*b*x + 2*a) - 4*I))/b^4`

### 3.6.8 Giac [F]

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan(bx + a)^2 dx$$

input `integrate(x^3*tan(b*x+a)^2,x, algorithm="giac")`

output `integrate(x^3*tan(b*x + a)^2, x)`

### 3.6.9 Mupad [F(-1)]

Timed out.

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan(a + bx)^2 dx$$

input `int(x^3*tan(a + b*x)^2,x)`

output `int(x^3*tan(a + b*x)^2, x)`

### 3.7 $\int x^2 \tan^2(a + bx) dx$

3.7.1	Optimal result . . . . .	79
3.7.2	Mathematica [B] (verified) . . . . .	79
3.7.3	Rubi [A] (verified) . . . . .	80
3.7.4	Maple [A] (verified) . . . . .	82
3.7.5	Fricas [B] (verification not implemented) . . . . .	82
3.7.6	Sympy [F] . . . . .	83
3.7.7	Maxima [B] (verification not implemented) . . . . .	83
3.7.8	Giac [F] . . . . .	84
3.7.9	Mupad [F(-1)] . . . . .	84

#### 3.7.1 Optimal result

Integrand size = 12, antiderivative size = 73

$$\int x^2 \tan^2(a + bx) dx = -\frac{ix^2}{b} - \frac{x^3}{3} + \frac{2x \log(1 + e^{2i(a+bx)})}{b^2} - \frac{i \text{PolyLog}(2, -e^{2i(a+bx)})}{b^3} + \frac{x^2 \tan(a + bx)}{b}$$

output `-I*x^2/b-1/3*x^3+2*x*ln(1+exp(2*I*(b*x+a)))/b^2-I*polylog(2,-exp(2*I*(b*x+a)))/b^3+x^2*tan(b*x+a)/b`

#### 3.7.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 160 vs. 2(73) = 146.

Time = 5.05 (sec) , antiderivative size = 160, normalized size of antiderivative = 2.19

$$\int x^2 \tan^2(a + bx) dx = -\frac{x^3}{3} + \frac{x^2 \sec(a) \sec(a + bx) \sin(bx)}{b} + \frac{ibx(\pi + 2 \arctan(\cot(a))) + \pi \log(1 + e^{-2ibx}) + 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))})}{b}$$

input `Integrate[x^2*Tan[a + b*x]^2,x]`



output 
$$-1/3*x^3 + (x^2*\text{Sec}[a]*\text{Sec}[a + b*x]*\text{Sin}[b*x])/b + (I*b*x*(\text{Pi} + 2*\text{ArcTan}[\text{Cot}[a]]) + \text{Pi}*\text{Log}[1 + E^{((-2*I)*b*x)}] + 2*(b*x - \text{ArcTan}[\text{Cot}[a]])*\text{Log}[1 - E^{(2*I)*(b*x - \text{ArcTan}[\text{Cot}[a]])}] - \text{Pi}*\text{Log}[\text{Cos}[b*x]] + 2*\text{ArcTan}[\text{Cot}[a]]*\text{Log}[\text{Sin}[b*x - \text{ArcTan}[\text{Cot}[a]]]] - I*\text{PolyLog}[2, E^{(2*I)*(b*x - \text{ArcTan}[\text{Cot}[a]])}]) + (b^2*x^2*\text{Sqrt}[\text{Csc}[a]^2*\text{Tan}[a])/E^{(I*\text{ArcTan}[\text{Cot}[a]])})/b^3$$

### 3.7.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.19, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 4203, 15, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \tan^2(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \tan(a + bx)^2 dx \\
 & \quad \downarrow \text{4203} \\
 & -\frac{2 \int x \tan(a + bx) dx}{b} - \int x^2 dx + \frac{x^2 \tan(a + bx)}{b} \\
 & \quad \downarrow \text{15} \\
 & -\frac{2 \int x \tan(a + bx) dx}{b} + \frac{x^2 \tan(a + bx)}{b} - \frac{x^3}{3} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2 \int x \tan(a + bx) dx}{b} + \frac{x^2 \tan(a + bx)}{b} - \frac{x^3}{3} \\
 & \quad \downarrow \text{4202} \\
 & -\frac{2\left(\frac{ix^2}{2} - 2i \int \frac{e^{2i(a+bx)}x}{1+e^{2i(a+bx)}} dx\right)}{b} + \frac{x^2 \tan(a + bx)}{b} - \frac{x^3}{3} \\
 & \quad \downarrow \text{2620} \\
 & -\frac{2\left(\frac{ix^2}{2} - 2i\left(\frac{i \int \log(1+e^{2i(a+bx)}) dx}{2b} - \frac{ix \log(1+e^{2i(a+bx)})}{2b}\right)\right)}{b} + \frac{x^2 \tan(a + bx)}{b} - \frac{x^3}{3} \\
 & \quad \downarrow \text{2715}
 \end{aligned}$$

$$-\frac{2\left(\frac{ix^2}{2} - 2i\left(\frac{\int e^{-2i(a+bx)} \log(1+e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b}\right)\right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3}$$

↓ 2838

$$-\frac{2\left(\frac{ix^2}{2} - 2i\left(-\frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b}\right)\right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3}$$

input `Int[x^2*Tan[a + b*x]^2,x]`

output `-1/3*x^3 - (2*((I/2)*x^2 - (2*I)*((-1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))]))/b - PolyLog[2, -E^((2*I)*(a + b*x))]/(4*b^2))/b + (x^2*Tan[a + b*x])/b`

### 3.7.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4202 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 4203 `Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Simp[b*d*(m/(f*(n - 1))) Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Simp[b^2 Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]`

### 3.7.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.48

method	result	size
risch	$-\frac{x^3}{3} + \frac{2ix^2}{b(e^{2i(bx+a)}+1)} - \frac{2ix^2}{b} - \frac{4iax}{b^2} - \frac{2ia^2}{b^3} + \frac{2x \ln(e^{2i(bx+a)}+1)}{b^2} - \frac{i \operatorname{Li}_2(-e^{2i(bx+a)})}{b^3} + \frac{4a \ln(e^{i(bx+a)})}{b^3}$	108

input `int(x^2*tan(b*x+a)^2,x,method=_RETURNVERBOSE)`

output  $-1/3*x^3+2*I*x^2/b/(\exp(2*I*(b*x+a))+1)-2*I/b*x^2-4*I/b^2*a*x-2*I/b^3*a^2+2*x*\ln(\exp(2*I*(b*x+a))+1)/b^2-I*\operatorname{polylog}(2,-\exp(2*I*(b*x+a)))/b^3+4/b^3*a*\ln(\exp(I*(b*x+a)))$

### 3.7.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 144 vs.  $2(62) = 124$ .

Time = 0.25 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.97

$$\int x^2 \tan^2(a + bx) dx = \frac{2b^3x^3 - 6b^2x^2 \tan(bx + a) - 6bx \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 6bx \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 3i \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 3i \operatorname{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right)}{6b^3}$$

input `integrate(x^2*tan(b*x+a)^2,x, algorithm="fricas")`

output 
$$-1/6*(2*b^3*x^3 - 6*b^2*x^2*\tan(b*x + a) - 6*b*x*\log(-2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 6*b*x*\log(-2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 3*I*dilog(2*(I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1) + 3*I*dilog(2*(-I*\tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1) + 1))/b^3$$

### 3.7.6 Sympy [F]

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan^2(a + bx) dx$$

input `integrate(x**2*tan(b*x+a)**2,x)`

output `Integral(x**2*tan(a + b*x)**2, x)`

### 3.7.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 257 vs.  $2(62) = 124$ .

Time = 0.63 (sec) , antiderivative size = 257, normalized size of antiderivative = 3.52

$$\int x^2 \tan^2(a + bx) dx = \frac{i b^3 x^3 + 6 (bx \cos(2bx + 2a) + i bx \sin(2bx + 2a) + bx) \arctan(\sin(2bx + 2a), \cos(2bx + 2a) + 1) + (I*b^3*x^3 + 6*(b*x*cos(2*b*x + 2*a) + I*b*x*sin(2*b*x + 2*a) + b*x)*arctan(2*(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + (I*b^3*x^3 - 6*b^2*x^2)*cos(2*b*x + 2*a) - 3*(cos(2*b*x + 2*a) + I*sin(2*b*x + 2*a) + 1)*dilog(-e^(2*I*b*x + 2*I*a)) - 3*(I*b*x*cos(2*b*x + 2*a) - b*x*sin(2*b*x + 2*a) + I*b*x)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - (b^3*x^3 + 6*I*b^2*x^2)*sin(2*b*x + 2*a))/(-3*I*b^3*cos(2*b*x + 2*a) + 3*b^3*sin(2*b*x + 2*a) - 3*I*b^3)}$$

input `integrate(x^2*tan(b*x+a)^2,x, algorithm="maxima")`

output 
$$(I*b^3*x^3 + 6*(b*x*cos(2*b*x + 2*a) + I*b*x*sin(2*b*x + 2*a) + b*x)*arctan(2*(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + (I*b^3*x^3 - 6*b^2*x^2)*cos(2*b*x + 2*a) - 3*(cos(2*b*x + 2*a) + I*sin(2*b*x + 2*a) + 1)*dilog(-e^(2*I*b*x + 2*I*a)) - 3*(I*b*x*cos(2*b*x + 2*a) - b*x*sin(2*b*x + 2*a) + I*b*x)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - (b^3*x^3 + 6*I*b^2*x^2)*sin(2*b*x + 2*a))/(-3*I*b^3*cos(2*b*x + 2*a) + 3*b^3*sin(2*b*x + 2*a) - 3*I*b^3)$$

### 3.7.8 Giac [F]

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan(bx + a)^2 dx$$

input `integrate(x^2*tan(b*x+a)^2,x, algorithm="giac")`

output `integrate(x^2*tan(b*x + a)^2, x)`

### 3.7.9 Mupad [F(-1)]

Timed out.

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan(a + bx)^2 dx$$

input `int(x^2*tan(a + b*x)^2,x)`

output `int(x^2*tan(a + b*x)^2, x)`

### 3.8 $\int x \tan^2(a + bx) dx$

3.8.1	Optimal result . . . . .	85
3.8.2	Mathematica [A] (verified) . . . . .	85
3.8.3	Rubi [A] (verified) . . . . .	86
3.8.4	Maple [A] (verified) . . . . .	87
3.8.5	Fricas [A] (verification not implemented) . . . . .	88
3.8.6	Sympy [A] (verification not implemented) . . . . .	88
3.8.7	Maxima [B] (verification not implemented) . . . . .	88
3.8.8	Giac [B] (verification not implemented) . . . . .	89
3.8.9	Mupad [B] (verification not implemented) . . . . .	89

#### 3.8.1 Optimal result

Integrand size = 10, antiderivative size = 30

$$\int x \tan^2(a + bx) dx = -\frac{x^2}{2} + \frac{\log(\cos(a + bx))}{b^2} + \frac{x \tan(a + bx)}{b}$$

output `-1/2*x^2+ln(cos(b*x+a))/b^2+x*tan(b*x+a)/b`

#### 3.8.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.43

$$\int x \tan^2(a + bx) dx = -\frac{x^2}{2} + \frac{\log(\cos(a + bx))}{b^2} + \frac{x \sec(a) \sec(a + bx) \sin(bx)}{b} + \frac{x \tan(a)}{b}$$

input `Integrate[x*Tan[a + b*x]^2,x]`

output `-1/2*x^2 + Log[Cos[a + b*x]]/b^2 + (x*Sec[a]*Sec[a + b*x]*Sin[b*x])/b + (x*Tan[a])/b`

### 3.8.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4203, 15, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tan^2(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \tan(a + bx)^2 dx \\
 & \quad \downarrow \text{4203} \\
 & -\frac{\int \tan(a + bx) dx}{b} - \int x dx + \frac{x \tan(a + bx)}{b} \\
 & \quad \downarrow \text{15} \\
 & -\frac{\int \tan(a + bx) dx}{b} + \frac{x \tan(a + bx)}{b} - \frac{x^2}{2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int \tan(a + bx) dx}{b} + \frac{x \tan(a + bx)}{b} - \frac{x^2}{2} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\log(\cos(a + bx))}{b^2} + \frac{x \tan(a + bx)}{b} - \frac{x^2}{2}
 \end{aligned}$$

input `Int[x*Tan[a + b*x]^2,x]`

output `-1/2*x^2 + Log[Cos[a + b*x]]/b^2 + (x*Tan[a + b*x])/b`

## 3.8.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4203 `Int[((c_.) + (d_.)*(x_)^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Simp[b*d*(m/(f*(n - 1))) Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Simp[b^2 Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]`

## 3.8.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

method	result	size
norman	$\frac{x \tan(bx+a)}{b} - \frac{x^2}{2} - \frac{\ln(1+\tan^2(bx+a))}{2b^2}$	34
parallelrisch	$-\frac{x^2 b^2 - 2x \tan(bx+a)b + \ln(1+\tan^2(bx+a))}{2b^2}$	35
default	$-\frac{x^2}{2} + \frac{(bx+a)\tan(bx+a) + \ln(\cos(bx+a)) - a \tan(bx+a)}{b^2}$	40
risch	$-\frac{x^2}{2} - \frac{2ix}{b} - \frac{2ia}{b^2} + \frac{2ix}{b(e^{2i(bx+a)}+1)} + \frac{\ln(e^{2i(bx+a)}+1)}{b^2}$	57

input `int(x*tan(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `x*tan(b*x+a)/b-1/2*x^2-1/2/b^2*ln(1+tan(b*x+a)^2)`



### 3.8.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.27

$$\int x \tan^2(a + bx) dx = -\frac{b^2 x^2 - 2bx \tan(bx + a) - \log\left(\frac{1}{\tan(bx+a)^2 + 1}\right)}{2b^2}$$

input `integrate(x*tan(b*x+a)^2,x, algorithm="fricas")`

output `-1/2*(b^2*x^2 - 2*b*x*tan(b*x + a) - log(1/(tan(b*x + a)^2 + 1)))/b^2`

### 3.8.6 Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.37

$$\int x \tan^2(a + bx) dx = \begin{cases} -\frac{x^2}{2} + \frac{x \tan(a+bx)}{b} - \frac{\log(\tan^2(a+bx)+1)}{2b^2} & \text{for } b \neq 0 \\ \frac{x^2 \tan^2(a)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*tan(b*x+a)**2,x)`

output `Piecewise((-x**2/2 + x*tan(a + b*x)/b - log(tan(a + b*x)**2 + 1)/(2*b**2), Ne(b, 0)), (x**2*tan(a)**2/2, True))`

### 3.8.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 214 vs. 2(28) = 56.

Time = 0.52 (sec) , antiderivative size = 214, normalized size of antiderivative = 7.13

$$\int x \tan^2(a + bx) dx = \frac{2(bx + a - \tan(bx + a))a - \frac{(bx+a)^2 \cos(2bx+2a)^2 + (bx+a)^2 \sin(2bx+2a)^2 + 2(bx+a)^2 \cos(2bx+2a) + (bx+a)^2 - (\cos(2bx+2a))^2 + \sin(2bx+2a)^2}{\cos(2bx+2a)^2 + \sin(2bx+2a)^2}}{2b^2}$$

input `integrate(x*tan(b*x+a)^2,x, algorithm="maxima")`

output  $\frac{1}{2}*(2*(b*x + a - \tan(b*x + a))*a - ((b*x + a)^2*\cos(2*b*x + 2*a)^2 + (b*x + a)^2*\sin(2*b*x + 2*a)^2 + 2*(b*x + a)^2*\cos(2*b*x + 2*a) + (b*x + a)^2 - (\cos(2*b*x + 2*a)^2 + \sin(2*b*x + 2*a)^2 + 2*\cos(2*b*x + 2*a) + 1)*\log(\cos(2*b*x + 2*a)^2 + \sin(2*b*x + 2*a)^2 + 2*\cos(2*b*x + 2*a) + 1) - 4*(b*x + a)*\sin(2*b*x + 2*a))/(\cos(2*b*x + 2*a)^2 + \sin(2*b*x + 2*a)^2 + 2*\cos(2*b*x + 2*a) + 1))/b^2$

### 3.8.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs.  $2(28) = 56$ .

Time = 0.60 (sec) , antiderivative size = 162, normalized size of antiderivative = 5.40

$$\int x \tan^2(a + bx) dx =$$

$$\frac{b^2 x^2 \tan(bx) \tan(a) - b^2 x^2 + 2bx \tan(bx) + 2bx \tan(a) - \log\left(\frac{4(\tan(bx)^2 \tan(a)^2 - 2 \tan(bx) \tan(a) + 1)}{\tan(bx)^2 \tan(a)^2 + \tan(bx)^2 + \tan(a)^2 + 1}\right) \tan(a)}{2(b^2 \tan(bx) \tan(a) - b^2)}$$

input `integrate(x*tan(b*x+a)^2,x, algorithm="giac")`

output  $\frac{-1/2*(b^2*x^2*\tan(b*x)*\tan(a) - b^2*x^2 + 2*b*x*\tan(b*x) + 2*b*x*\tan(a) - \log(4*(\tan(b*x)^2*\tan(a)^2 - 2*\tan(b*x)*\tan(a) + 1)/(\tan(b*x)^2*\tan(a)^2 + \tan(b*x)^2 + \tan(a)^2 + 1))*\tan(b*x)*\tan(a) + \log(4*(\tan(b*x)^2*\tan(a)^2 - 2*\tan(b*x)*\tan(a) + 1)/(\tan(b*x)^2*\tan(a)^2 + \tan(b*x)^2 + \tan(a)^2 + 1))}{(b^2*\tan(b*x)*\tan(a) - b^2)}$

### 3.8.9 Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int x \tan^2(a + bx) dx = -\frac{\ln(\tan(a+bx)^2+1)}{2} - \frac{bx \tan(a + bx)}{b^2} - \frac{x^2}{2}$$

input `int(x*tan(a + b*x)^2,x)`

output  $-(\log(\tan(a + b*x)^2 + 1)/2 - b*x*\tan(a + b*x))/b^2 - x^2/2$

### 3.9 $\int \frac{\tan^2(a+bx)}{x} dx$

3.9.1	Optimal result . . . . .	90
3.9.2	Mathematica [N/A] . . . . .	90
3.9.3	Rubi [N/A] . . . . .	91
3.9.4	Maple [N/A] (verified) . . . . .	92
3.9.5	Fricas [N/A] . . . . .	92
3.9.6	Sympy [N/A] . . . . .	92
3.9.7	Maxima [N/A] . . . . .	93
3.9.8	Giac [N/A] . . . . .	93
3.9.9	Mupad [N/A] . . . . .	93

#### 3.9.1 Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^2(a+bx)}{x} dx = \text{Int}\left(\frac{\tan^2(a+bx)}{x}, x\right)$$

output `Unintegrable(tan(b*x+a)^2/x,x)`

#### 3.9.2 Mathematica [N/A]

Not integrable

Time = 3.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a+bx)}{x} dx = \int \frac{\tan^2(a+bx)}{x} dx$$

input `Integrate[Tan[a + b*x]^2/x,x]`

output `Integrate[Tan[a + b*x]^2/x, x]`

### 3.9.3 Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(a + bx)}{x} dx$$

↓ 3042

$$\int \frac{\tan(a + bx)^2}{x} dx$$

↓ 4222

$$\int \frac{\tan^2(a + bx)}{x} dx$$

input `Int[Tan[a + b*x]^2/x,x]`

output `$Aborted`

#### 3.9.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4222 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], I^n*Unintegrable[(c + d*x)^m*Coth[(-I)*(e - Pi/2) - I*f*x]^n, x], I^n*Unintegrable[(c + d*x)^m*Tanh[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], (-1)^n*Unintegrable[(c + d*x)^m*Cot[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*Tan[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]`

**3.9.4 Maple [N/A] (verified)**

Not integrable

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(bx + a)}{x} dx$$

input `int(tan(b*x+a)^2/x,x)`output `int(tan(b*x+a)^2/x,x)`**3.9.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

input `integrate(tan(b*x+a)^2/x,x, algorithm="fricas")`output `integral(tan(b*x + a)^2/x, x)`**3.9.6 Sympy [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(a + bx)}{x} dx$$

input `integrate(tan(b*x+a)**2/x,x)`output `Integral(tan(a + b*x)**2/x, x)`

### 3.9.7 Maxima [N/A]

Not integrable

Time = 0.96 (sec) , antiderivative size = 243, normalized size of antiderivative = 20.25

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

input `integrate(tan(b*x+a)^2/x,x, algorithm="maxima")`

output `-(b*x*cos(2*b*x + 2*a)^2*log(x) + b*x*log(x)*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a)*log(x) + b*x*log(x) - 2*(b^2*x*cos(2*b*x + 2*a)^2 + b^2*x*sin(2*b*x + 2*a)^2 + 2*b^2*x*cos(2*b*x + 2*a) + b^2*x)*integrate(sin(2*b*x + 2*a)/(b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(2*b*x + 2*a)^2 + 2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2), x) - 2*sin(2*b*x + 2*a))/(b*x*cos(2*b*x + 2*a)^2 + b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + b*x)`

### 3.9.8 Giac [N/A]

Not integrable

Time = 0.63 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

input `integrate(tan(b*x+a)^2/x,x, algorithm="giac")`

output `integrate(tan(b*x + a)^2/x, x)`

### 3.9.9 Mupad [N/A]

Not integrable

Time = 3.57 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(a + bx)^2}{x} dx$$

input `int(tan(a + b*x)^2/x,x)`

output `int(tan(a + b*x)^2/x, x)`

### 3.10 $\int \frac{\tan^2(a+bx)}{x^2} dx$

3.10.1	Optimal result	95
3.10.2	Mathematica [N/A]	95
3.10.3	Rubi [N/A]	96
3.10.4	Maple [N/A] (verified)	97
3.10.5	Fricas [N/A]	97
3.10.6	Sympy [N/A]	97
3.10.7	Maxima [N/A]	98
3.10.8	Giac [N/A]	98
3.10.9	Mupad [N/A]	98

#### 3.10.1 Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \text{Int}\left(\frac{\tan^2(a + bx)}{x^2}, x\right)$$

output `Unintegrable(tan(b*x+a)^2/x^2,x)`

#### 3.10.2 Mathematica [N/A]

Not integrable

Time = 2.44 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(a + bx)}{x^2} dx$$

input `Integrate[Tan[a + b*x]^2/x^2,x]`

output `Integrate[Tan[a + b*x]^2/x^2, x]`



### 3.10.3 Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(a + bx)}{x^2} dx$$

↓ 3042

$$\int \frac{\tan(a + bx)^2}{x^2} dx$$

↓ 4222

$$\int \frac{\tan^2(a + bx)}{x^2} dx$$

input `Int[Tan[a + b*x]^2/x^2,x]`

output `$Aborted`

#### 3.10.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4222 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], I^n*Unintegrable[(c + d*x)^m*Coth[(-I)*(e - Pi/2) - I*f*x]^n, x], I^n*Unintegrable[(c + d*x)^m*Tanh[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], (-1)^n*Unintegrable[(c + d*x)^m*Cot[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*Tan[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]`

**3.10.4 Maple [N/A] (verified)**

Not integrable

Time = 0.31 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(bx + a)}{x^2} dx$$

input `int(tan(b*x+a)^2/x^2,x)`output `int(tan(b*x+a)^2/x^2,x)`**3.10.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

input `integrate(tan(b*x+a)^2/x^2,x, algorithm="fricas")`output `integral(tan(b*x + a)^2/x^2, x)`**3.10.6 Sympy [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(a + bx)}{x^2} dx$$

input `integrate(tan(b*x+a)**2/x**2,x)`output `Integral(tan(a + b*x)**2/x**2, x)`

**3.10.7 Maxima [N/A]**

Not integrable

Time = 0.45 (sec) , antiderivative size = 250, normalized size of antiderivative = 20.83

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

```
input integrate(tan(b*x+a)^2/x^2,x, algorithm="maxima")
```

```
output (b*x*cos(2*b*x + 2*a)^2 + b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a)
+ b*x + 4*(b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(2*b*x + 2*a)^2 + 2*b^2
*x^2*cos(2*b*x + 2*a) + b^2*x^2)*integrate(sin(2*b*x + 2*a)/(b^2*x^3*cos(2
*b*x + 2*a)^2 + b^2*x^3*sin(2*b*x + 2*a)^2 + 2*b^2*x^3*cos(2*b*x + 2*a) +
b^2*x^3), x) + 2*sin(2*b*x + 2*a))/(b*x^2*cos(2*b*x + 2*a)^2 + b*x^2*sin(2
*b*x + 2*a)^2 + 2*b*x^2*cos(2*b*x + 2*a) + b*x^2)
```

**3.10.8 Giac [N/A]**

Not integrable

Time = 0.67 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

```
input integrate(tan(b*x+a)^2/x^2,x, algorithm="giac")
```

```
output integrate(tan(b*x + a)^2/x^2, x)
```

**3.10.9 Mupad [N/A]**

Not integrable

Time = 2.93 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)^2}{x^2} dx$$

input `int(tan(a + b*x)^2/x^2,x)`

output `int(tan(a + b*x)^2/x^2, x)`

### 3.11 $\int x^3 \tan^3(a + bx) dx$

3.11.1	Optimal result . . . . .	100
3.11.2	Mathematica [A] (verified) . . . . .	101
3.11.3	Rubi [A] (verified) . . . . .	101
3.11.4	Maple [A] (verified) . . . . .	107
3.11.5	Fricas [B] (verification not implemented) . . . . .	107
3.11.6	Sympy [F] . . . . .	108
3.11.7	Maxima [B] (verification not implemented) . . . . .	108
3.11.8	Giac [F] . . . . .	109
3.11.9	Mupad [F(-1)] . . . . .	110

#### 3.11.1 Optimal result

Integrand size = 12, antiderivative size = 205

$$\begin{aligned} \int x^3 \tan^3(a + bx) dx = & \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} \\ & + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^4} \\ & - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} + \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} \\ & + \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} - \frac{3x^2 \tan(a + bx)}{2b^2} + \frac{x^3 \tan^2(a + bx)}{2b} \end{aligned}$$

output  $3/2*I*x^2/b^2+1/2*x^3/b-1/4*I*x^4-3*x*\ln(1+\exp(2*I*(b*x+a)))/b^3+x^3*\ln(1+\exp(2*I*(b*x+a)))/b+3/2*I*polylog(2,-\exp(2*I*(b*x+a)))/b^4-3/2*I*x^2*polylog(2,-\exp(2*I*(b*x+a)))/b^2+3/2*x*polylog(3,-\exp(2*I*(b*x+a)))/b^3+3/4*I*polylog(4,-\exp(2*I*(b*x+a)))/b^4-3/2*x^2*\tan(b*x+a)/b^2+1/2*x^3*\tan(b*x+a)^2/b$

### 3.11.2 Mathematica [A] (verified)

Time = 6.54 (sec) , antiderivative size = 370, normalized size of antiderivative = 1.80

$$\int x^3 \tan^3(a + bx) dx$$

$$= \frac{ie^{ia}(2b^4e^{-2ia}x^4 - 4ib^3(1 + e^{-2ia})x^3 \log(1 + e^{-2i(a+bx)}) + 6b^2(1 + e^{-2ia})x^2 \text{PolyLog}(2, -e^{-2i(a+bx)}) - 6ib}{8b^4}$$

$$+ \frac{x^3 \sec^2(a + bx)}{2b}$$

$$- \frac{3 \csc(a) \left( b^2 e^{-i \arctan(\cot(a))} x^2 - \frac{\cot(a)(ibx - \pi - 2 \arctan(\cot(a))) - \pi \log(1 + e^{-2ibx}) - 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))})}{\sqrt{2b^4 \sqrt{\csc^2(a) (\cos^2(a) + \sin^2(a))}}} \right)}{2b^2}$$

$$- \frac{3x^2 \sec(a) \sec(a + bx) \sin(bx)}{2b^2} - \frac{1}{4} x^4 \tan(a)$$

input `Integrate[x^3*Tan[a + b*x]^3,x]`

output `((I/8)*E^(I*a)*((2*b^4*x^4)/E^((2*I)*a) - (4*I)*b^3*(1 + E^((-2*I)*a))*x^3 *Log[1 + E^((-2*I)*(a + b*x))] + 6*b^2*(1 + E^((-2*I)*a))*x^2*PolyLog[2, - E^((-2*I)*(a + b*x))] - (6*I)*b*(1 + E^((-2*I)*a))*x*PolyLog[3, -E^((-2*I)*(a + b*x))] - 3*(1 + E^((-2*I)*a))*PolyLog[4, -E^((-2*I)*(a + b*x))]*Sec [a])/b^4 + (x^3*Sec[a + b*x]^2)/(2*b) - (3*Csc[a]*((b^2*x^2)/E^(I*ArcTan[C ot[a]]) - (Cot[a]*(I*b*x*(-Pi - 2*ArcTan[Cot[a]]) - Pi*Log[1 + E^((-2*I)*b *x]) - 2*(b*x - ArcTan[Cot[a]])*Log[1 - E^((2*I)*(b*x - ArcTan[Cot[a]])]) + Pi*Log[Cos[b*x]] - 2*ArcTan[Cot[a]]*Log[Sin[b*x - ArcTan[Cot[a]]]]) + I*P olyLog[2, E^((2*I)*(b*x - ArcTan[Cot[a]])])))/Sqrt[1 + Cot[a]^2])*Sec[a]/ (2*b^4*Sqrt[Csc[a]^2*(Cos[a]^2 + Sin[a]^2)]) - (3*x^2*Sec[a]*Sec[a + b*x]* Sin[b*x])/(2*b^2) - (x^4*Tan[a])/4`

### 3.11.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.20, number of steps used = 17, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.333$ , Rules used = {3042, 4203, 3042, 4202, 2620, 3011, 4203, 15, 3042, 4202, 2620, 2715, 2838, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.11.  $\int x^3 \tan^3(a + bx) dx$

$$\begin{aligned}
& \int x^3 \tan^3(a + bx) dx \\
& \quad \downarrow \text{3042} \\
& \int x^3 \tan(a + bx)^3 dx \\
& \quad \downarrow \text{4203} \\
& - \int x^3 \tan(a + bx) dx - \frac{3 \int x^2 \tan^2(a + bx) dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} \\
& \quad \downarrow \text{3042} \\
& - \int x^3 \tan(a + bx) dx - \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} \\
& \quad \downarrow \text{4202} \\
& 2i \int \frac{e^{2i(a+bx)} x^3}{1 + e^{2i(a+bx)}} dx - \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{2620} \\
& - \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + 2i \left( \frac{3i \int x^2 \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) + \\
& \quad \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{3011} \\
& 2i \left( \frac{3i \left( \frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{4203} \\
& 2i \left( \frac{3i \left( \frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \left( -\frac{2 \int x \tan(a+bx) dx}{b} - \int x^2 dx + \frac{x^2 \tan(a+bx)}{b} \right)}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{15}
\end{aligned}$$

$$\begin{aligned}
& 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad 3 \left( -\frac{2 \int x \tan(a+bx) dx}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{3042} \\
& 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad 3 \left( -\frac{2 \int x \tan(a+bx) dx}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{4202} \\
& 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad 3 \left( -\frac{2 \left( \frac{ix^2}{2} - 2i \int \frac{e^{2i(a+bx)} x}{1 + e^{2i(a+bx)}} dx \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{2620} \\
& 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad 3 \left( -\frac{2 \left( \frac{ix^2}{2} - 2i \left( \frac{i \int \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{2715} \\
& 3 \left( -\frac{2 \left( \frac{ix^2}{2} - 2i \left( \frac{\int e^{-2i(a+bx)} \log(1 + e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \\
& \quad - \frac{2b}{2b} \\
& 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) + \\
& \quad \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4}
\end{aligned}$$



$$\begin{aligned} & \downarrow \text{2838} \\ & 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\ & 3 \left( -\frac{2 \left( \frac{ix^2}{2} - 2i \left( -\frac{\operatorname{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) \\ & \frac{\hspace{10em}}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{7163} \\ & 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left( \frac{\int \operatorname{PolyLog}(3, -e^{2i(a+bx)}) dx}{2b} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\ & 3 \left( -\frac{2 \left( \frac{ix^2}{2} - 2i \left( -\frac{\operatorname{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) \\ & \frac{\hspace{10em}}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{2720} \\ & 2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left( \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(3, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\ & 3 \left( -\frac{2 \left( \frac{ix^2}{2} - 2i \left( -\frac{\operatorname{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) \\ & \frac{\hspace{10em}}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \end{aligned}$$

$$\downarrow \text{7143}$$

$$2i \left( \frac{3i \left( \frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left( \frac{\operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) -$$

$$3 \left( - \frac{2 \left( \frac{ix^2}{2} - 2i \left( - \frac{\operatorname{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4}$$

input `Int[x^3*Tan[a + b*x]^3,x]`

output `(-1/4*I)*x^4 + (2*I)*((( -1/2*I)*x^3*Log[1 + E^((2*I)*(a + b*x))])/b + ((3*I)/2)*(((I/2)*x^2*PolyLog[2, -E^((2*I)*(a + b*x))])/b - (I*((( -1/2*I)*x*PolyLog[3, -E^((2*I)*(a + b*x))])/b + PolyLog[4, -E^((2*I)*(a + b*x))]/(4*b^2))))/b)/b + (x^3*Tan[a + b*x]^2)/(2*b) - (3*(-1/3*x^3 - (2*((I/2)*x^2 - (2*I)*((( -1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))])/b - PolyLog[2, -E^((2*I)*(a + b*x))]/(4*b^2)))))/b + (x^2*Tan[a + b*x])/b)/(2*b)`

### 3.11.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n/(b*c*n*Log[F])], x) + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4202 `Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x)))], x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 4203 `Int[((c_) + (d_)*(x_)^(m_))*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Simp[b*d*(m/(f*(n - 1))) Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Simp[b^2 Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### 3.11.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.22

method	result
risch	$\frac{3i \operatorname{Li}_2(-e^{2i(bx+a)})}{2b^4} + \frac{x^2(2bx e^{2i(bx+a)} - 3i e^{2i(bx+a)} - 3i)}{b^2(e^{2i(bx+a)} + 1)^2} + \frac{3ix^2}{b^2} - \frac{ix^4}{4} + \frac{3i \operatorname{Li}_4(-e^{2i(bx+a)})}{4b^4} - \frac{3ia^4}{2b^4} - \frac{3ix^2 \operatorname{Li}_2(-e^{2i(bx+a)})}{2b^2}$

input `int(x^3*tan(b*x+a)^3,x,method=_RETURNVERBOSE)`

output  $\frac{3}{2}I \operatorname{polylog}(2, -\exp(2I(bx+a)))/b^4 + x^2(2b^2 \exp(2I(bx+a)) - 3I \exp(2I(bx+a)) - 3I)/b^2 + (\exp(2I(bx+a)) + 1)^2 + 3I/b^2 x^2 - 1/4 I x^4 + 3/4 I \operatorname{polylog}(4, -\exp(2I(bx+a)))/b^4 - 3/2 I/b^4 a^4 - 3/2 I x^2 \operatorname{polylog}(2, -\exp(2I(bx+a)))/b^2 + 6I/b^3 a x - 2I/b^3 a^3 x - 6/b^4 a \ln(\exp(I(bx+a))) + 2/b^4 a^3 \ln(\exp(I(bx+a))) + 3I/b^4 a^2 + x^3 \ln(\exp(2I(bx+a)) + 1)/b + 3/2 x \operatorname{polylog}(3, -\exp(2I(bx+a)))/b^3 - 3x \ln(\exp(2I(bx+a)) + 1)/b^3$

### 3.11.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 344 vs.  $2(163) = 326$ .

Time = 0.27 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.68

$$\int x^3 \tan^3(a + bx) dx = \frac{4b^3 x^3 \tan(bx + a)^2 + 4b^3 x^3 - 12b^2 x^2 \tan(bx + a) + 6bx \operatorname{polylog}\left(3, \frac{\tan(bx+a)^2 + 2i \tan(bx+a) - 1}{\tan(bx+a)^2 + 1}\right) + 6bx \operatorname{polylog}}$$

input `integrate(x^3*tan(b*x+a)^3,x, algorithm="fricas")`

output `1/8*(4*b^3*x^3*tan(b*x + a)^2 + 4*b^3*x^3 - 12*b^2*x^2*tan(b*x + a) + 6*b*x*polylog(3, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 6*b*x*polylog(3, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 6*(-I*b^2*x^2 + I)*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 6*(I*b^2*x^2 - I)*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) + 4*(b^3*x^3 - 3*b*x)*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 4*(b^3*x^3 - 3*b*x)*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 3*I*polylog(4, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 3*I*polylog(4, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)))/b^4`

### 3.11.6 Sympy [F]

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan^3(a + bx) dx$$

input `integrate(x**3*tan(b*x+a)**3,x)`

output `Integral(x**3*tan(a + b*x)**3, x)`

### 3.11.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1205 vs.  $2(163) = 326$ .

Time = 0.47 (sec) , antiderivative size = 1205, normalized size of antiderivative = 5.88

$$\int x^3 \tan^3(a + bx) dx = \text{Too large to display}$$

input `integrate(x^3*tan(b*x+a)^3,x, algorithm="maxima")`

output

```

1/2*(a^3*(1/(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2 - 1)) - 2*(3*(b*x +
a)^4 - 12*(b*x + a)^3*a + 18*(b*x + a)^2*a^2 + 36*a^2 - 4*(4*(b*x + a)^3 -
9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + (4*(b*x + a)^3 - 9*(b*x + a)^2*
a + 9*(a^2 - 1)*(b*x + a) + 9*a)*cos(4*b*x + 4*a) + 2*(4*(b*x + a)^3 - 9*(
b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + 9*a)*cos(2*b*x + 2*a) - (-4*I*(b*x
+ a)^3 + 9*I*(b*x + a)^2*a + 9*(-I*a^2 + I)*(b*x + a) - 9*I*a)*sin(4*b*x +
4*a) - 2*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a + 9*(-I*a^2 + I)*(b*x + a)
- 9*I*a)*sin(2*b*x + 2*a) + 9*a)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*
a) + 1) + 3*((b*x + a)^4 - 4*(b*x + a)^3*a + 6*(a^2 - 2)*(b*x + a)^2 + 24*
(b*x + a)*a)*cos(4*b*x + 4*a) + 6*((b*x + a)^4 - 4*(b*x + a)^3*(a - I) + 6
*(a^2 - 2*I*a - 1)*(b*x + a)^2 + 12*(I*a^2 + a)*(b*x + a) + 6*a^2)*cos(2*b
*x + 2*a) + 6*(4*(b*x + a)^2 - 6*(b*x + a)*a + 3*a^2 + (4*(b*x + a)^2 - 6*
(b*x + a)*a + 3*a^2 - 3)*cos(4*b*x + 4*a) + 2*(4*(b*x + a)^2 - 6*(b*x + a)
*a + 3*a^2 - 3)*cos(2*b*x + 2*a) + (4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*
I*a^2 - 3*I)*sin(4*b*x + 4*a) + 2*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I
*a^2 - 3*I)*sin(2*b*x + 2*a) - 3)*dilog(-e^(2*I*b*x + 2*I*a)) + 2*(4*I*(b*
x + a)^3 - 9*I*(b*x + a)^2*a + 9*(I*a^2 - I)*(b*x + a) + (4*I*(b*x + a)^3
- 9*I*(b*x + a)^2*a + 9*(I*a^2 - I)*(b*x + a) + 9*I*a)*cos(4*b*x + 4*a) +
2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*(I*a^2 - I)*(b*x + a) + 9*I*a)*
cos(2*b*x + 2*a) - (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x ...

```

### 3.11.8 Giac [F]

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan(bx + a)^3 dx$$

input `integrate(x^3*tan(b*x+a)^3,x, algorithm="giac")`

output `integrate(x^3*tan(b*x + a)^3, x)`

**3.11.9 Mupad [F(-1)]**

Timed out.

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan(a + bx)^3 dx$$

input `int(x^3*tan(a + b*x)^3,x)`output `int(x^3*tan(a + b*x)^3, x)`

### 3.12 $\int x^2 \tan^3(a + bx) dx$

3.12.1	Optimal result . . . . .	111
3.12.2	Mathematica [A] (verified) . . . . .	111
3.12.3	Rubi [A] (verified) . . . . .	112
3.12.4	Maple [A] (verified) . . . . .	115
3.12.5	Fricas [B] (verification not implemented) . . . . .	116
3.12.6	Sympy [F] . . . . .	116
3.12.7	Maxima [B] (verification not implemented) . . . . .	117
3.12.8	Giac [F] . . . . .	117
3.12.9	Mupad [F(-1)] . . . . .	118

#### 3.12.1 Optimal result

Integrand size = 12, antiderivative size = 128

$$\int x^2 \tan^3(a + bx) dx = \frac{x^2}{2b} - \frac{ix^3}{3} + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} - \frac{\log(\cos(a + bx))}{b^3} - \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} + \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b}$$

output `1/2*x^2/b-1/3*I*x^3+x^2*ln(1+exp(2*I*(b*x+a)))/b-ln(cos(b*x+a))/b^3-I*x*polylog(2,-exp(2*I*(b*x+a)))/b^2+1/2*polylog(3,-exp(2*I*(b*x+a)))/b^3-x*tan(b*x+a)/b^2+1/2*x^2*tan(b*x+a)^2/b`

#### 3.12.2 Mathematica [A] (verified)

Time = 2.28 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.34

$$\int x^2 \tan^3(a + bx) dx = \frac{e^{-ia} (2b^2 x^2 (2ibx + 3(1 + e^{2ia}) \log(1 + e^{-2i(a+bx)})) + 6ib(1 + e^{2ia}) x \operatorname{PolyLog}(2, -e^{-2i(a+bx)}) + 3(1 + e^{2ia}))}{b^3}$$

input `Integrate[x^2*Tan[a + b*x]^3,x]`



```
output (((2*b^2*x^2*((2*I)*b*x + 3*(1 + E^((2*I)*a)))*Log[1 + E^((-2*I)*(a + b*x))
]) + (6*I)*b*(1 + E^((2*I)*a))*x*PolyLog[2, -E^((-2*I)*(a + b*x))] + 3*(1
+ E^((2*I)*a))*PolyLog[3, -E^((-2*I)*(a + b*x))])*Sec[a])/E^(I*a) + 6*b^2*
x^2*Sec[a + b*x]^2 - 12*b*x*Sec[a]*Sec[a + b*x]*Sin[b*x] - 4*b^3*x^3*Tan[a
] - 12*(Log[Cos[a + b*x]] + b*x*Tan[a]))/(12*b^3)
```

### 3.12.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.16, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 4203, 3042, 4202, 2620, 3011, 2720, 4203, 15, 3042, 3956, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \tan^3(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \tan(a + bx)^3 dx \\
 & \quad \downarrow \text{4203} \\
 & - \int x^2 \tan(a + bx) dx - \frac{\int x \tan^2(a + bx) dx}{b} + \frac{x^2 \tan^2(a + bx)}{2b} \\
 & \quad \downarrow \text{3042} \\
 & - \int x^2 \tan(a + bx) dx - \frac{\int x \tan(a + bx)^2 dx}{b} + \frac{x^2 \tan^2(a + bx)}{2b} \\
 & \quad \downarrow \text{4202} \\
 & 2i \int \frac{e^{2i(a+bx)} x^2}{1 + e^{2i(a+bx)}} dx - \frac{\int x \tan(a + bx)^2 dx}{b} + \frac{x^2 \tan^2(a + bx)}{2b} - \frac{ix^3}{3} \\
 & \quad \downarrow \text{2620} \\
 & 2i \left( \frac{i \int x \log(1 + e^{2i(a+bx)}) dx}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \frac{\int x \tan(a + bx)^2 dx}{b} + \\
 & \quad \frac{x^2 \tan^2(a + bx)}{2b} - \frac{ix^3}{3} \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$\begin{aligned}
& 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{\int x \tan(a+bx)^2 dx}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{2720} \\
& 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{\int x \tan(a+bx)^2 dx}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{4203} \\
& 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{-\frac{\int \tan(a+bx) dx}{b} - \int x dx + \frac{x \tan(a+bx)}{b}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{15} \\
& 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{-\frac{\int \tan(a+bx) dx}{b} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{3042} \\
& 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{-\frac{\int \tan(a+bx) dx}{b} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{3956} \\
& 2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{\frac{\log(\cos(a+bx))}{b^2} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{7143}
\end{aligned}$$

$$2i \left( \frac{i \left( \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \frac{\frac{\log(\cos(a+bx))}{b^2} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3}$$

input `Int[x^2*Tan[a + b*x]^3,x]`

output `(-1/3*I)*x^3 + (2*I)*((( -1/2*I)*x^2*Log[1 + E^((2*I)*(a + b*x))])/b + (I*((I/2)*x*PolyLog[2, -E^((2*I)*(a + b*x))])/b - PolyLog[3, -E^((2*I)*(a + b*x))]/(4*b^2)))/b + (x^2*Tan[a + b*x]^2)/(2*b) - (-1/2*x^2 + Log[Cos[a + b*x]]/b^2 + (x*Tan[a + b*x])/b)/b`

### 3.12.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2620 `Int[(((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.))*((c_.) + (d_.)*(x_)^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_)^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4202 `Int[((c_.) + (d_.)*(x_.))^(m_.)*tan[(e_.) + (f_.)*(x_.)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 4203 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Simp[b*d*(m/(f*(n - 1))) Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Simp[b^2 Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.12.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.41

method	result
risch	$-\frac{ix^3}{3} + \frac{2x(bx e^{2i(bx+a)} - i e^{2i(bx+a)} - i)}{b^2(e^{2i(bx+a)} + 1)^2} - \frac{2a^2 \ln(e^{i(bx+a)})}{b^3} + \frac{4ia^3}{3b^3} + \frac{2ia^2 x}{b^2} + \frac{x^2 \ln(e^{2i(bx+a)} + 1)}{b} - \frac{ix \operatorname{Li}_2(-e^{2i(bx+a)})}{b^2}$

input `int(x^2*tan(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `-1/3*I*x^3+2*x*(b*x*exp(2*I*(b*x+a))-I*exp(2*I*(b*x+a))-I)/b^2/(exp(2*I*(b*x+a))+1)^2-2/b^3*a^2*ln(exp(I*(b*x+a)))+4/3*I/b^3*a^3+2*I/b^2*a^2*x+x^2*ln(exp(2*I*(b*x+a))+1)/b-I*x*polylog(2,-exp(2*I*(b*x+a)))/b^2+1/2*polylog(3,-exp(2*I*(b*x+a)))/b^3-1/b^3*ln(exp(2*I*(b*x+a))+1)+2/b^3*ln(exp(I*(b*x+a)))`

---

3.12.  $\int x^2 \tan^3(a + bx) dx$

### 3.12.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 240 vs.  $2(109) = 218$ .

Time = 0.26 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.88

$$\int x^2 \tan^3(a + bx) dx$$

$$= \frac{2b^2x^2 \tan(bx + a)^2 + 2b^2x^2 + 2i bx \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - 2i bx \operatorname{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - 4bx \tan(bx + a)}{b^3}$$

input `integrate(x^2*tan(b*x+a)^3,x, algorithm="fracas")`

output `1/4*(2*b^2*x^2*tan(b*x + a)^2 + 2*b^2*x^2 + 2*I*b*x*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 2*I*b*x*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 4*b*x*tan(b*x + a) + 2*(b^2*x^2 - 1)*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*(b^2*x^2 - 1)*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1))) + polylog(3, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + polylog(3, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)))/b^3`

### 3.12.6 Sympy [F]

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan^3(a + bx) dx$$

input `integrate(x**2*tan(b*x+a)**3,x)`

output `Integral(x**2*tan(a + b*x)**3, x)`

### 3.12.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 736 vs.  $2(109) = 218$ .

Time = 0.66 (sec) , antiderivative size = 736, normalized size of antiderivative = 5.75

$$\int x^2 \tan^3(a + bx) dx = \text{Too large to display}$$

input `integrate(x^2*tan(b*x+a)^3,x, algorithm="maxima")`

output

```
-1/2*(a^2*(1/(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2 - 1)) + 2*(2*(b*x +
a)^3 - 6*(b*x + a)^2*a - 6*((b*x + a)^2 - 2*(b*x + a)*a + ((b*x + a)^2 -
2*(b*x + a)*a - 1)*cos(4*b*x + 4*a) + 2*((b*x + a)^2 - 2*(b*x + a)*a - 1)*
cos(2*b*x + 2*a) - (-I*(b*x + a)^2 + 2*I*(b*x + a)*a + I)*sin(4*b*x + 4*a)
- 2*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a + I)*sin(2*b*x + 2*a) - 1)*arctan2(
sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 2*((b*x + a)^3 - 3*(b*x + a)^2*a
- 6*b*x - 6*a)*cos(4*b*x + 4*a) + 4*((b*x + a)^3 - 3*(b*x + a)^2*(a - I)
+ 3*(b*x + a)*(-2*I*a - 1) - 3*a)*cos(2*b*x + 2*a) + 6*(b*x*cos(4*b*x + 4*
a) + 2*b*x*cos(2*b*x + 2*a) + I*b*x*sin(4*b*x + 4*a) + 2*I*b*x*sin(2*b*x +
2*a) + b*x)*dilog(-e^(2*I*b*x + 2*I*a)) + 3*(I*(b*x + a)^2 - 2*I*(b*x + a
)*a + (I*(b*x + a)^2 - 2*I*(b*x + a)*a - I)*cos(4*b*x + 4*a) + 2*(I*(b*x +
a)^2 - 2*I*(b*x + a)*a - I)*cos(2*b*x + 2*a) - ((b*x + a)^2 - 2*(b*x + a)
*a - 1)*sin(4*b*x + 4*a) - 2*((b*x + a)^2 - 2*(b*x + a)*a - 1)*sin(2*b*x +
2*a) - I)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a)
) + 1) + 3*(I*cos(4*b*x + 4*a) + 2*I*cos(2*b*x + 2*a) - sin(4*b*x + 4*a) -
2*sin(2*b*x + 2*a) + I)*polylog(3, -e^(2*I*b*x + 2*I*a)) + 2*(I*(b*x + a)
^3 - 3*I*(b*x + a)^2*a - 6*I*b*x - 6*I*a)*sin(4*b*x + 4*a) + 4*(I*(b*x + a)
)^3 + 3*(b*x + a)^2*(-I*a - 1) + 3*(b*x + a)*(2*a - I) - 3*I*a)*sin(2*b*x
+ 2*a) - 12*a)/(-6*I*cos(4*b*x + 4*a) - 12*I*cos(2*b*x + 2*a) + 6*sin(4*b*
x + 4*a) + 12*sin(2*b*x + 2*a) - 6*I))/b^3
```

### 3.12.8 Giac [F]

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan(bx + a)^3 dx$$

input `integrate(x^2*tan(b*x+a)^3,x, algorithm="giac")`

output `integrate(x^2*tan(b*x + a)^3, x)`

**3.12.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan(a + bx)^3 dx$$

input `int(x^2*tan(a + b*x)^3,x)`output `int(x^2*tan(a + b*x)^3, x)`

### 3.13 $\int x \tan^3(a + bx) dx$

3.13.1	Optimal result . . . . .	119
3.13.2	Mathematica [A] (verified) . . . . .	119
3.13.3	Rubi [A] (verified) . . . . .	120
3.13.4	Maple [A] (verified) . . . . .	122
3.13.5	Fricas [B] (verification not implemented) . . . . .	123
3.13.6	Sympy [F] . . . . .	123
3.13.7	Maxima [B] (verification not implemented) . . . . .	123
3.13.8	Giac [F] . . . . .	124
3.13.9	Mupad [F(-1)] . . . . .	124

#### 3.13.1 Optimal result

Integrand size = 10, antiderivative size = 90

$$\int x \tan^3(a + bx) dx = \frac{x}{2b} - \frac{ix^2}{2} + \frac{x \log(1 + e^{2i(a+bx)})}{b} - \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{\tan(a + bx)}{2b^2} + \frac{x \tan^2(a + bx)}{2b}$$

output `1/2*x/b-1/2*I*x^2+x*ln(1+exp(2*I*(b*x+a)))/b-1/2*I*polylog(2,-exp(2*I*(b*x+a)))/b^2-1/2*tan(b*x+a)/b^2+1/2*x*tan(b*x+a)^2/b`

#### 3.13.2 Mathematica [A] (verified)

Time = 4.58 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.90

$$\int x \tan^3(a + bx) dx = \frac{ibx(\pi + 2 \arctan(\cot(a))) + \pi \log(1 + e^{-2ibx}) + 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))}) - \pi}{1}$$

input `Integrate[x*Tan[a + b*x]^3,x]`



```
output (I*b*x*(Pi + 2*ArcTan[Cot[a]]) + Pi*Log[1 + E^((-2*I)*b*x)] + 2*(b*x - Arc
Tan[Cot[a]])*Log[1 - E^((2*I)*(b*x - ArcTan[Cot[a]]))] - Pi*Log[Cos[b*x]]
+ 2*ArcTan[Cot[a]]*Log[Sin[b*x - ArcTan[Cot[a]]]] - I*PolyLog[2, E^((2*I)*
(b*x - ArcTan[Cot[a]]))] + b*x*Sec[a + b*x]^2 - Sec[a]*Sec[a + b*x]*Sin[b*
x] - b^2*x^2*Tan[a] + (b^2*x^2*sqrt[Csc[a]^2]*Tan[a])/E^(I*ArcTan[Cot[a]]
)/(2*b^2)
```

### 3.13.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.09, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {3042, 4203, 3042, 3954, 24, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tan^3(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \tan(a + bx)^3 dx \\
 & \quad \downarrow \text{4203} \\
 & -\frac{\int \tan^2(a + bx) dx}{2b} - \int x \tan(a + bx) dx + \frac{x \tan^2(a + bx)}{2b} \\
 & \quad \downarrow \text{3042} \\
 & -\int x \tan(a + bx) dx - \frac{\int \tan(a + bx)^2 dx}{2b} + \frac{x \tan^2(a + bx)}{2b} \\
 & \quad \downarrow \text{3954} \\
 & -\frac{\frac{\tan(a+bx)}{b} - \int 1 dx}{2b} - \int x \tan(a + bx) dx + \frac{x \tan^2(a + bx)}{2b} \\
 & \quad \downarrow \text{24} \\
 & -\int x \tan(a + bx) dx + \frac{x \tan^2(a + bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} \\
 & \quad \downarrow \text{4202} \\
 & 2i \int \frac{e^{2i(a+bx)} x}{1 + e^{2i(a+bx)}} dx + \frac{x \tan^2(a + bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 2620 \\
2i \left( \frac{i \int \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) + \frac{x \tan^2(a+bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2} \\
& \downarrow 2715 \\
2i \left( \frac{\int e^{-2i(a+bx)} \log(1 + e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) + \frac{x \tan^2(a+bx)}{2b} - \\
\frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2} \\
& \downarrow 2838 \\
2i \left( -\frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) + \frac{x \tan^2(a+bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2}
\end{aligned}$$

input `Int[x*Tan[a + b*x]^3,x]`

output `(-1/2*I)*x^2 + (2*I)*((( -1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))])/b - PolyLog[2, -E^((2*I)*(a + b*x))]/(4*b^2)) + (x*Tan[a + b*x]^2)/(2*b) - (-x + Tan[a + b*x]/b)/(2*b)`

### 3.13.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4202 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 4203 `Int[((c_.) + (d_.)*(x_)^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Simp[b*d*(m/(f*(n - 1))) Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Simp[b^2 Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]`

### 3.13.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.36

method	result	size
risch	$-\frac{ix^2}{2} + \frac{2bx e^{2i(bx+a)} - ie^{2i(bx+a)} - i}{b^2 (e^{2i(bx+a)} + 1)^2} - \frac{2iax}{b} - \frac{ia^2}{b^2} + \frac{x \ln(e^{2i(bx+a)} + 1)}{b} - \frac{i \operatorname{Li}_2(-e^{2i(bx+a)})}{2b^2} + \frac{2a \ln(e^{i(bx+a)})}{b^2}$	122

input `int(x*tan(b*x+a)^3,x,method=_RETURNVERBOSE)`

output 
$$-1/2*I*x^2 + (2*b*x*\exp(2*I*(b*x+a)) - I*\exp(2*I*(b*x+a)) - I)/b^2 / (\exp(2*I*(b*x+a)) + 1)^2 - 2*I/b*a*x - I/b^2*a^2 + x*\ln(\exp(2*I*(b*x+a)) + 1)/b - 1/2*I*\operatorname{polylog}(2, -\exp(2*I*(b*x+a))) / b^2 + 2/b^2*a*\ln(\exp(I*(b*x+a)))$$

### 3.13.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 146 vs.  $2(71) = 142$ .

Time = 0.26 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.62

$$\int x \tan^3(a + bx) dx = \frac{2bx \tan^2(bx + a) + 2bx \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2bx \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2bx + i \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right)}{4b^2}$$

input `integrate(x*tan(b*x+a)^3,x, algorithm="fricas")`

output `1/4*(2*b*x*tan(b*x + a)^2 + 2*b*x*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*b*x*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*b*x + I*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - I*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 2*tan(b*x + a))/b^2`

### 3.13.6 Sympy [F]

$$\int x \tan^3(a + bx) dx = \int x \tan^3(a + bx) dx$$

input `integrate(x*tan(b*x+a)**3,x)`

output `Integral(x*tan(a + b*x)**3, x)`

### 3.13.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 386 vs.  $2(71) = 142$ .

Time = 0.78 (sec) , antiderivative size = 386, normalized size of antiderivative = 4.29

$$\int x \tan^3(a + bx) dx = \frac{b^2 x^2 \cos(4bx + 4a) + i b^2 x^2 \sin(4bx + 4a) + b^2 x^2 - 2(bx \cos(4bx + 4a) + 2bx \cos(2bx + 2a) + i bx \sin(4bx + 4a) + 2bx \sin(2bx + 2a))}{4b^2}$$

input `integrate(x*tan(b*x+a)^3,x, algorithm="maxima")`

output `-(b^2*x^2*cos(4*b*x + 4*a) + I*b^2*x^2*sin(4*b*x + 4*a) + b^2*x^2 - 2*(b*x*cos(4*b*x + 4*a) + 2*b*x*cos(2*b*x + 2*a) + I*b*x*sin(4*b*x + 4*a) + 2*I*b*x*sin(2*b*x + 2*a) + b*x)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 2*(b^2*x^2 + 2*I*b*x + 1)*cos(2*b*x + 2*a) + (cos(4*b*x + 4*a) + 2*cos(2*b*x + 2*a) + I*sin(4*b*x + 4*a) + 2*I*sin(2*b*x + 2*a) + 1)*dilog(-e^(2*I*b*x + 2*I*a)) - (-I*b*x*cos(4*b*x + 4*a) - 2*I*b*x*cos(2*b*x + 2*a) + b*x*sin(4*b*x + 4*a) + 2*b*x*sin(2*b*x + 2*a) - I*b*x)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 2*(I*b^2*x^2 - 2*b*x + I)*sin(2*b*x + 2*a) + 2)/(-2*I*b^2*cos(4*b*x + 4*a) - 4*I*b^2*cos(2*b*x + 2*a) + 2*b^2*sin(4*b*x + 4*a) + 4*b^2*sin(2*b*x + 2*a) - 2*I*b^2)`

### 3.13.8 Giac [F]

$$\int x \tan^3(a + bx) dx = \int x \tan(bx + a)^3 dx$$

input `integrate(x*tan(b*x+a)^3,x, algorithm="giac")`

output `integrate(x*tan(b*x + a)^3, x)`

### 3.13.9 Mupad [F(-1)]

Timed out.

$$\int x \tan^3(a + bx) dx = \int x \tan(a + bx)^3 dx$$

input `int(x*tan(a + b*x)^3,x)`

output `int(x*tan(a + b*x)^3, x)`

### 3.14 $\int \frac{\tan^3(a+bx)}{x} dx$

3.14.1	Optimal result	125
3.14.2	Mathematica [N/A]	125
3.14.3	Rubi [N/A]	126
3.14.4	Maple [N/A] (verified)	127
3.14.5	Fricas [N/A]	127
3.14.6	Sympy [N/A]	127
3.14.7	Maxima [N/A]	128
3.14.8	Giac [N/A]	128
3.14.9	Mupad [N/A]	129

#### 3.14.1 Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^3(a+bx)}{x} dx = \text{Int}\left(\frac{\tan^3(a+bx)}{x}, x\right)$$

output `Unintegrable(tan(b*x+a)^3/x,x)`

#### 3.14.2 Mathematica [N/A]

Not integrable

Time = 4.85 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a+bx)}{x} dx = \int \frac{\tan^3(a+bx)}{x} dx$$

input `Integrate[Tan[a + b*x]^3/x,x]`

output `Integrate[Tan[a + b*x]^3/x, x]`

### 3.14.3 Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(a + bx)}{x} dx$$

↓ 3042

$$\int \frac{\tan(a + bx)^3}{x} dx$$

↓ 4222

$$\int \frac{\tan^3(a + bx)}{x} dx$$

input `Int[Tan[a + b*x]^3/x,x]`

output `$Aborted`

#### 3.14.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4222 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], I^n*Unintegrable[(c + d*x)^m*Coth[(-I)*(e - Pi/2) - I*f*x]^n, x], I^n*Unintegrable[(c + d*x)^m*Tanh[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], (-1)^n*Unintegrable[(c + d*x)^m*Cot[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*Tan[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]`

**3.14.4 Maple [N/A] (verified)**

Not integrable

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^3(bx + a)}{x} dx$$

input `int(tan(b*x+a)^3/x,x)`output `int(tan(b*x+a)^3/x,x)`**3.14.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

input `integrate(tan(b*x+a)^3/x,x, algorithm="fricas")`output `integral(tan(b*x + a)^3/x, x)`**3.14.6 Sympy [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(a + bx)}{x} dx$$

input `integrate(tan(b*x+a)**3/x,x)`output `Integral(tan(a + b*x)**3/x, x)`



**3.14.7 Maxima [N/A]**

Not integrable

Time = 0.91 (sec) , antiderivative size = 534, normalized size of antiderivative = 44.50

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

input `integrate(tan(b*x+a)^3/x,x, algorithm="maxima")`

```
output (4*b*x*cos(2*b*x + 2*a)^2 + 4*b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + (2*b*x*cos(2*b*x + 2*a) - sin(2*b*x + 2*a))*cos(4*b*x + 4*a) - (b^2*x^2*cos(4*b*x + 4*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(4*b*x + 4*a)^2 + 4*b^2*x^2*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^2*sin(2*b*x + 2*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2 + 2*(2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2)*cos(4*b*x + 4*a))*integrate(2*(b^2*x^2 - 1)*sin(2*b*x + 2*a)/(b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(2*b*x + 2*a)^2 + 2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3), x) + (2*b*x*sin(2*b*x + 2*a) + cos(2*b*x + 2*a) + 1)*sin(4*b*x + 4*a) + sin(2*b*x + 2*a))/(b^2*x^2*cos(4*b*x + 4*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(4*b*x + 4*a)^2 + 4*b^2*x^2*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^2*sin(2*b*x + 2*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2 + 2*(2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2)*cos(4*b*x + 4*a))
```

**3.14.8 Giac [N/A]**

Not integrable

Time = 0.92 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

input `integrate(tan(b*x+a)^3/x,x, algorithm="giac")`output `integrate(tan(b*x + a)^3/x, x)`

**3.14.9 Mupad [N/A]**

Not integrable

Time = 3.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(a + bx)^3}{x} dx$$

input `int(tan(a + b*x)^3/x,x)`

output `int(tan(a + b*x)^3/x, x)`

### 3.15 $\int \frac{\tan^3(a+bx)}{x^2} dx$

3.15.1 Optimal result . . . . .	130
3.15.2 Mathematica [N/A] . . . . .	130
3.15.3 Rubi [N/A] . . . . .	131
3.15.4 Maple [N/A] (verified) . . . . .	132
3.15.5 Fricas [N/A] . . . . .	132
3.15.6 Sympy [N/A] . . . . .	132
3.15.7 Maxima [N/A] . . . . .	133
3.15.8 Giac [N/A] . . . . .	133
3.15.9 Mupad [N/A] . . . . .	134

#### 3.15.1 Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \text{Int}\left(\frac{\tan^3(a + bx)}{x^2}, x\right)$$

output `Unintegrable(tan(b*x+a)^3/x^2,x)`

#### 3.15.2 Mathematica [N/A]

Not integrable

Time = 2.58 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan^3(a + bx)}{x^2} dx$$

input `Integrate[Tan[a + b*x]^3/x^2,x]`

output `Integrate[Tan[a + b*x]^3/x^2, x]`

### 3.15.3 Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(a + bx)}{x^2} dx$$

↓ 3042

$$\int \frac{\tan(a + bx)^3}{x^2} dx$$

↓ 4222

$$\int \frac{\tan^3(a + bx)}{x^2} dx$$

input `Int [Tan[a + b*x]^3/x^2,x]`

output `$Aborted`

#### 3.15.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4222 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], I^n*Unintegrable[(c + d*x)^m*Coth[(-I)*(e - Pi/2) - I*f*x]^n, x], I^n*Unintegrable[(c + d*x)^m*Tanh[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], (-1)^n*Unintegrable[(c + d*x)^m*Cot[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*Tan[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]`

**3.15.4 Maple [N/A] (verified)**

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^3(bx + a)}{x^2} dx$$

input `int(tan(b*x+a)^3/x^2,x)`output `int(tan(b*x+a)^3/x^2,x)`**3.15.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

input `integrate(tan(b*x+a)^3/x^2,x, algorithm="fricas")`output `integral(tan(b*x + a)^3/x^2, x)`**3.15.6 Sympy [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan^3(a + bx)}{x^2} dx$$

input `integrate(tan(b*x+a)**3/x**2,x)`output `Integral(tan(a + b*x)**3/x**2, x)`

**3.15.7 Maxima [N/A]**

Not integrable

Time = 0.71 (sec) , antiderivative size = 536, normalized size of antiderivative = 44.67

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

```
input integrate(tan(b*x+a)^3/x^2,x, algorithm="maxima")
```

```
output (4*b*x*cos(2*b*x + 2*a)^2 + 4*b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + 2*(b*x*cos(2*b*x + 2*a) - sin(2*b*x + 2*a))*cos(4*b*x + 4*a) - (b^2*x^3*cos(4*b*x + 4*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(4*b*x + 4*a)^2 + 4*b^2*x^3*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^3*sin(2*b*x + 2*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3 + 2*(2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3)*cos(4*b*x + 4*a))*integrate(2*(b^2*x^2 - 3)*sin(2*b*x + 2*a)/(b^2*x^4*cos(2*b*x + 2*a)^2 + b^2*x^4*sin(2*b*x + 2*a)^2 + 2*b^2*x^4*cos(2*b*x + 2*a) + b^2*x^4), x) + 2*(b*x*sin(2*b*x + 2*a) + cos(2*b*x + 2*a) + 1)*sin(4*b*x + 4*a) + 2*sin(2*b*x + 2*a))/(b^2*x^3*cos(4*b*x + 4*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(4*b*x + 4*a)^2 + 4*b^2*x^3*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^3*sin(2*b*x + 2*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3 + 2*(2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3)*cos(4*b*x + 4*a))
```

**3.15.8 Giac [N/A]**

Not integrable

Time = 0.96 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

```
input integrate(tan(b*x+a)^3/x^2,x, algorithm="giac")
```

```
output integrate(tan(b*x + a)^3/x^2, x)
```

**3.15.9 Mupad [N/A]**

Not integrable

Time = 2.83 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)^3}{x^2} dx$$

input `int(tan(a + b*x)^3/x^2,x)`

output `int(tan(a + b*x)^3/x^2, x)`

**3.16** 
$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2 \sqrt{\tan(a+bx)} \right) dx$$

3.16.1	Optimal result	135
3.16.2	Mathematica [A] (verified)	135
3.16.3	Rubi [A] (verified)	136
3.16.4	Maple [F]	136
3.16.5	Fricas [A] (verification not implemented)	137
3.16.6	Sympy [F]	137
3.16.7	Maxima [F]	137
3.16.8	Giac [F]	138
3.16.9	Mupad [B] (verification not implemented)	138

**3.16.1 Optimal result**

Integrand size = 45, antiderivative size = 18

$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2 \sqrt{\tan(a+bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(a+bx)}}$$

output `-2*x^2/b/tan(b*x+a)^(1/2)`

**3.16.2 Mathematica [A] (verified)**

Time = 1.81 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2 \sqrt{\tan(a+bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(a+bx)}}$$

input `Integrate[x^2/Tan[a + b*x]^(3/2) - (4*x)/(b*Sqrt[Tan[a + b*x]]) + x^2*Sqrt[Tan[a + b*x]],x]`

output `(-2*x^2)/(b*Sqrt[Tan[a + b*x]])`

---

3.16. 
$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2 \sqrt{\tan(a+bx)} \right) dx$$



### 3.16.3 Rubi [A] (verified)

Time = 2.91 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.022$ , Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} + x^2 \sqrt{\tan(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{2x^2}{b\sqrt{\tan(a+bx)}}$$

input `Int[x^2/Tan[a + b*x]^(3/2) - (4*x)/(b*Sqrt[Tan[a + b*x]]) + x^2*Sqrt[Tan[a + b*x]],x]`

output `(-2*x^2)/(b*Sqrt[Tan[a + b*x]])`

#### 3.16.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

### 3.16.4 Maple [F]

$$\int \left( -\frac{4x}{b\sqrt{\tan(bx+a)}} + x^2 \left( \sqrt{\tan(bx+a)} \right) + \frac{x^2}{\tan(bx+a)^{\frac{3}{2}}} \right) dx$$

input `int(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2),x)`

output `int(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2),x)`

---

3.16.  $\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2 \sqrt{\tan(a+bx)} \right) dx$

**3.16.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(bx+a)}}$$

input `integrate(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2),x, algorithm="fricas")`

output `-2*x^2/(b*sqrt(tan(b*x + a)))`

**3.16.6 Sympy [F]**

$$\begin{aligned} & \int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx \\ &= \frac{\int \left( -\frac{4x}{\sqrt{\tan(a+bx)}} \right) dx + \int \frac{bx^2}{\tan^{\frac{3}{2}}(a+bx)} dx + \int bx^2\sqrt{\tan(a+bx)} dx}{b} \end{aligned}$$

input `integrate(-4*x/b/tan(b*x+a)**(1/2)+x**2*tan(b*x+a)**(1/2)+x**2/tan(b*x+a)**(3/2),x)`

output `(Integral(-4*x/sqrt(tan(a + b*x)), x) + Integral(b*x**2/tan(a + b*x)**(3/2), x) + Integral(b*x**2*sqrt(tan(a + b*x)), x))/b`

**3.16.7 Maxima [F]**

$$\begin{aligned} & \int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx \\ &= \int x^2\sqrt{\tan(bx+a)} + \frac{x^2}{\tan(bx+a)^{\frac{3}{2}}} - \frac{4x}{b\sqrt{\tan(bx+a)}} dx \end{aligned}$$

---

3.16.  $\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$

input `integrate(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2),x, algorithm="maxima")`

output `integrate(x^2*sqrt(tan(b*x + a)) + x^2/tan(b*x + a)^(3/2) - 4*x/(b*sqrt(tan(b*x + a))), x)`

### 3.16.8 Giac [F]

$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$$

$$= \int x^2\sqrt{\tan(bx+a)} + \frac{x^2}{\tan(bx+a)^{\frac{3}{2}}} - \frac{4x}{b\sqrt{\tan(bx+a)}} dx$$

input `integrate(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2),x, algorithm="giac")`

output `integrate(x^2*sqrt(tan(b*x + a)) + x^2/tan(b*x + a)^(3/2) - 4*x/(b*sqrt(tan(b*x + a))), x)`

### 3.16.9 Mupad [B] (verification not implemented)

Time = 3.61 (sec) , antiderivative size = 50, normalized size of antiderivative = 2.78

$$\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$$

$$= -\frac{x^2 \sin(2a+2bx) \sqrt{\frac{\sin(2a+2bx)}{\cos(2a+2bx)+1}}}{b \sin(a+bx)^2}$$

input `int(x^2*tan(a + b*x)^(1/2) + x^2/tan(a + b*x)^(3/2) - (4*x)/(b*tan(a + b*x)^(1/2)),x)`

output `-(x^2*sin(2*a + 2*b*x)*(sin(2*a + 2*b*x)/(cos(2*a + 2*b*x) + 1))^(1/2))/(b*sin(a + b*x)^2)`

---

3.16.  $\int \left( \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$

$$3.17 \quad \int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$$

3.17.1	Optimal result	139
3.17.2	Mathematica [A] (verified)	139
3.17.3	Rubi [F]	140
3.17.4	Maple [F]	140
3.17.5	Fricas [A] (verification not implemented)	141
3.17.6	Sympy [F]	141
3.17.7	Maxima [F]	141
3.17.8	Giac [F]	142
3.17.9	Mupad [B] (verification not implemented)	142

### 3.17.1 Optimal result

Integrand size = 49, antiderivative size = 17

$$\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx = \frac{x\sqrt{\tan(a+bx^2)}}{b}$$

output `x*tan(b*x^2+a)^(1/2)/b`

### 3.17.2 Mathematica [A] (verified)

Time = 0.92 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx = \frac{x\sqrt{\tan(a+bx^2)}}{b}$$

input `Integrate[x^2/Sqrt[Tan[a + b*x^2]] + Sqrt[Tan[a + b*x^2]]/b + x^2*Tan[a + b*x^2]^(3/2), x]`

output `(x*Sqrt[Tan[a + b*x^2]])/b`

---

3.17.  $\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$

### 3.17.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left( x^2 \tan^{\frac{3}{2}}(a + bx^2) + \frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} \right) dx$$

↓ 2009

$$\int x^2 \tan^{\frac{3}{2}}(bx^2 + a) dx + \int \frac{x^2}{\sqrt{\tan(bx^2 + a)}} dx + \frac{\int \sqrt{\tan(bx^2 + a)} dx}{b}$$

input `Int[x^2/Sqrt[Tan[a + b*x^2]] + Sqrt[Tan[a + b*x^2]]/b + x^2*Tan[a + b*x^2]^(3/2),x]`

output `$Aborted`

#### 3.17.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

### 3.17.4 Maple [F]

$$\int \left( \frac{x^2}{\sqrt{\tan(x^2b + a)}} + \frac{\sqrt{\tan(x^2b + a)}}{b} + x^2 \left( \tan^{\frac{3}{2}}(x^2b + a) \right) \right) dx$$

input `int(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x)`

output `int(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x)`

---

3.17.  $\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx$

**3.17.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx = \frac{x\sqrt{\tan(bx^2+a)}}{b}$$

input `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="fricas")`

output `x*sqrt(tan(b*x^2 + a))/b`

**3.17.6 Sympy [F]**

$$\begin{aligned} & \int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx \\ &= \frac{\int \frac{bx^2}{\sqrt{\tan(a+bx^2)}} dx + \int bx^2 \tan^{\frac{3}{2}}(a+bx^2) dx + \int \sqrt{\tan(a+bx^2)} dx}{b} \end{aligned}$$

input `integrate(x**2/tan(b*x**2+a)**(1/2)+tan(b*x**2+a)**(1/2)/b+x**2*tan(b*x**2+a)**(3/2),x)`

output `(Integral(b*x**2/sqrt(tan(a + b*x**2)), x) + Integral(b*x**2*tan(a + b*x**2)**(3/2), x) + Integral(sqrt(tan(a + b*x**2)), x))/b`

**3.17.7 Maxima [F]**

$$\begin{aligned} & \int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx \\ &= \int x^2 \tan(bx^2+a)^{\frac{3}{2}} + \frac{x^2}{\sqrt{\tan(bx^2+a)}} + \frac{\sqrt{\tan(bx^2+a)}}{b} dx \end{aligned}$$

---

3.17.  $\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$

input `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate(x^2*tan(b*x^2 + a)^(3/2) + x^2/sqrt(tan(b*x^2 + a)) + sqrt(tan(b*x^2 + a))/b, x)`

### 3.17.8 Giac [F]

$$\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$$

$$= \int x^2 \tan(bx^2+a)^{\frac{3}{2}} + \frac{x^2}{\sqrt{\tan(bx^2+a)}} + \frac{\sqrt{\tan(bx^2+a)}}{b} dx$$

input `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="giac")`

output `integrate(x^2*tan(b*x^2 + a)^(3/2) + x^2/sqrt(tan(b*x^2 + a)) + sqrt(tan(b*x^2 + a))/b, x)`

### 3.17.9 Mupad [B] (verification not implemented)

Time = 5.45 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.65

$$\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx = \frac{x \sqrt{-\frac{e^{2i} b x^2 + a 2i 1i - i}{e^{2i} b x^2 + a 2i + 1}}}{b}$$

input `int(tan(a + b*x^2)^(1/2)/b + x^2/tan(a + b*x^2)^(1/2) + x^2*tan(a + b*x^2)^(3/2),x)`

output `(x*(-(exp(a*2i + b*x^2*2i)*1i - 1i)/(exp(a*2i + b*x^2*2i) + 1))^(1/2))/b`

---

3.17.  $\int \left( \frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$

### 3.18 $\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx$

3.18.1	Optimal result . . . . .	143
3.18.2	Mathematica [A] (verified) . . . . .	143
3.18.3	Rubi [A] (verified) . . . . .	144
3.18.4	Maple [A] (verified) . . . . .	146
3.18.5	Fricas [A] (verification not implemented) . . . . .	147
3.18.6	Sympy [A] (verification not implemented) . . . . .	147
3.18.7	Maxima [F(-2)] . . . . .	148
3.18.8	Giac [A] (verification not implemented) . . . . .	148
3.18.9	Mupad [B] (verification not implemented) . . . . .	149

#### 3.18.1 Optimal result

Integrand size = 23, antiderivative size = 189

$$\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx = \frac{3id^3x}{8af^3} - \frac{3d(c+dx)^2}{8af^2} - \frac{i(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^2(c+dx)}{8f^4(a+ia \tan(e+fx))} - \frac{3id^2(c+dx)}{4f^3(a+ia \tan(e+fx))} + \frac{3d(c+dx)^2}{4f^2(a+ia \tan(e+fx))} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))}$$

```
output 3/8*I*d^3*x/a/f^3-3/8*d*(d*x+c)^2/a/f^2-1/4*I*(d*x+c)^3/a/f+1/8*(d*x+c)^4/a/d-3/8*d^3/f^4/(a+I*a*tan(f*x+e))-3/4*I*d^2*(d*x+c)/f^3/(a+I*a*tan(f*x+e))+3/4*d*(d*x+c)^2/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)^3/f/(a+I*a*tan(f*x+e))
```

#### 3.18.2 Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.47

$$\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx = \frac{\sec(e+fx)(\cos(fx)+i \sin(fx))((4ic^3f^3+6c^2df^2(1+2ifx)+6cd^2f(-i+2fx+2if^2x^2)+d^3(-3-6i$$



input `Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x]),x]`

output `(Sec[e + f*x]*(Cos[f*x] + I*Sin[f*x])*(((4*I)*c^3*f^3 + 6*c^2*d*f^2*(1 + (2*I)*f*x) + 6*c*d^2*f*(-I + 2*f*x + (2*I)*f^2*x^2) + d^3*(-3 - (6*I)*f*x + 6*f^2*x^2 + (4*I)*f^3*x^3))*Cos[2*f*x]*(Cos[e] - I*Sin[e]) + 2*f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(Cos[e] + I*Sin[e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(-I + 2*f*x) + 6*c*d^2*f*(-1 - (2*I)*f*x + 2*f^2*x^2) + d^3*(3*I - 6*f*x - (6*I)*f^2*x^2 + 4*f^3*x^3))*(Cos[e] - I*Sin[e])*Sin[2*f*x]))/(16*f^4*(a + I*a*Tan[e + f*x]))`

### 3.18.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$ , Rules used = {3042, 4206, 3042, 4206, 3042, 4206, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx \\
 & \quad \downarrow \text{4206} \\
 & -\frac{3id \int \frac{(c+dx)^2}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3id \int \frac{(c+dx)^2}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
 & \quad \downarrow \text{4206} \\
 & -\frac{3id \left( -\frac{id \int \frac{c+dx}{i \tan(e+fx)a+a} dx}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.18.  $\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx$

$$\begin{aligned}
 & \frac{3id \left( -\frac{id \int \frac{c+dx}{i \tan(e+fx)a+a} dx}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
 & \quad \downarrow 4206 \\
 & \frac{3id \left( -\frac{id \left( -\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \\
 & \quad \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
 & \quad \downarrow 3042 \\
 & \frac{3id \left( -\frac{id \left( -\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \\
 & \quad \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
 & \quad \downarrow 3960 \\
 & \frac{3id \left( -\frac{id \left( -\frac{id \left( \frac{f1dx}{2a} + \frac{i}{2f(a+ia \tan(e+fx))} \right)}{2f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \\
 & \quad \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
 & \quad \downarrow 24 \\
 & \frac{3id \left( \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} - \frac{id \left( \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} - \frac{id \left( \frac{x}{2a} + \frac{i}{2f(a+ia \tan(e+fx))} \right)}{2f} \right)}{f} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \\
 & \quad \frac{(c+dx)^4}{8ad}
 \end{aligned}$$

input `Int[(c + d*x)^3/(a + I*a*Tan[e + f*x]),x]`

3.18.  $\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx$

```
output (c + d*x)^4/(8*a*d) + ((I/2)*(c + d*x)^3)/(f*(a + I*a*Tan[e + f*x])) - (((
3*I)/2)*d*((c + d*x)^3/(6*a*d) + ((I/2)*(c + d*x)^2)/(f*(a + I*a*Tan[e + f
*x]))) - (I*d*((c + d*x)^2/(4*a*d) + ((I/2)*(c + d*x))/(f*(a + I*a*Tan[e +
f*x]))) - ((I/2)*d*(x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))))/f)/f
```

### 3.18.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3960 Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a +
b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^
(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

```
rule 4206 Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Sy
mbol] := Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Simp[a*d*(m/(2*b*f))
Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m
/(2*b*f*(a + b*Tan[e + f*x]))], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[
a^2 + b^2, 0] && GtQ[m, 0]
```

### 3.18.4 Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.90

method	result
risch	$\frac{d^3 x^4}{8a} + \frac{d^2 c x^3}{2a} + \frac{3d c^2 x^2}{4a} + \frac{c^3 x}{2a} + \frac{c^4}{8ad} + \frac{i(4d^3 x^3 f^3 + 12c d^2 f^3 x^2 - 6id^3 f^2 x^2 + 12c^2 d f^3 x - 12ic d^2 f^2 x + 4c^3 f^3 - 6ic^2 d f^2 - 6d^3 f^3)}{16a f^4}$
default	Expression too large to display

```
input int((d*x+c)^3/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/8/a*d^3*x^4+1/2/a*d^2*c*x^3+3/4/a*d*c^2*x^2+1/2/a*c^3*x+1/8/a/d*c^4+1/16
*I*(4*d^3*x^3*f^3-6*I*d^3*f^2*x^2+12*c*d^2*f^3*x^2-12*I*c*d^2*f^2*x+12*c^2
*d*f^3*x-6*I*c^2*d*f^2+4*c^3*f^3-6*d^3*f*x+3*I*d^3-6*c*d^2*f)/a/f^4*exp(-2
*I*(f*x+e))
```

### 3.18.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.87

$$\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx$$

$$= \frac{(4i d^3 f^3 x^3 + 4i c^3 f^3 + 6 c^2 d f^2 - 6i c d^2 f - 3 d^3 - 6(-2i c d^2 f^3 - d^3 f^2)x^2 - 6(-2i c^2 d f^3 - 2 c d^2 f^2 + i d^3 f^3))}{16 a f^4}$$

```
input integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="fricas")
```

```
output 1/16*(4*I*d^3*f^3*x^3 + 4*I*c^3*f^3 + 6*c^2*d*f^2 - 6*I*c*d^2*f - 3*d^3 -
6*(-2*I*c*d^2*f^3 - d^3*f^2)*x^2 - 6*(-2*I*c^2*d*f^3 - 2*c*d^2*f^2 + I*d^3
*f)*x + 2*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*f^4*x)*
e^(2*I*f*x + 2*I*e))*e^(-2*I*f*x - 2*I*e)/(a*f^4)
```

### 3.18.6 Sympy [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.37

$$\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx$$

$$= \begin{cases} \frac{(4ic^3 f^3 + 12ic^2 d f^3 x + 6c^2 d f^2 + 12ic d^2 f^3 x^2 + 12c d^2 f^2 x - 6ic d^2 f + 4id^3 f^3 x^3 + 6d^3 f^2 x^2 - 6id^3 f x - 3d^3) e^{-2ie} e^{-2ifx}}{16af^4} & \text{for } af^4 e^{2ie} \neq 0 \\ \frac{c^3 x e^{-2ie}}{2a} + \frac{3c^2 d x^2 e^{-2ie}}{4a} + \frac{c d^2 x^3 e^{-2ie}}{2a} + \frac{d^3 x^4 e^{-2ie}}{8a} & \text{otherwise} \\ + \frac{c^3 x}{2a} + \frac{3c^2 d x^2}{4a} + \frac{c d^2 x^3}{2a} + \frac{d^3 x^4}{8a} & \end{cases}$$

```
input integrate((d*x+c)**3/(a+I*a*tan(f*x+e)),x)
```

```
output Piecewise(((4*I*c**3*f**3 + 12*I*c**2*d*f**3*x + 6*c**2*d*f**2 + 12*I*c*d*
*2*f**3*x**2 + 12*c*d**2*f**2*x - 6*I*c*d**2*f + 4*I*d**3*f**3*x**3 + 6*d*
*3*f**2*x**2 - 6*I*d**3*f*x - 3*d**3)*exp(-2*I*e)*exp(-2*I*f*x)/(16*a*f**4
), Ne(a*f**4*exp(2*I*e), 0)), (c**3*x*exp(-2*I*e)/(2*a) + 3*c**2*d*x**2*ex
p(-2*I*e)/(4*a) + c*d**2*x**3*exp(-2*I*e)/(2*a) + d**3*x**4*exp(-2*I*e)/(8
*a), True)) + c**3*x/(2*a) + 3*c**2*d*x**2/(4*a) + c*d**2*x**3/(2*a) + d**
3*x**4/(8*a)
```

### 3.18.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

### 3.18.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.99

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx = \frac{(2d^3 f^4 x^4 e^{(2i fx + 2i e)} + 8cd^2 f^4 x^3 e^{(2i fx + 2i e)} + 12c^2 d f^4 x^2 e^{(2i fx + 2i e)} + 4i d^3 f^3 x^3 + 8c^3 f^4 x e^{(2i fx + 2i e)} + 12i c^2 d f^3 x^2 + 12I c^2 d^2 f^3 x + 6d^3 f^2 x^2 + 4I c^3 f^3 + 12c d^2 f^2 x + 6c^2 d f^2 - 6I d^3 f x - 6I c d^2 f - 3d^3) * e^{(-2I f x - 2I e)}}{(a f^4)} + 16 a$$

```
input integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="giac")
```

```
output 1/16*(2*d^3*f^4*x^4*e^(2*I*f*x + 2*I*e) + 8*c*d^2*f^4*x^3*e^(2*I*f*x + 2*I
*e) + 12*c^2*d*f^4*x^2*e^(2*I*f*x + 2*I*e) + 4*I*d^3*f^3*x^3 + 8*c^3*f^4*x
*e^(2*I*f*x + 2*I*e) + 12*I*c*d^2*f^3*x^2 + 12*I*c^2*d*f^3*x + 6*d^3*f^2*x
^2 + 4*I*c^3*f^3 + 12*c*d^2*f^2*x + 6*c^2*d*f^2 - 6*I*d^3*f*x - 6*I*c*d^2*
f - 3*d^3)*e^(-2*I*f*x - 2*I*e)/(a*f^4)
```

**3.18.9 Mupad [B] (verification not implemented)**

Time = 4.12 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.24

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx$$

$$= \frac{8c^3 f^4 x - 3d^3 \cos(2e + 2fx) + 4c^3 f^3 \sin(2e + 2fx) + 2d^3 f^4 x^4 + 6c^2 d f^2 \cos(2e + 2fx) + 12c^2 d f^3 x^3 + 6c^2 d^2 f^2 \sin(2e + 2fx) + 12c^2 d^2 f^3 x^2 + 6c^2 d^3 f^2 x + 6c^2 d^3 f^3 \sin(2e + 2fx) + 12c^2 d^3 f^4 x^4 + 6c^2 d^4 f^2 \cos(2e + 2fx) + 12c^2 d^4 f^3 x^3 + 6c^2 d^4 f^4 x^4}{16a^4}$$

input `int((c + d*x)^3/(a + a*tan(e + f*x)*1i),x)`

output

```
(d^3*sin(2*e + 2*f*x)*3i - 3*d^3*cos(2*e + 2*f*x) + 8*c^3*f^4*x + c^3*f^3*cos(2*e + 2*f*x)*4i + 4*c^3*f^3*sin(2*e + 2*f*x) + 2*d^3*f^4*x^4 + 6*c^2*d*f^2*cos(2*e + 2*f*x) - c^2*d*f^2*sin(2*e + 2*f*x)*6i + 12*c^2*d*f^4*x^2 + 8*c*d^2*f^4*x^3 + 6*d^3*f^2*x^2*cos(2*e + 2*f*x) + d^3*f^3*x^3*cos(2*e + 2*f*x)*4i - d^3*f^2*x^2*sin(2*e + 2*f*x)*6i + 4*d^3*f^3*x^3*sin(2*e + 2*f*x) - c*d^2*f*cos(2*e + 2*f*x)*6i - 6*c*d^2*f*sin(2*e + 2*f*x) - d^3*f*x*cos(2*e + 2*f*x)*6i - 6*d^3*f*x*sin(2*e + 2*f*x) + 12*c*d^2*f^2*x*cos(2*e + 2*f*x) + c^2*d*f^3*x*cos(2*e + 2*f*x)*12i - c*d^2*f^2*x*sin(2*e + 2*f*x)*12i + 12*c^2*d*f^3*x*sin(2*e + 2*f*x) + c*d^2*f^3*x^2*cos(2*e + 2*f*x)*12i + 12*c*d^2*f^3*x^2*sin(2*e + 2*f*x))/(16*a*f^4)
```

### 3.19 $\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx$

3.19.1	Optimal result	150
3.19.2	Mathematica [A] (verified)	150
3.19.3	Rubi [A] (verified)	151
3.19.4	Maple [A] (verified)	153
3.19.5	Fricas [A] (verification not implemented)	153
3.19.6	Sympy [A] (verification not implemented)	154
3.19.7	Maxima [F(-2)]	154
3.19.8	Giac [A] (verification not implemented)	154
3.19.9	Mupad [B] (verification not implemented)	155

#### 3.19.1 Optimal result

Integrand size = 23, antiderivative size = 137

$$\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx = -\frac{d^2x}{4af^2} - \frac{i(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{id^2}{4f^3(a+ia \tan(e+fx))} + \frac{d(c+dx)}{2f^2(a+ia \tan(e+fx))} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))}$$

output `-1/4*d^2*x/a/f^2-1/4*I*(d*x+c)^2/a/f+1/6*(d*x+c)^3/a/d-1/4*I*d^2/f^3/(a+I*a*tan(f*x+e))+1/2*d*(d*x+c)/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)^2/f/(a+I*a*tan(f*x+e))`

#### 3.19.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.30

$$\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx = \frac{\sec(e+fx)(\cos(fx)+i \sin(fx))((d+(1+i)cf+(1+i)dfx)((1+i)cf+d(-i+(1+i)fx)) \cos(2fx))}{\dots}$$

input `Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x]),x]`

output  $(\text{Sec}[e + f*x] * (\text{Cos}[f*x] + I*\text{Sin}[f*x]) * ((d + (1 + I)*c*f + (1 + I)*d*f*x) * ((1 + I)*c*f + d*(-I + (1 + I)*f*x)) * \text{Cos}[2*f*x] * (\text{Cos}[e] - I*\text{Sin}[e]) + (4*f^3*x*(3*c^2 + 3*c*d*x + d^2*x^2) * (\text{Cos}[e] + I*\text{Sin}[e])) / 3 - I*(d + (1 + I)*c*f + (1 + I)*d*f*x) * ((1 + I)*c*f + d*(-I + (1 + I)*f*x)) * (\text{Cos}[e] - I*\text{Sin}[e]) * \text{Sin}[2*f*x])) / (8*f^3*(a + I*a*\text{Tan}[e + f*x]))$

### 3.19.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {3042, 4206, 3042, 4206, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx$$

↓ 4206

$$-\frac{id \int \frac{c+dx}{i \tan(e+fx)a+a} dx}{f} + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^3}{6ad}$$

↓ 3042

$$-\frac{id \int \frac{c+dx}{i \tan(e+fx)a+a} dx}{f} + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^3}{6ad}$$

↓ 4206

$$-\frac{id \left( -\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^3}{6ad}$$

↓ 3042

$$-\frac{id \left( -\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^3}{6ad}$$

↓ 3960

---

3.19.  $\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx$



$$\begin{aligned}
 & - \frac{id \left( -\frac{id \left( \frac{\int 1 dx}{2a} + \frac{i}{2f(a+ia \tan(e+fx))} \right) + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \\
 & \qquad \qquad \qquad \frac{(c+dx)^3}{6ad} \\
 & \qquad \qquad \qquad \downarrow 24 \\
 & \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} - \frac{id \left( \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} - \frac{id \left( \frac{x}{2a} + \frac{i}{2f(a+ia \tan(e+fx))} \right) \right)}{f} + \frac{(c+dx)^3}{6ad}
 \end{aligned}$$

```
input Int[(c + d*x)^2/(a + I*a*Tan[e + f*x]),x]
```

```
output (c + d*x)^3/(6*a*d) + ((I/2)*(c + d*x)^2)/(f*(a + I*a*Tan[e + f*x])) - (I*d*((c + d*x)^2/(4*a*d) + ((I/2)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])) - ((I/2)*d*(x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))))/f)/f
```

3.19.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3960 Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

```
rule 4206 Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Simp[a*d*(m/(2*b*f)) Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]
```

### 3.19.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.79

method	result
risch	$\frac{d^2 x^3}{6a} + \frac{dcx^2}{2a} + \frac{c^2 x}{2a} + \frac{c^3}{6ad} + \frac{i(2d^2 x^2 f^2 + 4cd f^2 x - 2id^2 f x + 2c^2 f^2 - 2icdf - d^2)e^{-2i(fx+e)}}{8a f^3}$
default	$\frac{\frac{ic^2(\cos^2(fx+e))}{2} - \frac{icde(\cos^2(fx+e))}{f} - \frac{2icd\left(-\frac{(fx+e)(\cos^2(fx+e))}{2} + \frac{\sin(fx+e)\cos(fx+e)}{4} + \frac{fx}{4} + \frac{e}{4}\right)}{f} + \frac{id^2 e^2(\cos^2(fx+e))}{2f^2} + \frac{2id^2 e\left(-\frac{(fx+e)}{2}\right)}{f^2}}$

input `int((d*x+c)^2/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/6/a*d^2*x^3+1/2/a*d*c*x^2+1/2/a*c^2*x+1/6/a/d*c^3+1/8*I*(2*d^2*x^2*f^2-2*I*d^2*f*x+4*c*d*f^2*x-2*I*c*d*f+2*c^2*f^2-d^2)/a/f^3*exp(-2*I*(f*x+e))`

### 3.19.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.77

$$\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx$$

$$= \frac{(6i d^2 f^2 x^2 + 6i c^2 f^2 + 6cdf - 3i d^2 - 6(-2i cdf^2 - d^2 f)x + 4(d^2 f^3 x^3 + 3cdf^3 x^2 + 3c^2 f^3 x)e^{(2i fx+2ie)})e^{(2i fx+2ie)}}{24 a f^3}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="fracas")`

output `1/24*(6*I*d^2*f^2*x^2 + 6*I*c^2*f^2 + 6*c*d*f - 3*I*d^2 - 6*(-2*I*c*d*f^2 - d^2*f)*x + 4*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 3*c^2*f^3*x)*e^(2*I*f*x + 2*I*e))*e^(-2*I*f*x - 2*I*e)/(a*f^3)`

### 3.19.6 Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.20

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx = \begin{cases} \frac{(2ic^2 f^2 + 4icdf^2 x + 2cdf + 2id^2 f^2 x^2 + 2d^2 fx - id^2) e^{-2ie} e^{-2ifx}}{8af^3} & \text{for } af^3 e^{2ie} \neq 0 \\ \frac{c^2 x e^{-2ie}}{2a} + \frac{cdx^2 e^{-2ie}}{2a} + \frac{d^2 x^3 e^{-2ie}}{6a} & \text{otherwise} \\ + \frac{c^2 x}{2a} + \frac{cdx^2}{2a} + \frac{d^2 x^3}{6a} \end{cases}$$

input `integrate((d*x+c)**2/(a+I*a*tan(f*x+e)),x)`

output `Piecewise(((2*I*c**2*f**2 + 4*I*c*d*f**2*x + 2*c*d*f + 2*I*d**2*f**2*x**2 + 2*d**2*f*x - I*d**2)*exp(-2*I*e)*exp(-2*I*f*x)/(8*a*f**3), Ne(a*f**3*exp(2*I*e), 0)), (c**2*x*exp(-2*I*e)/(2*a) + c*d*x**2*exp(-2*I*e)/(2*a) + d**2*x**3*exp(-2*I*e)/(6*a), True)) + c**2*x/(2*a) + c*d*x**2/(2*a) + d**2*x**3/(6*a)`

### 3.19.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

### 3.19.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.87

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx = \frac{(4d^2 f^3 x^3 e^{(2i f x + 2i e)} + 12cdf^3 x^2 e^{(2i f x + 2i e)} + 12c^2 f^3 x e^{(2i f x + 2i e)} + 6i d^2 f^2 x^2 + 12i cdf^2 x + 6i c^2 f^2 + 6d^2 f^2)}{24af^3}$$

---

3.19.  $\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `1/24*(4*d^2*f^3*x^3*e^(2*I*f*x + 2*I*e) + 12*c*d*f^3*x^2*e^(2*I*f*x + 2*I*e) + 12*c^2*f^3*x*e^(2*I*f*x + 2*I*e) + 6*I*d^2*f^2*x^2 + 12*I*c*d*f^2*x + 6*I*c^2*f^2 + 6*d^2*f*x + 6*c*d*f - 3*I*d^2)*e^(-2*I*f*x - 2*I*e)/(a*f^3)`

### 3.19.9 Mupad [B] (verification not implemented)

Time = 3.71 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.76

$$\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx$$

$$= \frac{12c^2 f^3 x - 3d^2 \sin(2e+2fx) + 6c^2 f^2 \sin(2e+2fx) + 4d^2 f^3 x^3 + 6cdf \cos(2e+2fx) + 6d^2 f^2}{24af^3}$$

input `int((c + d*x)^2/(a + a*tan(e + f*x)*1i),x)`

output `(12*c^2*f^3*x - 3*d^2*sin(2*e + 2*f*x) - d^2*cos(2*e + 2*f*x)*3i + c^2*f^2*cos(2*e + 2*f*x)*6i + 6*c^2*f^2*sin(2*e + 2*f*x) + 4*d^2*f^3*x^3 + 6*c*d*f*cos(2*e + 2*f*x) - c*d*f*sin(2*e + 2*f*x)*6i + d^2*f^2*x^2*cos(2*e + 2*f*x)*6i + 6*d^2*f^2*x^2*sin(2*e + 2*f*x) + 12*c*d*f^3*x^2 + 6*d^2*f*x*cos(2*e + 2*f*x) - d^2*f*x*sin(2*e + 2*f*x)*6i + c*d*f^2*x*cos(2*e + 2*f*x)*12i + 12*c*d*f^2*x*sin(2*e + 2*f*x))/(24*a*f^3)`

### 3.20 $\int \frac{c+dx}{a+ia \tan(e+fx)} dx$

3.20.1	Optimal result . . . . .	156
3.20.2	Mathematica [A] (verified) . . . . .	156
3.20.3	Rubi [A] (verified) . . . . .	157
3.20.4	Maple [A] (verified) . . . . .	158
3.20.5	Fricas [A] (verification not implemented) . . . . .	159
3.20.6	Sympy [A] (verification not implemented) . . . . .	159
3.20.7	Maxima [F(-2)] . . . . .	159
3.20.8	Giac [A] (verification not implemented) . . . . .	160
3.20.9	Mupad [B] (verification not implemented) . . . . .	160

#### 3.20.1 Optimal result

Integrand size = 21, antiderivative size = 84

$$\int \frac{c+dx}{a+ia \tan(e+fx)} dx = -\frac{idx}{4af} + \frac{(c+dx)^2}{4ad} + \frac{d}{4f^2(a+ia \tan(e+fx))} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))}$$

```
output -1/4*I*d*x/a/f+1/4*(d*x+c)^2/a/d+1/4*d/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)/f/(a+I*a*tan(f*x+e))
```

#### 3.20.2 Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.14

$$\int \frac{c+dx}{a+ia \tan(e+fx)} dx = \frac{-i(2cf(i+2fx) + d(1+2ifx+2f^2x^2)) + (2cf(-i+2fx) + d(-1-2ifx+2f^2x^2)) \tan(e+fx)}{8af^2(-i+\tan(e+fx))}$$

```
input Integrate[(c + d*x)/(a + I*a*Tan[e + f*x]),x]
```

```
output ((-I)*(2*c*f*(I + 2*f*x) + d*(1 + (2*I)*f*x + 2*f^2*x^2)) + (2*c*f*(-I + 2*f*x) + d*(-1 - (2*I)*f*x + 2*f^2*x^2))*Tan[e + f*x])/(8*a*f^2*(-I + Tan[e + f*x]))
```

### 3.20.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.07, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 4206, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{c + dx}{a + ia \tan(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{c + dx}{a + ia \tan(e + fx)} dx \\
 & \quad \downarrow \text{4206} \\
 & -\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c + dx)}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^2}{4ad} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c + dx)}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^2}{4ad} \\
 & \quad \downarrow \text{3960} \\
 & -\frac{id \left( \frac{\int 1 dx}{2a} + \frac{i}{2f(a + ia \tan(e + fx))} \right)}{2f} + \frac{i(c + dx)}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^2}{4ad} \\
 & \quad \downarrow \text{24} \\
 & \frac{i(c + dx)}{2f(a + ia \tan(e + fx))} + \frac{(c + dx)^2}{4ad} - \frac{id \left( \frac{x}{2a} + \frac{i}{2f(a + ia \tan(e + fx))} \right)}{2f}
 \end{aligned}$$

input `Int[(c + d*x)/(a + I*a*Tan[e + f*x]),x]`

output `(c + d*x)^2/(4*a*d) + ((I/2)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])) - ((I/2)*d*(x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))))/f`

### 3.20.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`
- rule 4206 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Simp[a*d*(m/(2*b*f)) Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))], x)) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]`

### 3.20.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.60

method	result
risch	$\frac{dx^2}{4a} + \frac{cx}{2a} + \frac{i(2dfx+2cf-id)e^{-2i(fx+e)}}{8af^2}$
norman	$\frac{\frac{dx^2}{4a} + \frac{2icf+d}{4af^2} + \frac{dx^2(\tan^2(fx+e))}{4a} + \frac{(2cf-id)\tan(fx+e)}{4af^2} + \frac{(2cf+id)x}{4af} + \frac{dx\tan(fx+e)}{2af} + \frac{(2cf-id)x(\tan^2(fx+e))}{4fa}}{1+\tan^2(fx+e)}$
default	$\frac{ic(\cos^2(fx+e))}{2} - \frac{ide(\cos^2(fx+e))}{2f} - \frac{id\left(-\frac{(fx+e)(\cos^2(fx+e))}{2} + \frac{\sin(fx+e)\cos(fx+e)}{4} + \frac{fx}{4} + \frac{e}{4}\right)}{f} + c\left(\frac{\sin(fx+e)\cos(fx+e)}{2} + \frac{fx}{2} + \frac{e}{2}\right) - \frac{de(\sin(fx+e))}{af}$

input `int((d*x+c)/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/4/a*d*x^2+1/2/a*c*x+1/8*I*(2*d*f*x-I*d+2*c*f)/a/f^2*exp(-2*I*(f*x+e))`

---

3.20.  $\int \frac{c+dx}{a+ia \tan(e+fx)} dx$

**3.20.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.65

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx = \frac{(2i dfx + 2i cf + 2(df^2x^2 + 2cf^2x)e^{(2i fx + 2i e)} + d)e^{(-2i fx - 2i e)}}{8af^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`output `1/8*(2*I*d*f*x + 2*I*c*f + 2*(d*f^2*x^2 + 2*c*f^2*x)*e^(2*I*f*x + 2*I*e) + d)*e^(-2*I*f*x - 2*I*e)/(a*f^2)`**3.20.6 Sympy [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.10

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx = \begin{cases} \frac{(2icf + 2idfx + d)e^{-2ie}e^{-2ifx}}{8af^2} & \text{for } af^2e^{2ie} \neq 0 \\ \frac{cxe^{-2ie}}{2a} + \frac{dx^2e^{-2ie}}{4a} & \text{otherwise} \end{cases} + \frac{cx}{2a} + \frac{dx^2}{4a}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x)`output `Piecewise(((2*I*c*f + 2*I*d*f*x + d)*exp(-2*I*e)*exp(-2*I*f*x)/(8*a*f**2), Ne(a*f**2*exp(2*I*e), 0)), (c*x*exp(-2*I*e)/(2*a) + d*x**2*exp(-2*I*e)/(4*a), True)) + c*x/(2*a) + d*x**2/(4*a)`**3.20.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`



**3.20.8 Giac [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.74

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx$$

$$= \frac{(2df^2x^2e^{(2ifx+2ie)} + 4cf^2xe^{(2ifx+2ie)} + 2idfx + 2icf + d)e^{(-2ifx-2ie)}}{8af^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="giac")`output `1/8*(2*d*f^2*x^2*e^(2*I*f*x + 2*I*e) + 4*c*f^2*x*e^(2*I*f*x + 2*I*e) + 2*I*d*f*x + 2*I*c*f + d)*e^(-2*I*f*x - 2*I*e)/(a*f^2)`**3.20.9 Mupad [B] (verification not implemented)**

Time = 3.45 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.25

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx$$

$$= \frac{d \cos(2e + 2fx) + 2df^2x^2 + 2cf \sin(2e + 2fx) + 4cf^2x + 2dfx \sin(2e + 2fx) - d \sin(2e + 2fx)}{8af^2}$$

input `int((c + d*x)/(a + a*tan(e + f*x)*1i),x)`output `(d*cos(2*e + 2*f*x) - d*sin(2*e + 2*f*x)*1i + 2*d*f^2*x^2 + c*f*cos(2*e + 2*f*x)*2i + 2*c*f*sin(2*e + 2*f*x) + 4*c*f^2*x + d*f*x*cos(2*e + 2*f*x)*2i + 2*d*f*x*sin(2*e + 2*f*x))/(8*a*f^2)`

### 3.21 $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx$

3.21.1	Optimal result	161
3.21.2	Mathematica [A] (verified)	161
3.21.3	Rubi [A] (verified)	162
3.21.4	Maple [A] (verified)	164
3.21.5	Fricas [A] (verification not implemented)	165
3.21.6	Sympy [F]	165
3.21.7	Maxima [A] (verification not implemented)	165
3.21.8	Giac [A] (verification not implemented)	166
3.21.9	Mupad [F(-1)]	166

#### 3.21.1 Optimal result

Integrand size = 23, antiderivative size = 161

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx = \frac{\cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{2ad} + \frac{\log(c+dx)}{2ad} - \frac{i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{2ad} - \frac{i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2ad} - \frac{\sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2ad}$$

output `1/2*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a/d+1/2*ln(d*x+c)/a/d-1/2*I*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a/d+1/2*I*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d+1/2*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d`

#### 3.21.2 Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.03

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx = \frac{\sec(e+fx) \left( -i \cos\left(f\left(\frac{c}{d}+x\right)\right) + \sin\left(f\left(\frac{c}{d}+x\right)\right) \right) \left( \operatorname{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) \left( \cos\left(e - \frac{cf}{d}\right) - i \sin\left(e - \frac{cf}{d}\right) \right) \right)}{2ad(-i + \tan(e+fx))}$$

input `Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])),x]`

output `(Sec[e + f*x]*((-I)*Cos[f*(c/d + x)] + Sin[f*(c/d + x)])*(CosIntegral[(2*f*(c + d*x))/d]*(Cos[e - (c*f)/d] - I*Sin[e - (c*f)/d]) + Log[f*(c + d*x)]*(Cos[e - (c*f)/d] + I*Sin[e - (c*f)/d]) + ((-I)*Cos[e - (c*f)/d] - Sin[e - (c*f)/d])*SinIntegral[(2*f*(c + d*x))/d))/(2*a*d*(-I + Tan[e + f*x]))`

### 3.21.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {3042, 4209, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx \\
 & \quad \downarrow \text{4209} \\
 & -\frac{i \int \frac{\sin(2e+2fx)}{c+dx} dx}{2a} + \frac{\int \frac{\cos(2e+2fx)}{c+dx} dx}{2a} + \frac{\log(c+dx)}{2ad} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{i \int \frac{\sin(2e+2fx)}{c+dx} dx}{2a} + \frac{\int \frac{\sin(2e+2fx+\frac{\pi}{2})}{c+dx} dx}{2a} + \frac{\log(c+dx)}{2ad} \\
 & \quad \downarrow \text{3784} \\
 & \frac{i \left( \sin \left( 2e - \frac{2cf}{d} \right) \int \frac{\cos \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx + \cos \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx \right)}{2a} + \\
 & \frac{\cos \left( 2e - \frac{2cf}{d} \right) \int \frac{\cos \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx - \sin \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx}{2a} + \frac{\log(c+dx)}{2ad} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.21.  $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx$

$$\begin{aligned}
& \frac{\cos\left(2e - \frac{2cf}{d}\right) \int \frac{\sin\left(2xf + \frac{2cf}{d} + \frac{\pi}{2}\right)}{c+dx} dx - \sin\left(2e - \frac{2cf}{d}\right) \int \frac{\sin\left(2xf + \frac{2cf}{d}\right)}{c+dx} dx}{2a} - \\
& \frac{i\left(\sin\left(2e - \frac{2cf}{d}\right) \int \frac{\sin\left(2xf + \frac{2cf}{d} + \frac{\pi}{2}\right)}{c+dx} dx + \cos\left(2e - \frac{2cf}{d}\right) \int \frac{\sin\left(2xf + \frac{2cf}{d}\right)}{c+dx} dx\right)}{2a} + \frac{\log(c+dx)}{2ad} \\
& \quad \downarrow \text{3780} \\
& \frac{\cos\left(2e - \frac{2cf}{d}\right) \int \frac{\sin\left(2xf + \frac{2cf}{d} + \frac{\pi}{2}\right)}{c+dx} dx - \frac{\sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{d}}{2a} - \\
& \frac{i\left(\sin\left(2e - \frac{2cf}{d}\right) \int \frac{\sin\left(2xf + \frac{2cf}{d} + \frac{\pi}{2}\right)}{c+dx} dx + \frac{\cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{d}\right)}{2a} + \frac{\log(c+dx)}{2ad} \\
& \quad \downarrow \text{3783} \\
& \frac{i\left(\frac{\text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{d} + \frac{\cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{d}\right)}{2a} + \\
& \frac{\frac{\text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{d} - \frac{\sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{d}}{2a} + \frac{\log(c+dx)}{2ad}
\end{aligned}$$

input `Int[1/((c + d*x)*(a + I*a*Tan[e + f*x])),x]`

output `Log[c + d*x]/(2*a*d) - ((I/2)*((CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/d + (Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/d)/a + ((Cos[2*e - (2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x])/d - (Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/d)/(2*a)`

### 3.21.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3780 `Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 4209 `Int[1/(((c_.) + (d_.)*(x_))*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[Log[c + d*x]/(2*a*d), x] + (Simp[1/(2*a) Int[Cos[2*e + 2*f*x]/(c + d*x), x], x] + Simp[1/(2*b) Int[Sin[2*e + 2*f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0]`

### 3.21.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.40

method	result
risch	$\frac{\ln(dx+c)}{2ad} - \frac{e^{\frac{2i(cf-de)}{d}} \operatorname{Ei}_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{2ad}$
default	$-\frac{if\left(\frac{2\operatorname{Si}\left(2fx+2e+\frac{2cf-2de}{d}\right)\cos\left(\frac{2cf-2de}{d}\right)-2\operatorname{Ci}\left(2fx+2e+\frac{2cf-2de}{d}\right)\sin\left(\frac{2cf-2de}{d}\right)}{4}\right)}{af} + \frac{f\left(\frac{2\operatorname{Si}\left(2fx+2e+\frac{2cf-2de}{d}\right)\sin\left(\frac{2cf-2de}{d}\right)+2\operatorname{Ci}\left(2fx+2e+\frac{2cf-2de}{d}\right)\cos\left(\frac{2cf-2de}{d}\right)}{4}\right)}{af}$

input `int(1/(d*x+c)/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/2*ln(d*x+c)/a/d-1/2/a/d*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)`

---

3.21.  $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx$

**3.21.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.32

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx = \frac{\operatorname{Ei}\left(-\frac{2(idfx+icf)}{d}\right) e^{\left(-\frac{2(de-icf)}{d}\right)} + \log\left(\frac{dx+c}{d}\right)}{2ad}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`output `1/2*(Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + log((d*x + c)/d))/a`**3.21.6 Sympy [F]**

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx = -\frac{i \int \frac{1}{c \tan(e+fx) - ic + dx \tan(e+fx) - idx} dx}{a}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x)`output `-I*Integral(1/(c*tan(e + f*x) - I*c + d*x*tan(e + f*x) - I*d*x), x)/a`**3.21.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.70

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx = \frac{f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) + i f E_1\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) - f \log\left(\frac{dx+c}{d}\right)}{2adf}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`output `-1/2*(f*cos(-2*(d*e - c*f)/d)*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + I*f*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) - f*log((f*x + e)*d - d*e + c*f)/(a*d*f)`

---

3.21.  $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx$

**3.21.8 Giac [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.86

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx$$

$$= \frac{\cos\left(\frac{2cf}{d}\right) \text{Ci}\left(-\frac{2(df x+cf)}{d}\right) + \cos(2e) \log(dx+c) + i \log(dx+c) \sin(2e) + i \text{Ci}\left(-\frac{2(df x+cf)}{d}\right) \sin\left(\frac{2cf}{d}\right) - 2(ad \cos(2e) + i ad \sin(2e))}{2(ad \cos(2e) + i ad \sin(2e))}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="giac")`output `1/2*(cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + cos(2*e)*log(d*x + c) + I*log(d*x + c)*sin(2*e) + I*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - I*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d))/(a*d*cos(2*e) + I*a*d*sin(2*e))`**3.21.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx = \int \frac{1}{(a+a \tan(e+fx) \text{li})(c+dx)} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)),x)`output `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)), x)`

### 3.22 $\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$

3.22.1 Optimal result . . . . .	167
3.22.2 Mathematica [A] (verified) . . . . .	168
3.22.3 Rubi [A] (verified) . . . . .	168
3.22.4 Maple [A] (verified) . . . . .	171
3.22.5 Fricas [A] (verification not implemented) . . . . .	171
3.22.6 Sympy [F] . . . . .	172
3.22.7 Maxima [A] (verification not implemented) . . . . .	172
3.22.8 Giac [B] (verification not implemented) . . . . .	172
3.22.9 Mupad [F(-1)] . . . . .	173

#### 3.22.1 Optimal result

Integrand size = 23, antiderivative size = 168

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx = -\frac{if \cos(2e - \frac{2cf}{d}) \text{CosIntegral}(\frac{2cf}{d} + 2fx)}{ad^2} - \frac{f \text{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{ad^2} - \frac{f \cos(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{ad^2} + \frac{if \sin(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{ad^2} - \frac{1}{d(c+dx)(a+ia \tan(e+fx))}$$

output

```
-I*f*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a/d^2-f*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a/d^2+f*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^2-I*f*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^2-1/d/(d*x+c)/(a+I*a*tan(f*x+e))
```



### 3.22.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.33

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$$

$$= \frac{\sec(e+fx) \left( \cos\left(\frac{cf}{d}\right) + i \sin\left(\frac{cf}{d}\right) \right) \left( d(i \cos(e+f(-\frac{c}{d}+x))) + i \cos(e+f(\frac{c}{d}+x)) - \sin(e+f(-\frac{c}{d}+x)) \right)}{(c+dx)^2(a+ia \tan(e+fx))}$$

input `Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])),x]`

output `(Sec[e + f*x]*(Cos[(c*f)/d] + I*Sin[(c*f)/d])*(d*(I*Cos[e + f*(-(c/d) + x)] + I*Cos[e + f*(c/d + x)] - Sin[e + f*(-(c/d) + x)] + Sin[e + f*(c/d + x)]) - 2*f*(c + d*x)*CosIntegral[(2*f*(c + d*x))/d]*(Cos[e - (f*(c + d*x))/d] - I*Sin[e - (f*(c + d*x))/d]) + 2*f*(c + d*x)*(I*Cos[e - (f*(c + d*x))/d] + Sin[e - (f*(c + d*x))/d])*SinIntegral[(2*f*(c + d*x))/d])/(2*a*d^2*(c + d*x)*(-I + Tan[e + f*x]))`

### 3.22.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {3042, 4207, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$$

$$\downarrow 3042$$

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$$

$$\downarrow 4207$$

$$-\frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{if \int \frac{\cos(2e+2fx)}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia \tan(e+fx))}$$

$$\downarrow 3042$$

---

3.22.  $\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$

$$\begin{aligned}
& \frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{if \int \frac{\sin(2e+2fx+\frac{\pi}{2})}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia \tan(e+fx))} \\
& \quad \downarrow \text{3784} \\
& \frac{f \left( \sin \left( 2e - \frac{2cf}{d} \right) \int \frac{\cos \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx + \cos \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx \right)}{ad} \\
& \frac{if \left( \cos \left( 2e - \frac{2cf}{d} \right) \int \frac{\cos \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx - \sin \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx \right)}{ad} \\
& \quad \downarrow \text{3042} \\
& \frac{if \left( \cos \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} + \frac{\pi}{2} \right)}{c+dx} dx - \sin \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx \right)}{ad} \\
& \frac{f \left( \sin \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} + \frac{\pi}{2} \right)}{c+dx} dx + \cos \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} \right)}{c+dx} dx \right)}{ad} \\
& \quad \downarrow \text{3780} \\
& \frac{if \left( \cos \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} + \frac{\pi}{2} \right)}{c+dx} dx - \frac{\sin \left( 2e - \frac{2cf}{d} \right) \text{Si} \left( 2xf + \frac{2cf}{d} \right)}{d} \right)}{ad} \\
& \frac{f \left( \sin \left( 2e - \frac{2cf}{d} \right) \int \frac{\sin \left( 2xf + \frac{2cf}{d} + \frac{\pi}{2} \right)}{c+dx} dx + \frac{\cos \left( 2e - \frac{2cf}{d} \right) \text{Si} \left( 2xf + \frac{2cf}{d} \right)}{d} \right)}{ad} - \frac{1}{d(c+dx)(a+ia \tan(e+fx))} \\
& \quad \downarrow \text{3783} \\
& \frac{f \left( \frac{\text{CosIntegral} \left( 2xf + \frac{2cf}{d} \right) \sin \left( 2e - \frac{2cf}{d} \right)}{d} + \frac{\cos \left( 2e - \frac{2cf}{d} \right) \text{Si} \left( 2xf + \frac{2cf}{d} \right)}{d} \right)}{ad} \\
& \frac{if \left( \frac{\text{CosIntegral} \left( 2xf + \frac{2cf}{d} \right) \cos \left( 2e - \frac{2cf}{d} \right)}{d} - \frac{\sin \left( 2e - \frac{2cf}{d} \right) \text{Si} \left( 2xf + \frac{2cf}{d} \right)}{d} \right)}{ad} - \frac{1}{d(c+dx)(a+ia \tan(e+fx))}
\end{aligned}$$

input `Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])),x]`

$$3.22. \quad \int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$$

```
output -((f*((CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/d + (Cos[2*e -
(2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/d))/(a*d) - (I*f*((Cos[2*e -
(2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x])/d - (Sin[2*e - (2*c*f)/d]*SinIn
tegral[(2*c*f)/d + 2*f*x])/d))/(a*d) - 1/(d*(c + d*x)*(a + I*a*Tan[e + f*x
]))
```

### 3.22.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3780 Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinInte
gral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

```
rule 3783 Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosInte
gral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) -
c*f, 0]
```

```
rule 3784 Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[Cos[(d*
e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*
f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x]
&& NeQ[d*e - c*f, 0]
```

```
rule 4207 Int[1/(((c_.) + (d_.)*(x_))^2*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])), x_Sy
mbol] :> -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))^(-1), x] + (-Simp[f/(a*d)
Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Simp[f/(b*d) Int[Cos[2*e + 2*
f*x]/(c + d*x), x], x)) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0
]
```

### 3.22.4 Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.57

method	result
risch	$-\frac{1}{2d(dx+c)a} - \frac{f e^{-2i(fx+e)}}{2a(dfx+cf)d} + \frac{i f e^{\frac{2i(cf-de)}{d}} \operatorname{Ei}_1\left(2i fx+2ie+\frac{2i(cf-de)}{d}\right)}{a d^2}$
default	$-\frac{i f^2 \left( -\frac{2 \sin(2fx+2e)}{(cf-de+d(fx+e))d} + \frac{4 \operatorname{Si}\left(2fx+2e+\frac{2cf-2de}{d}\right) \sin\left(\frac{2cf-2de}{d}\right)}{d} + \frac{4 \operatorname{Ci}\left(2fx+2e+\frac{2cf-2de}{d}\right) \cos\left(\frac{2cf-2de}{d}\right)}{d} \right)}{4} + \frac{f^2 \left( -\frac{2 \cos(2fx+2e)}{(cf-de+d(fx+e))} \right)}{af}$

input `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `-1/2/d/(d*x+c)/a-1/2/a*f*exp(-2*I*(f*x+e))/(d*f*x+c*f)/d+I/a*f/d^2*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)`

### 3.22.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.49

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$$

$$= -\frac{\left( \left( 2(i d f x + i c f) \operatorname{Ei}\left(-\frac{2(i d f x + i c f)}{d}\right) e^{\left(-\frac{2(i d e - i c f)}{d}\right)} + d \right) e^{(2i f x + 2i e)} + d \right) e^{(-2i f x - 2i e)}}{2(ad^3x + acd^2)}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output `-1/2*((2*(I*d*f*x + I*c*f)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + d)*e^(2*I*f*x + 2*I*e) + d)*e^(-2*I*f*x - 2*I*e)/(a*d^3*x + a*c*d^2)`

### 3.22.6 Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$$

$$= -\frac{i \int \frac{1}{c^2 \tan(e+fx) - ic^2 + 2cdx \tan(e+fx) - 2icdx + d^2x^2 \tan(e+fx) - id^2x^2} dx}{a}$$

input `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e)),x)`

output `-I*Integral(1/(c**2*tan(e + f*x) - I*c**2 + 2*c*d*x*tan(e + f*x) - 2*I*c*d*x + d**2*x**2*tan(e + f*x) - I*d**2*x**2), x)/a`

### 3.22.7 Maxima [A] (verification not implemented)

Time = 0.58 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.71

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx =$$

$$\frac{f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) + i f^2 E_2\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) + f^2}{2((fx+e)ad^2 - ad^2e + acdf)f}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `-1/2*(f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + f^2)/(((f*x + e)*a*d^2 - a*d^2*e + a*c*d*f)*f)`

### 3.22.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1013 vs.  $2(161) = 322$ .

Time = 2.30 (sec) , antiderivative size = 1013, normalized size of antiderivative = 6.03

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx = \text{Too large to display}$$

---

3.22.  $\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `1/2*(-2*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*I*d*e*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 2*d*e*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 2*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*d*e*f^2*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*c*f^3*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*I*d*e*f^2*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*c*f^3*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x...`

### 3.22.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx = \int \frac{1}{(a+a \tan(e+fx) 1i)(c+dx)^2} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^2),x)`

output `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^2), x)`

### 3.23 $\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$

3.23.1	Optimal result . . . . .	174
3.23.2	Mathematica [A] (verified) . . . . .	175
3.23.3	Rubi [A] (verified) . . . . .	175
3.23.4	Maple [A] (verified) . . . . .	178
3.23.5	Fricas [A] (verification not implemented) . . . . .	179
3.23.6	Sympy [F] . . . . .	179
3.23.7	Maxima [A] (verification not implemented) . . . . .	179
3.23.8	Giac [B] (verification not implemented) . . . . .	180
3.23.9	Mupad [F(-1)] . . . . .	181

#### 3.23.1 Optimal result

Integrand size = 23, antiderivative size = 227

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx = -\frac{if}{2ad^2(c+dx)} - \frac{f^2 \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{ad^3} + \frac{if^2 \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{ad^3} + \frac{if^2 \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^3} + \frac{f^2 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^3} - \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))} + \frac{if}{d^2(c+dx)(a+ia \tan(e+fx))}$$

output

```
-1/2*I*f/a/d^2/(d*x+c)-f^2*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a/d^3+I*f^2*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a/d^3-I*f^2*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^3-f^2*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^3-1/2/d/(d*x+c)^2/(a+I*a*tan(f*x+e))+I*f/d^2/(d*x+c)/(a+I*a*tan(f*x+e))
```

### 3.23.2 Mathematica [A] (verified)

Time = 1.00 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.26

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$$


---


$$= \frac{\sec(e+fx) \left( \cos\left(\frac{cf}{d}\right) + i \sin\left(\frac{cf}{d}\right) \right) \left( d(i d \cos(e+f(-\frac{c}{d}+x))) + (id+2cf+2dfx) \cos(e+f(\frac{c}{d}+x)) \right) - \dots}{\dots}$$

input `Integrate[1/((c + d*x)^3*(a + I*a*Tan[e + f*x])),x]`

output `(Sec[e + f*x]*(Cos[(c*f)/d] + I*Sin[(c*f)/d])*(d*(I*d*Cos[e + f*(-(c/d) + x)] + (I*d + 2*c*f + 2*d*f*x)*Cos[e + f*(c/d + x)] - d*Sin[e + f*(-(c/d) + x)] + d*Sin[e + f*(c/d + x)] - (2*I)*c*f*Sin[e + f*(c/d + x)] - (2*I)*d*f*x*Sin[e + f*(c/d + x)]) + 4*f^2*(c + d*x)^2*CosIntegral[(2*f*(c + d*x))/d]*(I*Cos[e - (f*(c + d*x))/d] + Sin[e - (f*(c + d*x))/d]) + 4*f^2*(c + d*x)^2*(Cos[e - (f*(c + d*x))/d] - I*Sin[e - (f*(c + d*x))/d])*SinIntegral[(2*f*(c + d*x))/d]))/(4*a*d^3*(c + d*x)^2*(-I + Tan[e + f*x]))`

### 3.23.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$ , Rules used = {3042, 4208, 3042, 4207, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$$

↓ 3042

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$$

↓ 4208

$$-\frac{if \int \frac{1}{(c+dx)^2(i \tan(e+fx)a+a)} dx}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))}$$

↓ 3042

---

3.23.  $\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$



$$\begin{aligned}
 & \frac{if \int \frac{1}{(c+dx)^2(i \tan(e+fx)a+a)} dx}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))} \\
 & \quad \downarrow 4207 \\
 & \frac{if \left( -\frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{if \int \frac{\cos(2e+2fx)}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia \tan(e+fx))} \right)}{d} - \frac{if}{2ad^2(c+dx)} - \\
 & \quad \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))} \\
 & \quad \downarrow 3042 \\
 & \frac{if \left( -\frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{if \int \frac{\sin(2e+2fx+\frac{\pi}{2})}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia \tan(e+fx))} \right)}{d} - \frac{if}{2ad^2(c+dx)} - \\
 & \quad \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))} \\
 & \quad \downarrow 3784 \\
 & \frac{if \left( -\frac{f \left( \sin(2e-\frac{2cf}{d}) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx + \cos(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} - \frac{if \left( \cos(2e-\frac{2cf}{d}) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx - \sin(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} \right)}{d} \\
 & \quad \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))} \\
 & \quad \downarrow 3042 \\
 & \frac{if \left( -\frac{if \left( \cos(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx - \sin(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} - \frac{f \left( \sin(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx + \cos(2e-\frac{2cf}{d}) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} \right)}{d} \\
 & \quad \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))} \\
 & \quad \downarrow 3780 \\
 & \frac{if \left( -\frac{if \left( \cos(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx - \frac{\sin(2e-\frac{2cf}{d}) \text{Si}(2xf+\frac{2cf}{d})}{d} \right)}{ad} - \frac{f \left( \sin(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx + \frac{\cos(2e-\frac{2cf}{d}) \text{Si}(2xf+\frac{2cf}{d})}{d} \right)}{ad} \right)}{d} \\
 & \quad \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))}
 \end{aligned}$$

3.23.  $\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$

$$\int \frac{if \left( \frac{\text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right) + \cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{ad} \right) - if \left( \frac{\text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right) - \sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{ad} \right)}{2ad^2(c+dx) - \frac{1}{2d(c+dx)^2(a+ia \tan(e+fx))}} dx$$

input `Int[1/((c + d*x)^3*(a + I*a*Tan[e + f*x])),x]`

output `((-1/2*I)*f)/(a*d^2*(c + d*x)) - 1/(2*d*(c + d*x)^2*(a + I*a*Tan[e + f*x]) - (I*f*(-((f*((CosIntegral[(2*c*f)/d + 2*f*x])*Sin[2*e - (2*c*f)/d])/d + (Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/d))/(a*d)) - (I*f*((Cos[2*e - (2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x])/d - (Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/d))/(a*d) - 1/(d*(c + d*x)*(a + I*a*Tan[e + f*x])))/d`

### 3.23.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3780 `Int[sin[(e_) + (f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 `Int[sin[(e_) + (f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

rule 3784 `Int[sin[(e_) + (f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

```
rule 4207 Int[1/(((c_.) + (d_.)*(x_.))^2*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol]
:> -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))^(-1), x] + (-Simp[f/(a*d)
  Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Simp[f/(b*d) Int[Cos[2*e + 2*
  f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0]
]
```

```
rule 4208 Int[(((c_.) + (d_.)*(x_.))^m)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol]
:> Simp[f*((c + d*x)^(m + 2)/(b*d^2*(m + 1)*(m + 2))), x] + (Simp[2*b*(
  f/(a*d*(m + 1))) Int[(c + d*x)^(m + 1)/(a + b*Tan[e + f*x]), x], x] + Si
  mp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + b*Tan[e + f*x])), x]) /; FreeQ[{a, b,
  c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[m, -1] && NeQ[m, -2]
```

### 3.23.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.95

method	result
risch	$-\frac{1}{4ad(dx+c)^2} + \frac{if^3e^{-2i(fx+e)}x}{2ad(d^2x^2f^2+2cdf^2x+c^2f^2)} - \frac{f^2e^{-2i(fx+e)}}{4ad(d^2x^2f^2+2cdf^2x+c^2f^2)} + \frac{if^3e^{-2i(fx+e)}c}{2ad^2(d^2x^2f^2+2cdf^2x+c^2f^2)} + \frac{f^2e^{\frac{2i(cf-d)}{d}}}{2ad^2(d^2x^2f^2+2cdf^2x+c^2f^2)}$ $if^3 \left( -\frac{\sin(2fx+2e)}{(cf-de+d(fx+e))^2d} + \frac{2\cos(2fx+2e)}{(cf-de+d(fx+e))d} - \frac{2\left( \frac{2\operatorname{Si}\left(2fx+2e+\frac{2cf-2de}{d}\right)\cos\left(\frac{2cf-2de}{d}\right)}{d} - \frac{2\operatorname{Ci}\left(2fx+2e+\frac{2cf-2de}{d}\right)\sin\left(\frac{2cf-2de}{d}\right)}{d} \right)}{d} \right)$
default	4

```
input int(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output -1/4/a/d/(d*x+c)^2+1/2*I/a*f^3*exp(-2*I*(f*x+e))/d/(d^2*f^2*x^2+2*c*d*f^2*
x+c^2*f^2)*x-1/4/a*f^2*exp(-2*I*(f*x+e))/d/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^
2)+1/2*I/a*f^3*exp(-2*I*(f*x+e))/d^2/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2)*c+1
/a*f^2/d^3*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)
```

$$3.23. \int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$$

### 3.23.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.57

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx = \frac{\left(2i d^2 f x + 2i c d f - d^2 - \left(4(d^2 f^2 x^2 + 2 c d f^2 x + c^2 f^2) \operatorname{Ei}\left(-\frac{2(i d f x + i c f)}{d}\right) e^{\left(-\frac{2(i d e - i c f)}{d}\right)} + d^2\right) e^{(2i f x + 2i e)}\right)}{4(ad^5 x^2 + 2acd^4 x + ac^2 d^3)}$$

input `integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output `1/4*(2*I*d^2*f*x + 2*I*c*d*f - d^2 - (4*(d^2*f^2*x^2 + 2*c*d*f^2*x + c^2*f^2)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + d^2)*e^(2*I*f*x + 2*I*e)*e^(-2*I*f*x - 2*I*e)/(a*d^5*x^2 + 2*a*c*d^4*x + a*c^2*d^3)`

### 3.23.6 Sympy [F]

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx = -\frac{i \int \frac{1}{c^3 \tan(e+fx) - ic^3 + 3c^2 dx \tan(e+fx) - 3ic^2 dx + 3cd^2 x^2 \tan(e+fx) - 3icd^2 x^2 + d^3 x^3 \tan(e+fx) - id^3 x^3} dx}{a}$$

input `integrate(1/(d*x+c)**3/(a+I*a*tan(f*x+e)),x)`

output `-I*Integral(1/(c**3*tan(e + f*x) - I*c**3 + 3*c**2*d*x*tan(e + f*x) - 3*I*c**2*d*x + 3*c*d**2*x**2*tan(e + f*x) - 3*I*c*d**2*x**2 + d**3*x**3*tan(e + f*x) - I*d**3*x**3), x)/a`

### 3.23.7 Maxima [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.70

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx = \frac{2 f^3 \cos\left(-\frac{2(de-cf)}{d}\right) E_3\left(-\frac{2(-i(fx+e)d+i de-icf)}{d}\right) + 2i f^3 E_3\left(-\frac{2(-i(fx+e)d+i de-icf)}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) + f^3}{4((fx+e)^2 ad^3 + ad^3 e^2 - 2acd^2 ef + ac^2 df^2 - 2(ad^3 e - acd^2 f)(fx+e)) f}$$

---

3.23.  $\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$

input `integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `-1/4*(2*f^3*cos(-2*(d*e - c*f)/d)*exp_integral_e(3, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f^3*exp_integral_e(3, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + f^3/(((f*x + e)^2*a*d^3 + a*d^3*e^2 - 2*a*c*d^2*e*f + a*c^2*d*f^2 - 2*(a*d^3*e - a*c*d^2*f)*(f*x + e))*f)`

### 3.23.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 532 vs.  $2(212) = 424$ .

Time = 0.44 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.34

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx = \frac{4d^2 f^2 x^2 \cos\left(\frac{2cf}{d}\right) \text{Ci}\left(-\frac{2(dfx+cf)}{d}\right) + 4i d^2 f^2 x^2 \text{Ci}\left(-\frac{2(dfx+cf)}{d}\right) \sin\left(\frac{2cf}{d}\right) - 4i d^2 f^2 x^2 \cos\left(\frac{2cf}{d}\right) \text{Si}\left(\frac{2(dfx+cf)}{d}\right)}{\dots}$$

input `integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `-1/4*(4*d^2*f^2*x^2*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + 4*I*d^2*f^2*x^2*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 4*I*d^2*f^2*x^2*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 4*d^2*f^2*x^2*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 8*c*d*f^2*x*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + 8*I*c*d*f^2*x*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 8*I*c*d*f^2*x*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 8*c*d*f^2*x*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 4*c^2*f^2*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + 4*I*c^2*f^2*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 4*I*c^2*f^2*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 4*c^2*f^2*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) - 2*I*d^2*f*x*cos(2*f*x) - 2*d^2*f*x*sin(2*f*x) - 2*I*c*d*f*cos(2*f*x) - 2*c*d*f*sin(2*f*x) + d^2*cos(2*f*x) + d^2*cos(2*e) - I*d^2*sin(2*f*x) + I*d^2*sin(2*e))/(a*d^5*x^2*cos(2*e) + I*a*d^5*x^2*sin(2*e) + 2*a*c*d^4*x*cos(2*e) + 2*I*a*c*d^4*x*sin(2*e) + a*c^2*d^3*cos(2*e) + I*a*c^2*d^3*sin(2*e))`

**3.23.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx = \int \frac{1}{(a+a \tan(e+fx) \text{ li}) (c+dx)^3} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^3),x)`output `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^3), x)`

### 3.24 $\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^2} dx$

3.24.1	Optimal result	182
3.24.2	Mathematica [A] (verified)	183
3.24.3	Rubi [A] (verified)	183
3.24.4	Maple [A] (verified)	185
3.24.5	Fricas [A] (verification not implemented)	185
3.24.6	Sympy [A] (verification not implemented)	186
3.24.7	Maxima [F(-2)]	186
3.24.8	Giac [A] (verification not implemented)	187
3.24.9	Mupad [B] (verification not implemented)	187

#### 3.24.1 Optimal result

Integrand size = 23, antiderivative size = 270

$$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^2} dx = -\frac{3d^3 e^{-2ie-2ifx}}{16a^2 f^4} - \frac{3d^3 e^{-4ie-4ifx}}{512a^2 f^4} - \frac{3id^2 e^{-2ie-2ifx}(c+dx)}{8a^2 f^3}$$

$$- \frac{3id^2 e^{-4ie-4ifx}(c+dx)}{128a^2 f^3} + \frac{3de^{-2ie-2ifx}(c+dx)^2}{8a^2 f^2}$$

$$+ \frac{3de^{-4ie-4ifx}(c+dx)^2}{64a^2 f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^3}{4a^2 f}$$

$$+ \frac{ie^{-4ie-4ifx}(c+dx)^3}{16a^2 f} + \frac{(c+dx)^4}{16a^2 d}$$

output `-3/16*d^3*exp(-2*I*e-2*I*f*x)/a^2/f^4-3/512*d^3*exp(-4*I*e-4*I*f*x)/a^2/f^4-3/8*I*d^2*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^2/f^3-3/128*I*d^2*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^2/f^3+3/8*d*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^2/f^2+3/64*d*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^2/f^2+1/4*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^3/a^2/f+1/16*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^3/a^2/f+1/16*(d*x+c)^4/a^2/d`

### 3.24.2 Mathematica [A] (verified)

Time = 2.54 (sec) , antiderivative size = 473, normalized size of antiderivative = 1.75

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{\sec^2(e + fx)(\cos(fx) + i \sin(fx))^2 ((4ic^3 f^3 + 6c^2 df^2(1 + 2ifx) + 6cd^2 f(-i + 2fx + 2if^2 x^2) + d^3(-3 -$$

input `Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x])^2,x]`

output `(Sec[e + f*x]^2*(Cos[f*x] + I*Sin[f*x])^2*(((4*I)*c^3*f^3 + 6*c^2*d*f^2*(1 + (2*I)*f*x) + 6*c*d^2*f*(-I + 2*f*x + (2*I)*f^2*x^2) + d^3*(-3 - (6*I)*f*x + 6*f^2*x^2 + (4*I)*f^3*x^3))*Cos[2*f*x] + (((32*I)*c^3*f^3 + 24*c^2*d*f^2*(1 + (4*I)*f*x) + 12*c*d^2*f*(-I + 4*f*x + (8*I)*f^2*x^2) + d^3*(-3 - (12*I)*f*x + 24*f^2*x^2 + (32*I)*f^3*x^3))*Cos[4*f*x]*(Cos[2*e] - I*Sin[2*e]))/32 + f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(Cos[2*e] + I*Sin[2*e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(-I + 2*f*x) + 6*c*d^2*f*(-1 - (2*I)*f*x + 2*f^2*x^2) + d^3*(3*I - 6*f*x - (6*I)*f^2*x^2 + 4*f^3*x^3))*Sin[2*f*x] + ((32*c^3*f^3 + 24*c^2*d*f^2*(-I + 4*f*x) + 12*c*d^2*f*(-1 - (4*I)*f*x + 8*f^2*x^2) + d^3*(3*I - 12*f*x - (24*I)*f^2*x^2 + 32*f^3*x^3))*(Cos[2*e] - I*Sin[2*e])*Sin[4*f*x])/32))/(16*f^4*(a + I*a*Tan[e + f*x])^2)`

### 3.24.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$\downarrow \text{4212}$$



$$\int \left( \frac{(c+dx)^3 e^{-2ie-2ifx}}{2a^2} + \frac{(c+dx)^3 e^{-4ie-4ifx}}{4a^2} + \frac{(c+dx)^3}{4a^2} \right) dx$$

↓ 2009

$$-\frac{3id^2(c+dx)e^{-2ie-2ifx}}{8a^2f^3} - \frac{3id^2(c+dx)e^{-4ie-4ifx}}{128a^2f^3} + \frac{3d(c+dx)^2e^{-2ie-2ifx}}{8a^2f^2} + \frac{3d(c+dx)^2e^{-4ie-4ifx}}{64a^2f^2} + \frac{i(c+dx)^3e^{-2ie-2ifx}}{4a^2f} + \frac{i(c+dx)^3e^{-4ie-4ifx}}{16a^2f} + \frac{(c+dx)^4}{16a^2d} - \frac{3d^3e^{-2ie-2ifx}}{16a^2f^4} - \frac{3d^3e^{-4ie-4ifx}}{512a^2f^4}$$

input `Int[(c + d*x)^3/(a + I*a*Tan[e + f*x])^2,x]`

output `(-3*d^3*E^((-2*I)*e - (2*I)*f*x))/(16*a^2*f^4) - (3*d^3*E^((-4*I)*e - (4*I)*f*x))/(512*a^2*f^4) - (((3*I)/8)*d^2*E^((-2*I)*e - (2*I)*f*x)*(c + d*x))/(a^2*f^3) - (((3*I)/128)*d^2*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))/(a^2*f^3) + (3*d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^2)/(8*a^2*f^2) + (3*d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^2)/(64*a^2*f^2) + ((I/4)*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^3)/(a^2*f) + ((I/16)*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^3)/(a^2*f) + (c + d*x)^4/(16*a^2*d)`

### 3.24.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4212 `Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]`

### 3.24.4 Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.05

method	result
risch	$\frac{d^3 x^4}{16a^2} + \frac{d^2 c x^3}{4a^2} + \frac{3d c^2 x^2}{8a^2} + \frac{c^3 x}{4a^2} + \frac{c^4}{16a^2 d} + \frac{i(4d^3 x^3 f^3 + 12c d^2 f^3 x^2 - 6i d^3 f^2 x^2 + 12c^2 d f^3 x - 12i c d^2 f^2 x + 4c^3 f^3 - 6i c^2 d f^2 - 6c^4)}{16a^2 f^4}$

input `int((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output  $\frac{1}{16/a^2 d^3 x^4 + 1/4/a^2 d^2 c x^3 + 3/8/a^2 d c^2 x^2 + 1/4/a^2 c^3 x + 1/16/a^2 d c^4 + 1/16 * I * (4 * d^3 x^3 f^3 - 6 * I * d^3 f^2 x^2 + 12 * c * d^2 f^3 x^2 - 12 * I * c * d^2 f^2 x + 12 * c^2 * d * f^3 x - 6 * I * c^2 * d * f^2 + 4 * c^3 * f^3 - 6 * d^3 * f * x + 3 * I * d^3 - 6 * c * d^2 * f) / a^2 / f^4 * \exp(-2 * I * (f * x + e)) + 1/512 * I * (32 * d^3 x^3 f^3 - 24 * I * d^3 f^2 x^2 + 96 * c * d^2 f^3 x^2 - 48 * I * c * d^2 f^2 x + 96 * c^2 * d * f^3 x - 24 * I * c^2 * d * f^2 + 32 * c^3 * f^3 - 12 * d^3 * f * x + 3 * I * d^3 - 12 * c * d^2 * f) / a^2 / f^4 * \exp(-4 * I * (f * x + e))$

### 3.24.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.99

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(32i d^3 f^3 x^3 + 32i c^3 f^3 + 24 c^2 d f^2 - 12i c d^2 f - 3 d^3 - 24(-4i c d^2 f^3 - d^3 f^2)x^2 - 12(-8i c^2 d f^3 - 4 c d^2 f^2))}{(a + ia \tan(e + fx))^2}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="fracas")`

output  $\frac{1}{512} * (32 * I * d^3 * f^3 * x^3 + 32 * I * c^3 * f^3 + 24 * c^2 * d * f^2 - 12 * I * c * d^2 * f - 3 * d^3 - 24 * (-4 * I * c * d^2 * f^3 - d^3 * f^2) * x^2 - 12 * (-8 * I * c^2 * d * f^3 - 4 * c * d^2 * f^2 + I * d^3 * f) * x + 32 * (d^3 * f^4 * x^4 + 4 * c * d^2 * f^4 * x^3 + 6 * c^2 * d * f^4 * x^2 + 4 * c^3 * f^4 * x) * e^{(4 * I * f * x + 4 * I * e)} - 32 * (-4 * I * d^3 * f^3 * x^3 - 4 * I * c^3 * f^3 - 6 * c^2 * d * f^2 + 6 * I * c * d^2 * f + 3 * d^3 + 6 * (-2 * I * c * d^2 * f^3 - d^3 * f^2) * x^2 + 6 * (-2 * I * c^2 * d * f^3 - 2 * c * d^2 * f^2 + I * d^3 * f) * x) * e^{(2 * I * f * x + 2 * I * e)}) * e^{(-4 * I * f * x - 4 * I * e)} / (a^2 * f^4)$

### 3.24.6 Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 665, normalized size of antiderivative = 2.46

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$= \left\{ \frac{((512ia^2c^3f^7e^{2ie} + 1536ia^2c^2df^7xe^{2ie} + 384a^2c^2df^6e^{2ie} + 1536ia^2cd^2f^7x^2e^{2ie} + 768a^2cd^2f^6xe^{2ie} - 192ia^2cd^2f^5e^{2ie} + 512ia^2d^3f^7x^3e^{2ie} + 384a^2d^3f^6xe^{2ie} - 192ia^2d^3f^5xe^{2ie} + 512ia^2d^3f^4e^{2ie} + 384a^2d^3f^3e^{2ie} + 192ia^2d^3f^2e^{2ie} + 192ia^2d^3fe^{2ie} + 192ia^2d^3e^{2ie}))e^{-4ie}}{16a^2} + \frac{x^3 \cdot (2cd^2e^{2ie} + cd^2)e^{-4ie}}{4a^2} + \frac{x^2 \cdot (6c^2de^{2ie} + 3c^2d)e^{-4ie}}{8a^2} + \frac{x(2c^3e^{2ie} + c^3)e^{-4ie}}{4a^2} \right.$$

$$\left. + \frac{c^3x}{4a^2} + \frac{3c^2dx^2}{8a^2} + \frac{cd^2x^3}{4a^2} + \frac{d^3x^4}{16a^2} \right.$$

input `integrate((d*x+c)**3/(a+I*a*tan(f*x+e))**2,x)`

output `Piecewise((((512*I*a**2*c**3*f**7*exp(2*I*e) + 1536*I*a**2*c**2*d*f**7*x*exp(2*I*e) + 384*a**2*c**2*d*f**6*exp(2*I*e) + 1536*I*a**2*c*d**2*f**7*x**2*exp(2*I*e) + 768*a**2*c*d**2*f**6*x*exp(2*I*e) - 192*I*a**2*c*d**2*f**5*exp(2*I*e) + 512*I*a**2*d**3*f**7*x**3*exp(2*I*e) + 384*a**2*d**3*f**6*x**2*exp(2*I*e) - 192*I*a**2*d**3*f**5*x*exp(2*I*e) - 48*a**2*d**3*f**4*exp(2*I*e))*exp(-4*I*f*x) + (2048*I*a**2*c**3*f**7*exp(4*I*e) + 6144*I*a**2*c**2*d*f**7*x*exp(4*I*e) + 3072*a**2*c**2*d*f**6*exp(4*I*e) + 6144*I*a**2*c*d**2*f**7*x**2*exp(4*I*e) + 6144*a**2*c*d**2*f**6*x*exp(4*I*e) - 3072*I*a**2*c*d**2*f**5*exp(4*I*e) + 2048*I*a**2*d**3*f**7*x**3*exp(4*I*e) + 3072*a**2*d**3*f**6*x**2*exp(4*I*e) - 3072*I*a**2*d**3*f**5*x*exp(4*I*e) - 1536*a**2*d**3*f**4*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(8192*a**4*f**8), Ne(a**4*f**8*exp(6*I*e), 0)), (x**4*(2*d**3*exp(2*I*e) + d**3)*exp(-4*I*e)/(16*a**2) + x**3*(2*c*d**2*exp(2*I*e) + c*d**2)*exp(-4*I*e)/(4*a**2) + x**2*(6*c**2*d*exp(2*I*e) + 3*c**2*d)*exp(-4*I*e)/(8*a**2) + x*(2*c**3*exp(2*I*e) + c**3)*exp(-4*I*e)/(4*a**2), True)) + c**3*x/(4*a**2) + 3*c**2*d*x**2/(8*a**2) + c*d**2*x**3/(4*a**2) + d**3*x**4/(16*a**2)`

### 3.24.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

### 3.24.8 Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.36

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(32 d^3 f^4 x^4 e^{(4i fx + 4i e)} + 128 cd^2 f^4 x^3 e^{(4i fx + 4i e)} + 192 c^2 d f^4 x^2 e^{(4i fx + 4i e)} + 128i d^3 f^3 x^3 e^{(2i fx + 2i e)} + 32i d^3 f^3 x^3 e^{(2i fx + 2i e)} + \dots)}{(a^2 f^4 + \dots)}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `1/512*(32*d^3*f^4*x^4*e^(4*I*f*x + 4*I*e) + 128*c*d^2*f^4*x^3*e^(4*I*f*x + 4*I*e) + 192*c^2*d*f^4*x^2*e^(4*I*f*x + 4*I*e) + 128*I*d^3*f^3*x^3*e^(2*I*f*x + 2*I*e) + 32*I*d^3*f^3*x^3 + 128*c^3*f^4*x*e^(4*I*f*x + 4*I*e) + 384*I*c*d^2*f^3*x^2*e^(2*I*f*x + 2*I*e) + 96*I*c*d^2*f^3*x^2 + 384*I*c^2*d*f^3*x*e^(2*I*f*x + 2*I*e) + 192*d^3*f^2*x^2*e^(2*I*f*x + 2*I*e) + 96*I*c^2*d*f^3*x + 24*d^3*f^2*x^2 + 128*I*c^3*f^3*e^(2*I*f*x + 2*I*e) + 384*c*d^2*f^2*x*e^(2*I*f*x + 2*I*e) + 32*I*c^3*f^3 + 48*c*d^2*f^2*x + 192*c^2*d*f^2*e^(2*I*f*x + 2*I*e) - 192*I*d^3*f*x*e^(2*I*f*x + 2*I*e) + 24*c^2*d*f^2 - 12*I*d^3*f*x - 192*I*c*d^2*f*e^(2*I*f*x + 2*I*e) - 12*I*c*d^2*f - 96*d^3*e^(2*I*f*x + 2*I*e) - 3*d^3)*e^(-4*I*f*x - 4*I*e)/(a^2*f^4)`

### 3.24.9 Mupad [B] (verification not implemented)

Time = 4.30 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx = e^{-e 2i - f x 2i} \left( \frac{(4c^3 f^3 - c^2 d f^2 6i - 6cd^2 f + d^3 3i) 1i}{16a^2 f^4} + \frac{d^3 x^3 1i}{4a^2 f} - \frac{dx(-2c^2 f^2 + cdf 2i + d^2) 3i}{8a^2 f^3} - \frac{d^2 x^2(-2cf + d 1i) 3i}{8a^2 f^2} \right)$$

$$+ e^{-e 4i - f x 4i} \left( \frac{(32c^3 f^3 - c^2 d f^2 24i - 12cd^2 f + d^3 3i) 1i}{512a^2 f^4} + \frac{d^3 x^3 1i}{16a^2 f} - \frac{dx(-8c^2 f^2 + cdf 4i + d^2) 3i}{128a^2 f^3} - \frac{d^2 x^2(-4cf + d 1i) 3i}{64a^2 f^2} \right) + \frac{c^3 x}{4a^2} + \frac{d^3 x^4}{16a^2} + \frac{3c^2 dx^2}{8a^2} + \frac{cd^2 x^3}{4a^2}$$

3.24.  $\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^2} dx$

input `int((c + d*x)^3/(a + a*tan(e + f*x)*1i)^2,x)`

output `exp(- e*2i - f*x*2i)*(((d^3*3i + 4*c^3*f^3 - c^2*d*f^2*6i - 6*c*d^2*f)*1i) / (16*a^2*f^4) + (d^3*x^3*1i)/(4*a^2*f) - (d*x*(d^2 - 2*c^2*f^2 + c*d*f*2i) *3i)/(8*a^2*f^3) - (d^2*x^2*(d*1i - 2*c*f)*3i)/(8*a^2*f^2)) + exp(- e*4i - f*x*4i)*(((d^3*3i + 32*c^3*f^3 - c^2*d*f^2*24i - 12*c*d^2*f)*1i)/(512*a^2 *f^4) + (d^3*x^3*1i)/(16*a^2*f) - (d*x*(d^2 - 8*c^2*f^2 + c*d*f*4i)*3i)/(1 28*a^2*f^3) - (d^2*x^2*(d*1i - 4*c*f)*3i)/(64*a^2*f^2)) + (c^3*x)/(4*a^2) + (d^3*x^4)/(16*a^2) + (3*c^2*d*x^2)/(8*a^2) + (c*d^2*x^3)/(4*a^2)`

### 3.25 $\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^2} dx$

3.25.1	Optimal result . . . . .	189
3.25.2	Mathematica [A] (verified) . . . . .	189
3.25.3	Rubi [A] (verified) . . . . .	190
3.25.4	Maple [A] (verified) . . . . .	191
3.25.5	Fricas [A] (verification not implemented) . . . . .	192
3.25.6	Sympy [A] (verification not implemented) . . . . .	192
3.25.7	Maxima [F(-2)] . . . . .	193
3.25.8	Giac [A] (verification not implemented) . . . . .	193
3.25.9	Mupad [B] (verification not implemented) . . . . .	194

#### 3.25.1 Optimal result

Integrand size = 23, antiderivative size = 202

$$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^2} dx = -\frac{id^2e^{-2ie-2ifx}}{8a^2f^3} - \frac{id^2e^{-4ie-4ifx}}{128a^2f^3} + \frac{de^{-2ie-2ifx}(c+dx)}{4a^2f^2}$$

$$+ \frac{de^{-4ie-4ifx}(c+dx)}{32a^2f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^2}{4a^2f}$$

$$+ \frac{ie^{-4ie-4ifx}(c+dx)^2}{16a^2f} + \frac{(c+dx)^3}{12a^2d}$$

output

```
-1/8*I*d^2*exp(-2*I*e-2*I*f*x)/a^2/f^3-1/128*I*d^2*exp(-4*I*e-4*I*f*x)/a^2/f^3+1/4*d*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^2/f^2+1/32*d*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^2/f^2+1/4*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^2/f+1/16*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^2/f+1/12*(d*x+c)^3/a^2/d
```

#### 3.25.2 Mathematica [A] (verified)

Time = 1.61 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.40

$$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^2} dx$$

$$= \frac{\sec^2(e+fx)(\cos(fx)+i \sin(fx))^2((d+(1+i)cf+(1+i)dfx)((1+i)cf+d(-i+(1+i)fx)) \cos(2fx))}{(a+ia \tan(e+fx))^2}$$

input `Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x])^2,x]`

output `(Sec[e + f*x]^2*(Cos[f*x] + I*Sin[f*x])^2*((d + (1 + I)*c*f + (1 + I)*d*f*x)*((1 + I)*c*f + d*(-I + (1 + I)*f*x))*Cos[2*f*x] + ((d + (2 + 2*I)*c*f + (2 + 2*I)*d*f*x)*((2 + 2*I)*c*f + d*(-I + (2 + 2*I)*f*x))*Cos[4*f*x]*(Cos[2*e] - I*Sin[2*e]))/16 + (2*f^3*x*(3*c^2 + 3*c*d*x + d^2*x^2)*(Cos[2*e] + I*Sin[2*e]))/3 - I*(d + (1 + I)*c*f + (1 + I)*d*f*x)*((1 + I)*c*f + d*(-I + (1 + I)*f*x))*Sin[2*f*x] - (I/16)*(d + (2 + 2*I)*c*f + (2 + 2*I)*d*f*x)*((2 + 2*I)*c*f + d*(-I + (2 + 2*I)*f*x))*(Cos[2*e] - I*Sin[2*e])*Sin[4*f*x]]/(8*f^3*(a + I*a*Tan[e + f*x])^2)`

### 3.25.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4212} \\
 & \int \left( \frac{(c + dx)^2 e^{-2ie - 2ifx}}{2a^2} + \frac{(c + dx)^2 e^{-4ie - 4ifx}}{4a^2} + \frac{(c + dx)^2}{4a^2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{d(c + dx)e^{-2ie - 2ifx}}{4a^2 f^2} + \frac{d(c + dx)e^{-4ie - 4ifx}}{32a^2 f^2} + \frac{i(c + dx)^2 e^{-2ie - 2ifx}}{4a^2 f} + \frac{i(c + dx)^2 e^{-4ie - 4ifx}}{16a^2 f} + \\
 & \quad \frac{(c + dx)^3}{12a^2 d} - \frac{id^2 e^{-2ie - 2ifx}}{8a^2 f^3} - \frac{4a^2 f}{id^2 e^{-4ie - 4ifx}} - \frac{16a^2 f}{128a^2 f^3} +
 \end{aligned}$$

input `Int[(c + d*x)^2/(a + I*a*Tan[e + f*x])^2,x]`

```
output ((-1/8*I)*d^2*E^((-2*I)*e - (2*I)*f*x)/(a^2*f^3) - ((I/128)*d^2*E^((-4*I)
*e - (4*I)*f*x)/(a^2*f^3) + (d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x))/(4*a^2
*f^2) + (d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))/(32*a^2*f^2) + ((I/4)*E^((-
2*I)*e - (2*I)*f*x)*(c + d*x)^2)/(a^2*f) + ((I/16)*E^((-4*I)*e - (4*I)*f*x
)*(c + d*x)^2)/(a^2*f) + (c + d*x)^3/(12*a^2*d)
```

### 3.25.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4212 Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*
x)))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2
, 0] && ILtQ[n, 0]
```

### 3.25.4 Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.86

method	result
risch	$\frac{d^2 x^3}{12a^2} + \frac{dcx^2}{4a^2} + \frac{c^2x}{4a^2} + \frac{c^3}{12a^2d} + \frac{i(2d^2x^2f^2+4cdf^2x-2id^2fx+2c^2f^2-2icdf-d^2)e^{-2i(fx+e)}}{8a^2f^3} + \frac{i(8d^2x^2f^2+16cdf^2x-4id^2f^2x)}{12a^2d}$

```
input int((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 1/12/a^2*d^2*x^3+1/4/a^2*d*c*x^2+1/4/a^2*c^2*x+1/12/a^2/d*c^3+1/8*I*(2*d^2
*x^2*f^2-2*I*d^2*f*x+4*c*d*f^2*x-2*I*c*d*f+2*c^2*f^2-d^2)/a^2/f^3*exp(-2*I
*(f*x+e))+1/128*I*(8*d^2*x^2*f^2-4*I*d^2*f*x+16*c*d*f^2*x-4*I*c*d*f+8*c^2*
f^2-d^2)/a^2/f^3*exp(-4*I*(f*x+e))
```



### 3.25.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.81

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(24i d^2 f^2 x^2 + 24i c^2 f^2 + 12 cdf - 3i d^2 - 12(-4i cdf^2 - d^2 f)x + 32(d^2 f^3 x^3 + 3cdf^3 x^2 + 3c^2 f^3 x)e^{4ifx} + 384 a^2 f^3)}{384 a^2 f^3}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="fracas")`

output `1/384*(24*I*d^2*f^2*x^2 + 24*I*c^2*f^2 + 12*c*d*f - 3*I*d^2 - 12*(-4*I*c*d*f^2 - d^2*f)*x + 32*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 3*c^2*f^3*x)*e^(4*I*f*x + 4*I*e) - 48*(-2*I*d^2*f^2*x^2 - 2*I*c^2*f^2 - 2*c*d*f + I*d^2 + 2*(-2*I*c*d*f^2 - d^2*f)*x)*e^(2*I*f*x + 2*I*e))*e^(-4*I*f*x - 4*I*e)/(a^2*f^3)`

### 3.25.6 Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 418, normalized size of antiderivative = 2.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx$$

$$= \left\{ \frac{((64ia^2 c^2 f^5 e^{2ie} + 128ia^2 cdf^5 x e^{2ie} + 32a^2 cdf^4 e^{2ie} + 64ia^2 d^2 f^5 x^2 e^{2ie} + 32a^2 d^2 f^4 x e^{2ie} - 8ia^2 d^2 f^3 e^{2ie})e^{-4ifx} + (256ia^2 c^2 f^5 e^{4ie} + 512ia^2 cdf^5 x e^{4ie})}{1024a^4 f^6} \right.$$

$$\left. + \frac{c^2 x}{4a^2} + \frac{cdx^2}{4a^2} + \frac{d^2 x^3}{12a^2} \right.$$

input `integrate((d*x+c)**2/(a+I*a*tan(f*x+e))**2,x)`

output `Piecewise((((64*I*a**2*c**2*f**5*exp(2*I*e) + 128*I*a**2*c*d*f**5*x*exp(2*I*e) + 32*a**2*c*d*f**4*exp(2*I*e) + 64*I*a**2*d**2*f**5*x**2*exp(2*I*e) + 32*a**2*d**2*f**4*x*exp(2*I*e) - 8*I*a**2*d**2*f**3*exp(2*I*e))*exp(-4*I*f*x) + (256*I*a**2*c**2*f**5*exp(4*I*e) + 512*I*a**2*c*d*f**5*x*exp(4*I*e) + 256*a**2*c*d*f**4*exp(4*I*e) + 256*I*a**2*d**2*f**5*x**2*exp(4*I*e) + 256*a**2*d**2*f**4*x*exp(4*I*e) - 128*I*a**2*d**2*f**3*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(1024*a**4*f**6), Ne(a**4*f**6*exp(6*I*e), 0)), (x**3*(2*d**2*exp(2*I*e) + d**2)*exp(-4*I*e)/(12*a**2) + x**2*(2*c*d*exp(2*I*e) + c*d)*exp(-4*I*e)/(4*a**2) + x*(2*c**2*exp(2*I*e) + c**2)*exp(-4*I*e)/(4*a**2), True)) + c**2*x/(4*a**2) + c*d*x**2/(4*a**2) + d**2*x**3/(12*a**2)`

---

3.25.  $\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^2} dx$

### 3.25.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

### 3.25.8 Giac [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(32 d^2 f^3 x^3 e^{(4i f x + 4i e)} + 96 c d f^3 x^2 e^{(4i f x + 4i e)} + 96 c^2 f^3 x e^{(4i f x + 4i e)} + 96 i d^2 f^2 x^2 e^{(2i f x + 2i e)} + 24 i d^2 f^2 x^2 + 1}{(a + ia \tan(e + fx))^2}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `1/384*(32*d^2*f^3*x^3*e^(4*I*f*x + 4*I*e) + 96*c*d*f^3*x^2*e^(4*I*f*x + 4*I*e) + 96*c^2*f^3*x*e^(4*I*f*x + 4*I*e) + 96*I*d^2*f^2*x^2*e^(2*I*f*x + 2*I*e) + 24*I*d^2*f^2*x^2 + 192*I*c*d*f^2*x*e^(2*I*f*x + 2*I*e) + 48*I*c*d*f^2*x + 96*I*c^2*f^2*e^(2*I*f*x + 2*I*e) + 96*d^2*f*x*e^(2*I*f*x + 2*I*e) + 24*I*c^2*f^2 + 12*d^2*f*x + 96*c*d*f*e^(2*I*f*x + 2*I*e) + 12*c*d*f - 48*I*d^2*e^(2*I*f*x + 2*I*e) - 3*I*d^2)*e^(-4*I*f*x - 4*I*e)/(a^2*f^3)`

**3.25.9 Mupad [B] (verification not implemented)**

Time = 3.65 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.91

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx = \frac{c^2 x}{4 a^2} - e^{-e 4i - f x 4i} \left( \frac{(-8 c^2 f^2 + c d f 4i + d^2) li}{128 a^2 f^3} - \frac{d^2 x^2 li}{16 a^2 f} + \frac{d x (-4 c f + d li) li}{32 a^2 f^2} \right) - e^{-e 2i - f x 2i} \left( \frac{(-2 c^2 f^2 + c d f 2i + d^2) li}{8 a^2 f^3} - \frac{d^2 x^2 li}{4 a^2 f} + \frac{d x (-2 c f + d li) li}{4 a^2 f^2} \right) + \frac{d^2 x^3}{12 a^2} + \frac{c d x^2}{4 a^2}$$

input `int((c + d*x)^2/(a + a*tan(e + f*x)*1i)^2,x)`output `(c^2*x)/(4*a^2) - exp(- e*4i - f*x*4i)*(((d^2 - 8*c^2*f^2 + c*d*f*4i)*1i)/(128*a^2*f^3) - (d^2*x^2*1i)/(16*a^2*f) + (d*x*(d*1i - 4*c*f)*1i)/(32*a^2*f^2)) - exp(- e*2i - f*x*2i)*(((d^2 - 2*c^2*f^2 + c*d*f*2i)*1i)/(8*a^2*f^3) - (d^2*x^2*1i)/(4*a^2*f) + (d*x*(d*1i - 2*c*f)*1i)/(4*a^2*f^2)) + (d^2*x^3)/(12*a^2) + (c*d*x^2)/(4*a^2)`

### 3.26 $\int \frac{c+dx}{(a+ia \tan(e+fx))^2} dx$

3.26.1	Optimal result . . . . .	195
3.26.2	Mathematica [A] (verified) . . . . .	195
3.26.3	Rubi [A] (verified) . . . . .	196
3.26.4	Maple [A] (verified) . . . . .	197
3.26.5	Fricas [A] (verification not implemented) . . . . .	197
3.26.6	Sympy [A] (verification not implemented) . . . . .	198
3.26.7	Maxima [F(-2)] . . . . .	198
3.26.8	Giac [A] (verification not implemented) . . . . .	199
3.26.9	Mupad [B] (verification not implemented) . . . . .	199

#### 3.26.1 Optimal result

Integrand size = 21, antiderivative size = 151

$$\int \frac{c+dx}{(a+ia \tan(e+fx))^2} dx = -\frac{3idx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2}$$

$$+ \frac{d}{16f^2(a+ia \tan(e+fx))^2} + \frac{i(c+dx)}{4f(a+ia \tan(e+fx))^2}$$

$$+ \frac{3d}{16f^2(a^2+ia^2 \tan(e+fx))} + \frac{i(c+dx)}{4f(a^2+ia^2 \tan(e+fx))}$$

output `-3/16*I*d*x/a^2/f-1/8*d*x^2/a^2+1/4*x*(d*x+c)/a^2+1/16*d/f^2/(a+I*a*tan(f*x+e))^2+1/4*I*(d*x+c)/f/(a+I*a*tan(f*x+e))^2+3/16*d/f^2/(a^2+I*a^2*tan(f*x+e))+1/4*I*(d*x+c)/f/(a^2+I*a^2*tan(f*x+e))`

#### 3.26.2 Mathematica [A] (verified)

Time = 1.07 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.86

$$\int \frac{c+dx}{(a+ia \tan(e+fx))^2} dx = \frac{\sec^2(e+fx)(8(d+2icf+2idfx)+(4cf(i+4fx)+d(1+4ifx+8f^2x^2))\cos(2(e+fx)))+(4cf(1+64a^2f^2(-i+\tan(e+fx))^2$$

input `Integrate[(c+d*x)/(a+I*a*Tan[e+f*x])^2,x]`

output `-1/64*(Sec[e + f*x]^2*(8*(d + (2*I)*c*f + (2*I)*d*f*x) + (4*c*f*(I + 4*f*x) + d*(1 + (4*I)*f*x + 8*f^2*x^2))*Cos[2*(e + f*x)] + (4*c*f*(1 + (4*I)*f*x) + d*(-I + 4*f*x + (8*I)*f^2*x^2))*Sin[2*(e + f*x)]))/(a^2*f^2*(-I + Tan[e + f*x])^2)`

### 3.26.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {3042, 4213, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx$$

↓ 4213

$$-d \int \left( \frac{x}{4a^2} + \frac{i}{4f(i \tan(e + fx)a^2 + a^2)} + \frac{i}{4f(i \tan(e + fx)a + a)^2} \right) dx +$$

$$\frac{i(c + dx)}{4f(a^2 + ia^2 \tan(e + fx))} + \frac{x(c + dx)}{4a^2} + \frac{i(c + dx)}{4f(a + ia \tan(e + fx))^2}$$

↓ 2009

$$d \left( -\frac{3}{16f^2(a^2 + ia^2 \tan(e + fx))} + \frac{3ix}{16a^2f} + \frac{x^2}{8a^2} - \frac{1}{16f^2(a + ia \tan(e + fx))^2} \right) +$$

$$\frac{i(c + dx)}{4f(a + ia \tan(e + fx))^2}$$

input `Int[(c + d*x)/(a + I*a*Tan[e + f*x])^2, x]`

output `(x*(c + d*x))/(4*a^2) + ((I/4)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])^2) + ((I/4)*(c + d*x))/(f*(a^2 + I*a^2*Tan[e + f*x])) - d*(((3*I)/16)*x)/(a^2*f) + x^2/(8*a^2) - 1/(16*f^2*(a + I*a*Tan[e + f*x])^2) - 3/(16*f^2*(a^2 + I*a^2*Tan[e + f*x]))`

---

3.26.  $\int \frac{c+dx}{(a+ia \tan(e+fx))^2} dx$

### 3.26.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4213 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := With[{u = IntHide[(a + b*Tan[e + f*x])^n, x]}, Simp[(c + d*x)^m u, x] - Simp[d*m Int[(c + d*x)^(m - 1) u, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, -1] && GtQ[m, 0]`

### 3.26.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.54

method	result	size
risch	$\frac{dx^2}{8a^2} + \frac{cx}{4a^2} + \frac{i(2dfx+2cf-id)e^{-2i(fx+e)}}{8a^2f^2} + \frac{i(4dfx+4cf-id)e^{-4i(fx+e)}}{64a^2f^2}$	82

input `int((d*x+c)/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `1/8*d*x^2/a^2+1/4/a^2*c*x+1/8*I*(2*d*f*x-I*d+2*c*f)/a^2/f^2*exp(-2*I*(f*x+e))+1/64*I*(4*d*f*x-I*d+4*c*f)/a^2/f^2*exp(-4*I*(f*x+e))`

### 3.26.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.52

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(4i dfx + 4i cf + 8(df^2x^2 + 2cf^2x))e^{(4i fx + 4i e)} - 8(-2i dfx - 2i cf - d)e^{(2i fx + 2i e)} + d)e^{(-4i fx - 4i e)}}{64 a^2 f^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output  $\frac{1}{64} \cdot (4 \cdot I \cdot d \cdot f \cdot x + 4 \cdot I \cdot c \cdot f + 8 \cdot (d \cdot f^2 \cdot x^2 + 2 \cdot c \cdot f^2 \cdot x)) \cdot e^{(4 \cdot I \cdot f \cdot x + 4 \cdot I \cdot e)} - 8 \cdot (-2 \cdot I \cdot d \cdot f \cdot x - 2 \cdot I \cdot c \cdot f - d) \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + d \cdot e^{(-4 \cdot I \cdot f \cdot x - 4 \cdot I \cdot e)} / (a^2 \cdot f^2)$

### 3.26.6 Sympy [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.50

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{((32ia^2cf^3e^{2ie} + 32ia^2df^3xe^{2ie} + 8a^2df^2e^{2ie})e^{-4ifx} + (128ia^2cf^3e^{4ie} + 128ia^2df^3xe^{4ie} + 64a^2df^2e^{4ie})e^{-2ifx})e^{-6ie}}{512a^4f^4} & \text{for } a^4f^4e^{6ie} \neq 0 \\ \frac{x^2 \cdot (2de^{2ie} + d)e^{-4ie}}{8a^2} + \frac{x(2ce^{2ie} + c)e^{-4ie}}{4a^2} & \text{otherwise} \\ + \frac{cx}{4a^2} + \frac{dx^2}{8a^2} \end{cases}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))**2,x)`

output `Piecewise((((32*I*a**2*c*f**3*exp(2*I*e) + 32*I*a**2*d*f**3*x*exp(2*I*e) + 8*a**2*d*f**2*exp(2*I*e))*exp(-4*I*f*x) + (128*I*a**2*c*f**3*exp(4*I*e) + 128*I*a**2*d*f**3*x*exp(4*I*e) + 64*a**2*d*f**2*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(512*a**4*f**4), Ne(a**4*f**4*exp(6*I*e), 0)), (x**2*(2*d*exp(2*I*e) + d)*exp(-4*I*e)/(8*a**2) + x*(2*c*exp(2*I*e) + c)*exp(-4*I*e)/(4*a**2), True)) + c*x/(4*a**2) + d*x**2/(8*a**2)`

### 3.26.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.26.8 Giac [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.67

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(8df^2x^2e^{(4ifx+4ie)} + 16cf^2xe^{(4ifx+4ie)} + 16idfxe^{(2ifx+2ie)} + 4idfx + 16icfe^{(2ifx+2ie)} + 4icf + 8de^{(2ifx+2ie)})}{64a^2f^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`output `1/64*(8*d*f^2*x^2*e^(4*I*f*x + 4*I*e) + 16*c*f^2*x*e^(4*I*f*x + 4*I*e) + 16*I*d*f*x*e^(2*I*f*x + 2*I*e) + 4*I*d*f*x + 16*I*c*f*e^(2*I*f*x + 2*I*e) + 4*I*c*f + 8*d*e^(2*I*f*x + 2*I*e) + d)*e^(-4*I*f*x - 4*I*e)/(a^2*f^2)`**3.26.9 Mupad [B] (verification not implemented)**

Time = 3.13 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.68

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx = \frac{dx^2}{8a^2} - e^{-e4i-fx4i} \left( \frac{(-4cf + dli) li}{64a^2f^2} - \frac{dx li}{16a^2f} \right)$$

$$- e^{-e2i-fx2i} \left( \frac{(-2cf + dli) li}{8a^2f^2} - \frac{dx li}{4a^2f} \right) + \frac{cx}{4a^2}$$

input `int((c + d*x)/(a + a*tan(e + f*x)*1i)^2,x)`output `(d*x^2)/(8*a^2) - exp(- e*4i - f*x*4i)*(((d*1i - 4*c*f)*1i)/(64*a^2*f^2) - (d*x*1i)/(16*a^2*f)) - exp(- e*2i - f*x*2i)*(((d*1i - 2*c*f)*1i)/(8*a^2*f^2) - (d*x*1i)/(4*a^2*f)) + (c*x)/(4*a^2)`



### 3.27 $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$

3.27.1	Optimal result	200
3.27.2	Mathematica [A] (verified)	201
3.27.3	Rubi [A] (verified)	201
3.27.4	Maple [A] (verified)	203
3.27.5	Fricas [A] (verification not implemented)	203
3.27.6	Sympy [F]	204
3.27.7	Maxima [A] (verification not implemented)	204
3.27.8	Giac [A] (verification not implemented)	205
3.27.9	Mupad [F(-1)]	205

#### 3.27.1 Optimal result

Integrand size = 23, antiderivative size = 305

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx = \frac{\cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{2a^2d} + \frac{\cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{4a^2d} + \frac{\log(c+dx)}{4a^2d} - \frac{i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{4a^2d} - \frac{i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{2a^2d} - \frac{i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{\sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{4a^2d} - \frac{\sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{4a^2d}$$

output  $\frac{1}{4} \text{Ci}(4cf/d+4fx) \cos(-4e+4cf/d)/a^{2/d} + \frac{1}{2} \text{Ci}(2cf/d+2fx) \cos(-2e+2cf/d)/a^{2/d} + \frac{1}{4} \ln(dx+c)/a^{2/d} - \frac{1}{2} I \cos(-2e+2cf/d) \text{Si}(2cf/d+2fx)/a^{2/d} - \frac{1}{4} I \cos(-4e+4cf/d) \text{Si}(4cf/d+4fx)/a^{2/d} + \frac{1}{4} I \text{Ci}(4cf/d+4fx) \sin(-4e+4cf/d)/a^{2/d} + \frac{1}{4} \text{Si}(4cf/d+4fx) \sin(-4e+4cf/d)/a^{2/d} + \frac{1}{2} I \text{Ci}(2cf/d+2fx) \sin(-2e+2cf/d)/a^{2/d} + \frac{1}{2} \text{Si}(2cf/d+2fx) \sin(-2e+2cf/d)/a^{2/d}$

### 3.27.2 Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.69

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$$

$$= \frac{(\cos(2e - \frac{2cf}{d}) - i \sin(2e - \frac{2cf}{d})) \left( 2 \text{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) + \cos(2e - \frac{2cf}{d}) \log(f(c+dx)) + \text{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) \right)}{(c+dx)(a+ia \tan(e+fx))^2}$$

input `Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])^2),x]`

output  $((\text{Cos}[2e - (2cf)/d] - I \text{Sin}[2e - (2cf)/d]) * (2 \text{CosIntegral}[(2f*(c + d*x))/d] + \text{Cos}[2e - (2cf)/d] * \text{Log}[f*(c + d*x)] + \text{CosIntegral}[(4f*(c + d*x))/d] * (\text{Cos}[2e - (2cf)/d] - I \text{Sin}[2e - (2cf)/d]) + I * \text{Log}[f*(c + d*x)]) * \text{Sin}[2e - (2cf)/d] - (2I) * \text{SinIntegral}[(2f*(c + d*x))/d] - I * \text{Cos}[2e - (2cf)/d] * \text{SinIntegral}[(4f*(c + d*x))/d] - \text{Sin}[2e - (2cf)/d] * \text{SinIntegral}[(4f*(c + d*x))/d])) / (4*a^2*d)$

### 3.27.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$$

↓ 3042

---

3.27.  $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$$

↓ 4211

$$\int \left( -\frac{\sin^2(2e+2fx)}{4a^2(c+dx)} - \frac{i \sin(2e+2fx)}{2a^2(c+dx)} - \frac{i \sin(4e+4fx)}{4a^2(c+dx)} + \frac{\cos^2(2e+2fx)}{4a^2(c+dx)} + \frac{\cos(2e+2fx)}{2a^2(c+dx)} + \frac{1}{4a^2(c+dx)} \right) dx$$

↓ 2009

$$\begin{aligned} & -\frac{i \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{2a^2d} - \frac{i \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{4a^2d} + \\ & \frac{\operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{2a^2d} + \frac{\operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{4a^2d} - \\ & \frac{\sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{2a^2d} - \frac{\sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{4a^2d} - \\ & \frac{i \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{2a^2d} - \frac{i \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{4a^2d} + \frac{\log(c+dx)}{4a^2d} \end{aligned}$$

input `Int[1/((c + d*x)*(a + I*a*Tan[e + f*x])^2),x]`

output `(Cos[2*e - (2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x])/(2*a^2*d) + (Cos[4*e - (4*c*f)/d]*CosIntegral[(4*c*f)/d + 4*f*x])/(4*a^2*d) + Log[c + d*x]/(4*a^2*d) - ((I/4)*CosIntegral[(4*c*f)/d + 4*f*x]*Sin[4*e - (4*c*f)/d])/(a^2*d) - ((I/2)*CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/(a^2*d) - ((I/2)*Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(a^2*d) - (Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(2*a^2*d) - ((I/4)*Cos[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(a^2*d) - (Sin[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(4*a^2*d)`

### 3.27.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4211 Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(
2*a) + Sin[2*e + 2*f*x]/(2*b))^(n), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

### 3.27.4 Maple [A] (verified)

Time = 0.87 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.37

method	result	size
risch	$\frac{\ln(dx+c)}{4a^2d} - \frac{e^{\frac{4i(cf-de)}{d}} \operatorname{Ei}_1\left(4ifx+4ie+\frac{4i(cf-de)}{d}\right)}{4a^2d} - \frac{e^{\frac{2i(cf-de)}{d}} \operatorname{Ei}_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{2a^2d}$	114

```
input int(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 1/4*ln(d*x+c)/a^2/d-1/4/a^2/d*exp(4*I*(c*f-d*e)/d)*Ei(1,4*I*f*x+4*I*e+4*I*
(c*f-d*e)/d)-1/2/a^2/d*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*
e)/d)
```

### 3.27.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.28

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$$

$$= \frac{2 \operatorname{Ei}\left(-\frac{2(i dfx+icf)}{d}\right) e^{\left(-\frac{2(i de-icf)}{d}\right)} + \operatorname{Ei}\left(-\frac{4(i dfx+icf)}{d}\right) e^{\left(-\frac{4(i de-icf)}{d}\right)} + \log\left(\frac{dx+c}{d}\right)}{4a^2d}$$

```
input integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="fracas")
```

```
output 1/4*(2*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + Ei(-4*(I*d*f*
x + I*c*f)/d)*e^(-4*(I*d*e - I*c*f)/d) + log((d*x + c)/d))/(a^2*d)
```

## 3.27.6 Sympy [F]

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$$

$$= -\frac{\int \frac{1}{c \tan^2(e+fx) - 2ic \tan(e+fx) - c + dx \tan^2(e+fx) - 2idx \tan(e+fx) - dx} dx}{a^2}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**2,x)`

output `-Integral(1/(c*tan(e + f*x)**2 - 2*I*c*tan(e + f*x) - c + d*x*tan(e + f*x)**2 - 2*I*d*x*tan(e + f*x) - d*x), x)/a**2`

## 3.27.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.64

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx =$$

$$\frac{2f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) + f \cos\left(-\frac{4(de-cf)}{d}\right) E_1\left(-\frac{4(-i(fx+e)d+ide-icf)}{d}\right) + 2if E_1\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right)}{a^2}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `-1/4*(2*f*cos(-2*(d*e - c*f)/d)*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f*cos(-4*(d*e - c*f)/d)*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + I*f*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) - f*log((f*x + e)*d - d*e + c*f))/(a^2*d*f)`

**3.27.8 Giac [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 404, normalized size of antiderivative = 1.32

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$$

$$= \frac{2 \cos(2e) \cos\left(\frac{2cf}{d}\right) \operatorname{Ci}\left(-\frac{2(dfx+cf)}{d}\right) + \cos(2e)^2 \log(dx+c) + 2i \cos\left(\frac{2cf}{d}\right) \operatorname{Ci}\left(-\frac{2(dfx+cf)}{d}\right) \sin(2e) + 2i \sin(2e) \log(dx+c)}{(a^2 d \cos(2e)^2 + 2I a^2 d \cos(2e) \sin(2e) - a^2 d \sin(2e)^2)}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

```
output 1/4*(2*cos(2*e)*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + cos(2*e)^2
*log(d*x + c) + 2*I*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e)
+ 2*I*cos(2*e)*log(d*x + c)*sin(2*e) - log(d*x + c)*sin(2*e)^2 + 2*I*cos(
2*e)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 2*cos_integral(-2*(d*
f*x + c*f)/d)*sin(2*e)*sin(2*c*f/d) - 2*I*cos(2*e)*cos(2*c*f/d)*sin_integr
al(2*(d*f*x + c*f)/d) + 2*cos(2*c*f/d)*sin(2*e)*sin_integral(2*(d*f*x + c*
f)/d) + 2*cos(2*e)*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 2*I*sin(
2*e)*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + cos(4*c*f/d)*cos_integ
ral(-4*(d*f*x + c*f)/d) + I*cos_integral(-4*(d*f*x + c*f)/d)*sin(4*c*f/d)
- I*cos(4*c*f/d)*sin_integral(4*(d*f*x + c*f)/d) + sin(4*c*f/d)*sin_integr
al(4*(d*f*x + c*f)/d))/(a^2*d*cos(2*e)^2 + 2*I*a^2*d*cos(2*e)*sin(2*e) - a
^2*d*sin(2*e)^2)
```

**3.27.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx = \int \frac{1}{(a+a \tan(e+fx) li)^2 (c+dx)} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)),x)`output `int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)), x)`

### 3.28 $\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx$

3.28.1 Optimal result . . . . .	206
3.28.2 Mathematica [A] (verified) . . . . .	207
3.28.3 Rubi [A] (verified) . . . . .	208
3.28.4 Maple [A] (verified) . . . . .	209
3.28.5 Fricas [A] (verification not implemented) . . . . .	210
3.28.6 Sympy [F] . . . . .	210
3.28.7 Maxima [A] (verification not implemented) . . . . .	211
3.28.8 Giac [B] (verification not implemented) . . . . .	211
3.28.9 Mupad [F(-1)] . . . . .	212

#### 3.28.1 Optimal result

Integrand size = 23, antiderivative size = 436

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx = -\frac{1}{4a^2d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2d(c+dx)}$$

$$- \frac{if \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{a^2d^2}$$

$$- \frac{if \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{a^2d^2}$$

$$- \frac{f \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{a^2d^2}$$

$$- \frac{f \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{a^2d^2}$$

$$+ \frac{i \sin(2e+2fx)}{2a^2d(c+dx)} + \frac{\sin^2(2e+2fx)}{4a^2d(c+dx)}$$

$$+ \frac{i \sin(4e+4fx)}{4a^2d(c+dx)} - \frac{f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2}$$

$$+ \frac{if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2}$$

$$- \frac{f \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2}$$

$$+ \frac{if \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2}$$

output 
$$-1/4/a^2/d/(d*x+c)-I*f*Ci(4*c*f/d+4*f*x)*cos(-4*e+4*c*f/d)/a^2/d^2-I*f*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a^2/d^2-1/2*cos(2*f*x+2*e)/a^2/d/(d*x+c)-1/4*cos(2*f*x+2*e)^2/a^2/d/(d*x+c)-f*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^2/d^2-f*cos(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^2/d^2+f*Ci(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a^2/d^2-I*f*Si(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a^2/d^2+f*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^2/d^2-I*f*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^2/d^2+1/2*I*sin(2*f*x+2*e)/a^2/d/(d*x+c)+1/4*sin(2*f*x+2*e)^2/a^2/d/(d*x+c)+1/4*I*sin(4*f*x+4*e)/a^2/d/(d*x+c)$$

### 3.28.2 Mathematica [A] (verified)

Time = 1.67 (sec) , antiderivative size = 467, normalized size of antiderivative = 1.07

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx = \frac{(\cos(2(e+f(-\frac{c}{d}+x))) - i \sin(2(e+f(-\frac{c}{d}+x)))) (2d \cos(\frac{2cf}{d}) + d \cos(2(e+f(-\frac{c}{d}+x)))) + d \cos(2(e+f(-\frac{c}{d}+x)))}{(c+dx)^2(a+ia \tan(e+fx))^2}$$

input `Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^2),x]`

output 
$$-1/4*((\text{Cos}[2*(e + f*(-(c/d) + x))] - I*\text{Sin}[2*(e + f*(-(c/d) + x))])*(2*d*\text{Cos}[(2*c*f)/d] + d*\text{Cos}[2*(e + f*(-(c/d) + x))] + d*\text{Cos}[2*(e + f*(c/d + x))] - (2*I)*d*\text{Sin}[(2*c*f)/d] + (4*I)*f*(c + d*x)*\text{CosIntegral}[(2*f*(c + d*x))/d]*(\text{Cos}[2*f*x] + I*\text{Sin}[2*f*x]) + I*d*\text{Sin}[2*(e + f*(-(c/d) + x))] - I*d*\text{Sin}[2*(e + f*(c/d + x))] + 4*f*(c + d*x)*\text{CosIntegral}[(4*f*(c + d*x))/d]*(I*\text{Cos}[2*e - (2*f*(c + d*x))/d] + \text{Sin}[2*e - (2*f*(c + d*x))/d] + 4*c*f*\text{Cos}[2*f*x]*\text{SinIntegral}[(2*f*(c + d*x))/d] + 4*d*f*x*\text{Cos}[2*f*x]*\text{SinIntegral}[(2*f*(c + d*x))/d] + (4*I)*c*f*\text{Sin}[2*f*x]*\text{SinIntegral}[(2*f*(c + d*x))/d] + (4*I)*d*f*x*\text{Sin}[2*f*x]*\text{SinIntegral}[(2*f*(c + d*x))/d] + 4*c*f*\text{Cos}[2*e - (2*f*(c + d*x))/d]*\text{SinIntegral}[(4*f*(c + d*x))/d] + 4*d*f*x*\text{Cos}[2*e - (2*f*(c + d*x))/d]*\text{SinIntegral}[(4*f*(c + d*x))/d] - (4*I)*c*f*\text{Sin}[2*e - (2*f*(c + d*x))/d]*\text{SinIntegral}[(4*f*(c + d*x))/d] - (4*I)*d*f*x*\text{Sin}[2*e - (2*f*(c + d*x))/d]*\text{SinIntegral}[(4*f*(c + d*x))/d]))/(a^2*d^2*(c + d*x))$$



### 3.28.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx \\
 & \quad \downarrow \text{4211} \\
 & \int \left( -\frac{\sin^2(2e+2fx)}{4a^2(c+dx)^2} - \frac{i \sin(2e+2fx)}{2a^2(c+dx)^2} - \frac{i \sin(4e+4fx)}{4a^2(c+dx)^2} + \frac{\cos^2(2e+2fx)}{4a^2(c+dx)^2} + \frac{\cos(2e+2fx)}{2a^2(c+dx)^2} + \frac{1}{4a^2(c+dx)^2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{f \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{a^2 d^2} - \frac{f \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{a^2 d^2} - \\
 & \frac{if \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{a^2 d^2} - \frac{if \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{a^2 d^2} + \\
 & \frac{if \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{a^2 d^2} + \frac{if \sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{a^2 d^2} - \\
 & \frac{f \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{a^2 d^2} - \frac{f \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{a^2 d^2} + \frac{\sin^2(2e+2fx)}{4a^2 d(c+dx)} + \\
 & \frac{i \sin(2e+2fx)}{2a^2 d(c+dx)} + \frac{i \sin(4e+4fx)}{4a^2 d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2 d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2 d(c+dx)} - \frac{1}{4a^2 d(c+dx)}
 \end{aligned}$$

input `Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^2),x]`

```
output -1/4*1/(a^2*d*(c + d*x)) - Cos[2*e + 2*f*x]/(2*a^2*d*(c + d*x)) - Cos[2*e
+ 2*f*x]^2/(4*a^2*d*(c + d*x)) - (I*f*Cos[2*e - (2*c*f)/d]*CosIntegral[(2*
c*f)/d + 2*f*x])/(a^2*d^2) - (I*f*Cos[4*e - (4*c*f)/d]*CosIntegral[(4*c*f)
/d + 4*f*x])/(a^2*d^2) - (f*CosIntegral[(4*c*f)/d + 4*f*x]*Sin[4*e - (4*c*
f)/d])/(a^2*d^2) - (f*CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])
/(a^2*d^2) + ((I/2)*Sin[2*e + 2*f*x])/(a^2*d*(c + d*x)) + Sin[2*e + 2*f*x]
^2/(4*a^2*d*(c + d*x)) + ((I/4)*Sin[4*e + 4*f*x])/(a^2*d*(c + d*x)) - (f*C
os[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(a^2*d^2) + (I*f*Sin[2
*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(a^2*d^2) - (f*Cos[4*e - (
4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(a^2*d^2) + (I*f*Sin[4*e - (4*c*
f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(a^2*d^2)
```

### 3.28.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4211 Int[((c_.) + (d_.)*(x_))^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(
2*a) + Sin[2*e + 2*f*x]/(2*b))^(n), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

### 3.28.4 Maple [A] (verified)

Time = 0.89 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.40

method	result
risch	$-\frac{1}{4a^2d(dx+c)} - \frac{f e^{-4i(fx+e)}}{4a^2(dfx+cf)d} + \frac{i f e^{\frac{4i(cf-de)}{d}} \operatorname{Ei}_1\left(\frac{4ifx+4ie+\frac{4i(cf-de)}{d}}{d}\right)}{a^2d^2} - \frac{f e^{-2i(fx+e)}}{2a^2(dfx+cf)d} + \frac{i f e^{\frac{2i(cf-de)}{d}} \operatorname{Ei}_1\left(\frac{2ifx+2ie+\frac{2i(cf-de)}{d}}{d}\right)}{a^2d^2}$

```
input int(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

---

3.28. 
$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx$$

output 
$$-1/4/a^2/d/(d*x+c)-1/4/a^2*f*\exp(-4*I*(f*x+e))/(d*f*x+c*f)/d+I/a^2*f/d^2*\exp(4*I*(c*f-d*e)/d)*\text{Ei}(1,4*I*f*x+4*I*e+4*I*(c*f-d*e)/d)-1/2/a^2*f*\exp(-2*I*(f*x+e))/(d*f*x+c*f)/d+I/a^2*f/d^2*\exp(2*I*(c*f-d*e)/d)*\text{Ei}(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)$$

### 3.28.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.33

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx = \frac{\left( \left( 4(i dfx + icf) \text{Ei} \left( -\frac{2(i dfx + icf)}{d} \right) e^{\left( -\frac{2(i de - icf)}{d} \right)} + 4(i dfx + icf) \text{Ei} \left( -\frac{4(i dfx + icf)}{d} \right) e^{\left( -\frac{4(i de - icf)}{d} \right)} + d \right) \right)}{4(a^2 d^3 x + a^2 c d^2)}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="fracas")`

output 
$$-1/4*((4*(I*d*f*x + I*c*f)*\text{Ei}(-2*(I*d*f*x + I*c*f)/d)*e^{(-2*(I*d*e - I*c*f)/d)} + 4*(I*d*f*x + I*c*f)*\text{Ei}(-4*(I*d*f*x + I*c*f)/d)*e^{(-4*(I*d*e - I*c*f)/d)} + d)*e^{4*I*f*x + 4*I*e} + 2*d*e^{(2*I*f*x + 2*I*e)} + d)*e^{(-4*I*f*x - 4*I*e)/(a^2*d^3*x + a^2*c*d^2)}$$

### 3.28.6 Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx = \frac{\int \frac{1}{c^2 \tan^2(e+fx) - 2ic^2 \tan(e+fx) - c^2 + 2cdx \tan^2(e+fx) - 4icdx \tan(e+fx) - 2cdx + d^2 x^2 \tan^2(e+fx) - 2id^2 x^2 \tan(e+fx) - d^2 x^2} dx}{a^2}$$

input `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e))**2,x)`

output 
$$-\text{Integral}(1/(c**2*\tan(e + f*x)**2 - 2*I*c**2*\tan(e + f*x) - c**2 + 2*c*d*x*\tan(e + f*x)**2 - 4*I*c*d*x*\tan(e + f*x) - 2*c*d*x + d**2*x**2*\tan(e + f*x)**2 - 2*I*d**2*x**2*\tan(e + f*x) - d**2*x**2), x)/a**2$$

---

3.28. 
$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx$$

**3.28.7 Maxima [A] (verification not implemented)**

Time = 0.48 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.48

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx = \frac{2f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) + f^2 \cos\left(-\frac{4(de-cf)}{d}\right) E_2\left(-\frac{4(-i(fx+e)d+ide-icf)}{d}\right) + 2if^2 \cos\left(-\frac{2(de-cf)}{d}\right) \operatorname{Ei}\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) + 2if^2 \cos\left(-\frac{4(de-cf)}{d}\right) \operatorname{Ei}\left(-\frac{4(-i(fx+e)d+ide-icf)}{d}\right)}{4((fx+e)a^2d^2 - a^2d^2)}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `-1/4*(2*f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f^2*cos(-4*(d*e - c*f)/d)*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + I*f^2*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + f^2)/(((f*x + e)^2*a^2*d^2 - a^2*d^2*e + a^2*c*d*f)*f)`

**3.28.8 Giac [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1967 vs. 2(410) = 820.

Time = 11.30 (sec) , antiderivative size = 1967, normalized size of antiderivative = 4.51

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx = \text{Too large to display}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `1/4*(-4*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 4*I*d*e*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 4*I*c*f^3*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 4*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 4*I*d*e*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 4*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 4*d*e*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 4*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-4*(d*e - c*f)/d) - 4*d*e*f^2*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-4*(d*e - c*f)/d) + 4*c*f^3*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) ...`

### 3.28.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^2 (c + dx)^2} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)^2),x)`

output `int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)^2), x)`

### 3.29 $\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx$

3.29.1	Optimal result . . . . .	213
3.29.2	Mathematica [A] (verified) . . . . .	214
3.29.3	Rubi [A] (verified) . . . . .	214
3.29.4	Maple [A] (verified) . . . . .	216
3.29.5	Fricas [A] (verification not implemented) . . . . .	216
3.29.6	Sympy [A] (verification not implemented) . . . . .	217
3.29.7	Maxima [F(-2)] . . . . .	218
3.29.8	Giac [A] (verification not implemented) . . . . .	219
3.29.9	Mupad [B] (verification not implemented) . . . . .	220

#### 3.29.1 Optimal result

Integrand size = 23, antiderivative size = 396

$$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx = -\frac{9d^3e^{-2ie-2ifx}}{64a^3f^4} - \frac{9d^3e^{-4ie-4ifx}}{1024a^3f^4} - \frac{d^3e^{-6ie-6ifx}}{1728a^3f^4} - \frac{9id^2e^{-2ie-2ifx}(c+dx)}{32a^3f^3} - \frac{9id^2e^{-4ie-4ifx}(c+dx)}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}(c+dx)}{288a^3f^3} + \frac{9de^{-2ie-2ifx}(c+dx)^2}{32a^3f^2} + \frac{9de^{-4ie-4ifx}(c+dx)^2}{128a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)^2}{96a^3f^2} + \frac{3ie^{-2ie-2ifx}(c+dx)^3}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^3}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^3}{48a^3f} + \frac{(c+dx)^4}{32a^3d}$$

output

```
-9/64*d^3*exp(-2*I*e-2*I*f*x)/a^3/f^4-9/1024*d^3*exp(-4*I*e-4*I*f*x)/a^3/f^4-1/1728*d^3*exp(-6*I*e-6*I*f*x)/a^3/f^4-9/32*I*d^2*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^3/f^3-9/256*I*d^2*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^3/f^3-1/288*I*d^2*exp(-6*I*e-6*I*f*x)*(d*x+c)/a^3/f^3+9/32*d*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^3/f^2+9/128*d*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^3/f^2+1/96*d*exp(-6*I*e-6*I*f*x)*(d*x+c)^2/a^3/f^2+3/16*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^3/a^3/f+3/32*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^3/a^3/f+1/48*I*exp(-6*I*e-6*I*f*x)*(d*x+c)^3/a^3/f+1/32*(d*x+c)^4/a^3/d
```

### 3.29.2 Mathematica [A] (verified)

Time = 3.76 (sec) , antiderivative size = 667, normalized size of antiderivative = 1.68

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{i \sec^3(e + fx) (243(32ic^3 f^3 + 8c^2 df^2(5 + 12ifx) + 4cd^2 f(-9i + 20fx + 24if^2 x^2) + d^3(-17 - 36ifx + 40f^2 x^2 + (32i)f^3 x^3)) \cos[e + fx] + 16(36c^3 f^3 (I + 6fx) + 18c^2 d f^2 (1 + (6I)fx + 18f^2 x^2) + 6c d^2 f (-I + 6fx + (18I)f^2 x^2 + 36f^3 x^3) + d^3 (-1 - (6I)fx + 18f^2 x^2 + (36I)f^3 x^3 + 54f^4 x^4)) \cos[3(e + fx)] - (3645I) d^3 \sin[e + fx] + 6804 c d^2 f f \sin[e + fx] + (5832I) c^2 d f^2 \sin[e + fx] - 2592 c^3 f^3 \sin[e + fx] + 6804 d^3 f x \sin[e + fx] + (11664I) c d^2 f^2 x \sin[e + fx] - 7776 c^2 d f^3 x \sin[e + fx] + (5832I) d^3 f^2 x^2 \sin[e + fx] - 7776 c d^2 f^3 x^2 \sin[e + fx] - 2592 d^3 f^3 x^3 \sin[e + fx] + (16I) d^3 \sin[3(e + fx)] - 96 c d^2 f f \sin[3(e + fx)] - (288I) c^2 d f^2 \sin[3(e + fx)] + 576 c^3 f^3 \sin[3(e + fx)] - 96 d^3 f x \sin[3(e + fx)] - (576I) c d^2 f^2 x \sin[3(e + fx)] + 1728 c^2 d f^3 x \sin[3(e + fx)] + (3456I) c^3 f^4 x \sin[3(e + fx)] - (288I) d^3 f^2 x^2 \sin[3(e + fx)] + 1728 c d^2 f^3 x^2 \sin[3(e + fx)] + (5184I) c^2 d f^4 x^2 \sin[3(e + fx)] + 576 d^3 f^3 x^3 \sin[3(e + fx)] + (3456I) c d^2 f^4 x^3 \sin[3(e + fx)] + (864I) d^3 f^4 x^4 \sin[3(e + fx)])) / (a^3 f^4 (-I + \tan[e + fx])^3)$$

input `Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x])^3,x]`

output `((I/27648)*Sec[e + f*x]^3*(243*((32*I)*c^3*f^3 + 8*c^2*d*f^2*(5 + (12*I)*f*x) + 4*c*d^2*f*(-9*I + 20*f*x + (24*I)*f^2*x^2) + d^3*(-17 - (36*I)*f*x + 40*f^2*x^2 + (32*I)*f^3*x^3))*Cos[e + f*x] + 16*(36*c^3*f^3*(I + 6*f*x) + 18*c^2*d*f^2*(1 + (6*I)*f*x + 18*f^2*x^2) + 6*c*d^2*f*(-I + 6*f*x + (18*I)*f^2*x^2 + 36*f^3*x^3) + d^3*(-1 - (6*I)*f*x + 18*f^2*x^2 + (36*I)*f^3*x^3 + 54*f^4*x^4))*Cos[3*(e + f*x)] - (3645*I)*d^3*Sin[e + f*x] + 6804*c*d^2*f*f*Sin[e + f*x] + (5832*I)*c^2*d*f^2*Sin[e + f*x] - 2592*c^3*f^3*Sin[e + f*x] + 6804*d^3*f*x*Sin[e + f*x] + (11664*I)*c*d^2*f^2*x*Sin[e + f*x] - 7776*c^2*d*f^3*x*Sin[e + f*x] + (5832*I)*d^3*f^2*x^2*Sin[e + f*x] - 7776*c*d^2*f^3*x^2*Sin[e + f*x] - 2592*d^3*f^3*x^3*Sin[e + f*x] + (16*I)*d^3*Sin[3*(e + f*x)] - 96*c*d^2*f*f*Sin[3*(e + f*x)] - (288*I)*c^2*d*f^2*Sin[3*(e + f*x)] + 576*c^3*f^3*Sin[3*(e + f*x)] - 96*d^3*f*x*Sin[3*(e + f*x)] - (576*I)*c*d^2*f^2*x*Sin[3*(e + f*x)] + 1728*c^2*d*f^3*x*Sin[3*(e + f*x)] + (3456*I)*c^3*f^4*x*Sin[3*(e + f*x)] - (288*I)*d^3*f^2*x^2*Sin[3*(e + f*x)] + 1728*c*d^2*f^3*x^2*Sin[3*(e + f*x)] + (5184*I)*c^2*d*f^4*x^2*Sin[3*(e + f*x)] + 576*d^3*f^3*x^3*Sin[3*(e + f*x)] + (3456*I)*c*d^2*f^4*x^3*Sin[3*(e + f*x)] + (864*I)*d^3*f^4*x^4*Sin[3*(e + f*x)])))/(a^3*f^4*(-I + Tan[e + f*x])^3)`

### 3.29.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.29.  $\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx$

$$\begin{aligned}
& \int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx \\
& \quad \downarrow \text{4212} \\
& \int \left( \frac{3(c+dx)^3 e^{-2ie-2ifx}}{8a^3} + \frac{3(c+dx)^3 e^{-4ie-4ifx}}{8a^3} + \frac{(c+dx)^3 e^{-6ie-6ifx}}{8a^3} + \frac{(c+dx)^3}{8a^3} \right) dx \\
& \quad \downarrow \text{2009} \\
& \frac{9id^2(c+dx)e^{-2ie-2ifx}}{32a^3 f^3} - \frac{9id^2(c+dx)e^{-4ie-4ifx}}{256a^3 f^3} - \frac{id^2(c+dx)e^{-6ie-6ifx}}{288a^3 f^3} + \frac{9d(c+dx)^2 e^{-2ie-2ifx}}{32a^3 f^2} + \\
& \frac{9d(c+dx)^2 e^{-4ie-4ifx}}{128a^3 f^2} + \frac{d(c+dx)^2 e^{-6ie-6ifx}}{96a^3 f^2} + \frac{3i(c+dx)^3 e^{-2ie-2ifx}}{16a^3 f} + \frac{3i(c+dx)^3 e^{-4ie-4ifx}}{32a^3 f} + \\
& \frac{i(c+dx)^3 e^{-6ie-6ifx}}{48a^3 f} + \frac{(c+dx)^4}{32a^3 d} - \frac{9d^3 e^{-2ie-2ifx}}{64a^3 f^4} - \frac{9d^3 e^{-4ie-4ifx}}{1024a^3 f^4} - \frac{d^3 e^{-6ie-6ifx}}{1728a^3 f^4}
\end{aligned}$$

input `Int[(c + d*x)^3/(a + I*a*Tan[e + f*x])^3,x]`

output `(-9*d^3*E^((-2*I)*e - (2*I)*f*x))/(64*a^3*f^4) - (9*d^3*E^((-4*I)*e - (4*I)*f*x))/(1024*a^3*f^4) - (d^3*E^((-6*I)*e - (6*I)*f*x))/(1728*a^3*f^4) - ((9*I)/32)*d^2*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)/(a^3*f^3) - (((9*I)/256)*d^2*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))/(a^3*f^3) - ((I/288)*d^2*E^((-6*I)*e - (6*I)*f*x)*(c + d*x))/(a^3*f^3) + (9*d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^2)/(32*a^3*f^2) + (9*d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^2)/(128*a^3*f^2) + (d*E^((-6*I)*e - (6*I)*f*x)*(c + d*x)^2)/(96*a^3*f^2) + (((3*I)/16)*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^3)/(a^3*f) + (((3*I)/32)*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^3)/(a^3*f) + ((I/48)*E^((-6*I)*e - (6*I)*f*x)*(c + d*x)^3)/(a^3*f) + (c + d*x)^4/(32*a^3*d)`

### 3.29.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



```
rule 4212 Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]
```

### 3.29.4 Maple [A] (verified)

Time = 1.21 (sec), antiderivative size = 396, normalized size of antiderivative = 1.00

method	result
risch	$\frac{d^3 x^4}{32a^3} + \frac{d^2 c x^3}{8a^3} + \frac{3dc^2 x^2}{16a^3} + \frac{c^3 x}{8a^3} + \frac{c^4}{32a^3 d} + \frac{3i(4d^3 x^3 f^3 + 12cd^2 f^3 x^2 - 6id^3 f^2 x^2 + 12c^2 d f^3 x - 12ic d^2 f^2 x + 4c^3 f^3 - 6ic^2 d f^2 - 6icd^2 f^2 - 6ic^2 d f^2 - 6icd^2 f^2)}{64a^3 f^4}$

```
input int((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/32/a^3*d^3*x^4+1/8/a^3*d^2*c*x^3+3/16/a^3*d*c^2*x^2+1/8/a^3*c^3*x+1/32/a^3/d*c^4+3/64*I*(4*d^3*x^3*f^3-6*I*d^3*f^2*x^2+12*c*d^2*f^3*x^2-12*I*c*d^2*f^2*x+12*c^2*d*f^3*x-6*I*c^2*d*f^2+4*c^3*f^3-6*d^3*f*x+3*I*d^3-6*c*d^2*f)/a^3/f^4*exp(-2*I*(f*x+e))+3/1024*I*(32*d^3*x^3*f^3-24*I*d^3*f^2*x^2+96*c*d^2*f^3*x^2-48*I*c*d^2*f^2*x+96*c^2*d*f^3*x-24*I*c^2*d*f^2+32*c^3*f^3-12*d^3*f*x+3*I*d^3-12*c*d^2*f)/a^3/f^4*exp(-4*I*(f*x+e))+1/1728*I*(36*d^3*x^3*f^3-18*I*d^3*f^2*x^2+108*c*d^2*f^3*x^2-36*I*c*d^2*f^2*x+108*c^2*d*f^3*x-18*I*c^2*d*f^2+36*c^3*f^3-6*d^3*f*x+I*d^3-6*c*d^2*f)/a^3/f^4*exp(-6*I*(f*x+e))
```

### 3.29.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec), antiderivative size = 370, normalized size of antiderivative = 0.93

$$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx$$

$$= \frac{(576i d^3 f^3 x^3 + 576i c^3 f^3 + 288 c^2 d f^2 - 96i c d^2 f - 16 d^3 - 288 (-6i c d^2 f^3 - d^3 f^2) x^2 - 96 (-18i c^2 d f^3 - 6i c d^2 f^2 - 6i c d^2 f^2 - 6i c d^2 f^2 - 6i c d^2 f^2 - 6i c d^2 f^2))}{(a+ia \tan(e+fx))^3}$$

```
input integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x, algorithm="fracas")
```

```
output 1/27648*(576*I*d^3*f^3*x^3 + 576*I*c^3*f^3 + 288*c^2*d*f^2 - 96*I*c*d^2*f
- 16*d^3 - 288*(-6*I*c*d^2*f^3 - d^3*f^2)*x^2 - 96*(-18*I*c^2*d*f^3 - 6*c*
d^2*f^2 + I*d^3*f)*x + 864*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^
2 + 4*c^3*f^4*x)*e^(6*I*f*x + 6*I*e) - 1296*(-4*I*d^3*f^3*x^3 - 4*I*c^3*f^
3 - 6*c^2*d*f^2 + 6*I*c*d^2*f + 3*d^3 + 6*(-2*I*c*d^2*f^3 - d^3*f^2)*x^2 +
6*(-2*I*c^2*d*f^3 - 2*c*d^2*f^2 + I*d^3*f)*x)*e^(4*I*f*x + 4*I*e) - 81*(-
32*I*d^3*f^3*x^3 - 32*I*c^3*f^3 - 24*c^2*d*f^2 + 12*I*c*d^2*f + 3*d^3 + 24
*(-4*I*c*d^2*f^3 - d^3*f^2)*x^2 + 12*(-8*I*c^2*d*f^3 - 4*c*d^2*f^2 + I*d^3
*f)*x)*e^(2*I*f*x + 2*I*e))*e^(-6*I*f*x - 6*I*e)/(a^3*f^4)
```

### 3.29.6 Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 945, normalized size of antiderivative = 2.39

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx$$

$$= \left\{ \frac{((2359296ia^6c^3f^{11}e^{6ie} + 7077888ia^6c^2df^{11}xe^{6ie} + 1179648a^6c^2df^{10}e^{6ie} + 7077888ia^6cd^2f^{11}x^2e^{6ie} + 2359296a^6cd^2f^{10}xe^{6ie} - 393216ia^6cd^2f^9e^{6ie} - 393216ia^6cd^2f^9e^{6ie} - 393216ia^6cd^2f^9e^{6ie} - 393216ia^6cd^2f^9e^{6ie})}{32a^3} + \frac{x^3 \cdot (3cd^2e^{4ie} + 3cd^2e^{2ie} + cd^2)e^{-6ie}}{8a^3} + \frac{x^2 \cdot (9c^2de^{4ie} + 9c^2de^{2ie} + 3c^2d)e^{-6ie}}{16a^3} + \frac{x(3c^3e^{4ie} + 3c^3e^{2ie} + c^3)e^{-6ie}}{8a^3} \right.$$

$$\left. + \frac{c^3x}{8a^3} + \frac{3c^2dx^2}{16a^3} + \frac{cd^2x^3}{8a^3} + \frac{d^3x^4}{32a^3} \right.$$

```
input integrate((d*x+c)**3/(a+I*a*tan(f*x+e))**3,x)
```

```
output Piecewise((((2359296*I**6*c**3*f**11*exp(6*I*e) + 7077888*I**6*c**2*d*
f**11*x*exp(6*I*e) + 1179648*a**6*c**2*d*f**10*exp(6*I*e) + 7077888*I**6
*c*d**2*f**11*x**2*exp(6*I*e) + 2359296*a**6*c*d**2*f**10*x*exp(6*I*e) - 3
93216*I**6*c*d**2*f**9*exp(6*I*e) + 2359296*I**6*d**3*f**11*x**3*exp(6
*I*e) + 1179648*a**6*d**3*f**10*x**2*exp(6*I*e) - 393216*I**6*d**3*f**9*
x*exp(6*I*e) - 65536*a**6*d**3*f**8*exp(6*I*e))*exp(-6*I*f*x) + (10616832*
I**6*c**3*f**11*exp(8*I*e) + 31850496*I**6*c**2*d*f**11*x*exp(8*I*e) +
7962624*a**6*c**2*d*f**10*exp(8*I*e) + 31850496*I**6*c*d**2*f**11*x**2*
exp(8*I*e) + 15925248*a**6*c*d**2*f**10*x*exp(8*I*e) - 3981312*I**6*c*d
**2*f**9*exp(8*I*e) + 10616832*I**6*d**3*f**11*x**3*exp(8*I*e) + 7962624*
a**6*d**3*f**10*x**2*exp(8*I*e) - 3981312*I**6*d**3*f**9*x*exp(8*I*e) -
995328*a**6*d**3*f**8*exp(8*I*e))*exp(-4*I*f*x) + (21233664*I**6*c**3*f*
**11*exp(10*I*e) + 63700992*I**6*c**2*d*f**11*x*exp(10*I*e) + 31850496*a
**6*c**2*d*f**10*exp(10*I*e) + 63700992*I**6*c*d**2*f**11*x**2*exp(10*I*
e) + 63700992*a**6*c*d**2*f**10*x*exp(10*I*e) - 31850496*I**6*c*d**2*f**9
*exp(10*I*e) + 21233664*I**6*d**3*f**11*x**3*exp(10*I*e) + 31850496*a**6
*d**3*f**10*x**2*exp(10*I*e) - 31850496*I**6*d**3*f**9*x*exp(10*I*e) - 1
5925248*a**6*d**3*f**8*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(113246208
*a**9*f**12), Ne(a**9*f**12*exp(12*I*e), 0)), (x**4*(3*d**3*exp(4*I*e) + 3
*d**3*exp(2*I*e) + d**3)*exp(-6*I*e)/(32*a**3) + x**3*(3*c*d**2*exp(4*I...
```

### 3.29.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

**3.29.8 Giac [A] (verification not implemented)**

Time = 0.58 (sec) , antiderivative size = 548, normalized size of antiderivative = 1.38

$$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx$$

$$= \frac{(864 d^3 f^4 x^4 e^{(6i fx+6i e)} + 3456 cd^2 f^4 x^3 e^{(6i fx+6i e)} + 5184 c^2 d f^4 x^2 e^{(6i fx+6i e)} + 5184 i d^3 f^3 x^3 e^{(4i fx+4i e)} + 2592 c^2 d^2 f^4 x^2 e^{(6i fx+6i e)} + 15552 I c d^2 f^3 x^2 e^{(4i fx+4i e)} + 7776 I c d^2 f^3 x^2 e^{(2i fx+2i e)} + 1728 I c d^2 f^3 x^2 + 15552 I c^2 d f^3 x e^{(4i fx+4i e)} + 7776 d^3 f^2 x^2 e^{(4i fx+4i e)} + 7776 I c^2 d f^3 x e^{(2i fx+2i e)} + 1944 d^3 f^2 x^2 e^{(2i fx+2i e)} + 1728 I c^2 d f^3 x + 288 d^3 f^2 x^2 + 5184 I c^3 f^3 e^{(4i fx+4i e)} + 15552 c d^2 f^2 x e^{(4i fx+4i e)} + 2592 I c^3 f^3 e^{(2i fx+2i e)} + 3888 c d^2 f^2 x e^{(2i fx+2i e)} + 576 I c^3 f^3 + 576 c d^2 f^2 x + 7776 c^2 d f^2 e^{(4i fx+4i e)} - 7776 I d^3 f^3 x e^{(4i fx+4i e)} + 1944 c^2 d f^2 e^{(2i fx+2i e)} - 972 I d^3 f^3 x e^{(2i fx+2i e)} + 288 c^2 d f^2 - 96 I d^3 f^3 x - 7776 I c d^2 f e^{(4i fx+4i e)} - 972 I c d^2 f e^{(2i fx+2i e)} - 96 I c d^2 f - 3888 d^3 e^{(4i fx+4i e)} - 243 d^3 e^{(2i fx+2i e)} - 16 d^3) e^{(-6i fx-6i e)}}{(a^3 f^4)}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output

```
1/27648*(864*d^3*f^4*x^4*e^(6*I*f*x + 6*I*e) + 3456*c*d^2*f^4*x^3*e^(6*I*f*x + 6*I*e) + 5184*c^2*d*f^4*x^2*e^(6*I*f*x + 6*I*e) + 5184*I*d^3*f^3*x^3*e^(4*I*f*x + 4*I*e) + 2592*I*d^3*f^3*x^3*e^(2*I*f*x + 2*I*e) + 576*I*d^3*f^3*x^3 + 3456*c^3*f^4*x*e^(6*I*f*x + 6*I*e) + 15552*I*c*d^2*f^3*x^2*e^(4*I*f*x + 4*I*e) + 7776*I*c*d^2*f^3*x^2*e^(2*I*f*x + 2*I*e) + 1728*I*c*d^2*f^3*x^2 + 15552*I*c^2*d*f^3*x*e^(4*I*f*x + 4*I*e) + 7776*d^3*f^2*x^2*e^(4*I*f*x + 4*I*e) + 7776*I*c^2*d*f^3*x*e^(2*I*f*x + 2*I*e) + 1944*d^3*f^2*x^2*e^(2*I*f*x + 2*I*e) + 1728*I*c^2*d*f^3*x + 288*d^3*f^2*x^2 + 5184*I*c^3*f^3*e^(4*I*f*x + 4*I*e) + 15552*c*d^2*f^2*x*e^(4*I*f*x + 4*I*e) + 2592*I*c^3*f^3*e^(2*I*f*x + 2*I*e) + 3888*c*d^2*f^2*x*e^(2*I*f*x + 2*I*e) + 576*I*c^3*f^3 + 576*c*d^2*f^2*x + 7776*c^2*d*f^2*e^(4*I*f*x + 4*I*e) - 7776*I*d^3*f^3*x*e^(4*I*f*x + 4*I*e) + 1944*c^2*d*f^2*e^(2*I*f*x + 2*I*e) - 972*I*d^3*f^3*x*e^(2*I*f*x + 2*I*e) + 288*c^2*d*f^2 - 96*I*d^3*f^3*x - 7776*I*c*d^2*f*e^(4*I*f*x + 4*I*e) - 972*I*c*d^2*f*e^(2*I*f*x + 2*I*e) - 96*I*c*d^2*f - 3888*d^3*e^(4*I*f*x + 4*I*e) - 243*d^3*e^(2*I*f*x + 2*I*e) - 16*d^3)*e^(-6*I*f*x - 6*I*e)/(a^3*f^4)
```

**3.29.9 Mupad [B] (verification not implemented)**

Time = 4.25 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.04

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx = e^{-e 2i - f x 2i} \left( \frac{(12 c^3 f^3 - c^2 d f^2 18i - 18 c d^2 f + d^3 9i) 1i}{64 a^3 f^4} \right. \\ \left. + \frac{d^3 x^3 3i}{16 a^3 f} - \frac{dx (-2 c^2 f^2 + c d f 2i + d^2) 9i}{32 a^3 f^3} - \frac{d^2 x^2 (-2 c f + d 1i) 9i}{32 a^3 f^2} \right) \\ + e^{-e 4i - f x 4i} \left( \frac{(96 c^3 f^3 - c^2 d f^2 72i - 36 c d^2 f + d^3 9i) 1i}{1024 a^3 f^4} \right. \\ \left. + \frac{d^3 x^3 3i}{32 a^3 f} - \frac{dx (-8 c^2 f^2 + c d f 4i + d^2) 9i}{256 a^3 f^3} - \frac{d^2 x^2 (-4 c f + d 1i) 9i}{128 a^3 f^2} \right) \\ + e^{-e 6i - f x 6i} \left( \frac{(36 c^3 f^3 - c^2 d f^2 18i - 6 c d^2 f + d^3 1i) 1i}{1728 a^3 f^4} \right. \\ \left. + \frac{d^3 x^3 1i}{48 a^3 f} - \frac{dx (-18 c^2 f^2 + c d f 6i + d^2) 1i}{288 a^3 f^3} - \frac{d^2 x^2 (-6 c f + d 1i) 1i}{96 a^3 f^2} \right) + \frac{c^3 x}{8 a^3} + \frac{d^3 x^4}{32 a^3} + \frac{3 c^2 d x^2}{16 a^3} + \frac{c d^2 x^3}{8 a^3}$$

input `int((c + d*x)^3/(a + a*tan(e + f*x)*1i)^3,x)`

```
output exp(- e*2i - f*x*2i)*(((d^3*9i + 12*c^3*f^3 - c^2*d*f^2*18i - 18*c*d^2*f)*
1i)/(64*a^3*f^4) + (d^3*x^3*3i)/(16*a^3*f) - (d*x*(d^2 - 2*c^2*f^2 + c*d*f
*2i)*9i)/(32*a^3*f^3) - (d^2*x^2*(d*1i - 2*c*f)*9i)/(32*a^3*f^2)) + exp(-
e*4i - f*x*4i)*(((d^3*9i + 96*c^3*f^3 - c^2*d*f^2*72i - 36*c*d^2*f)*1i)/(1
024*a^3*f^4) + (d^3*x^3*3i)/(32*a^3*f) - (d*x*(d^2 - 8*c^2*f^2 + c*d*f*4i)
*9i)/(256*a^3*f^3) - (d^2*x^2*(d*1i - 4*c*f)*9i)/(128*a^3*f^2)) + exp(- e
6i - f*x*6i)*(((d^3*1i + 36*c^3*f^3 - c^2*d*f^2*18i - 6*c*d^2*f)*1i)/(1728
*a^3*f^4) + (d^3*x^3*1i)/(48*a^3*f) - (d*x*(d^2 - 18*c^2*f^2 + c*d*f*6i)*1
i)/(288*a^3*f^3) - (d^2*x^2*(d*1i - 6*c*f)*1i)/(96*a^3*f^2)) + (c^3*x)/(8*
a^3) + (d^3*x^4)/(32*a^3) + (3*c^2*d*x^2)/(16*a^3) + (c*d^2*x^3)/(8*a^3)
```

### 3.30 $\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^3} dx$

3.30.1	Optimal result . . . . .	221
3.30.2	Mathematica [A] (verified) . . . . .	222
3.30.3	Rubi [A] (verified) . . . . .	222
3.30.4	Maple [A] (verified) . . . . .	224
3.30.5	Fricas [A] (verification not implemented) . . . . .	224
3.30.6	Sympy [A] (verification not implemented) . . . . .	225
3.30.7	Maxima [F(-2)] . . . . .	225
3.30.8	Giac [A] (verification not implemented) . . . . .	226
3.30.9	Mupad [B] (verification not implemented) . . . . .	226

#### 3.30.1 Optimal result

Integrand size = 23, antiderivative size = 294

$$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^3} dx = -\frac{3id^2e^{-2ie-2ifx}}{32a^3f^3} - \frac{3id^2e^{-4ie-4ifx}}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}}{864a^3f^3} + \frac{3de^{-2ie-2ifx}(c+dx)}{16a^3f^2} + \frac{3de^{-4ie-4ifx}(c+dx)}{64a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)}{144a^3f^2} + \frac{3ie^{-2ie-2ifx}(c+dx)^2}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^2}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^2}{48a^3f} + \frac{(c+dx)^3}{24a^3d}$$

output

```
-3/32*I*d^2*exp(-2*I*e-2*I*f*x)/a^3/f^3-3/256*I*d^2*exp(-4*I*e-4*I*f*x)/a^3/f^3-1/864*I*d^2*exp(-6*I*e-6*I*f*x)/a^3/f^3+3/16*d*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^3/f^2+3/64*d*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^3/f^2+1/144*d*exp(-6*I*e-6*I*f*x)*(d*x+c)/a^3/f^2+3/16*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^3/f+3/32*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^3/f+1/48*I*exp(-6*I*e-6*I*f*x)*(d*x+c)^2/a^3/f+1/24*(d*x+c)^3/a^3/d
```

### 3.30.2 Mathematica [A] (verified)

Time = 2.74 (sec) , antiderivative size = 405, normalized size of antiderivative = 1.38

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{i \sec^3(e + fx) (81(24ic^2 f^2 + 4cdf(5 + 12ifx) + d^2(-9i + 20fx + 24if^2 x^2)) \cos(e + fx) + 8(18c^2 f^2(i + 6$$

input `Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x])^3,x]`

output `((I/6912)*Sec[e + f*x]^3*(81*((24*I)*c^2*f^2 + 4*c*d*f*(5 + (12*I)*f*x) + d^2*(-9*I + 20*f*x + (24*I)*f^2*x^2))*Cos[e + f*x] + 8*(18*c^2*f^2*(I + 6*f*x) + 6*c*d*f*(1 + (6*I)*f*x + 18*f^2*x^2) + d^2*(-I + 6*f*x + (18*I)*f^2*x^2 + 36*f^3*x^3))*Cos[3*(e + f*x)] + 567*d^2*Sin[e + f*x] + (972*I)*c*d*f*Sin[e + f*x] - 648*c^2*f^2*Sin[e + f*x] + (972*I)*d^2*f*x*Sin[e + f*x] - 1296*c*d*f^2*x*Sin[e + f*x] - 648*d^2*f^2*x^2*Sin[e + f*x] - 8*d^2*Sin[3*(e + f*x)] - (48*I)*c*d*f*Sin[3*(e + f*x)] + 144*c^2*f^2*Sin[3*(e + f*x)] - (48*I)*d^2*f*x*Sin[3*(e + f*x)] + 288*c*d*f^2*x*Sin[3*(e + f*x)] + (864*I)*c^2*f^3*x*Sin[3*(e + f*x)] + 144*d^2*f^2*x^2*Sin[3*(e + f*x)] + (864*I)*c*d*f^3*x^2*Sin[3*(e + f*x)] + (288*I)*d^2*f^3*x^3*Sin[3*(e + f*x)]))/(a^3*f^3*(-I + Tan[e + f*x])^3)`

### 3.30.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx$$

$$\downarrow \text{4212}$$

$$\int \left( \frac{3(c+dx)^2 e^{-2ie-2ifx}}{8a^3} + \frac{3(c+dx)^2 e^{-4ie-4ifx}}{8a^3} + \frac{(c+dx)^2 e^{-6ie-6ifx}}{8a^3} + \frac{(c+dx)^2}{8a^3} \right) dx$$

↓ 2009

$$\frac{3d(c+dx)e^{-2ie-2ifx}}{16a^3 f^2} + \frac{3d(c+dx)e^{-4ie-4ifx}}{64a^3 f^2} + \frac{d(c+dx)e^{-6ie-6ifx}}{144a^3 f^2} + \frac{3i(c+dx)^2 e^{-2ie-2ifx}}{16a^3 f} + \frac{3i(c+dx)^2 e^{-4ie-4ifx}}{32a^3 f} + \frac{i(c+dx)^2 e^{-6ie-6ifx}}{48a^3 f} + \frac{(c+dx)^3}{24a^3 d} - \frac{3id^2 e^{-2ie-2ifx}}{32a^3 f^3} - \frac{3id^2 e^{-4ie-4ifx}}{256a^3 f^3} - \frac{id^2 e^{-6ie-6ifx}}{864a^3 f^3}$$

input `Int[(c + d*x)^2/(a + I*a*Tan[e + f*x])^3,x]`

output `(((-3*I)/32)*d^2*E^((-2*I)*e - (2*I)*f*x))/(a^3*f^3) - (((3*I)/256)*d^2*E^((-4*I)*e - (4*I)*f*x))/(a^3*f^3) - ((I/864)*d^2*E^((-6*I)*e - (6*I)*f*x))/(a^3*f^3) + (3*d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x))/(16*a^3*f^2) + (3*d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))/(64*a^3*f^2) + (d*E^((-6*I)*e - (6*I)*f*x)*(c + d*x))/(144*a^3*f^2) + (((3*I)/16)*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^2)/(a^3*f) + (((3*I)/32)*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^2)/(a^3*f) + ((I/48)*E^((-6*I)*e - (6*I)*f*x)*(c + d*x)^2)/(a^3*f) + (c + d*x)^3/(24*a^3*d)`

### 3.30.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4212 `Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]`



### 3.30.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.81

method	result
risch	$\frac{d^2 x^3}{24a^3} + \frac{dcx^2}{8a^3} + \frac{c^2 x}{8a^3} + \frac{c^3}{24a^3 d} + \frac{3i(2d^2 x^2 f^2 + 4cd f^2 x - 2id^2 f x + 2c^2 f^2 - 2icdf - d^2)e^{-2i(fx+e)}}{32a^3 f^3} + \frac{3i(8d^2 x^2 f^2 + 16cd f^2 x - 4id^2 f x + 2c^2 f^2 - 2icdf - d^2)e^{-2i(fx+e)}}{32a^3 f^3}$

input `int((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output  $\frac{1}{24} \frac{d^2 x^3}{a^3} + \frac{1}{8} \frac{dcx^2}{a^3} + \frac{1}{8} \frac{c^2 x}{a^3} + \frac{1}{24} \frac{c^3}{a^3 d} + \frac{3}{32} \frac{I(2d^2 x^2 f^2 - 2I d^2 f x + 4c d f^2 x - 2I c d f + 2c^2 f^2 - d^2)}{a^3 f^3} \exp(-2I(fx+e)) + \frac{3}{256} \frac{I(8d^2 x^2 f^2 - 4I d^2 f x + 16c d f^2 x - 4I c d f + 8c^2 f^2 - d^2)}{a^3 f^3} \exp(-4I(fx+e)) + \frac{1}{864} \frac{I(18d^2 x^2 f^2 - 6I d^2 f x + 36c d f^2 x - 6I c d f + 18c^2 f^2 - d^2)}{a^3 f^3} \exp(-6I(fx+e))$

### 3.30.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.75

$$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^3} dx$$

$$= \frac{(144i d^2 f^2 x^2 + 144i c^2 f^2 + 48cdf - 8i d^2 - 48(-6icdf^2 - d^2 f)x + 288(d^2 f^3 x^3 + 3cdf^3 x^2 + 3c^2 f^3 x))e^{6i(fx+e)}}{(a+ia \tan(e+fx))^3}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="fracas")`

output  $\frac{1}{6912} (144I d^2 f^2 x^2 + 144I c^2 f^2 + 48c d f - 8I d^2 - 48(-6I c d f^2 - d^2 f) x + 288(d^2 f^3 x^3 + 3c d f^3 x^2 + 3c^2 f^3 x)) e^{6I(fx+e)} - \frac{648(-2I d^2 f^2 x^2 - 2I c^2 f^2 - 2c d f + I d^2 + 2(-2I c d f^2 - d^2 f) x) e^{4I(fx+e)} - 81(-8I d^2 f^2 x^2 - 8I c^2 f^2 - 4c d f + I d^2 + 4(-4I c d f^2 - d^2 f) x) e^{2I(fx+e)}}{(a^3 f^3)}$

### 3.30.6 Sympy [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 588, normalized size of antiderivative = 2.00

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx$$

$$= \left\{ \frac{((147456ia^6c^2f^8e^{6ie} + 294912ia^6cdf^8xe^{6ie} + 49152a^6cdf^7e^{6ie} + 147456ia^6d^2f^8x^2e^{6ie} + 49152a^6d^2f^7xe^{6ie} - 8192ia^6d^2f^6e^{6ie})e^{-6ifx} + (663552x^3 \cdot (3d^2e^{4ie} + 3d^2e^{2ie} + d^2)e^{-6ie} + \frac{x^2 \cdot (3cde^{4ie} + 3cde^{2ie} + cd)e^{-6ie}}{8a^3} + \frac{x(3c^2e^{4ie} + 3c^2e^{2ie} + c^2)e^{-6ie}}{8a^3})}{24a^3} + \frac{c^2x}{8a^3} + \frac{cdx^2}{8a^3} + \frac{d^2x^3}{24a^3} \right.$$

input `integrate((d*x+c)**2/(a+I*a*tan(f*x+e))**3,x)`

output `Piecewise((((147456*I*a**6*c**2*f**8*exp(6*I*e) + 294912*I*a**6*c*d*f**8*x*exp(6*I*e) + 49152*a**6*c*d*f**7*exp(6*I*e) + 147456*I*a**6*d**2*f**8*x**2*exp(6*I*e) + 49152*a**6*d**2*f**7*x*exp(6*I*e) - 8192*I*a**6*d**2*f**6*exp(6*I*e))*exp(-6*I*f*x) + (663552*I*a**6*c**2*f**8*exp(8*I*e) + 1327104*I*a**6*c*d*f**8*x*exp(8*I*e) + 331776*a**6*c*d*f**7*exp(8*I*e) + 663552*I*a**6*d**2*f**8*x**2*exp(8*I*e) + 331776*a**6*d**2*f**7*x*exp(8*I*e) - 82944*I*a**6*d**2*f**6*exp(8*I*e))*exp(-4*I*f*x) + (1327104*I*a**6*c**2*f**8*exp(10*I*e) + 2654208*I*a**6*c*d*f**8*x*exp(10*I*e) + 1327104*a**6*c*d*f**7*exp(10*I*e) + 1327104*I*a**6*d**2*f**8*x**2*exp(10*I*e) + 1327104*a**6*d**2*f**7*x*exp(10*I*e) - 663552*I*a**6*d**2*f**6*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(7077888*a**9*f**9), Ne(a**9*f**9*exp(12*I*e), 0)), (x**3*(3*d**2*exp(4*I*e) + 3*d**2*exp(2*I*e) + d**2)*exp(-6*I*e)/(24*a**3) + x**2*(3*c*d*exp(4*I*e) + 3*c*d*exp(2*I*e) + c*d)*exp(-6*I*e)/(8*a**3) + x*(3*c**2*exp(4*I*e) + 3*c**2*exp(2*I*e) + c**2)*exp(-6*I*e)/(8*a**3), True)) + c**2*x/(8*a**3) + c*d*x**2/(8*a**3) + d**2*x**3/(24*a**3)`

### 3.30.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

### 3.30.8 Giac [A] (verification not implemented)

Time = 0.62 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(288 d^2 f^3 x^3 e^{(6i fx + 6i e)} + 864 c d f^3 x^2 e^{(6i fx + 6i e)} + 864 c^2 f^3 x e^{(6i fx + 6i e)} + 1296 i d^2 f^2 x^2 e^{(4i fx + 4i e)} + 648 i d^2 f^2 x e^{(4i fx + 4i e)} + 648 i d^2 f^2 x^2 e^{(2i fx + 2i e)} + 144 i d^2 f^2 x^2 e^{(2i fx + 2i e)} + 144 i d^2 f^2 x^2 e^{(2i fx + 2i e)} + 2592 i c d f^2 x^2 e^{(4i fx + 4i e)} + 1296 i c d f^2 x^2 e^{(2i fx + 2i e)} + 288 i c d f^2 x^2 e^{(2i fx + 2i e)} + 1296 i c^2 f^2 x^2 e^{(4i fx + 4i e)} + 1296 i c^2 f^2 x^2 e^{(4i fx + 4i e)} + 648 i c^2 f^2 x^2 e^{(2i fx + 2i e)} + 324 i c^2 f^2 x^2 e^{(2i fx + 2i e)} + 144 i c^2 f^2 x^2 e^{(2i fx + 2i e)} + 48 i c^2 f^2 x^2 e^{(2i fx + 2i e)} + 1296 i c d f x e^{(4i fx + 4i e)} + 1296 i c d f x e^{(4i fx + 4i e)} + 324 i c d f x e^{(2i fx + 2i e)} + 324 i c d f x e^{(2i fx + 2i e)} + 48 i c d f x e^{(2i fx + 2i e)} - 648 i d^2 x e^{(4i fx + 4i e)} + 648 i d^2 x e^{(4i fx + 4i e)} - 81 i d^2 x e^{(2i fx + 2i e)} - 81 i d^2 x e^{(2i fx + 2i e)} - 8 i d^2 x e^{(-6i fx - 6i e)})}{(a^3 f^3)}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `1/6912*(288*d^2*f^3*x^3*e^(6*I*f*x + 6*I*e) + 864*c*d*f^3*x^2*e^(6*I*f*x + 6*I*e) + 864*c^2*f^3*x*e^(6*I*f*x + 6*I*e) + 1296*I*d^2*f^2*x^2*e^(4*I*f*x + 4*I*e) + 648*I*d^2*f^2*x^2*e^(2*I*f*x + 2*I*e) + 144*I*d^2*f^2*x^2 + 2592*I*c*d*f^2*x^2*e^(4*I*f*x + 4*I*e) + 1296*I*c*d*f^2*x^2*e^(2*I*f*x + 2*I*e) + 288*I*c*d*f^2*x^2 + 1296*I*c^2*f^2*x^2*e^(4*I*f*x + 4*I*e) + 1296*d^2*f*x*e^(4*I*f*x + 4*I*e) + 648*I*c^2*f^2*x^2*e^(2*I*f*x + 2*I*e) + 324*d^2*f*x*e^(2*I*f*x + 2*I*e) + 144*I*c^2*f^2 + 48*d^2*f*x + 1296*c*d*f*x*e^(4*I*f*x + 4*I*e) + 324*c*d*f*x*e^(2*I*f*x + 2*I*e) + 48*c*d*f - 648*I*d^2*x*e^(4*I*f*x + 4*I*e) - 81*I*d^2*x*e^(2*I*f*x + 2*I*e) - 8*I*d^2)*e^(-6*I*f*x - 6*I*e)/(a^3*f^3)`

### 3.30.9 Mupad [B] (verification not implemented)

Time = 3.95 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.89

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx = \frac{c^2 x}{8 a^3} - e^{-e 2i - f x 2i} \left( \frac{(-6 c^2 f^2 + c d f 6i + 3 d^2) li}{32 a^3 f^3} - \frac{d^2 x^2 3i}{16 a^3 f} + \frac{d x (-2 c f + d li) 3i}{16 a^3 f^2} \right)$$

$$- e^{-e 4i - f x 4i} \left( \frac{(-24 c^2 f^2 + c d f 12i + 3 d^2) li}{256 a^3 f^3} - \frac{d^2 x^2 3i}{32 a^3 f} + \frac{d x (-4 c f + d li) 3i}{64 a^3 f^2} \right)$$

$$- e^{-e 6i - f x 6i} \left( \frac{(-18 c^2 f^2 + c d f 6i + d^2) li}{864 a^3 f^3} - \frac{d^2 x^2 li}{48 a^3 f} + \frac{d x (-6 c f + d li) li}{144 a^3 f^2} \right) + \frac{d^2 x^3}{24 a^3} + \frac{c d x^2}{8 a^3}$$

3.30.  $\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^3} dx$

input `int((c + d*x)^2/(a + a*tan(e + f*x)*1i)^3,x)`

output `(c^2*x)/(8*a^3) - exp(- e*2i - f*x*2i)*(((3*d^2 - 6*c^2*f^2 + c*d*f*6i)*1i)/(32*a^3*f^3) - (d^2*x^2*3i)/(16*a^3*f) + (d*x*(d*1i - 2*c*f)*3i)/(16*a^3*f^2)) - exp(- e*4i - f*x*4i)*(((3*d^2 - 24*c^2*f^2 + c*d*f*12i)*1i)/(256*a^3*f^3) - (d^2*x^2*3i)/(32*a^3*f) + (d*x*(d*1i - 4*c*f)*3i)/(64*a^3*f^2)) - exp(- e*6i - f*x*6i)*(((d^2 - 18*c^2*f^2 + c*d*f*6i)*1i)/(864*a^3*f^3) - (d^2*x^2*1i)/(48*a^3*f) + (d*x*(d*1i - 6*c*f)*1i)/(144*a^3*f^2)) + (d^2*x^3)/(24*a^3) + (c*d*x^2)/(8*a^3)`

### 3.31 $\int \frac{c+dx}{(a+ia \tan(e+fx))^3} dx$

3.31.1	Optimal result . . . . .	228
3.31.2	Mathematica [A] (verified) . . . . .	229
3.31.3	Rubi [A] (verified) . . . . .	229
3.31.4	Maple [A] (verified) . . . . .	230
3.31.5	Fricas [A] (verification not implemented) . . . . .	231
3.31.6	Sympy [A] (verification not implemented) . . . . .	231
3.31.7	Maxima [F(-2)] . . . . .	232
3.31.8	Giac [A] (verification not implemented) . . . . .	232
3.31.9	Mupad [B] (verification not implemented) . . . . .	233

#### 3.31.1 Optimal result

Integrand size = 21, antiderivative size = 209

$$\int \frac{c+dx}{(a+ia \tan(e+fx))^3} dx = -\frac{11idx}{96a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3}$$

$$+ \frac{d}{36f^2(a+ia \tan(e+fx))^3} + \frac{i(c+dx)}{6f(a+ia \tan(e+fx))^3}$$

$$+ \frac{5d}{96af^2(a+ia \tan(e+fx))^2} + \frac{i(c+dx)}{8af(a+ia \tan(e+fx))^2}$$

$$+ \frac{11d}{96f^2(a^3+ia^3 \tan(e+fx))} + \frac{i(c+dx)}{8f(a^3+ia^3 \tan(e+fx))}$$

output `-11/96*I*d*x/a^3/f-1/16*d*x^2/a^3+1/8*x*(d*x+c)/a^3+1/36*d/f^2/(a+I*a*tan(f*x+e))^3+1/6*I*(d*x+c)/f/(a+I*a*tan(f*x+e))^3+5/96*d/a/f^2/(a+I*a*tan(f*x+e))^2+1/8*I*(d*x+c)/a/f/(a+I*a*tan(f*x+e))^2+11/96*d/f^2/(a^3+I*a^3*tan(f*x+e))+1/8*I*(d*x+c)/f/(a^3+I*a^3*tan(f*x+e))`

### 3.31.2 Mathematica [A] (verified)

Time = 1.52 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.98

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{i \sec^3(e + fx) (27(12icf + d(5 + 12ifx)) \cos(e + fx) + 4(6cf(i + 6fx) + d(1 + 6ifx + 18f^2x^2)) \cos(3(e$$

input `Integrate[(c + d*x)/(a + I*a*Tan[e + f*x])^3,x]`

output `((I/1152)*Sec[e + f*x]^3*(27*((12*I)*c*f + d*(5 + (12*I)*f*x))*Cos[e + f*x] + 4*(6*c*f*(I + 6*f*x) + d*(1 + (6*I)*f*x + 18*f^2*x^2))*Cos[3*(e + f*x)] + (81*I)*d*Sin[e + f*x] - 108*c*f*Sin[e + f*x] - 108*d*f*x*Sin[e + f*x] - (4*I)*d*Sin[3*(e + f*x)] + 24*c*f*Sin[3*(e + f*x)] + 24*d*f*x*Sin[3*(e + f*x)] + (144*I)*c*f^2*x*Sin[3*(e + f*x)] + (72*I)*d*f^2*x^2*Sin[3*(e + f*x)]))/(a^3*f^2*(-I + Tan[e + f*x])^3)`

### 3.31.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {3042, 4213, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx$$

$$\downarrow \text{4213}$$

$$-d \int \left( \frac{x}{8a^3} + \frac{i}{8f(i \tan(e + fx)a^3 + a^3)} + \frac{i}{8af(i \tan(e + fx)a + a)^2} + \frac{i}{6f(i \tan(e + fx)a + a)^3} \right) dx +$$

$$\frac{i(c + dx)}{8f(a^3 + ia^3 \tan(e + fx))} + \frac{x(c + dx)}{8a^3} + \frac{i(c + dx)}{8af(a + ia \tan(e + fx))^2} + \frac{i(c + dx)}{6f(a + ia \tan(e + fx))^3}$$

$$\downarrow \text{2009}$$

---

3.31.  $\int \frac{c+dx}{(a+ia \tan(e+fx))^3} dx$

$$d\left(\frac{11}{96f^2(a^3 + ia^3 \tan(e + fx))} + \frac{11ix}{96a^3 f} + \frac{x^2}{16a^3} - \frac{5}{96af^2(a + ia \tan(e + fx))^2} - \frac{1}{36f^2(a + ia \tan(e + fx))^3} + \frac{i(c + dx)}{8f(a^3 + ia^3 \tan(e + fx))} + \frac{x(c + dx)}{8a^3} - \frac{i(c + dx)}{8af(a + ia \tan(e + fx))^2} + \frac{i(c + dx)}{6f(a + ia \tan(e + fx))^3}\right)$$

input `Int[(c + d*x)/(a + I*a*Tan[e + f*x])^3,x]`

output `(x*(c + d*x))/(8*a^3) + ((I/6)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])^3) + ((I/8)*(c + d*x))/(a*f*(a + I*a*Tan[e + f*x])^2) + ((I/8)*(c + d*x))/(f*(a^3 + I*a^3*Tan[e + f*x])) - d*(((11*I)/96)*x)/(a^3*f) + x^2/(16*a^3) - 1/(36*f^2*(a + I*a*Tan[e + f*x])^3) - 5/(96*a*f^2*(a + I*a*Tan[e + f*x])^2) - 11/(96*f^2*(a^3 + I*a^3*Tan[e + f*x]))`

### 3.31.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4213 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := With[{u = IntHide[(a + b*Tan[e + f*x])^n, x]}, Simp[(c + d*x)^m u, x] - Simp[d*m Int[(c + d*x)^(m - 1) u, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, -1] && GtQ[m, 0]`

### 3.31.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.55

method	result	size
risch	$\frac{dx^2}{16a^3} + \frac{cx}{8a^3} + \frac{3i(2dfx+2cf-id)e^{-2i(fx+e)}}{32a^3f^2} + \frac{3i(4dfx+4cf-id)e^{-4i(fx+e)}}{128a^3f^2} + \frac{i(6dfx+6cf-id)e^{-6i(fx+e)}}{288a^3f^2}$	114

input `int((d*x+c)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output  $\frac{1}{16}d^2x^2/a^3 + 1/8/a^3cx + 3/32I*(2d^2fx - Id + 2cf)/a^3/f^2 \exp(-2I*(fx+e)) + 3/128I*(4d^2fx - Id + 4cf)/a^3/f^2 \exp(-4I*(fx+e)) + 1/288I*(6d^2fx - Id + 6cf)/a^3/f^2 \exp(-6I*(fx+e))$

### 3.31.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.50

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(24i dfx + 24i cf + 72(df^2x^2 + 2cf^2x)e^{(6ifx+6ie)} - 108(-2i dfx - 2i cf - d)e^{(4ifx+4ie)} - 27(-4i dfx - 4i cf - d)e^{(2ifx+2ie)} + 4d)e^{(-6ifx - 6ie)}}{1152a^3f^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output  $\frac{1}{1152}*(24*I*d*f*x + 24*I*c*f + 72*(d*f^2*x^2 + 2*c*f^2*x)*e^{(6*I*f*x + 6*I*e)} - 108*(-2*I*d*f*x - 2*I*c*f - d)*e^{(4*I*f*x + 4*I*e)} - 27*(-4*I*d*f*x - 4*I*c*f - d)*e^{(2*I*f*x + 2*I*e)} + 4*d)*e^{(-6*I*f*x - 6*I*e)}/(a^3*f^2)$

### 3.31.6 Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.49

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx$$

$$= \left\{ \frac{((24576ia^6cf^5e^{6ie} + 24576ia^6df^5xe^{6ie} + 4096a^6df^4e^{6ie})e^{-6ifx} + (110592ia^6cf^5e^{8ie} + 110592ia^6df^5xe^{8ie} + 27648a^6df^4e^{8ie})e^{-4ifx} + (221184ia^6cf^5e^{10ie} + 221184ia^6df^5xe^{10ie} + 110592a^6df^4e^{10ie})e^{-2ifx} + 4d)e^{-6ie}}{1179648a^9f^6} \right.$$

$$+ \frac{x^2 \cdot (3de^{4ie} + 3de^{2ie} + d)e^{-6ie}}{16a^3} + \frac{x(3ce^{4ie} + 3ce^{2ie} + c)e^{-6ie}}{8a^3}$$

$$+ \frac{cx}{8a^3} + \frac{dx^2}{16a^3}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))**3,x)`



```
output Piecewise((((24576*I*a**6*c*f**5*exp(6*I*e) + 24576*I*a**6*d*f**5*x*exp(6*
I*e) + 4096*a**6*d*f**4*exp(6*I*e))*exp(-6*I*f*x) + (110592*I*a**6*c*f**5*
exp(8*I*e) + 110592*I*a**6*d*f**5*x*exp(8*I*e) + 27648*a**6*d*f**4*exp(8*I
*e))*exp(-4*I*f*x) + (221184*I*a**6*c*f**5*exp(10*I*e) + 221184*I*a**6*d*f
**5*x*exp(10*I*e) + 110592*a**6*d*f**4*exp(10*I*e))*exp(-2*I*f*x))*exp(-12
*I*e)/(1179648*a**9*f**6), Ne(a**9*f**6*exp(12*I*e), 0)), (x**2*(3*d*exp(4
*I*e) + 3*d*exp(2*I*e) + d)*exp(-6*I*e)/(16*a**3) + x*(3*c*exp(4*I*e) + 3*
c*exp(2*I*e) + c)*exp(-6*I*e)/(8*a**3), True)) + c*x/(8*a**3) + d*x**2/(16
*a**3)
```

### 3.31.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

### 3.31.8 Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.68

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \frac{(72df^2x^2e^{(6i fx+6ie)} + 144cf^2xe^{(6i fx+6ie)} + 216idfxe^{(4i fx+4ie)} + 108idfxe^{(2i fx+2ie)} + 24idfx + 216icfe)}{1152a^3f^2}$$

```
input integrate((d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
```

```
output 1/1152*(72*d*f^2*x^2*e^(6*I*f*x + 6*I*e) + 144*c*f^2*x*e^(6*I*f*x + 6*I*e)
+ 216*I*d*f*x*e^(4*I*f*x + 4*I*e) + 108*I*d*f*x*e^(2*I*f*x + 2*I*e) + 24*
I*d*f*x + 216*I*c*f*e^(4*I*f*x + 4*I*e) + 108*I*c*f*e^(2*I*f*x + 2*I*e) +
24*I*c*f + 108*d*e^(4*I*f*x + 4*I*e) + 27*d*e^(2*I*f*x + 2*I*e) + 4*d)*e^(
-6*I*f*x - 6*I*e)/(a^3*f^2)
```

---

3.31.  $\int \frac{c+dx}{(a+ia \tan(e+fx))^3} dx$

**3.31.9 Mupad [B] (verification not implemented)**

Time = 3.94 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.70

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \frac{dx^2}{16a^3} - e^{-e4i - fx4i} \left( \frac{(-12cf + d3i) li}{128a^3 f^2} - \frac{dx3i}{32a^3 f} \right) - e^{-e6i - fx6i} \left( \frac{(-6cf + d1i) li}{288a^3 f^2} - \frac{dx1i}{48a^3 f} \right) - e^{-e2i - fx2i} \left( \frac{(-6cf + d3i) li}{32a^3 f^2} - \frac{dx3i}{16a^3 f} \right) + \frac{cx}{8a^3}$$

input `int((c + d*x)/(a + a*tan(e + f*x)*1i)^3,x)`output `(d*x^2)/(16*a^3) - exp(- e*4i - f*x*4i)*(((d*3i - 12*c*f)*1i)/(128*a^3*f^2) - (d*x*3i)/(32*a^3*f)) - exp(- e*6i - f*x*6i)*(((d*1i - 6*c*f)*1i)/(288*a^3*f^2) - (d*x*1i)/(48*a^3*f)) - exp(- e*2i - f*x*2i)*(((d*3i - 6*c*f)*1i)/(32*a^3*f^2) - (d*x*3i)/(16*a^3*f)) + (c*x)/(8*a^3)`

$$\mathbf{3.32} \quad \int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$$

3.32.1	Optimal result . . . . .	235
3.32.2	Mathematica [A] (verified) . . . . .	236
3.32.3	Rubi [A] (verified) . . . . .	236
3.32.4	Maple [A] (verified) . . . . .	238
3.32.5	Fricas [A] (verification not implemented) . . . . .	238
3.32.6	Sympy [F] . . . . .	239
3.32.7	Maxima [A] (verification not implemented) . . . . .	239
3.32.8	Giac [A] (verification not implemented) . . . . .	240
3.32.9	Mupad [F(-1)] . . . . .	240

### 3.32.1 Optimal result

Integrand size = 23, antiderivative size = 449

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx = \frac{3 \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{8a^3d} + \frac{3 \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{8a^3d} + \frac{\cos(6e - \frac{6cf}{d}) \operatorname{CosIntegral}(\frac{6cf}{d} + 6fx)}{8a^3d} + \frac{\log(c+dx)}{8a^3d} - \frac{i \operatorname{CosIntegral}(\frac{6cf}{d} + 6fx) \sin(6e - \frac{6cf}{d})}{8a^3d} - \frac{3i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{8a^3d} - \frac{3i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{8a^3d} - \frac{3i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{8a^3d} - \frac{3 \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{8a^3d} - \frac{i \cos(6e - \frac{6cf}{d}) \operatorname{Si}(\frac{6cf}{d} + 6fx)}{8a^3d} - \frac{\sin(6e - \frac{6cf}{d}) \operatorname{Si}(\frac{6cf}{d} + 6fx)}{8a^3d}$$

output  $1/8*\operatorname{Ci}(6*c*f/d+6*f*x)*\cos(-6*e+6*c*f/d)/a^3/d+3/8*\operatorname{Ci}(4*c*f/d+4*f*x)*\cos(-4*e+4*c*f/d)/a^3/d+3/8*\operatorname{Ci}(2*c*f/d+2*f*x)*\cos(-2*e+2*c*f/d)/a^3/d+1/8*\ln(d*x+c)/a^3/d-3/8*I*\cos(-2*e+2*c*f/d)*\operatorname{Si}(2*c*f/d+2*f*x)/a^3/d-3/8*I*\cos(-4*e+4*c*f/d)*\operatorname{Si}(4*c*f/d+4*f*x)/a^3/d-1/8*I*\cos(-6*e+6*c*f/d)*\operatorname{Si}(6*c*f/d+6*f*x)/a^3/d+1/8*I*\operatorname{Ci}(6*c*f/d+6*f*x)*\sin(-6*e+6*c*f/d)/a^3/d+1/8*\operatorname{Si}(6*c*f/d+6*f*x)*\sin(-6*e+6*c*f/d)/a^3/d+3/8*I*\operatorname{Ci}(4*c*f/d+4*f*x)*\sin(-4*e+4*c*f/d)/a^3/d+3/8*\operatorname{Si}(4*c*f/d+4*f*x)*\sin(-4*e+4*c*f/d)/a^3/d+3/8*I*\operatorname{Ci}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a^3/d+3/8*\operatorname{Si}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a^3/d$

### 3.32.2 Mathematica [A] (verified)

Time = 1.14 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.75

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$$

$$= \frac{\sec^3(e+fx)(\cos(fx)+i \sin(fx))^3 \left( \cos(3e) \log(f(c+dx)) + i \log(f(c+dx)) \sin(3e) + \left( \cos\left(e - \frac{4cf}{d}\right) - \right. \right.}{\left. \left. \right. \right)}$$

input `Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])^3),x]`

output `(Sec[e + f*x]^3*(Cos[f*x] + I*Sin[f*x])^3*(Cos[3*e]*Log[f*(c + d*x)] + I*Log[f*(c + d*x)]*Sin[3*e] + (Cos[e - (4*c*f)/d] - I*Sin[e - (4*c*f)/d]))*(3*CosIntegral[(4*f*(c + d*x))/d] + Cos[2*e - (2*c*f)/d]*CosIntegral[(6*f*(c + d*x))/d] + 3*CosIntegral[(2*f*(c + d*x))/d]*(Cos[2*e - (2*c*f)/d] + I*Sin[2*e - (2*c*f)/d]) - I*CosIntegral[(6*f*(c + d*x))/d]*Sin[2*e - (2*c*f)/d] - (3*I)*Cos[2*e - (2*c*f)/d]*SinIntegral[(2*f*(c + d*x))/d] + 3*Sin[2*e - (2*c*f)/d]*SinIntegral[(2*f*(c + d*x))/d] - (3*I)*SinIntegral[(4*f*(c + d*x))/d] - I*Cos[2*e - (2*c*f)/d]*SinIntegral[(6*f*(c + d*x))/d] - Sin[2*e - (2*c*f)/d]*SinIntegral[(6*f*(c + d*x))/d]))/(8*d*(a + I*a*Tan[e + f*x])^3)`

### 3.32.3 Rubi [A] (verified)

Time = 1.98 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$$

$$\downarrow \text{4211}$$

$$\int \left( \frac{i \sin^3(2e + 2fx)}{8a^3(c + dx)} - \frac{3 \sin^2(2e + 2fx)}{8a^3(c + dx)} - \frac{3i \sin(2e + 2fx)}{8a^3(c + dx)} - \frac{3 \sin(2e + 2fx) \sin(4e + 4fx)}{16a^3(c + dx)} - \frac{3i \sin(4e + 4fx)}{8a^3(c + dx)} \right) dx$$

↓ 2009

$$\frac{3i \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{8a^3d} - \frac{i \operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \sin\left(6e - \frac{6cf}{d}\right)}{8a^3d} - \frac{3i \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{8a^3d} + \frac{3 \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{8a^3d} + \frac{3 \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{8a^3d} + \frac{\operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \cos\left(6e - \frac{6cf}{d}\right)}{8a^3d} - \frac{3 \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{8a^3d} - \frac{3 \sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{8a^3d} - \frac{\sin\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{8a^3d} - \frac{3i \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{8a^3d} - \frac{3i \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{8a^3d} - \frac{i \cos\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{8a^3d} + \frac{\log(c + dx)}{8a^3d}$$

input `Int[1/((c + d*x)*(a + I*a*Tan[e + f*x])^3),x]`

output `(3*Cos[2*e - (2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x])/(8*a^3*d) + (3*Cos[4*e - (4*c*f)/d]*CosIntegral[(4*c*f)/d + 4*f*x])/(8*a^3*d) + (Cos[6*e - (6*c*f)/d]*CosIntegral[(6*c*f)/d + 6*f*x])/(8*a^3*d) + Log[c + d*x]/(8*a^3*d) - ((I/8)*CosIntegral[(6*c*f)/d + 6*f*x]*Sin[6*e - (6*c*f)/d])/(a^3*d) - (((3*I)/8)*CosIntegral[(4*c*f)/d + 4*f*x]*Sin[4*e - (4*c*f)/d])/(a^3*d) - (((3*I)/8)*CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/(a^3*d) - (((3*I)/8)*Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(a^3*d) - (3*SIN[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(8*a^3*d) - (((3*I)/8)*Cos[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(a^3*d) - (3*Sin[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(8*a^3*d) - ((I/8)*Cos[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(a^3*d) - (Sin[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(8*a^3*d)`

### 3.32.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4211 `Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]`

### 3.32.4 Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.36

method	result
risch	$\frac{\ln(dx+c)}{8a^3d} - \frac{e^{\frac{6i(cf-de)}{d}} \operatorname{Ei}_1\left(\frac{6ifx+6ie+\frac{6i(cf-de)}{d}}{8a^3d}\right)}{8a^3d} - \frac{3e^{\frac{4i(cf-de)}{d}} \operatorname{Ei}_1\left(\frac{4ifx+4ie+\frac{4i(cf-de)}{d}}{8a^3d}\right)}{8a^3d} - \frac{3e^{\frac{2i(cf-de)}{d}} \operatorname{Ei}_1\left(\frac{2ifx+2ie+\frac{2i(cf-de)}{d}}{8a^3d}\right)}{8a^3d}$

input `int(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `1/8*ln(d*x+c)/a^3/d-1/8/a^3/d*exp(6*I*(c*f-d*e)/d)*Ei(1,6*I*f*x+6*I*e+6*I*(c*f-d*e)/d)-3/8/a^3/d*exp(4*I*(c*f-d*e)/d)*Ei(1,4*I*f*x+4*I*e+4*I*(c*f-d*e)/d)-3/8/a^3/d*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)`

### 3.32.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.26

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx = \frac{3 \operatorname{Ei}\left(-\frac{2(i dfx+icf)}{d}\right) e^{\left(-\frac{2(i de-icf)}{d}\right)} + 3 \operatorname{Ei}\left(-\frac{4(i dfx+icf)}{d}\right) e^{\left(-\frac{4(i de-icf)}{d}\right)} + \operatorname{Ei}\left(-\frac{6(i dfx+icf)}{d}\right) e^{\left(-\frac{6(i de-icf)}{d}\right)} + \dots}{8a^3d}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="fracas")`

3.32.  $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$

output  $1/8*(3*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + 3*Ei(-4*(I*d*f*x + I*c*f)/d)*e^(-4*(I*d*e - I*c*f)/d) + Ei(-6*(I*d*f*x + I*c*f)/d)*e^(-6*(I*d*e - I*c*f)/d) + \log((d*x + c)/d)/(a^3*d)$

### 3.32.6 Sympy [F]

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$$

$$= \frac{i \int \frac{1}{c \tan^3(e+fx) - 3ic \tan^2(e+fx) - 3c \tan(e+fx) + ic + dx \tan^3(e+fx) - 3idx \tan^2(e+fx) - 3dx \tan(e+fx) + idx} dx}{a^3}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**3,x)`

output `I*Integral(1/(c*tan(e + f*x)**3 - 3*I*c*tan(e + f*x)**2 - 3*c*tan(e + f*x) + I*c + d*x*tan(e + f*x)**3 - 3*I*d*x*tan(e + f*x)**2 - 3*d*x*tan(e + f*x) + I*d*x), x)/a**3`

### 3.32.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.61

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx =$$

$$\frac{3f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+ide-icf)}{d}\right) + 3f \cos\left(-\frac{4(de-cf)}{d}\right) E_1\left(-\frac{4(-i(fx+e)d+ide-icf)}{d}\right) + f \cos\left(-\frac{6(de-cf)}{d}\right) E_1\left(-\frac{6(-i(fx+e)d+ide-icf)}{d}\right)}{a^3}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `-1/8*(3*f*cos(-2*(d*e - c*f)/d)*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*f*cos(-4*(d*e - c*f)/d)*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f*cos(-6*(d*e - c*f)/d)*exp_integral_e(1, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*I*f*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + 3*I*f*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + I*f*exp_integral_e(1, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-6*(d*e - c*f)/d) - f*log((f*x + e)*d - d*e + c*f)/(a^3*d*f)`

---

3.32.  $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$



**3.32.8 Giac [A] (verification not implemented)**

Time = 0.58 (sec) , antiderivative size = 810, normalized size of antiderivative = 1.80

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx = \text{Too large to display}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

```
output 1/8*(3*cos(2*e)^2*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + cos(2*e)
^3*log(d*x + c) + 6*I*cos(2*e)*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/
d)*sin(2*e) + 3*I*cos(2*e)^2*log(d*x + c)*sin(2*e) - 3*cos(2*c*f/d)*cos_in
tegral(-2*(d*f*x + c*f)/d)*sin(2*e)^2 - 3*cos(2*e)*log(d*x + c)*sin(2*e)^2
- I*log(d*x + c)*sin(2*e)^3 + 3*I*cos(2*e)^2*cos_integral(-2*(d*f*x + c*f
)/d)*sin(2*c*f/d) - 6*cos(2*e)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e)*s
in(2*c*f/d) - 3*I*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e)^2*sin(2*c*f/d)
- 3*I*cos(2*e)^2*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 6*cos(2*e
)*cos(2*c*f/d)*sin(2*e)*sin_integral(2*(d*f*x + c*f)/d) + 3*I*cos(2*c*f/d)
*sin(2*e)^2*sin_integral(2*(d*f*x + c*f)/d) + 3*cos(2*e)^2*sin(2*c*f/d)*si
n_integral(2*(d*f*x + c*f)/d) + 6*I*cos(2*e)*sin(2*e)*sin(2*c*f/d)*sin_int
egral(2*(d*f*x + c*f)/d) - 3*sin(2*e)^2*sin(2*c*f/d)*sin_integral(2*(d*f*x
+ c*f)/d) + 3*cos(2*e)*cos(4*c*f/d)*cos_integral(-4*(d*f*x + c*f)/d) + 3*
I*cos(4*c*f/d)*cos_integral(-4*(d*f*x + c*f)/d)*sin(2*e) + 3*I*cos(2*e)*co
s_integral(-4*(d*f*x + c*f)/d)*sin(4*c*f/d) - 3*cos_integral(-4*(d*f*x + c
*f)/d)*sin(2*e)*sin(4*c*f/d) - 3*I*cos(2*e)*cos(4*c*f/d)*sin_integral(4*(d
*f*x + c*f)/d) + 3*cos(4*c*f/d)*sin(2*e)*sin_integral(4*(d*f*x + c*f)/d) +
3*cos(2*e)*sin(4*c*f/d)*sin_integral(4*(d*f*x + c*f)/d) + 3*I*sin(2*e)*si
n(4*c*f/d)*sin_integral(4*(d*f*x + c*f)/d) + cos(6*c*f/d)*cos_integral(-6*
(d*f*x + c*f)/d) + I*cos_integral(-6*(d*f*x + c*f)/d)*sin(6*c*f/d) - I*...
```

**3.32.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx = \int \frac{1}{(a+a \tan(e+fx) li)^3 (c+dx)} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)),x)`output `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)), x)`

---

3.32.  $\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$

$$\mathbf{3.33} \quad \int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx$$

3.33.1	Optimal result	242
3.33.2	Mathematica [A] (verified)	243
3.33.3	Rubi [A] (verified)	244
3.33.4	Maple [A] (verified)	246
3.33.5	Fricas [A] (verification not implemented)	246
3.33.6	Sympy [F]	247
3.33.7	Maxima [A] (verification not implemented)	247
3.33.8	Giac [B] (verification not implemented)	248
3.33.9	Mupad [F(-1)]	249

### 3.33.1 Optimal result

Integrand size = 23, antiderivative size = 712

$$\begin{aligned}
 \int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx = & -\frac{1}{8a^3d(c+dx)} - \frac{9 \cos(2e+2fx)}{32a^3d(c+dx)} - \frac{3 \cos^2(2e+2fx)}{8a^3d(c+dx)} \\
 & - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \cos(6e+6fx)}{32a^3d(c+dx)} \\
 & - \frac{3if \cos\left(2e - \frac{2cf}{d}\right) \operatorname{CosIntegral}\left(\frac{2cf}{d} + 2fx\right)}{4a^3d^2} \\
 & - \frac{3if \cos\left(4e - \frac{4cf}{d}\right) \operatorname{CosIntegral}\left(\frac{4cf}{d} + 4fx\right)}{2a^3d^2} \\
 & - \frac{3if \cos\left(6e - \frac{6cf}{d}\right) \operatorname{CosIntegral}\left(\frac{6cf}{d} + 6fx\right)}{4a^3d^2} \\
 & - \frac{3f \operatorname{CosIntegral}\left(\frac{6cf}{d} + 6fx\right) \sin\left(6e - \frac{6cf}{d}\right)}{4a^3d^2} \\
 & - \frac{3f \operatorname{CosIntegral}\left(\frac{4cf}{d} + 4fx\right) \sin\left(4e - \frac{4cf}{d}\right)}{2a^3d^2} \\
 & - \frac{3f \operatorname{CosIntegral}\left(\frac{2cf}{d} + 2fx\right) \sin\left(2e - \frac{2cf}{d}\right)}{4a^3d^2} \\
 & + \frac{15i \sin(2e+2fx)}{32a^3d(c+dx)} \\
 & + \frac{3 \sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i \sin^3(2e+2fx)}{8a^3d(c+dx)} \\
 & + \frac{3i \sin(4e+4fx)}{8a^3d(c+dx)} + \frac{3i \sin(6e+6fx)}{32a^3d(c+dx)} \\
 & - \frac{3f \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(\frac{2cf}{d} + 2fx\right)}{4a^3d^2} \\
 & + \frac{3if \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(\frac{2cf}{d} + 2fx\right)}{4a^3d^2} \\
 & - \frac{3f \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(\frac{4cf}{d} + 4fx\right)}{2a^3d^2} \\
 & + \frac{3if \sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(\frac{4cf}{d} + 4fx\right)}{2a^3d^2} \\
 & - \frac{3f \cos\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(\frac{6cf}{d} + 6fx\right)}{4a^3d^2} \\
 & + \frac{3if \sin\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(\frac{6cf}{d} + 6fx\right)}{4a^3d^2}
 \end{aligned}$$

output

```

-1/8/a^3/d/(d*x+c)-3/4*I*f*Si(6*c*f/d+6*f*x)*sin(-6*e+6*c*f/d)/a^3/d^2+15/
32*I*sin(2*f*x+2*e)/a^3/d/(d*x+c)-1/8*I*sin(2*f*x+2*e)^3/a^3/d/(d*x+c)-9/3
2*cos(2*f*x+2*e)/a^3/d/(d*x+c)-3/8*cos(2*f*x+2*e)^2/a^3/d/(d*x+c)-1/8*cos(
2*f*x+2*e)^3/a^3/d/(d*x+c)-3/32*cos(6*f*x+6*e)/a^3/d/(d*x+c)-3/4*f*cos(-2*
e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^3/d^2-3/2*f*cos(-4*e+4*c*f/d)*Si(4*c*f/d+4*
f*x)/a^3/d^2-3/4*f*cos(-6*e+6*c*f/d)*Si(6*c*f/d+6*f*x)/a^3/d^2+3/4*f*Ci(6*
c*f/d+6*f*x)*sin(-6*e+6*c*f/d)/a^3/d^2-3/2*I*f*Si(4*c*f/d+4*f*x)*sin(-4*e+
4*c*f/d)/a^3/d^2+3/2*f*Ci(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a^3/d^2+3/32*I*
sin(6*f*x+6*e)/a^3/d/(d*x+c)+3/4*f*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^3
/d^2-3/2*I*f*Ci(4*c*f/d+4*f*x)*cos(-4*e+4*c*f/d)/a^3/d^2-3/4*I*f*Ci(6*c*f/
d+6*f*x)*cos(-6*e+6*c*f/d)/a^3/d^2+3/8*sin(2*f*x+2*e)^2/a^3/d/(d*x+c)+3/8*
I*sin(4*f*x+4*e)/a^3/d/(d*x+c)-3/4*I*f*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)
/a^3/d^2-3/4*I*f*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^3/d^2

```

### 3.33.2 Mathematica [A] (verified)

Time = 2.85 (sec) , antiderivative size = 833, normalized size of antiderivative = 1.17

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx$$

$$= \frac{\sec^3(e+fx) \left(-i \cos\left(\frac{3cf}{d}\right) + \sin\left(\frac{3cf}{d}\right)\right) \left(3d \cos\left(e+f\left(-\frac{3c}{d}+x\right)\right) + d \cos\left(3\left(e+f\left(-\frac{c}{d}+x\right)\right)\right)\right) + d \cos\left(3\left(e+f\left(-\frac{c}{d}+x\right)\right)\right)}{\dots}$$

input `Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^3),x]`

```
output (Sec[e + f*x]^3*((-I)*Cos[(3*c*f)/d] + Sin[(3*c*f)/d])*(3*d*Cos[e + f*((-3
*c)/d + x)] + d*Cos[3*(e + f*(-(c/d) + x))] + d*Cos[3*(e + f*(c/d + x))] +
3*d*Cos[e + f*((3*c)/d + x)] + (6*I)*c*f*Cos[3*e - (3*f*(c + d*x))/d]*Cos
Integral[(6*f*(c + d*x))/d] + (6*I)*d*f*x*Cos[3*e - (3*f*(c + d*x))/d]*Cos
Integral[(6*f*(c + d*x))/d] + (6*I)*f*(c + d*x)*CosIntegral[(2*f*(c + d*x)
)/d]*(Cos[e - (c*f)/d + 3*f*x] + I*Sin[e - (c*f)/d + 3*f*x]) + (3*I)*d*Sin
[e + f*((-3*c)/d + x)] + I*d*Sin[3*(e + f*(-(c/d) + x))] - I*d*Sin[3*(e +
f*(c/d + x))] - (3*I)*d*Sin[e + f*((3*c)/d + x)] + 6*c*f*CosIntegral[(6*f*
(c + d*x))/d]*Sin[3*e - (3*f*(c + d*x))/d] + 6*d*f*x*CosIntegral[(6*f*(c +
d*x))/d]*Sin[3*e - (3*f*(c + d*x))/d] + 12*f*(c + d*x)*CosIntegral[(4*f*(
c + d*x))/d]*(I*Cos[e - (f*(c + 3*d*x))/d] + Sin[e - (f*(c + 3*d*x))/d]) +
6*c*f*Cos[e - (c*f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + 6*d*f*x*C
os[e - (c*f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + (6*I)*c*f*Sin[e -
(c*f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + (6*I)*d*f*x*Sin[e - (c*
f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + 12*c*f*Cos[e - (f*(c + 3*d*
x))/d]*SinIntegral[(4*f*(c + d*x))/d] + 12*d*f*x*Cos[e - (f*(c + 3*d*x))/d
]*SinIntegral[(4*f*(c + d*x))/d] - (12*I)*c*f*Sin[e - (f*(c + 3*d*x))/d]*S
inIntegral[(4*f*(c + d*x))/d] - (12*I)*d*f*x*Sin[e - (f*(c + 3*d*x))/d]*Si
nIntegral[(4*f*(c + d*x))/d] + 6*c*f*Cos[3*e - (3*f*(c + d*x))/d]*SinIntegr
al[(6*f*(c + d*x))/d] + 6*d*f*x*Cos[3*e - (3*f*(c + d*x))/d]*SinIntegr...
```

### 3.33.3 Rubi [A] (verified)

Time = 2.06 (sec) , antiderivative size = 712, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^3} dx$$

↓ 4211

$$\int \left( \frac{i \sin^3(2e + 2fx)}{8a^3(c + dx)^2} - \frac{3 \sin^2(2e + 2fx)}{8a^3(c + dx)^2} - \frac{3i \sin(2e + 2fx)}{8a^3(c + dx)^2} - \frac{3 \sin(2e + 2fx) \sin(4e + 4fx)}{16a^3(c + dx)^2} - \frac{3i \sin(4e + 4fx)}{8a^3(c + dx)^2} \right) dx$$

↓ 2009

---

3.33.  $\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx$

$$\begin{aligned}
& \frac{3f \operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \sin\left(6e - \frac{6cf}{d}\right)}{4a^3d^2} - \frac{3f \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{2a^3d^2} - \\
& \frac{3f \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{4a^3d^2} - \frac{3if \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{4a^3d^2} - \\
& \frac{3if \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{2a^3d^2} - \frac{3if \operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \cos\left(6e - \frac{6cf}{d}\right)}{4a^3d^2} + \\
& \frac{3if \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{4a^3d^2} + \frac{3if \sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{2a^3d^2} + \\
& \frac{3if \sin\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{4a^3d^2} - \frac{3f \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{4a^3d^2} - \\
& \frac{3f \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{2a^3d^2} - \frac{3f \cos\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{4a^3d^2} - \frac{i \sin^3(2e + 2fx)}{8a^3d(c + dx)} + \\
& \frac{3 \sin^2(2e + 2fx)}{8a^3d(c + dx)} + \frac{15i \sin(2e + 2fx)}{32a^3d(c + dx)} + \frac{3i \sin(4e + 4fx)}{8a^3d(c + dx)} + \frac{3i \sin(6e + 6fx)}{32a^3d(c + dx)} - \frac{\cos^3(2e + 2fx)}{8a^3d(c + dx)} - \\
& \frac{3 \cos^2(2e + 2fx)}{8a^3d(c + dx)} - \frac{9 \cos(2e + 2fx)}{32a^3d(c + dx)} - \frac{3 \cos(6e + 6fx)}{32a^3d(c + dx)} - \frac{1}{8a^3d(c + dx)}
\end{aligned}$$

input `Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^3),x]`

output

```

-1/8*1/(a^3*d*(c + d*x)) - (9*Cos[2*e + 2*f*x])/(32*a^3*d*(c + d*x)) - (3*
Cos[2*e + 2*f*x]^2)/(8*a^3*d*(c + d*x)) - Cos[2*e + 2*f*x]^3/(8*a^3*d*(c +
d*x)) - (3*Cos[6*e + 6*f*x])/(32*a^3*d*(c + d*x)) - (((3*I)/4)*f*Cos[2*e
- (2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x])/(a^3*d^2) - (((3*I)/2)*f*Cos[
4*e - (4*c*f)/d]*CosIntegral[(4*c*f)/d + 4*f*x])/(a^3*d^2) - (((3*I)/4)*f*
Cos[6*e - (6*c*f)/d]*CosIntegral[(6*c*f)/d + 6*f*x])/(a^3*d^2) - (3*f*CosI
ntegral[(6*c*f)/d + 6*f*x]*Sin[6*e - (6*c*f)/d])/(4*a^3*d^2) - (3*f*CosInt
egral[(4*c*f)/d + 4*f*x]*Sin[4*e - (4*c*f)/d])/(2*a^3*d^2) - (3*f*CosInteg
ral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/(4*a^3*d^2) + (((15*I)/32)*Si
n[2*e + 2*f*x])/(a^3*d*(c + d*x)) + (3*Sin[2*e + 2*f*x]^2)/(8*a^3*d*(c + d
*x)) - ((I/8)*Sin[2*e + 2*f*x]^3)/(a^3*d*(c + d*x)) + (((3*I)/8)*Sin[4*e +
4*f*x])/(a^3*d*(c + d*x)) + (((3*I)/32)*Sin[6*e + 6*f*x])/(a^3*d*(c + d*x
)) - (3*f*Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(4*a^3*d^2)
+ (((3*I)/4)*f*Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(a^3*
d^2) - (3*f*Cos[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(2*a^3*d^
2) + (((3*I)/2)*f*Sin[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(a^
3*d^2) - (3*f*Cos[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(4*a^3*
d^2) + (((3*I)/4)*f*Sin[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(
a^3*d^2)

```

### 3.33.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4211 `Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]`

### 3.33.4 Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.36

method	result
risch	$-\frac{1}{8a^3d(dx+c)} - \frac{fe^{-6i(fx+e)}}{8a^3(dfx+cf)d} + \frac{3ife^{\frac{6i(cf-de)}{d}} \operatorname{Ei}_1\left(\frac{6ifx+6ie+\frac{6i(cf-de)}{d}}{d}\right)}{4a^3d^2} - \frac{3fe^{-4i(fx+e)}}{8a^3(dfx+cf)d} + \frac{3ife^{\frac{4i(cf-de)}{d}} \operatorname{Ei}_1\left(\frac{4ifx+4ie+\frac{4i(cf-de)}{d}}{d}\right)}{2a^3d^2}$

input `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output 
$$-\frac{1}{8/a^3/d/(d*x+c)} - \frac{1}{8/a^3*f*\exp(-6*I*(f*x+e))/(d*f*x+c*f)/d} + \frac{3}{4*I/a^3*f/d^2*\exp(6*I*(c*f-d*e)/d)*\operatorname{Ei}\left(1, \frac{6*I*f*x+6*I*e+6*I*(c*f-d*e)}{d}\right)} - \frac{3}{8/a^3*f*\exp(-4*I*(f*x+e))/(d*f*x+c*f)/d} + \frac{3}{2*I/a^3*f/d^2*\exp(4*I*(c*f-d*e)/d)*\operatorname{Ei}\left(1, \frac{4*I*f*x+4*I*e+4*I*(c*f-d*e)}{d}\right)} - \frac{3}{8/a^3*f*\exp(-2*I*(f*x+e))/(d*f*x+c*f)/d} + \frac{3}{4*I/a^3*f/d^2*\exp(2*I*(c*f-d*e)/d)*\operatorname{Ei}\left(1, \frac{2*I*f*x+2*I*e+2*I*(c*f-d*e)}{d}\right)}$$

### 3.33.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.28

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx = \frac{\left( \left( 6(i dfx + i cf) \operatorname{Ei}\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + 12(i dfx + i cf) \operatorname{Ei}\left(-\frac{4(i dfx + i cf)}{d}\right) e^{\left(-\frac{4(i de - i cf)}{d}\right)} + 6(i dfx + i cf) \operatorname{Ei}\left(-\frac{6(i dfx + i cf)}{d}\right) e^{\left(-\frac{6(i de - i cf)}{d}\right)} \right)}{8(a^3c)}$$

3.33. 
$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output 
$$\begin{aligned} & -1/8*((6*(I*d*f*x + I*c*f)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^{(-2*(I*d*e - I*c*f)/d)} + 12*(I*d*f*x + I*c*f)*Ei(-4*(I*d*f*x + I*c*f)/d)*e^{(-4*(I*d*e - I*c*f)/d)} + 6*(I*d*f*x + I*c*f)*Ei(-6*(I*d*f*x + I*c*f)/d)*e^{(-6*(I*d*e - I*c*f)/d)} + d)*e^{(6*I*f*x + 6*I*e)} + 3*d*e^{(4*I*f*x + 4*I*e)} + 3*d*e^{(2*I*f*x + 2*I*e)} + d)*e^{(-6*I*f*x - 6*I*e)}/(a^3*d^3*x + a^3*c*d^2) \end{aligned}$$

### 3.33.6 Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx = \frac{i \int \frac{1}{c^2 \tan^3(e+fx) - 3ic^2 \tan^2(e+fx) - 3c^2 \tan(e+fx) + ic^2 + 2cdx \tan^3(e+fx) - 6icdx \tan^2(e+fx) - 6cdx \tan(e+fx) + 2icdx + d^2x^2 \tan^3(e+fx)}}{a^3} dx$$

input `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e))**3,x)`

output `I*Integral(1/(c**2*tan(e + f*x)**3 - 3*I*c**2*tan(e + f*x)**2 - 3*c**2*tan(e + f*x) + I*c**2 + 2*c*d*x*tan(e + f*x)**3 - 6*I*c*d*x*tan(e + f*x)**2 - 6*c*d*x*tan(e + f*x) + 2*I*c*d*x + d**2*x**2*tan(e + f*x)**3 - 3*I*d**2*x**2*tan(e + f*x)**2 - 3*d**2*x**2*tan(e + f*x) + I*d**2*x**2), x)/a**3`

### 3.33.7 Maxima [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.42

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx = \frac{3 f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+i de-icf)}{d}\right) + 3 f^2 \cos\left(-\frac{4(de-cf)}{d}\right) E_2\left(-\frac{4(-i(fx+e)d+i de-icf)}{d}\right) + f^2 \cos\left(-\frac{6(de-cf)}{d}\right) E_2\left(-\frac{6(-i(fx+e)d+i de-icf)}{d}\right)}{a^3 d^3}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`



```
output -1/8*(3*f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I
*d*e - I*c*f)/d) + 3*f^2*cos(-4*(d*e - c*f)/d)*exp_integral_e(2, -4*(-I*(f
*x + e)*d + I*d*e - I*c*f)/d) + f^2*cos(-6*(d*e - c*f)/d)*exp_integral_e(2
, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*I*f^2*exp_integral_e(2, -2*(-
I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + 3*I*f^2*exp_inte
gral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + I
*f^2*exp_integral_e(2, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-6*(d*e
- c*f)/d) + f^2)/(((f*x + e)*a^3*d^2 - a^3*d^2*e + a^3*c*d*f)*f)
```

### 3.33.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2915 vs.  $2(648) = 1296$ .

Time = 30.48 (sec) , antiderivative size = 2915, normalized size of antiderivative = 4.09

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx = \text{Too large to display}$$

```
input integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
```

```
output 1/8*(-6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-6*(d*e -
c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*
e + c*f)/d) + 6*I*d*e*f^2*cos(-6*(d*e - c*f)/d)*cos_integral(6*((d*x + c)*
(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*cos(-6*(d*
e - c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)
- d*e + c*f)/d) - 12*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*c
os(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x +
c) + f) - d*e + c*f)/d) + 12*I*d*e*f^2*cos(-4*(d*e - c*f)/d)*cos_integral
(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*I*c
*f^3*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/
(d*x + c) + f) - d*e + c*f)/d) - 6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x +
c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c
) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*I*d*e*f^2*cos(-2*(d*e - c*f)/d)
*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f
)/d) - 6*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x
+ c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*(d*x + c)*(d*e/(d*x + c) -
c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x
+ c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 6*d*e*f^2*cos_integral(
2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d
*e - c*f)/d) + 6*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(...
```

**3.33.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx = \int \frac{1}{(a+a \tan(e+fx) 1i)^3 (c+dx)^2} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)^2),x)`output `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)^2), x)`

### 3.34 $\int (c + dx)^m (a + ia \tan(e + fx))^2 dx$

3.34.1	Optimal result	250
3.34.2	Mathematica [N/A]	250
3.34.3	Rubi [N/A]	251
3.34.4	Maple [N/A] (verified)	252
3.34.5	Fricas [N/A]	252
3.34.6	Sympy [N/A]	252
3.34.7	Maxima [N/A]	253
3.34.8	Giac [N/A]	253
3.34.9	Mupad [N/A]	254

#### 3.34.1 Optimal result

Integrand size = 23, antiderivative size = 23

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \text{Int}((c + dx)^m (a + ia \tan(e + fx))^2, x)$$

output `Unintegrable((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x)`

#### 3.34.2 Mathematica [N/A]

Not integrable

Time = 42.84 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

input `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2,x]`

output `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2, x]`

**3.34.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

↓ 3042

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

↓ 4223

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

input `Int[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2,x]`

output `$Aborted`

**3.34.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.34.4 Maple [N/A] (verified)**

Not integrable

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int (dx + c)^m (a + ia \tan(fx + e))^2 dx$$

input `int((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x)`output `int((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x)`**3.34.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 5.00

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (ia \tan(fx + e) + a)^2 (dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`output `(-2*I*(d*x + c)^m*a^2 + (f*e^(2*I*f*x + 2*I*e) + f)*integral(-2*(-I*a^2*d*m - 2*(a^2*d*f*x + a^2*c*f)*e^(2*I*f*x + 2*I*e))*(d*x + c)^m/(d*f*x + c*f + (d*f*x + c*f)*e^(2*I*f*x + 2*I*e)), x))/(f*e^(2*I*f*x + 2*I*e) + f)`**3.34.6 Sympy [N/A]**

Not integrable

Time = 5.32 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.22

$$\begin{aligned} \int (c + dx)^m (a + ia \tan(e + fx))^2 dx = & -a^2 \left( \int (c + dx)^m \tan^2(e + fx) dx \right. \\ & + \int (-2i(c + dx)^m \tan(e + fx)) dx \\ & \left. + \int (-(c + dx)^m) dx \right) \end{aligned}$$

input `integrate((d*x+c)**m*(a+I*a*tan(f*x+e))**2,x)`

output `-a**2*(Integral((c + d*x)**m*tan(e + f*x)**2, x) + Integral(-2*I*(c + d*x)**m*tan(e + f*x), x) + Integral(-(c + d*x)**m, x))`

### 3.34.7 Maxima [N/A]

Not integrable

Time = 1.83 (sec) , antiderivative size = 520, normalized size of antiderivative = 22.61

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (ia \tan(fx + e) + a)^2 (dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `(d*x + c)^(m + 1)*a^2/(d*(m + 1)) + integrate((3*(d*x + c)^m*a^2*cos(4*f*x + 4*e)^2 - 4*(d*x + c)^m*a^2*cos(2*f*x + 2*e)^2 + 3*(d*x + c)^m*a^2*sin(4*f*x + 4*e)^2 + 4*(d*x + c)^m*a^2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) - 4*(d*x + c)^m*a^2*sin(2*f*x + 2*e)^2 - 4*(d*x + c)^m*a^2*cos(2*f*x + 2*e) - (d*x + c)^m*a^2 + 2*(2*(d*x + c)^m*a^2*cos(2*f*x + 2*e) + (d*x + c)^m*a^2)*cos(4*f*x + 4*e))/(2*(2*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 4*cos(2*f*x + 2*e)^2 + sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) + 1), x) + I*integrate(-4*(2*(d*x + c)^m*a^2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e) - (2*(d*x + c)^m*a^2*cos(2*f*x + 2*e) + (d*x + c)^m*a^2)*sin(4*f*x + 4*e))/(2*(2*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 4*cos(2*f*x + 2*e)^2 + sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) + 1), x)`

### 3.34.8 Giac [N/A]

Not integrable

Time = 0.64 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (ia \tan(fx + e) + a)^2 (dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((I*a*tan(f*x + e) + a)^2*(d*x + c)^m, x)`

### 3.34.9 Mupad [N/A]

Not integrable

Time = 3.37 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (a + a \tan(e + fx) i)^2 (c + dx)^m dx$$

input `int((a + a*tan(e + f*x)*1i)^2*(c + d*x)^m,x)`

output `int((a + a*tan(e + f*x)*1i)^2*(c + d*x)^m, x)`

### 3.35 $\int (c + dx)^m (a + ia \tan(e + fx)) dx$

3.35.1	Optimal result	255
3.35.2	Mathematica [N/A]	255
3.35.3	Rubi [N/A]	256
3.35.4	Maple [N/A] (verified)	257
3.35.5	Fricas [N/A]	257
3.35.6	Sympy [N/A]	257
3.35.7	Maxima [N/A]	258
3.35.8	Giac [N/A]	258
3.35.9	Mupad [N/A]	258

#### 3.35.1 Optimal result

Integrand size = 21, antiderivative size = 21

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \text{Int}((c + dx)^m (a + ia \tan(e + fx)), x)$$

output `Unintegrable((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`

#### 3.35.2 Mathematica [N/A]

Not integrable

Time = 9.86 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (c + dx)^m (a + ia \tan(e + fx)) dx$$

input `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x]),x]`

output `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x]), x]`



### 3.35.3 Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx$$

↓ 3042

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx$$

↓ 4223

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx$$

input `Int[(c + d*x)^m*(a + I*a*Tan[e + f*x]),x]`

output `$Aborted`

#### 3.35.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.35.4 Maple [N/A] (verified)**

Not integrable

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int (dx + c)^m (a + ia \tan(fx + e)) dx$$

input `int((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`output `int((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`**3.35.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="fricas")`output `integral(2*(d*x + c)^m*a*e^(2*I*f*x + 2*I*e)/(e^(2*I*f*x + 2*I*e) + 1), x)`**3.35.6 Sympy [N/A]**

Not integrable

Time = 2.40 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.38

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = ia \left( \int (-i(c + dx)^m) dx + \int (c + dx)^m \tan(e + fx) dx \right)$$

input `integrate((d*x+c)**m*(a+I*a*tan(f*x+e)),x)`output `I*a*(Integral(-I*(c + d*x)**m, x) + Integral((c + d*x)**m*tan(e + f*x), x))`

**3.35.7 Maxima [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 79, normalized size of antiderivative = 3.76

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

```
input integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="maxima")
```

```
output 2*I*a*integrate((d*x + c)^m*sin(2*f*x + 2*e)/(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1), x) + (d*x + c)^(m + 1)*a/(d*(m + 1))
```

**3.35.8 Giac [N/A]**

Not integrable

Time = 0.36 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

```
input integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="giac")
```

```
output integrate((I*a*tan(f*x + e) + a)*(d*x + c)^m, x)
```

**3.35.9 Mupad [N/A]**

Not integrable

Time = 3.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (a + a \tan(e + fx) li) (c + dx)^m dx$$

```
input int((a + a*tan(e + f*x)*1i)*(c + d*x)^m,x)
```

```
output int((a + a*tan(e + f*x)*1i)*(c + d*x)^m, x)
```

### 3.36 $\int \frac{(c+dx)^m}{a+ia \tan(e+fx)} dx$

3.36.1	Optimal result	259
3.36.2	Mathematica [A] (verified)	259
3.36.3	Rubi [A] (verified)	260
3.36.4	Maple [F]	261
3.36.5	Fricas [A] (verification not implemented)	261
3.36.6	Sympy [F]	262
3.36.7	Maxima [F]	262
3.36.8	Giac [F]	262
3.36.9	Mupad [F(-1)]	263

#### 3.36.1 Optimal result

Integrand size = 23, antiderivative size = 98

$$\int \frac{(c+dx)^m}{a+ia \tan(e+fx)} dx = \frac{(c+dx)^{1+m}}{2ad(1+m)} + \frac{i2^{-2-m}e^{-2i(e-\frac{cf}{d})}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{af}$$

output `1/2*(d*x+c)^(1+m)/a/d/(1+m)+I*2^(-2-m)*(d*x+c)^m*GAMMA(1+m,2*I*f*(d*x+c)/d)/a/exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)`

#### 3.36.2 Mathematica [A] (verified)

Time = 1.77 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.36

$$\int \frac{(c+dx)^m}{a+ia \tan(e+fx)} dx = \frac{e^{-ie}(c+dx)^m \left(\frac{2e^{2ie}f(c+dx)}{d(1+m)} + i2^{-m}e^{\frac{2icf}{d}} \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)\right) \sec(e+fx)(\cos(fx) + i \sin(fx))}{4f(a+ia \tan(e+fx))}$$

input `Integrate[(c + d*x)^m/(a + I*a*Tan[e + f*x]),x]`

output  $((c + dx)^m * ((2 * E^{((2 * I) * e) * f * (c + dx)) / (d * (1 + m)) + (I * E^{((2 * I) * c * f) / d}) * \text{Gamma}[1 + m, ((2 * I) * f * (c + dx)) / d]) / (2^m * ((I * f * (c + dx)) / d)^m)) * \text{Sec}[e + f * x] * (\text{Cos}[f * x] + I * \text{Sin}[f * x]) / (4 * E^{(I * e) * f * (a + I * a * \text{Tan}[e + f * x])})$

### 3.36.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4210, 2612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{4210} \\ & \frac{(c + dx)^{m+1}}{2ad(m + 1)} + \frac{\int e^{-2i(e+fx)} (c + dx)^m dx}{2a} \\ & \quad \downarrow \text{2612} \\ & \frac{(c + dx)^{m+1}}{2ad(m + 1)} + \frac{i 2^{-m-2} e^{-2i\left(e - \frac{cf}{d}\right)} (c + dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m + 1, \frac{2if(c+dx)}{d}\right)}{af} \end{aligned}$$

input  $\text{Int}[(c + dx)^m / (a + I * a * \text{Tan}[e + f * x]), x]$

output  $(c + dx)^{(1 + m)} / (2 * a * d * (1 + m)) + (I * 2^{(-2 - m)} * (c + dx)^m * \text{Gamma}[1 + m, ((2 * I) * f * (c + dx)) / d]) / (a * E^{((2 * I) * (e - (c * f) / d))} * f * ((I * f * (c + dx)) / d)^m)$

## 3.36.3.1 Defintions of rubi rules used

```
rule 2612 Int[(F_)^((g_)*((e_) + (f_)*(x_)))*((c_) + (d_)*(x_))^(m_), x_Symbol]
:> Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*((-f)*g*(Log[F]/d)
)^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m]))*Gamma[m + 1,
((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4210 Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Sym
bol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + Simp[1/(2*a) Int[(c
+ d*x)^m*E^(2*(a/b)*(e + f*x)), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] &
& EqQ[a^2 + b^2, 0] && !IntegerQ[m]
```

## 3.36.4 Maple [F]

$$\int \frac{(dx + c)^m}{a + ia \tan(fx + e)} dx$$

```
input int((d*x+c)^m/(a+I*a*tan(f*x+e)),x)
```

```
output int((d*x+c)^m/(a+I*a*tan(f*x+e)),x)
```

## 3.36.5 Fracas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.87

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx$$

$$= \frac{(i dm + i d) e^{\left(-\frac{dm \log\left(\frac{2if}{d}\right) + 2ide - 2icf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idf x - icf)}{d}\right) + 2(df x + cf)(dx + c)^m}{4(adfm + adf)}$$

```
input integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="fracas")
```

output `1/4*((I*d*m + I*d)*e^(-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d)*gamma(m + 1, -2*(-I*d*f*x - I*c*f)/d) + 2*(d*f*x + c*f)*(d*x + c)^m)/(a*d*f*m + a*d*f)`

### 3.36.6 Sympy [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = -\frac{i \int \frac{(c+dx)^m}{\tan(e+fx)-i} dx}{a}$$

input `integrate((d*x+c)**m/(a+I*a*tan(f*x+e)),x)`

output `-I*Integral((c + d*x)**m/(tan(e + f*x) - I), x)/a`

### 3.36.7 Maxima [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(dx + c)^m}{i a \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `1/2*((d*m + d)*integrate((d*x + c)^m*cos(2*f*x + 2*e), x) - (I*d*m + I*d)*integrate((d*x + c)^m*sin(2*f*x + 2*e), x) + e^(m*log(d*x + c) + log(d*x + c)))/(a*d*m + a*d)`

### 3.36.8 Giac [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(dx + c)^m}{i a \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^m/(I*a*tan(f*x + e) + a), x)`

**3.36.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(c + dx)^m}{a + a \tan(e + fx)} \operatorname{li} dx$$

input `int((c + d*x)^m/(a + a*tan(e + f*x)*1i),x)`output `int((c + d*x)^m/(a + a*tan(e + f*x)*1i), x)`



### 3.37 $\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^2} dx$

3.37.1	Optimal result	264
3.37.2	Mathematica [A] (verified)	264
3.37.3	Rubi [A] (verified)	265
3.37.4	Maple [F]	266
3.37.5	Fricas [A] (verification not implemented)	266
3.37.6	Sympy [F]	267
3.37.7	Maxima [F]	267
3.37.8	Giac [F]	268
3.37.9	Mupad [F(-1)]	268

#### 3.37.1 Optimal result

Integrand size = 23, antiderivative size = 171

$$\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^2} dx$$

$$= \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{i2^{-2-m}e^{-2i\left(\frac{e-cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^2f}$$

$$+ \frac{i4^{-2-m}e^{-4i\left(\frac{e-cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^2f}$$

output  $\frac{1}{4}*(d*x+c)^{(1+m)}/a^2/d/(1+m)+I*2^{(-2-m)}*(d*x+c)^m*\text{GAMMA}(1+m,2*I*f*(d*x+c)/d)/a^2/\exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+I*4^{(-2-m)}*(d*x+c)^m*\text{GAMMA}(1+m,4*I*f*(d*x+c)/d)/a^2/\exp(4*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)$

#### 3.37.2 Mathematica [A] (verified)

Time = 4.49 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.12

$$\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^2} dx$$

$$= \frac{(c+dx)^m \left( \frac{4e^{2ie} f(c+dx)}{d(1+m)} + i2^{2-m} e^{\frac{2icf}{d}} \left( \frac{if(c+dx)}{d} \right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right) + i4^{-m} e^{-2ie+\frac{4icf}{d}} \left( \frac{if(c+dx)}{d} \right)^{-m} \Gamma\left(1+m, \frac{4if(c+dx)}{d}\right) \right)}{16f(a+ia \tan(e+fx))^2}$$

input `Integrate[(c + d*x)^m/(a + I*a*Tan[e + f*x])^2,x]`

output `((c + d*x)^m*((4*E^((2*I)*e))*f*(c + d*x))/(d*(1 + m)) + (I*2^(2 - m)*E^(((2*I)*c*f)/d)*Gamma[1 + m, ((2*I)*f*(c + d*x))/d])/((I*f*(c + d*x))/d)^m + (I*E^((-2*I)*e + ((4*I)*c*f)/d)*Gamma[1 + m, ((4*I)*f*(c + d*x))/d])/(4^m*((I*f*(c + d*x))/d)^m)*Sec[e + f*x]^2*(Cos[f*x] + I*Sin[f*x])^2/(16*f*(a + I*a*Tan[e + f*x])^2)`

### 3.37.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4212} \\
 & \int \left( \frac{e^{-2ie-2ifx}(c + dx)^m}{2a^2} + \frac{e^{-4ie-4ifx}(c + dx)^m}{4a^2} + \frac{(c + dx)^m}{4a^2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{i2^{-m-2}e^{-2i\left(e-\frac{cf}{d}\right)}(c + dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m + 1, \frac{2if(c+dx)}{d}\right)}{a^2 f} + \\
 & \frac{i4^{-m-2}e^{-4i\left(e-\frac{cf}{d}\right)}(c + dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m + 1, \frac{4if(c+dx)}{d}\right)}{a^2 f} + \frac{(c + dx)^{m+1}}{4a^2 d(m + 1)}
 \end{aligned}$$

input `Int[(c + d*x)^m/(a + I*a*Tan[e + f*x])^2,x]`

```
output (c + d*x)^(1 + m)/(4*a^2*d*(1 + m)) + (I*2^(-2 - m)*(c + d*x)^m*Gamma[1 +
m, ((2*I)*f*(c + d*x))/d])/(a^2*E^((2*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))
/d)^m) + (I*4^(-2 - m)*(c + d*x)^m*Gamma[1 + m, ((4*I)*f*(c + d*x))/d])/(a
^2*E^((4*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m)
```

### 3.37.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4212 Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*
x)))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2
, 0] && ILtQ[n, 0]
```

### 3.37.4 Maple [F]

$$\int \frac{(dx + c)^m}{(a + ia \tan(fx + e))^2} dx$$

```
input int((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x)
```

```
output int((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x)
```

### 3.37.5 Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.84

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx =$$

$$\frac{4(-i dm - i d)e^{\left(-\frac{dm \log\left(\frac{2if}{d}\right) + 2ide - 2icf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idf x - icf)}{d}\right) - (idm + id)e^{\left(-\frac{dm \log\left(\frac{4if}{d}\right) + 4ide - 4icf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idf x - icf)}{d}\right)}{16(a^2dfm + a^2df)}$$

---

3.37.  $\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^2} dx$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output `-1/16*(4*(-I*d*m - I*d)*e^(-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d)*gamma(m + 1, -2*(-I*d*f*x - I*c*f)/d) - (I*d*m + I*d)*e^(-(d*m*log(4*I*f/d) + 4*I*d*e - 4*I*c*f)/d)*gamma(m + 1, -4*(-I*d*f*x - I*c*f)/d) - 4*(d*f*x + c*f)*(d*x + c)^m/(a^2*d*f*m + a^2*d*f)`

### 3.37.6 Sympy [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = -\frac{\int \frac{(c+dx)^m}{\tan^2(e+fx) - 2i \tan(e+fx) - 1} dx}{a^2}$$

input `integrate((d*x+c)**m/(a+I*a*tan(f*x+e))**2,x)`

output `-Integral((c + d*x)**m/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x)/a**2`

### 3.37.7 Maxima [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(dx + c)^m}{(ia \tan(fx + e) + a)^2} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `1/4*((d*m + d)*integrate((d*x + c)^m*cos(4*f*x + 4*e), x) + 2*(d*m + d)*integrate((d*x + c)^m*cos(2*f*x + 2*e), x) - (I*d*m + I*d)*integrate((d*x + c)^m*sin(4*f*x + 4*e), x) + 2*(-I*d*m - I*d)*integrate((d*x + c)^m*sin(2*f*x + 2*e), x) + e^(m*log(d*x + c) + log(d*x + c))/(a^2*d*m + a^2*d)`

**3.37.8 Giac [F]**

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(dx + c)^m}{(ia \tan(fx + e) + a)^2} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)^m/(I*a*tan(f*x + e) + a)^2, x)`

**3.37.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(c + dx)^m}{(a + a \tan(e + fx) li)^2} dx$$

input `int((c + d*x)^m/(a + a*tan(e + f*x)*li)^2,x)`

output `int((c + d*x)^m/(a + a*tan(e + f*x)*li)^2, x)`

### 3.38 $\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^3} dx$

3.38.1	Optimal result	269
3.38.2	Mathematica [A] (verified)	270
3.38.3	Rubi [A] (verified)	270
3.38.4	Maple [F]	272
3.38.5	Fricas [A] (verification not implemented)	272
3.38.6	Sympy [F]	272
3.38.7	Maxima [F]	273
3.38.8	Giac [F]	273
3.38.9	Mupad [F(-1)]	273

#### 3.38.1 Optimal result

Integrand size = 23, antiderivative size = 251

$$\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^3} dx$$

$$= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} + \frac{3i2^{-4-m}e^{-2i(e-\frac{cf}{d})}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^3f}$$

$$+ \frac{3i2^{-5-2m}e^{-4i(e-\frac{cf}{d})}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^3f}$$

$$+ \frac{i2^{-4-m}3^{-1-m}e^{-6i(e-\frac{cf}{d})}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{6if(c+dx)}{d}\right)}{a^3f}$$

output

```
1/8*(d*x+c)^(1+m)/a^3/d/(1+m)+3*I*2^(-4-m)*(d*x+c)^m*GAMMA(1+m,2*I*f*(d*x+c)/d)/a^3/exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+3*I*2^(-5-2*m)*(d*x+c)^m*GAMMA(1+m,4*I*f*(d*x+c)/d)/a^3/exp(4*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+I*2^(-4-m)*3^(-1-m)*(d*x+c)^m*GAMMA(1+m,6*I*f*(d*x+c)/d)/a^3/exp(6*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)
```

### 3.38.2 Mathematica [A] (verified)

Time = 13.03 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{2^{-5-2m} 3^{-1-m} e^{-3ie} (c + dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \left(12^{1+m} e^{6ie} f(c + dx) \left(\frac{if(c+dx)}{d}\right)^m + i2^{1+m} 3^{2+m} d e^{2i\left(2e + \frac{cf}{d}\right)} (1 + \right.$$

input `Integrate[(c + d*x)^m/(a + I*a*Tan[e + f*x])^3,x]`

output  $(2^{(-5 - 2*m)} 3^{(-1 - m)} (c + d*x)^m (12^{(1 + m)} E^{((6*I)*e)} * f*(c + d*x) * ((I*f*(c + d*x))/d)^m + I*2^{(1 + m)} 3^{(2 + m)} * d * E^{((2*I)*(2*e + (c*f)/d)}) * (1 + m) * \text{Gamma}[1 + m, ((2*I)*f*(c + d*x))/d] + I*3^{(2 + m)} * d * E^{((2*I)*e + ((4*I)*c*f)/d} * (1 + m) * \text{Gamma}[1 + m, ((4*I)*f*(c + d*x))/d] + I*2^{(1 + m)} * d * E^{(((6*I)*c*f)/d} * (1 + m) * \text{Gamma}[1 + m, ((6*I)*f*(c + d*x))/d]) * \text{Sec}[e + f*x]^3 * (\text{Cos}[f*x] + I*\text{Sin}[f*x])^3) / (d * E^{((3*I)*e)} * f * (1 + m) * ((I*f*(c + d*x))/d)^m * (a + I*a*Tan[e + f*x])^3)$

### 3.38.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx$$

$$\downarrow 3042$$

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx$$

$$\downarrow 4212$$

$$\int \left( \frac{3e^{-2ie-2ifx}(c + dx)^m}{8a^3} + \frac{3e^{-4ie-4ifx}(c + dx)^m}{8a^3} + \frac{e^{-6ie-6ifx}(c + dx)^m}{8a^3} + \frac{(c + dx)^m}{8a^3} \right) dx$$

$$\downarrow 2009$$

$$\frac{3i2^{-m-4}e^{-2i\left(\frac{e-cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{2if(c+dx)}{d}\right)}{a^3f} +$$

$$\frac{3i2^{-2m-5}e^{-4i\left(\frac{e-cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{4if(c+dx)}{d}\right)}{a^3f} +$$

$$\frac{i2^{-m-4}3^{-m-1}e^{-6i\left(\frac{e-cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{6if(c+dx)}{d}\right)}{a^3f} + \frac{(c+dx)^{m+1}}{8a^3d(m+1)}$$

input `Int[(c + d*x)^m/(a + I*a*Tan[e + f*x])^3,x]`

output `(c + d*x)^(1 + m)/(8*a^3*d*(1 + m)) + ((3*I)*2^(-4 - m)*(c + d*x)^m*Gamma[1 + m, ((2*I)*f*(c + d*x))/d])/(a^3*E^((2*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m) + ((3*I)*2^(-5 - 2*m)*(c + d*x)^m*Gamma[1 + m, ((4*I)*f*(c + d*x))/d])/(a^3*E^((4*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m) + (I*2^(-4 - m)*3^(-1 - m)*(c + d*x)^m*Gamma[1 + m, ((6*I)*f*(c + d*x))/d])/(a^3*E^((6*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m)`

### 3.38.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4212 `Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a)]^(-n), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]`



### 3.38.4 Maple [F]

$$\int \frac{(dx + c)^m}{(a + ia \tan(fx + e))^3} dx$$

input `int((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x)`

output `int((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x)`

### 3.38.5 Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.79

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx =$$

$$\frac{18(-i dm - i d)e^{\left(-\frac{dm \log\left(\frac{2i f}{d}\right) + 2i de - 2i cf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idf x - icf)}{d}\right) + 9(-i dm - i d)e^{\left(-\frac{dm \log\left(\frac{4i f}{d}\right) + 4i de - 4i cf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idf x - icf)}{d}\right) + \dots}{\dots}$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output `-1/96*(18*(-I*d*m - I*d)*e^(-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d)*gamma(m + 1, -2*(-I*d*f*x - I*c*f)/d) + 9*(-I*d*m - I*d)*e^(-(d*m*log(4*I*f/d) + 4*I*d*e - 4*I*c*f)/d)*gamma(m + 1, -4*(-I*d*f*x - I*c*f)/d) + 2*(-I*d*m - I*d)*e^(-(d*m*log(6*I*f/d) + 6*I*d*e - 6*I*c*f)/d)*gamma(m + 1, -6*(-I*d*f*x - I*c*f)/d) - 12*(d*f*x + c*f)*(d*x + c)^m/(a^3*d*f*m + a^3*d*f)`

### 3.38.6 Sympy [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \frac{i \int \frac{(c+dx)^m}{\tan^3(e+fx) - 3i \tan^2(e+fx) - 3 \tan(e+fx) + i} dx}{a^3}$$

input `integrate((d*x+c)**m/(a+I*a*tan(f*x+e))**3,x)`

output `I*Integral((c + d*x)**m/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I), x)/a**3`

---

3.38.  $\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^3} dx$

**3.38.7 Maxima [F]**

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(dx + c)^m}{(ia \tan(fx + e) + a)^3} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `1/8*((d*m + d)*integrate((d*x + c)^m*cos(6*f*x + 6*e), x) + 3*(d*m + d)*integrate((d*x + c)^m*cos(4*f*x + 4*e), x) + 3*(d*m + d)*integrate((d*x + c)^m*cos(2*f*x + 2*e), x) - (I*d*m + I*d)*integrate((d*x + c)^m*sin(6*f*x + 6*e), x) + 3*(-I*d*m - I*d)*integrate((d*x + c)^m*sin(4*f*x + 4*e), x) + 3*(-I*d*m - I*d)*integrate((d*x + c)^m*sin(2*f*x + 2*e), x) + e^(m*log(d*x + c) + log(d*x + c))/(a^3*d*m + a^3*d)`

**3.38.8 Giac [F]**

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(dx + c)^m}{(ia \tan(fx + e) + a)^3} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((d*x + c)^m/(I*a*tan(f*x + e) + a)^3, x)`

**3.38.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(c + dx)^m}{(a + a \tan(e + fx) li)^3} dx$$

input `int((c + d*x)^m/(a + a*tan(e + f*x)*li)^3,x)`

output `int((c + d*x)^m/(a + a*tan(e + f*x)*li)^3, x)`

### 3.39 $\int (c + dx)^3 (a + b \tan(e + fx)) dx$

3.39.1	Optimal result . . . . .	274
3.39.2	Mathematica [B] (verified) . . . . .	275
3.39.3	Rubi [A] (verified) . . . . .	276
3.39.4	Maple [B] (verified) . . . . .	277
3.39.5	Fricas [B] (verification not implemented) . . . . .	277
3.39.6	Sympy [F] . . . . .	278
3.39.7	Maxima [B] (verification not implemented) . . . . .	278
3.39.8	Giac [F] . . . . .	279
3.39.9	Mupad [F(-1)] . . . . .	279

#### 3.39.1 Optimal result

Integrand size = 18, antiderivative size = 152

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \frac{a(c + dx)^4}{4d} + \frac{ib(c + dx)^4}{4d} - \frac{b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} + \frac{3ibd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3bd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{3ibd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}$$

output  $\frac{1}{4}a*(d*x+c)^4/d + \frac{1}{4}I*b*(d*x+c)^4/d - \frac{b*(d*x+c)^3*\ln(1+\exp(2*I*(f*x+e)))}{f} + \frac{3}{2}I*b*d*(d*x+c)^2*\text{polylog}(2, -\exp(2*I*(f*x+e)))/f^2 - \frac{3}{2}b*d^2*(d*x+c)*\text{polylog}(3, -\exp(2*I*(f*x+e)))/f^3 - \frac{3}{4}I*b*d^3*\text{polylog}(4, -\exp(2*I*(f*x+e)))/f^4$

### 3.39.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 342 vs.  $2(152) = 304$ .

Time = 0.15 (sec) , antiderivative size = 342, normalized size of antiderivative = 2.25

$$\int (c + dx)^3(a + b \tan(e + fx)) dx = ac^3x + \frac{3}{2}ac^2dx^2 + \frac{3}{2}ibc^2dx^2 + acd^2x^3 + ibcd^2x^3 + \frac{1}{4}ad^3x^4 + \frac{1}{4}ibd^3x^4 - \frac{3bc^2dx \log(1 + e^{2i(e+fx)})}{f} - \frac{3bcd^2x^2 \log(1 + e^{2i(e+fx)})}{f} - \frac{bd^3x^3 \log(1 + e^{2i(e+fx)})}{f} - \frac{bc^3 \log(\cos(e + fx))}{f} + \frac{3ibc^2d \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} + \frac{3ibcd^2x \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \frac{3ibd^3x^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3bcd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{3bd^3x \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{3ibd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}$$

input `Integrate[(c + d*x)^3*(a + b*Tan[e + f*x]),x]`

output `a*c^3*x + (3*a*c^2*d*x^2)/2 + ((3*I)/2)*b*c^2*d*x^2 + a*c*d^2*x^3 + I*b*c*d^2*x^3 + (a*d^3*x^4)/4 + (I/4)*b*d^3*x^4 - (3*b*c^2*d*x*Log[1 + E^((2*I)*(e + f*x))])/f - (3*b*c*d^2*x^2*Log[1 + E^((2*I)*(e + f*x))])/f - (b*d^3*x^3*Log[1 + E^((2*I)*(e + f*x))])/f - (b*c^3*Log[Cos[e + f*x]])/f + (((3*I)/2)*b*c^2*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + ((3*I)*b*c*d^2*x*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (((3*I)/2)*b*d^3*x^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (3*b*c*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (3*b*d^3*x*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (((3*I)/4)*b*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4`

### 3.39.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx$$

↓ 3042

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx$$

↓ 4205

$$\int (a(c + dx)^3 + b(c + dx)^3 \tan(e + fx)) dx$$

↓ 2009

$$\frac{a(c + dx)^4}{4d} - \frac{3bd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{3ibd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} + \frac{ib(c + dx)^4}{4d} - \frac{3ibd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}$$

input `Int[(c + d*x)^3*(a + b*Tan[e + f*x]),x]`

output `(a*(c + d*x)^4)/(4*d) + ((I/4)*b*(c + d*x)^4)/d - (b*(c + d*x)^3*Log[1 + E^((2*I)*(e + f*x))])/f + (((3*I)/2)*b*d*(c + d*x)^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (3*b*d^2*(c + d*x)*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (((3*I)/4)*b*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4`

#### 3.39.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4205 Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

### 3.39.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 499 vs.  $2(131) = 262$ .

Time = 0.64 (sec) , antiderivative size = 500, normalized size of antiderivative = 3.29

method	result
risch	$-\frac{ibc^4}{4d} - ibc^3x + \frac{id^3bx^4}{4} - \frac{bc^3 \ln(e^{2i(fx+e)}+1)}{f} + \frac{2bc^3 \ln(e^{i(fx+e)})}{f} - \frac{bd^3 \ln(e^{2i(fx+e)}+1)x^3}{f} - \frac{3bd^3 \operatorname{Li}_3(-e^{2i(fx+e)})}{2f^3}$

```
input int((d*x+c)^3*(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 3*I/f^2*b*d^2*c*polylog(2,-exp(2*I*(f*x+e)))*x-6*I/f^2*b*d^2*c*e^2*x+6*I/f
*b*d*c^2*e*x+I*d^2*b*c*x^3-1/4*I/d*b*c^4+d^2*a*c*x^3+3/2*d*a*c^2*x^2+a*c^3
*x-I*b*c^3*x+1/4*I*d^3*b*x^4-1/f*b*c^3*ln(exp(2*I*(f*x+e))+1)+2/f*b*c^3*ln
(exp(I*(f*x+e)))-3/f*b*d^2*c*ln(exp(2*I*(f*x+e))+1)*x^2-3/f*b*d*c^2*ln(exp
(2*I*(f*x+e))+1)*x+3/2*I/f^2*b*d^3*polylog(2,-exp(2*I*(f*x+e)))*x^2+3*I/f^
2*b*d*c^2*e^2-4*I/f^3*b*d^2*c*e^3+2*I/f^3*b*d^3*e^3*x+3/2*I/f^2*b*d*c^2*po
lylog(2,-exp(2*I*(f*x+e)))+1/4*d^3*a*x^4+1/4/d*a*c^4+6/f^3*b*e^2*d^2*c*ln(
exp(I*(f*x+e)))-6/f^2*b*e*d*c^2*ln(exp(I*(f*x+e)))-1/f*b*d^3*ln(exp(2*I*(f
*x+e))+1)*x^3-3/2/f^3*b*d^3*polylog(3,-exp(2*I*(f*x+e)))*x-2/f^4*b*e^3*d^3
*ln(exp(I*(f*x+e)))-3/2/f^3*b*d^2*c*polylog(3,-exp(2*I*(f*x+e)))+3/2*I/f^4
*b*d^3*e^4+3/2*I*d*b*c^2*x^2-3/4*I*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4
```

### 3.39.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 500 vs.  $2(127) = 254$ .

Time = 0.27 (sec) , antiderivative size = 500, normalized size of antiderivative = 3.29

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx$$

$$= \frac{2ad^3 f^4 x^4 + 8acd^2 f^4 x^3 + 12ac^2 df^4 x^2 + 8ac^3 f^4 x + 3ibd^3 \operatorname{polylog}\left(4, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3ibd^3 \operatorname{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3ibd^3 \operatorname{polylog}\left(2, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3ibd^3 \operatorname{polylog}\left(1, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right)}{f^4}$$

---

3.39.  $\int (c + dx)^3 (a + b \tan(e + fx)) dx$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="fricas")`

output 
$$\begin{aligned} & 1/8*(2*a*d^3*f^4*x^4 + 8*a*c*d^2*f^4*x^3 + 12*a*c^2*d*f^4*x^2 + 8*a*c^3*f^4*x \\ & + 3*I*b*d^3*polylog(4, (\tan(f*x + e))^2 + 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 3*I*b*d^3*polylog(4, (\tan(f*x + e))^2 - 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 6*(I*b*d^3*f^2*x^2 + 2*I*b*c*d^2*f^2*x + I*b*c^2*d*f^2)*dilog(2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) - 6*(-I*b*d^3*f^2*x^2 - 2*I*b*c*d^2*f^2*x - I*b*c^2*d*f^2)*dilog(2*(-I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*c^3*f^3)*log(-2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*c^3*f^3)*log(-2*(-I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 6*(b*d^3*f*x + b*c*d^2*f)*polylog(3, (\tan(f*x + e))^2 + 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 6*(b*d^3*f*x + b*c*d^2*f)*polylog(3, (\tan(f*x + e))^2 - 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)))/f^4 \end{aligned}$$

### 3.39.6 Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^3 dx$$

input `integrate((d*x+c)**3*(a+b*tan(f*x+e)),x)`

output `Integral((a + b*tan(e + f*x))*(c + d*x)**3, x)`

### 3.39.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 672 vs.  $2(127) = 254$ .

Time = 0.72 (sec) , antiderivative size = 672, normalized size of antiderivative = 4.42

$$\begin{aligned} & \int (c + dx)^3 (a + b \tan(e + fx)) dx \\ & = \frac{12 (fx + e)ac^3 + \frac{3(fx+e)^4 ad^3}{f^3} - \frac{12(fx+e)^3 ad^3 e}{f^3} + \frac{18(fx+e)^2 ad^3 e^2}{f^3} - \frac{12(fx+e) ad^3 e^3}{f^3} + \frac{12(fx+e)^3 acd^2}{f^2} - \frac{36(fx+e)^2 acd^2 e}{f^2}}{1} \end{aligned}$$

---

3.39.  $\int (c + dx)^3 (a + b \tan(e + fx)) dx$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="maxima")`

output `1/12*(12*(f*x + e)*a*c^3 + 3*(f*x + e)^4*a*d^3/f^3 - 12*(f*x + e)^3*a*d^3*e/f^3 + 18*(f*x + e)^2*a*d^3*e^2/f^3 - 12*(f*x + e)*a*d^3*e^3/f^3 + 12*(f*x + e)^3*a*c*d^2/f^2 - 36*(f*x + e)^2*a*c*d^2*e/f^2 + 36*(f*x + e)*a*c*d^2*e^2/f^2 + 18*(f*x + e)^2*a*c^2*d/f - 36*(f*x + e)*a*c^2*d*e/f + 12*b*c^3*log(sec(f*x + e)) - 12*b*d^3*e^3*log(sec(f*x + e))/f^3 + 36*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 36*b*c^2*d*e*log(sec(f*x + e))/f - (-3*I*(f*x + e)^4*b*d^3 + 12*I*b*d^3*polylog(4, -e^(2*I*f*x + 2*I*e)) - 12*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e)^3 - 18*(I*b*d^3*e^2 - 2*I*b*c*d^2*e*f + I*b*c^2*d*f^2)*(f*x + e)^2 - 4*(-4*I*(f*x + e)^3*b*d^3 + 9*(I*b*d^3*e - I*b*c*d^2*f)*(f*x + e)^2 + 9*(-I*b*d^3*e^2 + 2*I*b*c*d^2*e*f - I*b*c^2*d*f^2)*(f*x + e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 6*(4*I*(f*x + e)^2*b*d^3 + 3*I*b*d^3*e^2 - 6*I*b*c*d^2*e*f + 3*I*b*c^2*d*f^2 + 6*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e))*dilog(-e^(2*I*f*x + 2*I*e)) + 2*(4*(f*x + e)^3*b*d^3 - 9*(b*d^3*e - b*c*d^2*f)*(f*x + e)^2 + 9*(b*d^3*e^2 - 2*b*c*d^2*e*f + b*c^2*d*f^2)*(f*x + e))*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) + 6*(4*(f*x + e)*b*d^3 - 3*b*d^3*e + 3*b*c*d^2*f)*polylog(3, -e^(2*I*f*x + 2*I*e)))/f^3)/f`

### 3.39.8 Giac [F]

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \int (dx + c)^3 (b \tan(fx + e) + a) dx$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^3*(b*tan(f*x + e) + a), x)`

### 3.39.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^3 dx$$

input `int((a + b*tan(e + f*x))*(c + d*x)^3,x)`



output `int((a + b*tan(e + f*x))*(c + d*x)^3, x)`

### 3.40 $\int (c + dx)^2 (a + b \tan(e + fx)) dx$

3.40.1	Optimal result . . . . .	281
3.40.2	Mathematica [A] (verified) . . . . .	282
3.40.3	Rubi [A] (verified) . . . . .	282
3.40.4	Maple [B] (verified) . . . . .	283
3.40.5	Fricas [B] (verification not implemented) . . . . .	284
3.40.6	Sympy [F] . . . . .	285
3.40.7	Maxima [B] (verification not implemented) . . . . .	285
3.40.8	Giac [F] . . . . .	286
3.40.9	Mupad [F(-1)] . . . . .	286

#### 3.40.1 Optimal result

Integrand size = 18, antiderivative size = 115

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx = \frac{a(c + dx)^3}{3d} + \frac{ib(c + dx)^3}{3d} - \frac{b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} + \frac{ibd(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{bd^2 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3}$$

```
output 1/3*a*(d*x+c)^3/d+1/3*I*b*(d*x+c)^3/d-b*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f
+I*b*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-1/2*b*d^2*polylog(3,-exp(2
*I*(f*x+e)))/f^3
```

### 3.40.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.66

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx = ac^2x + acdx^2 + ibcdx^2 + \frac{1}{3}ad^2x^3 + \frac{1}{3}ibd^2x^3 - \frac{2bcdx \log(1 + e^{2i(e+fx)})}{f} - \frac{bd^2x^2 \log(1 + e^{2i(e+fx)})}{f} - \frac{bc^2 \log(\cos(e + fx))}{f} + \frac{ibcd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \frac{ibd^2x \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3}$$

input `Integrate[(c + d*x)^2*(a + b*Tan[e + f*x]),x]`

output `a*c^2*x + a*c*d*x^2 + I*b*c*d*x^2 + (a*d^2*x^3)/3 + (I/3)*b*d^2*x^3 - (2*b*c*d*x*Log[1 + E^((2*I)*(e + f*x))])/f - (b*d^2*x^2*Log[1 + E^((2*I)*(e + f*x))])/f - (b*c^2*Log[Cos[e + f*x]])/f + (I*b*c*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (I*b*d^2*x*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3)`

### 3.40.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx$$

↓ 3042

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx$$

↓ 4205

$$\int (a(c + dx)^2 + b(c + dx)^2 \tan(e + fx)) dx$$

↓ 2009

$$\frac{a(c + dx)^3}{3d} + \frac{ibd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} + \frac{ib(c + dx)^3}{3d} - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3}$$

input `Int[(c + d*x)^2*(a + b*Tan[e + f*x]),x]`

output `(a*(c + d*x)^3)/(3*d) + ((I/3)*b*(c + d*x)^3)/d - (b*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f + (I*b*d*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3)`

### 3.40.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

### 3.40.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 313 vs. 2(101) = 202.

Time = 0.58 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.73

method	result
risch	$\frac{id^2bx^3}{3} - ibc^2x + \frac{d^2ax^3}{3} + \frac{4ibdce}{f} + dacx^2 + \frac{2ibdce^2}{f^2} + ac^2x + \frac{ac^3}{3d} - \frac{bc^2 \ln(e^{2i(fx+e)}+1)}{f} - \frac{2bdc \ln(e^{2i(fx+e)})}{f}$

```
input int((d*x+c)^2*(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/3*I*d^2*b*x^3-I*b*c^2*x+1/3*d^2*a*x^3+4*I/f*b*d*c*e*x+d*a*c*x^2+2*I/f^2*
b*d*c*e^2+a*c^2*x+1/3/d*a*c^3-1/f*b*c^2*ln(exp(2*I*(f*x+e))+1)-2/f*b*d*c*ln
n(exp(2*I*(f*x+e))+1)*x+I*d*b*c*x^2+2/f^3*b*d^2*e^2*ln(exp(I*(f*x+e)))-2*I
/f^2*b*d^2*e^2*x-1/f*b*d^2*ln(exp(2*I*(f*x+e))+1)*x^2-1/2*b*d^2*polylog(3,
-exp(2*I*(f*x+e)))/f^3+I/f^2*b*d*c*polylog(2,-exp(2*I*(f*x+e)))+I/f^2*b*d^
2*polylog(2,-exp(2*I*(f*x+e)))*x-1/3*I/d*b*c^3-4/3*I/f^3*b*d^2*e^3-4/f^2*b
*c*d*e*ln(exp(I*(f*x+e)))+2/f*b*c^2*ln(exp(I*(f*x+e)))
```

### 3.40.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 317 vs.  $2(98) = 196$ .

Time = 0.26 (sec) , antiderivative size = 317, normalized size of antiderivative = 2.76

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx$$

$$= \frac{4ad^2f^3x^3 + 12acdf^3x^2 + 12ac^2f^3x - 3bd^2 \operatorname{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3bd^2 \operatorname{polylog}\left(3, \frac{\tan(fx+e)}{\tan(fx+e)^2 + 1}\right)}{f^3}$$

```
input integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="fricas")
```

```
output 1/12*(4*a*d^2*f^3*x^3 + 12*a*c*d*f^3*x^2 + 12*a*c^2*f^3*x - 3*b*d^2*polylo
g(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*b*d
^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)
) - 6*(I*b*d^2*f*x + I*b*c*d*f)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)
^2 + 1) + 1) - 6*(-I*b*d^2*f*x - I*b*c*d*f)*dilog(2*(-I*tan(f*x + e) - 1)/
(tan(f*x + e)^2 + 1) + 1) - 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x + b*c^2*f^2)*
log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*(b*d^2*f^2*x^2 + 2*b
*c*d*f^2*x + b*c^2*f^2)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)
)/f^3
```

### 3.40.6 Sympy [F]

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^2 dx$$

input `integrate((d*x+c)**2*(a+b*tan(f*x+e)),x)`

output `Integral((a + b*tan(e + f*x))*(c + d*x)**2, x)`

### 3.40.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 373 vs.  $2(98) = 196$ .

Time = 0.52 (sec) , antiderivative size = 373, normalized size of antiderivative = 3.24

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx$$

$$= \frac{6(fx + e)ac^2 + \frac{2(fx+e)^3ad^2}{f^2} - \frac{6(fx+e)^2ad^2e}{f^2} + \frac{6(fx+e)ad^2e^2}{f^2} + \frac{6(fx+e)^2acd}{f} - \frac{12(fx+e)acde}{f} + 6bc^2 \log(\sec(fx + e))}{1}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="maxima")`

output `1/6*(6*(f*x + e)*a*c^2 + 2*(f*x + e)^3*a*d^2/f^2 - 6*(f*x + e)^2*a*d^2*e/f^2 + 6*(f*x + e)*a*d^2*e^2/f^2 + 6*(f*x + e)^2*a*c*d/f - 12*(f*x + e)*a*c*d*e/f + 6*b*c^2*log(sec(f*x + e)) + 6*b*d^2*e^2*log(sec(f*x + e))/f^2 - 12*b*c*d*e*log(sec(f*x + e))/f - (-2*I*(f*x + e)^3*b*d^2 + 3*b*d^2*polylog(3, -e^(2*I*f*x + 2*I*e))) - 6*(-I*b*d^2*e + I*b*c*d*f)*(f*x + e)^2 - 6*(-I*(f*x + e)^2*b*d^2 + 2*(I*b*d^2*e - I*b*c*d*f)*(f*x + e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 6*(I*(f*x + e)*b*d^2 - I*b*d^2*e + I*b*c*d*f)*dilog(-e^(2*I*f*x + 2*I*e)) + 3*((f*x + e)^2*b*d^2 - 2*(b*d^2*e - b*c*d*f)*(f*x + e))*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1))/f^2/f`

**3.40.8 Giac [F]**

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx = \int (dx + c)^2 (b \tan(fx + e) + a) dx$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^2*(b*tan(f*x + e) + a), x)`

**3.40.9 Mupad [F(-1)]**

Timed out.

$$\int (c + dx)^2 (a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^2 dx$$

input `int((a + b*tan(e + f*x))*(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))*(c + d*x)^2, x)`

### 3.41 $\int (c + dx)(a + b \tan(e + fx)) dx$

3.41.1	Optimal result . . . . .	287
3.41.2	Mathematica [A] (verified) . . . . .	287
3.41.3	Rubi [A] (verified) . . . . .	288
3.41.4	Maple [A] (verified) . . . . .	289
3.41.5	Fricas [B] (verification not implemented) . . . . .	289
3.41.6	Sympy [F] . . . . .	290
3.41.7	Maxima [A] (verification not implemented) . . . . .	290
3.41.8	Giac [F] . . . . .	290
3.41.9	Mupad [B] (verification not implemented) . . . . .	291

#### 3.41.1 Optimal result

Integrand size = 16, antiderivative size = 84

$$\int (c + dx)(a + b \tan(e + fx)) dx = \frac{a(c + dx)^2}{2d} + \frac{ib(c + dx)^2}{2d} - \frac{b(c + dx) \log(1 + e^{2i(e+fx)})}{f} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2}$$

output  $1/2*a*(d*x+c)^2/d+1/2*I*b*(d*x+c)^2/d-b*(d*x+c)*\ln(1+\exp(2*I*(f*x+e)))/f+1/2*I*b*d*\operatorname{polylog}(2,-\exp(2*I*(f*x+e)))/f^2$

#### 3.41.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.04

$$\int (c + dx)(a + b \tan(e + fx)) dx = acx + \frac{1}{2}adx^2 + \frac{1}{2}ibdx^2 - \frac{bdx \log(1 + e^{2i(e+fx)})}{f} - \frac{bc \log(\cos(e + fx))}{f} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2}$$

input `Integrate[(c + d*x)*(a + b*Tan[e + f*x]),x]`

output  $a*c*x + (a*d*x^2)/2 + (I/2)*b*d*x^2 - (b*d*x*\operatorname{Log}[1 + E^((2*I)*(e + f*x))])/f - (b*c*\operatorname{Log}[\operatorname{Cos}[e + f*x]])/f + ((I/2)*b*d*\operatorname{PolyLog}[2, -E^((2*I)*(e + f*x))])/f^2$



### 3.41.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c + dx)(a + b \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (c + dx)(a + b \tan(e + fx)) dx \\ & \quad \downarrow \text{4205} \\ & \int (a(c + dx) + b(c + dx) \tan(e + fx)) dx \\ & \quad \downarrow \text{2009} \\ & \frac{a(c + dx)^2}{2d} - \frac{b(c + dx) \log(1 + e^{2i(e+fx)})}{f} + \frac{ib(c + dx)^2}{2d} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \end{aligned}$$

input `Int[(c + d*x)*(a + b*Tan[e + f*x]),x]`

output `(a*(c + d*x)^2)/(2*d) + ((I/2)*b*(c + d*x)^2)/d - (b*(c + d*x)*Log[1 + E^((2*I)*(e + f*x))])/f + ((I/2)*b*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2`

#### 3.41.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

### 3.41.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.70

method	result
risch	$\frac{ibd x^2}{2} + \frac{adx^2}{2} - ibcx + acx - \frac{bc \ln(e^{2i(fx+e)}+1)}{f} + \frac{2bc \ln(e^{i(fx+e)})}{f} + \frac{2ibdex}{f} + \frac{ibde^2}{f^2} - \frac{bd \ln(e^{2i(fx+e)}+1)x}{f} + \dots$

input `int((d*x+c)*(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output  $\frac{1}{2}I*b*d*x^2 + \frac{1}{2}*a*d*x^2 - I*b*c*x + a*c*x - \frac{1}{f}*b*c*\ln(\exp(2*I*(f*x+e))+1) + \frac{2}{f}*b*c*\ln(\exp(I*(f*x+e)))+2*I/f*b*d*e*x + I/f^2*b*d*e^2 - \frac{1}{f}*b*d*\ln(\exp(2*I*(f*x+e))+1)*x + \frac{1}{2}*I*b*d*\text{polylog}(2, -\exp(2*I*(f*x+e)))/f^2 - \frac{2}{f^2}*b*d*e*\ln(\exp(I*(f*x+e)))$

### 3.41.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 160 vs.  $2(69) = 138$ .

Time = 0.26 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.90

$$\int (c + dx)(a + b \tan(e + fx)) dx$$

$$= \frac{2adf^2x^2 + 4acf^2x - i bd \text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + i bd \text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) - 2(bdfx + bcf) \log\left(\frac{-1 - i \tan(fx+e)}{1 + i \tan(fx+e)}\right)}{4f^2}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="fracas")`

output  $\frac{1}{4}*(2*a*d*f^2*x^2 + 4*a*c*f^2*x - I*b*d*dilog(2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) + I*b*d*dilog(2*(-I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) - 2*(b*d*f*x + b*c*f)*\log(-2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 2*(b*d*f*x + b*c*f)*\log(-2*(-I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)))/f^2$

**3.41.6 Sympy [F]**

$$\int (c + dx)(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx))(c + dx) dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x)`

output `Integral((a + b*tan(e + f*x))*(c + d*x), x)`

**3.41.7 Maxima [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.55

$$\int (c + dx)(a + b \tan(e + fx)) dx$$

$$= \frac{(a + ib)df^2x^2 + 2(a + ib)cf^2x + ibd\text{Li}_2(-e^{(2ifx+2ie)}) + 2(-ibdfx - ibcf) \arctan(\sin(2fx + 2e), \cos(2fx + 2e))}{2f^2}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="maxima")`

output `1/2*((a + I*b)*d*f^2*x^2 + 2*(a + I*b)*c*f^2*x + I*b*d*dilog(-e^(2*I*f*x + 2*I*e)) + 2*(-I*b*d*f*x - I*b*c*f)*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - (b*d*f*x + b*c*f)*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1))/f^2`

**3.41.8 Giac [F]**

$$\int (c + dx)(a + b \tan(e + fx)) dx = \int (dx + c)(b \tan(fx + e) + a) dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)*(b*tan(f*x + e) + a), x)`

**3.41.9 Mupad [B] (verification not implemented)**

Time = 3.60 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.92

$$\int (c + dx)(a + b \tan(e + fx)) dx = \frac{ax(2c + dx)}{2} - \frac{bd(\pi \ln(\cos(fx)) - \pi \ln(e^{fx^{2i}} + 1) + \text{polylog}(2, -e^{-e^{2i}} e^{-fx^{2i}}) 1i - \pi \ln(e^{-e^{2i}} e^{-fx^{2i}} + 1) + 2e \ln(e^{-e^{2i}} e^{-fx^{2i}} + 1))}{2f^2} + \frac{bc \ln(\tan(e + fx)^2 + 1)}{2f}$$

input `int((a + b*tan(e + f*x))*(c + d*x),x)`output `(a*x*(2*c + d*x))/2 - (b*d*(polylog(2, -exp(-e*2i)*exp(-f*x*2i))*1i - pi*log(exp(f*x*2i) + 1) - log(cos(e + f*x))*(2*e - pi) - pi*log(exp(-e*2i)*exp(-f*x*2i) + 1) + 2*e*log(exp(-e*2i)*exp(-f*x*2i) + 1) + pi*log(cos(f*x)) + f^2*x^2*1i + 2*f*x*log(exp(-e*2i)*exp(-f*x*2i) + 1) + e*f*x*2i))/(2*f^2) + (b*c*log(tan(e + f*x)^2 + 1))/(2*f)`

### 3.42 $\int \frac{a+b \tan(e+fx)}{c+dx} dx$

3.42.1	Optimal result . . . . .	292
3.42.2	Mathematica [N/A] . . . . .	292
3.42.3	Rubi [N/A] . . . . .	293
3.42.4	Maple [N/A] (verified) . . . . .	294
3.42.5	Fricas [N/A] . . . . .	294
3.42.6	Sympy [N/A] . . . . .	294
3.42.7	Maxima [N/A] . . . . .	295
3.42.8	Giac [N/A] . . . . .	295
3.42.9	Mupad [N/A] . . . . .	295

#### 3.42.1 Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \text{Int}\left(\frac{a + b \tan(e + fx)}{c + dx}, x\right)$$

output `Unintegrable((a+b*tan(f*x+e))/(d*x+c),x)`

#### 3.42.2 Mathematica [N/A]

Not integrable

Time = 1.91 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

input `Integrate[(a + b*Tan[e + f*x])/(c + d*x),x]`

output `Integrate[(a + b*Tan[e + f*x])/(c + d*x), x]`

### 3.42.3 Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx$$

↓ 3042

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx$$

↓ 4223

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx$$

input `Int[(a + b*Tan[e + f*x])/(c + d*x),x]`

output `$Aborted`

#### 3.42.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.42.4 Maple [N/A] (verified)**

Not integrable

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tan(fx + e)}{dx + c} dx$$

input `int((a+b*tan(f*x+e))/(d*x+c),x)`output `int((a+b*tan(f*x+e))/(d*x+c),x)`**3.42.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="fricas")`output `integral((b*tan(f*x + e) + a)/(d*x + c), x)`**3.42.6 Sympy [N/A]**

Not integrable

Time = 0.66 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x)`output `Integral((a + b*tan(e + f*x))/(c + d*x), x)`

**3.42.7 Maxima [N/A]**

Not integrable

Time = 0.44 (sec) , antiderivative size = 86, normalized size of antiderivative = 4.78

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="maxima")`output `(2*b*d*integrate(sin(2*f*x + 2*e)/((d*x + c)*cos(2*f*x + 2*e)^2 + (d*x + c)*sin(2*f*x + 2*e)^2 + d*x + 2*(d*x + c)*cos(2*f*x + 2*e) + c), x) + a*log(d*x + c))/d`**3.42.8 Giac [N/A]**

Not integrable

Time = 0.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="giac")`output `integrate((b*tan(f*x + e) + a)/(d*x + c), x)`**3.42.9 Mupad [N/A]**

Not integrable

Time = 3.53 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

input `int((a + b*tan(e + f*x))/(c + d*x),x)`output `int((a + b*tan(e + f*x))/(c + d*x), x)`



### 3.43 $\int \frac{a+b \tan(e+fx)}{(c+dx)^2} dx$

3.43.1	Optimal result	296
3.43.2	Mathematica [N/A]	296
3.43.3	Rubi [N/A]	297
3.43.4	Maple [N/A] (verified)	298
3.43.5	Fricas [N/A]	298
3.43.6	Sympy [N/A]	298
3.43.7	Maxima [N/A]	299
3.43.8	Giac [N/A]	299
3.43.9	Mupad [N/A]	299

#### 3.43.1 Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \text{Int}\left(\frac{a + b \tan(e + fx)}{(c + dx)^2}, x\right)$$

output `Unintegrable((a+b*tan(f*x+e))/(d*x+c)^2,x)`

#### 3.43.2 Mathematica [N/A]

Not integrable

Time = 5.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

input `Integrate[(a + b*Tan[e + f*x])/(c + d*x)^2,x]`

output `Integrate[(a + b*Tan[e + f*x])/(c + d*x)^2, x]`

### 3.43.3 Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

↓ 3042

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

↓ 4223

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

input `Int[(a + b*Tan[e + f*x])/(c + d*x)^2,x]`

output `$Aborted`

#### 3.43.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.43.4 Maple [N/A] (verified)**

Not integrable

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tan(fx + e)}{(dx + c)^2} dx$$

input `int((a+b*tan(f*x+e))/(d*x+c)^2,x)`output `int((a+b*tan(f*x+e))/(d*x+c)^2,x)`**3.43.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.72

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="fricas")`output `integral((b*tan(f*x + e) + a)/(d^2*x^2 + 2*c*d*x + c^2), x)`**3.43.6 Sympy [N/A]**

Not integrable

Time = 2.49 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)**2,x)`output `Integral((a + b*tan(e + f*x))/(c + d*x)**2, x)`

**3.43.7 Maxima [N/A]**

Not integrable

Time = 0.46 (sec) , antiderivative size = 142, normalized size of antiderivative = 7.89

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="maxima")`

output `(2*(b*d^2*x + b*c*d)*integrate(sin(2*f*x + 2*e)/(d^2*x^2 + 2*c*d*x + (d^2*x^2 + 2*c*d*x + c^2)*cos(2*f*x + 2*e)^2 + (d^2*x^2 + 2*c*d*x + c^2)*sin(2*f*x + 2*e)^2 + c^2 + 2*(d^2*x^2 + 2*c*d*x + c^2)*cos(2*f*x + 2*e)), x) - a)/(d^2*x + c*d)`

**3.43.8 Giac [N/A]**

Not integrable

Time = 4.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)/(d*x + c)^2, x)`

**3.43.9 Mupad [N/A]**

Not integrable

Time = 3.68 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

input `int((a + b*tan(e + f*x))/(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))/(c + d*x)^2, x)`

### 3.44 $\int (c + dx)^3 (a + b \tan(e + fx))^2 dx$

3.44.1	Optimal result	300
3.44.2	Mathematica [B] (warning: unable to verify)	301
3.44.3	Rubi [A] (verified)	302
3.44.4	Maple [B] (verified)	303
3.44.5	Fricas [B] (verification not implemented)	304
3.44.6	Sympy [F]	305
3.44.7	Maxima [B] (verification not implemented)	305
3.44.8	Giac [F]	306
3.44.9	Mupad [F(-1)]	307

#### 3.44.1 Optimal result

Integrand size = 20, antiderivative size = 300

$$\begin{aligned}
 \int (c + dx)^3 (a + b \tan(e + fx))^2 dx = & -\frac{ib^2(c + dx)^3}{f} + \frac{a^2(c + dx)^4}{4d} + \frac{iab(c + dx)^4}{2d} \\
 & - \frac{b^2(c + dx)^4}{4d} + \frac{3b^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\
 & - \frac{2ab(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
 & - \frac{3ib^2d^2(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
 & + \frac{3iabd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & + \frac{3b^2d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \\
 & - \frac{3abd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\
 & - \frac{3iabd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{2f^4} \\
 & + \frac{b^2(c + dx)^3 \tan(e + fx)}{f}
 \end{aligned}$$

output 
$$-I*b^2*(d*x+c)^3/f+1/4*a^2*(d*x+c)^4/d+1/2*I*a*b*(d*x+c)^4/d-1/4*b^2*(d*x+c)^4/d+3*b^2*d*(d*x+c)^2*\ln(1+\exp(2*I*(f*x+e)))/f^2-2*a*b*(d*x+c)^3*\ln(1+\exp(2*I*(f*x+e)))/f-3*I*b^2*d^2*(d*x+c)*\text{polylog}(2,-\exp(2*I*(f*x+e)))/f^3+3*I*a*b*d*(d*x+c)^2*\text{polylog}(2,-\exp(2*I*(f*x+e)))/f^2+3/2*b^2*d^3*\text{polylog}(3,-\exp(2*I*(f*x+e)))/f^4-3*a*b*d^2*(d*x+c)*\text{polylog}(3,-\exp(2*I*(f*x+e)))/f^3-3/2*I*a*b*d^3*\text{polylog}(4,-\exp(2*I*(f*x+e)))/f^4+b^2*(d*x+c)^3*\tan(f*x+e)/f$$

### 3.44.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1337 vs.  $2(300) = 600$ .

Time = 7.33 (sec) , antiderivative size = 1337, normalized size of antiderivative = 4.46

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

input `Integrate[(c + d*x)^3*(a + b*Tan[e + f*x])^2,x]`

output 
$$\begin{aligned} & ((I/4)*b^2*d^3*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^{((2*I)*e)}))*\text{Log}[1 + E^{((-2*I)*(e + f*x))}] + 6*(1 + E^{((2*I)*e)})*f*x*\text{PolyLog}[2, -E^{((-2*I)*(e + f*x))}] \\ & - (3*I)*(1 + E^{((2*I)*e)})*\text{PolyLog}[3, -E^{((-2*I)*(e + f*x))}])*\text{Sec}[e])/(E^{(I*e)*f^4} - ((I/2)*a*b*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^{((2*I)*e)}))* \\ & \text{Log}[1 + E^{((-2*I)*(e + f*x))}] + 6*(1 + E^{((2*I)*e)})*f*x*\text{PolyLog}[2, -E^{((-2*I)*(e + f*x))}] \\ & - (3*I)*(1 + E^{((2*I)*e)})*\text{PolyLog}[3, -E^{((-2*I)*(e + f*x))}])*\text{Sec}[e])/(E^{(I*e)*f^3} - ((I/4)*a*b*d^3*E^{(I*e)}*((2*f^4*x^4)/E^{((2*I)*e)} \\ & - (4*I)*(1 + E^{((-2*I)*e)})*f^3*x^3*\text{Log}[1 + E^{((-2*I)*(e + f*x))}] + 6*(1 + E^{((-2*I)*e)})*f^2*x^2*\text{PolyLog}[2, -E^{((-2*I)*(e + f*x))}] \\ & - (6*I)*(1 + E^{((-2*I)*e)})*f*x*\text{PolyLog}[3, -E^{((-2*I)*(e + f*x))}] - 3*(1 + E^{((-2*I)*e)})*\text{PolyLog}[4, -E^{((-2*I)*(e + f*x))}])*\text{Sec}[e])/f^4 + (3*b^2*c^2*d*\text{Sec}[e]*(\text{Cos}[e] \\ & *\text{Log}[\text{Cos}[e]*\text{Cos}[f*x] - \text{Sin}[e]*\text{Sin}[f*x] + f*x*\text{Sin}[e]))/(f^2*(\text{Cos}[e]^2 + \text{Sin}[e]^2)) - (2*a*b*c^3*\text{Sec}[e]*(\text{Cos}[e]*\text{Log}[\text{Cos}[e]*\text{Cos}[f*x] - \text{Sin}[e]*\text{Sin}[f*x] \\ & + f*x*\text{Sin}[e]))/(f*(\text{Cos}[e]^2 + \text{Sin}[e]^2)) + (3*b^2*c*d^2*\text{Csc}[e]*((f^2*x^2)/E^{(I*\text{ArcTan}[\text{Cot}[e]])} - (\text{Cot}[e]*(I*f*x*(-\text{Pi} - 2*\text{ArcTan}[\text{Cot}[e]]) - \text{Pi}*\text{Log}[ \\ & 1 + E^{((-2*I)*f*x}] - 2*(f*x - \text{ArcTan}[\text{Cot}[e]])*\text{Log}[1 - E^{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]])}])) + \text{Pi}*\text{Log}[\text{Cos}[f*x] - 2*\text{ArcTan}[\text{Cot}[e]]*\text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]] \\ & + I*\text{PolyLog}[2, E^{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]])}])))/\text{Sqrt}[1 + \text{Cot}[e]^2])* \text{Sec}[e])/(f^3*\text{Sqrt}[\text{Csc}[e]^2*(\text{Cos}[e]^2 + \text{Sin}[e]^2)]) - (3*a*b*c^2*d*\text{Csc}[e]*((f^2*x^2)/E^{(I*\text{ArcTan}[\text{Cot}[e]])} - (\text{Cot}[e]*(I*f*x*(-\text{Pi} - 2*\text{ArcTan}[\dots \end{aligned}$$

### 3.44.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^3 (a + b \tan(e + fx))^2 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c + dx)^3 (a + b \tan(e + fx))^2 dx \\
 & \quad \downarrow \text{4205} \\
 & \int (a^2(c + dx)^3 + 2ab(c + dx)^3 \tan(e + fx) + b^2(c + dx)^3 \tan^2(e + fx)) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2(c + dx)^4}{4d} - \frac{3abd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \frac{3iabd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \\
 & \frac{2ab(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} + \frac{iab(c + dx)^4}{2d} - \frac{3iabd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{2f^4} - \\
 & \frac{3ib^2d^2(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{3b^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{2f^4} + \\
 & \frac{b^2(c + dx)^3 \tan(e + fx)}{f} - \frac{ib^2(c + dx)^3}{f} - \frac{b^2(c + dx)^4}{4d} + \frac{3b^2d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4}
 \end{aligned}$$

input `Int[(c + d*x)^3*(a + b*Tan[e + f*x])^2,x]`

output `((-I)*b^2*(c + d*x)^3)/f + (a^2*(c + d*x)^4)/(4*d) + ((I/2)*a*b*(c + d*x)^4)/d - (b^2*(c + d*x)^4)/(4*d) + (3*b^2*d*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f^2 - (2*a*b*(c + d*x)^3*Log[1 + E^((2*I)*(e + f*x))])/f - ((3*I)*b^2*d^2*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^3 + ((3*I)*a*b*d*(c + d*x)^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (3*b^2*d^3*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^4) - (3*a*b*d^2*(c + d*x)*PolyLog[3, -E^((2*I)*(e + f*x))])/f^3 - (((3*I)/2)*a*b*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4 + (b^2*(c + d*x)^3*Tan[e + f*x])/f`

## 3.44.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

## 3.44.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 951 vs.  $2(271) = 542$ .

Time = 1.60 (sec) , antiderivative size = 952, normalized size of antiderivative = 3.17

method	result	size
risch	Expression too large to display	952

input `int((d*x+c)^3*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`



output

```

-12*I/f^2*b*e^2*d^2*c*a*x+6*I/f^2*b*a*c*d^2*polylog(2,-exp(2*I*(f*x+e)))*x
+12*I/f*b*d*c^2*a*e*x+2*I*b^2*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)/f/(exp(2
*I*(f*x+e))+1)+3/f^2*b^2*c^2*d*ln(exp(2*I*(f*x+e))+1)-6/f^2*b^2*c^2*d*ln(e
xp(I*(f*x+e)))+3/f^2*b^2*d^3*ln(exp(2*I*(f*x+e))+1)*x^2-2/f*b*a*c^3*ln(exp
(2*I*(f*x+e))+1)+4/f*b*a*c^3*ln(exp(I*(f*x+e)))-6/f^4*b^2*e^2*d^3*ln(exp(I
*(f*x+e)))-2*I/f*b^2*d^3*x^3+4*I/f^4*b^2*d^3*e^3+1/2*I*d^3*a*b*x^4-3/2*d*b
^2*c^2*x^2+1/4*d^3*a^2*x^4+1/4/d*a^2*c^4-1/4*d^3*b^2*x^4-b^2*c^3*x-1/4/d*b
^2*c^4-2*I*a*b*c^3*x-1/2*I/d*a*b*c^4+d^2*a^2*c*x^3+3/2*d*a^2*c^2*x^2+a^2*c
^3*x-d^2*b^2*c*x^3-3/2*I*a*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4-4/f^4*b
e^3*a*d^3*ln(exp(I*(f*x+e)))-3/f^3*b*d^3*a*polylog(3,-exp(2*I*(f*x+e)))*x+
6/f^2*b^2*d^2*c*ln(exp(2*I*(f*x+e))+1)*x-2/f*b*d^3*a*ln(exp(2*I*(f*x+e))+1
)*x^3-3/f^3*b*a*c*d^2*polylog(3,-exp(2*I*(f*x+e)))+12/f^3*b^2*e*c*d^2*ln(e
xp(I*(f*x+e)))-3*I/f^3*b^2*d^3*polylog(2,-exp(2*I*(f*x+e)))*x+3*I/f^4*b*e^
4*a*d^3-6*I/f*b^2*d^2*c*x^2-6*I/f^3*b^2*d^2*c*e^2-3*I/f^3*b^2*d^2*c*polylo
g(2,-exp(2*I*(f*x+e)))+6*I/f^3*b^2*d^3*e^2*x+2*I*d^2*a*b*c*x^3+3*I*d*a*b*c
^2*x^2+3/2*b^2*d^3*polylog(3,-exp(2*I*(f*x+e)))/f^4-12/f^2*b*e*a*c^2*d*ln(
exp(I*(f*x+e)))-6/f*b*d*c^2*a*ln(exp(2*I*(f*x+e))+1)*x-6/f*b*a*c*d^2*ln(ex
p(2*I*(f*x+e))+1)*x^2+12/f^3*b*e^2*a*c*d^2*ln(exp(I*(f*x+e)))+6*I/f^2*b*d*
c^2*a*e^2+3*I/f^2*b*d*c^2*a*polylog(2,-exp(2*I*(f*x+e)))+3*I/f^2*b*d^3*a*p
olylog(2,-exp(2*I*(f*x+e)))*x^2+4*I/f^3*b*e^3*a*d^3*x-8*I/f^3*b*e^3*d^2...

```

### 3.44.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 759 vs.  $2(264) = 528$ .

Time = 0.28 (sec) , antiderivative size = 759, normalized size of antiderivative = 2.53

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx$$

$$= \frac{(a^2 - b^2)d^3 f^4 x^4 + 4(a^2 - b^2)cd^2 f^4 x^3 + 6(a^2 - b^2)c^2 df^4 x^2 + 4(a^2 - b^2)c^3 f^4 x + 3iabd^3 \text{polylog}\left(4, \frac{\tan(fx+e)}{\tan(e)}\right)}{1}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output

```

1/4*((a^2 - b^2)*d^3*f^4*x^4 + 4*(a^2 - b^2)*c*d^2*f^4*x^3 + 6*(a^2 - b^2)
*c^2*d*f^4*x^2 + 4*(a^2 - b^2)*c^3*f^4*x + 3*I*a*b*d^3*polylog(4, (tan(f*x
+ e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*I*a*b*d^3*polylo
g(4, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*(I*
a*b*d^3*f^2*x^2 + I*a*b*c^2*d*f^2 - I*b^2*c*d^2*f + I*(2*a*b*c*d^2*f^2 - b
^2*d^3*f)*x)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(-
I*a*b*d^3*f^2*x^2 - I*a*b*c^2*d*f^2 + I*b^2*c*d^2*f - I*(2*a*b*c*d^2*f^2 -
b^2*d^3*f)*x)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 2
*(2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3 - 3*b^2*c^2*d*f^2 + 3*(2*a*b*c*d^2*f^3
- b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f^3 - b^2*c*d^2*f^2)*x)*log(-2*(I*tan(f
*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 2*(2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3
- 3*b^2*c^2*d*f^2 + 3*(2*a*b*c*d^2*f^3 - b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f
^3 - b^2*c*d^2*f^2)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1))
- 3*(2*a*b*d^3*f*x + 2*a*b*c*d^2*f - b^2*d^3)*polylog(3, (tan(f*x + e)^2 +
2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(2*a*b*d^3*f*x + 2*a*b*c*
d^2*f - b^2*d^3)*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f
*x + e)^2 + 1)) + 4*(b^2*d^3*f^3*x^3 + 3*b^2*c*d^2*f^3*x^2 + 3*b^2*c^2*d*f
^3*x + b^2*c^3*f^3)*tan(f*x + e))/f^4

```

### 3.44.6 Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^3 dx$$

input `integrate((d*x+c)**3*(a+b*tan(f*x+e))**2,x)`

output `Integral((a + b*tan(e + f*x))**2*(c + d*x)**3, x)`

### 3.44.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2525 vs.  $2(264) = 528$ .

Time = 1.73 (sec) , antiderivative size = 2525, normalized size of antiderivative = 8.42

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output

$$\begin{aligned} & 1/4*(4*(f*x + e)*a^2*c^3 + (f*x + e)^4*a^2*d^3/f^3 - 4*(f*x + e)^3*a^2*d^3 \\ & *e/f^3 + 6*(f*x + e)^2*a^2*d^3*e^2/f^3 - 4*(f*x + e)*a^2*d^3*e^3/f^3 + 4*( \\ & f*x + e)^3*a^2*c*d^2/f^2 - 12*(f*x + e)^2*a^2*c*d^2*e/f^2 + 12*(f*x + e)*a \\ & ^2*c*d^2*e^2/f^2 + 6*(f*x + e)^2*a^2*c^2*d/f - 12*(f*x + e)*a^2*c^2*d*e/f \\ & + 8*a*b*c^3*\log(\sec(f*x + e)) - 8*a*b*d^3*e^3*\log(\sec(f*x + e))/f^3 + 24*a \\ & *b*c*d^2*e^2*\log(\sec(f*x + e))/f^2 - 24*a*b*c^2*d*e*\log(\sec(f*x + e))/f + \\ & 4*(3*(2*a*b + I*b^2)*(f*x + e)^4*d^3 - 24*b^2*d^3*e^3 + 72*b^2*c*d^2*e^2*f \\ & - 72*b^2*c^2*d*e*f^2 + 24*b^2*c^3*f^3 - 12*((2*a*b + I*b^2)*d^3*e - (2*a* \\ & b + I*b^2)*c*d^2*f)*(f*x + e)^3 + 18*((2*a*b + I*b^2)*d^3*e^2 - 2*(2*a*b + \\ & I*b^2)*c*d^2*e*f + (2*a*b + I*b^2)*c^2*d*f^2)*(f*x + e)^2 + 12*(-I*b^2*d^ \\ & 3*e^3 + 3*I*b^2*c*d^2*e^2*f - 3*I*b^2*c^2*d*e*f^2 + I*b^2*c^3*f^3)*(f*x + \\ & e) - 4*(8*(f*x + e)^3*a*b*d^3 - 9*b^2*d^3*e^2 + 18*b^2*c*d^2*e*f - 9*b^2*c \\ & ^2*d*f^2 - 9*(2*a*b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e)^2 + 18*(a*b \\ & *d^3*e^2 + a*b*c^2*d*f^2 + b^2*d^3*e - (2*a*b*c*d^2*e + b^2*c*d^2)*f)*(f*x \\ & + e) + (8*(f*x + e)^3*a*b*d^3 - 9*b^2*d^3*e^2 + 18*b^2*c*d^2*e*f - 9*b^2* \\ & c^2*d*f^2 - 9*(2*a*b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e)^2 + 18*(a* \\ & b*d^3*e^2 + a*b*c^2*d*f^2 + b^2*d^3*e - (2*a*b*c*d^2*e + b^2*c*d^2)*f)*(f* \\ & x + e))*\cos(2*f*x + 2*e) - (-8*I*(f*x + e)^3*a*b*d^3 + 9*I*b^2*d^3*e^2 - 1 \\ & 8*I*b^2*c*d^2*e*f + 9*I*b^2*c^2*d*f^2 + 9*(2*I*a*b*d^3*e - 2*I*a*b*c*d^2*f \\ & + I*b^2*d^3)*(f*x + e)^2 + 18*(-I*a*b*d^3*e^2 - I*a*b*c^2*d*f^2 - I*b^2*d^3) \end{aligned}$$

### 3.44.8 Giac [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (dx + c)^3 (b \tan(fx + e) + a)^2 dx$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)^3*(b*tan(f*x + e) + a)^2, x)`

**3.44.9 Mupad [F(-1)]**

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^3 dx$$

input `int((a + b*tan(e + f*x))^2*(c + d*x)^3,x)`output `int((a + b*tan(e + f*x))^2*(c + d*x)^3, x)`

### 3.45 $\int (c + dx)^2 (a + b \tan(e + fx))^2 dx$

3.45.1	Optimal result	308
3.45.2	Mathematica [B] (warning: unable to verify)	309
3.45.3	Rubi [A] (verified)	310
3.45.4	Maple [B] (verified)	311
3.45.5	Fricas [B] (verification not implemented)	312
3.45.6	Sympy [F]	313
3.45.7	Maxima [B] (verification not implemented)	313
3.45.8	Giac [F]	314
3.45.9	Mupad [F(-1)]	315

#### 3.45.1 Optimal result

Integrand size = 20, antiderivative size = 229

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = -\frac{ib^2(c + dx)^2}{f} + \frac{a^2(c + dx)^3}{3d} + \frac{2iab(c + dx)^3}{3d} - \frac{b^2(c + dx)^3}{3d} + \frac{2b^2d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} - \frac{2ab(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} - \frac{ib^2d^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{2iabd(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{abd^2 \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \frac{b^2(c + dx)^2 \tan(e + fx)}{f}$$

output

```
-I*b^2*(d*x+c)^2/f+1/3*a^2*(d*x+c)^3/d+2/3*I*a*b*(d*x+c)^3/d-1/3*b^2*(d*x+c)^3/d+2*b^2*d*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^2-2*a*b*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f-I*b^2*d^2*polylog(2,-exp(2*I*(f*x+e)))/f^3+2*I*a*b*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-a*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+b^2*(d*x+c)^2*tan(f*x+e)/f
```

### 3.45.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 649 vs.  $2(229) = 458$ .

Time = 6.83 (sec) , antiderivative size = 649, normalized size of antiderivative = 2.83

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx =$$

$$\frac{iabd^2 e^{-ie} (2f^2 x^2 (2fx - 3i(1 + e^{2ie}) \log(1 + e^{-2i(e+fx)})) + 6(1 + e^{2ie}) fx \operatorname{PolyLog}(2, -e^{-2i(e+fx)}) - 3i)}{6f^3}$$

$$+ \frac{1}{3} x (3c^2 + 3cdx + d^2 x^2) \sec(e) (a^2 \cos(e) - b^2 \cos(e) + 2ab \sin(e))$$

$$+ \frac{2b^2 cd \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + fx \sin(e))}{f^2 (\cos^2(e) + \sin^2(e))}$$

$$- \frac{2abc^2 \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + fx \sin(e))}{f (\cos^2(e) + \sin^2(e))}$$

$$+ \frac{b^2 d^2 \csc(e) \left( e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (ifx(-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2ifx}) - 2(fx - \arctan(\cot(e))) \log(1 - e^{2i(fx - \arctan(\cot(e))))}{f^3 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)}{f^3 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}}$$

$$- \frac{2abcd \csc(e) \left( e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (ifx(-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2ifx}) - 2(fx - \arctan(\cot(e))) \log(1 - e^{2i(fx - \arctan(\cot(e))))}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}}$$

$$+ \frac{\sec(e) \sec(e + fx) (b^2 c^2 \sin(fx) + 2b^2 cdx \sin(fx) + b^2 d^2 x^2 \sin(fx))}{f}$$

input `Integrate[(c + d*x)^2*(a + b*Tan[e + f*x])^2,x]`

output

```
((-1/6*I)*a*b*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))])*Sec[e])/
(E^(I*e)*f^3) + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*Sec[e]*(a^2*Cos[e] - b^2*Cos[e] + 2*a*b*Sin[e]))/3 + (2*b^2*c*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^2*(Cos[e]^2 + Sin[e]^2)) - (2*a*b*c^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (b^2*d^2*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]]) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]]) - Pi*Log[1 + E^((-2*I)*f*x]) - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]])]) + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]]) + I*PolyLog[2, E^((2*I)*(f*x - ArcTan[Cot[e]])])))/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^3*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) - (2*a*b*c*d*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]]) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]]) - Pi*Log[1 + E^((-2*I)*f*x]) - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]])]) + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]]) + I*PolyLog[2, E^((2*I)*(f*x - ArcTan[Cot[e]])])))/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^2*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) + (Sec[e]*Sec[e + f*x]*(b^2*c^2*Sin[f*x] + 2*b^2*c*d*x*Sin[f*x] + b^2*d^2*x^2*Sin[f*x]))/f
```

### 3.45.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx$$

$$\downarrow \text{3042}$$

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx$$

$$\downarrow \text{4205}$$

$$\int (a^2 (c + dx)^2 + 2ab(c + dx)^2 \tan(e + fx) + b^2 (c + dx)^2 \tan^2(e + fx)) dx$$

$$\downarrow \text{2009}$$

$$\frac{a^2(c+dx)^3}{3d} + \frac{2iabd(c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} +$$

$$\frac{2iab(c+dx)^3}{3d} - \frac{abd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \frac{2b^2d(c+dx) \log(1+e^{2i(e+fx)})}{f^2} +$$

$$\frac{b^2(c+dx)^2 \tan(e+fx)}{f} - \frac{ib^2(c+dx)^2}{f} - \frac{b^2(c+dx)^3}{3d} - \frac{ib^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3}$$

input `Int[(c + d*x)^2*(a + b*Tan[e + f*x])^2,x]`

output `((-I)*b^2*(c + d*x)^2)/f + (a^2*(c + d*x)^3)/(3*d) + (((2*I)/3)*a*b*(c + d*x)^3)/d - (b^2*(c + d*x)^3)/(3*d) + (2*b^2*d*(c + d*x)*Log[1 + E^((2*I)*(e + f*x))])/f^2 - (2*a*b*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f - (I*b^2*d^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^3 + ((2*I)*a*b*d*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (a*b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/f^3 + (b^2*(c + d*x)^2*Tan[e + f*x])/f`

### 3.45.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

### 3.45.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 574 vs. 2(209) = 418.

Time = 1.39 (sec) , antiderivative size = 575, normalized size of antiderivative = 2.51

method	result
risch	$\frac{8ibdcaex}{f} + \frac{d^2a^2x^3}{3} + \frac{a^2c^3}{3d} - \frac{d^2b^2x^3}{3} - b^2c^2x - \frac{b^2c^3}{3d} + \frac{4b^2ed^2 \ln(e^{i(fx+e)})}{f^3} + da^2cx^2 + a^2c^2x - db^2cx^2 -$

---

3.45.  $\int (c + dx)^2(a + b \tan(e + fx))^2 dx$



```
input int((d*x+c)^2*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 8*I/f*b*d*c*a*e*x+1/3*d^2*a^2*x^3+1/3/d*a^2*c^3-1/3*d^2*b^2*x^3-b^2*c^2*x-
1/3/d*b^2*c^3+4/f^3*b^2*e*d^2*ln(exp(I*(f*x+e)))-2/f*b*a*c^2*ln(exp(2*I*(f
*x+e))+1)+d*a^2*c*x^2+a^2*c^2*x-d*b^2*c*x^2-2*I*a*b*c^2*x-2/3*I/d*a*b*c^3+
2*I*b^2*(d^2*x^2+2*c*d*x+c^2)/f/(exp(2*I*(f*x+e))+1)+4/f*b*a*c^2*ln(exp(I*
(f*x+e)))+2/f^2*b^2*c*d*ln(exp(2*I*(f*x+e))+1)-4/f^2*b^2*c*d*ln(exp(I*(f*x
+e)))+2/f^2*b^2*d^2*ln(exp(2*I*(f*x+e))+1)*x-2*I/f*b^2*d^2*x^2-2*I/f^3*b^2
*d^2*e^2+2/3*I*d^2*a*b*x^3-a*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+2*I*d*
a*b*c*x^2-8/f^2*b*e*a*c*d*ln(exp(I*(f*x+e)))-4/f*b*d*c*a*ln(exp(2*I*(f*x+e
))+1)*x-4*I/f^2*b*a*d^2*e^2*x+2*I/f^2*b*a*d^2*polylog(2,-exp(2*I*(f*x+e)))
*x+2*I/f^2*b*d*c*a*polylog(2,-exp(2*I*(f*x+e)))+4*I/f^2*b*d*c*a*e^2-I*b^2*
d^2*polylog(2,-exp(2*I*(f*x+e)))/f^3+4/f^3*b*e^2*a*d^2*ln(exp(I*(f*x+e)))-
2/f*b*a*d^2*ln(exp(2*I*(f*x+e))+1)*x^2-8/3*I/f^3*b*a*d^2*e^3-4*I/f^2*b^2*d
^2*e*x
```

### 3.45.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 450 vs.  $2(203) = 406$ .

Time = 0.26 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.97

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx$$

$$= \frac{2(a^2 - b^2)d^2 f^3 x^3 + 6(a^2 - b^2)cdf^3 x^2 + 6(a^2 - b^2)c^2 f^3 x - 3abd^2 \operatorname{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3}{}$$

```
input integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="fracas")
```

```
output 1/6*(2*(a^2 - b^2)*d^2*f^3*x^3 + 6*(a^2 - b^2)*c*d*f^3*x^2 + 6*(a^2 - b^2)
*c^2*f^3*x - 3*a*b*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/
(tan(f*x + e)^2 + 1)) - 3*a*b*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x
+ e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(2*I*a*b*d^2*f*x + 2*I*a*b*c*d*f - I*
b^2*d^2)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 3*(-2*I*
a*b*d^2*f*x - 2*I*a*b*c*d*f + I*b^2*d^2)*dilog(2*(-I*tan(f*x + e) - 1)/(ta
n(f*x + e)^2 + 1) + 1) - 6*(a*b*d^2*f^2*x^2 + a*b*c^2*f^2 - b^2*c*d*f + (2
*a*b*c*d*f^2 - b^2*d^2*f)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 +
1)) - 6*(a*b*d^2*f^2*x^2 + a*b*c^2*f^2 - b^2*c*d*f + (2*a*b*c*d*f^2 - b^2
*d^2*f)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(b^2*d^2
*f^2*x^2 + 2*b^2*c*d*f^2*x + b^2*c^2*f^2)*tan(f*x + e))/f^3
```

### 3.45.6 Sympy [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^2 dx$$

```
input integrate((d*x+c)**2*(a+b*tan(f*x+e))**2,x)
```

```
output Integral((a + b*tan(e + f*x))**2*(c + d*x)**2, x)
```

### 3.45.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1263 vs.  $2(203) = 406$ .

Time = 0.91 (sec) , antiderivative size = 1263, normalized size of antiderivative = 5.52

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

```
input integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

output

```

1/3*(3*(f*x + e)*a^2*c^2 + (f*x + e)^3*a^2*d^2/f^2 - 3*(f*x + e)^2*a^2*d^2
*e/f^2 + 3*(f*x + e)*a^2*d^2*e^2/f^2 + 3*(f*x + e)^2*a^2*c*d/f - 6*(f*x +
e)*a^2*c*d*e/f + 6*a*b*c^2*log(sec(f*x + e)) + 6*a*b*d^2*e^2*log(sec(f*x +
e))/f^2 - 12*a*b*c*d*e*log(sec(f*x + e))/f + 3*((2*a*b + I*b^2)*(f*x + e)
^3*d^2 + 6*b^2*d^2*e^2 - 12*b^2*c*d*e*f + 6*b^2*c^2*f^2 - 3*((2*a*b + I*b^
2)*d^2*e - (2*a*b + I*b^2)*c*d*f)*(f*x + e)^2 + 3*(I*b^2*d^2*e^2 - 2*I*b^2
*c*d*e*f + I*b^2*c^2*f^2)*(f*x + e) - 6*((f*x + e)^2*a*b*d^2 + b^2*d^2*e -
b^2*c*d*f - (2*a*b*d^2*e - 2*a*b*c*d*f + b^2*d^2)*(f*x + e) + ((f*x + e)^
2*a*b*d^2 + b^2*d^2*e - b^2*c*d*f - (2*a*b*d^2*e - 2*a*b*c*d*f + b^2*d^2)*
(f*x + e))*cos(2*f*x + 2*e) - (-I*(f*x + e)^2*a*b*d^2 - I*b^2*d^2*e + I*b^
2*c*d*f + (2*I*a*b*d^2*e - 2*I*a*b*c*d*f + I*b^2*d^2)*(f*x + e))*sin(2*f*x
+ 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + ((2*a*b + I*b^2
)*(f*x + e)^3*d^2 - 3*(2*b^2*d^2 + (2*a*b + I*b^2)*d^2*e - (2*a*b + I*b^2)
*c*d*f)*(f*x + e)^2 + 3*(I*b^2*d^2*e^2 + I*b^2*c^2*f^2 + 4*b^2*d^2*e + 2*(
-I*b^2*c*d*e - 2*b^2*c*d)*f)*(f*x + e))*cos(2*f*x + 2*e) + 3*(2*(f*x + e)*
a*b*d^2 - 2*a*b*d^2*e + 2*a*b*c*d*f - b^2*d^2 + (2*(f*x + e)*a*b*d^2 - 2*a
*b*d^2*e + 2*a*b*c*d*f - b^2*d^2)*cos(2*f*x + 2*e) + (2*I*(f*x + e)*a*b*d^
2 - 2*I*a*b*d^2*e + 2*I*a*b*c*d*f - I*b^2*d^2)*sin(2*f*x + 2*e))*dilog(-e^
(2*I*f*x + 2*I*e)) + 3*(I*(f*x + e)^2*a*b*d^2 + I*b^2*d^2*e - I*b^2*c*d*f
+ (-2*I*a*b*d^2*e + 2*I*a*b*c*d*f - I*b^2*d^2)*(f*x + e) + (I*(f*x + e)...

```

### 3.45.8 Giac [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (dx + c)^2 (b \tan(fx + e) + a)^2 dx$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)^2*(b*tan(f*x + e) + a)^2, x)`

**3.45.9 Mupad [F(-1)]**

Timed out.

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^2 dx$$

input `int((a + b*tan(e + f*x))^2*(c + d*x)^2,x)`output `int((a + b*tan(e + f*x))^2*(c + d*x)^2, x)`

### 3.46 $\int (c + dx)(a + b \tan(e + fx))^2 dx$

3.46.1	Optimal result . . . . .	316
3.46.2	Mathematica [A] (verified) . . . . .	316
3.46.3	Rubi [A] (verified) . . . . .	317
3.46.4	Maple [A] (verified) . . . . .	318
3.46.5	Fricas [A] (verification not implemented) . . . . .	319
3.46.6	Sympy [F] . . . . .	319
3.46.7	Maxima [B] (verification not implemented) . . . . .	319
3.46.8	Giac [F] . . . . .	320
3.46.9	Mupad [F(-1)] . . . . .	320

#### 3.46.1 Optimal result

Integrand size = 18, antiderivative size = 136

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = -b^2cx - \frac{1}{2}b^2dx^2 + \frac{a^2(c + dx)^2}{2d} + \frac{iab(c + dx)^2}{d} - \frac{2ab(c + dx) \log(1 + e^{2i(e+fx)})}{f} + \frac{b^2d \log(\cos(e + fx))}{f^2} + \frac{iabd \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \frac{b^2(c + dx) \tan(e + fx)}{f}$$

output

```
-b^2*c*x-1/2*b^2*d*x^2+1/2*a^2*(d*x+c)^2/d+I*a*b*(d*x+c)^2/d-2*a*b*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f+b^2*d*ln(cos(f*x+e))/f^2+I*a*b*d*polylog(2,-exp(2*I*(f*x+e)))/f^2+b^2*(d*x+c)*tan(f*x+e)/f
```

#### 3.46.2 Mathematica [A] (verified)

Time = 7.41 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.47

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \frac{\cos(e + fx) (\cos(e + fx) (-((e + fx) (-2iabd(e + fx) + a^2(de - 2cf - dfx) + b^2(-de + 2cf + dfx))) -$$

input `Integrate[(c + d*x)*(a + b*Tan[e + f*x])^2,x]`

output `(Cos[e + f*x]*(Cos[e + f*x]*(-(e + f*x)*((-2*I)*a*b*d*(e + f*x) + a^2*(d*e - 2*c*f - d*f*x) + b^2*(-(d*e) + 2*c*f + d*f*x))) - 4*a*b*d*(e + f*x)*Log[1 + E^((2*I)*(e + f*x))] + 2*b*(b*d + 2*a*d*e - 2*a*c*f)*Log[Cos[e + f*x]]) + (2*I)*a*b*d*Cos[e + f*x]*PolyLog[2, -E^((2*I)*(e + f*x))] + 2*b^2*f*(c + d*x)*Sin[e + f*x]*(a + b*Tan[e + f*x])^2)/(2*f^2*(a*Cos[e + f*x] + b*SIN[e + f*x])^2)`

### 3.46.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c + dx)(a + b \tan(e + fx))^2 dx \\ & \quad \downarrow \text{3042} \\ & \int (c + dx)(a + b \tan(e + fx))^2 dx \\ & \quad \downarrow \text{4205} \\ & \int (a^2(c + dx) + 2ab(c + dx) \tan(e + fx) + b^2(c + dx) \tan^2(e + fx)) dx \\ & \quad \downarrow \text{2009} \\ & \frac{a^2(c + dx)^2}{2d} - \frac{2ab(c + dx) \log(1 + e^{2i(e+fx)})}{f} + \frac{iab(c + dx)^2}{d} + \frac{iabd \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \\ & \quad \frac{b^2(c + dx) \tan(e + fx)}{f} - \frac{b^2(c + dx)^2}{2d} + \frac{b^2d \log(\cos(e + fx))}{f^2} \end{aligned}$$

input `Int[(c + d*x)*(a + b*Tan[e + f*x])^2,x]`

```
output (a^2*(c + d*x)^2)/(2*d) + (I*a*b*(c + d*x)^2)/d - (b^2*(c + d*x)^2)/(2*d)
- (2*a*b*(c + d*x)*Log[1 + E^((2*I)*(e + f*x))])/f + (b^2*d*Log[Cos[e + f*
x]])/f^2 + (I*a*b*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (b^2*(c + d*x)
*Tan[e + f*x])/f
```

### 3.46.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4205 Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

### 3.46.4 Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.75

method	result
risch	$-\frac{b^2 dx^2}{2} + \frac{4ibadex}{f} + \frac{a^2 dx^2}{2} - b^2 cx - 2iabcx + a^2 cx + \frac{2ibade^2}{f^2} + \frac{b^2 d \ln(e^{2i(fx+e)}+1)}{f^2} - \frac{2b^2 d \ln(e^{i(fx+e)})}{f^2} - \dots$

```
input int((d*x+c)*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output -1/2*b^2*d*x^2+4*I/f*b*a*d*e*x+1/2*a^2*d*x^2-b^2*c*x-2*I*a*b*c*x+a^2*c*x+2
*I/f^2*b*a*d*e^2+1/f^2*b^2*d*ln(exp(2*I*(f*x+e))+1)-2/f^2*b^2*d*ln(exp(I*(
f*x+e)))-2/f*b*a*c*ln(exp(2*I*(f*x+e))+1)+4/f*b*a*c*ln(exp(I*(f*x+e)))-4/f
^2*b*e*a*d*ln(exp(I*(f*x+e)))+2*I*b^2*(d*x+c)/f/(exp(2*I*(f*x+e))+1)+I*a*b
*d*polylog(2,-exp(2*I*(f*x+e)))/f^2+I*a*b*d*x^2-2/f*b*a*d*ln(exp(2*I*(f*x+
e))+1)*x
```

**3.46.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.61

$$\int (c + dx)(a + b \tan(e + fx))^2 dx$$

$$= \frac{(a^2 - b^2)df^2x^2 + 2(a^2 - b^2)cf^2x - iabd\text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + iabd\text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) - (2ab$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output `1/2*((a^2 - b^2)*d*f^2*x^2 + 2*(a^2 - b^2)*c*f^2*x - I*a*b*d*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) + I*a*b*d*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - (2*a*b*d*f*x + 2*a*b*c*f - b^2*d)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - (2*a*b*d*f*x + 2*a*b*c*f - b^2*d)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 2*(b^2*d*f*x + b^2*c*f)*tan(f*x + e))/f^2`

**3.46.6 Sympy [F]**

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx) dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))**2,x)`

output `Integral((a + b*tan(e + f*x))**2*(c + d*x), x)`

**3.46.7 Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 527 vs.  $2(123) = 246$ .

Time = 0.41 (sec) , antiderivative size = 527, normalized size of antiderivative = 3.88

$$\int (c + dx)(a + b \tan(e + fx))^2 dx$$

$$= \frac{2(fx + e)a^2c + \frac{(fx+e)^2a^2d}{f} - \frac{2(fx+e)a^2de}{f} + 4abc \log(\sec(fx + e)) - \frac{4abde \log(\sec(fx+e))}{f} + \frac{2((2ab+ib^2)(fx+e)^2d$$

---

3.46.  $\int (c + dx)(a + b \tan(e + fx))^2 dx$



input `integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output `1/2*(2*(f*x + e)*a^2*c + (f*x + e)^2*a^2*d/f - 2*(f*x + e)*a^2*d*e/f + 4*a*b*c*log(sec(f*x + e)) - 4*a*b*d*e*log(sec(f*x + e))/f + 2*((2*a*b + I*b^2)*(f*x + e)^2*d - 4*b^2*d*e + 4*b^2*c*f + 2*(-I*b^2*d*e + I*b^2*c*f)*(f*x + e) - 2*(2*(f*x + e)*a*b*d - b^2*d + (2*(f*x + e)*a*b*d - b^2*d)*cos(2*f*x + 2*e) - (-2*I*(f*x + e)*a*b*d + I*b^2*d)*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + ((2*a*b + I*b^2)*(f*x + e)^2*d + 2*(-I*b^2*d*e + I*b^2*c*f - 2*b^2*d)*(f*x + e))*cos(2*f*x + 2*e) + 2*(a*b*d*cos(2*f*x + 2*e) + I*a*b*d*sin(2*f*x + 2*e) + a*b*d)*dilog(-e^(2*I*f*x + 2*I*e)) - (-2*I*(f*x + e)*a*b*d + I*b^2*d + (-2*I*(f*x + e)*a*b*d + I*b^2*d)*cos(2*f*x + 2*e) + (2*(f*x + e)*a*b*d - b^2*d)*sin(2*f*x + 2*e))*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) - ((-2*I*a*b + b^2)*(f*x + e)^2*d - 2*(b^2*d*e - b^2*c*f - 2*I*b^2*d)*(f*x + e))*sin(2*f*x + 2*e))/(-2*I*f*cos(2*f*x + 2*e) + 2*f*sin(2*f*x + 2*e) - 2*I*f)/f`

### 3.46.8 Giac [F]

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (dx + c)(b \tan(fx + e) + a)^2 dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)*(b*tan(f*x + e) + a)^2, x)`

### 3.46.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx) dx$$

input `int((a + b*tan(e + f*x))^2*(c + d*x),x)`

output `int((a + b*tan(e + f*x))^2*(c + d*x), x)`

**3.47**  $\int \frac{(a+b \tan(e+fx))^2}{c+dx} dx$

3.47.1 Optimal result . . . . . 321  
 3.47.2 Mathematica [N/A] . . . . . 321  
 3.47.3 Rubi [N/A] . . . . . 322  
 3.47.4 Maple [N/A] (verified) . . . . . 323  
 3.47.5 Fricas [N/A] . . . . . 323  
 3.47.6 Sympy [N/A] . . . . . 323  
 3.47.7 Maxima [N/A] . . . . . 324  
 3.47.8 Giac [N/A] . . . . . 324  
 3.47.9 Mupad [N/A] . . . . . 325

**3.47.1 Optimal result**

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^2}{c + dx}, x\right)$$

output `Unintegrable((a+b*tan(f*x+e))^2/(d*x+c),x)`

**3.47.2 Mathematica [N/A]**

Not integrable

Time = 19.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

input `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x),x]`

output `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x), x]`

### 3.47.3 Rubi [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

↓ 4223

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

input `Int[(a + b*Tan[e + f*x])^2/(c + d*x),x]`

output `$Aborted`

#### 3.47.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

**3.47.4 Maple [N/A] (verified)**

Not integrable

Time = 0.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^2}{dx + c} dx$$

input `int((a+b*tan(f*x+e))^2/(d*x+c),x)`output `int((a+b*tan(f*x+e))^2/(d*x+c),x)`**3.47.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^2}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="fricas")`output `integral((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(d*x + c), x)`**3.47.6 Sympy [N/A]**

Not integrable

Time = 1.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

input `integrate((a+b*tan(f*x+e))**2/(d*x+c),x)`output `Integral((a + b*tan(e + f*x))**2/(c + d*x), x)`

**3.47.7 Maxima [N/A]**

Not integrable

Time = 0.83 (sec) , antiderivative size = 491, normalized size of antiderivative = 24.55

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^2}{dx + c} dx$$

```
input integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="maxima")
```

```
output (((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e)^2*log(d*x + c) + 2
*b^2*d*sin(2*f*x + 2*e) + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*log(d*x +
c)*sin(2*f*x + 2*e)^2 + 2*((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x
+ 2*e)*log(d*x + c) + (d^2*f*x + c*d*f + (d^2*f*x + c*d*f)*cos(2*f*x + 2*e
)^2 + (d^2*f*x + c*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^2*f*x + c*d*f)*cos(2*f*x
+ 2*e))*integrate(2*(2*a*b*d*f*x + 2*a*b*c*f + b^2*d)*sin(2*f*x + 2*e)/(d
^2*f*x^2 + 2*c*d*f*x + c^2*f + (d^2*f*x^2 + 2*c*d*f*x + c^2*f)*cos(2*f*x +
2*e)^2 + (d^2*f*x^2 + 2*c*d*f*x + c^2*f)*sin(2*f*x + 2*e)^2 + 2*(d^2*f*x^
2 + 2*c*d*f*x + c^2*f)*cos(2*f*x + 2*e)), x) + ((a^2 - b^2)*d*f*x + (a^2 -
b^2)*c*f)*log(d*x + c))/(d^2*f*x + c*d*f + (d^2*f*x + c*d*f)*cos(2*f*x +
2*e)^2 + (d^2*f*x + c*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^2*f*x + c*d*f)*cos(2*
f*x + 2*e))
```

**3.47.8 Giac [N/A]**

Not integrable

Time = 0.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^2}{dx + c} dx$$

```
input integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="giac")
```

```
output integrate((b*tan(f*x + e) + a)^2/(d*x + c), x)
```

**3.47.9 Mupad [N/A]**

Not integrable

Time = 4.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + f x))^2}{c + d x} dx = \int \frac{(a + b \tan(e + f x))^2}{c + d x} dx$$

input `int((a + b*tan(e + f*x))^2/(c + d*x), x)`output `int((a + b*tan(e + f*x))^2/(c + d*x), x)`

$$3.48 \quad \int \frac{(a+b \tan(e+fx))^2}{(c+dx)^2} dx$$

3.48.1	Optimal result	326
3.48.2	Mathematica [N/A]	326
3.48.3	Rubi [N/A]	327
3.48.4	Maple [N/A] (verified)	328
3.48.5	Fricas [N/A]	328
3.48.6	Sympy [N/A]	328
3.48.7	Maxima [N/A]	329
3.48.8	Giac [N/A]	329
3.48.9	Mupad [N/A]	330

### 3.48.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^2}{(c + dx)^2}, x\right)$$

output `Unintegrable((a+b*tan(f*x+e))^2/(d*x+c)^2,x)`

### 3.48.2 Mathematica [N/A]

Not integrable

Time = 14.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

input `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x)^2,x]`

output `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x)^2, x]`

### 3.48.3 Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

↓ 4223

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

input `Int[(a + b*Tan[e + f*x])^2/(c + d*x)^2,x]`

output `$Aborted`

#### 3.48.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`



**3.48.4 Maple [N/A] (verified)**

Not integrable

Time = 0.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^2}{(dx + c)^2} dx$$

input `int((a+b*tan(f*x+e))^2/(d*x+c)^2,x)`output `int((a+b*tan(f*x+e))^2/(d*x+c)^2,x)`**3.48.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="fracas")`output `integral((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(d^2*x^2 + 2*c*d*x + c^2), x)`**3.48.6 Sympy [N/A]**

Not integrable

Time = 2.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

input `integrate((a+b*tan(f*x+e))**2/(d*x+c)**2,x)`output `Integral((a + b*tan(e + f*x))**2/(c + d*x)**2, x)`

---

3.48.  $\int \frac{(a+b \tan(e+fx))^2}{(c+dx)^2} dx$

**3.48.7 Maxima [N/A]**

Not integrable

Time = 1.68 (sec) , antiderivative size = 609, normalized size of antiderivative = 30.45

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

```
input integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="maxima")
```

```
output -((a^2 - b^2)*d*f*x - 2*b^2*d*sin(2*f*x + 2*e) + (a^2 - b^2)*c*f + ((a^2 -
  b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e)^2 + ((a^2 - b^2)*d*f*x + (
  a^2 - b^2)*c*f)*sin(2*f*x + 2*e)^2 + 2*((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*
  f)*cos(2*f*x + 2*e) - (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f + (d^3*f*x^2 + 2*
  c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d
  *f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x +
  2*e))*integrate(4*(a*b*d*f*x + a*b*c*f + b^2*d)*sin(2*f*x + 2*e)/(d^3*f*x
  ^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f + (d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*
  c^2*d*f*x + c^3*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2
  *d*f*x + c^3*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*
  d*f*x + c^3*f)*cos(2*f*x + 2*e)), x)/(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f +
  (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^2 + 2*c
  *d^2*f*x + c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^2 + 2*c*d^2*f*x + c^2*
  d*f)*cos(2*f*x + 2*e))
```

**3.48.8 Giac [N/A]**

Not integrable

Time = 15.43 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

```
input integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="giac")
```

```
output integrate((b*tan(f*x + e) + a)^2/(d*x + c)^2, x)
```

**3.48.9 Mupad [N/A]**

Not integrable

Time = 3.64 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

input `int((a + b*tan(e + f*x))^2/(c + d*x)^2,x)`output `int((a + b*tan(e + f*x))^2/(c + d*x)^2, x)`

### 3.49 $\int (c + dx)^3 (a + b \tan(e + fx))^3 dx$

3.49.1	Optimal result . . . . .	332
3.49.2	Mathematica [B] (warning: unable to verify) . . . . .	333
3.49.3	Rubi [A] (verified) . . . . .	334
3.49.4	Maple [B] (verified) . . . . .	336
3.49.5	Fricas [B] (verification not implemented) . . . . .	337
3.49.6	Sympy [F] . . . . .	338
3.49.7	Maxima [B] (verification not implemented) . . . . .	339
3.49.8	Giac [F] . . . . .	339
3.49.9	Mupad [F(-1)] . . . . .	340

### 3.49.1 Optimal result

Integrand size = 20, antiderivative size = 612

$$\begin{aligned}
 \int (c + dx)^3 (a + b \tan(e + fx))^3 dx = & \frac{3ib^3 d(c + dx)^2}{2f^2} - \frac{3iab^2(c + dx)^3}{f} + \frac{b^3(c + dx)^3}{2f} \\
 & + \frac{a^3(c + dx)^4}{4d} + \frac{3ia^2b(c + dx)^4}{4d} - \frac{3ab^2(c + dx)^4}{4d} \\
 & - \frac{ib^3(c + dx)^4}{4d} - \frac{3b^3 d^2(c + dx) \log(1 + e^{2i(e+fx)})}{f^3} \\
 & + \frac{9ab^2 d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2 b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
 & + \frac{b^3(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
 & + \frac{3ib^3 d^3 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} \\
 & - \frac{9iab^2 d^2(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
 & + \frac{9ia^2 b d(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
 & - \frac{3ib^3 d(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
 & + \frac{9ab^2 d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \\
 & - \frac{9a^2 b d^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & + \frac{3b^3 d^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & - \frac{9ia^2 b d^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
 & + \frac{3ib^3 d^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
 & - \frac{3b^3 d(c + dx)^2 \tan(e + fx)}{2f^2} \\
 & + \frac{3ab^2(c + dx)^3 \tan(e + fx)}{f} \\
 & + \frac{b^3(c + dx)^3 \tan^2(e + fx)}{2f}
 \end{aligned}$$

output 
$$\begin{aligned} & -1/4*I*b^3*(d*x+c)^4/d+3/4*I*b^3*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4+1/2* \\ & b^3*(d*x+c)^3/f+1/4*a^3*(d*x+c)^4/d+3/4*I*a^2*b*(d*x+c)^4/d-3/4*a*b^2*(d*x \\ & +c)^4/d-3*I*a*b^2*(d*x+c)^3/f-3*b^3*d^2*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^3 \\ & +9*a*b^2*d*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f^2-3*a^2*b*(d*x+c)^3*ln(1+exp \\ & (2*I*(f*x+e)))/f+b^3*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f+3/2*I*b^3*d^3*poly \\ & log(2,-exp(2*I*(f*x+e)))/f^4-9/4*I*a^2*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/ \\ & f^4+9/2*I*a^2*b*d*(d*x+c)^2*polylog(2,-exp(2*I*(f*x+e)))/f^2-9*I*a*b^2*d^2 \\ & *(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^3+9/2*a*b^2*d^3*polylog(3,-exp(2*I \\ & *(f*x+e)))/f^4-9/2*a^2*b*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3+3/2* \\ & b^3*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3-3/2*I*b^3*d*(d*x+c)^2*pol \\ & ylog(2,-exp(2*I*(f*x+e)))/f^2+3/2*I*b^3*d*(d*x+c)^2/f^2-3/2*b^3*d*(d*x+c)^ \\ & 2*tan(f*x+e)/f^2+3*a*b^2*(d*x+c)^3*tan(f*x+e)/f+1/2*b^3*(d*x+c)^3*tan(f*x+ \\ & e)^2/f \end{aligned}$$

### 3.49.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2594 vs.  $2(612) = 1224$ .

Time = 7.57 (sec) , antiderivative size = 2594, normalized size of antiderivative = 4.24

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \text{Result too large to show}$$

input `Integrate[(c + d*x)^3*(a + b*Tan[e + f*x])^3,x]`

output

```

(((3*I)/4)*a*b^2*d^3*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))])*Sec[e])/(E^(I*e)*f^4) - (((3*I)/4)*a^2*b*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))])*Sec[e])/(E^(I*e)*f^3) + ((I/4)*b^3*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))])*Sec[e])/(E^(I*e)*f^3) - (((3*I)/8)*a^2*b*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))] + 6*(1 + E^((-2*I)*e))*f^2*x^2*PolyLog[2, -E^((-2*I)*(e + f*x))] - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))])*Sec[e])/f^4 + ((I/8)*b^3*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))] + 6*(1 + E^((-2*I)*e))*f^2*x^2*PolyLog[2, -E^((-2*I)*(e + f*x))] - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))])*Sec[e])/f^4 + ((b^3*c^3 + 3*b^3*c^2*d*x + 3*b^3*c*d^2*x^2 + b^3*d^3*x^3)*Sec[e + f*x]^2)/(2*f) - (3*b^3*c*d^2*Sec[e]*(Cos[e]*Log[Cos[e...

```

### 3.49.3 Rubi [A] (verified)

Time = 1.28 (sec) , antiderivative size = 612, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^3 (a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c + dx)^3 (a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \text{4205} \\
 & \int (a^3 (c + dx)^3 + 3a^2 b (c + dx)^3 \tan(e + fx) + 3ab^2 (c + dx)^3 \tan^2(e + fx) + b^3 (c + dx)^3 \tan^3(e + fx)) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.49.  $\int (c + dx)^3 (a + b \tan(e + fx))^3 dx$

$$\begin{aligned}
& \frac{a^3(c+dx)^4}{4d} - \frac{9a^2bd^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{9ia^2bd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \\
& \frac{3a^2b(c+dx)^3 \log(1 + e^{2i(e+fx)})}{f} + \frac{3ia^2b(c+dx)^4}{4d} - \frac{9ia^2bd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} - \\
& \frac{9iab^2d^2(c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{9ab^2d(c+dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} + \\
& \frac{3ab^2(c+dx)^3 \tan(e+fx)}{f} - \frac{3iab^2(c+dx)^3}{f} - \frac{3ab^2(c+dx)^4}{4d} + \frac{9ab^2d^3 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} + \\
& \frac{3b^3d^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{3b^3d^2(c+dx) \log(1 + e^{2i(e+fx)})}{f^3} - \\
& \frac{3ib^3d(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3b^3d(c+dx)^2 \tan(e+fx)}{2f^2} + \\
& \frac{b^3(c+dx)^3 \log(1 + e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^3 \tan^2(e+fx)}{2f} + \frac{3ib^3d(c+dx)^2}{2f^2} + \frac{b^3(c+dx)^3}{2f} - \\
& \frac{ib^3(c+dx)^4}{4d} + \frac{3ib^3d^3 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} + \frac{3ib^3d^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}
\end{aligned}$$

input `Int[(c + d*x)^3*(a + b*Tan[e + f*x])^3,x]`

output

```

(((3*I)/2)*b^3*d*(c + d*x)^2)/f^2 - ((3*I)*a*b^2*(c + d*x)^3)/f + (b^3*(c
+ d*x)^3)/(2*f) + (a^3*(c + d*x)^4)/(4*d) + (((3*I)/4)*a^2*b*(c + d*x)^4)/
d - (3*a*b^2*(c + d*x)^4)/(4*d) - ((I/4)*b^3*(c + d*x)^4)/d - (3*b^3*d^2*(
c + d*x)*Log[1 + E^((2*I)*(e + f*x))])/f^3 + (9*a*b^2*d*(c + d*x)^2*Log[1
+ E^((2*I)*(e + f*x))])/f^2 - (3*a^2*b*(c + d*x)^3*Log[1 + E^((2*I)*(e + f
*x))])/f + (b^3*(c + d*x)^3*Log[1 + E^((2*I)*(e + f*x))])/f + (((3*I)/2)*b
^3*d^3*PolyLog[2, -E^((2*I)*(e + f*x))])/f^4 - ((9*I)*a*b^2*d^2*(c + d*x)*
PolyLog[2, -E^((2*I)*(e + f*x))])/f^3 + (((9*I)/2)*a^2*b*d*(c + d*x)^2*Pol
yLog[2, -E^((2*I)*(e + f*x))])/f^2 - (((3*I)/2)*b^3*d*(c + d*x)^2*PolyLog[
2, -E^((2*I)*(e + f*x))])/f^2 + (9*a*b^2*d^3*PolyLog[3, -E^((2*I)*(e + f*x
))])/(2*f^4) - (9*a^2*b*d^2*(c + d*x)*PolyLog[3, -E^((2*I)*(e + f*x))])/(2
*f^3) + (3*b^3*d^2*(c + d*x)*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (
((9*I)/4)*a^2*b*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4 + (((3*I)/4)*b^3
*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4 - (3*b^3*d*(c + d*x)^2*Tan[e +
f*x])/(2*f^2) + (3*a*b^2*(c + d*x)^3*Tan[e + f*x])/f + (b^3*(c + d*x)^3*Ta
n[e + f*x]^2)/(2*f)

```



## 3.49.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

## 3.49.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1929 vs.  $2(544) = 1088$ .

Time = 1.21 (sec) , antiderivative size = 1930, normalized size of antiderivative = 3.15

method	result	size
risch	Expression too large to display	1930

input `int((d*x+c)^3*(a+b*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output

```

-1/4*I*b^3*d^3*x^4+d^2*a^3*c*x^3+3/2*d*a^3*c^2*x^2+a^3*c^3*x-3/4*d^3*a*b^2
*x^4-3*a*b^2*c^3*x-3/4/d*a*b^2*c^4+I*b^3*c^3*x+1/4*I/d*b^3*c^4+1/4*d^3*a^3
*x^4+1/4/d*c^4*a^3-36*I/f^2*b^2*d^2*c*a*e*x+3/4*I*b^3*d^3*polylog(4,-exp(2
*I*(f*x+e)))/f^4+1/f*b^3*c^3*ln(exp(2*I*(f*x+e))+1)-2/f*b^3*c^3*ln(exp(I*(
f*x+e))+b^2*(-6*I*c*d^2*x*b-6*I*b*c*d^2*x*exp(2*I*(f*x+e))+2*b*d^3*f*x^3*
exp(2*I*(f*x+e))+18*I*a*c*d^2*f*x^2-3*I*c^2*d*b-3*I*d^3*x^2*b+6*b*c*d^2*f*
x^2*exp(2*I*(f*x+e))-3*I*b*d^3*x^2*exp(2*I*(f*x+e))+18*I*a*c*d^2*f*x^2*exp
(2*I*(f*x+e))+6*I*a*d^3*f*x^3*exp(2*I*(f*x+e))+6*b*c^2*d*f*x*exp(2*I*(f*x+
e))-3*I*b*c^2*d*exp(2*I*(f*x+e))+6*I*a*c^3*f*exp(2*I*(f*x+e))+18*I*a*c^2*d
*f*x+2*b*c^3*f*exp(2*I*(f*x+e))+18*I*a*c^2*d*f*x*exp(2*I*(f*x+e))+6*I*a*d^
3*f*x^3+6*I*a*c^3*f)/f^2/(exp(2*I*(f*x+e))+1)^2+3/4*I*d^3*a^2*b*x^4-3/2*I*
b^3*d*c^2*x^2-18*I/f^2*b*a^2*c*d^2*e^2*x+9*I/f^2*b*a^2*c*d^2*polylog(2,-ex
p(2*I*(f*x+e)))*x+18*I/f*b*d*c^2*a^2*e*x-6/f^4*b^3*e*d^3*ln(exp(I*(f*x+e))
)+3*I/f^2*b^3*d^3*x^2+3*I/f^4*b^3*d^3*e^2-3/2*I/f^4*b^3*d^3*e^4-9/4*I*a^2*
b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4-I*d^2*b^3*c*x^3-3*d^2*a*b^2*c*x^3-9
/2*d*a*b^2*c^2*x^2-3*I*a^2*b*c^3*x-3/4*I/d*a^2*b*c^4+3/2*I*b^3*d^3*polylog
(2,-exp(2*I*(f*x+e)))/f^4-3/f^3*b^3*c*d^2*ln(exp(2*I*(f*x+e))+1)+6/f^3*b^3
*c*d^2*ln(exp(I*(f*x+e)))+1/f*b^3*d^3*ln(exp(2*I*(f*x+e))+1)*x^3+3/2/f^3*b
^3*c*d^2*polylog(3,-exp(2*I*(f*x+e)))-3/f*b*a^2*c^3*ln(exp(2*I*(f*x+e))+1)
+6/f*b*a^2*c^3*ln(exp(I*(f*x+e)))-3/f^3*b^3*d^3*ln(exp(2*I*(f*x+e))+1)*...

```

### 3.49.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1177 vs.  $2(530) = 1060$ .

Time = 0.27 (sec) , antiderivative size = 1177, normalized size of antiderivative = 1.92

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

output `1/8*(2*(a^3 - 3*a*b^2)*d^3*f^4*x^4 + 3*I*(3*a^2*b - b^3)*d^3*polylog(4, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*I*(3*a^2*b - b^3)*d^3*polylog(4, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 4*(b^3*d^3*f^3 + 2*(a^3 - 3*a*b^2)*c*d^2*f^4)*x^3 + 12*(b^3*c*d^2*f^3 + (a^3 - 3*a*b^2)*c^2*d*f^4)*x^2 + 4*(b^3*d^3*f^3*x^3 + 3*b^3*c*d^2*f^3*x^2 + 3*b^3*c^2*d*f^3*x + b^3*c^3*f^3)*tan(f*x + e)^2 + 4*(3*b^3*c^2*d*f^3 + 2*(a^3 - 3*a*b^2)*c^3*f^4)*x - 6*(I*(3*a^2*b - b^3)*d^3*f^2*x^2 - 6*I*a*b^2*c*d^2*f + I*b^3*d^3 + I*(3*a^2*b - b^3)*c^2*d*f^2 - 2*I*(3*a*b^2*d^3*f - (3*a^2*b - b^3)*c*d^2*f^2)*x)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(-I*(3*a^2*b - b^3)*d^3*f^2*x^2 + 6*I*a*b^2*c*d^2*f - I*b^3*d^3 - I*(3*a^2*b - b^3)*c^2*d*f^2 + 2*I*(3*a*b^2*d^3*f - (3*a^2*b - b^3)*c*d^2*f^2)*x)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 4*((3*a^2*b - b^3)*d^3*f^3*x^3 - 9*a*b^2*c^2*d*f^2 + 3*b^3*c*d^2*f + (3*a^2*b - b^3)*c^3*f^3 - 3*(3*a*b^2*d^3*f^2 - (3*a^2*b - b^3)*c*d^2*f^3)*x^2 - 3*(6*a*b^2*c*d^2*f^2 - b^3*d^3*f - (3*a^2*b - b^3)*c^2*d*f^3)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 4*((3*a^2*b - b^3)*d^3*f^3*x^3 - 9*a*b^2*c^2*d*f^2 + 3*b^3*c*d^2*f + (3*a^2*b - b^3)*c^3*f^3 - 3*(3*a*b^2*d^3*f^2 - (3*a^2*b - b^3)*c*d^2*f^3)*x^2 - 3*(6*a*b^2*c*d^2*f^2 - b^3*d^3*f - (3*a^2*b - b^3)*c^2*d*f^3)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(3*a*b^2*d^3 - (3*a^2*b - b^3)*d^3*f*x - (3*a...`

### 3.49.6 Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^3 dx$$

input `integrate((d*x+c)**3*(a+b*tan(f*x+e))**3,x)`

output `Integral((a + b*tan(e + f*x))**3*(c + d*x)**3, x)`

### 3.49.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 6861 vs.  $2(530) = 1060$ .

Time = 10.88 (sec) , antiderivative size = 6861, normalized size of antiderivative = 11.21

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output

```

1/4*(4*(f*x + e)*a^3*c^3 + (f*x + e)^4*a^3*d^3/f^3 - 4*(f*x + e)^3*a^3*d^3
*e/f^3 + 6*(f*x + e)^2*a^3*d^3*e^2/f^3 - 4*(f*x + e)*a^3*d^3*e^3/f^3 + 4*(
f*x + e)^3*a^3*c*d^2/f^2 - 12*(f*x + e)^2*a^3*c*d^2*e/f^2 + 12*(f*x + e)*a
^3*c*d^2*e^2/f^2 + 6*(f*x + e)^2*a^3*c^2*d/f - 12*(f*x + e)*a^3*c^2*d*e/f
+ 12*a^2*b*c^3*log(sec(f*x + e)) - 12*a^2*b*d^3*e^3*log(sec(f*x + e))/f^3
+ 36*a^2*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 36*a^2*b*c^2*d*e*log(sec(f*x
+ e))/f - 4*(72*a*b^2*d^3*e^3 - 72*a*b^2*c^3*f^3 - 3*(3*a^2*b + 3*I*a*b^2
- b^3)*(f*x + e)^4*d^3 + 36*b^3*d^3*e^2 + 12*((3*a^2*b + 3*I*a*b^2 - b^3)*
d^3*e - (3*a^2*b + 3*I*a*b^2 - b^3)*c*d^2*f)*(f*x + e)^3 - 18*((3*a^2*b +
3*I*a*b^2 - b^3)*d^3*e^2 - 2*(3*a^2*b + 3*I*a*b^2 - b^3)*c*d^2*e*f + (3*a^
2*b + 3*I*a*b^2 - b^3)*c^2*d*f^2)*(f*x + e)^2 + 36*(6*a*b^2*c^2*d*e + b^3*
c^2*d)*f^2 + 12*((3*I*a*b^2 - b^3)*d^3*e^3 + 3*(-3*I*a*b^2 + b^3)*c*d^2*e^
2*f + 3*(3*I*a*b^2 - b^3)*c^2*d*e*f^2 + (-3*I*a*b^2 + b^3)*c^3*f^3)*(f*x +
e) - 72*(3*a*b^2*c*d^2*e^2 + b^3*c*d^2*e)*f + 4*(3*b^3*d^3*e^3 - 3*b^3*c^
3*f^3 - 27*a*b^2*d^3*e^2 + 4*(3*a^2*b - b^3)*(f*x + e)^3*d^3 - 9*b^3*d^3*e
- 9*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3*e - (3*a^2*b - b^3)*c*d^2*f)*(f*x
+ e)^2 + 9*(b^3*c^2*d*e - 3*a*b^2*c^2*d)*f^2 + 9*(6*a*b^2*d^3*e + b^3*d^3
+ (3*a^2*b - b^3)*d^3*e^2 + (3*a^2*b - b^3)*c^2*d*f^2 - 2*(3*a*b^2*c*d^2 +
(3*a^2*b - b^3)*c*d^2*e)*f)*(f*x + e) - 9*(b^3*c*d^2*e^2 - 6*a*b^2*c*d^2*
e - b^3*c*d^2)*f + (3*b^3*d^3*e^3 - 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + ...

```

### 3.49.8 Giac [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \int (dx + c)^3 (b \tan(fx + e) + a)^3 dx$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((d*x + c)^3*(b*tan(f*x + e) + a)^3, x)`

**3.49.9 Mupad [F(-1)]**

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^3 dx$$

input `int((a + b*tan(e + f*x))^3*(c + d*x)^3,x)`output `int((a + b*tan(e + f*x))^3*(c + d*x)^3, x)`

### 3.50 $\int (c + dx)^2 (a + b \tan(e + fx))^3 dx$

3.50.1	Optimal result	342
3.50.2	Mathematica [B] (warning: unable to verify)	343
3.50.3	Rubi [A] (verified)	344
3.50.4	Maple [B] (verified)	346
3.50.5	Fricas [A] (verification not implemented)	347
3.50.6	Sympy [F]	347
3.50.7	Maxima [B] (verification not implemented)	348
3.50.8	Giac [F]	348
3.50.9	Mupad [F(-1)]	349

### 3.50.1 Optimal result

Integrand size = 20, antiderivative size = 436

$$\begin{aligned}
 \int (c + dx)^2 (a + b \tan(e + fx))^3 dx = & \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3iab^2(c + dx)^2}{f} + \frac{a^3(c + dx)^3}{3d} \\
 & + \frac{ia^2b(c + dx)^3}{d} - \frac{ab^2(c + dx)^3}{d} - \frac{ib^3(c + dx)^3}{3d} \\
 & + \frac{6ab^2d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\
 & + \frac{b^3(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\
 & - \frac{b^3d^2 \log(\cos(e + fx))}{f^3} \\
 & - \frac{3iab^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
 & + \frac{3ia^2bd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & - \frac{ib^3d(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & + \frac{b^3d^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & - \frac{b^3d(c + dx) \tan(e + fx)}{f^2} \\
 & + \frac{3ab^2(c + dx)^2 \tan(e + fx)}{f} \\
 & + \frac{b^3(c + dx)^2 \tan^2(e + fx)}{2f}
 \end{aligned}$$

output  $b^3 c d x / f + 1/2 b^3 d^2 x^2 / f + 3 I a^2 b d (d x + c) \operatorname{polylog}(2, -\exp(2 I (f x + e))) / f^2 + 1/3 a^3 (d x + c)^3 / d - 3 I a b^2 d^2 \operatorname{polylog}(2, -\exp(2 I (f x + e))) / f^3 - a b^2 (d x + c)^3 / d + I a^2 b (d x + c)^3 / d + 6 a b^2 d (d x + c) \ln(1 + \exp(2 I (f x + e))) / f^2 - 3 a^2 b (d x + c)^2 \ln(1 + \exp(2 I (f x + e))) / f + b^3 (d x + c)^2 \ln(1 + \exp(2 I (f x + e))) / f - b^3 d^2 \ln(\cos(f x + e)) / f^3 - 3 I a b^2 (d x + c)^2 / f - I b^3 d (d x + c) \operatorname{polylog}(2, -\exp(2 I (f x + e))) / f^2 - 1/3 I b^3 (d x + c)^3 / d - 3/2 a^2 b d^2 \operatorname{polylog}(3, -\exp(2 I (f x + e))) / f^3 + 1/2 b^3 d^2 \operatorname{polylog}(3, -\exp(2 I (f x + e))) / f^3 - b^3 d (d x + c) \tan(f x + e) / f^2 + 3 a b^2 (d x + c)^2 \tan(f x + e) / f + 1/2 b^3 (d x + c)^2 \tan(f x + e)^2 / f$

### 3.50.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1846 vs.  $2(436) = 872$ .

Time = 7.16 (sec) , antiderivative size = 1846, normalized size of antiderivative = 4.23

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `Integrate[(c + d*x)^2*(a + b*Tan[e + f*x])^3,x]`



output

```

((-1/4*I)*a^2*b*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^
((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e +
f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))])*Sec[e]
)/(E^(I*e)*f^3) + ((I/12)*b^3*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*
e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E
^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e +
f*x))])*Sec[e]/(E^(I*e)*f^3) - (b^3*d^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x]
] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^3*(Cos[e]^2 + Sin[e]^2)) + (6*a*b^2
*c*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/
(f^2*(Cos[e]^2 + Sin[e]^2)) - (3*a^2*b*c^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f
*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (b^3*c^2
*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(
Cos[e]^2 + Sin[e]^2)) + (3*a*b^2*d^2*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]]
) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]]) - Pi*Log[1 + E^((-2*I)*f*x)] -
2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]])])) + Pi*L
og[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]]) + I*PolyLog
[2, E^((2*I)*(f*x - ArcTan[Cot[e]])]))/Sqrt[1 + Cot[e]^2])*Sec[e]/(f^3*S
qrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)) - (3*a^2*b*c*d*Csc[e]*((f^2*x^2)/E^(I
*ArcTan[Cot[e]]) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]]) - Pi*Log[1 + E^
((-2*I)*f*x)] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan...

```

### 3.50.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 427, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^2 (a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c + dx)^2 (a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \text{4205} \\
 & \int (a^3 (c + dx)^2 + 3a^2 b (c + dx)^2 \tan(e + fx) + 3ab^2 (c + dx)^2 \tan^2(e + fx) + b^3 (c + dx)^2 \tan^3(e + fx)) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.50.  $\int (c + dx)^2 (a + b \tan(e + fx))^3 dx$

$$\begin{aligned} & \frac{a^3(c+dx)^3}{3d} + \frac{3ia^2bd(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{3a^2b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} + \\ & \frac{ia^2b(c+dx)^3}{d} - \frac{3a^2bd^2\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{6ab^2d(c+dx)\log(1+e^{2i(e+fx)})}{f^2} + \\ & \frac{3ab^2(c+dx)^2 \tan(e+fx)}{f} - \frac{3iab^2(c+dx)^2}{f} - \frac{ab^2(c+dx)^3}{d} - \frac{3iab^2d^2\text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} - \\ & \frac{ib^3d(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{b^3d(c+dx)\tan(e+fx)}{f} + \frac{b^3(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^3} + \\ & \frac{b^3(c+dx)^2 \tan^2(e+fx)}{2f} + \frac{b^3(c+dx)^2}{2f} - \frac{ib^3(c+dx)^3}{3d} + \frac{b^3d^2\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \\ & \frac{b^3d^2 \log(\cos(e+fx))}{f^3} \end{aligned}$$

input `Int[(c + d*x)^2*(a + b*Tan[e + f*x])^3,x]`

output `((-3*I)*a*b^2*(c + d*x)^2)/f + (b^3*(c + d*x)^2)/(2*f) + (a^3*(c + d*x)^3)/(3*d) + (I*a^2*b*(c + d*x)^3)/d - (a*b^2*(c + d*x)^3)/d - ((I/3)*b^3*(c + d*x)^3)/d + (6*a*b^2*d*(c + d*x)*Log[1 + E^((2*I)*(e + f*x))])/f^2 - (3*a^2*b*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f + (b^3*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f - (b^3*d^2*Log[Cos[e + f*x]])/f^3 - ((3*I)*a*b^2*d^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^3 + ((3*I)*a^2*b*d*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (I*b^3*d*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (3*a^2*b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) + (b^3*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (b^3*d*(c + d*x)*Tan[e + f*x])/f^2 + (3*a*b^2*(c + d*x)^2*Tan[e + f*x])/f + (b^3*(c + d*x)^2*Tan[e + f*x]^2)/(2*f)`

### 3.50.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

### 3.50.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1137 vs.  $2(402) = 804$ .

Time = 1.06 (sec) , antiderivative size = 1138, normalized size of antiderivative = 2.61

method	result	size
risch	Expression too large to display	1138

input `int((d*x+c)^2*(a+b*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output

```

3*I*d*a^2*b*c*x^2+12*I/f*b*a^2*c*d*e*x+6/f^2*b^2*a*c*d*ln(exp(2*I*(f*x+e))
+1)-12/f^2*b^2*a*c*d*ln(exp(I*(f*x+e)))+4/f^2*b^3*e*d*c*ln(exp(I*(f*x+e)))
+6/f^2*b^2*a*d^2*ln(exp(2*I*(f*x+e))+1)*x+2/f*b^3*c*d*ln(exp(2*I*(f*x+e))+
1)*x+6/f^3*b*e^2*d^2*a^2*ln(exp(I*(f*x+e)))-3/f*b*d^2*a^2*ln(exp(2*I*(f*x+
e))+1)*x^2+12/f^3*b^2*e*a*d^2*ln(exp(I*(f*x+e)))-6*I/f^3*b^2*a*d^2*e^2-I/f
^2*b^3*d^2*polylog(2,-exp(2*I*(f*x+e)))*x-2*I/f^2*b^3*c*d*e^2-I/f^2*b^3*c
d*polylog(2,-exp(2*I*(f*x+e)))+2*I/f^2*b^3*d^2*e^2*x-6*I/f*b^2*a*d^2*x^2-4
*I/f^3*b*d^2*a^2*e^3+1/f*b^3*d^2*ln(exp(2*I*(f*x+e))+1)*x^2-3/f*b*a^2*c^2*
ln(exp(2*I*(f*x+e))+1)+6/f*b*a^2*c^2*ln(exp(I*(f*x+e)))-2/f^3*b^3*e^2*d^2*
ln(exp(I*(f*x+e)))+4/3*I/f^3*b^3*d^2*e^3-I*d*b^3*c*x^2-3*d*a*b^2*c*x^2+I*d
^2*a^2*b*x^3-3*I*a^2*b*c^2*x-I/d*a^2*b*c^3+d*a^3*c*x^2+a^3*c^2*x-d^2*a*b^2
*x^3-3*a*b^2*c^2*x-1/d*a*b^2*c^3+I*b^3*c^2*x+1/3*I/d*b^3*c^3+1/3*d^2*a^3*x
^3+1/3/d*c^3*a^3-1/3*I*b^3*d^2*x^3+1/f*b^3*c^2*ln(exp(2*I*(f*x+e))+1)-2/f*
b^3*c^2*ln(exp(I*(f*x+e)))-1/f^3*b^3*d^2*ln(exp(2*I*(f*x+e))+1)+2/f^3*b^3*
d^2*ln(exp(I*(f*x+e)))+2*b^2*(3*I*a*d^2*f*x^2*exp(2*I*(f*x+e))+6*I*a*c*d*f
*x*exp(2*I*(f*x+e))+b*d^2*f*x^2*exp(2*I*(f*x+e))+3*I*a*c^2*f*exp(2*I*(f*x+
e))+3*I*a*d^2*f*x^2-I*b*d^2*x*exp(2*I*(f*x+e))+2*b*c*d*f*x*exp(2*I*(f*x+e)
)+6*I*a*c*d*f*x-I*b*c*d*exp(2*I*(f*x+e))+b*c^2*f*exp(2*I*(f*x+e))+3*I*a*c^
2*f-I*d^2*x*b-I*c*d*b)/f^2/(exp(2*I*(f*x+e))+1)^2-3*I*a*b^2*d^2*polylog(2,
-exp(2*I*(f*x+e)))/f^3-3/2*a^2*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+1...

```

### 3.50.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 688, normalized size of antiderivative = 1.58

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx$$

$$= \frac{4(a^3 - 3ab^2)d^2 f^3 x^3 - 3(3a^2b - b^3)d^2 \operatorname{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3(3a^2b - b^3)d^2 \operatorname{polylog}\left(3, \frac{\tan(fx+e)^2 - 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) + 6(b^3 d^2 f^2 + 2(a^3 - 3ab^2)c d f^3)x^2 + 6(b^3 d^2 f^2 x^2 + 2b^3 c d f^2 x + b^3 c^2 f^2) \tan(fx + e)^2 + 12(b^3 c d f^2 + (a^3 - 3ab^2)c^2 f^3)x - 6(-3I a b^2 d^2 + I(3a^2 b - b^3)d^2 f x + I(3a^2 b - b^3)c d f) \operatorname{dilog}\left(2 \frac{I \tan(fx + e) - 1}{\tan(fx + e)^2 + 1} + 1\right) - 6(3I a b^2 d^2 - I(3a^2 b - b^3)d^2 f x - I(3a^2 b - b^3)c d f) \operatorname{dilog}\left(2 \frac{-I \tan(fx + e) - 1}{\tan(fx + e)^2 + 1} + 1\right) - 6((3a^2 b - b^3)d^2 f^2 x^2 - 6a b^2 c d f + b^3 d^2 + (3a^2 b - b^3)c^2 f^2 - 2(3a b^2 d^2 f - (3a^2 b - b^3)c d f^2)x) \log(-2 \frac{I \tan(fx + e) - 1}{\tan(fx + e)^2 + 1}) - 6((3a^2 b - b^3)d^2 f^2 x^2 - 6a b^2 c d f + b^3 d^2 + (3a^2 b - b^3)c^2 f^2 - 2(3a b^2 d^2 f - (3a^2 b - b^3)c d f^2)x) \log(-2 \frac{-I \tan(fx + e) - 1}{\tan(fx + e)^2 + 1}) + 12(3a b^2 d^2 f^2 x^2 + 3a b^2 c^2 f^2 - b^3 c d f + (6a b^2 c d f^2 - b^3 d^2 f)x) \tan(fx + e)}{f^3}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="fracas")`

output

```
1/12*(4*(a^3 - 3*a*b^2)*d^2*f^3*x^3 - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(b^3*d^2*f^2 + 2*(a^3 - 3*a*b^2)*c*d*f^3)*x^2 + 6*(b^3*d^2*f^2*x^2 + 2*b^3*c*d*f^2*x + b^3*c^2*f^2)*tan(f*x + e)^2 + 12*(b^3*c*d*f^2 + (a^3 - 3*a*b^2)*c^2*f^3)*x - 6*(-3*I*a*b^2*d^2 + I*(3*a^2*b - b^3)*d^2*f*x + I*(3*a^2*b - b^3)*c*d*f)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(3*I*a*b^2*d^2 - I*(3*a^2*b - b^3)*d^2*f*x - I*(3*a^2*b - b^3)*c*d*f)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*((3*a^2*b - b^3)*d^2*f^2*x^2 - 6*a*b^2*c*d*f + b^3*d^2 + (3*a^2*b - b^3)*c^2*f^2 - 2*(3*a*b^2*d^2*f - (3*a^2*b - b^3)*c*d*f^2)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*((3*a^2*b - b^3)*d^2*f^2*x^2 - 6*a*b^2*c*d*f + b^3*d^2 + (3*a^2*b - b^3)*c^2*f^2 - 2*(3*a*b^2*d^2*f - (3*a^2*b - b^3)*c*d*f^2)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 12*(3*a*b^2*d^2*f^2*x^2 + 3*a*b^2*c^2*f^2 - b^3*c*d*f + (6*a*b^2*c*d*f^2 - b^3*d^2*f)*x)*tan(f*x + e)/f^3
```

### 3.50.6 Sympy [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^2 dx$$

input `integrate((d*x+c)**2*(a+b*tan(f*x+e))**3,x)`

output `Integral((a + b*tan(e + f*x))**3*(c + d*x)**2, x)`

### 3.50.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3407 vs.  $2(393) = 786$ .

Time = 1.97 (sec) , antiderivative size = 3407, normalized size of antiderivative = 7.81

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output

```
1/3*(3*(f*x + e)*a^3*c^2 + (f*x + e)^3*a^3*d^2/f^2 - 3*(f*x + e)^2*a^3*d^2
*e/f^2 + 3*(f*x + e)*a^3*d^2*e^2/f^2 + 3*(f*x + e)^2*a^3*c*d/f - 6*(f*x +
e)*a^3*c*d*e/f + 9*a^2*b*c^2*log(sec(f*x + e)) + 9*a^2*b*d^2*e^2*log(sec(f
*x + e))/f^2 - 18*a^2*b*c*d*e*log(sec(f*x + e))/f + 3*(36*a*b^2*d^2*e^2 +
36*a*b^2*c^2*f^2 + 2*(3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^3*d^2 + 12*b^3*
d^2*e - 6*((3*a^2*b + 3*I*a*b^2 - b^3)*d^2*e - (3*a^2*b + 3*I*a*b^2 - b^3)
*c*d*f)*(f*x + e)^2 - 6*((-3*I*a*b^2 + b^3)*d^2*e^2 + 2*(3*I*a*b^2 - b^3)*
c*d*e*f + (-3*I*a*b^2 + b^3)*c^2*f^2)*(f*x + e) - 12*(6*a*b^2*c*d*e + b^3*
c*d)*f + 6*(b^3*d^2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f
*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*
b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f + (b^3*d^2*e^2 +
b^3*c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 +
2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e)
- 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*cos(4*f*x + 4*e) + 2*(b^3*d^2*e^2 + b^3*
c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3
*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*
(b^3*c*d*e - 3*a*b^2*c*d)*f)*cos(2*f*x + 2*e) - (-I*b^3*d^2*e^2 - I*b^3*c^
2*f^2 + 6*I*a*b^2*d^2*e + (3*I*a^2*b - I*b^3)*(f*x + e)^2*d^2 + I*b^3*d^2
+ 2*(-3*I*a*b^2*d^2 + (-3*I*a^2*b + I*b^3)*d^2*e + (3*I*a^2*b - I*b^3)*c*d
*f)*(f*x + e) + 2*(I*b^3*c*d*e - 3*I*a*b^2*c*d)*f)*sin(4*f*x + 4*e) - 2...
```

### 3.50.8 Giac [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \int (dx + c)^2 (b \tan(fx + e) + a)^3 dx$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((d*x + c)^2*(b*tan(f*x + e) + a)^3, x)`

---

3.50.  $\int (c + dx)^2 (a + b \tan(e + fx))^3 dx$

**3.50.9 Mupad [F(-1)]**

Timed out.

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^2 dx$$

input `int((a + b*tan(e + f*x))^3*(c + d*x)^2,x)`output `int((a + b*tan(e + f*x))^3*(c + d*x)^2, x)`

### 3.51 $\int (c + dx)(a + b \tan(e + fx))^3 dx$

3.51.1	Optimal result . . . . .	350
3.51.2	Mathematica [A] (verified) . . . . .	351
3.51.3	Rubi [A] (verified) . . . . .	351
3.51.4	Maple [A] (verified) . . . . .	352
3.51.5	Fricas [A] (verification not implemented) . . . . .	353
3.51.6	Sympy [F] . . . . .	354
3.51.7	Maxima [B] (verification not implemented) . . . . .	354
3.51.8	Giac [F] . . . . .	355
3.51.9	Mupad [F(-1)] . . . . .	355

#### 3.51.1 Optimal result

Integrand size = 18, antiderivative size = 277

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx))^3 dx = & -3ab^2cx + \frac{b^3dx}{2f} - \frac{3}{2}ab^2dx^2 + \frac{a^3(c + dx)^2}{2d} \\ & + \frac{3ia^2b(c + dx)^2}{2d} - \frac{ib^3(c + dx)^2}{2d} \\ & - \frac{3a^2b(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{b^3(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{3ab^2d \log(\cos(e + fx))}{f^2} \\ & + \frac{3ia^2bd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & - \frac{ib^3d \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{b^3d \tan(e + fx)}{2f^2} \\ & + \frac{3ab^2(c + dx) \tan(e + fx)}{f} + \frac{b^3(c + dx) \tan^2(e + fx)}{2f} \end{aligned}$$

output

```
-3*a*b^2*c*x+1/2*b^3*d*x/f-3/2*a*b^2*d*x^2+1/2*a^3*(d*x+c)^2/d+3/2*I*a^2*b
*(d*x+c)^2/d-1/2*I*b^3*(d*x+c)^2/d-3*a^2*b*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/
f+b^3*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f+3*a*b^2*d*ln(cos(f*x+e))/f^2+3/2*I*
a^2*b*d*polylog(2,-exp(2*I*(f*x+e)))/f^2-1/2*I*b^3*d*polylog(2,-exp(2*I*(f
*x+e)))/f^2-1/2*b^3*d*tan(f*x+e)/f^2+3*a*b^2*(d*x+c)*tan(f*x+e)/f+1/2*b^3*
(d*x+c)*tan(f*x+e)^2/f
```

### 3.51.2 Mathematica [A] (verified)

Time = 8.32 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.00

$$\int (c + dx)(a + b \tan(e + fx))^3 dx$$

$$= \frac{\cos(e + fx) (\cos^2(e + fx) (-((e + fx) (-3ia^2bd(e + fx) + ib^3d(e + fx) + 3ab^2(-de + 2cf + dfx) + a^3($$

input `Integrate[(c + d*x)*(a + b*Tan[e + f*x])^3,x]`

output `(Cos[e + f*x]*(Cos[e + f*x]^2*(-((e + f*x)*((-3*I)*a^2*b*d*(e + f*x) + I*b^3*d*(e + f*x) + 3*a*b^2*(-(d*e) + 2*c*f + d*f*x) + a^3*(-2*c*f + d*(e - f*x)))) + 2*b*(-3*a^2 + b^2)*d*(e + f*x)*Log[1 + E^((2*I)*(e + f*x))] + 2*b*(3*a*b*d + 3*a^2*(d*e - c*f) + b^2*(-(d*e) + c*f))*Log[Cos[e + f*x]]) - I*b*(-3*a^2 + b^2)*d*Cos[e + f*x]^2*PolyLog[2, -E^((2*I)*(e + f*x))] + (b^2*(2*b*f*(c + d*x) + (-b*d) + 6*a*f*(c + d*x))*Sin[2*(e + f*x)]))/2*(a + b*Tan[e + f*x])^3)/(2*f^2*(a*Cos[e + f*x] + b*SIN[e + f*x])^3)`

### 3.51.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)(a + b \tan(e + fx))^3 dx$$

$$\downarrow \text{3042}$$

$$\int (c + dx)(a + b \tan(e + fx))^3 dx$$

$$\downarrow \text{4205}$$

$$\int (a^3(c + dx) + 3a^2b(c + dx) \tan(e + fx) + 3ab^2(c + dx) \tan^2(e + fx) + b^3(c + dx) \tan^3(e + fx)) dx$$

$$\downarrow \text{2009}$$



$$\frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)\log(1+e^{2i(e+fx)})}{f} + \frac{3ia^2b(c+dx)^2}{2d} + \frac{3ia^2bd\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} + \frac{3ab^2(c+dx)\tan(e+fx)}{f} - \frac{3ab^2(c+dx)^2}{2d} + \frac{3ab^2d\log(\cos(e+fx))}{f^2} + \frac{b^3(c+dx)\log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)\tan^2(e+fx)}{2f} - \frac{ib^3(c+dx)^2}{2d} - \frac{ib^3d\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{b^3d\tan(e+fx)}{2f^2} + \frac{b^3dx}{2f}$$

input `Int[(c + d*x)*(a + b*Tan[e + f*x])^3,x]`

output  $(b^3d*x)/(2*f) + (a^3*(c + d*x)^2)/(2*d) + (((3*I)/2)*a^2*b*(c + d*x)^2)/d - (3*a*b^2*(c + d*x)^2)/(2*d) - ((I/2)*b^3*(c + d*x)^2)/d - (3*a^2*b*(c + d*x)*\text{Log}[1 + E^{((2*I)*(e + f*x))}])/f + (b^3*(c + d*x)*\text{Log}[1 + E^{((2*I)*(e + f*x))}])/f + (3*a*b^2*d*\text{Log}[\text{Cos}[e + f*x]])/f^2 + (((3*I)/2)*a^2*b*d*\text{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 - ((I/2)*b^3*d*\text{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 - (b^3*d*\text{Tan}[e + f*x])/(2*f^2) + (3*a*b^2*(c + d*x)*\text{Tan}[e + f*x])/f + (b^3*(c + d*x)*\text{Tan}[e + f*x]^2)/(2*f)$

### 3.51.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

### 3.51.4 Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 493, normalized size of antiderivative = 1.78

method	result
risch	$-3ia^2bcx + \frac{b^3d\ln(e^{2i(fx+e)}+1)x}{f} + \frac{3b^2ad\ln(e^{2i(fx+e)}+1)}{f^2} - \frac{6b^2ad\ln(e^{i(fx+e)})}{f^2} - \frac{3ba^2c\ln(e^{2i(fx+e)}+1)}{f} + \frac{6ba^2c\ln(e^{i(fx+e)})}{f}$

3.51.  $\int (c+dx)(a+b\tan(e+fx))^3 dx$

```
input int((d*x+c)*(a+b*tan(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/2*a^3*d*x^2+a^3*c*x+I*b^3*c*x-3*I*a^2*b*c*x+1/f*b^3*c*ln(exp(2*I*(f*x+e)
)+1)-2/f*b^3*c*ln(exp(I*(f*x+e)))-1/2*I*b^3*d*x^2+1/f*b^3*d*ln(exp(2*I*(f*
x+e))+1)*x+3/f^2*b^2*a*d*ln(exp(2*I*(f*x+e))+1)-6/f^2*b^2*a*d*ln(exp(I*(f*
x+e)))-3/f*b*a^2*c*ln(exp(2*I*(f*x+e))+1)+6/f*b*a^2*c*ln(exp(I*(f*x+e)))+2
/f^2*b^3*e*d*ln(exp(I*(f*x+e)))-I/f^2*b^3*d*e^2+3/2*I*a^2*b*d*x^2+b^2*(6*I
*a*d*f*x*exp(2*I*(f*x+e))+6*I*a*c*f*exp(2*I*(f*x+e))+2*b*d*f*x*exp(2*I*(f*
x+e))+6*I*a*d*f*x-I*b*d*exp(2*I*(f*x+e))+2*b*c*f*exp(2*I*(f*x+e))+6*I*a*c*
f-I*b*d)/f^2/(exp(2*I*(f*x+e))+1)^2+3/2*I*a^2*b*d*polylog(2,-exp(2*I*(f*x+
e)))/f^2-1/2*I*b^3*d*polylog(2,-exp(2*I*(f*x+e)))/f^2-3*a*b^2*c*x-3/2*a*b^
2*d*x^2-6/f^2*b*e*a^2*d*ln(exp(I*(f*x+e)))-3/f*b*a^2*d*ln(exp(2*I*(f*x+e)
+1))*x-2*I/f*b^3*d*e*x+3*I/f^2*b*a^2*d*e^2+6*I/f*b*a^2*d*e*x
```

### 3.51.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.18

$$\int (c + dx)(a + b \tan(e + fx))^3 dx$$

$$= \frac{2(a^3 - 3ab^2)df^2x^2 - i(3a^2b - b^3)d\text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + i(3a^2b - b^3)d\text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + \dots}{1}$$

```
input integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="fracas")
```

```
output 1/4*(2*(a^3 - 3*a*b^2)*d*f^2*x^2 - I*(3*a^2*b - b^3)*d*dilog(2*(I*tan(f*x
+ e) - 1)/(tan(f*x + e)^2 + 1) + 1) + I*(3*a^2*b - b^3)*d*dilog(2*(-I*tan(
f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) + 2*(b^3*d*f*x + b^3*c*f)*tan(f*x
+ e)^2 + 2*(b^3*d*f + 2*(a^3 - 3*a*b^2)*c*f^2)*x + 2*(3*a*b^2*d - (3*a^2*b
- b^3)*d*f*x - (3*a^2*b - b^3)*c*f)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x
+ e)^2 + 1)) + 2*(3*a*b^2*d - (3*a^2*b - b^3)*d*f*x - (3*a^2*b - b^3)*c*f)
*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 2*(6*a*b^2*d*f*x + 6
*a*b^2*c*f - b^3*d)*tan(f*x + e))/f^2
```

### 3.51.6 Sympy [F]

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx) dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))**3,x)`

output `Integral((a + b*tan(e + f*x))**3*(c + d*x), x)`

### 3.51.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1319 vs.  $2(241) = 482$ .

Time = 0.67 (sec) , antiderivative size = 1319, normalized size of antiderivative = 4.76

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output `1/2*(2*(f*x + e)*a^3*c + (f*x + e)^2*a^3*d/f - 2*(f*x + e)*a^3*d*e/f + 6*a^2*b*c*log(sec(f*x + e)) - 6*a^2*b*d*e*log(sec(f*x + e))/f - 2*(12*a*b^2*d*e - 12*a*b^2*c*f - (3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d + 2*b^3*d + 2*((3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f)*(f*x + e) + 2*(b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d + (b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*cos(4*f*x + 4*e) + 2*(b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*cos(2*f*x + 2*e) + (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*sin(4*f*x + 4*e) + 2*(I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)) + 1) - ((3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d - 2*(6*a*b^2*d + (3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f)*(f*x + e))*cos(4*f*x + 4*e) - 2*((3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d - b^3*d - 2*(3*a*b^2 - I*b^3)*d*e + 2*(3*a*b^2 - I*b^3)*c*f - 2*((3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f + (3*a*b^2 + I*b^3)*d)*(f*x + e))*cos(2*f*x + 2*e) - ((3*a^2*b - b^3)*d*cos(4*f*x + 4*e) + 2*(3*a^2*b - b^3)*d*cos(2*f*x + 2*e) + (3*I*a^2*b - I*b^3)*d*sin(4*f*x + 4*e) - 2*(-3*I*a^2*b + I*b^3)*d*sin(2*f*x + 2*e) + (3*a^2*b - b^3)*d)*dilog(-e^(2*I*f*x + 2*I*e)) - (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d + (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*cos(4*f*x + 4*e) - 2*(-I*b...`

**3.51.8 Giac [F]**

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (dx + c)(b \tan(fx + e) + a)^3 dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((d*x + c)*(b*tan(f*x + e) + a)^3, x)`

**3.51.9 Mupad [F(-1)]**

Timed out.

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx) dx$$

input `int((a + b*tan(e + f*x))^3*(c + d*x),x)`

output `int((a + b*tan(e + f*x))^3*(c + d*x), x)`

$$3.52 \quad \int \frac{(a+b \tan(e+fx))^3}{c+dx} dx$$

3.52.1	Optimal result	356
3.52.2	Mathematica [N/A]	356
3.52.3	Rubi [N/A]	357
3.52.4	Maple [N/A] (verified)	358
3.52.5	Fricas [N/A]	358
3.52.6	Sympy [N/A]	358
3.52.7	Maxima [N/A]	359
3.52.8	Giac [N/A]	359
3.52.9	Mupad [N/A]	360

### 3.52.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a+b \tan(e+fx))^3}{c+dx} dx = \text{Int}\left(\frac{(a+b \tan(e+fx))^3}{c+dx}, x\right)$$

output `Unintegrable((a+b*tan(f*x+e))^3/(d*x+c), x)`

### 3.52.2 Mathematica [N/A]

Not integrable

Time = 14.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a+b \tan(e+fx))^3}{c+dx} dx = \int \frac{(a+b \tan(e+fx))^3}{c+dx} dx$$

input `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x), x]`

output `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x), x]`

### 3.52.3 Rubi [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

↓ 4223

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

input `Int[(a + b*Tan[e + f*x])^3/(c + d*x),x]`

output `$Aborted`

#### 3.52.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

**3.52.4 Maple [N/A] (verified)**

Not integrable

Time = 0.62 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^3}{dx + c} dx$$

input `int((a+b*tan(f*x+e))^3/(d*x+c),x)`output `int((a+b*tan(f*x+e))^3/(d*x+c),x)`**3.52.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.60

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^3}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))^3/(d*x+c),x, algorithm="fricas")`output `integral((b^3*tan(f*x + e)^3 + 3*a*b^2*tan(f*x + e)^2 + 3*a^2*b*tan(f*x + e) + a^3)/(d*x + c), x)`**3.52.6 Sympy [N/A]**

Not integrable

Time = 1.33 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

input `integrate((a+b*tan(f*x+e))**3/(d*x+c),x)`output `Integral((a + b*tan(e + f*x))**3/(c + d*x), x)`





input `integrate((a+b*tan(f*x+e))^3/(d*x+c),x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)^3/(d*x + c), x)`

### 3.52.9 Mupad [N/A]

Not integrable

Time = 3.91 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + f x))^3}{c + d x} dx = \int \frac{(a + b \tan(e + f x))^3}{c + d x} dx$$

input `int((a + b*tan(e + f*x))^3/(c + d*x),x)`

output `int((a + b*tan(e + f*x))^3/(c + d*x), x)`

$$3.53 \quad \int \frac{(a+b \tan(e+fx))^3}{(c+dx)^2} dx$$

3.53.1	Optimal result	361
3.53.2	Mathematica [N/A]	361
3.53.3	Rubi [N/A]	362
3.53.4	Maple [N/A] (verified)	363
3.53.5	Fricas [N/A]	363
3.53.6	Sympy [N/A]	363
3.53.7	Maxima [N/A]	364
3.53.8	Giac [N/A]	364
3.53.9	Mupad [N/A]	365

### 3.53.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a+b \tan(e+fx))^3}{(c+dx)^2} dx = \text{Int}\left(\frac{(a+b \tan(e+fx))^3}{(c+dx)^2}, x\right)$$

output `Unintegrable((a+b*tan(f*x+e))^3/(d*x+c)^2,x)`

### 3.53.2 Mathematica [N/A]

Not integrable

Time = 17.93 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a+b \tan(e+fx))^3}{(c+dx)^2} dx = \int \frac{(a+b \tan(e+fx))^3}{(c+dx)^2} dx$$

input `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x)^2,x]`

output `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x)^2, x]`

### 3.53.3 Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

↓ 4223

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

input `Int[(a + b*Tan[e + f*x])^3/(c + d*x)^2,x]`

output `$Aborted`

#### 3.53.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.53.4 Maple [N/A] (verified)**

Not integrable

Time = 0.76 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^3}{(dx + c)^2} dx$$

input `int((a+b*tan(f*x+e))^3/(d*x+c)^2,x)`output `int((a+b*tan(f*x+e))^3/(d*x+c)^2,x)`**3.53.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 3.15

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="fracas")`output `integral((b^3*tan(f*x + e)^3 + 3*a*b^2*tan(f*x + e)^2 + 3*a^2*b*tan(f*x + e) + a^3)/(d^2*x^2 + 2*c*d*x + c^2), x)`**3.53.6 Sympy [N/A]**

Not integrable

Time = 2.44 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

input `integrate((a+b*tan(f*x+e))**3/(d*x+c)**2,x)`output `Integral((a + b*tan(e + f*x))**3/(c + d*x)**2, x)`

**3.53.7 Maxima [N/A]**

Not integrable

Time = 9.08 (sec) , antiderivative size = 2194, normalized size of antiderivative = 109.70

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

```
input integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="maxima")
```

```
output -((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2 + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*cos(4*f*x + 4*e)^2 + 4*((a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + (a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 2*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e)^2 + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*sin(4*f*x + 4*e)^2 + 4*((a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + (a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 2*(a^3 - 3*a*b^2)*c*d*f^2)*x)*sin(2*f*x + 2*e)^2 + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2 + (2*(a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + 2*(a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 4*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e) + (3*a*b^2*d^2*f*x + 3*a*b^2*c*d*f + b^3*d^2)*sin(2*f*x + 2*e))*cos(4*f*x + 4*e) + 2*(2*(a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + 2*(a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 4*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e) + (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2 + (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(2*f*x + 2*e)^2 + (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*...
```

**3.53.8 Giac [N/A]**

Not integrable

Time = 18.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)^3/(d*x + c)^2, x)`

### 3.53.9 Mupad [N/A]

Not integrable

Time = 4.93 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + f x))^3}{(c + d x)^2} dx = \int \frac{(a + b \tan(e + f x))^3}{(c + d x)^2} dx$$

input `int((a + b*tan(e + f*x))^3/(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))^3/(c + d*x)^2, x)`

### 3.54 $\int \frac{(c+dx)^3}{a+b \tan(e+fx)} dx$

3.54.1	Optimal result	366
3.54.2	Mathematica [A] (verified)	367
3.54.3	Rubi [A] (verified)	367
3.54.4	Maple [B] (verified)	371
3.54.5	Fricas [B] (verification not implemented)	372
3.54.6	Sympy [F]	372
3.54.7	Maxima [B] (verification not implemented)	373
3.54.8	Giac [F]	374
3.54.9	Mupad [F(-1)]	374

#### 3.54.1 Optimal result

Integrand size = 20, antiderivative size = 243

$$\int \frac{(c+dx)^3}{a+b \tan(e+fx)} dx = \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} - \frac{3ibd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^2} + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^3} + \frac{3ibd^3 \text{PolyLog}\left(4, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{4(a^2+b^2)f^4}$$

output  $\frac{1}{4}*(d*x+c)^4/(a+I*b)/d+b*(d*x+c)^3*\ln(1+(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-3/2*I*b*d*(d*x+c)^2*\text{polylog}(2,-(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2+3/2*b*d^2*(d*x+c)*\text{polylog}(3,-(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^3+3/4*I*b*d^3*\text{polylog}(4,-(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^4$

### 3.54.2 Mathematica [A] (verified)

Time = 1.68 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.22

$$\int \frac{(c+dx)^3}{a+b \tan(e+fx)} dx$$

$$= \frac{1}{4} b \left( -\frac{2(c+dx)^4}{(ia+b)d(-ib(-1+e^{2ie})+a(1+e^{2ie}))} + \frac{4(c+dx)^3 \log\left(1 + \frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2+b^2)f} \right)$$

$$+ \frac{3d\left(2if^2(c+dx)^2 \text{PolyLog}\left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right) + d\left(2f(c+dx) \text{PolyLog}\left(3, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right) - id \text{PolyLog}\left(4, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)\right)\right)}{(a^2+b^2)f^4}$$

$$+ \frac{x(4c^3+6c^2dx+4cd^2x^2+d^3x^3)\cos(e)}{4(a\cos(e)+b\sin(e))}$$

input `Integrate[(c + d*x)^3/(a + b*Tan[e + f*x]),x]`

output `(b*((-2*(c + d*x)^4)/((I*a + b)*d*((-I)*b*(-1 + E^((2*I)*e))) + a*(1 + E^((2*I)*e)))) + (4*(c + d*x)^3*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f) + (3*d*((2*I)*f^2*(c + d*x)^2*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] + d*(2*f*(c + d*x)*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - I*d*PolyLog[4, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a^2 + b^2)*f^4))/4 + (x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cos[e])/(4*(a*Cos[e] + b*Sin[e]))`

### 3.54.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$ , Rules used = {3042, 4215, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c+dx)^3}{a+b \tan(e+fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c+dx)^3}{a+b \tan(e+fx)} dx$$



$$\begin{aligned}
 & \downarrow \text{4215} \\
 & 2ib \int \frac{e^{2i(e+fx)}(c+dx)^3}{(a+ib)^2 + (a^2+b^2)e^{2i(e+fx)}} dx + \frac{(c+dx)^4}{4d(a+ib)} \\
 & \downarrow \text{2620} \\
 & 2ib \left( \frac{3id \int (c+dx)^2 \log \left( \frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1 \right) dx}{2f(a^2+b^2)} - \frac{i(c+dx)^3 \log \left( 1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \\
 & \qquad \qquad \qquad \frac{(c+dx)^4}{4d(a+ib)} \\
 & \downarrow \text{3011} \\
 & 2ib \left( \frac{3id \left( \frac{i(c+dx)^2 \text{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \int (c+dx) \text{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{f} \right)}{2f(a^2+b^2)} - \frac{i(c+dx)^3 \log \left( 1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \\
 & \qquad \qquad \qquad \frac{(c+dx)^4}{4d(a+ib)} \\
 & \downarrow \text{7163} \\
 & 2ib \left( \frac{3id \left( \frac{i(c+dx)^2 \text{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \left( \frac{id \int \text{PolyLog} \left( 3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{2f} - \frac{i(c+dx) \text{PolyLog} \left( 3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} \right)}{f} \right)}{2f(a^2+b^2)} \right) + \\
 & \qquad \qquad \qquad \frac{(c+dx)^4}{4d(a+ib)} \\
 & \downarrow \text{2720}
 \end{aligned}$$

$$\left. \begin{array}{l} 3id \\ 2ib \end{array} \right\} \left( \frac{i(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f} - \frac{id \left( \frac{d f e^{-2i(e+fx)} \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right) d e^{2i(e+fx)}}{4f^2} - \frac{i(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f} \right)}{f} \right)$$

$$2f(a^2 + b^2)$$

$$\frac{(c+dx)^4}{4d(a+ib)}$$

↓ 7143

$$\left. \begin{array}{l} 3id \\ 2ib \end{array} \right\} \left( \frac{i(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f} - \frac{id \left( \frac{d \operatorname{PolyLog}\left(4, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{4f^2} - \frac{i(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f} \right)}{f} \right)$$

$$2f(a^2 + b^2)$$

$$\frac{(c+dx)^4}{4d(a+ib)}$$

input `Int[(c + d*x)^3/(a + b*Tan[e + f*x]),x]`

output `(c + d*x)^4/(4*(a + I*b)*d) + (2*I)*b*((( -1/2*I)*(c + d*x)^3*Log[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b]^2)]/((a^2 + b^2)*f) + (((3*I)/2)*d*(((I/2)*(c + d*x)^2*PolyLog[2, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/f - (I*d*((( -1/2*I)*(c + d*x)*PolyLog[3, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/f) + (d*PolyLog[4, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/(4*f^2)))/f)/((a^2 + b^2)*f))`

$$3.54. \int \frac{(c+dx)^3}{a+b \tan(e+fx)} dx$$

## 3.54.3.1 Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4215 `Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Simp[2*I*b Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

```
rule 7163 Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### 3.54.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1467 vs.  $2(220) = 440$ .

Time = 0.78 (sec) , antiderivative size = 1468, normalized size of antiderivative = 6.04

method	result	size
risch	Expression too large to display	1468

```
input int((d*x+c)^3/(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 3*I/f^3/(I*a+b)*b/(-I*b-a)*e^2*c*d^2*ln(1-(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a
))-3*I/f/(I*a+b)*b/(-I*b-a)*d*c^2*ln(1-(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a))*
x-3*I/f^2/(I*a+b)*b/(-I*b-a)*d*c^2*ln(1-(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a)
)*e-3*I/f^2/(I*a+b)*b*e*d*c^2/(a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*
x+e))-I*b-a)-6*I/f^3/(I*a+b)*b*e^2*c*d^2/(a+I*b)*ln(exp(I*(f*x+e)))+3*I/f^
3/(I*a+b)*b*e^2*c*d^2/(a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I
*b-a)-1/2/(I*a+b)*b/(-I*b-a)*d^3*x^4-3/2/f^4/(I*a+b)*b/(-I*b-a)*d^3*e^4+3/
4/f^4/(I*a+b)*b/(-I*b-a)*d^3*polylog(4,(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a))-
3/(I*a+b)*b/(-I*b-a)*d*c^2*x^2-2/(I*a+b)*b/(-I*b-a)*d^2*c*x^3-3*I/f/(I*a+b
)*b/(-I*b-a)*d^2*c*ln(1-(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a))*x^2+6*I/f^2/(I*
a+b)*b*e*d*c^2/(a+I*b)*ln(exp(I*(f*x+e)))-1/4/d/(I*b-a)*c^4-d^2/(I*b-a)*c*
x^3-3/2*d/(I*b-a)*c^2*x^2-1/4*d^3/(I*b-a)*x^4-1/(I*b-a)*c^3*x+6/f^2/(I*a+b
)*b/(-I*b-a)*e^2*c*d^2*x-3/f^2/(I*a+b)*b/(-I*b-a)*d^2*c*polylog(2,(a-I*b)*
exp(2*I*(f*x+e)))/(-I*b-a))*x-6/f/(I*a+b)*b/(-I*b-a)*d*c^2*e*x-I/f/(I*a+b)*
b/(-I*b-a)*d^3*ln(1-(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a))*x^3-I/f^4/(I*a+b)*b
/(-I*b-a)*e^3*d^3*ln(1-(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a))-3/2*I/f^3/(I*a+b
)*b/(-I*b-a)*d^3*polylog(3,(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a))*x-3/2*I/f^3/
(I*a+b)*b/(-I*b-a)*d^2*c*polylog(3,(a-I*b)*exp(2*I*(f*x+e)))/(-I*b-a))+2*I/
f^4/(I*a+b)*b*e^3*d^3/(a+I*b)*ln(exp(I*(f*x+e)))-I/f^4/(I*a+b)*b*e^3*d^3/(
a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I*b-a)-2/f^3/(I*a+b)*...
```

### 3.54.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1183 vs.  $2(212) = 424$ .

Time = 0.28 (sec) , antiderivative size = 1183, normalized size of antiderivative = 4.87

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

```
input integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="fracas")
```

```
output 1/8*(2*a*d^3*f^4*x^4 + 8*a*c*d^2*f^4*x^3 + 12*a*c^2*d*f^4*x^2 + 8*a*c^3*f^4*x - 3*I*b*d^3*polylog(4, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 3*I*b*d^3*polylog(4, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 6*(-I*b*d^3*f^2*x^2 - 2*I*b*c*d^2*f^2*x - I*b*c^2*d*f^2)*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*b*d^3*f^2*x^2 + 2*I*b*c*d^2*f^2*x + I*b*c^2*d*f^2)*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - 4*(b*d^3*e^3 - 3*b*c...
```

### 3.54.6 Sympy [F]

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx$$

```
input integrate((d*x+c)**3/(a+b*tan(f*x+e)),x)
```

output `Integral((c + d*x)**3/(a + b*tan(e + f*x)), x)`

### 3.54.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 983 vs.  $2(212) = 424$ .

Time = 0.57 (sec) , antiderivative size = 983, normalized size of antiderivative = 4.05

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output

```
-1/12*(18*c^2*d*e*(2*(f*x + e)*a/((a^2 + b^2)*f) + 2*b*log(b*tan(f*x + e)
+ a)/((a^2 + b^2)*f) - b*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*f)) - 6*(2*(
f*x + e)*a/(a^2 + b^2) + 2*b*log(b*tan(f*x + e) + a)/(a^2 + b^2) - b*log(t
an(f*x + e)^2 + 1)/(a^2 + b^2))*c^3 - (3*(f*x + e)^4*(a - I*b)*d^3 + 12*I*
b*d^3*polylog(4, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 12*((a - I*b)
*d^3*e - (a - I*b)*c*d^2*f)*(f*x + e)^3 + 18*((a - I*b)*d^3*e^2 - 2*(a - I
*b)*c*d^2*e*f + (a - I*b)*c^2*d*f^2)*(f*x + e)^2 - 12*((a - I*b)*d^3*e^3 -
3*(a - I*b)*c*d^2*e^2*f)*(f*x + e) - 12*(I*b*d^3*e^3 - 3*I*b*c*d^2*e^2*f)
*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e)
+ b*sin(2*f*x + 2*e) + a) - 4*(4*I*(f*x + e)^3*b*d^3 + 9*(-I*b*d^3*e + I*b
*c*d^2*f)*(f*x + e)^2 + 9*(I*b*d^3*e^2 - 2*I*b*c*d^2*e*f + I*b*c^2*d*f^2)*
(f*x + e)*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))
/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x
+ 2*e))/(a^2 + b^2)) - 6*(4*I*(f*x + e)^2*b*d^3 + 3*I*b*d^3*e^2 - 6*I*b*c*
d^2*e*f + 3*I*b*c^2*d*f^2 + 6*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e))*dilog(
(I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 6*(b*d^3*e^3 - 3*b*c*d^2*e^2*f
)*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2
)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 2*(4*
(f*x + e)^3*b*d^3 - 9*(b*d^3*e - b*c*d^2*f)*(f*x + e)^2 + 9*(b*d^3*e^2 - 2
*b*c*d^2*e*f + b*c^2*d*f^2)*(f*x + e))*log(((a^2 + b^2)*cos(2*f*x + 2*e...
```

**3.54.8 Giac [F]**

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \int \frac{(dx + c)^3}{b \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^3/(b*tan(f*x + e) + a), x)`

**3.54.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx$$

input `int((c + d*x)^3/(a + b*tan(e + f*x)),x)`

output `int((c + d*x)^3/(a + b*tan(e + f*x)), x)`

### 3.55 $\int \frac{(c+dx)^2}{a+b \tan(e+fx)} dx$

3.55.1	Optimal result	375
3.55.2	Mathematica [A] (verified)	376
3.55.3	Rubi [A] (verified)	376
3.55.4	Maple [B] (verified)	379
3.55.5	Fricas [B] (verification not implemented)	380
3.55.6	Sympy [F]	381
3.55.7	Maxima [B] (verification not implemented)	381
3.55.8	Giac [F]	382
3.55.9	Mupad [F(-1)]	382

#### 3.55.1 Optimal result

Integrand size = 20, antiderivative size = 181

$$\int \frac{(c+dx)^2}{a+b \tan(e+fx)} dx = \frac{(c+dx)^3}{3(a+ib)d} + \frac{b(c+dx)^2 \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} - \frac{ibd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^3}$$

output `1/3*(d*x+c)^3/(a+I*b)/d+b*(d*x+c)^2*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-I*b*d*(d*x+c)*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2+1/2*b*d^2*polylog(3,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^3`



### 3.55.2 Mathematica [A] (verified)

Time = 1.43 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.30

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

$$= \frac{1}{6}b \left( -\frac{4(c + dx)^3}{(ia + b)d(-ib(-1 + e^{2ie}) + a(1 + e^{2ie}))} + \frac{6(c + dx)^2 \log\left(1 + \frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2 + b^2)f} \right.$$

$$\left. + \frac{3d\left(2if(c + dx) \text{PolyLog}\left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right) + d \text{PolyLog}\left(3, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)\right)}{(a^2 + b^2)f^3} \right)$$

$$+ \frac{x(3c^2 + 3cdx + d^2x^2) \cos(e)}{3(a \cos(e) + b \sin(e))}$$

input `Integrate[(c + d*x)^2/(a + b*Tan[e + f*x]),x]`

output `(b*((-4*(c + d*x)^3)/((I*a + b)*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))) + (6*(c + d*x)^2*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x))]))/((a^2 + b^2)*f) + (3*d*((2*I)*f*(c + d*x)*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] + d*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a^2 + b^2)*f^3))/6 + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cos[e])/(3*(a*Cos[e] + b*Sin[e]))`

### 3.55.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {3042, 4215, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

$$\downarrow \text{4215}$$

$$\begin{aligned}
& 2ib \int \frac{e^{2i(e+fx)}(c+dx)^2}{(a+ib)^2 + (a^2+b^2)e^{2i(e+fx)}} dx + \frac{(c+dx)^3}{3d(a+ib)} \\
& \quad \downarrow \text{2620} \\
& 2ib \left( \frac{id \int (c+dx) \log \left( \frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1 \right) dx}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left( 1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \frac{(c+dx)^3}{3d(a+ib)} \\
& \quad \downarrow \text{3011} \\
& 2ib \left( \frac{id \left( \frac{i(c+dx) \operatorname{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \int \operatorname{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{2f} \right)}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left( 1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) \\
& \quad \quad \quad \frac{(c+dx)^3}{3d(a+ib)} \\
& \quad \quad \quad \downarrow \text{2720} \\
& 2ib \left( \frac{id \left( \frac{i(c+dx) \operatorname{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{d \int e^{-2i(e+fx)} \operatorname{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) de^{2i(e+fx)}}{4f^2} \right)}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left( 1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) \\
& \quad \quad \quad \frac{(c+dx)^3}{3d(a+ib)} \\
& \quad \quad \quad \downarrow \text{7143} \\
& 2ib \left( \frac{id \left( \frac{i(c+dx) \operatorname{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{d \operatorname{PolyLog} \left( 3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{4f^2} \right)}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left( 1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) \\
& \quad \quad \quad \frac{(c+dx)^3}{3d(a+ib)}
\end{aligned}$$

---

3.55.  $\int \frac{(c+dx)^2}{a+b \tan(e+fx)} dx$

input `Int[(c + d*x)^2/(a + b*Tan[e + f*x]),x]`

output `(c + d*x)^3/(3*(a + I*b)*d) + (2*I)*b*(((1/2*I)*(c + d*x)^2*Log[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/(a^2 + b^2)*f) + (I*d*(((I/2)*(c + d*x)*PolyLog[2, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/f - (d*PolyLog[3, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/(4*f^2)))/(a^2 + b^2)*f)`

### 3.55.3.1 Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)^v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4215 `Int[((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Simp[2*I*b Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

rule 7143 `Int [PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp [PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.55.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 939 vs.  $2(166) = 332$ .

Time = 0.69 (sec) , antiderivative size = 940, normalized size of antiderivative = 5.19

method	result
risch	$-\frac{d^2x^3}{3(ib-a)} - \frac{dcx^2}{ib-a} - \frac{c^2x}{ib-a} - \frac{c^3}{3d(ib-a)} + \frac{4be^3d^2}{3f^3(ia+b)(-ib-a)} - \frac{2ibcd \ln\left(1 - \frac{(-ib+a)e^{2i(fx+e)}}{-ib-a}\right)x}{f(ia+b)(-ib-a)} + \frac{2be^2d^2x}{f^2(ia+b)(-ib-a)} -$

input `int((d*x+c)^2/(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/3*d^2/(I*b-a)*x^3-d/(I*b-a)*c*x^2-1/(I*b-a)*c^2*x-1/3/d/(I*b-a)*c^3+4/3 \\ & /f^3/(I*a+b)*b/(-I*b-a)*e^3*d^2-2*I/f/(I*a+b)*b/(-I*b-a)*c*d*\ln(1-(a-I*b)* \\ & \exp(2*I*(f*x+e))/(-I*b-a))*x+2/f^2/(I*a+b)*b/(-I*b-a)*e^2*d^2*x-1/f^2/(I*a \\ & +b)*b/(-I*b-a)*d^2*polylog(2,(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))*x-2*I/f^2/ \\ & (I*a+b)*b/(-I*b-a)*c*d*\ln(1-(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))*e-2*I/f^3/( \\ & I*a+b)*b*d^2*e^2/(a+I*b)*\ln(\exp(I*(f*x+e)))-2/(I*a+b)*b/(-I*b-a)*c*d*x^2-4 \\ & /f/(I*a+b)*b/(-I*b-a)*c*d*e*x-2/f^2/(I*a+b)*b/(-I*b-a)*c*d*e^2-1/f^2/(I*a+ \\ & b)*b/(-I*b-a)*c*d*polylog(2,(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))-1/2*I/f^3/( \\ & I*a+b)*b/(-I*b-a)*d^2*polylog(3,(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))-2*I/f^2 \\ & / (I*a+b)*b*c*d*e/(a+I*b)*\ln(I*\exp(2*I*(f*x+e))*b-a*\exp(2*I*(f*x+e))-I*b-a) \\ & -2*I/f/(I*a+b)*b*c^2/(a+I*b)*\ln(\exp(I*(f*x+e)))+I/f/(I*a+b)*b*c^2/(a+I*b)* \\ & \ln(I*\exp(2*I*(f*x+e))*b-a*\exp(2*I*(f*x+e))-I*b-a)+4*I/f^2/(I*a+b)*b*c*d*e/ \\ & (a+I*b)*\ln(\exp(I*(f*x+e)))+I/f^3/(I*a+b)*b/(-I*b-a)*e^2*d^2*\ln(1-(a-I*b)*e \\ & \exp(2*I*(f*x+e))/(-I*b-a))-2/3/(I*a+b)*b/(-I*b-a)*d^2*x^3-I/f/(I*a+b)*b/(-I \\ & *b-a)*d^2*\ln(1-(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))*x^2+I/f^3/(I*a+b)*b*d^2* \\ & e^2/(a+I*b)*\ln(I*\exp(2*I*(f*x+e))*b-a*\exp(2*I*(f*x+e))-I*b-a) \end{aligned}$$

### 3.55.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 834 vs.  $2(160) = 320$ .

Time = 0.27 (sec) , antiderivative size = 834, normalized size of antiderivative = 4.61

$$\int \frac{(c+dx)^2}{a+b \tan(e+fx)} dx$$

$$4ad^2 f^3 x^3 + 12acd f^3 x^2 + 12ac^2 f^3 x + 3bd^2 \text{polylog}\left(3, \frac{(a^2+2iab-b^2) \tan(fx+e)^2 - a^2 - 2iab + b^2 - 2(-ia^2+2ab+ib^2) \tan(fx+e)}{(a^2+b^2) \tan(fx+e)^2 + a^2 + b^2}\right)$$


---

input `integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="fracas")`

output

```
1/12*(4*a*d^2*f^3*x^3 + 12*a*c*d*f^3*x^2 + 12*a*c^2*f^3*x + 3*b*d^2*polylog(3, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 3*b*d^2*polylog(3, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 6*(-I*b*d^2*f*x - I*b*c*d*f)*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*b*d^2*f*x + I*b*c*d*f)*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + 6*(b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)))/((a^2 + b^2)*f^3)
```

**3.55.6 Sympy [F]**

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

input `integrate((d*x+c)**2/(a+b*tan(f*x+e)),x)`

output `Integral((c + d*x)**2/(a + b*tan(e + f*x)), x)`

**3.55.7 Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 715 vs.  $2(160) = 320$ .

Time = 0.55 (sec) , antiderivative size = 715, normalized size of antiderivative = 3.95

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx =$$

$$6cde \left( \frac{2(fx+e)a}{(a^2+b^2)f} + \frac{2b \log(b \tan(fx+e)+a)}{(a^2+b^2)f} - \frac{b \log(\tan(fx+e)^2+1)}{(a^2+b^2)f} \right) - 3 \left( \frac{2(fx+e)a}{a^2+b^2} + \frac{2b \log(b \tan(fx+e)+a)}{a^2+b^2} - \frac{b \log(\tan(fx+e)^2+1)}{a^2+b^2} \right)$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output

```
-1/6*(6*c*d*e*(2*(f*x + e)*a/((a^2 + b^2)*f) + 2*b*log(b*tan(f*x + e) + a)
/((a^2 + b^2)*f) - b*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*f)) - 3*(2*(f*x
+ e)*a/(a^2 + b^2) + 2*b*log(b*tan(f*x + e) + a)/(a^2 + b^2) - b*log(tan(f
*x + e)^2 + 1)/(a^2 + b^2))*c^2 - (2*(f*x + e)^3*(a - I*b)*d^2 + 6*(f*x +
e)*(a - I*b)*d^2*e^2 + 6*I*b*d^2*e^2*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2
*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) + 3*b*d^2*e^
2*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2
)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 3*b*d
^2*polylog(3, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 6*((a - I*b)*d^2
*e - (a - I*b)*c*d*f)*(f*x + e)^2 - 6*(I*(f*x + e)^2*b*d^2 + 2*(-I*b*d^2*e
+ I*b*c*d*f)*(f*x + e))*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin
(2*f*x + 2*e))/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b
^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) - 6*(I*(f*x + e)*b*d^2 - I*b*d^2*e + I*
b*c*d*f)*dilog((I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) + 3*((f*x + e)^2*
b*d^2 - 2*(b*d^2*e - b*c*d*f)*(f*x + e))*log(((a^2 + b^2)*cos(2*f*x + 2*e)
^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 +
2*(a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)))/((a^2 + b^2)*f^2))/f
```

### 3.55.8 Giac [F]

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \int \frac{(dx + c)^2}{b \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^2/(b*tan(f*x + e) + a), x)`

### 3.55.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

input `int((c + d*x)^2/(a + b*tan(e + f*x)),x)`

output `int((c + d*x)^2/(a + b*tan(e + f*x)), x)`

---

3.55.  $\int \frac{(c+dx)^2}{a+b \tan(e+fx)} dx$

### 3.56 $\int \frac{c+dx}{a+b \tan(e+fx)} dx$

3.56.1	Optimal result . . . . .	383
3.56.2	Mathematica [A] (verified) . . . . .	383
3.56.3	Rubi [A] (verified) . . . . .	384
3.56.4	Maple [B] (verified) . . . . .	386
3.56.5	Fricas [B] (verification not implemented) . . . . .	386
3.56.6	Sympy [F] . . . . .	387
3.56.7	Maxima [B] (verification not implemented) . . . . .	387
3.56.8	Giac [F] . . . . .	388
3.56.9	Mupad [F(-1)] . . . . .	388

#### 3.56.1 Optimal result

Integrand size = 18, antiderivative size = 125

$$\int \frac{c+dx}{a+b \tan(e+fx)} dx = \frac{(c+dx)^2}{2(a+ib)d} + \frac{b(c+dx) \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} - \frac{ibd \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^2}$$

output  $\frac{1}{2}*(d*x+c)^2/(a+I*b)/d+b*(d*x+c)*\ln(1+(a^2+b^2)*\exp(2*I*(f*x+e)))/(a+I*b)^2)/(a^2+b^2)/f-1/2*I*b*d*\operatorname{polylog}(2,-(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2$

#### 3.56.2 Mathematica [A] (verified)

Time = 1.56 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.42

$$\int \frac{c+dx}{a+b \tan(e+fx)} dx = \frac{b(c+dx)^2}{(a-ib)d(b-be^{2ie}-ia(1+e^{2ie}))} + \frac{b(c+dx) \log\left(1 + \frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2+b^2)f} + \frac{ibd \operatorname{PolyLog}\left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)}{2(a^2+b^2)f^2} + \frac{x(2c+dx) \cos(e)}{2(a \cos(e) + b \sin(e))}$$



input `Integrate[(c + d*x)/(a + b*Tan[e + f*x]),x]`

output `(b*(c + d*x)^2)/((a - I*b)*d*(b - b*E^((2*I)*e) - I*a*(1 + E^((2*I)*e)))) + (b*(c + d*x)*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/((a^2 + b^2)*f) + ((I/2)*b*d*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/((a^2 + b^2)*f^2) + (x*(2*c + d*x)*Cos[e])/(2*(a*Cos[e] + b*Sin[e]))`

### 3.56.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$ , Rules used = {3042, 4215, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{c + dx}{a + b \tan(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{c + dx}{a + b \tan(e + fx)} dx \\
 & \quad \downarrow \text{4215} \\
 & 2ib \int \frac{e^{2i(e+fx)}(c + dx)}{(a + ib)^2 + (a^2 + b^2) e^{2i(e+fx)}} dx + \frac{(c + dx)^2}{2d(a + ib)} \\
 & \quad \downarrow \text{2620} \\
 & 2ib \left( \frac{id \int \log \left( \frac{e^{2i(e+fx)}(a^2 + b^2)}{(a + ib)^2} + 1 \right) dx}{2f(a^2 + b^2)} - \frac{i(c + dx) \log \left( 1 + \frac{(a^2 + b^2) e^{2i(e+fx)}}{(a + ib)^2} \right)}{2f(a^2 + b^2)} \right) + \frac{(c + dx)^2}{2d(a + ib)} \\
 & \quad \downarrow \text{2715} \\
 & 2ib \left( \frac{d \int e^{-2i(e+fx)} \log \left( \frac{e^{2i(e+fx)}(a^2 + b^2)}{(a + ib)^2} + 1 \right) de^{2i(e+fx)}}{4f^2(a^2 + b^2)} - \frac{i(c + dx) \log \left( 1 + \frac{(a^2 + b^2) e^{2i(e+fx)}}{(a + ib)^2} \right)}{2f(a^2 + b^2)} \right) + \\
 & \quad \frac{(c + dx)^2}{2d(a + ib)} \\
 & \quad \downarrow \text{2838}
 \end{aligned}$$

$$2ib \left( -\frac{i(c+dx) \log \left( 1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} - \frac{d \operatorname{PolyLog} \left( 2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{4f^2(a^2+b^2)} \right) + \frac{(c+dx)^2}{2d(a+ib)}$$

input `Int[(c + d*x)/(a + b*Tan[e + f*x]),x]`

output `(c + d*x)^2/(2*(a + I*b)*d) + (2*I)*b*((( -1/2*I)*(c + d*x)*Log[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) - (d*PolyLog[2, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/(4*(a^2 + b^2)*f^2))`

### 3.56.3.1 Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4215 `Int[((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Simp[2*I*b Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

### 3.56.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 461 vs.  $2(113) = 226$ .

Time = 0.64 (sec) , antiderivative size = 462, normalized size of antiderivative = 3.70

method	result
risch	$-\frac{dx^2}{2(ib-a)} - \frac{cx}{ib-a} + \frac{bc \ln(ie^{2i(fx+e)}b - ae^{2i(fx+e)} - ib - a)}{f(-ib+a)(ib+a)} - \frac{2bc \ln(e^{i(fx+e)})}{f(-ib+a)(ib+a)} - \frac{bd \ln\left(1 - \frac{(-ib+a)e^{2i(fx+e)}}{-ib-a}\right)x}{f(-ib+a)(-ib-a)} + \frac{ibd}{(-ib+a)}$

input `int((d*x+c)/(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/2/(I*b-a)*d*x^2-1/(I*b-a)*c*x+1/f*b/(a-I*b)*c/(a+I*b)*\ln(I*\exp(2*I*(f*x \\ & +e))*b-a*\exp(2*I*(f*x+e))-I*b-a-2/f*b/(a-I*b)*c/(a+I*b)*\ln(\exp(I*(f*x+e)) \\ & )-1/f*b/(a-I*b)/(-I*b-a)*d*\ln(1-(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))*x+I*b/( \\ & a-I*b)/(-I*b-a)*d*x^2-1/f^2*b/(a-I*b)/(-I*b-a)*d*\ln(1-(a-I*b)*\exp(2*I*(f*x \\ & +e))/(-I*b-a))*e+2*I/f*b/(a-I*b)/(-I*b-a)*d*e*x+I/f^2*b/(a-I*b)/(-I*b-a)*d \\ & *e^2+1/2*I/f^2*b/(a-I*b)/(-I*b-a)*d*polylog(2,(a-I*b)*\exp(2*I*(f*x+e))/(-I \\ & *b-a))-1/f^2*b/(a-I*b)*d*e/(a+I*b)*\ln(I*\exp(2*I*(f*x+e))*b-a*\exp(2*I*(f*x+ \\ & e))-I*b-a)+2/f^2*b/(a-I*b)*d*e/(a+I*b)*\ln(\exp(I*(f*x+e))) \end{aligned}$$

### 3.56.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 527 vs.  $2(108) = 216$ .

Time = 0.27 (sec) , antiderivative size = 527, normalized size of antiderivative = 4.22

$$\int \frac{c+dx}{a+b \tan(e+fx)} dx$$

$$= \frac{2adf^2x^2 + 4acf^2x + i b d \operatorname{Li}_2\left(\frac{2\left((iab-b^2)\tan(fx+e)^2 - a^2 - iab + (ia^2 - 2ab - ib^2)\tan(fx+e)\right)}{(a^2+b^2)\tan(fx+e)^2 + a^2 + b^2} + 1\right) - i b d \operatorname{Li}_2\left(\frac{2\left((-iab-b^2)\tan(fx+e)^2 - a^2 - iab + (ia^2 - 2ab - ib^2)\tan(fx+e)\right)}{(a^2+b^2)\tan(fx+e)^2 + a^2 + b^2} + 1\right)}{2adf^2x^2 + 4acf^2x + i b d \operatorname{Li}_2\left(\frac{2\left((iab-b^2)\tan(fx+e)^2 - a^2 - iab + (ia^2 - 2ab - ib^2)\tan(fx+e)\right)}{(a^2+b^2)\tan(fx+e)^2 + a^2 + b^2} + 1\right) - i b d \operatorname{Li}_2\left(\frac{2\left((-iab-b^2)\tan(fx+e)^2 - a^2 - iab + (ia^2 - 2ab - ib^2)\tan(fx+e)\right)}{(a^2+b^2)\tan(fx+e)^2 + a^2 + b^2} + 1\right)}$$

input `integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="fracas")`

```
output 1/4*(2*a*d*f^2*x^2 + 4*a*c*f^2*x + I*b*d*dilog(2*((I*a*b - b^2)*tan(f*x +
e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*ta
n(f*x + e)^2 + a^2 + b^2) + 1) - I*b*d*dilog(2*((-I*a*b - b^2)*tan(f*x + e
)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*ta
n(f*x + e)^2 + a^2 + b^2) + 1) + 2*(b*d*f*x + b*d*e)*log(-2*((I*a*b - b^2)
*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^
2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(b*d*f*x + b*d*e)*log(-2*((-I*a*
b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x +
e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 2*(b*d*e - b*c*f)*log(((I
*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/((
tan(f*x + e)^2 + 1)) - 2*(b*d*e - b*c*f)*log(((I*a*b - b^2)*tan(f*x + e)^2
+ a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)))/((a^
2 + b^2)*f^2)
```

### 3.56.6 Sympy [F]

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{c + dx}{a + b \tan(e + fx)} dx$$

```
input integrate((d*x+c)/(a+b*tan(f*x+e)),x)
```

```
output Integral((c + d*x)/(a + b*tan(e + f*x)), x)
```

### 3.56.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 399 vs.  $2(108) = 216$ .

Time = 0.60 (sec) , antiderivative size = 399, normalized size of antiderivative = 3.19

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx$$

$$= \frac{(a - ib)df^2x^2 + 2(a - ib)cf^2x - 2ibdfx \arctan\left(\frac{2ab \cos(2fx+2e) - (a^2 - b^2) \sin(2fx+2e)}{a^2 + b^2}\right), \frac{2ab \sin(2fx+2e) + a^2 + b^2}{a^2 + b^2}}$$

```
input integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="maxima")
```

---

3.56.  $\int \frac{c+dx}{a+b \tan(e+fx)} dx$

output `1/2*((a - I*b)*d*f^2*x^2 + 2*(a - I*b)*c*f^2*x - 2*I*b*d*f*x*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) + b*d*f*x*log(((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) + 2*I*b*c*f*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) + b*c*f*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) - I*b*d*dilog((I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)))/(a^2 + b^2)*f^2)`

### 3.56.8 Giac [F]

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{dx + c}{b \tan(fx + e) + a} dx$$

input `integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)/(b*tan(f*x + e) + a), x)`

### 3.56.9 Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{c + dx}{a + b \tan(e + fx)} dx$$

input `int((c + d*x)/(a + b*tan(e + f*x)),x)`

output `int((c + d*x)/(a + b*tan(e + f*x)), x)`

$$3.57 \quad \int \frac{1}{(c+dx)(a+b \tan(e+fx))} dx$$

3.57.1	Optimal result	389
3.57.2	Mathematica [N/A]	389
3.57.3	Rubi [N/A]	390
3.57.4	Maple [N/A] (verified)	391
3.57.5	Fricas [N/A]	391
3.57.6	Sympy [N/A]	391
3.57.7	Maxima [N/A]	392
3.57.8	Giac [N/A]	392
3.57.9	Mupad [N/A]	393

### 3.57.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b \tan(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b \tan(e+fx))}, x\right)$$

output `Unintegrable(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

### 3.57.2 Mathematica [N/A]

Not integrable

Time = 2.41 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b \tan(e+fx))} dx = \int \frac{1}{(c+dx)(a+b \tan(e+fx))} dx$$

input `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])),x]`

output `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])), x]`

### 3.57.3 Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx$$

↓ 3042

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx$$

↓ 4223

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx$$

input `Int[1/((c + d*x)*(a + b*Tan[e + f*x])),x]`

output `$Aborted`

#### 3.57.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.57.4 Maple [N/A] (verified)**

Not integrable

Time = 0.42 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \tan(fx + e))} dx$$

input `int(1/(d*x+c)/(a+b*tan(f*x+e)),x)`output `int(1/(d*x+c)/(a+b*tan(f*x+e)),x)`**3.57.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.35

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="fricas")`output `integral(1/(a*d*x + a*c + (b*d*x + b*c)*tan(f*x + e)), x)`**3.57.6 Sympy [N/A]**

Not integrable

Time = 1.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx))(c + dx)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x)`output `Integral(1/((a + b*tan(e + f*x))*(c + d*x)), x)`



**3.57.7 Maxima [N/A]**

Not integrable

Time = 0.89 (sec) , antiderivative size = 279, normalized size of antiderivative = 13.95

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx = \int \frac{1}{(dx+c)(b\tan(fx+e)+a)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output `(2*(a^2*b + b^3)*d*integrate((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/((a^4 + 2*a^2*b^2 + b^4)*d*x + ((a^4 + 2*a^2*b^2 + b^4)*d*x + (a^4 + 2*a^2*b^2 + b^4)*c)*cos(2*f*x + 2*e)^2 + ((a^4 + 2*a^2*b^2 + b^4)*d*x + (a^4 + 2*a^2*b^2 + b^4)*c)*sin(2*f*x + 2*e)^2 + (a^4 + 2*a^2*b^2 + b^4)*c + 2*((a^4 - b^4)*d*x + (a^4 - b^4)*c)*cos(2*f*x + 2*e) + 4*((a^3*b + a*b^3)*d*x + (a^3*b + a*b^3)*c)*sin(2*f*x + 2*e)), x) + a*log(d*x + c))/((a^2 + b^2)*d)`

**3.57.8 Giac [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx = \int \frac{1}{(dx+c)(b\tan(fx+e)+a)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate(1/((d*x + c)*(b*tan(f*x + e) + a)), x)`

**3.57.9 Mupad [N/A]**

Not integrable

Time = 3.96 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx = \int \frac{1}{(a+b\tan(e+fx))(c+dx)} dx$$

input `int(1/((a + b*tan(e + f*x))*(c + d*x)),x)`output `int(1/((a + b*tan(e + f*x))*(c + d*x)), x)`

**3.58**  $\int \frac{1}{(c+dx)^2(a+b \tan(e+fx))} dx$

3.58.1 Optimal result . . . . . 394  
 3.58.2 Mathematica [N/A] . . . . . 394  
 3.58.3 Rubi [N/A] . . . . . 395  
 3.58.4 Maple [N/A] (verified) . . . . . 396  
 3.58.5 Fricas [N/A] . . . . . 396  
 3.58.6 Sympy [N/A] . . . . . 396  
 3.58.7 Maxima [N/A] . . . . . 397  
 3.58.8 Giac [N/A] . . . . . 397  
 3.58.9 Mupad [N/A] . . . . . 398

**3.58.1 Optimal result**

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \text{Int}\left(\frac{1}{(c + dx)^2(a + b \tan(e + fx))}, x\right)$$

output `Unintegrable(1/(d*x+c)^2/(a+b*tan(f*x+e)),x)`

**3.58.2 Mathematica [N/A]**

Not integrable

Time = 4.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx$$

input `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])),x]`

output `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])), x]`

### 3.58.3 Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$$

↓ 3042

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$$

↓ 4223

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$$

input `Int[1/((c + d*x)^2*(a + b*Tan[e + f*x])),x]`

output `$Aborted`

#### 3.58.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.58.4 Maple [N/A] (verified)**

Not integrable

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)^2 (a + b \tan(fx + e))} dx$$

input `int(1/(d*x+c)^2/(a+b*tan(f*x+e)),x)`output `int(1/(d*x+c)^2/(a+b*tan(f*x+e)),x)`**3.58.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.55

$$\int \frac{1}{(c + dx)^2 (a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)^2 (b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="fricas")`output `integral(1/(a*d^2*x^2 + 2*a*c*d*x + a*c^2 + (b*d^2*x^2 + 2*b*c*d*x + b*c^2)*tan(f*x + e)), x)`**3.58.6 Sympy [N/A]**

Not integrable

Time = 1.76 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)^2 (a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx)) (c + dx)^2} dx$$

input `integrate(1/(d*x+c)**2/(a+b*tan(f*x+e)),x)`output `Integral(1/((a + b*tan(e + f*x))*(c + d*x)**2), x)`

**3.58.7 Maxima [N/A]**

Not integrable

Time = 1.73 (sec) , antiderivative size = 424, normalized size of antiderivative = 21.20

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \int \frac{1}{(dx+c)^2(b\tan(fx+e)+a)} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output `(2*((a^2*b + b^3)*d^2*x + (a^2*b + b^3)*c*d)*integrate((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2 + ((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2)*cos(2*f*x + 2*e)^2 + ((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2)*sin(2*f*x + 2*e)^2 + 2*((a^4 - b^4)*d^2*x^2 + 2*(a^4 - b^4)*c*d*x + (a^4 - b^4)*c^2)*cos(2*f*x + 2*e) + 4*((a^3*b + a*b^3)*d^2*x^2 + 2*(a^3*b + a*b^3)*c*d*x + (a^3*b + a*b^3)*c^2)*sin(2*f*x + 2*e)), x) - a)/((a^2 + b^2)*d^2*x + (a^2 + b^2)*c*d)`

**3.58.8 Giac [N/A]**

Not integrable

Time = 4.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \int \frac{1}{(dx+c)^2(b\tan(fx+e)+a)} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate(1/((d*x + c)^2*(b*tan(f*x + e) + a)), x)`

**3.58.9 Mupad [N/A]**

Not integrable

Time = 4.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \int \frac{1}{(a+b\tan(e+fx))(c+dx)^2} dx$$

input `int(1/((a + b*tan(e + f*x))*(c + d*x)^2),x)`output `int(1/((a + b*tan(e + f*x))*(c + d*x)^2), x)`

$$3.59 \quad \int \frac{(c+dx)^3}{(a+b \tan(e+fx))^2} dx$$

3.59.1	Optimal result	400
3.59.2	Mathematica [B] (warning: unable to verify)	401
3.59.3	Rubi [A] (verified)	402
3.59.4	Maple [B] (verified)	404
3.59.5	Fricas [B] (verification not implemented)	404
3.59.6	Sympy [F]	405
3.59.7	Maxima [B] (verification not implemented)	406
3.59.8	Giac [F]	406
3.59.9	Mupad [F(-1)]	407



### 3.59.1 Optimal result

Integrand size = 20, antiderivative size = 848

$$\begin{aligned}
 \int \frac{(c+dx)^3}{(a+b \tan(e+fx))^2} dx = & -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a+ib)(ia+b)^2 (ia-b+(ia+b)e^{2ie+2ifx}) f} \\
 & + \frac{(c+dx)^4}{4(a-ib)^2 d} + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} \\
 & + \frac{3b^2 d(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
 & + \frac{2b(c+dx)^3 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} \\
 & - \frac{2ib^2(c+dx)^3 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} \\
 & - \frac{3ib^2 d^2(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\
 & + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
 & - \frac{3b^2 d(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
 & + \frac{3b^2 d^3 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2 f^4} \\
 & + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} \\
 & - \frac{3ib^2 d^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\
 & - \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(ia-b)(a-ib)^2 f^4} \\
 & + \frac{3b^2 d^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2 f^4}
 \end{aligned}$$

output

```

-3*I*b^2*d^2*(d*x+c)*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b
^2)^2/f^3+2*b^2*(d*x+c)^3/(a+I*b)/(I*a+b)^2/(I*a-b+(I*a+b)*exp(2*I*e+2*I*f
*x))/f+1/4*(d*x+c)^4/(a-I*b)^2/d+b*(d*x+c)^4/(I*a-b)/(a-I*b)^2/d-b^2*(d*x+
c)^4/(a^2+b^2)^2/d+3*b^2*d*(d*x+c)^2*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*
b))/(a^2+b^2)^2/f^2+2*b*(d*x+c)^3*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))
/(a-I*b)^2/(a+I*b)/f-3*I*b^2*d^2*(d*x+c)*polylog(2,-(a-I*b)*exp(2*I*e+2*I*
f*x)/(a+I*b))/(a^2+b^2)^2/f^3-2*I*b^2*(d*x+c)^3/(a^2+b^2)^2/f+3*b*d*(d*x+c
)^2*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(I*a-b)/(a-I*b)^2/f^2-3
*b^2*d*(d*x+c)^2*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^
2/f^2+3/2*b^2*d^3*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)
^2/f^4+3*b*d^2*(d*x+c)*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I
*b)^2/(a+I*b)/f^3-2*I*b^2*(d*x+c)^3*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b
))/(a^2+b^2)^2/f-3/2*b*d^3*polylog(4,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/
(I*a-b)/(a-I*b)^2/f^4+3/2*b^2*d^3*polylog(4,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a
+I*b))/(a^2+b^2)^2/f^4

```

### 3.59.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1724 vs.  $2(848) = 1696$ .

Time = 9.64 (sec) , antiderivative size = 1724, normalized size of antiderivative = 2.03

$$\int \frac{(c+dx)^3}{(a+b \tan(e+fx))^2} dx = \text{Too large to display}$$

input `Integrate[(c + d*x)^3/(a + b*Tan[e + f*x])^2,x]`

```
output (b*((-4*c^2*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(3*b*d + 2*a
*c*f)*x)/(a^2 + b^2) + (4*b*(c + d*x)^3)/(a - I*b) + (2*a*f*(c + d*x)^4)/(
(a - I*b)*d) + (12*c*d*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(
b*d + a*c*f)*x*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/((a + I
*b)*(I*a + b)*f) + (6*d^2*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e))
)*(b*d + 2*a*c*f)*x^2*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/
((a + I*b)*(I*a + b)*f) + (4*a*d^3*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^
(2*I)*e))*x^3*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/((a + I
*b)*(I*a + b)) + (2*c^2*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*
(3*b*d + 2*a*c*f)*Log[a + I*b + (a - I*b)*E^((2*I)*(e + f*x))]/((a + I*b)
*(I*a + b)*f) + (6*c*d*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(
b*d + a*c*f)*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/((a^2
+ b^2)*f^2) + (3*d^2*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b
*d + 2*a*c*f)*(2*f*x*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x))
] - I*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2
)*f^3) + (3*a*d^3*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(2*f^2
*x^2*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - (2*I)*f*x*Po
lyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - PolyLog[4, (-a - I*
b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f^3))/(2*(a - I*b)*(a
+ I*b)*(b - b*E^((2*I)*e) - I*a*(1 + E^((2*I)*e))*f) + (3*x^2*(a*c^2*d...
```

### 3.59.3 Rubi [A] (verified)

Time = 2.25 (sec) , antiderivative size = 848, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {3042, 4217, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx$$

↓ 4217

$$\int \left( -\frac{4b^2(c + dx)^3}{(b + ia)^2 (ia (1 - \frac{ib}{a}) e^{2ie+2ifx} + ia (1 + \frac{ib}{a}))^2} + \frac{4b(c + dx)^3}{(a - ib)^2 (ia (1 - \frac{ib}{a}) e^{2ie+2ifx} + ia (1 + \frac{ib}{a}))} + \frac{(c + dx)^3}{(a - ib)^2} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{b(c+dx)^4}{(ia-b)(a-ib)^2d} + \frac{(c+dx)^4}{4(a-ib)^2d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2d} + \frac{2b \log\left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1\right)(c+dx)^3}{(a-ib)^2(a+ib)f} - \\
& \frac{2ib^2 \log\left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1\right)(c+dx)^3}{(a^2+b^2)^2f} + \frac{2b^2(c+dx)^3}{(a+ib)(ia+b)^2(ia+(ia+b)e^{2ie+2ifx}-b)f} - \\
& \frac{2ib^2(c+dx)^3}{(a^2+b^2)^2f} + \frac{3b^2d \log\left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1\right)(c+dx)^2}{(a^2+b^2)^2f^2} + \\
& \frac{3bd \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)(c+dx)^2}{(ia-b)(a-ib)^2f^2} - \frac{3b^2d \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)(c+dx)^2}{(a^2+b^2)^2f^2} - \\
& \frac{3ib^2d^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)(c+dx)}{(a^2+b^2)^2f^3} + \frac{3bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)(c+dx)}{(a-ib)^2(a+ib)f^3} - \\
& \frac{3ib^2d^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)(c+dx)}{(a^2+b^2)^2f^3} + \frac{3b^2d^3 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2f^4} - \\
& \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(ia-b)(a-ib)^2f^4} + \frac{3b^2d^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2f^4}
\end{aligned}$$

input `Int[(c + d*x)^3/(a + b*Tan[e + f*x])^2,x]`

output

```

((-2*I)*b^2*(c + d*x)^3)/((a^2 + b^2)^2*f) + (2*b^2*(c + d*x)^3)/((a + I*b)
)*(I*a + b)^2*(I*a - b + (I*a + b)*E^((2*I)*e + (2*I)*f*x))*f) + (c + d*x)
^4/(4*(a - I*b)^2*d) + (b*(c + d*x)^4)/((I*a - b)*(a - I*b)^2*d) - (b^2*(c
+ d*x)^4)/((a^2 + b^2)^2*d) + (3*b^2*d*(c + d*x)^2*Log[1 + ((a - I*b)*E^
(2*I)*e + (2*I)*f*x))/(a + I*b)]/((a^2 + b^2)^2*f^2) + (2*b*(c + d*x)^3*L
og[1 + ((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((a - I*b)^2*(a + I
*b)*f) - ((2*I)*b^2*(c + d*x)^3*Log[1 + ((a - I*b)*E^((2*I)*e + (2*I)*f*x)
)/(a + I*b)]/((a^2 + b^2)^2*f) - ((3*I)*b^2*d^2*(c + d*x)*PolyLog[2, -(((
a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((a^2 + b^2)^2*f^3) + (3*b*
d*(c + d*x)^2*PolyLog[2, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]
)/((I*a - b)*(a - I*b)^2*f^2) - (3*b^2*d*(c + d*x)^2*PolyLog[2, -(((a - I*
b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((a^2 + b^2)^2*f^2) + (3*b^2*d^3*
PolyLog[3, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((2*(a^2 + b^
2)^2*f^4) + (3*b*d^2*(c + d*x)*PolyLog[3, -(((a - I*b)*E^((2*I)*e + (2*I)*
f*x))/(a + I*b)]/((a - I*b)^2*(a + I*b)*f^3) - ((3*I)*b^2*d^2*(c + d*x)*
PolyLog[3, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((a^2 + b^2)
^2*f^3) - (3*b*d^3*PolyLog[4, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I
*b)]/((2*(I*a - b)*(a - I*b)^2*f^4) + (3*b^2*d^3*PolyLog[4, -(((a - I*b)*
E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((2*(a^2 + b^2)^2*f^4)

```

### 3.59.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4217 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 + b^2 + (a - I*b)^2*E^(2*I*(e + f*x)))))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IGtQ[m, 0]`

### 3.59.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 5958 vs.  $2(763) = 1526$ .

Time = 0.97 (sec) , antiderivative size = 5959, normalized size of antiderivative = 7.03

method	result	size
risch	Expression too large to display	5959

input `int((d*x+c)^3/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `result too large to display`

### 3.59.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2537 vs.  $2(695) = 1390$ .

Time = 0.33 (sec) , antiderivative size = 2537, normalized size of antiderivative = 2.99

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="fracas")`

```

output 1/4*((a^3 - a*b^2)*d^3*f^4*x^4 - 4*b^3*c^3*f^3 - 4*(b^3*d^3*f^3 - (a^3 - a
*b^2)*c*d^2*f^4)*x^3 - 6*(2*b^3*c*d^2*f^3 - (a^3 - a*b^2)*c^2*d*f^4)*x^2 -
4*(3*b^3*c^2*d*f^3 - (a^3 - a*b^2)*c^3*f^4)*x - 6*(-I*a^2*b*d^3*f^2*x^2 -
I*a^2*b*c^2*d*f^2 - I*a*b^2*c*d^2*f - I*(2*a^2*b*c*d^2*f^2 + a*b^2*d^3*f)
*x + (-I*a*b^2*d^3*f^2*x^2 - I*a*b^2*c^2*d*f^2 - I*b^3*c*d^2*f - I*(2*a*b^
2*c*d^2*f^2 + b^3*d^3*f)*x)*tan(f*x + e))*dilog(2*((I*a*b - b^2)*tan(f*x +
e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*t
an(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*a^2*b*d^3*f^2*x^2 + I*a^2*b*c^2*d*f
^2 + I*a*b^2*c*d^2*f + I*(2*a^2*b*c*d^2*f^2 + a*b^2*d^3*f)*x + (I*a*b^2*d^
3*f^2*x^2 + I*a*b^2*c^2*d*f^2 + I*b^3*c*d^2*f + I*(2*a*b^2*c*d^2*f^2 + b^3
*d^3*f)*x)*tan(f*x + e))*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*
a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 +
a^2 + b^2) + 1) + 2*(2*a^2*b*d^3*f^3*x^3 + 2*a^2*b*d^3*e^3 + 6*a^2*b*c^2*
d*e*f^2 - 3*a*b^2*d^3*e^2 + 3*(2*a^2*b*c*d^2*f^3 + a*b^2*d^3*f^2)*x^2 - 6*
(a^2*b*c*d^2*e^2 - a*b^2*c*d^2*e)*f + 6*(a^2*b*c^2*d*f^3 + a*b^2*c*d^2*f^2
)*x + (2*a*b^2*d^3*f^3*x^3 + 2*a*b^2*d^3*e^3 + 6*a*b^2*c^2*d*e*f^2 - 3*b^3
*d^3*e^2 + 3*(2*a*b^2*c*d^2*f^3 + b^3*d^3*f^2)*x^2 - 6*(a*b^2*c*d^2*e^2 -
b^3*c*d^2*e)*f + 6*(a*b^2*c^2*d*f^3 + b^3*c*d^2*f^2)*x)*tan(f*x + e))*log(
-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*t
an(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(2*a^2*b*d^3...

```

### 3.59.6 Sympy [F]

$$\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx = \int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx$$

```
input integrate((d*x+c)**3/(a+b*tan(f*x+e))**2,x)
```

```
output Integral((c + d*x)**3/(a + b*tan(e + f*x))**2, x)
```

### 3.59.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 4631 vs.  $2(695) = 1390$ .

Time = 2.41 (sec) , antiderivative size = 4631, normalized size of antiderivative = 5.46

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

```
input integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output -1/12*(36*c^2*d*e*(2*a*b*log(b*tan(f*x + e) + a)/((a^4 + 2*a^2*b^2 + b^4)*
f) - a*b*log(tan(f*x + e)^2 + 1)/((a^4 + 2*a^2*b^2 + b^4)*f) - b/((a^2*b +
b^3)*f*tan(f*x + e) + (a^3 + a*b^2)*f) + (a^2 - b^2)*(f*x + e)/((a^4 + 2*
a^2*b^2 + b^4)*f)) - 12*(2*a*b*log(b*tan(f*x + e) + a)/(a^4 + 2*a^2*b^2 +
b^4) - a*b*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + (a^2 - b^2)*(
f*x + e)/(a^4 + 2*a^2*b^2 + b^4) - b/(a^3 + a*b^2 + (a^2*b + b^3)*tan(f*x
+ e)))*c^3 - (3*(a^3 - I*a^2*b + a*b^2 - I*b^3)*(f*x + e)^4*d^3 - 24*(I*a*
b^2 - b^3)*d^3*e^3 - 72*(-I*a*b^2 + b^3)*c*d^2*e^2*f - 12*((a^3 - I*a^2*b
+ a*b^2 - I*b^3)*d^3*e - (a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d^2*f)*(f*x +
e)^3 + 18*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e^2 - 2*(a^3 - I*a^2*b + a*b
^2 - I*b^3)*c*d^2*e*f + (a^3 - I*a^2*b + a*b^2 - I*b^3)*c^2*d*f^2)*(f*x +
e)^2 - 12*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e^3 - 3*(a^3 - I*a^2*b + a*
b^2 - I*b^3)*c*d^2*e^2*f)*(f*x + e) - 12*(2*(I*a^2*b - a*b^2)*d^3*e^3 + 3*
(-I*a*b^2 + b^3)*d^3*e^2 + 3*(-I*a*b^2 + b^3)*c^2*d*f^2 + 6*((-I*a^2*b + a
*b^2)*c*d^2*e^2 + (I*a*b^2 - b^3)*c*d^2*e)*f + (2*(I*a^2*b + a*b^2)*d^3*e^
3 + 3*(-I*a*b^2 - b^3)*d^3*e^2 + 3*(-I*a*b^2 - b^3)*c^2*d*f^2 + 6*((-I*a^2
*b - a*b^2)*c*d^2*e^2 + (I*a*b^2 + b^3)*c*d^2*e)*f)*cos(2*f*x + 2*e) - (2*
(a^2*b - I*a*b^2)*d^3*e^3 - 3*(a*b^2 - I*b^3)*d^3*e^2 - 3*(a*b^2 - I*b^3)*
c^2*d*f^2 - 6*((a^2*b - I*a*b^2)*c*d^2*e^2 - (a*b^2 - I*b^3)*c*d^2*e)*f)*s
in(2*f*x + 2*e))*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, ...
```

### 3.59.8 Giac [F]

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \int \frac{(dx + c)^3}{(b \tan(fx + e) + a)^2} dx$$

```
input integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="giac")
```

---

3.59.  $\int \frac{(c+dx)^3}{(a+b \tan(e+fx))^2} dx$

output `integrate((d*x + c)^3/(b*tan(f*x + e) + a)^2, x)`

### 3.59.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx$$

input `int((c + d*x)^3/(a + b*tan(e + f*x))^2,x)`

output `int((c + d*x)^3/(a + b*tan(e + f*x))^2, x)`



$$\mathbf{3.60} \quad \int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx$$

3.60.1	Optimal result . . . . .	409
3.60.2	Mathematica [A] (verified) . . . . .	410
3.60.3	Rubi [A] (verified) . . . . .	410
3.60.4	Maple [B] (verified) . . . . .	412
3.60.5	Fricas [B] (verification not implemented) . . . . .	413
3.60.6	Sympy [F] . . . . .	414
3.60.7	Maxima [B] (verification not implemented) . . . . .	415
3.60.8	Giac [F] . . . . .	415
3.60.9	Mupad [F(-1)] . . . . .	416

### 3.60.1 Optimal result

Integrand size = 20, antiderivative size = 654

$$\int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx = -\frac{2ib^2(c+dx)^2}{(a^2+b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a+ib)(ia+b)^2 (ia-b+(a+b)e^{2ie+2ifx}) f}$$

$$+ \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{4b(c+dx)^3}{3(ia-b)(a-ib)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d}$$

$$+ \frac{2b^2 d(c+dx) \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2}$$

$$+ \frac{2b(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2 (a+ib) f}$$

$$- \frac{2ib^2(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f}$$

$$- \frac{ib^2 d^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3}$$

$$+ \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2}$$

$$- \frac{2b^2 d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2}$$

$$+ \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2 (a+ib) f^3}$$

$$- \frac{ib^2 d^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3}$$

output

```
-2*I*b^2*(d*x+c)^2/(a^2+b^2)^2/f+2*b^2*(d*x+c)^2/(a+I*b)/(I*a+b)^2/(I*a-b+
(I*a+b)*exp(2*I*e+2*I*f*x))/f+1/3*(d*x+c)^3/(a-I*b)^2/d+4/3*b*(d*x+c)^3/(I
*a-b)/(a-I*b)^2/d-4/3*b^2*(d*x+c)^3/(a^2+b^2)^2/d+2*b^2*d*(d*x+c)*ln(1+(a-
I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2+2*b*(d*x+c)^2*ln(1+(a-
I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f-2*I*b^2*(d*x+c)^2*ln(1+(
a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f-I*b^2*d^2*polylog(2,-(a-I
*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3+2*b*d*(d*x+c)*polylog(2,-(
a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(I*a-b)/(a-I*b)^2/f^2-2*b^2*d*(d*x+c)*p
olylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2+b*d^2*polylo
g(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f^3-I*b^2*d^2*p
olylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3
```

$$3.60. \quad \int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx$$

### 3.60.2 Mathematica [A] (verified)

Time = 7.53 (sec) , antiderivative size = 761, normalized size of antiderivative = 1.16

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx$$

$$2b \left( \frac{12c(bd+acf)x}{a-ib} - \frac{12c(-ib(-1+e^{2ie})+a(1+e^{2ie}))(bd+acf)x}{a^2+b^2} + \frac{6d(bd+2acf)x^2}{a-ib} + \frac{4ad^2fx^3}{a-ib} + \frac{6d(-ib(-1+e^{2ie})+a(1+e^{2ie}))(bd+2acf)x \log\left(1 + \frac{(a+ib)e^{-ix}}{a-ib}\right)}{(a+ib)(ia+b)f} \right)$$


---

input `Integrate[(c + d*x)^2/(a + b*Tan[e + f*x])^2,x]`

output

```
((2*b*((12*c*(b*d + a*c*f)*x)/(a - I*b) - (12*c*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e))))*(b*d + a*c*f)*x)/(a^2 + b^2) + (6*d*(b*d + 2*a*c*f)*x^2)/(a - I*b) + (4*a*d^2*f*x^3)/(a - I*b) + (6*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e))))*(b*d + 2*a*c*f)*x*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)*f) + (6*a*d^2*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e))))*x^2*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)) + (6*c*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e))))*(b*d + a*c*f)*Log[a + I*b + (a - I*b)*E^((2*I)*(e + f*x)))]/((a + I*b)*(I*a + b)*f) + (3*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e))))*(b*d + 2*a*c*f)*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/((a^2 + b^2)*f^2) + (3*a*d^2*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e))))*(2*f*x*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - I*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f^2))/((b - b*E^((2*I)*e) - I*a*(1 + E^((2*I)*e))) + ((a^2 - b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cos[f*x] + (a^2 + b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cos[2*e + f*x] + 2*b*(3*b*(c + d*x)^2 + a*f*x*(3*c^2 + 3*c*d*x + d^2*x^2))*Sin[f*x])/((a*Cos[e] + b*SIN[e])*(a*Cos[e + f*x] + b*SIN[e + f*x])))/(6*(a^2 + b^2)*f)
```

### 3.60.3 Rubi [A] (verified)

Time = 1.81 (sec) , antiderivative size = 654, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {3042, 4217, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.60.  $\int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx$

$$\begin{aligned}
& \int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx \\
& \quad \downarrow \text{4217} \\
& \int \left( -\frac{4b^2(c+dx)^2}{(b+ia)^2 \left( ia \left( 1 - \frac{ib}{a} \right) e^{2ie+2ifx} + ia \left( 1 + \frac{ib}{a} \right) \right)^2} + \frac{4b(c+dx)^2}{(a-ib)^2 \left( ia \left( 1 - \frac{ib}{a} \right) e^{2ie+2ifx} + ia \left( 1 + \frac{ib}{a} \right) \right)} + \frac{(c+dx)^2}{(a-ib)^2} \right) dx \\
& \quad \downarrow \text{2009} \\
& -\frac{2b^2 d(c+dx) \operatorname{PolyLog} \left( 2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f^2 (a^2+b^2)^2} + \frac{2b^2 d(c+dx) \log \left( 1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f^2 (a^2+b^2)^2} - \\
& \frac{2ib^2(c+dx)^2 \log \left( 1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f(a^2+b^2)^2} - \frac{2ib^2(c+dx)^2}{f(a^2+b^2)^2} - \frac{4b^2(c+dx)^3}{3d(a^2+b^2)^2} - \\
& \frac{ib^2 d^2 \operatorname{PolyLog} \left( 2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f^3 (a^2+b^2)^2} - \frac{ib^2 d^2 \operatorname{PolyLog} \left( 3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f^3 (a^2+b^2)^2} + \\
& \frac{2b^2(c+dx)^2}{f(a+ib)(b+ia)^2 \left( (b+ia)e^{2ie+2ifx} + ia - b \right)} + \frac{2bd(c+dx) \operatorname{PolyLog} \left( 2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f^2(-b+ia)(a-ib)^2} + \\
& \frac{2b(c+dx)^2 \log \left( 1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f(a-ib)^2(a+ib)} + \frac{4b(c+dx)^3}{3d(-b+ia)(a-ib)^2} + \frac{(c+dx)^3}{3d(a-ib)^2} + \\
& \frac{bd^2 \operatorname{PolyLog} \left( 3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{f^3(a-ib)^2(a+ib)}
\end{aligned}$$

input `Int[(c + d*x)^2/(a + b*Tan[e + f*x])^2,x]`

```
output ((-2*I)*b^2*(c + d*x)^2)/((a^2 + b^2)^2*f) + (2*b^2*(c + d*x)^2)/((a + I*b)
)*(I*a + b)^2*(I*a - b + (I*a + b)*E^((2*I)*e + (2*I)*f*x))*f) + (c + d*x)
^3/(3*(a - I*b)^2*d) + (4*b*(c + d*x)^3)/(3*(I*a - b)*(a - I*b)^2*d) - (4*
b^2*(c + d*x)^3)/(3*(a^2 + b^2)^2*d) + (2*b^2*d*(c + d*x)*Log[1 + ((a - I*
b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((a^2 + b^2)^2*f^2) + (2*b*(c + d
*x)^2*Log[1 + ((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)]/((a - I*b)^2*
(a + I*b)*f) - ((2*I)*b^2*(c + d*x)^2*Log[1 + ((a - I*b)*E^((2*I)*e + (2*I)
)*f*x))/(a + I*b)]/((a^2 + b^2)^2*f) - (I*b^2*d^2*PolyLog[2, -(((a - I*b)
)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)))/((a^2 + b^2)^2*f^3) + (2*b*d*(c + d
*x)*PolyLog[2, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)))]/((I*a -
b)*(a - I*b)^2*f^2) - (2*b^2*d*(c + d*x)*PolyLog[2, -(((a - I*b)*E^((2*I)*
e + (2*I)*f*x))/(a + I*b)))]/((a^2 + b^2)^2*f^2) + (b*d^2*PolyLog[3, -(((a
- I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)))]/((a - I*b)^2*(a + I*b)*f^3)
- (I*b^2*d^2*PolyLog[3, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)))]
/((a^2 + b^2)^2*f^3)
```

### 3.60.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4217 Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 +
b^2 + (a - I*b)^2*E^(2*I*(e + f*x))))^(-n), x], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IGtQ[m, 0]
```

### 3.60.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3900 vs.  $2(588) = 1176$ .

Time = 0.97 (sec) , antiderivative size = 3901, normalized size of antiderivative = 5.96

method	result	size
risch	Expression too large to display	3901

---

3.60.  $\int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx$

```
input int((d*x+c)^2/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 2/(I*a+b)^2/f^2/(b-I*a)*b^2*e*d*a*c/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e)
)+b^2*exp(4*I*(f*x+e))+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b
^2)+I/(I*a+b)^2/f^3/(b-I*a)*b^2*e*d^2/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+
e))+b^2*exp(4*I*(f*x+e))+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2
+b^2)*a+4/(I*a+b)^2/f^2/(b-I*a)*b*e*d*a^2*c/(a+I*b)/(I*b-a)*arctan(1/2/a*e
xp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)-4/(I*a+b)^2/f^
2/(b-I*a)*b*e*d*a^2*c/(a+I*b)/(I*b-a)*arctan(1/b*a)+2/(I*a+b)^2/f^2/(b-I*a
)*b^2*c*d/(a+I*b)/(I*b-a)*arctan(1/b*a)*a-2/(I*a+b)^2/f^2/(b-I*a)*b^2*c*d/
(a+I*b)/(I*b-a)*arctan(1/2/a*exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f
*x+e))+1/2/b*a)*a-2/(I*a+b)^2/f^3/(b-I*a)*b*e^2*a^2*d^2/(a+I*b)/(I*b-a)*ar
ctan(1/2/a*exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)-4*I
/(I*a+b)^2/f^3/(b-I*a)*b*e^2*a*d^2/(a+I*b)*ln(exp(I*(f*x+e)))+2*I/(I*a+b)
^2/f/(b-I*a)*b^2*a*c^2/(a+I*b)/(I*b-a)*arctan(1/2/a*exp(2*I*(f*x+e))*b-1/2
*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)-2*I/(I*a+b)^2/f/(b-I*a)*b^2*a*c^2/(
a+I*b)/(I*b-a)*arctan(1/b*a)-2*I/(I*a+b)^2/f^3/(b-I*a)*b/(a+I*b)*e^2*a*d^2
*ln(1-(I*b-a)*exp(2*I*(f*x+e)))/(a+I*b))-2*I/(I*a+b)^2/f^3/(b-I*a)*b^3*e*d^
2/(a+I*b)/(I*b-a)*arctan(1/2/a*exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*
(f*x+e))+1/2/b*a)+2*I/(I*a+b)^2/f^3/(b-I*a)*b^3*e*d^2/(a+I*b)/(I*b-a)*arct
an(1/b*a)+2*I/(I*a+b)^2/f/(b-I*a)*b/(a+I*b)*d^2*a*ln(1-(I*b-a)*exp(2*I*(f*
x+e)))/(a+I*b))*x^2-2*I/(I*a+b)^2/f^2/(b-I*a)*b^3*c*d/(a+I*b)/(I*b-a)*ar...
```

### 3.60.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1576 vs.  $2(534) = 1068$ .

Time = 0.29 (sec) , antiderivative size = 1576, normalized size of antiderivative = 2.41

$$\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx = \text{Too large to display}$$

```
input integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="fracas")
```

```

output 1/6*(2*(a^3 - a*b^2)*d^2*f^3*x^3 - 6*b^3*c^2*f^2 - 6*(b^3*d^2*f^2 - (a^3 -
a*b^2)*c*d*f^3)*x^2 - 6*(2*b^3*c*d*f^2 - (a^3 - a*b^2)*c^2*f^3)*x - 3*(-2
*I*a^2*b*d^2*f*x - 2*I*a^2*b*c*d*f - I*a*b^2*d^2 + (-2*I*a*b^2*d^2*f*x - 2
*I*a*b^2*c*d*f - I*b^3*d^2)*tan(f*x + e))*dilog(2*((I*a*b - b^2)*tan(f*x +
e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*t
an(f*x + e)^2 + a^2 + b^2) + 1) - 3*(2*I*a^2*b*d^2*f*x + 2*I*a^2*b*c*d*f +
I*a*b^2*d^2 + (2*I*a*b^2*d^2*f*x + 2*I*a*b^2*c*d*f + I*b^3*d^2)*tan(f*x +
e))*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*
b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 6
*(a^2*b*d^2*f^2*x^2 - a^2*b*d^2*e^2 + 2*a^2*b*c*d*e*f + a*b^2*d^2*e + (2*a
^2*b*c*d*f^2 + a*b^2*d^2*f)*x + (a*b^2*d^2*f^2*x^2 - a*b^2*d^2*e^2 + 2*a*b
^2*c*d*e*f + b^3*d^2*e + (2*a*b^2*c*d*f^2 + b^3*d^2*f)*x)*tan(f*x + e))*lo
g(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)
*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(a^2*b*d^2*f^
2*x^2 - a^2*b*d^2*e^2 + 2*a^2*b*c*d*e*f + a*b^2*d^2*e + (2*a^2*b*c*d*f^2 +
a*b^2*d^2*f)*x + (a*b^2*d^2*f^2*x^2 - a*b^2*d^2*e^2 + 2*a*b^2*c*d*e*f + b
^3*d^2*e + (2*a*b^2*c*d*f^2 + b^3*d^2*f)*x)*tan(f*x + e))*log(-2*((-I*a*b
- b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e
))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(a^2*b*d^2*e^2 + a^2*b*c^
2*f^2 - a*b^2*d^2*e - (2*a^2*b*c*d*e - a*b^2*c*d)*f + (a*b^2*d^2*e^2 + ...

```

### 3.60.6 Sympy [F]

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx$$

```
input integrate((d*x+c)**2/(a+b*tan(f*x+e))**2,x)
```

```
output Integral((c + d*x)**2/(a + b*tan(e + f*x))**2, x)
```

### 3.60.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2530 vs.  $2(534) = 1068$ .

Time = 1.02 (sec) , antiderivative size = 2530, normalized size of antiderivative = 3.87

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

```
input integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output -1/3*(6*c*d*e*(2*a*b*log(b*tan(f*x + e) + a)/((a^4 + 2*a^2*b^2 + b^4)*f) -
a*b*log(tan(f*x + e)^2 + 1)/((a^4 + 2*a^2*b^2 + b^4)*f) - b/((a^2*b + b^3)
)*f*tan(f*x + e) + (a^3 + a*b^2)*f) + (a^2 - b^2)*(f*x + e)/((a^4 + 2*a^2*
b^2 + b^4)*f)) - 3*(2*a*b*log(b*tan(f*x + e) + a)/(a^4 + 2*a^2*b^2 + b^4)
- a*b*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + (a^2 - b^2)*(f*x +
e)/(a^4 + 2*a^2*b^2 + b^4) - b/(a^3 + a*b^2 + (a^2*b + b^3)*tan(f*x + e))
)*c^2 - ((a^3 - I*a^2*b + a*b^2 - I*b^3)*(f*x + e)^3*d^2 + 3*(a^3 - I*a^2*
b + a*b^2 - I*b^3)*(f*x + e)*d^2*e^2 - 6*(-I*a*b^2 + b^3)*d^2*e^2 - 3*((a^
3 - I*a^2*b + a*b^2 - I*b^3)*d^2*e - (a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d*f
)*(f*x + e)^2 - 6*((-I*a^2*b + a*b^2)*d^2*e^2 + (I*a*b^2 - b^3)*d^2*e + (-
I*a*b^2 + b^3)*c*d*f + ((-I*a^2*b - a*b^2)*d^2*e^2 + (I*a*b^2 + b^3)*d^2*e
+ (-I*a*b^2 - b^3)*c*d*f)*cos(2*f*x + 2*e) + ((a^2*b - I*a*b^2)*d^2*e^2 -
(a*b^2 - I*b^3)*d^2*e + (a*b^2 - I*b^3)*c*d*f)*sin(2*f*x + 2*e))*arctan2(
-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2
*f*x + 2*e) + a) - 6*((I*a^2*b - a*b^2)*(f*x + e)^2*d^2 + (2*(-I*a^2*b + a
*b^2)*d^2*e + 2*(I*a^2*b - a*b^2)*c*d*f + (I*a*b^2 - b^3)*d^2)*(f*x + e) +
((I*a^2*b + a*b^2)*(f*x + e)^2*d^2 + (2*(-I*a^2*b - a*b^2)*d^2*e + 2*(I*a
^2*b + a*b^2)*c*d*f + (I*a*b^2 + b^3)*d^2)*(f*x + e))*cos(2*f*x + 2*e) - (
a^2*b - I*a*b^2)*(f*x + e)^2*d^2 - (2*(a^2*b - I*a*b^2)*d^2*e - 2*(a^2*b
- I*a*b^2)*c*d*f - (a*b^2 - I*b^3)*d^2)*(f*x + e))*sin(2*f*x + 2*e))*ar...
```

### 3.60.8 Giac [F]

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \int \frac{(dx + c)^2}{(b \tan(fx + e) + a)^2} dx$$

```
input integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="giac")
```

---

3.60.  $\int \frac{(c+dx)^2}{(a+b \tan(e+fx))^2} dx$



output `integrate((d*x + c)^2/(b*tan(f*x + e) + a)^2, x)`

### 3.60.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx$$

input `int((c + d*x)^2/(a + b*tan(e + f*x))^2,x)`

output `int((c + d*x)^2/(a + b*tan(e + f*x))^2, x)`

### 3.61 $\int \frac{c+dx}{(a+b \tan(e+fx))^2} dx$

3.61.1	Optimal result . . . . .	417
3.61.2	Mathematica [B] (verified) . . . . .	418
3.61.3	Rubi [A] (verified) . . . . .	419
3.61.4	Maple [B] (verified) . . . . .	422
3.61.5	Fricas [B] (verification not implemented) . . . . .	423
3.61.6	Sympy [F] . . . . .	424
3.61.7	Maxima [B] (verification not implemented) . . . . .	424
3.61.8	Giac [F] . . . . .	425
3.61.9	Mupad [F(-1)] . . . . .	426

#### 3.61.1 Optimal result

Integrand size = 18, antiderivative size = 214

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = -\frac{(c + dx)^2}{2(a^2 + b^2)d} + \frac{(bd + 2acf + 2adf x)^2}{4a(a + ib)(a^2 + b^2)df^2}$$

$$+ \frac{b(bd + 2acf + 2adf x) \log\left(1 + \frac{(a^2 + b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2 + b^2)^2 f^2}$$

$$- \frac{iabd \operatorname{PolyLog}\left(2, -\frac{(a^2 + b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2 + b^2)^2 f^2}$$

$$- \frac{b(c + dx)}{(a^2 + b^2) f(a + b \tan(e + fx))}$$

output

```
-1/2*(d*x+c)^2/(a^2+b^2)/d+1/4*(2*a*d*f*x+2*a*c*f+b*d)^2/a/(a+I*b)/(a^2+b^2)/d/f^2+b*(2*a*d*f*x+2*a*c*f+b*d)*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)^2/f^2-I*a*b*d*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)^2/f^2-b*(d*x+c)/(a^2+b^2)/f/(a+b*tan(f*x+e))
```

### 3.61.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 745 vs.  $2(214) = 428$ .

Time = 7.89 (sec) , antiderivative size = 745, normalized size of antiderivative = 3.48

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{(e + fx)(-2de + 2cf + d(e + fx)) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{2(a - ib)(a + ib)f^2(a + b \tan(e + fx))^2}$$

$$+ \frac{b^2 d(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{a(a - ib)(a + ib)(a^2 + b^2)f^2(a + b \tan(e + fx))^2}$$

$$- \frac{2bde(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{(a - ib)(a + ib)(a^2 + b^2)f^2(a + b \tan(e + fx))^2}$$

$$+ \frac{2bc(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{(a - ib)(a + ib)(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

$$d \left( e^{i \arctan(\frac{a}{b})} (e + fx)^2 + \frac{a \left( i(e + fx)(-\pi + 2 \arctan(\frac{a}{b})) - \pi \log(1 + e^{-2i(e + fx)}) - 2(e + fx + \arctan(\frac{a}{b})) \log\left(1 - e^{2i(e + fx + \arctan(\frac{a}{b}))}\right)\right)}{\sqrt{1 + \frac{a^2}{b^2}}}$$


---


$$+ \frac{\sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))(-b^2 d e \sin(e + fx) + b^2 c f \sin(e + fx) + b^2 d(e + fx) \sin(e + fx))}{a(a - ib)(a + ib)f^2(a + b \tan(e + fx))^2}$$

input `Integrate[(c + d*x)/(a + b*Tan[e + f*x])^2,x]`

output

```

((e + f*x)*(-2*d*e + 2*c*f + d*(e + f*x))*Sec[e + f*x]^2*(a*Cos[e + f*x] +
b*Sin[e + f*x])^2)/(2*(a - I*b)*(a + I*b)*f^2*(a + b*Tan[e + f*x])^2) + (
b^2*d*(-(b*(e + f*x)) + a*Log[a*Cos[e + f*x] + b*Sin[e + f*x]])*Sec[e + f*
x]^2*(a*Cos[e + f*x] + b*Sin[e + f*x])^2)/(a*(a - I*b)*(a + I*b)*(a^2 + b^
2)*f^2*(a + b*Tan[e + f*x])^2) - (2*b*d*e*(-(b*(e + f*x)) + a*Log[a*Cos[e
+ f*x] + b*Sin[e + f*x]])*Sec[e + f*x]^2*(a*Cos[e + f*x] + b*Sin[e + f*x])
^2)/((a - I*b)*(a + I*b)*(a^2 + b^2)*f^2*(a + b*Tan[e + f*x])^2) + (2*b*c*
(-(b*(e + f*x)) + a*Log[a*Cos[e + f*x] + b*Sin[e + f*x]])*Sec[e + f*x]^2*(
a*Cos[e + f*x] + b*Sin[e + f*x])^2)/((a - I*b)*(a + I*b)*(a^2 + b^2)*f*(a
+ b*Tan[e + f*x])^2) - (d*(E^(I*ArcTan[a/b]))*(e + f*x)^2 + (a*(I*(e + f*x)
*(-Pi + 2*ArcTan[a/b]) - Pi*Log[1 + E^((-2*I)*(e + f*x))] - 2*(e + f*x + A
rcTan[a/b])*Log[1 - E^((2*I)*(e + f*x + ArcTan[a/b]))] + Pi*Log[Cos[e + f*
x]] + 2*ArcTan[a/b]*Log[Sin[e + f*x + ArcTan[a/b]]] + I*PolyLog[2, E^((2*I)
*(e + f*x + ArcTan[a/b]))])))/(Sqrt[1 + a^2/b^2]*b)*Sec[e + f*x]^2*(a*Cos
[e + f*x] + b*Sin[e + f*x])^2)/((a - I*b)*(a + I*b)*Sqrt[(a^2 + b^2)/b^2]*
f^2*(a + b*Tan[e + f*x])^2) + (Sec[e + f*x]^2*(a*Cos[e + f*x] + b*Sin[e +
f*x])*(-(b^2*d*e*Sin[e + f*x]) + b^2*c*f*Sin[e + f*x] + b^2*d*(e + f*x)*Si
n[e + f*x]))/(a*(a - I*b)*(a + I*b)*f^2*(a + b*Tan[e + f*x])^2)

```

### 3.61.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$ , Rules used = {3042, 4216, 3042, 4215, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4216} \\
 & \frac{\int \frac{bd+2afxd+2acf}{a+b \tan(e+fx)} dx}{f(a^2 + b^2)} - \frac{b(c + dx)}{f(a^2 + b^2)(a + b \tan(e + fx))} - \frac{(c + dx)^2}{2d(a^2 + b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{bd+2afxd+2acf}{a+b \tan(e+fx)} dx}{f(a^2 + b^2)} - \frac{b(c + dx)}{f(a^2 + b^2)(a + b \tan(e + fx))} - \frac{(c + dx)^2}{2d(a^2 + b^2)}
 \end{aligned}$$

---

3.61.  $\int \frac{c+dx}{(a+b \tan(e+fx))^2} dx$

$$\begin{aligned}
& \downarrow 4215 \\
& \frac{2ib \int \frac{e^{2i(e+fx)}(bd+2afx+2acf)}{(a+ib)^2+(a^2+b^2)e^{2i(e+fx)}} dx + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)} \\
& \downarrow 2620 \\
& \frac{2ib \left( \frac{iad \int \log\left(\frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1\right) dx}{a^2+b^2} - \frac{i(2acf+2adf+bd) \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f(a^2+b^2)} \right) + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \\
& \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)} \\
& \downarrow 2715 \\
& \frac{2ib \left( \frac{ad \int e^{-2i(e+fx)} \log\left(\frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1\right) de^{2i(e+fx)}}{2f(a^2+b^2)} - \frac{i(2acf+2adf+bd) \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f(a^2+b^2)} \right) + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \\
& \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)} \\
& \downarrow 2838 \\
& \frac{2ib \left( -\frac{i(2acf+2adf+bd) \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f(a^2+b^2)} - \frac{ad \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f(a^2+b^2)} \right) + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \\
& \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)}
\end{aligned}$$

input `Int[(c + d*x)/(a + b*Tan[e + f*x])^2,x]`

output `-1/2*(c + d*x)^2/((a^2 + b^2)*d) + ((b*d + 2*a*c*f + 2*a*d*f*x)^2/(4*a*(a + I*b)*d*f) + (2*I)*b*(((1/2*I)*(b*d + 2*a*c*f + 2*a*d*f*x)*Log[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) - (a*d*PolyLog[2, -((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/(2*(a^2 + b^2)*f))/((a^2 + b^2)*f) - (b*(c + d*x))/((a^2 + b^2)*f*(a + b*Tan[e + f*x]))`

## 3.61.3.1 Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_)]], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4215 `Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Simp[2*I*b Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

rule 4216 `Int[((c_) + (d_)*(x_))/((a_) + (b_)*tan[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[-(c + d*x)^2/(2*d*(a^2 + b^2)), x] + (Simp[1/(f*(a^2 + b^2)) Int[(b*d + 2*a*c*f + 2*a*d*f*x)/(a + b*Tan[e + f*x]), x], x] - Simp[b*((c + d*x)/(f*(a^2 + b^2)*(a + b*Tan[e + f*x]))], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]`

### 3.61.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2000 vs.  $2(202) = 404$ .

Time = 0.86 (sec) , antiderivative size = 2001, normalized size of antiderivative = 9.35

method	result	size
risch	Expression too large to display	2001

input `int((d*x+c)/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & 4*I/(I*a+b)^2/f^2/(b-I*a)*b*a*d*e/(a+I*b)*\ln(\exp(I*(f*x+e)))+2*I/(I*a+b)^2 \\
 & /f/(b-I*a)*b^2*a*c/(a+I*b)/(I*b-a)*\arctan(1/2/a*\exp(2*I*(f*x+e))*b-1/2*b/a \\
 & +1/2/b*a*\exp(2*I*(f*x+e))+1/2/b*a)-2*I/(I*a+b)^2/f/(b-I*a)*b^2*a*c/(a+I*b) \\
 & / (I*b-a)*\arctan(1/b*a)+2*I/(I*a+b)^2/f^2/(b-I*a)*b/(a+I*b)*a*d*\ln(1-(I*b-a) \\
 & )*\exp(2*I*(f*x+e))/(a+I*b))*e-1/2*I/(I*a+b)^2/f^2/(b-I*a)*b^2*d/(a+I*b)/(I \\
 & *b-a)*\ln(a^2*\exp(4*I*(f*x+e))+b^2*\exp(4*I*(f*x+e))+2*a^2*\exp(2*I*(f*x+e))- \\
 & 2*b^2*\exp(2*I*(f*x+e))+a^2+b^2)*a-I/(I*a+b)^2/f/(b-I*a)*b*a^2*c/(a+I*b)/(I \\
 & *b-a)*\ln(a^2*\exp(4*I*(f*x+e))+b^2*\exp(4*I*(f*x+e))+2*a^2*\exp(2*I*(f*x+e))- \\
 & 2*b^2*\exp(2*I*(f*x+e))+a^2+b^2)+2*I/(I*a+b)^2/f/(b-I*a)*b/(a+I*b)*a*d*\ln(1 \\
 & -(I*b-a)*\exp(2*I*(f*x+e))/(a+I*b))*x-1/2/(2*I*a*b-a^2+b^2)*d*x^2-1/(2*I*a* \\
 & b-a^2+b^2)*c*x+1/(I*a+b)^2/f^2/(b-I*a)*b/(a+I*b)*a*d*polylog(2,(I*b-a)*\exp \\
 & (2*I*(f*x+e))/(a+I*b))-1/2/(I*a+b)^2/f^2/(b-I*a)*b^3*d/(a+I*b)/(I*b-a)*\ln( \\
 & a^2*\exp(4*I*(f*x+e))+b^2*\exp(4*I*(f*x+e))+2*a^2*\exp(2*I*(f*x+e))-2*b^2*\exp \\
 & (2*I*(f*x+e))+a^2+b^2)+2/(I*a+b)^2/f^2/(b-I*a)*b/(a+I*b)*a*d*e^2-2*I/(I*a+ \\
 & b)^2/f^2/(b-I*a)*b^2*d/(a+I*b)*\ln(\exp(I*(f*x+e)))-2/(I*a+b)^2/f^2/(b-I*a)* \\
 & b*a^2*d*e/(a+I*b)/(I*b-a)*\arctan(1/b*a)+2/(I*a+b)^2/f^2/(b-I*a)*b*a^2*d*e/ \\
 & (a+I*b)/(I*b-a)*\arctan(1/2/a*\exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*\exp(2*I*(f \\
 & *x+e))+1/2/b*a)+1/(I*a+b)^2/f^2/(b-I*a)*b^2*a*d*e/(a+I*b)/(I*b-a)*\ln(a^2*e \\
 & xp(4*I*(f*x+e))+b^2*\exp(4*I*(f*x+e))+2*a^2*\exp(2*I*(f*x+e))-2*b^2*\exp(2*I* \\
 & (f*x+e))+a^2+b^2)+4/(I*a+b)^2/f/(b-I*a)*b/(a+I*b)*a*d*e*x-1/(I*a+b)^2/f...
 \end{aligned}$$

### 3.61.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 843 vs.  $2(197) = 394$ .

Time = 0.27 (sec) , antiderivative size = 843, normalized size of antiderivative = 3.94

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{(a^3 - ab^2)df^2x^2 - 2b^3cf - 2(b^3df - (a^3 - ab^2)cf^2)x + (iab^2d \tan(fx + e) + ia^2bd) \operatorname{Li}_2\left(\frac{2((iab - b^2) \tan(fx + e) + a^2)}{(a^2 + b^2) \tan(fx + e)^2 + a^2 + b^2}\right)}{(a^2 + b^2) \tan(fx + e)^2 + a^2 + b^2}$$

input `integrate((d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="fracas")`

output

```

1/2*((a^3 - a*b^2)*d*f^2*x^2 - 2*b^3*c*f - 2*(b^3*d*f - (a^3 - a*b^2)*c*f^
2)*x + (I*a*b^2*d*tan(f*x + e) + I*a^2*b*d)*dilog(2*((I*a*b - b^2)*tan(f*x
+ e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)
*tan(f*x + e)^2 + a^2 + b^2) + 1) + (-I*a*b^2*d*tan(f*x + e) - I*a^2*b*d)*
dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I
*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 2*(a^2
*b*d*f*x + a^2*b*d*e + (a*b^2*d*f*x + a*b^2*d*e)*tan(f*x + e))*log(-2*((I*
a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x
+ e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(a^2*b*d*f*x + a^2*b*d
*e + (a*b^2*d*f*x + a*b^2*d*e)*tan(f*x + e))*log(-2*((-I*a*b - b^2)*tan(f*
x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^
2)*tan(f*x + e)^2 + a^2 + b^2)) - (2*a^2*b*d*e - 2*a^2*b*c*f - a*b^2*d + (
2*a*b^2*d*e - 2*a*b^2*c*f - b^3*d)*tan(f*x + e))*log(((I*a*b + b^2)*tan(f*
x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1
)) - (2*a^2*b*d*e - 2*a^2*b*c*f - a*b^2*d + (2*a*b^2*d*e - 2*a*b^2*c*f - b
^3*d)*tan(f*x + e))*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a
^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + ((a^2*b - b^3)*d*f^2*x^2
+ 2*a*b^2*c*f + 2*(a*b^2*d*f + (a^2*b - b^3)*c*f^2)*x)*tan(f*x + e))/((a^
4*b + 2*a^2*b^3 + b^5)*f^2*tan(f*x + e) + (a^5 + 2*a^3*b^2 + a*b^4)*f^2)

```



**3.61.6 Sympy [F]**

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx$$

input `integrate((d*x+c)/(a+b*tan(f*x+e))**2,x)`

output `Integral((c + d*x)/(a + b*tan(e + f*x))**2, x)`

**3.61.7 Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1156 vs.  $2(197) = 394$ .

Time = 0.71 (sec) , antiderivative size = 1156, normalized size of antiderivative = 5.40

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output

```

1/2*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d*f^2*x^2 + 2*(a^3 - I*a^2*b + a*b^2
- I*b^3)*c*f^2*x - 4*(-I*a*b^2 + b^3)*c*f - 2*(2*(-I*a^2*b + a*b^2)*c*f +
(-I*a*b^2 + b^3)*d + (2*(-I*a^2*b - a*b^2)*c*f + (-I*a*b^2 - b^3)*d)*cos(2
*f*x + 2*e) + (2*(a^2*b - I*a*b^2)*c*f + (a*b^2 - I*b^3)*d)*sin(2*f*x + 2*
e))*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*
e) + b*sin(2*f*x + 2*e) + a) - 4*((I*a^2*b + a*b^2)*d*f*x*cos(2*f*x + 2*e)
- (a^2*b - I*a*b^2)*d*f*x*sin(2*f*x + 2*e) + (I*a^2*b - a*b^2)*d*f*x)*arc
tan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2),
(2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 +
b^2)) + ((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*d*f^2*x^2 + 2*((a^3 - 3*I*a^
2*b - 3*a*b^2 + I*b^3)*c*f^2 - 2*(I*a*b^2 + b^3)*d*f)*x*cos(2*f*x + 2*e)
- 2*((I*a^2*b + a*b^2)*d*cos(2*f*x + 2*e) - (a^2*b - I*a*b^2)*d*sin(2*f*x
+ 2*e) + (I*a^2*b - a*b^2)*d)*dilog((I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a +
b)) + (2*(a^2*b + I*a*b^2)*c*f + (a*b^2 + I*b^3)*d + (2*(a^2*b - I*a*b^2)*
c*f + (a*b^2 - I*b^3)*d)*cos(2*f*x + 2*e) - (2*(-I*a^2*b - a*b^2)*c*f - (I
*a*b^2 + b^3)*d)*sin(2*f*x + 2*e))*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*
a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2
- b^2)*cos(2*f*x + 2*e)) + 2*((a^2*b - I*a*b^2)*d*f*x*cos(2*f*x + 2*e) -
(-I*a^2*b - a*b^2)*d*f*x*sin(2*f*x + 2*e) + (a^2*b + I*a*b^2)*d*f*x)*log((
a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*s...

```

### 3.61.8 Giac [F]

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{dx + c}{(b \tan(fx + e) + a)^2} dx$$

input `integrate((d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)/(b*tan(f*x + e) + a)^2, x)`

**3.61.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx$$

input `int((c + d*x)/(a + b*tan(e + f*x))^2,x)`output `int((c + d*x)/(a + b*tan(e + f*x))^2, x)`

### 3.62 $\int \frac{1}{(c+dx)(a+b \tan(e+fx))^2} dx$

3.62.1	Optimal result	427
3.62.2	Mathematica [N/A]	427
3.62.3	Rubi [N/A]	428
3.62.4	Maple [N/A] (verified)	429
3.62.5	Fricas [N/A]	429
3.62.6	Sympy [N/A]	429
3.62.7	Maxima [N/A]	430
3.62.8	Giac [N/A]	430
3.62.9	Mupad [N/A]	431

#### 3.62.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b \tan(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b \tan(e+fx))^2}, x\right)$$

output `Unintegrable(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)`

#### 3.62.2 Mathematica [N/A]

Not integrable

Time = 13.94 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b \tan(e+fx))^2} dx = \int \frac{1}{(c+dx)(a+b \tan(e+fx))^2} dx$$

input `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])^2),x]`

output `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])^2), x]`

### 3.62.3 Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$$

↓ 3042

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$$

↓ 4223

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$$

input `Int[1/((c + d*x)*(a + b*Tan[e + f*x])^2),x]`

output `$Aborted`

#### 3.62.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.62.4 Maple [N/A] (verified)**

Not integrable

Time = 0.55 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx+c)(a+b\tan(fx+e))^2} dx$$

input `int(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)`output `int(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)`**3.62.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(dx+c)(b\tan(fx+e)+a)^2} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`output `integral(1/(a^2*d*x + a^2*c + (b^2*d*x + b^2*c)*tan(f*x + e)^2 + 2*(a*b*d*x + a*b*c)*tan(f*x + e)), x)`**3.62.6 Sympy [N/A]**

Not integrable

Time = 1.67 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(a+b\tan(e+fx))^2(c+dx)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e))**2,x)`output `Integral(1/((a + b*tan(e + f*x))**2*(c + d*x)), x)`

**3.62.7 Maxima [N/A]**

Not integrable

Time = 8.73 (sec) , antiderivative size = 1498, normalized size of antiderivative = 74.90

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(dx+c)(b\tan(fx+e)+a)^2} dx$$

```
input integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output (((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*cos(2*f*x + 2*e)^2*log(d*x + c) + (
(a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*log(d*x + c)*sin(2*f*x + 2*e)^2 - 2*(
2*a*b^3*d - ((a^4 - 2*a^2*b^2 + b^4)*d*f*x + (a^4 - 2*a^2*b^2 + b^4)*c*f)*
log(d*x + c))*cos(2*f*x + 2*e) + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*
f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f + ((a^6 + 3*a^4*b^2 + 3*a^
2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f)*cos(2*f*
x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b
^2 + 3*a^2*b^4 + b^6)*c*d*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*
b^4 - b^6)*d^2*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d*f)*cos(2*f*x + 2*
e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^2*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*
c*d*f)*sin(2*f*x + 2*e))*integrate(2*(2*(2*a^2*b^2*d*f*x + 2*a^2*b^2*c*f -
a*b^3*d)*cos(2*f*x + 2*e) - (2*(a^3*b - a*b^3)*d*f*x + 2*(a^3*b - a*b^3)*
c*f - (a^2*b^2 - b^4)*d)*sin(2*f*x + 2*e))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 +
b^6)*d^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3
*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d
^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3*a^4*b^
2 + 3*a^2*b^4 + b^6)*c^2*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2
*b^4 + b^6)*d^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a
^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^
4*b^2 - a^2*b^4 - b^6)*d^2*f*x^2 + 2*(a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*...
```

**3.62.8 Giac [N/A]**

Not integrable

Time = 0.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(dx+c)(b\tan(fx+e)+a)^2} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate(1/((d*x + c)*(b*tan(f*x + e) + a)^2), x)`

### 3.62.9 Mupad [N/A]

Not integrable

Time = 4.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(a+b\tan(e+fx))^2 (c+dx)} dx$$

input `int(1/((a + b*tan(e + f*x))^2*(c + d*x)),x)`

output `int(1/((a + b*tan(e + f*x))^2*(c + d*x)), x)`



### 3.63 $\int \frac{1}{(c+dx)^2(a+b \tan(e+fx))^2} dx$

3.63.1	Optimal result . . . . .	432
3.63.2	Mathematica [N/A] . . . . .	432
3.63.3	Rubi [N/A] . . . . .	433
3.63.4	Maple [N/A] (verified) . . . . .	434
3.63.5	Fricas [N/A] . . . . .	434
3.63.6	Sympy [N/A] . . . . .	434
3.63.7	Maxima [N/A] . . . . .	435
3.63.8	Giac [N/A] . . . . .	435
3.63.9	Mupad [N/A] . . . . .	436

#### 3.63.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b \tan(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b \tan(e+fx))^2}, x\right)$$

output `Unintegrable(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)`

#### 3.63.2 Mathematica [N/A]

Not integrable

Time = 15.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b \tan(e+fx))^2} dx = \int \frac{1}{(c+dx)^2(a+b \tan(e+fx))^2} dx$$

input `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2),x]`

output `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2), x]`

### 3.63.3 Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c + dx)^2 (a + b \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{1}{(c + dx)^2 (a + b \tan(e + fx))^2} dx$$

↓ 4223

$$\int \frac{1}{(c + dx)^2 (a + b \tan(e + fx))^2} dx$$

input `Int[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2),x]`

output `$Aborted`

#### 3.63.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; Free Q[{a, b, c, d, e, f, m, n}, x]`

**3.63.4 Maple [N/A] (verified)**

Not integrable

Time = 0.65 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx+c)^2 (a+b \tan (fx+e))^2} dx$$

input `int(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)`output `int(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)`**3.63.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 96, normalized size of antiderivative = 4.80

$$\int \frac{1}{(c+dx)^2 (a+b \tan (e+fx))^2} dx = \int \frac{1}{(dx+c)^2 (b \tan (fx+e)+a)^2} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`output `integral(1/(a^2*d^2*x^2 + 2*a^2*c*d*x + a^2*c^2 + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*tan(f*x + e)^2 + 2*(a*b*d^2*x^2 + 2*a*b*c*d*x + a*b*c^2)*tan(f*x + e)), x)`**3.63.6 Sympy [N/A]**

Not integrable

Time = 3.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c+dx)^2 (a+b \tan (e+fx))^2} dx = \int \frac{1}{(a+b \tan (e+fx))^2 (c+dx)^2} dx$$

input `integrate(1/(d*x+c)**2/(a+b*tan(f*x+e))**2,x)`output `Integral(1/((a + b*tan(e + f*x))**2*(c + d*x)**2), x)`

**3.63.7 Maxima [N/A]**

Not integrable

Time = 27.22 (sec) , antiderivative size = 1977, normalized size of antiderivative = 98.85

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \int \frac{1}{(dx+c)^2(b\tan(fx+e)+a)^2} dx$$

```
input integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output -(a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*cos(2*f*x + 2*e)^2 + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*sin(2*f*x + 2*e)^2 + 2*(2*a*b^3*d + (a^4 - 2*a^2*b^2 + b^4)*d*f*x + (a^4 - 2*a^2*b^2 + b^4)*c*f)*cos(2*f*x + 2*e) - ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^3*f*x^2 + 2*(a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d^2*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c^2*d*f)*cos(2*f*x + 2*e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^3*f*x^2 + 2*(a^5*b + 2*a^3*b^3 + a*b^5)*c*d^2*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*c^2*d*f)*sin(2*f*x + 2*e))*integrate(4*(2*(a^2*b^2*d*f*x + a^2*b^2*c*f - a*b^3*d)*cos(2*f*x + 2*e) - ((a^3*b - a*b^3)*d*f*x + (a^3*b - a*b^3)*c*f - (a^2*b^2 - b^4)*d)*sin(2*f*x + 2*e))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^3 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x^2 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^3*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^3 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x^2 + 3*...
```

**3.63.8 Giac [N/A]**

Not integrable

Time = 22.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \int \frac{1}{(dx+c)^2(b\tan(fx+e)+a)^2} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate(1/((d*x + c)^2*(b*tan(f*x + e) + a)^2), x)`

### 3.63.9 Mupad [N/A]

Not integrable

Time = 3.99 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \int \frac{1}{(a+b\tan(e+fx))^2(c+dx)^2} dx$$

input `int(1/((a + b*tan(e + f*x))^2*(c + d*x)^2),x)`

output `int(1/((a + b*tan(e + f*x))^2*(c + d*x)^2), x)`

## APPENDIX

4.1 Listing of Grading functions . . . . .	437
--	-----

## 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

### 4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```



```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

### 4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(" ,
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

### 4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

*#main function*



```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

#### 4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```