

# Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.5-Secant/115-4.5.0-a-sec-<sup>m</sup>-b-trg-<sup>n</sup>

Nasser M. Abbasi

December 8, 2023

Compiled on December 8, 2023 at 8:02pm

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>detailed summary tables of results</b>	<b>20</b>
<b>3</b>	<b>Listing of integrals</b>	<b>112</b>
<b>4</b>	<b>Appendix</b>	<b>1747</b>

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	3
1.2	Results . . . . .	4
1.3	Time and leaf size Performance . . . . .	7
1.4	Performance based on number of rules Rubi used . . . . .	9
1.5	Performance based on number of steps Rubi used . . . . .	10
1.6	Solved integrals histogram based on leaf size of result . . . . .	11
1.7	Solved integrals histogram based on CPU time used . . . . .	12
1.8	Leaf size vs. CPU time used . . . . .	13
1.9	list of integrals with no known antiderivative . . . . .	14
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	14
1.11	list of integrals solved by CAS but failed verification . . . . .	14
1.12	Timing . . . . .	15
1.13	Verification . . . . .	15
1.14	Important notes about some of the results . . . . .	15
1.15	Design of the test system . . . . .	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 299 ]. This is test number [ 115 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.



## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 299 )	0.00 ( 0 )
Mathematica	100.00 ( 299 )	0.00 ( 0 )
Maple	75.92 ( 227 )	24.08 ( 72 )
Fricas	72.91 ( 218 )	27.09 ( 81 )
Maxima	31.10 ( 93 )	68.90 ( 206 )
Mupad	26.09 ( 78 )	73.91 ( 221 )
Giac	13.04 ( 39 )	86.96 ( 260 )
Sympy	9.70 ( 29 )	90.30 ( 270 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	89.632	1.003	9.365	0.000
Maple	39.130	8.361	28.428	24.080
Fricas	33.110	2.341	37.458	27.090
Maxima	23.077	8.027	0.000	68.896
Giac	10.368	2.676	0.000	86.957
Sympy	9.365	0.334	0.000	90.301
Mupad	0.000	26.087	0.000	73.913

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Maple	72	100.00	0.00	0.00
Fricas	81	100.00	0.00	0.00
Maxima	206	100.00	0.00	0.00
Mupad	221	0.00	100.00	0.00
Giac	260	99.23	0.00	0.77
Sympy	270	63.33	36.67	0.00

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.23
Rubi	0.33
Giac	0.34
Maxima	0.40
Mathematica	0.72
Maple	5.83
Mupad	9.45
Sympy	17.47

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Giac	40.05	1.36	35.00	1.03
Sympy	59.97	1.24	46.00	1.08
Mupad	61.22	1.19	46.00	1.05
Mathematica	62.39	0.89	56.00	0.86
Rubi	73.17	0.98	69.00	1.00
Fricas	160.93	1.59	87.00	1.27
Maxima	168.53	2.44	42.00	0.96
Maple	173.61	2.13	105.00	1.42

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution



## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

**Rubi** {}

**Mathematica** {}

**Maple** {}

**Maxima** {}

**Fricas** {}

**Sympy** {}

**Giac** {}

**Mupad** {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

**Rubi** {}

**Mathematica** {280, 281, 282, 283}

**Maple** {39, 40, 41, 42, 43, 44, 45, 46, 235, 237, 244, 253, 266, 268, 269, 270, 273, 277, 279}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### 1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### 1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### 1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

### 1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
June 27, 2023  
Design v0.6



# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	21
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	27
2.3	Detailed conclusion table specific for Rubi results . . . . .	102

## 2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi . . . . .	21
2.1.2	Mma . . . . .	22
2.1.3	Maple . . . . .	22
2.1.4	Fricas . . . . .	23
2.1.5	Maxima . . . . .	23
2.1.6	Giac . . . . .	24
2.1.7	Mupad . . . . .	25
2.1.8	Sympy . . . . .	25

### 2.1.1 Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

**B grade** { }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### 2.1.2 Mma

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 231, 233, 235, 237, 238, 240, 242, 244, 247, 249, 251, 253, 254, 256, 258, 260, 262, 264, 266, 268, 270, 271, 273, 275, 277, 279, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 296, 297, 298, 299 }

**B grade** { 42, 294, 295 }

**C grade** { 230, 232, 234, 236, 239, 241, 243, 245, 246, 248, 250, 252, 255, 257, 259, 261, 263, 265, 267, 269, 272, 274, 276, 278, 280, 281, 282, 283 }

**F normal fail** { }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.3 Maple

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 47, 48, 49, 50, 51, 52, 53, 54, 61, 62, 63, 64, 65, 66, 67, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 229, 230, 231, 232, 233, 234, 235, 237, 238, 239, 240, 242, 246, 247, 248, 249, 250, 251, 253, 254, 256, 258, 260, 262, 263, 264, 265, 267, 268, 270, 271, 275, 277, 279, 287, 288, 289 }

**B grade** { 9, 10, 11, 12, 13, 14, 15, 16, 228, 236, 241, 243, 244, 245, 252, 255, 257, 259, 261, 266, 272, 273, 274, 276, 278 }

**C grade** { 17, 18, 19, 20, 21, 22, 23, 24, 39, 40, 41, 42, 43, 44, 45, 46, 55, 56, 57, 58, 59, 60, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 269 }

**F normal fail** { 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 68, 69, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208,

209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 280, 281, 282, 283, 284, 285, 286, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

### 2.1.4 Fricas

**A grade { 2, 4, 5, 6, 7, 8, 39, 40, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 61, 62, 63, 64, 65, 66, 67, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 228, 229, 231, 233, 238, 240, 242, 247, 249, 254, 256, 262, 287, 288, 289 }**

**B grade { 1, 3, 41, 42, 251, 264, 271 }**

**C grade { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 55, 56, 57, 58, 59, 60, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 230, 232, 234, 235, 237, 239, 241, 243, 244, 246, 248, 250, 252, 253, 255, 257, 258, 260, 263, 265, 266, 268, 270, 272, 273, 275, 277, 279 }**

**F normal fail { 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 68, 69, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 236, 245, 259, 261, 267, 269, 274, 276, 278, 280, 281, 282, 283, 284, 285, 286, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }**

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

### 2.1.5 Maxima

**A grade { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 42, 43, 44, 45, 46, 50, 51, 52, 53, 54, 61, 62, 63, 64, 65, 66, 67, 135, 137, 138, 139, 140, 141, 145, 147, 148, 149, 150, 151, 155, 157, 158, 159, 160, 164, 165, 166, 167, 171, 172, 173, 174, 175, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 228, 229, 287, 288, 289 }**

**B grade { 47, 48, 49, 132, 133, 134, 136, 142, 143, 144, 146, 152, 153, 154, 156, 161, 162, 163, 168, 169, 170, 176, 177, 178 }**

**C grade { }**

**F normal fail** { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 55, 56, 57, 58, 59, 60, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.6 Giac

**A grade** { 2, 3, 4, 5, 6, 7, 8, 39, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 61, 62, 63, 64, 65, 138, 148, 225, 226, 228, 229 }

**B grade** { 1, 40, 41, 42, 139, 223, 224, 227 }

**C grade** { }

**F normal fail** { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 55, 56, 57, 58, 59, 60, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 140, 141, 142, 143, 144, 145, 146, 147, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { 66, 67 }

### 2.1.7 Mupad

**A grade** { }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 12, 20, 43, 51, 61, 62, 63, 64, 74, 133, 135, 137, 138, 139, 140, 141, 143, 145, 147, 148, 149, 150, 151, 152, 153, 155, 157, 158, 159, 160, 162, 164, 165, 166, 167, 169, 171, 172, 173, 174, 175, 177, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 228, 229, 231, 233, 238, 240, 242, 247, 249, 251, 254, 256, 262, 264, 271, 287, 288, 289 }

**C grade** { }

**F normal fail** { }

**F(-1) timeout fail** { 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 65, 66, 67, 68, 69, 70, 71, 72, 73, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 134, 136, 142, 144, 146, 154, 156, 161, 163, 168, 170, 176, 178, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 230, 232, 234, 235, 236, 237, 239, 241, 243, 244, 245, 246, 248, 250, 252, 253, 255, 257, 258, 259, 260, 261, 263, 265, 266, 267, 268, 269, 270, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

**F(-2) exception fail** { }

### 2.1.8 Sympy

**A grade** { 43, 44, 45, 46, 51, 52, 53, 54, 137, 138, 139, 147, 148, 157, 164, 165, 166, 167, 171, 172, 173, 174, 175, 179, 180, 181, 182, 183 }

**B grade** { 1 }

**C grade** { }

**F normal fail** { 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 48, 49, 50, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 80, 81, 82, 83, 84, 90, 91, 92, 101, 102, 103, 104, 105, 106, 107, 108, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 135, 136, 144, 145, 146, 163, 184, 185, 186, 187, 188, 189, 190, 191, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 218, 219, 220, 221, 222, 225, 226, 227, 234, 235, 236, 243, 244, 257, 258, 259, 260, 266, 267, 268, 280, 281, 282, 283, 284, 285, 286, 287, 291, 292, 293, 294, 295, 297, 298, 299 }

**F(-1) timeout fail** { 9, 17, 39, 47, 61, 77, 79, 85, 86, 87, 88, 89, 93, 94, 95, 96, 97, 98, 99, 100, 109, 120, 132, 133, 134, 140, 141, 142, 143, 149, 150, 151, 152, 153, 154, 155, 156, 158, 159, 160,

161, 162, 168, 169, 170, 176, 177, 178, 192, 193, 204, 216, 217, 223, 224, 228, 229, 230, 231, 232, 233, 237, 238, 239, 240, 241, 242, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 261, 262, 263, 264, 265, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 288, 289, 290, 296 }

**F(-2) exception fail { }**

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column N.S. means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	19	18	28	36	44	11
N.S.	1	1.00	1.00	1.73	1.64	2.55	3.27	4.00	1.00
time (sec)	N/A	0.151	0.005	1.705	0.255	0.284	0.819	0.299	0.112

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	10	18	0	10	10
N.S.	1	1.00	1.00	1.10	1.00	1.80	0.00	1.00	1.00
time (sec)	N/A	0.162	0.007	0.297	0.241	0.257	0.000	0.270	12.778

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	34	36	46	61	0	48	36
N.S.	1	1.00	1.00	1.06	1.35	1.79	0.00	1.41	1.06
time (sec)	N/A	0.224	0.014	0.315	0.259	0.291	0.000	0.274	0.069



Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	23	24	22	31	0	22	21
N.S.	1	1.00	0.88	0.92	0.85	1.19	0.00	0.85	0.81
time (sec)	N/A	0.180	0.089	0.276	0.237	0.262	0.000	0.273	12.668

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	60	55	49	71	74	0	63	58
N.S.	1	1.09	1.00	0.89	1.29	1.35	0.00	1.15	1.05
time (sec)	N/A	0.308	0.017	0.371	0.266	0.297	0.000	0.276	0.103

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	38	35	34	34	41	0	34	31
N.S.	1	0.93	0.85	0.83	0.83	1.00	0.00	0.83	0.76
time (sec)	N/A	0.187	0.179	0.314	0.245	0.276	0.000	0.305	12.525

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	86	76	59	91	84	0	73	79
N.S.	1	1.13	1.00	0.78	1.20	1.11	0.00	0.96	1.04
time (sec)	N/A	0.437	0.018	0.729	0.256	0.277	0.000	0.304	0.130

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	48	43	44	44	51	0	44	39
N.S.	1	0.91	0.81	0.83	0.83	0.96	0.00	0.83	0.74
time (sec)	N/A	0.209	0.286	0.447	0.275	0.268	0.000	0.281	12.301

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	86	59	358	0	110	0	0	0
N.S.	1	1.01	0.69	4.21	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	0.408	0.209	8.778	0.000	0.099	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	46	213	0	88	0	0	0
N.S.	1	1.00	0.74	3.44	0.00	1.42	0.00	0.00	0.00
time (sec)	N/A	0.310	0.071	4.511	0.000	0.106	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	45	182	0	73	0	0	0
N.S.	1	1.00	0.78	3.14	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.304	0.041	6.000	0.000	0.092	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	36	133	0	51	0	0	33
N.S.	1	1.00	1.00	3.69	0.00	1.42	0.00	0.00	0.92
time (sec)	N/A	0.224	0.023	3.855	0.000	0.087	0.000	0.000	0.099

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	36	133	0	57	0	0	0
N.S.	1	1.00	1.00	3.69	0.00	1.58	0.00	0.00	0.00
time (sec)	N/A	0.223	0.029	4.807	0.000	0.092	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	49	179	0	68	0	0	0
N.S.	1	1.00	0.79	2.89	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	0.305	0.052	6.182	0.000	0.099	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	55	202	0	74	0	0	0
N.S.	1	1.00	0.89	3.26	0.00	1.19	0.00	0.00	0.00
time (sec)	N/A	0.296	0.073	5.997	0.000	0.098	0.000	0.000	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	90	61	199	0	87	0	0	0
N.S.	1	1.06	0.72	2.34	0.00	1.02	0.00	0.00	0.00
time (sec)	N/A	0.398	0.125	6.195	0.000	0.103	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	100	62	415	0	125	0	0	0
N.S.	1	1.02	0.63	4.23	0.00	1.28	0.00	0.00	0.00
time (sec)	N/A	0.439	0.241	13.581	0.000	0.104	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	51	144	0	101	0	0	0
N.S.	1	1.00	0.73	2.06	0.00	1.44	0.00	0.00	0.00
time (sec)	N/A	0.323	0.099	4.363	0.000	0.091	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	48	394	0	84	0	0	0
N.S.	1	1.00	0.73	5.97	0.00	1.27	0.00	0.00	0.00
time (sec)	N/A	0.321	0.056	2.917	0.000	0.110	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	77	0	57	0	0	35
N.S.	1	1.00	1.00	2.03	0.00	1.50	0.00	0.00	0.92
time (sec)	N/A	0.227	0.039	2.873	0.000	0.090	0.000	0.000	0.159

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	299	0	66	0	0	0
N.S.	1	1.00	1.00	7.87	0.00	1.74	0.00	0.00	0.00
time (sec)	N/A	0.226	0.049	2.959	0.000	0.120	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	59	144	0	87	0	0	0
N.S.	1	1.00	0.82	2.00	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.322	0.084	2.934	0.000	0.089	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	60	420	0	95	0	0	0
N.S.	1	1.00	0.83	5.83	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	0.322	0.098	4.153	0.000	0.096	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	108	66	160	0	100	0	0	0
N.S.	1	1.08	0.66	1.60	0.00	1.00	0.00	0.00	0.00
time (sec)	N/A	0.428	0.122	3.013	0.000	0.102	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	51	51	55	0	0	0	0	0	0
N.S.	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.239	0.068	0.000	0.000	0.000	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	51	51	55	0	0	0	0	0	0
N.S.	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.240	0.042	0.000	0.000	0.000	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.235	0.040	0.000	0.000	0.000	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.233	0.090	0.000	0.000	0.000	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	53	53	55	0	0	0	0	0	0
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.233	0.089	0.000	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	53	53	55	0	0	0	0	0	0
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.235	0.074	0.000	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	54	54	57	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.252	0.050	0.000	0.000	0.000	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	54	54	57	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.263	0.043	0.000	0.000	0.000	0.000	0.000	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	0
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.254	0.042	0.000	0.000	0.000	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	0
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.252	0.061	0.000	0.000	0.000	0.000	0.000	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	57	0	0	0	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.256	0.055	0.000	0.000	0.000	0.000	0.000	0.000



Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	57	0	0	0	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.263	0.075	0.000	0.000	0.000	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	70	70	61	0	0	0	0	0	0
N.S.	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.261	0.056	0.000	0.000	0.000	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	73	73	61	0	0	0	0	0	0
N.S.	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.278	0.045	0.000	0.000	0.000	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	<b>F(-1)</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	50	66	39	51	42	49	0	59	0
N.S.	1	1.32	0.78	1.02	0.84	0.98	0.00	1.18	0.00
time (sec)	N/A	0.205	0.096	1.970	0.304	0.263	0.000	0.276	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	36	45	33	43	30	43	0	53	0
N.S.	1	1.25	0.92	1.19	0.83	1.19	0.00	1.47	0.00
time (sec)	N/A	0.182	0.048	1.797	0.342	0.275	0.000	0.287	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	22	24	23	32	18	34	0	44	0
N.S.	1	1.09	1.05	1.45	0.82	1.55	0.00	2.00	0.00
time (sec)	N/A	0.173	0.021	0.250	0.361	0.282	0.000	0.291	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	C	A	B	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	3	3	14	14	3	17	0	35	0
N.S.	1	1.00	4.67	4.67	1.00	5.67	0.00	11.67	0.00
time (sec)	N/A	0.157	0.004	0.375	0.331	0.267	0.000	0.275	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	B
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	11	13	11	7	11	4	10	6	12
N.S.	1	1.18	1.00	0.64	1.00	0.36	0.91	0.55	1.09
time (sec)	N/A	0.172	0.006	0.358	0.233	0.260	0.191	0.273	12.448

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	29	33	28	18	25	10	27	16	0
N.S.	1	1.14	0.97	0.62	0.86	0.34	0.93	0.55	0.00
time (sec)	N/A	0.179	0.022	0.371	0.240	0.271	0.320	0.280	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	43	54	28	22	37	18	44	25	0
N.S.	1	1.26	0.65	0.51	0.86	0.42	1.02	0.58	0.00
time (sec)	N/A	0.184	0.037	0.315	0.264	0.279	1.276	0.282	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	57	75	34	28	49	24	60	34	0
N.S.	1	1.32	0.60	0.49	0.86	0.42	1.05	0.60	0.00
time (sec)	N/A	0.191	0.040	0.429	0.227	0.267	12.268	0.284	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F(-1)</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	99	44	61	2175	65	0	79	0
N.S.	1	1.18	0.52	0.73	25.89	0.77	0.00	0.94	0.00
time (sec)	N/A	0.215	0.107	2.041	2.450	0.273	0.000	0.282	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	75	37	53	1111	56	0	67	0
N.S.	1	1.15	0.57	0.82	17.09	0.86	0.00	1.03	0.00
time (sec)	N/A	0.209	0.055	1.988	0.526	0.284	0.000	0.280	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	51	24	40	324	39	0	42	0
N.S.	1	1.11	0.52	0.87	7.04	0.85	0.00	0.91	0.00
time (sec)	N/A	0.199	0.041	0.340	0.372	0.280	0.000	0.289	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	27	16	21	38	55	0	31	0
N.S.	1	1.08	0.64	0.84	1.52	2.20	0.00	1.24	0.00
time (sec)	N/A	0.190	0.007	0.354	0.395	0.280	0.000	0.271	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	15	13	12	6	16	12	11	15
N.S.	1	1.15	1.00	0.92	0.46	1.23	0.92	0.85	1.15
time (sec)	N/A	0.179	0.029	0.197	0.390	0.263	0.245	0.290	12.719

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	45	25	24	14	24	31	19	0
N.S.	1	1.25	0.69	0.67	0.39	0.67	0.86	0.53	0.00
time (sec)	N/A	0.193	0.031	0.092	0.377	0.254	0.368	0.305	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	74	33	30	22	32	49	27	0
N.S.	1	1.35	0.60	0.55	0.40	0.58	0.89	0.49	0.00
time (sec)	N/A	0.205	0.027	0.203	0.520	0.272	1.340	0.275	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	103	39	36	28	38	66	33	0
N.S.	1	1.39	0.53	0.49	0.38	0.51	0.89	0.45	0.00
time (sec)	N/A	0.213	0.031	0.140	0.484	0.259	12.648	0.273	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	103	59	297	0	102	0	0	0
N.S.	1	0.88	0.50	2.54	0.00	0.87	0.00	0.00	0.00
time (sec)	N/A	0.606	0.132	26.888	0.000	0.104	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	71	43	100	0	74	0	0	0
N.S.	1	1.09	0.66	1.54	0.00	1.14	0.00	0.00	0.00
time (sec)	N/A	0.414	0.063	4.270	0.000	0.091	0.000	0.000	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	49	32	244	0	57	0	0	0
N.S.	1	1.17	0.76	5.81	0.00	1.36	0.00	0.00	0.00
time (sec)	N/A	0.331	0.035	1.755	0.000	0.093	0.000	0.000	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	53	31	92	0	58	0	0	0
N.S.	1	1.20	0.70	2.09	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	0.335	0.065	3.762	0.000	0.089	0.000	0.000	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	73	73	43	273	0	73	0	0	0
N.S.	1	1.00	0.59	3.74	0.00	1.00	0.00	0.00	0.00
time (sec)	N/A	0.418	0.132	5.079	0.000	0.098	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	107	59	117	0	79	0	0	0
N.S.	1	0.91	0.50	1.00	0.00	0.68	0.00	0.00	0.00
time (sec)	N/A	0.586	0.126	5.570	0.000	0.102	0.000	0.000	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	68	54	58	61	76	0	67	589
N.S.	1	0.42	0.33	0.36	0.37	0.47	0.00	0.41	3.61
time (sec)	N/A	0.266	0.270	97.124	0.428	0.289	0.000	0.287	17.504

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	56	42	46	43	58	0	49	119
N.S.	1	0.48	0.36	0.39	0.37	0.50	0.00	0.42	1.02
time (sec)	N/A	0.257	0.145	95.848	0.406	0.275	0.000	0.297	15.129

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	38	33	31	25	34	0	22	36
N.S.	1	0.62	0.54	0.51	0.41	0.56	0.00	0.36	0.59
time (sec)	N/A	0.252	0.044	0.384	0.453	0.274	0.000	0.284	0.556

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	6	13	0	6	6
N.S.	1	1.00	1.00	0.93	0.40	0.87	0.00	0.40	0.40
time (sec)	N/A	0.228	0.008	0.369	0.480	0.271	0.000	0.270	0.124

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	29	23	20	25	27	0	39	0
N.S.	1	0.81	0.64	0.56	0.69	0.75	0.00	1.08	0.00
time (sec)	N/A	0.219	0.045	0.229	0.500	0.284	0.000	0.280	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	62	38	40	58	43	0	0	0
N.S.	1	0.72	0.44	0.47	0.67	0.50	0.00	0.00	0.00
time (sec)	N/A	0.336	0.065	0.184	0.647	0.279	0.000	0.000	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	92	55	56	88	55	0	0	0
N.S.	1	0.70	0.42	0.42	0.67	0.42	0.00	0.00	0.00
time (sec)	N/A	0.470	0.121	0.497	0.338	0.298	0.000	0.000	0.000



Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	81	81	69	0	0	0	0	0	0
N.S.	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.364	0.108	0.000	0.000	0.000	0.000	0.000	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	83	83	71	0	0	0	0	0	0
N.S.	1	1.00	0.86	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.340	0.080	0.000	0.000	0.000	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	108	69	158	0	114	0	0	0
N.S.	1	1.11	0.71	1.63	0.00	1.18	0.00	0.00	0.00
time (sec)	N/A	0.459	0.404	14.252	0.000	0.095	0.000	0.000	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	104	69	412	0	120	0	0	0
N.S.	1	1.09	0.73	4.34	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.465	0.279	3.132	0.000	0.101	0.000	0.000	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	74	51	141	0	98	0	0	0
N.S.	1	1.07	0.74	2.04	0.00	1.42	0.00	0.00	0.00
time (sec)	N/A	0.355	0.149	1.761	0.000	0.094	0.000	0.000	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	70	47	393	0	83	0	0	0
N.S.	1	1.11	0.75	6.24	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	0.350	0.071	2.800	0.000	0.088	0.000	0.000	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	77	0	57	0	0	35
N.S.	1	1.00	1.00	2.03	0.00	1.50	0.00	0.00	0.92
time (sec)	N/A	0.230	0.037	2.829	0.000	0.088	0.000	0.000	12.989

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	39	162	0	63	0	0	0
N.S.	1	1.00	1.00	4.15	0.00	1.62	0.00	0.00	0.00
time (sec)	N/A	0.290	0.050	3.031	0.000	0.102	0.000	0.000	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	76	51	147	0	84	0	0	0
N.S.	1	1.13	0.76	2.19	0.00	1.25	0.00	0.00	0.00
time (sec)	N/A	0.366	0.239	2.945	0.000	0.092	0.000	0.000	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	76	57	172	0	92	0	0	0
N.S.	1	1.09	0.81	2.46	0.00	1.31	0.00	0.00	0.00
time (sec)	N/A	0.368	0.188	4.161	0.000	0.093	0.000	0.000	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	112	63	163	0	97	0	0	0
N.S.	1	1.18	0.66	1.72	0.00	1.02	0.00	0.00	0.00
time (sec)	N/A	0.476	0.219	3.165	0.000	0.110	0.000	0.000	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	112	71	184	0	105	0	0	0
N.S.	1	1.14	0.72	1.88	0.00	1.07	0.00	0.00	0.00
time (sec)	N/A	0.484	0.399	4.659	0.000	0.105	0.000	0.000	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	108	64	159	0	117	0	0	0
N.S.	1	1.14	0.67	1.67	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.467	0.365	0.518	0.000	0.094	0.000	0.000	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	104	64	413	0	121	0	0	0
N.S.	1	1.06	0.65	4.21	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.457	0.292	0.753	0.000	0.100	0.000	0.000	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	74	49	140	0	99	0	0	0
N.S.	1	1.10	0.73	2.09	0.00	1.48	0.00	0.00	0.00
time (sec)	N/A	0.347	0.094	0.558	0.000	0.090	0.000	0.000	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	48	394	0	84	0	0	0
N.S.	1	1.00	0.73	5.97	0.00	1.27	0.00	0.00	0.00
time (sec)	N/A	0.316	0.011	0.625	0.000	0.102	0.000	0.000	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	39	78	0	57	0	0	0
N.S.	1	1.00	1.00	2.00	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.279	0.009	0.383	0.000	0.098	0.000	0.000	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	161	0	63	0	0	0
N.S.	1	1.00	1.00	3.93	0.00	1.54	0.00	0.00	0.00
time (sec)	N/A	0.266	0.018	0.529	0.000	0.087	0.000	0.000	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	76	52	147	0	85	0	0	0
N.S.	1	1.09	0.74	2.10	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.369	0.168	0.562	0.000	0.091	0.000	0.000	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	76	58	175	0	93	0	0	0
N.S.	1	1.06	0.81	2.43	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	0.369	0.171	0.603	0.000	0.106	0.000	0.000	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	112	64	164	0	99	0	0	0
N.S.	1	1.14	0.65	1.67	0.00	1.01	0.00	0.00	0.00
time (sec)	N/A	0.482	0.252	0.619	0.000	0.101	0.000	0.000	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	112	72	185	0	107	0	0	0
N.S.	1	1.12	0.72	1.85	0.00	1.07	0.00	0.00	0.00
time (sec)	N/A	0.493	0.468	0.709	0.000	0.105	0.000	0.000	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	108	61	161	0	121	0	0	0
N.S.	1	1.10	0.62	1.64	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.453	0.370	4.767	0.000	0.094	0.000	0.000	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	104	61	415	0	125	0	0	0
N.S.	1	1.07	0.63	4.28	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	0.459	0.230	1.643	0.000	0.100	0.000	0.000	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	51	144	0	101	0	0	0
N.S.	1	1.00	0.73	2.06	0.00	1.44	0.00	0.00	0.00
time (sec)	N/A	0.323	0.012	0.992	0.000	0.089	0.000	0.000	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	50	396	0	86	0	0	0
N.S.	1	1.00	0.74	5.82	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.366	0.017	6.803	0.000	0.091	0.000	0.000	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	80	0	57	0	0	0
N.S.	1	1.00	1.00	1.95	0.00	1.39	0.00	0.00	0.00
time (sec)	N/A	0.269	0.017	31.782	0.000	0.092	0.000	0.000	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	38	165	0	63	0	0	0
N.S.	1	1.00	0.93	4.02	0.00	1.54	0.00	0.00	0.00
time (sec)	N/A	0.313	0.188	123.017	0.000	0.089	0.000	0.000	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	76	54	150	0	87	0	0	0
N.S.	1	1.06	0.75	2.08	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.371	0.194	4.258	0.000	0.091	0.000	0.000	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	76	60	175	0	95	0	0	0
N.S.	1	1.06	0.83	2.43	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	0.366	0.178	4.253	0.000	0.093	0.000	0.000	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	112	66	166	0	103	0	0	0
N.S.	1	1.12	0.66	1.66	0.00	1.03	0.00	0.00	0.00
time (sec)	N/A	0.482	0.302	2.893	0.000	0.092	0.000	0.000	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	112	74	187	0	111	0	0	0
N.S.	1	1.12	0.74	1.87	0.00	1.11	0.00	0.00	0.00
time (sec)	N/A	0.485	0.509	4.464	0.000	0.105	0.000	0.000	0.000



Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	100	62	415	0	125	0	0	0
N.S.	1	1.02	0.63	4.23	0.00	1.28	0.00	0.00	0.00
time (sec)	N/A	0.427	0.056	1.113	0.000	0.103	0.000	0.000	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	108	69	163	0	117	0	0	0
N.S.	1	1.08	0.69	1.63	0.00	1.17	0.00	0.00	0.00
time (sec)	N/A	0.459	0.290	0.739	0.000	0.093	0.000	0.000	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	104	61	417	0	123	0	0	0
N.S.	1	1.07	0.63	4.30	0.00	1.27	0.00	0.00	0.00
time (sec)	N/A	0.468	0.346	0.781	0.000	0.098	0.000	0.000	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	74	51	147	0	101	0	0	0
N.S.	1	1.03	0.71	2.04	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	0.352	0.189	0.503	0.000	0.084	0.000	0.000	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	70	48	389	0	86	0	0	0
N.S.	1	1.08	0.74	5.98	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	0.344	0.198	0.621	0.000	0.104	0.000	0.000	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	69	0	60	0	0	0
N.S.	1	1.00	1.00	1.68	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.251	0.018	0.538	0.000	0.088	0.000	0.000	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	299	0	66	0	0	0
N.S.	1	1.00	1.00	7.87	0.00	1.74	0.00	0.00	0.00
time (sec)	N/A	0.225	0.020	0.597	0.000	0.090	0.000	0.000	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	74	60	141	0	87	0	0	0
N.S.	1	1.07	0.87	2.04	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.354	0.086	0.606	0.000	0.095	0.000	0.000	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	76	60	417	0	95	0	0	0
N.S.	1	1.13	0.90	6.22	0.00	1.42	0.00	0.00	0.00
time (sec)	N/A	0.369	0.122	0.620	0.000	0.106	0.000	0.000	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	112	66	157	0	100	0	0	0
N.S.	1	1.15	0.68	1.62	0.00	1.03	0.00	0.00	0.00
time (sec)	N/A	0.472	0.247	0.652	0.000	0.098	0.000	0.000	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	112	70	451	0	108	0	0	0
N.S.	1	1.18	0.74	4.75	0.00	1.14	0.00	0.00	0.00
time (sec)	N/A	0.475	0.345	0.582	0.000	0.107	0.000	0.000	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	108	69	166	0	117	0	0	0
N.S.	1	1.08	0.69	1.66	0.00	1.17	0.00	0.00	0.00
time (sec)	N/A	0.454	0.402	0.677	0.000	0.097	0.000	0.000	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	104	64	420	0	123	0	0	0
N.S.	1	1.04	0.64	4.20	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.451	0.226	0.564	0.000	0.093	0.000	0.000	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	74	56	150	0	101	0	0	0
N.S.	1	1.03	0.78	2.08	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	0.344	0.227	0.512	0.000	0.103	0.000	0.000	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	70	51	392	0	86	0	0	0
N.S.	1	1.03	0.75	5.76	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.340	0.217	0.608	0.000	0.096	0.000	0.000	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	72	0	60	0	0	0
N.S.	1	1.00	1.00	1.76	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.251	0.011	0.592	0.000	0.094	0.000	0.000	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	305	0	66	0	0	0
N.S.	1	1.00	1.00	7.44	0.00	1.61	0.00	0.00	0.00
time (sec)	N/A	0.247	0.022	0.309	0.000	0.093	0.000	0.000	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	59	144	0	87	0	0	0
N.S.	1	1.00	0.82	2.00	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.323	0.044	0.524	0.000	0.096	0.000	0.000	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	74	60	420	0	95	0	0	0
N.S.	1	1.07	0.87	6.09	0.00	1.38	0.00	0.00	0.00
time (sec)	N/A	0.356	0.067	0.550	0.000	0.096	0.000	0.000	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	112	66	160	0	100	0	0	0
N.S.	1	1.14	0.67	1.63	0.00	1.02	0.00	0.00	0.00
time (sec)	N/A	0.481	0.140	0.600	0.000	0.096	0.000	0.000	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	112	73	454	0	108	0	0	0
N.S.	1	1.15	0.75	4.68	0.00	1.11	0.00	0.00	0.00
time (sec)	N/A	0.482	0.324	0.598	0.000	0.124	0.000	0.000	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	108	64	166	0	117	0	0	0
N.S.	1	1.08	0.64	1.66	0.00	1.17	0.00	0.00	0.00
time (sec)	N/A	0.452	0.473	0.627	0.000	0.105	0.000	0.000	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	104	64	420	0	123	0	0	0
N.S.	1	1.04	0.64	4.20	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.452	0.354	0.623	0.000	0.101	0.000	0.000	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	74	51	149	0	101	0	0	0
N.S.	1	1.03	0.71	2.07	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	0.347	0.214	0.566	0.000	0.091	0.000	0.000	0.000

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	70	51	392	0	86	0	0	0
N.S.	1	1.03	0.75	5.76	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.350	0.245	0.575	0.000	0.089	0.000	0.000	0.000

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	72	0	60	0	0	0
N.S.	1	1.00	1.00	1.76	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.260	0.015	0.626	0.000	0.086	0.000	0.000	0.000

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	38	305	0	66	0	0	0
N.S.	1	1.00	0.93	7.44	0.00	1.61	0.00	0.00	0.00
time (sec)	N/A	0.255	0.213	0.291	0.000	0.091	0.000	0.000	0.000

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	76	62	144	0	87	0	0	0
N.S.	1	1.06	0.86	2.00	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.353	0.033	0.523	0.000	0.096	0.000	0.000	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	60	420	0	95	0	0	0
N.S.	1	1.00	0.83	5.83	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	0.322	0.040	0.537	0.000	0.099	0.000	0.000	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	110	66	160	0	100	0	0	0
N.S.	1	1.13	0.68	1.65	0.00	1.03	0.00	0.00	0.00
time (sec)	N/A	0.474	0.037	0.573	0.000	0.099	0.000	0.000	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	112	73	454	0	108	0	0	0
N.S.	1	1.14	0.74	4.63	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	0.475	0.172	0.564	0.000	0.112	0.000	0.000	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	108	66	160	0	100	0	0	0
N.S.	1	1.08	0.66	1.60	0.00	1.00	0.00	0.00	0.00
time (sec)	N/A	0.427	0.016	0.500	0.000	0.105	0.000	0.000	0.000



Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	83	64	104	1656	229	0	0	0
N.S.	1	0.78	0.60	0.97	15.48	2.14	0.00	0.00	0.00
time (sec)	N/A	0.372	0.172	1.022	0.483	0.320	0.000	0.000	0.000

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	48	45	48	294	43	0	0	126
N.S.	1	0.69	0.64	0.69	4.20	0.61	0.00	0.00	1.80
time (sec)	N/A	0.231	0.101	0.662	0.398	0.269	0.000	0.000	15.010

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	57	50	92	661	199	0	0	0
N.S.	1	0.79	0.69	1.28	9.18	2.76	0.00	0.00	0.00
time (sec)	N/A	0.282	0.061	0.574	0.461	0.298	0.000	0.000	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	35	54	30	0	0	46
N.S.	1	1.00	1.00	1.09	1.69	0.94	0.00	0.00	1.44
time (sec)	N/A	0.215	0.022	0.661	0.439	0.272	0.000	0.000	0.295

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	46	65	111	0	0	0
N.S.	1	1.00	1.00	1.39	1.97	3.36	0.00	0.00	0.00
time (sec)	N/A	0.208	0.016	0.579	0.378	0.296	0.000	0.000	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	28	26	98	22	0	24
N.S.	1	1.00	1.00	1.17	1.08	4.08	0.92	0.00	1.00
time (sec)	N/A	0.149	0.038	0.537	0.319	0.295	1.805	0.000	12.522

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	13	30	46	38	32
N.S.	1	1.00	1.00	0.91	0.41	0.94	1.44	1.19	1.00
time (sec)	N/A	0.215	0.212	0.625	0.387	0.268	3.653	0.857	0.275

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	48	45	45	25	158	107	140	41
N.S.	1	0.76	0.71	0.71	0.40	2.51	1.70	2.22	0.65
time (sec)	N/A	0.207	0.195	0.657	0.445	0.315	9.813	1.323	12.696

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	48	45	47	42	48	0	0	45
N.S.	1	0.69	0.64	0.67	0.60	0.69	0.00	0.00	0.64
time (sec)	N/A	0.223	0.132	0.731	0.406	0.266	0.000	0.000	12.834

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	74	55	64	49	202	0	0	52
N.S.	1	0.76	0.56	0.65	0.50	2.06	0.00	0.00	0.53
time (sec)	N/A	0.275	0.313	0.716	0.429	0.314	0.000	0.000	12.775

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	84	64	105	1742	236	0	0	0
N.S.	1	0.76	0.58	0.95	15.84	2.15	0.00	0.00	0.00
time (sec)	N/A	0.375	0.153	0.639	0.487	0.307	0.000	0.000	0.000

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	49	45	49	299	44	0	0	127
N.S.	1	0.68	0.62	0.68	4.15	0.61	0.00	0.00	1.76
time (sec)	N/A	0.229	0.104	0.533	0.398	0.282	0.000	0.000	13.840

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	58	50	93	691	202	0	0	0
N.S.	1	0.78	0.68	1.26	9.34	2.73	0.00	0.00	0.00
time (sec)	N/A	0.285	0.077	0.541	0.435	0.295	0.000	0.000	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	32	36	54	31	0	0	47
N.S.	1	1.00	0.97	1.09	1.64	0.94	0.00	0.00	1.42
time (sec)	N/A	0.220	0.025	0.588	0.393	0.267	0.000	0.000	0.216

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	33	47	68	112	0	0	0
N.S.	1	1.00	0.97	1.38	2.00	3.29	0.00	0.00	0.00
time (sec)	N/A	0.202	0.024	0.547	0.462	0.295	0.000	0.000	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	29	26	99	22	0	25
N.S.	1	1.00	1.00	1.16	1.04	3.96	0.88	0.00	1.00
time (sec)	N/A	0.152	0.051	0.539	0.367	0.304	3.464	0.000	0.097

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	32	30	13	31	46	38	33
N.S.	1	1.00	0.97	0.91	0.39	0.94	1.39	1.15	1.00
time (sec)	N/A	0.193	0.201	0.561	0.406	0.275	18.084	0.868	0.242

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	49	45	46	28	161	0	0	42
N.S.	1	0.75	0.69	0.71	0.43	2.48	0.00	0.00	0.65
time (sec)	N/A	0.208	0.238	0.535	0.410	0.311	0.000	0.000	0.370

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	49	45	48	45	51	0	0	46
N.S.	1	0.68	0.62	0.67	0.62	0.71	0.00	0.00	0.64
time (sec)	N/A	0.226	0.151	0.607	0.412	0.269	0.000	0.000	13.279

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	75	55	65	53	207	0	0	53
N.S.	1	0.74	0.54	0.64	0.52	2.05	0.00	0.00	0.52
time (sec)	N/A	0.283	0.480	0.444	0.401	0.327	0.000	0.000	0.521

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	63	57	61	705	63	0	0	205
N.S.	1	0.54	0.49	0.53	6.08	0.54	0.00	0.00	1.77
time (sec)	N/A	0.233	0.238	24.605	0.474	0.279	0.000	0.000	18.012

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	51	45	51	311	48	0	0	129
N.S.	1	0.67	0.59	0.67	4.09	0.63	0.00	0.00	1.70
time (sec)	N/A	0.235	0.120	0.570	0.440	0.276	0.000	0.000	1.295

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	60	50	95	747	208	0	0	0
N.S.	1	0.77	0.64	1.22	9.58	2.67	0.00	0.00	0.00
time (sec)	N/A	0.286	0.094	0.616	0.419	0.316	0.000	0.000	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	38	54	33	0	0	66
N.S.	1	1.00	0.91	1.09	1.54	0.94	0.00	0.00	1.89
time (sec)	N/A	0.220	0.037	0.539	0.416	0.273	0.000	0.000	13.538

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	33	49	72	114	0	0	0
N.S.	1	1.00	0.92	1.36	2.00	3.17	0.00	0.00	0.00
time (sec)	N/A	0.208	0.041	0.520	0.418	0.297	0.000	0.000	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	24	31	26	101	22	0	27
N.S.	1	1.00	0.89	1.15	0.96	3.74	0.81	0.00	1.00
time (sec)	N/A	0.152	0.060	0.553	0.390	0.296	47.272	0.000	0.111

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	32	13	33	0	0	35
N.S.	1	1.00	0.91	0.91	0.37	0.94	0.00	0.00	1.00
time (sec)	N/A	0.198	0.291	0.540	0.411	0.272	0.000	0.000	13.795

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	51	45	48	32	167	0	0	44
N.S.	1	0.74	0.65	0.70	0.46	2.42	0.00	0.00	0.64
time (sec)	N/A	0.215	0.334	0.539	0.437	0.307	0.000	0.000	0.353

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	51	45	50	49	55	0	0	48
N.S.	1	0.67	0.59	0.66	0.64	0.72	0.00	0.00	0.63
time (sec)	N/A	0.232	0.209	0.385	0.438	0.273	0.000	0.000	13.886

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	57	50	92	661	205	0	0	0
N.S.	1	0.79	0.69	1.28	9.18	2.85	0.00	0.00	0.00
time (sec)	N/A	0.282	0.065	0.619	0.410	0.312	0.000	0.000	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	35	59	33	0	0	51
N.S.	1	1.00	1.00	1.09	1.84	1.03	0.00	0.00	1.59
time (sec)	N/A	0.216	0.027	0.587	0.416	0.262	0.000	0.000	0.270

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	46	65	114	0	0	0
N.S.	1	1.00	1.00	1.39	1.97	3.45	0.00	0.00	0.00
time (sec)	N/A	0.205	0.017	0.555	0.403	0.297	0.000	0.000	0.000



Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	28	26	101	22	0	27
N.S.	1	1.00	1.00	1.17	1.08	4.21	0.92	0.00	1.12
time (sec)	N/A	0.145	0.037	0.523	0.332	0.295	3.446	0.000	0.250

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	13	33	46	0	35
N.S.	1	1.00	1.00	0.91	0.41	1.03	1.44	0.00	1.09
time (sec)	N/A	0.202	0.150	0.521	0.383	0.261	6.527	0.000	13.733

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	48	45	45	25	165	107	0	44
N.S.	1	0.76	0.71	0.71	0.40	2.62	1.70	0.00	0.70
time (sec)	N/A	0.209	0.158	0.515	0.420	0.297	7.532	0.000	13.911

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	48	45	47	42	51	82	0	48
N.S.	1	0.69	0.64	0.67	0.60	0.73	1.17	0.00	0.69
time (sec)	N/A	0.220	0.094	0.522	0.405	0.281	23.975	0.000	13.914

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	60	50	95	670	205	0	0	0
N.S.	1	0.77	0.64	1.22	8.59	2.63	0.00	0.00	0.00
time (sec)	N/A	0.285	0.079	0.584	0.418	0.295	0.000	0.000	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	38	67	33	0	0	51
N.S.	1	1.00	0.91	1.09	1.91	0.94	0.00	0.00	1.46
time (sec)	N/A	0.213	0.042	0.522	0.404	0.292	0.000	0.000	0.267

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	33	49	65	114	0	0	0
N.S.	1	1.00	0.92	1.36	1.81	3.17	0.00	0.00	0.00
time (sec)	N/A	0.210	0.030	0.574	0.397	0.299	0.000	0.000	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	27	31	26	101	22	0	27
N.S.	1	1.00	1.00	1.15	0.96	3.74	0.81	0.00	1.00
time (sec)	N/A	0.148	0.045	0.592	0.336	0.300	5.178	0.000	0.258

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	32	13	33	46	0	39
N.S.	1	1.00	0.91	0.91	0.37	0.94	1.31	0.00	1.11
time (sec)	N/A	0.194	0.219	0.603	0.403	0.266	3.928	0.000	13.283

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	51	45	48	25	165	107	0	44
N.S.	1	0.74	0.65	0.70	0.36	2.39	1.55	0.00	0.64
time (sec)	N/A	0.206	0.185	0.566	0.422	0.298	8.198	0.000	13.178

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	51	45	50	42	51	82	0	48
N.S.	1	0.67	0.59	0.66	0.55	0.67	1.08	0.00	0.63
time (sec)	N/A	0.223	0.106	0.659	0.411	0.272	14.022	0.000	13.138

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	77	55	67	49	208	184	0	55
N.S.	1	0.72	0.51	0.63	0.46	1.94	1.72	0.00	0.51
time (sec)	N/A	0.285	0.267	0.543	0.427	0.316	105.669	0.000	13.575

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	60	53	95	688	205	0	0	0
N.S.	1	0.77	0.68	1.22	8.82	2.63	0.00	0.00	0.00
time (sec)	N/A	0.292	0.053	0.336	0.429	0.305	0.000	0.000	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	38	67	33	0	0	51
N.S.	1	1.00	0.91	1.09	1.91	0.94	0.00	0.00	1.46
time (sec)	N/A	0.219	0.038	0.526	0.403	0.273	0.000	0.000	0.263

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	33	49	65	114	0	0	0
N.S.	1	1.00	0.92	1.36	1.81	3.17	0.00	0.00	0.00
time (sec)	N/A	0.210	0.028	0.546	0.404	0.315	0.000	0.000	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	27	31	26	101	22	0	27
N.S.	1	1.00	1.00	1.15	0.96	3.74	0.81	0.00	1.00
time (sec)	N/A	0.151	0.040	0.535	0.377	0.294	49.360	0.000	13.200

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	35	32	13	33	46	0	39
N.S.	1	1.00	1.00	0.91	0.37	0.94	1.31	0.00	1.11
time (sec)	N/A	0.197	0.257	0.530	0.412	0.282	18.693	0.000	13.237

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	51	48	48	25	165	107	0	64
N.S.	1	0.74	0.70	0.70	0.36	2.39	1.55	0.00	0.93
time (sec)	N/A	0.207	0.165	0.593	0.556	0.301	10.097	0.000	12.993

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	51	48	50	42	51	82	0	48
N.S.	1	0.67	0.63	0.66	0.55	0.67	1.08	0.00	0.63
time (sec)	N/A	0.225	0.065	0.638	0.536	0.274	24.402	0.000	13.784

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	77	58	67	49	208	184	0	55
N.S.	1	0.72	0.54	0.63	0.46	1.94	1.72	0.00	0.51
time (sec)	N/A	0.289	0.216	0.624	0.541	0.337	112.023	0.000	13.358

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.292	0.078	0.000	0.000	0.000	0.000	0.000	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	53	53	60	0	0	0	0	0	0
N.S.	1	1.00	1.13	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.279	0.054	0.000	0.000	0.000	0.000	0.000	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	0
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.263	0.044	0.000	0.000	0.000	0.000	0.000	0.000

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.318	0.069	0.000	0.000	0.000	0.000	0.000	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	59	0	0	0	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.336	0.165	0.000	0.000	0.000	0.000	0.000	0.000

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.300	0.084	0.000	0.000	0.000	0.000	0.000	0.000

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	55	55	60	0	0	0	0	0	0
N.S.	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.271	0.052	0.000	0.000	0.000	0.000	0.000	0.000

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	54	54	57	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.257	0.001	0.000	0.000	0.000	0.000	0.000	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	56	0	0	0	0	0	0
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.299	0.003	0.000	0.000	0.000	0.000	0.000	0.000

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.289	0.008	0.000	0.000	0.000	0.000	0.000	0.000

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.283	0.062	0.000	0.000	0.000	0.000	0.000	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	53	53	60	0	0	0	0	0	0
N.S.	1	1.00	1.13	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.275	0.049	0.000	0.000	0.000	0.000	0.000	0.000



Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	0
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.260	0.057	0.000	0.000	0.000	0.000	0.000	0.000

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.340	0.089	0.000	0.000	0.000	0.000	0.000	0.000

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.364	0.076	0.000	0.000	0.000	0.000	0.000	0.000

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	60	0	0	0	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.277	0.003	0.000	0.000	0.000	0.000	0.000	0.000

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	55	55	58	0	0	0	0	0	0
N.S.	1	1.00	1.05	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.274	0.012	0.000	0.000	0.000	0.000	0.000	0.000

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	57	0	0	0	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.260	0.001	0.000	0.000	0.000	0.000	0.000	0.000

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.289	0.015	0.000	0.000	0.000	0.000	0.000	0.000

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.335	0.101	0.000	0.000	0.000	0.000	0.000	0.000

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	81	81	83	0	0	0	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.322	0.105	0.000	0.000	0.000	0.000	0.000	0.000

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.332	0.145	0.000	0.000	0.000	0.000	0.000	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.330	0.141	0.000	0.000	0.000	0.000	0.000	0.000

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.332	0.134	0.000	0.000	0.000	0.000	0.000	0.000

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.337	0.135	0.000	0.000	0.000	0.000	0.000	0.000

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	85	85	83	0	0	0	0	0	0
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.347	0.183	0.000	0.000	0.000	0.000	0.000	0.000

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	89	89	76	0	0	0	0	0	0
N.S.	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.334	0.074	0.000	0.000	0.000	0.000	0.000	0.000

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	72	72	71	0	0	0	0	0	0
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.317	0.049	0.000	0.000	0.000	0.000	0.000	0.000

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	61	61	65	0	0	0	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.292	0.041	0.000	0.000	0.000	0.000	0.000	0.000

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	73	73	61	0	0	0	0	0	0
N.S.	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.291	0.047	0.000	0.000	0.000	0.000	0.000	0.000

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	75	75	68	0	0	0	0	0	0
N.S.	1	1.00	0.91	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.328	0.101	0.000	0.000	0.000	0.000	0.000	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	75	75	71	0	0	0	0	0	0
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.337	0.078	0.000	0.000	0.000	0.000	0.000	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	75	75	73	0	0	0	0	0	0
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.334	0.108	0.000	0.000	0.000	0.000	0.000	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.337	0.104	0.000	0.000	0.000	0.000	0.000	0.000

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.325	0.093	0.000	0.000	0.000	0.000	0.000	0.000

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.326	0.102	0.000	0.000	0.000	0.000	0.000	0.000

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.327	0.118	0.000	0.000	0.000	0.000	0.000	0.000

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.333	0.141	0.000	0.000	0.000	0.000	0.000	0.000

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.336	0.177	0.000	0.000	0.000	0.000	0.000	0.000

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	23	28	0	33	77
N.S.	1	1.00	1.00	0.85	1.15	1.40	0.00	1.65	3.85
time (sec)	N/A	0.200	0.276	3.168	0.206	0.275	0.000	0.281	14.501

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	23	28	0	33	39
N.S.	1	1.00	1.00	0.85	1.15	1.40	0.00	1.65	1.95
time (sec)	N/A	0.201	0.215	3.370	0.235	0.282	0.000	0.278	0.238

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	23	18	0	25	18
N.S.	1	1.00	1.00	0.94	1.28	1.00	0.00	1.39	1.00
time (sec)	N/A	0.196	0.191	0.399	0.235	0.258	0.000	0.281	0.093

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	23	23	0	22	23
N.S.	1	1.00	1.00	0.94	1.28	1.28	0.00	1.22	1.28
time (sec)	N/A	0.191	0.207	0.445	0.217	0.256	0.000	0.272	13.069

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	23	28	0	33	28
N.S.	1	1.00	1.00	0.85	1.15	1.40	0.00	1.65	1.40
time (sec)	N/A	0.196	0.179	0.471	0.347	0.268	0.000	0.280	13.118



Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	40	32	319	36	42	0	49	50
N.S.	1	0.98	0.78	7.78	0.88	1.02	0.00	1.20	1.22
time (sec)	N/A	0.238	0.542	67.426	0.250	0.256	0.000	0.308	13.462

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	42	42	35	38	44	0	49	95
N.S.	1	0.98	0.98	0.81	0.88	1.02	0.00	1.14	2.21
time (sec)	N/A	0.237	0.526	60.719	0.301	0.258	0.000	0.289	17.166

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	134	122	152	0	177	0	0	0
N.S.	1	1.05	0.95	1.19	0.00	1.38	0.00	0.00	0.00
time (sec)	N/A	0.734	18.609	7.016	0.000	0.107	0.000	0.000	0.000

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	56	53	0	67	0	0	85
N.S.	1	1.00	0.81	0.77	0.00	0.97	0.00	0.00	1.23
time (sec)	N/A	0.343	0.900	1.241	0.000	0.282	0.000	0.000	14.058

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	93	93	109	132	0	119	0	0	0
N.S.	1	1.00	1.17	1.42	0.00	1.28	0.00	0.00	0.00
time (sec)	N/A	0.500	1.501	1.071	0.000	0.097	0.000	0.000	0.000

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	33	0	36	0	0	36
N.S.	1	1.00	1.00	1.06	0.00	1.16	0.00	0.00	1.16
time (sec)	N/A	0.204	0.615	1.028	0.000	0.279	0.000	0.000	13.018

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	68	113	0	58	0	0	0
N.S.	1	1.00	1.28	2.13	0.00	1.09	0.00	0.00	0.00
time (sec)	N/A	0.362	1.261	1.092	0.000	0.099	0.000	0.000	0.000

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	270	171	121	361	0	1135	0	0	0
N.S.	1	0.63	0.45	1.34	0.00	4.20	0.00	0.00	0.00
time (sec)	N/A	0.469	2.058	36.147	0.000	0.514	0.000	0.000	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	93	93	80	239	0	0	0	0	0
N.S.	1	1.00	0.86	2.57	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.491	1.388	2.197	0.000	0.000	0.000	0.000	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	322	212	157	464	0	1290	0	0	0
N.S.	1	0.66	0.49	1.44	0.00	4.01	0.00	0.00	0.00
time (sec)	N/A	0.575	2.326	7.354	0.000	0.617	0.000	0.000	0.000

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	110	57	60	0	85	0	0	110
N.S.	1	1.06	0.55	0.58	0.00	0.82	0.00	0.00	1.06
time (sec)	N/A	0.452	1.111	1.508	0.000	0.302	0.000	0.000	14.548

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	167	114	232	0	264	0	0	0
N.S.	1	1.01	0.69	1.40	0.00	1.59	0.00	0.00	0.00
time (sec)	N/A	0.859	11.950	1.153	0.000	0.114	0.000	0.000	0.000

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	45	48	0	58	0	0	61
N.S.	1	1.00	0.65	0.70	0.00	0.84	0.00	0.00	0.88
time (sec)	N/A	0.326	0.750	1.017	0.000	0.282	0.000	0.000	12.933

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	126	99	354	0	158	0	0	0
N.S.	1	1.01	0.79	2.83	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.665	1.063	1.165	0.000	0.109	0.000	0.000	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	33	0	36	0	0	36
N.S.	1	1.00	1.00	1.06	0.00	1.16	0.00	0.00	1.16
time (sec)	N/A	0.207	0.566	1.152	0.000	0.265	0.000	0.000	12.519

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	66	355	0	155	0	0	0
N.S.	1	1.00	0.74	3.99	0.00	1.74	0.00	0.00	0.00
time (sec)	N/A	0.510	0.851	1.222	0.000	0.103	0.000	0.000	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	327	211	142	685	0	1310	0	0	0
N.S.	1	0.65	0.43	2.09	0.00	4.01	0.00	0.00	0.00
time (sec)	N/A	0.603	2.530	6.831	0.000	0.608	0.000	0.000	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	94	69	396	0	0	0	0	0
N.S.	1	1.00	0.73	4.21	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.513	1.242	1.061	0.000	0.000	0.000	0.000	0.000

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	173	92	164	0	225	0	0	0
N.S.	1	1.04	0.55	0.99	0.00	1.36	0.00	0.00	0.00
time (sec)	N/A	0.832	2.388	1.877	0.000	0.113	0.000	0.000	0.000

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	110	57	68	0	89	0	0	112
N.S.	1	1.04	0.54	0.64	0.00	0.84	0.00	0.00	1.06
time (sec)	N/A	0.453	0.843	1.162	0.000	0.299	0.000	0.000	15.303

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	132	87	150	0	160	0	0	0
N.S.	1	1.01	0.66	1.15	0.00	1.22	0.00	0.00	0.00
time (sec)	N/A	0.670	1.368	1.124	0.000	0.099	0.000	0.000	0.000

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	45	47	0	58	0	0	64
N.S.	1	1.00	0.65	0.68	0.00	0.84	0.00	0.00	0.93
time (sec)	N/A	0.316	0.792	1.052	0.000	0.274	0.000	0.000	13.523

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	93	93	68	207	0	119	0	0	0
N.S.	1	1.00	0.73	2.23	0.00	1.28	0.00	0.00	0.00
time (sec)	N/A	0.492	1.158	1.123	0.000	0.096	0.000	0.000	0.000

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	35	0	58	0	0	66
N.S.	1	1.00	1.00	1.06	0.00	1.76	0.00	0.00	2.00
time (sec)	N/A	0.204	0.600	1.016	0.000	0.282	0.000	0.000	13.581

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	70	243	0	122	0	0	0
N.S.	1	1.00	0.71	2.48	0.00	1.24	0.00	0.00	0.00
time (sec)	N/A	0.522	1.207	1.229	0.000	0.105	0.000	0.000	0.000

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	329	213	143	487	0	1371	0	0	0
N.S.	1	0.65	0.43	1.48	0.00	4.17	0.00	0.00	0.00
time (sec)	N/A	0.581	2.397	8.795	0.000	0.668	0.000	0.000	0.000

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	45	56	0	79	0	0	99
N.S.	1	1.00	0.65	0.81	0.00	1.14	0.00	0.00	1.43
time (sec)	N/A	0.315	1.110	1.461	0.000	0.294	0.000	0.000	14.529

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	130	104	383	0	248	0	0	0
N.S.	1	1.02	0.81	2.99	0.00	1.94	0.00	0.00	0.00
time (sec)	N/A	0.650	1.768	1.417	0.000	0.114	0.000	0.000	0.000

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	35	0	51	0	0	49
N.S.	1	1.00	1.00	1.06	0.00	1.55	0.00	0.00	1.48
time (sec)	N/A	0.208	0.702	0.970	0.000	0.270	0.000	0.000	13.914

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	80	363	0	150	0	0	0
N.S.	1	1.00	0.90	4.08	0.00	1.69	0.00	0.00	0.00
time (sec)	N/A	0.499	1.009	1.205	0.000	0.114	0.000	0.000	0.000

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	270	171	123	361	0	1147	0	0	0
N.S.	1	0.63	0.46	1.34	0.00	4.25	0.00	0.00	0.00
time (sec)	N/A	0.456	2.192	1.215	0.000	0.541	0.000	0.000	0.000

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	66	393	0	0	0	0	0
N.S.	1	1.00	1.25	7.42	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.362	0.811	1.314	0.000	0.000	0.000	0.000	0.000



Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	322	212	156	434	0	1268	0	0	0
N.S.	1	0.66	0.48	1.35	0.00	3.94	0.00	0.00	0.00
time (sec)	N/A	0.601	2.671	1.727	0.000	0.673	0.000	0.000	0.000

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	84	410	0	0	0	0	0
N.S.	1	1.00	0.88	4.32	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.505	1.063	1.235	0.000	0.000	0.000	0.000	0.000

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	115	57	61	0	88	0	0	125
N.S.	1	1.05	0.52	0.55	0.00	0.80	0.00	0.00	1.14
time (sec)	N/A	0.497	1.729	1.265	0.000	0.318	0.000	0.000	16.493

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	139	119	231	0	183	0	0	0
N.S.	1	1.03	0.88	1.71	0.00	1.36	0.00	0.00	0.00
time (sec)	N/A	0.721	2.610	1.033	0.000	0.114	0.000	0.000	0.000

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	45	40	0	59	0	0	70
N.S.	1	1.00	1.36	1.21	0.00	1.79	0.00	0.00	2.12
time (sec)	N/A	0.212	1.065	1.013	0.000	0.281	0.000	0.000	13.736

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	105	208	0	122	0	0	0
N.S.	1	1.00	1.07	2.12	0.00	1.24	0.00	0.00	0.00
time (sec)	N/A	0.522	1.495	1.156	0.000	0.102	0.000	0.000	0.000

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	327	211	160	644	0	1286	0	0	0
N.S.	1	0.65	0.49	1.97	0.00	3.93	0.00	0.00	0.00
time (sec)	N/A	0.618	1.485	13.138	0.000	0.586	0.000	0.000	0.000

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	84	205	0	0	0	0	0
N.S.	1	1.00	0.91	2.23	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.498	1.119	8.125	0.000	0.000	0.000	0.000	0.000

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	322	212	157	438	0	1247	0	0	0
N.S.	1	0.66	0.49	1.36	0.00	3.87	0.00	0.00	0.00
time (sec)	N/A	0.582	2.203	48.780	0.000	0.651	0.000	0.000	0.000

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	135	135	89	1762	0	0	0	0	0
N.S.	1	1.00	0.66	13.05	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.670	1.221	58.144	0.000	0.000	0.000	0.000	0.000

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	371	255	168	551	0	1341	0	0	0
N.S.	1	0.69	0.45	1.49	0.00	3.61	0.00	0.00	0.00
time (sec)	N/A	0.743	2.659	50.575	0.000	1.149	0.000	0.000	0.000

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	45	40	0	67	0	0	93
N.S.	1	1.00	1.36	1.21	0.00	2.03	0.00	0.00	2.82
time (sec)	N/A	0.204	1.539	8.306	0.000	0.297	0.000	0.000	14.472

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	135	101	385	0	254	0	0	0
N.S.	1	1.00	0.75	2.85	0.00	1.88	0.00	0.00	0.00
time (sec)	N/A	0.675	2.887	7.816	0.000	0.135	0.000	0.000	0.000

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	329	213	154	730	0	1396	0	0	0
N.S.	1	0.65	0.47	2.22	0.00	4.24	0.00	0.00	0.00
time (sec)	N/A	0.595	2.788	57.000	0.000	0.692	0.000	0.000	0.000

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	94	80	377	0	0	0	0	0
N.S.	1	1.00	0.85	4.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.508	1.419	6.583	0.000	0.000	0.000	0.000	0.000

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	322	212	152	442	0	1309	0	0	0
N.S.	1	0.66	0.47	1.37	0.00	4.07	0.00	0.00	0.00
time (sec)	N/A	0.586	2.823	55.584	0.000	0.636	0.000	0.000	0.000

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	79	409	0	0	0	0	0
N.S.	1	1.00	0.83	4.31	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.500	1.044	6.607	0.000	0.000	0.000	0.000	0.000

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	371	255	165	517	0	1375	0	0	0
N.S.	1	0.69	0.44	1.39	0.00	3.71	0.00	0.00	0.00
time (sec)	N/A	0.751	3.091	48.996	0.000	1.299	0.000	0.000	0.000

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	138	90	425	0	0	0	0	0
N.S.	1	1.02	0.67	3.15	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.673	1.448	8.060	0.000	0.000	0.000	0.000	0.000

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	406	298	175	613	0	1409	0	0	0
N.S.	1	0.73	0.43	1.51	0.00	3.47	0.00	0.00	0.00
time (sec)	N/A	0.974	3.378	44.537	0.000	2.654	0.000	0.000	0.000

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	81	81	278	0	0	0	0	0	0
N.S.	1	1.00	3.43	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.344	2.659	0.000	0.000	0.000	0.000	0.000	0.000

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	86	86	280	0	0	0	0	0	0
N.S.	1	1.00	3.26	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.341	1.029	0.000	0.000	0.000	0.000	0.000	0.000

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	84	84	281	0	0	0	0	0	0
N.S.	1	1.00	3.35	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.345	1.037	0.000	0.000	0.000	0.000	0.000	0.000

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	89	89	283	0	0	0	0	0	0
N.S.	1	1.00	3.18	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.338	1.225	0.000	0.000	0.000	0.000	0.000	0.000

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	48	48	51	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.221	0.064	0.000	0.000	0.000	0.000	0.000	0.000

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	49	49	51	0	0	0	0	0	0
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.232	0.045	0.000	0.000	0.000	0.000	0.000	0.000

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	48	48	51	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.216	0.037	0.000	0.000	0.000	0.000	0.000	0.000

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	23	31	29	29	0	0	28
N.S.	1	1.00	0.96	1.29	1.21	1.21	0.00	0.00	1.17
time (sec)	N/A	0.197	0.026	1.430	0.262	0.294	0.000	0.000	13.182

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	50	45	67	58	49	0	0	66
N.S.	1	0.96	0.87	1.29	1.12	0.94	0.00	0.00	1.27
time (sec)	N/A	0.246	0.146	7.047	0.223	0.296	0.000	0.000	13.556

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	76	81	104	86	73	0	0	134
N.S.	1	0.97	1.04	1.33	1.10	0.94	0.00	0.00	1.72
time (sec)	N/A	0.259	0.589	5.014	0.232	0.270	0.000	0.000	15.071

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	72	72	77	0	0	0	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.323	1.883	0.000	0.000	0.000	0.000	0.000	0.000

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	72	72	77	0	0	0	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.316	1.530	0.000	0.000	0.000	0.000	0.000	0.000



Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	72	72	98	0	0	0	0	0	0
N.S.	1	1.00	1.36	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.326	2.654	0.000	0.000	0.000	0.000	0.000	0.000

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	72	72	65	0	0	0	0	0	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.282	0.144	0.000	0.000	0.000	0.000	0.000	0.000

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	72	72	165	0	0	0	0	0	0
N.S.	1	1.00	2.29	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.312	0.899	0.000	0.000	0.000	0.000	0.000	0.000

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	72	72	246	0	0	0	0	0	0
N.S.	1	1.00	3.42	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.306	1.246	0.000	0.000	0.000	0.000	0.000	0.000

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	81	81	92	0	0	0	0	0	0
N.S.	1	1.00	1.14	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.360	11.398	0.000	0.000	0.000	0.000	0.000	0.000

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	81	81	90	0	0	0	0	0	0
N.S.	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.352	11.377	0.000	0.000	0.000	0.000	0.000	0.000

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	81	81	95	0	0	0	0	0	0
N.S.	1	1.00	1.17	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.344	31.339	0.000	0.000	0.000	0.000	0.000	0.000

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	81	81	115	0	0	0	0	0	0
N.S.	1	1.00	1.42	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.370	11.697	0.000	0.000	0.000	0.000	0.000	0.000

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [55] had the largest ratio of [1.39999999999999991]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	2	1.00	6	0.333
2	A	4	3	1.00	8	0.375
3	A	4	4	1.00	8	0.500
4	A	4	3	1.00	8	0.375
5	A	6	6	1.09	8	0.750
6	A	4	3	0.93	8	0.375
7	A	8	8	1.13	8	1.000
8	A	4	3	0.91	8	0.375
9	A	8	8	1.01	10	0.800
10	A	6	6	1.00	10	0.600
11	A	6	6	1.00	10	0.600
12	A	4	4	1.00	10	0.400
13	A	4	4	1.00	10	0.400
14	A	6	6	1.00	10	0.600
15	A	6	6	1.00	10	0.600
16	A	8	8	1.06	10	0.800
17	A	8	8	1.02	12	0.667
18	A	6	6	1.00	12	0.500
19	A	6	6	1.00	12	0.500
20	A	4	4	1.00	12	0.333
21	A	4	4	1.00	12	0.333
22	A	6	6	1.00	12	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	6	6	1.00	12	0.500
24	A	8	8	1.08	12	0.667
25	A	4	4	1.00	10	0.400
26	A	4	4	1.00	10	0.400
27	A	4	4	1.00	10	0.400
28	A	4	4	1.00	10	0.400
29	A	4	4	1.00	10	0.400
30	A	4	4	1.00	10	0.400
31	A	4	4	1.00	12	0.333
32	A	4	4	1.00	12	0.333
33	A	4	4	1.00	12	0.333
34	A	4	4	1.00	12	0.333
35	A	4	4	1.00	12	0.333
36	A	4	4	1.00	12	0.333
37	A	4	4	1.00	8	0.500
38	A	4	4	1.00	10	0.400
39	A	7	6	1.32	8	0.750
40	A	6	5	1.25	8	0.625
41	A	5	4	1.09	8	0.500
42	A	4	3	1.00	8	0.375
43	A	4	3	1.18	8	0.375
44	A	5	4	1.14	8	0.500
45	A	6	5	1.26	8	0.625
46	A	7	6	1.32	8	0.750
47	A	8	7	1.18	10	0.700
48	A	7	6	1.15	10	0.600
49	A	6	5	1.11	10	0.500
50	A	5	4	1.08	10	0.400
51	A	4	3	1.15	10	0.300
52	A	5	4	1.25	10	0.400
53	A	6	5	1.35	10	0.500
54	A	7	6	1.39	10	0.600
55	A	14	14	0.88	10	1.400
56	A	10	10	1.09	10	1.000

Continued on next page

## 2.3. Detailed conclusion table specific for Rubi results

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	8	8	1.17	10	0.800
58	A	8	8	1.20	10	0.800
59	A	10	10	1.00	10	1.000
60	A	14	14	0.91	10	1.400
61	A	6	5	0.42	10	0.500
62	A	6	5	0.48	10	0.500
63	A	6	5	0.62	10	0.500
64	A	6	5	1.00	10	0.500
65	A	5	5	0.81	10	0.500
66	A	9	9	0.72	10	0.900
67	A	13	13	0.70	10	1.300
68	A	6	6	1.00	12	0.500
69	A	6	6	1.00	14	0.429
70	A	9	9	1.11	21	0.429
71	A	9	9	1.09	21	0.429
72	A	7	7	1.07	21	0.333
73	A	7	7	1.11	19	0.368
74	A	4	4	1.00	12	0.333
75	A	5	5	1.00	19	0.263
76	A	7	7	1.13	21	0.333
77	A	7	7	1.09	21	0.333
78	A	9	9	1.18	21	0.429
79	A	9	9	1.14	21	0.429
80	A	9	9	1.14	21	0.429
81	A	9	9	1.06	21	0.429
82	A	7	7	1.10	19	0.368
83	A	6	6	1.00	12	0.500
84	A	5	5	1.00	19	0.263
85	A	5	5	1.00	21	0.238
86	A	7	7	1.09	21	0.333
87	A	7	7	1.06	21	0.333
88	A	9	9	1.14	21	0.429
89	A	9	9	1.12	21	0.429
90	A	9	9	1.10	21	0.429

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
91	A	9	9	1.07	19	0.474
92	A	6	6	1.00	12	0.500
93	A	7	7	1.00	19	0.368
94	A	5	5	1.00	21	0.238
95	A	5	5	1.00	21	0.238
96	A	7	7	1.06	21	0.333
97	A	7	7	1.06	21	0.333
98	A	9	9	1.12	21	0.429
99	A	9	9	1.12	21	0.429
100	A	8	8	1.02	12	0.667
101	A	9	9	1.08	21	0.429
102	A	9	9	1.07	21	0.429
103	A	7	7	1.03	21	0.333
104	A	7	7	1.08	21	0.333
105	A	5	5	1.00	19	0.263
106	A	4	4	1.00	12	0.333
107	A	7	7	1.07	19	0.368
108	A	7	7	1.13	21	0.333
109	A	9	9	1.15	21	0.429
110	A	9	9	1.18	21	0.429
111	A	9	9	1.08	21	0.429
112	A	9	9	1.04	21	0.429
113	A	7	7	1.03	21	0.333
114	A	7	7	1.03	21	0.333
115	A	5	5	1.00	21	0.238
116	A	5	5	1.00	19	0.263
117	A	6	6	1.00	12	0.500
118	A	7	7	1.07	19	0.368
119	A	9	9	1.14	21	0.429
120	A	9	9	1.15	21	0.429
121	A	9	9	1.08	21	0.429
122	A	9	9	1.04	21	0.429
123	A	7	7	1.03	21	0.333
124	A	7	7	1.03	21	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
125	A	5	5	1.00	21	0.238
126	A	5	5	1.00	21	0.238
127	A	7	7	1.06	19	0.368
128	A	6	6	1.00	12	0.500
129	A	9	9	1.13	19	0.474
130	A	9	9	1.14	21	0.429
131	A	8	8	1.08	12	0.667
132	A	7	7	0.78	23	0.304
133	A	5	4	0.69	23	0.174
134	A	5	5	0.79	23	0.217
135	A	5	4	1.00	23	0.174
136	A	3	3	1.00	23	0.130
137	A	2	2	1.00	23	0.087
138	A	3	3	1.00	23	0.130
139	A	4	4	0.76	23	0.174
140	A	5	4	0.69	23	0.174
141	A	6	6	0.76	23	0.261
142	A	7	7	0.76	23	0.304
143	A	5	4	0.68	23	0.174
144	A	5	5	0.78	23	0.217
145	A	5	4	1.00	23	0.174
146	A	3	3	1.00	23	0.130
147	A	2	2	1.00	23	0.087
148	A	3	3	1.00	23	0.130
149	A	4	4	0.75	23	0.174
150	A	5	4	0.68	23	0.174
151	A	6	6	0.74	23	0.261
152	A	5	4	0.54	23	0.174
153	A	5	4	0.67	23	0.174
154	A	5	5	0.77	23	0.217
155	A	5	4	1.00	23	0.174
156	A	3	3	1.00	23	0.130
157	A	2	2	1.00	23	0.087
158	A	3	3	1.00	23	0.130

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
159	A	4	4	0.74	23	0.174
160	A	5	4	0.67	23	0.174
161	A	5	5	0.79	23	0.217
162	A	5	4	1.00	23	0.174
163	A	3	3	1.00	23	0.130
164	A	2	2	1.00	23	0.087
165	A	3	3	1.00	23	0.130
166	A	4	4	0.76	23	0.174
167	A	5	4	0.69	23	0.174
168	A	5	5	0.77	23	0.217
169	A	5	4	1.00	23	0.174
170	A	3	3	1.00	23	0.130
171	A	2	2	1.00	23	0.087
172	A	3	3	1.00	23	0.130
173	A	4	4	0.74	23	0.174
174	A	5	4	0.67	23	0.174
175	A	6	6	0.72	23	0.261
176	A	5	5	0.77	23	0.217
177	A	5	4	1.00	23	0.174
178	A	3	3	1.00	23	0.130
179	A	2	2	1.00	23	0.087
180	A	3	3	1.00	23	0.130
181	A	4	4	0.74	23	0.174
182	A	5	4	0.67	23	0.174
183	A	6	6	0.72	23	0.261
184	A	5	5	1.00	21	0.238
185	A	5	5	1.00	19	0.263
186	A	4	4	1.00	12	0.333
187	A	5	5	1.00	19	0.263
188	A	5	5	1.00	21	0.238
189	A	5	5	1.00	21	0.238
190	A	5	5	1.00	19	0.263
191	A	4	4	1.00	12	0.333
192	A	5	5	1.00	19	0.263

Continued on next page



Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
193	A	5	5	1.00	21	0.238
194	A	5	5	1.00	21	0.238
195	A	5	5	1.00	19	0.263
196	A	4	4	1.00	12	0.333
197	A	5	5	1.00	19	0.263
198	A	5	5	1.00	21	0.238
199	A	5	5	1.00	21	0.238
200	A	5	5	1.00	19	0.263
201	A	4	4	1.00	12	0.333
202	A	5	5	1.00	19	0.263
203	A	5	5	1.00	21	0.238
204	A	5	5	1.00	21	0.238
205	A	5	5	1.00	21	0.238
206	A	5	5	1.00	21	0.238
207	A	5	5	1.00	21	0.238
208	A	5	5	1.00	21	0.238
209	A	5	5	1.00	21	0.238
210	A	5	5	1.00	19	0.263
211	A	5	5	1.00	19	0.263
212	A	5	5	1.00	17	0.294
213	A	4	4	1.00	10	0.400
214	A	5	5	1.00	17	0.294
215	A	5	5	1.00	19	0.263
216	A	5	5	1.00	19	0.263
217	A	5	5	1.00	21	0.238
218	A	5	5	1.00	21	0.238
219	A	5	5	1.00	21	0.238
220	A	5	5	1.00	21	0.238
221	A	5	5	1.00	21	0.238
222	A	5	5	1.00	21	0.238
223	A	4	3	1.00	19	0.158
224	A	4	3	1.00	19	0.158
225	A	4	3	1.00	19	0.158
226	A	4	3	1.00	19	0.158

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
227	A	4	3	1.00	19	0.158
228	A	7	6	0.98	21	0.286
229	A	7	6	0.98	21	0.286
230	A	10	10	1.05	25	0.400
231	A	4	4	1.00	25	0.160
232	A	8	8	1.00	25	0.320
233	A	2	2	1.00	25	0.080
234	A	6	6	1.00	25	0.240
235	A	14	13	0.63	25	0.520
236	A	8	8	1.00	25	0.320
237	A	16	15	0.66	25	0.600
238	A	6	6	1.06	25	0.240
239	A	12	12	1.01	25	0.480
240	A	4	4	1.00	25	0.160
241	A	10	10	1.01	25	0.400
242	A	2	2	1.00	25	0.080
243	A	8	8	1.00	25	0.320
244	A	16	15	0.65	25	0.600
245	A	8	8	1.00	25	0.320
246	A	12	12	1.04	25	0.480
247	A	6	6	1.04	25	0.240
248	A	10	10	1.01	25	0.400
249	A	4	4	1.00	25	0.160
250	A	8	8	1.00	25	0.320
251	A	2	2	1.00	25	0.080
252	A	8	8	1.00	25	0.320
253	A	16	15	0.65	25	0.600
254	A	4	4	1.00	25	0.160
255	A	10	10	1.02	25	0.400
256	A	2	2	1.00	25	0.080
257	A	8	8	1.00	25	0.320
258	A	14	13	0.63	25	0.520
259	A	6	6	1.00	25	0.240
260	A	16	15	0.66	25	0.600

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
261	A	8	8	1.00	25	0.320
262	A	6	6	1.05	25	0.240
263	A	10	10	1.03	25	0.400
264	A	2	2	1.00	25	0.080
265	A	8	8	1.00	25	0.320
266	A	16	15	0.65	25	0.600
267	A	8	8	1.00	25	0.320
268	A	16	15	0.66	25	0.600
269	A	10	10	1.00	25	0.400
270	A	18	17	0.69	25	0.680
271	A	2	2	1.00	25	0.080
272	A	10	10	1.00	25	0.400
273	A	16	15	0.65	25	0.600
274	A	8	8	1.00	25	0.320
275	A	16	15	0.66	25	0.600
276	A	8	8	1.00	25	0.320
277	A	18	17	0.69	25	0.680
278	A	10	10	1.02	25	0.400
279	A	20	19	0.73	25	0.760
280	A	4	4	1.00	17	0.235
281	A	4	4	1.00	19	0.211
282	A	4	4	1.00	19	0.211
283	A	4	4	1.00	21	0.190
284	A	6	5	1.00	19	0.263
285	A	5	4	1.00	19	0.211
286	A	6	5	1.00	17	0.294
287	A	4	3	1.00	17	0.176
288	A	7	6	0.96	19	0.316
289	A	6	5	0.97	19	0.263
290	A	4	4	1.00	19	0.211
291	A	4	4	1.00	19	0.211
292	A	4	4	1.00	19	0.211
293	A	4	4	1.00	10	0.400
294	A	4	4	1.00	19	0.211

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
295	A	4	4	1.00	19	0.211
296	A	4	4	1.00	23	0.174
297	A	4	4	1.00	23	0.174
298	A	4	4	1.00	23	0.174
299	A	4	4	1.00	23	0.174

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int \sec(a + bx) dx$	121
3.2	$\int \sec^2(a + bx) dx$	126
3.3	$\int \sec^3(a + bx) dx$	131
3.4	$\int \sec^4(a + bx) dx$	136
3.5	$\int \sec^5(a + bx) dx$	141
3.6	$\int \sec^6(a + bx) dx$	146
3.7	$\int \sec^7(a + bx) dx$	151
3.8	$\int \sec^8(a + bx) dx$	157
3.9	$\int \sec^{\frac{7}{2}}(a + bx) dx$	162
3.10	$\int \sec^{\frac{5}{2}}(a + bx) dx$	168
3.11	$\int \sec^{\frac{3}{2}}(a + bx) dx$	173
3.12	$\int \sqrt{\sec(a + bx)} dx$	178
3.13	$\int \frac{1}{\sqrt{\sec(a+bx)}} dx$	183
3.14	$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx$	188
3.15	$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$	193
3.16	$\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx$	198
3.17	$\int (c \sec(a + bx))^{7/2} dx$	204
3.18	$\int (c \sec(a + bx))^{5/2} dx$	210
3.19	$\int (c \sec(a + bx))^{3/2} dx$	215
3.20	$\int \sqrt{c \sec(a + bx)} dx$	220
3.21	$\int \frac{1}{\sqrt{c \sec(a+bx)}} dx$	225
3.22	$\int \frac{1}{(c \sec(a+bx))^{3/2}} dx$	230
3.23	$\int \frac{1}{(c \sec(a+bx))^{5/2}} dx$	235
3.24	$\int \frac{1}{(c \sec(a+bx))^{7/2}} dx$	240
3.25	$\int \sec^{\frac{4}{3}}(a + bx) dx$	246
3.26	$\int \sec^{\frac{2}{3}}(a + bx) dx$	251
3.27	$\int \sqrt[3]{\sec(a + bx)} dx$	256

3.28	$\int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx$	261
3.29	$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx$	266
3.30	$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx$	271
3.31	$\int (c \sec(a+bx))^{4/3} dx$	276
3.32	$\int (c \sec(a+bx))^{2/3} dx$	281
3.33	$\int \sqrt[3]{c \sec(a+bx)} dx$	286
3.34	$\int \frac{1}{\sqrt[3]{c \sec(a+bx)}} dx$	291
3.35	$\int \frac{1}{(c \sec(a+bx))^{2/3}} dx$	296
3.36	$\int \frac{1}{(c \sec(a+bx))^{4/3}} dx$	301
3.37	$\int \sec^n(a+bx) dx$	306
3.38	$\int (c \sec(a+bx))^n dx$	310
3.39	$\int \sec^2(x)^{7/2} dx$	315
3.40	$\int \sec^2(x)^{5/2} dx$	320
3.41	$\int \sec^2(x)^{3/2} dx$	325
3.42	$\int \sqrt{\sec^2(x)} dx$	330
3.43	$\int \frac{1}{\sqrt{\sec^2(x)}} dx$	334
3.44	$\int \frac{1}{\sec^2(x)^{3/2}} dx$	338
3.45	$\int \frac{1}{\sec^2(x)^{5/2}} dx$	343
3.46	$\int \frac{1}{\sec^2(x)^{7/2}} dx$	348
3.47	$\int (a \sec^2(x))^{7/2} dx$	353
3.48	$\int (a \sec^2(x))^{5/2} dx$	359
3.49	$\int (a \sec^2(x))^{3/2} dx$	365
3.50	$\int \sqrt{a \sec^2(x)} dx$	370
3.51	$\int \frac{1}{\sqrt{a \sec^2(x)}} dx$	375
3.52	$\int \frac{1}{(a \sec^2(x))^{3/2}} dx$	379
3.53	$\int \frac{1}{(a \sec^2(x))^{5/2}} dx$	384
3.54	$\int \frac{1}{(a \sec^2(x))^{7/2}} dx$	389
3.55	$\int (a \sec^3(x))^{5/2} dx$	394
3.56	$\int (a \sec^3(x))^{3/2} dx$	401
3.57	$\int \sqrt{a \sec^3(x)} dx$	407
3.58	$\int \frac{1}{\sqrt{a \sec^3(x)}} dx$	413
3.59	$\int \frac{1}{(a \sec^3(x))^{3/2}} dx$	419
3.60	$\int \frac{1}{(a \sec^3(x))^{5/2}} dx$	425
3.61	$\int (a \sec^4(x))^{7/2} dx$	432
3.62	$\int (a \sec^4(x))^{5/2} dx$	437
3.63	$\int (a \sec^4(x))^{3/2} dx$	442

3.64	$\int \sqrt{a \sec^4(x)} dx$	447
3.65	$\int \frac{1}{\sqrt{a \sec^4(x)}} dx$	452
3.66	$\int \frac{1}{(a \sec^4(x))^{3/2}} dx$	457
3.67	$\int \frac{1}{(a \sec^4(x))^{5/2}} dx$	462
3.68	$\int ((b \sec(c + dx))^p)^n dx$	468
3.69	$\int (a(b \sec(c + dx))^p)^n dx$	473
3.70	$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx$	478
3.71	$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx$	484
3.72	$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx$	490
3.73	$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx$	495
3.74	$\int \sqrt{b \sec(c + dx)} dx$	501
3.75	$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx$	506
3.76	$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx$	511
3.77	$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx$	516
3.78	$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx$	521
3.79	$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx$	527
3.80	$\int \sec^3(c + dx) (b \sec(c + dx))^{3/2} dx$	533
3.81	$\int \sec^2(c + dx) (b \sec(c + dx))^{3/2} dx$	539
3.82	$\int \sec(c + dx) (b \sec(c + dx))^{3/2} dx$	545
3.83	$\int (b \sec(c + dx))^{3/2} dx$	551
3.84	$\int \cos(c + dx) (b \sec(c + dx))^{3/2} dx$	556
3.85	$\int \cos^2(c + dx) (b \sec(c + dx))^{3/2} dx$	561
3.86	$\int \cos^3(c + dx) (b \sec(c + dx))^{3/2} dx$	566
3.87	$\int \cos^4(c + dx) (b \sec(c + dx))^{3/2} dx$	572
3.88	$\int \cos^5(c + dx) (b \sec(c + dx))^{3/2} dx$	577
3.89	$\int \cos^6(c + dx) (b \sec(c + dx))^{3/2} dx$	583
3.90	$\int \sec^2(c + dx) (b \sec(c + dx))^{5/2} dx$	589
3.91	$\int \sec(c + dx) (b \sec(c + dx))^{5/2} dx$	595
3.92	$\int (b \sec(c + dx))^{5/2} dx$	601
3.93	$\int \cos(c + dx) (b \sec(c + dx))^{5/2} dx$	606
3.94	$\int \cos^2(c + dx) (b \sec(c + dx))^{5/2} dx$	612
3.95	$\int \cos^3(c + dx) (b \sec(c + dx))^{5/2} dx$	617
3.96	$\int \cos^4(c + dx) (b \sec(c + dx))^{5/2} dx$	622
3.97	$\int \cos^5(c + dx) (b \sec(c + dx))^{5/2} dx$	628
3.98	$\int \cos^6(c + dx) (b \sec(c + dx))^{5/2} dx$	633
3.99	$\int \cos^7(c + dx) (b \sec(c + dx))^{5/2} dx$	639
3.100	$\int (b \sec(c + dx))^{7/2} dx$	645
3.101	$\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	651
3.102	$\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	657

3.103	$\int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	663
3.104	$\int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	668
3.105	$\int \frac{\sec(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	674
3.106	$\int \frac{1}{\sqrt{b \sec(c+dx)}} dx$	679
3.107	$\int \frac{\cos(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	684
3.108	$\int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	690
3.109	$\int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	696
3.110	$\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	702
3.111	$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	708
3.112	$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	714
3.113	$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	720
3.114	$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	725
3.115	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	730
3.116	$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	735
3.117	$\int \frac{1}{(b \sec(c+dx))^{3/2}} dx$	740
3.118	$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	745
3.119	$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	750
3.120	$\int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	756
3.121	$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	762
3.122	$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	768
3.123	$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	774
3.124	$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	779
3.125	$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	784
3.126	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	789
3.127	$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	794
3.128	$\int \frac{1}{(b \sec(c+dx))^{5/2}} dx$	799
3.129	$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	804
3.130	$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	810
3.131	$\int \frac{1}{(b \sec(c+dx))^{7/2}} dx$	816
3.132	$\int \sec^{\frac{9}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	822
3.133	$\int \sec^{\frac{7}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	828
3.134	$\int \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	833
3.135	$\int \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	839



3.136	$\int \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} dx$	844
3.137	$\int \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx$	849
3.138	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx$	854
3.139	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx$	859
3.140	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx$	865
3.141	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$	870
3.142	$\int \sec^{\frac{7}{2}}(c+dx) (b \sec(c+dx))^{3/2} dx$	875
3.143	$\int \sec^{\frac{5}{2}}(c+dx) (b \sec(c+dx))^{3/2} dx$	881
3.144	$\int \sec^{\frac{3}{2}}(c+dx) (b \sec(c+dx))^{3/2} dx$	886
3.145	$\int \sqrt{\sec(c+dx)} (b \sec(c+dx))^{3/2} dx$	892
3.146	$\int \frac{(b \sec(c+dx))^{3/2}}{\sqrt{\sec(c+dx)}} dx$	897
3.147	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{3}{2}}(c+dx)} dx$	902
3.148	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{5}{2}}(c+dx)} dx$	906
3.149	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{7}{2}}(c+dx)} dx$	911
3.150	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{9}{2}}(c+dx)} dx$	916
3.151	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{11}{2}}(c+dx)} dx$	921
3.152	$\int \sec^{\frac{7}{2}}(c+dx) (b \sec(c+dx))^{5/2} dx$	926
3.153	$\int \sec^{\frac{5}{2}}(c+dx) (b \sec(c+dx))^{5/2} dx$	932
3.154	$\int \sqrt{\sec(c+dx)} (b \sec(c+dx))^{5/2} dx$	937
3.155	$\int \frac{(b \sec(c+dx))^{5/2}}{\sqrt{\sec(c+dx)}} dx$	943
3.156	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{3}{2}}(c+dx)} dx$	948
3.157	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{5}{2}}(c+dx)} dx$	953
3.158	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{7}{2}}(c+dx)} dx$	957
3.159	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{9}{2}}(c+dx)} dx$	962
3.160	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{11}{2}}(c+dx)} dx$	967
3.161	$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	972
3.162	$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	978
3.163	$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	983
3.164	$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx$	988
3.165	$\int \frac{1}{\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}} dx$	993

3.166	$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx$	998
3.167	$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx$	1003
3.168	$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx$	1008
3.169	$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx$	1014
3.170	$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx$	1019
3.171	$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx$	1024
3.172	$\int \frac{\sqrt{\sec(c+dx)}}{(b\sec(c+dx))^{\frac{3}{2}}} dx$	1028
3.173	$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{\frac{3}{2}}} dx$	1033
3.174	$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{\frac{3}{2}}} dx$	1038
3.175	$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b\sec(c+dx))^{\frac{3}{2}}} dx$	1043
3.176	$\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}} dx$	1048
3.177	$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}} dx$	1054
3.178	$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}} dx$	1059
3.179	$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}} dx$	1064
3.180	$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}} dx$	1068
3.181	$\int \frac{\sqrt{\sec(c+dx)}}{(b\sec(c+dx))^{\frac{5}{2}}} dx$	1073
3.182	$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{\frac{5}{2}}} dx$	1078
3.183	$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{\frac{5}{2}}} dx$	1083
3.184	$\int \sec^2(c+dx)\sqrt[3]{b\sec(c+dx)} dx$	1088
3.185	$\int \sec(c+dx)\sqrt[3]{b\sec(c+dx)} dx$	1093
3.186	$\int \sqrt[3]{b\sec(c+dx)} dx$	1098
3.187	$\int \cos(c+dx)\sqrt[3]{b\sec(c+dx)} dx$	1103
3.188	$\int \cos^2(c+dx)\sqrt[3]{b\sec(c+dx)} dx$	1108
3.189	$\int \sec^2(c+dx)(b\sec(c+dx))^{\frac{4}{3}} dx$	1113
3.190	$\int \sec(c+dx)(b\sec(c+dx))^{\frac{4}{3}} dx$	1118
3.191	$\int (b\sec(c+dx))^{\frac{4}{3}} dx$	1123
3.192	$\int \cos(c+dx)(b\sec(c+dx))^{\frac{4}{3}} dx$	1128
3.193	$\int \cos^2(c+dx)(b\sec(c+dx))^{\frac{4}{3}} dx$	1133
3.194	$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b\sec(c+dx)}} dx$	1138
3.195	$\int \frac{\sec(c+dx)}{\sqrt[3]{b\sec(c+dx)}} dx$	1143
3.196	$\int \frac{1}{\sqrt[3]{b\sec(c+dx)}} dx$	1148

3.197	$\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	1153
3.198	$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	1158
3.199	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	1163
3.200	$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	1168
3.201	$\int \frac{1}{(b \sec(c+dx))^{4/3}} dx$	1173
3.202	$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	1178
3.203	$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	1183
3.204	$\int \sec^m(c+dx)(b \sec(c+dx))^{4/3} dx$	1188
3.205	$\int \sec^m(c+dx)(b \sec(c+dx))^{2/3} dx$	1193
3.206	$\int \sec^m(c+dx)\sqrt[3]{b \sec(c+dx)} dx$	1198
3.207	$\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	1203
3.208	$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx$	1208
3.209	$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	1213
3.210	$\int \sec^m(c+dx)(b \sec(c+dx))^n dx$	1218
3.211	$\int \sec^2(c+dx)(b \sec(c+dx))^n dx$	1223
3.212	$\int \sec(c+dx)(b \sec(c+dx))^n dx$	1228
3.213	$\int (b \sec(c+dx))^n dx$	1233
3.214	$\int \cos(c+dx)(b \sec(c+dx))^n dx$	1238
3.215	$\int \cos^2(c+dx)(b \sec(c+dx))^n dx$	1243
3.216	$\int \cos^3(c+dx)(b \sec(c+dx))^n dx$	1248
3.217	$\int \sec^{5/2}(c+dx)(b \sec(c+dx))^n dx$	1253
3.218	$\int \sec^{3/2}(c+dx)(b \sec(c+dx))^n dx$	1258
3.219	$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^n dx$	1263
3.220	$\int \frac{(b \sec(c+dx))^n}{\sqrt{\sec(c+dx)}} dx$	1268
3.221	$\int \frac{(b \sec(c+dx))^n}{\sec^{3/2}(c+dx)} dx$	1273
3.222	$\int \frac{(b \sec(c+dx))^n}{\sec^{5/2}(c+dx)} dx$	1278
3.223	$\int (d \sec(a+bx))^{7/2} \sin(a+bx) dx$	1283
3.224	$\int (d \sec(a+bx))^{5/2} \sin(a+bx) dx$	1288
3.225	$\int (d \sec(a+bx))^{3/2} \sin(a+bx) dx$	1293
3.226	$\int \sqrt{d \sec(a+bx)} \sin(a+bx) dx$	1298
3.227	$\int \frac{\sin(a+bx)}{\sqrt{d \sec(a+bx)}} dx$	1303
3.228	$\int (d \sec(a+bx))^{5/2} \sin^3(a+bx) dx$	1308
3.229	$\int (d \sec(a+bx))^{9/2} \sin^3(a+bx) dx$	1313
3.230	$\int (d \csc(a+bx))^{9/2} \sqrt{c \sec(a+bx)} dx$	1318
3.231	$\int (d \csc(a+bx))^{7/2} \sqrt{c \sec(a+bx)} dx$	1324
3.232	$\int (d \csc(a+bx))^{5/2} \sqrt{c \sec(a+bx)} dx$	1329

3.233	$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx$	1335
3.234	$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx$	1339
3.235	$\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx$	1344
3.236	$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx$	1352
3.237	$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{5/2}} dx$	1358
3.238	$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx$	1367
3.239	$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx$	1372
3.240	$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx$	1379
3.241	$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx$	1384
3.242	$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx$	1391
3.243	$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx$	1395
3.244	$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx$	1401
3.245	$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx$	1410
3.246	$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx$	1416
3.247	$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx$	1423
3.248	$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx$	1428
3.249	$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx$	1435
3.250	$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx$	1440
3.251	$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx$	1446
3.252	$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx$	1450
3.253	$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx$	1456
3.254	$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx$	1465
3.255	$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx$	1470
3.256	$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx$	1476
3.257	$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx$	1480
3.258	$\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx$	1486
3.259	$\int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx$	1494
3.260	$\int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx$	1499
3.261	$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx$	1508
3.262	$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx$	1514
3.263	$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx$	1519
3.264	$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx$	1526
3.265	$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx$	1530
3.266	$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx$	1536

3.267	$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx$	1545
3.268	$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx$	1551
3.269	$\int \frac{1}{(d \csc(a+bx))^{3/2}(c \sec(a+bx))^{3/2}} dx$	1560
3.270	$\int \frac{1}{(d \csc(a+bx))^{5/2}(c \sec(a+bx))^{3/2}} dx$	1567
3.271	$\int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{5/2}} dx$	1578
3.272	$\int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{5/2}} dx$	1582
3.273	$\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{5/2}} dx$	1589
3.274	$\int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{5/2}} dx$	1598
3.275	$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx$	1604
3.276	$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{5/2}} dx$	1613
3.277	$\int \frac{1}{(d \csc(a+bx))^{3/2}(c \sec(a+bx))^{5/2}} dx$	1619
3.278	$\int \frac{1}{(d \csc(a+bx))^{5/2}(c \sec(a+bx))^{5/2}} dx$	1629
3.279	$\int \frac{1}{(d \csc(a+bx))^{7/2}(c \sec(a+bx))^{5/2}} dx$	1636
3.280	$\int \csc^n(e + fx) \sec^m(e + fx) dx$	1647
3.281	$\int \csc^n(e + fx)(a \sec(e + fx))^m dx$	1652
3.282	$\int (b \csc(e + fx))^n \sec^m(e + fx) dx$	1657
3.283	$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx$	1662
3.284	$\int (b \csc(e + fx))^n \sec^5(e + fx) dx$	1667
3.285	$\int (b \csc(e + fx))^n \sec^3(e + fx) dx$	1672
3.286	$\int (b \csc(e + fx))^n \sec(e + fx) dx$	1677
3.287	$\int \cos(e + fx)(b \csc(e + fx))^n dx$	1682
3.288	$\int \cos^3(e + fx)(b \csc(e + fx))^n dx$	1687
3.289	$\int \cos^5(e + fx)(b \csc(e + fx))^n dx$	1692
3.290	$\int (b \csc(e + fx))^n \sec^6(e + fx) dx$	1697
3.291	$\int (b \csc(e + fx))^n \sec^4(e + fx) dx$	1702
3.292	$\int (b \csc(e + fx))^n \sec^2(e + fx) dx$	1707
3.293	$\int (b \csc(e + fx))^n dx$	1712
3.294	$\int \cos^2(e + fx)(b \csc(e + fx))^n dx$	1717
3.295	$\int \cos^4(e + fx)(b \csc(e + fx))^n dx$	1722
3.296	$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx$	1727
3.297	$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$	1732
3.298	$\int \frac{(b \csc(e+fx))^n}{\sqrt{c \sec(e+fx)}} dx$	1737
3.299	$\int \frac{(b \csc(e+fx))^n}{(c \sec(e+fx))^{3/2}} dx$	1742

### 3.1 $\int \sec(a + bx) dx$

3.1.1	Optimal result . . . . .	121
3.1.2	Mathematica [A] (verified) . . . . .	121
3.1.3	Rubi [A] (verified) . . . . .	122
3.1.4	Maple [A] (verified) . . . . .	123
3.1.5	Fricas [B] (verification not implemented) . . . . .	123
3.1.6	Sympy [B] (verification not implemented) . . . . .	124
3.1.7	Maxima [A] (verification not implemented) . . . . .	124
3.1.8	Giac [B] (verification not implemented) . . . . .	124
3.1.9	Mupad [B] (verification not implemented) . . . . .	125

#### 3.1.1 Optimal result

Integrand size = 6, antiderivative size = 11

$$\int \sec(a + bx) dx = \frac{\operatorname{arctanh}(\sin(a + bx))}{b}$$

output `arctanh(sin(b*x+a))/b`

#### 3.1.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \sec(a + bx) dx = \frac{\operatorname{arctanh}(\sin(a + bx))}{b}$$

input `Integrate[Sec[a + b*x],x]`

output `ArcTanh[Sin[a + b*x]]/b`

### 3.1.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sec(a + bx) dx \\ \downarrow \text{3042} \\ \int \csc\left(a + bx + \frac{\pi}{2}\right) dx \\ \downarrow \text{4257} \\ \frac{\operatorname{arctanh}(\sin(a + bx))}{b} \end{array}$$

input `Int[Sec[a + b*x],x]`

output `ArcTanh[Sin[a + b*x]]/b`

#### 3.1.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.1.4 Maple [A] (verified)

Time = 1.70 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

method	result	size
derivativedivides	$\frac{\ln(\sec(bx+a)+\tan(bx+a))}{b}$	19
default	$\frac{\ln(\sec(bx+a)+\tan(bx+a))}{b}$	19
parallelrisc	$\frac{-\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)+\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{b}$	32
norman	$\frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)}{b}$	35
risc	$-\frac{\ln(e^{i(bx+a)}-i)}{b} + \frac{\ln(e^{i(bx+a)}+i)}{b}$	37

input `int(sec(b*x+a),x,method=_RETURNVERBOSE)`

output `1/b*ln(sec(b*x+a)+tan(b*x+a))`

### 3.1.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 28 vs. 2(11) = 22.

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.55

$$\int \sec(a + bx) dx = \frac{\log(\sin(bx + a) + 1) - \log(-\sin(bx + a) + 1)}{2b}$$

input `integrate(sec(b*x+a),x, algorithm="fricas")`

output `1/2*(log(sin(b*x + a) + 1) - log(-sin(b*x + a) + 1))/b`



### 3.1.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs.  $2(8) = 16$ .

Time = 0.82 (sec) , antiderivative size = 36, normalized size of antiderivative = 3.27

$$\int \sec(a + bx) dx = \begin{cases} \frac{\log(\tan(a+bx) + \sec(a+bx))}{b} & \text{for } b \neq 0 \\ \frac{x(\tan(a)\sec(a) + \sec^2(a))}{\tan(a) + \sec(a)} & \text{otherwise} \end{cases}$$

input `integrate(sec(b*x+a),x)`

output `Piecewise((log(tan(a + b*x) + sec(a + b*x))/b, Ne(b, 0)), (x*(tan(a)*sec(a) + sec(a)**2)/(tan(a) + sec(a)), True))`

### 3.1.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.64

$$\int \sec(a + bx) dx = \frac{\log(\sec(bx + a) + \tan(bx + a))}{b}$$

input `integrate(sec(b*x+a),x, algorithm="maxima")`

output `log(sec(b*x + a) + tan(b*x + a))/b`

### 3.1.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs.  $2(11) = 22$ .

Time = 0.30 (sec) , antiderivative size = 44, normalized size of antiderivative = 4.00

$$\int \sec(a + bx) dx = \frac{\log\left(\left|\frac{1}{\sin(bx+a)} + \sin(bx+a) + 2\right|\right) - \log\left(\left|\frac{1}{\sin(bx+a)} + \sin(bx+a) - 2\right|\right)}{4b}$$

input `integrate(sec(b*x+a),x, algorithm="giac")`

output `1/4*(log(abs(1/sin(b*x + a) + sin(b*x + a) + 2)) - log(abs(1/sin(b*x + a) + sin(b*x + a) - 2)))/b`

**3.1.9 Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \sec(a + bx) dx = \frac{\operatorname{atanh}(\sin(a + bx))}{b}$$

input `int(1/cos(a + b*x),x)`

output `atanh(sin(a + b*x))/b`

## 3.2 $\int \sec^2(a + bx) dx$

3.2.1	Optimal result . . . . .	126
3.2.2	Mathematica [A] (verified) . . . . .	126
3.2.3	Rubi [A] (verified) . . . . .	127
3.2.4	Maple [A] (verified) . . . . .	128
3.2.5	Fricas [A] (verification not implemented) . . . . .	128
3.2.6	Sympy [F] . . . . .	129
3.2.7	Maxima [A] (verification not implemented) . . . . .	129
3.2.8	Giac [A] (verification not implemented) . . . . .	129
3.2.9	Mupad [B] (verification not implemented) . . . . .	130

### 3.2.1 Optimal result

Integrand size = 8, antiderivative size = 10

$$\int \sec^2(a + bx) dx = \frac{\tan(a + bx)}{b}$$

output `tan(b*x+a)/b`

### 3.2.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \sec^2(a + bx) dx = \frac{\tan(a + bx)}{b}$$

input `Integrate[Sec[a + b*x]^2,x]`

output `Tan[a + b*x]/b`

### 3.2.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec^2(a + bx) dx \\
 \downarrow 3042 \\
 \int \csc\left(a + bx + \frac{\pi}{2}\right)^2 dx \\
 \downarrow 4254 \\
 \int \frac{1d(-\tan(a + bx))}{b} \\
 \downarrow 24 \\
 \frac{\tan(a + bx)}{b}
 \end{array}$$

input `Int[Sec[a + b*x]^2,x]`

output `Tan[a + b*x]/b`

#### 3.2.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

### 3.2.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
derivativedivides	$\frac{\tan(bx+a)}{b}$	11
default	$\frac{\tan(bx+a)}{b}$	11
risch	$\frac{2i}{b(e^{2i(bx+a)}+1)}$	20
norman	$-\frac{2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1\right)}$	30
parallelrisch	$-\frac{2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1\right)}$	30

input `int(sec(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `tan(b*x+a)/b`

### 3.2.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80

$$\int \sec^2(a + bx) dx = \frac{\sin(bx + a)}{b \cos(bx + a)}$$

input `integrate(sec(b*x+a)^2,x, algorithm="fricas")`

output `sin(b*x + a)/(b*cos(b*x + a))`

### 3.2.6 Sympy [F]

$$\int \sec^2(a + bx) dx = \int \sec^2(a + bx) dx$$

input `integrate(sec(b*x+a)**2,x)`

output `Integral(sec(a + b*x)**2, x)`

### 3.2.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \sec^2(a + bx) dx = \frac{\tan(bx + a)}{b}$$

input `integrate(sec(b*x+a)^2,x, algorithm="maxima")`

output `tan(b*x + a)/b`

### 3.2.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \sec^2(a + bx) dx = \frac{\tan(bx + a)}{b}$$

input `integrate(sec(b*x+a)^2,x, algorithm="giac")`

output `tan(b*x + a)/b`

**3.2.9 Mupad [B] (verification not implemented)**

Time = 12.78 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \sec^2(a + bx) dx = \frac{\tan(a + bx)}{b}$$

input `int(1/cos(a + b*x)^2,x)`

output `tan(a + b*x)/b`

### 3.3 $\int \sec^3(a + bx) dx$

3.3.1	Optimal result . . . . .	131
3.3.2	Mathematica [A] (verified) . . . . .	131
3.3.3	Rubi [A] (verified) . . . . .	132
3.3.4	Maple [A] (verified) . . . . .	133
3.3.5	Fricas [B] (verification not implemented) . . . . .	133
3.3.6	Sympy [F] . . . . .	134
3.3.7	Maxima [A] (verification not implemented) . . . . .	134
3.3.8	Giac [A] (verification not implemented) . . . . .	134
3.3.9	Mupad [B] (verification not implemented) . . . . .	135

#### 3.3.1 Optimal result

Integrand size = 8, antiderivative size = 34

$$\int \sec^3(a + bx) dx = \frac{\operatorname{arctanh}(\sin(a + bx))}{2b} + \frac{\sec(a + bx) \tan(a + bx)}{2b}$$

output `1/2*arctanh(sin(b*x+a))/b+1/2*sec(b*x+a)*tan(b*x+a)/b`

#### 3.3.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \sec^3(a + bx) dx = \frac{\operatorname{arctanh}(\sin(a + bx))}{2b} + \frac{\sec(a + bx) \tan(a + bx)}{2b}$$

input `Integrate[Sec[a + b*x]^3,x]`

output `ArcTanh[Sin[a + b*x]]/(2*b) + (Sec[a + b*x]*Tan[a + b*x])/(2*b)`



### 3.3.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{2} \int \sec(a + bx) dx + \frac{\tan(a + bx) \sec(a + bx)}{2b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \csc\left(a + bx + \frac{\pi}{2}\right) dx + \frac{\tan(a + bx) \sec(a + bx)}{2b} \\
 & \quad \downarrow \text{4257} \\
 & \frac{\operatorname{arctanh}(\sin(a + bx))}{2b} + \frac{\tan(a + bx) \sec(a + bx)}{2b}
 \end{aligned}$$

input `Int[Sec[a + b*x]^3,x]`

output `ArcTanh[Sin[a + b*x]]/(2*b) + (Sec[a + b*x]*Tan[a + b*x])/(2*b)`

#### 3.3.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.3.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

method	result	size
derivativedivides	$\frac{\frac{\sec(bx+a) \tan(bx+a)}{2} + \frac{\ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$	36
default	$\frac{\frac{\sec(bx+a) \tan(bx+a)}{2} + \frac{\ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$	36
risch	$-\frac{i(e^{3i(bx+a)} - e^{i(bx+a)})}{b(e^{2i(bx+a)} + 1)^2} - \frac{\ln(e^{i(bx+a)} - i)}{2b} + \frac{\ln(e^{i(bx+a)} + i)}{2b}$	78
parallelrisch	$\frac{(-1 - \cos(2bx+2a)) \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right) + (1 + \cos(2bx+2a)) \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right) + 2 \sin(bx+a)}{2b(1 + \cos(2bx+2a))}$	78
norman	$\frac{\frac{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} + \frac{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^3}{b}}{\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1\right)^2} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{2b} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{2b}$	81

input `int(sec(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `1/b*(1/2*sec(b*x+a)*tan(b*x+a)+1/2*ln(sec(b*x+a)+tan(b*x+a)))`

### 3.3.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs.  $2(30) = 60$ .

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.79

$$\int \sec^3(a + bx) dx$$

$$= \frac{\cos(bx + a)^2 \log(\sin(bx + a) + 1) - \cos(bx + a)^2 \log(-\sin(bx + a) + 1) + 2 \sin(bx + a)}{4b \cos(bx + a)^2}$$

input `integrate(sec(b*x+a)^3,x, algorithm="fricas")`

output `1/4*(cos(b*x + a)^2*log(sin(b*x + a) + 1) - cos(b*x + a)^2*log(-sin(b*x + a) + 1) + 2*sin(b*x + a))/(b*cos(b*x + a)^2)`

### 3.3.6 Sympy [F]

$$\int \sec^3(a + bx) dx = \int \sec^3(a + bx) dx$$

input `integrate(sec(b*x+a)**3,x)`

output `Integral(sec(a + b*x)**3, x)`

### 3.3.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.35

$$\int \sec^3(a + bx) dx = -\frac{\frac{2 \sin(bx+a)}{\sin(bx+a)^2-1} - \log(\sin(bx+a) + 1) + \log(\sin(bx+a) - 1)}{4b}$$

input `integrate(sec(b*x+a)^3,x, algorithm="maxima")`

output `-1/4*(2*sin(b*x + a)/(sin(b*x + a)^2 - 1) - log(sin(b*x + a) + 1) + log(sin(b*x + a) - 1))/b`

### 3.3.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.41

$$\int \sec^3(a + bx) dx = -\frac{\frac{2 \sin(bx+a)}{\sin(bx+a)^2-1} - \log(|\sin(bx+a) + 1|) + \log(|\sin(bx+a) - 1|)}{4b}$$

input `integrate(sec(b*x+a)^3,x, algorithm="giac")`

output `-1/4*(2*sin(b*x + a)/(sin(b*x + a)^2 - 1) - log(abs(sin(b*x + a) + 1)) + log(abs(sin(b*x + a) - 1)))/b`

**3.3.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \sec^3(a + bx) dx = \frac{\operatorname{atanh}(\sin(a + bx))}{2b} - \frac{\sin(a + bx)}{2b (\sin(a + bx)^2 - 1)}$$

input `int(1/cos(a + b*x)^3,x)`

output `atanh(sin(a + b*x))/(2*b) - sin(a + b*x)/(2*b*(sin(a + b*x)^2 - 1))`

### 3.4 $\int \sec^4(a + bx) dx$

3.4.1	Optimal result . . . . .	136
3.4.2	Mathematica [A] (verified) . . . . .	136
3.4.3	Rubi [A] (verified) . . . . .	137
3.4.4	Maple [A] (verified) . . . . .	138
3.4.5	Fricas [A] (verification not implemented) . . . . .	138
3.4.6	Sympy [F] . . . . .	139
3.4.7	Maxima [A] (verification not implemented) . . . . .	139
3.4.8	Giac [A] (verification not implemented) . . . . .	139
3.4.9	Mupad [B] (verification not implemented) . . . . .	140

#### 3.4.1 Optimal result

Integrand size = 8, antiderivative size = 26

$$\int \sec^4(a + bx) dx = \frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b}$$

output `tan(b*x+a)/b+1/3*tan(b*x+a)^3/b`

#### 3.4.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.88

$$\int \sec^4(a + bx) dx = \frac{\tan(a + bx) + \frac{1}{3} \tan^3(a + bx)}{b}$$

input `Integrate[Sec[a + b*x]^4,x]`

output `(Tan[a + b*x] + Tan[a + b*x]^3/3)/b`

### 3.4.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec^4(a + bx) dx \\
 \downarrow \text{3042} \\
 \int \csc\left(a + bx + \frac{\pi}{2}\right)^4 dx \\
 \downarrow \text{4254} \\
 -\frac{\int (\tan^2(a + bx) + 1) d(-\tan(a + bx))}{b} \\
 \downarrow \text{2009} \\
 -\frac{\frac{1}{3}\tan^3(a + bx) - \tan(a + bx)}{b}
 \end{array}$$

input `Int[Sec[a + b*x]^4,x]`

output `-((-Tan[a + b*x] - Tan[a + b*x]^3/3)/b)`

#### 3.4.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

### 3.4.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$-\frac{\left(-\frac{2}{3}-\frac{\sec(bx+a)^2}{3}\right)\tan(bx+a)}{b}$	24
default	$-\frac{\left(-\frac{2}{3}-\frac{\sec(bx+a)^2}{3}\right)\tan(bx+a)}{b}$	24
risch	$\frac{4i(3e^{2i(bx+a)}+1)}{3b(e^{2i(bx+a)}+1)^3}$	33
norman	$-\frac{2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{b}+\frac{4\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^3}{3b}-\frac{2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^5}{b}$ $\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1\right)^3$	64
parallelrisch	$\frac{-6\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^5+4\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^3-6\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{3b\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)^3\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)^3}$	70

input `int(sec(b*x+a)^4,x,method=_RETURNVERBOSE)`

output `-1/b*(-2/3-1/3*sec(b*x+a)^2)*tan(b*x+a)`

### 3.4.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.19

$$\int \sec^4(a+bx) dx = \frac{(2 \cos(bx+a)^2 + 1) \sin(bx+a)}{3b \cos(bx+a)^3}$$

input `integrate(sec(b*x+a)^4,x, algorithm="fricas")`

output `1/3*(2*cos(b*x + a)^2 + 1)*sin(b*x + a)/(b*cos(b*x + a)^3)`

### 3.4.6 Sympy [F]

$$\int \sec^4(a + bx) dx = \int \sec^4(a + bx) dx$$

input `integrate(sec(b*x+a)**4,x)`

output `Integral(sec(a + b*x)**4, x)`

### 3.4.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \sec^4(a + bx) dx = \frac{\tan(bx + a)^3 + 3 \tan(bx + a)}{3b}$$

input `integrate(sec(b*x+a)^4,x, algorithm="maxima")`

output `1/3*(tan(b*x + a)^3 + 3*tan(b*x + a))/b`

### 3.4.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \sec^4(a + bx) dx = \frac{\tan(bx + a)^3 + 3 \tan(bx + a)}{3b}$$

input `integrate(sec(b*x+a)^4,x, algorithm="giac")`

output `1/3*(tan(b*x + a)^3 + 3*tan(b*x + a))/b`



**3.4.9 Mupad [B] (verification not implemented)**

Time = 12.67 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \sec^4(a + bx) dx = \frac{\tan(a + bx) (\tan(a + bx)^2 + 3)}{3b}$$

input `int(1/cos(a + b*x)^4,x)`

output `(tan(a + b*x)*(tan(a + b*x)^2 + 3))/(3*b)`

### 3.5 $\int \sec^5(a + bx) dx$

3.5.1	Optimal result . . . . .	141
3.5.2	Mathematica [A] (verified) . . . . .	141
3.5.3	Rubi [A] (verified) . . . . .	142
3.5.4	Maple [A] (verified) . . . . .	143
3.5.5	Fricas [A] (verification not implemented) . . . . .	144
3.5.6	Sympy [F] . . . . .	144
3.5.7	Maxima [A] (verification not implemented) . . . . .	144
3.5.8	Giac [A] (verification not implemented) . . . . .	145
3.5.9	Mupad [B] (verification not implemented) . . . . .	145

#### 3.5.1 Optimal result

Integrand size = 8, antiderivative size = 55

$$\int \sec^5(a + bx) dx = \frac{3\arctanh(\sin(a + bx))}{8b} + \frac{3 \sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec^3(a + bx) \tan(a + bx)}{4b}$$

output `3/8*arctanh(sin(b*x+a))/b+3/8*sec(b*x+a)*tan(b*x+a)/b+1/4*sec(b*x+a)^3*tan(b*x+a)/b`

#### 3.5.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00

$$\int \sec^5(a + bx) dx = \frac{3\arctanh(\sin(a + bx))}{8b} + \frac{3 \sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec^3(a + bx) \tan(a + bx)}{4b}$$

input `Integrate[Sec[a + b*x]^5,x]`

output `(3*ArcTanh[Sin[a + b*x]])/(8*b) + (3*Sec[a + b*x]*Tan[a + b*x])/(8*b) + (Sec[a + b*x]^3*Tan[a + b*x])/(4*b)`

### 3.5.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^5(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^5 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{4} \int \sec^3(a + bx) dx + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int \csc\left(a + bx + \frac{\pi}{2}\right)^3 dx + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{4} \left( \frac{1}{2} \int \sec(a + bx) dx + \frac{\tan(a + bx) \sec(a + bx)}{2b} \right) + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \left( \frac{1}{2} \int \csc\left(a + bx + \frac{\pi}{2}\right) dx + \frac{\tan(a + bx) \sec(a + bx)}{2b} \right) + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \\
 & \quad \downarrow \text{4257} \\
 & \frac{3}{4} \left( \frac{\operatorname{arctanh}(\sin(a + bx))}{2b} + \frac{\tan(a + bx) \sec(a + bx)}{2b} \right) + \frac{\tan(a + bx) \sec^3(a + bx)}{4b}
 \end{aligned}$$

input `Int[Sec[a + b*x]^5,x]`

output `(Sec[a + b*x]^3*Tan[a + b*x])/(4*b) + (3*(ArcTanh[Sin[a + b*x]]/(2*b) + Sec[a + b*x]*Tan[a + b*x])/(2*b))/4`

## 3.5.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

## 3.5.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{-\left(\frac{\sec(bx+a)^3}{4} - \frac{3\sec(bx+a)}{8}\right)\tan(bx+a) + \frac{3\ln(\sec(bx+a)+\tan(bx+a))}{8}}{b}$
default	$\frac{-\left(\frac{\sec(bx+a)^3}{4} - \frac{3\sec(bx+a)}{8}\right)\tan(bx+a) + \frac{3\ln(\sec(bx+a)+\tan(bx+a))}{8}}{b}$
risch	$-\frac{i(3e^{7i(bx+a)}+11e^{5i(bx+a)}-11e^{3i(bx+a)}-3e^{i(bx+a)})}{4b(e^{2i(bx+a)}+1)^4} + \frac{3\ln(e^{i(bx+a)}+i)}{8b} - \frac{3\ln(e^{i(bx+a)}-i)}{8b}$
norman	$\frac{\frac{5\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{4b} + \frac{3\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^3}{4b} + \frac{3\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^5}{4b} + \frac{5\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^7}{4b}}{\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1\right)^4} - \frac{3\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)}{8b} + \frac{3\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{8b}$
parallelrisc	$\frac{(-12\cos(2bx+2a)-3\cos(4bx+4a)-9)\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)+(12\cos(2bx+2a)+3\cos(4bx+4a)+9)\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{8b(\cos(4bx+4a)+4\cos(2bx+2a)+3)}$

input `int(sec(b*x+a)^5,x,method=_RETURNVERBOSE)`

output `1/b*(-(-1/4*sec(b*x+a)^3-3/8*sec(b*x+a))*tan(b*x+a)+3/8*ln(sec(b*x+a)+tan(b*x+a)))`

### 3.5.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.35

$$\int \sec^5(a + bx) dx = \frac{3 \cos(bx + a)^4 \log(\sin(bx + a) + 1) - 3 \cos(bx + a)^4 \log(-\sin(bx + a) + 1) + 2(3 \cos(bx + a)^2 + 2) \sin(bx + a)}{16 b \cos(bx + a)^4}$$

input `integrate(sec(b*x+a)^5,x, algorithm="fricas")`

output `1/16*(3*cos(b*x + a)^4*log(sin(b*x + a) + 1) - 3*cos(b*x + a)^4*log(-sin(b*x + a) + 1) + 2*(3*cos(b*x + a)^2 + 2)*sin(b*x + a))/(b*cos(b*x + a)^4)`

### 3.5.6 Sympy [F]

$$\int \sec^5(a + bx) dx = \int \sec^5(a + bx) dx$$

input `integrate(sec(b*x+a)**5,x)`

output `Integral(sec(a + b*x)**5, x)`

### 3.5.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.29

$$\int \sec^5(a + bx) dx = -\frac{2(3 \sin(bx+a)^3 - 5 \sin(bx+a))}{\sin(bx+a)^4 - 2 \sin(bx+a)^2 + 1} - \frac{3 \log(\sin(bx + a) + 1) + 3 \log(\sin(bx + a) - 1)}{16 b}$$

input `integrate(sec(b*x+a)^5,x, algorithm="maxima")`

output `-1/16*(2*(3*sin(b*x + a)^3 - 5*sin(b*x + a))/(sin(b*x + a)^4 - 2*sin(b*x + a)^2 + 1) - 3*log(sin(b*x + a) + 1) + 3*log(sin(b*x + a) - 1))/b`

**3.5.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.15

$$\int \sec^5(a + bx) dx = -\frac{\frac{2(3 \sin(bx+a)^3 - 5 \sin(bx+a))}{(\sin(bx+a)^2 - 1)^2} - 3 \log(|\sin(bx+a) + 1|) + 3 \log(|\sin(bx+a) - 1|)}{16b}$$

input `integrate(sec(b*x+a)^5,x, algorithm="giac")`

output `-1/16*(2*(3*sin(b*x + a)^3 - 5*sin(b*x + a))/(sin(b*x + a)^2 - 1)^2 - 3*log(abs(sin(b*x + a) + 1)) + 3*log(abs(sin(b*x + a) - 1)))/b`

**3.5.9 Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.05

$$\int \sec^5(a + bx) dx = \frac{3 \operatorname{atanh}(\sin(a + bx))}{8b} + \frac{\frac{5 \sin(a+bx)}{8} - \frac{3 \sin(a+bx)^3}{8}}{b (\sin(a + bx)^4 - 2 \sin(a + bx)^2 + 1)}$$

input `int(1/cos(a + b*x)^5,x)`

output `(3*atanh(sin(a + b*x)))/(8*b) + ((5*sin(a + b*x))/8 - (3*sin(a + b*x)^3)/8)/(b*(sin(a + b*x)^4 - 2*sin(a + b*x)^2 + 1))`

### 3.6 $\int \sec^6(a + bx) dx$

3.6.1	Optimal result . . . . .	146
3.6.2	Mathematica [A] (verified) . . . . .	146
3.6.3	Rubi [A] (verified) . . . . .	147
3.6.4	Maple [A] (verified) . . . . .	148
3.6.5	Fricas [A] (verification not implemented) . . . . .	148
3.6.6	Sympy [F] . . . . .	149
3.6.7	Maxima [A] (verification not implemented) . . . . .	149
3.6.8	Giac [A] (verification not implemented) . . . . .	149
3.6.9	Mupad [B] (verification not implemented) . . . . .	150

#### 3.6.1 Optimal result

Integrand size = 8, antiderivative size = 41

$$\int \sec^6(a + bx) dx = \frac{\tan(a + bx)}{b} + \frac{2 \tan^3(a + bx)}{3b} + \frac{\tan^5(a + bx)}{5b}$$

output `tan(b*x+a)/b+2/3*tan(b*x+a)^3/b+1/5*tan(b*x+a)^5/b`

#### 3.6.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.85

$$\int \sec^6(a + bx) dx = \frac{\tan(a + bx) + \frac{2}{3} \tan^3(a + bx) + \frac{1}{5} \tan^5(a + bx)}{b}$$

input `Integrate[Sec[a + b*x]^6,x]`

output `(Tan[a + b*x] + (2*Tan[a + b*x]^3)/3 + Tan[a + b*x]^5/5)/b`

### 3.6.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec^6(a + bx) dx \\
 \downarrow \text{3042} \\
 \int \csc\left(a + bx + \frac{\pi}{2}\right)^6 dx \\
 \downarrow \text{4254} \\
 \frac{\int (\tan^4(a + bx) + 2 \tan^2(a + bx) + 1) d(-\tan(a + bx))}{b} \\
 \downarrow \text{2009} \\
 \frac{-\frac{1}{5} \tan^5(a + bx) - \frac{2}{3} \tan^3(a + bx) - \tan(a + bx)}{b}
 \end{array}$$

input `Int[Sec[a + b*x]^6,x]`

output `-((-Tan[a + b*x] - (2*Tan[a + b*x]^3)/3 - Tan[a + b*x]^5/5)/b)`

#### 3.6.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`



### 3.6.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{\left(-\frac{8}{15}-\frac{\sec(bx+a)^4}{5}-\frac{4\sec(bx+a)^2}{15}\right)\tan(bx+a)}{b}$	34
default	$-\frac{\left(-\frac{8}{15}-\frac{\sec(bx+a)^4}{5}-\frac{4\sec(bx+a)^2}{15}\right)\tan(bx+a)}{b}$	34
risch	$\frac{16i(10e^{4i(bx+a)}+5e^{2i(bx+a)}+1)}{15b(e^{2i(bx+a)}+1)^5}$	44
norman	$-\frac{2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{b}+\frac{8\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^3}{3b}-\frac{116\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^5}{15b}+\frac{8\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^7}{3b}-\frac{2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^9}{b}$	96
paralelrisch	$\frac{-2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^9+\frac{8\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^7}{3}-\frac{116\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^5}{15}+\frac{8\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^3}{3}-2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{b\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)^5\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)^5}$	96

input `int(sec(b*x+a)^6,x,method=_RETURNVERBOSE)`

output `-1/b*(-8/15-1/5*sec(b*x+a)^4-4/15*sec(b*x+a)^2)*tan(b*x+a)`

### 3.6.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \sec^6(a+bx) dx = \frac{(8 \cos(bx+a)^4 + 4 \cos(bx+a)^2 + 3) \sin(bx+a)}{15b \cos(bx+a)^5}$$

input `integrate(sec(b*x+a)^6,x, algorithm="fricas")`

output `1/15*(8*cos(b*x + a)^4 + 4*cos(b*x + a)^2 + 3)*sin(b*x + a)/(b*cos(b*x + a)^5)`

### 3.6.6 Sympy [F]

$$\int \sec^6(a + bx) dx = \int \sec^6(a + bx) dx$$

input `integrate(sec(b*x+a)**6,x)`

output `Integral(sec(a + b*x)**6, x)`

### 3.6.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \sec^6(a + bx) dx = \frac{3 \tan^5(bx + a) + 10 \tan^3(bx + a) + 15 \tan(bx + a)}{15b}$$

input `integrate(sec(b*x+a)^6,x, algorithm="maxima")`

output `1/15*(3*tan(b*x + a)^5 + 10*tan(b*x + a)^3 + 15*tan(b*x + a))/b`

### 3.6.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \sec^6(a + bx) dx = \frac{3 \tan^5(bx + a) + 10 \tan^3(bx + a) + 15 \tan(bx + a)}{15b}$$

input `integrate(sec(b*x+a)^6,x, algorithm="giac")`

output `1/15*(3*tan(b*x + a)^5 + 10*tan(b*x + a)^3 + 15*tan(b*x + a))/b`

**3.6.9 Mupad [B] (verification not implemented)**

Time = 12.53 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \sec^6(a + bx) dx = \frac{\tan(a+bx)^5}{5} + \frac{2\tan(a+bx)^3}{3} + \frac{\tan(a + bx)}{b}$$

input `int(1/cos(a + b*x)^6,x)`

output `(tan(a + b*x) + (2*tan(a + b*x)^3)/3 + tan(a + b*x)^5/5)/b`

### 3.7 $\int \sec^7(a + bx) dx$

3.7.1	Optimal result . . . . .	151
3.7.2	Mathematica [A] (verified) . . . . .	151
3.7.3	Rubi [A] (verified) . . . . .	152
3.7.4	Maple [A] (verified) . . . . .	154
3.7.5	Fricas [A] (verification not implemented) . . . . .	154
3.7.6	Sympy [F] . . . . .	155
3.7.7	Maxima [A] (verification not implemented) . . . . .	155
3.7.8	Giac [A] (verification not implemented) . . . . .	155
3.7.9	Mupad [B] (verification not implemented) . . . . .	156

#### 3.7.1 Optimal result

Integrand size = 8, antiderivative size = 76

$$\int \sec^7(a + bx) dx = \frac{5\operatorname{arctanh}(\sin(a + bx))}{16b} + \frac{5 \sec(a + bx) \tan(a + bx)}{16b} + \frac{5 \sec^3(a + bx) \tan(a + bx)}{24b} + \frac{\sec^5(a + bx) \tan(a + bx)}{6b}$$

output `5/16*arctanh(sin(b*x+a))/b+5/16*sec(b*x+a)*tan(b*x+a)/b+5/24*sec(b*x+a)^3*tan(b*x+a)/b+1/6*sec(b*x+a)^5*tan(b*x+a)/b`

#### 3.7.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00

$$\int \sec^7(a + bx) dx = \frac{5\operatorname{arctanh}(\sin(a + bx))}{16b} + \frac{5 \sec(a + bx) \tan(a + bx)}{16b} + \frac{5 \sec^3(a + bx) \tan(a + bx)}{24b} + \frac{\sec^5(a + bx) \tan(a + bx)}{6b}$$

input `Integrate[Sec[a + b*x]^7,x]`

output `(5*ArcTanh[Sin[a + b*x]])/(16*b) + (5*Sec[a + b*x]*Tan[a + b*x])/(16*b) + (5*Sec[a + b*x]^3*Tan[a + b*x])/(24*b) + (Sec[a + b*x]^5*Tan[a + b*x])/(6*b)`

### 3.7.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.13, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 4255, 3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^7(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^7 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{5}{6} \int \sec^5(a + bx) dx + \frac{\tan(a + bx) \sec^5(a + bx)}{6b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \int \csc\left(a + bx + \frac{\pi}{2}\right)^5 dx + \frac{\tan(a + bx) \sec^5(a + bx)}{6b} \\
 & \quad \downarrow \text{4255} \\
 & \frac{5}{6} \left( \frac{3}{4} \int \sec^3(a + bx) dx + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \right) + \frac{\tan(a + bx) \sec^5(a + bx)}{6b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \left( \frac{3}{4} \int \csc\left(a + bx + \frac{\pi}{2}\right)^3 dx + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \right) + \frac{\tan(a + bx) \sec^5(a + bx)}{6b} \\
 & \quad \downarrow \text{4255} \\
 & \frac{5}{6} \left( \frac{3}{4} \left( \frac{1}{2} \int \sec(a + bx) dx + \frac{\tan(a + bx) \sec(a + bx)}{2b} \right) + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \right) + \\
 & \quad \frac{\tan(a + bx) \sec^5(a + bx)}{6b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \left( \frac{3}{4} \left( \frac{1}{2} \int \csc\left(a + bx + \frac{\pi}{2}\right) dx + \frac{\tan(a + bx) \sec(a + bx)}{2b} \right) + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} \right) + \\
 & \quad \frac{\tan(a + bx) \sec^5(a + bx)}{6b} \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

$$\frac{5}{6} \left( \frac{3}{4} \left( \frac{\operatorname{arctanh}(\sin(a+bx))}{2b} + \frac{\tan(a+bx)\sec(a+bx)}{2b} \right) + \frac{\tan(a+bx)\sec^3(a+bx)}{4b} \right) + \frac{\tan(a+bx)\sec^5(a+bx)}{6b}$$

input `Int[Sec[a + b*x]^7,x]`

output `(Sec[a + b*x]^5*Tan[a + b*x])/(6*b) + (5*((Sec[a + b*x]^3*Tan[a + b*x])/(4*b) + (3*(ArcTanh[Sin[a + b*x]]/(2*b) + (Sec[a + b*x]*Tan[a + b*x])/(2*b)))/4))/6`

### 3.7.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.7.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{-\left(-\frac{\sec(bx+a)^5}{6}-\frac{5\sec(bx+a)^3}{24}-\frac{5\sec(bx+a)}{16}\right)\tan(bx+a)+\frac{5\ln(\sec(bx+a)+\tan(bx+a))}{16}}{b}$
default	$\frac{-\left(-\frac{\sec(bx+a)^5}{6}-\frac{5\sec(bx+a)^3}{24}-\frac{5\sec(bx+a)}{16}\right)\tan(bx+a)+\frac{5\ln(\sec(bx+a)+\tan(bx+a))}{16}}{b}$
risch	$-\frac{i(15e^{11i(bx+a)}+85e^{9i(bx+a)}+198e^{7i(bx+a)}-198e^{5i(bx+a)}-85e^{3i(bx+a)}-15e^{i(bx+a)})}{24b(e^{2i(bx+a)}+1)^6}-\frac{5\ln(e^{i(bx+a)}-i)}{16b}+\frac{5\ln(e^{i(bx+a)}+i)}{16b}$
norman	$\frac{\frac{11\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{8b}+\frac{5\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^3}{24b}+\frac{15\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^5}{4b}-\frac{15\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^7}{4b}+\frac{5\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^9}{24b}+\frac{11\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^{11}}{8b}}{\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1\right)^6}-\frac{5\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{16b}$
parallelrisc	$\frac{(-225\cos(2bx+2a)-90\cos(4bx+4a)-15\cos(6bx+6a)-150)\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)+(225\cos(2bx+2a)+90\cos(4bx+4a)+150)\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{48b(10+\cos(6bx+6a))+6\cos(4bx+4a)+150}$

input `int(sec(b*x+a)^7,x,method=_RETURNVERBOSE)`

output `1/b*(-(-1/6*sec(b*x+a)^5-5/24*sec(b*x+a)^3-5/16*sec(b*x+a))*tan(b*x+a)+5/16*ln(sec(b*x+a)+tan(b*x+a)))`

### 3.7.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.11

$$\int \sec^7(a + bx) dx = \frac{15 \cos(bx + a)^6 \log(\sin(bx + a) + 1) - 15 \cos(bx + a)^6 \log(-\sin(bx + a) + 1) + 2(15 \cos(bx + a)^4 + 10 \cos(bx + a)^2 + 8) \sin(bx + a)}{96 b \cos(bx + a)^6}$$

input `integrate(sec(b*x+a)^7,x, algorithm="fracas")`

output `1/96*(15*cos(b*x + a)^6*log(sin(b*x + a) + 1) - 15*cos(b*x + a)^6*log(-sin(b*x + a) + 1) + 2*(15*cos(b*x + a)^4 + 10*cos(b*x + a)^2 + 8)*sin(b*x + a))/(b*cos(b*x + a)^6)`

### 3.7.6 Sympy [F]

$$\int \sec^7(a + bx) dx = \int \sec^7(a + bx) dx$$

input `integrate(sec(b*x+a)**7,x)`

output `Integral(sec(a + b*x)**7, x)`

### 3.7.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.20

$$\int \sec^7(a + bx) dx = \frac{2 \left( 15 \sin(bx+a)^5 - 40 \sin(bx+a)^3 + 33 \sin(bx+a) \right)}{\sin(bx+a)^6 - 3 \sin(bx+a)^4 + 3 \sin(bx+a)^2 - 1} - 15 \log(\sin(bx+a) + 1) + 15 \log(\sin(bx+a) - 1)}{96b}$$

input `integrate(sec(b*x+a)^7,x, algorithm="maxima")`

output `-1/96*(2*(15*sin(b*x + a)^5 - 40*sin(b*x + a)^3 + 33*sin(b*x + a))/(sin(b*x + a)^6 - 3*sin(b*x + a)^4 + 3*sin(b*x + a)^2 - 1) - 15*log(sin(b*x + a) + 1) + 15*log(sin(b*x + a) - 1))/b`

### 3.7.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.96

$$\int \sec^7(a + bx) dx = \frac{2 \left( 15 \sin(bx+a)^5 - 40 \sin(bx+a)^3 + 33 \sin(bx+a) \right)}{\left( \sin(bx+a)^2 - 1 \right)^3} - 15 \log(|\sin(bx+a) + 1|) + 15 \log(|\sin(bx+a) - 1|)}{96b}$$

input `integrate(sec(b*x+a)^7,x, algorithm="giac")`



output  $-1/96*(2*(15*\sin(b*x + a)^5 - 40*\sin(b*x + a)^3 + 33*\sin(b*x + a))/(\sin(b*x + a)^2 - 1)^3 - 15*\log(\text{abs}(\sin(b*x + a) + 1)) + 15*\log(\text{abs}(\sin(b*x + a) - 1)))/b$

### 3.7.9 Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.04

$$\int \sec^7(a + bx) dx = \frac{5 \operatorname{atanh}(\sin(a + bx))}{16b} - \frac{\frac{5 \sin(a+bx)^5}{16} - \frac{5 \sin(a+bx)^3}{6} + \frac{11 \sin(a+bx)}{16}}{b (\sin(a + bx)^6 - 3 \sin(a + bx)^4 + 3 \sin(a + bx)^2 - 1)}$$

input `int(1/cos(a + b*x)^7,x)`

output  $(5*\operatorname{atanh}(\sin(a + b*x)))/(16*b) - ((11*\sin(a + b*x))/16 - (5*\sin(a + b*x)^3)/6 + (5*\sin(a + b*x)^5)/16)/(b*(3*\sin(a + b*x)^2 - 3*\sin(a + b*x)^4 + \sin(a + b*x)^6 - 1))$

### 3.8 $\int \sec^8(a + bx) dx$

3.8.1	Optimal result . . . . .	157
3.8.2	Mathematica [A] (verified) . . . . .	157
3.8.3	Rubi [A] (verified) . . . . .	158
3.8.4	Maple [A] (verified) . . . . .	159
3.8.5	Fricas [A] (verification not implemented) . . . . .	159
3.8.6	Sympy [F] . . . . .	160
3.8.7	Maxima [A] (verification not implemented) . . . . .	160
3.8.8	Giac [A] (verification not implemented) . . . . .	160
3.8.9	Mupad [B] (verification not implemented) . . . . .	161

#### 3.8.1 Optimal result

Integrand size = 8, antiderivative size = 53

$$\int \sec^8(a + bx) dx = \frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{b} + \frac{3 \tan^5(a + bx)}{5b} + \frac{\tan^7(a + bx)}{7b}$$

output `tan(b*x+a)/b+tan(b*x+a)^3/b+3/5*tan(b*x+a)^5/b+1/7*tan(b*x+a)^7/b`

#### 3.8.2 Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.81

$$\int \sec^8(a + bx) dx = \frac{\tan(a + bx) + \tan^3(a + bx) + \frac{3}{5} \tan^5(a + bx) + \frac{1}{7} \tan^7(a + bx)}{b}$$

input `Integrate[Sec[a + b*x]^8,x]`

output `(Tan[a + b*x] + Tan[a + b*x]^3 + (3*Tan[a + b*x]^5)/5 + Tan[a + b*x]^7/7)/b`

### 3.8.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.91, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^8(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^8 dx \\
 & \quad \downarrow \text{4254} \\
 & \frac{\int (\tan^6(a + bx) + 3 \tan^4(a + bx) + 3 \tan^2(a + bx) + 1) d(-\tan(a + bx))}{b} \\
 & \quad \downarrow \text{2009} \\
 & \frac{-\frac{1}{7} \tan^7(a + bx) - \frac{3}{5} \tan^5(a + bx) - \tan^3(a + bx) - \tan(a + bx)}{b}
 \end{aligned}$$

input `Int[Sec[a + b*x]^8,x]`

output `-((-Tan[a + b*x] - Tan[a + b*x]^3 - (3*Tan[a + b*x]^5)/5 - Tan[a + b*x]^7/7)/b)`

#### 3.8.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

### 3.8.4 Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{\left(-\frac{16}{35}-\frac{\sec(bx+a)^6}{7}-\frac{6\sec(bx+a)^4}{35}-\frac{8\sec(bx+a)^2}{35}\right)\tan(bx+a)}{b}$
default	$\frac{\left(-\frac{16}{35}-\frac{\sec(bx+a)^6}{7}-\frac{6\sec(bx+a)^4}{35}-\frac{8\sec(bx+a)^2}{35}\right)\tan(bx+a)}{b}$
risch	$\frac{32i(35e^{6i(bx+a)}+21e^{4i(bx+a)}+7e^{2i(bx+a)}+1)}{35b(e^{2i(bx+a)}+1)^7}$
parallelrisch	$\frac{2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^{12}-2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^{10}+\frac{43\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^8}{5}-\frac{212\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^6}{35}+\frac{43\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^4}{5}-2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^2+2\right)}{b\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1\right)^7}$
norman	$\frac{-\frac{2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{b}+\frac{4\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^3}{b}-\frac{86\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^5}{5b}+\frac{424\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^7}{35b}-\frac{86\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^9}{5b}+\frac{4\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^{11}}{b}-\frac{2\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^{13}}{b}}{\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1\right)^7}$

input `int(sec(b*x+a)^8,x,method=_RETURNVERBOSE)`

output `-1/b*(-16/35-1/7*sec(b*x+a)^6-6/35*sec(b*x+a)^4-8/35*sec(b*x+a)^2)*tan(b*x+a)`

### 3.8.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.96

$$\int \sec^8(a + bx) dx = \frac{(16 \cos(bx + a)^6 + 8 \cos(bx + a)^4 + 6 \cos(bx + a)^2 + 5) \sin(bx + a)}{35 b \cos(bx + a)^7}$$

input `integrate(sec(b*x+a)^8,x, algorithm="fricas")`

output `1/35*(16*cos(b*x + a)^6 + 8*cos(b*x + a)^4 + 6*cos(b*x + a)^2 + 5)*sin(b*x + a)/(b*cos(b*x + a)^7)`

### 3.8.6 Sympy [F]

$$\int \sec^8(a + bx) dx = \int \sec^8(a + bx) dx$$

input `integrate(sec(b*x+a)**8,x)`

output `Integral(sec(a + b*x)**8, x)`

### 3.8.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.83

$$\int \sec^8(a + bx) dx = \frac{5 \tan^7(bx + a) + 21 \tan^5(bx + a) + 35 \tan^3(bx + a) + 35 \tan(bx + a)}{35 b}$$

input `integrate(sec(b*x+a)^8,x, algorithm="maxima")`

output `1/35*(5*tan(b*x + a)^7 + 21*tan(b*x + a)^5 + 35*tan(b*x + a)^3 + 35*tan(b*x + a))/b`

### 3.8.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.83

$$\int \sec^8(a + bx) dx = \frac{5 \tan^7(bx + a) + 21 \tan^5(bx + a) + 35 \tan^3(bx + a) + 35 \tan(bx + a)}{35 b}$$

input `integrate(sec(b*x+a)^8,x, algorithm="giac")`

output `1/35*(5*tan(b*x + a)^7 + 21*tan(b*x + a)^5 + 35*tan(b*x + a)^3 + 35*tan(b*x + a))/b`

**3.8.9 Mupad [B] (verification not implemented)**

Time = 12.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.74

$$\int \sec^8(a + bx) dx = \frac{\frac{\tan(a+bx)^7}{7} + \frac{3\tan(a+bx)^5}{5} + \tan(a + bx)^3 + \tan(a + bx)}{b}$$

input `int(1/cos(a + b*x)^8,x)`

output `(tan(a + b*x) + tan(a + b*x)^3 + (3*tan(a + b*x)^5)/5 + tan(a + b*x)^7/7)/b`

### 3.9 $\int \sec^{\frac{7}{2}}(a + bx) dx$

3.9.1	Optimal result . . . . .	162
3.9.2	Mathematica [A] (verified) . . . . .	162
3.9.3	Rubi [A] (verified) . . . . .	163
3.9.4	Maple [B] (verified) . . . . .	165
3.9.5	Fricas [C] (verification not implemented) . . . . .	165
3.9.6	Sympy [F(-1)] . . . . .	166
3.9.7	Maxima [F] . . . . .	166
3.9.8	Giac [F] . . . . .	166
3.9.9	Mupad [F(-1)] . . . . .	167

#### 3.9.1 Optimal result

Integrand size = 10, antiderivative size = 85

$$\int \sec^{\frac{7}{2}}(a + bx) dx = -\frac{6\sqrt{\cos(a + bx)}E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{\sec(a + bx)}}{5b} + \frac{6\sqrt{\sec(a + bx)} \sin(a + bx)}{5b} + \frac{2 \sec^{\frac{5}{2}}(a + bx) \sin(a + bx)}{5b}$$

output `2/5*sec(b*x+a)^(5/2)*sin(b*x+a)/b+6/5*sin(b*x+a)*sec(b*x+a)^(1/2)/b-6/5*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.9.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.69

$$\int \sec^{\frac{7}{2}}(a + bx) dx = \frac{\sec^{\frac{5}{2}}(a + bx) \left( -12 \cos^{\frac{5}{2}}(a + bx) E\left(\frac{1}{2}(a + bx) \mid 2\right) + 7 \sin(a + bx) + 3 \sin(3(a + bx)) \right)}{10b}$$

input `Integrate[Sec[a + b*x]^(7/2),x]`

output `(Sec[a + b*x]^(5/2)*(-12*Cos[a + b*x]^(5/2)*EllipticE[(a + b*x)/2, 2] + 7*Sin[a + b*x] + 3*Sin[3*(a + b*x)]))/(10*b)`

### 3.9.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$ , Rules used = {3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{7}{2}}(a+bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a+bx+\frac{\pi}{2}\right)^{7/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{5} \int \sec^{\frac{3}{2}}(a+bx) dx + \frac{2 \sin(a+bx) \sec^{\frac{5}{2}}(a+bx)}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} \int \csc\left(a+bx+\frac{\pi}{2}\right)^{3/2} dx + \frac{2 \sin(a+bx) \sec^{\frac{5}{2}}(a+bx)}{5b} \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{5} \left( \frac{2 \sin(a+bx) \sqrt{\sec(a+bx)}}{b} - \int \frac{1}{\sqrt{\sec(a+bx)}} dx \right) + \frac{2 \sin(a+bx) \sec^{\frac{5}{2}}(a+bx)}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} \left( \frac{2 \sin(a+bx) \sqrt{\sec(a+bx)}}{b} - \int \frac{1}{\sqrt{\csc\left(a+bx+\frac{\pi}{2}\right)}} dx \right) + \frac{2 \sin(a+bx) \sec^{\frac{5}{2}}(a+bx)}{5b} \\
 & \quad \downarrow \text{4258} \\
 & \frac{3}{5} \left( \frac{2 \sin(a+bx) \sqrt{\sec(a+bx)}}{b} - \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \int \sqrt{\cos(a+bx)} dx \right) + \\
 & \quad \frac{2 \sin(a+bx) \sec^{\frac{5}{2}}(a+bx)}{5b} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$



$$\frac{3}{5} \left( \frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} \int \sqrt{\sin\left(a + bx + \frac{\pi}{2}\right)} dx \right) +$$

$$\frac{2 \sin(a + bx) \sec^{\frac{5}{2}}(a + bx)}{5b}$$

↓ 3119

$$\frac{2 \sin(a + bx) \sec^{\frac{5}{2}}(a + bx)}{5b} +$$

$$\frac{3}{5} \left( \frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \frac{2 \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b} \right)$$

input `Int[Sec[a + b*x]^(7/2),x]`

output `(2*Sec[a + b*x]^(5/2)*Sin[a + b*x])/(5*b) + (3*((-2*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/b + (2*Sqrt[Sec[a + b*x]]*Sin[a + b*x])/b))/5`

### 3.9.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.9.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 357 vs. 2(97) = 194.

Time = 8.78 (sec) , antiderivative size = 358, normalized size of antiderivative = 4.21

method	result
default	$-\frac{2\sqrt{-\left(-2\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^2+1\right)\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^2\left(24\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^6\cos\left(\frac{bx}{2}+\frac{a}{2}\right)-12\text{EllipticE}\left(\cos\left(\frac{bx}{2}+\frac{a}{2}\right),\sqrt{2}\right)\sqrt{2\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1}\sqrt{\dots}}{\dots}$

input `int(sec(b*x+a)^(7/2),x,method=_RETURNVERBOSE)`

output

$$-\frac{2}{5}\frac{\left(-2\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2+1\right)\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\left(8\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^6-12\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^4+6\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-1\right)}{\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^3\left(24\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^6\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)-12\text{EllipticE}\left(\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right),2^{\frac{1}{2}}\right)\right)\left(2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-1\right)^{\frac{1}{2}}\left(\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\right)^{\frac{1}{2}}\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^4-24\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^4\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)+12\text{EllipticE}\left(\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right),2^{\frac{1}{2}}\right)\left(2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-1\right)^{\frac{1}{2}}\left(\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\right)^{\frac{1}{2}}\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2+8\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)-3\text{EllipticE}\left(\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right),2^{\frac{1}{2}}\right)\left(2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-1\right)^{\frac{1}{2}}\left(\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\right)^{\frac{1}{2}}\left(-2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^4+\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\right)^{\frac{1}{2}}}{\left(2\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-1\right)^{\frac{1}{2}}}\frac{1}{b}$$

### 3.9.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.29

$$\int \sec^{\frac{7}{2}}(a+bx) dx$$

$$= \frac{-3i\sqrt{2}\cos^2(bx+a)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))) + 3}{\dots}$$

input `integrate(sec(b*x+a)^(7/2),x, algorithm="fricas")`

output `1/5*(-3*I*sqrt(2)*cos(b*x + a)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + 3*I*sqrt(2)*cos(b*x + a)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) + 2*(3*cos(b*x + a)^2 + 1)*sin(b*x + a)/sqrt(cos(b*x + a))/(b*cos(b*x + a)^2)`

### 3.9.6 Sympy [F(-1)]

Timed out.

$$\int \sec^{\frac{7}{2}}(a + bx) dx = \text{Timed out}$$

input `integrate(sec(b*x+a)**(7/2),x)`

output `Timed out`

### 3.9.7 Maxima [F]

$$\int \sec^{\frac{7}{2}}(a + bx) dx = \int \sec(bx + a)^{\frac{7}{2}} dx$$

input `integrate(sec(b*x+a)^(7/2),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(7/2), x)`

### 3.9.8 Giac [F]

$$\int \sec^{\frac{7}{2}}(a + bx) dx = \int \sec(bx + a)^{\frac{7}{2}} dx$$

input `integrate(sec(b*x+a)^(7/2),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(7/2), x)`

**3.9.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{7}{2}}(a + bx) dx = \int \left( \frac{1}{\cos(a + bx)} \right)^{7/2} dx$$

input `int((1/cos(a + b*x))^(7/2),x)`output `int((1/cos(a + b*x))^(7/2), x)`

### 3.10 $\int \sec^{\frac{5}{2}}(a + bx) dx$

3.10.1	Optimal result . . . . .	168
3.10.2	Mathematica [A] (verified) . . . . .	168
3.10.3	Rubi [A] (verified) . . . . .	169
3.10.4	Maple [B] (verified) . . . . .	170
3.10.5	Fricas [C] (verification not implemented) . . . . .	171
3.10.6	Sympy [F] . . . . .	171
3.10.7	Maxima [F] . . . . .	171
3.10.8	Giac [F] . . . . .	172
3.10.9	Mupad [F(-1)] . . . . .	172

#### 3.10.1 Optimal result

Integrand size = 10, antiderivative size = 62

$$\int \sec^{\frac{5}{2}}(a + bx) dx = \frac{2\sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{\sec(a + bx)}}{3b} + \frac{2 \sec^{\frac{3}{2}}(a + bx) \sin(a + bx)}{3b}$$

output `2/3*sec(b*x+a)^(3/2)*sin(b*x+a)/b+2/3*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.10.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

$$\int \sec^{\frac{5}{2}}(a + bx) dx = \frac{2 \sec^{\frac{3}{2}}(a + bx) \left( \cos^{\frac{3}{2}}(a + bx) \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) + \sin(a + bx) \right)}{3b}$$

input `Integrate[Sec[a + b*x]^(5/2),x]`

output `(2*Sec[a + b*x]^(3/2)*(Cos[a + b*x]^(3/2)*EllipticF[(a + b*x)/2, 2] + Sin[a + b*x]))/(3*b)`

### 3.10.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{5}{2}}(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^{\frac{5}{2}} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{3} \int \sqrt{\sec(a + bx)} dx + \frac{2 \sin(a + bx) \sec^{\frac{3}{2}}(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \sqrt{\csc\left(a + bx + \frac{\pi}{2}\right)} dx + \frac{2 \sin(a + bx) \sec^{\frac{3}{2}}(a + bx)}{3b} \\
 & \quad \downarrow \text{4258} \\
 & \frac{1}{3} \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} \int \frac{1}{\sqrt{\cos(a + bx)}} dx + \frac{2 \sin(a + bx) \sec^{\frac{3}{2}}(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} \int \frac{1}{\sqrt{\sin\left(a + bx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(a + bx) \sec^{\frac{3}{2}}(a + bx)}{3b} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2 \sin(a + bx) \sec^{\frac{3}{2}}(a + bx)}{3b} + \frac{2 \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} \text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{3b}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(5/2), x]`

output `(2*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*sqrt[Sec[a + b*x]])/(3*b) + (2*Sec[a + b*x]^(3/2)*Sin[a + b*x])/(3*b)`

## 3.10.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.10.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs.  $2(78) = 156$ .

Time = 4.51 (sec) , antiderivative size = 213, normalized size of antiderivative = 3.44

method	result
default	$\frac{2 \left( -2\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1} \operatorname{EllipticF}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right) \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 2 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2 \cos\left(\frac{bx}{2} + \frac{a}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \right)}{3 \sqrt{-2 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2} \left( 2 \cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1 \right)}$

input `int(sec(b*x+a)^(5/2), x, method=_RETURNVERBOSE)`

output `-2/3*(-2*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))*sin(1/2*b*x+1/2*a)^2-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2)))*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/(2*cos(1/2*b*x+1/2*a)^2-1)^(3/2)/sin(1/2*b*x+1/2*a)/b`

### 3.10.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.42

$$\int \sec^{\frac{5}{2}}(a + bx) dx$$

$$= \frac{-i \sqrt{2} \cos(bx + a) \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i \sqrt{2} \cos(bx + a) \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)) + 2 \sin(bx + a) / \sqrt{\cos(bx + a)}}{3 b \cos(bx + a)}$$

input `integrate(sec(b*x+a)^(5/2),x, algorithm="fricas")`

output `1/3*(-I*sqrt(2)*cos(b*x + a)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*cos(b*x + a)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*sin(b*x + a)/sqrt(cos(b*x + a)))/(b*cos(b*x + a))`

### 3.10.6 Sympy [F]

$$\int \sec^{\frac{5}{2}}(a + bx) dx = \int \sec^{\frac{5}{2}}(a + bx) dx$$

input `integrate(sec(b*x+a)**(5/2),x)`

output `Integral(sec(a + b*x)**(5/2), x)`

### 3.10.7 Maxima [F]

$$\int \sec^{\frac{5}{2}}(a + bx) dx = \int \sec(bx + a)^{\frac{5}{2}} dx$$

input `integrate(sec(b*x+a)^(5/2),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(5/2), x)`



**3.10.8 Giac [F]**

$$\int \sec^{\frac{5}{2}}(a + bx) dx = \int \sec (bx + a)^{\frac{5}{2}} dx$$

input `integrate(sec(b*x+a)^(5/2),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(5/2), x)`

**3.10.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{5}{2}}(a + bx) dx = \int \left( \frac{1}{\cos(a + bx)} \right)^{5/2} dx$$

input `int((1/cos(a + b*x))^(5/2),x)`

output `int((1/cos(a + b*x))^(5/2), x)`

### 3.11 $\int \sec^{\frac{3}{2}}(a + bx) dx$

3.11.1	Optimal result . . . . .	173
3.11.2	Mathematica [A] (verified) . . . . .	173
3.11.3	Rubi [A] (verified) . . . . .	174
3.11.4	Maple [B] (verified) . . . . .	175
3.11.5	Fricas [C] (verification not implemented) . . . . .	176
3.11.6	Sympy [F] . . . . .	176
3.11.7	Maxima [F] . . . . .	176
3.11.8	Giac [F] . . . . .	177
3.11.9	Mupad [F(-1)] . . . . .	177

#### 3.11.1 Optimal result

Integrand size = 10, antiderivative size = 58

$$\int \sec^{\frac{3}{2}}(a + bx) dx = -\frac{2\sqrt{\cos(a + bx)}E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{\sec(a + bx)}}{b} + \frac{2\sqrt{\sec(a + bx)} \sin(a + bx)}{b}$$

output `2*sin(b*x+a)*sec(b*x+a)^(1/2)/b-2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.11.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.78

$$\int \sec^{\frac{3}{2}}(a + bx) dx = \frac{2\sqrt{\sec(a + bx)}\left(-\sqrt{\cos(a + bx)}E\left(\frac{1}{2}(a + bx) \mid 2\right) + \sin(a + bx)\right)}{b}$$

input `Integrate[Sec[a + b*x]^(3/2),x]`

output `(2*Sqrt[Sec[a + b*x]]*(-(Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]) + Sin[a + b*x]))/b`

### 3.11.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{3}{2}}(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^{3/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \int \frac{1}{\sqrt{\sec(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \int \frac{1}{\sqrt{\csc\left(a + bx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} \int \sqrt{\cos(a + bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} \int \sqrt{\sin\left(a + bx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3119} \\
 & \frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \frac{2 \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(3/2),x]`

output `(-2*sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]*sqrt[Sec[a + b*x]])/b + (2*sqrt[Sec[a + b*x]]*Sin[a + b*x])/b`

## 3.11.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^n, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.11.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 181 vs. 2(78) = 156.

Time = 6.00 (sec) , antiderivative size = 182, normalized size of antiderivative = 3.14

method	result
default	$- \frac{2 \left( -2 \sqrt{-2 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2} \cos\left(\frac{bx}{2} + \frac{a}{2}\right) \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1} \sqrt{-2 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2} \right)}{\sqrt{-2 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2} \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{2 \cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1} b}$

input `int(sec(b*x+a)^(3/2), x, method=_RETURNVERBOSE)`

output `-2*(-2*(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^2+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2)))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b`

### 3.11.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.26

$$\int \sec^{\frac{3}{2}}(a + bx) dx$$

$$= \frac{-i \sqrt{2} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) + i \sqrt{2} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a))) + 2 \sin(bx + a) / \sqrt{\cos(bx + a)}}{b}$$

input `integrate(sec(b*x+a)^(3/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) + 2*sin(b*x + a)/sqrt(cos(b*x + a)))/b`

### 3.11.6 Sympy [F]

$$\int \sec^{\frac{3}{2}}(a + bx) dx = \int \sec^{\frac{3}{2}}(a + bx) dx$$

input `integrate(sec(b*x+a)**(3/2),x)`

output `Integral(sec(a + b*x)**(3/2), x)`

### 3.11.7 Maxima [F]

$$\int \sec^{\frac{3}{2}}(a + bx) dx = \int \sec(bx + a)^{\frac{3}{2}} dx$$

input `integrate(sec(b*x+a)^(3/2),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(3/2), x)`

**3.11.8 Giac [F]**

$$\int \sec^{\frac{3}{2}}(a + bx) dx = \int \sec(bx + a)^{\frac{3}{2}} dx$$

input `integrate(sec(b*x+a)^(3/2),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(3/2), x)`

**3.11.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{3}{2}}(a + bx) dx = \int \left( \frac{1}{\cos(a + bx)} \right)^{\frac{3}{2}} dx$$

input `int((1/cos(a + b*x))^(3/2),x)`

output `int((1/cos(a + b*x))^(3/2), x)`

### 3.12 $\int \sqrt{\sec(a + bx)} dx$

3.12.1	Optimal result . . . . .	178
3.12.2	Mathematica [A] (verified) . . . . .	178
3.12.3	Rubi [A] (verified) . . . . .	179
3.12.4	Maple [B] (verified) . . . . .	180
3.12.5	Fricas [C] (verification not implemented) . . . . .	180
3.12.6	Sympy [F] . . . . .	181
3.12.7	Maxima [F] . . . . .	181
3.12.8	Giac [F] . . . . .	181
3.12.9	Mupad [B] (verification not implemented) . . . . .	182

#### 3.12.1 Optimal result

Integrand size = 10, antiderivative size = 36

$$\int \sqrt{\sec(a + bx)} dx = \frac{2\sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{\sec(a + bx)}}{b}$$

output `2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x), 2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.12.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \sqrt{\sec(a + bx)} dx = \frac{2\sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{\sec(a + bx)}}{b}$$

input `Integrate[Sqrt[Sec[a + b*x]], x]`

output `(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/b`

### 3.12.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc\left(a+bx+\frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4258} \\
 & \sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)} \int \frac{1}{\sqrt{\sin\left(a+bx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3120} \\
 & \frac{2\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b}
 \end{aligned}$$

input `Int[Sqrt[Sec[a + b*x]],x]`

output `(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/b`

#### 3.12.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`



rule 4258 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.12.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. 2(58) = 116.

Time = 3.86 (sec) , antiderivative size = 133, normalized size of antiderivative = 3.69

method	result	size
default	$-\frac{2\sqrt{\left(2\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1\right)\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^2}\sqrt{\frac{1}{2}-\frac{\cos(bx+a)}{2}}\sqrt{-2\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^2+1}\operatorname{EllipticF}\left(\cos\left(\frac{bx}{2}+\frac{a}{2}\right),\sqrt{2}\right)}{\sqrt{-2\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^4+\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^2}\sin\left(\frac{bx}{2}+\frac{a}{2}\right)\sqrt{2\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1}b}$	133

input `int(sec(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b`

### 3.12.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.42

$$\int \sqrt{\sec(a + bx)} dx = \frac{-i\sqrt{2}\operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i\sin(bx + a)) + i\sqrt{2}\operatorname{weierstrassPInverse}(-4, 0, \cos(bx - a) + i\sin(bx - a))}{b}$$

input `integrate(sec(b*x+a)^(1/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/b`

**3.12.6 Sympy [F]**

$$\int \sqrt{\sec(a + bx)} dx = \int \sqrt{\sec(a + bx)} dx$$

input `integrate(sec(b*x+a)**(1/2),x)`

output `Integral(sqrt(sec(a + b*x)), x)`

**3.12.7 Maxima [F]**

$$\int \sqrt{\sec(a + bx)} dx = \int \sqrt{\sec(bx + a)} dx$$

input `integrate(sec(b*x+a)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(sec(b*x + a)), x)`

**3.12.8 Giac [F]**

$$\int \sqrt{\sec(a + bx)} dx = \int \sqrt{\sec(bx + a)} dx$$

input `integrate(sec(b*x+a)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(sec(b*x + a)), x)`

**3.12.9 Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \sqrt{\sec(a + bx)} dx = \frac{2 \sqrt{\cos(a + bx)} \sqrt{\frac{1}{\cos(a + bx)}} F\left(\frac{a}{2} + \frac{bx}{2} \mid 2\right)}{b}$$

input `int((1/cos(a + b*x))^(1/2),x)`

output `(2*cos(a + b*x)^(1/2)*(1/cos(a + b*x))^(1/2)*ellipticF(a/2 + (b*x)/2, 2))/b`

### 3.13 $\int \frac{1}{\sqrt{\sec(a+bx)}} dx$

3.13.1	Optimal result	183
3.13.2	Mathematica [A] (verified)	183
3.13.3	Rubi [A] (verified)	184
3.13.4	Maple [B] (verified)	185
3.13.5	Fricas [C] (verification not implemented)	185
3.13.6	Sympy [F]	186
3.13.7	Maxima [F]	186
3.13.8	Giac [F]	186
3.13.9	Mupad [F(-1)]	187

#### 3.13.1 Optimal result

Integrand size = 10, antiderivative size = 36

$$\int \frac{1}{\sqrt{\sec(a+bx)}} dx = \frac{2\sqrt{\cos(a+bx)}E\left(\frac{1}{2}(a+bx)|2\right)\sqrt{\sec(a+bx)}}{b}$$

output `2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.13.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{\sec(a+bx)}} dx = \frac{2E\left(\frac{1}{2}(a+bx)|2\right)}{b\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)}}$$

input `Integrate[1/Sqrt[Sec[a + b*x]],x]`

output `(2*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]]*Sqrt[Sec[a + b*x]])`

### 3.13.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\sec(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{\csc(a+bx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4258} \\
 & \sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)} \int \sqrt{\cos(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)} \int \sqrt{\sin(a+bx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3119} \\
 & \frac{2\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)}E(\frac{1}{2}(a+bx)|2)}{b}
 \end{aligned}$$

input `Int[1/Sqrt[Sec[a + b*x]],x]`

output `(2*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/b`

#### 3.13.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

---

3.13.  $\int \frac{1}{\sqrt{\sec(a+bx)}} dx$

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_], x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.13.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. 2(58) = 116.

Time = 4.81 (sec) , antiderivative size = 133, normalized size of antiderivative = 3.69

method	result
default	$\frac{2\sqrt{\left(2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1\right)\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2}\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{-2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + 1}\operatorname{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right)}{\sqrt{-2\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1}b}$
risch	$-\frac{i\sqrt{2}}{b\sqrt{\frac{e^{i(bx+a)}}{e^{2i(bx+a)}+1}}} - \frac{i\left(-\frac{2(e^{2i(bx+a)}+1)}{\sqrt{e^{i(bx+a)}(e^{2i(bx+a)}+1)}} + \frac{i\sqrt{-i(e^{i(bx+a)}+i)}\sqrt{2}\sqrt{i(e^{i(bx+a)}-i)}\sqrt{ie^{i(bx+a)}}}{\sqrt{e^{3i(bx+a)}+e^{i(bx+a)}}}\right)\operatorname{EllipticE}\left(\sqrt{-i(e^{i(bx+a)}+i)}, \sqrt{2}\right)}{b\sqrt{\frac{e^{i(bx+a)}}{e^{2i(bx+a)}+1}}(e^{2i(bx+a)}+1)}$

input `int(1/sec(b*x+a)^(1/2), x, method=_RETURNVERBOSE)`

output `2*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b`

### 3.13.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.58

$$\int \frac{1}{\sqrt{\sec(a+bx)}} dx = \frac{i\sqrt{2}\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + i\sin(bx+a))) - i\sqrt{2}\operatorname{weierstrassZeta}(-4, 0, \cos(bx+a) - i\sin(bx+a))}{b}$$

input `integrate(1/sec(b*x+a)^(1/2), x, algorithm="fracas")`

output `(I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))))/b`

### 3.13.6 Sympy [F]

$$\int \frac{1}{\sqrt{\sec(a + bx)}} dx = \int \frac{1}{\sqrt{\sec(a + bx)}} dx$$

input `integrate(1/sec(b*x+a)**(1/2), x)`

output `Integral(1/sqrt(sec(a + b*x)), x)`

### 3.13.7 Maxima [F]

$$\int \frac{1}{\sqrt{\sec(a + bx)}} dx = \int \frac{1}{\sqrt{\sec(bx + a)}} dx$$

input `integrate(1/sec(b*x+a)^(1/2), x, algorithm="maxima")`

output `integrate(1/sqrt(sec(b*x + a)), x)`

### 3.13.8 Giac [F]

$$\int \frac{1}{\sqrt{\sec(a + bx)}} dx = \int \frac{1}{\sqrt{\sec(bx + a)}} dx$$

input `integrate(1/sec(b*x+a)^(1/2), x, algorithm="giac")`

output `integrate(1/sqrt(sec(b*x + a)), x)`

**3.13.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{\sec(a+bx)}} dx = \int \frac{1}{\sqrt{\frac{1}{\cos(a+bx)}}} dx$$

input `int(1/(1/cos(a + b*x))^(1/2),x)`output `int(1/(1/cos(a + b*x))^(1/2), x)`



### 3.14 $\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx$

3.14.1	Optimal result . . . . .	188
3.14.2	Mathematica [A] (verified) . . . . .	188
3.14.3	Rubi [A] (verified) . . . . .	189
3.14.4	Maple [B] (verified) . . . . .	190
3.14.5	Fricas [C] (verification not implemented) . . . . .	191
3.14.6	Sympy [F] . . . . .	191
3.14.7	Maxima [F] . . . . .	191
3.14.8	Giac [F] . . . . .	192
3.14.9	Mupad [F(-1)] . . . . .	192

#### 3.14.1 Optimal result

Integrand size = 10, antiderivative size = 62

$$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx = \frac{2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) \sqrt{\sec(a+bx)}}{3b} + \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}}$$

output `2/3*sin(b*x+a)/b/sec(b*x+a)^(1/2)+2/3*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.14.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx = \frac{\sqrt{\sec(a+bx)} \left( 2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) + \sin(2(a+bx)) \right)}{3b}$$

input `Integrate[Sec[a + b*x]^(-3/2), x]`

output `(Sqrt[Sec[a + b*x]]*(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Sin[2*(a + b*x)]))/(3*b)`

### 3.14.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(a+bx+\frac{\pi}{2})^{3/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{1}{3} \int \sqrt{\sec(a+bx)} dx + \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \sqrt{\csc(a+bx+\frac{\pi}{2})} dx + \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{1}{3} \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx + \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \int \frac{1}{\sqrt{\sin(a+bx+\frac{\pi}{2})}} dx + \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} + \frac{2\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3b}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(-3/2), x]`

output `(2*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*sqrt[Sec[a + b*x]])/(3*b) + (2*Sin[a + b*x])/(3*b*sqrt[Sec[a + b*x]])`

---

3.14.  $\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx$

## 3.14.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.14.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 178 vs. 2(78) = 156.

Time = 6.18 (sec) , antiderivative size = 179, normalized size of antiderivative = 2.89

method	result
default	$\frac{2\sqrt{\left(2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1\right)\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2} \left(4\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 \cos\left(\frac{bx}{2} + \frac{a}{2}\right) - 2\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2 \cos\left(\frac{bx}{2} + \frac{a}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2\sin\left(\frac{bx}{2} + \frac{a}{2}\right)}\right)}{3\sqrt{-2\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2} \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1} b}$

input `int(1/sec(b*x+a)^(3/2),x,method=_RETURNVERBOSE)`

output `-2/3*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(4*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2)))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b`

**3.14.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.10

$$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx$$

$$= \frac{2\sqrt{\cos(bx+a)}\sin(bx+a) - i\sqrt{2}\text{weierstrassPInverse}(-4, 0, \cos(bx+a) + i\sin(bx+a)) + i\sqrt{2}\text{weierstrassPInverse}(-4, 0, \cos(bx+a) - i\sin(bx+a))}{3b}$$

input `integrate(1/sec(b*x+a)^(3/2),x, algorithm="fricas")`

output `1/3*(2*sqrt(cos(b*x + a))*sin(b*x + a) - I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/b`

**3.14.6 Sympy [F]**

$$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx$$

input `integrate(1/sec(b*x+a)**(3/2), x)`

output `Integral(sec(a + b*x)**(-3/2), x)`

**3.14.7 Maxima [F]**

$$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx = \int \frac{1}{\sec(bx+a)^{\frac{3}{2}}} dx$$

input `integrate(1/sec(b*x+a)^(3/2),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(-3/2), x)`

**3.14.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx = \int \frac{1}{\sec(bx+a)^{\frac{3}{2}}} dx$$

input `integrate(1/sec(b*x+a)^(3/2),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(-3/2), x)`

**3.14.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx = \int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{3/2}} dx$$

input `int(1/(1/cos(a + b*x))^(3/2),x)`

output `int(1/(1/cos(a + b*x))^(3/2), x)`

### 3.15 $\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$

3.15.1 Optimal result . . . . .	193
3.15.2 Mathematica [A] (verified) . . . . .	193
3.15.3 Rubi [A] (verified) . . . . .	194
3.15.4 Maple [B] (verified) . . . . .	195
3.15.5 Fricas [C] (verification not implemented) . . . . .	196
3.15.6 Sympy [F] . . . . .	196
3.15.7 Maxima [F] . . . . .	196
3.15.8 Giac [F] . . . . .	197
3.15.9 Mupad [F(-1)] . . . . .	197

#### 3.15.1 Optimal result

Integrand size = 10, antiderivative size = 62

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx = \frac{6\sqrt{\cos(a+bx)}E\left(\frac{1}{2}(a+bx)\middle|2\right)\sqrt{\sec(a+bx)}}{5b} + \frac{2\sin(a+bx)}{5b\sec^{\frac{3}{2}}(a+bx)}$$

output `2/5*sin(b*x+a)/b/sec(b*x+a)^(3/2)+6/5*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.15.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx = \frac{\sqrt{\sec(a+bx)}\left(12\sqrt{\cos(a+bx)}E\left(\frac{1}{2}(a+bx)\middle|2\right) + \sin(a+bx) + \sin(3(a+bx))\right)}{10b}$$

input `Integrate[Sec[a + b*x]^(-5/2),x]`

output `(Sqrt[Sec[a + b*x]]*(12*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + Sin[a + b*x] + Sin[3*(a + b*x)]))/(10*b)`

### 3.15.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(a+bx+\frac{\pi}{2})^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{3}{5} \int \frac{1}{\sqrt{\sec(a+bx)}} dx + \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} \int \frac{1}{\sqrt{\csc(a+bx+\frac{\pi}{2})}} dx + \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} \\
 & \quad \downarrow \text{4258} \\
 & \frac{3}{5} \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \int \sqrt{\cos(a+bx)} dx + \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \int \sqrt{\sin(a+bx+\frac{\pi}{2})} dx + \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} + \frac{6 \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} E(\frac{1}{2}(a+bx) | 2)}{5b}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(-5/2), x]`

output `(6*sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]*sqrt[Sec[a + b*x]])/(5*b) + (2*Sin[a + b*x])/(5*b*Sec[a + b*x]^(3/2))`

---

3.15.  $\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$

## 3.15.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.15.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(78) = 156.

Time = 6.00 (sec) , antiderivative size = 202, normalized size of antiderivative = 3.26

method	result
default	$\frac{2\sqrt{\left(2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2 - 1\right)\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2\left(-8\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^6\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + 8\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4\cos\left(\frac{bx}{2} + \frac{a}{2}\right) - 2\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{5\sqrt{-2\sin\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^2}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^2}}$

input `int(1/sec(b*x+a)^(5/2),x,method=_RETURNVERBOSE)`

output `-2/5*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(-8*sin(1/2*b*x+1/2*a)^6*cos(1/2*b*x+1/2*a)+8*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a)-3*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b`

---

3.15.  $\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$



### 3.15.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$$

$$= \frac{2 \cos(bx+a)^{\frac{3}{2}} \sin(bx+a) + 3i \sqrt{2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx+a) + i \sin(bx+a))) - 3i \sqrt{2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx+a) - i \sin(bx+a)))}{5b}$$

input `integrate(1/sec(b*x+a)^(5/2),x, algorithm="fricas")`

output `1/5*(2*cos(b*x + a)^(3/2)*sin(b*x + a) + 3*I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - 3*I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))))/b`

### 3.15.6 Sympy [F]

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$$

input `integrate(1/sec(b*x+a)**(5/2),x)`

output `Integral(sec(a + b*x)**(-5/2), x)`

### 3.15.7 Maxima [F]

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx = \int \frac{1}{\sec(bx+a)^{\frac{5}{2}}} dx$$

input `integrate(1/sec(b*x+a)^(5/2),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(-5/2), x)`

---

3.15.  $\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$

**3.15.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{5}{2}}(bx+a)} dx$$

input `integrate(1/sec(b*x+a)^(5/2),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(-5/2), x)`

**3.15.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx = \int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{5}{2}}} dx$$

input `int(1/(1/cos(a + b*x))^(5/2),x)`

output `int(1/(1/cos(a + b*x))^(5/2), x)`

### 3.16 $\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx$

3.16.1	Optimal result	198
3.16.2	Mathematica [A] (verified)	198
3.16.3	Rubi [A] (verified)	199
3.16.4	Maple [B] (verified)	201
3.16.5	Fricas [C] (verification not implemented)	201
3.16.6	Sympy [F]	202
3.16.7	Maxima [F]	202
3.16.8	Giac [F]	202
3.16.9	Mupad [F(-1)]	203

#### 3.16.1 Optimal result

Integrand size = 10, antiderivative size = 85

$$\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx = \frac{10\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) \sqrt{\sec(a+bx)}}{21b} + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{10 \sin(a+bx)}{21b \sqrt{\sec(a+bx)}}$$

output `2/7*sin(b*x+a)/b/sec(b*x+a)^(5/2)+10/21*sin(b*x+a)/b/sec(b*x+a)^(1/2)+10/21*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b`

#### 3.16.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.72

$$\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx = \frac{\sqrt{\sec(a+bx)} \left( 40\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) + 26 \sin(2(a+bx)) + 3 \sin(4(a+bx)) \right)}{84b}$$

input `Integrate[Sec[a + b*x]^(-7/2), x]`

output  $(\text{Sqrt}[\text{Sec}[a + b*x]]*(40*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2] + 26*\text{Sin}[2*(a + b*x)] + 3*\text{Sin}[4*(a + b*x)]))/(84*b)$

### 3.16.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$ , Rules used = {3042, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc\left(a+bx+\frac{\pi}{2}\right)^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{5}{7} \int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{7} \int \frac{1}{\csc\left(a+bx+\frac{\pi}{2}\right)^{3/2}} dx + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} \\
 & \quad \downarrow \text{4256} \\
 & \frac{5}{7} \left( \frac{1}{3} \int \sqrt{\sec(a+bx)} dx + \frac{2 \sin(a+bx)}{3b \sqrt{\sec(a+bx)}} \right) + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{7} \left( \frac{1}{3} \int \sqrt{\csc\left(a+bx+\frac{\pi}{2}\right)} dx + \frac{2 \sin(a+bx)}{3b \sqrt{\sec(a+bx)}} \right) + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} \\
 & \quad \downarrow \text{4258} \\
 & \frac{5}{7} \left( \frac{1}{3} \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx + \frac{2 \sin(a+bx)}{3b \sqrt{\sec(a+bx)}} \right) + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{5}{7} \left( \frac{1}{3} \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \int \frac{1}{\sqrt{\sin(a+bx + \frac{\pi}{2})}} dx + \frac{2 \sin(a+bx)}{3b \sqrt{\sec(a+bx)}} \right) + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)}$$

↓ 3120

$$\frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{5}{7} \left( \frac{2 \sin(a+bx)}{3b \sqrt{\sec(a+bx)}} + \frac{2 \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3b} \right)$$

input `Int[Sec[a + b*x]^(-7/2), x]`

output `(2*Sin[a + b*x])/(7*b*Sec[a + b*x]^(5/2)) + (5*((2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/(3*b) + (2*Sin[a + b*x])/(3*b*Sqrt[Sec[a + b*x]]))/7`

### 3.16.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.16.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. 2(97) = 194.

Time = 6.20 (sec) , antiderivative size = 199, normalized size of antiderivative = 2.34

method	result
default	$\frac{2\sqrt{\left(2\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^2-1\right)\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^2\left(48\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^9-120\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^7+128\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^5-72\cos\left(\frac{bx}{2}+\frac{a}{2}\right)^3+5\sqrt{\frac{1}{2}-\cos\left(\frac{bx}{2}+\frac{a}{2}\right)}\right)}{21\sqrt{-2\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^4+\sin\left(\frac{bx}{2}+\frac{a}{2}\right)^2}\sin\left(\frac{bx}{2}+\frac{a}{2}\right)\sqrt{2\cos\left(\frac{bx}{2}+\frac{a}{2}\right)}}$

input `int(1/sec(b*x+a)^(7/2),x,method=_RETURNVERBOSE)`

output `-2/21*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(48*cos(1/2*b*x+1/2*a)^9-120*cos(1/2*b*x+1/2*a)^7+128*cos(1/2*b*x+1/2*a)^5-72*cos(1/2*b*x+1/2*a)^3+5*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))+16*cos(1/2*b*x+1/2*a))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b`

### 3.16.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.02

$$\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx$$

$$= \frac{2\left(3\cos(bx+a)^3+5\cos(bx+a)\right)\sin(bx+a)}{\sqrt{\cos(bx+a)}} - \frac{5i\sqrt{2}\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))+5i\sqrt{2}}{21b}$$

input `integrate(1/sec(b*x+a)^(7/2),x,algorithm="fracas")`

output `1/21*(2*(3*cos(b*x + a)^3 + 5*cos(b*x + a))*sin(b*x + a)/sqrt(cos(b*x + a)) - 5*I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + 5*I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/b`

**3.16.6 Sympy [F]**

$$\int \frac{1}{\sec^{\frac{7}{2}}(a + bx)} dx = \int \frac{1}{\sec^{\frac{7}{2}}(a + bx)} dx$$

input `integrate(1/sec(b*x+a)**(7/2), x)`

output `Integral(sec(a + b*x)**(-7/2), x)`

**3.16.7 Maxima [F]**

$$\int \frac{1}{\sec^{\frac{7}{2}}(a + bx)} dx = \int \frac{1}{\sec^{\frac{7}{2}}(bx + a)} dx$$

input `integrate(1/sec(b*x+a)^(7/2), x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(-7/2), x)`

**3.16.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{7}{2}}(a + bx)} dx = \int \frac{1}{\sec^{\frac{7}{2}}(bx + a)} dx$$

input `integrate(1/sec(b*x+a)^(7/2), x, algorithm="giac")`

output `integrate(sec(b*x + a)^(-7/2), x)`

**3.16.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx = \int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{7/2}} dx$$

input `int(1/(1/cos(a + b*x))^(7/2),x)`output `int(1/(1/cos(a + b*x))^(7/2), x)`



### 3.17 $\int (c \sec(a + bx))^{7/2} dx$

3.17.1	Optimal result	204
3.17.2	Mathematica [A] (verified)	204
3.17.3	Rubi [A] (verified)	205
3.17.4	Maple [C] (verified)	207
3.17.5	Fricas [C] (verification not implemented)	207
3.17.6	Sympy [F(-1)]	208
3.17.7	Maxima [F]	208
3.17.8	Giac [F]	208
3.17.9	Mupad [F(-1)]	209

#### 3.17.1 Optimal result

Integrand size = 12, antiderivative size = 98

$$\int (c \sec(a + bx))^{7/2} dx = -\frac{6c^4 E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{6c^3 \sqrt{c \sec(a + bx)} \sin(a + bx)}{5b} + \frac{2c(c \sec(a + bx))^{5/2} \sin(a + bx)}{5b}$$

output  $2/5*c*(c*\sec(b*x+a))^(5/2)*\sin(b*x+a)/b-6/5*c^4*(\cos(1/2*a+1/2*b*x)^2)^(1/2)/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^(1/2))/b/\cos(b*x+a)^(1/2)/(c*\sec(b*x+a))^(1/2)+6/5*c^3*\sin(b*x+a)*(c*\sec(b*x+a))^(1/2)/b$

#### 3.17.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.63

$$\int (c \sec(a + bx))^{7/2} dx = \frac{c(c \sec(a + bx))^{5/2} \left( -12 \cos^{5/2}(a + bx) E\left(\frac{1}{2}(a + bx) \mid 2\right) + 7 \sin(a + bx) + 3 \sin(3(a + bx)) \right)}{10b}$$

input `Integrate[(c*Sec[a + b*x])^(7/2),x]`

output  $(c*(c*\text{Sec}[a + b*x])^(5/2)*(-12*\text{Cos}[a + b*x]^(5/2)*\text{EllipticE}[(a + b*x)/2, 2] + 7*\text{Sin}[a + b*x] + 3*\text{Sin}[3*(a + b*x)]))/(10*b)$

**3.17.3 Rubi [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( c \csc \left( a + bx + \frac{\pi}{2} \right) \right)^{7/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{5} c^2 \int (c \sec(a + bx))^{3/2} dx + \frac{2c \sin(a + bx)(c \sec(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} c^2 \int \left( c \csc \left( a + bx + \frac{\pi}{2} \right) \right)^{3/2} dx + \frac{2c \sin(a + bx)(c \sec(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{5} c^2 \left( \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - c^2 \int \frac{1}{\sqrt{c \sec(a + bx)}} dx \right) + \frac{2c \sin(a + bx)(c \sec(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} c^2 \left( \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - c^2 \int \frac{1}{\sqrt{c \csc \left( a + bx + \frac{\pi}{2} \right)}} dx \right) + \\
 & \quad \frac{2c \sin(a + bx)(c \sec(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{4258} \\
 & \frac{3}{5} c^2 \left( \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - \frac{c^2 \int \sqrt{\cos(a + bx)} dx}{\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} \right) + \\
 & \quad \frac{2c \sin(a + bx)(c \sec(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{3}{5}c^2 \left( \frac{2c \sin(a+bx) \sqrt{c \sec(a+bx)}}{b} - \frac{c^2 \int \sqrt{\sin(a+bx + \frac{\pi}{2})} dx}{\sqrt{\cos(a+bx)} \sqrt{c \sec(a+bx)}} \right) + \frac{2c \sin(a+bx) (c \sec(a+bx))^{5/2}}{5b}$$

↓ 3119

$$\frac{3}{5}c^2 \left( \frac{2c \sin(a+bx) \sqrt{c \sec(a+bx)}}{b} - \frac{2c^2 E(\frac{1}{2}(a+bx)|2)}{b \sqrt{\cos(a+bx)} \sqrt{c \sec(a+bx)}} \right) + \frac{2c \sin(a+bx) (c \sec(a+bx))^{5/2}}{5b}$$

input `Int[(c*Sec[a + b*x])^(7/2),x]`

output `(2*c*(c*Sec[a + b*x])^(5/2)*Sin[a + b*x])/(5*b) + (3*c^2*((-2*c^2*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]]*Sqrt[c*Sec[a + b*x]]) + (2*c*Sqrt[c*Sec[a + b*x]]*Sin[a + b*x])/b))/5`

### 3.17.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.17.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 13.58 (sec) , antiderivative size = 415, normalized size of antiderivative = 4.23

method	result
default	$2\sqrt{c\sec(bx+a)}c^3\left(-3i\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\operatorname{EllipticF}(i(\cot(bx+a)-\csc(bx+a)),i)\cos(bx+a)^2+3i\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\right)$

input `int((c*sec(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 2/5/b*(c*\sec(b*x+a))^{(1/2)}*c^3/(\cos(b*x+a)+1)*(-3*I*(1/(\cos(b*x+a)+1))^{(1/2)} \\ & *(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}*\operatorname{EllipticF}(I*(\cot(b*x+a)-\csc(b*x+a)),I) \\ & )*\cos(b*x+a)^2+3*I*(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)} \\ & *\operatorname{EllipticE}(I*(\cot(b*x+a)-\csc(b*x+a)),I)*\cos(b*x+a)^2-6*I*(1/(\cos(b*x+a)+1))^{(1/2)} \\ & *(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}*\operatorname{EllipticF}(I*(\cot(b*x+a)-\csc(b*x+a)),I) \\ & )*\cos(b*x+a)+6*I*(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)} \\ & *\operatorname{EllipticE}(I*(\cot(b*x+a)-\csc(b*x+a)),I)*\cos(b*x+a)-3*I*\operatorname{EllipticF}(I*(\cot(b*x+a)-\csc(b*x+a)),I) \\ & *(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}+3*I*\operatorname{EllipticE}(I*(\cot(b*x+a)-\csc(b*x+a)),I) \\ & *(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}+3*\sin(b*x+a)+\tan(b*x+a)+\sec(b*x+a) \\ & )*\tan(b*x+a) \end{aligned}$$

### 3.17.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.28

$$\int (c\sec(a+bx))^{7/2} dx = \frac{-3i\sqrt{2}c^{7/2}\cos(bx+a)^2\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))^{7/2}}{\dots}$$

input `integrate((c*sec(b*x+a))^(7/2),x, algorithm="fricas")`

output `1/5*(-3*I*sqrt(2)*c^(7/2)*cos(b*x + a)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + 3*I*sqrt(2)*c^(7/2)*cos(b*x + a)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) + 2*(3*c^3*cos(b*x + a)^2 + c^3)*sqrt(c/cos(b*x + a))*sin(b*x + a)/(b*cos(b*x + a)^2)`

### 3.17.6 Sympy [F(-1)]

Timed out.

$$\int (c \sec(a + bx))^{7/2} dx = \text{Timed out}$$

input `integrate((c*sec(b*x+a))**(7/2),x)`

output `Timed out`

### 3.17.7 Maxima [F]

$$\int (c \sec(a + bx))^{7/2} dx = \int (c \sec(bx + a))^{\frac{7}{2}} dx$$

input `integrate((c*sec(b*x+a))^(7/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(7/2), x)`

### 3.17.8 Giac [F]

$$\int (c \sec(a + bx))^{7/2} dx = \int (c \sec(bx + a))^{\frac{7}{2}} dx$$

input `integrate((c*sec(b*x+a))^(7/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(7/2), x)`

**3.17.9 Mupad [F(-1)]**

Timed out.

$$\int (c \sec(a + bx))^{7/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{7/2} dx$$

input `int((c/cos(a + b*x))^(7/2),x)`output `int((c/cos(a + b*x))^(7/2), x)`

### 3.18 $\int (c \sec(a + bx))^{5/2} dx$

3.18.1	Optimal result . . . . .	210
3.18.2	Mathematica [A] (verified) . . . . .	210
3.18.3	Rubi [A] (verified) . . . . .	211
3.18.4	Maple [C] (verified) . . . . .	212
3.18.5	Fricas [C] (verification not implemented) . . . . .	213
3.18.6	Sympy [F] . . . . .	213
3.18.7	Maxima [F] . . . . .	213
3.18.8	Giac [F] . . . . .	214
3.18.9	Mupad [F(-1)] . . . . .	214

#### 3.18.1 Optimal result

Integrand size = 12, antiderivative size = 70

$$\int (c \sec(a + bx))^{5/2} dx = \frac{2c^2 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{c \sec(a + bx)}}{3b} + \frac{2c(c \sec(a + bx))^{3/2} \sin(a + bx)}{3b}$$

output `2/3*c*(c*sec(b*x+a))^(3/2)*sin(b*x+a)/b+2/3*c^2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*(c*sec(b*x+a))^(1/2)/b`

#### 3.18.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.73

$$\int (c \sec(a + bx))^{5/2} dx = \frac{2c^2 \sqrt{c \sec(a + bx)} \left( \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) + \tan(a + bx) \right)}{3b}$$

input `Integrate[(c*Sec[a + b*x])^(5/2),x]`

output `(2*c^2*sqrt[c*Sec[a + b*x]]*(sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Tan[a + b*x]))/(3*b)`

### 3.18.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( c \csc \left( a + bx + \frac{\pi}{2} \right) \right)^{5/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{3} c^2 \int \sqrt{c \sec(a + bx)} dx + \frac{2c \sin(a + bx)(c \sec(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} c^2 \int \sqrt{c \csc \left( a + bx + \frac{\pi}{2} \right)} dx + \frac{2c \sin(a + bx)(c \sec(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{4258} \\
 & \frac{1}{3} c^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)} \int \frac{1}{\sqrt{\cos(a + bx)}} dx + \frac{2c \sin(a + bx)(c \sec(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} c^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)} \int \frac{1}{\sqrt{\sin \left( a + bx + \frac{\pi}{2} \right)}} dx + \frac{2c \sin(a + bx)(c \sec(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2c^2 \sqrt{\cos(a + bx)} \operatorname{EllipticF} \left( \frac{1}{2}(a + bx), 2 \right) \sqrt{c \sec(a + bx)}}{3b} + \frac{2c \sin(a + bx)(c \sec(a + bx))^{3/2}}{3b}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(5/2),x]`

output `(2*c^2*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*sqrt[c*Sec[a + b*x]])/(3*b) + (2*c*(c*Sec[a + b*x])^(3/2)*Sin[a + b*x])/(3*b)`



## 3.18.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.18.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.36 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.06

method	result
default	$-\frac{2\sqrt{c\sec(bx+a)}c^2\left(i\operatorname{EllipticF}\left(i(-\cot(bx+a)+\csc(bx+a)),i\right)\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\cos(bx+a)+i\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\right)}{3b}$

input `int((c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/3/b*(c*sec(b*x+a))^(1/2)*c^2*(I*EllipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*cos(b*x+a)+I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)-tan(b*x+a))`

### 3.18.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.44

$$\int (c \sec(a + bx))^{5/2} dx = \frac{-i \sqrt{2} c^{5/2} \cos(bx + a) \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i \sqrt{2} c^{5/2} \cos(bx + a) \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)) + 2c^2 \sqrt{c/\cos(bx + a)} \sin(bx + a)}{3b \cos(bx + a)}$$

input `integrate((c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

output `1/3*(-I*sqrt(2)*c^(5/2)*cos(b*x + a)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*c^(5/2)*cos(b*x + a)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*c^2*sqrt(c/cos(b*x + a))*sin(b*x + a))/(b*cos(b*x + a))`

### 3.18.6 Sympy [F]

$$\int (c \sec(a + bx))^{5/2} dx = \int (c \sec(a + bx))^{5/2} dx$$

input `integrate((c*sec(b*x+a))**(5/2),x)`

output `Integral((c*sec(a + b*x))**(5/2), x)`

### 3.18.7 Maxima [F]

$$\int (c \sec(a + bx))^{5/2} dx = \int (c \sec(bx + a))^{5/2} dx$$

input `integrate((c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(5/2), x)`

**3.18.8 Giac [F]**

$$\int (c \sec(a + bx))^{5/2} dx = \int (c \sec(bx + a))^{\frac{5}{2}} dx$$

input `integrate((c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(5/2), x)`

**3.18.9 Mupad [F(-1)]**

Timed out.

$$\int (c \sec(a + bx))^{5/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{5/2} dx$$

input `int((c/cos(a + b*x))^(5/2),x)`

output `int((c/cos(a + b*x))^(5/2), x)`

### 3.19 $\int (c \sec(a + bx))^{3/2} dx$

3.19.1	Optimal result . . . . .	215
3.19.2	Mathematica [A] (verified) . . . . .	215
3.19.3	Rubi [A] (verified) . . . . .	216
3.19.4	Maple [C] (verified) . . . . .	217
3.19.5	Fricas [C] (verification not implemented) . . . . .	218
3.19.6	Sympy [F] . . . . .	218
3.19.7	Maxima [F] . . . . .	219
3.19.8	Giac [F] . . . . .	219
3.19.9	Mupad [F(-1)] . . . . .	219

#### 3.19.1 Optimal result

Integrand size = 12, antiderivative size = 66

$$\int (c \sec(a + bx))^{3/2} dx = -\frac{2c^2 E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{2c \sqrt{c \sec(a + bx)} \sin(a + bx)}{b}$$

output `-2*c^2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))/b/cos(b*x+a)^(1/2)/(c*sec(b*x+a))^(1/2)+2*c*sin(b*x+a)*(c*sec(b*x+a))^(1/2)/b`

#### 3.19.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.73

$$\int (c \sec(a + bx))^{3/2} dx = \frac{2c \sqrt{c \sec(a + bx)} \left( -\sqrt{\cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right) + \sin(a + bx) \right)}{b}$$

input `Integrate[(c*Sec[a + b*x])^(3/2),x]`

output `(2*c*Sqrt[c*Sec[a + b*x]]*(-(Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]) + Sin[a + b*x]))/b`

### 3.19.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( c \csc \left( a + bx + \frac{\pi}{2} \right) \right)^{3/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - c^2 \int \frac{1}{\sqrt{c \sec(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - c^2 \int \frac{1}{\sqrt{c \csc(a + bx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - \frac{c^2 \int \sqrt{\cos(a + bx)} dx}{\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - \frac{c^2 \int \sqrt{\sin(a + bx + \frac{\pi}{2})} dx}{\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - \frac{2c^2 E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(3/2),x]`

output `(-2*c^2*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]]*Sqrt[c*Sec[a + b*x]]) + (2*c*Sqrt[c*Sec[a + b*x]]*Sin[a + b*x])/b`

## 3.19.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.19.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.92 (sec) , antiderivative size = 394, normalized size of antiderivative = 5.97

method	result
default	$-\frac{2 \left( i \operatorname{EllipticE} \left( i \left( -\cot (bx+a) + \csc (bx+a) \right), i \right) \sqrt{\frac{1}{\cos (bx+a)+1}} \sqrt{\frac{\cos (bx+a)}{\cos (bx+a)+1}} \cos (bx+a)^2 - i \operatorname{EllipticF} \left( i \left( -\cot (bx+a) + \csc (bx+a) \right), i \right) \right)}{\dots}$

input `int((c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -2/b*(I*EllipticE(I*(-cot(b*x+a)+csc(b*x+a)),I)*(1/(cos(b*x+a)+1))^(1/2)*( \\ & \cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*\cos(b*x+a)^2-I*EllipticF(I*(-cot(b*x+a)+ \\ & sc(b*x+a)),I)*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*c \\ & \cos(b*x+a)^2+2*I*EllipticE(I*(-cot(b*x+a)+csc(b*x+a)),I)*(1/(cos(b*x+a)+1)) \\ & ^{(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*\cos(b*x+a)-2*I*EllipticF(I*(-cot( \\ & b*x+a)+csc(b*x+a)),I)*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1)) \\ & ^{(1/2)*\cos(b*x+a)+I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^( \\ & 1/2)*EllipticE(I*(-cot(b*x+a)+csc(b*x+a)),I)-I*(1/(cos(b*x+a)+1))^(1/2)*(c \\ & \cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)-si \\ & n(b*x+a))*(c*sec(b*x+a))^(1/2)*c/(cos(b*x+a)+1) \end{aligned}$$

### 3.19.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.27

$$\int (c \sec(a + bx))^{3/2} dx = \frac{-i \sqrt{2} c^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) + i \sqrt{2} c^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)))}{2c \sqrt{c/\cos(bx + a) \sin(bx + a)}}$$

input `integrate((c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

output 
$$\begin{aligned} & (-I*\sqrt{2}*c^(3/2)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos( \\ & b*x + a) + I*\sin(b*x + a))) + I*\sqrt{2}*c^(3/2)*\text{weierstrassZeta}(-4, 0, \text{wei} \\ & erstrassPInverse(-4, 0, \cos(b*x + a) - I*\sin(b*x + a))) + 2*c*\sqrt{c/\cos(b \\ & *x + a)*\sin(b*x + a))/b \end{aligned}$$

### 3.19.6 Sympy [F]

$$\int (c \sec(a + bx))^{3/2} dx = \int (c \sec(a + bx))^{\frac{3}{2}} dx$$

input `integrate((c*sec(b*x+a))**(3/2),x)`

output `Integral((c*sec(a + b*x))**(3/2), x)`

**3.19.7 Maxima [F]**

$$\int (c \sec(a + bx))^{3/2} dx = \int (c \sec(bx + a))^{\frac{3}{2}} dx$$

input `integrate((c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(3/2), x)`

**3.19.8 Giac [F]**

$$\int (c \sec(a + bx))^{3/2} dx = \int (c \sec(bx + a))^{\frac{3}{2}} dx$$

input `integrate((c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(3/2), x)`

**3.19.9 Mupad [F(-1)]**

Timed out.

$$\int (c \sec(a + bx))^{3/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{3/2} dx$$

input `int((c/cos(a + b*x))^(3/2),x)`

output `int((c/cos(a + b*x))^(3/2), x)`



## 3.20 $\int \sqrt{c \sec(a + bx)} dx$

3.20.1	Optimal result . . . . .	220
3.20.2	Mathematica [A] (verified) . . . . .	220
3.20.3	Rubi [A] (verified) . . . . .	221
3.20.4	Maple [C] (verified) . . . . .	222
3.20.5	Fricas [C] (verification not implemented) . . . . .	222
3.20.6	Sympy [F] . . . . .	223
3.20.7	Maxima [F] . . . . .	223
3.20.8	Giac [F] . . . . .	223
3.20.9	Mupad [B] (verification not implemented) . . . . .	224

### 3.20.1 Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \sqrt{c \sec(a + bx)} dx = \frac{2\sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{c \sec(a + bx)}}{b}$$

output `2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x), 2^(1/2))*cos(b*x+a)^(1/2)*(c*sec(b*x+a))^(1/2)/b`

### 3.20.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \sqrt{c \sec(a + bx)} dx = \frac{2\sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{c \sec(a + bx)}}{b}$$

input `Integrate[Sqrt[c*Sec[a + b*x]], x]`

output `(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[c*Sec[a + b*x]])/b`

### 3.20.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{c \sec(a + bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{c \csc\left(a + bx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4258} \\
 & \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)} \int \frac{1}{\sqrt{\cos(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)} \int \frac{1}{\sqrt{\sin\left(a + bx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3120} \\
 & \frac{2\sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{c \sec(a + bx)}}{b}
 \end{aligned}$$

input `Int[Sqrt[c*Sec[a + b*x]],x]`

output `(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[c*Sec[a + b*x]])/b`

#### 3.20.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

### 3.20.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.87 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.03

method	result	size
default	$-\frac{2i(\cos(bx+a)+1)\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\text{EllipticF}(i(-\cot(bx+a)+\csc(bx+a)),i)\sqrt{c\sec(bx+a)}}{b}$	77

```
input int((c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2*I/b*(cos(b*x+a)+1)*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))
^(1/2)*EllipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)*(c*sec(b*x+a))^(1/2)
```

### 3.20.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.50

$$\int \sqrt{c \sec(a + bx)} dx$$

$$= \frac{-i\sqrt{2}\sqrt{c}\text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i\sqrt{2}\sqrt{c}\text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a))}{b}$$

```
input integrate((c*sec(b*x+a))^(1/2),x, algorithm="fricas")
```

```
output (-I*sqrt(2)*sqrt(c)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x +
a)) + I*sqrt(2)*sqrt(c)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*
x + a)))/b
```

**3.20.6 Sympy [F]**

$$\int \sqrt{c \sec(a + bx)} dx = \int \sqrt{c \sec(a + bx)} dx$$

input `integrate((c*sec(b*x+a))**(1/2),x)`

output `Integral(sqrt(c*sec(a + b*x)), x)`

**3.20.7 Maxima [F]**

$$\int \sqrt{c \sec(a + bx)} dx = \int \sqrt{c \sec(bx + a)} dx$$

input `integrate((c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(c*sec(b*x + a)), x)`

**3.20.8 Giac [F]**

$$\int \sqrt{c \sec(a + bx)} dx = \int \sqrt{c \sec(bx + a)} dx$$

input `integrate((c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(c*sec(b*x + a)), x)`

**3.20.9 Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

$$\int \sqrt{c \sec(a + bx)} dx = \frac{2 \sqrt{\cos(a + bx)} \sqrt{\frac{c}{\cos(a + bx)}} F\left(\frac{a}{2} + \frac{bx}{2} \mid 2\right)}{b}$$

input `int((c/cos(a + b*x))^(1/2),x)`

output `(2*cos(a + b*x)^(1/2)*(c/cos(a + b*x))^(1/2)*ellipticF(a/2 + (b*x)/2, 2))/b`

### 3.21 $\int \frac{1}{\sqrt{c \sec(a+bx)}} dx$

3.21.1	Optimal result	225
3.21.2	Mathematica [A] (verified)	225
3.21.3	Rubi [A] (verified)	226
3.21.4	Maple [C] (verified)	227
3.21.5	Fricas [C] (verification not implemented)	228
3.21.6	Sympy [F]	228
3.21.7	Maxima [F]	228
3.21.8	Giac [F]	229
3.21.9	Mupad [F(-1)]	229

#### 3.21.1 Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \frac{1}{\sqrt{c \sec(a+bx)}} dx = \frac{2E\left(\frac{1}{2}(a+bx) \mid 2\right)}{b\sqrt{\cos(a+bx)}\sqrt{c \sec(a+bx)}}$$

output `2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x), 2^(1/2))/b/cos(b*x+a)^(1/2)/(c*sec(b*x+a))^(1/2)`

#### 3.21.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{c \sec(a+bx)}} dx = \frac{2E\left(\frac{1}{2}(a+bx) \mid 2\right)}{b\sqrt{\cos(a+bx)}\sqrt{c \sec(a+bx)}}$$

input `Integrate[1/Sqrt[c*Sec[a + b*x]],x]`

output `(2*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]]*Sqrt[c*Sec[a + b*x]])`

### 3.21.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{c \sec(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{c \csc(a+bx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{\int \sqrt{\cos(a+bx)} dx}{\sqrt{\cos(a+bx)} \sqrt{c \sec(a+bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{\sin(a+bx+\frac{\pi}{2})} dx}{\sqrt{\cos(a+bx)} \sqrt{c \sec(a+bx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2E(\frac{1}{2}(a+bx)|2)}{b\sqrt{\cos(a+bx)}\sqrt{c \sec(a+bx)}}
 \end{aligned}$$

input `Int[1/Sqrt[c*Sec[a + b*x]],x]`

output `(2*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]]*Sqrt[c*Sec[a + b*x]])`

### 3.21.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.21.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.96 (sec) , antiderivative size = 299, normalized size of antiderivative = 7.87

method	result
risch	$-\frac{i\sqrt{2}}{b\sqrt{\frac{ce^{i(bx+a)}}{e^{2i(bx+a)}+1}}} - \frac{i\left(-\frac{2(c e^{2i(bx+a)}+c)}{c\sqrt{e^{i(bx+a)}(c e^{2i(bx+a)}+c)}} + \frac{i\sqrt{-i(e^{i(bx+a)}+i)}\sqrt{2}\sqrt{i(e^{i(bx+a)}-i)}\sqrt{ie^{i(bx+a)}}}{\sqrt{c e^{3i(bx+a)}+c e^{i(bx+a)}}}\right)}{b\sqrt{\frac{ce^{i(bx+a)}}{e^{2i(bx+a)}+1}}(e^{2i(bx+a)}+1)}$
default	$\frac{2i \operatorname{EllipticE}(i(-\cot(bx+a)+\csc(bx+a)), i)\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\cos(bx+a)-2i \operatorname{EllipticF}(i(-\cot(bx+a)+\csc(bx+a)), i)\sqrt{\frac{1}{\cos(bx+a)+1}}}{\dots}$

input `int(1/(c*sec(b*x+a))^(1/2), x, method=_RETURNVERBOSE)`

output `-I/b*2^(1/2)/(c*exp(I*(b*x+a))/(exp(I*(b*x+a))^2+1))^(1/2)-I/b*(-2*(c*exp(I*(b*x+a))^2+c)/c/(exp(I*(b*x+a))*(c*exp(I*(b*x+a))^2+c))^(1/2)+I*(-I*(exp(I*(b*x+a))+I))^(1/2)*2^(1/2)*(I*(exp(I*(b*x+a))-I))^(1/2)*(I*exp(I*(b*x+a)))^(1/2)/(c*exp(I*(b*x+a))^3+c*exp(I*(b*x+a)))^(1/2)*(-2*I*EllipticE((-I*(exp(I*(b*x+a))+I))^(1/2), 1/2*2^(1/2))+I*EllipticF((-I*(exp(I*(b*x+a))+I))^(1/2), 1/2*2^(1/2))))*2^(1/2)/(c*exp(I*(b*x+a))/(exp(I*(b*x+a))^2+1))^(1/2)*(c*exp(I*(b*x+a))*(exp(I*(b*x+a))^2+1))^(1/2)/(exp(I*(b*x+a))^2+1)`



### 3.21.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.74

$$\int \frac{1}{\sqrt{c \sec(a + bx)}} dx$$

$$= \frac{i \sqrt{2} \sqrt{c} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) - i \sqrt{2} \sqrt{c} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)))}{bc}$$

input `integrate(1/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`

output `(I*sqrt(2)*sqrt(c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - I*sqrt(2)*sqrt(c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/(b*c)`

### 3.21.6 Sympy [F]

$$\int \frac{1}{\sqrt{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt{c \sec(a + bx)}} dx$$

input `integrate(1/(c*sec(b*x+a))**(1/2),x)`

output `Integral(1/sqrt(c*sec(a + b*x)), x)`

### 3.21.7 Maxima [F]

$$\int \frac{1}{\sqrt{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate(1/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(c*sec(b*x + a)), x)`

**3.21.8 Giac [F]**

$$\int \frac{1}{\sqrt{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate(1/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(c*sec(b*x + a)), x)`

**3.21.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

input `int(1/(c/cos(a + b*x))^(1/2),x)`

output `int(1/(c/cos(a + b*x))^(1/2), x)`

## 3.22 $\int \frac{1}{(c \sec(a+bx))^{3/2}} dx$

3.22.1	Optimal result	230
3.22.2	Mathematica [A] (verified)	230
3.22.3	Rubi [A] (verified)	231
3.22.4	Maple [C] (verified)	232
3.22.5	Fricas [C] (verification not implemented)	233
3.22.6	Sympy [F]	233
3.22.7	Maxima [F]	233
3.22.8	Giac [F]	234
3.22.9	Mupad [F(-1)]	234

### 3.22.1 Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{1}{(c \sec(a+bx))^{3/2}} dx = \frac{2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) \sqrt{c \sec(a+bx)}}{3bc^2} + \frac{2 \sin(a+bx)}{3bc\sqrt{c \sec(a+bx)}}$$

output `2/3*sin(b*x+a)/b/c/(c*sec(b*x+a))^(1/2)+2/3*(cos(1/2*a+1/2*b*x)^2)^(1/2)/c  
os(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*(  
c*sec(b*x+a))^(1/2)/b/c^2`

### 3.22.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

$$\int \frac{1}{(c \sec(a+bx))^{3/2}} dx = \frac{\sec^2(a+bx) \left( 2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) + \sin(2(a+bx)) \right)}{3b(c \sec(a+bx))^{3/2}}$$

input `Integrate[(c*Sec[a + b*x])^(-3/2),x]`

output `(Sec[a + b*x]^2*(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Sin[2*(a  
+ b*x)]))/(3*b*(c*Sec[a + b*x])^(3/2))`

### 3.22.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \csc(a + bx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{\int \sqrt{c \sec(a + bx)} dx}{3c^2} + \frac{2 \sin(a + bx)}{3bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{c \csc(a + bx + \frac{\pi}{2})} dx}{3c^2} + \frac{2 \sin(a + bx)}{3bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)} \int \frac{1}{\sqrt{\cos(a + bx)}} dx}{3c^2} + \frac{2 \sin(a + bx)}{3bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)} \int \frac{1}{\sqrt{\sin(a + bx + \frac{\pi}{2})}} dx}{3c^2} + \frac{2 \sin(a + bx)}{3bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{c \sec(a + bx)}}{3bc^2} + \frac{2 \sin(a + bx)}{3bc \sqrt{c \sec(a + bx)}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(-3/2),x]`

output `(2*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*sqrt[c*Sec[a + b*x]])/(3*b*c^2) + (2*Sin[a + b*x])/(3*b*c*sqrt[c*Sec[a + b*x]])`

---

3.22.  $\int \frac{1}{(c \sec(a + bx))^{3/2}} dx$

## 3.22.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.22.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.93 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.00

method	result
default	$-\frac{2\left(i\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\operatorname{EllipticF}\left(i(-\cot(bx+a)+\csc(bx+a)),i\right)+i\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\operatorname{EllipticF}\left(i(-\cot(bx+a)+\csc(bx+a)),i\right)\right)}{3b\sqrt{c\sec(bx+a)}c}$

input `int(1/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/3/b/(c*sec(b*x+a))^(1/2)/c*(I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)+I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)*sec(b*x+a)-sin(b*x+a))`

**3.22.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.21

$$\int \frac{1}{(c \sec(a + bx))^{3/2}} dx = \frac{2 \sqrt{\frac{c}{\cos(bx+a)}} \cos(bx + a) \sin(bx + a) - i \sqrt{2} \sqrt{c} \text{weierstrassPInverse}(-4, 0, \cos(bx + a))}{(b^2 c^2)}$$

input `integrate(1/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

output `1/3*(2*sqrt(c/cos(b*x + a))*cos(b*x + a)*sin(b*x + a) - I*sqrt(2)*sqrt(c)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*sqrt(c)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/(b*c^2)`

**3.22.6 Sympy [F]**

$$\int \frac{1}{(c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(c \sec(a + bx))^{\frac{3}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))**(3/2),x)`

output `Integral((c*sec(a + b*x))**(-3/2), x)`

**3.22.7 Maxima [F]**

$$\int \frac{1}{(c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(3/2), x)`

**3.22.8 Giac [F]**

$$\int \frac{1}{(c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(3/2), x)`

**3.22.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c \sec(a + bx))^{3/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

input `int(1/(c/cos(a + b*x))^(3/2),x)`

output `int(1/(c/cos(a + b*x))^(3/2), x)`

### 3.23 $\int \frac{1}{(c \sec(a+bx))^{5/2}} dx$

3.23.1	Optimal result	235
3.23.2	Mathematica [A] (verified)	235
3.23.3	Rubi [A] (verified)	236
3.23.4	Maple [C] (verified)	237
3.23.5	Fricas [C] (verification not implemented)	238
3.23.6	Sympy [F]	238
3.23.7	Maxima [F]	239
3.23.8	Giac [F]	239
3.23.9	Mupad [F(-1)]	239

#### 3.23.1 Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{1}{(c \sec(a+bx))^{5/2}} dx = \frac{6E(\frac{1}{2}(a+bx)|2)}{5bc^2 \sqrt{\cos(a+bx)} \sqrt{c \sec(a+bx)}} + \frac{2 \sin(a+bx)}{5bc(c \sec(a+bx))^{3/2}}$$

output `2/5*sin(b*x+a)/b/c/(c*sec(b*x+a))^(3/2)+6/5*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))/b/c^2/cos(b*x+a)^(1/2)/(c*sec(b*x+a))^(1/2)`

#### 3.23.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.83

$$\int \frac{1}{(c \sec(a+bx))^{5/2}} dx = \frac{\sqrt{c \sec(a+bx)} \left( 12 \sqrt{\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right) + \sin(a+bx) + \sin(3(a+bx)) \right)}{10bc^3}$$

input `Integrate[(c*Sec[a + b*x])^(-5/2),x]`

output `(Sqrt[c*Sec[a + b*x]]*(12*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + Sin[a + b*x] + Sin[3*(a + b*x)]))/(10*b*c^3)`



### 3.23.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \csc(a + bx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{3 \int \frac{1}{\sqrt{c \sec(a+bx)}} dx}{5c^2} + \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \frac{1}{\sqrt{c \csc(a+bx+\frac{\pi}{2})}} dx}{5c^2} + \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{3 \int \sqrt{\cos(a + bx)} dx}{5c^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \sqrt{\sin(a + bx + \frac{\pi}{2})} dx}{5c^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{6E(\frac{1}{2}(a + bx) | 2)}{5bc^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(-5/2),x]`

output `(6*EllipticE[(a + b*x)/2, 2])/(5*b*c^2*Sqrt[Cos[a + b*x]]*Sqrt[c*Sec[a + b*x]]) + (2*Sin[a + b*x])/(5*b*c*(c*Sec[a + b*x])^(3/2))`

---

3.23.  $\int \frac{1}{(c \sec(a+bx))^{5/2}} dx$

## 3.23.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.23.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.15 (sec) , antiderivative size = 420, normalized size of antiderivative = 5.83

method	result
default	$\frac{6i \operatorname{EllipticE}(i(-\cot(bx+a)+\csc(bx+a)), i) \sqrt{\frac{1}{\cos(bx+a)+1}} \sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}} \cos(bx+a) - 6i \operatorname{EllipticF}(i(-\cot(bx+a)+\csc(bx+a)), i) \sqrt{\frac{1}{\cos(bx+a)+1}} \sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}}{5}$

input `int(1/(c*sec(b*x+a))^(5/2), x, method=_RETURNVERBOSE)`

```
output 2/5/b/(cos(b*x+a)+1)/(c*sec(b*x+a))^(1/2)/c^2*(3*I*(1/(cos(b*x+a)+1))^(1/2)
)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticE(I*(-cot(b*x+a)+csc(b*x+a)),I
)*cos(b*x+a)-3*I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)
)*EllipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)*cos(b*x+a)+6*I*(1/(cos(b*x+a)+1)
)^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticE(I*(-cot(b*x+a)+csc(b*x
+a)),I)-6*I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*Ell
ipticF(I*(-cot(b*x+a)+csc(b*x+a)),I)+3*I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x
+a)/(cos(b*x+a)+1))^(1/2)*EllipticE(I*(-cot(b*x+a)+csc(b*x+a)),I)*sec(b*x+
a)-3*I*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*Elliptic
F(I*(-cot(b*x+a)+csc(b*x+a)),I)*sec(b*x+a)+cos(b*x+a)^2*sin(b*x+a)+cos(b*x
+a)*sin(b*x+a)+3*sin(b*x+a))
```

### 3.23.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.32

$$\int \frac{1}{(c \sec(a + bx))^{5/2}} dx = \frac{2 \sqrt{\frac{c}{\cos(bx+a)}} \cos(bx+a)^2 \sin(bx+a) + 3i \sqrt{2} \sqrt{c} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx+a) + I \sin(bx+a))) - 3I \sqrt{2} \sqrt{c} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx+a) - I \sin(bx+a)))}{(b*c^3)}$$

```
input integrate(1/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")
```

```
output 1/5*(2*sqrt(c/cos(b*x + a))*cos(b*x + a)^2*sin(b*x + a) + 3*I*sqrt(2)*sqrt
(c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin
(b*x + a))) - 3*I*sqrt(2)*sqrt(c)*weierstrassZeta(-4, 0, weierstrassPInver
se(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/(b*c^3)
```

### 3.23.6 Sympy [F]

$$\int \frac{1}{(c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(c \sec(a + bx))^{\frac{5}{2}}} dx$$

```
input integrate(1/(c*sec(b*x+a))**(5/2),x)
```

```
output Integral((c*sec(a + b*x))**(-5/2), x)
```

---

3.23.  $\int \frac{1}{(c \sec(a + bx))^{5/2}} dx$

**3.23.7 Maxima [F]**

$$\int \frac{1}{(c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(-5/2), x)`

**3.23.8 Giac [F]**

$$\int \frac{1}{(c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(-5/2), x)`

**3.23.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

input `int(1/(c/cos(a + b*x))^(5/2),x)`

output `int(1/(c/cos(a + b*x))^(5/2), x)`

### 3.24 $\int \frac{1}{(c \sec(a+bx))^{7/2}} dx$

3.24.1	Optimal result	240
3.24.2	Mathematica [A] (verified)	240
3.24.3	Rubi [A] (verified)	241
3.24.4	Maple [C] (verified)	243
3.24.5	Fricas [C] (verification not implemented)	243
3.24.6	Sympy [F]	244
3.24.7	Maxima [F]	244
3.24.8	Giac [F]	244
3.24.9	Mupad [F(-1)]	245

#### 3.24.1 Optimal result

Integrand size = 12, antiderivative size = 100

$$\int \frac{1}{(c \sec(a + bx))^{7/2}} dx = \frac{10 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) \sqrt{c \sec(a + bx)}}{21bc^4} + \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} + \frac{10 \sin(a + bx)}{21bc^3 \sqrt{c \sec(a + bx)}}$$

output `2/7*sin(b*x+a)/b/c/(c*sec(b*x+a))^(5/2)+10/21*sin(b*x+a)/b/c^3/(c*sec(b*x+a))^(1/2)+10/21*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)*(c*sec(b*x+a))^(1/2)/b/c^4`

#### 3.24.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.66

$$\int \frac{1}{(c \sec(a + bx))^{7/2}} dx = \frac{\sqrt{c \sec(a + bx)} \left( 40 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) + 26 \sin(2(a + bx)) + 3 \sin(4(a + bx)) \right)}{84bc^4}$$

input `Integrate[(c*Sec[a + b*x])^(-7/2),x]`

output `(Sqrt[c*Sec[a + b*x]]*(40*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + 26*Sin[2*(a + b*x)] + 3*Sin[4*(a + b*x)]))/(84*b*c^4)`

### 3.24.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \csc(a + bx + \frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{5 \int \frac{1}{(c \sec(a+bx))^{3/2}} dx}{7c^2} + \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \int \frac{1}{(c \csc(a+bx+\frac{\pi}{2}))^{3/2}} dx}{7c^2} + \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{5 \left( \frac{\int \sqrt{c \sec(a+bx)} dx}{3c^2} + \frac{2 \sin(a+bx)}{3bc \sqrt{c \sec(a+bx)}} \right)}{7c^2} + \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \left( \frac{\int \sqrt{c \csc(a+bx+\frac{\pi}{2})} dx}{3c^2} + \frac{2 \sin(a+bx)}{3bc \sqrt{c \sec(a+bx)}} \right)}{7c^2} + \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{5 \left( \frac{\sqrt{\cos(a+bx)} \sqrt{c \sec(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{3c^2} + \frac{2 \sin(a+bx)}{3bc \sqrt{c \sec(a+bx)}} \right)}{7c^2} + \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{5 \left( \frac{\sqrt{\cos(a+bx)} \sqrt{c \sec(a+bx)} \int \frac{1}{\sqrt{\sin\left(a+bx+\frac{\pi}{2}\right)}} dx}{3c^2} + \frac{2 \sin(a+bx)}{3bc \sqrt{c \sec(a+bx)}} \right)}{7c^2} + \frac{2 \sin(a+bx)}{7bc(c \sec(a+bx))^{5/2}}$$

↓ 3120

$$\frac{5 \left( \frac{2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2, \sqrt{c \sec(a+bx)}\right)}{3bc^2} + \frac{2 \sin(a+bx)}{3bc \sqrt{c \sec(a+bx)}} \right)}{7c^2} + \frac{2 \sin(a+bx)}{7bc(c \sec(a+bx))^{5/2}}$$

input `Int[(c*Sec[a + b*x])^(-7/2), x]`

output `(2*Sin[a + b*x])/(7*b*c*(c*Sec[a + b*x])^(5/2)) + (5*((2*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*sqrt[c*Sec[a + b*x]])/(3*b*c^2) + (2*Sin[a + b*x])/(3*b*c*sqrt[c*Sec[a + b*x]])))/(7*c^2)`

### 3.24.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.24.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.01 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.60

method	result
default	$-\frac{2\left(5i\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\operatorname{EllipticF}(i(-\cot(bx+a)+\csc(bx+a)),i)+5i\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\operatorname{EllipticF}(i(-\cot(bx+a)+\csc(bx+a)),i)\right)}{21b\sqrt{c\sec(bx+a)}c^3}$

input `int(1/(c*sec(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

output 
$$-2/21/b/(c*\sec(b*x+a))^{(1/2)}/c^3*(5*I*(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(b*x+a)+\csc(b*x+a)),I)+5*I*(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(b*x+a)+\csc(b*x+a)),I)*\sec(b*x+a)-3*\cos(b*x+a)^2*\sin(b*x+a)-5*\sin(b*x+a))$$

### 3.24.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c\sec(a+bx))^{7/2}} dx = \frac{2(3\cos(bx+a)^3 + 5\cos(bx+a))\sqrt{\frac{c}{\cos(bx+a)}}\sin(bx+a) - 5i\sqrt{2}\sqrt{c}\operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + I\sin(bx+a))}{(b*c^4)}$$

input `integrate(1/(c*sec(b*x+a))^(7/2),x, algorithm="fricas")`

output 
$$1/21*(2*(3*\cos(b*x + a)^3 + 5*\cos(b*x + a))*\sqrt{c/\cos(b*x + a)}*\sin(b*x + a) - 5*I*\sqrt{2}*\sqrt{c}*\operatorname{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I*\sin(b*x + a)) + 5*I*\sqrt{2}*\sqrt{c}*\operatorname{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I*\sin(b*x + a)))/(b*c^4)$$



**3.24.6 Sympy [F]**

$$\int \frac{1}{(c \sec(a + bx))^{7/2}} dx = \int \frac{1}{(c \sec(a + bx))^{\frac{7}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))**(7/2), x)`

output `Integral((c*sec(a + b*x))**(-7/2), x)`

**3.24.7 Maxima [F]**

$$\int \frac{1}{(c \sec(a + bx))^{7/2}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{7}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(7/2), x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(7/2), x)`

**3.24.8 Giac [F]**

$$\int \frac{1}{(c \sec(a + bx))^{7/2}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{7}{2}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(7/2), x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(7/2), x)`

**3.24.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c \sec(a + bx))^{7/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{7/2}} dx$$

input `int(1/(c/cos(a + b*x))^(7/2),x)`output `int(1/(c/cos(a + b*x))^(7/2), x)`

### 3.25 $\int \sec^{\frac{4}{3}}(a + bx) dx$

3.25.1	Optimal result . . . . .	246
3.25.2	Mathematica [A] (verified) . . . . .	246
3.25.3	Rubi [A] (verified) . . . . .	247
3.25.4	Maple [F] . . . . .	248
3.25.5	Fricas [F] . . . . .	248
3.25.6	Sympy [F] . . . . .	249
3.25.7	Maxima [F] . . . . .	249
3.25.8	Giac [F] . . . . .	249
3.25.9	Mupad [F(-1)] . . . . .	250

#### 3.25.1 Optimal result

Integrand size = 10, antiderivative size = 51

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a + bx)\right) \sqrt[3]{\sec(a + bx)} \sin(a + bx)}{b \sqrt{\sin^2(a + bx)}}$$

output `3*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*sec(b*x+a)^(1/3)*sin(b*x+a)/b/(sin(b*x+a)^2)^(1/2)`

#### 3.25.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.08

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \frac{3 \operatorname{csc}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \sec^2(a + bx)\right) \sqrt[3]{\sec(a + bx)} \sqrt{-\tan^2(a + bx)}}{4b}$$

input `Integrate[Sec[a + b*x]^(4/3), x]`

output `(3*Csc[a + b*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[a + b*x]^2]*Sec[a + b*x]^(1/3)*Sqrt[-Tan[a + b*x]^2])/(4*b)`

### 3.25.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{4}{3}}(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^{\frac{4}{3}} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\cos(a + bx)} \sqrt[3]{\sec(a + bx)} \int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\cos(a + bx)} \sqrt[3]{\sec(a + bx)} \int \frac{1}{\sin\left(a + bx + \frac{\pi}{2}\right)^{\frac{4}{3}}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(a + bx) \sqrt[3]{\sec(a + bx)} \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)}}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(4/3),x]`

output `(3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sec[a + b*x]^(1/3)*Sin[a + b*x])/(b*Sqrt[Sin[a + b*x]^2])`

## 3.25.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.25.4 Maple [F]

$$\int \sec (bx + a)^{\frac{4}{3}} dx$$

input `int(sec(b*x+a)^(4/3),x)`

output `int(sec(b*x+a)^(4/3),x)`

## 3.25.5 Fracas [F]

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \int \sec (bx + a)^{\frac{4}{3}} dx$$

input `integrate(sec(b*x+a)^(4/3),x, algorithm="fricas")`

output `integral(sec(b*x + a)^(4/3), x)`

**3.25.6 Sympy [F]**

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \int \sec^{\frac{4}{3}}(a + bx) dx$$

input `integrate(sec(b*x+a)**(4/3),x)`

output `Integral(sec(a + b*x)**(4/3), x)`

**3.25.7 Maxima [F]**

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \int \sec (bx + a)^{\frac{4}{3}} dx$$

input `integrate(sec(b*x+a)^(4/3),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(4/3), x)`

**3.25.8 Giac [F]**

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \int \sec (bx + a)^{\frac{4}{3}} dx$$

input `integrate(sec(b*x+a)^(4/3),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(4/3), x)`

**3.25.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \int \left( \frac{1}{\cos(a + bx)} \right)^{\frac{4}{3}} dx$$

input `int((1/cos(a + b*x))^(4/3),x)`output `int((1/cos(a + b*x))^(4/3), x)`

## 3.26 $\int \sec^{\frac{2}{3}}(a + bx) dx$

3.26.1	Optimal result	251
3.26.2	Mathematica [A] (verified)	251
3.26.3	Rubi [A] (verified)	252
3.26.4	Maple [F]	253
3.26.5	Fricas [F]	253
3.26.6	Sympy [F]	254
3.26.7	Maxima [F]	254
3.26.8	Giac [F]	254
3.26.9	Mupad [F(-1)]	255

### 3.26.1 Optimal result

Integrand size = 10, antiderivative size = 51

$$\int \sec^{\frac{2}{3}}(a + bx) dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{b \sqrt[3]{\sec(a + bx)} \sqrt{\sin^2(a + bx)}}$$

output `-3*hypergeom([1/6, 1/2],[7/6],cos(b*x+a)^2)*sin(b*x+a)/b/sec(b*x+a)^(1/3)/  
(sin(b*x+a)^2)^(1/2)`

### 3.26.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.08

$$\int \sec^{\frac{2}{3}}(a + bx) dx = \frac{3 \operatorname{csc}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \sec^2(a + bx)\right) \sqrt{-\tan^2(a + bx)}}{2b \sqrt[3]{\sec(a + bx)}}$$

input `Integrate[Sec[a + b*x]^(2/3),x]`

output `(3*Csc[a + b*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[a + b*x]^2]*Sqrt[-Tan  
[a + b*x]^2])/(2*b*Sec[a + b*x]^(1/3))`



**3.26.3 Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{2}{3}}(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^{2/3} dx \\
 & \quad \downarrow \text{4259} \\
 & \cos^{\frac{2}{3}}(a + bx) \sec^{\frac{2}{3}}(a + bx) \int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \cos^{\frac{2}{3}}(a + bx) \sec^{\frac{2}{3}}(a + bx) \int \frac{1}{\sin\left(a + bx + \frac{\pi}{2}\right)^{2/3}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)} \sqrt[3]{\sec(a + bx)}}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(2/3),x]`

output `(-3*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[a + b*x]^2]*Sin[a + b*x])/(b*Sec[a + b*x]^(1/3)*Sqrt[Sin[a + b*x]^2])`

## 3.26.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.26.4 Maple [F]

$$\int \sec (bx + a)^{\frac{2}{3}} dx$$

```
input int(sec(b*x+a)^(2/3),x)
```

```
output int(sec(b*x+a)^(2/3),x)
```

## 3.26.5 Fricas [F]

$$\int \sec^{\frac{2}{3}}(a + bx) dx = \int \sec (bx + a)^{\frac{2}{3}} dx$$

```
input integrate(sec(b*x+a)^(2/3),x, algorithm="fricas")
```

```
output integral(sec(b*x + a)^(2/3), x)
```

**3.26.6 Sympy [F]**

$$\int \sec^{\frac{2}{3}}(a + bx) dx = \int \sec^{\frac{2}{3}}(a + bx) dx$$

input `integrate(sec(b*x+a)**(2/3),x)`

output `Integral(sec(a + b*x)**(2/3), x)`

**3.26.7 Maxima [F]**

$$\int \sec^{\frac{2}{3}}(a + bx) dx = \int \sec (bx + a)^{\frac{2}{3}} dx$$

input `integrate(sec(b*x+a)^(2/3),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(2/3), x)`

**3.26.8 Giac [F]**

$$\int \sec^{\frac{2}{3}}(a + bx) dx = \int \sec (bx + a)^{\frac{2}{3}} dx$$

input `integrate(sec(b*x+a)^(2/3),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(2/3), x)`

**3.26.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{2}{3}}(a + bx) dx = \int \left( \frac{1}{\cos(a + bx)} \right)^{\frac{2}{3}} dx$$

input `int((1/cos(a + b*x))^(2/3),x)`output `int((1/cos(a + b*x))^(2/3), x)`

### 3.27 $\int \sqrt[3]{\sec(a + bx)} dx$

3.27.1	Optimal result . . . . .	256
3.27.2	Mathematica [A] (verified) . . . . .	256
3.27.3	Rubi [A] (verified) . . . . .	257
3.27.4	Maple [F] . . . . .	258
3.27.5	Fricas [F] . . . . .	258
3.27.6	Sympy [F] . . . . .	259
3.27.7	Maxima [F] . . . . .	259
3.27.8	Giac [F] . . . . .	259
3.27.9	Mupad [F(-1)] . . . . .	260

#### 3.27.1 Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \sqrt[3]{\sec(a + bx)} dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{2b \sec^{\frac{2}{3}}(a + bx) \sqrt{\sin^2(a + bx)}}$$

output `-3/2*hypergeom([1/3, 1/2],[4/3],cos(b*x+a)^2)*sin(b*x+a)/b/sec(b*x+a)^(2/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.27.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \sqrt[3]{\sec(a + bx)} dx = \frac{3 \operatorname{csc}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \sec^2(a + bx)\right) \sqrt{-\tan^2(a + bx)}}{b \sec^{\frac{2}{3}}(a + bx)}$$

input `Integrate[Sec[a + b*x]^(1/3),x]`

output `(3*Csc[a + b*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(b*Sec[a + b*x]^(2/3))`

### 3.27.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{\sec(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt[3]{\csc\left(a+bx+\frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\cos(a+bx)} \sqrt[3]{\sec(a+bx)} \int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\cos(a+bx)} \sqrt[3]{\sec(a+bx)} \int \frac{1}{\sqrt[3]{\sin\left(a+bx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a+bx)\right)}{2b \sqrt{\sin^2(a+bx)} \sec^{\frac{2}{3}}(a+bx)}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(1/3),x]`

output `(-3*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[a + b*x]^2]*Sin[a + b*x])/(2*b*Sec[a + b*x]^(2/3)*Sqrt[Sin[a + b*x]^2])`

## 3.27.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.27.4 Maple [F]

$$\int \sec (bx + a)^{\frac{1}{3}} dx$$

input `int(sec(b*x+a)^(1/3),x)`

output `int(sec(b*x+a)^(1/3),x)`

## 3.27.5 Fracas [F]

$$\int \sqrt[3]{\sec(a + bx)} dx = \int \sec (bx + a)^{\frac{1}{3}} dx$$

input `integrate(sec(b*x+a)^(1/3),x, algorithm="fricas")`

output `integral(sec(b*x + a)^(1/3), x)`

**3.27.6 Sympy [F]**

$$\int \sqrt[3]{\sec(a + bx)} dx = \int \sqrt[3]{\sec(a + bx)} dx$$

input `integrate(sec(b*x+a)**(1/3),x)`

output `Integral(sec(a + b*x)**(1/3), x)`

**3.27.7 Maxima [F]**

$$\int \sqrt[3]{\sec(a + bx)} dx = \int \sec(bx + a)^{\frac{1}{3}} dx$$

input `integrate(sec(b*x+a)^(1/3),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(1/3), x)`

**3.27.8 Giac [F]**

$$\int \sqrt[3]{\sec(a + bx)} dx = \int \sec(bx + a)^{\frac{1}{3}} dx$$

input `integrate(sec(b*x+a)^(1/3),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(1/3), x)`



**3.27.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt[3]{\sec(a + bx)} dx = \int \left( \frac{1}{\cos(a + bx)} \right)^{1/3} dx$$

input `int((1/cos(a + b*x))^(1/3),x)`output `int((1/cos(a + b*x))^(1/3), x)`

### 3.28 $\int \frac{1}{\sqrt[3]{\sec(a + bx)}} dx$

3.28.1	Optimal result	261
3.28.2	Mathematica [A] (verified)	261
3.28.3	Rubi [A] (verified)	262
3.28.4	Maple [F]	263
3.28.5	Fricas [F]	263
3.28.6	Sympy [F]	264
3.28.7	Maxima [F]	264
3.28.8	Giac [F]	264
3.28.9	Mupad [F(-1)]	265

#### 3.28.1 Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \frac{1}{\sqrt[3]{\sec(a + bx)}} dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{4b \sec^{\frac{4}{3}}(a + bx) \sqrt{\sin^2(a + bx)}}$$

output `-3/4*hypergeom([1/2, 2/3], [5/3], cos(b*x+a)^2)*sin(b*x+a)/b/sec(b*x+a)^(4/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.28.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt[3]{\sec(a + bx)}} dx = -\frac{3 \operatorname{csc}(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \sec^2(a + bx)\right) \sqrt{-\tan^2(a + bx)}}{b \sec^{\frac{4}{3}}(a + bx)}$$

input `Integrate[Sec[a + b*x]^(-1/3), x]`

output `(-3*Csc[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(b*Sec[a + b*x]^(4/3))`

### 3.28.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{\csc\left(a+bx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{4259} \\
 & \cos^{\frac{2}{3}}(a+bx) \sec^{\frac{2}{3}}(a+bx) \int \sqrt[3]{\cos(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \cos^{\frac{2}{3}}(a+bx) \sec^{\frac{2}{3}}(a+bx) \int \sqrt[3]{\sin\left(a+bx+\frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a+bx)\right)}{4b \sqrt{\sin^2(a+bx)} \sec^{\frac{4}{3}}(a+bx)}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(-1/3),x]`

output `(-3*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[a + b*x]^2]*Sin[a + b*x])/(4*b*Sec[a + b*x]^(4/3)*Sqrt[Sin[a + b*x]^2])`

## 3.28.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.28.4 Maple [F]

$$\int \frac{1}{\sec(bx + a)^{\frac{1}{3}}} dx$$

input `int(1/sec(b*x+a)^(1/3),x)`

output `int(1/sec(b*x+a)^(1/3),x)`

## 3.28.5 Fricas [F]

$$\int \frac{1}{\sqrt[3]{\sec(a + bx)}} dx = \int \frac{1}{\sec(bx + a)^{\frac{1}{3}}} dx$$

input `integrate(1/sec(b*x+a)^(1/3),x, algorithm="fricas")`

output `integral(sec(b*x + a)^(-1/3), x)`

**3.28.6 Sympy [F]**

$$\int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx = \int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx$$

input `integrate(1/sec(b*x+a)**(1/3),x)`

output `Integral(sec(a + b*x)**(-1/3), x)`

**3.28.7 Maxima [F]**

$$\int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx = \int \frac{1}{\sec(bx+a)^{\frac{1}{3}}} dx$$

input `integrate(1/sec(b*x+a)^(1/3),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(-1/3), x)`

**3.28.8 Giac [F]**

$$\int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx = \int \frac{1}{\sec(bx+a)^{\frac{1}{3}}} dx$$

input `integrate(1/sec(b*x+a)^(1/3),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(-1/3), x)`

**3.28.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx = \int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{1/3}} dx$$

input `int(1/(1/cos(a + b*x))^(1/3),x)`output `int(1/(1/cos(a + b*x))^(1/3), x)`

### 3.29 $\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx$

3.29.1	Optimal result	266
3.29.2	Mathematica [A] (verified)	266
3.29.3	Rubi [A] (verified)	267
3.29.4	Maple [F]	268
3.29.5	Fricas [F]	268
3.29.6	Sympy [F]	269
3.29.7	Maxima [F]	269
3.29.8	Giac [F]	269
3.29.9	Mupad [F(-1)]	270

#### 3.29.1 Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{5b \sec^{\frac{5}{3}}(a+bx) \sqrt{\sin^2(a+bx)}}$$

output `-3/5*hypergeom([1/2, 5/6], [11/6], cos(b*x+a)^2)*sin(b*x+a)/b/sec(b*x+a)^(5/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.29.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.04

$$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx = -\frac{3 \operatorname{csc}(a+bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \sec^2(a+bx)\right) \sqrt{-\tan^2(a+bx)}}{2b \sec^{\frac{5}{3}}(a+bx)}$$

input `Integrate[Sec[a + b*x]^(-2/3), x]`

output `(-3*Csc[a + b*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(2*b*Sec[a + b*x]^(5/3))`

### 3.29.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc\left(a+bx+\frac{\pi}{2}\right)^{2/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\cos(a+bx)} \sqrt[3]{\sec(a+bx)} \int \cos^{\frac{2}{3}}(a+bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\cos(a+bx)} \sqrt[3]{\sec(a+bx)} \int \sin\left(a+bx+\frac{\pi}{2}\right)^{2/3} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3 \sin(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a+bx)\right)}{5b \sqrt{\sin^2(a+bx)} \sec^{\frac{5}{3}}(a+bx)}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(-2/3),x]`

output `(-3*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[a + b*x]^2]*Sin[a + b*x])/(5*b*Sec[a + b*x]^(5/3)*Sqrt[Sin[a + b*x]^2])`



## 3.29.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.29.4 Maple [F]

$$\int \frac{1}{\sec^{\frac{2}{3}}(bx + a)} dx$$

input `int(1/sec(b*x+a)^(2/3),x)`

output `int(1/sec(b*x+a)^(2/3),x)`

## 3.29.5 Fricas [F]

$$\int \frac{1}{\sec^{\frac{2}{3}}(a + bx)} dx = \int \frac{1}{\sec^{\frac{2}{3}}(bx + a)} dx$$

input `integrate(1/sec(b*x+a)^(2/3),x, algorithm="fricas")`

output `integral(sec(b*x + a)^(-2/3), x)`

**3.29.6 Sympy [F]**

$$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx$$

input `integrate(1/sec(b*x+a)**(2/3),x)`

output `Integral(sec(a + b*x)**(-2/3), x)`

**3.29.7 Maxima [F]**

$$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{2}{3}}(bx+a)} dx$$

input `integrate(1/sec(b*x+a)^(2/3),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(-2/3), x)`

**3.29.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{2}{3}}(bx+a)} dx$$

input `integrate(1/sec(b*x+a)^(2/3),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(-2/3), x)`

**3.29.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx = \int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{2}{3}}} dx$$

input `int(1/(1/cos(a + b*x))^(2/3),x)`output `int(1/(1/cos(a + b*x))^(2/3), x)`

### 3.30 $\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx$

3.30.1	Optimal result	271
3.30.2	Mathematica [A] (verified)	271
3.30.3	Rubi [A] (verified)	272
3.30.4	Maple [F]	273
3.30.5	Fricas [F]	273
3.30.6	Sympy [F]	274
3.30.7	Maxima [F]	274
3.30.8	Giac [F]	274
3.30.9	Mupad [F(-1)]	275

#### 3.30.1 Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{7b \sec^{\frac{7}{3}}(a+bx) \sqrt{\sin^2(a+bx)}}$$

output `-3/7*hypergeom([1/2, 7/6], [13/6], cos(b*x+a)^2)*sin(b*x+a)/b/sec(b*x+a)^(7/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.30.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.04

$$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx = -\frac{3 \operatorname{csc}(a+bx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \sec^2(a+bx)\right) \sqrt{-\tan^2(a+bx)}}{4b \sec^{\frac{7}{3}}(a+bx)}$$

input `Integrate[Sec[a + b*x]^(-4/3), x]`

output `(-3*Csc[a + b*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(4*b*Sec[a + b*x]^(7/3))`

### 3.30.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc\left(a+bx+\frac{\pi}{2}\right)^{\frac{4}{3}}} dx \\
 & \quad \downarrow \text{4259} \\
 & \cos^{\frac{2}{3}}(a+bx) \sec^{\frac{2}{3}}(a+bx) \int \cos^{\frac{4}{3}}(a+bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \cos^{\frac{2}{3}}(a+bx) \sec^{\frac{2}{3}}(a+bx) \int \sin\left(a+bx+\frac{\pi}{2}\right)^{\frac{4}{3}} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3 \sin(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a+bx)\right)}{7b \sqrt{\sin^2(a+bx)} \sec^{\frac{7}{3}}(a+bx)}
 \end{aligned}$$

input `Int[Sec[a + b*x]^(-4/3),x]`

output `(-3*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[a + b*x]^2]*Sin[a + b*x])/(7*b*Sec[a + b*x]^(7/3)*Sqrt[Sin[a + b*x]^2])`

## 3.30.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.30.4 Maple [F]

$$\int \frac{1}{\sec^{\frac{4}{3}}(bx + a)} dx$$

input `int(1/sec(b*x+a)^(4/3),x)`

output `int(1/sec(b*x+a)^(4/3),x)`

## 3.30.5 Fracas [F]

$$\int \frac{1}{\sec^{\frac{4}{3}}(a + bx)} dx = \int \frac{1}{\sec^{\frac{4}{3}}(bx + a)} dx$$

input `integrate(1/sec(b*x+a)^(4/3),x, algorithm="fricas")`

output `integral(sec(b*x + a)^(-4/3), x)`

**3.30.6 Sympy [F]**

$$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx$$

input `integrate(1/sec(b*x+a)**(4/3),x)`

output `Integral(sec(a + b*x)**(-4/3), x)`

**3.30.7 Maxima [F]**

$$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{4}{3}}(bx+a)} dx$$

input `integrate(1/sec(b*x+a)^(4/3),x, algorithm="maxima")`

output `integrate(sec(b*x + a)^(-4/3), x)`

**3.30.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx = \int \frac{1}{\sec^{\frac{4}{3}}(bx+a)} dx$$

input `integrate(1/sec(b*x+a)^(4/3),x, algorithm="giac")`

output `integrate(sec(b*x + a)^(-4/3), x)`

**3.30.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx = \int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{4}{3}}} dx$$

input `int(1/(1/cos(a + b*x))^(4/3),x)`output `int(1/(1/cos(a + b*x))^(4/3), x)`



### 3.31 $\int (c \sec(a + bx))^{4/3} dx$

3.31.1	Optimal result	276
3.31.2	Mathematica [A] (verified)	276
3.31.3	Rubi [A] (verified)	277
3.31.4	Maple [F]	278
3.31.5	Fricas [F]	278
3.31.6	Sympy [F]	279
3.31.7	Maxima [F]	279
3.31.8	Giac [F]	279
3.31.9	Mupad [F(-1)]	280

#### 3.31.1 Optimal result

Integrand size = 12, antiderivative size = 54

$$\int (c \sec(a + bx))^{4/3} dx = \frac{3c \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a + bx)\right) \sqrt[3]{c \sec(a + bx)} \sin(a + bx)}{b \sqrt{\sin^2(a + bx)}}$$

output `3*c*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*(c*sec(b*x+a))^(1/3)*sin(b*x+a)/b/(sin(b*x+a)^2)^(1/2)`

#### 3.31.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06

$$\int (c \sec(a + bx))^{4/3} dx = \frac{3 \cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \sec^2(a + bx)\right) (c \sec(a + bx))^{4/3} \sqrt{-\tan^2(a + bx)}}{4b}$$

input `Integrate[(c*Sec[a + b*x])^(4/3), x]`

output `(3*Cot[a + b*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[a + b*x]^2]*(c*Sec[a + b*x])^(4/3)*Sqrt[-Tan[a + b*x]^2])/(4*b)`

**3.31.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( c \csc \left( a + bx + \frac{\pi}{2} \right) \right)^{4/3} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \frac{1}{\left( \frac{\cos(a + bx)}{c} \right)^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \frac{1}{\left( \frac{\sin(a + bx + \frac{\pi}{2})}{c} \right)^{4/3}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3c \sin(a + bx) \sqrt[3]{c \sec(a + bx)} \operatorname{Hypergeometric2F1} \left( -\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a + bx) \right)}{b \sqrt{\sin^2(a + bx)}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(4/3),x]`

output `(3*c*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*(c*Sec[a + b*x])^(1/3)*Sin[a + b*x])/(b*Sqrt[Sin[a + b*x]^2])`

## 3.31.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.31.4 Maple [F]

$$\int (c \sec (bx + a))^{\frac{4}{3}} dx$$

input `int((c*sec(b*x+a))^(4/3),x)`

output `int((c*sec(b*x+a))^(4/3),x)`

## 3.31.5 Fricas [F]

$$\int (c \sec (a + bx))^{\frac{4}{3}} dx = \int (c \sec (bx + a))^{\frac{4}{3}} dx$$

input `integrate((c*sec(b*x+a))^(4/3),x, algorithm="fricas")`

output `integral((c*sec(b*x + a))^(1/3)*c*sec(b*x + a), x)`

**3.31.6 Sympy [F]**

$$\int (c \sec(a + bx))^{4/3} dx = \int (c \sec(a + bx))^{\frac{4}{3}} dx$$

input `integrate((c*sec(b*x+a))**(4/3),x)`

output `Integral((c*sec(a + b*x))**(4/3), x)`

**3.31.7 Maxima [F]**

$$\int (c \sec(a + bx))^{4/3} dx = \int (c \sec(bx + a))^{\frac{4}{3}} dx$$

input `integrate((c*sec(b*x+a))^(4/3),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(4/3), x)`

**3.31.8 Giac [F]**

$$\int (c \sec(a + bx))^{4/3} dx = \int (c \sec(bx + a))^{\frac{4}{3}} dx$$

input `integrate((c*sec(b*x+a))^(4/3),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(4/3), x)`

**3.31.9 Mupad [F(-1)]**

Timed out.

$$\int (c \sec(a + bx))^{4/3} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{4/3} dx$$

input `int((c/cos(a + b*x))^(4/3),x)`output `int((c/cos(a + b*x))^(4/3), x)`

### 3.32 $\int (c \sec(a + bx))^{2/3} dx$

3.32.1	Optimal result	281
3.32.2	Mathematica [A] (verified)	281
3.32.3	Rubi [A] (verified)	282
3.32.4	Maple [F]	283
3.32.5	Fricas [F]	283
3.32.6	Sympy [F]	284
3.32.7	Maxima [F]	284
3.32.8	Giac [F]	284
3.32.9	Mupad [F(-1)]	285

#### 3.32.1 Optimal result

Integrand size = 12, antiderivative size = 54

$$\int (c \sec(a + bx))^{2/3} dx = -\frac{3c \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{b \sqrt[3]{c \sec(a + bx)} \sqrt{\sin^2(a + bx)}}$$

output `-3*c*hypergeom([1/6, 1/2],[7/6],cos(b*x+a)^2)*sin(b*x+a)/b/(c*sec(b*x+a))^(1/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.32.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06

$$\int (c \sec(a + bx))^{2/3} dx = \frac{3 \cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \sec^2(a + bx)\right) (c \sec(a + bx))^{2/3} \sqrt{-\tan^2(a + bx)}}{2b}$$

input `Integrate[(c*Sec[a + b*x])^(2/3),x]`

output `(3*Cot[a + b*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[a + b*x]^2]*(c*Sec[a + b*x])^(2/3)*Sqrt[-Tan[a + b*x]^2])/(2*b)`

**3.32.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{2/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( c \csc \left( a + bx + \frac{\pi}{2} \right) \right)^{2/3} dx \\
 & \quad \downarrow \text{4259} \\
 & \left( \frac{\cos(a + bx)}{c} \right)^{2/3} (c \sec(a + bx))^{2/3} \int \frac{1}{\left( \frac{\cos(a + bx)}{c} \right)^{2/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \left( \frac{\cos(a + bx)}{c} \right)^{2/3} (c \sec(a + bx))^{2/3} \int \frac{1}{\left( \frac{\sin(a + bx + \frac{\pi}{2})}{c} \right)^{2/3}} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3c \sin(a + bx) \operatorname{Hypergeometric2F1} \left( \frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a + bx) \right)}{b \sqrt{\sin^2(a + bx)} \sqrt[3]{c \sec(a + bx)}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(2/3),x]`

output `(-3*c*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[a + b*x]^2]*Sin[a + b*x])/(b*(c*Sec[a + b*x])^(1/3)*Sqrt[Sin[a + b*x]^2])`

## 3.32.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.32.4 Maple [F]

$$\int (c \sec (bx + a))^{\frac{2}{3}} dx$$

input `int((c*sec(b*x+a))^(2/3),x)`

output `int((c*sec(b*x+a))^(2/3),x)`

## 3.32.5 Fricas [F]

$$\int (c \sec (a + bx))^{2/3} dx = \int (c \sec (bx + a))^{\frac{2}{3}} dx$$

input `integrate((c*sec(b*x+a))^(2/3),x, algorithm="fricas")`

output `integral((c*sec(b*x + a))^(2/3), x)`



**3.32.6 Sympy [F]**

$$\int (c \sec(a + bx))^{2/3} dx = \int (c \sec(a + bx))^{\frac{2}{3}} dx$$

input `integrate((c*sec(b*x+a))**(2/3),x)`

output `Integral((c*sec(a + b*x))**(2/3), x)`

**3.32.7 Maxima [F]**

$$\int (c \sec(a + bx))^{2/3} dx = \int (c \sec(bx + a))^{\frac{2}{3}} dx$$

input `integrate((c*sec(b*x+a))^(2/3),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(2/3), x)`

**3.32.8 Giac [F]**

$$\int (c \sec(a + bx))^{2/3} dx = \int (c \sec(bx + a))^{\frac{2}{3}} dx$$

input `integrate((c*sec(b*x+a))^(2/3),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(2/3), x)`

**3.32.9 Mupad [F(-1)]**

Timed out.

$$\int (c \sec(a + bx))^{2/3} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{2/3} dx$$

input `int((c/cos(a + b*x))^(2/3),x)`output `int((c/cos(a + b*x))^(2/3), x)`

### 3.33 $\int \sqrt[3]{c \sec(a + bx)} dx$

3.33.1	Optimal result . . . . .	286
3.33.2	Mathematica [A] (verified) . . . . .	286
3.33.3	Rubi [A] (verified) . . . . .	287
3.33.4	Maple [F] . . . . .	288
3.33.5	Fricas [F] . . . . .	288
3.33.6	Sympy [F] . . . . .	289
3.33.7	Maxima [F] . . . . .	289
3.33.8	Giac [F] . . . . .	289
3.33.9	Mupad [F(-1)] . . . . .	290

#### 3.33.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \sqrt[3]{c \sec(a + bx)} dx = -\frac{3c \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{2b(c \sec(a + bx))^{2/3} \sqrt{\sin^2(a + bx)}}$$

output `-3/2*c*hypergeom([1/3, 1/2],[4/3],cos(b*x+a)^2)*sin(b*x+a)/b/(c*sec(b*x+a))^(2/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.33.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98

$$\int \sqrt[3]{c \sec(a + bx)} dx = \frac{3 \cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \sec^2(a + bx)\right) \sqrt[3]{c \sec(a + bx)} \sqrt{-\tan^2(a + bx)}}{b}$$

input `Integrate[(c*Sec[a + b*x])^(1/3),x]`

output `(3*Cot[a + b*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[a + b*x]^2]*(c*Sec[a + b*x])^(1/3)*Sqrt[-Tan[a + b*x]^2])/b`

### 3.33.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{c \sec(a + bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt[3]{c \csc\left(a + bx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \frac{1}{\sqrt[3]{\frac{\cos(a + bx)}{c}}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \frac{1}{\sqrt[3]{\frac{\sin\left(a + bx + \frac{\pi}{2}\right)}{c}}} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3c \sin(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right)}{2b \sqrt{\sin^2(a + bx)} (c \sec(a + bx))^{2/3}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(1/3),x]`

output `(-3*c*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[a + b*x]^2]*Sin[a + b*x])/(2*b*(c*Sec[a + b*x])^(2/3)*Sqrt[Sin[a + b*x]^2])`

## 3.33.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.33.4 Maple [F]

$$\int (c \sec (bx + a))^{\frac{1}{3}} dx$$

input `int((c*sec(b*x+a))^(1/3),x)`

output `int((c*sec(b*x+a))^(1/3),x)`

## 3.33.5 Fricas [F]

$$\int \sqrt[3]{c \sec (a + bx)} dx = \int (c \sec (bx + a))^{\frac{1}{3}} dx$$

input `integrate((c*sec(b*x+a))^(1/3),x, algorithm="fricas")`

output `integral((c*sec(b*x + a))^(1/3), x)`

**3.33.6 Sympy [F]**

$$\int \sqrt[3]{c \sec(a + bx)} dx = \int \sqrt[3]{c \sec(a + bx)} dx$$

input `integrate((c*sec(b*x+a))**(1/3),x)`

output `Integral((c*sec(a + b*x))**(1/3), x)`

**3.33.7 Maxima [F]**

$$\int \sqrt[3]{c \sec(a + bx)} dx = \int (c \sec(bx + a))^{\frac{1}{3}} dx$$

input `integrate((c*sec(b*x+a))^(1/3),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(1/3), x)`

**3.33.8 Giac [F]**

$$\int \sqrt[3]{c \sec(a + bx)} dx = \int (c \sec(bx + a))^{\frac{1}{3}} dx$$

input `integrate((c*sec(b*x+a))^(1/3),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(1/3), x)`

**3.33.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt[3]{c \sec(a + bx)} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{1/3} dx$$

input `int((c/cos(a + b*x))^(1/3),x)`output `int((c/cos(a + b*x))^(1/3), x)`

### 3.34 $\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx$

3.34.1	Optimal result	291
3.34.2	Mathematica [A] (verified)	291
3.34.3	Rubi [A] (verified)	292
3.34.4	Maple [F]	293
3.34.5	Fricas [F]	293
3.34.6	Sympy [F]	294
3.34.7	Maxima [F]	294
3.34.8	Giac [F]	294
3.34.9	Mupad [F(-1)]	295

#### 3.34.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx = -\frac{3c \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{4b(c \sec(a + bx))^{4/3} \sqrt{\sin^2(a + bx)}}$$

output `-3/4*c*hypergeom([1/2, 2/3], [5/3], cos(b*x+a)^2)*sin(b*x+a)/b/(c*sec(b*x+a))^(4/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.34.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx = -\frac{3 \cot(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \sec^2(a + bx)\right) \sqrt{-\tan^2(a + bx)}}{b \sqrt[3]{c \sec(a + bx)}}$$

input `Integrate[(c*Sec[a + b*x])^(-1/3), x]`

output `(-3*Cot[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(b*(c*Sec[a + b*x])^(1/3))`



### 3.34.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{c \sec(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{c \csc\left(a+bx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{4259} \\
 & \left(\frac{\cos(a+bx)}{c}\right)^{2/3} (c \sec(a+bx))^{2/3} \int \sqrt[3]{\frac{\cos(a+bx)}{c}} dx \\
 & \quad \downarrow \text{3042} \\
 & \left(\frac{\cos(a+bx)}{c}\right)^{2/3} (c \sec(a+bx))^{2/3} \int \sqrt[3]{\frac{\sin\left(a+bx+\frac{\pi}{2}\right)}{c}} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3c \sin(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a+bx)\right)}{4b \sqrt{\sin^2(a+bx)} (c \sec(a+bx))^{4/3}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(-1/3),x]`

output `(-3*c*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[a + b*x]^2]*Sin[a + b*x])/(4*b*(c*Sec[a + b*x])^(4/3)*Sqrt[Sin[a + b*x]^2])`

## 3.34.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.34.4 Maple [F]

$$\int \frac{1}{(c \sec(bx + a))^{\frac{1}{3}}} dx$$

input `int(1/(c*sec(b*x+a))^(1/3),x)`

output `int(1/(c*sec(b*x+a))^(1/3),x)`

## 3.34.5 Fricas [F]

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{1}{3}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(1/3),x, algorithm="fricas")`

output `integral((c*sec(b*x + a))^(2/3)/(c*sec(b*x + a)), x)`

**3.34.6 Sympy [F]**

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx$$

input `integrate(1/(c*sec(b*x+a))**(1/3),x)`

output `Integral((c*sec(a + b*x))**(-1/3), x)`

**3.34.7 Maxima [F]**

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{1}{3}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(1/3),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(1/3), x)`

**3.34.8 Giac [F]**

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{1}{3}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(1/3),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(1/3), x)`

**3.34.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{1/3}} dx$$

input `int(1/(c/cos(a + b*x))^(1/3),x)`output `int(1/(c/cos(a + b*x))^(1/3), x)`

### 3.35 $\int \frac{1}{(c \sec(a+bx))^{2/3}} dx$

3.35.1	Optimal result	296
3.35.2	Mathematica [A] (verified)	296
3.35.3	Rubi [A] (verified)	297
3.35.4	Maple [F]	298
3.35.5	Fricas [F]	298
3.35.6	Sympy [F]	299
3.35.7	Maxima [F]	299
3.35.8	Giac [F]	299
3.35.9	Mupad [F(-1)]	300

#### 3.35.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(c \sec(a+bx))^{2/3}} dx = -\frac{3c \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{5b(c \sec(a+bx))^{5/3} \sqrt{\sin^2(a+bx)}}$$

output `-3/5*c*hypergeom([1/2, 5/6],[11/6],cos(b*x+a)^2)*sin(b*x+a)/b/(c*sec(b*x+a))^(5/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.35.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.02

$$\int \frac{1}{(c \sec(a+bx))^{2/3}} dx = \frac{3 \cot(a+bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \sec^2(a+bx)\right) \sqrt{-\tan^2(a+bx)}}{2b(c \sec(a+bx))^{2/3}}$$

input `Integrate[(c*Sec[a + b*x])^(-2/3),x]`

output `(-3*Cot[a + b*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(2*b*(c*Sec[a + b*x])^(2/3))`

**3.35.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{2/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \csc(a + bx + \frac{\pi}{2}))^{2/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \left( \frac{\cos(a + bx)}{c} \right)^{2/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \left( \frac{\sin(a + bx + \frac{\pi}{2})}{c} \right)^{2/3} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3c \sin(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right)}{5b \sqrt{\sin^2(a + bx)} (c \sec(a + bx))^{5/3}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(-2/3),x]`

output `(-3*c*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[a + b*x]^2]*Sin[a + b*x])/(5*b*(c*Sec[a + b*x])^(5/3)*Sqrt[Sin[a + b*x]^2])`

## 3.35.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.35.4 Maple [F]

$$\int \frac{1}{(c \sec(bx + a))^{\frac{2}{3}}} dx$$

input `int(1/(c*sec(b*x+a))^(2/3),x)`

output `int(1/(c*sec(b*x+a))^(2/3),x)`

## 3.35.5 Fricas [F]

$$\int \frac{1}{(c \sec(a + bx))^{2/3}} dx = \int \frac{1}{(c \sec(bx + a))^{\frac{2}{3}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(2/3),x, algorithm="fricas")`

output `integral((c*sec(b*x + a))^(1/3)/(c*sec(b*x + a)), x)`

**3.35.6 Sympy [F]**

$$\int \frac{1}{(c \sec(a + bx))^{2/3}} dx = \int \frac{1}{(c \sec(a + bx))^{2/3}} dx$$

input `integrate(1/(c*sec(b*x+a))**(2/3), x)`

output `Integral((c*sec(a + b*x))**(-2/3), x)`

**3.35.7 Maxima [F]**

$$\int \frac{1}{(c \sec(a + bx))^{2/3}} dx = \int \frac{1}{(c \sec(bx + a))^{2/3}} dx$$

input `integrate(1/(c*sec(b*x+a))^(2/3), x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(2/3), x)`

**3.35.8 Giac [F]**

$$\int \frac{1}{(c \sec(a + bx))^{2/3}} dx = \int \frac{1}{(c \sec(bx + a))^{2/3}} dx$$

input `integrate(1/(c*sec(b*x+a))^(2/3), x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(2/3), x)`



**3.35.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c \sec(a + bx))^{2/3}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{2/3}} dx$$

input `int(1/(c/cos(a + b*x))^(2/3),x)`output `int(1/(c/cos(a + b*x))^(2/3), x)`

### 3.36 $\int \frac{1}{(c \sec(a+bx))^{4/3}} dx$

3.36.1	Optimal result	301
3.36.2	Mathematica [A] (verified)	301
3.36.3	Rubi [A] (verified)	302
3.36.4	Maple [F]	303
3.36.5	Fricas [F]	303
3.36.6	Sympy [F]	304
3.36.7	Maxima [F]	304
3.36.8	Giac [F]	304
3.36.9	Mupad [F(-1)]	305

#### 3.36.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(c \sec(a+bx))^{4/3}} dx = -\frac{3c \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{7b(c \sec(a+bx))^{7/3} \sqrt{\sin^2(a+bx)}}$$

output `-3/7*c*hypergeom([1/2, 7/6],[13/6],cos(b*x+a)^2)*sin(b*x+a)/b/(c*sec(b*x+a))^(7/3)/(sin(b*x+a)^2)^(1/2)`

#### 3.36.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.02

$$\int \frac{1}{(c \sec(a+bx))^{4/3}} dx = \frac{3 \cot(a+bx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \sec^2(a+bx)\right) \sqrt{-\tan^2(a+bx)}}{4b(c \sec(a+bx))^{4/3}}$$

input `Integrate[(c*Sec[a + b*x])^(-4/3),x]`

output `(-3*Cot[a + b*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(4*b*(c*Sec[a + b*x])^(4/3))`

**3.36.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \csc(a + bx + \frac{\pi}{2}))^{4/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & \left(\frac{\cos(a + bx)}{c}\right)^{2/3} (c \sec(a + bx))^{2/3} \int \left(\frac{\cos(a + bx)}{c}\right)^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \left(\frac{\cos(a + bx)}{c}\right)^{2/3} (c \sec(a + bx))^{2/3} \int \left(\frac{\sin(a + bx + \frac{\pi}{2})}{c}\right)^{4/3} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3c \sin(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right)}{7b \sqrt{\sin^2(a + bx)} (c \sec(a + bx))^{7/3}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^(-4/3),x]`

output `(-3*c*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[a + b*x]^2]*Sin[a + b*x])/(7*b*(c*Sec[a + b*x])^(7/3)*Sqrt[Sin[a + b*x]^2])`

## 3.36.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.36.4 Maple [F]

$$\int \frac{1}{(c \sec (bx + a))^{\frac{4}{3}}} dx$$

input `int(1/(c*sec(b*x+a))^(4/3),x)`

output `int(1/(c*sec(b*x+a))^(4/3),x)`

## 3.36.5 Fricas [F]

$$\int \frac{1}{(c \sec (a + bx))^{\frac{4}{3}}} dx = \int \frac{1}{(c \sec (bx + a))^{\frac{4}{3}}} dx$$

input `integrate(1/(c*sec(b*x+a))^(4/3),x, algorithm="fricas")`

output `integral((c*sec(b*x + a))^(2/3)/(c^2*sec(b*x + a)^2), x)`

**3.36.6 Sympy [F]**

$$\int \frac{1}{(c \sec(a + bx))^{4/3}} dx = \int \frac{1}{(c \sec(a + bx))^{4/3}} dx$$

input `integrate(1/(c*sec(b*x+a))**(4/3), x)`

output `Integral((c*sec(a + b*x))**(-4/3), x)`

**3.36.7 Maxima [F]**

$$\int \frac{1}{(c \sec(a + bx))^{4/3}} dx = \int \frac{1}{(c \sec(bx + a))^{4/3}} dx$$

input `integrate(1/(c*sec(b*x+a))^(4/3), x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(4/3), x)`

**3.36.8 Giac [F]**

$$\int \frac{1}{(c \sec(a + bx))^{4/3}} dx = \int \frac{1}{(c \sec(bx + a))^{4/3}} dx$$

input `integrate(1/(c*sec(b*x+a))^(4/3), x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(4/3), x)`

**3.36.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(c \sec(a + bx))^{4/3}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{4/3}} dx$$

input `int(1/(c/cos(a + b*x))^(4/3),x)`output `int(1/(c/cos(a + b*x))^(4/3), x)`

### 3.37 $\int \sec^n(a + bx) dx$

3.37.1	Optimal result . . . . .	306
3.37.2	Mathematica [A] (verified) . . . . .	306
3.37.3	Rubi [A] (verified) . . . . .	307
3.37.4	Maple [F] . . . . .	308
3.37.5	Fricas [F] . . . . .	308
3.37.6	Sympy [F] . . . . .	308
3.37.7	Maxima [F] . . . . .	309
3.37.8	Giac [F] . . . . .	309
3.37.9	Mupad [F(-1)] . . . . .	309

#### 3.37.1 Optimal result

Integrand size = 8, antiderivative size = 70

$$\int \sec^n(a + bx) dx = \frac{\text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \cos^2(a + bx)\right) \sec^{-1+n}(a + bx) \sin(a + bx)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

output `-hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*sec(b*x+a)^(-1+n)*sin(b*x+a)/b/(1-n)/(sin(b*x+a)^2)^(1/2)`

#### 3.37.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.87

$$\int \sec^n(a + bx) dx = \frac{\csc(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n}{2}, \frac{2+n}{2}, \sec^2(a + bx)\right) \sec^{-1+n}(a + bx) \sqrt{-\tan^2(a + bx)}}{bn}$$

input `Integrate[Sec[a + b*x]^n,x]`

output `(Csc[a + b*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Sec[a + b*x]^2]*Sec[a + b*x]^(-1 + n)*Sqrt[-Tan[a + b*x]^2])/(b*n)`

### 3.37.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^n(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(a + bx + \frac{\pi}{2}\right)^n dx \\
 & \quad \downarrow \text{4259} \\
 & \cos^n(a + bx) \sec^n(a + bx) \int \cos^{-n}(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \cos^n(a + bx) \sec^n(a + bx) \int \sin\left(a + bx + \frac{\pi}{2}\right)^{-n} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{\sin(a + bx) \sec^{n-1}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \cos^2(a + bx)\right)}{b(1-n)\sqrt{\sin^2(a + bx)}}
 \end{aligned}$$

input `Int[Sec[a + b*x]^n,x]`

output `-((Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*Sec[a + b*x]^(-1 + n)*Sin[a + b*x])/(b*(1 - n)*Sqrt[Sin[a + b*x]^2]))`

#### 3.37.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`



rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### 3.37.4 Maple [F]

$$\int \sec (bx + a)^n dx$$

input `int(sec(b*x+a)^n,x)`

output `int(sec(b*x+a)^n,x)`

### 3.37.5 Fricas [F]

$$\int \sec^n(a + bx) dx = \int \sec (bx + a)^n dx$$

input `integrate(sec(b*x+a)^n,x, algorithm="fricas")`

output `integral(sec(b*x + a)^n, x)`

### 3.37.6 Sympy [F]

$$\int \sec^n(a + bx) dx = \int \sec^n (a + bx) dx$$

input `integrate(sec(b*x+a)**n,x)`

output `Integral(sec(a + b*x)**n, x)`

**3.37.7 Maxima [F]**

$$\int \sec^n(a + bx) dx = \int \sec (bx + a)^n dx$$

input `integrate(sec(b*x+a)^n,x, algorithm="maxima")`

output `integrate(sec(b*x + a)^n, x)`

**3.37.8 Giac [F]**

$$\int \sec^n(a + bx) dx = \int \sec (bx + a)^n dx$$

input `integrate(sec(b*x+a)^n,x, algorithm="giac")`

output `integrate(sec(b*x + a)^n, x)`

**3.37.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^n(a + bx) dx = \int \left( \frac{1}{\cos(a + bx)} \right)^n dx$$

input `int((1/cos(a + b*x))^n,x)`

output `int((1/cos(a + b*x))^n, x)`

### 3.38 $\int (c \sec(a + bx))^n dx$

3.38.1	Optimal result . . . . .	310
3.38.2	Mathematica [A] (verified) . . . . .	310
3.38.3	Rubi [A] (verified) . . . . .	311
3.38.4	Maple [F] . . . . .	312
3.38.5	Fricas [F] . . . . .	312
3.38.6	Sympy [F] . . . . .	313
3.38.7	Maxima [F] . . . . .	313
3.38.8	Giac [F] . . . . .	313
3.38.9	Mupad [F(-1)] . . . . .	314

#### 3.38.1 Optimal result

Integrand size = 10, antiderivative size = 73

$$\int (c \sec(a + bx))^n dx = -\frac{c \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \cos^2(a + bx)\right) (c \sec(a + bx))^{-1+n} \sin(a + bx)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

```
output -c*hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*(c*sec(b*x+a))^( -1+n)*sin(b*x+a)/b/(1-n)/(sin(b*x+a)^2)^(1/2)
```

#### 3.38.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.84

$$\int (c \sec(a + bx))^n dx = \frac{\cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n}{2}, \frac{2+n}{2}, \sec^2(a + bx)\right) (c \sec(a + bx))^n \sqrt{-\tan^2(a + bx)}}{bn}$$

```
input Integrate[(c*Sec[a + b*x])^n,x]
```

```
output (Cot[a + b*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Sec[a + b*x]^2]*(c*Sec[a + b*x])^n*Sqrt[-Tan[a + b*x]^2])/(b*n)
```

**3.38.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( c \csc \left( a + bx + \frac{\pi}{2} \right) \right)^n dx \\
 & \quad \downarrow \text{4259} \\
 & \left( \frac{\cos(a + bx)}{c} \right)^n (c \sec(a + bx))^n \int \left( \frac{\cos(a + bx)}{c} \right)^{-n} dx \\
 & \quad \downarrow \text{3042} \\
 & \left( \frac{\cos(a + bx)}{c} \right)^n (c \sec(a + bx))^n \int \left( \frac{\sin(a + bx + \frac{\pi}{2})}{c} \right)^{-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{c \sin(a + bx) (c \sec(a + bx))^{n-1} \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \cos^2(a + bx) \right)}{b(1-n) \sqrt{\sin^2(a + bx)}}
 \end{aligned}$$

input `Int[(c*Sec[a + b*x])^n,x]`

output `-((c*Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*(c*Sec[a + b*x])^(-1 + n)*Sin[a + b*x])/(b*(1 - n)*Sqrt[Sin[a + b*x]^2]))`

## 3.38.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.38.4 Maple [F]

$$\int (c \sec (bx + a))^n dx$$

input `int((c*sec(b*x+a))^n,x)`

output `int((c*sec(b*x+a))^n,x)`

## 3.38.5 Fricas [F]

$$\int (c \sec (a + bx))^n dx = \int (c \sec (bx + a))^n dx$$

input `integrate((c*sec(b*x+a))^n,x, algorithm="fricas")`

output `integral((c*sec(b*x + a))^n, x)`

**3.38.6 Sympy [F]**

$$\int (c \sec(a + bx))^n dx = \int (c \sec(a + bx))^n dx$$

input `integrate((c*sec(b*x+a))**n,x)`

output `Integral((c*sec(a + b*x))**n, x)`

**3.38.7 Maxima [F]**

$$\int (c \sec(a + bx))^n dx = \int (c \sec(bx + a))^n dx$$

input `integrate((c*sec(b*x+a))^n,x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^n, x)`

**3.38.8 Giac [F]**

$$\int (c \sec(a + bx))^n dx = \int (c \sec(bx + a))^n dx$$

input `integrate((c*sec(b*x+a))^n,x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^n, x)`

**3.38.9 Mupad [F(-1)]**

Timed out.

$$\int (c \sec(a + bx))^n dx = \int \left( \frac{c}{\cos(a + bx)} \right)^n dx$$

input `int((c/cos(a + b*x))^n,x)`output `int((c/cos(a + b*x))^n, x)`

### 3.39 $\int \sec^2(x)^{7/2} dx$

3.39.1	Optimal result . . . . .	315
3.39.2	Mathematica [A] (verified) . . . . .	315
3.39.3	Rubi [A] (verified) . . . . .	316
3.39.4	Maple [C] (warning: unable to verify) . . . . .	317
3.39.5	Fricas [A] (verification not implemented) . . . . .	318
3.39.6	Sympy [F(-1)] . . . . .	318
3.39.7	Maxima [A] (verification not implemented) . . . . .	318
3.39.8	Giac [A] (verification not implemented) . . . . .	319
3.39.9	Mupad [F(-1)] . . . . .	319

#### 3.39.1 Optimal result

Integrand size = 8, antiderivative size = 50

$$\int \sec^2(x)^{7/2} dx = \frac{5}{16} \operatorname{arcsinh}(\tan(x)) + \frac{5}{16} \sqrt{\sec^2(x)} \tan(x) + \frac{5}{24} \sec^2(x)^{3/2} \tan(x) + \frac{1}{6} \sec^2(x)^{5/2} \tan(x)$$

output `5/16*arcsinh(tan(x))+5/24*(sec(x)^2)^(3/2)*tan(x)+1/6*(sec(x)^2)^(5/2)*tan(x)+5/16*(sec(x)^2)^(1/2)*tan(x)`

#### 3.39.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.78

$$\int \sec^2(x)^{7/2} dx = \frac{\sec(x) (15 \operatorname{arctanh}(\sin(x)) + \sec(x) (15 + 10 \sec^2(x) + 8 \sec^4(x)) \tan(x))}{48 \sqrt{\sec^2(x)}}$$

input `Integrate[(Sec[x]^2)^(7/2),x]`

output `(Sec[x]*(15*ArcTanh[Sin[x]] + Sec[x]*(15 + 10*Sec[x]^2 + 8*Sec[x]^4)*Tan[x]))/(48*sqrt[Sec[x]^2])`



**3.39.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.32, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 4610, 211, 211, 211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(x)^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (\sec(x)^2)^{7/2} dx \\
 & \quad \downarrow \text{4610} \\
 & \int (\tan^2(x) + 1)^{5/2} d \tan(x) \\
 & \quad \downarrow \text{211} \\
 & \frac{5}{6} \int (\tan^2(x) + 1)^{3/2} d \tan(x) + \frac{1}{6} \tan(x) (\tan^2(x) + 1)^{5/2} \\
 & \quad \downarrow \text{211} \\
 & \frac{5}{6} \left( \frac{3}{4} \int \sqrt{\tan^2(x) + 1} d \tan(x) + \frac{1}{4} \tan(x) (\tan^2(x) + 1)^{3/2} \right) + \frac{1}{6} \tan(x) (\tan^2(x) + 1)^{5/2} \\
 & \quad \downarrow \text{211} \\
 & \frac{5}{6} \left( \frac{3}{4} \left( \frac{1}{2} \int \frac{1}{\sqrt{\tan^2(x) + 1}} d \tan(x) + \frac{1}{2} \sqrt{\tan^2(x) + 1} \tan(x) \right) + \frac{1}{4} \tan(x) (\tan^2(x) + 1)^{3/2} \right) + \\
 & \quad \frac{1}{6} \tan(x) (\tan^2(x) + 1)^{5/2} \\
 & \quad \downarrow \text{222} \\
 & \frac{5}{6} \left( \frac{3}{4} \left( \frac{1}{2} \operatorname{arcsinh}(\tan(x)) + \frac{1}{2} \tan(x) \sqrt{\tan^2(x) + 1} \right) + \frac{1}{4} \tan(x) (\tan^2(x) + 1)^{3/2} \right) + \\
 & \quad \frac{1}{6} \tan(x) (\tan^2(x) + 1)^{5/2}
 \end{aligned}$$

input `Int[(Sec[x]^2)^(7/2),x]`

output `(Tan[x]*(1 + Tan[x]^2)^(5/2))/6 + (5*((Tan[x]*(1 + Tan[x]^2)^(3/2))/4 + (3*(ArcSinh[Tan[x]]/2 + (Tan[x]*Sqrt[1 + Tan[x]^2])/2))/4))/6`

## 3.39.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

## 3.39.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.97 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.02

method	result
default	$\frac{\operatorname{csgn}(\sec(x)) \left( 15 \ln(-\cot(x) + \csc(x) + 1) - 15 \ln(-\cot(x) + \csc(x) - 1) + 15 \sec(x) \tan(x) + 10 \tan(x) \sec(x)^3 + 8 \tan(x) \sec(x)^5 \right)}{48}$
risch	$-\frac{i \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} (15 e^{10ix} + 85 e^{8ix} + 198 e^{6ix} - 198 e^{4ix} - 85 e^{2ix} - 15)}{24(e^{2ix}+1)^5} - \frac{5 \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix}-i) \cos(x)}{8} + \frac{5 \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix}+i)}{8}$

input `int((sec(x)^2)^(7/2), x, method=_RETURNVERBOSE)`

output `1/48*csgn(sec(x))*(15*ln(-cot(x)+csc(x)+1)-15*ln(-cot(x)+csc(x)-1)+15*sec(x)*tan(x)+10*tan(x)*sec(x)^3+8*tan(x)*sec(x)^5)`

**3.39.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.98

$$\int \sec^2(x)^{7/2} dx = \frac{15 \cos(x)^6 \log(\sin(x) + 1) - 15 \cos(x)^6 \log(-\sin(x) + 1) + 2(15 \cos(x)^4 + 10 \cos(x)^2 + 8) \sin(x)}{96 \cos(x)^6}$$

input `integrate((sec(x)^2)^(7/2),x, algorithm="fracas")`

output `-1/96*(15*cos(x)^6*log(sin(x) + 1) - 15*cos(x)^6*log(-sin(x) + 1) + 2*(15*cos(x)^4 + 10*cos(x)^2 + 8)*sin(x))/cos(x)^6`

**3.39.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^2(x)^{7/2} dx = \text{Timed out}$$

input `integrate((sec(x)**2)**(7/2),x)`

output `Timed out`

**3.39.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.84

$$\int \sec^2(x)^{7/2} dx = \frac{1}{6} (\tan(x)^2 + 1)^{5/2} \tan(x) + \frac{5}{24} (\tan(x)^2 + 1)^{3/2} \tan(x) + \frac{5}{16} \sqrt{\tan(x)^2 + 1} \tan(x) + \frac{5}{16} \operatorname{arsinh}(\tan(x))$$

input `integrate((sec(x)^2)^(7/2),x, algorithm="maxima")`

output `1/6*(tan(x)^2 + 1)^(5/2)*tan(x) + 5/24*(tan(x)^2 + 1)^(3/2)*tan(x) + 5/16*sqrt(tan(x)^2 + 1)*tan(x) + 5/16*arcsinh(tan(x))`

**3.39.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.18

$$\int \sec^2(x)^{7/2} dx = \frac{5 \log(\sin(x) + 1)}{32 \operatorname{sgn}(\cos(x))} - \frac{5 \log(-\sin(x) + 1)}{32 \operatorname{sgn}(\cos(x))} - \frac{15 \sin(x)^5 - 40 \sin(x)^3 + 33 \sin(x)}{48 (\sin(x)^2 - 1)^3 \operatorname{sgn}(\cos(x))}$$

input `integrate((sec(x)^2)^(7/2),x, algorithm="giac")`

output `5/32*log(sin(x) + 1)/sgn(cos(x)) - 5/32*log(-sin(x) + 1)/sgn(cos(x)) - 1/48*(15*sin(x)^5 - 40*sin(x)^3 + 33*sin(x))/((sin(x)^2 - 1)^3*sgn(cos(x)))`

**3.39.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(x)^{7/2} dx = \int \left( \frac{1}{\cos(x)^2} \right)^{7/2} dx$$

input `int((1/cos(x)^2)^(7/2),x)`

output `int((1/cos(x)^2)^(7/2), x)`

### 3.40 $\int \sec^2(x)^{5/2} dx$

3.40.1	Optimal result . . . . .	320
3.40.2	Mathematica [A] (verified) . . . . .	320
3.40.3	Rubi [A] (verified) . . . . .	321
3.40.4	Maple [C] (warning: unable to verify) . . . . .	322
3.40.5	Fricas [A] (verification not implemented) . . . . .	323
3.40.6	Sympy [F] . . . . .	323
3.40.7	Maxima [A] (verification not implemented) . . . . .	323
3.40.8	Giac [B] (verification not implemented) . . . . .	324
3.40.9	Mupad [F(-1)] . . . . .	324

#### 3.40.1 Optimal result

Integrand size = 8, antiderivative size = 36

$$\int \sec^2(x)^{5/2} dx = \frac{3}{8} \operatorname{arcsinh}(\tan(x)) + \frac{3}{8} \sqrt{\sec^2(x)} \tan(x) + \frac{1}{4} \sec^2(x)^{3/2} \tan(x)$$

output `3/8*arcsinh(tan(x))+1/4*(sec(x)^2)^(3/2)*tan(x)+3/8*(sec(x)^2)^(1/2)*tan(x)`

#### 3.40.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \sec^2(x)^{5/2} dx = \frac{\sec(x) (3 \operatorname{arctanh}(\sin(x)) + \sec(x) (3 + 2 \sec^2(x)) \tan(x))}{8 \sqrt{\sec^2(x)}}$$

input `Integrate[(Sec[x]^2)^(5/2),x]`

output `(Sec[x]*(3*ArcTanh[Sin[x]] + Sec[x]*(3 + 2*Sec[x]^2)*Tan[x]))/(8*Sqrt[Sec[x]^2])`

**3.40.3 Rubi [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.25, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 4610, 211, 211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(x)^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (\sec(x)^2)^{5/2} dx \\
 & \quad \downarrow \text{4610} \\
 & \int (\tan^2(x) + 1)^{3/2} d \tan(x) \\
 & \quad \downarrow \text{211} \\
 & \frac{3}{4} \int \sqrt{\tan^2(x) + 1} d \tan(x) + \frac{1}{4} \tan(x) (\tan^2(x) + 1)^{3/2} \\
 & \quad \downarrow \text{211} \\
 & \frac{3}{4} \left( \frac{1}{2} \int \frac{1}{\sqrt{\tan^2(x) + 1}} d \tan(x) + \frac{1}{2} \sqrt{\tan^2(x) + 1} \tan(x) \right) + \frac{1}{4} \tan(x) (\tan^2(x) + 1)^{3/2} \\
 & \quad \downarrow \text{222} \\
 & \frac{3}{4} \left( \frac{1}{2} \operatorname{arcsinh}(\tan(x)) + \frac{1}{2} \tan(x) \sqrt{\tan^2(x) + 1} \right) + \frac{1}{4} \tan(x) (\tan^2(x) + 1)^{3/2}
 \end{aligned}$$

input `Int[(Sec[x]^2)^(5/2),x]`

output `(Tan[x]*(1 + Tan[x]^2)^(3/2))/4 + (3*(ArcSinh[Tan[x]]/2 + (Tan[x]*Sqrt[1 + Tan[x]^2])/2))/4`

## 3.40.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

## 3.40.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.80 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.19

method	result	size
default	$\frac{\text{csgn}(\sec(x)) \left( 3 \ln(-\cot(x) + \csc(x) + 1) - 3 \ln(-\cot(x) + \csc(x) - 1) + 3 \sec(x) \tan(x) + 2 \tan(x) \sec(x)^3 \right)}{8}$	43
risch	$-\frac{i \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} (3e^{6ix} + 11e^{4ix} - 11e^{2ix} - 3)}{4(e^{2ix}+1)^3} - \frac{3 \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix} - i) \cos(x)}{4} + \frac{3 \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix} + i) \cos(x)}{4}$	114

input `int((sec(x)^2)^(5/2), x, method=_RETURNVERBOSE)`

output `1/8*csgn(sec(x))*(3*ln(-cot(x)+csc(x)+1)-3*ln(-cot(x)+csc(x)-1)+3*sec(x)*tan(x)+2*tan(x)*sec(x)^3)`

**3.40.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.19

$$\int \sec^2(x)^{5/2} dx = \frac{3 \cos(x)^4 \log(\sin(x) + 1) - 3 \cos(x)^4 \log(-\sin(x) + 1) + 2(3 \cos(x)^2 + 2) \sin(x)}{16 \cos(x)^4}$$

input `integrate((sec(x)^2)^(5/2),x, algorithm="fracas")`

output `-1/16*(3*cos(x)^4*log(sin(x) + 1) - 3*cos(x)^4*log(-sin(x) + 1) + 2*(3*cos(x)^2 + 2)*sin(x))/cos(x)^4`

**3.40.6 Sympy [F]**

$$\int \sec^2(x)^{5/2} dx = \int (\sec^2(x))^{\frac{5}{2}} dx$$

input `integrate((sec(x)**2)**(5/2),x)`

output `Integral((sec(x)**2)**(5/2), x)`

**3.40.7 Maxima [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \sec^2(x)^{5/2} dx = \frac{1}{4} (\tan(x)^2 + 1)^{\frac{3}{2}} \tan(x) + \frac{3}{8} \sqrt{\tan(x)^2 + 1} \tan(x) + \frac{3}{8} \operatorname{arsinh}(\tan(x))$$

input `integrate((sec(x)^2)^(5/2),x, algorithm="maxima")`

output `1/4*(tan(x)^2 + 1)^(3/2)*tan(x) + 3/8*sqrt(tan(x)^2 + 1)*tan(x) + 3/8*arcsinh(tan(x))`



**3.40.8 Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 53 vs.  $2(26) = 52$ .

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.47

$$\int \sec^2(x)^{5/2} dx = \frac{3 \log(\sin(x) + 1)}{16 \operatorname{sgn}(\cos(x))} - \frac{3 \log(-\sin(x) + 1)}{16 \operatorname{sgn}(\cos(x))} - \frac{3 \sin(x)^3 - 5 \sin(x)}{8 (\sin(x)^2 - 1)^2 \operatorname{sgn}(\cos(x))}$$

input `integrate((sec(x)^2)^(5/2),x, algorithm="giac")`

output `3/16*log(sin(x) + 1)/sgn(cos(x)) - 3/16*log(-sin(x) + 1)/sgn(cos(x)) - 1/8*(3*sin(x)^3 - 5*sin(x))/((sin(x)^2 - 1)^2*sgn(cos(x)))`

**3.40.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(x)^{5/2} dx = \int \left( \frac{1}{\cos(x)^2} \right)^{5/2} dx$$

input `int((1/cos(x)^2)^(5/2),x)`

output `int((1/cos(x)^2)^(5/2), x)`

### 3.41 $\int \sec^2(x)^{3/2} dx$

3.41.1	Optimal result . . . . .	325
3.41.2	Mathematica [A] (verified) . . . . .	325
3.41.3	Rubi [A] (verified) . . . . .	326
3.41.4	Maple [C] (warning: unable to verify) . . . . .	327
3.41.5	Fricas [B] (verification not implemented) . . . . .	327
3.41.6	Sympy [F] . . . . .	328
3.41.7	Maxima [A] (verification not implemented) . . . . .	328
3.41.8	Giac [B] (verification not implemented) . . . . .	328
3.41.9	Mupad [F(-1)] . . . . .	329

#### 3.41.1 Optimal result

Integrand size = 8, antiderivative size = 22

$$\int \sec^2(x)^{3/2} dx = \frac{1}{2} \operatorname{arcsinh}(\tan(x)) + \frac{1}{2} \sqrt{\sec^2(x)} \tan(x)$$

output `1/2*arcsinh(tan(x))+1/2*(sec(x)^2)^(1/2)*tan(x)`

#### 3.41.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.05

$$\int \sec^2(x)^{3/2} dx = \frac{\sec(x)(\operatorname{arctanh}(\sin(x)) + \sec(x) \tan(x))}{2\sqrt{\sec^2(x)}}$$

input `Integrate[(Sec[x]^2)^(3/2),x]`

output `(Sec[x]*(ArcTanh[Sin[x]] + Sec[x]*Tan[x]))/(2*Sqrt[Sec[x]^2])`

### 3.41.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4610, 211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(x)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (\sec(x)^2)^{3/2} dx \\
 & \quad \downarrow \text{4610} \\
 & \int \sqrt{\tan^2(x) + 1} d \tan(x) \\
 & \quad \downarrow \text{211} \\
 & \frac{1}{2} \int \frac{1}{\sqrt{\tan^2(x) + 1}} d \tan(x) + \frac{1}{2} \sqrt{\tan^2(x) + 1} \tan(x) \\
 & \quad \downarrow \text{222} \\
 & \frac{1}{2} \operatorname{arcsinh}(\tan(x)) + \frac{1}{2} \tan(x) \sqrt{\tan^2(x) + 1}
 \end{aligned}$$

input `Int[(Sec[x]^2)^(3/2),x]`

output `ArcSinh[Tan[x]]/2 + (Tan[x]*Sqrt[1 + Tan[x]^2])/2`

#### 3.41.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4610 Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

### 3.41.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.45

method	result	size
default	$\frac{\text{csgn}(\sec(x))(\ln(-\cot(x)+\csc(x)+1)-\ln(-\cot(x)+\csc(x)-1)+\sec(x)\tan(x))}{2}$	32
risch	$-\frac{i\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}-1)}{e^{2ix}+1} - \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix}-i)\cos(x) + \sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix}+i)\cos(x)$	97

```
input int((sec(x)^2)^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/2*csgn(sec(x))*(ln(-cot(x)+csc(x)+1)-ln(-cot(x)+csc(x)-1)+sec(x)*tan(x))
```

### 3.41.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs.  $2(16) = 32$ .

Time = 0.28 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55

$$\int \sec^2(x)^{3/2} dx = -\frac{\cos(x)^2 \log(\sin(x) + 1) - \cos(x)^2 \log(-\sin(x) + 1) + 2 \sin(x)}{4 \cos(x)^2}$$

```
input integrate((sec(x)^2)^(3/2),x, algorithm="fracas")
```

```
output -1/4*(cos(x)^2*log(sin(x) + 1) - cos(x)^2*log(-sin(x) + 1) + 2*sin(x))/cos
(x)^2
```

### 3.41.6 Sympy [F]

$$\int \sec^2(x)^{3/2} dx = \int (\sec^2(x))^{\frac{3}{2}} dx$$

input `integrate((sec(x)**2)**(3/2),x)`

output `Integral((sec(x)**2)**(3/2), x)`

### 3.41.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \sec^2(x)^{3/2} dx = \frac{1}{2} \sqrt{\tan(x)^2 + 1} \tan(x) + \frac{1}{2} \operatorname{arsinh}(\tan(x))$$

input `integrate((sec(x)^2)^(3/2),x, algorithm="maxima")`

output `1/2*sqrt(tan(x)^2 + 1)*tan(x) + 1/2*arcsinh(tan(x))`

### 3.41.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. 2(16) = 32.

Time = 0.29 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.00

$$\int \sec^2(x)^{3/2} dx = \frac{\log(\sin(x) + 1)}{4 \operatorname{sgn}(\cos(x))} - \frac{\log(-\sin(x) + 1)}{4 \operatorname{sgn}(\cos(x))} - \frac{\sin(x)}{2(\sin(x)^2 - 1) \operatorname{sgn}(\cos(x))}$$

input `integrate((sec(x)^2)^(3/2),x, algorithm="giac")`

output `1/4*log(sin(x) + 1)/sgn(cos(x)) - 1/4*log(-sin(x) + 1)/sgn(cos(x)) - 1/2*sin(x)/((sin(x)^2 - 1)*sgn(cos(x)))`

**3.41.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(x)^{3/2} dx = \int \left( \frac{1}{\cos(x)^2} \right)^{3/2} dx$$

input `int((1/cos(x)^2)^(3/2),x)`output `int((1/cos(x)^2)^(3/2), x)`

### 3.42 $\int \sqrt{\sec^2(x)} dx$

3.42.1	Optimal result . . . . .	330
3.42.2	Mathematica [B] (verified) . . . . .	330
3.42.3	Rubi [A] (verified) . . . . .	331
3.42.4	Maple [C] (warning: unable to verify) . . . . .	332
3.42.5	Fricas [B] (verification not implemented) . . . . .	332
3.42.6	Sympy [F] . . . . .	332
3.42.7	Maxima [A] (verification not implemented) . . . . .	333
3.42.8	Giac [B] (verification not implemented) . . . . .	333
3.42.9	Mupad [F(-1)] . . . . .	333

#### 3.42.1 Optimal result

Integrand size = 8, antiderivative size = 3

$$\int \sqrt{\sec^2(x)} dx = \operatorname{arcsinh}(\tan(x))$$

output `arcsinh(tan(x))`

#### 3.42.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 14 vs. 2(3) = 6.

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 4.67

$$\int \sqrt{\sec^2(x)} dx = \operatorname{arctanh}(\sin(x)) \cos(x) \sqrt{\sec^2(x)}$$

input `Integrate[Sqrt[Sec[x]^2], x]`

output `ArcTanh[Sin[x]]*Cos[x]*Sqrt[Sec[x]^2]`

### 3.42.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 4610, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sqrt{\sec^2(x)} dx \\
 \downarrow \text{3042} \\
 \int \sqrt{\sec(x)^2} dx \\
 \downarrow \text{4610} \\
 \int \frac{1}{\sqrt{\tan^2(x) + 1}} d \tan(x) \\
 \downarrow \text{222} \\
 \operatorname{arcsinh}(\tan(x))
 \end{array}$$

input `Int[Sqrt[Sec[x]^2], x]`

output `ArcSinh[Tan[x]]`

#### 3.42.3.1 Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^2]^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`



**3.42.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.38 (sec) , antiderivative size = 14, normalized size of antiderivative = 4.67

method	result	size
default	$-2 \operatorname{csgn}(\sec(x)) \operatorname{arctanh}(\cot(x) - \csc(x))$	14
risch	$2\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix} + i) \cos(x) - 2\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix} - i) \cos(x)$	62

input `int((sec(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*csgn(sec(x))*arctanh(cot(x)-csc(x))`

**3.42.5 Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 17 vs.  $2(3) = 6$ .

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 5.67

$$\int \sqrt{\sec^2(x)} dx = -\frac{1}{2} \log(\sin(x) + 1) + \frac{1}{2} \log(-\sin(x) + 1)$$

input `integrate((sec(x)^2)^(1/2),x, algorithm="fricas")`

output `-1/2*log(sin(x) + 1) + 1/2*log(-sin(x) + 1)`

**3.42.6 Sympy [F]**

$$\int \sqrt{\sec^2(x)} dx = \int \sqrt{\sec^2(x)} dx$$

input `integrate((sec(x)**2)**(1/2),x)`

output `Integral(sqrt(sec(x)**2), x)`

**3.42.7 Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \sqrt{\sec^2(x)} dx = \operatorname{arsinh}(\tan(x))$$

input `integrate((sec(x)^2)^(1/2),x, algorithm="maxima")`

output `arcsinh(tan(x))`

**3.42.8 Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. 2(3) = 6.

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 11.67

$$\int \sqrt{\sec^2(x)} dx = \frac{\log\left(\left|\frac{1}{\sin(x)} + \sin(x) + 2\right|\right)}{4 \operatorname{sgn}(\cos(x))} - \frac{\log\left(\left|\frac{1}{\sin(x)} + \sin(x) - 2\right|\right)}{4 \operatorname{sgn}(\cos(x))}$$

input `integrate((sec(x)^2)^(1/2),x, algorithm="giac")`

output `1/4*log(abs(1/sin(x) + sin(x) + 2))/sgn(cos(x)) - 1/4*log(abs(1/sin(x) + sin(x) - 2))/sgn(cos(x))`

**3.42.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{\sec^2(x)} dx = \int \sqrt{\frac{1}{\cos(x)^2}} dx$$

input `int((1/cos(x)^2)^(1/2),x)`

output `int((1/cos(x)^2)^(1/2), x)`

### 3.43 $\int \frac{1}{\sqrt{\sec^2(x)}} dx$

3.43.1	Optimal result	334
3.43.2	Mathematica [A] (verified)	334
3.43.3	Rubi [A] (verified)	335
3.43.4	Maple [C] (warning: unable to verify)	336
3.43.5	Fricas [A] (verification not implemented)	336
3.43.6	Sympy [A] (verification not implemented)	336
3.43.7	Maxima [A] (verification not implemented)	337
3.43.8	Giac [A] (verification not implemented)	337
3.43.9	Mupad [B] (verification not implemented)	337

#### 3.43.1 Optimal result

Integrand size = 8, antiderivative size = 11

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = \frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

output `tan(x)/(sec(x)^2)^(1/2)`

#### 3.43.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = \frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

input `Integrate[1/Sqrt[Sec[x]^2], x]`

output `Tan[x]/Sqrt[Sec[x]^2]`

### 3.43.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 4610, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{\sqrt{\sec^2(x)}} dx \\
 \downarrow \text{3042} \\
 \int \frac{1}{\sqrt{\sec(x)^2}} dx \\
 \downarrow \text{4610} \\
 \int \frac{1}{(\tan^2(x) + 1)^{3/2}} d \tan(x) \\
 \downarrow \text{208} \\
 \frac{\tan(x)}{\sqrt{\tan^2(x) + 1}}
 \end{array}$$

input `Int[1/Sqrt[Sec[x]^2], x]`

output `Tan[x]/Sqrt[1 + Tan[x]^2]`

#### 3.43.3.1 Defintions of rubi rules used

rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

---

3.43.  $\int \frac{1}{\sqrt{\sec^2(x)}} dx$

**3.43.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.36 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.64

method	result	size
default	$\text{csgn}(\sec(x)) \sin(x)$	7
risch	$-\frac{ie^{2ix}}{2\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)}} + \frac{i}{2\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)}}$	65

input `int(1/(sec(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `csgn(sec(x))*sin(x)`

**3.43.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.36

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = -\sin(x)$$

input `integrate(1/(sec(x)^2)^(1/2),x, algorithm="fracas")`

output `-sin(x)`

**3.43.6 Sympy [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = \frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

input `integrate(1/(sec(x)**2)**(1/2),x)`

output `tan(x)/sqrt(sec(x)**2)`

**3.43.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = \frac{\tan(x)}{\sqrt{\tan(x)^2 + 1}}$$

input `integrate(1/(sec(x)^2)^(1/2),x, algorithm="maxima")`output `tan(x)/sqrt(tan(x)^2 + 1)`**3.43.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = \operatorname{sgn}(\cos(x)) \sin(x)$$

input `integrate(1/(sec(x)^2)^(1/2),x, algorithm="giac")`output `sgn(cos(x))*sin(x)`**3.43.9 Mupad [B] (verification not implemented)**

Time = 12.45 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = \frac{\sqrt{2} \sin(2x)}{2 \sqrt{2 \cos(x)^2}}$$

input `int(1/(1/cos(x)^2)^(1/2),x)`output `(2^(1/2)*sin(2*x))/(2*(2*cos(x)^2)^(1/2))`

### 3.44 $\int \frac{1}{\sec^2(x)^{3/2}} dx$

3.44.1	Optimal result	338
3.44.2	Mathematica [A] (verified)	338
3.44.3	Rubi [A] (verified)	339
3.44.4	Maple [C] (warning: unable to verify)	340
3.44.5	Fricas [A] (verification not implemented)	340
3.44.6	Sympy [A] (verification not implemented)	341
3.44.7	Maxima [A] (verification not implemented)	341
3.44.8	Giac [A] (verification not implemented)	341
3.44.9	Mupad [F(-1)]	342

#### 3.44.1 Optimal result

Integrand size = 8, antiderivative size = 29

$$\int \frac{1}{\sec^2(x)^{3/2}} dx = \frac{\tan(x)}{3 \sec^2(x)^{3/2}} + \frac{2 \tan(x)}{3 \sqrt{\sec^2(x)}}$$

output  $1/3*\tan(x)/(\sec(x)^2)^{(3/2)}+2/3*\tan(x)/(\sec(x)^2)^{(1/2)}$

#### 3.44.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{1}{\sec^2(x)^{3/2}} dx = \frac{\tan(x)}{\sqrt{\sec^2(x)}} - \frac{\tan^3(x)}{3 \sec^2(x)^{3/2}}$$

input `Integrate[(Sec[x]^2)^(-3/2), x]`

output `Tan[x]/Sqrt[Sec[x]^2] - Tan[x]^3/(3*(Sec[x]^2)^(3/2))`

**3.44.3 Rubi [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4610, 209, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^2(x)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\sec(x)^2)^{3/2}} dx \\
 & \quad \downarrow \text{4610} \\
 & \int \frac{1}{(\tan^2(x) + 1)^{5/2}} d \tan(x) \\
 & \quad \downarrow \text{209} \\
 & \frac{2}{3} \int \frac{1}{(\tan^2(x) + 1)^{3/2}} d \tan(x) + \frac{\tan(x)}{3(\tan^2(x) + 1)^{3/2}} \\
 & \quad \downarrow \text{208} \\
 & \frac{2 \tan(x)}{3\sqrt{\tan^2(x) + 1}} + \frac{\tan(x)}{3(\tan^2(x) + 1)^{3/2}}
 \end{aligned}$$

input `Int[(Sec[x]^2)^(-3/2), x]`

output `Tan[x]/(3*(1 + Tan[x]^2)^(3/2)) + (2*Tan[x])/(3*Sqrt[1 + Tan[x]^2])`

**3.44.3.1 Defintions of rubi rules used**

rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] :> Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`



```
rule 209 Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)
/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1)
), x], x] /; FreeQ[{a, b}, x] && ILtQ[p + 3/2, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4610 Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

### 3.44.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.37 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.62

method	result	size
default	$\frac{\text{csgn}(\sec(x))(\sin(x)\cos(x)^2 + 2\sin(x))}{3}$	18
risch	$-\frac{ie^{4ix}}{24(e^{2ix}+1)\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}} - \frac{3ie^{2ix}}{8\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)} + \frac{3i}{8\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)} + \frac{ie^{-2ix}}{24(e^{2ix}+1)\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}}$	133

```
input int(1/(sec(x)^2)^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/3*csgn(sec(x))*(sin(x)*cos(x)^2+2*sin(x))
```

### 3.44.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.34

$$\int \frac{1}{\sec^2(x)^{3/2}} dx = -\frac{1}{3} (\cos(x)^2 + 2) \sin(x)$$

```
input integrate(1/(sec(x)^2)^(3/2),x, algorithm="fricas")
```

```
output -1/3*(cos(x)^2 + 2)*sin(x)
```

**3.44.6 Sympy [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{1}{\sec^2(x)^{3/2}} dx = \frac{2 \tan^3(x)}{3 (\sec^2(x))^{\frac{3}{2}}} + \frac{\tan(x)}{(\sec^2(x))^{\frac{3}{2}}}$$

input `integrate(1/(sec(x)**2)**(3/2),x)`output `2*tan(x)**3/(3*(sec(x)**2)**(3/2)) + tan(x)/(sec(x)**2)**(3/2)`**3.44.7 Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sec^2(x)^{3/2}} dx = \frac{2 \tan(x)}{3 \sqrt{\tan(x)^2 + 1}} + \frac{\tan(x)}{3 (\tan(x)^2 + 1)^{\frac{3}{2}}}$$

input `integrate(1/(sec(x)^2)^(3/2),x, algorithm="maxima")`output `2/3*tan(x)/sqrt(tan(x)^2 + 1) + 1/3*tan(x)/(tan(x)^2 + 1)^(3/2)`**3.44.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.55

$$\int \frac{1}{\sec^2(x)^{3/2}} dx = -\frac{1}{3} \operatorname{sgn}(\cos(x)) \sin(x)^3 + \operatorname{sgn}(\cos(x)) \sin(x)$$

input `integrate(1/(sec(x)^2)^(3/2),x, algorithm="giac")`output `-1/3*sgn(cos(x))*sin(x)^3 + sgn(cos(x))*sin(x)`

**3.44.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^2(x)^{3/2}} dx = \int \frac{1}{\left(\frac{1}{\cos(x)^2}\right)^{3/2}} dx$$

input `int(1/(1/cos(x)^2)^(3/2),x)`output `int(1/(1/cos(x)^2)^(3/2), x)`

### 3.45 $\int \frac{1}{\sec^2(x)^{5/2}} dx$

3.45.1	Optimal result . . . . .	343
3.45.2	Mathematica [A] (verified) . . . . .	343
3.45.3	Rubi [A] (verified) . . . . .	344
3.45.4	Maple [C] (warning: unable to verify) . . . . .	345
3.45.5	Fricas [A] (verification not implemented) . . . . .	346
3.45.6	Sympy [A] (verification not implemented) . . . . .	346
3.45.7	Maxima [A] (verification not implemented) . . . . .	346
3.45.8	Giac [A] (verification not implemented) . . . . .	347
3.45.9	Mupad [F(-1)] . . . . .	347

#### 3.45.1 Optimal result

Integrand size = 8, antiderivative size = 43

$$\int \frac{1}{\sec^2(x)^{5/2}} dx = \frac{\tan(x)}{5 \sec^2(x)^{5/2}} + \frac{4 \tan(x)}{15 \sec^2(x)^{3/2}} + \frac{8 \tan(x)}{15 \sqrt{\sec^2(x)}}$$

output  $1/5*\tan(x)/(\sec(x)^2)^{(5/2)}+4/15*\tan(x)/(\sec(x)^2)^{(3/2)}+8/15*\tan(x)/(\sec(x)^2)^{(1/2)}$

#### 3.45.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.65

$$\int \frac{1}{\sec^2(x)^{5/2}} dx = \frac{(15 - 10 \sin^2(x) + 3 \sin^4(x)) \tan(x)}{15 \sqrt{\sec^2(x)}}$$

input `Integrate[(Sec[x]^2)^(-5/2), x]`

output  $((15 - 10*\sin[x]^2 + 3*\sin[x]^4)*\tan[x])/(15*\text{Sqrt}[\sec[x]^2])$

### 3.45.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.26, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 4610, 209, 209, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^2(x)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\sec(x)^2)^{5/2}} dx \\
 & \quad \downarrow \text{4610} \\
 & \int \frac{1}{(\tan^2(x) + 1)^{7/2}} d \tan(x) \\
 & \quad \downarrow \text{209} \\
 & \frac{4}{5} \int \frac{1}{(\tan^2(x) + 1)^{5/2}} d \tan(x) + \frac{\tan(x)}{5(\tan^2(x) + 1)^{5/2}} \\
 & \quad \downarrow \text{209} \\
 & \frac{4}{5} \left( \frac{2}{3} \int \frac{1}{(\tan^2(x) + 1)^{3/2}} d \tan(x) + \frac{\tan(x)}{3(\tan^2(x) + 1)^{3/2}} \right) + \frac{\tan(x)}{5(\tan^2(x) + 1)^{5/2}} \\
 & \quad \downarrow \text{208} \\
 & \frac{\tan(x)}{5(\tan^2(x) + 1)^{5/2}} + \frac{4}{5} \left( \frac{2 \tan(x)}{3\sqrt{\tan^2(x) + 1}} + \frac{\tan(x)}{3(\tan^2(x) + 1)^{3/2}} \right)
 \end{aligned}$$

input `Int[(Sec[x]^2)^(-5/2),x]`

output `Tan[x]/(5*(1 + Tan[x]^2)^(5/2)) + (4*(Tan[x]/(3*(1 + Tan[x]^2)^(3/2)) + (2*Tan[x])/(3*Sqrt[1 + Tan[x]^2]))) / 5`

## 3.45.3.1 Defintions of rubi rules used

- rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`
- rule 209 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && ILtQ[p + 3/2, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

## 3.45.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.51

method	result
default	$\frac{\text{csgn}(\sec(x))(3 \cos(x)^4 + 4 \cos(x)^2 + 8) \sin(x)}{15}$
risch	$-\frac{ie^{6ix}}{160(e^{2ix}+1)\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}} - \frac{5ie^{2ix}}{16\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)} + \frac{5i}{16\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)} + \frac{5ie^{-2ix}}{96(e^{2ix}+1)\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}} - \frac{11i}{240\sqrt{\frac{e^2}{(e^{2ix}+1)^2}}}$

input `int(1/(sec(x)^2)^(5/2),x,method=_RETURNVERBOSE)`

output `1/15*csgn(sec(x))*(3*cos(x)^4+4*cos(x)^2+8)*sin(x)`

**3.45.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.42

$$\int \frac{1}{\sec^2(x)^{5/2}} dx = -\frac{1}{15} (3 \cos(x)^4 + 4 \cos(x)^2 + 8) \sin(x)$$

input `integrate(1/(sec(x)^2)^(5/2),x, algorithm="fricas")`output `-1/15*(3*cos(x)^4 + 4*cos(x)^2 + 8)*sin(x)`**3.45.6 Sympy [A] (verification not implemented)**

Time = 1.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.02

$$\int \frac{1}{\sec^2(x)^{5/2}} dx = \frac{8 \tan^5(x)}{15 (\sec^2(x))^{\frac{5}{2}}} + \frac{4 \tan^3(x)}{3 (\sec^2(x))^{\frac{5}{2}}} + \frac{\tan(x)}{(\sec^2(x))^{\frac{5}{2}}}$$

input `integrate(1/(sec(x)**2)**(5/2),x)`output `8*tan(x)**5/(15*(sec(x)**2)**(5/2)) + 4*tan(x)**3/(3*(sec(x)**2)**(5/2)) + tan(x)/(sec(x)**2)**(5/2)`**3.45.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sec^2(x)^{5/2}} dx = \frac{8 \tan(x)}{15 \sqrt{\tan(x)^2 + 1}} + \frac{4 \tan(x)}{15 (\tan(x)^2 + 1)^{\frac{3}{2}}} + \frac{\tan(x)}{5 (\tan(x)^2 + 1)^{\frac{5}{2}}}$$

input `integrate(1/(sec(x)^2)^(5/2),x, algorithm="maxima")`output `8/15*tan(x)/sqrt(tan(x)^2 + 1) + 4/15*tan(x)/(tan(x)^2 + 1)^(3/2) + 1/5*tan(x)/(tan(x)^2 + 1)^(5/2)`

**3.45.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.58

$$\int \frac{1}{\sec^2(x)^{5/2}} dx = \frac{1}{5} \operatorname{sgn}(\cos(x)) \sin(x)^5 - \frac{2}{3} \operatorname{sgn}(\cos(x)) \sin(x)^3 + \operatorname{sgn}(\cos(x)) \sin(x)$$

input `integrate(1/(sec(x)^2)^(5/2),x, algorithm="giac")`

output `1/5*sgn(cos(x))*sin(x)^5 - 2/3*sgn(cos(x))*sin(x)^3 + sgn(cos(x))*sin(x)`

**3.45.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^2(x)^{5/2}} dx = \int \frac{1}{\left(\frac{1}{\cos(x)^2}\right)^{5/2}} dx$$

input `int(1/(1/cos(x)^2)^(5/2),x)`

output `int(1/(1/cos(x)^2)^(5/2), x)`



### 3.46 $\int \frac{1}{\sec^2(x)^{7/2}} dx$

3.46.1	Optimal result . . . . .	348
3.46.2	Mathematica [A] (verified) . . . . .	348
3.46.3	Rubi [A] (verified) . . . . .	349
3.46.4	Maple [C] (warning: unable to verify) . . . . .	350
3.46.5	Fricas [A] (verification not implemented) . . . . .	351
3.46.6	Sympy [A] (verification not implemented) . . . . .	351
3.46.7	Maxima [A] (verification not implemented) . . . . .	351
3.46.8	Giac [A] (verification not implemented) . . . . .	352
3.46.9	Mupad [F(-1)] . . . . .	352

#### 3.46.1 Optimal result

Integrand size = 8, antiderivative size = 57

$$\int \frac{1}{\sec^2(x)^{7/2}} dx = \frac{\tan(x)}{7 \sec^2(x)^{7/2}} + \frac{6 \tan(x)}{35 \sec^2(x)^{5/2}} + \frac{8 \tan(x)}{35 \sec^2(x)^{3/2}} + \frac{16 \tan(x)}{35 \sqrt{\sec^2(x)}}$$

output `1/7*tan(x)/(sec(x)^2)^(7/2)+6/35*tan(x)/(sec(x)^2)^(5/2)+8/35*tan(x)/(sec(x)^2)^(3/2)+16/35*tan(x)/(sec(x)^2)^(1/2)`

#### 3.46.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.60

$$\int \frac{1}{\sec^2(x)^{7/2}} dx = \frac{(35 - 35 \sin^2(x) + 21 \sin^4(x) - 5 \sin^6(x)) \tan(x)}{35 \sqrt{\sec^2(x)}}$$

input `Integrate[(Sec[x]^2)^(-7/2), x]`

output `((35 - 35*Sin[x]^2 + 21*Sin[x]^4 - 5*Sin[x]^6)*Tan[x])/(35*Sqrt[Sec[x]^2])`

**3.46.3 Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.32, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 4610, 209, 209, 209, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^2(x)^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\sec(x)^2)^{7/2}} dx \\
 & \quad \downarrow \text{4610} \\
 & \int \frac{1}{(\tan^2(x) + 1)^{9/2}} d \tan(x) \\
 & \quad \downarrow \text{209} \\
 & \frac{6}{7} \int \frac{1}{(\tan^2(x) + 1)^{7/2}} d \tan(x) + \frac{\tan(x)}{7(\tan^2(x) + 1)^{7/2}} \\
 & \quad \downarrow \text{209} \\
 & \frac{6}{7} \left( \frac{4}{5} \int \frac{1}{(\tan^2(x) + 1)^{5/2}} d \tan(x) + \frac{\tan(x)}{5(\tan^2(x) + 1)^{5/2}} \right) + \frac{\tan(x)}{7(\tan^2(x) + 1)^{7/2}} \\
 & \quad \downarrow \text{209} \\
 & \frac{6}{7} \left( \frac{4}{5} \left( \frac{2}{3} \int \frac{1}{(\tan^2(x) + 1)^{3/2}} d \tan(x) + \frac{\tan(x)}{3(\tan^2(x) + 1)^{3/2}} \right) + \frac{\tan(x)}{5(\tan^2(x) + 1)^{5/2}} \right) + \\
 & \quad \frac{\tan(x)}{7(\tan^2(x) + 1)^{7/2}} \\
 & \quad \downarrow \text{208} \\
 & \frac{\tan(x)}{7(\tan^2(x) + 1)^{7/2}} + \frac{6}{7} \left( \frac{\tan(x)}{5(\tan^2(x) + 1)^{5/2}} + \frac{4}{5} \left( \frac{2 \tan(x)}{3\sqrt{\tan^2(x) + 1}} + \frac{\tan(x)}{3(\tan^2(x) + 1)^{3/2}} \right) \right)
 \end{aligned}$$

input `Int[(Sec[x]^2)^(-7/2),x]`

output  $\frac{\tan(x)}{7(1 + \tan(x)^2)^{7/2}} + \frac{6(\tan(x)/(5(1 + \tan(x)^2)^{5/2}) + 4(\tan(x)/(3(1 + \tan(x)^2)^{3/2}) + (2\tan(x))/(3\sqrt{1 + \tan(x)^2})))}{5}$

### 3.46.3.1 Defintions of rubi rules used

rule 208  $\text{Int}[(a + (b \cdot x)^2)^{-3/2}, x\_Symbol] \rightarrow \text{Simp}[x/(a\sqrt{a + b \cdot x^2}), x] /; \text{FreeQ}\{a, b, x\}$

rule 209  $\text{Int}[(a + (b \cdot x)^2)^p, x\_Symbol] \rightarrow \text{Simp}[(-x) \cdot (a + b \cdot x^2)^{p+1} / (2 \cdot a \cdot (p+1)), x] + \text{Simp}[(2p+3)/(2 \cdot a \cdot (p+1)) \cdot \text{Int}[(a + b \cdot x^2)^{p+1}], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{ILtQ}[p + 3/2, 0]$

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4610  $\text{Int}[(b \cdot \sec(e + f \cdot x) + (f \cdot x)^2)^p, x\_Symbol] \rightarrow \text{With}\{\text{ff} = \text{FreeFactors}[\tan[e + f \cdot x], x]\}, \text{Simp}[b \cdot (\text{ff}/f) \cdot \text{Subst}[\text{Int}[(b + b \cdot \text{ff}^2 \cdot x^2)^{p-1}], x, \tan[e + f \cdot x]/\text{ff}], x] /; \text{FreeQ}\{b, e, f, p, x\} \ \&\& \ \text{!IntegerQ}[p]$

### 3.46.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.43 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.49

method	result
default	$\frac{\text{csgn}(\sec(x)) (5 \cos(x)^6 + 6 \cos(x)^4 + 8 \cos(x)^2 + 16) \sin(x)}{35}$
risch	$-\frac{ie^{8ix}}{896(e^{2ix}+1)\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}} - \frac{35ie^{2ix}}{128\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)} + \frac{35i}{128\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)} + \frac{7ie^{-2ix}}{128(e^{2ix}+1)\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}} - \frac{1120}{\sqrt{\frac{e^{2ix}}{(e^{2ix}+1)^2}}}$

input `int(1/(sec(x)^2)^(7/2),x,method=_RETURNVERBOSE)`

output  $1/35 \cdot \text{csgn}(\sec(x)) \cdot (5 \cos(x)^6 + 6 \cos(x)^4 + 8 \cos(x)^2 + 16) \cdot \sin(x)$

---

3.46.  $\int \frac{1}{\sec^2(x)^{7/2}} dx$

**3.46.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.42

$$\int \frac{1}{\sec^2(x)^{7/2}} dx = -\frac{1}{35} (5 \cos(x)^6 + 6 \cos(x)^4 + 8 \cos(x)^2 + 16) \sin(x)$$

input `integrate(1/(sec(x)^2)^(7/2),x, algorithm="fricas")`output `-1/35*(5*cos(x)^6 + 6*cos(x)^4 + 8*cos(x)^2 + 16)*sin(x)`**3.46.6 Sympy [A] (verification not implemented)**

Time = 12.27 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.05

$$\int \frac{1}{\sec^2(x)^{7/2}} dx = \frac{16 \tan^7(x)}{35 (\sec^2(x))^{7/2}} + \frac{8 \tan^5(x)}{5 (\sec^2(x))^{7/2}} + \frac{2 \tan^3(x)}{(\sec^2(x))^{7/2}} + \frac{\tan(x)}{(\sec^2(x))^{7/2}}$$

input `integrate(1/(sec(x)**2)**(7/2),x)`output `16*tan(x)**7/(35*(sec(x)**2)**(7/2)) + 8*tan(x)**5/(5*(sec(x)**2)**(7/2)) + 2*tan(x)**3/(sec(x)**2)**(7/2) + tan(x)/(sec(x)**2)**(7/2)`**3.46.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sec^2(x)^{7/2}} dx = \frac{16 \tan(x)}{35 \sqrt{\tan(x)^2 + 1}} + \frac{8 \tan(x)}{35 (\tan(x)^2 + 1)^{3/2}} + \frac{6 \tan(x)}{35 (\tan(x)^2 + 1)^{5/2}} + \frac{\tan(x)}{7 (\tan(x)^2 + 1)^{7/2}}$$

input `integrate(1/(sec(x)^2)^(7/2),x, algorithm="maxima")`output `16/35*tan(x)/sqrt(tan(x)^2 + 1) + 8/35*tan(x)/(tan(x)^2 + 1)^(3/2) + 6/35*tan(x)/(tan(x)^2 + 1)^(5/2) + 1/7*tan(x)/(tan(x)^2 + 1)^(7/2)`

**3.46.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.60

$$\int \frac{1}{\sec^2(x)^{7/2}} dx = -\frac{1}{7} \operatorname{sgn}(\cos(x)) \sin(x)^7 + \frac{3}{5} \operatorname{sgn}(\cos(x)) \sin(x)^5 - \operatorname{sgn}(\cos(x)) \sin(x)^3 + \operatorname{sgn}(\cos(x)) \sin(x)$$

input `integrate(1/(sec(x)^2)^(7/2),x, algorithm="giac")`

output `-1/7*sgn(cos(x))*sin(x)^7 + 3/5*sgn(cos(x))*sin(x)^5 - sgn(cos(x))*sin(x)^3 + sgn(cos(x))*sin(x)`

**3.46.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sec^2(x)^{7/2}} dx = \int \frac{1}{\left(\frac{1}{\cos(x)^2}\right)^{7/2}} dx$$

input `int(1/(1/cos(x)^2)^(7/2),x)`

output `int(1/(1/cos(x)^2)^(7/2), x)`

### 3.47 $\int (a \sec^2(x))^{7/2} dx$

3.47.1 Optimal result . . . . .	353
3.47.2 Mathematica [A] (verified) . . . . .	353
3.47.3 Rubi [A] (verified) . . . . .	354
3.47.4 Maple [A] (verified) . . . . .	356
3.47.5 Fricas [A] (verification not implemented) . . . . .	356
3.47.6 Sympy [F(-1)] . . . . .	357
3.47.7 Maxima [B] (verification not implemented) . . . . .	357
3.47.8 Giac [A] (verification not implemented) . . . . .	358
3.47.9 Mupad [F(-1)] . . . . .	358

#### 3.47.1 Optimal result

Integrand size = 10, antiderivative size = 84

$$\int (a \sec^2(x))^{7/2} dx = \frac{5}{16} a^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{5}{16} a^3 \sqrt{a \sec^2(x)} \tan(x) \\ + \frac{5}{24} a^2 (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{6} a (a \sec^2(x))^{5/2} \tan(x)$$

output  $5/16*a^{(7/2)}*\operatorname{arctanh}(a^{(1/2)}*\tan(x)/(a*\sec(x)^2)^{(1/2)})+5/24*a^2*(a*\sec(x)^2)^{(3/2)}*\tan(x)+1/6*a*(a*\sec(x)^2)^{(5/2)}*\tan(x)+5/16*a^3*(a*\sec(x)^2)^{(1/2)}*\tan(x)$

#### 3.47.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.52

$$\int (a \sec^2(x))^{7/2} dx = \frac{1}{48} a^3 \cos(x) \sqrt{a \sec^2(x)} (15 \operatorname{arctanh}(\sin(x)) \\ + \sec(x) (15 + 10 \sec^2(x) + 8 \sec^4(x)) \tan(x))$$

input  $\operatorname{Integrate}[(a*\operatorname{Sec}[x]^2)^{(7/2)}, x]$

output  $(a^3*\operatorname{Cos}[x]*\operatorname{Sqrt}[a*\operatorname{Sec}[x]^2]*(15*\operatorname{ArcTanh}[\operatorname{Sin}[x]] + \operatorname{Sec}[x]*(15 + 10*\operatorname{Sec}[x]^2 + 8*\operatorname{Sec}[x]^4)*\operatorname{Tan}[x]))/48$

**3.47.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.18, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$ , Rules used = {3042, 4610, 211, 211, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^2(x))^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^2)^{7/2} dx \\
 & \quad \downarrow \text{4610} \\
 & a \int (a \tan^2(x) + a)^{5/2} d \tan(x) \\
 & \quad \downarrow \text{211} \\
 & a \left( \frac{5}{6} a \int (a \tan^2(x) + a)^{3/2} d \tan(x) + \frac{1}{6} \tan(x) (a \tan^2(x) + a)^{5/2} \right) \\
 & \quad \downarrow \text{211} \\
 & a \left( \frac{5}{6} a \left( \frac{3}{4} a \int \sqrt{a \tan^2(x) + a} d \tan(x) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right) + \frac{1}{6} \tan(x) (a \tan^2(x) + a)^{5/2} \right) \\
 & \quad \downarrow \text{211} \\
 & a \left( \frac{5}{6} a \left( \frac{3}{4} a \left( \frac{1}{2} a \int \frac{1}{\sqrt{a \tan^2(x) + a}} d \tan(x) + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right) + \frac{1}{6} \tan(x) (a \tan^2(x) + a)^{5/2} \right) \\
 & \quad \downarrow \text{224} \\
 & a \left( \frac{5}{6} a \left( \frac{3}{4} a \left( \frac{1}{2} a \int \frac{1}{1 - \frac{a \tan^2(x)}{a \tan^2(x) + a}} d \frac{\tan(x)}{\sqrt{a \tan^2(x) + a}} + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right) + \frac{1}{6} \tan(x) (a \tan^2(x) + a)^{5/2} \right) \\
 & \quad \downarrow \text{219} \\
 & a \left( \frac{5}{6} a \left( \frac{3}{4} a \left( \frac{1}{2} \sqrt{a} \operatorname{arctanh} \left( \frac{\sqrt{a} \tan(x)}{\sqrt{a \tan^2(x) + a}} \right) + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right) + \frac{1}{6} \tan(x) (a \tan^2(x) + a)^{5/2} \right)
 \end{aligned}$$

input `Int[(a*Sec[x]^2)^(7/2),x]`

output `a*((Tan[x]*(a + a*Tan[x]^2)^(5/2))/6 + (5*a*((Tan[x]*(a + a*Tan[x]^2)^(3/2)))/4 + (3*a*((Sqrt[a]*ArcTanh[(Sqrt[a]*Tan[x])/Sqrt[a + a*Tan[x]^2]])/2 + (Tan[x]*Sqrt[a + a*Tan[x]^2])/2)/4))/6)`

### 3.47.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`



**3.47.4 Maple [A] (verified)**

Time = 2.04 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.73

method	result
default	$\frac{\sqrt{a \sec(x)^2} a^3 (15 \tan(x) + 10 \sec(x)^2 \tan(x) + 8 \tan(x) \sec(x)^4 + 15 \cos(x) \ln(-\cot(x) + \csc(x) + 1) - 15 \cos(x) \ln(-\cot(x) + \csc(x) - 1))}{48}$
risch	$-\frac{ia^3 \sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}} (15e^{10ix} + 85e^{8ix} + 198e^{6ix} - 198e^{4ix} - 85e^{2ix} - 15)}{24(e^{2ix}+1)^5} + \frac{5a^3 \sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix}+i) \cos(x)}{8} - \frac{5a^3 \sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}}{8}$

input `int((a*sec(x)^2)^(7/2),x,method=_RETURNVERBOSE)`output `1/48*(a*sec(x)^2)^(1/2)*a^3*(15*tan(x)+10*sec(x)^2*tan(x)+8*tan(x)*sec(x)^4+15*cos(x)*ln(-cot(x)+csc(x)+1)-15*cos(x)*ln(-cot(x)+csc(x)-1))`**3.47.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.77

$$\int (a \sec^2(x))^{7/2} dx = \frac{\left(15 a^3 \cos(x)^6 \log\left(-\frac{\sin(x)-1}{\sin(x)+1}\right) - 2(15 a^3 \cos(x)^4 + 10 a^3 \cos(x)^2 + 8 a^3) \sin(x)\right) \sqrt{\frac{a}{\cos(x)^2}}}{96 \cos(x)^5}$$

input `integrate((a*sec(x)^2)^(7/2),x, algorithm="fracas")`output `-1/96*(15*a^3*cos(x)^6*log(-(sin(x) - 1)/(sin(x) + 1)) - 2*(15*a^3*cos(x)^4 + 10*a^3*cos(x)^2 + 8*a^3)*sin(x))*sqrt(a/cos(x)^2)/cos(x)^5`

### 3.47.6 Sympy [F(-1)]

Timed out.

$$\int (a \sec^2(x))^{7/2} dx = \text{Timed out}$$

input `integrate((a*sec(x)**2)**(7/2),x)`

output `Timed out`

### 3.47.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2175 vs. 2(64) = 128.

Time = 2.45 (sec) , antiderivative size = 2175, normalized size of antiderivative = 25.89

$$\int (a \sec^2(x))^{7/2} dx = \text{Too large to display}$$

input `integrate((a*sec(x)^2)^(7/2),x, algorithm="maxima")`

output

```

1/96*(2040*a^3*cos(3*x)*sin(2*x) + 360*a^3*cos(x)*sin(2*x) - 360*a^3*cos(2
*x)*sin(x) - 60*a^3*sin(x) + 4*(15*a^3*sin(11*x) + 85*a^3*sin(9*x) + 198*a
^3*sin(7*x) - 198*a^3*sin(5*x) - 85*a^3*sin(3*x) - 15*a^3*sin(x))*cos(12*x
) - 60*(6*a^3*sin(10*x) + 15*a^3*sin(8*x) + 20*a^3*sin(6*x) + 15*a^3*sin(4
*x) + 6*a^3*sin(2*x))*cos(11*x) + 24*(85*a^3*sin(9*x) + 198*a^3*sin(7*x) -
198*a^3*sin(5*x) - 85*a^3*sin(3*x) - 15*a^3*sin(x))*cos(10*x) - 340*(15*a
^3*sin(8*x) + 20*a^3*sin(6*x) + 15*a^3*sin(4*x) + 6*a^3*sin(2*x))*cos(9*x)
+ 60*(198*a^3*sin(7*x) - 198*a^3*sin(5*x) - 85*a^3*sin(3*x) - 15*a^3*sin(
x))*cos(8*x) - 792*(20*a^3*sin(6*x) + 15*a^3*sin(4*x) + 6*a^3*sin(2*x))*co
s(7*x) - 80*(198*a^3*sin(5*x) + 85*a^3*sin(3*x) + 15*a^3*sin(x))*cos(6*x)
+ 2376*(5*a^3*sin(4*x) + 2*a^3*sin(2*x))*cos(5*x) - 300*(17*a^3*sin(3*x) +
3*a^3*sin(x))*cos(4*x) + 15*(a^3*cos(12*x)^2 + 36*a^3*cos(10*x)^2 + 225*a
^3*cos(8*x)^2 + 400*a^3*cos(6*x)^2 + 225*a^3*cos(4*x)^2 + 36*a^3*cos(2*x)^
2 + a^3*sin(12*x)^2 + 36*a^3*sin(10*x)^2 + 225*a^3*sin(8*x)^2 + 400*a^3*si
n(6*x)^2 + 225*a^3*sin(4*x)^2 + 180*a^3*sin(4*x)*sin(2*x) + 36*a^3*sin(2*x
)^2 + 12*a^3*cos(2*x) + a^3 + 2*(6*a^3*cos(10*x) + 15*a^3*cos(8*x) + 20*a^
3*cos(6*x) + 15*a^3*cos(4*x) + 6*a^3*cos(2*x) + a^3)*cos(12*x) + 12*(15*a^
3*cos(8*x) + 20*a^3*cos(6*x) + 15*a^3*cos(4*x) + 6*a^3*cos(2*x) + a^3)*cos
(10*x) + 30*(20*a^3*cos(6*x) + 15*a^3*cos(4*x) + 6*a^3*cos(2*x) + a^3)*cos
(8*x) + 40*(15*a^3*cos(4*x) + 6*a^3*cos(2*x) + a^3)*cos(6*x) + 30*(6*a^...

```

**3.47.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.94

$$\int (a \sec^2(x))^{7/2} dx = \frac{1}{96} \left( 15 a^3 \log(\sin(x) + 1) \operatorname{sgn}(\cos(x)) - 15 a^3 \log(-\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \frac{2}{1} \right)$$

input `integrate((a*sec(x)^2)^(7/2),x, algorithm="giac")`

output `1/96*(15*a^3*log(sin(x) + 1)*sgn(cos(x)) - 15*a^3*log(-sin(x) + 1)*sgn(cos(x)) - 2*(15*a^3*sgn(cos(x))*sin(x)^5 - 40*a^3*sgn(cos(x))*sin(x)^3 + 33*a^3*sgn(cos(x))*sin(x))/(sin(x)^2 - 1)^3)*sqrt(a)`

**3.47.9 Mupad [F(-1)]**

Timed out.

$$\int (a \sec^2(x))^{7/2} dx = \int \left( \frac{a}{\cos(x)^2} \right)^{7/2} dx$$

input `int((a/cos(x)^2)^(7/2),x)`

output `int((a/cos(x)^2)^(7/2), x)`

## 3.48 $\int (a \sec^2(x))^{5/2} dx$

3.48.1	Optimal result	359
3.48.2	Mathematica [A] (verified)	359
3.48.3	Rubi [A] (verified)	360
3.48.4	Maple [A] (verified)	361
3.48.5	Fricas [A] (verification not implemented)	362
3.48.6	Sympy [F]	362
3.48.7	Maxima [B] (verification not implemented)	362
3.48.8	Giac [A] (verification not implemented)	363
3.48.9	Mupad [F(-1)]	364

### 3.48.1 Optimal result

Integrand size = 10, antiderivative size = 65

$$\int (a \sec^2(x))^{5/2} dx = \frac{3}{8} a^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{3}{8} a^2 \sqrt{a \sec^2(x)} \tan(x) + \frac{1}{4} a (a \sec^2(x))^{3/2} \tan(x)$$

output  $3/8*a^{(5/2)}*\operatorname{arctanh}(a^{(1/2)}*\tan(x)/(a*\sec(x)^2)^{(1/2)})+1/4*a*(a*\sec(x)^2)^{(3/2)}*\tan(x)+3/8*a^2*(a*\sec(x)^2)^{(1/2)}*\tan(x)$

### 3.48.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.57

$$\int (a \sec^2(x))^{5/2} dx = \frac{1}{8} \cos(x) (a \sec^2(x))^{5/2} (3 \operatorname{arctanh}(\sin(x)) \cos^4(x) + (2 + 3 \cos^2(x)) \sin(x))$$

input  $\operatorname{Integrate}[(a*\operatorname{Sec}[x]^2)^{(5/2)},x]$

output  $(\operatorname{Cos}[x]*(a*\operatorname{Sec}[x]^2)^{(5/2)}*(3*\operatorname{ArcTanh}[\operatorname{Sin}[x]]*\operatorname{Cos}[x]^4 + (2 + 3*\operatorname{Cos}[x]^2)*\operatorname{Sin}[x]))/8$

**3.48.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.15, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4610, 211, 211, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^2(x))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^2)^{5/2} dx \\
 & \quad \downarrow \text{4610} \\
 & a \int (a \tan^2(x) + a)^{3/2} d \tan(x) \\
 & \quad \downarrow \text{211} \\
 & a \left( \frac{3}{4} a \int \sqrt{a \tan^2(x) + a} d \tan(x) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right) \\
 & \quad \downarrow \text{211} \\
 & a \left( \frac{3}{4} a \left( \frac{1}{2} a \int \frac{1}{\sqrt{a \tan^2(x) + a}} d \tan(x) + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right) \\
 & \quad \downarrow \text{224} \\
 & a \left( \frac{3}{4} a \left( \frac{1}{2} a \int \frac{1}{1 - \frac{a \tan^2(x)}{a \tan^2(x) + a}} d \frac{\tan(x)}{\sqrt{a \tan^2(x) + a}} + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right) \\
 & \quad \downarrow \text{219} \\
 & a \left( \frac{3}{4} a \left( \frac{1}{2} \sqrt{a} \operatorname{arctanh} \left( \frac{\sqrt{a} \tan(x)}{\sqrt{a \tan^2(x) + a}} \right) + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) + \frac{1}{4} \tan(x) (a \tan^2(x) + a)^{3/2} \right)
 \end{aligned}$$

input `Int[(a*Sec[x]^2)^(5/2),x]`

output  $a*((\text{Tan}[x]*(a + a*\text{Tan}[x]^2)^{(3/2)})/4 + (3*a*((\text{Sqrt}[a]*\text{ArcTanh}[(\text{Sqrt}[a]*\text{Tan}[x])/(\text{Sqrt}[a + a*\text{Tan}[x]^2])])/2 + (\text{Tan}[x]*\text{Sqrt}[a + a*\text{Tan}[x]^2])/2))/4$

### 3.48.3.1 Defintions of rubi rules used

rule 211  $\text{Int}[(a + b*x^2)^p, x\_Symbol] \rightarrow \text{Simp}[x*(a + b*x^2)^{p/(2*p + 1)}, x] + \text{Simp}[2*a*(p/(2*p + 1)) \text{Int}[(a + b*x^2)^{p - 1}, x], x] /;$  FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4\*p] || IntegerQ[6\*p])

rule 219  $\text{Int}[(a + b*x^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /;$  FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

rule 224  $\text{Int}[1/\text{Sqrt}[a + b*x^2], x\_Symbol] \rightarrow \text{Subst}[\text{Int}[1/(1 - b*x^2), x], x, x/\text{Sqrt}[a + b*x^2]] /;$  FreeQ[{a, b}, x] && !GtQ[a, 0]

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$  FunctionOfTrigOfLinearQ[u, x]

rule 4610  $\text{Int}[(b*sec[e + f*x] + (f*x)^2)^p, x\_Symbol] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Simp}[b*(ff/f) \text{Subst}[\text{Int}[(b + b*ff^2*x^2)^{p - 1}, x], x, \text{Tan}[e + f*x]/ff], x] /;$  FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

### 3.48.4 Maple [A] (verified)

Time = 1.99 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\sqrt{a \sec(x)^2} a^2 (3 \cos(x) \ln(-\cot(x) + \csc(x) + 1) - 3 \cos(x) \ln(-\cot(x) + \csc(x) - 1) + 3 \tan(x) + 2 \sec(x)^2 \tan(x))}{8}$	53
risch	$-\frac{ia^2 \sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}} (3e^{6ix} + 11e^{4ix} - 11e^{2ix} - 3)}{4(e^{2ix}+1)^3} + \frac{3a^2 \sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix}+i) \cos(x)}{4} - \frac{3a^2 \sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix}-i) \cos(x)}{4}$	12

input `int((a*sec(x)^2)^(5/2),x,method=_RETURNVERBOSE)`

3.48.  $\int (a \sec^2(x))^{5/2} dx$

output  $1/8*(a*\sec(x)^2)^{(1/2)}*a^2*(3*\cos(x)*\ln(-\cot(x)+\csc(x)+1)-3*\cos(x)*\ln(-\cot(x)+\csc(x)-1)+3*\tan(x)+2*\sec(x)^2*\tan(x))$

### 3.48.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.86

$$\int (a \sec^2(x))^{5/2} dx = -\frac{\left(3 a^2 \cos(x)^4 \log\left(-\frac{\sin(x)-1}{\sin(x)+1}\right) - 2(3 a^2 \cos(x)^2 + 2 a^2) \sin(x)\right) \sqrt{\frac{a}{\cos(x)^2}}}{16 \cos(x)^3}$$

input `integrate((a*sec(x)^2)^(5/2),x, algorithm="fricas")`

output  $-1/16*(3*a^2*\cos(x)^4*\log(-(\sin(x) - 1)/(\sin(x) + 1)) - 2*(3*a^2*\cos(x)^2 + 2*a^2)*\sin(x))*\sqrt{a/\cos(x)^2}/\cos(x)^3$

### 3.48.6 Sympy [F]

$$\int (a \sec^2(x))^{5/2} dx = \int (a \sec^2(x))^{\frac{5}{2}} dx$$

input `integrate((a*sec(x)**2)**(5/2),x)`

output `Integral((a*sec(x)**2)**(5/2), x)`

### 3.48.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1111 vs.  $2(49) = 98$ .

Time = 0.53 (sec) , antiderivative size = 1111, normalized size of antiderivative = 17.09

$$\int (a \sec^2(x))^{5/2} dx = \text{Too large to display}$$

input `integrate((a*sec(x)^2)^(5/2),x, algorithm="maxima")`

output `1/16*(176*a^2*cos(3*x)*sin(2*x) + 48*a^2*cos(x)*sin(2*x) - 48*a^2*cos(2*x)*sin(x) - 12*a^2*sin(x) + 4*(3*a^2*sin(7*x) + 11*a^2*sin(5*x) - 11*a^2*sin(3*x) - 3*a^2*sin(x))*cos(8*x) - 24*(2*a^2*sin(6*x) + 3*a^2*sin(4*x) + 2*a^2*sin(2*x))*cos(7*x) + 16*(11*a^2*sin(5*x) - 11*a^2*sin(3*x) - 3*a^2*sin(x))*cos(6*x) - 88*(3*a^2*sin(4*x) + 2*a^2*sin(2*x))*cos(5*x) - 24*(11*a^2*sin(3*x) + 3*a^2*sin(x))*cos(4*x) + 3*(a^2*cos(8*x)^2 + 16*a^2*cos(6*x)^2 + 36*a^2*cos(4*x)^2 + 16*a^2*cos(2*x)^2 + a^2*sin(8*x)^2 + 16*a^2*sin(6*x)^2 + 36*a^2*sin(4*x)^2 + 48*a^2*sin(4*x)*sin(2*x) + 16*a^2*sin(2*x)^2 + 8*a^2*cos(2*x) + a^2 + 2*(4*a^2*cos(6*x) + 6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(8*x) + 8*(6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(6*x) + 12*(4*a^2*cos(2*x) + a^2)*cos(4*x) + 4*(2*a^2*sin(6*x) + 3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(8*x) + 16*(3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(6*x))*log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - 3*(a^2*cos(8*x)^2 + 16*a^2*cos(6*x)^2 + 36*a^2*cos(4*x)^2 + 16*a^2*cos(2*x)^2 + a^2*sin(8*x)^2 + 16*a^2*sin(6*x)^2 + 36*a^2*sin(4*x)^2 + 48*a^2*sin(4*x)*sin(2*x) + 16*a^2*sin(2*x)^2 + 8*a^2*cos(2*x) + a^2 + 2*(4*a^2*cos(6*x) + 6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(8*x) + 8*(6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(6*x) + 12*(4*a^2*cos(2*x) + a^2)*cos(4*x) + 4*(2*a^2*sin(6*x) + 3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(8*x) + 16*(3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(6*x))*log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1) - 4*(3*a^2*cos(7*x) + 11*a^2*cos(...`

### 3.48.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.03

$$\int (a \sec^2(x))^{5/2} dx = \frac{1}{16} \left( 3a^2 \log(\sin(x) + 1) \operatorname{sgn}(\cos(x)) - 3a^2 \log(-\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \frac{2(3a^2}{16} \right)$$

input `integrate((a*sec(x)^2)^(5/2),x, algorithm="giac")`

output `1/16*(3*a^2*log(sin(x) + 1)*sgn(cos(x)) - 3*a^2*log(-sin(x) + 1)*sgn(cos(x))) - 2*(3*a^2*sgn(cos(x))*sin(x)^3 - 5*a^2*sgn(cos(x))*sin(x))/(sin(x)^2 - 1)^2)*sqrt(a)`



**3.48.9 Mupad [F(-1)]**

Timed out.

$$\int (a \sec^2(x))^{5/2} dx = \int \left( \frac{a}{\cos(x)^2} \right)^{5/2} dx$$

input `int((a/cos(x)^2)^(5/2),x)`output `int((a/cos(x)^2)^(5/2), x)`

### 3.49 $\int (a \sec^2(x))^{3/2} dx$

3.49.1	Optimal result . . . . .	365
3.49.2	Mathematica [A] (verified) . . . . .	365
3.49.3	Rubi [A] (verified) . . . . .	366
3.49.4	Maple [A] (verified) . . . . .	367
3.49.5	Fricas [A] (verification not implemented) . . . . .	368
3.49.6	Sympy [F] . . . . .	368
3.49.7	Maxima [B] (verification not implemented) . . . . .	368
3.49.8	Giac [A] (verification not implemented) . . . . .	369
3.49.9	Mupad [F(-1)] . . . . .	369

#### 3.49.1 Optimal result

Integrand size = 10, antiderivative size = 46

$$\int (a \sec^2(x))^{3/2} dx = \frac{1}{2} a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{1}{2} a \sqrt{a \sec^2(x)} \tan(x)$$

output `1/2*a^(3/2)*arctanh(a^(1/2)*tan(x)/(a*sec(x)^2)^(1/2))+1/2*a*(a*sec(x)^2)^(1/2)*tan(x)`

#### 3.49.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int (a \sec^2(x))^{3/2} dx = \frac{1}{2} a \sqrt{a \sec^2(x)} (\operatorname{arctanh}(\sin(x)) \cos(x) + \tan(x))$$

input `Integrate[(a*Sec[x]^2)^(3/2),x]`

output `(a*Sqrt[a*Sec[x]^2]*(ArcTanh[Sin[x]]*Cos[x] + Tan[x]))/2`

**3.49.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4610, 211, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^2)^{3/2} dx \\
 & \quad \downarrow \text{4610} \\
 & a \int \sqrt{a \tan^2(x) + a} d \tan(x) \\
 & \quad \downarrow \text{211} \\
 & a \left( \frac{1}{2} a \int \frac{1}{\sqrt{a \tan^2(x) + a}} d \tan(x) + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) \\
 & \quad \downarrow \text{224} \\
 & a \left( \frac{1}{2} a \int \frac{1}{1 - \frac{a \tan^2(x)}{a \tan^2(x) + a}} d \frac{\tan(x)}{\sqrt{a \tan^2(x) + a}} + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right) \\
 & \quad \downarrow \text{219} \\
 & a \left( \frac{1}{2} \sqrt{a} \operatorname{arctanh} \left( \frac{\sqrt{a} \tan(x)}{\sqrt{a \tan^2(x) + a}} \right) + \frac{1}{2} \tan(x) \sqrt{a \tan^2(x) + a} \right)
 \end{aligned}$$

input `Int[(a*Sec[x]^2)^(3/2),x]`

output `a*((Sqrt[a]*ArcTanh[(Sqrt[a]*Tan[x])/Sqrt[a + a*Tan[x]^2]])/2 + (Tan[x]*Sqrt[a + a*Tan[x]^2])/2)`

### 3.49.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

### 3.49.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{\sqrt{a \sec^2(x)} a (\cos(x) \ln(-\cot(x) + \csc(x) + 1) - \cos(x) \ln(-\cot(x) + \csc(x) - 1) + \tan(x))}{2}$	40
risch	$-\frac{ia \sqrt{\frac{a e^{2ix}}{(e^{2ix} + 1)^2}} (e^{2ix} - 1)}{e^{2ix} + 1} + a \sqrt{\frac{a e^{2ix}}{(e^{2ix} + 1)^2}} \ln(e^{ix} + i) \cos(x) - a \sqrt{\frac{a e^{2ix}}{(e^{2ix} + 1)^2}} \ln(e^{ix} - i) \cos(x)$	103

input `int((a*sec(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `1/2*(a*sec(x)^2)^(1/2)*a*(cos(x)*ln(-cot(x)+csc(x)+1)-cos(x)*ln(-cot(x)+csc(x)-1)+tan(x))`

**3.49.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

$$\int (a \sec^2(x))^{3/2} dx = -\frac{\left(a \cos(x)^2 \log\left(-\frac{\sin(x)-1}{\sin(x)+1}\right) - 2 a \sin(x)\right) \sqrt{\frac{a}{\cos(x)^2}}}{4 \cos(x)}$$

input `integrate((a*sec(x)^2)^(3/2),x, algorithm="fracas")`

output `-1/4*(a*cos(x)^2*log(-(sin(x) - 1)/(sin(x) + 1)) - 2*a*sin(x))*sqrt(a/cos(x)^2)/cos(x)`

**3.49.6 Sympy [F]**

$$\int (a \sec^2(x))^{3/2} dx = \int (a \sec^2(x))^{\frac{3}{2}} dx$$

input `integrate((a*sec(x)**2)**(3/2),x)`

output `Integral((a*sec(x)**2)**(3/2), x)`

**3.49.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 324 vs.  $2(34) = 68$ .

Time = 0.37 (sec) , antiderivative size = 324, normalized size of antiderivative = 7.04

$$\int (a \sec^2(x))^{3/2} dx = \frac{(8 a \cos(3 x) \sin(2 x) - 8 a \cos(x) \sin(2 x) + 8 a \cos(2 x) \sin(x) - 4 (a \sin(3 x) - a \sin(x)) \cos(4 x) - (a$$

input `integrate((a*sec(x)^2)^(3/2),x, algorithm="maxima")`

```
output -1/4*(8*a*cos(3*x)*sin(2*x) - 8*a*cos(x)*sin(2*x) + 8*a*cos(2*x)*sin(x) -
4*(a*sin(3*x) - a*sin(x))*cos(4*x) - (a*cos(4*x)^2 + 4*a*cos(2*x)^2 + a*si
n(4*x)^2 + 4*a*sin(4*x)*sin(2*x) + 4*a*sin(2*x)^2 + 2*(2*a*cos(2*x) + a)*c
os(4*x) + 4*a*cos(2*x) + a)*log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) + (a*c
os(4*x)^2 + 4*a*cos(2*x)^2 + a*sin(4*x)^2 + 4*a*sin(4*x)*sin(2*x) + 4*a*si
n(2*x)^2 + 2*(2*a*cos(2*x) + a)*cos(4*x) + 4*a*cos(2*x) + a)*log(cos(x)^2
+ sin(x)^2 - 2*sin(x) + 1) + 4*(a*cos(3*x) - a*cos(x))*sin(4*x) - 4*(2*a*c
os(2*x) + a)*sin(3*x) + 4*a*sin(x))*sqrt(a)/(2*(2*cos(2*x) + 1)*cos(4*x) +
cos(4*x)^2 + 4*cos(2*x)^2 + sin(4*x)^2 + 4*sin(4*x)*sin(2*x) + 4*sin(2*x)
^2 + 4*cos(2*x) + 1)
```

### 3.49.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.91

$$\int (a \sec^2(x))^{3/2} dx = \frac{1}{4} \left( \log(\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \log(-\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \frac{2 \operatorname{sgn}(\cos(x)) \sin(x)}{\sin(x)^2 - 1} \right)$$

```
input integrate((a*sec(x)^2)^(3/2),x, algorithm="giac")
```

```
output 1/4*(log(sin(x) + 1)*sgn(cos(x)) - log(-sin(x) + 1)*sgn(cos(x)) - 2*sgn(co
s(x))*sin(x)/(sin(x)^2 - 1))*a^(3/2)
```

### 3.49.9 Mupad [F(-1)]

Timed out.

$$\int (a \sec^2(x))^{3/2} dx = \int \left( \frac{a}{\cos(x)^2} \right)^{3/2} dx$$

```
input int((a/cos(x)^2)^(3/2),x)
```

```
output int((a/cos(x)^2)^(3/2), x)
```

### 3.50 $\int \sqrt{a \sec^2(x)} dx$

3.50.1	Optimal result . . . . .	370
3.50.2	Mathematica [A] (verified) . . . . .	370
3.50.3	Rubi [A] (verified) . . . . .	371
3.50.4	Maple [A] (verified) . . . . .	372
3.50.5	Fricas [A] (verification not implemented) . . . . .	372
3.50.6	Sympy [F] . . . . .	373
3.50.7	Maxima [A] (verification not implemented) . . . . .	373
3.50.8	Giac [A] (verification not implemented) . . . . .	373
3.50.9	Mupad [F(-1)] . . . . .	374

#### 3.50.1 Optimal result

Integrand size = 10, antiderivative size = 25

$$\int \sqrt{a \sec^2(x)} dx = \sqrt{a} \operatorname{arctanh} \left( \frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}} \right)$$

output `arctanh(a^(1/2)*tan(x)/(a*sec(x)^2)^(1/2))*a^(1/2)`

#### 3.50.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.64

$$\int \sqrt{a \sec^2(x)} dx = \operatorname{arctanh}(\sin(x)) \cos(x) \sqrt{a \sec^2(x)}$$

input `Integrate[Sqrt[a*Sec[x]^2],x]`

output `ArcTanh[Sin[x]]*Cos[x]*Sqrt[a*Sec[x]^2]`

### 3.50.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4610, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \sec^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \sec(x)^2} dx \\
 & \quad \downarrow \text{4610} \\
 & a \int \frac{1}{\sqrt{a \tan^2(x) + a}} d \tan(x) \\
 & \quad \downarrow \text{224} \\
 & a \int \frac{1}{1 - \frac{a \tan^2(x)}{a \tan^2(x) + a}} d \frac{\tan(x)}{\sqrt{a \tan^2(x) + a}} \\
 & \quad \downarrow \text{219} \\
 & \sqrt{a} \operatorname{arctanh} \left( \frac{\sqrt{a} \tan(x)}{\sqrt{a \tan^2(x) + a}} \right)
 \end{aligned}$$

input `Int[Sqrt[a*Sec[x]^2],x]`

output `Sqrt[a]*ArcTanh[(Sqrt[a]*Tan[x])/Sqrt[a + a*Tan[x]^2]]`

#### 3.50.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`



rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

### 3.50.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

method	result	size
default	$-2 \cos(x) \sqrt{a \sec^2(x)} \operatorname{arctanh}(\cot(x) - \csc(x))$	21
risch	$-2 \sqrt{\frac{a e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix} - i) \cos(x) + 2 \sqrt{\frac{a e^{2ix}}{(e^{2ix}+1)^2}} \ln(e^{ix} + i) \cos(x)$	64

input `int((a*sec(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*cos(x)*(a*sec(x)^2)^(1/2)*arctanh(cot(x)-csc(x))`

### 3.50.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.20

$$\int \sqrt{a \sec^2(x)} dx = \left[ -\frac{1}{2} \sqrt{\frac{a}{\cos(x)^2}} \cos(x) \log\left(-\frac{\sin(x) - 1}{\sin(x) + 1}\right), -\sqrt{-a} \operatorname{arctan}\left(\frac{\sqrt{-a} \sqrt{\frac{a}{\cos(x)^2}} \cos(x) \sin(x)}{a}\right) \right]$$

input `integrate((a*sec(x)^2)^(1/2),x, algorithm="fracas")`

output `[-1/2*sqrt(a/cos(x)^2)*cos(x)*log(-(sin(x) - 1)/(sin(x) + 1)), -sqrt(-a)*arctan(sqrt(-a)*sqrt(a/cos(x)^2)*cos(x)*sin(x)/a)]`

**3.50.6 Sympy [F]**

$$\int \sqrt{a \sec^2(x)} dx = \int \sqrt{a \sec^2(x)} dx$$

input `integrate((a*sec(x)**2)**(1/2),x)`

output `Integral(sqrt(a*sec(x)**2), x)`

**3.50.7 Maxima [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.52

$$\begin{aligned} & \int \sqrt{a \sec^2(x)} dx \\ &= \frac{1}{2} \sqrt{a} (\log(\cos(x)^2 + \sin(x)^2 + 2 \sin(x) + 1) - \log(\cos(x)^2 + \sin(x)^2 - 2 \sin(x) + 1)) \end{aligned}$$

input `integrate((a*sec(x)^2)^(1/2),x, algorithm="maxima")`

output `1/2*sqrt(a)*(log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1))`

**3.50.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.24

$$\begin{aligned} & \int \sqrt{a \sec^2(x)} dx \\ &= \frac{1}{4} \sqrt{a} \left( \log \left( \left| \frac{1}{\sin(x)} + \sin(x) + 2 \right| \right) - \log \left( \left| \frac{1}{\sin(x)} + \sin(x) - 2 \right| \right) \right) \operatorname{sgn}(\cos(x)) \end{aligned}$$

input `integrate((a*sec(x)^2)^(1/2),x, algorithm="giac")`

output `1/4*sqrt(a)*(log(abs(1/sin(x) + sin(x) + 2)) - log(abs(1/sin(x) + sin(x) - 2)))*sgn(cos(x))`

**3.50.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \sec^2(x)} dx = \int \sqrt{\frac{a}{\cos(x)^2}} dx$$

input `int((a/cos(x)^2)^(1/2),x)`output `int((a/cos(x)^2)^(1/2), x)`

### 3.51 $\int \frac{1}{\sqrt{a \sec^2(x)}} dx$

3.51.1	Optimal result	375
3.51.2	Mathematica [A] (verified)	375
3.51.3	Rubi [A] (verified)	376
3.51.4	Maple [A] (verified)	377
3.51.5	Fricas [A] (verification not implemented)	377
3.51.6	Sympy [A] (verification not implemented)	377
3.51.7	Maxima [A] (verification not implemented)	378
3.51.8	Giac [A] (verification not implemented)	378
3.51.9	Mupad [B] (verification not implemented)	378

#### 3.51.1 Optimal result

Integrand size = 10, antiderivative size = 13

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = \frac{\tan(x)}{\sqrt{a \sec^2(x)}}$$

output `tan(x)/(a*sec(x)^2)^(1/2)`

#### 3.51.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = \frac{\tan(x)}{\sqrt{a \sec^2(x)}}$$

input `Integrate[1/Sqrt[a*Sec[x]^2],x]`

output `Tan[x]/Sqrt[a*Sec[x]^2]`

### 3.51.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {3042, 4610, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{\sqrt{a \sec^2(x)}} dx \\
 \downarrow 3042 \\
 \int \frac{1}{\sqrt{a \sec(x)^2}} dx \\
 \downarrow 4610 \\
 a \int \frac{1}{(a \tan^2(x) + a)^{3/2}} d \tan(x) \\
 \downarrow 208 \\
 \frac{\tan(x)}{\sqrt{a \tan^2(x) + a}}
 \end{array}$$

input `Int[1/Sqrt[a*Sec[x]^2],x]`

output `Tan[x]/Sqrt[a + a*Tan[x]^2]`

#### 3.51.3.1 Defintions of rubi rules used

rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

**3.51.4 Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\frac{\tan(x)}{\sqrt{a \sec(x)^2}}$	12
risch	$-\frac{ie^{2ix}}{2\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)} + \frac{i}{2\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}(e^{2ix}+1)}$	67

input `int(1/(a*sec(x)^2)^(1/2),x,method=_RETURNVERBOSE)`output `tan(x)/(a*sec(x)^2)^(1/2)`**3.51.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.23

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = \frac{\sqrt{\frac{a}{\cos(x)^2}} \cos(x) \sin(x)}{a}$$

input `integrate(1/(a*sec(x)^2)^(1/2),x, algorithm="fricas")`output `sqrt(a/cos(x)^2)*cos(x)*sin(x)/a`**3.51.6 Sympy [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = \frac{\tan(x)}{\sqrt{a \sec^2(x)}}$$

input `integrate(1/(a*sec(x)**2)**(1/2),x)`output `tan(x)/sqrt(a*sec(x)**2)`

**3.51.7 Maxima [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.46

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = \frac{\sin(x)}{\sqrt{a}}$$

input `integrate(1/(a*sec(x)^2)^(1/2),x, algorithm="maxima")`output `sin(x)/sqrt(a)`**3.51.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = \frac{\sin(x)}{\sqrt{a} \operatorname{sgn}(\cos(x))}$$

input `integrate(1/(a*sec(x)^2)^(1/2),x, algorithm="giac")`output `sin(x)/(sqrt(a)*sgn(cos(x)))`**3.51.9 Mupad [B] (verification not implemented)**

Time = 12.72 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = \frac{\sqrt{2} \sin(2x)}{2 \sqrt{a} \sqrt{2 \cos(x)^2}}$$

input `int(1/(a/cos(x)^2)^(1/2),x)`output `(2^(1/2)*sin(2*x))/(2*a^(1/2)*(2*cos(x)^2)^(1/2))`

### 3.52 $\int \frac{1}{(a \sec^2(x))^{3/2}} dx$

3.52.1	Optimal result . . . . .	379
3.52.2	Mathematica [A] (verified) . . . . .	379
3.52.3	Rubi [A] (verified) . . . . .	380
3.52.4	Maple [A] (verified) . . . . .	381
3.52.5	Fricas [A] (verification not implemented) . . . . .	382
3.52.6	Sympy [A] (verification not implemented) . . . . .	382
3.52.7	Maxima [A] (verification not implemented) . . . . .	382
3.52.8	Giac [A] (verification not implemented) . . . . .	383
3.52.9	Mupad [F(-1)] . . . . .	383

#### 3.52.1 Optimal result

Integrand size = 10, antiderivative size = 36

$$\int \frac{1}{(a \sec^2(x))^{3/2}} dx = \frac{\tan(x)}{3(a \sec^2(x))^{3/2}} + \frac{2 \tan(x)}{3a\sqrt{a \sec^2(x)}}$$

output `1/3*tan(x)/(a*sec(x)^2)^(3/2)+2/3*tan(x)/a/(a*sec(x)^2)^(1/2)`

#### 3.52.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.69

$$\int \frac{1}{(a \sec^2(x))^{3/2}} dx = \frac{(5 + \cos(2x)) \tan(x)}{6a\sqrt{a \sec^2(x)}}$$

input `Integrate[(a*Sec[x]^2)^(-3/2), x]`

output `((5 + Cos[2*x])*Tan[x])/(6*a*Sqrt[a*Sec[x]^2])`



### 3.52.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.25, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4610, 209, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sec^2(x))^{3/2}} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{1}{(a \sec(x)^2)^{3/2}} dx \\
 & \quad \downarrow 4610 \\
 & a \int \frac{1}{(a \tan^2(x) + a)^{5/2}} d \tan(x) \\
 & \quad \downarrow 209 \\
 & a \left( \frac{2 \int \frac{1}{(a \tan^2(x) + a)^{3/2}} d \tan(x)}{3a} + \frac{\tan(x)}{3a (a \tan^2(x) + a)^{3/2}} \right) \\
 & \quad \downarrow 208 \\
 & a \left( \frac{2 \tan(x)}{3a^2 \sqrt{a \tan^2(x) + a}} + \frac{\tan(x)}{3a (a \tan^2(x) + a)^{3/2}} \right)
 \end{aligned}$$

input `Int [(a*Sec [x]^2)^(-3/2), x]`

output `a*(Tan [x]/(3*a*(a + a*Tan [x]^2)^(3/2)) + (2*Tan [x])/(3*a^2*Sqrt [a + a*Tan [x]^2]))`

## 3.52.3.1 Defintions of rubi rules used

- rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`
- rule 209 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && ILtQ[p + 3/2, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

## 3.52.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{\sin(x) \cos(x) + 2 \tan(x)}{3 \sqrt{a \sec(x)^2 a}}$	24
risch	$-\frac{ie^{4ix}}{24a(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} - \frac{3ie^{2ix}}{8a(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} + \frac{3i}{8a(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} + \frac{ie^{-2ix}}{24a(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}}$	149

input `int(1/(a*sec(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `1/3/(a*sec(x)^2)^(1/2)/a*(sin(x)*cos(x)+2*tan(x))`

**3.52.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.67

$$\int \frac{1}{(a \sec^2(x))^{3/2}} dx = \frac{(\cos(x)^3 + 2 \cos(x)) \sqrt{\frac{a}{\cos(x)^2}} \sin(x)}{3 a^2}$$

input `integrate(1/(a*sec(x)^2)^(3/2),x, algorithm="fricas")`output `1/3*(cos(x)^3 + 2*cos(x))*sqrt(a/cos(x)^2)*sin(x)/a^2`**3.52.6 Sympy [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a \sec^2(x))^{3/2}} dx = \frac{2 \tan^3(x)}{3 (a \sec^2(x))^{3/2}} + \frac{\tan(x)}{(a \sec^2(x))^{3/2}}$$

input `integrate(1/(a*sec(x)**2)**(3/2),x)`output `2*tan(x)**3/(3*(a*sec(x)**2)**(3/2)) + tan(x)/(a*sec(x)**2)**(3/2)`**3.52.7 Maxima [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.39

$$\int \frac{1}{(a \sec^2(x))^{3/2}} dx = \frac{\sin(3x) + 9 \sin(x)}{12 a^{3/2}}$$

input `integrate(1/(a*sec(x)^2)^(3/2),x, algorithm="maxima")`output `1/12*(sin(3*x) + 9*sin(x))/a^(3/2)`

**3.52.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.53

$$\int \frac{1}{(a \sec^2(x))^{3/2}} dx = -\frac{\sin(x)^3 - 3 \sin(x)}{3 a^{3/2} \operatorname{sgn}(\cos(x))}$$

input `integrate(1/(a*sec(x)^2)^(3/2),x, algorithm="giac")`

output `-1/3*(sin(x)^3 - 3*sin(x))/(a^(3/2)*sgn(cos(x)))`

**3.52.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \sec^2(x))^{3/2}} dx = \int \frac{1}{\left(\frac{a}{\cos(x)^2}\right)^{3/2}} dx$$

input `int(1/(a/cos(x)^2)^(3/2),x)`

output `int(1/(a/cos(x)^2)^(3/2), x)`

### 3.53 $\int \frac{1}{(a \sec^2(x))^{5/2}} dx$

3.53.1	Optimal result . . . . .	384
3.53.2	Mathematica [A] (verified) . . . . .	384
3.53.3	Rubi [A] (verified) . . . . .	385
3.53.4	Maple [A] (verified) . . . . .	386
3.53.5	Fricas [A] (verification not implemented) . . . . .	387
3.53.6	Sympy [A] (verification not implemented) . . . . .	387
3.53.7	Maxima [A] (verification not implemented) . . . . .	387
3.53.8	Giac [A] (verification not implemented) . . . . .	388
3.53.9	Mupad [F(-1)] . . . . .	388

#### 3.53.1 Optimal result

Integrand size = 10, antiderivative size = 55

$$\int \frac{1}{(a \sec^2(x))^{5/2}} dx = \frac{\tan(x)}{5 (a \sec^2(x))^{5/2}} + \frac{4 \tan(x)}{15a (a \sec^2(x))^{3/2}} + \frac{8 \tan(x)}{15a^2 \sqrt{a \sec^2(x)}}$$

output `1/5*tan(x)/(a*sec(x)^2)^(5/2)+4/15*tan(x)/a/(a*sec(x)^2)^(3/2)+8/15*tan(x)/a^2/(a*sec(x)^2)^(1/2)`

#### 3.53.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.60

$$\int \frac{1}{(a \sec^2(x))^{5/2}} dx = \frac{(15 - 10 \sin^2(x) + 3 \sin^4(x)) \tan(x)}{15a^2 \sqrt{a \sec^2(x)}}$$

input `Integrate[(a*Sec[x]^2)^(-5/2),x]`

output `((15 - 10*Sin[x]^2 + 3*Sin[x]^4)*Tan[x])/(15*a^2*Sqrt[a*Sec[x]^2])`

### 3.53.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.35, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4610, 209, 209, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sec^2(x))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sec(x)^2)^{5/2}} dx \\
 & \quad \downarrow \text{4610} \\
 & a \int \frac{1}{(a \tan^2(x) + a)^{7/2}} d \tan(x) \\
 & \quad \downarrow \text{209} \\
 & a \left( \frac{4 \int \frac{1}{(a \tan^2(x) + a)^{5/2}} d \tan(x)}{5a} + \frac{\tan(x)}{5a (a \tan^2(x) + a)^{5/2}} \right) \\
 & \quad \downarrow \text{209} \\
 & a \left( \frac{4 \left( \frac{2 \int \frac{1}{(a \tan^2(x) + a)^{3/2}} d \tan(x)}{3a} + \frac{\tan(x)}{3a (a \tan^2(x) + a)^{3/2}} \right)}{5a} + \frac{\tan(x)}{5a (a \tan^2(x) + a)^{5/2}} \right) \\
 & \quad \downarrow \text{208} \\
 & a \left( \frac{4 \left( \frac{2 \tan(x)}{3a^2 \sqrt{a \tan^2(x) + a}} + \frac{\tan(x)}{3a (a \tan^2(x) + a)^{3/2}} \right)}{5a} + \frac{\tan(x)}{5a (a \tan^2(x) + a)^{5/2}} \right)
 \end{aligned}$$

input `Int[(a*Sec[x]^2)^(-5/2),x]`

output `a*(Tan[x]/(5*a*(a + a*Tan[x]^2)^(5/2)) + (4*(Tan[x]/(3*a*(a + a*Tan[x]^2)^(3/2)) + (2*Tan[x])/(3*a^2*Sqrt[a + a*Tan[x]^2])))/(5*a))`

---

3.53.  $\int \frac{1}{(a \sec^2(x))^{5/2}} dx$

## 3.53.3.1 Defintions of rubi rules used

- rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`
- rule 209 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && ILtQ[p + 3/2, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

## 3.53.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.55

method	result
default	$\frac{(3 \cos(x)^4 + 4 \cos(x)^2 + 8) \tan(x)}{15 \sqrt{a \sec(x)^2} a^2}$
risch	$-\frac{ie^{6ix}}{160a^2(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} - \frac{5ie^{2ix}}{16a^2(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} + \frac{5i}{16a^2(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} + \frac{5ie^{-2ix}}{96a^2(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} - \frac{5i}{96a^2(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}}$

input `int(1/(a*sec(x)^2)^(5/2),x,method=_RETURNVERBOSE)`

output `1/15*(3*cos(x)^4+4*cos(x)^2+8)/(a*sec(x)^2)^(1/2)/a^2*tan(x)`

**3.53.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.58

$$\int \frac{1}{(a \sec^2(x))^{5/2}} dx = \frac{(3 \cos(x)^5 + 4 \cos(x)^3 + 8 \cos(x)) \sqrt{\frac{a}{\cos(x)^2}} \sin(x)}{15 a^3}$$

input `integrate(1/(a*sec(x)^2)^(5/2),x, algorithm="fricas")`output `1/15*(3*cos(x)^5 + 4*cos(x)^3 + 8*cos(x))*sqrt(a/cos(x)^2)*sin(x)/a^3`**3.53.6 Sympy [A] (verification not implemented)**

Time = 1.34 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89

$$\int \frac{1}{(a \sec^2(x))^{5/2}} dx = \frac{8 \tan^5(x)}{15 (a \sec^2(x))^{5/2}} + \frac{4 \tan^3(x)}{3 (a \sec^2(x))^{5/2}} + \frac{\tan(x)}{(a \sec^2(x))^{5/2}}$$

input `integrate(1/(a*sec(x)**2)**(5/2),x)`output `8*tan(x)**5/(15*(a*sec(x)**2)**(5/2)) + 4*tan(x)**3/(3*(a*sec(x)**2)**(5/2)) + tan(x)/(a*sec(x)**2)**(5/2)`**3.53.7 Maxima [A] (verification not implemented)**

Time = 0.52 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.40

$$\int \frac{1}{(a \sec^2(x))^{5/2}} dx = \frac{3 \sin(5x) + 25 \sin(3x) + 150 \sin(x)}{240 a^{5/2}}$$

input `integrate(1/(a*sec(x)^2)^(5/2),x, algorithm="maxima")`output `1/240*(3*sin(5*x) + 25*sin(3*x) + 150*sin(x))/a^(5/2)`



**3.53.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.49

$$\int \frac{1}{(a \sec^2(x))^{5/2}} dx = \frac{3 \sin(x)^5 - 10 \sin(x)^3 + 15 \sin(x)}{15 a^{5/2} \operatorname{sgn}(\cos(x))}$$

input `integrate(1/(a*sec(x)^2)^(5/2),x, algorithm="giac")`

output `1/15*(3*sin(x)^5 - 10*sin(x)^3 + 15*sin(x))/(a^(5/2)*sgn(cos(x)))`

**3.53.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \sec^2(x))^{5/2}} dx = \int \frac{1}{\left(\frac{a}{\cos(x)^2}\right)^{5/2}} dx$$

input `int(1/(a/cos(x)^2)^(5/2),x)`

output `int(1/(a/cos(x)^2)^(5/2), x)`

### 3.54 $\int \frac{1}{(a \sec^2(x))^{7/2}} dx$

3.54.1	Optimal result . . . . .	389
3.54.2	Mathematica [A] (verified) . . . . .	389
3.54.3	Rubi [A] (verified) . . . . .	390
3.54.4	Maple [A] (verified) . . . . .	392
3.54.5	Fricas [A] (verification not implemented) . . . . .	392
3.54.6	Sympy [A] (verification not implemented) . . . . .	392
3.54.7	Maxima [A] (verification not implemented) . . . . .	393
3.54.8	Giac [A] (verification not implemented) . . . . .	393
3.54.9	Mupad [F(-1)] . . . . .	393

#### 3.54.1 Optimal result

Integrand size = 10, antiderivative size = 74

$$\int \frac{1}{(a \sec^2(x))^{7/2}} dx = \frac{\tan(x)}{7(a \sec^2(x))^{7/2}} + \frac{6 \tan(x)}{35a(a \sec^2(x))^{5/2}} + \frac{8 \tan(x)}{35a^2(a \sec^2(x))^{3/2}} + \frac{16 \tan(x)}{35a^3 \sqrt{a \sec^2(x)}}$$

output `1/7*tan(x)/(a*sec(x)^2)^(7/2)+6/35*tan(x)/a/(a*sec(x)^2)^(5/2)+8/35*tan(x)/a^2/(a*sec(x)^2)^(3/2)+16/35*tan(x)/a^3/(a*sec(x)^2)^(1/2)`

#### 3.54.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.53

$$\int \frac{1}{(a \sec^2(x))^{7/2}} dx = \frac{(35 - 35 \sin^2(x) + 21 \sin^4(x) - 5 \sin^6(x)) \tan(x)}{35a^3 \sqrt{a \sec^2(x)}}$$

input `Integrate[(a*Sec[x]^2)^(-7/2),x]`

output `((35 - 35*Sin[x]^2 + 21*Sin[x]^4 - 5*Sin[x]^6)*Tan[x])/(35*a^3*Sqrt[a*Sec[x]^2])`

**3.54.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.39, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4610, 209, 209, 209, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sec^2(x))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sec(x)^2)^{7/2}} dx \\
 & \quad \downarrow \text{4610} \\
 & a \int \frac{1}{(a \tan^2(x) + a)^{9/2}} d \tan(x) \\
 & \quad \downarrow \text{209} \\
 & a \left( \frac{6 \int \frac{1}{(a \tan^2(x) + a)^{7/2}} d \tan(x)}{7a} + \frac{\tan(x)}{7a (a \tan^2(x) + a)^{7/2}} \right) \\
 & \quad \downarrow \text{209} \\
 & a \left( \frac{6 \left( \frac{4 \int \frac{1}{(a \tan^2(x) + a)^{5/2}} d \tan(x)}{5a} + \frac{\tan(x)}{5a (a \tan^2(x) + a)^{5/2}} \right)}{7a} + \frac{\tan(x)}{7a (a \tan^2(x) + a)^{7/2}} \right) \\
 & \quad \downarrow \text{209} \\
 & a \left( \frac{6 \left( \frac{4 \left( \frac{2 \int \frac{1}{(a \tan^2(x) + a)^{3/2}} d \tan(x)}{3a} + \frac{\tan(x)}{3a (a \tan^2(x) + a)^{3/2}} \right)}{5a} + \frac{\tan(x)}{5a (a \tan^2(x) + a)^{5/2}} \right)}{7a} + \frac{\tan(x)}{7a (a \tan^2(x) + a)^{7/2}} \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 208 \\
 a \left( \frac{6 \left( \frac{4 \left( \frac{2 \tan(x)}{3a^2 \sqrt{a \tan^2(x)+a}} + \frac{\tan(x)}{3a (a \tan^2(x)+a)^{3/2}} \right)}{5a} + \frac{\tan(x)}{5a (a \tan^2(x)+a)^{5/2}} \right)}{7a} + \frac{\tan(x)}{7a (a \tan^2(x)+a)^{7/2}} \right)
 \end{array}$$

```
input Int[(a*Sec[x]^2)^(-7/2),x]
```

```
output a*(Tan[x]/(7*a*(a + a*Tan[x]^2)^(7/2)) + (6*(Tan[x]/(5*a*(a + a*Tan[x]^2)^(5/2)) + (4*(Tan[x]/(3*a*(a + a*Tan[x]^2)^(3/2)) + (2*Tan[x]/(3*a^2*Sqrt[a + a*Tan[x]^2])))/(5*a)))/(7*a))
```

**3.54.3.1 Defintions of rubi rules used**

```
rule 208 Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]
```

```
rule 209 Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && ILtQ[p + 3/2, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4610 Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

### 3.54.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.49

method	result
default	$\frac{(5 \cos(x)^6 + 6 \cos(x)^4 + 8 \cos(x)^2 + 16) \tan(x)}{35 \sqrt{a \sec(x)^2} a^3}$
risch	$-\frac{ie^{8ix}}{896a^3(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} - \frac{35ie^{2ix}}{128a^3(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} + \frac{35i}{128a^3(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}} + \frac{7ie^{-2ix}}{128a^3(e^{2ix}+1)\sqrt{\frac{ae^{2ix}}{(e^{2ix}+1)^2}}}$

input `int(1/(a*sec(x)^2)^(7/2),x,method=_RETURNVERBOSE)`

output `1/35*(5*cos(x)^6+6*cos(x)^4+8*cos(x)^2+16)/(a*sec(x)^2)^(1/2)/a^3*tan(x)`

### 3.54.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.51

$$\int \frac{1}{(a \sec^2(x))^{7/2}} dx = \frac{(5 \cos(x)^7 + 6 \cos(x)^5 + 8 \cos(x)^3 + 16 \cos(x)) \sqrt{\frac{a}{\cos(x)^2}} \sin(x)}{35 a^4}$$

input `integrate(1/(a*sec(x)^2)^(7/2),x, algorithm="fricas")`

output `1/35*(5*cos(x)^7 + 6*cos(x)^5 + 8*cos(x)^3 + 16*cos(x))*sqrt(a/cos(x)^2)*sin(x)/a^4`

### 3.54.6 Sympy [A] (verification not implemented)

Time = 12.65 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.89

$$\int \frac{1}{(a \sec^2(x))^{7/2}} dx = \frac{16 \tan^7(x)}{35 (a \sec^2(x))^{7/2}} + \frac{8 \tan^5(x)}{5 (a \sec^2(x))^{7/2}} + \frac{2 \tan^3(x)}{(a \sec^2(x))^{7/2}} + \frac{\tan(x)}{(a \sec^2(x))^{7/2}}$$

input `integrate(1/(a*sec(x)**2)**(7/2),x)`

output `16*tan(x)**7/(35*(a*sec(x)**2)**(7/2)) + 8*tan(x)**5/(5*(a*sec(x)**2)**(7/2)) + 2*tan(x)**3/(a*sec(x)**2)**(7/2) + tan(x)/(a*sec(x)**2)**(7/2)`

---

3.54.  $\int \frac{1}{(a \sec^2(x))^{7/2}} dx$

**3.54.7 Maxima [A] (verification not implemented)**

Time = 0.48 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.38

$$\int \frac{1}{(a \sec^2(x))^{7/2}} dx = \frac{5 \sin(7x) + 49 \sin(5x) + 245 \sin(3x) + 1225 \sin(x)}{2240 a^{7/2}}$$

input `integrate(1/(a*sec(x)^2)^(7/2),x, algorithm="maxima")`output `1/2240*(5*sin(7*x) + 49*sin(5*x) + 245*sin(3*x) + 1225*sin(x))/a^(7/2)`**3.54.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.45

$$\int \frac{1}{(a \sec^2(x))^{7/2}} dx = -\frac{5 \sin(x)^7 - 21 \sin(x)^5 + 35 \sin(x)^3 - 35 \sin(x)}{35 a^{7/2} \operatorname{sgn}(\cos(x))}$$

input `integrate(1/(a*sec(x)^2)^(7/2),x, algorithm="giac")`output `-1/35*(5*sin(x)^7 - 21*sin(x)^5 + 35*sin(x)^3 - 35*sin(x))/(a^(7/2)*sgn(cos(x)))`**3.54.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \sec^2(x))^{7/2}} dx = \int \frac{1}{\left(\frac{a}{\cos(x)^2}\right)^{7/2}} dx$$

input `int(1/(a/cos(x)^2)^(7/2),x)`output `int(1/(a/cos(x)^2)^(7/2), x)`

### 3.55 $\int (a \sec^3(x))^{5/2} dx$

3.55.1	Optimal result . . . . .	394
3.55.2	Mathematica [A] (verified) . . . . .	394
3.55.3	Rubi [A] (verified) . . . . .	395
3.55.4	Maple [C] (verified) . . . . .	398
3.55.5	Fricas [C] (verification not implemented) . . . . .	398
3.55.6	Sympy [F] . . . . .	399
3.55.7	Maxima [F] . . . . .	399
3.55.8	Giac [F] . . . . .	399
3.55.9	Mupad [F(-1)] . . . . .	400

#### 3.55.1 Optimal result

Integrand size = 10, antiderivative size = 117

$$\begin{aligned} \int (a \sec^3(x))^{5/2} dx &= -\frac{154}{195} a^2 \cos^3(x) E\left(\frac{x}{2} \middle| 2\right) \sqrt{a \sec^3(x)} \\ &+ \frac{154}{195} a^2 \cos(x) \sqrt{a \sec^3(x)} \sin(x) + \frac{154}{585} a^2 \sqrt{a \sec^3(x)} \tan(x) \\ &+ \frac{22}{117} a^2 \sec^2(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{2}{13} a^2 \sec^4(x) \sqrt{a \sec^3(x)} \tan(x) \end{aligned}$$

output `-154/195*a^2*cos(x)^(3/2)*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticE(sin(1/2*x),2^(1/2))*(a*sec(x)^3)^(1/2)+154/195*a^2*cos(x)*sin(x)*(a*sec(x)^3)^(1/2)+154/585*a^2*(a*sec(x)^3)^(1/2)*tan(x)+22/117*a^2*sec(x)^2*(a*sec(x)^3)^(1/2)*tan(x)+2/13*a^2*sec(x)^4*(a*sec(x)^3)^(1/2)*tan(x)`

#### 3.55.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.50

$$\begin{aligned} \int (a \sec^3(x))^{5/2} dx &= \\ &-\frac{2}{585} a \sec(x) (a \sec^3(x))^{3/2} \left( 231 \cos^{\frac{11}{2}}(x) E\left(\frac{x}{2} \middle| 2\right) - 55 \cos(x) \sin(x) - 77 \cos^3(x) \sin(x) - 231 \cos^5(x) \sin(x) \right) \end{aligned}$$

input `Integrate[(a*Sec[x]^3)^(5/2),x]`

output  $(-2*a*\text{Sec}[x]*(a*\text{Sec}[x]^3)^{(3/2)}*(231*\text{Cos}[x]^{(11/2)}*\text{EllipticE}[x/2, 2] - 55*\text{Cos}[x]*\text{Sin}[x] - 77*\text{Cos}[x]^3*\text{Sin}[x] - 231*\text{Cos}[x]^5*\text{Sin}[x] - 45*\text{Tan}[x]))/585$

### 3.55.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.88, number of steps used = 14, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.400$ , Rules used = {3042, 4611, 3042, 4255, 3042, 4255, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^3(x))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^3)^{5/2} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{a^2 \sqrt{a \sec^3(x)} \int \sec^{15/2}(x) dx}{\sec^{3/2}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a^2 \sqrt{a \sec^3(x)} \int \csc(x + \frac{\pi}{2})^{15/2} dx}{\sec^{3/2}(x)} \\
 & \quad \downarrow \text{4255} \\
 & \frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \int \sec^{11/2}(x) dx + \frac{2}{13} \sin(x) \sec^{13/2}(x) \right)}{\sec^{3/2}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \int \csc(x + \frac{\pi}{2})^{11/2} dx + \frac{2}{13} \sin(x) \sec^{13/2}(x) \right)}{\sec^{3/2}(x)} \\
 & \quad \downarrow \text{4255} \\
 & \frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \int \sec^{7/2}(x) dx + \frac{2}{9} \sin(x) \sec^{9/2}(x) \right) + \frac{2}{13} \sin(x) \sec^{13/2}(x) \right)}{\sec^{3/2}(x)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$



$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \int \csc \left( x + \frac{\pi}{2} \right)^{7/2} dx + \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) \right) + \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 4255

$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \left( \frac{3}{5} \int \sec^{\frac{3}{2}}(x) dx + \frac{2}{5} \sin(x) \sec^{\frac{5}{2}}(x) \right) + \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) \right) + \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 3042

$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \left( \frac{3}{5} \int \csc \left( x + \frac{\pi}{2} \right)^{3/2} dx + \frac{2}{5} \sin(x) \sec^{\frac{5}{2}}(x) \right) + \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) \right) + \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 4255

$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \left( \frac{3}{5} \left( 2 \sin(x) \sqrt{\sec(x)} - \int \frac{1}{\sqrt{\sec(x)}} dx \right) + \frac{2}{5} \sin(x) \sec^{\frac{5}{2}}(x) \right) + \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) \right) + \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 3042

$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \left( \frac{3}{5} \left( 2 \sin(x) \sqrt{\sec(x)} - \int \frac{1}{\sqrt{\csc \left( x + \frac{\pi}{2} \right)}} dx \right) + \frac{2}{5} \sin(x) \sec^{\frac{5}{2}}(x) \right) + \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) \right) + \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 4258

$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \left( \frac{3}{5} \left( 2 \sin(x) \sqrt{\sec(x)} - \sqrt{\cos(x)} \sqrt{\sec(x)} \int \sqrt{\cos(x)} dx \right) + \frac{2}{5} \sin(x) \sec^{\frac{5}{2}}(x) \right) + \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) \right) + \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 3042

$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{11}{13} \left( \frac{7}{9} \left( \frac{3}{5} \left( 2 \sin(x) \sqrt{\sec(x)} - \sqrt{\cos(x)} \sqrt{\sec(x)} \int \sqrt{\sin \left( x + \frac{\pi}{2} \right)} dx \right) + \frac{2}{5} \sin(x) \sec^{\frac{5}{2}}(x) \right) + \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) \right) + \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 3119

$$\frac{a^2 \sqrt{a \sec^3(x)} \left( \frac{2}{13} \sin(x) \sec^{\frac{13}{2}}(x) + \frac{11}{13} \left( \frac{2}{9} \sin(x) \sec^{\frac{9}{2}}(x) + \frac{7}{9} \left( \frac{2}{5} \sin(x) \sec^{\frac{5}{2}}(x) + \frac{3}{5} \left( 2 \sin(x) \sqrt{\sec(x)} - 2 \sqrt{\cos(x)} \int \sqrt{\cos(x)} dx \right) \right) \right) \right)}{\sec^{\frac{3}{2}}(x)}$$

input `Int[(a*Sec[x]^3)^(5/2),x]`

output `(a^2*Sqrt[a*Sec[x]^3]*((2*Sec[x]^(13/2)*Sin[x])/13 + (11*((2*Sec[x]^(9/2)*Sin[x])/9 + (7*((2*Sec[x]^(5/2)*Sin[x])/5 + (3*(-2*Sqrt[Cos[x]]*EllipticE[x/2, 2]*Sqrt[Sec[x]] + 2*Sqrt[Sec[x]]*Sin[x]))/5))/9))/13))/Sec[x]^(3/2)`

### 3.55.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] :=> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] :=> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] :=> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=> Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

### 3.55.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 26.89 (sec) , antiderivative size = 297, normalized size of antiderivative = 2.54

method	result
default	$-\frac{2i\sqrt{a\sec(x)^3}a^2\left(231\operatorname{EllipticE}(i(\csc(x)-\cot(x)),i)\sqrt{\frac{\cos(x)}{\cos(x)+1}}\sqrt{\frac{1}{\cos(x)+1}}\cos(x)^3-231\operatorname{EllipticF}(i(\csc(x)-\cot(x)),i)\sqrt{\frac{\cos(x)}{\cos(x)+1}}\right)}{\dots}$

input `int((a*sec(x)^3)^(5/2),x,method=_RETURNVERBOSE)`

output

```
-2/585*I*(a*sec(x)^3)^(1/2)*a^2/(cos(x)+1)*(231*EllipticE(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)^3-231*EllipticF(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)^3+462*EllipticE(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)^2-462*EllipticF(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)^2+231*EllipticE(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)-231*EllipticF(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)+231*I*cos(x)*sin(x)+77*I*sin(x)+77*I*tan(x)+55*I*sec(x)*tan(x)+55*I*sec(x)^2*tan(x)+45*I*tan(x)*sec(x)^3+45*I*tan(x)*sec(x)^4)
```

### 3.55.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.87

$$\int (a \sec^3(x))^{5/2} dx = \frac{231i \sqrt{2} a^{5/2} \cos(x)^5 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x)))}{\dots}$$

input `integrate((a*sec(x)^3)^(5/2),x, algorithm="fricas")`

output

```
1/585*(231*I*sqrt(2)*a^(5/2)*cos(x)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) + I*sin(x))) - 231*I*sqrt(2)*a^(5/2)*cos(x)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) - I*sin(x))) + 2*(231*a^2*cos(x)^6 + 77*a^2*cos(x)^4 + 55*a^2*cos(x)^2 + 45*a^2)*sqrt(a/cos(x)^3)*sin(x))/cos(x)^5
```

**3.55.6 Sympy [F]**

$$\int (a \sec^3(x))^{5/2} dx = \int (a \sec^3(x))^{\frac{5}{2}} dx$$

input `integrate((a*sec(x)**3)**(5/2),x)`

output `Integral((a*sec(x)**3)**(5/2), x)`

**3.55.7 Maxima [F]**

$$\int (a \sec^3(x))^{5/2} dx = \int (a \sec(x)^3)^{\frac{5}{2}} dx$$

input `integrate((a*sec(x)^3)^(5/2),x, algorithm="maxima")`

output `integrate((a*sec(x)^3)^(5/2), x)`

**3.55.8 Giac [F]**

$$\int (a \sec^3(x))^{5/2} dx = \int (a \sec(x)^3)^{\frac{5}{2}} dx$$

input `integrate((a*sec(x)^3)^(5/2),x, algorithm="giac")`

output `integrate((a*sec(x)^3)^(5/2), x)`

**3.55.9 Mupad [F(-1)]**

Timed out.

$$\int (a \sec^3(x))^{5/2} dx = \int \left( \frac{a}{\cos(x)^3} \right)^{5/2} dx$$

input `int((a/cos(x)^3)^(5/2),x)`output `int((a/cos(x)^3)^(5/2), x)`

### 3.56 $\int (a \sec^3(x))^{3/2} dx$

3.56.1	Optimal result . . . . .	401
3.56.2	Mathematica [A] (verified) . . . . .	401
3.56.3	Rubi [A] (verified) . . . . .	402
3.56.4	Maple [C] (verified) . . . . .	404
3.56.5	Fricas [C] (verification not implemented) . . . . .	404
3.56.6	Sympy [F] . . . . .	405
3.56.7	Maxima [F] . . . . .	405
3.56.8	Giac [F] . . . . .	405
3.56.9	Mupad [F(-1)] . . . . .	406

#### 3.56.1 Optimal result

Integrand size = 10, antiderivative size = 65

$$\int (a \sec^3(x))^{3/2} dx = \frac{10}{21} a \cos^{\frac{3}{2}}(x) \operatorname{EllipticF}\left(\frac{x}{2}, 2\right) \sqrt{a \sec^3(x)} + \frac{10}{21} a \sqrt{a \sec^3(x)} \sin(x) + \frac{2}{7} a \sec(x) \sqrt{a \sec^3(x)} \tan(x)$$

```
output 10/21*a*cos(x)^(3/2)*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticF(sin(1/2*x),
2^(1/2))*(a*sec(x)^3)^(1/2)+10/21*a*sin(x)*(a*sec(x)^3)^(1/2)+2/7*a*sec(x)
*(a*sec(x)^3)^(1/2)*tan(x)
```

#### 3.56.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.66

$$\int (a \sec^3(x))^{3/2} dx = \frac{2}{21} a \sec(x) \sqrt{a \sec^3(x)} \left( 5 \cos^{\frac{5}{2}}(x) \operatorname{EllipticF}\left(\frac{x}{2}, 2\right) + 5 \cos(x) \sin(x) + 3 \tan(x) \right)$$

```
input Integrate[(a*Sec[x]^3)^(3/2),x]
```

```
output (2*a*Sec[x]*Sqrt[a*Sec[x]^3]*(5*Cos[x]^(5/2)*EllipticF[x/2, 2] + 5*Cos[x]*
Sin[x] + 3*Tan[x]))/21
```

**3.56.3 Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.09, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 4611, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^3(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^3)^{3/2} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{a \sqrt{a \sec^3(x)} \int \sec^{\frac{9}{2}}(x) dx}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \sqrt{a \sec^3(x)} \int \csc(x + \frac{\pi}{2})^{9/2} dx}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{4255} \\
 & \frac{a \sqrt{a \sec^3(x)} \left( \frac{5}{7} \int \sec^{\frac{5}{2}}(x) dx + \frac{2}{7} \sin(x) \sec^{\frac{7}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \sqrt{a \sec^3(x)} \left( \frac{5}{7} \int \csc(x + \frac{\pi}{2})^{5/2} dx + \frac{2}{7} \sin(x) \sec^{\frac{7}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{4255} \\
 & \frac{a \sqrt{a \sec^3(x)} \left( \frac{5}{7} \left( \frac{1}{3} \int \sqrt{\sec(x)} dx + \frac{2}{3} \sin(x) \sec^{\frac{3}{2}}(x) \right) + \frac{2}{7} \sin(x) \sec^{\frac{7}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \sqrt{a \sec^3(x)} \left( \frac{5}{7} \left( \frac{1}{3} \int \sqrt{\csc(x + \frac{\pi}{2})} dx + \frac{2}{3} \sin(x) \sec^{\frac{3}{2}}(x) \right) + \frac{2}{7} \sin(x) \sec^{\frac{7}{2}}(x) \right)}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$\frac{a\sqrt{a\sec^3(x)}\left(\frac{5}{7}\left(\frac{1}{3}\sqrt{\cos(x)}\sqrt{\sec(x)}\int\frac{1}{\sqrt{\cos(x)}}dx+\frac{2}{3}\sin(x)\sec^{\frac{3}{2}}(x)\right)+\frac{2}{7}\sin(x)\sec^{\frac{7}{2}}(x)\right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 3042

$$\frac{a\sqrt{a\sec^3(x)}\left(\frac{5}{7}\left(\frac{1}{3}\sqrt{\cos(x)}\sqrt{\sec(x)}\int\frac{1}{\sqrt{\sin(x+\frac{\pi}{2})}}dx+\frac{2}{3}\sin(x)\sec^{\frac{3}{2}}(x)\right)+\frac{2}{7}\sin(x)\sec^{\frac{7}{2}}(x)\right)}{\sec^{\frac{3}{2}}(x)}$$

↓ 3120

$$\frac{a\sqrt{a\sec^3(x)}\left(\frac{2}{7}\sin(x)\sec^{\frac{7}{2}}(x)+\frac{5}{7}\left(\frac{2}{3}\sin(x)\sec^{\frac{3}{2}}(x)+\frac{2}{3}\sqrt{\cos(x)}\sqrt{\sec(x)}\operatorname{EllipticF}\left(\frac{x}{2},2\right)\right)\right)}{\sec^{\frac{3}{2}}(x)}$$

input `Int[(a*Sec[x]^3)^(3/2),x]`

output `(a*Sqrt[a*Sec[x]^3]*((2*Sec[x]^(7/2)*Sin[x])/7 + (5*((2*Sqrt[Cos[x]]*EllipticF[x/2, 2]*Sqrt[Sec[x]])/3 + (2*Sec[x]^(3/2)*Sin[x])/3))/7)/Sec[x]^(3/2)`

### 3.56.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Csc[c + d*x])^(n-1)/(d*(n-1)), x] + Simp[b^2*(n-2)/(n-1)Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`



```
rule 4611 Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :> Simp[b^
IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart
[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x]
&& !IntegerQ[p]
```

### 3.56.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.27 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.54

method	result
default	$-\frac{2i\sqrt{a\sec(x)^3}a\left(5\operatorname{EllipticF}\left(i(\csc(x)-\cot(x)),i\right)\sqrt{\frac{\cos(x)}{\cos(x)+1}}\sqrt{\frac{1}{\cos(x)+1}}\cos(x)^2+5\operatorname{EllipticF}\left(i(\csc(x)-\cot(x)),i\right)\sqrt{\frac{\cos(x)}{\cos(x)+1}}\sqrt{\frac{1}{\cos(x)+1}}\right)}{21}$

```
input int((a*sec(x)^3)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2/21*I*(a*sec(x)^3)^(1/2)*a*(5*EllipticF(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)^2+5*EllipticF(I*(csc(x)-cot(x)),I)*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*cos(x)+5*I*sin(x)+3*I*sec(x)*tan(x))
```

### 3.56.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.14

$$\int (a \sec^3(x))^{3/2} dx = \frac{5i\sqrt{2}a^{3/2}\cos(x)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x)) - 5i\sqrt{2}a^{3/2}\cos(x)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(x) - i \sin(x))}{21 \cos(x)}$$

```
input integrate((a*sec(x)^3)^(3/2),x, algorithm="fracas")
```

```
output 1/21*(5*I*sqrt(2)*a^(3/2)*cos(x)^2*weierstrassPInverse(-4, 0, cos(x) + I*sin(x)) - 5*I*sqrt(2)*a^(3/2)*cos(x)^2*weierstrassPInverse(-4, 0, cos(x) - I*sin(x)) + 2*(5*a*cos(x)^2 + 3*a)*sqrt(a/cos(x)^3)*sin(x)/cos(x)^2
```

**3.56.6 Sympy [F]**

$$\int (a \sec^3(x))^{3/2} dx = \int (a \sec^3(x))^{\frac{3}{2}} dx$$

input `integrate((a*sec(x)**3)**(3/2),x)`

output `Integral((a*sec(x)**3)**(3/2), x)`

**3.56.7 Maxima [F]**

$$\int (a \sec^3(x))^{3/2} dx = \int (a \sec(x)^3)^{\frac{3}{2}} dx$$

input `integrate((a*sec(x)^3)^(3/2),x, algorithm="maxima")`

output `integrate((a*sec(x)^3)^(3/2), x)`

**3.56.8 Giac [F]**

$$\int (a \sec^3(x))^{3/2} dx = \int (a \sec(x)^3)^{\frac{3}{2}} dx$$

input `integrate((a*sec(x)^3)^(3/2),x, algorithm="giac")`

output `integrate((a*sec(x)^3)^(3/2), x)`

**3.56.9 Mupad [F(-1)]**

Timed out.

$$\int (a \sec^3(x))^{3/2} dx = \int \left( \frac{a}{\cos(x)^3} \right)^{3/2} dx$$

input `int((a/cos(x)^3)^(3/2),x)`output `int((a/cos(x)^3)^(3/2), x)`

### 3.57 $\int \sqrt{a \sec^3(x)} dx$

3.57.1	Optimal result	407
3.57.2	Mathematica [A] (verified)	407
3.57.3	Rubi [A] (verified)	408
3.57.4	Maple [C] (verified)	410
3.57.5	Fricas [C] (verification not implemented)	410
3.57.6	Sympy [F]	411
3.57.7	Maxima [F]	411
3.57.8	Giac [F]	411
3.57.9	Mupad [F(-1)]	412

#### 3.57.1 Optimal result

Integrand size = 10, antiderivative size = 42

$$\int \sqrt{a \sec^3(x)} dx = -2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right) \sqrt{a \sec^3(x)} + 2 \cos(x) \sqrt{a \sec^3(x)} \sin(x)$$

output `-2*cos(x)^(3/2)*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticE(sin(1/2*x),2^(1/2))*(a*sec(x)^3)^(1/2)+2*cos(x)*sin(x)*(a*sec(x)^3)^(1/2)`

#### 3.57.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.76

$$\int \sqrt{a \sec^3(x)} dx = 2 \cos(x) \sqrt{a \sec^3(x)} \left( -\sqrt{\cos(x)} E\left(\frac{x}{2} \middle| 2\right) + \sin(x) \right)$$

input `Integrate[Sqrt[a*Sec[x]^3],x]`

output `2*Cos[x]*Sqrt[a*Sec[x]^3]*(-(Sqrt[Cos[x]]*EllipticE[x/2, 2]) + Sin[x])`

**3.57.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.17, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$ , Rules used = {3042, 4611, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \sec^3(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \sec(x)^3} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{\sqrt{a \sec^3(x)} \int \sec^{\frac{3}{2}}(x) dx}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a \sec^3(x)} \int \csc(x + \frac{\pi}{2})^{3/2} dx}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\sqrt{a \sec^3(x)} \left( 2 \sin(x) \sqrt{\sec(x)} - \int \frac{1}{\sqrt{\sec(x)}} dx \right)}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a \sec^3(x)} \left( 2 \sin(x) \sqrt{\sec(x)} - \int \frac{1}{\sqrt{\csc(x + \frac{\pi}{2})}} dx \right)}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sqrt{a \sec^3(x)} \left( 2 \sin(x) \sqrt{\sec(x)} - \sqrt{\cos(x)} \sqrt{\sec(x)} \int \sqrt{\cos(x)} dx \right)}{\sec^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a \sec^3(x)} \left( 2 \sin(x) \sqrt{\sec(x)} - \sqrt{\cos(x)} \sqrt{\sec(x)} \int \sqrt{\sin(x + \frac{\pi}{2})} dx \right)}{\sec^{\frac{3}{2}}(x)}
 \end{aligned}$$

$$\frac{\sqrt{a \sec^3(x)} \left( 2 \sin(x) \sqrt{\sec(x)} - 2 \sqrt{\cos(x)} \sqrt{\sec(x)} E\left(\frac{x}{2} \mid 2\right) \right)}{\sec^{\frac{3}{2}}(x)} \quad \downarrow \quad 3119$$

input `Int[Sqrt[a*Sec[x]^3], x]`

output `(Sqrt[a*Sec[x]^3]*(-2*Sqrt[Cos[x]]*EllipticE[x/2, 2]*Sqrt[Sec[x]] + 2*Sqrt[Sec[x]]*Sin[x]))/Sec[x]^(3/2)`

### 3.57.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

### 3.57.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.76 (sec) , antiderivative size = 244, normalized size of antiderivative = 5.81

method	result
default	$2\left(i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticF}(i(\csc(x)-\cot(x)),i)\cos(x)^2-i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticE}(i(\csc(x)-\cot(x)),i)\cos(x)^2+2i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticF}(i(\csc(x)-\cot(x)),i)\cos(x)-i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticE}(i(\csc(x)-\cot(x)),i)\cos(x)\right)$

input `int((a*sec(x)^3)^(1/2),x,method=_RETURNVERBOSE)`

output `2*(I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*cos(x)^2-I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)*cos(x)^2+2*I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*cos(x)-2*I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)*cos(x)+I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)-I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)+sin(x))*(a*sec(x)^3)^(1/2)*cos(x)/(cos(x)+1)`

### 3.57.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.36

$$\int \sqrt{a \sec^3(x)} dx = 2 \sqrt{\frac{a}{\cos(x)^3}} \cos(x) \sin(x) + i \sqrt{2} \sqrt{a} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x))) - i \sqrt{2} \sqrt{a} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(x) - i \sin(x)))$$

input `integrate((a*sec(x)^3)^(1/2),x, algorithm="fricas")`

output `2*sqrt(a/cos(x)^3)*cos(x)*sin(x) + I*sqrt(2)*sqrt(a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) + I*sin(x))) - I*sqrt(2)*sqrt(a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) - I*sin(x)))`

**3.57.6 Sympy [F]**

$$\int \sqrt{a \sec^3(x)} dx = \int \sqrt{a \sec^3(x)} dx$$

input `integrate((a*sec(x)**3)**(1/2),x)`

output `Integral(sqrt(a*sec(x)**3), x)`

**3.57.7 Maxima [F]**

$$\int \sqrt{a \sec^3(x)} dx = \int \sqrt{a \sec(x)^3} dx$$

input `integrate((a*sec(x)^3)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(a*sec(x)^3), x)`

**3.57.8 Giac [F]**

$$\int \sqrt{a \sec^3(x)} dx = \int \sqrt{a \sec(x)^3} dx$$

input `integrate((a*sec(x)^3)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(a*sec(x)^3), x)`



**3.57.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \sec^3(x)} dx = \int \sqrt{\frac{a}{\cos(x)^3}} dx$$

input `int((a/cos(x)^3)^(1/2),x)`output `int((a/cos(x)^3)^(1/2), x)`

$$3.58 \quad \int \frac{1}{\sqrt{a \sec^3(x)}} dx$$

3.58.1	Optimal result	413
3.58.2	Mathematica [A] (verified)	413
3.58.3	Rubi [A] (verified)	414
3.58.4	Maple [C] (verified)	416
3.58.5	Fricas [C] (verification not implemented)	416
3.58.6	Sympy [F]	417
3.58.7	Maxima [F]	417
3.58.8	Giac [F]	417
3.58.9	Mupad [F(-1)]	418

### 3.58.1 Optimal result

Integrand size = 10, antiderivative size = 44

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx = \frac{2 \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{3 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}} + \frac{2 \tan(x)}{3 \sqrt{a \sec^3(x)}}$$

output  $2/3*(\cos(1/2*x)^2)^{(1/2)}/\cos(1/2*x)*\operatorname{EllipticF}(\sin(1/2*x), 2^{(1/2)})/\cos(x)^{(3/2)}/(a*\sec(x)^3)^{(1/2)}+2/3*\tan(x)/(a*\sec(x)^3)^{(1/2)}$

### 3.58.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.70

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx = \frac{2 \left( \frac{\operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{\cos^{\frac{3}{2}}(x)} + \tan(x) \right)}{3 \sqrt{a \sec^3(x)}}$$

input `Integrate[1/Sqrt[a*Sec[x]^3], x]`

output  $(2*(\operatorname{EllipticF}[x/2, 2]/\operatorname{Cos}[x]^{(3/2)} + \operatorname{Tan}[x]))/(3*\operatorname{Sqrt}[a*\operatorname{Sec}[x]^3])$

**3.58.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.20, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$ , Rules used = {3042, 4611, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \sec^3(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a \sec(x)^3}} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\sec^{\frac{3}{2}}(x)} dx}{\sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\csc(x+\frac{\pi}{2})^{3/2}} dx}{\sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{1}{3} \int \sqrt{\sec(x)} dx + \frac{2 \sin(x)}{3\sqrt{\sec(x)}} \right)}{\sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{1}{3} \int \sqrt{\csc(x+\frac{\pi}{2})} dx + \frac{2 \sin(x)}{3\sqrt{\sec(x)}} \right)}{\sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{1}{3} \sqrt{\cos(x)} \sqrt{\sec(x)} \int \frac{1}{\sqrt{\cos(x)}} dx + \frac{2 \sin(x)}{3\sqrt{\sec(x)}} \right)}{\sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{1}{3} \sqrt{\cos(x)} \sqrt{\sec(x)} \int \frac{1}{\sqrt{\sin(x+\frac{\pi}{2})}} dx + \frac{2 \sin(x)}{3\sqrt{\sec(x)}} \right)}{\sqrt{a \sec^3(x)}}
 \end{aligned}$$

$$\frac{\sec^{\frac{3}{2}}(x) \left( \frac{2 \sin(x)}{3 \sqrt{\sec(x)}} + \frac{2}{3} \sqrt{\cos(x)} \sqrt{\sec(x)} \operatorname{EllipticF}\left(\frac{x}{2}, 2\right) \right)}{\sqrt{a \sec^3(x)}} \quad \downarrow \quad 3120$$

input `Int[1/Sqrt[a*Sec[x]^3],x]`

output `(Sec[x]^(3/2)*((2*sqrt[Cos[x]]*EllipticF[x/2, 2]*sqrt[Sec[x]])/3 + (2*Sin[x])/(3*sqrt[Sec[x]])))/sqrt[a*Sec[x]^3]`

### 3.58.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

### 3.58.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.76 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.09

method	result
default	$-\frac{2\left(i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticF}(i(\csc(x)-\cot(x)),i)\sec(x)+i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticF}(i(\csc(x)-\cot(x)),i)\sec(x)^2-\tan(x)\right)}{3\sqrt{a\sec(x)^3}}$

input `int(1/(a*sec(x)^3)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/3/(a*sec(x)^3)^(1/2)*(I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*sec(x)+I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*sec(x)^2-tan(x))`

### 3.58.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.32

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx = \frac{2\sqrt{\frac{a}{\cos(x)^3}}\cos(x)^2\sin(x) + i\sqrt{2}\sqrt{a}\operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i\sin(x)) - i\sqrt{2}\sqrt{a}\operatorname{weierstrassPInverse}(-4, 0, \cos(x) - i\sin(x))}{3a}$$

input `integrate(1/(a*sec(x)^3)^(1/2),x, algorithm="fricas")`

output `1/3*(2*sqrt(a/cos(x)^3)*cos(x)^2*sin(x) + I*sqrt(2)*sqrt(a)*weierstrassPInverse(-4, 0, cos(x) + I*sin(x)) - I*sqrt(2)*sqrt(a)*weierstrassPInverse(-4, 0, cos(x) - I*sin(x)))/a`

**3.58.6 Sympy [F]**

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx = \int \frac{1}{\sqrt{a \sec^3(x)}} dx$$

input `integrate(1/(a*sec(x)**3)**(1/2), x)`

output `Integral(1/sqrt(a*sec(x)**3), x)`

**3.58.7 Maxima [F]**

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx = \int \frac{1}{\sqrt{a \sec(x)^3}} dx$$

input `integrate(1/(a*sec(x)^3)^(1/2), x, algorithm="maxima")`

output `integrate(1/sqrt(a*sec(x)^3), x)`

**3.58.8 Giac [F]**

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx = \int \frac{1}{\sqrt{a \sec(x)^3}} dx$$

input `integrate(1/(a*sec(x)^3)^(1/2), x, algorithm="giac")`

output `integrate(1/sqrt(a*sec(x)^3), x)`

**3.58.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx = \int \frac{1}{\sqrt{\frac{a}{\cos(x)^3}}} dx$$

input `int(1/(a/cos(x)^3)^(1/2),x)`output `int(1/(a/cos(x)^3)^(1/2), x)`

### 3.59 $\int \frac{1}{(a \sec^3(x))^{3/2}} dx$

3.59.1	Optimal result	419
3.59.2	Mathematica [A] (verified)	419
3.59.3	Rubi [A] (verified)	420
3.59.4	Maple [C] (verified)	422
3.59.5	Fricas [C] (verification not implemented)	423
3.59.6	Sympy [F]	423
3.59.7	Maxima [F]	423
3.59.8	Giac [F]	424
3.59.9	Mupad [F(-1)]	424

#### 3.59.1 Optimal result

Integrand size = 10, antiderivative size = 73

$$\int \frac{1}{(a \sec^3(x))^{3/2}} dx = \frac{14E\left(\frac{x}{2} \mid 2\right)}{15a \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}} + \frac{14 \sin(x)}{45a \sqrt{a \sec^3(x)}} + \frac{2 \cos^2(x) \sin(x)}{9a \sqrt{a \sec^3(x)}}$$

output `14/15*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticE(sin(1/2*x),2^(1/2))/a/cos(x)^(3/2)/(a*sec(x)^3)^(1/2)+14/45*sin(x)/a/(a*sec(x)^3)^(1/2)+2/9*cos(x)^2*sin(x)/a/(a*sec(x)^3)^(1/2)`

#### 3.59.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.59

$$\int \frac{1}{(a \sec^3(x))^{3/2}} dx = \frac{\frac{84E\left(\frac{x}{2} \mid 2\right)}{\cos^{\frac{3}{2}}(x)} + 33 \sin(x) + 5 \sin(3x)}{90a \sqrt{a \sec^3(x)}}$$

input `Integrate[(a*Sec[x]^3)^(-3/2),x]`

output `((84*EllipticE[x/2, 2])/Cos[x]^(3/2) + 33*Sin[x] + 5*Sin[3*x])/(90*a*Sqrt[a*Sec[x]^3])`



**3.59.3 Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 4611, 3042, 4256, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sec^3(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sec(x)^3)^{3/2}} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\sec^{\frac{9}{2}}(x)} dx}{a \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\csc(x+\frac{\pi}{2})^{9/2}} dx}{a \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{7}{9} \int \frac{1}{\sec^{\frac{5}{2}}(x)} dx + \frac{2 \sin(x)}{9 \sec^{\frac{7}{2}}(x)} \right)}{a \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{7}{9} \int \frac{1}{\csc(x+\frac{\pi}{2})^{5/2}} dx + \frac{2 \sin(x)}{9 \sec^{\frac{7}{2}}(x)} \right)}{a \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{7}{9} \left( \frac{3}{5} \int \frac{1}{\sqrt{\sec(x)}} dx + \frac{2 \sin(x)}{5 \sec^{\frac{3}{2}}(x)} \right) + \frac{2 \sin(x)}{9 \sec^{\frac{7}{2}}(x)} \right)}{a \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{\sec^{\frac{3}{2}}(x) \left( \frac{7}{9} \left( \frac{3}{5} \int \frac{1}{\sqrt{\csc(x+\frac{\pi}{2})}} dx + \frac{2 \sin(x)}{5 \sec^{\frac{3}{2}}(x)} \right) + \frac{2 \sin(x)}{9 \sec^{\frac{7}{2}}(x)} \right)}{a \sqrt{a \sec^3(x)}}$$

↓ 4258

$$\frac{\sec^{\frac{3}{2}}(x) \left( \frac{7}{9} \left( \frac{3}{5} \sqrt{\cos(x)} \sqrt{\sec(x)} \int \sqrt{\cos(x)} dx + \frac{2 \sin(x)}{5 \sec^{\frac{3}{2}}(x)} \right) + \frac{2 \sin(x)}{9 \sec^{\frac{7}{2}}(x)} \right)}{a \sqrt{a \sec^3(x)}}$$

↓ 3042

$$\frac{\sec^{\frac{3}{2}}(x) \left( \frac{7}{9} \left( \frac{3}{5} \sqrt{\cos(x)} \sqrt{\sec(x)} \int \sqrt{\sin(x+\frac{\pi}{2})} dx + \frac{2 \sin(x)}{5 \sec^{\frac{3}{2}}(x)} \right) + \frac{2 \sin(x)}{9 \sec^{\frac{7}{2}}(x)} \right)}{a \sqrt{a \sec^3(x)}}$$

↓ 3119

$$\frac{\sec^{\frac{3}{2}}(x) \left( \frac{2 \sin(x)}{9 \sec^{\frac{7}{2}}(x)} + \frac{7}{9} \left( \frac{2 \sin(x)}{5 \sec^{\frac{3}{2}}(x)} + \frac{6}{5} \sqrt{\cos(x)} \sqrt{\sec(x)} E\left(\frac{x}{2} \mid 2\right) \right) \right)}{a \sqrt{a \sec^3(x)}}$$

input `Int[(a*Sec[x]^3)^(-3/2),x]`

output `(Sec[x]^(3/2)*((2*Sin[x])/(9*Sec[x]^(7/2)) + (7*((6*sqrt[Cos[x]]*EllipticE[x/2, 2]*sqrt[Sec[x]])/5 + (2*Sin[x])/(5*Sec[x]^(3/2))))/9))/(a*sqrt[a*Sec[x]^3])`

### 3.59.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x]
)^(n)*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

```
rule 4611 Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :> Simp[b^
IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart
[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x]
&& !IntegerQ[p]
```

### 3.59.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 5.08 (sec) , antiderivative size = 273, normalized size of antiderivative = 3.74

method	result
default	$-\frac{2\left(21i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticF}(i(\csc(x)-\cot(x)),i)-21i\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\operatorname{EllipticE}(i(\csc(x)-\cot(x)),i)-5\cos(x)^3\sin(x)\right)}{\dots}$

```
input int(1/(a*sec(x)^3)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2/45/(cos(x)+1)/(a*sec(x)^3)^(1/2)/a*(21*I*(1/(cos(x)+1))^(1/2)*(cos(x)/(
cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)-21*I*(1/(cos(x)+1))^(1/2)*
(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)-5*cos(x)^3*sin(x)
+42*I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-c
ot(x)),I)*sec(x)-42*I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*Ellip
ticE(I*(csc(x)-cot(x)),I)*sec(x)-5*sin(x)*cos(x)^2+21*I*(1/(cos(x)+1))^(1/
2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*sec(x)^2-21*I*
(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x))
,I)*sec(x)^2-7*sin(x)*cos(x)-7*sin(x)-21*tan(x))
```

**3.59.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a \sec^3(x))^{3/2}} dx = \frac{2(5 \cos(x)^5 + 7 \cos(x)^3) \sqrt{\frac{a}{\cos(x)^3}} \sin(x) - 21i \sqrt{2} \sqrt{a} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(x) + I \sin(x))) + 21i \sqrt{2} \sqrt{a} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(x) - I \sin(x)))}{a^2}$$

input `integrate(1/(a*sec(x)^3)^(3/2),x, algorithm="fricas")`

output `1/45*(2*(5*cos(x)^5 + 7*cos(x)^3)*sqrt(a/cos(x)^3)*sin(x) - 21*I*sqrt(2)*sqrt(a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) + I*sin(x))) + 21*I*sqrt(2)*sqrt(a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) - I*sin(x))))/a^2`

**3.59.6 Sympy [F]**

$$\int \frac{1}{(a \sec^3(x))^{3/2}} dx = \int \frac{1}{(a \sec^3(x))^{\frac{3}{2}}} dx$$

input `integrate(1/(a*sec(x)**3)**(3/2),x)`

output `Integral((a*sec(x)**3)**(-3/2), x)`

**3.59.7 Maxima [F]**

$$\int \frac{1}{(a \sec^3(x))^{3/2}} dx = \int \frac{1}{(a \sec(x)^3)^{\frac{3}{2}}} dx$$

input `integrate(1/(a*sec(x)^3)^(3/2),x, algorithm="maxima")`

output `integrate((a*sec(x)^3)^(-3/2), x)`

**3.59.8 Giac [F]**

$$\int \frac{1}{(a \sec^3(x))^{3/2}} dx = \int \frac{1}{(a \sec(x)^3)^{3/2}} dx$$

input `integrate(1/(a*sec(x)^3)^(3/2),x, algorithm="giac")`

output `integrate((a*sec(x)^3)^(-3/2), x)`

**3.59.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \sec^3(x))^{3/2}} dx = \int \frac{1}{\left(\frac{a}{\cos(x)^3}\right)^{3/2}} dx$$

input `int(1/(a/cos(x)^3)^(3/2),x)`

output `int(1/(a/cos(x)^3)^(3/2), x)`

### 3.60 $\int \frac{1}{(a \sec^3(x))^{5/2}} dx$

3.60.1	Optimal result . . . . .	425
3.60.2	Mathematica [A] (verified) . . . . .	425
3.60.3	Rubi [A] (verified) . . . . .	426
3.60.4	Maple [C] (verified) . . . . .	429
3.60.5	Fricas [C] (verification not implemented) . . . . .	429
3.60.6	Sympy [F] . . . . .	430
3.60.7	Maxima [F] . . . . .	430
3.60.8	Giac [F] . . . . .	430
3.60.9	Mupad [F(-1)] . . . . .	431

#### 3.60.1 Optimal result

Integrand size = 10, antiderivative size = 117

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx = \frac{26 \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{77a^2 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}} + \frac{78 \cos(x) \sin(x)}{385a^2 \sqrt{a \sec^3(x)}} \\ + \frac{26 \cos^3(x) \sin(x)}{165a^2 \sqrt{a \sec^3(x)}} + \frac{2 \cos^5(x) \sin(x)}{15a^2 \sqrt{a \sec^3(x)}} + \frac{26 \tan(x)}{77a^2 \sqrt{a \sec^3(x)}}$$

output `26/77*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticF(sin(1/2*x), 2^(1/2))/a^2/cos(x)^(3/2)/(a*sec(x)^3)^(1/2)+78/385*cos(x)*sin(x)/a^2/(a*sec(x)^3)^(1/2)+26/165*cos(x)^3*sin(x)/a^2/(a*sec(x)^3)^(1/2)+2/15*cos(x)^5*sin(x)/a^2/(a*sec(x)^3)^(1/2)+26/77*tan(x)/a^2/(a*sec(x)^3)^(1/2)`

#### 3.60.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.50

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx = \frac{\cos(x) \sqrt{a \sec^3(x)} \left( 24960 \sqrt{\cos(x)} \operatorname{EllipticF}\left(\frac{x}{2}, 2\right) + 19122 \sin(2x) + 4406 \sin(4x) + 826 \sin(6x) + 77 \sin(8x) \right)}{73920a^3}$$

input `Integrate[(a*Sec[x]^3)^(-5/2), x]`

output `(Cos[x]*Sqrt[a*Sec[x]^3]*(24960*Sqrt[Cos[x]]*EllipticF[x/2, 2] + 19122*Sin[2*x] + 4406*Sin[4*x] + 826*Sin[6*x] + 77*Sin[8*x]))/(73920*a^3)`

**3.60.3 Rubi [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.91, number of steps used = 14, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.400$ , Rules used = {3042, 4611, 3042, 4256, 3042, 4256, 3042, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sec^3(x))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sec(x)^3)^{5/2}} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\sec^{\frac{15}{2}}(x)} dx}{a^2 \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\csc(x+\frac{\pi}{2})^{15/2}} dx}{a^2 \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \int \frac{1}{\sec^{\frac{11}{2}}(x)} dx + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \int \frac{1}{\csc(x+\frac{\pi}{2})^{11/2}} dx + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \int \frac{1}{\sec^{\frac{7}{2}}(x)} dx + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \int \frac{1}{\csc(x+\frac{\pi}{2})^{7/2}} dx + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
& \quad \downarrow 4256 \\
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \left( \frac{5}{7} \int \frac{1}{\sec^{\frac{3}{2}}(x)} dx + \frac{2 \sin(x)}{7 \sec^{\frac{5}{2}}(x)} \right) + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
& \quad \downarrow 3042 \\
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \left( \frac{5}{7} \int \frac{1}{\csc(x+\frac{\pi}{2})^{3/2}} dx + \frac{2 \sin(x)}{7 \sec^{\frac{5}{2}}(x)} \right) + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
& \quad \downarrow 4256 \\
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \left( \frac{5}{7} \left( \frac{1}{3} \int \sqrt{\sec(x)} dx + \frac{2 \sin(x)}{3 \sqrt{\sec(x)}} \right) + \frac{2 \sin(x)}{7 \sec^{\frac{5}{2}}(x)} \right) + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
& \quad \downarrow 3042 \\
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \left( \frac{5}{7} \left( \frac{1}{3} \int \sqrt{\csc(x+\frac{\pi}{2})} dx + \frac{2 \sin(x)}{3 \sqrt{\sec(x)}} \right) + \frac{2 \sin(x)}{7 \sec^{\frac{5}{2}}(x)} \right) + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
& \quad \downarrow 4258 \\
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \left( \frac{5}{7} \left( \frac{1}{3} \sqrt{\cos(x)} \sqrt{\sec(x)} \int \frac{1}{\sqrt{\cos(x)}} dx + \frac{2 \sin(x)}{3 \sqrt{\sec(x)}} \right) + \frac{2 \sin(x)}{7 \sec^{\frac{5}{2}}(x)} \right) + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
& \quad \downarrow 3042 \\
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{13}{15} \left( \frac{9}{11} \left( \frac{5}{7} \left( \frac{1}{3} \sqrt{\cos(x)} \sqrt{\sec(x)} \int \frac{1}{\sqrt{\sin(x+\frac{\pi}{2})}} dx + \frac{2 \sin(x)}{3 \sqrt{\sec(x)}} \right) + \frac{2 \sin(x)}{7 \sec^{\frac{5}{2}}(x)} \right) + \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} \right) + \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} \right)}{a^2 \sqrt{a \sec^3(x)}} \\
& \quad \downarrow 3120 \\
& \frac{\sec^{\frac{3}{2}}(x) \left( \frac{2 \sin(x)}{15 \sec^{\frac{13}{2}}(x)} + \frac{13}{15} \left( \frac{2 \sin(x)}{11 \sec^{\frac{9}{2}}(x)} + \frac{9}{11} \left( \frac{2 \sin(x)}{7 \sec^{\frac{5}{2}}(x)} + \frac{5}{7} \left( \frac{2 \sin(x)}{3 \sqrt{\sec(x)}} + \frac{2}{3} \sqrt{\cos(x)} \sqrt{\sec(x)} \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)\right) \right) \right)}{a^2 \sqrt{a \sec^3(x)}}
\end{aligned}$$



input `Int[(a*Sec[x]^3)^(-5/2),x]`

output `(Sec[x]^(3/2)*((2*Sin[x])/(15*Sec[x]^(13/2)) + (13*((2*Sin[x])/(11*Sec[x]^(9/2)) + (9*((2*Sin[x])/(7*Sec[x]^(5/2)) + (5*((2*Sqrt[Cos[x]]*EllipticF[x/2, 2]*Sqrt[Sec[x]]))/3 + (2*Sin[x])/(3*Sqrt[Sec[x]]))))/7)/11)/15)/(a^2*Sqrt[a*Sec[x]^3])`

### 3.60.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

### 3.60.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 5.57 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00

method	result
default	$-\frac{2(-77 \sin(x) \cos(x)^5 - 91 \cos(x)^3 \sin(x) + 195i \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} \operatorname{EllipticF}(i(\csc(x) - \cot(x)), i) \sec(x) + 195i \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}})}{1155 \sqrt{a \sec(x)^3} a^2}$

input `int(1/(a*sec(x)^3)^(5/2),x,method=_RETURNVERBOSE)`

output `-2/1155/(a*sec(x)^3)^(1/2)/a^2*(-77*sin(x)*cos(x)^5-91*cos(x)^3*sin(x)+195*I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*sec(x)+195*I*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*sec(x)^2-117*sin(x)*cos(x)-195*tan(x))`

### 3.60.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.68

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx = \frac{2(77 \cos(x)^8 + 91 \cos(x)^6 + 117 \cos(x)^4 + 195 \cos(x)^2) \sqrt{\frac{a}{\cos(x)^3}} \sin(x) + 195i \sqrt{2}}{(a \sec^3(x))^{5/2}}$$

input `integrate(1/(a*sec(x)^3)^(5/2),x, algorithm="fricas")`

output `1/1155*(2*(77*cos(x)^8 + 91*cos(x)^6 + 117*cos(x)^4 + 195*cos(x)^2)*sqrt(a/cos(x)^3)*sin(x) + 195*I*sqrt(2)*sqrt(a)*weierstrassPInverse(-4, 0, cos(x) + I*sin(x)) - 195*I*sqrt(2)*sqrt(a)*weierstrassPInverse(-4, 0, cos(x) - I*sin(x)))/a^3`

**3.60.6 Sympy [F]**

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx = \int \frac{1}{(a \sec^3(x))^{\frac{5}{2}}} dx$$

input `integrate(1/(a*sec(x)**3)**(5/2), x)`

output `Integral((a*sec(x)**3)**(-5/2), x)`

**3.60.7 Maxima [F]**

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx = \int \frac{1}{(a \sec(x)^3)^{\frac{5}{2}}} dx$$

input `integrate(1/(a*sec(x)^3)^(5/2), x, algorithm="maxima")`

output `integrate((a*sec(x)^3)^(-5/2), x)`

**3.60.8 Giac [F]**

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx = \int \frac{1}{(a \sec(x)^3)^{\frac{5}{2}}} dx$$

input `integrate(1/(a*sec(x)^3)^(5/2), x, algorithm="giac")`

output `integrate((a*sec(x)^3)^(-5/2), x)`

**3.60.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx = \int \frac{1}{\left(\frac{a}{\cos(x)^3}\right)^{5/2}} dx$$

input `int(1/(a/cos(x)^3)^(5/2),x)`output `int(1/(a/cos(x)^3)^(5/2), x)`

### 3.61 $\int (a \sec^4(x))^{7/2} dx$

3.61.1	Optimal result . . . . .	432
3.61.2	Mathematica [A] (verified) . . . . .	432
3.61.3	Rubi [A] (verified) . . . . .	433
3.61.4	Maple [A] (verified) . . . . .	434
3.61.5	Fricas [A] (verification not implemented) . . . . .	435
3.61.6	Sympy [F(-1)] . . . . .	435
3.61.7	Maxima [A] (verification not implemented) . . . . .	435
3.61.8	Giac [A] (verification not implemented) . . . . .	436
3.61.9	Mupad [B] (verification not implemented) . . . . .	436

#### 3.61.1 Optimal result

Integrand size = 10, antiderivative size = 163

$$\begin{aligned} \int (a \sec^4(x))^{7/2} dx &= a^3 \cos(x) \sqrt{a \sec^4(x)} \sin(x) \\ &+ 2a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan(x) + 3a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan^3(x) \\ &+ \frac{20}{7} a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan^5(x) + \frac{5}{3} a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan^7(x) \\ &+ \frac{6}{11} a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan^9(x) + \frac{1}{13} a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan^{11}(x) \end{aligned}$$

```
output a^3*cos(x)*sin(x)*(a*sec(x)^4)^(1/2)+2*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan
(x)+3*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^3+20/7*a^3*sin(x)^2*(a*sec
(x)^4)^(1/2)*tan(x)^5+5/3*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^7+6/11*a^3*s
in(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^9+1/13*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*t
an(x)^11
```

#### 3.61.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.33

$$\int (a \sec^4(x))^{7/2} dx = \frac{\cos(x)(2048 + 2380 \cos(2x) + 1093 \cos(4x) + 378 \cos(6x) + 92 \cos(8x) + 14 \cos(10x))}{6006}$$

```
input Integrate[(a*Sec[x]^4)^(7/2),x]
```

output  $(\text{Cos}[x]*(2048 + 2380*\text{Cos}[2*x] + 1093*\text{Cos}[4*x] + 378*\text{Cos}[6*x] + 92*\text{Cos}[8*x] + 14*\text{Cos}[10*x] + \text{Cos}[12*x])*(a*\text{Sec}[x]^4)^{(7/2)}*\text{Sin}[x])/6006$

### 3.61.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.42, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4611, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^4(x))^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^4)^{7/2} dx \\
 & \quad \downarrow \text{4611} \\
 & a^3 \cos^2(x) \sqrt{a \sec^4(x)} \int \sec^{14}(x) dx \\
 & \quad \downarrow \text{3042} \\
 & a^3 \cos^2(x) \sqrt{a \sec^4(x)} \int \csc\left(x + \frac{\pi}{2}\right)^{14} dx \\
 & \quad \downarrow \text{4254} \\
 & -a^3 \cos^2(x) \sqrt{a \sec^4(x)} \int (\tan^{12}(x) + 6 \tan^{10}(x) + 15 \tan^8(x) + 20 \tan^6(x) + 15 \tan^4(x) + 6 \tan^2(x) + 1) d(-\tan(x)) \\
 & \quad \downarrow \text{2009} \\
 & -a^3 \cos^2(x) \left( -\frac{1}{13} \tan^{13}(x) - \frac{6 \tan^{11}(x)}{11} - \frac{5 \tan^9(x)}{3} - \frac{20 \tan^7(x)}{7} - 3 \tan^5(x) - 2 \tan^3(x) - \tan(x) \right) \sqrt{a \sec^4(x)}
 \end{aligned}$$

input  $\text{Int}[(a*\text{Sec}[x]^4)^{(7/2)}, x]$

output  $-(a^3*\text{Cos}[x]^2*\text{Sqrt}[a*\text{Sec}[x]^4]*(-\text{Tan}[x] - 2*\text{Tan}[x]^3 - 3*\text{Tan}[x]^5 - (20*\text{Tan}[x]^7)/7 - (5*\text{Tan}[x]^9)/3 - (6*\text{Tan}[x]^11)/11 - \text{Tan}[x]^13/13))\text{Sqrt}[a*\text{Sec}[x]^4]$

---

3.61.  $\int (a \sec^4(x))^{7/2} dx$

## 3.61.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.61.4 Maple [A] (verified)

Time = 97.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.36

method	result	size
default	$\frac{\sqrt{a \sec(x)^4} a^3 (1024 \cos(x)^{12} + 512 \cos(x)^{10} + 384 \cos(x)^8 + 320 \cos(x)^6 + 280 \cos(x)^4 + 252 \cos(x)^2 + 231) \tan(x) \sec(x)^{10}}{3003}$	58
risch	$\frac{2048ia^3 \sqrt{\frac{a e^{4ix}}{(e^{2ix}+1)^4}} (1716 e^{10ix} + 1287 e^{8ix} + 715 e^{6ix} + 286 e^{4ix} + 13 + 79 \cos(2x) + 77i \sin(2x))}{3003(e^{2ix}+1)^{11}}$	77

input `int((a*sec(x)^4)^(7/2),x,method=_RETURNVERBOSE)`

output `1/3003*(a*sec(x)^4)^(1/2)*a^3*(1024*cos(x)^12+512*cos(x)^10+384*cos(x)^8+320*cos(x)^6+280*cos(x)^4+252*cos(x)^2+231)*tan(x)*sec(x)^10`

**3.61.5 Fracas [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.47

$$\int (a \sec^4(x))^{7/2} dx = \frac{(1024 a^3 \cos(x)^{12} + 512 a^3 \cos(x)^{10} + 384 a^3 \cos(x)^8 + 320 a^3 \cos(x)^6 + 280 a^3 \cos(x)^4 + 252 a^3 \cos(x)^2 + 231 a^3) \sqrt{a/\cos(x)} \sin(x)}{3003 \cos(x)^{11}}$$

input `integrate((a*sec(x)^4)^(7/2),x, algorithm="fricas")`output `1/3003*(1024*a^3*cos(x)^12 + 512*a^3*cos(x)^10 + 384*a^3*cos(x)^8 + 320*a^3*cos(x)^6 + 280*a^3*cos(x)^4 + 252*a^3*cos(x)^2 + 231*a^3)*sqrt(a/cos(x)^4)*sin(x)/cos(x)^11`**3.61.6 Sympy [F(-1)]**

Timed out.

$$\int (a \sec^4(x))^{7/2} dx = \text{Timed out}$$

input `integrate((a*sec(x)**4)**(7/2),x)`output `Timed out`**3.61.7 Maxima [A] (verification not implemented)**

Time = 0.43 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.37

$$\int (a \sec^4(x))^{7/2} dx = \frac{1}{13} a^{7/2} \tan(x)^{13} + \frac{6}{11} a^{7/2} \tan(x)^{11} + \frac{5}{3} a^{7/2} \tan(x)^9 + \frac{20}{7} a^{7/2} \tan(x)^7 + 3 a^{7/2} \tan(x)^5 + 2 a^{7/2} \tan(x)^3 + a^{7/2} \tan(x)$$

input `integrate((a*sec(x)^4)^(7/2),x, algorithm="maxima")`output `1/13*a^(7/2)*tan(x)^13 + 6/11*a^(7/2)*tan(x)^11 + 5/3*a^(7/2)*tan(x)^9 + 20/7*a^(7/2)*tan(x)^7 + 3*a^(7/2)*tan(x)^5 + 2*a^(7/2)*tan(x)^3 + a^(7/2)*tan(x)`



**3.61.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.41

$$\int (a \sec^4(x))^{7/2} dx = \frac{1}{3003} (231 a^3 \tan(x)^{13} + 1638 a^3 \tan(x)^{11} + 5005 a^3 \tan(x)^9 + 8580 a^3 \tan(x)^7 + 9009 a^3 \tan(x)^5 + 6006 a^3 \tan(x)^3 + 3003 a^3 \tan(x)) \operatorname{sqrt}(a)$$

input `integrate((a*sec(x)^4)^(7/2),x, algorithm="giac")`output `1/3003*(231*a^3*tan(x)^13 + 1638*a^3*tan(x)^11 + 5005*a^3*tan(x)^9 + 8580*a^3*tan(x)^7 + 9009*a^3*tan(x)^5 + 6006*a^3*tan(x)^3 + 3003*a^3*tan(x))*sqrt(a)`**3.61.9 Mupad [B] (verification not implemented)**

Time = 17.50 (sec) , antiderivative size = 589, normalized size of antiderivative = 3.61

$$\int (a \sec^4(x))^{7/2} dx = \text{Too large to display}$$

input `int((a/cos(x)^4)^(7/2),x)`output `(a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*2048i)/(7*(exp(x*2i) + 1)^7*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*1536i)/((exp(x*2i) + 1)^8*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*10240i)/(3*(exp(x*2i) + 1)^9*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*4096i)/((exp(x*2i) + 1)^10*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*30720i)/(11*(exp(x*2i) + 1)^11*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*1024i)/((exp(x*2i) + 1)^12*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*2048i)/(13*(exp(x*2i) + 1)^13*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i)))`

### 3.62 $\int (a \sec^4(x))^{5/2} dx$

3.62.1	Optimal result . . . . .	437
3.62.2	Mathematica [A] (verified) . . . . .	437
3.62.3	Rubi [A] (verified) . . . . .	438
3.62.4	Maple [A] (verified) . . . . .	439
3.62.5	Fricas [A] (verification not implemented) . . . . .	440
3.62.6	Sympy [F] . . . . .	440
3.62.7	Maxima [A] (verification not implemented) . . . . .	440
3.62.8	Giac [A] (verification not implemented) . . . . .	441
3.62.9	Mupad [B] (verification not implemented) . . . . .	441

#### 3.62.1 Optimal result

Integrand size = 10, antiderivative size = 117

$$\int (a \sec^4(x))^{5/2} dx = a^2 \cos(x) \sqrt{a \sec^4(x)} \sin(x) + \frac{4}{3} a^2 \sqrt{a \sec^4(x)} \sin^2(x) \tan(x) + \frac{6}{5} a^2 \sqrt{a \sec^4(x)} \sin^2(x) \tan^3(x) + \frac{4}{7} a^2 \sqrt{a \sec^4(x)} \sin^2(x) \tan^5(x) + \frac{1}{9} a^2 \sqrt{a \sec^4(x)} \sin^2(x) \tan^7(x)$$

output `a^2*cos(x)*sin(x)*(a*sec(x)^4)^(1/2)+4/3*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)+6/5*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^3+4/7*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^5+1/9*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^7`

#### 3.62.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.36

$$\int (a \sec^4(x))^{5/2} dx = \frac{1}{315} \cos(x)(128 + 130 \cos(2x) + 46 \cos(4x) + 10 \cos(6x) + \cos(8x)) (a \sec^4(x))^{5/2} \sin(x)$$

input `Integrate[(a*Sec[x]^4)^(5/2),x]`

output `(Cos[x]*(128 + 130*Cos[2*x] + 46*Cos[4*x] + 10*Cos[6*x] + Cos[8*x]))*(a*Sec[x]^4)^(5/2)*Sin[x])/315`

**3.62.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.48, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4611, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^4(x))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^4)^{5/2} dx \\
 & \quad \downarrow \text{4611} \\
 & a^2 \cos^2(x) \sqrt{a \sec^4(x)} \int \sec^{10}(x) dx \\
 & \quad \downarrow \text{3042} \\
 & a^2 \cos^2(x) \sqrt{a \sec^4(x)} \int \csc\left(x + \frac{\pi}{2}\right)^{10} dx \\
 & \quad \downarrow \text{4254} \\
 & -a^2 \cos^2(x) \sqrt{a \sec^4(x)} \int (\tan^8(x) + 4 \tan^6(x) + 6 \tan^4(x) + 4 \tan^2(x) + 1) d(-\tan(x)) \\
 & \quad \downarrow \text{2009} \\
 & -a^2 \cos^2(x) \left( -\frac{1}{9} \tan^9(x) - \frac{4 \tan^7(x)}{7} - \frac{6 \tan^5(x)}{5} - \frac{4 \tan^3(x)}{3} - \tan(x) \right) \sqrt{a \sec^4(x)}
 \end{aligned}$$

input `Int[(a*Sec[x]^4)^(5/2),x]`

output `-(a^2*Cos[x]^2*Sqrt[a*Sec[x]^4]*(-Tan[x] - (4*Tan[x]^3)/3 - (6*Tan[x]^5)/5 - (4*Tan[x]^7)/7 - Tan[x]^9/9))`

## 3.62.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.62.4 Maple [A] (verified)

Time = 95.85 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.39

method	result	size
default	$\frac{a^2 \sqrt{a \sec(x)^4} (128 \cos(x)^8 + 64 \cos(x)^6 + 48 \cos(x)^4 + 40 \cos(x)^2 + 35) \tan(x) \sec(x)^6}{315}$	46
risch	$\frac{256ia^2 \sqrt{\frac{ae^{4ix}}{(e^{2ix}+1)^4}} (126 e^{6ix} + 84 e^{4ix} + 9 + 37 \cos(2x) + 35i \sin(2x))}{315(e^{2ix}+1)^7}$	63

input `int((a*sec(x)^4)^(5/2),x,method=_RETURNVERBOSE)`

output `1/315*a^2*(a*sec(x)^4)^(1/2)*(128*cos(x)^8+64*cos(x)^6+48*cos(x)^4+40*cos(x)^2+35)*tan(x)*sec(x)^6`

**3.62.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.50

$$\int (a \sec^4(x))^{5/2} dx = \frac{(128 a^2 \cos(x)^8 + 64 a^2 \cos(x)^6 + 48 a^2 \cos(x)^4 + 40 a^2 \cos(x)^2 + 35 a^2) \sqrt{\frac{a}{\cos(x)^4}} \sin(x)}{315 \cos(x)^7}$$

input `integrate((a*sec(x)^4)^(5/2),x, algorithm="fricas")`

output `1/315*(128*a^2*cos(x)^8 + 64*a^2*cos(x)^6 + 48*a^2*cos(x)^4 + 40*a^2*cos(x)^2 + 35*a^2)*sqrt(a/cos(x)^4)*sin(x)/cos(x)^7`

**3.62.6 Sympy [F]**

$$\int (a \sec^4(x))^{5/2} dx = \int (a \sec^4(x))^{\frac{5}{2}} dx$$

input `integrate((a*sec(x)**4)**(5/2),x)`

output `Integral((a*sec(x)**4)**(5/2), x)`

**3.62.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.37

$$\int (a \sec^4(x))^{5/2} dx = \frac{1}{9} a^{\frac{5}{2}} \tan(x)^9 + \frac{4}{7} a^{\frac{5}{2}} \tan(x)^7 + \frac{6}{5} a^{\frac{5}{2}} \tan(x)^5 + \frac{4}{3} a^{\frac{5}{2}} \tan(x)^3 + a^{\frac{5}{2}} \tan(x)$$

input `integrate((a*sec(x)^4)^(5/2),x, algorithm="maxima")`

output `1/9*a^(5/2)*tan(x)^9 + 4/7*a^(5/2)*tan(x)^7 + 6/5*a^(5/2)*tan(x)^5 + 4/3*a^(5/2)*tan(x)^3 + a^(5/2)*tan(x)`

**3.62.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.42

$$\int (a \sec^4(x))^{5/2} dx = \frac{1}{315} (35 a^2 \tan(x)^9 + 180 a^2 \tan(x)^7 + 378 a^2 \tan(x)^5 + 420 a^2 \tan(x)^3 + 315 a^2 \tan(x)) \sqrt{a}$$

input `integrate((a*sec(x)^4)^(5/2),x, algorithm="giac")`output `1/315*(35*a^2*tan(x)^9 + 180*a^2*tan(x)^7 + 378*a^2*tan(x)^5 + 420*a^2*tan(x)^3 + 315*a^2*tan(x))*sqrt(a)`**3.62.9 Mupad [B] (verification not implemented)**

Time = 15.13 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.02

$$\int (a \sec^4(x))^{5/2} dx = \frac{128 a^{5/2} (e^{x 46i} 1i + e^{x 48i} 9i + e^{x 50i} 36i + e^{x 52i} 84i + e^{x 54i} 126i)}{315 \left( \frac{e^{-x 2i}}{2} + \frac{e^{x 2i}}{2} + 1 \right) (e^{x 48i} + 7 e^{x 50i} + 21 e^{x 52i} + 35 e^{x 54i} + 35 e^{x 56i} + 21 e^{x 58i} + 7 e^{x 60i} + e^{x 62i})}$$

input `int((a/cos(x)^4)^(5/2),x)`output `(128*a^(5/2)*(exp(x*46i)*1i + exp(x*48i)*9i + exp(x*50i)*36i + exp(x*52i)*84i + exp(x*54i)*126i))/(315*(exp(-x*2i)/2 + exp(x*2i)/2 + 1)*(exp(x*48i) + 7*exp(x*50i) + 21*exp(x*52i) + 35*exp(x*54i) + 35*exp(x*56i) + 21*exp(x*58i) + 7*exp(x*60i) + exp(x*62i)))`

### 3.63 $\int (a \sec^4(x))^{3/2} dx$

3.63.1	Optimal result . . . . .	442
3.63.2	Mathematica [A] (verified) . . . . .	442
3.63.3	Rubi [A] (verified) . . . . .	443
3.63.4	Maple [A] (verified) . . . . .	444
3.63.5	Fricas [A] (verification not implemented) . . . . .	445
3.63.6	Sympy [F] . . . . .	445
3.63.7	Maxima [A] (verification not implemented) . . . . .	445
3.63.8	Giac [A] (verification not implemented) . . . . .	446
3.63.9	Mupad [B] (verification not implemented) . . . . .	446

#### 3.63.1 Optimal result

Integrand size = 10, antiderivative size = 61

$$\int (a \sec^4(x))^{3/2} dx = a \cos(x) \sqrt{a \sec^4(x)} \sin(x) + \frac{2}{3} a \sqrt{a \sec^4(x)} \sin^2(x) \tan(x) + \frac{1}{5} a \sqrt{a \sec^4(x)} \sin^2(x) \tan^3(x)$$

output `a*cos(x)*sin(x)*(a*sec(x)^4)^(1/2)+2/3*a*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)+1/5*a*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^3`

#### 3.63.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.54

$$\int (a \sec^4(x))^{3/2} dx = \frac{1}{15} a \cos(x) \sqrt{a \sec^4(x)} \sin(x) (15 + 10 \tan^2(x) + 3 \tan^4(x))$$

input `Integrate[(a*Sec[x]^4)^(3/2),x]`

output `(a*Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x]*(15 + 10*Tan[x]^2 + 3*Tan[x]^4))/15`

**3.63.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.62, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4611, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec^4(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sec(x)^4)^{3/2} dx \\
 & \quad \downarrow \text{4611} \\
 & a \cos^2(x) \sqrt{a \sec^4(x)} \int \sec^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & a \cos^2(x) \sqrt{a \sec^4(x)} \int \csc\left(x + \frac{\pi}{2}\right)^6 dx \\
 & \quad \downarrow \text{4254} \\
 & -a \cos^2(x) \sqrt{a \sec^4(x)} \int (\tan^4(x) + 2 \tan^2(x) + 1) d(-\tan(x)) \\
 & \quad \downarrow \text{2009} \\
 & -a \cos^2(x) \left( -\frac{1}{5} \tan^5(x) - \frac{2 \tan^3(x)}{3} - \tan(x) \right) \sqrt{a \sec^4(x)}
 \end{aligned}$$

input `Int[(a*Sec[x]^4)^(3/2),x]`

output `-(a*cos[x]^2*sqrt[a*sec[x]^4]*(-tan[x] - (2*tan[x]^3)/3 - tan[x]^5/5))`



## 3.63.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.63.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.51

method	result	size
default	$\frac{a\sqrt{a\sec(x)^4} (8\sin(x)\cos(x)+4\tan(x)+3\sec(x)^2\tan(x))}{15}$	31
risch	$\frac{16ia\sqrt{\frac{ae^{4ix}}{(e^{2ix}+1)^4}} (5+11\cos(2x)+9i\sin(2x))}{15(e^{2ix}+1)^3}$	47

input `int((a*sec(x)^4)^(3/2),x,method=_RETURNVERBOSE)`

output `1/15*a*(a*sec(x)^4)^(1/2)*(8*sin(x)*cos(x)+4*tan(x)+3*sec(x)^2*tan(x))`

**3.63.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.56

$$\int (a \sec^4(x))^{3/2} dx = \frac{(8a \cos(x)^4 + 4a \cos(x)^2 + 3a) \sqrt{\frac{a}{\cos(x)^4}} \sin(x)}{15 \cos(x)^3}$$

input `integrate((a*sec(x)^4)^(3/2),x, algorithm="fracas")`

output `1/15*(8*a*cos(x)^4 + 4*a*cos(x)^2 + 3*a)*sqrt(a/cos(x)^4)*sin(x)/cos(x)^3`

**3.63.6 Sympy [F]**

$$\int (a \sec^4(x))^{3/2} dx = \int (a \sec^4(x))^{\frac{3}{2}} dx$$

input `integrate((a*sec(x)**4)**(3/2),x)`

output `Integral((a*sec(x)**4)**(3/2), x)`

**3.63.7 Maxima [A] (verification not implemented)**

Time = 0.45 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.41

$$\int (a \sec^4(x))^{3/2} dx = \frac{1}{5} a^{\frac{3}{2}} \tan(x)^5 + \frac{2}{3} a^{\frac{3}{2}} \tan(x)^3 + a^{\frac{3}{2}} \tan(x)$$

input `integrate((a*sec(x)^4)^(3/2),x, algorithm="maxima")`

output `1/5*a^(3/2)*tan(x)^5 + 2/3*a^(3/2)*tan(x)^3 + a^(3/2)*tan(x)`

**3.63.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.36

$$\int (a \sec^4(x))^{3/2} dx = \frac{1}{15} (3 \tan(x)^5 + 10 \tan(x)^3 + 15 \tan(x)) a^{3/2}$$

input `integrate((a*sec(x)^4)^(3/2),x, algorithm="giac")`output `1/15*(3*tan(x)^5 + 10*tan(x)^3 + 15*tan(x))*a^(3/2)`**3.63.9 Mupad [B] (verification not implemented)**

Time = 0.56 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.59

$$\int (a \sec^4(x))^{3/2} dx = \frac{4 a^{3/2} \sin(x)}{5 \cos(x)^3} + \frac{a^{3/2} \sin(x)}{5 \cos(x)^5} - \frac{8 a^{3/2} \sin(x)^3}{15 \cos(x)^3}$$

input `int((a/cos(x)^4)^(3/2),x)`output `(4*a^(3/2)*sin(x))/(5*cos(x)^3) + (a^(3/2)*sin(x))/(5*cos(x)^5) - (8*a^(3/2)*sin(x)^3)/(15*cos(x)^3)`

## 3.64 $\int \sqrt{a \sec^4(x)} dx$

3.64.1	Optimal result . . . . .	447
3.64.2	Mathematica [A] (verified) . . . . .	447
3.64.3	Rubi [A] (verified) . . . . .	448
3.64.4	Maple [A] (verified) . . . . .	449
3.64.5	Fricas [A] (verification not implemented) . . . . .	450
3.64.6	Sympy [F] . . . . .	450
3.64.7	Maxima [A] (verification not implemented) . . . . .	450
3.64.8	Giac [A] (verification not implemented) . . . . .	451
3.64.9	Mupad [B] (verification not implemented) . . . . .	451

### 3.64.1 Optimal result

Integrand size = 10, antiderivative size = 15

$$\int \sqrt{a \sec^4(x)} dx = \cos(x) \sqrt{a \sec^4(x)} \sin(x)$$

output `cos(x)*sin(x)*(a*sec(x)^4)^(1/2)`

### 3.64.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \sqrt{a \sec^4(x)} dx = \cos(x) \sqrt{a \sec^4(x)} \sin(x)$$

input `Integrate[Sqrt[a*Sec[x]^4],x]`

output `Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x]`

### 3.64.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4611, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \sec^4(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \sec(x)^4} dx \\
 & \quad \downarrow \text{4611} \\
 & \cos^2(x) \sqrt{a \sec^4(x)} \int \sec^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \cos^2(x) \sqrt{a \sec^4(x)} \int \csc\left(x + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{4254} \\
 & -\cos^2(x) \sqrt{a \sec^4(x)} \int 1 d(-\tan(x)) \\
 & \quad \downarrow \text{24} \\
 & \sin(x) \cos(x) \sqrt{a \sec^4(x)}
 \end{aligned}$$

input `Int[Sqrt[a*Sec[x]^4],x]`

output `Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x]`

## 3.64.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.64.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$\cos(x) \sin(x) \sqrt{a \sec(x)^4}$	14
risch	$2i \sqrt{\frac{a e^{4ix}}{(e^{2ix}+1)^4}} (1 + e^{-2ix})$	29

input `int((a*sec(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `cos(x)*sin(x)*(a*sec(x)^4)^(1/2)`

**3.64.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \sqrt{a \sec^4(x)} dx = \sqrt{\frac{a}{\cos(x)^4}} \cos(x) \sin(x)$$

input `integrate((a*sec(x)^4)^(1/2),x, algorithm="fracas")`output `sqrt(a/cos(x)^4)*cos(x)*sin(x)`**3.64.6 Sympy [F]**

$$\int \sqrt{a \sec^4(x)} dx = \int \sqrt{a \sec^4(x)} dx$$

input `integrate((a*sec(x)**4)**(1/2),x)`output `Integral(sqrt(a*sec(x)**4), x)`**3.64.7 Maxima [A] (verification not implemented)**

Time = 0.48 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.40

$$\int \sqrt{a \sec^4(x)} dx = \sqrt{a} \tan(x)$$

input `integrate((a*sec(x)^4)^(1/2),x, algorithm="maxima")`output `sqrt(a)*tan(x)`

**3.64.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.40

$$\int \sqrt{a \sec^4(x)} dx = \sqrt{a} \tan(x)$$

input `integrate((a*sec(x)^4)^(1/2),x, algorithm="giac")`

output `sqrt(a)*tan(x)`

**3.64.9 Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.40

$$\int \sqrt{a \sec^4(x)} dx = \sqrt{a} \tan(x)$$

input `int((a/cos(x)^4)^(1/2),x)`

output `a^(1/2)*tan(x)`



$$3.65 \quad \int \frac{1}{\sqrt{a \sec^4(x)}} dx$$

3.65.1	Optimal result	452
3.65.2	Mathematica [A] (verified)	452
3.65.3	Rubi [A] (verified)	453
3.65.4	Maple [A] (verified)	454
3.65.5	Fricas [A] (verification not implemented)	455
3.65.6	Sympy [F]	455
3.65.7	Maxima [A] (verification not implemented)	455
3.65.8	Giac [A] (verification not implemented)	456
3.65.9	Mupad [F(-1)]	456

### 3.65.1 Optimal result

Integrand size = 10, antiderivative size = 36

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx = \frac{x \sec^2(x)}{2\sqrt{a \sec^4(x)}} + \frac{\tan(x)}{2\sqrt{a \sec^4(x)}}$$

output `1/2*x*sec(x)^2/(a*sec(x)^4)^(1/2)+1/2*tan(x)/(a*sec(x)^4)^(1/2)`

### 3.65.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx = \frac{x \sec^2(x) + \tan(x)}{2\sqrt{a \sec^4(x)}}$$

input `Integrate[1/Sqrt[a*Sec[x]^4],x]`

output `(x*Sec[x]^2 + Tan[x])/(2*sqrt[a*Sec[x]^4])`

**3.65.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.81, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4611, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \sec^4(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a \sec(x)^4}} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{\sec^2(x) \int \cos^2(x) dx}{\sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^2(x) \int \sin(x + \frac{\pi}{2})^2 dx}{\sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sec^2(x) \left( \int \frac{1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right)}{\sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sec^2(x) \left( \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right)}{\sqrt{a \sec^4(x)}}
 \end{aligned}$$

input `Int [1/Sqrt [a*Sec [x]^4] ,x]`

output `(Sec [x]^2*(x/2 + (Cos [x]*Sin [x])/2))/Sqrt [a*Sec [x]^4]`

## 3.65.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.65.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

method	result	size
default	$\frac{\tan(x)+x \sec(x)^2}{2\sqrt{a \sec(x)^4}}$	20
risch	$\frac{e^{2ix}x}{2\sqrt{\frac{ae^{4ix}}{(e^{2ix}+1)^4}}(e^{2ix}+1)^2} - \frac{ie^{4ix}}{8\sqrt{\frac{ae^{4ix}}{(e^{2ix}+1)^4}}(e^{2ix}+1)^2} + \frac{i}{8\sqrt{\frac{ae^{4ix}}{(e^{2ix}+1)^4}}(e^{2ix}+1)^2}$	102

input `int(1/(a*sec(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2/(a*sec(x)^4)^(1/2)*(tan(x)+x*sec(x)^2)`

**3.65.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx = \frac{(\cos(x)^3 \sin(x) + x \cos(x)^2) \sqrt{\frac{a}{\cos(x)^4}}}{2a}$$

input `integrate(1/(a*sec(x)^4)^(1/2),x, algorithm="fricas")`output `1/2*(cos(x)^3*sin(x) + x*cos(x)^2)*sqrt(a/cos(x)^4)/a`**3.65.6 Sympy [F]**

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx = \int \frac{1}{\sqrt{a \sec^4(x)}} dx$$

input `integrate(1/(a*sec(x)**4)**(1/2),x)`output `Integral(1/sqrt(a*sec(x)**4), x)`**3.65.7 Maxima [A] (verification not implemented)**

Time = 0.50 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.69

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx = \frac{x}{2\sqrt{a}} + \frac{\tan(x)}{2(\sqrt{a} \tan(x)^2 + \sqrt{a})}$$

input `integrate(1/(a*sec(x)^4)^(1/2),x, algorithm="maxima")`output `1/2*x/sqrt(a) + 1/2*tan(x)/(sqrt(a)*tan(x)^2 + sqrt(a))`

**3.65.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx = -\frac{1}{2} \sqrt{a} \left( \frac{\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor - x}{a} - \frac{\tan(x)}{(\tan(x)^2 + 1)a} \right)$$

input `integrate(1/(a*sec(x)^4)^(1/2),x, algorithm="giac")`

output `-1/2*sqrt(a)*((pi*floor(x/pi + 1/2) - x)/a - tan(x)/((tan(x)^2 + 1)*a))`

**3.65.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx = \int \frac{1}{\sqrt{\frac{a}{\cos(x)^4}}} dx$$

input `int(1/(a/cos(x)^4)^(1/2),x)`

output `int(1/(a/cos(x)^4)^(1/2), x)`

### 3.66 $\int \frac{1}{(a \sec^4(x))^{3/2}} dx$

3.66.1	Optimal result	457
3.66.2	Mathematica [A] (verified)	457
3.66.3	Rubi [A] (verified)	458
3.66.4	Maple [A] (verified)	460
3.66.5	Fricas [A] (verification not implemented)	460
3.66.6	Sympy [F]	460
3.66.7	Maxima [A] (verification not implemented)	461
3.66.8	Giac [F(-2)]	461
3.66.9	Mupad [F(-1)]	461

#### 3.66.1 Optimal result

Integrand size = 10, antiderivative size = 86

$$\int \frac{1}{(a \sec^4(x))^{3/2}} dx = \frac{5x \sec^2(x)}{16a \sqrt{a \sec^4(x)}} + \frac{5 \cos(x) \sin(x)}{24a \sqrt{a \sec^4(x)}} + \frac{\cos^3(x) \sin(x)}{6a \sqrt{a \sec^4(x)}} + \frac{5 \tan(x)}{16a \sqrt{a \sec^4(x)}}$$

output `5/16*x*sec(x)^2/a/(a*sec(x)^4)^(1/2)+5/24*cos(x)*sin(x)/a/(a*sec(x)^4)^(1/2)+1/6*cos(x)^3*sin(x)/a/(a*sec(x)^4)^(1/2)+5/16*tan(x)/a/(a*sec(x)^4)^(1/2)`

#### 3.66.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.44

$$\int \frac{1}{(a \sec^4(x))^{3/2}} dx = \frac{\sec^6(x)(60x + 45 \sin(2x) + 9 \sin(4x) + \sin(6x))}{192 (a \sec^4(x))^{3/2}}$$

input `Integrate[(a*Sec[x]^4)^(-3/2),x]`

output `(Sec[x]^6*(60*x + 45*Sin[2*x] + 9*Sin[4*x] + Sin[6*x]))/(192*(a*Sec[x]^4)^(3/2))`

**3.66.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.72, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {3042, 4611, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sec^4(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sec(x)^4)^{3/2}} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{\sec^2(x) \int \cos^6(x) dx}{a \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^2(x) \int \sin(x + \frac{\pi}{2})^6 dx}{a \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sec^2(x) (\frac{5}{6} \int \cos^4(x) dx + \frac{1}{6} \sin(x) \cos^5(x))}{a \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^2(x) (\frac{5}{6} \int \sin(x + \frac{\pi}{2})^4 dx + \frac{1}{6} \sin(x) \cos^5(x))}{a \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sec^2(x) (\frac{5}{6} (\frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x)) + \frac{1}{6} \sin(x) \cos^5(x))}{a \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^2(x) (\frac{5}{6} (\frac{3}{4} \int \sin(x + \frac{\pi}{2})^2 dx + \frac{1}{4} \sin(x) \cos^3(x)) + \frac{1}{6} \sin(x) \cos^5(x))}{a \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3115}
 \end{aligned}$$

$$\frac{\sec^2(x) \left( \frac{5}{6} \left( \frac{3}{4} \left( \frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \right)}{a \sqrt{a \sec^4(x)}}$$

↓ 24

$$\frac{\sec^2(x) \left( \frac{1}{6} \sin(x) \cos^5(x) + \frac{5}{6} \left( \frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left( \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) \right) \right)}{a \sqrt{a \sec^4(x)}}$$

input `Int[(a*Sec[x]^4)^(-3/2),x]`

output `(Sec[x]^2*((Cos[x]^5*Sin[x])/6 + (5*((Cos[x]^3*Sin[x])/4 + (3*(x/2 + (Cos[x]*Sin[x])/2))/4))/6))/(a*Sqrt[a*Sec[x]^4])`

### 3.66.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`



**3.66.4 Maple [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.47

method	result
default	$\frac{8 \cos(x)^3 \sin(x) + 10 \sin(x) \cos(x) + 15 \tan(x) + 15x \sec(x)^2}{48 \sqrt{a \sec(x)^4} a}$
risch	$\frac{5 e^{2ix} x}{16a(e^{2ix}+1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix}+1)^4}}} - \frac{i e^{8ix}}{384a(e^{2ix}+1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix}+1)^4}}} - \frac{3i e^{6ix}}{128a(e^{2ix}+1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix}+1)^4}}} - \frac{15i e^{4ix}}{128a(e^{2ix}+1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix}+1)^4}}} + \dots$

input `int(1/(a*sec(x)^4)^(3/2),x,method=_RETURNVERBOSE)`output `1/48/(a*sec(x)^4)^(1/2)/a*(8*cos(x)^3*sin(x)+10*sin(x)*cos(x)+15*tan(x)+15*x*sec(x)^2)`**3.66.5 Fricas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.50

$$\int \frac{1}{(a \sec^4(x))^{3/2}} dx = \frac{(15x \cos(x)^2 + (8 \cos(x)^7 + 10 \cos(x)^5 + 15 \cos(x)^3) \sin(x)) \sqrt{\frac{a}{\cos(x)^4}}}{48 a^2}$$

input `integrate(1/(a*sec(x)^4)^(3/2),x, algorithm="fricas")`output `1/48*(15*x*cos(x)^2 + (8*cos(x)^7 + 10*cos(x)^5 + 15*cos(x)^3)*sin(x))*sqrt(a/cos(x)^4)/a^2`**3.66.6 Sympy [F]**

$$\int \frac{1}{(a \sec^4(x))^{3/2}} dx = \int \frac{1}{(a \sec^4(x))^{\frac{3}{2}}} dx$$

input `integrate(1/(a*sec(x)**4)**(3/2),x)`output `Integral((a*sec(x)**4)**(-3/2), x)`

---

3.66.  $\int \frac{1}{(a \sec^4(x))^{3/2}} dx$

**3.66.7 Maxima [A] (verification not implemented)**

Time = 0.65 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.67

$$\int \frac{1}{(a \sec^4(x))^{3/2}} dx = \frac{15 \tan(x)^5 + 40 \tan(x)^3 + 33 \tan(x)}{48 \left( a^{3/2} \tan(x)^6 + 3 a^{3/2} \tan(x)^4 + 3 a^{3/2} \tan(x)^2 + a^{3/2} \right)} + \frac{5x}{16 a^{3/2}}$$

input `integrate(1/(a*sec(x)^4)^(3/2),x, algorithm="maxima")`

output `1/48*(15*tan(x)^5 + 40*tan(x)^3 + 33*tan(x))/(a^(3/2)*tan(x)^6 + 3*a^(3/2)*tan(x)^4 + 3*a^(3/2)*tan(x)^2 + a^(3/2)) + 5/16*x/a^(3/2)`

**3.66.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{(a \sec^4(x))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(a*sec(x)^4)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.66.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \sec^4(x))^{3/2}} dx = \int \frac{1}{\left( \frac{a}{\cos(x)^4} \right)^{3/2}} dx$$

input `int(1/(a/cos(x)^4)^(3/2),x)`

output `int(1/(a/cos(x)^4)^(3/2), x)`

### 3.67 $\int \frac{1}{(a \sec^4(x))^{5/2}} dx$

3.67.1	Optimal result . . . . .	462
3.67.2	Mathematica [A] (verified) . . . . .	462
3.67.3	Rubi [A] (verified) . . . . .	463
3.67.4	Maple [A] (verified) . . . . .	465
3.67.5	Fricas [A] (verification not implemented) . . . . .	466
3.67.6	Sympy [F] . . . . .	466
3.67.7	Maxima [A] (verification not implemented) . . . . .	466
3.67.8	Giac [F(-2)] . . . . .	467
3.67.9	Mupad [F(-1)] . . . . .	467

#### 3.67.1 Optimal result

Integrand size = 10, antiderivative size = 132

$$\int \frac{1}{(a \sec^4(x))^{5/2}} dx = \frac{63x \sec^2(x)}{256a^2 \sqrt{a \sec^4(x)}} + \frac{21 \cos(x) \sin(x)}{128a^2 \sqrt{a \sec^4(x)}} + \frac{21 \cos^3(x) \sin(x)}{160a^2 \sqrt{a \sec^4(x)}} + \frac{9 \cos^5(x) \sin(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{\cos^7(x) \sin(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{63 \tan(x)}{256a^2 \sqrt{a \sec^4(x)}}$$

output `63/256*x*sec(x)^2/a^2/(a*sec(x)^4)^(1/2)+21/128*cos(x)*sin(x)/a^2/(a*sec(x)^4)^(1/2)+21/160*cos(x)^3*sin(x)/a^2/(a*sec(x)^4)^(1/2)+9/80*cos(x)^5*sin(x)/a^2/(a*sec(x)^4)^(1/2)+1/10*cos(x)^7*sin(x)/a^2/(a*sec(x)^4)^(1/2)+63/256*tan(x)/a^2/(a*sec(x)^4)^(1/2)`

#### 3.67.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.42

$$\int \frac{1}{(a \sec^4(x))^{5/2}} dx = \frac{\cos^2(x) \sqrt{a \sec^4(x)} (2520x + 2100 \sin(2x) + 600 \sin(4x) + 150 \sin(6x) + 25 \sin(8x) + 2 \sin(10x))}{10240a^3}$$

input `Integrate[(a*Sec[x]^4)^(-5/2),x]`

output `(Cos[x]^2*Sqrt[a*Sec[x]^4]*(2520*x + 2100*Sin[2*x] + 600*Sin[4*x] + 150*Sin[6*x] + 25*Sin[8*x] + 2*Sin[10*x]))/(10240*a^3)`

**3.67.3 Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.70, number of steps used = 13, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.300$ , Rules used = {3042, 4611, 3042, 3115, 3042, 3115, 3042, 3115, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sec^4(x))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sec(x)^4)^{5/2}} dx \\
 & \quad \downarrow \text{4611} \\
 & \frac{\sec^2(x) \int \cos^{10}(x) dx}{a^2 \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^2(x) \int \sin(x + \frac{\pi}{2})^{10} dx}{a^2 \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sec^2(x) (\frac{9}{10} \int \cos^8(x) dx + \frac{1}{10} \sin(x) \cos^9(x))}{a^2 \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^2(x) (\frac{9}{10} \int \sin(x + \frac{\pi}{2})^8 dx + \frac{1}{10} \sin(x) \cos^9(x))}{a^2 \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sec^2(x) (\frac{9}{10} (\frac{7}{8} \int \cos^6(x) dx + \frac{1}{8} \sin(x) \cos^7(x)) + \frac{1}{10} \sin(x) \cos^9(x))}{a^2 \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^2(x) (\frac{9}{10} (\frac{7}{8} \int \sin(x + \frac{\pi}{2})^6 dx + \frac{1}{8} \sin(x) \cos^7(x)) + \frac{1}{10} \sin(x) \cos^9(x))}{a^2 \sqrt{a \sec^4(x)}} \\
 & \quad \downarrow \text{3115}
 \end{aligned}$$

$$\frac{\sec^2(x) \left( \frac{9}{10} \left( \frac{7}{8} \left( \frac{5}{6} \int \cos^4(x) dx + \frac{1}{6} \sin(x) \cos^5(x) \right) + \frac{1}{8} \sin(x) \cos^7(x) \right) + \frac{1}{10} \sin(x) \cos^9(x) \right)}{a^2 \sqrt{a \sec^4(x)}}$$

↓ 3042

$$\frac{\sec^2(x) \left( \frac{9}{10} \left( \frac{7}{8} \left( \frac{5}{6} \int \sin \left( x + \frac{\pi}{2} \right)^4 dx + \frac{1}{6} \sin(x) \cos^5(x) \right) + \frac{1}{8} \sin(x) \cos^7(x) \right) + \frac{1}{10} \sin(x) \cos^9(x) \right)}{a^2 \sqrt{a \sec^4(x)}}$$

↓ 3115

$$\frac{\sec^2(x) \left( \frac{9}{10} \left( \frac{7}{8} \left( \frac{5}{6} \left( \frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \right) + \frac{1}{8} \sin(x) \cos^7(x) \right) + \frac{1}{10} \sin(x) \cos^9(x) \right)}{a^2 \sqrt{a \sec^4(x)}}$$

↓ 3042

$$\frac{\sec^2(x) \left( \frac{9}{10} \left( \frac{7}{8} \left( \frac{5}{6} \left( \frac{3}{4} \int \sin \left( x + \frac{\pi}{2} \right)^2 dx + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \right) + \frac{1}{8} \sin(x) \cos^7(x) \right) + \frac{1}{10} \sin(x) \cos^9(x) \right)}{a^2 \sqrt{a \sec^4(x)}}$$

↓ 3115

$$\frac{\sec^2(x) \left( \frac{9}{10} \left( \frac{7}{8} \left( \frac{5}{6} \left( \frac{3}{4} \left( \frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \right) + \frac{1}{8} \sin(x) \cos^7(x) \right) + \frac{1}{10} \sin(x) \cos^9(x) \right)}{a^2 \sqrt{a \sec^4(x)}}$$

↓ 24

$$\frac{\sec^2(x) \left( \frac{1}{10} \sin(x) \cos^9(x) + \frac{9}{10} \left( \frac{1}{8} \sin(x) \cos^7(x) + \frac{7}{8} \left( \frac{1}{6} \sin(x) \cos^5(x) + \frac{5}{6} \left( \frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left( \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) \right) \right) \right)}{a^2 \sqrt{a \sec^4(x)}}$$

input `Int[(a*Sec[x]^4)^(-5/2),x]`

output `(Sec[x]^2*((Cos[x]^9*Sin[x])/10 + (9*((Cos[x]^7*Sin[x])/8 + (7*((Cos[x]^5*Sin[x])/6 + (5*((Cos[x]^3*Sin[x])/4 + (3*(x/2 + (Cos[x]*Sin[x])/2))/4))/6))/8))/10))/(a^2*Sqrt[a*Sec[x]^4])`

## 3.67.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.67.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.42

method	result
default	$\frac{128 \sin(x) \cos(x)^7 + 144 \sin(x) \cos(x)^5 + 168 \cos(x)^3 \sin(x) + 210 \sin(x) \cos(x) + 315 \tan(x) + 315x \sec(x)^2}{1280 \sqrt{a \sec(x)^4} a^2}$
risch	$\frac{63 e^{2ix} x}{256 a^2 (e^{2ix} + 1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix} + 1)^4}}} - \frac{ie^{12ix}}{10240 a^2 (e^{2ix} + 1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix} + 1)^4}}} - \frac{5ie^{10ix}}{4096 a^2 (e^{2ix} + 1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix} + 1)^4}}} - \frac{105ie^{4ix}}{1024 a^2 (e^{2ix} + 1)^2 \sqrt{\frac{a e^{4ix}}{(e^{2ix} + 1)^4}}}$

input `int(1/(a*sec(x)^4)^(5/2), x, method=_RETURNVERBOSE)`

output `1/1280/(a*sec(x)^4)^(1/2)/a^2*(128*sin(x)*cos(x)^7+144*sin(x)*cos(x)^5+168*cos(x)^3*sin(x)+210*sin(x)*cos(x)+315*tan(x)+315*x*sec(x)^2)`

**3.67.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.42

$$\int \frac{1}{(a \sec^4(x))^{5/2}} dx = \frac{(315 x \cos(x)^2 + (128 \cos(x)^{11} + 144 \cos(x)^9 + 168 \cos(x)^7 + 210 \cos(x)^5 + 315 \cos(x)^3) \sin(x) \sqrt{a/\cos(x)^4})}{1280 a^3}$$

input `integrate(1/(a*sec(x)^4)^(5/2),x, algorithm="fricas")`output `1/1280*(315*x*cos(x)^2 + (128*cos(x)^11 + 144*cos(x)^9 + 168*cos(x)^7 + 210*cos(x)^5 + 315*cos(x)^3)*sin(x))*sqrt(a/cos(x)^4)/a^3`**3.67.6 Sympy [F]**

$$\int \frac{1}{(a \sec^4(x))^{5/2}} dx = \int \frac{1}{(a \sec^4(x))^{\frac{5}{2}}} dx$$

input `integrate(1/(a*sec(x)**4)**(5/2),x)`output `Integral((a*sec(x)**4)**(-5/2), x)`**3.67.7 Maxima [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.67

$$\int \frac{1}{(a \sec^4(x))^{5/2}} dx = \frac{315 \tan(x)^9 + 1470 \tan(x)^7 + 2688 \tan(x)^5 + 2370 \tan(x)^3 + 965 \tan(x)}{1280 (a^{\frac{5}{2}} \tan(x)^{10} + 5 a^{\frac{5}{2}} \tan(x)^8 + 10 a^{\frac{5}{2}} \tan(x)^6 + 10 a^{\frac{5}{2}} \tan(x)^4 + 5 a^{\frac{5}{2}} \tan(x)^2 + a^{\frac{5}{2}})} + \frac{63 x}{256 a^{\frac{5}{2}}}$$

input `integrate(1/(a*sec(x)^4)^(5/2),x, algorithm="maxima")`output `1/1280*(315*tan(x)^9 + 1470*tan(x)^7 + 2688*tan(x)^5 + 2370*tan(x)^3 + 965*tan(x))/(a^(5/2)*tan(x)^10 + 5*a^(5/2)*tan(x)^8 + 10*a^(5/2)*tan(x)^6 + 10*a^(5/2)*tan(x)^4 + 5*a^(5/2)*tan(x)^2 + a^(5/2)) + 63/256*x/a^(5/2)`

---

3.67.  $\int \frac{1}{(a \sec^4(x))^{5/2}} dx$

**3.67.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{(a \sec^4(x))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(a*sec(x)^4)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.67.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \sec^4(x))^{5/2}} dx = \int \frac{1}{\left(\frac{a}{\cos(x)^4}\right)^{5/2}} dx$$

input `int(1/(a/cos(x)^4)^(5/2),x)`

output `int(1/(a/cos(x)^4)^(5/2), x)`



### 3.68 $\int ((b \sec(c + dx))^p)^n dx$

3.68.1	Optimal result	468
3.68.2	Mathematica [A] (verified)	468
3.68.3	Rubi [A] (verified)	469
3.68.4	Maple [F]	470
3.68.5	Fricas [F]	470
3.68.6	Sympy [F]	471
3.68.7	Maxima [F]	471
3.68.8	Giac [F]	471
3.68.9	Mupad [F(-1)]	472

#### 3.68.1 Optimal result

Integrand size = 12, antiderivative size = 81

$$\int ((b \sec(c + dx))^p)^n dx = \frac{\cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 - np), \frac{1}{2}(3 - np), \cos^2(c + dx)\right) ((b \sec(c + dx))^p)^n \sin(c + dx)}{d(1 - np) \sqrt{\sin^2(c + dx)}}$$

output `-cos(d*x+c)*hypergeom([1/2, -1/2*n*p+1/2], [-1/2*n*p+3/2], cos(d*x+c)^2)*((b*sec(d*x+c))^p)^n*sin(d*x+c)/d/(-n*p+1)/(sin(d*x+c)^2)^(1/2)`

#### 3.68.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.85

$$\int ((b \sec(c + dx))^p)^n dx = \frac{\cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{np}{2}, \frac{1}{2}(2 + np), \sec^2(c + dx)\right) ((b \sec(c + dx))^p)^n \sqrt{-\tan^2(c + dx)}}{dnp}$$

input `Integrate[((b*Sec[c + d*x])^p)^n,x]`

output `(Cot[c + d*x]*Hypergeometric2F1[1/2, (n*p)/2, (2 + n*p)/2, Sec[c + d*x]^2]*((b*Sec[c + d*x])^p)^n*sqrt[-Tan[c + d*x]^2])/(d*n*p)`

### 3.68.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4611, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int ((b \sec(c + dx))^p)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int ((b \sec(c + dx))^p)^n dx \\
 & \quad \downarrow \text{4611} \\
 & (b \sec(c + dx))^{-np} ((b \sec(c + dx))^p)^n \int (b \sec(c + dx))^{np} dx \\
 & \quad \downarrow \text{3042} \\
 & (b \sec(c + dx))^{-np} ((b \sec(c + dx))^p)^n \int \left( b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^{np} dx \\
 & \quad \downarrow \text{4259} \\
 & \left( \frac{\cos(c + dx)}{b} \right)^{np} ((b \sec(c + dx))^p)^n \int \left( \frac{\cos(c + dx)}{b} \right)^{-np} dx \\
 & \quad \downarrow \text{3042} \\
 & \left( \frac{\cos(c + dx)}{b} \right)^{np} ((b \sec(c + dx))^p)^n \int \left( \frac{\sin\left(c + dx + \frac{\pi}{2}\right)}{b} \right)^{-np} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{\sin(c + dx) \cos(c + dx) ((b \sec(c + dx))^p)^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 - np), \frac{1}{2}(3 - np), \cos^2(c + dx)\right)}{d(1 - np) \sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[((b*Sec[c + d*x])^p)^n,x]`

output `-((Cos[c + d*x]*Hypergeometric2F1[1/2, (1 - n*p)/2, (3 - n*p)/2, Cos[c + d*x]^2]*((b*Sec[c + d*x])^p)^n*Sin[c + d*x])/(d*(1 - n*p)*Sqrt[Sin[c + d*x]^2])`

## 3.68.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.68.4 Maple [F]

$$\int ((b \sec(dx + c))^p)^n dx$$

input `int(((b*sec(d*x+c))^p)^n,x)`

output `int(((b*sec(d*x+c))^p)^n,x)`

## 3.68.5 Fricas [F]

$$\int ((b \sec(c + dx))^p)^n dx = \int ((b \sec(dx + c))^p)^n dx$$

input `integrate(((b*sec(d*x+c))^p)^n,x, algorithm="fricas")`

output `integral(((b*sec(d*x + c))^p)^n, x)`

**3.68.6 Sympy [F]**

$$\int ((b \sec(c + dx))^p)^n dx = \int ((b \sec(c + dx))^p)^n dx$$

input `integrate(((b*sec(d*x+c))**p)**n,x)`

output `Integral(((b*sec(c + d*x))**p)**n, x)`

**3.68.7 Maxima [F]**

$$\int ((b \sec(c + dx))^p)^n dx = \int ((b \sec(dx + c))^p)^n dx$$

input `integrate(((b*sec(d*x+c))^p)^n,x, algorithm="maxima")`

output `integrate(((b*sec(d*x + c))^p)^n, x)`

**3.68.8 Giac [F]**

$$\int ((b \sec(c + dx))^p)^n dx = \int ((b \sec(dx + c))^p)^n dx$$

input `integrate(((b*sec(d*x+c))^p)^n,x, algorithm="giac")`

output `integrate(((b*sec(d*x + c))^p)^n, x)`

**3.68.9 Mupad [F(-1)]**

Timed out.

$$\int ((b \sec(c + dx))^p)^n dx = \int \left( \left( \frac{b}{\cos(c + dx)} \right)^p \right)^n dx$$

input `int(((b/cos(c + d*x))^p)^n,x)`output `int(((b/cos(c + d*x))^p)^n, x)`

### 3.69 $\int (a(b \sec(c + dx))^p)^n dx$

3.69.1	Optimal result . . . . .	473
3.69.2	Mathematica [A] (verified) . . . . .	473
3.69.3	Rubi [A] (verified) . . . . .	474
3.69.4	Maple [F] . . . . .	475
3.69.5	Fricas [F] . . . . .	475
3.69.6	Sympy [F] . . . . .	476
3.69.7	Maxima [F] . . . . .	476
3.69.8	Giac [F] . . . . .	476
3.69.9	Mupad [F(-1)] . . . . .	477

#### 3.69.1 Optimal result

Integrand size = 14, antiderivative size = 83

$$\int (a(b \sec(c + dx))^p)^n dx = \frac{\cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 - np), \frac{1}{2}(3 - np), \cos^2(c + dx)\right) (a(b \sec(c + dx))^p)^n \sin(c + dx)}{d(1 - np)\sqrt{\sin^2(c + dx)}}$$

output `-cos(d*x+c)*hypergeom([1/2, -1/2*n*p+1/2],[-1/2*n*p+3/2],cos(d*x+c)^2)*(a*(b*sec(d*x+c))^p)^n*sin(d*x+c)/d/(-n*p+1)/(sin(d*x+c)^2)^(1/2)`

#### 3.69.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.86

$$\int (a(b \sec(c + dx))^p)^n dx = \frac{\cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{np}{2}, \frac{1}{2}(2 + np), \sec^2(c + dx)\right) (a(b \sec(c + dx))^p)^n \sqrt{-\tan^2(c + dx)}}{dnp}$$

input `Integrate[(a*(b*Sec[c + d*x]))^p]^n,x]`

output `(Cot[c + d*x]*Hypergeometric2F1[1/2, (n*p)/2, (2 + n*p)/2, Sec[c + d*x]^2]*(a*(b*Sec[c + d*x]))^p)^n*sqrt[-Tan[c + d*x]^2]/(d*n*p)`

### 3.69.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 4611, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a(b \sec(c + dx))^p)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a(b \sec(c + dx))^p)^n dx \\
 & \quad \downarrow \text{4611} \\
 & (b \sec(c + dx))^{-np} (a(b \sec(c + dx))^p)^n \int (b \sec(c + dx))^{np} dx \\
 & \quad \downarrow \text{3042} \\
 & (b \sec(c + dx))^{-np} (a(b \sec(c + dx))^p)^n \int \left( b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^{np} dx \\
 & \quad \downarrow \text{4259} \\
 & \left( \frac{\cos(c + dx)}{b} \right)^{np} (a(b \sec(c + dx))^p)^n \int \left( \frac{\cos(c + dx)}{b} \right)^{-np} dx \\
 & \quad \downarrow \text{3042} \\
 & \left( \frac{\cos(c + dx)}{b} \right)^{np} (a(b \sec(c + dx))^p)^n \int \left( \frac{\sin\left(c + dx + \frac{\pi}{2}\right)}{b} \right)^{-np} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{\sin(c + dx) \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 - np), \frac{1}{2}(3 - np), \cos^2(c + dx)\right) (a(b \sec(c + dx))^p)^n}{d(1 - np) \sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[(a*(b*Sec[c + d*x])^p)^n,x]`

output `-((Cos[c + d*x]*Hypergeometric2F1[1/2, (1 - n*p)/2, (3 - n*p)/2, Cos[c + d*x]^2]*(a*(b*Sec[c + d*x])^p)^n*Sin[c + d*x])/(d*(1 - n*p)*Sqrt[Sin[c + d*x]^2])`

## 3.69.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4611 `Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p])) Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

## 3.69.4 Maple [F]

$$\int (a(b \sec(dx + c))^p)^n dx$$

input `int((a*(b*sec(d*x+c))^p)^n,x)`

output `int((a*(b*sec(d*x+c))^p)^n,x)`

## 3.69.5 Fricas [F]

$$\int (a(b \sec(c + dx))^p)^n dx = \int ((b \sec(dx + c))^p a)^n dx$$

input `integrate((a*(b*sec(d*x+c))^p)^n,x, algorithm="fricas")`

output `integral(((b*sec(d*x + c))^p*a)^n, x)`

---

3.69.  $\int (a(b \sec(c + dx))^p)^n dx$



**3.69.6 Sympy [F]**

$$\int (a(b \sec(c + dx))^p)^n dx = \int (a(b \sec(c + dx))^p)^n dx$$

input `integrate((a*(b*sec(d*x+c))**p)**n,x)`

output `Integral((a*(b*sec(c + d*x))**p)**n, x)`

**3.69.7 Maxima [F]**

$$\int (a(b \sec(c + dx))^p)^n dx = \int ((b \sec(dx + c))^p a)^n dx$$

input `integrate((a*(b*sec(d*x+c))^p)^n,x, algorithm="maxima")`

output `integrate(((b*sec(d*x + c))^p*a)^n, x)`

**3.69.8 Giac [F]**

$$\int (a(b \sec(c + dx))^p)^n dx = \int ((b \sec(dx + c))^p a)^n dx$$

input `integrate((a*(b*sec(d*x+c))^p)^n,x, algorithm="giac")`

output `integrate(((b*sec(d*x + c))^p*a)^n, x)`

**3.69.9 Mupad [F(-1)]**

Timed out.

$$\int (a(b \sec(c + dx))^p)^n dx = \int \left( a \left( \frac{b}{\cos(c + dx)} \right)^p \right)^n dx$$

input `int((a*(b/cos(c + d*x))^p)^n,x)`output `int((a*(b/cos(c + d*x))^p)^n, x)`

### 3.70 $\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx$

3.70.1	Optimal result . . . . .	478
3.70.2	Mathematica [A] (verified) . . . . .	478
3.70.3	Rubi [A] (verified) . . . . .	479
3.70.4	Maple [C] (verified) . . . . .	481
3.70.5	Fricas [C] (verification not implemented) . . . . .	481
3.70.6	Sympy [F] . . . . .	482
3.70.7	Maxima [F] . . . . .	482
3.70.8	Giac [F] . . . . .	482
3.70.9	Mupad [F(-1)] . . . . .	483

#### 3.70.1 Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{10 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21bd} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^3d}$$

output `10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^3/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.70.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.71

$$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sec^2(c + dx) \sqrt{b \sec(c + dx)} \left( 10 \cos^{\frac{5}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 5 \sin(2(c + dx)) + 6 \tan(c + dx) \right)}{21d}$$

input `Integrate[Sec[c + d*x]^4*Sqrt[b*Sec[c + d*x]],x]`

output  $(\text{Sec}[c + d*x]^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*(10*\text{Cos}[c + d*x]^{(5/2)}*\text{EllipticF}[(c + d*x)/2, 2] + 5*\text{Sin}[2*(c + d*x)] + 6*\text{Tan}[c + d*x]))/(21*d)$

### 3.70.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c + dx))^{9/2} dx}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{9/2} dx}{b^4} \\
 & \quad \downarrow \text{4255} \\
 & \quad \frac{\frac{5}{7} b^2 \int (b \sec(c + dx))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\frac{5}{7} b^2 \int (b \csc(c + dx + \frac{\pi}{2}))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^4} \\
 & \quad \downarrow \text{4255} \\
 & \quad \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^4} \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^4}$$

↓ 3042

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^4}$$

↓ 3120

$$\frac{\frac{5}{7}b^2 \left( \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^4}$$

input `Int[Sec[c + d*x]^4*Sqrt[b*Sec[c + d*x]],x]`

output `((2*b*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*d) + (5*b^2*((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d)))/7)/b^4`

### 3.70.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Simp[b^2*((n-2)/(n-1)) Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.70.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 14.25 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.63

method	result
default	$\frac{2i\sqrt{b}\sec(dx+c)\left(5\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(\cot(dx+c)-\csc(dx+c)),i)\cos(dx+c)+5\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\right)}{21d}$

input `int(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `2/21*I/d*(b*sec(d*x+c))^(1/2)*(5*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)+5*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)-5*I*tan(d*x+c)-3*I*tan(d*x+c)*sec(d*x+c)^2)`

### 3.70.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.18

$$\int \sec^4(c + dx)\sqrt{b\sec(c + dx)} dx = \frac{-5i\sqrt{2}\sqrt{b}\cos(dx+c)^3\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+5i\sqrt{2}\sqrt{b}\cos(dx+c)^3}{21d\cos(dx+c)}$$

input `integrate(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="fracas")`

output `1/21*(-5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(5*cos(d*x + c)^2 + 3)*sqrt(b/cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)`

**3.70.6 Sympy [F]**

$$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \sec^4(c + dx) dx$$

input `integrate(sec(d*x+c)**4*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*sec(c + d*x)**4, x)`

**3.70.7 Maxima [F]**

$$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^4 dx$$

input `integrate(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^4, x)`

**3.70.8 Giac [F]**

$$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^4 dx$$

input `integrate(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^4, x)`

**3.70.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c + dx)^4} dx$$

input `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^4,x)`output `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^4, x)`



### 3.71 $\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx$

3.71.1	Optimal result . . . . .	484
3.71.2	Mathematica [A] (verified) . . . . .	484
3.71.3	Rubi [A] (verified) . . . . .	485
3.71.4	Maple [C] (verified) . . . . .	487
3.71.5	Fricas [C] (verification not implemented) . . . . .	487
3.71.6	Sympy [F] . . . . .	488
3.71.7	Maxima [F] . . . . .	488
3.71.8	Giac [F] . . . . .	488
3.71.9	Mupad [F(-1)] . . . . .	489

#### 3.71.1 Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx = -\frac{6bE\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{6\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^2d}$$

output  $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/b^2/d-6/5*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

#### 3.71.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.73

$$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sec^2(c + dx) \sqrt{b \sec(c + dx)} \left( -12 \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 7 \sin(c + dx) + 3 \sin(3(c + dx)) \right)}{10d}$$

input `Integrate[Sec[c + d*x]^3*Sqrt[b*Sec[c + d*x]],x]`

output  $(\text{Sec}[c + d*x]^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*(-12*\text{Cos}[c + d*x]^{(5/2)}*\text{EllipticE}[(c + d*x)/2, 2] + 7*\text{Sin}[c + d*x] + 3*\text{Sin}[3*(c + d*x)]))/(10*d)$

### 3.71.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c + dx))^{7/2} dx}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{7/2} dx}{b^3} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \int (b \sec(c + dx))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \int (b \csc(c + dx + \frac{\pi}{2}))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^3} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx + \frac{\pi}{2})}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^3} \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

---

3.71.  $\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx$

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}}{b^3}$$

↓ 3042

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}}{b^3}$$

↓ 3119

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}}{b^3}$$

input `Int[Sec[c + d*x]^3*Sqrt[b*Sec[c + d*x]],x]`

output `((2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5)/b^3`

### 3.71.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n)*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

### 3.71.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.13 (sec) , antiderivative size = 412, normalized size of antiderivative = 4.34

method	result
default	$\frac{2\sqrt{b\sec(dx+c)} \left( 3i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)^2 - 3i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) \right)}{\dots}$

```
input int(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2), x, method=_RETURNVERBOSE)
```

```
output 2/5/d*(b*sec(d*x+c))^(1/2)/(cos(d*x+c)+1)*(3*I*EllipticF(I*(-cot(d*x+c)+cs
c(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*co
s(d*x+c)^2-3*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(
1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2+6*I*EllipticF(I*(-cot
(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1
))^(1/2)*cos(d*x+c)-6*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x
+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+3*I*(1/(cos(d*x
+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+c
sc(d*x+c)), I)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/
2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)+3*sin(d*x+c)+tan(d*x+c)+sec(d*x
+c)*tan(d*x+c))
```

### 3.71.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.26

$$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{-3i \sqrt{2} \sqrt{b} \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)))}{\dots}$$

input `integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*cos(d*x + c)^2 + 1)*sqrt(b/cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)`

### 3.71.6 Sympy [F]

$$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \sec^3(c + dx) dx$$

input `integrate(sec(d*x+c)**3*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*sec(c + d*x)**3, x)`

### 3.71.7 Maxima [F]

$$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^3 dx$$

input `integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^3, x)`

### 3.71.8 Giac [F]

$$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^3 dx$$

input `integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^3, x)`

**3.71.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx = \int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c + dx)^3} dx$$

input `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^3,x)`output `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^3, x)`

### 3.72 $\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx$

3.72.1	Optimal result . . . . .	490
3.72.2	Mathematica [A] (verified) . . . . .	490
3.72.3	Rubi [A] (verified) . . . . .	491
3.72.4	Maple [C] (verified) . . . . .	492
3.72.5	Fricas [C] (verification not implemented) . . . . .	493
3.72.6	Sympy [F] . . . . .	493
3.72.7	Maxima [F] . . . . .	494
3.72.8	Giac [F] . . . . .	494
3.72.9	Mupad [F(-1)] . . . . .	494

#### 3.72.1 Optimal result

Integrand size = 21, antiderivative size = 69

$$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3bd}$$

output `2/3*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b/d+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.72.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.74

$$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{2(b \sec(c + dx))^{3/2} \left( \cos^{3/2}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(c + dx) \right)}{3bd}$$

input `Integrate[Sec[c + d*x]^2*Sqrt[b*Sec[c + d*x]],x]`

output `(2*(b*Sec[c + d*x])^(3/2)*(Cos[c + d*x]^(3/2)*EllipticF[(c + d*x)/2, 2] + Sin[c + d*x]))/(3*b*d)`

**3.72.3 Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{5/2} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx + \frac{\pi}{2}))^{5/2} dx}{b^2} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{1}{3} b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3} b^2 \int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^2} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^2} \\
 & \quad \downarrow \text{3120} \\
 & \frac{\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^2}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2*Sqrt[b*Sec[c + d*x]],x]`

---

3.72.  $\int \sec^2(c+dx) \sqrt{b \sec(c+dx)} dx$



output 
$$\frac{((2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])}{(3*d) + (2*b*(b*\text{Sec}[c + d*x])^{3/2}*\text{Sin}[c + d*x])/(3*d)}/b^2$$

### 3.72.3.1 Defintions of rubi rules used

rule 2030 
$$\text{Int}[(\text{Fx}_.)*(v_.)^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x\_Symbol] \text{ :> } \text{Simp}[1/b^m \text{ Int}[(b*v)^{(m+n)*\text{Fx}, x}], x] \text{ ; FreeQ}\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$$

rule 3042 
$$\text{Int}[u_, x\_Symbol] \text{ :> } \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3120 
$$\text{Int}[1/\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_.)]], x\_Symbol] \text{ :> } \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] \text{ ; FreeQ}\{c, d\}, x]$$

rule 4255 
$$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x\_Symbol] \text{ :> } \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n-1)}/(d*(n-1))), x] + \text{Simp}[b^2*((n-2)/(n-1)) \text{ Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] \text{ ; FreeQ}\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$$

rule 4258 
$$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x\_Symbol] \text{ :> } \text{Simp}[(b*\text{Csc}[c + d*x])^{n-1}*\text{Sin}[c + d*x]^n \text{ Int}[1/\text{Sin}[c + d*x]^n, x], x] \text{ ; FreeQ}\{b, c, d\}, x] \ \&\& \ \text{EqQ}[n^2, 1/4]$$

### 3.72.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.76 (sec) , antiderivative size = 141, normalized size of antiderivative = 2.04

method	result
default	$-\frac{2\sqrt{b\sec(dx+c)} \left( i \text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) + i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \right)}{3d}$

input 
$$\text{int}(\sec(d*x+c)^2*(b*\sec(d*x+c))^{(1/2)}, x, \text{method}=\_RETURNVERBOSE)$$

output  $-2/3/d*(b*\sec(d*x+c))^{1/2}*(I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*EllipticF(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*\cos(d*x+c)+I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*EllipticF(I*(-\cot(d*x+c)+\csc(d*x+c)),I)-\tan(d*x+c))$

### 3.72.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.42

$$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2 \sqrt{b/\cos(dx + c)} \sin(dx + c)}{3 d \cos(dx + c)}$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output  $1/3*(-I*\sqrt{2}*\sqrt{b}*\cos(d*x + c)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + I*\sqrt{2}*\sqrt{b}*\cos(d*x + c)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + 2*\sqrt{b/\cos(d*x + c)}*\sin(d*x + c))/(d*\cos(d*x + c))$

### 3.72.6 Sympy [F]

$$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*sec(c + d*x)**2, x)`

**3.72.7 Maxima [F]**

$$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^2, x)`

**3.72.8 Giac [F]**

$$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^2, x)`

**3.72.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c + dx)^2} dx$$

input `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^2,x)`

output `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^2, x)`

### 3.73 $\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx$

3.73.1	Optimal result . . . . .	495
3.73.2	Mathematica [A] (verified) . . . . .	495
3.73.3	Rubi [A] (verified) . . . . .	496
3.73.4	Maple [C] (verified) . . . . .	497
3.73.5	Fricas [C] (verification not implemented) . . . . .	498
3.73.6	Sympy [F] . . . . .	499
3.73.7	Maxima [F] . . . . .	499
3.73.8	Giac [F] . . . . .	499
3.73.9	Mupad [F(-1)] . . . . .	500

#### 3.73.1 Optimal result

Integrand size = 19, antiderivative size = 63

$$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx = -\frac{2bE\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{d}$$

output `-2*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d`

#### 3.73.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.75

$$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{2\sqrt{b \sec(c + dx)} \left( -\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d}$$

input `Integrate[Sec[c + d*x]*Sqrt[b*Sec[c + d*x]],x]`

output `(2*Sqrt[b*Sec[c + d*x]]*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/d`

**3.73.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec(c+dx)\sqrt{b\sec(c+dx)} dx \\
 \downarrow \text{2030} \\
 \frac{\int (b\sec(c+dx))^{3/2} dx}{b} \\
 \downarrow \text{3042} \\
 \frac{\int (b\csc(c+dx+\frac{\pi}{2}))^{3/2} dx}{b} \\
 \downarrow \text{4255} \\
 \frac{\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b\sec(c+dx)}} dx}{b} \\
 \downarrow \text{3042} \\
 \frac{\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b\csc(c+dx+\frac{\pi}{2})}} dx}{b} \\
 \downarrow \text{4258} \\
 \frac{\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}}{b} \\
 \downarrow \text{3042} \\
 \frac{\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}}{b} \\
 \downarrow \text{3119} \\
 \frac{\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}}{b}
 \end{array}$$

input `Int[Sec[c + d*x]*Sqrt[b*Sec[c + d*x]],x]`

output `((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d)/b`

### 3.73.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] :> Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.73.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.80 (sec) , antiderivative size = 393, normalized size of antiderivative = 6.24

method	result
default	$-\frac{2 \left( i \operatorname{EllipticE} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)^2 - i \operatorname{EllipticF} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \right)}{\dots}$

---

3.73.  $\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx$

```
input int(sec(d*x+c)*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2/d*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(
cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2-I*EllipticF(I*(-cot(d*x+c)+c
sc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*c
os(d*x+c)^2+2*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))
^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-2*I*EllipticF(I*(-cot(
d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))
^(1/2)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(
1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-I*(1/(cos(d*x+c)+1))^(1/2)*(c
os(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-si
n(d*x+c))*(b*sec(d*x+c))^(1/2)/(cos(d*x+c)+1)
```

### 3.73.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.32

$$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 \sqrt{b} \sin(dx + c)}{d}$$

```
input integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output (-I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(
d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, wei
erstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b/cos(d*x
+ c))*sin(d*x + c))/d
```

**3.73.6 Sympy [F]**

$$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \sec(c + dx) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*sec(c + d*x), x)`

**3.73.7 Maxima [F]**

$$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c), x)`

**3.73.8 Giac [F]**

$$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c), x)`



**3.73.9 Mupad [F(-1)]**

Timed out.

$$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx = \int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c + dx)} dx$$

input `int((b/cos(c + d*x))^(1/2)/cos(c + d*x), x)`output `int((b/cos(c + d*x))^(1/2)/cos(c + d*x), x)`

### 3.74 $\int \sqrt{b \sec(c + dx)} dx$

3.74.1	Optimal result . . . . .	501
3.74.2	Mathematica [A] (verified) . . . . .	501
3.74.3	Rubi [A] (verified) . . . . .	502
3.74.4	Maple [C] (verified) . . . . .	503
3.74.5	Fricas [C] (verification not implemented) . . . . .	503
3.74.6	Sympy [F] . . . . .	504
3.74.7	Maxima [F] . . . . .	504
3.74.8	Giac [F] . . . . .	504
3.74.9	Mupad [B] (verification not implemented) . . . . .	505

#### 3.74.1 Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \sqrt{b \sec(c + dx)} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{d}$$

output `2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.74.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \sqrt{b \sec(c + dx)} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{d}$$

input `Integrate[Sqrt[b*Sec[c + d*x]],x]`

output `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d`

### 3.74.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{b \sec(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{b \csc\left(c + dx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4258} \\
 & \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3120} \\
 & \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{d}
 \end{aligned}$$

input `Int[Sqrt[b*Sec[c + d*x]],x]`

output `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d`

#### 3.74.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

### 3.74.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.83 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.03

method	result	size
default	$-\frac{2i(\cos(dx+c)+1)\operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{b\sec(dx+c)}\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}{d}$	77

```
input int((b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2*I/d*(cos(d*x+c)+1)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(b*sec(d*x+c)
)^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)
```

### 3.74.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.50

$$\int \sqrt{b \sec(c + dx)} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

```
input integrate((b*sec(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output (-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x +
c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*
x + c)))/d
```

**3.74.6 Sympy [F]**

$$\int \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} dx$$

input `integrate((b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x)), x)`

**3.74.7 Maxima [F]**

$$\int \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c)), x)`

**3.74.8 Giac [F]**

$$\int \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c)), x)`

**3.74.9 Mupad [B] (verification not implemented)**

Time = 12.99 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

$$\int \sqrt{b \sec(c + dx)} dx = \frac{2 \sqrt{\cos(c + dx)} \sqrt{\frac{b}{\cos(c + dx)}} F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

input `int((b/cos(c + d*x))^(1/2),x)`

output `(2*cos(c + d*x)^(1/2)*(b/cos(c + d*x))^(1/2)*ellipticF(c/2 + (d*x)/2, 2))/d`

### 3.75 $\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx$

3.75.1	Optimal result . . . . .	506
3.75.2	Mathematica [A] (verified) . . . . .	506
3.75.3	Rubi [A] (verified) . . . . .	507
3.75.4	Maple [C] (verified) . . . . .	508
3.75.5	Fricas [C] (verification not implemented) . . . . .	509
3.75.6	Sympy [F] . . . . .	509
3.75.7	Maxima [F] . . . . .	509
3.75.8	Giac [F] . . . . .	510
3.75.9	Mupad [F(-1)] . . . . .	510

#### 3.75.1 Optimal result

Integrand size = 19, antiderivative size = 39

$$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{2bE\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

output `2*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`

#### 3.75.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{2bE\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

input `Integrate[Cos[c + d*x]*Sqrt[b*Sec[c + d*x]],x]`

output `(2*b*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

**3.75.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx) \sqrt{b \sec(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{b \csc(c + dx + \frac{\pi}{2})}}{\csc(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \frac{1}{\sqrt{b \csc(\frac{1}{2}(2c + \pi) + dx)}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{b \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2bE(\frac{1}{2}(c + dx) | 2)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}
 \end{aligned}$$

input `Int[Cos[c + d*x]*Sqrt[b*Sec[c + d*x]],x]`

output `(2*b*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`



## 3.75.3.1 Defintions of rubi rules used

rule 2030  $\text{Int}[(F x_{.}) * (v_{.})^{(m_{.})} * ((b_{.}) * (v_{.}))^{(n_{.})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[1/b^{m_{.}} \text{Int}[(b * v)^{(m + n) * F x, x}], x] /;$   $\text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

rule 3042  $\text{Int}[u_{.}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3119  $\text{Int}[\text{Sqrt}[\sin[(c_{.}) + (d_{.}) * (x_{.})]], x_{\text{Symbol}}] \rightarrow \text{Simp}[(2/d) * \text{EllipticE}[(1/2) * (c - \text{Pi}/2 + d * x), 2], x] /;$   $\text{FreeQ}[\{c, d\}, x]$

rule 4258  $\text{Int}[(\text{csc}[(c_{.}) + (d_{.}) * (x_{.})] * (b_{.}))^{(n_{.})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(b * \text{Csc}[c + d * x])^{n_{.}} * \text{Sin}[c + d * x]^{n_{.}} \text{Int}[1/\text{Sin}[c + d * x]^{n_{.}}, x], x] /;$   $\text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{EqQ}[n^2, 1/4]$

## 3.75.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.03 (sec) , antiderivative size = 162, normalized size of antiderivative = 4.15

method	result
default	$\frac{2\sqrt{b\sec(dx+c)} \left( i(\cos(dx+c)+1)\sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \text{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) + i(-\cos(dx+c)-1)\sqrt{\frac{1}{\cos(dx+c)+1}} \right)}{d}$
risch	$-\frac{i(e^{2i(dx+c)}+1)\sqrt{2}\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} e^{-i(dx+c)}}{d} - i \left( -\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}}{\sqrt{2}} \sqrt{i(e^{i(dx+c)}-i)} \sqrt{i e^{i(dx+c)}} \right)$

input `int(cos(d*x+c)*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output  $2/d * (b * \sec(d * x + c))^{(1/2)} * (I * (\cos(d * x + c) + 1) * (1 / (\cos(d * x + c) + 1))^{(1/2)} * (\cos(d * x + c) / (\cos(d * x + c) + 1))^{(1/2)} * \text{EllipticE}(I * (-\cot(d * x + c) + \csc(d * x + c)), I) + I * (-\cos(d * x + c) - 1) * (1 / (\cos(d * x + c) + 1))^{(1/2)} * (\cos(d * x + c) / (\cos(d * x + c) + 1))^{(1/2)} * \text{EllipticF}(I * (-\cot(d * x + c) + \csc(d * x + c)), I) + (1 - \cos(d * x + c)) * \cot(d * x + c))$

**3.75.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.62

$$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{d}$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `(I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

**3.75.6 Sympy [F]**

$$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \cos(c + dx) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*cos(c + d*x), x)`

**3.75.7 Maxima [F]**

$$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c), x)`

**3.75.8 Giac [F]**

$$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c), x)`

**3.75.9 Mupad [F(-1)]**

Timed out.

$$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx = \int \cos(c + dx) \sqrt{\frac{b}{\cos(c + dx)}} dx$$

input `int(cos(c + d*x)*(b/cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)*(b/cos(c + d*x))^(1/2), x)`

### 3.76 $\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx$

3.76.1	Optimal result	511
3.76.2	Mathematica [A] (verified)	511
3.76.3	Rubi [A] (verified)	512
3.76.4	Maple [C] (verified)	513
3.76.5	Fricas [C] (verification not implemented)	514
3.76.6	Sympy [F]	514
3.76.7	Maxima [F]	515
3.76.8	Giac [F]	515
3.76.9	Mupad [F(-1)]	515

#### 3.76.1 Optimal result

Integrand size = 21, antiderivative size = 67

$$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}}$$

output `2/3*b*sin(d*x+c)/d/(b*sec(d*x+c))^(1/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.76.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.76

$$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} \left( 2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx)) \right)}{3d}$$

input `Integrate[Cos[c + d*x]^2*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[b*Sec[c + d*x]]*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*d)`

**3.76.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.13, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{b \csc(c+dx+\frac{\pi}{2})}}{\csc(c+dx+\frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^2 \left( \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b^2 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right) \\
 & \quad \downarrow \text{3120} \\
 & b^2 \left( \frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)
 \end{aligned}$$



input `int(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/3/d*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-cos(d*x+c)*sin(d*x+c))*(b*sec(d*x+c))^(1/2)`

### 3.76.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.25

$$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) - i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c)) + i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i \sin(dx+c))}{3d}$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/3*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)*sin(d*x + c) - I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

### 3.76.6 Sympy [F]

$$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \cos^2(c + dx) dx$$

input `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*cos(c + d*x)**2, x)`

**3.76.7 Maxima [F]**

$$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^2, x)`

**3.76.8 Giac [F]**

$$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^2, x)`

**3.76.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx = \int \cos(c + dx)^2 \sqrt{\frac{b}{\cos(c + dx)}} dx$$

input `int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2), x)`



### 3.77 $\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx$

3.77.1	Optimal result . . . . .	516
3.77.2	Mathematica [A] (verified) . . . . .	516
3.77.3	Rubi [A] (verified) . . . . .	517
3.77.4	Maple [C] (verified) . . . . .	518
3.77.5	Fricas [C] (verification not implemented) . . . . .	519
3.77.6	Sympy [F(-1)] . . . . .	519
3.77.7	Maxima [F] . . . . .	520
3.77.8	Giac [F] . . . . .	520
3.77.9	Mupad [F(-1)] . . . . .	520

#### 3.77.1 Optimal result

Integrand size = 21, antiderivative size = 70

$$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{6bE\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b^2 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}}$$

output  $2/5*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^(3/2)+6/5*b*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*EllipticE(\sin(1/2*d*x+1/2*c),2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)$

#### 3.77.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.81

$$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} \left( 12\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) + \sin(3(c + dx)) \right)}{10d}$$

input `Integrate[Cos[c + d*x]^3*Sqrt[b*Sec[c + d*x]],x]`

output  $(\text{Sqrt}[b*\text{Sec}[c + d*x]]*(12*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2] + \text{Sin}[c + d*x] + \text{Sin}[3*(c + d*x)]))/(10*d)$

**3.77.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{b \csc(c+dx+\frac{\pi}{2})}}{\csc(c+dx+\frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{2030} \\
 & b^3 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^3 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b^3 \left( \frac{3 \int \frac{\sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3119} \\
 & b^3 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)
 \end{aligned}$$

input `Int[Cos[c + d*x]^3*Sqrt[b*Sec[c + d*x]],x]`

output `b^3*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2)))`

### 3.77.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.77.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.16 (sec) , antiderivative size = 172, normalized size of antiderivative = 2.46

method	result
default	$-\frac{2\sqrt{b}\sec(dx+c)\left(i(-3\cos(dx+c)-3)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)+i(3\cos(dx+c)+3)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\right)}{5d}$

3.77.  $\int \cos^3(c + dx)\sqrt{b\sec(c + dx)} dx$

input `int(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/5/d*(b*sec(d*x+c))^(1/2)*(I*(-3*cos(d*x+c)-3)*(1/(cos(d*x+c)+1))^(1/2)*  
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+  
I*(3*cos(d*x+c)+3)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1  
/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+(cos(d*x+c)^3+2*cos(d*x+c)-3)*  
cot(d*x+c))`

### 3.77.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.31

$$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^2 \sin(dx+c) + 3i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c)))}{d}$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/5*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^2*sin(d*x + c) + 3*I*sqrt(2)*sqrt  
(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin  
(d*x + c))) - 3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInver  
se(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

### 3.77.6 Sympy [F(-1)]

Timed out.

$$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(b*sec(d*x+c))**(1/2),x)`

output `Timed out`

**3.77.7 Maxima [F]**

$$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^3, x)`

**3.77.8 Giac [F]**

$$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^3, x)`

**3.77.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx = \int \cos(c + dx)^3 \sqrt{\frac{b}{\cos(c + dx)}} dx$$

input `int(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2), x)`

### 3.78 $\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx$

3.78.1	Optimal result	521
3.78.2	Mathematica [A] (verified)	521
3.78.3	Rubi [A] (verified)	522
3.78.4	Maple [C] (verified)	524
3.78.5	Fricas [C] (verification not implemented)	524
3.78.6	Sympy [F]	525
3.78.7	Maxima [F]	525
3.78.8	Giac [F]	525
3.78.9	Mupad [F(-1)]	526

#### 3.78.1 Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{10 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{2b^3 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b \sin(c + dx)}{21d \sqrt{b \sec(c + dx)}}$$

output `2/7*b^3*sin(d*x+c)/d/(b*sec(d*x+c))^(5/2)+10/21*b*sin(d*x+c)/d/(b*sec(d*x+c))^(1/2)+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.78.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.66

$$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} \left( 40 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) \right)}{84d}$$

input `Integrate[Cos[c + d*x]^4*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*d)`

**3.78.3 Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{b \csc(c+dx+\frac{\pi}{2})}}{\csc(c+dx+\frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{2030} \\
 & b^4 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^4 \left( \frac{5 \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{5 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^4 \left( \frac{5 \left( \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{5 \left( \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$b^4 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd (b \sec(c+dx))^{5/2}} \right)$$

↓ 3042

$$b^4 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd (b \sec(c+dx))^{5/2}} \right)$$

↓ 3120

$$b^4 \left( \frac{5 \left( \frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd (b \sec(c+dx))^{5/2}} \right)$$

input `Int[Cos[c + d*x]^4*Sqrt[b*Sec[c + d*x]],x]`

output `b^4*((2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/(7*b^2))`

### 3.78.3.1 Defintions of rubi rules used

rule 2030 `Int[(F*x_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F*x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`



rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.78.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.16 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.72

method	result
default	$-\frac{2 \left( 5i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) + 5i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF} \right)}{21d}$

input `int(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2), x, method=_RETURNVERBOSE)`

output `-2/21/d*(5*I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)-3*cos(d*x+c)^3*sin(d*x+c)-5*cos(d*x+c)*sin(d*x+c))*(b*sec(d*x+c))^(1/2)`

### 3.78.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.02

$$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx$$


---


$$= \frac{2 \left( 3 \cos(dx + c)^3 + 5 \cos(dx + c) \right) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx + c) - 5i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c))}{21d}$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/21*(2*(3*cos(d*x + c)^3 + 5*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c) - 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

### 3.78.6 Sympy [F]

$$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \cos^4(c + dx) dx$$

input `integrate(cos(d*x+c)**4*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*cos(c + d*x)**4, x)`

### 3.78.7 Maxima [F]

$$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^4 dx$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^4, x)`

### 3.78.8 Giac [F]

$$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^4 dx$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^4, x)`

**3.78.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx = \int \cos(c + dx)^4 \sqrt{\frac{b}{\cos(c + dx)}} dx$$

input `int(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2),x)`output `int(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2), x)`

### 3.79 $\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx$

3.79.1	Optimal result	527
3.79.2	Mathematica [A] (verified)	527
3.79.3	Rubi [A] (verified)	528
3.79.4	Maple [C] (verified)	530
3.79.5	Fricas [C] (verification not implemented)	530
3.79.6	Sympy [F(-1)]	531
3.79.7	Maxima [F]	531
3.79.8	Giac [F]	531
3.79.9	Mupad [F(-1)]	532

#### 3.79.1 Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{14bE\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b^4 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}$$

output  $2/9*b^4*\sin(d*x+c)/d/(b*\sec(d*x+c))^(7/2)+14/45*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^(3/2)+14/15*b*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*EllipticE(\sin(1/2*d*x+1/2*c),2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)$

#### 3.79.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.72

$$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} \left( 84 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \cos^2(c + dx) (33 \sin(c + dx) + 5 \sin(3(c + dx))) \right)}{90d}$$

input `Integrate[Cos[c + d*x]^5*Sqrt[b*Sec[c + d*x]],x]`

output  $(\text{Sqrt}[b*\text{Sec}[c + d*x]]*(84*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2] + \text{Cos}[c + d*x]^2*(33*\text{Sin}[c + d*x] + 5*\text{Sin}[3*(c + d*x)])))/(90*d)$

**3.79.3 Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^5(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{b \csc(c+dx+\frac{\pi}{2})}}{\csc(c+dx+\frac{\pi}{2})^5} dx \\
 & \quad \downarrow \text{2030} \\
 & b^5 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{9/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^5 \left( \frac{7 \int \frac{1}{(b \sec(c+dx))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^5 \left( \frac{7 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^5 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^5 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 4258 \\
 b^5 \left( \frac{7 \left( \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 \downarrow 3042 \\
 b^5 \left( \frac{7 \left( \frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 \downarrow 3119 \\
 b^5 \left( \frac{7 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{array}$$

input `Int[Cos[c + d*x]^5*Sqrt[b*Sec[c + d*x]],x]`

output `b^5*((2*Sin[c + d*x])/(9*b*d*(b*Sec[c + d*x])^(7/2)) + (7*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))))/(9*b^2)`

### 3.79.3.1 Defintions of rubi rules used

rule 2030 `Int[(F*_)(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F_x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.79.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.66 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.88

method	result
default	$-\frac{2\sqrt{b\sec(dx+c)}\left(i(-21\cos(dx+c)-21)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)+i(21\cos(dx+c)+21)\right)}{\dots}$

input `int(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/45/d*(b*sec(d*x+c))^(1/2)*(I*(-21*cos(d*x+c)-21)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(21*cos(d*x+c)+21)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+(5*cos(d*x+c)^5+2*cos(d*x+c)^3+14*cos(d*x+c)-21)*cot(d*x+c)`

### 3.79.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.07

$$\int \cos^5(c + dx)\sqrt{b\sec(c + dx)} dx$$

$$= \frac{2\left(5\cos(dx+c)^4 + 7\cos(dx+c)^2\right)\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c) + 21i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassZeta}(\dots))}{\dots}$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/45*(2*(5*cos(d*x + c)^4 + 7*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c) + 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

### 3.79.6 Sympy [F(-1)]

Timed out.

$$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**5*(b*sec(d*x+c))**(1/2),x)`

output `Timed out`

### 3.79.7 Maxima [F]

$$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^5 dx$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^5, x)`

### 3.79.8 Giac [F]

$$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \cos(dx + c)^5 dx$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^5, x)`



**3.79.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx = \int \cos(c + dx)^5 \sqrt{\frac{b}{\cos(c + dx)}} dx$$

input `int(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2),x)`output `int(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2), x)`

### 3.80 $\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx$

3.80.1	Optimal result . . . . .	533
3.80.2	Mathematica [A] (verified) . . . . .	533
3.80.3	Rubi [A] (verified) . . . . .	534
3.80.4	Maple [C] (verified) . . . . .	536
3.80.5	Fricas [C] (verification not implemented) . . . . .	536
3.80.6	Sympy [F] . . . . .	537
3.80.7	Maxima [F] . . . . .	537
3.80.8	Giac [F] . . . . .	537
3.80.9	Mupad [F(-1)] . . . . .	538

#### 3.80.1 Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{10b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^2d}$$

```
output 10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^2/d+10/21*b*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d
```

#### 3.80.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.67

$$\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{(b \sec(c + dx))^{5/2} \left( 10 \cos^{\frac{5}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 5 \sin(2(c + dx)) + 6 \tan(c + dx) \right)}{21bd}$$

```
input Integrate[Sec[c + d*x]^3*(b*Sec[c + d*x])^(3/2),x]
```

output  $((b*\text{Sec}[c + d*x])^{5/2}*(10*\text{Cos}[c + d*x]^{5/2}*\text{EllipticF}[(c + d*x)/2, 2] + 5*\text{Sin}[2*(c + d*x)] + 6*\text{Tan}[c + d*x]))/(21*b*d)$

### 3.80.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c + dx))^{9/2} dx}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{9/2} dx}{b^3} \\
 & \quad \downarrow \text{4255} \\
 & \quad \frac{\frac{5}{7}b^2 \int (b \sec(c + dx))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\frac{5}{7}b^2 \int (b \csc(c + dx + \frac{\pi}{2}))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^3} \\
 & \quad \downarrow \text{4255} \\
 & \quad \frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^3} \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^3}$$

↓ 3042

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^3}$$

↓ 3120

$$\frac{\frac{5}{7}b^2 \left( \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^3}$$

input `Int[Sec[c + d*x]^3*(b*Sec[c + d*x])^(3/2),x]`

output `((2*b*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*d) + (5*b^2*((2*b^2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d)))/7)/b^3`

### 3.80.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Simp[b^2*((n-2)/(n-1)) Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.80.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.67

method	result
default	$-\frac{2ib\sqrt{b}\sec(dx+c)}{21d} \left( 5\sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \cos(dx+c) + 5\sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \right)$

input `int(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2), x, method=_RETURNVERBOSE)`

output 
$$-2/21*I/d*b*(b*\sec(d*x+c))^{(1/2)}*(5*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)), I)*\cos(d*x+c)+5*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)), I)+5*I*\tan(d*x+c)+3*I*\tan(d*x+c)*\sec(d*x+c)^2)$$

### 3.80.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.23

$$\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{-5i\sqrt{2}b^{3/2}\cos(dx+c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + 5i\sqrt{2}b^{3/2}\cos(dx+c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c)) + 2*(5*b*\cos(d*x + c)^2 + 3*b)*\sqrt{b/\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c)^3)}{21d}$$

input `integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")`

output 
$$1/21*(-5*I*\sqrt{2}*b^{(3/2)}*\cos(d*x + c)^3*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + 5*I*\sqrt{2}*b^{(3/2)}*\cos(d*x + c)^3*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + 2*(5*b*\cos(d*x + c)^2 + 3*b)*\sqrt{b/\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c)^3)$$

---

3.80.  $\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx$

**3.80.6 Sympy [F]**

$$\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(c + dx))^{\frac{3}{2}} \sec^3(c + dx) dx$$

input `integrate(sec(d*x+c)**3*(b*sec(d*x+c))**(3/2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*sec(c + d*x)**3, x)`

**3.80.7 Maxima [F]**

$$\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^3 dx$$

input `integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^3, x)`

**3.80.8 Giac [F]**

$$\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^3 dx$$

input `integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^3, x)`

**3.80.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\cos(c + dx)^3} dx$$

input `int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^3,x)`output `int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^3, x)`

### 3.81 $\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx$

3.81.1	Optimal result . . . . .	539
3.81.2	Mathematica [A] (verified) . . . . .	539
3.81.3	Rubi [A] (verified) . . . . .	540
3.81.4	Maple [C] (verified) . . . . .	542
3.81.5	Fricas [C] (verification not implemented) . . . . .	542
3.81.6	Sympy [F] . . . . .	543
3.81.7	Maxima [F] . . . . .	543
3.81.8	Giac [F] . . . . .	543
3.81.9	Mupad [F(-1)] . . . . .	544

#### 3.81.1 Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx = -\frac{6b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{6b \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5bd}$$

output `2/5*(b*sec(d*x+c))^(5/2)*sin(d*x+c)/b/d-6/5*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+6/5*b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d`

#### 3.81.2 Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

$$\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{(b \sec(c + dx))^{5/2} \left( -12 \cos^{5/2}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 7 \sin(c + dx) + 3 \sin(3(c + dx)) \right)}{10bd}$$

input `Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(3/2),x]`

output `((b*Sec[c + d*x])^(5/2)*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*b*d)`



### 3.81.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec^2(c+dx)(b \sec(c+dx))^{3/2} dx \\
 \downarrow \text{2030} \\
 \frac{\int (b \sec(c+dx))^{7/2} dx}{b^2} \\
 \downarrow \text{3042} \\
 \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{7/2} dx}{b^2} \\
 \downarrow \text{4255} \\
 \frac{\frac{3}{5}b^2 \int (b \sec(c+dx))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^2} \\
 \downarrow \text{3042} \\
 \frac{\frac{3}{5}b^2 \int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^2} \\
 \downarrow \text{4255} \\
 \frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^2} \\
 \downarrow \text{3042} \\
 \frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^2} \\
 \downarrow \text{4258} \\
 \frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^2} \\
 \downarrow \text{3042}
 \end{array}$$

---

3.81.  $\int \sec^2(c+dx)(b \sec(c+dx))^{3/2} dx$

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^2}$$

↓ 3119

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^2}$$

input `Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(3/2),x]`

output `((2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5)/b^2`

### 3.81.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b.)*(v.))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

---

3.81.  $\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx$

### 3.81.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 413, normalized size of antiderivative = 4.21

method	result
default	$\frac{2b\sqrt{b\sec(dx+c)} \left( 3i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)^2 - 3i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) \right)}{\dots}$

input `int(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2), x, method=_RETURNVERBOSE)`

output 
$$\frac{2}{5} \frac{d \cdot b \cdot (b \cdot \sec(dx+c))^{1/2}}{(\cos(dx+c)+1)} \cdot \left( 3i \operatorname{EllipticF}(I \cdot (-\cot(dx+c)+\csc(dx+c)), I) \cdot \frac{1}{(\cos(dx+c)+1)^{1/2}} \cdot \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cdot \cos(dx+c)^2 - 3i \operatorname{EllipticE}(I \cdot (-\cot(dx+c)+\csc(dx+c)), I) \cdot \frac{1}{(\cos(dx+c)+1)^{1/2}} \cdot \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cdot \cos(dx+c) \right) + 6i \operatorname{EllipticF}(I \cdot (-\cot(dx+c)+\csc(dx+c)), I) \cdot \frac{1}{(\cos(dx+c)+1)^{1/2}} \cdot \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cdot \cos(dx+c) - 6i \operatorname{EllipticE}(I \cdot (-\cot(dx+c)+\csc(dx+c)), I) \cdot \frac{1}{(\cos(dx+c)+1)^{1/2}} \cdot \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cdot \cos(dx+c) + 3i \cdot \frac{1}{(\cos(dx+c)+1)^{1/2}} \cdot \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cdot \operatorname{EllipticF}(I \cdot (-\cot(dx+c)+\csc(dx+c)), I) - 3i \cdot \frac{1}{(\cos(dx+c)+1)^{1/2}} \cdot \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cdot \operatorname{EllipticE}(I \cdot (-\cot(dx+c)+\csc(dx+c)), I) + 3 \sin(dx+c) + \tan(dx+c) + \sec(dx+c) \cdot \tan(dx+c)$$

### 3.81.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.23

$$\int \sec^2(c+dx) (b \sec(c+dx))^{3/2} dx = \frac{-3i \sqrt{2} b^{3/2} \cos(dx+c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c)))^{3/2}}{\dots}$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")`

output `1/5*(-3*I*sqrt(2)*b^(3/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*b^(3/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*b*cos(d*x + c)^2 + b)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^2)`

### 3.81.6 Sympy [F]

$$\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(c + dx))^{\frac{3}{2}} \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(3/2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*sec(c + d*x)**2, x)`

### 3.81.7 Maxima [F]

$$\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^2, x)`

### 3.81.8 Giac [F]

$$\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^2, x)`

**3.81.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\cos(c + dx)^2} dx$$

input `int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^2,x)`output `int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^2, x)`

### 3.82 $\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx$

3.82.1	Optimal result . . . . .	545
3.82.2	Mathematica [A] (verified) . . . . .	545
3.82.3	Rubi [A] (verified) . . . . .	546
3.82.4	Maple [C] (verified) . . . . .	548
3.82.5	Fricas [C] (verification not implemented) . . . . .	548
3.82.6	Sympy [F] . . . . .	549
3.82.7	Maxima [F] . . . . .	549
3.82.8	Giac [F] . . . . .	549
3.82.9	Mupad [F(-1)] . . . . .	550

#### 3.82.1 Optimal result

Integrand size = 19, antiderivative size = 67

$$\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d}$$

output

```
2/3*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/d+2/3*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d
```

#### 3.82.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.73

$$\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b\sqrt{b \sec(c + dx)} \left( \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx) \right)}{3d}$$

input

```
Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(3/2),x]
```

output  $(2*b*\text{Sqrt}[b*\text{Sec}[c + d*x]]*(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2] + \text{Tan}[c + d*x]))/(3*d)$

### 3.82.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.10, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c + dx)(b \sec(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c + dx))^{5/2} dx}{b} \\
 & \quad \quad \downarrow \text{3042} \\
 & \quad \quad \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{5/2} dx}{b} \\
 & \quad \quad \quad \downarrow \text{4255} \\
 & \quad \quad \quad \frac{\frac{1}{3}b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d}}{b} \\
 & \quad \quad \quad \quad \downarrow \text{3042} \\
 & \quad \quad \quad \quad \frac{\frac{1}{3}b^2 \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d}}{b} \\
 & \quad \quad \quad \quad \quad \downarrow \text{4258} \\
 & \quad \quad \quad \quad \quad \frac{\frac{1}{3}b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d}}{b} \\
 & \quad \quad \quad \quad \quad \quad \downarrow \text{3042} \\
 & \quad \quad \quad \quad \quad \quad \frac{\frac{1}{3}b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d}}{b} \\
 & \quad \quad \quad \quad \quad \quad \quad \downarrow \text{3120}
 \end{aligned}$$

$$\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d}$$

$b$

input `Int[Sec[c + d*x]*(b*Sec[c + d*x])^(3/2), x]`

output `((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d))/b`

### 3.82.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx._)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`



### 3.82.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.09

method	result
default	$\frac{2b\sqrt{b\sec(dx+c)} \left( -i \operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)), i\right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \right)}{3d}$

input `int(sec(d*x+c)*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2/3/d*b*(b*sec(d*x+c))^(1/2)*(-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I))+tan(d*x+c)`

### 3.82.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.48

$$\int \sec(c+dx)(b\sec(c+dx))^{3/2} dx = \frac{-i\sqrt{2}b^{3/2}\cos(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+i\sqrt{2}b^{3/2}\cos(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))+2*b*\sqrt{b/\cos(dx+c)}*\sin(dx+c)}{3d\cos(dx+c)}$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="fracas")`

output `1/3*(-I*sqrt(2)*b^(3/2)*cos(d*x+c)*weierstrassPInverse(-4,0,cos(d*x+c)+I*sin(d*x+c))+I*sqrt(2)*b^(3/2)*cos(d*x+c)*weierstrassPInverse(-4,0,cos(d*x+c)-I*sin(d*x+c))+2*b*sqrt(b/cos(d*x+c))*sin(d*x+c))/(d*cos(d*x+c))`

**3.82.6 Sympy [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(c + dx))^{\frac{3}{2}} \sec(c + dx) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))**(3/2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*sec(c + d*x), x)`

**3.82.7 Maxima [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c), x)`

**3.82.8 Giac [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c), x)`

**3.82.9 Mupad [F(-1)]**

Timed out.

$$\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\cos(c + dx)} dx$$

input `int((b/cos(c + d*x))^(3/2)/cos(c + d*x),x)`output `int((b/cos(c + d*x))^(3/2)/cos(c + d*x), x)`

### 3.83 $\int (b \sec(c + dx))^{3/2} dx$

3.83.1	Optimal result . . . . .	551
3.83.2	Mathematica [A] (verified) . . . . .	551
3.83.3	Rubi [A] (verified) . . . . .	552
3.83.4	Maple [C] (verified) . . . . .	553
3.83.5	Fricas [C] (verification not implemented) . . . . .	554
3.83.6	Sympy [F] . . . . .	554
3.83.7	Maxima [F] . . . . .	555
3.83.8	Giac [F] . . . . .	555
3.83.9	Mupad [F(-1)] . . . . .	555

#### 3.83.1 Optimal result

Integrand size = 12, antiderivative size = 66

$$\int (b \sec(c + dx))^{3/2} dx = -\frac{2b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b \sqrt{b \sec(c + dx)} \sin(c + dx)}{d}$$

output `-2*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d`

#### 3.83.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.73

$$\int (b \sec(c + dx))^{3/2} dx = \frac{2b \sqrt{b \sec(c + dx)} \left( -\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d}$$

input `Integrate[(b*Sec[c + d*x])^(3/2),x]`

output `(2*b*sqrt[b*Sec[c + d*x]]*(-(sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/d`

### 3.83.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( b \csc \left( c + dx + \frac{\pi}{2} \right) \right)^{3/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(3/2),x]`

output `(-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d`

## 3.83.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.83.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 394, normalized size of antiderivative = 5.97

method	result
default	$-\frac{2 \left( i \operatorname{EllipticE} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)^2 - i \operatorname{EllipticF} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \right)}{\dots}$

input `int((b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/d*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2+2*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-2*I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-sin(d*x+c)*(b*sec(d*x+c))^(1/2)*b/(cos(d*x+c)+1)`

### 3.83.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.27

$$\int (b \sec(c + dx))^{3/2} dx = \frac{-i \sqrt{2} b^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{2 \sqrt{2} b^{3/2}}$$

input `integrate((b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*b*sqrt(b/cos(d*x + c))*sin(d*x + c)/d`

### 3.83.6 Sympy [F]

$$\int (b \sec(c + dx))^{3/2} dx = \int (b \sec(c + dx))^{\frac{3}{2}} dx$$

input `integrate((b*sec(d*x+c))**(3/2),x)`

output `Integral((b*sec(c + d*x))**(3/2), x)`

**3.83.7 Maxima [F]**

$$\int (b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} dx$$

input `integrate((b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2), x)`

**3.83.8 Giac [F]**

$$\int (b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} dx$$

input `integrate((b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2), x)`

**3.83.9 Mupad [F(-1)]**

Timed out.

$$\int (b \sec(c + dx))^{3/2} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int((b/cos(c + d*x))^(3/2),x)`

output `int((b/cos(c + d*x))^(3/2), x)`



### 3.84 $\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx$

3.84.1	Optimal result . . . . .	556
3.84.2	Mathematica [A] (verified) . . . . .	556
3.84.3	Rubi [A] (verified) . . . . .	557
3.84.4	Maple [C] (verified) . . . . .	558
3.84.5	Fricas [C] (verification not implemented) . . . . .	559
3.84.6	Sympy [F] . . . . .	559
3.84.7	Maxima [F] . . . . .	559
3.84.8	Giac [F] . . . . .	560
3.84.9	Mupad [F(-1)] . . . . .	560

#### 3.84.1 Optimal result

Integrand size = 19, antiderivative size = 39

$$\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{d}$$

output `2*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.84.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{d}$$

input `Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(3/2),x]`

output `(2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d`

### 3.84.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c+dx)(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}}{\csc(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \sqrt{b \csc\left(\frac{1}{2}(2c+\pi)+dx\right)} dx \\
 & \quad \downarrow \text{4258} \\
 & b\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & b\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3120} \\
 & \frac{2b\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]*(b*Sec[c + d*x])^(3/2), x]`

output `(2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d`

## 3.84.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.84.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.00

method	result	size
default	$-\frac{2ib(\cos(dx+c)+1)\operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{b\sec(dx+c)}\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}{d}$	78

input `int(cos(d*x+c)*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-2*I*b/d*(cos(d*x+c)+1)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(b*sec(d*x+c))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)`

**3.84.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.46

$$\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{-i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

**3.84.6 Sympy [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(c + dx))^{3/2} \cos(c + dx) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))**(3/2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*cos(c + d*x), x)`

**3.84.7 Maxima [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{3/2} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c), x)`

**3.84.8 Giac [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c), x)`

**3.84.9 Mupad [F(-1)]**

Timed out.

$$\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx = \int \cos(c + dx) \left( \frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int(cos(c + d*x)*(b/cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)*(b/cos(c + d*x))^(3/2), x)`

### 3.85 $\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx$

3.85.1	Optimal result . . . . .	561
3.85.2	Mathematica [A] (verified) . . . . .	561
3.85.3	Rubi [A] (verified) . . . . .	562
3.85.4	Maple [C] (verified) . . . . .	563
3.85.5	Fricas [C] (verification not implemented) . . . . .	564
3.85.6	Sympy [F(-1)] . . . . .	564
3.85.7	Maxima [F] . . . . .	564
3.85.8	Giac [F] . . . . .	565
3.85.9	Mupad [F(-1)] . . . . .	565

#### 3.85.1 Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

output `2*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`

#### 3.85.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

input `Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(3/2),x]`

output `(2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

### 3.85.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 2030, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c+dx)(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}}{\csc(c+dx+\frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{\sqrt{b \csc(\frac{1}{2}(2c+\pi)+dx)}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^(3/2),x]`

output `(2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

3.85.3.1 Defintions of rubi rules used

```
rule 2030 Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] :> Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

3.85.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 161, normalized size of antiderivative = 3.93

method	result
default	$-\frac{2\sqrt{b}\sec(dx+c)b\left(i(-\cos(dx+c)-1)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)+i(\cos(dx+c)+1)\sqrt{\frac{1}{\cos(dx+c)+1}}\right)}{d}$
risch	$-\frac{i(e^{2i(dx+c)}+1)\sqrt{2}b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}e^{-i(dx+c)}}{d} - i\left(\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}}\right)$

```
input int(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2/d*(b*sec(d*x+c))^(1/2)*b*(I*(-cos(d*x+c)-1)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(cos(d*x+c)+1)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+(cos(d*x+c)-1)*cot(d*x+c))
```

---

3.85.  $\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx$



**3.85.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.54

$$\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{i \sqrt{2} b^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i \sqrt{2} b^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{d}$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `(I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

**3.85.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(3/2),x)`

output `Timed out`

**3.85.7 Maxima [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{3/2} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^2, x)`

**3.85.8 Giac [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^2, x)`

**3.85.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx = \int \cos(c + dx)^2 \left( \frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2), x)`

### 3.86 $\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx$

3.86.1	Optimal result . . . . .	566
3.86.2	Mathematica [A] (verified) . . . . .	566
3.86.3	Rubi [A] (verified) . . . . .	567
3.86.4	Maple [C] (verified) . . . . .	569
3.86.5	Fricas [C] (verification not implemented) . . . . .	569
3.86.6	Sympy [F(-1)] . . . . .	570
3.86.7	Maxima [F] . . . . .	570
3.86.8	Giac [F] . . . . .	570
3.86.9	Mupad [F(-1)] . . . . .	571

#### 3.86.1 Optimal result

Integrand size = 21, antiderivative size = 70

$$\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b^2 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}}$$

```
output 2/3*b^2*sin(d*x+c)/d/(b*sec(d*x+c))^(1/2)+2/3*b*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d
```

#### 3.86.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.74

$$\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{b\sqrt{b \sec(c + dx)} \left(2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx))\right)}{3d}$$

```
input Integrate[Cos[c + d*x]^3*(b*Sec[c + d*x])^(3/2),x]
```

output  $(b\sqrt{b\sec(c+dx)})(2\sqrt{\cos(c+dx)}\text{EllipticF}[(c+dx)/2, 2] + \text{Sin}[2(c+dx)])/(3d)$

### 3.86.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c+dx)(b\sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b\csc(c+dx+\frac{\pi}{2}))^{3/2}}{\csc(c+dx+\frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{2030} \\
 & b^3 \int \frac{1}{(b\csc(\frac{1}{2}(2c+\pi)+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^3 \left( \frac{\int \sqrt{b\sec(c+dx)} dx}{3b^2} + \frac{2\sin(c+dx)}{3bd\sqrt{b\sec(c+dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{\int \sqrt{b\csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2\sin(c+dx)}{3bd\sqrt{b\sec(c+dx)}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b^3 \left( \frac{\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2\sin(c+dx)}{3bd\sqrt{b\sec(c+dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3b^2} + \frac{2\sin(c+dx)}{3bd\sqrt{b\sec(c+dx)}} \right)
 \end{aligned}$$

$$b^3 \left( \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)$$

input `Int[Cos[c + d*x]^3*(b*Sec[c + d*x])^(3/2), x]`

output `b^3*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]]))`

### 3.86.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d*n)), x] + Simp[(n+1)/(b^2*n) Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.86.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 147, normalized size of antiderivative = 2.10

method	result
default	$\frac{2\left(-i \operatorname{EllipticF}\left(i\left(-\cot(dx+c)+\csc(dx+c)\right), i\right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}\left(i\left(-\cot(dx+c)+\csc(dx+c)\right), i\right) \cos(dx+c)\right)}{3d}$

input `int(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2), x, method=_RETURNVERBOSE)`

output `2/3/d*(-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)+cos(d*x+c)*sin(d*x+c))*(b*sec(d*x+c))^(1/2)*b`

### 3.86.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.21

$$\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) - i \sqrt{2} b^{3/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c))}{3d}$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")`

output `1/3*(2*b*sqrt(b/cos(d*x + c))*cos(d*x + c)*sin(d*x + c) - I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

**3.86.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(b*sec(d*x+c))**(3/2),x)`output `Timed out`**3.86.7 Maxima [F]**

$$\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^3, x)`**3.86.8 Giac [F]**

$$\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^3, x)`

**3.86.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx = \int \cos(c + dx)^3 \left( \frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2),x)`output `int(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2), x)`



### 3.87 $\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx$

3.87.1	Optimal result	572
3.87.2	Mathematica [A] (verified)	572
3.87.3	Rubi [A] (verified)	573
3.87.4	Maple [C] (verified)	574
3.87.5	Fricas [C] (verification not implemented)	575
3.87.6	Sympy [F(-1)]	575
3.87.7	Maxima [F]	576
3.87.8	Giac [F]	576
3.87.9	Mupad [F(-1)]	576

#### 3.87.1 Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{6b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b^3 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}}$$

output  $2/5*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^(3/2)+6/5*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)$

#### 3.87.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.81

$$\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{b\sqrt{b \sec(c + dx)}\left(12\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) + \sin(3(c + dx))\right)}{10d}$$

input `Integrate[Cos[c + d*x]^4*(b*Sec[c + d*x])^(3/2),x]`

output  $(b*\text{Sqrt}[b*\text{Sec}[c + d*x]]*(12*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2] + \text{Sin}[c + d*x] + \text{Sin}[3*(c + d*x)]))/(10*d)$

**3.87.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(c+dx)(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}}{\csc(c+dx+\frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{2030} \\
 & b^4 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^4 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b^4 \left( \frac{3 \int \frac{\sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3119} \\
 & b^4 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)
 \end{aligned}$$

input `Int[Cos[c + d*x]^4*(b*Sec[c + d*x])^(3/2),x]`

output `b^4*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2)))`

### 3.87.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.87.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.60 (sec) , antiderivative size = 175, normalized size of antiderivative = 2.43

method	result
default	$\frac{2\sqrt{b \sec(dx+c)} b \left( i(3 \cos(dx+c)+3) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) + i(-3 \cos(dx+c)-3) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \right)}{5d}$

3.87.  $\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx$

input `int(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2/5/d*(b*sec(d*x+c))^(1/2)*b*(I*(3*cos(d*x+c)+3)*(1/(cos(d*x+c)+1))^(1/2)*  
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+  
I*(-3*cos(d*x+c)-3)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(  
1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+(-cos(d*x+c)^3-2*cos(d*x+c)+3  
)cot(d*x+c))`

### 3.87.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.29

$$\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^2 \sin(dx+c) + 3i \sqrt{2} b^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + I \sin(dx+c))) - 3i \sqrt{2} b^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - I \sin(dx+c)))}{d}$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/5*(2*b*sqrt(b/cos(d*x + c))*cos(d*x + c)^2*sin(d*x + c) + 3*I*sqrt(2)*b^(  
(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*s  
in(d*x + c))) - 3*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInv  
erse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

### 3.87.6 Sympy [F(-1)]

Timed out.

$$\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**4*(b*sec(d*x+c))**(3/2),x)`

output `Timed out`

**3.87.7 Maxima [F]**

$$\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^4 dx$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^4, x)`

**3.87.8 Giac [F]**

$$\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^4 dx$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^4, x)`

**3.87.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx = \int \cos(c + dx)^4 \left( \frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2), x)`

### 3.88 $\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx$

3.88.1	Optimal result . . . . .	577
3.88.2	Mathematica [A] (verified) . . . . .	577
3.88.3	Rubi [A] (verified) . . . . .	578
3.88.4	Maple [C] (verified) . . . . .	580
3.88.5	Fricas [C] (verification not implemented) . . . . .	581
3.88.6	Sympy [F(-1)] . . . . .	581
3.88.7	Maxima [F] . . . . .	581
3.88.8	Giac [F] . . . . .	582
3.88.9	Mupad [F(-1)] . . . . .	582

#### 3.88.1 Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{10b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{2b^4 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^2 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}}$$

output  $2/7*b^4*\sin(d*x+c)/d/(b*\sec(d*x+c))^(5/2)+10/21*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^(1/2)+10/21*b*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)*(b*\sec(d*x+c))^(1/2)/d$

#### 3.88.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

$$\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{b\sqrt{b \sec(c + dx)}\left(40\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 26 \sin(2(c + dx)) + 3 \sin(4(c + dx))\right)}{84d}$$

input `Integrate[Cos[c + d*x]^5*(b*Sec[c + d*x])^(3/2), x]`

output  $(b\sqrt{b\sec[c + dx]}*(40\sqrt{\cos[c + dx]}*EllipticF[(c + dx)/2, 2] + 26*\sin[2*(c + dx)] + 3*\sin[4*(c + dx)]))/(84*d)$

### 3.88.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}}{\csc(c + dx + \frac{\pi}{2})^5} dx \\
 & \quad \downarrow \text{2030} \\
 & b^5 \int \frac{1}{(b \csc(\frac{1}{2}(2c + \pi) + dx))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^5 \left( \frac{5 \int \frac{1}{(b \sec(c + dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^5 \left( \frac{5 \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^5 \left( \frac{5 \left( \frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd\sqrt{b \sec(c + dx)}} \right)}{7b^2} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& b^5 \left( \frac{5 \left( \frac{\int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 4258 \\
& b^5 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 3042 \\
& b^5 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 3120 \\
& b^5 \left( \frac{5 \left( \frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)
\end{aligned}$$

input `Int[Cos[c + d*x]^5*(b*Sec[c + d*x])^(3/2), x]`

output `b^5*((2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/(7*b^2))`

### 3.88.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx._)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.88.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.67

method	result
default	$\frac{2 \left( -5i \operatorname{EllipticF} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) + 3 \cos(dx+c)^3 \sin(dx+c) - 5i \sqrt{\frac{1}{\cos(dx+c)+1}} \right)}{21d}$

input `int(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2), x, method=_RETURNVERBOSE)`

output `2/21/d*(-5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)+3*cos(d*x+c)^3*sin(d*x+c)-5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)+5*cos(d*x+c)*sin(d*x+c))*(b*sec(d*x+c))^(1/2)*b`

**3.88.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.01

$$\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{-5i \sqrt{2} b^{3/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} b^{3/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2(3b \cos(dx + c)^3 + 5b \cos(dx + c)) \sqrt{b/\cos(dx + c)} \sin(dx + c)}{d}$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/21*(-5*I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(3*b*cos(d*x + c)^3 + 5*b*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c))/d`

**3.88.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**5*(b*sec(d*x+c))**(3/2),x)`

output `Timed out`

**3.88.7 Maxima [F]**

$$\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{3/2} \cos(dx + c)^5 dx$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^5, x)`

**3.88.8 Giac [F]**

$$\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^5 dx$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^5, x)`

**3.88.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx = \int \cos(c + dx)^5 \left( \frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2), x)`

### 3.89 $\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx$

3.89.1	Optimal result . . . . .	583
3.89.2	Mathematica [A] (verified) . . . . .	583
3.89.3	Rubi [A] (verified) . . . . .	584
3.89.4	Maple [C] (verified) . . . . .	586
3.89.5	Fricas [C] (verification not implemented) . . . . .	586
3.89.6	Sympy [F(-1)] . . . . .	587
3.89.7	Maxima [F] . . . . .	587
3.89.8	Giac [F] . . . . .	587
3.89.9	Mupad [F(-1)] . . . . .	588

#### 3.89.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{14b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b^5 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}$$

```
output 2/9*b^5*sin(d*x+c)/d/(b*sec(d*x+c))^(7/2)+14/45*b^3*sin(d*x+c)/d/(b*sec(d*x+c))^(3/2)+14/15*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)
```

#### 3.89.2 Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.72

$$\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{b\sqrt{b \sec(c + dx)}\left(84\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) + \cos^2(c + dx)(33 \sin(c + dx) + 5 \sin(3(c + dx)))\right)}{90d}$$

```
input Integrate[Cos[c + d*x]^6*(b*Sec[c + d*x])^(3/2),x]
```

```
output (b*Sqrt[b*Sec[c + d*x]]*(84*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]^2*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)])))/(90*d)
```

**3.89.3 Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^6(c+dx)(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}}{\csc(c+dx+\frac{\pi}{2})^6} dx \\
 & \quad \downarrow \text{2030} \\
 & b^6 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{9/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^6 \left( \frac{7 \int \frac{1}{(b \sec(c+dx))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^6 \left( \frac{7 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^6 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^6 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 4258 \\
 & b^6 \left( \frac{7 \left( \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \downarrow 3042 \\
 & b^6 \left( \frac{7 \left( \frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \downarrow 3119 \\
 & b^6 \left( \frac{7 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{aligned}$$

input `Int[Cos[c + d*x]^6*(b*Sec[c + d*x])^(3/2),x]`

output `b^6*((2*Sin[c + d*x])/(9*b*d*(b*Sec[c + d*x])^(7/2)) + (7*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))))/(9*b^2)`

### 3.89.3.1 Defintions of rubi rules used

rule 2030 `Int[(F*_)(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F_x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.89.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.71 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.85

method	result
default	$\frac{2\sqrt{b\sec(dx+c)}b\left(i(21\cos(dx+c)+21)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)+i(-21\cos(dx+c)-21)\right)}{\dots}$

input `int(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{2}{45}d*(b*\sec(d*x+c))^{1/2}*b*(I*(21*\cos(d*x+c)+21)*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\operatorname{EllipticE}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+I*(-21*\cos(d*x+c)-21)*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I))+(-5*\cos(d*x+c)^5-2*\cos(d*x+c)^3-14*\cos(d*x+c)+21)*\cot(d*x+c)$$

### 3.89.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.07

$$\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{21i\sqrt{2}b^{3/2}\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 21i\sqrt{2}b^{3/2}\operatorname{weierstrassZeta}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{\dots}$$

input `integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/45*(21*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(5*b*cos(d*x + c)^4 + 7*b*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/d`

### 3.89.6 Sympy [F(-1)]

Timed out.

$$\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**6*(b*sec(d*x+c))**(3/2),x)`

output Timed out

### 3.89.7 Maxima [F]

$$\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^6 dx$$

input `integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^6, x)`

### 3.89.8 Giac [F]

$$\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^6 dx$$

input `integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^6, x)`



**3.89.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx = \int \cos(c + dx)^6 \left( \frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2),x)`output `int(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2), x)`

### 3.90 $\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx$

3.90.1	Optimal result . . . . .	589
3.90.2	Mathematica [A] (verified) . . . . .	589
3.90.3	Rubi [A] (verified) . . . . .	590
3.90.4	Maple [C] (verified) . . . . .	592
3.90.5	Fricas [C] (verification not implemented) . . . . .	592
3.90.6	Sympy [F] . . . . .	593
3.90.7	Maxima [F] . . . . .	593
3.90.8	Giac [F] . . . . .	593
3.90.9	Mupad [F(-1)] . . . . .	594

#### 3.90.1 Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{10b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{10b(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7bd}$$

output

```
10/21*b*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b/d+10/21*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d
```

#### 3.90.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.62

$$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{(b \sec(c + dx))^{5/2} \left( 10 \cos^{\frac{5}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 5 \sin(2(c + dx)) + 6 \tan(c + dx) \right)}{21d}$$

input

```
Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(5/2),x]
```

output  $((b*\text{Sec}[c + d*x])^{5/2}*(10*\text{Cos}[c + d*x]^{5/2}*\text{EllipticF}[(c + d*x)/2, 2] + 5*\text{Sin}[2*(c + d*x)] + 6*\text{Tan}[c + d*x]))/(21*d)$

### 3.90.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.10, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c + dx))^{9/2} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{9/2} dx}{b^2} \\
 & \quad \downarrow \text{4255} \\
 & \quad \frac{\frac{5}{7}b^2 \int (b \sec(c + dx))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\frac{5}{7}b^2 \int (b \csc(c + dx + \frac{\pi}{2}))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^2} \\
 & \quad \downarrow \text{4255} \\
 & \quad \frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^2} \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^2}$$

↓ 3042

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^2}$$

↓ 3120

$$\frac{\frac{5}{7}b^2 \left( \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^2}$$

input `Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(5/2),x]`

output `((2*b*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*d) + (5*b^2*((2*b^2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d)))/7)/b^2`

### 3.90.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Simp[b^2*((n-2)/(n-1)) Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_], x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.90.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.77 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.64

method	result
default	$-\frac{2ib^2\sqrt{b\sec(dx+c)}\left(5\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\cos(dx+c)+5\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\right)}{21d}$

input `int(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$-2/21*I/d*b^2*(b*\sec(d*x+c))^{1/2}*(5*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*\cos(d*x+c)+5*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+5*I*\tan(d*x+c)+3*I*\tan(d*x+c)*\sec(d*x+c)^2)$$

### 3.90.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.23

$$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{-5i\sqrt{2}b^{5/2}\cos(dx+c)^3\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+5i\sqrt{2}b^{5/2}\cos(dx+c)^3\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))+2*(5*b^2*\cos(dx+c)^2+3*b^2)*\sqrt{b/\cos(dx+c)}*\sin(dx+c)/(d*\cos(dx+c)^3)}{21d}$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output 
$$1/21*(-5*I*\sqrt{2}*b^{5/2}*\cos(d*x+c)^3*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+5*I*\sqrt{2}*b^{5/2}*\cos(d*x+c)^3*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))+2*(5*b^2*\cos(d*x+c)^2+3*b^2)*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c)/(d*\cos(d*x+c)^3)$$

---

3.90.  $\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx$

**3.90.6 Sympy [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(c + dx))^{\frac{5}{2}} \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(5/2),x)`

output `Integral((b*sec(c + d*x))**(5/2)*sec(c + d*x)**2, x)`

**3.90.7 Maxima [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^2, x)`

**3.90.8 Giac [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^2, x)`

**3.90.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}}{\cos(c + dx)^2} dx$$

input `int((b/cos(c + d*x))^(5/2)/cos(c + d*x)^2,x)`output `int((b/cos(c + d*x))^(5/2)/cos(c + d*x)^2, x)`

### 3.91 $\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx$

3.91.1	Optimal result . . . . .	595
3.91.2	Mathematica [A] (verified) . . . . .	595
3.91.3	Rubi [A] (verified) . . . . .	596
3.91.4	Maple [C] (verified) . . . . .	598
3.91.5	Fricas [C] (verification not implemented) . . . . .	598
3.91.6	Sympy [F] . . . . .	599
3.91.7	Maxima [F] . . . . .	599
3.91.8	Giac [F] . . . . .	599
3.91.9	Mupad [F(-1)] . . . . .	600

#### 3.91.1 Optimal result

Integrand size = 19, antiderivative size = 97

$$\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx = -\frac{6b^3 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{6b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d}$$

output `2/5*(b*sec(d*x+c))^(5/2)*sin(d*x+c)/d-6/5*b^3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+6/5*b^2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d`

#### 3.91.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.63

$$\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{(b \sec(c + dx))^{5/2} \left( -12 \cos^{5/2}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 7 \sin(c + dx) + 3 \sin(3(c + dx)) \right)}{10d}$$

input `Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(5/2),x]`

output `((b*Sec[c + d*x])^(5/2)*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*d)`



### 3.91.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.07, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec(c+dx)(b \sec(c+dx))^{5/2} dx \\
 \downarrow \text{2030} \\
 \frac{\int (b \sec(c+dx))^{7/2} dx}{b} \\
 \downarrow \text{3042} \\
 \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{7/2} dx}{b} \\
 \downarrow \text{4255} \\
 \frac{\frac{3}{5}b^2 \int (b \sec(c+dx))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b} \\
 \downarrow \text{3042} \\
 \frac{\frac{3}{5}b^2 \int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b} \\
 \downarrow \text{4255} \\
 \frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b} \\
 \downarrow \text{3042} \\
 \frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b} \\
 \downarrow \text{4258} \\
 \frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b} \\
 \downarrow \text{3042}
 \end{array}$$

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b}$$

↓ 3119

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b}$$

input `Int[Sec[c + d*x]*(b*Sec[c + d*x])^(5/2),x]`

output `((2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5)/b`

### 3.91.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b.)*(v.))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.91.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.64 (sec) , antiderivative size = 415, normalized size of antiderivative = 4.28

method	result
default	$2b^2 \sqrt{b \sec(dx+c)} \left( 3i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)^2 - 3i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) \right)$

input `int(sec(d*x+c)*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 2/5/d*b^2*(b*\sec(d*x+c))^{(1/2)}/(\cos(d*x+c)+1)*(3*I*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)} \\ & *\cos(d*x+c)^2-3*I*\operatorname{EllipticE}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\cos(d*x+c)^2+6*I*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I) \\ & *(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\cos(d*x+c)-6*I*\operatorname{EllipticE}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\cos(d*x+c) \\ & +3*I*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)-3*I*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)} \\ & *\operatorname{EllipticE}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+3*\sin(d*x+c)+\tan(d*x+c)+\sec(d*x+c)*\tan(d*x+c) \end{aligned}$$

### 3.91.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.29

$$\int \sec(c+dx)(b \sec(c+dx))^{5/2} dx = \frac{-3i \sqrt{2} b^{5/2} \cos(dx+c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c)))}{\cos(dx+c)}$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/5*(-3*I*sqrt(2)*b^(5/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*b^(5/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*b^2*cos(d*x + c)^2 + b^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^2)`

### 3.91.6 Sympy [F]

$$\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(c + dx))^{\frac{5}{2}} \sec(c + dx) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))**(5/2),x)`

output `Integral((b*sec(c + d*x))**(5/2)*sec(c + d*x), x)`

### 3.91.7 Maxima [F]

$$\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c), x)`

### 3.91.8 Giac [F]

$$\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c), x)`

**3.91.9 Mupad [F(-1)]**

Timed out.

$$\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}}{\cos(c + dx)} dx$$

input `int((b/cos(c + d*x))^(5/2)/cos(c + d*x),x)`output `int((b/cos(c + d*x))^(5/2)/cos(c + d*x), x)`

### 3.92 $\int (b \sec(c + dx))^{5/2} dx$

3.92.1	Optimal result . . . . .	601
3.92.2	Mathematica [A] (verified) . . . . .	601
3.92.3	Rubi [A] (verified) . . . . .	602
3.92.4	Maple [C] (verified) . . . . .	603
3.92.5	Fricas [C] (verification not implemented) . . . . .	604
3.92.6	Sympy [F] . . . . .	604
3.92.7	Maxima [F] . . . . .	604
3.92.8	Giac [F] . . . . .	605
3.92.9	Mupad [F(-1)] . . . . .	605

#### 3.92.1 Optimal result

Integrand size = 12, antiderivative size = 70

$$\int (b \sec(c + dx))^{5/2} dx = \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d}$$

```
output 2/3*b*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/d+2/3*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d
```

#### 3.92.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.73

$$\int (b \sec(c + dx))^{5/2} dx = \frac{2b^2 \sqrt{b \sec(c + dx)} \left( \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx) \right)}{3d}$$

```
input Integrate[(b*Sec[c + d*x])^(5/2),x]
```

```
output (2*b^2*sqrt[b*Sec[c + d*x]]*(sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d)
```

**3.92.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( b \csc \left( c + dx + \frac{\pi}{2} \right) \right)^{5/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{3} b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} b^2 \int \sqrt{b \csc \left( c + dx + \frac{\pi}{2} \right)} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{4258} \\
 & \frac{1}{3} b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin \left( c + dx + \frac{\pi}{2} \right)}} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF} \left( \frac{1}{2}(c + dx), 2 \right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(5/2),x]`

output `(2*b^2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d)`

## 3.92.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.92.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.99 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.06

method	result
default	$-\frac{2\sqrt{b\sec(dx+c)}b^2\left(i\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)+i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\right)}{3d}$

input `int((b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/3/d*(b*sec(d*x+c))^(1/2)*b^2*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-tan(d*x+c))`



**3.92.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.44

$$\int (b \sec(c + dx))^{5/2} dx = \frac{-i \sqrt{2} b^{5/2} \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{5/2} \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2 b^2 \sqrt{b/\cos(dx + c)} \sin(dx + c)}{3 d \cos(dx + c)}$$

input `integrate((b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/3*(-I*sqrt(2)*b^(5/2)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*b^(5/2)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*b^2*sqrt(b/cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c))`

**3.92.6 Sympy [F]**

$$\int (b \sec(c + dx))^{5/2} dx = \int (b \sec(c + dx))^{\frac{5}{2}} dx$$

input `integrate((b*sec(d*x+c))**(5/2),x)`

output `Integral((b*sec(c + d*x))**(5/2), x)`

**3.92.7 Maxima [F]**

$$\int (b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} dx$$

input `integrate((b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2), x)`

**3.92.8 Giac [F]**

$$\int (b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} dx$$

input `integrate((b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2), x)`

**3.92.9 Mupad [F(-1)]**

Timed out.

$$\int (b \sec(c + dx))^{5/2} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int((b/cos(c + d*x))^(5/2),x)`

output `int((b/cos(c + d*x))^(5/2), x)`

### 3.93 $\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx$

3.93.1	Optimal result . . . . .	606
3.93.2	Mathematica [A] (verified) . . . . .	606
3.93.3	Rubi [A] (verified) . . . . .	607
3.93.4	Maple [C] (verified) . . . . .	608
3.93.5	Fricas [C] (verification not implemented) . . . . .	609
3.93.6	Sympy [F(-1)] . . . . .	610
3.93.7	Maxima [F] . . . . .	610
3.93.8	Giac [F] . . . . .	610
3.93.9	Mupad [F(-1)] . . . . .	611

#### 3.93.1 Optimal result

Integrand size = 19, antiderivative size = 68

$$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx = -\frac{2b^3 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{d}$$

output

```
-2*b^3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*b^2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d
```

#### 3.93.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.74

$$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{2b^2 \sqrt{b \sec(c + dx)} \left( -\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d}$$

input

```
Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(5/2),x]
```

output

```
(2*b^2*Sqrt[b*Sec[c + d*x]]*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/d
```

**3.93.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {3042, 2030, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c+dx)(b \sec(c+dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}}{\csc(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \left( b \csc\left(\frac{1}{2}(2c+\pi)+dx\right) \right)^{3/2} dx \\
 & \quad \downarrow \text{4255} \\
 & b \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right) \\
 & \quad \downarrow \text{3042} \\
 & b \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right) \\
 & \quad \downarrow \text{4258} \\
 & b \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) \\
 & \quad \downarrow \text{3119} \\
 & b \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)
 \end{aligned}$$

input `Int[Cos[c + d*x]*(b*Sec[c + d*x])^(5/2),x]`

output `b*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d)`

### 3.93.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx._)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] :> Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.93.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 6.80 (sec) , antiderivative size = 396, normalized size of antiderivative = 5.82

method	result
default	$-\frac{2b^2 \left( i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)^2 - i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i) \right)}{d}$

---

3.93.  $\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx$

```
input int(cos(d*x+c)*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2*b^2/d*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*
(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2+2*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*
(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-2*I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*
(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-I*(1/(cos(d*x+c)+1))^(1/2)*
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-sin(d*x+c)*(b*sec(d*x+c))^(1/2)/(cos(d*x+c)+1)
```

### 3.93.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.26

$$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{-i \sqrt{2} b^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{2} + 2*b^2*\sqrt{b/\cos(d*x + c)}*\sin(d*x + c)/d$$

```
input integrate(cos(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output (-I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*b^2*sqrt(b/cos(d*x + c))*sin(d*x + c)/d
```

**3.93.6 Sympy [F(-1)]**

Timed out.

$$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))**(5/2),x)`output `Timed out`**3.93.7 Maxima [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c), x)`**3.93.8 Giac [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c), x)`

**3.93.9 Mupad [F(-1)]**

Timed out.

$$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx = \int \cos(c + dx) \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int(cos(c + d*x)*(b/cos(c + d*x))^(5/2), x)`output `int(cos(c + d*x)*(b/cos(c + d*x))^(5/2), x)`



### 3.94 $\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx$

3.94.1	Optimal result . . . . .	612
3.94.2	Mathematica [A] (verified) . . . . .	612
3.94.3	Rubi [A] (verified) . . . . .	613
3.94.4	Maple [C] (verified) . . . . .	614
3.94.5	Fricas [C] (verification not implemented) . . . . .	615
3.94.6	Sympy [F(-1)] . . . . .	615
3.94.7	Maxima [F] . . . . .	615
3.94.8	Giac [F] . . . . .	616
3.94.9	Mupad [F(-1)] . . . . .	616

#### 3.94.1 Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \cos^2(c+dx)(b \sec(c+dx))^{5/2} dx = \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{d}$$

output `2*b^2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.94.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \cos^2(c+dx)(b \sec(c+dx))^{5/2} dx = \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{d}$$

input `Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(5/2),x]`

output `(2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d`

**3.94.3 Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 2030, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c+dx)(b \sec(c+dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}}{\csc(c+dx+\frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \sqrt{b \csc\left(\frac{1}{2}(2c+\pi)+dx\right)} dx \\
 & \quad \downarrow \text{4258} \\
 & b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3120} \\
 & \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^(5/2),x]`

output `(2*b^2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/d`

## 3.94.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx._)*(v-)^(m-)*((b-)*(v-))^(n-), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c-) + (d-)*(x-)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c-) + (d-)*(x-)]*(b-))^(n-), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.94.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 31.78 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.95

method	result	size
default	$-\frac{2ib^2(\cos(dx+c)+1)\operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{b\sec(dx+c)}\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}{d}$	80

input `int(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `-2*I*b^2/d*(cos(d*x+c)+1)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(b*sec(d*x+c))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)`

**3.94.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.39

$$\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{-i \sqrt{2} b^{5/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{5/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

**3.94.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

**3.94.7 Maxima [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^2, x)`

**3.94.8 Giac [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^2, x)`

**3.94.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx = \int \cos(c + dx)^2 \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2),x)`

output `int(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2), x)`

### 3.95 $\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx$

3.95.1	Optimal result	617
3.95.2	Mathematica [A] (verified)	617
3.95.3	Rubi [A] (verified)	618
3.95.4	Maple [C] (verified)	619
3.95.5	Fricas [C] (verification not implemented)	620
3.95.6	Sympy [F(-1)]	620
3.95.7	Maxima [F]	620
3.95.8	Giac [F]	621
3.95.9	Mupad [F(-1)]	621

#### 3.95.1 Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{2b^3 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

output `2*b^3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`

#### 3.95.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93

$$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{2 \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) (b \sec(c + dx))^{5/2}}{d}$$

input `Integrate[Cos[c + d*x]^3*(b*Sec[c + d*x])^(5/2),x]`

output `(2*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2]*(b*Sec[c + d*x])^(5/2))/d`

**3.95.3 Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 2030, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c+dx)(b \sec(c+dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}}{\csc(c+dx+\frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{2030} \\
 & b^3 \int \frac{1}{\sqrt{b \csc(\frac{1}{2}(2c+\pi)+dx)}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{b^3 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2b^3 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^3*(b*Sec[c + d*x])^(5/2),x]`

output `(2*b^3*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

## 3.95.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx._)*(v.)^(m.)*((b.)*(v.)^(n.), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c.) + (d.)*(x.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c.) + (d.)*(x.)]*(b.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.95.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 123.02 (sec) , antiderivative size = 165, normalized size of antiderivative = 4.02

method	result
default	$\frac{2\sqrt{b\sec(dx+c)}b^2\left(i(\cos(dx+c)+1)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)+i(-\cos(dx+c)-1)\sqrt{\frac{1}{\cos(dx+c)+1}}\right)}{d}$
risch	$-\frac{i(e^{2i(dx+c)}+1)\sqrt{2}b^2e^{-i(dx+c)}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}}{d} - i\left(\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}}\right)$

input `int(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `2/d*(b*sec(d*x+c))^(1/2)*b^2*(I*(cos(d*x+c)+1)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(-cos(d*x+c)-1)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+(1-cos(d*x+c))*cot(d*x+c)`



**3.95.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.54

$$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{i \sqrt{2} b^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i \sqrt{2} b^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{d}$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `(I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

**3.95.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

**3.95.7 Maxima [F]**

$$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^3, x)`

**3.95.8 Giac [F]**

$$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^3, x)`

**3.95.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx = \int \cos(c + dx)^3 \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2),x)`

output `int(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2), x)`

### 3.96 $\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx$

3.96.1	Optimal result . . . . .	622
3.96.2	Mathematica [A] (verified) . . . . .	622
3.96.3	Rubi [A] (verified) . . . . .	623
3.96.4	Maple [C] (verified) . . . . .	625
3.96.5	Fricas [C] (verification not implemented) . . . . .	625
3.96.6	Sympy [F(-1)] . . . . .	626
3.96.7	Maxima [F] . . . . .	626
3.96.8	Giac [F] . . . . .	626
3.96.9	Mupad [F(-1)] . . . . .	627

#### 3.96.1 Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b^3 \sin(c + dx)}{3d \sqrt{b \sec(c + dx)}}$$

output  $2/3*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^(1/2)+2/3*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)*(b*\sec(d*x+c))^(1/2)/d$

#### 3.96.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.75

$$\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{b^2 \sqrt{b \sec(c + dx)} \left( 2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx)) \right)}{3d}$$

input `Integrate[Cos[c + d*x]^4*(b*Sec[c + d*x])^(5/2), x]`

```
output (b^2*Sqrt[b*Sec[c + d*x]]*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]
+ Sin[2*(c + d*x)]))/(3*d)
```

### 3.96.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}}{\csc(c + dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{2030} \\
 & b^4 \int \frac{1}{(b \csc(\frac{1}{2}(2c + \pi) + dx))^{3/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^4 \left( \frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{\int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b^4 \left( \frac{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right)
 \end{aligned}$$

$$b^4 \left( \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)$$

input `Int[Cos[c + d*x]^4*(b*Sec[c + d*x])^(5/2), x]`

output `b^4*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])`

### 3.96.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d*n), x] + Simp[(n+1)/(b^2*n) Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

**3.96.4 Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 4.26 (sec) , antiderivative size = 150, normalized size of antiderivative = 2.08

$$2 \left( i \operatorname{EllipticF} \left( i \left( -\cot(dx+c) + \csc(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) + i \sqrt{\frac{1}{\cos(dx+c)+1}} \right)$$

input `int(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2),x)`

output `-2/3/d*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-cos(d*x+c)*sin(d*x+c))*(b*sec(d*x+c))^(1/2)*b^2`

**3.96.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.21

$$\int \cos^4(c+dx)(b \sec(c+dx))^{5/2} dx = \frac{2b^2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) - i \sqrt{2} b^{5/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c))}{3d}$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/3*(2*b^2*sqrt(b/cos(d*x+c))*cos(d*x+c)*sin(d*x+c) - I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x+c) + I*sin(d*x+c)) + I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x+c) - I*sin(d*x+c)))/d`

**3.96.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**4*(b*sec(d*x+c))**(5/2),x)`output `Timed out`**3.96.7 Maxima [F]**

$$\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^4 dx$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^4, x)`**3.96.8 Giac [F]**

$$\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^4 dx$$

input `integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^4, x)`

**3.96.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx = \int \cos(c + dx)^4 \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2),x)`output `int(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2), x)`



### 3.97 $\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx$

3.97.1	Optimal result . . . . .	628
3.97.2	Mathematica [A] (verified) . . . . .	628
3.97.3	Rubi [A] (verified) . . . . .	629
3.97.4	Maple [C] (verified) . . . . .	630
3.97.5	Fricas [C] (verification not implemented) . . . . .	631
3.97.6	Sympy [F(-1)] . . . . .	631
3.97.7	Maxima [F] . . . . .	632
3.97.8	Giac [F] . . . . .	632
3.97.9	Mupad [F(-1)] . . . . .	632

#### 3.97.1 Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{6b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b^4 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}}$$

```
output 2/5*b^4*sin(d*x+c)/d/(b*sec(d*x+c))^(3/2)+6/5*b^3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)
```

#### 3.97.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.83

$$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{b^2 \sqrt{b \sec(c + dx)} \left( 12 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) + \sin(c + dx) + \sin(3(c + dx)) \right)}{10d}$$

```
input Integrate[Cos[c + d*x]^5*(b*Sec[c + d*x])^(5/2),x]
```

```
output (b^2*Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*d)
```

**3.97.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^5(c+dx)(b \sec(c+dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}}{\csc(c+dx+\frac{\pi}{2})^5} dx \\
 & \quad \downarrow \text{2030} \\
 & b^5 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^5 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^5 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b^5 \left( \frac{3 \int \frac{\sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^5 \left( \frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3119} \\
 & b^5 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)
 \end{aligned}$$

input `Int[Cos[c + d*x]^5*(b*Sec[c + d*x])^(5/2),x]`

output `b^5*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2)))`

### 3.97.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.97.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.25 (sec) , antiderivative size = 175, normalized size of antiderivative = 2.43

$$\frac{2\sqrt{b \sec(dx+c)} b^2 \left( i(-3 \cos(dx+c) - 3) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticE}(i(-\cot(dx+c) + \csc(dx+c))) \right)}{1}$$

input `int(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2),x)`

---

3.97.  $\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx$

output 
$$-2/5/d*(b*\sec(d*x+c))^{(1/2)}*b^2*(I*(-3*\cos(d*x+c)-3)*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*EllipticE(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+I*(3*\cos(d*x+c)+3)*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*EllipticF(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+(\cos(d*x+c)^3+2*\cos(d*x+c)-3)*\cot(d*x+c))$$

### 3.97.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.32

$$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{2b^2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^2 \sin(dx+c) + 3i \sqrt{2} b^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - I \sin(dx+c)))}{d}$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output 
$$1/5*(2*b^2*\sqrt{b/\cos(d*x+c)}*\cos(d*x+c)^2*\sin(d*x+c) + 3*I*\sqrt{2}*b^{(5/2)}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x+c) + I*\sin(d*x+c))) - 3*I*\sqrt{2}*b^{(5/2)}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x+c) - I*\sin(d*x+c))))/d$$

### 3.97.6 Sympy [F(-1)]

Timed out.

$$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**5*(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

**3.97.7 Maxima [F]**

$$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^5 dx$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^5, x)`

**3.97.8 Giac [F]**

$$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^5 dx$$

input `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^5, x)`

**3.97.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx = \int \cos(c + dx)^5 \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2),x)`

output `int(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2), x)`

### 3.98 $\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx$

3.98.1	Optimal result . . . . .	633
3.98.2	Mathematica [A] (verified) . . . . .	633
3.98.3	Rubi [A] (verified) . . . . .	634
3.98.4	Maple [C] (verified) . . . . .	636
3.98.5	Fricas [C] (verification not implemented) . . . . .	636
3.98.6	Sympy [F(-1)] . . . . .	637
3.98.7	Maxima [F] . . . . .	637
3.98.8	Giac [F] . . . . .	637
3.98.9	Mupad [F(-1)] . . . . .	638

#### 3.98.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{10b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{2b^5 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^3 \sin(c + dx)}{21d \sqrt{b \sec(c + dx)}}$$

output  $2/7*b^5*\sin(d*x+c)/d/(b*\sec(d*x+c))^(5/2)+10/21*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^(1/2)+10/21*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)*(b*\sec(d*x+c))^(1/2)/d$

#### 3.98.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.66

$$\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{b^2 \sqrt{b \sec(c + dx)} \left( 40 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) \right)}{84d}$$

input `Integrate[Cos[c + d*x]^6*(b*Sec[c + d*x])^(5/2), x]`

output  $(b^2 \sqrt{b \sec[c + dx]} (40 \sqrt{\cos[c + dx]} \text{EllipticF}[(c + dx)/2, 2] + 26 \sin[2(c + dx)] + 3 \sin[4(c + dx)])) / (84d)$

### 3.98.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^6(c + dx) (b \sec(c + dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}}{\csc(c + dx + \frac{\pi}{2})^6} dx \\
 & \quad \downarrow \text{2030} \\
 & b^6 \int \frac{1}{(b \csc(\frac{1}{2}(2c + \pi) + dx))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^6 \left( \frac{5 \int \frac{1}{(b \sec(c + dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^6 \left( \frac{5 \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^6 \left( \frac{5 \left( \frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right)}{7b^2} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& b^6 \left( \frac{5 \left( \frac{\int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 4258 \\
& b^6 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 3042 \\
& b^6 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 3120 \\
& b^6 \left( \frac{5 \left( \frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)
\end{aligned}$$

input `Int[Cos[c + d*x]^6*(b*Sec[c + d*x])^(5/2), x]`

output `b^6*((2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/(7*b^2))`

### 3.98.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)*(b_.)]^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)*(b_.)]^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.98.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.89 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.66

$$\frac{2 \left( 5i \operatorname{EllipticF} \left( i \left( -\cot(dx+c) + \csc(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) + 5i \sqrt{\frac{1}{\cos(dx+c)+1}} \right)}{}$$

input `int(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2),x)`

output `-2/21/d*(5*I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-3*cos(d*x+c)^3*sin(d*x+c)-5*cos(d*x+c)*sin(d*x+c))*(b*sec(d*x+c))^(1/2)*b^2`

### 3.98.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.03

$$\int \cos^6(c+dx)(b \sec(c+dx))^{5/2} dx = \frac{-5i \sqrt{2} b^{5/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c)) + 5i \sqrt{2} b^{5/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) - i \sin(dx+c))}{}$$

3.98.  $\int \cos^6(c+dx)(b \sec(c+dx))^{5/2} dx$

input `integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/21*(-5*I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(3*b^2*cos(d*x + c)^3 + 5*b^2*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c))/d`

### 3.98.6 Sympy [F(-1)]

Timed out.

$$\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**6*(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

### 3.98.7 Maxima [F]

$$\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^6 dx$$

input `integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^6, x)`

### 3.98.8 Giac [F]

$$\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^6 dx$$

input `integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^6, x)`

**3.98.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx = \int \cos(c + dx)^6 \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2),x)`output `int(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2), x)`

### 3.99 $\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx$

3.99.1	Optimal result . . . . .	639
3.99.2	Mathematica [A] (verified) . . . . .	639
3.99.3	Rubi [A] (verified) . . . . .	640
3.99.4	Maple [C] (verified) . . . . .	642
3.99.5	Fricas [C] (verification not implemented) . . . . .	642
3.99.6	Sympy [F(-1)] . . . . .	643
3.99.7	Maxima [F] . . . . .	643
3.99.8	Giac [F] . . . . .	643
3.99.9	Mupad [F(-1)] . . . . .	644

#### 3.99.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{14b^3 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b^6 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}$$

output  $2/9*b^6*\sin(d*x+c)/d/(b*\sec(d*x+c))^(7/2)+14/45*b^4*\sin(d*x+c)/d/(b*\sec(d*x+c))^(3/2)+14/15*b^3*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*EllipticE(\sin(1/2*d*x+1/2*c),2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)$

#### 3.99.2 Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.74

$$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{b^2 \sqrt{b \sec(c + dx)} \left( 84 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \cos^2(c + dx)(33 \sin(c + dx) + 5 \sin(3(c + dx))) \right)}{90d}$$

input `Integrate[Cos[c + d*x]^7*(b*Sec[c + d*x])^(5/2),x]`

output  $(b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*(84*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2] + \text{Cos}[c + d*x]^2*(33*\text{Sin}[c + d*x] + 5*\text{Sin}[3*(c + d*x)])))/(90*d)$

**3.99.3 Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^7(c+dx)(b \sec(c+dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}}{\csc(c+dx+\frac{\pi}{2})^7} dx \\
 & \quad \downarrow \text{2030} \\
 & b^7 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{9/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^7 \left( \frac{7 \int \frac{1}{(b \sec(c+dx))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^7 \left( \frac{7 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^7 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^7 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 4258 \\
 b^7 \left( \frac{7 \left( \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 \downarrow 3042 \\
 b^7 \left( \frac{7 \left( \frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 \downarrow 3119 \\
 b^7 \left( \frac{7 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{array}$$

input `Int[Cos[c + d*x]^7*(b*Sec[c + d*x])^(5/2),x]`

output `b^7*((2*Sin[c + d*x])/(9*b*d*(b*Sec[c + d*x])^(7/2)) + (7*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))))/(9*b^2)`

### 3.99.3.1 Defintions of rubi rules used

rule 2030 `Int[(F*_)(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F_x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.99.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.46 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.87

$$2\sqrt{b \sec(dx + c)} b^2 \left( i(-21 \cos(dx + c) - 21) \sqrt{\frac{1}{\cos(dx + c) + 1}} \sqrt{\frac{\cos(dx + c)}{\cos(dx + c) + 1}} \operatorname{EllipticE}(i(-\cot(dx + c) + \csc(dx + c))) \right)$$

input `int(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2), x)`

output `-2/45/d*(b*sec(d*x+c))^(1/2)*b^2*(I*(-21*cos(d*x+c)-21)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)+I*(21*cos(d*x+c)+21)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)+(5*cos(d*x+c)^5+2*cos(d*x+c)^3+14*cos(d*x+c)-21)*cot(d*x+c))`

### 3.99.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.11

$$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{21i \sqrt{2} b^{5/2} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 21i \sqrt{2} b^{5/2} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{2}$$

input `integrate(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")`

---

3.99.  $\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx$

output `1/45*(21*I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(5*b^2*cos(d*x + c)^4 + 7*b^2*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c))/d`

### 3.99.6 Sympy [F(-1)]

Timed out.

$$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**7*(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

### 3.99.7 Maxima [F]

$$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^7 dx$$

input `integrate(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^7, x)`

### 3.99.8 Giac [F]

$$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \cos(dx + c)^7 dx$$

input `integrate(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^7, x)`



**3.99.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx = \int \cos(c + dx)^7 \left( \frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2),x)`output `int(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2), x)`

### 3.100 $\int (b \sec(c + dx))^{7/2} dx$

3.100.1 Optimal result . . . . .	645
3.100.2 Mathematica [A] (verified) . . . . .	645
3.100.3 Rubi [A] (verified) . . . . .	646
3.100.4 Maple [C] (verified) . . . . .	648
3.100.5 Fricas [C] (verification not implemented) . . . . .	648
3.100.6 Sympy [F(-1)] . . . . .	649
3.100.7 Maxima [F] . . . . .	649
3.100.8 Giac [F] . . . . .	649
3.100.9 Mupad [F(-1)] . . . . .	650

#### 3.100.1 Optimal result

Integrand size = 12, antiderivative size = 98

$$\int (b \sec(c + dx))^{7/2} dx = -\frac{6b^4 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{6b^3 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2b(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d}$$

```
output 2/5*b*(b*sec(d*x+c))^(5/2)*sin(d*x+c)/d-6/5*b^4*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+6/5*b^3*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d
```

#### 3.100.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.63

$$\int (b \sec(c + dx))^{7/2} dx = \frac{b(b \sec(c + dx))^{5/2} \left( -12 \cos^{5/2}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 7 \sin(c + dx) + 3 \sin(3(c + dx)) \right)}{10d}$$

```
input Integrate[(b*Sec[c + d*x])^(7/2),x]
```

```
output (b*(b*Sec[c + d*x])^(5/2)*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*d)
```

**3.100.3 Rubi [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( b \csc \left( c + dx + \frac{\pi}{2} \right) \right)^{7/2} dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{5} b^2 \int (b \sec(c + dx))^{3/2} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} b^2 \int \left( b \csc \left( c + dx + \frac{\pi}{2} \right) \right)^{3/2} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{5} b^2 \left( \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \right) + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{5} b^2 \left( \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc \left( c + dx + \frac{\pi}{2} \right)}} dx \right) + \\
 & \quad \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \\
 & \quad \downarrow \text{4258} \\
 & \frac{3}{5} b^2 \left( \frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) + \\
 & \quad \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}$$

↓ 3119

$$\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}$$

input `Int[(b*Sec[c + d*x])^(7/2),x]`

output `(2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5`

### 3.100.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.100.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.11 (sec) , antiderivative size = 415, normalized size of antiderivative = 4.23

method	result
default	$2\sqrt{b\sec(dx+c)}b^3\left(3i\operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)^2-3i\operatorname{EllipticE}(i(-\cot(dx+c))\right)$

input `int((b*sec(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 2/5/d*(b*\sec(d*x+c))^{(1/2)}*b^3/(\cos(d*x+c)+1)*(3*I*\operatorname{EllipticF}(I*(-\cot(d*x+c) \\ & )+\csc(d*x+c)),I)*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)} \\ & )*\cos(d*x+c)^2-3*I*\operatorname{EllipticE}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*(1/(\cos(d*x+c)+ \\ & 1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\cos(d*x+c)^2+6*I*\operatorname{EllipticF}(I*( \\ & -\cot(d*x+c)+\csc(d*x+c)),I)*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c) \\ & )+1))^{(1/2)}*\cos(d*x+c)-6*I*\operatorname{EllipticE}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*(1/(\cos \\ & (d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\cos(d*x+c)+3*I*(1/(\cos \\ & (d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(d*x+ \\ & c)+\csc(d*x+c)),I)-3*I*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1)) \\ & )^{(1/2)}*\operatorname{EllipticE}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+3*\sin(d*x+c)+\tan(d*x+c)+\sec \\ & (d*x+c)*\tan(d*x+c) \end{aligned}$$

### 3.100.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.28

$$\int (b\sec(c + dx))^{7/2} dx = \frac{-3i\sqrt{2}b^{7/2}\cos(dx+c)^2\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c)))}{\cos(dx+c)+1}$$

input `integrate((b*sec(d*x+c))^(7/2),x, algorithm="fricas")`

output `1/5*(-3*I*sqrt(2)*b^(7/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*b^(7/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*b^3*cos(d*x + c)^2 + b^3)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^2)`

### 3.100.6 Sympy [F(-1)]

Timed out.

$$\int (b \sec(c + dx))^{7/2} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(7/2),x)`

output `Timed out`

### 3.100.7 Maxima [F]

$$\int (b \sec(c + dx))^{7/2} dx = \int (b \sec(dx + c))^{\frac{7}{2}} dx$$

input `integrate((b*sec(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(7/2), x)`

### 3.100.8 Giac [F]

$$\int (b \sec(c + dx))^{7/2} dx = \int (b \sec(dx + c))^{\frac{7}{2}} dx$$

input `integrate((b*sec(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(7/2), x)`

**3.100.9 Mupad [F(-1)]**

Timed out.

$$\int (b \sec(c + dx))^{7/2} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{7/2} dx$$

input `int((b/cos(c + d*x))^(7/2),x)`output `int((b/cos(c + d*x))^(7/2), x)`

### 3.101 $\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.101.1 Optimal result . . . . .	651
3.101.2 Mathematica [A] (verified) . . . . .	651
3.101.3 Rubi [A] (verified) . . . . .	652
3.101.4 Maple [C] (verified) . . . . .	654
3.101.5 Fracas [C] (verification not implemented) . . . . .	654
3.101.6 Sympy [F] . . . . .	655
3.101.7 Maxima [F] . . . . .	655
3.101.8 Giac [F] . . . . .	655
3.101.9 Mupad [F(-1)] . . . . .	656

#### 3.101.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{21bd} + \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^2d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^4d}$$

output `10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^2/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^4/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b/d`

#### 3.101.2 Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.69

$$\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\sec^3(c+dx) \left( 10 \cos^{\frac{5}{2}}(c+dx) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 5 \sin(2(c+dx)) + 6 \tan(c+dx) \right)}{21d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^5/Sqrt[b*Sec[c + d*x]],x]`



output  $(\text{Sec}[c + d*x]^3*(10*\text{Cos}[c + d*x]^{(5/2)}*\text{EllipticF}[(c + d*x)/2, 2] + 5*\text{Sin}[2*(c + d*x)] + 6*\text{Tan}[c + d*x]))/(21*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

### 3.101.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.08, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{9/2} dx}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx + \frac{\pi}{2}))^{9/2} dx}{b^5} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{5}{7} b^2 \int (b \sec(c+dx))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{5}{7} b^2 \int (b \csc(c+dx + \frac{\pi}{2}))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^5} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^5} \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

---

3.101.  $\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^5}$$

↓ 3042

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^5}$$

↓ 3120

$$\frac{\frac{5}{7}b^2 \left( \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^5}$$

input `Int[Sec[c + d*x]^5/Sqrt[b*Sec[c + d*x]],x]`

output `((2*b*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*d) + (5*b^2*((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d)))/7)/b^5`

### 3.101.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx.)*(v.)^(m.)*((b.)*(v.))^(n.), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c.) + (d.)*(x.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c.) + (d.)*(x.)]*(b.))^(n.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Simp[b^2*((n-2)/(n-1)) Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)]^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.101.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.63

method	result
default	$-\frac{2\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)\right)}{21d\sqrt{b}\sec(dx+c)}$

input `int(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/21/d/(b*sec(d*x+c))^(1/2)*(5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-5*sec(d*x+c)*tan(d*x+c)-3*tan(d*x+c)*sec(d*x+c)^3)`

### 3.101.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.17

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{-5i\sqrt{2}\sqrt{b}\cos(dx+c)^3\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+5i\sqrt{2}\sqrt{b}\cos(dx+c)^3}{21bd\cos(dx+c)}$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/21*(-5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(5*cos(d*x + c)^2 + 3)*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^3)`

3.101.  $\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

**3.101.6 Sympy [F]**

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec^5(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate(sec(d*x+c)**5/(b*sec(d*x+c))**(1/2),x)`

output `Integral(sec(c + d*x)**5/sqrt(b*sec(c + d*x)), x)`

**3.101.7 Maxima [F]**

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^5}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^5/sqrt(b*sec(d*x + c)), x)`

**3.101.8 Giac [F]**

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^5}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^5/sqrt(b*sec(d*x + c)), x)`

**3.101.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{1}{\cos(c+dx)^5 \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2)),x)`output `int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2)), x)`

### 3.102 $\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.102.1 Optimal result . . . . .	657
3.102.2 Mathematica [A] (verified) . . . . .	657
3.102.3 Rubi [A] (verified) . . . . .	658
3.102.4 Maple [C] (verified) . . . . .	660
3.102.5 Fricas [C] (verification not implemented) . . . . .	660
3.102.6 Sympy [F] . . . . .	661
3.102.7 Maxima [F] . . . . .	661
3.102.8 Giac [F] . . . . .	661
3.102.9 Mupad [F(-1)] . . . . .	662

#### 3.102.1 Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx = -\frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5bd} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^3d}$$

output  $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/b^3/d-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/b/d$

#### 3.102.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.63

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{-\frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right)}{\sqrt{\cos(c+dx)}} + 2(3 + \sec^2(c+dx)) \tan(c+dx)}{5d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^4/Sqrt[b*Sec[c + d*x]],x]`

output  $((-6*\text{EllipticE}[(c+d*x)/2, 2])/Sqrt[Cos[c+d*x]] + 2*(3 + Sec[c+d*x]^2)*Tan[c+d*x])/(5*d*Sqrt[b*Sec[c+d*x]])$

**3.102.3 Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.07, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{7/2} dx}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{7/2} dx}{b^4} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \int (b \sec(c+dx))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^4} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx)}{d} \sqrt{b \sec(c+dx)} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx)}{d} \sqrt{b \sec(c+dx)} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^4} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx)}{d} \sqrt{b \sec(c+dx)} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^4} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}}{b^4}$$

↓ 3119

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c+dx)|2\right)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}}{b^4}$$

input `Int[Sec[c + d*x]^4/Sqrt[b*Sec[c + d*x]],x]`

output `((2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5)/b^4`

### 3.102.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`



### 3.102.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 417, normalized size of antiderivative = 4.30

method	result
default	$\frac{6i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - 6i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{5}$

input `int(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2), x, method=_RETURNVERBOSE)`

output

```
2/5/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)+3*tan(d*x+c)+sec(d*x+c)*tan(d*x+c)+tan(d*x+c)*sec(d*x+c)^2)
```

### 3.102.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.27

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$


---


$$-3i \sqrt{2} \sqrt{b} \cos(dx+c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c)))$$


---

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")`

output `1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*cos(d*x + c)^2 + 1)*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^2)`

### 3.102.6 Sympy [F]

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate(sec(d*x+c)**4/(b*sec(d*x+c))**(1/2),x)`

output `Integral(sec(c + d*x)**4/sqrt(b*sec(c + d*x)), x)`

### 3.102.7 Maxima [F]

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^4}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^4/sqrt(b*sec(d*x + c)), x)`

### 3.102.8 Giac [F]

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^4}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^4/sqrt(b*sec(d*x + c)), x)`

**3.102.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{1}{\cos(c+dx)^4 \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2)),x)`output `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2)), x)`

### 3.103 $\int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.103.1 Optimal result . . . . .	663
3.103.2 Mathematica [A] (verified) . . . . .	663
3.103.3 Rubi [A] (verified) . . . . .	664
3.103.4 Maple [C] (verified) . . . . .	665
3.103.5 Fricas [C] (verification not implemented) . . . . .	666
3.103.6 Sympy [F] . . . . .	666
3.103.7 Maxima [F] . . . . .	667
3.103.8 Giac [F] . . . . .	667
3.103.9 Mupad [F(-1)] . . . . .	667

#### 3.103.1 Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3bd} + \frac{2(b \sec(c+dx))^{3/2} \sin(c+dx)}{3b^2d}$$

```
output 2/3*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^2/d+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)
*(b*sec(d*x+c))^(1/2)/b/d
```

#### 3.103.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{2\sqrt{b \sec(c+dx)} \left( \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \tan(c+dx) \right)}{3bd}$$

```
input Integrate[Sec[c + d*x]^3/Sqrt[b*Sec[c + d*x]],x]
```

```
output (2*Sqrt[b*Sec[c + d*x]]*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*b*d)
```

**3.103.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{5/2} dx}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{5/2} dx}{b^3} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{1}{3}b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3}b^2 \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^3} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^3} \\
 & \quad \downarrow \text{3120} \\
 & \frac{\frac{2b^2 \sqrt{\cos(c+dx)} \text{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^3}
 \end{aligned}$$

input `Int[Sec[c + d*x]^3/Sqrt[b*Sec[c + d*x]], x]`

3.103.  $\int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

```
output ((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])
/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d))/b^3
```

### 3.103.3.1 Defintions of rubi rules used

```
rule 2030 Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)
^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 4255 Int[(csc[(c_.) + (d_)*(x_)])*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_.) + (d_)*(x_)])*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

### 3.103.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 147, normalized size of antiderivative = 2.04

method	result
default	$-\frac{2\left(i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)+i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\right)}{3d\sqrt{b}\sec(dx+c)}$

```
input int(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

$$3.103. \quad \int \frac{\sec^3(c+dx)}{\sqrt{b\sec(c+dx)}} dx$$

output 
$$-2/3/d/(b*\sec(d*x+c))^{1/2}*(I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*EllipticF(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*EllipticF(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*\sec(d*x+c)-\sec(d*x+c)*\tan(d*x+c))$$

### 3.103.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.08 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.40

$$\int \frac{\sec^3(c+dx)}{\sqrt{b\sec(c+dx)}} dx$$

$$= \frac{-i\sqrt{2}\sqrt{b}\cos(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+i\sqrt{2}\sqrt{b}\cos(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))+2\sqrt{b/\cos(dx+c)}\sin(dx+c)}{3bd\cos(dx+c)}$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output 
$$1/3*(-I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))+2*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c))/(b*d*\cos(d*x+c))$$

### 3.103.6 SymPy [F]

$$\int \frac{\sec^3(c+dx)}{\sqrt{b\sec(c+dx)}} dx = \int \frac{\sec^3(c+dx)}{\sqrt{b\sec(c+dx)}} dx$$

input `integrate(sec(d*x+c)**3/(b*sec(d*x+c))**(1/2),x)`

output `Integral(sec(c+d*x)**3/sqrt(b*sec(c+d*x)),x)`

**3.103.7 Maxima [F]**

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^3}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^3/sqrt(b*sec(d*x + c)), x)`

**3.103.8 Giac [F]**

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^3}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^3/sqrt(b*sec(d*x + c)), x)`

**3.103.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^3 \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2)),x)`

output `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2)), x)`



### 3.104 $\int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.104.1 Optimal result	668
3.104.2 Mathematica [A] (verified)	668
3.104.3 Rubi [A] (verified)	669
3.104.4 Maple [C] (verified)	670
3.104.5 Fricas [C] (verification not implemented)	671
3.104.6 Sympy [F]	672
3.104.7 Maxima [F]	672
3.104.8 Giac [F]	672
3.104.9 Mupad [F(-1)]	673

#### 3.104.1 Optimal result

Integrand size = 21, antiderivative size = 65

$$\int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx = -\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{2\sqrt{b \sec(c+dx)} \sin(c+dx)}{bd}$$

output `-2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/b/d`

#### 3.104.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.74

$$\int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{-\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{\sqrt{\cos(c+dx)}} + 2 \tan(c+dx)}{d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^2/Sqrt[b*Sec[c + d*x]], x]`

output `((-2*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*Tan[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])`

**3.104.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{3/2} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx}{b^2} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{b^2} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^2} \\
 & \quad \downarrow \text{3119} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^2}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2/Sqrt[b*Sec[c + d*x]],x]`

output `((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d)/b^2`

### 3.104.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.104.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 389, normalized size of antiderivative = 5.98

method	result
default	$-\frac{2\left(i\operatorname{EllipticE}\left(i\left(-\cot(dx+c)+\csc(dx+c)\right),i\right)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)-i\operatorname{EllipticF}\left(i\left(-\cot(dx+c)+\csc(dx+c)\right),i\right)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\right)}{b^2}$

3.104. 
$$\int \frac{\sec^2(c+dx)}{\sqrt{b\sec(c+dx)}} dx$$

```
input int(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(I*EllipticE(I*(-cot(d*x+c)+csc(d
*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d
*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Ellipti
cF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+2*I*(1/(cos(d*x+c)+1))^(1/2)*(
cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-2
*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*
(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*
x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-I*(1/(co
s(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x
+c)+csc(d*x+c)),I)*sec(d*x+c)-tan(d*x+c))
```

### 3.104.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.32

$$\int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c))) + i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i \sin(dx+c)))}{bd}$$

```
input integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output (-I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(
d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, wei
erstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b/cos(d*x
+ c))*sin(d*x + c)/(b*d)
```

**3.104.6 Sympy [F]**

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(1/2),x)`

output `Integral(sec(c + d*x)**2/sqrt(b*sec(c + d*x)), x)`

**3.104.7 Maxima [F]**

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^2/sqrt(b*sec(d*x + c)), x)`

**3.104.8 Giac [F]**

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^2/sqrt(b*sec(d*x + c)), x)`

**3.104.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{1}{\cos(c+dx)^2 \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2)),x)`output `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2)), x)`

### 3.105 $\int \frac{\sec(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.105.1 Optimal result . . . . .	674
3.105.2 Mathematica [A] (verified) . . . . .	674
3.105.3 Rubi [A] (verified) . . . . .	675
3.105.4 Maple [C] (verified) . . . . .	676
3.105.5 Fracas [C] (verification not implemented) . . . . .	677
3.105.6 Sympy [F] . . . . .	677
3.105.7 Maxima [F] . . . . .	677
3.105.8 Giac [F] . . . . .	678
3.105.9 Mupad [F(-1)] . . . . .	678

#### 3.105.1 Optimal result

Integrand size = 19, antiderivative size = 41

$$\int \frac{\sec(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{bd}$$

output `2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b/d`

#### 3.105.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\sec(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{bd}$$

input `Integrate[Sec[c + d*x]/Sqrt[b*Sec[c + d*x]],x]`

output `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b*d)`

**3.105.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2030, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int \sqrt{b \sec(c+dx)} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{b} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{bd}
 \end{aligned}$$

input `Int[Sec[c + d*x]/Sqrt[b*Sec[c + d*x]],x]`

output `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b*d)`



## 3.105.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.105.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.68

method	result	size
default	$-\frac{2i\sqrt{\frac{1}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)), i)}{d\sqrt{b\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}$	69

input `int(sec(d*x+c)/(b*sec(d*x+c))^(1/2), x, method=_RETURNVERBOSE)`

output `-2*I/d*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)/(b*sec(d*x+c))^(1/2)/(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)`

**3.105.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \frac{\sec(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{bd}$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b*d)`

**3.105.6 Sympy [F]**

$$\int \frac{\sec(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))**(1/2),x)`

output `Integral(sec(c + d*x)/sqrt(b*sec(c + d*x)), x)`

**3.105.7 Maxima [F]**

$$\int \frac{\sec(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)/sqrt(b*sec(d*x + c)), x)`

**3.105.8 Giac [F]**

$$\int \frac{\sec(c+dx)}{\sqrt{b\sec(c+dx)}} dx = \int \frac{\sec(dx+c)}{\sqrt{b\sec(dx+c)}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)/sqrt(b*sec(d*x + c)), x)`

**3.105.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec(c+dx)}{\sqrt{b\sec(c+dx)}} dx = \int \frac{1}{\cos(c+dx) \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/2)),x)`

output `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/2)), x)`

### 3.106 $\int \frac{1}{\sqrt{b \sec(c+dx)}} dx$

3.106.1 Optimal result	679
3.106.2 Mathematica [A] (verified)	679
3.106.3 Rubi [A] (verified)	680
3.106.4 Maple [C] (verified)	681
3.106.5 Fricas [C] (verification not implemented)	682
3.106.6 Sympy [F]	682
3.106.7 Maxima [F]	682
3.106.8 Giac [F]	683
3.106.9 Mupad [F(-1)]	683

#### 3.106.1 Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \frac{1}{\sqrt{b \sec(c+dx)}} dx = \frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

output `2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`

#### 3.106.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{b \sec(c+dx)}} dx = \frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

input `Integrate[1/Sqrt[b*Sec[c + d*x]],x]`

output `(2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

**3.106.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \\ & \quad \downarrow \text{4258} \\ & \frac{\int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\ & \quad \downarrow \text{3119} \\ & \frac{2E(\frac{1}{2}(c+dx)|2)}{d\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \end{aligned}$$

input `Int[1/Sqrt[b*Sec[c + d*x]],x]`

output `(2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

### 3.106.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.106.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.60 (sec) , antiderivative size = 299, normalized size of antiderivative = 7.87

method	result
risch	$-\frac{i\sqrt{2}}{d\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}} - \frac{i\left(-\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}\left(-2i\text{EllipticE}\left(\sqrt{-i(e^{i(dx+c)}}\right)}\right)}{\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}}\right)}{d\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}(e^{2i(dx+c)}+1)}$
default	$\frac{2i\text{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)-2i\text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}}{\dots}$

input `int(1/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-I/d*2^(1/2)/(b*exp(I*(d*x+c))/(exp(I*(d*x+c))^2+1))^(1/2)-I/d*(-2*(b*exp(I*(d*x+c))^2+b)/b/(exp(I*(d*x+c))*(b*exp(I*(d*x+c))^2+b))^(1/2)+I*(-I*(exp(I*(d*x+c))+I))^(1/2)*2^(1/2)*(I*(exp(I*(d*x+c))-I))^(1/2)*(I*exp(I*(d*x+c)))^(1/2)/(b*exp(I*(d*x+c))^3+b*exp(I*(d*x+c)))^(1/2)*(-2*I*EllipticE((-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))+I*EllipticF((-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))))*2^(1/2)/(b*exp(I*(d*x+c))/(exp(I*(d*x+c))^2+1))^(1/2)*(b*exp(I*(d*x+c))*(exp(I*(d*x+c))^2+1))^(1/2)/(exp(I*(d*x+c))^2+1)`

**3.106.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.74

$$\int \frac{1}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{bd}$$

input `integrate(1/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `(I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b*d)`

**3.106.6 Sympy [F]**

$$\int \frac{1}{\sqrt{b \sec(c + dx)}} dx = \int \frac{1}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate(1/(b*sec(d*x+c))**(1/2),x)`

output `Integral(1/sqrt(b*sec(c + d*x)), x)`

**3.106.7 Maxima [F]**

$$\int \frac{1}{\sqrt{b \sec(c + dx)}} dx = \int \frac{1}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(1/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*sec(d*x + c)), x)`

**3.106.8 Giac [F]**

$$\int \frac{1}{\sqrt{b \sec(c + dx)}} dx = \int \frac{1}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(1/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(b*sec(d*x + c)), x)`

**3.106.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{b \sec(c + dx)}} dx = \int \frac{1}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(1/(b/cos(c + d*x))^(1/2),x)`

output `int(1/(b/cos(c + d*x))^(1/2), x)`



### 3.107 $\int \frac{\cos(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.107.1 Optimal result . . . . .	684
3.107.2 Mathematica [A] (verified) . . . . .	684
3.107.3 Rubi [A] (verified) . . . . .	685
3.107.4 Maple [C] (verified) . . . . .	687
3.107.5 Fracas [C] (verification not implemented) . . . . .	687
3.107.6 Sympy [F] . . . . .	688
3.107.7 Maxima [F] . . . . .	688
3.107.8 Giac [F] . . . . .	688
3.107.9 Mupad [F(-1)] . . . . .	689

#### 3.107.1 Optimal result

Integrand size = 19, antiderivative size = 69

$$\int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3bd} + \frac{2 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}}$$

output `2/3*sin(d*x+c)/d/(b*sec(d*x+c))^(1/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b/d`

#### 3.107.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.87

$$\int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{b \sec^2(c + dx) \left( 2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx)) \right)}{3d(b \sec(c + dx))^{3/2}}$$

input `Integrate[Cos[c + d*x]/Sqrt[b*Sec[c + d*x]],x]`

output  $(b \sec[c + d*x]^2 (2 \sqrt{\cos[c + d*x]} \text{EllipticF}[(c + d*x)/2, 2] + \sin[2*(c + d*x)]) / (3*d*(b \sec[c + d*x])^{3/2})$

### 3.107.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{1}{\csc(c + dx + \frac{\pi}{2}) \sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow 2030 \\
 & b \int \frac{1}{(b \csc(\frac{1}{2}(2c + \pi) + dx))^{3/2}} dx \\
 & \quad \downarrow 4256 \\
 & b \left( \frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right) \\
 & \quad \downarrow 3042 \\
 & b \left( \frac{\int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right) \\
 & \quad \downarrow 4258 \\
 & b \left( \frac{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right) \\
 & \quad \downarrow 3042 \\
 & b \left( \frac{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right)
 \end{aligned}$$

$$b \left( \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)$$

input `Int[Cos[c + d*x]/Sqrt[b*Sec[c + d*x]],x]`

output `b*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]]))`

### 3.107.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d*n)), x] + Simp[(n+1)/(b^2*n) Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.107.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 141, normalized size of antiderivative = 2.04

method	result
default	$-\frac{2\left(i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)+i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)\right)}{3d\sqrt{b}\sec(dx+c)}$

input `int(cos(d*x+c)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/3/d/(b*sec(d*x+c))^(1/2)*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-sin(d*x+c))`

### 3.107.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.26

$$\int \frac{\cos(c+dx)}{\sqrt{b}\sec(c+dx)} dx$$

$$= \frac{2\sqrt{\frac{b}{\cos(dx+c)}}\cos(dx+c)\sin(dx+c) - i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c)) + i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))}{3bd}$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/3*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)*sin(d*x + c) - I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b*d)`

**3.107.6 Sympy [F]**

$$\int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))**(1/2),x)`

output `Integral(cos(c + d*x)/sqrt(b*sec(c + d*x)), x)`

**3.107.7 Maxima [F]**

$$\int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(cos(d*x + c)/sqrt(b*sec(d*x + c)), x)`

**3.107.8 Giac [F]**

$$\int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(cos(d*x + c)/sqrt(b*sec(d*x + c)), x)`

**3.107.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\cos(c+dx)}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(cos(c + d*x)/(b/cos(c + d*x))^(1/2), x)`output `int(cos(c + d*x)/(b/cos(c + d*x))^(1/2), x)`

### 3.108 $\int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.108.1 Optimal result . . . . .	690
3.108.2 Mathematica [A] (verified) . . . . .	690
3.108.3 Rubi [A] (verified) . . . . .	691
3.108.4 Maple [C] (verified) . . . . .	692
3.108.5 Fricas [C] (verification not implemented) . . . . .	693
3.108.6 Sympy [F] . . . . .	694
3.108.7 Maxima [F] . . . . .	694
3.108.8 Giac [F] . . . . .	694
3.108.9 Mupad [F(-1)] . . . . .	695

#### 3.108.1 Optimal result

Integrand size = 21, antiderivative size = 67

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{6E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}}$$

```
output 2/5*b*sin(d*x+c)/d/(b*sec(d*x+c))^(3/2)+6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/c
os(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)
/(b*sec(d*x+c))^(1/2)
```

#### 3.108.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.90

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{\sqrt{b \sec(c + dx)} \left( 12\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) + \sin(3(c + dx)) \right)}{10bd}$$

```
input Integrate[Cos[c + d*x]^2/Sqrt[b*Sec[c + d*x]],x]
```

```
output (Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + S
in[c + d*x] + Sin[3*(c + d*x)]))/(10*b*d)
```

**3.108.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.13, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^2 \sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^2 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b^2 \left( \frac{3 \int \frac{\sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3119} \\
 & b^2 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)
 \end{aligned}$$



input `Int[Cos[c + d*x]^2/Sqrt[b*Sec[c + d*x]],x]`

output `b^2*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2)))`

### 3.108.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.108.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 417, normalized size of antiderivative = 6.22

method	result
default	$\frac{6i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c) - 6i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)}{5}$

3.108.  $\int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

input `int(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `2/5/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(3*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-3*I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+cos(d*x+c)^2*sin(d*x+c)+cos(d*x+c)*sin(d*x+c)+3*sin(d*x+c))`

### 3.108.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.42

$$\int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$


---


$$= 2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^2 \sin(dx+c) + 3i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c)))$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="fracas")`

output `1/5*(2*sqrt(b/cos(d*x+c))*cos(d*x+c)^2*sin(d*x+c)+3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4,0,weierstrassPInverse(-4,0,cos(d*x+c))+I*sin(d*x+c))-3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4,0,weierstrassPInverse(-4,0,cos(d*x+c)-I*sin(d*x+c))))/(b*d)`

**3.108.6 Sympy [F]**

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(1/2),x)`

output `Integral(cos(c + d*x)**2/sqrt(b*sec(c + d*x)), x)`

**3.108.7 Maxima [F]**

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(cos(d*x + c)^2/sqrt(b*sec(d*x + c)), x)`

**3.108.8 Giac [F]**

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(cos(d*x + c)^2/sqrt(b*sec(d*x + c)), x)`

**3.108.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\cos(c+dx)^2}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/2),x)`output `int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/2), x)`

### 3.109 $\int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.109.1 Optimal result . . . . .	696
3.109.2 Mathematica [A] (verified) . . . . .	696
3.109.3 Rubi [A] (verified) . . . . .	697
3.109.4 Maple [C] (verified) . . . . .	699
3.109.5 Fracas [C] (verification not implemented) . . . . .	700
3.109.6 Sympy [F(-1)] . . . . .	700
3.109.7 Maxima [F] . . . . .	700
3.109.8 Giac [F] . . . . .	701
3.109.9 Mupad [F(-1)] . . . . .	701

#### 3.109.1 Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{21bd} + \frac{2b^2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21d\sqrt{b \sec(c+dx)}}$$

output `2/7*b^2*sin(d*x+c)/d/(b*sec(d*x+c))^(5/2)+10/21*sin(d*x+c)/d/(b*sec(d*x+c))^(1/2)+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b/d`

#### 3.109.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.68

$$\int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\sqrt{b \sec(c+dx)} \left( 40\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 26 \sin(2(c+dx)) + 3 \sin(4(c+dx)) \right)}{84bd}$$

input `Integrate[Cos[c + d*x]^3/Sqrt[b*Sec[c + d*x]],x]`

output  $(\text{Sqrt}[b \cdot \text{Sec}[c + d \cdot x]] \cdot (40 \cdot \text{Sqrt}[\text{Cos}[c + d \cdot x]] \cdot \text{EllipticF}[(c + d \cdot x)/2, 2] + 2 \cdot 6 \cdot \text{Sin}[2 \cdot (c + d \cdot x)] + 3 \cdot \text{Sin}[4 \cdot (c + d \cdot x)])) / (84 \cdot b \cdot d)$

### 3.109.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.15, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^3 \sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^3 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^3 \left( \frac{5 \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{5 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^3 \left( \frac{5 \left( \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& b^3 \left( \frac{5 \left( \frac{\int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 4258 \\
& b^3 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 3042 \\
& b^3 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow 3120 \\
& b^3 \left( \frac{5 \left( \frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)
\end{aligned}$$

input `Int[Cos[c + d*x]^3/Sqrt[b*Sec[c + d*x]],x]`

output `b^3*((2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/(7*b^2))`

### 3.109.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx._)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.109.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.65 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.62

method	result
default	$-\frac{2\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\right)}{21d\sqrt{b\sec(dx+c)}}$

input `int(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/21/d/(b*sec(d*x+c))^(1/2)*(5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*cos(d*x+c)^2*sin(d*x+c)-5*sin(d*x+c))`



**3.109.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.03

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{2(3 \cos(dx + c)^3 + 5 \cos(dx + c)) \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c) - 5i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c))}{21bd}$$

input `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/21*(2*(3*cos(d*x + c)^3 + 5*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c) - 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b*d)`

**3.109.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3/(b*sec(d*x+c))**(1/2),x)`

output `Timed out`

**3.109.7 Maxima [F]**

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)^3}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(cos(d*x + c)^3/sqrt(b*sec(d*x + c)), x)`

---

3.109.  $\int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

**3.109.8 Giac [F]**

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)^3}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(cos(d*x + c)^3/sqrt(b*sec(d*x + c)), x)`

**3.109.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(c + dx)^3}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(cos(c + d*x)^3/(b/cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^3/(b/cos(c + d*x))^(1/2), x)`

### 3.110 $\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.110.1 Optimal result . . . . .	702
3.110.2 Mathematica [A] (verified) . . . . .	702
3.110.3 Rubi [A] (verified) . . . . .	703
3.110.4 Maple [C] (verified) . . . . .	705
3.110.5 Fricas [C] (verification not implemented) . . . . .	706
3.110.6 Sympy [F] . . . . .	706
3.110.7 Maxima [F] . . . . .	706
3.110.8 Giac [F] . . . . .	707
3.110.9 Mupad [F(-1)] . . . . .	707

#### 3.110.1 Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{14E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{2b^3 \sin(c+dx)}{9d(b \sec(c+dx))^{7/2}} + \frac{14b \sin(c+dx)}{45d(b \sec(c+dx))^{3/2}}$$

output  $2/9*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^(7/2)+14/45*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^(3/2)+14/15*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)$

#### 3.110.2 Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.74

$$\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{336E\left(\frac{1}{2}(c+dx) \mid 2\right)}{\sqrt{\cos(c+dx)}} + \frac{4 \cos(c+dx)(33 \sin(c+dx) + 5 \sin(3(c+dx)))}{360d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Cos[c + d*x]^4/Sqrt[b*Sec[c + d*x]],x]`

output  $((336*\text{EllipticE}[(c+d*x)/2, 2])/Sqrt[\text{Cos}[c+d*x]] + 4*\text{Cos}[c+d*x]*(33*\text{Sin}[c+d*x] + 5*\text{Sin}[3*(c+d*x)]))/(360*d*Sqrt[b*\text{Sec}[c+d*x]])$

**3.110.3 Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^4 \sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^4 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{9/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^4 \left( \frac{7 \int \frac{1}{(b \sec(c+dx))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{7 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^4 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^4 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 4258 \\
 b^4 \left( \frac{7 \left( \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 \downarrow 3042 \\
 b^4 \left( \frac{7 \left( \frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 \downarrow 3119 \\
 b^4 \left( \frac{7 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)
 \end{array}$$

input `Int[Cos[c + d*x]^4/Sqrt[b*Sec[c + d*x]],x]`

output `b^4*((2*Sin[c + d*x])/(9*b*d*(b*Sec[c + d*x])^(7/2)) + (7*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))))/(9*b^2)`

### 3.110.3.1 Defintions of rubi rules used

rule 2030 `Int[(F*_)(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F_x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 4256 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_], x_Symbol] := Simp[Cos[c + d*x]*((
b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c
+ d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_], x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

### 3.110.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 451, normalized size of antiderivative = 4.75

method	result
default	$-\frac{2(21i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - 21i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)))$

```
input int(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2), x, method=_RETURNVERBOSE)
```

```
output -2/45/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(21*I*(1/(cos(d*x+c)+1))^(1/2)
*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)
*cos(d*x+c)-21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)
)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)-5*cos(d*x+c)^4*sin(d*
x+c)+42*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Ellip
ticF(I*(-cot(d*x+c)+csc(d*x+c)), I)-42*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+
c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)-5*cos(d*x
+c)^3*sin(d*x+c)+21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))
^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)-21*I*(1/(cos(d*x
+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+c
sc(d*x+c)), I)*sec(d*x+c)-7*cos(d*x+c)^2*sin(d*x+c)-7*cos(d*x+c)*sin(d*x+c)
-21*sin(d*x+c))
```

**3.110.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.14

$$\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$


---


$$= \frac{2(5 \cos(dx+c)^4 + 7 \cos(dx+c)^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c) + 21i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + I \sin(dx+c))) - 21i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - I \sin(dx+c)))}{(b*d)}$$

input `integrate(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/45*(2*(5*cos(d*x + c)^4 + 7*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c) + 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b*d)`

**3.110.6 Sympy [F]**

$$\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

input `integrate(cos(d*x+c)**4/(b*sec(d*x+c))**(1/2),x)`

output `Integral(cos(c + d*x)**4/sqrt(b*sec(c + d*x)), x)`

**3.110.7 Maxima [F]**

$$\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\cos^4(dx+c)}{\sqrt{b \sec(dx+c)}} dx$$

input `integrate(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(cos(d*x + c)^4/sqrt(b*sec(d*x + c)), x)`

---

3.110.  $\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

**3.110.8 Giac [F]**

$$\int \frac{\cos^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)^4}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(cos(d*x + c)^4/sqrt(b*sec(d*x + c)), x)`

**3.110.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\cos(c + dx)^4}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int(cos(c + d*x)^4/(b/cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^4/(b/cos(c + d*x))^(1/2), x)`



### 3.111 $\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.111.1 Optimal result . . . . .	708
3.111.2 Mathematica [A] (verified) . . . . .	708
3.111.3 Rubi [A] (verified) . . . . .	709
3.111.4 Maple [C] (verified) . . . . .	711
3.111.5 Fricas [C] (verification not implemented) . . . . .	711
3.111.6 Sympy [F] . . . . .	712
3.111.7 Maxima [F] . . . . .	712
3.111.8 Giac [F] . . . . .	712
3.111.9 Mupad [F(-1)] . . . . .	713

#### 3.111.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{21b^2d} + \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^3d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^5d}$$

output `10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^3/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^5/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d`

#### 3.111.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.69

$$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\sec^4(c+dx) \left( 10 \cos^{5/2}(c+dx) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 5 \sin(2(c+dx)) + 6 \tan(c+dx) \right)}{21d(b \sec(c+dx))^{3/2}}$$

input `Integrate[Sec[c + d*x]^6/(b*Sec[c + d*x])^(3/2),x]`

output `(Sec[c + d*x]^4*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d*(b*Sec[c + d*x])^(3/2))`

**3.111.3 Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.08, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{9/2} dx}{b^6} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx + \frac{\pi}{2}))^{9/2} dx}{b^6} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{5}{7} b^2 \int (b \sec(c+dx))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^6} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{5}{7} b^2 \int (b \csc(c+dx + \frac{\pi}{2}))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^6} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^6} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^6} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^6} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.111.  $\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^6}$$

↓ 3120

$$\frac{\frac{5}{7}b^2 \left( \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^6}$$

input `Int[Sec[c + d*x]^6/(b*Sec[c + d*x])^(3/2),x]`

output `((2*b*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*d) + (5*b^2*((2*b^2*Sqrt[Cos[c + d*x])*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d))/7)/b^6`

### 3.111.3.1 Defintions of rubi rules used

rule 2030 `Int[(F*x_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F*x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Simp[b^2*((n-2)/(n-1)) Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.111.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.66

method	result
default	$-\frac{2\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)\right)}{21d\sqrt{b\sec(dx+c)}b}$

input `int(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -2/21/d/(b*\sec(d*x+c))^{(1/2)}/b*(5*I*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\operatorname{csc}(d*x+c)),I)+5*I*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\operatorname{csc}(d*x+c)),I)*\sec(d*x+c)-5*\sec(d*x+c)*\tan(d*x+c)-3*\tan(d*x+c)*\sec(d*x+c)^3) \end{aligned}$$

### 3.111.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.17

$$\int \frac{\sec^6(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{-5i\sqrt{2}\sqrt{b}\cos(dx+c)^3\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))}{(b\sec(c+dx))^{3/2}}$$

input `integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output 
$$\begin{aligned} & 1/21*(-5*I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)^3*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+5*I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)^3*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))+2*(5*\cos(d*x+c)^2+3)*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c))/(b^2*d*\cos(d*x+c)^3) \end{aligned}$$

**3.111.6 Sympy [F]**

$$\int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)**6/(b*sec(d*x+c))**(3/2),x)`

output `Integral(sec(c + d*x)**6/(b*sec(c + d*x))**(3/2), x)`

**3.111.7 Maxima [F]**

$$\int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^6}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(3/2), x)`

**3.111.8 Giac [F]**

$$\int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^6}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(3/2), x)`

**3.111.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^6(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{1}{\cos(c+dx)^6 \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2)),x)`output `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2)), x)`

### 3.112 $\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.112.1 Optimal result . . . . .	714
3.112.2 Mathematica [A] (verified) . . . . .	714
3.112.3 Rubi [A] (verified) . . . . .	715
3.112.4 Maple [C] (verified) . . . . .	717
3.112.5 Fricas [C] (verification not implemented) . . . . .	717
3.112.6 Sympy [F] . . . . .	718
3.112.7 Maxima [F] . . . . .	718
3.112.8 Giac [F] . . . . .	718
3.112.9 Mupad [F(-1)] . . . . .	719

#### 3.112.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx = -\frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5b^2d} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^4d}$$

output `2/5*(b*sec(d*x+c))^(5/2)*sin(d*x+c)/b^4/d-6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+6/5*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/b^2/d`

#### 3.112.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.64

$$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{-\frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right)}{\sqrt{\cos(c+dx)}} + 2(3 + \sec^2(c+dx)) \tan(c+dx)}{5bd\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^5/(b*Sec[c + d*x])^(3/2),x]`

output `((-6*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*(3 + Sec[c + d*x]^2)*Tan[c + d*x])/(5*b*d*Sqrt[b*Sec[c + d*x]])`

**3.112.3 Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{7/2} dx}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{7/2} dx}{b^5} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \int (b \sec(c+dx))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^5} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^5} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^5} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.112.  $\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx$



$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^5}$$

↓ 3119

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^5}$$

input `Int[Sec[c + d*x]^5/(b*Sec[c + d*x])^(3/2), x]`

output `((2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5)/b^5`

### 3.112.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.112.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 420, normalized size of antiderivative = 4.20

method	result
default	$\frac{6i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{5} - \frac{6i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}{5}$

input `int(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output

```
2/5/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b*(3*I*(1/(cos(d*x+c)+1))^(1/2)*
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*
cos(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*
EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1))^(
1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)
)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Ellip
ticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)
)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)
-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(
I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+3*tan(d*x+c)+sec(d*x+c)*tan(d*x+c)
)+tan(d*x+c)*sec(d*x+c)^2)
```

### 3.112.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.23

$$\int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{-3i\sqrt{2}\sqrt{b}\cos(dx+c)^2 \operatorname{weierstrassZeta}(-4,0, \operatorname{weierstrassPInverse}(-4,0, \cos(dx+c)))}{5}$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output

```
1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstras
sPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos
(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c)
- I*sin(d*x + c))) + 2*(3*cos(d*x + c)^2 + 1)*sqrt(b/cos(d*x + c))*sin(d*
x + c))/(b^2*d*cos(d*x + c)^2)
```

---

3.112.  $\int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{3/2}} dx$

**3.112.6 Sympy [F]**

$$\int \frac{\sec^5(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec^5(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)**5/(b*sec(d*x+c))**(3/2),x)`

output `Integral(sec(c + d*x)**5/(b*sec(c + d*x))**(3/2), x)`

**3.112.7 Maxima [F]**

$$\int \frac{\sec^5(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^5}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(3/2), x)`

**3.112.8 Giac [F]**

$$\int \frac{\sec^5(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^5}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(3/2), x)`

**3.112.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{1}{\cos(c+dx)^5 \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2)),x)`output `int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2)), x)`

### 3.113 $\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.113.1 Optimal result . . . . .	720
3.113.2 Mathematica [A] (verified) . . . . .	720
3.113.3 Rubi [A] (verified) . . . . .	721
3.113.4 Maple [C] (verified) . . . . .	722
3.113.5 Fricas [C] (verification not implemented) . . . . .	723
3.113.6 Sympy [F] . . . . .	723
3.113.7 Maxima [F] . . . . .	724
3.113.8 Giac [F] . . . . .	724
3.113.9 Mupad [F(-1)] . . . . .	724

#### 3.113.1 Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2d} + \frac{2(b \sec(c+dx))^{3/2} \sin(c+dx)}{3b^3d}$$

output  $2/3*(b*\sec(d*x+c))^(3/2)*\sin(d*x+c)/b^3/d+2/3*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)*(b*\sec(d*x+c))^(1/2)/b^2/d$

#### 3.113.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.78

$$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2 \sec^3(c+dx) \left( \cos^{3/2}(c+dx) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \sin(c+dx) \right)}{3d(b \sec(c+dx))^{3/2}}$$

input  $\operatorname{Integrate}[\operatorname{Sec}[c + d*x]^4/(b*\operatorname{Sec}[c + d*x])^(3/2), x]$

output  $(2*\operatorname{Sec}[c + d*x]^3*(\operatorname{Cos}[c + d*x]^(3/2)*\operatorname{EllipticF}[(c + d*x)/2, 2] + \operatorname{Sin}[c + d*x]))/(3*d*(b*\operatorname{Sec}[c + d*x])^(3/2))$

**3.113.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{5/2} dx}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{5/2} dx}{b^4} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{1}{3}b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3}b^2 \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^4} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^4} \\
 & \quad \downarrow \text{3120} \\
 & \frac{\frac{2b^2 \sqrt{\cos(c+dx)} \text{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^4}
 \end{aligned}$$

input `Int[Sec[c + d*x]^4/(b*Sec[c + d*x])^(3/2), x]`

$$3.113. \quad \int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

output  $((2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*b*(b*\text{Sec}[c + d*x])^(3/2)*\text{Sin}[c + d*x])/(3*d))/b^4$

### 3.113.3.1 Defintions of rubi rules used

rule 2030  $\text{Int}[(\text{Fx}_.)*(v_.)^(m_.)*((b_.)*(v_.))^(n_.), x\_Symbol] \rightarrow \text{Simp}[1/b^m \text{Int}[(b*v)^(m+n)*\text{Fx}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3120  $\text{Int}[1/\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_.)]], x\_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4255  $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x\_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^(n-1)/(d*(n-1))), x] + \text{Simp}[b^2*((n-2)/(n-1))\text{Int}[(b*\text{Csc}[c + d*x])^(n-2), x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

rule 4258  $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x\_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{EqQ}[n^2, 1/4]$

### 3.113.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.51 (sec) , antiderivative size = 150, normalized size of antiderivative = 2.08

method	result
default	$-\frac{2\left(i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticF}\left(i(-\cot(dx+c)+\text{csc}(dx+c)),i\right)+i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticF}\left(i(-\cot(dx+c)+\text{csc}(dx+c)),i\right)\right)}{3d\sqrt{b}\sec(dx+c)b}$

input  $\text{int}(\sec(d*x+c)^4/(b*\sec(d*x+c))^(3/2), x, \text{method}=\_RETURNVERBOSE)$

$$3.113. \quad \int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{3/2}} dx$$

output 
$$-2/3/d/(b*\sec(d*x+c))^{1/2}/b*(I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*EllipticF(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*EllipticF(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*\sec(d*x+c)-\sec(d*x+c)*\tan(d*x+c))$$

### 3.113.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.40

$$\int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\cos(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+i\sqrt{2}\sqrt{b}\sin(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+2\sqrt{b}\cos(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))+2\sqrt{b}\sin(dx+c)\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))}{(b^2d\cos(dx+c))^{3/2}}$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output 
$$1/3*(-I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))+2*\sqrt{b}*\cos(d*x+c)*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+2*\sqrt{b}*\sin(d*x+c)*\operatorname{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c)))/(b^2*d*\cos(d*x+c))^{3/2}$$

### 3.113.6 Sympy [F]

$$\int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{3/2}} dx$$

input `integrate(sec(d*x+c)**4/(b*sec(d*x+c))**(3/2),x)`

output `Integral(sec(c+d*x)**4/(b*sec(c+d*x))**(3/2),x)`



**3.113.7 Maxima [F]**

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^4}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(3/2), x)`

**3.113.8 Giac [F]**

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^4}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(3/2), x)`

**3.113.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx)^4 \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2)),x)`

output `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2)), x)`

### 3.114 $\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.114.1 Optimal result . . . . .	725
3.114.2 Mathematica [A] (verified) . . . . .	725
3.114.3 Rubi [A] (verified) . . . . .	726
3.114.4 Maple [C] (verified) . . . . .	727
3.114.5 Fracas [C] (verification not implemented) . . . . .	728
3.114.6 Sympy [F] . . . . .	728
3.114.7 Maxima [F] . . . . .	729
3.114.8 Giac [F] . . . . .	729
3.114.9 Mupad [F(-1)] . . . . .	729

#### 3.114.1 Optimal result

Integrand size = 21, antiderivative size = 68

$$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx = -\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{2\sqrt{b \sec(c+dx)} \sin(c+dx)}{b^2d}$$

output `-2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/b^2/d`

#### 3.114.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.75

$$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{-\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{\sqrt{\cos(c+dx)}} + 2 \tan(c+dx)}{bd\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^3/(b*Sec[c + d*x])^(3/2),x]`

output `((-2*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*Tan[c + d*x])/(b*d*Sqrt[b*Sec[c + d*x]])`

**3.114.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{3/2} dx}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx}{b^3} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{b^3} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^3} \\
 & \quad \downarrow \text{3119} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^3}
 \end{aligned}$$

---

3.114.  $\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

input `Int[Sec[c + d*x]^3/(b*Sec[c + d*x])^(3/2),x]`

output `((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d)/b^3`

### 3.114.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.114.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 392, normalized size of antiderivative = 5.76

method	result
default	$-\frac{2 \left( i \operatorname{EllipticE} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - i \operatorname{EllipticF} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \right)}{\dots}$

$$3.114. \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

input `int(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/d/(cos(d*x+c)+1)/b/(b*sec(d*x+c))^(1/2)*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-tan(d*x+c)`

### 3.114.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.26

$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)))}{(b\sec(c+dx))^{3/2}}$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2),x, algorithm="fracas")`

output `(-I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b^2*d)`

### 3.114.6 Sympy [F]

$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)**3/(b*sec(d*x+c))**(3/2),x)`

output `Integral(sec(c + d*x)**3/(b*sec(c + d*x))**(3/2), x)`

---

3.114.  $\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{3/2}} dx$

**3.114.7 Maxima [F]**

$$\int \frac{\sec^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^3}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)`

**3.114.8 Giac [F]**

$$\int \frac{\sec^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^3}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)`

**3.114.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx)^3 \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2)),x)`

output `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2)), x)`

### 3.115 $\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.115.1 Optimal result . . . . .	730
3.115.2 Mathematica [A] (verified) . . . . .	730
3.115.3 Rubi [A] (verified) . . . . .	731
3.115.4 Maple [C] (verified) . . . . .	732
3.115.5 Fracas [C] (verification not implemented) . . . . .	733
3.115.6 Sympy [F] . . . . .	733
3.115.7 Maxima [F] . . . . .	733
3.115.8 Giac [F] . . . . .	734
3.115.9 Mupad [F(-1)] . . . . .	734

#### 3.115.1 Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{b^2 d}$$

output `2*(cos(1/2*d*x+1/2*c)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d`

#### 3.115.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{b^2 d}$$

input `Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(3/2), x]`

output `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b^2*d)`

**3.115.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2030, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int \sqrt{b \sec(c+dx)} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{b^2} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b^2} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{b^2 d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(3/2),x]`

output `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b^2*d)`



## 3.115.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.115.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.76

method	result	size
default	$-\frac{2i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}}}{db \sqrt{b \sec(dx+c)} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}$	72

input `int(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2), x, method=_RETURNVERBOSE)`

output `-2*I/d*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)/b/(b*sec(d*x+c))^(1/2)/(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)`

**3.115.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \frac{\sec^2(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c))}{b^2d}$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2),x, algorithm="fracas")`

output `(-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^2*d)`

**3.115.6 Sympy [F]**

$$\int \frac{\sec^2(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{\sec^2(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(3/2),x)`

output `Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(3/2), x)`

**3.115.7 Maxima [F]**

$$\int \frac{\sec^2(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{\sec(dx+c)^2}{(b\sec(dx+c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(3/2), x)`

**3.115.8 Giac [F]**

$$\int \frac{\sec^2(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{\sec(dx+c)^2}{(b\sec(dx+c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(3/2), x)`

**3.115.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^2(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{1}{\cos(c+dx)^2 \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2)),x)`

output `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2)), x)`

### 3.116 $\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.116.1 Optimal result . . . . .	735
3.116.2 Mathematica [A] (verified) . . . . .	735
3.116.3 Rubi [A] (verified) . . . . .	736
3.116.4 Maple [C] (verified) . . . . .	737
3.116.5 Fricas [C] (verification not implemented) . . . . .	738
3.116.6 Sympy [F] . . . . .	738
3.116.7 Maxima [F] . . . . .	739
3.116.8 Giac [F] . . . . .	739
3.116.9 Mupad [F(-1)] . . . . .	739

#### 3.116.1 Optimal result

Integrand size = 19, antiderivative size = 41

$$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2E(\frac{1}{2}(c+dx)|2)}{bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

```
output 2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)
```

#### 3.116.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2E(\frac{1}{2}(c+dx)|2)}{bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

```
input Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(3/2),x]
```

```
output (2*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])
```

**3.116.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2030, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)}{(b\sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int \frac{1}{\sqrt{b\sec(c+dx)}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{1}{\sqrt{b\csc(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\int \sqrt{\cos(c+dx)} dx}{b\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2E(\frac{1}{2}(c+dx)|2)}{bd\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]/(b*Sec[c + d*x])^(3/2), x]`

output `(2*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

3.116.3.1 Defintions of rubi rules used

```
rule 2030 Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 4258 Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

3.116.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 305, normalized size of antiderivative = 7.44

method	result
risch	$-\frac{i\sqrt{2}}{db\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}} - \frac{i\left(-\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}(-2i\text{EllipticE}\left(\sqrt{-i(e^{i(dx+c)}+i)}\right)}{\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}}\right)}{db(e^{2i(dx+c)}+1)\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}}$
default	$\frac{2i\text{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)-2i\text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}}{db(e^{2i(dx+c)}+1)\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}}$

```
input int(sec(d*x+c)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

3.116.  $\int \frac{\sec(c+dx)}{(b\sec(c+dx))^{3/2}} dx$

output `-I/d*2^(1/2)/b/(b*exp(I*(d*x+c))/(exp(I*(d*x+c))^2+1))^(1/2)-I/d*(-2*(b*exp(I*(d*x+c))^2+b)/b/(exp(I*(d*x+c))*(b*exp(I*(d*x+c))^2+b))^(1/2)+I*(-I*(exp(I*(d*x+c))+I))^(1/2)*2^(1/2)*(I*(exp(I*(d*x+c))-I))^(1/2)*(I*exp(I*(d*x+c)))^(1/2)/(b*exp(I*(d*x+c))^3+b*exp(I*(d*x+c)))^(1/2)*(-2*I*EllipticE((-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))+I*EllipticF((-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))))*2^(1/2)/b/(exp(I*(d*x+c))^2+1)/(b*exp(I*(d*x+c))/(exp(I*(d*x+c))^2+1))^(1/2)*(b*exp(I*(d*x+c))*(exp(I*(d*x+c))^2+1))^(1/2)`

### 3.116.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.61

$$\int \frac{\sec(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c))) - I\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c)))}{(b^2d)}$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `(I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b^2*d)`

### 3.116.6 Sympy [F]

$$\int \frac{\sec(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \int \frac{\sec(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))**(3/2),x)`

output `Integral(sec(c + d*x)/(b*sec(c + d*x))**(3/2), x)`

**3.116.7 Maxima [F]**

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{3/2}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(3/2), x)`

**3.116.8 Giac [F]**

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{3/2}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(3/2), x)`

**3.116.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx) \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(3/2)),x)`

output `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(3/2)), x)`



### 3.117 $\int \frac{1}{(b \sec(c+dx))^{3/2}} dx$

3.117.1 Optimal result . . . . .	740
3.117.2 Mathematica [A] (verified) . . . . .	740
3.117.3 Rubi [A] (verified) . . . . .	741
3.117.4 Maple [C] (verified) . . . . .	742
3.117.5 Fricas [C] (verification not implemented) . . . . .	743
3.117.6 Sympy [F] . . . . .	743
3.117.7 Maxima [F] . . . . .	743
3.117.8 Giac [F] . . . . .	744
3.117.9 Mupad [F(-1)] . . . . .	744

#### 3.117.1 Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{1}{(b \sec(c + dx))^{3/2}} dx = \frac{2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3b^2d} + \frac{2 \sin(c + dx)}{3bd\sqrt{b \sec(c + dx)}}$$

output `2/3*sin(d*x+c)/b/d/(b*sec(d*x+c))^(1/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d`

#### 3.117.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

$$\int \frac{1}{(b \sec(c + dx))^{3/2}} dx = \frac{\sec^2(c + dx) \left( 2\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx)) \right)}{3d(b \sec(c + dx))^{3/2}}$$

input `Integrate[(b*Sec[c + d*x])^(-3/2),x]`

output `(Sec[c + d*x]^2*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*d*(b*Sec[c + d*x])^(3/2))`

**3.117.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3b^2 d} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(-3/2),x]`

output `(2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*sqrt[b*Sec[c + d*x]])`

---

3.117.  $\int \frac{1}{(b \sec(c + dx))^{3/2}} dx$

## 3.117.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.117.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.00

method	result
default	$-\frac{2\left(i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)+i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\right)}{3d\sqrt{b}\sec(dx+c)b}$

input `int(1/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/3/d/(b*sec(d*x+c))^(1/2)/b*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-sin(d*x+c))`

**3.117.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.21

$$\int \frac{1}{(b \sec(c + dx))^{3/2}} dx = \frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) - i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx+c))}{(b \sec(c + dx))^{3/2}}$$

input `integrate(1/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/3*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)*sin(d*x + c) - I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^2*d)`

**3.117.6 Sympy [F]**

$$\int \frac{1}{(b \sec(c + dx))^{3/2}} dx = \int \frac{1}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

input `integrate(1/(b*sec(d*x+c))**(3/2),x)`

output `Integral((b*sec(c + d*x))**(-3/2), x)`

**3.117.7 Maxima [F]**

$$\int \frac{1}{(b \sec(c + dx))^{3/2}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(3/2), x)`

**3.117.8 Giac [F]**

$$\int \frac{1}{(b \sec(c + dx))^{3/2}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2), x)`

**3.117.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(b \sec(c + dx))^{3/2}} dx = \int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(1/(b/cos(c + d*x))^(3/2),x)`

output `int(1/(b/cos(c + d*x))^(3/2), x)`

### 3.118 $\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.118.1 Optimal result . . . . .	745
3.118.2 Mathematica [A] (verified) . . . . .	745
3.118.3 Rubi [A] (verified) . . . . .	746
3.118.4 Maple [C] (verified) . . . . .	747
3.118.5 Fricas [C] (verification not implemented) . . . . .	748
3.118.6 Sympy [F] . . . . .	748
3.118.7 Maxima [F] . . . . .	749
3.118.8 Giac [F] . . . . .	749
3.118.9 Mupad [F(-1)] . . . . .	749

#### 3.118.1 Optimal result

Integrand size = 19, antiderivative size = 69

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{6E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5bd\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}}$$

```
output 2/5*sin(d*x+c)/d/(b*sec(d*x+c))^(3/2)+6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos
(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)
/(b*sec(d*x+c))^(1/2)
```

#### 3.118.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.87

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{\sqrt{b \sec(c + dx)} \left( 12 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) + \sin(3(c + dx)) \right)}{10b^2d}$$

```
input Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(3/2),x]
```

```
output (Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + S
in[c + d*x] + Sin[3*(c + d*x)]))/(10*b^2*d)
```

**3.118.3 Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {3042, 2030, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2}) (b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{4258} \\
 & b \left( \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b \left( \frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3119} \\
 & b \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)
 \end{aligned}$$

input `Int[Cos[c + d*x]/(b*Sec[c + d*x])^(3/2),x]`

output `b*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2)))`

### 3.118.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_)*(x_)])*(b_))^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_)*(x_)])*(b_))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.118.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.55 (sec) , antiderivative size = 420, normalized size of antiderivative = 6.09

method	result
default	$\frac{6i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)}{5} - \frac{6i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)}{5}$

3.118.  $\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx$



input `int(cos(d*x+c)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2/5/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b*(3*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-3*I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+cos(d*x+c)^2*sin(d*x+c)+cos(d*x+c)*sin(d*x+c)+3*sin(d*x+c))`

### 3.118.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.38

$$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^2 \sin(dx+c) + 3i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + I \sin(dx+c))) - 3i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - I \sin(dx+c)))}{b^2 d}$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/5*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^2*sin(d*x + c) + 3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/b^2*d`

### 3.118.6 Sympy [F]

$$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))**(3/2),x)`

3.118.  $\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

output `Integral(cos(c + d*x)/(b*sec(c + d*x))**(3/2), x)`

### 3.118.7 Maxima [F]

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(3/2), x)`

### 3.118.8 Giac [F]

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(3/2), x)`

### 3.118.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(cos(c + d*x)/(b/cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)/(b/cos(c + d*x))^(3/2), x)`

**3.119**       $\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.119.1 Optimal result . . . . .	750
3.119.2 Mathematica [A] (verified) . . . . .	750
3.119.3 Rubi [A] (verified) . . . . .	751
3.119.4 Maple [C] (verified) . . . . .	753
3.119.5 Fracas [C] (verification not implemented) . . . . .	753
3.119.6 Sympy [F] . . . . .	754
3.119.7 Maxima [F] . . . . .	754
3.119.8 Giac [F] . . . . .	754
3.119.9 Mupad [F(-1)] . . . . .	755

**3.119.1 Optimal result**

Integrand size = 21, antiderivative size = 98

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{21b^2d} + \frac{2b \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21bd\sqrt{b \sec(c+dx)}}$$

output `2/7*b*sin(d*x+c)/d/(b*sec(d*x+c))^(5/2)+10/21*sin(d*x+c)/b/d/(b*sec(d*x+c))^(1/2)+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d`

**3.119.2 Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.67

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\sqrt{b \sec(c+dx)} \left( 40\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 26 \sin(2(c+dx)) + 3 \sin(4(c+dx)) \right)}{84b^2d}$$

input `Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(3/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*b^2*d)`

---

3.119.       $\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

**3.119.3 Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^2 (b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^2 \left( \frac{5 \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{5 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^2 \left( \frac{5 \left( \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{5 \left( \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$b^2 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd (b \sec(c+dx))^{5/2}} \right)$$

↓ 3042

$$b^2 \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd (b \sec(c+dx))^{5/2}} \right)$$

↓ 3120

$$b^2 \left( \frac{5 \left( \frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd (b \sec(c+dx))^{5/2}} \right)$$

input `Int[Cos[c + d*x]^2/(b*Sec[c + d*x])^(3/2),x]`

output `b^2*((2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/(7*b^2))`

### 3.119.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.119.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.60 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.63

method	result
default	$-\frac{2\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\right)}{21d\sqrt{b\sec(dx+c)}b}$

input `int(cos(d*x+c)^2/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/21/d/(b*sec(d*x+c))^(1/2)/b*(5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*cos(d*x+c)^2*sin(d*x+c)-5*sin(d*x+c))`

### 3.119.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.02

$$\int \frac{\cos^2(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{2(3\cos(dx+c)^3 + 5\cos(dx+c))\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c) - 5i\sqrt{2}\sqrt{b}\operatorname{weierstrass}}{21d\sqrt{b\sec(dx+c)}b}$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(3/2),x, algorithm="fracas")`

3.119. 
$$\int \frac{\cos^2(c+dx)}{(b\sec(c+dx))^{3/2}} dx$$

output `1/21*(2*(3*cos(d*x + c)^3 + 5*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c) - 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^2*d)`

### 3.119.6 Sympy [F]

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(3/2), x)`

output `Integral(cos(c + d*x)**2/(b*sec(c + d*x))**(3/2), x)`

### 3.119.7 Maxima [F]

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(3/2), x, algorithm="maxima")`

output `integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(3/2), x)`

### 3.119.8 Giac [F]

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(3/2), x, algorithm="giac")`

output `integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(3/2), x)`

---

3.119.  $\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

**3.119.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^2}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(cos(c + d*x)^2/(b/cos(c + d*x))^(3/2), x)`output `int(cos(c + d*x)^2/(b/cos(c + d*x))^(3/2), x)`



### 3.120 $\int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.120.1 Optimal result . . . . .	756
3.120.2 Mathematica [A] (verified) . . . . .	756
3.120.3 Rubi [A] (verified) . . . . .	757
3.120.4 Maple [C] (verified) . . . . .	759
3.120.5 Fricas [C] (verification not implemented) . . . . .	759
3.120.6 Sympy [F(-1)] . . . . .	760
3.120.7 Maxima [F] . . . . .	760
3.120.8 Giac [F] . . . . .	760
3.120.9 Mupad [F(-1)] . . . . .	761

#### 3.120.1 Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{14E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{2b^2 \sin(c+dx)}{9d(b \sec(c+dx))^{7/2}} + \frac{14 \sin(c+dx)}{45d(b \sec(c+dx))^{3/2}}$$

output `2/9*b^2*sin(d*x+c)/d/(b*sec(d*x+c))^(7/2)+14/45*sin(d*x+c)/d/(b*sec(d*x+c))^(3/2)+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`

#### 3.120.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.75

$$\int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{84E\left(\frac{1}{2}(c+dx) \mid 2\right) + \cos^{\frac{3}{2}}(c+dx)(33 \sin(c+dx) + 5 \sin(3(c+dx)))}{90d \cos^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{3/2}}$$

input `Integrate[Cos[c + d*x]^3/(b*Sec[c + d*x])^(3/2),x]`

output `(84*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]^(3/2)*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)]))/(90*d*Cos[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2))`

**3.120.3 Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.15, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^3 (b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^3 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{9/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^3 \left( \frac{7 \int \frac{1}{(b \sec(c+dx))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{7 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^3 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$b^3 \left( \frac{7 \left( \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)$$

↓ 3042

$$b^3 \left( \frac{7 \left( \frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)$$

↓ 3119

$$b^3 \left( \frac{7 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)$$

input `Int[Cos[c + d*x]^3/(b*Sec[c + d*x])^(3/2),x]`

output `b^3*((2*Sin[c + d*x])/(9*b*d*(b*Sec[c + d*x])^(7/2)) + (7*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))))/(9*b^2)`

### 3.120.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.120.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.60 (sec) , antiderivative size = 454, normalized size of antiderivative = 4.68

method	result
default	$-\frac{2(21i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - 21i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)))$

input `int(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2), x, method=_RETURNVERBOSE)`

output `-2/45/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b*(21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)-21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)-5*cos(d*x+c)^4*sin(d*x+c)+42*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)-42*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)-5*cos(d*x+c)^3*sin(d*x+c)+21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)-21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)-7*cos(d*x+c)^2*sin(d*x+c)-7*cos(d*x+c)*sin(d*x+c)-21*sin(d*x+c)`

### 3.120.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.11

$$\int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2(5 \cos(dx+c)^4 + 7 \cos(dx+c)^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c) + 21i \sqrt{2} \sqrt{b} \operatorname{weierstrass}}{(b \sec(c+dx))^{3/2}}$$

input `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")`

$$3.120. \quad \int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

output  $1/45*(2*(5*\cos(d*x + c)^4 + 7*\cos(d*x + c)^2)*\sqrt{b/\cos(d*x + c)}*\sin(d*x + c) + 21*I*\sqrt{2}*\sqrt{b}*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - 21*I*\sqrt{2}*\sqrt{b}*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/(b^2*d)$

### 3.120.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3/(b*sec(d*x+c))**(3/2),x)`

output Timed out

### 3.120.7 Maxima [F]

$$\int \frac{\cos^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^3}{(b \sec(dx + c))^{3/2}} dx$$

input `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate(cos(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)`

### 3.120.8 Giac [F]

$$\int \frac{\cos^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^3}{(b \sec(dx + c))^{3/2}} dx$$

input `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(cos(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)`

**3.120.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^3}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(cos(c + d*x)^3/(b/cos(c + d*x))^(3/2), x)`output `int(cos(c + d*x)^3/(b/cos(c + d*x))^(3/2), x)`

### 3.121 $\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.121.1 Optimal result . . . . .	762
3.121.2 Mathematica [A] (verified) . . . . .	762
3.121.3 Rubi [A] (verified) . . . . .	763
3.121.4 Maple [C] (verified) . . . . .	765
3.121.5 Fricas [C] (verification not implemented) . . . . .	765
3.121.6 Sympy [F] . . . . .	766
3.121.7 Maxima [F] . . . . .	766
3.121.8 Giac [F] . . . . .	766
3.121.9 Mupad [F(-1)] . . . . .	767

#### 3.121.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{21b^3d} + \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^4d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^6d}$$

output `10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^4/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^6/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d`

#### 3.121.2 Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.64

$$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{(b \sec(c+dx))^{5/2} \left(10 \cos^{5/2}(c+dx) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 5 \sin(2(c+dx))\right) + 6 \tan(c+dx)}{21b^5d}$$

input `Integrate[Sec[c + d*x]^7/(b*Sec[c + d*x])^(5/2),x]`

output `((b*Sec[c + d*x])^(5/2)*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*b^5*d)`

**3.121.3 Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.08, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{9/2} dx}{b^7} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{9/2} dx}{b^7} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{5}{7} b^2 \int (b \sec(c+dx))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^7} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{5}{7} b^2 \int (b \csc(c+dx+\frac{\pi}{2}))^{5/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^7} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^7} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^7} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{5}{7} b^2 \left( \frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^7} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.121.  $\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx$



$$\frac{\frac{5}{7}b^2 \left( \frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^7}$$

↓ 3120

$$\frac{\frac{5}{7}b^2 \left( \frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{7/2}}{7d}}{b^7}$$

input `Int[Sec[c + d*x]^7/(b*Sec[c + d*x])^(5/2),x]`

output `((2*b*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*d) + (5*b^2*((2*b^2*Sqrt[Cos[c + d*x])*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d))/7)/b^7`

### 3.121.3.1 Defintions of rubi rules used

rule 2030 `Int[(F*x_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F*x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Simp[b^2*((n-2)/(n-1)) Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

**3.121.4 Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.63 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.66

method	result
default	$-\frac{10i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i)\sec(dx+c)}{21} + \frac{2\tan(dx+c)\sec(dx+c)^3}{d\sqrt{b\sec(dx+c)}b^2} - \frac{10i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}{d\sqrt{b\sec(dx+c)}b^2}$

input `int(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `2/21/d/(b*sec(d*x+c))^(1/2)/b^2*(-5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+3*tan(d*x+c)*sec(d*x+c)^3-5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+5*sec(d*x+c)*tan(d*x+c))`

**3.121.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.17

$$\int \frac{\sec^7(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \frac{-5i\sqrt{2}\sqrt{b}\cos(dx+c)^3\operatorname{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))}{(b\sec(c+dx))^{5/2}}$$

input `integrate(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2),x, algorithm="fracas")`

output `1/21*(-5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(5*cos(d*x + c)^2 + 3)*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b^3*d*cos(d*x + c)^3)`

**3.121.6 Sympy [F]**

$$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)**7/(b*sec(d*x+c))**(5/2),x)`

output `Integral(sec(c + d*x)**7/(b*sec(c + d*x))**(5/2), x)`

**3.121.7 Maxima [F]**

$$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^7}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^7/(b*sec(d*x + c))^(5/2), x)`

**3.121.8 Giac [F]**

$$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^7}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^7/(b*sec(d*x + c))^(5/2), x)`

**3.121.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \int \frac{1}{\cos(c+dx)^7 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(1/(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2)),x)`output `int(1/(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2)), x)`

### 3.122 $\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.122.1 Optimal result . . . . .	768
3.122.2 Mathematica [A] (verified) . . . . .	768
3.122.3 Rubi [A] (verified) . . . . .	769
3.122.4 Maple [C] (verified) . . . . .	771
3.122.5 Fracas [C] (verification not implemented) . . . . .	771
3.122.6 Sympy [F] . . . . .	772
3.122.7 Maxima [F] . . . . .	772
3.122.8 Giac [F] . . . . .	772
3.122.9 Mupad [F(-1)] . . . . .	773

#### 3.122.1 Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{5/2}} dx = -\frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5b^2d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5b^3d} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^5d}$$

output  $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/b^5/d-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/b^3/d$

#### 3.122.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.64

$$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{-\frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{\sqrt{\cos(c+dx)}} + 2(3 + \sec^2(c+dx)) \tan(c+dx)}{5b^2d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^6/(b*Sec[c + d*x])^(5/2),x]`

output  $((-6*\text{EllipticE}[(c+d*x)/2, 2])/ \text{Sqrt}[\text{Cos}[c+d*x]] + 2*(3 + \text{Sec}[c+d*x]^2)*\text{Tan}[c+d*x])/(5*b^2*d*\text{Sqrt}[b*\text{Sec}[c+d*x]])$

**3.122.3 Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{7/2} dx}{b^6} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{7/2} dx}{b^6} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \int (b \sec(c+dx))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^6} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^6} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^6} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^6} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{3}{5} b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx)(b \sec(c+dx))^{5/2}}{5d}}{b^6} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}}{b^6}$$

↓ 3119

$$\frac{\frac{3}{5}b^2 \left( \frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c+dx)|2\right)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right) + \frac{2b \sin(c+dx) (b \sec(c+dx))^{5/2}}{5d}}{b^6}$$

input `Int[Sec[c + d*x]^6/(b*Sec[c + d*x])^(5/2), x]`

output `((2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5)/b^6`

### 3.122.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b.)*(v.))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d.)*(x_)])*(b.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d.)*(x_)])*(b.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.122.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 420, normalized size of antiderivative = 4.20

method	result
default	$\frac{6i \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)}{5} - \frac{6i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}{5}$

```
input int(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2/5/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^2*(3*I*(1/(cos(d*x+c)+1))^(1/2)
)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I
)*cos(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)
)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1)
)^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x
+c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Ell
ipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x
+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+
c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Elliptic
E(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+3*tan(d*x+c)+sec(d*x+c)*tan(d*x
+c)+tan(d*x+c)*sec(d*x+c)^2)
```

### 3.122.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.23

$$\int \frac{\sec^6(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \frac{-3i\sqrt{2}\sqrt{b}\cos(dx+c)^2 \operatorname{weierstrassZeta}(-4,0, \operatorname{weierstrassPInverse}(-4,0, \cos(dx+c)))}{(b\sec(c+dx))^{5/2}}$$

```
input integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output 1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstras
sPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos
(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c)
- I*sin(d*x + c))) + 2*(3*cos(d*x + c)^2 + 1)*sqrt(b/cos(d*x + c))*sin(d*
x + c))/(b^3*d*cos(d*x + c)^2)
```

---

3.122.  $\int \frac{\sec^6(c+dx)}{(b\sec(c+dx))^{5/2}} dx$



**3.122.6 Sympy [F]**

$$\int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)**6/(b*sec(d*x+c))**(5/2),x)`

output `Integral(sec(c + d*x)**6/(b*sec(c + d*x))**(5/2), x)`

**3.122.7 Maxima [F]**

$$\int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^6}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(5/2), x)`

**3.122.8 Giac [F]**

$$\int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^6}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(5/2), x)`

**3.122.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^6(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{1}{\cos(c+dx)^6 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2)),x)`output `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2)), x)`

### 3.123 $\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.123.1 Optimal result . . . . .	774
3.123.2 Mathematica [A] (verified) . . . . .	774
3.123.3 Rubi [A] (verified) . . . . .	775
3.123.4 Maple [C] (verified) . . . . .	776
3.123.5 Fricas [C] (verification not implemented) . . . . .	777
3.123.6 Sympy [F] . . . . .	777
3.123.7 Maxima [F] . . . . .	778
3.123.8 Giac [F] . . . . .	778
3.123.9 Mupad [F(-1)] . . . . .	778

#### 3.123.1 Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^3d} + \frac{2(b \sec(c+dx))^{3/2} \sin(c+dx)}{3b^4d}$$

output `2/3*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^4/d+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d`

#### 3.123.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2\sqrt{b \sec(c+dx)} \left( \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \tan(c+dx) \right)}{3b^3d}$$

input `Integrate[Sec[c + d*x]^5/(b*Sec[c + d*x])^(5/2),x]`

output `(2*Sqrt[b*Sec[c + d*x]]*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*b^3*d)`

**3.123.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{5/2} dx}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{5/2} dx}{b^5} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{1}{3}b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3}b^2 \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^5} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3}b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^5} \\
 & \quad \downarrow \text{3120} \\
 & \frac{\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}}{b^5}
 \end{aligned}$$

input `Int[Sec[c + d*x]^5/(b*Sec[c + d*x])^(5/2), x]`

$$3.123. \quad \int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

output  $((2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) / (3*d) + (2*b*(b*\text{Sec}[c + d*x])^{3/2}*\text{Sin}[c + d*x]) / (3*d)) / b^5$

### 3.123.3.1 Defintions of rubi rules used

rule 2030  $\text{Int}[(\text{Fx}_.)*(v_.)^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[1/b^m \text{Int}[(b*v)^{(m+n)*\text{Fx}, x}], x] /;$   $\text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3120  $\text{Int}[1/\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_.)]], x\_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$   $\text{FreeQ}[\{c, d\}, x]$

rule 4255  $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n-1)} / (d*(n-1))), x] + \text{Simp}[b^2*((n-2)/(n-1)) \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /;$   $\text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

rule 4258  $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n * \text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /;$   $\text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{EqQ}[n^2, 1/4]$

### 3.123.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 149, normalized size of antiderivative = 2.07

method	result
default	$\frac{-\frac{2i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticF}(i(-\cot(dx+c)+\text{csc}(dx+c)),i)\sec(dx+c)}{3} - \frac{2i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticF}(i(-\cot(dx+c))}{3}}{d\sqrt{b\sec(dx+c)}b^2}$

input `int(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2), x, method=_RETURNVERBOSE)`

$$3.123. \int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{5/2}} dx$$

output  $2/3/d/(b*\sec(d*x+c))^{(1/2)}/b^2*(-I*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\text{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*\sec(d*x+c)-I*(1/(\cos(d*x+c)+1))^{(1/2)}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{(1/2)}*\text{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I))+\sec(d*x+c)*\tan(d*x+c))$

### 3.123.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.40

$$\int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\cos(dx+c)\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+i\sqrt{2}\sqrt{b}\cos(dx+c)\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))+2\sqrt{b/\cos(dx+c)}\sin(dx+c)}{(b^3d\cos(dx+c))^{5/2}}$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output  $1/3*(-I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))+2*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c))/(b^3*d*\cos(d*x+c))^{5/2}$

### 3.123.6 Sympy [F]

$$\int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{5/2}} dx$$

input `integrate(sec(d*x+c)**5/(b*sec(d*x+c))**(5/2),x)`

output `Integral(sec(c+d*x)**5/(b*sec(c+d*x))**(5/2),x)`

**3.123.7 Maxima [F]**

$$\int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^5}{(b\sec(dx+c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(5/2), x)`

**3.123.8 Giac [F]**

$$\int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^5}{(b\sec(dx+c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(5/2), x)`

**3.123.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^5(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{1}{\cos(c+dx)^5 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2)),x)`

output `int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2)), x)`

### 3.124 $\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.124.1 Optimal result . . . . .	779
3.124.2 Mathematica [A] (verified) . . . . .	779
3.124.3 Rubi [A] (verified) . . . . .	780
3.124.4 Maple [C] (verified) . . . . .	781
3.124.5 Fracas [C] (verification not implemented) . . . . .	782
3.124.6 Sympy [F] . . . . .	782
3.124.7 Maxima [F] . . . . .	783
3.124.8 Giac [F] . . . . .	783
3.124.9 Mupad [F(-1)] . . . . .	783

#### 3.124.1 Optimal result

Integrand size = 21, antiderivative size = 68

$$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx = -\frac{2E(\frac{1}{2}(c+dx)|2)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2\sqrt{b \sec(c+dx)} \sin(c+dx)}{b^3 d}$$

output `-2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/b^3/d`

#### 3.124.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.75

$$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{-\frac{2E(\frac{1}{2}(c+dx)|2)}{\sqrt{\cos(c+dx)}} + 2 \tan(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^4/(b*Sec[c + d*x])^(5/2),x]`

output `((-2*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*Tan[c + d*x])/(b^2*d*Sqrt[b*Sec[c + d*x]])`



**3.124.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2030, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{3/2} dx}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx}{b^4} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{b^4} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^4} \\
 & \quad \downarrow \text{3119} \\
 & \frac{\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}}{b^4}
 \end{aligned}$$

---

3.124.  $\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

input `Int[Sec[c + d*x]^4/(b*Sec[c + d*x])^(5/2),x]`

output `((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d)/b^4`

### 3.124.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.124.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 392, normalized size of antiderivative = 5.76

method	result
default	$-\frac{2 \left( i \operatorname{EllipticE} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - i \operatorname{EllipticF} \left( i \left( -\cot(dx+c) + \operatorname{csc}(dx+c) \right), i \right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \right)}{\dots}$

3.124. 
$$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

input `int(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^2*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-tan(d*x+c))`

### 3.124.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.26

$$\int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)))}{(b\sec(c+dx))^{5/2}}$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b^3*d)`

### 3.124.6 Sympy [F]

$$\int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{5/2}} dx$$

input `integrate(sec(d*x+c)**4/(b*sec(d*x+c))**(5/2),x)`

output `Integral(sec(c + d*x)**4/(b*sec(c + d*x))**(5/2), x)`

---

3.124.  $\int \frac{\sec^4(c+dx)}{(b\sec(c+dx))^{5/2}} dx$

**3.124.7 Maxima [F]**

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^4}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(5/2), x)`

**3.124.8 Giac [F]**

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^4}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(5/2), x)`

**3.124.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{1}{\cos(c + dx)^4 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2)),x)`

output `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2)), x)`

### 3.125 $\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.125.1 Optimal result	784
3.125.2 Mathematica [A] (verified)	784
3.125.3 Rubi [A] (verified)	785
3.125.4 Maple [C] (verified)	786
3.125.5 Fricas [C] (verification not implemented)	787
3.125.6 Sympy [F]	787
3.125.7 Maxima [F]	787
3.125.8 Giac [F]	788
3.125.9 Mupad [F(-1)]	788

#### 3.125.1 Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{b^3 d}$$

output  $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b^3/d$

#### 3.125.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{b^3 d}$$

input  $\operatorname{Integrate}[\operatorname{Sec}[c + d*x]^3/(b*\operatorname{Sec}[c + d*x])^{(5/2)}, x]$

output  $(2*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2]*\operatorname{Sqrt}[b*\operatorname{Sec}[c + d*x]])/(b^3*d)$

**3.125.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2030, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int \sqrt{b \sec(c+dx)} dx}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{b^3} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b^3} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{b^3 d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^3/(b*Sec[c + d*x])^(5/2), x]`

output `(2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/(b^3*d)`

## 3.125.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

## 3.125.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.63 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.76

method	result	size
default	$-\frac{2i\sqrt{\frac{1}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i)}{db^2\sqrt{b\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}$	72

input `int(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2), x, method=_RETURNVERBOSE)`

output `-2*I/d*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)/b^2/(b*sec(d*x+c))^(1/2)/(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)`

---

3.125. 
$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{5/2}} dx$$

**3.125.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c))}{b^3d}$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^3*d)`

**3.125.6 Sympy [F]**

$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{5/2}} dx$$

input `integrate(sec(d*x+c)**3/(b*sec(d*x+c))**(5/2),x)`

output `Integral(sec(c + d*x)**3/(b*sec(c + d*x))**(5/2), x)`

**3.125.7 Maxima [F]**

$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^3}{(b\sec(dx+c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(5/2), x)`



**3.125.8 Giac [F]**

$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^3}{(b\sec(dx+c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(5/2), x)`

**3.125.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^3(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{1}{\cos(c+dx)^3 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2)),x)`

output `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2)), x)`

### 3.126 $\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.126.1 Optimal result . . . . .	789
3.126.2 Mathematica [A] (verified) . . . . .	789
3.126.3 Rubi [A] (verified) . . . . .	790
3.126.4 Maple [C] (verified) . . . . .	791
3.126.5 Fracas [C] (verification not implemented) . . . . .	792
3.126.6 Sympy [F] . . . . .	792
3.126.7 Maxima [F] . . . . .	793
3.126.8 Giac [F] . . . . .	793
3.126.9 Mupad [F(-1)] . . . . .	793

#### 3.126.1 Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

output `2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`

#### 3.126.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93

$$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d \cos^{\frac{5}{2}}(c+dx) (b \sec(c+dx))^{5/2}}$$

input `Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(5/2),x]`

output `(2*EllipticE[(c + d*x)/2, 2])/(d*Cos[c + d*x]^(5/2)*(b*Sec[c + d*x])^(5/2))`

**3.126.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2030, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4258} \\
 & \frac{\int \sqrt{\cos(c+dx)} dx}{b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2E(\frac{1}{2}(c+dx)|2)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(5/2),x]`

output `(2*EllipticE[(c + d*x)/2, 2])/(b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

3.126.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

3.126.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 305, normalized size of antiderivative = 7.44

method	result
risch	$-\frac{i\sqrt{2}}{db^2\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}}+1}} - \frac{i\left(-\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}(-2i\text{EllipticE}\left(\sqrt{-i(e^{i(dx+c)}}+i)}\right))\right)}{\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}}$
default	$\frac{2i\text{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)-2i\text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}}{db^2(e^{2i(dx+c)}+1)\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}}+1}}$

input `int(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2), x, method=_RETURNVERBOSE)`

```
output -I/d*2^(1/2)/b^2/(b*exp(I*(d*x+c))/(exp(I*(d*x+c))^2+1))^(1/2)-I/d*(-2*(b*
exp(I*(d*x+c))^2+b)/b/(exp(I*(d*x+c))*(b*exp(I*(d*x+c))^2+b))^(1/2)+I*(-I*
(exp(I*(d*x+c))+I))^(1/2)*2^(1/2)*(I*(exp(I*(d*x+c))-I))^(1/2)*(I*exp(I*(d
*x+c)))^(1/2)/(b*exp(I*(d*x+c))^3+b*exp(I*(d*x+c)))^(1/2)*(-2*I*EllipticE(
(-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))+I*EllipticF((-I*(exp(I*(d*x+c))
+I))^(1/2),1/2*2^(1/2))))*2^(1/2)/b^2/(exp(I*(d*x+c))^2+1)/(b*exp(I*(d*x+c
)))/(exp(I*(d*x+c))^2+1))^(1/2)*(b*exp(I*(d*x+c))*(exp(I*(d*x+c))^2+1))^(1/
2)
```

### 3.126.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.61

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{b^3 d}$$

```
input integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output (I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d
*x + c) + I*sin(d*x + c))) - I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weie
rstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/b^3*d
```

### 3.126.6 Sympy [F]

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx$$

```
input integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(5/2),x)
```

```
output Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(5/2), x)
```

**3.126.7 Maxima [F]**

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(5/2), x)`

**3.126.8 Giac [F]**

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(5/2), x)`

**3.126.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{1}{\cos(c + dx)^2 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2)),x)`

output `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2)), x)`

### 3.127 $\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.127.1 Optimal result . . . . .	794
3.127.2 Mathematica [A] (verified) . . . . .	794
3.127.3 Rubi [A] (verified) . . . . .	795
3.127.4 Maple [C] (verified) . . . . .	796
3.127.5 Fracas [C] (verification not implemented) . . . . .	797
3.127.6 Sympy [F] . . . . .	797
3.127.7 Maxima [F] . . . . .	798
3.127.8 Giac [F] . . . . .	798
3.127.9 Mupad [F(-1)] . . . . .	798

#### 3.127.1 Optimal result

Integrand size = 19, antiderivative size = 72

$$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^3d} + \frac{2 \sin(c+dx)}{3b^2d\sqrt{b \sec(c+dx)}}$$

output `2/3*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(1/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d`

#### 3.127.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.86

$$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\sec^2(c+dx) \left( 2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \sin(2(c+dx)) \right)}{3bd(b \sec(c+dx))^{3/2}}$$

input `Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(5/2),x]`

output `(Sec[c + d*x]^2*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*b*d*(b*Sec[c + d*x])^(3/2))`

**3.127.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {2030, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \int \frac{1}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2 \sqrt{\cos(c+dx)} \text{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \\
 & \frac{\dots}{b}
 \end{aligned}$$

---

3.127.  $\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx$



input `Int[Sec[c + d*x]/(b*Sec[c + d*x])^(5/2),x]`

output `((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]]))/b`

### 3.127.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m + n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.127.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.00

method	result
default	$-\frac{2\left(i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)+i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)\right)}{3d\sqrt{b}\sec(dx+c)b^2}$

3.127.  $\int \frac{\sec(c+dx)}{(b\sec(c+dx))^{5/2}} dx$

input `int(sec(d*x+c)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/3/d/(b*sec(d*x+c))^(1/2)/b^2*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-sin(d*x+c))`

### 3.127.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.21

$$\int \frac{\sec(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \frac{2\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) - i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx+c))}{(b\sec(c+dx))^{5/2}}$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/3*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)*sin(d*x + c) - I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^3*d)`

### 3.127.6 Sympy [F]

$$\int \frac{\sec(c+dx)}{(b\sec(c+dx))^{5/2}} dx = \int \frac{\sec(c+dx)}{(b\sec(c+dx))^{5/2}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))**(5/2),x)`

output `Integral(sec(c + d*x)/(b*sec(c + d*x))**(5/2), x)`

**3.127.7 Maxima [F]**

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(5/2), x)`

**3.127.8 Giac [F]**

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(5/2), x)`

**3.127.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{1}{\cos(c + dx) \left(\frac{b}{\cos(c + dx)}\right)^{5/2}} dx$$

input `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(5/2)),x)`

output `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(5/2)), x)`

### 3.128 $\int \frac{1}{(b \sec(c+dx))^{5/2}} dx$

3.128.1 Optimal result . . . . .	799
3.128.2 Mathematica [A] (verified) . . . . .	799
3.128.3 Rubi [A] (verified) . . . . .	800
3.128.4 Maple [C] (verified) . . . . .	801
3.128.5 Fricas [C] (verification not implemented) . . . . .	802
3.128.6 Sympy [F] . . . . .	802
3.128.7 Maxima [F] . . . . .	803
3.128.8 Giac [F] . . . . .	803
3.128.9 Mupad [F(-1)] . . . . .	803

#### 3.128.1 Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{1}{(b \sec(c + dx))^{5/2}} dx = \frac{6E(\frac{1}{2}(c + dx) | 2)}{5b^2d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}}$$

```
output 2/5*sin(d*x+c)/b/d/(b*sec(d*x+c))^(3/2)+6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/c
os(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/d/cos(d*x+c)^(
1/2)/(b*sec(d*x+c))^(1/2)
```

#### 3.128.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.83

$$\int \frac{1}{(b \sec(c + dx))^{5/2}} dx = \frac{\sqrt{b \sec(c + dx)} \left( 12 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) + \sin(3(c + dx)) \right)}{10b^3d}$$

```
input Integrate[(b*Sec[c + d*x])^(-5/2),x]
```

```
output (Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + S
in[c + d*x] + Sin[3*(c + d*x)]))/(10*b^3*d)
```

**3.128.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{3 \int \frac{\sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}}{5b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}}{5b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{6E(\frac{1}{2}(c + dx) | 2)}{5b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(-5/2),x]`

output `(6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))`

3.128.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

3.128.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 420, normalized size of antiderivative = 5.83

method	result
default	$\frac{{}_6F_5 \text{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{5} - \frac{{}_6F_5 \text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{5}$

input `int(1/(b*sec(d*x+c))^(5/2), x, method=_RETURNVERBOSE)`

```
output 2/5/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^2*(3*I*EllipticE(I*(-cot(d*x+c)
)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2
)*cos(d*x+c)-3*I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1)
)^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1)
)^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x
+c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Ell
ipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x
+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+
c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Elliptic
F(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+cos(d*x+c)^2*sin(d*x+c)+cos(d*x
+c)*sin(d*x+c)+3*sin(d*x+c)
```

### 3.128.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.32

$$\int \frac{1}{(b \sec(c + dx))^{5/2}} dx = \frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^2 \sin(dx+c) + 3i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstra}}$$

```
input integrate(1/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output 1/5*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^2*sin(d*x + c) + 3*I*sqrt(2)*sqrt
(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin
(d*x + c))) - 3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInver
se(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/b^3*d
```

### 3.128.6 Sympy [F]

$$\int \frac{1}{(b \sec(c + dx))^{5/2}} dx = \int \frac{1}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

```
input integrate(1/(b*sec(d*x+c))**(5/2),x)
```

```
output Integral((b*sec(c + d*x))**(-5/2), x)
```

---

3.128.  $\int \frac{1}{(b \sec(c+dx))^{5/2}} dx$

**3.128.7 Maxima [F]**

$$\int \frac{1}{(b \sec(c + dx))^{5/2}} dx = \int \frac{1}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(1/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(5/2), x)`

**3.128.8 Giac [F]**

$$\int \frac{1}{(b \sec(c + dx))^{5/2}} dx = \int \frac{1}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(1/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2), x)`

**3.128.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(b \sec(c + dx))^{5/2}} dx = \int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(1/(b/cos(c + d*x))^(5/2),x)`

output `int(1/(b/cos(c + d*x))^(5/2), x)`



### 3.129 $\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.129.1 Optimal result . . . . .	804
3.129.2 Mathematica [A] (verified) . . . . .	804
3.129.3 Rubi [A] (verified) . . . . .	805
3.129.4 Maple [C] (verified) . . . . .	807
3.129.5 Fricas [C] (verification not implemented) . . . . .	807
3.129.6 Sympy [F] . . . . .	808
3.129.7 Maxima [F] . . . . .	808
3.129.8 Giac [F] . . . . .	808
3.129.9 Mupad [F(-1)] . . . . .	809

#### 3.129.1 Optimal result

Integrand size = 19, antiderivative size = 97

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{10 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21b^3d} + \frac{2 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21b^2d \sqrt{b \sec(c + dx)}}$$

output `2/7*sin(d*x+c)/d/(b*sec(d*x+c))^(5/2)+10/21*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(1/2)+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d`

#### 3.129.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.68

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{\sqrt{b \sec(c + dx)} \left( 40 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) \right)}{84b^3d}$$

input `Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(5/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*b^3*d)`

**3.129.3 Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.13, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b \left( \frac{5 \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b \left( \frac{5 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b \left( \frac{5 \left( \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b \left( \frac{5 \left( \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$\begin{aligned}
& b \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow \text{3042} \\
& b \left( \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
& \quad \downarrow \text{3120} \\
& b \left( \frac{5 \left( \frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)
\end{aligned}$$

input `Int[Cos[c + d*x]/(b*Sec[c + d*x])^(5/2), x]`

output `b*((2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/(7*b^2))`

### 3.129.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.129.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.65

method	result
default	$\frac{-\frac{10i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i)}{21} + \frac{2\cos(dx+c)^2\sin(dx+c)}{7} - \frac{10i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i)}{21}}{d\sqrt{b\sec(dx+c)}b^2}$

input `int(cos(d*x+c)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `2/21/d/(b*sec(d*x+c))^(1/2)/b^2*(-5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*cos(d*x+c)^2*sin(d*x+c)-5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+5*sin(d*x+c))`

### 3.129.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.03

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{2(3 \cos(dx + c)^3 + 5 \cos(dx + c))\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx + c) - 5i\sqrt{2}\sqrt{b}\operatorname{weierstrass}}{d}$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/21*(2*(3*cos(d*x + c)^3 + 5*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c) - 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^3*d)`

### 3.129.6 Sympy [F]

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))**(5/2),x)`

output `Integral(cos(c + d*x)/(b*sec(c + d*x))**(5/2), x)`

### 3.129.7 Maxima [F]

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(5/2), x)`

### 3.129.8 Giac [F]

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(5/2), x)`

**3.129.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(cos(c + d*x)/(b/cos(c + d*x))^(5/2), x)`output `int(cos(c + d*x)/(b/cos(c + d*x))^(5/2), x)`

### 3.130 $\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.130.1 Optimal result . . . . .	810
3.130.2 Mathematica [A] (verified) . . . . .	810
3.130.3 Rubi [A] (verified) . . . . .	811
3.130.4 Maple [C] (verified) . . . . .	813
3.130.5 Fricas [C] (verification not implemented) . . . . .	813
3.130.6 Sympy [F] . . . . .	814
3.130.7 Maxima [F] . . . . .	814
3.130.8 Giac [F] . . . . .	814
3.130.9 Mupad [F(-1)] . . . . .	815

#### 3.130.1 Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{14E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15b^2d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{2b \sin(c+dx)}{9d(b \sec(c+dx))^{7/2}} + \frac{14 \sin(c+dx)}{45bd(b \sec(c+dx))^{3/2}}$$

output  $2/9*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(7/2)}+14/45*\sin(d*x+c)/b/d/(b*\sec(d*x+c))^{(3/2)}+14/15*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

#### 3.130.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.74

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\frac{336E\left(\frac{1}{2}(c+dx) \mid 2\right)}{\sqrt{\cos(c+dx)}} + 4 \cos(c+dx)(33 \sin(c+dx) + 5 \sin(3(c+dx)))}{360b^2d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(5/2),x]`

output  $((336*\text{EllipticE}[(c+d*x)/2, 2])/Sqrt[\text{Cos}[c+d*x]] + 4*\text{Cos}[c+d*x]*(33*\text{Sin}[c+d*x] + 5*\text{Sin}[3*(c+d*x)]))/(360*b^2*d*Sqrt[b*\text{Sec}[c+d*x]])$

**3.130.3 Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 2030, 4256, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^2 (b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{9/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & b^2 \left( \frac{7 \int \frac{1}{(b \sec(c+dx))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{7 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4256} \\
 & b^2 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{7 \left( \frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right) \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$



$$b^2 \left( \frac{7 \left( \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)$$

↓ 3042

$$b^2 \left( \frac{7 \left( \frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)$$

↓ 3119

$$b^2 \left( \frac{7 \left( \frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2 \sin(c+dx)}{9bd(b \sec(c+dx))^{7/2}} \right)$$

input `Int[Cos[c + d*x]^2/(b*Sec[c + d*x])^(5/2),x]`

output `b^2*((2*Sin[c + d*x])/(9*b*d*(b*Sec[c + d*x])^(7/2)) + (7*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))))/(9*b^2))`

### 3.130.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d*n), x] + Simp[(n+1)/(b^2*n) Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_], x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

### 3.130.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 454, normalized size of antiderivative = 4.63

method	result
default	$\frac{2 \cos(dx+c)^4 \sin(dx+c)}{9} - \frac{14i \operatorname{EllipticF}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{15} + \frac{14i \operatorname{EllipticE}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{15}$

input `int(cos(d*x+c)^2/(b*sec(d*x+c))^(5/2), x, method=_RETURNVERBOSE)`

output `2/45/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^2*(5*cos(d*x+c)^4*sin(d*x+c)-21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*cos(d*x+c)+21*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+5*cos(d*x+c)^3*sin(d*x+c)-42*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)+42*I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)+7*cos(d*x+c)^2*sin(d*x+c)-21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)+21*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)+7*cos(d*x+c)*sin(d*x+c)+21*sin(d*x+c))`

### 3.130.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2(5 \cos(dx+c)^4 + 7 \cos(dx+c)^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c) + 21i \sqrt{2} \sqrt{b} \operatorname{weierstrass}(\dots)}{(b \sec(c+dx))^{5/2}}$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")`

3.130. 
$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

output  $1/45*(2*(5*\cos(dx + c)^4 + 7*\cos(dx + c)^2)*\sqrt{b/\cos(dx + c)}*\sin(dx + c) + 21*I*\sqrt{2}*\sqrt{b}*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, \cos(dx + c) + I*\sin(dx + c))) - 21*I*\sqrt{2}*\sqrt{b}*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, \cos(dx + c) - I*\sin(dx + c))))/(b^3*d)$

### 3.130.6 Sympy [F]

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx$$

input `integrate(cos(dx+c)**2/(b*sec(dx+c))**(5/2), x)`

output `Integral(cos(c + dx)**2/(b*sec(c + dx))**(5/2), x)`

### 3.130.7 Maxima [F]

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(cos(dx+c)^2/(b*sec(dx+c))^(5/2), x, algorithm="maxima")`

output `integrate(cos(dx + c)^2/(b*sec(dx + c))^(5/2), x)`

### 3.130.8 Giac [F]

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate(cos(dx+c)^2/(b*sec(dx+c))^(5/2), x, algorithm="giac")`

output `integrate(cos(dx + c)^2/(b*sec(dx + c))^(5/2), x)`

**3.130.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^2}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int(cos(c + d*x)^2/(b/cos(c + d*x))^(5/2), x)`output `int(cos(c + d*x)^2/(b/cos(c + d*x))^(5/2), x)`

### 3.131 $\int \frac{1}{(b \sec(c+dx))^{7/2}} dx$

3.131.1 Optimal result . . . . .	816
3.131.2 Mathematica [A] (verified) . . . . .	816
3.131.3 Rubi [A] (verified) . . . . .	817
3.131.4 Maple [C] (verified) . . . . .	819
3.131.5 Fricas [C] (verification not implemented) . . . . .	819
3.131.6 Sympy [F] . . . . .	820
3.131.7 Maxima [F] . . . . .	820
3.131.8 Giac [F] . . . . .	820
3.131.9 Mupad [F(-1)] . . . . .	821

#### 3.131.1 Optimal result

Integrand size = 12, antiderivative size = 100

$$\int \frac{1}{(b \sec(c + dx))^{7/2}} dx = \frac{10 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21b^4d} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21b^3d \sqrt{b \sec(c + dx)}}$$

output `2/7*sin(d*x+c)/b/d/(b*sec(d*x+c))^(5/2)+10/21*sin(d*x+c)/b^3/d/(b*sec(d*x+c))^(1/2)+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^4/d`

#### 3.131.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.66

$$\int \frac{1}{(b \sec(c + dx))^{7/2}} dx = \frac{\sqrt{b \sec(c + dx)} \left( 40 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 26 \sin(2(c + dx)) + 3 \right)}{84b^4d}$$

input `Integrate[(b*Sec[c + d*x])^(-7/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*b^4*d)`

**3.131.3 Rubi [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 4256, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(b \sec(c+dx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & \frac{5 \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{5 \left( \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \left( \frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{5 \left( \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{5 \left( \frac{\int \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}{3b^2} \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \\
& \quad \downarrow \text{3120} \\
& \frac{5 \left( \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}}
\end{aligned}$$

input `Int[(b*Sec[c + d*x])^(-7/2), x]`

output `(2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*sqrt[b*Sec[c + d*x]])))/(7*b^2)`

### 3.131.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

**3.131.4 Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.60

method	result
default	$-\frac{2\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\right)}{21d\sqrt{b\sec(dx+c)}b^3}$

input `int(1/(b*sec(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{-2/21/d/(b*\sec(d*x+c))^{1/2}/b^3*(5*I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+5*I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\operatorname{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I))*\sec(d*x+c)-3*\cos(d*x+c)^2*\sin(d*x+c)-5*\sin(d*x+c)}$$

**3.131.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00

$$\int \frac{1}{(b\sec(c+dx))^{7/2}} dx = \frac{2(3\cos(dx+c)^3 + 5\cos(dx+c))\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c) - 5i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + I\sin(dx+c))}{(b^4 d)}$$

input `integrate(1/(b*sec(d*x+c))^(7/2),x,algorithm="fricas")`

output 
$$\frac{1/21*(2*(3*\cos(d*x+c)^3 + 5*\cos(d*x+c))*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c) - 5*I*\sqrt{2}*\sqrt{b}*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x+c) + I*\sin(d*x+c)) + 5*I*\sqrt{2}*\sqrt{b}*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x+c) - I*\sin(d*x+c)))/(b^4*d)}$$



**3.131.6 Sympy [F]**

$$\int \frac{1}{(b \sec(c + dx))^{7/2}} dx = \int \frac{1}{(b \sec(c + dx))^{\frac{7}{2}}} dx$$

input `integrate(1/(b*sec(d*x+c))**(7/2), x)`

output `Integral((b*sec(c + d*x))**(-7/2), x)`

**3.131.7 Maxima [F]**

$$\int \frac{1}{(b \sec(c + dx))^{7/2}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{7}{2}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(7/2), x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(7/2), x)`

**3.131.8 Giac [F]**

$$\int \frac{1}{(b \sec(c + dx))^{7/2}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{7}{2}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(7/2), x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(7/2), x)`

**3.131.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(b \sec(c + dx))^{7/2}} dx = \int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{7/2}} dx$$

input `int(1/(b/cos(c + d*x))^(7/2),x)`output `int(1/(b/cos(c + d*x))^(7/2), x)`

### 3.132 $\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

3.132.1 Optimal result . . . . .	822
3.132.2 Mathematica [A] (verified) . . . . .	822
3.132.3 Rubi [A] (verified) . . . . .	823
3.132.4 Maple [A] (verified) . . . . .	824
3.132.5 Fricas [A] (verification not implemented) . . . . .	825
3.132.6 Sympy [F(-1)] . . . . .	826
3.132.7 Maxima [B] (verification not implemented) . . . . .	826
3.132.8 Giac [F] . . . . .	827
3.132.9 Mupad [F(-1)] . . . . .	827

#### 3.132.1 Optimal result

Integrand size = 23, antiderivative size = 107

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{3 \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{8d \sqrt{\sec(c + dx)}} + \frac{3 \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin(c + dx)}{8d} + \frac{\sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin(c + dx)}{4d}$$

output `3/8*sec(d*x+c)^(3/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+1/4*sec(d*x+c)^(7/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+3/8*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

#### 3.132.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.60

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} (3 \operatorname{arctanh}(\sin(c + dx)) + \sec(c + dx) (3 + 2 \sec^2(c + dx)) \tan(c + dx))}{8d \sqrt{\sec(c + dx)}}$$

input `Integrate[Sec[c + d*x]^(9/2)*Sqrt[b*Sec[c + d*x]],x]`

output  $(\text{Sqrt}[b \cdot \text{Sec}[c + d \cdot x]] \cdot (3 \cdot \text{ArcTanh}[\text{Sin}[c + d \cdot x]] + \text{Sec}[c + d \cdot x] \cdot (3 + 2 \cdot \text{Sec}[c + d \cdot x]^2) \cdot \text{Tan}[c + d \cdot x])) / (8 \cdot d \cdot \text{Sqrt}[\text{Sec}[c + d \cdot x]])$

### 3.132.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.78, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {2031, 3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{b \sec(c + dx)} \int \sec^5(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c + dx)} \int \csc(c + dx + \frac{\pi}{2})^5 dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \int \sec^3(c + dx) dx + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \int \csc(c + dx + \frac{\pi}{2})^3 dx + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \left( \frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \left( \frac{1}{2} \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

---

3.132.  $\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

$$\frac{\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right)}{\sqrt{\sec(c + dx)}}$$

input `Int[Sec[c + d*x]^(9/2)*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[b*Sec[c + d*x]]*((Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (3*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/4))/Sqrt[Sec[c + d*x]]`

### 3.132.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.132.4 Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.97

method	result
default	$\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^{\frac{9}{2}} \left( (3 \cos(dx+c)^2 + 2) \sin(dx+c) \cos(dx+c) + 3 \cos(dx+c)^5 \ln(\csc(dx+c) - \cot(dx+c) + 1) - 3 \cos(dx+c)^5 \ln(\dots) \right)}{8d}$
risch	$-\frac{i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (3 e^{6i(dx+c)} + 11 e^{4i(dx+c)} - 11 e^{2i(dx+c)} - 3)}{4(e^{2i(dx+c)}+1)^3 d} + \frac{3 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \ln(e^{i(dx+c)}+i) \cos(\dots)}{4d}$

3.132.  $\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

input `int(sec(d*x+c)^(9/2)*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/8/d*(b*sec(d*x+c))^(1/2)*sec(d*x+c)^(9/2)*((3*cos(d*x+c)^2+2)*sin(d*x+c)*cos(d*x+c)+3*cos(d*x+c)^5*ln(csc(d*x+c)-cot(d*x+c)+1)-3*cos(d*x+c)^5*ln(-cot(d*x+c)+csc(d*x+c)-1))`

### 3.132.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 229, normalized size of antiderivative = 2.14

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \left[ \frac{3 \sqrt{b} \cos(dx + c)^3 \log \left( -\frac{b \cos(dx+c)^2 - 2 \sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right) + \frac{2(3 \cos(dx+c)^2 + 2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{16 d \cos(dx + c)^3} - \frac{3 \sqrt{-b} \arctan \left( \frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b} \right) \cos(dx + c)^3 - \frac{(3 \cos(dx+c)^2 + 2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{8 d \cos(dx + c)^3} \right]$$

input `integrate(sec(d*x+c)^(9/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fracas")`

output `[1/16*(3*sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c)^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*(3*cos(d*x + c)^2 + 2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^3), -1/8*(3*sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c)^3 - (3*cos(d*x + c)^2 + 2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^3)]`

**3.132.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \text{Timed out}$$

```
input integrate(sec(d*x+c)**(9/2)*(b*sec(d*x+c))**(1/2),x)
```

```
output Timed out
```

**3.132.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1656 vs. 2(89) = 178.

Time = 0.48 (sec) , antiderivative size = 1656, normalized size of antiderivative = 15.48

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \text{Too large to display}$$

```
input integrate(sec(d*x+c)^(9/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")
```

```
output -1/16*(12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*
sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4
4*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*
x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(sin(8
*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c)
)*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(sin(8*d*x + 8
*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(1/
2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(2*(4*cos(6*d*x + 6*c)
+ 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*
x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6
*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c)
+ 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) +
3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8
*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16
*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*
x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arc
tan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c))) + 1) + 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d
*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + ...
```

**3.132.8 Giac [F]**

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^{\frac{9}{2}} dx$$

input `integrate(sec(d*x+c)^(9/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^(9/2), x)`

**3.132.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{\frac{b}{\cos(c + dx)}} \left( \frac{1}{\cos(c + dx)} \right)^{9/2} dx$$

input `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(9/2),x)`

output `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(9/2), x)`



### 3.133 $\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

3.133.1 Optimal result . . . . .	828
3.133.2 Mathematica [A] (verified) . . . . .	828
3.133.3 Rubi [A] (verified) . . . . .	829
3.133.4 Maple [A] (verified) . . . . .	830
3.133.5 Fricas [A] (verification not implemented) . . . . .	831
3.133.6 Sympy [F(-1)] . . . . .	831
3.133.7 Maxima [B] (verification not implemented) . . . . .	831
3.133.8 Giac [F] . . . . .	832
3.133.9 Mupad [B] (verification not implemented) . . . . .	832

#### 3.133.1 Optimal result

Integrand size = 23, antiderivative size = 70

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} \sin(c + dx)}{d} + \frac{\sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin^3(c + dx)}{3d}$$

output `1/3*sec(d*x+c)^(5/2)*sin(d*x+c)^3*(b*sec(d*x+c))^(1/2)/d+sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.133.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} (\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d \sqrt{\sec(c + dx)}}$$

input `Integrate[Sec[c + d*x]^(7/2)*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[b*Sec[c + d*x]]*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sqrt[Sec[c + d*x]])`

**3.133.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{7}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \sec^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\sqrt{b \sec(c+dx)} \int (\tan^2(c+dx)+1) d(-\tan(c+dx))}{d\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\left(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx)\right) \sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(7/2)*Sqrt[b*Sec[c + d*x]],x]`

output `-((Sqrt[b*Sec[c + d*x]]*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/(d*Sqrt[Sec[c + d*x]]))`

## 3.133.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

## 3.133.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69

method	result	size
default	$\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^{\frac{7}{2}} (2 \cos(dx+c)^2+1) \sin(dx+c) \cos(dx+c)}{3d}$	48
risch	$\frac{4i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} (4 \cos(dx+c)+2i \sin(dx+c))}{3(e^{2i(dx+c)+1})^2 d}$	89

input `int(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/3/d*(b*sec(d*x+c))^(1/2)*sec(d*x+c)^(7/2)*(2*cos(d*x+c)^2+1)*sin(d*x+c)*cos(d*x+c)`

**3.133.5 Fricas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.61

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{(2 \cos(dx + c)^2 + 1) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx + c)}{3 d \cos(dx + c)^{\frac{5}{2}}}$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/3*(2*cos(d*x + c)^2 + 1)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(5/2))`

**3.133.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(7/2)*(b*sec(d*x+c))**(1/2),x)`

output `Timed out`

**3.133.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. 2(60) = 120.

Time = 0.40 (sec) , antiderivative size = 294, normalized size of antiderivative = 4.20

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{4((3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + \cos(6 dx + 6 c)^2 + 6(3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + 3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + \cos(6 dx + 6 c)^2 + 6(3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + 3 \cos(2 dx + 2 c) + 1}{3(2(3 \cos(4 dx + 4 c) + 3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + \cos(6 dx + 6 c)^2 + 6(3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + 3 \cos(2 dx + 2 c) + 1)}$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output  $4/3*((3*\cos(2*d*x + 2*c) + 1)*\sin(6*d*x + 6*c) + 3*(3*\cos(2*d*x + 2*c) + 1)*\sin(4*d*x + 4*c) - 3*\cos(6*d*x + 6*c)*\sin(2*d*x + 2*c) - 9*\cos(4*d*x + 4*c)*\sin(2*d*x + 2*c))*\sqrt{b}/((2*(3*\cos(4*d*x + 4*c) + 3*\cos(2*d*x + 2*c) + 1)*\cos(6*d*x + 6*c) + \cos(6*d*x + 6*c)^2 + 6*(3*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + 9*\cos(4*d*x + 4*c)^2 + 9*\cos(2*d*x + 2*c)^2 + 6*(\sin(4*d*x + 4*c) + \sin(2*d*x + 2*c))*\sin(6*d*x + 6*c) + \sin(6*d*x + 6*c)^2 + 9*\sin(4*d*x + 4*c)^2 + 18*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 9*\sin(2*d*x + 2*c)^2 + 6*\cos(2*d*x + 2*c) + 1)*d)$

### 3.133.8 Giac [F]

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^{\frac{7}{2}} dx$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^(7/2), x)`

### 3.133.9 Mupad [B] (verification not implemented)

Time = 15.01 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.80

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{2 \cos(c + dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} (4 \sin(c + dx) + 5 \sin(3c + 3dx) + \sin(5c + 5dx) + \cos(c + dx))}{3d (10 \cos(c + dx) + 5 \cos(3c + 3dx) + \cos(5c + 5dx))}$$

input `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(7/2),x)`

output `(2*cos(c + d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*(cos(c + d*x)*10i + 4*sin(c + d*x) + cos(3*c + 3*d*x)*5i + cos(5*c + 5*d*x)*1i + 5*sin(3*c + 3*d*x) + sin(5*c + 5*d*x)))/(3*d*(10*cos(c + d*x) + 5*cos(3*c + 3*d*x) + cos(5*c + 5*d*x)))`

### 3.134 $\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

3.134.1 Optimal result . . . . .	833
3.134.2 Mathematica [A] (verified) . . . . .	833
3.134.3 Rubi [A] (verified) . . . . .	834
3.134.4 Maple [A] (verified) . . . . .	835
3.134.5 Fricas [A] (verification not implemented) . . . . .	836
3.134.6 Sympy [F(-1)] . . . . .	836
3.134.7 Maxima [B] (verification not implemented) . . . . .	837
3.134.8 Giac [F] . . . . .	838
3.134.9 Mupad [F(-1)] . . . . .	838

#### 3.134.1 Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{2d \sqrt{\sec(c + dx)}} + \frac{\sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin(c + dx)}{2d}$$

output `1/2*sec(d*x+c)^(3/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+1/2*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

#### 3.134.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.69

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} (\operatorname{arctanh}(\sin(c + dx)) + \sec(c + dx) \tan(c + dx))}{2d \sqrt{\sec(c + dx)}}$$

input `Integrate[Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[b*Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sqrt[Sec[c + d*x]])`

**3.134.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.79, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {2031, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[b*Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/Sqrt[Sec[c + d*x]]`

## 3.134.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_)+(d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

## 3.134.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.28

method	result
default	$-\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^{\frac{5}{2}} \left( \cos(dx+c)^3 \ln(-\cot(dx+c)+\csc(dx+c)-1) - \cos(dx+c)^3 \ln(\csc(dx+c)-\cot(dx+c)+1) - \cos(dx+c) \sin(dx+c) \right)}{2d}$
risch	$-i \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} (e^{2i(dx+c)} - 1)}{(e^{2i(dx+c)+1})d} - \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \ln(e^{i(dx+c)} - i) \cos(dx+c)}{d} + \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}$

input `int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2/d*(b*sec(d*x+c))^(1/2)*sec(d*x+c)^(5/2)*(cos(d*x+c)^3*ln(-cot(d*x+c)+csc(d*x+c)-1)-cos(d*x+c)^3*ln(csc(d*x+c)-cot(d*x+c)+1)-cos(d*x+c)*sin(d*x+c))`



**3.134.5 Fricas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 199, normalized size of antiderivative = 2.76

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \left[ \frac{\sqrt{b} \cos(dx + c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4 d \cos(dx + c)}, \right.$$

$$\left. - \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right) \cos(dx + c) - \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{2 d \cos(dx + c)} \right]$$

input `integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`output `[1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))]`**3.134.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(5/2)*(b*sec(d*x+c))**(1/2),x)`output `Timed out`

**3.134.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 661 vs.  $2(60) = 120$ .

Time = 0.46 (sec) , antiderivative size = 661, normalized size of antiderivative = 9.18

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx =$$

$$\frac{\left(4(\sin(4dx + 4c) + 2\sin(2dx + 2c)) \cos\left(\frac{3}{2} \arctan\left(\frac{\sin(2dx + 2c)}{\cos(2dx + 2c)}\right)\right) - 4(\sin(4dx + 4c) + 2\sin(2dx + 2c)) \cos\left(\frac{1}{2} \arctan\left(\frac{\sin(2dx + 2c)}{\cos(2dx + 2c)}\right)\right) - (2(2\cos(2dx + 2c) + 1)\cos(4dx + 4c) + \cos(4dx + 4c)^2 + 4\cos(2dx + 2c)^2 + \sin(4dx + 4c)^2 + 4\sin(4dx + 4c)\sin(2dx + 2c) + 4\sin(2dx + 2c)^2 + 4\cos(2dx + 2c) + 1) \log\left(\cos\left(\frac{1}{2} \arctan\left(\frac{\sin(2dx + 2c)}{\cos(2dx + 2c)}\right)\right), \cos(2dx + 2c)\right) + 1) + (2(2\cos(2dx + 2c) + 1)\cos(4dx + 4c) + \cos(4dx + 4c)^2 + 4\cos(2dx + 2c)^2 + \sin(4dx + 4c)^2 + 4\sin(4dx + 4c)\sin(2dx + 2c) + 4\sin(2dx + 2c)^2 + 4\cos(2dx + 2c) + 1) \log\left(\cos\left(\frac{1}{2} \arctan\left(\frac{\sin(2dx + 2c)}{\cos(2dx + 2c)}\right)\right), \cos(2dx + 2c)\right) + 1) - 4(\cos(4dx + 4c) + 2\cos(2dx + 2c) + 1)\sin\left(\frac{3}{2} \arctan\left(\frac{\sin(2dx + 2c)}{\cos(2dx + 2c)}\right)\right) + 4(\cos(4dx + 4c) + 2\cos(2dx + 2c) + 1)\sin\left(\frac{1}{2} \arctan\left(\frac{\sin(2dx + 2c)}{\cos(2dx + 2c)}\right)\right) \sqrt{b}}{(2(2\cos(2dx + 2c) + 1)\cos(4dx + 4c) + \cos(4dx + 4c)^2 + 4\cos(2dx + 2c)^2 + \sin(4dx + 4c)^2 + 4\sin(4dx + 4c)\sin(2dx + 2c) + 4\sin(2dx + 2c)^2 + 4\cos(2dx + 2c) + 1)dx}$$

```
input integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")
```

```
output -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)
```

**3.134.8 Giac [F]**

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^{\frac{5}{2}} dx$$

input `integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^(5/2), x)`

**3.134.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{\frac{b}{\cos(c + dx)}} \left( \frac{1}{\cos(c + dx)} \right)^{\frac{5}{2}} dx$$

input `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(5/2),x)`

output `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(5/2), x)`

### 3.135 $\int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

3.135.1 Optimal result . . . . .	839
3.135.2 Mathematica [A] (verified) . . . . .	839
3.135.3 Rubi [A] (verified) . . . . .	840
3.135.4 Maple [A] (verified) . . . . .	841
3.135.5 Fricas [A] (verification not implemented) . . . . .	841
3.135.6 Sympy [F] . . . . .	842
3.135.7 Maxima [A] (verification not implemented) . . . . .	842
3.135.8 Giac [F] . . . . .	842
3.135.9 Mupad [B] (verification not implemented) . . . . .	843

#### 3.135.1 Optimal result

Integrand size = 23, antiderivative size = 32

$$\int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} \sin(c + dx)}{d}$$

output `sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.135.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} \sin(c + dx)}{d}$$

input `Integrate[Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d`

**3.135.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \sec^2(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\sqrt{b \sec(c+dx)} \int 1d(-\tan(c+dx))}{d\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sin(c+dx) \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}{d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d`

**3.135.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

---

3.135.  $\int \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

### 3.135.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{\sec(dx+c)^{\frac{3}{2}} \sqrt{b \sec(dx+c)} \cos(dx+c) \sin(dx+c)}{d}$	35
risch	$\frac{2i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-i(dx+c)}}{d}$	67

```
input int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/d*sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2)*cos(d*x+c)*sin(d*x+c)
```

### 3.135.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94

$$\int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx + c)}{d \sqrt{\cos(dx + c)}}$$

```
input integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fracas")
```

```
output sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))
```

**3.135.6 Sympy [F]**

$$\int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(c + dx)} \sec^{\frac{3}{2}}(c + dx) dx$$

input `integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*sec(c + d*x)**(3/2), x)`

**3.135.7 Maxima [A] (verification not implemented)**

Time = 0.44 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.69

$$\begin{aligned} \int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx \\ = \frac{2\sqrt{b} \sin(2dx + 2c)}{(\cos(2dx + 2c)^2 + \sin(2dx + 2c)^2 + 2\cos(2dx + 2c) + 1)d} \end{aligned}$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `2*sqrt(b)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)`

**3.135.8 Giac [F]**

$$\int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx = \int \sqrt{b \sec(dx + c)} \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^(3/2), x)`

**3.135.9 Mupad [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.44

$$\int \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$$

$$= \frac{(\cos(dx) - \sin(dx) \operatorname{li}) (\sin(c) + \cos(c) \operatorname{li}) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}}}{d}$$

input `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(3/2),x)`output `((cos(d*x) - sin(d*x)*1i)*(cos(c)*1i + sin(c))*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2))/d`



### 3.136 $\int \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} dx$

3.136.1 Optimal result . . . . .	844
3.136.2 Mathematica [A] (verified) . . . . .	844
3.136.3 Rubi [A] (verified) . . . . .	845
3.136.4 Maple [A] (verified) . . . . .	846
3.136.5 Fricas [A] (verification not implemented) . . . . .	846
3.136.6 Sympy [F] . . . . .	847
3.136.7 Maxima [B] (verification not implemented) . . . . .	847
3.136.8 Giac [F] . . . . .	847
3.136.9 Mupad [F(-1)] . . . . .	848

#### 3.136.1 Optimal result

Integrand size = 23, antiderivative size = 33

$$\int \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} dx = \frac{\operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}}$$

output `arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

#### 3.136.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} dx = \frac{\operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}}$$

input `Integrate[Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]],x]`

output `(ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]])/(d*Sqrt[Sec[c + d*x]])`

### 3.136.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} dx$$

$$\downarrow \text{2031}$$

$$\frac{\sqrt{b \sec(c+dx)} \int \sec(c+dx) dx}{\sqrt{\sec(c+dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{b \sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2}) dx}{\sqrt{\sec(c+dx)}}$$

$$\downarrow \text{4257}$$

$$\frac{\operatorname{arctanh}(\sin(c+dx)) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}$$

input `Int[Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]],x]`

output `(ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]])/(d*Sqrt[Sec[c + d*x]])`

#### 3.136.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx.)*((a.)*(v.))(m.)((b.)*(v.))(n.), x_Symbol] := Simp[a(m + 1/2)*b(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c.) + (d.)*(x.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

**3.136.4 Maple [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

method	result	size
default	$-\frac{2\sqrt{b\sec(dx+c)} \operatorname{arctanh}(\cot(dx+c)-\csc(dx+c)) \cos(dx+c) \sqrt{\sec(dx+c)}}{d}$	46
risch	$\frac{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}} \ln(e^{i(dx+c)+i}) \cos(dx+c)}{d} - \frac{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}} \ln(e^{i(dx+c)-i}) \cos(dx+c)}{d}$	152

input `int(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`output `-2/d*(b*sec(d*x+c))^(1/2)*arctanh(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)*sec(d*x+c)^(1/2)`**3.136.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 111, normalized size of antiderivative = 3.36

$$\int \sqrt{\sec(c+dx)} \sqrt{b\sec(c+dx)} dx$$

$$= \left[ \frac{\sqrt{b} \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2d}, \right.$$

$$\left. - \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{d} \right]$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`output `[1/2*sqrt(b)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/d, -sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/d]`

**3.136.6 Sympy [F]**

$$\int \sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)} dx = \int \sqrt{b\sec(c+dx)}\sqrt{\sec(c+dx)} dx$$

input `integrate(sec(d*x+c)**(1/2)*(b*sec(d*x+c))**(1/2),x)`

output `Integral(sqrt(b*sec(c + d*x))*sqrt(sec(c + d*x)), x)`

**3.136.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs.  $2(29) = 58$ .

Time = 0.38 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.97

$$\int \sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)} dx$$

$$= \frac{\sqrt{b}(\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2\sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2\sin(dx+c) + 1))}{2d}$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `1/2*sqrt(b)*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/d`

**3.136.8 Giac [F]**

$$\int \sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)} dx = \int \sqrt{b\sec(dx+c)}\sqrt{\sec(dx+c)} dx$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))*sqrt(sec(d*x + c)), x)`

**3.136.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} dx = \int \sqrt{\frac{b}{\cos(c + dx)}} \sqrt{\frac{1}{\cos(c + dx)}} dx$$

input `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2),x)`output `int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2), x)`

$$3.137 \quad \int \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx$$

3.137.1 Optimal result . . . . .	849
3.137.2 Mathematica [A] (verified) . . . . .	849
3.137.3 Rubi [A] (verified) . . . . .	850
3.137.4 Maple [A] (verified) . . . . .	851
3.137.5 Fricas [A] (verification not implemented) . . . . .	851
3.137.6 Sympy [A] (verification not implemented) . . . . .	852
3.137.7 Maxima [A] (verification not implemented) . . . . .	852
3.137.8 Giac [F] . . . . .	852
3.137.9 Mupad [B] (verification not implemented) . . . . .	853

### 3.137.1 Optimal result

Integrand size = 23, antiderivative size = 24

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx = \frac{x \sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

output `x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

### 3.137.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx = \frac{x \sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]/Sqrt[Sec[c + d*x]],x]`

output `(x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]`

**3.137.3 Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2031, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}} dx$$

↓ 2031

$$\frac{\sqrt{b \sec(c + dx)} \int 1 dx}{\sqrt{\sec(c + dx)}}$$

↓ 24

$$\frac{x \sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}}$$

input `Int[Sqrt[b*Sec[c + d*x]]/Sqrt[Sec[c + d*x]],x]`

output `(x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]`

**3.137.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

**3.137.4 Maple [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{(dx+c)\sqrt{b\sec(dx+c)}}{d\sqrt{\sec(dx+c)}}$	28
risch	$\frac{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} x}}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}$	54

input `int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`output `1/d*(d*x+c)*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`**3.137.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 98, normalized size of antiderivative = 4.08

$$\int \frac{\sqrt{b\sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx$$

$$= \left[ \frac{\sqrt{-b} \log \left( -2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b \right)}{2d}, \frac{\sqrt{b} \arctan \left( \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{b} \sqrt{\cos(dx+c)}} \right)}{d} \right]$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorithm="fracas")`output `[1/2*sqrt(-b)*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/d, sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/d]`



**3.137.6 Sympy [A] (verification not implemented)**

Time = 1.81 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}} dx = \frac{x \sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}}$$

input `integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(1/2),x)`output `x*sqrt(b*sec(c + d*x))/sqrt(sec(c + d*x))`**3.137.7 Maxima [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}} dx = \frac{2 \sqrt{b} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`output `2*sqrt(b)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d`**3.137.8 Giac [F]**

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}} dx = \int \frac{\sqrt{b \sec(dx + c)}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorithm="giac")`output `integrate(sqrt(b*sec(d*x + c))/sqrt(sec(d*x + c)), x)`

**3.137.9 Mupad [B] (verification not implemented)**

Time = 12.52 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}} dx = \frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{\sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(1/2),x)`

output `(x*(b/cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2)`

**3.138**  $\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx$

3.138.1 Optimal result . . . . . 854  
 3.138.2 Mathematica [A] (verified) . . . . . 854  
 3.138.3 Rubi [A] (verified) . . . . . 855  
 3.138.4 Maple [A] (verified) . . . . . 856  
 3.138.5 Fricas [A] (verification not implemented) . . . . . 856  
 3.138.6 Sympy [A] (verification not implemented) . . . . . 857  
 3.138.7 Maxima [A] (verification not implemented) . . . . . 857  
 3.138.8 Giac [A] (verification not implemented) . . . . . 857  
 3.138.9 Mupad [B] (verification not implemented) . . . . . 858

**3.138.1 Optimal result**

Integrand size = 23, antiderivative size = 32

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sec(c + dx)}}$$

output `sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

**3.138.2 Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{\sqrt{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sec(c + dx)}}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(3/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])`

**3.138.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx$$

$$\downarrow \text{2031}$$

$$\frac{\sqrt{b \sec(c+dx)} \int \cos(c+dx) dx}{\sqrt{\sec(c+dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{b \sec(c+dx)} \int \sin\left(c+dx+\frac{\pi}{2}\right) dx}{\sqrt{\sec(c+dx)}}$$

$$\downarrow \text{3117}$$

$$\frac{\sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}$$

input `Int[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(3/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])`

**3.138.3.1 Defintions of rubi rules used**

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3117 Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

### 3.138.4 Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sqrt{b \sec(dx+c)} \tan(dx+c)}{d \sec(dx+c)^{\frac{3}{2}}}$	29
risch	$-\frac{i \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} e^{i(dx+c)}}}{2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{i \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} e^{-i(dx+c)}}}{2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}$	134

```
input int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/d*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2)*tan(d*x+c)
```

### 3.138.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{d}$$

```
input integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorithm="fricas")
```

```
output sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/d
```

**3.138.6 Sympy [A] (verification not implemented)**

Time = 3.65 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.44

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx = \begin{cases} \frac{\sqrt{b \sec(c+dx)} \tan(c+dx)}{d \sec^{\frac{3}{2}}(c+dx)} & \text{for } d \neq 0 \\ \frac{x \sqrt{b \sec(c)}}{\sec^{\frac{3}{2}}(c)} & \text{otherwise} \end{cases}$$

input `integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(3/2),x)`output `Piecewise((sqrt(b*sec(c + d*x))*tan(c + d*x)/(d*sec(c + d*x)**(3/2)), Ne(d, 0)), (x*sqrt(b*sec(c))/sec(c)**(3/2), True))`**3.138.7 Maxima [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.41

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx = \frac{\sqrt{b} \sin(dx+c)}{d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")`output `sqrt(b)*sin(d*x + c)/d`**3.138.8 Giac [A] (verification not implemented)**

Time = 0.86 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.19

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx = \frac{2\sqrt{b} \operatorname{sgn}(\cos(dx+c)) \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{d \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorithm="giac")`output `2*sqrt(b)*sgn(cos(d*x + c))*tan(1/2*d*x + 1/2*c)/(d*tan(1/2*d*x + 1/2*c)^2 + d)`

**3.138.9 Mupad [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx = \frac{\sin(c+dx) \sqrt{\frac{b}{\cos(c+dx)}}}{d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(3/2),x)`output `(sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(d*(1/cos(c + d*x))^(1/2))`

$$3.139 \quad \int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx$$

3.139.1 Optimal result . . . . .	859
3.139.2 Mathematica [A] (verified) . . . . .	859
3.139.3 Rubi [A] (verified) . . . . .	860
3.139.4 Maple [A] (verified) . . . . .	861
3.139.5 Fricas [A] (verification not implemented) . . . . .	862
3.139.6 Sympy [A] (verification not implemented) . . . . .	862
3.139.7 Maxima [A] (verification not implemented) . . . . .	863
3.139.8 Giac [B] (verification not implemented) . . . . .	863
3.139.9 Mupad [B] (verification not implemented) . . . . .	864

### 3.139.1 Optimal result

Integrand size = 23, antiderivative size = 63

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx = \frac{x \sqrt{b \sec(c+dx)}}{2 \sqrt{\sec(c+dx)}} + \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{2d \sec^{\frac{3}{2}}(c+dx)}$$

output `1/2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+1/2*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

### 3.139.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx = \frac{\sqrt{b \sec(c+dx)}(2(c+dx) + \sin(2(c+dx)))}{4d \sqrt{\sec(c+dx)}}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(5/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sqrt[Sec[c + d*x]])`

---

3.139.  $\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx$



**3.139.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.76, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \cos^2(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^2 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right)}{\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(5/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/Sqrt[Sec[c + d*x]]`

3.139.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_)*((a_)*(v_)^(m_))*((b_)*(v_)^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.139.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

method	result	size
default	$\frac{\sqrt{b \sec(dx+c)} \left( \tan(dx+c) + (dx+c) \sec(dx+c)^2 \right)}{2d \sec(dx+c)^{\frac{5}{2}}}$	45
risch	$\frac{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} x}{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{i\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{2i(dx+c)}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}} + \frac{i\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-2i(dx+c)}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}}$	188

input `int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2),x,method=_RETURNVERBOSE)`

output `1/2/d*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2)*(tan(d*x+c)+(d*x+c)*sec(d*x+c)^2)`

**3.139.5 Fricas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 158, normalized size of antiderivative = 2.51

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{\left[ 2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + \sqrt{-b} \log \left( -2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c) \right) \right]}{4d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2),x, algorithm="fricas")`output `[1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(-b)*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/d]`**3.139.6 Sympy [A] (verification not implemented)**

Time = 9.81 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.70

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx$$

$$= \begin{cases} \frac{x \sqrt{b \sec(c+dx)} \tan^2(c+dx)}{2 \sec^{\frac{5}{2}}(c+dx)} + \frac{x \sqrt{b \sec(c+dx)}}{2 \sec^{\frac{5}{2}}(c+dx)} + \frac{\sqrt{b \sec(c+dx)} \tan(c+dx)}{2d \sec^{\frac{5}{2}}(c+dx)} & \text{for } d \neq 0 \\ \frac{x \sqrt{b \sec(c)}}{\sec^{\frac{5}{2}}(c)} & \text{otherwise} \end{cases}$$

input `integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(5/2),x)`output `Piecewise((x*sqrt(b*sec(c + d*x))*tan(c + d*x)**2/(2*sec(c + d*x)**(5/2)) + x*sqrt(b*sec(c + d*x))/(2*sec(c + d*x)**(5/2)) + sqrt(b*sec(c + d*x))*tan(c + d*x)/(2*d*sec(c + d*x)**(5/2)), Ne(d, 0)), (x*sqrt(b*sec(c))/sec(c)**(5/2), True))`

**3.139.7 Maxima [A] (verification not implemented)**

Time = 0.44 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.40

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{(2 dx + 2 c + \sin(2 dx + 2 c))\sqrt{b}}{4 d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2),x, algorithm="maxima")`output `1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))*sqrt(b)/d`**3.139.8 Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(51) = 102.

Time = 1.32 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.22

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sec^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{b} dx \operatorname{sgn}(\cos(dx + c)) \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 2 \sqrt{b} dx \operatorname{sgn}(\cos(dx + c)) \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 2 \sqrt{b} \operatorname{sgn}(\cos(dx + c))}{2 \left(d \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 2 d \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + d\right)}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2),x, algorithm="giac")`output `1/2*(sqrt(b)*d*x*sgn(cos(d*x + c))*tan(1/2*d*x + 1/2*c)^4 + 2*sqrt(b)*d*x*sgn(cos(d*x + c))*tan(1/2*d*x + 1/2*c)^2 - 2*sqrt(b)*sgn(cos(d*x + c))*tan(1/2*d*x + 1/2*c)^3 + sqrt(b)*d*x*sgn(cos(d*x + c)) + 2*sqrt(b)*sgn(cos(d*x + c))*tan(1/2*d*x + 1/2*c))/(d*tan(1/2*d*x + 1/2*c)^4 + 2*d*tan(1/2*d*x + 1/2*c)^2 + d)`

**3.139.9 Mupad [B] (verification not implemented)**

Time = 12.70 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.65

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx = \frac{(\sin(2c+2dx) + 2dx) \sqrt{\frac{b}{\cos(c+dx)}}}{4d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(5/2),x)`output `((sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*d*(1/cos(c + d*x))^(1/2))`

**3.140**       $\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx$

3.140.1 Optimal result . . . . .	865
3.140.2 Mathematica [A] (verified) . . . . .	865
3.140.3 Rubi [A] (verified) . . . . .	866
3.140.4 Maple [A] (verified) . . . . .	867
3.140.5 Fricas [A] (verification not implemented) . . . . .	868
3.140.6 Sympy [F(-1)] . . . . .	868
3.140.7 Maxima [A] (verification not implemented) . . . . .	868
3.140.8 Giac [F] . . . . .	869
3.140.9 Mupad [B] (verification not implemented) . . . . .	869

**3.140.1 Optimal result**

Integrand size = 23, antiderivative size = 70

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx = \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{d \sqrt{\sec(c+dx)}} - \frac{\sqrt{b \sec(c+dx)} \sin^3(c+dx)}{3d \sqrt{\sec(c+dx)}}$$

output `sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)-1/3*sin(d*x+c)^3*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

**3.140.2 Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx = \frac{(5 + \cos(2(c+dx)))\sqrt{b \sec(c+dx)} \sin(c+dx)}{6d \sqrt{\sec(c+dx)}}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(7/2),x]`

output `((5 + Cos[2*(c + d*x)])*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(6*d*Sqrt[Sec[c + d*x]])`

**3.140.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \cos^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^3 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3113} \\
 & - \frac{\sqrt{b \sec(c+dx)} \int (1 - \sin^2(c+dx)) d(-\sin(c+dx))}{d\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{2009} \\
 & - \frac{(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx)) \sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(7/2),x]`

output `-((Sqrt[b*Sec[c + d*x]]*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/(d*Sqrt[Sec[c + d*x]]))`

## 3.140.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

## 3.140.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{\sqrt{b \sec(dx+c)} (\tan(dx+c) + 2 \tan(dx+c) \sec(dx+c)^2)}{3d \sec(dx+c)^{\frac{7}{2}}}$	47
risch	$-\frac{3i \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{3i \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sin(3dx+3c)}{12 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}$	199

input `int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2),x,method=_RETURNVERBOSE)`

output `1/3/d*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2)*(tan(d*x+c)+2*tan(d*x+c)*sec(d*x+c)^2)`



**3.140.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sec^{\frac{7}{2}}(c + dx)} dx = \frac{(\cos(dx + c))^3 + 2 \cos(dx + c)}{3d\sqrt{\cos(dx + c)}} \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c)$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2),x, algorithm="fricas")`output `1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))`**3.140.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sec^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(7/2),x)`output `Timed out`**3.140.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.60

$$\int \frac{\sqrt{b \sec(c + dx)}}{\sec^{\frac{7}{2}}(c + dx)} dx = \frac{\sqrt{b}(\sin(3dx + 3c) + 9 \sin(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))))}{12d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2),x, algorithm="maxima")`output `1/12*sqrt(b)*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/d`

---

3.140.  $\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx$

**3.140.8 Giac [F]**

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx = \int \frac{\sqrt{b \sec(dx+c)}}{\sec^{\frac{7}{2}}(dx+c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate(sqrt(b*sec(d*x + c))/sec(d*x + c)^(7/2), x)`

**3.140.9 Mupad [B] (verification not implemented)**

Time = 12.83 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx = \frac{(9 \sin(c+dx) + \sin(3c+3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(7/2),x)`

output `((9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*d*(1/cos(c + d*x))^(1/2))`

**3.141** 
$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$$

3.141.1 Optimal result . . . . . 870  
 3.141.2 Mathematica [A] (verified) . . . . . 870  
 3.141.3 Rubi [A] (verified) . . . . . 871  
 3.141.4 Maple [A] (verified) . . . . . 872  
 3.141.5 Fricas [A] (verification not implemented) . . . . . 873  
 3.141.6 Sympy [F(-1)] . . . . . 873  
 3.141.7 Maxima [A] (verification not implemented) . . . . . 874  
 3.141.8 Giac [F] . . . . . 874  
 3.141.9 Mupad [B] (verification not implemented) . . . . . 874

**3.141.1 Optimal result**

Integrand size = 23, antiderivative size = 98

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx = \frac{3x\sqrt{b \sec(c+dx)}}{8\sqrt{\sec(c+dx)}} + \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{4d \sec^{\frac{7}{2}}(c+dx)} + \frac{3\sqrt{b \sec(c+dx)} \sin(c+dx)}{8d \sec^{\frac{3}{2}}(c+dx)}$$

output `1/4*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(7/2)+3/8*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+3/8*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

**3.141.2 Mathematica [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.56

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx = \frac{\sqrt{b \sec(c+dx)}(12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx)))}{32d\sqrt{\sec(c+dx)}}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(9/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)]))/(32*d*Sqrt[Sec[c + d*x]])`

---

3.141. 
$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$$

**3.141.3 Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.76, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {2031, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \cos^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^4 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \int \cos^2(c+dx) dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \left( \frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sqrt{b \sec(c+dx)} \left( \frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left( \frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right)}{\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(9/2),x]`

---

3.141.  $\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$

output  $(\text{Sqrt}[b \cdot \text{Sec}[c + d \cdot x]] \cdot ((\text{Cos}[c + d \cdot x]^3 \cdot \text{Sin}[c + d \cdot x]) / (4 \cdot d) + (3 \cdot (x/2 + (\text{Cos}[c + d \cdot x] \cdot \text{Sin}[c + d \cdot x]) / (2 \cdot d)))) / 4) / \text{Sqrt}[\text{Sec}[c + d \cdot x]]$

### 3.141.3.1 Defintions of rubi rules used

rule 24  $\text{Int}[a\_ , x\_ \text{Symbol}] \text{:>} \text{Simp}[a \cdot x, x] \text{ /; FreeQ}[a, x]$

rule 2031  $\text{Int}[(\text{Fx}\_.) \cdot ((a\_.) \cdot (v\_))^m \cdot ((b\_.) \cdot (v\_))^n, x\_ \text{Symbol}] \text{:>} \text{Simp}[a^{m+1/2} \cdot b^{n-1/2} \cdot (\text{Sqrt}[b \cdot v] / \text{Sqrt}[a \cdot v]) \text{ Int}[v^{m+n} \cdot \text{Fx}, x], x] \text{ /; FreeQ}[\{a, b, m\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n + 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

rule 3042  $\text{Int}[u\_ , x\_ \text{Symbol}] \text{:>} \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3115  $\text{Int}[(b\_.) \cdot \text{sin}[c\_.] + (d\_.) \cdot (x\_)]^n, x\_ \text{Symbol}] \text{:>} \text{Simp}[(-b) \cdot \text{Cos}[c + d \cdot x] \cdot ((b \cdot \text{Sin}[c + d \cdot x])^{n-1} / (d \cdot n)), x] + \text{Simp}[b^{2 \cdot (n-1)/n} \text{ Int}[(b \cdot \text{Sin}[c + d \cdot x])^{n-2}, x], x] \text{ /; FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2 \cdot n]$

### 3.141.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{\sqrt{b \sec(dx+c)} \left( 2 \tan(dx+c) + 3 \tan(dx+c) \sec(dx+c)^2 + 3(dx+c) \sec(dx+c)^4 \right)}{8d \sec(dx+c)^{\frac{9}{2}}}$	64
risch	$\frac{3 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} x}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{i \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{2i(dx+c)}}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}} + \frac{i \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-2i(dx+c)}}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}} + \frac{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sin(4dx+4c)}}{32 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}}$	253

input  $\text{int}((b \cdot \text{sec}(d \cdot x + c))^{1/2} / \text{sec}(d \cdot x + c)^{9/2}, x, \text{method} = \_ \text{RETURNVERBOSE})$

output  $1/8/d \cdot (b \cdot \text{sec}(d \cdot x + c))^{1/2} / \text{sec}(d \cdot x + c)^{9/2} \cdot (2 \cdot \tan(d \cdot x + c) + 3 \cdot \tan(d \cdot x + c) \cdot \text{sec}(d \cdot x + c)^2 + 3 \cdot (d \cdot x + c) \cdot \text{sec}(d \cdot x + c)^4)$

3.141.  $\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$

**3.141.5 Fricas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 202, normalized size of antiderivative = 2.06

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{2 \left( 2 \cos(dx+c)^4 + 3 \cos(dx+c)^2 \right) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c) + 3 \sqrt{-b} \log \left( -2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2 \right)}{16 d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(9/2),x, algorithm="fricas")`

output `[1/16*(2*(2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) + 3*sqrt(-b)*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/8*((2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) + 3*sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/d]`

**3.141.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(9/2),x)`

output `Timed out`

**3.141.7 Maxima [A] (verification not implemented)**

Time = 0.43 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.50

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx = \frac{(12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(\frac{1}{2} \arctan(\sin(4 dx + 4 c), \cos(4 dx + 4 c)))) \sqrt{b}}{32 d}$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(9/2),x, algorithm="maxima")`output `1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d`**3.141.8 Giac [F]**

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx = \int \frac{\sqrt{b \sec(dx+c)}}{\sec^{\frac{9}{2}}(dx+c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(9/2),x, algorithm="giac")`output `integrate(sqrt(b*sec(d*x + c))/sec(d*x + c)^(9/2), x)`**3.141.9 Mupad [B] (verification not implemented)**

Time = 12.77 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.53

$$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx = \frac{\sqrt{\frac{b}{\cos(c+dx)}} (8 \sin(2c + 2dx) + \sin(4c + 4dx) + 12dx)}{32d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(9/2),x)`output `((b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))/(32*d*(1/cos(c + d*x))^(1/2))`

---

3.141.  $\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$

### 3.142 $\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx$

3.142.1 Optimal result . . . . .	875
3.142.2 Mathematica [A] (verified) . . . . .	875
3.142.3 Rubi [A] (verified) . . . . .	876
3.142.4 Maple [A] (verified) . . . . .	877
3.142.5 Fricas [A] (verification not implemented) . . . . .	878
3.142.6 Sympy [F(-1)] . . . . .	878
3.142.7 Maxima [B] (verification not implemented) . . . . .	879
3.142.8 Giac [F] . . . . .	879
3.142.9 Mupad [F(-1)] . . . . .	880

#### 3.142.1 Optimal result

Integrand size = 23, antiderivative size = 110

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{3b \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{8d \sqrt{\sec(c + dx)}} + \frac{3b \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin(c + dx)}{8d} + \frac{b \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin(c + dx)}{4d}$$

```
output 3/8*b*sec(d*x+c)^(3/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+1/4*b*sec(d*x+c)^(7/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+3/8*b*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)
```

#### 3.142.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.58

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{(b \sec(c + dx))^{3/2} (3 \operatorname{arctanh}(\sin(c + dx)) + \sec(c + dx) (3 + 2 \sec^2(c + dx)) \tan(c + dx))}{8d \sec^{\frac{3}{2}}(c + dx)}$$

```
input Integrate[Sec[c + d*x]^(7/2)*(b*Sec[c + d*x])^(3/2),x]
```

```
output ((b*Sec[c + d*x])^(3/2)*(3*ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*(3 + 2*Sec[c + d*x]^2)*Tan[c + d*x]))/(8*d*Sec[c + d*x]^(3/2))
```



**3.142.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.76, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {2031, 3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{7}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \sec^5(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2})^5 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \int \sec^3(c+dx) dx + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \int \csc(c+dx + \frac{\pi}{2})^3 dx + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \left( \frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \left( \frac{1}{2} \int \csc(c+dx + \frac{\pi}{2}) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{3}{4} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right)}{\sqrt{\sec(c+dx)}}
 \end{aligned}$$

---

3.142.  $\int \sec^{\frac{7}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx$

input `Int[Sec[c + d*x]^(7/2)*(b*Sec[c + d*x])^(3/2),x]`

output `(b*Sqrt[b*Sec[c + d*x]]*((Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (3*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/4))/Sqrt[Sec[c + d*x]]`

### 3.142.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.142.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.95

method	result
default	$\frac{\sqrt{b \sec(dx+c)} b \sec(dx+c)^{\frac{9}{2}} \left( (3 \cos(dx+c)^2 + 2) \sin(dx+c) \cos(dx+c) + 3 \cos(dx+c)^5 \ln(\csc(dx+c) - \cot(dx+c) + 1) - 3 \cos(dx+c)^5 \ln \right)}{8d}$
risch	$-\frac{ib \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (3e^{6i(dx+c)} + 11e^{4i(dx+c)} - 11e^{2i(dx+c)} - 3)}{4(e^{2i(dx+c)}+1)^3 d} + \frac{3b \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \ln(e^{i(dx+c)}+i)}{4d}$

input `int(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

$$3.142. \quad \int \sec^{\frac{7}{2}}(c + dx) (b \sec(c + dx))^{3/2} dx$$

output  $1/8/d*(b*\sec(d*x+c))^{(1/2)}*b*\sec(d*x+c)^{(9/2)}*((3*\cos(d*x+c)^2+2)*\sin(d*x+c)*\cos(d*x+c)+3*\cos(d*x+c)^5*\ln(\csc(d*x+c)-\cot(d*x+c)+1)-3*\cos(d*x+c)^5*\ln(-\cot(d*x+c)+\csc(d*x+c)-1))$

### 3.142.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 236, normalized size of antiderivative = 2.15

$$\int \sec^{\frac{7}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx = \left[ \frac{3 b^{\frac{3}{2}} \cos(dx+c)^3 \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b}\sqrt{\frac{b}{\cos(dx+c)}}\sqrt{\cos(dx+c)}\sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2(3b \cos(dx+c)^2 + 2b)\sqrt{\cos(dx+c)}}{\sqrt{\cos(dx+c)}}}{16 d \cos(dx+c)^3} - \frac{3\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}}\sqrt{\cos(dx+c)}\sin(dx+c)}{b}\right) \cos(dx+c)^3 - \frac{(3b \cos(dx+c)^2 + 2b)\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{\sqrt{\cos(dx+c)}}}{8 d \cos(dx+c)^3} \right]$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output  $[1/16*(3*b^{(3/2)}*\cos(d*x+c)^3*\log(-(b*\cos(d*x+c)^2-2*\sqrt{b})*\sqrt{b/\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-2*b)/\cos(d*x+c)^2)+2*(3*b*\cos(d*x+c)^2+2*b)*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c)/\sqrt{\cos(d*x+c)}}/(d*\cos(d*x+c)^3), -1/8*(3*\sqrt{-b})*b*\arctan(\sqrt{-b}*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c)/b)*\cos(d*x+c)^3 - (3*b*\cos(d*x+c)^2+2*b)*\sqrt{b/\cos(d*x+c)}*\sin(d*x+c)/\sqrt{\cos(d*x+c)}}/(d*\cos(d*x+c)^3)]$

### 3.142.6 Sympy [F(-1)]

Timed out.

$$\int \sec^{\frac{7}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(7/2)*(b*sec(d*x+c))**(3/2),x)`

output Timed out

---

3.142.  $\int \sec^{\frac{7}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx$

**3.142.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1742 vs.  $2(92) = 184$ .

Time = 0.49 (sec) , antiderivative size = 1742, normalized size of antiderivative = 15.84

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \text{Too large to display}$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output

```
-1/16*(12*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c)
+ 4*b*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*
c))) + 44*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c)
+ 4*b*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*
c))) - 44*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c)
+ 4*b*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*
c))) - 12*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c)
+ 4*b*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*
c))) - 3*(b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36*b*cos(4*d*x
+ 4*c)^2 + 16*b*cos(2*d*x + 2*c)^2 + b*sin(8*d*x + 8*c)^2 + 16*b*sin(6*d*x
+ 6*c)^2 + 36*b*sin(4*d*x + 4*c)^2 + 48*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*
c) + 16*b*sin(2*d*x + 2*c)^2 + 2*(4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*
*c) + 4*b*cos(2*d*x + 2*c) + b)*cos(8*d*x + 8*c) + 8*(6*b*cos(4*d*x + 4*c)
+ 4*b*cos(2*d*x + 2*c) + b)*cos(6*d*x + 6*c) + 12*(4*b*cos(2*d*x + 2*c) +
b)*cos(4*d*x + 4*c) + 8*b*cos(2*d*x + 2*c) + 4*(2*b*sin(6*d*x + 6*c) + 3*
b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b*sin(
4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b*log(cos(1/2*arc
tan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c)))) + 1) + 3*(b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36...
```

**3.142.8 Giac [F]**

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^{\frac{7}{2}} dx$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(7/2), x)`

---

3.142.  $\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx$

**3.142.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{3/2} \left( \frac{1}{\cos(c + dx)} \right)^{7/2} dx$$

input `int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(7/2),x)`output `int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(7/2), x)`

### 3.143 $\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx$

3.143.1 Optimal result . . . . .	881
3.143.2 Mathematica [A] (verified) . . . . .	881
3.143.3 Rubi [A] (verified) . . . . .	882
3.143.4 Maple [A] (verified) . . . . .	883
3.143.5 Fricas [A] (verification not implemented) . . . . .	884
3.143.6 Sympy [F(-1)] . . . . .	884
3.143.7 Maxima [B] (verification not implemented) . . . . .	884
3.143.8 Giac [F] . . . . .	885
3.143.9 Mupad [B] (verification not implemented) . . . . .	885

#### 3.143.1 Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{b\sqrt{\sec(c + dx)}\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} + \frac{b \sec^{\frac{5}{2}}(c + dx)\sqrt{b \sec(c + dx)} \sin^3(c + dx)}{3d}$$

output `1/3*b*sec(d*x+c)^(5/2)*sin(d*x+c)^3*(b*sec(d*x+c))^(1/2)/d+b*sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.143.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{(b \sec(c + dx))^{3/2} (\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d \sec^{\frac{3}{2}}(c + dx)}$$

input `Integrate[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2),x]`

output `((b*Sec[c + d*x])^(3/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sec[c + d*x]^(3/2))`

**3.143.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \sec^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2})^4 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & -\frac{b\sqrt{b \sec(c+dx)} \int (\tan^2(c+dx) + 1) d(-\tan(c+dx))}{d\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{b(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx)) \sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2),x]`

output `-((b*Sqrt[b*Sec[c + d*x]]*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/(d*Sqrt[Sec[c + d*x]]))`

## 3.143.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

## 3.143.4 Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.68

method	result	size
default	$\frac{b \sec(dx+c)^{\frac{7}{2}} \sqrt{b \sec(dx+c)} (2 \cos(dx+c)^2 + 1) \sin(dx+c) \cos(dx+c)}{3d}$	49
risch	$\frac{4ib \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (4 \cos(dx+c) + 2i \sin(dx+c))}{3(e^{2i(dx+c)}+1)^2 d}$	90

input `int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `1/3/d*b*sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2)*(2*cos(d*x+c)^2+1)*sin(d*x+c)*cos(d*x+c)`



**3.143.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.61

$$\int \sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx = \frac{(2b \cos(dx+c)^2 + b) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{3d \cos(dx+c)^{\frac{5}{2}}}$$

input `integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2),x, algorithm="fracas")`

output `1/3*(2*b*cos(d*x + c)^2 + b)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(5/2))`

**3.143.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(5/2)*(b*sec(d*x+c))**(3/2),x)`

output `Timed out`

**3.143.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 299 vs.  $2(62) = 124$ .

Time = 0.40 (sec) , antiderivative size = 299, normalized size of antiderivative = 4.15

$$\int \sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx =$$

$$\frac{4(3b \cos(6dx) - 3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) -$$

input `integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output 
$$-4/3*(3*b*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) + 9*b*cos(4*d*x + 4*c)*sin(2*d*x + 2*c) - (3*b*cos(2*d*x + 2*c) + b)*sin(6*d*x + 6*c) - 3*(3*b*cos(2*d*x + 2*c) + b)*sin(4*d*x + 4*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d$$

### 3.143.8 Giac [F]

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^{\frac{5}{2}} dx$$

input `integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(5/2), x)`

### 3.143.9 Mupad [B] (verification not implemented)

Time = 13.84 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.76

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \frac{2b \cos(c + dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} (4 \sin(c + dx) + 5 \sin(3c + 3dx) + \sin(5c + 5dx))}{3d(10 \cos(c + dx) + 5 \cos(3c + 3dx) + \cos(5c + 5dx))}$$

input `int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(5/2),x)`

output 
$$(2*b*cos(c + d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*(cos(c + d*x)*10i + 4*sin(c + d*x) + cos(3*c + 3*d*x)*5i + cos(5*c + 5*d*x)*1i + 5*sin(3*c + 3*d*x) + sin(5*c + 5*d*x)))/(3*d*(10*cos(c + d*x) + 5*cos(3*c + 3*d*x) + cos(5*c + 5*d*x)))$$

### 3.144 $\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{\frac{3}{2}} dx$

3.144.1 Optimal result . . . . .	886
3.144.2 Mathematica [A] (verified) . . . . .	886
3.144.3 Rubi [A] (verified) . . . . .	887
3.144.4 Maple [A] (verified) . . . . .	888
3.144.5 Fricas [A] (verification not implemented) . . . . .	889
3.144.6 Sympy [F] . . . . .	889
3.144.7 Maxima [B] (verification not implemented) . . . . .	890
3.144.8 Giac [F] . . . . .	890
3.144.9 Mupad [F(-1)] . . . . .	891

#### 3.144.1 Optimal result

Integrand size = 23, antiderivative size = 74

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{\frac{3}{2}} dx = \frac{\operatorname{arctanh}(\sin(c + dx))\sqrt{b \sec(c + dx)}}{2d\sqrt{\sec(c + dx)}} + \frac{b \sec^{\frac{3}{2}}(c + dx)\sqrt{b \sec(c + dx)} \sin(c + dx)}{2d}$$

output `1/2*b*sec(d*x+c)^(3/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+1/2*b*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

#### 3.144.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.68

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{\frac{3}{2}} dx = \frac{(b \sec(c + dx))^{\frac{3}{2}}(\operatorname{arctanh}(\sin(c + dx)) + \sec(c + dx) \tan(c + dx))}{2d \sec^{\frac{3}{2}}(c + dx)}$$

input `Integrate[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2),x]`

output `((b*Sec[c + d*x])^(3/2)*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sec[c + d*x]^(3/2))`

**3.144.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.78, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {2031, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{b\sqrt{b \sec(c+dx)} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2),x]`

output `(b*Sqrt[b*Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/Sqrt[Sec[c + d*x]]`

### 3.144.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.144.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.26

method	result
default	$-\frac{\sqrt{b \sec(dx+c)} b \sec(dx+c)^{\frac{5}{2}} \left( \cos(dx+c)^3 \ln(-\cot(dx+c)+\csc(dx+c)-1) - \cos(dx+c)^3 \ln(\csc(dx+c)-\cot(dx+c)+1) - \cos(dx+c) \sin(dx+c) \right)}{2d}$
risch	$-\frac{ib \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} (e^{2i(dx+c)}-1)}{(e^{2i(dx+c)}+1)d} + \frac{b \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \ln(e^{i(dx+c)}+i) \cos(dx+c)}{d} - \frac{b \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}{\sqrt{e^{2i(dx+c)}+1}}$

input `int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-1/2/d*(b*sec(d*x+c))^(1/2)*b*sec(d*x+c)^(5/2)*(cos(d*x+c)^3*ln(-cot(d*x+c)+csc(d*x+c)-1)-cos(d*x+c)^3*ln(csc(d*x+c)-cot(d*x+c)+1)-cos(d*x+c)*sin(d*x+c))`

**3.144.5 Fricas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 202, normalized size of antiderivative = 2.73

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{\frac{3}{2}} dx = \left[ \frac{b^{\frac{3}{2}} \cos(dx + c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b}\sqrt{\frac{b}{\cos(dx+c)}}\sqrt{\cos(dx+c)}\sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2b\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4d \cos(dx + c)}, \right. \\ \left. - \frac{\sqrt{-bb} \arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}}\sqrt{\cos(dx+c)}\sin(dx+c)}{b}\right) \cos(dx + c) - \frac{b\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{\sqrt{\cos(dx+c)}}}{2d \cos(dx + c)} \right]$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`output `[1/4*(b^(3/2)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*b*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)), -1/2*(sqrt(-b)*b*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - b*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))]`**3.144.6 Sympy [F]**

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{\frac{3}{2}} dx = \int (b \sec(c + dx))^{\frac{3}{2}} \sec^{\frac{3}{2}}(c + dx) dx$$

input `integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**(3/2),x)`output `Integral((b*sec(c + d*x))**(3/2)*sec(c + d*x)**(3/2), x)`

**3.144.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 691 vs.  $2(62) = 124$ .

Time = 0.44 (sec) , antiderivative size = 691, normalized size of antiderivative = 9.34

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \text{Too large to display}$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output

```
-1/4*(4*(b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (b*cos(4*d*x + 4*c)^2 + 4*b*cos(2*d*x + 2*c)^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + (b*cos(4*d*x + 4*c)^2 + 4*b*cos(2*d*x + 2*c)^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 4*(b*cos(4*d*x + 4*c) + 2*b*cos(2*d*x + 2*c) + b)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(b*cos(4*d*x + 4*c) + 2*b*cos(2*d*x + 2*c) + b)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)
```

**3.144.8 Giac [F]**

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(3/2), x)`

**3.144.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{3/2} \left( \frac{1}{\cos(c + dx)} \right)^{3/2} dx$$

input `int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(3/2),x)`output `int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(3/2), x)`



### 3.145 $\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{3/2} dx$

3.145.1 Optimal result . . . . .	892
3.145.2 Mathematica [A] (verified) . . . . .	892
3.145.3 Rubi [A] (verified) . . . . .	893
3.145.4 Maple [A] (verified) . . . . .	894
3.145.5 Fracas [A] (verification not implemented) . . . . .	894
3.145.6 Sympy [F] . . . . .	895
3.145.7 Maxima [A] (verification not implemented) . . . . .	895
3.145.8 Giac [F] . . . . .	895
3.145.9 Mupad [B] (verification not implemented) . . . . .	896

#### 3.145.1 Optimal result

Integrand size = 23, antiderivative size = 33

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{3/2} dx = \frac{b\sqrt{\sec(c + dx)}\sqrt{b \sec(c + dx)} \sin(c + dx)}{d}$$

output `b*sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.145.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{3/2} dx = \frac{(b \sec(c + dx))^{3/2} \sin(c + dx)}{d\sqrt{\sec(c + dx)}}$$

input `Integrate[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2),x]`

output `((b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])`

**3.145.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \sec^2(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & -\frac{b\sqrt{b \sec(c+dx)} \int 1d(-\tan(c+dx))}{d\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{b \sin(c+dx) \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}{d}
 \end{aligned}$$

input `Int[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2),x]`

output `(b*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d`

**3.145.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

---

3.145.  $\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2} dx$

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

### 3.145.4 Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{\sec(dx+c)^{\frac{3}{2}} b \sqrt{b \sec(dx+c)} \cos(dx+c) \sin(dx+c)}{d}$	36
risch	$\frac{2ib \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} e^{-i(dx+c)} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}}}{d}$	68

```
input int(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/d*sec(d*x+c)^(3/2)*b*(b*sec(d*x+c))^(1/2)*cos(d*x+c)*sin(d*x+c)
```

### 3.145.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \sqrt{\sec(c+dx)} (b \sec(c+dx))^{3/2} dx = \frac{b \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{d \sqrt{\cos(dx+c)}}$$

```
input integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2),x, algorithm="fracas")
```

```
output b*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))
```

**3.145.6 Sympy [F]**

$$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2} dx = \int (b \sec(c+dx))^{\frac{3}{2}} \sqrt{\sec(c+dx)} dx$$

input `integrate(sec(d*x+c)**(1/2)*(b*sec(d*x+c))**(3/2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*sqrt(sec(c + d*x)), x)`

**3.145.7 Maxima [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.64

$$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2} dx = \frac{2 b^{\frac{3}{2}} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c)^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1)d}$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `2*b^(3/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)`

**3.145.8 Giac [F]**

$$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2} dx = \int (b \sec(dx+c))^{\frac{3}{2}} \sqrt{\sec(dx+c)} dx$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)*sqrt(sec(d*x + c)), x)`

**3.145.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.42

$$\int \sqrt{\sec(c + dx)} (b \sec(c + dx))^{3/2} dx = \frac{b(\cos(dx) - \sin(dx) \operatorname{li}(\sin(c) + \cos(c) \operatorname{li}(\frac{b}{\cos(c+dx)})) \sqrt{\frac{1}{\cos(c+dx)}})}{d}$$

input `int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(1/2),x)`output `(b*(cos(d*x) - sin(d*x)*1i)*(cos(c)*1i + sin(c))*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2))/d`

**3.146**  $\int \frac{(b \sec(c+dx))^{3/2}}{\sqrt{\sec(c+dx)}} dx$

3.146.1 Optimal result . . . . . 897  
 3.146.2 Mathematica [A] (verified) . . . . . 897  
 3.146.3 Rubi [A] (verified) . . . . . 898  
 3.146.4 Maple [A] (verified) . . . . . 899  
 3.146.5 Fricas [A] (verification not implemented) . . . . . 899  
 3.146.6 Sympy [F] . . . . . 900  
 3.146.7 Maxima [B] (verification not implemented) . . . . . 900  
 3.146.8 Giac [F] . . . . . 900  
 3.146.9 Mupad [F(-1)] . . . . . 901

**3.146.1 Optimal result**

Integrand size = 23, antiderivative size = 34

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx = \frac{b \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}}$$

output `b*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

**3.146.2 Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.97

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx = \frac{\operatorname{arctanh}(\sin(c + dx))(b \sec(c + dx))^{3/2}}{d \sec^{3/2}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)/Sqrt[Sec[c + d*x]],x]`

output `(ArcTanh[Sin[c + d*x]]*(b*Sec[c + d*x])^(3/2))/(d*Sec[c + d*x]^(3/2))`

### 3.146.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{2031}$$

$$\frac{b\sqrt{b \sec(c + dx)} \int \sec(c + dx) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{b\sqrt{b \sec(c + dx)} \int \csc(c + dx + \frac{\pi}{2}) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{4257}$$

$$\frac{\text{arctanh}(\sin(c + dx))\sqrt{b \sec(c + dx)}}{d\sqrt{\sec(c + dx)}}$$

input `Int[(b*Sec[c + d*x])^(3/2)/Sqrt[Sec[c + d*x]],x]`

output `(b*ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]])/(d*Sqrt[Sec[c + d*x]])`

#### 3.146.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.146.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.38

method	result	size
default	$\frac{-2\sqrt{\sec(dx+c)} \cos(dx+c) \operatorname{arctanh}(\cot(dx+c) - \csc(dx+c))\sqrt{b\sec(dx+c)} b}{d}$	47
risch	$-\frac{b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)}-i)}}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} d}} + \frac{b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)}+i)}}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} d}}$	141

input `int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/d*sec(d*x+c)^(1/2)*cos(d*x+c)*arctanh(cot(d*x+c)-csc(d*x+c))*(b*sec(d*x+c))^(1/2)*b`

### 3.146.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 112, normalized size of antiderivative = 3.29

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx = \left[ \frac{b^{3/2} \log \left( -\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right)}{2d}, \right. \\ \left. -\frac{\sqrt{-b} b \arctan \left( \frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b} \right)}{d} \right]$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `[1/2*b^(3/2)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/d, -sqrt(-b)*b*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/d]`

---

3.146.  $\int \frac{(b \sec(c+dx))^{3/2}}{\sqrt{\sec(c+dx)}} dx$



**3.146.6 Sympy [F]**

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx = \int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx$$

input `integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(1/2),x)`

output `Integral((b*sec(c + d*x))**(3/2)/sqrt(sec(c + d*x)), x)`

**3.146.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. 2(30) = 60.

Time = 0.46 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.00

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx = \frac{(b \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - b \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1)) \sqrt{b}}{2d}$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/2*(b*log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - b*log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))*sqrt(b)/d`

**3.146.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx = \int \frac{(b \sec(dx + c))^{3/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)/sqrt(sec(d*x + c)), x)`

**3.146.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sqrt{\sec(c + dx)}} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\sqrt{\frac{1}{\cos(c+dx)}}} dx$$

input `int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(1/2),x)`output `int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(1/2), x)`

**3.147**  $\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{3}{2}}(c+dx)} dx$

3.147.1 Optimal result . . . . . 902  
 3.147.2 Mathematica [A] (verified) . . . . . 902  
 3.147.3 Rubi [A] (verified) . . . . . 903  
 3.147.4 Maple [A] (verified) . . . . . 904  
 3.147.5 Fricas [A] (verification not implemented) . . . . . 904  
 3.147.6 Sympy [A] (verification not implemented) . . . . . 904  
 3.147.7 Maxima [A] (verification not implemented) . . . . . 905  
 3.147.8 Giac [F] . . . . . 905  
 3.147.9 Mupad [B] (verification not implemented) . . . . . 905

**3.147.1 Optimal result**

Integrand size = 23, antiderivative size = 25

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{bx \sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}}$$

output `b*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

**3.147.2 Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{bx \sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(3/2),x]`

output `(b*x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]`

**3.147.3 Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2031, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{3/2}(c + dx)} dx$$

↓ 2031

$$\frac{b \sqrt{b \sec(c + dx)} \int 1 dx}{\sqrt{\sec(c + dx)}}$$

↓ 24

$$\frac{bx \sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}}$$

input `Int[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(3/2),x]`

output `(b*x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]`

**3.147.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fv_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

**3.147.4 Maple [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

method	result	size
default	$\frac{(dx+c)b\sqrt{b\sec(dx+c)}}{d\sqrt{\sec(dx+c)}}$	29
risch	$\frac{b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}x}}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}$	55

input `int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`output `1/d*(d*x+c)*b/sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)`**3.147.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 99, normalized size of antiderivative = 3.96

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \left[ \frac{\sqrt{-bb} \log \left( -2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b \right)}{2d} \right]$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algorithm="fracas")`output `[1/2*sqrt(-b)*b*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/d, b^(3/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/d]`**3.147.6 Sympy [A] (verification not implemented)**

Time = 3.46 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{x(b \sec(c + dx))^{\frac{3}{2}}}{\sec^{\frac{3}{2}}(c + dx)}$$

input `integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(3/2),x)`

output `x*(b*sec(c + d*x))**(3/2)/sec(c + d*x)**(3/2)`

### 3.147.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{2 b^{\frac{3}{2}} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `2*b^(3/2)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d`

### 3.147.8 Giac [F]

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{\frac{3}{2}}}{\sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(3/2), x)`

### 3.147.9 Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{b x \sqrt{\frac{b}{\cos(c+dx)}}}{\sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(3/2),x)`

output `(b*x*(b/cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2)`

---

3.147.  $\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{3}{2}}(c+dx)} dx$

**3.148**      
$$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{5/2}(c+dx)} dx$$

3.148.1 Optimal result . . . . .	906
3.148.2 Mathematica [A] (verified) . . . . .	906
3.148.3 Rubi [A] (verified) . . . . .	907
3.148.4 Maple [A] (verified) . . . . .	908
3.148.5 Fracas [A] (verification not implemented) . . . . .	908
3.148.6 Sympy [A] (verification not implemented) . . . . .	909
3.148.7 Maxima [A] (verification not implemented) . . . . .	909
3.148.8 Giac [A] (verification not implemented) . . . . .	909
3.148.9 Mupad [B] (verification not implemented) . . . . .	910

**3.148.1 Optimal result**

Integrand size = 23, antiderivative size = 33

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx = \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{d\sqrt{\sec(c + dx)}}$$

output `b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

**3.148.2 Mathematica [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx = \frac{(b \sec(c + dx))^{3/2} \sin(c + dx)}{d \sec^{3/2}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(5/2),x]`

output `((b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sec[c + d*x]^(3/2))`

### 3.148.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx$$

$$\downarrow \text{2031}$$

$$\frac{b\sqrt{b \sec(c + dx)} \int \cos(c + dx) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{b\sqrt{b \sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2}) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{3117}$$

$$\frac{b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}}$$

input `Int[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(5/2),x]`

output `(b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])`

#### 3.148.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



```
rule 3117 Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

### 3.148.4 Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{b\sqrt{b\sec(dx+c)}\tan(dx+c)}{d\sec(dx+c)^{\frac{3}{2}}}$	30
risch	$-\frac{ib\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}e^{i(dx+c)}}}{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}d} + \frac{ib\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}e^{-i(dx+c)}}}{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}d}$	136

```
input int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/d*b/sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2)*tan(d*x+c)
```

### 3.148.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \frac{(b\sec(c+dx))^{3/2}}{\sec^{\frac{5}{2}}(c+dx)} dx = \frac{b\sqrt{\frac{b}{\cos(dx+c)}}\sqrt{\cos(dx+c)}\sin(dx+c)}{d}$$

```
input integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2),x, algorithm="fracas")
```

```
output b*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/d
```

**3.148.6 Sympy [A] (verification not implemented)**

Time = 18.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx = \begin{cases} \frac{(b \sec(c + dx))^{3/2} \tan(c + dx)}{d \sec^{5/2}(c + dx)} & \text{for } d \neq 0 \\ \frac{x(b \sec(c))^{3/2}}{\sec^{5/2}(c)} & \text{otherwise} \end{cases}$$

input `integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(5/2),x)`output `Piecewise(((b*sec(c + d*x))**(3/2)*tan(c + d*x)/(d*sec(c + d*x)**(5/2)), Ne(d, 0)), (x*(b*sec(c))**(3/2)/sec(c)**(5/2), True))`**3.148.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.39

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx = \frac{b^{3/2} \sin(dx + c)}{d}$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2),x, algorithm="maxima")`output `b^(3/2)*sin(d*x + c)/d`**3.148.8 Giac [A] (verification not implemented)**

Time = 0.87 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.15

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx = \frac{2 b^{3/2} \operatorname{sgn}(\cos(dx + c)) \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{d \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + d}$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2),x, algorithm="giac")`output `2*b^(3/2)*sgn(cos(d*x + c))*tan(1/2*d*x + 1/2*c)/(d*tan(1/2*d*x + 1/2*c)^2 + d)`

---

3.148.  $\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx$

**3.148.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{5/2}(c + dx)} dx = \frac{b \sin(c + dx) \sqrt{\frac{b}{\cos(c+dx)}}}{d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(5/2),x)`output `(b*sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(d*(1/cos(c + d*x))^(1/2))`

**3.149** 
$$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{7/2}(c+dx)} dx$$

3.149.1 Optimal result . . . . . 911  
 3.149.2 Mathematica [A] (verified) . . . . . 911  
 3.149.3 Rubi [A] (verified) . . . . . 912  
 3.149.4 Maple [A] (verified) . . . . . 913  
 3.149.5 Fricas [A] (verification not implemented) . . . . . 914  
 3.149.6 Sympy [F(-1)] . . . . . 914  
 3.149.7 Maxima [A] (verification not implemented) . . . . . 914  
 3.149.8 Giac [F] . . . . . 915  
 3.149.9 Mupad [B] (verification not implemented) . . . . . 915

**3.149.1 Optimal result**

Integrand size = 23, antiderivative size = 65

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx = \frac{bx \sqrt{b \sec(c + dx)}}{2 \sqrt{\sec(c + dx)}} + \frac{b \sqrt{b \sec(c + dx)} \sin(c + dx)}{2d \sec^{3/2}(c + dx)}$$

output `1/2*b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+1/2*b*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

**3.149.2 Mathematica [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.69

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx = \frac{(b \sec(c + dx))^{3/2}(2(c + dx) + \sin(2(c + dx)))}{4d \sec^{3/2}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(7/2),x]`

output `((b*Sec[c + d*x])^(3/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sec[c + d*x]^(3/2))`

**3.149.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.75, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b\sqrt{b \sec(c + dx)} \int \cos^2(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2})^2 dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{b\sqrt{b \sec(c + dx)} \left( \int \frac{1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{b\sqrt{b \sec(c + dx)} \left( \frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right)}{\sqrt{\sec(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(7/2),x]`

output `(b*Sqrt[b*Sec[c + d*x]]*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/Sqrt[Sec[c + d*x]]`

3.149.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.149.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.71

method	result	size
default	$\frac{\sqrt{b \sec(dx+c)} b (\tan(dx+c) + (dx+c) \sec(dx+c)^2)}{2d \sec(dx+c)^{\frac{5}{2}}}$	46
risch	$\frac{b \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} x}{2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{ib \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{2i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}} + \frac{ib \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-2i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}}$	191

input `int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2),x,method=_RETURNVERBOSE)`

output `1/2/d*(b*sec(d*x+c))^(1/2)*b/sec(d*x+c)^(5/2)*(tan(d*x+c)+(d*x+c)*sec(d*x+c)^2)`

**3.149.5 Fracas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 161, normalized size of antiderivative = 2.48

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx = \left[ \frac{2b \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{3/2} \sin(dx+c) + \sqrt{-bb} \log\left(-2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)\right)}{4d} \right]$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2),x, algorithm="fracas")`output `[1/4*(2*b*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(-b)*b*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/2*(b*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + b^(3/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/d]`**3.149.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(7/2),x)`output `Timed out`**3.149.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.43

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx = \frac{(2(dx+c)b + b \sin(2dx + 2c))\sqrt{b}}{4d}$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2),x, algorithm="maxima")`output `1/4*(2*(d*x + c)*b + b*sin(2*d*x + 2*c))*sqrt(b)/d`

---

3.149.  $\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{7/2}(c+dx)} dx$

**3.149.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{3/2}}{\sec^{7/2}(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(7/2), x)`

**3.149.9 Mupad [B] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{7/2}(c + dx)} dx = \frac{b (\sin(2c + 2dx) + 2dx) \sqrt{\frac{b}{\cos(c+dx)}}}{4d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(7/2),x)`

output `(b*(sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*d*(1/cos(c + d*x))^(1/2))`



$$3.150 \quad \int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{9}{2}}(c+dx)} dx$$

3.150.1 Optimal result . . . . .	916
3.150.2 Mathematica [A] (verified) . . . . .	916
3.150.3 Rubi [A] (verified) . . . . .	917
3.150.4 Maple [A] (verified) . . . . .	918
3.150.5 Fricas [A] (verification not implemented) . . . . .	919
3.150.6 Sympy [F(-1)] . . . . .	919
3.150.7 Maxima [A] (verification not implemented) . . . . .	919
3.150.8 Giac [F] . . . . .	920
3.150.9 Mupad [B] (verification not implemented) . . . . .	920

### 3.150.1 Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{9}{2}}(c+dx)} dx = \frac{b\sqrt{b \sec(c+dx)} \sin(c+dx)}{d\sqrt{\sec(c+dx)}} - \frac{b\sqrt{b \sec(c+dx)} \sin^3(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

output `b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)-1/3*b*sin(d*x+c)^3*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

### 3.150.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{9}{2}}(c+dx)} dx = \frac{(5 + \cos(2(c+dx)))(b \sec(c+dx))^{3/2} \sin(c+dx)}{6d \sec^{\frac{3}{2}}(c+dx)}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(9/2),x]`

output `((5 + Cos[2*(c + d*x)])*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(6*d*Sec[c + d*x]^(3/2))`

---

3.150.  $\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{9}{2}}(c+dx)} dx$

**3.150.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{9}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b\sqrt{b \sec(c + dx)} \int \cos^3(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2})^3 dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3113} \\
 & -\frac{b\sqrt{b \sec(c + dx)} \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{b(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx)) \sqrt{b \sec(c + dx)}}{d\sqrt{\sec(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(9/2),x]`

output `-((b*Sqrt[b*Sec[c + d*x]]*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/(d*Sqrt[Sec[c + d*x]]))`

3.150.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

3.150.4 Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{b\sqrt{b\sec(dx+c)} (\tan(dx+c)+2\tan(dx+c)\sec(dx+c)^2)}{3d\sec(dx+c)^{\frac{7}{2}}}$	48
risch	$-\frac{3ib\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}d + \frac{3ib\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}d + \frac{b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}{12\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}d$	202

input `int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2),x,method=_RETURNVERBOSE)`

output `1/3/d*b*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2)*(tan(d*x+c)+2*tan(d*x+c)*sec(d*x+c)^2)`

3.150. 
$$\int \frac{(b\sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx$$

**3.150.5 Fricas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \frac{(b \cos(dx + c)^3 + 2b \cos(dx + c)) \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c)}{3d \sqrt{\cos(dx + c)}}$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2),x, algorithm="fricas")`output `1/3*(b*cos(d*x + c)^3 + 2*b*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))`**3.150.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(9/2),x)`output `Timed out`**3.150.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \frac{(b \sin(3dx + 3c) + 9b \sin(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c)))) \sqrt{b}}{12d}$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2),x, algorithm="maxima")`output `1/12*(b*sin(3*d*x + 3*c) + 9*b*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*sqrt(b)/d`

**3.150.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{\frac{3}{2}}}{\sec(dx + c)^{\frac{9}{2}}} dx$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(9/2), x)`

**3.150.9 Mupad [B] (verification not implemented)**

Time = 13.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.64

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \frac{b(9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(9/2),x)`

output `(b*(9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*d*(1/cos(c + d*x))^(1/2))`

**3.151** 
$$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{11}{2}}(c+dx)} dx$$

3.151.1 Optimal result . . . . . 921  
 3.151.2 Mathematica [A] (verified) . . . . . 921  
 3.151.3 Rubi [A] (verified) . . . . . 922  
 3.151.4 Maple [A] (verified) . . . . . 923  
 3.151.5 Fricas [A] (verification not implemented) . . . . . 924  
 3.151.6 Sympy [F(-1)] . . . . . 924  
 3.151.7 Maxima [A] (verification not implemented) . . . . . 924  
 3.151.8 Giac [F] . . . . . 925  
 3.151.9 Mupad [B] (verification not implemented) . . . . . 925

**3.151.1 Optimal result**

Integrand size = 23, antiderivative size = 101

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{11}{2}}(c + dx)} dx = \frac{3bx\sqrt{b \sec(c + dx)}}{8\sqrt{\sec(c + dx)}} + \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{4d \sec^{\frac{7}{2}}(c + dx)} + \frac{3b\sqrt{b \sec(c + dx)} \sin(c + dx)}{8d \sec^{\frac{3}{2}}(c + dx)}$$

output `1/4*b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(7/2)+3/8*b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+3/8*b*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

**3.151.2 Mathematica [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.54

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{\frac{11}{2}}(c + dx)} dx = \frac{(b \sec(c + dx))^{3/2}(12(c + dx) + 8 \sin(2(c + dx)) + \sin(4(c + dx)))}{32d \sec^{\frac{3}{2}}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(11/2),x]`

output `((b*Sec[c + d*x])^(3/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)]))/(32*d*Sec[c + d*x]^(3/2))`

---

3.151. 
$$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{\frac{11}{2}}(c+dx)} dx$$

**3.151.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.74, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {2031, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \sec(c + dx))^{3/2}}{\sec^{11/2}(c + dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b\sqrt{b \sec(c + dx)} \int \cos^4(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2})^4 dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{b\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \int \cos^2(c + dx) dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \int \sin(c + dx + \frac{\pi}{2})^2 dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{b\sqrt{b \sec(c + dx)} \left( \frac{3}{4} \left( \int \frac{1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{b\sqrt{b \sec(c + dx)} \left( \frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left( \frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right)}{\sqrt{\sec(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(11/2),x]`

---

3.151.  $\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{11/2}(c+dx)} dx$

output  $(b\sqrt{b\sec[c + dx]})*((\cos[c + dx]^3\sin[c + dx])/(4d) + (3*(x/2 + (\cos[c + dx]*\sin[c + dx]))/(2*d)))/4)/\sqrt{\sec[c + dx]}$

### 3.151.3.1 Defintions of rubi rules used

rule 24  $\text{Int}[a_, x\_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 2031  $\text{Int}[(F x_.)*((a_.)*(v_.))^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]) \text{Int}[v^{(m + n)}*F x, x], x] /; \text{FreeQ}\{a, b, m\}, x\} \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n + 1/2, 0] \&\& \text{IntegerQ}[m + n]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3115  $\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_)]^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c + dx]*((b*\sin[c + dx])^{(n - 1)}/(d*n)), x] + \text{Simp}[b^{2*(n - 1)}/n \text{Int}[(b*\sin[c + dx])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x\} \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

### 3.151.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.64

method	result	size
default	$\frac{b\sqrt{b\sec(dx+c)}(2\tan(dx+c)+3\tan(dx+c)\sec(dx+c)^2+3(dx+c)\sec(dx+c)^4)}{8d\sec(dx+c)^{\frac{9}{2}}}$	65
risch	$\frac{3b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}x}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{ib\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}e^{2i(dx+c)}}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}d}} + \frac{ib\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}e^{-2i(dx+c)}}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}d}} + \frac{b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}\sin(4dx+4c)}}{32\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}d}}$	257

input  $\text{int}((b*\sec(dx+c))^{(3/2)}/\sec(dx+c)^{(11/2)},x,\text{method}=\_RETURNVERBOSE)$

output  $1/8*b/d*(b*\sec(dx+c))^{(1/2)}/\sec(dx+c)^{(9/2)}*(2*\tan(dx+c)+3*\tan(dx+c)*\sec(dx+c)^2+3*(dx+c)*\sec(dx+c)^4)$

3.151.  $\int \frac{(b\sec(c+dx))^{3/2}}{\sec^{\frac{11}{2}}(c+dx)} dx$



**3.151.5 Fracas [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.05

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{11/2}(c + dx)} dx = \frac{3 \sqrt{-bb} \log \left( -2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{3/2} \sin(dx+c) + 2b \cos(dx+c)^2 - b \right)}{16d}$$

```
input integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(11/2),x, algorithm="fricas")
```

```
output [1/16*(3*sqrt(-b)*b*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)
)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b) + 2*(2*b*cos(d*x + c)^4 + 3*b*cos
(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d, 1/8*
(3*b^(3/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x
+ c)))) + (2*b*cos(d*x + c)^4 + 3*b*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*s
in(d*x + c)/sqrt(cos(d*x + c)))/d]
```

**3.151.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{11/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(11/2),x)
```

```
output Timed out
```

**3.151.7 Maxima [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.52

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{11/2}(c + dx)} dx = \frac{(12(dx+c)b + b \sin(4dx+4c) + 8b \sin(\frac{1}{2} \arctan(\sin(4dx+4c)), \cos(4dx+4c))}{32d}$$

```
input integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(11/2),x, algorithm="maxima")
```

---

3.151.  $\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{11/2}(c+dx)} dx$

output  $1/32*(12*(d*x + c)*b + b*\sin(4*d*x + 4*c) + 8*b*\sin(1/2*\arctan2(\sin(4*d*x + 4*c), \cos(4*d*x + 4*c))))*\sqrt{b}/d$

### 3.151.8 Giac [F]

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{11/2}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{3/2}}{\sec(dx + c)^{11/2}} dx$$

input `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(11/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(11/2), x)`

### 3.151.9 Mupad [B] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.52

$$\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{11/2}(c + dx)} dx = \frac{b \sqrt{\frac{b}{\cos(c+dx)}} (8 \sin(2c + 2dx) + \sin(4c + 4dx) + 12dx)}{32d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(11/2),x)`

output `(b*(b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))/(32*d*(1/cos(c + d*x))^(1/2))`

### 3.152 $\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx$

3.152.1 Optimal result . . . . .	926
3.152.2 Mathematica [A] (verified) . . . . .	926
3.152.3 Rubi [A] (verified) . . . . .	927
3.152.4 Maple [A] (verified) . . . . .	928
3.152.5 Fricas [A] (verification not implemented) . . . . .	929
3.152.6 Sympy [F(-1)] . . . . .	929
3.152.7 Maxima [B] (verification not implemented) . . . . .	929
3.152.8 Giac [F] . . . . .	930
3.152.9 Mupad [B] (verification not implemented) . . . . .	930

#### 3.152.1 Optimal result

Integrand size = 23, antiderivative size = 116

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{b^2 \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} \sin(c + dx)}{d} + \frac{2b^2 \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin^3(c + dx)}{3d} + \frac{b^2 \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin^5(c + dx)}{5d}$$

output `2/3*b^2*sec(d*x+c)^(5/2)*sin(d*x+c)^3*(b*sec(d*x+c))^(1/2)/d+1/5*b^2*sec(d*x+c)^(9/2)*sin(d*x+c)^5*(b*sec(d*x+c))^(1/2)/d+b^2*sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.152.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.49

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{(b \sec(c + dx))^{5/2} (\tan(c + dx) + \frac{2}{3} \tan^3(c + dx) + \frac{1}{5} \tan^5(c + dx))}{d \sec^{\frac{5}{2}}(c + dx)}$$

input `Integrate[Sec[c + d*x]^(7/2)*(b*Sec[c + d*x])^(5/2),x]`

output  $((b*\text{Sec}[c + d*x])^{5/2}*(\text{Tan}[c + d*x] + (2*\text{Tan}[c + d*x]^3)/3 + \text{Tan}[c + d*x]^5/5))/(d*\text{Sec}[c + d*x]^{5/2})$

### 3.152.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.54, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx \\ & \quad \downarrow \text{2031} \\ & \frac{b^2 \sqrt{b \sec(c + dx)} \int \sec^6(c + dx) dx}{\sqrt{\sec(c + dx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{b^2 \sqrt{b \sec(c + dx)} \int \csc(c + dx + \frac{\pi}{2})^6 dx}{\sqrt{\sec(c + dx)}} \\ & \quad \downarrow \text{4254} \\ & \frac{b^2 \sqrt{b \sec(c + dx)} \int (\tan^4(c + dx) + 2 \tan^2(c + dx) + 1) d(-\tan(c + dx))}{d \sqrt{\sec(c + dx)}} \\ & \quad \downarrow \text{2009} \\ & \frac{b^2 (-\frac{1}{5} \tan^5(c + dx) - \frac{2}{3} \tan^3(c + dx) - \tan(c + dx)) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}} \end{aligned}$$

input  $\text{Int}[\text{Sec}[c + d*x]^{7/2}*(b*\text{Sec}[c + d*x])^{5/2}, x]$

output  $-((b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*(-\text{Tan}[c + d*x] - (2*\text{Tan}[c + d*x]^3)/3 - \text{Tan}[c + d*x]^5/5))/(d*\text{Sqrt}[\text{Sec}[c + d*x]]))$

## 3.152.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

## 3.152.4 Maple [A] (verified)

Time = 24.60 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.53

method	result	size
default	$\frac{b^2 \sec(dx+c)^{\frac{11}{2}} \sqrt{b \sec(dx+c)} (8 \cos(dx+c)^4 + 4 \cos(dx+c)^2 + 3) \sin(dx+c) \cos(dx+c)}{15d}$	61
risch	$\frac{16ib^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (10 e^{3i(dx+c)} + 6 \cos(dx+c) + 4i \sin(dx+c))}{15(e^{2i(dx+c)}+1)^4 d}$	103

input `int(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `1/15/d*b^2*sec(d*x+c)^(11/2)*(b*sec(d*x+c))^(1/2)*(8*cos(d*x+c)^4+4*cos(d*x+c)^2+3)*sin(d*x+c)*cos(d*x+c)`

**3.152.5 Fricas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.54

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{\frac{5}{2}} dx = \frac{(8b^2 \cos(dx + c)^4 + 4b^2 \cos(dx + c)^2 + 3b^2) \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c)}{15d \cos(dx + c)^{\frac{9}{2}}}$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/15*(8*b^2*cos(d*x + c)^4 + 4*b^2*cos(d*x + c)^2 + 3*b^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(9/2))`

**3.152.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{\frac{5}{2}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(7/2)*(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

**3.152.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 705 vs. 2(100) = 200.

Time = 0.47 (sec) , antiderivative size = 705, normalized size of antiderivative = 6.08

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{\frac{5}{2}} dx = \frac{15(2(5 \cos(8dx + 8c) + 10 \cos(6dx + 6c) + 10 \cos(4dx + 4c) + 5 \cos(2dx + 2c) + 1) \cos(10dx + 10c) + \dots}{\dots}$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

3.152.  $\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{\frac{5}{2}} dx$

output

```
-16/15*(5*(2*b^2*sin(4*d*x + 4*c) + b^2*sin(2*d*x + 2*c))*cos(10*d*x + 10*c) + 25*(2*b^2*sin(4*d*x + 4*c) + b^2*sin(2*d*x + 2*c))*cos(8*d*x + 8*c) + 50*(2*b^2*sin(4*d*x + 4*c) + b^2*sin(2*d*x + 2*c))*cos(6*d*x + 6*c) - (10*b^2*cos(4*d*x + 4*c) + 5*b^2*cos(2*d*x + 2*c) + b^2)*sin(10*d*x + 10*c) - 5*(10*b^2*cos(4*d*x + 4*c) + 5*b^2*cos(2*d*x + 2*c) + b^2)*sin(8*d*x + 8*c) - 10*(10*b^2*cos(4*d*x + 4*c) + 5*b^2*cos(2*d*x + 2*c) + b^2)*sin(6*d*x + 6*c))*sqrt(b)/((2*(5*cos(8*d*x + 8*c) + 10*cos(6*d*x + 6*c) + 10*cos(4*d*x + 4*c) + 5*cos(2*d*x + 2*c) + 1)*cos(10*d*x + 10*c) + cos(10*d*x + 10*c)^2 + 10*(10*cos(6*d*x + 6*c) + 10*cos(4*d*x + 4*c) + 5*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + 25*cos(8*d*x + 8*c)^2 + 20*(10*cos(4*d*x + 4*c) + 5*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 100*cos(6*d*x + 6*c)^2 + 20*(5*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 100*cos(4*d*x + 4*c)^2 + 25*cos(2*d*x + 2*c)^2 + 10*(sin(8*d*x + 8*c) + 2*sin(6*d*x + 6*c) + 2*sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(10*d*x + 10*c) + sin(10*d*x + 10*c)^2 + 50*(2*sin(6*d*x + 6*c) + 2*sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 25*sin(8*d*x + 8*c)^2 + 100*(2*sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 100*sin(6*d*x + 6*c)^2 + 100*sin(4*d*x + 4*c)^2 + 100*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 25*sin(2*d*x + 2*c)^2 + 10*cos(2*d*x + 2*c) + 1)*d)
```

### 3.152.8 Giac [F]

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c)^{\frac{7}{2}} dx$$

input `integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^(7/2), x)`

### 3.152.9 Mupad [B] (verification not implemented)

Time = 18.01 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.77

$$\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx =$$

$$\sqrt{-\frac{b}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}} \left( \frac{b^2 \sqrt{-\frac{1}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}} 8i}{15d} + \frac{b^2 \sqrt{-\frac{1}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}} (-2 \sin(2c + 2dx)^2 + \sin(4c + 4dx) \operatorname{li}(1)) 16i}{3d} + \frac{b^2 \sqrt{-\frac{1}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}}}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1} \right)$$

$$16 (\sin(c + dx))^2 - 1$$

3.152.  $\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx$

input `int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(7/2),x)`

output `-((-b/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*((b^2*(-1/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*8i)/(15*d) + (b^2*(-1/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*(sin(4*c + 4*d*x)*1i - 2*sin(2*c + 2*d*x)^2 + 1)*16i)/(3*d) + (b^2*(-1/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*(sin(2*c + 2*d*x)*1i - 2*sin(c + d*x)^2 + 1)*8i)/(3*d))*(sin(5*c + 5*d*x)*1i + 2*sin((5*c)/2 + (5*d*x)/2)^2 - 1))/(16*(sin(c + d*x)^2 - 1)^2)`



### 3.153 $\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx$

3.153.1 Optimal result . . . . .	932
3.153.2 Mathematica [A] (verified) . . . . .	932
3.153.3 Rubi [A] (verified) . . . . .	933
3.153.4 Maple [A] (verified) . . . . .	934
3.153.5 Fricas [A] (verification not implemented) . . . . .	935
3.153.6 Sympy [F(-1)] . . . . .	935
3.153.7 Maxima [B] (verification not implemented) . . . . .	935
3.153.8 Giac [F] . . . . .	936
3.153.9 Mupad [B] (verification not implemented) . . . . .	936

#### 3.153.1 Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{b^2 \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} \sin(c + dx)}{d} + \frac{b^2 \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} \sin^3(c + dx)}{3d}$$

output `1/3*b^2*sec(d*x+c)^(5/2)*sin(d*x+c)^3*(b*sec(d*x+c))^(1/2)/d+b^2*sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

#### 3.153.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{(b \sec(c + dx))^{5/2} (\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d \sec^{\frac{5}{2}}(c + dx)}$$

input `Integrate[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(5/2),x]`

output `((b*Sec[c + d*x])^(5/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sec[c + d*x]^(5/2))`

**3.153.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.67, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \int \sec^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2})^4 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \int (\tan^2(c+dx) + 1) d(-\tan(c+dx))}{d \sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{b^2 (-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx)) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(5/2),x]`

output `-((b^2*Sqrt[b*Sec[c + d*x]]*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/(d*Sqrt[Sec[c + d*x]]))`

## 3.153.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

## 3.153.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{b^2 \sec(dx+c)^{\frac{7}{2}} \sqrt{b \sec(dx+c)} (2 \cos(dx+c)^2 + 1) \cos(dx+c) \sin(dx+c)}{3d}$	51
risch	$\frac{4ib^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (4 \cos(dx+c) + 2i \sin(dx+c))}{3(e^{2i(dx+c)}+1)^2 d}$	92

input `int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `1/3/d*b^2*sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2)*(2*cos(d*x+c)^2+1)*cos(d*x+c)*sin(d*x+c)`

**3.153.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.63

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{(2b^2 \cos(dx + c)^2 + b^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx + c)}{3d \cos(dx + c)^{\frac{5}{2}}}$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2),x, algorithm="fracas")`

output `1/3*(2*b^2*cos(d*x + c)^2 + b^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(5/2))`

**3.153.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

**3.153.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 311 vs.  $2(66) = 132$ .

Time = 0.44 (sec) , antiderivative size = 311, normalized size of antiderivative = 4.09

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{4(3b^2 \cos(6dx + 6c) + 3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2) + 6(3 \cos(2dx + 2c) - 1) \cos(6dx + 6c)}{3d \cos(dx + c)^{\frac{5}{2}}}$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output 
$$-4/3*(3*b^2*\cos(6*d*x + 6*c)*\sin(2*d*x + 2*c) + 9*b^2*\cos(4*d*x + 4*c)*\sin(2*d*x + 2*c) - (3*b^2*\cos(2*d*x + 2*c) + b^2)*\sin(6*d*x + 6*c) - 3*(3*b^2*\cos(2*d*x + 2*c) + b^2)*\sin(4*d*x + 4*c))*\sqrt{b}/((2*(3*\cos(4*d*x + 4*c) + 3*\cos(2*d*x + 2*c) + 1)*\cos(6*d*x + 6*c) + \cos(6*d*x + 6*c)^2 + 6*(3*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + 9*\cos(4*d*x + 4*c)^2 + 9*\cos(2*d*x + 2*c)^2 + 6*(\sin(4*d*x + 4*c) + \sin(2*d*x + 2*c))*\sin(6*d*x + 6*c) + \sin(6*d*x + 6*c)^2 + 9*\sin(4*d*x + 4*c)^2 + 18*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 9*\sin(2*d*x + 2*c)^2 + 6*\cos(2*d*x + 2*c) + 1)*d)$$

### 3.153.8 Giac [F]

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^(3/2), x)`

### 3.153.9 Mupad [B] (verification not implemented)

Time = 1.30 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.70

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx = \frac{2b^2 \cos(c + dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} (4 \sin(c + dx) + 5 \sin(3c + 3dx) + \sin(5c + 5dx))}{3d(10 \cos(c + dx) + 5 \cos(3c + 3dx) + \cos(5c + 5dx))}$$

input `int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(3/2),x)`

output 
$$(2*b^2*\cos(c + d*x)*(b/\cos(c + d*x))^(1/2)*(1/\cos(c + d*x))^(1/2)*(\cos(c + d*x)*10i + 4*\sin(c + d*x) + \cos(3*c + 3*d*x)*5i + \cos(5*c + 5*d*x)*1i + 5*\sin(3*c + 3*d*x) + \sin(5*c + 5*d*x)))/(3*d*(10*\cos(c + d*x) + 5*\cos(3*c + 3*d*x) + \cos(5*c + 5*d*x)))$$

### 3.154 $\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{5/2} dx$

3.154.1 Optimal result . . . . .	937
3.154.2 Mathematica [A] (verified) . . . . .	937
3.154.3 Rubi [A] (verified) . . . . .	938
3.154.4 Maple [A] (verified) . . . . .	939
3.154.5 Fricas [A] (verification not implemented) . . . . .	940
3.154.6 Sympy [F(-1)] . . . . .	940
3.154.7 Maxima [B] (verification not implemented) . . . . .	941
3.154.8 Giac [F] . . . . .	941
3.154.9 Mupad [F(-1)] . . . . .	942

#### 3.154.1 Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{5/2} dx = \frac{b^2 \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{2d \sqrt{\sec(c + dx)}} + \frac{b^2 \sec^{3/2}(c + dx) \sqrt{b \sec(c + dx)} \sin(c + dx)}{2d}$$

output `1/2*b^2*sec(d*x+c)^(3/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+1/2*b^2*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

#### 3.154.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.64

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{5/2} dx = \frac{(b \sec(c + dx))^{5/2} (\operatorname{arctanh}(\sin(c + dx)) + \sec(c + dx) \tan(c + dx))}{2d \sec^{5/2}(c + dx)}$$

input `Integrate[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(5/2),x]`

output `((b*Sec[c + d*x])^(5/2)*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sec[c + d*x]^(5/2))`

**3.154.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.77, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {2031, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(c+dx)} (b \sec(c+dx))^{5/2} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2})^3 dx}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \left( \frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \left( \frac{1}{2} \int \csc(c+dx + \frac{\pi}{2}) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{b^2 \sqrt{b \sec(c+dx)} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(5/2),x]`

output `(b^2*Sqrt[b*Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/Sqrt[Sec[c + d*x]]`

## 3.154.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

## 3.154.4 Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.22

method	result
default	$-\frac{\sqrt{b \sec(dx+c)} b^2 \sec(dx+c)^{\frac{5}{2}} \left( \cos(dx+c)^3 \ln(-\cot(dx+c) + \csc(dx+c) - 1) - \cos(dx+c)^3 \ln(\csc(dx+c) - \cot(dx+c) + 1) - \cos(dx+c) \right)}{2d}$
risch	$-\frac{ib^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (e^{2i(dx+c)}-1)}{(e^{2i(dx+c)}+1)d} + \frac{b^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \ln(e^{i(dx+c)}+i) \cos(dx+c)}{d} - \frac{b^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}}}{e^{2i(dx+c)}}$

input `int(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `-1/2/d*(b*sec(d*x+c))^(1/2)*b^2*sec(d*x+c)^(5/2)*(cos(d*x+c)^3*ln(-cot(d*x+c)+csc(d*x+c)-1)-cos(d*x+c)^3*ln(csc(d*x+c)-cot(d*x+c)+1)-cos(d*x+c)*sin(d*x+c))`



**3.154.5 Fricas [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 208, normalized size of antiderivative = 2.67

$$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{5/2} dx = \left[ \frac{b^{5/2} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2b^2 \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4d \cos(dx+c)}, \right. \\ \left. - \frac{\sqrt{-b} b^2 \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right) \cos(dx+c) - \frac{b^2 \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{2d \cos(dx+c)} \right]$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`output `[1/4*(b^(5/2)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*b^2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)), -1/2*(sqrt(-b)*b^2*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - b^2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))]`**3.154.6 Sympy [F(-1)]**

Timed out.

$$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{5/2} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(1/2)*(b*sec(d*x+c))**(5/2),x)`output `Timed out`

**3.154.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 747 vs.  $2(66) = 132$ .

Time = 0.42 (sec) , antiderivative size = 747, normalized size of antiderivative = 9.58

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{5/2} dx = \text{Too large to display}$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output

```
-1/4*(4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 4*(b^2*cos(4*d*x + 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(b^2*cos(4*d*x + 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)
```

**3.154.8 Giac [F]**

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^{5/2} dx = \int (b \sec(dx + c))^{5/2} \sqrt{\sec(dx + c)} dx$$

input `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)*sqrt(sec(d*x + c)), x)`

**3.154.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{\sec(c + dx)} (b \sec(c + dx))^{5/2} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{5/2} \sqrt{\frac{1}{\cos(c + dx)}} dx$$

input `int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(1/2),x)`output `int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(1/2), x)`

$$3.155 \quad \int \frac{(b \sec(c+dx))^{5/2}}{\sqrt{\sec(c+dx)}} dx$$

3.155.1 Optimal result . . . . .	943
3.155.2 Mathematica [A] (verified) . . . . .	943
3.155.3 Rubi [A] (verified) . . . . .	944
3.155.4 Maple [A] (verified) . . . . .	945
3.155.5 Fricas [A] (verification not implemented) . . . . .	945
3.155.6 Sympy [F(-1)] . . . . .	946
3.155.7 Maxima [A] (verification not implemented) . . . . .	946
3.155.8 Giac [F] . . . . .	946
3.155.9 Mupad [B] (verification not implemented) . . . . .	947

### 3.155.1 Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{(b \sec(c+dx))^{5/2}}{\sqrt{\sec(c+dx)}} dx = \frac{b^2 \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} \sin(c+dx)}{d}$$

output `b^2*sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d`

### 3.155.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{(b \sec(c+dx))^{5/2}}{\sqrt{\sec(c+dx)}} dx = \frac{(b \sec(c+dx))^{5/2} \sin(c+dx)}{d \sec^{3/2}(c+dx)}$$

input `Integrate[(b*Sec[c + d*x])^(5/2)/Sqrt[Sec[c + d*x]],x]`

output `((b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(d*Sec[c + d*x]^(3/2))`

**3.155.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \sec(c + dx))^{5/2}}{\sqrt{\sec(c + dx)}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \int \sec^2(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \int \csc(c + dx + \frac{\pi}{2})^2 dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{4254} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \int 1 d(-\tan(c + dx))}{d \sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{b^2 \sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(5/2)/Sqrt[Sec[c + d*x]],x]`

output `(b^2*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d`

**3.155.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] :-> Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] :-> Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

---

3.155.  $\int \frac{(b \sec(c+dx))^{5/2}}{\sqrt{\sec(c+dx)}} dx$

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

### 3.155.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{b^2 \sec(dx+c)^{\frac{3}{2}} \sqrt{b \sec(dx+c)} \cos(dx+c) \sin(dx+c)}{d}$	38
risch	$\frac{2ib^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d(e^{2i(dx+c)+1})}}$	74

```
input int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/d*b^2*sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2)*cos(d*x+c)*sin(d*x+c)
```

### 3.155.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sqrt{\sec(c + dx)}} dx = \frac{b^2 \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx + c)}{d \sqrt{\cos(dx + c)}}$$

```
input integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="fracas")
```

```
output b^2*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))
```

**3.155.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(1/2),x)`output `Timed out`**3.155.7 Maxima [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sqrt{\sec(c + dx)}} dx = \frac{2 b^{5/2} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c))^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1} d$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`output `2*b^(5/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)`**3.155.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sqrt{\sec(c + dx)}} dx = \int \frac{(b \sec(dx + c))^{5/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(5/2)/sqrt(sec(d*x + c)), x)`

**3.155.9 Mupad [B] (verification not implemented)**

Time = 13.54 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.89

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sqrt{\sec(c + dx)}} dx = \frac{b^2 \sqrt{\frac{b}{\cos(c+dx)}} (\cos(2c + 2dx) \operatorname{li} + \sin(2c + 2dx) + \operatorname{li})}{d (\cos(2c + 2dx) + 1) \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(1/2),x)`output `(b^2*(b/cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*li + sin(2*c + 2*d*x) + li)) / (d*(cos(2*c + 2*d*x) + 1)*(1/cos(c + d*x))^(1/2))`



**3.156** 
$$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.156.1 Optimal result . . . . . 948  
 3.156.2 Mathematica [A] (verified) . . . . . 948  
 3.156.3 Rubi [A] (verified) . . . . . 949  
 3.156.4 Maple [A] (verified) . . . . . 950  
 3.156.5 Fricas [A] (verification not implemented) . . . . . 950  
 3.156.6 Sympy [F(-1)] . . . . . 951  
 3.156.7 Maxima [B] (verification not implemented) . . . . . 951  
 3.156.8 Giac [F] . . . . . 951  
 3.156.9 Mupad [F(-1)] . . . . . 952

**3.156.1 Optimal result**

Integrand size = 23, antiderivative size = 36

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{b^2 \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}}$$

output `b^2*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

**3.156.2 Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{\operatorname{arctanh}(\sin(c + dx))(b \sec(c + dx))^{5/2}}{d \sec^{\frac{5}{2}}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(3/2),x]`

output `(ArcTanh[Sin[c + d*x]]*(b*Sec[c + d*x])^(5/2))/(d*Sec[c + d*x]^(5/2))`

### 3.156.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{3/2}(c + dx)} dx$$

$$\downarrow \text{2031}$$

$$\frac{b^2 \sqrt{b \sec(c + dx)} \int \sec(c + dx) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{b^2 \sqrt{b \sec(c + dx)} \int \csc(c + dx + \frac{\pi}{2}) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{4257}$$

$$\frac{b^2 \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}}$$

input `Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(3/2),x]`

output `(b^2*ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]])/(d*Sqrt[Sec[c + d*x]])`

#### 3.156.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.156.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.36

method	result	size
default	$\frac{-2 \cos(dx+c) \sqrt{\sec(dx+c)} \sqrt{b \sec(dx+c)} \operatorname{arctanh}(\cot(dx+c) - \csc(dx+c)) b^2}{d}$	49
risch	$\frac{b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)+i})}}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} d}} - \frac{b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)-i})}}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} d}}$	145

input `int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `-2/d*cos(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)*arctanh(cot(d*x+c)-csc(d*x+c))*b^2`

### 3.156.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.17

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{3}{2}}(c + dx)} dx = \left[ \frac{b^{\frac{5}{2}} \log \left( -\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right)}{2d}, \right. \\ \left. - \frac{\sqrt{-bb^2} \arctan \left( \frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b} \right)}{d} \right]$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algorithm="fracas")`

output `[1/2*b^(5/2)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/d, -sqrt(-b)*b^2*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/d]`

---

3.156.  $\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{3}{2}}(c+dx)} dx$

**3.156.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(3/2),x)`

output `Timed out`

**3.156.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(32) = 64.

Time = 0.42 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.00

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{3/2}(c + dx)} dx = \frac{(b^2 \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - b^2 \log(\cos(dx + c)^2)}{2d}$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `1/2*(b^2*log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - b^2*log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))*sqrt(b)/d`

**3.156.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{3/2}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{5/2}}{\sec^{3/2}(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(3/2), x)`

**3.156.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{3/2}(c + dx)} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(3/2),x)`output `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(3/2), x)`

**3.157** 
$$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{5}{2}}(c+dx)} dx$$

3.157.1 Optimal result . . . . .	953
3.157.2 Mathematica [A] (verified) . . . . .	953
3.157.3 Rubi [A] (verified) . . . . .	954
3.157.4 Maple [A] (verified) . . . . .	955
3.157.5 Fricas [A] (verification not implemented) . . . . .	955
3.157.6 Sympy [A] (verification not implemented) . . . . .	955
3.157.7 Maxima [A] (verification not implemented) . . . . .	956
3.157.8 Giac [F] . . . . .	956
3.157.9 Mupad [B] (verification not implemented) . . . . .	956

**3.157.1 Optimal result**

Integrand size = 23, antiderivative size = 27

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{b^2 x \sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}}$$

output `b^2*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

**3.157.2 Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{x(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(5/2),x]`

output `(x*(b*Sec[c + d*x])^(5/2))/Sec[c + d*x]^(5/2)`

**3.157.3 Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2031, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{5/2}(c + dx)} dx$$

↓ 2031

$$\frac{b^2 \sqrt{b \sec(c + dx)} \int 1 dx}{\sqrt{\sec(c + dx)}}$$

↓ 24

$$\frac{b^2 x \sqrt{b \sec(c + dx)}}{\sqrt{\sec(c + dx)}}$$

input `Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(5/2),x]`

output `(b^2*x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]`

**3.157.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fv_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

**3.157.4 Maple [A] (verified)**

Time = 0.55 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

method	result	size
default	$\frac{(dx+c)b^2\sqrt{b\sec(dx+c)}}{d\sqrt{\sec(dx+c)}}$	31
risch	$\frac{b^2\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}x}{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}$	57

input `int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x,method=_RETURNVERBOSE)`output `1/d*(d*x+c)*b^2/sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)`**3.157.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 101, normalized size of antiderivative = 3.74

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)} dx = \left[ \frac{\sqrt{-bb^2} \log\left(-2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b\right)}{2d} \right]$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algorithm="fracas")`output `[1/2*sqrt(-b)*b^2*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/d, b^(5/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/d]`**3.157.6 Sympy [A] (verification not implemented)**

Time = 47.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{x(b \sec(c + dx))^{\frac{5}{2}}}{\sec^{\frac{5}{2}}(c + dx)}$$



input `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(5/2),x)`

output `x*(b*sec(c + d*x))**(5/2)/sec(c + d*x)**(5/2)`

### 3.157.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{2 b^{\frac{5}{2}} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `2*b^(5/2)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d`

### 3.157.8 Giac [F]

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{\frac{5}{2}}}{\sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(5/2), x)`

### 3.157.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{b^2 x \sqrt{\frac{b}{\cos(c+dx)}}}{\sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(5/2),x)`

output `(b^2*x*(b/cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2)`

---

3.157.  $\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{5}{2}}(c+dx)} dx$

**3.158** 
$$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{7/2}(c+dx)} dx$$

3.158.1 Optimal result . . . . . 957  
 3.158.2 Mathematica [A] (verified) . . . . . 957  
 3.158.3 Rubi [A] (verified) . . . . . 958  
 3.158.4 Maple [A] (verified) . . . . . 959  
 3.158.5 Fricas [A] (verification not implemented) . . . . . 959  
 3.158.6 Sympy [F(-1)] . . . . . 960  
 3.158.7 Maxima [A] (verification not implemented) . . . . . 960  
 3.158.8 Giac [F] . . . . . 960  
 3.158.9 Mupad [B] (verification not implemented) . . . . . 961

**3.158.1 Optimal result**

Integrand size = 23, antiderivative size = 35

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{7/2}(c + dx)} dx = \frac{b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sec(c + dx)}}$$

output `b^2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

**3.158.2 Mathematica [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{7/2}(c + dx)} dx = \frac{(b \sec(c + dx))^{5/2} \sin(c + dx)}{d \sec^{5/2}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(7/2),x]`

output `((b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(d*Sec[c + d*x]^(5/2))`

**3.158.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^2(c + dx)} dx$$

$$\downarrow \text{2031}$$

$$\frac{b^2 \sqrt{b \sec(c + dx)} \int \cos(c + dx) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{b^2 \sqrt{b \sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2}) dx}{\sqrt{\sec(c + dx)}}$$

$$\downarrow \text{3117}$$

$$\frac{b^2 \sin(c + dx) \sqrt{b \sec(c + dx)}}{d \sqrt{\sec(c + dx)}}$$

input `Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(7/2),x]`

output `(b^2*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])`

**3.158.3.1 Defintions of rubi rules used**

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

---

3.158.  $\int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx$

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`  
`FreeQ[{c, d}, x]`

### 3.158.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{b^2 \sqrt{b \sec(dx+c)} \tan(dx+c)}{d \sec(dx+c)^{\frac{3}{2}}}$	32
risch	$-\frac{ib^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} e^{i(dx+c)}}}{2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{ib^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} e^{-i(dx+c)}}}{2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}$	140

input `int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2),x,method=_RETURNVERBOSE)`

output `1/d*b^2/sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2)*tan(d*x+c)`

### 3.158.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{7}{2}}(c + dx)} dx = \frac{b^2 \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{d}$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2),x, algorithm="fracas")`

output `b^2*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/d`

**3.158.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(7/2),x)`output `Timed out`**3.158.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.37

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{7/2}(c + dx)} dx = \frac{b^{5/2} \sin(dx + c)}{d}$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2),x, algorithm="maxima")`output `b^(5/2)*sin(d*x + c)/d`**3.158.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{7/2}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{5/2}}{\sec(dx + c)^{7/2}} dx$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(7/2), x)`

**3.158.9 Mupad [B] (verification not implemented)**

Time = 13.79 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{7/2}(c + dx)} dx = \frac{b^2 \sin(c + dx) \sqrt{\frac{b}{\cos(c+dx)}}}{d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(7/2),x)`output `(b^2*sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(d*(1/cos(c + d*x))^(1/2))`

**3.159** 
$$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{9}{2}}(c+dx)} dx$$

3.159.1 Optimal result . . . . .	962
3.159.2 Mathematica [A] (verified) . . . . .	962
3.159.3 Rubi [A] (verified) . . . . .	963
3.159.4 Maple [A] (verified) . . . . .	964
3.159.5 Fricas [A] (verification not implemented) . . . . .	965
3.159.6 Sympy [F(-1)] . . . . .	965
3.159.7 Maxima [A] (verification not implemented) . . . . .	965
3.159.8 Giac [F] . . . . .	966
3.159.9 Mupad [B] (verification not implemented) . . . . .	966

**3.159.1 Optimal result**

Integrand size = 23, antiderivative size = 69

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \frac{b^2 x \sqrt{b \sec(c + dx)}}{2 \sqrt{\sec(c + dx)}} + \frac{b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{2d \sec^{\frac{3}{2}}(c + dx)}$$

output `1/2*b^2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+1/2*b^2*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)`

**3.159.2 Mathematica [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \frac{(b \sec(c + dx))^{5/2}(2(c + dx) + \sin(2(c + dx)))}{4d \sec^{\frac{5}{2}}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(9/2),x]`

output `((b*Sec[c + d*x])^(5/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sec[c + d*x]^(5/2))`

**3.159.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.74, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{9}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \int \cos^2(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2})^2 dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \left( \frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right)}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \left( \frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right)}{\sqrt{\sec(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(9/2),x]`

output `(b^2*sqrt[b*Sec[c + d*x]]*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/sqrt[Sec[c + d*x]]`



## 3.159.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

## 3.159.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\sqrt{b \sec(dx+c)} b^2 (\tan(dx+c) + (dx+c) \sec(dx+c)^2)}{2d \sec(dx+c)^{\frac{5}{2}}}$	48
risch	$\frac{b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} x}{2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{i b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{2i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} d + \frac{i b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-2i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} d$	197

input `int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2),x,method=_RETURNVERBOSE)`

output `1/2/d*(b*sec(d*x+c))^(1/2)*b^2/sec(d*x+c)^(5/2)*(tan(d*x+c)+(d*x+c)*sec(d*x+c)^2)`

**3.159.5 Fracas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 167, normalized size of antiderivative = 2.42

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \left[ \frac{2b^2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + \sqrt{-bb^2} \log\left(-2\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)\right)}{4d} \right]$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2),x, algorithm="fricas")`output `[1/4*(2*b^2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(-b)*b^2*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/2*(b^2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + b^(5/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/d]`**3.159.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(9/2),x)`output `Timed out`**3.159.7 Maxima [A] (verification not implemented)**

Time = 0.44 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.46

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{\frac{9}{2}}(c + dx)} dx = \frac{(2(dx+c)b^2 + b^2 \sin(2dx+2c))\sqrt{b}}{4d}$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2),x, algorithm="maxima")`output `1/4*(2*(d*x + c)*b^2 + b^2*sin(2*d*x + 2*c))*sqrt(b)/d`

---

3.159.  $\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{\frac{9}{2}}(c+dx)} dx$

**3.159.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^2(c + dx)} dx = \int \frac{(b \sec(dx + c))^{5/2}}{\sec^2(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(9/2), x)`

**3.159.9 Mupad [B] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.64

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^2(c + dx)} dx = \frac{b^2 (\sin(2c + 2dx) + 2dx) \sqrt{\frac{b}{\cos(c+dx)}}}{4d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(9/2),x)`

output `(b^2*(sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*d*(1/cos(c + d*x))^(1/2))`

**3.160** 
$$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{11/2}(c+dx)} dx$$

3.160.1 Optimal result . . . . . 967  
 3.160.2 Mathematica [A] (verified) . . . . . 967  
 3.160.3 Rubi [A] (verified) . . . . . 968  
 3.160.4 Maple [A] (verified) . . . . . 969  
 3.160.5 Fricas [A] (verification not implemented) . . . . . 970  
 3.160.6 Sympy [F(-1)] . . . . . 970  
 3.160.7 Maxima [A] (verification not implemented) . . . . . 970  
 3.160.8 Giac [F] . . . . . 971  
 3.160.9 Mupad [B] (verification not implemented) . . . . . 971

**3.160.1 Optimal result**

Integrand size = 23, antiderivative size = 76

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx = \frac{b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sec(c + dx)}} - \frac{b^2 \sqrt{b \sec(c + dx)} \sin^3(c + dx)}{3d \sqrt{\sec(c + dx)}}$$

output `b^2*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)-1/3*b^2*sin(d*x+c)^3*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)`

**3.160.2 Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx = \frac{(5 + \cos(2(c + dx)))(b \sec(c + dx))^{5/2} \sin(c + dx)}{6d \sec^{5/2}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(11/2),x]`

output `((5 + Cos[2*(c + d*x)])*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(6*d*Sec[c + d*x]^(5/2))`

**3.160.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.67, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \int \cos^3(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 \sqrt{b \sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2})^3 dx}{\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3113} \\
 & - \frac{b^2 \sqrt{b \sec(c + dx)} \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{2009} \\
 & - \frac{b^2 (\frac{1}{3} \sin^3(c + dx) - \sin(c + dx)) \sqrt{b \sec(c + dx)}}{d\sqrt{\sec(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(11/2),x]`

output `-((b^2*sqrt[b*Sec[c + d*x]]*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/(d*sqrt[Sec[c + d*x]]))`

## 3.160.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

## 3.160.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.66

method	result	size
default	$\frac{b^2 \sqrt{b \sec(dx+c)} \left( \tan(dx+c) + 2 \tan(dx+c) \sec(dx+c)^2 \right)}{3d \sec(dx+c)^{\frac{7}{2}}}$	50
risch	$-\frac{3ib^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{3ib^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-i(dx+c)}}{8 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \sin(3dx+3c)}{12 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}$	208

input `int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2),x,method=_RETURNVERBOSE)`

output `1/3*b^2/d*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2)*(tan(d*x+c)+2*tan(d*x+c)*sec(d*x+c)^2)`

**3.160.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.72

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx = \frac{(b^2 \cos(dx + c)^3 + 2b^2 \cos(dx + c)) \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c)}{3d\sqrt{\cos(dx + c)}}$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2),x, algorithm="fracas")`output `1/3*(b^2*cos(d*x + c)^3 + 2*b^2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))`**3.160.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(11/2),x)`output `Timed out`**3.160.7 Maxima [A] (verification not implemented)**

Time = 0.44 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.64

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx = \frac{(b^2 \sin(3dx + 3c) + 9b^2 \sin(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c)))) \sqrt{b}}{12d}$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2),x, algorithm="maxima")`output `1/12*(b^2*sin(3*d*x + 3*c) + 9*b^2*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*sqrt(b)/d`

**3.160.8 Giac [F]**

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx = \int \frac{(b \sec(dx + c))^{5/2}}{\sec^{11/2}(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(11/2), x)`

**3.160.9 Mupad [B] (verification not implemented)**

Time = 13.89 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.63

$$\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx = \frac{b^2 (9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(11/2),x)`

output `(b^2*(9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*d*(1/cos(c + d*x))^(1/2))`



$$\mathbf{3.161} \quad \int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

3.161.1 Optimal result . . . . .	972
3.161.2 Mathematica [A] (verified) . . . . .	972
3.161.3 Rubi [A] (verified) . . . . .	973
3.161.4 Maple [A] (verified) . . . . .	974
3.161.5 Fricas [A] (verification not implemented) . . . . .	975
3.161.6 Sympy [F(-1)] . . . . .	975
3.161.7 Maxima [B] (verification not implemented) . . . . .	976
3.161.8 Giac [F] . . . . .	977
3.161.9 Mupad [F(-1)] . . . . .	977

### 3.161.1 Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\sec(c+dx)}}{2d\sqrt{b \sec(c+dx)}} + \frac{\sec^{\frac{5}{2}}(c+dx)\sin(c+dx)}{2d\sqrt{b \sec(c+dx)}}$$

output `1/2*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(b*sec(d*x+c))^(1/2)+1/2*arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/d/(b*sec(d*x+c))^(1/2)`

### 3.161.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.69

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\sqrt{\sec(c+dx)}(\operatorname{arctanh}(\sin(c+dx)) + \sec(c+dx)\tan(c+dx))}{2d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^(7/2)/Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sqrt[b*Sec[c + d*x]])`

---


$$3.161. \quad \int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

**3.161.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.79, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {2031, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx}{\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{\sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(7/2)/Sqrt[b*Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/Sqrt[b*Sec[c + d*x]]`

---

3.161.  $\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.161.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.161.4 Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.28

method	result	S
default	$-\frac{\sec(dx+c)^{\frac{7}{2}} \left( \cos(dx+c)^3 \ln(-\cot(dx+c)+\csc(dx+c)-1) - \cos(dx+c)^3 \ln(\csc(dx+c)-\cot(dx+c)+1) - \cos(dx+c) \sin(dx+c) \right)}{2d\sqrt{b \sec(dx+c)}}$	9
risch	$-i \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (e^{3i(dx+c)} - e^{i(dx+c)})}{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} d (e^{2i(dx+c)}+1)^2} - \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \ln(e^{i(dx+c)}-i)}{2\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} d} + \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \ln(e^{i(dx+c)}+i)}{2\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} d}$	2

input `int(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2/d*sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2)*(cos(d*x+c)^3*ln(-cot(d*x+c)+csc(d*x+c)-1)-cos(d*x+c)^3*ln(csc(d*x+c)-cot(d*x+c)+1)-cos(d*x+c)*sin(d*x+c))`

3.161. 
$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

**3.161.5 Fracas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.85

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

$$= \left[ \frac{\sqrt{b} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4bd \cos(dx+c)}, \right.$$

$$\left. - \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right) \cos(dx+c) - \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{2bd \cos(dx+c)} \right]$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fracas")`output `[1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b*d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b*d*cos(d*x + c))]`**3.161.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(7/2)/(b*sec(d*x+c))**(1/2),x)`output `Timed out`

**3.161.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 661 vs. 2(60) = 120.

Time = 0.41 (sec) , antiderivative size = 661, normalized size of antiderivative = 9.18

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx =$$

$$\frac{4(\sin(4dx+4c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right) - 4(\sin(4dx+2c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right) + 4(\sin(4dx+4c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right) - 4(\sin(4dx+2c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right) + \dots}{\sqrt{b \sec(c+dx)}}$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output

```
-1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x
+ 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos
(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4
*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2
+ 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*s
in(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x
+ 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2
+ sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x +
2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^
2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4
*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c))) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*a
rctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))/((2*(2*cos(2*d*x + 2*c) + 1)*
cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x +
4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*c
os(2*d*x + 2*c) + 1)*sqrt(b)*d)
```

**3.161.8 Giac [F]**

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\sec(dx+c)^{\frac{7}{2}}}{\sqrt{b \sec(dx+c)}} dx$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(7/2)/sqrt(b*sec(d*x + c)), x)`

**3.161.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{7/2}}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(1/2),x)`

output `int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(1/2), x)`

$$3.162 \quad \int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

3.162.1 Optimal result . . . . .	978
3.162.2 Mathematica [A] (verified) . . . . .	978
3.162.3 Rubi [A] (verified) . . . . .	979
3.162.4 Maple [A] (verified) . . . . .	980
3.162.5 Fricas [A] (verification not implemented) . . . . .	981
3.162.6 Sympy [F(-1)] . . . . .	981
3.162.7 Maxima [B] (verification not implemented) . . . . .	981
3.162.8 Giac [F] . . . . .	982
3.162.9 Mupad [B] (verification not implemented) . . . . .	982

### 3.162.1 Optimal result

Integrand size = 23, antiderivative size = 32

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{d \sqrt{b \sec(c+dx)}}$$

output `sec(d*x+c)^(3/2)*sin(d*x+c)/d/(b*sec(d*x+c))^(1/2)`

### 3.162.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{d \sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^(5/2)/Sqrt[b*Sec[c + d*x]],x]`

output `(Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])`

**3.162.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sec^2(c+dx) dx}{\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx}{\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\sqrt{\sec(c+dx)} \int 1 d(-\tan(c+dx))}{d\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d\sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(5/2)/Sqrt[b*Sec[c + d*x]],x]`

output `(Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])`



## 3.162.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

## 3.162.4 Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{\sec(dx+c)^{\frac{5}{2}} \cos(dx+c) \sin(dx+c)}{d \sqrt{b \sec(dx+c)}}$	35
risch	$\frac{2i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}}}{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} d (e^{2i(dx+c)}+1)}$	71

input `int(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/d*sec(d*x+c)^(5/2)*cos(d*x+c)*sin(d*x+c)/(b*sec(d*x+c))^(1/2)`

**3.162.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.03

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b}\sec(c+dx)} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{bd\sqrt{\cos(dx+c)}}$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fracas")`

output `sqrt(b/cos(d*x + c))*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))`

**3.162.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b}\sec(c+dx)} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(1/2),x)`

output `Timed out`

**3.162.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 59 vs.  $2(28) = 56$ .

Time = 0.42 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.84

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b}\sec(c+dx)} dx = \frac{2\sqrt{b}\sin(2dx+2c)}{(b\cos(2dx+2c)^2 + b\sin(2dx+2c)^2 + 2b\cos(2dx+2c) + b)d}$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `2*sqrt(b)*sin(2*d*x + 2*c)/((b*cos(2*d*x + 2*c)^2 + b*sin(2*d*x + 2*c)^2 + 2*b*cos(2*d*x + 2*c) + b)*d)`

**3.162.8 Giac [F]**

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\sec(dx+c)^{\frac{5}{2}}}{\sqrt{b \sec(dx+c)}} dx$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(5/2)/sqrt(b*sec(d*x + c)), x)`

**3.162.9 Mupad [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.59

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

$$= \frac{(\cos(dx) - \sin(dx) \operatorname{li}) (\cos(c) - \sin(c) \operatorname{li}) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} \operatorname{li}}{bd}$$

input `int((1/cos(c + d*x))^(5/2)/(b/cos(c + d*x))^(1/2),x)`

output `((cos(d*x) - sin(d*x)*1i)*(cos(c) - sin(c)*1i)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*1i)/(b*d)`

**3.163** 
$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

3.163.1 Optimal result . . . . . 983  
 3.163.2 Mathematica [A] (verified) . . . . . 983  
 3.163.3 Rubi [A] (verified) . . . . . 984  
 3.163.4 Maple [A] (verified) . . . . . 985  
 3.163.5 Fricas [A] (verification not implemented) . . . . . 985  
 3.163.6 Sympy [F] . . . . . 986  
 3.163.7 Maxima [B] (verification not implemented) . . . . . 986  
 3.163.8 Giac [F] . . . . . 986  
 3.163.9 Mupad [F(-1)] . . . . . 987

**3.163.1 Optimal result**

Integrand size = 23, antiderivative size = 33

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}}$$

output `arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/d/(b*sec(d*x+c))^(1/2)`

**3.163.2 Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^(3/2)/Sqrt[b*Sec[c + d*x]],x]`

output `(ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(d*Sqrt[b*Sec[c + d*x]])`

**3.163.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

$$\downarrow \text{2031}$$

$$\frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{\sqrt{b \sec(c+dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{\sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx}{\sqrt{b \sec(c+dx)}}$$

$$\downarrow \text{4257}$$

$$\frac{\sqrt{\sec(c+dx)} \operatorname{arctanh}(\sin(c+dx))}{d\sqrt{b \sec(c+dx)}}$$

input `Int[Sec[c + d*x]^(3/2)/Sqrt[b*Sec[c + d*x]],x]`

output `(ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(d*Sqrt[b*Sec[c + d*x]])`

**3.163.3.1 Defintions of rubi rules used**

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] :> Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.163.4 Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

method	result	size
default	$-\frac{2 \cos(dx+c) \sec(dx+c)^{\frac{3}{2}} \operatorname{arctanh}(\cot(dx+c) - \csc(dx+c))}{d \sqrt{b \sec(dx+c)}}$	46
risch	$-\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)} - i)}}{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} d}} + \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)} + i)}}{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}} d}}$	139

input `int(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/d*cos(d*x+c)*sec(d*x+c)^(3/2)*arctanh(cot(d*x+c)-csc(d*x+c))/(b*sec(d*x+c))^(1/2)`

### 3.163.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.45

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \left[ \frac{\log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2\sqrt{bd}}, \right. \\ \left. -\frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{bd} \right]$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `[1/2*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2/(sqrt(b)*d), -sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/(b*d)]`

---

3.163.  $\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

**3.163.6 Sympy [F]**

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

input `integrate(sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(1/2),x)`

output `Integral(sec(c + d*x)**(3/2)/sqrt(b*sec(c + d*x)), x)`

**3.163.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs.  $2(29) = 58$ .

Time = 0.40 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.97

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

$$= \frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2\sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2\sin(dx+c) + 1)}{2\sqrt{bd}}$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(sqrt(b)*d)`

**3.163.8 Giac [F]**

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\sec^{\frac{3}{2}}(dx+c)}{\sqrt{b \sec(dx+c)}} dx$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(3/2)/sqrt(b*sec(d*x + c)), x)`

---

3.163.  $\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

**3.163.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int((1/cos(c + d*x))^(3/2)/(b/cos(c + d*x))^(1/2),x)`output `int((1/cos(c + d*x))^(3/2)/(b/cos(c + d*x))^(1/2), x)`



**3.164**       $\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx$

3.164.1 Optimal result . . . . .	988
3.164.2 Mathematica [A] (verified) . . . . .	988
3.164.3 Rubi [A] (verified) . . . . .	989
3.164.4 Maple [A] (verified) . . . . .	990
3.164.5 Fricas [A] (verification not implemented) . . . . .	990
3.164.6 Sympy [A] (verification not implemented) . . . . .	991
3.164.7 Maxima [A] (verification not implemented) . . . . .	991
3.164.8 Giac [F] . . . . .	991
3.164.9 Mupad [B] (verification not implemented) . . . . .	992

**3.164.1 Optimal result**

Integrand size = 23, antiderivative size = 24

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx = \frac{x \sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}}$$

output `x*sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`

**3.164.2 Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx = \frac{x \sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sqrt[Sec[c + d*x]]/Sqrt[b*Sec[c + d*x]],x]`

output `(x*Sqrt[Sec[c + d*x]])/Sqrt[b*Sec[c + d*x]]`

**3.164.3 Rubi [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2031, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx$$

↓ 2031

$$\frac{\sqrt{\sec(c+dx)} \int 1 dx}{\sqrt{b \sec(c+dx)}}$$

↓ 24

$$\frac{x \sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}}$$

input `Int[Sqrt[Sec[c + d*x]]/Sqrt[b*Sec[c + d*x]],x]`

output `(x*Sqrt[Sec[c + d*x]])/Sqrt[b*Sec[c + d*x]]`

**3.164.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fv_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

**3.164.4 Maple [A] (verified)**

Time = 0.52 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{(dx+c)\sqrt{\sec(dx+c)}}{d\sqrt{b\sec(dx+c)}}$	28
risch	$\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} x}}{\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}$	54

input `int(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`output `1/d*(d*x+c)*sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)`**3.164.5 Fracas [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 101, normalized size of antiderivative = 4.21

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b\sec(c+dx)}} dx$$

$$= \left[ \frac{\sqrt{-b} \log \left( 2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b \right)}{2bd}, \frac{\arctan \left( \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{b} \sqrt{\cos(dx+c)}} \right)}{\sqrt{bd}} \right]$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fracas")`output `[-1/2*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/(b*d), arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))/(sqrt(b)*d)]`

**3.164.6 Sympy [A] (verification not implemented)**

Time = 3.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b\sec(c+dx)}} dx = \frac{x\sqrt{\sec(c+dx)}}{\sqrt{b\sec(c+dx)}}$$

input `integrate(sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(1/2),x)`output `x*sqrt(sec(c + d*x))/sqrt(b*sec(c + d*x))`**3.164.7 Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b\sec(c+dx)}} dx = \frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{\sqrt{bd}}$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`output `2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(sqrt(b)*d)`**3.164.8 Giac [F]**

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b\sec(c+dx)}} dx = \int \frac{\sqrt{\sec(dx+c)}}{\sqrt{b\sec(dx+c)}} dx$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`output `integrate(sqrt(sec(d*x + c))/sqrt(b*sec(d*x + c)), x)`

**3.164.9 Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12

$$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx = \frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{b \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((1/cos(c + d*x))^(1/2)/(b/cos(c + d*x))^(1/2),x)`

output `(x*(b/cos(c + d*x))^(1/2))/(b*(1/cos(c + d*x))^(1/2))`

**3.165**  $\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}} dx$

3.165.1 Optimal result . . . . . 993  
 3.165.2 Mathematica [A] (verified) . . . . . 993  
 3.165.3 Rubi [A] (verified) . . . . . 994  
 3.165.4 Maple [A] (verified) . . . . . 995  
 3.165.5 Fricas [A] (verification not implemented) . . . . . 995  
 3.165.6 Sympy [A] (verification not implemented) . . . . . 996  
 3.165.7 Maxima [A] (verification not implemented) . . . . . 996  
 3.165.8 Giac [F] . . . . . 996  
 3.165.9 Mupad [B] (verification not implemented) . . . . . 997

**3.165.1 Optimal result**

Integrand size = 23, antiderivative size = 32

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{d\sqrt{b \sec(c+dx)}}$$

output `sin(d*x+c)*sec(d*x+c)^(1/2)/d/(b*sec(d*x+c))^(1/2)`

**3.165.2 Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{d\sqrt{b \sec(c+dx)}}$$

input `Integrate[1/(Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]),x]`

output `(Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])`

**3.165.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2032, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} dx$$

↓ 2032

$$\frac{\sqrt{\sec(c+dx)} \int \cos(c+dx) dx}{\sqrt{b\sec(c+dx)}}$$

↓ 3042

$$\frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2}) dx}{\sqrt{b\sec(c+dx)}}$$

↓ 3117

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{d\sqrt{b\sec(c+dx)}}$$

input `Int[1/(Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]),x]`

output `(Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])`

**3.165.3.1 Defintions of rubi rules used**

rule 2032 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3117 Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

### 3.165.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\tan(dx+c)}{d\sqrt{\sec(dx+c)}\sqrt{b\sec(dx+c)}}$	29
risch	$-\frac{ie^{2i(dx+c)}}{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}d + \frac{i}{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}d$	151

```
input int(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)*tan(d*x+c)
```

### 3.165.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.03

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}}\sqrt{\cos(dx+c)}\sin(dx+c)}{bd}$$

```
input integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b*d)
```



**3.165.6 Sympy [A] (verification not implemented)**

Time = 6.53 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.44

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} dx = \begin{cases} \frac{\tan(c+dx)}{d\sqrt{b\sec(c+dx)}\sqrt{\sec(c+dx)}} & \text{for } d \neq 0 \\ \frac{x}{\sqrt{b\sec(c)}\sqrt{\sec(c)}} & \text{otherwise} \end{cases}$$

input `integrate(1/sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(1/2),x)`output `Piecewise((tan(c + d*x)/(d*sqrt(b*sec(c + d*x))*sqrt(sec(c + d*x))), Ne(d, 0)), (x/(sqrt(b*sec(c))*sqrt(sec(c))), True))`**3.165.7 Maxima [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.41

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} dx = \frac{\sin(dx+c)}{\sqrt{bd}}$$

input `integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`output `sin(d*x + c)/(sqrt(b)*d)`**3.165.8 Giac [F]**

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} dx = \int \frac{1}{\sqrt{b\sec(dx+c)}\sqrt{\sec(dx+c)}} dx$$

input `integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`output `integrate(1/(sqrt(b*sec(d*x + c))*sqrt(sec(d*x + c))), x)`

**3.165.9 Mupad [B] (verification not implemented)**

Time = 13.73 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.09

$$\int \frac{1}{\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} dx = \frac{\sin(c+dx) \sqrt{\frac{b}{\cos(c+dx)}}}{bd \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)),x)`output `(sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(b*d*(1/cos(c + d*x))^(1/2))`

**3.166** 
$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b \sec(c+dx)}} dx$$

3.166.1 Optimal result . . . . .	998
3.166.2 Mathematica [A] (verified) . . . . .	998
3.166.3 Rubi [A] (verified) . . . . .	999
3.166.4 Maple [A] (verified) . . . . .	1000
3.166.5 Fracas [A] (verification not implemented) . . . . .	1001
3.166.6 Sympy [A] (verification not implemented) . . . . .	1001
3.166.7 Maxima [A] (verification not implemented) . . . . .	1002
3.166.8 Giac [F] . . . . .	1002
3.166.9 Mupad [B] (verification not implemented) . . . . .	1002

**3.166.1 Optimal result**

Integrand size = 23, antiderivative size = 63

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b \sec(c+dx)}} dx = \frac{x\sqrt{\sec(c+dx)}}{2\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2d\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}}$$

output  $\frac{1}{2}*\sin(d*x+c)/d/\sec(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+1/2*x*\sec(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

**3.166.2 Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b \sec(c+dx)}} dx = \frac{\sqrt{\sec(c+dx)}(2(c+dx) + \sin(2(c+dx)))}{4d\sqrt{b \sec(c+dx)}}$$

input `Integrate[1/(Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]]),x]`

output  $(\text{Sqrt}[\text{Sec}[c + d*x]]*(2*(c + d*x) + \text{Sin}[2*(c + d*x)]))/(4*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

**3.166.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.76, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2032, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx \\
 & \quad \downarrow \text{2032} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^2(c+dx) dx}{\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^2 dx}{\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\int 1 dx}{2} + \frac{\sin(c+dx)\cos(c+dx)}{2d} \right)}{\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\sin(c+dx)\cos(c+dx)}{2d} + \frac{x}{2} \right)}{\sqrt{b\sec(c+dx)}}
 \end{aligned}$$

input `Int[1/(Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]]),x]`

output `(Sqrt[Sec[c + d*x]]*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/Sqrt[b*Sec[c + d*x]]`

## 3.166.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2032 `Int[(Fx_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Simp[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

## 3.166.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

method	result
default	$\frac{\tan(dx+c)+(dx+c)\sec(dx+c)^2}{2d\sec(dx+c)^{\frac{3}{2}}\sqrt{b\sec(dx+c)}}$
risch	$\frac{e^{i(dx+c)}x}{2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}(e^{2i(dx+c)+1})}} - \frac{ie^{3i(dx+c)}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}(e^{2i(dx+c)+1})}d} + \frac{ie^{-i(dx+c)}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}(e^{2i(dx+c)+1})}}$

input `int(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/2/d/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2)*(tan(d*x+c)+(d*x+c)*sec(d*x+c)^2)`

**3.166.5 Fricas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 165, normalized size of antiderivative = 2.62

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx$$

$$= \frac{2\sqrt{\frac{b}{\cos(dx+c)}}\cos(dx+c)^{\frac{3}{2}}\sin(dx+c) - \sqrt{-b}\log\left(2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}}\cos(dx+c)^{\frac{3}{2}}\sin(dx+c) + 2b\cos(dx+c)\right)}{4bd}$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`output `[1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) - sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b*d), 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b*d)]`**3.166.6 Sympy [A] (verification not implemented)**

Time = 7.53 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.70

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx$$

$$= \begin{cases} \frac{x \tan^2(c+dx)}{2\sqrt{b\sec(c+dx)}\sec^{\frac{3}{2}}(c+dx)} + \frac{x}{2\sqrt{b\sec(c+dx)}\sec^{\frac{3}{2}}(c+dx)} + \frac{\tan(c+dx)}{2d\sqrt{b\sec(c+dx)}\sec^{\frac{3}{2}}(c+dx)} & \text{for } d \neq 0 \\ \frac{x}{\sqrt{b\sec(c)}\sec^{\frac{3}{2}}(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(1/2),x)`output `Piecewise((x*tan(c + d*x)**2/(2*sqrt(b*sec(c + d*x))*sec(c + d*x)**(3/2)) + x/(2*sqrt(b*sec(c + d*x))*sec(c + d*x)**(3/2)) + tan(c + d*x)/(2*d*sqrt(b*sec(c + d*x))*sec(c + d*x)**(3/2)), Ne(d, 0)), (x/(sqrt(b*sec(c))*sec(c)**(3/2)), True))`

**3.166.7 Maxima [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.40

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \frac{2dx + 2c + \sin(2dx + 2c)}{4\sqrt{bd}}$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`output `1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(sqrt(b)*d)`**3.166.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \int \frac{1}{\sqrt{b\sec(dx+c)}\sec(dx+c)^{\frac{3}{2}}} dx$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`output `integrate(1/(sqrt(b*sec(d*x + c))*sec(d*x + c)^(3/2)), x)`**3.166.9 Mupad [B] (verification not implemented)**

Time = 13.91 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \frac{(\sin(2c + 2dx) + 2dx)\sqrt{\frac{b}{\cos(c+dx)}}}{4bd\sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(3/2)),x)`output `((sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*b*d*(1/cos(c + d*x))^(1/2))`

**3.167**  $\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b \sec(c+dx)}} dx$

3.167.1 Optimal result . . . . . 1003  
 3.167.2 Mathematica [A] (verified) . . . . . 1003  
 3.167.3 Rubi [A] (verified) . . . . . 1004  
 3.167.4 Maple [A] (verified) . . . . . 1005  
 3.167.5 Fracas [A] (verification not implemented) . . . . . 1006  
 3.167.6 Sympy [A] (verification not implemented) . . . . . 1006  
 3.167.7 Maxima [A] (verification not implemented) . . . . . 1006  
 3.167.8 Giac [F] . . . . . 1007  
 3.167.9 Mupad [B] (verification not implemented) . . . . . 1007

**3.167.1 Optimal result**

Integrand size = 23, antiderivative size = 70

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b \sec(c+dx)}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{d\sqrt{b \sec(c+dx)}} - \frac{\sqrt{\sec(c+dx)} \sin^3(c+dx)}{3d\sqrt{b \sec(c+dx)}}$$

output `sin(d*x+c)*sec(d*x+c)^(1/2)/d/(b*sec(d*x+c))^(1/2)-1/3*sin(d*x+c)^3*sec(d*x+c)^(1/2)/d/(b*sec(d*x+c))^(1/2)`

**3.167.2 Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b \sec(c+dx)}} dx = \frac{(5 + \cos(2(c+dx)))\sqrt{\sec(c+dx)} \sin(c+dx)}{6d\sqrt{b \sec(c+dx)}}$$

input `Integrate[1/(Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]]),x]`

output `((5 + Cos[2*(c + d*x)])*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(6*d*Sqrt[b*Sec[c + d*x]])`



**3.167.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2032, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx \\
 & \quad \downarrow \text{2032} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^3(c+dx) dx}{\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^3 dx}{\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3113} \\
 & -\frac{\sqrt{\sec(c+dx)} \int (1 - \sin^2(c+dx)) d(-\sin(c+dx))}{d\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{(\frac{1}{3}\sin^3(c+dx) - \sin(c+dx))\sqrt{\sec(c+dx)}}{d\sqrt{b\sec(c+dx)}}
 \end{aligned}$$

input `Int[1/(Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]]),x]`

output `-((Sqrt[Sec[c + d*x]]*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/(d*Sqrt[b*Sec[c + d*x]]))`

## 3.167.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2032 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

## 3.167.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.67

method	result
default	$\frac{\tan(dx+c)+2\tan(dx+c)\sec(dx+c)^2}{3d\sec(dx+c)^{\frac{5}{2}}\sqrt{b\sec(dx+c)}}$
risch	$-\frac{ie^{4i(dx+c)}}{24\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}d - \frac{3ie^{2i(dx+c)}}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}d + \frac{3i}{8\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}}$

input `int(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/3/d/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2)*(tan(d*x+c)+2*tan(d*x+c)*sec(d*x+c)^2)`

**3.167.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.73

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \frac{(\cos(dx+c)^3 + 2\cos(dx+c))\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{3bd\sqrt{\cos(dx+c)}}$$

input `integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`output `1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))`**3.167.6 Sympy [A] (verification not implemented)**

Time = 23.97 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \begin{cases} \frac{2\tan^3(c+dx)}{3d\sqrt{b\sec(c+dx)}\sec^{\frac{5}{2}}(c+dx)} + \frac{\tan(c+dx)}{d\sqrt{b\sec(c+dx)}\sec^{\frac{5}{2}}(c+dx)} & \text{for } d \neq 0 \\ \frac{x}{\sqrt{b\sec(c)}\sec^{\frac{5}{2}}(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(1/2),x)`output `Piecewise((2*tan(c + d*x)**3/(3*d*sqrt(b*sec(c + d*x))*sec(c + d*x)**(5/2)) + tan(c + d*x)/(d*sqrt(b*sec(c + d*x))*sec(c + d*x)**(5/2)), Ne(d, 0)), (x/(sqrt(b*sec(c))*sec(c)**(5/2)), True))`**3.167.7 Maxima [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.60

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \frac{\sin(3dx+3c) + 9\sin\left(\frac{1}{3}\arctan\left(\frac{\sin(3dx+3c)}{\cos(3dx+3c)}\right)\right)}{12\sqrt{bd}}$$

input `integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(sqrt(b)*d)`

### 3.167.8 Giac [F]

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \int \frac{1}{\sqrt{b\sec(dx+c)}\sec(dx+c)^{\frac{5}{2}}} dx$$

input `integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(b*sec(d*x + c))*sec(d*x + c)^(5/2)), x)`

### 3.167.9 Mupad [B] (verification not implemented)

Time = 13.91 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \frac{(9 \sin(c+dx) + \sin(3c+3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12bd \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(5/2)),x)`

output `((9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*b*d*(1/cos(c + d*x))^(1/2))`

$$3.168 \quad \int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

3.168.1 Optimal result . . . . .	1008
3.168.2 Mathematica [A] (verified) . . . . .	1008
3.168.3 Rubi [A] (verified) . . . . .	1009
3.168.4 Maple [A] (verified) . . . . .	1010
3.168.5 Fricas [A] (verification not implemented) . . . . .	1011
3.168.6 Sympy [F(-1)] . . . . .	1011
3.168.7 Maxima [B] (verification not implemented) . . . . .	1012
3.168.8 Giac [F] . . . . .	1013
3.168.9 Mupad [F(-1)] . . . . .	1013

### 3.168.1 Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\sec(c+dx)}}{2bd\sqrt{b \sec(c+dx)}} + \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2bd\sqrt{b \sec(c+dx)}}$$

output `1/2*sec(d*x+c)^(5/2)*sin(d*x+c)/b/d/(b*sec(d*x+c))^(1/2)+1/2*arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/b/d/(b*sec(d*x+c))^(1/2)`

### 3.168.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.64

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\sec^{\frac{3}{2}}(c+dx)(\operatorname{arctanh}(\sin(c+dx)) + \sec(c+dx) \tan(c+dx))}{2d(b \sec(c+dx))^{3/2}}$$

input `Integrate[Sec[c + d*x]^(9/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(Sec[c + d*x]^(3/2)*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*(b*Sec[c + d*x])^(3/2))`

---


$$3.168. \quad \int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

**3.168.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.77, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {2031, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sec^3(c+dx) dx}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{b\sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(9/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(Sqrt[Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/(b*Sqrt[b*Sec[c + d*x]])`

---

3.168.  $\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.168.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.168.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.22

method	result
default	$-\frac{\sec(dx+c)^{\frac{7}{2}} \left( \cos(dx+c)^3 \ln(-\cot(dx+c)+\csc(dx+c)-1) - \cos(dx+c)^3 \ln(\csc(dx+c)-\cot(dx+c)+1) - \cos(dx+c) \sin(dx+c) \right)}{2db\sqrt{b \sec(dx+c)}}$
risch	$-\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \left( -e^{4i(dx+c)} \ln(e^{i(dx+c)}+i) + e^{4i(dx+c)} \ln(e^{i(dx+c)}-i) + 2ie^{3i(dx+c)} - 2ie^{i(dx+c)} - 2e^{2i(dx+c)} \ln(e^{i(dx+c)}+i) + 2e^{2i(dx+c)} \ln(e^{i(dx+c)}-i) \right)}{2b\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}} d(e^{2i(dx+c)}+1)^2}$

input `int(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-1/2/d*sec(d*x+c)^(7/2)/b/(b*sec(d*x+c))^(1/2)*(cos(d*x+c)^3*ln(-cot(d*x+c)+csc(d*x+c)-1)-cos(d*x+c)^3*ln(csc(d*x+c)-cot(d*x+c)+1)-cos(d*x+c)*sin(d*x+c))`

3.168.  $\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

**3.168.5 Fracas [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.63

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \left[ \frac{\sqrt{b} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}}}{\sqrt{\cos(dx+c)}}}{4b^2 d \cos(dx+c)} - \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right) \cos(dx+c) - \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{2b^2 d \cos(dx+c)} \right]$$

input `integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`output `[1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^2*d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^2*d*cos(d*x + c))]`**3.168.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(9/2)/(b*sec(d*x+c))**(3/2),x)`output `Timed out`



**3.168.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 670 vs. 2(66) = 132.

Time = 0.42 (sec) , antiderivative size = 670, normalized size of antiderivative = 8.59

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}} dx =$$


---


$$4(\sin(4dx+4c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan(\sin(2dx+2c), \cos(2dx+2c))\right) - 4(\sin(4dx+4c)$$

input `integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output

```
-1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x
+ 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos
(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4
*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2
+ 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*s
in(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x
+ 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2
+ sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x +
2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^
2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4
*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c))) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*a
rctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))/((b*cos(4*d*x + 4*c)^2 + 4*b*
cos(2*d*x + 2*c)^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x
+ 2*c) + 4*b*sin(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x
+ 4*c) + 4*b*cos(2*d*x + 2*c) + b)*sqrt(b)*d)
```

**3.168.8 Giac [F]**

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\sec(dx+c)^{\frac{9}{2}}}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(9/2)/(b*sec(d*x + c))^(3/2), x)`

**3.168.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{9/2}}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int((1/cos(c + d*x))^(9/2)/(b/cos(c + d*x))^(3/2),x)`

output `int((1/cos(c + d*x))^(9/2)/(b/cos(c + d*x))^(3/2), x)`

**3.169**  $\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.169.1 Optimal result . . . . . 1014  
 3.169.2 Mathematica [A] (verified) . . . . . 1014  
 3.169.3 Rubi [A] (verified) . . . . . 1015  
 3.169.4 Maple [A] (verified) . . . . . 1016  
 3.169.5 Fricas [A] (verification not implemented) . . . . . 1017  
 3.169.6 Sympy [F(-1)] . . . . . 1017  
 3.169.7 Maxima [B] (verification not implemented) . . . . . 1017  
 3.169.8 Giac [F] . . . . . 1018  
 3.169.9 Mupad [B] (verification not implemented) . . . . . 1018

**3.169.1 Optimal result**

Integrand size = 23, antiderivative size = 35

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{bd\sqrt{b \sec(c+dx)}}$$

output `sec(d*x+c)^(3/2)*sin(d*x+c)/b/d/(b*sec(d*x+c))^(1/2)`

**3.169.2 Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{d(b \sec(c+dx))^{3/2}}$$

input `Integrate[Sec[c + d*x]^(7/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(Sec[c + d*x]^(5/2)*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(3/2))`

**3.169.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sec^2(c+dx) dx}{b \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2})^2 dx}{b \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & - \frac{\sqrt{\sec(c+dx)} \int 1 d(-\tan(c+dx))}{bd \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{bd \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(7/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(Sec[c + d*x]^(3/2)*Sin[c + d*x])/(b*d*Sqrt[b*Sec[c + d*x]])`

## 3.169.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

## 3.169.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{\sec(dx+c)^{\frac{5}{2}} \cos(dx+c) \sin(dx+c)}{db \sqrt{b \sec(dx+c)}}$	38
risch	$\frac{2i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}{b \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} d(e^{2i(dx+c)+1})$	74

input `int(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `1/d*sec(d*x+c)^(5/2)*cos(d*x+c)*sin(d*x+c)/b/(b*sec(d*x+c))^(1/2)`

**3.169.5 Fricas [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{b^2 d \sqrt{\cos(dx+c)}}$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output `sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^2*d*sqrt(cos(d*x + c)))`

**3.169.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(7/2)/(b*sec(d*x+c))**(3/2),x)`

output `Timed out`

**3.169.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 67 vs.  $2(31) = 62$ .

Time = 0.40 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.91

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2\sqrt{b} \sin(2dx+2c)}{(b^2 \cos(2dx+2c)^2 + b^2 \sin(2dx+2c)^2 + 2b^2 \cos(2dx+2c) + b^2)d}$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `2*sqrt(b)*sin(2*d*x + 2*c)/((b^2*cos(2*d*x + 2*c)^2 + b^2*sin(2*d*x + 2*c)^2 + 2*b^2*cos(2*d*x + 2*c) + b^2)*d)`

**3.169.8 Giac [F]**

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\sec(dx+c)^{\frac{7}{2}}}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(7/2)/(b*sec(d*x + c))^(3/2), x)`

**3.169.9 Mupad [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.46

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{(\cos(dx) - \sin(dx) \operatorname{li}) (\cos(c) - \sin(c) \operatorname{li}) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} \operatorname{li}}{b^2 d}$$

input `int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(3/2),x)`

output `((cos(d*x) - sin(d*x)*1i)*(cos(c) - sin(c)*1i)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*1i)/(b^2*d)`

**3.170**  $\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.170.1 Optimal result . . . . . 1019  
 3.170.2 Mathematica [A] (verified) . . . . . 1019  
 3.170.3 Rubi [A] (verified) . . . . . 1020  
 3.170.4 Maple [A] (verified) . . . . . 1021  
 3.170.5 Fricas [A] (verification not implemented) . . . . . 1021  
 3.170.6 Sympy [F(-1)] . . . . . 1022  
 3.170.7 Maxima [B] (verification not implemented) . . . . . 1022  
 3.170.8 Giac [F] . . . . . 1022  
 3.170.9 Mupad [F(-1)] . . . . . 1023

**3.170.1 Optimal result**

Integrand size = 23, antiderivative size = 36

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\sec(c+dx)}}{bd\sqrt{b \sec(c+dx)}}$$

output `arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/b/d/(b*sec(d*x+c))^(1/2)`

**3.170.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{d(b \sec(c+dx))^{3/2}}$$

input `Integrate[Sec[c + d*x]^(5/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(ArcTanh[Sin[c + d*x]]*Sec[c + d*x]^(3/2))/(d*(b*Sec[c + d*x])^(3/2))`



**3.170.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

$$\downarrow \text{2031}$$

$$\frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{b \sqrt{b \sec(c+dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{\sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2}) dx}{b \sqrt{b \sec(c+dx)}}$$

$$\downarrow \text{4257}$$

$$\frac{\sqrt{\sec(c+dx)} \operatorname{arctanh}(\sin(c+dx))}{bd \sqrt{b \sec(c+dx)}}$$

input `Int[Sec[c + d*x]^(5/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(b*d*Sqrt[b*Sec[c + d*x]])`

**3.170.3.1 Defintions of rubi rules used**

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.170.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.36

method	result	size
default	$-\frac{2 \sec(dx+c)^{\frac{3}{2}} \cos(dx+c) \operatorname{arctanh}(\cot(dx+c) - \csc(dx+c))}{db \sqrt{b \sec(dx+c)}}$	49
risch	$\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)+i})}}{b \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} d - \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)-i})}}{b \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} d$	145

input `int(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output 
$$-2/d*\sec(d*x+c)^(3/2)*\cos(d*x+c)*\operatorname{arctanh}(\cot(d*x+c)-\csc(d*x+c))/b/(b*\sec(d*x+c))^(1/2)$$

### 3.170.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.17

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx = \left[ \frac{\log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2b^{\frac{3}{2}}d}, \right. \\ \left. - \frac{\sqrt{-b} \operatorname{arctan}\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{b^2 d} \right]$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output 
$$[1/2*\log(-(b*\cos(d*x+c))^2 - 2*\sqrt{b}*\sqrt{b/\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c) - 2*b)/\cos(d*x+c)^2/(b^(3/2)*d), -\sqrt{-b}*\operatorname{arctan}(\sqrt{-b}*\sqrt{b/\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)/b)/(b^2*d)]$$

---

3.170. 
$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx$$

**3.170.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(3/2),x)`output `Timed out`**3.170.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(32) = 64.

Time = 0.40 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.81

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx = \frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1)}{2 b^{\frac{3}{2}} d}$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`output `1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(b^(3/2)*d)`**3.170.8 Giac [F]**

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx = \int \frac{\sec(dx+c)^{\frac{5}{2}}}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`output `integrate(sec(d*x + c)^(5/2)/(b*sec(d*x + c))^(3/2), x)`

**3.170.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int((1/cos(c + d*x))^(5/2)/(b/cos(c + d*x))^(3/2),x)`output `int((1/cos(c + d*x))^(5/2)/(b/cos(c + d*x))^(3/2), x)`

**3.171** 
$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

3.171.1 Optimal result . . . . . 1024  
 3.171.2 Mathematica [A] (verified) . . . . . 1024  
 3.171.3 Rubi [A] (verified) . . . . . 1025  
 3.171.4 Maple [A] (verified) . . . . . 1026  
 3.171.5 Fricas [A] (verification not implemented) . . . . . 1026  
 3.171.6 Sympy [A] (verification not implemented) . . . . . 1026  
 3.171.7 Maxima [A] (verification not implemented) . . . . . 1027  
 3.171.8 Giac [F] . . . . . 1027  
 3.171.9 Mupad [B] (verification not implemented) . . . . . 1027

**3.171.1 Optimal result**

Integrand size = 23, antiderivative size = 27

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{x \sqrt{\sec(c + dx)}}{b \sqrt{b \sec(c + dx)}}$$

output `x*sec(d*x+c)^(1/2)/b/(b*sec(d*x+c))^(1/2)`

**3.171.2 Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{x \sqrt{b \sec(c + dx)}}{b^2 \sqrt{\sec(c + dx)}}$$

input `Integrate[Sec[c + d*x]^(3/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(x*Sqrt[b*Sec[c + d*x]])/(b^2*Sqrt[Sec[c + d*x]])`

**3.171.3 Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2031, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

↓ 2031

$$\frac{\sqrt{\sec(c+dx)} \int 1 dx}{b \sqrt{b \sec(c+dx)}}$$

↓ 24

$$\frac{x \sqrt{\sec(c+dx)}}{b \sqrt{b \sec(c+dx)}}$$

input `Int[Sec[c + d*x]^(3/2)/(b*Sec[c + d*x])^(3/2),x]`

output `(x*Sqrt[Sec[c + d*x]])/(b*Sqrt[b*Sec[c + d*x]])`

**3.171.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

**3.171.4 Maple [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

method	result	size
default	$\frac{(dx+c)\sqrt{\sec(dx+c)}}{db\sqrt{b\sec(dx+c)}}$	31
risch	$\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}x}}{b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}$	57

input `int(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`output `1/d*(d*x+c)*sec(d*x+c)^(1/2)/b/(b*sec(d*x+c))^(1/2)`**3.171.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 101, normalized size of antiderivative = 3.74

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \left[ -\frac{\sqrt{-b} \log\left(2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b\right)}{2b^2d}, \right]$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fracas")`output `[-1/2*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/(b^2*d), arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))/(b^(3/2)*d)]`**3.171.6 Sympy [A] (verification not implemented)**

Time = 5.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b\sec(c+dx))^{3/2}} dx = \frac{x \sec^{\frac{3}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{3}{2}}}$$

input `integrate(sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(3/2),x)`

output `x*sec(c + d*x)**(3/2)/(b*sec(c + d*x))**(3/2)`

### 3.171.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx = \frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{3}{2}}d}$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(b^(3/2)*d)`

### 3.171.8 Giac [F]

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx = \int \frac{\sec(dx + c)^{\frac{3}{2}}}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(3/2)/(b*sec(d*x + c))^(3/2), x)`

### 3.171.9 Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx = \frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{b^2 \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((1/cos(c + d*x))^(3/2)/(b/cos(c + d*x))^(3/2),x)`

output `(x*(b/cos(c + d*x))^(1/2))/(b^2*(1/cos(c + d*x))^(1/2))`

---

3.171.  $\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx$



**3.172**       $\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx$

3.172.1 Optimal result . . . . . 1028  
 3.172.2 Mathematica [A] (verified) . . . . . 1028  
 3.172.3 Rubi [A] (verified) . . . . . 1029  
 3.172.4 Maple [A] (verified) . . . . . 1030  
 3.172.5 Fricas [A] (verification not implemented) . . . . . 1030  
 3.172.6 Sympy [A] (verification not implemented) . . . . . 1031  
 3.172.7 Maxima [A] (verification not implemented) . . . . . 1031  
 3.172.8 Giac [F] . . . . . 1031  
 3.172.9 Mupad [B] (verification not implemented) . . . . . 1032

**3.172.1 Optimal result**

Integrand size = 23, antiderivative size = 35

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{bd\sqrt{b \sec(c+dx)}}$$

output `sin(d*x+c)*sec(d*x+c)^(1/2)/b/d/(b*sec(d*x+c))^(1/2)`

**3.172.2 Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx = \frac{\sec^{3/2}(c+dx) \sin(c+dx)}{d(b \sec(c+dx))^{3/2}}$$

input `Integrate[Sqrt[Sec[c + d*x]]/(b*Sec[c + d*x])^(3/2),x]`

output `(Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(3/2))`

**3.172.3 Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx$$

$$\downarrow \text{2031}$$

$$\frac{\sqrt{\sec(c+dx)} \int \cos(c+dx) dx}{b \sqrt{b \sec(c+dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2}) dx}{b \sqrt{b \sec(c+dx)}}$$

$$\downarrow \text{3117}$$

$$\frac{\sin(c+dx) \sqrt{\sec(c+dx)}}{bd \sqrt{b \sec(c+dx)}}$$

input `Int[Sqrt[Sec[c + d*x]]/(b*Sec[c + d*x])^(3/2),x]`

output `(Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[b*Sec[c + d*x]])`

**3.172.3.1 Defintions of rubi rules used**

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3117 Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

### 3.172.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\tan(dx+c)}{d\sqrt{\sec(dx+c)}b\sqrt{b\sec(dx+c)}}$	32
risch	$-\frac{i\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}}e^{i(dx+c)}}{2b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}d} + \frac{i\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}}e^{-i(dx+c)}}{2b\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}d}$	140

```
input int(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/d/sec(d*x+c)^(1/2)/b/(b*sec(d*x+c))^(1/2)*tan(d*x+c)
```

### 3.172.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{\sec(c+dx)}}{(b\sec(c+dx))^{3/2}} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}}\sqrt{\cos(dx+c)}\sin(dx+c)}{b^2d}$$

```
input integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fracas")
```

```
output sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b^2*d)
```

**3.172.6 Sympy [A] (verification not implemented)**

Time = 3.93 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx = \begin{cases} \frac{\tan(c+dx)\sqrt{\sec(c+dx)}}{d(b \sec(c+dx))^{3/2}} & \text{for } d \neq 0 \\ \frac{x\sqrt{\sec(c)}}{(b \sec(c))^{3/2}} & \text{otherwise} \end{cases}$$

input `integrate(sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(3/2),x)`output `Piecewise((tan(c + d*x)*sqrt(sec(c + d*x))/(d*(b*sec(c + d*x))**(3/2)), Ne(d, 0)), (x*sqrt(sec(c))/(b*sec(c))**(3/2), True))`**3.172.7 Maxima [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.37

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx = \frac{\sin(dx+c)}{b^{3/2}d}$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`output `sin(d*x + c)/(b^(3/2)*d)`**3.172.8 Giac [F]**

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx = \int \frac{\sqrt{\sec(dx+c)}}{(b \sec(dx+c))^{3/2}} dx$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`output `integrate(sqrt(sec(d*x + c))/(b*sec(d*x + c))^(3/2), x)`

**3.172.9 Mupad [B] (verification not implemented)**

Time = 13.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.11

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx = \frac{\sin(2c+2dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}}}{2b^2 d}$$

input `int((1/cos(c + d*x))^(1/2)/(b/cos(c + d*x))^(3/2),x)`

output `(sin(2*c + 2*d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2))/(2*b^2*d)`

**3.173**  $\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2}} dx$

3.173.1 Optimal result . . . . . 1033  
 3.173.2 Mathematica [A] (verified) . . . . . 1033  
 3.173.3 Rubi [A] (verified) . . . . . 1034  
 3.173.4 Maple [A] (verified) . . . . . 1035  
 3.173.5 Fricas [A] (verification not implemented) . . . . . 1036  
 3.173.6 Sympy [A] (verification not implemented) . . . . . 1036  
 3.173.7 Maxima [A] (verification not implemented) . . . . . 1037  
 3.173.8 Giac [F] . . . . . 1037  
 3.173.9 Mupad [B] (verification not implemented) . . . . . 1037

**3.173.1 Optimal result**

Integrand size = 23, antiderivative size = 69

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2}} dx = \frac{x \sqrt{\sec(c+dx)}}{2b \sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2bd \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}$$

output `1/2*sin(d*x+c)/b/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+1/2*x*sec(d*x+c)^(1/2)/b/(b*sec(d*x+c))^(1/2)`

**3.173.2 Mathematica [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2}} dx = \frac{\sec^{3/2}(c+dx)(2(c+dx) + \sin(2(c+dx)))}{4d(b \sec(c+dx))^{3/2}}$$

input `Integrate[1/(Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2)),x]`

output `(Sec[c + d*x]^(3/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*(b*Sec[c + d*x])^(3/2))`

**3.173.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.74, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2032, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2032} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^2(c+dx) dx}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^2 dx}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right)}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right)}{b\sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[1/(Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2)),x]`

output `(Sqrt[Sec[c + d*x]]*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/(b*Sqrt[b*Sec[c + d*x]])`

3.173.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2032 `Int[(Fx_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Simp[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.173.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.70

method	result
default	$\frac{\tan(dx+c)+(dx+c)\sec(dx+c)^2}{2d\sqrt{b\sec(dx+c)}\sec(dx+c)^{\frac{3}{2}}b}$
risch	$\frac{e^{i(dx+c)}x}{2b\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}(e^{2i(dx+c)+1})\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}} - \frac{ie^{3i(dx+c)}}{8b\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}(e^{2i(dx+c)+1})\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}d + \frac{ie^{-i(dx+c)}}{8b\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}(e^{2i(dx+c)+1})$

input `int(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `1/2/d/(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2)/b*(tan(d*x+c)+(d*x+c)*sec(d*x+c)^2)`



**3.173.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 165, normalized size of antiderivative = 2.39

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2}} dx = \left[ \frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) - \sqrt{-b} \log \left( 2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \right)}{4 b^2 d} \right]$$

```
input integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")
```

```
output [1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) - sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^2*d), 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b^2*d)]
```

**3.173.6 Sympy [A] (verification not implemented)**

Time = 8.20 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.55

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2}} dx = \left\{ \begin{array}{l} \frac{x \tan^2(c+dx)}{2(b \sec(c+dx))^{\frac{3}{2}} \sqrt{\sec(c+dx)}} + \frac{x}{2(b \sec(c+dx))^{\frac{3}{2}} \sqrt{\sec(c+dx)}} + \frac{\tan(c+dx)}{2d(b \sec(c+dx))^{\frac{3}{2}} \sqrt{\sec(c+dx)}} \\ \frac{x}{(b \sec(c))^{\frac{3}{2}} \sqrt{\sec(c)}} \end{array} \right.$$

```
input integrate(1/sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(3/2),x)
```

```
output Piecewise((x*tan(c + d*x)**2/(2*(b*sec(c + d*x))**(3/2)*sqrt(sec(c + d*x))) + x/(2*(b*sec(c + d*x))**(3/2)*sqrt(sec(c + d*x))) + tan(c + d*x)/(2*d*(b*sec(c + d*x))**(3/2)*sqrt(sec(c + d*x))), Ne(d, 0)), (x/((b*sec(c))**(3/2)*sqrt(sec(c))), True))
```

**3.173.7 Maxima [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.36

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{3/2}} dx = \frac{2dx + 2c + \sin(2dx + 2c)}{4b^{3/2}d}$$

input `integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`output `1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(3/2)*d)`**3.173.8 Giac [F]**

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{3/2}} dx = \int \frac{1}{(b\sec(dx+c))^{3/2} \sqrt{\sec(dx+c)}} dx$$

input `integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`output `integrate(1/((b*sec(d*x + c))^(3/2)*sqrt(sec(d*x + c))), x)`**3.173.9 Mupad [B] (verification not implemented)**

Time = 13.18 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.64

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{3/2}} dx = \frac{(\sin(2c + 2dx) + 2dx) \sqrt{\frac{b}{\cos(c+dx)}}}{4b^2 d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(1/2)),x)`output `((sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*b^2*d*(1/cos(c + d*x))^(1/2))`

**3.174**  $\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx$

3.174.1 Optimal result . . . . . 1038  
 3.174.2 Mathematica [A] (verified) . . . . . 1038  
 3.174.3 Rubi [A] (verified) . . . . . 1039  
 3.174.4 Maple [A] (verified) . . . . . 1040  
 3.174.5 Fricas [A] (verification not implemented) . . . . . 1041  
 3.174.6 Sympy [A] (verification not implemented) . . . . . 1041  
 3.174.7 Maxima [A] (verification not implemented) . . . . . 1041  
 3.174.8 Giac [F] . . . . . 1042  
 3.174.9 Mupad [B] (verification not implemented) . . . . . 1042

**3.174.1 Optimal result**

Integrand size = 23, antiderivative size = 76

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{bd\sqrt{b \sec(c+dx)}} - \frac{\sqrt{\sec(c+dx)} \sin^3(c+dx)}{3bd\sqrt{b \sec(c+dx)}}$$

output `sin(d*x+c)*sec(d*x+c)^(1/2)/b/d/(b*sec(d*x+c))^(1/2)-1/3*sin(d*x+c)^3*sec(d*x+c)^(1/2)/b/d/(b*sec(d*x+c))^(1/2)`

**3.174.2 Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx = \frac{(5 + \cos(2(c+dx))) \sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{6d(b \sec(c+dx))^{3/2}}$$

input `Integrate[1/(Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2)),x]`

output `((5 + Cos[2*(c + d*x)])*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(6*d*(b*Sec[c + d*x])^(3/2))`

**3.174.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.67, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2032, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{\frac{3}{2}}} dx \\
 & \quad \downarrow \text{2032} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^3(c+dx) dx}{b\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^3 dx}{b\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3113} \\
 & -\frac{\sqrt{\sec(c+dx)} \int (1 - \sin^2(c+dx)) d(-\sin(c+dx))}{bd\sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{(\frac{1}{3}\sin^3(c+dx) - \sin(c+dx))\sqrt{\sec(c+dx)}}{bd\sqrt{b\sec(c+dx)}}
 \end{aligned}$$

input `Int[1/(Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2)),x]`

output `-((Sqrt[Sec[c + d*x]]*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/(b*d*Sqrt[b*Sec[c + d*x]]))`

### 3.174.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2032 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

### 3.174.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.66

method	result
default	$\frac{\tan(dx+c)+2\tan(dx+c)\sec(dx+c)^2}{3d\sec(dx+c)^{\frac{5}{2}}\sqrt{b\sec(dx+c)}b}$
risch	$-\frac{ie^{4i(dx+c)}}{24b\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}d}} - \frac{3ie^{2i(dx+c)}}{8b\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}d}} + \frac{8b\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}}{8b\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}(e^{2i(dx+c)+1})}}$

input `int(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `1/3/d/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2)/b*(tan(d*x+c)+2*tan(d*x+c)*sec(d*x+c)^2)`

**3.174.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \frac{(\cos(dx+c)^3 + 2\cos(dx+c))\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{3b^2d\sqrt{\cos(dx+c)}}$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`output `1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^2*d*sqrt(cos(d*x + c)))`**3.174.6 Sympy [A] (verification not implemented)**

Time = 14.02 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.08

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \begin{cases} \frac{2\tan^3(c+dx)}{3d(b\sec(c+dx))^{\frac{3}{2}}\sec^{\frac{3}{2}}(c+dx)} + \frac{\tan(c+dx)}{d(b\sec(c+dx))^{\frac{3}{2}}\sec^{\frac{3}{2}}(c+dx)} & \text{for } d \neq 0 \\ \frac{x}{(b\sec(c))^{\frac{3}{2}}\sec^{\frac{3}{2}}(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(3/2),x)`output `Piecewise((2*tan(c + d*x)**3/(3*d*(b*sec(c + d*x))**(3/2)*sec(c + d*x)**(3/2)) + tan(c + d*x)/(d*(b*sec(c + d*x))**(3/2)*sec(c + d*x)**(3/2)), Ne(d, 0)), (x/((b*sec(c))**(3/2)*sec(c)**(3/2)), True))`**3.174.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \frac{\sin(3dx+3c) + 9\sin\left(\frac{1}{3}\arctan\left(\frac{\sin(3dx+3c)}{\cos(3dx+3c)}\right)\right)}{12b^{\frac{3}{2}}d}$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`output `1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(b^(3/2)*d)`

---

3.174.  $\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx$

**3.174.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \int \frac{1}{(b\sec(dx+c))^{\frac{3}{2}} \sec(dx+c)^{\frac{3}{2}}} dx$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate(1/((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(3/2)), x)`

**3.174.9 Mupad [B] (verification not implemented)**

Time = 13.14 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.63

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \frac{(9 \sin(c+dx) + \sin(3c+3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12b^2 d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(3/2)),x)`

output `((9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*b^2*d*(1/cos(c + d*x))^(1/2))`

**3.175** 
$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx$$

3.175.1 Optimal result . . . . . 1043  
 3.175.2 Mathematica [A] (verified) . . . . . 1043  
 3.175.3 Rubi [A] (verified) . . . . . 1044  
 3.175.4 Maple [A] (verified) . . . . . 1045  
 3.175.5 Fricas [A] (verification not implemented) . . . . . 1046  
 3.175.6 Sympy [A] (verification not implemented) . . . . . 1046  
 3.175.7 Maxima [A] (verification not implemented) . . . . . 1047  
 3.175.8 Giac [F] . . . . . 1047  
 3.175.9 Mupad [B] (verification not implemented) . . . . . 1047

**3.175.1 Optimal result**

Integrand size = 23, antiderivative size = 107

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx = \frac{3x\sqrt{\sec(c+dx)}}{8b\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{4bd \sec^{\frac{5}{2}}(c+dx)\sqrt{b \sec(c+dx)}} + \frac{3 \sin(c+dx)}{8bd\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}}$$

output `1/4*sin(d*x+c)/b/d/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2)+3/8*sin(d*x+c)/b/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+3/8*x*sec(d*x+c)^(1/2)/b/(b*sec(d*x+c))^(1/2)`

**3.175.2 Mathematica [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.51

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx = \frac{\sec^{\frac{3}{2}}(c+dx)(12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx)))}{32d(b \sec(c+dx))^{3/2}}$$

input `Integrate[1/(Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2)),x]`

output `(Sec[c + d*x]^(3/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])/(32*d*(b*Sec[c + d*x])^(3/2))`

---

3.175. 
$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx$$



**3.175.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {2032, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2032} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^4(c+dx) dx}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^4 dx}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{3}{4} \int \cos^2(c+dx) dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{3}{4} \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{3}{4} \left( \frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right)}{b\sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left( \frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right)}{b\sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[1/(Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2)),x]`

```
output (Sqrt[Sec[c + d*x]]*((Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (3*(x/2 + (Cos[
c + d*x]*Sin[c + d*x])/(2*d))))/4)/(b*Sqrt[b*Sec[c + d*x]])
```

### 3.175.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 2032 Int[(Fv_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[a^(m - 1/
2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a
, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3115 Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

### 3.175.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.63

method	result
default	$\frac{2 \tan(dx+c)+3 \tan(dx+c) \sec(dx+c)^2+3(dx+c) \sec(dx+c)^4}{8d \sec(dx+c)^{\frac{7}{2}} \sqrt{b \sec(dx+c)} b}$
risch	$\frac{3 e^{i(dx+c)} x}{8b \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (e^{2i(dx+c)}+1) \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}}} - \frac{i e^{5i(dx+c)}}{64b (e^{2i(dx+c)}+1) \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)}+1}}} d + \frac{i e^{-i(dx+c)}}{8b \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)}+1}} (e^{2i(dx+c)}+1)}$

```
input int(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/8/d/sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2)/b*(2*tan(d*x+c)+3*tan(d*x+c)*s
ec(d*x+c)^2+3*(d*x+c)*sec(d*x+c)^4)
```

---

3.175. 
$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx$$

**3.175.5 Fracas [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.94

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \left[ \frac{2 \left( 2 \cos(dx+c)^4 + 3 \cos(dx+c)^2 \right) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}} - 3 \sqrt{-b} \log \left( 2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \right) \right] / (16 b^2 d)$$

input `integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`output `[1/16*(2*(2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) - 3*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^2*d), 1/8*((2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) + 3*sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b^2*d)]`**3.175.6 Sympy [A] (verification not implemented)**

Time = 105.67 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.72

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \left\{ \begin{array}{l} \frac{3x \tan^4(c+dx)}{8(b\sec(c+dx))^{\frac{3}{2}} \sec^{\frac{5}{2}}(c+dx)} + \frac{3x \tan^2(c+dx)}{4(b\sec(c+dx))^{\frac{3}{2}} \sec^{\frac{5}{2}}(c+dx)} + \frac{3x}{8(b\sec(c+dx))^{\frac{3}{2}} \sec^{\frac{5}{2}}(c+dx)} \\ \frac{x}{(b\sec(c))^{\frac{3}{2}} \sec^{\frac{5}{2}}(c)} \end{array} \right.$$

input `integrate(1/sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(3/2),x)`output `Piecewise((3*x*tan(c + d*x)**4/(8*(b*sec(c + d*x))**(3/2)*sec(c + d*x)**(5/2)) + 3*x*tan(c + d*x)**2/(4*(b*sec(c + d*x))**(3/2)*sec(c + d*x)**(5/2)) + 3*x/(8*(b*sec(c + d*x))**(3/2)*sec(c + d*x)**(5/2)) + 3*tan(c + d*x)**3/(8*d*(b*sec(c + d*x))**(3/2)*sec(c + d*x)**(5/2)) + 5*tan(c + d*x)/(8*d*(b*sec(c + d*x))**(3/2)*sec(c + d*x)**(5/2)), Ne(d, 0)), (x/((b*sec(c))**(3/2)*sec(c)**(5/2)), True))`

**3.175.7 Maxima [A] (verification not implemented)**

Time = 0.43 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.46

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \frac{12dx + 12c + \sin(4dx + 4c) + 8\sin\left(\frac{1}{2}\arctan(\sin(4dx + 4c)), \cos(4dx + 4c)\right)}{32b^{\frac{3}{2}}d}$$

input `integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`output `1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(3/2)*d)`**3.175.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \int \frac{1}{(b\sec(dx+c))^{\frac{3}{2}}\sec(dx+c)^{\frac{5}{2}}} dx$$

input `integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`output `integrate(1/((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(5/2)), x)`**3.175.9 Mupad [B] (verification not implemented)**

Time = 13.57 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.51

$$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx = \frac{\sqrt{\frac{b}{\cos(c+dx)}}(8\sin(2c+2dx) + \sin(4c+4dx) + 12dx)}{32b^2d\sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(5/2)),x)`output `((b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))/(32*b^2*d*(1/cos(c + d*x))^(1/2))`

**3.176**  $\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.176.1 Optimal result . . . . . 1048  
 3.176.2 Mathematica [A] (verified) . . . . . 1048  
 3.176.3 Rubi [A] (verified) . . . . . 1049  
 3.176.4 Maple [A] (verified) . . . . . 1050  
 3.176.5 Fricas [A] (verification not implemented) . . . . . 1051  
 3.176.6 Sympy [F(-1)] . . . . . 1051  
 3.176.7 Maxima [B] (verification not implemented) . . . . . 1052  
 3.176.8 Giac [F] . . . . . 1052  
 3.176.9 Mupad [F(-1)] . . . . . 1053

**3.176.1 Optimal result**

Integrand size = 23, antiderivative size = 78

$$\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\sec(c+dx)}}{2b^2d\sqrt{b \sec(c+dx)}} + \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2b^2d\sqrt{b \sec(c+dx)}}$$

output `1/2*sec(d*x+c)^(5/2)*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(1/2)+1/2*arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/b^2/d/(b*sec(d*x+c))^(1/2)`

**3.176.2 Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.68

$$\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\sec(c+dx)}(\operatorname{arctanh}(\sin(c+dx)) + \sec(c+dx) \tan(c+dx))}{2b^2d\sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^(11/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(Sqrt[Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*b^2*d*Sqrt[b*Sec[c + d*x]])`

---

3.176.  $\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

**3.176.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.77, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {2031, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sec^3(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4255} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right)}{b^2 \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(11/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(Sqrt[Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/(b^2*Sqrt[b*Sec[c + d*x]])`

---

3.176.  $\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

## 3.176.3.1 Defintions of rubi rules used

rule 2031 `Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

## 3.176.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.22

method	result
default	$-\frac{\sec(dx+c)^{\frac{7}{2}} \left( \cos(dx+c)^3 \ln(-\cot(dx+c)+\csc(dx+c)-1) - \cos(dx+c)^3 \ln(\csc(dx+c)-\cot(dx+c)+1) - \cos(dx+c) \sin(dx+c) \right)}{2b^2 d \sqrt{b \sec(dx+c)}}$
risch	$-\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} \left( -e^{4i(dx+c)} \ln(e^{i(dx+c)+i}) + e^{4i(dx+c)} \ln(e^{i(dx+c)-i}) + 2ie^{3i(dx+c)} - 2ie^{i(dx+c)} - 2e^{2i(dx+c)} \ln(e^{i(dx+c)+i}) + 2e^{2i(dx+c)} \ln(e^{i(dx+c)-i}) \right)}{2b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d (e^{2i(dx+c)+1})^2}$

input `int(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `-1/2/b^2/d*sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2)*(cos(d*x+c)^3*ln(-cot(d*x+c)+csc(d*x+c)-1)-cos(d*x+c)^3*ln(csc(d*x+c)-cot(d*x+c)+1)-cos(d*x+c)*sin(d*x+c))`

---

3.176.  $\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

**3.176.5 Fracas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.63

$$\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \left[ \frac{\sqrt{b} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}}}{\sqrt{\cos(dx+c)}}}{4b^3 d \cos(dx+c)} - \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right) \cos(dx+c) - \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{2b^3 d \cos(dx+c)} \right]$$

input `integrate(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`output `[1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^3*d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^3*d*cos(d*x + c))]`**3.176.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(11/2)/(b*sec(d*x+c))**(5/2),x)`output `Timed out`



**3.176.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 688 vs.  $2(66) = 132$ .

Time = 0.43 (sec) , antiderivative size = 688, normalized size of antiderivative = 8.82

$$\int \frac{\sec^{\frac{11}{2}}(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output

```
-1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x
+ 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos
(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4
*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2
+ 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*s
in(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x
+ 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2
+ sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x +
2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^
2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4
*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c))) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*a
rctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))/(b^2*cos(4*d*x + 4*c)^2 + 4*
b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*s
in(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2
+ 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*sqrt(b)*d
```

**3.176.8 Giac [F]**

$$\int \frac{\sec^{\frac{11}{2}}(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^{\frac{11}{2}}}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(11/2)/(b*sec(d*x + c))^(5/2), x)`

---

3.176.  $\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

**3.176.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{11/2}}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int((1/cos(c + d*x))^(11/2)/(b/cos(c + d*x))^(5/2), x)`output `int((1/cos(c + d*x))^(11/2)/(b/cos(c + d*x))^(5/2), x)`

$$3.177 \quad \int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

3.177.1 Optimal result . . . . .	1054
3.177.2 Mathematica [A] (verified) . . . . .	1054
3.177.3 Rubi [A] (verified) . . . . .	1055
3.177.4 Maple [A] (verified) . . . . .	1056
3.177.5 Fricas [A] (verification not implemented) . . . . .	1057
3.177.6 Sympy [F(-1)] . . . . .	1057
3.177.7 Maxima [B] (verification not implemented) . . . . .	1057
3.177.8 Giac [F] . . . . .	1058
3.177.9 Mupad [B] (verification not implemented) . . . . .	1058

### 3.177.1 Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}}$$

output `sec(d*x+c)^(3/2)*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(1/2)`

### 3.177.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\sec^{\frac{7}{2}}(c+dx) \sin(c+dx)}{d(b \sec(c+dx))^{5/2}}$$

input `Integrate[Sec[c + d*x]^(9/2)/(b*Sec[c + d*x])^(5/2), x]`

output `(Sec[c + d*x]^(7/2)*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(5/2))`

---


$$3.177. \quad \int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

**3.177.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sec^2(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\sqrt{\sec(c+dx)} \int 1 d(-\tan(c+dx))}{b^2 d \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(9/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(Sec[c + d*x]^(3/2)*Sin[c + d*x])/(b^2*d*Sqrt[b*Sec[c + d*x]])`

## 3.177.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

## 3.177.4 Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{\cos(dx+c) \sin(dx+c) \sec(dx+c)^{\frac{5}{2}}}{d b^2 \sqrt{b \sec(dx+c)}}$	38
risch	$\frac{2i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}}{b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d (e^{2i(dx+c)+1})}$	74

input `int(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `1/d*cos(d*x+c)*sin(d*x+c)*sec(d*x+c)^(5/2)/b^2/(b*sec(d*x+c))^(1/2)`

**3.177.5 Fricas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{b^3 d \sqrt{\cos(dx+c)}}$$

input `integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))`

**3.177.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(9/2)/(b*sec(d*x+c))**(5/2),x)`

output `Timed out`

**3.177.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 67 vs.  $2(31) = 62$ .

Time = 0.40 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.91

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \frac{2\sqrt{b} \sin(2dx+2c)}{(b^3 \cos(2dx+2c)^2 + b^3 \sin(2dx+2c)^2 + 2b^3 \cos(2dx+2c) + b^3)d}$$

input `integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `2*sqrt(b)*sin(2*d*x + 2*c)/((b^3*cos(2*d*x + 2*c)^2 + b^3*sin(2*d*x + 2*c)^2 + 2*b^3*cos(2*d*x + 2*c) + b^3)*d)`

**3.177.8 Giac [F]**

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \int \frac{\sec(dx+c)^{\frac{9}{2}}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(9/2)/(b*sec(d*x + c))^(5/2), x)`

**3.177.9 Mupad [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.46

$$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \frac{(\cos(dx) - \sin(dx) \operatorname{li}) (\cos(c) - \sin(c) \operatorname{li}) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} \operatorname{li}}{b^3 d}$$

input `int((1/cos(c + d*x))^(9/2)/(b/cos(c + d*x))^(5/2),x)`

output `((cos(d*x) - sin(d*x)*1i)*(cos(c) - sin(c)*1i)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*1i)/(b^3*d)`

**3.178** 
$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

3.178.1 Optimal result . . . . . 1059  
 3.178.2 Mathematica [A] (verified) . . . . . 1059  
 3.178.3 Rubi [A] (verified) . . . . . 1060  
 3.178.4 Maple [A] (verified) . . . . . 1061  
 3.178.5 Fricas [A] (verification not implemented) . . . . . 1061  
 3.178.6 Sympy [F(-1)] . . . . . 1062  
 3.178.7 Maxima [B] (verification not implemented) . . . . . 1062  
 3.178.8 Giac [F] . . . . . 1062  
 3.178.9 Mupad [F(-1)] . . . . . 1063

**3.178.1 Optimal result**

Integrand size = 23, antiderivative size = 36

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}}$$

output `arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/b^2/d/(b*sec(d*x+c))^(1/2)`

**3.178.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{d(b \sec(c+dx))^{5/2}}$$

input `Integrate[Sec[c + d*x]^(7/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(ArcTanh[Sin[c + d*x]]*Sec[c + d*x]^(5/2))/(d*(b*Sec[c + d*x])^(5/2))`



**3.178.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

↓ 2031

$$\frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}}$$

↓ 3042

$$\frac{\sqrt{\sec(c+dx)} \int \csc(c+dx + \frac{\pi}{2}) dx}{b^2 \sqrt{b \sec(c+dx)}}$$

↓ 4257

$$\frac{\sqrt{\sec(c+dx)} \operatorname{arctanh}(\sin(c+dx))}{b^2 d \sqrt{b \sec(c+dx)}}$$

input `Int[Sec[c + d*x]^(7/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(b^2*d*Sqrt[b*Sec[c + d*x]])`

**3.178.3.1 Defintions of rubi rules used**

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

### 3.178.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.36

method	result	size
default	$-\frac{2 \cos(dx+c) \sec(dx+c)^{\frac{3}{2}} \operatorname{arctanh}(\cot(dx+c) - \csc(dx+c))}{d b^2 \sqrt{b \sec(dx+c)}}$	49
risch	$-\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)} - i)}}{b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}} \ln(e^{i(dx+c)} + i)}}{b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}$	145

input `int(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$-2/d*\cos(d*x+c)*\sec(d*x+c)^(3/2)*\operatorname{arctanh}(\cot(d*x+c)-\csc(d*x+c))/b^2/(b*\sec(d*x+c))^(1/2)$$

### 3.178.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.17

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \left[ \frac{\log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2b^{\frac{5}{2}}d}, \right. \\ \left. -\frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{b^3 d} \right]$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output 
$$[1/2*\log(-(b*\cos(d*x+c))^2 - 2*\sqrt{b}*\sqrt{b/\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c) - 2*b)/\cos(d*x+c)^2/(b^(5/2)*d), -\sqrt{-b}*\arctan(\sqrt{-b}*\sqrt{b/\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)/b)/(b^3*d)]$$

---

3.178. 
$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$$

**3.178.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(7/2)/(b*sec(d*x+c))**(5/2),x)`output `Timed out`**3.178.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(32) = 64.

Time = 0.40 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.81

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 + 1)}{2 b^{\frac{5}{2}} d}$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`output `1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(b^(5/2)*d)`**3.178.8 Giac [F]**

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \int \frac{\sec(dx+c)^{\frac{7}{2}}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`output `integrate(sec(d*x + c)^(7/2)/(b*sec(d*x + c))^(5/2), x)`

**3.178.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{\frac{7}{2}}}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{5}{2}}} dx$$

input `int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(5/2),x)`output `int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(5/2), x)`

$$3.179 \quad \int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

3.179.1 Optimal result . . . . .	1064
3.179.2 Mathematica [A] (verified) . . . . .	1064
3.179.3 Rubi [A] (verified) . . . . .	1065
3.179.4 Maple [A] (verified) . . . . .	1066
3.179.5 Fricas [A] (verification not implemented) . . . . .	1066
3.179.6 Sympy [A] (verification not implemented) . . . . .	1066
3.179.7 Maxima [A] (verification not implemented) . . . . .	1067
3.179.8 Giac [F] . . . . .	1067
3.179.9 Mupad [B] (verification not implemented) . . . . .	1067

### 3.179.1 Optimal result

Integrand size = 23, antiderivative size = 27

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{x \sqrt{\sec(c+dx)}}{b^2 \sqrt{b \sec(c+dx)}}$$

output `x*sec(d*x+c)^(1/2)/b^2/(b*sec(d*x+c))^(1/2)`

### 3.179.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{x \sqrt{b \sec(c+dx)}}{b^3 \sqrt{\sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^(5/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(x*sqrt[b*Sec[c + d*x]])/(b^3*sqrt[Sec[c + d*x]])`

---


$$3.179. \quad \int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

**3.179.3 Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {2031, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$$

↓ 2031

$$\frac{\sqrt{\sec(c+dx)} \int 1 dx}{b^2 \sqrt{b \sec(c+dx)}}$$

↓ 24

$$\frac{x \sqrt{\sec(c+dx)}}{b^2 \sqrt{b \sec(c+dx)}}$$

input `Int[Sec[c + d*x]^(5/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(x*Sqrt[Sec[c + d*x]])/(b^2*Sqrt[b*Sec[c + d*x]])`

**3.179.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fv_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fv, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

**3.179.4 Maple [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

method	result	size
default	$\frac{(dx+c)\sqrt{\sec(dx+c)}}{db^2\sqrt{b\sec(dx+c)}}$	31
risch	$\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}x}}{b^2\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}}$	57

input `int(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`output `1/d*(d*x+c)*sec(d*x+c)^(1/2)/b^2/(b*sec(d*x+c))^(1/2)`**3.179.5 Fracas [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 101, normalized size of antiderivative = 3.74

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}} dx = \left[ -\frac{\sqrt{-b} \log\left(2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b\right)}{2b^3d}, \right]$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fracas")`output `[-1/2*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/(b^3*d), arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))/(b^(5/2)*d)]`**3.179.6 Sympy [A] (verification not implemented)**

Time = 49.36 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}} dx = \frac{x \sec^{\frac{5}{2}}(c+dx)}{(b\sec(c+dx))^{\frac{5}{2}}}$$

input `integrate(sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(5/2),x)`

output `x*sec(c + d*x)**(5/2)/(b*sec(c + d*x))**(5/2)`

### 3.179.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx = \frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{5}{2}}d}$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(b^(5/2)*d)`

### 3.179.8 Giac [F]

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx = \int \frac{\sec(dx + c)^{\frac{5}{2}}}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(sec(d*x + c)^(5/2)/(b*sec(d*x + c))^(5/2), x)`

### 3.179.9 Mupad [B] (verification not implemented)

Time = 13.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx = \frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{b^3 \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((1/cos(c + d*x))^(5/2)/(b/cos(c + d*x))^(5/2),x)`

output `(x*(b/cos(c + d*x))^(1/2))/(b^3*(1/cos(c + d*x))^(1/2))`

---

3.179.  $\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$



**3.180** 
$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

3.180.1 Optimal result . . . . . 1068  
 3.180.2 Mathematica [A] (verified) . . . . . 1068  
 3.180.3 Rubi [A] (verified) . . . . . 1069  
 3.180.4 Maple [A] (verified) . . . . . 1070  
 3.180.5 Fricas [A] (verification not implemented) . . . . . 1070  
 3.180.6 Sympy [A] (verification not implemented) . . . . . 1071  
 3.180.7 Maxima [A] (verification not implemented) . . . . . 1071  
 3.180.8 Giac [F] . . . . . 1071  
 3.180.9 Mupad [B] (verification not implemented) . . . . . 1072

**3.180.1 Optimal result**

Integrand size = 23, antiderivative size = 35

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}}$$

output `sin(d*x+c)*sec(d*x+c)^(1/2)/b^2/d/(b*sec(d*x+c))^(1/2)`

**3.180.2 Mathematica [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^(3/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(b^2*d*Sqrt[b*Sec[c + d*x]])`

**3.180.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {2031, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

↓ 2031

$$\frac{\sqrt{\sec(c+dx)} \int \cos(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}}$$

↓ 3042

$$\frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2}) dx}{b^2 \sqrt{b \sec(c+dx)}}$$

↓ 3117

$$\frac{\sin(c+dx) \sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}}$$

input `Int[Sec[c + d*x]^(3/2)/(b*Sec[c + d*x])^(5/2),x]`

output `(Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(b^2*d*Sqrt[b*Sec[c + d*x]])`

**3.180.3.1 Defintions of rubi rules used**

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3117 Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

### 3.180.4 Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\tan(dx+c)}{db^2 \sqrt{\sec(dx+c)} \sqrt{b \sec(dx+c)}}$	32
risch	$-\frac{i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{i(dx+c)}}{2b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d} + \frac{i \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} e^{-i(dx+c)}}{2b^2 \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}} d}$	140

```
input int(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/d/b^2/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)*tan(d*x+c)
```

### 3.180.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b^3 d}$$

```
input integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b^3*d)
```

**3.180.6 Sympy [A] (verification not implemented)**

Time = 18.69 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.31

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \begin{cases} \frac{\tan(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(b \sec(c+dx))^{\frac{5}{2}}} & \text{for } d \neq 0 \\ \frac{x \sec^{\frac{3}{2}}(c)}{(b \sec(c))^{\frac{5}{2}}} & \text{otherwise} \end{cases}$$

input `integrate(sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(5/2),x)`output `Piecewise((tan(c + d*x)*sec(c + d*x)**(3/2)/(d*(b*sec(c + d*x))**(5/2)), N  
e(d, 0)), (x*sec(c)**(3/2)/(b*sec(c))**(5/2), True))`**3.180.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.37

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \frac{\sin(dx+c)}{b^{\frac{5}{2}}d}$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`output `sin(d*x + c)/(b^(5/2)*d)`**3.180.8 Giac [F]**

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \int \frac{\sec(dx+c)^{\frac{3}{2}}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

input `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`output `integrate(sec(d*x + c)^(3/2)/(b*sec(d*x + c))^(5/2), x)`

**3.180.9 Mupad [B] (verification not implemented)**

Time = 13.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.11

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx = \frac{\sin(2c+2dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}}}{2b^3 d}$$

input `int((1/cos(c + d*x))^(3/2)/(b/cos(c + d*x))^(5/2),x)`output `(sin(2*c + 2*d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2))/(2*b^3*d)`

**3.181** 
$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx$$

3.181.1 Optimal result . . . . . 1073  
 3.181.2 Mathematica [A] (verified) . . . . . 1073  
 3.181.3 Rubi [A] (verified) . . . . . 1074  
 3.181.4 Maple [A] (verified) . . . . . 1075  
 3.181.5 Fricas [A] (verification not implemented) . . . . . 1076  
 3.181.6 Sympy [A] (verification not implemented) . . . . . 1076  
 3.181.7 Maxima [A] (verification not implemented) . . . . . 1077  
 3.181.8 Giac [F] . . . . . 1077  
 3.181.9 Mupad [B] (verification not implemented) . . . . . 1077

**3.181.1 Optimal result**

Integrand size = 23, antiderivative size = 69

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx = \frac{x \sqrt{\sec(c+dx)}}{2b^2 \sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2b^2 d \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}$$

output `1/2*sin(d*x+c)/b^2/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+1/2*x*sec(d*x+c)^(1/2)/b^2/(b*sec(d*x+c))^(1/2)`

**3.181.2 Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\sec(c+dx)}(2(c+dx) + \sin(2(c+dx)))}{4b^2 d \sqrt{b \sec(c+dx)}}$$

input `Integrate[Sqrt[Sec[c + d*x]]/(b*Sec[c + d*x])^(5/2),x]`

output `(Sqrt[Sec[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*b^2*d*Sqrt[b*Sec[c + d*x]])`

**3.181.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.74, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2031, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2031} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^2(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^2 dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right)}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right)}{b^2 \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sqrt[Sec[c + d*x]]/(b*Sec[c + d*x])^(5/2),x]`

output `(Sqrt[Sec[c + d*x]]*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/(b^2*Sqrt[b*Sec[c + d*x]])`

## 3.181.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2031 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

## 3.181.4 Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\tan(dx+c)+(dx+c)\sec(dx+c)^2}{2d\sqrt{b\sec(dx+c)}\sec(dx+c)^{\frac{3}{2}}b^2}$	48
risch	$\frac{\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}x}{2b^2\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{i\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}e^{2i(dx+c)}}}{8b^2\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}d}} + \frac{i\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}e^{-2i(dx+c)}}}{8b^2\sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}d}}$	197

input `int(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `1/2/d/(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2)/b^2*(tan(d*x+c)+(d*x+c)*sec(d*x+c)^2)`



**3.181.5 Fricas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 165, normalized size of antiderivative = 2.39

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx = \left[ \frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) - \sqrt{-b} \log \left( 2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \right)}{4 b^3 d} \right]$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`output `[1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) - sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^3*d), 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b^3*d)]`**3.181.6 Sympy [A] (verification not implemented)**

Time = 10.10 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.55

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx = \begin{cases} \frac{x \tan^2(c+dx) \sqrt{\sec(c+dx)}}{2(b \sec(c+dx))^{\frac{5}{2}}} + \frac{x \sqrt{\sec(c+dx)}}{2(b \sec(c+dx))^{\frac{5}{2}}} + \frac{\tan(c+dx) \sqrt{\sec(c+dx)}}{2d(b \sec(c+dx))^{\frac{5}{2}}} & \text{for } d \neq 0 \\ \frac{x \sqrt{\sec(c)}}{(b \sec(c))^{\frac{5}{2}}} & \text{otherwise} \end{cases}$$

input `integrate(sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(5/2),x)`output `Piecewise((x*tan(c + d*x)**2*sqrt(sec(c + d*x))/(2*(b*sec(c + d*x))**(5/2)) + x*sqrt(sec(c + d*x))/(2*(b*sec(c + d*x))**(5/2)) + tan(c + d*x)*sqrt(sec(c + d*x))/(2*d*(b*sec(c + d*x))**(5/2)), Ne(d, 0)), (x*sqrt(sec(c))/(b*sec(c))**(5/2), True))`

**3.181.7 Maxima [A] (verification not implemented)**

Time = 0.56 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.36

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx = \frac{2dx + 2c + \sin(2dx + 2c)}{4b^{5/2}d}$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`output `1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(5/2)*d)`**3.181.8 Giac [F]**

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx = \int \frac{\sqrt{\sec(dx+c)}}{(b \sec(dx+c))^{5/2}} dx$$

input `integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`output `integrate(sqrt(sec(d*x + c))/(b*sec(d*x + c))^(5/2), x)`**3.181.9 Mupad [B] (verification not implemented)**

Time = 12.99 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\frac{b}{\cos(c+dx)}} (\sin(c+dx) + \sin(3c+3dx) + 4dx \cos(c+dx))}{8b^3 d \cos(c+dx) \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int((1/cos(c + d*x))^(1/2)/(b/cos(c + d*x))^(5/2),x)`output `((b/cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(8*b^3*d*cos(c + d*x)*(1/cos(c + d*x))^(1/2))`

**3.182**  $\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{5/2}} dx$

3.182.1 Optimal result . . . . . 1078  
 3.182.2 Mathematica [A] (verified) . . . . . 1078  
 3.182.3 Rubi [A] (verified) . . . . . 1079  
 3.182.4 Maple [A] (verified) . . . . . 1080  
 3.182.5 Fricas [A] (verification not implemented) . . . . . 1081  
 3.182.6 Sympy [A] (verification not implemented) . . . . . 1081  
 3.182.7 Maxima [A] (verification not implemented) . . . . . 1081  
 3.182.8 Giac [F] . . . . . 1082  
 3.182.9 Mupad [B] (verification not implemented) . . . . . 1082

**3.182.1 Optimal result**

Integrand size = 23, antiderivative size = 76

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}} - \frac{\sqrt{\sec(c+dx)} \sin^3(c+dx)}{3b^2 d \sqrt{b \sec(c+dx)}}$$

output `sin(d*x+c)*sec(d*x+c)^(1/2)/b^2/d/(b*sec(d*x+c))^(1/2)-1/3*sin(d*x+c)^3*sec(d*x+c)^(1/2)/b^2/d/(b*sec(d*x+c))^(1/2)`

**3.182.2 Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.63

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{5/2}} dx = \frac{(5 + \cos(2(c+dx)))\sqrt{\sec(c+dx)} \sin(c+dx)}{6b^2 d \sqrt{b \sec(c+dx)}}$$

input `Integrate[1/(Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(5/2)),x]`

output `((5 + Cos[2*(c + d*x)])*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(6*b^2*d*Sqrt[b*Sec[c + d*x]])`

**3.182.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.67, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {2032, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2032} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^3(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^3 dx}{b^2 \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3113} \\
 & - \frac{\sqrt{\sec(c+dx)} \int (1 - \sin^2(c+dx)) d(-\sin(c+dx))}{b^2 d \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{2009} \\
 & - \frac{(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx)) \sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[1/(Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(5/2)),x]`

output `-((Sqrt[Sec[c + d*x]]*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/(b^2*d*Sqrt[b*Sec[c + d*x]]))`

## 3.182.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2032 `Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

## 3.182.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.66

method	result
default	$\frac{\tan(dx+c)+2\tan(dx+c)\sec(dx+c)^2}{3d\sqrt{b\sec(dx+c)}\sec(dx+c)^{\frac{5}{2}}b^2}$
risch	$-\frac{ie^{4i(dx+c)}}{24b^2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}(e^{2i(dx+c)+1})d} - \frac{3ie^{2i(dx+c)}}{8b^2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}(e^{2i(dx+c)+1})d} + \frac{1}{8b^2\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)+1}}}(e^{2i(dx+c)+1})d}}$

input `int(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `1/3/d/(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2)/b^2*(tan(d*x+c)+2*tan(d*x+c)*sec(d*x+c)^2)`

**3.182.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{5/2}} dx = \frac{(\cos(dx+c)^3 + 2\cos(dx+c))\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{3b^3d\sqrt{\cos(dx+c)}}$$

input `integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`output `1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))`**3.182.6 Sympy [A] (verification not implemented)**

Time = 24.40 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.08

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{5/2}} dx = \begin{cases} \frac{2\tan^3(c+dx)}{3d(b\sec(c+dx))^{\frac{5}{2}}\sqrt{\sec(c+dx)}} + \frac{\tan(c+dx)}{d(b\sec(c+dx))^{\frac{5}{2}}\sqrt{\sec(c+dx)}} & \text{for } d \neq 0 \\ \frac{x}{(b\sec(c))^{\frac{5}{2}}\sqrt{\sec(c)}} & \text{otherwise} \end{cases}$$

input `integrate(1/sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(5/2),x)`output `Piecewise((2*tan(c + d*x)**3/(3*d*(b*sec(c + d*x))**(5/2)*sqrt(sec(c + d*x))) + tan(c + d*x)/(d*(b*sec(c + d*x))**(5/2)*sqrt(sec(c + d*x))), Ne(d, 0)), (x/((b*sec(c))**(5/2)*sqrt(sec(c))), True))`**3.182.7 Maxima [A] (verification not implemented)**

Time = 0.54 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{5/2}} dx = \frac{\sin(3dx+3c) + 9\sin\left(\frac{1}{3}\arctan\left(\frac{\sin(3dx+3c)}{\cos(3dx+3c)}\right)\right)}{12b^{\frac{5}{2}}d}$$

input `integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`output `1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(b^(5/2)*d)`

**3.182.8 Giac [F]**

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{5/2}} dx = \int \frac{1}{(b\sec(dx+c))^{\frac{5}{2}} \sqrt{\sec(dx+c)}} dx$$

input `integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate(1/((b*sec(d*x + c))^(5/2)*sqrt(sec(d*x + c))), x)`

**3.182.9 Mupad [B] (verification not implemented)**

Time = 13.78 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.63

$$\int \frac{1}{\sqrt{\sec(c+dx)}(b\sec(c+dx))^{5/2}} dx = \frac{(9 \sin(c+dx) + \sin(3c+3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12b^3 d \sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(1/2)),x)`

output `((9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*b^3*d*(1/cos(c + d*x))^(1/2))`

**3.183** 
$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2}} dx$$

3.183.1 Optimal result . . . . . 1083  
 3.183.2 Mathematica [A] (verified) . . . . . 1083  
 3.183.3 Rubi [A] (verified) . . . . . 1084  
 3.183.4 Maple [A] (verified) . . . . . 1085  
 3.183.5 Fricas [A] (verification not implemented) . . . . . 1086  
 3.183.6 Sympy [A] (verification not implemented) . . . . . 1086  
 3.183.7 Maxima [A] (verification not implemented) . . . . . 1087  
 3.183.8 Giac [F] . . . . . 1087  
 3.183.9 Mupad [B] (verification not implemented) . . . . . 1087

**3.183.1 Optimal result**

Integrand size = 23, antiderivative size = 107

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2}} dx = \frac{3x \sqrt{\sec(c+dx)}}{8b^2 \sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}} + \frac{3 \sin(c+dx)}{8b^2 d \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}$$

output `1/4*sin(d*x+c)/b^2/d/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2)+3/8*sin(d*x+c)/b^2/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+3/8*x*sec(d*x+c)^(1/2)/b^2/(b*sec(d*x+c))^(1/2)`

**3.183.2 Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.54

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2}} dx = \frac{\sqrt{\sec(c+dx)}(12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx)))}{32b^2 d \sqrt{b \sec(c+dx)}}$$

input `Integrate[1/(Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(5/2)),x]`

output `(Sqrt[Sec[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])/(32*b^2*d*Sqrt[b*Sec[c + d*x]])`

---

3.183. 
$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2}} dx$$



**3.183.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {2032, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2032} \\
 & \frac{\sqrt{\sec(c+dx)} \int \cos^4(c+dx) dx}{b^2 \sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \int \sin(c+dx + \frac{\pi}{2})^4 dx}{b^2 \sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{3}{4} \int \cos^2(c+dx) dx + \frac{\sin(c+dx)\cos^3(c+dx)}{4d} \right)}{b^2 \sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{3}{4} \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{\sin(c+dx)\cos^3(c+dx)}{4d} \right)}{b^2 \sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{3}{4} \left( \frac{\int 1 dx}{2} + \frac{\sin(c+dx)\cos(c+dx)}{2d} \right) + \frac{\sin(c+dx)\cos^3(c+dx)}{4d} \right)}{b^2 \sqrt{b\sec(c+dx)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sqrt{\sec(c+dx)} \left( \frac{\sin(c+dx)\cos^3(c+dx)}{4d} + \frac{3}{4} \left( \frac{\sin(c+dx)\cos(c+dx)}{2d} + \frac{x}{2} \right) \right)}{b^2 \sqrt{b\sec(c+dx)}}
 \end{aligned}$$

input `Int[1/(Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(5/2)),x]`

output  $(\text{Sqrt}[\text{Sec}[c + d*x]]*((\text{Cos}[c + d*x]^3*\text{Sin}[c + d*x])/(4*d) + (3*(x/2 + (\text{Cos}[c + d*x]*\text{Sin}[c + d*x])/(2*d))))/4)/(b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

### 3.183.3.1 Defintions of rubi rules used

rule 24  $\text{Int}[a_, x\_Symbol] \text{ :> } \text{Simp}[a*x, x] \text{ /; } \text{FreeQ}[a, x]$

rule 2032  $\text{Int}[(\text{Fx}_.)*((a_.)*(v_.))^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x\_Symbol] \text{ :> } \text{Simp}[a^{(m - 1/2)}*b^{(n + 1/2)}*(\text{Sqrt}[a*v]/\text{Sqrt}[b*v]) \text{ Int}[v^{(m + n)}*\text{Fx}, x], x] \text{ /; } \text{FreeQ}[\{a, b, m\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n - 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

rule 3042  $\text{Int}[u_, x\_Symbol] \text{ :> } \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; } \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3115  $\text{Int}[(b_.)*\text{sin}[c_. + (d_.)*(x_.)]^{(n_.)}, x\_Symbol] \text{ :> } \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n - 1)})/(d*n), x] + \text{Simp}[b^2*((n - 1)/n) \text{ Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] \text{ /; } \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

### 3.183.4 Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.63

method	result
default	$\frac{2 \tan(dx+c)+3 \tan(dx+c) \sec(dx+c)^2+3(dx+c) \sec(dx+c)^4}{8d \sec(dx+c)^{\frac{7}{2}} \sqrt{b \sec(dx+c)} b^2}$
risch	$\frac{3 e^{i(dx+c)} x}{8b^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} (e^{2i(dx+c)+1}) \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{i e^{5i(dx+c)}}{64b^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} (e^{2i(dx+c)+1}) \sqrt{\frac{b e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} d + \frac{i e^{-i(dx+c)}}{8b^2 \sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}} (e^{2i(dx+c)+1})}$

input  $\text{int}(1/\text{sec}(d*x+c)^{(3/2)}/(b*\text{sec}(d*x+c))^{(5/2)}, x, \text{method}=\_RETURNVERBOSE)$

output  $1/8/d/\text{sec}(d*x+c)^{(7/2)}/(b*\text{sec}(d*x+c))^{(1/2)}/b^2*(2*\tan(d*x+c)+3*\tan(d*x+c)*\text{sec}(d*x+c)^2+3*(d*x+c)*\text{sec}(d*x+c)^4)$

---

3.183.  $\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2}} dx$

**3.183.5 Fricas [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.94

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{5/2}} dx = \left[ \frac{2 \left( 2 \cos(dx+c)^4 + 3 \cos(dx+c)^2 \right) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}} - 3 \sqrt{-b} \log \left( 2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \right) \right] / (16 b^3 d)$$

```
input integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output [1/16*(2*(2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) - 3*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^3*d), 1/8*((2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) + 3*sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b^3*d)]
```

**3.183.6 Sympy [A] (verification not implemented)**

Time = 112.02 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.72

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{5/2}} dx = \left\{ \begin{array}{l} \frac{3x \tan^4(c+dx)}{8(b\sec(c+dx))^{\frac{5}{2}} \sec^{\frac{3}{2}}(c+dx)} + \frac{3x \tan^2(c+dx)}{4(b\sec(c+dx))^{\frac{5}{2}} \sec^{\frac{3}{2}}(c+dx)} + \frac{3x}{8(b\sec(c+dx))^{\frac{5}{2}} \sec^{\frac{3}{2}}(c+dx)} \\ \frac{x}{(b\sec(c))^{\frac{5}{2}} \sec^{\frac{3}{2}}(c)} \end{array} \right.$$

```
input integrate(1/sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(5/2),x)
```

```
output Piecewise((3*x*tan(c + d*x)**4/(8*(b*sec(c + d*x))**(5/2)*sec(c + d*x)**(3/2)) + 3*x*tan(c + d*x)**2/(4*(b*sec(c + d*x))**(5/2)*sec(c + d*x)**(3/2)) + 3*x/(8*(b*sec(c + d*x))**(5/2)*sec(c + d*x)**(3/2)) + 3*tan(c + d*x)**3/(8*d*(b*sec(c + d*x))**(5/2)*sec(c + d*x)**(3/2)) + 5*tan(c + d*x)/(8*d*(b*sec(c + d*x))**(5/2)*sec(c + d*x)**(3/2)), Ne(d, 0)), (x/((b*sec(c))**(5/2)*sec(c)**(3/2)), True))
```

**3.183.7 Maxima [A] (verification not implemented)**

Time = 0.54 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.46

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{5/2}} dx = \frac{12dx + 12c + \sin(4dx + 4c) + 8\sin\left(\frac{1}{2}\arctan(\sin(4dx + 4c)), \cos(4dx + 4c)\right)}{32b^{\frac{5}{2}}d}$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`output `1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(5/2)*d)`**3.183.8 Giac [F]**

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{5/2}} dx = \int \frac{1}{(b\sec(dx+c))^{\frac{5}{2}}\sec(dx+c)^{\frac{3}{2}}} dx$$

input `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`output `integrate(1/((b*sec(d*x + c))^(5/2)*sec(d*x + c)^(3/2)), x)`**3.183.9 Mupad [B] (verification not implemented)**

Time = 13.36 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.51

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b\sec(c+dx))^{5/2}} dx = \frac{\sqrt{\frac{b}{\cos(c+dx)}}(8\sin(2c+2dx) + \sin(4c+4dx) + 12dx)}{32b^3d\sqrt{\frac{1}{\cos(c+dx)}}}$$

input `int(1/((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(3/2)),x)`output `((b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))/(32*b^3*d*(1/cos(c + d*x))^(1/2))`

### 3.184 $\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

3.184.1 Optimal result . . . . .	1088
3.184.2 Mathematica [A] (verified) . . . . .	1088
3.184.3 Rubi [A] (verified) . . . . .	1089
3.184.4 Maple [F] . . . . .	1090
3.184.5 Fracas [F] . . . . .	1090
3.184.6 Sympy [F] . . . . .	1091
3.184.7 Maxima [F] . . . . .	1091
3.184.8 Giac [F] . . . . .	1091
3.184.9 Mupad [F(-1)] . . . . .	1092

#### 3.184.1 Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) (b \sec(c + dx))^{4/3} \sin(c + dx)}{4bd \sqrt{\sin^2(c + dx)}}$$

output `3/4*hypergeom([-2/3, 1/2], [1/3], cos(d*x+c)^2)*(b*sec(d*x+c))^(4/3)*sin(d*x+c)/b/d/(sin(d*x+c)^2)^(1/2)`

#### 3.184.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= \frac{3 \csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \sec^2(c + dx)\right) (b \sec(c + dx))^{4/3} \sqrt{-\tan^2(c + dx)}}{7bd}$$

input `Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(1/3),x]`

output `(3*Csc[c + d*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/(7*b*d)`

**3.184.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx) \sqrt[3]{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c+dx))^{7/3} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{7/3} dx}{b^2} \\
 & \quad \downarrow \text{4259} \\
 & \quad \frac{\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{7/3}} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{7/3}} dx}{b^2} \\
 & \quad \downarrow \text{3122} \\
 & \quad \frac{3 \sin(c+dx) (b \sec(c+dx))^{4/3} \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c+dx)\right)}{4bd \sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(1/3),x]`

output `(3*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sin[c + d*x])/(4*b*d*Sqrt[Sin[c + d*x]^2])`

## 3.184.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.184.4 Maple [F]

$$\int \sec(dx + c)^2 (b \sec(dx + c))^{\frac{1}{3}} dx$$

input `int(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3),x)`

output `int(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3),x)`

## 3.184.5 Fracas [F]

$$\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x + c))^(1/3)*sec(d*x + c)^2, x)`

**3.184.6 Sympy [F]**

$$\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \sqrt[3]{b \sec(c + dx)} \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(1/3),x)`

output `Integral((b*sec(c + d*x))**(1/3)*sec(c + d*x)**2, x)`

**3.184.7 Maxima [F]**

$$\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^2, x)`

**3.184.8 Giac [F]**

$$\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^2, x)`



**3.184.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}}{\cos(c + dx)^2} dx$$

input `int((b/cos(c + d*x))^(1/3)/cos(c + d*x)^2,x)`output `int((b/cos(c + d*x))^(1/3)/cos(c + d*x)^2, x)`

### 3.185 $\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

3.185.1 Optimal result . . . . .	1093
3.185.2 Mathematica [A] (verified) . . . . .	1093
3.185.3 Rubi [A] (verified) . . . . .	1094
3.185.4 Maple [F] . . . . .	1095
3.185.5 Fracas [F] . . . . .	1095
3.185.6 Sympy [F] . . . . .	1096
3.185.7 Maxima [F] . . . . .	1096
3.185.8 Giac [F] . . . . .	1096
3.185.9 Mupad [F(-1)] . . . . .	1097

#### 3.185.1 Optimal result

Integrand size = 19, antiderivative size = 53

$$\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sin^2(c + dx)}}$$

output `3*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*(b*sec(d*x+c))^(1/3)*sin(d*x+c)/d/(sin(d*x+c)^2)^(1/2)`

#### 3.185.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.13

$$\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= \frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \sec^2(c + dx)\right) (b \sec(c + dx))^{4/3} \sqrt{-\tan^2(c + dx)}}{4bd}$$

input `Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(1/3),x]`

output `(3*Cot[c + d*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/(4*b*d)`

**3.185.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c + dx))^{4/3} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{4/3} dx}{b} \\
 & \quad \downarrow \text{4259} \\
 & \frac{\sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\left(\frac{\cos(c + dx)}{b}\right)^{4/3}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\left(\frac{\sin(c + dx + \frac{\pi}{2})}{b}\right)^{4/3}} dx}{b} \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(c + dx) \sqrt[3]{b \sec(c + dx)} \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]*(b*Sec[c + d*x])^(1/3),x]`

output `(3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sin[c + d*x])/(d*Sqrt[Sin[c + d*x]^2])`

## 3.185.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.185.4 Maple [F]

$$\int \sec(dx+c) (b \sec(dx+c))^{\frac{1}{3}} dx$$

input `int(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

output `int(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

## 3.185.5 Fracas [F]

$$\int \sec(c+dx) \sqrt[3]{b \sec(c+dx)} dx = \int (b \sec(dx+c))^{\frac{1}{3}} \sec(dx+c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^(1/3)*sec(d*x+c), x)`

**3.185.6 Sympy [F]**

$$\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \sqrt[3]{b \sec(c + dx)} \sec(c + dx) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))**(1/3),x)`

output `Integral((b*sec(c + d*x))**(1/3)*sec(c + d*x), x)`

**3.185.7 Maxima [F]**

$$\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c), x)`

**3.185.8 Giac [F]**

$$\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c), x)`

**3.185.9 Mupad [F(-1)]**

Timed out.

$$\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}}{\cos(c + dx)} dx$$

input `int((b/cos(c + d*x))^(1/3)/cos(c + d*x), x)`output `int((b/cos(c + d*x))^(1/3)/cos(c + d*x), x)`

### 3.186 $\int \sqrt[3]{b \sec(c + dx)} dx$

3.186.1 Optimal result . . . . .	1098
3.186.2 Mathematica [A] (verified) . . . . .	1098
3.186.3 Rubi [A] (verified) . . . . .	1099
3.186.4 Maple [F] . . . . .	1100
3.186.5 Fricas [F] . . . . .	1100
3.186.6 Sympy [F] . . . . .	1101
3.186.7 Maxima [F] . . . . .	1101
3.186.8 Giac [F] . . . . .	1101
3.186.9 Mupad [F(-1)] . . . . .	1102

#### 3.186.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \sqrt[3]{b \sec(c + dx)} dx = -\frac{3b \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \sec(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}}$$

output `-3/2*b*hypergeom([1/3, 1/2],[4/3],cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(2/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.186.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98

$$\int \sqrt[3]{b \sec(c + dx)} dx = \frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \sec^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sqrt{-\tan^2(c + dx)}}{d}$$

input `Integrate[(b*Sec[c + d*x])^(1/3),x]`

output `(3*Cot[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sqrt[-Tan[c + d*x]^2])/d`

**3.186.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{b \sec(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt[3]{b \csc\left(c + dx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\sqrt[3]{\frac{\cos(c + dx)}{b}}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\sqrt[3]{\frac{\sin\left(c + dx + \frac{\pi}{2}\right)}{b}}} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3b \sin(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right)}{2d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{2/3}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[c + d*x]^2]*Sin[c + d*x])/(2*d*(b*Sec[c + d*x])^(2/3)*Sqrt[Sin[c + d*x]^2])`



## 3.186.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.186.4 Maple [F]

$$\int (b \sec(dx + c))^{\frac{1}{3}} dx$$

input `int((b*sec(d*x+c))^(1/3),x)`

output `int((b*sec(d*x+c))^(1/3),x)`

## 3.186.5 Fracas [F]

$$\int \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} dx$$

input `integrate((b*sec(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(1/3), x)`

**3.186.6 Sympy [F]**

$$\int \sqrt[3]{b \sec(c + dx)} dx = \int \sqrt[3]{b \sec(c + dx)} dx$$

input `integrate((b*sec(d*x+c))**(1/3),x)`

output `Integral((b*sec(c + d*x))**(1/3), x)`

**3.186.7 Maxima [F]**

$$\int \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} dx$$

input `integrate((b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(1/3), x)`

**3.186.8 Giac [F]**

$$\int \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} dx$$

input `integrate((b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(1/3), x)`

**3.186.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt[3]{b \sec(c + dx)} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{1/3} dx$$

input `int((b/cos(c + d*x))^(1/3),x)`output `int((b/cos(c + d*x))^(1/3), x)`

### 3.187 $\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

3.187.1 Optimal result . . . . .	1103
3.187.2 Mathematica [A] (verified) . . . . .	1103
3.187.3 Rubi [A] (verified) . . . . .	1104
3.187.4 Maple [F] . . . . .	1105
3.187.5 Fricas [F] . . . . .	1105
3.187.6 Sympy [F] . . . . .	1106
3.187.7 Maxima [F] . . . . .	1106
3.187.8 Giac [F] . . . . .	1106
3.187.9 Mupad [F(-1)] . . . . .	1107

#### 3.187.1 Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= -\frac{3b^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5d(b \sec(c + dx))^{5/3} \sqrt{\sin^2(c + dx)}}$$

output `-3/5*b^2*hypergeom([1/2, 5/6], [11/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(5/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.187.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= -\frac{3b \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \sec^2(c + dx)\right) \sqrt{-\tan^2(c + dx)}}{2d(b \sec(c + dx))^{2/3}}$$

input `Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b*Cot[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(2*d*(b*Sec[c + d*x])^(2/3))`

**3.187.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c+dx) \sqrt[3]{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt[3]{b \csc\left(c+dx+\frac{\pi}{2}\right)}}{\csc\left(c+dx+\frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \frac{1}{\left(b \csc\left(\frac{1}{2}(2c+\pi)+dx\right)\right)^{2/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & b \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \left(\frac{\cos(c+dx)}{b}\right)^{2/3} dx \\
 & \quad \downarrow \text{3042} \\
 & b \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \left(\frac{\sin\left(c+dx+\frac{\pi}{2}\right)}{b}\right)^{2/3} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3b^2 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c+dx)\right)}{5d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{5/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]*(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b^2*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[c + d*x]^2]*Sin[c + d*x])/(5*d*(b*Sec[c + d*x])^(5/3)*Sqrt[Sin[c + d*x]^2])`

## 3.187.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.187.4 Maple [F]

$$\int \cos(dx + c) (b \sec(dx + c))^{\frac{1}{3}} dx$$

input `int(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

output `int(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

## 3.187.5 Fracas [F]

$$\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x + c))^(1/3)*cos(d*x + c), x)`

**3.187.6 Sympy [F]**

$$\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \sqrt[3]{b \sec(c + dx)} \cos(c + dx) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))**(1/3),x)`

output `Integral((b*sec(c + d*x))**(1/3)*cos(c + d*x), x)`

**3.187.7 Maxima [F]**

$$\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c), x)`

**3.187.8 Giac [F]**

$$\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c), x)`

**3.187.9 Mupad [F(-1)]**

Timed out.

$$\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \cos(c + dx) \left( \frac{b}{\cos(c + dx)} \right)^{1/3} dx$$

input `int(cos(c + d*x)*(b/cos(c + d*x))^(1/3), x)`output `int(cos(c + d*x)*(b/cos(c + d*x))^(1/3), x)`



### 3.188 $\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

3.188.1 Optimal result . . . . .	1108
3.188.2 Mathematica [A] (verified) . . . . .	1108
3.188.3 Rubi [A] (verified) . . . . .	1109
3.188.4 Maple [F] . . . . .	1110
3.188.5 Fricas [F] . . . . .	1110
3.188.6 Sympy [F] . . . . .	1111
3.188.7 Maxima [F] . . . . .	1111
3.188.8 Giac [F] . . . . .	1111
3.188.9 Mupad [F(-1)] . . . . .	1112

#### 3.188.1 Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= -\frac{3b^3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{8d(b \sec(c + dx))^{8/3} \sqrt{\sin^2(c + dx)}}$$

output `-3/8*b^3*hypergeom([1/2, 4/3], [7/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(8/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.188.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.02

$$\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$$

$$= \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \sec^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(2(c + dx))}{10d \sqrt{-\tan^2(c + dx)}}$$

input `Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(1/3),x]`

output `(3*Hypergeometric2F1[-5/6, 1/2, 1/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sin[2*(c + d*x)])/(10*d*Sqrt[-Tan[c + d*x]^2])`

**3.188.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c+dx) \sqrt[3]{b \sec(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt[3]{b \csc\left(c+dx+\frac{\pi}{2}\right)}}{\csc\left(c+dx+\frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{\left(b \csc\left(\frac{1}{2}(2c+\pi)+dx\right)\right)^{5/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & b^2 \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \left(\frac{\cos(c+dx)}{b}\right)^{5/3} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \left(\frac{\sin\left(c+dx+\frac{\pi}{2}\right)}{b}\right)^{5/3} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3b^3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c+dx)\right)}{8d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{8/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b^3*Hypergeometric2F1[1/2, 4/3, 7/3, Cos[c + d*x]^2]*Sin[c + d*x])/(8*d*(b*Sec[c + d*x])^(8/3)*Sqrt[Sin[c + d*x]^2])`

## 3.188.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.188.4 Maple [F]

$$\int \cos(dx+c)^2 (b \sec(dx+c))^{\frac{1}{3}} dx$$

input `int(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3),x)`

output `int(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3),x)`

## 3.188.5 Fracas [F]

$$\int \cos^2(c+dx) \sqrt[3]{b \sec(c+dx)} dx = \int (b \sec(dx+c))^{\frac{1}{3}} \cos(dx+c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^(1/3)*cos(d*x+c)^2, x)`

**3.188.6 Sympy [F]**

$$\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \sqrt[3]{b \sec(c + dx)} \cos^2(c + dx) dx$$

input `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(1/3),x)`

output `Integral((b*sec(c + d*x))**(1/3)*cos(c + d*x)**2, x)`

**3.188.7 Maxima [F]**

$$\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c)^2, x)`

**3.188.8 Giac [F]**

$$\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c)^2, x)`

**3.188.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \cos(c + dx)^2 \left( \frac{b}{\cos(c + dx)} \right)^{1/3} dx$$

input `int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3),x)`output `int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3), x)`

### 3.189 $\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx$

3.189.1 Optimal result . . . . .	1113
3.189.2 Mathematica [A] (verified) . . . . .	1113
3.189.3 Rubi [A] (verified) . . . . .	1114
3.189.4 Maple [F] . . . . .	1115
3.189.5 Fracas [F] . . . . .	1115
3.189.6 Sympy [F] . . . . .	1116
3.189.7 Maxima [F] . . . . .	1116
3.189.8 Giac [F] . . . . .	1116
3.189.9 Mupad [F(-1)] . . . . .	1117

#### 3.189.1 Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c + dx)\right) (b \sec(c + dx))^{7/3} \sin(c + dx)}{7bd \sqrt{\sin^2(c + dx)}}$$

output `3/7*hypergeom([-7/6, 1/2], [-1/6], cos(d*x+c)^2)*(b*sec(d*x+c))^(7/3)*sin(d*x+c)/b/d/(sin(d*x+c)^2)^(1/2)`

#### 3.189.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3 \operatorname{csc}(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \sec^2(c + dx)\right) (b \sec(c + dx))^{7/3} \sqrt{-\tan^2(c + dx)}}{10bd}$$

input `Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(4/3),x]`

output `(3*Csc[c + d*x]*Hypergeometric2F1[1/2, 5/3, 8/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(7/3)*Sqrt[-Tan[c + d*x]^2])/(10*b*d)`

**3.189.3 Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx)(b \sec(c+dx))^{4/3} dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c+dx))^{10/3} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c+dx + \frac{\pi}{2}))^{10/3} dx}{b^2} \\
 & \quad \downarrow \text{4259} \\
 & \quad \frac{\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{10/3}} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\left(\frac{\sin(c+dx + \frac{\pi}{2})}{b}\right)^{10/3}} dx}{b^2} \\
 & \quad \downarrow \text{3122} \\
 & \quad \frac{3 \sin(c+dx)(b \sec(c+dx))^{7/3} \text{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c+dx)\right)}{7bd\sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(4/3),x]`

output `(3*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(7/3)*Sin[c + d*x])/(7*b*d*Sqrt[Sin[c + d*x]^2])`

## 3.189.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.189.4 Maple [F]

$$\int \sec(dx + c)^2 (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `int(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3), x)`

output `int(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3), x)`

## 3.189.5 Fracas [F]

$$\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3), x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(1/3)*b*sec(d*x + c)^3, x)`



**3.189.6 Sympy [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(c + dx))^{4/3} \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(4/3),x)`

output `Integral((b*sec(c + d*x))**(4/3)*sec(c + d*x)**2, x)`

**3.189.7 Maxima [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{4/3} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^2, x)`

**3.189.8 Giac [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{4/3} \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^2, x)`

**3.189.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{4/3}}{\cos(c + dx)^2} dx$$

input `int((b/cos(c + d*x))^(4/3)/cos(c + d*x)^2,x)`output `int((b/cos(c + d*x))^(4/3)/cos(c + d*x)^2, x)`

### 3.190 $\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx$

3.190.1 Optimal result . . . . .	1118
3.190.2 Mathematica [A] (verified) . . . . .	1118
3.190.3 Rubi [A] (verified) . . . . .	1119
3.190.4 Maple [F] . . . . .	1120
3.190.5 Fricas [F] . . . . .	1120
3.190.6 Sympy [F] . . . . .	1121
3.190.7 Maxima [F] . . . . .	1121
3.190.8 Giac [F] . . . . .	1121
3.190.9 Mupad [F(-1)] . . . . .	1122

#### 3.190.1 Optimal result

Integrand size = 19, antiderivative size = 55

$$\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) (b \sec(c + dx))^{4/3} \sin(c + dx)}{4d \sqrt{\sin^2(c + dx)}}$$

output `3/4*hypergeom([-2/3, 1/2], [1/3], cos(d*x+c)^2)*(b*sec(d*x+c))^(4/3)*sin(d*x+c)/d/(sin(d*x+c)^2)^(1/2)`

#### 3.190.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.09

$$\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \sec^2(c + dx)\right) (b \sec(c + dx))^{7/3} \sqrt{-\tan^2(c + dx)}}{7bd}$$

input `Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(4/3),x]`

output `(3*Cot[c + d*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(7/3)*Sqrt[-Tan[c + d*x]^2])/(7*b*d)`

**3.190.3 Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c+dx)(b \sec(c+dx))^{4/3} dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c+dx))^{7/3} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{7/3} dx}{b} \\
 & \quad \downarrow \text{4259} \\
 & \quad \frac{\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{7/3}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{7/3}} dx}{b} \\
 & \quad \downarrow \text{3122} \\
 & \quad \frac{3 \sin(c+dx)(b \sec(c+dx))^{4/3} \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]*(b*Sec[c + d*x])^(4/3),x]`

output `(3*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sin[c + d*x])/(4*d*Sqrt[Sin[c + d*x]^2])`

## 3.190.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.190.4 Maple [F]

$$\int \sec(dx + c) (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `int(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x)`

output `int(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x)`

## 3.190.5 Fracas [F]

$$\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x, algorithm="fracas")`

output `integral((b*sec(d*x + c))^(1/3)*b*sec(d*x + c)^2, x)`

**3.190.6 Sympy [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(c + dx))^{4/3} \sec(c + dx) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))**(4/3),x)`

output `Integral((b*sec(c + d*x))**(4/3)*sec(c + d*x), x)`

**3.190.7 Maxima [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{4/3} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c), x)`

**3.190.8 Giac [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{4/3} \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c), x)`

**3.190.9 Mupad [F(-1)]**

Timed out.

$$\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{4/3}}{\cos(c + dx)} dx$$

input `int((b/cos(c + d*x))^(4/3)/cos(c + d*x),x)`output `int((b/cos(c + d*x))^(4/3)/cos(c + d*x), x)`

### 3.191 $\int (b \sec(c + dx))^{4/3} dx$

3.191.1 Optimal result . . . . .	1123
3.191.2 Mathematica [A] (verified) . . . . .	1123
3.191.3 Rubi [A] (verified) . . . . .	1124
3.191.4 Maple [F] . . . . .	1125
3.191.5 Fricas [F] . . . . .	1125
3.191.6 Sympy [F] . . . . .	1126
3.191.7 Maxima [F] . . . . .	1126
3.191.8 Giac [F] . . . . .	1126
3.191.9 Mupad [F(-1)] . . . . .	1127

#### 3.191.1 Optimal result

Integrand size = 12, antiderivative size = 54

$$\int (b \sec(c + dx))^{4/3} dx = \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sin^2(c + dx)}}$$

output `3*b*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*(b*sec(d*x+c))^(1/3)*sin(d*x+c)/d/(sin(d*x+c)^2)^(1/2)`

#### 3.191.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06

$$\int (b \sec(c + dx))^{4/3} dx = \frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \sec^2(c + dx)\right) (b \sec(c + dx))^{4/3} \sqrt{-\tan^2(c + dx)}}{4d}$$

input `Integrate[(b*Sec[c + d*x])^(4/3), x]`

output `(3*Cot[c + d*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/(4*d)`



**3.191.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( b \csc \left( c + dx + \frac{\pi}{2} \right) \right)^{4/3} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\left( \frac{\cos(c + dx)}{b} \right)^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\left( \frac{\sin(c + dx + \frac{\pi}{2})}{b} \right)^{4/3}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3b \sin(c + dx) \sqrt[3]{b \sec(c + dx)} \operatorname{Hypergeometric2F1} \left( -\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx) \right)}{d \sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(4/3),x]`

output `(3*b*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sin[c + d*x])/(d*Sqrt[Sin[c + d*x]^2])`

**3.191.3.1** Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.191.4** Maple **[F]**

$$\int (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `int((b*sec(d*x+c))^(4/3),x)`

output `int((b*sec(d*x+c))^(4/3),x)`

**3.191.5** Fracas **[F]**

$$\int (b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `integrate((b*sec(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(1/3)*b*sec(d*x + c), x)`

**3.191.6 Sympy [F]**

$$\int (b \sec(c + dx))^{4/3} dx = \int (b \sec(c + dx))^{\frac{4}{3}} dx$$

input `integrate((b*sec(d*x+c))**(4/3),x)`

output `Integral((b*sec(c + d*x))**(4/3), x)`

**3.191.7 Maxima [F]**

$$\int (b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `integrate((b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(4/3), x)`

**3.191.8 Giac [F]**

$$\int (b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `integrate((b*sec(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(4/3), x)`

**3.191.9 Mupad [F(-1)]**

Timed out.

$$\int (b \sec(c + dx))^{4/3} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{4/3} dx$$

input `int((b/cos(c + d*x))^(4/3),x)`output `int((b/cos(c + d*x))^(4/3), x)`

### 3.192 $\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx$

3.192.1 Optimal result . . . . .	1128
3.192.2 Mathematica [A] (verified) . . . . .	1128
3.192.3 Rubi [A] (verified) . . . . .	1129
3.192.4 Maple [F] . . . . .	1130
3.192.5 Fricas [F] . . . . .	1130
3.192.6 Sympy [F(-1)] . . . . .	1131
3.192.7 Maxima [F] . . . . .	1131
3.192.8 Giac [F] . . . . .	1131
3.192.9 Mupad [F(-1)] . . . . .	1132

#### 3.192.1 Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3b^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \sec(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}}$$

output `-3/2*b^2*hypergeom([1/3, 1/2],[4/3],cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(2/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.192.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.97

$$\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3b \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \sec^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sqrt{-\tan^2(c + dx)}}{d}$$

input `Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(4/3),x]`

output `(3*b*Cot[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sqrt[-Tan[c + d*x]^2])/d`

**3.192.3 Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c+dx)(b \sec(c+dx))^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{4/3}}{\csc(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \sqrt[3]{b \csc\left(\frac{1}{2}(2c+\pi)+dx\right)} dx \\
 & \quad \downarrow \text{4259} \\
 & b \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\sqrt[3]{\frac{\cos(c+dx)}{b}}} dx \\
 & \quad \downarrow \text{3042} \\
 & b \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \frac{1}{\sqrt[3]{\frac{\sin(c+dx+\frac{\pi}{2})}{b}}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3b^2 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c+dx)\right)}{2d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{2/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]*(b*Sec[c + d*x])^(4/3),x]`

output `(-3*b^2*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[c + d*x]^2]*Sin[c + d*x])/(2*d*(b*Sec[c + d*x])^(2/3)*Sqrt[Sin[c + d*x]^2])`

## 3.192.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.192.4 Maple [F]

$$\int \cos(dx + c) (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `int(cos(d*x+c)*(b*sec(d*x+c))^(4/3),x)`

output `int(cos(d*x+c)*(b*sec(d*x+c))^(4/3),x)`

## 3.192.5 Fracas [F]

$$\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c), x)`

**3.192.6 Sympy [F(-1)]**

Timed out.

$$\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))**(4/3),x)`output `Timed out`**3.192.7 Maxima [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c), x)`**3.192.8 Giac [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^(4/3),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c), x)`



**3.192.9 Mupad [F(-1)]**

Timed out.

$$\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx = \int \cos(c + dx) \left( \frac{b}{\cos(c + dx)} \right)^{4/3} dx$$

input `int(cos(c + d*x)*(b/cos(c + d*x))^(4/3), x)`output `int(cos(c + d*x)*(b/cos(c + d*x))^(4/3), x)`

### 3.193 $\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx$

3.193.1 Optimal result . . . . .	1133
3.193.2 Mathematica [A] (verified) . . . . .	1133
3.193.3 Rubi [A] (verified) . . . . .	1134
3.193.4 Maple [F] . . . . .	1135
3.193.5 Fricas [F] . . . . .	1135
3.193.6 Sympy [F(-1)] . . . . .	1136
3.193.7 Maxima [F] . . . . .	1136
3.193.8 Giac [F] . . . . .	1136
3.193.9 Mupad [F(-1)] . . . . .	1137

#### 3.193.1 Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3b^3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5d(b \sec(c + dx))^{5/3} \sqrt{\sin^2(c + dx)}}$$

output `-3/5*b^3*hypergeom([1/2, 5/6], [11/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(5/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.193.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3b^2 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \sec^2(c + dx)\right) \sqrt{-\tan^2(c + dx)}}{2d(b \sec(c + dx))^{2/3}}$$

input `Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(4/3),x]`

output `(-3*b^2*Cot[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(2*d*(b*Sec[c + d*x])^(2/3))`

**3.193.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c+dx)(b \sec(c+dx))^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^{4/3}}{\csc(c+dx+\frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{2/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & b^2 \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \left(\frac{\cos(c+dx)}{b}\right)^{2/3} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \int \left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{2/3} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3b^3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c+dx)\right)}{5d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{5/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^(4/3),x]`

output `(-3*b^3*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[c + d*x]^2]*Sin[c + d*x])/(5*d*(b*Sec[c + d*x])^(5/3)*Sqrt[Sin[c + d*x]^2])`

**3.193.3.1** Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.193.4** Maple **[F]**

$$\int \cos(dx + c)^2 (b \sec(dx + c))^{\frac{4}{3}} dx$$

input `int(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3), x)`

output `int(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3), x)`

**3.193.5** Fracas **[F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3), x, algorithm="fracas")`

output `integral((b*sec(d*x + c))^(1/3)*b*cos(d*x + c)^2*sec(d*x + c), x)`

**3.193.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(4/3),x)`output `Timed out`**3.193.7 Maxima [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c)^2, x)`**3.193.8 Giac [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c)^2, x)`

**3.193.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx = \int \cos(c + dx)^2 \left( \frac{b}{\cos(c + dx)} \right)^{4/3} dx$$

input `int(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3),x)`output `int(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3), x)`

**3.194** 
$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

3.194.1 Optimal result . . . . . 1138  
 3.194.2 Mathematica [A] (verified) . . . . . 1138  
 3.194.3 Rubi [A] (verified) . . . . . 1139  
 3.194.4 Maple [F] . . . . . 1140  
 3.194.5 Fricas [F] . . . . . 1140  
 3.194.6 Sympy [F] . . . . . 1141  
 3.194.7 Maxima [F] . . . . . 1141  
 3.194.8 Giac [F] . . . . . 1141  
 3.194.9 Mupad [F(-1)] . . . . . 1142

**3.194.1 Optimal result**

Integrand size = 21, antiderivative size = 58

$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{2bd \sqrt{\sin^2(c+dx)}}$$

output `3/2*hypergeom([-1/3, 1/2], [2/3], cos(d*x+c)^2)*(b*sec(d*x+c))^(2/3)*sin(d*x+c)/b/d/(sin(d*x+c)^2)^(1/2)`

**3.194.2 Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \frac{3 \operatorname{csc}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \sec^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sqrt{-\tan^2(c+dx)}}{5bd}$$

input `Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(1/3),x]`

output `(3*Csc[c + d*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/(5*b*d)`

---

3.194. 
$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

**3.194.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{5/3} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{5/3} dx}{b^2} \\
 & \quad \downarrow \text{4259} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{5/3}} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \frac{1}{\left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{5/3}} dx}{b^2} \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(c+dx) (b \sec(c+dx))^{2/3} \text{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c+dx)\right)}{2bd \sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(1/3),x]`

output `(3*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sin[c + d*x])/(2*b*d*Sqrt[Sin[c + d*x]^2])`

---

3.194.  $\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$



## 3.194.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.194.4 Maple [F]

$$\int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `int(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)`

output `int(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)`

## 3.194.5 Fracas [F]

$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)/b, x)`

---

3.194.  $\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

**3.194.6 Sympy [F]**

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

input `integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(1/3),x)`

output `Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(1/3), x)`

**3.194.7 Maxima [F]**

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)`

**3.194.8 Giac [F]**

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)`

**3.194.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b\sec(c+dx)}} dx = \int \frac{1}{\cos(c+dx)^2 \left(\frac{b}{\cos(c+dx)}\right)^{1/3}} dx$$

input `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3)),x)`output `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3)), x)`

**3.195**  $\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

3.195.1 Optimal result . . . . . 1143  
 3.195.2 Mathematica [A] (verified) . . . . . 1143  
 3.195.3 Rubi [A] (verified) . . . . . 1144  
 3.195.4 Maple [F] . . . . . 1145  
 3.195.5 Fracas [F] . . . . . 1145  
 3.195.6 Sympy [F] . . . . . 1146  
 3.195.7 Maxima [F] . . . . . 1146  
 3.195.8 Giac [F] . . . . . 1146  
 3.195.9 Mupad [F(-1)] . . . . . 1147

**3.195.1 Optimal result**

Integrand size = 19, antiderivative size = 53

$$\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{d \sqrt[3]{b \sec(c+dx)} \sqrt{\sin^2(c+dx)}}$$

output `-3*hypergeom([1/6, 1/2], [7/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)`

**3.195.2 Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.13

$$\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \frac{3 \cot(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \sec^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sqrt{-\tan^2(c+dx)}}{2bd}$$

input `Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(1/3), x]`

output `(3*Cot[c + d*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/(2*b*d)`

---

3.195.  $\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

**3.195.3 Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{2/3} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx+\frac{\pi}{2}))^{2/3} dx}{b} \\
 & \quad \downarrow \text{4259} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{2/3}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \frac{1}{\left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{2/3}} dx}{b} \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c+dx)\right)}{d \sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]/(b*Sec[c + d*x])^(1/3),x]`

output `(-3*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])`

---

3.195.  $\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

## 3.195.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.195.4 Maple [F]

$$\int \frac{\sec(dx+c)}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `int(sec(d*x+c)/(b*sec(d*x+c))^(1/3),x)`

output `int(sec(d*x+c)/(b*sec(d*x+c))^(1/3),x)`

## 3.195.5 Fracas [F]

$$\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \int \frac{\sec(dx+c)}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^(2/3)/b, x)`

---

3.195.  $\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

**3.195.6 Sympy [F]**

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))**(1/3),x)`

output `Integral(sec(c + d*x)/(b*sec(c + d*x))**(1/3), x)`

**3.195.7 Maxima [F]**

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(1/3), x)`

**3.195.8 Giac [F]**

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(1/3), x)`

**3.195.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec(c+dx)}{\sqrt[3]{b\sec(c+dx)}} dx = \int \frac{1}{\cos(c+dx) \left(\frac{b}{\cos(c+dx)}\right)^{1/3}} dx$$

input `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/3)),x)`output `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/3)), x)`



**3.196**  $\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx$

3.196.1 Optimal result . . . . . 1148  
 3.196.2 Mathematica [A] (verified) . . . . . 1148  
 3.196.3 Rubi [A] (verified) . . . . . 1149  
 3.196.4 Maple [F] . . . . . 1150  
 3.196.5 Fricas [F] . . . . . 1150  
 3.196.6 Sympy [F] . . . . . 1151  
 3.196.7 Maxima [F] . . . . . 1151  
 3.196.8 Giac [F] . . . . . 1151  
 3.196.9 Mupad [F(-1)] . . . . . 1152

**3.196.1 Optimal result**

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx = -\frac{3b \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d(b \sec(c + dx))^{4/3} \sqrt{\sin^2(c + dx)}}$$

output `-3/4*b*hypergeom([1/2, 2/3],[5/3],cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(4/3)/(sin(d*x+c)^2)^(1/2)`

**3.196.2 Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx = -\frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \sec^2(c + dx)\right) \sqrt{-\tan^2(c + dx)}}{d \sqrt[3]{b \sec(c + dx)}}$$

input `Integrate[(b*Sec[c + d*x])^(-1/3),x]`

output `(-3*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(d*(b*Sec[c + d*x])^(1/3))`

**3.196.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{b \csc\left(c+dx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{4259} \\
 & \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \sqrt[3]{\frac{\cos(c+dx)}{b}} dx \\
 & \quad \downarrow \text{3042} \\
 & \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \sqrt[3]{\frac{\sin\left(c+dx+\frac{\pi}{2}\right)}{b}} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3b \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{4/3}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(-1/3),x]`

output `(-3*b*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[c + d*x]^2]*Sin[c + d*x])/(4*d*(b*Sec[c + d*x])^(4/3)*Sqrt[Sin[c + d*x]^2])`

**3.196.3.1** Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.196.4** Maple **[F]**

$$\int \frac{1}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `int(1/(b*sec(d*x+c))^(1/3),x)`

output `int(1/(b*sec(d*x+c))^(1/3),x)`

**3.196.5** Fracas **[F]**

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(2/3)/(b*sec(d*x + c)), x)`

**3.196.6 Sympy [F]**

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx$$

input `integrate(1/(b*sec(d*x+c))**(1/3), x)`

output `Integral((b*sec(c + d*x))**(-1/3), x)`

**3.196.7 Maxima [F]**

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(1/3), x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(1/3), x)`

**3.196.8 Giac [F]**

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(1/3), x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(1/3), x)`

**3.196.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}} dx$$

input `int(1/(b/cos(c + d*x))^(1/3),x)`output `int(1/(b/cos(c + d*x))^(1/3), x)`

**3.197**  $\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

3.197.1 Optimal result . . . . . 1153  
 3.197.2 Mathematica [A] (verified) . . . . . 1153  
 3.197.3 Rubi [A] (verified) . . . . . 1154  
 3.197.4 Maple [F] . . . . . 1155  
 3.197.5 Fricas [F] . . . . . 1155  
 3.197.6 Sympy [F] . . . . . 1156  
 3.197.7 Maxima [F] . . . . . 1156  
 3.197.8 Giac [F] . . . . . 1156  
 3.197.9 Mupad [F(-1)] . . . . . 1157

**3.197.1 Optimal result**

Integrand size = 19, antiderivative size = 58

$$\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = -\frac{3b^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{7d(b \sec(c+dx))^{7/3} \sqrt{\sin^2(c+dx)}}$$

output `-3/7*b^2*hypergeom([1/2, 7/6], [13/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(7/3)/(sin(d*x+c)^2)^(1/2)`

**3.197.2 Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = -\frac{3b \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \sec^2(c+dx)\right) \sqrt{-\tan^2(c+dx)}}{4d(b \sec(c+dx))^{4/3}}$$

input `Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b*Cot[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(4*d*(b*Sec[c + d*x])^(4/3))`

---

3.197.  $\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

**3.197.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc\left(c+dx+\frac{\pi}{2}\right) \sqrt[3]{b \csc\left(c+dx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \frac{1}{\left(b \csc\left(\frac{1}{2}(2c+\pi)+dx\right)\right)^{4/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & b \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\cos(c+dx)}{b}\right)^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & b \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\sin\left(c+dx+\frac{\pi}{2}\right)}{b}\right)^{4/3} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3b^2 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c+dx)\right)}{7d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{7/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]/(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b^2*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[c + d*x]^2]*Sin[c + d*x])/(7*d*(b*Sec[c + d*x])^(7/3)*Sqrt[Sin[c + d*x]^2])`

---

3.197.  $\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

## 3.197.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.197.4 Maple [F]

$$\int \frac{\cos(dx+c)}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `int(cos(d*x+c)/(b*sec(d*x+c))^(1/3),x)`

output `int(cos(d*x+c)/(b*sec(d*x+c))^(1/3),x)`

## 3.197.5 Fracas [F]

$$\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \int \frac{\cos(dx+c)}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^(2/3)*cos(d*x+c)/(b*sec(d*x+c)), x)`

---

3.197.  $\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$



**3.197.6 Sympy [F]**

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\cos(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))**(1/3),x)`

output `Integral(cos(c + d*x)/(b*sec(c + d*x))**(1/3), x)`

**3.197.7 Maxima [F]**

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(1/3), x)`

**3.197.8 Giac [F]**

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(1/3), x)`

**3.197.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \int \frac{\cos(c+dx)}{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}} dx$$

input `int(cos(c + d*x)/(b/cos(c + d*x))^(1/3), x)`output `int(cos(c + d*x)/(b/cos(c + d*x))^(1/3), x)`

**3.198**  $\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

3.198.1 Optimal result . . . . . 1158  
 3.198.2 Mathematica [A] (verified) . . . . . 1158  
 3.198.3 Rubi [A] (verified) . . . . . 1159  
 3.198.4 Maple [F] . . . . . 1160  
 3.198.5 Fricas [F] . . . . . 1160  
 3.198.6 Sympy [F] . . . . . 1161  
 3.198.7 Maxima [F] . . . . . 1161  
 3.198.8 Giac [F] . . . . . 1161  
 3.198.9 Mupad [F(-1)] . . . . . 1162

**3.198.1 Optimal result**

Integrand size = 21, antiderivative size = 58

$$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = -\frac{3b^3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{10d(b \sec(c+dx))^{10/3} \sqrt{\sin^2(c+dx)}}$$

output `-3/10*b^3*hypergeom([1/2, 5/3],[8/3],cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(10/3)/(sin(d*x+c)^2)^(1/2)`

**3.198.2 Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = -\frac{3b^2 \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \sec^2(c+dx)\right) \sqrt{-\tan^2(c+dx)}}{7d(b \sec(c+dx))^{7/3}}$$

input `Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b^2*Cot[c + d*x]*Hypergeometric2F1[-7/6, 1/2, -1/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(7*d*(b*Sec[c + d*x])^(7/3))`

---

3.198.  $\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

**3.198.3 Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt[3]{b \csc\left(c+dx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{\left(b \csc\left(\frac{1}{2}(2c+\pi)+dx\right)\right)^{7/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & b^2 \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\cos(c+dx)}{b}\right)^{7/3} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\sin\left(c+dx+\frac{\pi}{2}\right)}{b}\right)^{7/3} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3b^3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c+dx)\right)}{10d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{10/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2/(b*Sec[c + d*x])^(1/3),x]`

output `(-3*b^3*Hypergeometric2F1[1/2, 5/3, 8/3, Cos[c + d*x]^2]*Sin[c + d*x])/(10*d*(b*Sec[c + d*x])^(10/3)*Sqrt[Sin[c + d*x]^2])`

---

3.198.  $\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

## 3.198.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.198.4 Maple [F]

$$\int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `int(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)`

output `int(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)`

## 3.198.5 Fracas [F]

$$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^(2/3)*cos(d*x+c)^2/(b*sec(d*x+c)), x)`

---

3.198.  $\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$

**3.198.6 Sympy [F]**

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\cos^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

input `integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(1/3),x)`

output `Integral(cos(c + d*x)**2/(b*sec(c + d*x))**(1/3), x)`

**3.198.7 Maxima [F]**

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)`

**3.198.8 Giac [F]**

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)`

**3.198.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\cos(c + dx)^2}{\left(\frac{b}{\cos(c + dx)}\right)^{1/3}} dx$$

input `int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/3),x)`output `int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/3), x)`

**3.199**       $\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$

3.199.1 Optimal result . . . . . 1163  
 3.199.2 Mathematica [A] (verified) . . . . . 1163  
 3.199.3 Rubi [A] (verified) . . . . . 1164  
 3.199.4 Maple [F] . . . . . 1165  
 3.199.5 Fricas [F] . . . . . 1165  
 3.199.6 Sympy [F] . . . . . 1166  
 3.199.7 Maxima [F] . . . . . 1166  
 3.199.8 Giac [F] . . . . . 1166  
 3.199.9 Mupad [F(-1)] . . . . . 1167

**3.199.1 Optimal result**

Integrand size = 21, antiderivative size = 56

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{bd\sqrt[3]{b \sec(c + dx)}\sqrt{\sin^2(c + dx)}}$$

output `-3*hypergeom([1/6, 1/2], [7/6], cos(d*x+c)^2)*sin(d*x+c)/b/d/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)`

**3.199.2 Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.07

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \sec^2(c + dx)\right) (b \sec(c + dx))^{2/3} \sqrt{-\tan^2(c + dx)}}{2b^2d}$$

input `Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(4/3),x]`

output `(3*Cot[c + d*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/(2*b^2*d)`



**3.199.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{\int (b \sec(c+dx))^{2/3} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \csc(c+dx + \frac{\pi}{2}))^{2/3} dx}{b^2} \\
 & \quad \downarrow \text{4259} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{2/3}} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \frac{1}{\left(\frac{\sin(c+dx + \frac{\pi}{2})}{b}\right)^{2/3}} dx}{b^2} \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(4/3),x]`

output `(-3*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[c + d*x]^2]*Sin[c + d*x])/(b*d*(b*Sec[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])`

**3.199.3.1** Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.199.4** Maple **[F]**

$$\int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

input `int(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3),x)`

output `int(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3),x)`

**3.199.5** Fracas **[F]**

$$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*sec(d*x+c))^(2/3)/b^2, x)`

**3.199.6 Sympy [F]**

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx$$

input `integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(4/3),x)`

output `Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(4/3), x)`

**3.199.7 Maxima [F]**

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)`

**3.199.8 Giac [F]**

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)`

**3.199.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \int \frac{1}{\cos(c+dx)^2 \left(\frac{b}{\cos(c+dx)}\right)^{4/3}} dx$$

input `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3)),x)`output `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3)), x)`

### 3.200 $\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{4/3}} dx$

3.200.1 Optimal result . . . . .	1168
3.200.2 Mathematica [A] (verified) . . . . .	1168
3.200.3 Rubi [A] (verified) . . . . .	1169
3.200.4 Maple [F] . . . . .	1170
3.200.5 Fricas [F] . . . . .	1170
3.200.6 Sympy [F] . . . . .	1171
3.200.7 Maxima [F] . . . . .	1171
3.200.8 Giac [F] . . . . .	1171
3.200.9 Mupad [F(-1)] . . . . .	1172

#### 3.200.1 Optimal result

Integrand size = 19, antiderivative size = 55

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{4/3}} dx = -\frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d(b \sec(c + dx))^{4/3} \sqrt{\sin^2(c + dx)}}$$

```
output -3/4*hypergeom([1/2, 2/3],[5/3],cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(4/3)/(sin(d*x+c)^2)^(1/2)
```

#### 3.200.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.05

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \sec^2(c + dx)\right) \sqrt{-\tan^2(c + dx)}}{bd \sqrt[3]{b \sec(c + dx)}}$$

```
input Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(4/3),x]
```

```
output (-3*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(b*d*(b*Sec[c + d*x])^(1/3))
```

**3.200.3 Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)}{(b \sec(c+dx))^{4/3}} dx \\
 & \quad \downarrow \text{2030} \\
 & \int \frac{1}{\sqrt[3]{b \sec(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{b \csc\left(c+dx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{4259} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \sqrt[3]{\frac{\cos(c+dx)}{b}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \sqrt[3]{\frac{\sin\left(c+dx+\frac{\pi}{2}\right)}{b}} dx}{b} \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{4/3}}
 \end{aligned}$$

input `Int[Sec[c + d*x]/(b*Sec[c + d*x])^(4/3),x]`

output `(-3*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[c + d*x]^2]*Sin[c + d*x])/(4*d*(b*Sec[c + d*x])^(4/3)*Sqrt[Sin[c + d*x]^2])`

## 3.200.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.200.4 Maple [F]

$$\int \frac{\sec(dx+c)}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

input `int(sec(d*x+c)/(b*sec(d*x+c))^(4/3),x)`

output `int(sec(d*x+c)/(b*sec(d*x+c))^(4/3),x)`

## 3.200.5 Fracas [F]

$$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \int \frac{\sec(dx+c)}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(4/3),x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^(2/3)/(b^2*sec(d*x+c)), x)`

**3.200.6 Sympy [F]**

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(c + dx)}{(b \sec(c + dx))^{4/3}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))**(4/3),x)`

output `Integral(sec(c + d*x)/(b*sec(c + d*x))**(4/3), x)`

**3.200.7 Maxima [F]**

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(4/3), x)`

**3.200.8 Giac [F]**

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(sec(d*x+c)/(b*sec(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate(sec(d*x + c)/(b*sec(d*x + c))^(4/3), x)`



**3.200.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec(c+dx)}{(b\sec(c+dx))^{4/3}} dx = \int \frac{1}{\cos(c+dx) \left(\frac{b}{\cos(c+dx)}\right)^{4/3}} dx$$

input `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(4/3)),x)`output `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(4/3)), x)`

### 3.201 $\int \frac{1}{(b \sec(c+dx))^{4/3}} dx$

3.201.1 Optimal result . . . . .	1173
3.201.2 Mathematica [A] (verified) . . . . .	1173
3.201.3 Rubi [A] (verified) . . . . .	1174
3.201.4 Maple [F] . . . . .	1175
3.201.5 Fricas [F] . . . . .	1175
3.201.6 Sympy [F] . . . . .	1176
3.201.7 Maxima [F] . . . . .	1176
3.201.8 Giac [F] . . . . .	1176
3.201.9 Mupad [F(-1)] . . . . .	1177

#### 3.201.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(b \sec(c+dx))^{4/3}} dx = -\frac{3b \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{7d(b \sec(c+dx))^{7/3} \sqrt{\sin^2(c+dx)}}$$

output `-3/7*b*hypergeom([1/2, 7/6],[13/6],cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(7/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.201.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.02

$$\int \frac{1}{(b \sec(c+dx))^{4/3}} dx = \frac{3 \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \sec^2(c+dx)\right) \sqrt{-\tan^2(c+dx)}}{4d(b \sec(c+dx))^{4/3}}$$

input `Integrate[(b*Sec[c + d*x])^(-4/3),x]`

output `(-3*Cot[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(4*d*(b*Sec[c + d*x])^(4/3))`

**3.201.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(b \sec(c + dx))^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{4/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & \left(\frac{\cos(c + dx)}{b}\right)^{2/3} (b \sec(c + dx))^{2/3} \int \left(\frac{\cos(c + dx)}{b}\right)^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \left(\frac{\cos(c + dx)}{b}\right)^{2/3} (b \sec(c + dx))^{2/3} \int \left(\frac{\sin(c + dx + \frac{\pi}{2})}{b}\right)^{4/3} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3b \sin(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right)}{7d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{7/3}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^(-4/3),x]`

output `(-3*b*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[c + d*x]^2]*Sin[c + d*x])/(7*d*(b*Sec[c + d*x])^(7/3)*Sqrt[Sin[c + d*x]^2])`

**3.201.3.1** Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.201.4** Maple **[F]**

$$\int \frac{1}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

input `int(1/(b*sec(d*x+c))^(4/3),x)`

output `int(1/(b*sec(d*x+c))^(4/3),x)`

**3.201.5** Fracas **[F]**

$$\int \frac{1}{(b \sec(c + dx))^{4/3}} dx = \int \frac{1}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

input `integrate(1/(b*sec(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(2/3)/(b^2*sec(d*x + c)^2), x)`

**3.201.6 Sympy [F]**

$$\int \frac{1}{(b \sec(c + dx))^{4/3}} dx = \int \frac{1}{(b \sec(c + dx))^{4/3}} dx$$

input `integrate(1/(b*sec(d*x+c))**(4/3), x)`

output `Integral((b*sec(c + d*x))**(-4/3), x)`

**3.201.7 Maxima [F]**

$$\int \frac{1}{(b \sec(c + dx))^{4/3}} dx = \int \frac{1}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(1/(b*sec(d*x+c))^(4/3), x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(4/3), x)`

**3.201.8 Giac [F]**

$$\int \frac{1}{(b \sec(c + dx))^{4/3}} dx = \int \frac{1}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(1/(b*sec(d*x+c))^(4/3), x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(4/3), x)`

**3.201.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(b \sec(c + dx))^{4/3}} dx = \int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{4/3}} dx$$

input `int(1/(b/cos(c + d*x))^(4/3),x)`output `int(1/(b/cos(c + d*x))^(4/3), x)`

### 3.202 $\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{4/3}} dx$

3.202.1 Optimal result . . . . .	1178
3.202.2 Mathematica [A] (verified) . . . . .	1178
3.202.3 Rubi [A] (verified) . . . . .	1179
3.202.4 Maple [F] . . . . .	1180
3.202.5 Fricas [F] . . . . .	1180
3.202.6 Sympy [F] . . . . .	1181
3.202.7 Maxima [F] . . . . .	1181
3.202.8 Giac [F] . . . . .	1181
3.202.9 Mupad [F(-1)] . . . . .	1182

#### 3.202.1 Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx = -\frac{3b^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{10d(b \sec(c + dx))^{10/3} \sqrt{\sin^2(c + dx)}}$$

```
output -3/10*b^2*hypergeom([1/2, 5/3], [8/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(10/3)/(sin(d*x+c)^2)^(1/2)
```

#### 3.202.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \frac{3b \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \sec^2(c + dx)\right) \sqrt{-\tan^2(c + dx)}}{7d(b \sec(c + dx))^{7/3}}$$

```
input Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(4/3), x]
```

```
output (-3*b*Cot[c + d*x]*Hypergeometric2F1[-7/6, 1/2, -1/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(7*d*(b*Sec[c + d*x])^(7/3))
```

**3.202.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})(b \csc(c+dx+\frac{\pi}{2}))^{4/3}} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{7/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & b \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\cos(c+dx)}{b}\right)^{7/3} dx \\
 & \quad \downarrow \text{3042} \\
 & b \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{7/3} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3b^2 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c+dx)\right)}{10d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{10/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]/(b*Sec[c + d*x])^(4/3),x]`

output `(-3*b^2*Hypergeometric2F1[1/2, 5/3, 8/3, Cos[c + d*x]^2]*Sin[c + d*x])/(10*d*(b*Sec[c + d*x])^(10/3)*Sqrt[Sin[c + d*x]^2])`



## 3.202.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.202.4 Maple [F]

$$\int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

input `int(cos(d*x+c)/(b*sec(d*x+c))^(4/3),x)`

output `int(cos(d*x+c)/(b*sec(d*x+c))^(4/3),x)`

## 3.202.5 Fracas [F]

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(2/3)*cos(d*x + c)/(b^2*sec(d*x + c)^2), x)`

**3.202.6 Sympy [F]**

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))**(4/3),x)`

output `Integral(cos(c + d*x)/(b*sec(c + d*x))**(4/3), x)`

**3.202.7 Maxima [F]**

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(4/3), x)`

**3.202.8 Giac [F]**

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(cos(d*x+c)/(b*sec(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate(cos(d*x + c)/(b*sec(d*x + c))^(4/3), x)`

**3.202.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos(c + dx)}{\left(\frac{b}{\cos(c+dx)}\right)^{4/3}} dx$$

input `int(cos(c + d*x)/(b/cos(c + d*x))^(4/3), x)`output `int(cos(c + d*x)/(b/cos(c + d*x))^(4/3), x)`

### 3.203 $\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$

3.203.1 Optimal result . . . . .	1183
3.203.2 Mathematica [A] (verified) . . . . .	1183
3.203.3 Rubi [A] (verified) . . . . .	1184
3.203.4 Maple [F] . . . . .	1185
3.203.5 Fracas [F] . . . . .	1185
3.203.6 Sympy [F] . . . . .	1186
3.203.7 Maxima [F] . . . . .	1186
3.203.8 Giac [F] . . . . .	1186
3.203.9 Mupad [F(-1)] . . . . .	1187

#### 3.203.1 Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx = -\frac{3b^3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{13}{6}, \frac{19}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{13d(b \sec(c+dx))^{13/3} \sqrt{\sin^2(c+dx)}}$$

output `-3/13*b^3*hypergeom([1/2, 13/6], [19/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(13/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.203.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx = -\frac{3b^2 \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{3}, \frac{1}{2}, -\frac{2}{3}, \sec^2(c+dx)\right) \sqrt{-\tan^2(c+dx)}}{10d(b \sec(c+dx))^{10/3}}$$

input `Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(4/3),x]`

output `(-3*b^2*Cot[c + d*x]*Hypergeometric2F1[-5/3, 1/2, -2/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(10*d*(b*Sec[c + d*x])^(10/3))`

**3.203.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^2 (b \csc(c+dx+\frac{\pi}{2}))^{4/3}} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \frac{1}{(b \csc(\frac{1}{2}(2c+\pi)+dx))^{10/3}} dx \\
 & \quad \downarrow \text{4259} \\
 & b^2 \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\cos(c+dx)}{b}\right)^{10/3} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3} \int \left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{10/3} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3b^3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{13}{6}, \frac{19}{6}, \cos^2(c+dx)\right)}{13d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{13/3}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2/(b*Sec[c + d*x])^(4/3),x]`

output `(-3*b^3*Hypergeometric2F1[1/2, 13/6, 19/6, Cos[c + d*x]^2]*Sin[c + d*x])/ (13*d*(b*Sec[c + d*x])^(13/3)*Sqrt[Sin[c + d*x]^2])`

## 3.203.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.203.4 Maple [F]

$$\int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

input `int(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3),x)`

output `int(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3),x)`

## 3.203.5 Fracas [F]

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3),x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^(2/3)*cos(d*x+c)^2/(b^2*sec(d*x+c)^2), x)`

**3.203.6 Sympy [F]**

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx$$

input `integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(4/3),x)`

output `Integral(cos(c + d*x)**2/(b*sec(c + d*x))**(4/3), x)`

**3.203.7 Maxima [F]**

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)`

**3.203.8 Giac [F]**

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)`

**3.203.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \int \frac{\cos(c+dx)^2}{\left(\frac{b}{\cos(c+dx)}\right)^{4/3}} dx$$

input `int(cos(c + d*x)^2/(b/cos(c + d*x))^(4/3), x)`output `int(cos(c + d*x)^2/(b/cos(c + d*x))^(4/3), x)`



### 3.204 $\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx$

3.204.1 Optimal result . . . . .	1188
3.204.2 Mathematica [A] (verified) . . . . .	1188
3.204.3 Rubi [A] (verified) . . . . .	1189
3.204.4 Maple [F] . . . . .	1190
3.204.5 Fricas [F] . . . . .	1190
3.204.6 Sympy [F(-1)] . . . . .	1191
3.204.7 Maxima [F] . . . . .	1191
3.204.8 Giac [F] . . . . .	1191
3.204.9 Mupad [F(-1)] . . . . .	1192

#### 3.204.1 Optimal result

Integrand size = 21, antiderivative size = 81

$$\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{3b \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(-1 - 3m), \frac{1}{6}(5 - 3m), \cos^2(c + dx)\right) \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)}}{d(1 + 3m)\sqrt{\sin^2(c + dx)}}$$

output `3*b*hypergeom([1/2, -1/6-1/2*m], [5/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^m*(b*sec(d*x+c))^(1/3)*sin(d*x+c)/d/(1+3*m)/(sin(d*x+c)^2)^(1/2)`

#### 3.204.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.02

$$\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx = \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{4}{3} + m\right), \frac{1}{2}\left(\frac{10}{3} + m\right), \sec^2(c + dx)\right) \sec^{-1+m}(c + dx)(b \sec(c + dx))^{4/3}}{d\left(\frac{4}{3} + m\right)}$$

input `Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^(4/3),x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (4/3 + m)/2, (10/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/ (d*(4/3 + m))`

**3.204.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{4/3} \sec^m(c + dx) dx \\
 & \quad \downarrow \text{2034} \\
 & \frac{b \sqrt[3]{b \sec(c + dx)} \int \sec^{m+\frac{4}{3}}(c + dx) dx}{\sqrt[3]{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b \sqrt[3]{b \sec(c + dx)} \int \csc\left(c + dx + \frac{\pi}{2}\right)^{m+\frac{4}{3}} dx}{\sqrt[3]{\sec(c + dx)}} \\
 & \quad \downarrow \text{4259} \\
 & b \sqrt[3]{b \sec(c + dx)} \cos^{m+\frac{1}{3}}(c + dx) \sec^m(c + dx) \int \cos^{-m-\frac{4}{3}}(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & b \sqrt[3]{b \sec(c + dx)} \cos^{m+\frac{1}{3}}(c + dx) \sec^m(c + dx) \int \sin\left(c + dx + \frac{\pi}{2}\right)^{-m-\frac{4}{3}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3b \sin(c + dx) \sqrt[3]{b \sec(c + dx)} \sec^m(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(-3m - 1), \frac{1}{6}(5 - 3m), \cos^2(c + dx)\right)}{d(3m + 1) \sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^m*(b*Sec[c + d*x])^(4/3),x]`

output `(3*b*Hypergeometric2F1[1/2, (-1 - 3*m)/6, (5 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^m*(b*Sec[c + d*x])^(1/3)*Sin[c + d*x])/(d*(1 + 3*m)*Sqrt[Sin[c + d*x]^2])`

## 3.204.3.1 Defintions of rubi rules used

rule 2034 `Int[(Fv_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m+n)*Fv, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.204.4 Maple [F]

$$\int \sec(dx+c)^m (b \sec(dx+c))^{\frac{4}{3}} dx$$

input `int(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x)`

output `int(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x)`

## 3.204.5 Fracas [F]

$$\int \sec^m(c+dx)(b \sec(c+dx))^{4/3} dx = \int (b \sec(dx+c))^{\frac{4}{3}} \sec(dx+c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*sec(d*x+c))^(1/3)*b*sec(d*x+c)^m*sec(d*x+c), x)`

**3.204.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**m*(b*sec(d*x+c))**(4/3),x)`output `Timed out`**3.204.7 Maxima [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^m, x)`**3.204.8 Giac [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx = \int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x, algorithm="giac")`output `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^m, x)`

**3.204.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{4/3} \left( \frac{1}{\cos(c + dx)} \right)^m dx$$

input `int((b/cos(c + d*x))^(4/3)*(1/cos(c + d*x))^m,x)`output `int((b/cos(c + d*x))^(4/3)*(1/cos(c + d*x))^m, x)`

### 3.205 $\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx$

3.205.1 Optimal result . . . . .	1193
3.205.2 Mathematica [A] (verified) . . . . .	1193
3.205.3 Rubi [A] (verified) . . . . .	1194
3.205.4 Maple [F] . . . . .	1195
3.205.5 Fricas [F] . . . . .	1195
3.205.6 Sympy [F] . . . . .	1196
3.205.7 Maxima [F] . . . . .	1196
3.205.8 Giac [F] . . . . .	1196
3.205.9 Mupad [F(-1)] . . . . .	1197

#### 3.205.1 Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(1 - 3m), \frac{1}{6}(7 - 3m), \cos^2(c + dx)\right) \sec^{-1+m}(c + dx)(b \sec(c + dx))^{2/3} \sin(c + dx)}{d(1 - 3m)\sqrt{\sin^2(c + dx)}}$$

output `-3*hypergeom([1/2, 1/6-1/2*m], [7/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*(b*sec(d*x+c))^(2/3)*sin(d*x+c)/d/(1-3*m)/(sin(d*x+c)^2)^(1/2)`

#### 3.205.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.01

$$\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx = \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{2}{3} + m\right), \frac{1}{2}\left(\frac{8}{3} + m\right), \sec^2(c + dx)\right) \sec^{-1+m}(c + dx)(b \sec(c + dx))^{2/3}}{d\left(\frac{2}{3} + m\right)}$$

input `Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^(2/3),x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (2/3 + m)/2, (8/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/d*(2/3 + m)`

**3.205.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{2/3} \sec^m(c + dx) dx \\
 & \quad \downarrow \text{2034} \\
 & \frac{(b \sec(c + dx))^{2/3} \int \sec^{m+\frac{2}{3}}(c + dx) dx}{\sec^{\frac{2}{3}}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(b \sec(c + dx))^{2/3} \int \csc(c + dx + \frac{\pi}{2})^{m+\frac{2}{3}} dx}{\sec^{\frac{2}{3}}(c + dx)} \\
 & \quad \downarrow \text{4259} \\
 & (b \sec(c + dx))^{2/3} \cos^{m+\frac{2}{3}}(c + dx) \sec^m(c + dx) \int \cos^{-m-\frac{2}{3}}(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & (b \sec(c + dx))^{2/3} \cos^{m+\frac{2}{3}}(c + dx) \sec^m(c + dx) \int \sin(c + dx + \frac{\pi}{2})^{-m-\frac{2}{3}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(c + dx) (b \sec(c + dx))^{2/3} \sec^{m-1}(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(1 - 3m), \frac{1}{6}(7 - 3m), \cos^2(c + dx)\right)}{d(1 - 3m) \sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^m*(b*Sec[c + d*x])^(2/3),x]`

output `(-3*Hypergeometric2F1[1/2, (1 - 3*m)/6, (7 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(2/3)*Sin[c + d*x])/(d*(1 - 3*m)*Sqrt[Sin[c + d*x]^2])`

## 3.205.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fv_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fv, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_)+(d_)*(x_)])*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.205.4 Maple [F]

$$\int \sec(dx + c)^m (b \sec(dx + c))^{\frac{2}{3}} dx$$

```
input int(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x)
```

```
output int(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x)
```

## 3.205.5 Fracas [F]

$$\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx = \int (b \sec(dx + c))^{\frac{2}{3}} \sec(dx + c)^m dx$$

```
input integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x, algorithm="fricas")
```

```
output integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m, x)
```



**3.205.6 Sympy [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx = \int (b \sec(c + dx))^{2/3} \sec^m(c + dx) dx$$

input `integrate(sec(d*x+c)**m*(b*sec(d*x+c))**(2/3),x)`

output `Integral((b*sec(c + d*x))**(2/3)*sec(c + d*x)**m, x)`

**3.205.7 Maxima [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx = \int (b \sec(dx + c))^{2/3} \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m, x)`

**3.205.8 Giac [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx = \int (b \sec(dx + c))^{2/3} \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m, x)`

**3.205.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{2/3} \left( \frac{1}{\cos(c + dx)} \right)^m dx$$

input `int((b/cos(c + d*x))^(2/3)*(1/cos(c + d*x))^m,x)`output `int((b/cos(c + d*x))^(2/3)*(1/cos(c + d*x))^m, x)`

### 3.206 $\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

3.206.1 Optimal result . . . . .	1198
3.206.2 Mathematica [A] (verified) . . . . .	1198
3.206.3 Rubi [A] (verified) . . . . .	1199
3.206.4 Maple [F] . . . . .	1200
3.206.5 Fricas [F] . . . . .	1200
3.206.6 Sympy [F] . . . . .	1201
3.206.7 Maxima [F] . . . . .	1201
3.206.8 Giac [F] . . . . .	1201
3.206.9 Mupad [F(-1)] . . . . .	1202

#### 3.206.1 Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(2 - 3m), \frac{1}{6}(8 - 3m), \cos^2(c + dx)\right) \sec^{-1+m}(c + dx) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{d(2 - 3m) \sqrt{\sin^2(c + dx)}}$$

```
output -3*hypergeom([1/2, 1/3-1/2*m], [4/3-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*
(b*sec(d*x+c))^(1/3)*sin(d*x+c)/d/(2-3*m)/(sin(d*x+c)^2)^(1/2)
```

#### 3.206.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.01

$$\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{1}{3} + m\right), \frac{1}{2}\left(\frac{7}{3} + m\right), \sec^2(c + dx)\right) \sec^{-1+m}(c + dx) \sqrt[3]{b \sec(c + dx)}}{d\left(\frac{1}{3} + m\right)}$$

```
input Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^(1/3),x]
```

```
output (Csc[c + d*x]*Hypergeometric2F1[1/2, (1/3 + m)/2, (7/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(1/3)*Sqrt[-Tan[c + d*x]^2])/
(d*(1/3 + m))
```

**3.206.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{b \sec(c+dx)} \sec^m(c+dx) dx \\
 & \quad \downarrow \text{2034} \\
 & \frac{\sqrt[3]{b \sec(c+dx)} \int \sec^{m+\frac{1}{3}}(c+dx) dx}{\sqrt[3]{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{b \sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^{m+\frac{1}{3}} dx}{\sqrt[3]{\sec(c+dx)}} \\
 & \quad \downarrow \text{4259} \\
 & \sqrt[3]{b \sec(c+dx)} \cos^{m+\frac{1}{3}}(c+dx) \sec^m(c+dx) \int \cos^{-m-\frac{1}{3}}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt[3]{b \sec(c+dx)} \cos^{m+\frac{1}{3}}(c+dx) \sec^m(c+dx) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{-m-\frac{1}{3}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(c+dx) \sqrt[3]{b \sec(c+dx)} \sec^{m-1}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(2-3m), \frac{1}{6}(8-3m), \cos^2(c+dx)\right)}{d(2-3m) \sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^m*(b*Sec[c + d*x])^(1/3),x]`

output `(-3*Hypergeometric2F1[1/2, (2 - 3*m)/6, (8 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(1/3)*Sin[c + d*x])/(d*(2 - 3*m)*Sqrt[Sin[c + d*x]^2])`

## 3.206.3.1 Defintions of rubi rules used

rule 2034 `Int[(Fv_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m+n)*Fv, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)]^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.206.4 Maple [F]

$$\int \sec(dx+c)^m (b \sec(dx+c))^{\frac{1}{3}} dx$$

input `int(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x)`

output `int(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x)`

## 3.206.5 Fracas [F]

$$\int \sec^m(c+dx) \sqrt[3]{b \sec(c+dx)} dx = \int (b \sec(dx+c))^{\frac{1}{3}} \sec(dx+c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*sec(d*x+c))^(1/3)*sec(d*x+c)^m, x)`

**3.206.6 Sympy [F]**

$$\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \sqrt[3]{b \sec(c + dx)} \sec^m(c + dx) dx$$

input `integrate(sec(d*x+c)**m*(b*sec(d*x+c))**(1/3),x)`

output `Integral((b*sec(c + d*x))**(1/3)*sec(c + d*x)**m, x)`

**3.206.7 Maxima [F]**

$$\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^m, x)`

**3.206.8 Giac [F]**

$$\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^m, x)`

**3.206.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx = \int \left( \frac{b}{\cos(c + dx)} \right)^{1/3} \left( \frac{1}{\cos(c + dx)} \right)^m dx$$

input `int((b/cos(c + d*x))^(1/3)*(1/cos(c + d*x))^m,x)`output `int((b/cos(c + d*x))^(1/3)*(1/cos(c + d*x))^m, x)`

$$3.207 \quad \int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

3.207.1 Optimal result . . . . . 1203  
 3.207.2 Mathematica [A] (verified) . . . . . 1203  
 3.207.3 Rubi [A] (verified) . . . . . 1204  
 3.207.4 Maple [F] . . . . . 1205  
 3.207.5 Fricas [F] . . . . . 1206  
 3.207.6 Sympy [F] . . . . . 1206  
 3.207.7 Maxima [F] . . . . . 1206  
 3.207.8 Giac [F] . . . . . 1207  
 3.207.9 Mupad [F(-1)] . . . . . 1207

### 3.207.1 Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(4-3m), \frac{1}{6}(10-3m), \cos^2(c+dx)\right) \sec^{-1+m}(c+dx) \sin(c+dx)}{d(4-3m) \sqrt[3]{b \sec(c+dx)} \sqrt{\sin^2(c+dx)}}$$

output `-3*hypergeom([1/2, 2/3-1/2*m], [5/3-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*sin(d*x+c)/d/(4-3*m)/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)`

### 3.207.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.01

$$\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \frac{\csc(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{1}{3}+m\right), \frac{1}{2}\left(\frac{5}{3}+m\right), \sec^2(c+dx)\right) \sec^{-1+m}(c+dx) \sqrt{-\tan^2(c+dx)}}{d\left(-\frac{1}{3}+m\right) \sqrt[3]{b \sec(c+dx)}}$$

input `Integrate[Sec[c + d*x]^m/(b*Sec[c + d*x])^(1/3), x]`

---

3.207.  $\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$



output  $(\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[1/2, (-1/3 + m)/2, (5/3 + m)/2, \text{Sec}[c + d*x]^2]*\text{Sec}[c + d*x]^{(-1 + m)}*\text{Sqrt}[-\text{Tan}[c + d*x]^2])/(d*(-1/3 + m)*(b*\text{Sec}[c + d*x])^{(1/3)})$

### 3.207.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^m(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx \\
 & \quad \downarrow \text{2034} \\
 & \frac{\sqrt[3]{\sec(c + dx)} \int \sec^{m-\frac{1}{3}}(c + dx) dx}{\sqrt[3]{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{\sec(c + dx)} \int \csc\left(c + dx + \frac{\pi}{2}\right)^{m-\frac{1}{3}} dx}{\sqrt[3]{b \sec(c + dx)}} \\
 & \quad \downarrow \text{4259} \\
 & \frac{\cos^{m+\frac{2}{3}}(c + dx) \sec^{m+1}(c + dx) \int \cos^{\frac{1}{3}-m}(c + dx) dx}{\sqrt[3]{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cos^{m+\frac{2}{3}}(c + dx) \sec^{m+1}(c + dx) \int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{1}{3}-m} dx}{\sqrt[3]{b \sec(c + dx)}} \\
 & \quad \downarrow \text{3122} \\
 & \frac{3 \sin(c + dx) \sec^{m-1}(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(4 - 3m), \frac{1}{6}(10 - 3m), \cos^2(c + dx)\right)}{d(4 - 3m) \sqrt{\sin^2(c + dx)} \sqrt[3]{b \sec(c + dx)}}
 \end{aligned}$$

input  $\text{Int}[\text{Sec}[c + d*x]^m/(b*\text{Sec}[c + d*x])^{(1/3)}, x]$

---

3.207.  $\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c + dx)}} dx$

output  $(-3\text{Hypergeometric2F1}[1/2, (4 - 3m)/6, (10 - 3m)/6, \text{Cos}[c + dx]^2] \cdot \text{Sec}[c + dx]^{-1+m} \cdot \text{Sin}[c + dx]) / (d \cdot (4 - 3m) \cdot (b \cdot \text{Sec}[c + dx])^{1/3} \cdot \text{Sqrt}[\text{Sin}[c + dx]^2])$

### 3.207.3.1 Defintions of rubi rules used

rule 2034  $\text{Int}[(F x_{.}) \cdot ((a_{.}) \cdot (v_{.}))^{(m_{.})} \cdot ((b_{.}) \cdot (v_{.}))^{(n_{.})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[b^{\text{IntPart}[n]} \cdot ((b \cdot v)^{\text{FracPart}[n]} / (a^{\text{IntPart}[n]} \cdot (a \cdot v)^{\text{FracPart}[n]})) \cdot \text{Int}[(a \cdot v)^{(m+n)} \cdot F x, x], x] /;$   $\text{FreeQ}[\{a, b, m, n\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ !\text{IntegerQ}[n] \ \&\& \ !\text{IntegerQ}[m+n]$

rule 3042  $\text{Int}[u_{.}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3122  $\text{Int}[(b_{.}) \cdot \text{sin}[(c_{.}) + (d_{.}) \cdot (x_{.})]^{(n_{.})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[c + dx] \cdot ((b \cdot \text{Sin}[c + dx])^{(n+1)} / (b \cdot d \cdot (n+1) \cdot \text{Sqrt}[\text{Cos}[c + dx]^2])) \cdot \text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + dx]^2], x] /;$   $\text{FreeQ}[\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[2n]$

rule 4259  $\text{Int}[(\text{csc}[(c_{.}) + (d_{.}) \cdot (x_{.})] \cdot (b_{.}))^{(n_{.})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(b \cdot \text{Csc}[c + dx])^{(n-1)} \cdot ((\text{Sin}[c + dx] / b)^{(n-1)} \cdot \text{Int}[1 / (\text{Sin}[c + dx] / b)^n, x]), x] /;$   $\text{FreeQ}[\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[n]$

### 3.207.4 Maple [F]

$$\int \frac{\sec(dx+c)^m}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input  $\text{int}(\sec(dx+c)^m / (b \cdot \sec(dx+c))^{1/3}, x)$

output  $\text{int}(\sec(dx+c)^m / (b \cdot \sec(dx+c))^{1/3}, x)$

**3.207.5 Fracas [F]**

$$\int \frac{\sec^m(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m/(b*sec(d*x + c)), x)`

**3.207.6 Sympy [F]**

$$\int \frac{\sec^m(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec^m(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

input `integrate(sec(d*x+c)**m/(b*sec(d*x+c))**(1/3),x)`

output `Integral(sec(c + d*x)**m/(b*sec(c + d*x))**(1/3), x)`

**3.207.7 Maxima [F]**

$$\int \frac{\sec^m(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(1/3), x)`

**3.207.8 Giac [F]**

$$\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \int \frac{\sec(dx+c)^m}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

input `integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(1/3), x)`

**3.207.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^m}{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}} dx$$

input `int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(1/3),x)`

output `int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(1/3), x)`

### 3.208 $\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx$

3.208.1 Optimal result . . . . .	1208
3.208.2 Mathematica [A] (verified) . . . . .	1208
3.208.3 Rubi [A] (verified) . . . . .	1209
3.208.4 Maple [F] . . . . .	1210
3.208.5 Fricas [F] . . . . .	1210
3.208.6 Sympy [F] . . . . .	1211
3.208.7 Maxima [F] . . . . .	1211
3.208.8 Giac [F] . . . . .	1211
3.208.9 Mupad [F(-1)] . . . . .	1212

#### 3.208.1 Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(5-3m), \frac{1}{6}(11-3m), \cos^2(c+dx)\right) \sec^{-1+m}(c+dx) \sin(c+dx)}{d(5-3m)(b \sec(c+dx))^{2/3} \sqrt{\sin^2(c+dx)}}$$

output `-3*hypergeom([1/2, 5/6-1/2*m], [11/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*sin(d*x+c)/d/(5-3*m)/(b*sec(d*x+c))^(2/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.208.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.01

$$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx = \frac{\csc(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{2}{3}+m\right), \frac{1}{2}\left(\frac{4}{3}+m\right), \sec^2(c+dx)\right) \sec^{-1+m}(c+dx)}{d\left(-\frac{2}{3}+m\right)(b \sec(c+dx))^{2/3}}$$

input `Integrate[Sec[c + d*x]^m/(b*Sec[c + d*x])^(2/3),x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (-2/3 + m)/2, (4/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sqrt[-Tan[c + d*x]^2])/(d*(-2/3 + m)*(b*Sec[c + d*x])^(2/3))`

### 3.208.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx \\
 & \quad \downarrow \text{2034} \\
 & \frac{\sec^{\frac{2}{3}}(c+dx) \int \sec^{m-\frac{2}{3}}(c+dx) dx}{(b \sec(c+dx))^{2/3}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sec^{\frac{2}{3}}(c+dx) \int \csc(c+dx+\frac{\pi}{2})^{m-\frac{2}{3}} dx}{(b \sec(c+dx))^{2/3}} \\
 & \quad \downarrow \text{4259} \\
 & \frac{\cos^{m+\frac{1}{3}}(c+dx) \sec^{m+1}(c+dx) \int \cos^{\frac{2}{3}-m}(c+dx) dx}{(b \sec(c+dx))^{2/3}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cos^{m+\frac{1}{3}}(c+dx) \sec^{m+1}(c+dx) \int \sin(c+dx+\frac{\pi}{2})^{\frac{2}{3}-m} dx}{(b \sec(c+dx))^{2/3}} \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3 \sin(c+dx) \sec^{m-1}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(5-3m), \frac{1}{6}(11-3m), \cos^2(c+dx)\right)}{d(5-3m) \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{2/3}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^m/(b*Sec[c + d*x])^(2/3),x]`

output `(-3*Hypergeometric2F1[1/2, (5 - 3*m)/6, (11 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sin[c + d*x])/(d*(5 - 3*m)*(b*Sec[c + d*x])^(2/3)*Sqrt[Sin[c + d*x]^2])`

## 3.208.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fx_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.208.4 Maple [F]

$$\int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{2}{3}}} dx$$

```
input int(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3),x)
```

```
output int(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3),x)
```

## 3.208.5 Fracas [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{2}{3}}} dx$$

```
input integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3),x, algorithm="fracas")
```

output `integral((b*sec(d*x + c))^(1/3)*sec(d*x + c)^m/(b*sec(d*x + c)), x)`

### 3.208.6 Sympy [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{2/3}} dx = \int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{2/3}} dx$$

input `integrate(sec(d*x+c)**m/(b*sec(d*x+c))**(2/3), x)`

output `Integral(sec(c + d*x)**m/(b*sec(c + d*x))**(2/3), x)`

### 3.208.7 Maxima [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{2/3}} dx$$

input `integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3), x, algorithm="maxima")`

output `integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(2/3), x)`

### 3.208.8 Giac [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{2/3}} dx$$

input `integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3), x, algorithm="giac")`

output `integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(2/3), x)`



**3.208.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^m}{\left(\frac{b}{\cos(c+dx)}\right)^{2/3}} dx$$

input `int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(2/3),x)`output `int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(2/3), x)`

### 3.209 $\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx$

3.209.1 Optimal result . . . . .	1213
3.209.2 Mathematica [A] (verified) . . . . .	1213
3.209.3 Rubi [A] (verified) . . . . .	1214
3.209.4 Maple [F] . . . . .	1215
3.209.5 Fricas [F] . . . . .	1215
3.209.6 Sympy [F] . . . . .	1216
3.209.7 Maxima [F] . . . . .	1216
3.209.8 Giac [F] . . . . .	1216
3.209.9 Mupad [F(-1)] . . . . .	1217

#### 3.209.1 Optimal result

Integrand size = 21, antiderivative size = 85

$$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(7-3m), \frac{1}{6}(13-3m), \cos^2(c+dx)\right) \sec^{-2+m}(c+dx) \sin(c+dx)}{bd(7-3m) \sqrt[3]{b \sec(c+dx)} \sqrt{\sin^2(c+dx)}}$$

output `-3*hypergeom([1/2, 7/6-1/2*m], [13/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-2+m)*sin(d*x+c)/b/d/(7-3*m)/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)`

#### 3.209.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.98

$$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \frac{\csc(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{4}{3}+m\right), \frac{1}{2}\left(\frac{2}{3}+m\right), \sec^2(c+dx)\right) \sec^{-1+m}(c+dx)}{d\left(-\frac{4}{3}+m\right)(b \sec(c+dx))^{4/3}}$$

input `Integrate[Sec[c + d*x]^m/(b*Sec[c + d*x])^(4/3),x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (-4/3 + m)/2, (2/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sqrt[-Tan[c + d*x]^2])/(d*(-4/3 + m)*(b*Sec[c + d*x])^(4/3))`

**3.209.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx \\
 & \quad \downarrow \text{2034} \\
 & \frac{\sqrt[3]{\sec(c+dx)} \int \sec^{m-\frac{4}{3}}(c+dx) dx}{b \sqrt[3]{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{\sec(c+dx)} \int \csc\left(c+dx+\frac{\pi}{2}\right)^{m-\frac{4}{3}} dx}{b \sqrt[3]{b \sec(c+dx)}} \\
 & \quad \downarrow \text{4259} \\
 & \frac{\cos^{m+\frac{2}{3}}(c+dx) \sec^{m+1}(c+dx) \int \cos^{\frac{4}{3}-m}(c+dx) dx}{b \sqrt[3]{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cos^{m+\frac{2}{3}}(c+dx) \sec^{m+1}(c+dx) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{4}{3}-m} dx}{b \sqrt[3]{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3122} \\
 & -\frac{3 \sin(c+dx) \sec^{m-2}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(7-3m), \frac{1}{6}(13-3m), \cos^2(c+dx)\right)}{bd(7-3m) \sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^m/(b*Sec[c + d*x])^(4/3),x]`

output `(-3*Hypergeometric2F1[1/2, (7 - 3*m)/6, (13 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-2 + m)*Sin[c + d*x])/(b*d*(7 - 3*m)*(b*Sec[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])`

## 3.209.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.209.4 Maple [F]

$$\int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

```
input int(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3),x)
```

```
output int(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3),x)
```

## 3.209.5 Fracas [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

```
input integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3),x, algorithm="fracas")
```

output `integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m/(b^2*sec(d*x + c)^2), x)`

### 3.209.6 Sympy [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{4/3}} dx$$

input `integrate(sec(d*x+c)**m/(b*sec(d*x+c))**(4/3), x)`

output `Integral(sec(c + d*x)**m/(b*sec(c + d*x))**(4/3), x)`

### 3.209.7 Maxima [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3), x, algorithm="maxima")`

output `integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(4/3), x)`

### 3.209.8 Giac [F]

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{4/3}} dx$$

input `integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3), x, algorithm="giac")`

output `integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(4/3), x)`

**3.209.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^m}{\left(\frac{b}{\cos(c+dx)}\right)^{4/3}} dx$$

input `int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(4/3),x)`output `int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(4/3), x)`

### 3.210 $\int \sec^m(c + dx)(b \sec(c + dx))^n dx$

3.210.1 Optimal result . . . . .	1218
3.210.2 Mathematica [A] (verified) . . . . .	1218
3.210.3 Rubi [A] (verified) . . . . .	1219
3.210.4 Maple [F] . . . . .	1220
3.210.5 Fricas [F] . . . . .	1220
3.210.6 Sympy [F] . . . . .	1221
3.210.7 Maxima [F] . . . . .	1221
3.210.8 Giac [F] . . . . .	1221
3.210.9 Mupad [F(-1)] . . . . .	1222

#### 3.210.1 Optimal result

Integrand size = 19, antiderivative size = 89

$$\int \sec^m(c + dx)(b \sec(c + dx))^n dx = \frac{\text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 - m - n), \frac{1}{2}(3 - m - n), \cos^2(c + dx)\right) \sec^{-1+m}(c + dx)(b \sec(c + dx))^n \sin(c + dx)}{d(1 - m - n)\sqrt{\sin^2(c + dx)}}$$

output `-hypergeom([1/2, 1/2-1/2*m-1/2*n], [3/2-1/2*m-1/2*n], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(1-m-n)/(sin(d*x+c)^2)^(1/2)`

#### 3.210.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.85

$$\int \sec^m(c + dx)(b \sec(c + dx))^n dx = \frac{\csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+n}{2}, \frac{1}{2}(2 + m + n), \sec^2(c + dx)\right) \sec^{-1+m}(c + dx)(b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d(m + n)}$$

input `Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^n,x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (m + n)/2, (2 + m + n)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^n*sqrt[-Tan[c + d*x]^2])/d*(m + n)`

**3.210.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^m(c+dx)(b \sec(c+dx))^n dx \\
 & \quad \downarrow \text{2034} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \sec^{m+n}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \csc\left(c+dx+\frac{\pi}{2}\right)^{m+n} dx \\
 & \quad \downarrow \text{4259} \\
 & \sec^m(c+dx)(b \sec(c+dx))^n \cos^{m+n}(c+dx) \int \cos^{-m-n}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sec^m(c+dx)(b \sec(c+dx))^n \cos^{m+n}(c+dx) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{-m-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{\sin(c+dx) \sec^{m-1}(c+dx)(b \sec(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-m-n+1), \frac{1}{2}(-m-n+3), \cos^2(c+dx)\right)}{d(-m-n+1)\sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^m*(b*Sec[c + d*x])^n,x]`

output `-((Hypergeometric2F1[1/2, (1 - m - n)/2, (3 - m - n)/2, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(1 - m - n)*Sqrt[Sin[c + d*x]^2]))`



## 3.210.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fv_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fv, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.210.4 Maple [F]

$$\int \sec(dx + c)^m (b \sec(dx + c))^n dx$$

```
input int(sec(d*x+c)^m*(b*sec(d*x+c))^n,x)
```

```
output int(sec(d*x+c)^m*(b*sec(d*x+c))^n,x)
```

## 3.210.5 Fracas [F]

$$\int \sec^m(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^m dx$$

```
input integrate(sec(d*x+c)^m*(b*sec(d*x+c))^n,x, algorithm="fracas")
```

```
output integral((b*sec(d*x + c))^n*sec(d*x + c)^m, x)
```

**3.210.6 Sympy [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n \sec^m(c + dx) dx$$

input `integrate(sec(d*x+c)**m*(b*sec(d*x+c))**n,x)`

output `Integral((b*sec(c + d*x))**n*sec(c + d*x)**m, x)`

**3.210.7 Maxima [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^m, x)`

**3.210.8 Giac [F]**

$$\int \sec^m(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^m dx$$

input `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^m, x)`

**3.210.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^m(c + dx)(b \sec(c + dx))^n dx = \int \left( \frac{b}{\cos(c + dx)} \right)^n \left( \frac{1}{\cos(c + dx)} \right)^m dx$$

input `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^m,x)`output `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^m, x)`

### 3.211 $\int \sec^2(c + dx)(b \sec(c + dx))^n dx$

3.211.1 Optimal result . . . . .	1223
3.211.2 Mathematica [A] (verified) . . . . .	1223
3.211.3 Rubi [A] (verified) . . . . .	1224
3.211.4 Maple [F] . . . . .	1225
3.211.5 Fricas [F] . . . . .	1225
3.211.6 Sympy [F] . . . . .	1226
3.211.7 Maxima [F] . . . . .	1226
3.211.8 Giac [F] . . . . .	1226
3.211.9 Mupad [F(-1)] . . . . .	1227

#### 3.211.1 Optimal result

Integrand size = 19, antiderivative size = 72

$$\int \sec^2(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-1 - n), \frac{1-n}{2}, \cos^2(c + dx)\right) (b \sec(c + dx))^{1+n} \sin(c + dx)}{bd(1 + n)\sqrt{\sin^2(c + dx)}}$$

output `hypergeom([1/2, -1/2-1/2*n], [1/2-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^(1+n)*sin(d*x+c)/b/d/(1+n)/(sin(d*x+c)^2)^(1/2)`

#### 3.211.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.99

$$\int \sec^2(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+n}{2}, \frac{4+n}{2}, \sec^2(c + dx)\right) \sec(c + dx)(b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d(2 + n)}$$

input `Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^n,x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (2 + n)/2, (4 + n)/2, Sec[c + d*x]^2]*Sec[c + d*x]*(b*Sec[c + d*x])^n*sqrt[-Tan[c + d*x]^2])/(d*(2 + n))`

**3.211.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c + dx)(b \sec(c + dx))^n dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c + dx))^{n+2} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{n+2} dx}{b^2} \\
 & \quad \downarrow \text{4259} \\
 & \quad \frac{\left(\frac{\cos(c+dx)}{b}\right)^n (b \sec(c + dx))^n \int \left(\frac{\cos(c+dx)}{b}\right)^{-n-2} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\left(\frac{\cos(c+dx)}{b}\right)^n (b \sec(c + dx))^n \int \left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{-n-2} dx}{b^2} \\
 & \quad \downarrow \text{3122} \\
 & \quad \frac{\sin(c + dx)(b \sec(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-n - 1), \frac{1-n}{2}, \cos^2(c + dx)\right)}{bd(n + 1)\sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^n,x]`

output `(Hypergeometric2F1[1/2, (-1 - n)/2, (1 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(1 + n)*Sin[c + d*x]/(b*d*(1 + n)*Sqrt[Sin[c + d*x]^2])`

## 3.211.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.211.4 Maple [F]

$$\int \sec(dx + c)^2 (b \sec(dx + c))^n dx$$

input `int(sec(d*x+c)^2*(b*sec(d*x+c))^n,x)`

output `int(sec(d*x+c)^2*(b*sec(d*x+c))^n,x)`

## 3.211.5 Fracas [F]

$$\int \sec^2(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^n*sec(d*x + c)^2, x)`

**3.211.6 Sympy [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**n,x)`

output `Integral((b*sec(c + d*x))**n*sec(c + d*x)**2, x)`

**3.211.7 Maxima [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^2, x)`

**3.211.8 Giac [F]**

$$\int \sec^2(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^2 dx$$

input `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^2, x)`

**3.211.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^2(c + dx)(b \sec(c + dx))^n dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\cos(c + dx)^2} dx$$

input `int((b/cos(c + d*x))^n/cos(c + d*x)^2,x)`output `int((b/cos(c + d*x))^n/cos(c + d*x)^2, x)`



### 3.212 $\int \sec(c + dx)(b \sec(c + dx))^n dx$

3.212.1 Optimal result . . . . .	1228
3.212.2 Mathematica [A] (verified) . . . . .	1228
3.212.3 Rubi [A] (verified) . . . . .	1229
3.212.4 Maple [F] . . . . .	1230
3.212.5 Fracas [F] . . . . .	1230
3.212.6 Sympy [F] . . . . .	1231
3.212.7 Maxima [F] . . . . .	1231
3.212.8 Giac [F] . . . . .	1231
3.212.9 Mupad [F(-1)] . . . . .	1232

#### 3.212.1 Optimal result

Integrand size = 17, antiderivative size = 61

$$\int \sec(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\text{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{n}{2}, \frac{2-n}{2}, \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{dn \sqrt{\sin^2(c + dx)}}$$

output `hypergeom([1/2, -1/2*n], [1-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/n/(sin(d*x+c)^2)^(1/2)`

#### 3.212.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.07

$$\int \sec(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d(1 + n)}$$

input `Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^n,x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(1 + n))`

**3.212.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {2030, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c + dx)(b \sec(c + dx))^n dx \\
 & \quad \downarrow \text{2030} \\
 & \quad \frac{\int (b \sec(c + dx))^{n+1} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\int (b \csc(c + dx + \frac{\pi}{2}))^{n+1} dx}{b} \\
 & \quad \downarrow \text{4259} \\
 & \quad \frac{\left(\frac{\cos(c+dx)}{b}\right)^n (b \sec(c + dx))^n \int \left(\frac{\cos(c+dx)}{b}\right)^{-n-1} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{\left(\frac{\cos(c+dx)}{b}\right)^n (b \sec(c + dx))^n \int \left(\frac{\sin(c+dx+\frac{\pi}{2})}{b}\right)^{-n-1} dx}{b} \\
 & \quad \downarrow \text{3122} \\
 & \quad \frac{\sin(c + dx)(b \sec(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{n}{2}, \frac{2-n}{2}, \cos^2(c + dx)\right)}{dn \sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]*(b*Sec[c + d*x])^n,x]`

output `(Hypergeometric2F1[1/2, -1/2*n, (2 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*n*Sqrt[Sin[c + d*x]^2])`

## 3.212.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.212.4 Maple [F]

$$\int \sec(dx+c)(b \sec(dx+c))^n dx$$

input `int(sec(d*x+c)*(b*sec(d*x+c))^n,x)`

output `int(sec(d*x+c)*(b*sec(d*x+c))^n,x)`

## 3.212.5 Fracas [F]

$$\int \sec(c+dx)(b \sec(c+dx))^n dx = \int (b \sec(dx+c))^n \sec(dx+c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^n*sec(d*x+c), x)`

**3.212.6 Sympy [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n \sec(c + dx) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))**n,x)`

output `Integral((b*sec(c + d*x))**n*sec(c + d*x), x)`

**3.212.7 Maxima [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c), x)`

**3.212.8 Giac [F]**

$$\int \sec(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c) dx$$

input `integrate(sec(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c), x)`

**3.212.9 Mupad [F(-1)]**

Timed out.

$$\int \sec(c + dx)(b \sec(c + dx))^n dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\cos(c + dx)} dx$$

input `int((b/cos(c + d*x))^n/cos(c + d*x),x)`output `int((b/cos(c + d*x))^n/cos(c + d*x), x)`

### 3.213 $\int (b \sec(c + dx))^n dx$

3.213.1 Optimal result . . . . .	1233
3.213.2 Mathematica [A] (verified) . . . . .	1233
3.213.3 Rubi [A] (verified) . . . . .	1234
3.213.4 Maple [F] . . . . .	1235
3.213.5 Fricas [F] . . . . .	1235
3.213.6 Sympy [F] . . . . .	1236
3.213.7 Maxima [F] . . . . .	1236
3.213.8 Giac [F] . . . . .	1236
3.213.9 Mupad [F(-1)] . . . . .	1237

#### 3.213.1 Optimal result

Integrand size = 10, antiderivative size = 73

$$\int (b \sec(c + dx))^n dx = -\frac{b \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \cos^2(c + dx)\right) (b \sec(c + dx))^{-1+n} \sin(c + dx)}{d(1-n)\sqrt{\sin^2(c + dx)}}$$

output `-b*hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^-1+n)*sin(d*x+c)/d/(1-n)/(sin(d*x+c)^2)^(1/2)`

#### 3.213.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.84

$$\int (b \sec(c + dx))^n dx = \frac{\cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n}{2}, \frac{2+n}{2}, \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{dn}$$

input `Integrate[(b*Sec[c + d*x])^n,x]`

output `(Cot[c + d*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*n)`

**3.213.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^n dx \\
 & \quad \downarrow \text{4259} \\
 & \left( \frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \int \left( \frac{\cos(c + dx)}{b} \right)^{-n} dx \\
 & \quad \downarrow \text{3042} \\
 & \left( \frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \int \left( \frac{\sin\left(c + dx + \frac{\pi}{2}\right)}{b} \right)^{-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{b \sin(c + dx) (b \sec(c + dx))^{n-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \cos^2(c + dx)\right)}{d(1-n)\sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^n,x]`

output `-((b*Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(-1 + n)*Sin[c + d*x])/(d*(1 - n)*Sqrt[Sin[c + d*x]^2]))`

## 3.213.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.213.4 Maple [F]

$$\int (b \sec(dx + c))^n dx$$

input `int((b*sec(d*x+c))^n,x)`

output `int((b*sec(d*x+c))^n,x)`

## 3.213.5 Fracas [F]

$$\int (b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n dx$$

input `integrate((b*sec(d*x+c))^n,x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^n, x)`



**3.213.6 Sympy [F]**

$$\int (b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n dx$$

input `integrate((b*sec(d*x+c))**n,x)`

output `Integral((b*sec(c + d*x))**n, x)`

**3.213.7 Maxima [F]**

$$\int (b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n dx$$

input `integrate((b*sec(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n, x)`

**3.213.8 Giac [F]**

$$\int (b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n dx$$

input `integrate((b*sec(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n, x)`

**3.213.9 Mupad [F(-1)]**

Timed out.

$$\int (b \sec(c + dx))^n dx = \int \left( \frac{b}{\cos(c + dx)} \right)^n dx$$

input `int((b/cos(c + d*x))^n,x)`output `int((b/cos(c + d*x))^n, x)`

### 3.214 $\int \cos(c + dx)(b \sec(c + dx))^n dx$

3.214.1 Optimal result . . . . .	1238
3.214.2 Mathematica [A] (verified) . . . . .	1238
3.214.3 Rubi [A] (verified) . . . . .	1239
3.214.4 Maple [F] . . . . .	1240
3.214.5 Fracas [F] . . . . .	1240
3.214.6 Sympy [F] . . . . .	1241
3.214.7 Maxima [F] . . . . .	1241
3.214.8 Giac [F] . . . . .	1241
3.214.9 Mupad [F(-1)] . . . . .	1242

#### 3.214.1 Optimal result

Integrand size = 17, antiderivative size = 75

$$\int \cos(c + dx)(b \sec(c + dx))^n dx$$

$$= -\frac{b^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2-n}{2}, \frac{4-n}{2}, \cos^2(c + dx)\right) (b \sec(c + dx))^{-2+n} \sin(c + dx)}{d(2-n)\sqrt{\sin^2(c + dx)}}$$

output `-b^2*hypergeom([1/2, 1-1/2*n], [2-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^-2+n)*sin(d*x+c)/d/(2-n)/(sin(d*x+c)^2)^(1/2)`

#### 3.214.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.91

$$\int \cos(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{b \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-1 + n), \frac{1+n}{2}, \sec^2(c + dx)\right) (b \sec(c + dx))^{-1+n} \sqrt{-\tan^2(c + dx)}}{d(-1 + n)}$$

input `Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^n,x]`

output `(b*Cot[c + d*x]*Hypergeometric2F1[1/2, (-1 + n)/2, (1 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^-1 + n)*Sqrt[-Tan[c + d*x]^2]/(d*(-1 + n))`

**3.214.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx)(b \sec(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c + dx + \frac{\pi}{2}))^n}{\csc(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{2030} \\
 & b \int \left( b \csc\left(\frac{1}{2}(2c + \pi) + dx\right) \right)^{n-1} dx \\
 & \quad \downarrow \text{4259} \\
 & b \left( \frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \int \left( \frac{\cos(c + dx)}{b} \right)^{1-n} dx \\
 & \quad \downarrow \text{3042} \\
 & b \left( \frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \int \left( \frac{\sin(c + dx + \frac{\pi}{2})}{b} \right)^{1-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{b^2 \sin(c + dx)(b \sec(c + dx))^{n-2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2-n}{2}, \frac{4-n}{2}, \cos^2(c + dx)\right)}{d(2-n)\sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[Cos[c + d*x]*(b*Sec[c + d*x])^n,x]`

output `-((b^2*Hypergeometric2F1[1/2, (2 - n)/2, (4 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(-2 + n)*Sin[c + d*x])/(d*(2 - n)*Sqrt[Sin[c + d*x]^2]))`

## 3.214.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.214.4 Maple [F]

$$\int \cos(dx + c) (b \sec(dx + c))^n dx$$

input `int(cos(d*x+c)*(b*sec(d*x+c))^n,x)`

output `int(cos(d*x+c)*(b*sec(d*x+c))^n,x)`

## 3.214.5 Fracas [F]

$$\int \cos(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="fracas")`

output `integral((b*sec(d*x + c))^n*cos(d*x + c), x)`

**3.214.6 Sympy [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n \cos(c + dx) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))**n,x)`

output `Integral((b*sec(c + d*x))**n*cos(c + d*x), x)`

**3.214.7 Maxima [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n*cos(d*x + c), x)`

**3.214.8 Giac [F]**

$$\int \cos(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n*cos(d*x + c), x)`

**3.214.9 Mupad [F(-1)]**

Timed out.

$$\int \cos(c + dx)(b \sec(c + dx))^n dx = \int \cos(c + dx) \left( \frac{b}{\cos(c + dx)} \right)^n dx$$

input `int(cos(c + d*x)*(b/cos(c + d*x))^n,x)`output `int(cos(c + d*x)*(b/cos(c + d*x))^n, x)`

### 3.215 $\int \cos^2(c + dx)(b \sec(c + dx))^n dx$

3.215.1 Optimal result . . . . .	1243
3.215.2 Mathematica [A] (verified) . . . . .	1243
3.215.3 Rubi [A] (verified) . . . . .	1244
3.215.4 Maple [F] . . . . .	1245
3.215.5 Fricas [F] . . . . .	1245
3.215.6 Sympy [F] . . . . .	1246
3.215.7 Maxima [F] . . . . .	1246
3.215.8 Giac [F] . . . . .	1246
3.215.9 Mupad [F(-1)] . . . . .	1247

#### 3.215.1 Optimal result

Integrand size = 19, antiderivative size = 75

$$\int \cos^2(c + dx)(b \sec(c + dx))^n dx$$

$$= -\frac{b^3 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3-n}{2}, \frac{5-n}{2}, \cos^2(c + dx)\right) (b \sec(c + dx))^{-3+n} \sin(c + dx)}{d(3-n)\sqrt{\sin^2(c + dx)}}$$

output `-b^3*hypergeom([1/2, 3/2-1/2*n], [5/2-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^(  
-3+n)*sin(d*x+c)/d/(3-n)/(sin(d*x+c)^2)^(1/2)`

#### 3.215.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.95

$$\int \cos^2(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\cos^2(c + dx) \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-2 + n), \frac{n}{2}, \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d(-2 + n)}$$

input `Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^n,x]`

output `(Cos[c + d*x]^2*Cot[c + d*x]*Hypergeometric2F1[1/2, (-2 + n)/2, n/2, Sec[c  
+ d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(-2 + n))`



**3.215.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c + dx)(b \sec(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c + dx + \frac{\pi}{2}))^n}{\csc(c + dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{2030} \\
 & b^2 \int \left( b \csc\left(\frac{1}{2}(2c + \pi) + dx\right) \right)^{n-2} dx \\
 & \quad \downarrow \text{4259} \\
 & b^2 \left( \frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \int \left( \frac{\cos(c + dx)}{b} \right)^{2-n} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 \left( \frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \int \left( \frac{\sin(c + dx + \frac{\pi}{2})}{b} \right)^{2-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{b^3 \sin(c + dx)(b \sec(c + dx))^{n-3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3-n}{2}, \frac{5-n}{2}, \cos^2(c + dx)\right)}{d(3-n)\sqrt{\sin^2(c + dx)}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^n,x]`

output `-((b^3*Hypergeometric2F1[1/2, (3 - n)/2, (5 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(-3 + n)*Sin[c + d*x])/(d*(3 - n)*Sqrt[Sin[c + d*x]^2]))`

## 3.215.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.215.4 Maple [F]

$$\int \cos(dx+c)^2 (b \sec(dx+c))^n dx$$

input `int(cos(d*x+c)^2*(b*sec(d*x+c))^n,x)`

output `int(cos(d*x+c)^2*(b*sec(d*x+c))^n,x)`

## 3.215.5 Fracas [F]

$$\int \cos^2(c+dx)(b \sec(c+dx))^n dx = \int (b \sec(dx+c))^n \cos(dx+c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^n*cos(d*x+c)^2, x)`

**3.215.6 Sympy [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n \cos^2(c + dx) dx$$

input `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**n,x)`

output `Integral((b*sec(c + d*x))**n*cos(c + d*x)**2, x)`

**3.215.7 Maxima [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n*cos(d*x + c)^2, x)`

**3.215.8 Giac [F]**

$$\int \cos^2(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n*cos(d*x + c)^2, x)`

**3.215.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^2(c + dx)(b \sec(c + dx))^n dx = \int \cos(c + dx)^2 \left( \frac{b}{\cos(c + dx)} \right)^n dx$$

input `int(cos(c + d*x)^2*(b/cos(c + d*x))^n,x)`output `int(cos(c + d*x)^2*(b/cos(c + d*x))^n, x)`

### 3.216 $\int \cos^3(c + dx)(b \sec(c + dx))^n dx$

3.216.1 Optimal result . . . . .	1248
3.216.2 Mathematica [A] (verified) . . . . .	1248
3.216.3 Rubi [A] (verified) . . . . .	1249
3.216.4 Maple [F] . . . . .	1250
3.216.5 Fricas [F] . . . . .	1250
3.216.6 Sympy [F(-1)] . . . . .	1251
3.216.7 Maxima [F] . . . . .	1251
3.216.8 Giac [F] . . . . .	1251
3.216.9 Mupad [F(-1)] . . . . .	1252

#### 3.216.1 Optimal result

Integrand size = 19, antiderivative size = 75

$$\int \cos^3(c + dx)(b \sec(c + dx))^n dx$$

$$= -\frac{b^4 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4-n}{2}, \frac{6-n}{2}, \cos^2(c + dx)\right) (b \sec(c + dx))^{-4+n} \sin(c + dx)}{d(4-n)\sqrt{\sin^2(c + dx)}}$$

output `-b^4*hypergeom([1/2, 2-1/2*n], [3-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^-4+n)*sin(d*x+c)/d/(4-n)/(sin(d*x+c)^2)^(1/2)`

#### 3.216.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.97

$$\int \cos^3(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\cos^3(c + dx) \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-3 + n), \frac{1}{2}(-1 + n), \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-}}{d(-3 + n)}$$

input `Integrate[Cos[c + d*x]^3*(b*Sec[c + d*x])^n,x]`

output `(Cos[c + d*x]^3*Cot[c + d*x]*Hypergeometric2F1[1/2, (-3 + n)/2, (-1 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(-3 + n))`

**3.216.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 2030, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c+dx)(b \sec(c+dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(c+dx+\frac{\pi}{2}))^n}{\csc(c+dx+\frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{2030} \\
 & b^3 \int \left( b \csc\left(\frac{1}{2}(2c+\pi)+dx\right) \right)^{n-3} dx \\
 & \quad \downarrow \text{4259} \\
 & b^3 \left( \frac{\cos(c+dx)}{b} \right)^n (b \sec(c+dx))^n \int \left( \frac{\cos(c+dx)}{b} \right)^{3-n} dx \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left( \frac{\cos(c+dx)}{b} \right)^n (b \sec(c+dx))^n \int \left( \frac{\sin(c+dx+\frac{\pi}{2})}{b} \right)^{3-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{b^4 \sin(c+dx)(b \sec(c+dx))^{n-4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4-n}{2}, \frac{6-n}{2}, \cos^2(c+dx)\right)}{d(4-n)\sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Cos[c + d*x]^3*(b*Sec[c + d*x])^n,x]`

output `-((b^4*Hypergeometric2F1[1/2, (4 - n)/2, (6 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(-4 + n)*Sin[c + d*x])/(d*(4 - n)*Sqrt[Sin[c + d*x]^2]))`

## 3.216.3.1 Defintions of rubi rules used

rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.216.4 Maple [F]

$$\int \cos(dx+c)^3 (b \sec(dx+c))^n dx$$

input `int(cos(d*x+c)^3*(b*sec(d*x+c))^n,x)`

output `int(cos(d*x+c)^3*(b*sec(d*x+c))^n,x)`

## 3.216.5 Fracas [F]

$$\int \cos^3(c+dx)(b \sec(c+dx))^n dx = \int (b \sec(dx+c))^n \cos(dx+c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^n,x, algorithm="fracas")`

output `integral((b*sec(d*x+c))^n*cos(d*x+c)^3, x)`

**3.216.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^3(c + dx)(b \sec(c + dx))^n dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(b*sec(d*x+c))**n,x)`output `Timed out`**3.216.7 Maxima [F]**

$$\int \cos^3(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^n,x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^n*cos(d*x + c)^3, x)`**3.216.8 Giac [F]**

$$\int \cos^3(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \cos(dx + c)^3 dx$$

input `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^n,x, algorithm="giac")`output `integrate((b*sec(d*x + c))^n*cos(d*x + c)^3, x)`



**3.216.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^3(c + dx)(b \sec(c + dx))^n dx = \int \cos(c + dx)^3 \left( \frac{b}{\cos(c + dx)} \right)^n dx$$

input `int(cos(c + d*x)^3*(b/cos(c + d*x))^n,x)`output `int(cos(c + d*x)^3*(b/cos(c + d*x))^n, x)`

### 3.217 $\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx$

3.217.1 Optimal result . . . . .	1253
3.217.2 Mathematica [A] (verified) . . . . .	1253
3.217.3 Rubi [A] (verified) . . . . .	1254
3.217.4 Maple [F] . . . . .	1255
3.217.5 Fricas [F] . . . . .	1255
3.217.6 Sympy [F(-1)] . . . . .	1256
3.217.7 Maxima [F] . . . . .	1256
3.217.8 Giac [F] . . . . .	1256
3.217.9 Mupad [F(-1)] . . . . .	1257

#### 3.217.1 Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-3 - 2n), \frac{1}{4}(1 - 2n), \cos^2(c + dx)\right) \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n \sin(c + dx)}{d(3 + 2n)\sqrt{\sin^2(c + dx)}}$$

```
output 2*hypergeom([1/2, -3/4-1/2*n], [1/4-1/2*n], cos(d*x+c)^2)*sec(d*x+c)^(3/2)*(
b*sec(d*x+c))^n*sin(d*x+c)/d/(3+2*n)/(sin(d*x+c)^2)^(1/2)
```

#### 3.217.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{5}{2} + n\right), \frac{1}{2}\left(\frac{9}{2} + n\right), \sec^2(c + dx)\right) \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d\left(\frac{5}{2} + n\right)}$$

```
input Integrate[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^n,x]
```

```
output (Csc[c + d*x]*Hypergeometric2F1[1/2, (5/2 + n)/2, (9/2 + n)/2, Sec[c + d*x]
]^2)*Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2]/(d*(5/2
+ n))
```

**3.217.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^n dx \\
 & \quad \downarrow \text{2034} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \sec^{n+\frac{5}{2}}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \csc\left(c+dx+\frac{\pi}{2}\right)^{n+\frac{5}{2}} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt{\sec(c+dx)} \cos^{n+\frac{1}{2}}(c+dx)(b \sec(c+dx))^n \int \cos^{-n-\frac{5}{2}}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\sec(c+dx)} \cos^{n+\frac{1}{2}}(c+dx)(b \sec(c+dx))^n \int \sin\left(c+dx+\frac{\pi}{2}\right)^{-n-\frac{5}{2}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-2n-3), \frac{1}{4}(1-2n), \cos^2(c+dx)\right)}{d(2n+3)\sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^n,x]`

output `(2*Hypergeometric2F1[1/2, (-3 - 2*n)/4, (1 - 2*n)/4, Cos[c + d*x]^2]*Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(3 + 2*n)*Sqrt[Sin[c + d*x]^2])`

## 3.217.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.217.4 Maple [F]

$$\int \sec(dx + c)^{\frac{5}{2}} (b \sec(dx + c))^n dx$$

```
input int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x)
```

```
output int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x)
```

## 3.217.5 Fracas [F]

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^{\frac{5}{2}} dx$$

```
input integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x, algorithm="fricas")
```

```
output integral((b*sec(d*x + c))^n*sec(d*x + c)^(5/2), x)
```

---


$$3.217. \quad \int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx$$

**3.217.6 Sympy [F(-1)]**

Timed out.

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx = \text{Timed out}$$

input `integrate(sec(d*x+c)**(5/2)*(b*sec(d*x+c))**n,x)`output `Timed out`**3.217.7 Maxima [F]**

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^{\frac{5}{2}} dx$$

input `integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x, algorithm="maxima")`output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^(5/2), x)`**3.217.8 Giac [F]**

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^{\frac{5}{2}} dx$$

input `integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x, algorithm="giac")`output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^(5/2), x)`

**3.217.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx = \int \left( \frac{b}{\cos(c + dx)} \right)^n \left( \frac{1}{\cos(c + dx)} \right)^{5/2} dx$$

input `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(5/2),x)`output `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(5/2), x)`

### 3.218 $\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx$

3.218.1 Optimal result . . . . .	1258
3.218.2 Mathematica [A] (verified) . . . . .	1258
3.218.3 Rubi [A] (verified) . . . . .	1259
3.218.4 Maple [F] . . . . .	1260
3.218.5 Fricas [F] . . . . .	1260
3.218.6 Sympy [F] . . . . .	1261
3.218.7 Maxima [F] . . . . .	1261
3.218.8 Giac [F] . . . . .	1261
3.218.9 Mupad [F(-1)] . . . . .	1262

#### 3.218.1 Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-1 - 2n), \frac{1}{4}(3 - 2n), \cos^2(c + dx)\right) \sqrt{\sec(c + dx)}(b \sec(c + dx))^n \sin(c + dx)}{d(1 + 2n)\sqrt{\sin^2(c + dx)}}$$

```
output 2*hypergeom([1/2, -1/4-1/2*n], [3/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*
sin(d*x+c)*sec(d*x+c)^(1/2)/d/(1+2*n)/(sin(d*x+c)^2)^(1/2)
```

#### 3.218.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx$$

$$= \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{3}{2} + n\right), \frac{1}{2}\left(\frac{7}{2} + n\right), \sec^2(c + dx)\right) \sqrt{\sec(c + dx)}(b \sec(c + dx))^n \sqrt{-1}}{d\left(\frac{3}{2} + n\right)}$$

```
input Integrate[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n,x]
```

```
output (Csc[c + d*x]*Hypergeometric2F1[1/2, (3/2 + n)/2, (7/2 + n)/2, Sec[c + d*x]
]^2)*Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2]/(d*(3/2
+ n))
```

**3.218.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^n dx \\
 & \quad \downarrow \text{2034} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \sec^{n+\frac{3}{2}}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \csc\left(c+dx+\frac{\pi}{2}\right)^{n+\frac{3}{2}} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt{\sec(c+dx)} \cos^{n+\frac{1}{2}}(c+dx)(b \sec(c+dx))^n \int \cos^{-n-\frac{3}{2}}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\sec(c+dx)} \cos^{n+\frac{1}{2}}(c+dx)(b \sec(c+dx))^n \int \sin\left(c+dx+\frac{\pi}{2}\right)^{-n-\frac{3}{2}} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{2 \sin(c+dx) \sqrt{\sec(c+dx)} (b \sec(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-2n-1), \frac{1}{4}(3-2n), \cos^2(c+dx)\right)}{d(2n+1) \sqrt{\sin^2(c+dx)}}
 \end{aligned}$$

input `Int[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n,x]`

output `(2*Hypergeometric2F1[1/2, (-1 - 2*n)/4, (3 - 2*n)/4, Cos[c + d*x]^2]*Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(1 + 2*n)*Sqrt[Sin[c + d*x]^2])`



## 3.218.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.218.4 Maple [F]

$$\int \sec(dx + c)^{\frac{3}{2}} (b \sec(dx + c))^n dx$$

```
input int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x)
```

```
output int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x)
```

## 3.218.5 Fracas [F]

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^{\frac{3}{2}} dx$$

```
input integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x, algorithm="fricas")
```

```
output integral((b*sec(d*x + c))^n*sec(d*x + c)^(3/2), x)
```

---

3.218.  $\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx$

**3.218.6 Sympy [F]**

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n \sec^{\frac{3}{2}}(c + dx) dx$$

input `integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**n,x)`

output `Integral((b*sec(c + d*x))**n*sec(c + d*x)**(3/2), x)`

**3.218.7 Maxima [F]**

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^(3/2), x)`

**3.218.8 Giac [F]**

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n*sec(d*x + c)^(3/2), x)`

**3.218.9 Mupad [F(-1)]**

Timed out.

$$\int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n dx = \int \left( \frac{b}{\cos(c + dx)} \right)^n \left( \frac{1}{\cos(c + dx)} \right)^{3/2} dx$$

input `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(3/2),x)`output `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(3/2), x)`

### 3.219 $\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^n dx$

3.219.1 Optimal result . . . . .	1263
3.219.2 Mathematica [A] (verified) . . . . .	1263
3.219.3 Rubi [A] (verified) . . . . .	1264
3.219.4 Maple [F] . . . . .	1265
3.219.5 Fracas [F] . . . . .	1265
3.219.6 Sympy [F] . . . . .	1266
3.219.7 Maxima [F] . . . . .	1266
3.219.8 Giac [F] . . . . .	1266
3.219.9 Mupad [F(-1)] . . . . .	1267

#### 3.219.1 Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^n dx = \frac{2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(1 - 2n), \frac{1}{4}(5 - 2n), \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(1 - 2n) \sqrt{\sec(c + dx)} \sqrt{\sin^2(c + dx)}}$$

output

```
-2*hypergeom([1/2, 1/4-1/2*n], [5/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*
sin(d*x+c)/d/(1-2*n)/sec(d*x+c)^(1/2)/(sin(d*x+c)^2)^(1/2)
```

#### 3.219.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^n dx = \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{1}{2} + n\right), \frac{1}{2}\left(\frac{5}{2} + n\right), \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d\left(\frac{1}{2} + n\right) \sqrt{\sec(c + dx)}}$$

input

```
Integrate[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n,x]
```

output

```
(Csc[c + d*x]*Hypergeometric2F1[1/2, (1/2 + n)/2, (5/2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(1/2 + n)*Sqrt[Sec[c + d*x]])
```

**3.219.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(c+dx)}(b \sec(c+dx))^n dx \\
 & \quad \downarrow \text{2034} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \sec^{n+\frac{1}{2}}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sec^{-n}(c+dx)(b \sec(c+dx))^n \int \csc\left(c+dx+\frac{\pi}{2}\right)^{n+\frac{1}{2}} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt{\sec(c+dx)} \cos^{n+\frac{1}{2}}(c+dx)(b \sec(c+dx))^n \int \cos^{-n-\frac{1}{2}}(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\sec(c+dx)} \cos^{n+\frac{1}{2}}(c+dx)(b \sec(c+dx))^n \int \sin\left(c+dx+\frac{\pi}{2}\right)^{-n-\frac{1}{2}} dx \\
 & \quad \downarrow \text{3122} \\
 & -\frac{2 \sin(c+dx)(b \sec(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(1-2n), \frac{1}{4}(5-2n), \cos^2(c+dx)\right)}{d(1-2n)\sqrt{\sin^2(c+dx)}\sqrt{\sec(c+dx)}}
 \end{aligned}$$

input `Int[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n,x]`

output `(-2*Hypergeometric2F1[1/2, (1 - 2*n)/4, (5 - 2*n)/4, Cos[c + d*x]^2]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(1 - 2*n)*Sqrt[Sec[c + d*x]]*Sqrt[Sin[c + d*x]^2])`

## 3.219.3.1 Defintions of rubi rules used

rule 2034 `Int[(Fv_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m+n)*Fv, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c+d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c+d*x])^(n-1)*((Sin[c+d*x]/b)^(n-1) Int[1/(Sin[c+d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.219.4 Maple [F]

$$\int (b \sec(dx + c))^n \sqrt{\sec(dx + c)} dx$$

input `int((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x)`

output `int((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x)`

## 3.219.5 Fracas [F]

$$\int \sqrt{\sec(c + dx)} (b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sqrt{\sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `integral((b*sec(d*x + c))^n*sqrt(sec(d*x + c)), x)`

**3.219.6 Sympy [F]**

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^n dx = \int (b \sec(c + dx))^n \sqrt{\sec(c + dx)} dx$$

input `integrate((b*sec(d*x+c))**n*sec(d*x+c)**(1/2),x)`

output `Integral((b*sec(c + d*x))**n*sqrt(sec(c + d*x)), x)`

**3.219.7 Maxima [F]**

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sqrt{\sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))n*sec(d*x+c)(1/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))n*sqrt(sec(d*x + c)), x)`

**3.219.8 Giac [F]**

$$\int \sqrt{\sec(c + dx)}(b \sec(c + dx))^n dx = \int (b \sec(dx + c))^n \sqrt{\sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))n*sec(d*x+c)(1/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))n*sqrt(sec(d*x + c)), x)`

**3.219.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{\sec(c + dx)} (b \sec(c + dx))^n dx = \int \left( \frac{b}{\cos(c + dx)} \right)^n \sqrt{\frac{1}{\cos(c + dx)}} dx$$

input `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(1/2),x)`output `int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(1/2), x)`



### 3.220 $\int \frac{(b \sec(c+dx))^n}{\sqrt{\sec(c+dx)}} dx$

3.220.1 Optimal result . . . . .	1268
3.220.2 Mathematica [A] (verified) . . . . .	1268
3.220.3 Rubi [A] (verified) . . . . .	1269
3.220.4 Maple [F] . . . . .	1270
3.220.5 Fricas [F] . . . . .	1270
3.220.6 Sympy [F] . . . . .	1271
3.220.7 Maxima [F] . . . . .	1271
3.220.8 Giac [F] . . . . .	1271
3.220.9 Mupad [F(-1)] . . . . .	1272

#### 3.220.1 Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx = \frac{2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(3 - 2n), \frac{1}{4}(7 - 2n), \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(3 - 2n) \sec^{\frac{3}{2}}(c + dx) \sqrt{\sin^2(c + dx)}}$$

output

```
-2*hypergeom([1/2, 3/4-1/2*n], [7/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(3-2*n)/sec(d*x+c)^(3/2)/(sin(d*x+c)^2)^(1/2)
```

#### 3.220.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx = \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{1}{2} + n\right), \frac{1}{2}\left(\frac{3}{2} + n\right), \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d\left(-\frac{1}{2} + n\right) \sec^{\frac{3}{2}}(c + dx)}$$

input

```
Integrate[(b*Sec[c + d*x])^n/Sqrt[Sec[c + d*x]],x]
```

output  $(\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[1/2, (-1/2 + n)/2, (3/2 + n)/2, \text{Sec}[c + d*x]^2]*(b*\text{Sec}[c + d*x])^n*\text{Sqrt}[-\text{Tan}[c + d*x]^2])/(d*(-1/2 + n)*\text{Sec}[c + d*x]^{3/2})$

### 3.220.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx$$

↓ 2034

$$\sec^{-n}(c + dx)(b \sec(c + dx))^n \int \sec^{n-\frac{1}{2}}(c + dx) dx$$

↓ 3042

$$\sec^{-n}(c + dx)(b \sec(c + dx))^n \int \csc\left(c + dx + \frac{\pi}{2}\right)^{n-\frac{1}{2}} dx$$

↓ 4259

$$\sqrt{\sec(c + dx)} \cos^{n+\frac{1}{2}}(c + dx)(b \sec(c + dx))^n \int \cos^{\frac{1}{2}-n}(c + dx) dx$$

↓ 3042

$$\sqrt{\sec(c + dx)} \cos^{n+\frac{1}{2}}(c + dx)(b \sec(c + dx))^n \int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{1}{2}-n} dx$$

↓ 3122

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(3 - 2n), \frac{1}{4}(7 - 2n), \cos^2(c + dx)\right)}{d(3 - 2n)\sqrt{\sin^2(c + dx)} \sec^{\frac{3}{2}}(c + dx)}$$

input  $\text{Int}[(b*\text{Sec}[c + d*x])^n/\text{Sqrt}[\text{Sec}[c + d*x]], x]$

output  $(-2*\text{Hypergeometric2F1}[1/2, (3 - 2*n)/4, (7 - 2*n)/4, \text{Cos}[c + d*x]^2]*(b*\text{Sec}[c + d*x])^n*\text{Sin}[c + d*x])/(d*(3 - 2*n)*\text{Sec}[c + d*x]^{3/2}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

3.220.  $\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx$

## 3.220.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.220.4 Maple [F]

$$\int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

```
input int((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x)
```

```
output int((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x)
```

## 3.220.5 Fracas [F]

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx = \int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

```
input integrate((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x, algorithm="fricas")
```

output `integral((b*sec(d*x + c))^n/sqrt(sec(d*x + c)), x)`

### 3.220.6 Sympy [F]

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx = \int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)**(1/2),x)`

output `Integral((b*sec(c + d*x))^n/sqrt(sec(c + d*x)), x)`

### 3.220.7 Maxima [F]

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx = \int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n/sqrt(sec(d*x + c)), x)`

### 3.220.8 Giac [F]

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx = \int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n/sqrt(sec(d*x + c)), x)`

**3.220.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\sqrt{\frac{1}{\cos(c+dx)}}} dx$$

input `int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(1/2),x)`output `int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(1/2), x)`

**3.221** 
$$\int \frac{(b \sec(c+dx))^n}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.221.1 Optimal result . . . . . 1273  
 3.221.2 Mathematica [A] (verified) . . . . . 1273  
 3.221.3 Rubi [A] (verified) . . . . . 1274  
 3.221.4 Maple [F] . . . . . 1275  
 3.221.5 Fracas [F] . . . . . 1275  
 3.221.6 Sympy [F] . . . . . 1276  
 3.221.7 Maxima [F] . . . . . 1276  
 3.221.8 Giac [F] . . . . . 1276  
 3.221.9 Mupad [F(-1)] . . . . . 1277

**3.221.1 Optimal result**

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(5 - 2n), \frac{1}{4}(9 - 2n), \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(5 - 2n) \sec^{\frac{5}{2}}(c + dx) \sqrt{\sin^2(c + dx)}}$$

output `-2*hypergeom([1/2, 5/4-1/2*n], [9/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(5-2*n)/sec(d*x+c)^(5/2)/(sin(d*x+c)^2)^(1/2)`

**3.221.2 Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{3}{2} + n\right), \frac{1}{2}\left(\frac{1}{2} + n\right), \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d\left(-\frac{3}{2} + n\right) \sec^{\frac{5}{2}}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^n/Sec[c + d*x]^(3/2),x]`

output  $(\text{Csc}[c + d*x]*\text{Hypergeometric2F1}[1/2, (-3/2 + n)/2, (1/2 + n)/2, \text{Sec}[c + d*x]^2]*(b*\text{Sec}[c + d*x])^n*\text{Sqrt}[-\text{Tan}[c + d*x]^2])/(d*(-3/2 + n)*\text{Sec}[c + d*x]^{5/2})$

### 3.221.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{2034} \\ & \sec^{-n}(c + dx)(b \sec(c + dx))^n \int \sec^{n-\frac{3}{2}}(c + dx) dx \\ & \quad \downarrow \text{3042} \\ & \sec^{-n}(c + dx)(b \sec(c + dx))^n \int \csc\left(c + dx + \frac{\pi}{2}\right)^{n-\frac{3}{2}} dx \\ & \quad \downarrow \text{4259} \\ & \sqrt{\sec(c + dx)} \cos^{n+\frac{1}{2}}(c + dx)(b \sec(c + dx))^n \int \cos^{\frac{3}{2}-n}(c + dx) dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\sec(c + dx)} \cos^{n+\frac{1}{2}}(c + dx)(b \sec(c + dx))^n \int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}-n} dx \\ & \quad \downarrow \text{3122} \\ & \frac{2 \sin(c + dx)(b \sec(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(5 - 2n), \frac{1}{4}(9 - 2n), \cos^2(c + dx)\right)}{d(5 - 2n)\sqrt{\sin^2(c + dx)} \sec^{\frac{5}{2}}(c + dx)} \end{aligned}$$

input  $\text{Int}[(b*\text{Sec}[c + d*x])^n/\text{Sec}[c + d*x]^{3/2}, x]$

output  $(-2*\text{Hypergeometric2F1}[1/2, (5 - 2*n)/4, (9 - 2*n)/4, \text{Cos}[c + d*x]^2]*(b*\text{Sec}[c + d*x])^n*\text{Sin}[c + d*x])/(d*(5 - 2*n)*\text{Sec}[c + d*x]^{5/2}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

---

3.221.  $\int \frac{(b \sec(c+dx))^n}{\sec^{\frac{3}{2}}(c+dx)} dx$

## 3.221.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.221.4 Maple [F]

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{3}{2}}} dx$$

```
input int((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x)
```

```
output int((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x)
```

## 3.221.5 Fracas [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{3}{2}}} dx$$

```
input integrate((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x, algorithm="fracas")
```



output `integral((b*sec(d*x + c))^n/sec(d*x + c)^(3/2), x)`

### 3.221.6 Sympy [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)**(3/2),x)`

output `Integral((b*sec(c + d*x))^n/sec(c + d*x)**(3/2), x)`

### 3.221.7 Maxima [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^n}{\sec^{\frac{3}{2}}(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n/sec(d*x + c)^(3/2), x)`

### 3.221.8 Giac [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^n}{\sec^{\frac{3}{2}}(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n/sec(d*x + c)^(3/2), x)`

**3.221.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(3/2),x)`output `int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(3/2), x)`

**3.222**  $\int \frac{(b \sec(c+dx))^n}{\sec^{\frac{5}{2}}(c+dx)} dx$

3.222.1 Optimal result . . . . . 1278  
 3.222.2 Mathematica [A] (verified) . . . . . 1278  
 3.222.3 Rubi [A] (verified) . . . . . 1279  
 3.222.4 Maple [F] . . . . . 1280  
 3.222.5 Fracas [F] . . . . . 1280  
 3.222.6 Sympy [F] . . . . . 1281  
 3.222.7 Maxima [F] . . . . . 1281  
 3.222.8 Giac [F] . . . . . 1281  
 3.222.9 Mupad [F(-1)] . . . . . 1282

**3.222.1 Optimal result**

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(7 - 2n), \frac{1}{4}(11 - 2n), \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(7 - 2n) \sec^{\frac{7}{2}}(c + dx) \sqrt{\sin^2(c + dx)}}$$

output `-2*hypergeom([1/2, 7/4-1/2*n], [11/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*  
 sin(d*x+c)/d/(7-2*n)/sec(d*x+c)^(7/2)/(sin(d*x+c)^2)^(1/2)`

**3.222.2 Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx = \frac{\csc(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{5}{2} + n\right), \frac{1}{2}\left(-\frac{1}{2} + n\right), \sec^2(c + dx)\right) (b \sec(c + dx))^n \sqrt{-\tan^2(c + dx)}}{d\left(-\frac{5}{2} + n\right) \sec^{\frac{7}{2}}(c + dx)}$$

input `Integrate[(b*Sec[c + d*x])^n/Sec[c + d*x]^(5/2),x]`

output `(Csc[c + d*x]*Hypergeometric2F1[1/2, (-5/2 + n)/2, (-1/2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(-5/2 + n)*Sec[c + d*x]^(7/2))`

### 3.222.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {2034, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{2034} \\
 & \sec^{-n}(c + dx)(b \sec(c + dx))^n \int \sec^{n-\frac{5}{2}}(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sec^{-n}(c + dx)(b \sec(c + dx))^n \int \csc\left(c + dx + \frac{\pi}{2}\right)^{n-\frac{5}{2}} dx \\
 & \quad \downarrow \text{4259} \\
 & \sqrt{\sec(c + dx)} \cos^{n+\frac{1}{2}}(c + dx)(b \sec(c + dx))^n \int \cos^{\frac{5}{2}-n}(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\sec(c + dx)} \cos^{n+\frac{1}{2}}(c + dx)(b \sec(c + dx))^n \int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{2 \sin(c + dx)(b \sec(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(7 - 2n), \frac{1}{4}(11 - 2n), \cos^2(c + dx)\right)}{d(7 - 2n)\sqrt{\sin^2(c + dx)} \sec^{\frac{7}{2}}(c + dx)}
 \end{aligned}$$

input `Int[(b*Sec[c + d*x])^n/Sec[c + d*x]^(5/2),x]`

output `(-2*Hypergeometric2F1[1/2, (7 - 2*n)/4, (11 - 2*n)/4, Cos[c + d*x]^2]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(7 - 2*n)*Sec[c + d*x]^(7/2)*Sqrt[Sin[c + d*x]^2])`

---

3.222.  $\int \frac{(b \sec(c+dx))^n}{\sec^{\frac{5}{2}}(c+dx)} dx$

## 3.222.3.1 Defintions of rubi rules used

```
rule 2034 Int[(Fx_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 4259 Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

## 3.222.4 Maple [F]

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{5}{2}}} dx$$

```
input int((b*sec(d*x+c))^n/sec(d*x+c)^(5/2),x)
```

```
output int((b*sec(d*x+c))^n/sec(d*x+c)^(5/2),x)
```

## 3.222.5 Fracas [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{5}{2}}} dx$$

```
input integrate((b*sec(d*x+c))^n/sec(d*x+c)^(5/2),x, algorithm="fracas")
```

output `integral((b*sec(d*x + c))^n/sec(d*x + c)^(5/2), x)`

### 3.222.6 Sympy [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)**(5/2), x)`

output `Integral((b*sec(c + d*x))^n/sec(c + d*x)**(5/2), x)`

### 3.222.7 Maxima [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^n}{\sec^{\frac{5}{2}}(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)^(5/2), x, algorithm="maxima")`

output `integrate((b*sec(d*x + c))^n/sec(d*x + c)^(5/2), x)`

### 3.222.8 Giac [F]

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \sec(dx + c))^n}{\sec^{\frac{5}{2}}(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^n/sec(d*x+c)^(5/2), x, algorithm="giac")`

output `integrate((b*sec(d*x + c))^n/sec(d*x + c)^(5/2), x)`

**3.222.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(5/2),x)`output `int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(5/2), x)`

### 3.223 $\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx$

3.223.1 Optimal result . . . . .	1283
3.223.2 Mathematica [A] (verified) . . . . .	1283
3.223.3 Rubi [A] (verified) . . . . .	1284
3.223.4 Maple [A] (verified) . . . . .	1285
3.223.5 Fracas [A] (verification not implemented) . . . . .	1285
3.223.6 Sympy [F(-1)] . . . . .	1286
3.223.7 Maxima [A] (verification not implemented) . . . . .	1286
3.223.8 Giac [B] (verification not implemented) . . . . .	1286
3.223.9 Mupad [B] (verification not implemented) . . . . .	1287

#### 3.223.1 Optimal result

Integrand size = 19, antiderivative size = 20

$$\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx = \frac{2d(d \sec(a + bx))^{5/2}}{5b}$$

output `2/5*d*(d*sec(b*x+a))^(5/2)/b`

#### 3.223.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx = \frac{2d(d \sec(a + bx))^{5/2}}{5b}$$

input `Integrate[(d*Sec[a + b*x])^(7/2)*Sin[a + b*x],x]`

output `(2*d*(d*Sec[a + b*x])^(5/2))/(5*b)`



**3.223.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3102, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(a + bx)(d \sec(a + bx))^{7/2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(d \sec(a + bx))^{7/2}}{\csc(a + bx)} dx$$

$$\downarrow \text{3102}$$

$$\frac{d \int (d \sec(a + bx))^{3/2} d(d \sec(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{2d(d \sec(a + bx))^{5/2}}{5b}$$

input `Int[(d*Sec[a + b*x])^(7/2)*Sin[a + b*x],x]`

output `(2*d*(d*Sec[a + b*x])^(5/2))/(5*b)`

**3.223.3.1 Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3102 Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol]
:> Simp[1/(f*a^n) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

### 3.223.4 Maple [A] (verified)

Time = 3.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$\frac{2d(d \sec(bx+a))^{5/2}}{5b}$	17
default	$\frac{2d(d \sec(bx+a))^{5/2}}{5b}$	17

```
input int((d*sec(b*x+a))^(7/2)*sin(b*x+a),x,method=_RETURNVERBOSE)
```

```
output 2/5*d*(d*sec(b*x+a))^(5/2)/b
```

### 3.223.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.40

$$\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx = \frac{2 d^3 \sqrt{\frac{d}{\cos(bx+a)}}}{5 b \cos(bx + a)^2}$$

```
input integrate((d*sec(b*x+a))^(7/2)*sin(b*x+a),x, algorithm="fricas")
```

```
output 2/5*d^3*sqrt(d/cos(b*x + a))/(b*cos(b*x + a)^2)
```

**3.223.6 Sympy [F(-1)]**

Timed out.

$$\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx = \text{Timed out}$$

input `integrate((d*sec(b*x+a))**(7/2)*sin(b*x+a),x)`output `Timed out`**3.223.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx = \frac{2 \left( \frac{d}{\cos(bx+a)} \right)^{7/2} \cos(bx + a)}{5b}$$

input `integrate((d*sec(b*x+a))^(7/2)*sin(b*x+a),x, algorithm="maxima")`output `2/5*(d/cos(b*x + a))^(7/2)*cos(b*x + a)/b`**3.223.8 Giac [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 33 vs.  $2(16) = 32$ .

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.65

$$\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx = \frac{2 d^4 \operatorname{sgn}(\cos(bx + a))}{5 \sqrt{d \cos(bx + a)} b \cos(bx + a)^2}$$

input `integrate((d*sec(b*x+a))^(7/2)*sin(b*x+a),x, algorithm="giac")`output `2/5*d^4*sgn(cos(b*x + a))/(sqrt(d*cos(b*x + a))*b*cos(b*x + a)^2)`

**3.223.9 Mupad [B] (verification not implemented)**

Time = 14.50 (sec) , antiderivative size = 77, normalized size of antiderivative = 3.85

$$\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx = \frac{8 d^3 \sqrt{\frac{d}{\cos(a+bx)}} (4 \cos(2a + 2bx) + \cos(4a + 4bx) + 3)}{5b (15 \cos(2a + 2bx) + 6 \cos(4a + 4bx) + \cos(6a + 6bx) + 10)}$$

input `int(sin(a + b*x)*(d/cos(a + b*x))^(7/2),x)`output `(8*d^3*(d/cos(a + b*x))^(1/2)*(4*cos(2*a + 2*b*x) + cos(4*a + 4*b*x) + 3)) / (5*b*(15*cos(2*a + 2*b*x) + 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) + 10))`

### 3.224 $\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx$

3.224.1 Optimal result . . . . .	1288
3.224.2 Mathematica [A] (verified) . . . . .	1288
3.224.3 Rubi [A] (verified) . . . . .	1289
3.224.4 Maple [A] (verified) . . . . .	1290
3.224.5 Fricas [A] (verification not implemented) . . . . .	1290
3.224.6 Sympy [F(-1)] . . . . .	1291
3.224.7 Maxima [A] (verification not implemented) . . . . .	1291
3.224.8 Giac [B] (verification not implemented) . . . . .	1291
3.224.9 Mupad [B] (verification not implemented) . . . . .	1292

#### 3.224.1 Optimal result

Integrand size = 19, antiderivative size = 20

$$\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx = \frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

output `2/3*d*(d*sec(b*x+a))^(3/2)/b`

#### 3.224.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx = \frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

input `Integrate[(d*Sec[a + b*x])^(5/2)*Sin[a + b*x],x]`

output `(2*d*(d*Sec[a + b*x])^(3/2))/(3*b)`

**3.224.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3102, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(a + bx)(d \sec(a + bx))^{5/2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(d \sec(a + bx))^{5/2}}{\csc(a + bx)} dx$$

$$\downarrow \text{3102}$$

$$\frac{d \int \sqrt{d \sec(a + bx)} d(d \sec(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

input `Int[(d*Sec[a + b*x])^(5/2)*Sin[a + b*x],x]`

output `(2*d*(d*Sec[a + b*x])^(3/2))/(3*b)`

**3.224.3.1 Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3102 Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol]
:> Simp[1/(f*a^n) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

### 3.224.4 Maple [A] (verified)

Time = 3.37 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$\frac{2d(d \sec(bx+a))^{\frac{3}{2}}}{3b}$	17
default	$\frac{2d(d \sec(bx+a))^{\frac{3}{2}}}{3b}$	17

```
input int((d*sec(b*x+a))^(5/2)*sin(b*x+a),x,method=_RETURNVERBOSE)
```

```
output 2/3*d*(d*sec(b*x+a))^(3/2)/b
```

### 3.224.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.40

$$\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx = \frac{2 d^2 \sqrt{\frac{d}{\cos(bx+a)}}}{3 b \cos(bx + a)}$$

```
input integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a),x, algorithm="fracas")
```

```
output 2/3*d^2*sqrt(d/cos(b*x + a))/(b*cos(b*x + a))
```

**3.224.6 Sympy [F(-1)]**

Timed out.

$$\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx = \text{Timed out}$$

input `integrate((d*sec(b*x+a))**(5/2)*sin(b*x+a),x)`output `Timed out`**3.224.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx = \frac{2 \left( \frac{d}{\cos(bx+a)} \right)^{5/2} \cos(bx + a)}{3b}$$

input `integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a),x, algorithm="maxima")`output `2/3*(d/cos(b*x + a))^(5/2)*cos(b*x + a)/b`**3.224.8 Giac [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 33 vs.  $2(16) = 32$ .

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.65

$$\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx = \frac{2 d^3 \operatorname{sgn}(\cos(bx + a))}{3 \sqrt{d \cos(bx + a)} b \cos(bx + a)}$$

input `integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a),x, algorithm="giac")`output `2/3*d^3*sgn(cos(b*x + a))/(sqrt(d*cos(b*x + a))*b*cos(b*x + a))`



**3.224.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.95

$$\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx = \frac{4 d^2 \cos(a + bx) \sqrt{\frac{d}{\cos(a+bx)}}}{3b (\cos(2a + 2bx) + 1)}$$

input `int(sin(a + b*x)*(d/cos(a + b*x))^(5/2),x)`output `(4*d^2*cos(a + b*x)*(d/cos(a + b*x))^(1/2))/(3*b*(cos(2*a + 2*b*x) + 1))`

### 3.225 $\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx$

3.225.1 Optimal result . . . . .	1293
3.225.2 Mathematica [A] (verified) . . . . .	1293
3.225.3 Rubi [A] (verified) . . . . .	1294
3.225.4 Maple [A] (verified) . . . . .	1295
3.225.5 Fricas [A] (verification not implemented) . . . . .	1295
3.225.6 Sympy [F] . . . . .	1296
3.225.7 Maxima [A] (verification not implemented) . . . . .	1296
3.225.8 Giac [A] (verification not implemented) . . . . .	1296
3.225.9 Mupad [B] (verification not implemented) . . . . .	1297

#### 3.225.1 Optimal result

Integrand size = 19, antiderivative size = 18

$$\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx = \frac{2d\sqrt{d \sec(a + bx)}}{b}$$

output `2*d*(d*sec(b*x+a))^(1/2)/b`

#### 3.225.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx = \frac{2d\sqrt{d \sec(a + bx)}}{b}$$

input `Integrate[(d*Sec[a + b*x])^(3/2)*Sin[a + b*x],x]`

output `(2*d*Sqrt[d*Sec[a + b*x]])/b`

**3.225.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3102, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin(a + bx)(d \sec(a + bx))^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(d \sec(a + bx))^{3/2}}{\csc(a + bx)} dx \\ & \quad \downarrow \text{3102} \\ & \frac{d \int \frac{1}{\sqrt{d \sec(a + bx)}} d(d \sec(a + bx))}{b} \\ & \quad \downarrow \text{15} \\ & \frac{2d\sqrt{d \sec(a + bx)}}{b} \end{aligned}$$

input `Int[(d*Sec[a + b*x])^(3/2)*Sin[a + b*x],x]`

output `(2*d*Sqrt[d*Sec[a + b*x]])/b`

**3.225.3.1 Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3102 Int[csc[(e_.) + (f_.)*(x_.)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_.)]^(m_.), x_Symbol]
:> Simp[1/(f*a^n) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

### 3.225.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{2d\sqrt{d\sec(bx+a)}}{b}$	17
default	$\frac{2d\sqrt{d\sec(bx+a)}}{b}$	17

```
input int((d*sec(b*x+a))^(3/2)*sin(b*x+a),x,method=_RETURNVERBOSE)
```

```
output 2*d*(d*sec(b*x+a))^(1/2)/b
```

### 3.225.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx = \frac{2d\sqrt{\frac{d}{\cos(bx+a)}}}{b}$$

```
input integrate((d*sec(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="fracas")
```

```
output 2*d*sqrt(d/cos(b*x + a))/b
```

**3.225.6 Sympy [F]**

$$\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx = \int (d \sec(a + bx))^{\frac{3}{2}} \sin(a + bx) dx$$

input `integrate((d*sec(b*x+a))**(3/2)*sin(b*x+a),x)`

output `Integral((d*sec(a + b*x))**(3/2)*sin(a + b*x), x)`

**3.225.7 Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx = \frac{2 \left( \frac{d}{\cos(bx+a)} \right)^{\frac{3}{2}} \cos(bx + a)}{b}$$

input `integrate((d*sec(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="maxima")`

output `2*(d/cos(b*x + a))^(3/2)*cos(b*x + a)/b`

**3.225.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.39

$$\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx = \frac{2 d^2 \operatorname{sgn}(\cos(bx + a))}{\sqrt{d \cos(bx + a)} b}$$

input `integrate((d*sec(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="giac")`

output `2*d^2*sgn(cos(b*x + a))/(sqrt(d*cos(b*x + a))*b)`

**3.225.9 Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx = \frac{2d \sqrt{\frac{d}{\cos(a+bx)}}}{b}$$

input `int(sin(a + b*x)*(d/cos(a + b*x))^(3/2),x)`output `(2*d*(d/cos(a + b*x))^(1/2))/b`

### 3.226 $\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx$

3.226.1 Optimal result . . . . .	1298
3.226.2 Mathematica [A] (verified) . . . . .	1298
3.226.3 Rubi [A] (verified) . . . . .	1299
3.226.4 Maple [A] (verified) . . . . .	1300
3.226.5 Fricas [A] (verification not implemented) . . . . .	1300
3.226.6 Sympy [F] . . . . .	1301
3.226.7 Maxima [A] (verification not implemented) . . . . .	1301
3.226.8 Giac [A] (verification not implemented) . . . . .	1301
3.226.9 Mupad [B] (verification not implemented) . . . . .	1302

#### 3.226.1 Optimal result

Integrand size = 19, antiderivative size = 18

$$\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx = -\frac{2d}{b\sqrt{d \sec(a + bx)}}$$

output `-2*d/b/(d*sec(b*x+a))^(1/2)`

#### 3.226.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx = -\frac{2d}{b\sqrt{d \sec(a + bx)}}$$

input `Integrate[Sqrt[d*Sec[a + b*x]]*Sin[a + b*x],x]`

output `(-2*d)/(b*Sqrt[d*Sec[a + b*x]])`

**3.226.3 Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3102, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(a + bx) \sqrt{d \sec(a + bx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{d \sec(a + bx)}}{\csc(a + bx)} dx$$

$$\downarrow \text{3102}$$

$$\frac{d \int \frac{1}{(d \sec(a + bx))^{3/2}} d(d \sec(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$-\frac{2d}{b \sqrt{d \sec(a + bx)}}$$

input `Int[Sqrt[d*Sec[a + b*x]]*Sin[a + b*x],x]`

output `(-2*d)/(b*Sqrt[d*Sec[a + b*x]])`

**3.226.3.1 Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



```
rule 3102 Int[csc[(e_.) + (f_.)*(x_.)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_.)]^(m_.), x_Symbol]
:> Simp[1/(f*a^n) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

### 3.226.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$-\frac{2d}{b\sqrt{d\sec(bx+a)}}$	17
default	$-\frac{2d}{b\sqrt{d\sec(bx+a)}}$	17
risch	$-\frac{2\sqrt{2}\sqrt{\frac{de^{i(bx+a)}}{e^{2i(bx+a)}+1}}\cos(bx+a)}{b}$	41

```
input int((d*sec(b*x+a))^(1/2)*sin(b*x+a),x,method=_RETURNVERBOSE)
```

```
output -2*d/b/(d*sec(b*x+a))^(1/2)
```

### 3.226.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \sqrt{d\sec(a+bx)} \sin(a+bx) dx = -\frac{2\sqrt{\frac{d}{\cos(bx+a)}} \cos(bx+a)}{b}$$

```
input integrate((d*sec(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="fracas")
```

```
output -2*sqrt(d/cos(b*x + a))*cos(b*x + a)/b
```

**3.226.6 Sympy [F]**

$$\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx = \int \sqrt{d \sec(a + bx)} \sin(a + bx) dx$$

input `integrate((d*sec(b*x+a))**(1/2)*sin(b*x+a),x)`

output `Integral(sqrt(d*sec(a + b*x))*sin(a + b*x), x)`

**3.226.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx = -\frac{2 \sqrt{\frac{d}{\cos(bx+a)}} \cos(bx + a)}{b}$$

input `integrate((d*sec(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="maxima")`

output `-2*sqrt(d/cos(b*x + a))*cos(b*x + a)/b`

**3.226.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx = -\frac{2 \sqrt{d \cos(bx + a)} \operatorname{sgn}(\cos(bx + a))}{b}$$

input `integrate((d*sec(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="giac")`

output `-2*sqrt(d*cos(b*x + a))*sgn(cos(b*x + a))/b`

**3.226.9 Mupad [B] (verification not implemented)**

Time = 13.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx = -\frac{2 \cos(a + bx) \sqrt{\frac{d}{\cos(a + bx)}}}{b}$$

input `int(sin(a + b*x)*(d/cos(a + b*x))^(1/2),x)`

output `-(2*cos(a + b*x)*(d/cos(a + b*x))^(1/2))/b`

**3.227**      $\int \frac{\sin(a+bx)}{\sqrt{d \sec(a+bx)}} dx$

3.227.1 Optimal result . . . . . 1303  
 3.227.2 Mathematica [A] (verified) . . . . . 1303  
 3.227.3 Rubi [A] (verified) . . . . . 1304  
 3.227.4 Maple [A] (verified) . . . . . 1305  
 3.227.5 Fricas [A] (verification not implemented) . . . . . 1305  
 3.227.6 Sympy [F] . . . . . 1306  
 3.227.7 Maxima [A] (verification not implemented) . . . . . 1306  
 3.227.8 Giac [B] (verification not implemented) . . . . . 1306  
 3.227.9 Mupad [B] (verification not implemented) . . . . . 1307

**3.227.1 Optimal result**

Integrand size = 19, antiderivative size = 20

$$\int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx = -\frac{2d}{3b(d \sec(a + bx))^{3/2}}$$

output `-2/3*d/b/(d*sec(b*x+a))^(3/2)`

**3.227.2 Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx = -\frac{2d}{3b(d \sec(a + bx))^{3/2}}$$

input `Integrate[Sin[a + b*x]/Sqrt[d*Sec[a + b*x]],x]`

output `(-2*d)/(3*b*(d*Sec[a + b*x])^(3/2))`

**3.227.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3102, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin(a+bx)}{\sqrt{d \sec(a+bx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\csc(a+bx) \sqrt{d \sec(a+bx)}} dx \\ & \quad \downarrow \text{3102} \\ & \frac{d \int \frac{1}{(d \sec(a+bx))^{5/2}} d(d \sec(a+bx))}{b} \\ & \quad \downarrow \text{15} \\ & -\frac{2d}{3b(d \sec(a+bx))^{3/2}} \end{aligned}$$

input `Int[Sin[a + b*x]/Sqrt[d*Sec[a + b*x]],x]`

output `(-2*d)/(3*b*(d*Sec[a + b*x])^(3/2))`

**3.227.3.1 Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3102 Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol]
:> Simp[1/(f*a^n) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

### 3.227.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$-\frac{2d}{3b(d \sec(bx+a))^{\frac{3}{2}}}$	17
default	$-\frac{2d}{3b(d \sec(bx+a))^{\frac{3}{2}}}$	17

```
input int(sin(b*x+a)/(d*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2/3*d/b/(d*sec(b*x+a))^(3/2)
```

### 3.227.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.40

$$\int \frac{\sin(a+bx)}{\sqrt{d \sec(a+bx)}} dx = -\frac{2 \sqrt{\frac{d}{\cos(bx+a)}} \cos(bx+a)^2}{3bd}$$

```
input integrate(sin(b*x+a)/(d*sec(b*x+a))^(1/2),x, algorithm="fricas")
```

```
output -2/3*sqrt(d/cos(b*x + a))*cos(b*x + a)^2/(b*d)
```

**3.227.6 Sympy [F]**

$$\int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx = \int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx$$

input `integrate(sin(b*x+a)/(d*sec(b*x+a))**(1/2),x)`

output `Integral(sin(a + b*x)/sqrt(d*sec(a + b*x)), x)`

**3.227.7 Maxima [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx = -\frac{2 \cos(bx + a)}{3b \sqrt{\frac{d}{\cos(bx+a)}}}$$

input `integrate(sin(b*x+a)/(d*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `-2/3*cos(b*x + a)/(b*sqrt(d/cos(b*x + a)))`

**3.227.8 Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(16) = 32.

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.65

$$\int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx = -\frac{2 \sqrt{d \cos(bx + a)} \cos(bx + a)}{3bd \operatorname{sgn}(\cos(bx + a))}$$

input `integrate(sin(b*x+a)/(d*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `-2/3*sqrt(d*cos(b*x + a))*cos(b*x + a)/(b*d*sgn(cos(b*x + a)))`

**3.227.9 Mupad [B] (verification not implemented)**

Time = 13.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.40

$$\int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx = -\frac{2 \cos(a + bx)^2 \sqrt{\frac{d}{\cos(a + bx)}}}{3bd}$$

input `int(sin(a + b*x)/(d/cos(a + b*x))^(1/2),x)`

output `-(2*cos(a + b*x)^2*(d/cos(a + b*x))^(1/2))/(3*b*d)`



### 3.228 $\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx$

3.228.1 Optimal result . . . . .	1308
3.228.2 Mathematica [A] (verified) . . . . .	1308
3.228.3 Rubi [A] (verified) . . . . .	1309
3.228.4 Maple [B] (verified) . . . . .	1310
3.228.5 Fricas [A] (verification not implemented) . . . . .	1311
3.228.6 Sympy [F(-1)] . . . . .	1311
3.228.7 Maxima [A] (verification not implemented) . . . . .	1312
3.228.8 Giac [A] (verification not implemented) . . . . .	1312
3.228.9 Mupad [B] (verification not implemented) . . . . .	1312

#### 3.228.1 Optimal result

Integrand size = 21, antiderivative size = 41

$$\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx = \frac{2d^3}{b\sqrt{d \sec(a + bx)}} + \frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

output `2/3*d*(d*sec(b*x+a))^(3/2)/b+2*d^3/b/(d*sec(b*x+a))^(1/2)`

#### 3.228.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx = \frac{d(5 + 3 \cos(2(a + bx)))(d \sec(a + bx))^{3/2}}{3b}$$

input `Integrate[(d*Sec[a + b*x])^(5/2)*Sin[a + b*x]^3,x]`

output `(d*(5 + 3*Cos[2*(a + b*x)])*(d*Sec[a + b*x])^(3/2))/(3*b)`

**3.228.3 Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3102, 25, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^3(a + bx)(d \sec(a + bx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \sec(a + bx))^{5/2}}{\csc(a + bx)^3} dx \\
 & \quad \downarrow \text{3102} \\
 & \frac{d^3 \int -\frac{d^2 - d^2 \sec^2(a + bx)}{d^2 (d \sec(a + bx))^{3/2}} d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{25} \\
 & -\frac{d^3 \int \frac{d^2 - d^2 \sec^2(a + bx)}{d^2 (d \sec(a + bx))^{3/2}} d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{27} \\
 & -\frac{d \int \frac{d^2 - d^2 \sec^2(a + bx)}{(d \sec(a + bx))^{3/2}} d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{244} \\
 & -\frac{d \int \left( \frac{d^2}{(d \sec(a + bx))^{3/2}} - \sqrt{d \sec(a + bx)} \right) d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{d \left( -\frac{2d^2}{\sqrt{d \sec(a + bx)}} - \frac{2}{3} (d \sec(a + bx))^{3/2} \right)}{b}
 \end{aligned}$$

input `Int[(d*Sec[a + b*x])^(5/2)*Sin[a + b*x]^3,x]`

output `-((d*((-2*d^2)/Sqrt[d*Sec[a + b*x]] - (2*(d*Sec[a + b*x])^(3/2))/3))/b)`

## 3.228.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 244 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3102 `Int[csc[(e_) + (f_)*(x_)^(n_)]*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[1/(f*a^n) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

## 3.228.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 318 vs. 2(35) = 70.

Time = 67.43 (sec) , antiderivative size = 319, normalized size of antiderivative = 7.78

method	result
default	$\frac{\sqrt{d \sec(bx+a)} d^2 \left( 12 \sqrt{-\frac{\cos(bx+a)}{(\cos(bx+a)+1)^2}} \cos(bx+a)^2 + 12 \cos(bx+a) \sqrt{-\frac{\cos(bx+a)}{(\cos(bx+a)+1)^2}} + 3 \cos(bx+a) \ln \left( \frac{4 \cos(bx+a) \sqrt{-\frac{\cos(bx+a)}{(\cos(bx+a)+1)^2}}}{(\cos(bx+a)+1)} \right) \right)}{\dots}$

input `int((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x,method=_RETURNVERBOSE)`

output  $1/6/b*(d*\sec(b*x+a))^{(1/2)*d^2/(\cos(b*x+a)+1)/(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)*(12*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)*\cos(b*x+a)^2+12*\cos(b*x+a)*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)+3*\cos(b*x+a)*\ln(2*(2*\cos(b*x+a)*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)+2*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)-\cos(b*x+a)+1)/(\cos(b*x+a)+1))-3*\cos(b*x+a)*\ln((2*\cos(b*x+a)*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)+2*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)-\cos(b*x+a)+1)/(\cos(b*x+a)+1))+4*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)+4*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)*\sec(b*x+a))}$

### 3.228.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.02

$$\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx = \frac{2(3d^2 \cos(bx + a)^2 + d^2) \sqrt{\frac{d}{\cos(bx + a)}}}{3b \cos(bx + a)}$$

input `integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x, algorithm="fracas")`

output  $2/3*(3*d^2*\cos(b*x + a)^2 + d^2)*\text{sqrt}(d/\cos(b*x + a))/(b*\cos(b*x + a))$

### 3.228.6 Sympy [F(-1)]

Timed out.

$$\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx = \text{Timed out}$$

input `integrate((d*sec(b*x+a))**(5/2)*sin(b*x+a)**3,x)`

output `Timed out`

**3.228.7 Maxima [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.88

$$\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx = \frac{2 \left( \frac{3d^2}{\sqrt{\frac{d}{\cos(bx+a)}}} + \left( \frac{d}{\cos(bx+a)} \right)^{\frac{3}{2}} \right) d}{3b}$$

input `integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x, algorithm="maxima")`output `2/3*(3*d^2/sqrt(d/cos(b*x + a)) + (d/cos(b*x + a))^(3/2))*d/b`**3.228.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.20

$$\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx = \frac{2 \left( 3 \sqrt{d \cos(bx + a)} d + \frac{d^2}{\sqrt{d \cos(bx+a) \cos(bx+a)}} \right) d \operatorname{sgn}(\cos(bx + a))}{3b}$$

input `integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x, algorithm="giac")`output `2/3*(3*sqrt(d*cos(b*x + a))*d + d^2/(sqrt(d*cos(b*x + a))*cos(b*x + a)))*d*sgn(cos(b*x + a))/b`**3.228.9 Mupad [B] (verification not implemented)**

Time = 13.46 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.22

$$\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx = \frac{d^2 \sqrt{\frac{d}{\cos(a+bx)}} \left( \frac{13 \cos(a+bx)}{3} + \cos(3a + 3bx) \right)}{b (\cos(2a + 2bx) + 1)}$$

input `int(sin(a + b*x)^3*(d/cos(a + b*x))^(5/2),x)`output `(d^2*(d/cos(a + b*x))^(1/2)*((13*cos(a + b*x))/3 + cos(3*a + 3*b*x)))/(b*(cos(2*a + 2*b*x) + 1))`

### 3.229 $\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx$

3.229.1 Optimal result . . . . .	1313
3.229.2 Mathematica [A] (verified) . . . . .	1313
3.229.3 Rubi [A] (verified) . . . . .	1314
3.229.4 Maple [A] (verified) . . . . .	1315
3.229.5 Fricas [A] (verification not implemented) . . . . .	1316
3.229.6 Sympy [F(-1)] . . . . .	1316
3.229.7 Maxima [A] (verification not implemented) . . . . .	1316
3.229.8 Giac [A] (verification not implemented) . . . . .	1317
3.229.9 Mupad [B] (verification not implemented) . . . . .	1317

#### 3.229.1 Optimal result

Integrand size = 21, antiderivative size = 43

$$\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx = -\frac{2d^3 (d \sec(a + bx))^{3/2}}{3b} + \frac{2d (d \sec(a + bx))^{7/2}}{7b}$$

output `-2/3*d^3*(d*sec(b*x+a))^(3/2)/b+2/7*d*(d*sec(b*x+a))^(7/2)/b`

#### 3.229.2 Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx = -\frac{d^4 (1 + 7 \cos(2(a + bx))) \sec^3(a + bx) \sqrt{d \sec(a + bx)}}{21b}$$

input `Integrate[(d*Sec[a + b*x])^(9/2)*Sin[a + b*x]^3,x]`

output `-1/21*(d^4*(1 + 7*Cos[2*(a + b*x)])*Sec[a + b*x]^3*Sqrt[d*Sec[a + b*x]])/b`

**3.229.3 Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3102, 25, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^3(a + bx)(d \sec(a + bx))^{9/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \sec(a + bx))^{9/2}}{\csc(a + bx)^3} dx \\
 & \quad \downarrow \text{3102} \\
 & \frac{d^3 \int -\frac{\sqrt{d \sec(a + bx)}(d^2 - d^2 \sec^2(a + bx))}{d^2} d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{25} \\
 & \frac{d^3 \int \frac{\sqrt{d \sec(a + bx)}(d^2 - d^2 \sec^2(a + bx))}{d^2} d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{27} \\
 & \frac{d \int \sqrt{d \sec(a + bx)}(d^2 - d^2 \sec^2(a + bx)) d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{244} \\
 & \frac{d \int \left( d^2 \sqrt{d \sec(a + bx)} - (d \sec(a + bx))^{5/2} \right) d(d \sec(a + bx))}{b} \\
 & \quad \downarrow \text{2009} \\
 & \frac{d \left( \frac{2}{3} d^2 (d \sec(a + bx))^{3/2} - \frac{2}{7} (d \sec(a + bx))^{7/2} \right)}{b}
 \end{aligned}$$

input `Int[(d*Sec[a + b*x])^(9/2)*Sin[a + b*x]^3,x]`

output `-((d*((2*d^2*(d*Sec[a + b*x])^(3/2))/3 - (2*(d*Sec[a + b*x])^(7/2))/7))/b)`

## 3.229.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3102 `Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Simp[1/(f*a^n) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

## 3.229.4 Maple [A] (verified)

Time = 60.72 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{2d \left( \frac{(d \sec(bx+a))^{\frac{7}{2}}}{7} - \frac{d^2 (d \sec(bx+a))^{\frac{3}{2}}}{3} \right)}{b}$	35
default	$\frac{2d \left( \frac{(d \sec(bx+a))^{\frac{7}{2}}}{7} - \frac{d^2 (d \sec(bx+a))^{\frac{3}{2}}}{3} \right)}{b}$	35

input `int((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `2/b*d*(1/7*(d*sec(b*x+a))^(7/2)-1/3*d^2*(d*sec(b*x+a))^(3/2))`



**3.229.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.02

$$\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx = -\frac{2(7d^4 \cos(bx + a)^2 - 3d^4) \sqrt{\frac{d}{\cos(bx+a)}}}{21b \cos(bx + a)^3}$$

input `integrate((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x, algorithm="fricas")`output `-2/21*(7*d^4*cos(b*x + a)^2 - 3*d^4)*sqrt(d/cos(b*x + a))/(b*cos(b*x + a)^3)`**3.229.6 Sympy [F(-1)]**

Timed out.

$$\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx = \text{Timed out}$$

input `integrate((d*sec(b*x+a))**(9/2)*sin(b*x+a)**3,x)`output `Timed out`**3.229.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.88

$$\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx = -\frac{2 \left( 7d^2 \left( \frac{d}{\cos(bx+a)} \right)^{\frac{3}{2}} - 3 \left( \frac{d}{\cos(bx+a)} \right)^{\frac{7}{2}} \right) d}{21b}$$

input `integrate((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x, algorithm="maxima")`output `-2/21*(7*d^2*(d/cos(b*x + a))^(3/2) - 3*(d/cos(b*x + a))^(7/2))*d/b`

**3.229.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.14

$$\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx = -\frac{2(7d^5 \cos(bx + a)^2 - 3d^5) \operatorname{sgn}(\cos(bx + a))}{21 \sqrt{d \cos(bx + a)} b \cos(bx + a)^3}$$

input `integrate((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x, algorithm="giac")`output `-2/21*(7*d^5*cos(b*x + a)^2 - 3*d^5)*sgn(cos(b*x + a))/(sqrt(d*cos(b*x + a))*b*cos(b*x + a)^3)`**3.229.9 Mupad [B] (verification not implemented)**

Time = 17.17 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.21

$$\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx = -\frac{4d^4 e^{a1i+bx1i} \sqrt{\frac{e^{-a1i-bx1i}}{2} + \frac{e^{a1i+bx1i}}{2}} (2e^{a2i+bx2i} + 7e^{a4i+bx4i} + 7)}{21b(e^{a2i+bx2i} + 1)^3}$$

input `int(sin(a + b*x)^3*(d/cos(a + b*x))^(9/2),x)`output `-(4*d^4*exp(a*1i + b*x*1i)*(d/(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2)*(2*exp(a*2i + b*x*2i) + 7*exp(a*4i + b*x*4i) + 7))/(21*b*(exp(a*2i + b*x*2i) + 1)^3)`

### 3.230 $\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx$

3.230.1 Optimal result . . . . .	1318
3.230.2 Mathematica [C] (verified) . . . . .	1318
3.230.3 Rubi [A] (verified) . . . . .	1319
3.230.4 Maple [A] (verified) . . . . .	1321
3.230.5 Fracas [C] (verification not implemented) . . . . .	1322
3.230.6 Sympy [F(-1)] . . . . .	1322
3.230.7 Maxima [F] . . . . .	1323
3.230.8 Giac [F] . . . . .	1323
3.230.9 Mupad [F(-1)] . . . . .	1323

#### 3.230.1 Optimal result

Integrand size = 25, antiderivative size = 128

$$\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx = -\frac{4cd^3(d \csc(a + bx))^{3/2}}{7b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b\sqrt{c \sec(a + bx)}} + \frac{4d^4 \sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{7b}$$

output

```
-4/7*c*d^3*(d*csc(b*x+a))^(3/2)/b/(c*sec(b*x+a))^(1/2)-2/7*c*d*(d*csc(b*x+a))^(7/2)/b/(c*sec(b*x+a))^(1/2)-4/7*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b
```

#### 3.230.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 18.61 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.95

$$\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx = \frac{2d^4 \cos(2(a + bx)) \cot(a + bx) \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \left( (-2 + \cos(2(a + bx))) \right)}{7}$$

input `Integrate[(d*Csc[a + b*x])^(9/2)*Sqrt[c*Sec[a + b*x]],x]`

output  $(2*d^4*\text{Cos}[2*(a + b*x)]*\text{Cot}[a + b*x]*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*((-2 + \text{Cos}[2*(a + b*x)])*\text{Csc}[a + b*x]^4 - 2*(-\text{Cot}[a + b*x]^2)^(3/4)*\text{Hypergeometric2F1}[1/2, 3/4, 3/2, \text{Csc}[a + b*x]^2]*\text{Sec}[a + b*x]^2))/(7*b*(-2 + \text{Csc}[a + b*x]^2))$

### 3.230.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3105, 3042, 3105, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{9/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{9/2} dx \\ & \quad \downarrow \text{3105} \\ & \frac{6}{7} d^2 \int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx - \frac{2cd (d \csc(a + bx))^{7/2}}{7b \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{6}{7} d^2 \int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx - \frac{2cd (d \csc(a + bx))^{7/2}}{7b \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3105} \\ & \frac{6}{7} d^2 \left( \frac{2}{3} d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx - \frac{2cd (d \csc(a + bx))^{3/2}}{3b \sqrt{c \sec(a + bx)}} \right) - \frac{2cd (d \csc(a + bx))^{7/2}}{7b \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{6}{7} d^2 \left( \frac{2}{3} d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx - \frac{2cd (d \csc(a + bx))^{3/2}}{3b \sqrt{c \sec(a + bx)}} \right) - \frac{2cd (d \csc(a + bx))^{7/2}}{7b \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3110} \end{aligned}$$

$$\frac{6}{7}d^2 \left( \frac{2}{3}d^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx - \frac{2cd}{3b\sqrt{c \sec(a+bx)}} \right) - \frac{2cd(d \csc(a+bx))^{7/2}}{7b\sqrt{c \sec(a+bx)}}$$

↓ 3042

$$\frac{6}{7}d^2 \left( \frac{2}{3}d^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx - \frac{2cd}{3b\sqrt{c \sec(a+bx)}} \right) - \frac{2cd(d \csc(a+bx))^{7/2}}{7b\sqrt{c \sec(a+bx)}}$$

↓ 3053

$$\frac{6}{7}d^2 \left( \frac{2}{3}d^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx - \frac{2cd(d \csc(a+bx))^{3/2}}{3b\sqrt{c \sec(a+bx)}} \right) - \frac{2cd(d \csc(a+bx))^{7/2}}{7b\sqrt{c \sec(a+bx)}}$$

↓ 3042

$$\frac{6}{7}d^2 \left( \frac{2}{3}d^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx - \frac{2cd(d \csc(a+bx))^{3/2}}{3b\sqrt{c \sec(a+bx)}} \right) - \frac{2cd(d \csc(a+bx))^{7/2}}{7b\sqrt{c \sec(a+bx)}}$$

↓ 3120

$$\frac{6}{7}d^2 \left( \frac{2d^2 \sqrt{\sin(2a+2bx)} \operatorname{EllipticF}\left(a+bx - \frac{\pi}{4}, 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3b} - \frac{2cd(d \csc(a+bx))^{3/2}}{3b\sqrt{c \sec(a+bx)}} \right) - \frac{2cd(d \csc(a+bx))^{7/2}}{7b\sqrt{c \sec(a+bx)}}$$

input `Int[(d*Csc[a + b*x])^(9/2)*Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*(d*Csc[a + b*x])^(7/2))/(7*b*Sqrt[c*Sec[a + b*x]]) + (6*d^2*((-2*c*d*(d*Csc[a + b*x])^(3/2))/(3*b*Sqrt[c*Sec[a + b*x]]) + (2*d^2*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(3*b)))/7`

## 3.230.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

## 3.230.4 Maple [A] (verified)

Time = 7.02 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.19

method	result
default	$\frac{\sqrt{2}d^4\sqrt{c\sec(bx+a)}\sqrt{d\csc(bx+a)}\left((4\cos(bx+a)+4)\sqrt{1+\csc(bx+a)-\cot(bx+a)}\sqrt{\cot(bx+a)-\csc(bx+a)+1}\sqrt{\cot(bx+a)-\csc(bx+a)}\right)}{7b}$

input `int((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output  $1/7/b*2^{(1/2)}*d^4*(c*\sec(b*x+a))^{(1/2)}*(d*\csc(b*x+a))^{(1/2)}*((4*\cos(b*x+a)+4)*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})+2^{(1/2)}*(2*\cot(b*x+a)^3-3*\cot(b*x+a)*\csc(b*x+a)^2))$

### 3.230.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.38

$$\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx =$$

$$2 \left( (i d^4 \cos(bx + a)^2 - i d^4) \sqrt{-4i cd} F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) \sin(bx + a) + (-i d^4 \cos(bx + a)^2 + i d^4) \sqrt{4i cd} F(\arcsin(\cos(bx + a) - i \sin(bx + a)) | -1) \sin(bx + a) \right)$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`

output  $-2/7*((I*d^4*\cos(b*x + a)^2 - I*d^4)*\text{sqrt}(-4*I*c*d)*\text{elliptic\_f}(\arcsin(\cos(b*x + a) + I*\sin(b*x + a)), -1)*\sin(b*x + a) + (-I*d^4*\cos(b*x + a)^2 + I*d^4)*\text{sqrt}(4*I*c*d)*\text{elliptic\_f}(\arcsin(\cos(b*x + a) - I*\sin(b*x + a)), -1)*\sin(b*x + a) + (2*d^4*\cos(b*x + a)^3 - 3*d^4*\cos(b*x + a))*\text{sqrt}(c/\cos(b*x + a))*\text{sqrt}(d/\sin(b*x + a)))/((b*\cos(b*x + a)^2 - b)*\sin(b*x + a))$

### 3.230.6 Sympy [F(-1)]

Timed out.

$$\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(9/2)*(c*sec(b*x+a))**(1/2),x)`

output `Timed out`

**3.230.7 Maxima [F]**

$$\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{9/2} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(9/2)*sqrt(c*sec(b*x + a)), x)`

**3.230.8 Giac [F]**

$$\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{9/2} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(9/2)*sqrt(c*sec(b*x + a)), x)`

**3.230.9 Mupad [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx = \int \sqrt{\frac{c}{\cos(a + bx)}} \left( \frac{d}{\sin(a + bx)} \right)^{9/2} dx$$

input `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(9/2),x)`

output `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(9/2), x)`



### 3.231 $\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx$

3.231.1 Optimal result . . . . .	1324
3.231.2 Mathematica [A] (verified) . . . . .	1324
3.231.3 Rubi [A] (verified) . . . . .	1325
3.231.4 Maple [A] (verified) . . . . .	1326
3.231.5 Fricas [A] (verification not implemented) . . . . .	1327
3.231.6 Sympy [F(-1)] . . . . .	1327
3.231.7 Maxima [F] . . . . .	1327
3.231.8 Giac [F] . . . . .	1328
3.231.9 Mupad [B] (verification not implemented) . . . . .	1328

#### 3.231.1 Optimal result

Integrand size = 25, antiderivative size = 69

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = -\frac{8cd^3 \sqrt{d \csc(a + bx)}}{5b \sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b \sqrt{c \sec(a + bx)}}$$

output  $-2/5*c*d*(d*\csc(b*x+a))^{(5/2)}/b/(c*\sec(b*x+a))^{(1/2)}-8/5*c*d^3*(d*\csc(b*x+a))^{(1/2)}/b/(c*\sec(b*x+a))^{(1/2)}$

#### 3.231.2 Mathematica [A] (verified)

Time = 0.90 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = \frac{2d^3 \sqrt{d \csc(a + bx)} (4 \cos(a + bx) + \cot(a + bx) \csc(a + bx)) \sqrt{c \sec(a + bx)}}{5b}$$

input `Integrate[(d*Csc[a + b*x])^(7/2)*Sqrt[c*Sec[a + b*x]],x]`

output  $(-2*d^3*Sqrt[d*Csc[a + b*x]]*(4*Cos[a + b*x] + Cot[a + b*x]*Csc[a + b*x])*Sqrt[c*Sec[a + b*x]])/(5*b)$

**3.231.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {3042, 3105, 3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{7/2} dx \\ & \quad \downarrow \text{3105} \\ & \frac{4}{5} d^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx - \frac{2cd (d \csc(a + bx))^{5/2}}{5b \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{4}{5} d^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx - \frac{2cd (d \csc(a + bx))^{5/2}}{5b \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3099} \\ & -\frac{8cd^3 \sqrt{d \csc(a + bx)}}{5b \sqrt{c \sec(a + bx)}} - \frac{2cd (d \csc(a + bx))^{5/2}}{5b \sqrt{c \sec(a + bx)}} \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(7/2)*Sqrt[c*Sec[a + b*x]],x]`

output `(-8*c*d^3*Sqrt[d*Csc[a + b*x]])/(5*b*Sqrt[c*Sec[a + b*x]]) - (2*c*d*(d*Csc[a + b*x])^(5/2))/(5*b*Sqrt[c*Sec[a + b*x]])`

## 3.231.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

## 3.231.4 Maple [A] (verified)

Time = 1.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.77

method	result	size
default	$\frac{2d^3 \sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)} (4 \cos(bx+a)^2 - 5) \cot(bx+a) \csc(bx+a)}{5b}$	53

input `int((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2/5/b*d^3*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*(4*cos(b*x+a)^2-5)*cot(b*x+a)*csc(b*x+a)`

**3.231.5 Fricas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.97

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = \frac{2(4d^3 \cos(bx + a)^3 - 5d^3 \cos(bx + a)) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{5(b \cos(bx + a)^2 - b)}$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`output `-2/5*(4*d^3*cos(b*x + a)^3 - 5*d^3*cos(b*x + a))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*cos(b*x + a)^2 - b)`**3.231.6 Sympy [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(7/2)*(c*sec(b*x+a))**(1/2),x)`output `Timed out`**3.231.7 Maxima [F]**

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{7/2} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(7/2)*sqrt(c*sec(b*x + a)), x)`

**3.231.8 Giac [F]**

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{7/2} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(7/2)*sqrt(c*sec(b*x + a)), x)`

**3.231.9 Mupad [B] (verification not implemented)**

Time = 14.06 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.23

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = \frac{4 d^3 \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (3 \cos(a + bx) - 4 \cos(3a + 3bx) + \cos(5a + 5bx))}{5b (\cos(4a + 4bx) - 4 \cos(2a + 2bx) + 3)}$$

input `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(7/2),x)`

output `-(4*d^3*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(3*cos(a + b*x) - 4*cos(3*a + 3*b*x) + cos(5*a + 5*b*x)))/(5*b*(cos(4*a + 4*b*x) - 4*cos(2*a + 2*b*x) + 3))`

### 3.232 $\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx$

3.232.1 Optimal result . . . . .	1329
3.232.2 Mathematica [C] (verified) . . . . .	1329
3.232.3 Rubi [A] (verified) . . . . .	1330
3.232.4 Maple [A] (verified) . . . . .	1332
3.232.5 Fricas [C] (verification not implemented) . . . . .	1332
3.232.6 Sympy [F(-1)] . . . . .	1333
3.232.7 Maxima [F] . . . . .	1333
3.232.8 Giac [F] . . . . .	1333
3.232.9 Mupad [F(-1)] . . . . .	1334

#### 3.232.1 Optimal result

Integrand size = 25, antiderivative size = 93

$$\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx = -\frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} + \frac{2d^2 \sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{3b}$$

output

```
-2/3*c*d*(d*csc(b*x+a))^(3/2)/b/(c*sec(b*x+a))^(1/2)-2/3*d^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b
```

#### 3.232.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.50 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.17

$$\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx = \frac{d(\cos(a + bx) + \cos(3(a + bx)))(d \csc(a + bx))^{3/2} \left( \cot^2(a + bx) + (-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{\cot^2(a + bx)}{\csc^2(a + bx)}\right) \right)}{3b(-2 + \csc^2(a + bx))}$$

input

```
Integrate[(d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]],x]
```

output 
$$-1/3*(d*(\cos[a + b*x] + \cos[3*(a + b*x)])*(d*\csc[a + b*x])^{3/2}*(\cot[a + b*x]^2 + (-\cot[a + b*x]^2)^{3/4}*\text{Hypergeometric2F1}[1/2, 3/4, 3/2, \csc[a + b*x]^2])*Sec[a + b*x]^2*\text{sqrt}[c*Sec[a + b*x]])/(b*(-2 + \csc[a + b*x]^2))$$

### 3.232.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3105, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{5/2} dx \\ & \quad \downarrow \text{3105} \\ & \frac{2}{3} d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx - \frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{2}{3} d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx - \frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3110} \\ & \frac{2}{3} d^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx - \\ & \quad \frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{2}{3} d^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx - \\ & \quad \frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3053} \end{aligned}$$

$$\frac{2}{3}d^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx-\frac{2cd(d\csc(a+bx))^{3/2}}{3b\sqrt{c\sec(a+bx)}}$$

↓ 3042

$$\frac{2}{3}d^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx-\frac{2cd(d\csc(a+bx))^{3/2}}{3b\sqrt{c\sec(a+bx)}}$$

↓ 3120

$$\frac{2d^2\sqrt{\sin(2a+2bx)}\operatorname{EllipticF}\left(a+bx-\frac{\pi}{4},2\right)\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{3b}-\frac{2cd(d\csc(a+bx))^{3/2}}{3b\sqrt{c\sec(a+bx)}}$$

input `Int[(d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*(d*Csc[a + b*x])^(3/2))/(3*b*Sqrt[c*Sec[a + b*x]]) + (2*d^2*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(3*b)`

### 3.232.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`



```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.232.4 Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.42

method	result
default	$\frac{\sqrt{2} d^2 \sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)} \left( (2 \cos(bx+a)+2) \sqrt{1+\csc(bx+a)-\cot(bx+a)} \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)} \right)}{3b}$

```
input int((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/3/b*2^(1/2)*d^2*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*((2*cos(b*x+a)+2)*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))-2^(1/2)*cot(b*x+a))
```

### 3.232.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.28

$$\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx = \frac{-i \sqrt{-4i c d d^2} F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) \sin(bx + a) + i \sqrt{4i c d d^2}}{3b}$$

```
input integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x,algorithm="fricas")
```

output `1/3*(-I*sqrt(-4*I*c*d)*d^2*elliptic_f(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1)*sin(b*x + a) + I*sqrt(4*I*c*d)*d^2*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1)*sin(b*x + a) - 2*d^2*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a))/(b*sin(b*x + a))`

### 3.232.6 Sympy [F(-1)]

Timed out.

$$\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(5/2)*(c*sec(b*x+a))**(1/2),x)`

output `Timed out`

### 3.232.7 Maxima [F]

$$\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{5/2} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a)), x)`

### 3.232.8 Giac [F]

$$\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{5/2} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a)), x)`

**3.232.9 Mupad [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx = \int \sqrt{\frac{c}{\cos(a + bx)}} \left( \frac{d}{\sin(a + bx)} \right)^{5/2} dx$$

input `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2),x)`output `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2), x)`

### 3.233 $\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx$

3.233.1 Optimal result . . . . .	1335
3.233.2 Mathematica [A] (verified) . . . . .	1335
3.233.3 Rubi [A] (verified) . . . . .	1336
3.233.4 Maple [A] (verified) . . . . .	1337
3.233.5 Fricas [A] (verification not implemented) . . . . .	1337
3.233.6 Sympy [F(-1)] . . . . .	1337
3.233.7 Maxima [F] . . . . .	1338
3.233.8 Giac [F] . . . . .	1338
3.233.9 Mupad [B] (verification not implemented) . . . . .	1338

#### 3.233.1 Optimal result

Integrand size = 25, antiderivative size = 31

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = -\frac{2cd \sqrt{d \csc(a + bx)}}{b \sqrt{c \sec(a + bx)}}$$

output `-2*c*d*(d*csc(b*x+a))^(1/2)/b/(c*sec(b*x+a))^(1/2)`

#### 3.233.2 Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = -\frac{2cd \sqrt{d \csc(a + bx)}}{b \sqrt{c \sec(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*Sqrt[d*Csc[a + b*x]])/(b*Sqrt[c*Sec[a + b*x]])`

**3.233.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2} dx$$

↓ 3042

$$\int \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2} dx$$

↓ 3099

$$-\frac{2cd\sqrt{d \csc(a + bx)}}{b\sqrt{c \sec(a + bx)}}$$

input `Int[(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*Sqrt[d*Csc[a + b*x]])/(b*Sqrt[c*Sec[a + b*x]])`

**3.233.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

**3.233.4 Maple [A] (verified)**

Time = 1.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

method	result	size
default	$-\frac{2\sqrt{c\sec(bx+a)}\sqrt{d\csc(bx+a)}d\cos(bx+a)}{b}$	33

input `int((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`output `-2/b*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*d*cos(b*x+a)`**3.233.5 Fricas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = -\frac{2d \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx + a)}{b}$$

input `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`output `-2*d*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)/b`**3.233.6 Sympy [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(3/2)*(c*sec(b*x+a))**(1/2),x)`output `Timed out`

**3.233.7 Maxima [F]**

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a)), x)`

**3.233.8 Giac [F]**

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = \int (d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a)), x)`

**3.233.9 Mupad [B] (verification not implemented)**

Time = 13.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = -\frac{2 d \cos(a + bx) \sqrt{\frac{c}{\cos(a + bx)}} \sqrt{\frac{d}{\sin(a + bx)}}}{b}$$

input `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(3/2),x)`

output `-(2*d*cos(a + b*x)*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2))/b`

### 3.234 $\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx$

3.234.1 Optimal result . . . . .	1339
3.234.2 Mathematica [C] (verified) . . . . .	1339
3.234.3 Rubi [A] (verified) . . . . .	1340
3.234.4 Maple [A] (verified) . . . . .	1341
3.234.5 Fricas [C] (verification not implemented) . . . . .	1342
3.234.6 Sympy [F] . . . . .	1342
3.234.7 Maxima [F] . . . . .	1342
3.234.8 Giac [F] . . . . .	1343
3.234.9 Mupad [F(-1)] . . . . .	1343

#### 3.234.1 Optimal result

Integrand size = 25, antiderivative size = 53

$$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx$$

$$= \frac{\sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{b}$$

output  $-(\sin(a+1/4*\pi+b*x)^2)^{(1/2)}/\sin(a+1/4*\pi+b*x)*\operatorname{EllipticF}(\cos(a+1/4*\pi+b*x), 2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b$

#### 3.234.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.26 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.28

$$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx$$

$$= \frac{(-\cot^2(a + bx))^{7/4} \sqrt{d \csc(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a + bx)\right) \sqrt{c \sec(a + bx)} \tan^3(a + bx)}{b}$$

input `Integrate[Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]],x]`

output  $((-\operatorname{Cot}[a + b*x]^2)^{(7/4)}*\operatorname{Sqrt}[d*\operatorname{Csc}[a + b*x]]*\operatorname{Hypergeometric2F1}[1/2, 3/4, 3/2, \operatorname{Csc}[a + b*x]^2]*\operatorname{Sqrt}[c*\operatorname{Sec}[a + b*x]]*\operatorname{Tan}[a + b*x]^3)/b$



**3.234.3 Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} dx \\
 & \quad \downarrow \text{3110} \\
 & \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx \\
 & \quad \downarrow \text{3053} \\
 & \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx \\
 & \quad \downarrow \text{3120} \\
 & \frac{\sqrt{\sin(2a+2bx)} \operatorname{EllipticF}\left(a+bx-\frac{\pi}{4}, 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{b}
 \end{aligned}$$

input `Int[Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]],x]`

output `(Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/b`

## 3.234.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

## 3.234.4 Maple [A] (verified)

Time = 1.09 (sec) , antiderivative size = 113, normalized size of antiderivative = 2.13

method	result
default	$\frac{\sqrt{2}(\cos(bx+a)+1)\sqrt{c\sec(bx+a)}\sqrt{d\csc(bx+a)}\sqrt{1+\csc(bx+a)-\cot(bx+a)}\sqrt{\cot(bx+a)-\csc(bx+a)+1}\sqrt{\cot(bx+a)-\csc(bx+a)}}{b}$

input `int((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `1/b*2^(1/2)*(cos(b*x+a)+1)*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))`

**3.234.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.09

$$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx$$

$$= \frac{-i \sqrt{-4i cd} F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) + i \sqrt{4i cd} F(\arcsin(\cos(bx + a) - i \sin(bx + a)) | -1)}{2b}$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`

output `1/2*(-I*sqrt(-4*I*c*d)*elliptic_f(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) + I*sqrt(4*I*c*d)*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1)/b`

**3.234.6 Sympy [F]**

$$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx = \int \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)} dx$$

input `integrate((d*csc(b*x+a))**(1/2)*(c*sec(b*x+a))**(1/2),x)`

output `Integral(sqrt(c*sec(a + b*x))*sqrt(d*csc(a + b*x)), x)`

**3.234.7 Maxima [F]**

$$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx = \int \sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a)), x)`

**3.234.8 Giac [F]**

$$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx = \int \sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} dx$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a)), x)`

**3.234.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx = \int \sqrt{\frac{c}{\cos(a + bx)}} \sqrt{\frac{d}{\sin(a + bx)}} dx$$

input `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2),x)`

output `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2), x)`

**3.235** 
$$\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx$$

3.235.1 Optimal result . . . . . 1344  
 3.235.2 Mathematica [A] (verified) . . . . . 1345  
 3.235.3 Rubi [A] (verified) . . . . . 1345  
 3.235.4 Maple [A] (warning: unable to verify) . . . . . 1349  
 3.235.5 Fricas [C] (verification not implemented) . . . . . 1350  
 3.235.6 Sympy [F] . . . . . 1350  
 3.235.7 Maxima [F] . . . . . 1351  
 3.235.8 Giac [F] . . . . . 1351  
 3.235.9 Mupad [F(-1)] . . . . . 1351

**3.235.1 Optimal result**

Integrand size = 25, antiderivative size = 270

$$\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx = -\frac{\arctan\left(1 - \sqrt{2}\sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{\sqrt{2}b\sqrt{d \csc(a+bx)}\sqrt{\tan(a+bx)}} + \frac{\arctan\left(1 + \sqrt{2}\sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{\sqrt{2}b\sqrt{d \csc(a+bx)}\sqrt{\tan(a+bx)}} + \frac{\log\left(1 - \sqrt{2}\sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{c \sec(a+bx)}}{2\sqrt{2}b\sqrt{d \csc(a+bx)}\sqrt{\tan(a+bx)}} - \frac{\log\left(1 + \sqrt{2}\sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{c \sec(a+bx)}}{2\sqrt{2}b\sqrt{d \csc(a+bx)}\sqrt{\tan(a+bx)}}$$

```
output 1/2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b*2^(1/2)/(d*
csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+1/2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*
(c*sec(b*x+a))^(1/2)/b*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+1/4*ln
(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b*2^(1/2)/(d
*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)-1/4*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(
b*x+a))*(c*sec(b*x+a))^(1/2)/b*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/
2)
```

**3.235.2 Mathematica [A] (verified)**

Time = 2.06 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.45

$$\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx = \frac{\left( \arctan \left( \frac{-1 + \sqrt{\cot^2(a+bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a+bx)}} \right) + \operatorname{arctanh} \left( \frac{\sqrt{2} \sqrt[4]{\cot^2(a+bx)}}{1 + \sqrt{\cot^2(a+bx)}} \right) \right) \cot(a+bx) \sqrt{c \sec(a+bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a+bx)} \sqrt{d \csc(a+bx)}}$$

input `Integrate[Sqrt[c*Sec[a + b*x]]/Sqrt[d*Csc[a + b*x]],x]`output `-(((ArcTan[(-1 + Sqrt[Cot[a + b*x]^2)]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]) + ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])])*Cot[a + b*x]*Sqrt[c*Sec[a + b*x]])/(Sqrt[2]*b*(Cot[a + b*x]^2)^(1/4)*Sqrt[d*Csc[a + b*x]])`**3.235.3 Rubi [A] (verified)**Time = 0.47 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.63, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.520$ , Rules used = {3042, 3109, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx \\ & \quad \downarrow \text{3109} \\ & \frac{\sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.235.  $\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx$

$$\begin{aligned}
& \frac{\sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3957} \\
& \frac{\sqrt{c \sec(a+bx)} \int \frac{\sqrt{\tan(a+bx)}}{\tan^2(a+bx)+1} d \tan(a+bx)}{b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{266} \\
& \frac{2 \sqrt{c \sec(a+bx)} \int \frac{\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{826} \\
& \frac{2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{1476} \\
& \frac{2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right) \right)}{b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{1082} \\
& \frac{2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{1}{-\tan(a+bx)-1} \frac{d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \int \frac{1}{-\tan(a+bx)-1} \frac{d(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{217} \\
& \frac{2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{1479} \\
& \frac{2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) - \arctan(1-\sqrt{2}\sqrt{\tan(a+bx)}) \right) \right)}{b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{25}
\end{aligned}$$

---

3.235.  $\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx$

$$\begin{aligned}
& \frac{2\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} \right) \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) \right)}{b\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}} \\
& \quad \downarrow 27 \\
& \frac{2\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} \right) \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) \right)}{b\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}} \\
& \quad \downarrow 1103 \\
& \frac{2\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{\log(\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} - \frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} \right)}{b\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}}
\end{aligned}$$

input `Int[Sqrt[c*Sec[a + b*x]]/Sqrt[d*Csc[a + b*x]],x]`

output `(2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]))/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[c*Sec[a + b*x]]/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])`

### 3.235.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`



- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### 3.235.4 Maple [A] (warning: unable to verify)

Time = 36.15 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.34

method	result
default	$\frac{\sqrt{2} \sqrt{c \sec(bx+a)}}{\dots} \left( \ln \left( 2\sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \cot(bx+a) + 2\sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \csc(bx+a) - 2 \cot(bx+a) + 2 \right) + 2 \arctan \left( \frac{-\sin(bx+a) \sqrt{2}}{\cos(bx+a) + 1} \right) \right)$

input `int((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output

$$-1/4/b*2^(1/2)*(c*\sec(b*x+a))^(1/2)*(ln(2*2^(1/2)*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^(1/2)*\cot(b*x+a)+2*2^(1/2)*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^(1/2)*\csc(b*x+a)-2*\cot(b*x+a)+2)+2*\arctan((-sin(b*x+a)*2^(1/2)*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^(1/2)+\cos(b*x+a)-1)/(\cos(b*x+a)-1))-\ln(-2*2^(1/2)*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^(1/2)*\cot(b*x+a)-2*2^(1/2)*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^(1/2)*\csc(b*x+a)-2*\cot(b*x+a)+2)-2*\arctan((sin(b*x+a)*2^(1/2)*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^(1/2)+\cos(b*x+a)-1)/(\cos(b*x+a)-1)))*\cos(b*x+a)/(\cos(b*x+a)+1)/(d*csc(b*x+a))^(1/2)/(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^(1/2)$$

**3.235.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.51 (sec) , antiderivative size = 1135, normalized size of antiderivative = 4.20

$$\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx = \text{Too large to display}$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x, algorithm="fricas")`

output

```
-1/8*I*(-c^2/(b^4*d^2))^(1/4)*log(2*b^2*c*d*sqrt(-c^2/(b^4*d^2))*cos(b*x +
a)*sin(b*x + a) - 2*c^2*cos(b*x + a)^2 + c^2 - 2*(I*b*c*(-c^2/(b^4*d^2))^(
1/4)*cos(b*x + a)^2*sin(b*x + a) + (I*b^3*d*cos(b*x + a)^3 - I*b^3*d*cos(
b*x + a))*(-c^2/(b^4*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a
)) + 1/8*I*(-c^2/(b^4*d^2))^(1/4)*log(2*b^2*c*d*sqrt(-c^2/(b^4*d^2))*cos(b
*x + a)*sin(b*x + a) - 2*c^2*cos(b*x + a)^2 + c^2 - 2*(-I*b*c*(-c^2/(b^4*d
^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (-I*b^3*d*cos(b*x + a)^3 + I*b^3*
d*cos(b*x + a))*(-c^2/(b^4*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*
x + a))) + 1/8*(-c^2/(b^4*d^2))^(1/4)*log(-2*b^2*c*d*sqrt(-c^2/(b^4*d^2))*
cos(b*x + a)*sin(b*x + a) - 2*c^2*cos(b*x + a)^2 + c^2 + 2*(b*c*(-c^2/(b^4
*d^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*d*cos(b*x + a)^3 - b^3*d*c
os(b*x + a))*(-c^2/(b^4*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x +
a))) - 1/8*(-c^2/(b^4*d^2))^(1/4)*log(-2*b^2*c*d*sqrt(-c^2/(b^4*d^2))*cos
(b*x + a)*sin(b*x + a) - 2*c^2*cos(b*x + a)^2 + c^2 - 2*(b*c*(-c^2/(b^4*d^
2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*d*cos(b*x + a)^3 - b^3*d*cos(
b*x + a))*(-c^2/(b^4*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a
)) + 1/8*(-c^2/(b^4*d^2))^(1/4)*log(-c^2 + 2*(b*c*(-c^2/(b^4*d^2))^(1/4)*c
os(b*x + a)^2*sin(b*x + a) + (b^3*d*cos(b*x + a)^3 - b^3*d*cos(b*x + a))*(-
c^2/(b^4*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) - 1/8*(-
c^2/(b^4*d^2))^(1/4)*log(-c^2 - 2*(b*c*(-c^2/(b^4*d^2))^(1/4)*cos(b*x + ...
```

**3.235.6 Sympy [F]**

$$\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx$$

input `integrate((c*sec(b*x+a))**(1/2)/(d*csc(b*x+a))**(1/2),x)`

output `Integral(sqrt(c*sec(a + b*x))/sqrt(d*csc(a + b*x)), x)`

---

3.235.  $\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx$

**3.235.7 Maxima [F]**

$$\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{\sqrt{c \sec(bx + a)}}{\sqrt{d \csc(bx + a)}} dx$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(c*sec(b*x + a))/sqrt(d*csc(b*x + a)), x)`

**3.235.8 Giac [F]**

$$\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{\sqrt{c \sec(bx + a)}}{\sqrt{d \csc(bx + a)}} dx$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(c*sec(b*x + a))/sqrt(d*csc(b*x + a)), x)`

**3.235.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{\sqrt{\frac{c}{\cos(a+bx)}}}{\sqrt{\frac{d}{\sin(a+bx)}}} dx$$

input `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(1/2),x)`

output `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(1/2), x)`

**3.236**  $\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{3/2}} dx$

3.236.1 Optimal result . . . . . 1352  
 3.236.2 Mathematica [C] (verified) . . . . . 1352  
 3.236.3 Rubi [A] (verified) . . . . . 1353  
 3.236.4 Maple [B] (verified) . . . . . 1355  
 3.236.5 Fracas [F] . . . . . 1355  
 3.236.6 Sympy [F] . . . . . 1356  
 3.236.7 Maxima [F] . . . . . 1356  
 3.236.8 Giac [F] . . . . . 1356  
 3.236.9 Mupad [F(-1)] . . . . . 1357

**3.236.1 Optimal result**

Integrand size = 25, antiderivative size = 93

$$\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{3/2}} dx = -\frac{c}{bd\sqrt{d \csc(a+bx)}\sqrt{c \sec(a+bx)}} + \frac{\sqrt{d \csc(a+bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a+bx)} \sqrt{\sin(2a+2bx)}}{2bd^2}$$

output

```
-c/b/d/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)-1/2*(sin(a+1/4*Pi+b*x))^2
^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a
))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b/d^2
```

**3.236.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.39 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.86

$$\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{3/2}} dx = \frac{\left(1 + \cos(2(a+bx)) + (-\cot^2(a+bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a+bx)\right)\right) (c \sec(a+bx))^{3/2}}{2bcd\sqrt{d \csc(a+bx)}}$$

input `Integrate[Sqrt[c*Sec[a + b*x]]/(d*Csc[a + b*x])^(3/2),x]`

output `-1/2*((1 + Cos[2*(a + b*x)] + (-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])*(c*Sec[a + b*x])^(3/2))/(b*c*d*Sqrt[d*Csc[a + b*x]])`

### 3.236.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3107, 3042, 3110, 3042, 3110, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3107} \\
 & \frac{\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{2d^2} - \frac{c}{bd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{2d^2} - \frac{c}{bd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} \\
 & \quad \downarrow \text{3110} \\
 & \frac{\sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx}{2d_c^2} \\
 & \quad \frac{bd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{\quad} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx}{2d_c^2} \\
 & \quad \frac{bd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{\quad}
 \end{aligned}$$

---

3.236.  $\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx$

$$\begin{array}{c}
 \downarrow \text{3053} \\
 \frac{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx}{2d^2} - \frac{c}{bd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \\
 \downarrow \text{3042} \\
 \frac{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx}{2d^2} - \frac{c}{bd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \\
 \downarrow \text{3120} \\
 \frac{\sqrt{\sin(2a+2bx)}\operatorname{EllipticF}\left(a+bx-\frac{\pi}{4}, 2\right)\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{2bd^2} - \frac{c}{bd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}
 \end{array}$$

input `Int[Sqrt[c*Sec[a + b*x]]/(d*Csc[a + b*x])^(3/2),x]`

output `-(c/(b*d*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])) + (Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(2*b*d^2)`

### 3.236.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), x] + Int[1/Sqrt[Sin[2*e + 2*f*x]], x] /; FreeQ[{a, b, e, f}, x]`

rule 3107 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.236.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs.  $2(106) = 212$ .

Time = 2.20 (sec) , antiderivative size = 239, normalized size of antiderivative = 2.57

method	result
default	$\frac{\sqrt{2} \left( -\sqrt{1+\csc(bx+a)}-\cot(bx+a) \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)} \operatorname{EllipticF}\left(\sqrt{1+\csc(bx+a)}-\cot(bx+a), \frac{\sqrt{2}}{2}\right) \right)}{\dots}$

```
input int((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/2/b*2^(1/2)*(-(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)-(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+sin(b*x+a)*2^(1/2)*cos(b*x+a)*(c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2)/d/(cos(b*x+a)-1)/(cos(b*x+a)+1)*sin(b*x+a)
```

### 3.236.5 Fracas [F]

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{3/2}} dx$$

```
input integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2),x, algorithm="fricas")
```



output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(d^2*csc(b*x + a)^2), x)`

### 3.236.6 Sympy [F]

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{\frac{3}{2}}} dx$$

input `integrate((c*sec(b*x+a))**(1/2)/(d*csc(b*x+a))**(3/2), x)`

output `Integral(sqrt(c*sec(a + b*x))/(d*csc(a + b*x))**(3/2), x)`

### 3.236.7 Maxima [F]

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{\frac{3}{2}}} dx$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2), x, algorithm="maxima")`

output `integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(3/2), x)`

### 3.236.8 Giac [F]

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{\frac{3}{2}}} dx$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2), x, algorithm="giac")`

output `integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(3/2), x)`

**3.236.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{\sqrt{\frac{c}{\cos(a+bx)}}}{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

input `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(3/2),x)`output `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(3/2), x)`

**3.237**  $\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx$

3.237.1 Optimal result . . . . . 1358  
 3.237.2 Mathematica [A] (verified) . . . . . 1359  
 3.237.3 Rubi [A] (verified) . . . . . 1359  
 3.237.4 Maple [A] (warning: unable to verify) . . . . . 1364  
 3.237.5 Fricas [C] (verification not implemented) . . . . . 1364  
 3.237.6 Sympy [F(-1)] . . . . . 1365  
 3.237.7 Maxima [F] . . . . . 1366  
 3.237.8 Giac [F] . . . . . 1366  
 3.237.9 Mupad [F(-1)] . . . . . 1366

**3.237.1 Optimal result**

Integrand size = 25, antiderivative size = 322

$$\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx = -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} - \frac{3 \arctan\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2}bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} + \frac{3 \arctan\left(1 + \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2}bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} + \frac{3 \log\left(1 - \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{c \sec(a+bx)}}{8\sqrt{2}bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} - \frac{3 \log\left(1 + \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{c \sec(a+bx)}}{8\sqrt{2}bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}$$

output

```
-1/2*c/b/d/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2)+3/8*arctan(-1+2^(1/2)
*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)
/tan(b*x+a)^(1/2)+3/8*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1
/2)/b/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+3/16*ln(1-2^(1/2)*
tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/d^2*2^(1/2)/(d*csc(b*x
+a))^(1/2)/tan(b*x+a)^(1/2)-3/16*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))
*(c*sec(b*x+a))^(1/2)/b/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)
```

**3.237.2 Mathematica [A] (verified)**

Time = 2.33 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.49

$$\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx = \frac{\left(4 \cos^2(a+bx) + 3\sqrt{2} \arctan\left(\frac{-1 + \sqrt{\cot^2(a+bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a+bx)}}\right) \cot^2(a+bx)^{3/4} + 3\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \sqrt[4]{\cot^2(a+bx)}}{1 + \sqrt{\cot^2(a+bx)}}\right)\right)}{8bd^2 \sqrt{d \csc(a+bx)}}$$

input `Integrate[Sqrt[c*Sec[a + b*x]]/(d*Csc[a + b*x])^(5/2), x]`output `-1/8*((4*Cos[a + b*x]^2 + 3*Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2])]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4)))*(Cot[a + b*x]^2)^(3/4) + 3*Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(3/4))*Sqrt[c*Sec[a + b*x]]*Tan[a + b*x])/(b*d^2*Sqrt[d*Csc[a + b*x]])`**3.237.3 Rubi [A] (verified)**Time = 0.57 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.66, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3107, 3042, 3109, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx \\ & \quad \downarrow \text{3107} \\ & \frac{3 \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{4d^2} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.237.  $\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx$

$$\begin{aligned}
& \frac{3 \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{4d^2} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3109} \\
& \frac{3 \sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{4d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{4d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3957} \\
& \frac{3 \sqrt{c \sec(a+bx)} \int \frac{\sqrt{\tan(a+bx)}}{\tan^2(a+bx)+1} d \tan(a+bx)}{4bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{266} \\
& \frac{3 \sqrt{c \sec(a+bx)} \int \frac{\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{826} \\
& \frac{3 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{1476} \\
& \frac{3 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right) \right)}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{1082} \\
& \frac{3 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\int \frac{1}{-\tan(a+bx)-1} d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(a+bx)-1} d(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c}{2bd \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{217}
\end{aligned}$$

---

3.237.  $\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx$

$$\begin{aligned}
& \frac{3\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d\sqrt{\tan(a+bx)} \right)}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \frac{c}{2bd\sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{1479} \\
& \frac{3\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) - \arctan(1-\sqrt{2}\sqrt{\tan(a+bx)}) \right) \right)}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \frac{c}{2bd\sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{25} \\
& \frac{3\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) - \arctan(1-\sqrt{2}\sqrt{\tan(a+bx)}) \right) \right)}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \frac{c}{2bd\sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{3\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) - \arctan(1-\sqrt{2}\sqrt{\tan(a+bx)}) \right) \right)}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \frac{c}{2bd\sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{1103} \\
& \frac{3\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} - \frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} \right) \right)}{2bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
& \frac{c}{2bd\sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}}
\end{aligned}$$

input `Int[Sqrt[c*Sec[a + b*x]]/(d*Csc[a + b*x])^(5/2),x]`

```
output -1/2*c/(b*d*(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]]) + (3*((-(ArcTan[1
- Sqrt[2]*Sqrt[Tan[a + b*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a +
b*x]])/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2
*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2])
)/2)*Sqrt[c*Sec[a + b*x]])/(2*b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]
])
```

### 3.237.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 266 Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = De
nominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))
^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && I
ntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 826 Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^
4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{
a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]
&& AtomQ[SplitProduct[SumBaseQ, b]]))
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Fre
eQ[{a, b, c}, x]
```

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3107 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^n, x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`



**3.237.4 Maple [A] (warning: unable to verify)**

Time = 7.35 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.44

method	result
default	$\sqrt{2} \left( 4\sqrt{2} \cos(bx+a) \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sin(bx+a) + 4 \sin(bx+a) \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} - 3 \ln \left( -2\sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \right) \right)$

input `int((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 1/16/b*2^{(1/2)}*(4*2^{(1/2)}*\cos(b*x+a)*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1) \\ & )^{(1/2)}*\sin(b*x+a)+4*\sin(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b* \\ & x+a)+1)^2)^{(1/2)}-3*\ln(-2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2) \\ & )^{(1/2)}*\cot(b*x+a)-2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} \\ & )*\csc(b*x+a)-2*\cot(b*x+a)+2)+3*\ln(2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b \\ & *x+a)+1)^2)^{(1/2)}*\cot(b*x+a)+2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a) \\ & +1)^2)^{(1/2)}*\csc(b*x+a)-2*\cot(b*x+a)+2)+6*\arctan((-\sin(b*x+a)*2^{(1/2)}*(-\cos \\ & (b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}+\cos(b*x+a)-1)/(\cos(b*x+a)-1))- \\ & 6*\arctan((\sin(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} \\ & )+\cos(b*x+a)-1)/(\cos(b*x+a)-1)))*(c*sec(b*x+a))^{(1/2)}/(\cos(b*x+a)-1)/(\cos \\ & (b*x+a)+1)^2/(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}/(d*csc(b*x+a) \\ & )^{(1/2)}/d^2*\sin(b*x+a)^2*\cos(b*x+a) \end{aligned}$$

**3.237.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 1290, normalized size of antiderivative = 4.01

$$\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2),x, algorithm="fricas")`

output

```
-1/32*(3*I*b*d^3*(-c^2/(b^4*d^10))^(1/4)*log(54*b^2*c*d^5*sqrt(-c^2/(b^4*d^10))*cos(b*x + a)*sin(b*x + a) - 54*c^2*cos(b*x + a)^2 + 27*c^2 - 54*(I*b*c*d^2*(-c^2/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (I*b^3*d^7*cos(b*x + a)^3 - I*b^3*d^7*cos(b*x + a))*(-c^2/(b^4*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) - 3*I*b*d^3*(-c^2/(b^4*d^10))^(1/4)*log(54*b^2*c*d^5*sqrt(-c^2/(b^4*d^10))*cos(b*x + a)*sin(b*x + a) - 54*c^2*cos(b*x + a)^2 + 27*c^2 - 54*(-I*b*c*d^2*(-c^2/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (-I*b^3*d^7*cos(b*x + a)^3 + I*b^3*d^7*cos(b*x + a))*(-c^2/(b^4*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) - 3*b*d^3*(-c^2/(b^4*d^10))^(1/4)*log(-54*b^2*c*d^5*sqrt(-c^2/(b^4*d^10))*cos(b*x + a)*sin(b*x + a) - 54*c^2*cos(b*x + a)^2 + 27*c^2 + 54*(b*c*d^2*(-c^2/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*d^7*cos(b*x + a)^3 - b^3*d^7*cos(b*x + a))*(-c^2/(b^4*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) + 3*b*d^3*(-c^2/(b^4*d^10))^(1/4)*log(-54*b^2*c*d^5*sqrt(-c^2/(b^4*d^10))*cos(b*x + a)*sin(b*x + a) - 54*c^2*cos(b*x + a)^2 + 27*c^2 - 54*(b*c*d^2*(-c^2/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*d^7*cos(b*x + a)^3 - b^3*d^7*cos(b*x + a))*(-c^2/(b^4*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) - 3*b*d^3*(-c^2/(b^4*d^10))^(1/4)*log(-27*c^2 + 54*(b*c*d^2*(-c^2/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*d^7*cos(b*x + a)^3 - b^3*d^7*cos(b*x + a))*(-c^2/(b^4*d^10))^(3/4))*s...
```

### 3.237.6 Sympy [**F(-1)**]

Timed out.

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((c*sec(b*x+a))**(1/2)/(d*csc(b*x+a))**(5/2), x)`

output `Timed out`

**3.237.7 Maxima [F]**

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{5/2}} dx$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(5/2), x)`

**3.237.8 Giac [F]**

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{5/2}} dx$$

input `integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(5/2), x)`

**3.237.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{\sqrt{\frac{c}{\cos(a+bx)}}}{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

input `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(5/2),x)`

output `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(5/2), x)`

### 3.238 $\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx$

3.238.1 Optimal result . . . . .	1367
3.238.2 Mathematica [A] (verified) . . . . .	1367
3.238.3 Rubi [A] (verified) . . . . .	1368
3.238.4 Maple [A] (verified) . . . . .	1369
3.238.5 Fricas [A] (verification not implemented) . . . . .	1370
3.238.6 Sympy [F(-1)] . . . . .	1370
3.238.7 Maxima [F] . . . . .	1370
3.238.8 Giac [F] . . . . .	1371
3.238.9 Mupad [B] (verification not implemented) . . . . .	1371

#### 3.238.1 Optimal result

Integrand size = 25, antiderivative size = 104

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx = \frac{64cd^5 \sqrt{c \sec(a + bx)}}{21b \sqrt{d \csc(a + bx)}} - \frac{16cd^3 (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}}{21b} - \frac{2cd (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)}}{7b}$$

```
output -16/21*c*d^3*(d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2)/b-2/7*c*d*(d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2)/b+64/21*c*d^5*(c*sec(b*x+a))^(1/2)/b/(d*csc(b*x+a))^(1/2)
```

#### 3.238.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.55

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx = \frac{2cd^5 (-32 + 8 \csc^2(a + bx) + 3 \csc^4(a + bx)) \sqrt{c \sec(a + bx)}}{21b \sqrt{d \csc(a + bx)}}$$

```
input Integrate[(d*Csc[a + b*x])^(9/2)*(c*Sec[a + b*x])^(3/2),x]
```

```
output (-2*c*d^5*(-32 + 8*Csc[a + b*x]^2 + 3*Csc[a + b*x]^4)*Sqrt[c*Sec[a + b*x]])/(21*b*Sqrt[d*Csc[a + b*x]])
```

**3.238.3 Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {3042, 3105, 3042, 3105, 3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{9/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{9/2} dx \\
 & \quad \downarrow \text{3105} \\
 & \frac{8}{7} d^2 \int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{7/2}}{7b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{7} d^2 \int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{7/2}}{7b} \\
 & \quad \downarrow \text{3105} \\
 & \frac{8}{7} d^2 \left( \frac{4}{3} d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}}{3b} \right) - \\
 & \quad \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{7/2}}{7b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{7} d^2 \left( \frac{4}{3} d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}}{3b} \right) - \\
 & \quad \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{7/2}}{7b} \\
 & \quad \downarrow \text{3099} \\
 & \frac{8}{7} d^2 \left( \frac{8cd^3 \sqrt{c \sec(a + bx)}}{3b \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}}{3b} \right) - \\
 & \quad \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{7/2}}{7b}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(9/2)*(c*Sec[a + b*x])^(3/2),x]`

```
output (-2*c*d*(d*Csc[a + b*x])^(7/2)*Sqrt[c*Sec[a + b*x]]/(7*b) + (8*d^2*((8*c*
d^3*Sqrt[c*Sec[a + b*x]])/(3*b*Sqrt[d*Csc[a + b*x]]) - (2*c*d*(d*Csc[a + b
*x])^(3/2)*Sqrt[c*Sec[a + b*x]]/(3*b)))/7
```

### 3.238.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3099 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n
_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1
)/(f*(n - 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] &&
NeQ[n, 1]
```

```
rule 3105 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n
_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n
- 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*
x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[
m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]
```

### 3.238.4 Maple [A] (verified)

Time = 1.51 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.58

method	result	size
default	$\frac{2d^4c\sqrt{c\sec(bx+a)}\sqrt{d\csc(bx+a)}\left(32\cos(bx+a)^4-56\cos(bx+a)^2+21\right)\csc(bx+a)^3}{21b}$	60

```
input int((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 2/21/b*d^4*c*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*(32*cos(b*x+a)^4-56
*cos(b*x+a)^2+21)*csc(b*x+a)^3
```

**3.238.5 Fricas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.82

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx = \frac{2(32cd^4 \cos(bx + a)^4 - 56cd^4 \cos(bx + a)^2 + 21cd^4) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{21(b \cos(bx + a)^2 - b) \sin(bx + a)}$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`output `-2/21*(32*c*d^4*cos(b*x + a)^4 - 56*c*d^4*cos(b*x + a)^2 + 21*c*d^4)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/((b*cos(b*x + a)^2 - b)*sin(b*x + a))`**3.238.6 Sympy [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(9/2)*(c*sec(b*x+a))**(3/2),x)`output `Timed out`**3.238.7 Maxima [F]**

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{9/2} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(3/2), x)`

**3.238.8 Giac [F]**

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{9/2} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(3/2), x)`

**3.238.9 Mupad [B] (verification not implemented)**

Time = 14.55 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.06

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx = \frac{16 c d^4 \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (41 \sin(a + bx) - 29 \sin(3a + 3bx) + 12 \sin(5a + 5bx) - 2 \sin(7a + 7bx))}{21 b (15 \cos(2a + 2bx) - 6 \cos(4a + 4bx) + \cos(6a + 6bx) - 10)}$$

input `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(9/2),x)`

output `-(16*c*d^4*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(41*sin(a + b*x) - 29*sin(3*a + 3*b*x) + 12*sin(5*a + 5*b*x) - 2*sin(7*a + 7*b*x)))/(21*b*(15*cos(2*a + 2*b*x) - 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) - 10))`



### 3.239 $\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx$

3.239.1 Optimal result . . . . .	1372
3.239.2 Mathematica [C] (verified) . . . . .	1372
3.239.3 Rubi [A] (verified) . . . . .	1373
3.239.4 Maple [A] (verified) . . . . .	1376
3.239.5 Fricas [C] (verification not implemented) . . . . .	1376
3.239.6 Sympy [F(-1)] . . . . .	1377
3.239.7 Maxima [F] . . . . .	1377
3.239.8 Giac [F] . . . . .	1378
3.239.9 Mupad [F(-1)] . . . . .	1378

#### 3.239.1 Optimal result

Integrand size = 25, antiderivative size = 166

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx = \frac{24cd^5 \sqrt{c \sec(a + bx)}}{5b(d \csc(a + bx))^{3/2}} - \frac{12cd^3 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{5b} - \frac{2cd(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}}{5b} - \frac{24c^2 d^4 E(a - \frac{\pi}{4} + bx | 2)}{5b \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}$$

```
output 24/5*c*d^5*(c*sec(b*x+a))^(1/2)/b/(d*csc(b*x+a))^(3/2)-2/5*c*d*(d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2)/b-12/5*c*d^3*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)/b+24/5*c^2*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

#### 3.239.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 11.95 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.69

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx = \frac{2cd^3 \sqrt{d \csc(a + bx)} \left( \cot^2(a + bx) (6 \cos(2(a + bx)) + \csc^2(a + bx)) + 12 \cos^2(a + bx) \sqrt{-\cot^2(a + bx)} \operatorname{H}_2 \right)}{5b}$$

input `Integrate[(d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(3/2),x]`

output  $(-2*c*d^3*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(\text{Cot}[a + b*x]^2*(6*\text{Cos}[2*(a + b*x)] + \text{Csc}[a + b*x]^2) + 12*\text{Cos}[a + b*x]^2*(-\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Hypergeometric2F1}[-1/2, 1/4, 1/2, \text{Csc}[a + b*x]^2])*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Tan}[a + b*x]^2)/(5*b)$

### 3.239.3 Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.480$ , Rules used = {3042, 3105, 3042, 3105, 3042, 3106, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{7/2} dx \\
 & \quad \downarrow \text{3105} \\
 & \frac{6}{5} d^2 \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{6}{5} d^2 \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3105} \\
 & \frac{6}{5} d^2 \left( 2d^2 \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx - \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b} \right) - \\
 & \quad \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{6}{5} d^2 \left( 2d^2 \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx - \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b} \right) - \\
 & \quad \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{5/2}}{5b}
 \end{aligned}$$

↓ 3106

$$\frac{6}{5}d^2 \left( 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - 2c^2 \int \frac{1}{\sqrt{d\csc(a+bx)}\sqrt{c\sec(a+bx)}} dx \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \right) - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{5/2}}{5b}$$

↓ 3042

$$\frac{6}{5}d^2 \left( 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - 2c^2 \int \frac{1}{\sqrt{d\csc(a+bx)}\sqrt{c\sec(a+bx)}} dx \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \right) - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{5/2}}{5b}$$

↓ 3110

$$\frac{6}{5}d^2 \left( 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - \frac{2c^2 \int \sqrt{c\cos(a+bx)}\sqrt{d\sin(a+bx)} dx}{\sqrt{c\cos(a+bx)}\sqrt{c\sec(a+bx)}\sqrt{d\sin(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \right) - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{5/2}}{5b}$$

↓ 3042

$$\frac{6}{5}d^2 \left( 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - \frac{2c^2 \int \sqrt{c\cos(a+bx)}\sqrt{d\sin(a+bx)} dx}{\sqrt{c\cos(a+bx)}\sqrt{c\sec(a+bx)}\sqrt{d\sin(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \right) - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{5/2}}{5b}$$

↓ 3052

$$\frac{6}{5}d^2 \left( 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - \frac{2c^2 \int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \right) - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{5/2}}{5b}$$

↓ 3042

$$\frac{6}{5}d^2 \left( 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - \frac{2c^2 \int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \right) - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{5/2}}{5b}$$

↓ 3119

$$\frac{6}{5}d^2 \left( 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - \frac{2c^2 E(a+bx - \frac{\pi}{4} | 2)}{b\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \right) - \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{5b}$$

input `Int[(d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(3/2),x]`

output `(-2*c*d*(d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]]/(5*b) + (6*d^2*((-2*c*d*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])/b + 2*d^2*((2*c*d*Sqrt[c*Sec[a + b*x]])/(b*(d*Csc[a + b*x])^(3/2)) - (2*c^2*EllipticE[a - Pi/4 + b*x, 2])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])))/5`

### 3.239.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3106 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]`

```
rule 3110 Int[(csc[(e_) + (f_)*(x_)]*(a_.))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.239.4 Maple [A] (verified)

Time = 1.15 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.40

method	result
default	$-\frac{\sqrt{2} d^3 c \sqrt{c \sec(bx+a)} \sqrt{d \csc(bx+a)} \left( (-24 \cos(bx+a) - 24) \sqrt{1 + \csc(bx+a) - \cot(bx+a)} \sqrt{\cot(bx+a) - \csc(bx+a) + 1} \sqrt{\cot(bx+a) - \csc(bx+a)} \right)}{\dots}$

```
input int((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -1/5/b*2^(1/2)*d^3*c*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*((-24*cos(b*x+a)-24)*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2)))+(12*cos(b*x+a)+12)*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*(12*cos(b*x+a)-6+csc(b*x+a)^2))
```

### 3.239.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.59

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx = \frac{2 \left( 3 (cd^3 \cos(bx + a))^2 - cd^3 \right) \sqrt{-4i cdE}(\arcsin(\cos(bx + a) + i \sin(bx + a)))}{\dots}$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

output `2/5*(3*(c*d^3*cos(b*x + a)^2 - c*d^3)*sqrt(-4*I*c*d)*elliptic_e(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) + 3*(c*d^3*cos(b*x + a)^2 - c*d^3)*sqrt(4*I*c*d)*elliptic_e(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1) - 3*(c*d^3*cos(b*x + a)^2 - c*d^3)*sqrt(-4*I*c*d)*elliptic_f(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) - 3*(c*d^3*cos(b*x + a)^2 - c*d^3)*sqrt(4*I*c*d)*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1) - (12*c*d^3*cos(b*x + a)^4 - 18*c*d^3*cos(b*x + a)^2 + 5*c*d^3)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)))/(b*cos(b*x + a)^2 - b)`

### 3.239.6 Sympy [F(-1)]

Timed out.

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(7/2)*(c*sec(b*x+a))**(3/2),x)`

output `Timed out`

### 3.239.7 Maxima [F]

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{7/2} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(3/2), x)`

**3.239.8 Giac [F]**

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{7/2} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(3/2), x)`

**3.239.9 Mupad [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{3/2} \left( \frac{d}{\sin(a + bx)} \right)^{7/2} dx$$

input `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(7/2),x)`

output `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(7/2), x)`

### 3.240 $\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx$

3.240.1 Optimal result . . . . .	1379
3.240.2 Mathematica [A] (verified) . . . . .	1379
3.240.3 Rubi [A] (verified) . . . . .	1380
3.240.4 Maple [A] (verified) . . . . .	1381
3.240.5 Fracas [A] (verification not implemented) . . . . .	1381
3.240.6 Sympy [F(-1)] . . . . .	1382
3.240.7 Maxima [F] . . . . .	1382
3.240.8 Giac [F] . . . . .	1382
3.240.9 Mupad [B] (verification not implemented) . . . . .	1383

#### 3.240.1 Optimal result

Integrand size = 25, antiderivative size = 69

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx = \frac{8cd^3 \sqrt{c \sec(a + bx)}}{3b \sqrt{d \csc(a + bx)}} - \frac{2cd(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}}{3b}$$

output `-2/3*c*d*(d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2)/b+8/3*c*d^3*(c*sec(b*x+a))^(1/2)/b/(d*csc(b*x+a))^(1/2)`

#### 3.240.2 Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx = -\frac{2cd^3(-4 + \csc^2(a + bx)) \sqrt{c \sec(a + bx)}}{3b \sqrt{d \csc(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(3/2),x]`

output `(-2*c*d^3*(-4 + Csc[a + b*x]^2)*Sqrt[c*Sec[a + b*x]])/(3*b*Sqrt[d*Csc[a + b*x]])`



**3.240.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {3042, 3105, 3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{5/2} dx \\
 & \quad \downarrow \text{3105} \\
 & \frac{4}{3} d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4}{3} d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3099} \\
 & \frac{8cd^3 \sqrt{c \sec(a + bx)}}{3b \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}}{3b}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(3/2),x]`

output `(8*c*d^3*Sqrt[c*Sec[a + b*x]])/(3*b*Sqrt[d*Csc[a + b*x]]) - (2*c*d*(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]])/(3*b)`

**3.240.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

### 3.240.4 Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.70

method	result	size
default	$-\frac{2d^2c\sqrt{c\sec(bx+a)}\sqrt{d\csc(bx+a)}(4\cos(bx+a)^2-3)\csc(bx+a)}{3b}$	48

input `int((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/3/b*d^2*c*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*(4*cos(b*x+a)^2-3)*csc(b*x+a)`

### 3.240.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.84

$$\int (d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2} dx = -\frac{2(4cd^2 \cos(bx+a)^2 - 3cd^2) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{3b \sin(bx+a)}$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

output `-2/3*(4*c*d^2*cos(b*x + a)^2 - 3*c*d^2)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*sin(b*x + a))`

**3.240.6 Sympy [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(5/2)*(c*sec(b*x+a))**(3/2),x)`output `Timed out`**3.240.7 Maxima [F]**

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{5/2} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2), x)`**3.240.8 Giac [F]**

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{5/2} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")`output `integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2), x)`

**3.240.9 Mupad [B] (verification not implemented)**

Time = 12.93 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.88

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx = \frac{2 c d^2 (2 \sin(a + bx) - \sin(3a + 3bx)) \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}}}{3 b \sin(a + bx)^2}$$

input `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(5/2),x)`output `(2*c*d^2*(2*sin(a + b*x) - sin(3*a + 3*b*x))*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2))/(3*b*sin(a + b*x)^2)`

### 3.241 $\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx$

3.241.1 Optimal result . . . . .	1384
3.241.2 Mathematica [C] (verified) . . . . .	1384
3.241.3 Rubi [A] (verified) . . . . .	1385
3.241.4 Maple [B] (verified) . . . . .	1388
3.241.5 Fricas [C] (verification not implemented) . . . . .	1388
3.241.6 Sympy [F(-1)] . . . . .	1389
3.241.7 Maxima [F] . . . . .	1389
3.241.8 Giac [F] . . . . .	1389
3.241.9 Mupad [F(-1)] . . . . .	1390

#### 3.241.1 Optimal result

Integrand size = 25, antiderivative size = 125

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx = \frac{4cd^3 \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2cd \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{b} - \frac{4c^2 d^2 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{b \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}$$

```
output 4*c*d^3*(c*sec(b*x+a))^(1/2)/b/(d*csc(b*x+a))^(3/2)-2*c*d*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)/b+4*c^2*d^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

#### 3.241.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.06 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.79

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx = \frac{2cd \sqrt{d \csc(a + bx)} \left( \cos(2(a + bx)) \cot^2(a + bx) + 2 \cos^2(a + bx) \sqrt{-\cot^2(a + bx)} \operatorname{Hypergeometric2F1} \left( \dots \right) \right)}{b}$$

input `Integrate[(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2),x]`

output `(-2*c*d*Sqrt[d*Csc[a + b*x]]*(Cos[2*(a + b*x)]*Cot[a + b*x]^2 + 2*Cos[a + b*x]^2*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]]*Tan[a + b*x]^2)/b`

### 3.241.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3105, 3042, 3106, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2} dx \\
 & \quad \downarrow \text{3105} \\
 & 2d^2 \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx - \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b} \\
 & \quad \downarrow \text{3042} \\
 & 2d^2 \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx - \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b} \\
 & \quad \downarrow \text{3106} \\
 & 2d^2 \left( \frac{2cd \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - 2c^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx \right) - \\
 & \quad \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b} \\
 & \quad \downarrow \text{3042} \\
 & 2d^2 \left( \frac{2cd \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - 2c^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx \right) - \\
 & \quad \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3110} \\
 & 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{(b(d\csc(a+bx)))^{3/2}} - \frac{2c^2 \int \sqrt{c\cos(a+bx)}\sqrt{d\sin(a+bx)}dx}{\sqrt{c\cos(a+bx)}\sqrt{c\sec(a+bx)}\sqrt{d\sin(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \\
 & \qquad \qquad \qquad \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \\
 & \downarrow \text{3042} \\
 & 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{(b(d\csc(a+bx)))^{3/2}} - \frac{2c^2 \int \sqrt{c\cos(a+bx)}\sqrt{d\sin(a+bx)}dx}{\sqrt{c\cos(a+bx)}\sqrt{c\sec(a+bx)}\sqrt{d\sin(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \\
 & \qquad \qquad \qquad \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \\
 & \downarrow \text{3052} \\
 & 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{(b(d\csc(a+bx)))^{3/2}} - \frac{2c^2 \int \sqrt{\sin(2a+2bx)}dx}{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \\
 & \qquad \qquad \qquad \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \\
 & \downarrow \text{3042} \\
 & 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{(b(d\csc(a+bx)))^{3/2}} - \frac{2c^2 \int \sqrt{\sin(2a+2bx)}dx}{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \\
 & \qquad \qquad \qquad \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b} \\
 & \downarrow \text{3119} \\
 & 2d^2 \left( \frac{2cd\sqrt{c\sec(a+bx)}}{(b(d\csc(a+bx)))^{3/2}} - \frac{2c^2 E(a+bx - \frac{\pi}{4} | 2)}{b\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} \right) - \\
 & \qquad \qquad \qquad \frac{2cd\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{b}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2),x]`

output `(-2*c*d*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])/b + 2*d^2*((2*c*d*Sqrt[c*Sec[a + b*x]])/(b*(d*Csc[a + b*x])^(3/2)) - (2*c^2*EllipticE[a - Pi/4 + b*x, 2])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]]))`

## 3.241.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3106 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]] , x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`



### 3.241.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 353 vs. 2(136) = 272.

Time = 1.16 (sec) , antiderivative size = 354, normalized size of antiderivative = 2.83

method	result
default	$\frac{\sqrt{2} \left( -\frac{c((1-\cos(bx+a))^2 \csc(bx+a)^2 + 1)}{(1-\cos(bx+a))^2 \csc(bx+a)^2 - 1} \right)^{\frac{3}{2}} \left( (1-\cos(bx+a))^2 \csc(bx+a)^2 - 1 \right) \left( \frac{d((1-\cos(bx+a))^2 \csc(bx+a) + \sin(bx+a))}{1-\cos(bx+a)} \right)^{\frac{3}{2}}}{(1-\cos(bx+a))^2 \csc(bx+a)^2 - 1}$

input `int((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/b*2^{(1/2)}*(-c*((1-\cos(b*x+a))^2*\csc(b*x+a)^2+1)/((1-\cos(b*x+a))^2*\csc(b \\ & *x+a)^2-1))^{(3/2)}*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*(d/(1-\cos(b*x+a)))*((1- \\ & \cos(b*x+a))^2*\csc(b*x+a)+\sin(b*x+a))^{(3/2)}*(1-\cos(b*x+a))*(4*(1+\csc(b*x+a) \\ & )-\cot(b*x+a))^{(1/2)}*(2-2*\csc(b*x+a)+2*\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b* \\ & x+a))^{(1/2)}*\text{EllipticE}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})-2*(1+\cs \\ & c(b*x+a)-\cot(b*x+a))^{(1/2)}*(2-2*\csc(b*x+a)+2*\cot(b*x+a))^{(1/2)}*(\cot(b*x+a) \\ & -\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})+ \\ & 3*(1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)/((1-\cos(b*x+a))^2*\csc(b*x+a)^2+1)^3*\csc \\ & (b*x+a) \end{aligned}$$

### 3.241.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.26

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx = \frac{\sqrt{-4i} \operatorname{cd} \operatorname{cd} E(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) + \sqrt{4i} \operatorname{cd} \operatorname{cd} E(\arcsin(\cos(bx + a) - i \sin(bx + a)) | -1)}{2}$$

input `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

```
output (sqrt(-4*I*c*d)*c*d*elliptic_e(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1)
+ sqrt(4*I*c*d)*c*d*elliptic_e(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1)
- sqrt(-4*I*c*d)*c*d*elliptic_f(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1)
- sqrt(4*I*c*d)*c*d*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1)
- 2*(2*c*d*cos(b*x + a)^2 - c*d)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)
))/b
```

### 3.241.6 Sympy [F(-1)]

Timed out.

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx = \text{Timed out}$$

```
input integrate((d*csc(b*x+a))**(3/2)*(c*sec(b*x+a))**(3/2),x)
```

```
output Timed out
```

### 3.241.7 Maxima [F]

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{3/2} (c \sec(bx + a))^{3/2} dx$$

```
input integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")
```

```
output integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2), x)
```

### 3.241.8 Giac [F]

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx = \int (d \csc(bx + a))^{3/2} (c \sec(bx + a))^{3/2} dx$$

```
input integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")
```

```
output integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2), x)
```

**3.241.9 Mupad [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{3/2} \left( \frac{d}{\sin(a + bx)} \right)^{3/2} dx$$

input `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2),x)`output `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2), x)`

### 3.242 $\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx$

3.242.1 Optimal result . . . . .	1391
3.242.2 Mathematica [A] (verified) . . . . .	1391
3.242.3 Rubi [A] (verified) . . . . .	1392
3.242.4 Maple [A] (verified) . . . . .	1393
3.242.5 Fricas [A] (verification not implemented) . . . . .	1393
3.242.6 Sympy [F(-1)] . . . . .	1393
3.242.7 Maxima [F] . . . . .	1394
3.242.8 Giac [F] . . . . .	1394
3.242.9 Mupad [B] (verification not implemented) . . . . .	1394

#### 3.242.1 Optimal result

Integrand size = 25, antiderivative size = 31

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \frac{2cd\sqrt{c \sec(a + bx)}}{b\sqrt{d \csc(a + bx)}}$$

output `2*c*d*(c*sec(b*x+a))^(1/2)/b/(d*csc(b*x+a))^(1/2)`

#### 3.242.2 Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \frac{2cd\sqrt{c \sec(a + bx)}}{b\sqrt{d \csc(a + bx)}}$$

input `Integrate[Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2),x]`

output `(2*c*d*Sqrt[c*Sec[a + b*x]])/(b*Sqrt[d*Csc[a + b*x]])`

**3.242.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)} dx$$

↓ 3042

$$\int (c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)} dx$$

↓ 3099

$$\frac{2cd\sqrt{c \sec(a + bx)}}{b\sqrt{d \csc(a + bx)}}$$

input `Int[Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2),x]`

output `(2*c*d*Sqrt[c*Sec[a + b*x]])/(b*Sqrt[d*Csc[a + b*x]])`

**3.242.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

**3.242.4 Maple [A] (verified)**

Time = 1.15 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

method	result	size
default	$\frac{2\sqrt{c \sec(bx+a)} \sqrt{d \csc(bx+a)} c \sin(bx+a)}{b}$	33

input `int((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`output `2/b*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*c*sin(b*x+a)`**3.242.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \frac{2c \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \sin(bx+a)}{b}$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`output `2*c*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*sin(b*x + a)/b`**3.242.6 Sympy [F(-1)]**

Timed out.

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(1/2)*(c*sec(b*x+a))**(3/2),x)`output `Timed out`

**3.242.7 Maxima [F]**

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \int \sqrt{d \csc(bx + a)} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2), x)`

**3.242.8 Giac [F]**

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \int \sqrt{d \csc(bx + a)} (c \sec(bx + a))^{3/2} dx$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2), x)`

**3.242.9 Mupad [B] (verification not implemented)**

Time = 12.52 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \frac{2 c \sin(a + bx) \sqrt{\frac{c}{\cos(a + bx)}} \sqrt{\frac{d}{\sin(a + bx)}}}{b}$$

input `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(1/2),x)`

output `(2*c*sin(a + b*x)*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2))/b`

**3.243**  $\int \frac{(c \sec(a+bx))^{3/2}}{\sqrt{d \csc(a+bx)}} dx$

3.243.1 Optimal result . . . . . 1395  
 3.243.2 Mathematica [C] (verified) . . . . . 1395  
 3.243.3 Rubi [A] (verified) . . . . . 1396  
 3.243.4 Maple [B] (verified) . . . . . 1398  
 3.243.5 Fricas [C] (verification not implemented) . . . . . 1398  
 3.243.6 Sympy [F] . . . . . 1399  
 3.243.7 Maxima [F] . . . . . 1399  
 3.243.8 Giac [F] . . . . . 1399  
 3.243.9 Mupad [F(-1)] . . . . . 1400

**3.243.1 Optimal result**

Integrand size = 25, antiderivative size = 89

$$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx = \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2c^2 E(a - \frac{\pi}{4} + bx | 2)}{b\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}\sqrt{\sin(2a + 2bx)}}$$

output `2*c*d*(c*sec(b*x+a))^(1/2)/b/(d*csc(b*x+a))^(3/2)+2*c^2*(sin(a+1/4*Pi+b*x)  
 ^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/(d*csc(  
 b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)`

**3.243.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.85 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.74

$$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx = \frac{2cd \left( -1 + \sqrt{-\cot^2(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{4}, \frac{1}{2}, \csc^2(a + bx)\right)} \right) \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}}$$

input `Integrate[(c*Sec[a + b*x])^(3/2)/Sqrt[d*Csc[a + b*x]],x]`



output  $(-2*c*d*(-1 + (-\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Hypergeometric2F1}[-1/2, 1/4, 1/2, \text{Cs}[a + b*x]^2])*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(b*(d*\text{Csc}[a + b*x])^{(3/2)})$

### 3.243.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3106, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx \\ & \quad \downarrow \text{3106} \\ & \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - 2c^2 \int \frac{1}{\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}} dx \\ & \quad \downarrow \text{3042} \\ & \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - 2c^2 \int \frac{1}{\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}} dx \\ & \quad \downarrow \text{3110} \\ & \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2c^2 \int \sqrt{c \cos(a + bx)}\sqrt{d \sin(a + bx)} dx}{\sqrt{c \cos(a + bx)}\sqrt{c \sec(a + bx)}\sqrt{d \sin(a + bx)}\sqrt{d \csc(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2c^2 \int \sqrt{c \cos(a + bx)}\sqrt{d \sin(a + bx)} dx}{\sqrt{c \cos(a + bx)}\sqrt{c \sec(a + bx)}\sqrt{d \sin(a + bx)}\sqrt{d \csc(a + bx)}} \\ & \quad \downarrow \text{3052} \\ & \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2c^2 \int \sqrt{\sin(2a + 2bx)} dx}{\sqrt{\sin(2a + 2bx)}\sqrt{c \sec(a + bx)}\sqrt{d \csc(a + bx)}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.243.  $\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx$

$$\frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - \frac{2c^2 \int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}$$

↓ 3119

$$\frac{2cd\sqrt{c\sec(a+bx)}}{b(d\csc(a+bx))^{3/2}} - \frac{2c^2 E(a+bx - \frac{\pi}{4} | 2)}{b\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}$$

input `Int[(c*Sec[a + b*x])^(3/2)/Sqrt[d*Csc[a + b*x]],x]`

output `(2*c*d*Sqrt[c*Sec[a + b*x]])/(b*(d*Csc[a + b*x])^(3/2)) - (2*c^2*EllipticE[a - Pi/4 + b*x, 2])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])`

### 3.243.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3106 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]] , x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

**3.243.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 354 vs.  $2(104) = 208$ .

Time = 1.22 (sec) , antiderivative size = 355, normalized size of antiderivative = 3.99

method	result
default	$\frac{\sqrt{2} \left( -\frac{c((1-\cos(bx+a))^2 \csc(bx+a)^2 + 1)}{(1-\cos(bx+a))^2 \csc(bx+a)^2 - 1} \right)^{\frac{3}{2}} \left( (1-\cos(bx+a))^2 \csc(bx+a)^2 - 1 \right) \left( 2\sqrt{1+\csc(bx+a)-\cot(bx+a)} \sqrt{2-2\csc(bx+a)+2\cot(bx+a)} \right)}{\dots}$

input `int((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/b*2^(1/2)*(-c*((1-cos(b*x+a))^2*csc(b*x+a)^2+1)/((1-cos(b*x+a))^2*csc(b*x+a)^2-1))^(3/2)*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*(2*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(2-2*csc(b*x+a)+2*cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))-(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(2-2*csc(b*x+a)+2*cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+2*(1-cos(b*x+a))^2*csc(b*x+a)^2)/((1-cos(b*x+a))^2*csc(b*x+a)^2+1)/(d/(1-cos(b*x+a)))*((1-cos(b*x+a))^2*csc(b*x+a)+sin(b*x+a))^(1/2)/(1-cos(b*x+a))*sin(b*x+a)`

**3.243.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.74

$$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx = \frac{\sqrt{-4i} \operatorname{cd}cE(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) + \sqrt{4i} \operatorname{cd}cE(\arcsin(\cos(bx + a) - i \sin(bx + a)) | -1)}{\dots}$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x, algorithm="fricas")`

output `1/2*(sqrt(-4*I*c*d)*c*elliptic_e(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) + sqrt(4*I*c*d)*c*elliptic_e(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1) - sqrt(-4*I*c*d)*c*elliptic_f(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) - sqrt(4*I*c*d)*c*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1) - 4*(c*cos(b*x + a)^2 - c)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)))/(b*d)`

$$3.243. \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx$$

**3.243.6 Sympy [F]**

$$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx$$

input `integrate((c*sec(b*x+a))**(3/2)/(d*csc(b*x+a))**(1/2),x)`

output `Integral((c*sec(a + b*x))**(3/2)/sqrt(d*csc(a + b*x)), x)`

**3.243.7 Maxima [F]**

$$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{(c \sec(bx + a))^{3/2}}{\sqrt{d \csc(bx + a)}} dx$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(3/2)/sqrt(d*csc(b*x + a)), x)`

**3.243.8 Giac [F]**

$$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{(c \sec(bx + a))^{3/2}}{\sqrt{d \csc(bx + a)}} dx$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(3/2)/sqrt(d*csc(b*x + a)), x)`

**3.243.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}}{\sqrt{\frac{d}{\sin(a+bx)}}} dx$$

input `int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(1/2),x)`output `int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(1/2), x)`

**3.244**  $\int \frac{(c \sec(a+bx))^{3/2}}{(d \csc(a+bx))^{3/2}} dx$

3.244.1 Optimal result . . . . . 1401  
 3.244.2 Mathematica [A] (verified) . . . . . 1402  
 3.244.3 Rubi [A] (verified) . . . . . 1402  
 3.244.4 Maple [B] (warning: unable to verify) . . . . . 1407  
 3.244.5 Fracas [C] (verification not implemented) . . . . . 1407  
 3.244.6 Sympy [F] . . . . . 1408  
 3.244.7 Maxima [F] . . . . . 1409  
 3.244.8 Giac [F] . . . . . 1409  
 3.244.9 Mupad [F(-1)] . . . . . 1409

**3.244.1 Optimal result**

Integrand size = 25, antiderivative size = 327

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx = \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} + \frac{c^2 \arctan\left(1 - \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{\sqrt{2}bd^2 \sqrt{c \sec(a + bx)}} + \frac{c^2 \arctan\left(1 + \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{\sqrt{2}bd^2 \sqrt{c \sec(a + bx)}} + \frac{c^2 \sqrt{d \csc(a + bx)} \log\left(1 - \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{2\sqrt{2}bd^2 \sqrt{c \sec(a + bx)}} - \frac{c^2 \sqrt{d \csc(a + bx)} \log\left(1 + \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{2\sqrt{2}bd^2 \sqrt{c \sec(a + bx)}}$$

output

```
2*c*(c*sec(b*x+a))^(1/2)/b/d/(d*csc(b*x+a))^(1/2)-1/2*c^2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/2*c^2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)+1/4*c^2*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/4*c^2*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)
```

### 3.244.2 Mathematica [A] (verified)

Time = 2.53 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.43

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx = \frac{c \left( 4 + \sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}} \right) \sqrt[4]{\cot^2(a + bx)} - \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}{1 + \sqrt{\cot^2(a + bx)}} \right) \right)}{2bd \sqrt{d \csc(a + bx)}}$$

input `Integrate[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(3/2), x]`

output `(c*(4 + Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2)]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(1/4) - Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(1/4))*Sqrt[c*Sec[a + b*x]])/(2*b*d*Sqrt[d*Csc[a + b*x]])`

### 3.244.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.65, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3104, 3042, 3109, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx \\ & \quad \downarrow \text{3104} \\ & \frac{2c \sqrt{c \sec(a + bx)}}{bd \sqrt{d \csc(a + bx)}} - \frac{c^2 \int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx}{d^2} \\ & \quad \downarrow \text{3042} \\ & \frac{2c \sqrt{c \sec(a + bx)}}{bd \sqrt{d \csc(a + bx)}} - \frac{c^2 \int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx}{d^2} \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3109} \\
& \frac{2c\sqrt{c\sec(a+bx)}}{bd\sqrt{d\csc(a+bx)}} - \frac{c^2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}}dx}{d^2\sqrt{c\sec(a+bx)}} \\
& \downarrow \text{3042} \\
& \frac{2c\sqrt{c\sec(a+bx)}}{bd\sqrt{d\csc(a+bx)}} - \frac{c^2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}}dx}{d^2\sqrt{c\sec(a+bx)}} \\
& \downarrow \text{3957} \\
& \frac{2c\sqrt{c\sec(a+bx)}}{bd\sqrt{d\csc(a+bx)}} - \frac{c^2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}(\tan^2(a+bx)+1)}d\tan(a+bx)}{bd^2\sqrt{c\sec(a+bx)}} \\
& \downarrow \text{266} \\
& \frac{2c\sqrt{c\sec(a+bx)}}{bd\sqrt{d\csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}}{bd^2\sqrt{c\sec(a+bx)}} \\
& \downarrow \text{755} \\
& \frac{2c\sqrt{c\sec(a+bx)}}{bd\sqrt{d\csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)} + \frac{1}{2}\int\frac{\tan(a+bx)+1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}\right)}{bd^2\sqrt{c\sec(a+bx)}} \\
& \downarrow \text{1476} \\
& \frac{2c\sqrt{c\sec(a+bx)}}{bd\sqrt{d\csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)} + \frac{1}{2}\left(\frac{1}{2}\int\frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{bd^2\sqrt{c\sec(a+bx)}} \\
& \downarrow \text{1082} \\
& \frac{2c\sqrt{c\sec(a+bx)}}{bd\sqrt{d\csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)} + \frac{1}{2}\left(\frac{\int\frac{1}{-\tan(a+bx)-1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \frac{\int\frac{1}{-\tan(a+bx)+1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{bd^2\sqrt{c\sec(a+bx)}} \\
& \downarrow \text{217}
\end{aligned}$$

---

3.244.  $\int \frac{(c\sec(a+bx))^{3/2}}{(d\csc(a+bx))^{3/2}} dx$



$$\frac{2c\sqrt{c \sec(a+bx)}}{bd\sqrt{d \csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)} + \frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{bd^2\sqrt{c \sec(a+bx)}}$$

↓ 1479

$$\frac{2c\sqrt{c \sec(a+bx)}}{bd\sqrt{d \csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}\left(\frac{1}{2}\left(-\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)} - \int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{bd^2\sqrt{c \sec(a+bx)}}$$

↓ 25

$$\frac{2c\sqrt{c \sec(a+bx)}}{bd\sqrt{d \csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}\left(\frac{1}{2}\left(\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)} + \int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{bd^2\sqrt{c \sec(a+bx)}}$$

↓ 27

$$\frac{2c\sqrt{c \sec(a+bx)}}{bd\sqrt{d \csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}\left(\frac{1}{2}\left(\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)} + \frac{1}{2}\int\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{bd^2\sqrt{c \sec(a+bx)}}$$

↓ 1103

$$\frac{2c\sqrt{c \sec(a+bx)}}{bd\sqrt{d \csc(a+bx)}} - \frac{2c^2\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)}\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right) + \frac{1}{2}\left(\frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}}\right)\right)}{bd^2\sqrt{c \sec(a+bx)}}$$

input `Int[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(3/2), x]`

```
output (2*c*Sqrt[c*Sec[a + b*x]]/(b*d*Sqrt[d*Csc[a + b*x]]) - (2*c^2*Sqrt[d*Csc[
a + b*x]]*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 +
Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[a
+ b*x]] + Tan[a + b*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Ta
n[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[Tan[a + b*x]]/(b*d^2*Sqrt[c*Sec[a + b*x]
])
```

### 3.244.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 266 Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = De
nominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))
^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && I
ntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 755 Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4)
, x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Fre
eQ[{a, b, c}, x]
```

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3104 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n_, x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(f*a*(n - 1))), x] + Simp[b^2*((m + 1)/(a^2*(n - 1))) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[n, 1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n_, x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^n_, x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.244.4 Maple [B] (warning: unable to verify)**

Leaf count of result is larger than twice the leaf count of optimal. 684 vs.  $2(271) = 542$ .

Time = 6.83 (sec) , antiderivative size = 685, normalized size of antiderivative = 2.09

method	result
default	$\sqrt{2} \left( -\frac{c((1-\cos(bx+a))^2 \csc(bx+a)^2+1)}{(1-\cos(bx+a))^2 \csc(bx+a)^2-1} \right)^{\frac{3}{2}} \left( (1-\cos(bx+a))^2 \csc(bx+a)^2-1 \right) \left( \ln \left( \frac{(1-\cos(bx+a))^2 \csc(bx+a)+2\sqrt{(1-\cos(bx+a))((1-\cos(bx+a))^2 \csc(bx+a)^2-1)}}{(1-\cos(bx+a))^2 \csc(bx+a)^2-1} \right) \right)$

input `int((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output

```
1/4/b*2^(1/2)*(-c*((1-cos(b*x+a))^2*csc(b*x+a)^2+1)/((1-cos(b*x+a))^2*csc(b*x+a)^2-1))^(3/2)*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*(ln(1/(1-cos(b*x+a)))*((1-cos(b*x+a))^2*csc(b*x+a)+2*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)*sin(b*x+a)-2*cos(b*x+a)+2-sin(b*x+a)))*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)-2*arctan(1/(1-cos(b*x+a))*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)*sin(b*x+a)-cos(b*x+a)+1))*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)-ln(-1/(1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)+2*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)*sin(b*x+a)+2*cos(b*x+a)-2+sin(b*x+a)))*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)-2*arctan(1/(1-cos(b*x+a))*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)*sin(b*x+a)+cos(b*x+a)-1))*((1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)^2-1)*csc(b*x+a))^(1/2)+8*cot(b*x+a)-8*csc(b*x+a))/(1-cos(b*x+a))^2*sin(b*x+a)^2/(d/(1-cos(b*x+a))*((1-cos(b*x+a))^2*csc(b*x+a)+sin(b*x+a)))^(3/2)
```

**3.244.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 1310, normalized size of antiderivative = 4.01

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x, algorithm="fricas")`

output

```
-1/8*(b*d^2*(-c^6/(b^4*d^6))^(1/4)*log(1/2*c^5*cos(b*x + a)*sin(b*x + a) +
  1/2*(b*c^3*d*(-c^6/(b^4*d^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*d^
  4*cos(b*x + a)^3 - b^3*d^4*cos(b*x + a))*(-c^6/(b^4*d^6))^(3/4))*sqrt(c/co
  s(b*x + a))*sqrt(d/sin(b*x + a)) + 1/4*(2*b^2*c^2*d^3*cos(b*x + a)^2 - b^2
  *c^2*d^3)*sqrt(-c^6/(b^4*d^6))) - b*d^2*(-c^6/(b^4*d^6))^(1/4)*log(1/2*c^5
  *cos(b*x + a)*sin(b*x + a) - 1/2*(b*c^3*d*(-c^6/(b^4*d^6))^(1/4)*cos(b*x +
  a)^2*sin(b*x + a) + (b^3*d^4*cos(b*x + a)^3 - b^3*d^4*cos(b*x + a))*(-c^6
  /(b^4*d^6))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1/4*(2*b^2*
  c^2*d^3*cos(b*x + a)^2 - b^2*c^2*d^3)*sqrt(-c^6/(b^4*d^6))) + I*b*d^2*(-c^
  6/(b^4*d^6))^(1/4)*log(1/2*c^5*cos(b*x + a)*sin(b*x + a) + 1/2*(I*b*c^3*d*
  (-c^6/(b^4*d^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (I*b^3*d^4*cos(b*x +
  a)^3 - I*b^3*d^4*cos(b*x + a))*(-c^6/(b^4*d^6))^(3/4))*sqrt(c/cos(b*x + a)
  )*sqrt(d/sin(b*x + a)) - 1/4*(2*b^2*c^2*d^3*cos(b*x + a)^2 - b^2*c^2*d^3)*
  sqrt(-c^6/(b^4*d^6))) - I*b*d^2*(-c^6/(b^4*d^6))^(1/4)*log(1/2*c^5*cos(b*x
  + a)*sin(b*x + a) + 1/2*(-I*b*c^3*d*(-c^6/(b^4*d^6))^(1/4)*cos(b*x + a)^2
  *sin(b*x + a) - (-I*b^3*d^4*cos(b*x + a)^3 + I*b^3*d^4*cos(b*x + a))*(-c^6
  /(b^4*d^6))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 1/4*(2*b^2*
  c^2*d^3*cos(b*x + a)^2 - b^2*c^2*d^3)*sqrt(-c^6/(b^4*d^6))) - b*d^2*(-c^6/
  (b^4*d^6))^(1/4)*log(c^5 + 2*(b^3*d^4*(-c^6/(b^4*d^6))^(3/4)*cos(b*x + a)^
  2*sin(b*x + a) + (b*c^3*d*cos(b*x + a)^3 - b*c^3*d*cos(b*x + a))*(-c^6/...
```

### 3.244.6 Sympy [F]

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx$$

input `integrate((c*sec(b*x+a))**(3/2)/(d*csc(b*x+a))**(3/2), x)`

output `Integral((c*sec(a + b*x))**(3/2)/(d*csc(a + b*x))**(3/2), x)`

**3.244.7 Maxima [F]**

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{(c \sec(bx + a))^{3/2}}{(d \csc(bx + a))^{3/2}} dx$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(3/2), x)`

**3.244.8 Giac [F]**

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{(c \sec(bx + a))^{3/2}}{(d \csc(bx + a))^{3/2}} dx$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(3/2), x)`

**3.244.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}}{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

input `int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(3/2),x)`

output `int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(3/2), x)`

**3.245**  $\int \frac{(c \sec(a+bx))^{3/2}}{(d \csc(a+bx))^{5/2}} dx$

3.245.1 Optimal result . . . . . 1410  
 3.245.2 Mathematica [C] (verified) . . . . . 1410  
 3.245.3 Rubi [A] (verified) . . . . . 1411  
 3.245.4 Maple [B] (verified) . . . . . 1413  
 3.245.5 Fricas [F] . . . . . 1413  
 3.245.6 Sympy [F(-1)] . . . . . 1414  
 3.245.7 Maxima [F] . . . . . 1414  
 3.245.8 Giac [F] . . . . . 1414  
 3.245.9 Mupad [F(-1)] . . . . . 1415

**3.245.1 Optimal result**

Integrand size = 25, antiderivative size = 94

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx = \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{3c^2 E(a - \frac{\pi}{4} + bx | 2)}{bd^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}$$

output

```
2*c*(c*sec(b*x+a))^(1/2)/b/d/(d*csc(b*x+a))^(3/2)+3*c^2*(sin(a+1/4*Pi+b*x)
^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/d^2/(d*
csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

**3.245.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.24 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.73

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx = \frac{c \left( -2 + 3 \sqrt{-\cot^2(a + bx)} \operatorname{Hypergeometric2F1} \left( -\frac{1}{2}, \frac{1}{4}, \frac{1}{2}, \csc^2(a + bx) \right) \right) \sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}}$$

input `Integrate[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(5/2),x]`

output `-((c*(-2 + 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(b*d*(d*Csc[a + b*x])^(3/2)))`

### 3.245.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3104, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3104} \\ & \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{3c^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx}{d^2} \\ & \quad \downarrow \text{3042} \\ & \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{3c^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx}{d^2} \\ & \quad \downarrow \text{3110} \\ & \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{3c^2 \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{d^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{3c^2 \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{d^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} \\ & \quad \downarrow \text{3052} \\ & \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{3c^2 \int \sqrt{\sin(2a + 2bx)} dx}{d^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} \end{aligned}$$

---

3.245.  $\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx$



$$\begin{array}{c} \downarrow 3042 \\ \frac{2c\sqrt{c\sec(a+bx)}}{bd(d\csc(a+bx))^{3/2}} - \frac{3c^2 \int \sqrt{\sin(2a+2bx)} dx}{d^2 \sqrt{\sin(2a+2bx)} \sqrt{c\sec(a+bx)} \sqrt{d\csc(a+bx)}} \\ \downarrow 3119 \\ \frac{2c\sqrt{c\sec(a+bx)}}{bd(d\csc(a+bx))^{3/2}} - \frac{3c^2 E\left(a+bx - \frac{\pi}{4} \mid 2\right)}{bd^2 \sqrt{\sin(2a+2bx)} \sqrt{c\sec(a+bx)} \sqrt{d\csc(a+bx)}} \end{array}$$

input `Int[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(5/2),x]`

output `(2*c*Sqrt[c*Sec[a + b*x]])/(b*d*(d*Csc[a + b*x])^(3/2)) - (3*c^2*EllipticE[a - Pi/4 + b*x, 2])/(b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])`

### 3.245.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3104 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(f*a*(n - 1))), x] + Simp[b^2*((m + 1)/(a^2*(n - 1))) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[n, 1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

### 3.245.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 395 vs.  $2(109) = 218$ .

Time = 1.06 (sec) , antiderivative size = 396, normalized size of antiderivative = 4.21

method	result
default	$\frac{\sqrt{2} \sqrt{c \sec(bx+a)} c \left(6\sqrt{1+\csc(bx+a)} - \cot(bx+a)\right) \sqrt{\cot(bx+a) - \csc(bx+a)+1} \sqrt{\cot(bx+a) - \csc(bx+a)}}{\text{EllipticE}\left(\sqrt{1+\csc(bx+a)} - \cot(bx+a)\right)}$

input `int((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `1/2/b*2^(1/2)*(c*sec(b*x+a))^(1/2)*c*(6*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)-3*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)+6*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))-3*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*cos(b*x+a)^2-3*2^(1/2)*cos(b*x+a)+2*2^(1/2))/d^2/(d*csc(b*x+a))^(1/2)*csc(b*x+a)`

### 3.245.5 Fracas [F]

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{(c \sec(bx + a))^{\frac{3}{2}}}{(d \csc(bx + a))^{\frac{5}{2}}} dx$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x, algorithm="fricas")`

output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c*sec(b*x + a)/(d^3*csc(b*x + a)^3), x)`

---

3.245.  $\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx$

**3.245.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((c*sec(b*x+a))**(3/2)/(d*csc(b*x+a))**(5/2),x)`output `Timed out`**3.245.7 Maxima [F]**

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{(c \sec(bx + a))^{3/2}}{(d \csc(bx + a))^{5/2}} dx$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x, algorithm="maxima")`output `integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(5/2), x)`**3.245.8 Giac [F]**

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{(c \sec(bx + a))^{3/2}}{(d \csc(bx + a))^{5/2}} dx$$

input `integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x, algorithm="giac")`output `integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(5/2), x)`

**3.245.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}}{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

input `int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(5/2),x)`output `int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(5/2), x)`

### 3.246 $\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx$

3.246.1 Optimal result . . . . .	1416
3.246.2 Mathematica [C] (verified) . . . . .	1416
3.246.3 Rubi [A] (verified) . . . . .	1417
3.246.4 Maple [A] (verified) . . . . .	1420
3.246.5 Fricas [C] (verification not implemented) . . . . .	1420
3.246.6 Sympy [F(-1)] . . . . .	1421
3.246.7 Maxima [F] . . . . .	1421
3.246.8 Giac [F] . . . . .	1421
3.246.9 Mupad [F(-1)] . . . . .	1422

#### 3.246.1 Optimal result

Integrand size = 25, antiderivative size = 166

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx = \frac{40cd^5(c \sec(a + bx))^{3/2}}{21b\sqrt{d \csc(a + bx)}} - \frac{20cd^3(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}}{21b} - \frac{2cd(d \csc(a + bx))^{7/2}(c \sec(a + bx))^{3/2}}{7b} + \frac{40c^2d^4\sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{21b}$$

```
output -20/21*c*d^3*(d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2)/b-2/7*c*d*(d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(3/2)/b+40/21*c*d^5*(c*sec(b*x+a))^(3/2)/b/(d*csc(b*x+a))^(1/2)-40/21*c^2*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b
```

#### 3.246.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.39 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.55

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx = \frac{2cd^5 \left( -7 + \cot^2(a + bx) (13 + 3 \csc^2(a + bx)) + 20(-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a + bx)\right) \right)}{21b\sqrt{d \csc(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(9/2)*(c*Sec[a + b*x])^(5/2),x]`

output `(-2*c*d^5*(-7 + Cot[a + b*x]^2*(13 + 3*Csc[a + b*x]^2) + 20*(-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])*(c*Sec[a + b*x])^(3/2))/(21*b*Sqrt[d*Csc[a + b*x]])`

### 3.246.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.480$ , Rules used = {3042, 3105, 3042, 3105, 3042, 3106, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{9/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{9/2} dx \\
 & \quad \downarrow \text{3105} \\
 & \frac{10}{7} d^2 \int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{7/2}}{7b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{10}{7} d^2 \int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{7/2}}{7b} \\
 & \quad \downarrow \text{3105} \\
 & \frac{10}{7} d^2 \left( 2d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}{3b} \right) - \\
 & \quad \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{7/2}}{7b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{10}{7} d^2 \left( 2d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}{3b} \right) - \\
 & \quad \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{7/2}}{7b}
 \end{aligned}$$

↓ 3106

$$\frac{10}{7}d^2 \left( 2d^2 \left( \frac{2}{3}c^2 \int \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} dx + \frac{2cd(c \sec(a+bx))^{3/2}}{3b\sqrt{d \csc(a+bx)}} \right) - \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right)$$

↓ 3042

$$\frac{10}{7}d^2 \left( 2d^2 \left( \frac{2}{3}c^2 \int \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} dx + \frac{2cd(c \sec(a+bx))^{3/2}}{3b\sqrt{d \csc(a+bx)}} \right) - \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right)$$

↓ 3110

$$\frac{10}{7}d^2 \left( 2d^2 \left( \frac{2}{3}c^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx + \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right) - \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right)$$

↓ 3042

$$\frac{10}{7}d^2 \left( 2d^2 \left( \frac{2}{3}c^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx + \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right) - \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right)$$

↓ 3053

$$\frac{10}{7}d^2 \left( 2d^2 \left( \frac{2}{3}c^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx + \frac{2cd(c \sec(a+bx))^{3/2}}{3b\sqrt{d \csc(a+bx)}} \right) - \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right)$$

↓ 3042

$$\frac{10}{7}d^2 \left( 2d^2 \left( \frac{2}{3}c^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx + \frac{2cd(c \sec(a+bx))^{3/2}}{3b\sqrt{d \csc(a+bx)}} \right) - \frac{2cd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{7/2}}{3b} \right)$$

↓ 3120

$$\frac{10}{7}d^2 \left( 2d^2 \left( \frac{2c^2 \sqrt{\sin(2a + 2bx)} \operatorname{EllipticF}\left(a + bx - \frac{\pi}{4}, 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{3b} + \frac{2cd(c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}} \right) \right. \\ \left. + \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{7/2}}{7b} \right)$$

input `Int[(d*Csc[a + b*x])^(9/2)*(c*Sec[a + b*x])^(5/2),x]`

output `(-2*c*d*(d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(3/2))/(7*b) + (10*d^2*((-2*c*d*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2))/(3*b) + 2*d^2*((2*c*d*(c*Sec[a + b*x])^(3/2))/(3*b*Sqrt[d*Csc[a + b*x]])) + (2*c^2*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(3*b)))/7`

### 3.246.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3106 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]`



rule 3110 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

### 3.246.4 Maple [A] (verified)

Time = 1.88 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.99

$$\frac{\sqrt{2} d^4 c^2 \sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} \left( (40 \cos(bx + a) + 40) \sqrt{1 + \csc(bx + a) - \cot(bx + a)} \sqrt{\cot(bx + a)} \right)}{\dots}$$

input `int((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2),x)`

output `1/21/b*2^(1/2)*d^4*c^2*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*((40*cos(b*x+a)+40)*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2)))+(20*cos(b*x+a)^4-30*cos(b*x+a)^2+7)*2^(1/2)*sec(b*x+a)*csc(b*x+a)^3)`

### 3.246.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.36

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx = \frac{2 \left( 10 (i c^2 d^4 \cos(bx + a)^3 - i c^2 d^4 \cos(bx + a)) \sqrt{-4i cd} F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) \sin(\dots) \right)}{\dots}$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fracas")`

---

3.246.  $\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx$

output 
$$\frac{-2/21*(10*(I*c^2*d^4*\cos(b*x + a)^3 - I*c^2*d^4*\cos(b*x + a))*\sqrt{-4*I*c*d}*\text{elliptic\_f}(\arcsin(\cos(b*x + a) + I*\sin(b*x + a)), -1)*\sin(b*x + a) + 10*(-I*c^2*d^4*\cos(b*x + a)^3 + I*c^2*d^4*\cos(b*x + a))*\sqrt{4*I*c*d}*\text{elliptic\_f}(\arcsin(\cos(b*x + a) - I*\sin(b*x + a)), -1)*\sin(b*x + a) + (20*c^2*d^4*\cos(b*x + a)^4 - 30*c^2*d^4*\cos(b*x + a)^2 + 7*c^2*d^4)*\sqrt{c/\cos(b*x + a))*\sqrt{d/\sin(b*x + a)}}{(b*\cos(b*x + a)^3 - b*\cos(b*x + a))*\sin(b*x + a)}$$

### 3.246.6 Sympy [F(-1)]

Timed out.

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(9/2)*(c*sec(b*x+a))**(5/2), x)`

output Timed out

### 3.246.7 Maxima [F]

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{9/2} (c \sec(bx + a))^{5/2} dx$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2), x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(5/2), x)`

### 3.246.8 Giac [F]

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{9/2} (c \sec(bx + a))^{5/2} dx$$

input `integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2), x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(5/2), x)`

**3.246.9 Mupad [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{5/2} \left( \frac{d}{\sin(a + bx)} \right)^{9/2} dx$$

input `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(9/2),x)`output `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(9/2), x)`

### 3.247 $\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx$

3.247.1 Optimal result . . . . .	1423
3.247.2 Mathematica [A] (verified) . . . . .	1423
3.247.3 Rubi [A] (verified) . . . . .	1424
3.247.4 Maple [A] (verified) . . . . .	1425
3.247.5 Fricas [A] (verification not implemented) . . . . .	1426
3.247.6 Sympy [F(-1)] . . . . .	1426
3.247.7 Maxima [F] . . . . .	1427
3.247.8 Giac [F] . . . . .	1427
3.247.9 Mupad [B] (verification not implemented) . . . . .	1427

#### 3.247.1 Optimal result

Integrand size = 25, antiderivative size = 106

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx = -\frac{64c^3 d^3 \sqrt{d \csc(a + bx)}}{15b \sqrt{c \sec(a + bx)}} + \frac{16cd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{15b} - \frac{2cd (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}}{5b}$$

output `-2/5*c*d*(d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2)/b+16/15*c*d^3*(c*sec(b*x+a))^(3/2)*(d*csc(b*x+a))^(1/2)/b-64/15*c^3*d^3*(d*csc(b*x+a))^(1/2)/b/(c*sec(b*x+a))^(1/2)`

#### 3.247.2 Mathematica [A] (verified)

Time = 0.84 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.54

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx = \frac{2cd^3(-5 + 32 \cos^2(a + bx) + 3 \cot^2(a + bx)) \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{15b}$$

input `Integrate[(d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(5/2),x]`

output `(-2*c*d^3*(-5 + 32*Cos[a + b*x]^2 + 3*Cot[a + b*x]^2)*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2))/(15*b)`

**3.247.3 Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {3042, 3105, 3042, 3106, 3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{7/2} dx \\
 & \quad \downarrow \text{3105} \\
 & \frac{8}{5} d^2 \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{5} d^2 \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3106} \\
 & \frac{8}{5} d^2 \left( \frac{4}{3} c^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} \right) - \\
 & \quad \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{5} d^2 \left( \frac{4}{3} c^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} \right) - \\
 & \quad \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{5/2}}{5b} \\
 & \quad \downarrow \text{3099} \\
 & \frac{8}{5} d^2 \left( \frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} - \frac{8c^3 d \sqrt{d \csc(a + bx)}}{3b \sqrt{c \sec(a + bx)}} \right) - \\
 & \quad \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{5/2}}{5b}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(5/2),x]`

output 
$$\frac{-2cd(d\csc[a+bx])^{5/2}(c\sec[a+bx])^{3/2}}{5b} + \frac{(8d^2((-8c^3d\sqrt{d\csc[a+bx]}))/(3b\sqrt{c\sec[a+bx]})) + (2cd\sqrt{d\csc[a+bx]})(c\sec[a+bx])^{3/2}}{3b}}{5}$$

### 3.247.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_) + (f_)*(x_)]*(a_.))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

rule 3105 `Int[(csc[(e_) + (f_)*(x_)]*(a_.))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3106 `Int[(csc[(e_) + (f_)*(x_)]*(a_.))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]`

### 3.247.4 Maple [A] (verified)

Time = 1.16 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.64

$$\frac{2d^3c^2\sqrt{c\sec(bx+a)}\sqrt{d\csc(bx+a)}(32\cos(bx+a)^4 - 40\cos(bx+a)^2 + 5)\sec(bx+a)\csc(bx+a)^2}{15b}$$

input `int((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2),x)`

output  $2/15/b*d^3*c^2*(c*\sec(b*x+a))^{(1/2)}*(d*\csc(b*x+a))^{(1/2)}*(32*\cos(b*x+a)^4-40*\cos(b*x+a)^2+5)*\sec(b*x+a)*\csc(b*x+a)^2$

### 3.247.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.84

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx = \frac{2(32c^2d^3\cos(bx+a)^4 - 40c^2d^3\cos(bx+a)^2 + 5c^2d^3)\sqrt{\frac{c}{\cos(bx+a)}}\sqrt{\frac{d}{\sin(bx+a)}}}{15(b\cos(bx+a)^3 - b\cos(bx+a))}$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

output `-2/15*(32*c^2*d^3*cos(b*x + a)^4 - 40*c^2*d^3*cos(b*x + a)^2 + 5*c^2*d^3)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*cos(b*x + a)^3 - b*cos(b*x + a))`

### 3.247.6 Sympy [F(-1)]

Timed out.

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(7/2)*(c*sec(b*x+a))**(5/2),x)`

output `Timed out`

**3.247.7 Maxima [F]**

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{\frac{7}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2), x)`

**3.247.8 Giac [F]**

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{\frac{7}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2), x)`

**3.247.9 Mupad [B] (verification not implemented)**

Time = 15.30 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.06

$$\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx = \frac{16 c^2 d^3 \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (5 \cos(a + bx) - 3 \cos(3a + 3bx) - 4 \cos(5a + 5bx) + 2 \cos(7a + 7bx))}{15 b (\cos(2a + 2bx) + 2 \cos(4a + 4bx) - \cos(6a + 6bx) - 2)}$$

input `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(7/2),x)`

output `(16*c^2*d^3*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(5*cos(a + b*x) - 3*cos(3*a + 3*b*x) - 4*cos(5*a + 5*b*x) + 2*cos(7*a + 7*b*x)))/(15*b*(cos(2*a + 2*b*x) + 2*cos(4*a + 4*b*x) - cos(6*a + 6*b*x) - 2))`



### 3.248 $\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx$

3.248.1 Optimal result . . . . .	1428
3.248.2 Mathematica [C] (verified) . . . . .	1428
3.248.3 Rubi [A] (verified) . . . . .	1429
3.248.4 Maple [A] (verified) . . . . .	1432
3.248.5 Fricas [C] (verification not implemented) . . . . .	1432
3.248.6 Sympy [F(-1)] . . . . .	1433
3.248.7 Maxima [F] . . . . .	1433
3.248.8 Giac [F] . . . . .	1433
3.248.9 Mupad [F(-1)] . . . . .	1434

#### 3.248.1 Optimal result

Integrand size = 25, antiderivative size = 131

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx = \frac{4cd^3 (c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}} - \frac{2cd (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{3b} + \frac{4c^2 d^2 \sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{3b}$$

output  $-2/3*c*d*(d*\csc(b*x+a))^{(3/2)}*(c*\sec(b*x+a))^{(3/2)}/b+4/3*c*d^3*(c*\sec(b*x+a))^{(3/2)}/b/(d*\csc(b*x+a))^{(1/2)}-4/3*c^2*d^2*(\sin(a+1/4*\pi+b*x)^2)^{(1/2)}/\sin(a+1/4*\pi+b*x)*\operatorname{EllipticF}(\cos(a+1/4*\pi+b*x), 2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b$

#### 3.248.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.37 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.66

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx = \frac{2c^3 d (d \csc(a + bx))^{3/2} \left( -1 + \cot^2(a + bx) + 2(-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a + bx)\right) \right)}{3b \sqrt{c \sec(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(5/2),x]`

output `(-2*c^3*d*(d*Csc[a + b*x])^(3/2)*(-1 + Cot[a + b*x]^2 + 2*(-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])*Tan[a + b*x]^2)/(3*b*Sqrt[c*Sec[a + b*x]])`

### 3.248.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3105, 3042, 3106, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{5/2} dx \\
 & \quad \downarrow \text{3105} \\
 & 2d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & 2d^2 \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx - \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3106} \\
 & 2d^2 \left( \frac{2}{3} c^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}} \right) - \\
 & \quad \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & 2d^2 \left( \frac{2}{3} c^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}} \right) - \\
 & \quad \frac{2cd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}{3b}
 \end{aligned}$$

↓ 3110

$$2d^2 \left( \frac{2}{3} c^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx + \frac{2cd}{3b} \right) - \frac{2cd(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}}{3b}$$

↓ 3042

$$2d^2 \left( \frac{2}{3} c^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx + \frac{2cd}{3b} \right) - \frac{2cd(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}}{3b}$$

↓ 3053

$$2d^2 \left( \frac{2}{3} c^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx + \frac{2cd(c \sec(a+bx))^{3/2}}{3b \sqrt{d \csc(a+bx)}} \right) - \frac{2cd(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}}{3b}$$

↓ 3042

$$2d^2 \left( \frac{2}{3} c^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx + \frac{2cd(c \sec(a+bx))^{3/2}}{3b \sqrt{d \csc(a+bx)}} \right) - \frac{2cd(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}}{3b}$$

↓ 3120

$$2d^2 \left( \frac{2c^2 \sqrt{\sin(2a+2bx)} \operatorname{EllipticF}\left(a+bx - \frac{\pi}{4}, 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3b} + \frac{2cd(c \sec(a+bx))^{3/2}}{3b \sqrt{d \csc(a+bx)}} \right) - \frac{2cd(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}}{3b}$$

input `Int[(d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(5/2),x]`

output `(-2*c*d*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2))/(3*b) + 2*d^2*((2*c*d*(c*Sec[a + b*x])^(3/2))/(3*b*sqrt[d*Csc[a + b*x]]) + (2*c^2*sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*sqrt[c*Sec[a + b*x]]*sqrt[Sin[2*a + 2*b*x]])/(3*b))`

## 3.248.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3106 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

**3.248.4 Maple [A] (verified)**

Time = 1.12 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.15

$$\sqrt{2} d^2 c^2 \sqrt{c \sec (bx+a)} \sqrt{d \csc (bx+a)} \left( (4 \cos (bx+a)+4) \sqrt{1+\csc (bx+a)-\cot (bx+a)} \sqrt{\cot (bx+a)} \right)$$

input `int((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x)`

output `1/3/b*2^(1/2)*d^2*c^2*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*((4*cos(b*x+a)+4)*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*(-2*cot(b*x+a)+csc(b*x+a)*sec(b*x+a)))`

**3.248.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.22

$$\int (d \csc (a+bx))^{5/2} (c \sec (a+bx))^{5/2} dx =$$

$$2 \left( i \sqrt{-4i c d c^2 d^2} \cos (bx+a) F(\arcsin (\cos (bx+a)+i \sin (bx+a))|-1) \sin (bx+a) - i \sqrt{4i c d c^2 d^2} \cos (bx+a) \right)$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

output `-2/3*(I*sqrt(-4*I*c*d)*c^2*d^2*cos(b*x+a)*elliptic_f(arcsin(cos(b*x+a)+I*sin(b*x+a)), -1)*sin(b*x+a) - I*sqrt(4*I*c*d)*c^2*d^2*cos(b*x+a)*elliptic_f(arcsin(cos(b*x+a)-I*sin(b*x+a)), -1)*sin(b*x+a) + (2*c^2*d^2*cos(b*x+a)^2 - c^2*d^2)*sqrt(c/cos(b*x+a))*sqrt(d/sin(b*x+a)))/(b*cos(b*x+a)*sin(b*x+a))`

**3.248.6 Sympy [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(5/2)*(c*sec(b*x+a))**(5/2),x)`output `Timed out`**3.248.7 Maxima [F]**

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2), x)`**3.248.8 Giac [F]**

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x, algorithm="giac")`output `integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2), x)`

**3.248.9 Mupad [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{5/2} \left( \frac{d}{\sin(a + bx)} \right)^{5/2} dx$$

input `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2),x)`output `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2), x)`

### 3.249 $\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx$

3.249.1 Optimal result . . . . .	1435
3.249.2 Mathematica [A] (verified) . . . . .	1435
3.249.3 Rubi [A] (verified) . . . . .	1436
3.249.4 Maple [A] (verified) . . . . .	1437
3.249.5 Fracas [A] (verification not implemented) . . . . .	1437
3.249.6 Sympy [F(-1)] . . . . .	1438
3.249.7 Maxima [F] . . . . .	1438
3.249.8 Giac [F] . . . . .	1438
3.249.9 Mupad [B] (verification not implemented) . . . . .	1439

#### 3.249.1 Optimal result

Integrand size = 25, antiderivative size = 69

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx = -\frac{8c^3 d \sqrt{d \csc(a + bx)}}{3b \sqrt{c \sec(a + bx)}} + \frac{2cd \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{3b}$$

output `2/3*c*d*(c*sec(b*x+a))^(3/2)*(d*csc(b*x+a))^(1/2)/b-8/3*c^3*d*(d*csc(b*x+a))^(1/2)/b/(c*sec(b*x+a))^(1/2)`

#### 3.249.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx = -\frac{2cd(1 + 2 \cos(2(a + bx))) \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{3b}$$

input `Integrate[(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2),x]`

output `(-2*c*d*(1 + 2*Cos[2*(a + b*x)])*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2))/(3*b)`



**3.249.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {3042, 3106, 3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(a + bx))^{5/2} (d \csc(a + bx))^{3/2} dx \\
 & \quad \downarrow \text{3106} \\
 & \frac{4}{3} c^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4}{3} c^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} \\
 & \quad \downarrow \text{3099} \\
 & \frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} - \frac{8c^3 d \sqrt{d \csc(a + bx)}}{3b \sqrt{c \sec(a + bx)}}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2),x]`

output `(-8*c^3*d*Sqrt[d*Csc[a + b*x]])/(3*b*Sqrt[c*Sec[a + b*x]]) + (2*c*d*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2))/(3*b)`

**3.249.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3099 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1
))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] &&
NeQ[n, 1]
```

```
rule 3106 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(
n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n -
1))/(f*(n - 1)), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])
^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n,
1] && IntegersQ[2*m, 2*n]
```

### 3.249.4 Maple [A] (verified)

Time = 1.05 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.68

method	result	size
default	$-\frac{2dc^2\sqrt{d\csc(bx+a)}\sqrt{c\sec(bx+a)}(4\cos(bx+a)-\sec(bx+a))}{3b}$	47

```
input int((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2/3/b*d*c^2*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*(4*cos(b*x+a)-sec(b
*x+a))
```

### 3.249.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.84

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx = -\frac{2(4c^2d \cos(bx + a)^2 - c^2d) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{3b \cos(bx + a)}$$

```
input integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fracas")
```

```
output -2/3*(4*c^2*d*cos(b*x + a)^2 - c^2*d)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x
+ a))/(b*cos(b*x + a))
```

**3.249.6 Sympy [F(-1)]**

Timed out.

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(3/2)*(c*sec(b*x+a))**(5/2),x)`output `Timed out`**3.249.7 Maxima [F]**

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2), x)`**3.249.8 Giac [F]**

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx = \int (d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x, algorithm="giac")`output `integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2), x)`

**3.249.9 Mupad [B] (verification not implemented)**

Time = 13.52 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.93

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx =$$

$$\frac{4c^2 d \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (2 \cos(a + bx) + \cos(3a + 3bx))}{3b (\cos(2a + 2bx) + 1)}$$

input `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(3/2),x)`output `-(4*c^2*d*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(2*cos(a + b*x) + cos(3*a + 3*b*x)))/(3*b*(cos(2*a + 2*b*x) + 1))`

### 3.250 $\int \sqrt{d \csc(a + bx)}(c \sec(a + bx))^{5/2} dx$

3.250.1 Optimal result . . . . .	1440
3.250.2 Mathematica [C] (verified) . . . . .	1440
3.250.3 Rubi [A] (verified) . . . . .	1441
3.250.4 Maple [A] (verified) . . . . .	1443
3.250.5 Fricas [C] (verification not implemented) . . . . .	1443
3.250.6 Sympy [F(-1)] . . . . .	1444
3.250.7 Maxima [F] . . . . .	1444
3.250.8 Giac [F] . . . . .	1444
3.250.9 Mupad [F(-1)] . . . . .	1445

#### 3.250.1 Optimal result

Integrand size = 25, antiderivative size = 93

$$\int \sqrt{d \csc(a + bx)}(c \sec(a + bx))^{5/2} dx = \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} + \frac{2c^2 \sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{3b}$$

output

```
2/3*c*d*(c*sec(b*x+a))^(3/2)/b/(d*csc(b*x+a))^(1/2)-2/3*c^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b
```

#### 3.250.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.16 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.73

$$\int \sqrt{d \csc(a + bx)}(c \sec(a + bx))^{5/2} dx = \frac{2cd \left( -1 + (-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a + bx)\right) \right) (c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}}$$

input

```
Integrate[Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(5/2),x]
```

output  $(-2*c*d*(-1 + (-\text{Cot}[a + b*x]^2)^{3/4}*\text{Hypergeometric2F1}[1/2, 3/4, 3/2, \text{Csc}[a + b*x]^2])*(c*\text{Sec}[a + b*x])^{3/2})/(3*b*\text{Sqrt}[d*\text{Csc}[a + b*x]])$

### 3.250.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3106, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)} dx$$

$$\downarrow \text{3042}$$

$$\int (c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)} dx$$

$$\downarrow \text{3106}$$

$$\frac{2}{3}c^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}}$$

$$\downarrow \text{3042}$$

$$\frac{2}{3}c^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx + \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}}$$

$$\downarrow \text{3110}$$

$$\frac{2}{3}c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx + \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}}$$

$$\downarrow \text{3042}$$

$$\frac{2}{3}c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx + \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}}$$

$$\downarrow \text{3053}$$

$$\frac{2}{3}c^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx+\frac{2cd(c\sec(a+bx))^{3/2}}{3b\sqrt{d\csc(a+bx)}}$$

↓ 3042

$$\frac{2}{3}c^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx+\frac{2cd(c\sec(a+bx))^{3/2}}{3b\sqrt{d\csc(a+bx)}}$$

↓ 3120

$$\frac{2c^2\sqrt{\sin(2a+2bx)}\operatorname{EllipticF}\left(a+bx-\frac{\pi}{4},2\right)\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{3b}+\frac{2cd(c\sec(a+bx))^{3/2}}{3b\sqrt{d\csc(a+bx)}}$$

input `Int[Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(5/2),x]`

output `(2*c*d*(c*Sec[a + b*x])^(3/2))/(3*b*Sqrt[d*Csc[a + b*x]]) + (2*c^2*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(3*b)`

### 3.250.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3106 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Simp[b^2*((m + n - 2)/(n - 1)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]`

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.250.4 Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.23

method	result
default	$\frac{\sqrt{2} \sqrt{c \sec(bx+a)} \sqrt{d \csc(bx+a)} c^2 (2\sqrt{1+\csc(bx+a)-\cot(bx+a)} \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)}) \text{EllipticF}(\sqrt{\dots})}{\dots}$

```
input int((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/3/b*2^(1/2)*(c*sec(b*x+a))^(1/2)*(d*csc(b*x+a))^(1/2)*c^2*(2*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a)))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)+2*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*tan(b*x+a))
```

### 3.250.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.28

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx = \frac{-i \sqrt{-4i c d c^2} \cos(bx + a) F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) + i \sqrt{4i c d c^2} \cos(bx + a)}{3 b \cos(bx + a)}$$

```
input integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fricas")
```

---

3.250.  $\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx$



output  $1/3*(-I*\sqrt{-4*I*c*d})*c^2*\cos(b*x + a)*\text{elliptic\_f}(\arcsin(\cos(b*x + a) + I*\sin(b*x + a)), -1) + I*\sqrt{4*I*c*d})*c^2*\cos(b*x + a)*\text{elliptic\_f}(\arcsin(\cos(b*x + a) - I*\sin(b*x + a)), -1) + 2*c^2*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)}*\sin(b*x + a)/(b*\cos(b*x + a))$

### 3.250.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(1/2)*(c*sec(b*x+a))**(5/2), x)`

output Timed out

### 3.250.7 Maxima [F]

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx = \int \sqrt{d \csc(bx + a)} (c \sec(bx + a))^{5/2} dx$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2), x, algorithm="maxima")`

output `integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2), x)`

### 3.250.8 Giac [F]

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx = \int \sqrt{d \csc(bx + a)} (c \sec(bx + a))^{5/2} dx$$

input `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2), x, algorithm="giac")`

output `integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2), x)`

**3.250.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx = \int \left( \frac{c}{\cos(a + bx)} \right)^{5/2} \sqrt{\frac{d}{\sin(a + bx)}} dx$$

input `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2),x)`output `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2), x)`

**3.251**  $\int \frac{(c \sec(a+bx))^{5/2}}{\sqrt{d \csc(a+bx)}} dx$

3.251.1 Optimal result . . . . . 1446  
 3.251.2 Mathematica [A] (verified) . . . . . 1446  
 3.251.3 Rubi [A] (verified) . . . . . 1447  
 3.251.4 Maple [A] (verified) . . . . . 1448  
 3.251.5 Fricas [B] (verification not implemented) . . . . . 1448  
 3.251.6 Sympy [F(-1)] . . . . . 1448  
 3.251.7 Maxima [F] . . . . . 1449  
 3.251.8 Giac [F] . . . . . 1449  
 3.251.9 Mupad [B] (verification not implemented) . . . . . 1449

**3.251.1 Optimal result**

Integrand size = 25, antiderivative size = 33

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx = \frac{2cd(c \sec(a + bx))^{3/2}}{3b(d \csc(a + bx))^{3/2}}$$

output  $2/3*c*d*(c*\sec(b*x+a))^(3/2)/b/(d*\csc(b*x+a))^(3/2)$

**3.251.2 Mathematica [A] (verified)**

Time = 0.60 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx = \frac{2cd(c \sec(a + bx))^{3/2}}{3b(d \csc(a + bx))^{3/2}}$$

input `Integrate[(c*Sec[a + b*x])^(5/2)/Sqrt[d*Csc[a + b*x]],x]`

output  $(2*c*d*(c*\sec[a + b*x])^(3/2))/(3*b*(d*\csc[a + b*x])^(3/2))$

**3.251.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx$$

↓ 3042

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx$$

↓ 3099

$$\frac{2cd(c \sec(a + bx))^{3/2}}{3b(d \csc(a + bx))^{3/2}}$$

input `Int[(c*Sec[a + b*x])^(5/2)/Sqrt[d*Csc[a + b*x]],x]`

output `(2*c*d*(c*Sec[a + b*x])^(3/2))/(3*b*(d*Csc[a + b*x])^(3/2))`

**3.251.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

**3.251.4 Maple [A] (verified)**

Time = 1.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

method	result	size
default	$\frac{2\sqrt{c \sec(bx+a)} c^2 \tan(bx+a)}{3b\sqrt{d \csc(bx+a)}}$	35

input `int((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2/3/b*(c*sec(b*x+a))^(1/2)*c^2/(d*csc(b*x+a))^(1/2)*tan(b*x+a)`

**3.251.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 58 vs.  $2(27) = 54$ .

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.76

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx = -\frac{2(c^2 \cos(bx + a)^2 - c^2) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{3bd \cos(bx + a)}$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x, algorithm="fracas")`

output `-2/3*(c^2*cos(b*x + a)^2 - c^2)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*d*cos(b*x + a))`

**3.251.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx = \text{Timed out}$$

input `integrate((c*sec(b*x+a))**(5/2)/(d*csc(b*x+a))**(1/2),x)`

output `Timed out`

**3.251.7 Maxima [F]**

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{(c \sec(bx + a))^{5/2}}{\sqrt{d \csc(bx + a)}} dx$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(5/2)/sqrt(d*csc(b*x + a)), x)`

**3.251.8 Giac [F]**

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx = \int \frac{(c \sec(bx + a))^{5/2}}{\sqrt{d \csc(bx + a)}} dx$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(5/2)/sqrt(d*csc(b*x + a)), x)`

**3.251.9 Mupad [B] (verification not implemented)**

Time = 13.58 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.00

$$\int \frac{(c \sec(a + bx))^{5/2}}{\sqrt{d \csc(a + bx)}} dx = \frac{c^2 \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (\cos(a + bx) - \cos(3a + 3bx))}{3bd (\cos(2a + 2bx) + 1)}$$

input `int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(1/2),x)`

output `(c^2*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(cos(a + b*x) - cos(3*a + 3*b*x)))/(3*b*d*(cos(2*a + 2*b*x) + 1))`

**3.252**  $\int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{3/2}} dx$

3.252.1 Optimal result . . . . . 1450  
 3.252.2 Mathematica [C] (verified) . . . . . 1450  
 3.252.3 Rubi [A] (verified) . . . . . 1451  
 3.252.4 Maple [B] (verified) . . . . . 1453  
 3.252.5 Fricas [C] (verification not implemented) . . . . . 1453  
 3.252.6 Sympy [F(-1)] . . . . . 1454  
 3.252.7 Maxima [F] . . . . . 1454  
 3.252.8 Giac [F] . . . . . 1454  
 3.252.9 Mupad [F(-1)] . . . . . 1455

**3.252.1 Optimal result**

Integrand size = 25, antiderivative size = 98

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx = \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d \csc(a + bx)}} - \frac{c^2 \sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{3bd^2}$$

output  $2/3*c*(c*\sec(b*x+a))^(3/2)/b/d/(d*\csc(b*x+a))^(1/2)+1/3*c^2*(\sin(a+1/4*Pi+b*x)^2)^(1/2)/\sin(a+1/4*Pi+b*x)*\operatorname{EllipticF}(\cos(a+1/4*Pi+b*x), 2^(1/2))*(d*cs c(b*x+a))^(1/2)*(c*\sec(b*x+a))^(1/2)*\sin(2*b*x+2*a)^(1/2)/b/d^2$

**3.252.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.21 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.71

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx = \frac{c\left(2 + (-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a + bx)\right)\right)}{3bd\sqrt{d \csc(a + bx)}} (c \sec(a + bx))^{3/2}$$

input `Integrate[(c*Sec[a + b*x])^(5/2)/(d*Csc[a + b*x])^(3/2),x]`

output  $(c*(2 + (-\operatorname{Cot}[a + b*x]^2)^(3/4)*\operatorname{Hypergeometric2F1}[1/2, 3/4, 3/2, \operatorname{Csc}[a + b*x]^2]))*(c*\operatorname{Sec}[a + b*x])^(3/2)/(3*b*d*\operatorname{Sqrt}[d*\operatorname{Csc}[a + b*x]])$

3.252.  $\int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{3/2}} dx$

**3.252.3 Rubi [A] (verified)**

Time = 0.52 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3104, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3104} \\
 & \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d \csc(a + bx)}} - \frac{c^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{3d^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d \csc(a + bx)}} - \frac{c^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{3d^2} \\
 & \quad \downarrow \text{3110} \\
 & \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d \csc(a + bx)}} - \frac{c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx}{3d^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d \csc(a + bx)}} - \frac{c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx}{3d^2} \\
 & \quad \downarrow \text{3053} \\
 & \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d \csc(a + bx)}} - \frac{c^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{\sin(2a + 2bx)}} dx}{3d^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.252.  $\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx$



$$\frac{2c(c\sec(a+bx))^{3/2}}{3bd\sqrt{d\csc(a+bx)}} - \frac{c^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{3d^2} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx$$

↓ 3120

$$\frac{2c(c\sec(a+bx))^{3/2}}{3bd\sqrt{d\csc(a+bx)}} - \frac{c^2\sqrt{\sin(2a+2bx)}\operatorname{EllipticF}\left(a+bx-\frac{\pi}{4}, 2\right)\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{3bd^2}$$

input `Int[(c*Sec[a + b*x])^(5/2)/(d*Csc[a + b*x])^(3/2), x]`

output `(2*c*(c*Sec[a + b*x])^(3/2))/(3*b*d*Sqrt[d*Csc[a + b*x]]) - (c^2*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(3*b*d^2)`

### 3.252.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), x] Int[1/Sqrt[Sin[2*e + 2*f*x]], x, x] /; FreeQ[{a, b, e, f}, x]`

rule 3104 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(f*a*(n - 1))), x] + Simp[b^2*((m + 1)/(a^2*(n - 1))) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[n, 1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

### 3.252.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 242 vs.  $2(109) = 218$ .

Time = 1.23 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.48

method	result
default	$\frac{\sqrt{2} \left( \sqrt{1+\csc(bx+a)} - \cot(bx+a) \right) \sqrt{\cot(bx+a) - \csc(bx+a)+1} \sqrt{\cot(bx+a) - \csc(bx+a)}}{\sqrt{2} \left( \sqrt{1+\csc(bx+a)} - \cot(bx+a) \right) \sqrt{\cot(bx+a) - \csc(bx+a)+1} \sqrt{\cot(bx+a) - \csc(bx+a)}} \text{EllipticF}\left(\sqrt{1+\csc(bx+a)} - \cot(bx+a), \frac{\sqrt{2}}{2}\right)$

input `int((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `1/3/b*2^(1/2)*((1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)^2+(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)-2^(1/2)*sin(b*x+a)*(c*sec(b*x+a))^(1/2)*c^2/d/(d*csc(b*x+a))^(1/2)/(cos(b*x+a)-1)/(cos(b*x+a)+1)*tan(b*x+a)`

### 3.252.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.24

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx = \frac{i \sqrt{-4i c d c^2} \cos(bx + a) F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) - i \sqrt{4i c d c^2}}{\dots}$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x, algorithm="fricas")`

output `1/6*(I*sqrt(-4*I*c*d)*c^2*cos(b*x + a)*elliptic_f(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) - I*sqrt(4*I*c*d)*c^2*cos(b*x + a)*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1) + 4*c^2*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*sin(b*x + a)/(b*d^2*cos(b*x + a))`

---

3.252.  $\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx$

**3.252.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate((c*sec(b*x+a))**(5/2)/(d*csc(b*x+a))**(3/2),x)`output `Timed out`**3.252.7 Maxima [F]**

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{(c \sec(bx + a))^{5/2}}{(d \csc(bx + a))^{3/2}} dx$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x, algorithm="maxima")`output `integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(3/2), x)`**3.252.8 Giac [F]**

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{(c \sec(bx + a))^{5/2}}{(d \csc(bx + a))^{3/2}} dx$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x, algorithm="giac")`output `integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(3/2), x)`

**3.252.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx = \int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}}{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

input `int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(3/2),x)`output `int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(3/2), x)`

**3.253** 
$$\int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{5/2}} dx$$

3.253.1 Optimal result . . . . . 1456  
 3.253.2 Mathematica [A] (verified) . . . . . 1457  
 3.253.3 Rubi [A] (verified) . . . . . 1457  
 3.253.4 Maple [A] (warning: unable to verify) . . . . . 1462  
 3.253.5 Fricas [C] (verification not implemented) . . . . . 1462  
 3.253.6 Sympy [F(-1)] . . . . . 1463  
 3.253.7 Maxima [F] . . . . . 1464  
 3.253.8 Giac [F] . . . . . 1464  
 3.253.9 Mupad [F(-1)] . . . . . 1464

**3.253.1 Optimal result**

Integrand size = 25, antiderivative size = 329

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx = \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} + \frac{c^2 \arctan\left(1 - \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2}bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} - \frac{c^2 \arctan\left(1 + \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2}bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} - \frac{c^2 \log\left(1 - \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c \sec(a + bx)}}{2\sqrt{2}bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} + \frac{c^2 \log\left(1 + \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c \sec(a + bx)}}{2\sqrt{2}bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}$$

output

```
2/3*c*(c*sec(b*x+a))^(3/2)/b/d/(d*csc(b*x+a))^(3/2)-1/2*c^2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)-1/2*c^2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)-1/4*c^2*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+1/4*c^2*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)
```

**3.253.2 Mathematica [A] (verified)**

Time = 2.40 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.43

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx = \frac{c \left( 4 + 3\sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}} \right) \cot^2(a + bx)^{3/4} + 3\sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}{1 + \sqrt{\cot^2(a + bx)}} \right) \cot^2(a + bx)^{3/4} \right)}{6bd(d \csc(a + bx))^{3/2}}$$

input `Integrate[(c*Sec[a + b*x])^(5/2)/(d*Csc[a + b*x])^(5/2), x]`output `(c*(4 + 3*Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2)]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(3/4) + 3*Sqrt[2]*ArcTanh[Sqrt[2]*(Cot[a + b*x]^2)^(1/4)]/(1 + Sqrt[Cot[a + b*x]^2]))*(Cot[a + b*x]^2)^(3/4)*(c*Sec[a + b*x])^(3/2))/(6*b*d*(d*Csc[a + b*x])^(3/2))`**3.253.3 Rubi [A] (verified)**Time = 0.58 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.65, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3104, 3042, 3109, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3104} \\ & \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{c^2 \int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx}{d^2} \\ & \quad \downarrow \text{3042} \\ & \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{c^2 \int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx}{d^2} \end{aligned}$$

---

3.253.  $\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx$

$$\begin{array}{c}
\downarrow \text{3109} \\
\frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{c^2 \sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
\downarrow \text{3042} \\
\frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{c^2 \sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
\downarrow \text{3957} \\
\frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{c^2 \sqrt{c \sec(a+bx)} \int \frac{\sqrt{\tan(a+bx)}}{\tan^2(a+bx)+1} d \tan(a+bx)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
\downarrow \text{266} \\
\frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{2c^2 \sqrt{c \sec(a+bx)} \int \frac{\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
\downarrow \text{826} \\
\frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
\downarrow \text{1476} \\
\frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \\
\downarrow \text{1082} \\
\frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{1}{-\tan(a+bx)-1} d \left( \frac{1-\sqrt{2}\sqrt{\tan(a+bx)}}{\sqrt{2}} \right) - \int \frac{1}{-\tan(a+bx)-1} d \left( \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\sqrt{2}} \right) \right) \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \\
\downarrow \text{217}
\end{array}$$

---

3.253.  $\int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{5/2}} dx$

$$\frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{2c(c \sec(a+bx))^{3/2}}{3bd(d \csc(a+bx))^{3/2}} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d\sqrt{\tan(a+bx)} \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

↓ 1479

$$\frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{-\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} \right) \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

↓ 25

$$\frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} \right) \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

↓ 27

$$\frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} \right) \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

↓ 1103

$$\frac{2c^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log\left(\frac{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{2\sqrt{2}} - \frac{\log\left(\frac{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}\right)}{2\sqrt{2}} \right) \right)}{bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

input `Int[(c*Sec[a + b*x])^(5/2)/(d*Csc[a + b*x])^(5/2), x]`



output  $(2*c*(c*\text{Sec}[a + b*x])^{3/2})/(3*b*d*(d*\text{Csc}[a + b*x])^{3/2}) - (2*c^2*((-\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]/\text{Sqrt}[2]) + \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]/\text{Sqrt}[2])/2 + (\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]/(2*\text{Sqrt}[2]) - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]/(2*\text{Sqrt}[2])))*\text{Sqrt}[c*\text{Sec}[a + b*x]]/(b*d^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])$

### 3.253.3.1 Defintions of rubi rules used

rule 25  $\text{Int}[-(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$

rule 27  $\text{Int}[(a_)*(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, x]]$

rule 217  $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 266  $\text{Int}[(c_)*(x_)^m*(a_ + (b_)*(x_)^2)^{p_}, x\_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \quad \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(2*k)/c^2})^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 826  $\text{Int}[(x_)^2/((a_ + (b_)*(x_)^4), x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Simp}[1/(2*s) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 1082  $\text{Int}[(a_ + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3104 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n_, x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(f*a*(n - 1))), x] + Simp[b^2*((m + 1)/(a^2*(n - 1))) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[n, 1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n_, x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^n_, x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.253.4 Maple [A] (warning: unable to verify)**

Time = 8.80 (sec) , antiderivative size = 487, normalized size of antiderivative = 1.48

method	result
default	$\frac{\sqrt{2} c^2 \sqrt{c \sec(bx+a)} \left( 6 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \arctan \left( \frac{\sin(bx+a) \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2} + \cos(bx+a) - 1}}{\cos(bx+a) - 1} \right) \sin(bx+a) - 6 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \right)}{\dots}$

```
input int((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/12/b*2^(1/2)*c^2*(c*sec(b*x+a))^(1/2)*(6*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*arctan((sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+cos(b*x+a)-1)/(cos(b*x+a)-1))*sin(b*x+a)-6*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*arctan((-sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+cos(b*x+a)-1)/(cos(b*x+a)-1))*sin(b*x+a)-3*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*ln(2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)+2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2)*sin(b*x+a)+3*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*ln(-2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)-2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2)*sin(b*x+a)+4*2^(1/2)*tan(b*x+a)-4*2^(1/2)*sin(b*x+a))/(cos(b*x+a)-1)/d^2/(d*csc(b*x+a))^(1/2)
```

**3.253.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.67 (sec) , antiderivative size = 1371, normalized size of antiderivative = 4.17

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx = \text{Too large to display}$$

```
input integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```

1/24*(3*b*d^3*(-c^10/(b^4*d^10))^(1/4)*cos(b*x + a)*log(2*b^2*c^3*d^5*sqrt
(-c^10/(b^4*d^10))*cos(b*x + a)*sin(b*x + a) + 2*c^8*cos(b*x + a)^2 - c^8
+ 2*(b*c^5*d^2*(-c^10/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3
*d^7*cos(b*x + a)^3 - b^3*d^7*cos(b*x + a))*(-c^10/(b^4*d^10))^(3/4))*sqrt
(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) - 3*b*d^3*(-c^10/(b^4*d^10))^(1/4)*
cos(b*x + a)*log(2*b^2*c^3*d^5*sqrt(-c^10/(b^4*d^10))*cos(b*x + a)*sin(b*x
+ a) + 2*c^8*cos(b*x + a)^2 - c^8 - 2*(b*c^5*d^2*(-c^10/(b^4*d^10))^(1/4)
*cos(b*x + a)^2*sin(b*x + a) - (b^3*d^7*cos(b*x + a)^3 - b^3*d^7*cos(b*x +
a))*(-c^10/(b^4*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)))
- 3*I*b*d^3*(-c^10/(b^4*d^10))^(1/4)*cos(b*x + a)*log(-2*b^2*c^3*d^5*sqrt(
-c^10/(b^4*d^10))*cos(b*x + a)*sin(b*x + a) + 2*c^8*cos(b*x + a)^2 - c^8 -
2*(I*b*c^5*d^2*(-c^10/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (I*
b^3*d^7*cos(b*x + a)^3 - I*b^3*d^7*cos(b*x + a))*(-c^10/(b^4*d^10))^(3/4))
*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) + 3*I*b*d^3*(-c^10/(b^4*d^10))
^(1/4)*cos(b*x + a)*log(-2*b^2*c^3*d^5*sqrt(-c^10/(b^4*d^10))*cos(b*x + a)
*sin(b*x + a) + 2*c^8*cos(b*x + a)^2 - c^8 - 2*(-I*b*c^5*d^2*(-c^10/(b^4*d
^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (-I*b^3*d^7*cos(b*x + a)^3 + I*b
^3*d^7*cos(b*x + a))*(-c^10/(b^4*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d
/sin(b*x + a))) - 3*b*d^3*(-c^10/(b^4*d^10))^(1/4)*cos(b*x + a)*log(-c^8 +
2*(b*c^5*d^2*(-c^10/(b^4*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b...

```

### 3.253.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((c*sec(b*x+a))**(5/2)/(d*csc(b*x+a))**(5/2),x)`

output `Timed out`

**3.253.7 Maxima [F]**

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{(c \sec(bx + a))^{5/2}}{(d \csc(bx + a))^{5/2}} dx$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(5/2), x)`

**3.253.8 Giac [F]**

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{(c \sec(bx + a))^{5/2}}{(d \csc(bx + a))^{5/2}} dx$$

input `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(5/2), x)`

**3.253.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx = \int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}}{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

input `int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(5/2),x)`

output `int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(5/2), x)`

**3.254**  $\int \frac{(d \csc(a+bx))^{9/2}}{\sqrt{c \sec(a+bx)}} dx$

3.254.1 Optimal result . . . . . 1465  
 3.254.2 Mathematica [A] (verified) . . . . . 1465  
 3.254.3 Rubi [A] (verified) . . . . . 1466  
 3.254.4 Maple [A] (verified) . . . . . 1467  
 3.254.5 Fricas [A] (verification not implemented) . . . . . 1468  
 3.254.6 Sympy [F(-1)] . . . . . 1468  
 3.254.7 Maxima [F] . . . . . 1468  
 3.254.8 Giac [F] . . . . . 1469  
 3.254.9 Mupad [B] (verification not implemented) . . . . . 1469

**3.254.1 Optimal result**

Integrand size = 25, antiderivative size = 69

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{8cd^3(d \csc(a + bx))^{3/2}}{21b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b(c \sec(a + bx))^{3/2}}$$

output `-8/21*c*d^3*(d*csc(b*x+a))^(3/2)/b/(c*sec(b*x+a))^(3/2)-2/7*c*d*(d*csc(b*x+a))^(7/2)/b/(c*sec(b*x+a))^(3/2)`

**3.254.2 Mathematica [A] (verified)**

Time = 1.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = \frac{2cd(-5 + 2 \cos(2(a + bx)))(d \csc(a + bx))^{7/2}}{21b(c \sec(a + bx))^{3/2}}$$

input `Integrate[(d*Csc[a + b*x])^(9/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(2*c*d*(-5 + 2*Cos[2*(a + b*x)])*(d*Csc[a + b*x])^(7/2))/(21*b*(c*Sec[a + b*x])^(3/2))`

**3.254.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {3042, 3105, 3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx \\
 & \quad \downarrow \text{3105} \\
 & \frac{4}{7} d^2 \int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx - \frac{2cd(d \csc(a + bx))^{7/2}}{7b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4}{7} d^2 \int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx - \frac{2cd(d \csc(a + bx))^{7/2}}{7b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3099} \\
 & -\frac{8cd^3(d \csc(a + bx))^{3/2}}{21b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b(c \sec(a + bx))^{3/2}}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(9/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(-8*c*d^3*(d*Csc[a + b*x])^(3/2))/(21*b*(c*Sec[a + b*x])^(3/2)) - (2*c*d*(d*Csc[a + b*x])^(7/2))/(7*b*(c*Sec[a + b*x])^(3/2))`

## 3.254.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

## 3.254.4 Maple [A] (verified)

Time = 1.46 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{2d^4 \sqrt{d \csc(bx+a)} (4 \cot(bx+a)^3 - 7 \cot(bx+a) \csc(bx+a)^2)}{21b \sqrt{c \sec(bx+a)}}$	56

input `int((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2/21/b*d^4*(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)*(4*cot(b*x+a)^3-7*cot(b*x+a)*csc(b*x+a)^2)`



**3.254.5 Fricas [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.14

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{2(4d^4 \cos(bx + a)^4 - 7d^4 \cos(bx + a)^2) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{21(bc \cos(bx + a)^2 - bc) \sin(bx + a)}$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`output `-2/21*(4*d^4*cos(b*x + a)^4 - 7*d^4*cos(b*x + a)^2)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/((b*c*cos(b*x + a)^2 - b*c)*sin(b*x + a))`**3.254.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(9/2)/(c*sec(b*x+a))**(1/2),x)`output `Timed out`**3.254.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{9/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(9/2)/sqrt(c*sec(b*x + a)), x)`

**3.254.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{9/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(9/2)/sqrt(c*sec(b*x + a)), x)`

**3.254.9 Mupad [B] (verification not implemented)**

Time = 14.53 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.43

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = \frac{8 d^4 \sqrt{\frac{d}{\sin(a+bx)}} (11 \sin(2a + 2bx) - 7 \sin(4a + 4bx) + \sin(6a + 6bx))}{21 b \sqrt{\frac{c}{\cos(a+bx)}} (15 \cos(2a + 2bx) - 6 \cos(4a + 4bx) + \cos(6a + 6bx) - 10)}$$

input `int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(1/2),x)`

output `(8*d^4*(d/sin(a + b*x))^(1/2)*(11*sin(2*a + 2*b*x) - 7*sin(4*a + 4*b*x) + sin(6*a + 6*b*x)))/(21*b*(c/cos(a + b*x))^(1/2)*(15*cos(2*a + 2*b*x) - 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) - 10))`

**3.255**  $\int \frac{(d \csc(a+bx))^{7/2}}{\sqrt{c \sec(a+bx)}} dx$

3.255.1 Optimal result . . . . . 1470  
 3.255.2 Mathematica [C] (verified) . . . . . 1470  
 3.255.3 Rubi [A] (verified) . . . . . 1471  
 3.255.4 Maple [B] (verified) . . . . . 1473  
 3.255.5 Fricas [C] (verification not implemented) . . . . . 1474  
 3.255.6 Sympy [F(-1)] . . . . . 1474  
 3.255.7 Maxima [F] . . . . . 1475  
 3.255.8 Giac [F] . . . . . 1475  
 3.255.9 Mupad [F(-1)] . . . . . 1475

**3.255.1 Optimal result**

Integrand size = 25, antiderivative size = 128

$$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{4cd^3 \sqrt{d \csc(a + bx)}}{5b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} - \frac{4d^4 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{5b \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}$$

output

```
-2/5*c*d*(d*csc(b*x+a))^(5/2)/b/(c*sec(b*x+a))^(3/2)-4/5*c*d^3*(d*csc(b*x+a))^(1/2)/b/(c*sec(b*x+a))^(3/2)+4/5*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

**3.255.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.77 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.81

$$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx = \frac{2d^2(d \csc(a + bx))^{3/2} \left( -((-2 + \cos(2(a + bx))) \cot^3(a + bx)) + \sqrt{-\cot^2(a + bx)} \operatorname{Hypergeometric2F1}(-\dots) \right)}{5b \sqrt{c \sec(a + bx)}}$$

3.255.  $\int \frac{(d \csc(a+bx))^{7/2}}{\sqrt{c \sec(a+bx)}} dx$

input `Integrate[(d*Csc[a + b*x])^(7/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(-2*d^2*(d*Csc[a + b*x])^(3/2)*((-2 + Cos[2*(a + b*x)])*Cot[a + b*x]^3) + (-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2]*Sin[2*(a + b*x)]*Tan[a + b*x]^2)/(5*b*Sqrt[c*Sec[a + b*x]])`

### 3.255.3 Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3105, 3042, 3105, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx \\
 & \quad \downarrow \text{3105} \\
 & \frac{2}{5} d^2 \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{5} d^2 \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3105} \\
 & \frac{2}{5} d^2 \left( -2d^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx - \frac{2cd\sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \right) - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{5} d^2 \left( -2d^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx - \frac{2cd\sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \right) - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3110}
 \end{aligned}$$

$$\frac{2}{5}d^2 \left( -\frac{2d^2 \int \sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)} dx}{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right) - \frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{3/2}}$$

↓ 3042

$$\frac{2}{5}d^2 \left( -\frac{2d^2 \int \sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)} dx}{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right) - \frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{3/2}}$$

↓ 3052

$$\frac{2}{5}d^2 \left( -\frac{2d^2 \int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right) - \frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{3/2}}$$

↓ 3042

$$\frac{2}{5}d^2 \left( -\frac{2d^2 \int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right) - \frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{3/2}}$$

↓ 3119

$$\frac{2}{5}d^2 \left( -\frac{2d^2 E\left(a+bx-\frac{\pi}{4} \mid 2\right)}{b \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right) - \frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{3/2}}$$

input `Int[(d*Csc[a + b*x])^(7/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*(d*Csc[a + b*x])^(5/2))/(5*b*(c*Sec[a + b*x])^(3/2)) + (2*d^2*((-2*c*d*Sqrt[d*Csc[a + b*x]])/(b*(c*Sec[a + b*x])^(3/2)) - (2*d^2*EllipticE[a - Pi/4 + b*x, 2])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])))/5`

## 3.255.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

## 3.255.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 382 vs.  $2(133) = 266$ .

Time = 1.42 (sec) , antiderivative size = 383, normalized size of antiderivative = 2.99

method	result
default	$\frac{\sqrt{2} d^3 \sqrt{d \csc(bx+a)} \left( 4\sqrt{1+\csc(bx+a)} - \cot(bx+a) \sqrt{\cot(bx+a) - \csc(bx+a)+1} \sqrt{\cot(bx+a) - \csc(bx+a)} \right) \text{EllipticE}\left(\sqrt{1+\csc(bx+a)}\right)}{\dots}$

input `int((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

$$3.255. \int \frac{(d \csc(a+bx))^{7/2}}{\sqrt{c \sec(a+bx)}} dx$$

output  $1/5/b*2^{(1/2)}*d^3*(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}*(4*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticE}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})-2*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})+4*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticE}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*\sec(b*x+a)-2*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*\sec(b*x+a)-2*2^{(1/2)}-2^{(1/2)}*\cot(b*x+a)*\csc(b*x+a)$

### 3.255.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.94

$$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx = \frac{(d^3 \cos(bx + a)^2 - d^3) \sqrt{-4i cd} E(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) + (d^3 \cos(bx + a)^2 - d^3) \sqrt{4i cd} E(\arcsin(\cos(bx + a) - i \sin(bx + a)) | -1) - (d^3 \cos(bx + a)^2 - d^3) \sqrt{-4i cd} F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) - (d^3 \cos(bx + a)^2 - d^3) \sqrt{4i cd} F(\arcsin(\cos(bx + a) - i \sin(bx + a)) | -1) - 2*(2*d^3*\cos(b*x+a)^4 - 3*d^3*\cos(b*x+a)^2)*\sqrt{c/\cos(b*x+a)}*\sqrt{d/\sin(b*x+a)}}{(b*c*\cos(b*x+a)^2 - b*c)}$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fracas")`

output  $1/5*((d^3*\cos(b*x + a)^2 - d^3)*\sqrt{-4*I*c*d}*\text{elliptic\_e}(\arcsin(\cos(b*x + a) + I*\sin(b*x + a)), -1) + (d^3*\cos(b*x + a)^2 - d^3)*\sqrt{4*I*c*d}*\text{elliptic\_e}(\arcsin(\cos(b*x + a) - I*\sin(b*x + a)), -1) - (d^3*\cos(b*x + a)^2 - d^3)*\sqrt{-4*I*c*d}*\text{elliptic\_f}(\arcsin(\cos(b*x + a) + I*\sin(b*x + a)), -1) - (d^3*\cos(b*x + a)^2 - d^3)*\sqrt{4*I*c*d}*\text{elliptic\_f}(\arcsin(\cos(b*x + a) - I*\sin(b*x + a)), -1) - 2*(2*d^3*\cos(b*x + a)^4 - 3*d^3*\cos(b*x + a)^2)*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)})/(b*c*\cos(b*x + a)^2 - b*c)$

### 3.255.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(1/2),x)`

output Timed out

### 3.255.7 Maxima [F]

$$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{7/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(7/2)/sqrt(c*sec(b*x + a)), x)`

### 3.255.8 Giac [F]

$$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{7/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(7/2)/sqrt(c*sec(b*x + a)), x)`

### 3.255.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{7/2}}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

input `int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(1/2),x)`

output `int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(1/2), x)`



$$3.256 \quad \int \frac{(d \csc(a+bx))^{5/2}}{\sqrt{c \sec(a+bx)}} dx$$

3.256.1 Optimal result	1476
3.256.2 Mathematica [A] (verified)	1476
3.256.3 Rubi [A] (verified)	1477
3.256.4 Maple [A] (verified)	1478
3.256.5 Fricas [A] (verification not implemented)	1478
3.256.6 Sympy [F(-1)]	1478
3.256.7 Maxima [F]	1479
3.256.8 Giac [F]	1479
3.256.9 Mupad [B] (verification not implemented)	1479

### 3.256.1 Optimal result

Integrand size = 25, antiderivative size = 33

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{2cd(d \csc(a + bx))^{3/2}}{3b(c \sec(a + bx))^{3/2}}$$

output `-2/3*c*d*(d*csc(b*x+a))^(3/2)/b/(c*sec(b*x+a))^(3/2)`

### 3.256.2 Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{2cd(d \csc(a + bx))^{3/2}}{3b(c \sec(a + bx))^{3/2}}$$

input `Integrate[(d*Csc[a + b*x])^(5/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*(d*Csc[a + b*x])^(3/2))/(3*b*(c*Sec[a + b*x])^(3/2))`

**3.256.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx$$

↓ 3042

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx$$

↓ 3099

$$-\frac{2cd(d \csc(a + bx))^{3/2}}{3b(c \sec(a + bx))^{3/2}}$$

input `Int[(d*Csc[a + b*x])^(5/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*(d*Csc[a + b*x])^(3/2))/(3*b*(c*Sec[a + b*x])^(3/2))`

**3.256.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

**3.256.4 Maple [A] (verified)**

Time = 0.97 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

method	result	size
default	$-\frac{2\sqrt{d}\csc(bx+a)d^2\cot(bx+a)}{3b\sqrt{c}\sec(bx+a)}$	35

input `int((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`output `-2/3/b*(d*csc(b*x+a))^(1/2)*d^2/(c*sec(b*x+a))^(1/2)*cot(b*x+a)`**3.256.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.55

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{2 d^2 \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx + a)^2}{3 bc \sin(bx + a)}$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fracas")`output `-2/3*d^2*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)^2/(b*c*sin(b*x + a))`**3.256.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(1/2),x)`output `Timed out`

**3.256.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{5/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(5/2)/sqrt(c*sec(b*x + a)), x)`

**3.256.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{5/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(5/2)/sqrt(c*sec(b*x + a)), x)`

**3.256.9 Mupad [B] (verification not implemented)**

Time = 13.91 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.48

$$\int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{d^2 \sin(2a + 2bx) \sqrt{\frac{d}{\sin(a+bx)}}}{3b \sin(a + bx)^2 \sqrt{\frac{c}{\cos(a+bx)}}}$$

input `int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(1/2),x)`

output `-(d^2*sin(2*a + 2*b*x)*(d/sin(a + b*x))^(1/2))/(3*b*sin(a + b*x)^2*(c/cos(a + b*x))^(1/2))`

**3.257**       $\int \frac{(d \csc(a+bx))^{3/2}}{\sqrt{c \sec(a+bx)}} dx$

3.257.1 Optimal result . . . . . 1480  
 3.257.2 Mathematica [C] (verified) . . . . . 1480  
 3.257.3 Rubi [A] (verified) . . . . . 1481  
 3.257.4 Maple [B] (verified) . . . . . 1483  
 3.257.5 Fricas [C] (verification not implemented) . . . . . 1484  
 3.257.6 Sympy [F] . . . . . 1484  
 3.257.7 Maxima [F] . . . . . 1484  
 3.257.8 Giac [F] . . . . . 1485  
 3.257.9 Mupad [F(-1)] . . . . . 1485

**3.257.1 Optimal result**

Integrand size = 25, antiderivative size = 89

$$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{2cd\sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} - \frac{2d^2 E(a - \frac{\pi}{4} + bx | 2)}{b\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}\sqrt{\sin(2a + 2bx)}}$$

output

```
-2*c*d*(d*csc(b*x+a))^(1/2)/b/(c*sec(b*x+a))^(3/2)+2*d^2*(sin(a+1/4*Pi+b*x)
)^2^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/(d*csc
(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

**3.257.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.01 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.90

$$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx = \frac{2d^2 \left( \cot^2(a + bx) + \sqrt{-\cot^2(a + bx)} \operatorname{Hypergeometric2F1} \left( -\frac{1}{2}, \frac{1}{4}, \frac{1}{2}, \csc^2(a + bx) \right) \right) \tan(a + bx)}{b\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(3/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(-2*d^2*(Cot[a + b*x]^2 + (-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Tan[a + b*x])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])`

### 3.257.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3105, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx \\
 & \quad \downarrow \text{3105} \\
 & -2d^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -2d^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3110} \\
 & -\frac{2d^2 \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{\sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2d^2 \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{\sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3052}
 \end{aligned}$$

---

3.257.  $\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx$

$$\begin{aligned}
& -\frac{2d^2 \int \sqrt{\sin(2a + 2bx)} dx}{\sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& -\frac{2d^2 \int \sqrt{\sin(2a + 2bx)} dx}{\sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \\
& \quad \downarrow \text{3119} \\
& -\frac{2d^2 E(a + bx - \frac{\pi}{4} | 2)}{b \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}}
\end{aligned}$$

input `Int[(d*Csc[a + b*x])^(3/2)/Sqrt[c*Sec[a + b*x]],x]`

output `(-2*c*d*Sqrt[d*Csc[a + b*x]]/(b*(c*Sec[a + b*x])^(3/2)) - (2*d^2*EllipticE[a - Pi/4 + b*x, 2])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]]))`

### 3.257.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.257.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 362 vs.  $2(104) = 208$ .

Time = 1.20 (sec) , antiderivative size = 363, normalized size of antiderivative = 4.08

method	result
default	$\frac{\sqrt{2}d\sqrt{d\csc(bx+a)}\left(2\sqrt{1+\csc(bx+a)-\cot(bx+a)}\sqrt{\cot(bx+a)-\csc(bx+a)+1}\sqrt{\cot(bx+a)-\csc(bx+a)}\operatorname{EllipticE}\left(\sqrt{1+\csc(bx+a)}-\right.\right.$

```
input int((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/b*2^(1/2)*d*(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)*(2*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))-(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+2*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*sec(b*x+a)-(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*sec(b*x+a)-2^(1/2))
```



**3.257.5 Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.69

$$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx =$$

$$4 d \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx + a)^2 - \sqrt{-4i c d d} E(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) - \sqrt{4i c d d}$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`

output `-1/2*(4*d*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)^2 - sqrt(-4*I*c*d)*d*elliptic_e(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) - sqrt(4*I*c*d)*d*elliptic_e(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1) + sqrt(-4*I*c*d)*d*elliptic_f(arcsin(cos(b*x + a) + I*sin(b*x + a)), -1) + sqrt(4*I*c*d)*d*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1))/(b*c)`

**3.257.6 Sympy [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx$$

input `integrate((d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(1/2),x)`

output `Integral((d*csc(a + b*x))**(3/2)/sqrt(c*sec(a + b*x)), x)`

**3.257.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{3/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(3/2)/sqrt(c*sec(b*x + a)), x)`

---

3.257.  $\int \frac{(d \csc(a+bx))^{3/2}}{\sqrt{c \sec(a+bx)}} dx$

**3.257.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{(d \csc(bx + a))^{3/2}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(3/2)/sqrt(c*sec(b*x + a)), x)`

**3.257.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

input `int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(1/2),x)`

output `int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(1/2), x)`

**3.258**  $\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx$

3.258.1 Optimal result . . . . . 1486  
 3.258.2 Mathematica [A] (verified) . . . . . 1487  
 3.258.3 Rubi [A] (verified) . . . . . 1487  
 3.258.4 Maple [A] (verified) . . . . . 1491  
 3.258.5 Fracas [C] (verification not implemented) . . . . . 1492  
 3.258.6 Sympy [F] . . . . . 1492  
 3.258.7 Maxima [F] . . . . . 1493  
 3.258.8 Giac [F] . . . . . 1493  
 3.258.9 Mupad [F(-1)] . . . . . 1493

**3.258.1 Optimal result**

Integrand size = 25, antiderivative size = 270

$$\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx$$

$$= -\frac{\arctan\left(1 - \sqrt{2}\sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}{\sqrt{2}b\sqrt{c \sec(a+bx)}} + \frac{\arctan\left(1 + \sqrt{2}\sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}{\sqrt{2}b\sqrt{c \sec(a+bx)}} - \frac{\sqrt{d \csc(a+bx)} \log\left(1 - \sqrt{2}\sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{\tan(a+bx)}}{2\sqrt{2}b\sqrt{c \sec(a+bx)}} + \frac{\sqrt{d \csc(a+bx)} \log\left(1 + \sqrt{2}\sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{\tan(a+bx)}}{2\sqrt{2}b\sqrt{c \sec(a+bx)}}$$

```
output 1/2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)+1/2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/4*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)+1/4*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)
```

**3.258.2 Mathematica [A] (verified)**

Time = 2.19 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.46

$$\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx = \frac{\left( \arctan \left( \frac{-1 + \sqrt{\cot^2(a+bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a+bx)}} \right) - \operatorname{arctanh} \left( \frac{\sqrt{2} \sqrt[4]{\cot^2(a+bx)}}{1 + \sqrt{\cot^2(a+bx)}} \right) \right) \cot(a+bx) \sqrt{d \csc(a+bx)}}{\sqrt{2b \cot^2(a+bx)^{3/4} \sqrt{c \sec(a+bx)}}}$$

input `Integrate[Sqrt[d*Csc[a + b*x]]/Sqrt[c*Sec[a + b*x]],x]`output `-(((ArcTan[-1 + Sqrt[Cot[a + b*x]^2]]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))) - ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])])*Cot[a + b*x]*Sqrt[d*Csc[a + b*x]])/(Sqrt[2]*b*(Cot[a + b*x]^2)^(3/4)*Sqrt[c*Sec[a + b*x]]))`**3.258.3 Rubi [A] (verified)**Time = 0.46 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.63, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.520$ , Rules used = {3042, 3109, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx \\ & \quad \downarrow \text{3109} \\ & \frac{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{\sqrt{c \sec(a+bx)}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.258.  $\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx$

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}}dx}{\sqrt{c\sec(a+bx)}}$$

↓ 3957

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}(\tan^2(a+bx)+1)}d\tan(a+bx)}{b\sqrt{c\sec(a+bx)}}$$

↓ 266

$$\frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}}{b\sqrt{c\sec(a+bx)}}$$

↓ 755

$$\frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\int\frac{\tan(a+bx)+1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}\right)}{b\sqrt{c\sec(a+bx)}}$$

↓ 1476

$$\frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{1}{2}\int\frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{b\sqrt{c\sec(a+bx)}}$$

↓ 1082

$$\frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{\int\frac{1}{-\tan(a+bx)-1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}-\frac{\int\frac{1}{-\tan(a+bx)+1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{b\sqrt{c\sec(a+bx)}}$$

↓ 217

$$\frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\sqrt{2}}\right)}{\sqrt{2}}-\frac{\arctan\left(1-\sqrt{2}\sqrt{\tan(a+bx)+1}\right)}{\sqrt{2}}\right)\right)}{b\sqrt{c\sec(a+bx)}}$$

↓ 1479

$$\frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\int-\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}-\frac{\int-\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}\right)\right)}{b\sqrt{c\sec(a+bx)}}$$

↓ 25

---

3.258.  $\int\frac{\sqrt{d\csc(a+bx)}}{\sqrt{c\sec(a+bx)}}dx$

$$\begin{aligned}
& \frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}+\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}\right)\right)}{b\sqrt{c\sec(a+bx)}} \\
& \quad \downarrow 27 \\
& \frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\int\frac{\sqrt{2}\sqrt{\tan(a+bx)}+1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}\right)\right)}{b\sqrt{c\sec(a+bx)}} \\
& \quad \downarrow 1103 \\
& \frac{2\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)+\frac{1}{2}\left(\frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)})}{2\sqrt{2}}\right)}{b\sqrt{c\sec(a+bx)}}
\end{aligned}$$

input `Int[Sqrt[d*Csc[a + b*x]]/Sqrt[c*Sec[a + b*x]],x]`

output `(2*Sqrt[d*Csc[a + b*x]]*((-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[Tan[a + b*x]]/(b*Sqrt[c*Sec[a + b*x]]))`

### 3.258.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] & AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### 3.258.4 Maple [A] (verified)

Time = 1.22 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.34

method	result
default	$-\frac{\sqrt{2} \sin(bx+a)}{\left( \ln\left( 2\sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \cot(bx+a) + 2\sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \csc(bx+a) - 2 \cot(bx+a) + 2 \right) - 2 \arctan\left( \dots \right) \right)}$

input `int((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/4/b*2^{(1/2)}*\sin(b*x+a)*(\ln(2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}*\cot(b*x+a)+2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}*\csc(b*x+a)-2*\cot(b*x+a)+2)-2*\arctan((-\sin(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}+\cos(b*x+a)-1)/(\cos(b*x+a)-1))-\ln(-2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}*\cot(b*x+a)-2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}*\csc(b*x+a)-2*\cot(b*x+a)+2)+2*\arctan((\sin(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}+\cos(b*x+a)-1)/(\cos(b*x+a)-1)))*(d*csc(b*x+a))^(1/2)/(\cos(b*x+a)+1)/(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}/(c*sec(b*x+a))^(1/2)$$



### 3.258.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 1147, normalized size of antiderivative = 4.25

$$\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx = \text{Too large to display}$$

```
input integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fracas")
```

```
output -1/8*(-d^2/(b^4*c^2))^(1/4)*log(-1/2*d^2*cos(b*x + a)*sin(b*x + a) + 1/2*(
b*d*(-d^2/(b^4*c^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c*cos(b*x +
a)^3 - b^3*c*cos(b*x + a))*(-d^2/(b^4*c^2))^(3/4))*sqrt(c/cos(b*x + a))*sq
rt(d/sin(b*x + a)) - 1/4*(2*b^2*c*d*cos(b*x + a)^2 - b^2*c*d)*sqrt(-d^2/(b
^4*c^2))) + 1/8*(-d^2/(b^4*c^2))^(1/4)*log(-1/2*d^2*cos(b*x + a)*sin(b*x +
a) - 1/2*(b*d*(-d^2/(b^4*c^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c
*cos(b*x + a)^3 - b^3*c*cos(b*x + a))*(-d^2/(b^4*c^2))^(3/4))*sqrt(c/cos(b
*x + a))*sqrt(d/sin(b*x + a)) - 1/4*(2*b^2*c*d*cos(b*x + a)^2 - b^2*c*d)*s
qrt(-d^2/(b^4*c^2))) - 1/8*I*(-d^2/(b^4*c^2))^(1/4)*log(-1/2*d^2*cos(b*x +
a)*sin(b*x + a) + 1/2*(I*b*d*(-d^2/(b^4*c^2))^(1/4)*cos(b*x + a)^2*sin(b*
x + a) - (I*b^3*c*cos(b*x + a)^3 - I*b^3*c*cos(b*x + a))*(-d^2/(b^4*c^2))^(
3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1/4*(2*b^2*c*d*cos(b*x
+ a)^2 - b^2*c*d)*sqrt(-d^2/(b^4*c^2))) + 1/8*I*(-d^2/(b^4*c^2))^(1/4)*log
(-1/2*d^2*cos(b*x + a)*sin(b*x + a) + 1/2*(-I*b*d*(-d^2/(b^4*c^2))^(1/4)*c
os(b*x + a)^2*sin(b*x + a) - (-I*b^3*c*cos(b*x + a)^3 + I*b^3*c*cos(b*x +
a))*(-d^2/(b^4*c^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1/
4*(2*b^2*c*d*cos(b*x + a)^2 - b^2*c*d)*sqrt(-d^2/(b^4*c^2))) - 1/8*(-d^2/(
b^4*c^2))^(1/4)*log(d^2 + 2*(b^3*c*(-d^2/(b^4*c^2))^(3/4)*cos(b*x + a)^2*s
in(b*x + a) + (b*d*cos(b*x + a)^3 - b*d*cos(b*x + a))*(-d^2/(b^4*c^2))^(1/
4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) + 1/8*(-d^2/(b^4*c^2))^(...
```

### 3.258.6 Sympy [F]

$$\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx$$

```
input integrate((d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(1/2),x)
```

```
output Integral(sqrt(d*csc(a + b*x))/sqrt(c*sec(a + b*x)), x)
```

---

3.258.  $\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx$

**3.258.7 Maxima [F]**

$$\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{\sqrt{d \csc(bx + a)}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(d*csc(b*x + a))/sqrt(c*sec(b*x + a)), x)`

**3.258.8 Giac [F]**

$$\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{\sqrt{d \csc(bx + a)}}{\sqrt{c \sec(bx + a)}} dx$$

input `integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(d*csc(b*x + a))/sqrt(c*sec(b*x + a)), x)`

**3.258.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx = \int \frac{\sqrt{\frac{d}{\sin(a+bx)}}}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

input `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(1/2),x)`

output `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(1/2), x)`

**3.259**  $\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx$

3.259.1 Optimal result . . . . . 1494  
 3.259.2 Mathematica [C] (verified) . . . . . 1494  
 3.259.3 Rubi [A] (verified) . . . . . 1495  
 3.259.4 Maple [B] (verified) . . . . . 1496  
 3.259.5 Fricas [F] . . . . . 1497  
 3.259.6 Sympy [F] . . . . . 1497  
 3.259.7 Maxima [F] . . . . . 1498  
 3.259.8 Giac [F] . . . . . 1498  
 3.259.9 Mupad [F(-1)] . . . . . 1498

**3.259.1 Optimal result**

Integrand size = 25, antiderivative size = 53

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \frac{E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{b \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{\sin(2a+2bx)}}$$

output  $-(\sin(a+1/4*\pi+b*x)^2)^{(1/2)}/\sin(a+1/4*\pi+b*x)*\text{EllipticE}(\cos(a+1/4*\pi+b*x), 2^{(1/2)})/b/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

**3.259.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.81 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.25

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \frac{\sqrt[4]{-\cot^2(a+bx)} \text{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{4}, \frac{1}{2}, \csc^2(a+bx)\right) \tan(a+bx)}{b \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}}$$

input `Integrate[1/(Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]),x]`

output  $((-\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Hypergeometric2F1}[-1/2, 1/4, 1/2, \text{Csc}[a + b*x]^2]*\text{Tan}[a + b*x])/(b*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

**3.259.3 Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} dx \\
 & \quad \downarrow \text{3110} \\
 & \frac{\int \sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)} dx}{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)} dx}{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3052} \\
 & \frac{\int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{E\left(a+bx-\frac{\pi}{4} \mid 2\right)}{b \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}
 \end{aligned}$$

input `Int[1/(Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]),x]`

output `EllipticE[a - Pi/4 + b*x, 2]/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])`

## 3.259.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_)])^n , x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]] , x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

## 3.259.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 392 vs.  $2(73) = 146$ .

Time = 1.31 (sec) , antiderivative size = 393, normalized size of antiderivative = 7.42

method	result
default	$-\frac{\sqrt{2} \left( 2\sqrt{1+\csc(bx+a)}-\cot(bx+a) \right) \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)}}{\sqrt{1+\csc(bx+a)}-\cot(bx+a)} \operatorname{EllipticE}\left(\sqrt{1+\csc(bx+a)}-\cot(bx+a), \frac{\sqrt{2}}{2}\right)$

input `int(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/2/b*2^{(1/2)}*(2*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticE}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*\cos(b*x+a)-(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*\cos(b*x+a)+2*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticE}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})-(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})+2^{(1/2)}*\cos(b*x+a)^2-2^{(1/2)}*\cos(b*x+a))/(c*\sec(b*x+a))^{(1/2)}/(d*\csc(b*x+a))^{(1/2)}*\sec(b*x+a)*\csc(b*x+a)$$

### 3.259.5 Fracas [F]

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \int \frac{1}{\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fracas")`

output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c*d*csc(b*x + a)*sec(b*x + a)), x)`

### 3.259.6 Sympy [F]

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \int \frac{1}{\sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} dx$$

input `integrate(1/(d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(1/2),x)`

output `Integral(1/(sqrt(c*sec(a + b*x))*sqrt(d*csc(a + b*x))), x)`

**3.259.7 Maxima [F]**

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \int \frac{1}{\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))), x)`

**3.259.8 Giac [F]**

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \int \frac{1}{\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))), x)`

**3.259.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}}} dx$$

input `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)),x)`

output `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)), x)`

**3.260**  $\int \frac{1}{(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} dx$

3.260.1 Optimal result . . . . . 1499  
 3.260.2 Mathematica [A] (verified) . . . . . 1500  
 3.260.3 Rubi [A] (verified) . . . . . 1500  
 3.260.4 Maple [A] (verified) . . . . . 1505  
 3.260.5 Fricas [C] (verification not implemented) . . . . . 1506  
 3.260.6 Sympy [F] . . . . . 1506  
 3.260.7 Maxima [F] . . . . . 1507  
 3.260.8 Giac [F] . . . . . 1507  
 3.260.9 Mupad [F(-1)] . . . . . 1507

**3.260.1 Optimal result**

Integrand size = 25, antiderivative size = 322

$$\int \frac{1}{(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} dx = -\frac{c}{2bd \sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}}$$

$$-\frac{\arctan\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}{4\sqrt{2}bd^2 \sqrt{c \sec(a+bx)}}$$

$$+\frac{\arctan\left(1 + \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}{4\sqrt{2}bd^2 \sqrt{c \sec(a+bx)}}$$

$$-\frac{\sqrt{d \csc(a+bx)} \log\left(1 - \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{\tan(a+bx)}}{8\sqrt{2}bd^2 \sqrt{c \sec(a+bx)}}$$

$$+\frac{\sqrt{d \csc(a+bx)} \log\left(1 + \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{\tan(a+bx)}}{8\sqrt{2}bd^2 \sqrt{c \sec(a+bx)}}$$

output

```
-1/2*c/b/d/(c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2)+1/8*arctan(-1+2^(1/2)
*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*
sec(b*x+a))^(1/2)+1/8*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1
/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/16*ln(1-2^(1/2)*
tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2
^(1/2)/(c*sec(b*x+a))^(1/2)+1/16*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))
*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)
```



**3.260.2 Mathematica [A] (verified)**

Time = 2.67 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.48

$$\int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx =$$

$$\frac{\left(4 \cos^2(a + bx) + \sqrt{2} \arctan\left(\frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}\right) \sqrt[4]{\cot^2(a + bx)} - \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}{1 + \sqrt{\cot^2(a + bx)}}\right) \sqrt[4]{\cot^2(a + bx)}\right)}{8bd^2 \sqrt{c \sec(a + bx)}}$$

input `Integrate[1/((d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]]),x]`output `-1/8*((4*Cos[a + b*x]^2 + Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2])]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(1/4) - Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(1/4))*Sqrt[d*Csc[a + b*x]]*Tan[a + b*x])/(b*d^2*Sqrt[c*Sec[a + b*x]])`**3.260.3 Rubi [A] (verified)**Time = 0.60 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.66, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3107, 3042, 3109, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{\sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}} dx$$

$$\downarrow \text{3107}$$

$$\frac{\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx}{4d^2} - \frac{c}{2bd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}$$

$$\downarrow \text{3042}$$

---


$$3.260. \quad \int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx$$

$$\begin{aligned}
& \frac{\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4d^2} - \frac{c}{2bd(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3109} \\
& \frac{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{4d^2 \sqrt{c \sec(a+bx)}} - \frac{c}{2bd(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{4d^2 \sqrt{c \sec(a+bx)}} - \frac{c}{2bd(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3957} \\
& \frac{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)(\tan^2(a+bx)+1)}} d \tan(a+bx)}{4bd^2 \sqrt{c \sec(a+bx)}} - \frac{c}{2bd(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{266} \\
& \frac{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{2bd^2 \sqrt{c \sec(a+bx)}} - \frac{c}{2bd(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{755} \\
& \frac{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{2bd^2 \sqrt{c \sec(a+bx)}} - \frac{c}{2bd(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{1476} \\
& \frac{\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} + \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \right. \right. \\
& \quad \left. \left. \frac{c}{2bd^2 \sqrt{c \sec(a+bx)}} \right) \right)}{2bd(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{1082}
\end{aligned}$$

---

3.260.  $\int \frac{1}{(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} dx$

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{\int\frac{1}{-\tan(a+bx)-1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}-\frac{\int\frac{1}{-\tan(a+bx)}}{\sqrt{2}}\right)\right)}{2bd^2\sqrt{c\sec(a+bx)}}$$


---


$$\frac{c}{2bd(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 217

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{2bd^2\sqrt{c\sec(a+bx)}}$$


---


$$\frac{c}{2bd(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 1479

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(-\frac{\int-\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}-\frac{\int-\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}\right)\right)}{2bd^2\sqrt{c\sec(a+bx)}}$$


---


$$\frac{c}{2bd(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 25

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}+\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}\right)\right)+\frac{1}{2}}{2bd^2\sqrt{c\sec(a+bx)}}$$


---


$$\frac{c}{2bd(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 27

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}+\frac{1}{2}\int\frac{\sqrt{2}\sqrt{\tan(a+bx)}+1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}\right)\right)}{2bd^2\sqrt{c\sec(a+bx)}}$$


---


$$\frac{c}{2bd(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 1103

$$\frac{\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)+\frac{1}{2}\left(\frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)})}{2\sqrt{2}}\right)\right)}{2bd^2\sqrt{c\sec(a+bx)}}$$

$$\frac{c}{2bd(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

input `Int[1/((d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]]),x]`

output `-1/2*c/(b*d*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)) + (Sqrt[d*Csc[a + b*x]]*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[Tan[a + b*x]]/(2*b*d^2*Sqrt[c*Sec[a + b*x]])`

### 3.260.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3107 `Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n  
 ), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n  
 ) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !Integer  
 Q[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^n, x_Symbol] := Simp[b/d Subst[Int  
 [x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&  
 !IntegerQ[n]`

### 3.260.4 Maple [A] (verified)

Time = 1.73 (sec) , antiderivative size = 434, normalized size of antiderivative = 1.35

method	result
default	$\frac{\sqrt{2} \left( 4 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sqrt{2} \cos(bx+a)^2 + 4 \cos(bx+a) \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} - 2 \arctan \left( \frac{-\sin(bx+a) \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}}}{\cos(bx+a)-1} \right) \right)}{\dots}$

input `int(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/16/b*2^(1/2)*(4*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*2^(1/2)  
 *cos(b*x+a)^2+4*cos(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)  
 ^2)^(1/2)-2*arctan((-sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)  
 +1)^2)^(1/2)+cos(b*x+a)-1)/(cos(b*x+a)-1))+ln(2*2^(1/2)*(-cos(b*x+a)*sin(b  
 *x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)+2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)  
 /(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2)-ln(-2*2^(1/2)*(-cos(b*  
 x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)-2*2^(1/2)*(-cos(b*x+a)*  
 sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2)+2*arctan((si  
 n(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+cos(b*x+a  
 )-1)/(cos(b*x+a)-1)))/(cos(b*x+a)+1)/(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1  
 )^2)^(1/2)/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/d`

**3.260.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.67 (sec) , antiderivative size = 1268, normalized size of antiderivative = 3.94

$$\int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx = \text{Too large to display}$$

```
input integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fracas")
```

```
output -1/32*(b*c*d^2*(-1/(b^4*c^2*d^6))^(1/4)*log(2*(b^3*c*d^4*(-1/(b^4*c^2*d^6))^(3/4)*cos(b*x + a)^2*sin(b*x + a) + (b*d*cos(b*x + a)^3 - b*d*cos(b*x + a))*(-1/(b^4*c^2*d^6))^(1/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1) - b*c*d^2*(-1/(b^4*c^2*d^6))^(1/4)*log(-2*(b^3*c*d^4*(-1/(b^4*c^2*d^6))^(3/4)*cos(b*x + a)^2*sin(b*x + a) + (b*d*cos(b*x + a)^3 - b*d*cos(b*x + a))*(-1/(b^4*c^2*d^6))^(1/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1) + I*b*c*d^2*(-1/(b^4*c^2*d^6))^(1/4)*log(-2*(I*b^3*c*d^4*(-1/(b^4*c^2*d^6))^(3/4)*cos(b*x + a)^2*sin(b*x + a) + (-I*b*d*cos(b*x + a)^3 + I*b*d*cos(b*x + a))*(-1/(b^4*c^2*d^6))^(1/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1) - I*b*c*d^2*(-1/(b^4*c^2*d^6))^(1/4)*log(-2*(-I*b^3*c*d^4*(-1/(b^4*c^2*d^6))^(3/4)*cos(b*x + a)^2*sin(b*x + a) + (I*b*d*cos(b*x + a)^3 - I*b*d*cos(b*x + a))*(-1/(b^4*c^2*d^6))^(1/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1) + b*c*d^2*(-1/(b^4*c^2*d^6))^(1/4)*log(1/2*(b*d*(-1/(b^4*c^2*d^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c*d^4*cos(b*x + a)^3 - b^3*c*d^4*cos(b*x + a))*(-1/(b^4*c^2*d^6))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 1/2*cos(b*x + a)*sin(b*x + a) - 1/4*(2*b^2*c*d^3*cos(b*x + a)^2 - b^2*c*d^3)*sqrt(-1/(b^4*c^2*d^6))) - b*c*d^2*(-1/(b^4*c^2*d^6))^(1/4)*log(-1/2*(b*d*(-1/(b^4*c^2*d^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c*d^4*cos(b*x + a)^3 - b^3*c*d^4*cos(b*x + a))*(-1/(b^4*c^2*d^6))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 1/2*cos(b*x + a)*si...
```

**3.260.6 Sympy [F]**

$$\int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}} dx$$

```
input integrate(1/(d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(1/2),x)
```

output `Integral(1/(sqrt(c*sec(a + b*x))*(d*csc(a + b*x))**(3/2)), x)`

### 3.260.7 Maxima [F]

$$\int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a))), x)`

### 3.260.8 Giac [F]

$$\int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(1/((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a))), x)`

### 3.260.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}} \left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

input `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(3/2)),x)`

output `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(3/2)), x)`

---

3.260.  $\int \frac{1}{(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} dx$



**3.261**  $\int \frac{1}{(d \csc(a+bx))^{5/2} \sqrt{c \sec(a+bx)}} dx$

3.261.1 Optimal result . . . . . 1508  
 3.261.2 Mathematica [C] (verified) . . . . . 1508  
 3.261.3 Rubi [A] (verified) . . . . . 1509  
 3.261.4 Maple [B] (verified) . . . . . 1511  
 3.261.5 Fricas [F] . . . . . 1512  
 3.261.6 Sympy [F(-1)] . . . . . 1512  
 3.261.7 Maxima [F] . . . . . 1512  
 3.261.8 Giac [F] . . . . . 1513  
 3.261.9 Mupad [F(-1)] . . . . . 1513

**3.261.1 Optimal result**

Integrand size = 25, antiderivative size = 95

$$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx = -\frac{c}{3bd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}} + \frac{E(a - \frac{\pi}{4} + bx | 2)}{2bd^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}$$

output

```
-1/3*c/b/d/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2)-1/2*(sin(a+1/4*Pi+b*x)
)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/d^2/(d
*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

**3.261.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.06 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.88

$$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx = \frac{\left(1 + \cos(2(a + bx)) - 3\sqrt[4]{-\cot^2(a + bx)} \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{4}, \frac{1}{2}, \csc^2(a + bx)\right)\right) \tan(a + bx)}{6bd^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}$$

input `Integrate[1/((d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]]),x]`

output `-1/6*((1 + Cos[2*(a + b*x)] - 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Tan[a + b*x])/(b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])`

### 3.261.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3107, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{c \sec(a+bx)}(d \csc(a+bx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{c \sec(a+bx)}(d \csc(a+bx))^{5/2}} dx \\
 & \quad \downarrow \text{3107} \\
 & \frac{\int \frac{1}{\sqrt{d \csc(a+bx)}\sqrt{c \sec(a+bx)}} dx}{2d^2} - \frac{c}{3bd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{1}{\sqrt{d \csc(a+bx)}\sqrt{c \sec(a+bx)}} dx}{2d^2} - \frac{c}{3bd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow \text{3110} \\
 & \frac{\int \sqrt{c \cos(a+bx)}\sqrt{d \sin(a+bx)} dx}{2d^2 \sqrt{c \cos(a+bx)}\sqrt{c \sec(a+bx)}\sqrt{d \sin(a+bx)}\sqrt{d \csc(a+bx)}} - \frac{c}{3bd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{c \cos(a+bx)}\sqrt{d \sin(a+bx)} dx}{2d^2 \sqrt{c \cos(a+bx)}\sqrt{c \sec(a+bx)}\sqrt{d \sin(a+bx)}\sqrt{d \csc(a+bx)}} - \frac{c}{3bd(c \sec(a+bx))^{3/2}(d \csc(a+bx))^{3/2}}
 \end{aligned}$$

---

3.261.  $\int \frac{1}{(d \csc(a+bx))^{5/2} \sqrt{c \sec(a+bx)}} dx$

$$\begin{aligned} & \int \frac{\sqrt{\sin(2a + 2bx)} dx}{2d^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{c}{3bd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \\ & \qquad \qquad \qquad \downarrow \text{3052} \\ & \int \frac{\sqrt{\sin(2a + 2bx)} dx}{2d^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{c}{3bd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \\ & \qquad \qquad \qquad \downarrow \text{3042} \\ & \frac{E\left(a + bx - \frac{\pi}{4} \mid 2\right)}{2bd^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{c}{3bd(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \\ & \qquad \qquad \qquad \downarrow \text{3119} \end{aligned}$$

input `Int[1/((d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]]),x]`

output `-1/3*c/(b*d*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2)) + EllipticE[a - Pi/4 + b*x, 2]/(2*b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])`

### 3.261.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3107 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*SIN[e + f*x])^m*(b*cos[e + f*x])^n Int[1/((a*SIN[e + f*x])^m*(b*cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.261.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 409 vs.  $2(106) = 212$ .

Time = 1.24 (sec) , antiderivative size = 410, normalized size of antiderivative = 4.32

method	result
default	$-\frac{\sqrt{2} \left( 6\sqrt{1+\csc(bx+a)}-\cot(bx+a) \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)} \operatorname{EllipticE}\left(\sqrt{1+\csc(bx+a)}-\cot(bx+a), \frac{\sqrt{2}}{2}\right) \right)}{d^2 \sec(bx+a) \csc(bx+a)}$

```
input int(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/12/b*2^(1/2)*(6*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)-3*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)-2*cos(b*x+a)^4*2^(1/2)+6*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))-3*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+5*2^(1/2)*cos(b*x+a)^2-3*2^(1/2)*cos(b*x+a))/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/d^2*sec(b*x+a)*csc(b*x+a)
```

**3.261.5 Fracas [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{(d \csc(bx + a))^{5/2} \sqrt{c \sec(bx + a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")`

output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c*d^3*csc(b*x + a)^3*sec(b*x + a)), x)`

**3.261.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx = \text{Timed out}$$

input `integrate(1/(d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(1/2),x)`

output `Timed out`

**3.261.7 Maxima [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{(d \csc(bx + a))^{5/2} \sqrt{c \sec(bx + a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a))), x)`

**3.261.8 Giac [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} \sqrt{c \sec(bx + a)}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(1/((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a))), x)`

**3.261.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx = \int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}} \left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

input `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2)),x)`

output `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2)), x)`

**3.262**       $\int \frac{(d \csc(a+bx))^{11/2}}{(c \sec(a+bx))^{3/2}} dx$

3.262.1 Optimal result . . . . . 1514  
 3.262.2 Mathematica [A] (verified) . . . . . 1514  
 3.262.3 Rubi [A] (verified) . . . . . 1515  
 3.262.4 Maple [A] (verified) . . . . . 1516  
 3.262.5 Fricas [A] (verification not implemented) . . . . . 1517  
 3.262.6 Sympy [F(-1)] . . . . . 1517  
 3.262.7 Maxima [F] . . . . . 1517  
 3.262.8 Giac [F] . . . . . 1518  
 3.262.9 Mupad [B] (verification not implemented) . . . . . 1518

**3.262.1 Optimal result**

Integrand size = 25, antiderivative size = 110

$$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{8d^5 \sqrt{d \csc(a + bx)}}{45bc \sqrt{c \sec(a + bx)}} + \frac{2d^3 (d \csc(a + bx))^{5/2}}{45bc \sqrt{c \sec(a + bx)}} - \frac{2d (d \csc(a + bx))^{9/2}}{9bc \sqrt{c \sec(a + bx)}}$$

output `2/45*d^3*(d*csc(b*x+a))^(5/2)/b/c/(c*sec(b*x+a))^(1/2)-2/9*d*(d*csc(b*x+a))^(9/2)/b/c/(c*sec(b*x+a))^(1/2)+8/45*d^5*(d*csc(b*x+a))^(1/2)/b/c/(c*sec(b*x+a))^(1/2)`

**3.262.2 Mathematica [A] (verified)**

Time = 1.73 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.52

$$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{2d^3(-7 + 2 \cos(2(a + bx))) \cot^2(a + bx)(d \csc(a + bx))^{5/2}}{45bc \sqrt{c \sec(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(11/2)/(c*Sec[a + b*x])^(3/2),x]`

output `(2*d^3*(-7 + 2*Cos[2*(a + b*x)])*Cot[a + b*x]^2*(d*Csc[a + b*x])^(5/2))/(45*b*c*Sqrt[c*Sec[a + b*x]])`

**3.262.3 Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {3042, 3103, 3042, 3105, 3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3103} \\
 & -\frac{d^2 \int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx}{9c^2} - \frac{2d(d \csc(a + bx))^{9/2}}{9bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{d^2 \int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx}{9c^2} - \frac{2d(d \csc(a + bx))^{9/2}}{9bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3105} \\
 & -\frac{d^2 \left( \frac{4}{5} d^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx - \frac{2cd(d \csc(a + bx))^{5/2}}{5b \sqrt{c \sec(a + bx)}} \right)}{9c^2} - \frac{2d(d \csc(a + bx))^{9/2}}{9bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{d^2 \left( \frac{4}{5} d^2 \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx - \frac{2cd(d \csc(a + bx))^{5/2}}{5b \sqrt{c \sec(a + bx)}} \right)}{9c^2} - \frac{2d(d \csc(a + bx))^{9/2}}{9bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3099} \\
 & -\frac{d^2 \left( -\frac{8cd^3 \sqrt{d \csc(a + bx)}}{5b \sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b \sqrt{c \sec(a + bx)}} \right)}{9c^2} - \frac{2d(d \csc(a + bx))^{9/2}}{9bc \sqrt{c \sec(a + bx)}}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(11/2)/(c*Sec[a + b*x])^(3/2), x]`



```
output (-2*d*(d*Csc[a + b*x])^(9/2))/(9*b*c*Sqrt[c*Sec[a + b*x]]) - (d^2*((-8*c*d
^3*Sqrt[d*Csc[a + b*x]])/(5*b*Sqrt[c*Sec[a + b*x]]) - (2*c*d*(d*Csc[a + b*
x])^(5/2))/(5*b*Sqrt[c*Sec[a + b*x]])))/(9*c^2)
```

### 3.262.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3099 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n
_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1
)/(f*(n - 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] &&
NeQ[n, 1]
```

```
rule 3103 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n
_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n +
1)/(f*b*(m - 1))), x] + Simp[a^2*((n + 1)/(b^2*(m - 1))) Int[(a*Csc[e + f
*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] &&
GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]
```

```
rule 3105 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n
_), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n
- 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*
x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[
m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]
```

### 3.262.4 Maple [A] (verified)

Time = 1.26 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.55

method	result	size
default	$\frac{2d^5 \sqrt{d \csc(bx+a)} (4 \cot(bx+a)^4 - 9 \cot(bx+a)^2 \csc(bx+a)^2)}{45b \sqrt{c \sec(bx+a)} c}$	61

```
input int((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

output  $2/45/b*d^5*(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/c*(4*cot(b*x+a)^4-9*cot(b*x+a)^2*csc(b*x+a)^2)$

### 3.262.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.80

$$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{2(4d^5 \cos(bx + a)^5 - 9d^5 \cos(bx + a)^3) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{45(bc^2 \cos(bx + a)^4 - 2bc^2 \cos(bx + a)^2 + bc^2)}$$

input `integrate((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

output  $2/45*(4*d^5*\cos(b*x + a)^5 - 9*d^5*\cos(b*x + a)^3)*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)}/(b*c^2*\cos(b*x + a)^4 - 2*b*c^2*\cos(b*x + a)^2 + b*c^2)$

### 3.262.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(11/2)/(c*sec(b*x+a))**(3/2),x)`

output Timed out

### 3.262.7 Maxima [F]

$$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{\frac{11}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

input `integrate((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(11/2)/(c*sec(b*x + a))^(3/2), x)`

---

3.262.  $\int \frac{(d \csc(a+bx))^{11/2}}{(c \sec(a+bx))^{3/2}} dx$

**3.262.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{11/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(11/2)/(c*sec(b*x + a))^(3/2), x)`

**3.262.9 Mupad [B] (verification not implemented)**

Time = 16.49 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.14

$$\int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{8 d^5 \sqrt{\frac{d}{\sin(a+bx)}} (9 \cos(2a + 2bx) + 14 \cos(4a + 4bx) - 9 \cos(6a + 6bx) + \cos(8a + 8bx) - 15)}{45 b c \sqrt{\frac{c}{\cos(a+bx)}} (28 \cos(4a + 4bx) - 56 \cos(2a + 2bx) - 8 \cos(6a + 6bx) + \cos(8a + 8bx) + 35)}$$

input `int((d/sin(a + b*x))^(11/2)/(c/cos(a + b*x))^(3/2),x)`

output `(8*d^5*(d/sin(a + b*x))^(1/2)*(9*cos(2*a + 2*b*x) + 14*cos(4*a + 4*b*x) - 9*cos(6*a + 6*b*x) + cos(8*a + 8*b*x) - 15))/(45*b*c*(c/cos(a + b*x))^(1/2)*(28*cos(4*a + 4*b*x) - 56*cos(2*a + 2*b*x) - 8*cos(6*a + 6*b*x) + cos(8*a + 8*b*x) + 35))`

**3.263**  $\int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{3/2}} dx$

3.263.1 Optimal result . . . . . 1519  
 3.263.2 Mathematica [C] (verified) . . . . . 1519  
 3.263.3 Rubi [A] (verified) . . . . . 1520  
 3.263.4 Maple [A] (verified) . . . . . 1523  
 3.263.5 Fricas [C] (verification not implemented) . . . . . 1523  
 3.263.6 Sympy [F(-1)] . . . . . 1524  
 3.263.7 Maxima [F] . . . . . 1524  
 3.263.8 Giac [F] . . . . . 1524  
 3.263.9 Mupad [F(-1)] . . . . . 1525

**3.263.1 Optimal result**

Integrand size = 25, antiderivative size = 135

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{2d^3(d \csc(a + bx))^{3/2}}{21bc\sqrt{c \sec(a + bx)}} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc\sqrt{c \sec(a + bx)}} - \frac{2d^4\sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{21bc^2}$$

output

```
2/21*d^3*(d*csc(b*x+a))^(3/2)/b/c/(c*sec(b*x+a))^(1/2)-2/7*d*(d*csc(b*x+a))^(7/2)/b/c/(c*sec(b*x+a))^(1/2)+2/21*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b/c^2
```

**3.263.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.61 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.88

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{d^3 \cos(2(a + bx))(d \csc(a + bx))^{3/2} \left( (5 + \cos(2(a + bx))) \csc^4(a + bx) - 2(-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{\cos(2(a + bx))}{1 + \cos(2(a + bx))}\right) \right)}{21bc(-2 + \csc^2(a + bx))\sqrt{c \sec(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(9/2)/(c*Sec[a + b*x])^(3/2),x]`

output `-1/21*(d^3*Cos[2*(a + b*x)]*(d*Csc[a + b*x])^(3/2)*((5 + Cos[2*(a + b*x)])  
*Csc[a + b*x]^4 - 2*(-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/  
2, Csc[a + b*x]^2]*Sec[a + b*x]^2))/(b*c*(-2 + Csc[a + b*x]^2)*Sqrt[c*Sec[  
a + b*x]])`

### 3.263.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3103, 3042, 3105, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3103} \\
 & -\frac{d^2 \int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx}{7c^2} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{d^2 \int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx}{7c^2} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3105} \\
 & -\frac{d^2 \left( \frac{2}{3} d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx - \frac{2cd(d \csc(a + bx))^{3/2}}{3b \sqrt{c \sec(a + bx)}} \right)}{7c^2} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{d^2 \left( \frac{2}{3} d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx - \frac{2cd(d \csc(a + bx))^{3/2}}{3b \sqrt{c \sec(a + bx)}} \right)}{7c^2} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc \sqrt{c \sec(a + bx)}} \\
 & \quad \downarrow \text{3110}
 \end{aligned}$$

---

3.263.  $\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx$

$$\frac{d^2 \left( \frac{2}{3} d^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx - \frac{2cd(d \csc(a+bx))}{3b\sqrt{c \sec(a+bx)}} \right)}{\frac{7c^2}{7bc\sqrt{c \sec(a+bx)}} \frac{2d(d \csc(a+bx))^{7/2}}{7bc\sqrt{c \sec(a+bx)}}}$$

↓ 3042

$$\frac{d^2 \left( \frac{2}{3} d^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx - \frac{2cd(d \csc(a+bx))}{3b\sqrt{c \sec(a+bx)}} \right)}{\frac{7c^2}{7bc\sqrt{c \sec(a+bx)}} \frac{2d(d \csc(a+bx))^{7/2}}{7bc\sqrt{c \sec(a+bx)}}}$$

↓ 3053

$$\frac{d^2 \left( \frac{2}{3} d^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx - \frac{2cd(d \csc(a+bx))^{3/2}}{3b\sqrt{c \sec(a+bx)}} \right)}{\frac{7c^2}{7bc\sqrt{c \sec(a+bx)}} \frac{2d(d \csc(a+bx))^{7/2}}{7bc\sqrt{c \sec(a+bx)}}}$$

↓ 3042

$$\frac{d^2 \left( \frac{2}{3} d^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx - \frac{2cd(d \csc(a+bx))^{3/2}}{3b\sqrt{c \sec(a+bx)}} \right)}{\frac{7c^2}{7bc\sqrt{c \sec(a+bx)}} \frac{2d(d \csc(a+bx))^{7/2}}{7bc\sqrt{c \sec(a+bx)}}}$$

↓ 3120

$$\frac{d^2 \left( \frac{2d^2 \sqrt{\sin(2a+2bx)} \operatorname{EllipticF}\left(a+bx-\frac{\pi}{4}, 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3b} - \frac{2cd(d \csc(a+bx))^{3/2}}{3b\sqrt{c \sec(a+bx)}} \right)}{\frac{7c^2}{7bc\sqrt{c \sec(a+bx)}} \frac{2d(d \csc(a+bx))^{7/2}}{7bc\sqrt{c \sec(a+bx)}}}$$

input `Int[(d*Csc[a + b*x])^(9/2)/(c*Sec[a + b*x])^(3/2),x]`

output `(-2*d*(d*Csc[a + b*x])^(7/2))/(7*b*c*Sqrt[c*Sec[a + b*x]]) - (d^2*((-2*c*d*(d*Csc[a + b*x])^(3/2))/(3*b*Sqrt[c*Sec[a + b*x]])) + (2*d^2*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(3*b)))/(7*c^2)`

## 3.263.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3103 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Simp[a^2*((n + 1)/(b^2*(m - 1))) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]`

rule 3105 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

**3.263.4 Maple [A] (verified)**

Time = 1.03 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.71

method	result
default	$-\frac{\sqrt{2}d^4\sqrt{d\csc(bx+a)}\left(2\sqrt{1+\csc(bx+a)}-\cot(bx+a)\sqrt{\cot(bx+a)-\csc(bx+a)+1}\sqrt{\cot(bx+a)-\csc(bx+a)}\right)\text{EllipticF}\left(\sqrt{1+\csc(bx+a)}\right)}{\dots}$

```
input int((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -1/21/b*2^(1/2)*d^4*(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/c*(2*(1+csc(
b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b
*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+2*(1+c
sc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-cs
c(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*sec
(b*x+a)+2^(1/2)*cot(b*x+a)^2*csc(b*x+a)+2*2^(1/2)*csc(b*x+a)^3)
```

**3.263.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.36

$$\int \frac{(d\csc(a+bx))^{9/2}}{(c\sec(a+bx))^{3/2}} dx = \frac{(i d^4 \cos(bx+a)^2 - i d^4) \sqrt{-4i cd} F(\arcsin(\cos(bx+a) + i \sin(bx+a)) | -1) \sin(bx+a) + (-i d^4 \cos(bx+a)^2 + i d^4) \sqrt{4i cd} F(\arcsin(\cos(bx+a) - i \sin(bx+a)) | -1) \sin(bx+a) + 2(d^4 \cos(bx+a)^3 + 2d^4 \cos(bx+a)) \sqrt{c/\cos(bx+a)} \sqrt{d/\sin(bx+a)}}{(b^2 c^2 \cos(bx+a)^2 - b^2 c^2) \sin(bx+a)}$$

```
input integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")
```

```
output 1/21*((I*d^4*cos(b*x + a)^2 - I*d^4)*sqrt(-4*I*c*d)*elliptic_f(arcsin(cos(
b*x + a) + I*sin(b*x + a)), -1)*sin(b*x + a) + (-I*d^4*cos(b*x + a)^2 + I*
d^4)*sqrt(4*I*c*d)*elliptic_f(arcsin(cos(b*x + a) - I*sin(b*x + a)), -1)*s
in(b*x + a) + 2*(d^4*cos(b*x + a)^3 + 2*d^4*cos(b*x + a))*sqrt(c/cos(b*x +
a))*sqrt(d/sin(b*x + a)))/((b*c^2*cos(b*x + a)^2 - b*c^2)*sin(b*x + a))
```



**3.263.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(9/2)/(c*sec(b*x+a))**(3/2),x)`output `Timed out`**3.263.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{9/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(9/2)/(c*sec(b*x + a))^(3/2), x)`**3.263.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{9/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`output `integrate((d*csc(b*x + a))^(9/2)/(c*sec(b*x + a))^(3/2), x)`

**3.263.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{9/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

input `int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(3/2),x)`output `int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(3/2), x)`

**3.264**       $\int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{3/2}} dx$

3.264.1 Optimal result . . . . . 1526  
 3.264.2 Mathematica [A] (verified) . . . . . 1526  
 3.264.3 Rubi [A] (verified) . . . . . 1527  
 3.264.4 Maple [A] (verified) . . . . . 1528  
 3.264.5 Fricas [B] (verification not implemented) . . . . . 1528  
 3.264.6 Sympy [F(-1)] . . . . . 1528  
 3.264.7 Maxima [F] . . . . . 1529  
 3.264.8 Giac [F] . . . . . 1529  
 3.264.9 Mupad [B] (verification not implemented) . . . . . 1529

**3.264.1 Optimal result**

Integrand size = 25, antiderivative size = 33

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx = -\frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{5/2}}$$

output `-2/5*c*d*(d*csc(b*x+a))^(5/2)/b/(c*sec(b*x+a))^(5/2)`

**3.264.2 Mathematica [A] (verified)**

Time = 1.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.36

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx = -\frac{2d^3 \cot^2(a + bx) \sqrt{d \csc(a + bx)}}{5bc \sqrt{c \sec(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(7/2)/(c*Sec[a + b*x])^(3/2),x]`

output `(-2*d^3*Cot[a + b*x]^2*Sqrt[d*Csc[a + b*x]])/(5*b*c*Sqrt[c*Sec[a + b*x]])`

**3.264.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx$$

↓ 3099

$$-\frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{5/2}}$$

input `Int[(d*Csc[a + b*x])^(7/2)/(c*Sec[a + b*x])^(3/2),x]`

output `(-2*c*d*(d*Csc[a + b*x])^(5/2))/(5*b*(c*Sec[a + b*x])^(5/2))`

**3.264.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`

**3.264.4 Maple [A] (verified)**

Time = 1.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.21

method	result	size
default	$-\frac{2\sqrt{d\csc(bx+a)}d^3\cot(bx+a)^2}{5bc\sqrt{c\sec(bx+a)}}$	40

input `int((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/5/b*(d*csc(b*x+a))^(1/2)*d^3/c/(c*sec(b*x+a))^(1/2)*cot(b*x+a)^2`

**3.264.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 59 vs.  $2(27) = 54$ .

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.79

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{2 d^3 \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx + a)^3}{5 (bc^2 \cos(bx + a)^2 - bc^2)}$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fracas")`

output `2/5*d^3*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)^3/(b*c^2*cos(b*x + a)^2 - b*c^2)`

**3.264.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(3/2),x)`

output `Timed out`

**3.264.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{7/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(3/2), x)`

**3.264.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{7/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(3/2), x)`

**3.264.9 Mupad [B] (verification not implemented)**

Time = 13.74 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.12

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{2 d^3 (\cos(4a + 4bx) - 1) \sqrt{\frac{d}{\sin(a+bx)}}}{5 b c \sqrt{\frac{c}{\cos(a+bx)}} (\cos(4a + 4bx) - 4 \cos(2a + 2bx) + 3)}$$

input `int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(3/2),x)`

output `(2*d^3*(cos(4*a + 4*b*x) - 1)*(d/sin(a + b*x))^(1/2))/(5*b*c*(c/cos(a + b*x))^(1/2)*(cos(4*a + 4*b*x) - 4*cos(2*a + 2*b*x) + 3))`

**3.265**  $\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{3/2}} dx$

3.265.1 Optimal result . . . . . 1530  
 3.265.2 Mathematica [C] (verified) . . . . . 1530  
 3.265.3 Rubi [A] (verified) . . . . . 1531  
 3.265.4 Maple [A] (verified) . . . . . 1533  
 3.265.5 Fricas [C] (verification not implemented) . . . . . 1533  
 3.265.6 Sympy [F(-1)] . . . . . 1534  
 3.265.7 Maxima [F] . . . . . 1534  
 3.265.8 Giac [F] . . . . . 1534  
 3.265.9 Mupad [F(-1)] . . . . . 1535

**3.265.1 Optimal result**

Integrand size = 25, antiderivative size = 98

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx = -\frac{2d(d \csc(a + bx))^{3/2}}{3bc\sqrt{c \sec(a + bx)}} - \frac{d^2 \sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{3bc^2}$$

output `-2/3*d*(d*csc(b*x+a))^(3/2)/b/c/(c*sec(b*x+a))^(1/2)+1/3*d^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b/c^2`

**3.265.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.50 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.07

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{d \cos(2(a + bx))(d \csc(a + bx))^{3/2} \left(2 \cot^2(a + bx) - (-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a + bx)\right)\right)}{3b(-2 + \csc^2(a + bx))(c \sec(a + bx))^{3/2}}$$

input `Integrate[(d*Csc[a + b*x])^(5/2)/(c*Sec[a + b*x])^(3/2),x]`

3.265.  $\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{3/2}} dx$

output 
$$-1/3*(d*\text{Cos}[2*(a + b*x)]*(d*\text{Csc}[a + b*x])^{(3/2)}*(2*\text{Cot}[a + b*x]^2 - (-\text{Cot}[a + b*x]^2)^{(3/4)}*\text{Hypergeometric2F1}[1/2, 3/4, 3/2, \text{Csc}[a + b*x]^2])*\text{Sec}[a + b*x]^3)/(b*(-2 + \text{Csc}[a + b*x]^2)*(c*\text{Sec}[a + b*x])^{(3/2)})$$

### 3.265.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3103, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx \\ & \quad \downarrow \text{3103} \\ & -\frac{d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{3c^2} - \frac{2d(d \csc(a + bx))^{3/2}}{3bc \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & -\frac{d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{3c^2} - \frac{2d(d \csc(a + bx))^{3/2}}{3bc \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3110} \\ & -\frac{d^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx}{3c^2} - \frac{2d(d \csc(a + bx))^{3/2}}{3bc \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3042} \\ & -\frac{d^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)} \int \frac{1}{\sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)}} dx}{3c^2} - \frac{2d(d \csc(a + bx))^{3/2}}{3bc \sqrt{c \sec(a + bx)}} \\ & \quad \downarrow \text{3053} \end{aligned}$$

---

3.265.  $\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx$



$$\begin{aligned}
& -\frac{d^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx}{3c^2} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{3042} \\
& -\frac{d^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx}{3c^2} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{3120} \\
& -\frac{d^2 \sqrt{\sin(2a+2bx)} \operatorname{EllipticF}\left(a+bx-\frac{\pi}{4}, 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3bc^2} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc \sqrt{c \sec(a+bx)}}
\end{aligned}$$

input `Int[(d*Csc[a + b*x])^(5/2)/(c*Sec[a + b*x])^(3/2),x]`

output `(-2*d*(d*Csc[a + b*x])^(3/2))/(3*b*c*Sqrt[c*Sec[a + b*x]]) - (d^2*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(3*b*c^2)`

### 3.265.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3103 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m-1)*((b*Sec[e + f*x])^(n+1)/(f*b*(m-1))), x] + Simp[a^2*((n+1)/(b^2*(m-1))) Int[(a*Csc[e + f*x])^(m-2)*(b*Sec[e + f*x])^(n+2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]`

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.265.4 Maple [A] (verified)

Time = 1.16 (sec) , antiderivative size = 208, normalized size of antiderivative = 2.12

method	result
default	$-\frac{\sqrt{2} d^2 \sqrt{d \csc(bx+a)} \left( \sqrt{1+\csc(bx+a)-\cot(bx+a)} \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)} \right) \text{EllipticF}\left(\sqrt{1+\csc(bx+a)-\cot(bx+a)}, 2\right)}{2d^2}$

```
input int((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -1/3/b*2^(1/2)*d^2*(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/c*((1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+ (1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*sec(b*x+a)+2^(1/2)*csc(b*x+a))
```

### 3.265.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.24

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx = \frac{i \sqrt{-4i c d d^2} F(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1) \sin(bx + a) - i \sqrt{4i c d d^2}}{2d^2}$$

```
input integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fracas")
```

---

3.265.  $\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx$

output  $1/6*(I*\sqrt{-4*I*c*d}*d^2*\text{elliptic\_f}(\arcsin(\cos(b*x + a) + I*\sin(b*x + a)), -1)*\sin(b*x + a) - I*\sqrt{4*I*c*d}*d^2*\text{elliptic\_f}(\arcsin(\cos(b*x + a) - I*\sin(b*x + a)), -1)*\sin(b*x + a) - 4*d^2*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)}*\cos(b*x + a))/(b*c^2*\sin(b*x + a))$

### 3.265.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(3/2), x)`

output Timed out

### 3.265.7 Maxima [F]

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{5/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2), x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(3/2), x)`

### 3.265.8 Giac [F]

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{5/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2), x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(3/2), x)`

---

3.265.  $\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{3/2}} dx$

**3.265.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

input `int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(3/2),x)`output `int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(3/2), x)`

**3.266** 
$$\int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{3/2}} dx$$

3.266.1 Optimal result . . . . . 1536  
 3.266.2 Mathematica [A] (verified) . . . . . 1537  
 3.266.3 Rubi [A] (verified) . . . . . 1537  
 3.266.4 Maple [B] (warning: unable to verify) . . . . . 1542  
 3.266.5 Fracas [C] (verification not implemented) . . . . . 1542  
 3.266.6 Sympy [F] . . . . . 1543  
 3.266.7 Maxima [F] . . . . . 1544  
 3.266.8 Giac [F] . . . . . 1544  
 3.266.9 Mupad [F(-1)] . . . . . 1544

**3.266.1 Optimal result**

Integrand size = 25, antiderivative size = 327

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx = -\frac{2d\sqrt{d \csc(a + bx)}}{bc\sqrt{c \sec(a + bx)}} + \frac{d^2 \arctan\left(1 - \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2}bc^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} - \frac{d^2 \arctan\left(1 + \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2}bc^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} - \frac{d^2 \log\left(1 - \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c \sec(a + bx)}}{2\sqrt{2}bc^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} + \frac{d^2 \log\left(1 + \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c \sec(a + bx)}}{2\sqrt{2}bc^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}$$

output

```
-2*d*(d*csc(b*x+a))^(1/2)/b/c/(c*sec(b*x+a))^(1/2)-1/2*d^2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)-1/2*d^2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)-1/4*d^2*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+1/4*d^2*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)
```

**3.266.2 Mathematica [A] (verified)**

Time = 1.48 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.49

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx =$$

$$d \left( 4 \cot^2(a + bx) - \sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2}^4 \sqrt{\cot^2(a + bx)}} \right) \cot^2(a + bx)^{3/4} - \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{2}^4 \sqrt{\cot^2(a + bx)}}{1 + \sqrt{\cot^2(a + bx)}} \right) \right) \\ \hline 2bc \sqrt{c \sec(a + bx)}$$

input `Integrate[(d*Csc[a + b*x])^(3/2)/(c*Sec[a + b*x])^(3/2),x]`output `-1/2*(d*(4*Cot[a + b*x]^2 - Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2)]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(3/4) - Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2)]]*(Cot[a + b*x]^2)^(3/4))*Sqrt[d*Csc[a + b*x]]*Tan[a + b*x]^2)/(b*c*Sqrt[c*Sec[a + b*x]])`**3.266.3 Rubi [A] (verified)**Time = 0.62 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.65, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3103, 3042, 3109, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx \\ \downarrow 3042 \\ \int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx \\ \downarrow 3103 \\ \frac{d^2 \int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx}{c^2} - \frac{2d \sqrt{d \csc(a + bx)}}{bc \sqrt{c \sec(a + bx)}} \\ \downarrow 3042$$

$$\begin{aligned}
& \frac{d^2 \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{c^2} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{3109} \\
& \frac{d^2 \sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{c^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{d^2 \sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{c^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{3957} \\
& \frac{d^2 \sqrt{c \sec(a+bx)} \int \frac{\sqrt{\tan(a+bx)}}{\tan^2(a+bx)+1} d \tan(a+bx)}{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{266} \\
& \frac{2d^2 \sqrt{c \sec(a+bx)} \int \frac{\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{826} \\
& \frac{2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{1476} \\
& \frac{2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right)}{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}} \\
& \quad \downarrow \text{1082} \\
& \frac{2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{1}{-\tan(a+bx)-1} \frac{d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \int \frac{1}{-\tan(a+bx)-1} \frac{d(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2d \sqrt{d \csc(a+bx)}}{bc \sqrt{c \sec(a+bx)}}
\end{aligned}$$

---

3.266.  $\int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{3/2}} dx$

↓ 217

$$2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d\sqrt{\tan(a+bx)} \right)$$


---


$$\frac{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{2d \sqrt{d \csc(a+bx)}} \\ \frac{bc \sqrt{c \sec(a+bx)}}{bc \sqrt{c \sec(a+bx)}}$$

↓ 1479

$$2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan \left( \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} \right) - \arctan \left( \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} \right) \right) \right)$$


---


$$\frac{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{2d \sqrt{d \csc(a+bx)}} \\ \frac{bc \sqrt{c \sec(a+bx)}}{bc \sqrt{c \sec(a+bx)}}$$

↓ 25

$$2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( - \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan \left( \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} \right) - \arctan \left( \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} \right) \right) \right)$$


---


$$\frac{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{2d \sqrt{d \csc(a+bx)}} \\ \frac{bc \sqrt{c \sec(a+bx)}}{bc \sqrt{c \sec(a+bx)}}$$

↓ 27

$$2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( - \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan \left( \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} \right) - \arctan \left( \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} \right) \right) \right)$$


---


$$\frac{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{2d \sqrt{d \csc(a+bx)}} \\ \frac{bc \sqrt{c \sec(a+bx)}}{bc \sqrt{c \sec(a+bx)}}$$

↓ 1103

$$2d^2 \sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} - \frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} \right) \right)$$


---


$$\frac{bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{2d \sqrt{d \csc(a+bx)}} \\ \frac{bc \sqrt{c \sec(a+bx)}}{bc \sqrt{c \sec(a+bx)}}$$



input `Int[(d*Csc[a + b*x])^(3/2)/(c*Sec[a + b*x])^(3/2),x]`

output `(-2*d*Sqrt[d*Csc[a + b*x]]/(b*c*Sqrt[c*Sec[a + b*x]]) - (2*d^2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2])))/2)*Sqrt[c*Sec[a + b*x]])/(b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])`

### 3.266.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3103 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Simp[a^2*((n + 1)/(b^2*(m - 1))) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.266.4 Maple [B] (warning: unable to verify)**

Leaf count of result is larger than twice the leaf count of optimal. 643 vs.  $2(271) = 542$ .

Time = 13.14 (sec) , antiderivative size = 644, normalized size of antiderivative = 1.97

method	result
default	$\sqrt{2} \left( -\ln \left( \frac{(1-\cos(bx+a))^2 \csc(bx+a) + 2\sqrt{(1-\cos(bx+a))((1-\cos(bx+a))^2 \csc(bx+a)^2 - 1)} \csc(bx+a) \sin(bx+a) - 2\cos(bx+a) + 2 - \sin(bx+a)}}{1-\cos(bx+a)} \right) \right)$

input `int((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/4/b*2^{(1/2)}*(-\ln(1/(1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)+2*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)-2*\cos(b*x+a)+2-\sin(b*x+a)))*(-\cot(b*x+a)+\csc(b*x+a))-2*\arctan(1/(1-\cos(b*x+a)))*(((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)-\cos(b*x+a)+1))*(-\cot(b*x+a)+\csc(b*x+a))+\ln(-1/(1-\cos(b*x+a)))*(-(1-\cos(b*x+a))^2*\csc(b*x+a)+2*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)+2*\cos(b*x+a)-2+\sin(b*x+a)))*(-\cot(b*x+a)+\csc(b*x+a))-2*\arctan(1/(1-\cos(b*x+a)))*(((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)+\cos(b*x+a)-1))*(-\cot(b*x+a)+\csc(b*x+a))+4*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*(d/(1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)+\sin(b*x+a)))^{(3/2)}/((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}/((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)/(-c*((1-\cos(b*x+a))^2*\csc(b*x+a)^2+1)/((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1))^{(3/2)}*(-\cot(b*x+a)+\csc(b*x+a)) \end{aligned}$$
**3.266.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 1286, normalized size of antiderivative = 3.93

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

```

output 1/8*(b*c^2*(-d^6/(b^4*c^6))^(1/4)*log(2*b^2*c^3*d^2*sqrt(-d^6/(b^4*c^6))*c
os(b*x + a)*sin(b*x + a) + 2*d^5*cos(b*x + a)^2 - d^5 + 2*(b*c*d^3*(-d^6/(
b^4*c^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*c^4*cos(b*x + a)^3 - b^
3*c^4*cos(b*x + a))*(-d^6/(b^4*c^6))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/si
n(b*x + a))) - b*c^2*(-d^6/(b^4*c^6))^(1/4)*log(2*b^2*c^3*d^2*sqrt(-d^6/(b
^4*c^6))*cos(b*x + a)*sin(b*x + a) + 2*d^5*cos(b*x + a)^2 - d^5 - 2*(b*c*d
^3*(-d^6/(b^4*c^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*c^4*cos(b*x +
a)^3 - b^3*c^4*cos(b*x + a))*(-d^6/(b^4*c^6))^(3/4))*sqrt(c/cos(b*x + a))
*sqrt(d/sin(b*x + a))) - I*b*c^2*(-d^6/(b^4*c^6))^(1/4)*log(-2*b^2*c^3*d^2
*sqrt(-d^6/(b^4*c^6))*cos(b*x + a)*sin(b*x + a) + 2*d^5*cos(b*x + a)^2 - d
^5 - 2*(I*b*c*d^3*(-d^6/(b^4*c^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (I*
b^3*c^4*cos(b*x + a)^3 - I*b^3*c^4*cos(b*x + a))*(-d^6/(b^4*c^6))^(3/4))*s
qrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) + I*b*c^2*(-d^6/(b^4*c^6))^(1/4)
*log(-2*b^2*c^3*d^2*sqrt(-d^6/(b^4*c^6))*cos(b*x + a)*sin(b*x + a) + 2*d^5
*cos(b*x + a)^2 - d^5 - 2*(-I*b*c*d^3*(-d^6/(b^4*c^6))^(1/4)*cos(b*x + a)^
2*sin(b*x + a) + (-I*b^3*c^4*cos(b*x + a)^3 + I*b^3*c^4*cos(b*x + a))*(-d^
6/(b^4*c^6))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) - b*c^2*(-d
^6/(b^4*c^6))^(1/4)*log(-d^5 + 2*(b*c*d^3*(-d^6/(b^4*c^6))^(1/4)*cos(b*x +
a)^2*sin(b*x + a) + (b^3*c^4*cos(b*x + a)^3 - b^3*c^4*cos(b*x + a))*(-d^6
/(b^4*c^6))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))) + b*c^2*...

```

### 3.266.6 Sympy [F]

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx$$

```
input integrate((d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(3/2), x)
```

```
output Integral((d*csc(a + b*x))**(3/2)/(c*sec(a + b*x))**(3/2), x)
```

**3.266.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{3/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(3/2)/(c*sec(b*x + a))^(3/2), x)`

**3.266.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{(d \csc(bx + a))^{3/2}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(3/2)/(c*sec(b*x + a))^(3/2), x)`

**3.266.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

input `int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(3/2),x)`

output `int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(3/2), x)`

**3.267**  $\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx$

3.267.1 Optimal result . . . . . 1545  
 3.267.2 Mathematica [C] (verified) . . . . . 1545  
 3.267.3 Rubi [A] (verified) . . . . . 1546  
 3.267.4 Maple [A] (verified) . . . . . 1548  
 3.267.5 Fracas [F] . . . . . 1548  
 3.267.6 Sympy [F] . . . . . 1549  
 3.267.7 Maxima [F] . . . . . 1549  
 3.267.8 Giac [F] . . . . . 1549  
 3.267.9 Mupad [F(-1)] . . . . . 1550

**3.267.1 Optimal result**

Integrand size = 25, antiderivative size = 92

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx = \frac{d}{bc \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{\sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{2bc^2}$$

output `d/b/c/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)-1/2*(sin(a+1/4*Pi+b*x))^2^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b/c^2`

**3.267.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.12 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx = \frac{d(1 + \cos(2(a + bx)) - (-\cot^2(a + bx))^{3/4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a + bx)\right))}{2b \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}$$

input `Integrate[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(3/2),x]`

output  $(d*(1 + \text{Cos}[2*(a + b*x)] - (-\text{Cot}[a + b*x]^2)^{(3/4)}*\text{Hypergeometric2F1}[1/2, 3/4, 3/2, \text{Csc}[a + b*x]^2])*\text{Sec}[a + b*x]^3/(2*b*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(3/2)})$

### 3.267.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3108, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx \\
 & \quad \downarrow \text{3108} \\
 & \frac{\int \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3110} \\
 & \frac{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx}{2c^2} + \\
 & \quad \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx}{2c^2} + \\
 & \quad \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} \\
 & \quad \downarrow \text{3053}
 \end{aligned}$$

---

3.267.  $\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx$

$$\frac{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx}{2c^2} + \frac{d}{bc\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}$$

↓ 3042

$$\frac{\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\sin(2a+2bx)}}dx}{2c^2} + \frac{d}{bc\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}$$

↓ 3120

$$\frac{\sqrt{\sin(2a+2bx)}\operatorname{EllipticF}\left(a+bx-\frac{\pi}{4}, 2\right)\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{2bc^2} + \frac{d}{bc\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}$$

input `Int[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(3/2),x]`

output `d/(b*c*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]) + (Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(2*b*c^2)`

### 3.267.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3108 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`



```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.267.4 Maple [A] (verified)

Time = 8.12 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.23

method	result
default	$\frac{\sqrt{2} \sqrt{d \csc(bx+a)} \left( \sqrt{1+\csc(bx+a)-\cot(bx+a)} \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)} \right) \text{EllipticF}\left(\sqrt{1+\csc(bx+a)-\cot(bx+a)}, 2\right)}{2d}$

```
input int((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/2/b*2^(1/2)*(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/c*((1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*sec(b*x+a)+2^(1/2)*sin(b*x+a))
```

### 3.267.5 Fricas [F]

$$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx = \int \frac{\sqrt{d \csc(bx+a)}}{(c \sec(bx+a))^{3/2}} dx$$

```
input integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")
```

```
output integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^2*sec(b*x + a)^2), x)
```

---

3.267.  $\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx$

**3.267.6 Sympy [F]**

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(3/2),x)`

output `Integral(sqrt(d*csc(a + b*x))/(c*sec(a + b*x))**(3/2), x)`

**3.267.7 Maxima [F]**

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{\sqrt{d \csc(bx + a)}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(3/2), x)`

**3.267.8 Giac [F]**

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{\sqrt{d \csc(bx + a)}}{(c \sec(bx + a))^{3/2}} dx$$

input `integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(3/2), x)`

**3.267.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx = \int \frac{\sqrt{\frac{d}{\sin(a+bx)}}}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

input `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(3/2),x)`output `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(3/2), x)`

**3.268**  $\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx$

3.268.1 Optimal result . . . . . 1551  
 3.268.2 Mathematica [A] (verified) . . . . . 1552  
 3.268.3 Rubi [A] (verified) . . . . . 1552  
 3.268.4 Maple [A] (warning: unable to verify) . . . . . 1557  
 3.268.5 Fricas [C] (verification not implemented) . . . . . 1557  
 3.268.6 Sympy [F] . . . . . 1558  
 3.268.7 Maxima [F] . . . . . 1559  
 3.268.8 Giac [F] . . . . . 1559  
 3.268.9 Mupad [F(-1)] . . . . . 1559

**3.268.1 Optimal result**

Integrand size = 25, antiderivative size = 322

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx = \frac{d}{2bc(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} - \frac{\arctan\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2}bc^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} + \frac{\arctan\left(1 + \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2}bc^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} + \frac{\log\left(1 - \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{c \sec(a+bx)}}{8\sqrt{2}bc^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} + \frac{\log\left(1 + \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{c \sec(a+bx)}}{8\sqrt{2}bc^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}$$

output

```
1/2*d/b/c/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2)+1/8*arctan(-1+2^(1/2)*
tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/
tan(b*x+a)^(1/2)+1/8*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/
2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+1/16*ln(1-2^(1/2)*t
an(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+
a))^(1/2)/tan(b*x+a)^(1/2)-1/16*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*
(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)
```

**3.268.2 Mathematica [A] (verified)**

Time = 2.20 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.49

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx = \frac{d \left( 4 \cos^2(a+bx) - \sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a+bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a+bx)}} \right) \cot^2(a+bx) \right)}{8b(d \csc(a+bx))}$$

input `Integrate[1/(Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)),x]`output `(d*(4*Cos[a + b*x]^2 - Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2)]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))])*(Cot[a + b*x]^2)^(3/4) - Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(3/4))*Sec[a + b*x]^3)/(8*b*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2))`**3.268.3 Rubi [A] (verified)**Time = 0.58 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.66, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3108, 3042, 3109, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} dx \\ & \quad \downarrow \text{3108} \\ & \frac{\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{4c^2} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{4c^2} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3109} \\
& \frac{\sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{4c^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \downarrow \text{3042} \\
& \frac{\sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{4c^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \downarrow \text{3957} \\
& \frac{\sqrt{c \sec(a+bx)} \int \frac{\sqrt{\tan(a+bx)}}{\tan^2(a+bx)+1} d \tan(a+bx)}{4bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \downarrow \text{266} \\
& \frac{\sqrt{c \sec(a+bx)} \int \frac{\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \downarrow \text{826} \\
& \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \\
& \quad \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \downarrow \text{1476} \\
& \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \\
& \quad \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \downarrow \text{1082} \\
& \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\int \frac{1}{-\tan(a+bx)-1} d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(a+bx)-1} d(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \\
& \quad \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \\
& \downarrow \text{217}
\end{aligned}$$

$$\frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d\sqrt{\tan(a+bx)} \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} +$$

$$\frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}}$$

↓ 1479

$$\frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} + \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) - \arctan(1-\sqrt{2}\sqrt{\tan(a+bx)}) \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} +$$

$$\frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}}$$

↓ 25

$$\frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) - \arctan(1-\sqrt{2}\sqrt{\tan(a+bx)}) \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} +$$

$$\frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}}$$

↓ 27

$$\frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{\tan(a+bx)+1}) - \arctan(1-\sqrt{2}\sqrt{\tan(a+bx)}) \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} +$$

$$\frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}}$$

↓ 1103

$$\frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} - \frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} +$$

$$\frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}}$$

input `Int[1/(Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)),x]`

3.268.  $\int \frac{1}{\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} dx$

```
output d/(2*b*c*(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]]) + (((-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[c*Sec[a + b*x]])/(2*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])
```

### 3.268.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 266 Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 826 Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```



rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3108 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.268.4 Maple [A] (warning: unable to verify)**

Time = 48.78 (sec) , antiderivative size = 438, normalized size of antiderivative = 1.36

method	result
default	$\sqrt{2} \left( 4\sqrt{2} \cos(bx+a) \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sin(bx+a) + 4 \sin(bx+a) \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} + \ln \left( -2\sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \right) \right)$

input `int(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `1/16/b*2^(1/2)*(4*2^(1/2)*cos(b*x+a)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*sin(b*x+a)+4*sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+ln(-2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)-2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2)-ln(2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)+2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2)-2*arctan((-sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+cos(b*x+a)-1)/(cos(b*x+a)-1))+2*arctan((sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+cos(b*x+a)-1)/(cos(b*x+a)-1)))/(cos(b*x+a)+1)/(c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2)/(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)/c`

**3.268.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.65 (sec) , antiderivative size = 1247, normalized size of antiderivative = 3.87

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fracas")`

output

```

-1/32*(I*b*c^2*d*(-1/(b^4*c^6*d^2))^(1/4)*log(2*b^2*c^3*d*sqrt(-1/(b^4*c^6*d^2)))*cos(b*x + a)*sin(b*x + a) - 2*(I*b*c*(-1/(b^4*c^6*d^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (I*b^3*c^4*d*cos(b*x + a)^3 - I*b^3*c^4*d*cos(b*x + a))*(-1/(b^4*c^6*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) - I*b*c^2*d*(-1/(b^4*c^6*d^2))^(1/4)*log(2*b^2*c^3*d*sqrt(-1/(b^4*c^6*d^2)))*cos(b*x + a)*sin(b*x + a) - 2*(-I*b*c*(-1/(b^4*c^6*d^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (-I*b^3*c^4*d*cos(b*x + a)^3 + I*b^3*c^4*d*cos(b*x + a))*(-1/(b^4*c^6*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) - b*c^2*d*(-1/(b^4*c^6*d^2))^(1/4)*log(-2*b^2*c^3*d*sqrt(-1/(b^4*c^6*d^2)))*cos(b*x + a)*sin(b*x + a) + 2*(b*c*(-1/(b^4*c^6*d^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*c^4*d*cos(b*x + a)^3 - b^3*c^4*d*cos(b*x + a))*(-1/(b^4*c^6*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) + b*c^2*d*(-1/(b^4*c^6*d^2))^(1/4)*log(-2*b^2*c^3*d*sqrt(-1/(b^4*c^6*d^2)))*cos(b*x + a)*sin(b*x + a) - 2*(b*c*(-1/(b^4*c^6*d^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*c^4*d*cos(b*x + a)^3 - b^3*c^4*d*cos(b*x + a))*(-1/(b^4*c^6*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) - b*c^2*d*(-1/(b^4*c^6*d^2))^(1/4)*log(2*(b*c*(-1/(b^4*c^6*d^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*c^4*d*cos(b*x + a)^3 - b^3*c^4*d*cos(b*x + a))*(-1/(b^4*c^6*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) - b*c^2*d*(-1/(b^4*c^6*d^2))^(1/4)*log(2*(b*c*(-1/(b^4*c^6*d^2))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c^4*d*cos(b*x + a)^3 - b^3*c^4*d*cos(b*x + a))*(-1/(b^4*c^6*d^2))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))...

```

### 3.268.6 Sympy [F]

$$\int \frac{1}{\sqrt{d \csc(a + bx)}(c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(c \sec(a + bx))^{\frac{3}{2}} \sqrt{d \csc(a + bx)}} dx$$

input `integrate(1/(d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(3/2), x)`

output `Integral(1/((c*sec(a + b*x))**(3/2)*sqrt(d*csc(a + b*x))), x)`

**3.268.7 Maxima [F]**

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx = \int \frac{1}{\sqrt{d \csc(bx+a)}(c \sec(bx+a))^{3/2}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2)), x)`

**3.268.8 Giac [F]**

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx = \int \frac{1}{\sqrt{d \csc(bx+a)}(c \sec(bx+a))^{3/2}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2)), x)`

**3.268.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2} \sqrt{\frac{d}{\sin(a+bx)}} dx$$

input `int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(1/2)),x)`

output `int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(1/2)), x)`

**3.269**  $\int \frac{1}{(d \csc(a+bx))^{3/2}(c \sec(a+bx))^{3/2}} dx$

3.269.1 Optimal result . . . . . 1560  
 3.269.2 Mathematica [C] (verified) . . . . . 1560  
 3.269.3 Rubi [A] (verified) . . . . . 1561  
 3.269.4 Maple [C] (warning: unable to verify) . . . . . 1564  
 3.269.5 Fricas [F] . . . . . 1564  
 3.269.6 Sympy [F(-1)] . . . . . 1565  
 3.269.7 Maxima [F] . . . . . 1565  
 3.269.8 Giac [F] . . . . . 1565  
 3.269.9 Mupad [F(-1)] . . . . . 1566

**3.269.1 Optimal result**

Integrand size = 25, antiderivative size = 135

$$\int \frac{1}{(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}} dx =$$

$$-\frac{c}{3bd\sqrt{d \csc(a + bx)}(c \sec(a + bx))^{5/2}} + \frac{1}{6bcd\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}}$$

$$+ \frac{\sqrt{d \csc(a + bx)} \operatorname{EllipticF}\left(a - \frac{\pi}{4} + bx, 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{12bc^2d^2}$$

output

```
-1/3*c/b/d/(c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2)+1/6/b/c/d/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)-1/12*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b/c^2/d^2
```

**3.269.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.22 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.66

$$\int \frac{1}{(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}} dx = \frac{-2 \cos(2(a + bx)) + \frac{\csc^2(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \csc^2(a+bx)\right)}{\sqrt{-\cot^2(a + bx)}}}{12bcd\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}}$$

input `Integrate[1/((d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2)),x]`

output `(-2*Cos[2*(a + b*x)] + (Csc[a + b*x]^2*Hypergeometric2F1[1/2, 3/4, 3/2, Cs  
c[a + b*x]^2])/(-Cot[a + b*x]^2)^(1/4))/(12*b*c*d*sqrt[d*Csc[a + b*x]]*Sqr  
t[c*Sec[a + b*x]])`

### 3.269.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3107, 3042, 3108, 3042, 3110, 3042, 3053, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3107} \\
 & \frac{\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx}{6d^2} - \frac{c}{3bd(c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx}{6d^2} - \frac{c}{3bd(c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)}} \\
 & \quad \downarrow \text{3108} \\
 & \frac{\int \frac{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}}{6d^2} - \frac{c}{3bd(c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}}{6d^2} - \frac{c}{3bd(c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)}} \\
 & \quad \downarrow \text{3110}
 \end{aligned}$$

---

3.269.  $\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} dx$

$$\frac{\frac{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}}{\frac{6d^2}{c}} = \frac{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}}{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}} \downarrow 3042$$

$$\frac{\frac{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}}{\frac{6d^2}{c}} = \frac{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}}{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}} \downarrow 3053$$

$$\frac{\frac{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}}{\frac{6d^2}{c}} = \frac{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}}{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}} \downarrow 3042$$

$$\frac{\frac{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx}{2c^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}}{\frac{6d^2}{c}} = \frac{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}}{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}} \downarrow 3120$$

$$\frac{\frac{\sqrt{\sin(2a+2bx)} \operatorname{EllipticF}(a+bx-\frac{\pi}{4}, 2) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{2bc^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}}{\frac{6d^2}{c}} = \frac{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}}{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}}$$

input `Int[1/((d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2)),x]`

output `-1/3*c/(b*d*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(5/2)) + (d/(b*c*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])) + (Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(2*b*c^2)/(6*d^2)`

## 3.269.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3053 `Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]) Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3107 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3108 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`



### 3.269.4 Maple [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 58.14 (sec) , antiderivative size = 1762, normalized size of antiderivative = 13.05

method	result	size
default	Expression too large to display	1762

```
input int(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/48/b^2^(1/2)*(-6*I*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)
)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticPi((1+csc(b*x+a)-cot(b*x+
a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-6*I*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(
b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticPi((1+csc
(b*x+a)-cot(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*cos(b*x+a)+6*I*(1+csc(b*x
+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+
a))^(1/2)*EllipticPi((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2)
)-6*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b
*x+a)-csc(b*x+a))^(1/2)*EllipticPi((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2+1/2
*I,1/2*2^(1/2))*cos(b*x+a)+6*I*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)
-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticPi((1+csc(b*x+a)
)-cot(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*cos(b*x+a)-6*(1+csc(b*x+a)-cot(
b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2
)*EllipticPi((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*cos(b*
x+a)+8*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(co
t(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2
^(1/2))*cos(b*x+a)+8*2^(1/2)*cos(b*x+a)^3*sin(b*x+a)-6*(1+csc(b*x+a)-cot(b
*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)
*EllipticPi((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))-6*(1+cs
c(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-...
```

### 3.269.5 Fricas [F]

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(d \csc(bx + a))^{3/2} (c \sec(bx + a))^{3/2}} dx$$

```
input integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas
")
```

output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^2*d^2*csc(b*x + a)^2*sec(b*x + a)^2), x)`

### 3.269.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate(1/(d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(3/2), x)`

output `Timed out`

### 3.269.7 Maxima [F]

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2), x, algorithm="maxima")`

output `integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2)), x)`

### 3.269.8 Giac [F]

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2), x, algorithm="giac")`

output `integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2)), x)`

**3.269.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2} \left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

input `int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2)),x)`output `int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2)), x)`

**3.270**      $\int \frac{1}{(d \csc(a+bx))^{5/2}(c \sec(a+bx))^{3/2}} dx$

3.270.1 Optimal result . . . . . 1567  
 3.270.2 Mathematica [A] (verified) . . . . . 1568  
 3.270.3 Rubi [A] (verified) . . . . . 1568  
 3.270.4 Maple [A] (warning: unable to verify) . . . . . 1574  
 3.270.5 Fricas [C] (verification not implemented) . . . . . 1575  
 3.270.6 Sympy [F(-1)] . . . . . 1576  
 3.270.7 Maxima [F] . . . . . 1577  
 3.270.8 Giac [F] . . . . . 1577  
 3.270.9 Mupad [F(-1)] . . . . . 1577

**3.270.1 Optimal result**

Integrand size = 25, antiderivative size = 371

$$\int \frac{1}{(d \csc(a + bx))^{5/2}(c \sec(a + bx))^{3/2}} dx =$$

$$-\frac{c}{4bd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2}\sqrt{c \sec(a + bx)}}$$

$$-\frac{3 \arctan\left(1 - \sqrt{2}\sqrt{\tan(a + bx)}\right)\sqrt{c \sec(a + bx)}}{32\sqrt{2}bc^2d^2\sqrt{d \csc(a + bx)}\sqrt{\tan(a + bx)}}$$

$$+\frac{3 \arctan\left(1 + \sqrt{2}\sqrt{\tan(a + bx)}\right)\sqrt{c \sec(a + bx)}}{32\sqrt{2}bc^2d^2\sqrt{d \csc(a + bx)}\sqrt{\tan(a + bx)}}$$

$$+\frac{3 \log\left(1 - \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right)\sqrt{c \sec(a + bx)}}{64\sqrt{2}bc^2d^2\sqrt{d \csc(a + bx)}\sqrt{\tan(a + bx)}}$$

$$-\frac{3 \log\left(1 + \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right)\sqrt{c \sec(a + bx)}}{64\sqrt{2}bc^2d^2\sqrt{d \csc(a + bx)}\sqrt{\tan(a + bx)}}$$

output 
$$-1/4*c/b/d/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2)+3/16/b/c/d/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2)+3/64*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/c^2/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+3/64*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/c^2/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+3/128*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/c^2/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)-3/128*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/c^2/d^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)$$

### 3.270.2 Mathematica [A] (verified)

Time = 2.66 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.45

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}} dx = \frac{\left( 2(\cos(2(a + bx)) + \cos(4(a + bx))) + 3\sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}} \right) \cot^2(a + bx)^{3/4} + 3\sqrt{2} \operatorname{arctanh} \left( \frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}} \right) \right)}{64bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}$$

input `Integrate[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(3/2)),x]`

output 
$$-1/64*((2*(Cos[2*(a + b*x)] + Cos[4*(a + b*x)]) + 3*Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2])/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(3/4) + 3*Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(3/4))*Sec[a + b*x]^3)/(b*d*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2))$$

### 3.270.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.69, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.680$ , Rules used = {3042, 3107, 3042, 3108, 3042, 3109, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.270.  $\int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{1}{(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{1}{(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{5/2}} dx \\
& \quad \downarrow \text{3107} \\
& \frac{3 \int \frac{1}{\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} dx}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \int \frac{1}{\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} dx}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3108} \\
& \frac{3 \left( \frac{\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{4c^2} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \left( \frac{\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{4c^2} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3109} \\
& \frac{3 \left( \frac{\sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{4c^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d_c^2} - \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \left( \frac{\sqrt{c \sec(a+bx)} \int \sqrt{\tan(a+bx)} dx}{4c^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d_c^2} - \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3957}
\end{aligned}$$

---

3.270.  $\int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{3 \left( \frac{\sqrt{c \sec(a+bx)} \int \frac{\sqrt{\tan(a+bx)}}{\tan^2(a+bx)+1} d \tan(a+bx)}{4bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d_c^2} \\
 & \quad \frac{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow 266 \\
 & \frac{3 \left( \frac{\sqrt{c \sec(a+bx)} \int \frac{\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d_c^2} \\
 & \quad \frac{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow 826 \\
 & \frac{3 \left( \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d_c^2} \\
 & \quad \frac{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow 1476 \\
 & \frac{3 \left( \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \right)}{8d^2} \\
 & \quad \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow 1082 \\
 & \frac{3 \left( \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\int \frac{1}{-\tan(a+bx)-1} d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(a+bx)-1} d(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))^{3/2}} \right)}{8d^2} \\
 & \quad \frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \\
 & \quad \downarrow 217
 \end{aligned}$$

---

3.270.  $\int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2}} dx$

$$3 \left( \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d\sqrt{\tan(a+bx)} \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \right) + \frac{d}{2bc \sqrt{c \sec(a+bx)} (d \csc(a+bx))}$$


---


$$\frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \quad 8d^2$$

↓ 1479

$$3 \left( \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \right)$$


---


$$\frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \quad 8d^2$$

↓ 25

$$3 \left( \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \right)$$


---


$$\frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \quad 8d^2$$

↓ 27

$$3 \left( \frac{\sqrt{c \sec(a+bx)} \left( \frac{1}{2} \left( -\int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} \right) \right)}{2bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} \right)$$


---


$$\frac{c}{4bd(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} \quad 8d^2$$

↓ 1103

---

3.270.  $\int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2}} dx$



$$3 \left( \frac{\sqrt{c \operatorname{csc}(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(a+bx)}}{\sqrt{2}}\right)}{2bc^2\sqrt{\tan(a+bx)}\sqrt{d \operatorname{csc}(a+bx)}} \right) + \frac{1}{2} \left( \frac{\log\left(\frac{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}{2\sqrt{2}}\right) - \log\left(\frac{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}}{2\sqrt{2}}\right)}{8d^2} \right)}{4bd(c \operatorname{csc}(a+bx))^{5/2}(d \operatorname{csc}(a+bx))^{3/2}} \right)$$

input `Int[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(3/2)),x]`

output `-1/4*c/(b*d*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2)) + (3*(d/(2*b*c*(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]])) + (((-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[c*Sec[a + b*x]]/(2*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])))/(8*d^2)`

### 3.270.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntegerBinomialQ[a, b, c, 2, m, p, x]`

- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3107 `Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

```
rule 3108 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

```
rule 3109 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]
```

```
rule 3957 Int[((b_.)*tan[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

### 3.270.4 Maple [A] (warning: unable to verify)

Time = 50.58 (sec) , antiderivative size = 551, normalized size of antiderivative = 1.49

method	result
default	$\frac{\sqrt{2} \left( 16\sqrt{2} \cos(bx+a)^3 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sin(bx+a) + 16\sqrt{2} \cos(bx+a)^2 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sin(bx+a) - 12\sqrt{2} \cos(bx+a) \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sin(bx+a) \right)}{\dots}$

```
input int(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

---

3.270.  $\int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2}} dx$

output  $\frac{1}{128}b^2^{1/2} \cdot (16 \cdot 2^{1/2} \cos(bx+a)^3 (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} \sin(bx+a) + 16 \cdot 2^{1/2} \cos(bx+a)^2 (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} \sin(bx+a) - 12 \cdot 2^{1/2} \cos(bx+a) (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} \sin(bx+a) - 12 \sin(bx+a) \cdot 2^{1/2} (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} + 3 \ln(2 \cdot 2^{1/2} (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} \cot(bx+a) + 2 \cdot 2^{1/2} (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} \csc(bx+a) - 2 \cot(bx+a) + 2) - 6 \arctan((\sin(bx+a) \cdot 2^{1/2} (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} + \cos(bx+a) - 1) / (\cos(bx+a) - 1)) + 6 \arctan((- \sin(bx+a) \cdot 2^{1/2} (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} + \cos(bx+a) - 1) / (\cos(bx+a) - 1)) - 3 \ln(-2 \cdot 2^{1/2} (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} \cot(bx+a) - 2 \cdot 2^{1/2} (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} \csc(bx+a) - 2 \cot(bx+a) + 2) / (\cos(bx+a) - 1) / (\cos(bx+a) + 1)^2 / (c \sec(bx+a))^{1/2} / (-\cos(bx+a) \sin(bx+a) / (\cos(bx+a)+1)^2)^{1/2} / (d \csc(bx+a))^{1/2} / c / d^2 \sin(bx+a)^2$

### 3.270.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.15 (sec) , antiderivative size = 1341, normalized size of antiderivative = 3.61

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fracas")`

output

```
-1/256*(3*I*b*c^2*d^3*(-1/(b^4*c^6*d^10))^(1/4)*log(2*b^2*c^3*d^5*sqrt(-1/
(b^4*c^6*d^10))*cos(b*x + a)*sin(b*x + a) - 2*(I*b*c*d^2*(-1/(b^4*c^6*d^10
))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (I*b^3*c^4*d^7*cos(b*x + a)^3 - I*b
^3*c^4*d^7*cos(b*x + a))*(-1/(b^4*c^6*d^10))^(3/4))*sqrt(c/cos(b*x + a))*s
qrt(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) - 3*I*b*c^2*d^3*(-1/(b^4*c^6*d
^10))^(1/4)*log(2*b^2*c^3*d^5*sqrt(-1/(b^4*c^6*d^10))*cos(b*x + a)*sin(b*x
+ a) - 2*(-I*b*c*d^2*(-1/(b^4*c^6*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a
) + (-I*b^3*c^4*d^7*cos(b*x + a)^3 + I*b^3*c^4*d^7*cos(b*x + a))*(-1/(b^4*
c^6*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 2*cos(b*x +
a)^2 + 1) - 3*b*c^2*d^3*(-1/(b^4*c^6*d^10))^(1/4)*log(-2*b^2*c^3*d^5*sqrt(
-1/(b^4*c^6*d^10))*cos(b*x + a)*sin(b*x + a) + 2*(b*c*d^2*(-1/(b^4*c^6*d^1
0))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^3*c^4*d^7*cos(b*x + a)^3 - b^3*
c^4*d^7*cos(b*x + a))*(-1/(b^4*c^6*d^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt
(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) + 3*b*c^2*d^3*(-1/(b^4*c^6*d^10))
^(1/4)*log(-2*b^2*c^3*d^5*sqrt(-1/(b^4*c^6*d^10))*cos(b*x + a)*sin(b*x + a
) - 2*(b*c*d^2*(-1/(b^4*c^6*d^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (b^
3*c^4*d^7*cos(b*x + a)^3 - b^3*c^4*d^7*cos(b*x + a))*(-1/(b^4*c^6*d^10))^(
3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 2*cos(b*x + a)^2 + 1) -
3*b*c^2*d^3*(-1/(b^4*c^6*d^10))^(1/4)*log(2*(b*c*d^2*(-1/(b^4*c^6*d^10))^(
1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c^4*d^7*cos(b*x + a)^3 - b^3*c^...
```

### 3.270.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate(1/(d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(3/2), x)`

output `Timed out`

**3.270.7 Maxima [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2)), x)`

**3.270.8 Giac [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

output `integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2)), x)`

**3.270.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2} \left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

input `int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(5/2)),x)`

output `int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(5/2)), x)`

$$3.271 \quad \int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{5/2}} dx$$

3.271.1 Optimal result . . . . .	1578
3.271.2 Mathematica [A] (verified) . . . . .	1578
3.271.3 Rubi [A] (verified) . . . . .	1579
3.271.4 Maple [A] (verified) . . . . .	1580
3.271.5 Fricas [B] (verification not implemented) . . . . .	1580
3.271.6 Sympy [F(-1)] . . . . .	1580
3.271.7 Maxima [F] . . . . .	1581
3.271.8 Giac [F] . . . . .	1581
3.271.9 Mupad [B] (verification not implemented) . . . . .	1581

### 3.271.1 Optimal result

Integrand size = 25, antiderivative size = 33

$$\int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{5/2}} dx = -\frac{2cd(d \csc(a+bx))^{7/2}}{7b(c \sec(a+bx))^{7/2}}$$

output `-2/7*c*d*(d*csc(b*x+a))^(7/2)/b/(c*sec(b*x+a))^(7/2)`

### 3.271.2 Mathematica [A] (verified)

Time = 1.54 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.36

$$\int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{5/2}} dx = -\frac{2d^4 \cot^3(a+bx) \sqrt{d \csc(a+bx)}}{7bc^2 \sqrt{c \sec(a+bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(9/2)/(c*Sec[a + b*x])^(5/2), x]`

output `(-2*d^4*Cot[a + b*x]^3*Sqrt[d*Csc[a + b*x]])/(7*b*c^2*Sqrt[c*Sec[a + b*x]])`

**3.271.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {3042, 3099}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{5/2}} dx$$

↓ 3099

$$-\frac{2cd(d \csc(a + bx))^{7/2}}{7b(c \sec(a + bx))^{7/2}}$$

input `Int[(d*Csc[a + b*x])^(9/2)/(c*Sec[a + b*x])^(5/2),x]`

output `(-2*c*d*(d*Csc[a + b*x])^(7/2))/(7*b*(c*Sec[a + b*x])^(7/2))`

**3.271.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3099 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]`



**3.271.4 Maple [A] (verified)**

Time = 8.31 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.21

method	result	size
default	$-\frac{2\sqrt{d\csc(bx+a)}d^4\cot(bx+a)^3}{7b\sqrt{c\sec(bx+a)}c^2}$	40

input `int((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/7/b*(d*csc(b*x+a))^(1/2)*d^4/(c*sec(b*x+a))^(1/2)/c^2*cot(b*x+a)^3`

**3.271.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 67 vs.  $2(27) = 54$ .

Time = 0.30 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.03

$$\int \frac{(d\csc(a+bx))^{9/2}}{(c\sec(a+bx))^{5/2}} dx = \frac{2d^4\sqrt{\frac{c}{\cos(bx+a)}}\sqrt{\frac{d}{\sin(bx+a)}}\cos(bx+a)^4}{7(bc^3\cos(bx+a)^2 - bc^3)\sin(bx+a)}$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fracas")`

output `2/7*d^4*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)^4/((b*c^3*cos(b*x + a)^2 - b*c^3)*sin(b*x + a))`

**3.271.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d\csc(a+bx))^{9/2}}{(c\sec(a+bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(9/2)/(c*sec(b*x+a))**(5/2),x)`

output `Timed out`

**3.271.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{9/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(9/2)/(c*sec(b*x + a))^(5/2), x)`

**3.271.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{9/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(9/2)/(c*sec(b*x + a))^(5/2), x)`

**3.271.9 Mupad [B] (verification not implemented)**

Time = 14.47 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.82

$$\int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{5/2}} dx = \frac{2 d^4 \sqrt{\frac{d}{\sin(a+bx)}} (3 \sin(2a + 2bx) - \sin(6a + 6bx))}{7 b c^2 \sqrt{\frac{c}{\cos(a+bx)}} (15 \cos(2a + 2bx) - 6 \cos(4a + 4bx) + \cos(6a + 6bx) - 10)}$$

input `int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(5/2),x)`

output `(2*d^4*(d/sin(a + b*x))^(1/2)*(3*sin(2*a + 2*b*x) - sin(6*a + 6*b*x)))/(7*b*c^2*(c/cos(a + b*x))^(1/2)*(15*cos(2*a + 2*b*x) - 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) - 10))`

**3.272**       $\int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{5/2}} dx$

3.272.1 Optimal result . . . . . 1582  
 3.272.2 Mathematica [C] (verified) . . . . . 1582  
 3.272.3 Rubi [A] (verified) . . . . . 1583  
 3.272.4 Maple [B] (verified) . . . . . 1585  
 3.272.5 Fricas [C] (verification not implemented) . . . . . 1586  
 3.272.6 Sympy [F(-1)] . . . . . 1587  
 3.272.7 Maxima [F] . . . . . 1587  
 3.272.8 Giac [F] . . . . . 1587  
 3.272.9 Mupad [F(-1)] . . . . . 1588

**3.272.1 Optimal result**

Integrand size = 25, antiderivative size = 135

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx = \frac{6d^3 \sqrt{d \csc(a + bx)}}{5bc(c \sec(a + bx))^{3/2}} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} + \frac{6d^4 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{5bc^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}$$

output `-2/5*d*(d*csc(b*x+a))^(5/2)/b/c/(c*sec(b*x+a))^(3/2)+6/5*d^3*(d*csc(b*x+a))^(1/2)/b/c/(c*sec(b*x+a))^(3/2)-6/5*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/c^2/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)`

**3.272.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.89 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.75

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx = \frac{d^5 \left( (1 - 3 \cos(2(a + bx))) \cot^2(a + bx) \csc^2(a + bx) + 6 \sqrt{-\cot^2(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{1 - \cos(2(a + bx))}{1 + \cos(2(a + bx))}\right) \right)}{5bc^3(d \csc(a + bx))^{3/2}}$$

input `Integrate[(d*Csc[a + b*x])^(7/2)/(c*Sec[a + b*x])^(5/2),x]`

3.272.       $\int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{5/2}} dx$

output  $(d^5*((1 - 3*\text{Cos}[2*(a + b*x)])*\text{Cot}[a + b*x]^2*\text{Csc}[a + b*x]^2 + 6*(-\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Hypergeometric2F1}[-1/2, 1/4, 1/2, \text{Csc}[a + b*x]^2])*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(5*b*c^3*(d*\text{Csc}[a + b*x])^{(3/2)})$

### 3.272.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3103, 3042, 3105, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3103} \\ & -\frac{3d^2 \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx}{5c^2} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{3d^2 \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx}{5c^2} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} \\ & \quad \downarrow \text{3105} \\ & -\frac{3d^2 \left( -2d^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \right)}{5c^2} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{3d^2 \left( -2d^2 \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \right)}{5c^2} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} \\ & \quad \downarrow \text{3110} \\ & -\frac{3d^2 \left( -\frac{2d^2 \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{\sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2cd \sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} \right)}{5c^2} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} \end{aligned}$$

---

3.272.  $\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & -\frac{3d^2 \left( -\frac{2d^2 \int \sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)} dx}{\sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right)}{5c^2} - \frac{2d(d \csc(a+bx))^{5/2}}{5bc(c \sec(a+bx))^{3/2}} \\
 & \downarrow 3052 \\
 & -\frac{3d^2 \left( -\frac{2d^2 \int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right)}{5c^2} - \frac{2d(d \csc(a+bx))^{5/2}}{5bc(c \sec(a+bx))^{3/2}} \\
 & \downarrow 3042 \\
 & -\frac{3d^2 \left( -\frac{2d^2 \int \sqrt{\sin(2a+2bx)} dx}{\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right)}{5c^2} - \frac{2d(d \csc(a+bx))^{5/2}}{5bc(c \sec(a+bx))^{3/2}} \\
 & \downarrow 3119 \\
 & -\frac{3d^2 \left( -\frac{2d^2 E(a+bx - \frac{\pi}{4} | 2)}{b \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{2cd \sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}} \right)}{5c^2} - \frac{2d(d \csc(a+bx))^{5/2}}{5bc(c \sec(a+bx))^{3/2}}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(7/2)/(c*Sec[a + b*x])^(5/2),x]`

output `(-2*d*(d*Csc[a + b*x])^(5/2))/(5*b*c*(c*Sec[a + b*x])^(3/2)) - (3*d^2*((-2*c*d*Sqrt[d*Csc[a + b*x]])/(b*(c*Sec[a + b*x])^(3/2)) - (2*d^2*EllipticE[a - Pi/4 + b*x, 2])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])))/(5*c^2)`

### 3.272.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

```
rule 3103 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Simp[a^2*((n + 1)/(b^2*(m - 1))) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]
```

```
rule 3105 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Simp[a^2*((m + n - 2)/(m - 1)) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]
```

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.272.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 384 vs.  $2(140) = 280$ .

Time = 7.82 (sec) , antiderivative size = 385, normalized size of antiderivative = 2.85

method	result
default	$-\frac{\sqrt{2}d^3\sqrt{d\csc(bx+a)}\left(6\sqrt{1+\csc(bx+a)}-\cot(bx+a)\sqrt{\cot(bx+a)-\csc(bx+a)+1}\sqrt{\cot(bx+a)-\csc(bx+a)}\right)\text{EllipticE}\left(\sqrt{1+\csc(bx+a)}\right)}{2d}$

```
input int((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

output 
$$-1/5/b*2^{(1/2)}*d^3*(d*csc(b*x+a))^{(1/2)}/(c*sec(b*x+a))^{(1/2)}/c^2*(6*(1+csc(b*x+a)-cot(b*x+a))^{(1/2)}*(cot(b*x+a)-csc(b*x+a)+1)^{(1/2)}*(cot(b*x+a)-csc(b*x+a))^{(1/2)}*EllipticE((1+csc(b*x+a)-cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})-3*(1+csc(b*x+a)-cot(b*x+a))^{(1/2)}*(cot(b*x+a)-csc(b*x+a)+1)^{(1/2)}*(cot(b*x+a)-csc(b*x+a))^{(1/2)}*EllipticF((1+csc(b*x+a)-cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})+6*(1+csc(b*x+a)-cot(b*x+a))^{(1/2)}*(cot(b*x+a)-csc(b*x+a)+1)^{(1/2)}*(cot(b*x+a)-csc(b*x+a))^{(1/2)}*EllipticE((1+csc(b*x+a)-cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*sec(b*x+a)-3*(1+csc(b*x+a)-cot(b*x+a))^{(1/2)}*(cot(b*x+a)-csc(b*x+a)+1)^{(1/2)}*(cot(b*x+a)-csc(b*x+a))^{(1/2)}*EllipticF((1+csc(b*x+a)-cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*sec(b*x+a)-3*2^{(1/2)}+2^{(1/2)}*cot(b*x+a)*csc(b*x+a))$$

### 3.272.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.88

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx = 3(d^3 \cos(bx + a)^2 - d^3) \sqrt{-4i cdE(\arcsin(\cos(bx + a) + i \sin(bx + a)) | -1)} + 3(d^3 \cos(bx + a)^2 - d^3)$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

output 
$$-1/10*(3*(d^3*\cos(b*x + a)^2 - d^3)*\sqrt{-4*I*c*d}*elliptic\_e(\arcsin(\cos(b*x + a) + I*\sin(b*x + a)), -1) + 3*(d^3*\cos(b*x + a)^2 - d^3)*\sqrt{4*I*c*d}*elliptic\_e(\arcsin(\cos(b*x + a) - I*\sin(b*x + a)), -1) - 3*(d^3*\cos(b*x + a)^2 - d^3)*\sqrt{-4*I*c*d}*elliptic\_f(\arcsin(\cos(b*x + a) + I*\sin(b*x + a))), -1) - 3*(d^3*\cos(b*x + a)^2 - d^3)*\sqrt{4*I*c*d}*elliptic\_f(\arcsin(\cos(b*x + a) - I*\sin(b*x + a))), -1) - 4*(3*d^3*\cos(b*x + a)^4 - 2*d^3*\cos(b*x + a)^2)*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)})/(b*c^3*\cos(b*x + a)^2 - b*c^3)$$

**3.272.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(5/2),x)`output `Timed out`**3.272.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{7/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`output `integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(5/2), x)`**3.272.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{7/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`output `integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(5/2), x)`



**3.272.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{7/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

input `int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(5/2),x)`output `int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(5/2), x)`

### 3.273 $\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{5/2}} dx$

3.273.1 Optimal result . . . . .	1589
3.273.2 Mathematica [A] (verified) . . . . .	1590
3.273.3 Rubi [A] (verified) . . . . .	1590
3.273.4 Maple [B] (warning: unable to verify) . . . . .	1595
3.273.5 Fricas [C] (verification not implemented) . . . . .	1595
3.273.6 Sympy [F(-1)] . . . . .	1596
3.273.7 Maxima [F] . . . . .	1597
3.273.8 Giac [F] . . . . .	1597
3.273.9 Mupad [F(-1)] . . . . .	1597

#### 3.273.1 Optimal result

Integrand size = 25, antiderivative size = 329

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx = -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} + \frac{d^2 \arctan\left(1 - \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{\sqrt{2}bc^2 \sqrt{c \sec(a + bx)}} - \frac{d^2 \arctan\left(1 + \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{\sqrt{2}bc^2 \sqrt{c \sec(a + bx)}} + \frac{d^2 \sqrt{d \csc(a + bx)} \log\left(1 - \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{2\sqrt{2}bc^2 \sqrt{c \sec(a + bx)}} - \frac{d^2 \sqrt{d \csc(a + bx)} \log\left(1 + \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{2\sqrt{2}bc^2 \sqrt{c \sec(a + bx)}}$$

output

```
-2/3*d*(d*csc(b*x+a))^(3/2)/b/c/(c*sec(b*x+a))^(3/2)-1/2*d^2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/2*d^2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2*2^(1/2)/(c*sec(b*x+a))^(1/2)+1/4*d^2*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/4*d^2*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2*2^(1/2)/(c*sec(b*x+a))^(1/2)
```

**3.273.2 Mathematica [A] (verified)**

Time = 2.79 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.47

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx =$$

$$\frac{d^3 \left( 4 \cot^2(a + bx) - 3\sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}} \right) \sqrt[4]{\cot^2(a + bx)} + 3\sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}{1 + \sqrt{\cot^2(a + bx)}} \right) \right)}{6bc^3 \sqrt{d \csc(a + bx)}}$$

input `Integrate[(d*Csc[a + b*x])^(5/2)/(c*Sec[a + b*x])^(5/2),x]`output `-1/6*(d^3*(4*Cot[a + b*x]^2 - 3*Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2)]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(1/4) + 3*Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(1/4))*Sqrt[c*Sec[a + b*x]])/(b*c^3*Sqrt[d*Csc[a + b*x]])`**3.273.3 Rubi [A] (verified)**Time = 0.60 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.65, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3103, 3042, 3109, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx \\ & \quad \downarrow \text{3103} \\ & -\frac{d^2 \int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx}{c^2} - \frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.273.  $\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx$

$$\begin{aligned}
& \frac{d^2 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{c^2} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}} \\
& \quad \downarrow \text{3109} \\
& \frac{d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{c^2 \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{c^2 \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}} \\
& \quad \downarrow \text{3957} \\
& \frac{d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}(\tan^2(a+bx)+1)} d \tan(a+bx)}{bc^2 \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}} \\
& \quad \downarrow \text{266} \\
& \frac{2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{bc^2 \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}} \\
& \quad \downarrow \text{755} \\
& \frac{2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{bc^2 \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}} \\
& \quad \downarrow \text{1476} \\
& \frac{2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} + \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d \sqrt{\tan(a+bx)} \right) \right)}{bc^2 \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}} \\
& \quad \downarrow \text{1082} \\
& \frac{2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} + \frac{1}{2} \left( \frac{\int \frac{1}{-\tan(a+bx)-1} d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(a+bx)+1} d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) \right)}{bc^2 \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}}
\end{aligned}$$

---

3.273.  $\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{5/2}} dx$

↓ 217

$$2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d\sqrt{\tan(a+bx)} + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) \right)$$


---


$$\frac{bc^2 \sqrt{c \sec(a+bx)}}{2d(d \csc(a+bx))^{3/2}}$$

$$\frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}}$$

↓ 1479

$$2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \left( - \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) \right)$$


---


$$\frac{bc^2 \sqrt{c \sec(a+bx)}}{2d(d \csc(a+bx))^{3/2}}$$

$$\frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}}$$

↓ 25

$$2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) \right)$$


---


$$\frac{bc^2 \sqrt{c \sec(a+bx)}}{2d(d \csc(a+bx))^{3/2}}$$

$$\frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}}$$

↓ 27

$$2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) \right)$$


---


$$\frac{bc^2 \sqrt{c \sec(a+bx)}}{2d(d \csc(a+bx))^{3/2}}$$

$$\frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}}$$

↓ 1103

$$2d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} \right) \right)$$


---


$$\frac{bc^2 \sqrt{c \sec(a+bx)}}{2d(d \csc(a+bx))^{3/2}}$$

$$\frac{2d(d \csc(a+bx))^{3/2}}{3bc(c \sec(a+bx))^{3/2}}$$

---

3.273.  $\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{5/2}} dx$

input `Int[(d*Csc[a + b*x])^(5/2)/(c*Sec[a + b*x])^(5/2),x]`

output `(-2*d*(d*Csc[a + b*x])^(3/2))/(3*b*c*(c*Sec[a + b*x])^(3/2)) - (2*d^2*Sqrt[d*Csc[a + b*x]]*((-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]])/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[Tan[a + b*x]]/(b*c^2*Sqrt[c*Sec[a + b*x]])`

### 3.273.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3103 `Int[(csc[(e_) + (f_)*(x_)]*(a_.))^m_*((b_)*sec[(e_) + (f_)*(x_)])^n_, x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Simp[a^2*((n + 1)/(b^2*(m - 1))) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_) + (f_)*(x_)]*(a_.))^m_*((b_)*sec[(e_) + (f_)*(x_)])^n_, x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^n_, x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.273.4 Maple [B] (warning: unable to verify)**

Leaf count of result is larger than twice the leaf count of optimal. 729 vs.  $2(271) = 542$ .

Time = 57.00 (sec) , antiderivative size = 730, normalized size of antiderivative = 2.22

method	result
default	$\sqrt{2} \left( 3 \ln \left( \frac{(1-\cos(bx+a))^2 \csc(bx+a) + 2\sqrt{(1-\cos(bx+a))((1-\cos(bx+a))^2 \csc(bx+a)^2 - 1) \csc(bx+a) \sin(bx+a) - 2\cos(bx+a) + 2 - \sin(bx+a)}}{1-\cos(bx+a)} \right) \right) (1)$

input `int((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 1/12/b*x^2^{(1/2)}*(3*\ln(1/(1-\cos(b*x+a)))*((1-\cos(b*x+a))^2*\csc(b*x+a)+2*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)- \\ & 2*\cos(b*x+a)+2-\sin(b*x+a)))*(1-\cos(b*x+a))^2*\csc(b*x+a)^2-6*\arctan(1/(1-\cos(b*x+a))*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)-\cos(b*x+a)+1))*(1-\cos(b*x+a))^2*\csc(b*x+a)^2-3*\ln(-1/(1-\cos(b*x+a))*(-1-\cos(b*x+a))^2*\csc(b*x+a)+2*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)+2*\cos(b*x+a)-2+\sin(b*x+a))) \\ & *(1-\cos(b*x+a))^2*\csc(b*x+a)^2-6*\arctan(1/(1-\cos(b*x+a))*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*\sin(b*x+a)+\cos(b*x+a)-1))*(1-\cos(b*x+a))^2*\csc(b*x+a)^2+2*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*(1-\cos(b*x+a))^2*\csc(b*x+a)^2-2*((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}*(1-\cos(b*x+a))* \\ & (d/(1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)+\sin(b*x+a)))^{(5/2)}/((1-\cos(b*x+a))*((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)*\csc(b*x+a))^{(1/2)}/((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1)^2/(-c*((1-\cos(b*x+a))^2*\csc(b*x+a)^2+1)/((1-\cos(b*x+a))^2*\csc(b*x+a)^2-1))^{(5/2)}*\csc(b*x+a) \end{aligned}$$

**3.273.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.69 (sec) , antiderivative size = 1396, normalized size of antiderivative = 4.24

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

---

3.273. 
$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx$$



output

```
-1/24*(3*b*c^3*(-d^10/(b^4*c^10))^(1/4)*log(1/2*d^8*cos(b*x + a)*sin(b*x + a) + 1/2*(b*c^2*d^5*(-d^10/(b^4*c^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c^7*cos(b*x + a)^3 - b^3*c^7*cos(b*x + a))*(-d^10/(b^4*c^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1/4*(2*b^2*c^5*d^3*cos(b*x + a)^2 - b^2*c^5*d^3)*sqrt(-d^10/(b^4*c^10))*sin(b*x + a) - 3*b*c^3*(-d^10/(b^4*c^10))^(1/4)*log(1/2*d^8*cos(b*x + a)*sin(b*x + a) - 1/2*(b*c^2*d^5*(-d^10/(b^4*c^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c^7*cos(b*x + a)^3 - b^3*c^7*cos(b*x + a))*(-d^10/(b^4*c^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1/4*(2*b^2*c^5*d^3*cos(b*x + a)^2 - b^2*c^5*d^3)*sqrt(-d^10/(b^4*c^10))*sin(b*x + a) + 3*I*b*c^3*(-d^10/(b^4*c^10))^(1/4)*log(1/2*d^8*cos(b*x + a)*sin(b*x + a) + 1/2*(I*b*c^2*d^5*(-d^10/(b^4*c^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (I*b^3*c^7*cos(b*x + a)^3 - I*b^3*c^7*cos(b*x + a))*(-d^10/(b^4*c^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 1/4*(2*b^2*c^5*d^3*cos(b*x + a)^2 - b^2*c^5*d^3)*sqrt(-d^10/(b^4*c^10))*sin(b*x + a) - 3*I*b*c^3*(-d^10/(b^4*c^10))^(1/4)*log(1/2*d^8*cos(b*x + a)*sin(b*x + a) + 1/2*(-I*b*c^2*d^5*(-d^10/(b^4*c^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) - (-I*b^3*c^7*cos(b*x + a)^3 + I*b^3*c^7*cos(b*x + a))*(-d^10/(b^4*c^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 1/4*(2*b^2*c^5*d^3*cos(b*x + a)^2 - b^2*c^5*d^3)*sqrt(-d^10/(b^4*c^10))*sin(b*x + a) - 3*b*c^3*(-d^10/(b^4*c^10))^(1/4)*log(d^8 + 2*(b^3*c^7*(-...
```

### 3.273.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(5/2),x)`

output `Timed out`

**3.273.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{5/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(5/2), x)`

**3.273.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{5/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(5/2), x)`

**3.273.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

input `int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(5/2),x)`

output `int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(5/2), x)`

**3.274**       $\int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{5/2}} dx$

3.274.1 Optimal result . . . . . 1598  
 3.274.2 Mathematica [C] (verified) . . . . . 1598  
 3.274.3 Rubi [A] (verified) . . . . . 1599  
 3.274.4 Maple [B] (verified) . . . . . 1601  
 3.274.5 Fricas [F] . . . . . 1602  
 3.274.6 Sympy [F(-1)] . . . . . 1602  
 3.274.7 Maxima [F] . . . . . 1602  
 3.274.8 Giac [F] . . . . . 1603  
 3.274.9 Mupad [F(-1)] . . . . . 1603

**3.274.1 Optimal result**

Integrand size = 25, antiderivative size = 94

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx = -\frac{2d\sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} - \frac{3d^2 E(a - \frac{\pi}{4} + bx | 2)}{bc^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}$$

output

```
-2*d*(d*csc(b*x+a))^(1/2)/b/c/(c*sec(b*x+a))^(3/2)+3*d^2*(sin(a+1/4*Pi+b*x)
)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/c^2/(d
*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

**3.274.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.42 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.85

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx = \frac{d^3 \left( 2 \cot^2(a + bx) + 3 \sqrt{-\cot^2(a + bx)} \operatorname{Hypergeometric2F1} \left( -\frac{1}{2}, \frac{1}{4}, \frac{1}{2}, \csc^2(a + bx) \right) \right) \sqrt{c \sec(a + bx)}}{bc^3 (d \csc(a + bx))^{3/2}}$$

input `Integrate[(d*Csc[a + b*x])^(3/2)/(c*Sec[a + b*x])^(5/2),x]`

output `-((d^3*(2*Cot[a + b*x]^2 + 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(b*c^3*(d*Csc[a + b*x])^(3/2))`

### 3.274.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3103, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx \\
 & \quad \downarrow \text{3103} \\
 & -\frac{3d^2 \int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx}{c^2} - \frac{2d \sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3d^2 \int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx}{c^2} - \frac{2d \sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3110} \\
 & -\frac{3d^2 \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2d \sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3d^2 \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2d \sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3052}
 \end{aligned}$$

---

3.274.  $\int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{5/2}} dx$

$$\begin{aligned}
 & -\frac{3d^2 \int \sqrt{\sin(2a + 2bx)} dx}{c^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2d \sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3d^2 \int \sqrt{\sin(2a + 2bx)} dx}{c^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2d \sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3119} \\
 & -\frac{3d^2 E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{bc^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2d \sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}}
 \end{aligned}$$

input `Int[(d*Csc[a + b*x])^(3/2)/(c*Sec[a + b*x])^(5/2),x]`

output `(-2*d*Sqrt[d*Csc[a + b*x]]/(b*c*(c*Sec[a + b*x])^(3/2)) - (3*d^2*EllipticE[a - Pi/4 + b*x, 2])/(b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])`

### 3.274.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x, x] /; FreeQ[{a, b, e, f}, x]`

rule 3103 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Simp[a^2*((n + 1)/(b^2*(m - 1))) Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]`

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.274.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 376 vs.  $2(109) = 218$ .

Time = 6.58 (sec) , antiderivative size = 377, normalized size of antiderivative = 4.01

method	result
default	$\frac{\sqrt{2}d\sqrt{d\csc(bx+a)}\left(6\sqrt{1+\csc(bx+a)-\cot(bx+a)}\sqrt{\cot(bx+a)-\csc(bx+a)+1}\sqrt{\cot(bx+a)-\csc(bx+a)}\operatorname{EllipticE}\left(\sqrt{1+\csc(bx+a)-\cot(bx+a)}\right)\right)}{\dots}$

```
input int((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/2/b*2^(1/2)*d*(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/c^2*(6*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))-3*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+6*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*sec(b*x+a)-3*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*sec(b*x+a)+2^(1/2)*cos(b*x+a)-3*2^(1/2))
```

**3.274.5 Fricas [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{3/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*d*csc(b*x + a)/(c^3*sec(b*x + a)^3), x)`

**3.274.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(5/2),x)`

output `Timed out`

**3.274.7 Maxima [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{3/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate((d*csc(b*x + a))^(3/2)/(c*sec(b*x + a))^(5/2), x)`

**3.274.8 Giac [F]**

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{(d \csc(bx + a))^{3/2}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate((d*csc(b*x + a))^(3/2)/(c*sec(b*x + a))^(5/2), x)`

**3.274.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

input `int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(5/2),x)`

output `int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(5/2), x)`



**3.275**  $\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx$

3.275.1 Optimal result . . . . . 1604  
 3.275.2 Mathematica [A] (verified) . . . . . 1605  
 3.275.3 Rubi [A] (verified) . . . . . 1605  
 3.275.4 Maple [A] (verified) . . . . . 1610  
 3.275.5 Fricas [C] (verification not implemented) . . . . . 1610  
 3.275.6 Sympy [F(-1)] . . . . . 1611  
 3.275.7 Maxima [F] . . . . . 1612  
 3.275.8 Giac [F] . . . . . 1612  
 3.275.9 Mupad [F(-1)] . . . . . 1612

**3.275.1 Optimal result**

Integrand size = 25, antiderivative size = 322

$$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx = \frac{d}{2bc\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{3/2}}$$

$$- \frac{3 \arctan\left(1 - \sqrt{2}\sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)}\sqrt{\tan(a+bx)}}{4\sqrt{2}bc^2\sqrt{c \sec(a+bx)}}$$

$$+ \frac{3 \arctan\left(1 + \sqrt{2}\sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)}\sqrt{\tan(a+bx)}}{4\sqrt{2}bc^2\sqrt{c \sec(a+bx)}}$$

$$- \frac{3\sqrt{d \csc(a+bx)} \log\left(1 - \sqrt{2}\sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{\tan(a+bx)}}{8\sqrt{2}bc^2\sqrt{c \sec(a+bx)}}$$

$$+ \frac{3\sqrt{d \csc(a+bx)} \log\left(1 + \sqrt{2}\sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{\tan(a+bx)}}{8\sqrt{2}bc^2\sqrt{c \sec(a+bx)}}$$

```
output 1/2*d/b/c/(c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2)+3/8*arctan(-1+2^(1/2)*
tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2*2^(1/2)/(c*s
ec(b*x+a))^(1/2)+3/8*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/
2)*tan(b*x+a)^(1/2)/b/c^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-3/16*ln(1-2^(1/2)*t
an(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2*2^(
1/2)/(c*sec(b*x+a))^(1/2)+3/16*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*
(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2*2^(1/2)/(c*sec(b*x+a))^(1/2)
```

### 3.275.2 Mathematica [A] (verified)

Time = 2.82 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.47

$$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx = \frac{d \left( 4 \cos^2(a+bx) - 3\sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a+bx)}}{\sqrt{2} \sqrt{\cot^2(a+bx)}} \right) \right) \sqrt[4]{\cot^2(a+bx)} + 3\sqrt{2} \arctan \left( \frac{-1 + \sqrt{\cot^2(a+bx)}}{\sqrt{2} \sqrt{\cot^2(a+bx)}} \right) \sqrt[4]{\cot^2(a+bx)}}{8bc^3 \sqrt{d \csc(a+bx)}}$$

input `Integrate[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(5/2),x]`

output `(d*(4*Cos[a + b*x]^2 - 3*Sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2)]/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(1/4) + 3*Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(1/4))*Sqrt[c*Sec[a + b*x]])/(8*b*c^3*Sqrt[d*Csc[a + b*x]])`

### 3.275.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.66, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 3108, 3042, 3109, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx \\ & \quad \downarrow \text{3108} \\ & \frac{3 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4c^2} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{3 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4c^2} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} \end{aligned}$$

---

3.275.  $\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx$

$$\begin{aligned}
& \downarrow 3109 \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}}dx}{4c^2\sqrt{c\sec(a+bx)}} + \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \downarrow 3042 \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}}dx}{4c^2\sqrt{c\sec(a+bx)}} + \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \downarrow 3957 \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}(\tan^2(a+bx)+1)}d\tan(a+bx)}{4bc^2\sqrt{c\sec(a+bx)}} + \\
& \quad \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \downarrow 266 \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}}{2bc^2\sqrt{c\sec(a+bx)}} + \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \downarrow 755 \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)} + \frac{1}{2}\int\frac{\tan(a+bx)+1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}\right)}{2bc^2\sqrt{c\sec(a+bx)}} + \\
& \quad \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \downarrow 1476 \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)} + \frac{1}{2}\left(\int\frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} + \\
& \quad \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \downarrow 1082 \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)} + \frac{1}{2}\left(\frac{\int\frac{1}{-\tan(a+bx)-1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} - \frac{\int\frac{1}{-\tan(a+bx)+1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} + \\
& \quad \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}
\end{aligned}$$

---

3.275.  $\int \frac{\sqrt{d\csc(a+bx)}}{(c\sec(a+bx))^{5/2}} dx$

↓ 217

$$\frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\ \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 1479

$$\frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(-\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}-\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\ \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 25

$$\frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}+\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\ \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 27

$$\frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}+\frac{1}{2}\int\frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\ \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

↓ 1103

$$\frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)+\frac{1}{2}\left(\frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}}\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\ \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}}$$

---

3.275.  $\int \frac{\sqrt{d\csc(a+bx)}}{(c\sec(a+bx))^{5/2}} dx$

input `Int[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(5/2),x]`

output `d/(2*b*c*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)) + (3*Sqrt[d*Csc[a + b*x]]*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[Tan[a + b*x]]/(2*b*c^2*Sqrt[c*Sec[a + b*x]])`

### 3.275.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3108 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.275.4 Maple [A] (verified)**

Time = 55.58 (sec) , antiderivative size = 442, normalized size of antiderivative = 1.37

method	result
default	$\sqrt{2} \left( 4 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sqrt{2} \cos(bx+a)^2 + 4 \cos(bx+a) \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} + 3 \ln \left( -2 \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \cot(bx+a) \right) \right)$

input `int((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 1/16/b*2^{(1/2)}*(4*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^{(1/2)}*2^{(1/2)}* \\ & \cos(b*x+a)^2+4*\cos(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2 \\ & )^{(1/2)}+3*\ln(-2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^{(1/2)}*co \\ & t(b*x+a)-2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^{(1/2)}*csc(b*x \\ & +a)-2*\cot(b*x+a)+2)-6*\arctan((\sin(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/( \\ & \cos(b*x+a)+1)^{(1/2)}+\cos(b*x+a)-1)/(\cos(b*x+a)-1))+6*\arctan((-sin(b*x+a) \\ & *2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^{(1/2)}+\cos(b*x+a)-1)/(co \\ & s(b*x+a)-1))-3*\ln(2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^{(1/2)} \\ & )*\cot(b*x+a)+2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^{(1/2)}*csc \\ & (b*x+a)-2*\cot(b*x+a)+2))*(d*csc(b*x+a))^(1/2)/(c*\sec(b*x+a) \\ & )^(1/2)/(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^{(1/2)}/c^2*\sin(b*x+a) \end{aligned}$$

**3.275.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.64 (sec) , antiderivative size = 1309, normalized size of antiderivative = 4.07

$$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fracas")`

output

```
-1/32*(3*b*c^3*(-d^2/(b^4*c^10))^(1/4)*log(-27/2*d^2*cos(b*x + a)*sin(b*x
+ a) + 27/2*(b*c^2*d*(-d^2/(b^4*c^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) +
(b^3*c^7*cos(b*x + a)^3 - b^3*c^7*cos(b*x + a))*(-d^2/(b^4*c^10))^(3/4))*
sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) - 27/4*(2*b^2*c^5*d*cos(b*x + a)
^2 - b^2*c^5*d)*sqrt(-d^2/(b^4*c^10))) - 3*b*c^3*(-d^2/(b^4*c^10))^(1/4)*l
og(-27/2*d^2*cos(b*x + a)*sin(b*x + a) - 27/2*(b*c^2*d*(-d^2/(b^4*c^10))^(
1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c^7*cos(b*x + a)^3 - b^3*c^7*cos(b
*x + a))*(-d^2/(b^4*c^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)
) - 27/4*(2*b^2*c^5*d*cos(b*x + a)^2 - b^2*c^5*d)*sqrt(-d^2/(b^4*c^10))) -
3*I*b*c^3*(-d^2/(b^4*c^10))^(1/4)*log(-27/2*d^2*cos(b*x + a)*sin(b*x + a)
- 27/2*(I*b*c^2*d*(-d^2/(b^4*c^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (
-I*b^3*c^7*cos(b*x + a)^3 + I*b^3*c^7*cos(b*x + a))*(-d^2/(b^4*c^10))^(3/4
))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 27/4*(2*b^2*c^5*d*cos(b*x +
a)^2 - b^2*c^5*d)*sqrt(-d^2/(b^4*c^10))) + 3*I*b*c^3*(-d^2/(b^4*c^10))^(1
/4)*log(-27/2*d^2*cos(b*x + a)*sin(b*x + a) - 27/2*(-I*b*c^2*d*(-d^2/(b^4*
c^10))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (I*b^3*c^7*cos(b*x + a)^3 - I*b
^3*c^7*cos(b*x + a))*(-d^2/(b^4*c^10))^(3/4))*sqrt(c/cos(b*x + a))*sqrt(d/
sin(b*x + a)) + 27/4*(2*b^2*c^5*d*cos(b*x + a)^2 - b^2*c^5*d)*sqrt(-d^2/(b
^4*c^10))) + 3*b*c^3*(-d^2/(b^4*c^10))^(1/4)*log(27*d^2 + 54*(b^3*c^7*(-d^
2/(b^4*c^10))^(3/4)*cos(b*x + a)^2*sin(b*x + a) + (b*c^2*d*cos(b*x + a)...
```

### 3.275.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate((d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(5/2),x)`

output `Timed out`



**3.275.7 Maxima [F]**

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{\sqrt{d \csc(bx + a)}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(5/2), x)`

**3.275.8 Giac [F]**

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{\sqrt{d \csc(bx + a)}}{(c \sec(bx + a))^{5/2}} dx$$

input `integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(5/2), x)`

**3.275.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{5/2}} dx = \int \frac{\sqrt{\frac{d}{\sin(a+bx)}}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

input `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(5/2),x)`

output `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(5/2), x)`

**3.276**  $\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{5/2}} dx$

3.276.1 Optimal result . . . . . 1613  
 3.276.2 Mathematica [C] (verified) . . . . . 1613  
 3.276.3 Rubi [A] (verified) . . . . . 1614  
 3.276.4 Maple [B] (verified) . . . . . 1616  
 3.276.5 Fracas [F] . . . . . 1617  
 3.276.6 Sympy [F(-1)] . . . . . 1617  
 3.276.7 Maxima [F] . . . . . 1617  
 3.276.8 Giac [F] . . . . . 1618  
 3.276.9 Mupad [F(-1)] . . . . . 1618

**3.276.1 Optimal result**

Integrand size = 25, antiderivative size = 95

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{5/2}} dx = \frac{d}{3bc(d \csc(a+bx))^{3/2}(c \sec(a+bx))^{3/2}} + \frac{E(a - \frac{\pi}{4} + bx | 2)}{2bc^2 \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{\sin(2a+2bx)}}$$

output `1/3*d/b/c/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2)-1/2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/c^2/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)`

**3.276.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{d \csc(a+bx)}(c \sec(a+bx))^{5/2}} dx = \frac{d \left( 1 + \cos(2(a+bx)) + 3 \sqrt[4]{-\cot^2(a+bx)} \operatorname{Hypergeometric2F1}(\dots) \right)}{6bc^3(d \csc(a+bx))^{3/2}}$$

input `Integrate[1/(Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(5/2)),x]`

output  $(d*(1 + \text{Cos}[2*(a + b*x)] + 3*(-\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Hypergeometric2F1}[-1/2, 1/4, 1/2, \text{Csc}[a + b*x]^2])* \text{Sqrt}[c*\text{Sec}[a + b*x]])/(6*b*c^3*(d*\text{Csc}[a + b*x])^{(3/2)})$

### 3.276.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {3042, 3108, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \sec(a + bx))^{5/2} \sqrt{d \csc(a + bx)}} dx \\
 & \quad \downarrow \text{3108} \\
 & \frac{\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx}{2c^2} + \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx}{2c^2} + \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3110} \\
 & \frac{\int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{2c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} + \\
 & \quad \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{2c^2 \sqrt{c \cos(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)} \sqrt{d \csc(a + bx)}} + \\
 & \quad \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3052}
 \end{aligned}$$

---

3.276.  $\int \frac{1}{\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{5/2}} dx$

$$\frac{\int \sqrt{\sin(2a + 2bx)} dx}{2c^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} + \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}$$

↓ 3042

$$\frac{\int \sqrt{\sin(2a + 2bx)} dx}{2c^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} + \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}$$

↓ 3119

$$\frac{E\left(a + bx - \frac{\pi}{4} \mid 2\right)}{2bc^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} + \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}}$$

input `Int[1/(Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(5/2)),x]`

output `d/(3*b*c*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2)) + EllipticE[a - Pi/4 + b*x, 2]/(2*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])`

### 3.276.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3108 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

```
rule 3110 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Ssin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Ssin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

### 3.276.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 408 vs.  $2(106) = 212$ .

Time = 6.61 (sec) , antiderivative size = 409, normalized size of antiderivative = 4.31

method	result
default	$-\frac{\sqrt{2} \left( 2 \cos(bx+a)^4 \sqrt{2-3\sqrt{1+\csc(bx+a)-\cot(bx+a)}} \sqrt{\cot(bx+a)-\csc(bx+a)+1} \sqrt{\cot(bx+a)-\csc(bx+a)} \operatorname{EllipticF}\left(\sqrt{1+\csc(bx+a)-\cot(bx+a)}, \frac{1}{2}\right) \right)}{c^2 \sec(bx+a)^5}$

```
input int(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/12/b*2^(1/2)*(2*cos(b*x+a)^4*2^(1/2)-3*(1+csc(b*x+a)-cot(b*x+a))^(1/2)*
(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2)*EllipticF((1
+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)+6*(1+csc(b*x+a)-cot(
b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-csc(b*x+a))^(1/2
)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))*cos(b*x+a)-3*(1+c
sc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)-cs
c(b*x+a))^(1/2)*EllipticF((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+6*(
1+csc(b*x+a)-cot(b*x+a))^(1/2)*(cot(b*x+a)-csc(b*x+a)+1)^(1/2)*(cot(b*x+a)
-csc(b*x+a))^(1/2)*EllipticE((1+csc(b*x+a)-cot(b*x+a))^(1/2),1/2*2^(1/2))+
2^(1/2)*cos(b*x+a)^2-3*2^(1/2)*cos(b*x+a))/(c*sec(b*x+a))^(1/2)/(d*csc(b*x
+a))^(1/2)/c^2*sec(b*x+a)*csc(b*x+a)
```

**3.276.5 Fracas [F]**

$$\int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\sqrt{d \csc(bx + a)} (c \sec(bx + a))^{5/2}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^3*d*csc(b*x + a)*sec(b*x + a)^3), x)`

**3.276.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate(1/(d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(5/2),x)`

output `Timed out`

**3.276.7 Maxima [F]**

$$\int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\sqrt{d \csc(bx + a)} (c \sec(bx + a))^{5/2}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2)), x)`

**3.276.8 Giac [F]**

$$\int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\sqrt{d \csc(bx + a)} (c \sec(bx + a))^{5/2}} dx$$

input `integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2)), x)`

**3.276.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a + bx)}\right)^{5/2} \sqrt{\frac{d}{\sin(a + bx)}}} dx$$

input `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2)),x)`

output `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2)), x)`

**3.277**       $\int \frac{1}{(d \csc(a+bx))^{3/2}(c \sec(a+bx))^{5/2}} dx$

3.277.1 Optimal result . . . . . 1619  
 3.277.2 Mathematica [A] (verified) . . . . . 1620  
 3.277.3 Rubi [A] (verified) . . . . . 1620  
 3.277.4 Maple [A] (warning: unable to verify) . . . . . 1626  
 3.277.5 Fricas [C] (verification not implemented) . . . . . 1626  
 3.277.6 Sympy [F(-1)] . . . . . 1627  
 3.277.7 Maxima [F] . . . . . 1628  
 3.277.8 Giac [F] . . . . . 1628  
 3.277.9 Mupad [F(-1)] . . . . . 1628

**3.277.1 Optimal result**

Integrand size = 25, antiderivative size = 371

$$\int \frac{1}{(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{5/2}} dx =$$

$$\frac{c}{4bd\sqrt{d \csc(a + bx)}(c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d \csc(a + bx)}(c \sec(a + bx))^{3/2}}$$

$$- \frac{3 \arctan\left(1 - \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{32\sqrt{2}bc^2d^2\sqrt{c \sec(a + bx)}}$$

$$+ \frac{3 \arctan\left(1 + \sqrt{2}\sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{32\sqrt{2}bc^2d^2\sqrt{c \sec(a + bx)}}$$

$$- \frac{3\sqrt{d \csc(a + bx)} \log\left(1 - \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{64\sqrt{2}bc^2d^2\sqrt{c \sec(a + bx)}}$$

$$+ \frac{3\sqrt{d \csc(a + bx)} \log\left(1 + \sqrt{2}\sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{64\sqrt{2}bc^2d^2\sqrt{c \sec(a + bx)}}$$



output 
$$\begin{aligned} & -1/4*c/b/d/(c*\sec(b*x+a))^(7/2)/(d*csc(b*x+a))^(1/2)+1/16/b/c/d/(c*\sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2)+3/64*\arctan(-1+2^(1/2)*\tan(b*x+a)^(1/2))* \\ & (d*csc(b*x+a))^(1/2)*\tan(b*x+a)^(1/2)/b/c^2/d^2*2^(1/2)/(c*\sec(b*x+a))^(1/2) \\ & +3/64*\arctan(1+2^(1/2)*\tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*\tan(b*x+a)^(1/2) \\ & /b/c^2/d^2*2^(1/2)/(c*\sec(b*x+a))^(1/2)-3/128*\ln(1-2^(1/2)*\tan(b*x+a)^(1/2) \\ & +\tan(b*x+a))*(d*csc(b*x+a))^(1/2)*\tan(b*x+a)^(1/2)/b/c^2/d^2*2^(1/2) \\ & /c*\sec(b*x+a))^(1/2)+3/128*\ln(1+2^(1/2)*\tan(b*x+a)^(1/2)+\tan(b*x+a))*(d*csc(b*x+a))^(1/2) \\ & *\tan(b*x+a)^(1/2)/b/c^2/d^2*2^(1/2)/(c*\sec(b*x+a))^(1/2) \end{aligned}$$

### 3.277.2 Mathematica [A] (verified)

Time = 3.09 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.44

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx = \frac{\left(4 + 6 \cos(2(a + bx)) + 2 \cos(4(a + bx)) + 3\sqrt{2} \arctan\left(\frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}\right)\right) \sqrt[4]{\cot^2(a + bx)} - 3\sqrt{2} \arctan\left(\frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}\right)}{64bc^3d\sqrt{d \csc(a + bx)}}$$

input `Integrate[1/((d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2)),x]`

output 
$$\begin{aligned} & -1/64*((4 + 6*\cos[2*(a + b*x)] + 2*\cos[4*(a + b*x)] + 3*Sqrt[2]*ArcTan[(-1 \\ & + Sqrt[Cot[a + b*x]^2])/(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))]*(Cot[a + b*x]^2)^(1/4) \\ & - 3*Sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b*x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2]) \\ & ]*(Cot[a + b*x]^2)^(1/4))*Sqrt[c*Sec[a + b*x]])/(b*c^3*d*Sqrt[d*Csc[a + b*x]]) \end{aligned}$$

### 3.277.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.69, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.680$ , Rules used = {3042, 3107, 3042, 3108, 3042, 3109, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.277. 
$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx$$

$$\begin{aligned}
& \int \frac{1}{(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{1}{(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{3/2}} dx \\
& \quad \downarrow \text{3107} \\
& \frac{\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3108} \\
& \frac{3 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4c^2} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4c^2} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3109} \\
& \frac{3\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{4c^2 \sqrt{c \sec(a+bx)}} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{3\sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{4c^2 \sqrt{c \sec(a+bx)}} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \\
& \quad \downarrow \text{3957}
\end{aligned}$$

$$\begin{aligned}
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\sqrt{\tan(a+bx)}(\tan^2(a+bx)+1)}d\tan(a+bx)}{4bc^2\sqrt{c\sec(a+bx)}} + \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \qquad \qquad \qquad \frac{8d^2}{c} \\
& \qquad \qquad \qquad \frac{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}}{c} \\
& \qquad \qquad \qquad \downarrow \text{266} \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\int\frac{1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}}{2bc^2\sqrt{c\sec(a+bx)}} + \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \qquad \qquad \qquad \frac{8d^2}{c} \\
& \qquad \qquad \qquad \frac{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}}{c} \\
& \qquad \qquad \qquad \downarrow \text{755} \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\int\frac{\tan(a+bx)+1}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}\right)}{2bc^2\sqrt{c\sec(a+bx)}} + \frac{d}{2bc(c\sec(a+bx))^{3/2}\sqrt{d\csc(a+bx)}} \\
& \qquad \qquad \qquad \frac{8d^2}{c} \\
& \qquad \qquad \qquad \frac{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}}{c} \\
& \qquad \qquad \qquad \downarrow \text{1476} \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\int\frac{1}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}+\int\frac{1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}}d\sqrt{\tan(a+bx)}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\
& \qquad \qquad \qquad \frac{8d^2}{c} \\
& \qquad \qquad \qquad \frac{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}}{c} \\
& \qquad \qquad \qquad \downarrow \text{1082} \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{\int\frac{1}{-\tan(a+bx)-1}d(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}-\frac{\int\frac{1}{-\tan(a+bx)-1}d(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\
& \qquad \qquad \qquad \frac{8d^2}{c} \\
& \qquad \qquad \qquad \frac{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}}{c} \\
& \qquad \qquad \qquad \downarrow \text{217} \\
& \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\int\frac{1-\tan(a+bx)}{\tan^2(a+bx)+1}d\sqrt{\tan(a+bx)}+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} + \frac{d}{2bc(c\sec(a+bx))^{3/2}} \\
& \qquad \qquad \qquad \frac{8d^2}{c} \\
& \qquad \qquad \qquad \frac{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}}{c} \\
& \qquad \qquad \qquad \downarrow \text{1479}
\end{aligned}$$

---

3.277.  $\int \frac{1}{(d\csc(a+bx))^{3/2}(c\sec(a+bx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(-\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}-\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}\right)+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\
 & \frac{c}{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}} \quad 8d^2 \\
 & \quad \downarrow \quad 25 \\
 & \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}+\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}\right)+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\
 & \frac{c}{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}} \quad 8d^2 \\
 & \quad \downarrow \quad 27 \\
 & \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}}{2\sqrt{2}}+\frac{1}{2}\int\frac{\sqrt{2}\sqrt{\tan(a+bx)}+1}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1}d\sqrt{\tan(a+bx)}\right)+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\
 & \frac{c}{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}} \quad 8d^2 \\
 & \quad \downarrow \quad 1103 \\
 & \frac{3\sqrt{\tan(a+bx)}\sqrt{d\csc(a+bx)}\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)}+1)}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}}\right)+\frac{1}{2}\left(\frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)}+1)}{2\sqrt{2}}-\frac{\log(\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)})}{2\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(a+bx)}} \\
 & \frac{c}{4bd(c\sec(a+bx))^{7/2}\sqrt{d\csc(a+bx)}} \quad 8d^2
 \end{aligned}$$

input `Int[1/((d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2)),x]`

output `-1/4*c/(b*d*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(7/2)) + (d/(2*b*c*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)) + (3*Sqrt[d*Csc[a + b*x]]*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]])/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[Tan[a + b*x]])/(2*b*c^2*Sqrt[c*Sec[a + b*x]))/(8*d^2)`

---

3.277.  $\int \frac{1}{(d\csc(a+bx))^{3/2}(c\sec(a+bx))^{5/2}} dx$

## 3.277.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2)]^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3107 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3108 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**3.277.4 Maple [A] (warning: unable to verify)**

Time = 49.00 (sec) , antiderivative size = 517, normalized size of antiderivative = 1.39

method	result
default	$\frac{\sqrt{2} \left( 16\sqrt{2} \sqrt{\frac{-\cos(bx+a)\sin(bx+a)}{(\cos(bx+a)+1)^2}} \cos(bx+a)^4 + 16\sqrt{2} \sqrt{\frac{-\cos(bx+a)\sin(bx+a)}{(\cos(bx+a)+1)^2}} \cos(bx+a)^3 - 4 \sqrt{\frac{-\cos(bx+a)\sin(bx+a)}{(\cos(bx+a)+1)^2}} \sqrt{2} \cos(bx+a) \right)}{\dots}$

input `int(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/128/b*2^{(1/2)}*(16*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} \\ & * \cos(b*x+a)^4 + 16*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} \\ & * \cos(b*x+a)^3 - 4*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} * 2^{(1/2)} * \cos \\ & (b*x+a)^2 - 4*\cos(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} \\ & + 3*\ln(2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} * \cot(b \\ & *x+a) + 2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} * \csc(b*x+a) \\ & - 2*\cot(b*x+a) + 2) - 6*\arctan((-\sin(b*x+a)*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos \\ & (b*x+a)+1)^2)^{(1/2)} + \cos(b*x+a) - 1)/(\cos(b*x+a) - 1) + 6*\arctan((\sin(b*x+a)*2^{(1/2)} \\ & *(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} + \cos(b*x+a) - 1)/(\cos(b \\ & *x+a) - 1)) - 3*\ln(-2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} * \\ & \cot(b*x+a) - 2*2^{(1/2)}*(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)} * \csc(b \\ & *x+a) - 2*\cot(b*x+a) + 2))/(\cos(b*x+a)+1)/(c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2) \\ & /(-\cos(b*x+a)*\sin(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}/c^2/d \end{aligned}$$

**3.277.5 Fracas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 1.30 (sec) , antiderivative size = 1375, normalized size of antiderivative = 3.71

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fracas")`

output

```

-1/256*(3*b*c^3*d^2*(-1/(b^4*c^10*d^6))^(1/4)*log(2*(b^3*c^7*d^4*(-1/(b^4*
c^10*d^6)))^(3/4)*cos(b*x + a)^2*sin(b*x + a) + (b*c^2*d*cos(b*x + a)^3 - b
*c^2*d*cos(b*x + a))*(-1/(b^4*c^10*d^6))^(1/4))*sqrt(c/cos(b*x + a))*sqrt(
d/sin(b*x + a)) + 1) - 3*b*c^3*d^2*(-1/(b^4*c^10*d^6))^(1/4)*log(-2*(b^3*c
^7*d^4*(-1/(b^4*c^10*d^6)))^(3/4)*cos(b*x + a)^2*sin(b*x + a) + (b*c^2*d*co
s(b*x + a)^3 - b*c^2*d*cos(b*x + a))*(-1/(b^4*c^10*d^6))^(1/4))*sqrt(c/cos
(b*x + a))*sqrt(d/sin(b*x + a)) + 1) + 3*I*b*c^3*d^2*(-1/(b^4*c^10*d^6))^(
1/4)*log(-2*(I*b^3*c^7*d^4*(-1/(b^4*c^10*d^6)))^(3/4)*cos(b*x + a)^2*sin(b*
x + a) + (-I*b*c^2*d*cos(b*x + a)^3 + I*b*c^2*d*cos(b*x + a))*(-1/(b^4*c^1
0*d^6))^(1/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a)) + 1) - 3*I*b*c^3*
d^2*(-1/(b^4*c^10*d^6))^(1/4)*log(-2*(-I*b^3*c^7*d^4*(-1/(b^4*c^10*d^6)))^(
3/4)*cos(b*x + a)^2*sin(b*x + a) + (I*b*c^2*d*cos(b*x + a)^3 - I*b*c^2*d*c
os(b*x + a))*(-1/(b^4*c^10*d^6))^(1/4))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*
x + a)) + 1) + 3*b*c^3*d^2*(-1/(b^4*c^10*d^6))^(1/4)*log(1/2*(b*c^2*d*(-1/
(b^4*c^10*d^6))^(1/4)*cos(b*x + a)^2*sin(b*x + a) + (b^3*c^7*d^4*cos(b*x +
a)^3 - b^3*c^7*d^4*cos(b*x + a))*(-1/(b^4*c^10*d^6))^(3/4))*sqrt(c/cos(b*
x + a))*sqrt(d/sin(b*x + a)) - 1/2*cos(b*x + a)*sin(b*x + a) - 1/4*(2*b^2*
c^5*d^3*cos(b*x + a)^2 - b^2*c^5*d^3)*sqrt(-1/(b^4*c^10*d^6))) - 3*b*c^3*d
^2*(-1/(b^4*c^10*d^6))^(1/4)*log(-1/2*(b*c^2*d*(-1/(b^4*c^10*d^6))^(1/4)*c
os(b*x + a)^2*sin(b*x + a) + (b^3*c^7*d^4*cos(b*x + a)^3 - b^3*c^7*d^4*...

```

### 3.277.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate(1/(d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(5/2), x)`

output `Timed out`



**3.277.7 Maxima [F]**

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2)), x)`

**3.277.8 Giac [F]**

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2)), x)`

**3.277.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2} \left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

input `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(3/2)),x)`

output `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(3/2)), x)`

**3.278**  $\int \frac{1}{(d \csc(a+bx))^{5/2}(c \sec(a+bx))^{5/2}} dx$

3.278.1 Optimal result . . . . . 1629  
 3.278.2 Mathematica [C] (verified) . . . . . 1629  
 3.278.3 Rubi [A] (verified) . . . . . 1630  
 3.278.4 Maple [B] (verified) . . . . . 1633  
 3.278.5 Fricas [F] . . . . . 1633  
 3.278.6 Sympy [F(-1)] . . . . . 1634  
 3.278.7 Maxima [F] . . . . . 1634  
 3.278.8 Giac [F] . . . . . 1634  
 3.278.9 Mupad [F(-1)] . . . . . 1635

**3.278.1 Optimal result**

Integrand size = 25, antiderivative size = 135

$$\int \frac{1}{(d \csc(a + bx))^{5/2}(c \sec(a + bx))^{5/2}} dx =$$

$$-\frac{c}{5bd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{7/2}} + \frac{1}{10bcd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}}$$

$$+ \frac{3E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{20bc^2d^2\sqrt{d \csc(a + bx)}\sqrt{c \sec(a + bx)}\sqrt{\sin(2a + 2bx)}}$$

output

```
-1/5*c/b/d/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(7/2)+1/10/b/c/d/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2)-3/20*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))/b/c^2/d^2/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)/sin(2*b*x+2*a)^(1/2)
```

**3.278.2 Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.45 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.67

$$\int \frac{1}{(d \csc(a + bx))^{5/2}(c \sec(a + bx))^{5/2}} dx = \frac{\left(-2 \cos^2(a + bx) \cos(2(a + bx)) + 3\sqrt[4]{-\cot^2(a + bx)} \operatorname{Hypergeometric}\right)}{20bc^3d(d \csc(a + bx))^{5/2}(c \sec(a + bx))^{5/2}}$$

input `Integrate[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(5/2)),x]`

output `((-2*Cos[a + b*x]^2*Cos[2*(a + b*x)] + 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(20*b*c^3*d*(d*Csc[a + b*x])^(3/2))`

### 3.278.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3107, 3042, 3108, 3042, 3110, 3042, 3052, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \sec(a + bx))^{5/2} (d \csc(a + bx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(c \sec(a + bx))^{5/2} (d \csc(a + bx))^{5/2}} dx \\
 & \quad \downarrow \text{3107} \\
 & \frac{3 \int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} dx}{10d^2} - \frac{c}{5bd(c \sec(a + bx))^{7/2} (d \csc(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} dx}{10d^2} - \frac{c}{5bd(c \sec(a + bx))^{7/2} (d \csc(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3108} \\
 & \frac{3 \left( \frac{\int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx}{2c^2} + \frac{d}{3bc(c \sec(a + bx))^{3/2} (d \csc(a + bx))^{3/2}} \right)}{10d_c^2} - \\
 & \quad \frac{c}{5bd(c \sec(a + bx))^{7/2} (d \csc(a + bx))^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.278.  $\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2}} dx$

$$\begin{aligned}
& \frac{3 \left( \frac{\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx}{2c^2} + \frac{d}{3bc(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}} \right)}{10d^2 c} \\
& \quad \frac{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}}{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3110} \\
& \frac{3 \left( \frac{\int \frac{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}}{2c^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)}} dx}{10d^2 c} + \frac{d}{3bc(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}} \right)}{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \left( \frac{\int \frac{\sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)}}{2c^2 \sqrt{c \cos(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \sqrt{d \csc(a+bx)}} dx}{10d^2 c} + \frac{d}{3bc(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}} \right)}{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3052} \\
& \frac{3 \left( \frac{\int \frac{\sqrt{\sin(2a+2bx)}}{2c^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} dx}{10d^2 c} + \frac{d}{3bc(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}} \right)}{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \left( \frac{\int \frac{\sqrt{\sin(2a+2bx)}}{2c^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} dx}{10d^2 c} + \frac{d}{3bc(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}} \right)}{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}} \\
& \quad \downarrow \text{3119} \\
& \frac{3 \left( \frac{E(a+bx - \frac{\pi}{4} | 2)}{2bc^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{d}{3bc(c \sec(a+bx))^{3/2} (d \csc(a+bx))^{3/2}} \right)}{10d^2 c} \\
& \quad \frac{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}}{5bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{3/2}}
\end{aligned}$$

input `Int[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(5/2)),x]`

output `-1/5*c/(b*d*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(7/2)) + (3*(d/(3*b*c*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2)) + EllipticE[a - Pi/4 + b*x, 2]/(2*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])))/(10*d^2)`

## 3.278.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3052 `Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Simp[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]) Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

rule 3107 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3108 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3110 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

**3.278.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 424 vs.  $2(140) = 280$ .

Time = 8.06 (sec) , antiderivative size = 425, normalized size of antiderivative = 3.15

method	result
default	$-\frac{\sqrt{2} \left( -4 \cos(bx+a)^6 \sqrt{2} + 6 \sqrt{1 + \csc(bx+a) - \cot(bx+a)} \sqrt{\cot(bx+a) - \csc(bx+a) + 1} \sqrt{\cot(bx+a) - \csc(bx+a)} \right) \text{EllipticE}\left(\sqrt{1 + \csc(bx+a) - \cot(bx+a)}\right)}{\dots}$

input `int(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$-1/40/b*2^{(1/2)}*(-4*\cos(b*x+a)^6*2^{(1/2)}+6*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticE}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*\cos(b*x+a)-3*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})*\cos(b*x+a)+6*\cos(b*x+a)^4*2^{(1/2)}+6*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticE}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})-3*(1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a)+1)^{(1/2)}*(\cot(b*x+a)-\csc(b*x+a))^{(1/2)}*\text{EllipticF}((1+\csc(b*x+a)-\cot(b*x+a))^{(1/2)},1/2*2^{(1/2)})+2^{(1/2)}*\cos(b*x+a)^2-3*2^{(1/2)}*\cos(b*x+a))/(d*csc(b*x+a))^{(1/2)}/(c*sec(b*x+a))^{(1/2)}/c^2/d^2*sec(b*x+a)*csc(b*x+a)$$

**3.278.5 Fracas [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fracas")`

output `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^3*d^3*csc(b*x + a)^3*sec(b*x + a)^3), x)`

**3.278.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate(1/(d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(5/2),x)`

output `Timed out`

**3.278.7 Maxima [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2)), x)`

**3.278.8 Giac [F]**

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2)), x)`

**3.278.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2} \left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

input `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2)),x)`output `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2)), x)`



**3.279**      $\int \frac{1}{(d \csc(a+bx))^{7/2}(c \sec(a+bx))^{5/2}} dx$

3.279.1 Optimal result . . . . . 1636  
 3.279.2 Mathematica [A] (verified) . . . . . 1637  
 3.279.3 Rubi [A] (verified) . . . . . 1637  
 3.279.4 Maple [A] (warning: unable to verify) . . . . . 1644  
 3.279.5 Fricas [C] (verification not implemented) . . . . . 1644  
 3.279.6 Sympy [F(-1)] . . . . . 1645  
 3.279.7 Maxima [F] . . . . . 1646  
 3.279.8 Giac [F] . . . . . 1646  
 3.279.9 Mupad [F(-1)] . . . . . 1646

**3.279.1 Optimal result**

Integrand size = 25, antiderivative size = 406

$$\int \frac{1}{(d \csc(a + bx))^{7/2}(c \sec(a + bx))^{5/2}} dx =$$

$$\frac{c}{6bd(d \csc(a + bx))^{5/2}(c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)}(c \sec(a + bx))^{7/2}}$$

$$+ \frac{5}{192bcd^3 \sqrt{d \csc(a + bx)}(c \sec(a + bx))^{3/2}}$$

$$- \frac{5 \arctan\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{128\sqrt{2}bc^2d^4 \sqrt{c \sec(a + bx)}}$$

$$+ \frac{5 \arctan\left(1 + \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{128\sqrt{2}bc^2d^4 \sqrt{c \sec(a + bx)}}$$

$$- \frac{5 \sqrt{d \csc(a + bx)} \log\left(1 - \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{256\sqrt{2}bc^2d^4 \sqrt{c \sec(a + bx)}}$$

$$+ \frac{5 \sqrt{d \csc(a + bx)} \log\left(1 + \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{\tan(a + bx)}}{256\sqrt{2}bc^2d^4 \sqrt{c \sec(a + bx)}}$$

output 
$$\begin{aligned} & -1/6*c/b/d/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(7/2)-5/48*c/b/d^3/(c*sec(b \\ & *x+a))^(7/2)/(d*csc(b*x+a))^(1/2)+5/192/b/c/d^3/(c*sec(b*x+a))^(3/2)/(d*cs \\ & c(b*x+a))^(1/2)+5/256*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^( \\ & 1/2)*tan(b*x+a)^(1/2)/b/c^2/d^4*2^(1/2)/(c*sec(b*x+a))^(1/2)+5/256*arctan( \\ & 1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2/d^ \\ & 4*2^(1/2)/(c*sec(b*x+a))^(1/2)-5/512*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x \\ & +a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/c^2/d^4*2^(1/2)/(c*sec(b*x+a) \\ & )^(1/2)+5/512*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/ \\ & 2)*tan(b*x+a)^(1/2)/b/c^2/d^4*2^(1/2)/(c*sec(b*x+a))^(1/2) \end{aligned}$$

### 3.279.2 Mathematica [A] (verified)

Time = 3.38 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.43

$$\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx = \frac{\left( 28 + 34 \cos(2(a + bx)) + 2 \cos(4(a + bx)) - 4 \cos(6(a + bx)) + 15\sqrt{2} \arctan\left(\frac{-1 + \sqrt{\cot^2(a + bx)}}{\sqrt{2} \sqrt[4]{\cot^2(a + bx)}}\right) \sqrt[4]{\cot^2(a + bx)} \right) \sqrt[4]{\cot^2(a + bx)}}{768bc^3d^3 \sqrt{d \csc(a + bx)}}$$

input `Integrate[1/((d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(5/2)),x]`

output 
$$\begin{aligned} & -1/768*((28 + 34*Cos[2*(a + b*x)] + 2*Cos[4*(a + b*x)] - 4*Cos[6*(a + b*x) \\ & ] + 15*sqrt[2]*ArcTan[(-1 + Sqrt[Cot[a + b*x]^2])/(Sqrt[2]*(Cot[a + b*x]^2 \\ & )^(1/4))]*(Cot[a + b*x]^2)^(1/4) - 15*sqrt[2]*ArcTanh[(Sqrt[2]*(Cot[a + b* \\ & x]^2)^(1/4))/(1 + Sqrt[Cot[a + b*x]^2])]*(Cot[a + b*x]^2)^(1/4))*Sqrt[c*Se \\ & c[a + b*x]])/(b*c^3*d^3*sqrt[d*Csc[a + b*x])) \end{aligned}$$

### 3.279.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 298, normalized size of antiderivative = 0.73, number of steps used = 20, number of rules used = 19,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.760$ , Rules used = {3042, 3107, 3042, 3107, 3042, 3108, 3042, 3109, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.279.  $\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{1}{(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{7/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{1}{(c \sec(a+bx))^{5/2} (d \csc(a+bx))^{7/2}} dx \\
& \quad \downarrow \text{3107} \\
& \frac{5 \int \frac{1}{(d \csc(a+bx))^{3/2} (c \sec(a+bx))^{5/2}} dx}{12d^2} - \frac{c}{6bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{5 \int \frac{1}{(d \csc(a+bx))^{3/2} (c \sec(a+bx))^{5/2}} dx}{12d^2} - \frac{c}{6bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{5/2}} \\
& \quad \downarrow \text{3107} \\
& \frac{5 \left( \frac{\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \right)}{12d^2} - \frac{c}{6bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{5 \left( \frac{\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx}{8d^2} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \right)}{12d^2} - \frac{c}{6bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{5/2}} \\
& \quad \downarrow \text{3108} \\
& \frac{5 \left( \frac{3 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4c^2} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \right)}{12d^2} - \frac{c}{6bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{5 \left( \frac{3 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4c^2} + \frac{d}{2bc(c \sec(a+bx))^{3/2} \sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2} \sqrt{d \csc(a+bx)}} \right)}{12d^2} - \frac{c}{6bd(c \sec(a+bx))^{7/2} (d \csc(a+bx))^{5/2}} \\
& \quad \downarrow \text{3109}
\end{aligned}$$

---

3.279.  $\int \frac{1}{(d \csc(a+bx))^{7/2} (c \sec(a+bx))^{5/2}} dx$

$$\begin{array}{c}
5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{4c^2\sqrt{c \sec(a+bx)}} + \frac{d}{2bc(c \sec(a+bx))^{3/2}\sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2}\sqrt{d \csc(a+bx)}} \right) \\
\hline
\frac{12d^2}{c} \\
\frac{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}{3042} \\
\downarrow 3042 \\
5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{4c^2\sqrt{c \sec(a+bx)}} + \frac{d}{2bc(c \sec(a+bx))^{3/2}\sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2}\sqrt{d \csc(a+bx)}} \right) \\
\hline
\frac{12d^2}{c} \\
\frac{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}{3957} \\
\downarrow 3957 \\
5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \int \frac{1}{\sqrt{\tan(a+bx)}(\tan^2(a+bx)+1)} d \tan(a+bx)}{4bc^2\sqrt{c \sec(a+bx)}} + \frac{d}{2bc(c \sec(a+bx))^{3/2}\sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2}\sqrt{d \csc(a+bx)}} \right) \\
\hline
\frac{c}{12d^2} \\
\frac{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}{266} \\
\downarrow 266 \\
5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \int \frac{1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)}}{2bc^2\sqrt{c \sec(a+bx)}} + \frac{d}{2bc(c \sec(a+bx))^{3/2}\sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2}\sqrt{d \csc(a+bx)}} \right) \\
\hline
\frac{12d^2}{c} \\
\frac{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}{755} \\
\downarrow 755 \\
5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \left( \frac{1}{2} \int \frac{1-\tan(a+bx)}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{\tan(a+bx)+1}{\tan^2(a+bx)+1} d \sqrt{\tan(a+bx)} \right)}{2bc^2\sqrt{c \sec(a+bx)}} + \frac{d}{2bc(c \sec(a+bx))^{3/2}\sqrt{d \csc(a+bx)}} - \frac{c}{4bd(c \sec(a+bx))^{7/2}\sqrt{d \csc(a+bx)}} \right) \\
\hline
\frac{c}{12d^2} \\
\frac{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}{1476} \\
\downarrow 1476
\end{array}$$


---

3.279.  $\int \frac{1}{(d \csc(a+bx))^{7/2}(c \sec(a+bx))^{5/2}} dx$

$$5 \left( \frac{3\sqrt{\tan(ax+bx)}\sqrt{d\csc(ax+bx)}\left(\frac{1}{2}\int\frac{1-\tan(ax+bx)}{\tan^2(ax+bx)+1}d\sqrt{\tan(ax+bx)}+\frac{1}{2}\left(\frac{1}{2}\int\frac{1}{\tan(ax+bx)-\sqrt{2}\sqrt{\tan(ax+bx)+1}}d\sqrt{\tan(ax+bx)}+\frac{1}{2}\int\frac{1}{\tan(ax+bx)+\sqrt{2}\sqrt{\tan(ax+bx)+1}}d\sqrt{\tan(ax+bx)}\right)\right)}{2bc^2\sqrt{c\sec(ax+bx)}} \right) \frac{1}{8d^2}$$

$$\frac{c}{6bd(c\sec(ax+bx))^{7/2}(d\csc(ax+bx))^{5/2}} \quad 12d^2$$

↓ 1082

$$5 \left( \frac{3\sqrt{\tan(ax+bx)}\sqrt{d\csc(ax+bx)}\left(\frac{1}{2}\int\frac{1-\tan(ax+bx)}{\tan^2(ax+bx)+1}d\sqrt{\tan(ax+bx)}+\frac{1}{2}\left(\frac{\int\frac{1}{-\tan(ax+bx)-1}d(1-\sqrt{2}\sqrt{\tan(ax+bx)})}{\sqrt{2}}-\frac{\int\frac{1}{-\tan(ax+bx)-1}d(\sqrt{2}\sqrt{\tan(ax+bx)+1})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(ax+bx)}} \right) \frac{1}{8d^2}$$

$$\frac{c}{6bd(c\sec(ax+bx))^{7/2}(d\csc(ax+bx))^{5/2}} \quad 12d^2$$

↓ 217

$$5 \left( \frac{3\sqrt{\tan(ax+bx)}\sqrt{d\csc(ax+bx)}\left(\frac{1}{2}\int\frac{1-\tan(ax+bx)}{\tan^2(ax+bx)+1}d\sqrt{\tan(ax+bx)}+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(ax+bx)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\tan(ax+bx)})}{\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(ax+bx)}} \right) \frac{1}{8d^2} + \frac{d}{2bc(c\sec(ax+bx))^{3/2}\sqrt{d\csc(ax+bx)}}$$

$$\frac{c}{6bd(c\sec(ax+bx))^{7/2}(d\csc(ax+bx))^{5/2}} \quad 12d^2$$

↓ 1479

$$5 \left( \frac{3\sqrt{\tan(ax+bx)}\sqrt{d\csc(ax+bx)}\left(\frac{1}{2}\left(-\frac{\int-\frac{\sqrt{2}-2\sqrt{\tan(ax+bx)}}{\tan(ax+bx)-\sqrt{2}\sqrt{\tan(ax+bx)+1}}d\sqrt{\tan(ax+bx)}}{2\sqrt{2}}-\frac{\int-\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(ax+bx)+1})}{\tan(ax+bx)+\sqrt{2}\sqrt{\tan(ax+bx)+1}}d\sqrt{\tan(ax+bx)}}{2\sqrt{2}}\right)\right)}{2bc^2\sqrt{c\sec(ax+bx)}} \right) \frac{1}{8d^2} + \frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(ax+bx)+1})}{\sqrt{2}}\right)$$

$$\frac{c}{6bd(c\sec(ax+bx))^{7/2}(d\csc(ax+bx))^{5/2}} \quad 12d^2$$

↓ 25

---

3.279.  $\int \frac{1}{(d\csc(ax+bx))^{7/2}(c\sec(ax+bx))^{5/2}} dx$

$$5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) \right)}{2bc^2\sqrt{c \sec(a+bx)}} \right) \frac{1}{8d^2}$$

$12d^2$

$$\frac{c}{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}$$

↓ 27

$$5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \left( \frac{1}{2} \left( \int \frac{\sqrt{2}-2\sqrt{\tan(a+bx)}}{\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(a+bx)+1}}{\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1}} d\sqrt{\tan(a+bx)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) \right)}{2bc^2\sqrt{c \sec(a+bx)}} \right) \frac{1}{8d^2}$$

$12d^2$

$$\frac{c}{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}$$

↓ 1103

$$5 \left( \frac{3\sqrt{\tan(a+bx)}\sqrt{d \csc(a+bx)} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\tan(a+bx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(a+bx)})}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\tan(a+bx)+\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} - \frac{\log(\tan(a+bx)-\sqrt{2}\sqrt{\tan(a+bx)+1})}{2\sqrt{2}} \right) \right)}{2bc^2\sqrt{c \sec(a+bx)}} \right) \frac{1}{8d^2}$$

$12d^2$

$$\frac{c}{6bd(c \sec(a+bx))^{7/2}(d \csc(a+bx))^{5/2}}$$

input `Int[1/((d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(5/2)),x]`

output `-1/6*c/(b*d*(d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(7/2)) + (5*(-1/4*c/(b*d*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(7/2)) + (d/(2*b*c*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)) + (3*Sqrt[d*Csc[a + b*x]]*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]/(2*Sqrt[2]))/2)*Sqrt[Tan[a + b*x]]/(2*b*c^2*Sqrt[c*Sec[a + b*x]]))/(8*d^2))/(12*d^2)`

## 3.279.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3107 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Simp[(m + 1)/(a^2*(m + n)) Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3108 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Simp[(n + 1)/(b^2*(m + n)) Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3109 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*Csc[e + f*x])^m*((b*Sec[e + f*x])^n/Tan[e + f*x]^n) Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`



### 3.279.4 Maple [A] (warning: unable to verify)

Time = 44.54 (sec) , antiderivative size = 613, normalized size of antiderivative = 1.51

method	result
default	$\frac{\sqrt{2} \left( 128 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sqrt{2} \cos(bx+a)^6 + 128 \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \sqrt{2} \cos(bx+a)^5 - 208 \sqrt{2} \sqrt{-\frac{\cos(bx+a) \sin(bx+a)}{(\cos(bx+a)+1)^2}} \cos(bx+a)^4 \right)}{\dots}$

input `int(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output

```

-1/1536/b*2^(1/2)*(128*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*2^(1/2)*cos(b*x+a)^6+128*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*2^(1/2)*cos(b*x+a)^5-208*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cos(b*x+a)^4-208*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cos(b*x+a)^3+20*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*2^(1/2)*cos(b*x+a)^2+20*cos(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)-30*arctan((sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+cos(b*x+a)-1)/(cos(b*x+a)-1))+30*arctan((-sin(b*x+a)*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)+cos(b*x+a)-1)/(cos(b*x+a)-1))+15*ln(-2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)-2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2)-15*ln(2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*cot(b*x+a)+2*2^(1/2)*(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)*csc(b*x+a)-2*cot(b*x+a)+2))/(cos(b*x+a)-1)/(cos(b*x+a)+1)^2/(-cos(b*x+a)*sin(b*x+a)/(cos(b*x+a)+1)^2)^(1/2)/(c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2)/c^2/d^3*sin(b*x+a)^2
    
```

### 3.279.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.65 (sec) , antiderivative size = 1409, normalized size of antiderivative = 3.47

$$\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x,algorithm="fracas")`

output 
$$-1/3072*(15*b*c^3*d^4*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\log(2*(b^3*c^7*d^{10}*(-1/(b^4*c^{10}*d^{14}))^{(3/4)}*\cos(b*x + a)^2*\sin(b*x + a) + (b*c^2*d^3*\cos(b*x + a))^3 - b*c^2*d^3*\cos(b*x + a))*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)} + 1) - 15*b*c^3*d^4*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\log(-2*(b^3*c^7*d^{10}*(-1/(b^4*c^{10}*d^{14}))^{(3/4)}*\cos(b*x + a)^2*\sin(b*x + a) + (b*c^2*d^3*\cos(b*x + a))^3 - b*c^2*d^3*\cos(b*x + a))*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)} + 1) + 15*I*b*c^3*d^4*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\log(-2*(I*b^3*c^7*d^{10}*(-1/(b^4*c^{10}*d^{14}))^{(3/4)}*\cos(b*x + a)^2*\sin(b*x + a) + (-I*b*c^2*d^3*\cos(b*x + a))^3 + I*b*c^2*d^3*\cos(b*x + a))*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)} + 1) - 15*I*b*c^3*d^4*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\log(-2*(-I*b^3*c^7*d^{10}*(-1/(b^4*c^{10}*d^{14}))^{(3/4)}*\cos(b*x + a)^2*\sin(b*x + a) + (I*b*c^2*d^3*\cos(b*x + a))^3 - I*b*c^2*d^3*\cos(b*x + a))*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)} + 1) + 15*b*c^3*d^4*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\log(1/2*(b*c^2*d^3*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\cos(b*x + a)^2*\sin(b*x + a) + (b^3*c^7*d^{10}*\cos(b*x + a))^3 - b^3*c^7*d^{10}*\cos(b*x + a))*(-1/(b^4*c^{10}*d^{14}))^{(3/4)}*\sqrt{c/\cos(b*x + a)}*\sqrt{d/\sin(b*x + a)} - 1/2*\cos(b*x + a)*\sin(b*x + a) - 1/4*(2*b^2*c^5*d^7*\cos(b*x + a)^2 - b^2*c^5*d^7)*\sqrt{-1/(b^4*c^{10}*d^{14}))} - 15*b*c^3*d^4*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\log(-1/2*(b*c^2*d^3*(-1/(b^4*c^{10}*d^{14}))^{(1/4)}*\cos(b*x + a)^2*\sin(b*...$$

### 3.279.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate(1/(d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(5/2),x)`

output `Timed out`

**3.279.7 Maxima [F]**

$$\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{7}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2)), x)`

**3.279.8 Giac [F]**

$$\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{(d \csc(bx + a))^{\frac{7}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

output `integrate(1/((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2)), x)`

**3.279.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx = \int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2} \left(\frac{d}{\sin(a+bx)}\right)^{7/2}} dx$$

input `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(7/2)),x)`

output `int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(7/2)), x)`

### 3.280 $\int \csc^n(e + fx) \sec^m(e + fx) dx$

3.280.1 Optimal result . . . . .	.1647
3.280.2 Mathematica [C] (warning: unable to verify) . . . . .	.1647
3.280.3 Rubi [A] (verified) . . . . .	.1648
3.280.4 Maple [F] . . . . .	.1649
3.280.5 Fracas [F] . . . . .	.1650
3.280.6 Sympy [F] . . . . .	.1650
3.280.7 Maxima [F] . . . . .	.1650
3.280.8 Giac [F] . . . . .	.1651
3.280.9 Mupad [F(-1)] . . . . .	.1651

#### 3.280.1 Optimal result

Integrand size = 17, antiderivative size = 81

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \frac{\cos^2(e + fx)^{\frac{1+m}{2}} \csc^{-1+n}(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1+m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) \sec^{1+m}(e + fx)}{f(1-n)}$$

output `(cos(f*x+e)^2)^(1/2+1/2*m)*csc(f*x+e)^(-1+n)*hypergeom([1/2+1/2*m, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*sec(f*x+e)^(1+m)/f/(1-n)`

#### 3.280.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 2.66 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.43

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \frac{(-3+n)}{f(-1+n) \left( (-3+n) \operatorname{AppellF1}\left(\frac{1}{2} - \frac{n}{2}, m, 1-m-n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - \right)}$$

input `Integrate[Csc[e + f*x]^n*Sec[e + f*x]^m,x]`

output `-(((−3 + n)*AppellF1[1/2 − n/2, m, 1 − m − n, 3/2 − n/2, Tan[(e + f*x)/2]^2, −Tan[(e + f*x)/2]^2]*Csc[e + f*x]^(−1 + n)*Sec[e + f*x]^m)/(f*(−1 + n)*((−3 + n)*AppellF1[1/2 − n/2, m, 1 − m − n, 3/2 − n/2, Tan[(e + f*x)/2]^2, −Tan[(e + f*x)/2]^2] − 2*((−1 + m + n)*AppellF1[3/2 − n/2, m, 2 − m − n, 5/2 − n/2, Tan[(e + f*x)/2]^2, −Tan[(e + f*x)/2]^2] + m*AppellF1[3/2 − n/2, 1 + m, 1 − m − n, 5/2 − n/2, Tan[(e + f*x)/2]^2, −Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))`

### 3.280.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^m(e + fx) \csc^n(e + fx) dx \\ & \quad \downarrow \text{3042} \\ & \int \sec(e + fx)^m \csc(e + fx)^n dx \\ & \quad \downarrow \text{3111} \\ & \cos^{m+1}(e + fx) \sec^{m+1}(e + fx) \sin^{n-1}(e + fx) \csc^{n-1}(e + fx) \int \cos^{-m}(e + fx) \sin^{-n}(e + fx) dx \\ & \quad \downarrow \text{3042} \\ & \cos^{m+1}(e + fx) \sec^{m+1}(e + fx) \sin^{n-1}(e + fx) \csc^{n-1}(e + fx) \int \cos(e + fx)^{-m} \sin(e + fx)^{-n} dx \\ & \quad \downarrow \text{3057} \\ & \frac{\cos^2(e + fx)^{\frac{m+1}{2}} \sec^{m+1}(e + fx) \csc^{n-1}(e + fx) \text{Hypergeometric2F1}\left(\frac{m+1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)} \end{aligned}$$

input `Int[Csc[e + f*x]^n*Sec[e + f*x]^m,x]`

```
output ((Cos[e + f*x]^2)^((1 + m)/2)*Csc[e + f*x]^(-1 + n)*Hypergeometric2F1[(1 +
m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sec[e + f*x]^(1 + m))/(f*(1 -
n))
```

### 3.280.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3057 Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^ (n_)*((a_.)*sin[(e_.) + (f_.)*(x_)] )^ (m
_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*Frac
Part[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]
```

```
rule 3111 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^ (m_)*((b_.)*sec[(e_.) + (f_.)*(x_)] )^ (n
_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(
n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*Sin[e
+ f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !
SimplerQ[-m, -n]
```

### 3.280.4 Maple [F]

$$\int \csc(fx + e)^n \sec(fx + e)^m dx$$

```
input int(csc(f*x+e)^n*sec(f*x+e)^m,x)
```

```
output int(csc(f*x+e)^n*sec(f*x+e)^m,x)
```

**3.280.5 Fracas [F]**

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \int \csc(fx + e)^n \sec(fx + e)^m dx$$

input `integrate(csc(f*x+e)^n*sec(f*x+e)^m,x, algorithm="fricas")`

output `integral(csc(f*x + e)^n*sec(f*x + e)^m, x)`

**3.280.6 Sympy [F]**

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \int \csc^n(e + fx) \sec^m(e + fx) dx$$

input `integrate(csc(f*x+e)**n*sec(f*x+e)**m,x)`

output `Integral(csc(e + f*x)**n*sec(e + f*x)**m, x)`

**3.280.7 Maxima [F]**

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \int \csc(fx + e)^n \sec(fx + e)^m dx$$

input `integrate(csc(f*x+e)^n*sec(f*x+e)^m,x, algorithm="maxima")`

output `integrate(csc(f*x + e)^n*sec(f*x + e)^m, x)`

**3.280.8 Giac [F]**

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \int \csc(fx + e)^n \sec(fx + e)^m dx$$

input `integrate(csc(f*x+e)^n*sec(f*x+e)^m,x, algorithm="giac")`

output `integrate(csc(f*x + e)^n*sec(f*x + e)^m, x)`

**3.280.9 Mupad [F(-1)]**

Timed out.

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \int \left( \frac{1}{\cos(e + fx)} \right)^m \left( \frac{1}{\sin(e + fx)} \right)^n dx$$

input `int((1/cos(e + f*x))^m*(1/sin(e + f*x))^n,x)`

output `int((1/cos(e + f*x))^m*(1/sin(e + f*x))^n, x)`



### 3.281 $\int \csc^n(e + fx)(a \sec(e + fx))^m dx$

3.281.1 Optimal result . . . . .	1652
3.281.2 Mathematica [C] (warning: unable to verify) . . . . .	1652
3.281.3 Rubi [A] (verified) . . . . .	1653
3.281.4 Maple [F] . . . . .	1654
3.281.5 Fracas [F] . . . . .	1655
3.281.6 Sympy [F] . . . . .	1655
3.281.7 Maxima [F] . . . . .	1655
3.281.8 Giac [F] . . . . .	1656
3.281.9 Mupad [F(-1)] . . . . .	1656

#### 3.281.1 Optimal result

Integrand size = 19, antiderivative size = 86

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \frac{\cos^2(e + fx)^{\frac{1+m}{2}} \csc^{-1+n}(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1+m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) (a \sec(e + fx))^{1+m}}{af(1-n)}$$

output `(cos(f*x+e)^2)^(1/2+1/2*m)*csc(f*x+e)^(-1+n)*hypergeom([1/2+1/2*m, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*(a*sec(f*x+e))^(1+m)/a/f/(1-n)`

#### 3.281.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.03 (sec) , antiderivative size = 280, normalized size of antiderivative = 3.26

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \frac{(-3+n) f(-1+n) \left( (-3+n) \operatorname{AppellF1}\left(\frac{1}{2} - \frac{n}{2}, m, 1-m-n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - \right)}{}$$

input `Integrate[Csc[e + f*x]^n*(a*Sec[e + f*x])^m,x]`

output  $-\left(\left(-3+n\right) \operatorname{AppellF1}\left[\frac{1}{2}-\frac{n}{2}, m, 1-m-n, \frac{3}{2}-\frac{n}{2}, \tan\left(\frac{e+f x}{2}\right)\right]^2, -\tan\left(\frac{e+f x}{2}\right)^2\right) \operatorname{Csc}\left[e+f x\right]^{-1+n}\left(a \operatorname{Sec}\left[e+f x\right]\right)^m / \left(f\left(-1+n\right)\left(-3+n\right) \operatorname{AppellF1}\left[\frac{1}{2}-\frac{n}{2}, m, 1-m-n, \frac{3}{2}-\frac{n}{2}, \tan\left(\frac{e+f x}{2}\right)\right]^2, -\tan\left(\frac{e+f x}{2}\right)^2\right)-2\left(-1+m+n\right) \operatorname{AppellF1}\left[\frac{3}{2}-\frac{n}{2}, m, 2-m-n, \frac{5}{2}-\frac{n}{2}, \tan\left(\frac{e+f x}{2}\right)\right]^2, -\tan\left(\frac{e+f x}{2}\right)^2\right)+m \operatorname{AppellF1}\left[\frac{3}{2}-\frac{n}{2}, 1+m, 1-m-n, \frac{5}{2}-\frac{n}{2}, \tan\left(\frac{e+f x}{2}\right)\right]^2, -\tan\left(\frac{e+f x}{2}\right)^2\right) \tan\left(\frac{e+f x}{2}\right)^2\right)$

### 3.281.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \csc^n(e+fx)(a \sec(e+fx))^m dx$$

$$\downarrow 3042$$

$$\int \csc(e+fx)^n(a \sec(e+fx))^m dx$$

$$\downarrow 3111$$

$$\frac{\sin^{n-1}(e+fx) \csc^{n-1}(e+fx)(a \cos(e+fx))^{m+1}(a \sec(e+fx))^{m+1} \int (a \cos(e+fx))^{-m} \sin^{-n}(e+fx) dx}{a^2}$$

$$\downarrow 3042$$

$$\frac{\sin^{n-1}(e+fx) \csc^{n-1}(e+fx)(a \cos(e+fx))^{m+1}(a \sec(e+fx))^{m+1} \int (a \cos(e+fx))^{-m} \sin(e+fx)^{-n} dx}{a^2}$$

$$\downarrow 3057$$

$$\frac{\cos^2(e+fx)^{\frac{m+1}{2}} \csc^{n-1}(e+fx)(a \sec(e+fx))^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{m+1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e+fx)\right)}{af(1-n)}$$

input  $\operatorname{Int}\left[\operatorname{Csc}\left[e+f x\right]^n\left(a \operatorname{Sec}\left[e+f x\right]\right)^m, x\right]$

output  $((\cos[e + f*x]^2)^{((1 + m)/2)} * \csc[e + f*x]^{-1 + n} * \text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 - n)/2, \sin[e + f*x]^2] * (a * \sec[e + f*x])^{(1 + m)}) / (a * f * (1 - n))$

### 3.281.3.1 Defintions of rubi rules used

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear } Q[u, x]$

rule 3057  $\text{Int}[(\cos[(e_.) + (f_.)(x_)] * (b_.))^{(n_)} * ((a_.) * \sin[(e_.) + (f_.)(x_)])^{(m_)}], x\_Symbol] \rightarrow \text{Simp}[b^{(2 * \text{IntPart}[(n - 1)/2] + 1)} * (b * \cos[e + f*x])^{(2 * \text{FracPart}[(n - 1)/2])} * ((a * \sin[e + f*x])^{(m + 1)} / (a * f * (m + 1) * (\cos[e + f*x]^2)^{\text{FracPart}[(n - 1)/2]}) * \text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \sin[e + f*x]^2], x] \text{ ; FreeQ}\{a, b, e, f, m, n\}, x]$

rule 3111  $\text{Int}[(\csc[(e_.) + (f_.)(x_)] * (a_.))^{(m_)} * ((b_.) * \sec[(e_.) + (f_.)(x_)])^{(n_)}], x\_Symbol] \rightarrow \text{Simp}[(a^2/b^2) * (a * \csc[e + f*x])^{(m - 1)} * (b * \sec[e + f*x])^{(n + 1)} * (a * \sin[e + f*x])^{(m - 1)} * (b * \cos[e + f*x])^{(n + 1)} \text{ Int}[1 / ((a * \sin[e + f*x])^m * (b * \cos[e + f*x])^n), x], x] \text{ ; FreeQ}\{a, b, e, f, m, n\}, x] \&\amp; ! \text{SimplerQ}[-m, -n]$

### 3.281.4 Maple [F]

$$\int \csc(fx + e)^n (a \sec(fx + e))^m dx$$

input  $\text{int}(\csc(f*x+e)^n * (a * \sec(f*x+e))^m, x)$

output  $\text{int}(\csc(f*x+e)^n * (a * \sec(f*x+e))^m, x)$

**3.281.5 Fracas [F]**

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \int (a \sec(fx + e))^m \csc(fx + e)^n dx$$

input `integrate(csc(f*x+e)^n*(a*sec(f*x+e))^m,x, algorithm="fricas")`

output `integral((a*sec(f*x + e))^m*csc(f*x + e)^n, x)`

**3.281.6 Sympy [F]**

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \int (a \sec(e + fx))^m \csc^n(e + fx) dx$$

input `integrate(csc(f*x+e)**n*(a*sec(f*x+e))**m,x)`

output `Integral((a*sec(e + f*x))**m*csc(e + f*x)**n, x)`

**3.281.7 Maxima [F]**

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \int (a \sec(fx + e))^m \csc(fx + e)^n dx$$

input `integrate(csc(f*x+e)^n*(a*sec(f*x+e))^m,x, algorithm="maxima")`

output `integrate((a*sec(f*x + e))^m*csc(f*x + e)^n, x)`

**3.281.8 Giac [F]**

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \int (a \sec(fx + e))^m \csc(fx + e)^n dx$$

input `integrate(csc(f*x+e)^n*(a*sec(f*x+e))^m,x, algorithm="giac")`

output `integrate((a*sec(f*x + e))^m*csc(f*x + e)^n, x)`

**3.281.9 Mupad [F(-1)]**

Timed out.

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \int \left( \frac{a}{\cos(e + fx)} \right)^m \left( \frac{1}{\sin(e + fx)} \right)^n dx$$

input `int((a/cos(e + f*x))^m*(1/sin(e + f*x))^n,x)`

output `int((a/cos(e + f*x))^m*(1/sin(e + f*x))^n, x)`

### 3.282 $\int (b \csc(e + fx))^n \sec^m(e + fx) dx$

3.282.1 Optimal result . . . . .	1657
3.282.2 Mathematica [C] (warning: unable to verify) . . . . .	1657
3.282.3 Rubi [A] (verified) . . . . .	1658
3.282.4 Maple [F] . . . . .	1659
3.282.5 Fracas [F] . . . . .	1660
3.282.6 Sympy [F] . . . . .	1660
3.282.7 Maxima [F] . . . . .	1660
3.282.8 Giac [F] . . . . .	1661
3.282.9 Mupad [F(-1)] . . . . .	1661

#### 3.282.1 Optimal result

Integrand size = 19, antiderivative size = 84

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \frac{b \cos^2(e + fx)^{\frac{1+m}{2}} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{1+m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) \sec^{1+m}(e + fx)}{f(1-n)}$$

```
output b*(cos(f*x+e)^2)^(1/2+1/2*m)*(b*csc(f*x+e))^(1-n)*hypergeom([1/2+1/2*m, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*sec(f*x+e)^(1+m)/f/(1-n)
```

#### 3.282.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.04 (sec) , antiderivative size = 281, normalized size of antiderivative = 3.35

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \frac{b(-3+n) \text{AppellF1}\left(\frac{1-n}{2}, m, 1-m-n, \frac{3-n}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - 2 \left(\dots\right)}{f(-1+n)}$$

```
input Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^m,x]
```

output  $-\left((b*(-3+n)*\text{AppellF1}[(1-n)/2, m, 1-m-n, (3-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2]*(b*\text{Csc}[e+fx])^{(-1+n)}*\text{Sec}[e+fx]^m)/(f*(-1+n)*((-3+n)*\text{AppellF1}[(1-n)/2, m, 1-m-n, (3-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2) - 2*((-1+m+n)*\text{AppellF1}[(3-n)/2, m, 2-m-n, (5-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2) + m*\text{AppellF1}[(3-n)/2, 1+m, 1-m-n, (5-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2))*\text{Tan}[(e+fx)/2]^2\right)$

### 3.282.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^m(e+fx)(b \csc(e+fx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int \sec(e+fx)^m (b \csc(e+fx))^n dx \\ & \quad \downarrow \text{3111} \\ & b^2 \cos^{m+1}(e+fx) \sec^{m+1}(e+fx) (b \sin(e+fx))^{n-1} (b \csc(e+fx))^{n-1} \int \cos^{-m}(e+fx) (b \sin(e+fx))^{-n} dx \\ & \quad \downarrow \text{3042} \\ & b^2 \cos^{m+1}(e+fx) \sec^{m+1}(e+fx) (b \sin(e+fx))^{n-1} (b \csc(e+fx))^{n-1} \int \cos(e+fx)^{-m} (b \sin(e+fx))^{-n} dx \\ & \quad \downarrow \text{3057} \\ & \frac{b \cos^2(e+fx)^{\frac{m+1}{2}} \sec^{m+1}(e+fx) (b \csc(e+fx))^{n-1} \text{Hypergeometric2F1}\left(\frac{m+1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e+fx)\right)}{f(1-n)} \end{aligned}$$

input  $\text{Int}[(b*\text{Csc}[e+fx])^n*\text{Sec}[e+fx]^m, x]$

```
output (b*(Cos[e + f*x]^2)^((1 + m)/2)*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F
1[(1 + m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sec[e + f*x]^(1 + m))/(
f*(1 - n))
```

### 3.282.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3057 Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^ (n_)*((a_.)*sin[(e_.) + (f_.)*(x_)] )^ (m
_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*Frac
Part[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]
```

```
rule 3111 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^ (m_)*((b_.)*sec[(e_.) + (f_.)*(x_)] )^ (n
_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(
n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*Sin[e
+ f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !
SimplerQ[-m, -n]
```

### 3.282.4 Maple [F]

$$\int (b \csc(fx + e))^n \sec(fx + e)^m dx$$

```
input int((b*csc(f*x+e))^n*sec(f*x+e)^m,x)
```

```
output int((b*csc(f*x+e))^n*sec(f*x+e)^m,x)
```



**3.282.5 Fracas [F]**

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^m dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^m,x, algorithm="fricas")`

output `integral((b*csc(f*x + e))^n*sec(f*x + e)^m, x)`

**3.282.6 Sympy [F]**

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \int (b \csc(e + fx))^n \sec^m(e + fx) dx$$

input `integrate((b*csc(f*x+e))**n*sec(f*x+e)**m,x)`

output `Integral((b*csc(e + f*x))**n*sec(e + f*x)**m, x)`

**3.282.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^m dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^m,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^m, x)`

**3.282.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^m dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^m,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^m, x)`

**3.282.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \int \left( \frac{b}{\sin(e + fx)} \right)^n \left( \frac{1}{\cos(e + fx)} \right)^m dx$$

input `int((b/sin(e + f*x))^n*(1/cos(e + f*x))^m,x)`

output `int((b/sin(e + f*x))^n*(1/cos(e + f*x))^m, x)`

### 3.283 $\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx$

3.283.1 Optimal result . . . . .	1662
3.283.2 Mathematica [C] (warning: unable to verify) . . . . .	1662
3.283.3 Rubi [A] (verified) . . . . .	1663
3.283.4 Maple [F] . . . . .	1664
3.283.5 Fracas [F] . . . . .	1665
3.283.6 Sympy [F] . . . . .	1665
3.283.7 Maxima [F] . . . . .	1665
3.283.8 Giac [F] . . . . .	1666
3.283.9 Mupad [F(-1)] . . . . .	1666

#### 3.283.1 Optimal result

Integrand size = 21, antiderivative size = 89

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \frac{b \cos^2(e + fx)^{\frac{1+m}{2}} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{1+m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) (a \sec(e + fx))^{1+m}}{af(1-n)}$$

```
output b*(cos(f*x+e)^2)^(1/2+1/2*m)*(b*csc(f*x+e))^(1-n)*hypergeom([1/2+1/2*m, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*(a*sec(f*x+e))^(1+m)/a/f/(1-n)
```

#### 3.283.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.22 (sec) , antiderivative size = 283, normalized size of antiderivative = 3.18

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \frac{b(-3+n) \text{AppellF1}\left(\frac{1-n}{2}, m, 1-m-n, \frac{3-n}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - 2\left(\dots\right)}{f(-1+n) \left(\dots\right)}$$

```
input Integrate[(b*Csc[e + f*x])^n*(a*Sec[e + f*x])^m,x]
```

output  $-\left((b*(-3+n)*\text{AppellF1}[(1-n)/2, m, 1-m-n, (3-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2]*(b*\text{Csc}[e+fx])^{(-1+n)}*(a*\text{Sec}[e+fx])^m)/(f*(-1+n)*((-3+n)*\text{AppellF1}[(1-n)/2, m, 1-m-n, (3-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2] - 2*((-1+m+n)*\text{AppellF1}[(3-n)/2, m, 2-m-n, (5-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2] + m*\text{AppellF1}[(3-n)/2, 1+m, 1-m-n, (5-n)/2, \text{Tan}[(e+fx)/2]^2, -\text{Tan}[(e+fx)/2]^2])* \text{Tan}[(e+fx)/2]^2)\right)$

### 3.283.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sec(e+fx))^m (b \csc(e+fx))^n dx$$

$$\downarrow 3042$$

$$\int (a \sec(e+fx))^m (b \csc(e+fx))^n dx$$

$$\downarrow 3111$$

$$\frac{b^2 (a \cos(e+fx))^{m+1} (a \sec(e+fx))^{m+1} (b \sin(e+fx))^{n-1} (b \csc(e+fx))^{n-1} \int (a \cos(e+fx))^{-m} (b \sin(e+fx))^{-n} dx}{a^2}$$

$$\downarrow 3042$$

$$\frac{b^2 (a \cos(e+fx))^{m+1} (a \sec(e+fx))^{m+1} (b \sin(e+fx))^{n-1} (b \csc(e+fx))^{n-1} \int (a \cos(e+fx))^{-m} (b \sin(e+fx))^{-n} dx}{a^2}$$

$$\downarrow 3057$$

$$\frac{b \cos^2(e+fx)^{\frac{m+1}{2}} (a \sec(e+fx))^{m+1} (b \csc(e+fx))^{n-1} \text{Hypergeometric2F1}\left(\frac{m+1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e+fx)\right)}{af(1-n)}$$

input  $\text{Int}[(b*\text{Csc}[e+fx])^n*(a*\text{Sec}[e+fx])^m,x]$

output  $(b \cdot (\cos[e + f \cdot x]^2)^{((1 + m)/2)} \cdot (b \cdot \csc[e + f \cdot x])^{-1 + n} \cdot \text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 - n)/2, \sin[e + f \cdot x]^2] \cdot (a \cdot \sec[e + f \cdot x])^{(1 + m)}) / (a \cdot f \cdot (1 - n))$

### 3.283.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_) + (f_)*(x_)]*(b_.))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*SIn[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_) + (f_)*(x_)]*(a_.))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*SIn[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*SIn[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

### 3.283.4 Maple [F]

$$\int (b \csc(fx + e))^n (a \sec(fx + e))^m dx$$

input `int((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x)`

output `int((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x)`

**3.283.5 Fracas [F]**

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \int (b \csc(fx + e))^n (a \sec(fx + e))^m dx$$

input `integrate((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="fricas")`

output `integral((b*csc(f*x + e))^n*(a*sec(f*x + e))^m, x)`

**3.283.6 Sympy [F]**

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \int (a \sec(e + fx))^m (b \csc(e + fx))^n dx$$

input `integrate((b*csc(f*x+e))**n*(a*sec(f*x+e))**m,x)`

output `Integral((a*sec(e + f*x))**m*(b*csc(e + f*x))**n, x)`

**3.283.7 Maxima [F]**

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \int (b \csc(fx + e))^n (a \sec(fx + e))^m dx$$

input `integrate((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*(a*sec(f*x + e))^m, x)`

**3.283.8 Giac [F]**

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \int (b \csc(fx + e))^n (a \sec(fx + e))^m dx$$

input `integrate((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*(a*sec(f*x + e))^m, x)`

**3.283.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \int \left( \frac{a}{\cos(e + fx)} \right)^m \left( \frac{b}{\sin(e + fx)} \right)^n dx$$

input `int((a/cos(e + f*x))^m*(b/sin(e + f*x))^n,x)`

output `int((a/cos(e + f*x))^m*(b/sin(e + f*x))^n, x)`

### 3.284 $\int (b \csc(e + fx))^n \sec^5(e + fx) dx$

3.284.1 Optimal result . . . . .	1667
3.284.2 Mathematica [A] (verified) . . . . .	1667
3.284.3 Rubi [A] (verified) . . . . .	1668
3.284.4 Maple [F] . . . . .	1669
3.284.5 Fracas [F] . . . . .	1670
3.284.6 Sympy [F] . . . . .	1670
3.284.7 Maxima [F] . . . . .	1670
3.284.8 Giac [F] . . . . .	1671
3.284.9 Mupad [F(-1)] . . . . .	1671

#### 3.284.1 Optimal result

Integrand size = 19, antiderivative size = 48

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = \frac{(b \csc(e + fx))^{5+n} \operatorname{Hypergeometric2F1}\left(3, \frac{5+n}{2}, \frac{7+n}{2}, \csc^2(e + fx)\right)}{b^5 f(5+n)}$$

output  $(b*\csc(f*x+e))^{(5+n)}*\operatorname{hypergeom}([3, 5/2+1/2*n], [7/2+1/2*n], \csc(f*x+e)^2)/b^5/f/(5+n)$

#### 3.284.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = -\frac{b(b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(3, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(-1+n)}$$

input `Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^5,x]`

output  $-((b*(b*Csc[e + f*x])^{(-1+n)}*\operatorname{Hypergeometric2F1}[3, (1-n)/2, (3-n)/2, \operatorname{Sin}[e + f*x]^2])/(f*(-1+n)))$



**3.284.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 3101, 25, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^5(e + fx)(b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sec(e + fx)^5 (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3101} \\
 & - \frac{\int -\frac{b^6 (b \csc(e + fx))^{n+4}}{(b^2 - b^2 \csc^2(e + fx))^3} d(b \csc(e + fx))}{b^5 f} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{b^6 (b \csc(e + fx))^{n+4}}{(b^2 - b^2 \csc^2(e + fx))^3} d(b \csc(e + fx))}{b^5 f} \\
 & \quad \downarrow \text{27} \\
 & \frac{b \int \frac{(b \csc(e + fx))^{n+4}}{(b^2 - b^2 \csc^2(e + fx))^3} d(b \csc(e + fx))}{f} \\
 & \quad \downarrow \text{278} \\
 & \frac{(b \csc(e + fx))^{n+5} \text{Hypergeometric2F1}\left(3, \frac{n+5}{2}, \frac{n+7}{2}, \csc^2(e + fx)\right)}{b^5 f(n + 5)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^5,x]`

output `((b*Csc[e + f*x])^(5 + n)*Hypergeometric2F1[3, (5 + n)/2, (7 + n)/2, Csc[e + f*x]^2])/(b^5*f*(5 + n))`

## 3.284.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 278 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3101 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Simp[-(f*a^n)^(-1) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

## 3.284.4 Maple [F]

$$\int (b \csc(fx + e))^n \sec(fx + e)^5 dx$$

input `int((b*csc(f*x+e))^n*sec(f*x+e)^5,x)`

output `int((b*csc(f*x+e))^n*sec(f*x+e)^5,x)`

**3.284.5 Fracas [F]**

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^5 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^5,x, algorithm="fricas")`

output `integral((b*csc(f*x + e))^n*sec(f*x + e)^5, x)`

**3.284.6 Sympy [F]**

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = \int (b \csc(e + fx))^n \sec^5(e + fx) dx$$

input `integrate((b*csc(f*x+e))**n*sec(f*x+e)**5,x)`

output `Integral((b*csc(e + f*x))**n*sec(e + f*x)**5, x)`

**3.284.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^5 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^5,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^5, x)`

**3.284.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^5 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^5,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^5, x)`

**3.284.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = \int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^5} dx$$

input `int((b/sin(e + f*x))^n/cos(e + f*x)^5,x)`

output `int((b/sin(e + f*x))^n/cos(e + f*x)^5, x)`

### 3.285 $\int (b \csc(e + fx))^n \sec^3(e + fx) dx$

3.285.1 Optimal result . . . . .	1672
3.285.2 Mathematica [A] (verified) . . . . .	1672
3.285.3 Rubi [A] (verified) . . . . .	1673
3.285.4 Maple [F] . . . . .	1674
3.285.5 Fracas [F] . . . . .	1674
3.285.6 Sympy [F] . . . . .	1675
3.285.7 Maxima [F] . . . . .	1675
3.285.8 Giac [F] . . . . .	1675
3.285.9 Mupad [F(-1)] . . . . .	1676

#### 3.285.1 Optimal result

Integrand size = 19, antiderivative size = 49

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx$$

$$= -\frac{(b \csc(e + fx))^{3+n} \operatorname{Hypergeometric2F1}\left(2, \frac{3+n}{2}, \frac{5+n}{2}, \csc^2(e + fx)\right)}{b^3 f(3+n)}$$

output `-(b*csc(f*x+e))^(3+n)*hypergeom([2, 3/2+1/2*n], [5/2+1/2*n], csc(f*x+e)^2)/b^3/f/(3+n)`

#### 3.285.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.04

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx$$

$$= -\frac{b(b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(2, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(-1+n)}$$

input `Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^3,x]`

output `-((b*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(-1 + n)))`

**3.285.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3101, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(e+fx)(b \csc(e+fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sec(e+fx)^3 (b \csc(e+fx))^n dx \\
 & \quad \downarrow \text{3101} \\
 & - \frac{\int \frac{b^4 (b \csc(e+fx))^{n+2}}{(b^2 - b^2 \csc^2(e+fx))^2} d(b \csc(e+fx))}{b^3 f} \\
 & \quad \downarrow \text{27} \\
 & - \frac{b \int \frac{(b \csc(e+fx))^{n+2}}{(b^2 - b^2 \csc^2(e+fx))^2} d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{278} \\
 & - \frac{(b \csc(e+fx))^{n+3} \text{Hypergeometric2F1}\left(2, \frac{n+3}{2}, \frac{n+5}{2}, \csc^2(e+fx)\right)}{b^3 f(n+3)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^3,x]`

output `-(((b*Csc[e + f*x])^(3 + n)*Hypergeometric2F1[2, (3 + n)/2, (5 + n)/2, Csc[e + f*x]^2])/(b^3*f*(3 + n)))`

## 3.285.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 278 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3101 `Int[(csc[(e_) + (f_)*(x_)]*(a_.))^m_*sec[(e_) + (f_)*(x_)]^n_, x_Symbol] := Simp[-(f*a^n)^(-1) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

## 3.285.4 Maple [F]

$$\int (b \csc(fx + e))^n \sec(fx + e)^3 dx$$

input `int((b*csc(f*x+e))^n*sec(f*x+e)^3,x)`

output `int((b*csc(f*x+e))^n*sec(f*x+e)^3,x)`

## 3.285.5 Fracas [F]

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^3 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^3,x, algorithm="fricas")`

output `integral((b*csc(f*x + e))^n*sec(f*x + e)^3, x)`

**3.285.6 Sympy [F]**

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx = \int (b \csc(e + fx))^n \sec^3(e + fx) dx$$

input `integrate((b*csc(f*x+e))**n*sec(f*x+e)**3,x)`

output `Integral((b*csc(e + f*x))**n*sec(e + f*x)**3, x)`

**3.285.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^3 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^3,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^3, x)`

**3.285.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^3 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^3,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^3, x)`



**3.285.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx = \int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^3} dx$$

input `int((b/sin(e + f*x))^n/cos(e + f*x)^3,x)`output `int((b/sin(e + f*x))^n/cos(e + f*x)^3, x)`

### 3.286 $\int (b \csc(e + fx))^n \sec(e + fx) dx$

3.286.1 Optimal result . . . . .	1677
3.286.2 Mathematica [A] (verified) . . . . .	1677
3.286.3 Rubi [A] (verified) . . . . .	1678
3.286.4 Maple [F] . . . . .	1679
3.286.5 Fricas [F] . . . . .	1680
3.286.6 Sympy [F] . . . . .	1680
3.286.7 Maxima [F] . . . . .	1680
3.286.8 Giac [F] . . . . .	1681
3.286.9 Mupad [F(-1)] . . . . .	1681

#### 3.286.1 Optimal result

Integrand size = 17, antiderivative size = 48

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = \frac{(b \csc(e + fx))^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \csc^2(e + fx)\right)}{bf(1+n)}$$

output  $(b*\csc(f*x+e))^{(1+n)}*\operatorname{hypergeom}([1, 1/2+1/2*n], [3/2+1/2*n], \csc(f*x+e)^2)/b/f/(1+n)$

#### 3.286.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = -\frac{b(b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(-1+n)}$$

input  $\operatorname{Integrate}[(b*\operatorname{Csc}[e + f*x])^n*\operatorname{Sec}[e + f*x], x]$

output  $-((b*(b*\operatorname{Csc}[e + f*x])^{-1+n}*\operatorname{Hypergeometric2F1}[1, (1-n)/2, (3-n)/2, \operatorname{Sin}[e + f*x]^2])/(f*(-1+n)))$

**3.286.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {3042, 3101, 25, 27, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sec(e + fx)(b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3101} \\
 & - \frac{\int -\frac{b^2(b \csc(e+fx))^n}{b^2 - b^2 \csc^2(e+fx)} d(b \csc(e + fx))}{bf} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{b^2(b \csc(e+fx))^n}{b^2 - b^2 \csc^2(e+fx)} d(b \csc(e + fx))}{bf} \\
 & \quad \downarrow \text{27} \\
 & \frac{b \int \frac{(b \csc(e+fx))^n}{b^2 - b^2 \csc^2(e+fx)} d(b \csc(e + fx))}{f} \\
 & \quad \downarrow \text{278} \\
 & \frac{(b \csc(e + fx))^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, \csc^2(e + fx)\right)}{bf(n+1)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*Sec[e + f*x],x]`

output `((b*Csc[e + f*x])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Csc[e + f*x]^2])/(b*f*(1 + n))`

## 3.286.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 278 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3101 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Simp[-(f*a^n)^(-1) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

## 3.286.4 Maple [F]

$$\int (b \csc(fx + e))^n \sec(fx + e) dx$$

input `int((b*csc(f*x+e))^n*sec(f*x+e),x)`

output `int((b*csc(f*x+e))^n*sec(f*x+e),x)`

**3.286.5 Fracas [F]**

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e) dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e),x, algorithm="fricas")`

output `integral((b*csc(f*x + e))^n*sec(f*x + e), x)`

**3.286.6 Sympy [F]**

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = \int (b \csc(e + fx))^n \sec(e + fx) dx$$

input `integrate((b*csc(f*x+e))**n*sec(f*x+e),x)`

output `Integral((b*csc(e + f*x))**n*sec(e + f*x), x)`

**3.286.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e) dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e),x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e), x)`

**3.286.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e) dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e),x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e), x)`

**3.286.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = \int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e + fx)} dx$$

input `int((b/sin(e + f*x))^n/cos(e + f*x),x)`

output `int((b/sin(e + f*x))^n/cos(e + f*x), x)`

### 3.287 $\int \cos(e + fx)(b \csc(e + fx))^n dx$

3.287.1 Optimal result . . . . .	1682
3.287.2 Mathematica [A] (verified) . . . . .	1682
3.287.3 Rubi [A] (verified) . . . . .	1683
3.287.4 Maple [A] (verified) . . . . .	1684
3.287.5 Fracas [A] (verification not implemented) . . . . .	1684
3.287.6 Sympy [F] . . . . .	1685
3.287.7 Maxima [A] (verification not implemented) . . . . .	1685
3.287.8 Giac [F] . . . . .	1685
3.287.9 Mupad [B] (verification not implemented) . . . . .	1686

#### 3.287.1 Optimal result

Integrand size = 17, antiderivative size = 24

$$\int \cos(e + fx)(b \csc(e + fx))^n dx = \frac{b(b \csc(e + fx))^{-1+n}}{f(1-n)}$$

output `b*(b*csc(f*x+e))(-1+n)/f/(1-n)`

#### 3.287.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \cos(e + fx)(b \csc(e + fx))^n dx = -\frac{b(b \csc(e + fx))^{-1+n}}{f(-1+n)}$$

input `Integrate[Cos[e + f*x]*(b*Csc[e + f*x])n,x]`

output `-((b*(b*Csc[e + f*x])(-1 + n))/(f*(-1 + n)))`

**3.287.3 Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {3042, 3101, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \cos(e + fx)(b \csc(e + fx))^n dx \\
 \downarrow \text{3042} \\
 \int \frac{(b \csc(e + fx))^n}{\sec(e + fx)} dx \\
 \downarrow \text{3101} \\
 -\frac{b \int (b \csc(e + fx))^{n-2} d(b \csc(e + fx))}{f} \\
 \downarrow \text{15} \\
 \frac{b(b \csc(e + fx))^{n-1}}{f(1-n)}
 \end{array}$$

input `Int[Cos[e + f*x]*(b*Csc[e + f*x])^n,x]`

output `(b*(b*Csc[e + f*x])^(-1 + n))/(f*(1 - n))`

**3.287.3.1 Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



```
rule 3101 Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol]
:> Simp[-(f*a^n)^(-1) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

### 3.287.4 Maple [A] (verified)

Time = 1.43 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.29

method	result
derivativedivides	$-\frac{e^{n \ln(b \csc(fx+e))}}{f(-1+n) \csc(fx+e)}$
default	$-\frac{e^{n \ln(b \csc(fx+e))}}{f(-1+n) \csc(fx+e)}$
parallelrisch	$-\frac{2^{-n} \left( b \sec\left(\frac{fx}{2} + \frac{e}{2}\right) \csc\left(\frac{fx}{2} + \frac{e}{2}\right) \right)^n \sin(fx+e)}{f(-1+n)}$
norman	$-\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) e^{n \ln\left(\frac{b \left(1 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2}{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}\right)}}{f(-1+n) \left(1 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2}$
risch	$-\frac{(e^{2i(fx+e)} - 1)^{-n} \sin(fx+e) b^n 2^n (e^{i(fx+e)})^n e^{i\pi n \left(-\operatorname{csgn}\left(\frac{ie^{i(fx+e)}}{e^{2i(fx+e)} - 1}\right) + \operatorname{csgn}\left(\frac{ie^{i(fx+e)}}{e^{2i(fx+e)} - 1}\right)\right)^2 \operatorname{csgn}(ie^{i(fx+e)}) + \operatorname{csgn}(ie^{i(fx+e)})}}{f(-1+n)}$

```
input int(cos(f*x+e)*(b*csc(f*x+e))^n,x,method=_RETURNVERBOSE)
```

```
output -1/f/(-1+n)*exp(n*ln(b*csc(f*x+e)))/csc(f*x+e)
```

### 3.287.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \cos(e + fx)(b \csc(e + fx))^n dx = -\frac{\left(\frac{b}{\sin(fx+e)}\right)^n \sin(fx + e)}{fn - f}$$

```
input integrate(cos(f*x+e)*(b*csc(f*x+e))^n,x, algorithm="fricas")
```

```
output -(b/sin(f*x + e))^n*sin(f*x + e)/(f*n - f)
```

**3.287.6 Sympy [F]**

$$\int \cos(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(e + fx))^n \cos(e + fx) dx$$

input `integrate(cos(f*x+e)*(b*csc(f*x+e))**n,x)`

output `Integral((b*csc(e + f*x))**n*cos(e + f*x), x)`

**3.287.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \cos(e + fx)(b \csc(e + fx))^n dx = -\frac{b^n \sin(fx + e)^{-n} \sin(fx + e)}{f(n - 1)}$$

input `integrate(cos(f*x+e)*(b*csc(f*x+e))^n,x, algorithm="maxima")`

output `-b^n*sin(f*x + e)^(-n)*sin(f*x + e)/(f*(n - 1))`

**3.287.8 Giac [F]**

$$\int \cos(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e) dx$$

input `integrate(cos(f*x+e)*(b*csc(f*x+e))^n,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*cos(f*x + e), x)`

**3.287.9 Mupad [B] (verification not implemented)**

Time = 13.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \cos(e + fx)(b \csc(e + fx))^n dx = -\frac{\sin(e + fx) \left(\frac{b}{\sin(e + fx)}\right)^n}{f(n - 1)}$$

input `int(cos(e + f*x)*(b/sin(e + f*x))^n,x)`output `-(sin(e + f*x)*(b/sin(e + f*x))^n)/(f*(n - 1))`

### 3.288 $\int \cos^3(e + fx)(b \csc(e + fx))^n dx$

3.288.1 Optimal result . . . . .	1687
3.288.2 Mathematica [A] (verified) . . . . .	1687
3.288.3 Rubi [A] (verified) . . . . .	1688
3.288.4 Maple [A] (verified) . . . . .	1689
3.288.5 Fricas [A] (verification not implemented) . . . . .	1690
3.288.6 Sympy [F(-1)] . . . . .	1690
3.288.7 Maxima [A] (verification not implemented) . . . . .	1690
3.288.8 Giac [F] . . . . .	1691
3.288.9 Mupad [B] (verification not implemented) . . . . .	1691

#### 3.288.1 Optimal result

Integrand size = 19, antiderivative size = 52

$$\int \cos^3(e + fx)(b \csc(e + fx))^n dx = -\frac{b^3(b \csc(e + fx))^{-3+n}}{f(3-n)} + \frac{b(b \csc(e + fx))^{-1+n}}{f(1-n)}$$

output `-b^3*(b*csc(f*x+e))^(3-n)/f/(3-n)+b*(b*csc(f*x+e))^(1-n)/f/(1-n)`

#### 3.288.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.87

$$\int \cos^3(e + fx)(b \csc(e + fx))^n dx \\ = -\frac{b(-5 + n + (-1 + n) \cos(2(e + fx)))(b \csc(e + fx))^{-1+n}}{2f(-3 + n)(-1 + n)}$$

input `Integrate[Cos[e + f*x]^3*(b*Csc[e + f*x])^n,x]`

output `-1/2*(b*(-5 + n + (-1 + n)*Cos[2*(e + f*x)])*(b*Csc[e + f*x])^(1-n))/(f*(-3 + n)*(-1 + n))`

**3.288.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {3042, 3101, 25, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(e+fx)(b \csc(e+fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(e+fx))^n}{\sec(e+fx)^3} dx \\
 & \quad \downarrow \text{3101} \\
 & - \frac{b^3 \int -\frac{(b \csc(e+fx))^{n-4}(b^2-b^2 \csc^2(e+fx))}{b^2} d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{25} \\
 & \frac{b^3 \int \frac{(b \csc(e+fx))^{n-4}(b^2-b^2 \csc^2(e+fx))}{b^2} d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{b \int (b \csc(e+fx))^{n-4} (b^2 - b^2 \csc^2(e+fx)) d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{244} \\
 & \frac{b \int (b^2 (b \csc(e+fx))^{n-4} - (b \csc(e+fx))^{n-2}) d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{b \left( \frac{(b \csc(e+fx))^{n-1}}{1-n} - \frac{b^2 (b \csc(e+fx))^{n-3}}{3-n} \right)}{f}
 \end{aligned}$$

input `Int[Cos[e + f*x]^3*(b*Csc[e + f*x])^n,x]`

output `(b*(-((b^2*(b*Csc[e + f*x])^(-3 + n))/(3 - n)) + (b*Csc[e + f*x])^(-1 + n)/(1 - n)))/f`

## 3.288.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 244 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3101 `Int[(csc[(e_) + (f_)*(x_)])*(a_)^(m_)*sec[(e_) + (f_)*(x_)^(n_)], x_Symbol] := Simp[-(f*a^n)^(-1) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

## 3.288.4 Maple [A] (verified)

Time = 7.05 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.29

method	result	size
parallelrisch	$-\frac{((-1+n)\sin(3fx+3e)+\sin(fx+e)(n-9))(b\sec\left(\frac{fx}{2}+\frac{e}{2}\right)\csc\left(\frac{fx}{2}+\frac{e}{2}\right))^n 2^{-n}}{4f(n^2-4n+3)}$	67

input `int(cos(f*x+e)^3*(b*csc(f*x+e))^n,x,method=_RETURNVERBOSE)`

output `-1/4*((-1+n)*sin(3*f*x+3*e)+sin(f*x+e)*(n-9))*(b*sec(1/2*f*x+1/2*e)*csc(1/2*f*x+1/2*e))^n*2^(-n)/f/(n^2-4*n+3)`

**3.288.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.94

$$\int \cos^3(e + fx)(b \csc(e + fx))^n dx = -\frac{((n - 1) \cos^2(fx + e) - 2) \left(\frac{b}{\sin(fx + e)}\right)^n \sin(fx + e)}{fn^2 - 4fn + 3f}$$

input `integrate(cos(f*x+e)^3*(b*csc(f*x+e))^n,x, algorithm="fricas")`output `-(n - 1)*cos(f*x + e)^2 - 2)*(b/sin(f*x + e))^n*sin(f*x + e)/(f*n^2 - 4*f*n + 3*f)`**3.288.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^3(e + fx)(b \csc(e + fx))^n dx = \text{Timed out}$$

input `integrate(cos(f*x+e)**3*(b*csc(f*x+e))**n,x)`output `Timed out`**3.288.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.12

$$\int \cos^3(e + fx)(b \csc(e + fx))^n dx = \frac{\frac{b^n \sin(fx+e)^{-n} \sin(fx+e)^3}{n-3} - \frac{b^n \sin(fx+e)^{-n} \sin(fx+e)}{n-1}}{f}$$

input `integrate(cos(f*x+e)^3*(b*csc(f*x+e))^n,x, algorithm="maxima")`output `(b^n*sin(f*x + e)^(-n)*sin(f*x + e)^3/(n - 3) - b^n*sin(f*x + e)^(-n)*sin(f*x + e)/(n - 1))/f`

**3.288.8 Giac [F]**

$$\int \cos^3(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^3 dx$$

input `integrate(cos(f*x+e)^3*(b*csc(f*x+e))^n,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*cos(f*x + e)^3, x)`

**3.288.9 Mupad [B] (verification not implemented)**

Time = 13.56 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.27

$$\int \cos^3(e + fx)(b \csc(e + fx))^n dx$$

$$= \frac{\left(\frac{b}{\sin(e+fx)}\right)^n (9 \sin(e + fx) + \sin(3e + 3fx) - n \sin(e + fx) - n \sin(3e + 3fx))}{4f(n^2 - 4n + 3)}$$

input `int(cos(e + f*x)^3*(b/sin(e + f*x))^n,x)`

output `((b/sin(e + f*x))^n*(9*sin(e + f*x) + sin(3*e + 3*f*x) - n*sin(e + f*x) - n*sin(3*e + 3*f*x)))/(4*f*(n^2 - 4*n + 3))`



### 3.289 $\int \cos^5(e + fx)(b \csc(e + fx))^n dx$

3.289.1 Optimal result . . . . .	1692
3.289.2 Mathematica [A] (verified) . . . . .	1692
3.289.3 Rubi [A] (verified) . . . . .	1693
3.289.4 Maple [A] (verified) . . . . .	1694
3.289.5 Fricas [A] (verification not implemented) . . . . .	1695
3.289.6 Sympy [F(-1)] . . . . .	1695
3.289.7 Maxima [A] (verification not implemented) . . . . .	1695
3.289.8 Giac [F] . . . . .	1696
3.289.9 Mupad [B] (verification not implemented) . . . . .	1696

#### 3.289.1 Optimal result

Integrand size = 19, antiderivative size = 78

$$\int \cos^5(e + fx)(b \csc(e + fx))^n dx = \frac{b^5(b \csc(e + fx))^{-5+n}}{f(5 - n)} - \frac{2b^3(b \csc(e + fx))^{-3+n}}{f(3 - n)} + \frac{b(b \csc(e + fx))^{-1+n}}{f(1 - n)}$$

```
output b^5*(b*csc(f*x+e))^(5-n)/f/(5-n)-2*b^3*(b*csc(f*x+e))^(3-n)/f/(3-n)+b*(b*csc(f*x+e))^(1-n)/f/(1-n)
```

#### 3.289.2 Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.04

$$\int \cos^5(e + fx)(b \csc(e + fx))^n dx = \frac{(b \csc(e + fx))^n (3 - 4n + n^2 - 2(5 - 6n + n^2) \csc^2(e + fx) + (15 - 8n + n^2) \csc^4(e + fx)) \sin^5(e + fx)}{f(-5 + n)(-3 + n)(-1 + n)}$$

```
input Integrate[Cos[e + f*x]^5*(b*Csc[e + f*x])^n,x]
```

```
output -(((b*Csc[e + f*x])^n*(3 - 4*n + n^2 - 2*(5 - 6*n + n^2)*Csc[e + f*x]^2 + (15 - 8*n + n^2)*Csc[e + f*x]^4)*Sin[e + f*x]^5)/(f*(-5 + n)*(-3 + n)*(-1 + n)))
```

**3.289.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3042, 3101, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^5(e+fx)(b \csc(e+fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(e+fx))^n}{\sec(e+fx)^5} dx \\
 & \quad \downarrow \text{3101} \\
 & \frac{b^5 \int \frac{(b \csc(e+fx))^{n-6} (b^2 - b^2 \csc^2(e+fx))^2}{b^4} d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{b \int (b \csc(e+fx))^{n-6} (b^2 - b^2 \csc^2(e+fx))^2 d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{244} \\
 & \frac{b \int (b^4 (b \csc(e+fx))^{n-6} - 2b^2 (b \csc(e+fx))^{n-4} + (b \csc(e+fx))^{n-2}) d(b \csc(e+fx))}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{b \left( -\frac{b^4 (b \csc(e+fx))^{n-5}}{5-n} + \frac{2b^2 (b \csc(e+fx))^{n-3}}{3-n} - \frac{(b \csc(e+fx))^{n-1}}{1-n} \right)}{f}
 \end{aligned}$$

input `Int[Cos[e + f*x]^5*(b*Csc[e + f*x])^n,x]`

output `-((b*(-((b^4*(b*Csc[e + f*x])^(-5 + n))/(5 - n)) + (2*b^2*(b*Csc[e + f*x])^(-3 + n))/(3 - n) - (b*Csc[e + f*x])^(-1 + n)/(1 - n)))/f)`

## 3.289.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 244 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3101 `Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Simp[-(f*a^n)^(-1) Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

## 3.289.4 Maple [A] (verified)

Time = 5.01 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.33

method	result
parallelrisch	$-\frac{\left(\frac{3}{2}n^2-14n+\frac{25}{2}\right)\sin(3fx+3e)+\left(\frac{1}{2}n^2-2n+\frac{3}{2}\right)\sin(5fx+5e)+\sin(fx+e)(n^2-12n+75)}{8(-1+n)(-3+n)(-5+n)f}2^{-n}\left(b\sec\left(\frac{fx}{2}+\frac{e}{2}\right)\csc\left(\frac{fx}{2}+\frac{e}{2}\right)\right)^n$

input `int(cos(f*x+e)^5*(b*csc(f*x+e))^n,x,method=_RETURNVERBOSE)`

output `-1/8*((3/2*n^2-14*n+25/2)*sin(3*f*x+3*e)+(1/2*n^2-2*n+3/2)*sin(5*f*x+5*e)+sin(f*x+e)*(n^2-12*n+75))*2^(-n)*(b*sec(1/2*f*x+1/2*e)*csc(1/2*f*x+1/2*e))^n/(-1+n)/(-3+n)/(-5+n)/f`

**3.289.5 Fricas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.94

$$\int \cos^5(e + fx)(b \csc(e + fx))^n dx$$

$$= -\frac{((n^2 - 4n + 3) \cos(fx + e)^4 - 4(n - 1) \cos(fx + e)^2 + 8) \left(\frac{b}{\sin(fx + e)}\right)^n \sin(fx + e)}{fn^3 - 9fn^2 + 23fn - 15f}$$

input `integrate(cos(f*x+e)^5*(b*csc(f*x+e))^n,x, algorithm="fricas")`output `-((n^2 - 4*n + 3)*cos(f*x + e)^4 - 4*(n - 1)*cos(f*x + e)^2 + 8)*(b/sin(f*x + e))^n*sin(f*x + e)/(f*n^3 - 9*f*n^2 + 23*f*n - 15*f)`**3.289.6 Sympy [F(-1)]**

Timed out.

$$\int \cos^5(e + fx)(b \csc(e + fx))^n dx = \text{Timed out}$$

input `integrate(cos(f*x+e)**5*(b*csc(f*x+e))**n,x)`output `Timed out`**3.289.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.10

$$\int \cos^5(e + fx)(b \csc(e + fx))^n dx$$

$$= -\frac{\frac{b^n \sin(fx+e)^{-n} \sin(fx+e)^5}{n-5} - \frac{2b^n \sin(fx+e)^{-n} \sin(fx+e)^3}{n-3} + \frac{b^n \sin(fx+e)^{-n} \sin(fx+e)}{n-1}}{f}$$

input `integrate(cos(f*x+e)^5*(b*csc(f*x+e))^n,x, algorithm="maxima")`output `-(b^n*sin(f*x + e)^(-n)*sin(f*x + e)^5/(n - 5) - 2*b^n*sin(f*x + e)^(-n)*sin(f*x + e)^3/(n - 3) + b^n*sin(f*x + e)^(-n)*sin(f*x + e)/(n - 1))/f`

---

3.289.  $\int \cos^5(e + fx)(b \csc(e + fx))^n dx$

**3.289.8 Giac [F]**

$$\int \cos^5(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^5 dx$$

input `integrate(cos(f*x+e)^5*(b*csc(f*x+e))^n,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*cos(f*x + e)^5, x)`

**3.289.9 Mupad [B] (verification not implemented)**

Time = 15.07 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.72

$$\int \cos^5(e + fx)(b \csc(e + fx))^n dx = \frac{\left(\frac{b}{\sin(e+fx)}\right)^n (150 \sin(e + fx) + 25 \sin(3e + 3fx) + 3 \sin(5e + 5fx) + 3n^2 \sin(3e + 3fx) + n^2 \sin(5e + 5fx) - 24n \sin(e + fx) - 28n \sin(3e + 3fx) - 4n \sin(5e + 5fx) + 2n^2 \sin(e + fx))}{16f(n^3 - 9n^2)}$$

input `int(cos(e + f*x)^5*(b/sin(e + f*x))^n,x)`

output `-((b/sin(e + f*x))^n*(150*sin(e + f*x) + 25*sin(3*e + 3*f*x) + 3*sin(5*e + 5*f*x) + 3*n^2*sin(3*e + 3*f*x) + n^2*sin(5*e + 5*f*x) - 24*n*sin(e + f*x) - 28*n*sin(3*e + 3*f*x) - 4*n*sin(5*e + 5*f*x) + 2*n^2*sin(e + f*x)))/(16*f*(23*n - 9*n^2 + n^3 - 15))`

### 3.290 $\int (b \csc(e + fx))^n \sec^6(e + fx) dx$

3.290.1 Optimal result . . . . .	1697
3.290.2 Mathematica [A] (verified) . . . . .	1697
3.290.3 Rubi [A] (verified) . . . . .	1698
3.290.4 Maple [F] . . . . .	1699
3.290.5 Fricas [F] . . . . .	1699
3.290.6 Sympy [F(-1)] . . . . .	1700
3.290.7 Maxima [F] . . . . .	1700
3.290.8 Giac [F] . . . . .	1700
3.290.9 Mupad [F(-1)] . . . . .	1701

#### 3.290.1 Optimal result

Integrand size = 19, antiderivative size = 72

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx$$

$$= \frac{b \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(\frac{7}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) \sec(e + fx)}{f(1-n)}$$

```
output b*(b*csc(f*x+e))^( -1+n)*hypergeom([7/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*sec(f*x+e)*(cos(f*x+e)^2)^(1/2)/f/(1-n)
```

#### 3.290.2 Mathematica [A] (verified)

Time = 1.88 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.07

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx$$

$$= \frac{(b \csc(e + fx))^n \operatorname{Hypergeometric2F1}\left(-2 - \frac{n}{2}, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2(e + fx)\right) \sec^2(e + fx)^{-n/2} \tan(e + fx)}{f(1-n)}$$

```
input Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^6,x]
```

```
output ((b*Csc[e + f*x])^n*Hypergeometric2F1[-2 - n/2, 1/2 - n/2, 3/2 - n/2, -Tan[e + f*x]^2]*Tan[e + f*x])/(f*(1 - n)*(Sec[e + f*x]^2)^(n/2))
```

**3.290.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^6(e + fx)(b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sec(e + fx)^6 (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3111} \\
 & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \sec^6(e + fx) (b \sin(e + fx))^{-n} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \frac{(b \sin(e + fx))^{-n}}{\cos(e + fx)^6} dx \\
 & \quad \downarrow \text{3057} \\
 & \frac{b \sqrt{\cos^2(e + fx)} \sec(e + fx) (b \csc(e + fx))^{n-1} \operatorname{Hypergeometric2F1}\left(\frac{7}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^6,x]`

output `(b*Sqrt[Cos[e + f*x]^2]*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[7/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sec[e + f*x])/(f*(1 - n))`

**3.290.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*SIn[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*SIn[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*SIn[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && ! SimplifierQ[-m, -n]`

### 3.290.4 Maple [F]

$$\int (b \csc(fx + e))^n \sec(fx + e)^6 dx$$

input `int((b*csc(f*x+e))^n*sec(f*x+e)^6,x)`

output `int((b*csc(f*x+e))^n*sec(f*x+e)^6,x)`

### 3.290.5 Fracas [F]

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^6 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^6,x, algorithm="fracas")`

output `integral((b*csc(f*x + e))^n*sec(f*x + e)^6, x)`



**3.290.6 Sympy [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx = \text{Timed out}$$

input `integrate((b*csc(f*x+e))**n*sec(f*x+e)**6,x)`output `Timed out`**3.290.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^6 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^6,x, algorithm="maxima")`output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^6, x)`**3.290.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^6 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^6,x, algorithm="giac")`output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^6, x)`

**3.290.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx = \int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^6} dx$$

input `int((b/sin(e + f*x))^n/cos(e + f*x)^6,x)`output `int((b/sin(e + f*x))^n/cos(e + f*x)^6, x)`

### 3.291 $\int (b \csc(e + fx))^n \sec^4(e + fx) dx$

3.291.1 Optimal result . . . . .	1702
3.291.2 Mathematica [A] (verified) . . . . .	1702
3.291.3 Rubi [A] (verified) . . . . .	1703
3.291.4 Maple [F] . . . . .	1704
3.291.5 Fricas [F] . . . . .	1704
3.291.6 Sympy [F] . . . . .	1705
3.291.7 Maxima [F] . . . . .	1705
3.291.8 Giac [F] . . . . .	1705
3.291.9 Mupad [F(-1)] . . . . .	1706

#### 3.291.1 Optimal result

Integrand size = 19, antiderivative size = 72

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx$$

$$= \frac{b \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) \sec(e + fx)}{f(1-n)}$$

output `b*(b*csc(f*x+e))(-1+n)*hypergeom([5/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)2)*sec(f*x+e)*(cos(f*x+e)2)(1/2)/f/(1-n)`

#### 3.291.2 Mathematica [A] (verified)

Time = 1.53 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.07

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx$$

$$= \frac{(b \csc(e + fx))^n \operatorname{Hypergeometric2F1}\left(-1 - \frac{n}{2}, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2(e + fx)\right) \sec^2(e + fx)^{-n/2} \tan(e + fx)}{f(1-n)}$$

input `Integrate[(b*Csc[e + f*x])n*Sec[e + f*x]4,x]`

output `((b*Csc[e + f*x])n*Hypergeometric2F1[-1 - n/2, 1/2 - n/2, 3/2 - n/2, -Tan[e + f*x]2]*Tan[e + f*x])/(f*(1 - n)*(Sec[e + f*x]2)(n/2))`

**3.291.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(e + fx)(b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sec(e + fx)^4 (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3111} \\
 & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \sec^4(e + fx) (b \sin(e + fx))^{-n} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \frac{(b \sin(e + fx))^{-n}}{\cos(e + fx)^4} dx \\
 & \quad \downarrow \text{3057} \\
 & \frac{b \sqrt{\cos^2(e + fx)} \sec(e + fx) (b \csc(e + fx))^{n-1} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^4,x]`

output `(b*Sqrt[Cos[e + f*x]^2]*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[5/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sec[e + f*x])/(f*(1 - n))`

**3.291.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*SIN[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*SIN[e + f*x])^(m - 1)*(b*COS[e + f*x])^(n + 1) Int[1/((a*SIN[e + f*x])^m*(b*COS[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

### 3.291.4 Maple [F]

$$\int (b \csc(fx + e))^n \sec(fx + e)^4 dx$$

input `int((b*csc(f*x+e))^n*sec(f*x+e)^4,x)`

output `int((b*csc(f*x+e))^n*sec(f*x+e)^4,x)`

### 3.291.5 Fracas [F]

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^4 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^4,x, algorithm="fricas")`

output `integral((b*csc(f*x + e))^n*sec(f*x + e)^4, x)`

**3.291.6 Sympy [F]**

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx = \int (b \csc(e + fx))^n \sec^4(e + fx) dx$$

input `integrate((b*csc(f*x+e))**n*sec(f*x+e)**4,x)`

output `Integral((b*csc(e + f*x))**n*sec(e + f*x)**4, x)`

**3.291.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^4 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^4,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^4, x)`

**3.291.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^4 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^4,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^4, x)`

**3.291.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx = \int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^4} dx$$

input `int((b/sin(e + f*x))^n/cos(e + f*x)^4,x)`output `int((b/sin(e + f*x))^n/cos(e + f*x)^4, x)`

### 3.292 $\int (b \csc(e + fx))^n \sec^2(e + fx) dx$

3.292.1 Optimal result . . . . .	1707
3.292.2 Mathematica [A] (verified) . . . . .	1707
3.292.3 Rubi [A] (verified) . . . . .	1708
3.292.4 Maple [F] . . . . .	1709
3.292.5 Fricas [F] . . . . .	1709
3.292.6 Sympy [F] . . . . .	1710
3.292.7 Maxima [F] . . . . .	1710
3.292.8 Giac [F] . . . . .	1710
3.292.9 Mupad [F(-1)] . . . . .	1711

#### 3.292.1 Optimal result

Integrand size = 19, antiderivative size = 72

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \frac{b \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) \sec(e + fx)}{f(1-n)}$$

```
output b*(b*csc(f*x+e))^(−1+n)*hypergeom([3/2, 1/2−1/2*n], [3/2−1/2*n], sin(f*x+e)^2)*sec(f*x+e)*(cos(f*x+e)^2)^(1/2)/f/(1−n)
```

#### 3.292.2 Mathematica [A] (verified)

Time = 2.65 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.36

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \frac{2(b \csc(e + fx))^n \left( -\frac{n \operatorname{Hypergeometric2F1}\left(1-n, \frac{1}{2}-\frac{n}{2}, \frac{3}{2}-\frac{n}{2}, -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) \sec^2\left(\frac{1}{2}(e+fx)\right)^{-n} \tan\left(\frac{1}{2}(e+fx)\right)}{-1+n} + \frac{1}{2} \tan(e + fx) \right)}{f}$$

```
input Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^2,x]
```

```
output (2*(b*Csc[e + f*x])^n*(-((n*Hypergeometric2F1[1 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2]*Tan[(e + f*x)/2])/((-1 + n)*(Sec[(e + f*x)/2]^2)^n) + Tan[e + f*x]/2))/f
```



**3.292.3 Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(e + fx)(b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sec(e + fx)^2 (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3111} \\
 & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \sec^2(e + fx) (b \sin(e + fx))^{-n} dx \\
 & \quad \downarrow \text{3042} \\
 & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \frac{(b \sin(e + fx))^{-n}}{\cos(e + fx)^2} dx \\
 & \quad \downarrow \text{3057} \\
 & \frac{b \sqrt{\cos^2(e + fx)} \sec(e + fx) (b \csc(e + fx))^{n-1} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^2,x]`

output `(b*Sqrt[Cos[e + f*x]^2]*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[3/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sec[e + f*x])/(f*(1 - n))`

**3.292.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3057 Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*Frac
Part[(n - 1)/2])*((a*SIn[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]
```

```
rule 3111 Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] :> Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n
+ 1)*(a*SIn[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*SIn[e
+ f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !
SimplerQ[-m, -n]
```

### 3.292.4 Maple [F]

$$\int (b \csc(fx + e))^n \sec(fx + e)^2 dx$$

```
input int((b*csc(f*x+e))^n*sec(f*x+e)^2,x)
```

```
output int((b*csc(f*x+e))^n*sec(f*x+e)^2,x)
```

### 3.292.5 Fracas [F]

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^2 dx$$

```
input integrate((b*csc(f*x+e))^n*sec(f*x+e)^2,x, algorithm="fracas")
```

```
output integral((b*csc(f*x + e))^n*sec(f*x + e)^2, x)
```

**3.292.6 Sympy [F]**

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \int (b \csc(e + fx))^n \sec^2(e + fx) dx$$

input `integrate((b*csc(f*x+e))**n*sec(f*x+e)**2,x)`

output `Integral((b*csc(e + f*x))**n*sec(e + f*x)**2, x)`

**3.292.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^2 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^2,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^2, x)`

**3.292.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \int (b \csc(fx + e))^n \sec(fx + e)^2 dx$$

input `integrate((b*csc(f*x+e))^n*sec(f*x+e)^2,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*sec(f*x + e)^2, x)`

**3.292.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^2} dx$$

input `int((b/sin(e + f*x))^n/cos(e + f*x)^2,x)`output `int((b/sin(e + f*x))^n/cos(e + f*x)^2, x)`

### 3.293 $\int (b \csc(e + fx))^n dx$

3.293.1 Optimal result . . . . .	1712
3.293.2 Mathematica [A] (verified) . . . . .	1712
3.293.3 Rubi [A] (verified) . . . . .	1713
3.293.4 Maple [F] . . . . .	1714
3.293.5 Fricas [F] . . . . .	1714
3.293.6 Sympy [F] . . . . .	1715
3.293.7 Maxima [F] . . . . .	1715
3.293.8 Giac [F] . . . . .	1715
3.293.9 Mupad [F(-1)] . . . . .	1716

#### 3.293.1 Optimal result

Integrand size = 10, antiderivative size = 72

$$\int (b \csc(e + fx))^n dx = \frac{b \cos(e + fx) (b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

output `b*cos(f*x+e)*(b*csc(f*x+e))(-1+n)*hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)2)/f/(1-n)/(cos(f*x+e)2)(1/2)`

#### 3.293.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.90

$$\int (b \csc(e + fx))^n dx = \frac{\cos(e + fx) (b \csc(e + fx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3}{2}, \cos^2(e + fx)\right) \sin(e + fx) \sin^2(e + fx)^{\frac{1}{2}(-1+n)}}{f}$$

input `Integrate[(b*Csc[e + f*x])n,x]`

output `-((Cos[e + f*x]*(b*Csc[e + f*x])n*Hypergeometric2F1[1/2, (1 + n)/2, 3/2, Cos[e + f*x]2]*Sin[e + f*x]*(Sin[e + f*x]2)((-1 + n)/2))/f)`

**3.293.3 Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{4259} \\
 & \left(\frac{\sin(e + fx)}{b}\right)^n (b \csc(e + fx))^n \int \left(\frac{\sin(e + fx)}{b}\right)^{-n} dx \\
 & \quad \downarrow \text{3042} \\
 & \left(\frac{\sin(e + fx)}{b}\right)^n (b \csc(e + fx))^n \int \left(\frac{\sin(e + fx)}{b}\right)^{-n} dx \\
 & \quad \downarrow \text{3122} \\
 & \frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n,x]`

output `(b*cos[e + f*x]*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(1 - n)*Sqrt[Cos[e + f*x]^2])`

## 3.293.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## 3.293.4 Maple [F]

$$\int (b \csc (fx + e))^n dx$$

input `int((b*csc(f*x+e))^n,x)`

output `int((b*csc(f*x+e))^n,x)`

## 3.293.5 Fracas [F]

$$\int (b \csc (e + fx))^n dx = \int (b \csc (fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n,x, algorithm="fracas")`

output `integral((b*csc(f*x + e))^n, x)`

**3.293.6 Sympy [F]**

$$\int (b \csc(e + fx))^n dx = \int (b \csc(e + fx))^n dx$$

input `integrate((b*csc(f*x+e))**n,x)`

output `Integral((b*csc(e + f*x))**n, x)`

**3.293.7 Maxima [F]**

$$\int (b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n, x)`

**3.293.8 Giac [F]**

$$\int (b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n, x)`



**3.293.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n dx = \int \left( \frac{b}{\sin(e + fx)} \right)^n dx$$

input `int((b/sin(e + f*x))^n,x)`output `int((b/sin(e + f*x))^n, x)`

### 3.294 $\int \cos^2(e + fx)(b \csc(e + fx))^n dx$

3.294.1 Optimal result . . . . .	1717
3.294.2 Mathematica [B] (verified) . . . . .	1717
3.294.3 Rubi [A] (verified) . . . . .	1718
3.294.4 Maple [F] . . . . .	1719
3.294.5 Fracas [F] . . . . .	1719
3.294.6 Sympy [F] . . . . .	1720
3.294.7 Maxima [F] . . . . .	1720
3.294.8 Giac [F] . . . . .	1720
3.294.9 Mupad [F(-1)] . . . . .	1721

#### 3.294.1 Optimal result

Integrand size = 19, antiderivative size = 72

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \frac{b \cos(e + fx)(b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

```
output b*cos(f*x+e)*(b*csc(f*x+e))^(n-1)*hypergeom([-1/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)/f/(1-n)/(cos(f*x+e)^2)^(1/2)
```

#### 3.294.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 165 vs. 2(72) = 144.

Time = 0.90 (sec) , antiderivative size = 165, normalized size of antiderivative = 2.29

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \frac{2(b \csc(e + fx))^n \left(\operatorname{Hypergeometric2F1}\left(1 - n, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 4 \operatorname{Hypergeometric2F1}\left(1 - n, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)\right)}{f}$$

```
input Integrate[Cos[e + f*x]^2*(b*Csc[e + f*x])^n,x]
```

output  $(-2*(b*\text{Csc}[e + f*x])^n*(\text{Hypergeometric2F1}[1 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2] - 4*\text{Hypergeometric2F1}[2 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2] + 4*\text{Hypergeometric2F1}[3 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2])* \text{Tan}[(e + f*x)/2]) / (f*(-1 + n)*( \text{Sec}[(e + f*x)/2]^2)^n)$

### 3.294.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(e + fx)(b \csc(e + fx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(b \csc(e + fx))^n}{\sec(e + fx)^2} dx \\ & \quad \downarrow \text{3111} \\ & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \cos^2(e + fx)(b \sin(e + fx))^{-n} dx \\ & \quad \downarrow \text{3042} \\ & b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \cos(e + fx)^2 (b \sin(e + fx))^{-n} dx \\ & \quad \downarrow \text{3057} \\ & \frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} \text{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}} \end{aligned}$$

input  $\text{Int}[\text{Cos}[e + f*x]^2*(b*\text{Csc}[e + f*x])^n, x]$

output  $(b*\text{Cos}[e + f*x]*(b*\text{Csc}[e + f*x])^{-(1 + n)}*\text{Hypergeometric2F1}[-1/2, (1 - n)/2, (3 - n)/2, \text{Sin}[e + f*x]^2]) / (f*(1 - n)*\text{Sqrt}[\text{Cos}[e + f*x]^2])$

## 3.294.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

## 3.294.4 Maple [F]

$$\int \cos(fx + e)^2 (b \csc(fx + e))^n dx$$

input `int(cos(f*x+e)^2*(b*csc(f*x+e))^n,x)`

output `int(cos(f*x+e)^2*(b*csc(f*x+e))^n,x)`

## 3.294.5 Fracas [F]

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^2 dx$$

input `integrate(cos(f*x+e)^2*(b*csc(f*x+e))^n,x, algorithm="fracas")`

output `integral((b*csc(f*x + e))^n*cos(f*x + e)^2, x)`

**3.294.6 Sympy [F]**

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(e + fx))^n \cos^2(e + fx) dx$$

input `integrate(cos(f*x+e)**2*(b*csc(f*x+e))**n,x)`

output `Integral((b*csc(e + f*x))**n*cos(e + f*x)**2, x)`

**3.294.7 Maxima [F]**

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^2 dx$$

input `integrate(cos(f*x+e)^2*(b*csc(f*x+e))^n,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*cos(f*x + e)^2, x)`

**3.294.8 Giac [F]**

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^2 dx$$

input `integrate(cos(f*x+e)^2*(b*csc(f*x+e))^n,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*cos(f*x + e)^2, x)`

**3.294.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \int \cos(e + fx)^2 \left( \frac{b}{\sin(e + fx)} \right)^n dx$$

input `int(cos(e + f*x)^2*(b/sin(e + f*x))^n,x)`output `int(cos(e + f*x)^2*(b/sin(e + f*x))^n, x)`

### 3.295 $\int \cos^4(e + fx)(b \csc(e + fx))^n dx$

3.295.1 Optimal result . . . . .	1722
3.295.2 Mathematica [B] (verified) . . . . .	1722
3.295.3 Rubi [A] (verified) . . . . .	1723
3.295.4 Maple [F] . . . . .	1724
3.295.5 Fricas [F] . . . . .	1724
3.295.6 Sympy [F] . . . . .	1725
3.295.7 Maxima [F] . . . . .	1725
3.295.8 Giac [F] . . . . .	1725
3.295.9 Mupad [F(-1)] . . . . .	1726

#### 3.295.1 Optimal result

Integrand size = 19, antiderivative size = 72

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \frac{b \cos(e + fx)(b \csc(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

```
output b*cos(f*x+e)*(b*csc(f*x+e))^(n-1)*hypergeom([-3/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)/f/(1-n)/(cos(f*x+e)^2)^(1/2)
```

#### 3.295.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 246 vs. 2(72) = 144.

Time = 1.25 (sec) , antiderivative size = 246, normalized size of antiderivative = 3.42

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \frac{2(b \csc(e + fx))^n \left(\operatorname{Hypergeometric2F1}\left(1 - n, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 8\left(\operatorname{Hypergeometric2F1}\left(1 - n, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, \sin^2\left(\frac{1}{2}(e + fx)\right)\right)\right)\right)}{f}$$

```
input Integrate[Cos[e + f*x]^4*(b*Csc[e + f*x])^n,x]
```

output  $(-2*(b*\text{Csc}[e + f*x])^n*(\text{Hypergeometric2F1}[1 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2] - 8*(\text{Hypergeometric2F1}[2 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2] - 3*\text{Hypergeometric2F1}[3 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2] + 4*\text{Hypergeometric2F1}[4 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2] - 2*\text{Hypergeometric2F1}[5 - n, 1/2 - n/2, 3/2 - n/2, -\text{Tan}[(e + f*x)/2]^2]))*\text{Tan}[(e + f*x)/2])/(f*(-1 + n)*(Sec[(e + f*x)/2]^2)^n)$

### 3.295.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^4(e + fx)(b \csc(e + fx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(b \csc(e + fx))^n}{\sec(e + fx)^4} dx \\ & \quad \downarrow \text{3111} \\ & b^2(b \sin(e + fx))^{n-1}(b \csc(e + fx))^{n-1} \int \cos^4(e + fx)(b \sin(e + fx))^{-n} dx \\ & \quad \downarrow \text{3042} \\ & b^2(b \sin(e + fx))^{n-1}(b \csc(e + fx))^{n-1} \int \cos(e + fx)^4(b \sin(e + fx))^{-n} dx \\ & \quad \downarrow \text{3057} \\ & \frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} \text{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}} \end{aligned}$$

input  $\text{Int}[\text{Cos}[e + f*x]^4*(b*\text{Csc}[e + f*x])^n, x]$

output  $(b*\text{Cos}[e + f*x]*(b*\text{Csc}[e + f*x])^{-(1 + n)}*\text{Hypergeometric2F1}[-3/2, (1 - n)/2, (3 - n)/2, \text{Sin}[e + f*x]^2])/(f*(1 - n)*\text{Sqrt}[\text{Cos}[e + f*x]^2])$



## 3.295.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

## 3.295.4 Maple [F]

$$\int \cos(fx + e)^4 (b \csc(fx + e))^n dx$$

input `int(cos(f*x+e)^4*(b*csc(f*x+e))^n,x)`

output `int(cos(f*x+e)^4*(b*csc(f*x+e))^n,x)`

## 3.295.5 Fracas [F]

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^4 dx$$

input `integrate(cos(f*x+e)^4*(b*csc(f*x+e))^n,x, algorithm="fracas")`

output `integral((b*csc(f*x + e))^n*cos(f*x + e)^4, x)`

**3.295.6 Sympy [F]**

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(e + fx))^n \cos^4(e + fx) dx$$

input `integrate(cos(f*x+e)**4*(b*csc(f*x+e))**n,x)`

output `Integral((b*csc(e + f*x))**n*cos(e + f*x)**4, x)`

**3.295.7 Maxima [F]**

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^4 dx$$

input `integrate(cos(f*x+e)^4*(b*csc(f*x+e))^n,x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n*cos(f*x + e)^4, x)`

**3.295.8 Giac [F]**

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^4 dx$$

input `integrate(cos(f*x+e)^4*(b*csc(f*x+e))^n,x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n*cos(f*x + e)^4, x)`

**3.295.9 Mupad [F(-1)]**

Timed out.

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \int \cos(e + fx)^4 \left( \frac{b}{\sin(e + fx)} \right)^n dx$$

input `int(cos(e + f*x)^4*(b/sin(e + f*x))^n,x)`output `int(cos(e + f*x)^4*(b/sin(e + f*x))^n, x)`

### 3.296 $\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx$

3.296.1 Optimal result . . . . .	1727
3.296.2 Mathematica [A] (verified) . . . . .	1727
3.296.3 Rubi [A] (verified) . . . . .	1728
3.296.4 Maple [F] . . . . .	1729
3.296.5 Fricas [F] . . . . .	1729
3.296.6 Sympy [F(-1)] . . . . .	1730
3.296.7 Maxima [F] . . . . .	1730
3.296.8 Giac [F] . . . . .	1730
3.296.9 Mupad [F(-1)] . . . . .	1731

#### 3.296.1 Optimal result

Integrand size = 23, antiderivative size = 81

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \frac{b \cos^2(e + fx)^{5/4} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) (c \sec(e + fx))^{3/2}}{cf(1-n)}$$

output `b*(cos(f*x+e)^2)^(5/4)*(b*csc(f*x+e))^(n-1)*hypergeom([5/4, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*(c*sec(f*x+e))^(3/2)/c/f/(1-n)`

#### 3.296.2 Mathematica [A] (verified)

Time = 11.40 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.14

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \frac{2 \cot(e + fx) (b \csc(e + fx))^n \text{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{1}{4}(3 + 2n), \frac{1}{4}(7 + 2n), \sec^2(e + fx)\right) (c \sec(e + fx))^{3/2}}{f(3 + 2n)}$$

input `Integrate[(b*Csc[e + f*x])^n*(c*Sec[e + f*x])^(3/2),x]`

output `(2*Cot[e + f*x]*(b*Csc[e + f*x])^n*Hypergeometric2F1[(1 + n)/2, (3 + 2*n)/4, (7 + 2*n)/4, Sec[e + f*x]^2]*(c*Sec[e + f*x])^(3/2)*(-Tan[e + f*x]^2)^(1 + n/2))/(f*(3 + 2*n))`

**3.296.3 Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c \sec(e + fx))^{3/2} (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (c \sec(e + fx))^{3/2} (b \csc(e + fx))^n dx \\
 & \quad \downarrow \text{3111} \\
 & \frac{b^2 (c \cos(e + fx))^{5/2} (c \sec(e + fx))^{5/2} (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \frac{(b \sin(e + fx))^{-n}}{(c \cos(e + fx))^{3/2}} dx}{c^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 (c \cos(e + fx))^{5/2} (c \sec(e + fx))^{5/2} (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \frac{(b \sin(e + fx))^{-n}}{(c \cos(e + fx))^{3/2}} dx}{c^2} \\
 & \quad \downarrow \text{3057} \\
 & \frac{b \cos^2(e + fx)^{5/4} (c \sec(e + fx))^{5/2} (b \csc(e + fx))^{n-1} \text{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{cf(1-n)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*(c*Sec[e + f*x])^(3/2),x]`

output `(b*(Cos[e + f*x]^2)^(5/4)*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[5/4, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*(c*Sec[e + f*x])^(5/2))/(c*f*(1 - n))`

## 3.296.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^n_*((a_.)*sin[(e_.) + (f_.)*(x_)])^m_, x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*SIn[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n_, x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*SIn[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*SIn[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

## 3.296.4 Maple [F]

$$\int (b \csc(fx + e))^n (c \sec(fx + e))^{\frac{3}{2}} dx$$

input `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x)`

output `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x)`

## 3.296.5 Fracas [F]

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \int (c \sec(fx + e))^{\frac{3}{2}} (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x, algorithm="fracas")`

output `integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n*c*sec(f*x + e), x)`

**3.296.6 Sympy [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \text{Timed out}$$

input `integrate((b*csc(f*x+e))**n*(c*sec(f*x+e))**(3/2),x)`output `Timed out`**3.296.7 Maxima [F]**

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \int (c \sec(fx + e))^{\frac{3}{2}} (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x, algorithm="maxima")`output `integrate((c*sec(f*x + e))^(3/2)*(b*csc(f*x + e))^n, x)`**3.296.8 Giac [F]**

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \int (c \sec(fx + e))^{\frac{3}{2}} (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x, algorithm="giac")`output `integrate((c*sec(f*x + e))^(3/2)*(b*csc(f*x + e))^n, x)`

**3.296.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \int \left( \frac{c}{\cos(e + fx)} \right)^{3/2} \left( \frac{b}{\sin(e + fx)} \right)^n dx$$

input `int((c/cos(e + f*x))^(3/2)*(b/sin(e + f*x))^n,x)`output `int((c/cos(e + f*x))^(3/2)*(b/sin(e + f*x))^n, x)`



### 3.297 $\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$

3.297.1 Optimal result . . . . .	1732
3.297.2 Mathematica [A] (verified) . . . . .	1732
3.297.3 Rubi [A] (verified) . . . . .	1733
3.297.4 Maple [F] . . . . .	1734
3.297.5 Fricas [F] . . . . .	1734
3.297.6 Sympy [F] . . . . .	1735
3.297.7 Maxima [F] . . . . .	1735
3.297.8 Giac [F] . . . . .	1735
3.297.9 Mupad [F(-1)] . . . . .	1736

#### 3.297.1 Optimal result

Integrand size = 23, antiderivative size = 81

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$$

$$= \frac{b \cos^2(e + fx)^{3/4} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) (c \sec(e + fx))^{3/2}}{cf(1-n)}$$

```
output b*(cos(f*x+e)^2)^(3/4)*(b*csc(f*x+e))^(1-n)*hypergeom([3/4, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*(c*sec(f*x+e))^(3/2)/c/f/(1-n)
```

#### 3.297.2 Mathematica [A] (verified)

Time = 11.38 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.11

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$$

$$= \frac{2 \cot(e + fx) (b \csc(e + fx))^n \text{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{1}{4}(1+2n), \frac{1}{4}(5+2n), \sec^2(e + fx)\right) \sqrt{c \sec(e + fx)}}{f + 2fn}$$

```
input Integrate[(b*Csc[e + f*x])^n*Sqrt[c*Sec[e + f*x]],x]
```

```
output (2*Cot[e + f*x]*(b*Csc[e + f*x])^n*Hypergeometric2F1[(1 + n)/2, (1 + 2*n)/4, (5 + 2*n)/4, Sec[e + f*x]^2]*Sqrt[c*Sec[e + f*x]]*(-Tan[e + f*x]^2)^((1 + n)/2))/(f + 2*f*n)
```

**3.297.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{c \sec(e+fx)} (b \csc(e+fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{c \sec(e+fx)} (b \csc(e+fx))^n dx \\
 & \quad \downarrow \text{3111} \\
 & \frac{b^2 (c \cos(e+fx))^{3/2} (c \sec(e+fx))^{3/2} (b \sin(e+fx))^{n-1} (b \csc(e+fx))^{n-1} \int \frac{(b \sin(e+fx))^{-n}}{\sqrt{c \cos(e+fx)}} dx}{c^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 (c \cos(e+fx))^{3/2} (c \sec(e+fx))^{3/2} (b \sin(e+fx))^{n-1} (b \csc(e+fx))^{n-1} \int \frac{(b \sin(e+fx))^{-n}}{\sqrt{c \cos(e+fx)}} dx}{c^2} \\
 & \quad \downarrow \text{3057} \\
 & \frac{b \cos^2(e+fx)^{3/4} (c \sec(e+fx))^{3/2} (b \csc(e+fx))^{n-1} \text{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e+fx)\right)}{cf(1-n)}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n*Sqrt[c*Sec[e + f*x]],x]`

output `(b*(Cos[e + f*x]^2)^(3/4)*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[3/4, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*(c*Sec[e + f*x])^(3/2))/(c*f*(1 - n))`

## 3.297.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^n_*((a_.)*sin[(e_.) + (f_.)*(x_)])^m_, x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*SIn[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m_*((b_.)*sec[(e_.) + (f_.)*(x_)])^n_, x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*SIn[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*SIn[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

## 3.297.4 Maple [F]

$$\int (b \csc(fx + e))^n \sqrt{c \sec(fx + e)} dx$$

input `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x)`

output `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x)`

## 3.297.5 Fracas [F]

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx = \int \sqrt{c \sec(fx + e)} (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x, algorithm="fracas")`

output `integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n, x)`

**3.297.6 Sympy [F]**

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx = \int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$$

input `integrate((b*csc(f*x+e))**n*(c*sec(f*x+e))**(1/2),x)`

output `Integral((b*csc(e + f*x))**n*sqrt(c*sec(e + f*x)), x)`

**3.297.7 Maxima [F]**

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx = \int \sqrt{c \sec(fx + e)} (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n, x)`

**3.297.8 Giac [F]**

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx = \int \sqrt{c \sec(fx + e)} (b \csc(fx + e))^n dx$$

input `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n, x)`

**3.297.9 Mupad [F(-1)]**

Timed out.

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx = \int \sqrt{\frac{c}{\cos(e + fx)}} \left( \frac{b}{\sin(e + fx)} \right)^n dx$$

input `int((c/cos(e + f*x))^(1/2)*(b/sin(e + f*x))^n,x)`output `int((c/cos(e + f*x))^(1/2)*(b/sin(e + f*x))^n, x)`

**3.298**  $\int \frac{(b \csc(e+fx))^n}{\sqrt{c \sec(e+fx)}} dx$

3.298.1 Optimal result . . . . . 1737  
 3.298.2 Mathematica [A] (verified) . . . . . 1737  
 3.298.3 Rubi [A] (verified) . . . . . 1738  
 3.298.4 Maple [F] . . . . . 1739  
 3.298.5 Fracas [F] . . . . . 1739  
 3.298.6 Sympy [F] . . . . . 1740  
 3.298.7 Maxima [F] . . . . . 1740  
 3.298.8 Giac [F] . . . . . 1740  
 3.298.9 Mupad [F(-1)] . . . . . 1741

**3.298.1 Optimal result**

Integrand size = 23, antiderivative size = 81

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx$$

$$= \frac{b^4 \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right) \sqrt{c \sec(e + fx)}}{cf(1 - n)}$$

output `b*(cos(f*x+e)^2)^(1/4)*(b*csc(f*x+e))^(−1+n)*hypergeom([1/4, 1/2−1/2*n], [3/2−1/2*n], sin(f*x+e)^2)*(c*sec(f*x+e))^(1/2)/c/f/(1−n)`

**3.298.2 Mathematica [A] (verified)**

Time = 31.34 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.17

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx$$

$$= \frac{(b \csc(e + fx))^n \text{Hypergeometric2F1}\left(\frac{1}{2} - \frac{n}{2}, \frac{5}{4} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2(e + fx)\right) \sec^2(e + fx)^{\frac{1}{4} - \frac{n}{2}} \tan(e + fx)}{f(1 - n) \sqrt{c \sec(e + fx)}}$$

input `Integrate[(b*Csc[e + f*x])^n/Sqrt[c*Sec[e + f*x]],x]`

output `((b*Csc[e + f*x])^n*Hypergeometric2F1[1/2 - n/2, 5/4 - n/2, 3/2 - n/2, -Tan[e + f*x]^2]*(Sec[e + f*x]^2)^(1/4 - n/2)*Tan[e + f*x])/(f*(1 - n)*Sqrt[c*Sec[e + f*x]])`

---

3.298.  $\int \frac{(b \csc(e+fx))^n}{\sqrt{c \sec(e+fx)}} dx$

**3.298.3 Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx$$

↓ 3042

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx$$

↓ 3111

$$\frac{b^2 \sqrt{c \cos(e + fx)} \sqrt{c \sec(e + fx)} (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \sqrt{c \cos(e + fx)} (b \sin(e + fx))^{-n} dx}{c^2}$$

↓ 3042

$$\frac{b^2 \sqrt{c \cos(e + fx)} \sqrt{c \sec(e + fx)} (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int \sqrt{c \cos(e + fx)} (b \sin(e + fx))^{-n} dx}{c^2}$$

↓ 3057

$$\frac{b^4 \sqrt{\cos^2(e + fx)} \sqrt{c \sec(e + fx)} (b \csc(e + fx))^{n-1} \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{cf(1-n)}$$

input `Int[(b*Csc[e + f*x])^n/Sqrt[c*Sec[e + f*x]],x]`

output `(b*(Cos[e + f*x]^2)^(1/4)*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[1/4, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sqrt[c*Sec[e + f*x]])/(c*f*(1 - n))`

## 3.298.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

## 3.298.4 Maple [F]

$$\int \frac{(b \csc(fx + e))^n}{\sqrt{c \sec(fx + e)}} dx$$

input `int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x)`

output `int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x)`

## 3.298.5 Fracas [F]

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx = \int \frac{(b \csc(fx + e))^n}{\sqrt{c \sec(fx + e)}} dx$$

input `integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x, algorithm="fracas")`

output `integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n/(c*sec(f*x + e)), x)`



**3.298.6 Sympy [F]**

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx = \int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx$$

input `integrate((b*csc(f*x+e))**n/(c*sec(f*x+e))**(1/2),x)`

output `Integral((b*csc(e + f*x))**n/sqrt(c*sec(e + f*x)), x)`

**3.298.7 Maxima [F]**

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx = \int \frac{(b \csc(fx + e))^n}{\sqrt{c \sec(fx + e)}} dx$$

input `integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n/sqrt(c*sec(f*x + e)), x)`

**3.298.8 Giac [F]**

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx = \int \frac{(b \csc(fx + e))^n}{\sqrt{c \sec(fx + e)}} dx$$

input `integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n/sqrt(c*sec(f*x + e)), x)`

**3.298.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx = \int \frac{\left(\frac{b}{\sin(e + fx)}\right)^n}{\sqrt{\frac{c}{\cos(e + fx)}}} dx$$

input `int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(1/2),x)`output `int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(1/2), x)`

**3.299**  $\int \frac{(b \csc(e+fx))^n}{(c \sec(e+fx))^{3/2}} dx$

3.299.1 Optimal result . . . . .	1742
3.299.2 Mathematica [A] (verified) . . . . .	1742
3.299.3 Rubi [A] (verified) . . . . .	1743
3.299.4 Maple [F] . . . . .	1744
3.299.5 Fricas [F] . . . . .	1744
3.299.6 Sympy [F] . . . . .	1745
3.299.7 Maxima [F] . . . . .	1745
3.299.8 Giac [F] . . . . .	1745
3.299.9 Mupad [F(-1)] . . . . .	1746

**3.299.1 Optimal result**

Integrand size = 23, antiderivative size = 81

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx = \frac{b(b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{cf(1-n)\sqrt[4]{\cos^2(e + fx)}\sqrt{c \sec(e + fx)}}$$

output `b*(b*csc(f*x+e))(-1+n)*hypergeom([-1/4, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)2)/c/f/(1-n)/(cos(f*x+e)2)(1/4)/(c*sec(f*x+e))(1/2)`

**3.299.2 Mathematica [A] (verified)**

Time = 11.70 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.42

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx = \frac{2 \cos(2(e + fx)) \cot(e + fx) (b \csc(e + fx))^n \text{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{1}{4}(-3 + 2n), \frac{1}{4}(1 + 2n), \sec^2(e + fx)\right)}{c^2 f(-3 + 2n)(-2 + \sec^2(e + fx))}$$

input `Integrate[(b*Csc[e + f*x])n/(c*Sec[e + f*x])(3/2),x]`

output `(-2*Cos[2*(e + f*x)]*Cot[e + f*x]*(b*Csc[e + f*x])n*Hypergeometric2F1[(1 + n)/2, (-3 + 2*n)/4, (1 + 2*n)/4, Sec[e + f*x]2]*Sqrt[c*Sec[e + f*x]]*(-Tan[e + f*x]2)((1 + n)/2))/(c2*f*(-3 + 2*n)*(-2 + Sec[e + f*x]2))`

---

3.299.  $\int \frac{(b \csc(e+fx))^n}{(c \sec(e+fx))^{3/2}} dx$

**3.299.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {3042, 3111, 3042, 3057}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3111} \\
 & \frac{b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int (c \cos(e + fx))^{3/2} (b \sin(e + fx))^{-n} dx}{c^2 \sqrt{c \cos(e + fx)} \sqrt{c \sec(e + fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b^2 (b \sin(e + fx))^{n-1} (b \csc(e + fx))^{n-1} \int (c \cos(e + fx))^{3/2} (b \sin(e + fx))^{-n} dx}{c^2 \sqrt{c \cos(e + fx)} \sqrt{c \sec(e + fx)}} \\
 & \quad \downarrow \text{3057} \\
 & \frac{b (b \csc(e + fx))^{n-1} \text{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{c f (1-n) \sqrt[4]{\cos^2(e + fx)} \sqrt{c \sec(e + fx)}}
 \end{aligned}$$

input `Int[(b*Csc[e + f*x])^n/(c*Sec[e + f*x])^(3/2),x]`

output `(b*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[-1/4, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(c*f*(1 - n)*(Cos[e + f*x]^2)^(1/4)*Sqrt[c*Sec[e + f*x]])`

## 3.299.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3057 `Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

rule 3111 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a^2/b^2)*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1) Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]`

## 3.299.4 Maple [F]

$$\int \frac{(b \csc (fx + e))^n}{(c \sec (fx + e))^{\frac{3}{2}}} dx$$

input `int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2),x)`

output `int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2),x)`

## 3.299.5 Fracas [F]

$$\int \frac{(b \csc (e + fx))^n}{(c \sec (e + fx))^{3/2}} dx = \int \frac{(b \csc (fx + e))^n}{(c \sec (fx + e))^{\frac{3}{2}}} dx$$

input `integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2),x, algorithm="fracas")`

output `integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n/(c^2*sec(f*x + e)^2), x)`

**3.299.6 Sympy [F]**

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx = \int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{\frac{3}{2}}} dx$$

input `integrate((b*csc(f*x+e))**n/(c*sec(f*x+e))**(3/2),x)`

output `Integral((b*csc(e + f*x))**n/(c*sec(e + f*x))**(3/2), x)`

**3.299.7 Maxima [F]**

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx = \int \frac{(b \csc(fx + e))^n}{(c \sec(fx + e))^{\frac{3}{2}}} dx$$

input `integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((b*csc(f*x + e))^n/(c*sec(f*x + e))^(3/2), x)`

**3.299.8 Giac [F]**

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx = \int \frac{(b \csc(fx + e))^n}{(c \sec(fx + e))^{\frac{3}{2}}} dx$$

input `integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((b*csc(f*x + e))^n/(c*sec(f*x + e))^(3/2), x)`

**3.299.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx = \int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\left(\frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(3/2),x)`output `int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(3/2), x)`

## APPENDIX

4.1 Listing of Grading functions . . . . .	1747
--	------

## 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

### 4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```



```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

### 4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string),"$ vs. $2(",
                        convert(leaf_count_optimal,string),"=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

### 4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```



```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

*#main function*

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

#### 4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```



```
return grade, grade_annotation
```