

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.5-Secant/122-4.5.2.3-g-sec[^]p-a+b-sec[^]m-c+d-
sec[^]n

Nasser M. Abbasi

December 8, 2023

Compiled on December 8, 2023 at 11:54pm

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	107
4	Appendix	2098

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	15
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [286]. This is test number [122].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (286)	0.00 (0)
Mathematica	98.60 (282)	1.40 (4)
Maple	93.36 (267)	6.64 (19)
Fricas	83.22 (238)	16.78 (48)
Giac	70.63 (202)	29.37 (84)
Mupad	66.78 (191)	33.22 (95)
Maxima	58.04 (166)	41.96 (120)
Sympy	0.35 (1)	99.65 (285)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	99.301	0.699	0.000	0.000
Maple	79.021	8.741	5.594	6.643
Mathematica	74.476	5.944	18.182	1.399
Fricas	62.587	20.280	0.350	16.783
Giac	56.294	14.336	0.000	29.371
Maxima	23.776	34.266	0.000	41.958
Mupad	0.000	66.783	0.000	33.217
Sympy	0.000	0.350	0.000	99.650

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	4	100.00	0.00	0.00
Maple	19	100.00	0.00	0.00
Fricas	48	68.75	31.25	0.00
Giac	84	90.48	0.00	9.52
Mupad	95	0.00	100.00	0.00
Maxima	120	60.00	5.00	35.00
Sympy	285	85.96	14.04	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.51
Giac	0.58
Rubi	0.60
Sympy	0.85
Mathematica	2.56
Fricas	2.65
Maple	4.43
Mupad	15.97

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Sympy	51.00	3.00	51.00	3.00
Rubi	141.19	1.08	120.50	1.00
Maple	161.87	1.27	120.00	1.12
Giac	192.50	1.33	124.00	1.15
Fricas	361.42	2.60	176.50	1.72
Maxima	453.47	4.44	219.00	2.21
Mathematica	467.60	2.40	83.50	0.95
Mupad	803.47	4.72	158.00	1.46

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {176, 205, 206, 207, 216, 224, 232, 233, 265, 269, 275, 277}

Maple {129, 184, 234, 235, 237, 238, 239, 240, 241, 242, 243}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2013
Design v0.01

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	26
2.3	Detailed conclusion table specific for Rubi results	98

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	22
2.1.3	Maple	22
2.1.4	Fricas	23
2.1.5	Maxima	23
2.1.6	Giac	24
2.1.7	Mupad	24
2.1.8	Sympy	25

2.1.1 Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286 }
}

B grade { 197, 212 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 29, 30, 31, 32, 33, 37, 38, 39, 40, 41, 45, 46, 47, 48, 49, 50, 51, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 72, 73, 74, 75, 77, 79, 80, 81, 82, 83, 86, 87, 88, 89, 93, 94, 95, 96, 100, 101, 102, 103, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 156, 157, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 172, 173, 175, 177, 178, 179, 182, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 200, 201, 202, 203, 204, 209, 211, 217, 218, 221, 226, 227, 229, 230, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 254, 255, 256, 257, 259, 260, 261, 262, 263, 264, 266, 267, 268, 271, 275, 276, 279, 281, 282, 283, 286 }

B grade { 3, 171, 174, 180, 181, 210, 212, 213, 219, 220, 225, 228, 252, 253, 258, 273, 285 }

C grade { 15, 27, 28, 34, 35, 36, 42, 43, 44, 52, 53, 54, 70, 76, 78, 84, 85, 90, 91, 92, 97, 98, 99, 104, 105, 106, 158, 176, 183, 197, 198, 199, 205, 206, 207, 208, 214, 215, 216, 222, 223, 224, 231, 232, 233, 265, 269, 270, 272, 277, 280, 284 }

F normal fail { 154, 155, 274, 278 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.3 Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 13, 14, 15, 16, 17, 18, 19, 20, 22, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 71, 72, 73, 74, 75, 76, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 108, 109, 111, 112, 113, 114, 115, 116, 117, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 163, 164, 168, 169, 171, 172, 173, 179, 181, 182, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 244, 245, 246, 247, 248, 249, 250, 251, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 268, 269, 273, 275, 276, 279, 282, 285, 286 }

B grade { 69, 70, 77, 78, 107, 110, 118, 129, 170, 178, 180, 183, 184, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 252, 281 }

C grade { 10, 12, 21, 23, 24, 25, 267, 270, 271, 272, 274, 277, 278, 280, 283, 284 }

F normal fail { 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 165, 166, 167, 174, 175, 176, 177 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.4 Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 70, 71, 72, 73, 75, 76, 77, 78, 79, 80, 81, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 109, 113, 114, 115, 120, 121, 123, 124, 125, 132, 136, 138, 143, 144, 148, 149, 150, 156, 157, 158, 162, 163, 164, 168, 169, 171, 172, 173, 178, 179, 180, 181, 182, 185, 186, 187, 188, 189, 193, 194, 195, 196, 197, 202, 203, 204, 205, 210, 211, 213, 214, 217, 218, 220, 221, 225, 226, 227, 228, 229, 230, 235, 236, 237, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 255, 256, 262, 286 }

B grade { 17, 30, 56, 69, 74, 82, 107, 108, 111, 112, 116, 119, 122, 126, 130, 131, 137, 141, 142, 146, 147, 170, 183, 184, 190, 191, 192, 198, 199, 200, 201, 206, 207, 208, 209, 212, 215, 216, 219, 222, 223, 224, 231, 232, 233, 234, 238, 250, 251, 252, 253, 254, 257, 259, 260, 261, 263, 285 }

C grade { 277 }

F normal fail { 110, 117, 118, 127, 128, 129, 133, 134, 135, 139, 140, 145, 151, 152, 153, 154, 155, 159, 160, 161, 165, 166, 167, 174, 175, 176, 177, 266, 267, 268, 273, 275, 276 }

F(-1) timeout fail { 258, 264, 265, 269, 270, 271, 272, 274, 278, 279, 280, 281, 282, 283, 284 }

F(-2) exception fail { }

2.1.5 Maxima

A grade { 2, 3, 4, 7, 8, 9, 14, 26, 38, 39, 40, 41, 47, 48, 49, 50, 51, 57, 58, 59, 60, 61, 62, 63, 86, 87, 88, 93, 94, 95, 100, 101, 109, 110, 116, 118, 126, 129, 135, 136, 140, 141, 145, 147, 156, 157, 162, 163, 164, 168, 173, 183, 185, 186, 187, 188, 195, 196, 203, 221, 229, 230, 244, 245, 246, 247, 285, 286 }

B grade { 1, 5, 6, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 52, 53, 54, 55, 56, 89, 96, 102, 103, 107, 108, 111, 112, 113, 114, 115, 117, 119, 120, 121, 122, 123, 124, 125, 127, 128, 130, 131, 132, 133, 134, 137, 138, 139, 142, 143, 146, 148, 150, 158, 169, 170, 171, 172, 178, 180, 181, 182, 193, 194, 202, 204, 210, 211, 212, 213, 217, 218, 219, 220, 225, 226, 227, 228 }

C grade { }

F normal fail { 64, 65, 66, 67, 68, 69, 70, 72, 73, 74, 75, 76, 77, 82, 83, 84, 90, 91, 92, 97, 98, 99, 104, 105, 106, 151, 152, 153, 154, 155, 159, 160, 161, 165, 166, 167, 174, 175, 176, 177, 179, 184, }

234, 235, 236, 237, 238, 240, 241, 242, 243, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284 }

F(-1) timeout fail { 71, 78, 79, 80, 81, 85 }

F(-2) exception fail { 144, 149, 189, 190, 191, 192, 197, 198, 199, 200, 201, 205, 206, 207, 208, 209, 214, 215, 216, 222, 223, 224, 231, 232, 233, 239, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263 }

2.1.6 Giac

A grade { 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 168, 169, 170, 171, 172, 173, 179, 190, 195, 196, 199, 203, 204, 208, 210, 211, 212, 213, 214, 215, 216, 218, 219, 220, 221, 225, 226, 227, 229, 230, 248, 249, 255, 257, 261, 262, 263, 285, 286 }

B grade { 3, 14, 185, 186, 187, 188, 189, 191, 192, 193, 194, 197, 198, 200, 201, 202, 205, 206, 207, 209, 217, 222, 223, 224, 228, 231, 232, 233, 244, 245, 246, 247, 250, 251, 252, 253, 254, 256, 258, 259, 260 }

C grade { }

F normal fail { 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 162, 163, 164, 165, 166, 167, 174, 175, 178, 180, 181, 182, 184, 234, 235, 236, 237, 238, 239, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284 }

F(-1) timeout fail { }

F(-2) exception fail { 161, 176, 177, 183, 240, 241, 242, 243 }

2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 71, 72, 73, 74, 79, 80, 81, 82, 86, 87, 88, 89, 93, 94, 95, 96, 100, 101, 102, 103, 107, 108, 109, 111, 112, 113, 114, 115, 116, 119, 120, 121, 122, 123, 124, 125, 126, 130, 131, 132, 141, 146, 147, 157, 162, 163, 164, 168, 169, 170, 171, 172, 173, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224,

225, 226, 227, 228, 229, 230, 231, 232, 233, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254,
255, 256, 257, 258, 259, 260, 261, 262, 263, 285, 286 }

C grade { }

F normal fail { }

F(-1) timedout fail { 68, 69, 70, 75, 76, 77, 78, 83, 84, 85, 90, 91, 92, 97, 98, 99, 104, 105, 106,
110, 117, 118, 127, 128, 129, 133, 134, 135, 136, 137, 138, 139, 140, 142, 143, 144, 145, 148, 149,
150, 151, 152, 153, 154, 155, 156, 158, 159, 160, 161, 165, 166, 167, 174, 175, 176, 177, 178, 179,
180, 181, 182, 183, 184, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 264, 265, 266, 267, 268,
269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { }

B grade { 170 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25,
26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51,
52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 72, 73, 74, 75, 76, 77, 78, 80,
81, 82, 83, 84, 85, 87, 88, 89, 90, 91, 92, 95, 96, 97, 98, 99, 102, 103, 104, 105, 106, 108, 109, 110,
111, 112, 116, 117, 118, 119, 134, 135, 136, 137, 138, 140, 141, 142, 143, 146, 147, 148, 151, 152,
153, 154, 155, 158, 159, 160, 162, 163, 164, 166, 167, 168, 169, 171, 172, 173, 174, 175, 176, 177,
178, 179, 181, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199,
200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219,
220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239,
240, 241, 242, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260,
261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 279, 280, 281,
282, 283, 285, 286 }

F(-1) timedout fail { 64, 71, 79, 86, 93, 94, 100, 101, 107, 113, 114, 115, 120, 121, 122, 123, 124,
125, 126, 127, 128, 129, 130, 131, 132, 133, 139, 144, 145, 149, 150, 156, 157, 161, 165, 180, 182,
243, 278, 284 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	104	154	159	215	131	0	145	176
N.S.	1	0.99	1.47	1.51	2.05	1.25	0.00	1.38	1.68
time (sec)	N/A	0.385	5.132	6.241	0.242	0.286	0.000	0.337	18.912

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	144	111	133	117	0	128	146
N.S.	1	1.00	1.67	1.29	1.55	1.36	0.00	1.49	1.70
time (sec)	N/A	0.351	1.289	3.683	0.238	0.290	0.000	0.325	17.219

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	56	132	98	108	103	0	111	114
N.S.	1	0.92	2.16	1.61	1.77	1.69	0.00	1.82	1.87
time (sec)	N/A	0.296	0.571	4.149	0.213	0.275	0.000	0.311	15.641

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	58	68	67	0	55	77
N.S.	1	1.00	1.00	1.53	1.79	1.76	0.00	1.45	2.03
time (sec)	N/A	0.323	0.031	1.509	0.214	0.292	0.000	0.284	14.542

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	77	50	101	66	0	60	31
N.S.	1	1.00	1.83	1.19	2.40	1.57	0.00	1.43	0.74
time (sec)	N/A	0.293	0.202	0.745	0.220	0.268	0.000	0.297	13.764

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	23	21	97	51	0	20	20
N.S.	1	1.00	0.64	0.58	2.69	1.42	0.00	0.56	0.56
time (sec)	N/A	0.236	0.222	0.802	0.220	0.273	0.000	0.303	13.524

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	43	36	117	78	0	37	35
N.S.	1	1.00	0.57	0.47	1.54	1.03	0.00	0.49	0.46
time (sec)	N/A	0.386	0.399	0.832	0.232	0.258	0.000	0.316	14.247

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	120	55	49	177	104	0	51	61
N.S.	1	1.03	0.47	0.42	1.53	0.90	0.00	0.44	0.53
time (sec)	N/A	0.543	3.953	0.915	0.233	0.264	0.000	0.322	14.117

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	158	164	65	62	197	128	0	65	106
N.S.	1	1.04	0.41	0.39	1.25	0.81	0.00	0.41	0.67
time (sec)	N/A	0.723	5.136	0.899	0.220	0.263	0.000	0.362	13.796

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	157	102	209	368	177	0	197	251
N.S.	1	0.92	0.60	1.22	2.15	1.04	0.00	1.15	1.47
time (sec)	N/A	0.483	1.320	8.879	0.207	0.286	0.000	0.417	17.751

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	139	91	195	321	161	0	178	219
N.S.	1	0.93	0.61	1.30	2.14	1.07	0.00	1.19	1.46
time (sec)	N/A	0.444	0.893	6.697	0.215	0.294	0.000	0.363	17.241

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	81	82	143	227	145	0	159	187
N.S.	1	0.86	0.87	1.52	2.41	1.54	0.00	1.69	1.99
time (sec)	N/A	0.344	0.450	6.147	0.211	0.288	0.000	0.365	18.310

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	73	67	80	79	150	99	0	87	155
N.S.	1	0.92	1.10	1.08	2.05	1.36	0.00	1.19	2.12
time (sec)	N/A	0.440	0.038	2.635	0.212	0.274	0.000	0.321	16.965

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	56	45	96	108	103	0	111	113
N.S.	1	0.92	0.74	1.57	1.77	1.69	0.00	1.82	1.85
time (sec)	N/A	0.298	0.088	4.211	0.202	0.262	0.000	0.305	15.884

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	69	65	82	225	108	0	100	77
N.S.	1	0.93	0.88	1.11	3.04	1.46	0.00	1.35	1.04
time (sec)	N/A	0.471	0.586	0.933	0.207	0.276	0.000	0.310	13.771

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	109	65	201	128	0	84	63
N.S.	1	1.00	1.22	0.73	2.26	1.44	0.00	0.94	0.71
time (sec)	N/A	0.481	0.208	1.003	0.208	0.277	0.000	0.326	13.786

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	25	23	189	83	0	22	22
N.S.	1	1.00	0.66	0.61	4.97	2.18	0.00	0.58	0.58
time (sec)	N/A	0.261	0.140	0.876	0.216	0.279	0.000	0.325	13.543

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	80	47	38	270	114	0	41	37
N.S.	1	1.00	0.59	0.48	3.38	1.42	0.00	0.51	0.46
time (sec)	N/A	0.430	0.570	0.879	0.215	0.261	0.000	0.352	14.421

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	126	59	51	269	140	0	57	67
N.S.	1	1.04	0.49	0.42	2.22	1.16	0.00	0.47	0.55
time (sec)	N/A	0.622	0.772	1.002	0.225	0.258	0.000	0.373	14.170

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	172	69	64	389	168	0	73	108
N.S.	1	1.06	0.42	0.39	2.39	1.03	0.00	0.45	0.66
time (sec)	N/A	0.844	1.255	1.011	0.224	0.275	0.000	0.406	14.261

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	227	208	122	253	443	209	0	235	316
N.S.	1	0.92	0.54	1.11	1.95	0.92	0.00	1.04	1.39
time (sec)	N/A	0.543	3.241	9.323	0.227	0.295	0.000	0.419	17.264

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	206	190	111	239	408	193	0	216	284
N.S.	1	0.92	0.54	1.16	1.98	0.94	0.00	1.05	1.38
time (sec)	N/A	0.506	2.200	8.284	0.207	0.293	0.000	0.439	16.929

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	104	102	176	368	177	0	197	252
N.S.	1	0.86	0.84	1.45	3.04	1.46	0.00	1.63	2.08
time (sec)	N/A	0.370	0.917	8.492	0.213	0.284	0.000	0.398	16.854

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	94	125	142	244	115	0	103	220
N.S.	1	0.94	1.25	1.42	2.44	1.15	0.00	1.03	2.20
time (sec)	N/A	0.538	0.043	3.906	0.232	0.306	0.000	0.369	16.504

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	81	81	143	227	145	0	159	188
N.S.	1	0.86	0.86	1.52	2.41	1.54	0.00	1.69	2.00
time (sec)	N/A	0.339	0.494	6.603	0.217	0.271	0.000	0.361	17.534

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	70	111	133	117	0	128	146
N.S.	1	1.00	0.81	1.29	1.55	1.36	0.00	1.49	1.70
time (sec)	N/A	0.344	0.404	4.161	0.208	0.288	0.000	0.334	16.369

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	98	65	106	387	125	0	118	105
N.S.	1	0.98	0.65	1.06	3.87	1.25	0.00	1.18	1.05
time (sec)	N/A	0.610	0.574	1.111	0.207	0.279	0.000	0.338	14.682

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	117	65	97	349	165	0	116	93
N.S.	1	0.98	0.55	0.82	2.93	1.39	0.00	0.97	0.78
time (sec)	N/A	0.699	0.659	1.145	0.226	0.257	0.000	0.363	13.372

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	135	139	78	309	176	0	102	78
N.S.	1	1.02	1.05	0.59	2.34	1.33	0.00	0.77	0.59
time (sec)	N/A	0.694	0.215	1.077	0.225	0.266	0.000	0.368	13.101

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	25	23	356	111	0	22	22
N.S.	1	1.00	0.66	0.61	9.37	2.92	0.00	0.58	0.58
time (sec)	N/A	0.251	0.111	0.939	0.225	0.262	0.000	0.349	12.855

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	80	47	38	357	140	0	41	37
N.S.	1	1.00	0.59	0.48	4.46	1.75	0.00	0.51	0.46
time (sec)	N/A	0.429	0.350	1.045	0.233	0.262	0.000	0.406	12.926

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	126	59	51	518	168	0	57	67
N.S.	1	1.04	0.49	0.42	4.28	1.39	0.00	0.47	0.55
time (sec)	N/A	0.603	0.750	1.012	0.222	0.263	0.000	0.418	13.323

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	162	172	69	64	517	194	0	73	108
N.S.	1	1.06	0.43	0.40	3.19	1.20	0.00	0.45	0.67
time (sec)	N/A	0.846	5.005	1.167	0.242	0.262	0.000	0.503	13.230

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	116	53	138	591	153	0	132	112
N.S.	1	0.96	0.44	1.14	4.88	1.26	0.00	1.09	0.93
time (sec)	N/A	0.447	0.991	1.303	0.209	0.286	0.000	0.338	13.273

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	98	53	106	386	140	0	116	96
N.S.	1	0.98	0.53	1.06	3.86	1.40	0.00	1.16	0.96
time (sec)	N/A	0.632	0.478	1.104	0.199	0.270	0.000	0.326	13.594

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	70	62	80	224	119	0	97	77
N.S.	1	0.95	0.84	1.08	3.03	1.61	0.00	1.31	1.04
time (sec)	N/A	0.483	0.437	0.963	0.231	0.287	0.000	0.302	13.121

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	77	47	101	70	0	58	31
N.S.	1	1.00	1.88	1.15	2.46	1.71	0.00	1.41	0.76
time (sec)	N/A	0.294	0.153	0.796	0.197	0.269	0.000	0.287	13.267

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	17	18	18	0	18	18
N.S.	1	1.00	1.00	1.06	1.12	1.12	0.00	1.12	1.12
time (sec)	N/A	0.313	0.022	0.994	0.193	0.255	0.000	0.285	12.919

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	51	53	48	77	50	0	56	50
N.S.	1	0.86	0.90	0.81	1.31	0.85	0.00	0.95	0.85
time (sec)	N/A	0.340	0.461	0.787	0.196	0.286	0.000	0.294	13.096

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	74	61	59	97	74	0	69	63
N.S.	1	0.95	0.78	0.76	1.24	0.95	0.00	0.88	0.81
time (sec)	N/A	0.398	0.526	0.933	0.204	0.251	0.000	0.317	13.093

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	120	109	79	74	117	102	0	82	83
N.S.	1	0.91	0.66	0.62	0.98	0.85	0.00	0.68	0.69
time (sec)	N/A	0.466	1.526	0.888	0.209	0.256	0.000	0.330	13.438

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	162	75	155	765	210	0	156	170
N.S.	1	0.99	0.46	0.95	4.66	1.28	0.00	0.95	1.04
time (sec)	N/A	0.660	2.917	1.450	0.230	0.274	0.000	0.406	12.939

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	146	75	125	531	197	0	140	136
N.S.	1	0.97	0.50	0.83	3.54	1.31	0.00	0.93	0.91
time (sec)	N/A	0.879	1.964	1.267	0.212	0.291	0.000	0.349	12.991

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	118	75	95	341	178	0	121	104
N.S.	1	0.99	0.63	0.80	2.87	1.50	0.00	1.02	0.87
time (sec)	N/A	0.717	0.455	1.118	0.214	0.269	0.000	0.346	13.039

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	109	65	196	138	0	89	46
N.S.	1	1.00	1.24	0.74	2.23	1.57	0.00	1.01	0.52
time (sec)	N/A	0.494	0.156	0.827	0.219	0.280	0.000	0.311	13.037

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	23	21	94	53	0	20	20
N.S.	1	1.00	0.64	0.58	2.61	1.47	0.00	0.56	0.56
time (sec)	N/A	0.233	0.059	0.785	0.203	0.270	0.000	0.286	12.844

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	51	59	48	76	49	0	69	61
N.S.	1	0.86	1.00	0.81	1.29	0.83	0.00	1.17	1.03
time (sec)	N/A	0.339	0.707	0.728	0.210	0.266	0.000	0.304	12.871

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	32	33	28	31	50	0	31	28
N.S.	1	0.84	0.87	0.74	0.82	1.32	0.00	0.82	0.74
time (sec)	N/A	0.348	0.041	1.026	0.186	0.256	0.000	0.318	12.991

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	69	79	74	121	109	0	96	76
N.S.	1	0.86	0.99	0.92	1.51	1.36	0.00	1.20	0.95
time (sec)	N/A	0.356	0.828	0.665	0.202	0.282	0.000	0.327	13.465

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	90	89	87	140	120	0	109	89
N.S.	1	0.92	0.91	0.89	1.43	1.22	0.00	1.11	0.91
time (sec)	N/A	0.401	1.690	0.670	0.200	0.253	0.000	0.346	14.314

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	129	99	100	161	163	0	122	102
N.S.	1	0.91	0.70	0.71	1.14	1.16	0.00	0.87	0.72
time (sec)	N/A	0.469	3.868	0.851	0.214	0.269	0.000	0.391	15.563

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	215	210	75	168	935	263	0	176	193
N.S.	1	0.98	0.35	0.78	4.35	1.22	0.00	0.82	0.90
time (sec)	N/A	0.891	5.144	1.056	0.260	0.270	0.000	0.460	12.969

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	193	194	75	136	680	250	0	159	159
N.S.	1	1.01	0.39	0.70	3.52	1.30	0.00	0.82	0.82
time (sec)	N/A	1.144	3.394	0.984	0.214	0.288	0.000	0.412	12.956

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	166	75	108	470	231	0	141	126
N.S.	1	1.01	0.46	0.66	2.87	1.41	0.00	0.86	0.77
time (sec)	N/A	0.967	0.926	0.885	0.223	0.297	0.000	0.373	13.038

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	134	139	76	304	192	0	109	61
N.S.	1	1.02	1.06	0.58	2.32	1.47	0.00	0.83	0.47
time (sec)	N/A	0.702	0.187	0.725	0.211	0.272	0.000	0.355	13.042

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	25	23	185	82	0	22	22
N.S.	1	1.00	0.66	0.61	4.87	2.16	0.00	0.58	0.58
time (sec)	N/A	0.259	0.085	0.723	0.200	0.257	0.000	0.336	13.005

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	43	36	115	79	0	37	35
N.S.	1	1.00	0.57	0.47	1.51	1.04	0.00	0.49	0.46
time (sec)	N/A	0.385	0.196	0.659	0.204	0.262	0.000	0.320	13.132

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	74	67	59	95	76	0	87	74
N.S.	1	0.95	0.86	0.76	1.22	0.97	0.00	1.12	0.95
time (sec)	N/A	0.388	0.488	0.713	0.213	0.259	0.000	0.343	12.950

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	69	79	74	120	109	0	103	111
N.S.	1	0.86	0.99	0.92	1.50	1.36	0.00	1.29	1.39
time (sec)	N/A	0.348	0.733	0.703	0.205	0.258	0.000	0.359	13.239

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	41	50	41	41	76	0	41	38
N.S.	1	0.69	0.85	0.69	0.69	1.29	0.00	0.69	0.64
time (sec)	N/A	0.362	0.062	1.078	0.210	0.273	0.000	0.367	13.393

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	81	99	99	159	163	0	128	129
N.S.	1	0.82	1.00	1.00	1.61	1.65	0.00	1.29	1.30
time (sec)	N/A	0.366	3.107	0.789	0.199	0.263	0.000	0.401	14.146

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	120	110	109	110	181	190	0	142	109
N.S.	1	0.92	0.91	0.92	1.51	1.58	0.00	1.18	0.91
time (sec)	N/A	0.438	5.142	0.787	0.208	0.278	0.000	0.418	14.424

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	162	145	119	121	200	217	0	155	120
N.S.	1	0.90	0.73	0.75	1.23	1.34	0.00	0.96	0.74
time (sec)	N/A	0.495	5.141	0.889	0.216	0.253	0.000	0.426	14.679

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	171	70	77	0	119	0	107	483
N.S.	1	1.05	0.43	0.47	0.00	0.73	0.00	0.66	2.96
time (sec)	N/A	0.793	2.276	7.670	0.000	0.274	0.000	0.840	20.552

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	126	60	67	0	105	0	83	384
N.S.	1	1.03	0.49	0.55	0.00	0.86	0.00	0.68	3.15
time (sec)	N/A	0.601	0.678	7.285	0.000	0.275	0.000	0.829	17.242

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	81	50	55	0	82	0	56	120
N.S.	1	1.00	0.62	0.68	0.00	1.01	0.00	0.69	1.48
time (sec)	N/A	0.419	0.416	4.587	0.000	0.270	0.000	0.697	17.059

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	38	42	0	65	0	31	87
N.S.	1	1.00	0.97	1.08	0.00	1.67	0.00	0.79	2.23
time (sec)	N/A	0.249	0.170	3.961	0.000	0.264	0.000	0.664	14.442

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	77	78	80	0	272	0	61	0
N.S.	1	1.00	1.01	1.04	0.00	3.53	0.00	0.79	0.00
time (sec)	N/A	0.383	0.253	3.635	0.000	0.329	0.000	0.953	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	81	178	0	342	0	72	0
N.S.	1	1.00	1.07	2.34	0.00	4.50	0.00	0.95	0.00
time (sec)	N/A	0.382	0.777	4.181	0.000	0.339	0.000	1.004	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	117	60	269	0	405	0	104	0
N.S.	1	1.04	0.53	2.38	0.00	3.58	0.00	0.92	0.00
time (sec)	N/A	0.526	0.479	4.186	0.000	0.346	0.000	1.176	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-1)	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	179	88	79	0	147	0	109	606
N.S.	1	1.05	0.51	0.46	0.00	0.86	0.00	0.64	3.54
time (sec)	N/A	0.953	2.477	16.900	0.000	0.288	0.000	1.019	25.263

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	132	78	69	0	131	0	85	503
N.S.	1	1.03	0.61	0.54	0.00	1.02	0.00	0.66	3.93
time (sec)	N/A	0.746	1.662	15.644	0.000	0.268	0.000	0.833	20.220

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	66	57	0	105	0	58	384
N.S.	1	1.00	0.78	0.67	0.00	1.24	0.00	0.68	4.52
time (sec)	N/A	0.496	1.458	5.524	0.000	0.277	0.000	0.875	17.148

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	55	46	0	84	0	33	93
N.S.	1	1.00	1.34	1.12	0.00	2.05	0.00	0.80	2.27
time (sec)	N/A	0.282	0.602	4.711	0.000	0.273	0.000	0.757	17.857

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	123	107	108	0	343	0	82	0
N.S.	1	1.05	0.91	0.92	0.00	2.93	0.00	0.70	0.00
time (sec)	N/A	0.610	0.563	4.243	0.000	0.316	0.000	1.061	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	126	64	120	0	372	0	109	0
N.S.	1	1.12	0.57	1.06	0.00	3.29	0.00	0.96	0.00
time (sec)	N/A	0.620	0.423	4.841	0.000	0.314	0.000	1.230	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	127	150	228	0	429	0	106	0
N.S.	1	1.09	1.28	1.95	0.00	3.67	0.00	0.91	0.00
time (sec)	N/A	0.622	1.133	4.581	0.000	0.327	0.000	1.299	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	168	64	320	0	517	0	139	0
N.S.	1	1.02	0.39	1.95	0.00	3.15	0.00	0.85	0.00
time (sec)	N/A	0.776	0.614	5.486	0.000	0.345	0.000	1.114	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-1)	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	179	88	79	0	163	0	109	710
N.S.	1	1.05	0.51	0.46	0.00	0.95	0.00	0.64	4.15
time (sec)	N/A	0.941	3.768	41.034	0.000	0.290	0.000	1.064	27.136

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-1)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	132	78	69	0	147	0	85	607
N.S.	1	1.03	0.61	0.54	0.00	1.15	0.00	0.66	4.74
time (sec)	N/A	0.719	1.034	38.804	0.000	0.271	0.000	1.033	25.004

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-1)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	66	57	0	119	0	58	471
N.S.	1	1.00	0.78	0.67	0.00	1.40	0.00	0.68	5.54
time (sec)	N/A	0.483	0.679	6.830	0.000	0.276	0.000	0.862	21.874

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	55	46	0	97	0	33	375
N.S.	1	1.00	1.34	1.12	0.00	2.37	0.00	0.80	9.15
time (sec)	N/A	0.295	0.774	5.636	0.000	0.263	0.000	0.823	17.156

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	168	119	125	0	377	0	111	0
N.S.	1	1.02	0.73	0.76	0.00	2.30	0.00	0.68	0.00
time (sec)	N/A	0.840	0.844	6.383	0.000	0.344	0.000	1.177	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	171	64	141	0	432	0	124	0
N.S.	1	1.02	0.38	0.84	0.00	2.57	0.00	0.74	0.00
time (sec)	N/A	0.852	0.595	7.095	0.000	0.331	0.000	1.033	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-1)	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	174	176	64	223	0	441	0	133	0
N.S.	1	1.01	0.37	1.28	0.00	2.53	0.00	0.76	0.00
time (sec)	N/A	0.853	0.615	6.293	0.000	0.318	0.000	1.138	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	142	72	81	163	88	0	108	164
N.S.	1	1.00	0.51	0.57	1.15	0.62	0.00	0.76	1.15
time (sec)	N/A	0.692	0.850	5.281	0.316	0.266	0.000	0.830	17.983

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	107	62	71	137	77	0	84	125
N.S.	1	0.99	0.57	0.66	1.27	0.71	0.00	0.78	1.16
time (sec)	N/A	0.566	0.535	5.040	0.298	0.280	0.000	0.797	15.559

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	50	57	110	50	0	60	77
N.S.	1	1.00	0.69	0.79	1.53	0.69	0.00	0.83	1.07
time (sec)	N/A	0.443	0.312	2.822	0.300	0.267	0.000	0.702	13.685

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	29	28	84	45	0	59	40
N.S.	1	1.00	0.74	0.72	2.15	1.15	0.00	1.51	1.03
time (sec)	N/A	0.297	0.114	3.295	0.303	0.275	0.000	0.627	13.157

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	52	112	0	269	0	64	0
N.S.	1	1.00	0.58	1.26	0.00	3.02	0.00	0.72	0.00
time (sec)	N/A	0.456	0.292	3.001	0.000	0.324	0.000	0.437	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	124	58	192	0	329	0	97	0
N.S.	1	1.02	0.48	1.57	0.00	2.70	0.00	0.80	0.00
time (sec)	N/A	0.601	0.528	3.100	0.000	0.321	0.000	0.497	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	160	58	268	0	401	0	128	0
N.S.	1	1.03	0.37	1.72	0.00	2.57	0.00	0.82	0.00
time (sec)	N/A	0.770	0.586	3.310	0.000	0.324	0.000	0.544	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	155	72	89	188	103	0	121	188
N.S.	1	1.00	0.46	0.57	1.21	0.66	0.00	0.78	1.21
time (sec)	N/A	0.866	1.505	12.579	0.305	0.270	0.000	0.930	17.736

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	122	64	79	163	82	0	99	136
N.S.	1	0.99	0.52	0.64	1.33	0.67	0.00	0.80	1.11
time (sec)	N/A	0.700	0.899	12.524	0.319	0.269	0.000	0.808	17.268

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	88	54	53	110	72	0	60	134
N.S.	1	0.99	0.61	0.60	1.24	0.81	0.00	0.67	1.51
time (sec)	N/A	0.529	0.359	3.375	0.309	0.263	0.000	0.703	16.881

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	55	44	109	60	0	62	94
N.S.	1	1.00	1.34	1.07	2.66	1.46	0.00	1.51	2.29
time (sec)	N/A	0.310	0.118	3.705	0.315	0.267	0.000	0.648	17.175

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	137	60	184	0	331	0	93	0
N.S.	1	0.99	0.43	1.33	0.00	2.40	0.00	0.67	0.00
time (sec)	N/A	0.722	0.318	3.503	0.000	0.317	0.000	0.571	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	172	64	200	0	369	0	129	0
N.S.	1	1.02	0.38	1.18	0.00	2.18	0.00	0.76	0.00
time (sec)	N/A	0.886	0.488	3.612	0.000	0.334	0.000	0.517	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	203	208	64	281	0	487	0	160	0
N.S.	1	1.02	0.32	1.38	0.00	2.40	0.00	0.79	0.00
time (sec)	N/A	1.036	0.798	3.350	0.000	0.334	0.000	0.555	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	172	74	91	214	109	0	126	492
N.S.	1	1.02	0.44	0.54	1.27	0.64	0.00	0.75	2.91
time (sec)	N/A	0.954	0.911	36.524	0.324	0.284	0.000	1.024	22.304

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	138	64	71	189	104	0	90	456
N.S.	1	1.02	0.47	0.53	1.40	0.77	0.00	0.67	3.38
time (sec)	N/A	0.773	1.463	41.609	0.322	0.264	0.000	0.860	19.273

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	54	73	163	88	0	58	446
N.S.	1	1.00	0.61	0.83	1.85	1.00	0.00	0.66	5.07
time (sec)	N/A	0.515	0.504	3.442	0.334	0.272	0.000	0.738	18.939

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	55	46	136	74	0	62	441
N.S.	1	1.00	1.34	1.12	3.32	1.80	0.00	1.51	10.76
time (sec)	N/A	0.296	0.117	3.775	0.316	0.261	0.000	0.689	19.906

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	185	60	211	0	401	0	119	0
N.S.	1	1.02	0.33	1.17	0.00	2.22	0.00	0.66	0.00
time (sec)	N/A	0.956	0.753	3.408	0.000	0.340	0.000	0.474	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	212	220	64	275	0	483	0	154	0
N.S.	1	1.04	0.30	1.30	0.00	2.28	0.00	0.73	0.00
time (sec)	N/A	1.169	0.835	3.577	0.000	0.347	0.000	0.583	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	256	64	272	0	461	0	184	0
N.S.	1	1.04	0.26	1.11	0.00	1.87	0.00	0.75	0.00
time (sec)	N/A	1.377	0.999	3.538	0.000	0.366	0.000	0.713	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	70	88	638	93	0	0	136
N.S.	1	1.00	1.63	2.05	14.84	2.16	0.00	0.00	3.16
time (sec)	N/A	0.320	0.801	3.337	0.392	0.261	0.000	0.000	15.320

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	60	74	298	78	0	0	78
N.S.	1	1.00	1.40	1.72	6.93	1.81	0.00	0.00	1.81
time (sec)	N/A	0.331	0.617	3.461	0.366	0.272	0.000	0.000	14.006

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	51	47	55	56	0	0	47
N.S.	1	1.00	1.24	1.15	1.34	1.37	0.00	0.00	1.15
time (sec)	N/A	0.323	0.323	3.299	0.342	0.274	0.000	0.000	13.506

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	54	98	92	0	0	0	0
N.S.	1	1.00	1.06	1.92	1.80	0.00	0.00	0.00	0.00
time (sec)	N/A	0.336	0.322	3.268	0.337	0.000	0.000	0.000	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	42	50	514	79	0	0	118
N.S.	1	1.00	1.00	1.19	12.24	1.88	0.00	0.00	2.81
time (sec)	N/A	0.332	0.509	3.219	0.379	0.268	0.000	0.000	14.622

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	56	67	758	106	0	0	203
N.S.	1	1.00	1.30	1.56	17.63	2.47	0.00	0.00	4.72
time (sec)	N/A	0.336	0.534	3.367	0.463	0.265	0.000	0.000	18.108

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	84	107	1680	112	0	0	294
N.S.	1	1.00	0.94	1.20	18.88	1.26	0.00	0.00	3.30
time (sec)	N/A	0.591	2.933	3.603	0.409	0.283	0.000	0.000	17.665

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	87	96	1105	112	0	0	195
N.S.	1	1.00	0.98	1.08	12.42	1.26	0.00	0.00	2.19
time (sec)	N/A	0.580	1.619	3.543	0.393	0.280	0.000	0.000	17.210

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	64	83	550	82	0	0	108
N.S.	1	1.00	0.72	0.93	6.18	0.92	0.00	0.00	1.21
time (sec)	N/A	0.565	0.703	3.406	0.417	0.273	0.000	0.000	14.723

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	66	54	56	76	0	0	76
N.S.	1	1.00	1.53	1.26	1.30	1.77	0.00	0.00	1.77
time (sec)	N/A	0.319	0.408	3.799	0.328	0.259	0.000	0.000	14.088

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	62	128	275	0	0	0	0
N.S.	1	1.00	0.65	1.35	2.89	0.00	0.00	0.00	0.00
time (sec)	N/A	0.561	0.412	3.830	0.389	0.000	0.000	0.000	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	99	70	199	122	0	0	0	0
N.S.	1	1.00	0.71	2.01	1.23	0.00	0.00	0.00	0.00
time (sec)	N/A	0.571	0.588	3.399	0.318	0.000	0.000	0.000	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	42	65	533	95	0	0	165
N.S.	1	1.00	1.00	1.55	12.69	2.26	0.00	0.00	3.93
time (sec)	N/A	0.337	0.861	3.590	0.421	0.268	0.000	0.000	16.765

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	68	77	1559	133	0	0	273
N.S.	1	1.00	0.77	0.88	17.72	1.51	0.00	0.00	3.10
time (sec)	N/A	0.582	1.694	3.227	0.804	0.269	0.000	0.000	18.890

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	68	87	2608	158	0	0	340
N.S.	1	1.00	0.74	0.95	28.35	1.72	0.00	0.00	3.70
time (sec)	N/A	0.570	4.356	3.335	3.321	0.278	0.000	0.000	19.886

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	68	97	3906	184	0	0	407
N.S.	1	1.00	0.74	1.05	42.46	2.00	0.00	0.00	4.42
time (sec)	N/A	0.578	5.381	3.577	16.645	0.280	0.000	0.000	19.241

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	134	138	92	111	2454	152	0	0	307
N.S.	1	1.03	0.69	0.83	18.31	1.13	0.00	0.00	2.29
time (sec)	N/A	0.813	4.304	1.501	0.430	0.260	0.000	0.000	17.764

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	134	138	76	101	1526	106	0	0	215
N.S.	1	1.03	0.57	0.75	11.39	0.79	0.00	0.00	1.60
time (sec)	N/A	0.816	1.229	1.665	0.404	0.272	0.000	0.000	17.358

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	88	87	1106	112	0	0	195
N.S.	1	1.00	0.99	0.98	12.43	1.26	0.00	0.00	2.19
time (sec)	N/A	0.557	0.833	3.662	0.394	0.262	0.000	0.000	16.854

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	70	58	58	93	0	0	136
N.S.	1	1.00	1.63	1.35	1.35	2.16	0.00	0.00	3.16
time (sec)	N/A	0.323	0.431	3.794	0.322	0.258	0.000	0.000	14.985

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	142	76	148	737	0	0	0	0
N.S.	1	1.01	0.54	1.05	5.23	0.00	0.00	0.00	0.00
time (sec)	N/A	0.815	0.369	3.606	0.412	0.000	0.000	0.000	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	143	78	235	2035	0	0	0	0
N.S.	1	0.99	0.54	1.62	14.03	0.00	0.00	0.00	0.00
time (sec)	N/A	0.828	0.719	3.618	0.515	0.000	0.000	0.000	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	145	149	81	272	169	0	0	0	0
N.S.	1	1.03	0.56	1.88	1.17	0.00	0.00	0.00	0.00
time (sec)	N/A	0.821	1.301	3.595	0.310	0.000	0.000	0.000	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	42	71	1815	126	0	0	199
N.S.	1	1.00	1.00	1.69	43.21	3.00	0.00	0.00	4.74
time (sec)	N/A	0.326	1.808	3.612	0.414	0.272	0.000	0.000	17.870

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	78	81	2719	166	0	0	350
N.S.	1	1.00	0.89	0.92	30.90	1.89	0.00	0.00	3.98
time (sec)	N/A	0.575	3.977	3.522	3.264	0.271	0.000	0.000	18.646

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	138	78	91	4108	194	0	0	419
N.S.	1	1.04	0.59	0.68	30.89	1.46	0.00	0.00	3.15
time (sec)	N/A	0.841	5.425	3.619	17.083	0.284	0.000	0.000	18.618

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	F	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	139	141	74	137	737	0	0	160	0
N.S.	1	1.01	0.53	0.99	5.30	0.00	0.00	1.15	0.00
time (sec)	N/A	0.807	0.620	3.785	0.404	0.000	0.000	1.554	0.000

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	F	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	94	60	116	276	0	0	129	0
N.S.	1	1.00	0.64	1.23	2.94	0.00	0.00	1.37	0.00
time (sec)	N/A	0.549	0.485	3.616	0.393	0.000	0.000	1.341	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	F	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	50	50	75	64	0	0	59	0
N.S.	1	1.00	1.00	1.50	1.28	0.00	0.00	1.18	0.00
time (sec)	N/A	0.306	0.232	3.682	0.305	0.000	0.000	1.166	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	47	65	44	204	0	70	0
N.S.	1	1.00	1.00	1.38	0.94	4.34	0.00	1.49	0.00
time (sec)	N/A	0.386	0.289	3.239	0.360	0.311	0.000	1.619	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	75	115	406	382	0	92	0
N.S.	1	1.00	0.79	1.21	4.27	4.02	0.00	0.97	0.00
time (sec)	N/A	0.619	0.486	3.594	0.378	0.341	0.000	1.881	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	145	83	157	1201	456	0	122	0
N.S.	1	1.04	0.59	1.12	8.58	3.26	0.00	0.87	0.00
time (sec)	N/A	0.909	0.976	3.637	0.429	0.348	0.000	1.711	0.000

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	F	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	141	84	192	2035	0	0	161	0
N.S.	1	0.99	0.59	1.35	14.33	0.00	0.00	1.13	0.00
time (sec)	N/A	0.813	0.635	3.681	0.490	0.000	0.000	1.535	0.000

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	F	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	70	148	99	0	0	79	0
N.S.	1	1.00	0.74	1.56	1.04	0.00	0.00	0.83	0.00
time (sec)	N/A	0.560	0.421	3.376	0.306	0.000	0.000	1.495	0.000

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	42	56	54	78	0	36	50
N.S.	1	1.00	1.00	1.33	1.29	1.86	0.00	0.86	1.19
time (sec)	N/A	0.310	1.116	3.572	0.312	0.265	0.000	1.240	14.626

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	60	100	397	380	0	95	0
N.S.	1	1.00	0.63	1.05	4.18	4.00	0.00	1.00	0.00
time (sec)	N/A	0.644	0.352	3.585	0.383	0.318	0.000	1.900	0.000

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	77	64	115	567	402	0	120	0
N.S.	1	0.74	0.62	1.11	5.45	3.87	0.00	1.15	0.00
time (sec)	N/A	0.499	0.412	3.509	0.406	0.325	0.000	2.088	0.000

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	122	102	216	0	544	0	153	0
N.S.	1	0.84	0.70	1.48	0.00	3.73	0.00	1.05	0.00
time (sec)	N/A	0.779	0.623	3.616	0.000	0.351	0.000	1.919	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	F	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	145	88	218	133	0	0	114	0
N.S.	1	1.00	0.61	1.50	0.92	0.00	0.00	0.79	0.00
time (sec)	N/A	0.823	0.722	3.541	0.299	0.000	0.000	1.350	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	51	67	98	95	0	59	119
N.S.	1	1.00	1.21	1.60	2.33	2.26	0.00	1.40	2.83
time (sec)	N/A	0.341	0.522	2.385	0.316	0.264	0.000	1.535	15.261

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	71	58	58	104	0	37	120
N.S.	1	1.00	1.65	1.35	1.35	2.42	0.00	0.86	2.79
time (sec)	N/A	0.327	1.165	3.506	0.327	0.262	0.000	1.485	14.609

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	145	70	138	1191	456	0	124	0
N.S.	1	1.04	0.50	0.99	8.51	3.26	0.00	0.89	0.00
time (sec)	N/A	0.933	0.350	2.307	0.433	0.342	0.000	1.763	0.000

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	122	102	215	0	536	0	151	0
N.S.	1	0.84	0.70	1.47	0.00	3.67	0.00	1.03	0.00
time (sec)	N/A	0.777	0.520	2.474	0.000	0.350	0.000	1.896	0.000

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	102	77	158	1659	482	0	182	0
N.S.	1	0.64	0.48	0.99	10.37	3.01	0.00	1.14	0.00
time (sec)	N/A	0.604	0.564	2.289	0.552	0.351	0.000	1.860	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	101	101	95	0	0	0	0	0	0
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.320	0.677	0.000	0.000	0.000	0.000	0.000	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	92	92	89	0	0	0	0	0	0
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.315	0.285	0.000	0.000	0.000	0.000	0.000	0.000

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	90	90	85	0	0	0	0	0	0
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.271	0.191	0.000	0.000	0.000	0.000	0.000	0.000

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	90	90	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.313	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	92	92	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.309	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	160	159	109	0	228	191	0	0	0
N.S.	1	0.99	0.68	0.00	1.42	1.19	0.00	0.00	0.00
time (sec)	N/A	0.804	0.817	0.000	0.329	0.292	0.000	0.000	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	100	102	72	0	171	112	0	0	154
N.S.	1	1.02	0.72	0.00	1.71	1.12	0.00	0.00	1.54
time (sec)	N/A	0.503	0.516	0.000	0.350	0.274	0.000	0.000	15.508

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	46	46	163	0	114	70	0	0	0
N.S.	1	1.00	3.54	0.00	2.48	1.52	0.00	0.00	0.00
time (sec)	N/A	0.282	1.697	0.000	0.328	0.276	0.000	0.000	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	69	69	67	0	0	0	0	0	0
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.325	0.366	0.000	0.000	0.000	0.000	0.000	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	74	74	74	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.333	0.427	0.000	0.000	0.000	0.000	0.000	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	74	74	74	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.336	0.459	0.000	0.000	0.000	0.000	0.000	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	169	168	104	0	156	120	0	0	290
N.S.	1	0.99	0.62	0.00	0.92	0.71	0.00	0.00	1.72
time (sec)	N/A	0.817	2.473	0.000	0.310	0.287	0.000	0.000	22.279

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	106	76	97	107	93	0	0	145
N.S.	1	1.02	0.73	0.93	1.03	0.89	0.00	0.00	1.39
time (sec)	N/A	0.538	1.085	5.438	0.315	0.268	0.000	0.000	19.408

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	49	76	62	72	0	0	105
N.S.	1	1.00	1.04	1.62	1.32	1.53	0.00	0.00	2.23
time (sec)	N/A	0.302	0.793	5.239	0.324	0.265	0.000	0.000	14.551

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	101	101	104	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.336	0.541	0.000	0.000	0.000	0.000	0.000	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	99	99	105	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.344	1.289	0.000	0.000	0.000	0.000	0.000	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	101	101	105	0	0	0	0	0	0
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.360	2.694	0.000	0.000	0.000	0.000	0.000	0.000

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	104	68	137	172	131	0	145	175
N.S.	1	0.99	0.65	1.30	1.64	1.25	0.00	1.38	1.67
time (sec)	N/A	0.399	0.228	4.052	0.231	0.269	0.000	0.349	17.843

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	78	57	126	160	117	0	128	146
N.S.	1	0.91	0.66	1.47	1.86	1.36	0.00	1.49	1.70
time (sec)	N/A	0.363	0.135	3.808	0.225	0.272	0.000	0.319	16.592

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	34	36	35	51	15	15
N.S.	1	1.00	1.00	2.00	2.12	2.06	3.00	0.88	0.88
time (sec)	N/A	0.294	0.013	1.974	0.230	0.249	0.853	0.297	13.273

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	57	154	78	194	105	0	87	71
N.S.	1	1.02	2.75	1.39	3.46	1.88	0.00	1.55	1.27
time (sec)	N/A	0.477	0.599	0.726	0.246	0.275	0.000	0.302	13.561

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	73	48	60	144	123	0	81	44
N.S.	1	1.04	0.69	0.86	2.06	1.76	0.00	1.16	0.63
time (sec)	N/A	0.569	0.352	0.583	0.233	0.266	0.000	0.319	13.155

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	88	47	36	115	78	0	37	35
N.S.	1	1.02	0.55	0.42	1.34	0.91	0.00	0.43	0.41
time (sec)	N/A	0.519	0.155	0.748	0.225	0.268	0.000	0.323	13.288

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	140	138	325	0	0	0	0	0	0
N.S.	1	0.99	2.32	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.422	3.025	0.000	0.000	0.000	0.000	0.000	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	65	65	72	0	0	0	0	0	0
N.S.	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.318	0.248	0.000	0.000	0.000	0.000	0.000	0.000

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	180	181	3396	0	0	0	0	0	0
N.S.	1	1.01	18.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.720	14.787	0.000	0.000	0.000	0.000	0.000	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	226	229	187	0	0	0	0	0	0
N.S.	1	1.01	0.83	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.046	1.985	0.000	0.000	0.000	0.000	0.000	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	127	162	189	979	340	0	0	0
N.S.	1	1.22	1.56	1.82	9.41	3.27	0.00	0.00	0.00
time (sec)	N/A	0.450	4.330	5.285	0.455	0.343	0.000	0.000	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	99	73	107	0	260	0	132	0
N.S.	1	1.22	0.90	1.32	0.00	3.21	0.00	1.63	0.00
time (sec)	N/A	0.416	1.677	2.405	0.000	0.281	0.000	0.824	0.000

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	157	724	255	1310	462	0	0	0
N.S.	1	1.12	5.17	1.82	9.36	3.30	0.00	0.00	0.00
time (sec)	N/A	0.464	11.248	5.046	0.449	0.311	0.000	0.000	0.000

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	135	236	124	536	330	0	0	0
N.S.	1	1.16	2.03	1.07	4.62	2.84	0.00	0.00	0.00
time (sec)	N/A	0.481	3.920	3.067	0.418	0.276	0.000	0.000	0.000

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	179	198	328	222	1400	569	0	0	0
N.S.	1	1.11	1.83	1.24	7.82	3.18	0.00	0.00	0.00
time (sec)	N/A	0.530	3.071	5.196	0.453	0.379	0.000	0.000	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	A	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	94	101	56	255	0	0	0
N.S.	1	1.00	2.04	2.20	1.22	5.54	0.00	0.00	0.00
time (sec)	N/A	0.461	1.760	2.581	0.399	0.342	0.000	0.000	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	65	65	98	402	0	357	0	0	0
N.S.	1	1.00	1.51	6.18	0.00	5.49	0.00	0.00	0.00
time (sec)	N/A	0.340	0.592	18.878	0.000	0.423	0.000	0.000	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	252	153	237	379	281	0	566	361
N.S.	1	1.07	0.65	1.00	1.61	1.19	0.00	2.40	1.53
time (sec)	N/A	1.398	1.798	5.160	0.233	0.279	0.000	0.390	17.120

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	182	103	185	266	211	0	380	255
N.S.	1	1.06	0.60	1.08	1.56	1.23	0.00	2.22	1.49
time (sec)	N/A	1.023	0.521	4.002	0.223	0.284	0.000	0.356	17.051

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	114	75	118	165	150	0	232	196
N.S.	1	1.06	0.69	1.09	1.53	1.39	0.00	2.15	1.81
time (sec)	N/A	0.691	0.312	3.179	0.214	0.271	0.000	0.332	16.372

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	59	75	75	88	96	0	124	111
N.S.	1	1.05	1.34	1.34	1.57	1.71	0.00	2.21	1.98
time (sec)	N/A	0.425	0.027	2.195	0.207	0.267	0.000	0.303	14.433

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	107	90	0	255	0	127	195
N.S.	1	1.00	1.55	1.30	0.00	3.70	0.00	1.84	2.83
time (sec)	N/A	0.492	0.761	0.716	0.000	0.305	0.000	0.328	13.829

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	90	75	105	0	357	0	137	85
N.S.	1	1.14	0.95	1.33	0.00	4.52	0.00	1.73	1.08
time (sec)	N/A	0.474	0.769	0.659	0.000	0.286	0.000	0.319	13.473

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	149	167	178	0	736	0	263	171
N.S.	1	1.14	1.27	1.36	0.00	5.62	0.00	2.01	1.31
time (sec)	N/A	0.720	1.537	0.793	0.000	0.310	0.000	0.358	15.425

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	189	216	247	271	0	1278	0	449	321
N.S.	1	1.14	1.31	1.43	0.00	6.76	0.00	2.38	1.70
time (sec)	N/A	1.071	3.648	1.061	0.000	0.359	0.000	0.367	16.761

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	327	390	180	361	683	387	0	736	484
N.S.	1	1.19	0.55	1.10	2.09	1.18	0.00	2.25	1.48
time (sec)	N/A	0.547	6.186	5.925	0.219	0.278	0.000	0.414	17.084

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	242	302	140	269	469	294	0	506	394
N.S.	1	1.25	0.58	1.11	1.94	1.21	0.00	2.09	1.63
time (sec)	N/A	0.458	3.455	5.180	0.220	0.286	0.000	0.390	17.060

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	176	269	100	202	324	209	0	320	237
N.S.	1	1.53	0.57	1.15	1.84	1.19	0.00	1.82	1.35
time (sec)	N/A	0.404	1.675	4.360	0.214	0.272	0.000	0.351	17.204

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	92	63	120	167	138	0	178	161
N.S.	1	0.89	0.61	1.17	1.62	1.34	0.00	1.73	1.56
time (sec)	N/A	0.563	0.730	2.933	0.205	0.271	0.000	0.315	16.210

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	C	A	F(-2)	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	201	329	150	0	398	0	196	529
N.S.	1	2.12	3.46	1.58	0.00	4.19	0.00	2.06	5.57
time (sec)	N/A	0.406	2.848	0.807	0.000	0.415	0.000	0.352	14.258

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	230	312	157	0	567	0	230	2563
N.S.	1	1.97	2.67	1.34	0.00	4.85	0.00	1.97	21.91
time (sec)	N/A	0.415	2.440	0.809	0.000	0.418	0.000	0.363	16.422

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	224	249	167	0	622	0	211	158
N.S.	1	1.72	1.92	1.28	0.00	4.78	0.00	1.62	1.22
time (sec)	N/A	0.363	1.845	0.801	0.000	0.311	0.000	0.387	15.301

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	307	211	228	0	1234	0	403	286
N.S.	1	1.44	0.99	1.07	0.00	5.79	0.00	1.89	1.34
time (sec)	N/A	0.433	3.980	1.054	0.000	0.341	0.000	0.413	16.584

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	276	462	322	352	0	1908	0	710	438
N.S.	1	1.67	1.17	1.28	0.00	6.91	0.00	2.57	1.59
time (sec)	N/A	0.604	8.011	1.439	0.000	0.401	0.000	0.506	17.024

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	288	345	170	355	701	337	0	584	411
N.S.	1	1.20	0.59	1.23	2.43	1.17	0.00	2.03	1.43
time (sec)	N/A	0.477	6.094	6.190	0.242	0.282	0.000	0.441	17.051

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	257	309	130	257	459	245	0	376	287
N.S.	1	1.20	0.51	1.00	1.79	0.95	0.00	1.46	1.12
time (sec)	N/A	0.414	4.434	4.535	0.242	0.282	0.000	0.377	17.100

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	110	173	175	262	161	0	212	203
N.S.	1	0.88	1.38	1.40	2.10	1.29	0.00	1.70	1.62
time (sec)	N/A	0.410	1.044	3.983	0.227	0.289	0.000	0.338	16.880

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	F	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	153	265	419	224	0	532	0	285	1902
N.S.	1	1.73	2.74	1.46	0.00	3.48	0.00	1.86	12.43
time (sec)	N/A	0.491	3.378	0.870	0.000	0.534	0.000	0.367	14.363

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	161	284	455	216	0	859	0	317	3135
N.S.	1	1.76	2.83	1.34	0.00	5.34	0.00	1.97	19.47
time (sec)	N/A	0.477	5.178	1.076	0.000	0.563	0.000	0.366	16.914

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	188	330	393	227	0	1176	0	376	4131
N.S.	1	1.76	2.09	1.21	0.00	6.26	0.00	2.00	21.97
time (sec)	N/A	0.510	4.327	1.070	0.000	0.601	0.000	0.419	20.184

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	178	288	398	227	0	1012	0	307	264
N.S.	1	1.62	2.24	1.28	0.00	5.69	0.00	1.72	1.48
time (sec)	N/A	0.408	3.881	1.215	0.000	0.342	0.000	0.392	17.094

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	266	371	274	303	0	1714	0	601	385
N.S.	1	1.39	1.03	1.14	0.00	6.44	0.00	2.26	1.45
time (sec)	N/A	0.457	6.449	1.695	0.000	0.363	0.000	0.484	17.090

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	305	1243	275	596	297	0	344	211
N.S.	1	1.67	6.79	1.50	3.26	1.62	0.00	1.88	1.15
time (sec)	N/A	0.472	7.845	0.974	0.224	0.274	0.000	0.366	14.169

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	215	78	181	388	216	0	219	139
N.S.	1	1.84	0.67	1.55	3.32	1.85	0.00	1.87	1.19
time (sec)	N/A	0.426	1.604	0.832	0.223	0.284	0.000	0.333	13.678

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	B	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	169	237	103	223	155	0	136	85
N.S.	1	2.49	3.49	1.51	3.28	2.28	0.00	2.00	1.25
time (sec)	N/A	0.336	2.657	0.712	0.217	0.275	0.000	0.313	13.526

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	109	53	99	74	0	70	41
N.S.	1	1.00	2.53	1.23	2.30	1.72	0.00	1.63	0.95
time (sec)	N/A	0.353	0.896	0.664	0.200	0.270	0.000	0.304	13.865

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	154	160	74	0	353	0	110	110
N.S.	1	1.86	1.93	0.89	0.00	4.25	0.00	1.33	1.33
time (sec)	N/A	0.372	1.062	0.700	0.000	0.293	0.000	0.314	13.585

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	234	286	146	0	691	0	221	187
N.S.	1	1.61	1.97	1.01	0.00	4.77	0.00	1.52	1.29
time (sec)	N/A	0.411	3.139	0.795	0.000	0.321	0.000	0.312	13.544

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	A	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	207	328	1422	221	0	1331	0	362	379
N.S.	1	1.58	6.87	1.07	0.00	6.43	0.00	1.75	1.83
time (sec)	N/A	0.507	7.449	0.986	0.000	0.356	0.000	0.379	14.583

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	258	392	446	382	772	456	0	506	268
N.S.	1	1.52	1.73	1.48	2.99	1.77	0.00	1.96	1.04
time (sec)	N/A	0.590	7.419	1.269	0.236	0.309	0.000	0.439	13.554

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	193	302	310	275	536	361	0	359	193
N.S.	1	1.56	1.61	1.42	2.78	1.87	0.00	1.86	1.00
time (sec)	N/A	0.479	5.100	1.013	0.234	0.306	0.000	0.364	13.699

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	222	294	175	342	268	0	250	136
N.S.	1	1.67	2.21	1.32	2.57	2.02	0.00	1.88	1.02
time (sec)	N/A	0.436	3.648	0.843	0.249	0.284	0.000	0.364	13.521

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	178	181	84	195	155	0	158	89
N.S.	1	2.00	2.03	0.94	2.19	1.74	0.00	1.78	1.00
time (sec)	N/A	0.344	2.407	0.652	0.217	0.274	0.000	0.321	13.416

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	64	76	42	93	58	0	60	45
N.S.	1	0.98	1.17	0.65	1.43	0.89	0.00	0.92	0.69
time (sec)	N/A	0.341	0.797	0.694	0.212	0.267	0.000	0.288	13.581

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	129	219	209	122	0	598	0	249	168
N.S.	1	1.70	1.62	0.95	0.00	4.64	0.00	1.93	1.30
time (sec)	N/A	0.426	2.051	0.659	0.000	0.291	0.000	0.335	13.883

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	309	376	203	0	1242	0	474	314
N.S.	1	1.46	1.78	0.96	0.00	5.89	0.00	2.25	1.49
time (sec)	N/A	0.481	3.961	0.794	0.000	0.331	0.000	0.328	14.294

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	284	420	2220	280	0	2030	0	751	505
N.S.	1	1.48	7.82	0.99	0.00	7.15	0.00	2.64	1.78
time (sec)	N/A	0.598	8.182	1.080	0.000	0.363	0.000	0.403	14.200

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	363	501	1338	491	946	620	0	672	327
N.S.	1	1.38	3.69	1.35	2.61	1.71	0.00	1.85	0.90
time (sec)	N/A	0.700	9.806	1.472	0.251	0.305	0.000	0.475	13.625

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	287	400	439	360	689	502	0	504	252
N.S.	1	1.39	1.53	1.25	2.40	1.75	0.00	1.76	0.88
time (sec)	N/A	0.599	5.506	1.067	0.240	0.288	0.000	0.430	13.856

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	310	292	247	475	385	0	374	195
N.S.	1	1.51	1.42	1.20	2.32	1.88	0.00	1.82	0.95
time (sec)	N/A	0.503	3.859	0.936	0.230	0.280	0.000	0.377	13.540

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	225	295	117	307	248	0	259	147
N.S.	1	1.69	2.22	0.88	2.31	1.86	0.00	1.95	1.11
time (sec)	N/A	0.408	2.592	0.693	0.230	0.273	0.000	0.365	13.518

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	192	84	67	184	113	0	129	79
N.S.	1	1.67	0.73	0.58	1.60	0.98	0.00	1.12	0.69
time (sec)	N/A	0.361	0.793	0.639	0.220	0.265	0.000	0.332	13.752

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	99	135	56	115	93	0	75	66
N.S.	1	0.97	1.32	0.55	1.13	0.91	0.00	0.74	0.65
time (sec)	N/A	0.465	0.950	0.779	0.228	0.268	0.000	0.313	13.511

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	294	345	203	0	1001	0	471	228
N.S.	1	1.62	1.91	1.12	0.00	5.53	0.00	2.60	1.26
time (sec)	N/A	0.483	3.164	0.726	0.000	0.304	0.000	0.361	13.800

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	288	392	1772	284	0	1693	0	918	464
N.S.	1	1.36	6.15	0.99	0.00	5.88	0.00	3.19	1.61
time (sec)	N/A	0.590	7.962	0.893	0.000	0.341	0.000	0.391	13.882

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	368	508	1096	365	0	2677	0	1369	655
N.S.	1	1.38	2.98	0.99	0.00	7.27	0.00	3.72	1.78
time (sec)	N/A	0.699	9.054	1.208	0.000	0.395	0.000	0.468	14.089

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	61	61	102	278	0	307	0	0	0
N.S.	1	1.00	1.67	4.56	0.00	5.03	0.00	0.00	0.00
time (sec)	N/A	0.328	0.723	5.513	0.000	0.407	0.000	0.000	0.000

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	140	140	187	478	0	1048	0	0	0
N.S.	1	1.00	1.34	3.41	0.00	7.49	0.00	0.00	0.00
time (sec)	N/A	0.844	14.280	5.300	0.000	0.453	0.000	0.000	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	107	135	0	246	0	0	0
N.S.	1	1.00	1.37	1.73	0.00	3.15	0.00	0.00	0.00
time (sec)	N/A	0.332	0.790	2.860	0.000	0.328	0.000	0.000	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	141	141	171	366	0	1100	0	0	0
N.S.	1	1.00	1.21	2.60	0.00	7.80	0.00	0.00	0.00
time (sec)	N/A	0.871	0.851	5.157	0.000	0.512	0.000	0.000	0.000

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	61	61	94	424	0	343	0	0	0
N.S.	1	1.00	1.54	6.95	0.00	5.62	0.00	0.00	0.00
time (sec)	N/A	0.308	0.387	19.377	0.000	0.445	0.000	0.000	0.000

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	A	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	149	149	188	685	0	1126	0	0	0
N.S.	1	1.00	1.26	4.60	0.00	7.56	0.00	0.00	0.00
time (sec)	N/A	0.937	2.035	22.714	0.000	4.814	0.000	0.000	0.000

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	122	122	144	504	0	963	0	0	0
N.S.	1	1.00	1.18	4.13	0.00	7.89	0.00	0.00	0.00
time (sec)	N/A	0.657	1.433	19.783	0.000	0.488	0.000	0.000	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	124	124	141	504	0	1041	0	0	0
N.S.	1	1.00	1.14	4.06	0.00	8.40	0.00	0.00	0.00
time (sec)	N/A	0.696	0.717	18.953	0.000	0.572	0.000	0.000	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	167	167	198	557	0	1103	0	0	0
N.S.	1	1.00	1.19	3.34	0.00	6.60	0.00	0.00	0.00
time (sec)	N/A	0.948	1.756	20.660	0.000	0.800	0.000	0.000	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	231	245	155	796	0	1597	0	0	0
N.S.	1	1.06	0.67	3.45	0.00	6.91	0.00	0.00	0.00
time (sec)	N/A	1.413	1.489	22.869	0.000	52.752	0.000	0.000	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	250	266	201	237	379	281	0	850	555
N.S.	1	1.06	0.80	0.95	1.52	1.12	0.00	3.40	2.22
time (sec)	N/A	1.415	2.840	5.139	0.232	0.295	0.000	0.367	17.353

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	191	143	185	266	211	0	586	395
N.S.	1	1.06	0.79	1.03	1.48	1.17	0.00	3.26	2.19
time (sec)	N/A	1.047	0.775	3.814	0.255	0.276	0.000	0.361	17.202

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	121	88	118	165	150	0	294	227
N.S.	1	1.05	0.77	1.03	1.43	1.30	0.00	2.56	1.97
time (sec)	N/A	0.686	0.453	3.395	0.225	0.278	0.000	0.339	17.016

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	64	75	75	88	96	0	153	104
N.S.	1	1.05	1.23	1.23	1.44	1.57	0.00	2.51	1.70
time (sec)	N/A	0.429	0.018	2.275	0.229	0.291	0.000	0.308	14.561

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	112	92	0	316	0	127	573
N.S.	1	1.00	1.47	1.21	0.00	4.16	0.00	1.67	7.54
time (sec)	N/A	0.479	0.480	0.768	0.000	0.632	0.000	0.346	14.539

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	110	97	132	0	389	0	172	106
N.S.	1	1.11	0.98	1.33	0.00	3.93	0.00	1.74	1.07
time (sec)	N/A	0.500	0.427	0.770	0.000	0.306	0.000	0.313	13.878

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	191	172	236	0	752	0	399	250
N.S.	1	1.15	1.04	1.42	0.00	4.53	0.00	2.40	1.51
time (sec)	N/A	0.802	1.136	0.884	0.000	0.325	0.000	0.375	16.701

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	278	405	376	0	1238	0	693	439
N.S.	1	1.17	1.71	1.59	0.00	5.22	0.00	2.92	1.85
time (sec)	N/A	1.185	2.012	1.283	0.000	0.357	0.000	0.404	18.527

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	247	247	580	480	0	1093	0	606	9987
N.S.	1	1.00	2.35	1.94	0.00	4.43	0.00	2.45	40.43
time (sec)	N/A	0.722	5.682	1.437	0.000	102.849	0.000	0.405	23.128

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	170	389	289	0	779	0	339	6730
N.S.	1	1.00	2.29	1.70	0.00	4.58	0.00	1.99	39.59
time (sec)	N/A	0.622	3.112	1.038	0.000	22.065	0.000	0.397	21.040

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	103	135	165	0	518	0	195	3559
N.S.	1	1.00	1.31	1.60	0.00	5.03	0.00	1.89	34.55
time (sec)	N/A	0.539	2.509	0.946	0.000	3.298	0.000	0.351	18.706

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	112	92	0	309	0	127	571
N.S.	1	1.00	1.47	1.21	0.00	4.07	0.00	1.67	7.51
time (sec)	N/A	0.474	0.439	0.742	0.000	0.570	0.000	0.337	14.615

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	121	119	108	0	1040	0	522	2665
N.S.	1	1.00	0.98	0.89	0.00	8.60	0.00	4.31	22.02
time (sec)	N/A	0.543	0.444	1.025	0.000	1.855	0.000	0.397	15.932

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	187	221	229	208	0	2863	0	331	20827
N.S.	1	1.18	1.22	1.11	0.00	15.31	0.00	1.77	111.37
time (sec)	N/A	0.969	1.337	1.371	0.000	96.827	0.000	0.356	27.240

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	F(-1)	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	379	379	784	689	0	0	0	857	17256
N.S.	1	1.00	2.07	1.82	0.00	0.00	0.00	2.26	45.53
time (sec)	N/A	0.986	10.090	2.507	0.000	0.000	0.000	0.449	28.858

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	297	511	465	0	1925	0	551	12483
N.S.	1	1.00	1.72	1.57	0.00	6.48	0.00	1.86	42.03
time (sec)	N/A	0.820	5.400	1.725	0.000	170.550	0.000	0.390	26.195

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	228	362	314	0	1326	0	539	7958
N.S.	1	1.00	1.59	1.38	0.00	5.82	0.00	2.36	34.90
time (sec)	N/A	0.717	3.408	1.467	0.000	39.313	0.000	0.379	22.917

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	198	180	215	0	798	0	267	4926
N.S.	1	1.00	0.91	1.09	0.00	4.03	0.00	1.35	24.88
time (sec)	N/A	0.635	1.238	0.936	0.000	5.697	0.000	0.365	21.295

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	111	97	132	0	394	0	173	106
N.S.	1	1.11	0.97	1.32	0.00	3.94	0.00	1.73	1.06
time (sec)	N/A	0.490	0.436	0.648	0.000	0.280	0.000	0.315	13.753

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	218	176	210	0	2852	0	330	20827
N.S.	1	1.17	0.95	1.13	0.00	15.33	0.00	1.77	111.97
time (sec)	N/A	0.981	1.587	1.452	0.000	60.792	0.000	0.364	27.475

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	213	183	316	0	0	0	0	0
N.S.	1	1.00	0.86	1.48	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.714	4.103	9.154	0.000	0.000	0.000	0.000	0.000

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	196	196	44664	321	0	0	0	0	0
N.S.	1	1.00	227.88	1.64	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.433	37.864	12.835	0.000	0.000	0.000	0.000	0.000

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	192	192	233	191	0	0	0	0	0
N.S.	1	1.00	1.21	0.99	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.416	3.197	10.710	0.000	0.000	0.000	0.000	0.000

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	176	148	0	0	0	0	0
N.S.	1	1.00	1.60	1.35	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.326	2.645	7.537	0.000	0.000	0.000	0.000	0.000

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	176	142	0	0	0	0	0
N.S.	1	1.00	1.41	1.14	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.327	0.649	7.361	0.000	0.000	0.000	0.000	0.000

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	396	396	39359	262	0	0	0	0	0
N.S.	1	1.00	99.39	0.66	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.096	36.528	15.931	0.000	0.000	0.000	0.000	0.000

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	170	223	445	0	0	0	0	0
N.S.	1	1.00	1.31	2.62	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.052	25.506	8.494	0.000	0.000	0.000	0.000	0.000

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	83	83	223	0	0	0	0	0
N.S.	1	1.00	1.00	2.69	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.726	1.349	7.344	0.000	0.000	0.000	0.000	0.000

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	174	222	442	0	0	0	0	0
N.S.	1	1.04	1.32	2.63	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.331	25.117	8.237	0.000	0.000	0.000	0.000	0.000

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	264	123	0	0	0	0	0
N.S.	1	1.00	2.78	1.29	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.332	6.443	7.297	0.000	0.000	0.000	0.000	0.000

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	C	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	295	311	0	275	0	0	0	0	0
N.S.	1	1.05	0.00	0.93	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.755	0.000	6.872	0.000	0.000	0.000	0.000	0.000

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	209	209	375	191	0	0	0	0	0
N.S.	1	1.00	1.79	0.91	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.730	13.338	5.330	0.000	0.000	0.000	0.000	0.000

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	214	156	192	0	0	0	0	0
N.S.	1	1.00	0.73	0.90	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.799	5.477	8.774	0.000	0.000	0.000	0.000	0.000

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	229	230	1019	207	0	517	0	0	0
N.S.	1	1.00	4.45	0.90	0.00	2.26	0.00	0.00	0.00
time (sec)	N/A	1.268	8.387	4.700	0.000	0.118	0.000	0.000	0.000

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	C	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	312	310	0	323	0	0	0	0	0
N.S.	1	0.99	0.00	1.04	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.651	0.000	7.967	0.000	0.000	0.000	0.000	0.000

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	213	183	316	0	0	0	0	0
N.S.	1	1.00	0.86	1.48	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.704	0.236	8.320	0.000	0.000	0.000	0.000	0.000

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	170	223	445	0	0	0	0	0
N.S.	1	1.00	1.31	2.62	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.058	25.557	8.221	0.000	0.000	0.000	0.000	0.000

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	102	187	203	0	0	0	0	0
N.S.	1	1.00	1.83	1.99	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.342	8.833	7.645	0.000	0.000	0.000	0.000	0.000

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	209	165	203	0	0	0	0	0
N.S.	1	1.00	0.79	0.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.783	4.289	8.545	0.000	0.000	0.000	0.000	0.000

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	83	83	223	0	0	0	0	0
N.S.	1	1.00	1.00	2.69	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.723	1.378	7.462	0.000	0.000	0.000	0.000	0.000

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	166	246	313	0	0	0	0	0
N.S.	1	1.00	1.48	1.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.075	25.394	9.418	0.000	0.000	0.000	0.000	0.000

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	59	151	49	68	121	0	47	47
N.S.	1	0.88	2.25	0.73	1.01	1.81	0.00	0.70	0.70
time (sec)	N/A	0.417	4.031	11.267	0.218	0.312	0.000	1.104	13.517

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	75	175	62	88	146	0	60	60
N.S.	1	0.84	1.97	0.70	0.99	1.64	0.00	0.67	0.67
time (sec)	N/A	0.441	5.022	17.280	0.224	0.288	0.000	1.174	14.181

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [274] had the largest ratio of [.538461999999999996]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	0.99	30	0.100
2	A	3	3	1.00	30	0.100
3	A	3	3	0.92	30	0.100
4	A	6	6	1.00	28	0.214
5	A	4	4	1.00	30	0.133
6	A	2	2	1.00	30	0.067
7	A	4	4	1.00	30	0.133
8	A	6	6	1.03	30	0.200
9	A	8	8	1.04	30	0.267
10	A	3	3	0.92	32	0.094
11	A	3	3	0.93	32	0.094
12	A	3	3	0.86	32	0.094
13	A	8	8	0.92	32	0.250
14	A	3	3	0.92	30	0.100
15	A	9	8	0.93	32	0.250
16	A	6	6	1.00	32	0.188
17	A	2	2	1.00	32	0.062
18	A	4	4	1.00	32	0.125
19	A	6	6	1.04	32	0.188
20	A	8	8	1.06	32	0.250
21	A	3	3	0.92	32	0.094
22	A	3	3	0.92	32	0.094

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	3	3	0.86	32	0.094
24	A	10	10	0.94	32	0.312
25	A	3	3	0.86	32	0.094
26	A	3	3	1.00	30	0.100
27	A	11	10	0.98	32	0.312
28	A	11	10	0.98	32	0.312
29	A	8	8	1.02	32	0.250
30	A	2	2	1.00	32	0.062
31	A	4	4	1.00	32	0.125
32	A	6	6	1.04	32	0.188
33	A	8	8	1.06	32	0.250
34	A	5	5	0.96	32	0.156
35	A	11	10	0.98	32	0.312
36	A	9	8	0.95	32	0.250
37	A	4	4	1.00	30	0.133
38	A	7	6	1.00	32	0.188
39	A	3	3	0.86	32	0.094
40	A	3	3	0.95	32	0.094
41	A	3	3	0.91	32	0.094
42	A	7	7	0.99	32	0.219
43	A	13	12	0.97	32	0.375
44	A	11	10	0.99	32	0.312
45	A	6	6	1.00	32	0.188
46	A	2	2	1.00	30	0.067
47	A	3	3	0.86	32	0.094
48	A	7	6	0.84	32	0.188
49	A	3	3	0.86	32	0.094
50	A	3	3	0.92	32	0.094
51	A	3	3	0.91	32	0.094
52	A	9	9	0.98	32	0.281
53	A	15	14	1.01	32	0.438
54	A	13	12	1.01	32	0.375
55	A	8	8	1.02	32	0.250
56	A	2	2	1.00	32	0.062

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	4	4	1.00	30	0.133
58	A	3	3	0.95	32	0.094
59	A	3	3	0.86	32	0.094
60	A	8	7	0.69	32	0.219
61	A	3	3	0.82	32	0.094
62	A	3	3	0.92	32	0.094
63	A	3	3	0.90	32	0.094
64	A	8	8	1.05	32	0.250
65	A	6	6	1.03	32	0.188
66	A	4	4	1.00	32	0.125
67	A	2	2	1.00	32	0.062
68	A	6	5	1.00	32	0.156
69	A	6	5	1.00	32	0.156
70	A	8	7	1.04	32	0.219
71	A	8	8	1.05	34	0.235
72	A	6	6	1.03	34	0.176
73	A	4	4	1.00	34	0.118
74	A	2	2	1.00	34	0.059
75	A	8	7	1.05	34	0.206
76	A	8	7	1.12	34	0.206
77	A	8	7	1.09	34	0.206
78	A	10	9	1.02	34	0.265
79	A	8	8	1.05	34	0.235
80	A	6	6	1.03	34	0.176
81	A	4	4	1.00	34	0.118
82	A	2	2	1.00	34	0.059
83	A	10	9	1.02	34	0.265
84	A	10	9	1.02	34	0.265
85	A	10	9	1.01	34	0.265
86	A	8	8	1.00	34	0.235
87	A	6	6	0.99	34	0.176
88	A	4	4	1.00	34	0.118
89	A	2	2	1.00	34	0.059
90	A	6	5	1.00	34	0.147

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
91	A	8	7	1.02	34	0.206
92	A	10	9	1.03	34	0.265
93	A	8	8	1.00	34	0.235
94	A	6	6	0.99	34	0.176
95	A	4	4	0.99	34	0.118
96	A	2	2	1.00	34	0.059
97	A	8	7	0.99	34	0.206
98	A	10	9	1.02	34	0.265
99	A	12	11	1.02	34	0.324
100	A	8	8	1.02	34	0.235
101	A	6	6	1.02	34	0.176
102	A	4	4	1.00	34	0.118
103	A	2	2	1.00	34	0.059
104	A	10	9	1.02	34	0.265
105	A	12	11	1.04	34	0.324
106	A	14	13	1.04	34	0.382
107	A	2	2	1.00	36	0.056
108	A	2	2	1.00	36	0.056
109	A	2	2	1.00	36	0.056
110	A	2	2	1.00	36	0.056
111	A	2	2	1.00	36	0.056
112	A	2	2	1.00	36	0.056
113	A	4	4	1.00	36	0.111
114	A	4	4	1.00	36	0.111
115	A	4	4	1.00	36	0.111
116	A	2	2	1.00	36	0.056
117	A	4	4	1.00	36	0.111
118	A	4	4	1.00	36	0.111
119	A	2	2	1.00	36	0.056
120	A	4	4	1.00	36	0.111
121	A	4	4	1.00	36	0.111
122	A	4	4	1.00	36	0.111
123	A	6	6	1.03	36	0.167
124	A	6	6	1.03	36	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
125	A	4	4	1.00	36	0.111
126	A	2	2	1.00	36	0.056
127	A	6	6	1.01	36	0.167
128	A	6	6	0.99	36	0.167
129	A	6	6	1.03	36	0.167
130	A	2	2	1.00	36	0.056
131	A	4	4	1.00	36	0.111
132	A	6	6	1.04	36	0.167
133	A	6	6	1.01	36	0.167
134	A	4	4	1.00	36	0.111
135	A	2	2	1.00	36	0.056
136	A	5	5	1.00	36	0.139
137	A	7	7	1.00	36	0.194
138	A	9	9	1.04	36	0.250
139	A	6	6	0.99	36	0.167
140	A	4	4	1.00	36	0.111
141	A	2	2	1.00	36	0.056
142	A	7	7	1.00	36	0.194
143	A	7	7	0.74	36	0.194
144	A	9	9	0.84	36	0.250
145	A	6	6	1.00	36	0.167
146	A	2	2	1.00	36	0.056
147	A	2	2	1.00	36	0.056
148	A	9	9	1.04	36	0.250
149	A	9	9	0.84	36	0.250
150	A	9	9	0.64	36	0.250
151	A	5	4	1.00	32	0.125
152	A	5	4	1.00	32	0.125
153	A	5	4	1.00	30	0.133
154	A	5	4	1.00	32	0.125
155	A	5	4	1.00	32	0.125
156	A	6	6	0.99	34	0.176
157	A	4	4	1.02	34	0.118
158	A	2	2	1.00	34	0.059

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
159	A	5	4	1.00	34	0.118
160	A	5	4	1.00	34	0.118
161	A	5	4	1.00	34	0.118
162	A	6	6	0.99	36	0.167
163	A	4	4	1.02	36	0.111
164	A	2	2	1.00	36	0.056
165	A	5	4	1.00	34	0.118
166	A	5	4	1.00	36	0.111
167	A	5	4	1.00	36	0.111
168	A	3	3	0.99	32	0.094
169	A	3	3	0.91	32	0.094
170	A	6	5	1.00	30	0.167
171	A	10	9	1.02	32	0.281
172	A	8	8	1.04	32	0.250
173	A	7	7	1.02	32	0.219
174	A	3	3	0.99	34	0.088
175	A	4	4	1.00	32	0.125
176	A	8	8	1.01	34	0.235
177	A	10	10	1.01	34	0.294
178	A	6	5	1.22	40	0.125
179	A	7	6	1.22	36	0.167
180	A	11	10	1.12	38	0.263
181	A	7	6	1.16	40	0.150
182	A	10	9	1.11	40	0.225
183	A	7	6	1.00	38	0.158
184	A	4	3	1.00	34	0.088
185	A	15	14	1.07	29	0.483
186	A	13	12	1.06	29	0.414
187	A	11	10	1.06	29	0.345
188	A	9	8	1.05	27	0.296
189	A	9	8	1.00	29	0.276
190	A	10	9	1.14	29	0.310
191	A	13	12	1.14	29	0.414
192	A	15	14	1.14	29	0.483

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
193	A	14	13	1.19	31	0.419
194	A	11	10	1.25	31	0.323
195	A	11	10	1.53	31	0.323
196	A	11	10	0.89	29	0.345
197	B	10	9	2.12	31	0.290
198	A	10	9	1.97	31	0.290
199	A	7	6	1.72	31	0.194
200	A	8	7	1.44	31	0.226
201	A	14	13	1.67	31	0.419
202	A	12	11	1.20	31	0.355
203	A	12	11	1.20	31	0.355
204	A	5	5	0.88	29	0.172
205	A	13	12	1.73	31	0.387
206	A	12	11	1.76	31	0.355
207	A	12	11	1.76	31	0.355
208	A	8	7	1.62	31	0.226
209	A	9	8	1.39	31	0.258
210	A	12	11	1.67	31	0.355
211	A	9	8	1.84	31	0.258
212	B	8	7	2.49	31	0.226
213	A	5	5	1.00	29	0.172
214	A	6	5	1.86	31	0.161
215	A	9	8	1.61	31	0.258
216	A	11	10	1.58	31	0.323
217	A	14	13	1.52	31	0.419
218	A	11	10	1.56	31	0.323
219	A	9	8	1.67	31	0.258
220	A	8	7	2.00	31	0.226
221	A	4	4	0.98	29	0.138
222	A	10	9	1.70	31	0.290
223	A	12	11	1.46	31	0.355
224	A	14	13	1.48	31	0.419
225	A	16	15	1.38	31	0.484
226	A	13	12	1.39	31	0.387

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
227	A	11	10	1.51	31	0.323
228	A	9	8	1.69	31	0.258
229	A	7	6	1.67	31	0.194
230	A	6	6	0.97	29	0.207
231	A	13	12	1.62	31	0.387
232	A	15	14	1.36	31	0.452
233	A	17	16	1.38	31	0.516
234	A	4	3	1.00	35	0.086
235	A	8	7	1.00	35	0.200
236	A	4	3	1.00	35	0.086
237	A	8	7	1.00	37	0.189
238	A	4	3	1.00	33	0.091
239	A	8	7	1.00	39	0.179
240	A	8	7	1.00	33	0.212
241	A	8	7	1.00	35	0.200
242	A	8	7	1.00	39	0.179
243	A	12	11	1.06	39	0.282
244	A	15	14	1.06	29	0.483
245	A	13	12	1.06	29	0.414
246	A	11	10	1.05	29	0.345
247	A	9	8	1.05	27	0.296
248	A	9	8	1.00	29	0.276
249	A	10	9	1.11	29	0.310
250	A	12	11	1.15	29	0.379
251	A	15	14	1.17	29	0.483
252	A	5	5	1.00	31	0.161
253	A	5	5	1.00	31	0.161
254	A	5	5	1.00	31	0.161
255	A	9	8	1.00	29	0.276
256	A	8	7	1.00	31	0.226
257	A	11	10	1.18	31	0.323
258	A	5	5	1.00	31	0.161
259	A	5	5	1.00	31	0.161
260	A	5	5	1.00	31	0.161

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
261	A	5	5	1.00	31	0.161
262	A	10	9	1.11	29	0.310
263	A	11	10	1.17	31	0.323
264	A	5	5	1.00	33	0.152
265	A	2	2	1.00	35	0.057
266	A	2	2	1.00	35	0.057
267	A	2	2	1.00	35	0.057
268	A	2	2	1.00	35	0.057
269	A	5	5	1.00	37	0.135
270	A	13	13	1.00	39	0.333
271	A	6	6	1.00	39	0.154
272	A	15	15	1.04	39	0.385
273	A	2	2	1.00	33	0.061
274	A	21	21	1.05	39	0.538
275	A	5	5	1.00	33	0.152
276	A	5	5	1.00	35	0.143
277	A	14	14	1.00	39	0.359
278	A	21	21	0.99	39	0.538
279	A	5	5	1.00	33	0.152
280	A	13	13	1.00	39	0.333
281	A	2	2	1.00	33	0.061
282	A	5	5	1.00	35	0.143
283	A	6	6	1.00	39	0.154
284	A	13	13	1.00	39	0.333
285	A	6	5	0.88	28	0.179
286	A	6	5	0.84	28	0.179

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$	116
3.2	$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$	122
3.3	$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$	128
3.4	$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$	134
3.5	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c-c \sec(e+fx)} dx$	140
3.6	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^2} dx$	146
3.7	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^3} dx$	151
3.8	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^4} dx$	157
3.9	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^5} dx$	163
3.10	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$	170
3.11	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$	177
3.12	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx$	183
3.13	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$	189
3.14	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$	195
3.15	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c-c \sec(e+fx)} dx$	201
3.16	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^2} dx$	207
3.17	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^3} dx$	213
3.18	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^4} dx$	218
3.19	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^5} dx$	224
3.20	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^6} dx$	230
3.21	$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$	238
3.22	$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$	245
3.23	$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$	252
3.24	$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$	258
3.25	$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx$	265
3.26	$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$	271

3.27	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c-c \sec(e+fx)} dx$	277
3.28	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^2} dx$	285
3.29	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^3} dx$	293
3.30	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^4} dx$	300
3.31	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^5} dx$	305
3.32	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^6} dx$	311
3.33	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^7} dx$	318
3.34	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^4}{a+a \sec(e+fx)} dx$	326
3.35	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^3}{a+a \sec(e+fx)} dx$	333
3.36	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^2}{a+a \sec(e+fx)} dx$	341
3.37	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))}{a+a \sec(e+fx)} dx$	347
3.38	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))} dx$	352
3.39	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^2} dx$	357
3.40	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^3} dx$	362
3.41	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx$	367
3.42	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^5}{(a+a \sec(e+fx))^2} dx$	372
3.43	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^4}{(a+a \sec(e+fx))^2} dx$	380
3.44	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^3}{(a+a \sec(e+fx))^2} dx$	388
3.45	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^2}{(a+a \sec(e+fx))^2} dx$	396
3.46	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))}{(a+a \sec(e+fx))^2} dx$	402
3.47	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))} dx$	407
3.48	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^2} dx$	412
3.49	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^3} dx$	417
3.50	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^4} dx$	422
3.51	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^5} dx$	428
3.52	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$	434
3.53	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$	442
3.54	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^4}{(a+a \sec(e+fx))^3} dx$	452
3.55	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^3}{(a+a \sec(e+fx))^3} dx$	461
3.56	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^2}{(a+a \sec(e+fx))^3} dx$	468
3.57	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))}{(a+a \sec(e+fx))^3} dx$	473
3.58	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))} dx$	478
3.59	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^2} dx$	483

3.60	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^3} dx$	488
3.61	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^4} dx$	494
3.62	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^5} dx$	500
3.63	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^6} dx$	506
3.64	$\int \sec(e+fx)(a+a \sec(e+fx))(c-c \sec(e+fx))^{7/2} dx$	512
3.65	$\int \sec(e+fx)(a+a \sec(e+fx))(c-c \sec(e+fx))^{5/2} dx$	520
3.66	$\int \sec(e+fx)(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2} dx$	527
3.67	$\int \sec(e+fx)(a+a \sec(e+fx))\sqrt{c-c \sec(e+fx)} dx$	533
3.68	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{\sqrt{c-c \sec(e+fx)}} dx$	538
3.69	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{3/2}} dx$	544
3.70	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{5/2}} dx$	550
3.71	$\int \sec(e+fx)(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{7/2} dx$	557
3.72	$\int \sec(e+fx)(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{5/2} dx$	564
3.73	$\int \sec(e+fx)(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{3/2} dx$	571
3.74	$\int \sec(e+fx)(a+a \sec(e+fx))^2\sqrt{c-c \sec(e+fx)} dx$	578
3.75	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{\sqrt{c-c \sec(e+fx)}} dx$	584
3.76	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{3/2}} dx$	591
3.77	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{5/2}} dx$	598
3.78	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{7/2}} dx$	604
3.79	$\int \sec(e+fx)(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{7/2} dx$	612
3.80	$\int \sec(e+fx)(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{5/2} dx$	620
3.81	$\int \sec(e+fx)(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{3/2} dx$	627
3.82	$\int \sec(e+fx)(a+a \sec(e+fx))^3\sqrt{c-c \sec(e+fx)} dx$	633
3.83	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{\sqrt{c-c \sec(e+fx)}} dx$	639
3.84	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{3/2}} dx$	647
3.85	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{5/2}} dx$	655
3.86	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{7/2}}{a+a \sec(e+fx)} dx$	663
3.87	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{5/2}}{a+a \sec(e+fx)} dx$	669
3.88	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{3/2}}{a+a \sec(e+fx)} dx$	675
3.89	$\int \frac{\sec(e+fx)\sqrt{c-c \sec(e+fx)}}{a+a \sec(e+fx)} dx$	680
3.90	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))\sqrt{c-c \sec(e+fx)}} dx$	685
3.91	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2}} dx$	691
3.92	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{5/2}} dx$	697
3.93	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{7/2}}{(a+a \sec(e+fx))^2} dx$	704
3.94	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{5/2}}{(a+a \sec(e+fx))^2} dx$	710

3.95	$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx$	716
3.96	$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx$	721
3.97	$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2\sqrt{c-c\sec(e+fx)}} dx$	726
3.98	$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^{3/2}} dx$	732
3.99	$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^{5/2}} dx$	739
3.100	$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx$	748
3.101	$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx$	755
3.102	$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx$	762
3.103	$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx$	768
3.104	$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}} dx$	773
3.105	$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx$	780
3.106	$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx$	789
3.107	$\int \sec(e+fx)\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2} dx$	801
3.108	$\int \sec(e+fx)\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{3/2} dx$	806
3.109	$\int \sec(e+fx)\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)} dx$	811
3.110	$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx$	816
3.111	$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx$	821
3.112	$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx$	826
3.113	$\int \sec(e+fx)(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{7/2} dx$	832
3.114	$\int \sec(e+fx)(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{5/2} dx$	838
3.115	$\int \sec(e+fx)(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{3/2} dx$	844
3.116	$\int \sec(e+fx)(a+a\sec(e+fx))^{3/2}\sqrt{c-c\sec(e+fx)} dx$	850
3.117	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx$	855
3.118	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx$	860
3.119	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx$	866
3.120	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{7/2}} dx$	871
3.121	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx$	877
3.122	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx$	883
3.123	$\int \sec(e+fx)(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{7/2} dx$	889
3.124	$\int \sec(e+fx)(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{5/2} dx$	896
3.125	$\int \sec(e+fx)(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{3/2} dx$	903
3.126	$\int \sec(e+fx)(a+a\sec(e+fx))^{5/2}\sqrt{c-c\sec(e+fx)} dx$	909
3.127	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{\sqrt{c-c\sec(e+fx)}} dx$	914
3.128	$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{3/2}} dx$	921

3.129	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{5/2}} dx$	928
3.130	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{7/2}} dx$	934
3.131	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{9/2}} dx$	940
3.132	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{11/2}} dx$	946
3.133	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}} dx$	952
3.134	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}} dx$	959
3.135	$\int \frac{\sec(e+fx)\sqrt{c-c \sec(e+fx)}}{\sqrt{a+a \sec(e+fx)}} dx$	964
3.136	$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx$	969
3.137	$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}} dx$	975
3.138	$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{5/2}} dx$	982
3.139	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{5/2}}{(a+a \sec(e+fx))^{3/2}} dx$	989
3.140	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{3/2}}{(a+a \sec(e+fx))^{3/2}} dx$	996
3.141	$\int \frac{\sec(e+fx)\sqrt{c-c \sec(e+fx)}}{(a+a \sec(e+fx))^{3/2}} dx$	1001
3.142	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}\sqrt{c-c \sec(e+fx)}} dx$	1006
3.143	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2}} dx$	1013
3.144	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{5/2}} dx$	1019
3.145	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{5/2}}{(a+a \sec(e+fx))^{5/2}} dx$	1026
3.146	$\int \frac{\sec(e+fx)(c-c \sec(e+fx))^{3/2}}{(a+a \sec(e+fx))^{5/2}} dx$	1032
3.147	$\int \frac{\sec(e+fx)\sqrt{c-c \sec(e+fx)}}{(a+a \sec(e+fx))^{5/2}} dx$	1037
3.148	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}\sqrt{c-c \sec(e+fx)}} dx$	1042
3.149	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2}} dx$	1049
3.150	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2}} dx$	1056
3.151	$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^n dx$	1063
3.152	$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^2 dx$	1068
3.153	$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx)) dx$	1073
3.154	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{c-c \sec(e+fx)} dx$	1078
3.155	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^2} dx$	1083
3.156	$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{5/2} dx$	1088
3.157	$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{3/2} dx$	1094
3.158	$\int \sec(e+fx)(a+a \sec(e+fx))^m\sqrt{c-c \sec(e+fx)} dx$	1100
3.159	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{\sqrt{c-c \sec(e+fx)}} dx$	1105
3.160	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{3/2}} dx$	1110
3.161	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{5/2}} dx$	1115

3.162	$\int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx))^{-3-m} dx$	1120
3.163	$\int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx))^{-2-m} dx$	1126
3.164	$\int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx))^{-1-m} dx$	1132
3.165	$\int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx))^{-m} dx$	1137
3.166	$\int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx))^{1-m} dx$	1142
3.167	$\int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx))^{2-m} dx$	1147
3.168	$\int \sec^2(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$	1152
3.169	$\int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$	1158
3.170	$\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$	1164
3.171	$\int \frac{\sec^2(e+fx)(c-c \sec(e+fx))}{a+a \sec(e+fx)} dx$	1170
3.172	$\int \frac{\sec^2(e+fx)(c-c \sec(e+fx))}{(a+a \sec(e+fx))^2} dx$	1177
3.173	$\int \frac{\sec^2(e+fx)(c-c \sec(e+fx))}{(a+a \sec(e+fx))^3} dx$	1183
3.174	$\int (g \sec(e + fx))^p(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$	1189
3.175	$\int (g \sec(e + fx))^p(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$	1194
3.176	$\int \frac{(g \sec(e+fx))^p(c-c \sec(e+fx))}{a+a \sec(e+fx)} dx$	1199
3.177	$\int \frac{(g \sec(e+fx))^p(c-c \sec(e+fx))}{(a+a \sec(e+fx))^2} dx$	1206
3.178	$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c-c \sec(e+fx)} dx$	1213
3.179	$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))}} dx$	1220
3.180	$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))}} dx$	1226
3.181	$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))}} dx$	1236
3.182	$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))}} dx$	1243
3.183	$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} dx$	1252
3.184	$\int \frac{\sec(e+fx) \sqrt{a+a \sec(e+fx)}}{c-d \sec(e+fx)} dx$	1258
3.185	$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$	1263
3.186	$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$	1273
3.187	$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$	1282
3.188	$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$	1290
3.189	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c+d \sec(e+fx)} dx$	1297
3.190	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^2} dx$	1304
3.191	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^3} dx$	1311
3.192	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^4} dx$	1319
3.193	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$	1328
3.194	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx$	1339
3.195	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$	1349
3.196	$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$	1358
3.197	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c+d \sec(e+fx)} dx$	1365

3.198	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^2} dx$	1374
3.199	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^3} dx$	1384
3.200	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^4} dx$	1392
3.201	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^5} dx$	1401
3.202	$\int \sec(e+fx)(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3 dx$	1414
3.203	$\int \sec(e+fx)(a+a \sec(e+fx))^3(c+d \sec(e+fx))^2 dx$	1425
3.204	$\int \sec(e+fx)(a+a \sec(e+fx))^3(c+d \sec(e+fx)) dx$	1434
3.205	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c+d \sec(e+fx)} dx$	1441
3.206	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^2} dx$	1452
3.207	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^3} dx$	1463
3.208	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^4} dx$	1473
3.209	$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^5} dx$	1483
3.210	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+a \sec(e+fx)} dx$	1494
3.211	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+a \sec(e+fx)} dx$	1504
3.212	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+a \sec(e+fx)} dx$	1512
3.213	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+a \sec(e+fx)} dx$	1519
3.214	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx$	1525
3.215	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$	1531
3.216	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$	1539
3.217	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^2} dx$	1549
3.218	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^2} dx$	1560
3.219	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^2} dx$	1570
3.220	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^2} dx$	1579
3.221	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^2} dx$	1586
3.222	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))} dx$	1591
3.223	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2} dx$	1599
3.224	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^3} dx$	1609
3.225	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$	1621
3.226	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$	1634
3.227	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^3} dx$	1646
3.228	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^3} dx$	1655
3.229	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^3} dx$	1664
3.230	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^3} dx$	1671
3.231	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))} dx$	1677

3.232	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^2} dx$	1687
3.233	$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$	1699
3.234	$\int \frac{\sec(e+fx)\sqrt{a+a \sec(e+fx)}}{\sqrt{c+d \sec(e+fx)}} dx$	1713
3.235	$\int \frac{\sec(e+fx)\sqrt{c+d \sec(e+fx)}}{\sqrt{a+a \sec(e+fx)}} dx$	1718
3.236	$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$	1725
3.237	$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$	1730
3.238	$\int \frac{\sec(e+fx)\sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx$	1738
3.239	$\int \frac{(g \sec(e+fx))^{3/2}\sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx$	1743
3.240	$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$	1750
3.241	$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$	1757
3.242	$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$	1764
3.243	$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$	1771
3.244	$\int \sec(e+fx)(a+b \sec(e+fx))(c+d \sec(e+fx))^4 dx$	1780
3.245	$\int \sec(e+fx)(a+b \sec(e+fx))(c+d \sec(e+fx))^3 dx$	1790
3.246	$\int \sec(e+fx)(a+b \sec(e+fx))(c+d \sec(e+fx))^2 dx$	1799
3.247	$\int \sec(e+fx)(a+b \sec(e+fx))(c+d \sec(e+fx)) dx$	1807
3.248	$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{c+d \sec(e+fx)} dx$	1814
3.249	$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^2} dx$	1822
3.250	$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^3} dx$	1829
3.251	$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^4} dx$	1837
3.252	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+b \sec(e+fx)} dx$	1847
3.253	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+b \sec(e+fx)} dx$	1856
3.254	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+b \sec(e+fx)} dx$	1863
3.255	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+b \sec(e+fx)} dx$	1870
3.256	$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))} dx$	1878
3.257	$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))^2} dx$	1886
3.258	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+b \sec(e+fx))^2} dx$	1894
3.259	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+b \sec(e+fx))^2} dx$	1903
3.260	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+b \sec(e+fx))^2} dx$	1912
3.261	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+b \sec(e+fx))^2} dx$	1920
3.262	$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+b \sec(e+fx))^2} dx$	1927
3.263	$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))^2(c+d \sec(e+fx))} dx$	1934
3.264	$\int \frac{\sec(e+fx)\sqrt{a+b \sec(e+fx)}}{c+d \sec(e+fx)} dx$	1942

3.265	$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$	1948
3.266	$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx$	1953
3.267	$\int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx$	1958
3.268	$\int \frac{\sec(e+fx)}{\sqrt{4-5\sec(e+fx)}\sqrt{2+3\sec(e+fx)}} dx$	1963
3.269	$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx$	1968
3.270	$\int \frac{(g\sec(e+fx))^{3/2}\sqrt{c+d\sec(e+fx)}}{a+b\sec(e+fx)} dx$	1975
3.271	$\int \frac{(g\sec(e+fx))^{3/2}}{(a+b\sec(e+fx))\sqrt{c+d\sec(e+fx)}} dx$	1983
3.272	$\int \frac{\sqrt{g\sec(e+fx)}\sqrt{c+d\sec(e+fx)}}{a+b\cos(e+fx)} dx$	1989
3.273	$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx$	1997
3.274	$\int \frac{(g\sec(e+fx))^{3/2}\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx$	2002
3.275	$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$	2013
3.276	$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$	2019
3.277	$\int \frac{(g\sec(e+fx))^{3/2}}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$	2025
3.278	$\int \frac{(g\sec(e+fx))^{5/2}}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$	2035
3.279	$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$	2047
3.280	$\int \frac{(g\sec(e+fx))^{3/2}\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$	2053
3.281	$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx$	2061
3.282	$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx$	2066
3.283	$\int \frac{(g\sec(e+fx))^{3/2}}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx$	2072
3.284	$\int \frac{(g\sec(e+fx))^{5/2}}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx$	2078
3.285	$\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx$	2086
3.286	$\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^8} dx$	2092

3.1 $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$

3.1.1	Optimal result	116
3.1.2	Mathematica [A] (verified)	116
3.1.3	Rubi [A] (verified)	117
3.1.4	Maple [A] (verified)	118
3.1.5	Fricas [A] (verification not implemented)	119
3.1.6	Sympy [F]	119
3.1.7	Maxima [B] (verification not implemented)	120
3.1.8	Giac [A] (verification not implemented)	120
3.1.9	Mupad [B] (verification not implemented)	121

3.1.1 Optimal result

Integrand size = 30, antiderivative size = 105

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$$

$$= \frac{7ac^4 \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{ac^4 \sec(e + fx) \tan(e + fx)}{8f}$$

$$- \frac{3ac^4 \sec^3(e + fx) \tan(e + fx)}{4f} + \frac{4ac^4 \tan^3(e + fx)}{3f} + \frac{ac^4 \tan^5(e + fx)}{5f}$$

output $7/8*a*c^4*\operatorname{arctanh}(\sin(f*x+e))/f-1/8*a*c^4*\sec(f*x+e)*\tan(f*x+e)/f-3/4*a*c^4*\sec(f*x+e)^3*\tan(f*x+e)/f+4/3*a*c^4*\tan(f*x+e)^3/f+1/5*a*c^4*\tan(f*x+e)^5/f$

3.1.2 Mathematica [A] (verified)

Time = 5.13 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.47

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$$

$$= \frac{ac^{7/2} \left(-210 \arcsin \left(\frac{\sqrt{c - c \sec(e + fx)}}{\sqrt{2\sqrt{c}}} \right) \sqrt{c - c \sec(e + fx)} + \sqrt{c} \sqrt{1 + \sec(e + fx)} (136 - 121 \sec(e + fx)) - 120f(-1 + \sec(e + fx)) \sqrt{1 + \sec(e + fx)} \right)}{120f(-1 + \sec(e + fx)) \sqrt{1 + \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^4,x]`

output `(a*c^(7/2)*(-210*ArcSin[Sqrt[c - c*Sec[e + f*x]]/(Sqrt[2]*Sqrt[c]])*Sqrt[c - c*Sec[e + f*x]] + Sqrt[c]*Sqrt[1 + Sec[e + f*x]]*(136 - 121*Sec[e + f*x] - 127*Sec[e + f*x]^2 + 202*Sec[e + f*x]^3 - 114*Sec[e + f*x]^4 + 24*Sec[e + f*x]^5))*Tan[e + f*x])/(120*f*(-1 + Sec[e + f*x])*Sqrt[1 + Sec[e + f*x]])`

3.1.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^4 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^4 dx$$

$$\downarrow 4446$$

$$-ac \int \left(-c^3 \tan^2(e + fx) \sec^4(e + fx) + 3c^3 \tan^2(e + fx) \sec^3(e + fx) - 3c^3 \tan^2(e + fx) \sec^2(e + fx) + c^3 \tan^2(e + fx)\right) dx$$

$$\downarrow 2009$$

$$-ac \left(-\frac{7c^3 \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{c^3 \tan^5(e + fx)}{5f} - \frac{4c^3 \tan^3(e + fx)}{3f} + \frac{3c^3 \tan(e + fx) \sec^3(e + fx)}{4f} + \frac{c^3 \tan(e + fx)}{f} \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^4,x]`

output `-(a*c*((-7*c^3*ArcTanh[Sin[e + f*x]])/(8*f) + (c^3*Sec[e + f*x]*Tan[e + f*x])/(8*f) + (3*c^3*Sec[e + f*x]^3*Tan[e + f*x])/(4*f) - (4*c^3*Tan[e + f*x]^3)/(3*f) - (c^3*Tan[e + f*x]^5)/(5*f)))`

3.1. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$

3.1.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.1.4 Maple [A] (verified)

Time = 6.24 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.51

method	result
norman	$\frac{7a^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 49a^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 224a^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 79a^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7 - 7a^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^5} - \frac{7a^4 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)}{8}$
risch	$\frac{ia^4(15e^{9i(fx+e)} - 360e^{8i(fx+e)} + 390e^{7i(fx+e)} - 960e^{6i(fx+e)} - 400e^{4i(fx+e)} - 390e^{3i(fx+e)} - 320e^{2i(fx+e)} - 15e^{i(fx+e)})}{60f(1+e^{2i(fx+e)})^5}$
parts	$\frac{2a^4 \ln(\sec(fx+e) + \tan(fx+e))}{f} - \frac{a^4 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4\sec(fx+e)^2}{15}\right) \tan(fx+e)}{f} - \frac{3a^4 \tan(fx+e)}{f} + \frac{a^4 \sec(fx+e)}{f}$
parallelrisc	$- \frac{13a^4 \left(\left(\frac{35 \cos(fx+e)}{26} + \frac{35 \cos(3fx+3e)}{52} + \frac{7 \cos(5fx+5e)}{52}\right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \left(-\frac{35 \cos(fx+e)}{26} - \frac{35 \cos(3fx+3e)}{52} - \frac{7 \cos(5fx+5e)}{52}\right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2f(\cos(5fx+5e) + 5 \cos(3fx+3e))}$
derivativedivides	$-a^4 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4\sec(fx+e)^2}{15}\right) \tan(fx+e) - 3a^4 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8}\right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8}\right)$
default	$-a^4 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4\sec(fx+e)^2}{15}\right) \tan(fx+e) - 3a^4 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8}\right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8}\right)$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^4,x,method=_RETURNVERBOSE)
```

3.1. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$

output $(7/4*a*c^4/f*\tan(1/2*f*x+1/2*e)-49/6*a*c^4/f*\tan(1/2*f*x+1/2*e)^3+224/15*a*c^4/f*\tan(1/2*f*x+1/2*e)^5-79/6*a*c^4/f*\tan(1/2*f*x+1/2*e)^7-7/4*a*c^4/f*\tan(1/2*f*x+1/2*e)^9)/(\tan(1/2*f*x+1/2*e)^2-1)^5-7/8*a*c^4/f*\ln(\tan(1/2*f*x+1/2*e)-1)+7/8*a*c^4/f*\ln(\tan(1/2*f*x+1/2*e)+1)$

3.1.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.25

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$$

$$= \frac{105 ac^4 \cos(fx + e)^5 \log(\sin(fx + e) + 1) - 105 ac^4 \cos(fx + e)^5 \log(-\sin(fx + e) + 1) - 2(136 ac^4 \cos(fx + e)^4 + 15 a^2 c^4 \cos(fx + e)^3 - 112 a^2 c^4 \cos(fx + e)^2 + 90 a^2 c^4 \cos(fx + e) - 24 a^2 c^4 \sin(fx + e))}{240 f \cos(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^4,x, algorithm="fricas")`

output $1/240*(105*a*c^4*\cos(f*x + e)^5*\log(\sin(f*x + e) + 1) - 105*a*c^4*\cos(f*x + e)^5*\log(-\sin(f*x + e) + 1) - 2*(136*a*c^4*\cos(f*x + e)^4 + 15*a*c^4*\cos(f*x + e)^3 - 112*a*c^4*\cos(f*x + e)^2 + 90*a*c^4*\cos(f*x + e) - 24*a*c^4*\sin(f*x + e))/(f*\cos(f*x + e)^5)$

3.1.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$$

$$= ac^4 \left(\int \sec(e + fx) dx + \int (-3 \sec^2(e + fx)) dx + \int 2 \sec^3(e + fx) dx \right. \\ \left. + \int 2 \sec^4(e + fx) dx + \int (-3 \sec^5(e + fx)) dx + \int \sec^6(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))**4,x)`

output `a*c**4*(Integral(sec(e + f*x), x) + Integral(-3*sec(e + f*x)**2, x) + Integral(2*sec(e + f*x)**3, x) + Integral(2*sec(e + f*x)**4, x) + Integral(-3*sec(e + f*x)**5, x) + Integral(sec(e + f*x)**6, x))`

3.1. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$

3.1.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 215 vs. 2(95) = 190.

Time = 0.24 (sec) , antiderivative size = 215, normalized size of antiderivative = 2.05

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$$

$$= \frac{16 (3 \tan (fx + e)^5 + 10 \tan (fx + e)^3 + 15 \tan (fx + e)) ac^4 + 160 (\tan (fx + e)^3 + 3 \tan (fx + e)) ac^4}{120 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output `1/240*(16*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a*c^4 + 160*(tan(f*x + e)^3 + 3*tan(f*x + e))*a*c^4 + 45*a*c^4*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 120*a*c^4*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 240*a*c^4*log(sec(f*x + e) + tan(f*x + e)) - 720*a*c^4*tan(f*x + e))/f`

3.1.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.38

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$$

$$= \frac{105 ac^4 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + 1 \right| \right) - 105 ac^4 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(105 ac^4 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^9 + 790 ac^4 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^7 + 896 ac^4 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^5 + 490 ac^4 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 - 105 ac^4 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)}{120 f}}{120 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output `1/120*(105*a*c^4*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 105*a*c^4*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(105*a*c^4*tan(1/2*f*x + 1/2*e)^9 + 790*a*c^4*tan(1/2*f*x + 1/2*e)^7 - 896*a*c^4*tan(1/2*f*x + 1/2*e)^5 + 490*a*c^4*tan(1/2*f*x + 1/2*e)^3 - 105*a*c^4*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^5)/f`

3.1. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx$

3.1.9 Mupad [B] (verification not implemented)

Time = 18.91 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.68

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^4 dx = \frac{7 a c^4 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{4 f} - \frac{\frac{7 a c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{4} + \frac{79 a c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{6} - \frac{224 a c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{15} + \frac{49 a c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{6} - \frac{7 a c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1 \right)}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x))^4)/cos(e + f*x),x)`output `(7*a*c^4*atanh(tan(e/2 + (f*x)/2)))/(4*f) - ((49*a*c^4*tan(e/2 + (f*x)/2)^3)/6 - (7*a*c^4*tan(e/2 + (f*x)/2))/4 - (224*a*c^4*tan(e/2 + (f*x)/2)^5)/15 + (79*a*c^4*tan(e/2 + (f*x)/2)^7)/6 + (7*a*c^4*tan(e/2 + (f*x)/2)^9)/4)/(f*(5*tan(e/2 + (f*x)/2)^2 - 10*tan(e/2 + (f*x)/2)^4 + 10*tan(e/2 + (f*x)/2)^6 - 5*tan(e/2 + (f*x)/2)^8 + tan(e/2 + (f*x)/2)^10 - 1))`

3.2 $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$

3.2.1	Optimal result	122
3.2.2	Mathematica [A] (verified)	122
3.2.3	Rubi [A] (verified)	123
3.2.4	Maple [A] (verified)	124
3.2.5	Fricas [A] (verification not implemented)	125
3.2.6	Sympy [F]	125
3.2.7	Maxima [A] (verification not implemented)	126
3.2.8	Giac [A] (verification not implemented)	126
3.2.9	Mupad [B] (verification not implemented)	127

3.2.1 Optimal result

Integrand size = 30, antiderivative size = 86

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$$

$$= \frac{5ac^3 \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{3ac^3 \sec(e + fx) \tan(e + fx)}{8f}$$

$$- \frac{ac^3 \sec^3(e + fx) \tan(e + fx)}{4f} + \frac{2ac^3 \tan^3(e + fx)}{3f}$$

```
output 5/8*a*c^3*arctanh(sin(f*x+e))/f-3/8*a*c^3*sec(f*x+e)*tan(f*x+e)/f-1/4*a*c^
3*sec(f*x+e)^3*tan(f*x+e)/f+2/3*a*c^3*tan(f*x+e)^3/f
```

3.2.2 Mathematica [A] (verified)

Time = 1.29 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.67

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx =$$

$$\frac{ac^{5/2} \left(30 \arcsin \left(\frac{\sqrt{c - c \sec(e + fx)}}{\sqrt{2}\sqrt{c}} \right) \sqrt{c - c \sec(e + fx)} + \sqrt{c} \sqrt{1 + \sec(e + fx)} (-16 + 7 \sec(e + fx) + 25 \sec(e + fx)) \right)}{24f(-1 + \sec(e + fx))\sqrt{1 + \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^3,x]`

output `-1/24*(a*c^(5/2)*(30*ArcSin[Sqrt[c - c*Sec[e + f*x]]/(Sqrt[2]*Sqrt[c]])*Sqrt[c - c*Sec[e + f*x]] + Sqrt[c]*Sqrt[1 + Sec[e + f*x]]*(-16 + 7*Sec[e + f*x] + 25*Sec[e + f*x]^2 - 22*Sec[e + f*x]^3 + 6*Sec[e + f*x]^4))*Tan[e + f*x])/(f*(-1 + Sec[e + f*x])*Sqrt[1 + Sec[e + f*x]])`

3.2.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^3 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 dx$$

$$\downarrow \text{4446}$$

$$-ac \int (c^2 \tan^2(e + fx) \sec^3(e + fx) - 2c^2 \tan^2(e + fx) \sec^2(e + fx) + c^2 \tan^2(e + fx) \sec(e + fx)) dx$$

$$\downarrow \text{2009}$$

$$-ac \left(-\frac{5c^2 \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{2c^2 \tan^3(e + fx)}{3f} + \frac{c^2 \tan(e + fx) \sec^3(e + fx)}{4f} + \frac{3c^2 \tan(e + fx) \sec(e + fx)}{8f} \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^3,x]`

output `-(a*c*((-5*c^2*ArcTanh[Sin[e + f*x]])/(8*f) + (3*c^2*Sec[e + f*x]*Tan[e + f*x])/(8*f) + (c^2*Sec[e + f*x]^3*Tan[e + f*x])/(4*f) - (2*c^2*Tan[e + f*x]^3)/(3*f)))`

3.2. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$

3.2.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.2.4 Maple [A] (verified)

Time = 3.68 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.29

method	result
derivativedivides	$\frac{-a c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right) - 2a c^3 \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) - \frac{2a c^3 \ln(\sec(fx+e) + \tan(fx+e))}{f}}{f}$
default	$\frac{-a c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right) - 2a c^3 \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) - \frac{2a c^3 \ln(\sec(fx+e) + \tan(fx+e))}{f}}{f}$
parts	$\frac{a c^3 \ln(\sec(fx+e) + \tan(fx+e))}{f} - \frac{2a c^3 \tan(fx+e)}{f} - \frac{2a c^3 \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e)}{f} - \frac{a c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f}$
norman	$\frac{-\frac{5a c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f} + \frac{55a c^3 \tan^3\left(\frac{fx}{2} + \frac{e}{2}\right)}{12f} - \frac{73a c^3 \tan^5\left(\frac{fx}{2} + \frac{e}{2}\right)}{12f} - \frac{5a c^3 \tan^7\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^4} - \frac{5a c^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{8f} + \frac{5a c^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{8f}$
risch	$\frac{ia c^3 (9e^{7i(fx+e)} - 48e^{6i(fx+e)} + 33e^{5i(fx+e)} - 48e^{4i(fx+e)} - 33e^{3i(fx+e)} - 16e^{2i(fx+e)} - 9e^{i(fx+e)} - 16)}{12f(1+e^{2i(fx+e)})^4} + \frac{5a c^3 \ln(e^{i(fx+e)} + \tan(fx+e))}{8f}$
parallelrisc	$-\frac{3 \left(\left(\frac{10 \cos(2fx+2e)}{3} + \frac{5 \cos(4fx+4e)}{6} + \frac{5}{2} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \left(-\frac{10 \cos(2fx+2e)}{3} - \frac{5 \cos(4fx+4e)}{6} - \frac{5}{2} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \right)}{4f(3 + \cos(4fx+4e) + 4 \cos(2fx+2e))}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

3.2. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$

output $1/f*(-a*c^3*(-(-1/4*\sec(f*x+e)^3-3/8*\sec(f*x+e))*\tan(f*x+e)+3/8*\ln(\sec(f*x+e)+\tan(f*x+e)))-2*a*c^3*(-2/3-1/3*\sec(f*x+e)^2)*\tan(f*x+e)-2*a*c^3*\tan(f*x+e)+a*c^3*\ln(\sec(f*x+e)+\tan(f*x+e)))$

3.2.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.36

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$$

$$= \frac{15 ac^3 \cos(fx + e)^4 \log(\sin(fx + e) + 1) - 15 ac^3 \cos(fx + e)^4 \log(-\sin(fx + e) + 1) - 2(16 ac^3 \cos(fx + e)^3 + 9 a^2 c^3 \cos(fx + e)^2 - 16 a^2 c^3 \cos(fx + e) + 6 a^2 c^3) \sin(fx + e)}{48 f \cos(fx + e)^4}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^3,x, algorithm="fricas")`

output $1/48*(15*a*c^3*\cos(f*x + e)^4*\log(\sin(f*x + e) + 1) - 15*a*c^3*\cos(f*x + e)^4*\log(-\sin(f*x + e) + 1) - 2*(16*a*c^3*\cos(f*x + e)^3 + 9*a*c^3*\cos(f*x + e)^2 - 16*a*c^3*\cos(f*x + e) + 6*a*c^3)*\sin(f*x + e))/(f*\cos(f*x + e)^4)$

3.2.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$$

$$= -ac^3 \left(\int (-\sec(e + fx)) dx + \int 2 \sec^2(e + fx) dx + \int (-2 \sec^4(e + fx)) dx + \int \sec^5(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))**3,x)`

output `-a*c**3*(Integral(-sec(e + f*x), x) + Integral(2*sec(e + f*x)**2, x) + Integral(-2*sec(e + f*x)**4, x) + Integral(sec(e + f*x)**5, x))`

3.2.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.55

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$$

$$= \frac{32 (\tan (fx + e)^3 + 3 \tan (fx + e)) ac^3 + 3 ac^3 \left(\frac{2 (3 \sin (fx + e)^3 - 5 \sin (fx + e))}{\sin (fx + e)^4 - 2 \sin (fx + e)^2 + 1} - 3 \log (\sin (fx + e) + 1) + 3 \log (\sin (fx + e) - 1) \right)}{48 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/48*(32*(tan(f*x + e)^3 + 3*tan(f*x + e))*a*c^3 + 3*a*c^3*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) + 48*a*c^3*log(sec(f*x + e) + tan(f*x + e)) - 96*a*c^3*tan(f*x + e))/f`

3.2.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.49

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$$

$$= \frac{15 ac^3 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + 1 \right| \right) - 15 ac^3 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(15 ac^3 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^7 + 73 ac^3 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^5 - 55 ac^3 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + 15 ac^3 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)}{24 f}}{\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - 1}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `1/24*(15*a*c^3*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*a*c^3*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(15*a*c^3*tan(1/2*f*x + 1/2*e)^7 + 73*a*c^3*tan(1/2*f*x + 1/2*e)^5 - 55*a*c^3*tan(1/2*f*x + 1/2*e)^3 + 15*a*c^3*tan(1/2*f*x + 1/2*e)))/(tan(1/2*f*x + 1/2*e)^2 - 1)/f`

3.2.9 Mupad [B] (verification not implemented)

Time = 17.22 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.70

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^3 dx$$

$$= \frac{5ac^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{4f}$$

$$- \frac{\frac{5ac^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{4} + \frac{73ac^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{12} - \frac{55ac^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{12} + \frac{5ac^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1 \right)}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x))^3)/cos(e + f*x),x)`output `(5*a*c^3*atanh(tan(e/2 + (f*x)/2)))/(4*f) - ((5*a*c^3*tan(e/2 + (f*x)/2))/4 - (55*a*c^3*tan(e/2 + (f*x)/2)^3)/12 + (73*a*c^3*tan(e/2 + (f*x)/2)^5)/12 + (5*a*c^3*tan(e/2 + (f*x)/2)^7)/4)/(f*(6*tan(e/2 + (f*x)/2)^4 - 4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/2)^8 + 1))`

3.3 $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$

3.3.1	Optimal result	128
3.3.2	Mathematica [B] (verified)	128
3.3.3	Rubi [A] (verified)	129
3.3.4	Maple [A] (verified)	130
3.3.5	Fricas [A] (verification not implemented)	131
3.3.6	Sympy [F]	131
3.3.7	Maxima [A] (verification not implemented)	132
3.3.8	Giac [B] (verification not implemented)	132
3.3.9	Mupad [B] (verification not implemented)	133

3.3.1 Optimal result

Integrand size = 30, antiderivative size = 61

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$$

$$= \frac{ac^2 \operatorname{arctanh}(\sin(e + fx))}{2f} - \frac{ac^2 \sec(e + fx) \tan(e + fx)}{2f} + \frac{ac^2 \tan^3(e + fx)}{3f}$$

output `1/2*a*c^2*arctanh(sin(f*x+e))/f-1/2*a*c^2*sec(f*x+e)*tan(f*x+e)/f+1/3*a*c^2*tan(f*x+e)^3/f`

3.3.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 132 vs. 2(61) = 122.

Time = 0.57 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.16

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$$

$$= \frac{ac^{3/2} \left(-6 \arcsin \left(\frac{\sqrt{c - c \sec(e + fx)}}{\sqrt{2}\sqrt{c}} \right) \sqrt{c - c \sec(e + fx)} + \sqrt{c} \sqrt{1 + \sec(e + fx)} (2 + \sec(e + fx)) - 5 \sec^2(e + fx) \right)}{6f(-1 + \sec(e + fx))\sqrt{1 + \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^2,x]`

output $(a*c^{(3/2)}*(-6*ArcSin[Sqrt[c - c*Sec[e + f*x]]/(Sqrt[2]*Sqrt[c]])*Sqrt[c - c*Sec[e + f*x]] + Sqrt[c]*Sqrt[1 + Sec[e + f*x]]*(2 + Sec[e + f*x] - 5*Sec[e + f*x]^2 + 2*Sec[e + f*x]^3)*Tan[e + f*x])/(6*f*(-1 + Sec[e + f*x])*Sqrt[1 + Sec[e + f*x]])$

3.3.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^2 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 dx$$

$$\downarrow 4446$$

$$-ac \int (c \sec(e + fx) \tan^2(e + fx) - c \sec^2(e + fx) \tan^2(e + fx)) dx$$

$$\downarrow 2009$$

$$-ac \left(-\frac{\text{arctanh}(\sin(e + fx))}{2f} - \frac{c \tan^3(e + fx)}{3f} + \frac{c \tan(e + fx) \sec(e + fx)}{2f} \right)$$

input $\text{Int}[\text{Sec}[e + f*x]*(a + a*\text{Sec}[e + f*x])*(c - c*\text{Sec}[e + f*x])^2,x]$

output $-(a*c*(-1/2*(c*ArcTanh[Sin[e + f*x]]))/f + (c*Sec[e + f*x]*Tan[e + f*x])/(2*f) - (c*Tan[e + f*x]^3)/(3*f))$

3.3.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.3.4 Maple [A] (verified)

Time = 4.15 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.61

method	result
derivativedivides	$\frac{-a c^2 \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e) - a c^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2}\right)}{f} - a c^2 \tan(fx+e) + a c^2 \ln(\sec(fx+e) + \tan(fx+e))$
default	$\frac{-a c^2 \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e) - a c^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2}\right)}{f} - a c^2 \tan(fx+e) + a c^2 \ln(\sec(fx+e) + \tan(fx+e))$
risch	$\frac{ia c^2 (3e^{5i(fx+e)} - 6e^{4i(fx+e)} - 3e^{i(fx+e)} - 2)}{3f(1+e^{2i(fx+e)})^3} + \frac{a c^2 \ln(e^{i(fx+e)} + i)}{2f} - \frac{a c^2 \ln(e^{i(fx+e)} - i)}{2f}$
parts	$\frac{a c^2 \ln(\sec(fx+e) + \tan(fx+e))}{f} - \frac{a c^2 \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e)}{f} - \frac{a c^2 \tan(fx+e)}{f} - \frac{a c^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2}\right)}{f}$
norman	$\frac{a c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f} - \frac{8a c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3f} - \frac{a c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f} - \frac{a c^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2f} + \frac{a c^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2f}$
parallelrisch	$-\frac{a c^2 \left(\frac{3(\cos(3fx+3e) + 3 \cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2} + \frac{3(-\cos(3fx+3e) - 3 \cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} + \sin(3fx+3e)\right)}{3f(\cos(3fx+3e) + 3 \cos(fx+e))}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

3.3. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$

output `1/f*(-a*c^2*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)-a*c^2*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))-a*c^2*tan(f*x+e)+a*c^2*ln(sec(f*x+e)+tan(f*x+e)))`

3.3.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.69

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$$

$$= \frac{3ac^2 \cos(fx + e)^3 \log(\sin(fx + e) + 1) - 3ac^2 \cos(fx + e)^3 \log(-\sin(fx + e) + 1) - 2(2ac^2 \cos(fx + e)^2 + 3a^2c^2 \cos(fx + e) - 2a^2c^2) \sin(fx + e)}{12f \cos(fx + e)^3}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^2,x, algorithm="fricas")`

output `1/12*(3*a*c^2*cos(f*x + e)^3*log(sin(f*x + e) + 1) - 3*a*c^2*cos(f*x + e)^3*log(-sin(f*x + e) + 1) - 2*(2*a*c^2*cos(f*x + e)^2 + 3*a*c^2*cos(f*x + e) - 2*a*c^2)*sin(f*x + e))/(f*cos(f*x + e)^3)`

3.3.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$$

$$= ac^2 \left(\int \sec(e + fx) dx + \int (-\sec^2(e + fx)) dx + \int (-\sec^3(e + fx)) dx + \int \sec^4(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))**2,x)`

output `a*c**2*(Integral(sec(e + f*x), x) + Integral(-sec(e + f*x)**2, x) + Integral(-sec(e + f*x)**3, x) + Integral(sec(e + f*x)**4, x))`

3.3.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.77

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$$

$$= \frac{4(\tan(fx + e)^3 + 3 \tan(fx + e))ac^2 + 3ac^2 \left(\frac{2 \sin(fx + e)}{\sin(fx + e)^2 - 1} - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1) \right) + 12ac^2 \log(\sec(fx + e) + \tan(fx + e)) - 12ac^2 \tan(fx + e)}{12f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output `1/12*(4*(tan(f*x + e)^3 + 3*tan(f*x + e))*a*c^2 + 3*a*c^2*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 12*a*c^2*log(sec(f*x + e) + tan(f*x + e)) - 12*a*c^2*tan(f*x + e))/f`

3.3.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. 2(55) = 110.

Time = 0.31 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.82

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$$

$$= \frac{3ac^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|) - 3ac^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|) - \frac{2(3ac^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 8ac^2 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)}}{6f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `1/6*(3*a*c^2*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 3*a*c^2*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(3*a*c^2*tan(1/2*f*x + 1/2*e)^5 + 8*a*c^2*tan(1/2*f*x + 1/2*e)^3 - 3*a*c^2*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1))/f`

3.3.9 Mupad [B] (verification not implemented)

Time = 15.64 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.87

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^2 dx$$

$$= \frac{a c^2 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{f} - \frac{a c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \frac{8 a c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3} - a c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1 \right)}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x))^2)/cos(e + f*x),x)`

output `(a*c^2*atanh(tan(e/2 + (f*x)/2)))/f - ((8*a*c^2*tan(e/2 + (f*x)/2)^3)/3 - a*c^2*tan(e/2 + (f*x)/2) + a*c^2*tan(e/2 + (f*x)/2)^5)/(f*(3*tan(e/2 + (f*x)/2)^2 - 3*tan(e/2 + (f*x)/2)^4 + tan(e/2 + (f*x)/2)^6 - 1))`

3.4 $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$

3.4.1	Optimal result	134
3.4.2	Mathematica [A] (verified)	134
3.4.3	Rubi [A] (verified)	135
3.4.4	Maple [A] (verified)	136
3.4.5	Fricas [A] (verification not implemented)	137
3.4.6	Sympy [F]	137
3.4.7	Maxima [A] (verification not implemented)	138
3.4.8	Giac [A] (verification not implemented)	138
3.4.9	Mupad [B] (verification not implemented)	139

3.4.1 Optimal result

Integrand size = 28, antiderivative size = 38

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx \\ &= \frac{a \operatorname{arctanh}(\sin(e + fx))}{2f} - \frac{ac \sec(e + fx) \tan(e + fx)}{2f} \end{aligned}$$

output `1/2*a*c*arctanh(sin(f*x+e))/f-1/2*a*c*sec(f*x+e)*tan(f*x+e)/f`

3.4.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx \\ &= -ac \left(-\frac{\operatorname{arctanh}(\sin(e + fx))}{2f} + \frac{\sec(e + fx) \tan(e + fx)}{2f} \right) \end{aligned}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x]),x]`

output `-(a*c*(-1/2*ArcTanh[Sin[e + f*x]]/f + (Sec[e + f*x]*Tan[e + f*x])/(2*f)))`

3.4.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3042, 4446, 3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e+fx)(a \sec(e+fx)+a)(c-c \sec(e+fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e+fx+\frac{\pi}{2}\right)\left(a \csc\left(e+fx+\frac{\pi}{2}\right)+a\right)\left(c-c \csc\left(e+fx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4446} \\
 & -ac \int \sec(e+fx) \tan^2(e+fx) dx \\
 & \quad \downarrow \text{3042} \\
 & -ac \int \sec(e+fx) \tan(e+fx)^2 dx \\
 & \quad \downarrow \text{3091} \\
 & -ac \left(\frac{\tan(e+fx) \sec(e+fx)}{2f} - \frac{1}{2} \int \sec(e+fx) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & -ac \left(\frac{\tan(e+fx) \sec(e+fx)}{2f} - \frac{1}{2} \int \csc\left(e+fx+\frac{\pi}{2}\right) dx \right) \\
 & \quad \downarrow \text{4257} \\
 & -ac \left(\frac{\tan(e+fx) \sec(e+fx)}{2f} - \frac{\operatorname{arctanh}(\sin(e+fx))}{2f} \right)
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x]),x]`

output `-(a*c*(-1/2*ArcTanh[Sin[e + f*x]]/f + (Sec[e + f*x]*Tan[e + f*x])/(2*f)))`

3.4.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3091 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] & & NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.4.4 Maple [A] (verified)

Time = 1.51 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.53

method	result	size
derivativedivides	$\frac{-ac\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) + ac\ln(\sec(fx+e)+\tan(fx+e))}{f}$	58
default	$\frac{-ac\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) + ac\ln(\sec(fx+e)+\tan(fx+e))}{f}$	58
parts	$\frac{ac\ln(\sec(fx+e)+\tan(fx+e))}{f} - \frac{ac\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)}{f}$	60
parallelrisc	$\frac{\left((-1-\cos(2fx+2e))\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)+(1+\cos(2fx+2e))\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)-2\sin(fx+e)\right)ac}{2f(1+\cos(2fx+2e))}$	80
risc	$\frac{iac(e^{3i(fx+e)}-e^{i(fx+e)})}{f(1+e^{2i(fx+e)})^2} + \frac{ac\ln(e^{i(fx+e)}+i)}{2f} - \frac{ac\ln(e^{i(fx+e)}-i)}{2f}$	84
norman	$\frac{-\frac{ac\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{f} - \frac{ac\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{f}}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)^2} - \frac{ac\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{2f} + \frac{ac\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{2f}$	91

3.4. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/f*(-a*c*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+a*c*ln(sec(f*x+e)+tan(f*x+e)))`

3.4.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.76

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \frac{ac \cos(fx + e)^2 \log(\sin(fx + e) + 1) - ac \cos(fx + e)^2 \log(-\sin(fx + e) + 1) - 2ac \sin(fx + e)}{4f \cos(fx + e)^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `1/4*(a*c*cos(f*x + e)^2*log(sin(f*x + e) + 1) - a*c*cos(f*x + e)^2*log(-sin(f*x + e) + 1) - 2*a*c*sin(f*x + e))/(f*cos(f*x + e)^2)`

3.4.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= -ac \left(\int (-\sec(e + fx)) dx + \int \sec^3(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x)`

output `-a*c*(Integral(-sec(e + f*x), x) + Integral(sec(e + f*x)**3, x))`

3.4.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.79

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \frac{ac \left(\frac{2 \sin(fx+e)}{\sin(fx+e)^2-1} - \log(\sin(fx+e) + 1) + \log(\sin(fx+e) - 1) \right) + 4ac \log(\sec(fx+e) + \tan(fx+e))}{4f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `1/4*(a*c*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 4*a*c*log(sec(f*x + e) + tan(f*x + e)))/f`

3.4.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.45

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \frac{ac \log(|\sin(fx+e) + 1|) - ac \log(|\sin(fx+e) - 1|) + \frac{2ac \sin(fx+e)}{\sin(fx+e)^2-1}}{4f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="giac")`

output `1/4*(a*c*log(abs(sin(f*x + e) + 1)) - a*c*log(abs(sin(f*x + e) - 1)) + 2*a*c*sin(f*x + e)/(sin(f*x + e)^2 - 1))/f`

3.4.9 Mupad [B] (verification not implemented)

Time = 14.54 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.03

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \frac{a c \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{f} - \frac{a c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + a c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1\right)}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x)))/cos(e + f*x),x)`

output `(a*c*atanh(tan(e/2 + (f*x)/2)))/f - (a*c*tan(e/2 + (f*x)/2)^3 + a*c*tan(e/2 + (f*x)/2))/(f*(tan(e/2 + (f*x)/2)^4 - 2*tan(e/2 + (f*x)/2)^2 + 1))`

$$3.5 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c-c \sec(e+fx)} dx$$

3.5.1	Optimal result	140
3.5.2	Mathematica [A] (verified)	140
3.5.3	Rubi [A] (verified)	141
3.5.4	Maple [A] (verified)	142
3.5.5	Fricas [A] (verification not implemented)	143
3.5.6	Sympy [F]	143
3.5.7	Maxima [B] (verification not implemented)	144
3.5.8	Giac [A] (verification not implemented)	144
3.5.9	Mupad [B] (verification not implemented)	145

3.5.1 Optimal result

Integrand size = 30, antiderivative size = 42

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c-c \sec(e+fx)} dx = -\frac{a \operatorname{arctanh}(\sin(e+fx))}{cf} - \frac{2a \tan(e+fx)}{f(c-c \sec(e+fx))}$$

output `-a*arctanh(sin(f*x+e))/c/f-2*a*tan(f*x+e)/f/(c-c*sec(f*x+e))`

3.5.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.83

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c-c \sec(e+fx)} dx$$

$$= -\frac{a \left(-\frac{2 \cot(\frac{1}{2}(e+fx))}{f} - \frac{\log(\cos(\frac{1}{2}(e+fx)) - \sin(\frac{1}{2}(e+fx)))}{f} + \frac{\log(\cos(\frac{1}{2}(e+fx)) + \sin(\frac{1}{2}(e+fx)))}{f} \right)}{c}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x]),x]`

output `-((a*((-2*Cot[(e + f*x)/2])/f - Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]]/f + Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]]/f))/c`

$$3.5. \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c-c \sec(e+fx)} dx$$

3.5.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4445, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{c-c\sec(e+fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{c-c\csc(e+fx+\frac{\pi}{2})} dx \\ & \quad \downarrow \text{4445} \\ & -\frac{a \int \sec(e+fx) dx}{c} - \frac{2a \tan(e+fx)}{f(c-c\sec(e+fx))} \\ & \quad \downarrow \text{3042} \\ & -\frac{a \int \csc(e+fx+\frac{\pi}{2}) dx}{c} - \frac{2a \tan(e+fx)}{f(c-c\sec(e+fx))} \\ & \quad \downarrow \text{4257} \\ & -\frac{a \operatorname{arctanh}(\sin(e+fx))}{cf} - \frac{2a \tan(e+fx)}{f(c-c\sec(e+fx))} \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x]),x]`

output `-((a*ArcTanh[Sin[e + f*x]])/(c*f)) - (2*a*Tan[e + f*x])/(f*(c - c*Sec[e + f*x]))`

3.5.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_)*(csc[(e_.) + (f_.)*(x_)*(b_.) + (a_)^(m_)*(csc[(e_.) + (f_.)*(x_)*(d_.) + (c_)^(n_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.5.4 Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

method	result	size
derivativedivides	$\frac{2a \left(\frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2} \right)}{fc}$	50
default	$\frac{2a \left(\frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2} \right)}{fc}$	50
parallelrisc	$\frac{a \left(\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 2 \right)}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right) cf}$	67
risc	$\frac{4ia}{fc(e^{i(fx+e)} - 1)} + \frac{a \ln(e^{i(fx+e)} - i)}{cf} - \frac{a \ln(e^{i(fx+e)} + i)}{cf}$	68
norman	$\frac{-\frac{2a}{cf} + \frac{2a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)} + \frac{a \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{cf} - \frac{a \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{cf}$	100

input `int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

$$3.5. \int \frac{\sec(e+fx)(a+a\sec(e+fx))}{c-c\sec(e+fx)} dx$$

output $2/f*a/c*(1/\tan(1/2*f*x+1/2*e)-1/2*\ln(\tan(1/2*f*x+1/2*e)+1)+1/2*\ln(\tan(1/2*f*x+1/2*e)-1))$

3.5.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.57

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{c-c\sec(e+fx)} dx = \frac{a \log(\sin(fx+e)+1) \sin(fx+e) - a \log(-\sin(fx+e)+1) \sin(fx+e) - 4a \cos(fx+e) - 4a}{2cf \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x, algorithm="fricas")`

output $-1/2*(a*\log(\sin(f*x+e)+1)*\sin(f*x+e) - a*\log(-\sin(f*x+e)+1)*\sin(f*x+e) - 4*a*\cos(f*x+e) - 4*a)/(c*f*\sin(f*x+e))$

3.5.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{c-c\sec(e+fx)} dx = -\frac{a \left(\int \frac{\sec(e+fx)}{\sec(e+fx)-1} dx + \int \frac{\sec^2(e+fx)}{\sec(e+fx)-1} dx \right)}{c}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x)`

output $-a*(\text{Integral}(\sec(e+f*x)/(\sec(e+f*x)-1),x) + \text{Integral}(\sec(e+f*x)**2/(\sec(e+f*x)-1),x))/c$

3.5.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. $2(43) = 86$.

Time = 0.22 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.40

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{c - c \sec(e + fx)} dx$$

$$= - \frac{a \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{c} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{c} - \frac{\cos(fx+e)+1}{c \sin(fx+e)} \right) - \frac{a(\cos(fx+e)+1)}{c \sin(fx+e)}}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-(a*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/c - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/c - (cos(f*x + e) + 1)/(c*sin(f*x + e))) - a*(cos(f*x + e) + 1)/(c*sin(f*x + e)))/f`

3.5.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.43

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{c - c \sec(e + fx)} dx$$

$$= - \frac{\frac{a \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{c} - \frac{a \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{c} - \frac{2a}{c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x, algorithm="giac")`

output `-(a*log(abs(tan(1/2*f*x + 1/2*e) + 1))/c - a*log(abs(tan(1/2*f*x + 1/2*e) - 1))/c - 2*a/(c*tan(1/2*f*x + 1/2*e)))/f`

3.5.9 Mupad [B] (verification not implemented)

Time = 13.76 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{c - c \sec(e + fx)} dx = -\frac{2a \left(\operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) - \cot\left(\frac{e}{2} + \frac{fx}{2}\right) \right)}{cf}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))),x)`

output `-(2*a*(atanh(tan(e/2 + (f*x)/2)) - cot(e/2 + (f*x)/2)))/(c*f)`

3.6 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^2} dx$

3.6.1	Optimal result	146
3.6.2	Mathematica [A] (verified)	146
3.6.3	Rubi [A] (verified)	147
3.6.4	Maple [A] (verified)	148
3.6.5	Fricas [A] (verification not implemented)	148
3.6.6	Sympy [F]	149
3.6.7	Maxima [B] (verification not implemented)	149
3.6.8	Giac [A] (verification not implemented)	150
3.6.9	Mupad [B] (verification not implemented)	150

3.6.1 Optimal result

Integrand size = 30, antiderivative size = 36

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^2} dx = -\frac{(a+a \sec(e+fx)) \tan(e+fx)}{3f(c-c \sec(e+fx))^2}$$

output `-1/3*(a+a*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^2`

3.6.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^2} dx = -\frac{a \cot^3\left(\frac{1}{2}(e+fx)\right)}{3c^2 f}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^2,x]`

output `-1/3*(a*Cot[(e + f*x)/2]^3)/(c^2*f)`

3.6.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c-c\sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^2} dx$$

↓ 4438

$$-\frac{\tan(e+fx)(a\sec(e+fx)+a)}{3f(c-c\sec(e+fx))^2}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^2,x]`

output `-1/3*((a + a*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^2)`

3.6.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.6.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$-\frac{a}{3f c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}$	21
default	$-\frac{a}{3f c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}$	21
parallelrisch	$-\frac{a}{3f c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}$	21
risch	$\frac{2ia(3e^{2i(fx+e)}+1)}{3f c^2 (e^{i(fx+e)}-1)^3}$	37
norman	$\frac{\frac{a}{3cf} - \frac{a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{3cf}}{c\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}$	61

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
)
```

```
output -1/3/f*a/c^2/tan(1/2*f*x+1/2*e)^3
```

3.6.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.42

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^2} dx = \frac{a \cos(fx+e)^2 + 2a \cos(fx+e) + a}{3(c^2 f \cos(fx+e) - c^2 f) \sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x, algorithm="fracas")
```

```
output 1/3*(a*cos(f*x + e)^2 + 2*a*cos(f*x + e) + a)/((c^2*f*cos(f*x + e) - c^2*f)*sin(f*x + e))
```

3.6.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^2} dx$$

$$= \frac{a \left(\int \frac{\sec(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx + \int \frac{\sec^2(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx \right)}{c^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**2,x)`

output `a*(Integral(sec(e + f*x)/(sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**2/(sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x))/c**2`

3.6.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. $2(35) = 70$.

Time = 0.22 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.69

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^2} dx$$

$$= -\frac{a \left(\frac{3 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} + 1 \right) (\cos(fx+e)+1)^3}{c^2 \sin^3(fx+e)} - \frac{a \left(\frac{3 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - 1 \right) (\cos(fx+e)+1)^3}{c^2 \sin^3(fx+e)}$$

$$6f$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output `-1/6*(a*(3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)*(cos(f*x + e) + 1)^3/(c^2*sin(f*x + e)^3) - a*(3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 1)*(cos(f*x + e) + 1)^3/(c^2*sin(f*x + e)^3))/f`

3.6.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c - c \sec(e + fx))^2} dx = -\frac{a}{3c^2 f \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `-1/3*a/(c^2*f*tan(1/2*f*x + 1/2*e)^3)`

3.6.9 Mupad [B] (verification not implemented)

Time = 13.52 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c - c \sec(e + fx))^2} dx = -\frac{a \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3c^2 f}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^2),x)`

output `-(a*cot(e/2 + (f*x)/2)^3)/(3*c^2*f)`

$$3.7 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^3} dx$$

3.7.1	Optimal result	151
3.7.2	Mathematica [A] (verified)	151
3.7.3	Rubi [A] (verified)	152
3.7.4	Maple [A] (verified)	153
3.7.5	Fricas [A] (verification not implemented)	154
3.7.6	Sympy [F]	154
3.7.7	Maxima [A] (verification not implemented)	155
3.7.8	Giac [A] (verification not implemented)	155
3.7.9	Mupad [B] (verification not implemented)	156

3.7.1 Optimal result

Integrand size = 30, antiderivative size = 76

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^3} dx = -\frac{(a+a \sec(e+fx)) \tan(e+fx)}{5f(c-c \sec(e+fx))^3} - \frac{(a+a \sec(e+fx)) \tan(e+fx)}{15cf(c-c \sec(e+fx))^2}$$

output `-1/5*(a+a*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^3-1/15*(a+a*sec(f*x+e))*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^2`

3.7.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.57

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^3} dx = -\frac{a(-4+\sec(e+fx))(1+\sec(e+fx)) \tan(e+fx)}{15c^3f(-1+\sec(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^3,x]`

output `-1/15*(a*(-4 + Sec[e + f*x]))*(1 + Sec[e + f*x])*Tan[e + f*x]/(c^3*f*(-1 + Sec[e + f*x])^3)`

3.7.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c-c\sec(e+fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{4439} \\
 & \frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^2} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^2} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3} \\
 & \quad \downarrow \text{4438} \\
 & -\frac{\tan(e+fx)(a\sec(e+fx)+a)}{15cf(c-c\sec(e+fx))^2} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^3,x]`

output `-1/5*((a + a*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^3) - ((a + a*Sec[e + f*x])*Tan[e + f*x])/(15*c*f*(c - c*Sec[e + f*x])^2)`

3.7.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.7.4 Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.47

method	result	size
parallelsch	$\frac{a \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 \left(3 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 5\right)}{30c^3 f}$	36
derivativedivides	$a \left(-\frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} \right) \frac{1}{2f c^3}$	37
default	$a \left(-\frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} \right) \frac{1}{2f c^3}$	37
risch	$\frac{2ia(15e^{4i(fx+e)} - 15e^{3i(fx+e)} + 25e^{2i(fx+e)} - 5e^{i(fx+e)} + 4)}{15f c^3 (e^{i(fx+e)} - 1)^5}$	70
norman	$\frac{-\frac{a}{10cf} + \frac{4a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{15cf} - \frac{a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{6cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right) c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	81

3.7. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^3} dx$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/30*a*cot(1/2*f*x+1/2*e)^3*(3*cot(1/2*f*x+1/2*e)^2-5)/c^3/f
```

3.7.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.03

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^3} dx$$

$$= \frac{4a\cos(fx+e)^3 + 7a\cos(fx+e)^2 + 2a\cos(fx+e) - a}{15(c^3f\cos(fx+e)^2 - 2c^3f\cos(fx+e) + c^3f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x, algorithm="fricas")
```

```
output 1/15*(4*a*cos(f*x + e)^3 + 7*a*cos(f*x + e)^2 + 2*a*cos(f*x + e) - a)/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x + e))
```

3.7.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^3} dx$$

$$= -\frac{a\left(\int \frac{\sec(e+fx)}{\sec^3(e+fx)-3\sec^2(e+fx)+3\sec(e+fx)-1} dx + \int \frac{\sec^2(e+fx)}{\sec^3(e+fx)-3\sec^2(e+fx)+3\sec(e+fx)-1} dx\right)}{c^3}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**3,x)
```

```
output -a*(Integral(sec(e + f*x)/(sec(e + f*x)**3 - 3*sec(e + f*x)**2 + 3*sec(e + f*x) - 1), x) + Integral(sec(e + f*x)**2/(sec(e + f*x)**3 - 3*sec(e + f*x)**2 + 3*sec(e + f*x) - 1), x))/c**3
```

3.7.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.54

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^3} dx$$

$$= -\frac{a\left(\frac{10\sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{15\sin(fx+e)^4}{(\cos(fx+e)+1)^4} - 3\right)(\cos(fx+e)+1)^5}{c^3\sin(fx+e)^5} + \frac{3a\left(\frac{5\sin(fx+e)^4}{(\cos(fx+e)+1)^4} - 1\right)(\cos(fx+e)+1)^5}{c^3\sin(fx+e)^5}$$

$60f$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `-1/60*(a*(10*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 15*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 3)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5) + 3*a*(5*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 1)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5))/f`

3.7.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.49

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^3} dx = -\frac{5a\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 3a}{30c^3f\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `-1/30*(5*a*tan(1/2*f*x + 1/2*e)^2 - 3*a)/(c^3*f*tan(1/2*f*x + 1/2*e)^5)`

3.7.9 Mupad [B] (verification not implemented)

Time = 14.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.46

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c - c \sec(e + fx))^3} dx = \frac{a \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(3 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 5\right)}{30 c^3 f}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^3),x)`

output `(a*cot(e/2 + (f*x)/2)^3*(3*cot(e/2 + (f*x)/2)^2 - 5))/(30*c^3*f)`

3.8 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^4} dx$

3.8.1 Optimal result 157
 3.8.2 Mathematica [A] (verified) 157
 3.8.3 Rubi [A] (verified) 158
 3.8.4 Maple [A] (verified) 160
 3.8.5 Fricas [A] (verification not implemented) 160
 3.8.6 Sympy [F] 161
 3.8.7 Maxima [A] (verification not implemented) 161
 3.8.8 Giac [A] (verification not implemented) 162
 3.8.9 Mupad [B] (verification not implemented) 162

3.8.1 Optimal result

Integrand size = 30, antiderivative size = 116

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^4} dx = -\frac{(a+a \sec(e+fx)) \tan(e+fx)}{7f(c-c \sec(e+fx))^4} - \frac{2(a+a \sec(e+fx)) \tan(e+fx)}{35cf(c-c \sec(e+fx))^3} - \frac{2(a+a \sec(e+fx)) \tan(e+fx)}{105f(c^2-c^2 \sec(e+fx))^2}$$

output

```
-1/7*(a+a*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^4-2/35*(a+a*sec(f*x+e))
)*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^3-2/105*(a+a*sec(f*x+e))*tan(f*x+e)/f/(c
^2-c^2*sec(f*x+e))^2
```

3.8.2 Mathematica [A] (verified)

Time = 3.95 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.47

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^4} dx = -\frac{a(1+\sec(e+fx))(23-10 \sec(e+fx)+2 \sec^2(e+fx)) \tan(e+fx)}{105c^4f(-1+\sec(e+fx))^4}$$

input

```
Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^4,x]
```

output
$$\frac{-1/105*(a*(1 + \text{Sec}[e + f*x])*(23 - 10*\text{Sec}[e + f*x] + 2*\text{Sec}[e + f*x]^2)*\text{Tan}[e + f*x])}{(c^4*f*(-1 + \text{Sec}[e + f*x])^4)}$$

3.8.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c-c\sec(e+fx))^4} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx \\ & \quad \downarrow \text{4439} \\ & \frac{2 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} \\ & \quad \downarrow \text{3042} \\ & \frac{2 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} \\ & \quad \downarrow \text{4439} \\ & \frac{2 \left(\frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^2} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3} \right)}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} \\ & \quad \downarrow \text{3042} \\ & \frac{2 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^2} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3} \right)}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} \\ & \quad \downarrow \text{4438} \end{aligned}$$

3.8.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^4} dx$$

$$\frac{2\left(-\frac{\tan(e+fx)(a\sec(e+fx)+a)}{15cf(c-c\sec(e+fx))^2} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3}\right)}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^4,x]`

output `-1/7*((a + a*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^4) + (2*(-1/5*((a + a*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^3) - ((a + a*Sec[e + f*x])*Tan[e + f*x])/(15*c*f*(c - c*Sec[e + f*x])^2)))/(7*c)`

3.8.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.8.4 Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.42

method	result	size
parallelrisch	$-\frac{a \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 \left(15 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^4 - 42 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 + 35\right)}{420c^4 f}$	49
derivativedivides	$a \left(\frac{-\frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{2}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}}{4f c^4} \right)$	50
default	$a \left(\frac{-\frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{2}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}}{4f c^4} \right)$	50
risch	$\frac{2ia(105 e^{6i(fx+e)} - 210 e^{5i(fx+e)} + 455 e^{4i(fx+e)} - 350 e^{3i(fx+e)} + 273 e^{2i(fx+e)} - 56 e^{i(fx+e)} + 23)}{105f c^4 (e^{i(fx+e)} - 1)^7}$	92
norman	$\frac{\frac{a}{28cf} - \frac{19a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{140cf} + \frac{11a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{60cf} - \frac{a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{12cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right) c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$	101

input `int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x,method=_RETURNVERBOSE)`

output `-1/420*a*cot(1/2*f*x+1/2*e)^3*(15*cot(1/2*f*x+1/2*e)^4-42*cot(1/2*f*x+1/2*e)^2+35)/c^4/f`

3.8.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.90

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^4} dx$$

$$= \frac{23a \cos(fx+e)^4 + 36a \cos(fx+e)^3 + 5a \cos(fx+e)^2 - 6a \cos(fx+e) + 2a}{105(c^4 f \cos(fx+e)^3 - 3c^4 f \cos(fx+e)^2 + 3c^4 f \cos(fx+e) - c^4 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x, algorithm="fricas")`

output `1/105*(23*a*cos(f*x + e)^4 + 36*a*cos(f*x + e)^3 + 5*a*cos(f*x + e)^2 - 6*a*cos(f*x + e) + 2*a)/((c^4*f*cos(f*x + e)^3 - 3*c^4*f*cos(f*x + e)^2 + 3*c^4*f*cos(f*x + e) - c^4*f)*sin(f*x + e))`

3.8.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^4} dx$$

3.8.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^4} dx$$

$$= \frac{a \left(\int \frac{\sec(e+fx)}{\sec^4(e+fx)-4\sec^3(e+fx)+6\sec^2(e+fx)-4\sec(e+fx)+1} dx + \int \frac{\sec^2(e+fx)}{\sec^4(e+fx)-4\sec^3(e+fx)+6\sec^2(e+fx)-4\sec(e+fx)+1} dx \right)}{c^4}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**4,x)`

output `a*(Integral(sec(e + f*x)/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**2/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x))/c**4`

3.8.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.53

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^4} dx$$

$$= \frac{a \left(\frac{21 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{35 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{105 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - 15 \right) (\cos(fx+e)+1)^7}{c^4 \sin(fx+e)^7} + \frac{3a \left(\frac{21 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{35 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{35 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - 5 \right) (\cos(fx+e)+1)^7}{c^4 \sin(fx+e)^7}$$

840 f

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output `1/840*(a*(21*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 35*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 105*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 15)*(cos(f*x + e) + 1)^7/(c^4*sin(f*x + e)^7) + 3*a*(21*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 35*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 35*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 5)*(cos(f*x + e) + 1)^7/(c^4*sin(f*x + e)^7))/f`

3.8.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.44

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^4} dx$$

$$= -\frac{35a \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^4 - 42a \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 + 15a}{420c^4 f \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^7}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output `-1/420*(35*a*tan(1/2*f*x + 1/2*e)^4 - 42*a*tan(1/2*f*x + 1/2*e)^2 + 15*a)/(c^4*f*tan(1/2*f*x + 1/2*e)^7)`

3.8.9 Mupad [B] (verification not implemented)

Time = 14.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.53

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^4} dx = \frac{a \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{10c^4 f} - \frac{a \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{12c^4 f} - \frac{a \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{28c^4 f}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^4),x)`

output `(a*cot(e/2 + (f*x)/2)^5)/(10*c^4*f) - (a*cot(e/2 + (f*x)/2)^3)/(12*c^4*f) - (a*cot(e/2 + (f*x)/2)^7)/(28*c^4*f)`

3.9 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^5} dx$

3.9.1	Optimal result	163
3.9.2	Mathematica [A] (verified)	164
3.9.3	Rubi [A] (verified)	164
3.9.4	Maple [A] (verified)	166
3.9.5	Fricas [A] (verification not implemented)	167
3.9.6	Sympy [F]	167
3.9.7	Maxima [A] (verification not implemented)	168
3.9.8	Giac [A] (verification not implemented)	168
3.9.9	Mupad [B] (verification not implemented)	169

3.9.1 Optimal result

Integrand size = 30, antiderivative size = 158

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^5} dx = -\frac{(a+a \sec(e+fx)) \tan(e+fx)}{9f(c-c \sec(e+fx))^5} - \frac{(a+a \sec(e+fx)) \tan(e+fx)}{21cf(c-c \sec(e+fx))^4} - \frac{2(a+a \sec(e+fx)) \tan(e+fx)}{105c^2f(c-c \sec(e+fx))^3} - \frac{2(a+a \sec(e+fx)) \tan(e+fx)}{315cf(c^2-c^2 \sec(e+fx))^2}$$

```
output -1/9*(a+a*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^5-1/21*(a+a*sec(f*x+e))
)*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^4-2/105*(a+a*sec(f*x+e))*tan(f*x+e)/c^2/
f/(c-c*sec(f*x+e))^3-2/315*(a+a*sec(f*x+e))*tan(f*x+e)/c/f/(c^2-c^2*sec(f*
x+e))^2
```

3.9.2 Mathematica [A] (verified)

Time = 5.14 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.41

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx = \frac{a(1+\sec(e+fx))(-58+33\sec(e+fx)-12\sec^2(e+fx)+2\sec^3(e+fx))\tan(e+fx)}{315c^5f(-1+\sec(e+fx))^5}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^5,x]`

output `-1/315*(a*(1 + Sec[e + f*x])*(-58 + 33*Sec[e + f*x] - 12*Sec[e + f*x]^2 + 2*Sec[e + f*x]^3)*Tan[e + f*x])/(c^5*f*(-1 + Sec[e + f*x])^5)`

3.9.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4439, 3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c-c\sec(e+fx))^5} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^5} dx \\ & \quad \downarrow \text{4439} \\ & \frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^4} dx}{3c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{9f(c-c\sec(e+fx))^5} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx}{3c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{9f(c-c\sec(e+fx))^5} \\ & \quad \downarrow \text{4439} \end{aligned}$$

3.9. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx$

$$\begin{aligned}
& \frac{2 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{9f(c-c\sec(e+fx))^5} \\
& \quad \quad \quad \downarrow \quad 3042 \\
& \frac{2 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{9f(c-c\sec(e+fx))^5} \\
& \quad \quad \quad \downarrow \quad 4439 \\
& \frac{2 \left(\frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^2} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3} \right)}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} - \\
& \quad \quad \quad \frac{3c}{9f(c-c\sec(e+fx))^5} \frac{\tan(e+fx)(a\sec(e+fx)+a)}{\tan(e+fx)(a\sec(e+fx)+a)} \\
& \quad \quad \quad \downarrow \quad 3042 \\
& \frac{2 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^2} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3} \right)}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} - \\
& \quad \quad \quad \frac{3c}{9f(c-c\sec(e+fx))^5} \frac{\tan(e+fx)(a\sec(e+fx)+a)}{\tan(e+fx)(a\sec(e+fx)+a)} \\
& \quad \quad \quad \downarrow \quad 4438 \\
& \frac{2 \left(\frac{-\tan(e+fx)(a\sec(e+fx)+a)}{15cf(c-c\sec(e+fx))^2} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{5f(c-c\sec(e+fx))^3} \right)}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)}{7f(c-c\sec(e+fx))^4} - \\
& \quad \quad \quad \frac{3c}{9f(c-c\sec(e+fx))^5} \frac{\tan(e+fx)(a\sec(e+fx)+a)}{\tan(e+fx)(a\sec(e+fx)+a)}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^5,x]`

output `-1/9*((a + a*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^5) + (-1/7*((a + a*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^4) + (2*(-1/5*((a + a*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^3) - ((a + a*Sec[e + f*x])*Tan[e + f*x])/(15*c*f*(c - c*Sec[e + f*x])^2)))/(7*c))/(3*c)`

3.9.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)]`

3.9.4 Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.39

method	result
parallelrisch	$\frac{a \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 \left(35 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^6 - 135 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^4 + 189 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 105\right)}{2520c^5 f}$
derivativedivides	$\frac{a \left(\frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} - \frac{3}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{3}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}\right)}{8f c^5}$
default	$\frac{a \left(\frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} - \frac{3}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{3}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}\right)}{8f c^5}$
risch	$\frac{2ia(315 e^{8i(fx+e)} - 945 e^{7i(fx+e)} + 2625 e^{6i(fx+e)} - 3465 e^{5i(fx+e)} + 3843 e^{4i(fx+e)} - 2247 e^{3i(fx+e)} + 1143 e^{2i(fx+e)} - 20)}{315f c^5 (e^{i(fx+e)} - 1)^9}$
norman	$\frac{-\frac{a}{72cf} + \frac{17a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{252cf} - \frac{9a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{70cf} + \frac{7a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{60cf} - \frac{a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{24cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right) c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$

3.9. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^5} dx$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^5,x,method=_RETURNVERBOSE)
```

```
output 1/2520*a*cot(1/2*f*x+1/2*e)^3*(35*cot(1/2*f*x+1/2*e)^6-135*cot(1/2*f*x+1/2
*e)^4+189*cot(1/2*f*x+1/2*e)^2-105)/c^5/f
```

3.9.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.81

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx = \frac{58a\cos(fx+e)^5 + 83a\cos(fx+e)^4 + 4a\cos(fx+e)^3 - 11a\cos(fx+e)^2 + 8a\cos(fx+e) - 2a}{315(c^5f\cos(fx+e)^4 - 4c^5f\cos(fx+e)^3 + 6c^5f\cos(fx+e)^2 - 4c^5f\cos(fx+e) + c^5f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^5,x, algorithm="fricas")
```

```
output 1/315*(58*a*cos(f*x + e)^5 + 83*a*cos(f*x + e)^4 + 4*a*cos(f*x + e)^3 - 11
*a*cos(f*x + e)^2 + 8*a*cos(f*x + e) - 2*a)/((c^5*f*cos(f*x + e)^4 - 4*c^5
*f*cos(f*x + e)^3 + 6*c^5*f*cos(f*x + e)^2 - 4*c^5*f*cos(f*x + e) + c^5*f)
*sin(f*x + e))
```

3.9.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx = \frac{a\left(\int \frac{\sec(e+fx)}{\sec^5(e+fx)-5\sec^4(e+fx)+10\sec^3(e+fx)-10\sec^2(e+fx)+5\sec(e+fx)-1} dx + \int \frac{\sec^2(e+fx)}{\sec^5(e+fx)-5\sec^4(e+fx)+10\sec^3(e+fx)-10\sec^2(e+fx)+5\sec(e+fx)-1} dx\right)}{c^5}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**5,x)
```

```
output -a*(Integral(sec(e + f*x)/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec(e
+ f*x)**3 - 10*sec(e + f*x)**2 + 5*sec(e + f*x) - 1), x) + Integral(sec(e
+ f*x)**2/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec(e + f*x)**3 - 10*s
ec(e + f*x)**2 + 5*sec(e + f*x) - 1), x))/c**5
```

3.9. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx$

3.9.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.25

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx = \frac{a\left(\frac{180\sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{378\sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{420\sin(fx+e)^6}{(\cos(fx+e)+1)^6} - \frac{315\sin(fx+e)^8}{(\cos(fx+e)+1)^8} - 35\right)(\cos(fx+e)+1)^9}{c^5\sin(fx+e)^9} + \frac{5a\left(\frac{18\sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{42\sin(fx+e)^6}{(\cos(fx+e)+1)^6} + \frac{63\sin(fx+e)^8}{(\cos(fx+e)+1)^8} - 7\right)(\cos(fx+e)+1)^9}{c^5\sin(fx+e)^9}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^5,x, algorithm="maxima")`

output `-1/5040*(a*(180*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 378*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 420*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 315*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 - 35)*(cos(f*x + e) + 1)^9/(c^5*sin(f*x + e)^9) + 5*a*(18*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 42*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 63*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 - 7)*(cos(f*x + e) + 1)^9/(c^5*sin(f*x + e)^9))/f`

3.9.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.41

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx = -\frac{105a\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^6 - 189a\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^4 + 135a\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 35a}{2520c^5f\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^9}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^5,x, algorithm="giac")`

output `-1/2520*(105*a*tan(1/2*f*x + 1/2*e)^6 - 189*a*tan(1/2*f*x + 1/2*e)^4 + 135*a*tan(1/2*f*x + 1/2*e)^2 - 35*a)/(c^5*f*tan(1/2*f*x + 1/2*e)^9)`

3.9.9 Mupad [B] (verification not implemented)

Time = 13.80 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^5} dx$$

$$= \frac{a \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(35 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 135 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^4 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 189 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^2 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2520 c^5 f \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{2520 c^5 f \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^9}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^5),x)`

output `(a*cos(e/2 + (f*x)/2)^3*(35*cos(e/2 + (f*x)/2)^6 - 105*sin(e/2 + (f*x)/2)^6 + 189*cos(e/2 + (f*x)/2)^2*sin(e/2 + (f*x)/2)^4 - 135*cos(e/2 + (f*x)/2)^4*sin(e/2 + (f*x)/2)^2)/(2520*c^5*f*sin(e/2 + (f*x)/2)^9)`

3.10 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$

3.10.1	Optimal result	170
3.10.2	Mathematica [A] (verified)	171
3.10.3	Rubi [A] (verified)	171
3.10.4	Maple [C] (verified)	172
3.10.5	Fricas [A] (verification not implemented)	173
3.10.6	Sympy [F]	174
3.10.7	Maxima [B] (verification not implemented)	174
3.10.8	Giac [A] (verification not implemented)	175
3.10.9	Mupad [B] (verification not implemented)	176

3.10.1 Optimal result

Integrand size = 32, antiderivative size = 171

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$$

$$= \frac{9a^2c^5 \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{3a^2c^5 \sec(e + fx) \tan(e + fx)}{16f}$$

$$- \frac{3a^2c^5 \sec^3(e + fx) \tan(e + fx)}{8f} + \frac{a^2c^5 \sec(e + fx) \tan^3(e + fx)}{4f}$$

$$+ \frac{a^2c^5 \sec^3(e + fx) \tan^3(e + fx)}{2f} - \frac{4a^2c^5 \tan^5(e + fx)}{5f} - \frac{a^2c^5 \tan^7(e + fx)}{7f}$$

output `9/16*a^2*c^5*arctanh(sin(f*x+e))/f-3/16*a^2*c^5*sec(f*x+e)*tan(f*x+e)/f-3/8*a^2*c^5*sec(f*x+e)^3*tan(f*x+e)/f+1/4*a^2*c^5*sec(f*x+e)*tan(f*x+e)^3/f+1/2*a^2*c^5*sec(f*x+e)^3*tan(f*x+e)^3/f-4/5*a^2*c^5*tan(f*x+e)^5/f-1/7*a^2*c^5*tan(f*x+e)^7/f`

3.10.2 Mathematica [A] (verified)

Time = 1.32 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.60

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$$

$$= \frac{a^2 c^5 (10080 \operatorname{arctanh}(\sin(e + fx)) - \sec^7(e + fx)(2520 \sin(e + fx) - 455 \sin(2(e + fx)) - 616 \sin(3(e + fx) - 2380 \sin(4(e + fx)) + 392 \sin(5(e + fx)) + 245 \sin(6(e + fx)) + 184 \sin(7(e + fx))))}{17920 f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^5,x]`

output `(a^2*c^5*(10080*ArcTanh[Sin[e + f*x]] - Sec[e + f*x]^7*(2520*Sin[e + f*x] - 455*Sin[2*(e + f*x)] - 616*Sin[3*(e + f*x)] + 2380*Sin[4*(e + f*x)] - 392*Sin[5*(e + f*x)] + 245*Sin[6*(e + f*x)] + 184*Sin[7*(e + f*x)])))/(17920*f)`

3.10.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^5 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^5 dx$$

$$\downarrow \text{4446}$$

$$a^2 c^2 \int \left(-c^3 \sec^4(e + fx) \tan^4(e + fx) + 3c^3 \sec^3(e + fx) \tan^4(e + fx) - 3c^3 \sec^2(e + fx) \tan^4(e + fx) + c^3 \sec(e + fx) \tan^4(e + fx)\right) dx$$

$$\downarrow \text{2009}$$

$$a^2 c^2 \left(\frac{9c^3 \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{c^3 \tan^7(e + fx)}{7f} - \frac{4c^3 \tan^5(e + fx)}{5f} + \frac{c^3 \tan^3(e + fx) \sec^3(e + fx)}{2f} - \frac{3c^3 \tan(e + fx) \sec^3(e + fx)}{2f} \right)$$

3.10. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^5,x]`

output `a^2*c^2*((9*c^3*ArcTanh[Sin[e + f*x]])/(16*f) - (3*c^3*Sec[e + f*x]*Tan[e + f*x])/(16*f) - (3*c^3*Sec[e + f*x]^3*Tan[e + f*x])/(8*f) + (c^3*Sec[e + f*x]*Tan[e + f*x]^3)/(4*f) + (c^3*Sec[e + f*x]^3*Tan[e + f*x]^3)/(2*f) - (4*c^3*Tan[e + f*x]^5)/(5*f) - (c^3*Tan[e + f*x]^7)/(7*f))`

3.10.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.10.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 8.88 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.22

$$3.10. \quad \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$$

method	result
risch	$\frac{ic^5 a^2 (245 e^{13i(fx+e)} - 1680 e^{12i(fx+e)} + 2380 e^{11i(fx+e)} - 4480 e^{10i(fx+e)} - 455 e^{9i(fx+e)} - 3920 e^{8i(fx+e)} - 8960 e^{6i(fx+e)} + 455 e^{5i(fx+e)} - 3248 e^{4i(fx+e)} - 2380 e^{3i(fx+e)} - 896 e^{2i(fx+e)} - 245 e^{i(fx+e)} - 368)}{280 f (1 + e^{2i(fx+e)})^7}$
norman	$\frac{9c^5 a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{15c^5 a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{2f} + \frac{849c^5 a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{40f} - \frac{1152c^5 a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{35f} + \frac{1199c^5 a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{40f} + \frac{15c^5 a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{40f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^7}$
parallelrisc	$9a^2 c^5 \left(\frac{(\cos(7fx+7e) + 7\cos(5fx+5e) + 21\cos(3fx+3e) + 35\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{16} + \frac{(-\cos(7fx+7e) - 7\cos(5fx+5e) - 21\cos(3fx+3e) - 35\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{16} \right)$
derivativedivides	$c^5 a^2 \left(-\frac{16}{35} - \frac{\sec(fx+e)^6}{7} - \frac{6\sec(fx+e)^4}{35} - \frac{8\sec(fx+e)^2}{35} \right) \tan(fx+e) + 3c^5 a^2 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5\sec(fx+e)^3}{24} - \frac{5\sec(fx+e)}{16} \right) \right)$
default	$c^5 a^2 \left(-\frac{16}{35} - \frac{\sec(fx+e)^6}{7} - \frac{6\sec(fx+e)^4}{35} - \frac{8\sec(fx+e)^2}{35} \right) \tan(fx+e) + 3c^5 a^2 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5\sec(fx+e)^3}{24} - \frac{5\sec(fx+e)}{16} \right) \right)$
parts	$\frac{c^5 a^2 \ln(\sec(fx+e) + \tan(fx+e))}{f} + \frac{c^5 a^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right)}{f} - \frac{3c^5 a^2 \tan(fx+e)}{f} - \frac{5c^5 a^2}{f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^5,x,method=_RETURNVERBOSE)
```

```
output 1/280*I*c^5*a^2*(245*exp(13*I*(f*x+e))-1680*exp(12*I*(f*x+e))+2380*exp(11*I*(f*x+e))-4480*exp(10*I*(f*x+e))-455*exp(9*I*(f*x+e))-3920*exp(8*I*(f*x+e))-8960*exp(6*I*(f*x+e))+455*exp(5*I*(f*x+e))-3248*exp(4*I*(f*x+e))-2380*exp(3*I*(f*x+e))-896*exp(2*I*(f*x+e))-245*exp(I*(f*x+e))-368)/f/(1+exp(2*I*(f*x+e)))^7+9/16*c^5*a^2/f*ln(exp(I*(f*x+e))+I)-9/16*c^5*a^2/f*ln(exp(I*(f*x+e))-I)
```

3.10.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.04

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$$

$$= \frac{315 a^2 c^5 \cos(fx + e)^7 \log(\sin(fx + e) + 1) - 315 a^2 c^5 \cos(fx + e)^7 \log(-\sin(fx + e) + 1) - 2(368 a^2 c^5 \cos(fx + e)^7 \log(\tan(fx + e) + 1) - 368 a^2 c^5 \cos(fx + e)^7 \log(\tan(fx + e) - 1))}{16}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^5,x, algorithm="fricas")
```

3.10. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$

```
output 1/1120*(315*a^2*c^5*cos(f*x + e)^7*log(sin(f*x + e) + 1) - 315*a^2*c^5*cos
(f*x + e)^7*log(-sin(f*x + e) + 1) - 2*(368*a^2*c^5*cos(f*x + e)^6 + 245*a
^2*c^5*cos(f*x + e)^5 - 656*a^2*c^5*cos(f*x + e)^4 + 350*a^2*c^5*cos(f*x +
e)^3 + 208*a^2*c^5*cos(f*x + e)^2 - 280*a^2*c^5*cos(f*x + e) + 80*a^2*c^5
)*sin(f*x + e))/(f*cos(f*x + e)^7)
```

3.10.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx \\ &= -a^2c^5 \left(\int (-\sec(e + fx)) dx + \int 3\sec^2(e + fx) dx + \int (-\sec^3(e + fx)) dx \right. \\ & \quad \left. + \int (-5\sec^4(e + fx)) dx + \int 5\sec^5(e + fx) dx + \int \sec^6(e + fx) dx \right. \\ & \quad \left. + \int (-3\sec^7(e + fx)) dx + \int \sec^8(e + fx) dx \right) \end{aligned}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**5,x)
```

```
output -a**2*c**5*(Integral(-sec(e + f*x), x) + Integral(3*sec(e + f*x)**2, x) +
Integral(-sec(e + f*x)**3, x) + Integral(-5*sec(e + f*x)**4, x) + Integral
(5*sec(e + f*x)**5, x) + Integral(sec(e + f*x)**6, x) + Integral(-3*sec(e
+ f*x)**7, x) + Integral(sec(e + f*x)**8, x))
```

3.10.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 368 vs. $2(157) = 314$.

Time = 0.21 (sec) , antiderivative size = 368, normalized size of antiderivative = 2.15

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx = \\ & 96(5 \tan(fx + e)^7 + 21 \tan(fx + e)^5 + 35 \tan(fx + e)^3 + 35 \tan(fx + e))a^2c^5 + 224(3 \tan(fx + e) \end{aligned}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^5,x, algorithm="m
axima")
```

3.10. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$

output
$$\begin{aligned} & -1/3360*(96*(5*\tan(f*x + e))^7 + 21*\tan(f*x + e)^5 + 35*\tan(f*x + e)^3 + 35 \\ & * \tan(f*x + e))*a^2*c^5 + 224*(3*\tan(f*x + e)^5 + 10*\tan(f*x + e)^3 + 15*\tan \\ & \tan(f*x + e))*a^2*c^5 - 5600*(\tan(f*x + e)^3 + 3*\tan(f*x + e))*a^2*c^5 + 105 \\ & *a^2*c^5*(2*(15*\sin(f*x + e)^5 - 40*\sin(f*x + e)^3 + 33*\sin(f*x + e))/(\sin \\ & (f*x + e)^6 - 3*\sin(f*x + e)^4 + 3*\sin(f*x + e)^2 - 1) - 15*\log(\sin(f*x + \\ & e) + 1) + 15*\log(\sin(f*x + e) - 1)) - 1050*a^2*c^5*(2*(3*\sin(f*x + e)^3 - \\ & 5*\sin(f*x + e))/(\sin(f*x + e)^4 - 2*\sin(f*x + e)^2 + 1) - 3*\log(\sin(f*x + \\ & e) + 1) + 3*\log(\sin(f*x + e) - 1)) + 840*a^2*c^5*(2*\sin(f*x + e))/(\sin(f*x \\ & + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) - 3360*a^2*c^ \\ & 5*\log(\sec(f*x + e) + \tan(f*x + e)) + 10080*a^2*c^5*\tan(f*x + e))/f \end{aligned}$$

3.10.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.15

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$$

$$= \frac{315 a^2 c^5 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + 1 \right| \right) - 315 a^2 c^5 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(315 a^2 c^5 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)^{13} - 2100 a^2 c^5 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^{11} + 8393 a^2 c^5 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^9 - 9216 a^2 c^5 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^7 - 5943 a^2 c^5 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^5 + 2100 a^2 c^5 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 - 315 a^2 c^5 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)}{\left(\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)^2 - 1}}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^5,x, algorithm="giac")`

output
$$\begin{aligned} & 1/560*(315*a^2*c^5*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1)) - 315*a^2*c^5*\log(\text{abs} \\ & \text{abs}(\tan(1/2*f*x + 1/2*e) - 1)) - 2*(315*a^2*c^5*\tan(1/2*f*x + 1/2*e)^{13} - 21 \\ & 00*a^2*c^5*\tan(1/2*f*x + 1/2*e)^{11} - 8393*a^2*c^5*\tan(1/2*f*x + 1/2*e)^9 + \\ & 9216*a^2*c^5*\tan(1/2*f*x + 1/2*e)^7 - 5943*a^2*c^5*\tan(1/2*f*x + 1/2*e)^5 \\ & + 2100*a^2*c^5*\tan(1/2*f*x + 1/2*e)^3 - 315*a^2*c^5*\tan(1/2*f*x + 1/2*e)) \\ & /(\tan(1/2*f*x + 1/2*e)^2 - 1)^7)/f \end{aligned}$$

3.10.9 Mupad [B] (verification not implemented)

Time = 17.75 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.47

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx$$

$$= \frac{-\frac{9a^2c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{13}}{8} + \frac{15a^2c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}{2} + \frac{1199a^2c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{40} - \frac{1152a^2c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{35} + \frac{849a^2c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{40}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{14} - 7 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{12} + 21 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - 35 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 35 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 21 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1 \right)} + \frac{9a^2c^5 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{8f}$$

input `int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^5)/cos(e + f*x),x)`output `((849*a^2*c^5*tan(e/2 + (f*x)/2)^5)/40 - (15*a^2*c^5*tan(e/2 + (f*x)/2)^3)/2 - (1152*a^2*c^5*tan(e/2 + (f*x)/2)^7)/35 + (1199*a^2*c^5*tan(e/2 + (f*x)/2)^9)/40 + (15*a^2*c^5*tan(e/2 + (f*x)/2)^11)/2 - (9*a^2*c^5*tan(e/2 + (f*x)/2)^13)/8 + (9*a^2*c^5*tan(e/2 + (f*x)/2))/8)/(f*(7*tan(e/2 + (f*x)/2)^2 - 21*tan(e/2 + (f*x)/2)^4 + 35*tan(e/2 + (f*x)/2)^6 - 35*tan(e/2 + (f*x)/2)^8 + 21*tan(e/2 + (f*x)/2)^10 - 7*tan(e/2 + (f*x)/2)^12 + tan(e/2 + (f*x)/2)^14 - 1)) + (9*a^2*c^5*atanh(tan(e/2 + (f*x)/2)))/(8*f)`

3.11 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$

3.11.1	Optimal result	177
3.11.2	Mathematica [A] (verified)	178
3.11.3	Rubi [A] (verified)	178
3.11.4	Maple [A] (verified)	179
3.11.5	Fricas [A] (verification not implemented)	180
3.11.6	Sympy [F]	180
3.11.7	Maxima [B] (verification not implemented)	181
3.11.8	Giac [A] (verification not implemented)	182
3.11.9	Mupad [B] (verification not implemented)	182

3.11.1 Optimal result

Integrand size = 32, antiderivative size = 150

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx \\ &= \frac{7a^2c^4 \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{5a^2c^4 \sec(e + fx) \tan(e + fx)}{16f} \\ & \quad - \frac{a^2c^4 \sec^3(e + fx) \tan(e + fx)}{8f} + \frac{a^2c^4 \sec(e + fx) \tan^3(e + fx)}{4f} \\ & \quad + \frac{a^2c^4 \sec^3(e + fx) \tan^3(e + fx)}{6f} - \frac{2a^2c^4 \tan^5(e + fx)}{5f} \end{aligned}$$

output `7/16*a^2*c^4*arctanh(sin(f*x+e))/f-5/16*a^2*c^4*sec(f*x+e)*tan(f*x+e)/f-1/8*a^2*c^4*sec(f*x+e)^3*tan(f*x+e)/f+1/4*a^2*c^4*sec(f*x+e)*tan(f*x+e)^3/f+1/6*a^2*c^4*sec(f*x+e)^3*tan(f*x+e)^3/f-2/5*a^2*c^4*tan(f*x+e)^5/f`

3.11.2 Mathematica [A] (verified)

Time = 0.89 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.61

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$$

$$= \frac{a^2 c^4 (1680 \operatorname{arctanh}(\sin(e + fx)) + \sec^6(e + fx)(330 \sin(e + fx) - 240 \sin(2(e + fx)) - 445 \sin(3(e + fx)))}{3840 f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^4,x]`

output `(a^2*c^4*(1680*ArcTanh[Sin[e + f*x]] + Sec[e + f*x]^6*(330*Sin[e + f*x] - 240*Sin[2*(e + f*x)] - 445*Sin[3*(e + f*x)] + 192*Sin[4*(e + f*x)] - 135*Sin[5*(e + f*x)] - 48*Sin[6*(e + f*x)]))/(3840*f)`

3.11.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.93, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^4 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^4 dx$$

$$\downarrow 4446$$

$$a^2 c^2 \int (c^2 \sec^3(e + fx) \tan^4(e + fx) - 2c^2 \sec^2(e + fx) \tan^4(e + fx) + c^2 \sec(e + fx) \tan^4(e + fx)) dx$$

$$\downarrow 2009$$

$$a^2 c^2 \left(\frac{7c^2 \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{2c^2 \tan^5(e + fx)}{5f} + \frac{c^2 \tan^3(e + fx) \sec^3(e + fx)}{6f} - \frac{c^2 \tan(e + fx) \sec^3(e + fx)}{8f} \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^4,x]`

3.11. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$

output $a^2c^2((7c^2\text{ArcTanh}[\text{Sin}[e + fx]])/(16f) - (5c^2\text{Sec}[e + fx]\text{Tan}[e + fx])/(16f) - (c^2\text{Sec}[e + fx]^3\text{Tan}[e + fx])/(8f) + (c^2\text{Sec}[e + fx]\text{Tan}[e + fx]^3)/(4f) + (c^2\text{Sec}[e + fx]^3\text{Tan}[e + fx]^3)/(6f) - (2c^2\text{Tan}[e + fx]^5)/(5f))$

3.11.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.11.4 Maple [A] (verified)

Time = 6.70 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.30

method	result
norman	$\frac{-\frac{7c^4a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{8f} + \frac{119c^4a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{24f} - \frac{231c^4a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{20f} + \frac{281c^4a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{20f} + \frac{119c^4a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{24f} - \frac{7c^4a^2}{24f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^6}$
risch	$\frac{ic^4a^2(135e^{11i(fx+e)} - 480e^{10i(fx+e)} + 445e^{9i(fx+e)} - 480e^{8i(fx+e)} - 330e^{7i(fx+e)} - 960e^{6i(fx+e)} + 330e^{5i(fx+e)} - 960e^{4i(fx+e)} + 480e^{3i(fx+e)} - 135e^{2i(fx+e)} + 135e^{i(fx+e)} - 135)}{120f(1+e^{2i(fx+e)})^6}$
parallelrisc	$-\frac{2a^2c^4 \left(7\left(5 + \frac{\cos(6fx+6e)}{2}\right) + 3\cos(4fx+4e) + \frac{15\cos(2fx+2e)}{2} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 7\left(-5 - \frac{15\cos(2fx+2e)}{2} - 3\cos(4fx+4e) - \cos(6fx+6e)\right)}{16}$
derivativedivides	$c^4a^2 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5\sec(fx+e)^3}{24} - \frac{5\sec(fx+e)}{16} \right) \tan(fx+e) + \frac{5\ln(\sec(fx+e) + \tan(fx+e))}{16} \right) + 2c^4a^2 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} \right)$
default	$c^4a^2 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5\sec(fx+e)^3}{24} - \frac{5\sec(fx+e)}{16} \right) \tan(fx+e) + \frac{5\ln(\sec(fx+e) + \tan(fx+e))}{16} \right) + 2c^4a^2 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} \right)$
parts	$\frac{c^4a^2 \ln(\sec(fx+e) + \tan(fx+e))}{f} + \frac{c^4a^2 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5\sec(fx+e)^3}{24} - \frac{5\sec(fx+e)}{16} \right) \tan(fx+e) + \frac{5\ln(\sec(fx+e) + \tan(fx+e))}{16} \right)}{f}$

3.11. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^4,x,method=_RETURNVERBOSE)`

output $(-7/8*c^4*a^2/f*\tan(1/2*f*x+1/2*e)+119/24*c^4*a^2/f*\tan(1/2*f*x+1/2*e)^3-31/20*c^4*a^2/f*\tan(1/2*f*x+1/2*e)^5+281/20*c^4*a^2/f*\tan(1/2*f*x+1/2*e)^7+119/24*c^4*a^2/f*\tan(1/2*f*x+1/2*e)^9-7/8*c^4*a^2/f*\tan(1/2*f*x+1/2*e)^{11})/(\tan(1/2*f*x+1/2*e)^2-1)^6-7/16*c^4*a^2/f*\ln(\tan(1/2*f*x+1/2*e)-1)+7/16*c^4*a^2/f*\ln(\tan(1/2*f*x+1/2*e)+1)$

3.11.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.07

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$$

$$= \frac{105 a^2 c^4 \cos(fx + e)^6 \log(\sin(fx + e) + 1) - 105 a^2 c^4 \cos(fx + e)^6 \log(-\sin(fx + e) + 1) - 2(96 a^2 c^4 \cos(fx + e)^6 \log(\sin(fx + e) + 1) - 105 a^2 c^4 \cos(fx + e)^6 \log(-\sin(fx + e) + 1) - 2(96 a^2 c^4 \cos(fx + e)^5 + 135 a^2 c^4 \cos(fx + e)^4 - 192 a^2 c^4 \cos(fx + e)^3 + 10 a^2 c^4 \cos(fx + e)^2 + 96 a^2 c^4 \cos(fx + e) - 40 a^2 c^4) \sin(fx + e))}{(f \cos(fx + e))^6}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^4,x, algorithm="fricas")`

output $1/480*(105*a^2*c^4*\cos(f*x + e)^6*\log(\sin(f*x + e) + 1) - 105*a^2*c^4*\cos(f*x + e)^6*\log(-\sin(f*x + e) + 1) - 2*(96*a^2*c^4*\cos(f*x + e)^5 + 135*a^2*c^4*\cos(f*x + e)^4 - 192*a^2*c^4*\cos(f*x + e)^3 + 10*a^2*c^4*\cos(f*x + e)^2 + 96*a^2*c^4*\cos(f*x + e) - 40*a^2*c^4)*\sin(f*x + e))/(f*\cos(f*x + e)^6)$

3.11.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$$

$$= a^2 c^4 \left(\int \sec(e + fx) dx + \int (-2 \sec^2(e + fx)) dx + \int (-\sec^3(e + fx)) dx \right.$$

$$+ \int 4 \sec^4(e + fx) dx + \int (-\sec^5(e + fx)) dx + \int (-2 \sec^6(e + fx)) dx$$

$$\left. + \int \sec^7(e + fx) dx \right)$$

3.11. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**4,x)`

output `a**2*c**4*(Integral(sec(e + f*x), x) + Integral(-2*sec(e + f*x)**2, x) + Integral(-sec(e + f*x)**3, x) + Integral(4*sec(e + f*x)**4, x) + Integral(-sec(e + f*x)**5, x) + Integral(-2*sec(e + f*x)**6, x) + Integral(sec(e + f*x)**7, x))`

3.11.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 321 vs. $2(138) = 276$.

Time = 0.22 (sec) , antiderivative size = 321, normalized size of antiderivative = 2.14

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx =$$

$$64(3 \tan(fx + e)^5 + 10 \tan(fx + e)^3 + 15 \tan(fx + e))a^2c^4 - 640(\tan(fx + e)^3 + 3 \tan(fx + e))$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output `-1/480*(64*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^2*c^4 - 640*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^2*c^4 + 5*a^2*c^4*(2*(15*sin(f*x + e)^5 - 40*sin(f*x + e)^3 + 33*sin(f*x + e))/(sin(f*x + e)^6 - 3*sin(f*x + e)^4 + 3*sin(f*x + e)^2 - 1) - 15*log(sin(f*x + e) + 1) + 15*log(sin(f*x + e) - 1)) - 30*a^2*c^4*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 120*a^2*c^4*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 480*a^2*c^4*log(sec(f*x + e) + tan(f*x + e)) + 960*a^2*c^4*tan(f*x + e))/f`

3.11. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$

3.11.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.19

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$$

$$= \frac{105 a^2 c^4 \log\left(\left|\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) + 1\right|\right) - 105 a^2 c^4 \log\left(\left|\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) - 1\right|\right) - \frac{2\left(105 a^2 c^4 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11} - 595 a^2 c^4 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^9 + 1686 a^2 c^4 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^7 - 386 a^2 c^4 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 - 595 a^2 c^4 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + 105 a^2 c^4 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)\right)}{\left(\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 1\right)^6}{f} - 240 f$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^4,x, algorithm="giac")
```

```
output 1/240*(105*a^2*c^4*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 105*a^2*c^4*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(105*a^2*c^4*tan(1/2*f*x + 1/2*e)^11 - 595*a^2*c^4*tan(1/2*f*x + 1/2*e)^9 - 1686*a^2*c^4*tan(1/2*f*x + 1/2*e)^7 + 386*a^2*c^4*tan(1/2*f*x + 1/2*e)^5 - 595*a^2*c^4*tan(1/2*f*x + 1/2*e)^3 + 105*a^2*c^4*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^6)/f
```

3.11.9 Mupad [B] (verification not implemented)

Time = 17.24 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.46

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$$

$$= \frac{-\frac{7 a^2 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}{8} + \frac{119 a^2 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{24} + \frac{281 a^2 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{20} - \frac{231 a^2 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{20} + \frac{119 a^2 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{24} - \frac{7 a^2 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{24}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{12} - 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} + 15 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 20 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 15 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1 \right)} + \frac{7 a^2 c^4 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{8 f}$$

```
input int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^4)/cos(e + f*x),x)
```

```
output ((119*a^2*c^4*tan(e/2 + (f*x)/2)^3)/24 - (231*a^2*c^4*tan(e/2 + (f*x)/2)^5)/20 + (281*a^2*c^4*tan(e/2 + (f*x)/2)^7)/20 + (119*a^2*c^4*tan(e/2 + (f*x)/2)^9)/24 - (7*a^2*c^4*tan(e/2 + (f*x)/2)^11)/8 - (7*a^2*c^4*tan(e/2 + (f*x)/2))/8)/(f*(15*tan(e/2 + (f*x)/2)^4 - 6*tan(e/2 + (f*x)/2)^2 - 20*tan(e/2 + (f*x)/2)^6 + 15*tan(e/2 + (f*x)/2)^8 - 6*tan(e/2 + (f*x)/2)^10 + tan(e/2 + (f*x)/2)^12 + 1)) + (7*a^2*c^4*atanh(tan(e/2 + (f*x)/2)))/(8*f)
```

3.11. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^4 dx$

3.12 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx$

3.12.1	Optimal result	183
3.12.2	Mathematica [A] (verified)	183
3.12.3	Rubi [A] (verified)	184
3.12.4	Maple [C] (verified)	185
3.12.5	Fricas [A] (verification not implemented)	186
3.12.6	Sympy [F]	186
3.12.7	Maxima [B] (verification not implemented)	187
3.12.8	Giac [A] (verification not implemented)	187
3.12.9	Mupad [B] (verification not implemented)	188

3.12.1 Optimal result

Integrand size = 32, antiderivative size = 94

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx \\ &= \frac{3a^2c^3 \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{3a^2c^3 \sec(e + fx) \tan(e + fx)}{8f} \\ & \quad + \frac{a^2c^3 \sec(e + fx) \tan^3(e + fx)}{4f} - \frac{a^2c^3 \tan^5(e + fx)}{5f} \end{aligned}$$

output `3/8*a^2*c^3*arctanh(sin(f*x+e))/f-3/8*a^2*c^3*sec(f*x+e)*tan(f*x+e)/f+1/4*a^2*c^3*sec(f*x+e)*tan(f*x+e)^3/f-1/5*a^2*c^3*tan(f*x+e)^5/f`

3.12.2 Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.87

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx \\ &= \frac{a^2c^3(120 \operatorname{arctanh}(\sin(e + fx)) - \sec^5(e + fx)(40 \sin(e + fx) + 10 \sin(2(e + fx)) - 20 \sin(3(e + fx))) + 2}{320f} \end{aligned}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^3,x]`

output $(a^2 c^3 (120 \operatorname{ArcTanh}[\sin[e + f x]] - \sec[e + f x]^5 (40 \sin[e + f x] + 10 \sin[2(e + f x)] - 20 \sin[3(e + f x)] + 25 \sin[4(e + f x)] + 4 \sin[5(e + f x)])))/(320 f)$

3.12.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.86, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^3 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 dx$$

$$\downarrow 4446$$

$$a^2 c^2 \int (c \sec(e + fx) \tan^4(e + fx) - c \sec^2(e + fx) \tan^4(e + fx)) dx$$

$$\downarrow 2009$$

$$a^2 c^2 \left(\frac{3c \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{c \tan^5(e + fx)}{5f} + \frac{c \tan^3(e + fx) \sec(e + fx)}{4f} - \frac{3c \tan(e + fx) \sec(e + fx)}{8f} \right)$$

input $\operatorname{Int}[\sec[e + f x] * (a + a \sec[e + f x])^2 * (c - c \sec[e + f x])^3, x]$

output $a^2 c^2 * ((3 * c * \operatorname{ArcTanh}[\sin[e + f x]]) / (8 * f) - (3 * c * \sec[e + f x] * \tan[e + f x]) / (8 * f) + (c * \sec[e + f x] * \tan[e + f x]^3) / (4 * f) - (c * \tan[e + f x]^5) / (5 * f))$

3.12.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.12.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 6.15 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.52

method	result
risch	$\frac{ia^2c^3(25e^{9i(fx+e)} - 40e^{8i(fx+e)} + 10e^{7i(fx+e)} - 80e^{4i(fx+e)} - 10e^{3i(fx+e)} - 25e^{i(fx+e)} - 8)}{20f(1+e^{2i(fx+e)})^5} - \frac{3a^2c^3 \ln(e^{i(fx+e)} - i)}{8f}$
parts	$\frac{a^2c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f} - \frac{a^2c^3 \tan(fx+e)}{f} - \frac{a^2c^3 \sec(fx+e)}{f}$
norman	$\frac{3a^2c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f} - \frac{7a^2c^3 \tan^3\left(\frac{fx}{2} + \frac{e}{2}\right)}{2f} + \frac{32a^2c^3 \tan^5\left(\frac{fx}{2} + \frac{e}{2}\right)}{5f} + \frac{7a^2c^3 \tan^7\left(\frac{fx}{2} + \frac{e}{2}\right)}{2f} - \frac{3a^2c^3 \tan^9\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f} - \frac{3a^2c^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^5}$
parallelrisch	$-\frac{\left(\left(\frac{15 \cos(fx+e)}{2} + \frac{15 \cos(3fx+3e)}{4} + \frac{3 \cos(5fx+5e)}{4} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \left(-\frac{15 \cos(fx+e)}{2} - \frac{15 \cos(3fx+3e)}{4} - \frac{3 \cos(5fx+5e)}{4} \right) \right)}{2f(\cos(5fx+5e) + 5 \cos(3fx+3e))}$
derivativedivides	$\frac{a^2c^3 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) + a^2c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{\left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) + a^2c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}$
default	$\frac{a^2c^3 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) + a^2c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{\left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) + a^2c^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

$$3.12. \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx$$

output $\frac{1}{20}Ia^2c^3(25\exp(9I(fx+e))-40\exp(8I(fx+e))+10\exp(7I(fx+e))-80\exp(4I(fx+e))-10\exp(3I(fx+e))-25\exp(I(fx+e))-8)/f/(1+\exp(2I(fx+e)))^5-3/8a^2c^3/f*\ln(\exp(I(fx+e))-1)+3/8a^2c^3/f*\ln(\exp(I(fx+e))+1)$

3.12.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.54

$$\int \sec(e+fx)(a+a\sec(e+fx))^2(c-c\sec(e+fx))^3 dx$$

$$= \frac{15a^2c^3 \cos(fx+e)^5 \log(\sin(fx+e)+1) - 15a^2c^3 \cos(fx+e)^5 \log(-\sin(fx+e)+1) - 2(8a^2c^3 \cos(fx+e)^4 + 25a^2c^3 \cos(fx+e)^3 - 16a^2c^3 \cos(fx+e)^2 - 10a^2c^3 \cos(fx+e) + 8a^2c^3 \sin(fx+e))}{80f \cos(fx+e)^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^3,x, algorithm="fricas")`

output $\frac{1}{80}(15a^2c^3\cos(fx+e)^5*\log(\sin(fx+e)+1) - 15a^2c^3\cos(fx+e)^5*\log(-\sin(fx+e)+1) - 2*(8a^2c^3\cos(fx+e)^4 + 25a^2c^3\cos(fx+e)^3 - 16a^2c^3\cos(fx+e)^2 - 10a^2c^3\cos(fx+e) + 8a^2c^3*\sin(fx+e))/(f*\cos(fx+e)^5)$

3.12.6 Sympy [F]

$$\int \sec(e+fx)(a+a\sec(e+fx))^2(c-c\sec(e+fx))^3 dx$$

$$= -a^2c^3 \left(\int (-\sec(e+fx)) dx + \int \sec^2(e+fx) dx + \int 2\sec^3(e+fx) dx \right. \\ \left. + \int (-2\sec^4(e+fx)) dx + \int (-\sec^5(e+fx)) dx + \int \sec^6(e+fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**3,x)`

output `-a**2*c**3*(Integral(-sec(e + f*x), x) + Integral(sec(e + f*x)**2, x) + Integral(2*sec(e + f*x)**3, x) + Integral(-2*sec(e + f*x)**4, x) + Integral(-sec(e + f*x)**5, x) + Integral(sec(e + f*x)**6, x))`

3.12. $\int \sec(e+fx)(a+a\sec(e+fx))^2(c-c\sec(e+fx))^3 dx$

3.12.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(86) = 172.

Time = 0.21 (sec) , antiderivative size = 227, normalized size of antiderivative = 2.41

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx =$$

$$16(3 \tan(fx + e)^5 + 10 \tan(fx + e)^3 + 15 \tan(fx + e))a^2c^3 - 160(\tan(fx + e)^3 + 3 \tan(fx + e))$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `-1/240*(16*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^2*c^3 - 160*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^2*c^3 + 15*a^2*c^3*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 120*a^2*c^3*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 240*a^2*c^3*log(sec(f*x + e) + tan(f*x + e)) + 240*a^2*c^3*tan(f*x + e))/f`

3.12.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.69

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx$$

$$= \frac{15a^2c^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 15a^2c^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right) - \frac{2\left(15a^2c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^9 - 70a^2c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^7 + 128a^2c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 70a^2c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 - 15a^2c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)^2 - 1}{40f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `1/40*(15*a^2*c^3*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*a^2*c^3*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(15*a^2*c^3*tan(1/2*f*x + 1/2*e)^9 - 70*a^2*c^3*tan(1/2*f*x + 1/2*e)^7 - 128*a^2*c^3*tan(1/2*f*x + 1/2*e)^5 + 70*a^2*c^3*tan(1/2*f*x + 1/2*e)^3 - 15*a^2*c^3*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^5/f`

3.12. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx$

3.12.9 Mupad [B] (verification not implemented)

Time = 18.31 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.99

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^3 dx$$

$$= \frac{-\frac{3a^2c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{4} + \frac{7a^2c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{2} + \frac{32a^2c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{5} - \frac{7a^2c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{2} + \frac{3a^2c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1 \right)} + \frac{3a^2c^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{4f}$$

input `int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^3)/cos(e + f*x),x)`output `((32*a^2*c^3*tan(e/2 + (f*x)/2)^5)/5 - (7*a^2*c^3*tan(e/2 + (f*x)/2)^3)/2 + (7*a^2*c^3*tan(e/2 + (f*x)/2)^7)/2 - (3*a^2*c^3*tan(e/2 + (f*x)/2)^9)/4 + (3*a^2*c^3*tan(e/2 + (f*x)/2))/4)/(f*(5*tan(e/2 + (f*x)/2)^2 - 10*tan(e/2 + (f*x)/2)^4 + 10*tan(e/2 + (f*x)/2)^6 - 5*tan(e/2 + (f*x)/2)^8 + tan(e/2 + (f*x)/2)^10 - 1)) + (3*a^2*c^3*atanh(tan(e/2 + (f*x)/2)))/(4*f)`

3.13 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$

3.13.1	Optimal result	189
3.13.2	Mathematica [A] (verified)	189
3.13.3	Rubi [A] (verified)	190
3.13.4	Maple [A] (verified)	192
3.13.5	Fricas [A] (verification not implemented)	192
3.13.6	Sympy [F]	193
3.13.7	Maxima [B] (verification not implemented)	193
3.13.8	Giac [A] (verification not implemented)	194
3.13.9	Mupad [B] (verification not implemented)	194

3.13.1 Optimal result

Integrand size = 32, antiderivative size = 73

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx \\ &= \frac{3a^2c^2 \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{3a^2c^2 \sec(e + fx) \tan(e + fx)}{8f} \\ & \quad + \frac{a^2c^2 \sec(e + fx) \tan^3(e + fx)}{4f} \end{aligned}$$

output `3/8*a^2*c^2*arctanh(sin(f*x+e))/f-3/8*a^2*c^2*sec(f*x+e)*tan(f*x+e)/f+1/4*a^2*c^2*sec(f*x+e)*tan(f*x+e)^3/f`

3.13.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.10

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx \\ &= a^2c^2 \left(\frac{3 \operatorname{arctanh}(\sin(e + fx))}{8f} + \frac{3 \sec(e + fx) \tan(e + fx)}{8f} - \frac{3 \sec^3(e + fx) \tan(e + fx)}{4f} \right. \\ & \quad \left. + \frac{\sec(e + fx) \tan^3(e + fx)}{f} \right) \end{aligned}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^2,x]`

output $a^2c^2\left(\frac{3\text{ArcTanh}[\text{Sin}[e + f*x]]}{8f} + \frac{3\text{Sec}[e + f*x]\text{Tan}[e + f*x]}{8f} - \frac{3\text{Sec}[e + f*x]^3\text{Tan}[e + f*x]}{4f} + \frac{(\text{Sec}[e + f*x]\text{Tan}[e + f*x]^3)}{f}\right)$

3.13.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.92, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4446, 3042, 3091, 3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^2 dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 dx \\
 & \quad \downarrow \text{4446} \\
 & a^2c^2 \int \sec(e + fx) \tan^4(e + fx) dx \\
 & \quad \downarrow \text{3042} \\
 & a^2c^2 \int \sec(e + fx) \tan(e + fx)^4 dx \\
 & \quad \downarrow \text{3091} \\
 & a^2c^2 \left(\frac{\tan^3(e + fx) \sec(e + fx)}{4f} - \frac{3}{4} \int \sec(e + fx) \tan^2(e + fx) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & a^2c^2 \left(\frac{\tan^3(e + fx) \sec(e + fx)}{4f} - \frac{3}{4} \int \sec(e + fx) \tan(e + fx)^2 dx \right) \\
 & \quad \downarrow \text{3091} \\
 & a^2c^2 \left(\frac{\tan^3(e + fx) \sec(e + fx)}{4f} - \frac{3}{4} \left(\frac{\tan(e + fx) \sec(e + fx)}{2f} - \frac{1}{2} \int \sec(e + fx) dx \right) \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.13. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$

$$a^2 c^2 \left(\frac{\tan^3(e + fx) \sec(e + fx)}{4f} - \frac{3}{4} \left(\frac{\tan(e + fx) \sec(e + fx)}{2f} - \frac{1}{2} \int \csc \left(e + fx + \frac{\pi}{2} \right) dx \right) \right)$$

↓ 4257

$$a^2 c^2 \left(\frac{\tan^3(e + fx) \sec(e + fx)}{4f} - \frac{3}{4} \left(\frac{\tan(e + fx) \sec(e + fx)}{2f} - \frac{\operatorname{arctanh}(\sin(e + fx))}{2f} \right) \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^2,x]`

output `a^2*c^2*((Sec[e + f*x]*Tan[e + f*x]^3)/(4*f) - (3*(-1/2*ArcTanh[Sin[e + f*x]]/f + (Sec[e + f*x]*Tan[e + f*x])/(2*f)))/4)`

3.13.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3091 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.13.4 Maple [A] (verified)

Time = 2.64 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.08

method	result
parts	$\frac{a^2 c^2 \left(- \left(- \frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f} - \frac{a^2 c^2 \sec(fx+e) \tan(fx+e)}{f}$
derivativedivides	$\frac{a^2 c^2 \left(- \left(- \frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right) - 2a^2 c^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right)}{f}$
default	$\frac{a^2 c^2 \left(- \left(- \frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right) - 2a^2 c^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right)}{f}$
risch	$\frac{ia^2 c^2 (5 e^{7i(fx+e)} - 3 e^{5i(fx+e)} + 3 e^{3i(fx+e)} - 5 e^{i(fx+e)})}{4f(1+e^{2i(fx+e)})^4} - \frac{3a^2 c^2 \ln(e^{i(fx+e)} - i)}{8f} + \frac{3a^2 c^2 \ln(e^{i(fx+e)} + i)}{8f}$
parallelrisc	$-\frac{3a^2 \left(\left(\frac{3}{4} + \frac{\cos(4fx+4e)}{4} \right) + \cos(2fx+2e) \right) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) + \left(-\cos(2fx+2e) - \frac{\cos(4fx+4e)}{4} - \frac{3}{4} \right) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)}{2f(3 + \cos(4fx+4e) + 4 \cos(2fx+2e))}$
norman	$\frac{-\frac{3a^2 c^2 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)}{4f} + \frac{11a^2 c^2 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^3}{4f} + \frac{11a^2 c^2 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^5}{4f} - \frac{3a^2 c^2 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^7}{4f}}{\left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right)^2 - 1 \right)^4} - \frac{3a^2 c^2 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)}{8f} + \dots$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output a^2*c^2/f*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln(sec(f*x+e)+tan(f*x+e)))-a^2*c^2*sec(f*x+e)*tan(f*x+e)/f
```

3.13.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.36

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$$

$$= \frac{3 a^2 c^2 \cos (f x + e)^4 \log (\sin (f x + e) + 1) - 3 a^2 c^2 \cos (f x + e)^4 \log (-\sin (f x + e) + 1) - 2 \left(5 a^2 c^2 \cos (f x + e)^4 \log (\tan (f x + e) + 1) - 5 a^2 c^2 \cos (f x + e)^4 \log (\tan (f x + e) - 1) \right)}{16 f \cos (f x + e)^4}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^2,x, algorithm="fracas")
```

3.13. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$

output $1/16*(3*a^2*c^2*\cos(f*x + e)^4*\log(\sin(f*x + e) + 1) - 3*a^2*c^2*\cos(f*x + e)^4*\log(-\sin(f*x + e) + 1) - 2*(5*a^2*c^2*\cos(f*x + e)^2 - 2*a^2*c^2)*\sin(f*x + e))/(f*\cos(f*x + e)^4)$

3.13.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$$

$$= a^2 c^2 \left(\int \sec(e + fx) dx + \int (-2 \sec^3(e + fx)) dx + \int \sec^5(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**2,x)`

output `a**2*c**2*(Integral(sec(e + f*x), x) + Integral(-2*sec(e + f*x)**3, x) + Integral(sec(e + f*x)**5, x))`

3.13.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 150 vs. $2(67) = 134$.

Time = 0.21 (sec) , antiderivative size = 150, normalized size of antiderivative = 2.05

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx =$$

$$\frac{a^2 c^2 \left(\frac{2(3 \sin(fx+e)^3 - 5 \sin(fx+e))}{\sin(fx+e)^4 - 2 \sin(fx+e)^2 + 1} - 3 \log(\sin(fx + e) + 1) + 3 \log(\sin(fx + e) - 1) \right) - 8 a^2 c^2 \left(\frac{2 \sin(fx+e)}{\sin(fx+e)^2} \right)}{16 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output $-1/16*(a^2*c^2*(2*(3*\sin(f*x + e)^3 - 5*\sin(f*x + e))/(\sin(f*x + e)^4 - 2*\sin(f*x + e)^2 + 1) - 3*\log(\sin(f*x + e) + 1) + 3*\log(\sin(f*x + e) - 1)) - 8*a^2*c^2*(2*\sin(f*x + e)/(\sin(f*x + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) - 16*a^2*c^2*\log(\sec(f*x + e) + \tan(f*x + e)))/f$

3.13.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.19

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$$

$$= \frac{3a^2c^2 \log(|\sin(fx + e) + 1|) - 3a^2c^2 \log(|\sin(fx + e) - 1|) + \frac{2(5a^2c^2 \sin(fx+e)^3 - 3a^2c^2 \sin(fx+e))}{(\sin(fx+e)^2 - 1)^2}}{16f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^2,x, algorithm="giac")
```

```
output 1/16*(3*a^2*c^2*log(abs(sin(f*x + e) + 1)) - 3*a^2*c^2*log(abs(sin(f*x + e) - 1)) + 2*(5*a^2*c^2*sin(f*x + e)^3 - 3*a^2*c^2*sin(f*x + e))/(sin(f*x + e)^2 - 1)^2)/f
```

3.13.9 Mupad [B] (verification not implemented)

Time = 16.97 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.12

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^2 dx$$

$$= \frac{-\frac{3a^2c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{4} + \frac{11a^2c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{4} + \frac{11a^2c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{4} - \frac{3a^2c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1 \right)} + \frac{3a^2c^2 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{4f}$$

```
input int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^2)/cos(e + f*x),x)
```

```
output ((11*a^2*c^2*tan(e/2 + (f*x)/2)^3)/4 + (11*a^2*c^2*tan(e/2 + (f*x)/2)^5)/4 - (3*a^2*c^2*tan(e/2 + (f*x)/2)^7)/4 - (3*a^2*c^2*tan(e/2 + (f*x)/2))/4)/(f*(6*tan(e/2 + (f*x)/2)^4 - 4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/2)^8 + 1)) + (3*a^2*c^2*atanh(tan(e/2 + (f*x)/2)))/(4*f)
```

3.14 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$

3.14.1	Optimal result	195
3.14.2	Mathematica [A] (verified)	195
3.14.3	Rubi [A] (verified)	196
3.14.4	Maple [A] (verified)	197
3.14.5	Fricas [A] (verification not implemented)	198
3.14.6	Sympy [F]	198
3.14.7	Maxima [A] (verification not implemented)	199
3.14.8	Giac [B] (verification not implemented)	199
3.14.9	Mupad [B] (verification not implemented)	200

3.14.1 Optimal result

Integrand size = 30, antiderivative size = 61

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{a^2 c \operatorname{arctanh}(\sin(e + fx))}{2f} - \frac{a^2 c \sec(e + fx) \tan(e + fx)}{2f} - \frac{a^2 c \tan^3(e + fx)}{3f}$$

output `1/2*a^2*c*arctanh(sin(f*x+e))/f-1/2*a^2*c*sec(f*x+e)*tan(f*x+e)/f-1/3*a^2*c*tan(f*x+e)^3/f`

3.14.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.74

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{a^2 c (3 \operatorname{ArcTanh}[\sin(e + fx)] - 3 \sec(e + fx) \tan(e + fx) - 2 \tan^3(e + fx))}{6f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x]),x]`

output `(a^2*c*(3*ArcTanh[Sin[e + f*x]] - 3*Sec[e + f*x]*Tan[e + f*x] - 2*Tan[e + f*x]^3))/(6*f)`

3.14. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$

3.14.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx)) dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow 4446$$

$$-ac \int (a \sec^2(e + fx) \tan^2(e + fx) + a \sec(e + fx) \tan^2(e + fx)) dx$$

$$\downarrow 2009$$

$$-ac \left(-\frac{a \operatorname{arctanh}(\sin(e + fx))}{2f} + \frac{a \tan^3(e + fx)}{3f} + \frac{a \tan(e + fx) \sec(e + fx)}{2f} \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x]),x]`

output `-(a*c*(-1/2*(a*ArcTanh[Sin[e + f*x]])/f + (a*Sec[e + f*x]*Tan[e + f*x])/(2*f) + (a*Tan[e + f*x]^3)/(3*f))`

3.14.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.14.4 Maple [A] (verified)

Time = 4.21 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.57

method	result
derivativedivides	$\frac{a^2 c \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) - a^2 c \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) + a^2 c \tan(fx+e) + a^2 c \ln(\sec(fx+e) + \tan(fx+e))}{f}$
default	$\frac{a^2 c \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) - a^2 c \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) + a^2 c \tan(fx+e) + a^2 c \ln(\sec(fx+e) + \tan(fx+e))}{f}$
risch	$\frac{ia^2 c (3e^{5i(fx+e)} + 6e^{4i(fx+e)} - 3e^{i(fx+e)} + 2)}{3f(1+e^{2i(fx+e)})^3} + \frac{a^2 c \ln(e^{i(fx+e)} + i)}{2f} - \frac{a^2 c \ln(e^{i(fx+e)} - i)}{2f}$
parts	$\frac{a^2 c \ln(\sec(fx+e) + \tan(fx+e))}{f} + \frac{a^2 c \tan(fx+e)}{f} - \frac{a^2 c \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right)}{f} + \frac{a^2 c \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e)}{f}$
norman	$\frac{\frac{a^2 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f} + \frac{8a^2 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3f} - \frac{a^2 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^3} - \frac{a^2 c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2f} + \frac{a^2 c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2f}$
parallelrisc	$\frac{a^2 \left(\frac{3(-\cos(3fx+3e) - 3\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2} + \frac{3(\cos(3fx+3e) + 3\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} + \sin(3fx+3e) - 3\sin(fx+e) \right)}{3f(\cos(3fx+3e) + 3\cos(fx+e))}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE
)
```

```
output 1/f*(a^2*c*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)-a^2*c*(1/2*sec(f*x+e)*tan(f*
x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+a^2*c*tan(f*x+e)+a^2*c*ln(sec(f*x+e)+t
an(f*x+e)))
```

3.14. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$

3.14.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.69

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{3a^2c \cos(fx + e)^3 \log(\sin(fx + e) + 1) - 3a^2c \cos(fx + e)^3 \log(-\sin(fx + e) + 1) + 2(2a^2c \cos(fx + e) - 2a^2c) \sin(fx + e)}{12f \cos(fx + e)^3}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `1/12*(3*a^2*c*cos(f*x + e)^3*log(sin(f*x + e) + 1) - 3*a^2*c*cos(f*x + e)^3*log(-sin(f*x + e) + 1) + 2*(2*a^2*c*cos(f*x + e)^2 - 3*a^2*c*cos(f*x + e) - 2*a^2*c)*sin(f*x + e))/(f*cos(f*x + e)^3)`

3.14.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= -a^2c \left(\int (-\sec(e + fx)) dx + \int (-\sec^2(e + fx)) dx + \int \sec^3(e + fx) dx + \int \sec^4(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e)),x)`

output `-a**2*c*(Integral(-sec(e + f*x), x) + Integral(-sec(e + f*x)**2, x) + Integral(sec(e + f*x)**3, x) + Integral(sec(e + f*x)**4, x))`

3.14.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.77

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx =$$

$$\frac{4(\tan(fx + e)^3 + 3 \tan(fx + e))a^2c - 3a^2c\left(\frac{2 \sin(fx + e)}{\sin(fx + e)^2 - 1} - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1)\right) - 12a^2c \tan(fx + e)}{12f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-1/12*(4*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^2*c - 3*a^2*c*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 12*a^2*c*log(sec(f*x + e) + tan(f*x + e)) - 12*a^2*c*tan(f*x + e))/f`

3.14.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. 2(55) = 110.

Time = 0.31 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.82

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{3a^2c \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3a^2c \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right) - \frac{2\left(3a^2c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 8a^2c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)^2 - 1}}{6f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="giac")`

output `1/6*(3*a^2*c*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 3*a^2*c*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(3*a^2*c*tan(1/2*f*x + 1/2*e)^5 - 8*a^2*c*tan(1/2*f*x + 1/2*e)^3 - 3*a^2*c*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^3)/f`

3.14.9 Mupad [B] (verification not implemented)

Time = 15.88 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.85

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{-ca^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \frac{8ca^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3} + ca^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)} + \frac{a^2 c \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{f}$$

input `int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x)))/cos(e + f*x),x)`

output `(a^2*c*tan(e/2 + (f*x)/2) + (8*a^2*c*tan(e/2 + (f*x)/2)^3)/3 - a^2*c*tan(e/2 + (f*x)/2)^5)/(f*(3*tan(e/2 + (f*x)/2)^2 - 3*tan(e/2 + (f*x)/2)^4 + tan(e/2 + (f*x)/2)^6 - 1)) + (a^2*c*atanh(tan(e/2 + (f*x)/2)))/f`

3.15 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c-c \sec(e+fx)} dx$

3.15.1	Optimal result	201
3.15.2	Mathematica [C] (verified)	201
3.15.3	Rubi [A] (verified)	202
3.15.4	Maple [A] (verified)	204
3.15.5	Fricas [A] (verification not implemented)	204
3.15.6	Sympy [F]	205
3.15.7	Maxima [B] (verification not implemented)	205
3.15.8	Giac [A] (verification not implemented)	206
3.15.9	Mupad [B] (verification not implemented)	206

3.15.1 Optimal result

Integrand size = 32, antiderivative size = 74

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c-c \sec(e+fx)} dx = -\frac{3a^2 \operatorname{arctanh}(\sin(e+fx))}{cf} - \frac{3a^2 \tan(e+fx)}{cf} - \frac{2(a^2+a^2 \sec(e+fx)) \tan(e+fx)}{f(c-c \sec(e+fx))}$$

```
output -3*a^2*arctanh(sin(f*x+e))/c/f-3*a^2*tan(f*x+e)/c/f-2*(a^2+a^2*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))
```

3.15.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.59 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.88

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c-c \sec(e+fx)} dx = -\frac{a^2 \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{1}{2}(1+\sec(e+fx))\right) (1+\sec(e+fx))^2 \tan(e+fx)}{5cf \sqrt{2-2\sec(e+fx)}}$$

```
input Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x]),x]
```

```
output -1/5*(a^2*Hypergeometric2F1[3/2, 5/2, 7/2, (1 + Sec[e + f*x])/2]*(1 + Sec[e + f*x])^2*Tan[e + f*x])/(c*f*Sqrt[2 - 2*Sec[e + f*x]])
```

3.15.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4445, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{c-c\sec(e+fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{c-c\csc(e+fx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{3a \int \sec(e+fx)(\sec(e+fx)a+a) dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3a \int \csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a) dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{4274} \\
 & -\frac{3a(a \int \sec^2(e+fx) dx + a \int \sec(e+fx) dx)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3a(a \int \csc(e+fx+\frac{\pi}{2}) dx + a \int \csc(e+fx+\frac{\pi}{2})^2 dx)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{4254} \\
 & -\frac{3a(a \int \csc(e+fx+\frac{\pi}{2}) dx - \frac{a \int 1d(-\tan(e+fx))}{f})}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{24} \\
 & -\frac{3a(a \int \csc(e+fx+\frac{\pi}{2}) dx + \frac{a \tan(e+fx)}{f})}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

3.15. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c-c\sec(e+fx)} dx$

$$\frac{2 \tan(e + fx) (a^2 \sec(e + fx) + a^2)}{f(c - c \sec(e + fx))} - \frac{3a \left(\frac{a \operatorname{arctanh}(\frac{\sin(e+fx)}{f})}{f} + \frac{a \tan(e+fx)}{f} \right)}{c}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x]),x]`

output `(-2*(a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])) - (3*a*((a*ArcTanh[Sin[e + f*x]])/f + (a*Tan[e + f*x])/f))/c`

3.15.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.15.4 Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{4a^2 \left(\frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 4} - \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} + \frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 4} + \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} \right)}{fc}$
default	$\frac{4a^2 \left(\frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 4} - \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} + \frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 4} + \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} \right)}{fc}$
parallelrisc	$- \frac{a^2 \left(-3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) \cos(fx+e) + 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \cos(fx+e) - 5 \cot\left(\frac{fx}{2} + \frac{e}{2}\right) \cos(fx+e) + \cot\left(\frac{fx}{2} + \frac{e}{2}\right) \right)}{cf \cos(fx+e)}$
risc	$\frac{2ia^2 (4e^{2i(fx+e)} - e^{i(fx+e)} + 5)}{fc(1+e^{2i(fx+e)})(e^{i(fx+e)} - 1)} + \frac{3a^2 \ln(e^{i(fx+e)} - i)}{cf} - \frac{3a^2 \ln(e^{i(fx+e)} + i)}{cf}$
norman	$\frac{\frac{4a^2}{cf} - \frac{10a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{cf} + \frac{6a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)} + \frac{3a^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{cf} - \frac{3a^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{cf}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `4/f*a^2/c*(1/4/(tan(1/2*f*x+1/2*e)+1)-3/4*ln(tan(1/2*f*x+1/2*e)+1)+1/4/(tan(1/2*f*x+1/2*e)-1)+3/4*ln(tan(1/2*f*x+1/2*e)-1)+1/tan(1/2*f*x+1/2*e))`

3.15.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.46

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c-c\sec(e+fx)} dx = \frac{3a^2 \cos(fx+e) \log(\sin(fx+e)+1) \sin(fx+e) - 3a^2 \cos(fx+e) \log(-\sin(fx+e)+1) \sin(fx+e)}{2cf \cos(fx+e) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `-1/2*(3*a^2*cos(f*x + e)*log(sin(f*x + e) + 1)*sin(f*x + e) - 3*a^2*cos(f*x + e)*log(-sin(f*x + e) + 1)*sin(f*x + e) - 10*a^2*cos(f*x + e)^2 - 8*a^2*cos(f*x + e) + 2*a^2)/(c*f*cos(f*x + e)*sin(f*x + e))`

3.15. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c-c\sec(e+fx)} dx$

3.15.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c-c\sec(e+fx)} dx$$

$$= -\frac{a^2 \left(\int \frac{\sec(e+fx)}{\sec(e+fx)-1} dx + \int \frac{2\sec^2(e+fx)}{\sec(e+fx)-1} dx + \int \frac{\sec^3(e+fx)}{\sec(e+fx)-1} dx \right)}{c}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e)),x)`

output `-a**2*(Integral(sec(e + f*x)/(sec(e + f*x) - 1), x) + Integral(2*sec(e + f*x)**2/(sec(e + f*x) - 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x) - 1), x))/c`

3.15.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 225 vs. $2(75) = 150$.

Time = 0.21 (sec) , antiderivative size = 225, normalized size of antiderivative = 3.04

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c-c\sec(e+fx)} dx =$$

$$\frac{a^2 \left(\frac{\frac{3\sin(fx+e)^2}{(\cos(fx+e)+1)^2} - 1}{\frac{c\sin(fx+e)}{\cos(fx+e)+1} - \frac{c\sin(fx+e)^3}{(\cos(fx+e)+1)^3}} + \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{c} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{c} \right) + 2a^2 \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{c} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{c} \right)}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-(a^2*((3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 1)/(c*sin(f*x + e)/(cos(f*x + e) + 1) - c*sin(f*x + e)^3/(cos(f*x + e) + 1)^3) + log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/c - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/c) + 2*a^2*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/c - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/c - (cos(f*x + e) + 1)/(c*sin(f*x + e))) - a^2*(cos(f*x + e) + 1)/(c*sin(f*x + e)))/f`

3.15.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.35

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c-c\sec(e+fx)} dx$$

$$= -\frac{\frac{3a^2 \log(|\tan(\frac{1}{2}fx+\frac{1}{2}e)+1|)}{c} - \frac{3a^2 \log(|\tan(\frac{1}{2}fx+\frac{1}{2}e)-1|)}{c} - \frac{2(3a^2 \tan(\frac{1}{2}fx+\frac{1}{2}e)^2 - 2a^2)}{(\tan(\frac{1}{2}fx+\frac{1}{2}e)^3 - \tan(\frac{1}{2}fx+\frac{1}{2}e))c}}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x, algorithm="giac")`

output `-(3*a^2*log(abs(tan(1/2*f*x + 1/2*e) + 1))/c - 3*a^2*log(abs(tan(1/2*f*x + 1/2*e) - 1))/c - 2*(3*a^2*tan(1/2*f*x + 1/2*e)^2 - 2*a^2)/((tan(1/2*f*x + 1/2*e)^3 - tan(1/2*f*x + 1/2*e))*c))/f`

3.15.9 Mupad [B] (verification not implemented)

Time = 13.77 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.04

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c-c\sec(e+fx)} dx = \frac{6a^2 \tan(\frac{e}{2} + \frac{fx}{2})^2 - 4a^2}{cf \tan(\frac{e}{2} + \frac{fx}{2}) (\tan(\frac{e}{2} + \frac{fx}{2})^2 - 1)} - \frac{6a^2 \operatorname{atanh}(\tan(\frac{e}{2} + \frac{fx}{2}))}{cf}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))),x)`

output `(6*a^2*tan(e/2 + (f*x)/2)^2 - 4*a^2)/(c*f*tan(e/2 + (f*x)/2)*(tan(e/2 + (f*x)/2)^2 - 1)) - (6*a^2*atanh(tan(e/2 + (f*x)/2)))/(c*f)`

3.16 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^2} dx$

3.16.1 Optimal result 207
 3.16.2 Mathematica [A] (verified) 207
 3.16.3 Rubi [A] (verified) 208
 3.16.4 Maple [A] (verified) 209
 3.16.5 Fricas [A] (verification not implemented) 210
 3.16.6 Sympy [F] 211
 3.16.7 Maxima [B] (verification not implemented) 211
 3.16.8 Giac [A] (verification not implemented) 212
 3.16.9 Mupad [B] (verification not implemented) 212

3.16.1 Optimal result

Integrand size = 32, antiderivative size = 89

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^2} dx = \frac{a^2 \operatorname{arctanh}(\sin(e+fx))}{c^2 f} - \frac{2(a^2 + a^2 \sec(e+fx)) \tan(e+fx)}{3f(c-c \sec(e+fx))^2} + \frac{2a^2 \tan(e+fx)}{f(c^2 - c^2 \sec(e+fx))}$$

output `a^2*arctanh(sin(f*x+e))/c^2/f-2/3*(a^2+a^2*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^2+2*a^2*tan(f*x+e)/f/(c^2-c^2*sec(f*x+e))`

3.16.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.22

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^2} dx = \frac{a^2 \left(-\frac{4 \cot(\frac{1}{2}(e+fx))}{3f} - \frac{2 \cot(\frac{1}{2}(e+fx)) \operatorname{csc}^2(\frac{1}{2}(e+fx))}{3f} - \frac{\log(\cos(\frac{1}{2}(e+fx)) - \sin(\frac{1}{2}(e+fx)))}{f} + \frac{\log(\cos(\frac{1}{2}(e+fx)) + \sin(\frac{1}{2}(e+fx)))}{f} \right)}{c^2}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^2,x]`

output $(a^2 * ((-4 * \cot[(e + f*x)/2]) / (3*f) - (2 * \cot[(e + f*x)/2] * \csc[(e + f*x)/2]^2) / (3*f) - \log[\cos[(e + f*x)/2] - \sin[(e + f*x)/2]] / f + \log[\cos[(e + f*x)/2] + \sin[(e + f*x)/2]] / f) / c^2$

3.16.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4445, 3042, 4445, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a \sec(e+fx)+a)^2}{(c-c \sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a \csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c \csc(e+fx+\frac{\pi}{2}))^2} dx$$

↓ 4445

$$-\frac{a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{c-c \sec(e+fx)} dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c \sec(e+fx))^2}$$

↓ 3042

$$-\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{c-c \csc(e+fx+\frac{\pi}{2})} dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c \sec(e+fx))^2}$$

↓ 4445

$$-\frac{a \left(-\frac{a \int \sec(e+fx) dx}{c} - \frac{2a \tan(e+fx)}{f(c-c \sec(e+fx))} \right)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c \sec(e+fx))^2}$$

↓ 3042

$$-\frac{a \left(-\frac{a \int \csc(e+fx+\frac{\pi}{2}) dx}{c} - \frac{2a \tan(e+fx)}{f(c-c \sec(e+fx))} \right)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c \sec(e+fx))^2}$$

↓ 4257

$$-\frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c \sec(e+fx))^2} - \frac{a \left(-\frac{a \operatorname{arctanh}(\sin(e+fx))}{cf} - \frac{2a \tan(e+fx)}{f(c-c \sec(e+fx))} \right)}{c}$$

3.16. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^2} dx$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^2,x]`

output `(-2*(a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(3*f*(c - c*Sec[e + f*x])^2) - (a*(-((a*ArcTanh[Sin[e + f*x]])/(c*f)) - (2*a*Tan[e + f*x])/(f*(c - c*Sec[e + f*x]))))/c`

3.16.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^n, x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.16.4 Maple [A] (verified)

Time = 1.00 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.73

3.16. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^2} dx$

method	result
parallelrisch	$\frac{a^2 \left(-2 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) - 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - 6 \cot\left(\frac{fx}{2} + \frac{e}{2}\right) \right)}{3c^2 f}$
derivativdivides	$\frac{2a^2 \left(-\frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} - \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} \right)}{f c^2}$
default	$\frac{2a^2 \left(-\frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} - \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} \right)}{f c^2}$
risch	$\frac{8ia^2 (3e^{i(fx+e)} - 1)}{3f c^2 (e^{i(fx+e)} - 1)^3} + \frac{a^2 \ln(e^{i(fx+e)} + i)}{c^2 f} - \frac{a^2 \ln(e^{i(fx+e)} - i)}{c^2 f}$
norman	$\frac{-\frac{2a^2}{3cf} - \frac{2a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{3cf} + \frac{10a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{3cf} - \frac{2a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{a^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{c^2 f} - \frac{a^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{c^2 f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 1/3*a^2*(-2*cot(1/2*f*x+1/2*e)^3+3*ln(tan(1/2*f*x+1/2*e)+1)-3*ln(tan(1/2*f*x+1/2*e)-1)-6*cot(1/2*f*x+1/2*e))/c^2/f
```

3.16.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.44

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^2} dx = \frac{8a^2 \cos^2(fx + e) - 8a^2 \cos(fx + e) - 3(a^2 \cos(fx + e) - a^2) \log(\sin(fx + e) + 1) \sin(fx + e) + 3(6(c^2 f \cos(fx + e) - c^2 f) \sin(fx + e))}{6(c^2 f \cos(fx + e) - c^2 f) \sin(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x, algorithm="fricas")
```

```
output -1/6*(8*a^2*cos(f*x + e)^2 - 8*a^2*cos(f*x + e) - 3*(a^2*cos(f*x + e) - a^2)*log(sin(f*x + e) + 1)*sin(f*x + e) + 3*(a^2*cos(f*x + e) - a^2)*log(-sin(f*x + e) + 1)*sin(f*x + e) - 16*a^2)/((c^2*f*cos(f*x + e) - c^2*f)*sin(f*x + e))
```

3.16. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^2} dx$

3.16.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^2} dx$$

$$= \frac{a^2 \left(\int \frac{\sec(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx + \int \frac{2\sec^2(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx + \int \frac{\sec^3(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx \right)}{c^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**2,x)`

output `a**2*(Integral(sec(e+f*x)/(sec(e+f*x)**2-2*sec(e+f*x)+1),x)+Integral(2*sec(e+f*x)**2/(sec(e+f*x)**2-2*sec(e+f*x)+1),x)+Integral(sec(e+f*x)**3/(sec(e+f*x)**2-2*sec(e+f*x)+1),x))/c**2`

3.16.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(89) = 178.

Time = 0.21 (sec) , antiderivative size = 201, normalized size of antiderivative = 2.26

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^2} dx$$

$$= \frac{a^2 \left(\frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}+1\right)}{c^2} - \frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}-1\right)}{c^2} - \frac{\left(\frac{9 \sin^2(fx+e)}{(\cos(fx+e)+1)^2}+1\right)(\cos(fx+e)+1)^3}{c^2 \sin^3(fx+e)} \right) - \frac{2a^2 \left(\frac{3 \sin^2(fx+e)}{(\cos(fx+e)+1)^2}+1\right)(\cos(fx+e))}{c^2 \sin^3(fx+e)}}{6f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output `1/6*(a^2*(6*log(sin(f*x+e)/(cos(f*x+e)+1)+1)/c^2-6*log(sin(f*x+e)/(cos(f*x+e)+1)-1)/c^2-(9*sin(f*x+e)^2/(cos(f*x+e)+1)^2+1)*(cos(f*x+e)+1)^3/(c^2*sin(f*x+e)^3))-2*a^2*(3*sin(f*x+e)^2/(cos(f*x+e)+1)^2+1)*(cos(f*x+e)+1)^3/(c^2*sin(f*x+e)^3)+a^2*(3*sin(f*x+e)^2/(cos(f*x+e)+1)^2-1)*(cos(f*x+e)+1)^3/(c^2*sin(f*x+e)^3))/f`

3.16.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.94

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^2} dx$$

$$= \frac{\frac{3a^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{c^2} - \frac{3a^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{c^2} - \frac{2(3a^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 + a^2)}{c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3}}{3f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `1/3*(3*a^2*log(abs(tan(1/2*f*x + 1/2*e) + 1))/c^2 - 3*a^2*log(abs(tan(1/2*f*x + 1/2*e) - 1))/c^2 - 2*(3*a^2*tan(1/2*f*x + 1/2*e)^2 + a^2)/(c^2*tan(1/2*f*x + 1/2*e)^3))/f`

3.16.9 Mupad [B] (verification not implemented)

Time = 13.79 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.71

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^2} dx = \frac{2a^2 \operatorname{atanh}(\tan(\frac{e}{2} + \frac{fx}{2}))}{c^2 f} - \frac{2a^2 \tan(\frac{e}{2} + \frac{fx}{2})^2 + \frac{2a^2}{3}}{c^2 f \tan(\frac{e}{2} + \frac{fx}{2})^3}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^2),x)`

output `(2*a^2*atanh(tan(e/2 + (f*x)/2)))/(c^2*f) - (2*a^2*tan(e/2 + (f*x)/2)^2 + (2*a^2)/3)/(c^2*f*tan(e/2 + (f*x)/2)^3)`

$$3.17 \quad \int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx$$

3.17.1	Optimal result	213
3.17.2	Mathematica [A] (verified)	213
3.17.3	Rubi [A] (verified)	214
3.17.4	Maple [A] (verified)	215
3.17.5	Fricas [B] (verification not implemented)	215
3.17.6	Sympy [F]	216
3.17.7	Maxima [B] (verification not implemented)	216
3.17.8	Giac [A] (verification not implemented)	217
3.17.9	Mupad [B] (verification not implemented)	217

3.17.1 Optimal result

Integrand size = 32, antiderivative size = 38

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx = -\frac{(a+a\sec(e+fx))^2 \tan(e+fx)}{5f(c-c\sec(e+fx))^3}$$

output `-1/5*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^3`

3.17.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx = \frac{a^2 \cot^5\left(\frac{1}{2}(e+fx)\right)}{5c^3 f}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^3,x]`

output `(a^2*Cot[(e + f*x)/2]^5)/(5*c^3*f)`

3.17.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c-c\sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^3} dx$$

↓ 4438

$$\frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{5f(c-c\sec(e+fx))^3}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^3,x]`

output `-1/5*((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^3)`

3.17.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^n, x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.17.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.61

method	result	size
derivativedivides	$\frac{a^2}{5f c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	23
default	$\frac{a^2}{5f c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	23
parallelrisc	$\frac{a^2}{5f c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	23
risc	$\frac{2ia^2(5e^{4i(fx+e)}+10e^{2i(fx+e)}+1)}{5f c^3(e^{i(fx+e)}-1)^5}$	50
norman	$\frac{\frac{a^2}{5cf} - \frac{2a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{5cf} + \frac{a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{5cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	87

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `1/5/f*a^2/c^3/tan(1/2*f*x+1/2*e)^5`

3.17.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(37) = 74$.

Time = 0.28 (sec) , antiderivative size = 83, normalized size of antiderivative = 2.18

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx$$

$$= \frac{a^2 \cos(fx+e)^3 + 3a^2 \cos(fx+e)^2 + 3a^2 \cos(fx+e) + a^2}{5(c^3 f \cos(fx+e)^2 - 2c^3 f \cos(fx+e) + c^3 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x, algorithm="fracas")`

output `1/5*(a^2*cos(f*x + e)^3 + 3*a^2*cos(f*x + e)^2 + 3*a^2*cos(f*x + e) + a^2)/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x + e))`

3.17.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx = \frac{a^2 \left(\int \frac{\sec(e+fx)}{\sec^3(e+fx)-3\sec^2(e+fx)+3\sec(e+fx)-1} dx + \int \frac{2\sec^2(e+fx)}{\sec^3(e+fx)-3\sec^2(e+fx)+3\sec(e+fx)-1} dx + \int \frac{\sec^3}{\sec^3(e+fx)-3\sec^2} \right)}{c^3}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**3,x)`

output `-a**2*(Integral(sec(e + f*x)/(sec(e + f*x)**3 - 3*sec(e + f*x)**2 + 3*sec(e + f*x) - 1), x) + Integral(2*sec(e + f*x)**2/(sec(e + f*x)**3 - 3*sec(e + f*x)**2 + 3*sec(e + f*x) - 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x)**3 - 3*sec(e + f*x)**2 + 3*sec(e + f*x) - 1), x))/c**3`

3.17.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(37) = 74$.

Time = 0.22 (sec) , antiderivative size = 189, normalized size of antiderivative = 4.97

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx = \frac{a^2 \left(\frac{10 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{15 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + 3 \right) (\cos(fx+e)+1)^5}{c^3 \sin(fx+e)^5} - \frac{a^2 \left(\frac{10 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{15 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - 3 \right) (\cos(fx+e)+1)^5}{c^3 \sin(fx+e)^5} - \frac{6a^2 \left(\frac{5 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} \right)}{c^3 \sin(fx+e)^5}$$

60 f

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/60*(a^2*(10*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 15*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 3)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5) - a^2*(10*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 15*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 3)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5) - 6*a^2*(5*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 1)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5))/f`

3.17.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx = \frac{a^2}{5c^3 f \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `1/5*a^2/(c^3*f*tan(1/2*f*x + 1/2*e)^5)`

3.17.9 Mupad [B] (verification not implemented)

Time = 13.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^3} dx = \frac{a^2 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{5c^3 f}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^3),x)`

output `(a^2*cot(e/2 + (f*x)/2)^5)/(5*c^3*f)`

3.18 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^4} dx$

3.18.1	Optimal result	218
3.18.2	Mathematica [A] (verified)	218
3.18.3	Rubi [A] (verified)	219
3.18.4	Maple [A] (verified)	220
3.18.5	Fricas [A] (verification not implemented)	221
3.18.6	Sympy [F]	221
3.18.7	Maxima [B] (verification not implemented)	222
3.18.8	Giac [A] (verification not implemented)	222
3.18.9	Mupad [B] (verification not implemented)	223

3.18.1 Optimal result

Integrand size = 32, antiderivative size = 80

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^4} dx = -\frac{(a+a \sec(e+fx))^2 \tan(e+fx)}{7f(c-c \sec(e+fx))^4} - \frac{(a+a \sec(e+fx))^2 \tan(e+fx)}{35cf(c-c \sec(e+fx))^3}$$

output `-1/7*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^4-1/35*(a+a*sec(f*x+e))^2*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^3`

3.18.2 Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.59

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^4} dx = \frac{a^2(-6+\sec(e+fx))(1+\sec(e+fx))^2 \tan(e+fx)}{35c^4 f(-1+\sec(e+fx))^4}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^4,x]`

output `(a^2*(-6 + Sec[e + f*x])*(1 + Sec[e + f*x])^2*Tan[e + f*x])/(35*c^4*f*(-1 + Sec[e + f*x])^4)`

3.18.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c-c\sec(e+fx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx \\
 & \quad \downarrow \text{4439} \\
 & \frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{(c-c\sec(e+fx))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4} \\
 & \quad \downarrow \text{4438} \\
 & -\frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{35cf(c-c\sec(e+fx))^3} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^4,x]`

output `-1/7*((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^4) - ((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(35*c*f*(c - c*Sec[e + f*x])^3)`

3.18.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.18.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.48

method	result	size
parallelrisch	$-\frac{a^2 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 \left(5 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 7\right)}{70c^4 f}$	38
derivativedivides	$\frac{a^2 \left(\frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} \right)}{2f c^4}$	39
default	$\frac{a^2 \left(\frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} \right)}{2f c^4}$	39
risch	$\frac{2ia^2 (35 e^{6i(fx+e)} - 35 e^{5i(fx+e)} + 140 e^{4i(fx+e)} - 70 e^{3i(fx+e)} + 91 e^{2i(fx+e)} - 7 e^{i(fx+e)} + 6)}{35f c^4 (e^{i(fx+e)} - 1)^7}$	94
norman	$-\frac{\frac{a^2}{14cf} + \frac{17a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{70cf} - \frac{19a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{70cf} + \frac{a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{10cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$	109

3.18. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^4} dx$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x,method=_RETURNVERBOSE)`

output $-1/70*a^2*\cot(1/2*f*x+1/2*e)^5*(5*\cot(1/2*f*x+1/2*e)^2-7)/c^4/f$

3.18.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.42

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^4} dx$$

$$= \frac{6a^2 \cos^4(fx+e) + 17a^2 \cos^3(fx+e) + 15a^2 \cos^2(fx+e) + 3a^2 \cos(fx+e) - a^2}{35(c^4 f \cos^3(fx+e) - 3c^4 f \cos^2(fx+e) + 3c^4 f \cos(fx+e) - c^4 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x, algorithm="fricas")`

output $1/35*(6*a^2*\cos(f*x + e)^4 + 17*a^2*\cos(f*x + e)^3 + 15*a^2*\cos(f*x + e)^2 + 3*a^2*\cos(f*x + e) - a^2)/((c^4*f*\cos(f*x + e)^3 - 3*c^4*f*\cos(f*x + e)^2 + 3*c^4*f*\cos(f*x + e) - c^4*f)*\sin(f*x + e))$

3.18.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^4} dx$$

$$= \frac{a^2 \left(\int \frac{\sec(e+fx)}{\sec^4(e+fx) - 4\sec^3(e+fx) + 6\sec^2(e+fx) - 4\sec(e+fx) + 1} dx + \int \frac{2\sec^2(e+fx)}{\sec^4(e+fx) - 4\sec^3(e+fx) + 6\sec^2(e+fx) - 4\sec(e+fx) + 1} dx \right)}{c^4}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**4,x)`

output `a**2*(Integral(sec(e + f*x)/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x) + Integral(2*sec(e + f*x)**2/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x))/c**4`

3.18. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^4} dx$

3.18.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 270 vs. 2(78) = 156.

Time = 0.22 (sec) , antiderivative size = 270, normalized size of antiderivative = 3.38

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^4} dx$$

$$= \frac{2a^2 \left(\frac{21 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{35 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{105 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - 15 \right) (\cos(fx+e)+1)^7}{c^4 \sin(fx+e)^7} + \frac{3a^2 \left(\frac{21 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{35 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{35 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - 5 \right) (\cos(fx+e)+1)^7}{c^4 \sin(fx+e)^7}$$

840 f

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output `1/840*(2*a^2*(21*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 35*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 105*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 15)*(cos(f*x + e) + 1)^7/(c^4*sin(f*x + e)^7) + 3*a^2*(21*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 35*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 35*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 5)*(cos(f*x + e) + 1)^7/(c^4*sin(f*x + e)^7) - a^2*(21*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 35*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 105*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 15)*(cos(f*x + e) + 1)^7/(c^4*sin(f*x + e)^7))/f`

3.18.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.51

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^4} dx = \frac{7a^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 5a^2}{70c^4 f \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^7}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output `1/70*(7*a^2*tan(1/2*f*x + 1/2*e)^2 - 5*a^2)/(c^4*f*tan(1/2*f*x + 1/2*e)^7)`

3.18.9 Mupad [B] (verification not implemented)

Time = 14.42 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.46

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^4} dx = -\frac{a^2 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^5 \left(5 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 7\right)}{70 c^4 f}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^4),x)`

output `-(a^2*cot(e/2 + (f*x)/2)^5*(5*cot(e/2 + (f*x)/2)^2 - 7))/(70*c^4*f)`

3.19 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^5} dx$

3.19.1 Optimal result 224
 3.19.2 Mathematica [A] (verified) 224
 3.19.3 Rubi [A] (verified) 225
 3.19.4 Maple [A] (verified) 227
 3.19.5 Fricas [A] (verification not implemented) 227
 3.19.6 Sympy [F] 228
 3.19.7 Maxima [B] (verification not implemented) 228
 3.19.8 Giac [A] (verification not implemented) 229
 3.19.9 Mupad [B] (verification not implemented) 229

3.19.1 Optimal result

Integrand size = 32, antiderivative size = 121

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^5} dx = -\frac{(a+a \sec(e+fx))^2 \tan(e+fx)}{9f(c-c \sec(e+fx))^5} - \frac{2(a+a \sec(e+fx))^2 \tan(e+fx)}{63cf(c-c \sec(e+fx))^4} - \frac{2(a+a \sec(e+fx))^2 \tan(e+fx)}{315c^2f(c-c \sec(e+fx))^3}$$

output `-1/9*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^5-2/63*(a+a*sec(f*x+e))^2*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^4-2/315*(a+a*sec(f*x+e))^2*tan(f*x+e)/c^2/f/(c-c*sec(f*x+e))^3`

3.19.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.49

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^5} dx = \frac{a^2(1+\sec(e+fx))^2(47-14\sec(e+fx)+2\sec^2(e+fx))\tan(e+fx)}{315c^5f(-1+\sec(e+fx))^5}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^5,x]`

output $(a^2(1 + \sec[e + fx])^2(47 - 14\sec[e + fx] + 2\sec[e + fx]^2)\tan[e + fx]) / (315c^5f(-1 + \sec[e + fx])^5)$

3.19.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c-c\sec(e+fx))^5} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^5} dx$$

↓ 4439

$$\frac{2 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{(c-c\sec(e+fx))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5}$$

↓ 3042

$$\frac{2 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5}$$

↓ 4439

$$\frac{2 \left(\frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{(c-c\sec(e+fx))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4} \right)}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5}$$

↓ 3042

$$\frac{2 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^3} dx}{7c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4} \right)}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5}$$

↓ 4438

3.19. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^5} dx$

$$2 \left(\frac{-\tan(e+fx)(a \sec(e+fx)+a)^2}{35cf(c-c \sec(e+fx))^3} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^2}{7f(c-c \sec(e+fx))^4} \right) - \frac{\tan(e+fx)(a \sec(e+fx)+a)^2}{9f(c-c \sec(e+fx))^5}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^5,x]`

output `-1/9*((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^5) + 2*(-1/7*((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^4) - ((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(35*c*f*(c - c*Sec[e + f*x])^3))/9*c)`

3.19.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.19.4 Maple [A] (verified)

Time = 1.00 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.42

method	result
parallelrisch	$\frac{a^2 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 \left(35 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^4 - 90 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 + 63\right)}{1260c^5 f}$
derivativedivides	$\frac{a^2 \left(\frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{2}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}\right)}{4f c^5}$
default	$\frac{a^2 \left(\frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{2}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}\right)}{4f c^5}$
risch	$\frac{2ia^2 (315 e^{8i(fx+e)} - 630 e^{7i(fx+e)} + 2310 e^{6i(fx+e)} - 2520 e^{5i(fx+e)} + 3402 e^{4i(fx+e)} - 1638 e^{3i(fx+e)} + 1062 e^{2i(fx+e)} - 315)}{315 f c^5 (e^{i(fx+e)} - 1)^9}$
norman	$\frac{\frac{a^2}{36cf} - \frac{8a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{63cf} + \frac{139a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{630cf} - \frac{6a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{35cf} + \frac{a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{20cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x,method=_RETURNVERBOSE)`

output `1/1260*a^2*cot(1/2*f*x+1/2*e)^5*(35*cot(1/2*f*x+1/2*e)^4-90*cot(1/2*f*x+1/2*e)^2+63)/c^5/f`

3.19.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.16

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^5} dx$$

$$= \frac{47 a^2 \cos (fx+e)^5+127 a^2 \cos (fx+e)^4+101 a^2 \cos (fx+e)^3+11 a^2 \cos (fx+e)^2-8 a^2 \cos (fx+e)}{315\left(c^5 f \cos (fx+e)^4-4 c^5 f \cos (fx+e)^3+6 c^5 f \cos (fx+e)^2-4 c^5 f \cos (fx+e)+c^5 f\right) \sin (fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x, algorithm="fracas")`

output $1/315*(47*a^2*\cos(f*x + e)^5 + 127*a^2*\cos(f*x + e)^4 + 101*a^2*\cos(f*x + e)^3 + 11*a^2*\cos(f*x + e)^2 - 8*a^2*\cos(f*x + e) + 2*a^2)/((c^5*f*\cos(f*x + e)^4 - 4*c^5*f*\cos(f*x + e)^3 + 6*c^5*f*\cos(f*x + e)^2 - 4*c^5*f*\cos(f*x + e) + c^5*f)*\sin(f*x + e))$

3.19.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^5} dx =$$

$$\frac{a^2 \left(\int \frac{\sec(e + fx)}{\sec^5(e + fx) - 5 \sec^4(e + fx) + 10 \sec^3(e + fx) - 10 \sec^2(e + fx) + 5 \sec(e + fx) - 1} dx + \int \frac{2 \sec^2(e + fx)}{\sec^5(e + fx) - 5 \sec^4(e + fx) + 10 \sec^3(e + fx) - 10 \sec^2(e + fx) + 5 \sec(e + fx) - 1} dx \right)}{c^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**5,x)`

output `-a**2*(Integral(sec(e + f*x)/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec(e + f*x)**3 - 10*sec(e + f*x)**2 + 5*sec(e + f*x) - 1), x) + Integral(2*sec(e + f*x)**2/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec(e + f*x)**3 - 10*sec(e + f*x)**2 + 5*sec(e + f*x) - 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec(e + f*x)**3 - 10*sec(e + f*x)**2 + 5*sec(e + f*x) - 1), x))/c**5`

3.19.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 269 vs. $2(118) = 236$.

Time = 0.22 (sec) , antiderivative size = 269, normalized size of antiderivative = 2.22

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^5} dx =$$

$$\frac{a^2 \left(\frac{180 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{378 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{420 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{315 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} - 35 \right) (\cos(fx+e)+1)^9}{c^5 \sin^9(fx+e)} + \frac{10 a^2 \left(\frac{18 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{42 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{6}{(\cos(fx+e)+1)^6} \right)}{c^5 \sin^6(fx+e)}$$

5040 f

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x, algorithm="maxima")`

3.19. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^5} dx$

output
$$\frac{-1/5040*(a^2*(180*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 378*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 + 420*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6 - 315*\sin(f*x + e)^8/(\cos(f*x + e) + 1)^8 - 35)*(\cos(f*x + e) + 1)^9/(c^5*\sin(f*x + e)^9) + 10*a^2*(18*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 42*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6 + 63*\sin(f*x + e)^8/(\cos(f*x + e) + 1)^8 - 7)*(\cos(f*x + e) + 1)^9/(c^5*\sin(f*x + e)^9) + 7*a^2*(18*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 - 45*\sin(f*x + e)^8/(\cos(f*x + e) + 1)^8 - 5)*(\cos(f*x + e) + 1)^9/(c^5*\sin(f*x + e)^9))/f}$$

3.19.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.47

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^5} dx$$

$$= \frac{63 a^2 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 - 90 a^2 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 + 35 a^2}{1260 c^5 f \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^9}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x, algorithm="giac")`

output
$$1/1260*(63*a^2*\tan(1/2*f*x + 1/2*e)^4 - 90*a^2*\tan(1/2*f*x + 1/2*e)^2 + 35*a^2)/(c^5*f*\tan(1/2*f*x + 1/2*e)^9)$$

3.19.9 Mupad [B] (verification not implemented)

Time = 14.17 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.55

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^5} dx$$

$$= \frac{a^2 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{20 c^5 f} - \frac{a^2 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{14 c^5 f} + \frac{a^2 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{36 c^5 f}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^5),x)`

output
$$(a^2*\cot(e/2 + (f*x)/2)^5)/(20*c^5*f) - (a^2*\cot(e/2 + (f*x)/2)^7)/(14*c^5*f) + (a^2*\cot(e/2 + (f*x)/2)^9)/(36*c^5*f)$$

3.20
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^6} dx$$

3.20.1 Optimal result 230
 3.20.2 Mathematica [A] (verified) 231
 3.20.3 Rubi [A] (verified) 231
 3.20.4 Maple [A] (verified) 233
 3.20.5 Fricas [A] (verification not implemented) 234
 3.20.6 Sympy [F] 235
 3.20.7 Maxima [B] (verification not implemented) 235
 3.20.8 Giac [A] (verification not implemented) 236
 3.20.9 Mupad [B] (verification not implemented) 236

3.20.1 Optimal result

Integrand size = 32, antiderivative size = 163

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^6} dx = -\frac{(a+a \sec(e+fx))^2 \tan(e+fx)}{11f(c-c \sec(e+fx))^6} - \frac{(a+a \sec(e+fx))^2 \tan(e+fx)}{33cf(c-c \sec(e+fx))^5} - \frac{2(a+a \sec(e+fx))^2 \tan(e+fx)}{231c^2f(c-c \sec(e+fx))^4} - \frac{2(a+a \sec(e+fx))^2 \tan(e+fx)}{1155f(c^2-c^2 \sec(e+fx))^3}$$

```
output -1/11*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^6-1/33*(a+a*sec(f*x+e))^2*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^5-2/231*(a+a*sec(f*x+e))^2*tan(f*x+e)/c^2/f/(c-c*sec(f*x+e))^4-2/1155*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c^2-c^2*sec(f*x+e))^3
```

3.20.2 Mathematica [A] (verified)

Time = 1.25 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.42

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$$

$$= \frac{a^2(1+\sec(e+fx))^2(-152+61\sec(e+fx)-16\sec^2(e+fx)+2\sec^3(e+fx))\tan(e+fx)}{1155c^6f(-1+\sec(e+fx))^6}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^6,x]`

output `(a^2*(1 + Sec[e + f*x])^2*(-152 + 61*Sec[e + f*x] - 16*Sec[e + f*x]^2 + 2*Sec[e + f*x]^3)*Tan[e + f*x])/(1155*c^6*f*(-1 + Sec[e + f*x])^6)`

3.20.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4439, 3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c-c\sec(e+fx))^6} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^6} dx$$

↓ 4439

$$\frac{3 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{(c-c\sec(e+fx))^5} dx}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{11f(c-c\sec(e+fx))^6}$$

↓ 3042

$$\frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^5} dx}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{11f(c-c\sec(e+fx))^6}$$

↓ 4439

3.20. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$

$$\begin{aligned}
& \frac{3 \left(\frac{2 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2 dx}{(c-c\sec(e+fx))^4} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5}}{9c} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{11f(c-c\sec(e+fx))^6} \\
& \quad \downarrow 3042 \\
& \frac{3 \left(\frac{2 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5}}{9c} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{11f(c-c\sec(e+fx))^6} \\
& \quad \downarrow 4439 \\
& \frac{3 \left(\frac{2 \left(\frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2 dx}{(c-c\sec(e+fx))^3} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4} \right)}{9c} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5} \\
& \quad \downarrow 3042 \\
& \frac{3 \left(\frac{2 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2 dx}{(c-c\csc(e+fx+\frac{\pi}{2}))^3} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4} \right)}{9c} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5} \\
& \quad \downarrow 4438 \\
& \frac{3 \left(\frac{2 \left(-\frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{35cf(c-c\sec(e+fx))^3} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{7f(c-c\sec(e+fx))^4} \right)}{9c} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^2}{9f(c-c\sec(e+fx))^5} \\
& \quad \downarrow \\
& \frac{11c}{11f(c-c\sec(e+fx))^6} \tan(e+fx)(a\sec(e+fx)+a)^2
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^6,x]`

$$3.20. \quad \int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$$

```
output -1/11*((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^6) + (
3*(-1/9*((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^5) +
(2*(-1/7*((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^4)
- ((a + a*Sec[e + f*x])^2*Tan[e + f*x])/(35*c*f*(c - c*Sec[e + f*x])^3))
/(9*c)))/(11*c)
```

3.20.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4438 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]
*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; Fre
eQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] &
& EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]
```

```
rule 4439 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]
*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp
[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*
(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ
[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0
] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])
```

3.20.4 Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.39

$$3.20. \int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$$

method	result
parallelrisc	$\frac{a^2 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 \left(105 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^6 - 385 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^4 + 495 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 231\right)}{9240c^6 f}$
derivativdivides	$\frac{a^2 \left(\frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{3}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}\right)}{8f c^6}$
default	$\frac{a^2 \left(\frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{3}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}\right)}{8f c^6}$
risc	$\frac{2ia^2 (1155 e^{10i(fx+e)} - 3465 e^{9i(fx+e)} + 13860 e^{8i(fx+e)} - 23100 e^{7i(fx+e)} + 37422 e^{6i(fx+e)} - 32802 e^{5i(fx+e)} + 27060 e^{4i(fx+e)} - 15552 e^{3i(fx+e)} + 7560 e^{2i(fx+e)} - 1260 e^{i(fx+e)} + 126)}{1155 f c^6 (e^{i(fx+e)} - 1)^{11}}$
norman	$\frac{-\frac{a^2}{88cf} + \frac{17a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{264cf} - \frac{137a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{924cf} + \frac{73a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{420cf} - \frac{29a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{280cf} + \frac{a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{10}}{40cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^6,x,method=_RETURNVERBOSE)
```

```
output -1/9240*a^2*cot(1/2*f*x+1/2*e)^5*(105*cot(1/2*f*x+1/2*e)^6-385*cot(1/2*f*x+1/2*e)^4+495*cot(1/2*f*x+1/2*e)^2-231)/c^6/f
```

3.20.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.03

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$$

$$= \frac{152 a^2 \cos(fx+e)^6 + 395 a^2 \cos(fx+e)^5 + 289 a^2 \cos(fx+e)^4 + 15 a^2 \cos(fx+e)^3 - 19 a^2 \cos(fx+e)^2 + 10 a^2 \cos(fx+e) - 2 a^2}{1155 (c^6 f \cos(fx+e)^5 - 5 c^6 f \cos(fx+e)^4 + 10 c^6 f \cos(fx+e)^3 - 10 c^6 f \cos(fx+e)^2 + 5 c^6 f \cos(fx+e) - c^6 f) \sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^6,x, algorithm="fricas")
```

```
output 1/1155*(152*a^2*cos(f*x + e)^6 + 395*a^2*cos(f*x + e)^5 + 289*a^2*cos(f*x + e)^4 + 15*a^2*cos(f*x + e)^3 - 19*a^2*cos(f*x + e)^2 + 10*a^2*cos(f*x + e) - 2*a^2)/((c^6*f*cos(f*x + e)^5 - 5*c^6*f*cos(f*x + e)^4 + 10*c^6*f*cos(f*x + e)^3 - 10*c^6*f*cos(f*x + e)^2 + 5*c^6*f*cos(f*x + e) - c^6*f)*sin(f*x + e))
```

3.20. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$

3.20.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$$

$$= \frac{a^2 \left(\int \frac{\sec(e+fx)}{\sec^6(e+fx)-6\sec^5(e+fx)+15\sec^4(e+fx)-20\sec^3(e+fx)+15\sec^2(e+fx)-6\sec(e+fx)+1} dx + \int \frac{1}{\sec^6(e+fx)-6\sec^5(e+fx)+1} dx \right)}{c^6}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**6,x)`

output `a**2*(Integral(sec(e + f*x)/(sec(e + f*x)**6 - 6*sec(e + f*x)**5 + 15*sec(e + f*x)**4 - 20*sec(e + f*x)**3 + 15*sec(e + f*x)**2 - 6*sec(e + f*x) + 1), x) + Integral(2*sec(e + f*x)**2/(sec(e + f*x)**6 - 6*sec(e + f*x)**5 + 15*sec(e + f*x)**4 - 20*sec(e + f*x)**3 + 15*sec(e + f*x)**2 - 6*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x)**6 - 6*sec(e + f*x)**5 + 15*sec(e + f*x)**4 - 20*sec(e + f*x)**3 + 15*sec(e + f*x)**2 - 6*sec(e + f*x) + 1), x))/c**6`

3.20.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 389 vs. $2(159) = 318$.

Time = 0.22 (sec) , antiderivative size = 389, normalized size of antiderivative = 2.39

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^6} dx$$

$$= \frac{a^2 \left(\frac{385 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} + \frac{990 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} - \frac{1386 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{1155 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} + \frac{3465 \sin^{10}(fx+e)}{(\cos(fx+e)+1)^{10}} - 315 \right) (\cos(fx+e)+1)^{11}}{c^6 \sin^2(fx+e)} + \frac{6 a^2 \left(\frac{385 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{330 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{1386 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{1155 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} + \frac{3465 \sin^{10}(fx+e)}{(\cos(fx+e)+1)^{10}} - 315 \right) (\cos(fx+e)+1)^{11}}{c^6 \sin^2(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^6,x, algorithm="maxima")`

output $\frac{1}{110880} \cdot (a^2 \cdot (385 \sin(fx + e)^2 / (\cos(fx + e) + 1)^2 + 990 \sin(fx + e)^4 / (\cos(fx + e) + 1)^4 - 1386 \sin(fx + e)^6 / (\cos(fx + e) + 1)^6 - 1155 \sin(fx + e)^8 / (\cos(fx + e) + 1)^8 + 3465 \sin(fx + e)^{10} / (\cos(fx + e) + 1)^{10} - 315) \cdot (\cos(fx + e) + 1)^{11} / (c^6 \sin(fx + e)^{11}) + 6 \cdot a^2 \cdot (385 \sin(fx + e)^2 / (\cos(fx + e) + 1)^2 - 330 \sin(fx + e)^4 / (\cos(fx + e) + 1)^4 - 462 \sin(fx + e)^6 / (\cos(fx + e) + 1)^6 + 1155 \sin(fx + e)^8 / (\cos(fx + e) + 1)^8 - 1155 \sin(fx + e)^{10} / (\cos(fx + e) + 1)^{10} - 105) \cdot (\cos(fx + e) + 1)^{11} / (c^6 \sin(fx + e)^{11}) + 5 \cdot a^2 \cdot (385 \sin(fx + e)^2 / (\cos(fx + e) + 1)^2 - 990 \sin(fx + e)^4 / (\cos(fx + e) + 1)^4 + 1386 \sin(fx + e)^6 / (\cos(fx + e) + 1)^6 - 1155 \sin(fx + e)^8 / (\cos(fx + e) + 1)^8 + 693 \sin(fx + e)^{10} / (\cos(fx + e) + 1)^{10} - 63) \cdot (\cos(fx + e) + 1)^{11} / (c^6 \sin(fx + e)^{11})) / f$

3.20.8 Giac [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.45

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^6} dx$$

$$= \frac{231 a^2 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 - 495 a^2 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 + 385 a^2 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 105 a^2}{9240 c^6 f \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^6,x, algorithm="giac")`

output $\frac{1}{9240} \cdot (231 \cdot a^2 \cdot \tan(1/2 \cdot fx + 1/2 \cdot e)^6 - 495 \cdot a^2 \cdot \tan(1/2 \cdot fx + 1/2 \cdot e)^4 + 385 \cdot a^2 \cdot \tan(1/2 \cdot fx + 1/2 \cdot e)^2 - 105 \cdot a^2) / (c^6 \cdot f \cdot \tan(1/2 \cdot fx + 1/2 \cdot e)^{11})$

3.20.9 Mupad [B] (verification not implemented)

Time = 14.26 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^6} dx =$$

$$\frac{a^2 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^5 \left(105 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 385 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^4 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 495 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^2 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{9240 c^6 f \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}$$

3.20. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^6} dx$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^6),x)`

output
$$-(a^2 \cos(e/2 + (f*x)/2)^5 (105 \cos(e/2 + (f*x)/2)^6 - 231 \sin(e/2 + (f*x)/2)^6 + 495 \cos(e/2 + (f*x)/2)^2 \sin(e/2 + (f*x)/2)^4 - 385 \cos(e/2 + (f*x)/2)^4 \sin(e/2 + (f*x)/2)^2) / (9240 * c^6 * f * \sin(e/2 + (f*x)/2)^{11})$$

3.21 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$

3.21.1	Optimal result	238
3.21.2	Mathematica [A] (verified)	239
3.21.3	Rubi [A] (verified)	239
3.21.4	Maple [C] (verified)	240
3.21.5	Fricas [A] (verification not implemented)	241
3.21.6	Sympy [F]	242
3.21.7	Maxima [B] (verification not implemented)	242
3.21.8	Giac [A] (verification not implemented)	243
3.21.9	Mupad [B] (verification not implemented)	244

3.21.1 Optimal result

Integrand size = 32, antiderivative size = 227

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$$

$$= \frac{55a^3c^6 \operatorname{arctanh}(\sin(e + fx))}{128f} - \frac{25a^3c^6 \sec(e + fx) \tan(e + fx)}{128f}$$

$$- \frac{15a^3c^6 \sec^3(e + fx) \tan(e + fx)}{64f} + \frac{5a^3c^6 \sec(e + fx) \tan^3(e + fx)}{24f}$$

$$+ \frac{5a^3c^6 \sec^3(e + fx) \tan^3(e + fx)}{16f} - \frac{a^3c^6 \sec(e + fx) \tan^5(e + fx)}{6f}$$

$$- \frac{3a^3c^6 \sec^3(e + fx) \tan^5(e + fx)}{8f} + \frac{4a^3c^6 \tan^7(e + fx)}{7f} + \frac{a^3c^6 \tan^9(e + fx)}{9f}$$

```
output 55/128*a^3*c^6*arctanh(sin(f*x+e))/f-25/128*a^3*c^6*sec(f*x+e)*tan(f*x+e)/
f-15/64*a^3*c^6*sec(f*x+e)^3*tan(f*x+e)/f+5/24*a^3*c^6*sec(f*x+e)*tan(f*x+
e)^3/f+5/16*a^3*c^6*sec(f*x+e)^3*tan(f*x+e)^3/f-1/6*a^3*c^6*sec(f*x+e)*tan
(f*x+e)^5/f-3/8*a^3*c^6*sec(f*x+e)^3*tan(f*x+e)^5/f+4/7*a^3*c^6*tan(f*x+e)
^7/f+1/9*a^3*c^6*tan(f*x+e)^9/f
```

3.21.2 Mathematica [A] (verified)

Time = 3.24 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.54

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$$

$$= \frac{a^3 c^6 (443520 \operatorname{arctanh}(\sin(e + fx)) - \sec^9(e + fx)(-88704 \sin(e + fx) + 88074 \sin(2(e + fx)) + 37632 \sin(3(e + fx)) - 2142 \sin(4(e + fx)) + 2304 \sin(5(e + fx)) + 39858 \sin(6(e + fx)) - 7488 \sin(7(e + fx)) + 4599 \sin(8(e + fx)) + 1856 \sin(9(e + fx))))}{1032192 f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^6,x]`

output `(a^3*c^6*(443520*ArcTanh[Sin[e + f*x]] - Sec[e + f*x]^9*(-88704*Sin[e + f*x] + 88074*Sin[2*(e + f*x)] + 37632*Sin[3*(e + f*x)] - 2142*Sin[4*(e + f*x)]) + 2304*Sin[5*(e + f*x)] + 39858*Sin[6*(e + f*x)] - 7488*Sin[7*(e + f*x)] + 4599*Sin[8*(e + f*x)] + 1856*Sin[9*(e + f*x)]))/(1032192*f)`

3.21.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^6 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^6 dx$$

$$\downarrow \text{4446}$$

$$-a^3 c^3 \int \left(-c^3 \sec^4(e + fx) \tan^6(e + fx) + 3c^3 \sec^3(e + fx) \tan^6(e + fx) - 3c^3 \sec^2(e + fx) \tan^6(e + fx) + c^3 \sec(e + fx) \tan^6(e + fx)\right) dx$$

$$\downarrow \text{2009}$$

$$-a^3 c^3 \left(-\frac{55c^3 \operatorname{arctanh}(\sin(e + fx))}{128f} - \frac{c^3 \tan^9(e + fx)}{9f} - \frac{4c^3 \tan^7(e + fx)}{7f} + \frac{3c^3 \tan^5(e + fx) \sec^3(e + fx)}{8f} - \frac{5c^3 \tan^3(e + fx) \sec^3(e + fx)}{8f} \right)$$

3.21. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^6,x]`

output `-(a^3*c^3*((-55*c^3*ArcTanh[Sin[e + f*x]])/(128*f) + (25*c^3*Sec[e + f*x]*Tan[e + f*x])/(128*f) + (15*c^3*Sec[e + f*x]^3*Tan[e + f*x])/(64*f) - (5*c^3*Sec[e + f*x]*Tan[e + f*x]^3)/(24*f) - (5*c^3*Sec[e + f*x]^3*Tan[e + f*x]^3)/(16*f) + (c^3*Sec[e + f*x]*Tan[e + f*x]^5)/(6*f) + (3*c^3*Sec[e + f*x]^3*Tan[e + f*x]^5)/(8*f) - (4*c^3*Tan[e + f*x]^7)/(7*f) - (c^3*Tan[e + f*x]^9)/(9*f)))`

3.21.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.21.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 9.32 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.11

3.21. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$

method	result
risch	$ia^3c^6(4599e^{17i(fx+e)} - 24192e^{16i(fx+e)} + 39858e^{15i(fx+e)} - 64512e^{14i(fx+e)} - 2142e^{13i(fx+e)} - 118272e^{12i(fx+e)} + 88074e^{11i(fx+e)} - 322560e^{10i(fx+e)} - 145152e^{9i(fx+e)} - 88074e^{8i(fx+e)} - 193536e^{7i(fx+e)} + 2142e^{6i(fx+e)} + 24142e^{5i(fx+e)} - 69120e^{4i(fx+e)} - 39858e^{3i(fx+e)} - 9216e^{2i(fx+e)} - 4599e^{i(fx+e)} - 3712)/f/(1 + \exp(2i(fx+e)))^9 - 55/128*a^3*c^6/f*\ln(\exp(i(fx+e)) + I) + 55/128*a^3*c^6/f*\ln(\exp(i(fx+e)) - I)$
parallelrisch	$55a^3c^6((\cos(9fx+9e)+9\cos(7fx+7e)+36\cos(5fx+5e)+84\cos(3fx+3e)+126\cos(fx+e))\ln(\tan(\frac{fx}{2}+\frac{e}{2})-1))+(-$
derivativedivides	$-a^3c^6\left(-\frac{128}{315}-\frac{\sec(fx+e)^8}{9}-\frac{8\sec(fx+e)^6}{63}-\frac{16\sec(fx+e)^4}{105}-\frac{64\sec(fx+e)^2}{315}\right)\tan(fx+e)-3a^3c^6\tan(fx+e)+a^3c^6\ln(\sec(fx+e)+\tan(fx+e))$
default	$-a^3c^6\left(-\frac{128}{315}-\frac{\sec(fx+e)^8}{9}-\frac{8\sec(fx+e)^6}{63}-\frac{16\sec(fx+e)^4}{105}-\frac{64\sec(fx+e)^2}{315}\right)\tan(fx+e)-3a^3c^6\tan(fx+e)+a^3c^6\ln(\sec(fx+e)+\tan(fx+e))$
parts	$\frac{a^3c^6\ln(\sec(fx+e)+\tan(fx+e))}{f} - \frac{a^3c^6\left(-\frac{128}{315}-\frac{\sec(fx+e)^8}{9}-\frac{8\sec(fx+e)^6}{63}-\frac{16\sec(fx+e)^4}{105}-\frac{64\sec(fx+e)^2}{315}\right)\tan(fx+e)}{f}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^6,x,method=_RETURNVERBOSE)`

output `1/4032*I*a^3*c^6*(4599*exp(17*I*(f*x+e))-24192*exp(16*I*(f*x+e))+39858*exp(15*I*(f*x+e))-64512*exp(14*I*(f*x+e))-2142*exp(13*I*(f*x+e))-118272*exp(12*I*(f*x+e))+88074*exp(11*I*(f*x+e))-322560*exp(10*I*(f*x+e))-145152*exp(9*I*(f*x+e))-88074*exp(8*I*(f*x+e))-193536*exp(7*I*(f*x+e))+2142*exp(6*I*(f*x+e))+24142*exp(5*I*(f*x+e))-69120*exp(4*I*(f*x+e))-39858*exp(3*I*(f*x+e))-9216*exp(2*I*(f*x+e))-4599*exp(I*(f*x+e))-3712)/f/(1+exp(2*I*(f*x+e)))^9-55/128*a^3*c^6/f*ln(exp(I*(f*x+e))-I)+55/128*a^3*c^6/f*ln(exp(I*(f*x+e))+I)`

3.21.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.92

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$$

$$= \frac{3465 a^3 c^6 \cos(fx + e)^9 \log(\sin(fx + e) + 1) - 3465 a^3 c^6 \cos(fx + e)^9 \log(-\sin(fx + e) + 1) - 2(3712$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^6,x, algorithm="fracas")`

3.21. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$

```
output 1/16128*(3465*a^3*c^6*cos(f*x + e)^9*log(sin(f*x + e) + 1) - 3465*a^3*c^6*
cos(f*x + e)^9*log(-sin(f*x + e) + 1) - 2*(3712*a^3*c^6*cos(f*x + e)^8 + 4
599*a^3*c^6*cos(f*x + e)^7 - 10240*a^3*c^6*cos(f*x + e)^6 + 3066*a^3*c^6*c
os(f*x + e)^5 + 8448*a^3*c^6*cos(f*x + e)^4 - 7224*a^3*c^6*cos(f*x + e)^3
- 1024*a^3*c^6*cos(f*x + e)^2 + 3024*a^3*c^6*cos(f*x + e) - 896*a^3*c^6)*s
in(f*x + e))/(f*cos(f*x + e)^9)
```

3.21.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx \\ &= a^3 c^6 \left(\int \sec(e + fx) dx + \int (-3 \sec^2(e + fx)) dx + \int 8 \sec^4(e + fx) dx \right. \\ & \quad + \int (-6 \sec^5(e + fx)) dx + \int (-6 \sec^6(e + fx)) dx + \int 8 \sec^7(e + fx) dx \\ & \quad \left. + \int (-3 \sec^9(e + fx)) dx + \int \sec^{10}(e + fx) dx \right) \end{aligned}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**6,x)
```

```
output a**3*c**6*(Integral(sec(e + f*x), x) + Integral(-3*sec(e + f*x)**2, x) + I
ntegral(8*sec(e + f*x)**4, x) + Integral(-6*sec(e + f*x)**5, x) + Integral
(-6*sec(e + f*x)**6, x) + Integral(8*sec(e + f*x)**7, x) + Integral(-3*sec
(e + f*x)**9, x) + Integral(sec(e + f*x)**10, x))
```

3.21.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 443 vs. 2(209) = 418.

Time = 0.23 (sec) , antiderivative size = 443, normalized size of antiderivative = 1.95

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx \\ &= \frac{256 (35 \tan(fx + e)^9 + 180 \tan(fx + e)^7 + 378 \tan(fx + e)^5 + 420 \tan(fx + e)^3 + 315 \tan(fx + e))}{\dots} \end{aligned}$$

3.21. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^6,x, algorithm="maxima")`

output `1/80640*(256*(35*tan(f*x + e)^9 + 180*tan(f*x + e)^7 + 378*tan(f*x + e)^5 + 420*tan(f*x + e)^3 + 315*tan(f*x + e))*a^3*c^6 - 32256*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^3*c^6 + 215040*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c^6 + 315*a^3*c^6*(2*(105*sin(f*x + e)^7 - 385*sin(f*x + e)^5 + 511*sin(f*x + e)^3 - 279*sin(f*x + e)))/(sin(f*x + e)^8 - 4*sin(f*x + e)^6 + 6*sin(f*x + e)^4 - 4*sin(f*x + e)^2 + 1) - 105*log(sin(f*x + e) + 1) + 105*log(sin(f*x + e) - 1)) - 6720*a^3*c^6*(2*(15*sin(f*x + e)^5 - 40*sin(f*x + e)^3 + 33*sin(f*x + e)))/(sin(f*x + e)^6 - 3*sin(f*x + e)^4 + 3*sin(f*x + e)^2 - 1) - 15*log(sin(f*x + e) + 1) + 15*log(sin(f*x + e) - 1)) + 30240*a^3*c^6*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e)))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) + 80640*a^3*c^6*log(sec(f*x + e) + tan(f*x + e)) - 241920*a^3*c^6*tan(f*x + e))/f`

3.21.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.04

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$$

$$= \frac{3465 a^3 c^6 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + 1 \right| \right) - 3465 a^3 c^6 \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(3465 a^3 c^6 \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)^{17} - 3}{\dots}}{\dots}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^6,x, algorithm="giac")`

output `1/8064*(3465*a^3*c^6*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 3465*a^3*c^6*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(3465*a^3*c^6*tan(1/2*f*x + 1/2*e)^17 - 30030*a^3*c^6*tan(1/2*f*x + 1/2*e)^15 + 115038*a^3*c^6*tan(1/2*f*x + 1/2*e)^13 + 334602*a^3*c^6*tan(1/2*f*x + 1/2*e)^11 - 360448*a^3*c^6*tan(1/2*f*x + 1/2*e)^9 + 255222*a^3*c^6*tan(1/2*f*x + 1/2*e)^7 - 115038*a^3*c^6*tan(1/2*f*x + 1/2*e)^5 + 30030*a^3*c^6*tan(1/2*f*x + 1/2*e)^3 - 3465*a^3*c^6*tan(1/2*f*x + 1/2*e)))/(tan(1/2*f*x + 1/2*e)^2 - 1)^9)/f`

3.21. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx$

3.21.9 Mupad [B] (verification not implemented)

Time = 17.26 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.39

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^6 dx = \frac{55 a^3 c^6 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{64 f}$$

$$- \frac{55 a^3 c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{17}}{64} - \frac{715 a^3 c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{15}}{96} + \frac{913 a^3 c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{13}}{32} + \frac{18589 a^3 c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}{224} - \frac{5632 a^3 c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{63}$$

$$f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{18} - 9 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{16} + 36 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{14} - 84 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{12} + 126 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - \dots \right)$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^6)/cos(e + f*x),x)`output `(55*a^3*c^6*atanh(tan(e/2 + (f*x)/2)))/(64*f) - ((715*a^3*c^6*tan(e/2 + (f*x)/2)^3)/96 - (913*a^3*c^6*tan(e/2 + (f*x)/2)^5)/32 + (14179*a^3*c^6*tan(e/2 + (f*x)/2)^7)/224 - (5632*a^3*c^6*tan(e/2 + (f*x)/2)^9)/63 + (18589*a^3*c^6*tan(e/2 + (f*x)/2)^11)/224 + (913*a^3*c^6*tan(e/2 + (f*x)/2)^13)/32 - (715*a^3*c^6*tan(e/2 + (f*x)/2)^15)/96 + (55*a^3*c^6*tan(e/2 + (f*x)/2)^17)/64 - (55*a^3*c^6*tan(e/2 + (f*x)/2))/64/(f*(9*tan(e/2 + (f*x)/2)^2 - 36*tan(e/2 + (f*x)/2)^4 + 84*tan(e/2 + (f*x)/2)^6 - 126*tan(e/2 + (f*x)/2)^8 + 126*tan(e/2 + (f*x)/2)^10 - 84*tan(e/2 + (f*x)/2)^12 + 36*tan(e/2 + (f*x)/2)^14 - 9*tan(e/2 + (f*x)/2)^16 + tan(e/2 + (f*x)/2)^18 - 1))`

3.22 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$

3.22.1	Optimal result	245
3.22.2	Mathematica [A] (verified)	246
3.22.3	Rubi [A] (verified)	246
3.22.4	Maple [A] (verified)	247
3.22.5	Fricas [A] (verification not implemented)	248
3.22.6	Sympy [F]	249
3.22.7	Maxima [B] (verification not implemented)	249
3.22.8	Giac [A] (verification not implemented)	250
3.22.9	Mupad [B] (verification not implemented)	251

3.22.1 Optimal result

Integrand size = 32, antiderivative size = 206

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$$

$$= \frac{45a^3c^5 \operatorname{arctanh}(\sin(e + fx))}{128f} - \frac{35a^3c^5 \sec(e + fx) \tan(e + fx)}{128f}$$

$$- \frac{5a^3c^5 \sec^3(e + fx) \tan(e + fx)}{64f} + \frac{5a^3c^5 \sec(e + fx) \tan^3(e + fx)}{24f}$$

$$+ \frac{5a^3c^5 \sec^3(e + fx) \tan^3(e + fx)}{48f} - \frac{a^3c^5 \sec(e + fx) \tan^5(e + fx)}{6f}$$

$$- \frac{a^3c^5 \sec^3(e + fx) \tan^5(e + fx)}{8f} + \frac{2a^3c^5 \tan^7(e + fx)}{7f}$$

```
output 45/128*a^3*c^5*arctanh(sin(f*x+e))/f-35/128*a^3*c^5*sec(f*x+e)*tan(f*x+e)/
f-5/64*a^3*c^5*sec(f*x+e)^3*tan(f*x+e)/f+5/24*a^3*c^5*sec(f*x+e)*tan(f*x+e)
)^3/f+5/48*a^3*c^5*sec(f*x+e)^3*tan(f*x+e)^3/f-1/6*a^3*c^5*sec(f*x+e)*tan(
f*x+e)^5/f-1/8*a^3*c^5*sec(f*x+e)^3*tan(f*x+e)^5/f+2/7*a^3*c^5*tan(f*x+e)^
7/f
```

3.22.2 Mathematica [A] (verified)

Time = 2.20 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.54

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx =$$

$$\frac{a^3 c^5 (-20160 \operatorname{arctanh}(\sin(e + fx)) + \sec^8(e + fx)(5705 \sin(e + fx) - 1792 \sin(2(e + fx)) + 21 \sin(3(e + fx)) + 1792 \sin(4(e + fx)) + 2065 \sin(5(e + fx)) - 768 \sin(6(e + fx)) + 581 \sin(7(e + fx)) + 128 \sin(8(e + fx))))}{f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^5,x]`

output `-1/57344*(a^3*c^5*(-20160*ArcTanh[Sin[e + f*x]] + Sec[e + f*x]^8*(5705*Sin[e + f*x] - 1792*Sin[2*(e + f*x)] + 21*Sin[3*(e + f*x)] + 1792*Sin[4*(e + f*x)] + 2065*Sin[5*(e + f*x)] - 768*Sin[6*(e + f*x)] + 581*Sin[7*(e + f*x)] + 128*Sin[8*(e + f*x)])))/f`

3.22.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^5 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^5 dx$$

$$\downarrow \text{4446}$$

$$-a^3 c^3 \int (c^2 \sec^3(e + fx) \tan^6(e + fx) - 2c^2 \sec^2(e + fx) \tan^6(e + fx) + c^2 \sec(e + fx) \tan^6(e + fx)) dx$$

$$\downarrow \text{2009}$$

$$-a^3 c^3 \left(-\frac{45c^2 \operatorname{arctanh}(\sin(e + fx))}{128f} - \frac{2c^2 \tan^7(e + fx)}{7f} + \frac{c^2 \tan^5(e + fx) \sec^3(e + fx)}{8f} - \frac{5c^2 \tan^3(e + fx) \sec^3(e + fx)}{48f} \right)$$

3.22. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^5,x]`

output `-(a^3*c^3*((-45*c^2*ArcTanh[Sin[e + f*x]])/(128*f) + (35*c^2*Sec[e + f*x]*Tan[e + f*x])/(128*f) + (5*c^2*Sec[e + f*x]^3*Tan[e + f*x])/(64*f) - (5*c^2*Sec[e + f*x]*Tan[e + f*x]^3)/(24*f) - (5*c^2*Sec[e + f*x]^3*Tan[e + f*x]^3)/(48*f) + (c^2*Sec[e + f*x]*Tan[e + f*x]^5)/(6*f) + (c^2*Sec[e + f*x]^3*Tan[e + f*x]^5)/(8*f) - (2*c^2*Tan[e + f*x]^7)/(7*f)))`

3.22.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.22.4 Maple [A] (verified)

Time = 8.28 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.16

method	result
norman	$-\frac{45c^5 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{64f} + \frac{345c^5 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{64f} - \frac{1149c^5 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{64f} + \frac{15159c^5 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{448f} - \frac{17609c^5 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{448f}$
risch	$\frac{ic^5 a^3 (581 e^{15i(fx+e)} - 1792 e^{14i(fx+e)} + 2065 e^{13i(fx+e)} - 1792 e^{12i(fx+e)} + 21 e^{11i(fx+e)} - 8960 e^{10i(fx+e)} + 5705 e^{9i(fx+e)} - 1792 e^{8i(fx+e)} + 179 e^{7i(fx+e)} - 17 e^{6i(fx+e)} + 1 e^{5i(fx+e)})}{448f}$
parallelrisch	$-\frac{815 \left(9 \left(\frac{35}{2} + 28 \cos(2fx+2e) + 14 \cos(4fx+4e) + 4 \cos(6fx+6e) + \frac{\cos(8fx+8e)}{2} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 9 \left(-\frac{35}{2} - 28 \cos(2fx+2e) - 14 \cos(4fx+4e) - 4 \cos(6fx+6e) - \frac{\cos(8fx+8e)}{2} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \right)}{163}$
parts	$-\frac{2a^3 c^5 \tan(fx+e)}{f} - \frac{a^3 c^5 \sec(fx+e) \tan(fx+e)}{f} - \frac{6c^5 a^3 \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e)}{f} + \frac{6c^5 a^3 \left(-\frac{8}{15} - \frac{\sec(fx+e)}{5} \right) \tan(fx+e)}{f}$
derivativedivides	$-c^5 a^3 \left(-\left(-\frac{\sec(fx+e)^7}{8} - \frac{7 \sec(fx+e)^5}{48} - \frac{35 \sec(fx+e)^3}{192} - \frac{35 \sec(fx+e)}{128} \right) \tan(fx+e) + \frac{35 \ln(\sec(fx+e) + \tan(fx+e))}{128} \right) - 2c^5 a^3$
default	$-c^5 a^3 \left(-\left(-\frac{\sec(fx+e)^7}{8} - \frac{7 \sec(fx+e)^5}{48} - \frac{35 \sec(fx+e)^3}{192} - \frac{35 \sec(fx+e)}{128} \right) \tan(fx+e) + \frac{35 \ln(\sec(fx+e) + \tan(fx+e))}{128} \right) - 2c^5 a^3$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^5,x,method=_RETURNVERBOSE)
```

```
output (-45/64*c^5*a^3/f*tan(1/2*f*x+1/2*e)+345/64*c^5*a^3/f*tan(1/2*f*x+1/2*e)^3-1149/64*c^5*a^3/f*tan(1/2*f*x+1/2*e)^5+15159/448*c^5*a^3/f*tan(1/2*f*x+1/2*e)^7-17609/448*c^5*a^3/f*tan(1/2*f*x+1/2*e)^9-1149/64*c^5*a^3/f*tan(1/2*f*x+1/2*e)^11+345/64*c^5*a^3/f*tan(1/2*f*x+1/2*e)^13-45/64*c^5*a^3/f*tan(1/2*f*x+1/2*e)^15)/(tan(1/2*f*x+1/2*e)^2-1)^8-45/128*c^5*a^3/f*ln(tan(1/2*f*x+1/2*e)-1)+45/128*c^5*a^3/f*ln(tan(1/2*f*x+1/2*e)+1)
```

3.22.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.94

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$$

$$= \frac{315 a^3 c^5 \cos(fx + e)^8 \log(\sin(fx + e) + 1) - 315 a^3 c^5 \cos(fx + e)^8 \log(-\sin(fx + e) + 1) - 2(256 a^3 c^5 \cos(fx + e)^8 \log(\tan(fx + e) + 1) - 256 a^3 c^5 \cos(fx + e)^8 \log(\tan(fx + e) - 1))}{128}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^5,x, algorithm="fracas")
```

3.22. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$

output `1/1792*(315*a^3*c^5*cos(f*x + e)^8*log(sin(f*x + e) + 1) - 315*a^3*c^5*cos(f*x + e)^8*log(-sin(f*x + e) + 1) - 2*(256*a^3*c^5*cos(f*x + e)^7 + 581*a^3*c^5*cos(f*x + e)^6 - 768*a^3*c^5*cos(f*x + e)^5 - 210*a^3*c^5*cos(f*x + e)^4 + 768*a^3*c^5*cos(f*x + e)^3 - 168*a^3*c^5*cos(f*x + e)^2 - 256*a^3*c^5*cos(f*x + e) + 112*a^3*c^5)*sin(f*x + e))/(f*cos(f*x + e)^8)`

3.22.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx \\ &= -a^3c^5 \left(\int (-\sec(e + fx)) dx + \int 2\sec^2(e + fx) dx + \int 2\sec^3(e + fx) dx \right. \\ & \quad \left. + \int (-6\sec^4(e + fx)) dx + \int 6\sec^6(e + fx) dx + \int (-2\sec^7(e + fx)) dx \right. \\ & \quad \left. + \int (-2\sec^8(e + fx)) dx + \int \sec^9(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**5,x)`

output `-a**3*c**5*(Integral(-sec(e + f*x), x) + Integral(2*sec(e + f*x)**2, x) + Integral(2*sec(e + f*x)**3, x) + Integral(-6*sec(e + f*x)**4, x) + Integral(6*sec(e + f*x)**6, x) + Integral(-2*sec(e + f*x)**7, x) + Integral(-2*sec(e + f*x)**8, x) + Integral(sec(e + f*x)**9, x))`

3.22.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(190) = 380$.

Time = 0.21 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.98

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx \\ &= \frac{1536 (5 \tan(fx + e)^7 + 21 \tan(fx + e)^5 + 35 \tan(fx + e)^3 + 35 \tan(fx + e))a^3c^5 - 10752 (3 \tan(fx + e) + 1)}{\dots} \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^5,x, algorithm="maxima")`

3.22. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$

output `1/26880*(1536*(5*tan(f*x + e)^7 + 21*tan(f*x + e)^5 + 35*tan(f*x + e)^3 + 35*tan(f*x + e))*a^3*c^5 - 10752*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^3*c^5 + 53760*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c^5 + 35*a^3*c^5*(2*(105*sin(f*x + e)^7 - 385*sin(f*x + e)^5 + 511*sin(f*x + e)^3 - 279*sin(f*x + e))/(sin(f*x + e)^8 - 4*sin(f*x + e)^6 + 6*sin(f*x + e)^4 - 4*sin(f*x + e)^2 + 1) - 105*log(sin(f*x + e) + 1) + 105*log(sin(f*x + e) - 1)) - 560*a^3*c^5*(2*(15*sin(f*x + e)^5 - 40*sin(f*x + e)^3 + 33*sin(f*x + e))/(sin(f*x + e)^6 - 3*sin(f*x + e)^4 + 3*sin(f*x + e)^2 - 1) - 15*log(sin(f*x + e) + 1) + 15*log(sin(f*x + e) - 1)) + 13440*a^3*c^5*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 26880*a^3*c^5*log(sec(f*x + e) + tan(f*x + e)) - 53760*a^3*c^5*tan(f*x + e))/f`

3.22.8 Giac [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.05

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$$

$$= \frac{315 a^3 c^5 \log\left(\left|\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) + 1\right|\right) - 315 a^3 c^5 \log\left(\left|\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) - 1\right|\right) - \frac{2\left(315 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{15} - 2415 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{13} + 8043 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11} - 17609 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^9 - 15159 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^7 + 8043 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 - 2415 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + 315 a^3 c^5 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)\right)}{\left(\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 1\right)^8}}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^5,x, algorithm="giac")`

output `1/896*(315*a^3*c^5*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 315*a^3*c^5*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(315*a^3*c^5*tan(1/2*f*x + 1/2*e)^15 - 2415*a^3*c^5*tan(1/2*f*x + 1/2*e)^13 + 8043*a^3*c^5*tan(1/2*f*x + 1/2*e)^11 + 17609*a^3*c^5*tan(1/2*f*x + 1/2*e)^9 - 15159*a^3*c^5*tan(1/2*f*x + 1/2*e)^7 + 8043*a^3*c^5*tan(1/2*f*x + 1/2*e)^5 - 2415*a^3*c^5*tan(1/2*f*x + 1/2*e)^3 + 315*a^3*c^5*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^8)/f`

3.22. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$

3.22.9 Mupad [B] (verification not implemented)

Time = 16.93 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.38

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx = \frac{45 a^3 c^5 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{64 f} - \frac{45 a^3 c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{15}}{64} - \frac{345 a^3 c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{13}}{64} + \frac{1149 a^3 c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}{64} + \frac{17609 a^3 c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{448} - \frac{15159 a^3 c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{448} - f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{16} - 8 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{14} + 28 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{12} - 56 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} + 70 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 56 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 28 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 8 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1 \right)$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^5)/cos(e + f*x),x)`output `(45*a^3*c^5*atanh(tan(e/2 + (f*x)/2)))/(64*f) - ((1149*a^3*c^5*tan(e/2 + (f*x)/2)^5)/64 - (345*a^3*c^5*tan(e/2 + (f*x)/2)^3)/64 - (15159*a^3*c^5*tan(e/2 + (f*x)/2)^7)/448 + (17609*a^3*c^5*tan(e/2 + (f*x)/2)^9)/448 + (1149*a^3*c^5*tan(e/2 + (f*x)/2)^11)/64 - (345*a^3*c^5*tan(e/2 + (f*x)/2)^13)/64 + (45*a^3*c^5*tan(e/2 + (f*x)/2)^15)/64 + (45*a^3*c^5*tan(e/2 + (f*x)/2))/64)/(f*(28*tan(e/2 + (f*x)/2)^4 - 8*tan(e/2 + (f*x)/2)^2 - 56*tan(e/2 + (f*x)/2)^6 + 70*tan(e/2 + (f*x)/2)^8 - 56*tan(e/2 + (f*x)/2)^10 + 28*tan(e/2 + (f*x)/2)^12 - 8*tan(e/2 + (f*x)/2)^14 + tan(e/2 + (f*x)/2)^16 + 1))`

3.23 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$

3.23.1	Optimal result	252
3.23.2	Mathematica [A] (verified)	252
3.23.3	Rubi [A] (verified)	253
3.23.4	Maple [C] (verified)	254
3.23.5	Fricas [A] (verification not implemented)	255
3.23.6	Sympy [F]	255
3.23.7	Maxima [B] (verification not implemented)	256
3.23.8	Giac [A] (verification not implemented)	256
3.23.9	Mupad [B] (verification not implemented)	257

3.23.1 Optimal result

Integrand size = 32, antiderivative size = 121

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$$

$$= \frac{5a^3c^4 \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{5a^3c^4 \sec(e + fx) \tan(e + fx)}{16f}$$

$$+ \frac{5a^3c^4 \sec(e + fx) \tan^3(e + fx)}{24f} - \frac{a^3c^4 \sec(e + fx) \tan^5(e + fx)}{6f} + \frac{a^3c^4 \tan^7(e + fx)}{7f}$$

output `5/16*a^3*c^4*arctanh(sin(f*x+e))/f-5/16*a^3*c^4*sec(f*x+e)*tan(f*x+e)/f+5/24*a^3*c^4*sec(f*x+e)*tan(f*x+e)^3/f-1/6*a^3*c^4*sec(f*x+e)*tan(f*x+e)^5/f+1/7*a^3*c^4*tan(f*x+e)^7/f`

3.23.2 Mathematica [A] (verified)

Time = 0.92 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.84

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$$

$$= \frac{a^3c^4(3360 \operatorname{arctanh}(\sin(e + fx)) - \sec^7(e + fx)(-840 \sin(e + fx) + 595 \sin(2(e + fx))) + 504 \sin(3(e + fx)))}{10752f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^4,x]`

output `(a^3*c^4*(3360*ArcTanh[Sin[e + f*x]] - Sec[e + f*x]^7*(-840*Sin[e + f*x] + 595*Sin[2*(e + f*x)] + 504*Sin[3*(e + f*x)] + 196*Sin[4*(e + f*x)] - 168*Sin[5*(e + f*x)] + 231*Sin[6*(e + f*x)] + 24*Sin[7*(e + f*x)])))/(10752*f)`

3.23.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.86, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^4 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^4 dx$$

$$\downarrow 4446$$

$$-a^3 c^3 \int (c \sec(e + fx) \tan^6(e + fx) - c \sec^2(e + fx) \tan^6(e + fx)) dx$$

$$\downarrow 2009$$

$$-a^3 c^3 \left(-\frac{5c \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{c \tan^7(e + fx)}{7f} + \frac{c \tan^5(e + fx) \sec(e + fx)}{6f} - \frac{5c \tan^3(e + fx) \sec(e + fx)}{24f} + \dots \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^4,x]`

output `-(a^3*c^3*((-5*c*ArcTanh[Sin[e + f*x]])/(16*f) + (5*c*Sec[e + f*x]*Tan[e + f*x])/(16*f) - (5*c*Sec[e + f*x]*Tan[e + f*x]^3)/(24*f) + (c*Sec[e + f*x]*Tan[e + f*x]^5)/(6*f) - (c*Tan[e + f*x]^7)/(7*f)))`

3.23.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.23.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 8.49 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.45

method	result
risch	$\frac{ic^4 a^3 (231 e^{13i(fx+e)} - 336 e^{12i(fx+e)} + 196 e^{11i(fx+e)} + 595 e^{9i(fx+e)} - 1680 e^{8i(fx+e)} - 595 e^{5i(fx+e)} - 1008 e^{4i(fx+e)} - 168 f (1 + e^{2i(fx+e)})^7}{168 f (1 + e^{2i(fx+e)})^7}$
norman	$\frac{5c^4 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 25c^4 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + \frac{283c^4 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{24f} - \frac{128c^4 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{7f} - \frac{283c^4 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{24f} + \frac{25c^4 a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{24f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^7}$
parallelrisc	$5a^3 \left(\frac{(-\cos(7fx+7e) - 7\cos(5fx+5e) - 21\cos(3fx+3e) - 35\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{16} + \frac{(\cos(7fx+7e) + 7\cos(5fx+5e) + 21\cos(3fx+3e) + 35\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{16} \right)$
derivativedivides	$-c^4 a^3 \left(-\frac{16}{35} - \frac{\sec(fx+e)^6}{7} - \frac{6 \sec(fx+e)^4}{35} - \frac{8 \sec(fx+e)^2}{35} \right) \tan(fx+e) - c^4 a^3 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5 \sec(fx+e)^3}{24} - \frac{5 \sec(fx+e)}{16} \right) \tan(fx+e) \right)$
default	$-c^4 a^3 \left(-\frac{16}{35} - \frac{\sec(fx+e)^6}{7} - \frac{6 \sec(fx+e)^4}{35} - \frac{8 \sec(fx+e)^2}{35} \right) \tan(fx+e) - c^4 a^3 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5 \sec(fx+e)^3}{24} - \frac{5 \sec(fx+e)}{16} \right) \tan(fx+e) \right)$
parts	$\frac{c^4 a^3 \ln(\sec(fx+e) + \tan(fx+e))}{f} - \frac{c^4 a^3 \left(-\frac{16}{35} - \frac{\sec(fx+e)^6}{7} - \frac{6 \sec(fx+e)^4}{35} - \frac{8 \sec(fx+e)^2}{35} \right) \tan(fx+e)}{f} - \frac{c^4 a^3 \tan(fx+e)}{f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^4,x,method=_RETURNVERBOSE)
```

3.23. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$

output $\frac{1}{168}I^4c^4a^3(231\exp(13I(f*x+e))-336\exp(12I(f*x+e))+196\exp(11I(f*x+e))+595\exp(9I(f*x+e))-1680\exp(8I(f*x+e))-595\exp(5I(f*x+e))-1008\exp(4I(f*x+e))-196\exp(3I(f*x+e))-231\exp(I(f*x+e))-48)/f/(1+\exp(2I(f*x+e)))^7-5/16c^4a^3/f*\ln(\exp(I(f*x+e))-I)+5/16c^4a^3/f*\ln(\exp(I(f*x+e))+I)$

3.23.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.46

$$\int \sec(e+fx)(a+a\sec(e+fx))^3(c-c\sec(e+fx))^4 dx$$

$$= \frac{105a^3c^4 \cos(fx+e)^7 \log(\sin(fx+e)+1) - 105a^3c^4 \cos(fx+e)^7 \log(-\sin(fx+e)+1) - 2(48a^3c^4 \cos(fx+e)^6 + 231a^3c^4 \cos(fx+e)^5 - 144a^3c^4 \cos(fx+e)^4 - 182a^3c^4 \cos(fx+e)^3 + 144a^3c^4 \cos(fx+e)^2 + 56a^3c^4 \cos(fx+e) - 48a^3c^4) \sin(fx+e)}{f \cos(fx+e)^7}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^4,x, algorithm="fricas")`

output $\frac{1}{672}*(105*a^3*c^4*\cos(f*x+e)^7*\log(\sin(f*x+e)+1) - 105*a^3*c^4*\cos(f*x+e)^7*\log(-\sin(f*x+e)+1) - 2*(48*a^3*c^4*\cos(f*x+e)^6 + 231*a^3*c^4*\cos(f*x+e)^5 - 144*a^3*c^4*\cos(f*x+e)^4 - 182*a^3*c^4*\cos(f*x+e)^3 + 144*a^3*c^4*\cos(f*x+e)^2 + 56*a^3*c^4*\cos(f*x+e) - 48*a^3*c^4)*\sin(f*x+e))/(f*\cos(f*x+e)^7)$

3.23.6 Sympy [F]

$$\int \sec(e+fx)(a+a\sec(e+fx))^3(c-c\sec(e+fx))^4 dx$$

$$= a^3c^4 \left(\int \sec(e+fx) dx + \int (-\sec^2(e+fx)) dx + \int (-3\sec^3(e+fx)) dx \right. \\ \left. + \int 3\sec^4(e+fx) dx + \int 3\sec^5(e+fx) dx + \int (-3\sec^6(e+fx)) dx \right. \\ \left. + \int (-\sec^7(e+fx)) dx + \int \sec^8(e+fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**4,x)`

3.23. $\int \sec(e+fx)(a+a\sec(e+fx))^3(c-c\sec(e+fx))^4 dx$

output `a**3*c**4*(Integral(sec(e + f*x), x) + Integral(-sec(e + f*x)**2, x) + Integral(-3*sec(e + f*x)**3, x) + Integral(3*sec(e + f*x)**4, x) + Integral(3*sec(e + f*x)**5, x) + Integral(-3*sec(e + f*x)**6, x) + Integral(-sec(e + f*x)**7, x) + Integral(sec(e + f*x)**8, x))`

3.23.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 368 vs. $2(111) = 222$.

Time = 0.21 (sec) , antiderivative size = 368, normalized size of antiderivative = 3.04

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$$

$$= \frac{96(5 \tan(fx + e)^7 + 21 \tan(fx + e)^5 + 35 \tan(fx + e)^3 + 35 \tan(fx + e))a^3c^4 - 672(3 \tan(fx + e)^5}{}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output `1/3360*(96*(5*tan(f*x + e)^7 + 21*tan(f*x + e)^5 + 35*tan(f*x + e)^3 + 35*tan(f*x + e))*a^3*c^4 - 672*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^3*c^4 + 3360*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c^4 + 35*a^3*c^4*(2*(15*sin(f*x + e)^5 - 40*sin(f*x + e)^3 + 33*sin(f*x + e))/(sin(f*x + e)^6 - 3*sin(f*x + e)^4 + 3*sin(f*x + e)^2 - 1) - 15*log(sin(f*x + e) + 1) + 15*log(sin(f*x + e) - 1)) - 630*a^3*c^4*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) + 2520*a^3*c^4*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 3360*a^3*c^4*log(sec(f*x + e) + tan(f*x + e)) - 3360*a^3*c^4*tan(f*x + e))/f`

3.23.8 Giac [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.63

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$$

$$= \frac{105 a^3 c^4 \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + 1 \right| \right) - 105 a^3 c^4 \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(105 a^3 c^4 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^{13} - 700 a^3 c^4 \right)}{}$$

3.23. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output `1/336*(105*a^3*c^4*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 105*a^3*c^4*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(105*a^3*c^4*tan(1/2*f*x + 1/2*e)^13 - 700*a^3*c^4*tan(1/2*f*x + 1/2*e)^11 + 1981*a^3*c^4*tan(1/2*f*x + 1/2*e)^9 + 3072*a^3*c^4*tan(1/2*f*x + 1/2*e)^7 - 1981*a^3*c^4*tan(1/2*f*x + 1/2*e)^5 + 700*a^3*c^4*tan(1/2*f*x + 1/2*e)^3 - 105*a^3*c^4*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1/2*e)^2 - 1)^7)/f`

3.23.9 Mupad [B] (verification not implemented)

Time = 16.85 (sec) , antiderivative size = 252, normalized size of antiderivative = 2.08

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^4 dx = \frac{5 a^3 c^4 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{8 f} - \frac{5 a^3 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{13}}{8} - \frac{25 a^3 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}{6} + \frac{283 a^3 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{24} + \frac{128 a^3 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{7} - \frac{283 a^3 c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{24} + f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{14} - 7 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{12} + 21 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - 35 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 35 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 21 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 7 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1 \right)$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^4)/cos(e + f*x),x)`

output `(5*a^3*c^4*atanh(tan(e/2 + (f*x)/2)))/(8*f) - ((25*a^3*c^4*tan(e/2 + (f*x)/2)^3)/6 - (283*a^3*c^4*tan(e/2 + (f*x)/2)^5)/24 + (128*a^3*c^4*tan(e/2 + (f*x)/2)^7)/7 + (283*a^3*c^4*tan(e/2 + (f*x)/2)^9)/24 - (25*a^3*c^4*tan(e/2 + (f*x)/2)^11)/6 + (5*a^3*c^4*tan(e/2 + (f*x)/2)^13)/8 - (5*a^3*c^4*tan(e/2 + (f*x)/2))/8)/(f*(7*tan(e/2 + (f*x)/2)^2 - 21*tan(e/2 + (f*x)/2)^4 + 35*tan(e/2 + (f*x)/2)^6 - 35*tan(e/2 + (f*x)/2)^8 + 21*tan(e/2 + (f*x)/2)^10 - 7*tan(e/2 + (f*x)/2)^12 + tan(e/2 + (f*x)/2)^14 - 1))`

3.24 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$

3.24.1	Optimal result	258
3.24.2	Mathematica [A] (verified)	258
3.24.3	Rubi [A] (verified)	259
3.24.4	Maple [C] (verified)	261
3.24.5	Fricas [A] (verification not implemented)	262
3.24.6	Sympy [F]	262
3.24.7	Maxima [B] (verification not implemented)	263
3.24.8	Giac [A] (verification not implemented)	263
3.24.9	Mupad [B] (verification not implemented)	264

3.24.1 Optimal result

Integrand size = 32, antiderivative size = 100

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx \\ &= \frac{5a^3c^3 \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{5a^3c^3 \sec(e + fx) \tan(e + fx)}{16f} \\ &+ \frac{5a^3c^3 \sec(e + fx) \tan^3(e + fx)}{24f} - \frac{a^3c^3 \sec(e + fx) \tan^5(e + fx)}{6f} \end{aligned}$$

output $5/16*a^3*c^3*\operatorname{arctanh}(\sin(f*x+e))/f-5/16*a^3*c^3*\sec(f*x+e)*\tan(f*x+e)/f+5/24*a^3*c^3*\sec(f*x+e)*\tan(f*x+e)^3/f-1/6*a^3*c^3*\sec(f*x+e)*\tan(f*x+e)^5/f$

3.24.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.25

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx \\ &= -a^3c^3 \left(-\frac{5 \operatorname{arctanh}(\sin(e + fx))}{16f} - \frac{5 \sec(e + fx) \tan(e + fx)}{16f} \right. \\ &\quad \left. - \frac{5 \sec^3(e + fx) \tan(e + fx)}{24f} + \frac{5 \sec^5(e + fx) \tan(e + fx)}{6f} \right. \\ &\quad \left. - \frac{5 \sec^3(e + fx) \tan^3(e + fx)}{3f} + \frac{\sec(e + fx) \tan^5(e + fx)}{f} \right) \end{aligned}$$

3.24. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^3,x]`

output `-(a^3*c^3*((-5*ArcTanh[Sin[e + f*x]])/(16*f) - (5*Sec[e + f*x]*Tan[e + f*x])/ (16*f) - (5*Sec[e + f*x]^3*Tan[e + f*x])/(24*f) + (5*Sec[e + f*x]^5*Tan[e + f*x])/(6*f) - (5*Sec[e + f*x]^3*Tan[e + f*x]^3)/(3*f) + (Sec[e + f*x]*Tan[e + f*x]^5)/f))`

3.24.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.94, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4446, 3042, 3091, 3042, 3091, 3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^3 dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 dx \\
 & \quad \downarrow \text{4446} \\
 & -a^3 c^3 \int \sec(e + fx) \tan^6(e + fx) dx \\
 & \quad \downarrow \text{3042} \\
 & -a^3 c^3 \int \sec(e + fx) \tan(e + fx)^6 dx \\
 & \quad \downarrow \text{3091} \\
 & -a^3 c^3 \left(\frac{\tan^5(e + fx) \sec(e + fx)}{6f} - \frac{5}{6} \int \sec(e + fx) \tan^4(e + fx) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & -a^3 c^3 \left(\frac{\tan^5(e + fx) \sec(e + fx)}{6f} - \frac{5}{6} \int \sec(e + fx) \tan(e + fx)^4 dx \right) \\
 & \quad \downarrow \text{3091}
 \end{aligned}$$

$$\begin{aligned}
& -a^3 c^3 \left(\frac{\tan^5(e+fx) \sec(e+fx)}{6f} - \frac{5}{6} \left(\frac{\tan^3(e+fx) \sec(e+fx)}{4f} - \frac{3}{4} \int \sec(e+fx) \tan^2(e+fx) dx \right) \right) \\
& \quad \downarrow \text{3042} \\
& -a^3 c^3 \left(\frac{\tan^5(e+fx) \sec(e+fx)}{6f} - \frac{5}{6} \left(\frac{\tan^3(e+fx) \sec(e+fx)}{4f} - \frac{3}{4} \int \sec(e+fx) \tan(e+fx)^2 dx \right) \right) \\
& \quad \downarrow \text{3091} \\
& -a^3 c^3 \left(\frac{\tan^5(e+fx) \sec(e+fx)}{6f} - \frac{5}{6} \left(\frac{\tan^3(e+fx) \sec(e+fx)}{4f} - \frac{3}{4} \left(\frac{\tan(e+fx) \sec(e+fx)}{2f} - \frac{1}{2} \int \sec(e+fx) dx \right) \right) \right) \\
& \quad \downarrow \text{3042} \\
& -a^3 c^3 \left(\frac{\tan^5(e+fx) \sec(e+fx)}{6f} - \frac{5}{6} \left(\frac{\tan^3(e+fx) \sec(e+fx)}{4f} - \frac{3}{4} \left(\frac{\tan(e+fx) \sec(e+fx)}{2f} - \frac{1}{2} \int \csc(e+fx) dx \right) \right) \right) \\
& \quad \downarrow \text{4257} \\
& -a^3 c^3 \left(\frac{\tan^5(e+fx) \sec(e+fx)}{6f} - \frac{5}{6} \left(\frac{\tan^3(e+fx) \sec(e+fx)}{4f} - \frac{3}{4} \left(\frac{\tan(e+fx) \sec(e+fx)}{2f} - \frac{\operatorname{arctanh}(\sin(e+fx))}{2f} \right) \right) \right)
\end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^3,x]`

output `-(a^3*c^3*((Sec[e + f*x]*Tan[e + f*x]^5)/(6*f) - (5*((Sec[e + f*x]*Tan[e + f*x]^3)/(4*f) - (3*(-1/2*ArcTanh[Sin[e + f*x]]/f + (Sec[e + f*x]*Tan[e + f*x])/(2*f))))/4)/6))`

3.24.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3091 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

3.24. $\int \sec(e+fx)(a + a \sec(e+fx))^3(c - c \sec(e+fx))^3 dx$

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.24.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.91 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.42

method	result
risch	$\frac{ia^3c^3(33e^{11i(fx+e)} - 5e^{9i(fx+e)} + 90e^{7i(fx+e)} - 90e^{5i(fx+e)} + 5e^{3i(fx+e)} - 33e^{i(fx+e)})}{24f(1+e^{2i(fx+e)})^6} - \frac{5c^3a^3 \ln(e^{i(fx+e)} - i)}{16f} + \dots$
parallelrisch	$\frac{5a^3c^3 \left(\left(-\frac{45 \cos(2fx+2e)}{2} - 9 \cos(4fx+4e) - \frac{3 \cos(6fx+6e)}{2} - 15 \right) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) + \left(\frac{3 \cos(6fx+6e)}{2} + 9 \cos(4fx+4e) + \dots \right) \right)}{24f(6 \cos(4fx+4e) + 10 + 15 \cos(2fx+2e) + \dots)}$
derivativedivides	$-c^3a^3 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5 \sec(fx+e)^3}{24} - \frac{5 \sec(fx+e)}{16} \right) \tan(fx+e) + \frac{5 \ln(\sec(fx+e) + \tan(fx+e))}{16} \right) + 3c^3a^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} \right) \right)$
default	$-c^3a^3 \left(-\left(-\frac{\sec(fx+e)^5}{6} - \frac{5 \sec(fx+e)^3}{24} - \frac{5 \sec(fx+e)}{16} \right) \tan(fx+e) + \frac{5 \ln(\sec(fx+e) + \tan(fx+e))}{16} \right) + 3c^3a^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} \right) \right)$
parts	$\frac{c^3a^3 \ln(\sec(fx+e) + \tan(fx+e))}{f} - \frac{3c^3a^3 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right)}{f} + \frac{3c^3a^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} \right) \right)}{f}$
norman	$\frac{-\frac{5c^3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{8f} + \frac{85c^3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{24f} - \frac{33c^3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{4f} - \frac{33c^3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{4f} + \frac{85c^3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{24f} - \frac{5c^3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{8f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1 \right)^6}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `1/24*I*a^3*c^3/f/(1+exp(2*I*(f*x+e)))^6*(33*exp(11*I*(f*x+e))-5*exp(9*I*(f*x+e))+90*exp(7*I*(f*x+e))-90*exp(5*I*(f*x+e))+5*exp(3*I*(f*x+e))-33*exp(I*(f*x+e)))-5/16*c^3*a^3/f*ln(exp(I*(f*x+e))-I)+5/16*c^3*a^3/f*ln(exp(I*(f*x+e))+I)`

$$3.24. \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$$

3.24.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.15

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$$

$$= \frac{15 a^3 c^3 \cos(fx + e)^6 \log(\sin(fx + e) + 1) - 15 a^3 c^3 \cos(fx + e)^6 \log(-\sin(fx + e) + 1) - 2(33 a^3 c^3 \cos(fx + e)^4 - 26 a^3 c^3 \cos(fx + e)^2 + 8 a^3 c^3) \sin(fx + e)}{96 f \cos(fx + e)^6}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^3,x, algorithm="fricas")
```

```
output 1/96*(15*a^3*c^3*cos(f*x + e)^6*log(sin(f*x + e) + 1) - 15*a^3*c^3*cos(f*x + e)^6*log(-sin(f*x + e) + 1) - 2*(33*a^3*c^3*cos(f*x + e)^4 - 26*a^3*c^3*cos(f*x + e)^2 + 8*a^3*c^3)*sin(f*x + e))/(f*cos(f*x + e)^6)
```

3.24.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$$

$$= -a^3 c^3 \left(\int (-\sec(e + fx)) dx + \int 3 \sec^3(e + fx) dx + \int (-3 \sec^5(e + fx)) dx + \int \sec^7(e + fx) dx \right)$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**3,x)
```

```
output -a**3*c**3*(Integral(-sec(e + f*x), x) + Integral(3*sec(e + f*x)**3, x) + Integral(-3*sec(e + f*x)**5, x) + Integral(sec(e + f*x)**7, x))
```

3.24.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 244 vs. 2(92) = 184.

Time = 0.23 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.44

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$$

$$= \frac{a^3 c^3 \left(\frac{2(15 \sin(fx+e)^5 - 40 \sin(fx+e)^3 + 33 \sin(fx+e))}{\sin(fx+e)^6 - 3 \sin(fx+e)^4 + 3 \sin(fx+e)^2 - 1} - 15 \log(\sin(fx+e) + 1) + 15 \log(\sin(fx+e) - 1) \right) - 18 a^3 c^3 \log(\sec(fx+e) + \tan(fx+e))}{96 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/96*(a^3*c^3*(2*(15*sin(f*x + e)^5 - 40*sin(f*x + e)^3 + 33*sin(f*x + e))/(sin(f*x + e)^6 - 3*sin(f*x + e)^4 + 3*sin(f*x + e)^2 - 1) - 15*log(sin(f*x + e) + 1) + 15*log(sin(f*x + e) - 1)) - 18*a^3*c^3*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) + 72*a^3*c^3*(2*sin(f*x + e))/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 96*a^3*c^3*log(sec(f*x + e) + tan(f*x + e)))/f`

3.24.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.03

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$$

$$= \frac{15 a^3 c^3 \log(|\sin(fx+e) + 1|) - 15 a^3 c^3 \log(|\sin(fx+e) - 1|) + \frac{2(33 a^3 c^3 \sin(fx+e)^5 - 40 a^3 c^3 \sin(fx+e)^3 + 15 a^3 c^3 \sin(fx+e))}{(\sin(fx+e)^2 - 1)^3}}{96 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `1/96*(15*a^3*c^3*log(abs(sin(f*x + e) + 1)) - 15*a^3*c^3*log(abs(sin(f*x + e) - 1)) + 2*(33*a^3*c^3*sin(f*x + e)^5 - 40*a^3*c^3*sin(f*x + e)^3 + 15*a^3*c^3*sin(f*x + e))/(sin(f*x + e)^2 - 1)^3)/f`

3.24. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx$

3.24.9 Mupad [B] (verification not implemented)

Time = 16.50 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.20

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^3 dx = \frac{5 a^3 c^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{8 f} - \frac{5 a^3 c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}{8} - \frac{85 a^3 c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{24} + \frac{33 a^3 c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{4} + \frac{33 a^3 c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{4} - \frac{85 a^3 c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{24} + \frac{5 a^3 c^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{8} + \frac{1}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{12} - 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} + 15 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 20 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 15 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1 \right)}$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^3)/cos(e + f*x),x)`output `(5*a^3*c^3*atanh(tan(e/2 + (f*x)/2)))/(8*f) - ((33*a^3*c^3*tan(e/2 + (f*x)/2)^5)/4 - (85*a^3*c^3*tan(e/2 + (f*x)/2)^3)/24 + (33*a^3*c^3*tan(e/2 + (f*x)/2)^7)/4 - (85*a^3*c^3*tan(e/2 + (f*x)/2)^9)/24 + (5*a^3*c^3*tan(e/2 + (f*x)/2)^11)/8 + (5*a^3*c^3*tan(e/2 + (f*x)/2))/8)/(f*(15*tan(e/2 + (f*x)/2)^4 - 6*tan(e/2 + (f*x)/2)^2 - 20*tan(e/2 + (f*x)/2)^6 + 15*tan(e/2 + (f*x)/2)^8 - 6*tan(e/2 + (f*x)/2)^10 + tan(e/2 + (f*x)/2)^12 + 1))`

3.25 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx$

3.25.1	Optimal result	265
3.25.2	Mathematica [A] (verified)	265
3.25.3	Rubi [A] (verified)	266
3.25.4	Maple [C] (verified)	267
3.25.5	Fricas [A] (verification not implemented)	268
3.25.6	Sympy [F]	268
3.25.7	Maxima [B] (verification not implemented)	269
3.25.8	Giac [A] (verification not implemented)	269
3.25.9	Mupad [B] (verification not implemented)	270

3.25.1 Optimal result

Integrand size = 32, antiderivative size = 94

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx \\ &= \frac{3a^3c^2 \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{3a^3c^2 \sec(e + fx) \tan(e + fx)}{8f} \\ & \quad + \frac{a^3c^2 \sec(e + fx) \tan^3(e + fx)}{4f} + \frac{a^3c^2 \tan^5(e + fx)}{5f} \end{aligned}$$

output `3/8*a^3*c^2*arctanh(sin(f*x+e))/f-3/8*a^3*c^2*sec(f*x+e)*tan(f*x+e)/f+1/4*a^3*c^2*sec(f*x+e)*tan(f*x+e)^3/f+1/5*a^3*c^2*tan(f*x+e)^5/f`

3.25.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.86

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx \\ &= \frac{a^3c^2(120\operatorname{arctanh}(\sin(e + fx)) + \sec^5(e + fx)(40 \sin(e + fx) - 10 \sin(2(e + fx)) - 20 \sin(3(e + fx)) - 2 \sin(4(e + fx)))}{320f} \end{aligned}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^2,x]`

output $(a^3 c^2 (120 \operatorname{ArcTanh}[\sin[e + f x]] + \operatorname{Sec}[e + f x]^5 (40 \sin[e + f x] - 10 \sin[2(e + f x)] - 20 \sin[3(e + f x)] - 25 \sin[4(e + f x)] + 4 \sin[5(e + f x)])))/(320 f)$

3.25.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.86, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^2 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 dx$$

$$\downarrow 4446$$

$$a^2 c^2 \int (a \sec^2(e + fx) \tan^4(e + fx) + a \sec(e + fx) \tan^4(e + fx)) dx$$

$$\downarrow 2009$$

$$a^2 c^2 \left(\frac{3a \operatorname{arctanh}(\sin(e + fx))}{8f} + \frac{a \tan^5(e + fx)}{5f} + \frac{a \tan^3(e + fx) \sec(e + fx)}{4f} - \frac{3a \tan(e + fx) \sec(e + fx)}{8f} \right)$$

input $\operatorname{Int}[\operatorname{Sec}[e + f x] * (a + a * \operatorname{Sec}[e + f x])^3 * (c - c * \operatorname{Sec}[e + f x])^2, x]$

output $a^2 c^2 * ((3 * a * \operatorname{ArcTanh}[\sin[e + f x]]) / (8 * f) - (3 * a * \operatorname{Sec}[e + f x] * \operatorname{Tan}[e + f x]) / (8 * f) + (a * \operatorname{Sec}[e + f x] * \operatorname{Tan}[e + f x]^3) / (4 * f) + (a * \operatorname{Tan}[e + f x]^5) / (5 * f))$

3.25.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.25.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 6.60 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.52

method	result
risch	$\frac{ic^2a^3(25e^{9i(fx+e)}+40e^{8i(fx+e)}+10e^{7i(fx+e)}+80e^{4i(fx+e)}-10e^{3i(fx+e)}-25e^{i(fx+e)}+8)}{20f(1+e^{2i(fx+e)})^5} - \frac{3c^2a^3 \ln(e^{i(fx+e)}-i)}{8f}$
parts	$\frac{c^2a^3 \tan(fx+e)}{f} + \frac{c^2a^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f} - \frac{c^2a^3 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} \right)}{f}$
norman	$\frac{\frac{3c^2a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f} - \frac{7c^2a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{2f} - \frac{32c^2a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5f} + \frac{7c^2a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{2f} - \frac{3c^2a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{4f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^5} - \frac{3c^2a^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2f}$
parallelrisch	$- \frac{a^3c^2 \left(\left(\frac{15 \cos(fx+e)}{2} + \frac{15 \cos(3fx+3e)}{4} + \frac{3 \cos(5fx+5e)}{4} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \left(-\frac{15 \cos(fx+e)}{2} - \frac{15 \cos(3fx+3e)}{4} - \frac{3 \cos(5fx+5e)}{4} \right) \right)}{2f(\cos(5fx+5e) + 5 \cos(3fx+3e))} - \frac{c^2a^3 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) + c^2a^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f}$
derivativedivides	$- \frac{c^2a^3 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) + c^2a^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f}$
default	$- \frac{c^2a^3 \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) + c^2a^3 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

$$3.25. \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx$$

output $\frac{1}{20}I^2c^2a^3(25\exp(9I(fx+e))+40\exp(8I(fx+e))+10\exp(7I(fx+e))+80\exp(4I(fx+e))-10\exp(3I(fx+e))-25\exp(I(fx+e))+8)/f/(1+\exp(2I(fx+e)))^5-3/8c^2a^3/f\ln(\exp(I(fx+e))-1)+3/8c^2a^3/f\ln(\exp(I(fx+e))+1)$

3.25.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.54

$$\int \sec(e+fx)(a+a\sec(e+fx))^3(c-c\sec(e+fx))^2 dx$$

$$= \frac{15a^3c^2\cos(fx+e)^5\log(\sin(fx+e)+1)-15a^3c^2\cos(fx+e)^5\log(-\sin(fx+e)+1)+2(8a^3c^2\cos(fx+e)^4-25a^3c^2\cos(fx+e)^3-16a^3c^2\cos(fx+e)^2+10a^3c^2\cos(fx+e)+8a^3c^2)\sin(fx+e)}{80f\cos(fx+e)^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^2,x, algorithm="fricas")`

output $\frac{1}{80}(15a^3c^2\cos(fx+e)^5\log(\sin(fx+e)+1)-15a^3c^2\cos(fx+e)^5\log(-\sin(fx+e)+1)+2(8a^3c^2\cos(fx+e)^4-25a^3c^2\cos(fx+e)^3-16a^3c^2\cos(fx+e)^2+10a^3c^2\cos(fx+e)+8a^3c^2)\sin(fx+e))/(f\cos(fx+e)^5)$

3.25.6 Sympy [F]

$$\int \sec(e+fx)(a+a\sec(e+fx))^3(c-c\sec(e+fx))^2 dx$$

$$= a^3c^2\left(\int \sec(e+fx) dx + \int \sec^2(e+fx) dx + \int (-2\sec^3(e+fx)) dx + \int (-2\sec^4(e+fx)) dx + \int \sec^5(e+fx) dx + \int \sec^6(e+fx) dx\right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**2,x)`

output `a**3*c**2*(Integral(sec(e+f*x), x) + Integral(sec(e+f*x)**2, x) + Integral(-2*sec(e+f*x)**3, x) + Integral(-2*sec(e+f*x)**4, x) + Integral(sec(e+f*x)**5, x) + Integral(sec(e+f*x)**6, x))`

3.25. $\int \sec(e+fx)(a+a\sec(e+fx))^3(c-c\sec(e+fx))^2 dx$

3.25.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(86) = 172.

Time = 0.22 (sec) , antiderivative size = 227, normalized size of antiderivative = 2.41

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx$$

$$= \frac{16(3 \tan(fx + e)^5 + 10 \tan(fx + e)^3 + 15 \tan(fx + e))a^3c^2 - 160(\tan(fx + e)^3 + 3 \tan(fx + e))a^3c}{40f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output `1/240*(16*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^3*c^2 - 160*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c^2 - 15*a^3*c^2*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) + 120*a^3*c^2*(2*sin(f*x + e))/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 240*a^3*c^2*log(sec(f*x + e) + tan(f*x + e)) + 240*a^3*c^2*tan(f*x + e))/f`

3.25.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.69

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx$$

$$= \frac{15a^3c^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|) - 15a^3c^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|) - \frac{2(15a^3c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^9 - 70a^3c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^7 + 128a^3c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 70a^3c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 15a^3c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)^2 - 1}}{40f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `1/40*(15*a^3*c^2*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*a^3*c^2*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(15*a^3*c^2*tan(1/2*f*x + 1/2*e)^9 - 70*a^3*c^2*tan(1/2*f*x + 1/2*e)^7 + 128*a^3*c^2*tan(1/2*f*x + 1/2*e)^5 + 70*a^3*c^2*tan(1/2*f*x + 1/2*e)^3 - 15*a^3*c^2*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^5)/f`

3.25. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx$

3.25.9 Mupad [B] (verification not implemented)

Time = 17.53 (sec) , antiderivative size = 188, normalized size of antiderivative = 2.00

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^2 dx = \frac{3a^3 c^2 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{4f} - \frac{\frac{3a^3 c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{4} - \frac{7a^3 c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{2} + \frac{32a^3 c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{5} + \frac{7a^3 c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{2} - \frac{3a^3 c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1 \right)}$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^2)/cos(e + f*x),x)`output `(3*a^3*c^2*atanh(tan(e/2 + (f*x)/2)))/(4*f) - ((7*a^3*c^2*tan(e/2 + (f*x)/2)^3)/2 + (32*a^3*c^2*tan(e/2 + (f*x)/2)^5)/5 - (7*a^3*c^2*tan(e/2 + (f*x)/2)^7)/2 + (3*a^3*c^2*tan(e/2 + (f*x)/2)^9)/4 - (3*a^3*c^2*tan(e/2 + (f*x)/2))/4)/(f*(5*tan(e/2 + (f*x)/2)^2 - 10*tan(e/2 + (f*x)/2)^4 + 10*tan(e/2 + (f*x)/2)^6 - 5*tan(e/2 + (f*x)/2)^8 + tan(e/2 + (f*x)/2)^10 - 1))`

3.26 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$

3.26.1	Optimal result	271
3.26.2	Mathematica [A] (verified)	271
3.26.3	Rubi [A] (verified)	272
3.26.4	Maple [A] (verified)	273
3.26.5	Fricas [A] (verification not implemented)	274
3.26.6	Sympy [F]	274
3.26.7	Maxima [A] (verification not implemented)	275
3.26.8	Giac [A] (verification not implemented)	275
3.26.9	Mupad [B] (verification not implemented)	276

3.26.1 Optimal result

Integrand size = 30, antiderivative size = 86

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{5a^3c \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{3a^3c \sec(e + fx) \tan(e + fx)}{8f}$$

$$- \frac{a^3c \sec^3(e + fx) \tan(e + fx)}{4f} - \frac{2a^3c \tan^3(e + fx)}{3f}$$

```
output 5/8*a^3*c*arctanh(sin(f*x+e))/f-3/8*a^3*c*sec(f*x+e)*tan(f*x+e)/f-1/4*a^3*c*sec(f*x+e)^3*tan(f*x+e)/f-2/3*a^3*c*tan(f*x+e)^3/f
```

3.26.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.81

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{a^3c(60 \operatorname{arctanh}(\sin(e + fx)) - \sec^4(e + fx)(33 \sin(e + fx) + 16 \sin(2(e + fx)) + 9 \sin(3(e + fx))) - 8 \sin(e + fx))}{96f}$$

```
input Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x]),x]
```


output $(a^3c(60\text{ArcTanh}[\text{Sin}[e + fx]] - \text{Sec}[e + fx]^4(33\text{Sin}[e + fx] + 16\text{Sin}[2(e + fx)] + 9\text{Sin}[3(e + fx)] - 8\text{Sin}[4(e + fx)])))/(96f)$

3.26.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx)) dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow 4446$$

$$-ac \int (a^2 \tan^2(e + fx) \sec^3(e + fx) + 2a^2 \tan^2(e + fx) \sec^2(e + fx) + a^2 \tan^2(e + fx) \sec(e + fx)) dx$$

$$\downarrow 2009$$

$$-ac \left(-\frac{5a^2 \arctanh(\sin(e + fx))}{8f} + \frac{2a^2 \tan^3(e + fx)}{3f} + \frac{a^2 \tan(e + fx) \sec^3(e + fx)}{4f} + \frac{3a^2 \tan(e + fx) \sec(e + fx)}{8f} \right)$$

input $\text{Int}[\text{Sec}[e + fx]*(a + a*\text{Sec}[e + fx])^3*(c - c*\text{Sec}[e + fx]),x]$

output $-(a*c*((-5*a^2*\text{ArcTanh}[\text{Sin}[e + fx]])/(8*f) + (3*a^2*\text{Sec}[e + fx]*\text{Tan}[e + fx])/(8*f) + (a^2*\text{Sec}[e + fx]^3*\text{Tan}[e + fx])/(4*f) + (2*a^2*\text{Tan}[e + fx]^3)/(3*f))$

3.26.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.26.4 Maple [A] (verified)

Time = 4.16 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.29

method	result
derivativedivides	$\frac{-a^3c\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)+2a^3c\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+2}{f}$
default	$\frac{-a^3c\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)+2a^3c\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+2}{f}$
parts	$\frac{a^3c\ln(\sec(fx+e)+\tan(fx+e))}{f} + \frac{2a^3c\tan(fx+e)}{f} + \frac{2a^3c\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)}{f} - \frac{a^3c\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)+2a^3c\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+2}{f}$
norman	$\frac{-\frac{5a^3c\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{4f}-\frac{73a^3c\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{12f}+\frac{55a^3c\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{12f}-\frac{5a^3c\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{4f}}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)^4} - \frac{5a^3c\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{8f} + \frac{5a^3c\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)+2a^3c\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+2}{f}$
risch	$\frac{ia^3c(9e^{7i(fx+e)}+48e^{6i(fx+e)}+33e^{5i(fx+e)}+48e^{4i(fx+e)}-33e^{3i(fx+e)}+16e^{2i(fx+e)}-9e^{i(fx+e)}+16)}{12f(1+e^{2i(fx+e)})^4} - \frac{5a^3c\ln(e^{i(fx+e)}+1)}{8f}$
parallelrisc	$-\frac{3a^3c\left(\left(\frac{10\cos(2fx+2e)}{3}+\frac{5\cos(4fx+4e)}{6}+\frac{5}{2}\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)+\left(-\frac{10\cos(2fx+2e)}{3}-\frac{5\cos(4fx+4e)}{6}-\frac{5}{2}\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)\right)}{4f(3+\cos(4fx+4e)+4\cos(2fx+2e))}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

3.26. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$

output `1/f*(-a^3*c*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln(sec(f*x+e)+tan(f*x+e)))+2*a^3*c*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)+2*a^3*c*tan(f*x+e)+a^3*c*ln(sec(f*x+e)+tan(f*x+e)))`

3.26.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.36

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{15 a^3 c \cos(fx + e)^4 \log(\sin(fx + e) + 1) - 15 a^3 c \cos(fx + e)^4 \log(-\sin(fx + e) + 1) + 2(16 a^3 c \cos(fx + e)^3 - 9 a^3 c \cos(fx + e)^2 - 16 a^3 c \cos(fx + e) - 6 a^3 c) \sin(fx + e)}{48 f \cos(fx + e)^4}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `1/48*(15*a^3*c*cos(f*x + e)^4*log(sin(f*x + e) + 1) - 15*a^3*c*cos(f*x + e)^4*log(-sin(f*x + e) + 1) + 2*(16*a^3*c*cos(f*x + e)^3 - 9*a^3*c*cos(f*x + e)^2 - 16*a^3*c*cos(f*x + e) - 6*a^3*c)*sin(f*x + e))/(f*cos(f*x + e)^4)`

3.26.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= -a^3 c \left(\int (-\sec(e + fx)) dx + \int (-2 \sec^2(e + fx)) dx + \int 2 \sec^4(e + fx) dx + \int \sec^5(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e)),x)`

output `-a**3*c*(Integral(-sec(e + f*x), x) + Integral(-2*sec(e + f*x)**2, x) + Integral(2*sec(e + f*x)**4, x) + Integral(sec(e + f*x)**5, x))`

3.26.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.55

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx =$$

$$\frac{32 (\tan (fx + e)^3 + 3 \tan (fx + e)) a^3 c - 3 a^3 c \left(\frac{2 (3 \sin (fx + e)^3 - 5 \sin (fx + e))}{\sin (fx + e)^4 - 2 \sin (fx + e)^2 + 1} - 3 \log (\sin (fx + e) + 1) + 3 \log (\sin (fx + e) - 1) \right)}{48 f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x, algorithm="maxima")
```

```
output -1/48*(32*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c - 3*a^3*c*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 48*a^3*c*log(sec(f*x + e) + tan(f*x + e)) - 96*a^3*c*tan(f*x + e))/f
```

3.26.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.49

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{15 a^3 c \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + 1 \right| \right) - 15 a^3 c \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(15 a^3 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 - 55 a^3 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 + 73 a^3 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 + 15 a^3 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right)}{\left(\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 - 1 \right)^4}{24 f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x, algorithm="giac")
```

```
output 1/24*(15*a^3*c*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*a^3*c*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(15*a^3*c*tan(1/2*f*x + 1/2*e)^7 - 55*a^3*c*tan(1/2*f*x + 1/2*e)^5 + 73*a^3*c*tan(1/2*f*x + 1/2*e)^3 + 15*a^3*c*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^4)/f
```

3.26.9 Mupad [B] (verification not implemented)

Time = 16.37 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.70

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{5a^3 c \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{4f}$$

$$- \frac{\frac{5ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{4} - \frac{55ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{12} + \frac{73ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{12} + \frac{5ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1 \right)}$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x)))/cos(e + f*x),x)`output `(5*a^3*c*atanh(tan(e/2 + (f*x)/2)))/(4*f) - ((5*a^3*c*tan(e/2 + (f*x)/2))/4 + (73*a^3*c*tan(e/2 + (f*x)/2)^3)/12 - (55*a^3*c*tan(e/2 + (f*x)/2)^5)/12 + (5*a^3*c*tan(e/2 + (f*x)/2)^7)/4)/(f*(6*tan(e/2 + (f*x)/2)^4 - 4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/2)^8 + 1))`

$$3.27 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c-c \sec(e+fx)} dx$$

3.27.1	Optimal result	277
3.27.2	Mathematica [C] (verified)	277
3.27.3	Rubi [A] (verified)	278
3.27.4	Maple [A] (verified)	281
3.27.5	Fricas [A] (verification not implemented)	281
3.27.6	Sympy [F]	282
3.27.7	Maxima [B] (verification not implemented)	282
3.27.8	Giac [A] (verification not implemented)	283
3.27.9	Mupad [B] (verification not implemented)	283

3.27.1 Optimal result

Integrand size = 32, antiderivative size = 100

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c-c \sec(e+fx)} dx = -\frac{15a^3 \arctanh(\sin(e+fx))}{2cf} - \frac{10a^3 \tan(e+fx)}{cf} - \frac{5a^3 \sec(e+fx) \tan(e+fx)}{2cf} - \frac{2a(a+a \sec(e+fx))^2 \tan(e+fx)}{f(c-c \sec(e+fx))}$$

output

```
-15/2*a^3*arctanh(sin(f*x+e))/c/f-10*a^3*tan(f*x+e)/c/f-5/2*a^3*sec(f*x+e)*tan(f*x+e)/c/f-2*a*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))
```

3.27.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.57 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.65

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c-c \sec(e+fx)} dx = -\frac{a^3 \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{7}{2}, \frac{9}{2}, \frac{1}{2}(1+\sec(e+fx))\right) (1+\sec(e+fx))^3 \tan(e+fx)}{7cf \sqrt{2-2\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x]),x]`

output `-1/7*(a^3*Hypergeometric2F1[3/2, 7/2, 9/2, (1 + Sec[e + f*x])/2]*(1 + Sec[e + f*x])^3*Tan[e + f*x])/(c*f*Sqrt[2 - 2*Sec[e + f*x]])`

3.27.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4445, 3042, 4275, 3042, 4254, 24, 4534, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{c-c\sec(e+fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{c-c\csc(e+fx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{5a \int \sec(e+fx)(\sec(e+fx)a+a)^2 dx}{c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{5a \int \csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2 dx}{c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{4275} \\
 & -\frac{5a(2a^2 \int \sec^2(e+fx) dx + \int \sec(e+fx)(\sec^2(e+fx)a^2+a^2) dx)}{c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{f(c-c\sec(e+fx))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{5a\left(2a^2 \int \csc(e+fx+\frac{\pi}{2})^2 dx + \int \csc(e+fx+\frac{\pi}{2})\left(\csc(e+fx+\frac{\pi}{2})^2 a^2+a^2\right) dx\right)}{c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{f(c-c\sec(e+fx))}
 \end{aligned}$$

3.27. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c-c\sec(e+fx)} dx$

$$\begin{aligned}
& \downarrow 4254 \\
& \frac{5a \left(\int \csc \left(e + fx + \frac{\pi}{2} \right) \left(\csc \left(e + fx + \frac{\pi}{2} \right)^2 a^2 + a^2 \right) dx - \frac{2a^2 \int 1d(-\tan(e+fx))}{f} \right)}{\frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{f(c - c \sec(e+fx))}} \\
& \downarrow 24 \\
& \frac{5a \left(\int \csc \left(e + fx + \frac{\pi}{2} \right) \left(\csc \left(e + fx + \frac{\pi}{2} \right)^2 a^2 + a^2 \right) dx + \frac{2a^2 \tan(e+fx)}{f} \right)}{\frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{f(c - c \sec(e+fx))}} \\
& \downarrow 4534 \\
& \frac{5a \left(\frac{3}{2} a^2 \int \sec(e+fx) dx + \frac{2a^2 \tan(e+fx)}{f} + \frac{a^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{\frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{f(c - c \sec(e+fx))}} \\
& \downarrow 3042 \\
& \frac{5a \left(\frac{3}{2} a^2 \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{2a^2 \tan(e+fx)}{f} + \frac{a^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{\frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{f(c - c \sec(e+fx))}} \\
& \downarrow 4257 \\
& \frac{5a \left(\frac{3a^2 \operatorname{arctanh}(\sin(e+fx))}{2f} + \frac{2a^2 \tan(e+fx)}{f} + \frac{a^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{\frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{f(c - c \sec(e+fx))}}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x]),x]`

output `(-2*a*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])) - (5*a*((3*a^2*ArcTanh[Sin[e + f*x]])/(2*f) + (2*a^2*Tan[e + f*x])/f + (a^2*Sec[e + f*x]*Tan[e + f*x])/(2*f)))/c`

3.27.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4275 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^2, x_Symbol] := Simp[2*a*(b/d) Int[(d*Csc[e + f*x])^(n + 1), x], x] + Int[(d*Csc[e + f*x])^n*(a^2 + b^2*Csc[e + f*x]^2), x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`
- rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.27.4 Maple [A] (verified)

Time = 1.11 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.06

method	result
parallelrisch	$\frac{15 \left((1 + \cos(2fx + 2e)) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) + (-1 - \cos(2fx + 2e)) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right) - \frac{14 \left(\cos(fx + e) - \frac{12 \cos(2fx + 2e) - 11}{7} \right) \cot \left(\frac{1}{2} fx + \frac{1}{2} e \right)}{15}}{2cf(1 + \cos(2fx + 2e))}$
derivativedivides	$\frac{8a^3 \left(-\frac{1}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^2} + \frac{7}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)} + \frac{15 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)}{16} + \frac{1}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)^2} + \frac{7}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)} - \frac{15 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)}{16} \right)}{fc}$
default	$\frac{8a^3 \left(-\frac{1}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^2} + \frac{7}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)} + \frac{15 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)}{16} + \frac{1}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)^2} + \frac{7}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)} - \frac{15 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)}{16} \right)}{fc}$
risch	$\frac{ia^3 (17e^{4i(fx+e)} - 9e^{3i(fx+e)} + 39e^{2i(fx+e)} - 7e^{i(fx+e)} + 24)}{fc(e^{i(fx+e)} - 1)(1 + e^{2i(fx+e)})^2} + \frac{15a^3 \ln(e^{i(fx+e)} - i)}{2cf} - \frac{15a^3 \ln(e^{i(fx+e)} + i)}{2cf}$
norman	$\frac{-\frac{8a^3}{cf} + \frac{33a^3 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^2}{cf} - \frac{40a^3 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^4}{cf} + \frac{15a^3 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^6}{cf}}{\left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^3 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)} + \frac{15a^3 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)}{2cf} - \frac{15a^3 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)}{2cf}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `15/2*((1+cos(2*f*x+2*e))*ln(tan(1/2*f*x+1/2*e)-1)+(-1-cos(2*f*x+2*e))*ln(tan(1/2*f*x+1/2*e)+1)-14/15*(cos(f*x+e)-12/7*cos(2*f*x+2*e)-11/7)*cot(1/2*f*x+1/2*e))*a^3/c/f/(1+cos(2*f*x+2*e))`

3.27.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.25

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{c - c \sec(e + fx)} dx = \frac{15 a^3 \cos^2 (fx + e) \log (\sin (fx + e) + 1) \sin (fx + e) - 15 a^3 \cos^2 (fx + e) \log (-\sin (fx + e) + 1) \sin (fx + e)}{4 c f \cos (fx + e)^2 \sin (fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x, algorithm="fricas")`

output
$$\frac{-1/4*(15*a^3*\cos(f*x + e)^2*\log(\sin(f*x + e) + 1)*\sin(f*x + e) - 15*a^3*\cos(f*x + e)^2*\log(-\sin(f*x + e) + 1)*\sin(f*x + e) - 48*a^3*\cos(f*x + e)^3 - 34*a^3*\cos(f*x + e)^2 + 16*a^3*\cos(f*x + e) + 2*a^3)/(c*f*\cos(f*x + e)^2*\sin(f*x + e))}{c}$$

3.27.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{c - c \sec(e + fx)} dx$$

$$= \frac{a^3 \left(\int \frac{\sec(e+fx)}{\sec(e+fx)-1} dx + \int \frac{3\sec^2(e+fx)}{\sec(e+fx)-1} dx + \int \frac{3\sec^3(e+fx)}{\sec(e+fx)-1} dx + \int \frac{\sec^4(e+fx)}{\sec(e+fx)-1} dx \right)}{c}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e)),x)`

output
$$-a**3*(Integral(\sec(e + f*x)/(\sec(e + f*x) - 1), x) + Integral(3*\sec(e + f*x)**2/(\sec(e + f*x) - 1), x) + Integral(3*\sec(e + f*x)**3/(\sec(e + f*x) - 1), x) + Integral(\sec(e + f*x)**4/(\sec(e + f*x) - 1), x))/c$$

3.27.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 387 vs. $2(97) = 194$.

Time = 0.21 (sec) , antiderivative size = 387, normalized size of antiderivative = 3.87

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{c - c \sec(e + fx)} dx =$$

$$a^3 \left(\frac{2 \left(\frac{5 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{2 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} - 1 \right)}{\frac{c \sin(fx+e)}{\cos(fx+e)+1} - \frac{2c \sin^3(fx+e)}{(\cos(fx+e)+1)^3} + \frac{c \sin^5(fx+e)}{(\cos(fx+e)+1)^5}} + \frac{3 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{c} - \frac{3 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{c} \right) + 6a^3 \left(\frac{\frac{3 \sin(fx+e)}{(\cos(fx+e)+1)}}{\frac{c \sin(fx+e)}{\cos(fx+e)+1} - \frac{c}{(\cos(fx+e)+1)}} \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x, algorithm="maxima")`

output
$$-1/2*(a^3*(2*(5*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 2*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 - 1)/(c*\sin(f*x + e)/(\cos(f*x + e) + 1) - 2*c*\sin(f*x + e)^3/(\cos(f*x + e) + 1)^3 + c*\sin(f*x + e)^5/(\cos(f*x + e) + 1)^5) + 3*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/c - 3*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/c) + 6*a^3*((3*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 1)/(c*\sin(f*x + e)/(\cos(f*x + e) + 1) - c*\sin(f*x + e)^3/(\cos(f*x + e) + 1)^3) + \log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/c - \log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/c) + 6*a^3*(\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/c - \log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/c - (\cos(f*x + e) + 1)/(c*\sin(f*x + e))) - 2*a^3*(\cos(f*x + e) + 1)/(c*\sin(f*x + e)))/f$$

3.27.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.18

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{c - c \sec(e + fx)} dx =$$

$$\frac{15a^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{c} - \frac{15a^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{c} - \frac{16a^3}{c \tan(\frac{1}{2}fx + \frac{1}{2}e)} - \frac{2(7a^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 9a^3 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)^2 c}$$

$$2f$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x, algorithm="giac")`

output
$$-1/2*(15*a^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/c - 15*a^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/c - 16*a^3/(c*\tan(1/2*f*x + 1/2*e)) - 2*(7*a^3*\tan(1/2*f*x + 1/2*e)^3 - 9*a^3*\tan(1/2*f*x + 1/2*e))/((\tan(1/2*f*x + 1/2*e)^2 - 1)^2*c))/f$$

3.27.9 Mupad [B] (verification not implemented)

Time = 14.68 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.05

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{c - c \sec(e + fx)} dx$$

$$= \frac{15a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 25a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 8a^3}{f \left(c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 - 2c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + c \tan\left(\frac{e}{2} + \frac{fx}{2}\right) \right)} - \frac{15a^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{cf}$$

3.27.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c-c\sec(e+fx)} dx$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))),x)`

output `(15*a^3*tan(e/2 + (f*x)/2)^4 - 25*a^3*tan(e/2 + (f*x)/2)^2 + 8*a^3)/(f*(c*tan(e/2 + (f*x)/2) - 2*c*tan(e/2 + (f*x)/2)^3 + c*tan(e/2 + (f*x)/2)^5) - (15*a^3*atanh(tan(e/2 + (f*x)/2)))/(c*f)`

3.28 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^2} dx$

3.28.1	Optimal result	285
3.28.2	Mathematica [C] (verified)	285
3.28.3	Rubi [A] (verified)	286
3.28.4	Maple [A] (verified)	289
3.28.5	Fricas [A] (verification not implemented)	289
3.28.6	Sympy [F]	290
3.28.7	Maxima [B] (verification not implemented)	290
3.28.8	Giac [A] (verification not implemented)	291
3.28.9	Mupad [B] (verification not implemented)	291

3.28.1 Optimal result

Integrand size = 32, antiderivative size = 119

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^2} dx = \frac{5a^3 \operatorname{arctanh}(\sin(e+fx))}{c^2 f} + \frac{5a^3 \tan(e+fx)}{c^2 f} - \frac{2a(a+a \sec(e+fx))^2 \tan(e+fx)}{3f(c-c \sec(e+fx))^2} + \frac{10(a^3+a^3 \sec(e+fx)) \tan(e+fx)}{3f(c^2-c^2 \sec(e+fx))}$$

```
output 5*a^3*arctanh(sin(f*x+e))/c^2/f+5*a^3*tan(f*x+e)/c^2/f-2/3*a*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^2+10/3*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/f/(c^2-c^2*sec(f*x+e))
```

3.28.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.66 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.55

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^2} dx = -\frac{a^3 \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{1}{2}(1+\sec(e+fx))\right) (1+\sec(e+fx))^3 \tan(e+fx)}{14c^2 f \sqrt{2-2 \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^2,x]`

output `-1/14*(a^3*Hypergeometric2F1[5/2, 7/2, 9/2, (1 + Sec[e + f*x])/2]*(1 + Sec[e + f*x])^3*Tan[e + f*x])/(c^2*f*Sqrt[2 - 2*Sec[e + f*x]])`

3.28.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4445, 3042, 4445, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c-c\sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^2} dx$$

↓ 4445

$$-\frac{5a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{c-c\sec(e+fx)} dx}{3c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{3f(c-c\sec(e+fx))^2}$$

↓ 3042

$$-\frac{5a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{c-c\csc(e+fx+\frac{\pi}{2})} dx}{3c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{3f(c-c\sec(e+fx))^2}$$

↓ 4445

$$-\frac{5a \left(-\frac{3a \int \sec(e+fx)(\sec(e+fx)a+a) dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \right)}{3c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{3f(c-c\sec(e+fx))^2}$$

↓ 3042

$$-\frac{5a \left(-\frac{3a \int \csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a) dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \right)}{3c} - \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{3f(c-c\sec(e+fx))^2}$$

3.28. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^2} dx$

$$\begin{array}{c}
\downarrow 4274 \\
5a \left(-\frac{3a(a \int \sec^2(e+fx)dx + a \int \sec(e+fx)dx)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \right) \\
\hline
\frac{3c}{2a \tan(e+fx)(a \sec(e+fx) + a)^2} \\
\frac{3f(c - c\sec(e+fx))^2}{} \\
\downarrow 3042 \\
5a \left(-\frac{3a(a \int \csc(e+fx+\frac{\pi}{2})dx + a \int \csc(e+fx+\frac{\pi}{2})^2 dx)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \right) \\
\hline
\frac{3c}{2a \tan(e+fx)(a \sec(e+fx) + a)^2} \\
\frac{3f(c - c\sec(e+fx))^2}{} \\
\downarrow 4254 \\
5a \left(-\frac{3a(a \int \csc(e+fx+\frac{\pi}{2})dx - \frac{a \int 1d(-\tan(e+fx))}{f})}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \right) \\
\hline
\frac{3c}{2a \tan(e+fx)(a \sec(e+fx) + a)^2} \\
\frac{3f(c - c\sec(e+fx))^2}{} \\
\downarrow 24 \\
5a \left(-\frac{3a(a \int \csc(e+fx+\frac{\pi}{2})dx + \frac{a \tan(e+fx)}{f})}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} \right) \\
\hline
\frac{3c}{2a \tan(e+fx)(a \sec(e+fx) + a)^2} \\
\frac{3f(c - c\sec(e+fx))^2}{} \\
\downarrow 4257 \\
5a \left(-\frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{f(c-c\sec(e+fx))} - \frac{3a \left(\frac{a \operatorname{arctanh}(\sin(e+fx))}{f} + \frac{a \tan(e+fx)}{f} \right)}{c} \right) \\
\hline
\frac{3c}{2a \tan(e+fx)(a \sec(e+fx) + a)^2} \\
\frac{3f(c - c\sec(e+fx))^2}{}
\end{array}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^2,x]`

output `(-2*a*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(3*f*(c - c*Sec[e + f*x])^2) - (5*a*((-2*(a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])) - (3*a*((a*ArcTanh[Sin[e + f*x]])/f + (a*Tan[e + f*x])/f))/c))/(3*c)`

3.28. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^2} dx$

3.28.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.28.4 Maple [A] (verified)

Time = 1.14 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.82

method	result
derivativedivides	$4a^3 \left(-\frac{1}{4(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)} + \frac{5 \ln(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)}{4} - \frac{1}{4(\tan(\frac{fx}{2} + \frac{e}{2}) - 1)} - \frac{5 \ln(\tan(\frac{fx}{2} + \frac{e}{2}) - 1)}{4} - \frac{1}{3 \tan(\frac{fx}{2} + \frac{e}{2})^3} - \frac{2}{\tan(\frac{fx}{2} + \frac{e}{2})} \right) \frac{1}{f c^2}$
default	$4a^3 \left(-\frac{1}{4(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)} + \frac{5 \ln(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)}{4} - \frac{1}{4(\tan(\frac{fx}{2} + \frac{e}{2}) - 1)} - \frac{5 \ln(\tan(\frac{fx}{2} + \frac{e}{2}) - 1)}{4} - \frac{1}{3 \tan(\frac{fx}{2} + \frac{e}{2})^3} - \frac{2}{\tan(\frac{fx}{2} + \frac{e}{2})} \right) \frac{1}{f c^2}$
parallelrisch	$5a^3 \left(\ln(\tan(\frac{fx}{2} + \frac{e}{2}) - 1) \cos(fx+e) - \ln(\tan(\frac{fx}{2} + \frac{e}{2}) + 1) \cos(fx+e) + \frac{17(\cos(fx+e) - \frac{23 \cos(2fx+2e)}{68} - \frac{29}{68}) \operatorname{csc}(\frac{fx}{2} + \frac{e}{2})}{15} \right) \frac{1}{f \cos(fx+e) c^2}$
risch	$-\frac{2ia^3(12e^{4i(fx+e)} - 51e^{3i(fx+e)} + 41e^{2i(fx+e)} - 57e^{i(fx+e)} + 23)}{3f c^2(1+e^{2i(fx+e)})(e^{i(fx+e)}-1)^3} - \frac{5a^3 \ln(e^{i(fx+e)}-i)}{c^2 f} + \frac{5a^3 \ln(e^{i(fx+e)}+i)}{c^2 f}$
norman	$\frac{\frac{4a^3}{3cf} + \frac{4a^3 \tan(\frac{fx}{2} + \frac{e}{2})^2}{cf} - \frac{22a^3 \tan(\frac{fx}{2} + \frac{e}{2})^4}{cf} + \frac{80a^3 \tan(\frac{fx}{2} + \frac{e}{2})^6}{3cf} - \frac{10a^3 \tan(\frac{fx}{2} + \frac{e}{2})^8}{cf}}{\left(\tan(\frac{fx}{2} + \frac{e}{2})^2 - 1\right)^3 c \tan(\frac{fx}{2} + \frac{e}{2})^3} - \frac{5a^3 \ln(\tan(\frac{fx}{2} + \frac{e}{2}) - 1)}{c^2 f} + 5a^3 \ln(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `4/f*a^3/c^2*(-1/4/(tan(1/2*f*x+1/2*e)+1)+5/4*ln(tan(1/2*f*x+1/2*e)+1)-1/4/(tan(1/2*f*x+1/2*e)-1)-5/4*ln(tan(1/2*f*x+1/2*e)-1)-1/3/tan(1/2*f*x+1/2*e)^3-2/tan(1/2*f*x+1/2*e))`

3.28.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.39

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^2} dx = \frac{46a^3 \cos^3(fx+e) - 22a^3 \cos^2(fx+e) - 62a^3 \cos(fx+e) + 6a^3 - 15(a^3 \cos(fx+e)^2 - a^3 \cos(fx+e))}{6(c^2 f \cos(fx+e))}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x, algorithm="fracas")`

output
$$\frac{-1/6*(46*a^3*\cos(f*x + e)^3 - 22*a^3*\cos(f*x + e)^2 - 62*a^3*\cos(f*x + e) + 6*a^3 - 15*(a^3*\cos(f*x + e)^2 - a^3*\cos(f*x + e))*\log(\sin(f*x + e) + 1) * \sin(f*x + e) + 15*(a^3*\cos(f*x + e)^2 - a^3*\cos(f*x + e))*\log(-\sin(f*x + e) + 1)*\sin(f*x + e)}{(c^2*f*\cos(f*x + e)^2 - c^2*f*\cos(f*x + e))*\sin(f*x + e)}$$

3.28.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^2} dx$$

$$= \frac{a^3 \left(\int \frac{\sec(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx + \int \frac{3\sec^2(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx + \int \frac{3\sec^3(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx + \int \frac{\sec^4(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx \right)}{c^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**2,x)`

output `a**3*(Integral(sec(e + f*x)/(sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**2/(sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**3/(sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x))/c**2`

3.28.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 349 vs. $2(117) = 234$.

Time = 0.23 (sec) , antiderivative size = 349, normalized size of antiderivative = 2.93

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^2} dx =$$

$$\frac{a^3 \left(\frac{14 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{27 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + 1 - \frac{12 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{c^2} + \frac{12 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{c^2} \right) - 3a^3 \left(\frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{c^2} \right)}{c^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output
$$\begin{aligned} & -1/6*(a^3*((14*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 27*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 + 1)/(c^2*\sin(f*x + e)^3/(\cos(f*x + e) + 1)^3 - c^2*\sin(f*x + e)^5/(\cos(f*x + e) + 1)^5) - 12*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/c^2 + 12*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/c^2 - 3*a^3*(6*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/c^2 - 6*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/c^2 - (9*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 + 1)*(\cos(f*x + e) + 1)^3/(c^2*\sin(f*x + e)^3)) + 3*a^3*(3*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 + 1)*(\cos(f*x + e) + 1)^3/(c^2*\sin(f*x + e)^3) - a^3*(3*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 1)*(\cos(f*x + e) + 1)^3/(c^2*\sin(f*x + e)^3))/f \end{aligned}$$

3.28.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.97

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^2} dx$$

$$= \frac{15 a^3 \log(|\tan(\frac{1}{2} f x + \frac{1}{2} e) + 1|)}{c^2} - \frac{15 a^3 \log(|\tan(\frac{1}{2} f x + \frac{1}{2} e) - 1|)}{c^2} - \frac{6 a^3 \tan(\frac{1}{2} f x + \frac{1}{2} e)}{(\tan(\frac{1}{2} f x + \frac{1}{2} e)^2 - 1) c^2} - \frac{4 (6 a^3 \tan(\frac{1}{2} f x + \frac{1}{2} e)^2 + a^3)}{c^2 \tan(\frac{1}{2} f x + \frac{1}{2} e)^3}$$

$3 f$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output
$$\begin{aligned} & 1/3*(15*a^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/c^2 - 15*a^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/c^2 - 6*a^3*\tan(1/2*f*x + 1/2*e)/((\tan(1/2*f*x + 1/2*e)^2 - 1)*c^2) - 4*(6*a^3*\tan(1/2*f*x + 1/2*e)^2 + a^3)/(c^2*\tan(1/2*f*x + 1/2*e)^3))/f \end{aligned}$$

3.28.9 Mupad [B] (verification not implemented)

Time = 13.37 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.78

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^2} dx = \frac{10 a^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{c^2 f}$$

$$+ \frac{-10 a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + \frac{20 a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2}{3} + \frac{4 a^3}{3}}{c^2 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^2),x)`

output `(10*a^3*atanh(tan(e/2 + (f*x)/2)))/(c^2*f) + ((20*a^3*tan(e/2 + (f*x)/2)^2)/3 - 10*a^3*tan(e/2 + (f*x)/2)^4 + (4*a^3)/3)/(c^2*f*tan(e/2 + (f*x)/2)^3*(tan(e/2 + (f*x)/2)^2 - 1))`

3.29 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^3} dx$

3.29.1	Optimal result	293
3.29.2	Mathematica [A] (verified)	293
3.29.3	Rubi [A] (verified)	294
3.29.4	Maple [A] (verified)	296
3.29.5	Fricas [A] (verification not implemented)	297
3.29.6	Sympy [F]	297
3.29.7	Maxima [B] (verification not implemented)	298
3.29.8	Giac [A] (verification not implemented)	298
3.29.9	Mupad [B] (verification not implemented)	299

3.29.1 Optimal result

Integrand size = 32, antiderivative size = 132

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^3} dx = -\frac{a^3 \operatorname{arctanh}(\sin(e+fx))}{c^3 f} - \frac{2a(a+a \sec(e+fx))^2 \tan(e+fx)}{5f(c-c \sec(e+fx))^3} + \frac{2(a^3+a^3 \sec(e+fx)) \tan(e+fx)}{3cf(c-c \sec(e+fx))^2} - \frac{2a^3 \tan(e+fx)}{f(c^3-c^3 \sec(e+fx))}$$

output

```
-a^3*arctanh(sin(f*x+e))/c^3/f-2/5*a*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*
sec(f*x+e))^3+2/3*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^2-2
*a^3*tan(f*x+e)/f/(c^3-c^3*sec(f*x+e))
```

3.29.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.05

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^3} dx = \frac{a^3 \left(-\frac{26 \cot(\frac{1}{2}(e+fx))}{15f} + \frac{2 \cot(\frac{1}{2}(e+fx)) \operatorname{csc}^2(\frac{1}{2}(e+fx))}{15f} - \frac{2 \cot(\frac{1}{2}(e+fx)) \operatorname{csc}^4(\frac{1}{2}(e+fx))}{5f} - \frac{\log(\cos(\frac{1}{2}(e+fx))-\sin(\frac{1}{2}(e+fx)))}{f} \right)}{c^3} + \dots$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^3,x]`

output $-\left(\frac{a^3(-26\cot[(e + fx)/2])}{15f} + \frac{2\cot[(e + fx)/2]*\csc[(e + fx)/2]^2}{15f} - \frac{2\cot[(e + fx)/2]*\csc[(e + fx)/2]^4}{5f} - \log\left[\frac{\cos[(e + fx)/2] - \sin[(e + fx)/2]}{f} + \log\left[\frac{\cos[(e + fx)/2] + \sin[(e + fx)/2]}{f}\right]\right)/c^3\right)$

3.29.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4445, 3042, 4445, 3042, 4445, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(a \sec(e + fx) + a)^3}{(c - c \sec(e + fx))^3} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3}{\left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3} dx$$

↓ 4445

$$-\frac{a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{(c-c\sec(e+fx))^2} dx}{c} - \frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{5f(c - c \sec(e+fx))^3}$$

↓ 3042

$$-\frac{a \int \frac{\csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)^2}{\left(c-c\csc\left(e+fx+\frac{\pi}{2}\right)\right)^2} dx}{c} - \frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{5f(c - c \sec(e+fx))^3}$$

↓ 4445

$$-\frac{a \left(-\frac{a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{c-c\sec(e+fx)} dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^2} \right)}{c} - \frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{5f(c - c \sec(e+fx))^3}$$

↓ 3042

3.29. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^3} dx$

$$\begin{aligned}
 & a \left(\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{c-c\csc(e+fx+\frac{\pi}{2})} dx}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^2} \right) \\
 & \frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{5f(c-c\sec(e+fx))^3} \\
 & \quad \downarrow \text{4445} \\
 & a \left(\frac{a \left(-\frac{a \int \sec(e+fx) dx}{c} - \frac{2a \tan(e+fx)}{f(c-c\sec(e+fx))} \right)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^2} \right) \\
 & \frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{5f(c-c\sec(e+fx))^3} \\
 & \quad \downarrow \text{3042} \\
 & a \left(\frac{a \left(-\frac{a \int \csc(e+fx+\frac{\pi}{2}) dx}{c} - \frac{2a \tan(e+fx)}{f(c-c\sec(e+fx))} \right)}{c} - \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^2} \right) \\
 & \frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{5f(c-c\sec(e+fx))^3} \\
 & \quad \downarrow \text{4257} \\
 & a \left(-\frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^2} - \frac{a \left(-\frac{a \operatorname{arctanh}(\sin(e+fx))}{cf} - \frac{2a \tan(e+fx)}{f(c-c\sec(e+fx))} \right)}{c} \right) \\
 & \frac{2a \tan(e+fx)(a \sec(e+fx) + a)^2}{5f(c-c\sec(e+fx))^3}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^3,x]`

output `(-2*a*(a + a*Sec[e + f*x])^2*Tan[e + f*x]/(5*f*(c - c*Sec[e + f*x])^3) - (a*((-2*(a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(3*f*(c - c*Sec[e + f*x])^2) - (a*(-((a*ArcTanh[Sin[e + f*x]])/(c*f)) - (2*a*Tan[e + f*x])/(f*(c - c*Sec[e + f*x]))))/c))/c`

3.29.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.29.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.59

method	result
derivativedivides	$\frac{2a^3 \left(-\frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} + \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{f c^3} \right)}{f c^3}$
default	$\frac{2a^3 \left(-\frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} + \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{f c^3} \right)}{f c^3}$
parallelrisc	$\frac{a^3 \left(6 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 + 10 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 15 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - 15 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) + 30 \cot\left(\frac{fx}{2} + \frac{e}{2}\right) \right)}{15c^3 f}$
risc	$\frac{4ia^3 (15e^{4i(fx+e)} - 30e^{3i(fx+e)} + 100e^{2i(fx+e)} - 50e^{i(fx+e)} + 13)}{15f c^3 (e^{i(fx+e)} - 1)^5} + \frac{a^3 \ln(e^{i(fx+e)} - i)}{c^3 f} - \frac{a^3 \ln(e^{i(fx+e)} + i)}{c^3 f}$
norman	$\frac{-\frac{2a^3}{5cf} + \frac{8a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{15cf} - \frac{6a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{5cf} + \frac{22a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{5cf} - \frac{16a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{3cf} + \frac{2a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{10}}{cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^3 c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} + \frac{a^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{c^3 f}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

$$3.29. \int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^3} dx$$

output $\frac{2f a^3 c^3 (-1/2 \ln(\tan(1/2 f x + 1/2 e) + 1) + 1/5 \tan(1/2 f x + 1/2 e)^5 + 1/3 \tan(1/2 f x + 1/2 e)^3 + 1/\tan(1/2 f x + 1/2 e) + 1/2 \ln(\tan(1/2 f x + 1/2 e) - 1))}{c^3}$

3.29.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.33

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^3} dx = \frac{52 a^3 \cos(fx + e)^3 - 44 a^3 \cos(fx + e)^2 - 4 a^3 \cos(fx + e) + 92 a^3 - 15 (a^3 \cos(fx + e)^2 - 2 a^3 \cos(fx + e) + a^3) \log(\sin(fx + e) + 1) \sin(fx + e) + 15 (a^3 \cos(fx + e)^2 - 2 a^3 \cos(fx + e) + a^3) \log(-\sin(fx + e) + 1) \sin(fx + e)}{30 (c^3 f \cos(fx + e))}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x, algorithm="fricas")`

output $\frac{1/30 * (52 * a^3 * \cos(f * x + e)^3 - 44 * a^3 * \cos(f * x + e)^2 - 4 * a^3 * \cos(f * x + e) + 92 * a^3 - 15 * (a^3 * \cos(f * x + e)^2 - 2 * a^3 * \cos(f * x + e) + a^3) * \log(\sin(f * x + e) + 1) * \sin(f * x + e) + 15 * (a^3 * \cos(f * x + e)^2 - 2 * a^3 * \cos(f * x + e) + a^3) * \log(-\sin(f * x + e) + 1) * \sin(f * x + e))}{(c^3 * f * \cos(f * x + e)^2 - 2 * c^3 * f * \cos(f * x + e) + c^3 * f) * \sin(f * x + e)}$

3.29.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^3} dx = \frac{a^3 \left(\int \frac{\sec(e+fx)}{\sec^3(e+fx) - 3 \sec^2(e+fx) + 3 \sec(e+fx) - 1} dx + \int \frac{3 \sec^2(e+fx)}{\sec^3(e+fx) - 3 \sec^2(e+fx) + 3 \sec(e+fx) - 1} dx + \int \frac{3 \sec^5(e+fx)}{\sec^3(e+fx) - 3 \sec^2(e+fx) + 3 \sec(e+fx) - 1} dx \right)}{c^3}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**3,x)`

output $-a^{**3} * (\text{Integral}(\sec(e + f*x) / (\sec(e + f*x)^{**3} - 3 * \sec(e + f*x)^{**2} + 3 * \sec(e + f*x) - 1), x) + \text{Integral}(3 * \sec(e + f*x)^{**2} / (\sec(e + f*x)^{**3} - 3 * \sec(e + f*x)^{**2} + 3 * \sec(e + f*x) - 1), x) + \text{Integral}(3 * \sec(e + f*x)^{**3} / (\sec(e + f*x)^{**3} - 3 * \sec(e + f*x)^{**2} + 3 * \sec(e + f*x) - 1), x) + \text{Integral}(\sec(e + f*x)^{**4} / (\sec(e + f*x)^{**3} - 3 * \sec(e + f*x)^{**2} + 3 * \sec(e + f*x) - 1), x)) / c^{**3}$

3.29. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^3} dx$

3.29.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 309 vs. $2(131) = 262$.

Time = 0.23 (sec) , antiderivative size = 309, normalized size of antiderivative = 2.34

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^3} dx =$$

$$a^3 \left(\frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)+1}{c^3} - \frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)-1}{c^3} - \frac{\left(\frac{20 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{105 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + 3\right)(\cos(fx+e)+1)^5}{c^3 \sin(fx+e)^5} \right) - \frac{3a^3 \left(\frac{10 \sin(fx+e)}{\cos(fx+e)+1}\right)}{c^3 \sin(fx+e)^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `-1/60*(a^3*(60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/c^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/c^3 - (20*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 105*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 3)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5)) - 3*a^3*(10*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 15*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 3)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5) + a^3*(10*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 15*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 3)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5) + 9*a^3*(5*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 1)*(cos(f*x + e) + 1)^5/(c^3*sin(f*x + e)^5))/f`

3.29.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.77

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^3} dx =$$

$$\frac{15a^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{c^3} - \frac{15a^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{c^3} - \frac{2\left(15a^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^4 + 5a^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 + 3a^3\right)}{c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5}$$

15f

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `-1/15*(15*a^3*log(abs(tan(1/2*f*x + 1/2*e) + 1))/c^3 - 15*a^3*log(abs(tan(1/2*f*x + 1/2*e) - 1))/c^3 - 2*(15*a^3*tan(1/2*f*x + 1/2*e)^4 + 5*a^3*tan(1/2*f*x + 1/2*e)^2 + 3*a^3)/(c^3*tan(1/2*f*x + 1/2*e)^5))/f`

3.29. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^3} dx$

3.29.9 Mupad [B] (verification not implemented)

Time = 13.10 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.59

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^3} dx = \frac{2a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + \frac{2a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2}{3} + \frac{2a^3}{5}}{c^3 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5} - \frac{2a^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{c^3 f}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^3),x)`output `((2*a^3*tan(e/2 + (f*x)/2)^2)/3 + 2*a^3*tan(e/2 + (f*x)/2)^4 + (2*a^3)/5)/(c^3*f*tan(e/2 + (f*x)/2)^5) - (2*a^3*atanh(tan(e/2 + (f*x)/2)))/(c^3*f)`

$$3.30 \quad \int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^4} dx$$

3.30.1	Optimal result	300
3.30.2	Mathematica [A] (verified)	300
3.30.3	Rubi [A] (verified)	301
3.30.4	Maple [A] (verified)	302
3.30.5	Fricas [B] (verification not implemented)	302
3.30.6	Sympy [F]	303
3.30.7	Maxima [B] (verification not implemented)	303
3.30.8	Giac [A] (verification not implemented)	304
3.30.9	Mupad [B] (verification not implemented)	304

3.30.1 Optimal result

Integrand size = 32, antiderivative size = 38

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^4} dx = -\frac{(a+a\sec(e+fx))^3 \tan(e+fx)}{7f(c-c\sec(e+fx))^4}$$

output `-1/7*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^4`

3.30.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^4} dx = -\frac{a^3 \cot^7\left(\frac{1}{2}(e+fx)\right)}{7c^4 f}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^4,x]`

output `-1/7*(a^3*Cot[(e + f*x)/2]^7)/(c^4*f)`

3.30.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c-c\sec(e+fx))^4} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx$$

↓ 4438

$$-\frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{7f(c-c\sec(e+fx))^4}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^4,x]`

output `-1/7*((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^4)`

3.30.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^n, x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.30.4 Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.61

method	result	size
derivativedivides	$-\frac{a^3}{7f c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$	23
default	$-\frac{a^3}{7f c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$	23
parallelrisc	$-\frac{a^3}{7f c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$	23
risc	$\frac{2ia^3(7e^{6i(fx+e)} + 35e^{4i(fx+e)} + 21e^{2i(fx+e)} + 1)}{7f c^4 (e^{i(fx+e)} - 1)^7}$	61
norman	$\frac{\frac{a^3}{7cf} - \frac{3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{7cf} + \frac{3a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{7cf} - \frac{a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{7cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^3 c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$	109

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x,method=_RETURNVERBOSE)`

output `-1/7/f*a^3/c^4/tan(1/2*f*x+1/2*e)^7`

3.30.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(37) = 74$.

Time = 0.26 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.92

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^4} dx$$

$$= \frac{a^3 \cos^4(fx+e) + 4a^3 \cos^3(fx+e) + 6a^3 \cos^2(fx+e) + 4a^3 \cos(fx+e) + a^3}{7(c^4 f \cos^3(fx+e) - 3c^4 f \cos^2(fx+e) + 3c^4 f \cos(fx+e) - c^4 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x, algorithm="fricas")`

output `1/7*(a^3*cos(f*x + e)^4 + 4*a^3*cos(f*x + e)^3 + 6*a^3*cos(f*x + e)^2 + 4*a^3*cos(f*x + e) + a^3)/((c^4*f*cos(f*x + e)^3 - 3*c^4*f*cos(f*x + e)^2 + 3*c^4*f*cos(f*x + e) - c^4*f)*sin(f*x + e))`

3.30. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^4} dx$

3.30.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^4} dx$$

$$= \frac{a^3 \left(\int \frac{\sec(e+fx)}{\sec^4(e+fx)-4\sec^3(e+fx)+6\sec^2(e+fx)-4\sec(e+fx)+1} dx + \int \frac{3\sec^2(e+fx)}{\sec^4(e+fx)-4\sec^3(e+fx)+6\sec^2(e+fx)-4\sec(e+fx)+1} dx \right)}{c^4}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**4,x)`

output `a**3*(Integral(sec(e + f*x)/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**2/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**3/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x)**4 - 4*sec(e + f*x)**3 + 6*sec(e + f*x)**2 - 4*sec(e + f*x) + 1), x))/c**4`

3.30.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 356 vs. 2(37) = 74.

Time = 0.23 (sec) , antiderivative size = 356, normalized size of antiderivative = 9.37

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^4} dx =$$

$$\frac{a^3 \left(\frac{21 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} + \frac{35 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{35 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} + 5 \right) (\cos(fx+e)+1)^7}{c^4 \sin^7(fx+e)} - \frac{a^3 \left(\frac{21 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} + \frac{35 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} - \frac{105 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - 15 \right) (\cos(fx+e)+1)^7}{c^4 \sin^7(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output
$$\begin{aligned} & -1/280*(a^3*(21*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 + 35*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 + 35*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6 + 5)*(\cos(f*x + e) + 1)^7/(c^4*\sin(f*x + e)^7) - a^3*(21*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 + 35*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 - 105*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6 - 15)*(\cos(f*x + e) + 1)^7/(c^4*\sin(f*x + e)^7) - a^3*(21*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 35*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 + 35*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6 - 5)*(\cos(f*x + e) + 1)^7/(c^4*\sin(f*x + e)^7) + a^3*(21*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 35*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 - 105*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6 + 15)*(\cos(f*x + e) + 1)^7/(c^4*\sin(f*x + e)^7))/f \end{aligned}$$

3.30.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^4} dx = -\frac{a^3}{7c^4 f \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^7}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output
$$-1/7*a^3/(c^4*f*\tan(1/2*f*x + 1/2*e)^7)$$

3.30.9 Mupad [B] (verification not implemented)

Time = 12.85 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^4} dx = -\frac{a^3 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{7c^4 f}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^4),x)`

output
$$-(a^3*\cot(e/2 + (f*x)/2)^7)/(7*c^4*f)$$

3.31
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx$$

3.31.1	Optimal result	305
3.31.2	Mathematica [A] (verified)	305
3.31.3	Rubi [A] (verified)	306
3.31.4	Maple [A] (verified)	307
3.31.5	Fricas [A] (verification not implemented)	308
3.31.6	Sympy [F]	308
3.31.7	Maxima [B] (verification not implemented)	309
3.31.8	Giac [A] (verification not implemented)	310
3.31.9	Mupad [B] (verification not implemented)	310

3.31.1 Optimal result

Integrand size = 32, antiderivative size = 80

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx = -\frac{(a+a\sec(e+fx))^3 \tan(e+fx)}{9f(c-c\sec(e+fx))^5} - \frac{(a+a\sec(e+fx))^3 \tan(e+fx)}{63cf(c-c\sec(e+fx))^4}$$

output `-1/9*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^5-1/63*(a+a*sec(f*x+e))^3*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^4`

3.31.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.59

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx = -\frac{a^3(-8+\sec(e+fx))(1+\sec(e+fx))^3 \tan(e+fx)}{63c^5f(-1+\sec(e+fx))^5}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^5,x]`

output `-1/63*(a^3*(-8 + Sec[e + f*x])*(1 + Sec[e + f*x])^3*Tan[e + f*x])/(c^5*f*(-1 + Sec[e + f*x])^5)`

3.31.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx$$

3.31.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c-c\sec(e+fx))^5} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^5} dx \\
 & \quad \downarrow \text{4439} \\
 & \frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^3}{(c-c\sec(e+fx))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \\
 & \quad \downarrow \text{4438} \\
 & -\frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{63cf(c-c\sec(e+fx))^4} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^5,x]`

output `-1/9*((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^5) - ((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(63*c*f*(c - c*Sec[e + f*x])^4)`

3.31. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx$

3.31.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)]`

3.31.4 Maple [A] (verified)

Time = 1.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.48

method	result
parallelrisch	$\frac{a^3 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^7 \left(7 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 9\right)}{126c^5 f}$
derivativedivides	$\frac{a^3 \left(\frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} \right)}{2f c^5}$
default	$\frac{a^3 \left(\frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} \right)}{2f c^5}$
risch	$\frac{2ia^3 (63 e^{8i(fx+e)} - 63 e^{7i(fx+e)} + 483 e^{6i(fx+e)} - 315 e^{5i(fx+e)} + 693 e^{4i(fx+e)} - 189 e^{3i(fx+e)} + 225 e^{2i(fx+e)} - 9 e^{i(fx+e)})}{63f c^5 (e^{i(fx+e)} - 1)^9}$
norman	$\frac{-\frac{a^3}{18cf} + \frac{5a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{21cf} - \frac{8a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{21cf} + \frac{17a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{63cf} - \frac{a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{14cf}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^3 c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$

3.31. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x,method=_RETURNVERBOSE)`

output $1/126*a^3*\cot(1/2*f*x+1/2*e)^7*(7*\cot(1/2*f*x+1/2*e)^2-9)/c^5/f$

3.31.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.75

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx$$

$$= \frac{8a^3 \cos(fx+e)^5 + 31a^3 \cos(fx+e)^4 + 44a^3 \cos(fx+e)^3 + 26a^3 \cos(fx+e)^2 + 4a^3 \cos(fx+e) - a^3}{63(c^5 f \cos(fx+e)^4 - 4c^5 f \cos(fx+e)^3 + 6c^5 f \cos(fx+e)^2 - 4c^5 f \cos(fx+e) + c^5 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x, algorithm="fracas")`

output $1/63*(8*a^3*\cos(f*x + e)^5 + 31*a^3*\cos(f*x + e)^4 + 44*a^3*\cos(f*x + e)^3 + 26*a^3*\cos(f*x + e)^2 + 4*a^3*\cos(f*x + e) - a^3)/((c^5*f*\cos(f*x + e)^4 - 4*c^5*f*\cos(f*x + e)^3 + 6*c^5*f*\cos(f*x + e)^2 - 4*c^5*f*\cos(f*x + e) + c^5*f)*\sin(f*x + e))$

3.31.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx =$$

$$\frac{a^3 \left(\int \frac{\sec(e+fx)}{\sec^5(e+fx)-5\sec^4(e+fx)+10\sec^3(e+fx)-10\sec^2(e+fx)+5\sec(e+fx)-1} dx + \int \frac{3\sec^2(e+fx)}{\sec^5(e+fx)-5\sec^4(e+fx)+10\sec^3(e+fx)} dx \right)}{c^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**5,x)`

```
output -a**3*(Integral(sec(e + f*x)/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec
(e + f*x)**3 - 10*sec(e + f*x)**2 + 5*sec(e + f*x) - 1), x) + Integral(3*s
ec(e + f*x)**2/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec(e + f*x)**3 -
10*sec(e + f*x)**2 + 5*sec(e + f*x) - 1), x) + Integral(3*sec(e + f*x)**3
/(sec(e + f*x)**5 - 5*sec(e + f*x)**4 + 10*sec(e + f*x)**3 - 10*sec(e + f
*x)**2 + 5*sec(e + f*x) - 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x)**
5 - 5*sec(e + f*x)**4 + 10*sec(e + f*x)**3 - 10*sec(e + f*x)**2 + 5*sec(e
+ f*x) - 1), x))/c**5
```

3.31.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 357 vs. $2(78) = 156$.

Time = 0.23 (sec) , antiderivative size = 357, normalized size of antiderivative = 4.46

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^5} dx = \frac{a^3 \left(\frac{180 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{378 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{420 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{315 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} - 35 \right) (\cos(fx+e)+1)^9}{c^5 \sin^9(fx+e)} + \frac{15 a^3 \left(\frac{18 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{42 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{63 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{7 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} - 7 \right) (\cos(fx+e)+1)^9}{c^5 \sin^9(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x, algorithm="m
axima")
```

```
output -1/5040*(a^3*(180*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 378*sin(f*x + e)^4
/(cos(f*x + e) + 1)^4 + 420*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 315*sin(
f*x + e)^8/(cos(f*x + e) + 1)^8 - 35)*(cos(f*x + e) + 1)^9/(c^5*sin(f*x +
e)^9) + 15*a^3*(18*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 42*sin(f*x + e)^6
/(cos(f*x + e) + 1)^6 + 63*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 - 7)*(cos(f
*x + e) + 1)^9/(c^5*sin(f*x + e)^9) - 5*a^3*(18*sin(f*x + e)^2/(cos(f*x +
e) + 1)^2 - 42*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 63*sin(f*x + e)^8/(co
s(f*x + e) + 1)^8 + 7)*(cos(f*x + e) + 1)^9/(c^5*sin(f*x + e)^9) + 21*a^3*
(18*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 45*sin(f*x + e)^8/(cos(f*x + e)
+ 1)^8 - 5)*(cos(f*x + e) + 1)^9/(c^5*sin(f*x + e)^9))/f
```

3.31.8 Giac [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.51

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^5} dx = -\frac{9a^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 7a^3}{126c^5 f \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^9}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x, algorithm="giac")`

output `-1/126*(9*a^3*tan(1/2*f*x + 1/2*e)^2 - 7*a^3)/(c^5*f*tan(1/2*f*x + 1/2*e)^9)`

3.31.9 Mupad [B] (verification not implemented)

Time = 12.93 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.46

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^5} dx = \frac{a^3 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^7 \left(7 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 9\right)}{126c^5 f}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^5),x)`

output `(a^3*cot(e/2 + (f*x)/2)^7*(7*cot(e/2 + (f*x)/2)^2 - 9))/(126*c^5*f)`

3.32 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^6} dx$

3.32.1 Optimal result 311
 3.32.2 Mathematica [A] (verified) 311
 3.32.3 Rubi [A] (verified) 312
 3.32.4 Maple [A] (verified) 314
 3.32.5 Fricas [A] (verification not implemented) 314
 3.32.6 Sympy [F] 315
 3.32.7 Maxima [B] (verification not implemented) 315
 3.32.8 Giac [A] (verification not implemented) 316
 3.32.9 Mupad [B] (verification not implemented) 317

3.32.1 Optimal result

Integrand size = 32, antiderivative size = 121

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^6} dx = -\frac{(a+a \sec(e+fx))^3 \tan(e+fx)}{11f(c-c \sec(e+fx))^6} - \frac{2(a+a \sec(e+fx))^3 \tan(e+fx)}{99cf(c-c \sec(e+fx))^5} - \frac{2(a+a \sec(e+fx))^3 \tan(e+fx)}{693c^2f(c-c \sec(e+fx))^4}$$

output `-1/11*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^6-2/99*(a+a*sec(f*x+e))^3*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^5-2/693*(a+a*sec(f*x+e))^3*tan(f*x+e)/c^2/f/(c-c*sec(f*x+e))^4`

3.32.2 Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.49

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^6} dx = -\frac{a^3(1+\sec(e+fx))^3(79-18 \sec(e+fx)+2 \sec^2(e+fx)) \tan(e+fx)}{693c^6f(-1+\sec(e+fx))^6}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^6,x]`

output
$$-1/693*(a^3*(1 + \text{Sec}[e + f*x])^3*(79 - 18*\text{Sec}[e + f*x] + 2*\text{Sec}[e + f*x]^2)*\text{Tan}[e + f*x])/(c^6*f*(-1 + \text{Sec}[e + f*x])^6)$$

3.32.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c-c\sec(e+fx))^6} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^6} dx \\ & \quad \downarrow \text{4439} \\ & \frac{2 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^3}{(c-c\sec(e+fx))^5} dx}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \\ & \quad \downarrow \text{3042} \\ & \frac{2 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^5} dx}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \\ & \quad \downarrow \text{4439} \\ & \frac{2 \left(\frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^3}{(c-c\sec(e+fx))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \\ & \quad \downarrow \text{3042} \\ & \frac{2 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \\ & \quad \downarrow \text{4438} \end{aligned}$$

3.32.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^6} dx$$

$$2 \left(\frac{-\tan(e+fx)(a \sec(e+fx)+a)^3}{63cf(c-c\sec(e+fx))^4} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \right) - \frac{\tan(e+fx)(a \sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^6,x]`

output `-1/11*((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^6) + (2*(-1/9*((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^5) - ((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(63*c*f*(c - c*Sec[e + f*x])^4)))/(11*c)`

3.32.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.32.4 Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.42

method	result
parallelrisc	$-\frac{a^3 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^7 \left(63 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^4 - 154 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 + 99\right)}{2772c^6 f}$
derivativedivides	$a^3 \left(-\frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{2}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} \right) \frac{1}{4f c^6}$
default	$a^3 \left(-\frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{2}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} \right) \frac{1}{4f c^6}$
risc	$\frac{2ia^3 (693 e^{10i(fx+e)} - 1386 e^{9i(fx+e)} + 8085 e^{8i(fx+e)} - 10626 e^{7i(fx+e)} + 21252 e^{6i(fx+e)} - 15246 e^{5i(fx+e)} + 15444 e^{4i(fx+e)} - 10626 e^{3i(fx+e)} + 4085 e^{2i(fx+e)} - 1386 e^{i(fx+e)} + 693) e^{i(fx+e)}}{693 f c^6 (e^{i(fx+e)} - 1)^{11}}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x,method=_RETURNVERBOSE)`

output `-1/2772*a^3*cot(1/2*f*x+1/2*e)^7*(63*cot(1/2*f*x+1/2*e)^4-154*cot(1/2*f*x+1/2*e)^2+99)/c^6/f`

3.32.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.39

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^6} dx$$

$$= \frac{79 a^3 \cos(fx + e)^6 + 298 a^3 \cos(fx + e)^5 + 404 a^3 \cos(fx + e)^4 + 216 a^3 \cos(fx + e)^3 + 19 a^3 \cos(fx + e)^2 - 10 a^3 \cos(fx + e) + 2 a^3}{693 (c^6 f \cos(fx + e)^5 - 5 c^6 f \cos(fx + e)^4 + 10 c^6 f \cos(fx + e)^3 - 10 c^6 f \cos(fx + e)^2 + 5 c^6 f \cos(fx + e) - c^6 f) \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x, algorithm="fricas")`

output `1/693*(79*a^3*cos(f*x + e)^6 + 298*a^3*cos(f*x + e)^5 + 404*a^3*cos(f*x + e)^4 + 216*a^3*cos(f*x + e)^3 + 19*a^3*cos(f*x + e)^2 - 10*a^3*cos(f*x + e) + 2*a^3)/((c^6*f*cos(f*x + e)^5 - 5*c^6*f*cos(f*x + e)^4 + 10*c^6*f*cos(f*x + e)^3 - 10*c^6*f*cos(f*x + e)^2 + 5*c^6*f*cos(f*x + e) - c^6*f)*sin(f*x + e))`

3.32. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^6} dx$

3.32.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^6} dx$$

$$= \frac{a^3 \left(\int \frac{\sec(e+fx)}{\sec^6(e+fx)-6\sec^5(e+fx)+15\sec^4(e+fx)-20\sec^3(e+fx)+15\sec^2(e+fx)-6\sec(e+fx)+1} dx + \int \frac{1}{\sec^6(e+fx)-6\sec^5(e+fx)+1} dx \right)}{c^6}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**6,x)`

output `a**3*(Integral(sec(e + f*x)/(sec(e + f*x)**6 - 6*sec(e + f*x)**5 + 15*sec(e + f*x)**4 - 20*sec(e + f*x)**3 + 15*sec(e + f*x)**2 - 6*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**2/(sec(e + f*x)**6 - 6*sec(e + f*x)**5 + 15*sec(e + f*x)**4 - 20*sec(e + f*x)**3 + 15*sec(e + f*x)**2 - 6*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**3/(sec(e + f*x)**6 - 6*sec(e + f*x)**5 + 15*sec(e + f*x)**4 - 20*sec(e + f*x)**3 + 15*sec(e + f*x)**2 - 6*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x)**6 - 6*sec(e + f*x)**5 + 15*sec(e + f*x)**4 - 20*sec(e + f*x)**3 + 15*sec(e + f*x)**2 - 6*sec(e + f*x) + 1), x))/c**6`

3.32.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 518 vs. $2(118) = 236$.

Time = 0.22 (sec) , antiderivative size = 518, normalized size of antiderivative = 4.28

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^6} dx$$

$$= \frac{3a^3 \left(\frac{385 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} + \frac{990 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} - \frac{1386 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{1155 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} + \frac{3465 \sin^{10}(fx+e)}{(\cos(fx+e)+1)^{10}} - 315 \right) (\cos(fx+e)+1)^{11}}{c^6 \sin^6(fx+e)^{11}} + \frac{9a^3 \left(\frac{385 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{385 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{1386 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{1155 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} + \frac{3465 \sin^{10}(fx+e)}{(\cos(fx+e)+1)^{10}} - 315 \right) (\cos(fx+e)+1)^{11}}{c^6 \sin^6(fx+e)^{11}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x, algorithm="maxima")`

```
output 1/110880*(3*a^3*(385*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 990*sin(f*x + e)
)^4/(cos(f*x + e) + 1)^4 - 1386*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 1155
*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 3465*sin(f*x + e)^10/(cos(f*x + e)
+ 1)^10 - 315)*(cos(f*x + e) + 1)^11/(c^6*sin(f*x + e)^11) + 9*a^3*(385*si
n(f*x + e)^2/(cos(f*x + e) + 1)^2 - 330*sin(f*x + e)^4/(cos(f*x + e) + 1)^
4 - 462*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 1155*sin(f*x + e)^8/(cos(f*x
+ e) + 1)^8 - 1155*sin(f*x + e)^10/(cos(f*x + e) + 1)^10 - 105)*(cos(f*x
+ e) + 1)^11/(c^6*sin(f*x + e)^11) + 5*a^3*(385*sin(f*x + e)^2/(cos(f*x +
e) + 1)^2 - 990*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 1386*sin(f*x + e)^6/
(cos(f*x + e) + 1)^6 - 1155*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 693*sin(
f*x + e)^10/(cos(f*x + e) + 1)^10 - 63)*(cos(f*x + e) + 1)^11/(c^6*sin(f*x
+ e)^11) - a^3*(385*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 990*sin(f*x + e
)^4/(cos(f*x + e) + 1)^4 - 1386*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 1155
*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 3465*sin(f*x + e)^10/(cos(f*x + e)
+ 1)^10 + 315)*(cos(f*x + e) + 1)^11/(c^6*sin(f*x + e)^11))/f
```

3.32.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.47

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^6} dx$$

$$= -\frac{99 a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 - 154 a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 + 63 a^3}{2772 c^6 f \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11}}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x, algorithm="g
iac")
```

```
output -1/2772*(99*a^3*tan(1/2*f*x + 1/2*e)^4 - 154*a^3*tan(1/2*f*x + 1/2*e)^2 +
63*a^3)/(c^6*f*tan(1/2*f*x + 1/2*e)^11)
```

3.32.9 Mupad [B] (verification not implemented)

Time = 13.32 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.55

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^6} dx$$

$$= \frac{a^3 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{18c^6 f} - \frac{a^3 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{28c^6 f} - \frac{a^3 \cot\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}{44c^6 f}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^6),x)`output `(a^3*cot(e/2 + (f*x)/2)^9)/(18*c^6*f) - (a^3*cot(e/2 + (f*x)/2)^7)/(28*c^6*f) - (a^3*cot(e/2 + (f*x)/2)^11)/(44*c^6*f)`

3.33
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^7} dx$$

3.33.1	Optimal result	318
3.33.2	Mathematica [A] (verified)	319
3.33.3	Rubi [A] (verified)	319
3.33.4	Maple [A] (verified)	321
3.33.5	Fricas [A] (verification not implemented)	322
3.33.6	Sympy [F]	323
3.33.7	Maxima [B] (verification not implemented)	323
3.33.8	Giac [A] (verification not implemented)	324
3.33.9	Mupad [B] (verification not implemented)	325

3.33.1 Optimal result

Integrand size = 32, antiderivative size = 162

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^7} dx = -\frac{(a+a \sec(e+fx))^3 \tan(e+fx)}{13f(c-c \sec(e+fx))^7} - \frac{3(a+a \sec(e+fx))^3 \tan(e+fx)}{143cf(c-c \sec(e+fx))^6} - \frac{2(a+a \sec(e+fx))^3 \tan(e+fx)}{429c^2f(c-c \sec(e+fx))^5} - \frac{2(a+a \sec(e+fx))^3 \tan(e+fx)}{3003c^3f(c-c \sec(e+fx))^4}$$

```
output -1/13*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^7-3/143*(a+a*sec(f*x+e))^3*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^6-2/429*(a+a*sec(f*x+e))^3*tan(f*x+e)/c^2/f/(c-c*sec(f*x+e))^5-2/3003*(a+a*sec(f*x+e))^3*tan(f*x+e)/c^3/f/(c-c*sec(f*x+e))^4
```

3.33.2 Mathematica [A] (verified)

Time = 5.00 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.43

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^7} dx = \frac{a^3(1+\sec(e+fx))^3(-310+97\sec(e+fx)-20\sec^2(e+fx)+2\sec^3(e+fx))\tan(e+fx)}{3003c^7f(-1+\sec(e+fx))^7}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^7,x]`

output `-1/3003*(a^3*(1 + Sec[e + f*x])^3*(-310 + 97*Sec[e + f*x] - 20*Sec[e + f*x]^2 + 2*Sec[e + f*x]^3)*Tan[e + f*x])/(c^7*f*(-1 + Sec[e + f*x])^7)`

3.33.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4439, 3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c-c\sec(e+fx))^7} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^7} dx \\ & \quad \downarrow \text{4439} \\ & \frac{3 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^3}{(c-c\sec(e+fx))^6} dx}{13c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{13f(c-c\sec(e+fx))^7} \\ & \quad \downarrow \text{3042} \\ & \frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^6} dx}{13c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{13f(c-c\sec(e+fx))^7} \\ & \quad \downarrow \text{4439} \end{aligned}$$

3.33. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^7} dx$

$$\begin{aligned}
& \frac{3 \left(\frac{2 \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^3}{(c-c\sec(e+fx))^5} dx}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \right)}{13c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{13f(c-c\sec(e+fx))^7} \\
& \quad \downarrow 3042 \\
& \frac{3 \left(\frac{2 \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^5} dx}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \right)}{13c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{13f(c-c\sec(e+fx))^7} \\
& \quad \downarrow 4439 \\
& \frac{3 \left(\frac{2 \left(\frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^3}{(c-c\sec(e+fx))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \right)}{13c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{13f(c-c\sec(e+fx))^7} \\
& \quad \downarrow 3042 \\
& \frac{3 \left(\frac{2 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx}{9c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \right)}{13c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{13f(c-c\sec(e+fx))^7} \\
& \quad \downarrow 4438 \\
& \frac{3 \left(\frac{2 \left(-\frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{63cf(c-c\sec(e+fx))^4} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{9f(c-c\sec(e+fx))^5} \right)}{11c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{11f(c-c\sec(e+fx))^6} \right)}{13c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^3}{13f(c-c\sec(e+fx))^7}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^7,x]`

3.33. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^7} dx$

```
output -1/13*((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^7) + (
3*(-1/11*((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^6)
+ (2*(-1/9*((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^5
) - ((a + a*Sec[e + f*x])^3*Tan[e + f*x])/(63*c*f*(c - c*Sec[e + f*x])^4)
)/(11*c)))/(13*c)
```

3.33.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4438 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]
*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; Fre
eQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] &
& EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]
```

```
rule 4439 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]
*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp
[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*
(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ
[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0
] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])
```

3.33.4 Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.40

method	result
parallelrisch	$\frac{a^3 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^7 \left(231 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^6 - 819 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^4 + 1001 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 429\right)}{24024c^7 f}$
derivativedivides	$\frac{a^3 \left(\frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{3}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{1}{13 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{13}} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}\right)}{8f c^7}$
default	$\frac{a^3 \left(\frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} - \frac{3}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{1}{13 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{13}} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}\right)}{8f c^7}$
risch	$\frac{2ia^3(3003e^{12i(fx+e)} - 9009e^{11i(fx+e)} + 51051e^{10i(fx+e)} - 99099e^{9i(fx+e)} + 216216e^{8i(fx+e)} - 246246e^{7i(fx+e)} + 285717e^{6i(fx+e)} - 252252e^{5i(fx+e)} + 182182e^{4i(fx+e)} - 98919e^{3i(fx+e)} + 3003e^{2i(fx+e)} - 21e^{i(fx+e)} + 1)}{3003f c^7 (e^{i(fx+e)} - 1)}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^7,x,method=_RETURNVERBOSE)`

output $\frac{1}{24024} a^3 \cot\left(\frac{1}{2} f x + \frac{1}{2} e\right)^7 (231 \cot\left(\frac{1}{2} f x + \frac{1}{2} e\right)^6 - 819 \cot\left(\frac{1}{2} f x + \frac{1}{2} e\right)^4 + 1001 \cot\left(\frac{1}{2} f x + \frac{1}{2} e\right)^2 - 429) / c^7 / f$

3.33.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.20

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^7} dx$$

$$= \frac{310 a^3 \cos(fx+e)^7 + 1143 a^3 \cos(fx+e)^6 + 1492 a^3 \cos(fx+e)^5 + 736 a^3 \cos(fx+e)^4 + 34 a^3 \cos(fx+e)^3 - 29 a^3 \cos(fx+e)^2 + 12 a^3 \cos(fx+e) - 2 a^3}{3003 (c^7 f \cos(fx+e)^6 - 6 c^7 f \cos(fx+e)^5 + 15 c^7 f \cos(fx+e)^4 - 20 c^7 f \cos(fx+e)^3 + 15 c^7 f \cos(fx+e)^2 - 6 c^7 f \cos(fx+e) + c^7 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^7,x, algorithm="fricas")`

output $\frac{1}{3003} (310 a^3 \cos(fx+e)^7 + 1143 a^3 \cos(fx+e)^6 + 1492 a^3 \cos(fx+e)^5 + 736 a^3 \cos(fx+e)^4 + 34 a^3 \cos(fx+e)^3 - 29 a^3 \cos(fx+e)^2 + 12 a^3 \cos(fx+e) - 2 a^3) / ((c^7 f \cos(fx+e)^6 - 6 c^7 f \cos(fx+e)^5 + 15 c^7 f \cos(fx+e)^4 - 20 c^7 f \cos(fx+e)^3 + 15 c^7 f \cos(fx+e)^2 - 6 c^7 f \cos(fx+e) + c^7 f) \sin(fx+e))$

3.33.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^7} dx =$$

$$a^3 \left(\int \frac{\sec(e+fx)}{\sec^7(e+fx) - 7\sec^6(e+fx) + 21\sec^5(e+fx) - 35\sec^4(e+fx) + 35\sec^3(e+fx) - 21\sec^2(e+fx) + 7\sec(e+fx) - 1} dx + \int \frac{1}{\sec^7(e+fx)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**7,x)`

output

```
-a**3*(Integral(sec(e + f*x)/(sec(e + f*x)**7 - 7*sec(e + f*x)**6 + 21*sec(e + f*x)**5 - 35*sec(e + f*x)**4 + 35*sec(e + f*x)**3 - 21*sec(e + f*x)**2 + 7*sec(e + f*x) - 1), x) + Integral(3*sec(e + f*x)**2/(sec(e + f*x)**7 - 7*sec(e + f*x)**6 + 21*sec(e + f*x)**5 - 35*sec(e + f*x)**4 + 35*sec(e + f*x)**3 - 21*sec(e + f*x)**2 + 7*sec(e + f*x) - 1), x) + Integral(3*sec(e + f*x)**3/(sec(e + f*x)**7 - 7*sec(e + f*x)**6 + 21*sec(e + f*x)**5 - 35*sec(e + f*x)**4 + 35*sec(e + f*x)**3 - 21*sec(e + f*x)**2 + 7*sec(e + f*x) - 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x)**7 - 7*sec(e + f*x)**6 + 21*sec(e + f*x)**5 - 35*sec(e + f*x)**4 + 35*sec(e + f*x)**3 - 21*sec(e + f*x)**2 + 7*sec(e + f*x) - 1), x))/c**7
```

3.33.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 517 vs. $2(158) = 316$.

Time = 0.24 (sec) , antiderivative size = 517, normalized size of antiderivative = 3.19

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^7} dx =$$

$$\frac{a^3 \left(\frac{8190 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} + \frac{5005 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} - \frac{25740 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} + \frac{9009 \sin^8(fx+e)}{(\cos(fx+e)+1)^8} + \frac{30030 \sin^{10}(fx+e)}{(\cos(fx+e)+1)^{10}} - \frac{45045 \sin^{12}(fx+e)}{(\cos(fx+e)+1)^{12}} - 3465 \right) (\cos(fx+e)+1)^{13}}{c^7 \sin^2(fx+e)^{13}} +$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^7,x, algorithm="maxima")`

output

```
-1/960960*(a^3*(8190*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 5005*sin(f*x +
e)^4/(cos(f*x + e) + 1)^4 - 25740*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 90
09*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 30030*sin(f*x + e)^10/(cos(f*x +
e) + 1)^10 - 45045*sin(f*x + e)^12/(cos(f*x + e) + 1)^12 - 3465)*(cos(f*x
+ e) + 1)^13/(c^7*sin(f*x + e)^13) + 5*a^3*(1638*sin(f*x + e)^2/(cos(f*x +
e) + 1)^2 - 5005*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 8580*sin(f*x + e)^
6/(cos(f*x + e) + 1)^6 - 9009*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 6006*s
in(f*x + e)^10/(cos(f*x + e) + 1)^10 - 3003*sin(f*x + e)^12/(cos(f*x + e)
+ 1)^12 - 231)*(cos(f*x + e) + 1)^13/(c^7*sin(f*x + e)^13) + 35*a^3*(468*s
in(f*x + e)^2/(cos(f*x + e) + 1)^2 - 715*sin(f*x + e)^4/(cos(f*x + e) + 1)
^4 + 1287*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 - 1716*sin(f*x + e)^10/(cos(
f*x + e) + 1)^10 + 1287*sin(f*x + e)^12/(cos(f*x + e) + 1)^12 - 99)*(cos(f
*x + e) + 1)^13/(c^7*sin(f*x + e)^13) + 77*a^3*(65*sin(f*x + e)^4/(cos(f*x
+ e) + 1)^4 - 117*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 195*sin(f*x + e)^
12/(cos(f*x + e) + 1)^12 - 15)*(cos(f*x + e) + 1)^13/(c^7*sin(f*x + e)^13)
)/f
```

3.33.8 Giac [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.45

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^7} dx$$

$$= \frac{429 a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 - 1001 a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 + 819 a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 231 a^3}{24024 c^7 f \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{13}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^7,x, algorithm="giac")`

output

```
-1/24024*(429*a^3*tan(1/2*f*x + 1/2*e)^6 - 1001*a^3*tan(1/2*f*x + 1/2*e)^4
+ 819*a^3*tan(1/2*f*x + 1/2*e)^2 - 231*a^3)/(c^7*f*tan(1/2*f*x + 1/2*e)^1
3)
```

3.33.9 Mupad [B] (verification not implemented)

Time = 13.23 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^7} dx$$

$$= \frac{a^3 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^7 \left(231 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 819 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^4 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1001 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^2 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{24024 c^7 f \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^{13}}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^7),x)`

output `(a^3*cos(e/2 + (f*x)/2)^7*(231*cos(e/2 + (f*x)/2)^6 - 429*sin(e/2 + (f*x)/2)^6 + 1001*cos(e/2 + (f*x)/2)^2*sin(e/2 + (f*x)/2)^4 - 819*cos(e/2 + (f*x)/2)^4*sin(e/2 + (f*x)/2)^2)/(24024*c^7*f*sin(e/2 + (f*x)/2)^13)`

$$3.34 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx$$

3.34.1	Optimal result	326
3.34.2	Mathematica [C] (verified)	326
3.34.3	Rubi [A] (verified)	327
3.34.4	Maple [A] (verified)	329
3.34.5	Fricas [A] (verification not implemented)	329
3.34.6	Sympy [F]	330
3.34.7	Maxima [B] (verification not implemented)	330
3.34.8	Giac [A] (verification not implemented)	331
3.34.9	Mupad [B] (verification not implemented)	332

3.34.1 Optimal result

Integrand size = 32, antiderivative size = 121

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx = -\frac{35c^4 \operatorname{arctanh}(\sin(e+fx))}{2af} + \frac{28c^4 \tan(e+fx)}{af} - \frac{21c^4 \sec(e+fx) \tan(e+fx)}{2af} + \frac{2c(c-c\sec(e+fx))^3 \tan(e+fx)}{f(a+a\sec(e+fx))} + \frac{7c^4 \tan^3(e+fx)}{3af}$$

output `-35/2*c^4*arctanh(sin(f*x+e))/a/f+28*c^4*tan(f*x+e)/a/f-21/2*c^4*sec(f*x+e)*tan(f*x+e)/a/f+2*c*(c-c*sec(f*x+e))^3*tan(f*x+e)/f/(a+a*sec(f*x+e))+7/3*c^4*tan(f*x+e)^3/a/f`

3.34.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.99 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.44

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx = -\frac{16c^4 \cot(e+fx) \operatorname{Hypergeometric2F1}\left(-\frac{7}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)}}{af}$$

3.34. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^4)/(a + a*Sec[e + f*x]),x]`

output `(-16*c^4*Cot[e + f*x]*Hypergeometric2F1[-7/2, -1/2, 1/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]])/(a*f)`

3.34.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4445, 3042, 4278, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^4}{a\csc(e+fx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{4445} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{f(a\sec(e+fx)+a)} - \frac{7c\int\sec(e+fx)(c-c\sec(e+fx))^3dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{f(a\sec(e+fx)+a)} - \frac{7c\int\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^3dx}{a} \\
 & \quad \downarrow \text{4278} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{f(a\sec(e+fx)+a)} - \frac{7c\int(-c^3\sec^4(e+fx)+3c^3\sec^3(e+fx)-3c^3\sec^2(e+fx)+c^3\sec(e+fx))dx}{a} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{f(a\sec(e+fx)+a)} - \frac{7c\left(\frac{5c^3\operatorname{arctanh}(\sin(e+fx))}{2f} - \frac{c^3\tan^3(e+fx)}{3f} - \frac{4c^3\tan(e+fx)}{f} + \frac{3c^3\tan(e+fx)\sec(e+fx)}{2f}\right)}{a}
 \end{aligned}$$

3.34. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^4)/(a + a*Sec[e + f*x]),x]`

output `(2*c*(c - c*Sec[e + f*x])^3*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (7*c*((5*c^3*ArcTanh[Sin[e + f*x]])/(2*f) - (4*c^3*Tan[e + f*x])/f + (3*c^3*Sec[e + f*x]*Tan[e + f*x])/(2*f) - (c^3*Tan[e + f*x]^3)/(3*f)))/a`

3.34.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4278 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^{(n_.)}*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^{(m_.)}, x_Symbol] := Int[ExpandTrig[(a + b*csc[e + f*x])^m*(d*csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && IGtQ[m, 0] && RationalQ[n]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_)])*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^{(m_.)}*(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.)^{(n_.)}, x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.34.4 Maple [A] (verified)

Time = 1.30 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.14

method	result
parallelrisch	$105 \left(\left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3} \right) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) + \left(-\cos(fx+e) - \frac{\cos(3fx+3e)}{3} \right) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right) + \frac{446 \left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3} \right)}{2af(\cos(3fx+3e) + 3\cos(fx+e))} \right)$
derivativedivides	$16c^4 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - \frac{1}{48 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^3} - \frac{3}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^2} - \frac{29}{32 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)} + \frac{35 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)}{32} - \frac{1}{48 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)} \right) \frac{1}{fa}$
default	$16c^4 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - \frac{1}{48 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^3} - \frac{3}{16 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^2} - \frac{29}{32 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)} + \frac{35 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)}{32} - \frac{1}{48 \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)} \right) \frac{1}{fa}$
risch	$\frac{ic^4 \left(111 e^{6i(fx+e)} + 81 e^{5i(fx+e)} + 354 e^{4i(fx+e)} + 144 e^{3i(fx+e)} + 417 e^{2i(fx+e)} + 55 e^{i(fx+e)} + 166 \right)}{3af(1+e^{2i(fx+e)})^3(e^{i(fx+e)}+1)} + \frac{35c^4 \ln(e^{i(fx+e)}-1)}{2af}$
norman	$\frac{\frac{35c^4 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)}{af} - \frac{385c^4 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^3}{3af} + \frac{511c^4 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^5}{3af} - \frac{93c^4 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^7}{af} + \frac{16c^4 \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^9}{af}}{\left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)^4} + \frac{35c^4 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)}{2af}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `105/2*((cos(f*x+e)+1/3*cos(3*f*x+3*e))*ln(tan(1/2*f*x+1/2*e)-1)+(-cos(f*x+e)-1/3*cos(3*f*x+3*e))*ln(tan(1/2*f*x+1/2*e)+1)+446/315*(cos(f*x+e)+55/223*cos(2*f*x+2*e))+83/223*cos(3*f*x+3*e)+59/223)*tan(1/2*f*x+1/2*e)*c^4/a/f/(cos(3*f*x+3*e)+3*cos(f*x+e))`

3.34.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.26

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx = \frac{105(c^4 \cos(fx+e)^4 + c^4 \cos(fx+e)^3) \log(\sin(fx+e)+1) - 105(c^4 \cos(fx+e)^4 + c^4 \cos(fx+e)^3)}{12(af \cos(fx+e) + 3c^4)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e)),x, algorithm="fricas")`

3.34.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx$$

output
$$\frac{-1/12*(105*(c^4*\cos(f*x + e))^4 + c^4*\cos(f*x + e)^3)*\log(\sin(f*x + e) + 1) - 105*(c^4*\cos(f*x + e)^4 + c^4*\cos(f*x + e)^3)*\log(-\sin(f*x + e) + 1) - 2*(166*c^4*\cos(f*x + e)^3 + 55*c^4*\cos(f*x + e)^2 - 13*c^4*\cos(f*x + e) + 2*c^4)*\sin(f*x + e))/(a*f*\cos(f*x + e)^4 + a*f*\cos(f*x + e)^3)}$$

3.34.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^4}{a + a\sec(e + fx)} dx$$

$$= \frac{c^4 \left(\int \frac{\sec(e+fx)}{\sec(e+fx)+1} dx + \int \left(-\frac{4\sec^2(e+fx)}{\sec(e+fx)+1} \right) dx + \int \frac{6\sec^3(e+fx)}{\sec(e+fx)+1} dx + \int \left(-\frac{4\sec^4(e+fx)}{\sec(e+fx)+1} \right) dx + \int \frac{\sec^5(e+fx)}{\sec(e+fx)+1} dx \right)}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**4/(a+a*sec(f*x+e)),x)`

output `c**4*(Integral(sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(-4*sec(e + f*x)**2/(sec(e + f*x) + 1), x) + Integral(6*sec(e + f*x)**3/(sec(e + f*x) + 1), x) + Integral(-4*sec(e + f*x)**4/(sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**5/(sec(e + f*x) + 1), x))/a`

3.34.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 591 vs. $2(116) = 232$.

Time = 0.21 (sec) , antiderivative size = 591, normalized size of antiderivative = 4.88

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^4}{a + a\sec(e + fx)} dx$$

$$= \frac{c^4 \left(\frac{2 \left(\frac{9 \sin(fx+e)}{\cos(fx+e)+1} - \frac{16 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{15 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a - \frac{3a \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{3a \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{a \sin(fx+e)^6}{(\cos(fx+e)+1)^6}} - \frac{9 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a} + \frac{9 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a} + \frac{6 \sin(fx+e)}{a(\cos(fx+e)+1)} \right) + 12 \dots}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output
$$\frac{1}{6}c^4 \left(\frac{2(9\sin(fx+e))}{(\cos(fx+e)+1)} - 16\frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + 15\frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right) / (a - 3a\sin(fx+e)^2 / (\cos(fx+e)+1)^2 + 3a\sin(fx+e)^4 / (\cos(fx+e)+1)^4 - a\sin(fx+e)^6 / (\cos(fx+e)+1)^6) - 9\log(\sin(fx+e) / (\cos(fx+e)+1)) / a + 9\log(\sin(fx+e) / (\cos(fx+e)+1) - 1) / a + 6\sin(fx+e) / (a(\cos(fx+e)+1))) + 12c^4 \left(\frac{2(\sin(fx+e))}{(\cos(fx+e)+1)} - 3\frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right) / (a - 2a\sin(fx+e)^2 / (\cos(fx+e)+1)^2 + a\sin(fx+e)^4 / (\cos(fx+e)+1)^4) - 3\log(\sin(fx+e) / (\cos(fx+e)+1)) / a + 3\log(\sin(fx+e) / (\cos(fx+e)+1) - 1) / a + 2\sin(fx+e) / (a(\cos(fx+e)+1))) - 36c^4 \left(\log(\sin(fx+e) / (\cos(fx+e)+1)) + 1 \right) / a - \log(\sin(fx+e) / (\cos(fx+e)+1) - 1) / a - 2\sin(fx+e) / ((a - a\sin(fx+e)^2 / (\cos(fx+e)+1)^2) * (\cos(fx+e)+1)) - \sin(fx+e) / (a(\cos(fx+e)+1))) - 24c^4 \left(\log(\sin(fx+e) / (\cos(fx+e)+1)) + 1 \right) / a - \log(\sin(fx+e) / (\cos(fx+e)+1) - 1) / a - \sin(fx+e) / (a(\cos(fx+e)+1))) + 6c^4 \sin(fx+e) / (a(\cos(fx+e)+1))) / f$$

3.34.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.09

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx = \frac{105c^4 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a} - \frac{105c^4 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a} - \frac{96c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{a} + \frac{2(87c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 136c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 57c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)^3 a} \cdot \frac{1}{6f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e)),x, algorithm="giac")`

output
$$-1/6*(105*c^4*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a - 105*c^4*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a - 96*c^4*\tan(1/2*f*x + 1/2*e)/a + 2*(87*c^4*\tan(1/2*f*x + 1/2*e)^5 - 136*c^4*\tan(1/2*f*x + 1/2*e)^3 + 57*c^4*\tan(1/2*f*x + 1/2*e)) / ((\tan(1/2*f*x + 1/2*e)^2 - 1)^3*a)) / f$$

3.34.9 Mupad [B] (verification not implemented)

Time = 13.27 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.93

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{a+a\sec(e+fx)} dx$$

$$= \frac{16c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{af} - \frac{29c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 - \frac{136c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3} + 19c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{af \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)^3} - \frac{35c^4 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{af}$$

input `int((c - c/cos(e + f*x))^4/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`output `(16*c^4*tan(e/2 + (f*x)/2))/(a*f) - (29*c^4*tan(e/2 + (f*x)/2)^5 - (136*c^4*tan(e/2 + (f*x)/2)^3)/3 + 19*c^4*tan(e/2 + (f*x)/2))/(a*f*(tan(e/2 + (f*x)/2)^2 - 1)^3) - (35*c^4*atanh(tan(e/2 + (f*x)/2)))/(a*f)`

3.35 $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{a+a\sec(e+fx)} dx$

3.35.1 Optimal result 333
 3.35.2 Mathematica [C] (verified) 333
 3.35.3 Rubi [A] (verified) 334
 3.35.4 Maple [A] (verified) 337
 3.35.5 Fricas [A] (verification not implemented) 337
 3.35.6 Sympy [F] 338
 3.35.7 Maxima [B] (verification not implemented) 338
 3.35.8 Giac [A] (verification not implemented) 339
 3.35.9 Mupad [B] (verification not implemented) 339

3.35.1 Optimal result

Integrand size = 32, antiderivative size = 100

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{a+a\sec(e+fx)} dx = -\frac{15c^3 \operatorname{arctanh}(\sin(e+fx))}{2af} + \frac{10c^3 \tan(e+fx)}{af} - \frac{5c^3 \sec(e+fx) \tan(e+fx)}{2af} + \frac{2c(c-c\sec(e+fx))^2 \tan(e+fx)}{f(a+a\sec(e+fx))}$$

output

```
-15/2*c^3*arctanh(sin(f*x+e))/a/f+10*c^3*tan(f*x+e)/a/f-5/2*c^3*sec(f*x+e)*tan(f*x+e)/a/f+2*c*(c-c*sec(f*x+e))^2*tan(f*x+e)/f/(a+a*sec(f*x+e))
```

3.35.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.48 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.53

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{a+a\sec(e+fx)} dx = \frac{8c^3 \cot(e+fx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)}}{af}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^3)/(a + a*Sec[e + f*x]),x]`

output `(-8*c^3*Cot[e + f*x]*Hypergeometric2F1[-5/2, -1/2, 1/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]])/(a*f)`

3.35.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4445, 3042, 4275, 3042, 4254, 24, 4534, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^3}{a\csc(e+fx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{4445} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{f(a\sec(e+fx)+a)} - \frac{5c\int\sec(e+fx)(c-c\sec(e+fx))^2 dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{f(a\sec(e+fx)+a)} - \frac{5c\int\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2 dx}{a} \\
 & \quad \downarrow \text{4275} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{f(a\sec(e+fx)+a)} - \frac{5c(\int\sec(e+fx)(\sec^2(e+fx)c^2+c^2) dx - 2c^2\int\sec^2(e+fx) dx)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{f(a\sec(e+fx)+a)} - \frac{5c(\int\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})^2 c^2+c^2) dx - 2c^2\int\csc(e+fx+\frac{\pi}{2})^2 dx)}{a}
 \end{aligned}$$

3.35. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{a+a\sec(e+fx)} dx$

$$\begin{aligned}
& \downarrow 4254 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{f(a \sec(e+fx) + a)} - \\
& \frac{5c \left(\frac{2c^2 \int 1d(-\tan(e+fx))}{f} + \int \csc \left(e+fx + \frac{\pi}{2} \right) \left(\csc \left(e+fx + \frac{\pi}{2} \right)^2 c^2 + c^2 \right) dx \right)}{a} \\
& \downarrow 24 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{f(a \sec(e+fx) + a)} - \\
& \frac{5c \left(\int \csc \left(e+fx + \frac{\pi}{2} \right) \left(\csc \left(e+fx + \frac{\pi}{2} \right)^2 c^2 + c^2 \right) dx - \frac{2c^2 \tan(e+fx)}{f} \right)}{a} \\
& \downarrow 4534 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{f(a \sec(e+fx) + a)} - \frac{5c \left(\frac{3}{2} c^2 \int \sec(e+fx) dx - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \\
& \downarrow 3042 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{f(a \sec(e+fx) + a)} - \\
& \frac{5c \left(\frac{3}{2} c^2 \int \csc \left(e+fx + \frac{\pi}{2} \right) dx - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \\
& \downarrow 4257 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{f(a \sec(e+fx) + a)} - \\
& \frac{5c \left(\frac{3c^2 \operatorname{arctanh}(\sin(e+fx))}{2f} - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^3)/(a + a*Sec[e + f*x]),x]`

output `(2*c*(c - c*Sec[e + f*x])^2*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (5*c*((3*c^2*ArcTanh[Sin[e + f*x]])/(2*f) - (2*c^2*Tan[e + f*x])/f + (c^2*Sec[e + f*x]*Tan[e + f*x])/(2*f)))/a`

3.35.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4275 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^2, x_Symbol] := Simp[2*a*(b/d) Int[(d*Csc[e + f*x])^(n + 1), x], x] + Int[(d*Csc[e + f*x])^n*(a^2 + b^2*Csc[e + f*x]^2), x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`
- rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.35.4 Maple [A] (verified)

Time = 1.10 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.06

method	result
parallelrisch	$\frac{15 \left((-1 - \cos(2fx+2e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + (1 + \cos(2fx+2e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) - \frac{14 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) (\cos(fx+e) + \frac{12}{15})}{15} \right)}{2af(1 + \cos(2fx+2e))}$
derivativedivides	$\frac{8c^3 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1)^2} - \frac{9}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1)} + \frac{15 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{16} + \frac{1}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^2} - \frac{9}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)} \right)}{fa}$
default	$\frac{8c^3 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1)^2} - \frac{9}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1)} + \frac{15 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{16} + \frac{1}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^2} - \frac{9}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)} \right)}{fa}$
risch	$\frac{ic^3 (17e^{4i(fx+e)} + 9e^{3i(fx+e)} + 39e^{2i(fx+e)} + 7e^{i(fx+e)} + 24)}{fa(e^{i(fx+e)} + 1)(1 + e^{2i(fx+e)})^2} + \frac{15c^3 \ln(e^{i(fx+e)} - i)}{2af} - \frac{15c^3 \ln(e^{i(fx+e)} + i)}{2af}$
norman	$\frac{-\frac{15c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{40c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} - \frac{33c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{af} + \frac{8c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^3} + \frac{15c^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2af} - \frac{15c^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2af}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `-15/2*((-1-cos(2*f*x+2*e))*ln(tan(1/2*f*x+1/2*e)-1)+(1+cos(2*f*x+2*e))*ln(tan(1/2*f*x+1/2*e)+1)-14/15*tan(1/2*f*x+1/2*e)*(cos(f*x+e)+12/7*cos(2*f*x+2*e)+11/7))*c^3/a/f/(1+cos(2*f*x+2*e))`

3.35.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.40

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{a+a\sec(e+fx)} dx = \frac{15(c^3 \cos(fx+e)^3 + c^3 \cos(fx+e)^2) \log(\sin(fx+e)+1) - 15(c^3 \cos(fx+e)^3 + c^3 \cos(fx+e)^2)}{4(af \cos(fx+e)^3 + af \cos(fx+e))}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e)),x, algorithm="fracas")`

output
$$\frac{-1/4*(15*(c^3*\cos(f*x + e))^3 + c^3*\cos(f*x + e)^2)*\log(\sin(f*x + e) + 1) - 15*(c^3*\cos(f*x + e))^3 + c^3*\cos(f*x + e)^2)*\log(-\sin(f*x + e) + 1) - 2*(24*c^3*\cos(f*x + e)^2 + 7*c^3*\cos(f*x + e) - c^3)*\sin(f*x + e)/(a*f*\cos(f*x + e)^3 + a*f*\cos(f*x + e)^2)}{a}$$

3.35.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^3}{a + a \sec(e + fx)} dx$$

$$= \frac{c^3 \left(\int \left(-\frac{\sec(e+fx)}{\sec(e+fx)+1} \right) dx + \int \frac{3 \sec^2(e+fx)}{\sec(e+fx)+1} dx + \int \left(-\frac{3 \sec^3(e+fx)}{\sec(e+fx)+1} \right) dx + \int \frac{\sec^4(e+fx)}{\sec(e+fx)+1} dx \right)}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**3/(a+a*sec(f*x+e)),x)`

output
$$-c**3*(Integral(-sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**2/(sec(e + f*x) + 1), x) + Integral(-3*sec(e + f*x)**3/(sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x) + 1), x))/a$$

3.35.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(97) = 194$.

Time = 0.20 (sec) , antiderivative size = 386, normalized size of antiderivative = 3.86

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^3}{a + a \sec(e + fx)} dx$$

$$= \frac{c^3 \left(\frac{2 \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - \frac{3 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a - \frac{2a \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{a \sin(fx+e)^4}{(\cos(fx+e)+1)^4}} - \frac{3 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a} + \frac{3 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a} + \frac{2 \sin(fx+e)}{a(\cos(fx+e)+1)} \right) - 6c^3 \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)}{a} \right)}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output $\frac{1}{2}(c^3(2(\sin(fx + e))/(\cos(fx + e) + 1) - 3\sin(fx + e)^3/(\cos(fx + e) + 1)^3)/(a - 2a\sin(fx + e)^2/(\cos(fx + e) + 1)^2 + a\sin(fx + e)^4/(\cos(fx + e) + 1)^4) - 3\log(\sin(fx + e)/(\cos(fx + e) + 1) + 1)/a + 3\log(\sin(fx + e)/(\cos(fx + e) + 1) - 1)/a + 2\sin(fx + e)/(a(\cos(fx + e) + 1))) - 6c^3(\log(\sin(fx + e)/(\cos(fx + e) + 1) + 1)/a - \log(\sin(fx + e)/(\cos(fx + e) + 1) - 1)/a - 2\sin(fx + e)/((a - a\sin(fx + e)^2/(\cos(fx + e) + 1)^2)(\cos(fx + e) + 1)) - \sin(fx + e)/(a(\cos(fx + e) + 1))) - 6c^3(\log(\sin(fx + e)/(\cos(fx + e) + 1) + 1)/a - \log(\sin(fx + e)/(\cos(fx + e) + 1) - 1)/a - \sin(fx + e)/(a(\cos(fx + e) + 1))) + 2c^3\sin(fx + e)/(a(\cos(fx + e) + 1)))/f$

3.35.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.16

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^3}{a + a\sec(e + fx)} dx = \frac{15c^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a} - \frac{15c^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a} - \frac{16c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{a} + \frac{2(9c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 7c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)^2 a}$$

$2f$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e)),x, algorithm="giac")`

output $-1/2(15c^3\log(\text{abs}(\tan(1/2*fx + 1/2*e) + 1))/a - 15c^3\log(\text{abs}(\tan(1/2*fx + 1/2*e) - 1))/a - 16c^3\tan(1/2*fx + 1/2*e)/a + 2(9c^3\tan(1/2*fx + 1/2*e)^3 - 7c^3\tan(1/2*fx + 1/2*e))/((\tan(1/2*fx + 1/2*e)^2 - 1)^2 a))/f$

3.35.9 Mupad [B] (verification not implemented)

Time = 13.59 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.96

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^3}{a + a\sec(e + fx)} dx = \frac{8c^3 \tan(\frac{e}{2} + \frac{fx}{2})}{af} - \frac{9c^3 \tan(\frac{e}{2} + \frac{fx}{2})^3 - 7c^3 \tan(\frac{e}{2} + \frac{fx}{2})}{af \left(\tan(\frac{e}{2} + \frac{fx}{2})^2 - 1\right)^2} - \frac{15c^3 \operatorname{atanh}(\tan(\frac{e}{2} + \frac{fx}{2}))}{af}$$

3.35. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{a+a\sec(e+fx)} dx$

input `int((c - c/cos(e + f*x))^3/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output `(8*c^3*tan(e/2 + (f*x)/2))/(a*f) - (9*c^3*tan(e/2 + (f*x)/2)^3 - 7*c^3*tan(e/2 + (f*x)/2))/(a*f*(tan(e/2 + (f*x)/2)^2 - 1)^2 - (15*c^3*atanh(tan(e/2 + (f*x)/2)))/(a*f)`

3.36 $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx$

3.36.1 Optimal result 341
 3.36.2 Mathematica [C] (verified) 341
 3.36.3 Rubi [A] (verified) 342
 3.36.4 Maple [A] (verified) 344
 3.36.5 Fricas [A] (verification not implemented) 344
 3.36.6 Sympy [F] 345
 3.36.7 Maxima [B] (verification not implemented) 345
 3.36.8 Giac [A] (verification not implemented) 346
 3.36.9 Mupad [B] (verification not implemented) 346

3.36.1 Optimal result

Integrand size = 32, antiderivative size = 74

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx = -\frac{3c^2 \operatorname{arctanh}(\sin(e+fx))}{af} + \frac{3c^2 \tan(e+fx)}{af} + \frac{2(c^2 - c^2 \sec(e+fx)) \tan(e+fx)}{f(a+a\sec(e+fx))}$$

output `-3*c^2*arctanh(sin(f*x+e))/a/f+3*c^2*tan(f*x+e)/a/f+2*(c^2-c^2*sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))`

3.36.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.44 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx = \frac{4\sqrt{2}c^2 \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan\left(\frac{1}{2}(e+fx)\right)}{af\sqrt{1-\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^2)/(a + a*Sec[e + f*x]),x]`

output `(4*sqrt(2)*c^2*Hypergeometric2F1[-3/2, -1/2, 1/2, (1 + Sec[e + f*x])/2]*Tan[(e + f*x)/2])/(a*f*sqrt[1 - Sec[e + f*x]])`

3.36.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.95, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4445, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2}{a\csc(e+fx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{4445} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c\int\sec(e+fx)(c-c\sec(e+fx))dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c\int\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))dx}{a} \\
 & \quad \downarrow \text{4274} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c(c\int\sec(e+fx)dx - c\int\sec^2(e+fx)dx)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c(c\int\csc(e+fx+\frac{\pi}{2})dx - c\int\csc(e+fx+\frac{\pi}{2})^2dx)}{a} \\
 & \quad \downarrow \text{4254} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c\left(\frac{c\int 1d(-\tan(e+fx))}{f} + c\int\csc(e+fx+\frac{\pi}{2})dx\right)}{a} \\
 & \quad \downarrow \text{24} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c\left(c\int\csc(e+fx+\frac{\pi}{2})dx - \frac{c\tan(e+fx)}{f}\right)}{a} \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

$$\frac{2 \tan(e + fx) (c^2 - c^2 \sec(e + fx))}{f(a \sec(e + fx) + a)} - \frac{3c \left(\frac{\operatorname{arctanh}(\sin(e + fx))}{f} - \frac{c \tan(e + fx)}{f} \right)}{a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^2)/(a + a*Sec[e + f*x]),x]`

output `(2*(c^2 - c^2*Sec[e + f*x])*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (3*c*((c*ArcTanh[Sin[e + f*x]])/f - (c*Tan[e + f*x])/f))/a`

3.36.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.36.4 Maple [A] (verified)

Time = 0.96 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.08

method	result
derivativedivides	$\frac{4c^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} - \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} \right)}{fa}$
default	$\frac{4c^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} - \frac{3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} \right)}{fa}$
parallelrirsch	$\frac{c^2 \left(3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) \cos(fx+e) - 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \cos(fx+e) + 5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \cos(fx+e) + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \right)}{af \cos(fx+e)}$
risch	$\frac{2ic^2(4e^{2i(fx+e)} + e^{i(fx+e)} + 5)}{fa(1+e^{2i(fx+e)})(e^{i(fx+e)} + 1)} + \frac{3c^2 \ln(e^{i(fx+e)} - i)}{af} - \frac{3c^2 \ln(e^{i(fx+e)} + i)}{af}$
norman	$\frac{6c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{10c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} + \frac{4c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2} + \frac{3c^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{af} - \frac{3c^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{af}$

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 4/f/a*c^2*(tan(1/2*f*x+1/2*e)-1/4/(tan(1/2*f*x+1/2*e)-1)+3/4*ln(tan(1/2*f*x+1/2*e)-1)-1/4/(tan(1/2*f*x+1/2*e)+1)-3/4*ln(tan(1/2*f*x+1/2*e)+1))
```

3.36.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.61

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx = \frac{3(c^2 \cos^2(fx+e) + c^2 \cos(fx+e)) \log(\sin(fx+e) + 1) - 3(c^2 \cos^2(fx+e) + c^2 \cos(fx+e)) \log(-\sin(fx+e) + 1) - 2(5c^2 \cos(fx+e) + c^2) \sin(fx+e)}{2(af \cos^2(fx+e) + af \cos(fx+e))}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e)),x, algorithm="fricas")
```

```
output -1/2*(3*(c^2*cos(f*x + e)^2 + c^2*cos(f*x + e))*log(sin(f*x + e) + 1) - 3*(c^2*cos(f*x + e)^2 + c^2*cos(f*x + e))*log(-sin(f*x + e) + 1) - 2*(5*c^2*cos(f*x + e) + c^2)*sin(f*x + e))/(a*f*cos(f*x + e)^2 + a*f*cos(f*x + e))
```

3.36. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx$

3.36.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx$$

$$= \frac{c^2 \left(\int \frac{\sec(e+fx)}{\sec(e+fx)+1} dx + \int \left(-\frac{2\sec^2(e+fx)}{\sec(e+fx)+1} \right) dx + \int \frac{\sec^3(e+fx)}{\sec(e+fx)+1} dx \right)}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**2/(a+a*sec(f*x+e)),x)`

output `c**2*(Integral(sec(e+f*x)/(sec(e+f*x)+1),x)+Integral(-2*sec(e+f*x)**2/(sec(e+f*x)+1),x)+Integral(sec(e+f*x)**3/(sec(e+f*x)+1),x))/a`

3.36.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 224 vs. $2(75) = 150$.

Time = 0.23 (sec) , antiderivative size = 224, normalized size of antiderivative = 3.03

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx =$$

$$\frac{c^2 \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}+1\right)}{a} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}-1\right)}{a} - \frac{2\sin(fx+e)}{\left(a-\frac{a\sin(fx+e)^2}{(\cos(fx+e)+1)^2}\right)(\cos(fx+e)+1)} - \frac{\sin(fx+e)}{a(\cos(fx+e)+1)} \right) + 2c^2 \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)}{a} \right)}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `-(c^2*(log(sin(f*x+e)/(cos(f*x+e)+1)+1)/a - log(sin(f*x+e)/(cos(f*x+e)+1)-1)/a - 2*sin(f*x+e)/((a-a*sin(f*x+e)^2/(cos(f*x+e)+1)^2)*(cos(f*x+e)+1)) - sin(f*x+e)/(a*(cos(f*x+e)+1)))) + 2*c^2*(log(sin(f*x+e)/(cos(f*x+e)+1)+1)/a - log(sin(f*x+e)/(cos(f*x+e)+1)-1)/a - sin(f*x+e)/(a*(cos(f*x+e)+1))) - c^2*sin(f*x+e)/(a*(cos(f*x+e)+1)))/f`

3.36.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.31

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx$$

$$= \frac{\frac{3c^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a} - \frac{3c^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a} - \frac{4c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{a} + \frac{2c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)a}}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e)),x, algorithm="giac")`

output `-(3*c^2*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a - 3*c^2*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a - 4*c^2*tan(1/2*f*x + 1/2*e)/a + 2*c^2*tan(1/2*f*x + 1/2*e)/((tan(1/2*f*x + 1/2*e)^2 - 1)*a))/f`

3.36.9 Mupad [B] (verification not implemented)

Time = 13.12 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.04

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{a+a\sec(e+fx)} dx = \frac{4c^2 \tan(\frac{e}{2} + \frac{fx}{2})}{af} + \frac{2c^2 \tan(\frac{e}{2} + \frac{fx}{2})}{f(a - a \tan(\frac{e}{2} + \frac{fx}{2})^2)} - \frac{6c^2 \operatorname{atanh}(\tan(\frac{e}{2} + \frac{fx}{2}))}{af}$$

input `int((c - c/cos(e + f*x))^2/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output `(4*c^2*tan(e/2 + (f*x)/2))/(a*f) + (2*c^2*tan(e/2 + (f*x)/2))/(f*(a - a*tan(e/2 + (f*x)/2)^2)) - (6*c^2*atanh(tan(e/2 + (f*x)/2)))/(a*f)`

3.37 $\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$

3.37.1	Optimal result	347
3.37.2	Mathematica [A] (verified)	347
3.37.3	Rubi [A] (verified)	348
3.37.4	Maple [A] (verified)	349
3.37.5	Fricas [A] (verification not implemented)	350
3.37.6	Sympy [F]	350
3.37.7	Maxima [B] (verification not implemented)	350
3.37.8	Giac [A] (verification not implemented)	351
3.37.9	Mupad [B] (verification not implemented)	351

3.37.1 Optimal result

Integrand size = 30, antiderivative size = 41

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = -\frac{\operatorname{carctanh}(\sin(e+fx))}{af} + \frac{2c \tan(e+fx)}{f(a+a\sec(e+fx))}$$

output `-c*arctanh(sin(f*x+e))/a/f+2*c*tan(f*x+e)/f/(a+a*sec(f*x+e))`

3.37.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.88

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = -\frac{c\left(-\frac{\log(\cos(\frac{1}{2}(e+fx))-\sin(\frac{1}{2}(e+fx)))}{f} + \frac{\log(\cos(\frac{1}{2}(e+fx))+\sin(\frac{1}{2}(e+fx)))}{f} - \frac{2 \tan(\frac{1}{2}(e+fx))}{f}\right)}{a}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output `-((c*(-(Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]])/f) + Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]])/f - (2*Tan[(e + f*x)/2])/f)/a`

3.37.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4445, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a\sec(e+fx)+a} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))}{a\csc(e+fx+\frac{\pi}{2})+a} dx$$

↓ 4445

$$\frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)} - \frac{c\int\sec(e+fx)dx}{a}$$

↓ 3042

$$\frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)} - \frac{c\int\csc(e+fx+\frac{\pi}{2})dx}{a}$$

↓ 4257

$$\frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)} - \frac{\text{arctanh}(\sin(e+fx))}{af}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output `-((c*ArcTanh[Sin[e + f*x]])/(a*f)) + (2*c*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))`

3.37.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)*(d_.) + (c_)]^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.37.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.15

method	result	size
parallelrisch	$\frac{c\left(2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)\right)}{af}$	47
derivativedivides	$\frac{2c\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2}\right)}{fa}$	48
default	$\frac{2c\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2}\right)}{fa}$	48
risch	$\frac{4ic}{fa(e^{i(fx+e)}+1)} - \frac{c \ln(e^{i(fx+e)}+i)}{af} + \frac{c \ln(e^{i(fx+e)}-i)}{af}$	68
norman	$-\frac{2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} + \frac{c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{af} - \frac{c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{af}$	98

input `int(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `c*(2*tan(1/2*f*x+1/2*e)+ln(tan(1/2*f*x+1/2*e)-1)-ln(tan(1/2*f*x+1/2*e)+1))/a/f`

$$3.37. \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$$

3.37.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.71

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = \frac{(c\cos(fx+e)+c)\log(\sin(fx+e)+1) - (c\cos(fx+e)+c)\log(-\sin(fx+e)+1) - 4c\sin(fx+e)}{2(af\cos(fx+e)+af)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="fracas")`

output `-1/2*((c*cos(f*x + e) + c)*log(sin(f*x + e) + 1) - (c*cos(f*x + e) + c)*log(-sin(f*x + e) + 1) - 4*c*sin(f*x + e))/(a*f*cos(f*x + e) + a*f)`

3.37.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = -\frac{c\left(\int\left(-\frac{\sec(e+fx)}{\sec(e+fx)+1}\right) dx + \int\frac{\sec^2(e+fx)}{\sec(e+fx)+1} dx\right)}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x)`

output `-c*(Integral(-sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(sec(e + f*x)*2/(sec(e + f*x) + 1), x))/a`

3.37.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. $2(41) = 82$.

Time = 0.20 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.46

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = -\frac{c\left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}+1\right)}{a} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}-1\right)}{a} - \frac{\sin(fx+e)}{a(\cos(fx+e)+1)}\right) - \frac{c\sin(fx+e)}{a(\cos(fx+e)+1)}}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `-(c*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a - sin(f*x + e)/(a*(cos(f*x + e) + 1))) - c*sin(f*x + e)/(a*(cos(f*x + e) + 1)))/f`

3.37.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.41

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))}{a + a \sec(e + fx)} dx$$

$$= -\frac{\frac{c \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) + 1|)}{a} - \frac{c \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) - 1|)}{a} - \frac{2c \tan(\frac{1}{2} fx + \frac{1}{2} e)}{a}}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="giac")`

output `-(c*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a - c*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a - 2*c*tan(1/2*f*x + 1/2*e)/a)/f`

3.37.9 Mupad [B] (verification not implemented)

Time = 13.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))}{a + a \sec(e + fx)} dx = -\frac{2c (\operatorname{atanh}(\tan(\frac{e}{2} + \frac{fx}{2})) - \tan(\frac{e}{2} + \frac{fx}{2}))}{af}$$

input `int((c - c/cos(e + f*x))/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output `-(2*c*(atanh(tan(e/2 + (f*x)/2)) - tan(e/2 + (f*x)/2)))/(a*f)`

$$3.38 \quad \int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))} dx$$

3.38.1	Optimal result	352
3.38.2	Mathematica [A] (verified)	352
3.38.3	Rubi [A] (verified)	353
3.38.4	Maple [A] (verified)	354
3.38.5	Fricas [A] (verification not implemented)	355
3.38.6	Sympy [F]	355
3.38.7	Maxima [A] (verification not implemented)	355
3.38.8	Giac [A] (verification not implemented)	356
3.38.9	Mupad [B] (verification not implemented)	356

3.38.1 Optimal result

Integrand size = 32, antiderivative size = 16

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))} dx = \frac{\csc(e+fx)}{acf}$$

output `csc(f*x+e)/a/c/f`

3.38.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))} dx = \frac{\csc(e+fx)}{acf}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])),x]`

output `Csc[e + f*x]/(a*c*f)`

3.38.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4446, 3042, 25, 3086, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)(c-c \sec(e+fx))} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)(c-c \csc(e+fx+\frac{\pi}{2}))} dx$$

↓ 4446

$$-\frac{\int \cot(e+fx) \csc(e+fx) dx}{ac}$$

↓ 3042

$$-\frac{\int -\sec(e+fx-\frac{\pi}{2}) \tan(e+fx-\frac{\pi}{2}) dx}{ac}$$

↓ 25

$$\frac{\int \sec(\frac{1}{2}(2e-\pi)+fx) \tan(\frac{1}{2}(2e-\pi)+fx) dx}{ac}$$

↓ 3086

$$\frac{\int 1 d \csc(e+fx)}{acf}$$

↓ 24

$$\frac{\csc(e+fx)}{acf}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])),x]`

output `Csc[e + f*x]/(a*c*f)`

3.38. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))} dx$

3.38.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`
- rule 4446 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^(m) Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.38.4 Maple [A] (verified)

Time = 0.99 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
default	$\frac{\csc(fx+e)}{acf}$	17
parallelrisch	$\frac{\sec\left(\frac{fx}{2} + \frac{e}{2}\right) \csc\left(\frac{fx}{2} + \frac{e}{2}\right)}{2acf}$	30
norman	$\frac{\frac{1}{2acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{2acf}}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}$	47
risch	$\frac{2ie^{i(fx+e)}}{fac(e^{i(fx+e)}-1)(e^{i(fx+e)}+1)}$	48

input `int(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

3.38. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))} dx$

output `csc(f*x+e)/a/c/f`

3.38.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))} dx = \frac{1}{acf \sin(fx + e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `1/(a*c*f*sin(f*x + e))`

3.38.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))} dx = -\frac{\int \frac{\sec(e+fx)}{\sec^2(e+fx)-1} dx}{ac}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**2 - 1), x)/(a*c)`

3.38.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))} dx = \frac{1}{acf \sin(fx + e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `1/(a*c*f*sin(f*x + e))`

3.38. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))} dx$

3.38.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))} dx = \frac{1}{acf \sin(fx + e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e)),x, algorithm="giac")`

output `1/(a*c*f*sin(f*x + e))`

3.38.9 Mupad [B] (verification not implemented)

Time = 12.92 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))} dx = \frac{1}{acf \sin(e + fx)}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))),x)`

output `1/(a*c*f*sin(e + f*x))`

$$3.39 \quad \int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^2} dx$$

3.39.1	Optimal result	357
3.39.2	Mathematica [A] (verified)	357
3.39.3	Rubi [A] (verified)	358
3.39.4	Maple [A] (verified)	359
3.39.5	Fricas [A] (verification not implemented)	359
3.39.6	Sympy [F]	360
3.39.7	Maxima [A] (verification not implemented)	360
3.39.8	Giac [A] (verification not implemented)	360
3.39.9	Mupad [B] (verification not implemented)	361

3.39.1 Optimal result

Integrand size = 32, antiderivative size = 59

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^2} dx = -\frac{\cot^3(e+fx)}{3ac^2f} + \frac{\csc(e+fx)}{ac^2f} - \frac{\csc^3(e+fx)}{3ac^2f}$$

output `-1/3*cot(f*x+e)^3/a/c^2/f+csc(f*x+e)/a/c^2/f-1/3*csc(f*x+e)^3/a/c^2/f`

3.39.2 Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^2} dx$$

$$= -\frac{(-3+4\cos(e+fx)+\cos(2(e+fx)))\csc^3\left(\frac{1}{2}(e+fx)\right)\sec\left(\frac{1}{2}(e+fx)\right)}{24ac^2f}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^2),x]`

output `-1/24*((-3 + 4*Cos[e + f*x] + Cos[2*(e + f*x)])*Csc[(e + f*x)/2]^3*Sec[(e + f*x)/2])/(a*c^2*f)`

$$3.39. \quad \int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^2} dx$$

3.39.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.86, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)(c-c\sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a\csc(e+fx+\frac{\pi}{2})+a)(c-c\csc(e+fx+\frac{\pi}{2}))^2} dx$$

↓ 4446

$$\int \frac{(a\csc(e+fx)\cot^3(e+fx)+a\csc^2(e+fx)\cot^2(e+fx)) dx}{a^2c^2}$$

↓ 2009

$$\frac{-\frac{a\cot^3(e+fx)}{3f}-\frac{a\csc^3(e+fx)}{3f}+\frac{a\csc(e+fx)}{f}}{a^2c^2}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^2),x]`

output `(-1/3*(a*Cot[e + f*x]^3)/f + (a*Csc[e + f*x])/f - (a*Csc[e + f*x]^3)/(3*f))/(a^2*c^2)`

3.39.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.39.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{2}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{4fa^2c^2}$	48
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{2}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{4fa^2c^2}$	48
parallelrisc	$\frac{3 \sec\left(\frac{fx}{2} + \frac{e}{2}\right) \csc\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - 4 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{12fa^2c^2}$	48
norman	$\frac{-\frac{1}{12acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{2acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{4acf}}{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}$	72
risc	$\frac{2i(3e^{3i(fx+e)} - 3e^{2i(fx+e)} + e^{i(fx+e)} + 1)}{3fa^2c^2(e^{i(fx+e)} - 1)^3(e^{i(fx+e)} + 1)}$	72

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE
)
```

```
output 1/4/f/a/c^2*(tan(1/2*f*x+1/2*e)-1/3/tan(1/2*f*x+1/2*e)^3+2/tan(1/2*f*x+1/2
*e))
```

3.39.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.85

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^2} dx = \frac{\cos(fx + e)^2 + 2 \cos(fx + e) - 2}{3(ac^2 f \cos(fx + e) - ac^2 f) \sin(fx + e)}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x, algorithm="fri
cas")
```

3.39.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^2} dx$$

output $1/3*(\cos(f*x + e)^2 + 2*\cos(f*x + e) - 2)/((a*c^2*f*\cos(f*x + e) - a*c^2*f)*\sin(f*x + e))$

3.39.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^2} dx = \frac{\int \frac{\sec(e + fx)}{\sec^3(e + fx) - \sec^2(e + fx) - \sec(e + fx) + 1} dx}{ac^2}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**2,x)`

output `Integral(sec(e + f*x)/(sec(e + f*x)**3 - sec(e + f*x)**2 - sec(e + f*x) + 1), x)/(a*c**2)`

3.39.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.31

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^2} dx = \frac{\left(\frac{6 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - 1\right)(\cos(fx+e)+1)^3}{ac^2 \sin^3(fx+e)} + \frac{3 \sin(fx+e)}{ac^2(\cos(fx+e)+1)}{12 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output $1/12*((6*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - 1)*(\cos(f*x + e) + 1)^3/(a*c^2*\sin(f*x + e)^3) + 3*\sin(f*x + e)/(a*c^2*(\cos(f*x + e) + 1)))/f$

3.39.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^2} dx = \frac{\frac{3 \tan(\frac{1}{2} fx + \frac{1}{2} e)}{ac^2} + \frac{6 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 1}{ac^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3}}{12 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `1/12*(3*tan(1/2*f*x + 1/2*e)/(a*c^2) + (6*tan(1/2*f*x + 1/2*e)^2 - 1)/(a*c^2*tan(1/2*f*x + 1/2*e)^3))/f`

3.39.9 Mupad [B] (verification not implemented)

Time = 13.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.85

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^2} dx = \frac{3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1}{12 a c^2 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^2),x)`

output `(6*tan(e/2 + (f*x)/2)^2 + 3*tan(e/2 + (f*x)/2)^4 - 1)/(12*a*c^2*f*tan(e/2 + (f*x)/2)^3)`

3.40 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^3} dx$

3.40.1	Optimal result	362
3.40.2	Mathematica [A] (verified)	362
3.40.3	Rubi [A] (verified)	363
3.40.4	Maple [A] (verified)	364
3.40.5	Fricas [A] (verification not implemented)	364
3.40.6	Sympy [F]	365
3.40.7	Maxima [A] (verification not implemented)	365
3.40.8	Giac [A] (verification not implemented)	366
3.40.9	Mupad [B] (verification not implemented)	366

3.40.1 Optimal result

Integrand size = 32, antiderivative size = 78

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^3} dx = \frac{2 \cot^5(e+fx)}{5ac^3f} + \frac{\csc(e+fx)}{ac^3f} - \frac{\csc^3(e+fx)}{ac^3f} + \frac{2 \csc^5(e+fx)}{5ac^3f}$$

output `2/5*cot(f*x+e)^5/a/c^3/f+csc(f*x+e)/a/c^3/f-csc(f*x+e)^3/a/c^3/f+2/5*csc(f*x+e)^5/a/c^3/f`

3.40.2 Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.78

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^3} dx = \frac{(5 - 5 \cos(e+fx) + \cos(2(e+fx)) + \cos(3(e+fx))) \csc^5(\frac{1}{2}(e+fx)) \sec(\frac{1}{2}(e+fx))}{80ac^3f}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^3),x]`

output `((5 - 5*Cos[e + f*x] + Cos[2*(e + f*x)] + Cos[3*(e + f*x)])*Csc[(e + f*x)/2]^5*Sec[(e + f*x)/2])/(80*a*c^3*f)`

3.40. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^3} dx$

3.40.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)(c-c \sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)(c-c \csc(e+fx+\frac{\pi}{2}))^3} dx$$

↓ 4446

$$\frac{\int (a^2 \csc(e+fx) \cot^5(e+fx) + 2a^2 \csc^2(e+fx) \cot^4(e+fx) + a^2 \csc^3(e+fx) \cot^3(e+fx)) dx}{a^3 c^3}$$

↓ 2009

$$\frac{-\frac{2a^2 \cot^5(e+fx)}{5f} - \frac{2a^2 \csc^5(e+fx)}{5f} + \frac{a^2 \csc^3(e+fx)}{f} - \frac{a^2 \csc(e+fx)}{f}}{a^3 c^3}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^3),x]`

output `-(((-2*a^2*Cot[e + f*x]^5)/(5*f) - (a^2*Csc[e + f*x])/f + (a^2*Csc[e + f*x]^3)/f - (2*a^2*Csc[e + f*x]^5)/(5*f))/(a^3*c^3))`

3.40.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.40. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^3} dx$

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.40.4 Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.76

method	result	size
parallelsch	$\frac{\cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 5\cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 5\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 15\cot\left(\frac{fx}{2} + \frac{e}{2}\right)}{40fac^3}$	59
derivativdivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{5\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{3}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{8fac^3}$	61
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{5\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{3}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{8fac^3}$	61
risch	$\frac{2i(5e^{5i(fx+e)} - 10e^{4i(fx+e)} + 10e^{3i(fx+e)} - 3e^{i(fx+e)} + 2)}{5fac^3(e^{i(fx+e)} + 1)(e^{i(fx+e)} - 1)^5}$	85
norman	$\frac{\frac{1}{40acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{8acf} + \frac{3\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{8acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{8acf}}{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	94

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE
)
```

```
output 1/40*(cot(1/2*f*x+1/2*e)^5-5*cot(1/2*f*x+1/2*e)^3+5*tan(1/2*f*x+1/2*e)+15*
cot(1/2*f*x+1/2*e))/f/a/c^3
```

3.40.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^3} dx$$

$$= \frac{2 \cos(fx + e)^3 + \cos(fx + e)^2 - 4 \cos(fx + e) + 2}{5 (ac^3 f \cos(fx + e)^2 - 2ac^3 f \cos(fx + e) + ac^3 f) \sin(fx + e)}$$

3.40. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^3} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x, algorithm="fricas")`

output `1/5*(2*cos(f*x + e)^3 + cos(f*x + e)^2 - 4*cos(f*x + e) + 2)/((a*c^3*f*cos(f*x + e)^2 - 2*a*c^3*f*cos(f*x + e) + a*c^3*f)*sin(f*x + e))`

3.40.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^3} dx = -\frac{\int \frac{\sec(e+fx)}{\sec^4(e+fx)-2\sec^3(e+fx)+2\sec(e+fx)-1} dx}{ac^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**3,x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**4 - 2*sec(e + f*x)**3 + 2*sec(e + f*x) - 1), x)/(a*c**3)`

3.40.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.24

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^3} dx$$

$$= -\frac{\left(\frac{5\sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{15\sin^4(fx+e)}{(\cos(fx+e)+1)^4} - 1\right)(\cos(fx+e)+1)^5}{ac^3\sin^5(fx+e)} - \frac{5\sin(fx+e)}{ac^3(\cos(fx+e)+1)}$$

$$40f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `-1/40*((5*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 15*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 1)*(cos(f*x + e) + 1)^5/(a*c^3*sin(f*x + e)^5) - 5*sin(f*x + e)/(a*c^3*(cos(f*x + e) + 1)))/f`

3.40.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.88

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^3} dx$$

$$= \frac{\frac{5 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{ac^3} + \frac{15 \tan(\frac{1}{2}fx + \frac{1}{2}e)^4 - 5 \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 + 1}{ac^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5}}{40f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `1/40*(5*tan(1/2*f*x + 1/2*e)/(a*c^3) + (15*tan(1/2*f*x + 1/2*e)^4 - 5*tan(1/2*f*x + 1/2*e)^2 + 1)/(a*c^3*tan(1/2*f*x + 1/2*e)^5))/f`

3.40.9 Mupad [B] (verification not implemented)

Time = 13.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.81

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^3} dx$$

$$= \frac{5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 15 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1}{40 a c^3 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^3),x)`

output `(15*tan(e/2 + (f*x)/2)^4 - 5*tan(e/2 + (f*x)/2)^2 + 5*tan(e/2 + (f*x)/2)^6 + 1)/(40*a*c^3*f*tan(e/2 + (f*x)/2)^5)`

$$3.41 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx$$

3.41.1	Optimal result	367
3.41.2	Mathematica [A] (verified)	367
3.41.3	Rubi [A] (verified)	368
3.41.4	Maple [A] (verified)	369
3.41.5	Fricas [A] (verification not implemented)	370
3.41.6	Sympy [F]	370
3.41.7	Maxima [A] (verification not implemented)	370
3.41.8	Giac [A] (verification not implemented)	371
3.41.9	Mupad [B] (verification not implemented)	371

3.41.1 Optimal result

Integrand size = 32, antiderivative size = 120

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx = -\frac{\cot^5(e+fx)}{5ac^4f} - \frac{4 \cot^7(e+fx)}{7ac^4f} + \frac{\csc(e+fx)}{ac^4f} - \frac{2 \csc^3(e+fx)}{ac^4f} + \frac{9 \csc^5(e+fx)}{5ac^4f} - \frac{4 \csc^7(e+fx)}{7ac^4f}$$

output `-1/5*cot(f*x+e)^5/a/c^4/f-4/7*cot(f*x+e)^7/a/c^4/f+csc(f*x+e)/a/c^4/f-2*csc(f*x+e)^3/a/c^4/f+9/5*csc(f*x+e)^5/a/c^4/f-4/7*csc(f*x+e)^7/a/c^4/f`

3.41.2 Mathematica [A] (verified)

Time = 1.53 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx = \frac{(-13 + 4 \sec(e+fx) + 20 \sec^2(e+fx) - 24 \sec^3(e+fx) + 8 \sec^4(e+fx)) \tan(e+fx)}{35ac^4f(-1 + \sec(e+fx))^4(1 + \sec(e+fx))}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^4),x]`

3.41. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx$


```
output ((-13 + 4*Sec[e + f*x] + 20*Sec[e + f*x]^2 - 24*Sec[e + f*x]^3 + 8*Sec[e +
f*x]^4)*Tan[e + f*x])/(35*a*c^4*f*(-1 + Sec[e + f*x])^4*(1 + Sec[e + f*x]
))
```

3.41.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.91, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)(c - c \sec(e+fx))^4} dx$$

↓ 3042

$$\int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)(c - c \csc(e+fx + \frac{\pi}{2}))^4} dx$$

↓ 4446

$$\int \frac{(a^3 \csc(e+fx) \cot^7(e+fx) + 3a^3 \csc^2(e+fx) \cot^6(e+fx) + 3a^3 \csc^3(e+fx) \cot^5(e+fx) + a^3 \csc^4(e+fx))}{a^4 c^4} dx$$

↓ 2009

$$\frac{-\frac{4a^3 \cot^7(e+fx)}{7f} - \frac{a^3 \cot^5(e+fx)}{5f} - \frac{4a^3 \csc^7(e+fx)}{7f} + \frac{9a^3 \csc^5(e+fx)}{5f} - \frac{2a^3 \csc^3(e+fx)}{f} + \frac{a^3 \csc(e+fx)}{f}}{a^4 c^4}$$

```
input Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^4),x]
```

```
output (-1/5*(a^3*Cot[e + f*x]^5)/f - (4*a^3*Cot[e + f*x]^7)/(7*f) + (a^3*Csc[e +
f*x])/f - (2*a^3*Csc[e + f*x]^3)/f + (9*a^3*Csc[e + f*x]^5)/(5*f) - (4*a^
3*Csc[e + f*x]^7)/(7*f))/(a^4*c^4)
```

3.41. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx$

3.41.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4446 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.41.4 Maple [A] (verified)

Time = 0.89 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.62

method	result	size
derivativedivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{4}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{2}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{4}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}}{16fac^4}$	74
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{4}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{2}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{4}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}}{16fac^4}$	74
parallelrisch	$\frac{-5 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^7 + 28 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 70 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 140 \cot\left(\frac{fx}{2} + \frac{e}{2}\right) + 35 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{560fac^4}$	74
norman	$\frac{-\frac{1}{112acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{20acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{8acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{4acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{16acf}}{c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$	116
risch	$\frac{2i(35e^{7i(fx+e)} - 105e^{6i(fx+e)} + 175e^{5i(fx+e)} - 105e^{4i(fx+e)} - 7e^{3i(fx+e)} + 77e^{2i(fx+e)} - 43e^{i(fx+e)} + 13)}{35fac^4(e^{i(fx+e)} - 1)^7(e^{i(fx+e)} + 1)}$	118

input `int(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x,method=_RETURNVERBOSE)`

output `1/16/f/a/c^4*(tan(1/2*f*x+1/2*e)+4/5/tan(1/2*f*x+1/2*e)^5-2/tan(1/2*f*x+1/2*e)^3+4/tan(1/2*f*x+1/2*e)-1/7/tan(1/2*f*x+1/2*e)^7)`

3.41.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx$$

3.41.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.85

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^4} dx$$

$$= \frac{13 \cos(fx + e)^4 - 4 \cos(fx + e)^3 - 20 \cos(fx + e)^2 + 24 \cos(fx + e) - 8}{35 (ac^4 f \cos(fx + e)^3 - 3ac^4 f \cos(fx + e)^2 + 3ac^4 f \cos(fx + e) - ac^4 f) \sin(fx + e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x, algorithm="fricas")`

output `1/35*(13*cos(f*x + e)^4 - 4*cos(f*x + e)^3 - 20*cos(f*x + e)^2 + 24*cos(f*x + e) - 8)/((a*c^4*f*cos(f*x + e)^3 - 3*a*c^4*f*cos(f*x + e)^2 + 3*a*c^4*f*cos(f*x + e) - a*c^4*f)*sin(f*x + e))`

3.41.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^4} dx$$

$$= \int \frac{\sec(e + fx)}{\sec^5(e + fx) - 3 \sec^4(e + fx) + 2 \sec^3(e + fx) + 2 \sec^2(e + fx) - 3 \sec(e + fx) + 1} dx$$

$$ac^4$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**4,x)`

output `Integral(sec(e + f*x)/(sec(e + f*x)**5 - 3*sec(e + f*x)**4 + 2*sec(e + f*x)**3 + 2*sec(e + f*x)**2 - 3*sec(e + f*x) + 1), x)/(a*c**4)`

3.41.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.98

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^4} dx$$

$$= \frac{\left(\frac{28 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{70 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{140 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - 5 \right) (\cos(fx+e)+1)^7}{ac^4 \sin(fx+e)^7} + \frac{35 \sin(fx+e)}{ac^4 (\cos(fx+e)+1)}$$

$$560 f$$

3.41. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output $\frac{1}{560} * ((28 * \sin(f * x + e) ^ 2 / (\cos(f * x + e) + 1) ^ 2 - 70 * \sin(f * x + e) ^ 4 / (\cos(f * x + e) + 1) ^ 4 + 140 * \sin(f * x + e) ^ 6 / (\cos(f * x + e) + 1) ^ 6 - 5) * (\cos(f * x + e) + 1) ^ 7 / (a * c ^ 4 * \sin(f * x + e) ^ 7) + 35 * \sin(f * x + e) / (a * c ^ 4 * (\cos(f * x + e) + 1))) / f$

3.41.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.68

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^4} dx$$

$$= \frac{\frac{35 \tan(\frac{1}{2} fx + \frac{1}{2} e)}{ac^4} + \frac{140 \tan(\frac{1}{2} fx + \frac{1}{2} e)^6 - 70 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 + 28 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 5}{ac^4 \tan(\frac{1}{2} fx + \frac{1}{2} e)^7}}{560 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output $\frac{1}{560} * (35 * \tan(1/2 * f * x + 1/2 * e) / (a * c ^ 4) + (140 * \tan(1/2 * f * x + 1/2 * e) ^ 6 - 70 * \tan(1/2 * f * x + 1/2 * e) ^ 4 + 28 * \tan(1/2 * f * x + 1/2 * e) ^ 2 - 5) / (a * c ^ 4 * \tan(1/2 * f * x + 1/2 * e) ^ 7)) / f$

3.41.9 Mupad [B] (verification not implemented)

Time = 13.44 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.69

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^4} dx$$

$$= \frac{\tan(\frac{e}{2} + \frac{fx}{2})}{16 a c^4 f} + \frac{\frac{\tan(\frac{e}{2} + \frac{fx}{2})^6}{4} - \frac{\tan(\frac{e}{2} + \frac{fx}{2})^4}{8} + \frac{\tan(\frac{e}{2} + \frac{fx}{2})^2}{20} - \frac{1}{112}}{a c^4 f \tan(\frac{e}{2} + \frac{fx}{2})^7}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^4),x)`

output $\frac{\tan(e/2 + (f*x)/2)}{16 * a * c ^ 4 * f} + (\tan(e/2 + (f*x)/2) ^ 2 / 20 - \tan(e/2 + (f*x)/2) ^ 4 / 8 + \tan(e/2 + (f*x)/2) ^ 6 / 4 - 1 / 112) / (a * c ^ 4 * f * \tan(e/2 + (f*x)/2) ^ 7)$

3.41. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^4} dx$

$$3.42 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$$

3.42.1	Optimal result	372
3.42.2	Mathematica [C] (verified)	373
3.42.3	Rubi [A] (verified)	373
3.42.4	Maple [A] (verified)	375
3.42.5	Fricas [A] (verification not implemented)	376
3.42.6	Sympy [F]	376
3.42.7	Maxima [B] (verification not implemented)	377
3.42.8	Giac [A] (verification not implemented)	378
3.42.9	Mupad [B] (verification not implemented)	379

3.42.1 Optimal result

Integrand size = 32, antiderivative size = 164

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx = \frac{105c^5 \operatorname{arctanh}(\sin(e+fx))}{2a^2 f} - \frac{84c^5 \tan(e+fx)}{a^2 f} + \frac{63c^5 \sec(e+fx) \tan(e+fx)}{2a^2 f} - \frac{6c^2 (c-c\sec(e+fx))^3 \tan(e+fx)}{f(a^2+a^2\sec(e+fx))} + \frac{2c(c-c\sec(e+fx))^4 \tan(e+fx)}{3f(a+a\sec(e+fx))^2} - \frac{7c^5 \tan^3(e+fx)}{a^2 f}$$

output `105/2*c^5*arctanh(sin(f*x+e))/a^2/f-84*c^5*tan(f*x+e)/a^2/f+63/2*c^5*sec(f*x+e)*tan(f*x+e)/a^2/f-6*c^2*(c-c*sec(f*x+e))^3*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))+2/3*c*(c-c*sec(f*x+e))^4*tan(f*x+e)/f/(a+a*sec(f*x+e))^2-7*c^5*tan(f*x+e)^3/a^2/f`

3.42.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.92 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.46

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx = \frac{32c^5 \operatorname{Hypergeometric2F1}\left(-\frac{9}{2}, -\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)} \tan(e+fx)}{3a^2 f(-1+\sec(e+fx))(1+\sec(e+fx))^2}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^2,x]`

output `(-32*c^5*Hypergeometric2F1[-9/2, -3/2, -1/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]]*Tan[e + f*x])/(3*a^2*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^2)`

3.42.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4445, 3042, 4445, 3042, 4278, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a\sec(e+fx)+a)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^5}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx \\ & \quad \downarrow \text{4445} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^4}{3f(a\sec(e+fx)+a)^2} - \frac{3c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{\sec(e+fx)a+a} dx}{a} \\ & \quad \downarrow \text{3042} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^4}{3f(a\sec(e+fx)+a)^2} - \frac{3c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^4}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{a} \end{aligned}$$

3.42. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$

$$\begin{aligned}
 & \downarrow 4445 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{f(a \sec(e+fx)+a)} - \frac{7c \int \sec(e+fx)(c-c \sec(e+fx))^3 dx}{a} \right)}{a} \\
 & \downarrow 3042 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{f(a \sec(e+fx)+a)} - \frac{7c \int \csc(e+fx+\frac{\pi}{2})(c-c \csc(e+fx+\frac{\pi}{2}))^3 dx}{a} \right)}{a} \\
 & \downarrow 4278 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{f(a \sec(e+fx)+a)} - \frac{7c \int (-c^3 \sec^4(e+fx)+3c^3 \sec^3(e+fx)-3c^3 \sec^2(e+fx)+c^3 \sec(e+fx)) dx}{a} \right)}{a} \\
 & \downarrow 2009 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{f(a \sec(e+fx)+a)} - \frac{7c \left(\frac{5c^3 \operatorname{arctanh}(\sin(e+fx))}{2f} - \frac{c^3 \tan^3(e+fx)}{3f} - \frac{4c^3 \tan(e+fx)}{f} + \frac{3c^3 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \right)}{a}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^2,x]`

output `(2*c*(c - c*Sec[e + f*x])^4*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (3*c*((2*c*(c - c*Sec[e + f*x])^3*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (7*c*((5*c^3*ArcTanh[Sin[e + f*x]])/(2*f) - (4*c^3*Tan[e + f*x])/f + (3*c^3*Sec[e + f*x]*Tan[e + f*x])/(2*f) - (c^3*Tan[e + f*x]^3)/(3*f))))/a)/a`

3.42.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4278 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n_*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_], x_Symbol] := Int[ExpandTrig[(a + b*csc[e + f*x])^m*(d*csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && I GtQ[m, 0] && RationalQ[n]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^n_], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2*(-1)] && IntegerQ[2*m]`

3.42.4 Maple [A] (verified)

Time = 1.45 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.95

method	result
derivativedivides	$16c^5 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{48\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^3} - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{55}{32\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} + \frac{105 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{32} \right) f a^2$
default	$16c^5 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{48\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^3} - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{55}{32\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} + \frac{105 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{32} \right) f a^2$
parallelrisc	$1969 \left(\frac{630(\cos(3fx+3e)+3\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{1969} + \frac{630(-\cos(3fx+3e)-3\cos(fx+e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{1969} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \right) f a^2 (\cos(3fx+3e)+3\cos(fx+e))$
risc	$- \frac{ic^5 (309 e^{8i(fx+e)} + 969 e^{7i(fx+e)} + 1693 e^{6i(fx+e)} + 3027 e^{5i(fx+e)} + 2901 e^{4i(fx+e)} + 3247 e^{3i(fx+e)} + 1995 e^{2i(fx+e)} + 630 e^{i(fx+e)} + 105)}{3a^2 f (1+e^{2i(fx+e)})^3 (e^{i(fx+e)}+1)^3}$
norman	$\frac{105c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} - \frac{490c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} + \frac{896c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{af} - \frac{790c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{af} + \frac{965c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{3af} - \frac{112c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{3af} \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1 \right)^5 a$

3.42. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `16/f*c^5/a^2*(-1/3*tan(1/2*f*x+1/2*e)^3-4*tan(1/2*f*x+1/2*e)+1/48/(tan(1/2*f*x+1/2*e)+1)^3-1/4/(tan(1/2*f*x+1/2*e)+1)^2+55/32/(tan(1/2*f*x+1/2*e)+1)+105/32*ln(tan(1/2*f*x+1/2*e)+1)+1/48/(tan(1/2*f*x+1/2*e)-1)^3+1/4/(tan(1/2*f*x+1/2*e)-1)^2+55/32/(tan(1/2*f*x+1/2*e)-1)-105/32*ln(tan(1/2*f*x+1/2*e)-1))`

3.42.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.28

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx = \frac{315(c^5 \cos(fx+e)^5 + 2c^5 \cos(fx+e)^4 + c^5 \cos(fx+e)^3) \log(\sin(fx+e)+1) - 315(c^5 \cos(fx+e)^5)}{a^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x, algorithm="fricas")`

output `1/12*(315*(c^5*cos(f*x + e)^5 + 2*c^5*cos(f*x + e)^4 + c^5*cos(f*x + e)^3)*log(sin(f*x + e) + 1) - 315*(c^5*cos(f*x + e)^5 + 2*c^5*cos(f*x + e)^4 + c^5*cos(f*x + e)^3)*log(-sin(f*x + e) + 1) - 2*(494*c^5*cos(f*x + e)^4 + 679*c^5*cos(f*x + e)^3 + 102*c^5*cos(f*x + e)^2 - 17*c^5*cos(f*x + e) + 2*c^5)*sin(f*x + e))/(a^2*f*cos(f*x + e)^5 + 2*a^2*f*cos(f*x + e)^4 + a^2*f*cos(f*x + e)^3)`

3.42.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx = \frac{c^5 \left(\int \left(-\frac{\sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} \right) dx + \int \frac{5\sec^2(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \left(-\frac{10\sec^3(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} \right) dx + \dots \right)}{a^2}$$

3.42. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**5/(a+a*sec(f*x+e))**2,x)`

output `-c**5*(Integral(-sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) +
Integral(5*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + I
ntegral(-10*sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + I
ntegral(10*sec(e + f*x)**4/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + In
tegral(-5*sec(e + f*x)**5/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Int
egral(sec(e + f*x)**6/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.42.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 765 vs. $2(160) = 320$.

Time = 0.23 (sec) , antiderivative size = 765, normalized size of antiderivative = 4.66

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^5}{(a + a\sec(e + fx))^2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x, algorithm="m
axima")`

output

```

-1/6*(c^5*(4*(9*sin(f*x + e)/(cos(f*x + e) + 1) - 20*sin(f*x + e)^3/(cos(f
*x + e) + 1)^3 + 15*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^2 - 3*a^2*sin(
f*x + e)^2/(cos(f*x + e) + 1)^2 + 3*a^2*sin(f*x + e)^4/(cos(f*x + e) + 1)^
4 - a^2*sin(f*x + e)^6/(cos(f*x + e) + 1)^6) + (27*sin(f*x + e)/(cos(f*x +
e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 30*log(sin(f*x + e)/
(cos(f*x + e) + 1) + 1)/a^2 + 30*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/
a^2) + 5*c^5*(6*(3*sin(f*x + e)/(cos(f*x + e) + 1) - 5*sin(f*x + e)^3/(cos
(f*x + e) + 1)^3)/(a^2 - 2*a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^2*s
in(f*x + e)^4/(cos(f*x + e) + 1)^4) + (21*sin(f*x + e)/(cos(f*x + e) + 1)
+ sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 21*log(sin(f*x + e)/(cos(f*x
+ e) + 1) + 1)/a^2 + 21*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2) + 10
*c^5*((15*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) +
1)^3)/a^2 - 12*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 12*log(sin(
f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 + 12*sin(f*x + e)/((a^2 - a^2*sin(f*x
+ e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1))) + 10*c^5*((9*sin(f*x +
e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 6*log(s
in(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 6*log(sin(f*x + e)/(cos(f*x + e)
+ 1) - 1)/a^2) + 5*c^5*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^
3/(cos(f*x + e) + 1)^3)/a^2 - c^5*(3*sin(f*x + e)/(cos(f*x + e) + 1) - sin
(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f

```

3.42.8 Giac [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^5}{(a + a\sec(e + fx))^2} dx$$

$$= \frac{315c^5 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{a^2} - \frac{315c^5 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{a^2} + \frac{2\left(165c^5 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 280c^5 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + 123c^5 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 1\right)^3 a^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output

```

1/6*(315*c^5*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^2 - 315*c^5*log(abs(tan(
1/2*f*x + 1/2*e) - 1))/a^2 + 2*(165*c^5*tan(1/2*f*x + 1/2*e)^5 - 280*c^5*t
an(1/2*f*x + 1/2*e)^3 + 123*c^5*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1/2*
e)^2 - 1)^3*a^2) - 32*(a^4*c^5*tan(1/2*f*x + 1/2*e)^3 + 12*a^4*c^5*tan(1/2
*f*x + 1/2*e))/a^6)/f

```

3.42.9 Mupad [B] (verification not implemented)

Time = 12.94 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.04

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{55c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 - \frac{280c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3} + 41c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - a^2 \right)}$$

$$- \frac{64c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a^2 f} - \frac{16c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3a^2 f} + \frac{105c^5 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a^2 f}$$

input `int((c - c/cos(e + f*x))^5/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`output `(55*c^5*tan(e/2 + (f*x)/2)^5 - (280*c^5*tan(e/2 + (f*x)/2)^3)/3 + 41*c^5*tan(e/2 + (f*x)/2))/(f*(3*a^2*tan(e/2 + (f*x)/2)^2 - 3*a^2*tan(e/2 + (f*x)/2)^4 + a^2*tan(e/2 + (f*x)/2)^6 - a^2)) - (64*c^5*tan(e/2 + (f*x)/2))/(a^2*f) - (16*c^5*tan(e/2 + (f*x)/2)^3)/(3*a^2*f) + (105*c^5*atanh(tan(e/2 + (f*x)/2)))/(a^2*f)`

$$3.43 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

3.43.1	Optimal result	380
3.43.2	Mathematica [C] (verified)	380
3.43.3	Rubi [A] (verified)	381
3.43.4	Maple [A] (verified)	384
3.43.5	Fricas [A] (verification not implemented)	385
3.43.6	Sympy [F]	385
3.43.7	Maxima [B] (verification not implemented)	386
3.43.8	Giac [A] (verification not implemented)	386
3.43.9	Mupad [B] (verification not implemented)	387

3.43.1 Optimal result

Integrand size = 32, antiderivative size = 150

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx = \frac{35c^4 \operatorname{arctanh}(\sin(e+fx))}{2a^2 f} - \frac{70c^4 \tan(e+fx)}{3a^2 f} + \frac{35c^4 \sec(e+fx) \tan(e+fx)}{6a^2 f} + \frac{2c(c-c\sec(e+fx))^3 \tan(e+fx)}{3f(a+a\sec(e+fx))^2} - \frac{14(c^2-c^2\sec(e+fx))^2 \tan(e+fx)}{3f(a^2+a^2\sec(e+fx))}$$

output `35/2*c^4*arctanh(sin(f*x+e))/a^2/f-70/3*c^4*tan(f*x+e)/a^2/f+35/6*c^4*sec(f*x+e)*tan(f*x+e)/a^2/f+2/3*c*(c-c*sec(f*x+e))^3*tan(f*x+e)/f/(a+a*sec(f*x+e))^2-14/3*(c^2-c^2*sec(f*x+e))^2*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))`

3.43.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.96 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.50

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx = \frac{16c^4 \operatorname{Hypergeometric2F1}\left(-\frac{7}{2}, -\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)} \tan(e+fx)}{3a^2 f(-1+\sec(e+fx))(1+\sec(e+fx))^2}$$

3.43. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^2,x]`

output `(-16*c^4*Hypergeometric2F1[-7/2, -3/2, -1/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]]*Tan[e + f*x])/(3*a^2*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^2)`

3.43.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.97, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 4445, 3042, 4445, 3042, 4275, 3042, 4254, 24, 4534, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a\sec(e+fx)+a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^4}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx \\
 & \quad \downarrow \text{4445} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{3f(a\sec(e+fx)+a)^2} - \frac{7c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{\sec(e+fx)a+a} dx}{3a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{3f(a\sec(e+fx)+a)^2} - \frac{7c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^3}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} \\
 & \quad \downarrow \text{4445} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{3f(a\sec(e+fx)+a)^2} - \frac{7c \left(\frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{f(a\sec(e+fx)+a)} - \frac{5c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^2 dx}{a} \right)}{3a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^3}{3f(a\sec(e+fx)+a)^2} - \frac{7c \left(\frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{f(a\sec(e+fx)+a)} - \frac{5c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2 dx}{a} \right)}{3a}
 \end{aligned}$$

3.43. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

$$\begin{aligned}
 & \downarrow 4275 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \\
 & \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \int \sec(e+fx)(\sec^2(e+fx)c^2+c^2) dx - 2c^2 \int \sec^2(e+fx) dx}{a} \right)}{3a} \\
 & \downarrow 3042 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \\
 & \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\int \csc(e+fx+\frac{\pi}{2}) (\csc(e+fx+\frac{\pi}{2})^2 c^2+c^2) dx - 2c^2 \int \csc(e+fx+\frac{\pi}{2})^2 dx \right)}{a} \right)}{3a} \\
 & \downarrow 4254 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \\
 & \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\frac{2c^2 \int 1d(-\tan(e+fx))}{f} + \int \csc(e+fx+\frac{\pi}{2}) (\csc(e+fx+\frac{\pi}{2})^2 c^2+c^2) dx \right)}{a} \right)}{3a} \\
 & \downarrow 24 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \\
 & \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\int \csc(e+fx+\frac{\pi}{2}) (\csc(e+fx+\frac{\pi}{2})^2 c^2+c^2) dx - \frac{2c^2 \tan(e+fx)}{f} \right)}{a} \right)}{3a} \\
 & \downarrow 4534 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \\
 & \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\frac{3}{2} c^2 \int \sec(e+fx) dx - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \right)}{3a} \\
 & \downarrow 3042 \\
 & \frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \\
 & \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\frac{3}{2} c^2 \int \csc(e+fx+\frac{\pi}{2}) dx - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \right)}{3a}
 \end{aligned}$$

3.43. $\int \frac{\sec(e+fx)(c-c \sec(e+fx))^4}{(a+a \sec(e+fx))^2} dx$

$$\begin{array}{c}
 \downarrow 4257 \\
 \frac{2c \tan(e+fx)(c - c \sec(e+fx))^3}{3f(a \sec(e+fx) + a)^2} - \\
 7c \left(\frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{f(a \sec(e+fx) + a)} - \frac{5c \left(\frac{3c^2 \operatorname{arctanh}(\sin(e+fx))}{2f} - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \right) \\
 \hline
 3a
 \end{array}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^2,x]`

output `(2*c*(c - c*Sec[e + f*x])^3*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (7*c*((2*c*(c - c*Sec[e + f*x])^2*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (5*c*((3*c^2*ArcTanh[Sin[e + f*x]])/(2*f) - (2*c^2*Tan[e + f*x])/f + (c^2*Sec[e + f*x]*Tan[e + f*x])/(2*f))))/a)/(3*a)`

3.43.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4275 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^2, x_Symbol] := Simp[2*a*(b/d) Int[(d*Csc[e + f*x])^(n + 1), x], x] + Int[(d*Csc[e + f*x])^n*(a^2 + b^2*Csc[e + f*x]^2), x] /; FreeQ[{a, b, d, e, f, n}, x]`


```
rule 4445 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]
```

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

3.43.4 Maple [A] (verified)

Time = 1.27 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.83

method	result
derivativedivides	$8c^4 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{16\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^2} + \frac{13}{16\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} - \frac{35 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{16} - \frac{1}{16\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)} \right) \frac{1}{fa^2}$
default	$8c^4 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{16\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^2} + \frac{13}{16\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} - \frac{35 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{16} - \frac{1}{16\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)} \right) \frac{1}{fa^2}$
parallelrisch	$51 \left(\frac{35(1 + \cos(2fx + 2e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{51} + \frac{35(-1 - \cos(2fx + 2e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{51} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \sec\left(\frac{fx}{2} + \frac{e}{2}\right)^2 \left(\cos(fx + e)\right) \right) \frac{1}{2fa^2(1 + \cos(2fx + 2e))}$
risch	$-\frac{ic^4(99e^{6i(fx+e)} + 333e^{5i(fx+e)} + 434e^{4i(fx+e)} + 714e^{3i(fx+e)} + 487e^{2i(fx+e)} + 393e^{i(fx+e)} + 164)}{3fa^2(1 + e^{2i(fx+e)})^2(e^{i(fx+e)} + 1)^3} + \frac{35c^4 \ln(e^{i(fx+e)} + 1)}{2a^2 f}$
norman	$-\frac{35c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{385c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} - \frac{511c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{3af} + \frac{93c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{af} - \frac{40c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{3af} - \frac{8c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{3af} \frac{1}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^4 a}$

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

3.43. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

output $8/f*c^4/a^2*(-1/3*\tan(1/2*f*x+1/2*e)^3-3*\tan(1/2*f*x+1/2*e)+1/16/(\tan(1/2*f*x+1/2*e)-1)^2+13/16/(\tan(1/2*f*x+1/2*e)-1)-35/16*\ln(\tan(1/2*f*x+1/2*e)-1)-1/16/(\tan(1/2*f*x+1/2*e)+1)^2+13/16/(\tan(1/2*f*x+1/2*e)+1)+35/16*\ln(\tan(1/2*f*x+1/2*e)+1))$

3.43.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.31

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{105(c^4 \cos(fx+e)^4 + 2c^4 \cos(fx+e)^3 + c^4 \cos(fx+e)^2) \log(\sin(fx+e)+1) - 105(c^4 \cos(fx+e)^4 + 2c^4 \cos(fx+e)^3 + c^4 \cos(fx+e)^2) \log(\sin(fx+e)-1)}{12(a^2 f c)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x, algorithm="fracas")`

output $1/12*(105*(c^4*\cos(f*x + e)^4 + 2*c^4*\cos(f*x + e)^3 + c^4*\cos(f*x + e)^2)*\log(\sin(f*x + e) + 1) - 105*(c^4*\cos(f*x + e)^4 + 2*c^4*\cos(f*x + e)^3 + c^4*\cos(f*x + e)^2)*\log(-\sin(f*x + e) + 1) - 2*(164*c^4*\cos(f*x + e)^3 + 229*c^4*\cos(f*x + e)^2 + 30*c^4*\cos(f*x + e) - 3*c^4)*\sin(f*x + e))/(a^2*f*\cos(f*x + e)^4 + 2*a^2*f*\cos(f*x + e)^3 + a^2*f*\cos(f*x + e)^2)$

3.43.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{c^4 \left(\int \frac{\sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \left(-\frac{4\sec^2(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} \right) dx + \int \frac{6\sec^3(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \left(-\frac{4\sec^4(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} \right) dx \right)}{a^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**4/(a+a*sec(f*x+e))**2,x)`

output `c**4*(Integral(sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(-4*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(6*sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(-4*sec(e + f*x)**4/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**5/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.43. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

3.43.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 531 vs. $2(142) = 284$.

Time = 0.21 (sec) , antiderivative size = 531, normalized size of antiderivative = 3.54

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx =$$

$$c^4 \left(\frac{6 \left(\frac{3 \sin(fx+e)}{\cos(fx+e)+1} - \frac{5 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^2 - \frac{2a^2 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{a^2 \sin(fx+e)^4}{(\cos(fx+e)+1)^4}} + \frac{21 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} - \frac{21 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^2} + \frac{21 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^2} \right) +$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output

```
-1/6*(c^4*(6*(3*sin(f*x + e)/(cos(f*x + e) + 1) - 5*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^2 - 2*a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^2*sin(f*x + e)^4/(cos(f*x + e) + 1)^4) + (21*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 21*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 21*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 + 4*c^4*((15*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 12*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 12*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 + 12*sin(f*x + e)/((a^2 - a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1))) + 6*c^4*((9*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 6*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 6*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2) + 4*c^4*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - c^4*(3*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f
```

3.43.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.93

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{105c^4 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{a^2} - \frac{105c^4 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{a^2} + \frac{6\left(13c^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 - 11c^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 1\right)^2 a^2} - \frac{16\left(a^4 c^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{6f}$$

3.43. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output `1/6*(105*c^4*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^2 - 105*c^4*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a^2 + 6*(13*c^4*tan(1/2*f*x + 1/2*e)^3 - 11*c^4*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1/2*e)^2 - 1)^2*a^2) - 16*(a^4*c^4*tan(1/2*f*x + 1/2*e)^3 + 9*a^4*c^4*tan(1/2*f*x + 1/2*e))/a^6)/f`

3.43.9 Mupad [B] (verification not implemented)

Time = 12.99 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx = \frac{13c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 - 11c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + a^2 \right)} - \frac{24c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a^2 f} - \frac{8c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3a^2 f} + \frac{35c^4 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a^2 f}$$

input `int((c - c/cos(e + f*x))^4/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output `(13*c^4*tan(e/2 + (f*x)/2)^3 - 11*c^4*tan(e/2 + (f*x)/2))/(f*(a^2*tan(e/2 + (f*x)/2)^4 - 2*a^2*tan(e/2 + (f*x)/2)^2 + a^2)) - (24*c^4*tan(e/2 + (f*x)/2))/(a^2*f) - (8*c^4*tan(e/2 + (f*x)/2)^3)/(3*a^2*f) + (35*c^4*atanh(tan(e/2 + (f*x)/2)))/(a^2*f)`

3.44
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$$

3.44.1	Optimal result	388
3.44.2	Mathematica [C] (verified)	388
3.44.3	Rubi [A] (verified)	389
3.44.4	Maple [A] (verified)	392
3.44.5	Fricas [A] (verification not implemented)	392
3.44.6	Sympy [F]	393
3.44.7	Maxima [B] (verification not implemented)	393
3.44.8	Giac [A] (verification not implemented)	394
3.44.9	Mupad [B] (verification not implemented)	394

3.44.1 Optimal result

Integrand size = 32, antiderivative size = 119

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx = \frac{5c^3 \operatorname{arctanh}(\sin(e+fx))}{a^2 f} - \frac{5c^3 \tan(e+fx)}{a^2 f} + \frac{2c(c-c\sec(e+fx))^2 \tan(e+fx)}{3f(a+a\sec(e+fx))^2} - \frac{10(c^3-c^3\sec(e+fx)) \tan(e+fx)}{3f(a^2+a^2\sec(e+fx))}$$

output

```
5*c^3*arctanh(sin(f*x+e))/a^2/f-5*c^3*tan(f*x+e)/a^2/f+2/3*c*(c-c*sec(f*x+e))^2*tan(f*x+e)/f/(a+a*sec(f*x+e))^2-10/3*(c^3-c^3*sec(f*x+e))*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))
```

3.44.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.45 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.63

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx = \frac{8c^3 \operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, -\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)} \tan(e+fx)}{3a^2 f (-1+\sec(e+fx))(1+\sec(e+fx))^2}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^2,x]`

output `(-8*c^3*Hypergeometric2F1[-5/2, -3/2, -1/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]]*Tan[e + f*x])/(3*a^2*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^2)`

3.44.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.99, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4445, 3042, 4445, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a\sec(e+fx)+a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^3}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx \\
 & \quad \downarrow \text{4445} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{\sec(e+fx)a+a} dx}{3a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} \\
 & \quad \downarrow \text{4445} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \left(\frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c \int \frac{\sec(e+fx)(c-c\sec(e+fx)) dx}{a} \right)}{3a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \left(\frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2})) dx}{a} \right)}{3a}
 \end{aligned}$$

3.44. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$

$$\begin{aligned}
 & \downarrow 4274 \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \\
 & \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c(c \int \sec(e+fx)dx - c \int \sec^2(e+fx)dx)}{a} \right)}{3a} \\
 & \downarrow 3042 \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \\
 & \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c(c \int \csc(e+fx+\frac{\pi}{2})dx - c \int \csc(e+fx+\frac{\pi}{2})^2 dx)}{a} \right)}{3a} \\
 & \downarrow 4254 \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \\
 & \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c \left(\frac{c \int 1d(-\tan(e+fx))}{f} + c \int \csc(e+fx+\frac{\pi}{2})dx \right)}{a} \right)}{3a} \\
 & \downarrow 24 \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \\
 & \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c(c \int \csc(e+fx+\frac{\pi}{2})dx - \frac{c \tan(e+fx)}{f})}{a} \right)}{3a} \\
 & \downarrow 4257 \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \\
 & \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c \left(\frac{\operatorname{arctanh}(\sin(e+fx))}{f} - \frac{c \tan(e+fx)}{f} \right)}{a} \right)}{3a}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^2,x]`

output `(2*c*(c - c*Sec[e + f*x])^2*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (5*c*((2*(c^2 - c^2*Sec[e + f*x])*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (3*c*((c*ArcTanh[Sin[e + f*x]])/f - (c*Tan[e + f*x])/f))/a))/(3*a)`

3.44. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$

3.44.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4254 `Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4274 `Int[(csc[(e_) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4445 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.44.4 Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.80

method	result
derivativedivides	$\frac{4c^3 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 4} + \frac{5 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} + \frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 4} - \frac{5 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} \right)}{fa^2}$
default	$\frac{4c^3 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 4} + \frac{5 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} + \frac{1}{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 4} - \frac{5 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} \right)}{fa^2}$
parallelrisch	$\frac{5 \left(\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) \cos(fx+e) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \cos(fx+e) + \frac{17 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \left(\cos(fx+e) + \frac{23 \cos(2fx+2e)}{68} + \frac{29}{68} \right)}{15} \right)}{fa^2 \cos(fx+e)}$
risch	$-\frac{2ic^3 (12e^{4i(fx+e)} + 51e^{3i(fx+e)} + 41e^{2i(fx+e)} + 57e^{i(fx+e)} + 23)}{3fa^2 (e^{i(fx+e)} + 1)^3 (1 + e^{2i(fx+e)})} + \frac{5c^3 \ln(e^{i(fx+e)} + i)}{a^2 f} - \frac{5c^3 \ln(e^{i(fx+e)} - i)}{a^2 f}$
norman	$\frac{\frac{10c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} - \frac{80c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} + \frac{22c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{af} - \frac{4c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{af} - \frac{4c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{3af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^3 a} - \frac{5c^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{a^2 f}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `4/f/a^2*c^3*(-1/3*tan(1/2*f*x+1/2*e)^3-2*tan(1/2*f*x+1/2*e)+1/4/(tan(1/2*f*x+1/2*e)+1)+5/4*ln(tan(1/2*f*x+1/2*e)+1)+1/4/(tan(1/2*f*x+1/2*e)-1)-5/4*ln(tan(1/2*f*x+1/2*e)-1))`

3.44.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.50

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{15(c^3 \cos(fx+e)^3 + 2c^3 \cos(fx+e)^2 + c^3 \cos(fx+e)) \log(\sin(fx+e)+1) - 15(c^3 \cos(fx+e)^3 + \dots)}{6(a^2 f \cos(fx+e))^3 + \dots}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^2,x, algorithm="fracas")`

output $1/6*(15*(c^3*\cos(f*x + e)^3 + 2*c^3*\cos(f*x + e)^2 + c^3*\cos(f*x + e))*\log(\sin(f*x + e) + 1) - 15*(c^3*\cos(f*x + e)^3 + 2*c^3*\cos(f*x + e)^2 + c^3*\cos(f*x + e))*\log(-\sin(f*x + e) + 1) - 2*(23*c^3*\cos(f*x + e)^2 + 34*c^3*\cos(f*x + e) + 3*c^3)*\sin(f*x + e))/(a^2*f*\cos(f*x + e)^3 + 2*a^2*f*\cos(f*x + e)^2 + a^2*f*\cos(f*x + e))$

3.44.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^3}{(a + a \sec(e + fx))^2} dx = \frac{c^3 \left(\int \left(-\frac{\sec(e + fx)}{\sec^2(e + fx) + 2 \sec(e + fx) + 1} \right) dx + \int \frac{3 \sec^2(e + fx)}{\sec^2(e + fx) + 2 \sec(e + fx) + 1} dx + \int \left(-\frac{3 \sec^3(e + fx)}{\sec^2(e + fx) + 2 \sec(e + fx) + 1} \right) dx + \dots \right)}{a^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**3/(a+a*sec(f*x+e))**2,x)`

output `-c**3*(Integral(-sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(-3*sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.44.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 341 vs. $2(117) = 234$.

Time = 0.21 (sec) , antiderivative size = 341, normalized size of antiderivative = 2.87

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^3}{(a + a \sec(e + fx))^2} dx = \frac{c^3 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} - \frac{12 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^2} + \frac{12 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^2} + \frac{12 \sin(fx+e)}{\left(a^2 - \frac{a^2 \sin^2(fx+e)}{(\cos(fx+e)+1)^2}\right) (\cos(fx+e)+1)} \right)}{a^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output
$$\begin{aligned} & -1/6*(c^3*((15*\sin(f*x + e)/(\cos(f*x + e) + 1) + \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2 - 12*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/a^2 + 12*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/a^2 + 12*\sin(f*x + e)/((a^2 - a^2*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2)*(\cos(f*x + e) + 1))) + 3*c^3*((9*\sin(f*x + e)/(\cos(f*x + e) + 1) + \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2 - 6*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/a^2 + 6*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/a^2) + 3*c^3*(3*\sin(f*x + e)/(\cos(f*x + e) + 1) + \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2 - c^3*(3*\sin(f*x + e)/(\cos(f*x + e) + 1) - \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2)/f \end{aligned}$$

3.44.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.02

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^3}{(a + a \sec(e + fx))^2} dx$$

$$= \frac{15c^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a^2} - \frac{15c^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a^2} + \frac{6c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)a^2} - \frac{4(a^4c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + 6a^4c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{a^6}$$

$$= \frac{\hspace{15em}}{3f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output
$$\begin{aligned} & 1/3*(15*c^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a^2 - 15*c^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a^2 + 6*c^3*\tan(1/2*f*x + 1/2*e)/((\tan(1/2*f*x + 1/2*e)^2 - 1)*a^2) - 4*(a^4*c^3*\tan(1/2*f*x + 1/2*e)^3 + 6*a^4*c^3*\tan(1/2*f*x + 1/2*e))/a^6)/f \end{aligned}$$

3.44.9 Mupad [B] (verification not implemented)

Time = 13.04 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.87

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^3}{(a + a \sec(e + fx))^2} dx = \frac{10c^3 \operatorname{atanh}(\tan(\frac{e}{2} + \frac{fx}{2}))}{a^2 f} - \frac{4c^3 \tan(\frac{e}{2} + \frac{fx}{2})^3}{3a^2 f}$$

$$- \frac{8c^3 \tan(\frac{e}{2} + \frac{fx}{2})}{a^2 f} + \frac{2c^3 \tan(\frac{e}{2} + \frac{fx}{2})}{f(a^2 \tan(\frac{e}{2} + \frac{fx}{2})^2 - a^2)}$$

input `int((c - c/cos(e + f*x))^3/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output `(10*c^3*atanh(tan(e/2 + (f*x)/2)))/(a^2*f) - (4*c^3*tan(e/2 + (f*x)/2)^3)/(3*a^2*f) - (8*c^3*tan(e/2 + (f*x)/2))/(a^2*f) + (2*c^3*tan(e/2 + (f*x)/2))/(f*(a^2*tan(e/2 + (f*x)/2)^2 - a^2)`

3.45
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

3.45.1	Optimal result	396
3.45.2	Mathematica [A] (verified)	396
3.45.3	Rubi [A] (verified)	397
3.45.4	Maple [A] (verified)	398
3.45.5	Fricas [A] (verification not implemented)	399
3.45.6	Sympy [F]	399
3.45.7	Maxima [B] (verification not implemented)	400
3.45.8	Giac [A] (verification not implemented)	400
3.45.9	Mupad [B] (verification not implemented)	401

3.45.1 Optimal result

Integrand size = 32, antiderivative size = 88

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx = \frac{c^2 \operatorname{arctanh}(\sin(e+fx))}{a^2 f} - \frac{2c^2 \tan(e+fx)}{f(a^2+a^2\sec(e+fx))} + \frac{2(c^2-c^2\sec(e+fx))\tan(e+fx)}{3f(a+a\sec(e+fx))^2}$$

output `c^2*arctanh(sin(f*x+e))/a^2/f-2*c^2*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))+2/3*(c^2-c^2*sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^2`

3.45.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.24

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx = \frac{c^2 \left(-\frac{\log(\cos(\frac{1}{2}(e+fx))-\sin(\frac{1}{2}(e+fx)))}{f} + \frac{\log(\cos(\frac{1}{2}(e+fx))+\sin(\frac{1}{2}(e+fx)))}{f} - \frac{4\tan(\frac{1}{2}(e+fx))}{3f} - \frac{2\sec^2(\frac{1}{2}(e+fx))\tan(\frac{1}{2}(e+fx))}{3f} \right)}{a^2}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^2,x]`

output `(c^2*(-(Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]]/f) + Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]]/f - (4*Tan[(e + f*x)/2])/(3*f) - (2*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/(3*f))/a^2`

3.45.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

3.45.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4445, 3042, 4445, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a\sec(e+fx)+a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx \\
 & \quad \downarrow \text{4445} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{3f(a\sec(e+fx)+a)^2} - \frac{c \int \frac{\sec(e+fx)(c-c\sec(e+fx))}{\sec(e+fx)a+a} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{3f(a\sec(e+fx)+a)^2} - \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{a} \\
 & \quad \downarrow \text{4445} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{3f(a\sec(e+fx)+a)^2} - \frac{c \left(\frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)} - \frac{c \int \sec(e+fx) dx}{a} \right)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{3f(a\sec(e+fx)+a)^2} - \frac{c \left(\frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)} - \frac{c \int \csc(e+fx+\frac{\pi}{2}) dx}{a} \right)}{a} \\
 & \quad \downarrow \text{4257} \\
 & \frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{3f(a\sec(e+fx)+a)^2} - \frac{c \left(\frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)} - \frac{c \operatorname{arctanh}(\sin(e+fx))}{af} \right)}{a}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^2,x]`

3.45. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$

output $(2*(c^2 - c^2*\text{Sec}[e + f*x])*Tan[e + f*x])/(3*f*(a + a*\text{Sec}[e + f*x])^2) - (c*(-((c*\text{ArcTanh}[\text{Sin}[e + f*x]])/(a*f)) + (2*c*Tan[e + f*x])/(f*(a + a*\text{Sec}[e + f*x]))))/a$

3.45.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2*(-1)] && IntegerQ[2*m]`

3.45.4 Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.74

method	result
derivativedivides	$\frac{2c^2 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} \right)}{fa^2}$
default	$\frac{2c^2 \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} \right)}{fa^2}$
parallelrisc	$\frac{c^2 \left(-2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) - 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \right)}{3a^2 f}$
risc	$-\frac{8ic^2(3e^{i(fx+e)}+1)}{3fa^2(e^{i(fx+e)}+1)^3} + \frac{c^2 \ln(e^{i(fx+e)}+i)}{a^2 f} - \frac{c^2 \ln(e^{i(fx+e)}-i)}{a^2 f}$
norman	$\frac{-\frac{2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{10c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} - \frac{2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{3af} - \frac{2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{3af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 a} + \frac{c^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{a^2 f} - \frac{c^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{a^2 f}$

3.45. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 2/f/a^2*c^2*(-1/3*tan(1/2*f*x+1/2*e)^3-tan(1/2*f*x+1/2*e)-1/2*ln(tan(1/2*f*x+1/2*e)-1)+1/2*ln(tan(1/2*f*x+1/2*e)+1))
```

3.45.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.57

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{3(c^2 \cos(fx+e)^2 + 2c^2 \cos(fx+e) + c^2) \log(\sin(fx+e)+1) - 3(c^2 \cos(fx+e)^2 + 2c^2 \cos(fx+e) + c^2) \log(\sin(fx+e)-1) - 8c^2 \sin(fx+e)}{6(a^2 f \cos(fx+e)^2 + 2a^2 f \cos(fx+e) + a^2 f)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x, algorithm="fracas")
```

```
output 1/6*(3*(c^2*cos(f*x + e)^2 + 2*c^2*cos(f*x + e) + c^2)*log(sin(f*x + e) + 1) - 3*(c^2*cos(f*x + e)^2 + 2*c^2*cos(f*x + e) + c^2)*log(-sin(f*x + e) + 1) - 8*(c^2*cos(f*x + e) + 2*c^2)*sin(f*x + e))/(a^2*f*cos(f*x + e)^2 + 2*a^2*f*cos(f*x + e) + a^2*f)
```

3.45.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{c^2 \left(\int \frac{\sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \left(-\frac{2\sec^2(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} \right) dx + \int \frac{\sec^3(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx \right)}{a^2}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x)
```

```
output c**2*(Integral(sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(-2*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2
```

3.45. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$

3.45.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(87) = 174.

Time = 0.22 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.23

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx =$$

$$\frac{c^2 \left(\frac{9 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} - \frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^2} + \frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^2} \right) + \frac{2c^2 \left(\frac{3 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^2} - c^2}{6f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output `-1/6*(c^2*((9*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 6*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 6*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2) + 2*c^2*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - c^2*(3*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f`

3.45.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{3c^2 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3c^2 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right) - \frac{2\left(a^4c^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + 3a^4c^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{a^6}}{3f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output `1/3*(3*c^2*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^2 - 3*c^2*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a^2 - 2*(a^4*c^2*tan(1/2*f*x + 1/2*e)^3 + 3*a^4*c^2*tan(1/2*f*x + 1/2*e))/a^6)/f`

3.45.9 Mupad [B] (verification not implemented)

Time = 13.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.52

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

$$= -\frac{2c^2 \left(3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) - 3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) + \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \right)}{3a^2 f}$$

input `int((c - c/cos(e + f*x))^2/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`output `-(2*c^2*(3*tan(e/2 + (f*x)/2) - 3*atanh(tan(e/2 + (f*x)/2)) + tan(e/2 + (f*x)/2)^3))/(3*a^2*f)`

$$3.46 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$$

3.46.1	Optimal result	402
3.46.2	Mathematica [A] (verified)	402
3.46.3	Rubi [A] (verified)	403
3.46.4	Maple [A] (verified)	404
3.46.5	Fricas [A] (verification not implemented)	404
3.46.6	Sympy [F]	405
3.46.7	Maxima [B] (verification not implemented)	405
3.46.8	Giac [A] (verification not implemented)	406
3.46.9	Mupad [B] (verification not implemented)	406

3.46.1 Optimal result

Integrand size = 30, antiderivative size = 36

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx = \frac{(c-c\sec(e+fx))\tan(e+fx)}{3f(a+a\sec(e+fx))^2}$$

output `1/3*(c-c*sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^2`

3.46.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx = -\frac{c\tan^3\left(\frac{1}{2}(e+fx)\right)}{3a^2f}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2,x]`

output `-1/3*(c*Tan[(e + f*x)/2]^3)/(a^2*f)`

3.46.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a\sec(e+fx)+a)^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx$$

↓ 4438

$$\frac{\tan(e+fx)(c-c\sec(e+fx))}{3f(a\sec(e+fx)+a)^2}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2,x]`

output `((c - c*Sec[e + f*x])*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2)`

3.46.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.46.4 Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3fa^2}$	21
default	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3fa^2}$	21
parallelrisch	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3fa^2}$	21
risch	$\frac{2ic(3e^{2i(fx+e)}+1)}{3fa^2(e^{i(fx+e)}+1)^3}$	37
norman	$\frac{\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} - \frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{3af}}{a\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)}$	61

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
)
```

```
output -1/3/f/a^2*c*tan(1/2*f*x+1/2*e)^3
```

3.46.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.47

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx = \frac{(c \cos(fx+e) - c) \sin(fx+e)}{3(a^2 f \cos(fx+e)^2 + 2a^2 f \cos(fx+e) + a^2 f)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="fricas")
cas")
```

```
output 1/3*(c*cos(f*x + e) - c)*sin(f*x + e)/(a^2*f*cos(f*x + e)^2 + 2*a^2*f*cos(f*x + e) + a^2*f)
```

3.46.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$$

$$= -\frac{c\left(\int\left(-\frac{\sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1}\right) dx + \int\frac{\sec^2(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx\right)}{a^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**2,x)`

output `-c*(Integral(-sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.46.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(35) = 70$.

Time = 0.20 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.61

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$$

$$= -\frac{c\left(\frac{3\sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3}\right)}{a^2} - \frac{c\left(\frac{3\sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3}\right)}{a^2}$$

$6f$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output `-1/6*(c*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - c*(3*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f`

3.46.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx = -\frac{c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3}{3a^2f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output `-1/3*c*tan(1/2*f*x + 1/2*e)^3/(a^2*f)`

3.46.9 Mupad [B] (verification not implemented)

Time = 12.84 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx = -\frac{c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3a^2f}$$

input `int((c - c/cos(e + f*x))/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output `-(c*tan(e/2 + (f*x)/2)^3)/(3*a^2*f)`

3.47 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))} dx$

3.47.1 Optimal result 407
 3.47.2 Mathematica [A] (verified) 407
 3.47.3 Rubi [A] (verified) 408
 3.47.4 Maple [A] (verified) 409
 3.47.5 Fricas [A] (verification not implemented) 409
 3.47.6 Sympy [F] 410
 3.47.7 Maxima [A] (verification not implemented) 410
 3.47.8 Giac [A] (verification not implemented) 410
 3.47.9 Mupad [B] (verification not implemented) 411

3.47.1 Optimal result

Integrand size = 32, antiderivative size = 59

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))} dx = \frac{\cot^3(e+fx)}{3a^2cf} + \frac{\csc(e+fx)}{a^2cf} - \frac{\csc^3(e+fx)}{3a^2cf}$$

output `1/3*cot(f*x+e)^3/a^2/c/f+csc(f*x+e)/a^2/c/f-1/3*csc(f*x+e)^3/a^2/c/f`

3.47.2 Mathematica [A] (verified)

Time = 0.71 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))} dx = \frac{(-1+2 \sec(e+fx)+2 \sec^2(e+fx)) \tan(e+fx)}{3a^2cf(-1+\sec(e+fx))(1+\sec(e+fx))^2}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])),x]`

output `((-1 + 2*Sec[e + f*x] + 2*Sec[e + f*x]^2)*Tan[e + f*x])/(3*a^2*c*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^2)`

3.47.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.86, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^2 (c - c \sec(e+fx))} dx$$

↓ 3042

$$\int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^2 (c - c \csc(e+fx + \frac{\pi}{2}))} dx$$

↓ 4446

$$\int \frac{(c \cot^3(e+fx) \csc(e+fx) - c \cot^2(e+fx) \csc^2(e+fx)) dx}{a^2 c^2}$$

↓ 2009

$$\frac{\frac{c \cot^3(e+fx)}{3f} - \frac{c \csc^3(e+fx)}{3f} + \frac{c \csc(e+fx)}{f}}{a^2 c^2}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])),x]`

output `((c*Cot[e + f*x]^3)/(3*f) + (c*Csc[e + f*x])/f - (c*Csc[e + f*x]^3)/(3*f)) / (a^2*c^2)`

3.47.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.47.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{4fa^2c}$	48
default	$\frac{-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{4fa^2c}$	48
parallelrisc	$\frac{-\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 3 \cot\left(\frac{fx}{2} + \frac{e}{2}\right) + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{12a^2cf}$	48
norman	$\frac{\frac{1}{4acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{2acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{12acf}}{a \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}$	72
risc	$\frac{2i(3e^{3i(fx+e)} + 3e^{2i(fx+e)} + e^{i(fx+e)} - 1)}{3fa^2c(e^{i(fx+e)} + 1)^3(e^{i(fx+e)} - 1)}$	72

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE
)
```

```
output 1/4/f/a^2/c*(-1/3*tan(1/2*f*x+1/2*e)^3+2*tan(1/2*f*x+1/2*e)+1/tan(1/2*f*x+
1/2*e))
```

3.47.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.83

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))} dx = -\frac{\cos(fx + e)^2 - 2 \cos(fx + e) - 2}{3(a^2cf \cos(fx + e) + a^2cf) \sin(fx + e)}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x, algorithm="fri
cas")
```

3.47.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))} dx$$

output $-1/3*(\cos(f*x + e)^2 - 2*\cos(f*x + e) - 2)/((a^2*c*f*\cos(f*x + e) + a^2*c*f)*\sin(f*x + e))$

3.47.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))} dx = -\frac{\int \frac{\sec(e+fx)}{\sec^3(e+fx)+\sec^2(e+fx)-\sec(e+fx)-1} dx}{a^2 c}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e)),x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**3 + sec(e + f*x)**2 - sec(e + f*x) - 1), x)/(a**2*c)`

3.47.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.29

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))} dx = \frac{\frac{6 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3(\cos(fx+e)+1)}{a^2 c \sin(fx+e)}}{12 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `1/12*((6*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^2*c) + 3*(cos(f*x + e) + 1)/(a^2*c*sin(f*x + e)))/f`

3.47.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.17

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))} dx = \frac{\frac{3}{a^2 c \tan(\frac{1}{2} fx + \frac{1}{2} e)} - \frac{a^4 c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 - 6 a^4 c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)}{a^6 c^3}}{12 f}$$

3.47. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e)),x, algorithm="giac")`

output `1/12*(3/(a^2*c*tan(1/2*f*x + 1/2*e)) - (a^4*c^2*tan(1/2*f*x + 1/2*e)^3 - 6*a^4*c^2*tan(1/2*f*x + 1/2*e))/(a^6*c^3))/f`

3.47.9 Mupad [B] (verification not implemented)

Time = 12.87 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.03

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))} dx = -\frac{4 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 8 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1}{12 a^2 c f \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))),x)`

output `-(4*cos(e/2 + (f*x)/2)^4 - 8*cos(e/2 + (f*x)/2)^2 + 1)/(12*a^2*c*f*cos(e/2 + (f*x)/2)^3*sin(e/2 + (f*x)/2))`

$$3.48 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^2} dx$$

3.48.1	Optimal result	412
3.48.2	Mathematica [A] (verified)	412
3.48.3	Rubi [A] (verified)	413
3.48.4	Maple [A] (verified)	414
3.48.5	Fricas [A] (verification not implemented)	415
3.48.6	Sympy [F]	415
3.48.7	Maxima [A] (verification not implemented)	415
3.48.8	Giac [A] (verification not implemented)	416
3.48.9	Mupad [B] (verification not implemented)	416

3.48.1 Optimal result

Integrand size = 32, antiderivative size = 38

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^2} dx = \frac{\csc(e+fx)}{a^2c^2f} - \frac{\csc^3(e+fx)}{3a^2c^2f}$$

output `csc(f*x+e)/a^2/c^2/f-1/3*csc(f*x+e)^3/a^2/c^2/f`

3.48.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^2} dx = \frac{\frac{\csc(e+fx)}{f} - \frac{\csc^3(e+fx)}{3f}}{a^2c^2}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^2),x]`

output `(Csc[e + f*x]/f - Csc[e + f*x]^3/(3*f))/(a^2*c^2)`

3.48.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4446, 3042, 25, 3086, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)^2(c-c \csc(e+fx+\frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{4446} \\
 & \frac{\int \cot^3(e+fx) \csc(e+fx) dx}{a^2 c^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -\sec(e+fx-\frac{\pi}{2}) \tan(e+fx-\frac{\pi}{2})^3 dx}{a^2 c^2} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\int \sec(\frac{1}{2}(2e-\pi)+fx) \tan(\frac{1}{2}(2e-\pi)+fx)^3 dx}{a^2 c^2} \\
 & \quad \downarrow \text{3086} \\
 & -\frac{\int (\csc^2(e+fx)-1) d \csc(e+fx)}{a^2 c^2 f} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\frac{1}{3} \csc^3(e+fx) - \csc(e+fx)}{a^2 c^2 f}
 \end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^2),x]`

output `-((-Csc[e + f*x] + Csc[e + f*x]^3/3)/(a^2*c^2*f))`

3.48. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^2} dx$

3.48.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`
- rule 4446 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^(m) Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.48.4 Maple [A] (verified)

Time = 1.03 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{-\frac{\csc(fx+e)^3}{3} + \csc(fx+e)}{a^2 c^2 f}$	28
parallelrisch	$\frac{-\cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 9\cot\left(\frac{fx}{2} + \frac{e}{2}\right) + 9\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{24f a^2 c^2}$	61
risch	$\frac{2i(3e^{5i(fx+e)} - 2e^{3i(fx+e)} + 3e^{i(fx+e)})}{3f a^2 c^2 (e^{i(fx+e)} + 1)^3 (e^{i(fx+e)} - 1)^3}$	73
norman	$\frac{-\frac{1}{24acf} + \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{8acf} + \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{8acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{24acf}}{a c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}$	97

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

$$3.48. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^2} dx$$

output $1/a^2/c^2/f*(-1/3*\csc(f*x+e)^3+\csc(f*x+e))$

3.48.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.32

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^2} dx = \frac{3\cos(fx+e)^2-2}{3(a^2c^2f\cos(fx+e)^2-a^2c^2f)\sin(fx+e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x, algorithm="fricas")`

output $1/3*(3*\cos(f*x + e)^2 - 2)/((a^2*c^2*f*\cos(f*x + e)^2 - a^2*c^2*f)*\sin(f*x + e))$

3.48.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^2} dx = \frac{\int \frac{\sec(e+fx)}{\sec^4(e+fx)-2\sec^2(e+fx)+1} dx}{a^2c^2}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**2,x)`

output `Integral(sec(e + f*x)/(sec(e + f*x)**4 - 2*sec(e + f*x)**2 + 1), x)/(a**2*c**2)`

3.48.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^2} dx = \frac{3\sin(fx+e)^2-1}{3a^2c^2f\sin(fx+e)^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output $1/3*(3*\sin(f*x + e)^2 - 1)/(a^2*c^2*f*\sin(f*x + e)^3)$

3.48. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^2} dx$

3.48.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^2} dx = \frac{3 \sin(fx + e)^2 - 1}{3 a^2 c^2 f \sin(fx + e)^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `1/3*(3*sin(f*x + e)^2 - 1)/(a^2*c^2*f*sin(f*x + e)^3)`

3.48.9 Mupad [B] (verification not implemented)

Time = 12.99 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^2} dx = \frac{\sin(e + fx)^2 - \frac{1}{3}}{a^2 c^2 f \sin(e + fx)^3}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^2),x)`

output `(sin(e + f*x)^2 - 1/3)/(a^2*c^2*f*sin(e + f*x)^3)`

3.49
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^3} dx$$

3.49.1 Optimal result 417
 3.49.2 Mathematica [A] (verified) 417
 3.49.3 Rubi [A] (verified) 418
 3.49.4 Maple [A] (verified) 419
 3.49.5 Fricas [A] (verification not implemented) 419
 3.49.6 Sympy [F] 420
 3.49.7 Maxima [A] (verification not implemented) 420
 3.49.8 Giac [A] (verification not implemented) 421
 3.49.9 Mupad [B] (verification not implemented) 421

3.49.1 Optimal result

Integrand size = 32, antiderivative size = 80

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^3} dx$$

$$= \frac{\cot^5(e+fx)}{5a^2c^3f} + \frac{\csc(e+fx)}{a^2c^3f} - \frac{2 \csc^3(e+fx)}{3a^2c^3f} + \frac{\csc^5(e+fx)}{5a^2c^3f}$$

output `1/5*cot(f*x+e)^5/a^2/c^3/f+csc(f*x+e)/a^2/c^3/f-2/3*csc(f*x+e)^3/a^2/c^3/f
+1/5*csc(f*x+e)^5/a^2/c^3/f`

3.49.2 Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.99

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^3} dx$$

$$= \frac{(3 + 12 \sec(e+fx) - 12 \sec^2(e+fx) - 8 \sec^3(e+fx) + 8 \sec^4(e+fx)) \tan(e+fx)}{15a^2c^3f(-1 + \sec(e+fx))^3(1 + \sec(e+fx))^2}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^3),x]`

output `((3 + 12*Sec[e + f*x] - 12*Sec[e + f*x]^2 - 8*Sec[e + f*x]^3 + 8*Sec[e + f*x]^4)*Tan[e + f*x])/(15*a^2*c^3*f*(-1 + Sec[e + f*x])^3*(1 + Sec[e + f*x])^2)`

3.49.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^3} dx$$

3.49.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.86, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)^2(c-c \csc(e+fx+\frac{\pi}{2}))^3} dx$$

↓ 4446

$$-\frac{\int (a \csc(e+fx) \cot^5(e+fx) + a \csc^2(e+fx) \cot^4(e+fx)) dx}{a^3 c^3}$$

↓ 2009

$$-\frac{-\frac{a \cot^5(e+fx)}{5f} - \frac{a \csc^5(e+fx)}{5f} + \frac{2a \csc^3(e+fx)}{3f} - \frac{a \csc(e+fx)}{f}}{a^3 c^3}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^3),x]`

output `-((-1/5*(a*Cot[e + f*x]^5)/f - (a*Csc[e + f*x])/f + (2*a*Csc[e + f*x]^3)/(3*f) - (a*Csc[e + f*x]^5)/(5*f))/(a^3*c^3)`

3.49.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.49.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.92

method	result	size
parallelrisch	$\frac{3 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - 20 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 60 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 90 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)}{240 f a^2 c^3}$	74
derivativedivides	$-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{4}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} + \frac{6}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}$	76
default	$-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{4}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} + \frac{6}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}$	76
risch	$\frac{2i(15e^{7i(fx+e)} - 15e^{6i(fx+e)} - 5e^{5i(fx+e)} + 25e^{4i(fx+e)} + 13e^{3i(fx+e)} - 21e^{2i(fx+e)} + 9e^{i(fx+e)} + 3)}{15f a^2 c^3 (e^{i(fx+e)} + 1)^3 (e^{i(fx+e)} - 1)^5}$	118
norman	$\frac{\frac{1}{80acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{12acf} + \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{8acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{4acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{48acf}}{a c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	119

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x,method=_RETURNVERBO
SE)
```

```
output 1/240*(3*cot(1/2*f*x+1/2*e)^5-5*tan(1/2*f*x+1/2*e)^3-20*cot(1/2*f*x+1/2*e)
^3+60*tan(1/2*f*x+1/2*e)+90*cot(1/2*f*x+1/2*e))/f/a^2/c^3
```

3.49.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.36

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^3} dx$$

$$= \frac{3 \cos(fx + e)^4 + 12 \cos(fx + e)^3 - 12 \cos(fx + e)^2 - 8 \cos(fx + e) + 8}{15 (a^2 c^3 f \cos(fx + e)^3 - a^2 c^3 f \cos(fx + e)^2 - a^2 c^3 f \cos(fx + e) + a^2 c^3 f) \sin(fx + e)}$$

3.49. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))^3} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x, algorithm="fricas")`

output `1/15*(3*cos(f*x + e)^4 + 12*cos(f*x + e)^3 - 12*cos(f*x + e)^2 - 8*cos(f*x + e) + 8)/((a^2*c^3*f*cos(f*x + e)^3 - a^2*c^3*f*cos(f*x + e)^2 - a^2*c^3*f*cos(f*x + e) + a^2*c^3*f)*sin(f*x + e))`

3.49.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^3} dx$$

$$= - \frac{\int \frac{\sec(e+fx)}{\sec^5(e+fx) - \sec^4(e+fx) - 2\sec^3(e+fx) + 2\sec^2(e+fx) + \sec(e+fx) - 1} dx}{a^2 c^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**3,x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**5 - sec(e + f*x)**4 - 2*sec(e + f*x)**3 + 2*sec(e + f*x)**2 + sec(e + f*x) - 1), x)/(a**2*c**3)`

3.49.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^3} dx$$

$$= \frac{5 \left(\frac{12 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right) - \left(\frac{20 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{90 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - 3 \right) (\cos(fx+e)+1)^5}{a^2 c^3 \sin(fx+e)^5} \cdot \frac{1}{240 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/240*(5*(12*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^2*c^3) - (20*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 90*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 3)*(cos(f*x + e) + 1)^5/(a^2*c^3*sin(f*x + e)^5))/f`

3.49. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))^3} dx$

3.49.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.20

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^3} dx$$

$$= \frac{90 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 - 20 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 + 3}{a^2 c^3 \tan(\frac{1}{2} fx + \frac{1}{2} e)^5} - \frac{5 (a^4 c^6 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 - 12 a^4 c^6 \tan(\frac{1}{2} fx + \frac{1}{2} e))}{a^6 c^9}$$

$$240 f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `1/240*((90*tan(1/2*f*x + 1/2*e)^4 - 20*tan(1/2*f*x + 1/2*e)^2 + 3)/(a^2*c^3*tan(1/2*f*x + 1/2*e)^5) - 5*(a^4*c^6*tan(1/2*f*x + 1/2*e)^3 - 12*a^4*c^6*tan(1/2*f*x + 1/2*e))/(a^6*c^9))/f`

3.49.9 Mupad [B] (verification not implemented)

Time = 13.47 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^3} dx$$

$$= \frac{-5 \tan(\frac{e}{2} + \frac{fx}{2})^8 + 60 \tan(\frac{e}{2} + \frac{fx}{2})^6 + 90 \tan(\frac{e}{2} + \frac{fx}{2})^4 - 20 \tan(\frac{e}{2} + \frac{fx}{2})^2 + 3}{240 a^2 c^3 f \tan(\frac{e}{2} + \frac{fx}{2})^5}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^3),x)`

output `(90*tan(e/2 + (f*x)/2)^4 - 20*tan(e/2 + (f*x)/2)^2 + 60*tan(e/2 + (f*x)/2)^6 - 5*tan(e/2 + (f*x)/2)^8 + 3)/(240*a^2*c^3*f*tan(e/2 + (f*x)/2)^5)`

3.50 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^4} dx$

3.50.1 Optimal result 422
 3.50.2 Mathematica [A] (verified) 422
 3.50.3 Rubi [A] (verified) 423
 3.50.4 Maple [A] (verified) 424
 3.50.5 Fricas [A] (verification not implemented) 425
 3.50.6 Sympy [F] 425
 3.50.7 Maxima [A] (verification not implemented) 426
 3.50.8 Giac [A] (verification not implemented) 426
 3.50.9 Mupad [B] (verification not implemented) 427

3.50.1 Optimal result

Integrand size = 32, antiderivative size = 98

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^4} dx$$

$$= -\frac{2 \cot^7(e+fx)}{7a^2c^4f} + \frac{\csc(e+fx)}{a^2c^4f} - \frac{4 \csc^3(e+fx)}{3a^2c^4f} + \frac{\csc^5(e+fx)}{a^2c^4f} - \frac{2 \csc^7(e+fx)}{7a^2c^4f}$$

output `-2/7*cot(f*x+e)^7/a^2/c^4/f+csc(f*x+e)/a^2/c^4/f-4/3*csc(f*x+e)^3/a^2/c^4/f+csc(f*x+e)^5/a^2/c^4/f-2/7*csc(f*x+e)^7/a^2/c^4/f`

3.50.2 Mathematica [A] (verified)

Time = 1.69 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^4} dx$$

$$= \frac{(-6 - 9 \sec(e+fx) + 24 \sec^2(e+fx) - 4 \sec^3(e+fx) - 16 \sec^4(e+fx) + 8 \sec^5(e+fx)) \tan(e+fx)}{21a^2c^4f(-1 + \sec(e+fx))^4(1 + \sec(e+fx))^2}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^4),x]`

output `((-6 - 9*Sec[e + f*x] + 24*Sec[e + f*x]^2 - 4*Sec[e + f*x]^3 - 16*Sec[e + f*x]^4 + 8*Sec[e + f*x]^5)*Tan[e + f*x])/(21*a^2*c^4*f*(-1 + Sec[e + f*x])^4*(1 + Sec[e + f*x])^2)`

3.50. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^4} dx$

3.50.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^4} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)^2(c-c \csc(e+fx+\frac{\pi}{2}))^4} dx$$

↓ 4446

$$\int \frac{(a^2 \csc(e+fx) \cot^7(e+fx) + 2a^2 \csc^2(e+fx) \cot^6(e+fx) + a^2 \csc^3(e+fx) \cot^5(e+fx)) dx}{a^4 c^4}$$

↓ 2009

$$\frac{-\frac{2a^2 \cot^7(e+fx)}{7f} - \frac{2a^2 \csc^7(e+fx)}{7f} + \frac{a^2 \csc^5(e+fx)}{f} - \frac{4a^2 \csc^3(e+fx)}{3f} + \frac{a^2 \csc(e+fx)}{f}}{a^4 c^4}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^4),x]`

output `((-2*a^2*Cot[e + f*x]^7)/(7*f) + (a^2*Csc[e + f*x])/f - (4*a^2*Csc[e + f*x]^3)/(3*f) + (a^2*Csc[e + f*x]^5)/f - (2*a^2*Csc[e + f*x]^7)/(7*f))/(a^4*c^4)`

3.50.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.50. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^4} dx$


```
rule 4446 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.50.4 Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{10}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{10}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}}{32f c^4 a^2}$
default	$\frac{-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{10}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{10}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} - \frac{1}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}}{32f c^4 a^2}$
parallelrisch	$\frac{-3 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^7 + 21 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - 70 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 105 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 210 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)}{672f a^2 c^4}$
risch	$\frac{2i(21 e^{9i(fx+e)} - 42 e^{8i(fx+e)} + 28 e^{7i(fx+e)} + 56 e^{6i(fx+e)} - 42 e^{5i(fx+e)} - 28 e^{4i(fx+e)} + 76 e^{3i(fx+e)} - 24 e^{2i(fx+e)} - 3 e^{i(fx+e)})}{21f c^4 a^2 (e^{i(fx+e)} - 1)^7 (e^{i(fx+e)} + 1)^3}$
norman	$\frac{-\frac{1}{224acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{32acf} - \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{48acf} + \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{16acf} + \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{32acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{10}}{96acf}}{a^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x,method=_RETURNVERBO
SE)
```

```
output 1/32/f/c^4/a^2*(-1/3*tan(1/2*f*x+1/2*e)^3+5*tan(1/2*f*x+1/2*e)+10/tan(1/2*
f*x+1/2*e)-10/3/tan(1/2*f*x+1/2*e)^3-1/7/tan(1/2*f*x+1/2*e)^7+1/tan(1/2*f*
x+1/2*e)^5)
```

3.50.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^4} dx$$

3.50.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.22

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^4} dx$$

$$= \frac{6 \cos(fx + e)^5 + 9 \cos(fx + e)^4 - 24 \cos(fx + e)^3 + 4 \cos(fx + e)^2 + 16 \cos(fx + e) - 8}{21 (a^2 c^4 f \cos(fx + e)^4 - 2 a^2 c^4 f \cos(fx + e)^3 + 2 a^2 c^4 f \cos(fx + e) - a^2 c^4 f) \sin(fx + e)}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x, algorithm="fracas")
```

```
output 1/21*(6*cos(f*x + e)^5 + 9*cos(f*x + e)^4 - 24*cos(f*x + e)^3 + 4*cos(f*x + e)^2 + 16*cos(f*x + e) - 8)/((a^2*c^4*f*cos(f*x + e)^4 - 2*a^2*c^4*f*cos(f*x + e)^3 + 2*a^2*c^4*f*cos(f*x + e) - a^2*c^4*f)*sin(f*x + e))
```

3.50.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^4} dx$$

$$= \int \frac{\sec(e+fx)}{\sec^6(e+fx) - 2\sec^5(e+fx) - \sec^4(e+fx) + 4\sec^3(e+fx) - \sec^2(e+fx) - 2\sec(e+fx) + 1} dx$$

$$a^2 c^4$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**4,x)
```

```
output Integral(sec(e + f*x)/(sec(e + f*x)**6 - 2*sec(e + f*x)**5 - sec(e + f*x)**4 + 4*sec(e + f*x)**3 - sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x)/(a**2*c**4)
```

3.50.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.43

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^4} dx$$

$$= \frac{7 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^2 c^4} + \frac{\left(\frac{21 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{70 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{210 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - 3 \right) (\cos(fx+e)+1)^7}{a^2 c^4 \sin(fx+e)^7}$$

$$672 f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output `1/672*(7*(15*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^2*c^4) + (21*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 70*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 210*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 3)*(cos(f*x + e) + 1)^7/(a^2*c^4*sin(f*x + e)^7))/f`

3.50.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.11

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^4} dx$$

$$= \frac{210 \tan(\frac{1}{2} fx + \frac{1}{2} e)^6 - 70 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 + 21 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 3}{a^2 c^4 \tan(\frac{1}{2} fx + \frac{1}{2} e)^7} - \frac{7 (a^4 c^8 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 - 15 a^4 c^8 \tan(\frac{1}{2} fx + \frac{1}{2} e))}{a^6 c^{12}}$$

$$672 f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output `1/672*((210*tan(1/2*f*x + 1/2*e)^6 - 70*tan(1/2*f*x + 1/2*e)^4 + 21*tan(1/2*f*x + 1/2*e)^2 - 3)/(a^2*c^4*tan(1/2*f*x + 1/2*e)^7) - 7*(a^4*c^8*tan(1/2*f*x + 1/2*e)^3 - 15*a^4*c^8*tan(1/2*f*x + 1/2*e))/(a^6*c^12))/f`

3.50.9 Mupad [B] (verification not implemented)

Time = 14.31 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^4} dx$$

$$= \frac{-7 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} + 105 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 210 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 70 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 21 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 3}{672 a^2 c^4 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}$$

input `int(1/(cos(e + f*x))*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^4),x)`

output `(21*tan(e/2 + (f*x)/2)^2 - 70*tan(e/2 + (f*x)/2)^4 + 210*tan(e/2 + (f*x)/2)^6 + 105*tan(e/2 + (f*x)/2)^8 - 7*tan(e/2 + (f*x)/2)^10 - 3)/(672*a^2*c^4*f*tan(e/2 + (f*x)/2)^7)`

$$3.51 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^5} dx$$

3.51.1	Optimal result	428
3.51.2	Mathematica [A] (verified)	428
3.51.3	Rubi [A] (verified)	429
3.51.4	Maple [A] (verified)	430
3.51.5	Fricas [A] (verification not implemented)	431
3.51.6	Sympy [F]	431
3.51.7	Maxima [A] (verification not implemented)	432
3.51.8	Giac [A] (verification not implemented)	432
3.51.9	Mupad [B] (verification not implemented)	433

3.51.1 Optimal result

Integrand size = 32, antiderivative size = 141

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^5} dx$$

$$= \frac{\cot^7(e+fx)}{7a^2c^5f} + \frac{4 \cot^9(e+fx)}{9a^2c^5f} + \frac{\csc(e+fx)}{a^2c^5f} - \frac{7 \csc^3(e+fx)}{3a^2c^5f}$$

$$+ \frac{3 \csc^5(e+fx)}{a^2c^5f} - \frac{13 \csc^7(e+fx)}{7a^2c^5f} + \frac{4 \csc^9(e+fx)}{9a^2c^5f}$$

output `1/7*cot(f*x+e)^7/a^2/c^5/f+4/9*cot(f*x+e)^9/a^2/c^5/f+csc(f*x+e)/a^2/c^5/f
-7/3*csc(f*x+e)^3/a^2/c^5/f+3*csc(f*x+e)^5/a^2/c^5/f-13/7*csc(f*x+e)^7/a^2/
/c^5/f+4/9*csc(f*x+e)^9/a^2/c^5/f`

3.51.2 Mathematica [A] (verified)

Time = 3.87 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.70

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^5} dx$$

$$= \frac{(19 + 6 \sec(e+fx) - 66 \sec^2(e+fx) + 56 \sec^3(e+fx) + 24 \sec^4(e+fx) - 48 \sec^5(e+fx) + 16 \sec^6(e+fx))}{63a^2c^5f(-1 + \sec(e+fx))^5(1 + \sec(e+fx))^2}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^5),x]`

3.51. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^5} dx$

output $((19 + 6*\text{Sec}[e + f*x] - 66*\text{Sec}[e + f*x]^2 + 56*\text{Sec}[e + f*x]^3 + 24*\text{Sec}[e + f*x]^4 - 48*\text{Sec}[e + f*x]^5 + 16*\text{Sec}[e + f*x]^6)*\text{Tan}[e + f*x])/(63*a^2*c^5*f*(-1 + \text{Sec}[e + f*x])^5*(1 + \text{Sec}[e + f*x])^2)$

3.51.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.91, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^2 (c - c \sec(e+fx))^5} dx$$

↓ 3042

$$\int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^2 (c - c \csc(e+fx + \frac{\pi}{2}))^5} dx$$

↓ 4446

$$\frac{\int (a^3 \csc(e+fx) \cot^9(e+fx) + 3a^3 \csc^2(e+fx) \cot^8(e+fx) + 3a^3 \csc^3(e+fx) \cot^7(e+fx) + a^3 \csc^4(e+fx)) dx}{a^5 c^5}$$

↓ 2009

$$\frac{-\frac{4a^3 \cot^9(e+fx)}{9f} - \frac{a^3 \cot^7(e+fx)}{7f} - \frac{4a^3 \csc^9(e+fx)}{9f} + \frac{13a^3 \csc^7(e+fx)}{7f} - \frac{3a^3 \csc^5(e+fx)}{f} + \frac{7a^3 \csc^3(e+fx)}{3f} - \frac{a^3 \csc(e+fx)}{f}}{a^5 c^5}$$

input $\text{Int}[\text{Sec}[e + f*x]/((a + a*\text{Sec}[e + f*x])^2*(c - c*\text{Sec}[e + f*x])^5), x]$

output $-((-1/7*(a^3*\text{Cot}[e + f*x]^7)/f - (4*a^3*\text{Cot}[e + f*x]^9)/(9*f) - (a^3*\text{Csc}[e + f*x])/f + (7*a^3*\text{Csc}[e + f*x]^3)/(3*f) - (3*a^3*\text{Csc}[e + f*x]^5)/f + (13*a^3*\text{Csc}[e + f*x]^7)/(7*f) - (4*a^3*\text{Csc}[e + f*x]^9)/(9*f))/(a^5*c^5))$

3.51. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^5} dx$

3.51.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.51.4 Maple [A] (verified)

Time = 0.85 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.71

method	result
parallelrisch	$\frac{7 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^9 - 54 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^7 + 189 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 21 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - 420 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 378 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 945 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)}{4032 f a^2 c^5}$
derivativedivides	$-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{20}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{3}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{6}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{15}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} + \frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$
default	$-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{20}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{3}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{6}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{15}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} + \frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$
risch	$\frac{2i(63 e^{11i(fx+e)} - 189 e^{10i(fx+e)} + 273 e^{9i(fx+e)} + 63 e^{8i(fx+e)} - 378 e^{7i(fx+e)} + 294 e^{6i(fx+e)} + 306 e^{5i(fx+e)} - 450 e^{4i(fx+e)} - 189 e^{3i(fx+e)} + 189 e^{2i(fx+e)} - 189 e^{i(fx+e)} + 189)}{63 f c^5 a^2 (e^{i(fx+e)} - 1)^9 (e^{i(fx+e)} + 1)^3}$
norman	$\frac{\frac{1}{576 a c f} - \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{224 a c f} + \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{64 a c f} - \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{48 a c f} + \frac{15 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{64 a c f} + \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{10}}{32 a c f} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{12}}{192 a c f}}{a c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x,method=_RETURNVERBOSE)
```

```
output 1/4032*(7*cot(1/2*f*x+1/2*e)^9-54*cot(1/2*f*x+1/2*e)^7+189*cot(1/2*f*x+1/2*e)^5-21*tan(1/2*f*x+1/2*e)^3-420*cot(1/2*f*x+1/2*e)^3+378*tan(1/2*f*x+1/2*e)+945*cot(1/2*f*x+1/2*e))/f/a^2/c^5
```

$$3.51. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^5} dx$$

3.51.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.16

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^5} dx$$

$$= \frac{19 \cos(fx + e)^6 + 6 \cos(fx + e)^5 - 66 \cos(fx + e)^4 + 56 \cos(fx + e)^3 + 24 \cos(fx + e)^2 - 48 \cos(fx + e) + 16}{63 (a^2 c^5 f \cos(fx + e)^5 - 3 a^2 c^5 f \cos(fx + e)^4 + 2 a^2 c^5 f \cos(fx + e)^3 + 2 a^2 c^5 f \cos(fx + e)^2 - 3 a^2 c^5 f \cos(fx + e) + a^2 c^5 f) \sin(fx + e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x, algorithm="fricas")`

output `1/63*(19*cos(f*x + e)^6 + 6*cos(f*x + e)^5 - 66*cos(f*x + e)^4 + 56*cos(f*x + e)^3 + 24*cos(f*x + e)^2 - 48*cos(f*x + e) + 16)/((a^2*c^5*f*cos(f*x + e)^5 - 3*a^2*c^5*f*cos(f*x + e)^4 + 2*a^2*c^5*f*cos(f*x + e)^3 + 2*a^2*c^5*f*cos(f*x + e)^2 - 3*a^2*c^5*f*cos(f*x + e) + a^2*c^5*f)*sin(f*x + e))`

3.51.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^5} dx$$

$$= - \frac{\int \frac{\sec(e+fx)}{\sec^7(e+fx) - 3 \sec^6(e+fx) + \sec^5(e+fx) + 5 \sec^4(e+fx) - 5 \sec^3(e+fx) - \sec^2(e+fx) + 3 \sec(e+fx) - 1} dx}{a^2 c^5}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**5,x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**7 - 3*sec(e + f*x)**6 + sec(e + f*x)**5 + 5*sec(e + f*x)**4 - 5*sec(e + f*x)**3 - sec(e + f*x)**2 + 3*sec(e + f*x) - 1), x)/(a**2*c**5)`

3.51.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.14

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^5} dx$$

$$= \frac{21 \left(\frac{18 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right) - \left(\frac{54 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{189 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{420 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - \frac{945 \sin(fx+e)^8}{(\cos(fx+e)+1)^8} - 7 \right) (\cos(fx+e)+1)^9}{a^2 c^5 \sin(fx+e)^9} 4032 f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x, algorithm="maxima")`

output `1/4032*(21*(18*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^2*c^5) - (54*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 189*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 420*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 945*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 - 7)*(cos(f*x + e) + 1)^9/(a^2*c^5*sin(f*x + e)^9))/f`

3.51.8 Giac [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.87

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^5} dx$$

$$= \frac{945 \tan(\frac{1}{2} fx + \frac{1}{2} e)^8 - 420 \tan(\frac{1}{2} fx + \frac{1}{2} e)^6 + 189 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 - 54 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 + 7}{a^2 c^5 \tan(\frac{1}{2} fx + \frac{1}{2} e)^9} - \frac{21 (a^4 c^{10} \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 - 18 a^4 c^{10} \tan(\frac{1}{2} fx + \frac{1}{2} e))}{a^6 c^{15}} 4032 f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^5,x, algorithm="giac")`

output `1/4032*((945*tan(1/2*f*x + 1/2*e)^8 - 420*tan(1/2*f*x + 1/2*e)^6 + 189*tan(1/2*f*x + 1/2*e)^4 - 54*tan(1/2*f*x + 1/2*e)^2 + 7)/(a^2*c^5*tan(1/2*f*x + 1/2*e)^9) - 21*(a^4*c^10*tan(1/2*f*x + 1/2*e)^3 - 18*a^4*c^10*tan(1/2*f*x + 1/2*e)))/(a^6*c^15))/f`

3.51. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))^5} dx$

3.51.9 Mupad [B] (verification not implemented)

Time = 15.56 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.72

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^5} dx$$

$$= \frac{-21 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{12} + 378 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} + 945 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 420 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 189 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 54 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 21}{4032 a^2 c^5 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^5),x)`output `(189*tan(e/2 + (f*x)/2)^4 - 54*tan(e/2 + (f*x)/2)^2 - 420*tan(e/2 + (f*x)/2)^6 + 945*tan(e/2 + (f*x)/2)^8 + 378*tan(e/2 + (f*x)/2)^10 - 21*tan(e/2 + (f*x)/2)^12 + 7)/(4032*a^2*c^5*f*tan(e/2 + (f*x)/2)^9)`

3.52
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$$

3.52.1 Optimal result 434
 3.52.2 Mathematica [C] (verified) 435
 3.52.3 Rubi [A] (verified) 435
 3.52.4 Maple [A] (verified) 438
 3.52.5 Fricas [A] (verification not implemented) 438
 3.52.6 Sympy [F] 439
 3.52.7 Maxima [B] (verification not implemented) 439
 3.52.8 Giac [A] (verification not implemented) 440
 3.52.9 Mupad [B] (verification not implemented) 441

3.52.1 Optimal result

Integrand size = 32, antiderivative size = 215

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx = -\frac{231c^6 \operatorname{arctanh}(\sin(e+fx))}{2a^3 f} + \frac{924c^6 \tan(e+fx)}{5a^3 f} - \frac{693c^6 \sec(e+fx) \tan(e+fx)}{10a^3 f} - \frac{22c^2(c-c\sec(e+fx))^4 \tan(e+fx)}{15af(a+a\sec(e+fx))^2} + \frac{2c(c-c\sec(e+fx))^5 \tan(e+fx)}{5f(a+a\sec(e+fx))^3} + \frac{66(c^2-c^2\sec(e+fx))^3 \tan(e+fx)}{5f(a^3+a^3\sec(e+fx))} + \frac{77c^6 \tan^3(e+fx)}{5a^3 f}$$

```
output -231/2*c^6*arctanh(sin(f*x+e))/a^3/f+924/5*c^6*tan(f*x+e)/a^3/f-693/10*c^6
*sec(f*x+e)*tan(f*x+e)/a^3/f-22/15*c^2*(c-c*sec(f*x+e))^4*tan(f*x+e)/a/f/(
a+a*sec(f*x+e))^2+2/5*c*(c-c*sec(f*x+e))^5*tan(f*x+e)/f/(a+a*sec(f*x+e))^3
+66/5*(c^2-c^2*sec(f*x+e))^3*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))+77/5*c^6*ta
n(f*x+e)^3/a^3/f
```

3.52.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 5.14 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.35

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx = \frac{64c^6 \operatorname{Hypergeometric2F1}\left(-\frac{11}{2}, -\frac{5}{2}, -\frac{3}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)} \tan(e+fx)}{5a^3 f(-1+\sec(e+fx))(1+\sec(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^6)/(a + a*Sec[e + f*x])^3,x]`

output `(-64*c^6*Hypergeometric2F1[-11/2, -5/2, -3/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]]*Tan[e + f*x])/(5*a^3*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^3)`

3.52.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.98, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.281$, Rules used = {3042, 4445, 3042, 4445, 3042, 4445, 3042, 4278, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a\sec(e+fx)+a)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^6}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\ & \quad \downarrow \text{4445} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^5}{5f(a\sec(e+fx)+a)^3} - \frac{11c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(\sec(e+fx)a+a)^2} dx}{5a} \\ & \quad \downarrow \text{3042} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^5}{5f(a\sec(e+fx)+a)^3} - \frac{11c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^5}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a} \end{aligned}$$

3.52. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$

$$\begin{array}{c}
\downarrow 4445 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^5}{5f(a \sec(e+fx)+a)^3} - \frac{11c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \int \frac{\sec(e+fx)(c-c \sec(e+fx))^4 dx}{\sec(e+fx)a+a}}{a} \right)}{5a} \\
\downarrow 3042 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^5}{5f(a \sec(e+fx)+a)^3} - \\
\frac{11c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c \csc(e+fx+\frac{\pi}{2}))^4 dx}{\csc(e+fx+\frac{\pi}{2})a+a}}{a} \right)}{5a} \\
\downarrow 4445 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^5}{5f(a \sec(e+fx)+a)^3} - \\
\frac{11c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{f(a \sec(e+fx)+a)} - \frac{7c \int \sec(e+fx)(c-c \sec(e+fx))^3 dx}{a} \right)}{a} \right)}{5a} \\
\downarrow 3042 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^5}{5f(a \sec(e+fx)+a)^3} - \\
\frac{11c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{f(a \sec(e+fx)+a)} - \frac{7c \int \csc(e+fx+\frac{\pi}{2})(c-c \csc(e+fx+\frac{\pi}{2}))^3 dx}{a} \right)}{a} \right)}{5a} \\
\downarrow 4278 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^5}{5f(a \sec(e+fx)+a)^3} - \\
\frac{11c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{3f(a \sec(e+fx)+a)^2} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{f(a \sec(e+fx)+a)} - \frac{7c \int (-c^3 \sec^4(e+fx)+3c^3 \sec^3(e+fx)-3c^3 \sec^2(e+fx)+c^3 \sec(e+fx)) dx}{a} \right)}{a} \right)}{5a} \\
\downarrow 2009
\end{array}$$

3.52. $\int \frac{\sec(e+fx)(c-c \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$

$$11c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^4}{3f(a\sec(e+fx)+a)^2} - \frac{2c \tan(e+fx)(c-c\sec(e+fx))^5}{5f(a\sec(e+fx)+a)^3} - \frac{3c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{f(a\sec(e+fx)+a)} - \frac{7c \left(\frac{5c^3 \operatorname{arctanh}(\sin(e+fx))}{2f} - \frac{c^3 \tan^3(e+fx)}{3f} - \frac{4c^3 \tan(e+fx)}{f} + \frac{3c^3 \tan(e+fx)}{a} \right)}{a} \right)}{5a} \right)$$

```
input Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^6)/(a + a*Sec[e + f*x])^3,x]
```

```
output (2*c*(c - c*Sec[e + f*x])^5*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) - (11*c*((2*c*(c - c*Sec[e + f*x])^4*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (3*c*((2*c*(c - c*Sec[e + f*x])^3*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))) - (7*c*((5*c^3*ArcTanh[Sin[e + f*x]])/(2*f) - (4*c^3*Tan[e + f*x])/f + (3*c^3*Sec[e + f*x]*Tan[e + f*x])/(2*f) - (c^3*Tan[e + f*x]^3)/(3*f)))/a)/a))/(5*a)
```

3.52.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4278 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n_*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_], x_Symbol] := Int[ExpandTrig[(a + b*csc[e + f*x])^m*(d*csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && I GtQ[m, 0] && RationalQ[n]
```

```
rule 4445 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^n_], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]
```

3.52.4 Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.78

method	result
derivativedivides	$16c^6 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{48(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^3} + \frac{5}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^2} - \frac{89}{32(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)} \right) f a^3$
default	$16c^6 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{48(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^3} + \frac{5}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^2} - \frac{89}{32(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)} \right) f a^3$
parallelrisch	$2723c^6 \left(\frac{3960(\cos(3fx+3e)+3\cos(fx+e))\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{389} + \frac{3960(-\cos(3fx+3e)-3\cos(fx+e))\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{389} + \tan\left(\frac{fx}{2}\right) \right) 240f a^3 (\cos(3fx+e))$
risch	$\frac{ic^6 (3495 e^{10i(fx+e)} + 17205 e^{9i(fx+e)} + 44480 e^{8i(fx+e)} + 79450 e^{7i(fx+e)} + 120176 e^{6i(fx+e)} + 130340 e^{5i(fx+e)} + 127498 e^{4i(fx+e)} + 100000 e^{3i(fx+e)} + 60000 e^{2i(fx+e)} + 20000 e^{i(fx+e)} + 10000)}{15f a^3 (1+e^{2i(fx+e)})^3 (e^{i(fx+e)}+1)^5}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$16/f*c^6/a^3*(1/5*\tan(1/2*f*x+1/2*e)^5+4/3*\tan(1/2*f*x+1/2*e)^3+10*\tan(1/2*f*x+1/2*e)-1/48/(\tan(1/2*f*x+1/2*e)+1)^3+5/16/(\tan(1/2*f*x+1/2*e)+1)^2-89/32/(\tan(1/2*f*x+1/2*e)+1)-231/32*\ln(\tan(1/2*f*x+1/2*e)+1)-1/48/(\tan(1/2*f*x+1/2*e)-1)^3-5/16/(\tan(1/2*f*x+1/2*e)-1)^2-89/32/(\tan(1/2*f*x+1/2*e)-1)+231/32*\ln(\tan(1/2*f*x+1/2*e)-1))$$

3.52.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.22

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx = \frac{3465(c^6 \cos(fx+e)^6 + 3c^6 \cos(fx+e)^5 + 3c^6 \cos(fx+e)^4 + c^6 \cos(fx+e)^3) \log(\sin(fx+e)+1)}{15f a^3 (1+e^{2i(fx+e)})^3 (e^{i(fx+e)}+1)^5}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x, algorithm="fricas")`

```
output -1/60*(3465*(c^6*cos(f*x + e)^6 + 3*c^6*cos(f*x + e)^5 + 3*c^6*cos(f*x + e)^4 + c^6*cos(f*x + e)^3)*log(sin(f*x + e) + 1) - 3465*(c^6*cos(f*x + e)^6 + 3*c^6*cos(f*x + e)^5 + 3*c^6*cos(f*x + e)^4 + c^6*cos(f*x + e)^3)*log(-sin(f*x + e) + 1) - 2*(5446*c^6*cos(f*x + e)^5 + 12843*c^6*cos(f*x + e)^4 + 8711*c^6*cos(f*x + e)^3 + 815*c^6*cos(f*x + e)^2 - 105*c^6*cos(f*x + e) + 10*c^6)*sin(f*x + e))/(a^3*f*cos(f*x + e)^6 + 3*a^3*f*cos(f*x + e)^5 + 3*a^3*f*cos(f*x + e)^4 + a^3*f*cos(f*x + e)^3)
```

3.52.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{c^6 \left(\int \frac{\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \left(-\frac{6\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} \right) dx + \int \frac{15}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx \right)}{a^3}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))**6/(a+a*sec(f*x+e))**3,x)
```

```
output c**6*(Integral(sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(-6*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(15*sec(e + f*x)**3/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(-20*sec(e + f*x)**4/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(15*sec(e + f*x)**5/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(-6*sec(e + f*x)**6/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**7/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3
```

3.52.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 935 vs. $2(204) = 408$.

Time = 0.26 (sec) , antiderivative size = 935, normalized size of antiderivative = 4.35

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x, algorithm="maxima")
```

3.52. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$

output

```

1/60*(c^6*(20*(33*sin(f*x + e)/(cos(f*x + e) + 1) - 76*sin(f*x + e)^3/(cos
(f*x + e) + 1)^3 + 51*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^3 - 3*a^3*si
n(f*x + e)^2/(cos(f*x + e) + 1)^2 + 3*a^3*sin(f*x + e)^4/(cos(f*x + e) + 1
)^4 - a^3*sin(f*x + e)^6/(cos(f*x + e) + 1)^6) + (735*sin(f*x + e)/(cos(f*
x + e) + 1) + 50*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(c
os(f*x + e) + 1)^5)/a^3 - 690*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3
+ 690*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) + 6*c^6*(60*(5*sin(f*
x + e)/(cos(f*x + e) + 1) - 7*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^3 -
2*a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^3*sin(f*x + e)^4/(cos(f*x +
e) + 1)^4) + (465*sin(f*x + e)/(cos(f*x + e) + 1) + 40*sin(f*x + e)^3/(cos
(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 390*log(si
n(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 390*log(sin(f*x + e)/(cos(f*x + e
) + 1) - 1)/a^3) + 45*c^6*(40*sin(f*x + e)/((a^3 - a^3*sin(f*x + e)^2/(cos
(f*x + e) + 1)^2)*(cos(f*x + e) + 1)) + (85*sin(f*x + e)/(cos(f*x + e) + 1
) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + sin(f*x + e)^5/(cos(f*x + e)
+ 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin
(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) + 20*c^6*((105*sin(f*x + e)/(cos(f*
x + e) + 1) + 20*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(c
os(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3
+ 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) + 15*c^6*(15*sin(f*x...

```

3.52.8 Giac [A] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.82

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^6}{(a + a\sec(e + fx))^3} dx =$$

$$\frac{3465 c^6 \log\left(\left|\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) + 1\right|\right)}{a^3} - \frac{3465 c^6 \log\left(\left|\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) - 1\right|\right)}{a^3} + \frac{10 \left(267 c^6 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 - 472 c^6 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + 213 c^6 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)\right)}{\left(\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 1\right)^3 a^3}$$

30 f

input

```

integrate(sec(f*x+e)*(c-c*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x, algorithm="g
iac")

```

output

```

-1/30*(3465*c^6*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^3 - 3465*c^6*log(abs(
tan(1/2*f*x + 1/2*e) - 1))/a^3 + 10*(267*c^6*tan(1/2*f*x + 1/2*e)^5 - 472*
c^6*tan(1/2*f*x + 1/2*e)^3 + 213*c^6*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x +
1/2*e)^2 - 1)^3*a^3) - 32*(3*a^12*c^6*tan(1/2*f*x + 1/2*e)^5 + 20*a^12*c^
6*tan(1/2*f*x + 1/2*e)^3 + 150*a^12*c^6*tan(1/2*f*x + 1/2*e))/a^15)/f

```

3.52.9 Mupad [B] (verification not implemented)

Time = 12.97 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.90

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{160c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a^3 f}$$

$$- \frac{89c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 - \frac{472c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3} + 71c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - a^3 \right)}$$

$$+ \frac{64c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3a^3 f} + \frac{16c^6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{5a^3 f} - \frac{231c^6 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a^3 f}$$

input `int((c - c/cos(e + f*x))^6/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`output `(160*c^6*tan(e/2 + (f*x)/2))/(a^3*f) - (89*c^6*tan(e/2 + (f*x)/2)^5 - (472*c^6*tan(e/2 + (f*x)/2)^3)/3 + 71*c^6*tan(e/2 + (f*x)/2))/(f*(3*a^3*tan(e/2 + (f*x)/2)^2 - 3*a^3*tan(e/2 + (f*x)/2)^4 + a^3*tan(e/2 + (f*x)/2)^6 - a^3)) + (64*c^6*tan(e/2 + (f*x)/2)^3)/(3*a^3*f) + (16*c^6*tan(e/2 + (f*x)/2)^5)/(5*a^3*f) - (231*c^6*atanh(tan(e/2 + (f*x)/2)))/(a^3*f)`

$$3.53 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$$

3.53.1	Optimal result	442
3.53.2	Mathematica [C] (verified)	443
3.53.3	Rubi [A] (verified)	443
3.53.4	Maple [A] (verified)	447
3.53.5	Fricas [A] (verification not implemented)	448
3.53.6	Sympy [F]	449
3.53.7	Maxima [B] (verification not implemented)	449
3.53.8	Giac [A] (verification not implemented)	450
3.53.9	Mupad [B] (verification not implemented)	451

3.53.1 Optimal result

Integrand size = 32, antiderivative size = 193

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx = -\frac{63c^5 \operatorname{arctanh}(\sin(e+fx))}{2a^3 f} + \frac{42c^5 \tan(e+fx)}{a^3 f} - \frac{21c^5 \sec(e+fx) \tan(e+fx)}{2a^3 f} - \frac{6c^2 (c-c\sec(e+fx))^3 \tan(e+fx)}{5af(a+a\sec(e+fx))^2} + \frac{2c(c-c\sec(e+fx))^4 \tan(e+fx)}{5f(a+a\sec(e+fx))^3} + \frac{42c(c^2-c^2\sec(e+fx))^2 \tan(e+fx)}{5f(a^3+a^3\sec(e+fx))}$$

output `-63/2*c^5*arctanh(sin(f*x+e))/a^3/f+42*c^5*tan(f*x+e)/a^3/f-21/2*c^5*sec(f*x+e)*tan(f*x+e)/a^3/f-6/5*c^2*(c-c*sec(f*x+e))^3*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2+2/5*c*(c-c*sec(f*x+e))^4*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+42/5*c*(c^2-c^2*sec(f*x+e))^2*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))`

3.53.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.39 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.39

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx = \frac{32c^5 \operatorname{Hypergeometric2F1}\left(-\frac{9}{2}, -\frac{5}{2}, -\frac{3}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)} \tan(e+fx)}{5a^3 f(-1+\sec(e+fx))(1+\sec(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^3,x]`

output `(-32*c^5*Hypergeometric2F1[-9/2, -5/2, -3/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]]*Tan[e + f*x])/(5*a^3*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^3)`

3.53.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {3042, 4445, 3042, 4445, 3042, 4445, 3042, 4275, 3042, 4254, 24, 4534, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a\sec(e+fx)+a)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^5}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\ & \quad \downarrow \text{4445} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^4}{5f(a\sec(e+fx)+a)^3} - \frac{9c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(\sec(e+fx)a+a)^2} dx}{5a} \\ & \quad \downarrow \text{3042} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^4}{5f(a\sec(e+fx)+a)^3} - \frac{9c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^4}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a} \end{aligned}$$

3.53. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$

$$\begin{array}{c}
\downarrow 4445 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{7c \int \frac{\sec(e+fx)(c-c \sec(e+fx))^3}{\sec(e+fx)a+a} dx}{3a} \right)}{5a} \\
\downarrow 3042 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{7c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c \csc(e+fx+\frac{\pi}{2}))^3}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} \right)}{5a} \\
\downarrow 4445 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \int \sec(e+fx)(c-c \sec(e+fx))^2 dx}{a} \right)}{3a} \right)}{5a} \\
\downarrow 3042 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \int \csc(e+fx+\frac{\pi}{2})(c-c \csc(e+fx+\frac{\pi}{2}))^2 dx}{a} \right)}{3a} \right)}{5a} \\
\downarrow 4275 \\
\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\int \sec(e+fx)(\sec^2(e+fx)c^2+c^2) dx - 2c^2 \int \sec^2(e+fx) dx \right)}{a} \right)}{3a} \right)}{5a} \\
\downarrow 3042
\end{array}$$

3.53. $\int \frac{\sec(e+fx)(c-c \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$

$$9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\int \csc(e+fx+\frac{\pi}{2}) \left(\csc(e+fx+\frac{\pi}{2})^2 c^2 + c^2 \right) dx - 2c^2 \int \csc(e+fx+\frac{\pi}{2})^2 dx \right)}{a} \right)}{3a} \right)$$

5a

↓ 4254

$$9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\frac{2c^2 \int 1d(-\tan(e+fx))}{f} + \int \csc(e+fx+\frac{\pi}{2}) \left(\csc(e+fx+\frac{\pi}{2})^2 c^2 + c^2 \right) dx \right)}{a} \right)}{3a} \right)$$

5a

↓ 24

$$9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\int \csc(e+fx+\frac{\pi}{2}) \left(\csc(e+fx+\frac{\pi}{2})^2 c^2 + c^2 \right) dx - \frac{2c^2 \tan(e+fx)}{f} \right)}{a} \right)}{3a} \right)$$

5a

↓ 4534

$$9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\frac{3}{2} c^2 \int \sec(e+fx) dx - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \right)}{3a} \right)$$

5a

↓ 3042

3.53. $\int \frac{\sec(e+fx)(c-c \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$

$$\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\frac{3}{2}c^2 \int \csc(e+fx+\frac{\pi}{2}) dx - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \right)}{3a} \right)}{5a}$$

↓ 4257

$$\frac{2c \tan(e+fx)(c-c \sec(e+fx))^4}{5f(a \sec(e+fx)+a)^3} - \frac{9c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^3}{3f(a \sec(e+fx)+a)^2} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{f(a \sec(e+fx)+a)} - \frac{5c \left(\frac{3c^2 \operatorname{arctanh}(\sin(e+fx))}{2f} - \frac{2c^2 \tan(e+fx)}{f} + \frac{c^2 \tan(e+fx) \sec(e+fx)}{2f} \right)}{a} \right)}{3a} \right)}{5a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^3,x]`

output `(2*c*(c - c*Sec[e + f*x])^4*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) - (9*c*((2*c*(c - c*Sec[e + f*x])^3*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (7*c*((2*c*(c - c*Sec[e + f*x])^2*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (5*c*((3*c^2*ArcTanh[Sin[e + f*x]])/(2*f) - (2*c^2*Tan[e + f*x])/f + (c^2*Sec[e + f*x]*Tan[e + f*x])/(2*f))/a))/(3*a)))/(5*a)`

3.53.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.53. $\int \frac{\sec(e+fx)(c-c \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4275 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^2, x_Symbol] := Simp[2*a*(b/d) Int[(d*Csc[e + f*x])^(n + 1), x], x] + Int[(d*Csc[e + f*x])^n*(a^2 + b^2*Csc[e + f*x]^2), x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.53.4 Maple [A] (verified)

Time = 0.98 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.70

3.53. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$

method	result
derivativedivides	$8c^5 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^2} - \frac{17}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)} - \frac{63 \ln(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)}{16} \right) \frac{1}{fa^3}$
default	$8c^5 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)^2} - \frac{17}{16(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)} - \frac{63 \ln(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)}{16} \right) \frac{1}{fa^3}$
parallelrisc	$3749 \left(\frac{2520(1 + \cos(2fx + 2e)) \ln(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1)}{3749} + \frac{2520(-1 - \cos(2fx + 2e)) \ln(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1)}{3749} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \sec\left(\frac{fx}{2} + \frac{e}{2}\right)^4 \cos\left(\frac{fx}{2} + \frac{e}{2}\right) \right) \frac{1}{80fa^3(1 + \cos(2fx + 2e))}$
risc	$\frac{ic^5(325e^{8i(fx+e)} + 1545e^{7i(fx+e)} + 3805e^{6i(fx+e)} + 5545e^{5i(fx+e)} + 7351e^{4i(fx+e)} + 6115e^{3i(fx+e)} + 4407e^{2i(fx+e)} + 2107e^{i(fx+e)} + 1001)}{5fa^3(e^{i(fx+e)} + 1)^5(1 + e^{2i(fx+e)})^2}$
norman	$\frac{-\frac{63c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{294c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} - \frac{2688c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5af} + \frac{474c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{af} - \frac{193c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{af} + \frac{24c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^5 a^2}$

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 8/f*c^5/a^3*(1/5*tan(1/2*f*x+1/2*e)^5+tan(1/2*f*x+1/2*e)^3+6*tan(1/2*f*x+1/2*e)+1/16/(tan(1/2*f*x+1/2*e)+1)^2-17/16/(tan(1/2*f*x+1/2*e)+1)-63/16*ln(tan(1/2*f*x+1/2*e)+1)-1/16/(tan(1/2*f*x+1/2*e)-1)^2-17/16/(tan(1/2*f*x+1/2*e)-1)+63/16*ln(tan(1/2*f*x+1/2*e)-1))
```

3.53.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.30

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^5}{(a + a \sec(e + fx))^3} dx = \frac{315(c^5 \cos(fx + e))^5 + 3c^5 \cos(fx + e)^4 + 3c^5 \cos(fx + e)^3 + c^5 \cos(fx + e)^2 \log(\sin(fx + e) + 1)}{a^2}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x, algorithm="fracas")
```

3.53. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$

output
$$\begin{aligned} & -1/20*(315*(c^5*\cos(f*x + e)^5 + 3*c^5*\cos(f*x + e)^4 + 3*c^5*\cos(f*x + e) \\ & \quad \wedge 3 + c^5*\cos(f*x + e)^2)*\log(\sin(f*x + e) + 1) - 315*(c^5*\cos(f*x + e)^5 + \\ & \quad 3*c^5*\cos(f*x + e)^4 + 3*c^5*\cos(f*x + e)^3 + c^5*\cos(f*x + e)^2)*\log(-\sin \\ & \quad \text{n}(f*x + e) + 1) - 2*(496*c^5*\cos(f*x + e)^4 + 1163*c^5*\cos(f*x + e)^3 + 80 \\ & \quad 1*c^5*\cos(f*x + e)^2 + 65*c^5*\cos(f*x + e) - 5*c^5)*\sin(f*x + e))/(a^3*f*c \\ & \quad \text{os}(f*x + e)^5 + 3*a^3*f*\cos(f*x + e)^4 + 3*a^3*f*\cos(f*x + e)^3 + a^3*f*c \\ & \quad \text{s}(f*x + e)^2) \end{aligned}$$

3.53.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx =$$

$$c^5 \left(\int \left(-\frac{\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} \right) dx + \int \frac{5\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \left(-\frac{1}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} \right) dx \right)$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**5/(a+a*sec(f*x+e))**3,x)`

output
$$\begin{aligned} & -c**5*(Integral(-sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec \\ & \quad (e + f*x) + 1), x) + Integral(5*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e \\ & \quad + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(-10*sec(e + f*x)**3/(sec(e \\ & \quad + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(10*sec \\ & \quad (e + f*x)**4/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x \\ & \quad) + Integral(-5*sec(e + f*x)**5/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*s \\ & \quad \text{ec}(e + f*x) + 1), x) + Integral(sec(e + f*x)**6/(sec(e + f*x)**3 + 3*sec(e \\ & \quad + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3 \end{aligned}$$

3.53.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 680 vs. $2(186) = 372$.

Time = 0.21 (sec) , antiderivative size = 680, normalized size of antiderivative = 3.52

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$$

$$= c^5 \left(\frac{60 \left(\frac{5 \sin(fx+e)}{\cos(fx+e)+1} - \frac{7 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^3 - \frac{2a^3 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{a^3 \sin(fx+e)^4}{(\cos(fx+e)+1)^4}} + \frac{\frac{465 \sin(fx+e)}{\cos(fx+e)+1} + \frac{40 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5}}{a^3} - \frac{390 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^3} + \frac{390 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^3} \right)$$

3.53.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output
$$\frac{1}{60}c^5\left(\frac{60(5\sin(fx+e))}{(\cos(fx+e)+1)} - \frac{7\sin(fx+e)^3}{(\cos(fx+e)+1)^3}\right) / (a^3 - 2a^3\sin(fx+e)^2/(\cos(fx+e)+1)^2 + a^3\sin(fx+e)^4/(\cos(fx+e)+1)^4) + \frac{465\sin(fx+e)}{(\cos(fx+e)+1)} + \frac{40\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3\sin(fx+e)^5}{(\cos(fx+e)+1)^5} / a^3 - \frac{390\log(\sin(fx+e)/(\cos(fx+e)+1)+1)}{a^3} + \frac{390\log(\sin(fx+e)/(\cos(fx+e)+1)-1)}{a^3} + \frac{15c^5(40\sin(fx+e))}{(a^3 - a^3\sin(fx+e)^2/(\cos(fx+e)+1)^2)(\cos(fx+e)+1)} + \frac{85\sin(fx+e)}{(\cos(fx+e)+1)} + \frac{10\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} / a^3 - \frac{60\log(\sin(fx+e)/(\cos(fx+e)+1)+1)}{a^3} + \frac{60\log(\sin(fx+e)/(\cos(fx+e)+1)-1)}{a^3} + \frac{10c^5((105\sin(fx+e))}{(\cos(fx+e)+1)} + \frac{20\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3\sin(fx+e)^5}{(\cos(fx+e)+1)^5})}{a^3} - \frac{60\log(\sin(fx+e)/(\cos(fx+e)+1)+1)}{a^3} + \frac{60\log(\sin(fx+e)/(\cos(fx+e)+1)-1)}{a^3} + \frac{10c^5(15\sin(fx+e))}{(\cos(fx+e)+1)} + \frac{10\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3\sin(fx+e)^5}{(\cos(fx+e)+1)^5} / a^3 + \frac{c^5(15\sin(fx+e))}{(\cos(fx+e)+1)} - \frac{10\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3\sin(fx+e)^5}{(\cos(fx+e)+1)^5} / a^3 - \frac{15c^5(5\sin(fx+e))}{(\cos(fx+e)+1)} - \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} / a^3) / f$$

3.53.8 Giac [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.82

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx = \frac{315c^5 \log\left(\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+1\right|\right)}{a^3} - \frac{315c^5 \log\left(\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-1\right|\right)}{a^3} + \frac{10\left(17c^5 \tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3 - 15c^5 \tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)\right)}{\left(\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^2-1\right)^2 a^3} - \frac{16\left(a^{12}c^5 \tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)\right)}{10f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output
$$\begin{aligned} & -1/10*(315*c^5*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a^3 - 315*c^5*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a^3 + 10*(17*c^5*\tan(1/2*f*x + 1/2*e)^3 - 15*c^5*\tan(1/2*f*x + 1/2*e))/((\tan(1/2*f*x + 1/2*e)^2 - 1)^2*a^3) - 16*(a^{12}*c^5*\tan(1/2*f*x + 1/2*e)^5 + 5*a^{12}*c^5*\tan(1/2*f*x + 1/2*e)^3 + 30*a^{12}*c^5*\tan(1/2*f*x + 1/2*e))/a^{15})/f \end{aligned}$$

3.53.9 Mupad [B] (verification not implemented)

Time = 12.96 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.82

$$\begin{aligned} \int \frac{\sec(e+fx)(c-c\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx &= \frac{48c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a^3 f} \\ & - \frac{17c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 - 15c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + a^3 \right)} \\ & + \frac{8c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{a^3 f} + \frac{8c^5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{5a^3 f} \\ & - \frac{63c^5 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a^3 f} \end{aligned}$$

input $\text{int}((c - c/\cos(e + f*x))^5/(\cos(e + f*x)*(a + a/\cos(e + f*x))^3),x)$

output
$$\begin{aligned} & (48*c^5*\tan(e/2 + (f*x)/2))/(a^3*f) - (17*c^5*\tan(e/2 + (f*x)/2)^3 - 15*c^5*\tan(e/2 + (f*x)/2))/(f*(a^3*\tan(e/2 + (f*x)/2)^4 - 2*a^3*\tan(e/2 + (f*x)/2)^2 + a^3)) + (8*c^5*\tan(e/2 + (f*x)/2)^3)/(a^3*f) + (8*c^5*\tan(e/2 + (f*x)/2)^5)/(5*a^3*f) - (63*c^5*\operatorname{atanh}(\tan(e/2 + (f*x)/2)))/(a^3*f) \end{aligned}$$

$$3.54 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

3.54.1	Optimal result	452
3.54.2	Mathematica [C] (verified)	453
3.54.3	Rubi [A] (verified)	453
3.54.4	Maple [A] (verified)	457
3.54.5	Fricas [A] (verification not implemented)	457
3.54.6	Sympy [F]	458
3.54.7	Maxima [B] (verification not implemented)	458
3.54.8	Giac [A] (verification not implemented)	459
3.54.9	Mupad [B] (verification not implemented)	460

3.54.1 Optimal result

Integrand size = 32, antiderivative size = 164

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx = -\frac{7c^4 \operatorname{arctanh}(\sin(e+fx))}{a^3 f} + \frac{7c^4 \tan(e+fx)}{a^3 f} \\ + \frac{2c(c-c\sec(e+fx))^3 \tan(e+fx)}{5f(a+a\sec(e+fx))^3} \\ - \frac{14(c^2-c^2\sec(e+fx))^2 \tan(e+fx)}{15af(a+a\sec(e+fx))^2} \\ + \frac{14(c^4-c^4\sec(e+fx)) \tan(e+fx)}{3f(a^3+a^3\sec(e+fx))}$$

output

```
-7*c^4*arctanh(sin(f*x+e))/a^3/f+7*c^4*tan(f*x+e)/a^3/f+2/5*c*(c-c*sec(f*x
+e))^3*tan(f*x+e)/f/(a+a*sec(f*x+e))^3-14/15*(c^2-c^2*sec(f*x+e))^2*tan(f*
x+e)/a/f/(a+a*sec(f*x+e))^2+14/3*(c^4-c^4*sec(f*x+e))*tan(f*x+e)/f/(a^3+a^
3*sec(f*x+e))
```

3.54.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.93 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.46

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx = \frac{16c^4 \operatorname{Hypergeometric2F1}\left(-\frac{7}{2}, -\frac{5}{2}, -\frac{3}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \sqrt{2-2\sec(e+fx)} \tan(e+fx)}{5a^3 f(-1+\sec(e+fx))(1+\sec(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^3,x]`

output `(-16*c^4*Hypergeometric2F1[-7/2, -5/2, -3/2, (1 + Sec[e + f*x])/2]*Sqrt[2 - 2*Sec[e + f*x]]*Tan[e + f*x])/(5*a^3*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^3)`

3.54.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.01, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 4445, 3042, 4445, 3042, 4445, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a\sec(e+fx)+a)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c-c\csc\left(e+fx+\frac{\pi}{2}\right))^4}{(a\csc\left(e+fx+\frac{\pi}{2}\right)+a)^3} dx \\ & \quad \downarrow \text{4445} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{5f(a\sec(e+fx)+a)^3} - \frac{7c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(\sec(e+fx)a+a)^2} dx}{5a} \\ & \quad \downarrow \text{3042} \\ & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{5f(a\sec(e+fx)+a)^3} - \frac{7c \int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c-c\csc\left(e+fx+\frac{\pi}{2}\right))^3}{(\csc\left(e+fx+\frac{\pi}{2}\right)a+a)^2} dx}{5a} \end{aligned}$$

3.54. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$

$$\begin{array}{c}
\downarrow 4445 \\
\frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{5f(a\sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{\sec(e+fx)a+a} dx}{3a} \right)}{5a} \\
\downarrow 3042 \\
\frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{5f(a\sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} \right)}{5a} \\
\downarrow 4445 \\
\frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{5f(a\sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c \int \sec(e+fx)(c-c\sec(e+fx)) dx}{a} \right)}{3a} \right)}{5a} \\
\downarrow 3042 \\
\frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{5f(a\sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c \int \csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2})) dx}{a} \right)}{3a} \right)}{5a} \\
\downarrow 4274 \\
\frac{2c \tan(e+fx)(c-c\sec(e+fx))^3}{5f(a\sec(e+fx)+a)^3} - \frac{7c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^2}{3f(a\sec(e+fx)+a)^2} - \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2\sec(e+fx))}{f(a\sec(e+fx)+a)} - \frac{3c(c \int \sec(e+fx) dx - c \int \sec^2(e+fx) dx)}{a} \right)}{3a} \right)}{5a} \\
\downarrow 3042
\end{array}$$

3.54. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$

$$7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{3f(a \sec(e+fx)+a)^2} - \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2 \sec(e+fx))}{f(a \sec(e+fx)+a)} - \frac{3c \left(c \int \csc(e+fx+\frac{\pi}{2}) dx - c \int \csc(e+fx+\frac{\pi}{2})^2 dx \right)}{a} \right)}{3a} \right)$$

5a

↓ 4254

$$7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{3f(a \sec(e+fx)+a)^2} - \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2 \sec(e+fx))}{f(a \sec(e+fx)+a)} - \frac{3c \left(\frac{c \int 1d(-\tan(e+fx))}{f} + c \int \csc(e+fx+\frac{\pi}{2}) dx \right)}{a} \right)}{3a} \right)$$

5a

↓ 24

$$7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{3f(a \sec(e+fx)+a)^2} - \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2 \sec(e+fx))}{f(a \sec(e+fx)+a)} - \frac{3c \left(c \int \csc(e+fx+\frac{\pi}{2}) dx - \frac{c \tan(e+fx)}{f} \right)}{a} \right)}{3a} \right)$$

5a

↓ 4257

$$7c \left(\frac{2c \tan(e+fx)(c-c \sec(e+fx))^2}{3f(a \sec(e+fx)+a)^2} - \frac{5c \left(\frac{2 \tan(e+fx)(c^2-c^2 \sec(e+fx))}{f(a \sec(e+fx)+a)} - \frac{3c \left(\frac{\operatorname{arctanh}(\sin(e+fx))}{f} - \frac{c \tan(e+fx)}{f} \right)}{a} \right)}{3a} \right)$$

5a

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^3,x]`


```
output (2*c*(c - c*Sec[e + f*x])^3*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) - (
7*c*((2*c*(c - c*Sec[e + f*x])^2*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2
) - (5*c*((2*(c^2 - c^2*Sec[e + f*x])*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]
)) - (3*c*((c*ArcTanh[Sin[e + f*x]])/f - (c*Tan[e + f*x])/f)/a)/(3*a)))/
(5*a)
```

3.54.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

```
rule 4274 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d In
t[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]
```

```
rule 4445 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(cs
c[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[2*a*c*Cot[e +
f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))),
x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*
x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^
(-1)] && IntegerQ[2*m]
```

3.54.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.66

method	result
derivativedivides	$\frac{4c^4 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} - \frac{7 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{7 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} \right)}{fa^3}$
default	$\frac{4c^4 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} - \frac{7 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4} - \frac{1}{4\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{7 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4} \right)}{fa^3}$
parallelrirsch	$\frac{1609c^4 \left(\frac{1680 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) \cos(fx+e)}{1609} - \frac{1680 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \cos(fx+e)}{1609} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \sec\left(\frac{fx}{2} + \frac{e}{2}\right)^4 \left(\cos(fx+e) + \sin(fx+e)\right) \right)}{240fa^3 \cos(fx+e)}$
rirsch	$\frac{2ic^4 \left(120e^{6i(fx+e)} + 495e^{5i(fx+e)} + 1235e^{4i(fx+e)} + 1270e^{3i(fx+e)} + 1342e^{2i(fx+e)} + 715e^{i(fx+e)} + 167 \right)}{15fa^3(1+e^{2i(fx+e)})(e^{i(fx+e)}+1)^5} - \frac{7c^4 \ln(e^{i(fx+e)}+1)}{a^3}$
norman	$\frac{\frac{14c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} - \frac{154c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} + \frac{1022c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{15af} - \frac{186c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{5af} + \frac{92c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{15af} - \frac{8c^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{15af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^4 a^2}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `4/f*c^4/a^3*(1/5*tan(1/2*f*x+1/2*e)^5+2/3*tan(1/2*f*x+1/2*e)^3+3*tan(1/2*f*x+1/2*e)-1/4/(tan(1/2*f*x+1/2*e)+1)-7/4*ln(tan(1/2*f*x+1/2*e)+1)-1/4/(tan(1/2*f*x+1/2*e)-1)+7/4*ln(tan(1/2*f*x+1/2*e)-1))`

3.54.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.41

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx = \frac{105(c^4 \cos(fx+e)^4 + 3c^4 \cos(fx+e)^3 + 3c^4 \cos(fx+e)^2 + c^4 \cos(fx+e)) \log(\sin(fx+e)+1) - \dots}{\dots}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x, algorithm="fricas")`

output
$$\begin{aligned} & -1/30*(105*(c^4*\cos(f*x + e))^4 + 3*c^4*\cos(f*x + e)^3 + 3*c^4*\cos(f*x + e) \\ & \quad \wedge 2 + c^4*\cos(f*x + e))*\log(\sin(f*x + e) + 1) - 105*(c^4*\cos(f*x + e))^4 + 3 \\ & \quad *c^4*\cos(f*x + e)^3 + 3*c^4*\cos(f*x + e)^2 + c^4*\cos(f*x + e))*\log(-\sin(f* \\ & \quad x + e) + 1) - 2*(167*c^4*\cos(f*x + e)^3 + 381*c^4*\cos(f*x + e)^2 + 277*c^4 \\ & \quad *\cos(f*x + e) + 15*c^4)*\sin(f*x + e))/(a^3*f*\cos(f*x + e)^4 + 3*a^3*f*\cos(\\ & \quad f*x + e)^3 + 3*a^3*f*\cos(f*x + e)^2 + a^3*f*\cos(f*x + e)) \end{aligned}$$

3.54.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^4}{(a + a \sec(e + fx))^3} dx$$

$$= \frac{c^4 \left(\int \frac{\sec(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx + \int \left(-\frac{4 \sec^2(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} \right) dx + \int \frac{6}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx \right)}{a^3}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**4/(a+a*sec(f*x+e))**3,x)`

output
$$\begin{aligned} & c**4*(Integral(\sec(e + f*x)/(\sec(e + f*x)**3 + 3*\sec(e + f*x)**2 + 3*\sec(e \\ & \quad + f*x) + 1), x) + Integral(-4*\sec(e + f*x)**2/(\sec(e + f*x)**3 + 3*\sec(e \\ & \quad + f*x)**2 + 3*\sec(e + f*x) + 1), x) + Integral(6*\sec(e + f*x)**3/(\sec(e + \\ & \quad f*x)**3 + 3*\sec(e + f*x)**2 + 3*\sec(e + f*x) + 1), x) + Integral(-4*\sec(e \\ & \quad + f*x)**4/(\sec(e + f*x)**3 + 3*\sec(e + f*x)**2 + 3*\sec(e + f*x) + 1), x) + \\ & \quad Integral(\sec(e + f*x)**5/(\sec(e + f*x)**3 + 3*\sec(e + f*x)**2 + 3*\sec(e + \\ & \quad f*x) + 1), x))/a**3 \end{aligned}$$

3.54.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 470 vs. $2(161) = 322$.

Time = 0.22 (sec) , antiderivative size = 470, normalized size of antiderivative = 2.87

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^4}{(a + a \sec(e + fx))^3} dx$$

$$= \frac{3c^4 \left(\frac{40 \sin(fx+e)}{\left(a^3 - \frac{a^3 \sin^2(fx+e)}{(\cos(fx+e)+1)^2}\right)(\cos(fx+e)+1)} + \frac{\frac{85 \sin(fx+e)}{\cos(fx+e)+1} + \frac{10 \sin^3(fx+e)}{(\cos(fx+e)+1)^3} + \frac{\sin^5(fx+e)}{(\cos(fx+e)+1)^5}}{a^3} - \frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^3} + \frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^3} \right)}{a^3}$$

3.54.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output
$$\frac{1}{60} \cdot (3c^4 \cdot (40 \sin(fx + e) / ((a^3 - a^3 \sin(fx + e))^2 / (\cos(fx + e) + 1)^2) \cdot (\cos(fx + e) + 1)) + (85 \sin(fx + e) / (\cos(fx + e) + 1) + 10 \sin(fx + e)^3 / (\cos(fx + e) + 1)^3 + \sin(fx + e)^5 / (\cos(fx + e) + 1)^5) / a^3 - 60 \log(\sin(fx + e) / (\cos(fx + e) + 1) + 1) / a^3 + 60 \log(\sin(fx + e) / (\cos(fx + e) + 1) - 1) / a^3 + 4c^4 \cdot ((105 \sin(fx + e) / (\cos(fx + e) + 1) + 20 \sin(fx + e)^3 / (\cos(fx + e) + 1)^3 + 3 \sin(fx + e)^5 / (\cos(fx + e) + 1)^5) / a^3 - 60 \log(\sin(fx + e) / (\cos(fx + e) + 1) + 1) / a^3 + 60 \log(\sin(fx + e) / (\cos(fx + e) + 1) - 1) / a^3 + 6c^4 \cdot (15 \sin(fx + e) / (\cos(fx + e) + 1) + 10 \sin(fx + e)^3 / (\cos(fx + e) + 1)^3 + 3 \sin(fx + e)^5 / (\cos(fx + e) + 1)^5) / a^3 + c^4 \cdot (15 \sin(fx + e) / (\cos(fx + e) + 1) - 10 \sin(fx + e)^3 / (\cos(fx + e) + 1)^3 + 3 \sin(fx + e)^5 / (\cos(fx + e) + 1)^5) / a^3 - 12c^4 \cdot (5 \sin(fx + e) / (\cos(fx + e) + 1) - \sin(fx + e)^5 / (\cos(fx + e) + 1)^5) / a^3) / f$$

3.54.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.86

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^4}{(a + a \sec(e + fx))^3} dx = \frac{105c^4 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a^3} - \frac{105c^4 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a^3} + \frac{30c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)a^3} - \frac{4(3a^{12}c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 10a^{12}c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + 45a^{12}c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{15f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output
$$-1/15 \cdot (105c^4 \cdot \log(\text{abs}(\tan(1/2 \cdot fx + 1/2 \cdot e) + 1)) / a^3 - 105c^4 \cdot \log(\text{abs}(\tan(1/2 \cdot fx + 1/2 \cdot e) - 1)) / a^3 + 30c^4 \cdot \tan(1/2 \cdot fx + 1/2 \cdot e) / ((\tan(1/2 \cdot fx + 1/2 \cdot e)^2 - 1) \cdot a^3) - 4 \cdot (3a^{12}c^4 \cdot \tan(1/2 \cdot fx + 1/2 \cdot e)^5 + 10a^{12}c^4 \cdot \tan(1/2 \cdot fx + 1/2 \cdot e)^3 + 45a^{12}c^4 \cdot \tan(1/2 \cdot fx + 1/2 \cdot e)) / a^{15}) / f$$

3.54.9 Mupad [B] (verification not implemented)

Time = 13.04 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.77

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx = \frac{12c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a^3 f} + \frac{8c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3a^3 f}$$

$$+ \frac{4c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{5a^3 f} - \frac{14c^4 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a^3 f}$$

$$- \frac{2c^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - a^3\right)}$$

input `int((c - c/cos(e + f*x))^4/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`output `(12*c^4*tan(e/2 + (f*x)/2))/(a^3*f) + (8*c^4*tan(e/2 + (f*x)/2)^3)/(3*a^3*f) + (4*c^4*tan(e/2 + (f*x)/2)^5)/(5*a^3*f) - (14*c^4*atanh(tan(e/2 + (f*x)/2)))/(a^3*f) - (2*c^4*tan(e/2 + (f*x)/2))/(f*(a^3*tan(e/2 + (f*x)/2)^2 - a^3))`

3.55 $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$

3.55.1	Optimal result	461
3.55.2	Mathematica [A] (verified)	461
3.55.3	Rubi [A] (verified)	462
3.55.4	Maple [A] (verified)	464
3.55.5	Fricas [A] (verification not implemented)	465
3.55.6	Sympy [F]	465
3.55.7	Maxima [B] (verification not implemented)	466
3.55.8	Giac [A] (verification not implemented)	466
3.55.9	Mupad [B] (verification not implemented)	467

3.55.1 Optimal result

Integrand size = 32, antiderivative size = 131

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx = -\frac{c^3 \operatorname{arctanh}(\sin(e+fx))}{a^3 f} + \frac{2c^3 \tan(e+fx)}{f(a^3+a^3\sec(e+fx))} + \frac{2c(c-c\sec(e+fx))^2 \tan(e+fx)}{5f(a+a\sec(e+fx))^3} - \frac{2(c^3-c^3\sec(e+fx)) \tan(e+fx)}{3af(a+a\sec(e+fx))^2}$$

```
output -c^3*arctanh(sin(f*x+e))/a^3/f+2*c^3*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))+2/5
*c*(c-c*sec(f*x+e))^2*tan(f*x+e)/f/(a+a*sec(f*x+e))^3-2/3*(c^3-c^3*sec(f*x
+e))*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2
```

3.55.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.06

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx = \frac{c^3 \left(-\frac{\log(\cos(\frac{1}{2}(e+fx))-\sin(\frac{1}{2}(e+fx)))}{f} + \frac{\log(\cos(\frac{1}{2}(e+fx))+\sin(\frac{1}{2}(e+fx)))}{f} - \frac{26 \tan(\frac{1}{2}(e+fx))}{15f} + \frac{2 \sec^2(\frac{1}{2}(e+fx)) \tan(\frac{1}{2}(e+fx))}{15f} \right)}{a^3}$$

```
input Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^3,x]
```

output $-\left(\frac{c^3(-\log[\cos[(e+fx)/2] - \sin[(e+fx)/2]]/f) + \log[\cos[(e+fx)/2] + \sin[(e+fx)/2]]/f - (26\tan[(e+fx)/2])/(15f) + (2\sec[(e+fx)/2]^2\tan[(e+fx)/2])/(15f) - (2\sec[(e+fx)/2]^4\tan[(e+fx)/2])/(5f)}{a^3}\right)$

3.55.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4445, 3042, 4445, 3042, 4445, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a\sec(e+fx)+a)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^3}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\ & \quad \downarrow \text{4445} \\ & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{5f(a\sec(e+fx)+a)^3} - \frac{c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(\sec(e+fx)a+a)^2} dx}{a} \\ & \quad \downarrow \text{3042} \\ & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{5f(a\sec(e+fx)+a)^3} - \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{a} \\ & \quad \downarrow \text{4445} \\ & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{5f(a\sec(e+fx)+a)^3} - \frac{c \left(\frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{3f(a\sec(e+fx)+a)^2} - \frac{c \int \frac{\sec(e+fx)(c-c\sec(e+fx))}{\sec(e+fx)a+a} dx}{a} \right)}{a} \\ & \quad \downarrow \text{3042} \\ & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^2}{5f(a\sec(e+fx)+a)^3} - \frac{c \left(\frac{2\tan(e+fx)(c^2-c^2\sec(e+fx))}{3f(a\sec(e+fx)+a)^2} - \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{a} \right)}{a} \end{aligned}$$

3.55. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$

$$\begin{aligned}
& \downarrow 4445 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{5f(a \sec(e+fx) + a)^3} - \frac{c \left(\frac{2 \tan(e+fx)(c^2 - c^2 \sec(e+fx))}{3f(a \sec(e+fx) + a)^2} - \frac{c \left(\frac{2c \tan(e+fx)}{f(a \sec(e+fx) + a)} - \frac{c \int \sec(e+fx) dx}{a} \right)}{a} \right)}{a} \\
& \downarrow 3042 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{5f(a \sec(e+fx) + a)^3} - \frac{c \left(\frac{2 \tan(e+fx)(c^2 - c^2 \sec(e+fx))}{3f(a \sec(e+fx) + a)^2} - \frac{c \left(\frac{2c \tan(e+fx)}{f(a \sec(e+fx) + a)} - \frac{c \int \csc(e+fx + \frac{\pi}{2}) dx}{a} \right)}{a} \right)}{a} \\
& \downarrow 4257 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^2}{5f(a \sec(e+fx) + a)^3} - \frac{c \left(\frac{2 \tan(e+fx)(c^2 - c^2 \sec(e+fx))}{3f(a \sec(e+fx) + a)^2} - \frac{c \left(\frac{2c \tan(e+fx)}{f(a \sec(e+fx) + a)} - \frac{c \operatorname{arctanh}(\sin(e+fx))}{af} \right)}{a} \right)}{a}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^3,x]`

output `(2*c*(c - c*Sec[e + f*x])^2*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) - (c*((2*(c^2 - c^2*Sec[e + f*x])*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (c*(-((c*ArcTanh[Sin[e + f*x]])/(a*f)) + (2*c*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))))/a))/a`

3.55.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`


```
rule 4445 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e +
f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))),
x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*
x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^
(-1)] && IntegerQ[2*m]
```

3.55.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.58

method	result
derivativedivides	$\frac{2c^3 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} \right)}{fa^3}$
default	$\frac{2c^3 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2} \right)}{fa^3}$
parallelrisc	$\frac{c^3 \left(6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 + 10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 15 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - 15 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) + 30 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \right)}{15a^3f}$
risc	$\frac{4ic^3(15e^{4i(fx+e)} + 30e^{3i(fx+e)} + 100e^{2i(fx+e)} + 50e^{i(fx+e)} + 13)}{15fa^3(e^{i(fx+e)} + 1)^5} - \frac{c^3 \ln(e^{i(fx+e)} + i)}{a^3f} + \frac{c^3 \ln(e^{i(fx+e)} - i)}{a^3f}$
norman	$\frac{-\frac{2c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{16c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} - \frac{22c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5af} + \frac{6c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{5af} - \frac{8c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{15af} + \frac{2c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{5af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^3 a^2} + \dots$

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBO
SE)
```

```
output 2/f*c^3/a^3*(1/5*tan(1/2*f*x+1/2*e)^5+1/3*tan(1/2*f*x+1/2*e)^3+tan(1/2*f*x
+1/2*e)+1/2*ln(tan(1/2*f*x+1/2*e)-1)-1/2*ln(tan(1/2*f*x+1/2*e)+1))
```

3.55.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$$

3.55.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.47

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx = \frac{15(c^3 \cos(fx+e)^3 + 3c^3 \cos(fx+e)^2 + 3c^3 \cos(fx+e) + c^3) \log(\sin(fx+e)+1) - 15(c^3 \cos(fx+e)^3 + 3c^3 \cos(fx+e)^2 + 3c^3 \cos(fx+e) + c^3) \log(-\sin(fx+e)+1) - 4(13c^3 \cos(fx+e)^2 + 24c^3 \cos(fx+e) + 23c^3) \sin(fx+e)}{30(a^3 f \cos(fx+e)^3 + 3a^3 f \cos(fx+e)^2 + 3a^3 f \cos(fx+e) + a^3 f)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x, algorithm="fracas")`

output `-1/30*(15*(c^3*cos(f*x + e)^3 + 3*c^3*cos(f*x + e)^2 + 3*c^3*cos(f*x + e) + c^3)*log(sin(f*x + e) + 1) - 15*(c^3*cos(f*x + e)^3 + 3*c^3*cos(f*x + e)^2 + 3*c^3*cos(f*x + e) + c^3)*log(-sin(f*x + e) + 1) - 4*(13*c^3*cos(f*x + e)^2 + 24*c^3*cos(f*x + e) + 23*c^3)*sin(f*x + e))/(a^3*f*cos(f*x + e)^3 + 3*a^3*f*cos(f*x + e)^2 + 3*a^3*f*cos(f*x + e) + a^3*f)`

3.55.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx = \frac{c^3 \left(\int \left(-\frac{\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} \right) dx + \int \frac{3\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \left(-\frac{1}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} \right) dx \right)}{a^3}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**3/(a+a*sec(f*x+e))**3,x)`

output `-c**3*(Integral(-sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(3*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(-3*sec(e + f*x)**3/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**4/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3`

3.55.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(129) = 258$.

Time = 0.21 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.32

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{c^3 \left(\frac{\frac{105 \sin(fx+e)}{\cos(fx+e)+1} + \frac{20 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5}}{a^3} - \frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^3} + \frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^3} \right) + \frac{3c^3 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} + \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^3}}{60f}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x, algorithm="maxima")
```

```
output 1/60*(c^3*((105*sin(f*x + e)/(cos(f*x + e) + 1) + 20*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) + 3*c^3*(15*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 + c^3*(15*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 9*c^3*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3)/f
```

3.55.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.83

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx =$$

$$-\frac{15c^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{a^3} - \frac{15c^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{a^3} - \frac{2\left(3a^{12}c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 + 5a^{12}c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + 15a^{12}c^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{a^{15}}}{15f}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x, algorithm="giac")
```

```
output -1/15*(15*c^3*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^3 - 15*c^3*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a^3 - 2*(3*a^12*c^3*tan(1/2*f*x + 1/2*e)^5 + 5*a^12*c^3*tan(1/2*f*x + 1/2*e)^3 + 15*a^12*c^3*tan(1/2*f*x + 1/2*e))/a^15)/f
```

3.55. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$

3.55.9 Mupad [B] (verification not implemented)

Time = 13.04 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.47

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{2c^3 \left(15 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) - 15 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) + 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 \right)}{15a^3 f}$$

input `int((c - c/cos(e + f*x))^3/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`

output `(2*c^3*(15*tan(e/2 + (f*x)/2) - 15*atanh(tan(e/2 + (f*x)/2)) + 5*tan(e/2 + (f*x)/2)^3 + 3*tan(e/2 + (f*x)/2)^5)/(15*a^3*f)`

$$3.56 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

3.56.1	Optimal result	468
3.56.2	Mathematica [A] (verified)	468
3.56.3	Rubi [A] (verified)	469
3.56.4	Maple [A] (verified)	470
3.56.5	Fricas [B] (verification not implemented)	470
3.56.6	Sympy [F]	471
3.56.7	Maxima [B] (verification not implemented)	471
3.56.8	Giac [A] (verification not implemented)	472
3.56.9	Mupad [B] (verification not implemented)	472

3.56.1 Optimal result

Integrand size = 32, antiderivative size = 38

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx = \frac{(c-c\sec(e+fx))^2 \tan(e+fx)}{5f(a+a\sec(e+fx))^3}$$

output `1/5*(c-c*sec(f*x+e))^2*tan(f*x+e)/f/(a+a*sec(f*x+e))^3`

3.56.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx = \frac{c^2 \tan^5\left(\frac{1}{2}(e+fx)\right)}{5a^3 f}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^3,x]`

output `(c^2*Tan[(e + f*x)/2]^5)/(5*a^3*f)`

3.56.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a\sec(e+fx)+a)^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^2}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx$$

↓ 4438

$$\frac{\tan(e+fx)(c-c\sec(e+fx))^2}{5f(a\sec(e+fx)+a)^3}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^3,x]`

output `((c - c*Sec[e + f*x])^2*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3)`

3.56.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.56.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.61

method	result	size
derivativedivides	$\frac{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5f a^3}$	23
default	$\frac{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5f a^3}$	23
parallelrisch	$\frac{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5f a^3}$	23
risch	$\frac{2ic^2(5e^{4i(fx+e)} + 10e^{2i(fx+e)} + 1)}{5f a^3(e^{i(fx+e)} + 1)^5}$	50
norman	$\frac{\frac{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5af} - \frac{2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{5af} + \frac{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{5af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 a^2}$	87

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/5/f*c^2/a^3*tan(1/2*f*x+1/2*e)^5
```

3.56.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(37) = 74.

Time = 0.26 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.16

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{(c^2 \cos(fx+e)^2 - 2c^2 \cos(fx+e) + c^2) \sin(fx+e)}{5(a^3 f \cos(fx+e)^3 + 3a^3 f \cos(fx+e)^2 + 3a^3 f \cos(fx+e) + a^3 f)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x, algorithm="fracas")
```

```
output 1/5*(c^2*cos(f*x + e)^2 - 2*c^2*cos(f*x + e) + c^2)*sin(f*x + e)/(a^3*f*cos(f*x + e)^3 + 3*a^3*f*cos(f*x + e)^2 + 3*a^3*f*cos(f*x + e) + a^3*f)
```

3.56.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{c^2 \left(\int \frac{\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \left(-\frac{2\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} \right) dx + \int \frac{1}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx \right)}{a^3}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**2/(a+a*sec(f*x+e))**3,x)`

output `c**2*(Integral(sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(-2*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3`

3.56.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs. $2(37) = 74$.

Time = 0.20 (sec) , antiderivative size = 185, normalized size of antiderivative = 4.87

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{c^2 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} + \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3} + \frac{c^2 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} - \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{60 f a^3} - \frac{6 c^2 \left(\frac{5 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)}{(\cos(fx+e)+1)} \right)}{a^3}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/60*(c^2*(15*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 + c^2*(15*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 6*c^2*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)/(cos(f*x + e) + 1))/a^3/f`

3.56.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^2}{(a + a \sec(e + fx))^3} dx = \frac{c^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5}{5a^3 f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output `1/5*c^2*tan(1/2*f*x + 1/2*e)^5/(a^3*f)`

3.56.9 Mupad [B] (verification not implemented)

Time = 13.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^2}{(a + a \sec(e + fx))^3} dx = \frac{c^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{5a^3 f}$$

input `int((c - c/cos(e + f*x))^2/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`

output `(c^2*tan(e/2 + (f*x)/2)^5)/(5*a^3*f)`

$$3.57 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

3.57.1	Optimal result	473
3.57.2	Mathematica [A] (verified)	473
3.57.3	Rubi [A] (verified)	474
3.57.4	Maple [A] (verified)	475
3.57.5	Fricas [A] (verification not implemented)	476
3.57.6	Sympy [F]	476
3.57.7	Maxima [A] (verification not implemented)	477
3.57.8	Giac [A] (verification not implemented)	477
3.57.9	Mupad [B] (verification not implemented)	477

3.57.1 Optimal result

Integrand size = 30, antiderivative size = 76

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx = \frac{(c-c\sec(e+fx))\tan(e+fx)}{5f(a+a\sec(e+fx))^3} + \frac{(c-c\sec(e+fx))\tan(e+fx)}{15af(a+a\sec(e+fx))^2}$$

output `1/5*(c-c*sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+1/15*(c-c*sec(f*x+e))*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2`

3.57.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.57

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx = -\frac{c(-1+\sec(e+fx))(4+\sec(e+fx))\tan(e+fx)}{15a^3f(1+\sec(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^3,x]`

output `-1/15*(c*(-1 + Sec[e + f*x])*(4 + Sec[e + f*x])*Tan[e + f*x])/(a^3*f*(1 + Sec[e + f*x])^3)`

3.57. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$

3.57.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a\sec(e+fx)+a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\
 & \quad \downarrow \text{4439} \\
 & \frac{\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(\sec(e+fx)a+a)^2} dx}{5a} + \frac{\tan(e+fx)(c-c\sec(e+fx))}{5f(a\sec(e+fx)+a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a} + \frac{\tan(e+fx)(c-c\sec(e+fx))}{5f(a\sec(e+fx)+a)^3} \\
 & \quad \downarrow \text{4438} \\
 & \frac{\tan(e+fx)(c-c\sec(e+fx))}{15af(a\sec(e+fx)+a)^2} + \frac{\tan(e+fx)(c-c\sec(e+fx))}{5f(a\sec(e+fx)+a)^3}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^3,x]`

output `((c - c*Sec[e + f*x])*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) + ((c - c*Sec[e + f*x])*Tan[e + f*x])/(15*a*f*(a + a*Sec[e + f*x])^2)`

3.57.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)]`

3.57.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.47

method	result	size
parallelsch	$\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 \left(3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 5\right)}{30a^3 f}$	36
derivativedivides	$\frac{c \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3}\right)}{2f a^3}$	37
default	$\frac{c \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3}\right)}{2f a^3}$	37
risch	$\frac{2ic(15e^{4i(fx+e)} + 15e^{3i(fx+e)} + 25e^{2i(fx+e)} + 5e^{i(fx+e)} + 4)}{15fa^3(e^{i(fx+e)} + 1)^5}$	70
norman	$\frac{\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{6af} - \frac{4c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{15af} + \frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{10af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)a^2}$	81

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/30*c*tan(1/2*f*x+1/2*e)^3*(3*tan(1/2*f*x+1/2*e)^2-5)/a^3/f
```

3.57.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.04

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{(4c\cos(fx+e)^2 - 3c\cos(fx+e) - c)\sin(fx+e)}{15(a^3f\cos(fx+e)^3 + 3a^3f\cos(fx+e)^2 + 3a^3f\cos(fx+e) + a^3f)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="fricas")
```

```
output 1/15*(4*c*cos(f*x + e)^2 - 3*c*cos(f*x + e) - c)*sin(f*x + e)/(a^3*f*cos(f*x + e)^3 + 3*a^3*f*cos(f*x + e)^2 + 3*a^3*f*cos(f*x + e) + a^3*f)
```

3.57.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= -\frac{c\left(\int \left(-\frac{\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1}\right) dx + \int \frac{\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx\right)}{a^3}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**3,x)
```

```
output -c*(Integral(-sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3
```

3.57.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))}{(a + a \sec(e + fx))^3} dx$$

$$= \frac{c \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} - \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3} - \frac{3c \left(\frac{5 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3}$$

$$60f$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/60*(c*(15*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 3*c*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3)/f`

3.57.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.49

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))}{(a + a \sec(e + fx))^3} dx = \frac{3c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 5c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3}{30a^3f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output `1/30*(3*c*tan(1/2*f*x + 1/2*e)^5 - 5*c*tan(1/2*f*x + 1/2*e)^3)/(a^3*f)`

3.57.9 Mupad [B] (verification not implemented)

Time = 13.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.46

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))}{(a + a \sec(e + fx))^3} dx = \frac{c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 5\right)}{30a^3f}$$

input `int((c - c/cos(e + f*x))/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`

output `(c*tan(e/2 + (f*x)/2)^3*(3*tan(e/2 + (f*x)/2)^2 - 5))/(30*a^3*f)`

3.57. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$

3.58 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))} dx$

3.58.1	Optimal result	478
3.58.2	Mathematica [A] (verified)	478
3.58.3	Rubi [A] (verified)	479
3.58.4	Maple [A] (verified)	480
3.58.5	Fricas [A] (verification not implemented)	480
3.58.6	Sympy [F]	481
3.58.7	Maxima [A] (verification not implemented)	481
3.58.8	Giac [A] (verification not implemented)	482
3.58.9	Mupad [B] (verification not implemented)	482

3.58.1 Optimal result

Integrand size = 32, antiderivative size = 78

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))} dx = -\frac{2 \cot^5(e+fx)}{5a^3cf} + \frac{\csc(e+fx)}{a^3cf} - \frac{\csc^3(e+fx)}{a^3cf} + \frac{2 \csc^5(e+fx)}{5a^3cf}$$

output `-2/5*cot(f*x+e)^5/a^3/c/f+csc(f*x+e)/a^3/c/f-csc(f*x+e)^3/a^3/c/f+2/5*csc(f*x+e)^5/a^3/c/f`

3.58.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.86

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))} dx = \frac{(-2 + \sec(e+fx) + 4 \sec^2(e+fx) + 2 \sec^3(e+fx)) \tan(e+fx)}{5a^3cf(-1 + \sec(e+fx))(1 + \sec(e+fx))^3}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])),x]`

output `((-2 + Sec[e + f*x] + 4*Sec[e + f*x]^2 + 2*Sec[e + f*x]^3)*Tan[e + f*x])/ (5*a^3*c*f*(-1 + Sec[e + f*x])*(1 + Sec[e + f*x])^3)`

3.58. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))} dx$

3.58.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^3 (c - c \sec(e+fx))} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2}) + a)^3 (c - c \csc(e+fx+\frac{\pi}{2}))} dx$$

↓ 4446

$$\frac{\int (c^2 \csc(e+fx) \cot^5(e+fx) - 2c^2 \csc^2(e+fx) \cot^4(e+fx) + c^2 \csc^3(e+fx) \cot^3(e+fx)) dx}{a^3 c^3}$$

↓ 2009

$$\frac{\frac{2c^2 \cot^5(e+fx)}{5f} - \frac{2c^2 \csc^5(e+fx)}{5f} + \frac{c^2 \csc^3(e+fx)}{f} - \frac{c^2 \csc(e+fx)}{f}}{a^3 c^3}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])),x]`

output `-(((2*c^2*Cot[e + f*x]^5)/(5*f) - (c^2*Csc[e + f*x])/f + (c^2*Csc[e + f*x]^3)/f - (2*c^2*Csc[e + f*x]^5)/(5*f))/(a^3*c^3))`

3.58.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.58. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))} dx$


```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.58.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.76

method	result	size
parallelrisch	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 5 \cot\left(\frac{fx}{2} + \frac{e}{2}\right) + 15 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{40 f a^3 c}$	59
derivativedivides	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{8 f a^3 c}$	61
default	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}}{8 f a^3 c}$	61
risch	$\frac{2i(5e^{5i(fx+e)} + 10e^{4i(fx+e)} + 10e^{3i(fx+e)} - 3e^{i(fx+e)} - 2)}{5fa^3c(e^{i(fx+e)} + 1)^5(e^{i(fx+e)} - 1)}$	85
norman	$\frac{\frac{1}{8acf} + \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{8acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{8acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{40acf}}{a^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}$	94

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE
)
```

```
output 1/40*(tan(1/2*f*x+1/2*e)^5-5*tan(1/2*f*x+1/2*e)^3+5*cot(1/2*f*x+1/2*e)+15*
tan(1/2*f*x+1/2*e))/f/a^3/c
```

3.58.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.97

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))} dx$$

$$= -\frac{2 \cos^3(fx + e) - \cos^2(fx + e) - 4 \cos(fx + e) - 2}{5(a^3 c f \cos^2(fx + e) + 2 a^3 c f \cos(fx + e) + a^3 c f) \sin(fx + e)}$$

3.58.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `-1/5*(2*cos(f*x + e)^3 - cos(f*x + e)^2 - 4*cos(f*x + e) - 2)/((a^3*c*f*cos(f*x + e)^2 + 2*a^3*c*f*cos(f*x + e) + a^3*c*f)*sin(f*x + e))`

3.58.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))} dx = -\frac{\int \frac{\sec(e + fx)}{\sec^4(e + fx) + 2 \sec^3(e + fx) - 2 \sec(e + fx) - 1} dx}{a^3 c}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e)),x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**4 + 2*sec(e + f*x)**3 - 2*sec(e + f*x) - 1), x)/(a**3*c)`

3.58.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.22

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))} dx$$

$$= \frac{\frac{15 \sin(fx+e)}{\cos(fx+e)+1} - \frac{5 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} + \frac{5(\cos(fx+e)+1)}{a^3 c \sin(fx+e)}}{40 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `1/40*((15*sin(f*x + e)/(cos(f*x + e) + 1) - 5*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^3*c) + 5*(cos(f*x + e) + 1)/(a^3*c*sin(f*x + e)))/f`

3.58.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.12

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))} dx$$

$$= \frac{\frac{5}{a^3 c \tan(\frac{1}{2} fx + \frac{1}{2} e)} + \frac{a^{12} c^4 \tan(\frac{1}{2} fx + \frac{1}{2} e)^5 - 5 a^{12} c^4 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + 15 a^{12} c^4 \tan(\frac{1}{2} fx + \frac{1}{2} e)}{a^{15} c^5}}{40 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e)),x, algorithm="giac")`

output `1/40*(5/(a^3*c*tan(1/2*f*x + 1/2*e)) + (a^12*c^4*tan(1/2*f*x + 1/2*e)^5 - 5*a^12*c^4*tan(1/2*f*x + 1/2*e)^3 + 15*a^12*c^4*tan(1/2*f*x + 1/2*e))/(a^15*c^5))/f`

3.58.9 Mupad [B] (verification not implemented)

Time = 12.95 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))} dx$$

$$= -\frac{16 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 28 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 8 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1}{40 a^3 c f \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^5 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))),x)`

output `-(8*cos(e/2 + (f*x)/2)^2 - 28*cos(e/2 + (f*x)/2)^4 + 16*cos(e/2 + (f*x)/2)^6 - 1)/(40*a^3*c*f*cos(e/2 + (f*x)/2)^5*sin(e/2 + (f*x)/2))`

3.59
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^2} dx$$

3.59.1	Optimal result	483
3.59.2	Mathematica [A] (verified)	483
3.59.3	Rubi [A] (verified)	484
3.59.4	Maple [A] (verified)	485
3.59.5	Fricas [A] (verification not implemented)	485
3.59.6	Sympy [F]	486
3.59.7	Maxima [A] (verification not implemented)	486
3.59.8	Giac [A] (verification not implemented)	487
3.59.9	Mupad [B] (verification not implemented)	487

3.59.1 Optimal result

Integrand size = 32, antiderivative size = 80

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^2} dx$$

$$= -\frac{\cot^5(e+fx)}{5a^3c^2f} + \frac{\csc(e+fx)}{a^3c^2f} - \frac{2 \csc^3(e+fx)}{3a^3c^2f} + \frac{\csc^5(e+fx)}{5a^3c^2f}$$

output `-1/5*cot(f*x+e)^5/a^3/c^2/f+csc(f*x+e)/a^3/c^2/f-2/3*csc(f*x+e)^3/a^3/c^2/f+1/5*csc(f*x+e)^5/a^3/c^2/f`

3.59.2 Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.99

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^2} dx$$

$$= \frac{(3 - 12 \sec(e+fx) - 12 \sec^2(e+fx) + 8 \sec^3(e+fx) + 8 \sec^4(e+fx)) \tan(e+fx)}{15a^3c^2f(-1 + \sec(e+fx))^2(1 + \sec(e+fx))^3}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^2),x]`

output `((3 - 12*Sec[e + f*x] - 12*Sec[e + f*x]^2 + 8*Sec[e + f*x]^3 + 8*Sec[e + f*x]^4)*Tan[e + f*x])/(15*a^3*c^2*f*(-1 + Sec[e + f*x])^2*(1 + Sec[e + f*x])^3)`

3.59.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^2} dx$$

3.59.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.86, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^3 (c - c \sec(e+fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^3 (c - c \csc(e+fx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{4446} \\
 & - \frac{\int (c \cot^5(e+fx) \csc(e+fx) - c \cot^4(e+fx) \csc^2(e+fx)) dx}{a^3 c^3} \\
 & \quad \downarrow \text{2009} \\
 & - \frac{\frac{c \cot^5(e+fx)}{5f} - \frac{c \csc^5(e+fx)}{5f} + \frac{2c \csc^3(e+fx)}{3f} - \frac{c \csc(e+fx)}{f}}{a^3 c^3}
 \end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^2),x]`

output `-(((c*Cot[e + f*x]^5)/(5*f) - (c*Csc[e + f*x])/f + (2*c*Csc[e + f*x]^3)/(3*f) - (c*Csc[e + f*x]^5)/(5*f))/(a^3*c^3))`

3.59.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.59.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.92

method	result	size
parallelrisch	$\frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 20 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - 5 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 90 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 60 \cot\left(\frac{fx}{2} + \frac{e}{2}\right)}{240 f a^3 c^2}$	74
derivativedivides	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{4}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}}{16 f c^2 a^3}$	76
default	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{4}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)} - \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}}{16 f c^2 a^3}$	76
risch	$\frac{2i(15e^{7i(fx+e)} + 15e^{6i(fx+e)} - 5e^{5i(fx+e)} - 25e^{4i(fx+e)} + 13e^{3i(fx+e)} + 21e^{2i(fx+e)} + 9e^{i(fx+e)} - 3)}{15f c^2 a^3 (e^{i(fx+e)} + 1)^5 (e^{i(fx+e)} - 1)^3}$	118
norman	$\frac{-\frac{1}{48acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{4acf} + \frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{8acf} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{12acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{80acf}}{a^2 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}$	119

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x,method=_RETURNVERBO
SE)
```

```
output 1/240*(3*tan(1/2*f*x+1/2*e)^5-20*tan(1/2*f*x+1/2*e)^3-5*cot(1/2*f*x+1/2*e)
^3+90*tan(1/2*f*x+1/2*e)+60*cot(1/2*f*x+1/2*e))/f/a^3/c^2
```

3.59.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.36

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^2} dx$$

$$= \frac{3 \cos(fx + e)^4 - 12 \cos(fx + e)^3 - 12 \cos(fx + e)^2 + 8 \cos(fx + e) + 8}{15 (a^3 c^2 f \cos(fx + e)^3 + a^3 c^2 f \cos(fx + e)^2 - a^3 c^2 f \cos(fx + e) - a^3 c^2 f \sin(fx + e))}$$

3.59.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 (c-c \sec(e+fx))^2} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x, algorithm="fricas")`

output `-1/15*(3*cos(f*x + e)^4 - 12*cos(f*x + e)^3 - 12*cos(f*x + e)^2 + 8*cos(f*x + e) + 8)/((a^3*c^2*f*cos(f*x + e)^3 + a^3*c^2*f*cos(f*x + e)^2 - a^3*c^2*f*cos(f*x + e) - a^3*c^2*f)*sin(f*x + e))`

3.59.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^2} dx$$

$$= \frac{\int \frac{\sec(e+fx)}{\sec^5(e+fx) + \sec^4(e+fx) - 2\sec^3(e+fx) - 2\sec^2(e+fx) + \sec(e+fx) + 1} dx}{a^3 c^2}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**2,x)`

output `Integral(sec(e + f*x)/(sec(e + f*x)**5 + sec(e + f*x)**4 - 2*sec(e + f*x)**3 - 2*sec(e + f*x)**2 + sec(e + f*x) + 1), x)/(a**3*c**2)`

3.59.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.50

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^2} dx$$

$$= \frac{\frac{90 \sin(fx+e)}{\cos(fx+e)+1} - \frac{20 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5}}{a^3 c^2} + \frac{5 \left(\frac{12 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - 1 \right) (\cos(fx+e)+1)^3}{a^3 c^2 \sin(fx+e)^3}$$

$$= \frac{\hspace{15em}}{240 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output `1/240*((90*sin(f*x + e)/(cos(f*x + e) + 1) - 20*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^3*c^2) + 5*(12*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 1)*(cos(f*x + e) + 1)^3/(a^3*c^2*sin(f*x + e)^3))/f`

3.59.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.29

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^2} dx$$

$$= \frac{5 \left(12 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 1\right)}{a^3 c^2 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3} + \frac{3 a^{12} c^8 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 - 20 a^{12} c^8 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + 90 a^{12} c^8 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)}{a^{15} c^{10}}$$

$$= \frac{\hspace{15em}}{240 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `1/240*(5*(12*tan(1/2*f*x + 1/2*e)^2 - 1)/(a^3*c^2*tan(1/2*f*x + 1/2*e)^3) + (3*a^12*c^8*tan(1/2*f*x + 1/2*e)^5 - 20*a^12*c^8*tan(1/2*f*x + 1/2*e)^3 + 90*a^12*c^8*tan(1/2*f*x + 1/2*e))/(a^15*c^10))/f`

3.59.9 Mupad [B] (verification not implemented)

Time = 13.24 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.39

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^2} dx$$

$$= \frac{48 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 192 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 168 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 32 \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 3}{240 a^3 c^2 f \left(\cos\left(\frac{e}{2} + \frac{fx}{2}\right)^5 \sin\left(\frac{e}{2} + \frac{fx}{2}\right) - \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^7 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^2),x)`

output `(168*cos(e/2 + (f*x)/2)^4 - 32*cos(e/2 + (f*x)/2)^2 - 192*cos(e/2 + (f*x)/2)^6 + 48*cos(e/2 + (f*x)/2)^8 + 3)/(240*a^3*c^2*f*(cos(e/2 + (f*x)/2)^5*sin(e/2 + (f*x)/2) - cos(e/2 + (f*x)/2)^7*sin(e/2 + (f*x)/2))`

$$3.60 \quad \int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx$$

3.60.1	Optimal result	488
3.60.2	Mathematica [A] (verified)	488
3.60.3	Rubi [A] (verified)	489
3.60.4	Maple [A] (verified)	491
3.60.5	Fricas [A] (verification not implemented)	491
3.60.6	Sympy [F]	492
3.60.7	Maxima [A] (verification not implemented)	492
3.60.8	Giac [A] (verification not implemented)	492
3.60.9	Mupad [B] (verification not implemented)	493

3.60.1 Optimal result

Integrand size = 32, antiderivative size = 59

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx = \frac{\csc(e+fx)}{a^3c^3f} - \frac{2\csc^3(e+fx)}{3a^3c^3f} + \frac{\csc^5(e+fx)}{5a^3c^3f}$$

output `csc(f*x+e)/a^3/c^3/f-2/3*csc(f*x+e)^3/a^3/c^3/f+1/5*csc(f*x+e)^5/a^3/c^3/f`

3.60.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.85

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx = -\frac{-\frac{\csc(e+fx)}{f} + \frac{2\csc^3(e+fx)}{3f} - \frac{\csc^5(e+fx)}{5f}}{a^3c^3}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^3),x]`

output `-((- (Csc[e + f*x]/f) + (2*Csc[e + f*x]^3)/(3*f) - Csc[e + f*x]^5/(5*f))/(a^3*c^3))`

3.60. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx$

3.60.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.69, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4446, 3042, 25, 3086, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)^3(c-c \sec(e+fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)^3(c-c \csc(e+fx+\frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{4446} \\
 & \frac{\int \cot^5(e+fx) \csc(e+fx) dx}{a^3 c^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -\sec(e+fx-\frac{\pi}{2}) \tan(e+fx-\frac{\pi}{2})^5 dx}{a^3 c^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \sec(\frac{1}{2}(2e-\pi)+fx) \tan(\frac{1}{2}(2e-\pi)+fx)^5 dx}{a^3 c^3} \\
 & \quad \downarrow \text{3086} \\
 & \frac{\int (\csc^2(e+fx)-1)^2 d \csc(e+fx)}{a^3 c^3 f} \\
 & \quad \downarrow \text{210} \\
 & \frac{\int (\csc^4(e+fx)-2 \csc^2(e+fx)+1) d \csc(e+fx)}{a^3 c^3 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\frac{1}{5} \csc^5(e+fx) - \frac{2}{3} \csc^3(e+fx) + \csc(e+fx)}{a^3 c^3 f}
 \end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^3),x]`

3.60. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^3} dx$

output $(\text{Csc}[e + f*x] - (2*\text{Csc}[e + f*x]^3)/3 + \text{Csc}[e + f*x]^5/5)/(a^3*c^3*f)$

3.60.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$

rule 210 $\text{Int}[(a + (b \cdot x)^2)^p, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

rule 2009 $\text{Int}[u, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3086 $\text{Int}[(a \cdot \sec(e + f \cdot x) + (b \cdot x))^m \cdot ((c \cdot \tan(e + f \cdot x) + (d \cdot x))^n), x_Symbol] \rightarrow \text{Simp}[a/f \text{ Subst}[\text{Int}[(a*x)^{m-1} \cdot (-1 + x^2)^{(n-1)/2}], x], x, \text{Sec}[e + f*x], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n+1])$

rule 4446 $\text{Int}[\text{csc}(e + f \cdot x) \cdot (\text{csc}(e + f \cdot x) \cdot (b \cdot x) + a)^m \cdot (\text{csc}(e + f \cdot x) \cdot (d \cdot x) + c)^n, x_Symbol] \rightarrow \text{Simp}[((-a) \cdot c)^m \text{ Int}[\text{ExpandTrig}[\text{csc}[e + f*x] \cdot \text{cot}[e + f*x]^{(2*m)}, (c + d \cdot \text{csc}[e + f*x])^{(n-m)}], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{EqQ}[b \cdot c + a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{IntegersQ}[m, n] \ \&\& \ \text{GeQ}[n - m, 0] \ \&\& \ \text{GtQ}[m \cdot n, 0]$

3.60.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.69

method	result	size
default	$-\frac{\csc(fx+e)^5}{5} + \frac{2 \csc(fx+e)^3}{3} - \csc(fx+e)$	41
parallelrisc	$\frac{(15 \cos(4fx+4e) - 20 \cos(2fx+2e) + 29) \sec\left(\frac{fx}{2} + \frac{e}{2}\right)^5 \csc\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{3840f c^3 a^3}$	58
risc	$\frac{2i(15 e^{9i(fx+e)} - 20 e^{7i(fx+e)} + 58 e^{5i(fx+e)} - 20 e^{3i(fx+e)} + 15 e^{i(fx+e)})}{15f c^3 a^3 (e^{i(fx+e)} + 1)^5 (e^{i(fx+e)} - 1)^5}$	95
norman	$\frac{\frac{1}{160acf} - \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{96acf} + \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4}{16acf} + \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^6}{16acf} - \frac{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^8}{96acf} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{10}}{160acf}}{a^2 c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}$	141

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `-1/c^3/a^3/f*(-1/5*csc(f*x+e)^5+2/3*csc(f*x+e)^3-csc(f*x+e))`

3.60.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.29

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx$$

$$= \frac{15 \cos(fx+e)^4 - 20 \cos(fx+e)^2 + 8}{15(a^3 c^3 f \cos(fx+e)^4 - 2a^3 c^3 f \cos(fx+e)^2 + a^3 c^3 f \sin(fx+e))}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x, algorithm="fricas")`

output `1/15*(15*cos(f*x + e)^4 - 20*cos(f*x + e)^2 + 8)/((a^3*c^3*f*cos(f*x + e)^4 - 2*a^3*c^3*f*cos(f*x + e)^2 + a^3*c^3*f)*sin(f*x + e))`

3.60.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx = -\frac{\int \frac{\sec(e+fx)}{\sec^6(e+fx)-3\sec^4(e+fx)+3\sec^2(e+fx)-1} dx}{a^3c^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**3,x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**6 - 3*sec(e + f*x)**4 + 3*sec(e + f*x)**2 - 1), x)/(a**3*c**3)`

3.60.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.69

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx = \frac{15 \sin(fx+e)^4 - 10 \sin(fx+e)^2 + 3}{15a^3c^3f \sin(fx+e)^5}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/15*(15*sin(f*x + e)^4 - 10*sin(f*x + e)^2 + 3)/(a^3*c^3*f*sin(f*x + e)^5)`

3.60.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.69

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx = \frac{15 \sin(fx+e)^4 - 10 \sin(fx+e)^2 + 3}{15a^3c^3f \sin(fx+e)^5}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^3,x, algorithm="giac")`

output `1/15*(15*sin(f*x + e)^4 - 10*sin(f*x + e)^2 + 3)/(a^3*c^3*f*sin(f*x + e)^5)`

3.60.9 Mupad [B] (verification not implemented)

Time = 13.39 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.64

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^3} dx = \frac{\sin(e+fx)^4 - \frac{2\sin(e+fx)^2}{3} + \frac{1}{5}}{a^3 c^3 f \sin(e+fx)^5}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^3),x)`

output `(sin(e + f*x)^4 - (2*sin(e + f*x)^2)/3 + 1/5)/(a^3*c^3*f*sin(e + f*x)^5)`

3.61 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^4} dx$

3.61.1 Optimal result 494
 3.61.2 Mathematica [A] (verified) 494
 3.61.3 Rubi [A] (verified) 495
 3.61.4 Maple [A] (verified) 496
 3.61.5 Fricas [A] (verification not implemented) 497
 3.61.6 Sympy [F] 497
 3.61.7 Maxima [A] (verification not implemented) 498
 3.61.8 Giac [A] (verification not implemented) 498
 3.61.9 Mupad [B] (verification not implemented) 499

3.61.1 Optimal result

Integrand size = 32, antiderivative size = 99

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^4} dx$$

$$= -\frac{\cot^7(e+fx)}{7a^3c^4f} + \frac{\csc(e+fx)}{a^3c^4f} - \frac{\csc^3(e+fx)}{a^3c^4f} + \frac{3 \csc^5(e+fx)}{5a^3c^4f} - \frac{\csc^7(e+fx)}{7a^3c^4f}$$

output `-1/7*cot(f*x+e)^7/a^3/c^4/f+csc(f*x+e)/a^3/c^4/f-csc(f*x+e)^3/a^3/c^4/f+3/5*csc(f*x+e)^5/a^3/c^4/f-1/7*csc(f*x+e)^7/a^3/c^4/f`

3.61.2 Mathematica [A] (verified)

Time = 3.11 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^4} dx$$

$$= \frac{(-5 - 30 \sec(e+fx) + 30 \sec^2(e+fx) + 40 \sec^3(e+fx) - 40 \sec^4(e+fx) - 16 \sec^5(e+fx) + 16 \sec^6(e+fx)) \tan(e+fx)}{35a^3c^4f(-1 + \sec(e+fx))^4(1 + \sec(e+fx))^3}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^4),x]`

output `((-5 - 30*Sec[e + f*x] + 30*Sec[e + f*x]^2 + 40*Sec[e + f*x]^3 - 40*Sec[e + f*x]^4 - 16*Sec[e + f*x]^5 + 16*Sec[e + f*x]^6)*Tan[e + f*x])/(35*a^3*c^4*f*(-1 + Sec[e + f*x])^4*(1 + Sec[e + f*x])^3)`

3.61. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^4} dx$

3.61.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.82, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^4} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a\csc(e+fx+\frac{\pi}{2})+a)^3(c-c\csc(e+fx+\frac{\pi}{2}))^4} dx$$

↓ 4446

$$\int \frac{(a\csc(e+fx)\cot^7(e+fx) + a\csc^2(e+fx)\cot^6(e+fx)) dx}{a^4c^4}$$

↓ 2009

$$\frac{-\frac{a\cot^7(e+fx)}{7f} - \frac{a\csc^7(e+fx)}{7f} + \frac{3a\csc^5(e+fx)}{5f} - \frac{a\csc^3(e+fx)}{f} + \frac{a\csc(e+fx)}{f}}{a^4c^4}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^4),x]`

output `(-1/7*(a*Cot[e + f*x]^7)/f + (a*Csc[e + f*x])/f - (a*Csc[e + f*x]^3)/f + (3*a*Csc[e + f*x]^5)/(5*f) - (a*Csc[e + f*x]^7)/(7*f))/(a^4*c^4)`

3.61.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.61. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^4} dx$


```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[((-a)*c)^m
Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m
), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && Eq
Q[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.61.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00

method	result
parallelrisch	$-\frac{(-90 \cos(4fx+4e)+5 \cos(6fx+6e)+152 \cos(fx+e)+60 \cos(5fx+5e)-182-20 \cos(3fx+3e)+235 \cos(2fx+2e)) \sec\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{71680 f a^3 c^4}$
derivativedivides	$\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 + 15 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) - \frac{1}{7 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7} + \frac{20}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)} + \frac{6}{5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5} - \frac{5}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}}{64 f c^4 a^3}$
default	$\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 + 15 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) - \frac{1}{7 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7} + \frac{20}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)} + \frac{6}{5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5} - \frac{5}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}}{64 f c^4 a^3}$
risch	$\frac{2i(35 e^{11i(fx+e)} - 35 e^{10i(fx+e)} - 35 e^{9i(fx+e)} + 105 e^{8i(fx+e)} + 126 e^{7i(fx+e)} - 182 e^{6i(fx+e)} + 26 e^{5i(fx+e)} + 130 e^{4i(fx+e)} - 35 e^{3i(fx+e)} + 5 e^{2i(fx+e)} - 5 e^{i(fx+e)} + 1)}{35 f c^4 a^3 (e^{i(fx+e)} + 1)^5 (e^{i(fx+e)} - 1)^7}$
norman	$-\frac{\frac{1}{448acf} + \frac{3 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2}{160acf} - \frac{5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^4}{64acf} + \frac{5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^6}{16acf} + \frac{15 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^8}{64acf} - \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^{10}}{32acf} + \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^{12}}{320acf}}{a^2 c^3 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x,method=_RETURNVERBO
SE)
```

```
output -1/71680*(-90*cos(4*f*x+4*e)+5*cos(6*f*x+6*e)+152*cos(f*x+e)+60*cos(5*f*x+
5*e)-182-20*cos(3*f*x+3*e)+235*cos(2*f*x+2*e))*sec(1/2*f*x+1/2*e)^5*csc(1/
2*f*x+1/2*e)^7/f/a^3/c^4
```

$$3.61. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^4} dx$$

3.61.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.65

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^4} dx$$

$$= \frac{5 \cos(fx + e)^6 + 30 \cos(fx + e)^5 - 30 \cos(fx + e)^4 - 40 \cos(fx + e)^3 + 40 \cos(fx + e)^2 + 16 \cos(fx + e) - 16}{35 (a^3 c^4 f \cos(fx + e)^5 - a^3 c^4 f \cos(fx + e)^4 - 2 a^3 c^4 f \cos(fx + e)^3 + 2 a^3 c^4 f \cos(fx + e)^2 + a^3 c^4 f \cos(fx + e) - a^3 c^4 f) \sin(fx + e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x, algorithm="fracas")`

output `1/35*(5*cos(f*x + e)^6 + 30*cos(f*x + e)^5 - 30*cos(f*x + e)^4 - 40*cos(f*x + e)^3 + 40*cos(f*x + e)^2 + 16*cos(f*x + e) - 16)/((a^3*c^4*f*cos(f*x + e)^5 - a^3*c^4*f*cos(f*x + e)^4 - 2*a^3*c^4*f*cos(f*x + e)^3 + 2*a^3*c^4*f*cos(f*x + e)^2 + a^3*c^4*f*cos(f*x + e) - a^3*c^4*f)*sin(f*x + e))`

3.61.6 SymPy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^4} dx$$

$$= \frac{\int \frac{\sec(e+fx)}{\sec^7(e+fx) - \sec^6(e+fx) - 3\sec^5(e+fx) + 3\sec^4(e+fx) + 3\sec^3(e+fx) - 3\sec^2(e+fx) - \sec(e+fx) + 1} dx}{a^3 c^4}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**4,x)`

output `Integral(sec(e + f*x)/(sec(e + f*x)**7 - sec(e + f*x)**6 - 3*sec(e + f*x)**5 + 3*sec(e + f*x)**4 + 3*sec(e + f*x)**3 - 3*sec(e + f*x)**2 - sec(e + f*x) + 1), x)/(a**3*c**4)`

3.61.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.61

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^4} dx$$

$$= \frac{7 \left(\frac{75 \sin(fx+e)}{\cos(fx+e)+1} - \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3 c^4} + \frac{\left(\frac{42 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{175 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{700 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - 5 \right) (\cos(fx+e)+1)^7}{a^3 c^4 \sin(fx+e)^7}$$

$$= \frac{2240 f}{2240 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x, algorithm="maxima")`

output `1/2240*(7*(75*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^3*c^4) + (42*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 175*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 700*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 5)*(cos(f*x + e) + 1)^7/(a^3*c^4*sin(f*x + e)^7))/f`

3.61.8 Giac [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.29

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^4} dx$$

$$= \frac{700 \tan(\frac{1}{2} fx + \frac{1}{2} e)^6 - 175 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 + 42 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 5}{a^3 c^4 \tan(\frac{1}{2} fx + \frac{1}{2} e)^7} + \frac{7 (a^{12} c^{16} \tan(\frac{1}{2} fx + \frac{1}{2} e)^5 - 10 a^{12} c^{16} \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + 75 a^{12} c^{16} \tan(\frac{1}{2} fx + \frac{1}{2} e))}{a^{15} c^{20}}$$

$$= \frac{2240 f}{2240 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^4,x, algorithm="giac")`

output `1/2240*((700*tan(1/2*f*x + 1/2*e)^6 - 175*tan(1/2*f*x + 1/2*e)^4 + 42*tan(1/2*f*x + 1/2*e)^2 - 5)/(a^3*c^4*tan(1/2*f*x + 1/2*e)^7) + 7*(a^12*c^16*tan(1/2*f*x + 1/2*e)^5 - 10*a^12*c^16*tan(1/2*f*x + 1/2*e)^3 + 75*a^12*c^16*tan(1/2*f*x + 1/2*e))/(a^15*c^20))/f`

3.61.9 Mupad [B] (verification not implemented)

Time = 14.15 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.30

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^4} dx$$

$$= \frac{\left(2 \sin\left(\frac{e}{4} + \frac{fx}{4}\right)^2 - 1\right) \left(\frac{235 \sin(e+fx)^2}{16} - \frac{45 \sin(2e+2fx)^2}{8} + \frac{19 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^2}{2} + \frac{5 \sin(3e+3fx)^2}{16} - \frac{5 \sin\left(\frac{3e}{2} + \frac{3fx}{2}\right)^2}{4} + \frac{15 \sin\left(\frac{3e}{4} + \frac{3fx}{4}\right)^2}{4}\right)}{2240 a^3 c^4 f \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^7 \left(\sin\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)^3}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^4),x)`output `((2*sin(e/4 + (f*x)/4)^2 - 1)*((19*sin(e/2 + (f*x)/2)^2)/2 - (45*sin(2*e + 2*f*x)^2)/8 + (5*sin(3*e + 3*f*x)^2)/16 - (5*sin((3*e)/2 + (3*f*x)/2)^2)/4 + (15*sin((5*e)/2 + (5*f*x)/2)^2)/4 + (235*sin(e + f*x)^2)/16 - 5)/(2240*a^3*c^4*f*sin(e/2 + (f*x)/2)^7*(sin(e/2 + (f*x)/2)^2 - 1)^3)`

3.62
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^5} dx$$

3.62.1	Optimal result	500
3.62.2	Mathematica [A] (verified)	500
3.62.3	Rubi [A] (verified)	501
3.62.4	Maple [A] (verified)	502
3.62.5	Fricas [A] (verification not implemented)	503
3.62.6	Sympy [F]	503
3.62.7	Maxima [A] (verification not implemented)	504
3.62.8	Giac [A] (verification not implemented)	504
3.62.9	Mupad [B] (verification not implemented)	505

3.62.1 Optimal result

Integrand size = 32, antiderivative size = 120

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^5} dx \\ &= \frac{2 \cot^9(e+fx)}{9a^3c^5f} + \frac{\csc(e+fx)}{a^3c^5f} - \frac{5 \csc^3(e+fx)}{3a^3c^5f} \\ &+ \frac{9 \csc^5(e+fx)}{5a^3c^5f} - \frac{\csc^7(e+fx)}{a^3c^5f} + \frac{2 \csc^9(e+fx)}{9a^3c^5f} \end{aligned}$$

output $2/9*\cot(f*x+e)^9/a^3/c^5/f+\csc(f*x+e)/a^3/c^5/f-5/3*\csc(f*x+e)^3/a^3/c^5/f+9/5*\csc(f*x+e)^5/a^3/c^5/f-\csc(f*x+e)^7/a^3/c^5/f+2/9*\csc(f*x+e)^9/a^3/c^5/f$

3.62.2 Mathematica [A] (verified)

Time = 5.14 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.91

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^5} dx \\ &= \frac{(10 + 25 \sec(e+fx) - 60 \sec^2(e+fx) - 10 \sec^3(e+fx) + 80 \sec^4(e+fx) - 24 \sec^5(e+fx) - 32 \sec^6(e+fx))}{45a^3c^5f(-1 + \sec(e+fx))^5(1 + \sec(e+fx))^3} \end{aligned}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^5),x]`

3.62.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^5} dx$$

output $((10 + 25*\text{Sec}[e + f*x] - 60*\text{Sec}[e + f*x]^2 - 10*\text{Sec}[e + f*x]^3 + 80*\text{Sec}[e + f*x]^4 - 24*\text{Sec}[e + f*x]^5 - 32*\text{Sec}[e + f*x]^6 + 16*\text{Sec}[e + f*x]^7)*\text{Tan}[e + f*x])/(45*a^3*c^5*f*(-1 + \text{Sec}[e + f*x])^5*(1 + \text{Sec}[e + f*x])^3)$

3.62.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^5} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{(a \csc(e + fx + \frac{\pi}{2}) + a)^3 (c - c \csc(e + fx + \frac{\pi}{2}))^5} dx$$

↓ 4446

$$\int \frac{(a^2 \csc(e + fx) \cot^9(e + fx) + 2a^2 \csc^2(e + fx) \cot^8(e + fx) + a^2 \csc^3(e + fx) \cot^7(e + fx)) dx}{a^5 c^5}$$

↓ 2009

$$\frac{-\frac{2a^2 \cot^9(e+fx)}{9f} - \frac{2a^2 \csc^9(e+fx)}{9f} + \frac{a^2 \csc^7(e+fx)}{f} - \frac{9a^2 \csc^5(e+fx)}{5f} + \frac{5a^2 \csc^3(e+fx)}{3f} - \frac{a^2 \csc(e+fx)}{f}}{a^5 c^5}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^5),x]`

output $-(((-2*a^2*\text{Cot}[e + f*x]^9)/(9*f) - (a^2*\text{Csc}[e + f*x])/f + (5*a^2*\text{Csc}[e + f*x]^3)/(3*f) - (9*a^2*\text{Csc}[e + f*x]^5)/(5*f) + (a^2*\text{Csc}[e + f*x]^7)/f - (2*a^2*\text{Csc}[e + f*x]^9)/(9*f)))/(a^5*c^5)$

3.62.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.62.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.92

method	result
parallelrisch	$\frac{(-5 \cos(7fx+7e)-110 \cos(4fx+4e)-25 \cos(6fx+6e)+129 \cos(fx+e)+85 \cos(5fx+5e)-145 \cos(3fx+3e)+169 \cos(2fx+2e)-258) \sec(1/2fx+1/2e)^5 \csc(1/2fx+1/2e)^9 / f a^3 c^5}{184320 f a^3 c^5}$
derivativedivides	$\frac{\tan(\frac{fx}{2} + \frac{e}{2})^5}{5} - \frac{7 \tan(\frac{fx}{2} + \frac{e}{2})^3}{3} + 21 \tan(\frac{fx}{2} + \frac{e}{2}) + \frac{21}{5 \tan(\frac{fx}{2} + \frac{e}{2})^5} + \frac{35}{\tan(\frac{fx}{2} + \frac{e}{2})} - \frac{35}{3 \tan(\frac{fx}{2} + \frac{e}{2})^3} - \frac{1}{\tan(\frac{fx}{2} + \frac{e}{2})^7} + \frac{1}{9 \tan(\frac{fx}{2} + \frac{e}{2})^9}}{128 f c^5 a^3}$
default	$\frac{\tan(\frac{fx}{2} + \frac{e}{2})^5}{5} - \frac{7 \tan(\frac{fx}{2} + \frac{e}{2})^3}{3} + 21 \tan(\frac{fx}{2} + \frac{e}{2}) + \frac{21}{5 \tan(\frac{fx}{2} + \frac{e}{2})^5} + \frac{35}{\tan(\frac{fx}{2} + \frac{e}{2})} - \frac{35}{3 \tan(\frac{fx}{2} + \frac{e}{2})^3} - \frac{1}{\tan(\frac{fx}{2} + \frac{e}{2})^7} + \frac{1}{9 \tan(\frac{fx}{2} + \frac{e}{2})^9}}{128 f c^5 a^3}$
risch	$\frac{2i(45 e^{13i(fx+e)} - 90 e^{12i(fx+e)} + 30 e^{11i(fx+e)} + 240 e^{10i(fx+e)} - 69 e^{9i(fx+e)} - 354 e^{8i(fx+e)} + 516 e^{7i(fx+e)} + 96 e^{6i(fx+e)} - 45 f c^5 a^3 (e^{i(fx+e)} + 1)^5 (e^{i(fx+e)} - 1)^9)}{c^4 a^2 \tan(\frac{fx}{2} + \frac{e}{2})^9}$
norman	$\frac{\frac{1}{1152acf} - \frac{\tan(\frac{fx}{2} + \frac{e}{2})^2}{128acf} + \frac{21 \tan(\frac{fx}{2} + \frac{e}{2})^4}{640acf} - \frac{35 \tan(\frac{fx}{2} + \frac{e}{2})^6}{384acf} + \frac{35 \tan(\frac{fx}{2} + \frac{e}{2})^8}{128acf} + \frac{21 \tan(\frac{fx}{2} + \frac{e}{2})^{10}}{128acf} - \frac{7 \tan(\frac{fx}{2} + \frac{e}{2})^{12}}{384acf} + \frac{\tan(\frac{fx}{2} + \frac{e}{2})^{14}}{640acf}}{c^4 a^2 \tan(\frac{fx}{2} + \frac{e}{2})^9}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x,method=_RETURNVERBOSE)
```

```
output -1/184320*(-5*cos(7*f*x+7*e)-110*cos(4*f*x+4*e)-25*cos(6*f*x+6*e)+129*cos(f*x+e)+85*cos(5*f*x+5*e)-145*cos(3*f*x+3*e)+169*cos(2*f*x+2*e)-258)*sec(1/2*f*x+1/2*e)^5*csc(1/2*f*x+1/2*e)^9/f/a^3/c^5
```

$$3.62. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^5} dx$$

3.62.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.58

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^5} dx$$

$$= \frac{10 \cos(fx + e)^7 + 25 \cos(fx + e)^6 - 60 \cos(fx + e)^5 - 10 \cos(fx + e)^4 + 80 \cos(fx + e)^3}{45 (a^3 c^5 f \cos(fx + e)^6 - 2 a^3 c^5 f \cos(fx + e)^5 - a^3 c^5 f \cos(fx + e)^4 + 4 a^3 c^5 f \cos(fx + e)^3 - a^3 c^5 f \cos(fx + e)^2 + 4 a^3 c^5 f \cos(fx + e) - a^3 c^5 f)} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x, algorithm="fracas")`

output `1/45*(10*cos(f*x + e)^7 + 25*cos(f*x + e)^6 - 60*cos(f*x + e)^5 - 10*cos(f*x + e)^4 + 80*cos(f*x + e)^3 - 24*cos(f*x + e)^2 - 32*cos(f*x + e) + 16)/((a^3*c^5*f*cos(f*x + e)^6 - 2*a^3*c^5*f*cos(f*x + e)^5 - a^3*c^5*f*cos(f*x + e)^4 + 4*a^3*c^5*f*cos(f*x + e)^3 - a^3*c^5*f*cos(f*x + e)^2 - 2*a^3*c^5*f*cos(f*x + e) + a^3*c^5*f)*sin(f*x + e))`

3.62.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^5} dx$$

$$= - \frac{\int \frac{\sec(e+fx)}{\sec^8(e+fx) - 2\sec^7(e+fx) - 2\sec^6(e+fx) + 6\sec^5(e+fx) - 6\sec^3(e+fx) + 2\sec^2(e+fx) + 2\sec(e+fx) - 1} dx}{a^3 c^5}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**5,x)`

output `-Integral(sec(e + f*x)/(sec(e + f*x)**8 - 2*sec(e + f*x)**7 - 2*sec(e + f*x)**6 + 6*sec(e + f*x)**5 - 6*sec(e + f*x)**3 + 2*sec(e + f*x)**2 + 2*sec(e + f*x) - 1), x)/(a**3*c**5)`

3.62.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^5} dx$$

$$= \frac{3 \left(\frac{315 \sin(fx+e)}{\cos(fx+e)+1} - \frac{35 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right) - \left(\frac{45 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{189 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{525 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - \frac{1575 \sin(fx+e)^8}{(\cos(fx+e)+1)^8} - 5 \right) (\cos(fx+e)+1)^9}{a^3 c^5 \sin(fx+e)^9} 5760 f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x, algorithm="maxima")`

output `1/5760*(3*(315*sin(f*x + e)/(cos(f*x + e) + 1) - 35*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^3*c^5) - (45*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 189*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 525*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 1575*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 - 5)*(cos(f*x + e) + 1)^9/(a^3*c^5*sin(f*x + e)^9))/f`

3.62.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.18

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^5} dx$$

$$= \frac{1575 \tan(\frac{1}{2} fx + \frac{1}{2} e)^8 - 525 \tan(\frac{1}{2} fx + \frac{1}{2} e)^6 + 189 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 - 45 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 + 5}{a^3 c^5 \tan(\frac{1}{2} fx + \frac{1}{2} e)^9} + \frac{3 (3 a^{12} c^{20} \tan(\frac{1}{2} fx + \frac{1}{2} e)^5 - 35 a^{12} c^{20} \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + 315 a^{12} c^{20} \tan(\frac{1}{2} fx + \frac{1}{2} e))}{a^{15} c^{25}} 5760 f$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^5,x, algorithm="giac")`

output `1/5760*((1575*tan(1/2*f*x + 1/2*e)^8 - 525*tan(1/2*f*x + 1/2*e)^6 + 189*tan(1/2*f*x + 1/2*e)^4 - 45*tan(1/2*f*x + 1/2*e)^2 + 5)/(a^3*c^5*tan(1/2*f*x + 1/2*e)^9) + 3*(3*a^12*c^20*tan(1/2*f*x + 1/2*e)^5 - 35*a^12*c^20*tan(1/2*f*x + 1/2*e)^3 + 315*a^12*c^20*tan(1/2*f*x + 1/2*e))/(a^15*c^25))/f`

3.62.9 Mupad [B] (verification not implemented)

Time = 14.42 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^5} dx$$

$$= \frac{\frac{145 \cos(3e+3fx)}{32} - \frac{169 \cos(2e+2fx)}{32} - \frac{129 \cos(e+fx)}{32} + \frac{55 \cos(4e+4fx)}{16} - \frac{85 \cos(5e+5fx)}{32} + \frac{25 \cos(6e+6fx)}{32} + \frac{5 \cos(7e+7fx)}{32}}{5760 a^3 c^5 f \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^5 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^9}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^5),x)`

output `((145*cos(3*e + 3*f*x))/32 - (169*cos(2*e + 2*f*x))/32 - (129*cos(e + f*x))/32 + (55*cos(4*e + 4*f*x))/16 - (85*cos(5*e + 5*f*x))/32 + (25*cos(6*e + 6*f*x))/32 + (5*cos(7*e + 7*f*x))/32 + 129/16)/(5760*a^3*c^5*f*cos(e/2 + (f*x)/2)^5*sin(e/2 + (f*x)/2)^9)`

3.63 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^6} dx$

3.63.1	Optimal result	506
3.63.2	Mathematica [A] (verified)	506
3.63.3	Rubi [A] (verified)	507
3.63.4	Maple [A] (verified)	508
3.63.5	Fricas [A] (verification not implemented)	509
3.63.6	Sympy [F]	509
3.63.7	Maxima [A] (verification not implemented)	510
3.63.8	Giac [A] (verification not implemented)	510
3.63.9	Mupad [B] (verification not implemented)	511

3.63.1 Optimal result

Integrand size = 32, antiderivative size = 162

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^6} dx$$

$$= -\frac{\cot^9(e+fx)}{9a^3c^6f} - \frac{4 \cot^{11}(e+fx)}{11a^3c^6f} + \frac{\csc(e+fx)}{a^3c^6f} - \frac{8 \csc^3(e+fx)}{3a^3c^6f}$$

$$+ \frac{22 \csc^5(e+fx)}{5a^3c^6f} - \frac{4 \csc^7(e+fx)}{a^3c^6f} + \frac{17 \csc^9(e+fx)}{9a^3c^6f} - \frac{4 \csc^{11}(e+fx)}{11a^3c^6f}$$

output `-1/9*cot(f*x+e)^9/a^3/c^6/f-4/11*cot(f*x+e)^11/a^3/c^6/f+csc(f*x+e)/a^3/c^6/f-8/3*csc(f*x+e)^3/a^3/c^6/f+22/5*csc(f*x+e)^5/a^3/c^6/f-4*csc(f*x+e)^7/a^3/c^6/f+17/9*csc(f*x+e)^9/a^3/c^6/f-4/11*csc(f*x+e)^11/a^3/c^6/f`

3.63.2 Mathematica [A] (verified)

Time = 5.14 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.73

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^6} dx$$

$$= \frac{(-125 - 120 \sec(e+fx) + 680 \sec^2(e+fx) - 400 \sec^3(e+fx) - 720 \sec^4(e+fx) + 832 \sec^5(e+fx) + 495a^3c^6f(-1 + \sec(e+fx))^6(1 + \sec(e+fx))}{495a^3c^6f(-1 + \sec(e+fx))^6(1 + \sec(e+fx))}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^6),x]`

3.63. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^6} dx$

output $((-125 - 120*\text{Sec}[e + f*x] + 680*\text{Sec}[e + f*x]^2 - 400*\text{Sec}[e + f*x]^3 - 720*\text{Sec}[e + f*x]^4 + 832*\text{Sec}[e + f*x]^5 + 64*\text{Sec}[e + f*x]^6 - 384*\text{Sec}[e + f*x]^7 + 128*\text{Sec}[e + f*x]^8)*\text{Tan}[e + f*x])/(495*a^3*c^6*f*(-1 + \text{Sec}[e + f*x])^6*(1 + \text{Sec}[e + f*x])^3)$

3.63.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.90, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4446, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^6} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{(a \csc(e + fx + \frac{\pi}{2}) + a)^3 (c - c \csc(e + fx + \frac{\pi}{2}))^6} dx$$

↓ 4446

$$\frac{\int (a^3 \csc(e + fx) \cot^{11}(e + fx) + 3a^3 \csc^2(e + fx) \cot^{10}(e + fx) + 3a^3 \csc^3(e + fx) \cot^9(e + fx) + a^3 \csc^4(e + fx)) dx}{a^6 c^6}$$

↓ 2009

$$\frac{-\frac{4a^3 \cot^{11}(e+fx)}{11f} - \frac{a^3 \cot^9(e+fx)}{9f} - \frac{4a^3 \csc^{11}(e+fx)}{11f} + \frac{17a^3 \csc^9(e+fx)}{9f} - \frac{4a^3 \csc^7(e+fx)}{f} + \frac{22a^3 \csc^5(e+fx)}{5f} - \frac{8a^3 \csc^3(e+fx)}{3f} + \frac{a^3 \csc(e+fx)}{f}}{a^6 c^6}$$

input $\text{Int}[\text{Sec}[e + f*x]/((a + a*\text{Sec}[e + f*x])^3*(c - c*\text{Sec}[e + f*x])^6), x]$

output $(-1/9*(a^3*\text{Cot}[e + f*x]^9)/f - (4*a^3*\text{Cot}[e + f*x]^11)/(11*f) + (a^3*\text{Csc}[e + f*x])/f - (8*a^3*\text{Csc}[e + f*x]^3)/(3*f) + (22*a^3*\text{Csc}[e + f*x]^5)/(5*f) - (4*a^3*\text{Csc}[e + f*x]^7)/f + (17*a^3*\text{Csc}[e + f*x]^9)/(9*f) - (4*a^3*\text{Csc}[e + f*x]^11)/(11*f))/(a^6*c^6)$

3.63. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^6} dx$

3.63.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4446 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-a)*c]^m Int[ExpandTrig[csc[e + f*x]*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]
```

3.63.4 Maple [A] (verified)

Time = 0.89 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.75

method	result
parallelrisch	$\frac{(240 \cos(7fx+7e) - 1300 \cos(4fx+4e) - 1720 \cos(6fx+6e) + 9680 \cos(fx+e) + 4880 \cos(5fx+5e) - 5584 \cos(3fx+3e) + 8184 \cos(2fx+2e) - 8745 + 125 \cos(8fx+8e)) \sec(1/2fx+1/2e)^5 \csc(1/2fx+1/2e)^{11}}{16220160 f a^3 c^6}$
derivativedivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - \frac{8 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 28 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{8}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} + \frac{56}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{70}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{495}{495 f a^3 c^6}}{256 f a^3 c^6}$
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - \frac{8 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 28 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{8}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9} + \frac{56}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{70}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3} + \frac{495}{495 f a^3 c^6}}{256 f a^3 c^6}$
risch	$\frac{2i(495 e^{15i(fx+e)} - 1485 e^{14i(fx+e)} + 1815 e^{13i(fx+e)} + 2475 e^{12i(fx+e)} - 4917 e^{11i(fx+e)} - 33 e^{10i(fx+e)} + 11715 e^{9i(fx+e)} - 11715 e^{8i(fx+e)} + 495 e^{7i(fx+e)} - 1485 e^{6i(fx+e)} + 1815 e^{5i(fx+e)} - 2475 e^{4i(fx+e)} + 4917 e^{3i(fx+e)} - 33 e^{2i(fx+e)} + 11715 e^{i(fx+e)} - 11715)}{495 f a^3 c^6}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x,method=_RETURNVERBOSE)
```

```
output -1/16220160*(240*cos(7*f*x+7*e)-1300*cos(4*f*x+4*e)-1720*cos(6*f*x+6*e)+9680*cos(f*x+e)+4880*cos(5*f*x+5*e)-5584*cos(3*f*x+3*e)+8184*cos(2*f*x+2*e)-8745+125*cos(8*f*x+8*e))*sec(1/2*f*x+1/2*e)^5*csc(1/2*f*x+1/2*e)^11/f/a^3/c^6
```

3.63.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^6} dx$$

3.63.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.34

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^6} dx$$

$$= \frac{125 \cos(fx+e)^8 + 120 \cos(fx+e)^7 - 680 \cos(fx+e)^6 + 400 \cos(fx+e)^5 + 720 \cos(fx+e)^4 - 832 \cos(fx+e)^3 - 64 \cos(fx+e)^2 + 384 \cos(fx+e) - 128}{495 (a^3 c^6 f \cos(fx+e)^7 - 3 a^3 c^6 f \cos(fx+e)^6 + a^3 c^6 f \cos(fx+e)^5 + 5 a^3 c^6 f \cos(fx+e)^4 - 5 a^3 c^6 f \cos(fx+e)^3 - a^3 c^6 f \cos(fx+e)^2 + 3 a^3 c^6 f \cos(fx+e) - a^3 c^6 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x, algorithm="fracas")`

output `1/495*(125*cos(f*x + e)^8 + 120*cos(f*x + e)^7 - 680*cos(f*x + e)^6 + 400*cos(f*x + e)^5 + 720*cos(f*x + e)^4 - 832*cos(f*x + e)^3 - 64*cos(f*x + e)^2 + 384*cos(f*x + e) - 128)/((a^3*c^6*f*cos(f*x + e)^7 - 3*a^3*c^6*f*cos(f*x + e)^6 + a^3*c^6*f*cos(f*x + e)^5 + 5*a^3*c^6*f*cos(f*x + e)^4 - 5*a^3*c^6*f*cos(f*x + e)^3 - a^3*c^6*f*cos(f*x + e)^2 + 3*a^3*c^6*f*cos(f*x + e) - a^3*c^6*f)*sin(f*x + e))`

3.63.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^6} dx$$

$$= \frac{\int \frac{\sec(e+fx)}{\sec^9(e+fx)-3\sec^8(e+fx)+8\sec^6(e+fx)-6\sec^5(e+fx)-6\sec^4(e+fx)+8\sec^3(e+fx)-3\sec(e+fx)+1} dx}{a^3 c^6}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**6,x)`

output `Integral(sec(e + f*x)/(sec(e + f*x)**9 - 3*sec(e + f*x)**8 + 8*sec(e + f*x)**6 - 6*sec(e + f*x)**5 - 6*sec(e + f*x)**4 + 8*sec(e + f*x)**3 - 3*sec(e + f*x) + 1), x)/(a**3*c**6)`

3.63.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.23

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^6} dx$$

$$= \frac{33 \left(\frac{420 \sin(fx+e)}{\cos(fx+e)+1} - \frac{40 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right) + \frac{\left(\frac{440 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{1980 \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + \frac{5544 \sin(fx+e)^6}{(\cos(fx+e)+1)^6} - \frac{11550 \sin(fx+e)^8}{(\cos(fx+e)+1)^8} + \frac{27720 \sin(fx+e)^{10}}{(\cos(fx+e)+1)^{10}} - \frac{45 \sin(fx+e)^{11}}{(\cos(fx+e)+1)^{11}} \right)}{a^3 c^6 \sin(fx+e)^{11}}}{126720 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x, algorithm="maxima")`

output `1/126720*(33*(420*sin(f*x + e)/(cos(f*x + e) + 1) - 40*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^3*c^6) + (440*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 1980*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 5544*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 11550*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 27720*sin(f*x + e)^10/(cos(f*x + e) + 1)^10 - 45*(cos(f*x + e) + 1)^11/(a^3*c^6*sin(f*x + e)^11))/f`

3.63.8 Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.96

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^6} dx$$

$$= \frac{27720 \tan(\frac{1}{2} fx + \frac{1}{2} e)^{10} - 11550 \tan(\frac{1}{2} fx + \frac{1}{2} e)^8 + 5544 \tan(\frac{1}{2} fx + \frac{1}{2} e)^6 - 1980 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 + 440 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 45}{a^3 c^6 \tan(\frac{1}{2} fx + \frac{1}{2} e)^{11}} + \frac{33 (3 a^{12} c^{24} \tan(\frac{1}{2} fx + \frac{1}{2} e)^5 - 40 a^{12} c^{24} \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + 420 a^{12} c^{24} \tan(\frac{1}{2} fx + \frac{1}{2} e))}{126720 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^6,x, algorithm="giac")`

output `1/126720*((27720*tan(1/2*f*x + 1/2*e)^10 - 11550*tan(1/2*f*x + 1/2*e)^8 + 5544*tan(1/2*f*x + 1/2*e)^6 - 1980*tan(1/2*f*x + 1/2*e)^4 + 440*tan(1/2*f*x + 1/2*e)^2 - 45)/(a^3*c^6*tan(1/2*f*x + 1/2*e)^11) + 33*(3*a^12*c^24*tan(1/2*f*x + 1/2*e)^5 - 40*a^12*c^24*tan(1/2*f*x + 1/2*e)^3 + 420*a^12*c^24*tan(1/2*f*x + 1/2*e))/(a^15*c^30))/f`

3.63.9 Mupad [B] (verification not implemented)

Time = 14.68 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^6} dx =$$

$$-\frac{\frac{605 \cos(e+fx)}{8} + \frac{1023 \cos(2e+2fx)}{16} - \frac{349 \cos(3e+3fx)}{8} - \frac{325 \cos(4e+4fx)}{32} + \frac{305 \cos(5e+5fx)}{8} - \frac{215 \cos(6e+6fx)}{16} + \frac{15 \cos(7e+7fx)}{8}}{126720 a^3 c^6 f \cos\left(\frac{e}{2} + \frac{fx}{2}\right)^5 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^6),x)`output `-((605*cos(e + f*x))/8 + (1023*cos(2*e + 2*f*x))/16 - (349*cos(3*e + 3*f*x))/8 - (325*cos(4*e + 4*f*x))/32 + (305*cos(5*e + 5*f*x))/8 - (215*cos(6*e + 6*f*x))/16 + (15*cos(7*e + 7*f*x))/8 + (125*cos(8*e + 8*f*x))/128 - 8745/128)/(126720*a^3*c^6*f*cos(e/2 + (f*x)/2)^5*sin(e/2 + (f*x)/2)^11)`

3.64 $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx$

3.64.1	Optimal result	512
3.64.2	Mathematica [A] (verified)	513
3.64.3	Rubi [A] (verified)	513
3.64.4	Maple [A] (verified)	515
3.64.5	Fricas [A] (verification not implemented)	516
3.64.6	Sympy [F(-1)]	516
3.64.7	Maxima [F]	517
3.64.8	Giac [A] (verification not implemented)	517
3.64.9	Mupad [B] (verification not implemented)	518

3.64.1 Optimal result

Integrand size = 32, antiderivative size = 163

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx =$$

$$\frac{256c^4(a + a \sec(e + fx)) \tan(e + fx)}{315f \sqrt{c - c \sec(e + fx)}} - \frac{64c^3(a + a \sec(e + fx)) \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{105f} - \frac{8c^2(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} \tan(e + fx)}{21f} - \frac{2c(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} \tan(e + fx)}{9f}$$

output

```
-8/21*c^2*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f-2/9*c*(a+a*
sec(f*x+e))*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f-256/315*c^4*(a+a*sec(f*x+e
))*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-64/105*c^3*(a+a*sec(f*x+e))*(c-c*se
c(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.64.2 Mathematica [A] (verified)

Time = 2.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.43

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx = \frac{2ac^4(1 + \sec(e + fx))(-319 + 321 \sec(e + fx) - 165 \sec^2(e + fx) + 35 \sec^3(e + fx)) \tan(e + fx)}{315f \sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(7/2),x]`

output `(2*a*c^4*(1 + Sec[e + f*x])*(-319 + 321*Sec[e + f*x] - 165*Sec[e + f*x]^2 + 35*Sec[e + f*x]^3)*Tan[e + f*x])/(315*f*Sqrt[c - c*Sec[e + f*x]])`

3.64.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4443, 3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{7/2} dx \\ & \quad \downarrow \text{4443} \\ & \frac{4}{3}c \int \sec(e + fx)(\sec(e + fx)a + a)(c - c \sec(e + fx))^{5/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^{5/2}}{9f} \\ & \quad \downarrow \text{3042} \\ & \frac{4}{3}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^{5/2}}{9f} \end{aligned}$$

3.64. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx$

↓ 4443

$$\frac{4}{3}c \left(\frac{8}{7}c \int \sec(e+fx)(\sec(e+fx)a+a)(c-c\sec(e+fx))^{3/2} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}{7f} \right) - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}{9f}$$

↓ 3042

$$\frac{4}{3}c \left(\frac{8}{7}c \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a \right) \left(c-c\csc\left(e+fx+\frac{\pi}{2}\right) \right)^{3/2} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}{7f} \right) - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}{9f}$$

↓ 4443

$$\frac{4}{3}c \left(\frac{8}{7}c \left(\frac{4}{5}c \int \sec(e+fx)(\sec(e+fx)a+a)\sqrt{c-c\sec(e+fx)} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}{5f} \right) - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}{9f} \right)$$

↓ 3042

$$\frac{4}{3}c \left(\frac{8}{7}c \left(\frac{4}{5}c \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a \right) \sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}{5f} \right) - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}{9f} \right)$$

↓ 4441

$$\frac{4}{3}c \left(\frac{8}{7}c \left(-\frac{8c^2 \tan(e+fx)(a \sec(e+fx)+a)}{15f\sqrt{c-c\sec(e+fx)}} - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}{5f} \right) - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}{9f} \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(7/2),x]`

```
output (-2*c*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(9*f)
+ (4*c*((-2*c*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])
)/(7*f) + (8*c*((-8*c^2*(a + a*Sec[e + f*x])*Tan[e + f*x])/(15*f*Sqrt[c -
c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x])*Tan[
e + f*x])/(5*f)))/7)/3
```

3.64.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*Sq
rt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f
*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /
; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
&& NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.64.4 Maple [A] (verified)

Time = 7.67 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.47

method	result
default	$\frac{2a c^3 (319 \cos(fx+e)^3 - 321 \cos(fx+e)^2 + 165 \cos(fx+e) - 35) \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1)^2 \sec(fx+e)^4 \csc(fx+e)}{315 f}$
parts	$-\frac{2a(\sec(fx+e)-1)^3 (177 \cos(fx+e)^3 - 71 \cos(fx+e)^2 + 27 \cos(fx+e) - 5) c^3 \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1) \csc(fx+e)}{35 f (\cos(fx+e)-1)^3} + \dots$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(7/2), x, method=_RETURNVER
BOSE)
```

$$3.64. \int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx$$

output $2/315*a*c^3/f*(319*\cos(f*x+e)^3-321*\cos(f*x+e)^2+165*\cos(f*x+e)-35)*(-c*(\sec(f*x+e)-1))^{1/2}*(\cos(f*x+e)+1)^2*\sec(f*x+e)^4*\csc(f*x+e)$

3.64.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.73

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx = \frac{2(319ac^3 \cos(fx + e)^5 + 317ac^3 \cos(fx + e)^4 - 158ac^3 \cos(fx + e)^3 - 26ac^3 \cos(fx + e)^2 + 95ac^3 \cos(fx + e) - 35a^2c^3) \sqrt{(c \cos(fx + e) - c)/\cos(fx + e)}}{315 f \cos(fx + e)^4 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(7/2),x, algorithm="fricas")`

output $2/315*(319*a*c^3*\cos(f*x + e)^5 + 317*a*c^3*\cos(f*x + e)^4 - 158*a*c^3*\cos(f*x + e)^3 - 26*a*c^3*\cos(f*x + e)^2 + 95*a*c^3*\cos(f*x + e) - 35*a*c^3)*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^4*\sin(f*x + e))$

3.64.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))**(7/2),x)`

output Timed out

3.64.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx = \int (a \sec(fx + e) + a)(-c \sec(fx + e) + c)^{7/2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(7/2),x, algorithm="maxima")`

output `-2/315*(315*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/4)*(5*(a*c^3*f*cos(2*f*x + 2*e)^4 + a*c^3*f*sin(2*f*x + 2*e)^4 + 4*a*c^3*f*cos(2*f*x + 2*e)^3 + 6*a*c^3*f*cos(2*f*x + 2*e)^2 + 4*a*c^3*f*cos(2*f*x + 2*e) + a*c^3*f + 2*(a*c^3*f*cos(2*f*x + 2*e)^2 + 2*a*c^3*f*cos(2*f*x + 2*e) + a*c^3*f)*sin(2*f*x + 2*e)^2)*integrate((cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/4)*(((cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 2*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 2*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)))/((cos(2*f*x + 2*e)^6 + sin(2*f*x + 2*e)^6 ...`

3.64.8 Giac [A] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.66

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx = \frac{32\sqrt{2}\left(105\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^3 c^2 + 189\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^2 c^3 + 135\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)c^4\right)}{315\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{9}{2}}f}$$

3.64. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(7/2),x, algorithm="giac")`

output `32/315*sqrt(2)*(105*(c*tan(1/2*f*x + 1/2*e)^2 - c)^3*c^2 + 189*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^3 + 135*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^4 + 35*c^5)*a*c^3/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(9/2)*f)`

3.64.9 Mupad [B] (verification not implemented)

Time = 20.55 (sec) , antiderivative size = 483, normalized size of antiderivative = 2.96

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{7/2} dx = \frac{\sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li}-fx \operatorname{li}}}{2} + \frac{e^{e \operatorname{li}+fx \operatorname{li}}}{2}}} \left(\frac{ac^3 2i}{f} + \frac{ac^3 e^{e \operatorname{li}+fx \operatorname{li}} 638i}{315 f} \right)}{e^{e \operatorname{li}+fx \operatorname{li}} - 1} - \frac{\sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li}-fx \operatorname{li}}}{2} + \frac{e^{e \operatorname{li}+fx \operatorname{li}}}{2}}} \left(\frac{ac^3 32i}{9 f} + \frac{ac^3 e^{e \operatorname{li}+fx \operatorname{li}} 32i}{9 f} \right)}{(e^{e \operatorname{li}+fx \operatorname{li}} - 1) (e^{2e \operatorname{li}+2fx \operatorname{li}} + 1)^4} + \frac{\sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li}-fx \operatorname{li}}}{2} + \frac{e^{e \operatorname{li}+fx \operatorname{li}}}{2}}} \left(\frac{ac^3 96i}{7 f} + \frac{ac^3 e^{e \operatorname{li}+fx \operatorname{li}} 32i}{63 f} \right)}{(e^{e \operatorname{li}+fx \operatorname{li}} - 1) (e^{2e \operatorname{li}+2fx \operatorname{li}} + 1)^3} - \frac{\sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li}-fx \operatorname{li}}}{2} + \frac{e^{e \operatorname{li}+fx \operatorname{li}}}{2}}} \left(\frac{ac^3 64i}{5 f} - \frac{ac^3 e^{e \operatorname{li}+fx \operatorname{li}} 736i}{105 f} \right)}{(e^{e \operatorname{li}+fx \operatorname{li}} - 1) (e^{2e \operatorname{li}+2fx \operatorname{li}} + 1)^2} + \frac{\sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li}-fx \operatorname{li}}}{2} + \frac{e^{e \operatorname{li}+fx \operatorname{li}}}{2}}} \left(\frac{ac^3 8i}{3 f} - \frac{ac^3 e^{e \operatorname{li}+fx \operatorname{li}} 1256i}{315 f} \right)}{(e^{e \operatorname{li}+fx \operatorname{li}} - 1) (e^{2e \operatorname{li}+2fx \operatorname{li}} + 1)}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(7/2))/cos(e + f*x),x)`

output

$$\begin{aligned} & \left((c - c/\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2) \right)^{1/2} * \left((a*c^3*2i) \right. \\ & /f + (a*c^3*\exp(e*1i + f*x*1i)*638i)/(315*f) \left. \right) / (\exp(e*1i + f*x*1i) - 1) - \\ & \left((c - c/\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2) \right)^{1/2} * \left((a*c^3*32i) \right. \\ &) / (9*f) + (a*c^3*\exp(e*1i + f*x*1i)*32i)/(9*f) \left. \right) / ((\exp(e*1i + f*x*1i) - 1) \\ & * (\exp(e*2i + f*x*2i) + 1)^4) + \left((c - c/\exp(-e*1i - f*x*1i)/2 + \exp(e*1i \right. \\ & + f*x*1i)/2) \left. \right)^{1/2} * \left((a*c^3*96i)/(7*f) + (a*c^3*\exp(e*1i + f*x*1i)*32i)/(6 \right. \\ & 3*f) \left. \right) / ((\exp(e*1i + f*x*1i) - 1) * (\exp(e*2i + f*x*2i) + 1)^3) - \left((c - c/\exp(- \right. \\ & e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2) \left. \right)^{1/2} * \left((a*c^3*64i)/(5*f) - (\right. \\ & a*c^3*\exp(e*1i + f*x*1i)*736i)/(105*f) \left. \right) / ((\exp(e*1i + f*x*1i) - 1) * (\exp(e* \\ & 2i + f*x*2i) + 1)^2) + \left((c - c/\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i \right. \\ &)/2) \left. \right)^{1/2} * \left((a*c^3*8i)/(3*f) - (a*c^3*\exp(e*1i + f*x*1i)*1256i)/(315*f) \right) \\ & / ((\exp(e*1i + f*x*1i) - 1) * (\exp(e*2i + f*x*2i) + 1)) \end{aligned}$$

3.65 $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx$

3.65.1	Optimal result	520
3.65.2	Mathematica [A] (verified)	521
3.65.3	Rubi [A] (verified)	521
3.65.4	Maple [A] (verified)	523
3.65.5	Fricas [A] (verification not implemented)	523
3.65.6	Sympy [F]	524
3.65.7	Maxima [F]	524
3.65.8	Giac [A] (verification not implemented)	525
3.65.9	Mupad [B] (verification not implemented)	526

3.65.1 Optimal result

Integrand size = 32, antiderivative size = 122

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx =$$

$$-\frac{64c^3(a + a \sec(e + fx)) \tan(e + fx)}{105f \sqrt{c - c \sec(e + fx)}}$$

$$-\frac{16c^2(a + a \sec(e + fx)) \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{35f}$$

$$-\frac{2c(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} \tan(e + fx)}{7f}$$

output

```
-2/7*c*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f-64/105*c^3*(a+
a*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-16/35*c^2*(a+a*sec(f*x+e)
))*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.65.2 Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.49

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx =$$

$$\frac{2ac^3(1 + \sec(e + fx))(71 - 54 \sec(e + fx) + 15 \sec^2(e + fx)) \tan(e + fx)}{105f \sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2),x]`

output `(-2*a*c^3*(1 + Sec[e + f*x])*(71 - 54*Sec[e + f*x] + 15*Sec[e + f*x]^2)*Tan[e + f*x])/(105*f*Sqrt[c - c*Sec[e + f*x]])`

3.65.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^{5/2} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx$$

$$\downarrow \text{4443}$$

$$\frac{8}{7}c \int \sec(e + fx)(\sec(e + fx)a + a)(c - c \sec(e + fx))^{3/2} dx -$$

$$\frac{2c \tan(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^{3/2}}{7f}$$

$$\downarrow \text{3042}$$

$$\frac{8}{7}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right) a + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx -$$

$$\frac{2c \tan(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^{3/2}}{7f}$$

3.65. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx$

↓ 4443

$$\frac{8}{7}c \left(\frac{4}{5}c \int \sec(e+fx)(\sec(e+fx)a+a)\sqrt{c-c\sec(e+fx)}dx - \frac{2c\tan(e+fx)(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}{5f} \right. \\ \left. - \frac{2c\tan(e+fx)(a\sec(e+fx)+a)(c-c\sec(e+fx))^{3/2}}{7f} \right)$$

↓ 3042

$$\frac{8}{7}c \left(\frac{4}{5}c \int \csc\left(e+fx+\frac{\pi}{2}\right)\left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)\sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)}dx - \frac{2c\tan(e+fx)(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}{5f} \right. \\ \left. - \frac{2c\tan(e+fx)(a\sec(e+fx)+a)(c-c\sec(e+fx))^{3/2}}{7f} \right)$$

↓ 4441

$$\frac{8}{7}c \left(-\frac{8c^2 \tan(e+fx)(a\sec(e+fx)+a)}{15f\sqrt{c-c\sec(e+fx)}} - \frac{2c\tan(e+fx)(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}{5f} \right) - \\ \frac{2c\tan(e+fx)(a\sec(e+fx)+a)(c-c\sec(e+fx))^{3/2}}{7f}$$

```
input Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2),x]
```

```
output (-2*c*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(7*f) + (8*c*((-8*c^2*(a + a*Sec[e + f*x])*Tan[e + f*x])/(15*f*Sqrt[c - c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x])*Tan[e + f*x])/(5*f)))/7
```

3.65.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]
```

3.65. $\int \sec(e+fx)(a+a\sec(e+fx))(c-c\sec(e+fx))^{5/2} dx$

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.65.4 Maple [A] (verified)

Time = 7.28 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.55

method	result
default	$\frac{2a c^2 (71 \cos(fx+e)^2 - 54 \cos(fx+e) + 15) \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1)^2 \sec(fx+e)^3 \csc(fx+e)}{105f}$
parts	$\frac{2a(\sec(fx+e)-1)^2 (43 \cos(fx+e)^2 - 14 \cos(fx+e) + 3) c^2 \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1) \csc(fx+e)}{15f(\cos(fx+e)-1)^2} - \frac{2a(46 \cos(fx+e)^3 - 23 \cos(fx+e) + 1) \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1) \csc(fx+e)}{105f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(5/2), x, method=_RETURNVER
BOSE)
```

```
output 2/105*a*c^2/f*(71*cos(f*x+e)^2-54*cos(f*x+e)+15)*(-c*(sec(f*x+e)-1))^(1/2)
*(cos(f*x+e)+1)^2*sec(f*x+e)^3*csc(f*x+e)
```

3.65.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.86

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx = \frac{2(71ac^2 \cos(fx + e)^4 + 88ac^2 \cos(fx + e)^3 - 22ac^2 \cos(fx + e)^2 - 24ac^2 \cos(fx + e) + 15) \sqrt{-c(\sec(fx + e) - 1)} (\cos(fx + e) + 1) \csc(fx + e)}{105f \cos(fx + e)^3 \sin(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(5/2), x, algorithm=
"fracas")
```

output $2/105*(71*a*c^2*\cos(f*x + e)^4 + 88*a*c^2*\cos(f*x + e)^3 - 22*a*c^2*\cos(f*x + e)^2 - 24*a*c^2*\cos(f*x + e) + 15*a*c^2)*\text{sqrt}((c*\cos(f*x + e) - c)/\cos(f*x + e))/(f*\cos(f*x + e)^3*\sin(f*x + e))$

3.65.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx = a \left(\int c^2 \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx + \int \left(-c^2 \sqrt{-c \sec(e + fx) + c} \sec^2(e + fx) \right) dx + \int \left(-c^2 \sqrt{-c \sec(e + fx) + c} \sec^3(e + fx) \right) dx + \int c^2 \sqrt{-c \sec(e + fx) + c} \sec^4(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))**(5/2),x)`

output `a*(Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(-c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2, x) + Integral(-c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3, x) + Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4, x))`

3.65.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx = \int (a \sec(fx + e) + a)(-c \sec(fx + e) + c)^{5/2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(5/2),x, algorithm="maxima")`

```

output -2/105*(105*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e)
+ 1)^(3/4)*(3*(a*c^2*f*cos(2*f*x + 2*e)^2 + a*c^2*f*sin(2*f*x + 2*e)^2 + 2
*a*c^2*f*cos(2*f*x + 2*e) + a*c^2*f)*integrate((((cos(6*f*x + 6*e)*cos(2*f
*x + 2*e) + 2*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin
(6*f*x + 6*e)*sin(2*f*x + 2*e) + 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin
(2*f*x + 2*e)^2)*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + (c
os(2*f*x + 2*e)*sin(6*f*x + 6*e) + 2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - c
os(6*f*x + 6*e)*sin(2*f*x + 2*e) - 2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*si
n(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(5/2*arctan2(sin(2*
f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x + 2*e)*sin(6*f*x + 6*e) +
2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(6*f*x + 6*e)*sin(2*f*x + 2*e) -
2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(7/2*arctan2(sin(2*f*x + 2*e), co
s(2*f*x + 2*e))) - (cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 2*cos(4*f*x + 4*e)
*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(6*f*x + 6*e)*sin(2*f*x + 2*e)
+ 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*sin(7/2*arcta
n2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sin(5/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e) + 1)))/(((cos(2*f*x + 2*e)^4 + sin(2*f*x + 2*e)^4 + (cos
(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)*cos(6*f*x +
6*e)^2 + 4*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e)
+ 1)*cos(4*f*x + 4*e)^2 + 2*cos(2*f*x + 2*e)^3 + (cos(2*f*x + 2*e)^2 + ...

```

3.65.8 Giac [A] (verification not implemented)

Time = 0.83 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.68

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx = \frac{16\sqrt{2} \left(35 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right)^2 c^2 + 42 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right) c^3 + 15 c^4 \right) a c^2}{105 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right)^{7/2} f}$$

```

input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(5/2),x, algorithm=
"giac")

```

```

output 16/105*sqrt(2)*(35*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^2 + 42*(c*tan(1/2*f*
x + 1/2*e)^2 - c)*c^3 + 15*c^4)*a*c^2/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(7/2
)*f)

```

3.65. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx$

3.65.9 Mupad [B] (verification not implemented)

Time = 17.24 (sec) , antiderivative size = 384, normalized size of antiderivative = 3.15

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2} dx = \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{\frac{a c^2 2i}{f} + \frac{a c^2 e^{e \cdot 1i + f \cdot x \cdot 1i} 142i}{105 f}}}{e^{e \cdot 1i + f \cdot x \cdot 1i} - 1} + \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{\left(\frac{a c^2 16i}{7 f} - \frac{a c^2 e^{e \cdot 1i + f \cdot x \cdot 1i} 16i}{7 f}\right)}{\left(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1\right) \left(e^{e \cdot 2i + f \cdot x \cdot 2i} + 1\right)^3} - \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{\left(\frac{a c^2 8i}{5 f} - \frac{a c^2 e^{e \cdot 1i + f \cdot x \cdot 1i} 184i}{35 f}\right)}{\left(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1\right) \left(e^{e \cdot 2i + f \cdot x \cdot 2i} + 1\right)^2} - \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{\left(\frac{a c^2 4i}{3 f} + \frac{a c^2 e^{e \cdot 1i + f \cdot x \cdot 1i} 244i}{105 f}\right)}{\left(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1\right) \left(e^{e \cdot 2i + f \cdot x \cdot 2i} + 1\right)}$$

```
input int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x),x)
```

```
output ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a*c^2*2i)/f + (a*c^2*exp(e*1i + f*x*1i)*142i)/(105*f)))/(exp(e*1i + f*x*1i) - 1) + ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a*c^2*16i)/(7*f) - (a*c^2*exp(e*1i + f*x*1i)*16i)/(7*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^3) - ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a*c^2*8i)/(5*f) - (a*c^2*exp(e*1i + f*x*1i)*184i)/(35*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^2) - ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a*c^2*4i)/(3*f) + (a*c^2*exp(e*1i + f*x*1i)*244i)/(105*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1))
```

3.66 $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx$

3.66.1	Optimal result	527
3.66.2	Mathematica [A] (verified)	527
3.66.3	Rubi [A] (verified)	528
3.66.4	Maple [A] (verified)	529
3.66.5	Fricas [A] (verification not implemented)	530
3.66.6	Sympy [F]	530
3.66.7	Maxima [F]	531
3.66.8	Giac [A] (verification not implemented)	531
3.66.9	Mupad [B] (verification not implemented)	532

3.66.1 Optimal result

Integrand size = 32, antiderivative size = 81

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx =$$

$$-\frac{8c^2(a + a \sec(e + fx)) \tan(e + fx)}{15f \sqrt{c - c \sec(e + fx)}}$$

$$-\frac{2c(a + a \sec(e + fx)) \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{5f}$$

output

```
-8/15*c^2*(a+a*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-2/5*c*(a+a*
sec(f*x+e))*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.66.2 Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.62

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx =$$

$$\frac{2ac^2(1 + \sec(e + fx))(-7 + 3 \sec(e + fx)) \tan(e + fx)}{15f \sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2),x]`

output `(2*a*c^2*(1 + Sec[e + f*x])*(-7 + 3*Sec[e + f*x])*Tan[e + f*x])/(15*f*Sqrt[c - c*Sec[e + f*x]])`

3.66.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{4}{5}c \int \sec(e + fx)(\sec(e + fx)a + a)\sqrt{c - c \sec(e + fx)} dx - \\
 & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)\sqrt{c - c \sec(e + fx)}}{5f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4}{5}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right) a + a\right) \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx - \\
 & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)\sqrt{c - c \sec(e + fx)}}{5f} \\
 & \quad \downarrow \text{4441} \\
 & -\frac{8c^2 \tan(e + fx)(a \sec(e + fx) + a)}{15f \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)\sqrt{c - c \sec(e + fx)}}{5f}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2),x]`

output $(-8*c^2*(a + a*\text{Sec}[e + f*x])*Tan[e + f*x])/(15*f*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]) - (2*c*(a + a*\text{Sec}[e + f*x])*Sqrt[c - c*\text{Sec}[e + f*x]]*Tan[e + f*x])/(5*f)$

3.66.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.66.4 Maple [A] (verified)

Time = 4.59 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.68

method	result
default	$\frac{2ac(7 \cos(fx+e)-3)\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)^2 \sec(fx+e)^2 \csc(fx+e)}{15f}$
parts	$-\frac{2a(\sec(fx+e)-1)(5 \cos(fx+e)-1)c\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1) \csc(fx+e)}{3f(\cos(fx+e)-1)} + \frac{2a(6 \cos(fx+e)^2 - 3 \cos(fx+e) + 1)(\sec(fx+e))}{3f(\cos(fx+e)-1)}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output $2/15*a*c/f*(7*\cos(f*x+e)-3)*(-c*(\sec(f*x+e)-1))^(1/2)*(c - c*\sec(f*x+e))^2*\sec(f*x+e)^2*\csc(f*x+e)$

3.66. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx$

3.66.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.01

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx = \frac{2(7ac \cos(fx + e)^3 + 11ac \cos(fx + e)^2 + ac \cos(fx + e) - 3ac) \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}}}{15 f \cos(fx + e)^2 \sin(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(3/2),x, algorithm="fricas")
```

```
output 2/15*(7*a*c*cos(f*x + e)^3 + 11*a*c*cos(f*x + e)^2 + a*c*cos(f*x + e) - 3*a*c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)^2*sin(f*x + e))
```

3.66.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx = a \left(\int c \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx + \int (-c \sqrt{-c \sec(e + fx) + c} \sec^3(e + fx)) dx \right)$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))**(3/2),x)
```

```
output a*(Integral(c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3, x))
```

3.66.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx = \int (a \sec(fx + e) + a)(-c \sec(fx + e) + c)^{3/2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `-2/15*(15*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/4)*((a*c*f*cos(2*f*x + 2*e)^2 + a*c*f*sin(2*f*x + 2*e)^2 + 2*a*c*f*cos(2*f*x + 2*e) + a*c*f)*integrate((cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/4)*(((cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 2*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - (cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 2*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)))/((cos(2*f*x + 2*e)^4 + sin(2*f*x + 2*e)^4 + (cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)*cos(6*f*x + 6*e)^2 + 4*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)*cos(4*...`

3.66.8 Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.69

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx = \frac{8\sqrt{2}\left(5\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)c^3 + 3c^4\right)a}{15\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{5}{2}}f}$$

3.66. $\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `8/15*sqrt(2)*(5*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^3 + 3*c^4)*a/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2)*f)`

3.66.9 Mupad [B] (verification not implemented)

Time = 17.06 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.48

$$\int \sec(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2} dx = \frac{2ac(e^{e+fx} \operatorname{li} + 1)^3 \sqrt{c - \frac{e^{-e+fx} - 1}{2} + \frac{e^{e+fx} - 1}{2}} (7 + 7e^{2e+2fx} - 6e^{e+fx})}{15f(e^{e+fx} - 1)(e^{2e+2fx} + 1)^2}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(3/2))/cos(e + f*x),x)`

output `-(2*a*c*(exp(e*1i + f*x*1i)*1i + 1i)^3*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*(7*exp(e*2i + f*x*2i) - 6*exp(e*1i + f*x*1i) + 7))/(15*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^2)`

3.67 $\int \sec(e+fx)(a+a \sec(e+fx))\sqrt{c - c \sec(e + fx)} dx$

3.67.1	Optimal result	533
3.67.2	Mathematica [A] (verified)	533
3.67.3	Rubi [A] (verified)	534
3.67.4	Maple [A] (verified)	535
3.67.5	Fricas [A] (verification not implemented)	535
3.67.6	Sympy [F]	535
3.67.7	Maxima [F]	536
3.67.8	Giac [A] (verification not implemented)	537
3.67.9	Mupad [B] (verification not implemented)	537

3.67.1 Optimal result

Integrand size = 32, antiderivative size = 39

$$\int \sec(e+fx)(a+a \sec(e+fx))\sqrt{c - c \sec(e + fx)} dx = -\frac{2c(a + a \sec(e + fx)) \tan(e + fx)}{3f\sqrt{c - c \sec(e + fx)}}$$

output `-2/3*c*(a+a*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)`

3.67.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.97

$$\int \sec(e+fx)(a+a \sec(e+fx))\sqrt{c - c \sec(e + fx)} dx = -\frac{2ac(1 + \sec(e + fx)) \tan(e + fx)}{3f\sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]],x]`

output `(-2*a*c*(1 + Sec[e + f*x])*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e + f*x]])`

3.67.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)\sqrt{c - c \sec(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4441}$$

$$-\frac{2c \tan(e + fx)(a \sec(e + fx) + a)}{3f \sqrt{c - c \sec(e + fx)}}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]],x]`

output `(-2*c*(a + a*Sec[e + f*x])*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e + f*x]])`

3.67.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] :> Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.67.4 Maple [A] (verified)

Time = 3.96 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

method	result	size
default	$\frac{2a(\cos(fx+e)+1)^2 \sqrt{-c(\sec(fx+e)-1)} \sec(fx+e) \csc(fx+e)}{3f}$	42
parts	$-\frac{2a\sqrt{-c(\sec(fx+e)-1)} \sin(fx+e)}{f(\cos(fx+e)-1)} - \frac{2a\sqrt{-c(\sec(fx+e)-1)} (-\sec(fx+e) \csc(fx+e) + 2 \cot(fx+e) + \csc(fx+e))}{3f}$	85

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(1/2),x,method=_RETURNVER
BOSE)
```

```
output 2/3*a/f*(cos(f*x+e)+1)^2*(-c*(sec(f*x+e)-1))^(1/2)*sec(f*x+e)*csc(f*x+e)
```

3.67.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.67

$$\int \sec(e + fx)(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{2(a \cos(fx + e)^2 + 2a \cos(fx + e) + a) \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}}}{3f \cos(fx + e) \sin(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(1/2),x, algorithm=
"fricas")
```

```
output 2/3*(a*cos(f*x + e)^2 + 2*a*cos(f*x + e) + a)*sqrt((c*cos(f*x + e) - c)/co
s(f*x + e))/(f*cos(f*x + e)*sin(f*x + e))
```

3.67.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)} dx$$

$$= a \left(\int \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx + \int \sqrt{-c \sec(e + fx) + c} \sec^2(e + fx) dx \right)$$

3.67. $\int \sec(e + fx)(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))**(1/2),x)`

output `a*(Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2, x))`

3.67.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)} dx$$

$$= \int (a \sec(fx + e) + a)\sqrt{-c \sec(fx + e) + c \sec(fx + e)} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `2/3*(3*(a*f*integrate((((cos(6*f*x + 6*e))*cos(2*f*x + 2*e) + 2*cos(4*f*x + 4*e))*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 2*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)))/(((2*(2*cos(4*f*x + 4*e) + cos(2*f*x + 2*e))*cos(6*f*x + 6*e) + cos(6*f*x + 6*e)^2 + 4*cos(4*f*x + 4*e)^2 + 4*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + 2*(2*sin(4*f*x + 4*e) + sin(2*f*x + 2*e))*sin(6*f*x + 6*e) + sin(6*f*x + 6*e)^2 + 4*sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1))^2 + (2*(2*cos(4*f*x + 4*e) + cos(2*f*x + 2*e))*cos(6*f...`

3.67.8 Giac [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int \sec(e + fx)(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)} dx = \frac{4\sqrt{2}ac^2}{3 \left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c \right)^{\frac{3}{2}} f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `4/3*sqrt(2)*a*c^2/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*f)`

3.67.9 Mupad [B] (verification not implemented)

Time = 14.44 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.23

$$\int \sec(e + fx)(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{2a \sqrt{c - \frac{c}{\cos(e+fx)}} (2 \sin(2e + 2fx) - \sin(4e + 4fx))}{3f (8 \cos(2e + 2fx) - 12 \cos(e + fx) - 4 \cos(3e + 3fx) + \cos(4e + 4fx) + 7)}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x),x)`

output `(2*a*(c - c/cos(e + f*x))^(1/2)*(2*sin(2*e + 2*f*x) - sin(4*e + 4*f*x)))/(3*f*(8*cos(2*e + 2*f*x) - 12*cos(e + f*x) - 4*cos(3*e + 3*f*x) + cos(4*e + 4*f*x) + 7))`

3.68
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{\sqrt{c-c \sec(e+fx)}} dx$$

3.68.1	Optimal result	538
3.68.2	Mathematica [A] (verified)	538
3.68.3	Rubi [A] (verified)	539
3.68.4	Maple [A] (verified)	540
3.68.5	Fricas [A] (verification not implemented)	541
3.68.6	Sympy [F]	541
3.68.7	Maxima [F]	542
3.68.8	Giac [A] (verification not implemented)	542
3.68.9	Mupad [F(-1)]	543

3.68.1 Optimal result

Integrand size = 32, antiderivative size = 77

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{\sqrt{c-c \sec(e+fx)}} dx = -\frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{\sqrt{cf}} + \frac{2a \tan(e+fx)}{f\sqrt{c-c \sec(e+fx)}}$$

output `-2*a*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))*2^(1/2)/f/c^(1/2)+2*a*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)`

3.68.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.01

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{\sqrt{c-c \sec(e+fx)}} dx = \frac{2a\left(-\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1+\sec(e+fx)}}{\sqrt{2}}\right) + \sqrt{1+\sec(e+fx)}\right) \tan(e+fx)}{f\sqrt{1+\sec(e+fx)}\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/Sqrt[c - c*Sec[e + f*x]],x]`

output `(2*a*(-(Sqrt[2]*ArcTanh[Sqrt[1 + Sec[e + f*x]]/Sqrt[2]]) + Sqrt[1 + Sec[e + f*x]])*Tan[e + f*x])/(f*Sqrt[1 + Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.68.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{\sqrt{c-c \sec(e+fx)}} dx$$

3.68.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4444, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{\sqrt{c-c\sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4444} \\
 & 2a \int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & 2a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{4282} \\
 & \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{4a \int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c\sec(e+fx)} + 2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{f} \\
 & \quad \downarrow \text{216} \\
 & \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{c}\tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{c}f}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/Sqrt[c - c*Sec[e + f*x]],x]`

output `(-2*Sqrt[2]*a*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])])/(Sqrt[c]*f) + (2*a*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.68. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{\sqrt{c-c\sec(e+fx)}} dx$

3.68.3.1 Defintions of rubi rules used

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4282 Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] :> Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x])]], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]
```

```
rule 4444 Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.))/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] :> Simp[-2*d*Cot[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/(f*(2*n - 1)*Sqrt[a + b*Csc[e + f*x]]), x] + Simp[2*c*((2*n - 1)/(2*n - 1)) Int[Csc[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/Sqrt[a + b*Csc[e + f*x])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]
```

3.68.4 Maple [A] (verified)

Time = 3.64 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.04

method	result
default	$\frac{a\sqrt{2} \left(-2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) + \sqrt{2} \right) \tan(fx+e)}{f\sqrt{-c(\sec(fx+e)-1)}}$
parts	$\frac{a\sqrt{2} \sin(fx+e) \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right)}{f(\cos(fx+e)+1)\sqrt{-c(\sec(fx+e)-1)}\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} + \frac{a\sqrt{2} \left(-\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) + \sqrt{2} \right) \tan(fx+e)}{f\sqrt{-c(\sec(fx+e)-1)}}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2), x, method=_RETURNVERBOSE)
```

3.68. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{\sqrt{c-c\sec(e+fx)}} dx$

output $a/f*2^{(1/2)}*(-2*(-\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*\arctan(1/2*2^{(1/2)/(-\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}+2^{(1/2)})/(-c*(\sec(f*x+e)-1))^{(1/2)}*\tan(f*x+e)$

3.68.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 272, normalized size of antiderivative = 3.53

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{\sqrt{c-c\sec(e+fx)}} dx$$

$$= \frac{\left[\sqrt{2}ac\sqrt{-\frac{1}{c}} \log \left(-\frac{2\sqrt{2}(\cos(fx+e)^2+\cos(fx+e))\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\sqrt{-\frac{1}{c}}-(3\cos(fx+e)+1)\sin(fx+e)}{(\cos(fx+e)-1)\sin(fx+e)} \right) \sin(fx+e) - 2(a\cos(fx+e)+a)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}} \right]}{cf \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `[(sqrt(2)*a*c*sqrt(-1/c)*log(-(2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*sqrt(-1/c) - (3*cos(f*x + e) + 1)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) - 2*(a*cos(f*x + e) + a)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c*f*sin(f*x + e)), 2*(sqrt(2)*a*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - (a*cos(f*x + e) + a)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c*f*sin(f*x + e))]`

3.68.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{\sqrt{c-c\sec(e+fx)}} dx = a \left(\int \frac{\sec(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx + \int \frac{\sec^2(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**(1/2),x)`

output `a*(Integral(sec(e + f*x)/sqrt(-c*sec(e + f*x) + c), x) + Integral(sec(e + f*x)**2/sqrt(-c*sec(e + f*x) + c), x))`

3.68.7 Maxima [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{\sqrt{c - c \sec(e + fx)}} dx = \int \frac{(a \sec(fx + e) + a) \sec(fx + e)}{\sqrt{-c \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((a*sec(f*x + e) + a)*sec(f*x + e)/sqrt(-c*sec(f*x + e) + c), x)`

3.68.8 Giac [A] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.79

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{\sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{2a \left(\frac{\sqrt{2} \arctan\left(\frac{\sqrt{c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c}}{\sqrt{c}}\right)}{\sqrt{c}} + \frac{\sqrt{2}}{\sqrt{c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c}} \right)}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `2*a*(sqrt(2)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/sqrt(c) + sqrt(2)/sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c))/f`

3.68.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{a + \frac{a}{\cos(e+fx)}}{\cos(e+fx) \sqrt{c - \frac{c}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`output `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)), x)`

3.69 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{3/2}} dx$

3.69.1 Optimal result 544
 3.69.2 Mathematica [A] (verified) 544
 3.69.3 Rubi [A] (verified) 545
 3.69.4 Maple [B] (verified) 546
 3.69.5 Fracas [B] (verification not implemented) 547
 3.69.6 Sympy [F] 548
 3.69.7 Maxima [F] 548
 3.69.8 Giac [A] (verification not implemented) 548
 3.69.9 Mupad [F(-1)] 549

3.69.1 Optimal result

Integrand size = 32, antiderivative size = 76

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{3/2}} dx = \frac{a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{\sqrt{2}c^{3/2}f} - \frac{a \tan(e+fx)}{f(c-c \sec(e+fx))^{3/2}}$$

output `1/2*a*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/c^(3/2)/f*2^(1/2)-a*tan(f*x+e)/f/(c-c*sec(f*x+e))^(3/2)`

3.69.2 Mathematica [A] (verified)

Time = 0.78 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.07

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{3/2}} dx = \frac{a \left(\cot\left(\frac{1}{2}(e+fx)\right) + \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\sec(e+fx)}}{\sqrt{2}}\right)\sqrt{1+\sec(e+fx)} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{2}} \right)}{cf\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^(3/2),x]`

output `(a*(Cot[(e + f*x)/2] + (ArcTanh[Sqrt[1 + Sec[e + f*x]]/Sqrt[2]]*Sqrt[1 + Sec[e + f*x]]*Tan[(e + f*x)/2])/Sqrt[2]))/(c*f*Sqrt[c - c*Sec[e + f*x]])`

3.69.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4445, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c-c\sec(e+fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{a \int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx}{2c} - \frac{a \tan(e+fx)}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2c} - \frac{a \tan(e+fx)}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{4282} \\
 & \frac{a \int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c\sec(e+fx)}+2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{cf} - \frac{a \tan(e+fx)}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{216} \\
 & \frac{a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{2}c^{3/2}f} - \frac{a \tan(e+fx)}{f(c-c\sec(e+fx))^{3/2}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^(3/2),x]`

output `(a*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])])/(Sqrt[2]*c^(3/2)*f) - (a*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2))`

3.69.3.1 Defintions of rubi rules used

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4282 Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]
```

```
rule 4445 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]
```

3.69.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 177 vs. 2(66) = 132.

Time = 4.18 (sec) , antiderivative size = 178, normalized size of antiderivative = 2.34

method	result
default	$-\frac{a\sqrt{2} \left(\arctan\left(\frac{1}{\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}}\right) (1-\cos(fx+e))^2 \csc(fx+e) - \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \sin(fx+e) \right)}{2cf \sqrt{\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} (1-\cos(fx+e)) \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}}$
parts	$-\frac{a\sqrt{2} (1-\cos(fx+e)) \left(\left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{3}{2}} - \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} (1-\cos(fx+e))^2 \csc(fx+e)^2 - \arctan\left(\frac{1}{\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}}\right) (1-\cos(fx+e))^2 \csc(fx+e) \right)}{4f \left(\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \right)^{\frac{3}{2}} \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2), x, method=_RETURNVERBOSE)
```

3.69. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{3/2}} dx$

output
$$-1/2*a/c/f*2^(1/2)/(c*(1-\cos(f*x+e))^2/((1-\cos(f*x+e))^2*csc(f*x+e)^2-1)*csc(f*x+e)^2)^(1/2)/(1-\cos(f*x+e))/((1-\cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(arctan(1/((1-\cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2))*(1-\cos(f*x+e))^2*csc(f*x+e)-((1-\cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*\sin(f*x+e))$$

3.69.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 147 vs. $2(66) = 132$.

Time = 0.34 (sec) , antiderivative size = 342, normalized size of antiderivative = 4.50

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{3/2}} dx = \frac{\sqrt{2}(ac \cos(fx+e) - ac)\sqrt{-\frac{1}{c}} \log\left(\frac{2\sqrt{2}(\cos(fx+e)^2 + \cos(fx+e))\sqrt{\frac{c \cos(fx+e) - c}{\cos(fx+e)}}}{(\cos(fx+e))}\right)}{4(c^2 f \cos(fx+e) - c^2 f \sin(fx+e))} + \frac{\sqrt{2}(ac \cos(fx+e) - ac) \arctan\left(\frac{\sqrt{2}\sqrt{\frac{c \cos(fx+e) - c}{\cos(fx+e)}} \cos(fx+e)}{\sqrt{c} \sin(fx+e)}\right) \sin(fx+e)}{\sqrt{c}} - 2(a \cos(fx+e))^2 + a \cos(fx+e) \sqrt{\frac{c \cos(fx+e) - c}{\cos(fx+e)}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2),x, algorithm="fricas")`

output
$$[1/4*(\sqrt{2}*(a*c*\cos(f*x + e) - a*c)*\sqrt{-1/c}*\log((2*\sqrt{2}*(\cos(f*x + e)^2 + \cos(f*x + e))*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)})*\sqrt{-1/c} + (3*\cos(f*x + e) + 1)*\sin(f*x + e))/((\cos(f*x + e) - 1)*\sin(f*x + e)))*\sin(f*x + e) + 4*(a*\cos(f*x + e)^2 + a*\cos(f*x + e))*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e))}/((c^2*f*\cos(f*x + e) - c^2*f)*\sin(f*x + e)), -1/2*(\sqrt{2}*(a*c*\cos(f*x + e) - a*c)*\arctan(\sqrt{2}*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)})*\cos(f*x + e)/(\sqrt{c}*\sin(f*x + e)))*\sin(f*x + e)/\sqrt{c} - 2*(a*\cos(f*x + e)^2 + a*\cos(f*x + e))*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e))}/((c^2*f*\cos(f*x + e) - c^2*f)*\sin(f*x + e))]$$

3.69.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{3/2}} dx = a \left(\int \frac{\sec(e+fx)}{-c\sqrt{-c\sec(e+fx)+c}\sec(e+fx)+c\sqrt{-c\sec(e+fx)+c}} \right. \\ \left. + \int \frac{\sec^2(e+fx)}{-c\sqrt{-c\sec(e+fx)+c}\sec(e+fx)+c\sqrt{-c\sec(e+fx)+c}} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**(3/2),x)`

output `a*(Integral(sec(e + f*x)/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x) + Integral(sec(e + f*x)**2/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x))`

3.69.7 Maxima [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{(a\sec(fx+e)+a)\sec(fx+e)}{(-c\sec(fx+e)+c)^{3/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((a*sec(f*x + e) + a)*sec(f*x + e)/(-c*sec(f*x + e) + c)^(3/2), x)`

3.69.8 Giac [A] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{3/2}} dx = \frac{\sqrt{2} \left(\sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}} \right) + \frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\tan(\frac{1}{2}fx + \frac{1}{2}e)^2} \right) a}{2c^2f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `-1/2*sqrt(2)*(sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) + sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/tan(1/2*f*x + 1/2*e)^2)*a/(c^2*f)`

3.69.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{a + \frac{a}{\cos(e+fx)}}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),x)`

output `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)), x)`

3.70
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{5/2}} dx$$

3.70.1	Optimal result	550
3.70.2	Mathematica [C] (verified)	550
3.70.3	Rubi [A] (verified)	551
3.70.4	Maple [B] (verified)	553
3.70.5	Fricas [A] (verification not implemented)	554
3.70.6	Sympy [F]	554
3.70.7	Maxima [F]	555
3.70.8	Giac [A] (verification not implemented)	555
3.70.9	Mupad [F(-1)]	556

3.70.1 Optimal result

Integrand size = 32, antiderivative size = 113

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{5/2}} dx = \frac{a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{8\sqrt{2}c^{5/2}f} - \frac{a \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}} + \frac{a \tan(e+fx)}{8cf(c-c \sec(e+fx))^{3/2}}$$

output `1/16*a*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/c^(5/2)/f*2^(1/2)-1/2*a*tan(f*x+e)/f/(c-c*sec(f*x+e))^(5/2)+1/8*a*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(3/2)`

3.70.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.48 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.53

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c-c \sec(e+fx))^{5/2}} dx = \frac{a \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, 3, \frac{5}{2}, \frac{1}{2}(1+\sec(e+fx))\right) (1+\sec(e+fx)) \tan(e+fx)}{12c^2 f \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^(5/2),x]`

output $-1/12*(a*\text{Hypergeometric2F1}[3/2, 3, 5/2, (1 + \text{Sec}[e + f*x])/2]*(1 + \text{Sec}[e + f*x])*\text{Tan}[e + f*x])/(c^2*f*\text{Sqrt}[c - c*\text{Sec}[e + f*x]])$

3.70.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4445, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c-c\sec(e+fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{a \int \frac{\sec(e+fx)}{(c-c\sec(e+fx))^{3/2}} dx}{4c} - \frac{a \tan(e+fx)}{2f(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{4c} - \frac{a \tan(e+fx)}{2f(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4283} \\
 & -\frac{a \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)}{2f(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)}{2f(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4282}
 \end{aligned}$$

3.70. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{a \left(-\frac{\int \frac{1}{c^2 \tan^2(e+fx) + 2c} d \frac{c \tan(e+fx)}{\sqrt{c - c \sec(e+fx)}}}{2cf} - \frac{\tan(e+fx)}{2f(c - c \sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)}{2f(c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{216} \\
 & \frac{a \left(-\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c - c \sec(e+fx)}}\right)}{2\sqrt{2}c^{3/2}f} - \frac{\tan(e+fx)}{2f(c - c \sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)}{2f(c - c \sec(e+fx))^{5/2}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/2*(a*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(5/2)) - (a*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2))))/(4*c)`

3.70.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4283 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

```
rule 4445 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]
```

3.70.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. 2(94) = 188.

Time = 4.19 (sec) , antiderivative size = 269, normalized size of antiderivative = 2.38

method	result
default	$-\frac{a\sqrt{2} \left(\arctan \left(\frac{1}{\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} \right) (1-\cos(fx+e))^4 \csc(fx+e) - \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{3}{2}} (1-\cos(fx+e))^2 \right)}{16c^2 f \sqrt{\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} (1-\cos(fx+e))^2}}$
parts	$\frac{a\sqrt{2}(1-\cos(fx+e)) \left(\left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{5}{2}} (1-\cos(fx+e))^2 \csc(fx+e)^2 - \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{3}{2}} (1-\cos(fx+e))^2 \right)}{16c^2 f \sqrt{\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} (1-\cos(fx+e))^2}}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)
```

```
output -1/16*a/c^2/f*2^(1/2)/(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*csc(f*x+e)^2)^(1/2)/(1-cos(f*x+e))^3/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(arctan(1/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2))*(1-cos(f*x+e))^4*csc(f*x+e)-((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(3/2)*(1-cos(f*x+e))^2*sin(f*x+e))+((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(1-cos(f*x+e))^4*csc(f*x+e)-2*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(3/2)*sin(f*x+e)^3)
```

3.70. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{5/2}} dx$

3.70.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 405, normalized size of antiderivative = 3.58

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{5/2}} dx = \frac{\sqrt{2}(a\cos(fx+e)^2 - 2a\cos(fx+e) + a)\sqrt{-c}\log\left(-\frac{2\sqrt{2}(\cos(fx+e) - 1)\sin(fx+e)}{\cos(fx+e) - c}\right) + \sqrt{2}(a\cos(fx+e)^2 - 2a\cos(fx+e) + a)\sqrt{c}\arctan\left(\frac{\sqrt{2}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\cos(fx+e)}{\sqrt{c}\sin(fx+e)}\right)\sin(fx+e) - 2(3a\cos(fx+e)^3 + 4a\cos(fx+e)^2 + a\cos(fx+e))\sqrt{(c\cos(fx+e)-c)/\cos(fx+e)}}{16(c^3f\cos(fx+e)^2 - 2c^3f\cos(fx+e) + c^3f)\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2),x, algorithm="fracas")`

output `[-1/32*(sqrt(2)*(a*cos(f*x + e)^2 - 2*a*cos(f*x + e) + a)*sqrt(-c)*log(-(2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) - (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) - 4*(3*a*cos(f*x + e)^3 + 4*a*cos(f*x + e)^2 + a*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x + e)), -1/16*(sqrt(2)*(a*cos(f*x + e)^2 - 2*a*cos(f*x + e) + a)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(3*a*cos(f*x + e)^3 + 4*a*cos(f*x + e)^2 + a*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x + e))]`

3.70.6 SymPy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{5/2}} dx = a \left(\int \frac{\sec(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}} dx + \int \frac{\sec^2(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}+c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**(5/2),x)`

```
output a*(Integral(sec(e + f*x)/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 -
2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f*x)
+ c)), x) + Integral(sec(e + f*x)**2/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(
e + f*x)**2 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c
*sec(e + f*x) + c)), x))
```

3.70.7 Maxima [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c - c \sec(e + fx))^{5/2}} dx = \int \frac{(a \sec(fx + e) + a) \sec(fx + e)}{(-c \sec(fx + e) + c)^{5/2}} dx$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2),x, algorithm=
"maxima")
```

```
output integrate((a*sec(f*x + e) + a)*sec(f*x + e)/(-c*sec(f*x + e) + c)^(5/2), x
)
```

3.70.8 Giac [A] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.92

$$\frac{\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c - c \sec(e + fx))^{5/2}} dx = \sqrt{2} \left(a\sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}} \right) + \frac{(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)^{\frac{3}{2}} ac - \sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - cac^2}}{c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4} \right)}{16 c^3 f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2),x, algorithm=
"giac")
```

```
output -1/16*sqrt(2)*(a*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)
) + ((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*a*c - sqrt(c*tan(1/2*f*x + 1/2*e)
)^2 - c)*a*c^2)/(c^2*tan(1/2*f*x + 1/2*e)^4))/(c^3*f)
```

3.70.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{a + \frac{a}{\cos(e+fx)}}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x)`output `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)), x)`

3.71 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx$

3.71.1	Optimal result	557
3.71.2	Mathematica [A] (verified)	558
3.71.3	Rubi [A] (verified)	558
3.71.4	Maple [A] (verified)	560
3.71.5	Fricas [A] (verification not implemented)	561
3.71.6	Sympy [F(-1)]	561
3.71.7	Maxima [F(-1)]	562
3.71.8	Giac [A] (verification not implemented)	562
3.71.9	Mupad [B] (verification not implemented)	563

3.71.1 Optimal result

Integrand size = 34, antiderivative size = 171

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx =$$

$$\frac{256c^4(a + a \sec(e + fx))^2 \tan(e + fx)}{1155f \sqrt{c - c \sec(e + fx)}} - \frac{64c^3(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{231f} - \frac{8c^2(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} \tan(e + fx)}{33f} - \frac{2c(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} \tan(e + fx)}{11f}$$

output

```
-8/33*c^2*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f-2/11*c*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f-256/1155*c^4*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-64/231*c^3*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.71.2 Mathematica [A] (verified)

Time = 2.48 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.51

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx = \frac{2a^2c^3 \cos^4\left(\frac{1}{2}(e + fx)\right) (-1930 + 3419 \cos(e + fx) - 1510 \cos(2(e + fx)) + 533 \cos(3(e + fx)))}{1155f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(7/2),x]`

output `(2*a^2*c^3*Cos[(e + f*x)/2]^4*(-1930 + 3419*Cos[e + f*x] - 1510*Cos[2*(e + f*x)] + 533*Cos[3*(e + f*x)])*Cot[(e + f*x)/2]*Sec[e + f*x]^5*Sqrt[c - c*Sec[e + f*x]]/(1155*f)`

3.71.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4443, 3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{7/2} dx \\ & \quad \downarrow \text{4443} \\ & \frac{12}{11}c \int \sec(e + fx)(\sec(e + fx)a + a)^2(c - c \sec(e + fx))^{5/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^{5/2}}{11f} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.71. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx$

$$\frac{12}{11}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^2 \left(c - c\csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{5/2}}{11f}$$

↓ 4443

$$\frac{12}{11}c \left(\frac{8}{9}c \int \sec(e + fx)(\sec(e + fx)a + a)^2 (c - c \sec(e + fx))^{3/2} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{3/2}}{9f} \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{5/2}}{11f}$$

↓ 3042

$$\frac{12}{11}c \left(\frac{8}{9}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^2 \left(c - c\csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{3/2}}{9f} \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{5/2}}{11f}$$

↓ 4443

$$\frac{12}{11}c \left(\frac{8}{9}c \left(\frac{4}{7}c \int \sec(e + fx)(\sec(e + fx)a + a)^2 \sqrt{c - c \sec(e + fx)} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 \sqrt{c - c \sec(e + fx)}}{7f} \right) \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{5/2}}{11f}$$

↓ 3042

$$\frac{12}{11}c \left(\frac{8}{9}c \left(\frac{4}{7}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^2 \sqrt{c - c\csc\left(e + fx + \frac{\pi}{2}\right)} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 \sqrt{c - c \sec(e + fx)}}{7f} \right) \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{5/2}}{11f}$$

↓ 4441

$$\frac{12}{11}c \left(\frac{8}{9}c \left(-\frac{8c^2 \tan(e + fx)(a \sec(e + fx) + a)^2}{35f \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 \sqrt{c - c \sec(e + fx)}}{7f} \right) \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{5/2}}{11f}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(7/2),x]`

3.71. $\int \sec(e + fx)(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{7/2} dx$


```
output (-2*c*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(11*f) + (12*c*((-2*c*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(9*f) + (8*c*((-8*c^2*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(35*f*Sqrt[c - c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]])*Tan[e + f*x])/(7*f)))/9)/11
```

3.71.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.71.4 Maple [A] (verified)

Time = 16.90 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.46

method	result
default	$\frac{2a^2c^3 \left(533 \cos(fx+e)^3 - 755 \cos(fx+e)^2 + 455 \cos(fx+e) - 105 \right) \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1)^3 \sec(fx+e)^5 \csc(fx+e)}{1155f}$
parts	$-\frac{2a^2(\sec(fx+e)-1)^3 \left(177 \cos(fx+e)^3 - 71 \cos(fx+e)^2 + 27 \cos(fx+e) - 5 \right) c^3 \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1) \csc(fx+e)}{35f(\cos(fx+e)-1)^3} - \frac{2a^2c^3 \left(533 \cos(fx+e)^3 - 755 \cos(fx+e)^2 + 455 \cos(fx+e) - 105 \right) \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1)^3 \sec(fx+e)^5 \csc(fx+e)}{1155f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(7/2),x,method=_RETURNV ERBOSE)
```

3.71. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx$

output $2/1155*a^2*c^3/f*(533*\cos(f*x+e)^3-755*\cos(f*x+e)^2+455*\cos(f*x+e)-105)*(-c*(\sec(f*x+e)-1))^{(1/2)}*(\cos(f*x+e)+1)^3*\sec(f*x+e)^5*\csc(f*x+e)$

3.71.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.86

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx) + fx)^{7/2} dx = \frac{2(533 a^2 c^3 \cos(fx + e)^6 + 844 a^2 c^3 \cos(fx + e)^5 - 211 a^2 c^3 \cos(fx + e)^4 - 472 a^2 c^3 \cos(fx + e)^3 + 295 a^2 c^3 \cos(fx + e)^2 + 140 a^2 c^3 \cos(fx + e) - 105 a^2 c^3) \sqrt{(c \cos(fx + e) - c) / \cos(fx + e)}}{1155 f \cos(fx + e)^5 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(7/2),x, algorithm="fricas")`

output $2/1155*(533*a^2*c^3*\cos(f*x + e)^6 + 844*a^2*c^3*\cos(f*x + e)^5 - 211*a^2*c^3*\cos(f*x + e)^4 - 472*a^2*c^3*\cos(f*x + e)^3 + 295*a^2*c^3*\cos(f*x + e)^2 + 140*a^2*c^3*\cos(f*x + e) - 105*a^2*c^3)*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^5*\sin(f*x + e))$

3.71.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**(7/2),x)`

output `Timed out`

3.71.7 Maxima [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(7/2),x, algorithm m="maxima")`

output `Timed out`

3.71.8 Giac [A] (verification not implemented)

Time = 1.02 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.64

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx =$$

$$\frac{64\sqrt{2}\left(231\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^3c^3 + 495\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^2c^4 + 385\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)c^5 + 105c^6\right)a^2c^3}{1155\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{11}{2}}f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(7/2),x, algorithm m="giac")`

output `-64/1155*sqrt(2)*(231*(c*tan(1/2*f*x + 1/2*e)^2 - c)^3*c^3 + 495*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^4 + 385*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^5 + 105*c^6)*a^2*c^3/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(11/2)*f)`

3.71.9 Mupad [B] (verification not implemented)

Time = 25.26 (sec) , antiderivative size = 606, normalized size of antiderivative = 3.54

$$\begin{aligned}
& \int \sec(e + fx)(a + a \sec(e + fx))^2(c \\
& - c \sec(e + fx))^{7/2} dx = \frac{\left(\frac{a^2 c^3 2i}{f} + \frac{a^2 c^3 e^{e li + f x li} 1066i}{1155 f}\right) \sqrt{c - \frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}{e^{e li + f x li} - 1} \\
& + \frac{\left(\frac{a^2 c^3 64i}{11 f} - \frac{a^2 c^3 e^{e li + f x li} 64i}{11 f}\right) \sqrt{c - \frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^5} \\
& - \frac{\left(\frac{a^2 c^3 32i}{3 f} - \frac{a^2 c^3 e^{e li + f x li} 608i}{33 f}\right) \sqrt{c - \frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^4} \\
& - \frac{\left(\frac{a^2 c^3 4i}{f} + \frac{a^2 c^3 e^{e li + f x li} 2932i}{1155 f}\right) \sqrt{c - \frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)} \\
& + \frac{\left(\frac{a^2 c^3 16i}{5 f} + \frac{a^2 c^3 e^{e li + f x li} 4272i}{385 f}\right) \sqrt{c - \frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^2} \\
& + \frac{\left(\frac{a^2 c^3 32i}{7 f} - \frac{a^2 c^3 e^{e li + f x li} 4640i}{231 f}\right) \sqrt{c - \frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^3}
\end{aligned}$$

```
input int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(7/2))/cos(e + f*x),x)
```

```
output (((a^2*c^3*2i)/f + (a^2*c^3*exp(e*1i + f*x*1i)*1066i)/(1155*f))*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/(exp(e*1i + f*x*1i) - 1) + (((a^2*c^3*64i)/(11*f) - (a^2*c^3*exp(e*1i + f*x*1i)*64i)/(11*f))*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^5) - (((a^2*c^3*32i)/(3*f) - (a^2*c^3*exp(e*1i + f*x*1i)*608i)/(33*f))*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^4) - (((a^2*c^3*4i)/f + (a^2*c^3*exp(e*1i + f*x*1i)*2932i)/(1155*f))*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)) + (((a^2*c^3*16i)/(5*f) + (a^2*c^3*exp(e*1i + f*x*1i)*4272i)/(385*f))*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^2) + (((a^2*c^3*32i)/(7*f) - (a^2*c^3*exp(e*1i + f*x*1i)*4640i)/(231*f))*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^3)
```

3.71. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{7/2} dx$

3.72 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx$

3.72.1	Optimal result	564
3.72.2	Mathematica [A] (verified)	565
3.72.3	Rubi [A] (verified)	565
3.72.4	Maple [A] (verified)	567
3.72.5	Fricas [A] (verification not implemented)	567
3.72.6	Sympy [F]	568
3.72.7	Maxima [F]	568
3.72.8	Giac [A] (verification not implemented)	569
3.72.9	Mupad [B] (verification not implemented)	570

3.72.1 Optimal result

Integrand size = 34, antiderivative size = 128

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx =$$

$$-\frac{64c^3(a + a \sec(e + fx))^2 \tan(e + fx)}{315f \sqrt{c - c \sec(e + fx)}}$$

$$-\frac{16c^2(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{63f}$$

$$-\frac{2c(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} \tan(e + fx)}{9f}$$

output

```
-2/9*c*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f-64/315*c^3*(
a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-16/63*c^2*(a+a*sec(f
*x+e))^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.72.2 Mathematica [A] (verified)

Time = 1.66 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.61

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx = \frac{4a^2c^2 \cos^4\left(\frac{1}{2}(e + fx)\right) (177 - 220 \cos(e + fx) + 107 \cos(2(e + fx))) \cot\left(\frac{1}{2}(e + fx)\right) \sec^4(e + fx)}{315f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(5/2),x]`

output `(4*a^2*c^2*Cos[(e + f*x)/2]^4*(177 - 220*Cos[e + f*x] + 107*Cos[2*(e + f*x)])*Cot[(e + f*x)/2]*Sec[e + f*x]^4*Sqrt[c - c*Sec[e + f*x]]/(315*f)`

3.72.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx \\ & \quad \downarrow \text{4443} \\ & \frac{8}{9}c \int \sec(e + fx)(\sec(e + fx)a + a)^2(c - c \sec(e + fx))^{3/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^{3/2}}{9f} \\ & \quad \downarrow \text{3042} \\ & \frac{8}{9}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^{3/2}}{9f} \end{aligned}$$

3.72. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx$

↓ 4443

$$\frac{8}{9}c \left(\frac{4}{7}c \int \sec(e+fx)(\sec(e+fx)a+a)^2 \sqrt{c-c\sec(e+fx)} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}}{7f} \right. \\ \left. - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^2 (c-c\sec(e+fx))^{3/2}}{9f} \right)$$

↓ 3042

$$\frac{8}{9}c \left(\frac{4}{7}c \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)^2 \sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}}{7f} \right. \\ \left. - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^2 (c-c\sec(e+fx))^{3/2}}{9f} \right)$$

↓ 4441

$$\frac{8}{9}c \left(-\frac{8c^2 \tan(e+fx)(a \sec(e+fx)+a)^2}{35f\sqrt{c-c\sec(e+fx)}} - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}}{7f} \right) - \\ \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^2 (c-c\sec(e+fx))^{3/2}}{9f}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(5/2),x]`

output `(-2*c*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(9*f) + (8*c*((-8*c^2*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(35*f*Sqrt[c - c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x])*Tan[e + f*x])/(7*f)))/9`

3.72.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_.*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.72. $\int \sec(e+fx)(a+a\sec(e+fx))^2(c-c\sec(e+fx))^{5/2} dx$

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.72.4 Maple [A] (verified)

Time = 15.64 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.54

method	result
default	$\frac{2a^2c^2(107\cos(fx+e)^2-110\cos(fx+e)+35)\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)^3\sec(fx+e)^4\csc(fx+e)}{315f}$
parts	$\frac{2a^2(\sec(fx+e)-1)^2(43\cos(fx+e)^2-14\cos(fx+e)+3)c^2\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)\csc(fx+e)}{15f(\cos(fx+e)-1)^2} + \frac{2a^2(584\cos(fx+e)^4-...}{...}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 2/315*a^2*c^2/f*(107*cos(f*x+e)^2-110*cos(f*x+e)+35)*(-c*(sec(f*x+e)-1))^(
1/2)*(cos(f*x+e)+1)^3*sec(f*x+e)^4*csc(f*x+e)
```

3.72.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.02

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^5 dx = \frac{2(107a^2c^2 \cos(fx + e)^5 + 211a^2c^2 \cos(fx + e)^4 + 26a^2c^2 \cos(fx + e)^3 - 118a^2c^2 \cos(fx + e)^2 + 26a^2c^2 \cos(fx + e) - 118a^2c^2) \sin(fx + e)}{315f \cos(fx + e)^4 \sin(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(5/2),x, algorith
m="fricas")
```


output `2/315*(107*a^2*c^2*cos(f*x + e)^5 + 211*a^2*c^2*cos(f*x + e)^4 + 26*a^2*c^2*cos(f*x + e)^3 - 118*a^2*c^2*cos(f*x + e)^2 - 5*a^2*c^2*cos(f*x + e) + 35*a^2*c^2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)^4*sin(f*x + e))`

3.72.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx = a^2 \left(\int c^2 \sqrt{-c \sec(e + fx) + c \sec(e + fx)} dx + \int (-2c^2 \sqrt{-c \sec(e + fx) + c \sec^3(e + fx)}) dx + \int c^2 \sqrt{-c \sec(e + fx) + c \sec^5(e + fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**(5/2),x)`

output `a**2*(Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(-2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3, x) + Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**5, x))`

3.72.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx = \int (a \sec(fx + e) + a)^2(-c \sec(fx + e) + c)^{5/2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(5/2),x, algorithm="maxima")`

```
output -2/315*(315*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e)
+ 1)^(1/4)*((a^2*c^2*f*cos(2*f*x + 2*e)^4 + a^2*c^2*f*sin(2*f*x + 2*e)^4 +
4*a^2*c^2*f*cos(2*f*x + 2*e)^3 + 6*a^2*c^2*f*cos(2*f*x + 2*e)^2 + 4*a^2*c
^2*f*cos(2*f*x + 2*e) + a^2*c^2*f + 2*(a^2*c^2*f*cos(2*f*x + 2*e)^2 + 2*a^
2*c^2*f*cos(2*f*x + 2*e) + a^2*c^2*f)*sin(2*f*x + 2*e)^2)*integrate((((cos
(8*f*x + 8*e)*cos(2*f*x + 2*e) + 3*cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 3*c
os(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(8*f*x + 8*e)*s
in(2*f*x + 2*e) + 3*sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 3*sin(4*f*x + 4*e)
*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*cos(9/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + (cos(2*f*x + 2*e)*sin(8*f*x + 8*e) + 3*cos(2*f*x + 2*
e)*sin(6*f*x + 6*e) + 3*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(8*f*x + 8*
e)*sin(2*f*x + 2*e) - 3*cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 3*cos(4*f*x +
4*e)*sin(2*f*x + 2*e))*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))
))*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x
+ 2*e)*sin(8*f*x + 8*e) + 3*cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 3*cos(2*f*
x + 2*e)*sin(4*f*x + 4*e) - cos(8*f*x + 8*e)*sin(2*f*x + 2*e) - 3*cos(6*f*
x + 6*e)*sin(2*f*x + 2*e) - 3*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(9/2*a
rctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (cos(8*f*x + 8*e)*cos(2*f*x
+ 2*e) + 3*cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 3*cos(4*f*x + 4*e)*cos(2*f*
x + 2*e) + cos(2*f*x + 2*e)^2 + sin(8*f*x + 8*e)*sin(2*f*x + 2*e) + 3*s...
```

3.72.8 Giac [A] (verification not implemented)

Time = 0.83 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.66

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx =$$

$$\frac{32\sqrt{2}\left(63\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^2 c^3 + 90\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)c^4 + 35c^5\right)a^2 c^2}{315\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{9}{2}} f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(5/2),x, algorith
m="giac")
```

```
output -32/315*sqrt(2)*(63*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^3 + 90*(c*tan(1/2*f
*x + 1/2*e)^2 - c)*c^4 + 35*c^5)*a^2*c^2/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(
9/2)*f)
```

3.72. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{5/2} dx$

3.72.9 Mupad [B] (verification not implemented)

Time = 20.22 (sec) , antiderivative size = 503, normalized size of antiderivative = 3.93

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{5/2} dx = \frac{\left(\frac{a^2 c^2 2i}{f} + \frac{a^2 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 214i}{315 f}\right) \sqrt{c - \frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}{e^{e \operatorname{li} + f x \operatorname{li}} - 1} + \frac{\left(\frac{a^2 c^2 32i}{9 f} + \frac{a^2 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 32i}{9 f}\right) \sqrt{c - \frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)^4} - \frac{\left(\frac{a^2 c^2 64i}{7 f} + \frac{a^2 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 320i}{63 f}\right) \sqrt{c - \frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)^3} + \frac{\left(\frac{a^2 c^2 48i}{5 f} + \frac{a^2 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 368i}{105 f}\right) \sqrt{c - \frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)^2} - \frac{\left(\frac{a^2 c^2 16i}{3 f} + \frac{a^2 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 208i}{315 f}\right) \sqrt{c - \frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)}$$

input `int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x),x)`

output `((a^2*c^2*2i)/f + (a^2*c^2*exp(e*li + f*x*li)*214i)/(315*f))*(c - c/(exp(-e*li - f*x*li)/2 + exp(e*li + f*x*li)/2))^(1/2))/(exp(e*li + f*x*li) - 1) + (((a^2*c^2*32i)/(9*f) + (a^2*c^2*exp(e*li + f*x*li)*32i)/(9*f))*(c - c/(exp(-e*li - f*x*li)/2 + exp(e*li + f*x*li)/2))^(1/2))/((exp(e*li + f*x*li) - 1)*(exp(e*2i + f*x*2i) + 1)^4) - (((a^2*c^2*64i)/(7*f) + (a^2*c^2*exp(e*li + f*x*li)*320i)/(63*f))*(c - c/(exp(-e*li - f*x*li)/2 + exp(e*li + f*x*li)/2))^(1/2))/((exp(e*li + f*x*li) - 1)*(exp(e*2i + f*x*2i) + 1)^3) + (((a^2*c^2*48i)/(5*f) + (a^2*c^2*exp(e*li + f*x*li)*368i)/(105*f))*(c - c/(exp(-e*li - f*x*li)/2 + exp(e*li + f*x*li)/2))^(1/2))/((exp(e*li + f*x*li) - 1)*(exp(e*2i + f*x*2i) + 1)^2) - (((a^2*c^2*16i)/(3*f) + (a^2*c^2*exp(e*li + f*x*li)*208i)/(315*f))*(c - c/(exp(-e*li - f*x*li)/2 + exp(e*li + f*x*li)/2))^(1/2))/((exp(e*li + f*x*li) - 1)*(exp(e*2i + f*x*2i) + 1))`

3.73 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx$

3.73.1	Optimal result	571
3.73.2	Mathematica [A] (verified)	571
3.73.3	Rubi [A] (verified)	572
3.73.4	Maple [A] (verified)	573
3.73.5	Fricas [A] (verification not implemented)	574
3.73.6	Sympy [F]	574
3.73.7	Maxima [F]	575
3.73.8	Giac [A] (verification not implemented)	576
3.73.9	Mupad [B] (verification not implemented)	576

3.73.1 Optimal result

Integrand size = 34, antiderivative size = 85

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx =$$

$$-\frac{8c^2(a + a \sec(e + fx))^2 \tan(e + fx)}{35f \sqrt{c - c \sec(e + fx)}} - \frac{2c(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{7f}$$

output

```
-8/35*c^2*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-2/7*c*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.73.2 Mathematica [A] (verified)

Time = 1.46 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.78

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx = \frac{8a^2c \cos^4\left(\frac{1}{2}(e + fx)\right) (-5 + 9 \cos(e + fx)) \cot\left(\frac{1}{2}(e + fx)\right) \sec^3(e + fx) \sqrt{c - c \sec(e + fx)}}{35f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2),x]`

output `(8*a^2*c*Cos[(e + f*x)/2]^4*(-5 + 9*Cos[e + f*x])*Cot[(e + f*x)/2]*Sec[e + f*x]^3*sqrt[c - c*Sec[e + f*x]])/(35*f)`

3.73.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{4}{7}c \int \sec(e + fx)(\sec(e + fx)a + a)^2 \sqrt{c - c \sec(e + fx)} dx - \\
 & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 \sqrt{c - c \sec(e + fx)}}{7f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4}{7}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right) a + a\right)^2 \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx - \\
 & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 \sqrt{c - c \sec(e + fx)}}{7f} \\
 & \quad \downarrow \text{4441} \\
 & -\frac{8c^2 \tan(e + fx)(a \sec(e + fx) + a)^2}{35f \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2 \sqrt{c - c \sec(e + fx)}}{7f}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2),x]`

```
output (-8*c^2*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(35*f*Sqrt[c - c*Sec[e + f*x]
]) - (2*c*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(7
*f)
```

3.73.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f
*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /
; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
&& NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m*(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^n, x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.73.4 Maple [A] (verified)

Time = 5.52 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

method	result
default	$\frac{2a^2c(9 \cos(fx+e)-5)\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)^3 \sec(fx+e)^3 \csc(fx+e)}{35f}$
parts	$-\frac{2a^2(\sec(fx+e)-1)(5 \cos(fx+e)-1)c\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1) \csc(fx+e)}{3f(\cos(fx+e)-1)} - \frac{2a^2(104 \cos(fx+e)^3 - 52 \cos(fx+e)^2 + 3 \cos(fx+e) - 1)}{3f(\cos(fx+e)-1)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(3/2),x,method=_RETURNV
ERBOSE)
```

3.73. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx$

output $2/35*a^2*c/f*(9*\cos(f*x+e)-5)*(-c*(\sec(f*x+e)-1))^{(1/2)}*(\cos(f*x+e)+1)^3*\sec(f*x+e)^3*\csc(f*x+e)$

3.73.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.24

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx = \frac{2(9a^2c \cos(fx + e)^4 + 22a^2c \cos(fx + e)^3 + 12a^2c \cos(fx + e)^2 - 6a^2c \cos(fx + e) - 5a^2)}{35f \cos(fx + e)^3 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(3/2),x, algorithm="fricas")`

output $2/35*(9*a^2*c*\cos(f*x + e)^4 + 22*a^2*c*\cos(f*x + e)^3 + 12*a^2*c*\cos(f*x + e)^2 - 6*a^2*c*\cos(f*x + e) - 5*a^2*c)*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^3*\sin(f*x + e))$

3.73.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx \\ &= a^2 \left(\int c \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx \right. \\ &+ \int c \sqrt{-c \sec(e + fx) + c} \sec^2(e + fx) dx \\ &+ \int \left(-c \sqrt{-c \sec(e + fx) + c} \sec^3(e + fx) \right) dx \\ &+ \left. \int \left(-c \sqrt{-c \sec(e + fx) + c} \sec^4(e + fx) \right) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**(3/2),x)`

output `a**2*(Integral(c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2, x) + Integral(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3, x) + Integral(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4, x))`

3.73.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx = \int (a \sec(fx + e) + a)^2(-c \sec(fx + e) + c)^{3/2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `2/35*(35*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(3/4)*((a^2*c*f*cos(2*f*x + 2*e)^2 + a^2*c*f*sin(2*f*x + 2*e)^2 + 2*a^2*c*f*cos(2*f*x + 2*e) + a^2*c*f)*integrate((cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/4)*(((cos(8*f*x + 8*e)*cos(2*f*x + 2*e) + 3*cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 3*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(8*f*x + 8*e)*sin(2*f*x + 2*e) + 3*sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 3*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (cos(2*f*x + 2*e)*sin(8*f*x + 8*e) + 3*cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 3*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(8*f*x + 8*e)*sin(2*f*x + 2*e) - 3*cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 3*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x + 2*e)*sin(8*f*x + 8*e) + 3*cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 3*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(8*f*x + 8*e)*sin(2*f*x + 2*e) - 3*cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 3*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (cos(8*f*x + 8*e)*cos(2*f*x + 2*e) + 3*cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 3*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(8*f*x + 8*e)*sin(2*f*x + 2*e) + 3*sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 3*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*sin(7/2*...`

3.73.8 Giac [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.68

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx = \frac{16\sqrt{2}\left(7\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)c^4 + 5c^5\right)a^2}{35\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{7}{2}}f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(3/2),x, algorithm m="giac")`

output `-16/35*sqrt(2)*(7*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^4 + 5*c^5)*a^2/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(7/2)*f)`

3.73.9 Mupad [B] (verification not implemented)

Time = 17.15 (sec) , antiderivative size = 384, normalized size of antiderivative = 4.52

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx))^{3/2} dx = \frac{\sqrt{c - \frac{c}{\frac{e^{-e li - f x li} + e^{e li + f x li}}{2}}}}{e^{e li + f x li} - 1} \left(\frac{a^2 c 2i}{f} + \frac{a^2 c e^{e li + f x li} 18i}{35 f} \right) - \frac{\sqrt{c - \frac{c}{\frac{e^{-e li - f x li} + e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1)(e^{2i + f x 2i} + 1)^3} \left(\frac{a^2 c 16i}{7 f} - \frac{a^2 c e^{e li + f x li} 16i}{7 f} \right) - \frac{\sqrt{c - \frac{c}{\frac{e^{-e li - f x li} + e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1)(e^{2i + f x 2i} + 1)} \left(\frac{a^2 c 4i}{f} - \frac{a^2 c e^{e li + f x li} 44i}{35 f} \right) + \frac{\sqrt{c - \frac{c}{\frac{e^{-e li - f x li} + e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1)(e^{2i + f x 2i} + 1)^2} \left(\frac{a^2 c 24i}{5 f} - \frac{a^2 c e^{e li + f x li} 72i}{35 f} \right)$$

input `int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(3/2))/cos(e + f*x),x)`

output $((c - c/(\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2))^{1/2} * ((a^2*c*2i)/f + (a^2*c*\exp(e*1i + f*x*1i)*18i)/(35*f)))/(\exp(e*1i + f*x*1i) - 1) - ((c - c/(\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2))^{1/2} * ((a^2*c*16i)/(7*f) - (a^2*c*\exp(e*1i + f*x*1i)*16i)/(7*f)))/((\exp(e*1i + f*x*1i) - 1) * (\exp(e*2i + f*x*2i) + 1)^3) - ((c - c/(\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2))^{1/2} * ((a^2*c*4i)/f - (a^2*c*\exp(e*1i + f*x*1i)*44i)/(35*f)))/((\exp(e*1i + f*x*1i) - 1) * (\exp(e*2i + f*x*2i) + 1)) + ((c - c/(\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2))^{1/2} * ((a^2*c*24i)/(5*f) - (a^2*c*\exp(e*1i + f*x*1i)*72i)/(35*f)))/((\exp(e*1i + f*x*1i) - 1) * (\exp(e*2i + f*x*2i) + 1)^2)$

3.74 $\int \sec(e+fx)(a+a \sec(e+fx))^2 \sqrt{c - c \sec(e + fx)} dx$

3.74.1	Optimal result	578
3.74.2	Mathematica [A] (verified)	578
3.74.3	Rubi [A] (verified)	579
3.74.4	Maple [A] (verified)	580
3.74.5	Fricas [B] (verification not implemented)	580
3.74.6	Sympy [F]	581
3.74.7	Maxima [F]	581
3.74.8	Giac [A] (verification not implemented)	582
3.74.9	Mupad [B] (verification not implemented)	583

3.74.1 Optimal result

Integrand size = 34, antiderivative size = 41

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} dx$$

$$= -\frac{2c(a + a \sec(e + fx))^2 \tan(e + fx)}{5f \sqrt{c - c \sec(e + fx)}}$$

output `-2/5*c*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)`

3.74.2 Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.34

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{8a^2 \cos^4\left(\frac{1}{2}(e + fx)\right) \cot\left(\frac{1}{2}(e + fx)\right) \sec^2(e + fx) \sqrt{c - c \sec(e + fx)}}{5f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]],x]`

output `(8*a^2*Cos[(e + f*x)/2]^4*Cot[(e + f*x)/2]*Sec[e + f*x]^2*Sqrt[c - c*Sec[e + f*x]])/(5*f)`

3.74.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^2 \sqrt{c - c \sec(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4441}$$

$$\frac{2c \tan(e + fx)(a \sec(e + fx) + a)^2}{5f \sqrt{c - c \sec(e + fx)}}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]],x]`

output `(-2*c*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(5*f*Sqrt[c - c*Sec[e + f*x]])`

3.74.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.74.4 Maple [A] (verified)

Time = 4.71 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

method	result
default	$\frac{2a^2(\cos(fx+e)+1)^3\sqrt{-c(\sec(fx+e)-1)}\sec(fx+e)^2\csc(fx+e)}{5f}$
parts	$-\frac{2a^2\sqrt{-c(\sec(fx+e)-1)}\sin(fx+e)}{f(\cos(fx+e)-1)} + \frac{2a^2\sqrt{-c(\sec(fx+e)-1)}(3+8\cos(fx+e)^3+4\cos(fx+e)^2-\cos(fx+e))\sec(fx+e)^2\csc(fx+e)}{15f}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(1/2),x,method=_RETURNV
ERBOSE)`

output `2/5*a^2/f*(cos(f*x+e)+1)^3*(-c*(sec(f*x+e)-1))^(1/2)*sec(f*x+e)^2*csc(f*x+
e)`

3.74.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(37) = 74.

Time = 0.27 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.05

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{2(a^2 \cos(fx + e)^3 + 3a^2 \cos(fx + e)^2 + 3a^2 \cos(fx + e) + a^2) \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}}}{5f \cos(fx + e)^2 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(1/2),x, algorithm
m="fracas")`

output `2/5*(a^2*cos(f*x + e)^3 + 3*a^2*cos(f*x + e)^2 + 3*a^2*cos(f*x + e) + a^2)
*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)^2*sin(f*x + e))`

3.74.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} dx$$

$$= a^2 \left(\int \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx + \int 2\sqrt{-c \sec(e + fx) + c} \sec^2(e + fx) dx \right. \\ \left. + \int \sqrt{-c \sec(e + fx) + c} \sec^3(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e))**(1/2),x)`

output `a**2*(Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2, x) + Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3, x))`

3.74.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} dx$$

$$= \int (a \sec(fx + e) + a)^2 \sqrt{-c \sec(fx + e) + c} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

```
output 2/5*(5*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^
(1/4)*(3*(a^2*f*cos(2*f*x + 2*e)^2 + a^2*f*sin(2*f*x + 2*e)^2 + 2*a^2*f*co
s(2*f*x + 2*e) + a^2*f)*integrate((((cos(8*f*x + 8*e)*cos(2*f*x + 2*e) + 3
*cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + 3*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) +
cos(2*f*x + 2*e)^2 + sin(8*f*x + 8*e)*sin(2*f*x + 2*e) + 3*sin(6*f*x + 6*
e)*sin(2*f*x + 2*e) + 3*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*
e)^2)*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (cos(2*f*x +
2*e)*sin(8*f*x + 8*e) + 3*cos(2*f*x + 2*e)*sin(6*f*x + 6*e) + 3*cos(2*f*x
+ 2*e)*sin(4*f*x + 4*e) - cos(8*f*x + 8*e)*sin(2*f*x + 2*e) - 3*cos(6*f*x
+ 6*e)*sin(2*f*x + 2*e) - 3*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*sin(5/2*arc
tan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(1/2*arctan2(sin(2*f*x + 2*e
), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x + 2*e)*sin(8*f*x + 8*e) + 3*cos(2*
f*x + 2*e)*sin(6*f*x + 6*e) + 3*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(8*
f*x + 8*e)*sin(2*f*x + 2*e) - 3*cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 3*cos(
4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e))) - (cos(8*f*x + 8*e)*cos(2*f*x + 2*e) + 3*cos(6*f*x + 6*e)*cos(2*
f*x + 2*e) + 3*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + si
n(8*f*x + 8*e)*sin(2*f*x + 2*e) + 3*sin(6*f*x + 6*e)*sin(2*f*x + 2*e) + 3*
sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*x + 2*e)^2)*sin(5/2*arctan2(si
n(2*f*x + 2*e), cos(2*f*x + 2*e))))*sin(1/2*arctan2(sin(2*f*x + 2*e), c...
```

3.74.8 Giac [A] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} dx = -\frac{8\sqrt{2}a^2c^3}{5\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{5}{2}}f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e))^(1/2),x, algorith
m="giac")
```

```
output -8/5*sqrt(2)*a^2*c^3/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2)*f)
```

3.74.9 Mupad [B] (verification not implemented)

Time = 17.86 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.27

$$\int \sec(e + fx)(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{2a^2 (e^{e + fx} - 1)^5 \sqrt{c - \frac{c}{\frac{e^{-e - fx} - 1}{2} + \frac{e^{e + fx} - 1}{2}}}}{5f (e^{e + fx} - 1) (e^{2e + 2fx} + 1)^2}$$

input `int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x),x)`output `(2*a^2*(exp(e*1i + f*x*1i)*1i + 1i)^5*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/(5*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^2)`

$$3.75 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{\sqrt{c-c \sec(e+fx)}} dx$$

3.75.1	Optimal result	584
3.75.2	Mathematica [A] (verified)	584
3.75.3	Rubi [A] (verified)	585
3.75.4	Maple [A] (verified)	587
3.75.5	Fricas [A] (verification not implemented)	588
3.75.6	Sympy [F]	588
3.75.7	Maxima [F]	589
3.75.8	Giac [A] (verification not implemented)	589
3.75.9	Mupad [F(-1)]	590

3.75.1 Optimal result

Integrand size = 34, antiderivative size = 117

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{\sqrt{c-c \sec(e+fx)}} dx = -\frac{4\sqrt{2}a^2 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{\sqrt{cf}} + \frac{16a^2 \tan(e+fx)}{3f\sqrt{c-c \sec(e+fx)}} - \frac{2a^2\sqrt{c-c \sec(e+fx)} \tan(e+fx)}{3cf}$$

output $-4*a^2*\arctan(1/2*c^{(1/2)}*\tan(f*x+e)*2^{(1/2)/(c-c*\sec(f*x+e))^{(1/2)}}*2^{(1/2)}/f/c^{(1/2)}+16/3*a^2*\tan(f*x+e)/f/(c-c*\sec(f*x+e))^{(1/2)}-2/3*a^2*(c-c*\sec(f*x+e))^{(1/2)}*\tan(f*x+e)/c/f$

3.75.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{\sqrt{c-c \sec(e+fx)}} dx = \frac{2a^2\left(-6\sqrt{2}\sqrt{a}\operatorname{arctanh}\left(\frac{\sqrt{a(1+\sec(e+fx))}}{\sqrt{2}\sqrt{a}}\right) + \sqrt{a(1+\sec(e+fx))}(7+\sec(e+fx))\right) \tan(e+fx)}{3f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/Sqrt[c - c*Sec[e + f*x]],x]`

output `(2*a^2*(-6*Sqrt[2]*Sqrt[a]*ArcTanh[Sqrt[a*(1 + Sec[e + f*x])]/(Sqrt[2]*Sqrt[a])]) + Sqrt[a*(1 + Sec[e + f*x])]*(7 + Sec[e + f*x]))*Tan[e + f*x]/(3*f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.75.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4444, 3042, 4444, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{\sqrt{c-c\sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4444} \\
 & 2a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2\tan(e+fx)(a^2\sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & 2a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2\tan(e+fx)(a^2\sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{4444} \\
 & 2a \left(2a \int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2a\tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right) + \frac{2\tan(e+fx)(a^2\sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& 2a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\sec(e+fx+\frac{\pi}{2})}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right) + \\
& \quad \frac{2 \tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{4282} \\
& 2a \left(\frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{4a \int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c\sec(e+fx)} + 2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{f} \right) + \\
& \quad \frac{2 \tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{216} \\
& \quad \frac{2 \tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f\sqrt{c-c\sec(e+fx)}} + \\
& 2a \left(\frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{cf}} \right)
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/Sqrt[c - c*Sec[e + f*x]],x]`

output `(2*(a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e + f*x]]) + 2*a*((-2*Sqrt[2]*a*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])])/(Sqrt[c]*f) + (2*a*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]]))`

3.75.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4444 `Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.))/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*d*Cot[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/(f*(2*n - 1)*Sqrt[a + b*Csc[e + f*x]]), x] + Simp[2*c*((2*n - 1)/(2*n - 1)) Int[Csc[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/Sqrt[a + b*Csc[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`

3.75.4 Maple [A] (verified)

Time = 4.24 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.92

method	result
default	$-\frac{a^2\sqrt{2} \left(12 \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \tan(fx+e) - 7\sqrt{2} \tan(fx+e) - \sqrt{2} \tan(fx+e) \sec(fx+e) \right)}{3f\sqrt{-c(\sec(fx+e)-1)}}$
parts	$\frac{a^2\sqrt{2} \sin(fx+e) \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right)}{f(\cos(fx+e)+1)\sqrt{-c(\sec(fx+e)-1)}\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} + \frac{a^2\sqrt{2} \left(-3 \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \tan(fx+e) + \sqrt{2} \tan(fx+e) \sec(fx+e) \right)}{3f\sqrt{-c(\sec(fx+e)-1)}}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNV ERBOSE)`

output `-1/3*a^2/f*2^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)*(12*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*tan(f*x+e)-7*2^(1/2)*tan(f*x+e)-2^(1/2)*tan(f*x+e)*sec(f*x+e))`

3.75. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{\sqrt{c-c\sec(e+fx)}} dx$

3.75.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 343, normalized size of antiderivative = 2.93

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{\sqrt{c-c\sec(e+fx)}} dx$$

$$= \frac{2 \left(3\sqrt{2}a^2c\sqrt{-\frac{1}{c}} \cos(fx+e) \log \left(-\frac{2\sqrt{2}(\cos(fx+e)^2+\cos(fx+e))\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\sqrt{-\frac{1}{c}}-(3\cos(fx+e)+1)\sin(fx+e)}{(\cos(fx+e)-1)\sin(fx+e)} \right) \right) \sin(fx+e)}{3cf \cos(fx+e) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `[2/3*(3*sqrt(2)*a^2*c*sqrt(-1/c)*cos(f*x + e)*log(-(2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*sqrt(-1/c) - (3*cos(f*x + e) + 1)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) - (7*a^2*cos(f*x + e)^2 + 8*a^2*cos(f*x + e) + a^2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c*f*cos(f*x + e)*sin(f*x + e)), 2/3*(6*sqrt(2)*a^2*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*cos(f*x + e)*sin(f*x + e) - (7*a^2*cos(f*x + e)^2 + 8*a^2*cos(f*x + e) + a^2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c*f*cos(f*x + e)*sin(f*x + e))]`

3.75.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{\sqrt{c-c\sec(e+fx)}} dx = a^2 \left(\int \frac{\sec(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx + \int \frac{2\sec^2(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx + \int \frac{\sec^3(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**(1/2),x)`

output `a**2*(Integral(sec(e + f*x)/sqrt(-c*sec(e + f*x) + c), x) + Integral(2*sec(e + f*x)**2/sqrt(-c*sec(e + f*x) + c), x) + Integral(sec(e + f*x)**3/sqrt(-c*sec(e + f*x) + c), x))`

3.75.7 Maxima [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{(a\sec(fx+e)+a)^2 \sec(fx+e)}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^2*sec(f*x + e)/sqrt(-c*sec(f*x + e) + c), x)`

3.75.8 Giac [A] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.70

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{\sqrt{c-c\sec(e+fx)}} dx$$

$$= \frac{4a^2 \left(\frac{3\sqrt{2} \arctan\left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}}\right)}{\sqrt{c}} + \frac{\sqrt{2}(3c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 4c)}{(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^{\frac{3}{2}}} \right)}{3f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x, algorithm m="giac")`

output `4/3*a^2*(3*sqrt(2)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/sqrt(c) + sqrt(2)*(3*c*tan(1/2*f*x + 1/2*e)^2 - 4*c)/(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2))/f`

3.75.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^2}{\cos(e+fx) \sqrt{c - \frac{c}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)), x)`

$$3.76 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{3/2}} dx$$

3.76.1	Optimal result	591
3.76.2	Mathematica [C] (verified)	591
3.76.3	Rubi [A] (verified)	592
3.76.4	Maple [A] (verified)	594
3.76.5	Fricas [A] (verification not implemented)	595
3.76.6	Sympy [F]	595
3.76.7	Maxima [F]	596
3.76.8	Giac [A] (verification not implemented)	596
3.76.9	Mupad [F(-1)]	597

3.76.1 Optimal result

Integrand size = 34, antiderivative size = 113

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{3/2}} dx = \frac{3\sqrt{2}a^2 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{c^{3/2} f} - \frac{2a^2 \tan(e+fx)}{f(c-c \sec(e+fx))^{3/2}} - \frac{2a^2 \tan(e+fx)}{cf \sqrt{c-c \sec(e+fx)}}$$

```
output 3*a^2*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))*2^(1/2)
)/c^(3/2)/f-2*a^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(3/2)-2*a^2*tan(f*x+e)/c/f
/(c-c*sec(f*x+e))^(1/2)
```

3.76.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.42 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.57

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{3/2}} dx = \frac{a^2 \operatorname{Hypergeometric2F1}\left(2, \frac{5}{2}, \frac{7}{2}, \frac{1}{2}(1+\sec(e+fx))\right) (1+\sec(e+fx))^2 \tan(e+fx)}{10cf \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^(3/2),x]`

output `-1/10*(a^2*Hypergeometric2F1[2, 5/2, 7/2, (1 + Sec[e + f*x])/2]*(1 + Sec[e + f*x])^2*Tan[e + f*x])/(c*f*Sqrt[c - c*Sec[e + f*x]])`

3.76.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4445, 3042, 4444, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c-c\sec(e+fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{3a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{\sqrt{c-c\sec(e+fx)}} dx}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{4444} \\
 & -\frac{3a \left(2a \int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right)}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right)}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{f(c-c\sec(e+fx))^{3/2}}
 \end{aligned}$$

3.76. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{3/2}} dx$

$$\begin{aligned}
 & \downarrow 4282 \\
 & \frac{3a \left(\frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{4a \int \frac{1}{c^2 \tan^2(e+fx) + 2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{f} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{f(c-c\sec(e+fx))^{3/2}} \\
 & \downarrow 216 \\
 & - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{f(c-c\sec(e+fx))^{3/2}} - \frac{3a \left(\frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{cf}} \right)}{2c}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^(3/2),x]`

output `-(((a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2))) - (3*a*((-2*sqrt[2]*a*ArcTan[(sqrt[c]*Tan[e + f*x])/(sqrt[2]*sqrt[c - c*Sec[e + f*x]])])/(sqrt[c]*f) + (2*a*Tan[e + f*x])/(f*sqrt[c - c*Sec[e + f*x]])))/(2*c)`

3.76.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

```
rule 4444 Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.))/
Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] :> Simp[-2*d*Cot[e +
f*x]*((c + d*Csc[e + f*x])^(n - 1)/(f*(2*n - 1)*Sqrt[a + b*Csc[e + f*x]]),
x] + Simp[2*c*((2*n - 1)/(2*n - 1)) Int[Csc[e + f*x]*((c + d*Csc[e + f*x]
)^(n - 1)/Sqrt[a + b*Csc[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]
```

```
rule 4445 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m)*
(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[2*a*c*Cot[e +
f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))),
x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*
x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^
(-1)] && IntegerQ[2*m]
```

3.76.4 Maple [A] (verified)

Time = 4.84 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.06

method	result
default	$\frac{a^2\sqrt{2} \left(3 \arctan \left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} \right) \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \tan(fx+e) + 2\sqrt{2} \cot(fx+e) + \sqrt{2} \csc(fx+e) - \sqrt{2} \sec(fx+e) \csc(fx+e) \right)}{cf\sqrt{-c(\sec(fx+e)-1)}}$
parts	$-\frac{a^2\sqrt{2}(1-\cos(fx+e)) \left(\left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{3}{2}} - \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} (1-\cos(fx+e))^2 \csc(fx+e)^2 - \arccos \left(\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \right)^{\frac{3}{2}} \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right) \right)}{4f \left(\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \right)^{\frac{3}{2}} \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output a^2/c/f*2^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)*(3*arctan(1/2*2^(1/2)/(-cos(f*x+
e)/(cos(f*x+e)+1))^(1/2))*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*tan(f*x+e)+2*
2^(1/2)*cot(f*x+e)+2^(1/2)*csc(f*x+e)-2^(1/2)*sec(f*x+e)*csc(f*x+e))
```

$$3.76. \int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{3/2}} dx$$

3.76.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 372, normalized size of antiderivative = 3.29

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{3/2}} dx = \frac{3\sqrt{2}(a^2c\cos(fx+e)-a^2c)\sqrt{-\frac{1}{c}}\log\left(\frac{2\sqrt{2}(\cos(fx+e)^2+\cos(fx+e))\sqrt{\cos(fx+e)}}{\cos(fx+e)}\right) + 3\sqrt{2}(a^2c\cos(fx+e)-a^2c)\arctan\left(\frac{\sqrt{2}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\cos(fx+e)}{\sqrt{c}\sin(fx+e)}\right)\sin(fx+e)}{(c^2f\cos(fx+e)-c^2f)\sin(fx+e)} - 2(2a^2\cos(fx+e)^2+a^2\cos(fx+e)-a^2)\sqrt{\cos(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x, algorithm="fracas")`

output `[1/2*(3*sqrt(2)*(a^2*c*cos(f*x + e) - a^2*c)*sqrt(-1/c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*sqrt(-1/c) + (3*cos(f*x + e) + 1)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e))))*sin(f*x + e) + 4*(2*a^2*cos(f*x + e)^2 + a^2*cos(f*x + e) - a^2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^2*f*cos(f*x + e) - c^2*f)*sin(f*x + e)), -(3*sqrt(2)*(a^2*c*cos(f*x + e) - a^2*c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e)/sqrt(c) - 2*(2*a^2*cos(f*x + e)^2 + a^2*cos(f*x + e) - a^2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^2*f*cos(f*x + e) - c^2*f)*sin(f*x + e))]`

3.76.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{3/2}} dx = a^2 \left(\int \frac{\sec(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec(e+fx)+c}\sqrt{-c\sec(e+fx)+c}} dx + \int \frac{2\sec^2(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec(e+fx)+c}\sqrt{-c\sec(e+fx)+c}} dx + \int \frac{\sec^3(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec(e+fx)+c}\sqrt{-c\sec(e+fx)+c}} dx \right)$$

3.76. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{3/2}} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**(3/2),x)`

output `a**2*(Integral(sec(e + f*x)/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x) + Integral(2*sec(e + f*x)**2/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x) + Integral(sec(e + f*x)**3/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x))`

3.76.7 Maxima [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^{3/2}} dx = \int \frac{(a \sec(fx + e) + a)^2 \sec(fx + e)}{(-c \sec(fx + e) + c)^{3/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((a*sec(f*x + e) + a)^2*sec(f*x + e)/(-c*sec(f*x + e) + c)^(3/2), x)`

3.76.8 Giac [A] (verification not implemented)

Time = 1.23 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.96

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^{3/2}} dx = \frac{a^2 \left(\frac{3\sqrt{2} \arctan\left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}}\right)}{c^{3/2}} + \frac{\sqrt{2}(3c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)}{\left((c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c\right)^{3/2} + \sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}\right)c} \right)}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `-a^2*(3*sqrt(2)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/c^(3/2) + sqrt(2)*(3*c*tan(1/2*f*x + 1/2*e)^2 - c)/(((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2) + sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c))/f`

3.76. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{3/2}} dx$

3.76.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^2}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),x)`output `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)), x)`

$$3.77 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{5/2}} dx$$

3.77.1	Optimal result	598
3.77.2	Mathematica [A] (verified)	598
3.77.3	Rubi [A] (verified)	599
3.77.4	Maple [B] (verified)	601
3.77.5	Fricas [A] (verification not implemented)	601
3.77.6	Sympy [F]	602
3.77.7	Maxima [F]	603
3.77.8	Giac [A] (verification not implemented)	603
3.77.9	Mupad [F(-1)]	603

3.77.1 Optimal result

Integrand size = 34, antiderivative size = 117

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{5/2}} dx = -\frac{3a^2 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{4\sqrt{2}c^{5/2}f} - \frac{a^2 \tan(e+fx)}{f(c-c \sec(e+fx))^{5/2}} + \frac{5a^2 \tan(e+fx)}{4cf(c-c \sec(e+fx))^{3/2}}$$

output

```
-3/8*a^2*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/c^(5/2)/f*2^(1/2)-a^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(5/2)+5/4*a^2*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(3/2)
```

3.77.2 Mathematica [A] (verified)

Time = 1.13 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.28

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{5/2}} dx = \frac{a^{3/2} \left(\sqrt{a}(-1+4 \sec(e+fx)+5 \sec^2(e+fx)) + 6\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a(1+\sec(e+fx))}}{\sqrt{2}\sqrt{a}}\right) \sec^2(e+fx) \sqrt{a(1+\sec(e+fx))} \right)}{4c^2 f (-1+\sec(e+fx))^2 (1+\sec(e+fx)) \sqrt{c-c \sec(e+fx)}}$$

input

```
Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^(5/2),x]
```

3.77. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{5/2}} dx$

output
$$-1/4*(a^{(3/2)}*(\text{Sqrt}[a]*(-1 + 4*\text{Sec}[e + f*x] + 5*\text{Sec}[e + f*x]^2) + 6*\text{Sqrt}[2]*\text{ArcTanh}[\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]]/(\text{Sqrt}[2]*\text{Sqrt}[a]))*\text{Sec}[e + f*x]^2*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]*\text{Sin}[(e + f*x)/2]^4*\text{Tan}[e + f*x]/(c^2*f*(-1 + \text{Sec}[e + f*x])^2*(1 + \text{Sec}[e + f*x])*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]))$$

3.77.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.09, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4445, 3042, 4445, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c-c\sec(e+fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx \\ & \quad \downarrow \text{4445} \\ & -\frac{3a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^{3/2}} dx}{4c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{2f(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{3a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{4c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{2f(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow \text{4445} \\ & -\frac{3a \left(-\frac{a \int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx}{2c} - \frac{a \tan(e+fx)}{f(c-c\sec(e+fx))^{3/2}} \right)}{4c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{2f(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{3a \left(-\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2c} - \frac{a \tan(e+fx)}{f(c-c\sec(e+fx))^{3/2}} \right)}{4c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{2f(c-c\sec(e+fx))^{5/2}} \end{aligned}$$

3.77.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{5/2}} dx$$

$$\begin{array}{c}
 \downarrow 4282 \\
 3a \left(\frac{a \int \frac{1}{c^2 \tan^2(e+fx) + 2c} d \frac{c \tan(e+fx)}{\sqrt{c - c \sec(e+fx)}}}{cf} - \frac{a \tan(e+fx)}{f(c - c \sec(e+fx))^{3/2}} \right) - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{2f(c - c \sec(e+fx))^{5/2}} \\
 \hline
 \downarrow 216 \\
 - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{2f(c - c \sec(e+fx))^{5/2}} - \frac{3a \left(\frac{a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2} \sqrt{c - c \sec(e+fx)}}\right)}{\sqrt{2} c^{3/2} f} - \frac{a \tan(e+fx)}{f(c - c \sec(e+fx))^{3/2}} \right)}{4c}
 \end{array}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/2*((a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(5/2)) - (3*a*((a*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]))/(Sqrt[2]*c^(3/2)*f) - (a*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2)))/(4*c)`

3.77.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

```
rule 4445 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e +
f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))),
x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*
x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^
(-1)] && IntegerQ[2*m]
```

3.77.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(100) = 200.

Time = 4.58 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.95

method	result
default	$\frac{a^2\sqrt{2} \left(3 \arctan \left(\frac{1}{\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} \right) (1-\cos(fx+e))^4 \csc(fx+e) - 3\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} (1-\cos(fx+e))^2 \right)}{8c^2 f \sqrt{\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} (1-\cos(fx+e))^3 \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}}$
parts	Expression too large to display

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/8*a^2/c^2/f*2^(1/2)/(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e)^2-1
)*csc(f*x+e)^2)^(1/2)/(1-cos(f*x+e))^3/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(
1/2)*(3*arctan(1/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2))*(1-cos(f*x+e))^4
*csc(f*x+e)-3*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(1-cos(f*x+e))^2*sin
(f*x+e)-2*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*sin(f*x+e)^3)
```

3.77.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 429, normalized size of antiderivative = 3.67

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{5/2}} dx = \left[\frac{3\sqrt{2}(a^2 \cos(fx+e)^2 - 2a^2 \cos(fx+e) + a^2)\sqrt{-c} \log\left(\frac{2\sqrt{2}(c \dots}{\dots}\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x, algorithm="fricas")`

output `[-1/16*(3*sqrt(2)*(a^2*cos(f*x + e)^2 - 2*a^2*cos(f*x + e) + a^2)*sqrt(-c) *log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e))) *sin(f*x + e) + 4*(a^2*cos(f*x + e)^3 - 4*a^2*cos(f*x + e)^2 - 5*a^2*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x + e)), 1/8*(3 *sqrt(2)*(a^2*cos(f*x + e)^2 - 2*a^2*cos(f*x + e) + a^2)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(a^2*cos(f*x + e)^3 - 4*a^2*cos(f*x + e)^2 - 5*a^2*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x + e))]`

3.77.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{5/2}} dx = a^2 \left(\int \frac{\sec(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}+c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}{2\sec^2(e+fx)} \right. \\ \left. + \int \frac{\sec^3(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}+c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}} \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**(5/2),x)`

output `a**2*(Integral(sec(e + f*x)/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f*x) + c)), x) + Integral(2*sec(e + f*x)**2/(c**2*sqrt(-c*sec(e + f*x) + c) *sec(e + f*x)**2 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f*x) + c)), x) + Integral(sec(e + f*x)**3/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f*x) + c)), x))`

3.77.7 Maxima [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{(a\sec(fx+e)+a)^2 \sec(fx+e)}{(-c\sec(fx+e)+c)^{5/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^2*sec(f*x + e)/(-c*sec(f*x + e) + c)^(5/2), x)`

3.77.8 Giac [A] (verification not implemented)

Time = 1.30 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{5/2}} dx = \frac{\sqrt{2} \left(3\sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}} \right) + \frac{3(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^{\frac{3}{2}} c + 5\sqrt{c}}{c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)} \right)}{8c^3 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x, algorithm m="giac")`

output `1/8*sqrt(2)*(3*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) + (3*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c + 5*sqrt(c*tan(1/2*f*x + 1/2*e))^2 - c)*c^2)/(c^2*tan(1/2*f*x + 1/2*e)^4)*a^2/(c^3*f)`

3.77.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^2}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x)`

output `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)), x)`

3.77. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{5/2}} dx$

3.78 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{7/2}} dx$

3.78.1 Optimal result 604
 3.78.2 Mathematica [C] (verified) 604
 3.78.3 Rubi [A] (verified) 605
 3.78.4 Maple [B] (verified) 608
 3.78.5 Fricas [A] (verification not implemented) 609
 3.78.6 Sympy [F] 609
 3.78.7 Maxima [F(-1)] 610
 3.78.8 Giac [A] (verification not implemented) 610
 3.78.9 Mupad [F(-1)] 611

3.78.1 Optimal result

Integrand size = 34, antiderivative size = 164

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{7/2}} dx =$$

$$-\frac{a^2 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{16\sqrt{2}c^{7/2}f} - \frac{(a^2 + a^2 \sec(e+fx)) \tan(e+fx)}{3f(c-c \sec(e+fx))^{7/2}}$$

$$+ \frac{a^2 \tan(e+fx)}{4cf(c-c \sec(e+fx))^{5/2}} - \frac{a^2 \tan(e+fx)}{16c^2 f(c-c \sec(e+fx))^{3/2}}$$

output

```
-1/32*a^2*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/c^(7/2)/f*2^(1/2)-1/3*(a^2+a^2*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^(7/2)+1/4*a^2*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(5/2)-1/16*a^2*tan(f*x+e)/c^2/f/(c-c*sec(f*x+e))^(3/2)
```

3.78.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.61 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.39

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{7/2}} dx =$$

$$\frac{a^2 \text{Hypergeometric2F1}\left(\frac{5}{2}, 4, \frac{7}{2}, \frac{1}{2}(1 + \sec(e+fx))\right) (1 + \sec(e+fx))^2 \tan(e+fx)}{40c^3 f \sqrt{c-c \sec(e+fx)}}$$

3.78. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{7/2}} dx$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^(7/2),x]`

output `-1/40*(a^2*Hypergeometric2F1[5/2, 4, 7/2, (1 + Sec[e + f*x])/2]*(1 + Sec[e + f*x])^2*Tan[e + f*x])/(c^3*f*Sqrt[c - c*Sec[e + f*x]])`

3.78.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4445, 3042, 4445, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c-c\sec(e+fx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{(c-c\sec(e+fx))^{5/2}} dx}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{4445} \\
 & -\frac{a \left(-\frac{a \int \frac{\sec(e+fx)}{(c-c\sec(e+fx))^{3/2}} dx}{4c} - \frac{a \tan(e+fx)}{2f(c-c\sec(e+fx))^{5/2}} \right)}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{3f(c-c\sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.78. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{7/2}} dx$

$$\begin{aligned}
 & \frac{a \left(\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c \csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{4c} - \frac{a \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f(c-c \sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{4283} \\
 & \frac{a \left(\frac{a \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c-c \sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f(c-c \sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \left(\frac{a \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f(c-c \sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{4282} \\
 & \frac{a \left(\frac{a \left(\frac{\int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c \sec(e+fx)} + 2c} dx - \frac{d-c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}}}{2cf} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f(c-c \sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{216}
 \end{aligned}$$

3.78. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{7/2}} dx$

$$\frac{\tan(e+fx)(a^2 \sec(e+fx) + a^2)}{3f(c - c \sec(e+fx))^{7/2}} - \frac{a \left(\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right) - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}}}{4c} - \frac{a \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}} \right)}{2c}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c - c*Sec[e + f*x])^(7/2),x]`

output `-1/3*((a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(7/2)) - (a*(-1/2*(a*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(5/2)) - (a*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2))))/(4*c))/(2*c)`

3.78.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4283 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`


```
rule 4445 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]
```

3.78.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 319 vs. 2(141) = 282.

Time = 5.49 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.95

method	result
default	$\frac{a^2\sqrt{2} \left(3 \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{3}{2}} (1-\cos(fx+e))^4 \sin(fx+e) - 3 \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} (1-\cos(fx+e))^6 \csc(fx+e) \right)}{96c^3 f \sqrt{\dots}}$
parts	Expression too large to display

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(7/2),x,method=_RETURNV ERBOSE)
```

```
output -1/96*a^2/c^3/f*2^(1/2)/(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*csc(f*x+e)^2)^(1/2)/(1-cos(f*x+e))^5/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(3*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(3/2)*(1-cos(f*x+e))^4*sin(f*x+e)-3*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(1-cos(f*x+e))^6*csc(f*x+e)-3*arctan(1/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2))*(1-cos(f*x+e))^6*csc(f*x+e)+6*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(3/2)*(1-cos(f*x+e))^2*sin(f*x+e))^3+8*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(3/2)*sin(f*x+e)^5)
```

$$3.78. \int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{7/2}} dx$$

3.78.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 517, normalized size of antiderivative = 3.15

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{7/2}} dx = \left[\frac{3\sqrt{2}(a^2 \cos(fx+e)^3 - 3a^2 \cos(fx+e)^2 + 3a^2 \cos(fx+e) - a^2) \sqrt{-c} \log((2\sqrt{2})(\cos(fx+e)^2 + \cos(fx+e)) \sqrt{-c} \sqrt{(c\cos(fx+e) - c)/\cos(fx+e)} + (3c\cos(fx+e) + c) \sin(fx+e)) / ((\cos(fx+e) - 1) \sin(fx+e)) \sin(fx+e) - 4(7a^2 \cos(fx+e)^4 + 29a^2 \cos(fx+e)^3 + 25a^2 \cos(fx+e)^2 + 3a^2 \cos(fx+e)) \sqrt{(c\cos(fx+e) - c)/\cos(fx+e)}) / ((c^4 f \cos(fx+e)^3 - 3c^4 f \cos(fx+e)^2 + 3c^4 f \cos(fx+e) - c^4 f) \sin(fx+e))}{1/96 * (3\sqrt{2})(a^2 \cos(fx+e)^3 - 3a^2 \cos(fx+e)^2 + 3a^2 \cos(fx+e) - a^2) \sqrt{c} \arctan(\sqrt{2} \sqrt{(c\cos(fx+e) - c)/\cos(fx+e)}) \cos(fx+e) / (\sqrt{c} \sin(fx+e)) \sin(fx+e) + 2(7a^2 \cos(fx+e)^4 + 29a^2 \cos(fx+e)^3 + 25a^2 \cos(fx+e)^2 + 3a^2 \cos(fx+e)) \sqrt{(c\cos(fx+e) - c)/\cos(fx+e)}) / ((c^4 f \cos(fx+e)^3 - 3c^4 f \cos(fx+e)^2 + 3c^4 f \cos(fx+e) - c^4 f) \sin(fx+e))} \right]$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(7/2),x, algorithm="fracas")
```

```
output [-1/192*(3*sqrt(2)*(a^2*cos(f*x + e)^3 - 3*a^2*cos(f*x + e)^2 + 3*a^2*cos(f*x + e) - a^2)*sqrt(-c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e))*sin(f*x + e) - 4*(7*a^2*cos(f*x + e)^4 + 29*a^2*cos(f*x + e)^3 + 25*a^2*cos(f*x + e)^2 + 3*a^2*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c^4*f*cos(f*x + e)^3 - 3*c^4*f*cos(f*x + e)^2 + 3*c^4*f*cos(f*x + e) - c^4*f)*sin(f*x + e)), 1/96*(3*sqrt(2)*(a^2*cos(f*x + e)^3 - 3*a^2*cos(f*x + e)^2 + 3*a^2*cos(f*x + e) - a^2)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e))*sin(f*x + e) + 2*(7*a^2*cos(f*x + e)^4 + 29*a^2*cos(f*x + e)^3 + 25*a^2*cos(f*x + e)^2 + 3*a^2*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c^4*f*cos(f*x + e)^3 - 3*c^4*f*cos(f*x + e)^2 + 3*c^4*f*cos(f*x + e) - c^4*f)*sin(f*x + e)]]
```

3.78.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{7/2}} dx = a^2 \left(\int \frac{1}{-c^3 \sqrt{-c\sec(e+fx)} + c\sec^3(e+fx) + 3c^3 \sqrt{-c\sec(e+fx)}} dx \right. \\ \left. + \int \frac{2\sec^2(e+fx)}{-c^3 \sqrt{-c\sec(e+fx)} + c\sec^3(e+fx) + 3c^3 \sqrt{-c\sec(e+fx)} + c\sec^2(e+fx) - 3c^3 \sqrt{-c\sec(e+fx)}} dx \right. \\ \left. + \int \frac{\sec^3(e+fx)}{-c^3 \sqrt{-c\sec(e+fx)} + c\sec^3(e+fx) + 3c^3 \sqrt{-c\sec(e+fx)} + c\sec^2(e+fx) - 3c^3 \sqrt{-c\sec(e+fx)}} dx \right)$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**(7/2),x)
```

output `a**2*(Integral(sec(e + f*x)/(-c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3 + 3*c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 - 3*c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**3*sqrt(-c*sec(e + f*x) + c)), x) + Integral(2*sec(e + f*x)**2/(-c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3 + 3*c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 - 3*c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**3*sqrt(-c*sec(e + f*x) + c)), x) + Integral(sec(e + f*x)**3/(-c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3 + 3*c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 - 3*c**3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**3*sqrt(-c*sec(e + f*x) + c)), x))`

3.78.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^{7/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(7/2),x, algorithm="maxima")`

output Timed out

3.78.8 Giac [A] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.85

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c - c \sec(e + fx))^{7/2}} dx = \frac{\sqrt{2} \left(3 a^2 \sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}} \right) + \frac{3 \left(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c \right)^{\frac{5}{2}} a^2 c}{96 c^4 f} \right)}{96 c^4 f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(7/2),x, algorithm="giac")`

output `1/96*sqrt(2)*(3*a^2*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) + (3*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2)*a^2*c + 8*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*a^2*c^2 - 3*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*a^2*c^3)/(c^3*tan(1/2*f*x + 1/2*e)^6)/(c^4*f)`

3.78. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c-c \sec(e+fx))^{7/2}} dx$

3.78.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c-c\sec(e+fx))^{7/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^2}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{7/2}} dx$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(7/2)),x)`

output `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c - c/cos(e + f*x))^(7/2)), x)`

3.79 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx$

3.79.1	Optimal result	612
3.79.2	Mathematica [A] (verified)	613
3.79.3	Rubi [A] (verified)	613
3.79.4	Maple [A] (verified)	615
3.79.5	Fricas [A] (verification not implemented)	616
3.79.6	Sympy [F(-1)]	616
3.79.7	Maxima [F(-1)]	617
3.79.8	Giac [A] (verification not implemented)	617
3.79.9	Mupad [B] (verification not implemented)	618

3.79.1 Optimal result

Integrand size = 34, antiderivative size = 171

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx =$$

$$\frac{256c^4(a + a \sec(e + fx))^3 \tan(e + fx)}{3003f \sqrt{c - c \sec(e + fx)}} - \frac{64c^3(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{429f} - \frac{24c^2(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} \tan(e + fx)}{143f} - \frac{2c(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{5/2} \tan(e + fx)}{13f}$$

```
output -24/143*c^2*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f-2/13*c*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f-256/3003*c^4*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-64/429*c^3*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.79.2 Mathematica [A] (verified)

Time = 3.77 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.51

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx = \frac{4a^3c^3 \cos^6\left(\frac{1}{2}(e + fx)\right) (-3766 + 6285 \cos(e + fx) - 2842 \cos(2(e + fx)) + 835 \cos(3(e + fx)))}{3003f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(7/2),x]`

output `(4*a^3*c^3*Cos[(e + f*x)/2]^6*(-3766 + 6285*Cos[e + f*x] - 2842*Cos[2*(e + f*x)] + 835*Cos[3*(e + f*x)])*Cot[(e + f*x)/2]*Sec[e + f*x]^6*Sqrt[c - c*Sec[e + f*x]]/(3003*f)`

3.79.3 Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4443, 3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{7/2} dx \\ & \quad \downarrow \text{4443} \\ & \frac{12}{13}c \int \sec(e + fx)(\sec(e + fx)a + a)^3(c - c \sec(e + fx))^{5/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^{5/2}}{13f} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{12}{13}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^3 \left(c - c\csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{5/2}}{13f}$$

↓ 4443

$$\frac{12}{13}c \left(\frac{8}{11}c \int \sec(e + fx)(\sec(e + fx)a + a)^3 (c - c \sec(e + fx))^{3/2} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{3/2}}{11f} \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{5/2}}{13f}$$

↓ 3042

$$\frac{12}{13}c \left(\frac{8}{11}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^3 \left(c - c\csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{3/2}}{11f} \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{5/2}}{13f}$$

↓ 4443

$$\frac{12}{13}c \left(\frac{8}{11}c \left(\frac{4}{9}c \int \sec(e + fx)(\sec(e + fx)a + a)^3 \sqrt{c - c \sec(e + fx)} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 \sqrt{c - c \sec(e + fx)}}{9f} \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{5/2}}{13f} \right)$$

↓ 3042

$$\frac{12}{13}c \left(\frac{8}{11}c \left(\frac{4}{9}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^3 \sqrt{c - c\csc\left(e + fx + \frac{\pi}{2}\right)} dx - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 \sqrt{c - c \sec(e + fx)}}{9f} \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{5/2}}{13f} \right)$$

↓ 4441

$$\frac{12}{13}c \left(\frac{8}{11}c \left(-\frac{8c^2 \tan(e + fx)(a \sec(e + fx) + a)^3}{63f \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 \sqrt{c - c \sec(e + fx)}}{9f} \right) - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 (c - c \sec(e + fx))^{5/2}}{13f} \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(7/2),x]`

```
output (-2*c*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(13*f) + (12*c*((-2*c*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(11*f) + (8*c*((-8*c^2*(a + a*Sec[e + f*x])^3*Tan[e + f*x])/(63*f* Sqrt[c - c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x])*Tan[e + f*x])/(9*f)))/11)/13
```

3.79.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.79.4 Maple [A] (verified)

Time = 41.03 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.46

method	result
default	$\frac{2c^3a^3 \left(835 \cos(fx+e)^3 - 1421 \cos(fx+e)^2 + 945 \cos(fx+e) - 231 \right) \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1)^4 \sec(fx+e)^6 \csc(fx+e)}{3003f}$
parts	$-\frac{2a^3 (\sec(fx+e)-1)^3 \left(177 \cos(fx+e)^3 - 71 \cos(fx+e)^2 + 27 \cos(fx+e) - 5 \right) c^3 \sqrt{-c(\sec(fx+e)-1)} (\cos(fx+e)+1) \csc(fx+e)}{35f(\cos(fx+e)-1)^3} + \dots$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(7/2),x,method=_RETURNV ERBOSE)
```

$$3.79. \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx$$

output $\frac{2}{3003}c^3a^3/f*(835*\cos(f*x+e)^3-1421*\cos(f*x+e)^2+945*\cos(f*x+e)-231)*(-c*(\sec(f*x+e)-1))^{(1/2)}*(\cos(f*x+e)+1)^4*\sec(f*x+e)^6*\csc(f*x+e)$

3.79.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.95

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx = \frac{2(835 a^3 c^3 \cos(fx + e)^7 + 1919 a^3 c^3 \cos(fx + e)^6 + 271 a^3 c^3 \cos(fx + e)^5 - 1637 a^3 c^3 \cos(fx + e)^4 - 103 a^3 c^3 \cos(fx + e)^3 + 973 a^3 c^3 \cos(fx + e)^2 + 21 a^3 c^3 \cos(fx + e) - 231 a^3 c^3) \sqrt{(c \cos(fx + e) - c) / \cos(fx + e)}}{3003 f \cos(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(7/2),x, algorithm="fricas")`

output $\frac{2}{3003}*(835*a^3*c^3*\cos(f*x + e)^7 + 1919*a^3*c^3*\cos(f*x + e)^6 + 271*a^3*c^3*\cos(f*x + e)^5 - 1637*a^3*c^3*\cos(f*x + e)^4 - 103*a^3*c^3*\cos(f*x + e)^3 + 973*a^3*c^3*\cos(f*x + e)^2 + 21*a^3*c^3*\cos(f*x + e) - 231*a^3*c^3)*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^6*\sin(f*x + e))$

3.79.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**(7/2),x)`

output `Timed out`

3.79.7 Maxima [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(7/2),x, algorithm
m="maxima")`

output `Timed out`

3.79.8 Giac [A] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.64

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{7/2} dx = \frac{128 \sqrt{2} \left(429 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right)^3 c^4 + 1001 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right)^2 c^5 + 819 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right) c^6 + 231 c^7 \right) a^3 c^3}{3003 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right)^{\frac{13}{2}} f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(7/2),x, algorithm
m="giac")`

output `128/3003*sqrt(2)*(429*(c*tan(1/2*f*x + 1/2*e)^2 - c)^3*c^4 + 1001*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^5 + 819*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^6 + 231*c^7)*a^3*c^3/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(13/2)*f)`

3.79.9 Mupad [B] (verification not implemented)

Time = 27.14 (sec) , antiderivative size = 710, normalized size of antiderivative = 4.15

$$\begin{aligned}
& \int \sec(e + fx)(a + a \sec(e + fx))^3(c \\
& - c \sec(e + fx))^{7/2} dx = \frac{\left(\frac{a^3 c^3 2i}{f} + \frac{a^3 c^3 e^{e li + f x li} 1670i}{3003 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}}{e^{e li + f x li} - 1} \\
& + \frac{\left(\frac{a^3 c^3 128i}{13 f} + \frac{a^3 c^3 e^{e li + f x li} 128i}{13 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^6} \\
& - \frac{\left(\frac{a^3 c^3 384i}{11 f} + \frac{a^3 c^3 e^{e li + f x li} 3456i}{143 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^5} \\
& - \frac{\left(\frac{a^3 c^3 8i}{f} + \frac{a^3 c^3 e^{e li + f x li} 2168i}{3003 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)} \\
& + \frac{\left(\frac{a^3 c^3 24i}{f} + \frac{a^3 c^3 e^{e li + f x li} 5464i}{1001 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^2} \\
& + \frac{\left(\frac{a^3 c^3 160i}{3 f} + \frac{a^3 c^3 e^{e li + f x li} 11360i}{429 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^4} \\
& - \frac{\left(\frac{a^3 c^3 320i}{7 f} + \frac{a^3 c^3 e^{e li + f x li} 46400i}{3003 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e li - f x li}}{2} + \frac{e^{e li + f x li}}{2}}}}{(e^{e li + f x li} - 1) (e^{e 2i + f x 2i} + 1)^3}
\end{aligned}$$

```
input int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(7/2))/cos(e + f*x),x)
```

output

$$\begin{aligned}
& \left(\frac{a^3 c^3 e^{2i}}{f} + \frac{a^3 c^3 \exp(e^{1i} + f x^{1i}) 1670i}{3003 f} \right) \left(c - \frac{c}{\exp(-e^{1i} - f x^{1i})/2 + \exp(e^{1i} + f x^{1i})/2} \right)^{1/2} \left(\exp(e^{1i} + f x^{1i}) - 1 \right) \\
& + \left(\frac{a^3 c^3 128i}{13 f} + \frac{a^3 c^3 \exp(e^{1i} + f x^{1i}) 128i}{13 f} \right) \left(c - \frac{c}{\exp(-e^{1i} - f x^{1i})/2 + \exp(e^{1i} + f x^{1i})/2} \right)^{1/2} \left(\frac{\exp(e^{1i} + f x^{1i})}{2} \right)^{1/2} \\
& \left(\exp(e^{1i} + f x^{1i}) - 1 \right) \left(\exp(e^{2i} + f x^{2i}) + 1 \right)^6 - \left(\frac{a^3 c^3 384i}{11 f} + \frac{a^3 c^3 \exp(e^{1i} + f x^{1i}) 3456i}{143 f} \right) \left(c - \frac{c}{\exp(-e^{1i} - f x^{1i})/2 + \exp(e^{1i} + f x^{1i})/2} \right)^{1/2} \\
& \left(\exp(e^{1i} + f x^{1i}) - 1 \right) \left(\exp(e^{2i} + f x^{2i}) + 1 \right)^5 - \left(\frac{a^3 c^3 8i}{f} + \frac{a^3 c^3 \exp(e^{1i} + f x^{1i}) 2168i}{3003 f} \right) \left(c - \frac{c}{\exp(-e^{1i} - f x^{1i})/2 + \exp(e^{1i} + f x^{1i})/2} \right)^{1/2} \\
& \left(\frac{\exp(e^{1i} + f x^{1i})}{2} \right)^{1/2} \left(\exp(e^{1i} + f x^{1i}) - 1 \right) \left(\exp(e^{2i} + f x^{2i}) + 1 \right) + \left(\frac{a^3 c^3 24i}{f} + \frac{a^3 c^3 \exp(e^{1i} + f x^{1i}) 5464i}{1001 f} \right) \left(c - \frac{c}{\exp(-e^{1i} - f x^{1i})/2 + \exp(e^{1i} + f x^{1i})/2} \right)^{1/2} \\
& \left(\frac{\exp(e^{1i} + f x^{1i})}{2} \right)^{1/2} \left(\exp(e^{1i} + f x^{1i}) - 1 \right) \left(\exp(e^{2i} + f x^{2i}) + 1 \right)^2 + \left(\frac{a^3 c^3 160i}{3 f} + \frac{a^3 c^3 \exp(e^{1i} + f x^{1i}) 11360i}{429 f} \right) \left(c - \frac{c}{\exp(-e^{1i} - f x^{1i})/2 + \exp(e^{1i} + f x^{1i})/2} \right)^{1/2} \\
& \left(\frac{\exp(e^{1i} + f x^{1i})}{2} \right)^{1/2} \left(\exp(e^{1i} + f x^{1i}) - 1 \right) \left(\exp(e^{2i} + f x^{2i}) + 1 \right)^4 - \left(\frac{a^3 c^3 320i}{7 f} + \frac{a^3 c^3 \exp(e^{1i} + f x^{1i}) 46400i}{3003 f} \right) \left(c - \frac{c}{\exp(-e^{1i} - f x^{1i})/2 + \exp(e^{1i} + f x^{1i})/2} \right)^{1/2} \\
& \left(\frac{\exp(e^{1i} + f x^{1i})}{2} \right)^{1/2} \left(\exp(e^{1i} + f x^{1i}) - 1 \right) \left(\exp(e^{2i} + f x^{2i}) + 1 \right)^3
\end{aligned}$$

3.80 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{5/2} dx$

3.80.1	Optimal result	620
3.80.2	Mathematica [A] (verified)	621
3.80.3	Rubi [A] (verified)	621
3.80.4	Maple [A] (verified)	623
3.80.5	Fricas [A] (verification not implemented)	623
3.80.6	Sympy [F]	624
3.80.7	Maxima [F(-1)]	625
3.80.8	Giac [A] (verification not implemented)	625
3.80.9	Mupad [B] (verification not implemented)	626

3.80.1 Optimal result

Integrand size = 34, antiderivative size = 128

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{5/2} dx =$$

$$-\frac{64c^3(a + a \sec(e + fx))^3 \tan(e + fx)}{693f \sqrt{c - c \sec(e + fx)}}$$

$$-\frac{16c^2(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{99f}$$

$$-\frac{2c(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} \tan(e + fx)}{11f}$$

output

```
-2/11*c*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f-64/693*c^3*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-16/99*c^2*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.80.2 Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.61

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{5/2} dx = \frac{8a^3c^2 \cos^6\left(\frac{1}{2}(e + fx)\right) (277 - 364 \cos(e + fx) + 151 \cos(2(e + fx))) \cot\left(\frac{1}{2}(e + fx)\right) \sec^5(e + fx)}{693f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(5/2),x]`

output `(8*a^3*c^2*Cos[(e + f*x)/2]^6*(277 - 364*Cos[e + f*x] + 151*Cos[2*(e + f*x)])*Cot[(e + f*x)/2]*Sec[e + f*x]^5*Sqrt[c - c*Sec[e + f*x]]/(693*f)`

3.80.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx \\ & \quad \downarrow \text{4443} \\ & \frac{8}{11}c \int \sec(e + fx)(\sec(e + fx)a + a)^3(c - c \sec(e + fx))^{3/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^{3/2}}{11f} \\ & \quad \downarrow \text{3042} \\ & \frac{8}{11}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right) a + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx - \\ & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^{3/2}}{11f} \end{aligned}$$

3.80. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{5/2} dx$

↓ 4443

$$\frac{8}{11}c \left(\frac{4}{9}c \int \sec(e+fx)(\sec(e+fx)a+a)^3 \sqrt{c-c\sec(e+fx)} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{9f} \right. \\ \left. - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^3 (c-c\sec(e+fx))^{3/2}}{11f} \right)$$

↓ 3042

$$\frac{8}{11}c \left(\frac{4}{9}c \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)^3 \sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)} dx - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{9f} \right. \\ \left. - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^3 (c-c\sec(e+fx))^{3/2}}{11f} \right)$$

↓ 4441

$$\frac{8}{11}c \left(-\frac{8c^2 \tan(e+fx)(a \sec(e+fx)+a)^3}{63f \sqrt{c-c\sec(e+fx)}} - \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{9f} \right) - \\ \frac{2c \tan(e+fx)(a \sec(e+fx)+a)^3 (c-c\sec(e+fx))^{3/2}}{11f}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(5/2),x]`

output `(-2*c*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(11*f) + (8*c*((-8*c^2*(a + a*Sec[e + f*x])^3*Tan[e + f*x])/(63*f*Sqrt[c - c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(9*f)))/11`

3.80.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.80. $\int \sec(e+fx)(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2} dx$

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.80.4 Maple [A] (verified)

Time = 38.80 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.54

method	result
default	$\frac{2c^2a^3(151\cos(fx+e)^2-182\cos(fx+e)+63)\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)^4\sec(fx+e)^5\csc(fx+e)}{693f}$
parts	$\frac{2a^3(\sec(fx+e)-1)^2(43\cos(fx+e)^2-14\cos(fx+e)+3)c^2\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)\csc(fx+e)}{15f(\cos(fx+e)-1)^2} - \frac{2a^3(1136\cos(fx+e)^5}{\dots}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 2/693*c^2*a^3/f*(151*cos(f*x+e)^2-182*cos(f*x+e)+63)*(-c*(sec(f*x+e)-1))^(
1/2)*(cos(f*x+e)+1)^4*sec(f*x+e)^5*csc(f*x+e)
```

3.80.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.15

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx = \frac{2(151a^3c^2 \cos(fx + e)^6 + 422a^3c^2 \cos(fx + e)^5 + 241a^3c^2 \cos(fx + e)^4 - 236a^3c^2 \cos(fx + e)^3 + 1136a^3c^2 \cos(fx + e)^2 - 1136a^3c^2 \cos(fx + e) + 1136a^3c^2)}{693f \cos(fx + e)^5 \sin(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(5/2),x, algorithm
m="fracas")
```

3.80. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^5 dx$

output $2/693*(151*a^3*c^2*\cos(f*x + e)^6 + 422*a^3*c^2*\cos(f*x + e)^5 + 241*a^3*c^2*\cos(f*x + e)^4 - 236*a^3*c^2*\cos(f*x + e)^3 - 199*a^3*c^2*\cos(f*x + e)^2 + 70*a^3*c^2*\cos(f*x + e) + 63*a^3*c^2)*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^5*\sin(f*x + e))$

3.80.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c \\ & - c \sec(e + fx))^{5/2} dx = a^3 \left(\int c^2 \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx \right. \\ & + \int c^2 \sqrt{-c \sec(e + fx) + c} \sec^2(e + fx) dx \\ & + \int (-2c^2 \sqrt{-c \sec(e + fx) + c} \sec^3(e + fx)) dx \\ & + \int (-2c^2 \sqrt{-c \sec(e + fx) + c} \sec^4(e + fx)) dx \\ & + \int c^2 \sqrt{-c \sec(e + fx) + c} \sec^5(e + fx) dx \\ & \left. + \int c^2 \sqrt{-c \sec(e + fx) + c} \sec^6(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**(5/2),x)`

output `a**3*(Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2, x) + Integral(-2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3, x) + Integral(-2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4, x) + Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**5, x) + Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**6, x))`

3.80.7 Maxima [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{5/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(5/2),x, algorithm
m="maxima")`

output `Timed out`

3.80.8 Giac [A] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.66

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{5/2} dx = \frac{64\sqrt{2}\left(99\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^2 c^4 + 154\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)c^5 + 63c^6\right)a^3 c^2}{693\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{11}{2}} f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(5/2),x, algorithm
m="giac")`

output `64/693*sqrt(2)*(99*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^4 + 154*(c*tan(1/2*f
*x + 1/2*e)^2 - c)*c^5 + 63*c^6)*a^3*c^2/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(
11/2)*f)`

3.80.9 Mupad [B] (verification not implemented)

Time = 25.00 (sec) , antiderivative size = 607, normalized size of antiderivative = 4.74

$$\int \sec(e + fx)(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^{5/2} dx = \frac{\left(\frac{a^3 c^2 2i}{f} + \frac{a^3 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 302i}{693 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}}{e^{e \operatorname{li} + f x \operatorname{li}} - 1} - \frac{\left(\frac{a^3 c^2 64i}{11 f} - \frac{a^3 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 64i}{11 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)^5} + \frac{\left(\frac{a^3 c^2 16i}{f} - \frac{a^3 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 944i}{231 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)^2} + \frac{\left(\frac{a^3 c^2 160i}{9 f} - \frac{a^3 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 1120i}{99 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)^4} - \frac{\left(\frac{a^3 c^2 20i}{3 f} - \frac{a^3 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 844i}{693 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)} - \frac{\left(\frac{a^3 c^2 160i}{7 f} - \frac{a^3 c^2 e^{e \operatorname{li} + f x \operatorname{li}} 6880i}{693 f}\right) \sqrt{c - \frac{c}{\frac{e^{-e \operatorname{li} - f x \operatorname{li}}}{2} + \frac{e^{e \operatorname{li} + f x \operatorname{li}}}{2}}}}{(e^{e \operatorname{li} + f x \operatorname{li}} - 1) (e^{e 2i + f x 2i} + 1)^3}$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x),x)`

output `((a^3*c^2*2i)/f + (a^3*c^2*exp(e*1i + f*x*1i)*302i)/(693*f))*(c - c/(exp(-e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/(exp(e*1i + f*x*1i) - 1) - (((a^3*c^2*64i)/(11*f) - (a^3*c^2*exp(e*1i + f*x*1i)*64i)/(11*f))*(c - c/(exp(-e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^5) + (((a^3*c^2*16i)/f - (a^3*c^2*exp(e*1i + f*x*1i)*944i)/(231*f))*(c - c/(exp(-e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^2) + (((a^3*c^2*160i)/(9*f) - (a^3*c^2*exp(e*1i + f*x*1i)*1120i)/(99*f))*(c - c/(exp(-e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^4) - (((a^3*c^2*20i)/(3*f) - (a^3*c^2*exp(e*1i + f*x*1i)*844i)/(693*f))*(c - c/(exp(-e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)) - (((a^3*c^2*160i)/(7*f) - (a^3*c^2*exp(e*1i + f*x*1i)*6880i)/(693*f))*(c - c/(exp(-e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^3)`

3.80. $\int \sec(e + fx)(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^{5/2} dx$

3.81 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx$

3.81.1	Optimal result	627
3.81.2	Mathematica [A] (verified)	627
3.81.3	Rubi [A] (verified)	628
3.81.4	Maple [A] (verified)	629
3.81.5	Fricas [A] (verification not implemented)	630
3.81.6	Sympy [F]	630
3.81.7	Maxima [F(-1)]	631
3.81.8	Giac [A] (verification not implemented)	631
3.81.9	Mupad [B] (verification not implemented)	632

3.81.1 Optimal result

Integrand size = 34, antiderivative size = 85

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx =$$

$$-\frac{8c^2(a + a \sec(e + fx))^3 \tan(e + fx)}{63f \sqrt{c - c \sec(e + fx)}} - \frac{2c(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} \tan(e + fx)}{9f}$$

output `-8/63*c^2*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-2/9*c*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f`

3.81.2 Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.78

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx = \frac{16a^3 c \cos^6\left(\frac{1}{2}(e + fx)\right) (-7 + 11 \cos(e + fx)) \cot\left(\frac{1}{2}(e + fx)\right) \sec^4(e + fx) \sqrt{c - c \sec(e + fx)}}{63f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(3/2),x]`

output `(16*a^3*c*Cos[(e + f*x)/2]^6*(-7 + 11*Cos[e + f*x])*Cot[(e + f*x)/2]*Sec[e + f*x]^4*sqrt[c - c*Sec[e + f*x]])/(63*f)`

3.81.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{4}{9}c \int \sec(e + fx)(\sec(e + fx)a + a)^3 \sqrt{c - c \sec(e + fx)} dx - \\
 & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 \sqrt{c - c \sec(e + fx)}}{9f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4}{9}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right) a + a\right)^3 \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx - \\
 & \quad \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 \sqrt{c - c \sec(e + fx)}}{9f} \\
 & \quad \downarrow \text{4441} \\
 & -\frac{8c^2 \tan(e + fx)(a \sec(e + fx) + a)^3}{63f \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3 \sqrt{c - c \sec(e + fx)}}{9f}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(3/2),x]`

```
output (-8*c^2*(a + a*Sec[e + f*x])^3*Tan[e + f*x]/(63*f*Sqrt[c - c*Sec[e + f*x]
]) - (2*c*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x]/(9
*f)
```

3.81.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sq
rt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f
*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /
; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
&& NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.81.4 Maple [A] (verified)

Time = 6.83 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

method	result
default	$\frac{2a^3c(11\cos(fx+e)-7)\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)^4\sec(fx+e)^4\csc(fx+e)}{63f}$
parts	$-\frac{2a^3(\sec(fx+e)-1)(5\cos(fx+e)-1)c\sqrt{-c(\sec(fx+e)-1)}(\cos(fx+e)+1)\csc(fx+e)}{3f(\cos(fx+e)-1)} + \frac{2a^3(\sec(fx+e)-1)(272\cos(fx+e)^4-13c^2)}{3f(\cos(fx+e)-1)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(3/2),x,method=_RETURNV
ERBOSE)
```

$$3.81. \quad \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx$$

output $2/63*a^3*c/f*(11*\cos(f*x+e)-7)*(-c*(\sec(f*x+e)-1))^{(1/2)}*(\cos(f*x+e)+1)^4*\sec(f*x+e)^4*csc(f*x+e)$

3.81.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.40

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx = \frac{2(11a^3c \cos(fx + e)^5 + 37a^3c \cos(fx + e)^4 + 38a^3c \cos(fx + e)^3 + 2a^3c \cos(fx + e)^2 - 17a^3c \cos(fx + e) - 7a^3c) \sqrt{c \cos(fx + e) - c}}{63f \cos(fx + e)^4 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(3/2),x, algorithm="fricas")`

output $2/63*(11*a^3*c*\cos(f*x + e)^5 + 37*a^3*c*\cos(f*x + e)^4 + 38*a^3*c*\cos(f*x + e)^3 + 2*a^3*c*\cos(f*x + e)^2 - 17*a^3*c*\cos(f*x + e) - 7*a^3*c)*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^4*\sin(f*x + e))$

3.81.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx \\ &= a^3 \left(\int c \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx \right. \\ &+ \int 2c \sqrt{-c \sec(e + fx) + c} \sec^2(e + fx) dx \\ &+ \int (-2c \sqrt{-c \sec(e + fx) + c} \sec^4(e + fx)) dx \\ &\left. + \int (-c \sqrt{-c \sec(e + fx) + c} \sec^5(e + fx)) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**(3/2),x)`

output `a**3*(Integral(c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(2*c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2, x) + Integral(-2*c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4, x) + Integral(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**5, x))`

3.81.7 Maxima [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `Timed out`

3.81.8 Giac [A] (verification not implemented)

Time = 0.86 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.68

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx))^{3/2} dx = \frac{32\sqrt{2}\left(9\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)c^5 + 7c^6\right)a^3}{63\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{9}{2}}f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `32/63*sqrt(2)*(9*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^5 + 7*c^6)*a^3/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(9/2)*f)`

3.81.9 Mupad [B] (verification not implemented)

Time = 21.87 (sec) , antiderivative size = 471, normalized size of antiderivative = 5.54

$$\int \sec(e + fx)(a + a \sec(e + fx))^3 (c - c \sec(e + fx))^{3/2} dx = \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{e^{e \cdot 1i + f \cdot x \cdot 1i} - 1} \left(\frac{a^3 c 2i}{f} + \frac{a^3 c e^{e \cdot 1i + f \cdot x \cdot 1i} 22i}{63 f} \right) - \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{e \cdot 2i + f \cdot x \cdot 2i} + 1)^4} \left(\frac{a^3 c 32i}{9 f} + \frac{a^3 c e^{e \cdot 1i + f \cdot x \cdot 1i} 32i}{9 f} \right) - \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{e \cdot 2i + f \cdot x \cdot 2i} + 1)} \left(\frac{a^3 c 8i}{3 f} - \frac{a^3 c e^{e \cdot 1i + f \cdot x \cdot 1i} 200i}{63 f} \right) + \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{e \cdot 2i + f \cdot x \cdot 2i} + 1)^3} \left(\frac{a^3 c 32i}{7 f} + \frac{a^3 c e^{e \cdot 1i + f \cdot x \cdot 1i} 608i}{63 f} \right) - \frac{a^3 c e^{e \cdot 1i + f \cdot x \cdot 1i} \sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{21 f (e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{e \cdot 2i + f \cdot x \cdot 2i} + 1)^2} 160i$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(3/2))/cos(e + f*x),x)`

output `((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*c*2i)/f + (a^3*c*exp(e*1i + f*x*1i)*22i)/(63*f)))/(exp(e*1i + f*x*1i) - 1) - ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*c*32i)/(9*f) + (a^3*c*exp(e*1i + f*x*1i)*32i)/(9*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^4) - ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*c*8i)/(3*f) - (a^3*c*exp(e*1i + f*x*1i)*200i)/(63*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)) + ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*c*32i)/(7*f) + (a^3*c*exp(e*1i + f*x*1i)*608i)/(63*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^3) - (a^3*c*exp(e*1i + f*x*1i)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*160i)/(21*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^2)`

3.82 $\int \sec(e+fx)(a+a \sec(e+fx))^3 \sqrt{c - c \sec(e + fx)} dx$

3.82.1	Optimal result	633
3.82.2	Mathematica [A] (verified)	633
3.82.3	Rubi [A] (verified)	634
3.82.4	Maple [A] (verified)	635
3.82.5	Fricas [B] (verification not implemented)	635
3.82.6	Sympy [F]	636
3.82.7	Maxima [F]	636
3.82.8	Giac [A] (verification not implemented)	637
3.82.9	Mupad [B] (verification not implemented)	638

3.82.1 Optimal result

Integrand size = 34, antiderivative size = 41

$$\int \sec(e + fx)(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} dx$$

$$= -\frac{2c(a + a \sec(e + fx))^3 \tan(e + fx)}{7f \sqrt{c - c \sec(e + fx)}}$$

output `-2/7*c*(a+a*sec(f*x+e))^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)`

3.82.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.34

$$\int \sec(e + fx)(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{16a^3 \cos^6\left(\frac{1}{2}(e + fx)\right) \cot\left(\frac{1}{2}(e + fx)\right) \sec^3(e + fx) \sqrt{c - c \sec(e + fx)}}{7f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]],x]`

output `(16*a^3*Cos[(e + f*x)/2]^6*Cot[(e + f*x)/2]*Sec[e + f*x]^3*Sqrt[c - c*Sec[e + f*x]])/(7*f)`

3.82.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3 \sqrt{c - c \sec(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4441}$$

$$\frac{2c \tan(e + fx)(a \sec(e + fx) + a)^3}{7f \sqrt{c - c \sec(e + fx)}}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]],x]`

output `(-2*c*(a + a*Sec[e + f*x])^3*Tan[e + f*x])/(7*f*Sqrt[c - c*Sec[e + f*x]])`

3.82.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.82.4 Maple [A] (verified)

Time = 5.64 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

method	result
default	$\frac{2a^3(\cos(fx+e)+1)^4\sqrt{-c(\sec(fx+e)-1)}\sec(fx+e)^3\csc(fx+e)}{7f}$
parts	$-\frac{2a^3\sqrt{-c(\sec(fx+e)-1)}\sin(fx+e)}{f(\cos(fx+e)-1)} - \frac{2a^3\sqrt{-c(\sec(fx+e)-1)}(-5+16\cos(fx+e)^4+8\cos(fx+e)^3-2\cos(fx+e)^2+\cos(fx+e))}{35f}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(1/2),x,method=_RETURNV
ERBOSE)`

output `2/7*a^3/f*(cos(f*x+e)+1)^4*(-c*(sec(f*x+e)-1))^(1/2)*sec(f*x+e)^3*csc(f*x+
e)`

3.82.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. $2(37) = 74$.

Time = 0.26 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.37

$$\int \sec(e+fx)(a+a\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}dx$$

$$= \frac{2(a^3\cos(fx+e)^4+4a^3\cos(fx+e)^3+6a^3\cos(fx+e)^2+4a^3\cos(fx+e)+a^3)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{7f\cos(fx+e)^3\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(1/2),x, algorithm
m="fracas")`

output `2/7*(a^3*cos(f*x + e)^4 + 4*a^3*cos(f*x + e)^3 + 6*a^3*cos(f*x + e)^2 + 4*
a^3*cos(f*x + e) + a^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x
+ e)^3*sin(f*x + e))`

3.82.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} dx \\ &= a^3 \left(\int \sqrt{-c \sec(e + fx) + c} \sec(e + fx) dx + \int 3 \sqrt{-c \sec(e + fx) + c} \sec^2(e + fx) dx \right. \\ & \quad \left. + \int 3 \sqrt{-c \sec(e + fx) + c} \sec^3(e + fx) dx \right. \\ & \quad \left. + \int \sqrt{-c \sec(e + fx) + c} \sec^4(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e))**(1/2),x)`

output `a**3*(Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x), x) + Integral(3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2, x) + Integral(3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3, x) + Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4, x))`

3.82.7 Maxima [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} dx \\ &= \int (a \sec(fx + e) + a)^3 \sqrt{-c \sec(fx + e) + c} \sec(fx + e) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

```
output 2/7*(7*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^
(3/4)*(5*(a^3*f*cos(2*f*x + 2*e)^2 + a^3*f*sin(2*f*x + 2*e)^2 + 2*a^3*f*co
s(2*f*x + 2*e) + a^3*f)*integrate((((cos(10*f*x + 10*e)*cos(2*f*x + 2*e) +
4*cos(8*f*x + 8*e)*cos(2*f*x + 2*e) + 6*cos(6*f*x + 6*e)*cos(2*f*x + 2*e)
+ 4*cos(4*f*x + 4*e)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + sin(10*f*x +
10*e)*sin(2*f*x + 2*e) + 4*sin(8*f*x + 8*e)*sin(2*f*x + 2*e) + 6*sin(6*f*
x + 6*e)*sin(2*f*x + 2*e) + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + sin(2*f*
x + 2*e)^2)*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + (cos(2*
f*x + 2*e)*sin(10*f*x + 10*e) + 4*cos(2*f*x + 2*e)*sin(8*f*x + 8*e) + 6*co
s(2*f*x + 2*e)*sin(6*f*x + 6*e) + 4*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - co
s(10*f*x + 10*e)*sin(2*f*x + 2*e) - 4*cos(8*f*x + 8*e)*sin(2*f*x + 2*e) -
6*cos(6*f*x + 6*e)*sin(2*f*x + 2*e) - 4*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))
*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(1/2*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((cos(2*f*x + 2*e)*sin(10*f*x + 10
*e) + 4*cos(2*f*x + 2*e)*sin(8*f*x + 8*e) + 6*cos(2*f*x + 2*e)*sin(6*f*x +
6*e) + 4*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - cos(10*f*x + 10*e)*sin(2*f*x
+ 2*e) - 4*cos(8*f*x + 8*e)*sin(2*f*x + 2*e) - 6*cos(6*f*x + 6*e)*sin(2*f
*x + 2*e) - 4*cos(4*f*x + 4*e)*sin(2*f*x + 2*e))*cos(7/2*arctan2(sin(2*f*x
+ 2*e), cos(2*f*x + 2*e)))) - (cos(10*f*x + 10*e)*cos(2*f*x + 2*e) + 4*cos
(8*f*x + 8*e)*cos(2*f*x + 2*e) + 6*cos(6*f*x + 6*e)*cos(2*f*x + 2*e) + ...
```

3.82.8 Giac [A] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \sec(e + fx)(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} dx = \frac{16 \sqrt{2} a^3 c^4}{7 \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right)^{\frac{7}{2}} f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e))^(1/2),x, algorith
m="giac")
```

```
output 16/7*sqrt(2)*a^3*c^4/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(7/2)*f)
```

3.82.9 Mupad [B] (verification not implemented)

Time = 17.16 (sec) , antiderivative size = 375, normalized size of antiderivative = 9.15

$$\int \sec(e + fx)(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}} \left(\frac{a^3 \cdot 2i}{f} + \frac{a^3 e^{e \cdot 1i + f \cdot x \cdot 1i} \cdot 2i}{7f} \right)}{e^{e \cdot 1i + f \cdot x \cdot 1i} - 1}$$

$$- \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}} \left(\frac{a^3 \cdot 8i}{f} + \frac{a^3 e^{e \cdot 1i + f \cdot x \cdot 1i} \cdot 8i}{7f} \right)}{(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{2i + f \cdot x \cdot 2i} + 1)^2}$$

$$+ \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}} \left(\frac{a^3 \cdot 4i}{f} + \frac{a^3 e^{e \cdot 1i + f \cdot x \cdot 1i} \cdot 36i}{7f} \right)}{(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{2i + f \cdot x \cdot 2i} + 1)}$$

$$+ \frac{\sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}} \left(\frac{a^3 \cdot 16i}{7f} - \frac{a^3 e^{e \cdot 1i + f \cdot x \cdot 1i} \cdot 16i}{7f} \right)}{(e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{2i + f \cdot x \cdot 2i} + 1)^3}$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x),x)`

output `((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*2i)/f + (a^3*exp(e*1i + f*x*1i)*2i)/(7*f)))/(exp(e*1i + f*x*1i) - 1) - ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*8i)/f + (a^3*exp(e*1i + f*x*1i)*8i)/(7*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^2) + ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*4i)/f + (a^3*exp(e*1i + f*x*1i)*36i)/(7*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)) + ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((a^3*16i)/(7*f) - (a^3*exp(e*1i + f*x*1i)*16i)/(7*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*2i + f*x*2i) + 1)^3)`

3.83
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{\sqrt{c-c \sec(e+fx)}} dx$$

3.83.1	Optimal result	639
3.83.2	Mathematica [A] (verified)	640
3.83.3	Rubi [A] (verified)	640
3.83.4	Maple [A] (verified)	643
3.83.5	Fricas [A] (verification not implemented)	643
3.83.6	Sympy [F]	644
3.83.7	Maxima [F]	645
3.83.8	Giac [A] (verification not implemented)	645
3.83.9	Mupad [F(-1)]	646

3.83.1 Optimal result

Integrand size = 34, antiderivative size = 164

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{\sqrt{c-c \sec(e+fx)}} dx = -\frac{8\sqrt{2}a^3 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{\sqrt{c}f} + \frac{8a^3 \tan(e+fx)}{f\sqrt{c-c \sec(e+fx)}} + \frac{2a(a+a \sec(e+fx))^2 \tan(e+fx)}{5f\sqrt{c-c \sec(e+fx)}} + \frac{4(a^3+a^3 \sec(e+fx)) \tan(e+fx)}{3f\sqrt{c-c \sec(e+fx)}}$$

output

```
-8*a^3*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))*2^(1/2)/f/c^(1/2)+8*a^3*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)+2/5*a*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)+4/3*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)
```


3.83.2 Mathematica [A] (verified)

Time = 0.84 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.73

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{\sqrt{c-c\sec(e+fx)}} dx$$

$$= \frac{2a^3 \left(-60\sqrt{2}\sqrt{a}\operatorname{arctanh}\left(\frac{\sqrt{a(1+\sec(e+fx))}}{\sqrt{2}\sqrt{a}}\right) + \sqrt{a(1+\sec(e+fx))}(73+16\sec(e+fx)+3\sec^2(e+fx)) \right)}{15f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/Sqrt[c - c*Sec[e + f*x]],x]`

output `(2*a^3*(-60*Sqrt[2]*Sqrt[a]*ArcTanh[Sqrt[a*(1 + Sec[e + f*x])]/(Sqrt[2]*Sqrt[a])] + Sqrt[a*(1 + Sec[e + f*x])]*(73 + 16*Sec[e + f*x] + 3*Sec[e + f*x]^2))*Tan[e + f*x]/(15*f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.83.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4444, 3042, 4444, 3042, 4444, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{\sqrt{c-c\sec(e+fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx$$

$$\downarrow \text{4444}$$

$$2a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2a \tan(e+fx)(a\sec(e+fx)+a)^2}{5f\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow \text{3042}$$

3.83. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{\sqrt{c-c\sec(e+fx)}} dx$

$$\begin{aligned}
& 2a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2a \tan(e+fx)(a \sec(e+fx)+a)^2}{5f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{4444} \\
& 2a \left(2a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right) + \\
& \quad \frac{2a \tan(e+fx)(a \sec(e+fx)+a)^2}{5f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& 2a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right) + \\
& \quad \frac{2a \tan(e+fx)(a \sec(e+fx)+a)^2}{5f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{4444} \\
& 2a \left(2a \left(2a \int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right) + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right) + \\
& \quad \frac{2a \tan(e+fx)(a \sec(e+fx)+a)^2}{5f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& 2a \left(2a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right) + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right) + \\
& \quad \frac{2a \tan(e+fx)(a \sec(e+fx)+a)^2}{5f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{4282} \\
& 2a \left(2a \left(\frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{4a \int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c\sec(e+fx)} + 2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{f} \right) + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right) + \\
& \quad \frac{2a \tan(e+fx)(a \sec(e+fx)+a)^2}{5f\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{216}
\end{aligned}$$

$$2a \left(\frac{2 \tan(e+fx) (a^2 \sec(e+fx) + a^2)}{3f \sqrt{c - c \sec(e+fx)}} + 2a \left(\frac{2a \tan(e+fx)}{f \sqrt{c - c \sec(e+fx)}} - \frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c - c \sec(e+fx)}}\right)}{\sqrt{cf}} \right) \right) + \frac{2a \tan(e+fx) (a \sec(e+fx) + a)^2}{5f \sqrt{c - c \sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/Sqrt[c - c*Sec[e + f*x]],x]`

output `(2*a*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(5*f*Sqrt[c - c*Sec[e + f*x]]) + 2*a*((2*(a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e + f*x]]) + 2*a*((-2*Sqrt[2]*a*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])])/(Sqrt[c]*f) + (2*a*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]])))`

3.83.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x])/Sqrt[a + b*Csc[e + f*x]]], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4444 `Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.))/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*d*Cot[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/(f*(2*n - 1)*Sqrt[a + b*Csc[e + f*x]]), x] + Simp[2*c*((2*n - 1)/(2*n - 1)) Int[Csc[e + f*x]*((c + d*Csc[e + f*x])^(n - 1))/Sqrt[a + b*Csc[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`

3.83.4 Maple [A] (verified)

Time = 6.38 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.76

method	result
default	$\frac{a^3 \sqrt{2} \left(120 \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e)^2 \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) - 73\sqrt{2} \cos(fx+e)^2 - 16\sqrt{2} \cos(fx+e) - 3\sqrt{2} \right) \tan(fx+e) \sec(fx+e)}{15f \sqrt{-c(\sec(fx+e)-1)}}$
parts	$\frac{a^3 \sqrt{2} \sin(fx+e) \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right)}{f(\cos(fx+e)+1)\sqrt{-c(\sec(fx+e)-1)}\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} + \frac{a^3 \sqrt{2} \left(-15 \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e)^2 \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) + 1 \right)}{15f \sqrt{-c(\sec(fx+e)-1)}}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNV
ERBOSE)`

output `-1/15*a^3/f*2^(1/2)*(120*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2*a
rctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))-73*2^(1/2)*cos(f*x+e
)^2-16*2^(1/2)*cos(f*x+e)-3*2^(1/2))/(-c*(sec(f*x+e)-1))^(1/2)*tan(f*x+e)*
sec(f*x+e)^2`

3.83.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 377, normalized size of antiderivative = 2.30

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{\sqrt{c-c\sec(e+fx)}} dx$$

$$= \left[\frac{2 \left(30 \sqrt{2} a^3 c \sqrt{-\frac{1}{c}} \cos(fx+e)^2 \log \left(-\frac{2\sqrt{2}(\cos(fx+e)^2 + \cos(fx+e)) \sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}} \sqrt{-\frac{1}{c}} - (3\cos(fx+e)+1)\sin(fx+e)}{(\cos(fx+e)-1)\sin(fx+e)} \right)}{15cf \cos(fx+e)} \right) \right] s$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x, algorithm
m="fricas")`

```
output [2/15*(30*sqrt(2)*a^3*c*sqrt(-1/c)*cos(f*x + e)^2*log(-(2*sqrt(2))*(cos(f*x
+ e)^2 + cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*sqrt(-1/c)
- (3*cos(f*x + e) + 1)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*s
in(f*x + e) - (73*a^3*cos(f*x + e)^3 + 89*a^3*cos(f*x + e)^2 + 19*a^3*cos(
f*x + e) + 3*a^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c*f*cos(f*x +
e)^2*sin(f*x + e)), 2/15*(60*sqrt(2)*a^3*sqrt(c)*arctan(sqrt(2)*sqrt((c*co
s(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*cos(f*x
+ e)^2*sin(f*x + e) - (73*a^3*cos(f*x + e)^3 + 89*a^3*cos(f*x + e)^2 + 19
*a^3*cos(f*x + e) + 3*a^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(c*f*c
os(f*x + e)^2*sin(f*x + e))]
```

3.83.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{\sqrt{c-c\sec(e+fx)}} dx = a^3 \left(\int \frac{\sec(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx \right. \\ \left. + \int \frac{3\sec^2(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx \right. \\ \left. + \int \frac{3\sec^3(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx \right. \\ \left. + \int \frac{\sec^4(e+fx)}{\sqrt{-c\sec(e+fx)+c}} dx \right)$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**(1/2),x)
```

```
output a**3*(Integral(sec(e + f*x)/sqrt(-c*sec(e + f*x) + c), x) + Integral(3*sec
(e + f*x)**2/sqrt(-c*sec(e + f*x) + c), x) + Integral(3*sec(e + f*x)**3/sq
rt(-c*sec(e + f*x) + c), x) + Integral(sec(e + f*x)**4/sqrt(-c*sec(e + f*x
) + c), x))
```

3.83.7 Maxima [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{(a\sec(fx+e)+a)^3 \sec(fx+e)}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^3*sec(f*x + e)/sqrt(-c*sec(f*x + e) + c), x)`

3.83.8 Giac [A] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.68

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{\sqrt{c-c\sec(e+fx)}} dx$$

$$= \frac{8a^3 \left(\frac{15\sqrt{2} \arctan\left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}}\right)}{\sqrt{c}} + \frac{\sqrt{2} \left(15(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^2 - 5(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)c + 3c^2 \right)}{(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^{\frac{5}{2}}} \right)}{15f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x, algorithm m="giac")`

output `8/15*a^3*(15*sqrt(2)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/sqrt(c) + sqrt(2)*(15*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2 - 5*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c + 3*c^2)/(c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2))/f`

3.83.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^3}{\cos(e+fx) \sqrt{c - \frac{c}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)), x)`

$$3.84 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{3/2}} dx$$

3.84.1	Optimal result	647
3.84.2	Mathematica [C] (verified)	647
3.84.3	Rubi [A] (verified)	648
3.84.4	Maple [A] (verified)	651
3.84.5	Fricas [A] (verification not implemented)	651
3.84.6	Sympy [F]	652
3.84.7	Maxima [F]	653
3.84.8	Giac [A] (verification not implemented)	653
3.84.9	Mupad [F(-1)]	654

3.84.1 Optimal result

Integrand size = 34, antiderivative size = 168

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{3/2}} dx = \frac{10\sqrt{2}a^3 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{c^{3/2}f} - \frac{a(a+a \sec(e+fx))^2 \tan(e+fx)}{f(c-c \sec(e+fx))^{3/2}} - \frac{10a^3 \tan(e+fx)}{cf\sqrt{c-c \sec(e+fx)}} - \frac{5(a^3+a^3 \sec(e+fx)) \tan(e+fx)}{3cf\sqrt{c-c \sec(e+fx)}}$$

```
output 10*a^3*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))*2^(1/2)/c^(3/2)/f-a*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(3/2)-10*a^3*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(1/2)-5/3*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(1/2)
```

3.84.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.60 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.38

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{3/2}} dx = \frac{a^3 \operatorname{Hypergeometric2F1}\left(2, \frac{7}{2}, \frac{9}{2}, \frac{1}{2}(1+\sec(e+fx))\right) (1+\sec(e+fx))^3 \tan(e+fx)}{14cf\sqrt{c-c \sec(e+fx)}}$$

3.84. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{3/2}} dx$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^(3/2),x]`

output `-1/14*(a^3*Hypergeometric2F1[2, 7/2, 9/2, (1 + Sec[e + f*x])/2]*(1 + Sec[e + f*x])^3*Tan[e + f*x])/(c*f*Sqrt[c - c*Sec[e + f*x]])`

3.84.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4445, 3042, 4444, 3042, 4444, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c-c\sec(e+fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{5a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{\sqrt{c-c\sec(e+fx)}} dx}{2c} - \frac{a \tan(e+fx)(a\sec(e+fx)+a)^2}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{5a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2c} - \frac{a \tan(e+fx)(a\sec(e+fx)+a)^2}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{4444} \\
 & -\frac{5a \left(2a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right)}{2c} - \frac{a \tan(e+fx)(a\sec(e+fx)+a)^2}{f(c-c\sec(e+fx))^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.84. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{3/2}} dx$

$$\begin{aligned}
& \frac{5a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right)}{\frac{2c}{a \tan(e+fx)(a \sec(e+fx)+a)^2} f(c-c\sec(e+fx))^{3/2}} \\
& \quad \downarrow \text{4444} \\
& \frac{5a \left(2a \left(2a \int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right) + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right)}{\frac{2c}{a \tan(e+fx)(a \sec(e+fx)+a)^2} f(c-c\sec(e+fx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{5a \left(2a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} \right) + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right)}{\frac{2c}{a \tan(e+fx)(a \sec(e+fx)+a)^2} f(c-c\sec(e+fx))^{3/2}} \\
& \quad \downarrow \text{4282} \\
& \frac{5a \left(2a \left(\frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{4a \int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c\sec(e+fx)} + 2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{f} \right) + \frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} \right)}{\frac{2c}{a \tan(e+fx)(a \sec(e+fx)+a)^2} f(c-c\sec(e+fx))^{3/2}} \\
& \quad \downarrow \text{216} \\
& \frac{5a \left(\frac{2 \tan(e+fx)(a^2 \sec(e+fx)+a^2)}{3f\sqrt{c-c\sec(e+fx)}} + 2a \left(\frac{2a \tan(e+fx)}{f\sqrt{c-c\sec(e+fx)}} - \frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{c}f} \right) \right)}{\frac{2c}{a \tan(e+fx)(a \sec(e+fx)+a)^2} f(c-c\sec(e+fx))^{3/2}}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^(3/2),x]`

```
output -((a*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2)))
- (5*a*((2*(a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e +
f*x]]) + 2*a*((-2*Sqrt[2]*a*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c
- c*Sec[e + f*x]])])/(Sqrt[c]*f) + (2*a*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e
+ f*x]])))))/(2*c)
```

3.84.3.1 Defintions of rubi rules used

- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`
- rule 4444 `Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.))/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*d*Cot[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/(f*(2*n - 1)*Sqrt[a + b*Csc[e + f*x]]), x] + Simp[2*c*((2*n - 1)/(2*n - 1)) Int[Csc[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/Sqrt[a + b*Csc[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`
- rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

3.84.4 Maple [A] (verified)

Time = 7.10 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.84

method	result
default	$a^3 \sqrt{2} \left(30 \arctan \left(\frac{\sqrt{2}}{2 \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} \right) \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \tan(fx+e) + 19\sqrt{2} \cot(fx+e) + 7\sqrt{2} \csc(fx+e) - 13\sqrt{2} \sec(fx+e) \csc(fx+e) \right) \sqrt{-c(\sec(fx+e)-1)}$
parts	Expression too large to display

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x,method=_RETURNV ERBOSE)`

output `1/3*a^3/c/f*2^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)*(30*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*tan(f*x+e)+19*2^(1/2)*cot(f*x+e)+7*2^(1/2)*csc(f*x+e)-13*2^(1/2)*sec(f*x+e)*csc(f*x+e)-2^(1/2)*sec(f*x+e)^2*csc(f*x+e))`

3.84.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 432, normalized size of antiderivative = 2.57

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{3/2}} dx = \frac{15\sqrt{2}(a^3c\cos(fx+e)^2 - a^3c\cos(fx+e))\sqrt{-\frac{1}{c}} \log\left(\frac{2\sqrt{2}(\cos(fx+e)+1)}{\cos(fx+e)+1}\right)}{3(c^2f\cos(fx+e)^2 - c^2f\cos(fx+e))\sin(fx+e)} + \frac{2\left(\frac{15\sqrt{2}(a^3c\cos(fx+e)^2 - a^3c\cos(fx+e))\arctan\left(\frac{\sqrt{2}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\cos(fx+e)}{\sqrt{c}\sin(fx+e)}\right)\sin(fx+e)}{\sqrt{c}} - (19a^3\cos(fx+e)^3 + 7a^3\cos(fx+e))\right)}{3(c^2f\cos(fx+e)^2 - c^2f\cos(fx+e))\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x, algorithm="fricas")`

output `[1/3*(15*sqrt(2)*(a^3*c*cos(f*x + e)^2 - a^3*c*cos(f*x + e))*sqrt(-1/c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*sqrt(-1/c) + (3*cos(f*x + e) + 1)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) + 2*(19*a^3*cos(f*x + e)^3 + 7*a^3*cos(f*x + e)^2 - 13*a^3*cos(f*x + e) - a^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^2*f*cos(f*x + e)^2 - c^2*f*cos(f*x + e))*sin(f*x + e)), -2/3*(15*sqrt(2)*(a^3*c*cos(f*x + e)^2 - a^3*c*cos(f*x + e))*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e)/sqrt(c) - (19*a^3*cos(f*x + e)^3 + 7*a^3*cos(f*x + e)^2 - 13*a^3*cos(f*x + e) - a^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((c^2*f*cos(f*x + e)^2 - c^2*f*cos(f*x + e))*sin(f*x + e))]`

3.84.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{3/2}} dx = a^3 \left(\int \frac{\sec(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec(e+fx)+c\sqrt{-c\sec(e+fx)+c}} dx \right. \\ \left. + \int \frac{3\sec^2(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec(e+fx)+c\sqrt{-c\sec(e+fx)+c}} dx \right. \\ \left. + \int \frac{3\sec^3(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec(e+fx)+c\sqrt{-c\sec(e+fx)+c}} dx \right. \\ \left. + \int \frac{\sec^4(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec(e+fx)+c\sqrt{-c\sec(e+fx)+c}} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**(3/2),x)`

output `a**3*(Integral(sec(e + f*x)/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x) + Integral(3*sec(e + f*x)**2/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x) + Integral(3*sec(e + f*x)**3/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x) + Integral(sec(e + f*x)**4/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x))`

3.84.7 Maxima [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{(a\sec(fx+e)+a)^3 \sec(fx+e)}{(-c\sec(fx+e)+c)^{3/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^3*sec(f*x + e)/(-c*sec(f*x + e) + c)^(3/2), x)`

3.84.8 Giac [A] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{3/2}} dx = \frac{2a^3 \left(\frac{15\sqrt{2} \arctan\left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}}\right)}{c^{3/2}} + \frac{2\sqrt{2}(6c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 7c)}{(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^{3/2} c} + \frac{3\sqrt{2}\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^2} \right)}{3f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x, algorithm m="giac")`

output `-2/3*a^3*(15*sqrt(2)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/c^(3/2) + 2*sqrt(2)*(6*c*tan(1/2*f*x + 1/2*e)^2 - 7*c)/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c) + 3*sqrt(2)*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/(c^2*tan(1/2*f*x + 1/2*e)^2))/f`

3.84.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^3}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),x)`output `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)), x)`

3.85
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{5/2}} dx$$

3.85.1	Optimal result	655
3.85.2	Mathematica [C] (verified)	655
3.85.3	Rubi [A] (verified)	656
3.85.4	Maple [A] (verified)	659
3.85.5	Fricas [A] (verification not implemented)	659
3.85.6	Sympy [F]	660
3.85.7	Maxima [F(-1)]	661
3.85.8	Giac [A] (verification not implemented)	661
3.85.9	Mupad [F(-1)]	661

3.85.1 Optimal result

Integrand size = 34, antiderivative size = 174

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{5/2}} dx =$$

$$-\frac{15a^3 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{2\sqrt{2}c^{5/2}f} - \frac{a(a+a \sec(e+fx))^2 \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}}$$

$$+ \frac{5(a^3+a^3 \sec(e+fx)) \tan(e+fx)}{4cf(c-c \sec(e+fx))^{3/2}} + \frac{15a^3 \tan(e+fx)}{4c^2 f \sqrt{c-c \sec(e+fx)}}$$

```
output -15/4*a^3*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/c^(5/2)/f*2^(1/2)-1/2*a*(a+a*sec(f*x+e))^2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(5/2)+5/4*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(3/2)+15/4*a^3*tan(f*x+e)/c^2/f/(c-c*sec(f*x+e))^(1/2)
```

3.85.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.62 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.37

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{5/2}} dx =$$

$$\frac{a^3 \text{Hypergeometric2F1}\left(3, \frac{7}{2}, \frac{9}{2}, \frac{1}{2}(1+\sec(e+fx))\right) (1+\sec(e+fx))^3 \tan(e+fx)}{28c^2 f \sqrt{c-c \sec(e+fx)}}$$

3.85.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{5/2}} dx$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/28*(a^3*Hypergeometric2F1[3, 7/2, 9/2, (1 + Sec[e + f*x])/2]*(1 + Sec[e + f*x])^3*Tan[e + f*x])/(c^2*f*Sqrt[c - c*Sec[e + f*x]])`

3.85.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4445, 3042, 4445, 3042, 4444, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c-c\sec(e+fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^3}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4445} \\
 & -\frac{5a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^2}{(c-c\sec(e+fx))^{3/2}} dx}{4c} - \frac{a \tan(e+fx)(a\sec(e+fx)+a)^2}{2f(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{5a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^2}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{4c} - \frac{a \tan(e+fx)(a\sec(e+fx)+a)^2}{2f(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4445} \\
 & -\frac{5a \left(-\frac{3a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)}{\sqrt{c-c\sec(e+fx)}} dx}{2c} - \frac{\tan(e+fx)(a^2\sec(e+fx)+a^2)}{f(c-c\sec(e+fx))^{3/2}} \right)}{4c} - \frac{a \tan(e+fx)(a\sec(e+fx)+a)^2}{2f(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.85. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{5a \left(\frac{3a \int \frac{\csc(e+fx+\frac{\pi}{2}) (\csc(e+fx+\frac{\pi}{2}))^{a+a}}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx)+a^2)}{f(c-c \sec(e+fx))^{3/2}} \right)}{a \tan(e+fx) (a \sec(e+fx) + a)^2} \\
 & \qquad \qquad \qquad \frac{4c}{2f(c-c \sec(e+fx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{4444} \\
 & \frac{5a \left(\frac{3a \left(2a \int \frac{\sec(e+fx)}{\sqrt{c-c \sec(e+fx)}} dx + \frac{2a \tan(e+fx)}{f \sqrt{c-c \sec(e+fx)}} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx)+a^2)}{f(c-c \sec(e+fx))^{3/2}} \right)}{a \tan(e+fx) (a \sec(e+fx) + a)^2} \\
 & \qquad \qquad \qquad \frac{4c}{2f(c-c \sec(e+fx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{5a \left(\frac{3a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx + \frac{2a \tan(e+fx)}{f \sqrt{c-c \sec(e+fx)}} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx)+a^2)}{f(c-c \sec(e+fx))^{3/2}} \right)}{a \tan(e+fx) (a \sec(e+fx) + a)^2} \\
 & \qquad \qquad \qquad \frac{4c}{2f(c-c \sec(e+fx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{4282} \\
 & \frac{5a \left(\frac{3a \left(\frac{2a \tan(e+fx)}{f \sqrt{c-c \sec(e+fx)}} - \frac{4a \int \frac{1}{c^2 \tan^2(e+fx) + 2c} dx - \frac{c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}}}{f} \right)}{2c} - \frac{\tan(e+fx) (a^2 \sec(e+fx)+a^2)}{f(c-c \sec(e+fx))^{3/2}} \right)}{a \tan(e+fx) (a \sec(e+fx) + a)^2} \\
 & \qquad \qquad \qquad \frac{4c}{2f(c-c \sec(e+fx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{216} \\
 & \frac{5a \left(\frac{\tan(e+fx) (a^2 \sec(e+fx)+a^2)}{f(c-c \sec(e+fx))^{3/2}} - \frac{3a \left(\frac{2a \tan(e+fx)}{f \sqrt{c-c \sec(e+fx)}} - \frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{\sqrt{c}f} \right)}{2c} \right)}{a \tan(e+fx) (a \sec(e+fx) + a)^2} \\
 & \qquad \qquad \qquad \frac{4c}{2f(c-c \sec(e+fx))^{5/2}}
 \end{aligned}$$

3.85. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c-c \sec(e+fx))^{5/2}} dx$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/2*(a*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(5/2)) - (5*a*(-((a^2 + a^2*Sec[e + f*x])*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2))) - (3*a*((-2*sqrt[2]*a*ArcTan[(sqrt[c]*Tan[e + f*x])/(sqrt[2]*sqrt[c - c*Sec[e + f*x]])])/(sqrt[c]*f) + (2*a*Tan[e + f*x])/(f*sqrt[c - c*Sec[e + f*x]])))/(2*c)))/(4*c)`

3.85.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x])/sqrt[a + b*Csc[e + f*x]]], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4444 `Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.))/sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*d*Cot[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/(f*(2*n - 1)*sqrt[a + b*Csc[e + f*x]])), x] + Simp[2*c*((2*n - 1)/(2*n - 1)) Int[Csc[e + f*x]*((c + d*Csc[e + f*x])^(n - 1)/sqrt[a + b*Csc[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`

rule 4445 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0] && LtQ[m, -2*(-1)] && IntegerQ[2*m]`

3.85.4 Maple [A] (verified)

Time = 6.29 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.28

method	result
default	$\frac{a^3 \sqrt{2} \left(15 \arctan \left(\frac{1}{\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} \right) (1-\cos(fx+e))^4 \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \csc(fx+e) - 15(1-\cos(fx+e)) \right)}{4c^2 f \sqrt{\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} (1-\cos(fx+e))^3 \left((1-\cos(fx+e))^2 \csc(fx+e) \right)}$
parts	Expression too large to display

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x,method=_RETURNV
ERBOSE)`

output `1/4*a^3/c^2/f*2^(1/2)/(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)
)*csc(f*x+e)^2)^(1/2)/(1-cos(f*x+e))^3/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*(
15*arctan(1/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2))*(1-cos(f*x+e))^4*((1-
cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*csc(f*x+e)-15*(1-cos(f*x+e))^4*csc(f*x
+e)+5*(1-cos(f*x+e))^2*sin(f*x+e)+2*sin(f*x+e)^3)`

3.85.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.53

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{5/2}} dx = \left[\frac{15\sqrt{2}(a^3 \cos(fx+e)^2 - 2a^3 \cos(fx+e) + a^3) \sqrt{-c} \log\left(\frac{2\sqrt{2}}{\dots}\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x, algorithm
m="fracas")`

```
output [-1/8*(15*sqrt(2)*(a^3*cos(f*x + e)^2 - 2*a^3*cos(f*x + e) + a^3)*sqrt(-c)
*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x +
e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x +
e) - 1)*sin(f*x + e))) *sin(f*x + e) + 4*(9*a^3*cos(f*x + e)^3 - 8*a^3*cos(
f*x + e)^2 - 13*a^3*cos(f*x + e) + 4*a^3)*sqrt((c*cos(f*x + e) - c)/cos(f*
x + e)))/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x +
e)), 1/4*(15*sqrt(2)*(a^3*cos(f*x + e)^2 - 2*a^3*cos(f*x + e) + a^3)*sqrt(
c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sq
rt(c)*sin(f*x + e))) *sin(f*x + e) - 2*(9*a^3*cos(f*x + e)^3 - 8*a^3*cos(f*
x + e)^2 - 13*a^3*cos(f*x + e) + 4*a^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x
+ e)))/((c^3*f*cos(f*x + e)^2 - 2*c^3*f*cos(f*x + e) + c^3*f)*sin(f*x + e)
)]
```

3.85.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{5/2}} dx = a^3 \left(\int \frac{\sec(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}} \right. \\ + \int \frac{3\sec^2(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)+c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}} \\ + \int \frac{3\sec^3(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)+c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}} \\ \left. + \int \frac{\sec^4(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)+c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}} \right)$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**(5/2),x)
```

```
output a**3*(Integral(sec(e + f*x)/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**
2 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f
*x) + c)), x) + Integral(3*sec(e + f*x)**2/(c**2*sqrt(-c*sec(e + f*x) + c)
*sec(e + f*x)**2 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sq
rt(-c*sec(e + f*x) + c)), x) + Integral(3*sec(e + f*x)**3/(c**2*sqrt(-c*se
c(e + f*x) + c)*sec(e + f*x)**2 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e +
f*x) + c**2*sqrt(-c*sec(e + f*x) + c)), x) + Integral(sec(e + f*x)**4/(c*
**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 - 2*c**2*sqrt(-c*sec(e + f*x)
+ c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f*x) + c)), x))
```

3.85.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^{5/2}} dx = \text{Timed out}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x, algorithm
m="maxima")
```

```
output Timed out
```

3.85.8 Giac [A] (verification not implemented)

Time = 1.14 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.76

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^{5/2}} dx = \frac{a^3 \left(\frac{15\sqrt{2} \arctan\left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}}\right)}{c^{5/2}} + \frac{8\sqrt{2}}{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}} + \frac{7\sqrt{2}(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^{3/2}}{c^4} \right)}{4f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x, algorithm
m="giac")
```

```
output 1/4*a^3*(15*sqrt(2)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/c^(
5/2) + 8*sqrt(2)/(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^2) + (7*sqrt(2)*(c*
tan(1/2*f*x + 1/2*e)^2 - c)^(3/2) + 9*sqrt(2)*sqrt(c*tan(1/2*f*x + 1/2*e)^
2 - c)*c)/(c^4*tan(1/2*f*x + 1/2*e)^4)/f
```

3.85.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c - c \sec(e + fx))^{5/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e + fx)}\right)^3}{\cos(e + fx) \left(c - \frac{c}{\cos(e + fx)}\right)^{5/2}} dx$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x)`

output `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)), x)`

3.85. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c-c\sec(e+fx))^{5/2}} dx$

3.86 $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{a+a\sec(e+fx)} dx$

3.86.1	Optimal result	663
3.86.2	Mathematica [A] (verified)	663
3.86.3	Rubi [A] (verified)	664
3.86.4	Maple [A] (verified)	666
3.86.5	Fricas [A] (verification not implemented)	666
3.86.6	Sympy [F(-1)]	667
3.86.7	Maxima [A] (verification not implemented)	667
3.86.8	Giac [A] (verification not implemented)	668
3.86.9	Mupad [B] (verification not implemented)	668

3.86.1 Optimal result

Integrand size = 34, antiderivative size = 142

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{a+a\sec(e+fx)} dx = \frac{128c^4 \tan(e+fx)}{5af\sqrt{c-c\sec(e+fx)}} + \frac{32c^3\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{5af} + \frac{12c^2(c-c\sec(e+fx))^{3/2}\tan(e+fx)}{5af} + \frac{2c(c-c\sec(e+fx))^{5/2}\tan(e+fx)}{f(a+a\sec(e+fx))}$$

output `12/5*c^2*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/a/f+2*c*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))+128/5*c^4*tan(f*x+e)/a/f/(c-c*sec(f*x+e))^(1/2)+32/5*c^3*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/a/f`

3.86.2 Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.51

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{a+a\sec(e+fx)} dx = \frac{2c^4(91+43\sec(e+fx)-7\sec^2(e+fx)+\sec^3(e+fx))\tan(e+fx)}{5af(1+\sec(e+fx))\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(7/2))/(a + a*Sec[e + f*x]),x]`

output $(2*c^4*(91 + 43*Sec[e + f*x] - 7*Sec[e + f*x]^2 + Sec[e + f*x]^3)*Tan[e + f*x])/(5*a*f*(1 + Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]])$

3.86.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4442, 3042, 4280, 3042, 4280, 3042, 4279}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{7/2}}{a\csc(e+fx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow 4442 \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{f(a\sec(e+fx)+a)} - \frac{6c\int\sec(e+fx)(c-c\sec(e+fx))^{5/2}dx}{a} \\
 & \quad \downarrow 3042 \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{f(a\sec(e+fx)+a)} - \frac{6c\int\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}dx}{a} \\
 & \quad \downarrow 4280 \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{f(a\sec(e+fx)+a)} - \frac{6c\left(\frac{8}{5}c\int\sec(e+fx)(c-c\sec(e+fx))^{3/2}dx - \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{5f}\right)}{a} \\
 & \quad \downarrow 3042 \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{f(a\sec(e+fx)+a)} - \frac{6c\left(\frac{8}{5}c\int\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}dx - \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{5f}\right)}{a} \\
 & \quad \downarrow 4280
 \end{aligned}$$

$$\frac{\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{5/2}}{f(a \sec(e+fx) + a)} - 6c \left(\frac{8}{5}c \left(\frac{4}{3}c \int \sec(e+fx) \sqrt{c - c \sec(e+fx)} dx - \frac{2c \tan(e+fx) \sqrt{c - c \sec(e+fx)}}{3f} \right) - \frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{5f} \right)}{a}$$

↓ 3042

$$\frac{\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{5/2}}{f(a \sec(e+fx) + a)} - 6c \left(\frac{8}{5}c \left(\frac{4}{3}c \int \csc(e+fx + \frac{\pi}{2}) \sqrt{c - c \csc(e+fx + \frac{\pi}{2})} dx - \frac{2c \tan(e+fx) \sqrt{c - c \sec(e+fx)}}{3f} \right) - \frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{5f} \right)}{a}$$

↓ 4279

$$\frac{\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{5/2}}{f(a \sec(e+fx) + a)} - 6c \left(\frac{8}{5}c \left(-\frac{8c^2 \tan(e+fx)}{3f \sqrt{c - c \sec(e+fx)}} - \frac{2c \tan(e+fx) \sqrt{c - c \sec(e+fx)}}{3f} \right) - \frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{5f} \right)}{a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(7/2))/(a + a*Sec[e + f*x]),x]`

output `(2*c*(c - c*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (6*c*((-2*c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(5*f) + (8*c*((-8*c^2*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e + f*x]]) - (2*c*Sqrt[c - c*Sec[e + f*x]])*Tan[e + f*x])/(3*f))))/5)/a`

3.86.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4279 `Int[csc[(e_.) + (f_.)*(x_.)]*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] := Simp[-2*b*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]])), x] /; Free Q[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

```
rule 4280 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_
Symbol] :> Simp[(-b)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m - 1)/(f*m)), x]
+ Simp[a*((2*m - 1)/m) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m - 1), x],
x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && Intege
rQ[2*m]
```

```
rule 4442 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[2*a*c*Cot[e +
f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))),
x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*
x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[
m, -2^(-1)]
```

3.86.4 Maple [A] (verified)

Time = 5.28 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.57

method	result	size
default	$\frac{2c^3(\sec(fx+e)-1)^3\sqrt{-c(\sec(fx+e)-1)}(91\cos(fx+e)^3+43\cos(fx+e)^2-7\cos(fx+e)+1)\cot(fx+e)}{5af(\cos(fx+e)-1)^3}$	81

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e)),x,method=_RETURNVER
BOSE)
```

```
output 2/5/a/f*c^3*(sec(f*x+e)-1)^3*(-c*(sec(f*x+e)-1))^(1/2)*(91*cos(f*x+e)^3+43
*cos(f*x+e)^2-7*cos(f*x+e)+1)/(cos(f*x+e)-1)^3*cot(f*x+e)
```

3.86.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.62

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^{7/2}}{a + a\sec(e + fx)} dx =$$

$$\frac{2(91c^3\cos(fx + e)^3 + 43c^3\cos(fx + e)^2 - 7c^3\cos(fx + e) + c^3)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{5af\cos(fx + e)^2\sin(fx + e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e)),x, algorithm="fricas")`

output `-2/5*(91*c^3*cos(f*x + e)^3 + 43*c^3*cos(f*x + e)^2 - 7*c^3*cos(f*x + e) + c^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(a*f*cos(f*x + e)^2*sin(f*x + e))`

3.86.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{7/2}}{a + a \sec(e + fx)} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(7/2)/(a+a*sec(f*x+e)),x)`

output `Timed out`

3.86.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.15

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{7/2}}{a + a \sec(e + fx)} dx = \frac{8 \left(16 \sqrt{2} c^{7/2} - \frac{56 \sqrt{2} c^{7/2} \sin^2(fx+e)}{(\cos(fx+e)+1)^2} + \frac{70 \sqrt{2} c^{7/2} \sin^4(fx+e)}{(\cos(fx+e)+1)^4} - \frac{35 \sqrt{2} c^{7/2} \sin^6(fx+e)}{(\cos(fx+e)+1)^6} \right)}{5 a f \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right)^{7/2} \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)^{7/2}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `8/5*(16*sqrt(2)*c^(7/2) - 56*sqrt(2)*c^(7/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 70*sqrt(2)*c^(7/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 35*sqrt(2)*c^(7/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 5*sqrt(2)*c^(7/2)*sin(f*x + e)^8/(cos(f*x + e) + 1)^8)/(a*f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^(7/2)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^(7/2))`

3.86.8 Giac [A] (verification not implemented)

Time = 0.83 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.76

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{a+a\sec(e+fx)} dx =$$

$$\frac{8\sqrt{2}c^3 \left(\frac{5\sqrt{c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c}}{a} - \frac{15(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)^2 c+5(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)c^2+c^3}{(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)^{\frac{5}{2}}a} \right)}{5f}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e)),x, algorithm="giac")
```

```
output -8/5*sqrt(2)*c^3*(5*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/a - (15*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c + 5*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^2 + c^3)/((c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2)*a))/f
```

3.86.9 Mupad [B] (verification not implemented)

Time = 17.98 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.15

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{a+a\sec(e+fx)} dx =$$

$$\frac{2c^3 \sqrt{c - \frac{c}{\frac{e^{-e1i-fx1i}}{2} + \frac{e^{e1i+fx1i}}{2}}}}{5af (e^{e2i+fx2i} - 1) (e^{e2i+fx2i} + 1)^2} (e^{e1i+fx1i} 86i + e^{e2i+fx2i} 245i + e^{e3i+fx3i} 180i + e^{e4i+fx4i} 245i + e^{e5i+fx5i} 86i)$$

```
input int((c - c/cos(e + f*x))^(7/2)/(cos(e + f*x)*(a + a/cos(e + f*x))),x)
```

```
output -(2*c^3*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*(exp(e*1i + f*x*1i)*86i + exp(e*2i + f*x*2i)*245i + exp(e*3i + f*x*3i)*180i + exp(e*4i + f*x*4i)*245i + exp(e*5i + f*x*5i)*86i + exp(e*6i + f*x*6i)*91i + 91i))/(5*a*f*(exp(e*2i + f*x*2i) - 1)*(exp(e*2i + f*x*2i) + 1)^2)
```

3.87
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a+a\sec(e+fx)} dx$$

3.87.1	Optimal result	669
3.87.2	Mathematica [A] (verified)	669
3.87.3	Rubi [A] (verified)	670
3.87.4	Maple [A] (verified)	672
3.87.5	Fricas [A] (verification not implemented)	672
3.87.6	Sympy [F]	672
3.87.7	Maxima [A] (verification not implemented)	673
3.87.8	Giac [A] (verification not implemented)	673
3.87.9	Mupad [B] (verification not implemented)	674

3.87.1 Optimal result

Integrand size = 34, antiderivative size = 108

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a+a\sec(e+fx)} dx = \frac{32c^3 \tan(e+fx)}{3af\sqrt{c-c\sec(e+fx)}} + \frac{8c^2\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{3af} + \frac{2c(c-c\sec(e+fx))^{3/2}\tan(e+fx)}{f(a+a\sec(e+fx))}$$

output

```
2*c*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))+32/3*c^3*tan(f*x+e)/a/f/(c-c*sec(f*x+e))^(1/2)+8/3*c^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/a/f
```

3.87.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.57

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a+a\sec(e+fx)} dx = \frac{2c^3(-23-10\sec(e+fx)+\sec^2(e+fx))\tan(e+fx)}{3af(1+\sec(e+fx))\sqrt{c-c\sec(e+fx)}}$$

input

```
Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x]),x]
```

output $(-2*c^3*(-23 - 10*Sec[e + f*x] + Sec[e + f*x]^2)*Tan[e + f*x])/(3*a*f*(1 + Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]])$

3.87.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4442, 3042, 4280, 3042, 4279}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{a\csc(e+fx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{4442} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)} - \frac{4c\int\sec(e+fx)(c-c\sec(e+fx))^{3/2}dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)} - \frac{4c\int\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}dx}{a} \\
 & \quad \downarrow \text{4280} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)} - \frac{4c\left(\frac{4}{3}c\int\sec(e+fx)\sqrt{c-c\sec(e+fx)}dx - \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{3f}\right)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)} - \frac{4c\left(\frac{4}{3}c\int\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}dx - \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{3f}\right)}{a} \\
 & \quad \downarrow \text{4279}
 \end{aligned}$$

3.87. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a+a\sec(e+fx)} dx$

$$\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{f(a \sec(e+fx) + a)} - \frac{4c \left(-\frac{8c^2 \tan(e+fx)}{3f \sqrt{c - c \sec(e+fx)}} - \frac{2c \tan(e+fx) \sqrt{c - c \sec(e+fx)}}{3f} \right)}{a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x]),x]`

output `(2*c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (4*c*((-8*c^2*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e + f*x]]) - (2*c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(3*f)))/a`

3.87.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4279 `Int[csc[(e_.) + (f_.)*(x_.)]*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] := Simp[-2*b*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4280 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] := Simp[(-b)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m - 1)/(f*m)), x] + Simp[a*((2*m - 1)/m) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m - 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && IntegerQ[2*m]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.87.4 Maple [A] (verified)

Time = 5.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.66

method	result	size
default	$-\frac{2c^2(\sec(fx+e)-1)^2\sqrt{-c(\sec(fx+e)-1)}(23\cos(fx+e)^2+10\cos(fx+e)-1)\cot(fx+e)}{3af(\cos(fx+e)-1)^2}$	71

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$-2/3/a/f*c^2*(\sec(f*x+e)-1)^2*(-c*(\sec(f*x+e)-1))^(1/2)*(23*\cos(f*x+e)^2+10*\cos(f*x+e)-1)/(\cos(f*x+e)-1)^2*\cot(f*x+e)$$

3.87.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.71

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a+a\sec(e+fx)} dx = \frac{2(23c^2\cos(fx+e)^2+10c^2\cos(fx+e)-c^2)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{3af\cos(fx+e)\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e)),x, algorithm="fricas")`

output
$$-2/3*(23*c^2*\cos(f*x+e)^2+10*c^2*\cos(f*x+e)-c^2)*\sqrt{(c*\cos(f*x+e)-c)/\cos(f*x+e)}/(a*f*\cos(f*x+e)*\sin(f*x+e))$$

3.87.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a+a\sec(e+fx)} dx = \frac{\int \frac{c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}{\sec(e+fx)+1} dx + \int \left(-\frac{2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}{\sec(e+fx)+1}\right)}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(5/2)/(a+a*sec(f*x+e)),x)`

3.87.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{a+a\sec(e+fx)} dx$$

output `(Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(-2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2/(sec(e + f*x) + 1), x) + Integral(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3/(sec(e + f*x) + 1), x))/a`

3.87.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.27

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{a + a \sec(e + fx)} dx =$$

$$\frac{4 \left(8 \sqrt{2} c^{5/2} - \frac{20 \sqrt{2} c^{5/2} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{15 \sqrt{2} c^{5/2} \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{3 \sqrt{2} c^{5/2} \sin(fx+e)^6}{(\cos(fx+e)+1)^6} \right)}{3af \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right)^{5/2} \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)^{5/2}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `-4/3*(8*sqrt(2)*c^(5/2) - 20*sqrt(2)*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 15*sqrt(2)*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 3*sqrt(2)*c^(5/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6)/(a*f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^(5/2)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^(5/2))`

3.87.8 Giac [A] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.78

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{a + a \sec(e + fx)} dx =$$

$$\frac{4 \sqrt{2} c^2 \left(\frac{3 \sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{a} - \frac{6 (c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c) c + c^2}{(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)^{3/2} a} \right)}{3f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e)),x, algorithm="giac")`

output
$$-4/3\sqrt{2}*c^2*(3\sqrt{c*\tan(1/2*f*x + 1/2*e)^2 - c})/a - (6*(c*\tan(1/2*f*x + 1/2*e)^2 - c)*c + c^2)/((c*\tan(1/2*f*x + 1/2*e)^2 - c)^{(3/2)*a})/f$$

3.87.9 Mupad [B] (verification not implemented)

Time = 15.56 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.16

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{a + a \sec(e + fx)} dx = \frac{2c^2 \sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}} (2 \sin(e + fx) - 44 \sin(2e + 2fx) + 25 \sin(3e + 3fx) - 26 \sin(4e + 4fx) + 23 \sin(5e + 5fx))}{3af(\cos(3e + 3fx) - 2\cos(e + fx) - 2\cos(4e + 4fx) + \cos(5e + 5fx) + 2)}$$

input `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output
$$(2c^2*((c*(\cos(e + f*x) - 1))/\cos(e + f*x))^{(1/2)}*(2*\sin(e + f*x) - 44*\sin(2*e + 2*f*x) + 25*\sin(3*e + 3*f*x) - 26*\sin(4*e + 4*f*x) + 23*\sin(5*e + 5*f*x)))/(3*a*f*(\cos(3*e + 3*f*x) - 2*\cos(e + f*x) - 2*\cos(4*e + 4*f*x) + \cos(5*e + 5*f*x) + 2))$$

$$3.88 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a+a\sec(e+fx)} dx$$

3.88.1	Optimal result	675
3.88.2	Mathematica [A] (verified)	675
3.88.3	Rubi [A] (verified)	676
3.88.4	Maple [A] (verified)	677
3.88.5	Fricas [A] (verification not implemented)	678
3.88.6	Sympy [F]	678
3.88.7	Maxima [A] (verification not implemented)	678
3.88.8	Giac [A] (verification not implemented)	679
3.88.9	Mupad [B] (verification not implemented)	679

3.88.1 Optimal result

Integrand size = 34, antiderivative size = 72

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a+a\sec(e+fx)} dx = \frac{4c^2 \tan(e+fx)}{af\sqrt{c-c\sec(e+fx)}} + \frac{2c\sqrt{c-c\sec(e+fx)} \tan(e+fx)}{f(a+a\sec(e+fx))}$$

output `4*c^2*tan(f*x+e)/a/f/(c-c*sec(f*x+e))^(1/2)+2*c*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))`

3.88.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.69

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a+a\sec(e+fx)} dx = \frac{2c^2(3+\sec(e+fx))\tan(e+fx)}{af(1+\sec(e+fx))\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x]),x]`

output `(2*c^2*(3 + Sec[e + f*x])*Tan[e + f*x])/(a*f*(1 + Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]])`

3.88. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a+a\sec(e+fx)} dx$

3.88.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4442, 3042, 4279}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{a\csc(e+fx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{4442} \\
 & \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)} - \frac{2c\int\sec(e+fx)\sqrt{c-c\sec(e+fx)}dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)} - \frac{2c\int\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}dx}{a} \\
 & \quad \downarrow \text{4279} \\
 & \frac{4c^2\tan(e+fx)}{af\sqrt{c-c\sec(e+fx)}} + \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x]),x]`

output `(4*c^2*Tan[e + f*x])/(a*f*Sqrt[c - c*Sec[e + f*x]]) + (2*c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))`

3.88.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4279 `Int[csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[-2*b*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.88.4 Maple [A] (verified)

Time = 2.82 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{2c(\sec(fx+e)-1)\sqrt{-c(\sec(fx+e)-1)}(3\cos(fx+e)+1)\cot(fx+e)}{af(\cos(fx+e)-1)}$	57

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$\frac{2}{a} \frac{1}{f} c * (\sec(f*x+e)-1) * (-c * (\sec(f*x+e)-1))^{1/2} * (3 * \cos(f*x+e)+1) / (\cos(f*x+e)-1) * \cot(f*x+e)$$

3.88.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.69

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a+a\sec(e+fx)} dx = -\frac{2(3c\cos(fx+e)+c)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{af\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e)),x, algorithm="fracas")
```

```
output -2*(3*c*cos(f*x + e) + c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(a*f*sin(f*x + e))
```

3.88.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a+a\sec(e+fx)} dx = \frac{\int \frac{c\sqrt{-c\sec(e+fx)+c\sec(e+fx)}}{\sec(e+fx)+1} dx + \int \left(-\frac{c\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}{\sec(e+fx)+1} \right) dx}{a}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(3/2)/(a+a*sec(f*x+e)),x)
```

```
output (Integral(c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2/(sec(e + f*x) + 1), x))/a
```

3.88.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.53

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{a+a\sec(e+fx)} dx = \frac{2\left(2\sqrt{2}c^{\frac{3}{2}} - \frac{3\sqrt{2}c^{\frac{3}{2}}\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{\sqrt{2}c^{\frac{3}{2}}\sin(fx+e)^4}{(\cos(fx+e)+1)^4}\right)}{af\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)^{\frac{3}{2}}\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)^{\frac{3}{2}}}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e)),x, algorithm="maxima")
```

output $2*(2*\text{sqrt}(2)*c^{(3/2)} - 3*\text{sqrt}(2)*c^{(3/2)}*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 + \text{sqrt}(2)*c^{(3/2)}*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4)/(a*f*(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)^{(3/2)}*(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)^{(3/2)})$

3.88.8 Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.83

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{a + a \sec(e + fx)} dx = -\frac{2\sqrt{2}\left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - cc}}{a} - \frac{c^2}{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - ca}}\right)}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e)),x, algorithm="giac")`

output $-2*\text{sqrt}(2)*(\text{sqrt}(c*\tan(1/2*f*x + 1/2*e)^2 - c)*c/a - c^2/(\text{sqrt}(c*\tan(1/2*f*x + 1/2*e)^2 - c)*a))/f$

3.88.9 Mupad [B] (verification not implemented)

Time = 13.69 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.07

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{a + a \sec(e + fx)} dx = \frac{c \sqrt{c - \frac{c}{\cos(e + fx)}} (2 \sin(e + fx) + 6 \sin(2e + 2fx) + 2 \sin(3e + 3fx) + 3 \sin(4e + 4fx))}{af \sin(2e + 2fx)^2}$$

input `int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output $-(c*(c - c/\cos(e + f*x))^{(1/2)}*(2*\sin(e + f*x) + 6*\sin(2*e + 2*f*x) + 2*\sin(3*e + 3*f*x) + 3*\sin(4*e + 4*f*x)))/(a*f*\sin(2*e + 2*f*x)^2)$

$$3.89 \quad \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a+a\sec(e+fx)} dx$$

3.89.1	Optimal result	680
3.89.2	Mathematica [A] (verified)	680
3.89.3	Rubi [A] (verified)	681
3.89.4	Maple [A] (verified)	682
3.89.5	Fricas [A] (verification not implemented)	682
3.89.6	Sympy [F]	682
3.89.7	Maxima [B] (verification not implemented)	683
3.89.8	Giac [A] (verification not implemented)	683
3.89.9	Mupad [B] (verification not implemented)	684

3.89.1 Optimal result

Integrand size = 34, antiderivative size = 39

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a+a\sec(e+fx)} dx = \frac{2c \tan(e+fx)}{f(a+a\sec(e+fx))\sqrt{c-c\sec(e+fx)}}$$

output `2*c*tan(f*x+e)/f/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2)`

3.89.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a+a\sec(e+fx)} dx = -\frac{2 \cot(e+fx)\sqrt{c-c\sec(e+fx)}}{af}$$

input `Integrate[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x]),x]`

output `(-2*Cot[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a*f)`

3.89.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a\sec(e+fx)+a} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{a\csc(e+fx+\frac{\pi}{2})+a} dx$$

↓ 4441

$$\frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x]),x]`

output `(2*c*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]])`

3.89.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.89.4 Maple [A] (verified)

Time = 3.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72

method	result	size
default	$-\frac{2\sqrt{-c(\sec(fx+e)-1)} \cot(fx+e)}{af}$	28

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e)),x,method=_RETURNVER
BOSE)`

output `-2/a/f*(-c*(sec(f*x+e)-1))^(1/2)*cot(f*x+e)`

3.89.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.15

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a+a\sec(e+fx)} dx = -\frac{2\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\cos(fx+e)}{af\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e)),x, algorithm=
"fracas")`

output `-2*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(a*f*sin(f*x + e))`

3.89.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a+a\sec(e+fx)} dx = \frac{\int \frac{\sqrt{-c\sec(e+fx)+c\sec(e+fx)}}{\sec(e+fx)+1} dx}{a}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(1/2)/(a+a*sec(f*x+e)),x)`

output `Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)/(sec(e + f*x) + 1), x)/a`

3.89.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. $2(37) = 74$.

Time = 0.30 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.15

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a+a\sec(e+fx)} dx = -\frac{\sqrt{2}\sqrt{c} - \frac{\sqrt{2}\sqrt{c}\sin(fx+e)^2}{(\cos(fx+e)+1)^2}}{af\sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1}\sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `-(sqrt(2)*sqrt(c) - sqrt(2)*sqrt(c)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)/(a*f*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) + 1)*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) - 1))`

3.89.8 Giac [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{a+a\sec(e+fx)} dx = \frac{\sqrt{2}\sqrt{c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c}\operatorname{sgn}\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)\operatorname{sgn}(\cos(fx+e))}{af}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e)),x, algorithm="giac")`

output `-sqrt(2)*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))*sgn(cos(f*x + e))/(a*f)`

3.89.9 Mupad [B] (verification not implemented)

Time = 13.16 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.03

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{a + a \sec(e + fx)} dx = -\frac{\sin(2e + 2fx) \sqrt{c - \frac{c}{\cos(e + fx)}}}{af \sin(e + fx)^2}$$

input `int((c - c/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`output `-(sin(2*e + 2*f*x)*(c - c/cos(e + f*x))^(1/2))/(a*f*sin(e + f*x)^2)`

3.90
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))\sqrt{c-c \sec(e+fx)}} dx$$

3.90.1	Optimal result	685
3.90.2	Mathematica [C] (verified)	685
3.90.3	Rubi [A] (verified)	686
3.90.4	Maple [A] (verified)	687
3.90.5	Fricas [A] (verification not implemented)	688
3.90.6	Sympy [F]	688
3.90.7	Maxima [F]	689
3.90.8	Giac [A] (verification not implemented)	689
3.90.9	Mupad [F(-1)]	690

3.90.1 Optimal result

Integrand size = 34, antiderivative size = 89

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))\sqrt{c-c \sec(e+fx)}} dx = -\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{\sqrt{2}a\sqrt{c}f} + \frac{\tan(e+fx)}{f(a+a \sec(e+fx))\sqrt{c-c \sec(e+fx)}}$$

output `-1/2*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a/f*2^(1/2)/c^(1/2)+tan(f*x+e)/f/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2)`

3.90.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.29 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.58

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))\sqrt{c-c \sec(e+fx)}} dx = \frac{\text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan\left(\frac{1}{2}(e+fx)\right)}{af\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]]),x]`

3.90.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))\sqrt{c-c \sec(e+fx)}} dx$$

output `(Hypergeometric2F1[-1/2, 1, 1/2, (1 + Sec[e + f*x])/2]*Tan[(e + f*x)/2])/ (a*f*Sqrt[c - c*Sec[e + f*x]])`

3.90.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.147$, Rules used = {3042, 4448, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx) + a) \sqrt{c - c \sec(e+fx)}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2}) + a) \sqrt{c - c \csc(e+fx+\frac{\pi}{2})}} dx$$

↓ 4448

$$\frac{\int \frac{\sec(e+fx)}{\sqrt{c - c \sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a) \sqrt{c - c \sec(e+fx)}}$$

↓ 3042

$$\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c - c \csc(e+fx+\frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a) \sqrt{c - c \sec(e+fx)}}$$

↓ 4282

$$\frac{\tan(e+fx)}{f(a \sec(e+fx) + a) \sqrt{c - c \sec(e+fx)}} - \frac{\int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c - c \sec(e+fx)} + 2c} d \frac{c \tan(e+fx)}{\sqrt{c - c \sec(e+fx)}}}{af}$$

↓ 216

$$\frac{\tan(e+fx)}{f(a \sec(e+fx) + a) \sqrt{c - c \sec(e+fx)}} - \frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2} \sqrt{c - c \sec(e+fx)}}\right)}{\sqrt{2} a \sqrt{c} f}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]]),x]`

3.90. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx)) \sqrt{c - c \sec(e+fx)}} dx$

output $-\frac{\text{ArcTan}[\sqrt{c}\tan(e+fx)]}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}/\sqrt{2a\sqrt{c}f} + \frac{\tan(e+fx)}{f(a+a\sec(e+fx))\sqrt{c-c\sec(e+fx)}}$

3.90.3.1 Defintions of rubi rules used

rule 216 $\text{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 3042 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4282 $\text{Int}[\text{csc}[(e_+) + (f_+)(x_+)]/\sqrt{\text{csc}[(e_+) + (f_+)(x_+)]*(b_+) + (a_+)}, x_Symbol] \rightarrow \text{Simp}[-2/f \ \text{Subst}[\text{Int}[1/(2*a + x^2), x], x, b*(\text{Cot}[e + f*x]/\sqrt{a + b*\text{Csc}[e + f*x]})], x] /; \text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

rule 4448 $\text{Int}[\text{csc}[(e_+) + (f_+)(x_+)]*(\text{csc}[(e_+) + (f_+)(x_+)]*(b_+) + (a_+))^{(m_+)}*(\text{csc}[(e_+) + (f_+)(x_+)]*(d_+) + (c_+))^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[b*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^m*((c + d*\text{Csc}[e + f*x])^n/(a*f*(2*m + 1))), x] + \text{Simp}[(m + n + 1)/(a*(2*m + 1)) \ \text{Int}[\text{Csc}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m + 1)}*(c + d*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ ((\text{ILtQ}[m, 0] \ \&\& \ \text{ILtQ}[n - 1/2, 0]) \ || \ (\text{ILtQ}[m - 1/2, 0] \ \&\& \ \text{ILtQ}[n - 1/2, 0] \ \&\& \ \text{LtQ}[m, n]))$

3.90.4 Maple [A] (verified)

Time = 3.00 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.26

method	result	size
default	$\frac{\sqrt{2} \sin(fx+e) \left(\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} + \arctan \left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} \right) \right)}{2af(\cos(fx+e)+1)\sqrt{-c(\sec(fx+e)-1)}\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}$	112

input `int(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `1/2/a/f*2^(1/2)*sin(f*x+e)*(2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)+arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)))/(cos(f*x+e)+1)/(-c*(sec(f*x+e)-1))^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)`

3.90.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 269, normalized size of antiderivative = 3.02

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))\sqrt{c-c\sec(e+fx)}} dx$$

$$= \frac{\sqrt{2}c\sqrt{-\frac{1}{c}} \log\left(-\frac{2\sqrt{2}(\cos(fx+e)^2+\cos(fx+e))\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\sqrt{-\frac{1}{c}}-(3\cos(fx+e)+1)\sin(fx+e)}{(\cos(fx+e)-1)\sin(fx+e)}\right) \sin(fx+e) - 4\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{4acf \sin(fx+e)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `[1/4*(sqrt(2)*c*sqrt(-1/c)*log(-(2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*sqrt(-1/c) - (3*cos(f*x + e) + 1)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e))*sin(f*x + e) - 4*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e))/(a*c*f*sin(f*x + e)), 1/2*(sqrt(2)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e))/(a*c*f*sin(f*x + e))]`

3.90.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))\sqrt{c-c\sec(e+fx)}} dx = \frac{\int \frac{\sec(e+fx)}{\sqrt{-c\sec(e+fx)+c\sec(e+fx)+\sqrt{-c\sec(e+fx)+c}}} dx}{a}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x)`

3.90. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))\sqrt{c-c\sec(e+fx)}} dx$

output `Integral(sec(e + f*x)/(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + sqrt(-c*sec(e + f*x) + c)), x)/a`

3.90.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{(a \sec(fx + e) + a)\sqrt{-c \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)), x)`

3.90.8 Giac [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.72

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))\sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{\sqrt{2} \left(\frac{\arctan\left(\frac{\sqrt{c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c}}{\sqrt{c}}\right)}{\sqrt{c}} - \frac{\sqrt{c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c}}{c} \right)}{2af}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `1/2*sqrt(2)*(arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/sqrt(c) - sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/c)/(a*f)`

3.90.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))\sqrt{c-c\sec(e+fx)}} dx$$

$$= \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right) \sqrt{c - \frac{c}{\cos(e+fx)}}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(1/2)),x)`output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(1/2)), x)`

3.91
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2}} dx$$

3.91.1 Optimal result 691
 3.91.2 Mathematica [C] (verified) 691
 3.91.3 Rubi [A] (verified) 692
 3.91.4 Maple [A] (verified) 694
 3.91.5 Fricas [A] (verification not implemented) 694
 3.91.6 Sympy [F] 695
 3.91.7 Maxima [F] 695
 3.91.8 Giac [A] (verification not implemented) 696
 3.91.9 Mupad [F(-1)] 696

3.91.1 Optimal result

Integrand size = 34, antiderivative size = 122

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2}} dx = -\frac{3 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{4\sqrt{2}ac^{3/2}f} - \frac{3 \tan(e+fx)}{4af(c-c \sec(e+fx))^{3/2}} + \frac{\tan(e+fx)}{f(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2}}$$

output `-3/8*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a/c^(3/2)/f*2^(1/2)-3/4*tan(f*x+e)/a/f/(c-c*sec(f*x+e))^(3/2)+tan(f*x+e)/f/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2)`

3.91.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.53 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.48

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2}} dx = \frac{\text{Hypergeometric2F1}\left(-\frac{1}{2}, 2, \frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan\left(\frac{1}{2}\right)}{2acf\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2)),x]`

output `(Hypergeometric2F1[-1/2, 2, 1/2, (1 + Sec[e + f*x])/2]*Tan[(e + f*x)/2])/ (2*a*c*f*Sqrt[c - c*Sec[e + f*x]])`

3.91.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4448, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)(c - c \sec(e+fx))^{3/2}} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2}) + a)(c - c \csc(e+fx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow 4448 \\
 & \frac{3 \int \frac{\sec(e+fx)}{(c - c \sec(e+fx))^{3/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c - c \csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 4283 \\
 & \frac{3 \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c - c \sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c - c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{3 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c - c \csc(e+fx+\frac{\pi}{2})}} dx}{4c} - \frac{\tan(e+fx)}{2f(c - c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 4282
 \end{aligned}$$

3.91. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2}} dx$

$$\begin{aligned}
& 3 \left(\frac{\int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c \sec(e+fx)} + 2c} d \frac{c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}}}{2cf} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right) \\
& \quad + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c-c \sec(e+fx))^{3/2}} \\
& \quad \quad \quad \downarrow \text{216} \\
& 3 \left(\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{2\sqrt{2}c^{3/2}f} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right) \\
& \quad + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c-c \sec(e+fx))^{3/2}}
\end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2)),x]`

output `Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2)) + (3*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2)))/(2*a)`

3.91.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4283 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

```
rule 4448 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_), x_Symbol] :> Simp[b*Cot[e + f*x]*
(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[
(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(
c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (IL
tQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.91.4 Maple [A] (verified)

Time = 3.10 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.57

method	result
default	$-\frac{\sqrt{2} \left(\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e) + 3 \arctan \left(\frac{\sqrt{2}}{2 \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} \right) \cos(fx+e) - 3 \sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} - 3 \arctan \left(\frac{\sqrt{2}}{2 \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} \right) \right)}{8af(\cos(fx+e)+1)c(\sec(fx+e)-1)\sqrt{-c(\sec(fx+e)-1)}\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2),x,method=_RETURNVER
BOSE)
```

```
output -1/8/a/f*2^(1/2)*(2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)+3*
arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))*cos(f*x+e)-3*2^(1/2)
)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)-3*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(co
s(f*x+e)+1))^(1/2)))/(cos(f*x+e)+1)/c/(sec(f*x+e)-1)/(-c*(sec(f*x+e)-1))^(
1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*tan(f*x+e)
```

3.91.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 329, normalized size of antiderivative = 2.70

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2}} dx = \left[-\frac{3 \sqrt{2} \sqrt{-c} (\cos(fx + e) - 1) \log \left(\frac{2 \sqrt{2} (\cos(fx + e)^2 + \cos(fx + e) + 1)}{\dots} \right)}{\dots} \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2),x, algorithm=
"fricas")
```

3.91. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{3/2}} dx$

output `[-1/16*(3*sqrt(2)*sqrt(-c)*(cos(f*x + e) - 1)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e))*sin(f*x + e) + 4*(cos(f*x + e)^2 - 3*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a*c^2*f*cos(f*x + e) - a*c^2*f)*sin(f*x + e)), 1/8*(3*sqrt(2)*sqrt(c)*(cos(f*x + e) - 1)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(cos(f*x + e)^2 - 3*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a*c^2*f*cos(f*x + e) - a*c^2*f)*sin(f*x + e))]`

3.91.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2}} dx = \frac{\int \frac{\sec(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)+c\sqrt{-c\sec(e+fx)+c}} dx}{a}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))**(3/2), x)`

output `Integral(sec(e + f*x)/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 + c*sqrt(-c*sec(e + f*x) + c)), x)/a`

3.91.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2}} dx = \int \frac{\sec(fx + e)}{(a \sec(fx + e) + a)(-c \sec(fx + e) + c)^{3/2}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2), x, algorithm="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)*(-c*sec(f*x + e) + c)^(3/2)), x)`

3.91.8 Giac [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.80

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2}} dx = \frac{\sqrt{2} \left(3 \sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}} \right) - 2 \sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c} \right)}{8 a c^2 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `1/8*sqrt(2)*(3*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) - 2*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c) - sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/tan(1/2*f*x + 1/2*e)^2)/(a*c^2*f)`

3.91.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^{3/2}} dx = \int \frac{1}{\cos(e + fx) \left(a + \frac{a}{\cos(e + fx)} \right) \left(c - \frac{c}{\cos(e + fx)} \right)^{3/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(3/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(3/2)), x)`

$$3.92 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{5/2}} dx$$

3.92.1	Optimal result	697
3.92.2	Mathematica [C] (verified)	697
3.92.3	Rubi [A] (verified)	698
3.92.4	Maple [A] (verified)	701
3.92.5	Fricas [A] (verification not implemented)	701
3.92.6	Sympy [F]	702
3.92.7	Maxima [F]	702
3.92.8	Giac [A] (verification not implemented)	703
3.92.9	Mupad [F(-1)]	703

3.92.1 Optimal result

Integrand size = 34, antiderivative size = 156

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{5/2}} dx =$$

$$-\frac{15 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{32\sqrt{2}ac^{5/2}f} - \frac{5 \tan(e+fx)}{8af(c-c \sec(e+fx))^{5/2}}$$

$$+ \frac{\tan(e+fx)}{f(a+a \sec(e+fx))(c-c \sec(e+fx))^{5/2}} - \frac{15 \tan(e+fx)}{32acf(c-c \sec(e+fx))^{3/2}}$$

output `-15/64*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a/c^(5/2)/f*2^(1/2)-5/8*tan(f*x+e)/a/f/(c-c*sec(f*x+e))^(5/2)+tan(f*x+e)/f/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2)-15/32*tan(f*x+e)/a/c/f/(c-c*sec(f*x+e))^(3/2)`

3.92.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.59 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.37

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c-c \sec(e+fx))^{5/2}} dx = \frac{\text{Hypergeometric2F1}\left(-\frac{1}{2}, 3, \frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan\left(\frac{1}{2}\right)}{4ac^2 f \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2)),x
]`

output `(Hypergeometric2F1[-1/2, 3, 1/2, (1 + Sec[e + f*x])/2]*Tan[(e + f*x)/2])/(
4*a*c^2*f*Sqrt[c - c*Sec[e + f*x]])`

3.92.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4448, 3042, 4283, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)(c - c \sec(e+fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2}) + a)(c - c \csc(e+fx+\frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4448} \\
 & \frac{5 \int \frac{\sec(e+fx)}{(c - c \sec(e+fx))^{5/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c - c \csc(e+fx+\frac{\pi}{2}))^{5/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4283} \\
 & \frac{5 \left(\frac{3 \int \frac{\sec(e+fx)}{(c - c \sec(e+fx))^{3/2}} dx}{8c} - \frac{\tan(e+fx)}{4f(c - c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{5 \left(\frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c \csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c-c \sec(e+fx))^{5/2}} \\
& \quad \downarrow 4283 \\
& \frac{5 \left(\frac{3 \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c-c \sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \\
& \quad \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c-c \sec(e+fx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{5 \left(\frac{3 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \\
& \quad \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c-c \sec(e+fx))^{5/2}} \\
& \quad \downarrow 4282 \\
& \frac{5 \left(\frac{3 \left(\frac{\int \frac{1}{c^2 \tan^2(e+fx)} d \frac{c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}}}{2cf} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \\
& \quad \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c-c \sec(e+fx))^{5/2}} \\
& \quad \downarrow 216
\end{aligned}$$

$$\frac{5 \left(\frac{3 \left(-\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{2\sqrt{2}c^{3/2}f} - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c\sec(e+fx))^{5/2}} \right)}{\frac{2a \tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}} +$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2)),x]`

output `Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2)) + (5*(-1/4*Tan[e + f*x]/(f*(c - c*Sec[e + f*x])^(5/2)) + (3*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2))))/(8*c)))/(2*a)`

3.92.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4283 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

```
rule 4448 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.92.4 Maple [A] (verified)

Time = 3.31 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.72

method	result
default	$\frac{\sqrt{2} \left(3\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e)^2 + 20\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e) - 15 \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) \cos(fx+e)^2 - 15\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \right)}{64af(\cos(fx+e)+1)c^2(\sec(fx+e)-1)^2\sqrt{-c}}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)
```

```
output -1/64/a/f*2^(1/2)*(3*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2+20*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)-15*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))*cos(f*x+e)^2-15*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)+30*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))*cos(f*x+e)-15*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)))/(cos(f*x+e)+1)/c^2/(sec(f*x+e)-1)^2/(-c*(sec(f*x+e)-1))^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*tan(f*x+e)*sec(f*x+e)
```

3.92.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 401, normalized size of antiderivative = 2.57

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c - c \sec(e + fx))^{5/2}} dx = \left[-\frac{15\sqrt{2}(\cos(fx + e)^2 - 2\cos(fx + e) + 1)\sqrt{-c} \log\left(\frac{2}{\dots}\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2),x, algorithm="fricas")`

output `[-1/128*(15*sqrt(2)*(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)*sqrt(-c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e))) * sin(f*x + e) - 4*(3*cos(f*x + e)^3 + 20*cos(f*x + e)^2 - 15*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a*c^3*f*cos(f*x + e)^2 - 2*a*c^3*f*cos(f*x + e) + a*c^3*f)*sin(f*x + e)), 1/64*(15*sqrt(2)*(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e))) * sin(f*x + e) + 2*(3*cos(f*x + e)^3 + 20*cos(f*x + e)^2 - 15*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a*c^3*f*cos(f*x + e)^2 - 2*a*c^3*f*cos(f*x + e) + a*c^3*f)*sin(f*x + e))]`

3.92.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\sec(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^3(e+fx)-c^2}\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)-c^2}\sqrt{a}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2),x)`

output `Integral(sec(e + f*x)/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3 - c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 - c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f*x) + c)), x)/a`

3.92.7 Maxima [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\sec(fx+e)}{(a\sec(fx+e)+a)(-c\sec(fx+e)+c)^{5/2}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)*(-c*sec(f*x + e) + c)^(5/2)), x)`

3.92. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^{5/2}} dx$

3.92.8 Giac [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.82

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^{5/2}} dx = \frac{\sqrt{2} \left(15 \sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}} \right) - 8 \sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c} \right)}{64 a^3 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `1/64*sqrt(2)*(15*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) - 8*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c) - (9*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c + 7*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^2)/(c^2*tan(1/2*f*x + 1/2*e)^4)/(a*c^3*f)`

3.92.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c-c\sec(e+fx))^{5/2}} dx = \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)} \right) \left(c - \frac{c}{\cos(e+fx)} \right)^{5/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(5/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c - c/cos(e + f*x))^(5/2)), x)`

3.93
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx$$

3.93.1	Optimal result	704
3.93.2	Mathematica [A] (verified)	704
3.93.3	Rubi [A] (verified)	705
3.93.4	Maple [A] (verified)	707
3.93.5	Fricas [A] (verification not implemented)	707
3.93.6	Sympy [F(-1)]	708
3.93.7	Maxima [A] (verification not implemented)	708
3.93.8	Giac [A] (verification not implemented)	709
3.93.9	Mupad [B] (verification not implemented)	709

3.93.1 Optimal result

Integrand size = 34, antiderivative size = 155

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx =$$

$$-\frac{64c^4 \tan(e+fx)}{3a^2 f \sqrt{c-c\sec(e+fx)}} - \frac{16c^3 \sqrt{c-c\sec(e+fx)} \tan(e+fx)}{3a^2 f}$$

$$- \frac{4c^2(c-c\sec(e+fx))^{3/2} \tan(e+fx)}{f(a^2+a^2\sec(e+fx))} + \frac{2c(c-c\sec(e+fx))^{5/2} \tan(e+fx)}{3f(a+a\sec(e+fx))^2}$$

output `-4*c^2*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))+2/3*c*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^2-64/3*c^4*tan(f*x+e)/a^2/f/(c-c*sec(f*x+e))^(1/2)-16/3*c^3*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/a^2/f`

3.93.2 Mathematica [A] (verified)

Time = 1.51 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.46

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx = \frac{2c^4(-45-69\sec(e+fx)-15\sec^2(e+fx)+\sec^3(e+fx))\tan(e+fx)}{3a^2 f(1+\sec(e+fx))^2 \sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(7/2))/(a + a*Sec[e + f*x])^2, x]`

output $(2*c^4*(-45 - 69*Sec[e + f*x] - 15*Sec[e + f*x]^2 + Sec[e + f*x]^3)*Tan[e + f*x])/(3*a^2*f*(1 + Sec[e + f*x])^2*sqrt[c - c*Sec[e + f*x]])$

3.93.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4442, 3042, 4442, 3042, 4280, 3042, 4279}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a\sec(e+fx)+a)^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{7/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx$$

↓ 4442

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{3f(a\sec(e+fx)+a)^2} - \frac{2c\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sec(e+fx)a+a} dx}{a}$$

↓ 3042

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{3f(a\sec(e+fx)+a)^2} - \frac{2c\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{a}$$

↓ 4442

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{3f(a\sec(e+fx)+a)^2} - \frac{2c\left(\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)} - \frac{4c\int \sec(e+fx)(c-c\sec(e+fx))^{3/2} dx}{a}\right)}{a}$$

↓ 3042

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{5/2}}{3f(a\sec(e+fx)+a)^2} - \frac{2c\left(\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)} - \frac{4c\int \csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2} dx}{a}\right)}{a}$$

↓ 4280

3.93. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx$

$$\frac{2c \tan(e + fx)(c - c \sec(e + fx))^{5/2}}{3f(a \sec(e + fx) + a)^2} - \frac{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}}{f(a \sec(e + fx) + a)} - \frac{4c \left(\frac{4}{3} \int \sec(e + fx) \sqrt{c - c \sec(e + fx)} dx - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)}}{3f} \right)}{a}$$

\downarrow 3042

$$\frac{2c \tan(e + fx)(c - c \sec(e + fx))^{5/2}}{3f(a \sec(e + fx) + a)^2} - \frac{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}}{f(a \sec(e + fx) + a)} - \frac{4c \left(\frac{4}{3} \int \csc(e + fx + \frac{\pi}{2}) \sqrt{c - c \csc(e + fx + \frac{\pi}{2})} dx - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)}}{3f} \right)}{a}$$

\downarrow 4279

$$\frac{2c \tan(e + fx)(c - c \sec(e + fx))^{5/2}}{3f(a \sec(e + fx) + a)^2} - \frac{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}}{f(a \sec(e + fx) + a)} - \frac{4c \left(-\frac{8c^2 \tan(e + fx)}{3f \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)}}{3f} \right)}{a}$$

```
input Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(7/2))/(a + a*Sec[e + f*x])^2,x]
```

```
output (2*c*(c - c*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (2*c*((2*c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) - (4*c*((-8*c^2*Tan[e + f*x])/(3*f*Sqrt[c - c*Sec[e + f*x]]) - (2*c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(3*f))))/a)/a
```

3.93.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]
```

```
rule 4279 Int[csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*b*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]])), x] /; Free Q[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]
```

```
rule 4280 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_), x_
Symbol] :> Simp[(-b)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m - 1)/(f*m)), x]
+ Simp[a*((2*m - 1)/m) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m - 1), x],
x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && IntegerQ[2*m]
```

```
rule 4442 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc
[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[2*a*c*Cot[e +
f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))),
x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*
x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[
m, -2^(-1)]
```

3.93.4 Maple [A] (verified)

Time = 12.58 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.57

method	result	size
default	$\frac{2\sqrt{-c(\sec(fx+e)-1)}(\sec(fx+e)-1)^3 c^3(3\cos(fx+e)+1)(15\cos(fx+e)^2+18\cos(fx+e)-1)\cot(fx+e)^2 \csc(fx+e)}{3a^2 f(\cos(fx+e)-1)^2}$	89

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^2,x,method=_RETURNV
ERBOSE)
```

```
output 2/3/a^2/f*(-c*(sec(f*x+e)-1))^(1/2)*(sec(f*x+e)-1)^3*c^3*(3*cos(f*x+e)+1)*
(15*cos(f*x+e)^2+18*cos(f*x+e)-1)/(cos(f*x+e)-1)^2*cot(f*x+e)^2*csc(f*x+e)
```

3.93.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx = \frac{2(45c^3\cos(fx+e)^3+69c^3\cos(fx+e)^2+15c^3\cos(fx+e)-3(a^2f\cos(fx+e)^2+a^2f\cos(fx+e))\sin(fx+e))}{3(a^2f\cos(fx+e)^2+a^2f\cos(fx+e))\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^2,x, algorithm
m="fricas")
```

3.93.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx$$

output $2/3*(45*c^3*\cos(f*x + e)^3 + 69*c^3*\cos(f*x + e)^2 + 15*c^3*\cos(f*x + e) - c^3)*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)} / ((a^2*f*\cos(f*x + e)^2 + a^2*f*\cos(f*x + e))*\sin(f*x + e))$

3.93.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{7/2}}{(a + a \sec(e + fx))^2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(7/2)/(a+a*sec(f*x+e))**2,x)`

output Timed out

3.93.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.21

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{7/2}}{(a + a \sec(e + fx))^2} dx =$$

$$\frac{4 \left(16 \sqrt{2} c^{7/2} - \frac{56 \sqrt{2} c^{7/2} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{70 \sqrt{2} c^{7/2} \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{35 \sqrt{2} c^{7/2} \sin(fx+e)^6}{(\cos(fx+e)+1)^6} + \frac{4 \sqrt{2} c^{7/2} \sin(fx+e)^8}{(\cos(fx+e)+1)^8} + \frac{\sqrt{2} c^{7/2} \sin(fx+e)^{10}}{(\cos(fx+e)+1)^{10}} \right)}{3 a^2 f \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right)^{7/2} \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)^{7/2}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output $-4/3*(16*\sqrt{2}*c^{(7/2)} - 56*\sqrt{2}*c^{(7/2)}*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 + 70*\sqrt{2}*c^{(7/2)}*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 - 35*\sqrt{2}*c^{(7/2)}*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6 + 4*\sqrt{2}*c^{(7/2)}*\sin(f*x + e)^8/(\cos(f*x + e) + 1)^8 + \sqrt{2}*c^{(7/2)}*\sin(f*x + e)^{10}/(\cos(f*x + e) + 1)^{10}) / (a^2*f*(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)^{(7/2)}*(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)^{(7/2)})$

3.93.8 Giac [A] (verification not implemented)

Time = 0.93 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.78

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx = \frac{4\sqrt{2}c^3 \left(\frac{9(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)c+c^2}{(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)^{\frac{3}{2}}a^2} - \frac{(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)^{\frac{3}{2}}a^4c^2+9\sqrt{c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-ca^4c^3}}{a^6c^3} \right)}{3f}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^2,x, algorithm
m="giac")
```

```
output -4/3*sqrt(2)*c^3*((9*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c + c^2)/((c*tan(1/2*f
*x + 1/2*e)^2 - c)^(3/2)*a^2) - ((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*a^4*
c^2 + 9*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*a^4*c^3)/(a^6*c^3))/f
```

3.93.9 Mupad [B] (verification not implemented)

Time = 17.74 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.21

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^2} dx = \frac{2c^3 \sqrt{c - \frac{c}{\frac{e^{-e1i-fx1i}}{2} + \frac{e^{e1i+fx1i}}{2}}}}{3a^2 f (e^{e1i+fx1i} + 1)^3 (e^{e1i+fx1i} + e^{e2i+fx2i} + 195i + e^{e3i+fx3i} + 268i + e^{e4i+fx4i} + 195i + e^{e5i+fx5i} + 138i + e^{e6i+fx6i} + 45i)}$$

```
input int((c - c/cos(e + f*x))^(7/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)
```

```
output (2*c^3*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*(exp(
e*1i + f*x*1i)*138i + exp(e*2i + f*x*2i)*195i + exp(e*3i + f*x*3i)*268i +
exp(e*4i + f*x*4i)*195i + exp(e*5i + f*x*5i)*138i + exp(e*6i + f*x*6i)*45i
+ 45i))/(3*a^2*f*(exp(e*1i + f*x*1i) + 1)^3*(exp(e*1i + f*x*1i) - exp(e*2
i + f*x*2i) + exp(e*3i + f*x*3i) - 1))
```

3.94 $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx$

3.94.1	Optimal result	710
3.94.2	Mathematica [A] (verified)	710
3.94.3	Rubi [A] (verified)	711
3.94.4	Maple [A] (verified)	712
3.94.5	Fricas [A] (verification not implemented)	713
3.94.6	Sympy [F(-1)]	713
3.94.7	Maxima [A] (verification not implemented)	714
3.94.8	Giac [A] (verification not implemented)	714
3.94.9	Mupad [B] (verification not implemented)	715

3.94.1 Optimal result

Integrand size = 34, antiderivative size = 123

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx = -\frac{16c^3 \tan(e+fx)}{3a^2 f \sqrt{c-c\sec(e+fx)}} - \frac{8c^2 \sqrt{c-c\sec(e+fx)} \tan(e+fx)}{3f(a^2+a^2\sec(e+fx))} + \frac{2c(c-c\sec(e+fx))^{3/2} \tan(e+fx)}{3f(a+a\sec(e+fx))^2}$$

output `2/3*c*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^2-16/3*c^3*tan(f*x+e)/a^2/f/(c-c*sec(f*x+e))^(1/2)-8/3*c^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))`

3.94.2 Mathematica [A] (verified)

Time = 0.90 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.52

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx = -\frac{2c^3(11+18\sec(e+fx)+3\sec^2(e+fx))\tan(e+fx)}{3a^2 f(1+\sec(e+fx))^2 \sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^2, x]`

output $(-2*c^3*(11 + 18*\text{Sec}[e + f*x] + 3*\text{Sec}[e + f*x]^2)*\text{Tan}[e + f*x])/(3*a^2*f*(1 + \text{Sec}[e + f*x])^2*\text{Sqrt}[c - c*\text{Sec}[e + f*x]])$

3.94.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4442, 3042, 4442, 3042, 4279}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a\sec(e+fx)+a)^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx$$

↓ 4442

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{3f(a\sec(e+fx)+a)^2} - \frac{4c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sec(e+fx)a+a} dx}{3a}$$

↓ 3042

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{3f(a\sec(e+fx)+a)^2} - \frac{4c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a}$$

↓ 4442

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{3f(a\sec(e+fx)+a)^2} - \frac{4c \left(\frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)} - \frac{2c \int \sec(e+fx)\sqrt{c-c\sec(e+fx)} dx}{a} \right)}{3a}$$

↓ 3042

$$\frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{3f(a\sec(e+fx)+a)^2} - 4c \left(\frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)} - \frac{2c \int \csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})} dx}{a} \right)$$

↓ 4279

$$\frac{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}}{3f(a \sec(e + fx) + a)^2} - \frac{4c \left(\frac{4c^2 \tan(e + fx)}{af \sqrt{c - c \sec(e + fx)}} + \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)}}{f(a \sec(e + fx) + a)} \right)}{3a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^2,x]`

output `(2*c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (4*c*((4*c^2*Tan[e + f*x])/(a*f*Sqrt[c - c*Sec[e + f*x]]) + (2*c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))))/(3*a)`

3.94.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4279 `Int[csc[(e_.) + (f_.)*(x_.)]*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] := Simp[-2*b*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.94.4 Maple [A] (verified)

Time = 12.52 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.64

method	result	size
default	$-\frac{2\sqrt{-c(\sec(fx+e)-1)}(\sec(fx+e)-1)^2c^2(11\cos(fx+e)^2+18\cos(fx+e)+3)\cot(fx+e)^2\csc(fx+e)}{3a^2f(\cos(fx+e)-1)}$	79

3.94. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^2,x,method=_RETURNV
ERBOSE)`

output
$$-2/3/a^2/f*(-c*(\sec(f*x+e)-1))^{(1/2)*(\sec(f*x+e)-1)^2*c^2*(11*\cos(f*x+e)^2+18*\cos(f*x+e)+3)/(\cos(f*x+e)-1)*\cot(f*x+e)^2*\csc(f*x+e)}$$

3.94.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx = \frac{2(11c^2\cos^2(fx+e)+18c^2\cos(fx+e)+3c^2)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{3(a^2f\cos(fx+e)+a^2f)\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^2,x, algorithm
m="fricas")`

output
$$2/3*(11*c^2*\cos(f*x + e)^2 + 18*c^2*\cos(f*x + e) + 3*c^2)*\text{sqrt}((c*\cos(f*x + e) - c)/\cos(f*x + e))/((a^2*f*\cos(f*x + e) + a^2*f)*\sin(f*x + e))$$

3.94.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(5/2)/(a+a*sec(f*x+e))**2,x)`

output `Timed out`

3.94.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.33

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx = \frac{2 \left(8\sqrt{2}c^{5/2} - \frac{20\sqrt{2}c^{5/2}\sin^2(fx+e)}{(\cos(fx+e)+1)^2} + \frac{15\sqrt{2}c^{5/2}\sin^4(fx+e)}{(\cos(fx+e)+1)^4} - \frac{2\sqrt{2}c^{5/2}\sin^6(fx+e)}{(\cos(fx+e)+1)^6} \right)}{3a^2f \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right)^{5/2} \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)^{5/2}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^2,x, algorithm m="maxima")`

output `2/3*(8*sqrt(2)*c^(5/2) - 20*sqrt(2)*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 15*sqrt(2)*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 2*sqrt(2)*c^(5/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - sqrt(2)*c^(5/2)*sin(f*x + e)^8/(cos(f*x + e) + 1)^8)/(a^2*f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^(5/2)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^(5/2))`

3.94.8 Giac [A] (verification not implemented)

Time = 0.81 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.80

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx = \frac{2\sqrt{2}c^2 \left(\frac{3c}{\sqrt{c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-ca^2}} - \frac{(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)^{3/2}a^4c^2+6\sqrt{c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-ca^4c^3}}{a^6c^3} \right)}{3f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^2,x, algorithm m="giac")`

output `-2/3*sqrt(2)*c^2*(3*c/(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*a^2) - ((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*a^4*c^2 + 6*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*a^4*c^3)/(a^6*c^3))/f`

3.94.9 Mupad [B] (verification not implemented)

Time = 17.27 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.11

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^2} dx = \frac{2c^2 \sqrt{c - \frac{e^{-e-1i-fx1i}c}{2} + \frac{e^{e1i+fx1i}}{2}}}{3a^2 f (e^{e1i+fx1i} - 1) (e^{e1i+fx1i} + 1)} (e^{e1i+fx1i} 36i + e^{e2i+fx2i} 34i + e^{e3i+fx3i} 32i + e^{e4i+fx4i} 30i + e^{e5i+fx5i} 28i + e^{e6i+fx6i} 26i + e^{e7i+fx7i} 24i + e^{e8i+fx8i} 22i + e^{e9i+fx9i} 20i + e^{e10i+fx10i} 18i + e^{e11i+fx11i} 16i + e^{e12i+fx12i} 14i + e^{e13i+fx13i} 12i + e^{e14i+fx14i} 10i + e^{e15i+fx15i} 8i + e^{e16i+fx16i} 6i + e^{e17i+fx17i} 4i + e^{e18i+fx18i} 2i)$$

input `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output `(2*c^2*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*(exp(e*1i + f*x*1i)*36i + exp(e*2i + f*x*2i)*34i + exp(e*3i + f*x*3i)*32i + exp(e*4i + f*x*4i)*30i + exp(e*5i + f*x*5i)*28i + exp(e*6i + f*x*6i)*26i + exp(e*7i + f*x*7i)*24i + exp(e*8i + f*x*8i)*22i + exp(e*9i + f*x*9i)*20i + exp(e*10i + f*x*10i)*18i + exp(e*11i + f*x*11i)*16i + exp(e*12i + f*x*12i)*14i + exp(e*13i + f*x*13i)*12i + exp(e*14i + f*x*14i)*10i + exp(e*15i + f*x*15i)*8i + exp(e*16i + f*x*16i)*6i + exp(e*17i + f*x*17i)*4i + exp(e*18i + f*x*18i)*2i)/(3*a^2*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^3)`

3.95
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx$$

3.95.1	Optimal result	716
3.95.2	Mathematica [A] (verified)	716
3.95.3	Rubi [A] (verified)	717
3.95.4	Maple [A] (verified)	718
3.95.5	Fricas [A] (verification not implemented)	719
3.95.6	Sympy [F]	719
3.95.7	Maxima [A] (verification not implemented)	719
3.95.8	Giac [A] (verification not implemented)	720
3.95.9	Mupad [B] (verification not implemented)	720

3.95.1 Optimal result

Integrand size = 34, antiderivative size = 89

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx = -\frac{4c^2 \tan(e+fx)}{3f(a^2+a^2\sec(e+fx))\sqrt{c-c\sec(e+fx)}} + \frac{2c\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{3f(a+a\sec(e+fx))^2}$$

output `-4/3*c^2*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2)+2/3*c*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^2`

3.95.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.61

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx = -\frac{2c^2(1+3\sec(e+fx))\tan(e+fx)}{3a^2f(1+\sec(e+fx))^2\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x])^2, x]`

output `(-2*c^2*(1 + 3*Sec[e + f*x])*Tan[e + f*x])/(3*a^2*f*(1 + Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]])`

3.95.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx$$

3.95.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.99, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4442, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a\sec(e+fx)+a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx \\
 & \quad \downarrow \text{4442} \\
 & \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{3f(a\sec(e+fx)+a)^2} - \frac{2c \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sec(e+fx)a+a} dx}{3a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{3f(a\sec(e+fx)+a)^2} - \frac{2c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} \\
 & \quad \downarrow \text{4441} \\
 & \frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{3f(a\sec(e+fx)+a)^2} - \frac{4c^2 \tan(e+fx)}{3af(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x])^2,x]`

output `(-4*c^2*Tan[e + f*x])/(3*a*f*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]]) + (2*c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2)`

3.95.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.95.4 Maple [A] (verified)

Time = 3.38 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.60

method	result	size
default	$\frac{2\sqrt{-c(\sec(fx+e)-1)}(\sec(fx+e)-1)c(\cos(fx+e)+3)\cot(fx+e)^2\csc(fx+e)}{3a^2f}$	53

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^2,x,method=_RETURNV ERBOSE)`

output `2/3/a^2/f*(-c*(sec(f*x+e)-1))^(1/2)*(sec(f*x+e)-1)*c*(cos(f*x+e)+3)*cot(f*x+e)^2*csc(f*x+e)`

3.95.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx$$

3.95.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.81

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx = \frac{2(c\cos(fx+e)^2+3c\cos(fx+e))\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{3(a^2f\cos(fx+e)+a^2f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^2,x, algorithm
m="fricas")
```

```
output 2/3*(c*cos(f*x + e)^2 + 3*c*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*
x + e))/((a^2*f*cos(f*x + e) + a^2*f)*sin(f*x + e))
```

3.95.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx = \frac{\int \frac{c\sqrt{-c\sec(e+fx)+c\sec(e+fx)}}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \left(-\frac{c\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)}}{\sec^2(e+fx)+2\sec(e+fx)+1}\right) dx}{a^2}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(3/2)/(a+a*sec(f*x+e))**2,x)
```

```
output (Integral(c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)/(sec(e + f*x)**2 + 2*se
c(e + f*x) + 1), x) + Integral(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**
2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2
```

3.95.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.24

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^2} dx = -\frac{2\sqrt{2}c^{\frac{3}{2}} - \frac{3\sqrt{2}c^{\frac{3}{2}}\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{\sqrt{2}c^{\frac{3}{2}}\sin(fx+e)^6}{(\cos(fx+e)+1)^6}}{3a^2f\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)^{\frac{3}{2}}\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)^{\frac{3}{2}}}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^2,x, algorithm
m="maxima")
```


output
$$-1/3*(2*\sqrt{2}*c^{(3/2)} - 3*\sqrt{2}*c^{(3/2)}*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 + \sqrt{2}*c^{(3/2)}*\sin(f*x + e)^6/(\cos(f*x + e) + 1)^6)/(a^2*f*(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)^{(3/2)}*(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)^{(3/2)})$$

3.95.8 Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{(a + a \sec(e + fx))^2} dx = \frac{\sqrt{2}(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)^{\frac{3}{2}}}{a^2} + \frac{3\sqrt{2}\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - cc}}{3f a^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output
$$1/3*(\sqrt{2}*(c*\tan(1/2*f*x + 1/2*e)^2 - c)^{(3/2)}/a^2 + 3*\sqrt{2}*\sqrt{c*\tan(1/2*f*x + 1/2*e)^2 - c}*c/a^2)/f$$

3.95.9 Mupad [B] (verification not implemented)

Time = 16.88 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{(a + a \sec(e + fx))^2} dx = \frac{2c \sqrt{c - \frac{c}{\frac{e^{-e \cdot 1i - f \cdot x \cdot 1i}}{2} + \frac{e^{e \cdot 1i + f \cdot x \cdot 1i}}{2}}}}{3a^2 f (e^{e \cdot 1i + f \cdot x \cdot 1i} - 1) (e^{e \cdot 1i + f \cdot x \cdot 1i} + 1)^3} (e^{e \cdot 1i + f \cdot x \cdot 1i} \cdot 6i + e^{e \cdot 2i + f \cdot x \cdot 2i} \cdot 2i + e^{e \cdot 3i + f \cdot x \cdot 3i})$$

input `int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output
$$(2*c*(c - c/(\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2))^{(1/2)}*(\exp(e*1i + f*x*1i)*6i + \exp(e*2i + f*x*2i)*2i + \exp(e*3i + f*x*3i)*6i + \exp(e*4i + f*x*4i)*1i + 1i))/(3*a^2*f*(\exp(e*1i + f*x*1i) - 1)*(\exp(e*1i + f*x*1i) + 1)^3)$$

3.96
$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx$$

3.96.1 Optimal result 721
 3.96.2 Mathematica [A] (verified) 721
 3.96.3 Rubi [A] (verified) 722
 3.96.4 Maple [A] (verified) 723
 3.96.5 Fricas [A] (verification not implemented) 723
 3.96.6 Sympy [F] 723
 3.96.7 Maxima [B] (verification not implemented) 724
 3.96.8 Giac [A] (verification not implemented) 724
 3.96.9 Mupad [B] (verification not implemented) 725

3.96.1 Optimal result

Integrand size = 34, antiderivative size = 41

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx = \frac{2c \tan(e+fx)}{3f(a+a\sec(e+fx))^2 \sqrt{c-c\sec(e+fx)}}$$

output `2/3*c*tan(f*x+e)/f/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2)`

3.96.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.34

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx = -\frac{\cos^2(e+fx) \csc\left(\frac{1}{2}(e+fx)\right) \sec^3\left(\frac{1}{2}(e+fx)\right) \sqrt{c-c\sec(e+fx)}}{6a^2f}$$

input `Integrate[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^2,x]`

output `-1/6*(Cos[e + f*x]^2*Csc[(e + f*x)/2]*Sec[(e + f*x)/2]^3*Sqrt[c - c*Sec[e + f*x]])/(a^2*f)`

3.96.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a\sec(e+fx)+a)^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx$$

↓ 4441

$$\frac{2c\tan(e+fx)}{3f(a\sec(e+fx)+a)^2\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^2,x]`

output `(2*c*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]])`

3.96.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.96.4 Maple [A] (verified)

Time = 3.70 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.07

method	result	size
default	$-\frac{2\sqrt{-c(\sec(fx+e)-1)} \cos(fx+e) \cot(fx+e)}{3a^2 f (\cos(fx+e)+1)}$	44

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^2,x,method=_RETURNV
ERBOSE)
```

```
output -2/3/a^2/f*(-c*(sec(f*x+e)-1))^(1/2)/(cos(f*x+e)+1)*cos(f*x+e)*cot(f*x+e)
```

3.96.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx = -\frac{2\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}} \cos(fx+e)^2}{3(a^2 f \cos(fx+e) + a^2 f) \sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^2,x, algorithm
m="fracas")
```

```
output -2/3*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2/((a^2*f*cos(f*
x + e) + a^2*f)*sin(f*x + e))
```

3.96.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx = \frac{\int \frac{\sqrt{-c\sec(e+fx)+c\sec(e+fx)}}{\sec^2(e+fx)+2\sec(e+fx)+1} dx}{a^2}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(1/2)/(a+a*sec(f*x+e))**2,x)
```

```
output Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e
+ f*x) + 1), x)/a**2
```

3.96.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 109 vs. 2(37) = 74.

Time = 0.31 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.66

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx = -\frac{\sqrt{2}\sqrt{c} - \frac{2\sqrt{2}\sqrt{c}\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{\sqrt{2}\sqrt{c}\sin(fx+e)^4}{(\cos(fx+e)+1)^4}}{6a^2f\sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1}} + 1\sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output `-1/6*(sqrt(2)*sqrt(c) - 2*sqrt(2)*sqrt(c)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + sqrt(2)*sqrt(c)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4)/(a^2*f*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) + 1)*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) - 1))`

3.96.8 Giac [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx = \frac{\sqrt{2}\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{3}{2}}\operatorname{sgn}\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)\operatorname{sgn}(\cos(fx+e))}{6a^2cf}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output `1/6*sqrt(2)*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))*sgn(cos(f*x + e))/(a^2*c*f)`

3.96.9 Mupad [B] (verification not implemented)

Time = 17.18 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.29

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^2} dx = \frac{(e^{e^{2i+fx}2i} + 1)^2 \sqrt{c - \frac{e^{-e^{1i-fx}1i} + e^{e^{1i+fx}1i}}{2}}}{3a^2 f (e^{e^{1i+fx}1i} - 1) (e^{e^{1i+fx}1i} + 1)^3} 2i$$

input `int((c - c/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`output `((exp(e*2i + f*x*2i)*1i + 1i)^2*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*2i)/(3*a^2*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^3)`

$$3.97 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} dx$$

3.97.1	Optimal result	726
3.97.2	Mathematica [C] (verified)	726
3.97.3	Rubi [A] (verified)	727
3.97.4	Maple [A] (verified)	729
3.97.5	Fricas [A] (verification not implemented)	729
3.97.6	Sympy [F]	730
3.97.7	Maxima [F]	730
3.97.8	Giac [A] (verification not implemented)	731
3.97.9	Mupad [F(-1)]	731

3.97.1 Optimal result

Integrand size = 34, antiderivative size = 138

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} dx \\ &= -\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2} \sqrt{c-c \sec(e+fx)}}\right)}{2\sqrt{2}a^2\sqrt{cf}} + \frac{\tan(e+fx)}{3f(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} \\ & \quad + \frac{\tan(e+fx)}{2f(a^2+a^2 \sec(e+fx)) \sqrt{c-c \sec(e+fx)}} \end{aligned}$$

output
$$-1/4*\arctan(1/2*c^{(1/2)}*\tan(f*x+e)*2^{(1/2)}/(c-c*\sec(f*x+e))^{(1/2)})/a^2/f*2^{(1/2)}/c^{(1/2)}+1/3*\tan(f*x+e)/f/(a+a*\sec(f*x+e))^2/(c-c*\sec(f*x+e))^{(1/2)}+1/2*\tan(f*x+e)/f/(a^2+a^2*\sec(f*x+e))/(c-c*\sec(f*x+e))^{(1/2)}$$

3.97.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.32 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.43

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} dx \\ &= \frac{\text{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan(e+fx)}{3f(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} \end{aligned}$$

3.97.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} dx$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]]),x
]`

output `(Hypergeometric2F1[-3/2, 1, -1/2, (1 + Sec[e + f*x])/2]*Tan[e + f*x])/(3*f
*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]])`

3.97.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4448, 3042, 4448, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^2 \sqrt{c - c \sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^2 \sqrt{c - c \csc(e+fx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4448} \\
 & \frac{\int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)\sqrt{c-c\sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{3f(a \sec(e+fx) + a)^2 \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx + \frac{\pi}{2})}{(\csc(e+fx + \frac{\pi}{2})a+a)\sqrt{c-c\csc(e+fx + \frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{3f(a \sec(e+fx) + a)^2 \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{4448} \\
 & \frac{\int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)\sqrt{c - c \sec(e+fx)}} + \frac{\tan(e+fx)}{3f(a \sec(e+fx) + a)^2 \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.97. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} dx$

$$\begin{aligned}
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{4282} \\
& \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}} - \frac{\int \frac{1}{\frac{c^2 \tan^2(e+fx)}{c-c\sec(e+fx)}+2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{af} + \\
& \quad \frac{2a \tan(e+fx)}{3f(a\sec(e+fx)+a)^2\sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow \text{216} \\
& \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}} - \frac{\arctan\left(\frac{\sqrt{c}\tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{2a}\sqrt{cf}} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2\sqrt{c-c\sec(e+fx)}}
\end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]]),x]`

output `Tan[e + f*x]/(3*f*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]]) + (-ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*a*Sqrt[c]*f)) + Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]])/(2*a)`

3.97.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

3.97. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2\sqrt{c-c\sec(e+fx)}} dx$

```
rule 4448 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] :> Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.97.4 Maple [A] (verified)

Time = 3.50 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.33

method	result
default	$\frac{\sqrt{2} \left(-((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1)^{\frac{3}{2}} + 3 \arctan\left(\frac{1}{\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}}\right) + 3\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \right) (-\cos(fx+e))}{12a^2 f \sqrt{\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNV ERBOSE)
```

```
output 1/12/a^2/f*2^(1/2)/(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*csc(f*x+e)^2)^(1/2)*(-((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(3/2)+3*arctan(1/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2))+3*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)))/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(-cot(f*x+e)+csc(f*x+e))
```

3.97.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 331, normalized size of antiderivative = 2.40

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \left[-\frac{3\sqrt{2}\sqrt{-c}(\cos(fx + e) + 1) \log\left(\frac{2\sqrt{2}(\cos(fx+e)^2 + \cos(fx+e))\sqrt{-c}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)} + (3c\cos(fx+e)+c)\sin(fx+e)}}{(\cos(fx+e)-1)\sin(fx+e)}}\right) \sin(fx+e)}{24(a^2cf \cos(fx + e) + a^2cf) \sin(fx + e)} \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x, algorithm m="fricas")
```

3.97. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}} dx$

output `[-1/24*(3*sqrt(2)*sqrt(-c)*(cos(f*x + e) + 1)*log((2*sqrt(2)*(cos(f*x + e) ^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e))*sin(f*x + e) + 4*(5*cos(f*x + e)^2 + 3*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c*f*cos(f*x + e) + a^2*c*f)*sin(f*x + e)), 1/12*(3*sqrt(2)*sqrt(c)*(cos(f*x + e) + 1)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(5*cos(f*x + e)^2 + 3*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c*f*cos(f*x + e) + a^2*c*f)*sin(f*x + e))]`

3.97.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{\int \frac{\sec(e + fx)}{\sqrt{-c \sec(e + fx) + c \sec^2(e + fx) + 2\sqrt{-c \sec(e + fx) + c \sec(e + fx) + \sqrt{-c \sec(e + fx) + c}}}}{a^2} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 + 2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + sqrt(-c*sec(e + f*x) + c)), x)/a**2`

3.97.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)}} dx = \int \frac{\sec(fx + e)}{(a \sec(fx + e) + a)^2 \sqrt{-c \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)^2*sqrt(-c*sec(f*x + e) + c)), x)`

3.97.8 Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{\sqrt{2} \left(\frac{3 \arctan\left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}}\right)}{\sqrt{c}} + \frac{(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)^{\frac{3}{2}} c^4 - 3 \sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c} c^5}{c^6} \right)}{12 a^2 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `1/12*sqrt(2)*(3*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/sqrt(c) + ((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c^4 - 3*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^5)/c^6)/(a^2*f)`

3.97.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx) \left(a + \frac{a}{\cos(e + fx)}\right)^2 \sqrt{c - \frac{c}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(1/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(1/2)), x)`

$$3.98 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{3/2}} dx$$

3.98.1	Optimal result	732
3.98.2	Mathematica [C] (verified)	732
3.98.3	Rubi [A] (verified)	733
3.98.4	Maple [A] (verified)	736
3.98.5	Fricas [A] (verification not implemented)	736
3.98.6	Sympy [F]	737
3.98.7	Maxima [F]	737
3.98.8	Giac [A] (verification not implemented)	738
3.98.9	Mupad [F(-1)]	738

3.98.1 Optimal result

Integrand size = 34, antiderivative size = 169

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{3/2}} dx = -\frac{5 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{8\sqrt{2}a^2c^{3/2}f} - \frac{5 \tan(e+fx)}{8a^2f(c-c \sec(e+fx))^{3/2}} + \frac{\tan(e+fx)}{3f(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{3/2}} + \frac{5 \tan(e+fx)}{6f(a^2+a^2 \sec(e+fx))(c-c \sec(e+fx))^{3/2}}$$

```
output -5/16*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a^2/c^(3/2)/f*2^(1/2)-5/8*tan(f*x+e)/a^2/f/(c-c*sec(f*x+e))^(3/2)+1/3*tan(f*x+e)/f/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2)+5/6*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2)
```

3.98.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.49 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.38

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{3/2}} dx = \frac{\text{Hypergeometric2F1}\left(-\frac{3}{2}, 2, -\frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan(e+fx)}{6a^2cf(1+\sec(e+fx))^2\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2)),x]`

output `(Hypergeometric2F1[-3/2, 2, -1/2, (1 + Sec[e + f*x])/2]*Tan[e + f*x])/(6*a^2*c*f*(1 + Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]])`

3.98.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4448, 3042, 4448, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e + fx)}{(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e + fx + \frac{\pi}{2})}{(a \csc(e + fx + \frac{\pi}{2}) + a)^2 (c - c \csc(e + fx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4448} \\
 & \frac{5 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)(c-c\sec(e+fx))^{3/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{3/2}} \\
 & \quad \downarrow \text{4448} \\
 & \frac{5 \left(\frac{3 \int \frac{\sec(e+fx)}{(c-c\sec(e+fx))^{3/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c\sec(e+fx))^{3/2}} \right)}{6a} + \\
 & \quad \frac{\tan(e+fx)}{3f(a \sec(e + fx) + a)^2 (c - c \sec(e + fx))^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.98. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))^{3/2}} dx$

$$\begin{aligned}
 & 5 \left(\frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c \csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right) + \\
 & \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 4283 \\
 & 5 \left(\frac{3 \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c-c \sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right) + \\
 & \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & 5 \left(\frac{3 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right) + \\
 & \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 4282 \\
 & 5 \left(\frac{3 \left(\frac{\int \frac{1}{c^2 \tan^2(e+fx)+2c} d \frac{c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}}}{2cf} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right) + \\
 & \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}} \\
 & \quad \downarrow 216
 \end{aligned}$$

3.98. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{3/2}} dx$

$$5 \left(\frac{3 \left(-\frac{\arctan\left(\frac{\sqrt{c}\tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{2\sqrt{2}c^{3/2}f} - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{3/2}} \right) + \frac{6a \tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{3/2}}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2)),x]`

output `Tan[e + f*x]/(3*f*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2)) + (5*(Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2)) + (3*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2))))/(2*a)))/(6*a)`

3.98.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4283 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`


```
rule 4448 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_), x_Symbol] :> Simp[b*Cot[e + f*x]*
(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[
(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(
c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (IL
tQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.98.4 Maple [A] (verified)

Time = 3.61 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.18

method	result
default	$\frac{\sqrt{2} \left(-13\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e)^2 + 15 \sin(fx+e)^2 \arctan \left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} \right) + 10\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e) + 15\sqrt{2} \right)}{48a^2 f (\sec(fx+e)-1) \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \sqrt{-c(\sec(fx+e)-1)} c(\cos(fx+e)+1)^2}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/48/a^2/f*2^(1/2)*(-13*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x
+e)^2+15*sin(f*x+e)^2*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2
))+10*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)+15*2^(1/2)*(-c
os(f*x+e)/(cos(f*x+e)+1))^(1/2))/(sec(f*x+e)-1)/(-cos(f*x+e)/(cos(f*x+e)+1
))^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)/c/(cos(f*x+e)+1)^2*tan(f*x+e)
```

3.98.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 369, normalized size of antiderivative = 2.18

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{3/2}} dx = \left[\frac{15 \sqrt{2} (\cos(fx + e)^2 - 1) \sqrt{-c} \log \left(\frac{2 \sqrt{2} (\cos(fx + e)^2 + \cos(fx + e))}{\dots} \right)}{\dots} \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x, algorith
m="fricas")
```

3.98. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))^{3/2}} dx$

output `[-1/96*(15*sqrt(2)*(cos(f*x + e)^2 - 1)*sqrt(-c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) + 4*(13*cos(f*x + e)^3 - 10*cos(f*x + e)^2 - 15*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c^2*f*cos(f*x + e)^2 - a^2*c^2*f)*sin(f*x + e)), 1/48*(15*sqrt(2)*(cos(f*x + e)^2 - 1)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(13*cos(f*x + e)^3 - 10*cos(f*x + e)^2 - 15*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c^2*f*cos(f*x + e)^2 - a^2*c^2*f)*sin(f*x + e))]`

3.98.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{3/2}} dx = \frac{\int \frac{\sec(e + fx)}{-c\sqrt{-c \sec(e + fx) + c \sec^3(e + fx) - c\sqrt{-c \sec(e + fx) + c \sec^2(e + fx) + c\sqrt{-c \sec(e + fx) + c \sec^3(e + fx)}}} dx}{a^2}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**(3/2),x)`

output `Integral(sec(e + f*x)/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3 - c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 + c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x)/a**2`

3.98.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{3/2}} dx = \int \frac{\sec(fx + e)}{(a \sec(fx + e) + a)^2 (-c \sec(fx + e) + c)^{3/2}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)^2*(-c*sec(f*x + e) + c)^(3/2)), x)`

3.98. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))^{3/2}} dx$

3.98.8 Giac [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.76

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^{3/2}} dx = \frac{\sqrt{2} \left(15\sqrt{c} \arctan\left(\frac{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}}{\sqrt{c}}\right) - \frac{3\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)}}{\tan(\frac{1}{2}fx + \frac{1}{2}e)} \right)}{48a^2c}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2),x, algorithm m="giac")`

output `1/48*sqrt(2)*(15*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) - 3*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/tan(1/2*f*x + 1/2*e)^2 + 2*((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c^2 - 6*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^3)/c^3)/(a^2*c^2*f)`

3.98.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^{3/2}} dx = \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^2 \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(3/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(3/2)), x)`

3.99
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{5/2}} dx$$

3.99.1	Optimal result	739
3.99.2	Mathematica [C] (verified)	739
3.99.3	Rubi [A] (verified)	740
3.99.4	Maple [A] (verified)	744
3.99.5	Fricas [A] (verification not implemented)	745
3.99.6	Sympy [F]	745
3.99.7	Maxima [F]	746
3.99.8	Giac [A] (verification not implemented)	746
3.99.9	Mupad [F(-1)]	747

3.99.1 Optimal result

Integrand size = 34, antiderivative size = 203

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{5/2}} dx = -\frac{35 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{64\sqrt{2}a^2c^{5/2}f} - \frac{35 \tan(e+fx)}{48a^2f(c-c \sec(e+fx))^{5/2}} + \frac{\tan(e+fx)}{3f(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{5/2}} + \frac{7 \tan(e+fx)}{6f(a^2+a^2 \sec(e+fx))(c-c \sec(e+fx))^{5/2}} - \frac{35 \tan(e+fx)}{64a^2cf(c-c \sec(e+fx))^{3/2}}$$

```
output -35/128*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a^2/
c^(5/2)/f*2^(1/2)-35/48*tan(f*x+e)/a^2/f/(c-c*sec(f*x+e))^(5/2)+1/3*tan(f*
x+e)/f/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2)+7/6*tan(f*x+e)/f/(a^2+a^2
*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2)-35/64*tan(f*x+e)/a^2/c/f/(c-c*sec(f*x+
e))^(3/2)
```

3.99.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.80 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.32

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{5/2}} dx = \frac{\text{Hypergeometric2F1}\left(-\frac{3}{2}, 3, -\frac{1}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan(e+fx)}{12a^2c^2f(1+\sec(e+fx))^2\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(5/2)),x]`

output `(Hypergeometric2F1[-3/2, 3, -1/2, (1 + Sec[e + f*x])/2]*Tan[e + f*x])/(12*a^2*c^2*f*(1 + Sec[e + f*x])^2*sqrt[c - c*Sec[e + f*x]])`

3.99.3 Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4448, 3042, 4448, 3042, 4283, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^2 (c - c \sec(e+fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^2 (c - c \csc(e+fx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4448} \\
 & \frac{7 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)(c-c \sec(e+fx))^{5/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a \sec(e+fx) + a)^2 (c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{7 \int \frac{\csc(e+fx + \frac{\pi}{2})}{(\csc(e+fx + \frac{\pi}{2})a+a)(c-c \csc(e+fx + \frac{\pi}{2}))^{5/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a \sec(e+fx) + a)^2 (c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4448} \\
 & \frac{7 \left(\frac{5 \int \frac{\sec(e+fx)}{(c-c \sec(e+fx))^{5/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c - c \sec(e+fx))^{5/2}} \right)}{6a} + \\
 & \quad \frac{\tan(e+fx)}{3f(a \sec(e+fx) + a)^2 (c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.99. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c-c \sec(e+fx))^{5/2}} dx$

$$\begin{aligned}
 & 7 \left(\frac{5 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c \csc(e+fx+\frac{\pi}{2}))^{5/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow 4283 \\
 & 7 \left(\frac{5 \left(\frac{3 \int \frac{\sec(e+fx)}{(c-c \sec(e+fx))^{3/2}} dx}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow 3042 \\
 & 7 \left(\frac{5 \left(\frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c \csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow 4283 \\
 & 7 \left(\frac{5 \left(\frac{3 \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c-c \sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.99. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{5/2}} dx$

$$\left(\frac{5}{7} \left(\frac{3 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right) + \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}}$$

↓ 4282

$$\left(\frac{5}{7} \left(\frac{3 \left(\frac{\int \frac{\frac{1}{c^2 \tan^2(e+fx)} + 2c}{c-c \sec(e+fx)} dx - \frac{c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}}}{2cf} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right) + \frac{6a \tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}}$$

↓ 216

$$\frac{\left(\frac{5 \left(\frac{3 \left(-\frac{\arctan\left(\frac{\sqrt{c}\tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right)}{2\sqrt{2}c^{3/2}f} - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}} \right)}{8c} - \frac{\tan(e+fx)}{4f(c-c\sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}} \right)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{5/2}} + \frac{6a \tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{5/2}}$$

```
input Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(5/2)),x]
```

```
output Tan[e + f*x]/(3*f*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(5/2)) + (7*(Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2)) + (5*(-1/4*Tan[e + f*x]/(f*(c - c*Sec[e + f*x])^(5/2)) + (3*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])])/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2))))/(8*c)))/(2*a)))/(6*a)
```

3.99.3.1 Defintions of rubi rules used

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4282 Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]
```



```
rule 4283 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_
Symbol] :> Simp[b*Cot[e + f*x]**((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x]
+ Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m +
1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)
] && IntegerQ[2*m]
```

```
rule 4448 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_), x_Symbol] :> Simp[b*Cot[e + f*x]*
(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[
(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(
c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (IL
tQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.99.4 Maple [A] (verified)

Time = 3.35 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.38

method	result
default	$\frac{\sqrt{2} \left(-43 \cos(fx+e)^3 \sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} + 105 \sin(fx+e)^2 \arctan \left(\frac{\sqrt{2}}{2 \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}} \right) \cos(fx+e) + 161 \sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e) \right)}{384 a^2 f \sqrt{-c(\sec(fx+e)-1)} (\sec(fx+e)-1)}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/384/a^2/f*2^(1/2)*(-43*cos(f*x+e)^3*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1)
)^(1/2)+105*sin(f*x+e)^2*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(
1/2))*cos(f*x+e)+161*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)
^2-105*sin(f*x+e)^2*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))
+35*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)-105*2^(1/2)*(-co
s(f*x+e)/(cos(f*x+e)+1))^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)/(sec(f*x+e)-1)^2
/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)/c^2/(cos(f*x+e)+1)^2*tan(f*x+e)*sec(f*
x+e)
```

3.99.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 487, normalized size of antiderivative = 2.40

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^{5/2}} dx = \left[\frac{105\sqrt{2}(\cos(fx+e))^3 - \cos(fx+e)^2 - \cos(fx+e)}{\dots} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x, algorithm="fricas")`

output `[-1/768*(105*sqrt(2)*(cos(f*x + e)^3 - cos(f*x + e)^2 - cos(f*x + e) + 1)*sqrt(-c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) + 4*(43*cos(f*x + e)^4 - 161*cos(f*x + e)^3 - 35*cos(f*x + e)^2 + 105*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c^3*f*cos(f*x + e)^3 - a^2*c^3*f*cos(f*x + e)^2 - a^2*c^3*f*cos(f*x + e) + a^2*c^3*f)*sin(f*x + e)), 1/384*(105*sqrt(2)*(cos(f*x + e)^3 - cos(f*x + e)^2 - cos(f*x + e) + 1)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(43*cos(f*x + e)^4 - 161*cos(f*x + e)^3 - 35*cos(f*x + e)^2 + 105*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c^3*f*cos(f*x + e)^3 - a^2*c^3*f*cos(f*x + e)^2 - a^2*c^3*f*cos(f*x + e) + a^2*c^3*f)*sin(f*x + e))]`

3.99.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c-c\sec(e+fx))^{5/2}} dx = \frac{\int \frac{\sec(e+fx)}{c^2\sqrt{-c\sec(e+fx)+c\sec^4(e+fx)-2c^2\sqrt{-c\sec(e+fx)+c\sec^2(e+fx)+c^2}}}{a^2}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c-c*sec(f*x+e))**(5/2),x)`

output `Integral(sec(e + f*x)/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 + c**2*sqrt(-c*sec(e + f*x) + c)), x)/a**2`

3.99.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{5/2}} dx = \int \frac{\sec(fx + e)}{(a \sec(fx + e) + a)^2 (-c \sec(fx + e) + c)^{5/2}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x, algorithm
m="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)^2*(-c*sec(f*x + e) + c)^(5/2)
, x)`

3.99.8 Giac [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.79

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{5/2}} dx = \frac{\sqrt{2} \left(105 \sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}} \right) + 8 \left(c \tan(\frac{1}{2} fx + \frac{1}{2} e) \right)^{3/2} \right)}{(a \sec(fx + e) + a)^2 (-c \sec(fx + e) + c)^{5/2}}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2),x, algorithm
m="giac")`

output `1/384*sqrt(2)*(105*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) + 8*((c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c^2 - 9*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^3)/c^3 - 3*(13*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c + 11*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^2)/(c^2*tan(1/2*f*x + 1/2*e)^4)/(a^2*c^3*f)`

3.99.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c - c \sec(e + fx))^{5/2}} dx = \int \frac{1}{\cos(e + fx) \left(a + \frac{a}{\cos(e + fx)}\right)^2 \left(c - \frac{c}{\cos(e + fx)}\right)^{5/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(5/2)),x)`output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))^(5/2)), x)`

3.100
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx$$

3.100.1 Optimal result	748
3.100.2 Mathematica [A] (verified)	748
3.100.3 Rubi [A] (verified)	749
3.100.4 Maple [A] (verified)	751
3.100.5 Fricas [A] (verification not implemented)	752
3.100.6 Sympy [F(-1)]	752
3.100.7 Maxima [A] (verification not implemented)	752
3.100.8 Giac [A] (verification not implemented)	753
3.100.9 Mupad [B] (verification not implemented)	754

3.100.1 Optimal result

Integrand size = 34, antiderivative size = 169

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx = \frac{32c^4 \tan(e+fx)}{5a^3 f \sqrt{c-c\sec(e+fx)}} + \frac{16c^3 \sqrt{c-c\sec(e+fx)} \tan(e+fx)}{5f(a^3+a^3\sec(e+fx))} - \frac{4c^2(c-c\sec(e+fx))^{3/2} \tan(e+fx)}{5af(a+a\sec(e+fx))^2} + \frac{2c(c-c\sec(e+fx))^{5/2} \tan(e+fx)}{5f(a+a\sec(e+fx))^3}$$

output `-4/5*c^2*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2+2/5*c*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+32/5*c^4*tan(f*x+e)/a^3/f/(c-c*sec(f*x+e))^(1/2)+16/5*c^3*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))`

3.100.2 Mathematica [A] (verified)

Time = 0.91 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.44

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx = \frac{2c^4(23+55\sec(e+fx)+45\sec^2(e+fx)+5\sec^3(e+fx))\tan(e+fx)}{5a^3f(1+\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e+f*x]*(c-c*Sec[e+f*x])^(7/2))/(a+a*Sec[e+f*x])^3,x]`

3.100.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx$$

output $(2*c^4*(23 + 55*Sec[e + f*x] + 45*Sec[e + f*x]^2 + 5*Sec[e + f*x]^3)*Tan[e + f*x])/(5*a^3*f*(1 + Sec[e + f*x])^3*sqrt[c - c*Sec[e + f*x]])$

3.100.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4442, 3042, 4442, 3042, 4442, 3042, 4279}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a\sec(e+fx)+a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{7/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\
 & \quad \downarrow \text{4442} \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^{5/2}}{5f(a\sec(e+fx)+a)^3} - \frac{6c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(\sec(e+fx)a+a)^2} dx}{5a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^{5/2}}{5f(a\sec(e+fx)+a)^3} - \frac{6c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a} \\
 & \quad \downarrow \text{4442} \\
 & \frac{2c \tan(e+fx)(c-c\sec(e+fx))^{5/2}}{5f(a\sec(e+fx)+a)^3} - \\
 & \frac{6c \left(\frac{2c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{3f(a\sec(e+fx)+a)^2} - \frac{4c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sec(e+fx)a+a} dx \right)}{5a} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^{5/2}}{5f(a \sec(e+fx) + a)^3} - \\
& \frac{6c \left(\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{3f(a \sec(e+fx) + a)^2} - \frac{4c \int \frac{\csc(e+fx + \frac{\pi}{2})(c - c \csc(e+fx + \frac{\pi}{2}))^{3/2}}{\csc(e+fx + \frac{\pi}{2})a + a} dx}{3a} \right)}{5a} \\
& \quad \downarrow 4442 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^{5/2}}{5f(a \sec(e+fx) + a)^3} - \\
& \frac{6c \left(\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{3f(a \sec(e+fx) + a)^2} - \frac{4c \left(\frac{2c \tan(e+fx)\sqrt{c - c \sec(e+fx)}}{f(a \sec(e+fx) + a)} - \frac{2c \int \sec(e+fx)\sqrt{c - c \sec(e+fx)} dx}{a} \right)}{3a} \right)}{5a} \\
& \quad \downarrow 3042 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^{5/2}}{5f(a \sec(e+fx) + a)^3} - \\
& \frac{6c \left(\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{3f(a \sec(e+fx) + a)^2} - \frac{4c \left(\frac{2c \tan(e+fx)\sqrt{c - c \sec(e+fx)}}{f(a \sec(e+fx) + a)} - \frac{2c \int \csc(e+fx + \frac{\pi}{2})\sqrt{c - c \csc(e+fx + \frac{\pi}{2})} dx}{a} \right)}{3a} \right)}{5a} \\
& \quad \downarrow 4279 \\
& \frac{2c \tan(e+fx)(c - c \sec(e+fx))^{5/2}}{5f(a \sec(e+fx) + a)^3} - \\
& \frac{6c \left(\frac{2c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{3f(a \sec(e+fx) + a)^2} - \frac{4c \left(\frac{4c^2 \tan(e+fx)}{af\sqrt{c - c \sec(e+fx)}} + \frac{2c \tan(e+fx)\sqrt{c - c \sec(e+fx)}}{f(a \sec(e+fx) + a)} \right)}{3a} \right)}{5a}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(7/2))/(a + a*Sec[e + f*x])^3,x]`

output `(2*c*(c - c*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) - (6*c*((2*c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (4*c*((4*c^2*Tan[e + f*x])/(a*f*Sqrt[c - c*Sec[e + f*x]]) + (2*c*Sqrt[c - c*Sec[e + f*x])*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))))/(3*a)))/(5*a)`

3.100.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4279 `Int[csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[-2*b*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.100.4 Maple [A] (verified)

Time = 36.52 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.54

method	result	size
default	$\frac{2\sqrt{-c(\sec(fx+e)-1)}(\sec(fx+e)-1)^3c^3(23\cos(fx+e)^3+55\cos(fx+e)^2+45\cos(fx+e)+5)\cot(fx+e)^3\csc(fx+e)^2}{5a^3f(\cos(fx+e)-1)}$	91

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^3,x,method=_RETURNV ERBOSE)`

output
$$\frac{2}{5} \frac{1}{a^3 f} \frac{(-c(\sec(fx+e)-1))^{1/2} (\sec(fx+e)-1)^3 c^3 (23 \cos(fx+e)^3 + 55 \cos(fx+e)^2 + 45 \cos(fx+e) + 5)}{(\cos(fx+e)-1) \cot(fx+e)^3 \csc(fx+e)^2}$$

3.100.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.64

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx = \frac{2(23c^3\cos(fx+e)^3 + 55c^3\cos(fx+e)^2 + 45c^3\cos(fx+e) + 5c^3)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{5(a^3f\cos(fx+e)^2 + 2a^3f\cos(fx+e) + a^3f)\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^3,x, algorithm m="fricas")`

output `-2/5*(23*c^3*cos(f*x + e)^3 + 55*c^3*cos(f*x + e)^2 + 45*c^3*cos(f*x + e) + 5*c^3)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((a^3*f*cos(f*x + e)^2 + 2*a^3*f*cos(f*x + e) + a^3*f)*sin(f*x + e))`

3.100.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(7/2)/(a+a*sec(f*x+e))**3,x)`

output `Timed out`

3.100.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.27

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx = \frac{2\left(16\sqrt{2}c^{\frac{7}{2}} - \frac{56\sqrt{2}c^{\frac{7}{2}}\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{70\sqrt{2}c^{\frac{7}{2}}\sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{35\sqrt{2}c^{\frac{7}{2}}\sin(fx+e)^6}{(\cos(fx+e)+1)^6}\right)}{5a^3f\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)^{\frac{7}{2}}\left(\frac{c}{\cos(fx+e)+1} + 1\right)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^3,x, algorithm m="maxima")`

output
$$\frac{2}{5} * (16 * \sqrt{2} * c^{7/2} - 56 * \sqrt{2} * c^{7/2} * \sin(f*x + e)^2 / (\cos(f*x + e) + 1)^2 + 70 * \sqrt{2} * c^{7/2} * \sin(f*x + e)^4 / (\cos(f*x + e) + 1)^4 - 35 * \sqrt{2} * c^{7/2} * \sin(f*x + e)^6 / (\cos(f*x + e) + 1)^6 + 5 * \sqrt{2} * c^{7/2} * \sin(f*x + e)^8 / (\cos(f*x + e) + 1)^8 - \sqrt{2} * c^{7/2} * \sin(f*x + e)^{10} / (\cos(f*x + e) + 1)^{10} + \sqrt{2} * c^{7/2} * \sin(f*x + e)^{12} / (\cos(f*x + e) + 1)^{12}) / (a^3 * f * (\sin(f*x + e) / (\cos(f*x + e) + 1) + 1)^{7/2} * (\sin(f*x + e) / (\cos(f*x + e) + 1) - 1)^{7/2})$$

3.100.8 Giac [A] (verification not implemented)

Time = 1.02 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.75

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{7/2}}{(a + a \sec(e + fx))^3} dx = \frac{2\sqrt{2}c^3 \left(\frac{5c}{\sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - ca^3}} - \frac{(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^{5/2} a^{12} c^8 + 5(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^{3/2} a^{12} c^9 + 15 \sqrt{c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c} a^{12} c^{10}}{5f} \right)}{5f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^3,x, algorithm m="giac")`

output
$$\frac{2}{5} * \sqrt{2} * c^3 * (5 * c / (\sqrt{c * \tan(1/2 * f * x + 1/2 * e)^2 - c} * a^3) - ((c * \tan(1/2 * f * x + 1/2 * e)^2 - c)^{5/2} * a^{12} * c^8 + 5 * (c * \tan(1/2 * f * x + 1/2 * e)^2 - c)^{3/2} * a^{12} * c^9 + 15 * \sqrt{c * \tan(1/2 * f * x + 1/2 * e)^2 - c} * a^{12} * c^{10}) / (a^{15} * c^{10})) / f$$

3.100.9 Mupad [B] (verification not implemented)

Time = 22.30 (sec) , antiderivative size = 492, normalized size of antiderivative = 2.91

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{7/2}}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{\sqrt{c - \frac{c}{\frac{e^{-e^{1i}-fx^{1i}}}{2} + e^{e^{1i}+fx^{1i}}}} \left(\frac{c^3 46i}{5a^3 f} + \frac{c^3 e^{e^{1i}+fx^{1i}} 4i}{a^3 f} + \frac{c^3 e^{e^{2i}+fx^{2i}} 46i}{5a^3 f} \right)}{(e^{e^{1i}+fx^{1i}} - 1)(e^{e^{1i}+fx^{1i}} + 1)}$$

$$- \frac{c^3 (e^{e^{2i}+fx^{2i}} + 1) \sqrt{c - \frac{c}{\frac{e^{-e^{1i}-fx^{1i}}}{2} + e^{e^{1i}+fx^{1i}}}} 16i}{5a^3 f (e^{e^{1i}+fx^{1i}} - 1)(e^{e^{1i}+fx^{1i}} + 1)^2}$$

$$- \frac{c^3 (e^{e^{2i}+fx^{2i}} + 1) \sqrt{c - \frac{c}{\frac{e^{-e^{1i}-fx^{1i}}}{2} + e^{e^{1i}+fx^{1i}}}} 48i}{5a^3 f (e^{e^{1i}+fx^{1i}} - 1)(e^{e^{1i}+fx^{1i}} + 1)^3}$$

$$+ \frac{c^3 (e^{e^{2i}+fx^{2i}} + 1) \sqrt{c - \frac{c}{\frac{e^{-e^{1i}-fx^{1i}}}{2} + e^{e^{1i}+fx^{1i}}}} 128i}{5a^3 f (e^{e^{1i}+fx^{1i}} - 1)(e^{e^{1i}+fx^{1i}} + 1)^4}$$

$$- \frac{c^3 (e^{e^{2i}+fx^{2i}} + 1) \sqrt{c - \frac{c}{\frac{e^{-e^{1i}-fx^{1i}}}{2} + e^{e^{1i}+fx^{1i}}}} 64i}{5a^3 f (e^{e^{1i}+fx^{1i}} - 1)(e^{e^{1i}+fx^{1i}} + 1)^5}$$

```
input int((c - c/cos(e + f*x))^(7/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)
```

```
output (c^3*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*128i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^4) - (c^3*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*16i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^2) - (c^3*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*48i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^3) - ((c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*((c^3*46i)/(5*a^3*f) + (c^3*exp(e*1i + f*x*1i)*4i)/(a^3*f) + (c^3*exp(e*2i + f*x*2i)*46i)/(5*a^3*f)))/((exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)) - (c^3*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*64i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^5)
```

3.101
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx$$

3.101.1 Optimal result	755
3.101.2 Mathematica [A] (verified)	755
3.101.3 Rubi [A] (verified)	756
3.101.4 Maple [A] (verified)	758
3.101.5 Fricas [A] (verification not implemented)	758
3.101.6 Sympy [F(-1)]	759
3.101.7 Maxima [A] (verification not implemented)	759
3.101.8 Giac [A] (verification not implemented)	760
3.101.9 Mupad [B] (verification not implemented)	760

3.101.1 Optimal result

Integrand size = 34, antiderivative size = 135

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx = \frac{16c^3 \tan(e+fx)}{15f(a^3+a^3\sec(e+fx))\sqrt{c-c\sec(e+fx)}} - \frac{8c^2\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{15af(a+a\sec(e+fx))^2} + \frac{2c(c-c\sec(e+fx))^{3/2}\tan(e+fx)}{5f(a+a\sec(e+fx))^3}$$

output `2/5*c*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+16/15*c^3*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2)-8/15*c^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2`

3.101.2 Mathematica [A] (verified)

Time = 1.46 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.47

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx = \frac{2c^3(7+10\sec(e+fx)+15\sec^2(e+fx))\tan(e+fx)}{15a^3f(1+\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^3, x]`

output `(2*c^3*(7 + 10*Sec[e + f*x] + 15*Sec[e + f*x]^2)*Tan[e + f*x])/(15*a^3*f*(1 + Sec[e + f*x])^3*sqrt[c - c*Sec[e + f*x]])`

3.101.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx$$

3.101.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4442, 3042, 4442, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a\sec(e+fx)+a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\
 & \quad \downarrow \text{4442} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{5f(a\sec(e+fx)+a)^3} - \frac{4c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(\sec(e+fx)a+a)^2} dx}{5a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{5f(a\sec(e+fx)+a)^3} - \frac{4c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a} \\
 & \quad \downarrow \text{4442} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{5f(a\sec(e+fx)+a)^3} - \frac{4c \left(\frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{3f(a\sec(e+fx)+a)^2} - \frac{2c \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sec(e+fx)a+a} dx}{3a} \right)}{5a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{5f(a\sec(e+fx)+a)^3} - \frac{4c \left(\frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{3f(a\sec(e+fx)+a)^2} - \frac{2c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} \right)}{5a} \\
 & \quad \downarrow \text{4441}
 \end{aligned}$$

$$\frac{2c \tan(e+fx)(c - c \operatorname{sec}(e+fx))^{3/2}}{5f(a \operatorname{sec}(e+fx) + a)^3} - \frac{4c \left(\frac{2c \tan(e+fx) \sqrt{c - c \operatorname{sec}(e+fx)}}{3f(a \operatorname{sec}(e+fx) + a)^2} - \frac{4c^2 \tan(e+fx)}{3af(a \operatorname{sec}(e+fx) + a) \sqrt{c - c \operatorname{sec}(e+fx)}} \right)}{5a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^3,x]`

output `(2*c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) - (4*c*((-4*c^2*Tan[e + f*x])/(3*a*f*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]]) + (2*c*Sqrt[c - c*Sec[e + f*x])*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2)))/(5*a)`

3.101.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*Sqrt[csc[(e_) + (f_)*(x_)]*(d_) + (c_)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4442 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.101.4 Maple [A] (verified)

Time = 41.61 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.53

method	result	size
default	$-\frac{2\sqrt{-c(\sec(fx+e)-1)}(\sec(fx+e)-1)^2c^2(7\cos(fx+e)^2+10\cos(fx+e)+15)\cot(fx+e)^3\csc(fx+e)^2}{15a^3f}$	71

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^3,x,method=_RETURNV
ERBOSE)
```

```
output -2/15/a^3/f*(-c*(sec(f*x+e)-1))^(1/2)*(sec(f*x+e)-1)^2*c^2*(7*cos(f*x+e)^2
+10*cos(f*x+e)+15)*cot(f*x+e)^3*csc(f*x+e)^2
```

3.101.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.77

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{2(7c^2\cos(fx+e)^3+10c^2\cos(fx+e)^2+15c^2\cos(fx+e))\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{15(a^3f\cos(fx+e)^2+2a^3f\cos(fx+e)+a^3f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^3,x, algorithm
m="fricas")
```

```
output -2/15*(7*c^2*cos(f*x + e)^3 + 10*c^2*cos(f*x + e)^2 + 15*c^2*cos(f*x + e))
*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((a^3*f*cos(f*x + e)^2 + 2*a^3*f*
cos(f*x + e) + a^3*f)*sin(f*x + e))
```

3.101.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{(a + a \sec(e + fx))^3} dx = \text{Timed out}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(5/2)/(a+a*sec(f*x+e))**3,x)
```

```
output Timed out
```

3.101.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.40

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{(a + a \sec(e + fx))^3} dx =$$

$$\frac{8\sqrt{2}c^{\frac{5}{2}} - \frac{20\sqrt{2}c^{\frac{5}{2}}\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{15\sqrt{2}c^{\frac{5}{2}}\sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{5\sqrt{2}c^{\frac{5}{2}}\sin(fx+e)^6}{(\cos(fx+e)+1)^6} + \frac{5\sqrt{2}c^{\frac{5}{2}}\sin(fx+e)^8}{(\cos(fx+e)+1)^8} - \frac{3\sqrt{2}c^{\frac{5}{2}}\sin(fx+e)^{10}}{(\cos(fx+e)+1)^{10}}}{15a^3f\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)^{\frac{5}{2}}\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)^{\frac{5}{2}}}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^3,x, algorithm="maxima")
```

```
output -1/15*(8*sqrt(2)*c^(5/2) - 20*sqrt(2)*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 15*sqrt(2)*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 5*sqrt(2)*c^(5/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 5*sqrt(2)*c^(5/2)*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 - 3*sqrt(2)*c^(5/2)*sin(f*x + e)^10/(cos(f*x + e) + 1)^10)/(a^3*f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^(5/2)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^(5/2))
```


3.101.8 Giac [A] (verification not implemented)

Time = 0.86 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{15\sqrt{2}\sqrt{c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-cc^2}}{a^3} + \frac{3\sqrt{2}(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)^{5/2}+10\sqrt{2}(c\tan(\frac{1}{2}fx+\frac{1}{2}e)^2-c)^{3/2}c}{a^3}$$

$$\frac{\hspace{15em}}{15f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^3,x, algorithm m="giac")`

output `-1/15*(15*sqrt(2)*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^2/a^3 + (3*sqrt(2)*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2) + 10*sqrt(2)*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c)/a^3)/f`

3.101.9 Mupad [B] (verification not implemented)

Time = 19.27 (sec) , antiderivative size = 456, normalized size of antiderivative = 3.38

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{c^2 (e^{e^{2i+fx}2i} + 1) \sqrt{c - \frac{e^{-e^{li-fx}li}c}{2} + \frac{e^{e^{li+fx}li}}{2}}}{15 a^3 f (e^{e^{li+fx}li} - 1) (e^{e^{li+fx}li} + 1)} 14i$$

$$+ \frac{c^2 (e^{e^{2i+fx}2i} + 1) \sqrt{c - \frac{e^{-e^{li-fx}li}c}{2} + \frac{e^{e^{li+fx}li}}{2}}}{15 a^3 f (e^{e^{li+fx}li} - 1) (e^{e^{li+fx}li} + 1)^2} 16i$$

$$- \frac{c^2 (e^{e^{2i+fx}2i} + 1) \sqrt{c - \frac{e^{-e^{li-fx}li}c}{2} + \frac{e^{e^{li+fx}li}}{2}}}{15 a^3 f (e^{e^{li+fx}li} - 1) (e^{e^{li+fx}li} + 1)^3} 112i$$

$$+ \frac{c^2 (e^{e^{2i+fx}2i} + 1) \sqrt{c - \frac{e^{-e^{li-fx}li}c}{2} + \frac{e^{e^{li+fx}li}}{2}}}{5 a^3 f (e^{e^{li+fx}li} - 1) (e^{e^{li+fx}li} + 1)^4} 64i$$

$$- \frac{c^2 (e^{e^{2i+fx}2i} + 1) \sqrt{c - \frac{e^{-e^{li-fx}li}c}{2} + \frac{e^{e^{li+fx}li}}{2}}}{5 a^3 f (e^{e^{li+fx}li} - 1) (e^{e^{li+fx}li} + 1)^5} 32i$$

input `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`

output

```
(c^2*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*16i)/(15*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^2) - (c^2*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*14i)/(15*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)) - (c^2*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*112i)/(15*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^3) + (c^2*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*64i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^4) - (c^2*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*32i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^5)
```

$$3.102 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx$$

3.102.1 Optimal result	762
3.102.2 Mathematica [A] (verified)	762
3.102.3 Rubi [A] (verified)	763
3.102.4 Maple [A] (verified)	764
3.102.5 Fricas [A] (verification not implemented)	765
3.102.6 Sympy [F]	765
3.102.7 Maxima [B] (verification not implemented)	766
3.102.8 Giac [A] (verification not implemented)	766
3.102.9 Mupad [B] (verification not implemented)	767

3.102.1 Optimal result

Integrand size = 34, antiderivative size = 88

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx = -\frac{4c^2 \tan(e+fx)}{15af(a+a\sec(e+fx))^2 \sqrt{c-c\sec(e+fx)}} + \frac{2c\sqrt{c-c\sec(e+fx)} \tan(e+fx)}{5f(a+a\sec(e+fx))^3}$$

output `-4/15*c^2*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2)+2/5*c*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^3`

3.102.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.61

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx = -\frac{2c^2(-1+5\sec(e+fx)) \tan(e+fx)}{15a^3 f(1+\sec(e+fx))^3 \sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x])^3, x]`

output `(-2*c^2*(-1 + 5*Sec[e + f*x])*Tan[e + f*x])/(15*a^3*f*(1 + Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]])`

3.102. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx$

3.102.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4442, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a\sec(e+fx)+a)^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx$$

↓ 4442

$$\frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3} - \frac{2c\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(\sec(e+fx)a+a)^2} dx}{5a}$$

↓ 3042

$$\frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3} - \frac{2c\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a}$$

↓ 4441

$$\frac{2c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3} - \frac{4c^2\tan(e+fx)}{15af(a\sec(e+fx)+a)^2\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x])^3,x]`

output `(-4*c^2*Tan[e + f*x])/(15*a*f*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]]) + (2*c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3)`

3.102.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.102.4 Maple [A] (verified)

Time = 3.44 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{2(\sec(fx+e)-1)(\cos(fx+e)-5)\sqrt{-c(\sec(fx+e)-1)}c\cos(fx+e)^2\cot(fx+e)}{15a^3f(\cos(fx+e)+1)^2(\cos(fx+e)-1)}$	73

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^3,x,method=_RETURNV ERBOSE)`

output $\frac{2}{15}a^{-3}f*(\sec(f*x+e)-1)*(\cos(f*x+e)-5)*(-c*(\sec(f*x+e)-1))^{(1/2)}*c/(\cos(f*x+e)+1)^2/(\cos(f*x+e)-1)*\cos(f*x+e)^2*\cot(f*x+e)$

3.102.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx = \frac{2(c\cos(fx+e)^3 - 5c\cos(fx+e)^2)\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{15(a^3f\cos(fx+e)^2 + 2a^3f\cos(fx+e) + a^3f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^3,x, algorithm="fricas")
```

```
output -2/15*(c*cos(f*x + e)^3 - 5*c*cos(f*x + e)^2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((a^3*f*cos(f*x + e)^2 + 2*a^3*f*cos(f*x + e) + a^3*f)*sin(f*x + e))
```

3.102.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx = \int \frac{c\sqrt{-c\sec(e+fx)+c}\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \left(-\frac{c\sqrt{-c\sec(e+fx)+c}}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1}\right) dx$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(3/2)/(a+a*sec(f*x+e))**3,x)
```

```
output (Integral(c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3
```

3.102.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(80) = 160.

Time = 0.33 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.85

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{2\sqrt{2}c^{\frac{3}{2}} - \frac{3\sqrt{2}c^{\frac{3}{2}}\sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{3\sqrt{2}c^{\frac{3}{2}}\sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{7\sqrt{2}c^{\frac{3}{2}}\sin^6(fx+e)}{(\cos(fx+e)+1)^6} - \frac{3\sqrt{2}c^{\frac{3}{2}}\sin^8(fx+e)}{(\cos(fx+e)+1)^8}}{30a^3f\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)^{\frac{3}{2}}\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)^{\frac{3}{2}}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output `-1/30*(2*sqrt(2)*c^(3/2) - 3*sqrt(2)*c^(3/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 3*sqrt(2)*c^(3/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 7*sqrt(2)*c^(3/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 3*sqrt(2)*c^(3/2)*sin(f*x + e)^8/(cos(f*x + e) + 1)^8)/(a^3*f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^(3/2)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^(3/2))`

3.102.8 Giac [A] (verification not implemented)

Time = 0.74 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{\sqrt{2}\left(3\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{5}{2}} + 5\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^{\frac{3}{2}}c\right)}{30a^3cf}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output `-1/30*sqrt(2)*(3*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2) + 5*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c)/(a^3*c*f)`

3.102.9 Mupad [B] (verification not implemented)

Time = 18.94 (sec) , antiderivative size = 446, normalized size of antiderivative = 5.07

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{c(e^{e^{2i+fx^{2i}}+1})\sqrt{c-\frac{e^{-e^{li-fx^{li}}}}{2}+\frac{e^{e^{li+fx^{li}}}}{2}}}{15a^3 f(e^{e^{li+fx^{li}}}-1)(e^{e^{li+fx^{li}}}+1)} 2i$$

$$+ \frac{c(e^{e^{2i+fx^{2i}}+1})\sqrt{c-\frac{e^{-e^{li-fx^{li}}}}{2}+\frac{e^{e^{li+fx^{li}}}}{2}}}{15a^3 f(e^{e^{li+fx^{li}}}-1)(e^{e^{li+fx^{li}}}+1)^2} 28i$$

$$- \frac{c(e^{e^{2i+fx^{2i}}+1})\sqrt{c-\frac{e^{-e^{li-fx^{li}}}}{2}+\frac{e^{e^{li+fx^{li}}}}{2}}}{15a^3 f(e^{e^{li+fx^{li}}}-1)(e^{e^{li+fx^{li}}}+1)^3} 76i$$

$$+ \frac{c(e^{e^{2i+fx^{2i}}+1})\sqrt{c-\frac{e^{-e^{li-fx^{li}}}}{2}+\frac{e^{e^{li+fx^{li}}}}{2}}}{5a^3 f(e^{e^{li+fx^{li}}}-1)(e^{e^{li+fx^{li}}}+1)^4} 32i$$

$$- \frac{c(e^{e^{2i+fx^{2i}}+1})\sqrt{c-\frac{e^{-e^{li-fx^{li}}}}{2}+\frac{e^{e^{li+fx^{li}}}}{2}}}{5a^3 f(e^{e^{li+fx^{li}}}-1)(e^{e^{li+fx^{li}}}+1)^5} 16i$$

```
input int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)
```

```
output (c*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*28i)/(15*a^3*f*(exp(e*1i + f*x*1i) + 1)^2) - (c*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*2i)/(15*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^3) + (c*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*76i)/(15*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^3) + (c*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*32i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^4) - (c*(exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*16i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^5)
```


3.103 $\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx$

3.103.1 Optimal result	768
3.103.2 Mathematica [A] (verified)	768
3.103.3 Rubi [A] (verified)	769
3.103.4 Maple [A] (verified)	770
3.103.5 Fricas [A] (verification not implemented)	770
3.103.6 Sympy [F]	770
3.103.7 Maxima [B] (verification not implemented)	771
3.103.8 Giac [A] (verification not implemented)	771
3.103.9 Mupad [B] (verification not implemented)	772

3.103.1 Optimal result

Integrand size = 34, antiderivative size = 41

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx = \frac{2c \tan(e+fx)}{5f(a+a\sec(e+fx))^3 \sqrt{c-c\sec(e+fx)}}$$

output `2/5*c*tan(f*x+e)/f/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2)`

3.103.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.34

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx$$

$$= -\frac{\cos^3(e+fx) \csc\left(\frac{1}{2}(e+fx)\right) \sec^5\left(\frac{1}{2}(e+fx)\right) \sqrt{c-c\sec(e+fx)}}{20a^3 f}$$

input `Integrate[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^3,x]`

output `-1/20*(Cos[e + f*x]^3*Csc[(e + f*x)/2]*Sec[(e + f*x)/2]^5*Sqrt[c - c*Sec[e + f*x]])/(a^3*f)`

3.103.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a\sec(e+fx)+a)^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx$$

↓ 4441

$$\frac{2c\tan(e+fx)}{5f(a\sec(e+fx)+a)^3\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^3,x]`

output `(2*c*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]])`

3.103.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.103.4 Maple [A] (verified)

Time = 3.78 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

method	result	size
default	$-\frac{2\sqrt{-c(\sec(fx+e)-1)} \cos(fx+e)^2 \cot(fx+e)}{5a^3 f (\cos(fx+e)+1)^2}$	46

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^3,x,method=_RETURNV
ERBOSE)`

output `-2/5/a^3/f*(-c*(sec(f*x+e)-1))^(1/2)/(cos(f*x+e)+1)^2*cos(f*x+e)^2*cot(f*x
+e)`

3.103.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.80

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx$$

$$= -\frac{2\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}} \cos(fx+e)^3}{5(a^3 f \cos(fx+e)^2 + 2a^3 f \cos(fx+e) + a^3 f) \sin(fx+e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^3,x, algorithm
m="fricas")`

output `-2/5*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^3/((a^3*f*cos(f*
x + e)^2 + 2*a^3*f*cos(f*x + e) + a^3*f)*sin(f*x + e))`

3.103.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx = \frac{\int \frac{\sqrt{-c\sec(e+fx)+c\sec(e+fx)}}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx}{a^3}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(1/2)/(a+a*sec(f*x+e))**3,x)`

3.103. $\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx$

output `Integral(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x)/a**3`

3.103.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs. $2(37) = 74$.

Time = 0.32 (sec) , antiderivative size = 136, normalized size of antiderivative = 3.32

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{(a + a \sec(e + fx))^3} dx$$

$$= - \frac{\sqrt{2} \sqrt{c} - \frac{3 \sqrt{2} \sqrt{c} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{3 \sqrt{2} \sqrt{c} \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - \frac{\sqrt{2} \sqrt{c} \sin(fx+e)^6}{(\cos(fx+e)+1)^6}}{20 a^3 f \sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1} \sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output `-1/20*(sqrt(2)*sqrt(c) - 3*sqrt(2)*sqrt(c)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 3*sqrt(2)*sqrt(c)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - sqrt(2)*sqrt(c)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6)/(a^3*f*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) + 1)*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) - 1))`

3.103.8 Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{(a + a \sec(e + fx))^3} dx =$$

$$- \frac{\sqrt{2} \left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right)^{\frac{5}{2}} \operatorname{sgn} \left(\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right) \operatorname{sgn}(\cos(fx + e))}{20 a^3 c^2 f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output `-1/20*sqrt(2)*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2)*sgn(tan(1/2*f*x + 1/2*e))^3 + tan(1/2*f*x + 1/2*e)*sgn(cos(f*x + e))/(a^3*c^2*f)`

3.103.9 Mupad [B] (verification not implemented)

Time = 19.91 (sec) , antiderivative size = 441, normalized size of antiderivative = 10.76

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^3} dx = -\frac{(e^{2i+fx^2i}+1)\sqrt{c-\frac{e^{-e^{1i}-fx^{1i}}}{2}+\frac{e^{e^{1i}+fx^{1i}}}{2}}}{5a^3 f (e^{e^{1i}+fx^{1i}}-1)(e^{e^{1i}+fx^{1i}}+1)} 2i$$

$$+\frac{(e^{2i+fx^2i}+1)\sqrt{c-\frac{e^{-e^{1i}-fx^{1i}}}{2}+\frac{e^{e^{1i}+fx^{1i}}}{2}}}{5a^3 f (e^{e^{1i}+fx^{1i}}-1)(e^{e^{1i}+fx^{1i}}+1)^2} 8i$$

$$-\frac{(e^{2i+fx^2i}+1)\sqrt{c-\frac{e^{-e^{1i}-fx^{1i}}}{2}+\frac{e^{e^{1i}+fx^{1i}}}{2}}}{5a^3 f (e^{e^{1i}+fx^{1i}}-1)(e^{e^{1i}+fx^{1i}}+1)^3} 16i$$

$$+\frac{(e^{2i+fx^2i}+1)\sqrt{c-\frac{e^{-e^{1i}-fx^{1i}}}{2}+\frac{e^{e^{1i}+fx^{1i}}}{2}}}{5a^3 f (e^{e^{1i}+fx^{1i}}-1)(e^{e^{1i}+fx^{1i}}+1)^4} 16i$$

$$-\frac{(e^{2i+fx^2i}+1)\sqrt{c-\frac{e^{-e^{1i}-fx^{1i}}}{2}+\frac{e^{e^{1i}+fx^{1i}}}{2}}}{5a^3 f (e^{e^{1i}+fx^{1i}}-1)(e^{e^{1i}+fx^{1i}}+1)^5} 8i$$

```
input int((c - c/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)
```

```
output ((exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*8i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^2) - ((exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*2i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)) - ((exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*16i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^3) + ((exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*16i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^4) - ((exp(e*2i + f*x*2i) + 1)*(c - c/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(1/2)*8i)/(5*a^3*f*(exp(e*1i + f*x*1i) - 1)*(exp(e*1i + f*x*1i) + 1)^5)
```

3.104
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 \sqrt{c-c \sec(e+fx)}} dx$$

3.104.1 Optimal result 773
 3.104.2 Mathematica [C] (verified) 774
 3.104.3 Rubi [A] (verified) 774
 3.104.4 Maple [A] (verified) 777
 3.104.5 Fracas [A] (verification not implemented) 777
 3.104.6 Sympy [F] 778
 3.104.7 Maxima [F] 778
 3.104.8 Giac [A] (verification not implemented) 779
 3.104.9 Mupad [F(-1)] 779

3.104.1 Optimal result

Integrand size = 34, antiderivative size = 181

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 \sqrt{c-c \sec(e+fx)}} dx$$

$$= -\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2} \sqrt{c-c \sec(e+fx)}}\right)}{4\sqrt{2}a^3 \sqrt{c} f} + \frac{\tan(e+fx)}{5f(a+a \sec(e+fx))^3 \sqrt{c-c \sec(e+fx)}}$$

$$+ \frac{\tan(e+fx)}{6af(a+a \sec(e+fx))^2 \sqrt{c-c \sec(e+fx)}}$$

$$+ \frac{\tan(e+fx)}{4f(a^3+a^3 \sec(e+fx)) \sqrt{c-c \sec(e+fx)}}$$

```
output -1/8*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a^3/f*2
^(1/2)/c^(1/2)+1/5*tan(f*x+e)/f/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2)+
1/6*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(1/2)+1/4*tan(f*x+e
)/f/(a^3+a^3*sec(f*x+e))/(c-c*sec(f*x+e))^(1/2)
```

3.104.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.75 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.33

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}} dx$$

$$= \frac{\text{Hypergeometric2F1}\left(-\frac{5}{2}, 1, -\frac{3}{2}, \frac{1}{2}(1+\sec(e+fx))\right)\tan(e+fx)}{5f(a+a\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]]),x]`

output `(Hypergeometric2F1[-5/2, 1, -3/2, (1 + Sec[e + f*x])/2]*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]])`

3.104.3 Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4448, 3042, 4448, 3042, 4448, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)^3\sqrt{c-c\sec(e+fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)}{\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^3\sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)}} dx$$

$$\downarrow \text{4448}$$

$$\frac{\int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)^2\sqrt{c-c\sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow \text{3042}$$

3.104. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}} dx$

$$\begin{aligned}
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)^2 \sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow 4448 \\
& \frac{\int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)\sqrt{c-c\sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}} + \\
& \quad \frac{2a}{\tan(e+fx)} \\
& \quad \frac{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}} + \\
& \quad \frac{2a}{\tan(e+fx)} \\
& \quad \frac{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow 4448 \\
& \frac{\int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}} + \\
& \quad \frac{2a}{\tan(e+fx)} \\
& \quad \frac{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}} + \\
& \quad \frac{2a}{\tan(e+fx)} \\
& \quad \frac{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow 4282 \\
& \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)\sqrt{c-c\sec(e+fx)}} - \frac{\int \frac{1}{c^2 \tan^2(e+fx) + 2c} d \frac{c \tan(e+fx)}{\sqrt{c-c\sec(e+fx)}}}{af} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2 \sqrt{c-c\sec(e+fx)}} + \\
& \quad \frac{2a}{\tan(e+fx)} \\
& \quad \frac{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}}{5f(a\sec(e+fx)+a)^3 \sqrt{c-c\sec(e+fx)}} \\
& \quad \downarrow 216
\end{aligned}$$

3.104. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3 \sqrt{c-c\sec(e+fx)}} dx$

$$\frac{\frac{\tan(e+fx)}{f(a \sec(e+fx)+a)\sqrt{c-c \sec(e+fx)}} - \frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{\sqrt{2a}\sqrt{cf}}}{2a} + \frac{\tan(e+fx)}{3f(a \sec(e+fx)+a)^2\sqrt{c-c \sec(e+fx)}} + \frac{2a \tan(e+fx)}{5f(a \sec(e+fx)+a)^3\sqrt{c-c \sec(e+fx)}}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]]),x]`

output `Tan[e + f*x]/(5*f*(a + a*Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]]) + (Tan[e + f*x]/(3*f*(a + a*Sec[e + f*x])^2*Sqrt[c - c*Sec[e + f*x]]) + (-ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*a*Sqrt[c]*f)) + Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]])/(2*a))/(2*a)`

3.104.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4448 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))`

3.104.4 Maple [A] (verified)

Time = 3.41 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.17

method	result
default	$\frac{\sqrt{2} \left(3 \left((1 - \cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{5}{2}} - 5 \left((1 - \cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{3}{2}} + 15 \sqrt{(1 - \cos(fx+e))^2 \csc(fx+e)^2 - 1} + 15 \arctan \left(\frac{1}{\sqrt{(1 - \cos(fx+e))^2 \csc(fx+e)^2 - 1}} \right) \right)}{120 a^3 f \sqrt{\frac{c(1 - \cos(fx+e))^2 \csc(fx+e)^2}{(1 - \cos(fx+e))^2 \csc(fx+e)^2 - 1}} \sqrt{(1 - \cos(fx+e))^2 \csc(fx+e)^2 - 1}}$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNV
ERBOSE)`

output `1/120/a^3/f*2^(1/2)/(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*
csc(f*x+e)^2)^(1/2)*(3*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(5/2)-5*((1-cos(f
*x+e))^2*csc(f*x+e)^2-1)^(3/2)+15*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)+
15*arctan(1/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)))/((1-cos(f*x+e))^2*cs
c(f*x+e)^2-1)^(1/2)*(-cot(f*x+e)+csc(f*x+e))`

3.104.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 401, normalized size of antiderivative = 2.22

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \left[\frac{15 \sqrt{2} (\cos(fx + e)^2 + 2 \cos(fx + e) + 1) \sqrt{-c} \log \left(\frac{2 \sqrt{2} (\cos(fx + e)^2 + \cos(fx + e)) \sqrt{-c} \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}} + (3 c \cos(fx + e) + 2 \sqrt{2} (\cos(fx + e)^2 + \cos(fx + e) + 1) \sqrt{-c}}{(\cos(fx + e) - 1) \sin(fx + e)}} \right)}{240 (a^3 c f \cos(fx + e))^2 + 2 a^3 c f c} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x, algorithm
m="fricas")`

output `[-1/240*(15*sqrt(2)*(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)*sqrt(-c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) + 4*(37*cos(f*x + e)^3 + 40*cos(f*x + e)^2 + 15*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^3*c*f*cos(f*x + e)^2 + 2*a^3*c*f*cos(f*x + e) + a^3*c*f)*sin(f*x + e)), 1/120*(15*sqrt(2)*(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(37*cos(f*x + e)^3 + 40*cos(f*x + e)^2 + 15*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^3*c*f*cos(f*x + e)^2 + 2*a^3*c*f*cos(f*x + e) + a^3*c*f)*sin(f*x + e))]`

3.104.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{\int \frac{\sec(e+fx)}{\sqrt{-c \sec(e+fx)+c \sec^3(e+fx)+3\sqrt{-c \sec(e+fx)+c \sec^2(e+fx)+3\sqrt{-c \sec(e+fx)+c \sec(e+fx)+\sqrt{-c \sec(e+fx)+c}}} dx}}{a^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3 + 3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 + 3*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + sqrt(-c*sec(e + f*x) + c)), x)/a**3`

3.104.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)}} dx = \int \frac{\sec(fx + e)}{(a \sec(fx + e) + a)^3 \sqrt{-c \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)^3*sqrt(-c*sec(f*x + e) + c)), x)`

3.104. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 \sqrt{c-c \sec(e+fx)}} dx$

3.104.8 Giac [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.66

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{\sqrt{2} \left(\frac{15 \arctan\left(\frac{\sqrt{c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c}}{\sqrt{c}}\right)}{\sqrt{c}} - \frac{3 \left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c\right)^{\frac{5}{2}} c^{12-5} \left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c\right)^{\frac{3}{2}} c^{13+15} \sqrt{c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c c^{14}}}{c^{15}} \right)}{120 a^3 f}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(1/2),x, algorithm
m="giac")
```

```
output 1/120*sqrt(2)*(15*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c))/sqrt(
c) - (3*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2)*c^12 - 5*(c*tan(1/2*f*x + 1/2
*e)^2 - c)^(3/2)*c^13 + 15*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^14)/c^15)/
(a^3*f)
```

3.104.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 \sqrt{c - c \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx) \left(a + \frac{a}{\cos(e + fx)}\right)^3 \sqrt{c - \frac{c}{\cos(e + fx)}}} dx$$

```
input int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(1/2)),x)
```

```
output int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(1/2)), x)
```

3.105
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{3/2}} dx$$

3.105.1 Optimal result 780
 3.105.2 Mathematica [C] (verified) 781
 3.105.3 Rubi [A] (verified) 781
 3.105.4 Maple [A] (verified) 785
 3.105.5 Fricas [A] (verification not implemented) 786
 3.105.6 Sympy [F] 787
 3.105.7 Maxima [F] 787
 3.105.8 Giac [A] (verification not implemented) 787
 3.105.9 Mupad [F(-1)] 788

3.105.1 Optimal result

Integrand size = 34, antiderivative size = 212

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{3/2}} dx = -\frac{7 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{16\sqrt{2}a^3c^{3/2}f} - \frac{7 \tan(e+fx)}{16a^3f(c-c \sec(e+fx))^{3/2}} + \frac{\tan(e+fx)}{5f(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{3/2}} + \frac{7 \tan(e+fx)}{30af(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{3/2}} + \frac{7 \tan(e+fx)}{12f(a^3+a^3 \sec(e+fx))(c-c \sec(e+fx))^{3/2}}$$

output

```
-7/32*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a^3/c^(3/2)/f*2^(1/2)-7/16*tan(f*x+e)/a^3/f/(c-c*sec(f*x+e))^(3/2)+1/5*tan(f*x+e)/f/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2)+7/30*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(3/2)+7/12*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))/(c-c*sec(f*x+e))^(3/2)
```

3.105.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.84 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.30

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx = \frac{\text{Hypergeometric2F1}\left(-\frac{5}{2}, 2, -\frac{3}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan(e+fx)}{10a^3cf(1+\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(3/2)),x]`

output `(Hypergeometric2F1[-5/2, 2, -3/2, (1 + Sec[e + f*x])/2]*Tan[e + f*x])/(10*a^3*c*f*(1 + Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]])`

3.105.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4448, 3042, 4448, 3042, 4448, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a\csc(e+fx+\frac{\pi}{2})+a)^3(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx \\ & \quad \downarrow \text{4448} \\ & \frac{7 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)^2(c-c\sec(e+fx))^{3/2}} dx}{10a} + \frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{7 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)^2(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{10a} + \frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} \\ & \quad \downarrow \text{4448} \end{aligned}$$

3.105. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{7 \left(\frac{5 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)(c-c\sec(e+fx))^{3/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{3/2}} \right)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} + \\
 & \qquad \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{7 \left(\frac{5 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{3/2}} \right)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} + \\
 & \qquad \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow \text{4448} \\
 & \frac{7 \left(\frac{5 \left(\frac{3 \int \frac{\sec(e+fx)}{(c-c\sec(e+fx))^{3/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{3/2}} \right)}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{3/2}} \right)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} + \\
 & \qquad \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{7 \left(\frac{5 \left(\frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{3/2}} \right)}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{3/2}} \right)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} + \\
 & \qquad \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow \text{4283}
 \end{aligned}$$

3.105. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx$

$$7 \left(\frac{5 \left(\frac{3 \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c-c \sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right)}{6a} \right) + \frac{\tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}} \right) +$$

$$\frac{\tan(e+fx) 10a}{5f(a \sec(e+fx)+a)^3(c-c \sec(e+fx))^{3/2}}$$

↓ 3042

$$7 \left(\frac{5 \left(\frac{3 \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right)}{6a} \right) + \frac{\tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}} \right) +$$

$$\frac{\tan(e+fx) 10a}{5f(a \sec(e+fx)+a)^3(c-c \sec(e+fx))^{3/2}}$$

↓ 4282

$$7 \left(\frac{5 \left(\frac{3 \left(\int \frac{1}{c^2 \tan^2(e+fx) + 2c} dx - \frac{c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}} \right)}{2cf} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right)}{6a} \right) + \frac{\tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}}$$

$$\frac{10a \tan(e+fx)}{5f(a \sec(e+fx)+a)^3(c-c \sec(e+fx))^{3/2}}$$

↓ 216

$$7 \left(\frac{5 \left(\frac{3 \left(-\frac{\arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{2\sqrt{2}c^{3/2}f} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{3/2}} \right)}{6a} \right) + \frac{\tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{3/2}}$$

$$\frac{10a \tan(e+fx)}{5f(a \sec(e+fx)+a)^3(c-c \sec(e+fx))^{3/2}}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(3/2)),x]`

output `Tan[e + f*x]/(5*f*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(3/2)) + (7*(Tan[e + f*x]/(3*f*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(3/2)) + (5*(Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(3/2)) + (3*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])])/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2))))/(2*a)))/(6*a)))/(10*a)`

3.105. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{3/2}} dx$

3.105.3.1 Defintions of rubi rules used

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4282 Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]
```

```
rule 4283 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]
```

```
rule 4448 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.105.4 Maple [A] (verified)

Time = 3.58 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.30

method	result
default	$\frac{\sqrt{2} \left(139 \cos(fx+e)^3 \sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} + 21\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e)^2 - 105 \sin(fx+e)^2 \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) \cos(fx+e) \right)}{480a^3 f \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \sqrt{-c(\sec(fx+e))}}$

3.105. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{3/2}} dx$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x,method=_RETURNV
ERBOSE)`

output `-1/480/a^3/f*2^(1/2)*(139*cos(f*x+e)^3*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1)
)^(1/2)+21*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2-105*sin
(f*x+e)^2*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))*cos(f*x+e
)-175*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)-105*sin(f*x+e
)^2*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))-105*2^(1/2)*(-co
s(f*x+e)/(cos(f*x+e)+1))^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)/(-c*(se
c(f*x+e)-1))^(1/2)/(sec(f*x+e)-1)/c/(cos(f*x+e)+1)^3*tan(f*x+e)`

3.105.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 483, normalized size of antiderivative = 2.28

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx = \left[\frac{105\sqrt{2}(\cos(fx+e)^3 + \cos(fx+e)^2 - \cos(fx+e))}{\dots} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x, algorithm
m="fricas")`

output `[-1/960*(105*sqrt(2)*(cos(f*x + e)^3 + cos(f*x + e)^2 - cos(f*x + e) - 1)*
sqrt(-c)*log((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos
os(f*x + e) - c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((co
s(f*x + e) - 1)*sin(f*x + e)))*sin(f*x + e) + 4*(139*cos(f*x + e)^4 + 21*c
os(f*x + e)^3 - 175*cos(f*x + e)^2 - 105*cos(f*x + e))*sqrt((c*cos(f*x + e
) - c)/cos(f*x + e)))/((a^3*c^2*f*cos(f*x + e)^3 + a^3*c^2*f*cos(f*x + e)
^2 - a^3*c^2*f*cos(f*x + e) - a^3*c^2*f)*sin(f*x + e)), 1/480*(105*sqrt(2)*
(cos(f*x + e)^3 + cos(f*x + e)^2 - cos(f*x + e) - 1)*sqrt(c)*arctan(sqrt(2
)*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x +
e)))*sin(f*x + e) - 2*(139*cos(f*x + e)^4 + 21*cos(f*x + e)^3 - 175*cos(f*x
+ e)^2 - 105*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^
3*c^2*f*cos(f*x + e)^3 + a^3*c^2*f*cos(f*x + e)^2 - a^3*c^2*f*cos(f*x + e)
- a^3*c^2*f)*sin(f*x + e)]]`

3.105.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx = \frac{\int \frac{\sec(e+fx)}{-c\sqrt{-c\sec(e+fx)+c\sec^4(e+fx)-2c\sqrt{-c\sec(e+fx)+c\sec^3(e+fx)+2c}}}{a^3} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**(3/2),x)`

output `Integral(sec(e + f*x)/(-c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4 - 2*c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**3 + 2*c*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c*sqrt(-c*sec(e + f*x) + c)), x)/a**3`

3.105.7 Maxima [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\sec(fx+e)}{(a\sec(fx+e)+a)^3(-c\sec(fx+e)+c)^{3/2}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x, algorithm m="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)^3*(-c*sec(f*x + e) + c)^(3/2)), x)`

3.105.8 Giac [A] (verification not implemented)

Time = 0.58 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.73

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx = \frac{\sqrt{2} \left(105 \sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}} \right) - \frac{15 \sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)}}{\tan(\frac{1}{2} fx + \frac{1}{2} e)} \right)}{a^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(3/2),x, algorithm m="giac")`

3.105. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx$

output $1/480*\sqrt{2}*(105*\sqrt{c}*\arctan(\sqrt{c*\tan(1/2*f*x + 1/2*e)^2 - c})/\sqrt{c}) - 15*\sqrt{c*\tan(1/2*f*x + 1/2*e)^2 - c}/\tan(1/2*f*x + 1/2*e)^2 - 2*(3*(c*\tan(1/2*f*x + 1/2*e)^2 - c)^{(5/2)}*c^8 - 10*(c*\tan(1/2*f*x + 1/2*e)^2 - c)^{(3/2)}*c^9 + 45*\sqrt{c*\tan(1/2*f*x + 1/2*e)^2 - c}*c^{10})/c^{10}/(a^3*c^2*f)$

3.105.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{3/2}} dx = \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^3 \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(3/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(3/2)), x)`

3.106
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{5/2}} dx$$

3.106.1 Optimal result	789
3.106.2 Mathematica [C] (verified)	790
3.106.3 Rubi [A] (verified)	790
3.106.4 Maple [A] (verified)	797
3.106.5 Fricas [A] (verification not implemented)	798
3.106.6 Sympy [F]	798
3.106.7 Maxima [F]	799
3.106.8 Giac [A] (verification not implemented)	799
3.106.9 Mupad [F(-1)]	800

3.106.1 Optimal result

Integrand size = 34, antiderivative size = 246

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{5/2}} dx = -\frac{63 \arctan\left(\frac{\sqrt{c} \tan(e+fx)}{\sqrt{2}\sqrt{c-c \sec(e+fx)}}\right)}{128\sqrt{2}a^3c^{5/2}f} - \frac{21 \tan(e+fx)}{32a^3f(c-c \sec(e+fx))^{5/2}} + \frac{\tan(e+fx)}{5f(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{5/2}} + \frac{3 \tan(e+fx)}{10af(a+a \sec(e+fx))^2(c-c \sec(e+fx))^{5/2}} + \frac{21 \tan(e+fx)}{20f(a^3+a^3 \sec(e+fx))(c-c \sec(e+fx))^{5/2}} - \frac{63 \tan(e+fx)}{128a^3cf(c-c \sec(e+fx))^{3/2}}$$

output

```
-63/256*arctan(1/2*c^(1/2)*tan(f*x+e)*2^(1/2)/(c-c*sec(f*x+e))^(1/2))/a^3/c^(5/2)/f*2^(1/2)-21/32*tan(f*x+e)/a^3/f/(c-c*sec(f*x+e))^(5/2)+1/5*tan(f*x+e)/f/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2)+3/10*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2/(c-c*sec(f*x+e))^(5/2)+21/20*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))/(c-c*sec(f*x+e))^(5/2)-63/128*tan(f*x+e)/a^3/c/f/(c-c*sec(f*x+e))^(3/2)
```

3.106.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.00 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.26

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx = \frac{\text{Hypergeometric2F1}\left(-\frac{5}{2}, 3, -\frac{3}{2}, \frac{1}{2}(1+\sec(e+fx))\right) \tan(e+fx)}{20a^3c^2f(1+\sec(e+fx))^3\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(5/2)), x]`

output `(Hypergeometric2F1[-5/2, 3, -3/2, (1 + Sec[e + f*x])/2]*Tan[e + f*x])/(20*a^3*c^2*f*(1 + Sec[e + f*x])^3*Sqrt[c - c*Sec[e + f*x]])`

3.106.3 Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4448, 3042, 4448, 3042, 4448, 3042, 4283, 3042, 4283, 3042, 4282, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a\csc(e+fx+\frac{\pi}{2})+a)^3(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx \\ & \quad \downarrow \text{4448} \\ & \frac{9 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)^2(c-c\sec(e+fx))^{5/2}} dx}{10a} + \frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{9 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)^2(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx}{10a} + \frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow \text{4448} \end{aligned}$$

3.106. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx$

$$\begin{aligned}
 & 9 \left(\frac{7 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)(c-c\sec(e+fx))^{5/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & 9 \left(\frac{7 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4448} \\
 & 9 \left(\frac{7 \left(\frac{5 \int \frac{\sec(e+fx)}{(c-c\sec(e+fx))^{5/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}} \right)}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & 9 \left(\frac{7 \left(\frac{5 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}} \right)}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2(c-c\sec(e+fx))^{5/2}} \right) \\
 & \quad + \frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4283}
 \end{aligned}$$

3.106. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx$

$$9 \left(\frac{7 \left(\frac{5 \left(\frac{3 \int \frac{\sec(e+fx)}{(c-c \sec(e+fx))^{3/2}} dx}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right)}{6a} \right) + \frac{\tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}}$$

$$\frac{\tan(e+fx) 10a}{5f(a \sec(e+fx)+a)^3(c-c \sec(e+fx))^{5/2}}$$

↓ 3042

$$9 \left(\frac{7 \left(\frac{5 \left(\frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(c-c \csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{8c} - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right)}{2a} + \frac{\tan(e+fx)}{f(a \sec(e+fx)+a)(c-c \sec(e+fx))^{5/2}} \right)}{6a} \right) + \frac{\tan(e+fx)}{3f(a \sec(e+fx)+a)^2(c-c \sec(e+fx))^{5/2}}$$

$$\frac{\tan(e+fx) 10a}{5f(a \sec(e+fx)+a)^3(c-c \sec(e+fx))^{5/2}}$$

↓ 4283

$$\left(\frac{3 \left(\frac{\int \frac{\sec(e+fx)}{\sqrt{c-c\sec(e+fx)}} dx}{4c} - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}} \right) - \frac{\tan(e+fx)}{4f(c-c\sec(e+fx))^{5/2}}}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}} \right) + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2}$$

$$\frac{10a \tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}}$$

↓ 3042

$$\left(\left(\left(\left(\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx \right) - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}} \right) - \frac{\tan(e+fx)}{4f(c-c\sec(e+fx))^{5/2}} \right) - \frac{\tan(e+fx)}{4f(c-c\sec(e+fx))^{5/2}} \right) + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}} \right) + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}$$

$$\frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}}$$

↓ 4282

$$\left(\frac{ \left(\frac{ \int \frac{1}{c^2 \tan^2(e+fx) + 2c} dx \frac{c \tan(e+fx)}{\sqrt{c-c \sec(e+fx)}} - \frac{\tan(e+fx)}{2f(c-c \sec(e+fx))^{3/2}} \right) - \frac{\tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}} \right) + \frac{\tan(e+fx)}{f(a \sec(e+fx) + a)(c-c \sec(e+fx))^{5/2}} \right) + \frac{\tan(e+fx)}{3f}$$

$$\frac{\tan(e+fx)}{5f(a \sec(e+fx) + a)^3(c-c \sec(e+fx))^{5/2}}$$

↓ 216

3.106. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{5/2}} dx$

$$\frac{\left(\frac{\left(\frac{\arctan\left(\frac{\sqrt{c}\tan(e+fx)}{\sqrt{2}\sqrt{c-c\sec(e+fx)}}\right) - \frac{\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}}}{2\sqrt{2}c^{3/2}f} \right) - \frac{\tan(e+fx)}{4f(c-c\sec(e+fx))^{5/2}}}{8c} \right)}{2a} + \frac{\tan(e+fx)}{f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}} \right)}{6a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)(c-c\sec(e+fx))^{5/2}}$$

$$\frac{\tan(e+fx)}{5f(a\sec(e+fx)+a)^3(c-c\sec(e+fx))^{5/2}}$$

```
input Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(5/2)),x]
```

```
output Tan[e + f*x]/(5*f*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x])^(5/2)) + (9*(Tan[e + f*x]/(3*f*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x])^(5/2)) + (7*(Tan[e + f*x]/(f*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x])^(5/2)) + (5*(-1/4*Tan[e + f*x]/(f*(c - c*Sec[e + f*x])^(5/2)) + (3*(-1/2*ArcTan[(Sqrt[c]*Tan[e + f*x])/(Sqrt[2]*Sqrt[c - c*Sec[e + f*x]])]/(Sqrt[2]*c^(3/2)*f) - Tan[e + f*x]/(2*f*(c - c*Sec[e + f*x])^(3/2)))/(8*c)))/(2*a)))/(6*a)))/(10*a)
```

3.106.3.1 Defintions of rubi rules used

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.106. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx$

rule 4282 `Int[csc[(e_.) + (f_.)*(x_.)]/Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4283 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

rule 4448 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))`

3.106.4 Maple [A] (verified)

Time = 3.54 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.11

method	result
default	$\frac{\sqrt{2} \left(257\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \cos(fx+e)^4 - 354 \cos(fx+e)^3 \sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} + 315 \sin(fx+e)^4 \arctan\left(\frac{\sqrt{2}}{2\sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}}}\right) - 588\sqrt{2} \right)}{1280a^3 f \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \sqrt{-c(\sec(fx+e)-1)}}$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x,method=_RETURNV ERBOSE)`

output `1/1280/a^3/f*2^(1/2)*(257*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^4-354*cos(f*x+e)^3*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)+315*sin(f*x+e)^4*arctan(1/2*2^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))-588*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2+210*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)+315*2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2))/(-cos(f*x+e)/(cos(f*x+e)+1))^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)/(sec(f*x+e)-1)^2/c^2/(cos(f*x+e)+1)^3*tan(f*x+e)*sec(f*x+e)`

3.106. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c-c \sec(e+fx))^{5/2}} dx$

3.106.5 Fracas [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.87

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx = \left[\frac{315\sqrt{2}(\cos(fx+e)^4 - 2\cos(fx+e)^2 + 1)\sqrt{-c}\log}{\dots} \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x, algorithm
m="fricas")
```

```
output [-1/2560*(315*sqrt(2)*(cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*sqrt(-c)*log
((2*sqrt(2)*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(-c)*sqrt((c*cos(f*x + e)
- c)/cos(f*x + e)) + (3*c*cos(f*x + e) + c)*sin(f*x + e))/((cos(f*x + e) -
1)*sin(f*x + e)))*sin(f*x + e) + 4*(257*cos(f*x + e)^5 - 354*cos(f*x + e)
^4 - 588*cos(f*x + e)^3 + 210*cos(f*x + e)^2 + 315*cos(f*x + e))*sqrt((c*c
os(f*x + e) - c)/cos(f*x + e)))/((a^3*c^3*f*cos(f*x + e)^4 - 2*a^3*c^3*f*c
os(f*x + e)^2 + a^3*c^3*f)*sin(f*x + e)), 1/1280*(315*sqrt(2)*(cos(f*x + e)
)^4 - 2*cos(f*x + e)^2 + 1)*sqrt(c)*arctan(sqrt(2)*sqrt((c*cos(f*x + e) -
c)/cos(f*x + e))*cos(f*x + e)/(sqrt(c)*sin(f*x + e)))*sin(f*x + e) - 2*(25
7*cos(f*x + e)^5 - 354*cos(f*x + e)^4 - 588*cos(f*x + e)^3 + 210*cos(f*x +
e)^2 + 315*cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^3*c
^3*f*cos(f*x + e)^4 - 2*a^3*c^3*f*cos(f*x + e)^2 + a^3*c^3*f)*sin(f*x + e)
)]
```

3.106.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx = \int \frac{c^2\sqrt{-c\sec(e+fx)+c\sec^5(e+fx)+c^2\sqrt{-c\sec(e+fx)+c\sec^4(e+fx)-2c^2}}{\dots}}{\dots}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c-c*sec(f*x+e))**(5/2),x)
```

```
output Integral(sec(e + f*x)/(c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**5 + c
**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**4 - 2*c**2*sqrt(-c*sec(e + f*x)
+ c)*sec(e + f*x)**3 - 2*c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x)**2 +
c**2*sqrt(-c*sec(e + f*x) + c)*sec(e + f*x) + c**2*sqrt(-c*sec(e + f*x) +
c)), x)/a**3
```

3.106.7 Maxima [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\sec(fx+e)}{(a\sec(fx+e)+a)^3(-c\sec(fx+e)+c)^{5/2}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x, algorithm m="maxima")`

output `integrate(sec(f*x + e)/((a*sec(f*x + e) + a)^3*(-c*sec(f*x + e) + c)^(5/2)), x)`

3.106.8 Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.75

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx = \frac{\sqrt{2} \left(315 \sqrt{c} \arctan \left(\frac{\sqrt{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}}{\sqrt{c}} \right) - 5 \left(17 \left(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c \right)^{3/2} \right) \right)}{c^{10} (a^3 c^3 f)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c-c*sec(f*x+e))^(5/2),x, algorithm m="giac")`

output `1/1280*sqrt(2)*(315*sqrt(c)*arctan(sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)/sqrt(c)) - 5*(17*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c + 15*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^2)/(c^2*tan(1/2*f*x + 1/2*e)^4) - 8*((c*tan(1/2*f*x + 1/2*e)^2 - c)^(5/2)*c^8 - 5*(c*tan(1/2*f*x + 1/2*e)^2 - c)^(3/2)*c^9 + 30*sqrt(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^10)/c^10/(a^3*c^3*f)`

3.106.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c-c\sec(e+fx))^{5/2}} dx = \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^3 \left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(5/2)),x)`output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c - c/cos(e + f*x))^(5/2)), x)`

3.107 $\int \sec(e+fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e+fx))^{5/2} dx$

3.107.1 Optimal result	801
3.107.2 Mathematica [A] (verified)	801
3.107.3 Rubi [A] (verified)	802
3.107.4 Maple [B] (verified)	803
3.107.5 Fricas [B] (verification not implemented)	803
3.107.6 Sympy [F(-1)]	804
3.107.7 Maxima [B] (verification not implemented)	804
3.107.8 Giac [F]	805
3.107.9 Mupad [B] (verification not implemented)	805

3.107.1 Optimal result

Integrand size = 36, antiderivative size = 43

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{5/2} dx = \frac{a(c - c \sec(e + fx))^{5/2} \tan(e + fx)}{3f \sqrt{a + a \sec(e + fx)}}$$

```
output 1/3*a*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)
```

3.107.2 Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.63

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{5/2} dx = \frac{ac^3 \sec(e + fx) (3 - 3 \sec(e + fx) + \sec^2(e + fx)) \tan(e + fx)}{3f \sqrt{a(1 + \sec(e + fx))} \sqrt{c - c \sec(e + fx)}}$$

```
input Integrate[Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(5/2),x]
```

```
output -1/3*(a*c^3*Sec[e + f*x]*(3 - 3*Sec[e + f*x] + Sec[e + f*x]^2)*Tan[e + f*x])/
(f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])
```

3.107.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx) \sqrt{a \sec(e + fx) + a} (c - c \sec(e + fx))^{5/2} dx$$

↓ 3042

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{a \csc\left(e + fx + \frac{\pi}{2}\right) + a} \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx$$

↓ 4441

$$\frac{a \tan(e + fx) (c - c \sec(e + fx))^{5/2}}{3f \sqrt{a \sec(e + fx) + a}}$$

input `Int[Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(5/2),x]`

output `(a*(c - c*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(3*f*Sqrt[a + a*Sec[e + f*x]])`

3.107.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.107.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(37) = 74$.

Time = 3.34 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.05

method	result	size
default	$\frac{(\sec(fx+e)-1)^2(7\cos(fx+e)^2-4\cos(fx+e)+1)\sqrt{-c(\sec(fx+e)-1)}c^2\sqrt{a(\sec(fx+e)+1)}(\cos(fx+e)+1)\csc(fx+e)}{3f(\cos(fx+e)-1)^2}$	88
risch	$\frac{2ic^2\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}(3e^{5i(fx+e)}-6e^{4i(fx+e)}+10e^{3i(fx+e)}-6e^{2i(fx+e)}+3e^{i(fx+e)})}{3(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)(1+e^{2i(fx+e)})^2f}}$	165

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)*(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `1/3/f*(sec(f*x+e)-1)^2*(7*cos(f*x+e)^2-4*cos(f*x+e)+1)*(-c*(sec(f*x+e)-1))^(1/2)*c^2*(a*(sec(f*x+e)+1))^(1/2)*(cos(f*x+e)+1)/(cos(f*x+e)-1)^2*csc(f*x+e)`

3.107.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 93 vs. $2(37) = 74$.

Time = 0.26 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.16

$$\int \sec(e+fx)\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}dx = \frac{(3c^2\cos(fx+e)^2-3c^2\cos(fx+e)+c^2)\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{3f\cos(fx+e)^2\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)*(a+a*sec(f*x+e))^(1/2),x,algorithm="fracas")`

output `1/3*(3*c^2*cos(f*x+e)^2-3*c^2*cos(f*x+e)+c^2)*sqrt((a*cos(f*x+e)+a)/cos(f*x+e))*sqrt((c*cos(f*x+e)-c)/cos(f*x+e))/(f*cos(f*x+e))^2*sin(f*x+e)`

3.107.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{5/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(5/2)*(a+a*sec(f*x+e))**(1/2),x)`

output `Timed out`

3.107.7 Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 638 vs. $2(37) = 74$.

Time = 0.39 (sec) , antiderivative size = 638, normalized size of antiderivative = 14.84

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{5/2} dx = \frac{2(30c^2 \cos(3fx + 3e) \sin(2fx + 2e) - 9c^2 \cos(2fx + 2e) \sin(fx + e) - 3c^2 \sin(fx + e) \cos(5fx + 5e) - 3c^2 \sin(fx + e) \cos(4fx + 4e) + 10c^2 \sin(3fx + 3e) - 6c^2 \sin(2fx + 2e) + 3c^2 \sin(fx + e)) \cos(6fx + 6e) + 9(c^2 \sin(4fx + 4e) + c^2 \sin(2fx + 2e)) \cos(5fx + 5e) - 3(10c^2 \sin(3fx + 3e) + 3c^2 \sin(fx + e)) \cos(4fx + 4e) + (3c^2 \cos(5fx + 5e) - 6c^2 \cos(4fx + 4e) + 10c^2 \cos(3fx + 3e) - 6c^2 \cos(2fx + 2e) + 3c^2 \cos(fx + e)) \sin(6fx + 6e) - 3(3c^2 \cos(4fx + 4e) + 3c^2 \cos(2fx + 2e) + c^2) \sin(5fx + 5e) + 3(10c^2 \cos(3fx + 3e) + 3c^2 \cos(fx + e) + 2c^2) \sin(4fx + 4e) - 10(3c^2 \cos(2fx + 2e) + c^2) \sin(3fx + 3e) + 3(3c^2 \cos(fx + e) + 2c^2) \sin(2fx + 2e)) \sqrt{a} \sqrt{c}}{(2(3 \cos(4fx + 4e) + 3 \cos(2fx + 2e) + 1) \cos(6fx + 6e) + \cos(6fx + 6e)^2 + 6(3 \cos(2fx + 2e) + 1) \cos(4fx + 4e) + 9 \cos(4fx + 4e)^2 + 9 \cos(2fx + 2e)^2 + 6(\sin(4fx + 4e) + \sin(2fx + 2e)) \sin(6fx + 6e) + \sin(6fx + 6e)^2 + 9 \sin(4fx + 4e)^2 + 18 \sin(4fx + 4e) \sin(2fx + 2e) + 9 \sin(2fx + 2e)^2 + 6 \cos(2fx + 2e) + 1) f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)*(a+a*sec(f*x+e))^(1/2),x, algo rithm="maxima")`

output `2/3*(30*c^2*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) - 9*c^2*cos(2*f*x + 2*e)*sin(f*x + e) - 3*c^2*sin(f*x + e) - (3*c^2*sin(5*f*x + 5*e) - 6*c^2*sin(4*f*x + 4*e) + 10*c^2*sin(3*f*x + 3*e) - 6*c^2*sin(2*f*x + 2*e) + 3*c^2*sin(f*x + e))*cos(6*f*x + 6*e) + 9*(c^2*sin(4*f*x + 4*e) + c^2*sin(2*f*x + 2*e))*cos(5*f*x + 5*e) - 3*(10*c^2*sin(3*f*x + 3*e) + 3*c^2*sin(f*x + e))*cos(4*f*x + 4*e) + (3*c^2*cos(5*f*x + 5*e) - 6*c^2*cos(4*f*x + 4*e) + 10*c^2*cos(3*f*x + 3*e) - 6*c^2*cos(2*f*x + 2*e) + 3*c^2*cos(f*x + e))*sin(6*f*x + 6*e) - 3*(3*c^2*cos(4*f*x + 4*e) + 3*c^2*cos(2*f*x + 2*e) + c^2)*sin(5*f*x + 5*e) + 3*(10*c^2*cos(3*f*x + 3*e) + 3*c^2*cos(f*x + e) + 2*c^2)*sin(4*f*x + 4*e) - 10*(3*c^2*cos(2*f*x + 2*e) + c^2)*sin(3*f*x + 3*e) + 3*(3*c^2*cos(f*x + e) + 2*c^2)*sin(2*f*x + 2*e))*sqrt(a)*sqrt(c)/((2*(3*cos(4*f*x + 4*e) + 3*cos(2*f*x + 2*e) + 1)*cos(6*f*x + 6*e) + cos(6*f*x + 6*e)^2 + 6*(3*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + 9*cos(4*f*x + 4*e)^2 + 9*cos(2*f*x + 2*e)^2 + 6*(sin(4*f*x + 4*e) + sin(2*f*x + 2*e))*sin(6*f*x + 6*e) + sin(6*f*x + 6*e)^2 + 9*sin(4*f*x + 4*e)^2 + 18*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 9*sin(2*f*x + 2*e)^2 + 6*cos(2*f*x + 2*e) + 1)*f)`

3.107. $\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{5/2} dx$

3.107.8 Giac [F]

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{5/2} dx = \int \sqrt{a \sec(fx + e) + a} (-c \sec(fx + e) + c)^{5/2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)*(a+a*sec(f*x+e))^(1/2),x, algo
rithm="giac")`

output `sage0*x`

3.107.9 Mupad [B] (verification not implemented)

Time = 15.32 (sec) , antiderivative size = 136, normalized size of antiderivative = 3.16

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{5/2} dx = \frac{2c^2 \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}} \sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}} (10 \sin(e + fx) - 12 \sin(2e + 2fx) + 13 \sin(3e + 3fx) - 6 \sin(4e + 4fx) + 3 \sin(5e + 5fx))}{3f(\cos(2e + 2fx) - 2 \cos(4e + 4fx) - \cos(6e + 6fx) + 2)}$$

input `int(((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x),x
)`

output `(2*c^2*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2)*((c*(cos(e + f*x) - 1))
/cos(e + f*x))^(1/2)*(10*sin(e + f*x) - 12*sin(2*e + 2*f*x) + 13*sin(3*e +
3*f*x) - 6*sin(4*e + 4*f*x) + 3*sin(5*e + 5*f*x)))/(3*f*(cos(2*e + 2*f*x)
- 2*cos(4*e + 4*f*x) - cos(6*e + 6*f*x) + 2))`

3.108 $\int \sec(e+fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e+fx))^{3/2} dx$

3.108.1 Optimal result	806
3.108.2 Mathematica [A] (verified)	806
3.108.3 Rubi [A] (verified)	807
3.108.4 Maple [A] (verified)	808
3.108.5 Fricas [B] (verification not implemented)	808
3.108.6 Sympy [F]	809
3.108.7 Maxima [B] (verification not implemented)	809
3.108.8 Giac [F]	810
3.108.9 Mupad [B] (verification not implemented)	810

3.108.1 Optimal result

Integrand size = 36, antiderivative size = 43

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{3/2} dx = \frac{a(c - c \sec(e + fx))^{3/2} \tan(e + fx)}{2f \sqrt{a + a \sec(e + fx)}}$$

output `1/2*a*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)`

3.108.2 Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.40

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{3/2} dx = \frac{ac^2(-2 + \sec(e + fx)) \sec(e + fx) \tan(e + fx)}{2f \sqrt{a(1 + \sec(e + fx))} \sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(3/2),x]`

output `(a*c^2*(-2 + Sec[e + f*x])*Sec[e + f*x]*Tan[e + f*x])/(2*f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.108.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx) \sqrt{a \sec(e + fx) + a} (c - c \sec(e + fx))^{3/2} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{a \csc\left(e + fx + \frac{\pi}{2}\right) + a} \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx$$

$$\downarrow \text{4441}$$

$$\frac{a \tan(e + fx) (c - c \sec(e + fx))^{3/2}}{2f \sqrt{a \sec(e + fx) + a}}$$

input `Int[Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(3/2),x]`

output `(a*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(2*f*Sqrt[a + a*Sec[e + f*x]])`

3.108.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.108.4 Maple [A] (verified)

Time = 3.46 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.72

method	result	size
default	$-\frac{(\sec(fx+e)-1)(3\cos(fx+e)-1)\sqrt{-c(\sec(fx+e)-1)}c\sqrt{a(\sec(fx+e)+1)}(\cos(fx+e)+1)\csc(fx+e)}{2f(\cos(fx+e)-1)}$	74
risch	$\frac{2ic\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}(e^{3i(fx+e)}-e^{2i(fx+e)}+e^{i(fx+e)})}{(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)(1+e^{2i(fx+e)})f}}$	137

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2/f*(sec(f*x+e)-1)*(3*cos(f*x+e)-1)*(-c*(sec(f*x+e)-1))^(1/2)*c*(a*(sec(f*x+e)+1))^(1/2)*(cos(f*x+e)+1)/(cos(f*x+e)-1)*csc(f*x+e)`

3.108.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. $2(37) = 74$.

Time = 0.27 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.81

$$\int \sec(e+fx)\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{3/2}dx = \frac{(2c\cos(fx+e)-c)\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{2f\cos(fx+e)\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2),x,algorithm="fracas")`

output `1/2*(2*c*cos(f*x + e) - c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)*sin(f*x + e))`

3.108.6 Sympy [F]

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{3/2} dx = \int \sqrt{a(\sec(e + fx) + 1)} (-c(\sec(e + fx) - 1))^{3/2} \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(3/2)*(a+a*sec(f*x+e))**(1/2),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*(-c*(sec(e + f*x) - 1))**(3/2)*sec(e + f*x), x)`

3.108.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 298 vs. 2(37) = 74.

Time = 0.37 (sec) , antiderivative size = 298, normalized size of antiderivative = 6.93

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{3/2} dx = \frac{2(2c \cos(3fx + 3e) \sin(2fx + 2e) - 2c \cos(2fx + 2e) \sin(fx + e) - (c \sin(3fx + 3e) - c \sin(2fx + 2e) + c \sin(fx + e)) \cos(4fx + 4e) + (c \cos(3fx + 3e) - c \cos(2fx + 2e) + c \cos(fx + e)) \sin(4fx + 4e) - (2c \cos(2fx + 2e) + c) \sin(3fx + 3e) + (2c \cos(fx + e) + c) \sin(2fx + 2e) - c \sin(fx + e)) \sqrt{a} \sqrt{c}}{(2(2 \cos(2fx + 2e) + 1) \cos(4fx + 4e) + \cos(4fx + 4e)^2 + 4 \cos(2fx + 2e)^2 + \sin(4fx + 4e)^2 + 4 \sin(4fx + 4e) \sin(2fx + 2e) + 4 \sin(2fx + 2e)^2 + 4 \cos(2fx + 2e) + 1) * f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `2*(2*c*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) - 2*c*cos(2*f*x + 2*e)*sin(f*x + e) - (c*sin(3*f*x + 3*e) - c*sin(2*f*x + 2*e) + c*sin(f*x + e))*cos(4*f*x + 4*e) + (c*cos(3*f*x + 3*e) - c*cos(2*f*x + 2*e) + c*cos(f*x + e))*sin(4*f*x + 4*e) - (2*c*cos(2*f*x + 2*e) + c)*sin(3*f*x + 3*e) + (2*c*cos(f*x + e) + c)*sin(2*f*x + 2*e) - c*sin(f*x + e))*sqrt(a)*sqrt(c)/((2*(2*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 4*cos(2*f*x + 2*e)^2 + sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) + 1)*f)`

3.108.8 Giac [F]

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{3/2} dx = \int \sqrt{a \sec(fx + e) + a} (-c \sec(fx + e) + c)^{\frac{3}{2}} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2),x, algo
rithm="giac")`

output `sage0*x`

3.108.9 Mupad [B] (verification not implemented)

Time = 14.01 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.81

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} (c - c \sec(e + fx))^{3/2} dx = \frac{c \sqrt{c - \frac{c}{\cos(e+fx)}} \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}} (\sin(e + fx) - \sin(2e + 2fx) + \sin(3e + 3fx))}{f \sin(2e + 2fx)^2}$$

input `int(((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(3/2))/cos(e + f*x),x
)`

output `(c*(c - c/cos(e + f*x))^(1/2)*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2)*
(sin(e + f*x) - sin(2*e + 2*f*x) + sin(3*e + 3*f*x)))/(f*sin(2*e + 2*f*x)^
2)`

3.109 $\int \sec(e+fx) \sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}$

3.109.1 Optimal result	811
3.109.2 Mathematica [A] (verified)	811
3.109.3 Rubi [A] (verified)	812
3.109.4 Maple [A] (verified)	813
3.109.5 Fricas [A] (verification not implemented)	813
3.109.6 Sympy [F]	813
3.109.7 Maxima [A] (verification not implemented)	814
3.109.8 Giac [F]	814
3.109.9 Mupad [B] (verification not implemented)	815

3.109.1 Optimal result

Integrand size = 36, antiderivative size = 41

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)} dx$$

$$= -\frac{c \sqrt{a + a \sec(e + fx)} \tan(e + fx)}{f \sqrt{c - c \sec(e + fx)}}$$

output `-c*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)`

3.109.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.24

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)} dx$$

$$= -\frac{c \sec(e + fx) \sqrt{a(1 + \sec(e + fx))} \tan\left(\frac{1}{2}(e + fx)\right)}{f \sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]],x]`

output `-((c*Sec[e + f*x]*Sqrt[a*(1 + Sec[e + f*x])]*Tan[(e + f*x)/2])/(f*Sqrt[c - c*Sec[e + f*x]]))`

3.109.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx) \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{a \csc\left(e + fx + \frac{\pi}{2}\right) + a} \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4441}$$

$$-\frac{c \tan(e + fx) \sqrt{a \sec(e + fx) + a}}{f \sqrt{c - c \sec(e + fx)}}$$

input `Int[Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]],x]`

output `-((c*Sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]]))`

3.109.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.109.4 Maple [A] (verified)

Time = 3.30 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.15

method	result	size
default	$-\frac{\sqrt{a(\sec(fx+e)+1)}\sqrt{-c(\sec(fx+e)-1)}\sin(fx+e)}{f(\cos(fx+e)-1)}$	47
risch	$\frac{2i\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}e^{i(fx+e)}}{(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)f}$	102

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)*(c-c*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/f*(a*(sec(f*x+e)+1))^(1/2)*(-c*(sec(f*x+e)-1))^(1/2)*sin(f*x+e)/(cos(f*x+e)-1)
```

3.109.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.37

$$\int \sec(e+fx)\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}dx = \frac{\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{f\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)*(c-c*sec(f*x+e))^(1/2),x, algorithm="fracas")
```

```
output sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*sin(f*x + e))
```

3.109.6 Sympy [F]

$$\begin{aligned} & \int \sec(e+fx)\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}dx \\ &= \int \sqrt{a(\sec(e+fx)+1)}\sqrt{-c(\sec(e+fx)-1)}\sec(e+fx)dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(1/2)*(c-c*sec(f*x+e))**(1/2),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*sqrt(-c*(sec(e + f*x) - 1))*sec(e + f*x), x)`

3.109.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.34

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{2\sqrt{-a}\sqrt{c}}{f \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right) \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)*(c-c*sec(f*x+e))^(1/2),x, algorith="maxima")`

output `2*sqrt(-a)*sqrt(c)/(f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1))`

3.109.8 Giac [F]

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)} dx$$

$$= \int \sqrt{a \sec(fx + e) + a} \sqrt{-c \sec(fx + e) + c} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)*(c-c*sec(f*x+e))^(1/2),x, algorith="giac")`

output `sage0*x`

3.109.9 Mupad [B] (verification not implemented)

Time = 13.51 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.15

$$\int \sec(e + fx) \sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)} dx = \frac{\sqrt{c - \frac{c}{\cos(e+fx)}} \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}}}{f \sin(e + fx)}$$

input `int(((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x),x)`

output `((c - c/cos(e + f*x))^(1/2)*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2))/(f*sin(e + f*x))`

$$3.110 \quad \int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx$$

3.110.1 Optimal result	816
3.110.2 Mathematica [A] (verified)	816
3.110.3 Rubi [A] (verified)	817
3.110.4 Maple [B] (verified)	818
3.110.5 Fricas [F]	818
3.110.6 Sympy [F]	819
3.110.7 Maxima [A] (verification not implemented)	819
3.110.8 Giac [F]	819
3.110.9 Mupad [F(-1)]	820

3.110.1 Optimal result

Integrand size = 36, antiderivative size = 51

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx = \frac{a \log(1 - \sec(e+fx)) \tan(e+fx)}{f \sqrt{a+a\sec(e+fx)} \sqrt{c-c\sec(e+fx)}}$$

output `a*ln(1-sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.110.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.06

$$\begin{aligned} & \int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx \\ &= \frac{\log(1 - \sec(e+fx))\sqrt{a(1 + \sec(e+fx))} \tan\left(\frac{1}{2}(e+fx)\right)}{f \sqrt{c-c\sec(e+fx)}} \end{aligned}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/Sqrt[c - c*Sec[e + f*x]],x]`

output `(Log[1 - Sec[e + f*x]]*Sqrt[a*(1 + Sec[e + f*x]])*Tan[(e + f*x)/2])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.110. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx$

3.110.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a\sec(e+fx)+a}}{\sqrt{c-c\sec(e+fx)}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx$$

↓ 4440

$$\frac{a \tan(e+fx) \log(1-\sec(e+fx))}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/Sqrt[c - c*Sec[e + f*x]],x]`

output `(a*Log[1 - Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.110.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.110.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. $2(47) = 94$.

Time = 3.27 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.92

method	result
default	$\frac{\sqrt{a(\sec(fx+e)+1)}(\ln(-\cot(fx+e)+\csc(fx+e)-1)+\ln(-\cot(fx+e)+\csc(fx+e)+1)-2\ln(-\cot(fx+e)+\csc(fx+e)))(\cot(fx+e)-\csc(fx+e))}{f\sqrt{-c(\sec(fx+e)-1)}}$
risch	$-\frac{2i\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)\ln(e^{i(fx+e)}-1)}{(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}} + \frac{i\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)\ln(1+e^{2i(fx+e)})}{(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `1/f*(a*(sec(f*x+e)+1))^(1/2)*(ln(-cot(f*x+e)+csc(f*x+e)-1)+ln(-cot(f*x+e)+csc(f*x+e)+1)-2*ln(-cot(f*x+e)+csc(f*x+e)))/(-c*(sec(f*x+e)-1))^(1/2)*(cot(f*x+e)-csc(f*x+e))`

3.110.5 Fracas [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)*sec(f*x + e)/(c*sec(f*x + e) - c), x)`

3.110.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\sqrt{a(\sec(e+fx)+1)}\sec(e+fx)}{\sqrt{-c(\sec(e+fx)-1)}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(1/2)/(c-c*sec(f*x+e))**(1/2),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*sec(e + f*x)/sqrt(-c*(sec(e + f*x) - 1)), x)`

3.110.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.80

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx$$

$$= -\frac{\sqrt{-a}\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)}{\sqrt{c}} + \frac{\sqrt{-a}\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}-1\right)}{\sqrt{c}} - \frac{2\sqrt{-a}\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `-(sqrt(-a)*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/sqrt(c) + sqrt(-a)*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/sqrt(c) - 2*sqrt(-a)*log(sin(f*x + e)/(cos(f*x + e) + 1))/sqrt(c))/f`

3.110.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `sage0*x`

3.110. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx$

3.110.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\sqrt{a+\frac{a}{\cos(e+fx)}}}{\cos(e+fx)\sqrt{c-\frac{c}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

$$3.111 \quad \int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx$$

3.111.1 Optimal result	821
3.111.2 Mathematica [A] (verified)	821
3.111.3 Rubi [A] (verified)	822
3.111.4 Maple [A] (verified)	823
3.111.5 Fricas [B] (verification not implemented)	823
3.111.6 Sympy [F]	824
3.111.7 Maxima [B] (verification not implemented)	824
3.111.8 Giac [F]	825
3.111.9 Mupad [B] (verification not implemented)	825

3.111.1 Optimal result

Integrand size = 36, antiderivative size = 42

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx = -\frac{\sqrt{a+a\sec(e+fx)}\tan(e+fx)}{2f(c-c\sec(e+fx))^{3/2}}$$

output `-1/2*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(3/2)`

3.111.2 Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx = \frac{\cot(e+fx)\sqrt{a(1+\sec(e+fx))}}{cf\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c - c*Sec[e + f*x])^(3/2),x]`

output `(Cot[e + f*x]*Sqrt[a*(1 + Sec[e + f*x])])/(c*f*Sqrt[c - c*Sec[e + f*x]])`

3.111.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a\sec(e+fx)+a}}{(c-c\sec(e+fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx$$

↓ 4438

$$-\frac{\tan(e+fx)\sqrt{a\sec(e+fx)+a}}{2f(c-c\sec(e+fx))^{3/2}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c - c*Sec[e + f*x])^(3/2),x]`

output `-1/2*(Sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2))`

3.111.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.111.4 Maple [A] (verified)

Time = 3.22 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

method	result	size
default	$\frac{\sqrt{a(\sec(fx+e)+1)} \tan(fx+e)}{2fc(\sec(fx+e)-1)\sqrt{-c(\sec(fx+e)-1)}}$	50
risch	$\frac{2i\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} e^{i(fx+e)}}{c(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$	105

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/2/f*(a*(sec(f*x+e)+1))^(1/2)/c/(sec(f*x+e)-1)/(-c*(sec(f*x+e)-1))^(1/2)*
tan(f*x+e)
```

3.111.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. 2(36) = 72.

Time = 0.27 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.88

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx = \frac{\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}} \cos(fx+e)}{(c^2f\cos(fx+e)-c^2f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="fracas")
```

```
output sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x
+ e))*cos(f*x + e)/((c^2*f*cos(f*x + e) - c^2*f)*sin(f*x + e))
```


3.111.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\sqrt{a(\sec(e+fx)+1)}\sec(e+fx)}{(-c(\sec(e+fx)-1))^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(1/2)/(c-c*sec(f*x+e))**(3/2),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*sec(e + f*x)/(-c*(sec(e + f*x) - 1))**(3/2), x)`

3.111.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 514 vs. $2(36) = 72$.

Time = 0.38 (sec) , antiderivative size = 514, normalized size of antiderivative = 12.24

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx =$$

$$\frac{1}{(c^2 \cos(4fx + 4e)^2 + 4c^2 \cos(3fx + 3e)^2 + 4c^2 \cos(2fx + 2e)^2 + 4c^2 \cos(fx + e)^2 + c^2 \sin(4fx + 4e)^2 + \dots}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(3/2),x, algo rithm="maxima")`

output `-2*((sin(3*f*x + 3*e) + sin(f*x + e))*cos(4*f*x + 4*e) - (cos(3*f*x + 3*e) + cos(f*x + e))*sin(4*f*x + 4*e) + (2*cos(2*f*x + 2*e) + 1)*sin(3*f*x + 3*e) - 2*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) - 2*cos(f*x + e)*sin(2*f*x + 2*e) + 2*cos(2*f*x + 2*e)*sin(f*x + e) + sin(f*x + e))*sqrt(a)*sqrt(c)/((c^2*cos(4*f*x + 4*e)^2 + 4*c^2*cos(3*f*x + 3*e)^2 + 4*c^2*cos(2*f*x + 2*e)^2 + 4*c^2*cos(f*x + e)^2 + c^2*sin(4*f*x + 4*e)^2 + 4*c^2*sin(3*f*x + 3*e)^2 + 4*c^2*sin(2*f*x + 2*e)^2 - 8*c^2*sin(2*f*x + 2*e)*sin(f*x + e) + 4*c^2*sin(f*x + e)^2 - 4*c^2*cos(f*x + e) + c^2 - 2*(2*c^2*cos(3*f*x + 3*e) - 2*c^2*cos(2*f*x + 2*e) + 2*c^2*cos(f*x + e) - c^2)*cos(4*f*x + 4*e) - 4*(2*c^2*cos(2*f*x + 2*e) - 2*c^2*cos(f*x + e) + c^2)*cos(3*f*x + 3*e) - 4*(2*c^2*cos(f*x + e) - c^2)*cos(2*f*x + 2*e) - 4*(c^2*sin(3*f*x + 3*e) - c^2*sin(2*f*x + 2*e) + c^2*sin(f*x + e))*sin(4*f*x + 4*e) - 8*(c^2*sin(2*f*x + 2*e) - c^2*sin(f*x + e))*sin(3*f*x + 3*e))*f)`

3.111. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx$

3.111.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{(-c\sec(fx+e)+c)^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `sage0*x`

3.111.9 Mupad [B] (verification not implemented)

Time = 14.62 (sec) , antiderivative size = 118, normalized size of antiderivative = 2.81

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{3/2}} dx = \frac{2\sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}}\sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}}(\sin(e+fx)-2\sin(2e+2fx)+\sin(3e+3fx))}{c^2f(4\cos(e+fx)+4\cos(2e+2fx)-4\cos(3e+3fx)+\cos(4e+4fx)-5)}$$

input `int((a+a/cos(e+f*x))^(1/2)/(cos(e+f*x)*(c-c/cos(e+f*x))^(3/2)),x)`

output `-(2*((a*(cos(e+f*x)+1))/cos(e+f*x))^(1/2)*((c*(cos(e+f*x)-1))/cos(e+f*x))^(1/2)*(sin(e+f*x)-2*sin(2*e+2*f*x)+sin(3*e+3*f*x)))/(c^2*f*(4*cos(e+f*x)+4*cos(2*e+2*f*x)-4*cos(3*e+3*f*x)+cos(4*e+4*f*x)-5))`

$$3.112 \quad \int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx$$

3.112.1 Optimal result	826
3.112.2 Mathematica [A] (verified)	826
3.112.3 Rubi [A] (verified)	827
3.112.4 Maple [A] (verified)	828
3.112.5 Fricas [B] (verification not implemented)	828
3.112.6 Sympy [F]	829
3.112.7 Maxima [B] (verification not implemented)	829
3.112.8 Giac [F]	830
3.112.9 Mupad [B] (verification not implemented)	831

3.112.1 Optimal result

Integrand size = 36, antiderivative size = 43

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx = -\frac{a \tan(e+fx)}{2f\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}}$$

output `-1/2*a*tan(f*x+e)/f/(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2)`

3.112.2 Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.30

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx = \frac{a\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{2c^3f(-1+\sec(e+fx))^3\sqrt{a(1+\sec(e+fx))}}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c - c*Sec[e + f*x])^(5/2),x]`

output `(a*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/((2*c^3*f*(-1 + Sec[e + f*x])^3*Sqrt[a*(1 + Sec[e + f*x])])`

$$3.112. \quad \int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx$$

3.112.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a\sec(e+fx)+a}}{(c-c\sec(e+fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx$$

↓ 4441

$$\frac{a \tan(e+fx)}{2f\sqrt{a\sec(e+fx)+a}(c-c\sec(e+fx))^{5/2}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/2*(a*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(5/2))`

3.112.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.112.4 Maple [A] (verified)

Time = 3.37 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.56

method	result	size
default	$-\frac{\sqrt{a(\sec(fx+e)+1)}(3\tan(fx+e)-\sec(fx+e)\tan(fx+e))}{8f(\sec(fx+e)-1)^2\sqrt{-c(\sec(fx+e)-1)}c^2}$	67
risch	$\frac{2i\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{3i(fx+e)}-e^{2i(fx+e)}+e^{i(fx+e)})}{c^2(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)^3\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}$	126

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(5/2),x,method=_RET
URNVERBOSE)
```

```
output -1/8/f*(a*(sec(f*x+e)+1))^(1/2)/(sec(f*x+e)-1)^2/(-c*(sec(f*x+e)-1))^(1/2)
/c^2*(3*tan(f*x+e)-sec(f*x+e)*tan(f*x+e))
```

3.112.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. 2(37) = 74.

Time = 0.27 (sec) , antiderivative size = 106, normalized size of antiderivative = 2.47

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx = \frac{(2\cos(fx+e)^2 - \cos(fx+e))\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{2(c^3f\cos(fx+e)^2 - 2c^3f\cos(fx+e) + c^3f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(5/2),x, algo
rithm="fricas")
```

```
output 1/2*(2*cos(f*x + e)^2 - cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x +
e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((c^3*f*cos(f*x + e)^2 - 2*c^3
*f*cos(f*x + e) + c^3*f)*sin(f*x + e))
```

3.112.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\sqrt{a(\sec(e+fx)+1)}\sec(e+fx)}{(-c(\sec(e+fx)-1))^{5/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(1/2)/(c-c*sec(f*x+e))**(5/2),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*sec(e + f*x)/(-c*(sec(e + f*x) - 1))**(5/2), x)`

3.112.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 758 vs. $2(37) = 74$.

Time = 0.46 (sec) , antiderivative size = 758, normalized size of antiderivative = 17.63

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(5/2),x, algorith="maxima")`

output

```

2*((sin(4*f*x + 4*e) + 2*sin(2*f*x + 2*e))*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (sin(4*f*x + 4*e) + 2*sin(2*f*x + 2*e))*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - cos(2*f*x + 2*e)*sin(4*f*x + 4*e) + cos(4*f*x + 4*e)*sin(2*f*x + 2*e) - (cos(4*f*x + 4*e) + 2*cos(2*f*x + 2*e) + 1)*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (cos(4*f*x + 4*e) + 2*cos(2*f*x + 2*e) + 1)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + sin(2*f*x + 2*e))*sqrt(a)*sqrt(c)/((c^3*cos(4*f*x + 4*e))^2 + 36*c^3*cos(2*f*x + 2*e)^2 + 16*c^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 16*c^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + c^3*sin(4*f*x + 4*e)^2 + 12*c^3*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 36*c^3*sin(2*f*x + 2*e)^2 + 16*c^3*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 16*c^3*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 12*c^3*cos(2*f*x + 2*e) + c^3 + 2*(6*c^3*cos(2*f*x + 2*e) + c^3)*cos(4*f*x + 4*e) - 8*(c^3*cos(4*f*x + 4*e) + 6*c^3*cos(2*f*x + 2*e) - 4*c^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + c^3)*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 8*(c^3*cos(4*f*x + 4*e) + 6*c^3*cos(2*f*x + 2*e) + c^3)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 8*(c^3*sin(4*f*x + 4*e) + 6*c^3*sin(2*f*x + 2*e) - 4*c^3*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 8*(c^3*sin(4*f*x + 4*e) + 6*c^3*sin(2*f*...

```

3.112.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{(-c\sec(fx+e)+c)^{5/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `sage0*x`

3.112.9 Mupad [B] (verification not implemented)

Time = 18.11 (sec) , antiderivative size = 203, normalized size of antiderivative = 4.72

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{(c-c\sec(e+fx))^{5/2}} dx = \frac{\sqrt{c-\frac{c}{\cos(e+fx)}} \left(\frac{e^{e3i+fx3i} \sqrt{a+\frac{a}{\cos(e+fx)}} 4i}{c^3 f} + \frac{e^{e3i+fx3i} \cos(2e+2fx) \sqrt{a+\frac{a}{\cos(e+fx)}}}{c^3 f} \right)}{e^{e3i+fx3i} \sin(e+fx) 10i - e^{e3i+fx3i} \sin(2e+2fx) 8i}$$

```
input int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x
)
```

```
output ((c - c/cos(e + f*x))^(1/2)*((exp(e*3i + f*x*3i)*(a + a/cos(e + f*x))^(1/2)
)*4i)/(c^3*f) + (exp(e*3i + f*x*3i)*cos(2*e + 2*f*x)*(a + a/cos(e + f*x))^(
1/2)*4i)/(c^3*f) - (cos(e + f*x)*exp(e*3i + f*x*3i)*(a + a/cos(e + f*x))^(
1/2)*4i)/(c^3*f))/(exp(e*3i + f*x*3i)*sin(e + f*x)*10i - exp(e*3i + f*x*
3i)*sin(2*e + 2*f*x)*8i + exp(e*3i + f*x*3i)*sin(3*e + 3*f*x)*2i)
```


3.113 $\int \sec(e+fx)(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{7/2} dx$

3.113.1 Optimal result	832
3.113.2 Mathematica [A] (verified)	832
3.113.3 Rubi [A] (verified)	833
3.113.4 Maple [A] (verified)	834
3.113.5 Fricas [A] (verification not implemented)	835
3.113.6 Sympy [F(-1)]	835
3.113.7 Maxima [B] (verification not implemented)	836
3.113.8 Giac [F]	836
3.113.9 Mupad [B] (verification not implemented)	837

3.113.1 Optimal result

Integrand size = 36, antiderivative size = 89

$$\int \sec(e+fx)(a + a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{7/2} dx = \frac{a^2(c-c \sec(e+fx))^{7/2} \tan(e+fx)}{10f \sqrt{a+a \sec(e+fx)}} + \frac{a \sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{7/2} \tan(e+fx)}{5f}$$

```
output 1/10*a^2*(c-c*sec(f*x+e))^(7/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)+1/5*a*(c-c*sec(f*x+e))^(7/2)*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.113.2 Mathematica [A] (verified)

Time = 2.93 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.94

$$\int \sec(e+fx)(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{7/2} dx = \frac{a^2 c^4 \sec(e+fx) (-10 + 10 \sec(e+fx) - 5 \sec^3(e+fx) + 2 \sec^4(e+fx)) \tan(e+fx)}{10f \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(7/2),x]`

output `(a^2*c^4*Sec[e + f*x]*(-10 + 10*Sec[e + f*x] - 5*Sec[e + f*x]^3 + 2*Sec[e + f*x]^4)*Tan[e + f*x])/(10*f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.113.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^{3/2}(c - c \sec(e + fx))^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{7/2} dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{2}{5}a \int \sec(e + fx) \sqrt{\sec(e + fx)a + a}(c - c \sec(e + fx))^{7/2} dx + \\
 & \quad \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a}(c - c \sec(e + fx))^{7/2}}{5f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{5}a \int \csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{\csc\left(e + fx + \frac{\pi}{2}\right)a + a}(c - c \csc\left(e + fx + \frac{\pi}{2}\right))^{7/2} dx + \\
 & \quad \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a}(c - c \sec(e + fx))^{7/2}}{5f} \\
 & \quad \downarrow \text{4441} \\
 & \frac{a^2 \tan(e + fx)(c - c \sec(e + fx))^{7/2}}{10f \sqrt{a \sec(e + fx) + a}} + \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a}(c - c \sec(e + fx))^{7/2}}{5f}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(7/2),x]`

3.113. $\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{7/2} dx$

```
output (a^2*(c - c*Sec[e + f*x])^(7/2)*Tan[e + f*x])/(10*f*Sqrt[a + a*Sec[e + f*x
]]) + (a*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(7/2)*Tan[e + f*x])
/(5*f)
```

3.113.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sq
rt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f
*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /
; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
&& NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.113.4 Maple [A] (verified)

Time = 3.60 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.20

method	result
default	$-\frac{a(13 \cos(fx+e)^3 - 16 \cos(fx+e)^2 + 9 \cos(fx+e) - 2)(\sec(fx+e) - 1)^3 \sqrt{a(\sec(fx+e) + 1)} \sqrt{-c(\sec(fx+e) - 1)} c^3 (\cos(fx+e) + 1)^2}{10f(\cos(fx+e) - 1)^3}$
risch	$\frac{2ia c^3 \sqrt{\frac{a(e^{i(fx+e)} + 1)^2}{1 + e^{2i(fx+e)}}} \sqrt{\frac{c(e^{i(fx+e)} - 1)^2}{1 + e^{2i(fx+e)}}}}{5(1 + e^{2i(fx+e)})^4 (e^{i(fx+e)} + 1)(e^{i(fx+e)} - 1)f} (5e^{9i(fx+e)} - 10e^{8i(fx+e)} + 20e^{7i(fx+e)} - 10e^{6i(fx+e)} + 14e^{5i(fx+e)} - 10e^{4i(fx+e)} + 20e^{3i(fx+e)} - 10e^{2i(fx+e)} + 5e^{i(fx+e)} - 5)$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(7/2), x, method=_RET
URNVERBOSE)
```

output
$$-1/10/f*a*(13*\cos(f*x+e)^3-16*\cos(f*x+e)^2+9*\cos(f*x+e)-2)*(\sec(f*x+e)-1)^3*(a*(\sec(f*x+e)+1))^{(1/2)}*(-c*(\sec(f*x+e)-1))^{(1/2)}*c^3*(\cos(f*x+e)+1)^2/(\cos(f*x+e)-1)^3*\sec(f*x+e)*\csc(f*x+e)$$

3.113.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.26

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{7/2} dx = \frac{(10ac^3 \cos(fx + e)^4 - 10ac^3 \cos(fx + e)^3 + 5ac^3 \cos(fx + e) - 2ac^3) \sqrt{\frac{a \cos(fx + e)}{\cos(fx + e)}}}{10f \cos(fx + e)^4 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(7/2),x, algorithm="fricas")`

output
$$1/10*(10*a*c^3*\cos(f*x + e)^4 - 10*a*c^3*\cos(f*x + e)^3 + 5*a*c^3*\cos(f*x + e) - 2*a*c^3)*\sqrt{(a*\cos(f*x + e) + a)/\cos(f*x + e)}*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^4*\sin(f*x + e))$$

3.113.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)*(c-c*sec(f*x+e))**(7/2),x)`

output `Timed out`

3.113.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1680 vs. $2(77) = 154$.

Time = 0.41 (sec) , antiderivative size = 1680, normalized size of antiderivative = 18.88

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{7/2} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(7/2),x, algo
rithm="maxima")
```

```
output 2/5*(100*a*c^3*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) - 25*a*c^3*cos(2*f*x + 2*
e)*sin(f*x + e) - 5*a*c^3*sin(f*x + e) - (5*a*c^3*sin(9*f*x + 9*e) - 10*a*
c^3*sin(8*f*x + 8*e) + 20*a*c^3*sin(7*f*x + 7*e) - 10*a*c^3*sin(6*f*x + 6*
e) + 14*a*c^3*sin(5*f*x + 5*e) - 10*a*c^3*sin(4*f*x + 4*e) + 20*a*c^3*sin(
3*f*x + 3*e) - 10*a*c^3*sin(2*f*x + 2*e) + 5*a*c^3*sin(f*x + e))*cos(10*f*
x + 10*e) + 25*(a*c^3*sin(8*f*x + 8*e) + 2*a*c^3*sin(6*f*x + 6*e) + 2*a*c^
3*sin(4*f*x + 4*e) + a*c^3*sin(2*f*x + 2*e))*cos(9*f*x + 9*e) - 5*(20*a*c^
3*sin(7*f*x + 7*e) + 10*a*c^3*sin(6*f*x + 6*e) + 14*a*c^3*sin(5*f*x + 5*e)
+ 10*a*c^3*sin(4*f*x + 4*e) + 20*a*c^3*sin(3*f*x + 3*e) + 5*a*c^3*sin(f*x
+ e))*cos(8*f*x + 8*e) + 100*(2*a*c^3*sin(6*f*x + 6*e) + 2*a*c^3*sin(4*f*
x + 4*e) + a*c^3*sin(2*f*x + 2*e))*cos(7*f*x + 7*e) - 10*(14*a*c^3*sin(5*f
*x + 5*e) + 20*a*c^3*sin(3*f*x + 3*e) - 5*a*c^3*sin(2*f*x + 2*e) + 5*a*c^3
*sin(f*x + e))*cos(6*f*x + 6*e) + 70*(2*a*c^3*sin(4*f*x + 4*e) + a*c^3*sin
(2*f*x + 2*e))*cos(5*f*x + 5*e) - 50*(4*a*c^3*sin(3*f*x + 3*e) - a*c^3*sin
(2*f*x + 2*e) + a*c^3*sin(f*x + e))*cos(4*f*x + 4*e) + (5*a*c^3*cos(9*f*x
+ 9*e) - 10*a*c^3*cos(8*f*x + 8*e) + 20*a*c^3*cos(7*f*x + 7*e) - 10*a*c^3*
cos(6*f*x + 6*e) + 14*a*c^3*cos(5*f*x + 5*e) - 10*a*c^3*cos(4*f*x + 4*e) +
20*a*c^3*cos(3*f*x + 3*e) - 10*a*c^3*cos(2*f*x + 2*e) + 5*a*c^3*cos(f*x +
e))*sin(10*f*x + 10*e) - 5*(5*a*c^3*cos(8*f*x + 8*e) + 10*a*c^3*cos(6*f*x
+ 6*e) + 10*a*c^3*cos(4*f*x + 4*e) + 5*a*c^3*cos(2*f*x + 2*e) + a*c^3)...
```

3.113.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{7/2} dx = \int (a \sec(fx + e) + a)^{\frac{3}{2}}(-c \sec(fx + e) + c)^{\frac{7}{2}} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(7/2),x, algo
rithm="giac")`

output `sage0*x`

3.113.9 Mupad [B] (verification not implemented)

Time = 17.66 (sec) , antiderivative size = 294, normalized size of antiderivative = 3.30

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{7/2} dx = \frac{\sqrt{c - \frac{c}{\cos(e+fx)}} \left(\frac{a c^3 e^{e 5i + f x 5i} \sqrt{a + \frac{a}{\cos(e+fx)}} 28i}{5f} - \frac{a c^3 \cos(e+fx) e^{e 5i + f x 5i} \sqrt{a + \frac{a}{\cos(e+fx)}} 8i}{f} + \frac{a c^3}{e^{e 5i + f x 5i} \sin(e + fx) 4i + e^e} \right)}{e^{e 5i + f x 5i} \sin(e + fx) 4i + e^e}$$

input `int(((a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(7/2))/cos(e + f*x),x
)`

output `((c - c/cos(e + f*x))^(1/2)*((a*c^3*exp(e*5i + f*x*5i)*(a + a/cos(e + f*x))
)^(1/2)*28i)/(5*f) - (a*c^3*cos(e + f*x)*exp(e*5i + f*x*5i)*(a + a/cos(e +
f*x))^(1/2)*8i)/f + (a*c^3*exp(e*5i + f*x*5i)*cos(2*e + 2*f*x)*(a + a/cos
(e + f*x))^(1/2)*16i)/f - (a*c^3*exp(e*5i + f*x*5i)*cos(3*e + 3*f*x)*(a +
a/cos(e + f*x))^(1/2)*8i)/f + (a*c^3*exp(e*5i + f*x*5i)*cos(4*e + 4*f*x)*(
a + a/cos(e + f*x))^(1/2)*4i)/f)/(exp(e*5i + f*x*5i)*sin(e + f*x)*4i + ex
p(e*5i + f*x*5i)*sin(3*e + 3*f*x)*6i + exp(e*5i + f*x*5i)*sin(5*e + 5*f*x)
*2i)`

3.114 $\int \sec(e+fx)(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{5/2} dx$

3.114.1 Optimal result	838
3.114.2 Mathematica [A] (verified)	838
3.114.3 Rubi [A] (verified)	839
3.114.4 Maple [A] (verified)	840
3.114.5 Fricas [A] (verification not implemented)	841
3.114.6 Sympy [F(-1)]	841
3.114.7 Maxima [B] (verification not implemented)	842
3.114.8 Giac [F]	842
3.114.9 Mupad [B] (verification not implemented)	843

3.114.1 Optimal result

Integrand size = 36, antiderivative size = 89

$$\int \sec(e+fx)(a + a \sec(e+fx))^{3/2}(c - c \sec(e+fx))^{5/2} dx = \frac{a^2(c - c \sec(e+fx))^{5/2} \tan(e+fx)}{6f \sqrt{a + a \sec(e+fx)}} + \frac{a \sqrt{a + a \sec(e+fx)}(c - c \sec(e+fx))^{5/2} \tan(e+fx)}{4f}$$

```
output 1/6*a^2*(c-c*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)+1/4*a*(c-c*sec(f*x+e))^(5/2)*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.114.2 Mathematica [A] (verified)

Time = 1.62 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.98

$$\int \sec(e+fx)(a + a \sec(e+fx))^{3/2}(c - c \sec(e+fx))^{5/2} dx = \frac{ac^3(5 \cos(e+fx) - 3 \cos(2(e+fx)) + 3 \cos(3(e+fx))) \sec^4(e+fx) \sqrt{a(1 + \sec(e+fx))} \tan(\frac{1}{2}(e+fx))}{12f \sqrt{c - c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/12*(a*c^3*(5*Cos[e + f*x] - 3*Cos[2*(e + f*x)] + 3*Cos[3*(e + f*x)])*Sec[e + f*x]^4*Sqrt[a*(1 + Sec[e + f*x])]*Tan[(e + f*x)/2])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.114.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^{3/2}(c - c \sec(e + fx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{1}{2}a \int \sec(e + fx) \sqrt{\sec(e + fx)a + a(c - c \sec(e + fx))}^{5/2} dx + \\
 & \quad \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))}^{5/2}}{4f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}a \int \csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{\csc\left(e + fx + \frac{\pi}{2}\right) a + a\left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)}^{5/2} dx + \\
 & \quad \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))}^{5/2}}{4f} \\
 & \quad \downarrow \text{4441} \\
 & \frac{a^2 \tan(e + fx)(c - c \sec(e + fx))^{5/2}}{6f \sqrt{a \sec(e + fx) + a}} + \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))}^{5/2}}{4f}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(5/2),x]`

output $(a^2(c - c\sec[e + fx])^{5/2}\tan[e + fx]) / (6f\sqrt{a + a\sec[e + fx]}) + (a\sqrt{a + a\sec[e + fx]})(c - c\sec[e + fx])^{5/2}\tan[e + fx] / (4f)$

3.114.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*Sqrt[csc[(e_) + (f_)*(x_)]*(d_) + (c_)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4443 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.114.4 Maple [A] (verified)

Time = 3.54 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.08

method	result
default	$\frac{a(\sec(fx+e)-1)^2 \sqrt{a(\sec(fx+e)+1)} \sqrt{-c(\sec(fx+e)-1)} c^2 (\cos(fx+e)+1)^2 (11 \cot(fx+e) - 10 \csc(fx+e) + 3 \sec(fx+e) \csc(fx+e))}{12f(\cos(fx+e)-1)^2}$
risch	$\frac{2ia c^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} (3e^{7i(fx+e)} - 3e^{6i(fx+e)} + 5e^{5i(fx+e)} + 5e^{3i(fx+e)} - 3e^{2i(fx+e)} + 3e^{i(fx+e)})}{3(1+e^{2i(fx+e)})^3 (e^{i(fx+e)}+1)(e^{i(fx+e)}-1)} f$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(5/2),x,method=_RET URNVERBOSE)`

3.114. $\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{5/2} dx$

output $1/12/f*a*(\sec(f*x+e)-1)^2*(a*(\sec(f*x+e)+1))^{(1/2)}*(-c*(\sec(f*x+e)-1))^{(1/2)}*c^2*(\cos(f*x+e)+1)^2/(\cos(f*x+e)-1)^2*(11*\cot(f*x+e)-10*\csc(f*x+e)+3*\sec(f*x+e)*\csc(f*x+e))$

3.114.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.26

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{5/2} dx = \frac{(12 ac^2 \cos(fx + e))^3 - 6 ac^2 \cos(fx + e)^2 - 4 ac^2 \cos(fx + e) + 3 ac^2}{12 f \cos(fx + e)^3 \sin(fx + e)} \sqrt{\frac{a \cos(fx + e) + a}{\cos(fx + e)}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(5/2),x, algorithm="fricas")`

output $1/12*(12*a*c^2*\cos(f*x + e)^3 - 6*a*c^2*\cos(f*x + e)^2 - 4*a*c^2*\cos(f*x + e) + 3*a*c^2)*\sqrt{(a*\cos(f*x + e) + a)/\cos(f*x + e)}*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^3*\sin(f*x + e))$

3.114.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{5/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)*(c-c*sec(f*x+e))**(5/2),x)`

output **Timed out**

3.114.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1105 vs. $2(77) = 154$.

Time = 0.39 (sec) , antiderivative size = 1105, normalized size of antiderivative = 12.42

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{5/2} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(5/2),x, algo
rithm="maxima")
```

```
output 2/3*(20*a*c^2*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) - 12*a*c^2*cos(2*f*x + 2*e)
)*sin(f*x + e) - 3*a*c^2*sin(f*x + e) - (3*a*c^2*sin(7*f*x + 7*e) - 3*a*c^
2*sin(6*f*x + 6*e) + 5*a*c^2*sin(5*f*x + 5*e) + 5*a*c^2*sin(3*f*x + 3*e) -
3*a*c^2*sin(2*f*x + 2*e) + 3*a*c^2*sin(f*x + e))*cos(8*f*x + 8*e) + 6*(2*
a*c^2*sin(6*f*x + 6*e) + 3*a*c^2*sin(4*f*x + 4*e) + 2*a*c^2*sin(2*f*x + 2*
e))*cos(7*f*x + 7*e) - 2*(10*a*c^2*sin(5*f*x + 5*e) + 9*a*c^2*sin(4*f*x +
4*e) + 10*a*c^2*sin(3*f*x + 3*e) + 6*a*c^2*sin(f*x + e))*cos(6*f*x + 6*e)
+ 10*(3*a*c^2*sin(4*f*x + 4*e) + 2*a*c^2*sin(2*f*x + 2*e))*cos(5*f*x + 5*e)
- 6*(5*a*c^2*sin(3*f*x + 3*e) - 3*a*c^2*sin(2*f*x + 2*e) + 3*a*c^2*sin(f
*x + e))*cos(4*f*x + 4*e) + (3*a*c^2*cos(7*f*x + 7*e) - 3*a*c^2*cos(6*f*x
+ 6*e) + 5*a*c^2*cos(5*f*x + 5*e) + 5*a*c^2*cos(3*f*x + 3*e) - 3*a*c^2*cos
(2*f*x + 2*e) + 3*a*c^2*cos(f*x + e))*sin(8*f*x + 8*e) - 3*(4*a*c^2*cos(6*
f*x + 6*e) + 6*a*c^2*cos(4*f*x + 4*e) + 4*a*c^2*cos(2*f*x + 2*e) + a*c^2)*
sin(7*f*x + 7*e) + (20*a*c^2*cos(5*f*x + 5*e) + 18*a*c^2*cos(4*f*x + 4*e)
+ 20*a*c^2*cos(3*f*x + 3*e) + 12*a*c^2*cos(f*x + e) + 3*a*c^2)*sin(6*f*x +
6*e) - 5*(6*a*c^2*cos(4*f*x + 4*e) + 4*a*c^2*cos(2*f*x + 2*e) + a*c^2)*si
n(5*f*x + 5*e) + 6*(5*a*c^2*cos(3*f*x + 3*e) - 3*a*c^2*cos(2*f*x + 2*e) +
3*a*c^2*cos(f*x + e))*sin(4*f*x + 4*e) - 5*(4*a*c^2*cos(2*f*x + 2*e) + a*c
^2)*sin(3*f*x + 3*e) + 3*(4*a*c^2*cos(f*x + e) + a*c^2)*sin(2*f*x + 2*e))*
sqrt(a)*sqrt(c)/((2*(4*cos(6*f*x + 6*e) + 6*cos(4*f*x + 4*e) + 4*cos(2*...
```

3.114.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{5/2} dx = \int (a \sec(fx + e) + a)^{\frac{3}{2}}(-c \sec(fx + e) + c)^{\frac{5}{2}} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(5/2),x, algo
rithm="giac")`

output `sage0*x`

3.114.9 Mupad [B] (verification not implemented)

Time = 17.21 (sec) , antiderivative size = 195, normalized size of antiderivative = 2.19

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{5/2} dx = \frac{\sqrt{c - \frac{c}{\cos(e + fx)}} \left(\frac{a c^2 \cos(e + fx) e^{e 4i + f x 4i} \sqrt{a + \frac{a}{\cos(e + fx)}} 20i}{3 f} - \frac{a c^2 e^{e 4i + f x 4i} \cos(2e + 2fx) \sqrt{a + \frac{a}{\cos(e + fx)}}}{f} \right)}{e^{e 4i + f x 4i} \sin(2e + 2fx) 4i + e^{e 4i + f x 4i} \sin(4e + 4fx) 2i}$$

input `int(((a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x),x
)`

output `((c - c/cos(e + f*x))^(1/2)*((a*c^2*cos(e + f*x)*exp(e*4i + f*x*4i)*(a + a
/cos(e + f*x))^(1/2)*20i)/(3*f) - (a*c^2*exp(e*4i + f*x*4i)*cos(2*e + 2*f*
x)*(a + a/cos(e + f*x))^(1/2)*4i)/f + (a*c^2*exp(e*4i + f*x*4i)*cos(3*e +
3*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/f))/(exp(e*4i + f*x*4i)*sin(2*e + 2*
f*x)*4i + exp(e*4i + f*x*4i)*sin(4*e + 4*f*x)*2i)`

3.115 $\int \sec(e+fx)(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2} dx$

3.115.1 Optimal result	844
3.115.2 Mathematica [A] (verified)	844
3.115.3 Rubi [A] (verified)	845
3.115.4 Maple [A] (verified)	846
3.115.5 Fricas [A] (verification not implemented)	847
3.115.6 Sympy [F(-1)]	847
3.115.7 Maxima [B] (verification not implemented)	848
3.115.8 Giac [F]	848
3.115.9 Mupad [B] (verification not implemented)	849

3.115.1 Optimal result

Integrand size = 36, antiderivative size = 89

$$\int \sec(e+fx)(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2} dx =$$

$$\frac{c^2(a+a \sec(e+fx))^{3/2} \tan(e+fx)}{3f \sqrt{c-c \sec(e+fx)}} - \frac{c(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)} \tan(e+fx)}{3f}$$

```
output -1/3*c^2*(a+a*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-1/3*c*
(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.115.2 Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.72

$$\int \sec(e+fx)(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2} dx = \frac{a^2 c^2 \sec(e+fx) (-3 + \sec^2(e+fx)) \tan(e+fx)}{3f \sqrt{a(1 + \sec(e+fx))} \sqrt{c - c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(3/2),x]`

output `(a^2*c^2*Sec[e + f*x]*(-3 + Sec[e + f*x]^2)*Tan[e + f*x])/(3*f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.115.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^{3/2}(c - c \sec(e + fx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{2}{3}c \int \sec(e + fx)(\sec(e + fx)a + a)^{3/2} \sqrt{c - c \sec(e + fx)} dx - \\
 & \quad \frac{c \tan(e + fx)(a \sec(e + fx) + a)^{3/2} \sqrt{c - c \sec(e + fx)}}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{3}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right) a + a\right)^{3/2} \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx - \\
 & \quad \frac{c \tan(e + fx)(a \sec(e + fx) + a)^{3/2} \sqrt{c - c \sec(e + fx)}}{3f} \\
 & \quad \downarrow \text{4441} \\
 & \frac{c^2 \tan(e + fx)(a \sec(e + fx) + a)^{3/2}}{3f \sqrt{c - c \sec(e + fx)}} - \frac{c \tan(e + fx)(a \sec(e + fx) + a)^{3/2} \sqrt{c - c \sec(e + fx)}}{3f}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(3/2),x]`

3.115. $\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{3/2} dx$

```
output -1/3*(c^2*(a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]]) - (c*(a + a*Sec[e + f*x])^(3/2)*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(3*f)
```

3.115.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.115.4 Maple [A] (verified)

Time = 3.41 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.93

method	result	size
default	$\frac{-a(\sec(fx+e)-1)(2\cos(fx+e)-1)\sqrt{-c(\sec(fx+e)-1)}c\sqrt{a(\sec(fx+e)+1)}(\cos(fx+e)+1)^2\sec(fx+e)\csc(fx+e)}{3f(\cos(fx+e)-1)}$	83
risch	$\frac{2iac\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}(3e^{5i(fx+e)}+2e^{3i(fx+e)}+3e^{i(fx+e)})}{3(1+e^{2i(fx+e)})^2(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)f}}$	142

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

output $-1/3/f*a*(\sec(f*x+e)-1)*(2*\cos(f*x+e)-1)*(-c*(\sec(f*x+e)-1))^{(1/2)}*c*(a*(\sec(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)+1)^2/(\cos(f*x+e)-1)*\sec(f*x+e)*\csc(f*x+e)$

3.115.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.92

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{3/2} dx = \frac{(3ac \cos(fx + e)^2 - ac) \sqrt{\frac{a \cos(fx + e) + a}{\cos(fx + e)}} \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}}}{3f \cos(fx + e)^2 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(3/2),x, algo rithm="fricas")`

output $1/3*(3*a*c*\cos(f*x + e)^2 - a*c)*\sqrt{(a*\cos(f*x + e) + a)/\cos(f*x + e)}*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^2*\sin(f*x + e))$

3.115.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{3/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)*(c-c*sec(f*x+e))**(3/2),x)`

output `Timed out`

3.115.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 550 vs. $2(77) = 154$.

Time = 0.42 (sec) , antiderivative size = 550, normalized size of antiderivative = 6.18

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{3/2} dx = \frac{2(6ac \cos(3fx + 3e) \sin(2fx + 2e) + 9ac \cos(fx + e) \sin(2fx + 2e) - 9ac \cos($$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output `2/3*(6*a*c*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) + 9*a*c*cos(f*x + e)*sin(2*f*x + 2*e) - 9*a*c*cos(2*f*x + 2*e)*sin(f*x + e) - 3*a*c*sin(f*x + e) - (3*a*c*sin(5*f*x + 5*e) + 2*a*c*sin(3*f*x + 3*e) + 3*a*c*sin(f*x + e))*cos(6*f*x + 6*e) + 9*(a*c*sin(4*f*x + 4*e) + a*c*sin(2*f*x + 2*e))*cos(5*f*x + 5*e) - 3*(2*a*c*sin(3*f*x + 3*e) + 3*a*c*sin(f*x + e))*cos(4*f*x + 4*e) + (3*a*c*cos(5*f*x + 5*e) + 2*a*c*cos(3*f*x + 3*e) + 3*a*c*cos(f*x + e))*sin(6*f*x + 6*e) - 3*(3*a*c*cos(4*f*x + 4*e) + 3*a*c*cos(2*f*x + 2*e) + a*c)*sin(5*f*x + 5*e) + 3*(2*a*c*cos(3*f*x + 3*e) + 3*a*c*cos(f*x + e))*sin(4*f*x + 4*e) - 2*(3*a*c*cos(2*f*x + 2*e) + a*c)*sin(3*f*x + 3*e))*sqrt(a)*sqrt(c)/((2*(3*cos(4*f*x + 4*e) + 3*cos(2*f*x + 2*e) + 1)*cos(6*f*x + 6*e) + cos(6*f*x + 6*e)^2 + 6*(3*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + 9*cos(4*f*x + 4*e)^2 + 9*cos(2*f*x + 2*e)^2 + 6*(sin(4*f*x + 4*e) + sin(2*f*x + 2*e))*sin(6*f*x + 6*e) + sin(6*f*x + 6*e)^2 + 9*sin(4*f*x + 4*e)^2 + 18*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 9*sin(2*f*x + 2*e)^2 + 6*cos(2*f*x + 2*e) + 1)*f)`

3.115.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{3/2} dx = \int (a \sec(fx + e) + a)^{\frac{3}{2}}(-c \sec(fx + e) + c)^{\frac{3}{2}} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(3/2),x, algorithm="giac")`

output sage0*x

3.115.9 Mupad [B] (verification not implemented)

Time = 14.72 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.21

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2}(c - c \sec(e + fx))^{3/2} dx = \frac{2ac \sqrt{c - \frac{c}{\cos(e+fx)}} \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}} (2 \sin(e + fx) + 5 \sin(3e + 3fx) + 3 \sin(5e + 5fx))}{3f (\cos(2e + 2fx) - 2 \cos(4e + 4fx) - \cos(6e + 6fx) + 2)}$$

input `int(((a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(3/2))/cos(e + f*x),x)`

output `(2*a*c*(c - c/cos(e + f*x))^(1/2)*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2)*(2*sin(e + f*x) + 5*sin(3*e + 3*f*x) + 3*sin(5*e + 5*f*x)))/(3*f*(cos(2*e + 2*f*x) - 2*cos(4*e + 4*f*x) - cos(6*e + 6*f*x) + 2))`

3.116 $\int \sec(e+fx)(a+a \sec(e+fx))^{3/2} \sqrt{c - c \sec(e + fx)}$

3.116.1 Optimal result	850
3.116.2 Mathematica [A] (verified)	850
3.116.3 Rubi [A] (verified)	851
3.116.4 Maple [A] (verified)	852
3.116.5 Fricas [B] (verification not implemented)	852
3.116.6 Sympy [F]	853
3.116.7 Maxima [A] (verification not implemented)	853
3.116.8 Giac [F]	853
3.116.9 Mupad [B] (verification not implemented)	854

3.116.1 Optimal result

Integrand size = 36, antiderivative size = 43

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)} dx = \frac{c(a + a \sec(e + fx))^{3/2} \tan(e + fx)}{2f \sqrt{c - c \sec(e + fx)}}$$

output `-1/2*c*(a+a*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)`

3.116.2 Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.53

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)} dx = \frac{ac(1 + 2 \cos(e + fx)) \sec^2(e + fx) \sqrt{a(1 + \sec(e + fx))} \tan(\frac{1}{2}(e + fx))}{2f \sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*Sqrt[c - c*Sec[e + f*x]],x]`

output `-1/2*(a*c*(1 + 2*Cos[e + f*x])*Sec[e + f*x]^2*Sqrt[a*(1 + Sec[e + f*x])]*Tan[(e + f*x)/2])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.116.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^{3/2} \sqrt{c - c \sec(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{3/2} \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4441}$$

$$-\frac{c \tan(e + fx)(a \sec(e + fx) + a)^{3/2}}{2f \sqrt{c - c \sec(e + fx)}}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2)*Sqrt[c - c*Sec[e + f*x]],x]`

output `-1/2*(c*(a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.116.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.116.4 Maple [A] (verified)

Time = 3.80 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.26

method	result	size
default	$\frac{a(\cos(fx+e)+1)^2 \sqrt{a(\sec(fx+e)+1)} \sqrt{-c(\sec(fx+e)-1)} \sec(fx+e) \csc(fx+e)}{2f}$	54
risch	$\frac{2ia \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} (e^{3i(fx+e)}+e^{2i(fx+e)}+e^{i(fx+e)})}{(1+e^{2i(fx+e)})(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)} f$	135

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/2/f*a*(cos(f*x+e)+1)^2*(a*(sec(f*x+e)+1))^(1/2)*(-c*(sec(f*x+e)-1))^(1/2)*sec(f*x+e)*csc(f*x+e)
```

3.116.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(37) = 74.

Time = 0.26 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.77

$$\int \sec(e+fx)(a+a\sec(e+fx))^{3/2} \sqrt{c-c\sec(e+fx)} dx = \frac{(2a\cos(fx+e)+a) \sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{2f\cos(fx+e)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(1/2),x, algorithm="fracas")
```

```
output 1/2*(2*a*cos(f*x + e) + a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)*sin(f*x + e))
```

3.116.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)} dx = \int (a(\sec(e + fx) + 1))^{3/2} \sqrt{-c(\sec(e + fx) - 1)} \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)*(c-c*sec(f*x+e))**(1/2),x)`

output `Integral((a*(sec(e + f*x) + 1))**(3/2)*sqrt(-c*(sec(e + f*x) - 1))*sec(e + f*x), x)`

3.116.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.30

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)} dx =$$

$$-\frac{2\sqrt{-aa}\sqrt{c}}{f\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)^2\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `-2*sqrt(-a)*a*sqrt(c)/(f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^2*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^2)`

3.116.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)} dx = \int (a \sec(fx + e) + a)^{3/2} \sqrt{-c \sec(fx + e) + c \sec(fx + e)} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(1/2),x, algorith="giac")`

output `sage0*x`

3.116.9 Mupad [B] (verification not implemented)

Time = 14.09 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.77

$$\int \sec(e + fx)(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)} dx = \frac{a \sqrt{c - \frac{c}{\cos(e + fx)}} \sqrt{\frac{a(\cos(e + fx) + 1)}{\cos(e + fx)}} (\sin(e + fx) + \sin(2e + 2fx) + \sin(3e + 3fx))}{f \sin(2e + 2fx)^2}$$

input `int(((a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x),x)`

output `(a*(c - c/cos(e + f*x))^(1/2)*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2)*(sin(e + f*x) + sin(2*e + 2*f*x) + sin(3*e + 3*f*x)))/(f*sin(2*e + 2*f*x)^2)`

$$3.117 \quad \int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{\sqrt{c-c \sec(e+fx)}} dx$$

3.117.1 Optimal result	855
3.117.2 Mathematica [A] (verified)	855
3.117.3 Rubi [A] (verified)	856
3.117.4 Maple [A] (verified)	857
3.117.5 Fracas [F]	858
3.117.6 Sympy [F]	858
3.117.7 Maxima [B] (verification not implemented)	858
3.117.8 Giac [F]	859
3.117.9 Mupad [F(-1)]	859

3.117.1 Optimal result

Integrand size = 36, antiderivative size = 95

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{\sqrt{c-c \sec(e+fx)}} dx = \frac{2a^2 \log(1-\sec(e+fx)) \tan(e+fx)}{f \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} + \frac{a \sqrt{a+a \sec(e+fx)} \tan(e+fx)}{f \sqrt{c-c \sec(e+fx)}}$$

```
output 2*a^2*ln(1-sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)+a*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)
```

3.117.2 Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.65

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{\sqrt{c-c \sec(e+fx)}} dx = \frac{a^2(2 \log(1-\sec(e+fx)) + \sec(e+fx)) \tan(e+fx)}{f \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

```
input Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/Sqrt[c - c*Sec[e + f*x]],x]
```

```
output (a^2*(2*Log[1 - Sec[e + f*x]] + Sec[e + f*x])*Tan[e + f*x])/(f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])
```

3.117. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{\sqrt{c-c \sec(e+fx)}} dx$

3.117.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4443, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx$$

↓ 4443

$$2a \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{\sqrt{c-c\sec(e+fx)}} dx + \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}}$$

↓ 3042

$$2a \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}}$$

↓ 4440

$$\frac{2a^2 \tan(e+fx) \log(1-\sec(e+fx))}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} + \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/Sqrt[c - c*Sec[e + f*x]],x]`

output `(2*a^2*Log[1 - Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]) + (a*Sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.117.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.117.4 Maple [A] (verified)

Time = 3.83 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.35

method	result
default	$\frac{a\sqrt{a(\sec(fx+e)+1)}(4\ln(-\cot(fx+e)+\csc(fx+e))\sin(fx+e)-2\ln(-\cot(fx+e)+\csc(fx+e)-1)\sin(fx+e)-2\ln(-\cot(fx+e)+\csc(fx+e)+1))}{f(\cos(fx+e)+1)\sqrt{-c(\sec(fx+e)-1)}}$
risch	$-\frac{2ia\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(2\ln(e^{i(fx+e)}-1)e^{3i(fx+e)}-\ln(1+e^{2i(fx+e)})e^{3i(fx+e)}+2e^{i(fx+e)}\ln(e^{i(fx+e)}-1)-e^{i(fx+e)}\ln(1+e^{2i(fx+e)}))}{(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)}{1+e^{2i(fx+e)}}}}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `1/f*a*(a*(sec(f*x+e)+1))^(1/2)/(cos(f*x+e)+1)/(-c*(sec(f*x+e)-1))^(1/2)*(4*ln(-cot(f*x+e)+csc(f*x+e))*sin(f*x+e)-2*ln(-cot(f*x+e)+csc(f*x+e)-1)*sin(f*x+e)-2*ln(-cot(f*x+e)+csc(f*x+e)+1)*sin(f*x+e)+sin(f*x+e)+tan(f*x+e))`

$$3.117. \int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx$$

3.117.5 Fricas [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{(a\sec(fx+e)+a)^{3/2}\sec(fx+e)}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral(-(a*sec(f*x + e)^2 + a*sec(f*x + e))*sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(c*sec(f*x + e) - c), x)`

3.117.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{(a(\sec(e+fx)+1))^{3/2}\sec(e+fx)}{\sqrt{-c(\sec(e+fx)-1)}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(1/2),x)`

output `Integral((a*(sec(e + f*x) + 1))**(3/2)*sec(e + f*x)/sqrt(-c*(sec(e + f*x) - 1)), x)`

3.117.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. 2(87) = 174.

Time = 0.39 (sec) , antiderivative size = 275, normalized size of antiderivative = 2.89

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx =$$

$$\frac{2(a \cos(\frac{1}{2} \arctan(\sin(2fx+2e), \cos(2fx+2e))) \sin(2fx+2e) + (a \cos(2fx+2e))^2 + a \sin(2fx+2e))}{\sqrt{c-c\sec(e+fx)}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

3.117. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx$

output $-2*(a*\cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*\sin(2*f*x + 2*e) + (a*\cos(2*f*x + 2*e)^2 + a*\sin(2*f*x + 2*e)^2 + 2*a*\cos(2*f*x + 2*e) + a)*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e) + 1) - 2*(a*\cos(2*f*x + 2*e)^2 + a*\sin(2*f*x + 2*e)^2 + 2*a*\cos(2*f*x + 2*e) + a)*\arctan2(\sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))), \cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e)))) - 1) - (a*\cos(2*f*x + 2*e) + a)*\sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*\sqrt{a}*\sqrt{c}/((c*\cos(2*f*x + 2*e)^2 + c*\sin(2*f*x + 2*e)^2 + 2*c*\cos(2*f*x + 2*e) + c)*f)$

3.117.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{(a\sec(fx+e)+a)^{3/2}\sec(fx+e)}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `sage0*x`

3.117.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^{3/2}}{\cos(e+fx) \sqrt{c - \frac{c}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

3.118
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{3/2}} dx$$

3.118.1 Optimal result 860
 3.118.2 Mathematica [A] (verified) 860
 3.118.3 Rubi [A] (verified) 861
 3.118.4 Maple [B] (verified) 862
 3.118.5 Fricas [F] 863
 3.118.6 Sympy [F] 863
 3.118.7 Maxima [A] (verification not implemented) 864
 3.118.8 Giac [F] 864
 3.118.9 Mupad [F(-1)] 864

3.118.1 Optimal result

Integrand size = 36, antiderivative size = 99

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{3/2}} dx = -\frac{a\sqrt{a+a \sec(e+fx)} \tan(e+fx)}{f(c-c \sec(e+fx))^{3/2}} - \frac{a^2 \log(1-\sec(e+fx)) \tan(e+fx)}{cf\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}}$$

output `-a*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(3/2)-a^2*ln(1-sec(f*x+e))*tan(f*x+e)/c/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.118.2 Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.71

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{3/2}} dx = \frac{a^2 \left(\log(1-\sec(e+fx)) - \frac{2}{-1+\sec(e+fx)} \right) \tan(e+fx)}{cf\sqrt{a(1+\sec(e+fx))}\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(3/2),x]`

output $-\left((a^2 \cdot (\log[1 - \sec[e + f \cdot x]] - 2/(-1 + \sec[e + f \cdot x])) \cdot \tan[e + f \cdot x]) / (c \cdot f \cdot \sqrt{a \cdot (1 + \sec[e + f \cdot x])}) \cdot \sqrt{c - c \cdot \sec[e + f \cdot x]}\right)$

3.118.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4442, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(a \sec(e + fx) + a)^{3/2}}{(c - c \sec(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{3/2}}{\left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2}} dx$$

↓ 4442

$$\frac{a \int \frac{\sec(e + fx) \sqrt{\sec(e + fx) a + a}}{\sqrt{c - c \sec(e + fx)}} dx}{c} - \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a}}{f(c - c \sec(e + fx))^{3/2}}$$

↓ 3042

$$\frac{a \int \frac{\csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{\csc\left(e + fx + \frac{\pi}{2}\right) a + a}}{\sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)}} dx}{c} - \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a}}{f(c - c \sec(e + fx))^{3/2}}$$

↓ 4440

$$-\frac{a^2 \tan(e + fx) \log(1 - \sec(e + fx))}{c f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}} - \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a}}{f(c - c \sec(e + fx))^{3/2}}$$

input $\text{Int}[(\sec[e + f \cdot x] \cdot (a + a \cdot \sec[e + f \cdot x])^{3/2}) / (c - c \cdot \sec[e + f \cdot x])^{3/2}, x]$

output $-\left((a \cdot \sqrt{a + a \cdot \sec[e + f \cdot x]}) \cdot \tan[e + f \cdot x] / (f \cdot (c - c \cdot \sec[e + f \cdot x])^{3/2})\right) - \left(a^2 \cdot \log[1 - \sec[e + f \cdot x]] \cdot \tan[e + f \cdot x] / (c \cdot f \cdot \sqrt{a + a \cdot \sec[e + f \cdot x]}) \cdot \sqrt{c - c \cdot \sec[e + f \cdot x]}\right)$

3.118. $\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{3/2}}{(c - c \sec(e + fx))^{3/2}} dx$

3.118.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.118.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. 2(91) = 182.

Time = 3.40 (sec) , antiderivative size = 199, normalized size of antiderivative = 2.01

method	result
default	$\frac{a(\cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e)-1)+\cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e)+1)-2 \cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e)))}{f \sqrt{-c(\sec(fx+e)+1)}}$
risch	$\frac{ia \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (2e^{2i(fx+e)} \ln(e^{i(fx+e)}-1) - e^{2i(fx+e)} \ln(1+e^{2i(fx+e)}) - 4e^{i(fx+e)} \ln(e^{i(fx+e)}-1) + 2e^{i(fx+e)} \ln(1+e^{2i(fx+e)}))}{c(e^{i(fx+e)}+1)(e^{i(fx+e)}-1) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2), x, method=_RET URNVERBOSE)`

3.118.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx$$

output $-1/f*a*(\cos(f*x+e)*\ln(-\cot(f*x+e)+\csc(f*x+e)-1)+\cos(f*x+e)*\ln(-\cot(f*x+e)+\csc(f*x+e)+1)-2*\cos(f*x+e)*\ln(-\cot(f*x+e)+\csc(f*x+e))-\ln(-\cot(f*x+e)+\csc(f*x+e)-1)-\ln(-\cot(f*x+e)+\csc(f*x+e)+1)+2*\ln(-\cot(f*x+e)+\csc(f*x+e))-\cos(f*x+e)-1)*(a*(\sec(f*x+e)+1))^{1/2}/(-c*(\sec(f*x+e)-1))^{1/2}/(\sec(f*x+e)-1)/c/(\cos(f*x+e)+1)*\tan(f*x+e)$

3.118.5 Fricas [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{(a\sec(fx+e)+a)^{3/2}\sec(fx+e)}{(-c\sec(fx+e)+c)^{3/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2),x, algorith="fricas")`

output `integral((a*sec(f*x + e)^2 + a*sec(f*x + e))*sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(c^2*sec(f*x + e)^2 - 2*c^2*sec(f*x + e) + c^2), x)`

3.118.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{(a(\sec(e+fx)+1))^{3/2}\sec(e+fx)}{(-c(\sec(e+fx)-1))^{3/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(3/2),x)`

output `Integral((a*(sec(e + f*x) + 1))**(3/2)*sec(e + f*x)/(-c*(sec(e + f*x) - 1))**(3/2), x)`

3.118.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.23

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx = \frac{\sqrt{-aa} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}+1\right)}{c^{3/2}} + \frac{\sqrt{-aa} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}-1\right)}{c^{3/2}} - \frac{2\sqrt{-aa} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)}{c^{3/2}} \cdot f$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="maxima")
```

```
output (sqrt(-a)*a*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/c^(3/2) + sqrt(-a)*a*
log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/c^(3/2) - 2*sqrt(-a)*a*log(sin(f*
x + e)/(cos(f*x + e) + 1))/c^(3/2) + sqrt(-a)*a*(cos(f*x + e) + 1)^2/(c^(3
/2)*sin(f*x + e)^2))/f
```

3.118.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{(a\sec(fx+e)+a)^{3/2} \sec(fx+e)}{(-c\sec(fx+e)+c)^{3/2}} dx$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="giac")
```

```
output sage0*x
```

3.118.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^{3/2}}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

```
input int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),x
)
```

3.118. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx$

output `int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),
x)`

3.118. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{3/2}} dx$

3.119
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{5/2}} dx$$

3.119.1 Optimal result	866
3.119.2 Mathematica [A] (verified)	866
3.119.3 Rubi [A] (verified)	867
3.119.4 Maple [A] (verified)	868
3.119.5 Fricas [B] (verification not implemented)	868
3.119.6 Sympy [F]	869
3.119.7 Maxima [B] (verification not implemented)	869
3.119.8 Giac [F]	870
3.119.9 Mupad [B] (verification not implemented)	870

3.119.1 Optimal result

Integrand size = 36, antiderivative size = 42

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{5/2}} dx = -\frac{(a+a \sec(e+fx))^{3/2} \tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}}$$

output `-1/4*(a+a*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(5/2)`

3.119.2 Mathematica [A] (verified)

Time = 0.86 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{5/2}} dx = -\frac{(a(1+\sec(e+fx)))^{3/2} \tan(e+fx)}{4f(c-c \sec(e+fx))^{5/2}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/4*((a*(1 + Sec[e + f*x]))^(3/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(5/2))`

3.119.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx$$

↓ 4438

$$-\frac{\tan(e+fx)(a\sec(e+fx)+a)^{3/2}}{4f(c-c\sec(e+fx))^{5/2}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/4*((a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(5/2))`

3.119.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.119.4 Maple [A] (verified)

Time = 3.59 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.55

method	result	size
default	$-\frac{a\sqrt{a(\sec(fx+e)+1)}(\tan(fx+e)+\sec(fx+e)\tan(fx+e))}{4f(\sec(fx+e)-1)^2\sqrt{-c(\sec(fx+e)-1)}c^2}$	65
risch	$\frac{2ia\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{3i(fx+e)}+e^{i(fx+e)})}{c^2(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)^3\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}$	116

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2),x,method=_RET
URNVERBOSE)
```

```
output -1/4/f*a*(a*(sec(f*x+e)+1))^(1/2)/(sec(f*x+e)-1)^2/(-c*(sec(f*x+e)-1))^(1/
2)/c^2*(tan(f*x+e)+sec(f*x+e)*tan(f*x+e))
```

3.119.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(36) = 72.

Time = 0.27 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.26

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx = \frac{a\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\cos(fx+e)^2}{(c^3f\cos(fx+e))^2 - 2c^3f\cos(fx+e) + c^3f\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2),x, algo
rithm="fricas")
```

```
output a*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*
x + e))*cos(f*x + e)^2/((c^3*f*cos(f*x + e))^2 - 2*c^3*f*cos(f*x + e) + c^3
*f)*sin(f*x + e)
```

3.119.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{(a(\sec(e+fx)+1))^{3/2} \sec(e+fx)}{(-c(\sec(e+fx)-1))^{5/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(5/2),x)`

output `Integral((a*(sec(e + f*x) + 1))**(3/2)*sec(e + f*x)/(-c*(sec(e + f*x) - 1))**(5/2), x)`

3.119.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 533 vs. $2(36) = 72$.

Time = 0.42 (sec) , antiderivative size = 533, normalized size of antiderivative = 12.69

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx = \frac{\dots}{(c^3 \cos(4fx+4e)^2 + 16c^3 \cos(3fx+3e)^2 + 36c^3 \cos(2fx+2e)^2 + \dots)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="maxima")`

output `2*(6*a*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) + 6*a*cos(f*x + e)*sin(2*f*x + 2*e) - 6*a*cos(2*f*x + 2*e)*sin(f*x + e) - (a*sin(3*f*x + 3*e) + a*sin(f*x + e))*cos(4*f*x + 4*e) + (a*cos(3*f*x + 3*e) + a*cos(f*x + e))*sin(4*f*x + 4*e) - (6*a*cos(2*f*x + 2*e) + a)*sin(3*f*x + 3*e) - a*sin(f*x + e))*sqrt(a)*sqrt(c)/((c^3*cos(4*f*x + 4*e)^2 + 16*c^3*cos(3*f*x + 3*e)^2 + 36*c^3*cos(2*f*x + 2*e)^2 + 16*c^3*cos(f*x + e)^2 + c^3*sin(4*f*x + 4*e)^2 + 16*c^3*sin(3*f*x + 3*e)^2 + 36*c^3*sin(2*f*x + 2*e)^2 - 48*c^3*sin(2*f*x + 2*e)*sin(f*x + e) + 16*c^3*sin(f*x + e)^2 - 8*c^3*cos(f*x + e) + c^3 - 2*(4*c^3*cos(3*f*x + 3*e) - 6*c^3*cos(2*f*x + 2*e) + 4*c^3*cos(f*x + e) - c^3)*cos(4*f*x + 4*e) - 8*(6*c^3*cos(2*f*x + 2*e) - 4*c^3*cos(f*x + e) + c^3)*cos(3*f*x + 3*e) - 12*(4*c^3*cos(f*x + e) - c^3)*cos(2*f*x + 2*e) - 4*(2*c^3*sin(3*f*x + 3*e) - 3*c^3*sin(2*f*x + 2*e) + 2*c^3*sin(f*x + e))*sin(4*f*x + 4*e) - 16*(3*c^3*sin(2*f*x + 2*e) - 2*c^3*sin(f*x + e))*sin(3*f*x + 3*e))*f)`

3.119.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{(a\sec(fx+e)+a)^{\frac{3}{2}}\sec(fx+e)}{(-c\sec(fx+e)+c)^{\frac{5}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `sage0*x`

3.119.9 Mupad [B] (verification not implemented)

Time = 16.76 (sec) , antiderivative size = 165, normalized size of antiderivative = 3.93

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx = \frac{2a \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}} \sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}} (6 \sin(e+fx) - 8 \sin(2e+2fx) + 7 \sin(3e+3fx) - 4 \sin(4e+4fx) + \sin(5e+5fx))}{c^3 f (48 \cos(e+fx) + 15 \cos(2e+2fx) - 40 \cos(3e+3fx) + 26 \cos(4e+4fx) - 8 \cos(5e+5fx) + \cos(6e+6fx) - 42)}$$

input `int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x)`

output `-(2*a*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2)*((c*(cos(e + f*x) - 1))/cos(e + f*x))^(1/2)*(6*sin(e + f*x) - 8*sin(2*e + 2*f*x) + 7*sin(3*e + 3*f*x) - 4*sin(4*e + 4*f*x) + sin(5*e + 5*f*x)))/(c^3*f*(48*cos(e + f*x) + 15*cos(2*e + 2*f*x) - 40*cos(3*e + 3*f*x) + 26*cos(4*e + 4*f*x) - 8*cos(5*e + 5*f*x) + cos(6*e + 6*f*x) - 42))`

3.120
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{7/2}} dx$$

3.120.1 Optimal result 871
 3.120.2 Mathematica [A] (verified) 871
 3.120.3 Rubi [A] (verified) 872
 3.120.4 Maple [A] (verified) 873
 3.120.5 Fricas [A] (verification not implemented) 874
 3.120.6 Sympy [F(-1)] 874
 3.120.7 Maxima [B] (verification not implemented) 874
 3.120.8 Giac [F] 875
 3.120.9 Mupad [B] (verification not implemented) 876

3.120.1 Optimal result

Integrand size = 36, antiderivative size = 88

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{7/2}} dx = -\frac{(a+a \sec(e+fx))^{3/2} \tan(e+fx)}{6f(c-c \sec(e+fx))^{7/2}} - \frac{(a+a \sec(e+fx))^{3/2} \tan(e+fx)}{24cf(c-c \sec(e+fx))^{5/2}}$$

output `-1/6*(a+a*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(7/2)-1/24*(a+a*sec(f*x+e))^(3/2)*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(5/2)`

3.120.2 Mathematica [A] (verified)

Time = 1.69 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.77

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{7/2}} dx = \frac{a^2(1+3 \sec(e+fx)) \tan(e+fx)}{6c^3 f(-1+\sec(e+fx))^3 \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(7/2),x]`

output `(a^2*(1 + 3*Sec[e + f*x])*Tan[e + f*x])/((6*c^3*f*(-1 + Sec[e + f*x])^3*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.120.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{7/2}} dx$$

3.120.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a \sec(e+fx) + a)^{3/2}}{(c - c \sec(e+fx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a \csc(e+fx+\frac{\pi}{2}) + a)^{3/2}}{(c - c \csc(e+fx+\frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4439} \\
 & \frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^{3/2}}{(c-c\sec(e+fx))^{5/2}} dx}{6c} - \frac{\tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{6f(c - c \sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^{3/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx}{6c} - \frac{\tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{6f(c - c \sec(e+fx))^{7/2}} \\
 & \quad \downarrow \text{4438} \\
 & \frac{\tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{24cf(c - c \sec(e+fx))^{5/2}} - \frac{\tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{6f(c - c \sec(e+fx))^{7/2}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(7/2),x]`

output `-1/6*((a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(7/2)) - ((a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(24*c*f*(c - c*Sec[e + f*x])^(5/2))`

3.120.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.120.4 Maple [A] (verified)

Time = 3.23 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{a(5 \cos(fx+e)-1)\sqrt{a(\sec(fx+e)+1)}(\cos(fx+e)+1)\tan(fx+e)\sec(fx+e)^2}{24f(\sec(fx+e)-1)^3\sqrt{-c(\sec(fx+e)-1)}c^3}$	77
risch	$\frac{2ia\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(3e^{5i(fx+e)}-3e^{4i(fx+e)}+8e^{3i(fx+e)}-3e^{2i(fx+e)}+3e^{i(fx+e)})}{3c^3(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)^5\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}$	153

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`

output $\frac{1}{24}f*a*(5*\cos(f*x+e)-1)*(a*(\sec(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)+1)/(\sec(f*x+e)-1)^3/(-c*(\sec(f*x+e)-1))^{(1/2)}/c^3*\tan(f*x+e)*\sec(f*x+e)^2$

3.120.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{7/2}} dx = \frac{(6a\cos(fx+e)^3 - 3a\cos(fx+e)^2 + a\cos(fx+e))\sqrt{\frac{a\cos(fx+e)}{\cos(fx+e)}}}{6(c^4f\cos(fx+e)^3 - 3c^4f\cos(fx+e)^2 + 3c^4f\cos(fx+e) -$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(7/2),x, algo
rithm="fricas")`

output `1/6*(6*a*cos(f*x + e)^3 - 3*a*cos(f*x + e)^2 + a*cos(f*x + e))*sqrt((a*cos
(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((c^4
*f*cos(f*x + e)^3 - 3*c^4*f*cos(f*x + e)^2 + 3*c^4*f*cos(f*x + e) - c^4*f
*sin(f*x + e))`

3.120.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{7/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(7/2),x)`

output `Timed out`

3.120.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1559 vs. 2(76) = 152.

Time = 0.80 (sec) , antiderivative size = 1559, normalized size of antiderivative = 17.72

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{7/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(7/2),x, algo
rithm="maxima")`

output

```

2/3*(3*(a*sin(4*f*x + 4*e) + a*sin(2*f*x + 2*e))*cos(6*f*x + 6*e) + 3*(a*s
in(6*f*x + 6*e) + 9*a*sin(4*f*x + 4*e) + 9*a*sin(2*f*x + 2*e) - 4*a*sin(3/
2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(5/2*arctan2(sin(2*f*x
+ 2*e), cos(2*f*x + 2*e))) + 4*(2*a*sin(6*f*x + 6*e) + 15*a*sin(4*f*x + 4*
e) + 15*a*sin(2*f*x + 2*e) + 3*a*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e))))*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 3*(a*s
in(6*f*x + 6*e) + 9*a*sin(4*f*x + 4*e) + 9*a*sin(2*f*x + 2*e))*cos(1/2*arc
tan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 3*(a*cos(4*f*x + 4*e) + a*cos(
2*f*x + 2*e))*sin(6*f*x + 6*e) + 3*a*sin(4*f*x + 4*e) + 3*a*sin(2*f*x + 2*
e) - 3*(a*cos(6*f*x + 6*e) + 9*a*cos(4*f*x + 4*e) + 9*a*cos(2*f*x + 2*e) -
4*a*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + a)*sin(5/2*arc
tan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 4*(2*a*cos(6*f*x + 6*e) + 15*a
*cos(4*f*x + 4*e) + 15*a*cos(2*f*x + 2*e) + 3*a*cos(1/2*arctan2(sin(2*f*x
+ 2*e), cos(2*f*x + 2*e)))) + 2*a)*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*
f*x + 2*e))) - 3*(a*cos(6*f*x + 6*e) + 9*a*cos(4*f*x + 4*e) + 9*a*cos(2*f*
x + 2*e) + a)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a
)*sqrt(c)/((c^4*cos(6*f*x + 6*e)^2 + 225*c^4*cos(4*f*x + 4*e)^2 + 225*c^4*
cos(2*f*x + 2*e)^2 + 36*c^4*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e)))^2 + 400*c^4*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2
+ 36*c^4*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + c^4*s...

```

3.120.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{7/2}} dx = \int \frac{(a\sec(fx+e)+a)^{3/2}\sec(fx+e)}{(-c\sec(fx+e)+c)^{7/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(7/2),x, algo
rithm="giac")`

output `sage0*x`

3.120.9 Mupad [B] (verification not implemented)

Time = 18.89 (sec) , antiderivative size = 273, normalized size of antiderivative = 3.10

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{7/2}} dx = \frac{\sqrt{c-\frac{c}{\cos(e+fx)}} \left(\frac{a e^{e^{4i}+f x^{4i}} \sqrt{a+\frac{a}{\cos(e+fx)}}^{4i}}{c^4 f} - \frac{a \cos(e+fx) e^{e^{4i}+f x^{4i}} \sqrt{a+\frac{a}{\cos(e+fx)}}^{4i}}{3 c^4 f} \right)}{e^{e^{4i}+f x^{4i}} \sin(e+fx) 28i - e^{e^{4i}+f x^{4i}} \sin(2e+2fx)}$$

```
input int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(7/2)),x)
```

```
output ((c - c/cos(e + f*x))^(1/2)*((a*exp(e*4i + f*x*4i)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^4*f) - (a*cos(e + f*x)*exp(e*4i + f*x*4i)*(a + a/cos(e + f*x))^(1/2)*4i)/(3*c^4*f) + (a*exp(e*4i + f*x*4i)*cos(2*e + 2*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^4*f) - (a*exp(e*4i + f*x*4i)*cos(3*e + 3*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^4*f)))/(exp(e*4i + f*x*4i)*sin(e + f*x)*28i - exp(e*4i + f*x*4i)*sin(2*e + 2*f*x)*28i + exp(e*4i + f*x*4i)*sin(3*e + 3*f*x)*12i - exp(e*4i + f*x*4i)*sin(4*e + 4*f*x)*2i)
```

3.121
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{9/2}} dx$$

3.121.1 Optimal result 877
 3.121.2 Mathematica [A] (verified) 877
 3.121.3 Rubi [A] (verified) 878
 3.121.4 Maple [A] (verified) 879
 3.121.5 Fracas [A] (verification not implemented) 880
 3.121.6 Sympy [F(-1)] 880
 3.121.7 Maxima [B] (verification not implemented) 880
 3.121.8 Giac [F] 881
 3.121.9 Mupad [B] (verification not implemented) 882

3.121.1 Optimal result

Integrand size = 36, antiderivative size = 92

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{9/2}} dx = -\frac{a\sqrt{a+a \sec(e+fx)} \tan(e+fx)}{4f(c-c \sec(e+fx))^{9/2}} + \frac{a^2 \tan(e+fx)}{12cf\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{7/2}}$$

output `1/12*a^2*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(7/2)/(a+a*sec(f*x+e))^(1/2)-1/4*a*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(9/2)`

3.121.2 Mathematica [A] (verified)

Time = 4.36 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{9/2}} dx = \frac{a^2(1+2 \sec(e+fx)) \tan(e+fx)}{6c^4 f(-1+\sec(e+fx))^4 \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(9/2),x]`

output `-1/6*(a^2*(1 + 2*Sec[e + f*x])*Tan[e + f*x])/(c^4*f*(-1 + Sec[e + f*x])^4*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.121.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{9/2}} dx$$

3.121.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4442, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{9/2}} dx$$

↓ 4442

$$-\frac{a \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{(c-c\sec(e+fx))^{7/2}} dx}{4c} - \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{4f(c-c\sec(e+fx))^{9/2}}$$

↓ 3042

$$-\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{7/2}} dx}{4c} - \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{4f(c-c\sec(e+fx))^{9/2}}$$

↓ 4441

$$\frac{a^2 \tan(e+fx)}{12cf\sqrt{a\sec(e+fx)+a}(c-c\sec(e+fx))^{7/2}} - \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{4f(c-c\sec(e+fx))^{9/2}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(9/2),x]`

output `-1/4*(a*sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(9/2)) + (a^2*Tan[e + f*x])/(12*c*f*sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(7/2))`

3.121.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.121.4 Maple [A] (verified)

Time = 3.34 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.95

method	result	size
default	$-\frac{a(17 \cos(fx+e)^2 - 6 \cos(fx+e) + 1) \sqrt{a(\sec(fx+e)+1) (\cos(fx+e)+1) \tan(fx+e) \sec(fx+e)^3}}{96 f (\sec(fx+e)-1)^4 \sqrt{-c(\sec(fx+e)-1)} c^4}$	87
risch	$\frac{2ia \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (3e^{7i(fx+e)} - 6e^{6i(fx+e)} + 17e^{5i(fx+e)} - 16e^{4i(fx+e)} + 17e^{3i(fx+e)} - 6e^{2i(fx+e)} + 3e^{i(fx+e)})}{3c^4 (e^{i(fx+e)}+1) (e^{i(fx+e)}-1)^7 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$	175

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(9/2),x,method=_RETURNVERBOSE)`

output `-1/96/f*a*(17*cos(f*x+e)^2-6*cos(f*x+e)+1)*(a*(sec(f*x+e)+1))^(1/2)*(cos(f*x+e)+1)/(sec(f*x+e)-1)^4/(-c*(sec(f*x+e)-1))^(1/2)/c^4*tan(f*x+e)*sec(f*x+e)^3`

3.121.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx$$

3.121.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.72

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx = \frac{(6a\cos(fx+e)^4 - 6a\cos(fx+e)^3 + 4a\cos(fx+e)^2 - a\cos(fx+e))\sqrt{(a\cos(fx+e)+a)/\cos(fx+e)}\sqrt{(c\cos(fx+e)-c)/\cos(fx+e)}}{6(c^5f\cos(fx+e)^4 - 4c^5f\cos(fx+e)^3 + 6c^5f\cos(fx+e)^2 - 4c^5f\cos(fx+e) + c^5f)\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(9/2),x, algorith="fricas")`

output `1/6*(6*a*cos(f*x + e)^4 - 6*a*cos(f*x + e)^3 + 4*a*cos(f*x + e)^2 - a*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((c^5*f*cos(f*x + e)^4 - 4*c^5*f*cos(f*x + e)^3 + 6*c^5*f*cos(f*x + e)^2 - 4*c^5*f*cos(f*x + e) + c^5*f)*sin(f*x + e))`

3.121.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(9/2),x)`

output `Timed out`

3.121.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2608 vs. 2(80) = 160.

Time = 3.32 (sec) , antiderivative size = 2608, normalized size of antiderivative = 28.35

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(9/2),x, algorith="maxima")`

output `2/3*(28*a*cos(6*f*x + 6*e)*sin(4*f*x + 4*e) - 28*a*cos(4*f*x + 4*e)*sin(2*f*x + 2*e) + 2*(3*a*sin(6*f*x + 6*e) + 8*a*sin(4*f*x + 4*e) + 3*a*sin(2*f*x + 2*e))*cos(8*f*x + 8*e) + (3*a*sin(8*f*x + 8*e) + 36*a*sin(6*f*x + 6*e) + 82*a*sin(4*f*x + 4*e) + 36*a*sin(2*f*x + 2*e) - 32*a*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - 32*a*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (17*a*sin(8*f*x + 8*e) + 140*a*sin(6*f*x + 6*e) + 294*a*sin(4*f*x + 4*e) + 140*a*sin(2*f*x + 2*e) + 32*a*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (17*a*sin(8*f*x + 8*e) + 140*a*sin(6*f*x + 6*e) + 294*a*sin(4*f*x + 4*e) + 140*a*sin(2*f*x + 2*e) + 32*a*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (3*a*sin(8*f*x + 8*e) + 36*a*sin(6*f*x + 6*e) + 82*a*sin(4*f*x + 4*e) + 36*a*sin(2*f*x + 2*e))*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 2*(3*a*cos(6*f*x + 6*e) + 8*a*cos(4*f*x + 4*e) + 3*a*cos(2*f*x + 2*e))*sin(8*f*x + 8*e) - 2*(14*a*cos(4*f*x + 4*e) - 3*a)*sin(6*f*x + 6*e) + 4*(7*a*cos(2*f*x + 2*e) + 4*a)*sin(4*f*x + 4*e) + 6*a*sin(2*f*x + 2*e) - (3*a*cos(8*f*x + 8*e) + 36*a*cos(6*f*x + 6*e) + 82*a*cos(4*f*x + 4*e) + 36*a*cos(2*f*x + 2*e) - 32*a*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - 32*a*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 3*a)*sin(7/2*arctan2(s...`

3.121.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx = \int \frac{(a\sec(fx+e)+a)^{3/2}\sec(fx+e)}{(-c\sec(fx+e)+c)^{9/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(9/2),x, algo rithm="giac")`

output `sage0*x`

3.121.9 Mupad [B] (verification not implemented)

Time = 19.89 (sec) , antiderivative size = 340, normalized size of antiderivative = 3.70

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{9/2}} dx = \frac{\sqrt{c-\frac{c}{\cos(e+fx)}}}{e^{e^{5i+fx}5i} \sin(e+fx)} \left(\frac{a e^{e^{5i+fx}5i} \sqrt{a+\frac{a}{\cos(e+fx)}} 68i}{3c^5 f} - \frac{a \cos(e+fx) e^{e^{5i+fx}5i} \sqrt{a+\frac{a}{\cos(e+fx)}}}{3c^5 f} \right)$$

```
input int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(9/2)),x)
```

```
output ((c - c/cos(e + f*x))^(1/2)*((a*exp(e*5i + f*x*5i))*(a + a/cos(e + f*x))^(1/2)*68i)/(3*c^5*f) - (a*cos(e + f*x)*exp(e*5i + f*x*5i)*(a + a/cos(e + f*x))^(1/2)*88i)/(3*c^5*f) + (a*exp(e*5i + f*x*5i)*cos(2*e + 2*f*x)*(a + a/cos(e + f*x))^(1/2)*80i)/(3*c^5*f) - (a*exp(e*5i + f*x*5i)*cos(3*e + 3*f*x)*(a + a/cos(e + f*x))^(1/2)*8i)/(c^5*f) + (a*exp(e*5i + f*x*5i)*cos(4*e + 4*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^5*f))/(exp(e*5i + f*x*5i)*sin(e + f*x)*84i - exp(e*5i + f*x*5i)*sin(2*e + 2*f*x)*96i + exp(e*5i + f*x*5i)*sin(3*e + 3*f*x)*54i - exp(e*5i + f*x*5i)*sin(4*e + 4*f*x)*16i + exp(e*5i + f*x*5i)*sin(5*e + 5*f*x)*2i)
```

3.122
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{11/2}} dx$$

3.122.1 Optimal result	883
3.122.2 Mathematica [A] (verified)	883
3.122.3 Rubi [A] (verified)	884
3.122.4 Maple [A] (verified)	885
3.122.5 Fracas [B] (verification not implemented)	886
3.122.6 Sympy [F(-1)]	886
3.122.7 Maxima [B] (verification not implemented)	886
3.122.8 Giac [F]	887
3.122.9 Mupad [B] (verification not implemented)	888

3.122.1 Optimal result

Integrand size = 36, antiderivative size = 92

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{11/2}} dx = -\frac{a\sqrt{a+a \sec(e+fx)} \tan(e+fx)}{5f(c-c \sec(e+fx))^{11/2}} + \frac{a^2 \tan(e+fx)}{20cf\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{9/2}}$$

output `1/20*a^2*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(9/2)/(a+a*sec(f*x+e))^(1/2)-1/5*a*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(11/2)`

3.122.2 Mathematica [A] (verified)

Time = 5.38 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{11/2}} dx = \frac{a^2(3+5 \sec(e+fx)) \tan(e+fx)}{20c^5 f(-1+\sec(e+fx))^5 \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(11/2),x]`

output `(a^2*(3 + 5*Sec[e + f*x])*Tan[e + f*x])/(20*c^5*f*(-1 + Sec[e + f*x])^5*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.122.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{3/2}}{(c-c \sec(e+fx))^{11/2}} dx$$

3.122.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4442, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{11/2}} dx$$

↓ 4442

$$-\frac{a \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{(c-c\sec(e+fx))^{9/2}} dx}{5c} - \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{5f(c-c\sec(e+fx))^{11/2}}$$

↓ 3042

$$-\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{9/2}} dx}{5c} - \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{5f(c-c\sec(e+fx))^{11/2}}$$

↓ 4441

$$\frac{a^2 \tan(e+fx)}{20cf\sqrt{a\sec(e+fx)+a}(c-c\sec(e+fx))^{9/2}} - \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{5f(c-c\sec(e+fx))^{11/2}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(3/2))/(c - c*Sec[e + f*x])^(11/2), x]`

output `-1/5*(a*Sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(11/2)) + (a^2*Tan[e + f*x])/(20*c*f*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(9/2))`

3.122.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.122.4 Maple [A] (verified)

Time = 3.58 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.05

method	result
default	$\frac{a(49 \cos^3(fx+e) - 23 \cos^2(fx+e) + 7 \cos(fx+e) - 1) \sqrt{a(\sec(fx+e)+1)} (\cos(fx+e)+1) \tan(fx+e) \sec(fx+e)^4}{320 f (\sec(fx+e)-1)^5 \sqrt{-c(\sec(fx+e)-1)} c^5}$
risch	$\frac{2ia \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (5e^{9i(fx+e)} - 15e^{8i(fx+e)} + 50e^{7i(fx+e)} - 75e^{6i(fx+e)} + 102e^{5i(fx+e)} - 75e^{4i(fx+e)} + 50e^{3i(fx+e)} - 15e^{2i(fx+e)} - 5c^5 (e^{i(fx+e)}+1) (e^{i(fx+e)}-1)^9 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}{}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(11/2),x,method=_RETURNVERBOSE)`

output `1/320/f*a*(49*cos(f*x+e)^3-23*cos(f*x+e)^2+7*cos(f*x+e)-1)*(a*(sec(f*x+e)+1))^(1/2)*(cos(f*x+e)+1)/(sec(f*x+e)-1)^5/(-c*(sec(f*x+e)-1))^(1/2)/c^5*tan(f*x+e)*sec(f*x+e)^4`

3.122.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx$$

3.122.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. 2(80) = 160.

Time = 0.28 (sec) , antiderivative size = 184, normalized size of antiderivative = 2.00

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx = \frac{(20a\cos(fx+e)^5 - 30a\cos(fx+e)^4 + 30a\cos(fx+e)^3 - 15a^2\cos(fx+e)^2 + 3a^2\cos(fx+e))\sqrt{(a\cos(fx+e)+a)/\cos(fx+e)}\sqrt{(c\cos(fx+e)-c)/\cos(fx+e)}}{20(c^6f\cos(fx+e)^5 - 5c^6f\cos(fx+e)^4 + 10c^6f\cos(fx+e)^3 - 10c^6f\cos(fx+e)^2 + 5c^6f\cos(fx+e) - c^6f)\sin(fx+e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(11/2),x, algorithm="fracas")`

output `1/20*(20*a*cos(f*x + e)^5 - 30*a*cos(f*x + e)^4 + 30*a*cos(f*x + e)^3 - 15*a*cos(f*x + e)^2 + 3*a*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((c^6*f*cos(f*x + e)^5 - 5*c^6*f*cos(f*x + e)^4 + 10*c^6*f*cos(f*x + e)^3 - 10*c^6*f*cos(f*x + e)^2 + 5*c^6*f*cos(f*x + e) - c^6*f)*sin(f*x + e))`

3.122.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(11/2),x)`

output `Timed out`

3.122.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3906 vs. 2(80) = 160.

Time = 16.64 (sec) , antiderivative size = 3906, normalized size of antiderivative = 42.46

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(11/2),x, algorithm="maxima")`

output `-2/5*(225*a*cos(6*f*x + 6*e)*sin(2*f*x + 2*e) + 225*a*cos(4*f*x + 4*e)*sin(2*f*x + 2*e) - 15*(a*sin(8*f*x + 8*e) + 5*a*sin(6*f*x + 6*e) + 5*a*sin(4*f*x + 4*e) + a*sin(2*f*x + 2*e))*cos(10*f*x + 10*e) - 225*(a*sin(6*f*x + 6*e) + a*sin(4*f*x + 4*e))*cos(8*f*x + 8*e) - 5*(a*sin(10*f*x + 10*e) + 15*a*sin(8*f*x + 8*e) + 60*a*sin(6*f*x + 6*e) + 60*a*sin(4*f*x + 4*e) + 15*a*sin(2*f*x + 2*e) - 20*a*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - 48*a*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 20*a*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 10*(5*a*sin(10*f*x + 10*e) + 45*a*sin(8*f*x + 8*e) + 150*a*sin(6*f*x + 6*e) + 150*a*sin(4*f*x + 4*e) + 45*a*sin(2*f*x + 2*e) - 36*a*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 10*a*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 6*(17*a*sin(10*f*x + 10*e) + 135*a*sin(8*f*x + 8*e) + 420*a*sin(6*f*x + 6*e) + 420*a*sin(4*f*x + 4*e) + 135*a*sin(2*f*x + 2*e) + 60*a*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 40*a*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 50*(a*sin(10*f*x + 10*e) + 9*a*sin(8*f*x + 8*e) + 30*a*sin(6*f*x + 6*e) + 30*a*sin(4*f*x + 4*e) + 9*a*sin(2*f*x + 2*e) + 2*a*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 5*(a*sin(10*f*x ...`

3.122.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx = \int \frac{(a\sec(fx+e)+a)^{3/2}\sec(fx+e)}{(-c\sec(fx+e)+c)^{11/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(11/2),x, algorithm="giac")`

output `sage0*x`

3.122.9 Mupad [B] (verification not implemented)

Time = 19.24 (sec) , antiderivative size = 407, normalized size of antiderivative = 4.42

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{3/2}}{(c-c\sec(e+fx))^{11/2}} dx = \sqrt{c - \frac{c}{\cos(e+fx)}} \left(\frac{a e^{e6i+fx6i} \sqrt{a + \frac{a}{\cos(e+fx)}} 60i}{c^6 f} - \frac{a \cos(e+fx) e^{e6i+fx6i} \sqrt{a + \frac{a}{\cos(e+fx)}}}{5 c^6 f} \right) \frac{1}{e^{e6i+fx6i} \sin(e+fx) 264i - e^{e6i+fx6i} \sin(2e+2fx)}$$

```
input int((a + a/cos(e + f*x))^(3/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(11/2)),
x)
```

```
output ((c - c/cos(e + f*x))^(1/2)*((a*exp(e*6i + f*x*6i)*(a + a/cos(e + f*x))^(1/2)*60i)/(c^6*f) - (a*cos(e + f*x)*exp(e*6i + f*x*6i)*(a + a/cos(e + f*x))^(1/2)*608i)/(5*c^6*f) + (a*exp(e*6i + f*x*6i)*cos(2*e + 2*f*x)*(a + a/cos(e + f*x))^(1/2)*72i)/(c^6*f) - (a*exp(e*6i + f*x*6i)*cos(3*e + 3*f*x)*(a + a/cos(e + f*x))^(1/2)*44i)/(c^6*f) + (a*exp(e*6i + f*x*6i)*cos(4*e + 4*f*x)*(a + a/cos(e + f*x))^(1/2)*12i)/(c^6*f) - (a*exp(e*6i + f*x*6i)*cos(5*e + 5*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^6*f)))/(exp(e*6i + f*x*6i)*sin(e + f*x)*264i - exp(e*6i + f*x*6i)*sin(2*e + 2*f*x)*330i + exp(e*6i + f*x*6i)*sin(3*e + 3*f*x)*220i - exp(e*6i + f*x*6i)*sin(4*e + 4*f*x)*88i + exp(e*6i + f*x*6i)*sin(5*e + 5*f*x)*20i - exp(e*6i + f*x*6i)*sin(6*e + 6*f*x)*2i)
```

3.123 $\int \sec(e+fx)(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{7/2} dx$

3.123.1 Optimal result	889
3.123.2 Mathematica [A] (verified)	890
3.123.3 Rubi [A] (verified)	890
3.123.4 Maple [A] (verified)	892
3.123.5 Fricas [A] (verification not implemented)	893
3.123.6 Sympy [F(-1)]	893
3.123.7 Maxima [B] (verification not implemented)	893
3.123.8 Giac [F]	894
3.123.9 Mupad [B] (verification not implemented)	895

3.123.1 Optimal result

Integrand size = 36, antiderivative size = 134

$$\int \sec(e+fx)(a + a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{7/2} dx = \frac{a^3(c-c \sec(e+fx))^{7/2} \tan(e+fx)}{15f \sqrt{a+a \sec(e+fx)}} + \frac{2a^2 \sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{7/2} \tan(e+fx)}{15f} + \frac{a(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{7/2} \tan(e+fx)}{6f}$$

```
output 1/6*a*(a+a*sec(f*x+e))^(3/2)*(c-c*sec(f*x+e))^(7/2)*tan(f*x+e)/f+1/15*a^3*(c-c*sec(f*x+e))^(7/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)+2/15*a^2*(c-c*sec(f*x+e))^(7/2)*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.123.2 Mathematica [A] (verified)

Time = 4.30 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.69

$$\int \sec(e+fx)(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{7/2} dx = \frac{4a^2c^4(21+28\cos(e+fx)+11\cos(2(e+fx)))\sec^6(e+fx)\sqrt{a(1+\sec(e+fx))}\sin(e+fx)}{15f\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(7/2), x]`

output `(4*a^2*c^4*(21 + 28*Cos[e + f*x] + 11*Cos[2*(e + f*x)])*Sec[e + f*x]^6*Sin[e + f*x]*Sqrt[a*(1 + Sec[e + f*x])]*Sin[(e + f*x)/2]^8*Tan[(e + f*x)/2])/(15*f*Sqrt[c - c*Sec[e + f*x]])`

3.123.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e+fx)(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^{5/2}\left(c-c\csc\left(e+fx+\frac{\pi}{2}\right)\right)^{7/2} dx \\ & \quad \downarrow \text{4443} \\ & \frac{2}{3}a \int \sec(e+fx)(\sec(e+fx)a+a)^{3/2}(c-c\sec(e+fx))^{7/2} dx + \\ & \quad \frac{a \tan(e+fx)(a\sec(e+fx)+a)^{3/2}(c-c\sec(e+fx))^{7/2}}{6f} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{2}{3}a \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^{3/2} \left(c - c\csc\left(e + fx + \frac{\pi}{2}\right)\right)^{7/2} dx + \frac{a \tan(e + fx)(a \sec(e + fx) + a)^{3/2}(c - c \sec(e + fx))^{7/2}}{6f}$$

↓ 4443

$$\frac{2}{3}a \left(\frac{2}{5}a \int \sec(e + fx) \sqrt{\sec(e + fx)a + a} (c - c \sec(e + fx))^{7/2} dx + \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a} (c - c \sec(e + fx))^{7/2}}{5f} \right) + \frac{a \tan(e + fx)(a \sec(e + fx) + a)^{3/2}(c - c \sec(e + fx))^{7/2}}{6f}$$

↓ 3042

$$\frac{2}{3}a \left(\frac{2}{5}a \int \csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{\csc\left(e + fx + \frac{\pi}{2}\right)a + a} \left(c - c\csc\left(e + fx + \frac{\pi}{2}\right)\right)^{7/2} dx + \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a} (c - c \sec(e + fx))^{7/2}}{5f} \right) + \frac{a \tan(e + fx)(a \sec(e + fx) + a)^{3/2}(c - c \sec(e + fx))^{7/2}}{6f}$$

↓ 4441

$$\frac{2}{3}a \left(\frac{a^2 \tan(e + fx)(c - c \sec(e + fx))^{7/2}}{10f \sqrt{a \sec(e + fx) + a}} + \frac{a \tan(e + fx) \sqrt{a \sec(e + fx) + a} (c - c \sec(e + fx))^{7/2}}{5f} \right) + \frac{a \tan(e + fx)(a \sec(e + fx) + a)^{3/2}(c - c \sec(e + fx))^{7/2}}{6f}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(7/2),x]`

output `(a*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(7/2)*Tan[e + f*x])/(6*f) + (2*a*((a^2*(c - c*Sec[e + f*x])^(7/2)*Tan[e + f*x])/(10*f*Sqrt[a + a*Sec[e + f*x]]) + (a*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(7/2)*Tan[e + f*x])/(5*f)))/3`

3.123.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.123.4 Maple [A] (verified)

Time = 1.50 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.83

$$\frac{a^2(21 \cos (fx + e)^3 - 33 \cos (fx + e)^2 + 21 \cos (fx + e) - 5) (\sec (fx + e) - 1)^3 \sqrt{a(\sec (fx + e) + 1)} \sqrt{a(\sec (fx + e) - 1)}}{30 f (\cos (fx + e) - 1)^3}$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(7/2),x)`

output `-1/30/f*a^2*(21*cos(f*x+e)^3-33*cos(f*x+e)^2+21*cos(f*x+e)-5)*(sec(f*x+e)-1)^3*(a*(sec(f*x+e)+1))^(1/2)*(-c*(sec(f*x+e)-1))^(1/2)*c^3*(cos(f*x+e)+1)^3/(cos(f*x+e)-1)^3*sec(f*x+e)^2*csc(f*x+e)`

3.123.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.13

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{7/2} dx = \frac{(30 a^2 c^3 \cos(fx + e)^5 - 15 a^2 c^3 \cos(fx + e)^4 - 20 a^2 c^3 \cos(fx + e)^3 + 15 a^2 c^3 \cos(fx + e)^2 + 6 a^2 c^3 \cos(fx + e) - 5 a^2 c^3) \sqrt{(a \cos(fx + e) + a) / \cos(fx + e)} \sqrt{(c \cos(fx + e) - c) / \cos(fx + e)}}{30 f \cos(fx + e)^5 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(7/2),x, algorithm="fricas")`

output `1/30*(30*a^2*c^3*cos(f*x + e)^5 - 15*a^2*c^3*cos(f*x + e)^4 - 20*a^2*c^3*cos(f*x + e)^3 + 15*a^2*c^3*cos(f*x + e)^2 + 6*a^2*c^3*cos(f*x + e) - 5*a^2*c^3)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)^5*sin(f*x + e))`

3.123.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)*(c-c*sec(f*x+e))**(7/2),x)`

output `Timed out`

3.123.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2454 vs. 2(116) = 232.

Time = 0.43 (sec) , antiderivative size = 2454, normalized size of antiderivative = 18.31

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{7/2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(7/2),x, algo
rithm="maxima")`

output `2/15*(210*a^2*c^3*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) - 90*a^2*c^3*cos(2*f*x
+ 2*e)*sin(f*x + e) - 15*a^2*c^3*sin(f*x + e) - (15*a^2*c^3*sin(11*f*x +
11*e) - 15*a^2*c^3*sin(10*f*x + 10*e) + 35*a^2*c^3*sin(9*f*x + 9*e) + 78*a
^2*c^3*sin(7*f*x + 7*e) - 50*a^2*c^3*sin(6*f*x + 6*e) + 78*a^2*c^3*sin(5*f
*x + 5*e) + 35*a^2*c^3*sin(3*f*x + 3*e) - 15*a^2*c^3*sin(2*f*x + 2*e) + 15
*a^2*c^3*sin(f*x + e))*cos(12*f*x + 12*e) + 15*(6*a^2*c^3*sin(10*f*x + 10*
e) + 15*a^2*c^3*sin(8*f*x + 8*e) + 20*a^2*c^3*sin(6*f*x + 6*e) + 15*a^2*c
^3*sin(4*f*x + 4*e) + 6*a^2*c^3*sin(2*f*x + 2*e))*cos(11*f*x + 11*e) - 3*(7
0*a^2*c^3*sin(9*f*x + 9*e) + 75*a^2*c^3*sin(8*f*x + 8*e) + 156*a^2*c^3*sin
(7*f*x + 7*e) + 156*a^2*c^3*sin(5*f*x + 5*e) + 75*a^2*c^3*sin(4*f*x + 4*e)
+ 70*a^2*c^3*sin(3*f*x + 3*e) + 30*a^2*c^3*sin(f*x + e))*cos(10*f*x + 10*
e) + 35*(15*a^2*c^3*sin(8*f*x + 8*e) + 20*a^2*c^3*sin(6*f*x + 6*e) + 15*a^
2*c^3*sin(4*f*x + 4*e) + 6*a^2*c^3*sin(2*f*x + 2*e))*cos(9*f*x + 9*e) - 15
*(78*a^2*c^3*sin(7*f*x + 7*e) - 50*a^2*c^3*sin(6*f*x + 6*e) + 78*a^2*c^3*s
in(5*f*x + 5*e) + 35*a^2*c^3*sin(3*f*x + 3*e) - 15*a^2*c^3*sin(2*f*x + 2*e
) + 15*a^2*c^3*sin(f*x + e))*cos(8*f*x + 8*e) + 78*(20*a^2*c^3*sin(6*f*x +
6*e) + 15*a^2*c^3*sin(4*f*x + 4*e) + 6*a^2*c^3*sin(2*f*x + 2*e))*cos(7*f*
x + 7*e) - 10*(156*a^2*c^3*sin(5*f*x + 5*e) + 75*a^2*c^3*sin(4*f*x + 4*e)
+ 70*a^2*c^3*sin(3*f*x + 3*e) + 30*a^2*c^3*sin(f*x + e))*cos(6*f*x + 6*e)
+ 234*(5*a^2*c^3*sin(4*f*x + 4*e) + 2*a^2*c^3*sin(2*f*x + 2*e))*cos(5*f...`

3.123.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{7/2} dx = \int (a \sec(fx + e) + a)^{5/2}(-c \sec(fx + e) + c)^{7/2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(7/2),x, algo
rithm="giac")`

output `sage0*x`

3.123.9 Mupad [B] (verification not implemented)

Time = 17.76 (sec) , antiderivative size = 307, normalized size of antiderivative = 2.29

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{7/2} dx = \frac{\sqrt{c - \frac{c}{\cos(e + fx)}} \left(-\frac{a^2 c^3 e^{e 6i + f x 6i} \sqrt{a + \frac{a}{\cos(e + fx)}} 20i}{3f} + \frac{a^2 c^3 \cos(e + fx) e^{e 6i + f x 6i} \sqrt{a + \frac{a}{\cos(e + fx)}} 104i}{5f} \right)}{e^{e 6i + f x 6i} \sin(2e + 2fx) 10}$$

```
input int(((a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(7/2))/cos(e + f*x),x)
```

```
output ((c - c/cos(e + f*x))^(1/2)*((a^2*c^3*cos(e + f*x)*exp(e*6i + f*x*6i)*(a + a/cos(e + f*x))^(1/2)*104i)/(5*f) - (a^2*c^3*exp(e*6i + f*x*6i)*(a + a/cos(e + f*x))^(1/2)*20i)/(3*f) + (a^2*c^3*exp(e*6i + f*x*6i)*cos(3*e + 3*f*x)*(a + a/cos(e + f*x))^(1/2)*28i)/(3*f) - (a^2*c^3*exp(e*6i + f*x*6i)*cos(4*e + 4*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/f + (a^2*c^3*exp(e*6i + f*x*6i)*cos(5*e + 5*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/f))/(exp(e*6i + f*x*6i)*sin(2*e + 2*f*x)*10i + exp(e*6i + f*x*6i)*sin(4*e + 4*f*x)*8i + exp(e*6i + f*x*6i)*sin(6*e + 6*f*x)*2i)
```


3.124 $\int \sec(e+fx)(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2} dx$

3.124.1 Optimal result	896
3.124.2 Mathematica [A] (verified)	897
3.124.3 Rubi [A] (verified)	897
3.124.4 Maple [A] (verified)	899
3.124.5 Fricas [A] (verification not implemented)	899
3.124.6 Sympy [F(-1)]	900
3.124.7 Maxima [B] (verification not implemented)	900
3.124.8 Giac [F]	901
3.124.9 Mupad [B] (verification not implemented)	902

3.124.1 Optimal result

Integrand size = 36, antiderivative size = 134

$$\int \sec(e+fx)(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2} dx =$$

$$\frac{2c^3(a+a \sec(e+fx))^{5/2} \tan(e+fx)}{15f \sqrt{c-c \sec(e+fx)}} - \frac{c^2(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)} \tan(e+fx)}{5f} - \frac{c(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2} \tan(e+fx)}{5f}$$

```
output -1/5*c*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f-2/15*c^3
*(a+a*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-1/5*c^2*(a+a*s
ec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.124.2 Mathematica [A] (verified)

Time = 1.23 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.57

$$\int \sec(e+fx)(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{5/2} dx = \frac{a^3 c^3 \sec(e+fx)(15-10\sec^2(e+fx)+3\sec^4(e+fx))\tan(e+fx)}{15f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/15*(a^3*c^3*Sec[e + f*x]*(15 - 10*Sec[e + f*x]^2 + 3*Sec[e + f*x]^4)*Tan[e + f*x])/(f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.124.3 Rubi [A] (verified)Time = 0.82 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e+fx)(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^{5/2}\left(c-c\csc\left(e+fx+\frac{\pi}{2}\right)\right)^{5/2} dx \\ & \quad \downarrow \text{4443} \\ & \frac{4}{5}c \int \sec(e+fx)(\sec(e+fx)a+a)^{5/2}(c-c\sec(e+fx))^{3/2} dx - \\ & \quad \frac{c \tan(e+fx)(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{3/2}}{5f} \\ & \quad \downarrow \text{3042} \\ & \frac{4}{5}c \int \csc\left(e+fx+\frac{\pi}{2}\right)\left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)^{5/2}\left(c-c\csc\left(e+fx+\frac{\pi}{2}\right)\right)^{3/2} dx - \\ & \quad \frac{c \tan(e+fx)(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{3/2}}{5f} \end{aligned}$$

3.124. $\int \sec(e+fx)(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{5/2} dx$

↓ 4443

$$\frac{4}{5}c \left(\frac{1}{2}c \int \sec(e+fx)(\sec(e+fx)a+a)^{5/2} \sqrt{c-c\sec(e+fx)} dx - \frac{c \tan(e+fx)(a \sec(e+fx)+a)^{5/2} \sqrt{c-c\sec(e+fx)}}{4f} \right) - \frac{c \tan(e+fx)(a \sec(e+fx)+a)^{5/2} (c-c\sec(e+fx))^{3/2}}{5f}$$

↓ 3042

$$\frac{4}{5}c \left(\frac{1}{2}c \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)^{5/2} \sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)} dx - \frac{c \tan(e+fx)(a \sec(e+fx)+a)^{5/2} \sqrt{c-c\sec(e+fx)}}{4f} \right) - \frac{c \tan(e+fx)(a \sec(e+fx)+a)^{5/2} (c-c\sec(e+fx))^{3/2}}{5f}$$

↓ 4441

$$\frac{4}{5}c \left(-\frac{c^2 \tan(e+fx)(a \sec(e+fx)+a)^{5/2}}{6f \sqrt{c-c\sec(e+fx)}} - \frac{c \tan(e+fx)(a \sec(e+fx)+a)^{5/2} \sqrt{c-c\sec(e+fx)}}{4f} \right) - \frac{c \tan(e+fx)(a \sec(e+fx)+a)^{5/2} (c-c\sec(e+fx))^{3/2}}{5f}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/5*(c*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/f + (4*c*(-1/6*(c^2*(a + a*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]]) - (c*(a + a*Sec[e + f*x])^(5/2)*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(4*f)))/5`

3.124.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.124.4 Maple [A] (verified)

Time = 1.66 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.75

$$\frac{a^2(\sec(fx + e) - 1)^2 (8 \cos(fx + e)^2 - 9 \cos(fx + e) + 3) \sqrt{-c(\sec(fx + e) - 1)} c^2 \sqrt{a(\sec(fx + e) + 1)}}{15f(\cos(fx + e) - 1)^2}$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(5/2),x)`

output `1/15/f*a^2*(sec(f*x+e)-1)^2*(8*cos(f*x+e)^2-9*cos(f*x+e)+3)*(-c*(sec(f*x+e)-1))^(1/2)*c^2*(a*(sec(f*x+e)+1))^(1/2)*(cos(f*x+e)+1)^3/(cos(f*x+e)-1)^2*sec(f*x+e)^2*csc(f*x+e)`

3.124.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.79

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{5/2} dx = \frac{(15 a^2 c^2 \cos(fx + e)^4 - 10 a^2 c^2 \cos(fx + e)^2 + 3 a^2 c^2) \sqrt{\frac{a \cos(fx + e) + a}{\cos(fx + e)}} \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}}}{15 f \cos(fx + e)^4 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(5/2),x, algorithm="fracas")`

output `1/15*(15*a^2*c^2*cos(f*x + e)^4 - 10*a^2*c^2*cos(f*x + e)^2 + 3*a^2*c^2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)^4*sin(f*x + e))`

3.124. $\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{5/2} dx$

3.124.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{5/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)*(c-c*sec(f*x+e))**(5/2),x)`

output `Timed out`

3.124.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1526 vs. $2(116) = 232$.

Time = 0.40 (sec) , antiderivative size = 1526, normalized size of antiderivative = 11.39

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{5/2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(5/2),x, algorith="maxima")`

output

```

2/15*(100*a^2*c^2*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) + 75*a^2*c^2*cos(f*x +
e)*sin(2*f*x + 2*e) - 75*a^2*c^2*cos(2*f*x + 2*e)*sin(f*x + e) - 15*a^2*c
^2*sin(f*x + e) - (15*a^2*c^2*sin(9*f*x + 9*e) + 20*a^2*c^2*sin(7*f*x + 7*
e) + 58*a^2*c^2*sin(5*f*x + 5*e) + 20*a^2*c^2*sin(3*f*x + 3*e) + 15*a^2*c^
2*sin(f*x + e))*cos(10*f*x + 10*e) + 75*(a^2*c^2*sin(8*f*x + 8*e) + 2*a^2*
c^2*sin(6*f*x + 6*e) + 2*a^2*c^2*sin(4*f*x + 4*e) + a^2*c^2*sin(2*f*x + 2*
e))*cos(9*f*x + 9*e) - 5*(20*a^2*c^2*sin(7*f*x + 7*e) + 58*a^2*c^2*sin(5*f
*x + 5*e) + 20*a^2*c^2*sin(3*f*x + 3*e) + 15*a^2*c^2*sin(f*x + e))*cos(8*f
*x + 8*e) + 100*(2*a^2*c^2*sin(6*f*x + 6*e) + 2*a^2*c^2*sin(4*f*x + 4*e) +
a^2*c^2*sin(2*f*x + 2*e))*cos(7*f*x + 7*e) - 10*(58*a^2*c^2*sin(5*f*x + 5
*e) + 20*a^2*c^2*sin(3*f*x + 3*e) + 15*a^2*c^2*sin(f*x + e))*cos(6*f*x + 6
*e) + 290*(2*a^2*c^2*sin(4*f*x + 4*e) + a^2*c^2*sin(2*f*x + 2*e))*cos(5*f*
x + 5*e) - 50*(4*a^2*c^2*sin(3*f*x + 3*e) + 3*a^2*c^2*sin(f*x + e))*cos(4*
f*x + 4*e) + (15*a^2*c^2*cos(9*f*x + 9*e) + 20*a^2*c^2*cos(7*f*x + 7*e) +
58*a^2*c^2*cos(5*f*x + 5*e) + 20*a^2*c^2*cos(3*f*x + 3*e) + 15*a^2*c^2*cos
(f*x + e))*sin(10*f*x + 10*e) - 15*(5*a^2*c^2*cos(8*f*x + 8*e) + 10*a^2*c^
2*cos(6*f*x + 6*e) + 10*a^2*c^2*cos(4*f*x + 4*e) + 5*a^2*c^2*cos(2*f*x + 2
*e) + a^2*c^2)*sin(9*f*x + 9*e) + 5*(20*a^2*c^2*cos(7*f*x + 7*e) + 58*a^2*
c^2*cos(5*f*x + 5*e) + 20*a^2*c^2*cos(3*f*x + 3*e) + 15*a^2*c^2*cos(f*x +
e))*sin(8*f*x + 8*e) - 20*(10*a^2*c^2*cos(6*f*x + 6*e) + 10*a^2*c^2*cos...

```

3.124.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{5/2} dx = \int (a \sec(fx + e) + a)^{5/2}(-c \sec(fx + e) + c)^{5/2} \sec(fx + e) dx$$

input

```

integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(5/2),x, algo
rithm="giac")

```

output

```

sage0*x

```

3.124.9 Mupad [B] (verification not implemented)

Time = 17.36 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.60

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{5/2} dx = \frac{\sqrt{c - \frac{c}{\cos(e+fx)}} \left(\frac{a^2 c^2 e^{e 5i + f x 5i} \sqrt{a + \frac{a}{\cos(e+fx)}} 116i}{15 f} + \frac{a^2 c^2 e^{e 5i + f x 5i} \cos(2e + 2fx) \sqrt{a + \frac{a}{\cos(e+fx)}} 16i}{3 f} \right)}{e^{e 5i + f x 5i} \sin(e + fx) 4i + e^{e 5i + f x 5i} \sin(3e + 3fx) 6i + e^{e 5i + f x 5i} \sin(5e + 5fx) 2i}$$

```
input int(((a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x),x
)
```

```
output ((c - c/cos(e + f*x))^(1/2)*((a^2*c^2*exp(e*5i + f*x*5i)*(a + a/cos(e + f*
x))^(1/2)*116i)/(15*f) + (a^2*c^2*exp(e*5i + f*x*5i)*cos(2*e + 2*f*x)*(a +
a/cos(e + f*x))^(1/2)*16i)/(3*f) + (a^2*c^2*exp(e*5i + f*x*5i)*cos(4*e +
4*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/f))/(exp(e*5i + f*x*5i)*sin(e + f*x)
*4i + exp(e*5i + f*x*5i)*sin(3*e + 3*f*x)*6i + exp(e*5i + f*x*5i)*sin(5*e
+ 5*f*x)*2i)
```

3.125 $\int \sec(e+fx)(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2} dx$

3.125.1 Optimal result	903
3.125.2 Mathematica [A] (verified)	903
3.125.3 Rubi [A] (verified)	904
3.125.4 Maple [A] (verified)	905
3.125.5 Fricas [A] (verification not implemented)	906
3.125.6 Sympy [F(-1)]	906
3.125.7 Maxima [B] (verification not implemented)	907
3.125.8 Giac [F]	907
3.125.9 Mupad [B] (verification not implemented)	908

3.125.1 Optimal result

Integrand size = 36, antiderivative size = 89

$$\int \sec(e+fx)(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2} dx =$$

$$\frac{c^2(a+a \sec(e+fx))^{5/2} \tan(e+fx)}{6f \sqrt{c-c \sec(e+fx)}} - \frac{c(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)} \tan(e+fx)}{4f}$$

```
output -1/6*c^2*(a+a*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)-1/4*c*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f
```

3.125.2 Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.99

$$\int \sec(e+fx)(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2} dx =$$

$$\frac{a^2 c^2 (5 \cos(e+fx) + 3(\cos(2(e+fx)) + \cos(3(e+fx)))) \sec^4(e+fx) \sqrt{a(1 + \sec(e+fx))} \tan(\frac{1}{2}(e+fx))}{12f \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(3/2),x]`

output `-1/12*(a^2*c^2*(5*Cos[e + f*x] + 3*(Cos[2*(e + f*x)] + Cos[3*(e + f*x)]))*Sec[e + f*x]^4*sqrt[a*(1 + Sec[e + f*x])]*Tan[(e + f*x)/2]/(f*sqrt[c - c*Sec[e + f*x]])`

3.125.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^{5/2}(c - c \sec(e + fx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{1}{2}c \int \sec(e + fx)(\sec(e + fx)a + a)^{5/2} \sqrt{c - c \sec(e + fx)} dx - \\
 & \quad \frac{c \tan(e + fx)(a \sec(e + fx) + a)^{5/2} \sqrt{c - c \sec(e + fx)}}{4f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right) a + a\right)^{5/2} \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx - \\
 & \quad \frac{c \tan(e + fx)(a \sec(e + fx) + a)^{5/2} \sqrt{c - c \sec(e + fx)}}{4f} \\
 & \quad \downarrow \text{4441} \\
 & \frac{c^2 \tan(e + fx)(a \sec(e + fx) + a)^{5/2}}{6f \sqrt{c - c \sec(e + fx)}} - \frac{c \tan(e + fx)(a \sec(e + fx) + a)^{5/2} \sqrt{c - c \sec(e + fx)}}{4f}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(3/2),x]`

output
$$-1/6*(c^2*(a + a*\text{Sec}[e + f*x])^{5/2}*\text{Tan}[e + f*x])/(f*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]) - (c*(a + a*\text{Sec}[e + f*x])^{5/2}*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]*\text{Tan}[e + f*x])/ (4*f)$$

3.125.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.125.4 Maple [A] (verified)

Time = 3.66 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.98

method	result	size
default	$\frac{a^2(\sec(fx+e)-1)(5\cos(fx+e)-3)\sqrt{-c(\sec(fx+e)-1)}c\sqrt{a(\sec(fx+e)+1)}(\cos(fx+e)+1)^3\sec(fx+e)^2\csc(fx+e)}{12f(\cos(fx+e)-1)}$	87
risch	$\frac{2ia^2c\sqrt{\frac{a(e^{i(fx+e)}+1)}{1+e^{2i(fx+e)}}}}{\sqrt{\frac{c(e^{i(fx+e)}-1)}{1+e^{2i(fx+e)}}}}(3e^{7i(fx+e)}+3e^{6i(fx+e)}+5e^{5i(fx+e)}+5e^{3i(fx+e)}+3e^{2i(fx+e)}+3e^{i(fx+e)})}{3(1+e^{2i(fx+e)})^3(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)}f$	177

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

3.125.
$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{3/2} dx$$

output $-1/12/f*a^2*(\sec(f*x+e)-1)*(5*\cos(f*x+e)-3)*(-c*(\sec(f*x+e)-1))^{(1/2)}*c*(a*(\sec(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)+1)^3/(\cos(f*x+e)-1)*\sec(f*x+e)^2*csc(f*x+e)$

3.125.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.26

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{3/2} dx = \frac{(12 a^2 c \cos(fx + e)^3 + 6 a^2 c \cos(fx + e)^2 - 4 a^2 c \cos(fx + e) - 3 a^2 c) \sqrt{\frac{a \cos(fx + e) + 1}{\cos(fx + e)}}}{12 f \cos(fx + e)^3 \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(3/2),x, algorithm="fricas")`

output $1/12*(12*a^2*c*\cos(f*x + e)^3 + 6*a^2*c*\cos(f*x + e)^2 - 4*a^2*c*\cos(f*x + e) - 3*a^2*c)*\sqrt{(a*\cos(f*x + e) + a)/\cos(f*x + e)}*\sqrt{(c*\cos(f*x + e) - c)/\cos(f*x + e)}/(f*\cos(f*x + e)^3*\sin(f*x + e))$

3.125.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{3/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)*(c-c*sec(f*x+e))**(3/2),x)`

output `Timed out`

3.125.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1106 vs. $2(77) = 154$.

Time = 0.39 (sec) , antiderivative size = 1106, normalized size of antiderivative = 12.43

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{3/2} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(3/2),x, algo
rithm="maxima")
```

```
output 2/3*(20*a^2*c*cos(3*f*x + 3*e)*sin(2*f*x + 2*e) - 12*a^2*c*cos(2*f*x + 2*e)
)*sin(f*x + e) - 3*a^2*c*sin(f*x + e) - (3*a^2*c*sin(7*f*x + 7*e) + 3*a^2*
c*sin(6*f*x + 6*e) + 5*a^2*c*sin(5*f*x + 5*e) + 5*a^2*c*sin(3*f*x + 3*e) +
3*a^2*c*sin(2*f*x + 2*e) + 3*a^2*c*sin(f*x + e))*cos(8*f*x + 8*e) + 6*(2*
a^2*c*sin(6*f*x + 6*e) + 3*a^2*c*sin(4*f*x + 4*e) + 2*a^2*c*sin(2*f*x + 2*
e))*cos(7*f*x + 7*e) - 2*(10*a^2*c*sin(5*f*x + 5*e) - 9*a^2*c*sin(4*f*x +
4*e) + 10*a^2*c*sin(3*f*x + 3*e) + 6*a^2*c*sin(f*x + e))*cos(6*f*x + 6*e)
+ 10*(3*a^2*c*sin(4*f*x + 4*e) + 2*a^2*c*sin(2*f*x + 2*e))*cos(5*f*x + 5*e)
- 6*(5*a^2*c*sin(3*f*x + 3*e) + 3*a^2*c*sin(2*f*x + 2*e) + 3*a^2*c*sin(f
*x + e))*cos(4*f*x + 4*e) + (3*a^2*c*cos(7*f*x + 7*e) + 3*a^2*c*cos(6*f*x
+ 6*e) + 5*a^2*c*cos(5*f*x + 5*e) + 5*a^2*c*cos(3*f*x + 3*e) + 3*a^2*c*cos
(2*f*x + 2*e) + 3*a^2*c*cos(f*x + e))*sin(8*f*x + 8*e) - 3*(4*a^2*c*cos(6*
f*x + 6*e) + 6*a^2*c*cos(4*f*x + 4*e) + 4*a^2*c*cos(2*f*x + 2*e) + a^2*c)*
sin(7*f*x + 7*e) + (20*a^2*c*cos(5*f*x + 5*e) - 18*a^2*c*cos(4*f*x + 4*e)
+ 20*a^2*c*cos(3*f*x + 3*e) + 12*a^2*c*cos(f*x + e) - 3*a^2*c)*sin(6*f*x +
6*e) - 5*(6*a^2*c*cos(4*f*x + 4*e) + 4*a^2*c*cos(2*f*x + 2*e) + a^2*c)*si
n(5*f*x + 5*e) + 6*(5*a^2*c*cos(3*f*x + 3*e) + 3*a^2*c*cos(2*f*x + 2*e) +
3*a^2*c*cos(f*x + e))*sin(4*f*x + 4*e) - 5*(4*a^2*c*cos(2*f*x + 2*e) + a^2
*c)*sin(3*f*x + 3*e) + 3*(4*a^2*c*cos(f*x + e) - a^2*c)*sin(2*f*x + 2*e))*
sqrt(a)*sqrt(c)/((2*(4*cos(6*f*x + 6*e) + 6*cos(4*f*x + 4*e) + 4*cos(2*...
```

3.125.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{3/2} dx = \int (a \sec(fx + e) + a)^{\frac{5}{2}}(-c \sec(fx + e) + c)^{\frac{3}{2}} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(3/2),x, algo
rithm="giac")`

output `sage0*x`

3.125.9 Mupad [B] (verification not implemented)

Time = 16.85 (sec) , antiderivative size = 195, normalized size of antiderivative = 2.19

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{3/2} dx = \frac{\sqrt{c - \frac{c}{\cos(e + fx)}} \left(\frac{a^2 c \cos(e + fx) e^{e 4i + f x 4i} \sqrt{a + \frac{a}{\cos(e + fx)}} 20i}{3 f} + \frac{a^2 c e^{e 4i + f x 4i} \cos(2e + 2fx) \sqrt{a + \frac{a}{\cos(e + fx)}}}{f} \right)}{e^{e 4i + f x 4i} \sin(2e + 2fx) 4i + e^{e 4i + f x 4i} \sin(4e + 4fx) 2i}$$

input `int(((a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(3/2))/cos(e + f*x),x
)`

output `((c - c/cos(e + f*x))^(1/2)*((a^2*c*cos(e + f*x)*exp(e*4i + f*x*4i))*(a + a
/cos(e + f*x))^(1/2)*20i)/(3*f) + (a^2*c*exp(e*4i + f*x*4i)*cos(2*e + 2*f*
x)*(a + a/cos(e + f*x))^(1/2)*4i)/f + (a^2*c*exp(e*4i + f*x*4i)*cos(3*e +
3*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/f)/(exp(e*4i + f*x*4i)*sin(2*e + 2*
f*x)*4i + exp(e*4i + f*x*4i)*sin(4*e + 4*f*x)*2i)`

3.126 $\int \sec(e+fx)(a+a \sec(e+fx))^{5/2} \sqrt{c - c \sec(e + fx)}$

3.126.1 Optimal result	909
3.126.2 Mathematica [A] (verified)	909
3.126.3 Rubi [A] (verified)	910
3.126.4 Maple [A] (verified)	911
3.126.5 Fricas [B] (verification not implemented)	911
3.126.6 Sympy [F(-1)]	912
3.126.7 Maxima [A] (verification not implemented)	912
3.126.8 Giac [F]	912
3.126.9 Mupad [B] (verification not implemented)	913

3.126.1 Optimal result

Integrand size = 36, antiderivative size = 43

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)} dx = \frac{c(a + a \sec(e + fx))^{5/2} \tan(e + fx)}{3f \sqrt{c - c \sec(e + fx)}}$$

```
output -1/3*c*(a+a*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(1/2)
```

3.126.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.63

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)} dx = \frac{a^3 c \sec(e + fx) (3 + 3 \sec(e + fx) + \sec^2(e + fx)) \tan(e + fx)}{3f \sqrt{a(1 + \sec(e + fx))} \sqrt{c - c \sec(e + fx)}}$$

```
input Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*Sqrt[c - c*Sec[e + f*x]],x]
```

```
output -1/3*(a^3*c*Sec[e + f*x]*(3 + 3*Sec[e + f*x] + Sec[e + f*x]^2)*Tan[e + f*x])/
(f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])
```

3.126.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^{5/2} \sqrt{c - c \sec(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{5/2} \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4441}$$

$$-\frac{c \tan(e + fx)(a \sec(e + fx) + a)^{5/2}}{3f \sqrt{c - c \sec(e + fx)}}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2)*Sqrt[c - c*Sec[e + f*x]],x]`

output `-1/3*(c*(a + a*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.126.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.126.4 Maple [A] (verified)

Time = 3.79 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.35

method	result	size
default	$\frac{a^2(\cos(fx+e)+1)^3 \sqrt{a(\sec(fx+e)+1)} \sqrt{-c(\sec(fx+e)-1)} \sec(fx+e)^2 \csc(fx+e)}{3f}$	58
risch	$\frac{2ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} (3e^{5i(fx+e)}+6e^{4i(fx+e)}+10e^{3i(fx+e)}+6e^{2i(fx+e)}+3e^{i(fx+e)})}{3(1+e^{2i(fx+e)})^2(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)f}$	165

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 1/3/f*a^2*(cos(f*x+e)+1)^3*(a*(sec(f*x+e)+1))^(1/2)*(-c*(sec(f*x+e)-1))^(1
/2)*sec(f*x+e)^2*csc(f*x+e)
```

3.126.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 93 vs. 2(37) = 74.

Time = 0.26 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.16

$$\int \sec(e+fx)(a+a\sec(e+fx))^{5/2} \sqrt{c-c\sec(e+fx)} dx = \frac{(3a^2 \cos(fx+e)^2 + 3a^2 \cos(fx+e) + a^2) \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{c \cos(fx+e)-c}{\cos(fx+e)}}}{3f \cos(fx+e)^2 \sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(1/2),x, algo
rithm="fracas")
```

```
output 1/3*(3*a^2*cos(f*x + e)^2 + 3*a^2*cos(f*x + e) + a^2)*sqrt((a*cos(f*x + e)
+ a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(f*cos(f*x + e)
)^2*sin(f*x + e)
```


3.126.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)*(c-c*sec(f*x+e))**(1/2),x)`

output `Timed out`

3.126.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.35

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)} dx = \frac{8 \sqrt{-aa^2} \sqrt{c}}{3 f \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right)^3 \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)^3}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `8/3*sqrt(-a)*a^2*sqrt(c)/(f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^3*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^3)`

3.126.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)} dx = \int (a \sec(fx + e) + a)^{5/2} \sqrt{-c \sec(fx + e) + c \sec(fx + e)} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)*(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `sage0*x`

3.126. $\int \sec(e + fx)(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)} dx$

3.126.9 Mupad [B] (verification not implemented)

Time = 14.98 (sec) , antiderivative size = 136, normalized size of antiderivative = 3.16

$$\int \sec(e + fx)(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)} dx = \frac{2a^2 \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}} \sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}} (10 \sin(e + fx) + 12 \sin(2e + 2fx) + 13 \sin(3e + 3fx) + 6 \sin(4e + 4fx) + 3 \sin(5e + 5fx))}{3f (\cos(2e + 2fx) - 2 \cos(4e + 4fx) + \cos(6e + 6fx) + 2)}$$

input `int(((a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x),x)`

output `(2*a^2*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2)*((c*(cos(e + f*x) - 1))/cos(e + f*x))^(1/2)*(10*sin(e + f*x) + 12*sin(2*e + 2*f*x) + 13*sin(3*e + 3*f*x) + 6*sin(4*e + 4*f*x) + 3*sin(5*e + 5*f*x)))/(3*f*(cos(2*e + 2*f*x) - 2*cos(4*e + 4*f*x) - cos(6*e + 6*f*x) + 2))`

3.127
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{\sqrt{c-c \sec(e+fx)}} dx$$

3.127.1 Optimal result 914
 3.127.2 Mathematica [A] (verified) 914
 3.127.3 Rubi [A] (verified) 915
 3.127.4 Maple [A] (verified) 917
 3.127.5 Fracas [F] 917
 3.127.6 Sympy [F(-1)] 918
 3.127.7 Maxima [B] (verification not implemented) 918
 3.127.8 Giac [F] 919
 3.127.9 Mupad [F(-1)] 920

3.127.1 Optimal result

Integrand size = 36, antiderivative size = 141

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{\sqrt{c-c \sec(e+fx)}} dx = \frac{4a^3 \log(1-\sec(e+fx)) \tan(e+fx)}{f \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} + \frac{2a^2 \sqrt{a+a \sec(e+fx)} \tan(e+fx)}{f \sqrt{c-c \sec(e+fx)}} + \frac{a(a+a \sec(e+fx))^{3/2} \tan(e+fx)}{2f \sqrt{c-c \sec(e+fx)}}$$

output $\frac{1}{2} a (a + a \sec(fx + e))^{3/2} \tan(fx + e) / (c - c \sec(fx + e))^{1/2} + 4 a^3 \ln(1 - \sec(fx + e)) \tan(fx + e) / (a + a \sec(fx + e))^{1/2} / (c - c \sec(fx + e))^{1/2} + 2 a^2 \sqrt{a + a \sec(fx + e)} \tan(fx + e) / (c - c \sec(fx + e))^{1/2}$

3.127.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.54

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{\sqrt{c-c \sec(e+fx)}} dx = \frac{a^3(1+8 \log(1-\sec(e+fx))+6 \sec(e+fx)+\sec^2(e+fx)) \tan(e+fx)}{2f \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/Sqrt[c - c*Sec[e + f*x]],x]`

output $(a^3(1+8 \log[1-\sec[e+fx]]+6 \sec[e+fx]+\sec[e+fx]^2) \tan[e+fx]) / (2 f \sqrt{a(1+\sec[e+fx])} \sqrt{c-c \sec[e+fx]})$

3.127.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{\sqrt{c-c \sec(e+fx)}} dx$$

3.127.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4443, 3042, 4443, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^{5/2}}{\sqrt{c-c\sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^{5/2}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4443} \\
 & 2a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx + \frac{a \tan(e+fx)(a\sec(e+fx)+a)^{3/2}}{2f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & 2a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^{3/2}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{a \tan(e+fx)(a\sec(e+fx)+a)^{3/2}}{2f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{4443} \\
 & 2a \left(2a \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{\sqrt{c-c\sec(e+fx)}} dx + \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}} \right) + \\
 & \quad \frac{a \tan(e+fx)(a\sec(e+fx)+a)^{3/2}}{2f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & 2a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{a \tan(e+fx)\sqrt{a\sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}} \right) + \\
 & \quad \frac{a \tan(e+fx)(a\sec(e+fx)+a)^{3/2}}{2f\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{4440}
 \end{aligned}$$

$$2a \left(\frac{2a^2 \tan(e+fx) \log(1 - \sec(e+fx))}{f \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} + \frac{a \tan(e+fx) \sqrt{a \sec(e+fx) + a}}{f \sqrt{c - c \sec(e+fx)}} \right) + \frac{a \tan(e+fx) (a \sec(e+fx) + a)^{3/2}}{2f \sqrt{c - c \sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/Sqrt[c - c*Sec[e + f*x]],x]`

output `(a*(a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(2*f*Sqrt[c - c*Sec[e + f*x]]) + 2*a*((2*a^2*Log[1 - Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]])*Sqrt[c - c*Sec[e + f*x]]) + (a*Sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]])`

3.127.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.127.4 Maple [A] (verified)

Time = 3.61 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.05

method	result
default	$\frac{a^2 \sqrt{a(\sec(fx+e)+1)} (16 \ln(-\cot(fx+e)+\csc(fx+e)) \sin(fx+e) - 8 \ln(-\cot(fx+e)+\csc(fx+e)-1) \sin(fx+e) - 8 \ln(-\cot(fx+e)+\csc(fx+e)+1) \sin(fx+e) - 8 \ln(-\cot(fx+e)+\csc(fx+e)-1) \sin(fx+e))}{2f(\cos(fx+e)+1)\sqrt{-c(\sec(fx+e)-1)}}$
risch	$-\frac{2ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (3e^{2i(fx+e)}+e^{i(fx+e)}+3)(e^{2i(fx+e)}-e^{i(fx+e)})}{(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f(1+e^{2i(fx+e)})^2} - \frac{8ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1) \ln(e^{i(fx+e)}-1)}{(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x,method=_RET
URNVERBOSE)`

output `1/2/f*a^2*(a*(sec(f*x+e)+1))^(1/2)/(cos(f*x+e)+1)/(-c*(sec(f*x+e)-1))^(1/2
)*(16*ln(-cot(f*x+e)+csc(f*x+e))*sin(f*x+e)-8*ln(-cot(f*x+e)+csc(f*x+e)-1)
*sin(f*x+e)-8*ln(-cot(f*x+e)+csc(f*x+e)+1)*sin(f*x+e)+5*sin(f*x+e)+6*tan(f
*x+e)+sec(f*x+e)*tan(f*x+e))`

3.127.5 Fracas [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{(a\sec(fx+e)+a)^{5/2} \sec(fx+e)}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="fracas")`

output `integral(-(a^2*sec(f*x + e)^3 + 2*a^2*sec(f*x + e)^2 + a^2*sec(f*x + e))*s
qrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(c*sec(f*x + e) - c), x)`

3.127.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{\sqrt{c - c \sec(e + fx)}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(1/2),x)`

output `Timed out`

3.127.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 737 vs. 2(127) = 254.

Time = 0.41 (sec) , antiderivative size = 737, normalized size of antiderivative = 5.23

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{\sqrt{c - c \sec(e + fx)}} dx =$$

$$\frac{2(a^2 \cos(2fx + 2e) \sin(4fx + 4e) - a^2 \cos(4fx + 4e) \sin(2fx + 2e) - a^2 \sin(2fx + 2e) + 2(a^2 \cos($$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="maxima")`

output

```

-2*(a^2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - a^2*cos(4*f*x + 4*e)*sin(2*f*x
+ 2*e) - a^2*sin(2*f*x + 2*e) + 2*(a^2*cos(4*f*x + 4*e)^2 + 4*a^2*cos(2*f
*x + 2*e)^2 + a^2*sin(4*f*x + 4*e)^2 + 4*a^2*sin(4*f*x + 4*e)*sin(2*f*x +
2*e) + 4*a^2*sin(2*f*x + 2*e)^2 + 4*a^2*cos(2*f*x + 2*e) + a^2 + 2*(2*a^2*
cos(2*f*x + 2*e) + a^2)*cos(4*f*x + 4*e))*arctan2(sin(2*f*x + 2*e), cos(2*
f*x + 2*e) + 1) - 4*(a^2*cos(4*f*x + 4*e)^2 + 4*a^2*cos(2*f*x + 2*e)^2 + a
^2*sin(4*f*x + 4*e)^2 + 4*a^2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*a^2*si
n(2*f*x + 2*e)^2 + 4*a^2*cos(2*f*x + 2*e) + a^2 + 2*(2*a^2*cos(2*f*x + 2*e
) + a^2)*cos(4*f*x + 4*e))*arctan2(sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2
*f*x + 2*e))), cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - 1) +
3*(a^2*sin(4*f*x + 4*e) + 2*a^2*sin(2*f*x + 2*e))*cos(3/2*arctan2(sin(2*f
*x + 2*e), cos(2*f*x + 2*e))) + 3*(a^2*sin(4*f*x + 4*e) + 2*a^2*sin(2*f*x
+ 2*e))*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 3*(a^2*cos(
4*f*x + 4*e) + 2*a^2*cos(2*f*x + 2*e) + a^2)*sin(3/2*arctan2(sin(2*f*x + 2
*e), cos(2*f*x + 2*e))) - 3*(a^2*cos(4*f*x + 4*e) + 2*a^2*cos(2*f*x + 2*e)
+ a^2)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)*sqrt
(c)/((c*cos(4*f*x + 4*e)^2 + 4*c*cos(2*f*x + 2*e)^2 + c*sin(4*f*x + 4*e)^2
+ 4*c*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*c*sin(2*f*x + 2*e)^2 + 2*(2*c
*cos(2*f*x + 2*e) + c)*cos(4*f*x + 4*e) + 4*c*cos(2*f*x + 2*e) + c)*f)

```

3.127.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{(a\sec(fx+e)+a)^{5/2}\sec(fx+e)}{\sqrt{-c\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="giac")`

output `sage0*x`

3.127.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^{5/2}}{\cos(e+fx) \sqrt{c - \frac{c}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

3.128
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{3/2}} dx$$

3.128.1 Optimal result 921
 3.128.2 Mathematica [A] (verified) 921
 3.128.3 Rubi [A] (verified) 922
 3.128.4 Maple [A] (verified) 924
 3.128.5 Fricas [F] 924
 3.128.6 Sympy [F(-1)] 925
 3.128.7 Maxima [B] (verification not implemented) 925
 3.128.8 Giac [F] 926
 3.128.9 Mupad [F(-1)] 927

3.128.1 Optimal result

Integrand size = 36, antiderivative size = 145

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{3/2}} dx = -\frac{a(a+a \sec(e+fx))^{3/2} \tan(e+fx)}{f(c-c \sec(e+fx))^{3/2}} - \frac{4a^3 \log(1-\sec(e+fx)) \tan(e+fx)}{cf \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} - \frac{2a^2 \sqrt{a+a \sec(e+fx)} \tan(e+fx)}{cf \sqrt{c-c \sec(e+fx)}}$$

output

```
-a*(a+a*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(3/2)-4*a^3*ln(1-sec(f*x+e))*tan(f*x+e)/c/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-2*a^2*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(1/2)
```

3.128.2 Mathematica [A] (verified)

Time = 0.72 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.54

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{3/2}} dx = \frac{a^3 \left(4 \log(1-\sec(e+fx)) - \frac{4}{-1+\sec(e+fx)} + \sec(e+fx) \right) \tan(e+fx)}{cf \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input

```
Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(3/2),x]
```

3.128.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{3/2}} dx$$

output $-\left(\frac{a^3(4\log[1 - \sec[e + fx]] - 4/(-1 + \sec[e + fx]) + \sec[e + fx])\tan[e + fx]}{c f \sqrt{a(1 + \sec[e + fx])}} \sqrt{c - c \sec[e + fx]}\right)$

3.128.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4442, 3042, 4443, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(a \sec(e + fx) + a)^{5/2}}{(c - c \sec(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^{5/2}}{\left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2}} dx$$

↓ 4442

$$\frac{2a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^{3/2}}{\sqrt{c-c\sec(e+fx)}} dx}{c} - \frac{a \tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{f(c - c \sec(e + fx))^{3/2}}$$

↓ 3042

$$\frac{2a \int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^{3/2}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx}{c} - \frac{a \tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{f(c - c \sec(e + fx))^{3/2}}$$

↓ 4443

$$\frac{2a \left(2a \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{\sqrt{c-c\sec(e+fx)}} dx + \frac{a \tan(e+fx)\sqrt{a \sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}} \right)}{c} - \frac{a \tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{f(c - c \sec(e + fx))^{3/2}}$$

↓ 3042

$$\frac{2a \left(2a \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx + \frac{a \tan(e+fx)\sqrt{a \sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}} \right)}{c} - \frac{a \tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{f(c - c \sec(e + fx))^{3/2}}$$

3.128. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{3/2}} dx$

$$\begin{array}{c} \downarrow 4440 \\ 2a \left(\frac{2a^2 \tan(e+fx) \log(1-\sec(e+fx))}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} + \frac{a \tan(e+fx) \sqrt{a\sec(e+fx)+a}}{f\sqrt{c-c\sec(e+fx)}} \right) \\ \hline \frac{c}{a \tan(e+fx)(a\sec(e+fx)+a)^{3/2}} \\ \frac{f(c-c\sec(e+fx))^{3/2}}{f(c-c\sec(e+fx))^{3/2}} \end{array}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(3/2),x]`

output `-((a*(a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2))) - (2*a*((2*a^2*Log[1 - Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]) + (a*Sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[c - c*Sec[e + f*x]]))))/c`

3.128.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.128.4 Maple [A] (verified)

Time = 3.62 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.62

method	result
default	$-\frac{a^2 \sqrt{a(\sec(fx+e)+1)} (4 \ln(-\cot(fx+e)+\csc(fx+e)-1) \sin(fx+e)+4 \ln(-\cot(fx+e)+\csc(fx+e)+1) \sin(fx+e)-8 \ln(-\cot(fx+e)+\csc(fx+e)-1) \tan(fx+e)+8 \ln(-\cot(fx+e)+\csc(fx+e)+1) \tan(fx+e)-4 \ln(-\cot(fx+e)+\csc(fx+e)-1) \operatorname{atanh}(\frac{\sin(fx+e)}{\csc(fx+e)+1})+4 \ln(-\cot(fx+e)+\csc(fx+e)+1) \operatorname{atanh}(\frac{\sin(fx+e)}{\csc(fx+e)+1}))}{c^2 \sqrt{a(\sec(fx+e)+1)}}$
risch	$\frac{2ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (5e^{3i(fx+e)}-2e^{2i(fx+e)}+5e^{i(fx+e)})}{c(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f(1+e^{2i(fx+e)})} + \frac{8ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1) \ln(e^{i(fx+e)}-1)}{c(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f} - \frac{4ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}}{c(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2),x,method=_RET
URNVERBOSE)
```

```
output -1/f*a^2*(a*(sec(f*x+e)+1))^(1/2)/(sec(f*x+e)-1)/(-c*(sec(f*x+e)-1))^(1/2)
/c/(cos(f*x+e)+1)*(4*ln(-cot(f*x+e)+csc(f*x+e)-1)*sin(f*x+e)+4*ln(-cot(f*x
+e)+csc(f*x+e)+1)*sin(f*x+e)-8*ln(-cot(f*x+e)+csc(f*x+e))*sin(f*x+e)-4*ln(
-cot(f*x+e)+csc(f*x+e)-1)*tan(f*x+e)-4*ln(-cot(f*x+e)+csc(f*x+e)+1)*tan(f*
x+e)+8*ln(-cot(f*x+e)+csc(f*x+e))*tan(f*x+e)-3*sin(f*x+e)-2*tan(f*x+e)+sec
(f*x+e)*tan(f*x+e))
```

3.128.5 Fricas [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{(a\sec(fx+e)+a)^{5/2} \sec(fx+e)}{(-c\sec(fx+e)+c)^{3/2}} dx$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="fricas")
```

3.128. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{3/2}} dx$

output `integral((a^2*sec(f*x + e)^3 + 2*a^2*sec(f*x + e)^2 + a^2*sec(f*x + e))*sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(c^2*sec(f*x + e)^2 - 2*c^2*sec(f*x + e) + c^2), x)`

3.128.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(3/2),x)`

output `Timed out`

3.128.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2035 vs. 2(133) = 266.

Time = 0.51 (sec) , antiderivative size = 2035, normalized size of antiderivative = 14.03

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output

```

2*(8*a^2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)*cos(3/2*arctan2(s
in(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 8*a^2*arctan2(sin(2*f*x + 2*e), co
s(2*f*x + 2*e) + 1)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2
+ 8*a^2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)*sin(3/2*arctan2(s
in(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 8*a^2*arctan2(sin(2*f*x + 2*e), co
s(2*f*x + 2*e) + 1)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2
- 2*a^2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) + 2*a^2*cos(4*f*x + 4*e)*sin(2*
f*x + 2*e) + 2*a^2*sin(2*f*x + 2*e) + 2*(a^2*cos(4*f*x + 4*e)^2 + 4*a^2*co
s(2*f*x + 2*e)^2 + a^2*sin(4*f*x + 4*e)^2 + 4*a^2*sin(4*f*x + 4*e)*sin(2*f
*x + 2*e) + 4*a^2*sin(2*f*x + 2*e)^2 + 4*a^2*cos(2*f*x + 2*e) + a^2 + 2*(2
*a^2*cos(2*f*x + 2*e) + a^2)*cos(4*f*x + 4*e))*arctan2(sin(2*f*x + 2*e), c
os(2*f*x + 2*e) + 1) - 4*(a^2*cos(4*f*x + 4*e)^2 + 4*a^2*cos(2*f*x + 2*e)^
2 + 4*a^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 4*a^2*c
os(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + a^2*sin(4*f*x + 4*
e)^2 + 4*a^2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*a^2*sin(2*f*x + 2*e)^2
+ 4*a^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 4*a^2*sin
(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 4*a^2*cos(2*f*x + 2*
e) + a^2 + 2*(2*a^2*cos(2*f*x + 2*e) + a^2)*cos(4*f*x + 4*e) - 4*(a^2*cos(
4*f*x + 4*e) + 2*a^2*cos(2*f*x + 2*e) - 2*a^2*cos(1/2*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e)))) + a^2)*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2...

```

3.128.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{(a\sec(fx+e)+a)^{5/2}\sec(fx+e)}{(-c\sec(fx+e)+c)^{3/2}} dx$$

input

```

integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="giac")

```

output

```

sage0*x

```

3.128.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^{5/2}}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),x)`

output `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),x)`

3.129
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{5/2}} dx$$

3.129.1 Optimal result	928
3.129.2 Mathematica [A] (verified)	928
3.129.3 Rubi [A] (verified)	929
3.129.4 Maple [B] (warning: unable to verify)	931
3.129.5 Fricas [F]	932
3.129.6 Sympy [F(-1)]	932
3.129.7 Maxima [A] (verification not implemented)	932
3.129.8 Giac [F]	933
3.129.9 Mupad [F(-1)]	933

3.129.1 Optimal result

Integrand size = 36, antiderivative size = 145

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{5/2}} dx = -\frac{a(a+a \sec(e+fx))^{3/2} \tan(e+fx)}{2f(c-c \sec(e+fx))^{5/2}} + \frac{a^2 \sqrt{a+a \sec(e+fx)} \tan(e+fx)}{cf(c-c \sec(e+fx))^{3/2}} + \frac{a^3 \log(1-\sec(e+fx)) \tan(e+fx)}{c^2 f \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}}$$

output `-1/2*a*(a+a*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(5/2)+a^2*(a+a*sec(f*x+e))^(1/2)*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(3/2)+a^3*ln(1-sec(f*x+e))*tan(f*x+e)/c^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.129.2 Mathematica [A] (verified)

Time = 1.30 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.56

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{5/2}} dx = \frac{a^3 \left(-\log(1-\sec(e+fx)) + \frac{-2+4 \sec(e+fx)}{(-1+\sec(e+fx))^2} \right) \tan(e+fx)}{c^2 f \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(5/2),x]`

output $-\left(\left(a^3(-\text{Log}[1 - \text{Sec}[e + f*x]] + (-2 + 4*\text{Sec}[e + f*x])/(-1 + \text{Sec}[e + f*x])^2)*\text{Tan}[e + f*x]\right)/\left(c^2*f*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]\right)\right)$

3.129.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4442, 3042, 4442, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a \sec(e+fx) + a)^{5/2}}{(c - c \sec(e+fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(a \csc\left(e+fx+\frac{\pi}{2}\right) + a\right)^{5/2}}{\left(c - c \csc\left(e+fx+\frac{\pi}{2}\right)\right)^{5/2}} dx$$

↓ 4442

$$-\frac{a \int \frac{\sec(e+fx)(\sec(e+fx)a+a)^{3/2}}{(c-c \sec(e+fx))^{3/2}} dx}{c} - \frac{a \tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{2f(c - c \sec(e+fx))^{5/2}}$$

↓ 3042

$$-\frac{a \int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)^{3/2}}{\left(c - c \csc\left(e+fx+\frac{\pi}{2}\right)\right)^{3/2}} dx}{c} - \frac{a \tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{2f(c - c \sec(e+fx))^{5/2}}$$

↓ 4442

$$-\frac{a \left(-\frac{a \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{\sqrt{c-c \sec(e+fx)}} dx}{c} - \frac{a \tan(e+fx)\sqrt{a \sec(e+fx)+a}}{f(c-c \sec(e+fx))^{3/2}} \right)}{c} - \frac{a \tan(e+fx)(a \sec(e+fx) + a)^{3/2}}{2f(c - c \sec(e+fx))^{5/2}}$$

↓ 3042

$$\begin{aligned}
 & a \left(\frac{a \int \frac{\csc(e+fx+\frac{\pi}{2}) \sqrt{\csc(e+fx+\frac{\pi}{2}) a+a}}{\sqrt{c-c \csc(e+fx+\frac{\pi}{2})}} dx}{c} - \frac{a \tan(e+fx) \sqrt{a \sec(e+fx)+a}}{f(c-c \sec(e+fx))^{3/2}} \right) \\
 & \frac{c}{2f(c-c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow 4440 \\
 & a \left(-\frac{a^2 \tan(e+fx) \log(1-\sec(e+fx))}{cf \sqrt{a \sec(e+fx)+a} \sqrt{c-c \sec(e+fx)}} - \frac{a \tan(e+fx) \sqrt{a \sec(e+fx)+a}}{f(c-c \sec(e+fx))^{3/2}} \right) \\
 & \frac{c}{2f(c-c \sec(e+fx))^{5/2}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/2*(a*(a + a*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(5/2)) - (a*(-((a*Sqrt[a + a*Sec[e + f*x]]*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(3/2)))) - (a^2*Log[1 - Sec[e + f*x]]*Tan[e + f*x])/(c*f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])))/c`

3.129.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

```
rule 4442 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]
```

3.129.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 271 vs. 2(131) = 262.

Time = 3.60 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.88

method	result
default	$-\frac{\sqrt{2}a^2\sqrt{-\frac{2a}{(1-\cos(fx+e))^2\csc(fx+e)^2-1}}(1-\cos(fx+e))(2\ln(-\cot(fx+e)+\csc(fx+e)+1)(1-\cos(fx+e))^4\csc(fx+e)^4+2\ln(-\cot(fx+e)+\csc(fx+e)+1))}{4f((1-\cos(fx+e))^2)}$
risch	$\frac{8ia^2\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}e^{2i(fx+e)}}{c^2(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)^3\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f} - \frac{2ia^2\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)\ln(e^{i(fx+e)}-1)}}{c^2(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}} + \frac{ia^2\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}}{c^2(e^{i(fx+e)}+1)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/4/f*2^(1/2)*a^2*(-2*a/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^2/(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*csc(f*x+e)^2)^(5/2)*(1-cos(f*x+e))*(2*ln(-cot(f*x+e)+csc(f*x+e)+1)*(1-cos(f*x+e))^4*csc(f*x+e)^4+2*ln(-cot(f*x+e)+csc(f*x+e)-1)*(1-cos(f*x+e))^4*csc(f*x+e)^4-4*ln(-cot(f*x+e)+csc(f*x+e))*(1-cos(f*x+e))^4*csc(f*x+e)^4+2*(1-cos(f*x+e))^2*csc(f*x+e)^2+1)*csc(f*x+e)
```

$$3.129. \int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{5/2}} dx$$

3.129.5 Fricas [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{5/2}} dx = \int \frac{(a \sec(fx + e) + a)^{5/2} \sec(fx + e)}{(-c \sec(fx + e) + c)^{5/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="fricas")`

output `integral(-(a^2*sec(f*x + e)^3 + 2*a^2*sec(f*x + e)^2 + a^2*sec(f*x + e))*sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(c^3*sec(f*x + e)^3 - 3*c^3*sec(f*x + e)^2 + 3*c^3*sec(f*x + e) - c^3), x)`

3.129.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(5/2),x)`

output `Timed out`

3.129.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.17

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{5/2}} dx = \frac{\frac{2\sqrt{-aa^2} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{c^{5/2}} + \frac{2\sqrt{-aa^2} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{c^{5/2}} - \frac{4\sqrt{-aa^2} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)}{c^{5/2}} + \frac{\left(\sqrt{-aa^2}\sqrt{c} + \frac{2\sqrt{-aa^2}\sqrt{c}\sin(fx+e)^2}{(\cos(fx+e)+1)^2}\right)(\cos(fx+e))}{c^3 \sin(fx+e)^4}}{2f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="maxima")`

3.129. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{5/2}} dx$

output
$$-1/2*(2*\sqrt{-a}*a^2*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/c^{(5/2)} + 2*\sqrt{-a}*a^2*\log(\sin(f*x + e)/(\cos(f*x + e) - 1)/c^{(5/2)} - 4*\sqrt{-a}*a^2*\log(\sin(f*x + e)/(\cos(f*x + e) + 1))/c^{(5/2)} + (\sqrt{-a}*a^2*\sqrt{c} + 2*\sqrt{-a}*a^2*\sqrt{c}*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2*(\cos(f*x + e) + 1)^4/(c^3*\sin(f*x + e)^4))/f$$

3.129.8 Giac [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{5/2}} dx = \int \frac{(a \sec(fx + e) + a)^{5/2} \sec(fx + e)}{(-c \sec(fx + e) + c)^{5/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `sage0*x`

3.129.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{5/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e + fx)}\right)^{5/2}}{\cos(e + fx) \left(c - \frac{c}{\cos(e + fx)}\right)^{5/2}} dx$$

input `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x)`

output `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x)`

$$3.130 \quad \int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx$$

3.130.1 Optimal result	934
3.130.2 Mathematica [A] (verified)	934
3.130.3 Rubi [A] (verified)	935
3.130.4 Maple [A] (verified)	936
3.130.5 Fricas [B] (verification not implemented)	936
3.130.6 Sympy [F(-1)]	937
3.130.7 Maxima [B] (verification not implemented)	937
3.130.8 Giac [F]	938
3.130.9 Mupad [B] (verification not implemented)	939

3.130.1 Optimal result

Integrand size = 36, antiderivative size = 42

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx = -\frac{(a+a\sec(e+fx))^{5/2}\tan(e+fx)}{6f(c-c\sec(e+fx))^{7/2}}$$

output `-1/6*(a+a*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(7/2)`

3.130.2 Mathematica [A] (verified)

Time = 1.81 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx = -\frac{(a(1+\sec(e+fx)))^{5/2}\tan(e+fx)}{6f(c-c\sec(e+fx))^{7/2}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(7/2), x]`

output `-1/6*((a*(1 + Sec[e + f*x]))^(5/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(7/2))`

$$3.130. \quad \int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx$$

3.130.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^{5/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{7/2}} dx$$

↓ 4438

$$-\frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{6f(c-c\sec(e+fx))^{7/2}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(7/2),x]`

output `-1/6*((a + a*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(7/2))`

3.130.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.130.4 Maple [A] (verified)

Time = 3.61 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.69

method	result	size
default	$\frac{a^2(\cos(fx+e)+1)^2\sqrt{a(\sec(fx+e)+1)}\tan(fx+e)\sec(fx+e)^2}{6f(\sec(fx+e)-1)^3\sqrt{-c(\sec(fx+e)-1)}c^3}$	71
risch	$\frac{2ia^2\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(3e^{5i(fx+e)}+10e^{3i(fx+e)}+3e^{i(fx+e)})}{3c^3(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)^5\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}$	133

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(7/2),x,method=_RET
URNVERBOSE)
```

```
output 1/6/f*a^2*(cos(f*x+e)+1)^2*(a*(sec(f*x+e)+1))^(1/2)/(sec(f*x+e)-1)^3/(-c*(
sec(f*x+e)-1))^(1/2)/c^3*tan(f*x+e)*sec(f*x+e)^2
```

3.130.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 126 vs. 2(36) = 72.

Time = 0.27 (sec) , antiderivative size = 126, normalized size of antiderivative = 3.00

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx = \frac{(3a^2\cos(fx+e)^3+a^2\cos(fx+e))\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)+c}{\cos(fx+e)}}}{3(c^4f\cos(fx+e)^3-3c^4f\cos(fx+e)^2+3c^4f\cos(fx+e)-c^4f)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(7/2),x, algo
rithm="fracas")
```

```
output 1/3*(3*a^2*cos(f*x + e)^3 + a^2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/co
s(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((c^4*f*cos(f*x + e)^3
- 3*c^4*f*cos(f*x + e)^2 + 3*c^4*f*cos(f*x + e) - c^4*f)*sin(f*x + e))
```

3.130.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{7/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(7/2),x)`

output `Timed out`

3.130.7 Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 1815 vs. $2(36) = 72$.

Time = 0.41 (sec) , antiderivative size = 1815, normalized size of antiderivative = 43.21

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^{5/2}}{(c - c \sec(e + fx))^{7/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(7/2),x, algorithm="maxima")`

output

$$\begin{aligned} & \frac{2}{3}(208a^2\cos(3fx + 3e)\sin(2fx + 2e) + 48a^2\cos(fx + e)\sin(2fx + 2e) - 48a^2\cos(2fx + 2e)\sin(fx + e) - 3a^2\sin(fx + e) - \\ & (3a^2\sin(7fx + 7e) + 13a^2\sin(5fx + 5e) + 13a^2\sin(3fx + 3e) + 3a^2\sin(fx + e))\cos(8fx + 8e) + 6(8a^2\sin(6fx + 6e) + 15a^2\sin(4fx + 4e) + 8a^2\sin(2fx + 2e))\cos(7fx + 7e) - 16(13a^2\sin(5fx + 5e) + 13a^2\sin(3fx + 3e) + 3a^2\sin(fx + e))\cos(6fx + 6e) + 26(15a^2\sin(4fx + 4e) + 8a^2\sin(2fx + 2e))\cos(5fx + 5e) - 30(13a^2\sin(3fx + 3e) + 3a^2\sin(fx + e))\cos(4fx + 4e) + (3a^2\cos(7fx + 7e) + 13a^2\cos(5fx + 5e) + 13a^2\cos(3fx + 3e) + 3a^2\cos(fx + e))\sin(8fx + 8e) - 3(16a^2\cos(6fx + 6e) + 30a^2\cos(4fx + 4e) + 16a^2\cos(2fx + 2e) + a^2)\sin(7fx + 7e) + 16(13a^2\cos(5fx + 5e) + 13a^2\cos(3fx + 3e) + 3a^2\cos(fx + e))\sin(6fx + 6e) - 13(30a^2\cos(4fx + 4e) + 16a^2\cos(2fx + 2e) + a^2)\sin(5fx + 5e) + 30(13a^2\cos(3fx + 3e) + 3a^2\cos(fx + e))\sin(4fx + 4e) - 13(16a^2\cos(2fx + 2e) + a^2)\sin(3fx + 3e))\sqrt{a}\sqrt{c}/((c^4\cos(8fx + 8e))^2 + 36c^4\cos(7fx + 7e)^2 + 256c^4\cos(6fx + 6e)^2 + 676c^4\cos(5fx + 5e)^2 + 900c^4\cos(4fx + 4e)^2 + 676c^4\cos(3fx + 3e)^2 + 256c^4\cos(2fx + 2e)^2 + 36c^4\cos(fx + e)^2 + c^4\sin(8fx + 8e)^2 + 36c^4\sin(7fx + 7e)^2 + 256c^4\sin(6fx + 6e)^2 + 676c^4\sin(5fx + 5e)^2 + 900c^4\sin(4fx + 4e)^2 + 676c^4\sin(3fx + 3e)^2 + 256c^4\sin(2fx + 2e)^2 + 36c^4\sin(fx + e)^2) \end{aligned}$$

3.130.8 Giac [F]

$$\int \frac{\sec(e + fx)(a + a\sec(e + fx))^{5/2}}{(c - c\sec(e + fx))^{7/2}} dx = \int \frac{(a\sec(fx + e) + a)^{5/2} \sec(fx + e)}{(-c\sec(fx + e) + c)^{7/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(7/2),x, algo rithm="giac")`

output `sage0*x`

3.130.9 Mupad [B] (verification not implemented)

Time = 17.87 (sec) , antiderivative size = 199, normalized size of antiderivative = 4.74

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx =$$

$$\frac{\sqrt{c-\frac{c}{\cos(e+fx)}} \left(\frac{a^2 \cos(e+fx) e^{e4i+fx4i} \sqrt{a+\frac{a}{\cos(e+fx)}} 52i}{3c^4 f} + \frac{a^2 e^{e4i+fx4i} \cos(3e+3fx) \sqrt{a+\frac{a}{\cos(e+fx)}} 4i}{c^4 f} \right)}{e^{e4i+fx4i} \sin(e+fx) 28i - e^{e4i+fx4i} \sin(2e+2fx) 28i + e^{e4i+fx4i} \sin(3e+3fx) 12i - e^{e4i+fx4i} \sin(4e+4fx) 2i}$$

input `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(7/2)),x)`

output `-((c - c/cos(e + f*x))^(1/2))*((a^2*cos(e + f*x)*exp(e*4i + f*x*4i)*(a + a/cos(e + f*x))^(1/2)*52i)/(3*c^4*f) + (a^2*exp(e*4i + f*x*4i)*cos(3*e + 3*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^4*f))/(exp(e*4i + f*x*4i)*sin(e + f*x)*28i - exp(e*4i + f*x*4i)*sin(2*e + 2*f*x)*28i + exp(e*4i + f*x*4i)*sin(3*e + 3*f*x)*12i - exp(e*4i + f*x*4i)*sin(4*e + 4*f*x)*2i)`

3.131
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{9/2}} dx$$

3.131.1 Optimal result	940
3.131.2 Mathematica [A] (verified)	940
3.131.3 Rubi [A] (verified)	941
3.131.4 Maple [A] (verified)	942
3.131.5 Fracas [B] (verification not implemented)	943
3.131.6 Sympy [F(-1)]	943
3.131.7 Maxima [B] (verification not implemented)	943
3.131.8 Giac [F]	944
3.131.9 Mupad [B] (verification not implemented)	945

3.131.1 Optimal result

Integrand size = 36, antiderivative size = 88

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{9/2}} dx =$$

$$-\frac{(a+a \sec(e+fx))^{5/2} \tan(e+fx)}{8f(c-c \sec(e+fx))^{9/2}} - \frac{(a+a \sec(e+fx))^{5/2} \tan(e+fx)}{48cf(c-c \sec(e+fx))^{7/2}}$$

output
$$-1/8*(a+a*\sec(f*x+e))^(5/2)*\tan(f*x+e)/f/(c-c*\sec(f*x+e))^(9/2)-1/48*(a+a*\sec(f*x+e))^(5/2)*\tan(f*x+e)/c/f/(c-c*\sec(f*x+e))^(7/2)$$

3.131.2 Mathematica [A] (verified)

Time = 3.98 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.89

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{9/2}} dx =$$

$$-\frac{a^3(1+2 \sec(e+fx)+3 \sec^2(e+fx)) \tan(e+fx)}{6c^4 f(-1+\sec(e+fx))^4 \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input
$$\text{Integrate}[(\text{Sec}[e+f*x]*(a+a*\text{Sec}[e+f*x])^(5/2))/(c-c*\text{Sec}[e+f*x])^(9/2),x]$$

output
$$-1/6*(a^3*(1+2*\text{Sec}[e+f*x]+3*\text{Sec}[e+f*x]^2)*\text{Tan}[e+f*x])/(c^4*f*(-1+\text{Sec}[e+f*x])^4*\text{Sqrt}[a*(1+\text{Sec}[e+f*x])]*\text{Sqrt}[c-c*\text{Sec}[e+f*x]])$$

3.131.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{9/2}} dx$$

3.131.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a \sec(e+fx)+a)^{5/2}}{(c-c \sec(e+fx))^{9/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a \csc(e+fx+\frac{\pi}{2})+a)^{5/2}}{(c-c \csc(e+fx+\frac{\pi}{2}))^{9/2}} dx$$

↓ 4439

$$\frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^{5/2}}{(c-c \sec(e+fx))^{7/2}} dx}{8c} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^{5/2}}{8f(c-c \sec(e+fx))^{9/2}}$$

↓ 3042

$$\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^{5/2}}{(c-c \csc(e+fx+\frac{\pi}{2}))^{7/2}} dx}{8c} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^{5/2}}{8f(c-c \sec(e+fx))^{9/2}}$$

↓ 4438

$$\frac{\tan(e+fx)(a \sec(e+fx)+a)^{5/2}}{48cf(c-c \sec(e+fx))^{7/2}} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^{5/2}}{8f(c-c \sec(e+fx))^{9/2}}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(9/2),x]`

output `-1/8*((a + a*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^(9/2)) - ((a + a*Sec[e + f*x])^(5/2)*Tan[e + f*x])/(48*c*f*(c - c*Sec[e + f*x])^(7/2))`

3.131.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.131.4 Maple [A] (verified)

Time = 3.52 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.92

method	result	size
default	$-\frac{a^2(7 \cos(fx+e)-1)\sqrt{a(\sec(fx+e)+1)}(\cos(fx+e)+1)^2 \tan(fx+e) \sec(fx+e)^3}{48f(\sec(fx+e)-1)^4 \sqrt{-c(\sec(fx+e)-1)} c^4}$	81
risch	$\frac{2ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (3e^{7i(fx+e)}-3e^{6i(fx+e)}+17e^{5i(fx+e)}-10e^{4i(fx+e)}+17e^{3i(fx+e)}-3e^{2i(fx+e)}+3e^{i(fx+e)})}{3c^4(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)^7 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$	177

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(9/2), x, method=_RETURNVERBOSE)`

output `-1/48/f*a^2*(7*cos(f*x+e)-1)*(a*(sec(f*x+e)+1))^(1/2)*(cos(f*x+e)+1)^2/(sec(f*x+e)-1)^4/(-c*(sec(f*x+e)-1))^(1/2)/c^4*tan(f*x+e)*sec(f*x+e)^3`

3.131.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{9/2}} dx$$

3.131.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 166 vs. $2(76) = 152$.

Time = 0.27 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.89

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{9/2}} dx = \frac{(6a^2\cos(fx+e)^4 - 3a^2\cos(fx+e)^3 + 4a^2\cos(fx+e)^2 - a^2\cos(fx+e))\sqrt{(a\cos(fx+e)+a)/\cos(fx+e)}\sqrt{(c\cos(fx+e)+e-c)/\cos(fx+e)}}{6(c^5f\cos(fx+e)^4 - 4c^5f\cos(fx+e)^3 + 6c^5f\cos(fx+e)^2 - 4c^5f\cos(fx+e) + c^5f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(9/2),x, algo
rithm="fricas")
```

```
output 1/6*(6*a^2*cos(f*x + e)^4 - 3*a^2*cos(f*x + e)^3 + 4*a^2*cos(f*x + e)^2 -
a^2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x
+ e) - c)/cos(f*x + e))/((c^5*f*cos(f*x + e)^4 - 4*c^5*f*cos(f*x + e)^3 +
6*c^5*f*cos(f*x + e)^2 - 4*c^5*f*cos(f*x + e) + c^5*f)*sin(f*x + e))
```

3.131.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{9/2}} dx = \text{Timed out}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(9/2),x)
```

```
output Timed out
```

3.131.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2719 vs. $2(76) = 152$.

Time = 3.26 (sec) , antiderivative size = 2719, normalized size of antiderivative = 30.90

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{9/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(9/2),x, algo
rithm="maxima")`

output `2/3*(70*a^2*cos(6*f*x + 6*e)*sin(4*f*x + 4*e) - 70*a^2*cos(4*f*x + 4*e)*si
n(2*f*x + 2*e) + 3*a^2*sin(2*f*x + 2*e) + (3*a^2*sin(6*f*x + 6*e) + 10*a^2
*sin(4*f*x + 4*e) + 3*a^2*sin(2*f*x + 2*e))*cos(8*f*x + 8*e) + (3*a^2*sin(
8*f*x + 8*e) + 60*a^2*sin(6*f*x + 6*e) + 130*a^2*sin(4*f*x + 4*e) + 60*a^2
*sin(2*f*x + 2*e) - 32*a^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2
*e))) - 32*a^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos(7
/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (17*a^2*sin(8*f*x + 8*e)
+ 308*a^2*sin(6*f*x + 6*e) + 630*a^2*sin(4*f*x + 4*e) + 308*a^2*sin(2*f*x
+ 2*e) + 32*a^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*cos
(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (17*a^2*sin(8*f*x + 8*
e) + 308*a^2*sin(6*f*x + 6*e) + 630*a^2*sin(4*f*x + 4*e) + 308*a^2*sin(2*f*
*x + 2*e) + 32*a^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*c
os(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (3*a^2*sin(8*f*x + 8
*e) + 60*a^2*sin(6*f*x + 6*e) + 130*a^2*sin(4*f*x + 4*e) + 60*a^2*sin(2*f*
x + 2*e))*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (3*a^2*co
s(6*f*x + 6*e) + 10*a^2*cos(4*f*x + 4*e) + 3*a^2*cos(2*f*x + 2*e))*sin(8*f
*x + 8*e) - (70*a^2*cos(4*f*x + 4*e) - 3*a^2)*sin(6*f*x + 6*e) + 10*(7*a^2
*cos(2*f*x + 2*e) + a^2)*sin(4*f*x + 4*e) - (3*a^2*cos(8*f*x + 8*e) + 60*a
^2*cos(6*f*x + 6*e) + 130*a^2*cos(4*f*x + 4*e) + 60*a^2*cos(2*f*x + 2*e) -
32*a^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 32*a^2*c...`

3.131.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{9/2}} dx = \int \frac{(a\sec(fx+e)+a)^{5/2}\sec(fx+e)}{(-c\sec(fx+e)+c)^{9/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(9/2),x, algo
rithm="giac")`

output `sage0*x`

3.131.9 Mupad [B] (verification not implemented)

Time = 18.65 (sec) , antiderivative size = 350, normalized size of antiderivative = 3.98

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{9/2}} dx = \frac{\sqrt{c-\frac{c}{\cos(e+fx)}} \left(\frac{a^2 e^{e5i+fx5i} \sqrt{a+\frac{a}{\cos(e+fx)}} 68i}{3c^5 f} - \frac{a^2 \cos(e+fx) e^{e5i+fx5i} \sqrt{a+\frac{a}{\cos(e+fx)}}}{3c^5 f} \right)}{e^{e5i+fx5i} \sin(e+fx) 84i - e^{e5i+fx5i} \sin(2e+2fx) 96i + e^{e5i+fx5i} \sin(3e+3fx) 54i - e^{e5i+fx5i} \sin(4e+4fx) 16i + e^{e5i+fx5i} \sin(5e+5fx) 2i}$$

```
input int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(9/2)),x
)
```

```
output ((c - c/cos(e + f*x))^(1/2)*((a^2*exp(e*5i + f*x*5i)*(a + a/cos(e + f*x))^(1/2)*68i)/(3*c^5*f) - (a^2*cos(e + f*x)*exp(e*5i + f*x*5i)*(a + a/cos(e + f*x))^(1/2)*52i)/(3*c^5*f) + (a^2*exp(e*5i + f*x*5i)*cos(2*e + 2*f*x)*(a + a/cos(e + f*x))^(1/2)*80i)/(3*c^5*f) - (a^2*exp(e*5i + f*x*5i)*cos(3*e + 3*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^5*f) + (a^2*exp(e*5i + f*x*5i)*cos(4*e + 4*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^5*f)))/(exp(e*5i + f*x*5i)*sin(e + f*x)*84i - exp(e*5i + f*x*5i)*sin(2*e + 2*f*x)*96i + exp(e*5i + f*x*5i)*sin(3*e + 3*f*x)*54i - exp(e*5i + f*x*5i)*sin(4*e + 4*f*x)*16i + exp(e*5i + f*x*5i)*sin(5*e + 5*f*x)*2i)
```

3.132
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{11/2}} dx$$

3.132.1 Optimal result	946
3.132.2 Mathematica [A] (verified)	946
3.132.3 Rubi [A] (verified)	947
3.132.4 Maple [A] (verified)	948
3.132.5 Fricas [A] (verification not implemented)	949
3.132.6 Sympy [F(-1)]	949
3.132.7 Maxima [B] (verification not implemented)	950
3.132.8 Giac [F]	950
3.132.9 Mupad [B] (verification not implemented)	951

3.132.1 Optimal result

Integrand size = 36, antiderivative size = 133

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{11/2}} dx = -\frac{(a+a \sec(e+fx))^{5/2} \tan(e+fx)}{10f(c-c \sec(e+fx))^{11/2}} - \frac{(a+a \sec(e+fx))^{5/2} \tan(e+fx)}{40cf(c-c \sec(e+fx))^{9/2}} - \frac{(a+a \sec(e+fx))^{5/2} \tan(e+fx)}{240c^2f(c-c \sec(e+fx))^{7/2}}$$

```
output -1/10*(a+a*sec(f*x+e))^(5/2)*tan(f*x+e)/f/(c-c*sec(f*x+e))^(11/2)-1/40*(a+a*sec(f*x+e))^(5/2)*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(9/2)-1/240*(a+a*sec(f*x+e))^(5/2)*tan(f*x+e)/c^2/f/(c-c*sec(f*x+e))^(7/2)
```

3.132.2 Mathematica [A] (verified)

Time = 5.43 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.59

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{11/2}} dx = \frac{a^3(2+5 \sec(e+fx)+5 \sec^2(e+fx)) \tan(e+fx)}{15c^5f(-1+\sec(e+fx))^5 \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

```
input Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(11/2),x]
```

```
output (a^3*(2 + 5*Sec[e + f*x] + 5*Sec[e + f*x]^2)*Tan[e + f*x])/(15*c^5*f*(-1 + Sec[e + f*x])^5*sqrt[a*(1 + Sec[e + f*x])]*sqrt[c - c*Sec[e + f*x]])
```

3.132.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^{5/2}}{(c-c \sec(e+fx))^{11/2}} dx$$

3.132.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^{5/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{11/2}} dx \\
 & \quad \downarrow \text{4439} \\
 & \frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^{5/2}}{(c-c\sec(e+fx))^{9/2}} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{10f(c-c\sec(e+fx))^{11/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^{5/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{9/2}} dx}{5c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{10f(c-c\sec(e+fx))^{11/2}} \\
 & \quad \downarrow \text{4439} \\
 & \frac{\int \frac{\sec(e+fx)(\sec(e+fx)a+a)^{5/2}}{(c-c\sec(e+fx))^{7/2}} dx}{8c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{8f(c-c\sec(e+fx))^{9/2}} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{10f(c-c\sec(e+fx))^{11/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^{5/2}}{(c-c\csc(e+fx+\frac{\pi}{2}))^{7/2}} dx}{8c} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{8f(c-c\sec(e+fx))^{9/2}} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{10f(c-c\sec(e+fx))^{11/2}} \\
 & \quad \downarrow \text{4438} \\
 & -\frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{48cf(c-c\sec(e+fx))^{7/2}} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{8f(c-c\sec(e+fx))^{9/2}} - \frac{\tan(e+fx)(a\sec(e+fx)+a)^{5/2}}{10f(c-c\sec(e+fx))^{11/2}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^(5/2))/(c - c*Sec[e + f*x])^(11/2), x]`

$$3.132. \quad \int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx$$

output
$$-1/10*((a + a*\text{Sec}[e + f*x])^{(5/2)}*\text{Tan}[e + f*x])/(f*(c - c*\text{Sec}[e + f*x])^{(11/2)}) + (-1/8*((a + a*\text{Sec}[e + f*x])^{(5/2)}*\text{Tan}[e + f*x])/(f*(c - c*\text{Sec}[e + f*x])^{(9/2)}) - ((a + a*\text{Sec}[e + f*x])^{(5/2)}*\text{Tan}[e + f*x])/(48*c*f*(c - c*\text{Sec}[e + f*x])^{(7/2)})))/(5*c)$$

3.132.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.132.4 Maple [A] (verified)

Time = 3.62 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.68

method	result
default	$\frac{a^2 \left(31 \cos(fx+e)^2 - 8 \cos(fx+e) + 1 \right) \sqrt{a(\sec(fx+e)+1)} (\cos(fx+e)+1)^2 \tan(fx+e) \sec(fx+e)^4}{240 f (\sec(fx+e)-1)^5 \sqrt{-c(\sec(fx+e)-1)} c^5}$
risch	$\frac{2ia^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (15e^{9i(fx+e)} - 30e^{8i(fx+e)} + 140e^{7i(fx+e)} - 170e^{6i(fx+e)} + 282e^{5i(fx+e)} - 170e^{4i(fx+e)} + 140e^{3i(fx+e)} - 30e^{2i(fx+e)} + 15) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}{15c^5 (e^{i(fx+e)}+1) (e^{i(fx+e)}-1)^9 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$

3.132.
$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx$$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(11/2),x,method=_RE
TURNVERBOSE)
```

```
output 1/240/f*a^2*(31*cos(f*x+e)^2-8*cos(f*x+e)+1)*(a*(sec(f*x+e)+1))^(1/2)*(cos
(f*x+e)+1)^2/(sec(f*x+e)-1)^5/(-c*(sec(f*x+e)-1))^(1/2)/c^5*tan(f*x+e)*sec
(f*x+e)^4
```

3.132.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.46

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx = \frac{(15a^2 \cos(fx+e)^5 - 15a^2 \cos(fx+e)^4 + 20a^2 \cos(fx+e)^3 - 10a^2 \cos(fx+e)^2 + 2a^2 \cos(fx+e)) \sqrt{(a \cos(fx+e) + a) / \cos(fx+e)} \sqrt{(c \cos(fx+e) - c) / \cos(fx+e)}}{15(c^6 f \cos(fx+e)^5 - 5c^6 f \cos(fx+e)^4 + 10c^6 f \cos(fx+e)^3 - 10c^6 f \cos(fx+e)^2 + 5c^6 f \cos(fx+e) - c^6 f) \sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(11/2),x, alg
orithm="fricas")
```

```
output 1/15*(15*a^2*cos(f*x + e)^5 - 15*a^2*cos(f*x + e)^4 + 20*a^2*cos(f*x + e)^
3 - 10*a^2*cos(f*x + e)^2 + 2*a^2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/
cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((c^6*f*cos(f*x + e)
^5 - 5*c^6*f*cos(f*x + e)^4 + 10*c^6*f*cos(f*x + e)^3 - 10*c^6*f*cos(f*x +
e)^2 + 5*c^6*f*cos(f*x + e) - c^6*f)*sin(f*x + e))
```

3.132.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx = \text{Timed out}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(11/2),x)
```

```
output Timed out
```

3.132.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4108 vs. $2(115) = 230$.

Time = 17.08 (sec) , antiderivative size = 4108, normalized size of antiderivative = 30.89

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(11/2),x, algorithm="maxima")`

output

```
-2/15*(1350*a^2*cos(6*f*x + 6*e)*sin(2*f*x + 2*e) + 1350*a^2*cos(4*f*x + 4
*e)*sin(2*f*x + 2*e) - 30*a^2*sin(2*f*x + 2*e) - 10*(3*a^2*sin(8*f*x + 8*e
) + 17*a^2*sin(6*f*x + 6*e) + 17*a^2*sin(4*f*x + 4*e) + 3*a^2*sin(2*f*x +
2*e))*cos(10*f*x + 10*e) - 1350*(a^2*sin(6*f*x + 6*e) + a^2*sin(4*f*x + 4
e))*cos(8*f*x + 8*e) - 5*(3*a^2*sin(10*f*x + 10*e) + 75*a^2*sin(8*f*x + 8
e) + 290*a^2*sin(6*f*x + 6*e) + 290*a^2*sin(4*f*x + 4*e) + 75*a^2*sin(2*f
*x + 2*e) - 80*a^2*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - 1
92*a^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 80*a^2*sin(3
/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*cos(9/2*arctan2(sin(2*f*x
+ 2*e), cos(2*f*x + 2*e))) - 20*(7*a^2*sin(10*f*x + 10*e) + 135*a^2*sin(8
*f*x + 8*e) + 450*a^2*sin(6*f*x + 6*e) + 450*a^2*sin(4*f*x + 4*e) + 135*a^
2*sin(2*f*x + 2*e) - 72*a^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) + 20*a^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*cos(
7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 6*(47*a^2*sin(10*f*x +
10*e) + 855*a^2*sin(8*f*x + 8*e) + 2730*a^2*sin(6*f*x + 6*e) + 2730*a^2*si
n(4*f*x + 4*e) + 855*a^2*sin(2*f*x + 2*e) + 240*a^2*sin(3/2*arctan2(sin(2*
f*x + 2*e), cos(2*f*x + 2*e))) + 160*a^2*sin(1/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))))*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))
- 20*(7*a^2*sin(10*f*x + 10*e) + 135*a^2*sin(8*f*x + 8*e) + 450*a^2*sin(6*
f*x + 6*e) + 450*a^2*sin(4*f*x + 4*e) + 135*a^2*sin(2*f*x + 2*e) + 20*a...
```

3.132.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx = \int \frac{(a\sec(fx+e)+a)^{5/2}\sec(fx+e)}{(-c\sec(fx+e)+c)^{11/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(11/2),x, algorithm="giac")`

output `sage0*x`

3.132.9 Mupad [B] (verification not implemented)

Time = 18.62 (sec) , antiderivative size = 419, normalized size of antiderivative = 3.15

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^{5/2}}{(c-c\sec(e+fx))^{11/2}} dx = \frac{\sqrt{c-\frac{c}{\cos(e+fx)}} \left(\frac{a^2 e^{e6i+fx6i} \sqrt{a+\frac{a}{\cos(e+fx)}} 136i}{3c^6 f} - \frac{a^2 \cos(e+fx) e^{e6i+fx6i}}{15c^6 f} \right)}{e^{e6i+fx6i} \sin(e+fx) 264i - e^{e6i+fx6i} \sin(e+fx)}$$

input `int((a + a/cos(e + f*x))^(5/2)/(cos(e + f*x)*(c - c/cos(e + f*x))^(11/2)), x)`

output `((c - c/cos(e + f*x))^(1/2)*((a^2*exp(e*6i + f*x*6i)*(a + a/cos(e + f*x))^(1/2)*136i)/(3*c^6*f) - (a^2*cos(e + f*x)*exp(e*6i + f*x*6i)*(a + a/cos(e + f*x))^(1/2)*1688i)/(15*c^6*f) + (a^2*exp(e*6i + f*x*6i)*cos(2*e + 2*f*x)*(a + a/cos(e + f*x))^(1/2)*160i)/(3*c^6*f) - (a^2*exp(e*6i + f*x*6i)*cos(3*e + 3*f*x)*(a + a/cos(e + f*x))^(1/2)*124i)/(3*c^6*f) + (a^2*exp(e*6i + f*x*6i)*cos(4*e + 4*f*x)*(a + a/cos(e + f*x))^(1/2)*8i)/(c^6*f) - (a^2*exp(e*6i + f*x*6i)*cos(5*e + 5*f*x)*(a + a/cos(e + f*x))^(1/2)*4i)/(c^6*f)))/(exp(e*6i + f*x*6i)*sin(e + f*x)*264i - exp(e*6i + f*x*6i)*sin(2*e + 2*f*x)*330i + exp(e*6i + f*x*6i)*sin(3*e + 3*f*x)*220i - exp(e*6i + f*x*6i)*sin(4*e + 4*f*x)*88i + exp(e*6i + f*x*6i)*sin(5*e + 5*f*x)*20i - exp(e*6i + f*x*6i)*sin(6*e + 6*f*x)*2i)`

3.133
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a+a\sec(e+fx)}} dx$$

3.133.1 Optimal result	952
3.133.2 Mathematica [A] (verified)	952
3.133.3 Rubi [A] (verified)	953
3.133.4 Maple [A] (verified)	955
3.133.5 Fricas [F]	955
3.133.6 Sympy [F(-1)]	956
3.133.7 Maxima [B] (verification not implemented)	956
3.133.8 Giac [A] (verification not implemented)	957
3.133.9 Mupad [F(-1)]	958

3.133.1 Optimal result

Integrand size = 36, antiderivative size = 139

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a+a\sec(e+fx)}} dx = -\frac{4c^3 \log(1+\sec(e+fx)) \tan(e+fx)}{f\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} - \frac{2c^2 \sqrt{c-c\sec(e+fx)} \tan(e+fx)}{f\sqrt{a+a\sec(e+fx)}} - \frac{c(c-c\sec(e+fx))^{3/2} \tan(e+fx)}{2f\sqrt{a+a\sec(e+fx)}}$$

output `-1/2*c*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)-4*c^3*ln(1+sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-2*c^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)`

3.133.2 Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.53

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a+a\sec(e+fx)}} dx = \frac{c^3(1+8\log(1+\sec(e+fx))-6\sec(e+fx)+\sec^2(e+fx))\tan(e+fx)}{2f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e+f*x]*(c-c*Sec[e+f*x])^(5/2))/Sqrt[a+a*Sec[e+f*x]],x]`

output
$$-1/2*(c^3*(1 + 8*\text{Log}[1 + \text{Sec}[e + f*x]] - 6*\text{Sec}[e + f*x] + \text{Sec}[e + f*x]^2)*\text{Tan}[e + f*x])/(f*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]*\text{Sqrt}[c - c*\text{Sec}[e + f*x]])$$

3.133.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4443, 3042, 4443, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a\sec(e+fx)+a}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}} dx \\ & \quad \downarrow \text{4443} \\ & 2c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{\sec(e+fx)a+a}} dx - \frac{c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{2f\sqrt{a\sec(e+fx)+a}} \\ & \quad \downarrow \text{3042} \\ & 2c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx - \frac{c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{2f\sqrt{a\sec(e+fx)+a}} \\ & \quad \downarrow \text{4443} \\ & 2c \left(2c \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} dx - \frac{c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f\sqrt{a\sec(e+fx)+a}} \right) - \\ & \quad \frac{c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{2f\sqrt{a\sec(e+fx)+a}} \\ & \quad \downarrow \text{3042} \\ & 2c \left(2c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx - \frac{c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f\sqrt{a\sec(e+fx)+a}} \right) - \\ & \quad \frac{c\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{2f\sqrt{a\sec(e+fx)+a}} \end{aligned}$$

3.133.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a+a\sec(e+fx)}} dx$$

$$\downarrow 4440$$

$$2c \left(\frac{2c^2 \tan(e+fx) \log(\sec(e+fx)+1)}{f\sqrt{a \sec(e+fx)+a}\sqrt{c-c \sec(e+fx)}} - \frac{c \tan(e+fx) \sqrt{c-c \sec(e+fx)}}{f\sqrt{a \sec(e+fx)+a}} \right) - \frac{c \tan(e+fx) (c-c \sec(e+fx))^{3/2}}{2f\sqrt{a \sec(e+fx)+a}}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/Sqrt[a + a*Sec[e + f*x]],x]`

output `-1/2*(c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]) + 2*c*((-2*c^2*Log[1 + Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]])*Sqrt[c - c*Sec[e + f*x]]) - (c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]])`

3.133.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.133.4 Maple [A] (verified)

Time = 3.78 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99

method	result
default	$-\frac{c^2 \sqrt{a(\sec(fx+e)+1)} \sqrt{-c(\sec(fx+e)-1)} (\sec(fx+e)-1)^2 (8 \cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e)-1)+8 \cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e)+1)+7 \cos(fx+e)^2+6 \cos(fx+e)-1)}{2fa(\cos(fx+e)-1)^2}$
risch	$-\frac{2ic^2 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} (3e^{2i(fx+e)}-e^{i(fx+e)}+3)(e^{2i(fx+e)}+e^{i(fx+e)})}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f(1+e^{2i(fx+e)})^2} + \frac{8ic^2(e^{i(fx+e)}+1) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} \ln(e^{i(fx+e)}+1)}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2/f/a*c^2*(a*(sec(f*x+e)+1))^(1/2)*(-c*(sec(f*x+e)-1))^(1/2)*(sec(f*x+e)-1)^2*(8*cos(f*x+e)^2*ln(-cot(f*x+e)+csc(f*x+e)-1)+8*cos(f*x+e)^2*ln(-cot(f*x+e)+csc(f*x+e)+1)+7*cos(f*x+e)^2+6*cos(f*x+e)-1)/(cos(f*x+e)-1)^2*cot(f*x+e)`

3.133.5 Fracas [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{(-c\sec(fx+e)+c)^{5/2} \sec(fx+e)}{\sqrt{a\sec(fx+e)+a}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x,algorithm="fracas")`

output `integral((c^2*sec(f*x + e)^3 - 2*c^2*sec(f*x + e)^2 + c^2*sec(f*x + e))*sqrt(-c*sec(f*x + e) + c)/sqrt(a*sec(f*x + e) + a), x)`

3.133.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(5/2)/(a+a*sec(f*x+e))**(1/2),x)`

output `Timed out`

3.133.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 737 vs. 2(125) = 250.

Time = 0.40 (sec) , antiderivative size = 737, normalized size of antiderivative = 5.30

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}} dx = \frac{2(c^2 \cos(2fx + 2e) \sin(4fx + 4e) - c^2 \cos(4fx + 4e) \sin(2fx + 2e))}{\sqrt{a + a \sec(e + fx)}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output

```

2*(c^2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) - c^2*cos(4*f*x + 4*e)*sin(2*f*x
+ 2*e) - c^2*sin(2*f*x + 2*e) + 2*(c^2*cos(4*f*x + 4*e)^2 + 4*c^2*cos(2*f*
x + 2*e)^2 + c^2*sin(4*f*x + 4*e)^2 + 4*c^2*sin(4*f*x + 4*e)*sin(2*f*x + 2
*e) + 4*c^2*sin(2*f*x + 2*e)^2 + 4*c^2*cos(2*f*x + 2*e) + c^2 + 2*(2*c^2*c
os(2*f*x + 2*e) + c^2)*cos(4*f*x + 4*e))*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e) + 1) - 4*(c^2*cos(4*f*x + 4*e)^2 + 4*c^2*cos(2*f*x + 2*e)^2 + c^
2*sin(4*f*x + 4*e)^2 + 4*c^2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*c^2*sin
(2*f*x + 2*e)^2 + 4*c^2*cos(2*f*x + 2*e) + c^2 + 2*(2*c^2*cos(2*f*x + 2*e)
+ c^2)*cos(4*f*x + 4*e))*arctan2(sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*
f*x + 2*e))), cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) -
3*(c^2*sin(4*f*x + 4*e) + 2*c^2*sin(2*f*x + 2*e))*cos(3/2*arctan2(sin(2*f*
x + 2*e), cos(2*f*x + 2*e))) - 3*(c^2*sin(4*f*x + 4*e) + 2*c^2*sin(2*f*x +
2*e))*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 3*(c^2*cos(4
*f*x + 4*e) + 2*c^2*cos(2*f*x + 2*e) + c^2)*sin(3/2*arctan2(sin(2*f*x + 2*
e), cos(2*f*x + 2*e))) + 3*(c^2*cos(4*f*x + 4*e) + 2*c^2*cos(2*f*x + 2*e)
+ c^2)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)*sqrt(
c)/((a*cos(4*f*x + 4*e)^2 + 4*a*cos(2*f*x + 2*e)^2 + a*sin(4*f*x + 4*e)^2
+ 4*a*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*a*sin(2*f*x + 2*e)^2 + 2*(2*a*
cos(2*f*x + 2*e) + a)*cos(4*f*x + 4*e) + 4*a*cos(2*f*x + 2*e) + a)*f)

```

3.133.8 Giac [A] (verification not implemented)

Time = 1.55 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.15

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a+a\sec(e+fx)}} dx = \frac{2c^2 \left(\frac{2\sqrt{-acc} \log\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)}{a|c|} - \frac{3\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^2 \sqrt{-acc+4}\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)^2} \right)}{\sqrt{a+a\sec(e+fx)}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x, algo
rithm="giac")`

output

```

2*c^2*(2*sqrt(-a*c)*c*log(c*tan(1/2*f*x + 1/2*e)^2 - c)/(a*abs(c)) - (3*(c
*tan(1/2*f*x + 1/2*e)^2 - c)^2*sqrt(-a*c)*c + 4*(c*tan(1/2*f*x + 1/2*e)^2
- c)*sqrt(-a*c)*c^2 + sqrt(-a*c)*c^3)/((c*tan(1/2*f*x + 1/2*e)^2 - c)^2*a*
abs(c))*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))/f

```

3.133.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{\left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}}{\cos(e+fx) \sqrt{a + \frac{a}{\cos(e+fx)}}} dx$$

input `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

output `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

3.134
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx$$

3.134.1 Optimal result	959
3.134.2 Mathematica [A] (verified)	959
3.134.3 Rubi [A] (verified)	960
3.134.4 Maple [A] (verified)	961
3.134.5 Fracas [F]	962
3.134.6 Sympy [F]	962
3.134.7 Maxima [B] (verification not implemented)	962
3.134.8 Giac [A] (verification not implemented)	963
3.134.9 Mupad [F(-1)]	963

3.134.1 Optimal result

Integrand size = 36, antiderivative size = 94

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx = -\frac{2c^2 \log(1+\sec(e+fx)) \tan(e+fx)}{f\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} - \frac{c\sqrt{c-c\sec(e+fx)} \tan(e+fx)}{f\sqrt{a+a\sec(e+fx)}}$$

output `-2*c^2*ln(1+sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-c*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)`

3.134.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.64

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx = \frac{c^2(-2\log(1+\sec(e+fx))+\sec(e+fx))\tan(e+fx)}{f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/Sqrt[a + a*Sec[e + f*x]],x]`

output `(c^2*(-2*Log[1 + Sec[e + f*x]] + Sec[e + f*x])*Tan[e + f*x])/(f*Sqrt[a*(1 + Sec[e + f*x]])*Sqrt[c - c*Sec[e + f*x]])`

3.134.
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx$$

3.134.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4443, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a\sec(e+fx)+a}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}} dx$$

↓ 4443

$$2c \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} dx - \frac{c \tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f\sqrt{a\sec(e+fx)+a}}$$

↓ 3042

$$2c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx - \frac{c \tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f\sqrt{a\sec(e+fx)+a}}$$

↓ 4440

$$\frac{2c^2 \tan(e+fx) \log(\sec(e+fx)+1)}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} - \frac{c \tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f\sqrt{a\sec(e+fx)+a}}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/Sqrt[a + a*Sec[e + f*x]],x]`

output `(-2*c^2*Log[1 + Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]) - (c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]])`

3.134.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.134.4 Maple [A] (verified)

Time = 3.62 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.23

method	result
default	$\frac{c(\sec(fx+e)-1)\sqrt{-c(\sec(fx+e)-1)}\sqrt{a(\sec(fx+e)+1)}(2\cos(fx+e)\ln(-\cot(fx+e)+\csc(fx+e)-1)+2\cos(fx+e)\ln(-\cot(fx+e)-1))}{fa(\cos(fx+e)-1)}$
risch	$\frac{2ic\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}(2\ln(e^{i(fx+e)}+1)e^{3i(fx+e)}-\ln(1+e^{2i(fx+e)})e^{3i(fx+e)}+2e^{i(fx+e)}\ln(e^{i(fx+e)}+1)-e^{i(fx+e)}\ln(1+e^{2i(fx+e)}))}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `1/f/a*c*(sec(f*x+e)-1)*(-c*(sec(f*x+e)-1))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)*(2*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)-1)+2*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)+cos(f*x+e)+1)/(cos(f*x+e)-1)*cot(f*x+e)`

$$3.134. \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx$$

3.134.5 Fricas [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{(-c\sec(fx+e)+c)^{3/2} \sec(fx+e)}{\sqrt{a\sec(fx+e)+a}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral(-(c*sec(f*x + e)^2 - c*sec(f*x + e))*sqrt(-c*sec(f*x + e) + c)/sqrt(a*sec(f*x + e) + a), x)`

3.134.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{(-c(\sec(e+fx)-1))^{3/2} \sec(e+fx)}{\sqrt{a(\sec(e+fx)+1)}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(3/2)/(a+a*sec(f*x+e))**(1/2),x)`

output `Integral((-c*(sec(e + f*x) - 1))**(3/2)*sec(e + f*x)/sqrt(a*(sec(e + f*x) + 1)), x)`

3.134.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 276 vs. 2(86) = 172.

Time = 0.39 (sec) , antiderivative size = 276, normalized size of antiderivative = 2.94

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx =$$

$$\frac{2(c \cos(\frac{1}{2} \arctan(\sin(2fx+2e), \cos(2fx+2e))) \sin(2fx+2e) - (c \cos(2fx+2e))^2 + c \sin(2fx+2e))}{\sqrt{a+a\sec(e+fx)}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

3.134. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{a+a\sec(e+fx)}} dx$

output `-2*(c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*sin(2*f*x + 2*e) - (c*cos(2*f*x + 2*e)^2 + c*sin(2*f*x + 2*e)^2 + 2*c*cos(2*f*x + 2*e) + c)*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + 2*(c*cos(2*f*x + 2*e)^2 + c*sin(2*f*x + 2*e)^2 + 2*c*cos(2*f*x + 2*e) + c)*arctan2(sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) - (c*cos(2*f*x + 2*e) + c)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)*sqrt(c)/((a*cos(2*f*x + 2*e)^2 + a*sin(2*f*x + 2*e)^2 + 2*a*cos(2*f*x + 2*e) + a)*f)`

3.134.8 Giac [A] (verification not implemented)

Time = 1.34 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.37

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}} dx = \frac{2 \left(\frac{\sqrt{-acc^2} \log\left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c\right)}{a|c|} - \frac{\left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c\right) \sqrt{-acc^2 + \sqrt{-acc^3}}}{\left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c\right) a|c|} \right)}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x, algo rithm="giac")`

output `2*(sqrt(-a*c)*c^2*log(c*tan(1/2*f*x + 1/2*e)^2 - c)/(a*abs(c)) - ((c*tan(1/2*f*x + 1/2*e)^2 - c)*sqrt(-a*c)*c^2 + sqrt(-a*c)*c^3)/((c*tan(1/2*f*x + 1/2*e)^2 - c)*a*abs(c))*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))/f`

3.134.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}} dx = \int \frac{\left(c - \frac{c}{\cos(e + fx)}\right)^{3/2}}{\cos(e + fx) \sqrt{a + \frac{a}{\cos(e + fx)}}} dx$$

input `int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

output `int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

$$3.135 \quad \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx$$

3.135.1 Optimal result 964
 3.135.2 Mathematica [A] (verified) 964
 3.135.3 Rubi [A] (verified) 965
 3.135.4 Maple [A] (verified) 966
 3.135.5 Fricas [F] 966
 3.135.6 Sympy [F] 966
 3.135.7 Maxima [A] (verification not implemented) 967
 3.135.8 Giac [A] (verification not implemented) 967
 3.135.9 Mupad [F(-1)] 968

3.135.1 Optimal result

Integrand size = 36, antiderivative size = 50

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = -\frac{c \log(1+\sec(e+fx)) \tan(e+fx)}{f \sqrt{a+a\sec(e+fx)} \sqrt{c-c\sec(e+fx)}}$$

output `-c*ln(1+sec(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.135.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = -\frac{c \log(1+\sec(e+fx)) \tan(e+fx)}{f \sqrt{a(1+\sec(e+fx))} \sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/Sqrt[a + a*Sec[e + f*x]],x]`

output `-((c*Log[1 + Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a*(1 + Sec[e + f*x]])*Sqrt[c - c*Sec[e + f*x]]))`

3.135.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a\sec(e+fx)+a}} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)}}{\sqrt{a\csc\left(e+fx+\frac{\pi}{2}\right)+a}} dx$$

↓ 4440

$$-\frac{c\tan(e+fx)\log(\sec(e+fx)+1)}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/Sqrt[a + a*Sec[e + f*x]],x]`

output `-((c*Log[1 + Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]))`

3.135.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.135.4 Maple [A] (verified)

Time = 3.68 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.50

method	result	size
default	$-\frac{\sqrt{-c(\sec(fx+e)-1)}\sqrt{a(\sec(fx+e)+1)}(\ln(-\cot(fx+e)+\csc(fx+e)-1)+\ln(-\cot(fx+e)+\csc(fx+e)+1))\cot(fx+e)}{fa}$	75
risch	$\frac{2i(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}\ln(e^{i(fx+e)}+1)}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)f}} - \frac{i(e^{i(fx+e)}+1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}\ln(1+e^{2i(fx+e)})}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)f}}$	206

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -1/f/a*(-c*(sec(f*x+e)-1))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)*(ln(-cot(f*x+e)+
csc(f*x+e)-1)+ln(-cot(f*x+e)+csc(f*x+e)+1))*cot(f*x+e)
```

3.135.5 Fracas [F]

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{\sqrt{-c\sec(fx+e)+c\sec(fx+e)}}{\sqrt{a\sec(fx+e)+a}} dx$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x, algo
rithm="fracas")
```

```
output integral(sqrt(-c*sec(f*x + e) + c)*sec(f*x + e)/sqrt(a*sec(f*x + e) + a),
x)
```

3.135.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{\sqrt{-c(\sec(e+fx)-1)}\sec(e+fx)}{\sqrt{a(\sec(e+fx)+1)}} dx$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(1/2)/(a+a*sec(f*x+e))**(1/2),x)
```

3.135.
$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx$$

output `Integral(sqrt(-c*(sec(e + f*x) - 1))*sec(e + f*x)/sqrt(a*(sec(e + f*x) + 1)), x)`

3.135.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.28

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{\sqrt{a + a \sec(e + fx)}} dx = -\frac{\sqrt{c} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{\sqrt{-a}} + \frac{\sqrt{c} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `-(sqrt(c)*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/sqrt(-a) + sqrt(c)*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/sqrt(-a))/f`

3.135.8 Giac [A] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.18

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{\sqrt{a + a \sec(e + fx)}} dx = -\frac{c^2 \log\left(\left|c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c\right|\right) \operatorname{sgn}\left(\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)\right)}{\sqrt{-ac} f |c|}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `-c^2*log(abs(c*tan(1/2*f*x + 1/2*e)^2 - c))*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))/(sqrt(-a*c)*f*abs(c))`

3.135.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{\sqrt{c-\frac{c}{\cos(e+fx)}}}{\cos(e+fx)\sqrt{a+\frac{a}{\cos(e+fx)}}} dx$$

input `int((c - c/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

output `int((c - c/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

3.136
$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx$$

3.136.1 Optimal result 969
 3.136.2 Mathematica [A] (verified) 969
 3.136.3 Rubi [A] (verified) 970
 3.136.4 Maple [A] (verified) 971
 3.136.5 Fricas [A] (verification not implemented) 972
 3.136.6 Sympy [F] 972
 3.136.7 Maxima [A] (verification not implemented) 973
 3.136.8 Giac [A] (verification not implemented) 973
 3.136.9 Mupad [F(-1)] 974

3.136.1 Optimal result

Integrand size = 36, antiderivative size = 47

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx = -\frac{\operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{f\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}}$$

output `-arctanh(cos(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.136.2 Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx = -\frac{\operatorname{arctanh}(\sec(e+fx)) \tan(e+fx)}{f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]),x]`

output `-((ArcTanh[Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]]))`

3.136.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {3042, 4447, 25, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx$$

↓ 4447

$$-\frac{\tan(e+fx) \int -\csc(e+fx) dx}{\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

↓ 25

$$\frac{\tan(e+fx) \int \csc(e+fx) dx}{\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

↓ 3042

$$\frac{\tan(e+fx) \int \csc(e+fx) dx}{\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

↓ 4257

$$-\frac{\tan(e+fx)\operatorname{arctanh}(\cos(e+fx))}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

input `Int[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]),x]`

output `-((ArcTanh[Cos[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]))`

3.136. $\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} dx$

3.136.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4447 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(m_), x_Symbol] := Simp[((-a)*c)^(m + 1/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Int[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]`

3.136.4 Maple [A] (verified)

Time = 3.24 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.38

method	result	size
default	$-\frac{\sqrt{a(\sec(fx+e)+1)} \ln(-\cot(fx+e)+\csc(fx+e))(\cot(fx+e)-\csc(fx+e))}{fa\sqrt{-c(\sec(fx+e)-1)}}$	65
risch	$-\frac{i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)\ln(e^{i(fx+e)}-1)}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}(1+e^{2i(fx+e)})}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f} + \frac{i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)\ln(e^{i(fx+e)}+1)}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}(1+e^{2i(fx+e)})}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}$	228

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/f/a*(a*(sec(f*x+e)+1))^(1/2)*ln(-cot(f*x+e)+csc(f*x+e))/(-c*(sec(f*x+e)-1))^(1/2)*(cot(f*x+e)-csc(f*x+e))`

3.136.
$$\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} dx$$

3.136.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 204, normalized size of antiderivative = 4.34

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx$$

$$= \left[\frac{\sqrt{-ac} \log \left(-\frac{4 \left(2\sqrt{-ac} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{c \cos(fx+e)-c}{\cos(fx+e)}} \cos(fx+e)^2 + (ac \cos(fx+e)^2 + ac) \sin(fx+e) \right)}{(\cos(fx+e)^2 - 1) \sin(fx+e)} \right)}{2acf}, \sqrt{ac} \arctan \left(\frac{\sqrt{ac} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}}}{\sqrt{c - c \sec(e + fx)}} \right) \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="fricas")
```

```
output [-1/2*sqrt(-a*c)*log(-4*(2*sqrt(-a*c))*sqrt((a*cos(f*x + e) + a)/cos(f*x +
e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2 + (a*c*cos(f*x
+ e)^2 + a*c)*sin(f*x + e))/((cos(f*x + e)^2 - 1)*sin(f*x + e)))/(a*c*f),
sqrt(a*c)*arctan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c
*cos(f*x + e) - c)/cos(f*x + e))/(a*c*sin(f*x + e)))/(a*c*f)]
```

3.136.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx$$

$$= \int \frac{\sec(e + fx)}{\sqrt{a (\sec(e + fx) + 1)} \sqrt{-c (\sec(e + fx) - 1)}} dx$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(1/2)/(c-c*sec(f*x+e))**(1/2),x)
```

```
output Integral(sec(e + f*x)/(sqrt(a*(sec(e + f*x) + 1))*sqrt(-c*(sec(e + f*x) -
1))), x)
```

3.136.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.94

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{\arctan(\sin(fx + e), \cos(fx + e) + 1) - \arctan(\sin(fx + e), \cos(fx + e) - 1)}{\sqrt{a}\sqrt{c}f}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="maxima")
```

```
output -(arctan2(sin(f*x + e), cos(f*x + e) + 1) - arctan2(sin(f*x + e), cos(f*x
+ e) - 1))/(sqrt(a)*sqrt(c)*f)
```

3.136.8 Giac [A] (verification not implemented)

Time = 1.62 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.49

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx$$

$$= \frac{c^2 \left(\frac{\log(|c| \tan(\frac{1}{2} fx + \frac{1}{2} e)^2)}{c} - \frac{\log(|c|)}{c} \right)}{2 \sqrt{-ac} f |c| \operatorname{sgn} \left(\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="giac")
```

```
output -1/2*c^2*(log(abs(c)*tan(1/2*f*x + 1/2*e)^2)/c - log(abs(c))/c)/(sqrt(-a*c
)*f*abs(c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e)))
```

3.136.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx) \sqrt{a + \frac{a}{\cos(e + fx)}} \sqrt{c - \frac{c}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x))*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int(1/(cos(e + f*x))*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(1/2)), x)`

$$3.137 \quad \int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}} dx$$

3.137.1 Optimal result	975
3.137.2 Mathematica [A] (verified)	975
3.137.3 Rubi [A] (verified)	976
3.137.4 Maple [A] (verified)	978
3.137.5 Fracas [B] (verification not implemented)	978
3.137.6 Sympy [F]	979
3.137.7 Maxima [B] (verification not implemented)	979
3.137.8 Giac [A] (verification not implemented)	980
3.137.9 Mupad [F(-1)]	980

3.137.1 Optimal result

Integrand size = 36, antiderivative size = 95

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}} dx =$$

$$\frac{\tan(e+fx)}{2f\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}}$$

$$-\frac{\operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{2cf\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}}$$

output `-1/2*tan(f*x+e)/f/(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2)-1/2*arctanh(cos(f*x+e))*tan(f*x+e)/c/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.137.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.79

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}} dx =$$

$$\frac{c\left(\frac{\operatorname{arctanh}(\sec(e+fx))}{2c^2} + \frac{1}{2c^2(1-\sec(e+fx))}\right) \tan(e+fx)}{f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(3/2)),x]`

3.137. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}} dx$

output $-\left(\left(c \cdot \text{ArcTanh}[\text{Sec}[e + f \cdot x]] / (2 \cdot c^2) + 1 / (2 \cdot c^2 \cdot (1 - \text{Sec}[e + f \cdot x]))\right) \cdot \text{Tan}[e + f \cdot x]\right) / (f \cdot \text{Sqrt}[a \cdot (1 + \text{Sec}[e + f \cdot x])] \cdot \text{Sqrt}[c - c \cdot \text{Sec}[e + f \cdot x]])$

3.137.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 4448, 3042, 4447, 25, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{\sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))^{3/2}}} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a(c - c \csc(e + fx + \frac{\pi}{2}))^{3/2}}} dx$$

↓ 4448

$$\frac{\int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a+a\sqrt{c-c\sec(e+fx)}}} dx}{2c} - \frac{\tan(e + fx)}{2f \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))^{3/2}}}$$

↓ 3042

$$\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}} dx}{2c} - \frac{\tan(e + fx)}{2f \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))^{3/2}}}$$

↓ 4447

$$- \frac{\tan(e + fx) \int -\csc(e + fx) dx}{2c \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} - \frac{\tan(e + fx)}{2f \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))^{3/2}}}$$

↓ 25

$$\frac{\tan(e + fx) \int \csc(e + fx) dx}{2c \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} - \frac{\tan(e + fx)}{2f \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))^{3/2}}}$$

↓ 3042

$$\frac{\tan(e + fx) \int \csc(e + fx) dx}{2c \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} - \frac{\tan(e + fx)}{2f \sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))^{3/2}}}$$

↓ 4257

3.137. $\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)(c-c\sec(e+fx))^{3/2}}} dx$

$$\frac{\tan(e+fx)\operatorname{arctanh}(\cos(e+fx))}{2cf\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} - \frac{\tan(e+fx)}{2f\sqrt{a\sec(e+fx)+a}(c-c\sec(e+fx))^{3/2}}$$

input `Int[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(3/2)),x]`

output `-1/2*Tan[e + f*x]/(f*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(3/2))
- (ArcTanh[Cos[e + f*x]]*Tan[e + f*x]/(2*c*f*Sqrt[a + a*Sec[e + f*x]]*Sqr
t[c - c*Sec[e + f*x]))`

3.137.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]`

rule 4447 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(cs
c[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(m_), x_Symbol] := Simp[((-a)*c)^(m + 1
/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) In
t[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] &
& EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]`

rule 4448 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(cs
c[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*
(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[
(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*
(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (IL
tQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))`

3.137.4 Maple [A] (verified)

Time = 3.59 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.21

method	result
default	$-\frac{(2 \cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e))-2 \ln(-\cot(fx+e)+\csc(fx+e))-\cos(fx+e)-1)\sqrt{a(\sec(fx+e)+1)} \tan(fx+e)}{4fa\sqrt{-c(\sec(fx+e)-1)} c(\sec(fx+e)-1)(\cos(fx+e)+1)}$
risch	$i(\ln(e^{i(fx+e)}+1)e^{3i(fx+e)}-e^{3i(fx+e)} \ln(e^{i(fx+e)}-1)-e^{2i(fx+e)} \ln(e^{i(fx+e)}+1)+e^{2i(fx+e)} \ln(e^{i(fx+e)}-1)-e^{i(fx+e)} \ln(e^{i(fx+e)}+1))$ $2c\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}(1+e^{2i(fx+e)})(e^{i(fx+e)}-1)}\sqrt{\frac{c(e^{i(fx+e)}+1)}{1+e^{2i(fx+e)}}}$

input `int(sec(f*x+e)/(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4/f/a*(2*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e))-2*ln(-cot(f*x+e)+csc(f*x+e))-cos(f*x+e)-1)*(a*(sec(f*x+e)+1))^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)/c/(sec(f*x+e)-1)/(cos(f*x+e)+1)*tan(f*x+e)`

3.137.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. 2(83) = 166.

Time = 0.34 (sec) , antiderivative size = 382, normalized size of antiderivative = 4.02

$$\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{3/2}} dx = \left[-\frac{\sqrt{-ac}(\cos(fx+e)-1) \log\left(-\frac{4(2\sqrt{-ac}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c(e^{i(fx+e)}+1)}{1+e^{2i(fx+e)}}})\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x, algo rithm="fricas")`

output `[-1/4*(sqrt(-a*c)*(cos(f*x + e) - 1)*log(-4*(2*sqrt(-a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2 + (a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e)))/((cos(f*x + e)^2 - 1)*sin(f*x + e))*sin(f*x + e) - 2*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e))/((a*c^2*f*cos(f*x + e) - a*c^2*f)*sin(f*x + e)), 1/2*(sqrt(a*c)*(cos(f*x + e) - 1)*arctan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(a*c*sin(f*x + e))*sin(f*x + e) + sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e))/((a*c^2*f*cos(f*x + e) - a*c^2*f)*sin(f*x + e))]`

3.137.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))^{3/2}} dx = \int \frac{\sec(e + fx)}{\sqrt{a(\sec(e + fx) + 1)}(-c(\sec(e + fx) - 1))^{3/2}} dx$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))**(3/2)/(a+a*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(a*(sec(e + f*x) + 1))*(-c*(sec(e + f*x) - 1))**(3/2)), x)`

3.137.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 406 vs. $2(83) = 166$.

Time = 0.38 (sec) , antiderivative size = 406, normalized size of antiderivative = 4.27

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))^{3/2}} dx = \frac{((2(2 \cos(fx + e) - 1) \cos(2fx + 2e) - \cos(2fx + 2e))}{\dots}$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x, algo rithm="maxima")`

output $1/2*((2*(2*\cos(f*x + e) - 1)*\cos(2*f*x + 2*e) - \cos(2*f*x + 2*e)^2 - 4*\cos(f*x + e)^2 - \sin(2*f*x + 2*e)^2 + 4*\sin(2*f*x + 2*e)*\sin(f*x + e) - 4*\sin(f*x + e)^2 + 4*\cos(f*x + e) - 1)*\arctan2(\sin(f*x + e), \cos(f*x + e) + 1) - (2*(2*\cos(f*x + e) - 1)*\cos(2*f*x + 2*e) - \cos(2*f*x + 2*e)^2 - 4*\cos(f*x + e)^2 - \sin(2*f*x + 2*e)^2 + 4*\sin(2*f*x + 2*e)*\sin(f*x + e) - 4*\sin(f*x + e)^2 + 4*\cos(f*x + e) - 1)*\arctan2(\sin(f*x + e), \cos(f*x + e) - 1) + 2*\cos(f*x + e)*\sin(2*f*x + 2*e) - 2*\cos(2*f*x + 2*e)*\sin(f*x + e) - 2*\sin(f*x + e)*\sqrt{a}*\sqrt{c}/((a*c^2*\cos(2*f*x + 2*e)^2 + 4*a*c^2*\cos(f*x + e)^2 + a*c^2*\sin(2*f*x + 2*e)^2 - 4*a*c^2*\sin(2*f*x + 2*e)*\sin(f*x + e) + 4*a*c^2*\sin(f*x + e)^2 - 4*a*c^2*\cos(f*x + e) + a*c^2 - 2*(2*a*c^2*\cos(f*x + e) - a*c^2)*\cos(2*f*x + 2*e))*f)$

3.137.8 Giac [A] (verification not implemented)

Time = 1.88 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.97

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))^{3/2}} dx = \frac{\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2} - \log\left(|c| \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2\right) + \log(|c|)}{4 \sqrt{-ac} f |c| \operatorname{sgn}\left(\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)\right)}$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2),x, algorith="giac")`

output $1/4*((c*\tan(1/2*f*x + 1/2*e)^2 - c)/(c*\tan(1/2*f*x + 1/2*e)^2) - \log(\operatorname{abs}(c)*\tan(1/2*f*x + 1/2*e)^2) + \log(\operatorname{abs}(c)))/(\sqrt{-a*c}*f*\operatorname{abs}(c)*\operatorname{sgn}(\tan(1/2*f*x + 1/2*e)^3 + \tan(1/2*f*x + 1/2*e)))$

3.137.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))^{3/2}} dx = \int \frac{1}{\cos(e + fx) \sqrt{a + \frac{a}{\cos(e + fx)}} \left(c - \frac{c}{\cos(e + fx)}\right)^{3/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(3/2)),x)`

3.137. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}} dx$

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(3/2))
, x)`

3.137. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}} dx$

3.138
$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{5/2}} dx$$

3.138.1 Optimal result	982
3.138.2 Mathematica [A] (verified)	982
3.138.3 Rubi [A] (verified)	983
3.138.4 Maple [A] (verified)	985
3.138.5 Fricas [A] (verification not implemented)	986
3.138.6 Sympy [F]	986
3.138.7 Maxima [B] (verification not implemented)	987
3.138.8 Giac [A] (verification not implemented)	988
3.138.9 Mupad [F(-1)]	988

3.138.1 Optimal result

Integrand size = 36, antiderivative size = 140

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{5/2}} dx =$$

$$-\frac{\tan(e+fx)}{4f\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{5/2}}$$

$$-\frac{\tan(e+fx)}{4cf\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{3/2}}$$

$$-\frac{\operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{4c^2f\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}}$$

output

```
-1/4*tan(f*x+e)/f/(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2)-1/4*tan(f*x+e)/c/f/(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(1/2)-1/4*arctanh(cos(f*x+e))*tan(f*x+e)/c^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)
```

3.138.2 Mathematica [A] (verified)

Time = 0.98 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.59

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{5/2}} dx =$$

$$-\frac{(2 + \operatorname{arctanh}(\sec(e+fx))(-1 + \sec(e+fx))^2 - \sec(e+fx)) \tan(e+fx)}{4c^2f(-1 + \sec(e+fx))^2\sqrt{a(1 + \sec(e+fx))}\sqrt{c-c \sec(e+fx)}}$$

3.138.
$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{5/2}} dx$$

input `Integrate[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(5/2)),x]`

output `-1/4*((2 + ArcTanh[Sec[e + f*x]]*(-1 + Sec[e + f*x])^2 - Sec[e + f*x])*Tan[e + f*x])/(c^2*f*(-1 + Sec[e + f*x])^2*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.138.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4448, 3042, 4448, 3042, 4447, 25, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{\sqrt{a \sec(e+fx) + a(c - c \sec(e+fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})}{\sqrt{a \csc(e+fx + \frac{\pi}{2}) + a(c - c \csc(e+fx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4448} \\
 & \frac{\int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a + a(c - c \sec(e+fx))^{3/2}} dx}{2c} - \frac{\tan(e+fx)}{4f \sqrt{a \sec(e+fx) + a(c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx + \frac{\pi}{2})}{\sqrt{\csc(e+fx + \frac{\pi}{2})a + a(c - c \csc(e+fx + \frac{\pi}{2}))^{3/2}} dx}{2c} - \frac{\tan(e+fx)}{4f \sqrt{a \sec(e+fx) + a(c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{4448} \\
 & \frac{\int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a + a \sqrt{c - c \sec(e+fx)}} dx}{2c} - \frac{\tan(e+fx)}{2f \sqrt{a \sec(e+fx) + a(c - c \sec(e+fx))^{3/2}} - \\
 & \quad \frac{2c \tan(e+fx)}{4f \sqrt{a \sec(e+fx) + a(c - c \sec(e+fx))^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.138. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}} dx \\
& \frac{2c}{2f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{3/2}}} - \frac{\tan(e+fx)}{4f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{5/2}}} \\
& \quad \downarrow 4447 \\
& \frac{\tan(e+fx) \int -\csc(e+fx) dx}{2c\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}} - \frac{\tan(e+fx)}{2f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{3/2}}} \\
& \frac{2c}{4f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{5/2}}} \\
& \quad \downarrow 25 \\
& \frac{\tan(e+fx) \int \csc(e+fx) dx}{2c\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}} - \frac{\tan(e+fx)}{2f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{3/2}}} \\
& \frac{2c}{4f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{5/2}}} \\
& \quad \downarrow 3042 \\
& \frac{\tan(e+fx) \int \csc(e+fx) dx}{2c\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}} - \frac{\tan(e+fx)}{2f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{3/2}}} \\
& \frac{2c}{4f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{5/2}}} \\
& \quad \downarrow 4257 \\
& \frac{\tan(e+fx)\operatorname{arctanh}(\cos(e+fx))}{2cf\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}} - \frac{\tan(e+fx)}{2f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{3/2}}} \\
& \frac{2c}{4f\sqrt{a\sec(e+fx)+a(c-c\sec(e+fx))^{5/2}}}
\end{aligned}$$

input `Int[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(5/2)),x]`

output `-1/4*Tan[e + f*x]/(f*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(5/2)) + (-1/2*Tan[e + f*x]/(f*Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])^(3/2))) - (ArcTanh[Cos[e + f*x]]*Tan[e + f*x])/(2*c*f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])/(2*c)`

3.138. $\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}} dx$

3.138.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

```
rule 4447 Int[csc[(e_) + (f_)*(x_)*(csc[(e_) + (f_)*(x_)*(b_) + (a_)^(m_)*(cs
c[(e_) + (f_)*(x_)*(d_) + (c_)^(m_)], x_Symbol] := Simp[((-a)*c)^(m + 1
/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) In
t[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] &
& EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]
```

```
rule 4448 Int[csc[(e_) + (f_)*(x_)*(csc[(e_) + (f_)*(x_)*(b_) + (a_)^(m_)*(cs
c[(e_) + (f_)*(x_)*(d_) + (c_)^(n_)], x_Symbol] := Simp[b*Cot[e + f*x]*
(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[
(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(
c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (IL
tQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.138.4 Maple [A] (verified)

Time = 3.64 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.12

method	result
default	$\frac{(4 \cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e))-8 \cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e))-5 \cos(fx+e)^2+4 \ln(-\cot(fx+e)+\csc(fx+e)))}{16fa(\sec(fx+e)-1)^2 \sqrt{-c(\sec(fx+e)-1)} c^2(\cos(fx+e)+1)}$
risch	$\frac{i(3e^{2i(fx+e)}-4e^{i(fx+e)}+3)(e^{2i(fx+e)}+e^{i(fx+e)})}{2c^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)^3 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} (1+e^{2i(fx+e)})f} + \frac{i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1) \ln(e^{i(fx+e)}+1)}{4c^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} (1+e^{2i(fx+e)})}$

```
input int(sec(f*x+e)/(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x,method=_RET
URNVERBOSE)
```

$$3.138. \int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))^{5/2}} dx$$

output $1/16/f/a*(4*\cos(f*x+e)^2*\ln(-\cot(f*x+e)+\csc(f*x+e))-8*\cos(f*x+e)*\ln(-\cot(f*x+e)+\csc(f*x+e))-5*\cos(f*x+e)^2+4*\ln(-\cot(f*x+e)+\csc(f*x+e))-2*\cos(f*x+e)+3)*(a*(\sec(f*x+e)+1))^{(1/2)}/(\sec(f*x+e)-1)^{2}/(-c*(\sec(f*x+e)-1))^{(1/2)}/c^{2}/(\cos(f*x+e)+1)*\tan(f*x+e)*\sec(f*x+e)$

3.138.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 456, normalized size of antiderivative = 3.26

$$\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}} dx = \left[\frac{\sqrt{-ac}(\cos(fx+e)^2 - 2\cos(fx+e) + 1) \log\left(-\frac{4}{2}\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x, algo rithm="fricas")`

output `[-1/8*(sqrt(-a*c)*(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)*log(-4*(2*sqrt(-a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2 + (a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e))/((cos(f*x + e)^2 - 1)*sin(f*x + e))*sin(f*x + e) - 2*(3*cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a*c^3*f*cos(f*x + e)^2 - 2*a*c^3*f*cos(f*x + e) + a*c^3*f)*sin(f*x + e)), 1/4*(sqrt(a*c)*(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)*arc tan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(a*c*sin(f*x + e))*sin(f*x + e) + (3*cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a*c^3*f*cos(f*x + e)^2 - 2*a*c^3*f*cos(f*x + e) + a*c^3*f)*sin(f*x + e))]`

3.138.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\sec(e+fx)}{\sqrt{a(\sec(e+fx)+1)}(-c(\sec(e+fx)-1))^{5/2}} dx$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))**(5/2)/(a+a*sec(f*x+e))**(1/2),x)`

3.138. $\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}} dx$

output `Integral(sec(e + f*x)/(sqrt(a*(sec(e + f*x) + 1))*(-c*(sec(e + f*x) - 1))*
*(5/2)), x)`

3.138.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1201 vs. $2(122) = 244$.

Time = 0.43 (sec) , antiderivative size = 1201, normalized size of antiderivative = 8.58

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x, algo
rithm="maxima")`

output `1/4*((2*(4*cos(3*f*x + 3*e) - 6*cos(2*f*x + 2*e) + 4*cos(f*x + e) - 1)*cos
(4*f*x + 4*e) - cos(4*f*x + 4*e)^2 + 8*(6*cos(2*f*x + 2*e) - 4*cos(f*x + e
) + 1)*cos(3*f*x + 3*e) - 16*cos(3*f*x + 3*e)^2 + 12*(4*cos(f*x + e) - 1)*
cos(2*f*x + 2*e) - 36*cos(2*f*x + 2*e)^2 - 16*cos(f*x + e)^2 + 4*(2*sin(3*
f*x + 3*e) - 3*sin(2*f*x + 2*e) + 2*sin(f*x + e))*sin(4*f*x + 4*e) - sin(4
*f*x + 4*e)^2 + 16*(3*sin(2*f*x + 2*e) - 2*sin(f*x + e))*sin(3*f*x + 3*e)
- 16*sin(3*f*x + 3*e)^2 - 36*sin(2*f*x + 2*e)^2 + 48*sin(2*f*x + 2*e)*sin(
f*x + e) - 16*sin(f*x + e)^2 + 8*cos(f*x + e) - 1)*arctan2(sin(f*x + e), c
os(f*x + e) + 1) - (2*(4*cos(3*f*x + 3*e) - 6*cos(2*f*x + 2*e) + 4*cos(f*x
+ e) - 1)*cos(4*f*x + 4*e) - cos(4*f*x + 4*e)^2 + 8*(6*cos(2*f*x + 2*e) -
4*cos(f*x + e) + 1)*cos(3*f*x + 3*e) - 16*cos(3*f*x + 3*e)^2 + 12*(4*cos(
f*x + e) - 1)*cos(2*f*x + 2*e) - 36*cos(2*f*x + 2*e)^2 - 16*cos(f*x + e)^2
+ 4*(2*sin(3*f*x + 3*e) - 3*sin(2*f*x + 2*e) + 2*sin(f*x + e))*sin(4*f*x
+ 4*e) - sin(4*f*x + 4*e)^2 + 16*(3*sin(2*f*x + 2*e) - 2*sin(f*x + e))*sin
(3*f*x + 3*e) - 16*sin(3*f*x + 3*e)^2 - 36*sin(2*f*x + 2*e)^2 + 48*sin(2*f
*x + 2*e)*sin(f*x + e) - 16*sin(f*x + e)^2 + 8*cos(f*x + e) - 1)*arctan2(s
in(f*x + e), cos(f*x + e) - 1) - 2*(3*sin(3*f*x + 3*e) - 4*sin(2*f*x + 2*e
) + 3*sin(f*x + e))*cos(4*f*x + 4*e) + 2*(3*cos(3*f*x + 3*e) - 4*cos(2*f*x
+ 2*e) + 3*cos(f*x + e))*sin(4*f*x + 4*e) - 2*(2*cos(2*f*x + 2*e) + 3)*si
n(3*f*x + 3*e) + 4*(cos(f*x + e) + 2)*sin(2*f*x + 2*e) + 4*cos(3*f*x + ...`

3.138.8 Giac [A] (verification not implemented)

Time = 1.71 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.87

$$\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}} dx = \frac{3\left(c\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^2-c\right)^2+2\left(c\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^2-c\right)c}{c^2\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^4} - 2\log\left(|c|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)\right) - \frac{1}{16\sqrt{-accf}|c|\operatorname{sgn}\left(\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3+\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)\right)}$$

input `integrate(sec(f*x+e)/(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(1/2),x, algo rithm="giac")`

output `1/16*((3*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2 + 2*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c)/(c^2*tan(1/2*f*x + 1/2*e)^4) - 2*log(abs(c)*tan(1/2*f*x + 1/2*e)^2) + 2*log(abs(c)))/(sqrt(-a*c)*c*f*abs(c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e)))`

3.138.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))^{5/2}} dx = \int \frac{1}{\cos(e+fx)\sqrt{a+\frac{a}{\cos(e+fx)}}\left(c-\frac{c}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(5/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(5/2)), x)`

3.139
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx$$

3.139.1 Optimal result	989
3.139.2 Mathematica [A] (verified)	989
3.139.3 Rubi [A] (verified)	990
3.139.4 Maple [A] (verified)	992
3.139.5 Fricas [F]	992
3.139.6 Sympy [F(-1)]	993
3.139.7 Maxima [B] (verification not implemented)	993
3.139.8 Giac [A] (verification not implemented)	994
3.139.9 Mupad [F(-1)]	995

3.139.1 Optimal result

Integrand size = 36, antiderivative size = 142

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{4c^3 \log(1+\sec(e+fx)) \tan(e+fx)}{af\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} + \frac{2c^2\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{af\sqrt{a+a\sec(e+fx)}} + \frac{c(c-c\sec(e+fx))^{3/2}\tan(e+fx)}{f(a+a\sec(e+fx))^{3/2}}$$

output `c*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(3/2)+4*c^3*ln(1+sec(f*x+e))*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)+2*c^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^(1/2)`

3.139.2 Mathematica [A] (verified)

Time = 0.64 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.59

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{c\left(-4c^2 \log(1+\sec(e+fx)) + c^2 \sec(e+fx) - \frac{4c^2}{1+\sec(e+fx)}\right) \tan(e+fx)}{af\sqrt{a(1+\sec(e+fx))}\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^(3/2), x]`

output $-\left(\left(c*(-4*c^2*\text{Log}[1 + \text{Sec}[e + f*x]] + c^2*\text{Sec}[e + f*x] - (4*c^2)/(1 + \text{Sec}[e + f*x]))*\text{Tan}[e + f*x]\right)/\left(a*f*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]\right)*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]\right)$

3.139.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4442, 3042, 4443, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a\sec(e+fx)+a)^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}} dx$$

↓ 4442

$$\frac{c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)^{3/2}} - \frac{2c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{\sqrt{\sec(e+fx)a+a}} dx}{a}$$

↓ 3042

$$\frac{c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)^{3/2}} - \frac{2c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx}{a}$$

↓ 4443

$$\frac{c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)^{3/2}} - \frac{2c \left(2c \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} dx - \frac{c \tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f\sqrt{a\sec(e+fx)+a}} \right)}{a}$$

↓ 3042

$$\frac{c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{f(a\sec(e+fx)+a)^{3/2}} - \frac{2c \left(2c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx - \frac{c \tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f\sqrt{a\sec(e+fx)+a}} \right)}{a}$$

3.139. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx$

$$\frac{c \tan(e + fx)(c - c \sec(e + fx))^{3/2}}{f(a \sec(e + fx) + a)^{3/2}} - \frac{2c \left(-\frac{2c^2 \tan(e+fx) \log(\sec(e+fx)+1)}{f \sqrt{a \sec(e+fx)+a} \sqrt{c-c \sec(e+fx)}} - \frac{c \tan(e+fx) \sqrt{c-c \sec(e+fx)}}{f \sqrt{a \sec(e+fx)+a}} \right)}{a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^(3/2),x]`

output `(c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])^(3/2)) - (2*c*((-2*c^2*Log[1 + Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]])*Sqrt[c - c*Sec[e + f*x]]) - (c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]])))/a`

3.139.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`


```
rule 4443 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(c
sc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.139.4 Maple [A] (verified)

Time = 3.68 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.35

method	result
default	$-\left(4 \cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e)-1)+4 \cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e)+1)+\sin(fx+e)^2+4 \cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e)-1)+4 \cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e)+1)\right)$
risch	$\frac{2ic^2 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} (5e^{3i(fx+e)}+2e^{2i(fx+e)}+5e^{i(fx+e)})}{a(e^{i(fx+e)}+1) \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f(1+e^{2i(fx+e)})} - \frac{8ic^2(e^{i(fx+e)}+1) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} \ln(e^{i(fx+e)}+1)}{a \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f} + \frac{4ic^2(e^{i(fx+e)}-1) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} \ln(e^{i(fx+e)}-1)}{a \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f}$

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(3/2),x,method=_RET
URNVERBOSE)
```

```
output -1/f/a^2*(4*cos(f*x+e)^2*ln(-cot(f*x+e)+csc(f*x+e)-1)+4*cos(f*x+e)^2*ln(-c
ot(f*x+e)+csc(f*x+e)+1)+sin(f*x+e)^2+4*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e
)-1)+4*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)+4*cos(f*x+e))*(-c*(sec(f*x+
e)-1))^(1/2)*(sec(f*x+e)-1)^2*c^2*(a*(sec(f*x+e)+1))^(1/2)/(cos(f*x+e)-1)*
cot(f*x+e)^2*csc(f*x+e)
```

3.139.5 Fracas [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx = \int \frac{(-c\sec(fx+e)+c)^{5/2} \sec(fx+e)}{(a\sec(fx+e)+a)^{3/2}} dx$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(3/2),x, algo
rithm="fracas")
```

output `integral((c^2*sec(f*x + e)^3 - 2*c^2*sec(f*x + e)^2 + c^2*sec(f*x + e))*sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(a^2*sec(f*x + e)^2 + 2*a^2*sec(f*x + e) + a^2), x)`

3.139.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^{5/2}}{(a + a\sec(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(5/2)/(a+a*sec(f*x+e))**(3/2),x)`

output `Timed out`

3.139.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2035 vs. 2(130) = 260.

Time = 0.49 (sec) , antiderivative size = 2035, normalized size of antiderivative = 14.33

$$\int \frac{\sec(e + fx)(c - c\sec(e + fx))^{5/2}}{(a + a\sec(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(3/2),x, algorithm="maxima")`

output

```
-2*(8*c^2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)*cos(3/2*arctan2(
sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 8*c^2*arctan2(sin(2*f*x + 2*e), c
os(2*f*x + 2*e) + 1)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^
2 + 8*c^2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)*sin(3/2*arctan2(
sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 8*c^2*arctan2(sin(2*f*x + 2*e), c
os(2*f*x + 2*e) + 1)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^
2 - 2*c^2*cos(2*f*x + 2*e)*sin(4*f*x + 4*e) + 2*c^2*cos(4*f*x + 4*e)*sin(2
*f*x + 2*e) + 2*c^2*sin(2*f*x + 2*e) + 2*(c^2*cos(4*f*x + 4*e)^2 + 4*c^2*c
os(2*f*x + 2*e)^2 + c^2*sin(4*f*x + 4*e)^2 + 4*c^2*sin(4*f*x + 4*e)*sin(2*
f*x + 2*e) + 4*c^2*sin(2*f*x + 2*e)^2 + 4*c^2*cos(2*f*x + 2*e) + c^2 + 2*(
2*c^2*cos(2*f*x + 2*e) + c^2)*cos(4*f*x + 4*e))*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e) + 1) - 4*(c^2*cos(4*f*x + 4*e)^2 + 4*c^2*cos(2*f*x + 2*e)
^2 + 4*c^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 4*c^2*
cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + c^2*sin(4*f*x + 4
*e)^2 + 4*c^2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*c^2*sin(2*f*x + 2*e)^2
+ 4*c^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 4*c^2*si
n(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 4*c^2*cos(2*f*x + 2
*e) + c^2 + 2*(2*c^2*cos(2*f*x + 2*e) + c^2)*cos(4*f*x + 4*e) + 4*(c^2*cos
(4*f*x + 4*e) + 2*c^2*cos(2*f*x + 2*e) + 2*c^2*cos(1/2*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e)))) + c^2)*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(...
```

3.139.8 Giac [A] (verification not implemented)

Time = 1.53 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.13

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx =$$

$$\frac{2c^2 \left(\frac{2\sqrt{-acc} \log\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)}{a^2|c|} + \frac{\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)\sqrt{-ac}}{a^2|c|} - \frac{2\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)\sqrt{-acc} + \sqrt{-acc^2}}{\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)a^2|c|} \right) \operatorname{sgn}\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(3/2),x, algo
rithm="giac")`

output

```
-2*c^2*(2*sqrt(-a*c)*c*log(c*tan(1/2*f*x + 1/2*e)^2 - c)/(a^2*abs(c)) + (c
*tan(1/2*f*x + 1/2*e)^2 - c)*sqrt(-a*c)/(a^2*abs(c)) - (2*(c*tan(1/2*f*x +
1/2*e)^2 - c)*sqrt(-a*c)*c + sqrt(-a*c)*c^2)/((c*tan(1/2*f*x + 1/2*e)^2 -
c)*a^2*abs(c)))*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))/f
```

3.139. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx$

3.139.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{3/2}} dx = \int \frac{\left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)),x)`

output `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)),x)`

$$3.140 \quad \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx$$

3.140.1 Optimal result	996
3.140.2 Mathematica [A] (verified)	996
3.140.3 Rubi [A] (verified)	997
3.140.4 Maple [A] (verified)	998
3.140.5 Fricas [F]	999
3.140.6 Sympy [F]	999
3.140.7 Maxima [A] (verification not implemented)	999
3.140.8 Giac [A] (verification not implemented)	1000
3.140.9 Mupad [F(-1)]	1000

3.140.1 Optimal result

Integrand size = 36, antiderivative size = 95

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{c^2 \log(1+\sec(e+fx)) \tan(e+fx)}{af\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} + \frac{c\sqrt{c-c\sec(e+fx)} \tan(e+fx)}{f(a+a\sec(e+fx))^{3/2}}$$

output `c^2*ln(1+sec(f*x+e))*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)+c*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(3/2)`

3.140.2 Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{c\left(-c\log(1+\sec(e+fx)) - \frac{2c}{1+\sec(e+fx)}\right) \tan(e+fx)}{af\sqrt{a(1+\sec(e+fx))}\sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x])^(3/2), x]`

output $-\left(\left(c\left(-\left(c\log\left[1+\sec\left[e+fx\right]\right)\right)-\left(2c\right)/\left(1+\sec\left[e+fx\right]\right)\right)\tan\left[e+fx\right]\right)/\left(a\sqrt{a\sec\left[e+fx\right]}\sqrt{c-c\sec\left[e+fx\right]}\right)$

3.140.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4442, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a\sec(e+fx)+a)^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(c-c\csc\left(e+fx+\frac{\pi}{2}\right)\right)^{3/2}}{\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^{3/2}} dx$$

↓ 4442

$$\frac{c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)^{3/2}} - \frac{c\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} dx}{a}$$

↓ 3042

$$\frac{c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)^{3/2}} - \frac{c\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)}}{\sqrt{\csc\left(e+fx+\frac{\pi}{2}\right)a+a}} dx}{a}$$

↓ 4440

$$\frac{c^2\tan(e+fx)\log(\sec(e+fx)+1)}{af\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} + \frac{c\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)^{3/2}}$$

input $\text{Int}[(\sec[e+fx]*(c-c*\sec[e+fx])^(3/2))/(a+a*\sec[e+fx])^(3/2),x]$

output $(c^2*\log[1+\sec[e+fx]]*\tan[e+fx])/(a*\sqrt{a+a*\sec[e+fx]}*\sqrt{c-c*\sec[e+fx]})+(c*\sqrt{c-c*\sec[e+fx]}*\tan[e+fx])/(f*(a+a*\sec[e+fx])^(3/2))$

3.140. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx$

3.140.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4442 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^n, x_Symbol] := Simp[2*a*c*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))), x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[m, -2^(-1)]`

3.140.4 Maple [A] (verified)

Time = 3.38 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.56

method	result
default	$\frac{\sqrt{-c(\sec(fx+e)-1)}(\sec(fx+e)-1)c\sqrt{a(\sec(fx+e)+1)}(\cos(fx+e)\ln(-\cot(fx+e)+\csc(fx+e)-1)+\cos(fx+e)\ln(-\cot(fx+e)+\csc(fx+e)-1))}{fa^2}$
risch	$-\frac{ic\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}(2e^{2i(fx+e)}\ln(e^{i(fx+e)}+1)-e^{2i(fx+e)}\ln(1+e^{2i(fx+e)})+4e^{i(fx+e)}\ln(e^{i(fx+e)}+1)-2e^{i(fx+e)}\ln(1+e^{2i(fx+e)}))}{a(e^{i(fx+e)}+1)\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)f}$

input `int(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `1/f/a^2*(-c*(sec(f*x+e)-1))^(1/2)*(sec(f*x+e)-1)*c*(a*(sec(f*x+e)+1))^(1/2)*(cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)-1)+cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)-cos(f*x+e)+ln(-cot(f*x+e)+csc(f*x+e)-1)+ln(-cot(f*x+e)+csc(f*x+e)+1)+1)*cot(f*x+e)^2*csc(f*x+e)`

3.140. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx$

3.140.5 Fracas [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx = \int \frac{(-c\sec(fx+e)+c)^{3/2} \sec(fx+e)}{(a\sec(fx+e)+a)^{3/2}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(3/2),x, algorith="fricas")`

output `integral(-(c*sec(f*x + e)^2 - c*sec(f*x + e))*sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(a^2*sec(f*x + e)^2 + 2*a^2*sec(f*x + e) + a^2), x)`

3.140.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx = \int \frac{(-c(\sec(e+fx)-1))^{3/2} \sec(e+fx)}{(a(\sec(e+fx)+1))^{3/2}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(3/2)/(a+a*sec(f*x+e))**(3/2),x)`

output `Integral((-c*(sec(e + f*x) - 1))**(3/2)*sec(e + f*x)/(a*(sec(e + f*x) + 1))**(3/2), x)`

3.140.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.04

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{c^{3/2} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}+1\right)}{\sqrt{-aa}} + \frac{c^{3/2} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}-1\right)}{\sqrt{-aa}} + \frac{c^{3/2} \sin(fx+e)^2}{\sqrt{-aa}(\cos(fx+e)+1)^2}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(3/2),x, algorith="maxima")`

output `(c^(3/2)*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/(sqrt(-a)*a) + c^(3/2)*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/(sqrt(-a)*a) + c^(3/2)*sin(f*x + e)^2/(sqrt(-a)*a*(cos(f*x + e) + 1)^2))/f`

3.140. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx$

3.140.8 Giac [A] (verification not implemented)

Time = 1.50 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.83

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 + c \log\left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right) - c\right) c^2 \operatorname{sgn}\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)}{\sqrt{-aca}f|c|}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(3/2),x, algorithm="giac")`

output `(c*tan(1/2*f*x + 1/2*e)^2 + c*log(c*tan(1/2*f*x + 1/2*e)^2 - c) - c)*c^2*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))/(sqrt(-a*c)*a*f*abs(c))`

3.140.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{3/2}} dx = \int \frac{\left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)),x)`

output `int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)),x)`

3.141
$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{3/2}} dx$$

3.141.1 Optimal result 1001
 3.141.2 Mathematica [A] (verified) 1001
 3.141.3 Rubi [A] (verified) 1002
 3.141.4 Maple [A] (verified) 1003
 3.141.5 Fricas [B] (verification not implemented) 1003
 3.141.6 Sympy [F] 1004
 3.141.7 Maxima [A] (verification not implemented) 1004
 3.141.8 Giac [A] (verification not implemented) 1004
 3.141.9 Mupad [B] (verification not implemented) 1005

3.141.1 Optimal result

Integrand size = 36, antiderivative size = 42

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{2f(a+a\sec(e+fx))^{3/2}}$$

output `1/2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(3/2)`

3.141.2 Mathematica [A] (verified)

Time = 1.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{\csc(e+fx)\sqrt{c-c\sec(e+fx)}}{af\sqrt{a(1+\sec(e+fx))}}$$

input `Integrate[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^(3/2),x]`

output `(Csc[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a*f*Sqrt[a*(1 + Sec[e + f*x])])`

3.141.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a\sec(e+fx)+a)^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}} dx$$

↓ 4438

$$\frac{\tan(e+fx)\sqrt{c-c\sec(e+fx)}}{2f(a\sec(e+fx)+a)^{3/2}}$$

input `Int[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^(3/2),x]`

output `(Sqrt[c - c*Sec[e + f*x])*Tan[e + f*x]/(2*f*(a + a*Sec[e + f*x])^(3/2))`

3.141.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.141.4 Maple [A] (verified)

Time = 3.57 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.33

method	result	size
default	$\frac{\sin(fx+e)\sqrt{-c(\sec(fx+e)-1)}\sqrt{a(\sec(fx+e)+1)}\cos(fx+e)}{2fa^2(\cos(fx+e)+1)^2}$	56
risch	$\frac{2i\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}e^{i(fx+e)}}{a(e^{i(fx+e)}+1)\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)f}$	105

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/2/f/a^2*sin(f*x+e)*(-c*(sec(f*x+e)-1))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)*co
s(f*x+e)/(cos(f*x+e)+1)^2
```

3.141.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. 2(36) = 72.

Time = 0.26 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.86

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\cos(fx+e)}{(a^2f\cos(fx+e)+a^2f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(3/2),x, algo
rithm="fracas")
```

```
output sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x
+ e))*cos(f*x + e)/((a^2*f*cos(f*x + e) + a^2*f)*sin(f*x + e))
```

3.141.6 Sympy [F]

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{(a + a \sec(e + fx))^{3/2}} dx = \int \frac{\sqrt{-c(\sec(e + fx) - 1)} \sec(e + fx)}{(a(\sec(e + fx) + 1))^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(1/2)/(a+a*sec(f*x+e))**(3/2),x)`

output `Integral(sqrt(-c*(sec(e + f*x) - 1))*sec(e + f*x)/(a*(sec(e + f*x) + 1))**(3/2), x)`

3.141.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.29

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{(a + a \sec(e + fx))^{3/2}} dx = \frac{\sqrt{c} \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right) \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)}{2 \sqrt{-aaf}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(3/2),x, algo rithm="maxima")`

output `1/2*sqrt(c)*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/(sqrt(-a)*a*f)`

3.141.8 Giac [A] (verification not implemented)

Time = 1.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{\sec(e + fx) \sqrt{c - c \sec(e + fx)}}{(a + a \sec(e + fx))^{3/2}} dx = \frac{\left(c \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - c \right) c}{2 \sqrt{-aca} f |c|}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(3/2),x, algo rithm="giac")`

output `1/2*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c/(sqrt(-a*c)*a*f*abs(c))`

3.141. $\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{3/2}} dx$

3.141.9 Mupad [B] (verification not implemented)

Time = 14.63 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{3/2}} dx = \frac{\sqrt{c-\frac{c}{\cos(e+fx)}}}{af \sin(e+fx) \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}}}$$

input `int((c - c/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)),x)`

output `(c - c/cos(e + f*x))^(1/2)/(a*f*sin(e + f*x)*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2))`

$$3.142 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx$$

3.142.1 Optimal result	1006
3.142.2 Mathematica [A] (verified)	1006
3.142.3 Rubi [A] (verified)	1007
3.142.4 Maple [A] (verified)	1009
3.142.5 Fracas [B] (verification not implemented)	1009
3.142.6 Sympy [F]	1010
3.142.7 Maxima [B] (verification not implemented)	1010
3.142.8 Giac [A] (verification not implemented)	1011
3.142.9 Mupad [F(-1)]	1011

3.142.1 Optimal result

Integrand size = 36, antiderivative size = 95

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx = \frac{\tan(e+fx)}{2f(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} - \frac{\operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{2af \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}}$$

output `1/2*tan(f*x+e)/f/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2)-1/2*arctanh(cos(f*x+e))*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.142.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.63

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx = \frac{(-1 + \operatorname{arctanh}(\sec(e+fx))(1 + \sec(e+fx))) \tan(e+fx)}{2f(a(1 + \sec(e+fx)))^{3/2} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^(3/2)*Sqrt[c - c*Sec[e + f*x]]),x]`

output `-1/2*((-1 + ArcTanh[Sec[e + f*x]]*(1 + Sec[e + f*x]))*Tan[e + f*x])/(f*(a*(1 + Sec[e + f*x]))^(3/2)*Sqrt[c - c*Sec[e + f*x]])`

$$3.142. \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx$$

3.142.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 4448, 3042, 4447, 25, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^{3/2} \sqrt{c - c \csc(e+fx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4448} \\
 & \frac{\int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a+a\sqrt{c-c\sec(e+fx)}}} dx}{2a} + \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx + \frac{\pi}{2})}{\sqrt{\csc(e+fx + \frac{\pi}{2})a+a\sqrt{c-c\csc(e+fx + \frac{\pi}{2})}}} dx}{2a} + \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{4447} \\
 & \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} - \frac{\tan(e+fx) \int -\csc(e+fx) dx}{2a \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\tan(e+fx) \int \csc(e+fx) dx}{2a \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} + \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tan(e+fx) \int \csc(e+fx) dx}{2a \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} + \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{4257} \\
 & \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} - \frac{\tan(e+fx) \operatorname{arctanh}(\cos(e+fx))}{2af \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}
 \end{aligned}$$

3.142. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^(3/2)*Sqrt[c - c*Sec[e + f*x]]),x]`

output `Tan[e + f*x]/(2*f*(a + a*Sec[e + f*x])^(3/2)*Sqrt[c - c*Sec[e + f*x]]) - (ArcTanh[Cos[e + f*x]]*Tan[e + f*x])/(2*a*f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.142.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4447 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(m_), x_Symbol] := Simp[((-a)*c)^(m + 1/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Int[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]`

rule 4448 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))`

3.142.4 Maple [A] (verified)

Time = 3.58 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.05

method	result
default	$\frac{\sin(fx+e)(2\cos(fx+e)\ln(-\cot(fx+e)+\csc(fx+e))+2\ln(-\cot(fx+e)+\csc(fx+e))+\cos(fx+e)-1)\sqrt{a(\sec(fx+e)+1)}}{4fa^2(\cos(fx+e)+1)^2\sqrt{-c(\sec(fx+e)-1)}}$
risch	$i(\ln(e^{i(fx+e)}+1)e^{3i(fx+e)}-e^{3i(fx+e)}\ln(e^{i(fx+e)}-1)+e^{2i(fx+e)}\ln(e^{i(fx+e)}+1)-e^{2i(fx+e)}\ln(e^{i(fx+e)}-1)-e^{i(fx+e)}\ln(e^{i(fx+e)}+1)+e^{i(fx+e)}\ln(e^{i(fx+e)}-1))\sqrt{2a(e^{i(fx+e)}+1)}\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 1/4/f/a^2*sin(f*x+e)*(2*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e))+2*ln(-cot(f*
x+e)+csc(f*x+e))+cos(f*x+e)-1)*(a*(sec(f*x+e)+1))^(1/2)/(cos(f*x+e)+1)^2/(
-c*(sec(f*x+e)-1))^(1/2)
```

3.142.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(83) = 166.

Time = 0.32 (sec) , antiderivative size = 380, normalized size of antiderivative = 4.00

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}\sqrt{c-c\sec(e+fx)}} dx = \left[-\frac{\sqrt{-ac}(\cos(fx+e)+1)\log\left(-\frac{4\left(2\sqrt{-ac}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}\right)}{\dots}\right)}{\dots} \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="fracas")
```

3.142.
$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}\sqrt{c-c\sec(e+fx)}} dx$$

```
output [-1/4*(sqrt(-a*c)*(cos(f*x + e) + 1)*log(-4*(2*sqrt(-a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2 + (a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e)))/((cos(f*x + e)^2 - 1)*sin(f*x + e)))*sin(f*x + e) - 2*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e))/((a^2*c*f*cos(f*x + e) + a^2*c*f)*sin(f*x + e)), 1/2*(sqrt(a*c)*(cos(f*x + e) + 1)*arctan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(a*c*sin(f*x + e)))*sin(f*x + e) + sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e))/((a^2*c*f*cos(f*x + e) + a^2*c*f)*sin(f*x + e))]
```

3.142.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)}} dx = \int \frac{\sec(e + fx)}{(a(\sec(e + fx) + 1))^{3/2} \sqrt{-c(\sec(e + fx) - 1)}} dx$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(1/2),x)
```

```
output Integral(sec(e + f*x)/((a*(sec(e + f*x) + 1))**(3/2)*sqrt(-c*(sec(e + f*x) - 1))), x)
```

3.142.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. $2(83) = 166$.

Time = 0.38 (sec) , antiderivative size = 397, normalized size of antiderivative = 4.18

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)}} dx = \frac{((2(2 \cos(fx + e) + 1) \cos(2fx + 2e) + \cos(2fx + 2e))^2 + 4 \cos(fx + e)^2 + \sin(2fx + 2e)^2 + 4 \sin(2fx + 2e)^2)}{\dots}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x, algo
rithm="maxima")
```

3.142. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx$

output `-1/2*((2*(2*cos(f*x + e) + 1)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + 4*cos(f*x + e)^2 + sin(2*f*x + 2*e)^2 + 4*sin(2*f*x + 2*e)*sin(f*x + e) + 4*sin(f*x + e)^2 + 4*cos(f*x + e) + 1)*arctan2(sin(f*x + e), cos(f*x + e) + 1) - (2*(2*cos(f*x + e) + 1)*cos(2*f*x + 2*e) + cos(2*f*x + 2*e)^2 + 4*cos(f*x + e)^2 + sin(2*f*x + 2*e)^2 + 4*sin(2*f*x + 2*e)*sin(f*x + e) + 4*sin(f*x + e)^2 + 4*cos(f*x + e) + 1)*arctan2(sin(f*x + e), cos(f*x + e) - 1) - 2*cos(f*x + e)*sin(2*f*x + 2*e) + 2*cos(2*f*x + 2*e)*sin(f*x + e) + 2*sin(f*x + e))*sqrt(a)*sqrt(c)/((a^2*c*cos(2*f*x + 2*e)^2 + 4*a^2*c*cos(f*x + e)^2 + a^2*c*sin(2*f*x + 2*e)^2 + 4*a^2*c*sin(2*f*x + 2*e)*sin(f*x + e) + 4*a^2*c*sin(f*x + e)^2 + 4*a^2*c*cos(f*x + e) + a^2*c + 2*(2*a^2*c*cos(f*x + e) + a^2*c)*cos(2*f*x + 2*e))*f)`

3.142.8 Giac [A] (verification not implemented)

Time = 1.90 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)}} dx = \frac{c^2 \left(\frac{\log(|c| \tan(\frac{1}{2} fx + \frac{1}{2} e)^2)}{c} - \frac{\log(|c|)}{c} - \frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}{c^2} \right)}{4 \sqrt{-aca} f |c| \operatorname{sgn} \left(\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2),x, algorith="giac")`

output `-1/4*c^2*(log(abs(c)*tan(1/2*f*x + 1/2*e)^2)/c - log(abs(c))/c - (c*tan(1/2*f*x + 1/2*e)^2 - c)/c^2)/(sqrt(-a*c)*a*f*abs(c)*sgn(tan(1/2*f*x + 1/2*e))^3 + tan(1/2*f*x + 1/2*e))`

3.142.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} \sqrt{c - c \sec(e + fx)}} dx = \int \frac{1}{\cos(e + fx) \left(a + \frac{a}{\cos(e + fx)} \right)^{3/2} \sqrt{c - \frac{c}{\cos(e + fx)}}} dx$$

3.142. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(1/2))
,x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(1/2))
, x)`

3.142. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} dx$

$$3.143 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2}} dx$$

3.143.1 Optimal result	1013
3.143.2 Mathematica [A] (verified)	1013
3.143.3 Rubi [A] (verified)	1014
3.143.4 Maple [A] (verified)	1016
3.143.5 Fracas [A] (verification not implemented)	1016
3.143.6 Sympy [F]	1017
3.143.7 Maxima [B] (verification not implemented)	1017
3.143.8 Giac [A] (verification not implemented)	1018
3.143.9 Mupad [F(-1)]	1018

3.143.1 Optimal result

Integrand size = 36, antiderivative size = 104

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2}} dx = \frac{\csc(e+fx)}{2acf \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} - \frac{\operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{2acf \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}}$$

output `1/2*csc(f*x+e)/a/c/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-1/2*arc
tanh(cos(f*x+e))*tan(f*x+e)/a/c/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(
(1/2)`

3.143.2 Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.62

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2}} dx = \frac{\csc(e+fx) - \operatorname{arctanh}(\sec(e+fx)) \tan(e+fx)}{2acf \sqrt{a(1+\sec(e+fx))} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(3
(/2)),x]`

output `(Csc[e + f*x] - ArcTanh[Sec[e + f*x]]*Tan[e + f*x])/(2*a*c*f*Sqrt[a*(1 + S
ec[e + f*x]])*Sqrt[c - c*Sec[e + f*x]])`

$$3.143. \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{3/2}} dx$$

3.143.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.74, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 4447, 25, 3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)^{3/2}(c-c\sec(e+fx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx$$

$$\downarrow \text{4447}$$

$$\frac{\tan(e+fx) \int -\cot^2(e+fx) \csc(e+fx) dx}{ac\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow \text{25}$$

$$-\frac{\tan(e+fx) \int \cot^2(e+fx) \csc(e+fx) dx}{ac\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow \text{3042}$$

$$-\frac{\tan(e+fx) \int \sec(e+fx-\frac{\pi}{2}) \tan(e+fx-\frac{\pi}{2})^2 dx}{ac\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow \text{3091}$$

$$-\frac{\tan(e+fx) \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx)\csc(e+fx)}{2f} \right)}{ac\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow \text{3042}$$

$$-\frac{\tan(e+fx) \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx)\csc(e+fx)}{2f} \right)}{ac\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow \text{4257}$$

$$-\frac{\tan(e+fx) \left(\frac{\operatorname{arctanh}(\cos(e+fx))}{2f} - \frac{\cot(e+fx)\csc(e+fx)}{2f} \right)}{ac\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

3.143. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{3/2}} dx$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(3/2)),x
]`

output `-(((ArcTanh[Cos[e + f*x]]/(2*f) - (Cot[e + f*x]*Csc[e + f*x])/(2*f))*Tan[e
+ f*x])/(a*c*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]))`

3.143.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3091 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(
n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m
+ n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(
b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] &
& NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]`

rule 4447 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(m_), x_Symbol] := Simp[((-a)*c)^(m + 1
/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) In
t[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] &
& EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]`

3.143.4 Maple [A] (verified)

Time = 3.51 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.11

method	result
default	$-\frac{(\cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e))-\ln(-\cot(fx+e)+\csc(fx+e))-\cos(fx+e))\sqrt{a(\sec(fx+e)+1)}\tan(fx+e)}{2fa^2\sqrt{-c(\sec(fx+e)-1)}c(\sec(fx+e)-1)(\cos(fx+e)+1)^2}$
risch	$-\frac{i(e^{4i(fx+e)}\ln(e^{i(fx+e)}-1)-e^{4i(fx+e)}\ln(e^{i(fx+e)}+1))-2e^{2i(fx+e)}\ln(e^{i(fx+e)}-1)+2e^{2i(fx+e)}\ln(e^{i(fx+e)}+1)-2e^{3i(fx+e)}-2e^{i(fx+e)})}{2ac(e^{i(fx+e)}+1)\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}}f(1+e^{2i(fx+e)})$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2),x,method=_RET
URNVERBOSE)
```

```
output -1/2/f/a^2*(cos(f*x+e)^2*ln(-cot(f*x+e)+csc(f*x+e))-ln(-cot(f*x+e)+csc(f*x
+e))-cos(f*x+e))*(a*(sec(f*x+e)+1))^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)/c/(sec
(f*x+e)-1)/(cos(f*x+e)+1)^2*tan(f*x+e)
```

3.143.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 402, normalized size of antiderivative = 3.87

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{3/2}} dx = \left[-\frac{\sqrt{-ac}(\cos(fx+e)^2-1)\log\left(-\frac{4(2\sqrt{-ac}\sqrt{\frac{a\cos(fx+e)+1}{\cos(fx+e)}})}{\dots}\right)}{\dots} \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="fricas")
```

```
output [-1/4*(sqrt(-a*c)*(cos(f*x + e)^2 - 1)*log(-4*(2*sqrt(-a*c)*sqrt((a*cos(f*
x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x
+ e)^2 + (a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e))/((cos(f*x + e)^2 - 1)*si
n(f*x + e)))*sin(f*x + e) - 2*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt
((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2)/((a^2*c^2*f*cos(f*x +
e)^2 - a^2*c^2*f)*sin(f*x + e)), 1/2*(sqrt(a*c)*(cos(f*x + e)^2 - 1)*arcta
n(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) -
c)/cos(f*x + e))/(a*c*sin(f*x + e)))*sin(f*x + e) + sqrt((a*cos(f*x + e)
+ a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2)
/((a^2*c^2*f*cos(f*x + e)^2 - a^2*c^2*f)*sin(f*x + e))]
```

$$3.143. \int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{3/2}} dx$$

3.143.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\sec(e+fx)}{(a(\sec(e+fx)+1))^{\frac{3}{2}}(-c(\sec(e+fx)-1))^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(3/2),x)`

output `Integral(sec(e + f*x)/((a*(sec(e + f*x) + 1))**(3/2)*(-c*(sec(e + f*x) - 1))**(3/2)), x)`

3.143.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 567 vs. $2(92) = 184$.

Time = 0.41 (sec) , antiderivative size = 567, normalized size of antiderivative = 5.45

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{3/2}} dx = \frac{((2(2\cos(2fx+2e)-1)\cos(4fx+4e)-\cos(4fx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="maxima")`

output `1/2*((2*(2*cos(2*f*x + 2*e) - 1)*cos(4*f*x + 4*e) - cos(4*f*x + 4*e)^2 - 4
*cos(2*f*x + 2*e)^2 - sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x +
2*e) - 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) - 1)*arctan2(sin(f*x + e)
, cos(f*x + e) + 1) - (2*(2*cos(2*f*x + 2*e) - 1)*cos(4*f*x + 4*e) - cos(4
*f*x + 4*e)^2 - 4*cos(2*f*x + 2*e)^2 - sin(4*f*x + 4*e)^2 + 4*sin(4*f*x +
4*e)*sin(2*f*x + 2*e) - 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) - 1)*arc
tan2(sin(f*x + e), cos(f*x + e) - 1) - 2*(sin(3*f*x + 3*e) + sin(f*x + e))
*cos(4*f*x + 4*e) + 2*(cos(3*f*x + 3*e) + cos(f*x + e))*sin(4*f*x + 4*e) +
2*(2*cos(2*f*x + 2*e) - 1)*sin(3*f*x + 3*e) - 4*cos(3*f*x + 3*e)*sin(2*f*
x + 2*e) - 4*cos(f*x + e)*sin(2*f*x + 2*e) + 4*cos(2*f*x + 2*e)*sin(f*x +
e) - 2*sin(f*x + e))*sqrt(a)*sqrt(c)/((a^2*c^2*cos(4*f*x + 4*e)^2 + 4*a^2*
c^2*cos(2*f*x + 2*e)^2 + a^2*c^2*sin(4*f*x + 4*e)^2 - 4*a^2*c^2*sin(4*f*x
+ 4*e)*sin(2*f*x + 2*e) + 4*a^2*c^2*sin(2*f*x + 2*e)^2 - 4*a^2*c^2*cos(2*f
*x + 2*e) + a^2*c^2 - 2*(2*a^2*c^2*cos(2*f*x + 2*e) - a^2*c^2)*cos(4*f*x +
4*e))*f)`

3.143.8 Giac [A] (verification not implemented)

Time = 2.09 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.15

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} (c - c \sec(e + fx))^{3/2}} dx = \frac{\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}{c} + \frac{2c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c}{c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2} - 2 \log \left(|c| \tan \left(\frac{1}{2} \right)}{8 \sqrt{-aca} f |c| \operatorname{sgn} \left(\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + \tan \right)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="giac")`

output `1/8*((c*tan(1/2*f*x + 1/2*e)^2 - c)/c + (2*c*tan(1/2*f*x + 1/2*e)^2 - c)/(
c*tan(1/2*f*x + 1/2*e)^2) - 2*log(abs(c)*tan(1/2*f*x + 1/2*e)^2) + 2*log(a
bs(c)) - 1)/(sqrt(-a*c)*a*f*abs(c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*
x + 1/2*e)))`

3.143.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} (c - c \sec(e + fx))^{3/2}} dx = \int \frac{1}{\cos(e + fx) \left(a + \frac{a}{\cos(e + fx)} \right)^{3/2} \left(c - \frac{c}{\cos(e + fx)} \right)^{3/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(3/2))
,x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(3/2))
, x)`

$$3.144 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{5/2}} dx$$

3.144.1 Optimal result	1019
3.144.2 Mathematica [A] (verified)	1019
3.144.3 Rubi [A] (verified)	1020
3.144.4 Maple [A] (verified)	1022
3.144.5 Fricas [A] (verification not implemented)	1023
3.144.6 Sympy [F(-1)]	1024
3.144.7 Maxima [F(-2)]	1024
3.144.8 Giac [A] (verification not implemented)	1024
3.144.9 Mupad [F(-1)]	1025

3.144.1 Optimal result

Integrand size = 36, antiderivative size = 146

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{5/2}} dx = \frac{3 \csc(e+fx)}{8ac^2 f \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} - \frac{\tan(e+fx)}{4f(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{5/2}} - \frac{3 \arctanh(\cos(e+fx)) \tan(e+fx)}{8ac^2 f \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}}$$

output `3/8*csc(f*x+e)/a/c^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-1/4*tan(f*x+e)/f/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2)-3/8*arctanh(cos(f*x+e))*tan(f*x+e)/a/c^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.144.2 Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.70

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{5/2}} dx = \frac{(2+3 \sec(e+fx)-3 \sec^2(e+fx)+3 \arctanh(\sec(e+fx))(-1+\sec(e+fx))^2(1+\sec(e+fx))) \tan(e+fx)}{8c^2 f (-1+\sec(e+fx))^2 (a(1+\sec(e+fx)))^{3/2} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(5/2)),x]`

3.144. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2}(c-c \sec(e+fx))^{5/2}} dx$

output
$$-1/8*((2 + 3*\text{Sec}[e + f*x] - 3*\text{Sec}[e + f*x]^2 + 3*\text{ArcTanh}[\text{Sec}[e + f*x]]*(-1 + \text{Sec}[e + f*x])^2*(1 + \text{Sec}[e + f*x]))*\text{Tan}[e + f*x])/(c^2*f*(-1 + \text{Sec}[e + f*x])^2*(a*(1 + \text{Sec}[e + f*x]))^{3/2}*\text{Sqrt}[c - c*\text{Sec}[e + f*x]])$$

3.144.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.84, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4448, 3042, 4447, 25, 3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)^{3/2}(c-c\sec(e+fx))^{5/2}} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a\csc(e+fx+\frac{\pi}{2})+a)^{3/2}(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx \\ & \quad \downarrow 4448 \\ & \frac{3 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)^{3/2}(c-c\sec(e+fx))^{3/2}} dx}{4c} - \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{3/2}(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow 3042 \\ & \frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)^{3/2}(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{4c} - \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{3/2}(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow 4447 \\ & \frac{3 \tan(e+fx) \int -\cot^2(e+fx) \csc(e+fx) dx}{4ac^2 \sqrt{a\sec(e+fx)+a} \sqrt{c-c\sec(e+fx)}} - \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{3/2}(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow 25 \\ & \frac{3 \tan(e+fx) \int \cot^2(e+fx) \csc(e+fx) dx}{4ac^2 \sqrt{a\sec(e+fx)+a} \sqrt{c-c\sec(e+fx)}} - \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{3/2}(c-c\sec(e+fx))^{5/2}} \\ & \quad \downarrow 3042 \end{aligned}$$

3.144.
$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{5/2}} dx$$

$$\begin{aligned}
& - \frac{3 \tan(e+fx) \int \sec(e+fx - \frac{\pi}{2}) \tan(e+fx - \frac{\pi}{2})^2 dx}{4ac^2 \sqrt{a \sec(e+fx) + a\sqrt{c - c \sec(e+fx)}} \tan(e+fx)} \\
& \frac{4f(a \sec(e+fx) + a)^{3/2} (c - c \sec(e+fx))^{5/2}}{\downarrow \text{3091}} \\
& - \frac{3 \tan(e+fx) \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right)}{4ac^2 \sqrt{a \sec(e+fx) + a\sqrt{c - c \sec(e+fx)}} \tan(e+fx)} \\
& \frac{4f(a \sec(e+fx) + a)^{3/2} (c - c \sec(e+fx))^{5/2}}{\downarrow \text{3042}} \\
& - \frac{3 \tan(e+fx) \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right)}{4ac^2 \sqrt{a \sec(e+fx) + a\sqrt{c - c \sec(e+fx)}} \tan(e+fx)} \\
& \frac{4f(a \sec(e+fx) + a)^{3/2} (c - c \sec(e+fx))^{5/2}}{\downarrow \text{4257}} \\
& - \frac{3 \tan(e+fx) \left(\frac{\operatorname{arctanh}(\cos(e+fx))}{2f} - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right)}{4ac^2 \sqrt{a \sec(e+fx) + a\sqrt{c - c \sec(e+fx)}} \tan(e+fx)} \\
& \frac{4f(a \sec(e+fx) + a)^{3/2} (c - c \sec(e+fx))^{5/2}}{}
\end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(5/2)),x]`

output `-1/4*Tan[e + f*x]/(f*(a + a*Sec[e + f*x])^(3/2)*(c - c*Sec[e + f*x])^(5/2)) - (3*(ArcTanh[Cos[e + f*x]]/(2*f) - (Cot[e + f*x]*Csc[e + f*x])/(2*f))*Tan[e + f*x])/(4*a*c^2*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.144.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

3.144. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} (c-c \sec(e+fx))^{5/2}} dx$

rule 3091 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4447 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[((-a)*c)^(m + 1/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Int[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]`

rule 4448 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))`

3.144.4 Maple [A] (verified)

Time = 3.62 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.48

method	result
default	$\frac{\sqrt{2} \sqrt{-\frac{2a}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}} (1-\cos(fx+e)) (-2(1-\cos(fx+e))^6 \csc(fx+e)^6 + 12 \ln(-\cot(fx+e) + \csc(fx+e)) (1-\cos(fx+e)))}{64 f a^2 ((1-\cos(fx+e))^2 \csc(fx+e)^2-1)^2 \left(\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}\right)}$
risch	$\frac{i(5 e^{5i(fx+e)} - 2 e^{4i(fx+e)} + 2 e^{3i(fx+e)} - 2 e^{2i(fx+e)} + 5 e^{i(fx+e)})}{4 a c^2 (1+e^{2i(fx+e)}) (e^{i(fx+e)}+1) \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)^3 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f} - \frac{3i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1) \ln(e^{i(fx+e)})}{8 a c^2 (1+e^{2i(fx+e)}) \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}}$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2), x, method=_RET URNVERBOSE)`

$$3.144. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} (c-c \sec(e+fx))^{5/2}} dx$$

output $1/64/f*2^{(1/2)}/a^2*(-2*a/((1-\cos(f*x+e))^2*\csc(f*x+e)^2-1))^{(1/2)}/((1-\cos(f*x+e))^2*\csc(f*x+e)^2-1)^2/(c*(1-\cos(f*x+e))^2/((1-\cos(f*x+e))^2*\csc(f*x+e)^2-1)*\csc(f*x+e)^2)^{(5/2)}*(1-\cos(f*x+e))*(-2*(1-\cos(f*x+e))^6*\csc(f*x+e)^6+12*\ln(-\cot(f*x+e)+\csc(f*x+e))*(1-\cos(f*x+e))^4*\csc(f*x+e)^4+6*(1-\cos(f*x+e))^2*\csc(f*x+e)^2-1)*\csc(f*x+e)$

3.144.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 544, normalized size of antiderivative = 3.73

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{5/2}} dx = \left[\frac{3(\cos(fx+e)^3 - \cos(fx+e)^2 - \cos(fx+e) + 1)}{\dots} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="fracas")`

output `[-1/16*(3*(cos(f*x + e)^3 - cos(f*x + e)^2 - cos(f*x + e) + 1)*sqrt(-a*c)*log(-4*(2*sqrt(-a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2 + (a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e))/((cos(f*x + e)^2 - 1)*sin(f*x + e)))*sin(f*x + e) - 2*(5*cos(f*x + e)^3 - cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c^3*f*cos(f*x + e)^3 - a^2*c^3*f*cos(f*x + e)^2 - a^2*c^3*f*cos(f*x + e) + a^2*c^3*f)*sin(f*x + e)), 1/8*(3*(cos(f*x + e)^3 - cos(f*x + e)^2 - cos(f*x + e) + 1)*sqrt(a*c)*arctan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/(a*c*sin(f*x + e)))*sin(f*x + e) + (5*cos(f*x + e)^3 - cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^2*c^3*f*cos(f*x + e)^3 - a^2*c^3*f*cos(f*x + e)^2 - a^2*c^3*f*cos(f*x + e) + a^2*c^3*f)*sin(f*x + e))]`

3.144.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} (c - c \sec(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))**(5/2),x)`

output `Timed out`

3.144.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} (c - c \sec(e + fx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2),x, algo
rithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.`

3.144.8 Giac [A] (verification not implemented)

Time = 1.92 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.05

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{3/2} (c - c \sec(e + fx))^{5/2}} dx = \frac{2 \left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c \right)}{c} + \frac{9 \left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c \right)^2 + 12 \left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 \right)}{c^2 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4} \cdot \frac{1}{32 \sqrt{-acac} f |c| \operatorname{sgn} \left(\tan\left(\frac{1}{2} \right. \right)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(5/2),x, algo
rithm="giac")`

output `1/32*(2*(c*tan(1/2*f*x + 1/2*e)^2 - c)/c + (9*(c*tan(1/2*f*x + 1/2*e)^2 -
c)^2 + 12*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c + 4*c^2)/(c^2*tan(1/2*f*x + 1/2
*e)^4) - 6*log(abs(c)*tan(1/2*f*x + 1/2*e)^2) + 6*log(abs(c)) - 4)/(sqrt(-
a*c)*a*c*f*abs(c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e)))`

3.144. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{3/2} (c-c \sec(e+fx))^{5/2}} dx$

3.144.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{3/2}(c-c\sec(e+fx))^{5/2}} dx = \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^{3/2} \left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(5/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(3/2)*(c - c/cos(e + f*x))^(5/2)), x)`

3.145
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx$$

3.145.1 Optimal result	1026
3.145.2 Mathematica [A] (verified)	1026
3.145.3 Rubi [A] (verified)	1027
3.145.4 Maple [A] (verified)	1029
3.145.5 Fricas [F]	1029
3.145.6 Sympy [F(-1)]	1030
3.145.7 Maxima [A] (verification not implemented)	1030
3.145.8 Giac [A] (verification not implemented)	1031
3.145.9 Mupad [F(-1)]	1031

3.145.1 Optimal result

Integrand size = 36, antiderivative size = 145

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx = -\frac{c^3 \log(1+\sec(e+fx)) \tan(e+fx)}{a^2 f \sqrt{a+a\sec(e+fx)} \sqrt{c-c\sec(e+fx)}} - \frac{c^2 \sqrt{c-c\sec(e+fx)} \tan(e+fx)}{af(a+a\sec(e+fx))^{3/2}} + \frac{c(c-c\sec(e+fx))^{3/2} \tan(e+fx)}{2f(a+a\sec(e+fx))^{5/2}}$$

output `1/2*c*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(5/2)-c^3*ln(1+sec(f*x+e))*tan(f*x+e)/a^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-c^2*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^(3/2)`

3.145.2 Mathematica [A] (verified)

Time = 0.72 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.61

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx = \frac{c\left(c^2 \log(1+\sec(e+fx)) - \frac{2c^2}{(1+\sec(e+fx))^2} + \frac{4c^2}{1+\sec(e+fx)}\right) \tan(e+fx)}{a^2 f \sqrt{a(1+\sec(e+fx))} \sqrt{c-c\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^(5/2), x]`

output $-\left((c*(c^2*\text{Log}[1 + \text{Sec}[e + f*x]] - (2*c^2)/(1 + \text{Sec}[e + f*x])^2 + (4*c^2)/(1 + \text{Sec}[e + f*x]))*\text{Tan}[e + f*x])/(a^2*f*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]*\text{Sqrt}[c - c*\text{Sec}[e + f*x]])\right)$

3.145.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4442, 3042, 4442, 3042, 4440}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a\sec(e+fx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^{5/2}} dx$$

↓ 4442

$$\frac{c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{2f(a\sec(e+fx)+a)^{5/2}} - \frac{c \int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(\sec(e+fx)a+a)^{3/2}} dx}{a}$$

↓ 3042

$$\frac{c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{2f(a\sec(e+fx)+a)^{5/2}} - \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{(\csc(e+fx+\frac{\pi}{2})a+a)^{3/2}} dx}{a}$$

↓ 4442

$$\frac{c \tan(e+fx)(c-c\sec(e+fx))^{3/2}}{2f(a\sec(e+fx)+a)^{5/2}} - \frac{c \left(\frac{c \tan(e+fx)\sqrt{c-c\sec(e+fx)}}{f(a\sec(e+fx)+a)^{3/2}} - \frac{c \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} dx}{a} \right)}{a}$$

↓ 3042

3.145. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx$

$$\frac{c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{2f(a \sec(e+fx) + a)^{5/2}} - c \left(\frac{c \tan(e+fx) \sqrt{c - c \sec(e+fx)}}{f(a \sec(e+fx) + a)^{3/2}} - \frac{c \int \frac{\csc(e+fx + \frac{\pi}{2}) \sqrt{c - c \csc(e+fx + \frac{\pi}{2})}}{\sqrt{\csc(e+fx + \frac{\pi}{2}) a + a}} dx}{a} \right)$$

a
↓ 4440

$$\frac{c \tan(e+fx)(c - c \sec(e+fx))^{3/2}}{2f(a \sec(e+fx) + a)^{5/2}} - \frac{c \left(\frac{c^2 \tan(e+fx) \log(\sec(e+fx) + 1)}{af \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} + \frac{c \tan(e+fx) \sqrt{c - c \sec(e+fx)}}{f(a \sec(e+fx) + a)^{3/2}} \right)}{a}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(5/2))/(a + a*Sec[e + f*x])^(5/2),x]`

output `(c*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(2*f*(a + a*Sec[e + f*x])^(5/2)) - (c*((c^2*Log[1 + Sec[e + f*x]]*Tan[e + f*x])/(a*f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]) + (c*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])^(3/2))))/a`

3.145.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4440 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[a*c*Log[1 + (b/a)*Csc[e + f*x]]*(Cot[e + f*x]/(b*f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

```
rule 4442 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[2*a*c*Cot[e +
f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(b*f*(2*m + 1))),
x] - Simp[d*((2*n - 1)/(b*(2*m + 1))) Int[Csc[e + f*x]*(a + b*Csc[e + f*
x])^(m + 1)*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && LtQ[
m, -2^(-1)]
```

3.145.4 Maple [A] (verified)

Time = 3.54 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.50

method	result
default	$\frac{\sqrt{2} \sqrt{-\frac{2a}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}} \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^3 \left(\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \right)^{\frac{5}{2}} \sin(fx+e)^5 \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right)^{\frac{5}{2}}}{4f a^3 (1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}$
risch	$-\frac{8ic^2 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} e^{2i(fx+e)}}{a^2 (e^{i(fx+e)}+1)^3 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f} + \frac{2ic^2 (e^{i(fx+e)}+1) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} \ln(e^{i(fx+e)}+1)}{a^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f} - \frac{ic^2 (e^{i(fx+e)}+1)}{a^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)f}$

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 1/4/f*2^(1/2)/a^3*(-2*a/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)*((1-cos(f
*x+e))^2*csc(f*x+e)^2-1)^3*(c*(1-cos(f*x+e))^2/((1-cos(f*x+e))^2*csc(f*x+e
)^2-1)*csc(f*x+e)^2)^(5/2)/(1-cos(f*x+e))^5*sin(f*x+e)^5*((1-cos(f*x+e))^4
*csc(f*x+e)^4+2*(1-cos(f*x+e))^2*csc(f*x+e)^2+2*ln(-cot(f*x+e)+csc(f*x+e)-
1)+2*ln(-cot(f*x+e)+csc(f*x+e)+1))
```

3.145.5 Fracas [F]

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx = \int \frac{(-c\sec(fx+e)+c)^{5/2} \sec(fx+e)}{(a\sec(fx+e)+a)^{5/2}} dx$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(5/2),x, algo
rithm="fracas")
```

3.145. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx$

output `integral((c^2*sec(f*x + e)^3 - 2*c^2*sec(f*x + e)^2 + c^2*sec(f*x + e))*sqrt(a*sec(f*x + e) + a)*sqrt(-c*sec(f*x + e) + c)/(a^3*sec(f*x + e)^3 + 3*a^3*sec(f*x + e)^2 + 3*a^3*sec(f*x + e) + a^3), x)`

3.145.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{(a + a \sec(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(5/2)/(a+a*sec(f*x+e))**(5/2),x)`

output `Timed out`

3.145.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.92

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{5/2}}{(a + a \sec(e + fx))^{5/2}} dx =$$

$$\frac{2c^{5/2} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{\sqrt{-aa^2}} + \frac{2c^{5/2} \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{\sqrt{-aa^2}} - \frac{\frac{2\sqrt{-ac}^{5/2} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{\sqrt{-ac}^{5/2} \sin(fx+e)^4}{(\cos(fx+e)+1)^4}}{a^3}$$

$$\frac{\hspace{10em}}{2f}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(5/2),x, algorithm="maxima")`

output `-1/2*(2*c^(5/2)*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/(sqrt(-a)*a^2) + 2*c^(5/2)*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/(sqrt(-a)*a^2) - (2*sqrt(-a)*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + sqrt(-a)*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4)/a^3)/f`

3.145.8 Giac [A] (verification not implemented)

Time = 1.35 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.79

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx =$$

$$\frac{c^4 \left(\frac{(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)^2 c^2 + 4(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c)c^3}{c^4} + 2 \log(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c) \right) \operatorname{sgn}(\tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + \tan(\frac{1}{2}fx + \frac{1}{2}e))}{2\sqrt{-aca^2f|c|}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(5/2)/(a+a*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `-1/2*c^4*((c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^2 + 4*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^3)/c^4 + 2*log(c*tan(1/2*f*x + 1/2*e)^2 - c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e))/(sqrt(-a*c)*a^2*f*abs(c))`

3.145.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{5/2}}{(a+a\sec(e+fx))^{5/2}} dx = \int \frac{\left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)),x)`

output `int((c - c/cos(e + f*x))^(5/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)),x)`

3.146
$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{5/2}} dx$$

3.146.1 Optimal result 1032
 3.146.2 Mathematica [A] (verified) 1032
 3.146.3 Rubi [A] (verified) 1033
 3.146.4 Maple [A] (verified) 1034
 3.146.5 Fricas [B] (verification not implemented) 1034
 3.146.6 Sympy [F] 1035
 3.146.7 Maxima [B] (verification not implemented) 1035
 3.146.8 Giac [A] (verification not implemented) 1035
 3.146.9 Mupad [B] (verification not implemented) 1036

3.146.1 Optimal result

Integrand size = 36, antiderivative size = 42

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{5/2}} dx = \frac{(c-c\sec(e+fx))^{3/2} \tan(e+fx)}{4f(a+a\sec(e+fx))^{5/2}}$$

output `1/4*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(a+a*sec(f*x+e))^(5/2)`

3.146.2 Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.21

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{5/2}} dx = -\frac{c(-1+\sec(e+fx))\sqrt{c-c\sec(e+fx)}\tan(e+fx)}{4f(a(1+\sec(e+fx)))^{5/2}}$$

input `Integrate[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x])^(5/2),x]`

output `-1/4*(c*(-1 + Sec[e + f*x])*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*(a*(1 + Sec[e + f*x]))^(5/2))`

3.146.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a\sec(e+fx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}}{(a\csc(e+fx+\frac{\pi}{2})+a)^{5/2}} dx$$

↓ 4438

$$\frac{\tan(e+fx)(c-c\sec(e+fx))^{3/2}}{4f(a\sec(e+fx)+a)^{5/2}}$$

input `Int[(Sec[e + f*x]*(c - c*Sec[e + f*x])^(3/2))/(a + a*Sec[e + f*x])^(5/2),x]`

output `((c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(4*f*(a + a*Sec[e + f*x])^(5/2))`

3.146.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.146.4 Maple [A] (verified)

Time = 2.38 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.60

method	result	size
default	$-\frac{\sin(fx+e)(\sec(fx+e)-1)\sqrt{-c(\sec(fx+e)-1)}\sqrt{a(\sec(fx+e)+1)}c\cos(fx+e)^2}{4fa^3(\cos(fx+e)+1)^3}$	67
risch	$\frac{2ic\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}(e^{3i(fx+e)}+e^{i(fx+e)})}{a^2(e^{i(fx+e)}+1)^3\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)}f$	116

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(5/2),x,method=_RET
URNVERBOSE)
```

```
output -1/4/f/a^3*sin(f*x+e)*(sec(f*x+e)-1)*(-c*(sec(f*x+e)-1))^(1/2)*(a*(sec(f*x
+e)+1))^(1/2)*c*cos(f*x+e)^2/(cos(f*x+e)+1)^3
```

3.146.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(36) = 72.

Time = 0.26 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.26

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{5/2}} dx = \frac{c\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}\cos(fx+e)^2}{(a^3f\cos(fx+e))^2+2a^3f\cos(fx+e)+a^3f}\sin(fx+e)$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(5/2),x, algo
rithm="fracas")
```

```
output c*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*
x + e))*cos(f*x + e)^2/((a^3*f*cos(f*x + e))^2 + 2*a^3*f*cos(f*x + e) + a^3
*f)*sin(f*x + e)
```

3.146.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{(a + a \sec(e + fx))^{5/2}} dx = \int \frac{(-c(\sec(e + fx) - 1))^{\frac{3}{2}} \sec(e + fx)}{(a(\sec(e + fx) + 1))^{\frac{5}{2}}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(3/2)/(a+a*sec(f*x+e))**(5/2),x)`

output `Integral((-c*(sec(e + f*x) - 1))**(3/2)*sec(e + f*x)/(a*(sec(e + f*x) + 1))**(5/2), x)`

3.146.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 98 vs. $2(36) = 72$.

Time = 0.32 (sec) , antiderivative size = 98, normalized size of antiderivative = 2.33

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{(a + a \sec(e + fx))^{5/2}} dx = \frac{\sqrt{-ac^3} \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right) \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right) \sin(fx+e)^4}{4 \left(a^3 - \frac{a^3 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} \right) f(\cos(fx+e) + 1)^4}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(5/2),x, algorithm="maxima")`

output `-1/4*sqrt(-a)*c^(3/2)*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)*sin(f*x + e)^4/((a^3 - a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*f*(cos(f*x + e) + 1)^4)`

3.146.8 Giac [A] (verification not implemented)

Time = 1.54 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.40

$$\int \frac{\sec(e + fx)(c - c \sec(e + fx))^{3/2}}{(a + a \sec(e + fx))^{5/2}} dx = \frac{\left(\left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c \right)^2 + 2 \left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c \right) c \right)}{4 \sqrt{-aca^2 f} |c|}$$

3.146. $\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{5/2}} dx$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(3/2)/(a+a*sec(f*x+e))^(5/2),x, algo
rithm="giac")`

output `-1/4*((c*tan(1/2*f*x + 1/2*e)^2 - c)^2 + 2*(c*tan(1/2*f*x + 1/2*e)^2 - c)*
c)/(sqrt(-a*c)*a^2*f*abs(c))`

3.146.9 Mupad [B] (verification not implemented)

Time = 15.26 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.83

$$\int \frac{\sec(e+fx)(c-c\sec(e+fx))^{3/2}}{(a+a\sec(e+fx))^{5/2}} dx =$$

$$\frac{2c \sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}} (\sin(e+fx) + 2\sin(2e+2fx) + \sin(3e+3fx))}{a^2 f \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}} (4\cos(2e+2fx) - 4\cos(e+fx) + 4\cos(3e+3fx) + \cos(4e+4fx) - 5)}$$

input `int((c - c/cos(e + f*x))^(3/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)),x
)`

output `-(2*c*((c*(cos(e + f*x) - 1))/cos(e + f*x))^(1/2)*(sin(e + f*x) + 2*sin(2*
e + 2*f*x) + sin(3*e + 3*f*x)))/(a^2*f*((a*(cos(e + f*x) + 1))/cos(e + f*x
)^(1/2)*(4*cos(2*e + 2*f*x) - 4*cos(e + f*x) + 4*cos(3*e + 3*f*x) + cos(4
*e + 4*f*x) - 5))`

$$3.147 \quad \int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx$$

3.147.1 Optimal result	1037
3.147.2 Mathematica [A] (verified)	1037
3.147.3 Rubi [A] (verified)	1038
3.147.4 Maple [A] (verified)	1039
3.147.5 Fricas [B] (verification not implemented)	1039
3.147.6 Sympy [F]	1040
3.147.7 Maxima [A] (verification not implemented)	1040
3.147.8 Giac [A] (verification not implemented)	1040
3.147.9 Mupad [B] (verification not implemented)	1041

3.147.1 Optimal result

Integrand size = 36, antiderivative size = 43

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx = \frac{c \tan(e+fx)}{2f(a+a\sec(e+fx))^{5/2}\sqrt{c-c\sec(e+fx)}}$$

output `1/2*c*tan(f*x+e)/f/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2)`

3.147.2 Mathematica [A] (verified)

Time = 1.17 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.65

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx = \frac{(1+2\cos(e+fx))\csc(\frac{1}{2}(e+fx))\sec^3(\frac{1}{2}(e+fx))\sqrt{c-c\sec(e+fx)}}{8a^2f\sqrt{a(1+\sec(e+fx))}}$$

input `Integrate[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^(5/2),x]`

output `((1 + 2*Cos[e + f*x])*Csc[(e + f*x)/2]*Sec[(e + f*x)/2]^3*Sqrt[c - c*Sec[e + f*x]])/(8*a^2*f*Sqrt[a*(1 + Sec[e + f*x])])`

3.147.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a\sec(e+fx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}}{(a\csc(e+fx+\frac{\pi}{2})+a)^{5/2}} dx$$

↓ 4441

$$\frac{c\tan(e+fx)}{2f(a\sec(e+fx)+a)^{5/2}\sqrt{c-c\sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[c - c*Sec[e + f*x]])/(a + a*Sec[e + f*x])^(5/2),x]`

output `(c*Tan[e + f*x])/(2*f*(a + a*Sec[e + f*x])^(5/2)*Sqrt[c - c*Sec[e + f*x]])`

3.147.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

3.147.4 Maple [A] (verified)

Time = 3.51 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.35

method	result	size
default	$-\frac{\sqrt{-c(\sec(fx+e)-1)}\sqrt{a(\sec(fx+e)+1)}\cos(fx+e)^2\cot(fx+e)}{2fa^3(\cos(fx+e)+1)^2}$	58
risch	$\frac{2i\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}(e^{3i(fx+e)}+e^{2i(fx+e)}+e^{i(fx+e)})}{a^2(e^{i(fx+e)}+1)^3\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}(e^{i(fx+e)}-1)f}$	124

```
input int(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(5/2),x,method=_RET
URNVERBOSE)
```

```
output -1/2/f/a^3*(-c*(sec(f*x+e)-1))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)/(cos(f*x+e)+
1)^2*cos(f*x+e)^2*cot(f*x+e)
```

3.147.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs. 2(37) = 74.

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 2.42

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx = \frac{(2\cos(fx+e)^2 + \cos(fx+e))\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)-c}{\cos(fx+e)}}}{2(a^3f\cos(fx+e)^2 + 2a^3f\cos(fx+e) + a^3f)\sin(fx+e)}$$

```
input integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(5/2),x, algo
rithm="fricas")
```

```
output 1/2*(2*cos(f*x + e)^2 + cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x +
e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((a^3*f*cos(f*x + e)^2 + 2*a^3
*f*cos(f*x + e) + a^3*f)*sin(f*x + e))
```


3.147.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx = \int \frac{\sqrt{-c(\sec(e+fx)-1)}\sec(e+fx)}{(a(\sec(e+fx)+1))^{5/2}} dx$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))**(1/2)/(a+a*sec(f*x+e))**(5/2),x)`

output `Integral(sqrt(-c*(sec(e + f*x) - 1))*sec(e + f*x)/(a*(sec(e + f*x) + 1))**(5/2), x)`

3.147.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.35

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx = -\frac{\sqrt{c}\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)^2\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)^2}{8\sqrt{-aa^2f}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(5/2),x, algorithm="maxima")`

output `-1/8*sqrt(c)*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^2*(sin(f*x + e)/(cos(f*x + e) + 1) - 1)^2/(sqrt(-a)*a^2*f)`

3.147.8 Giac [A] (verification not implemented)

Time = 1.49 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.86

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx = -\frac{\left(c\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - c\right)^2}{8\sqrt{-aca^2f|c|}}$$

input `integrate(sec(f*x+e)*(c-c*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `-1/8*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2/(sqrt(-a*c)*a^2*f*abs(c))`

3.147. $\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx$

3.147.9 Mupad [B] (verification not implemented)

Time = 14.61 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.79

$$\int \frac{\sec(e+fx)\sqrt{c-c\sec(e+fx)}}{(a+a\sec(e+fx))^{5/2}} dx =$$

$$\frac{2(3\sin(e+fx) + 3\sin(2e+2fx) + \sin(3e+3fx))\sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}}}{a^2 f \sqrt{\frac{a(\cos(e+fx)+1)}{\cos(e+fx)}} (4\cos(2e+2fx) - 4\cos(e+fx) + 4\cos(3e+3fx) + \cos(4e+4fx) - 5)}$$

input `int((c - c/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)),x)`

output `-(2*(3*sin(e + f*x) + 3*sin(2*e + 2*f*x) + sin(3*e + 3*f*x))*((c*(cos(e + f*x) - 1))/cos(e + f*x))^(1/2))/(a^2*f*((a*(cos(e + f*x) + 1))/cos(e + f*x))^(1/2)*(4*cos(2*e + 2*f*x) - 4*cos(e + f*x) + 4*cos(3*e + 3*f*x) + cos(4*e + 4*f*x) - 5))`

$$3.148 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)}} dx$$

3.148.1 Optimal result	1042
3.148.2 Mathematica [A] (verified)	1042
3.148.3 Rubi [A] (verified)	1043
3.148.4 Maple [A] (verified)	1045
3.148.5 Fricas [A] (verification not implemented)	1046
3.148.6 Sympy [F]	1046
3.148.7 Maxima [B] (verification not implemented)	1047
3.148.8 Giac [A] (verification not implemented)	1048
3.148.9 Mupad [F(-1)]	1048

3.148.1 Optimal result

Integrand size = 36, antiderivative size = 140

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)}} dx = \frac{\tan(e+fx)}{4f(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)}} + \frac{\tan(e+fx)}{4af(a+a \sec(e+fx))^{3/2} \sqrt{c-c \sec(e+fx)}} - \frac{\operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{4a^2 f \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}}$$

```
output 1/4*tan(f*x+e)/f/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2)+1/4*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))^(1/2)-1/4*arctanh(cos(f*x+e))*tan(f*x+e)/a^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)
```

3.148.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.50

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)}} dx = \frac{(-2 - \sec(e+fx) + \operatorname{arctanh}(\sec(e+fx))(1 + \sec(e+fx))^2) \tan(e+fx)}{4f(a(1 + \sec(e+fx)))^{5/2} \sqrt{c-c \sec(e+fx)}}$$

```
input Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^(5/2)*Sqrt[c - c*Sec[e + f*x]]),x]
```

3.148. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)}} dx$

output
$$-1/4*((-2 - \text{Sec}[e + f*x] + \text{ArcTanh}[\text{Sec}[e + f*x]])*(1 + \text{Sec}[e + f*x])^2*\text{Tan}[e + f*x])/(f*(a*(1 + \text{Sec}[e + f*x]))^{5/2}*\text{Sqrt}[c - c*\text{Sec}[e + f*x]])$$

3.148.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4448, 3042, 4448, 3042, 4447, 25, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^{5/2} \sqrt{c - c \sec(e+fx)}} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^{5/2} \sqrt{c - c \csc(e+fx + \frac{\pi}{2})}} dx \\ & \quad \downarrow 4448 \\ & \frac{\int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)^{3/2} \sqrt{c - c \sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2} \sqrt{c - c \sec(e+fx)}} \\ & \quad \downarrow 3042 \\ & \frac{\int \frac{\csc(e+fx + \frac{\pi}{2})}{(\csc(e+fx + \frac{\pi}{2})a+a)^{3/2} \sqrt{c - c \csc(e+fx + \frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2} \sqrt{c - c \sec(e+fx)}} \\ & \quad \downarrow 4448 \\ & \frac{\int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a+a} \sqrt{c - c \sec(e+fx)}} dx}{2a} + \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} + \\ & \quad \frac{2a \tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2} \sqrt{c - c \sec(e+fx)}} \\ & \quad \downarrow 3042 \\ & \frac{\int \frac{\csc(e+fx + \frac{\pi}{2})}{\sqrt{\csc(e+fx + \frac{\pi}{2})a+a} \sqrt{c - c \csc(e+fx + \frac{\pi}{2})}} dx}{2a} + \frac{\tan(e+fx)}{2f(a \sec(e+fx) + a)^{3/2} \sqrt{c - c \sec(e+fx)}} + \\ & \quad \frac{2a \tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2} \sqrt{c - c \sec(e+fx)}} \end{aligned}$$

3.148.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} \sqrt{c - c \sec(e+fx)}} dx$$

$$\begin{aligned}
& \downarrow 4447 \\
& \frac{\frac{\tan(e+fx)}{2f(a \sec(e+fx)+a)^{3/2}\sqrt{c-c \sec(e+fx)}} - \frac{\tan(e+fx) \int -c \sec(e+fx) dx}{2a\sqrt{a \sec(e+fx)+a}\sqrt{c-c \sec(e+fx)}}}{\frac{\tan^2(e+fx)}{4f(a \sec(e+fx)+a)^{5/2}\sqrt{c-c \sec(e+fx)}}} + \\
& \downarrow 25 \\
& \frac{\frac{\tan(e+fx) \int c \sec(e+fx) dx}{2a\sqrt{a \sec(e+fx)+a}\sqrt{c-c \sec(e+fx)}} + \frac{\tan(e+fx)}{2f(a \sec(e+fx)+a)^{3/2}\sqrt{c-c \sec(e+fx)}}}{\frac{\tan^2(e+fx)}{4f(a \sec(e+fx)+a)^{5/2}\sqrt{c-c \sec(e+fx)}}} + \\
& \downarrow 3042 \\
& \frac{\frac{\tan(e+fx) \int c \sec(e+fx) dx}{2a\sqrt{a \sec(e+fx)+a}\sqrt{c-c \sec(e+fx)}} + \frac{\tan(e+fx)}{2f(a \sec(e+fx)+a)^{3/2}\sqrt{c-c \sec(e+fx)}}}{\frac{\tan^2(e+fx)}{4f(a \sec(e+fx)+a)^{5/2}\sqrt{c-c \sec(e+fx)}}} + \\
& \downarrow 4257 \\
& \frac{\frac{\tan(e+fx)}{2f(a \sec(e+fx)+a)^{3/2}\sqrt{c-c \sec(e+fx)}} - \frac{\tan(e+fx) \operatorname{arctanh}(\cos(e+fx))}{2af\sqrt{a \sec(e+fx)+a}\sqrt{c-c \sec(e+fx)}}}{\frac{\tan^2(e+fx)}{4f(a \sec(e+fx)+a)^{5/2}\sqrt{c-c \sec(e+fx)}}} +
\end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^(5/2)*Sqrt[c - c*Sec[e + f*x]]),x]`

output `Tan[e + f*x]/(4*f*(a + a*Sec[e + f*x])^(5/2)*Sqrt[c - c*Sec[e + f*x]]) + (Tan[e + f*x]/(2*f*(a + a*Sec[e + f*x])^(3/2)*Sqrt[c - c*Sec[e + f*x]]) - (ArcTanh[Cos[e + f*x]]*Tan[e + f*x])/(2*a*f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]))/(2*a)`

3.148. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}\sqrt{c-c \sec(e+fx)}} dx$

3.148.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4447 Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(m_), x_Symbol] := Simp[((-a)*c)^(m + 1/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Int[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]
```

```
rule 4448 Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))
```

3.148.4 Maple [A] (verified)

Time = 2.31 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.99

method	result
default	$\frac{\sin(fx+e) \left(4 \cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e))+8 \cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e))+5 \cos(fx+e)^2+4 \ln(-\cot(fx+e)) \right)}{16 f a^3 (\cos(fx+e)+1)^3 \sqrt{-c(\sec(fx+e)-1)}}$
risch	$\frac{i(3e^{2i(fx+e)}+4e^{i(fx+e)}+3)(e^{2i(fx+e)}-e^{i(fx+e)})}{2a^2(1+e^{2i(fx+e)})(e^{i(fx+e)}+1)^3 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f} - \frac{i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1) \ln(e^{i(fx+e)}-1)}{4a^2 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (1+e^{2i(fx+e)}) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x,method=_RET URNVERBOSE)
```

$$3.148. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} \sqrt{c-c \sec(e+fx)}} dx$$

output $1/16/f/a^3*\sin(f*x+e)*(4*\cos(f*x+e)^2*\ln(-\cot(f*x+e)+\csc(f*x+e))+8*\cos(f*x+e)*\ln(-\cot(f*x+e)+\csc(f*x+e))+5*\cos(f*x+e)^2+4*\ln(-\cot(f*x+e)+\csc(f*x+e))-2*\cos(f*x+e)-3)*(a*(\sec(f*x+e)+1))^{1/2}/(\cos(f*x+e)+1)^3/(-c*(\sec(f*x+e)-1))^{1/2}$

3.148.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 456, normalized size of antiderivative = 3.26

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{5/2}\sqrt{c-c\sec(e+fx)}} dx = \left[\frac{\sqrt{-ac}(\cos(fx+e)^2+2\cos(fx+e)+1)\log\left(-\frac{4}{2}\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x, algo rithm="fricas")`

output `[-1/8*(sqrt(-a*c)*(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)*log(-4*(2*sqrt(-a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2 + (a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e))/((cos(f*x + e)^2 - 1)*sin(f*x + e)))*sin(f*x + e) - 2*(3*cos(f*x + e)^2 + 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^3*c*f*cos(f*x + e)^2 + 2*a^3*c*f*cos(f*x + e) + a^3*c*f)*sin(f*x + e)), 1/4*(sqrt(a*c)*(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)*arc tan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(a*c*sin(f*x + e)))*sin(f*x + e) + (3*cos(f*x + e)^2 + 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^3*c*f*cos(f*x + e)^2 + 2*a^3*c*f*cos(f*x + e) + a^3*c*f)*sin(f*x + e))]`

3.148.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{5/2}\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\sec(e+fx)}{(a(\sec(e+fx)+1))^{5/2}\sqrt{-c(\sec(e+fx)-1)}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(1/2),x)`

3.148. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{5/2}\sqrt{c-c\sec(e+fx)}} dx$

output `Integral(sec(e + f*x)/((a*(sec(e + f*x) + 1))**(5/2)*sqrt(-c*(sec(e + f*x) - 1))), x)`

3.148.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1191 vs. $2(122) = 244$.

Time = 0.43 (sec) , antiderivative size = 1191, normalized size of antiderivative = 8.51

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x, algorith="maxima")`

output `-1/4*((2*(4*cos(3*f*x + 3*e) + 6*cos(2*f*x + 2*e) + 4*cos(f*x + e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 8*(6*cos(2*f*x + 2*e) + 4*cos(f*x + e) + 1)*cos(3*f*x + 3*e) + 16*cos(3*f*x + 3*e)^2 + 12*(4*cos(f*x + e) + 1)*cos(2*f*x + 2*e) + 36*cos(2*f*x + 2*e)^2 + 16*cos(f*x + e)^2 + 4*(2*sin(3*f*x + 3*e) + 3*sin(2*f*x + 2*e) + 2*sin(f*x + e))*sin(4*f*x + 4*e) + sin(4*f*x + 4*e)^2 + 16*(3*sin(2*f*x + 2*e) + 2*sin(f*x + e))*sin(3*f*x + 3*e) + 16*sin(3*f*x + 3*e)^2 + 36*sin(2*f*x + 2*e)^2 + 48*sin(2*f*x + 2*e)*sin(f*x + e) + 16*sin(f*x + e)^2 + 8*cos(f*x + e) + 1)*arctan2(sin(f*x + e), cos(f*x + e) + 1) - (2*(4*cos(3*f*x + 3*e) + 6*cos(2*f*x + 2*e) + 4*cos(f*x + e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 8*(6*cos(2*f*x + 2*e) + 4*cos(f*x + e) + 1)*cos(3*f*x + 3*e) + 16*cos(3*f*x + 3*e)^2 + 12*(4*cos(f*x + e) + 1)*cos(2*f*x + 2*e) + 36*cos(2*f*x + 2*e)^2 + 16*cos(f*x + e)^2 + 4*(2*sin(3*f*x + 3*e) + 3*sin(2*f*x + 2*e) + 2*sin(f*x + e))*sin(4*f*x + 4*e) + sin(4*f*x + 4*e)^2 + 16*(3*sin(2*f*x + 2*e) + 2*sin(f*x + e))*sin(3*f*x + 3*e) + 16*sin(3*f*x + 3*e)^2 + 36*sin(2*f*x + 2*e)^2 + 48*sin(2*f*x + 2*e)*sin(f*x + e) + 16*sin(f*x + e)^2 + 8*cos(f*x + e) + 1)*arctan2(sin(f*x + e), cos(f*x + e) - 1) + 2*(3*sin(3*f*x + 3*e) + 4*sin(2*f*x + 2*e) + 3*sin(f*x + e))*cos(4*f*x + 4*e) - 2*(3*cos(3*f*x + 3*e) + 4*cos(2*f*x + 2*e) + 3*cos(f*x + e))*sin(4*f*x + 4*e) + 2*(2*cos(2*f*x + 2*e) + 3)*sin(3*f*x + 3*e) - 4*(cos(f*x + e) - 2)*sin(2*f*x + 2*e) - 4*cos(3*f*x + ...`

3.148.8 Giac [A] (verification not implemented)

Time = 1.76 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.89

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)}} dx =$$

$$\frac{c^2 \left(\frac{2 \log(|c| \tan(\frac{1}{2} fx + \frac{1}{2} e)^2)}{c} - \frac{2 \log(|c|)}{c} + \frac{(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)^2 c^3 - 2 (c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c) c^4}{c^6} \right)}{16 \sqrt{-aca^2 f |c| \operatorname{sgn} \left(\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)}}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `-1/16*c^2*(2*log(abs(c)*tan(1/2*f*x + 1/2*e)^2)/c - 2*log(abs(c))/c + ((c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^3 - 2*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^4)/c^6)/(sqrt(-a*c)*a^2*f*abs(c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e)))`

3.148.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} \sqrt{c - c \sec(e + fx)}} dx = \int \frac{1}{\cos(e + fx) \left(a + \frac{a}{\cos(e + fx)} \right)^{5/2} \sqrt{c - \frac{c}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(1/2)), x)`

$$3.149 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2}} dx$$

3.149.1 Optimal result	1049
3.149.2 Mathematica [A] (verified)	1049
3.149.3 Rubi [A] (verified)	1050
3.149.4 Maple [A] (verified)	1052
3.149.5 Fricas [A] (verification not implemented)	1053
3.149.6 Sympy [F(-1)]	1054
3.149.7 Maxima [F(-2)]	1054
3.149.8 Giac [A] (verification not implemented)	1054
3.149.9 Mupad [F(-1)]	1055

3.149.1 Optimal result

Integrand size = 36, antiderivative size = 146

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2}} dx = \frac{3 \csc(e+fx)}{8a^2cf \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} + \frac{\tan(e+fx)}{4f(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2}} - \frac{3 \operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{8a^2cf \sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}}$$

output `3/8*csc(f*x+e)/a^2/c/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)+1/4*tan(f*x+e)/f/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2)-3/8*arctanh(cos(f*x+e))*tan(f*x+e)/a^2/c/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.149.2 Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.70

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2}} dx = \frac{(2-3 \sec(e+fx)-3 \sec^2(e+fx)+3 \operatorname{arctanh}(\sec(e+fx))(-1+\sec(e+fx))(1+\sec(e+fx))^2) \tan(e+fx)}{8cf(-1+\sec(e+fx))(a(1+\sec(e+fx)))^{5/2} \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(3/2)),x]`

3.149. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{3/2}} dx$

output
$$-1/8*((2 - 3*\text{Sec}[e + f*x] - 3*\text{Sec}[e + f*x]^2 + 3*\text{ArcTanh}[\text{Sec}[e + f*x]]*(-1 + \text{Sec}[e + f*x])*(1 + \text{Sec}[e + f*x])^2)*\text{Tan}[e + f*x])/(c*f*(-1 + \text{Sec}[e + f*x]))*(a*(1 + \text{Sec}[e + f*x]))^{(5/2)}*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]$$

3.149.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.84, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4448, 3042, 4447, 25, 3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{3/2}} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})}{(a\csc(e+fx+\frac{\pi}{2})+a)^{5/2}(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx \\ & \quad \downarrow 4448 \\ & \frac{3 \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)^{3/2}(c-c\sec(e+fx))^{3/2}} dx}{4a} + \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{3/2}} \\ & \quad \downarrow 3042 \\ & \frac{3 \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)^{3/2}(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx}{4a} + \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{3/2}} \\ & \quad \downarrow 4447 \\ & \frac{3 \tan(e+fx) \int -\cot^2(e+fx) \csc(e+fx) dx}{4a^2 c \sqrt{a\sec(e+fx)+a} \sqrt{c-c\sec(e+fx)}} + \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{3/2}} \\ & \quad \downarrow 25 \\ & \frac{\tan(e+fx)}{4f(a\sec(e+fx)+a)^{5/2}(c-c\sec(e+fx))^{3/2}} - \frac{3 \tan(e+fx) \int \cot^2(e+fx) \csc(e+fx) dx}{4a^2 c \sqrt{a\sec(e+fx)+a} \sqrt{c-c\sec(e+fx)}} \\ & \quad \downarrow 3042 \end{aligned}$$

3.149.
$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{3/2}} dx$$

$$\frac{\tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2}(c - c \sec(e+fx))^{3/2}} - \frac{3 \tan(e+fx) \int \sec(e+fx - \frac{\pi}{2}) \tan(e+fx - \frac{\pi}{2})^2 dx}{4a^2c\sqrt{a \sec(e+fx) + a}\sqrt{c - c \sec(e+fx)}}$$

↓ 3091

$$\frac{\tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2}(c - c \sec(e+fx))^{3/2}} - \frac{3 \tan(e+fx) \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right)}{4a^2c\sqrt{a \sec(e+fx) + a}\sqrt{c - c \sec(e+fx)}}$$

↓ 3042

$$\frac{\tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2}(c - c \sec(e+fx))^{3/2}} - \frac{3 \tan(e+fx) \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right)}{4a^2c\sqrt{a \sec(e+fx) + a}\sqrt{c - c \sec(e+fx)}}$$

↓ 4257

$$\frac{\tan(e+fx)}{4f(a \sec(e+fx) + a)^{5/2}(c - c \sec(e+fx))^{3/2}} - \frac{3 \tan(e+fx) \left(\frac{\operatorname{arctanh}(\cos(e+fx))}{2f} - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right)}{4a^2c\sqrt{a \sec(e+fx) + a}\sqrt{c - c \sec(e+fx)}}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(3/2)),x]`

output `Tan[e + f*x]/(4*f*(a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(3/2)) - (3*(ArcTanh[Cos[e + f*x]]/(2*f) - (Cot[e + f*x]*Csc[e + f*x])/(2*f))*Tan[e + f*x])/(4*a^2*c*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.149.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3091 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4447 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(m_), x_Symbol] := Simp[((-a)*c)^(m + 1/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Int[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]`

rule 4448 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ((ILtQ[m, 0] && ILtQ[n - 1/2, 0]) || (ILtQ[m - 1/2, 0] && ILtQ[n - 1/2, 0] && LtQ[m, n]))`

3.149.4 Maple [A] (verified)

Time = 2.47 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.47

method	result
default	$\frac{\sqrt{2} \sqrt{-\frac{2a}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}} (1-\cos(fx+e)) \left((1-\cos(fx+e))^6 \csc(fx+e)^6 - 6(1-\cos(fx+e))^4 \csc(fx+e)^4 + 12 \ln(-\cot(fx+e)) \right)}{64f a^3 \left((1-\cos(fx+e))^2 \csc(fx+e)^2 - 1 \right) \left(\frac{c(1-\cos(fx+e))^2 \csc(fx+e)^2}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1} \right)^{\frac{3}{2}}}$
risch	$\frac{i(5e^{5i(fx+e)} + 2e^{4i(fx+e)} + 2e^{3i(fx+e)} + 2e^{2i(fx+e)} + 5e^{i(fx+e)})}{4a^2c(1+e^{2i(fx+e)})(e^{i(fx+e)}+1)^3 \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1) \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f} + \frac{3i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1) \ln(e^{i(fx+e)})}{8a^2c(1+e^{2i(fx+e)}) \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f}$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2), x, method=_RET URNVERBOSE)`

$$3.149. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} (c-c \sec(e+fx))^{3/2}} dx$$

output $1/64/f*2^{(1/2)}/a^3*(-2*a/((1-\cos(f*x+e))^2*\csc(f*x+e)^2-1))^{(1/2)}/((1-\cos(f*x+e))^2*\csc(f*x+e)^2-1)/(c*(1-\cos(f*x+e))^2/((1-\cos(f*x+e))^2*\csc(f*x+e)^2-1)*\csc(f*x+e)^2)^{(3/2)}*(1-\cos(f*x+e))*((1-\cos(f*x+e))^6*\csc(f*x+e)^6-6*(1-\cos(f*x+e))^4*\csc(f*x+e)^4+12*\ln(-\cot(f*x+e)+\csc(f*x+e))*(1-\cos(f*x+e))^2*\csc(f*x+e)^2+2)*\csc(f*x+e)$

3.149.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 536, normalized size of antiderivative = 3.67

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{3/2}} dx = \left[\frac{3(\cos(fx+e)^3 + \cos(fx+e)^2 - \cos(fx+e) - 1)}{\dots} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2),x, algorithm="fracas")`

output $[-1/16*(3*(\cos(f*x+e))^3 + \cos(f*x+e)^2 - \cos(f*x+e) - 1)*\sqrt{-a*c}*\log(-4*(2*\sqrt{-a*c})*\sqrt{(a*\cos(f*x+e)+a)/\cos(f*x+e)}*\sqrt{(c*\cos(f*x+e)-c)/\cos(f*x+e)}*\cos(f*x+e)^2 + (a*c*\cos(f*x+e)^2 + a*c)*\sin(f*x+e))/((\cos(f*x+e)^2 - 1)*\sin(f*x+e)))*\sin(f*x+e) - 2*(5*\cos(f*x+e)^3 + \cos(f*x+e)^2 - 2*\cos(f*x+e))*\sqrt{(a*\cos(f*x+e)+a)/\cos(f*x+e)}*\sqrt{(c*\cos(f*x+e)-c)/\cos(f*x+e))}/((a^3*c^2*f*\cos(f*x+e)^3 + a^3*c^2*f*\cos(f*x+e)^2 - a^3*c^2*f*\cos(f*x+e) - a^3*c^2*f)*\sin(f*x+e)), 1/8*(3*(\cos(f*x+e))^3 + \cos(f*x+e)^2 - \cos(f*x+e) - 1)*\sqrt{a*c}*\arctan(\sqrt{a*c}*\sqrt{(a*\cos(f*x+e)+a)/\cos(f*x+e)}*\sqrt{(c*\cos(f*x+e)-c)/\cos(f*x+e)})/(a*c*\sin(f*x+e)))*\sin(f*x+e) + (5*\cos(f*x+e)^3 + \cos(f*x+e)^2 - 2*\cos(f*x+e))*\sqrt{(a*\cos(f*x+e)+a)/\cos(f*x+e)}*\sqrt{(c*\cos(f*x+e)-c)/\cos(f*x+e))}/((a^3*c^2*f*\cos(f*x+e)^3 + a^3*c^2*f*\cos(f*x+e)^2 - a^3*c^2*f*\cos(f*x+e) - a^3*c^2*f)*\sin(f*x+e))]$

3.149.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} (c - c \sec(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(3/2),x)`

output `Timed out`

3.149.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} (c - c \sec(e + fx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.`

3.149.8 Giac [A] (verification not implemented)

Time = 1.90 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.03

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} (c - c \sec(e + fx))^{3/2}} dx = \frac{2 \left(3 c \tan\left(\frac{1}{2} f x + \frac{1}{2} e\right)^2 - c \right)}{c \tan\left(\frac{1}{2} f x + \frac{1}{2} e\right)^2} - \frac{\left(c \tan\left(\frac{1}{2} f x + \frac{1}{2} e\right)^2 - c \right)^2 c^2 - 4 \left(c \tan\left(\frac{1}{2} f x + \frac{1}{2} e\right)^2 - c \right) c^4}{c^4} \frac{1}{32 \sqrt{-a c a^2 f} |c| \operatorname{sgn}\left(\tan\left(\frac{1}{2} f x + \frac{1}{2} e\right)\right)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(3/2),x, algo
rithm="giac")`

output `1/32*(2*(3*c*tan(1/2*f*x + 1/2*e)^2 - c)/(c*tan(1/2*f*x + 1/2*e)^2) - ((c*
tan(1/2*f*x + 1/2*e)^2 - c)^2*c^2 - 4*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^3)/
c^4 - 6*log(abs(c)*tan(1/2*f*x + 1/2*e)^2) + 6*log(abs(c)) - 4)/(sqrt(-a*c
) * a^2 * f * abs(c) * sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e)))`

3.149. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} (c-c \sec(e+fx))^{3/2}} dx$

3.149.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{3/2}} dx = \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^{5/2} \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(3/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(3/2)), x)`

3.150 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2}} dx$

3.150.1 Optimal result 1056
 3.150.2 Mathematica [A] (verified) 1056
 3.150.3 Rubi [A] (verified) 1057
 3.150.4 Maple [A] (verified) 1059
 3.150.5 Fricas [A] (verification not implemented) 1059
 3.150.6 Sympy [F(-1)] 1060
 3.150.7 Maxima [B] (verification not implemented) 1060
 3.150.8 Giac [A] (verification not implemented) 1061
 3.150.9 Mupad [F(-1)] 1062

3.150.1 Optimal result

Integrand size = 36, antiderivative size = 160

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2}} dx = \frac{3 \csc(e+fx)}{8a^2c^2f\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} - \frac{\cot^2(e+fx) \csc(e+fx)}{4a^2c^2f\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} - \frac{3\operatorname{arctanh}(\cos(e+fx)) \tan(e+fx)}{8a^2c^2f\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}}$$

output `3/8*csc(f*x+e)/a^2/c^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-1/4*cot(f*x+e)^2*csc(f*x+e)/a^2/c^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)-3/8*arctanh(cos(f*x+e))*tan(f*x+e)/a^2/c^2/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.150.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.48

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2}} dx = \frac{(3-2 \cot^2(e+fx)) \csc(e+fx) - 3\operatorname{arctanh}(\sec(e+fx))}{8a^2c^2f\sqrt{a(1+\sec(e+fx))}\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(5/2)),x]`

3.150. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2}} dx$

output $((3 - 2*\text{Cot}[e + f*x]^2)*\text{Csc}[e + f*x] - 3*\text{ArcTanh}[\text{Sec}[e + f*x]]*\text{Tan}[e + f*x]) / (8*a^2*c^2*f*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])] * \text{Sqrt}[c - c*\text{Sec}[e + f*x]])$

3.150.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.64, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4447, 25, 3042, 3091, 3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^{5/2} (c - c \sec(e+fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^{5/2} (c - c \csc(e+fx + \frac{\pi}{2}))^{5/2}} dx$$

↓ 4447

$$\frac{\tan(e+fx) \int -\cot^4(e+fx) \csc(e+fx) dx}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

↓ 25

$$\frac{\tan(e+fx) \int \cot^4(e+fx) \csc(e+fx) dx}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

↓ 3042

$$\frac{\tan(e+fx) \int \sec(e+fx - \frac{\pi}{2}) \tan(e+fx - \frac{\pi}{2})^4 dx}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

↓ 3091

$$\frac{\tan(e+fx) \left(-\frac{3}{4} \int \cot^2(e+fx) \csc(e+fx) dx - \frac{\cot^3(e+fx) \csc(e+fx)}{4f} \right)}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

↓ 3042

$$\frac{\tan(e+fx) \left(-\frac{3}{4} \int \sec(e+fx - \frac{\pi}{2}) \tan(e+fx - \frac{\pi}{2})^2 dx - \frac{\cot^3(e+fx) \csc(e+fx)}{4f} \right)}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

↓ 3091

3.150. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} (c-c \sec(e+fx))^{5/2}} dx$

$$\frac{\tan(e+fx) \left(-\frac{3}{4} \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right) - \frac{\cot^3(e+fx) \csc(e+fx)}{4f} \right)}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

↓ 3042

$$\frac{\tan(e+fx) \left(-\frac{3}{4} \left(-\frac{1}{2} \int \csc(e+fx) dx - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right) - \frac{\cot^3(e+fx) \csc(e+fx)}{4f} \right)}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

↓ 4257

$$\frac{\tan(e+fx) \left(-\frac{3}{4} \left(\frac{\operatorname{arctanh}(\cos(e+fx))}{2f} - \frac{\cot(e+fx) \csc(e+fx)}{2f} \right) - \frac{\cot^3(e+fx) \csc(e+fx)}{4f} \right)}{a^2 c^2 \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^(5/2)*(c - c*Sec[e + f*x])^(5/2)),x]`

output `((-1/4*(Cot[e + f*x]^3*Csc[e + f*x])/f - (3*(ArcTanh[Cos[e + f*x]]/(2*f) - (Cot[e + f*x]*Csc[e + f*x])/(2*f)))/4)*Tan[e + f*x])/(a^2*c^2*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.150.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3091 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4447 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(cs
c[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(m_), x_Symbol] :> Simp[((-a)*c)^(m +
1/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) In
t[Csc[e + f*x]*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f}, x] &
& EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]
```

3.150.4 Maple [A] (verified)

Time = 2.29 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.99

method	result
default	$\frac{(3 \cos(fx+e)^4 \ln(-\cot(fx+e)+\csc(fx+e))-6 \cos(fx+e)^2 \ln(-\cot(fx+e)+\csc(fx+e))-5 \cos(fx+e)^3+3 \ln(-\cot(fx+e)+\csc(fx+e)))}{8 f a^3 \sqrt{-c(\sec(fx+e)-1)}(\sec(fx+e)-1)^2 c^2(\cos(fx+e)+1)^3}$
risch	$\frac{i(5 e^{7i(fx+e)}+3 e^{5i(fx+e)}+3 e^{3i(fx+e)}+5 e^{i(fx+e)})}{4 a^2 c^2(1+e^{2i(fx+e)})(e^{i(fx+e)}+1)^3} \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}} (e^{i(fx+e)}-1)^3 \sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}} f + \frac{3i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1) \ln}{8 a^2 c^2(1+e^{2i(fx+e)}) \sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}}}$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 1/8/f/a^3*(3*cos(f*x+e)^4*ln(-cot(f*x+e)+csc(f*x+e))-6*cos(f*x+e)^2*ln(-co
t(f*x+e)+csc(f*x+e))-5*cos(f*x+e)^3+3*ln(-cot(f*x+e)+csc(f*x+e))+3*cos(f*x
+e))*(a*(sec(f*x+e)+1))^(1/2)/(-c*(sec(f*x+e)-1))^(1/2)/(sec(f*x+e)-1)^2/c
^2/(cos(f*x+e)+1)^3*tan(f*x+e)*sec(f*x+e)
```

3.150.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 482, normalized size of antiderivative = 3.01

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2}(c - c \sec(e + fx))^{5/2}} dx = \left[-\frac{3(\cos(fx + e)^4 - 2 \cos(fx + e)^2 + 1) \sqrt{-ac} \log\left(-\right)}{\dots} \right]$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x, algo
rithm="fracas")
```

3.150.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2}(c-c \sec(e+fx))^{5/2}} dx$$

output `[-1/16*(3*(cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*sqrt(-a*c)*log(-4*(2*sqrt(-a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)^2 + (a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e))/((cos(f*x + e)^2 - 1)*sin(f*x + e)))*sin(f*x + e) - 2*(5*cos(f*x + e)^4 - 3*cos(f*x + e)^2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^3*c^3*f*cos(f*x + e)^4 - 2*a^3*c^3*f*cos(f*x + e)^2 + a^3*c^3*f)*sin(f*x + e)), 1/8*(3*(cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*sqrt(a*c)*arctan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/(a*c*sin(f*x + e)))*sin(f*x + e) + (5*cos(f*x + e)^4 - 3*cos(f*x + e)^2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)))/((a^3*c^3*f*cos(f*x + e)^4 - 2*a^3*c^3*f*cos(f*x + e)^2 + a^3*c^3*f)*sin(f*x + e))]`

3.150.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} (c - c \sec(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))**(5/2),x)`

output `Timed out`

3.150.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1659 vs. 2(142) = 284.

Time = 0.55 (sec) , antiderivative size = 1659, normalized size of antiderivative = 10.37

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} (c - c \sec(e + fx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="maxima")`

output `1/8*(3*(2*(4*cos(6*f*x + 6*e) - 6*cos(4*f*x + 4*e) + 4*cos(2*f*x + 2*e) - 1)*cos(8*f*x + 8*e) - cos(8*f*x + 8*e)^2 + 8*(6*cos(4*f*x + 4*e) - 4*cos(2*f*x + 2*e) + 1)*cos(6*f*x + 6*e) - 16*cos(6*f*x + 6*e)^2 + 12*(4*cos(2*f*x + 2*e) - 1)*cos(4*f*x + 4*e) - 36*cos(4*f*x + 4*e)^2 - 16*cos(2*f*x + 2*e)^2 + 4*(2*sin(6*f*x + 6*e) - 3*sin(4*f*x + 4*e) + 2*sin(2*f*x + 2*e))*sin(8*f*x + 8*e) - sin(8*f*x + 8*e)^2 + 16*(3*sin(4*f*x + 4*e) - 2*sin(2*f*x + 2*e))*sin(6*f*x + 6*e) - 16*sin(6*f*x + 6*e)^2 - 36*sin(4*f*x + 4*e)^2 + 48*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) - 16*sin(2*f*x + 2*e)^2 + 8*cos(2*f*x + 2*e) - 1)*arctan2(sin(f*x + e), cos(f*x + e) + 1) - 3*(2*(4*cos(6*f*x + 6*e) - 6*cos(4*f*x + 4*e) + 4*cos(2*f*x + 2*e) - 1)*cos(8*f*x + 8*e) - cos(8*f*x + 8*e)^2 + 8*(6*cos(4*f*x + 4*e) - 4*cos(2*f*x + 2*e) + 1)*cos(6*f*x + 6*e) - 16*cos(6*f*x + 6*e)^2 + 12*(4*cos(2*f*x + 2*e) - 1)*cos(4*f*x + 4*e) - 36*cos(4*f*x + 4*e)^2 - 16*cos(2*f*x + 2*e)^2 + 4*(2*sin(6*f*x + 6*e) - 3*sin(4*f*x + 4*e) + 2*sin(2*f*x + 2*e))*sin(8*f*x + 8*e) - sin(8*f*x + 8*e)^2 + 16*(3*sin(4*f*x + 4*e) - 2*sin(2*f*x + 2*e))*sin(6*f*x + 6*e) - 16*sin(6*f*x + 6*e)^2 - 36*sin(4*f*x + 4*e)^2 + 48*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) - 16*sin(2*f*x + 2*e)^2 + 8*cos(2*f*x + 2*e) - 1)*arctan2(sin(f*x + e), cos(f*x + e) - 1) - 2*(5*sin(7*f*x + 7*e) + 3*sin(5*f*x + 5*e) + 3*sin(3*f*x + 3*e) + 5*sin(f*x + e))*cos(8*f*x + 8*e) - 20*(2*sin(6*f*x + 6*e) - 3*sin(4*f*x + 4*e) + 2*sin(2*f*x + 2*e))*cos(7*f*x + 7*e) ...`

3.150.8 Giac [A] (verification not implemented)

Time = 1.86 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.14

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^{5/2} (c - c \sec(e + fx))^{5/2}} dx = \frac{(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)^2 c^2 - 6 (c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c) c^3}{c^4} - \frac{18 (c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c)^2 + 28 (c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c) c + 11 c^2}{c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4} + 12 \log(|c| \tan(\frac{1}{2} fx + \frac{1}{2} e)) - \frac{64 \sqrt{-aca^2 cf} |c| \operatorname{sgn} \left(\tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + \tan(\frac{1}{2} fx + \frac{1}{2} e) \right)}{64 \sqrt{-aca^2 cf} |c| \operatorname{sgn} \left(\tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + \tan(\frac{1}{2} fx + \frac{1}{2} e) \right)}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `-1/64*(((c*tan(1/2*f*x + 1/2*e)^2 - c)^2*c^2 - 6*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c^3)/c^4 - (18*(c*tan(1/2*f*x + 1/2*e)^2 - c)^2 + 28*(c*tan(1/2*f*x + 1/2*e)^2 - c)*c + 11*c^2)/(c^2*tan(1/2*f*x + 1/2*e)^4) + 12*log(abs(c)*tan(1/2*f*x + 1/2*e)^2) - 12*log(abs(c)) + 11)/(sqrt(-a*c)*a^2*c*f*abs(c)*sgn(tan(1/2*f*x + 1/2*e)^3 + tan(1/2*f*x + 1/2*e)))`

3.150. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^{5/2} (c-c \sec(e+fx))^{5/2}} dx$

3.150.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^{5/2}(c-c\sec(e+fx))^{5/2}} dx = \int \frac{1}{\cos(e+fx) \left(a + \frac{a}{\cos(e+fx)}\right)^{5/2} \left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(5/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(5/2)*(c - c/cos(e + f*x))^(5/2)), x)`

3.151 $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx$

3.151.1 Optimal result	1063
3.151.2 Mathematica [A] (verified)	1063
3.151.3 Rubi [A] (verified)	1064
3.151.4 Maple [F]	1065
3.151.5 Fricas [F]	1066
3.151.6 Sympy [F]	1066
3.151.7 Maxima [F]	1066
3.151.8 Giac [F]	1067
3.151.9 Mupad [F(-1)]	1067

3.151.1 Optimal result

Integrand size = 32, antiderivative size = 101

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx = \frac{2^{\frac{1}{2}+n} c \operatorname{Hypergeometric2F1}\left(\frac{1}{2} + m, \frac{1}{2} - n, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e + fx))\right) (1 - \sec(e + fx))^{\frac{1}{2}-n} (a + a \sec(e + fx))^m}{f(1 + 2m)}$$

```
output -2^(1/2+n)*c*hypergeom([1/2-n, 1/2+m], [3/2+m], 1/2+1/2*sec(f*x+e))*(1-sec(f*x+e))^(1/2-n)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(-1+n)*tan(f*x+e)/f/(1+2*m)
```

3.151.2 Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.94

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx = \frac{2^{\frac{1}{2}+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2} - m, \frac{1}{2} + n, \frac{3}{2} + n, \frac{1}{2}(1 - \sec(e + fx))\right) (1 + \sec(e + fx))^{-\frac{1}{2}-m} (a(1 + \sec(e + fx)))^m}{f + 2fn}$$

```
input Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^n,x]
```


output $(2^{(1/2 + m)} \text{Hypergeometric2F1}[1/2 - m, 1/2 + n, 3/2 + n, (1 - \text{Sec}[e + f*x])/2] * (1 + \text{Sec}[e + f*x])^{(-1/2 - m)} * (a * (1 + \text{Sec}[e + f*x]))^m * (c - c * \text{Sec}[e + f*x])^n * \text{Tan}[e + f*x]) / (f + 2*f*n)$

3.151.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^n dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^n dx$$

$$\downarrow 4449$$

$$\frac{a \tan(e + fx) \int (\sec(e + fx)a + a)^{m-\frac{1}{2}} (c - c \sec(e + fx))^{n-\frac{1}{2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}}$$

$$\downarrow 80$$

$$\frac{a c 2^{n-\frac{1}{2}} \tan(e + fx) (1 - \sec(e + fx))^{\frac{1}{2}-n} (c - c \sec(e + fx))^{n-1} \int \left(\frac{1}{2} - \frac{1}{2} \sec(e + fx)\right)^{n-\frac{1}{2}} (\sec(e + fx)a + a)^{m-\frac{1}{2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a}}$$

$$\downarrow 79$$

$$\frac{c 2^{n+\frac{1}{2}} \tan(e + fx) (1 - \sec(e + fx))^{\frac{1}{2}-n} (a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{n-1} \text{Hypergeometric2F1}\left(m + \frac{1}{2}, \frac{1}{2} - n, \frac{3}{2} + m, \frac{1 - \sec(e + fx)}{2}\right)}{f(2m + 1)}$$

input $\text{Int}[\text{Sec}[e + f*x] * (a + a * \text{Sec}[e + f*x])^m * (c - c * \text{Sec}[e + f*x])^n, x]$

output $-((2^{(1/2 + n)} * c * \text{Hypergeometric2F1}[1/2 + m, 1/2 - n, 3/2 + m, (1 + \text{Sec}[e + f*x])/2] * (1 - \text{Sec}[e + f*x])^{(1/2 - n)} * (a + a * \text{Sec}[e + f*x])^m * (c - c * \text{Sec}[e + f*x])^{(-1 + n)} * \text{Tan}[e + f*x]) / (f * (1 + 2 * m)))$

3.151. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx$

3.151.3.1 Defintions of rubi rules used

- rule 79 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`
- rule 80 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4449 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.151.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (c - c \sec(fx + e))^n dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^n,x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^n,x)`

3.151.5 Fricas [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^n \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^n,x, algorithm="fricas")`

output `integral((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^n*sec(f*x + e), x)`

3.151.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx$$

$$= \int (a(\sec(e + fx) + 1))^m (-c(\sec(e + fx) - 1))^n \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^n,x)`

output `Integral((a*(sec(e + f*x) + 1))^m*(-c*(sec(e + f*x) - 1))^n*sec(e + f*x), x)`

3.151.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^n \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^n,x, algorithm="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^n*sec(f*x + e), x)`

3.151.8 Giac [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx \\ &= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^n \sec(fx + e) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^n,x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^n*sec(f*x + e), x)`

3.151.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^n dx \\ &= \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m \left(c - \frac{c}{\cos(e+fx)}\right)^n}{\cos(e + fx)} dx \end{aligned}$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^n)/cos(e + f*x),x)`

output `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^n)/cos(e + f*x), x)`

3.152 $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^2 dx$

3.152.1 Optimal result	1068
3.152.2 Mathematica [A] (verified)	1068
3.152.3 Rubi [A] (verified)	1069
3.152.4 Maple [F]	1070
3.152.5 Fricas [F]	1071
3.152.6 Sympy [F]	1071
3.152.7 Maxima [F]	1072
3.152.8 Giac [F]	1072
3.152.9 Mupad [F(-1)]	1072

3.152.1 Optimal result

Integrand size = 32, antiderivative size = 92

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^2 dx$$

$$= \frac{2^{\frac{1}{2}+m} a \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{1}{2}-m, \frac{7}{2}, \frac{1}{2}(1-\sec(e+fx))\right) (1+\sec(e+fx))^{\frac{1}{2}-m} (a+a \sec(e+fx))^{-1}}{5f}$$

output

```
1/5*2^(1/2+m)*a*hypergeom([5/2, 1/2-m], [7/2], 1/2-1/2*sec(f*x+e))*(1+sec(f*x+e))^(1/2-m)*(a+a*sec(f*x+e))^(-1+m)*(c-c*sec(f*x+e))^2*tan(f*x+e)/f
```

3.152.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.97

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^2 dx$$

$$= \frac{2^{\frac{1}{2}+m} c^2 \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{1}{2}-m, \frac{7}{2}, \frac{1}{2}(1-\sec(e+fx))\right) (-1+\sec(e+fx))^2 (1+\sec(e+fx))^{-\frac{1}{2}-m}}{5f}$$

input

```
Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^2,x]
```

output $(2^{(1/2 + m)}c^2\text{Hypergeometric2F1}[5/2, 1/2 - m, 7/2, (1 - \text{Sec}[e + f*x])/2] * (-1 + \text{Sec}[e + f*x])^{-2} * (1 + \text{Sec}[e + f*x])^{(-1/2 - m)} * (a * (1 + \text{Sec}[e + f*x]))^m * \text{Tan}[e + f*x]) / (5*f)$

3.152.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(c - c \sec(e + fx))^2 (a \sec(e + fx) + a)^m dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m dx$$

$$\downarrow 4449$$

$$-\frac{ac \tan(e + fx) \int (\sec(e + fx)a + a)^{m-\frac{1}{2}} (c - c \sec(e + fx))^{3/2} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}}$$

$$\downarrow 80$$

$$-\frac{ac2^{m-\frac{1}{2}} \tan(e + fx) (\sec(e + fx) + 1)^{\frac{1}{2}-m} (a \sec(e + fx) + a)^{m-1} \int \left(\frac{1}{2} \sec(e + fx) + \frac{1}{2}\right)^{m-\frac{1}{2}} (c - c \sec(e + fx))^{3/2} d \sec(e + fx)}{f \sqrt{c - c \sec(e + fx)}}$$

$$\downarrow 79$$

$$\frac{a2^{m+\frac{1}{2}} \tan(e + fx) (c - c \sec(e + fx))^2 (\sec(e + fx) + 1)^{\frac{1}{2}-m} (a \sec(e + fx) + a)^{m-1} \text{Hypergeometric2F1}\left(\frac{5}{2}, \frac{1}{2} - m, \frac{7}{2}, \frac{1 - \sec(e + fx)}{2}\right)}{5f}$$

input $\text{Int}[\text{Sec}[e + f*x] * (a + a * \text{Sec}[e + f*x])^m * (c - c * \text{Sec}[e + f*x])^2, x]$

output $(2^{(1/2 + m)} * a * \text{Hypergeometric2F1}[5/2, 1/2 - m, 7/2, (1 - \text{Sec}[e + f*x])/2] * (1 + \text{Sec}[e + f*x])^{(1/2 - m)} * (a + a * \text{Sec}[e + f*x])^{(-1 + m)} * (c - c * \text{Sec}[e + f*x])^2 * \text{Tan}[e + f*x]) / (5*f)$

3.152. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^2 dx$

3.152.3.1 Defintions of rubi rules used

- rule 79 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`
- rule 80 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4449 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.152.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (c - c \sec(fx + e))^2 dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^2,x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^2,x)`

3.152.5 Fracas [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^2 dx$$

$$= \int (c \sec(fx + e) - c)^2 (a \sec(fx + e) + a)^m \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^2,x, algorithm="fracas")`

output `integral((c^2*sec(f*x + e)^3 - 2*c^2*sec(f*x + e)^2 + c^2*sec(f*x + e))*(a*sec(f*x + e) + a)^m, x)`

3.152.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^2 dx$$

$$= c^2 \left(\int (a \sec(e + fx) + a)^m \sec(e + fx) dx \right.$$

$$\quad \left. + \int (-2(a \sec(e + fx) + a)^m \sec^2(e + fx)) dx \right.$$

$$\quad \left. + \int (a \sec(e + fx) + a)^m \sec^3(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**m*(c-c*sec(f*x+e))**2,x)`

output `c**2*(Integral((a*sec(e + f*x) + a)**m*sec(e + f*x), x) + Integral(-2*(a*sec(e + f*x) + a)**m*sec(e + f*x)**2, x) + Integral((a*sec(e + f*x) + a)**m*sec(e + f*x)**3, x))`

3.152.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^2 dx$$

$$= \int (c \sec(fx + e) - c)^2 (a \sec(fx + e) + a)^m \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output `integrate((c*sec(f*x + e) - c)^2*(a*sec(f*x + e) + a)^m*sec(f*x + e), x)`

3.152.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^2 dx$$

$$= \int (c \sec(fx + e) - c)^2 (a \sec(fx + e) + a)^m \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `integrate((c*sec(f*x + e) - c)^2*(a*sec(f*x + e) + a)^m*sec(f*x + e), x)`

3.152.9 Mupad [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^2 dx$$

$$= \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m \left(c - \frac{c}{\cos(e+fx)}\right)^2}{\cos(e + fx)} dx$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^2)/cos(e + f*x),x)`

output `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^2)/cos(e + f*x), x)`

3.153 $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx)) dx$

3.153.1 Optimal result	1073
3.153.2 Mathematica [A] (verified)	1073
3.153.3 Rubi [A] (verified)	1074
3.153.4 Maple [F]	1075
3.153.5 Fracas [F]	1076
3.153.6 Sympy [F]	1076
3.153.7 Maxima [F]	1076
3.153.8 Giac [F]	1077
3.153.9 Mupad [F(-1)]	1077

3.153.1 Optimal result

Integrand size = 30, antiderivative size = 90

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx)) dx = \frac{2^{\frac{1}{2}+m} a \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1}{2}-m, \frac{5}{2}, \frac{1}{2}(1-\sec(e+fx))\right) (1+\sec(e+fx))^{\frac{1}{2}-m} (a+a \sec(e+fx))^{-1}}{3f}$$

```
output 1/3*2^(1/2+m)*a*hypergeom([3/2, 1/2-m], [5/2], 1/2-1/2*sec(f*x+e))*(1+sec(f*x+e))^(1/2-m)*(a+a*sec(f*x+e))^(1-m)*(c-c*sec(f*x+e))*tan(f*x+e)/f
```

3.153.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.94

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx)) dx = \frac{2^{\frac{1}{2}+m} c \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1}{2}-m, \frac{5}{2}, \frac{1}{2}(1-\sec(e+fx))\right) (-1+\sec(e+fx))(1+\sec(e+fx))^{-\frac{1}{2}-m}}{3f}$$

```
input Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x]),x]
```

output $-1/3*(2^{(1/2 + m)*c*Hypergeometric2F1[3/2, 1/2 - m, 5/2, (1 - \text{Sec}[e + f*x])/2]*(-1 + \text{Sec}[e + f*x])*(1 + \text{Sec}[e + f*x])^{(-1/2 - m)*(a*(1 + \text{Sec}[e + f*x]))^m*\text{Tan}[e + f*x])/f$

3.153.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(c - c \sec(e + fx))(a \sec(e + fx) + a)^m dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m dx$$

$$\downarrow 4449$$

$$-\frac{ac \tan(e + fx) \int (\sec(e + fx)a + a)^{m-\frac{1}{2}} \sqrt{c - c \sec(e + fx)} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}}$$

$$\downarrow 80$$

$$-\frac{ac2^{m-\frac{1}{2}} \tan(e + fx) (\sec(e + fx) + 1)^{\frac{1}{2}-m} (a \sec(e + fx) + a)^{m-1} \int \left(\frac{1}{2} \sec(e + fx) + \frac{1}{2}\right)^{m-\frac{1}{2}} \sqrt{c - c \sec(e + fx)} dx}{f \sqrt{c - c \sec(e + fx)}}$$

$$\downarrow 79$$

$$\frac{a2^{m+\frac{1}{2}} \tan(e + fx) (c - c \sec(e + fx)) (\sec(e + fx) + 1)^{\frac{1}{2}-m} (a \sec(e + fx) + a)^{m-1} \text{Hypergeometric2F1}\left(\frac{3}{2}, \frac{1}{2} - m, \frac{5}{2}, \frac{1 - \sec(e + fx)}{2}\right)}{3f}$$

input $\text{Int}[\text{Sec}[e + f*x]*(a + a*\text{Sec}[e + f*x])^m*(c - c*\text{Sec}[e + f*x]),x]$

output $(2^{(1/2 + m)*a*Hypergeometric2F1[3/2, 1/2 - m, 5/2, (1 - \text{Sec}[e + f*x])/2]*(1 + \text{Sec}[e + f*x])^{(1/2 - m)*(a + a*\text{Sec}[e + f*x])^{(-1 + m)*(c - c*\text{Sec}[e + f*x])*Tan}[e + f*x])/(3*f)$

3.153. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx)) dx$

3.153.3.1 Defintions of rubi rules used

```
rule 79 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((
a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1
, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x]
&& !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m]
|| !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

```
rule 80 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c
+ d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))
^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)
), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !Integ
erQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4449 Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(c
sc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f
*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a +
b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a,
b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
```

3.153.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (c - c \sec(fx + e)) dx$$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e)),x)
```

```
output int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e)),x)
```

3.153.5 Fricas [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx)) dx \\ &= \int -(c \sec(fx + e) - c)(a \sec(fx + e) + a)^m \sec(fx + e) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `integral(-(c*sec(f*x + e)^2 - c*sec(f*x + e))*(a*sec(f*x + e) + a)^m, x)`

3.153.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx)) dx \\ &= -c \left(\int (-(a \sec(e + fx) + a)^m \sec(e + fx)) dx \right. \\ & \quad \left. + \int (a \sec(e + fx) + a)^m \sec^2(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**m*(c-c*sec(f*x+e)),x)`

output `-c*(Integral(-(a*sec(e + f*x) + a)**m*sec(e + f*x), x) + Integral((a*sec(e + f*x) + a)**m*sec(e + f*x)**2, x))`

3.153.7 Maxima [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx)) dx \\ &= \int -(c \sec(fx + e) - c)(a \sec(fx + e) + a)^m \sec(fx + e) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-integrate((c*sec(f*x + e) - c)*(a*sec(f*x + e) + a)^m*sec(f*x + e), x)`

3.153.8 Giac [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx)) dx \\ &= \int -(c \sec(fx + e) - c)(a \sec(fx + e) + a)^m \sec(fx + e) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e)),x, algorithm="giac")`

output `integrate(-(c*sec(f*x + e) - c)*(a*sec(f*x + e) + a)^m*sec(f*x + e), x)`

3.153.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m(c - c \sec(e + fx)) dx \\ &= \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m \left(c - \frac{c}{\cos(e+fx)}\right)}{\cos(e + fx)} dx \end{aligned}$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x)))/cos(e + f*x),x)`

output `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x)))/cos(e + f*x), x)`

3.154
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{c-c \sec(e+fx)} dx$$

3.154.1 Optimal result 1078
 3.154.2 Mathematica [F] 1078
 3.154.3 Rubi [A] (verified) 1079
 3.154.4 Maple [F] 1080
 3.154.5 Fricas [F] 1081
 3.154.6 Sympy [F] 1081
 3.154.7 Maxima [F] 1081
 3.154.8 Giac [F] 1082
 3.154.9 Mupad [F(-1)] 1082

3.154.1 Optimal result

Integrand size = 32, antiderivative size = 90

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{c-c \sec(e+fx)} dx = \frac{2^{\frac{1}{2}+m} a \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{2}-m, \frac{1}{2}, \frac{1}{2}(1-\sec(e+fx))\right) (1+\sec(e+fx))^{\frac{1}{2}-m} (a+a \sec(e+fx))}{f(c-c \sec(e+fx))}$$

output `-2^(1/2+m)*a*hypergeom([-1/2, 1/2-m], [1/2], 1/2-1/2*sec(f*x+e))*(1+sec(f*x+e))^(1/2-m)*(a+a*sec(f*x+e))^(1+m)*tan(f*x+e)/f/(c-c*sec(f*x+e))`

3.154.2 Mathematica [F]

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{c-c \sec(e+fx)} dx = \int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{c-c \sec(e+fx)} dx$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x]),x]`

output `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x]), x]`

3.154.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^m}{c-c\sec(e+fx)} dx$$

$$\downarrow 3042$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^m}{c-c\csc\left(e+fx+\frac{\pi}{2}\right)} dx$$

$$\downarrow 4449$$

$$\frac{a\csc\left(e+fx+\frac{\pi}{2}\right)\int\frac{\left(\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^{m-\frac{1}{2}}d\csc\left(e+fx+\frac{\pi}{2}\right)}{\csc\left(e+fx+\frac{\pi}{2}\right)}}{f\sqrt{a\csc\left(e+fx+\frac{\pi}{2}\right)+a}\sqrt{c-c\csc\left(e+fx+\frac{\pi}{2}\right)}}$$

$$\downarrow 80$$

$$\frac{a2^{m-\frac{1}{2}}\tan(e+fx)(\sec(e+fx)+1)^{\frac{1}{2}-m}(a\sec(e+fx)+a)^{m-1}\int\frac{\left(\frac{1}{2}\sec(e+fx)+\frac{1}{2}\right)^{m-\frac{1}{2}}d\sec(e+fx)}{(c-c\sec(e+fx))^{\frac{3}{2}}}}{f\sqrt{c-c\sec(e+fx)}}$$

$$\downarrow 79$$

$$\frac{a2^{m+\frac{1}{2}}\tan(e+fx)(\sec(e+fx)+1)^{\frac{1}{2}-m}(a\sec(e+fx)+a)^{m-1}\text{Hypergeometric2F1}\left(-\frac{1}{2},\frac{1}{2}-m,\frac{1}{2},\frac{1}{2}(1-\sec(e+fx))\right)}{f(c-c\sec(e+fx))}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x]),x]`

output `-((2^(1/2 + m)*a*Hypergeometric2F1[-1/2, 1/2 - m, 1/2, (1 - Sec[e + f*x])/2]*(1 + Sec[e + f*x])^(1/2 - m)*(a + a*Sec[e + f*x])^(-1 + m)*Tan[e + f*x])/f*(c - c*Sec[e + f*x]))`

3.154.3.1 Defintions of rubi rules used

```
rule 79 Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

```
rule 80 Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4449 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
```

3.154.4 Maple [F]

$$\int \frac{\sec(fx + e)(a + a \sec(fx + e))^m}{c - c \sec(fx + e)} dx$$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e)),x)
```

```
output int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e)),x)
```

3.154.5 Fricas [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{c-c\sec(e+fx)} dx = \int -\frac{(a\sec(fx+e)+a)^m \sec(fx+e)}{c\sec(fx+e)-c} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `integral(-(a*sec(f*x + e) + a)^m*sec(f*x + e)/(c*sec(f*x + e) - c), x)`

3.154.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{c-c\sec(e+fx)} dx = -\frac{\int \frac{(a\sec(e+fx)+a)^m \sec(e+fx)}{\sec(e+fx)-1} dx}{c}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e)),x)`

output `-Integral((a*sec(e + f*x) + a)**m*sec(e + f*x)/(sec(e + f*x) - 1), x)/c`

3.154.7 Maxima [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{c-c\sec(e+fx)} dx = \int -\frac{(a\sec(fx+e)+a)^m \sec(fx+e)}{c\sec(fx+e)-c} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(c*sec(f*x + e) - c), x)`

3.154.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{c-c\sec(e+fx)} dx = \int -\frac{(a\sec(fx+e)+a)^m \sec(fx+e)}{c\sec(fx+e)-c} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e)),x, algorithm="giac")`

output `integrate(-(a*sec(f*x + e) + a)^m*sec(f*x + e)/(c*sec(f*x + e) - c), x)`

3.154.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{c-c\sec(e+fx)} dx = -\int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m}{c-c\cos(e+fx)} dx$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))),x)`

output `-int((a + a/cos(e + f*x))^m/(c - c*cos(e + f*x)), x)`

3.155
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^2} dx$$

3.155.1 Optimal result 1083
 3.155.2 Mathematica [F] 1083
 3.155.3 Rubi [A] (verified) 1084
 3.155.4 Maple [F] 1085
 3.155.5 Fricas [F] 1086
 3.155.6 Sympy [F] 1086
 3.155.7 Maxima [F] 1086
 3.155.8 Giac [F] 1087
 3.155.9 Mupad [F(-1)] 1087

3.155.1 Optimal result

Integrand size = 32, antiderivative size = 92

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^2} dx = \frac{2^{\frac{1}{2}+m} a \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, \frac{1}{2}-m, -\frac{1}{2}, \frac{1}{2}(1-\sec(e+fx))\right) (1+\sec(e+fx))^{\frac{1}{2}-m} (a+a \sec(e+fx))}{3f(c-c \sec(e+fx))^2}$$

output `-1/3*2^(1/2+m)*a*hypergeom([-3/2, 1/2-m], [-1/2], 1/2-1/2*sec(f*x+e))*(1+sec(f*x+e))^(1/2-m)*(a+a*sec(f*x+e))^(-1+m)*tan(f*x+e)/f/(c-c*sec(f*x+e))^2`

3.155.2 Mathematica [F]

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^2} dx = \int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^2} dx$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^2, x]`

output `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^2, x]`

3.155.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^m}{(c-c\sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^m}{(c-c\csc(e+fx+\frac{\pi}{2}))^2} dx$$

↓ 4449

$$\frac{a \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{m-\frac{1}{2}}}{(c-c\sec(e+fx))^{5/2}} d\sec(e+fx)}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}}$$

↓ 80

$$\frac{ac^{2^{m-\frac{1}{2}}} \tan(e+fx)(\sec(e+fx)+1)^{\frac{1}{2}-m} (a\sec(e+fx)+a)^{m-1} \int \frac{(\frac{1}{2}\sec(e+fx)+\frac{1}{2})^{m-\frac{1}{2}}}{(c-c\sec(e+fx))^{5/2}} d\sec(e+fx)}{f\sqrt{c-c\sec(e+fx)}}$$

↓ 79

$$\frac{a2^{m+\frac{1}{2}} \tan(e+fx)(\sec(e+fx)+1)^{\frac{1}{2}-m} (a\sec(e+fx)+a)^{m-1} \text{Hypergeometric2F1}(-\frac{3}{2}, \frac{1}{2}-m, -\frac{1}{2}, \frac{1}{2}(1-\sec(e+fx)))}{3f(c-c\sec(e+fx))^2}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^2,x]`

output `-1/3*(2^(1/2 + m)*a*Hypergeometric2F1[-3/2, 1/2 - m, -1/2, (1 - Sec[e + f*x])/2]*(1 + Sec[e + f*x])^(1/2 - m)*(a + a*Sec[e + f*x])^(-1 + m)*Tan[e + f*x])/(f*(c - c*Sec[e + f*x])^2)`

3.155.3.1 Defintions of rubi rules used

```
rule 79 Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

```
rule 80 Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4449 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
```

3.155.4 Maple [F]

$$\int \frac{\sec(fx + e)(a + a \sec(fx + e))^m}{(c - c \sec(fx + e))^2} dx$$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^2,x)
```

```
output int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^2,x)
```

3.155.5 Fricas [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{(c-c\sec(e+fx))^2} dx = \int \frac{(a\sec(fx+e)+a)^m \sec(fx+e)}{(c\sec(fx+e)-c)^2} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^2,x, algorithm="fricas")`

output `integral((a*sec(f*x + e) + a)^m*sec(f*x + e)/(c^2*sec(f*x + e)^2 - 2*c^2*sec(f*x + e) + c^2), x)`

3.155.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{(c-c\sec(e+fx))^2} dx = \frac{\int \frac{(a\sec(e+fx)+a)^m \sec(e+fx)}{\sec^2(e+fx)-2\sec(e+fx)+1} dx}{c^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^2,x)`

output `Integral((a*sec(e + f*x) + a)**m*sec(e + f*x)/(sec(e + f*x)**2 - 2*sec(e + f*x) + 1), x)/c**2`

3.155.7 Maxima [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{(c-c\sec(e+fx))^2} dx = \int \frac{(a\sec(fx+e)+a)^m \sec(fx+e)}{(c\sec(fx+e)-c)^2} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^2,x, algorithm="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(c*sec(f*x + e) - c)^2, x)`

3.155.8 Giac [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{(c-c\sec(e+fx))^2} dx = \int \frac{(a\sec(fx+e)+a)^m \sec(fx+e)}{(c\sec(fx+e)-c)^2} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^2,x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(c*sec(f*x + e) - c)^2, x)`

3.155.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{(c-c\sec(e+fx))^2} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^2} dx$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^2),x)`

output `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^2), x)`

3.156 $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{5/2} dx$

3.156.1 Optimal result	1088
3.156.2 Mathematica [A] (verified)	1089
3.156.3 Rubi [A] (verified)	1089
3.156.4 Maple [F]	1091
3.156.5 Fricas [A] (verification not implemented)	1091
3.156.6 Sympy [F(-1)]	1092
3.156.7 Maxima [A] (verification not implemented)	1092
3.156.8 Giac [F]	1093
3.156.9 Mupad [F(-1)]	1093

3.156.1 Optimal result

Integrand size = 34, antiderivative size = 160

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{5/2} dx =$$

$$\frac{64c^3(a+a \sec(e+fx))^m \tan(e+fx)}{f(5+2m)(3+8m+4m^2)\sqrt{c-c \sec(e+fx)}}$$

$$- \frac{16c^2(a+a \sec(e+fx))^m \sqrt{c-c \sec(e+fx)} \tan(e+fx)}{f(15+16m+4m^2)}$$

$$- \frac{2c(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{3/2} \tan(e+fx)}{f(5+2m)}$$

output

```
-2*c*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3/2)*tan(f*x+e)/f/(5+2*m)-64*c^3
*(a+a*sec(f*x+e))^m*tan(f*x+e)/f/(5+2*m)/(4*m^2+8*m+3)/(c-c*sec(f*x+e))^(1
/2)-16*c^2*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(4*m^2+1
6*m+15)
```

3.156.2 Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.68

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{5/2} dx = \frac{2c^3(a(1 + \sec(e + fx)))^m (43 + 24m + 4m^2 - 2(7 + 16m + 4m^2) \sec(e + fx) + (3 + 8m + 4m^2) \sec^2(e + fx))}{f(1 + 2m)(3 + 2m)(5 + 2m)\sqrt{c - c \sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(5/2),x]`

output `(-2*c^3*(a*(1 + Sec[e + f*x]))^m*(43 + 24*m + 4*m^2 - 2*(7 + 16*m + 4*m^2)*Sec[e + f*x] + (3 + 8*m + 4*m^2)*Sec[e + f*x]^2)*Tan[e + f*x]/(f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*Sqrt[c - c*Sec[e + f*x]])`

3.156.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4443, 3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e + fx)(c - c \sec(e + fx))^{5/2}(a \sec(e + fx) + a)^m dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{5/2} \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m dx \\ & \quad \downarrow \text{4443} \\ & \frac{8c \int \sec(e + fx)(\sec(e + fx)a + a)^m (c - c \sec(e + fx))^{3/2} dx}{2m + 5} - \frac{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}(a \sec(e + fx) + a)^m}{f(2m + 5)} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{8c \int \csc(e + fx + \frac{\pi}{2}) (\csc(e + fx + \frac{\pi}{2}) a + a)^m (c - c \csc(e + fx + \frac{\pi}{2}))^{3/2} dx}{\frac{2m + 5}{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}(a \sec(e + fx) + a)^m} f(2m + 5)}$$

↓ 4443

$$\frac{8c \left(\frac{4c \int \sec(e + fx)(\sec(e + fx)a + a)^m \sqrt{c - c \sec(e + fx)} dx}{2m + 3} - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)} (a \sec(e + fx) + a)^m}{f(2m + 3)} \right)}{\frac{2m + 5}{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}(a \sec(e + fx) + a)^m} f(2m + 5)}$$

↓ 3042

$$\frac{8c \left(\frac{4c \int \csc(e + fx + \frac{\pi}{2}) (\csc(e + fx + \frac{\pi}{2}) a + a)^m \sqrt{c - c \csc(e + fx + \frac{\pi}{2})} dx}{2m + 3} - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)} (a \sec(e + fx) + a)^m}{f(2m + 3)} \right)}{\frac{2m + 5}{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}(a \sec(e + fx) + a)^m} f(2m + 5)}$$

↓ 4441

$$\frac{8c \left(-\frac{8c^2 \tan(e + fx)(a \sec(e + fx) + a)^m}{f(2m + 1)(2m + 3) \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)} (a \sec(e + fx) + a)^m}{f(2m + 3)} \right)}{\frac{2m + 5}{2c \tan(e + fx)(c - c \sec(e + fx))^{3/2}(a \sec(e + fx) + a)^m} f(2m + 5)}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(5/2),x]`

output `(-2*c*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(3/2)*Tan[e + f*x])/(f*(5 + 2*m)) + (8*c*((-8*c^2*(a + a*Sec[e + f*x])^m*Tan[e + f*x])/(f*(1 + 2*m))*(3 + 2*m)*Sqrt[c - c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])^m*Sqrt[c - c*Sec[e + f*x]]*Tan[e + f*x])/(f*(3 + 2*m)))/(5 + 2*m)`

3.156.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4441 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]`

rule 4443 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[(-d)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] + Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c + d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)] && !(IGtQ[m - 1/2, 0] && LtQ[m, n])`

3.156.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (c - c \sec(fx + e))^{\frac{5}{2}} dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(5/2),x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(5/2),x)`

3.156.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.19

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{\frac{5}{2}} dx = \frac{2(4c^2m^2 + (4c^2m^2 + 24c^2m + 43c^2)\cos(fx + e)^3 + 8c^2m - (4c^2m^2 + 8c^2m - 29c^2)\cos(fx + e))^{\frac{5}{2}}}{(8fm^3 + 36fm^2 + 46fm + 15f)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(5/2),x, algorithm="fricas")`

output `2*(4*c^2*m^2 + (4*c^2*m^2 + 24*c^2*m + 43*c^2)*cos(f*x + e)^3 + 8*c^2*m - (4*c^2*m^2 + 8*c^2*m - 29*c^2)*cos(f*x + e)^2 + 3*c^2 - (4*c^2*m^2 + 24*c^2*m + 11*c^2)*cos(f*x + e))*((a*cos(f*x + e) + a)/cos(f*x + e))^m*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*cos(f*x + e)^2*sin(f*x + e))`

3.156.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{5/2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(5/2),x)`

output Timed out

3.156.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.42

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{5/2} dx = \frac{2 \left(\frac{\sqrt{2}(2^{m+5}m+5\cdot 2^{m+4})(-a)^m c^{\frac{5}{2}} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{\sqrt{2}(2^{m+4}m^2+2^{m+6}m+15\cdot 2^{m+2})(-a)^m c^{\frac{5}{2}} \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - 2^{m+\frac{11}{2}}(-a)^m c^{\frac{5}{2}} \right) e^{(-m)}}{(8m^3 + 36m^2 + 46m + 15)f \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right)^{\frac{5}{2}} \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1 \right)^{\frac{5}{2}}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(5/2),x, algorithm="maxima")`

output
$$-2*(\text{sqrt}(2)*(2^{(m+5)}*m + 5*2^{(m+4)})*(-a)^m*c^{(5/2)}*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2 - \text{sqrt}(2)*(2^{(m+4)}*m^2 + 2^{(m+6)}*m + 15*2^{(m+2)})*(-a)^m*c^{(5/2)}*\sin(f*x + e)^4/(\cos(f*x + e) + 1)^4 - 2^{(m+11/2)}*(-a)^m*c^{(5/2)})*e^{(-m*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1) - m*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1))/((8*m^3 + 36*m^2 + 46*m + 15)*f*(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)^{(5/2)}*(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)^{(5/2)})}$$

3.156.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{5/2} dx = \int (-c \sec(fx + e) + c)^{5/2} (a \sec(fx + e) + a)^m \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((-c*sec(f*x + e) + c)^(5/2)*(a*sec(f*x + e) + a)^m*sec(f*x + e), x)`

3.156.9 Mupad [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{5/2} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m \left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}}{\cos(e + fx)} dx$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x),x)`

output `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(5/2))/cos(e + f*x), x)`

3.157 $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{3/2} dx$

3.157.1 Optimal result	1094
3.157.2 Mathematica [A] (verified)	1094
3.157.3 Rubi [A] (verified)	1095
3.157.4 Maple [F]	1096
3.157.5 Fricas [A] (verification not implemented)	1097
3.157.6 Sympy [F(-1)]	1097
3.157.7 Maxima [A] (verification not implemented)	1097
3.157.8 Giac [F]	1098
3.157.9 Mupad [B] (verification not implemented)	1098

3.157.1 Optimal result

Integrand size = 34, antiderivative size = 100

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{3/2} dx =$$

$$\frac{8c^2(a+a \sec(e+fx))^m \tan(e+fx)}{f(3+8m+4m^2)\sqrt{c-c \sec(e+fx)}}$$

$$- \frac{2c(a+a \sec(e+fx))^m \sqrt{c-c \sec(e+fx)} \tan(e+fx)}{f(3+2m)}$$

output

```
-8*c^2*(a+a*sec(f*x+e))^m*tan(f*x+e)/f/(4*m^2+8*m+3)/(c-c*sec(f*x+e))^(1/2)
)-2*c*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1/2)*tan(f*x+e)/f/(3+2*m)
```

3.157.2 Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.72

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{3/2} dx =$$

$$\frac{2c^2(a(1+\sec(e+fx)))^m(-5-2m+(1+2m)\sec(e+fx))\tan(e+fx)}{f(1+2m)(3+2m)\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(3/2),x]`

output `(2*c^2*(a*(1 + Sec[e + f*x]))^m*(-5 - 2*m + (1 + 2*m)*Sec[e + f*x])*Tan[e + f*x])/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sec[e + f*x]])`

3.157.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4443, 3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(c - c \sec(e + fx))^{3/2}(a \sec(e + fx) + a)^m dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{3/2} \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m dx \\
 & \quad \downarrow \text{4443} \\
 & \frac{4c \int \sec(e + fx)(\sec(e + fx)a + a)^m \sqrt{c - c \sec(e + fx)} dx}{2m + 3} - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)} (a \sec(e + fx) + a)^m}{f(2m + 3)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4c \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^m \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} dx}{2m + 3} - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)} (a \sec(e + fx) + a)^m}{f(2m + 3)} \\
 & \quad \downarrow \text{4441} \\
 & \frac{8c^2 \tan(e + fx)(a \sec(e + fx) + a)^m}{f(2m + 1)(2m + 3) \sqrt{c - c \sec(e + fx)}} - \frac{2c \tan(e + fx) \sqrt{c - c \sec(e + fx)} (a \sec(e + fx) + a)^m}{f(2m + 3)}
 \end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(3/2),x]`


```
output (-8*c^2*(a + a*Sec[e + f*x])^m*Tan[e + f*x])/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c
- c*Sec[e + f*x]]) - (2*c*(a + a*Sec[e + f*x])^m*Sqrt[c - c*Sec[e + f*x]]
*Tan[e + f*x])/(f*(3 + 2*m))
```

3.157.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m_*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] := Simp[2*a*c*Cot[e + f
*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] /
; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
&& NeQ[m, -2^(-1)]
```

```
rule 4443 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m*(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^n, x_Symbol] := Simp[(-d)*Cot[e + f
*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^(n - 1)/(f*(m + n))), x] +
Simp[c*((2*n - 1)/(m + n)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m*(c +
d*Csc[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b
*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0] && !LtQ[m, -2^(-1)]
&& !(IGtQ[m - 1/2, 0] && LtQ[m, n])
```

3.157.4 Maple [F]

$$\int \sec(fx + e)(a + a \sec(fx + e))^m (c - c \sec(fx + e))^{\frac{3}{2}} dx$$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3/2),x)
```

```
output int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3/2),x)
```

3.157.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.12

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{3/2} dx = \frac{2 \left((2cm + 5c) \cos(fx + e)^2 - 2cm + 4c \cos(fx + e) - c \right) \left(\frac{a \cos(fx + e) + a}{\cos(fx + e)} \right)^m \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}}}{(4fm^2 + 8fm + 3f) \cos(fx + e) \sin(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3/2),x, algorithm
m="fricas")
```

```
output 2*((2*c*m + 5*c)*cos(f*x + e)^2 - 2*c*m + 4*c*cos(f*x + e) - c)*((a*cos(f*
x + e) + a)/cos(f*x + e))^m*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))/((4*f*
m^2 + 8*f*m + 3*f)*cos(f*x + e)*sin(f*x + e))
```

3.157.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{3/2} dx = \text{Timed out}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3/2),x)
```

```
output Timed out
```

3.157.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.71

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{3/2} dx = \frac{2 \left(\sqrt{2} 2^{m+2} (-a)^m c^{\frac{3}{2}} - \frac{\sqrt{2} (2^{m+2} m + 3 \cdot 2^{m+1}) (-a)^m c^{\frac{3}{2}} \sin(fx + e)^2}{(\cos(fx + e) + 1)^2} \right) e^{\left(-m \log\left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} + 1 \right) - m \log\left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} - 1 \right) \right)}}{(4m^2 + 8m + 3) f \left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} + 1 \right)^{\frac{3}{2}} \left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} - 1 \right)^{\frac{3}{2}}}$$

3.157. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{3/2} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3/2),x, algorithm
m="maxima")`

output `-2*(sqrt(2)*2^(m + 2)*(-a)^m*c^(3/2) - sqrt(2)*(2^(m + 2)*m + 3*2^(m + 1))
*(-a)^m*c^(3/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*e^(-m*log(sin(f*x + e)
)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/((
4*m^2 + 8*m + 3)*f*(sin(f*x + e)/(cos(f*x + e) + 1) + 1)^(3/2)*(sin(f*x +
e)/(cos(f*x + e) + 1) - 1)^(3/2))`

3.157.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{3/2} dx = \int (-c \sec(fx + e) + c)^{3/2} (a \sec(fx + e) + a)^m \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3/2),x, algorithm
m="giac")`

output `integrate((-c*sec(f*x + e) + c)^(3/2)*(a*sec(f*x + e) + a)^m*sec(f*x + e),
x)`

3.157.9 Mupad [B] (verification not implemented)

Time = 15.51 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.54

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{3/2} dx =$$

$$\frac{2c \left(\frac{a(\cos(e+fx)+1)}{\cos(e+fx)} \right)^m \sqrt{\frac{c(\cos(e+fx)-1)}{\cos(e+fx)}} (5 \sin(e + fx) - 2 \sin(2e + 2fx) + 5 \sin(3e + 3fx) + 2m \sin(e + fx))}{f(4m^2 + 8m + 3)(3 \cos(e + fx) - 2 \cos(2e + 2fx) + \cos(3e + 3fx))}$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(3/2))/cos(e + f*x),x)`

output $-(2*c*((a*(\cos(e + f*x) + 1))/\cos(e + f*x))^m*((c*(\cos(e + f*x) - 1))/\cos(e + f*x))^{1/2}*(5*\sin(e + f*x) - 2*\sin(2*e + 2*f*x) + 5*\sin(3*e + 3*f*x) + 2*m*\sin(e + f*x) - 4*m*\sin(2*e + 2*f*x) + 2*m*\sin(3*e + 3*f*x)))/(f*(8*m + 4*m^2 + 3)*(3*\cos(e + f*x) - 2*\cos(2*e + 2*f*x) + \cos(3*e + 3*f*x) - 2))$

3.158 $\int \sec(e+fx)(a+a \sec(e+fx))^m \sqrt{c - c \sec(e + fx)} dx$

3.158.1 Optimal result	1100
3.158.2 Mathematica [C] (verified)	1100
3.158.3 Rubi [A] (verified)	1101
3.158.4 Maple [F]	1102
3.158.5 Fricas [A] (verification not implemented)	1102
3.158.6 Sympy [F]	1103
3.158.7 Maxima [B] (verification not implemented)	1103
3.158.8 Giac [F]	1104
3.158.9 Mupad [F(-1)]	1104

3.158.1 Optimal result

Integrand size = 34, antiderivative size = 46

$$\int \sec(e + fx)(a + a \sec(e + fx))^m \sqrt{c - c \sec(e + fx)} dx$$

$$= -\frac{2c(a + a \sec(e + fx))^m \tan(e + fx)}{f(1 + 2m)\sqrt{c - c \sec(e + fx)}}$$

output `-2*c*(a+a*sec(f*x+e))^m*tan(f*x+e)/f/(1+2*m)/(c-c*sec(f*x+e))^(1/2)`

3.158.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.70 (sec) , antiderivative size = 163, normalized size of antiderivative = 3.54

$$\int \sec(e + fx)(a + a \sec(e + fx))^m \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{\sqrt{2}e^{-\frac{1}{2}i(e+fx)}(1 + e^{i(e+fx)}) \sqrt{\frac{e^{i(e+fx)}}{1+e^{2i(e+fx)}}} \left(\frac{(1+e^{i(e+fx)})^2}{1+e^{2i(e+fx)}}\right)^m \csc\left(\frac{1}{2}(e + fx)\right) (1 + \sec(e + fx))^{-m} (a(1 + \sec(e + fx)))^m}{(f + 2fm)\sqrt{\sec(e + fx)}}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*Sqrt[c - c*Sec[e + f*x]],x]`

output $(\text{Sqrt}[2]*(1 + E^{(I*(e + f*x))})*\text{Sqrt}[E^{(I*(e + f*x))}/(1 + E^{((2*I)*(e + f*x))})])*((1 + E^{(I*(e + f*x))})^2/(1 + E^{((2*I)*(e + f*x))}))^m*\text{Csc}[(e + f*x)/2]*(\text{a}*(1 + \text{Sec}[e + f*x]))^m*\text{Sqrt}[c - c*\text{Sec}[e + f*x]]/(E^{(I/2)*(e + f*x)}*(f + 2*f*m)*\text{Sqrt}[\text{Sec}[e + f*x]]*(1 + \text{Sec}[e + f*x])^m)$

3.158.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {3042, 4441}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx) \sqrt{c - c \sec(e + fx)} (a \sec(e + fx) + a)^m dx$$

↓ 3042

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \sqrt{c - c \csc\left(e + fx + \frac{\pi}{2}\right)} \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m dx$$

↓ 4441

$$\frac{2c \tan(e + fx) (a \sec(e + fx) + a)^m}{f(2m + 1) \sqrt{c - c \sec(e + fx)}}$$

input $\text{Int}[\text{Sec}[e + f*x]*(a + a*\text{Sec}[e + f*x])^m*\text{Sqrt}[c - c*\text{Sec}[e + f*x]],x]$

output $(-2*c*(a + a*\text{Sec}[e + f*x])^m*\text{Tan}[e + f*x])/(f*(1 + 2*m)*\text{Sqrt}[c - c*\text{Sec}[e + f*x]])$

3.158.3.1 Defintions of rubi rules used

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

```
rule 4441 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)], x_Symbol] :> Simp[2*a*c*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1)*Sqrt[c + d*Csc[e + f*x]])), x] / ; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[m, -2^(-1)]
```

3.158.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m \sqrt{c - c \sec(fx + e)} dx$$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1/2),x)
```

```
output int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1/2),x)
```

3.158.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.52

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m \sqrt{c - c \sec(e + fx)} dx \\ &= \frac{2 \left(\frac{a \cos(fx + e) + a}{\cos(fx + e)} \right)^m \sqrt{\frac{c \cos(fx + e) - c}{\cos(fx + e)}} (\cos(fx + e) + 1)}{(2fm + f) \sin(fx + e)} \end{aligned}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1/2),x, algorithm m="fricas")
```

```
output 2*((a*cos(f*x + e) + a)/cos(f*x + e))^m*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*(cos(f*x + e) + 1)/((2*f*m + f)*sin(f*x + e))
```

3.158.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m \sqrt{c - c \sec(e + fx)} dx$$

$$= \int (a(\sec(e + fx) + 1))^m \sqrt{-c(\sec(e + fx) - 1)} \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**m*(c-c*sec(f*x+e))**(1/2),x)`

output `Integral((a*(sec(e + f*x) + 1))**m*sqrt(-c*(sec(e + f*x) - 1))*sec(e + f*x), x)`

3.158.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(44) = 88$.

Time = 0.33 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.48

$$\int \sec(e + fx)(a + a \sec(e + fx))^m \sqrt{c - c \sec(e + fx)} dx$$

$$= \frac{2^{m+\frac{3}{2}}(-a)^m \sqrt{c} e^{(-m \log(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1) - m \log(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1))}}{f(2m+1) \sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1} \sqrt{\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1/2),x, algorithm m="maxima")`

output `2^(m + 3/2)*(-a)^m*sqrt(c)*e^(-m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1))/(f*(2*m + 1)*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) + 1)*sqrt(sin(f*x + e)/(cos(f*x + e) + 1) - 1))`

3.158.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m \sqrt{c - c \sec(e + fx)} dx$$

$$= \int \sqrt{-c \sec(fx + e) + c} (a \sec(fx + e) + a)^m \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1/2),x, algorithm m="giac")`

output `integrate(sqrt(-c*sec(f*x + e) + c)*(a*sec(f*x + e) + a)^m*sec(f*x + e), x)`

3.158.9 Mupad [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^m \sqrt{c - c \sec(e + fx)} dx$$

$$= \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m \sqrt{c - \frac{c}{\cos(e+fx)}}}{\cos(e + fx)} dx$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x),x)`

output `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(1/2))/cos(e + f*x), x)`

3.159 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{\sqrt{c-c \sec(e+fx)}} dx$

3.159.1 Optimal result 1105
 3.159.2 Mathematica [A] (verified) 1105
 3.159.3 Rubi [A] (verified) 1106
 3.159.4 Maple [F] 1107
 3.159.5 Fracas [F] 1107
 3.159.6 Sympy [F] 1108
 3.159.7 Maxima [F] 1108
 3.159.8 Giac [F] 1108
 3.159.9 Mupad [F(-1)] 1109

3.159.1 Optimal result

Integrand size = 34, antiderivative size = 69

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{\sqrt{c-c \sec(e+fx)}} dx =$$

$$\frac{\text{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e+fx))\right) (a+a \sec(e+fx))^m \tan(e+fx)}{f(1+2m)\sqrt{c-c \sec(e+fx)}}$$

output `-hypergeom([1, 1/2+m], [3/2+m], 1/2+1/2*sec(f*x+e))*(a+a*sec(f*x+e))^m*tan(f*x+e)/f/(1+2*m)/(c-c*sec(f*x+e))^(1/2)`

3.159.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.97

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{\sqrt{c-c \sec(e+fx)}} dx =$$

$$\frac{\text{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e+fx))\right) (a(1 + \sec(e+fx)))^m \tan(e+fx)}{(f+2fm)\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/Sqrt[c - c*Sec[e + f*x]],x]`

output `-((Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sec[e + f*x])/2]*(a*(1 + Sec[e + f*x]))^m*Tan[e + f*x])/((f + 2*f*m)*Sqrt[c - c*Sec[e + f*x]]))`

3.159. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{\sqrt{c-c \sec(e+fx)}} dx$

3.159.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4449, 27, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^m}{\sqrt{c-c\sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^m}{\sqrt{c-c\csc(e+fx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4449} \\
 & -\frac{a \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{m-\frac{1}{2}}}{c(1-\sec(e+fx))} d\sec(e+fx)}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{27} \\
 & -\frac{a \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{m-\frac{1}{2}}}{1-\sec(e+fx)} d\sec(e+fx)}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{78} \\
 & -\frac{\tan(e+fx)(a\sec(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}(\sec(e+fx)+1)\right)}{f(2m+1)\sqrt{c-c\sec(e+fx)}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/Sqrt[c - c*Sec[e + f*x]],x]`

output `-((Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sec[e + f*x])/2]*(a + a*Sec[e + f*x])^m*Tan[e + f*x])/(f*(1 + 2*m)*Sqrt[c - c*Sec[e + f*x]]))`

3.159.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 78 `Int[((a_) + (b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4449 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.159.4 Maple [F]

$$\int \frac{\sec(fx + e)(a + a \sec(fx + e))^m}{\sqrt{c - c \sec(fx + e)}} dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(1/2),x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(1/2),x)`

3.159.5 Fracas [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{\sqrt{c - c \sec(e + fx)}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{\sqrt{-c \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(1/2),x, algorithm m="fracas")`

output `integral(-sqrt(-c*sec(f*x + e) + c)*(a*sec(f*x + e) + a)^m*sec(f*x + e)/(c*sec(f*x + e) - c), x)`

3.159.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{\sqrt{c - c \sec(e + fx)}} dx = \int \frac{(a(\sec(e + fx) + 1))^m \sec(e + fx)}{\sqrt{-c(\sec(e + fx) - 1)}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(1/2),x)`

output `Integral((a*(sec(e + f*x) + 1))^m*sec(e + f*x)/sqrt(-c*(sec(e + f*x) - 1)), x)`

3.159.7 Maxima [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{\sqrt{c - c \sec(e + fx)}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{\sqrt{-c \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(1/2),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/sqrt(-c*sec(f*x + e) + c), x)`

3.159.8 Giac [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{\sqrt{c - c \sec(e + fx)}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{\sqrt{-c \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(1/2),x, algorithm m="giac")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/sqrt(-c*sec(f*x + e) + c), x)`

3.159.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{\sqrt{c-c\sec(e+fx)}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m}{\cos(e+fx) \sqrt{c - \frac{c}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(1/2)), x)`

3.160
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{3/2}} dx$$

3.160.1 Optimal result 1110
 3.160.2 Mathematica [A] (verified) 1110
 3.160.3 Rubi [A] (verified) 1111
 3.160.4 Maple [F] 1112
 3.160.5 Fracas [F] 1112
 3.160.6 Sympy [F] 1113
 3.160.7 Maxima [F] 1113
 3.160.8 Giac [F] 1113
 3.160.9 Mupad [F(-1)] 1114

3.160.1 Optimal result

Integrand size = 34, antiderivative size = 74

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{3/2}} dx = \frac{\text{Hypergeometric2F1}\left(2, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e+fx))\right) (a+a \sec(e+fx))^m \tan(e+fx)}{2cf(1+2m)\sqrt{c-c \sec(e+fx)}}$$

output `-1/2*hypergeom([2, 1/2+m],[3/2+m],1/2+1/2*sec(f*x+e))*(a+a*sec(f*x+e))^m*tan(f*x+e)/c/f/(1+2*m)/(c-c*sec(f*x+e))^(1/2)`

3.160.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{3/2}} dx = \frac{\text{Hypergeometric2F1}\left(2, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e+fx))\right) (a(1 + \sec(e+fx)))^m \tan(e+fx)}{4cf\left(\frac{1}{2} + m\right)\sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^(3/2),x]`

output `-1/4*(Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sec[e + f*x])/2]*(a*(1 + Sec[e + f*x]))^m*Tan[e + f*x])/(c*f*(1/2 + m)*Sqrt[c - c*Sec[e + f*x]])`

3.160.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{3/2}} dx$$

3.160.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4449, 27, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^m}{(c-c\sec(e+fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^m}{(c-c\csc(e+fx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4449} \\
 & \frac{a \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{m-\frac{1}{2}}}{c^2(1-\sec(e+fx))^2} d\sec(e+fx)}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{a \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{m-\frac{1}{2}}}{(1-\sec(e+fx))^2} d\sec(e+fx)}{cf\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{78} \\
 & \frac{\tan(e+fx)(a\sec(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(2, m+\frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}(\sec(e+fx)+1)\right)}{2cf(2m+1)\sqrt{c-c\sec(e+fx)}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^(3/2),x]`

output `-1/2*(Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sec[e + f*x])/2]*(a + a*Sec[e + f*x])^m*Tan[e + f*x])/(c*f*(1 + 2*m)*Sqrt[c - c*Sec[e + f*x]])`

3.160.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4449 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.160.4 Maple [F]

$$\int \frac{\sec(fx + e)(a + a \sec(fx + e))^m}{(c - c \sec(fx + e))^{\frac{3}{2}}} dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(3/2),x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(3/2),x)`

3.160.5 Fracas [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{\frac{3}{2}}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(3/2),x, algorithm m="fracas")`

output `integral(sqrt(-c*sec(f*x + e) + c)*(a*sec(f*x + e) + a)^m*sec(f*x + e)/(c^2*sec(f*x + e)^2 - 2*c^2*sec(f*x + e) + c^2), x)`

3.160.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{3/2}} dx = \int \frac{(a(\sec(e + fx) + 1))^m \sec(e + fx)}{(-c(\sec(e + fx) - 1))^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(3/2),x)`

output `Integral((a*(sec(e + f*x) + 1))^m*sec(e + f*x)/(-c*(sec(e + f*x) - 1))^(3/2), x)`

3.160.7 Maxima [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{3/2}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(3/2),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(-c*sec(f*x + e) + c)^(3/2), x)`

3.160.8 Giac [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{3/2}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^{\frac{3}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(3/2),x, algorithm m="giac")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(-c*sec(f*x + e) + c)^(3/2), x)`

3.160.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{(c-c\sec(e+fx))^{3/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)),x)`output `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(3/2)), x)`

3.161
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{5/2}} dx$$

3.161.1 Optimal result 1115
 3.161.2 Mathematica [A] (verified) 1115
 3.161.3 Rubi [A] (verified) 1116
 3.161.4 Maple [F] 1117
 3.161.5 Fracas [F] 1117
 3.161.6 Sympy [F(-1)] 1118
 3.161.7 Maxima [F] 1118
 3.161.8 Giac [F(-2)] 1118
 3.161.9 Mupad [F(-1)] 1119

3.161.1 Optimal result

Integrand size = 34, antiderivative size = 74

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{5/2}} dx = \frac{\text{Hypergeometric2F1}\left(3, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e+fx))\right) (a+a \sec(e+fx))^m \tan(e+fx)}{4c^2 f(1+2m) \sqrt{c-c \sec(e+fx)}}$$

output `-1/4*hypergeom([3, 1/2+m], [3/2+m], 1/2+1/2*sec(f*x+e))*(a+a*sec(f*x+e))^m*tan(f*x+e)/c^2/f/(1+2*m)/(c-c*sec(f*x+e))^(1/2)`

3.161.2 Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{5/2}} dx = \frac{\text{Hypergeometric2F1}\left(3, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e+fx))\right) (a(1 + \sec(e+fx)))^m \tan(e+fx)}{8c^2 f\left(\frac{1}{2} + m\right) \sqrt{c-c \sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^(5/2), x]`

output `-1/8*(Hypergeometric2F1[3, 1/2 + m, 3/2 + m, (1 + Sec[e + f*x])/2]*(a*(1 + Sec[e + f*x]))^m*Tan[e + f*x])/(c^2*f*(1/2 + m)*Sqrt[c - c*Sec[e + f*x]])`

3.161.
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^m}{(c-c \sec(e+fx))^{5/2}} dx$$

3.161.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4449, 27, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^m}{(c-c\sec(e+fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^m}{(c-c\csc(e+fx+\frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4449} \\
 & \frac{a \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{m-\frac{1}{2}}}{c^3(1-\sec(e+fx))^3} d\sec(e+fx)}{f\sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{a \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{m-\frac{1}{2}}}{(1-\sec(e+fx))^3} d\sec(e+fx)}{c^2 f \sqrt{a\sec(e+fx)+a}\sqrt{c-c\sec(e+fx)}} \\
 & \quad \downarrow \text{78} \\
 & \frac{\tan(e+fx)(a\sec(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(3, m+\frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}(\sec(e+fx)+1)\right)}{4c^2 f(2m+1)\sqrt{c-c\sec(e+fx)}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^(5/2),x]`

output `-1/4*(Hypergeometric2F1[3, 1/2 + m, 3/2 + m, (1 + Sec[e + f*x])/2]*(a + a*Sec[e + f*x])^m*Tan[e + f*x])/(c^2*f*(1 + 2*m)*Sqrt[c - c*Sec[e + f*x]])`

3.161.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4449 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.161.4 Maple [F]

$$\int \frac{\sec(fx + e)(a + a \sec(fx + e))^m}{(c - c \sec(fx + e))^{\frac{5}{2}}} dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(5/2),x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(5/2),x)`

3.161.5 Fracas [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{\frac{5}{2}}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^{\frac{5}{2}}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(5/2),x, algorithm m="fracas")`

output `integral(-sqrt(-c*sec(f*x + e) + c)*(a*sec(f*x + e) + a)^m*sec(f*x + e)/(c^3*sec(f*x + e)^3 - 3*c^3*sec(f*x + e)^2 + 3*c^3*sec(f*x + e) - c^3), x)`

3.161.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(5/2),x)`

output Timed out

3.161.7 Maxima [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{5/2}} dx = \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^{5/2}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(5/2),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(-c*sec(f*x + e) + c)^(5/2), x)`

3.161.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^m}{(c - c \sec(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/(c-c*sec(f*x+e))^(5/2),x, algorithm m="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [0,4,1,0]%%}+%%{2, [0,2,1,1]%%}+%%{1, [0,0,1,2]%%} / %%{1, [0,

3.161.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^m}{(c-c\sec(e+fx))^{5/2}} dx = \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m}{\cos(e+fx) \left(c - \frac{c}{\cos(e+fx)}\right)^{5/2}} dx$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)),x)`

output `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(5/2)), x)`

3.162 $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx$

3.162.1 Optimal result	1120
3.162.2 Mathematica [A] (verified)	1121
3.162.3 Rubi [A] (verified)	1121
3.162.4 Maple [F]	1123
3.162.5 Fricas [A] (verification not implemented)	1123
3.162.6 Sympy [F]	1124
3.162.7 Maxima [A] (verification not implemented)	1124
3.162.8 Giac [F]	1125
3.162.9 Mupad [B] (verification not implemented)	1125

3.162.1 Optimal result

Integrand size = 36, antiderivative size = 169

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx \\ &= -\frac{(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} \tan(e + fx)}{f(1 + 2m)} \\ & \quad + \frac{2(a + a \sec(e + fx))^{1+m} (c - c \sec(e + fx))^{-3-m} \tan(e + fx)}{af(3 + 8m + 4m^2)} \\ & \quad - \frac{2(a + a \sec(e + fx))^{2+m} (c - c \sec(e + fx))^{-3-m} \tan(e + fx)}{a^2 f(1 + 2m)(15 + 16m + 4m^2)} \end{aligned}$$

```
output -(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3+m)*tan(f*x+e)/f/(1+2*m)+2*(a+a*sec(f*x+e))^(1+m)*(c-c*sec(f*x+e))^(3+m)*tan(f*x+e)/a/f/(4*m^2+8*m+3)-2*(a+a*sec(f*x+e))^(2+m)*(c-c*sec(f*x+e))^(3+m)*tan(f*x+e)/a^2/f/(1+2*m)/(4*m^2+16*m+15)
```

3.162.2 Mathematica [A] (verified)

Time = 2.47 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.62

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx$$

$$= \frac{(a(1 + \sec(e + fx)))^m (c - c \sec(e + fx))^{-m} (7 + 12m + 4m^2 - 2(3 + 2m) \sec(e + fx) + 2 \sec^2(e + fx))}{c^3 f (1 + 2m)(3 + 2m)(5 + 2m)(-1 + \sec(e + fx))^3}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(-3 - m),x]`

output `((a*(1 + Sec[e + f*x]))^m*(7 + 12*m + 4*m^2 - 2*(3 + 2*m)*Sec[e + f*x] + 2*Sec[e + f*x]^2)*Tan[e + f*x])/(c^3*f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(-1 + Sec[e + f*x])^3*(c - c*Sec[e + f*x])^m)`

3.162.3 Rubi [A] (verified)Time = 0.82 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4439, 3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-3} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{-m-3} dx$$

$$\downarrow \text{4439}$$

$$-\frac{2 \int \sec(e + fx)(\sec(e + fx)a + a)^{m+1} (c - c \sec(e + fx))^{-m-3} dx}{a(2m + 1)}$$

$$\frac{\tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-3}}{f(2m + 1)}$$

$$\downarrow \text{3042}$$

3.162. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx$

$$\begin{aligned}
& \frac{2 \int \csc(e + fx + \frac{\pi}{2}) (\csc(e + fx + \frac{\pi}{2}) a + a)^{m+1} (c - c \csc(e + fx + \frac{\pi}{2}))^{-m-3} dx}{a(2m+1)} \\
& \frac{\tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-3}}{f(2m+1)} \\
& \quad \downarrow \text{4439} \\
& \frac{2 \left(-\frac{\int \sec(e+fx)(\sec(e+fx)a+a)^{m+2}(c-c\sec(e+fx))^{-m-3} dx}{a(2m+3)} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^{m+1}(c-c\sec(e+fx))^{-m-3}}{f(2m+3)} \right)}{a(2m+1)} \\
& \frac{\tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-3}}{f(2m+1)} \\
& \quad \downarrow \text{3042} \\
& \frac{2 \left(-\frac{\int \csc(e+fx+\frac{\pi}{2})(\csc(e+fx+\frac{\pi}{2})a+a)^{m+2}(c-c\csc(e+fx+\frac{\pi}{2}))^{-m-3} dx}{a(2m+3)} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^{m+1}(c-c\sec(e+fx))^{-m-3}}{f(2m+3)} \right)}{a(2m+1)} \\
& \frac{\tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-3}}{f(2m+1)} \\
& \quad \downarrow \text{4438} \\
& \frac{\tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-3}}{f(2m+1)} \\
& \frac{2 \left(\frac{\tan(e+fx)(a \sec(e+fx)+a)^{m+2}(c-c\sec(e+fx))^{-m-3}}{af(2m+3)(2m+5)} - \frac{\tan(e+fx)(a \sec(e+fx)+a)^{m+1}(c-c\sec(e+fx))^{-m-3}}{f(2m+3)} \right)}{a(2m+1)}
\end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(-3 - m),x]`

output `-(((a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(-3 - m)*Tan[e + f*x])/(f*(1 + 2*m))) - (2*(-(((a + a*Sec[e + f*x])^(1 + m)*(c - c*Sec[e + f*x])^(-3 - m)*Tan[e + f*x])/(f*(3 + 2*m)))) + ((a + a*Sec[e + f*x])^(2 + m)*(c - c*Sec[e + f*x])^(-3 - m)*Tan[e + f*x])/(a*f*(3 + 2*m)*(5 + 2*m))))/(a*(1 + 2*m))`

3.162.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

rule 4439 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])`

3.162.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (c - c \sec(fx + e))^{-3-m} dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3-m),x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3-m),x)`

3.162.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.71

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx = \frac{((4m^2 + 12m + 7) \cos(fx + e)^2 - 2(2m + 3) \cos(fx + e) + 2) \left(\frac{a \cos(fx + e) + a}{\cos(fx + e)}\right)^m \left(\frac{c \cos(fx + e) - c}{\cos(fx + e)}\right)^{-m-3}}{(8fm^3 + 36fm^2 + 46fm + 15f) \cos(fx + e)^3}$$

3.162. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3-m),x, algorithm="fricas")`

output
$$-\left((4m^2 + 12m + 7)\cos(fx + e)^2 - 2(2m + 3)\cos(fx + e) + 2\right)\left(\frac{a\cos(fx + e) + a}{\cos(fx + e)}\right)^m \left(\frac{c\cos(fx + e) - c}{\cos(fx + e)}\right)^{-m-3} \sin(fx + e) / \left((8f^3m^3 + 36f^2m^2 + 46f^2m + 15f)\cos(fx + e)^3\right)$$

3.162.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx \\ &= \int (a(\sec(e + fx) + 1))^m (-c(\sec(e + fx) - 1))^{-m-3} \sec(e + fx) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3-m),x)`

output `Integral((a*(sec(e + f*x) + 1))^m*(-c*(sec(e + f*x) - 1))^(3-m)*sec(e + f*x), x)`

3.162.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.92

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx \\ &= \frac{\left((4m^2 + 8m + 3)(-a)^m - \frac{2(4m^2 + 12m + 5)(-a)^m \sin^2(fx + e)}{(\cos(fx + e) + 1)^2} + \frac{(4m^2 + 16m + 15)(-a)^m \sin^4(fx + e)}{(\cos(fx + e) + 1)^4} \right) c^{-m-3} (\cos(fx + e) + 1)^{-2m}}{4(8m^3 + 36m^2 + 46m + 15)f \left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} \right)^{2m} \sin(fx + e)^5} \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3-m),x, algorithm="maxima")`

output
$$\frac{1}{4} \left((4m^2 + 8m + 3)(-a)^m - \frac{2(4m^2 + 12m + 5)(-a)^m \sin^2(fx + e)}{(\cos(fx + e) + 1)^2} + \frac{(4m^2 + 16m + 15)(-a)^m \sin^4(fx + e)}{(\cos(fx + e) + 1)^4} \right) c^{-m-3} (\cos(fx + e) + 1)^5 / \left((8m^3 + 36m^2 + 46m + 15) f \left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} \right)^{2m} \sin(fx + e)^5 \right)$$

3.162. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx$

3.162.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m-3} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(3-m),x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m - 3)*sec(f*x + e), x)`

3.162.9 Mupad [B] (verification not implemented)

Time = 22.28 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.72

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx =$$

$$\frac{(\cos(3e + 3fx) - \sin(3e + 3fx) i) \left(\frac{\sin(e+fx) \left(a + \frac{a}{\cos(e+fx)} \right)^m (\cos(3e+3fx) + \sin(3e+3fx) i) (4m^2 + 12m + 15) 2i}{f(m^3 8i + m^2 36i + m 46i + 15i)} \right)}{8 \cos(e + fx)}$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(m + 3)),x)`

output `-((cos(3*e + 3*f*x) - sin(3*e + 3*f*x)*1i)*((sin(e + f*x)*(a + a/cos(e + f*x))^m*(cos(3*e + 3*f*x) + sin(3*e + 3*f*x)*1i)*(12*m + 4*m^2 + 15)*2i)/(f*(m*46i + m^2*36i + m^3*8i + 15i)) - (sin(2*e + 2*f*x)*(8*m + 12)*(a + a/cos(e + f*x))^m*(cos(3*e + 3*f*x) + sin(3*e + 3*f*x)*1i)*2i)/(f*(m*46i + m^2*36i + m^3*8i + 15i)) + (sin(3*e + 3*f*x)*(a + a/cos(e + f*x))^m*(cos(3*e + 3*f*x) + sin(3*e + 3*f*x)*1i)*(12*m + 4*m^2 + 7)*2i)/(f*(m*46i + m^2*36i + m^3*8i + 15i))))/(8*cos(e + f*x)^3*(c - c/cos(e + f*x))^(m + 3))`

3.162. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-3-m} dx$

3.163 $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-2-m} dx$

3.163.1 Optimal result	1126
3.163.2 Mathematica [A] (verified)	1126
3.163.3 Rubi [A] (verified)	1127
3.163.4 Maple [A] (verified)	1128
3.163.5 Fricas [A] (verification not implemented)	1129
3.163.6 Sympy [F]	1129
3.163.7 Maxima [A] (verification not implemented)	1130
3.163.8 Giac [F]	1130
3.163.9 Mupad [B] (verification not implemented)	1131

3.163.1 Optimal result

Integrand size = 36, antiderivative size = 104

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-2-m} dx$$

$$= -\frac{(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-2-m} \tan(e+fx)}{f(1+2m)} + \frac{(a+a \sec(e+fx))^{1+m}(c-c \sec(e+fx))^{-2-m} \tan(e+fx)}{af(3+8m+4m^2)}$$

```
output -(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2+m)*tan(f*x+e)/f/(1+2*m)+(a+a*sec(f*x+e))^(1+m)*(c-c*sec(f*x+e))^(2+m)*tan(f*x+e)/a/f/(4*m^2+8*m+3)
```

3.163.2 Mathematica [A] (verified)

Time = 1.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.73

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-2-m} dx$$

$$= \frac{(a(1+\sec(e+fx)))^m(-2(1+m)+\sec(e+fx))(c-c \sec(e+fx))^{-m} \tan(e+fx)}{c^2 f(1+2m)(3+2m)(-1+\sec(e+fx))^2}$$

```
input Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(2 + m),x]
```

output $((a*(1 + \text{Sec}[e + f*x]))^m*(-2*(1 + m) + \text{Sec}[e + f*x])* \text{Tan}[e + f*x]) / (c^2*f*(1 + 2*m)*(3 + 2*m)*(-1 + \text{Sec}[e + f*x])^2*(c - c*\text{Sec}[e + f*x])^m)$

3.163.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4439, 3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-2} dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{-m-2} dx \\ & \quad \downarrow \text{4439} \\ & \frac{\int \sec(e + fx)(\sec(e + fx)a + a)^{m+1} (c - c \sec(e + fx))^{-m-2} dx}{\frac{a(2m+1) \tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-2}}{f(2m+1)}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^{m+1} \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{-m-2} dx}{\frac{a(2m+1) \tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-2}}{f(2m+1)}} \\ & \quad \downarrow \text{4438} \\ & \frac{\tan(e + fx)(a \sec(e + fx) + a)^{m+1} (c - c \sec(e + fx))^{-m-2}}{af(2m+1)(2m+3)} - \frac{\tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-2}}{f(2m+1)} \end{aligned}$$

input $\text{Int}[\text{Sec}[e + f*x]*(a + a*\text{Sec}[e + f*x])^m*(c - c*\text{Sec}[e + f*x])^{-(2 + m)}, x]$

3.163. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-2-m} dx$


```
output -(((a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(-2 - m)*Tan[e + f*x])/(f*(1 + 2*m))) + ((a + a*Sec[e + f*x])^(1 + m)*(c - c*Sec[e + f*x])^(-2 - m)*Tan[e + f*x])/(a*f*(1 + 2*m)*(3 + 2*m))
```

3.163.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4438 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]
```

```
rule 4439 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(m + n + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*(c + d*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && ILtQ[m + n + 1, 0] && NeQ[2*m + 1, 0] && !LtQ[n, 0] && !(IGtQ[n + 1/2, 0] && LtQ[n + 1/2, -(m + n)])
```

3.163.4 Maple [A] (verified)

Time = 5.44 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.93

method	result	size
parallelrisch	$-\frac{\left(-\frac{2a}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1}\right)^m \cot\left(\frac{fx}{2} + \frac{e}{2}\right) \left(-\frac{1}{2} + \cos(fx+e)(m+1)\right) \csc\left(\frac{fx}{2} + \frac{e}{2}\right)^2 \left(\frac{c(\cos(fx+e)-1)}{\cos(fx+e)}\right)^{-m}}{f(3+2m)(1+2m)c^2}$	97

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(-2-m), x, method=_RETURN VERBOSE)
```

output
$$-\left(-\frac{2}{\tan\left(\frac{1}{2}f*x+\frac{1}{2}e\right)^2-1}\right)*a^m*\cot\left(\frac{1}{2}f*x+\frac{1}{2}e\right)*\left(-\frac{1}{2}+\cos(f*x+e)\right)*(m+1)*\csc\left(\frac{1}{2}f*x+\frac{1}{2}e\right)^2*\left(\frac{c*\cos(f*x+e)-1}{\cos(f*x+e)}\right)^{-m}/f/(3+2*m)/(1+2*m)/c^2$$

3.163.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.89

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-2-m} dx$$

$$= -\frac{(2(m+1)\cos(fx+e)-1)\left(\frac{a\cos(fx+e)+a}{\cos(fx+e)}\right)^m \left(\frac{c\cos(fx+e)-c}{\cos(fx+e)}\right)^{-m-2} \sin(fx+e)}{(4fm^2 + 8fm + 3f)\cos(fx+e)^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2-m),x, algorithm="fricas")`

output
$$-(2*(m+1)*\cos(f*x+e)-1)*\left(\frac{a*\cos(f*x+e)+a}{\cos(f*x+e)}\right)^m*\left(\frac{c*\cos(f*x+e)-c}{\cos(f*x+e)}\right)^{-m-2}*\sin(f*x+e)/((4*f*m^2+8*f*m+3*f)*\cos(f*x+e)^2)$$

3.163.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-2-m} dx$$

$$= \int (a(\sec(e + fx) + 1))^m (-c(\sec(e + fx) - 1))^{-m-2} \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2-m),x)`

output `Integral((a*(sec(e + f*x) + 1))^m*(-c*(sec(e + f*x) - 1))^(2-m)*sec(e + f*x), x)`

3.163.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.03

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-2-m} dx$$

$$= -\frac{\left((-a)^m (2m + 1) - \frac{(-a)^m (2m + 3) \sin(fx + e)^2}{(\cos(fx + e) + 1)^2}\right) c^{-m-2} (\cos(fx + e) + 1)^3}{2(4m^2 + 8m + 3)f \left(\frac{\sin(fx + e)}{\cos(fx + e) + 1}\right)^{2m} \sin(fx + e)^3}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^-2-m),x, algorithm="maxima")`

output `-1/2*((-a)^(2*m + 1) - (-a)^(2*m + 3)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*c^(-m - 2)*(cos(f*x + e) + 1)^3/((4*m^2 + 8*m + 3)*f*(sin(f*x + e)/(cos(f*x + e) + 1))^(2*m)*sin(f*x + e)^3)`

3.163.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-2-m} dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m-2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^-2-m),x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m - 2)*sec(f*x + e), x)`

3.163.9 Mupad [B] (verification not implemented)

Time = 19.41 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.39

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-2-m} dx$$

$$= \frac{\sin(e + fx) \left(a + \frac{a}{\cos(e+fx)}\right)^m \operatorname{li}}{f \cos(e + fx)^2 \left(c - \frac{c}{\cos(e+fx)}\right)^{m+2} (m^2 4i + m 8i + 3i)}$$

$$- \frac{\sin(2e + 2fx) (2m + 2) \left(a + \frac{a}{\cos(e+fx)}\right)^m \operatorname{li}}{2 f \cos(e + fx)^2 \left(c - \frac{c}{\cos(e+fx)}\right)^{m+2} (m^2 4i + m 8i + 3i)}$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(m + 2)),x)`output `(sin(e + f*x)*(a + a/cos(e + f*x))^m*li)/(f*cos(e + f*x)^2*(c - c/cos(e + f*x))^(m + 2)*(m*8i + m^2*4i + 3i)) - (sin(2*e + 2*f*x)*(2*m + 2)*(a + a/cos(e + f*x))^m*li)/(2*f*cos(e + f*x)^2*(c - c/cos(e + f*x))^(m + 2)*(m*8i + m^2*4i + 3i))`

3.164 $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-1-m} dx$

3.164.1 Optimal result	1132
3.164.2 Mathematica [A] (verified)	1132
3.164.3 Rubi [A] (verified)	1133
3.164.4 Maple [A] (verified)	1134
3.164.5 Fricas [A] (verification not implemented)	1134
3.164.6 Sympy [F]	1135
3.164.7 Maxima [A] (verification not implemented)	1135
3.164.8 Giac [F]	1135
3.164.9 Mupad [B] (verification not implemented)	1136

3.164.1 Optimal result

Integrand size = 36, antiderivative size = 47

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-1-m} dx$$

$$= -\frac{(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-1-m} \tan(e+fx)}{f(1+2m)}$$

output `-(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^{-1-m}*tan(f*x+e)/f/(1+2*m)`

3.164.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.04

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-1-m} dx$$

$$= -\frac{(a(1+\sec(e+fx)))^m(c-c \sec(e+fx))^{-1-m} \tan(e+fx)}{2f(\frac{1}{2}+m)}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^{-1 - m}, x]`

output `-1/2*((a*(1 + Sec[e + f*x]))^m*(c - c*Sec[e + f*x])^{-1 - m}*Tan[e + f*x])/(f*(1/2 + m))`

3.164. $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-1-m} dx$

3.164.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {3042, 4438}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-1} dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{-m-1} dx$$

$$\downarrow \text{4438}$$

$$\frac{\tan(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m-1}}{f(2m + 1)}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(-1 - m),x]`

output `-(((a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(-1 - m)*Tan[e + f*x])/(f*(1 + 2*m)))`

3.164.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4438 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^n, x_Symbol] := Simp[b*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((c + d*Csc[e + f*x])^n/(a*f*(2*m + 1))), x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && EqQ[m + n + 1, 0] && NeQ[2*m + 1, 0]`

3.164.4 Maple [A] (verified)

Time = 5.24 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.62

method	result	size
parallelrisc	$\frac{\left(\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 c}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 - 1}\right)^{-m} \left(-\frac{a}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 - 1}\right)^m \cot\left(\frac{fx}{2}+\frac{e}{2}\right)}{f(1+2m)c}$	76

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x,method=_RETURN
VERBOSE)
```

```
output 1/f/(1+2*m)/c*(1/(tan(1/2*f*x+1/2*e)^2-1)*tan(1/2*f*x+1/2*e)^2*c)^(-m)*(-1
/(tan(1/2*f*x+1/2*e)^2-1)*a)^m*cot(1/2*f*x+1/2*e)
```

3.164.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.53

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-1-m} dx$$

$$= -\frac{\left(\frac{a \cos(fx+e)+a}{\cos(fx+e)}\right)^m \left(\frac{c \cos(fx+e)-c}{\cos(fx+e)}\right)^{-m-1} \sin(fx+e)}{(2fm+f)\cos(fx+e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x, algorit
hm="fricas")
```

```
output -((a*cos(f*x + e) + a)/cos(f*x + e))^m*((c*cos(f*x + e) - c)/cos(f*x + e))
^(-m - 1)*sin(f*x + e)/((2*f*m + f)*cos(f*x + e))
```

3.164.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-1-m} dx$$

$$= \int (a(\sec(e + fx) + 1))^m (-c(\sec(e + fx) - 1))^{-m-1} \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**m*(c-c*sec(f*x+e))**(-1-m),x)`

output `Integral((a*(sec(e + f*x) + 1))**m*(-c*(sec(e + f*x) - 1))**(-m - 1)*sec(e + f*x), x)`

3.164.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.32

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-1-m} dx$$

$$= \frac{(-a)^m c^{-m-1} (\cos(fx + e) + 1)}{f(2m + 1) \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} \right)^{2m} \sin(fx + e)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^-1-m,x, algorithm="maxima")`

output `(-a)^m*c^(-m - 1)*(cos(f*x + e) + 1)/(f*(2*m + 1)*(sin(f*x + e)/(cos(f*x + e) + 1))^(2*m)*sin(f*x + e))`

3.164.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-1-m} dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m-1} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(1-m)*sec(f*x + e), x)`

3.164.9 Mupad [B] (verification not implemented)

Time = 14.55 (sec) , antiderivative size = 105, normalized size of antiderivative = 2.23

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-1-m} dx =$$

$$\frac{(\sin(e + fx) + \sin(3e + 3fx)) \left(\frac{a(\cos(e+fx)+1)}{\cos(e+fx)} \right)^m}{cf(2m+1) \left(\frac{c(\cos(e+fx)-1)}{\cos(e+fx)} \right)^m (3 \cos(e + fx) - 2 \cos(2e + 2fx) + \cos(3e + 3fx) - 2)}$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^(m + 1)),x)`

output `-((sin(e + f*x) + sin(3*e + 3*f*x))*((a*(cos(e + f*x) + 1))/cos(e + f*x))^m)/(c*f*(2*m + 1)*((c*(cos(e + f*x) - 1))/cos(e + f*x))^m*(3*cos(e + f*x) - 2*cos(2*e + 2*f*x) + cos(3*e + 3*f*x) - 2))`

3.165 $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-m} dx$

3.165.1 Optimal result	1137
3.165.2 Mathematica [A] (verified)	1137
3.165.3 Rubi [A] (verified)	1138
3.165.4 Maple [F]	1139
3.165.5 Fricas [F]	1140
3.165.6 Sympy [F(-1)]	1140
3.165.7 Maxima [F]	1140
3.165.8 Giac [F]	1141
3.165.9 Mupad [F(-1)]	1141

3.165.1 Optimal result

Integrand size = 34, antiderivative size = 101

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-m} dx = \frac{2^{\frac{1}{2}-m}c \operatorname{Hypergeometric2F1}\left(\frac{1}{2}+m, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sec(e+fx))\right) (1-\sec(e+fx))^{\frac{1}{2}+m} (a+a \sec(e+fx))^m}{f(1+2m)}$$

output

```
-2^(1/2-m)*c*hypergeom([1/2+m, 1/2+m], [3/2+m], 1/2+1/2*sec(f*x+e))*(1-sec(f*x+e))^(1/2+m)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m)*tan(f*x+e)/f/(1+2*m)
```

3.165.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.03

$$\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-m} dx = \frac{2^{\frac{1}{2}+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-m, \frac{1}{2}-m, \frac{3}{2}-m, \frac{1}{2}(1-\sec(e+fx))\right) (1+\sec(e+fx))^{-\frac{1}{2}-m} (a(1+\sec(e+fx)))^m}{f(-1+2m)}$$

input

```
Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^m)/(c - c*Sec[e + f*x])^m,x]
```

output $-\left(\frac{2^{1/2+m} \text{Hypergeometric2F1}\left[\frac{1}{2}-m, \frac{1}{2}-m, \frac{3}{2}-m, (1-\text{Sec}[e+fx])/2\right]}{2}\right) \cdot (1+\text{Sec}[e+fx])^{-1/2-m} \cdot (a \cdot (1+\text{Sec}[e+fx]))^m \cdot \text{Tan}[e+fx] / (f \cdot (-1+2m) \cdot (c-c \cdot \text{Sec}[e+fx])^m)$

3.165.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e+fx)(a \sec(e+fx)+a)^m(c-c \sec(e+fx))^{-m} dx$$

$$\downarrow 3042$$

$$\int \csc\left(e+fx+\frac{\pi}{2}\right)\left(a \csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^m\left(c-c \csc\left(e+fx+\frac{\pi}{2}\right)\right)^{-m} dx$$

$$\downarrow 4449$$

$$\frac{a c \tan(e+fx) \int (\sec(e+fx)a+a)^{m-\frac{1}{2}}(c-c \sec(e+fx))^{-m-\frac{1}{2}} d \sec(e+fx)}{f \sqrt{a \sec(e+fx)+a} \sqrt{c-c \sec(e+fx)}}$$

$$\downarrow 80$$

$$\frac{a c 2^{-m-\frac{1}{2}} \tan(e+fx)(1-\sec(e+fx))^{m+\frac{1}{2}}(c-c \sec(e+fx))^{-m-1} \int \left(\frac{1}{2}-\frac{1}{2} \sec(e+fx)\right)^{-m-\frac{1}{2}}(\sec(e+fx)a+)}{f \sqrt{a \sec(e+fx)+a}}$$

$$\downarrow 79$$

$$\frac{c 2^{\frac{1}{2}-m} \tan(e+fx)(1-\sec(e+fx))^{m+\frac{1}{2}}(a \sec(e+fx)+a)^m(c-c \sec(e+fx))^{-m-1} \text{Hypergeometric2F1}\left(m+\frac{1}{2}, \frac{1}{2}+m, \frac{3}{2}+m, \frac{(1+\text{Sec}[e+fx])/2}{a \sec(e+fx)+a}\right)}{f(2m+1)}$$

input $\text{Int}[(\text{Sec}[e+fx] \cdot (a+a \cdot \text{Sec}[e+fx])^m) / (c-c \cdot \text{Sec}[e+fx])^m, x]$

output $-\left(\frac{2^{1/2-m} c \text{Hypergeometric2F1}\left[\frac{1}{2}+m, \frac{1}{2}+m, \frac{3}{2}+m, (1+\text{Sec}[e+fx])/2\right]}{2}\right) \cdot (1-\text{Sec}[e+fx])^{1/2+m} \cdot (a+a \cdot \text{Sec}[e+fx])^m \cdot (c-c \cdot \text{Sec}[e+fx])^{-1-m} \cdot \text{Tan}[e+fx] / (f \cdot (1+2m))$

3.165. $\int \sec(e+fx)(a+a \sec(e+fx))^m(c-c \sec(e+fx))^{-m} dx$

3.165.3.1 Defintions of rubi rules used

rule 79 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`

rule 80 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4449 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.165.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (-c(\sec(fx + e) - 1))^{-m} dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/((c-c*sec(f*x+e))^m),x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m/((c-c*sec(f*x+e))^m),x)`

3.165.5 Fracas [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-m} dx$$

$$= \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^m} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/((c-c*sec(f*x+e))^m),x, algorithm="fricas")`

output `integral((a*sec(f*x + e) + a)^m*sec(f*x + e)/(-c*sec(f*x + e) + c)^m, x)`

3.165.6 Sympy [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-m} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**m/((c-c*sec(f*x+e))**m),x)`

output `Timed out`

3.165.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-m} dx$$

$$= \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^m} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/((c-c*sec(f*x+e))^m),x, algorithm="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(-c*sec(f*x + e) + c)^m, x)`

3.165.8 Giac [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-m} dx$$

$$= \int \frac{(a \sec(fx + e) + a)^m \sec(fx + e)}{(-c \sec(fx + e) + c)^m} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m/((c-c*sec(f*x+e))^m),x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*sec(f*x + e)/(-c*sec(f*x + e) + c)^m, x)`

3.165.9 Mupad [F(-1)]

Timed out.

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{-m} dx$$

$$= \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m}{\cos(e + fx) \left(c - \frac{c}{\cos(e+fx)}\right)^m} dx$$

input `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^m),x)`

output `int((a + a/cos(e + f*x))^m/(cos(e + f*x)*(c - c/cos(e + f*x))^m), x)`

3.166 $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx$

3.166.1 Optimal result	1142
3.166.2 Mathematica [A] (verified)	1142
3.166.3 Rubi [A] (verified)	1143
3.166.4 Maple [F]	1144
3.166.5 Fracas [F]	1145
3.166.6 Sympy [F]	1145
3.166.7 Maxima [F]	1145
3.166.8 Giac [F]	1146
3.166.9 Mupad [F(-1)]	1146

3.166.1 Optimal result

Integrand size = 36, antiderivative size = 99

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx = \frac{2^{\frac{3}{2}-m} c \operatorname{Hypergeometric2F1}\left(-\frac{1}{2} + m, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e + fx))\right) (1 - \sec(e + fx))^{-\frac{1}{2}+m} (a + a \sec(e + fx))}{f(1 + 2m)}$$

```
output -2^(3/2-m)*c*hypergeom([-1/2+m, 1/2+m],[3/2+m],1/2+1/2*sec(f*x+e))*(1-sec(f*x+e))^(1/2+m)*(a+a*sec(f*x+e))^m*tan(f*x+e)/f/(1+2*m)/((c-c*sec(f*x+e))^m)
```

3.166.2 Mathematica [A] (verified)

Time = 1.29 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.06

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx = \frac{2^{\frac{1}{2}+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2} - m, \frac{3}{2} - m, \frac{5}{2} - m, \frac{1}{2}(1 - \sec(e + fx))\right) (1 + \sec(e + fx))^{-\frac{1}{2}-m} (a(1 + \sec(e + fx)) + c)}{f(3 - 2m)}$$

```
input Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(1 - m),x]
```

output $(2^{(1/2 + m)} \text{Hypergeometric2F1}[1/2 - m, 3/2 - m, 5/2 - m, (1 - \text{Sec}[e + f*x])/2] * (1 + \text{Sec}[e + f*x])^{(-1/2 - m)} * (a * (1 + \text{Sec}[e + f*x]))^m * (c - c * \text{Sec}[e + f*x])^{(1 - m)} * \text{Tan}[e + f*x]) / (f * (3 - 2 * m))$

3.166.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{1-m} dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{1-m} dx$$

$$\downarrow 4449$$

$$\frac{a c \tan(e + fx) \int (\sec(e + fx)a + a)^{m-\frac{1}{2}} (c - c \sec(e + fx))^{\frac{1}{2}-m} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}}$$

$$\downarrow 80$$

$$\frac{a c 2^{\frac{1}{2}-m} \tan(e + fx) (1 - \sec(e + fx))^{m-\frac{1}{2}} (c - c \sec(e + fx))^{-m} \int \left(\frac{1}{2} - \frac{1}{2} \sec(e + fx)\right)^{\frac{1}{2}-m} (\sec(e + fx)a + a)^m}{f \sqrt{a \sec(e + fx) + a}}$$

$$\downarrow 79$$

$$\frac{c 2^{\frac{3}{2}-m} \tan(e + fx) (1 - \sec(e + fx))^{m-\frac{1}{2}} (a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m} \text{Hypergeometric2F1}\left(m - \frac{1}{2}, \dots\right)}{f(2m + 1)}$$

input $\text{Int}[\text{Sec}[e + f*x] * (a + a * \text{Sec}[e + f*x])^m * (c - c * \text{Sec}[e + f*x])^{(1 - m)}, x]$

output $-((2^{(3/2 - m)} * c * \text{Hypergeometric2F1}[-1/2 + m, 1/2 + m, 3/2 + m, (1 + \text{Sec}[e + f*x])/2] * (1 - \text{Sec}[e + f*x])^{(-1/2 + m)} * (a + a * \text{Sec}[e + f*x])^m * \text{Tan}[e + f*x]) / (f * (1 + 2 * m) * (c - c * \text{Sec}[e + f*x])^m)$

3.166.3.1 Defintions of rubi rules used

- rule 79 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`
- rule 80 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/(b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n] Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4449 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.166.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (c - c \sec(fx + e))^{1-m} dx$$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x)`

output `int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x)`

3.166.5 Fracas [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m+1} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x, algorithm m="fricas")`

output `integral((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m + 1)*sec(f*x + e), x)`

3.166.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx$$

$$= \int (a(\sec(e + fx) + 1))^m (-c(\sec(e + fx) - 1))^{1-m} \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x)`

output `Integral((a*(sec(e + f*x) + 1))^m*(-c*(sec(e + f*x) - 1))^(1 - m)*sec(e + f*x), x)`

3.166.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m+1} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m + 1)*sec(f*x + e), x)`

3.166.8 Giac [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx \\ &= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m+1} \sec(fx + e) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(1-m),x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m + 1)*sec(f*x + e), x)`

3.166.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{1-m} dx \\ &= \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m \left(c - \frac{c}{\cos(e+fx)}\right)^{1-m}}{\cos(e + fx)} dx \end{aligned}$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(1 - m))/cos(e + f*x),x)`

output `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(1 - m))/cos(e + f*x), x)`

3.167 $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx$

3.167.1 Optimal result	1147
3.167.2 Mathematica [A] (verified)	1147
3.167.3 Rubi [A] (verified)	1148
3.167.4 Maple [F]	1149
3.167.5 Fricas [F]	1150
3.167.6 Sympy [F]	1150
3.167.7 Maxima [F]	1150
3.167.8 Giac [F]	1151
3.167.9 Mupad [F(-1)]	1151

3.167.1 Optimal result

Integrand size = 36, antiderivative size = 101

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx = \frac{2^{\frac{5}{2}-m} c^2 \operatorname{Hypergeometric2F1}\left(-\frac{3}{2} + m, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sec(e + fx))\right) (1 - \sec(e + fx))^{-\frac{1}{2}+m} (a + c \sec(e + fx))}{f(1 + 2m)}$$

```
output -2^(5/2-m)*c^2*hypergeom([1/2+m, -3/2+m],[3/2+m],1/2+1/2*sec(f*x+e))*(1-sec(f*x+e))^(1/2+m)*(a+a*sec(f*x+e))^m*tan(f*x+e)/f/(1+2*m)/((c-c*sec(f*x+e))^m)
```

3.167.2 Mathematica [A] (verified)

Time = 2.69 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.04

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx = \frac{2^{\frac{1}{2}+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2} - m, \frac{5}{2} - m, \frac{7}{2} - m, \frac{1}{2}(1 - \sec(e + fx))\right) (1 + \sec(e + fx))^{-\frac{1}{2}-m} (a(1 + \sec(e + fx)) + c)}{f(5 - 2m)}$$

```
input Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^m*(c - c*Sec[e + f*x])^(2 - m),x]
```

output $(2^{(1/2 + m)} \text{Hypergeometric2F1}[1/2 - m, 5/2 - m, 7/2 - m, (1 - \text{Sec}[e + f*x])/2] * (1 + \text{Sec}[e + f*x])^{(-1/2 - m)} * (a * (1 + \text{Sec}[e + f*x]))^m * (c - c * \text{Sec}[e + f*x])^{(2 - m)} * \text{Tan}[e + f*x]) / (f * (5 - 2 * m))$

3.167.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4449, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{2-m} dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^m \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^{2-m} dx$$

$$\downarrow 4449$$

$$\frac{a c \tan(e + fx) \int (\sec(e + fx)a + a)^{m-\frac{1}{2}} (c - c \sec(e + fx))^{\frac{3}{2}-m} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}}$$

$$\downarrow 80$$

$$\frac{a c^2 2^{\frac{3}{2}-m} \tan(e + fx) (1 - \sec(e + fx))^{m-\frac{1}{2}} (c - c \sec(e + fx))^{-m} \int \left(\frac{1}{2} - \frac{1}{2} \sec(e + fx)\right)^{\frac{3}{2}-m} (\sec(e + fx)a + a)^m d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a}}$$

$$\downarrow 79$$

$$\frac{c^2 2^{\frac{5}{2}-m} \tan(e + fx) (1 - \sec(e + fx))^{m-\frac{1}{2}} (a \sec(e + fx) + a)^m (c - c \sec(e + fx))^{-m} \text{Hypergeometric2F1}(m - \frac{1}{2}, \frac{3}{2} - m, \frac{5}{2} - m, \frac{1 - \sec(e + fx)}{2})}{f(2m + 1)}$$

input $\text{Int}[\text{Sec}[e + f*x] * (a + a * \text{Sec}[e + f*x])^m * (c - c * \text{Sec}[e + f*x])^{(2 - m)}, x]$

output $-((2^{(5/2 - m)} * c^2 * \text{Hypergeometric2F1}[-3/2 + m, 1/2 + m, 3/2 + m, (1 + \text{Sec}[e + f*x])/2] * (1 - \text{Sec}[e + f*x])^{(-1/2 + m)} * (a + a * \text{Sec}[e + f*x])^m * \text{Tan}[e + f*x]) / (f * (1 + 2 * m) * (c - c * \text{Sec}[e + f*x])^m))$

3.167. $\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx$

3.167.3.1 Defintions of rubi rules used

```
rule 79 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

```
rule 80 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4449 Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a*c*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
```

3.167.4 Maple [F]

$$\int \sec(fx + e) (a + a \sec(fx + e))^m (c - c \sec(fx + e))^{2-m} dx$$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2-m),x)
```

```
output int(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2-m),x)
```

3.167.5 Fracas [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m+2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2-m),x, algorithm m="fricas")`

output `integral((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m + 2)*sec(f*x + e), x)`

3.167.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx$$

$$= \int (a(\sec(e + fx) + 1))^m (-c(\sec(e + fx) - 1))^{2-m} \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**m*(c-c*sec(f*x+e))**(2-m),x)`

output `Integral((a*(sec(e + f*x) + 1))**m*(-c*(sec(e + f*x) - 1))**(2 - m)*sec(e + f*x), x)`

3.167.7 Maxima [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx$$

$$= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m+2} \sec(fx + e) dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2-m),x, algorithm m="maxima")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m + 2)*sec(f*x + e), x)`

3.167.8 Giac [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx \\ &= \int (a \sec(fx + e) + a)^m (-c \sec(fx + e) + c)^{-m+2} \sec(fx + e) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^m*(c-c*sec(f*x+e))^(2-m),x, algorithm="giac")`

output `integrate((a*sec(f*x + e) + a)^m*(-c*sec(f*x + e) + c)^(-m + 2)*sec(f*x + e), x)`

3.167.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^m (c - c \sec(e + fx))^{2-m} dx \\ &= \int \frac{\left(a + \frac{a}{\cos(e+fx)}\right)^m \left(c - \frac{c}{\cos(e+fx)}\right)^{2-m}}{\cos(e + fx)} dx \end{aligned}$$

input `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(2 - m))/cos(e + f*x),x)`

output `int(((a + a/cos(e + f*x))^m*(c - c/cos(e + f*x))^(2 - m))/cos(e + f*x), x)`

3.168 $\int \sec^2(e+fx)(a+a \sec(e+fx))^3(c-c \sec(e+fx)) dx$

3.168.1 Optimal result	1152
3.168.2 Mathematica [A] (verified)	1152
3.168.3 Rubi [A] (verified)	1153
3.168.4 Maple [A] (verified)	1154
3.168.5 Fricas [A] (verification not implemented)	1155
3.168.6 Sympy [F]	1155
3.168.7 Maxima [A] (verification not implemented)	1156
3.168.8 Giac [A] (verification not implemented)	1156
3.168.9 Mupad [B] (verification not implemented)	1157

3.168.1 Optimal result

Integrand size = 32, antiderivative size = 105

$$\int \sec^2(e+fx)(a+a \sec(e+fx))^3(c-c \sec(e+fx)) dx$$

$$= \frac{a^3 c \operatorname{arctanh}(\sin(e+fx))}{4f} + \frac{a^3 c \sec(e+fx) \tan(e+fx)}{4f}$$

$$- \frac{a^3 c \sec^3(e+fx) \tan(e+fx)}{2f} - \frac{2a^3 c \tan^3(e+fx)}{3f} - \frac{a^3 c \tan^5(e+fx)}{5f}$$

```
output 1/4*a^3*c*arctanh(sin(f*x+e))/f+1/4*a^3*c*sec(f*x+e)*tan(f*x+e)/f-1/2*a^3*
c*sec(f*x+e)^3*tan(f*x+e)/f-2/3*a^3*c*tan(f*x+e)^3/f-1/5*a^3*c*tan(f*x+e)^
5/f
```

3.168.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.65

$$\int \sec^2(e+fx)(a+a \sec(e+fx))^3(c-c \sec(e+fx)) dx$$

$$= \frac{a^3 c (15 \operatorname{arctanh}(\sin(e+fx)) - \tan(e+fx) (-15 \sec(e+fx) + 30 \sec^3(e+fx) + 40 \tan^2(e+fx) + 12 \tan^4(e+fx)))}{60f}$$

input `Integrate[Sec[e + f*x]^2*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x]),x]`

output `(a^3*c*(15*ArcTanh[Sin[e + f*x]] - Tan[e + f*x]*(-15*Sec[e + f*x] + 30*Sec[e + f*x]^3 + 40*Tan[e + f*x]^2 + 12*Tan[e + f*x]^4)))/(60*f)`

3.168.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4450, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(e + fx)(a \sec(e + fx) + a)^3(c - c \sec(e + fx)) dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right)^2 \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow 4450$$

$$-ac \int (a^2 \tan^2(e + fx) \sec^4(e + fx) + 2a^2 \tan^2(e + fx) \sec^3(e + fx) + a^2 \tan^2(e + fx) \sec^2(e + fx)) dx$$

$$\downarrow 2009$$

$$-ac \left(-\frac{a^2 \operatorname{arctanh}(\sin(e + fx))}{4f} + \frac{a^2 \tan^5(e + fx)}{5f} + \frac{2a^2 \tan^3(e + fx)}{3f} + \frac{a^2 \tan(e + fx) \sec^3(e + fx)}{2f} - \frac{a^2 \tan(e + fx)}{2f} \right)$$

input `Int[Sec[e + f*x]^2*(a + a*Sec[e + f*x])^3*(c - c*Sec[e + f*x]),x]`

output `-(a*c*(-1/4*(a^2*ArcTanh[Sin[e + f*x]]))/f - (a^2*Sec[e + f*x]*Tan[e + f*x])/(4*f) + (a^2*Sec[e + f*x]^3*Tan[e + f*x])/(2*f) + (2*a^2*Tan[e + f*x]^3)/(3*f) + (a^2*Tan[e + f*x]^5)/(5*f))`

3.168.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4450 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[(g*csc[e + f*x])^p*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.168.4 Maple [A] (verified)

Time = 4.05 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.30

method	result
derivativedivides	$\frac{a^3 c \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) - 2a^3 c \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8}}{f}$
default	$\frac{a^3 c \left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15} \right) \tan(fx+e) - 2a^3 c \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8}}{f}$
parts	$\frac{a^3 c \tan(fx+e)}{f} + \frac{a^3 c \sec(fx+e) \tan(fx+e)}{f} + \frac{a^3 c \ln(\sec(fx+e) + \tan(fx+e))}{f} - \frac{2a^3 c \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8}}{f}$
norman	$\frac{a^3 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2f} + \frac{25a^3 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3f} - \frac{64a^3 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{15f} + \frac{7a^3 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{3f} - \frac{a^3 c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{2f} - \frac{a^3 c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)}{4f} \frac{1}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 - 1}$
risch	$-\frac{ia^3 c (15 e^{9i(fx+e)} - 60 e^{8i(fx+e)} - 90 e^{7i(fx+e)} - 240 e^{6i(fx+e)} - 40 e^{4i(fx+e)} + 90 e^{3i(fx+e)} - 80 e^{2i(fx+e)} - 15 e^{i(fx+e)} - 15)}{30 f (1 + e^{2i(fx+e)})^5}$
parallelrisc	$-\frac{3 \left(\left(\frac{5 \cos(fx+e)}{6} + \frac{5 \cos(3fx+3e)}{12} + \frac{\cos(5fx+5e)}{12} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right) - 1 \right) + \left(-\frac{5 \cos(fx+e)}{6} - \frac{5 \cos(3fx+3e)}{12} - \frac{\cos(5fx+5e)}{12} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)}{f (\cos(5fx+5e) + 5 \cos(3fx+3e) + 10)}$

input `int(sec(f*x+e)^2*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

3.168. $\int \sec^2(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$

output $1/f*(a^3*c*(-8/15-1/5*\sec(f*x+e)^4-4/15*\sec(f*x+e)^2)*\tan(f*x+e)-2*a^3*c*(-(-1/4*\sec(f*x+e)^3-3/8*\sec(f*x+e))*\tan(f*x+e)+3/8*\ln(\sec(f*x+e)+\tan(f*x+e))))+2*a^3*c*(1/2*\sec(f*x+e)*\tan(f*x+e)+1/2*\ln(\sec(f*x+e)+\tan(f*x+e)))+a^3*c*\tan(f*x+e)$

3.168.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.25

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{15 a^3 c \cos(fx + e)^5 \log(\sin(fx + e) + 1) - 15 a^3 c \cos(fx + e)^5 \log(-\sin(fx + e) + 1) + 2(28 a^3 c \cos(fx + e)^4 + 15 a^3 c \cos(fx + e)^3 - 16 a^3 c \cos(fx + e)^2 - 30 a^3 c \cos(fx + e) - 12 a^3 c) \sin(fx + e)}{120 f \cos(fx + e)}$$

input `integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x, algorithm="fricas")`

output $1/120*(15*a^3*c*\cos(f*x + e)^5*\log(\sin(f*x + e) + 1) - 15*a^3*c*\cos(f*x + e)^5*\log(-\sin(f*x + e) + 1) + 2*(28*a^3*c*\cos(f*x + e)^4 + 15*a^3*c*\cos(f*x + e)^3 - 16*a^3*c*\cos(f*x + e)^2 - 30*a^3*c*\cos(f*x + e) - 12*a^3*c)*\sin(f*x + e))/(f*\cos(f*x + e)^5)$

3.168.6 Sympy [F]

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= -a^3 c \left(\int (-\sec^2(e + fx)) dx + \int (-2 \sec^3(e + fx)) dx + \int 2 \sec^5(e + fx) dx + \int \sec^6(e + fx) dx \right)$$

input `integrate(sec(f*x+e)**2*(a+a*sec(f*x+e))**3*(c-c*sec(f*x+e)),x)`

output $-a^3*c*(\text{Integral}(-\sec(e + f*x)**2, x) + \text{Integral}(-2*\sec(e + f*x)**3, x) + \text{Integral}(2*\sec(e + f*x)**5, x) + \text{Integral}(\sec(e + f*x)**6, x))$

3.168.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.64

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx =$$

$$\frac{8(3 \tan(fx + e)^5 + 10 \tan(fx + e)^3 + 15 \tan(fx + e))a^3c - 15a^3c \left(\frac{2(3 \sin(fx+e)^3 - 5 \sin(fx+e))}{\sin(fx+e)^4 - 2 \sin(fx+e)^2 + 1} - 3 \log \right)}{60f}$$

```
input integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x, algorithm="maxima")
```

```
output -1/120*(8*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^3*c -
15*a^3*c*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f
*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) + 60*a
^3*c*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(si
n(f*x + e) - 1)) - 120*a^3*c*tan(f*x + e))/f
```

3.168.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.38

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{15a^3c \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + 1 \right| \right) - 15a^3c \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2(15a^3c \tan(\frac{1}{2} fx + \frac{1}{2} e)^9 - 70a^3c \tan(\frac{1}{2} fx + \frac{1}{2} e)^7 + 128a^3c \tan(\frac{1}{2} fx + \frac{1}{2} e)^5 - 250a^3c \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 - 15a^3c \tan(\frac{1}{2} fx + \frac{1}{2} e))}{(\tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 1)^5}}{60f}$$

```
input integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))^3*(c-c*sec(f*x+e)),x, algorithm="giac")
```

```
output 1/60*(15*a^3*c*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*a^3*c*log(abs(tan(1
/2*f*x + 1/2*e) - 1)) - 2*(15*a^3*c*tan(1/2*f*x + 1/2*e)^9 - 70*a^3*c*tan(
1/2*f*x + 1/2*e)^7 + 128*a^3*c*tan(1/2*f*x + 1/2*e)^5 - 250*a^3*c*tan(1/2*
f*x + 1/2*e)^3 - 15*a^3*c*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 -
1)^5)/f
```

3.168.9 Mupad [B] (verification not implemented)

Time = 17.84 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.67

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^3(c - c \sec(e + fx)) dx$$

$$= \frac{-\frac{ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{2} + \frac{7ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{3} - \frac{64ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{15} + \frac{25ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{3} + \frac{ca^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{2}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1 \right)} + \frac{a^3 c \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{2f}$$

input `int(((a + a/cos(e + f*x))^3*(c - c/cos(e + f*x)))/cos(e + f*x)^2,x)`output `((a^3*c*tan(e/2 + (f*x)/2))/2 + (25*a^3*c*tan(e/2 + (f*x)/2)^3)/3 - (64*a^3*c*tan(e/2 + (f*x)/2)^5)/15 + (7*a^3*c*tan(e/2 + (f*x)/2)^7)/3 - (a^3*c*tan(e/2 + (f*x)/2)^9)/2)/(f*(5*tan(e/2 + (f*x)/2)^2 - 10*tan(e/2 + (f*x)/2)^4 + 10*tan(e/2 + (f*x)/2)^6 - 5*tan(e/2 + (f*x)/2)^8 + tan(e/2 + (f*x)/2)^10 - 1)) + (a^3*c*atanh(tan(e/2 + (f*x)/2)))/(2*f)`

3.169 $\int \sec^2(e+fx)(a+a \sec(e+fx))^2(c-c \sec(e+fx)) dx$

3.169.1 Optimal result	1158
3.169.2 Mathematica [A] (verified)	1158
3.169.3 Rubi [A] (verified)	1159
3.169.4 Maple [A] (verified)	1160
3.169.5 Fricas [A] (verification not implemented)	1161
3.169.6 Sympy [F]	1161
3.169.7 Maxima [B] (verification not implemented)	1162
3.169.8 Giac [A] (verification not implemented)	1162
3.169.9 Mupad [B] (verification not implemented)	1163

3.169.1 Optimal result

Integrand size = 32, antiderivative size = 86

$$\int \sec^2(e+fx)(a+a \sec(e+fx))^2(c-c \sec(e+fx)) dx$$

$$= \frac{a^2 c \arctanh(\sin(e+fx))}{8f} + \frac{a^2 c \sec(e+fx) \tan(e+fx)}{8f}$$

$$- \frac{a^2 c \sec^3(e+fx) \tan(e+fx)}{4f} - \frac{a^2 c \tan^3(e+fx)}{3f}$$

output `1/8*a^2*c*arctanh(sin(f*x+e))/f+1/8*a^2*c*sec(f*x+e)*tan(f*x+e)/f-1/4*a^2*c*sec(f*x+e)^3*tan(f*x+e)/f-1/3*a^2*c*tan(f*x+e)^3/f`

3.169.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.66

$$\int \sec^2(e+fx)(a+a \sec(e+fx))^2(c-c \sec(e+fx)) dx$$

$$= \frac{a^2 c (3 \arctanh(\sin(e+fx)) + \tan(e+fx) (3 \sec(e+fx) - 6 \sec^3(e+fx) - 8 \tan^2(e+fx)))}{24f}$$

input `Integrate[Sec[e + f*x]^2*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x]),x]`

output $(a^2*c*(3*ArcTanh[\text{Sin}[e + f*x]] + \text{Tan}[e + f*x]*(3*\text{Sec}[e + f*x] - 6*\text{Sec}[e + f*x]^3 - 8*\text{Tan}[e + f*x]^2)))/(24*f)$

3.169.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.91, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {3042, 4450, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(e + fx)(a \sec(e + fx) + a)^2(c - c \sec(e + fx)) dx$$

↓ 3042

$$\int \csc\left(e + fx + \frac{\pi}{2}\right)^2 \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx$$

↓ 4450

$$-ac \int (a \tan^2(e + fx) \sec^3(e + fx) + a \tan^2(e + fx) \sec^2(e + fx)) dx$$

↓ 2009

$$-ac \left(-\frac{a \operatorname{arctanh}(\sin(e + fx))}{8f} + \frac{a \tan^3(e + fx)}{3f} + \frac{a \tan(e + fx) \sec^3(e + fx)}{4f} - \frac{a \tan(e + fx) \sec(e + fx)}{8f} \right)$$

input $\text{Int}[\text{Sec}[e + f*x]^2*(a + a*\text{Sec}[e + f*x])^2*(c - c*\text{Sec}[e + f*x]),x]$

output $-(a*c*(-1/8*(a*ArcTanh[\text{Sin}[e + f*x]]))/f - (a*\text{Sec}[e + f*x]*\text{Tan}[e + f*x])/(8*f) + (a*\text{Sec}[e + f*x]^3*\text{Tan}[e + f*x])/(4*f) + (a*\text{Tan}[e + f*x]^3)/(3*f))$

3.169.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4450 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[(g*csc[e + f*x])^p*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.169.4 Maple [A] (verified)

Time = 3.81 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.47

method	result
derivativedivides	$-a^2c \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8} \right) + a^2c \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) + \frac{a^2c}{f}$
default	$-a^2c \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8} \right) + a^2c \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) + \frac{a^2c}{f}$
parts	$\frac{a^2c \tan(fx+e)}{f} + \frac{a^2c \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2} \right)}{f} + \frac{a^2c \left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e)}{f} - \frac{a^2c}{f}$
norman	$\frac{-\frac{a^2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f} - \frac{53a^2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{12f} + \frac{11a^2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{12f} - \frac{a^2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{4f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^4} - \frac{a^2c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{8f} + \frac{a^2c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{8f}$
risch	$-\frac{ia^2c(3e^{7i(fx+e)} - 24e^{6i(fx+e)} - 21e^{5i(fx+e)} - 24e^{4i(fx+e)} + 21e^{3i(fx+e)} - 8e^{2i(fx+e)} - 3e^{i(fx+e)} - 8)}{12f(1+e^{2i(fx+e)})^4} + \frac{a^2c \ln(e^{i(fx+e)} - 1)}{8f}$
parallelrisc	$\frac{a^2c \left(\left(-2 \cos(2fx+2e) - \frac{\cos(4fx+4e)}{2} - \frac{3}{2} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \left(2 \cos(2fx+2e) + \frac{\cos(4fx+4e)}{2} + \frac{3}{2} \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \right)}{4f(3+\cos(4fx+4e)+4\cos(2fx+2e))}$

input `int(sec(f*x+e)^2*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

$$3.169. \int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

output `1/f*(-a^2*c*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln(sec(f*x+e)+tan(f*x+e)))+a^2*c*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)+a^2*c*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+a^2*c*tan(f*x+e))`

3.169.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.36

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{3a^2c \cos(fx + e)^4 \log(\sin(fx + e) + 1) - 3a^2c \cos(fx + e)^4 \log(-\sin(fx + e) + 1) + 2(8a^2c \cos(fx + e)^3 - 8a^2c \cos(fx + e) - 6a^2c) \sin(fx + e)}{48f \cos(fx + e)^4}$$

input `integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `1/48*(3*a^2*c*cos(f*x + e)^4*log(sin(f*x + e) + 1) - 3*a^2*c*cos(f*x + e)^4*log(-sin(f*x + e) + 1) + 2*(8*a^2*c*cos(f*x + e)^3 + 3*a^2*c*cos(f*x + e)^2 - 8*a^2*c*cos(f*x + e) - 6*a^2*c)*sin(f*x + e))/(f*cos(f*x + e)^4)`

3.169.6 Sympy [F]

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= -a^2c \left(\int (-\sec^2(e + fx)) dx + \int (-\sec^3(e + fx)) dx + \int \sec^4(e + fx) dx + \int \sec^5(e + fx) dx \right)$$

input `integrate(sec(f*x+e)**2*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e)),x)`

output `-a**2*c*(Integral(-sec(e + f*x)**2, x) + Integral(-sec(e + f*x)**3, x) + Integral(sec(e + f*x)**4, x) + Integral(sec(e + f*x)**5, x))`

3.169.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 160 vs. 2(78) = 156.

Time = 0.23 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.86

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx =$$

$$\frac{16 (\tan (fx + e))^3 + 3 \tan (fx + e) a^2 c - 3 a^2 c \left(\frac{2 (3 \sin (fx + e)^3 - 5 \sin (fx + e))}{\sin (fx + e)^4 - 2 \sin (fx + e)^2 + 1} - 3 \log (\sin (fx + e) + 1) + 3 \log (\sin (fx + e) - 1) \right)}{24 f}$$

input `integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-1/48*(16*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^2*c - 3*a^2*c*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) + 12*a^2*c*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 48*a^2*c*tan(f*x + e))/f`

3.169.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.49

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{3 a^2 c \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + 1 \right| \right) - 3 a^2 c \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(3 a^2 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 - 11 a^2 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 + 53 a^2 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 + 3 a^2 c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right)}{\left(\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right)^2 - 1}}{24 f}$$

input `integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="giac")`

output `1/24*(3*a^2*c*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 3*a^2*c*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(3*a^2*c*tan(1/2*f*x + 1/2*e)^7 - 11*a^2*c*tan(1/2*f*x + 1/2*e)^5 + 53*a^2*c*tan(1/2*f*x + 1/2*e)^3 + 3*a^2*c*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^4)/f`

3.169. $\int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$

3.169.9 Mupad [B] (verification not implemented)

Time = 16.59 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.70

$$\int \sec^2(e + fx)(a + a \sec(e + fx))^2(c - c \sec(e + fx)) dx$$

$$= \frac{a^2 c \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{4f}$$

$$- \frac{\frac{ca^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7}{4} - \frac{11ca^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{12} + \frac{53ca^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{12} + \frac{ca^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1 \right)}$$

input `int(((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x)))/cos(e + f*x)^2,x)`output `(a^2*c*atanh(tan(e/2 + (f*x)/2)))/(4*f) - ((a^2*c*tan(e/2 + (f*x)/2))/4 + (53*a^2*c*tan(e/2 + (f*x)/2)^3)/12 - (11*a^2*c*tan(e/2 + (f*x)/2)^5)/12 + (a^2*c*tan(e/2 + (f*x)/2)^7)/4)/(f*(6*tan(e/2 + (f*x)/2)^4 - 4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/2)^8 + 1))`

3.170 $\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$

3.170.1 Optimal result	1164
3.170.2 Mathematica [A] (verified)	1164
3.170.3 Rubi [A] (verified)	1165
3.170.4 Maple [B] (verified)	1166
3.170.5 Fricas [B] (verification not implemented)	1167
3.170.6 Sympy [B] (verification not implemented)	1168
3.170.7 Maxima [B] (verification not implemented)	1168
3.170.8 Giac [A] (verification not implemented)	1169
3.170.9 Mupad [B] (verification not implemented)	1169

3.170.1 Optimal result

Integrand size = 30, antiderivative size = 17

$$\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx = -\frac{a c \tan^3(e + fx)}{3f}$$

output `-1/3*a*c*tan(f*x+e)^3/f`

3.170.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx = -\frac{a c \tan^3(e + fx)}{3f}$$

input `Integrate[Sec[e + f*x]^2*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x]),x]`

output `-1/3*(a*c*Tan[e + f*x]^3)/f`

3.170.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4450, 3042, 3087, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(e + fx)(a \sec(e + fx) + a)(c - c \sec(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right)^2 \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4450} \\
 & -ac \int \sec^2(e + fx) \tan^2(e + fx) dx \\
 & \quad \downarrow \text{3042} \\
 & -ac \int \sec(e + fx)^2 \tan(e + fx)^2 dx \\
 & \quad \downarrow \text{3087} \\
 & -\frac{ac \int \tan^2(e + fx) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{15} \\
 & -\frac{ac \tan^3(e + fx)}{3f}
 \end{aligned}$$

input `Int[Sec[e + f*x]^2*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x]),x]`

output `-1/3*(a*c*Tan[e + f*x]^3)/f`

3.170.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

rule 4450 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[(g*csc[e + f*x])^p*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.170.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(15) = 30$.

Time = 1.97 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.00

method	result	size
norman	$\frac{8ac \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3f \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^3}$	34
risch	$\frac{2iac(3e^{4i(fx+e)}+1)}{3f(1+e^{2i(fx+e)})^3}$	35
derivativedivides	$\frac{ac\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e) + ac \tan(fx+e)}{f}$	36
default	$\frac{ac\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e) + ac \tan(fx+e)}{f}$	36
parts	$\frac{ac \tan(fx+e)}{f} + \frac{ac\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e)}{f}$	38
parallelrisc	$\frac{8ac \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3f \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^3 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^3}$	45

```
input int(sec(f*x+e)^2*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x,method=_RETURNVERBOSE
)
```

```
output 8/3*a*c/f*tan(1/2*f*x+1/2*e)^3/(tan(1/2*f*x+1/2*e)^2-1)^3
```

3.170.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(15) = 30$.

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.06

$$\int \sec^2(e+fx)(a+a\sec(e+fx))(c-c\sec(e+fx)) dx = \frac{(ac \cos(fx+e)^2 - ac) \sin(fx+e)}{3f \cos(fx+e)^3}$$

```
input integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="fri
cas")
```

```
output 1/3*(a*c*cos(f*x + e)^2 - a*c)*sin(f*x + e)/(f*cos(f*x + e)^3)
```


3.170.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. $2(15) = 30$.

Time = 0.85 (sec) , antiderivative size = 51, normalized size of antiderivative = 3.00

$$\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \begin{cases} \frac{-ac \left(\frac{\tan^3(e+fx)}{3} + \tan(e+fx) \right) + ac \tan(e+fx)}{f} & \text{for } f \neq 0 \\ x(a \sec(e) + a)(-c \sec(e) + c) \sec^2(e) & \text{otherwise} \end{cases}$$

input `integrate(sec(f*x+e)**2*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x)`

output `Piecewise(((-a*c*(tan(e + f*x)**3/3 + tan(e + f*x)) + a*c*tan(e + f*x))/f, Ne(f, 0)), (x*(a*sec(e) + a)*(-c*sec(e) + c)*sec(e)**2, True))`

3.170.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(15) = 30$.

Time = 0.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.12

$$\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= -\frac{(\tan(fx + e))^3 + 3 \tan(fx + e)ac - 3ac \tan(fx + e)}{3f}$$

input `integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-1/3*((tan(f*x + e)^3 + 3*tan(f*x + e))*a*c - 3*a*c*tan(f*x + e))/f`

3.170.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx = -\frac{a c \tan(fx + e)^3}{3f}$$

input `integrate(sec(f*x+e)^2*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="giac")`

output `-1/3*a*c*tan(f*x + e)^3/f`

3.170.9 Mupad [B] (verification not implemented)

Time = 13.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \sec^2(e + fx)(a + a \sec(e + fx))(c - c \sec(e + fx)) dx = -\frac{a c \tan(e + fx)^3}{3f}$$

input `int(((a + a/cos(e + f*x))*(c - c/cos(e + f*x)))/cos(e + f*x)^2,x)`

output `-(a*c*tan(e + f*x)^3)/(3*f)`

3.171 $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$

3.171.1 Optimal result 1170
 3.171.2 Mathematica [B] (verified) 1170
 3.171.3 Rubi [A] (verified) 1171
 3.171.4 Maple [A] (verified) 1173
 3.171.5 Fricas [A] (verification not implemented) 1174
 3.171.6 Sympy [F] 1174
 3.171.7 Maxima [B] (verification not implemented) 1174
 3.171.8 Giac [A] (verification not implemented) 1175
 3.171.9 Mupad [B] (verification not implemented) 1175

3.171.1 Optimal result

Integrand size = 32, antiderivative size = 56

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = \frac{2c\operatorname{arctanh}(\sin(e+fx))}{af} - \frac{c\tan(e+fx)}{af} - \frac{2c\tan(e+fx)}{f(a+a\sec(e+fx))}$$

output `2*c*arctanh(sin(f*x+e))/a/f-c*tan(f*x+e)/a/f-2*c*tan(f*x+e)/f/(a+a*sec(f*x+e))`

3.171.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 154 vs. 2(56) = 112.

Time = 0.60 (sec) , antiderivative size = 154, normalized size of antiderivative = 2.75

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = \frac{c\left(\frac{2\log(\cos(\frac{1}{2}(e+fx))-\sin(\frac{1}{2}(e+fx)))}{f} - \frac{2\log(\cos(\frac{1}{2}(e+fx))+\sin(\frac{1}{2}(e+fx)))}{f}\right) + \frac{\sin(\frac{1}{2}(e+fx))}{f(\cos(\frac{1}{2}(e+fx))-\sin(\frac{1}{2}(e+fx)))} + \frac{\sin(\frac{1}{2}(e+fx))}{f(\cos(\frac{1}{2}(e+fx))+\sin(\frac{1}{2}(e+fx)))}}{a}$$

input `Integrate[(Sec[e + f*x]^2*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output $-\left(\frac{c\left(2\log\left[\cos\left(\frac{e+fx}{2}\right)-\sin\left(\frac{e+fx}{2}\right)\right]}{f}-\frac{2\log\left[\cos\left(\frac{e+fx}{2}\right)+\sin\left(\frac{e+fx}{2}\right)\right]}{f}+\frac{\sin\left(\frac{e+fx}{2}\right)}{f\left(\cos\left(\frac{e+fx}{2}\right)-\sin\left(\frac{e+fx}{2}\right)\right)}+\frac{\sin\left(\frac{e+fx}{2}\right)}{f\left(\cos\left(\frac{e+fx}{2}\right)+\sin\left(\frac{e+fx}{2}\right)\right)}+\frac{2\tan\left(\frac{e+fx}{2}\right)}{f}\right)/a$

3.171.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.281$, Rules used = {3042, 4496, 25, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a\sec(e+fx)+a} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)^2(c-c\csc\left(e+fx+\frac{\pi}{2}\right))}{a\csc\left(e+fx+\frac{\pi}{2}\right)+a} dx$$

$$\downarrow \text{4496}$$

$$\frac{\int -\sec(e+fx)(2ac-ac\sec(e+fx))dx}{a^2} - \frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)}$$

$$\downarrow \text{25}$$

$$\frac{\int \sec(e+fx)(2ac-ac\sec(e+fx))dx}{a^2} - \frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)}$$

$$\downarrow \text{3042}$$

$$\frac{\int \csc\left(e+fx+\frac{\pi}{2}\right)(2ac-ac\csc\left(e+fx+\frac{\pi}{2}\right))dx}{a^2} - \frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)}$$

$$\downarrow \text{4274}$$

$$\frac{2ac\int \sec(e+fx)dx-ac\int \sec^2(e+fx)dx}{a^2} - \frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)}$$

$$\downarrow \text{3042}$$

$$\frac{2ac\int \csc\left(e+fx+\frac{\pi}{2}\right)dx-ac\int \csc\left(e+fx+\frac{\pi}{2}\right)^2dx}{a^2} - \frac{2c\tan(e+fx)}{f(a\sec(e+fx)+a)}$$

$$\downarrow \text{4254}$$

3.171. $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$

$$\frac{\frac{ac \int 1d(-\tan(e+fx))}{f} + 2ac \int \csc(e+fx + \frac{\pi}{2}) dx}{a^2} - \frac{2c \tan(e+fx)}{f(a \sec(e+fx) + a)}$$

↓ 24

$$\frac{2ac \int \csc(e+fx + \frac{\pi}{2}) dx - \frac{ac \tan(e+fx)}{f}}{a^2} - \frac{2c \tan(e+fx)}{f(a \sec(e+fx) + a)}$$

↓ 4257

$$\frac{\frac{2ac \operatorname{arctanh}(\sin(e+fx))}{f} - \frac{ac \tan(e+fx)}{f}}{a^2} - \frac{2c \tan(e+fx)}{f(a \sec(e+fx) + a)}$$

input `Int[(Sec[e + f*x]^2*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output `(-2*c*Tan[e + f*x])/(f*(a + a*Sec[e + f*x])) + ((2*a*c*ArcTanh[Sin[e + f*x]])/f - (a*c*Tan[e + f*x])/f)/a^2`

3.171.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4496 `Int[csc[(e_.) + (f_.)*(x_)]^2*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-A*b - a*B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1))), x] + Simp[1/(b^2*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[A*b*m - a*B*m + b*B*(2*m + 1)*Csc[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`

3.171.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.39

method	result
derivativedivides	$\frac{2c \left(-\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 2} - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) + \frac{1}{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 2} \right)}{fa}$
default	$\frac{2c \left(-\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \frac{1}{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 2} - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) + \frac{1}{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 2} \right)}{fa}$
parallelrisc	$-\frac{c \left(2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) \cos(fx+e) - 2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \cos(fx+e) + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \cos(fx+e) + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \right)}{af \cos(fx+e)}$
risc	$-\frac{2ic(2e^{2i(fx+e)} + e^{i(fx+e)} + 3)}{fa(e^{i(fx+e)} + 1)(1 + e^{2i(fx+e)})} + \frac{2c \ln(e^{i(fx+e)} + i)}{af} - \frac{2c \ln(e^{i(fx+e)} - i)}{af}$
norman	$-\frac{4c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{6c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} - \frac{2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{af} - \frac{2c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{af} + \frac{2c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{af}$

input `int(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `2/f*c/a*(-tan(1/2*f*x+1/2*e)+1/2/(tan(1/2*f*x+1/2*e)-1)-ln(tan(1/2*f*x+1/2*e)-1)+ln(tan(1/2*f*x+1/2*e)+1)+1/2/(tan(1/2*f*x+1/2*e)+1))`

3.171.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.88

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$$

$$= \frac{(c\cos(fx+e)^2+c\cos(fx+e))\log(\sin(fx+e)+1) - (c\cos(fx+e)^2+c\cos(fx+e))\log(-\sin(fx+e))}{af\cos(fx+e)^2+af\cos(fx+e)}$$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="fricas")`

output `((c*cos(f*x + e)^2 + c*cos(f*x + e))*log(sin(f*x + e) + 1) - (c*cos(f*x + e)^2 + c*cos(f*x + e))*log(-sin(f*x + e) + 1) - (3*c*cos(f*x + e) + c)*sin(f*x + e))/(a*f*cos(f*x + e)^2 + a*f*cos(f*x + e))`

3.171.6 Sympy [F]

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx = -\frac{c\left(\int\left(-\frac{\sec^2(e+fx)}{\sec(e+fx)+1}\right) dx + \int\frac{\sec^3(e+fx)}{\sec(e+fx)+1} dx\right)}{a}$$

input `integrate(sec(f*x+e)**2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x)`

output `-c*(Integral(-sec(e + f*x)**2/(sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x) + 1), x))/a`

3.171.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. 2(56) = 112.

Time = 0.25 (sec) , antiderivative size = 194, normalized size of antiderivative = 3.46

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$$

$$= \frac{c\left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}+1\right)}{a} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}-1\right)}{a} - \frac{2\sin(fx+e)}{\left(a-\frac{a\sin(fx+e)^2}{(\cos(fx+e)+1)^2}\right)(\cos(fx+e)+1)} - \frac{\sin(fx+e)}{a(\cos(fx+e)+1)}\right) + c\left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}+1\right)}{a}\right)}{f}$$

3.171. $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `(c*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a - 2*sin(f*x + e)/((a - a*sin(f*x + e)^2/(cos(f*x + e) + 1)^2*(cos(f*x + e) + 1)) - sin(f*x + e)/(a*(cos(f*x + e) + 1))) + c*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a - sin(f*x + e)/(a*(cos(f*x + e) + 1))))/f`

3.171.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.55

$$\int \frac{\sec^2(e + fx)(c - c \sec(e + fx))}{a + a \sec(e + fx)} dx$$

$$= \frac{2 \left(\frac{c \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) + 1|)}{a} - \frac{c \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) - 1|)}{a} - \frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e)}{a} + \frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e)}{(\tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 1)a} \right)}{f}$$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="giac")`

output `2*(c*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a - c*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a - c*tan(1/2*f*x + 1/2*e)/a + c*tan(1/2*f*x + 1/2*e)/((tan(1/2*f*x + 1/2*e)^2 - 1)*a))/f`

3.171.9 Mupad [B] (verification not implemented)

Time = 13.56 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.27

$$\int \frac{\sec^2(e + fx)(c - c \sec(e + fx))}{a + a \sec(e + fx)} dx = \frac{4c \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{af} - \frac{2c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a - a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2\right)} - \frac{2c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{af}$$

input `int((c - c/cos(e + f*x))/(cos(e + f*x)^2*(a + a/cos(e + f*x))),x)`

3.171. $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$

output $(4*c*atanh(\tan(e/2 + (f*x)/2)))/(a*f) - (2*c*\tan(e/2 + (f*x)/2))/(f*(a - a$
 $*\tan(e/2 + (f*x)/2)^2) - (2*c*\tan(e/2 + (f*x)/2))/(a*f)$

3.171. $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{a+a\sec(e+fx)} dx$

3.172 $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$

3.172.1 Optimal result 1177
 3.172.2 Mathematica [A] (verified) 1177
 3.172.3 Rubi [A] (verified) 1178
 3.172.4 Maple [A] (verified) 1180
 3.172.5 Fricas [A] (verification not implemented) 1180
 3.172.6 Sympy [F] 1181
 3.172.7 Maxima [B] (verification not implemented) 1181
 3.172.8 Giac [A] (verification not implemented) 1182
 3.172.9 Mupad [B] (verification not implemented) 1182

3.172.1 Optimal result

Integrand size = 32, antiderivative size = 70

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx = -\frac{\operatorname{arctanh}(\sin(e+fx))}{a^2 f} + \frac{7c \tan(e+fx)}{3a^2 f(1+\sec(e+fx))} - \frac{2c \tan(e+fx)}{3f(a+a\sec(e+fx))^2}$$

output `-c*arctanh(sin(f*x+e))/a^2/f+7/3*c*tan(f*x+e)/a^2/f/(1+sec(f*x+e))-2/3*c*tan(f*x+e)/f/(a+a*sec(f*x+e))^2`

3.172.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx = \frac{c\left(-3\operatorname{arctanh}(\sin(e+fx)) + \frac{(5+7\sec(e+fx))\tan(e+fx)}{(1+\sec(e+fx))^2}\right)}{3a^2 f}$$

input `Integrate[(Sec[e + f*x]^2*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2,x]`

output `(c*(-3*ArcTanh[Sin[e + f*x]] + ((5 + 7*Sec[e + f*x])*Tan[e + f*x])/(1 + Sec[e + f*x]^2)))/(3*a^2*f)`

3.172.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4496, 25, 3042, 4486, 3042, 4257, 4281}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a\sec(e+fx)+a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})^2(c-c\csc(e+fx+\frac{\pi}{2}))}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx \\
 & \quad \downarrow \text{4496} \\
 & -\frac{\int -\frac{\sec(e+fx)(4ac-3ac\sec(e+fx))}{\sec(e+fx)a+a} dx}{3a^2} - \frac{2c\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{\sec(e+fx)(4ac-3ac\sec(e+fx))}{\sec(e+fx)a+a} dx}{3a^2} - \frac{2c\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(4ac-3ac\csc(e+fx+\frac{\pi}{2}))}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a^2} - \frac{2c\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \\
 & \quad \downarrow \text{4486} \\
 & \frac{7ac \int \frac{\sec(e+fx)}{\sec(e+fx)a+a} dx - 3c \int \sec(e+fx) dx}{3a^2} - \frac{2c\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{7ac \int \frac{\csc(e+fx+\frac{\pi}{2})}{\csc(e+fx+\frac{\pi}{2})a+a} dx - 3c \int \csc(e+fx+\frac{\pi}{2}) dx}{3a^2} - \frac{2c\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \\
 & \quad \downarrow \text{4257} \\
 & \frac{7ac \int \frac{\csc(e+fx+\frac{\pi}{2})}{\csc(e+fx+\frac{\pi}{2})a+a} dx - \frac{3c\operatorname{arctanh}(\sin(e+fx))}{f}}{3a^2} - \frac{2c\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \\
 & \quad \downarrow \text{4281}
 \end{aligned}$$

3.172. $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$

$$\frac{\frac{7a \tan(e+fx)}{f(a \sec(e+fx)+a)} - \frac{3c \operatorname{arctanh}(\sin(e+fx))}{f}}{3a^2} - \frac{2c \tan(e+fx)}{3f(a \sec(e+fx)+a)^2}$$

input `Int[(Sec[e + f*x]^2*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2,x]`

output `(-2*c*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) + ((-3*c*ArcTanh[Sin[e + f*x]])/f + (7*a*c*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))) / (3*a^2)`

3.172.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4281 `Int[csc[(e_) + (f_)*(x_) / (csc[(e_) + (f_)*(x_)]*(b_) + (a_))], x_Symbol] := Simp[-Cot[e + f*x] / (f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4486 `Int[(csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(B_) + (A_))) / (csc[(e_) + (f_)*(x_)]*(b_) + (a_)), x_Symbol] := Simp[B/b Int[Csc[e + f*x], x], x] + Simp[(A*b - a*B) / b Int[Csc[e + f*x] / (a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0]`

rule 4496 `Int[csc[(e_) + (f_)*(x_)]^2*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(B_) + (A_)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*((a + b*Csc[e + f*x])^m / (b*f*(2*m + 1))), x] + Simp[1 / (b^2*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[A*b*m - a*B*m + b*B*(2*m + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`

3.172.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{c \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \right)}{f a^2}$
default	$\frac{c \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \right)}{f a^2}$
parallelrisch	$\frac{c \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - 3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) \right)}{3 a^2 f}$
risch	$\frac{2ic(3e^{2i(fx+e)}+12e^{i(fx+e)}+5)}{3fa^2(e^{i(fx+e)}+1)^3} + \frac{c \ln(e^{i(fx+e)}-i)}{a^2 f} - \frac{c \ln(e^{i(fx+e)}+i)}{a^2 f}$
norman	$\frac{\frac{2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} - \frac{11c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} + \frac{4c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{3af} + \frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{3af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 a} + \frac{c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{a^2 f} - \frac{c \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{a^2 f}$

input `int(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `1/f/a^2*c*(1/3*tan(1/2*f*x+1/2*e)^3+2*tan(1/2*f*x+1/2*e)+ln(tan(1/2*f*x+1/2*e)-1)-ln(tan(1/2*f*x+1/2*e)+1))`

3.172.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.76

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx =$$

$$\frac{3(c\cos(fx+e)^2+2c\cos(fx+e)+c)\log(\sin(fx+e)+1)-3(c\cos(fx+e)^2+2c\cos(fx+e)+c)}{6(a^2f\cos(fx+e)^2+2a^2f\cos(fx+e)+c)}$$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="fracas")`

output
$$\frac{-1/6*(3*(c*\cos(f*x + e)^2 + 2*c*\cos(f*x + e) + c)*\log(\sin(f*x + e) + 1) - 3*(c*\cos(f*x + e)^2 + 2*c*\cos(f*x + e) + c)*\log(-\sin(f*x + e) + 1) - 2*(5*c*\cos(f*x + e) + 7*c)*\sin(f*x + e))/(a^2*f*\cos(f*x + e)^2 + 2*a^2*f*\cos(f*x + e) + a^2*f)}{a^2}$$

3.172.6 Sympy [F]

$$\int \frac{\sec^2(e + fx)(c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx$$

$$= -\frac{c \left(\int \left(-\frac{\sec^2(e + fx)}{\sec^2(e + fx) + 2 \sec(e + fx) + 1} \right) dx + \int \frac{\sec^3(e + fx)}{\sec^2(e + fx) + 2 \sec(e + fx) + 1} dx \right)}{a^2}$$

input `integrate(sec(f*x+e)**2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**2,x)`

output
$$-c*(\text{Integral}(-\sec(e + fx)**2/(\sec(e + fx)**2 + 2*\sec(e + fx) + 1), x) + \text{Integral}(\sec(e + fx)**3/(\sec(e + fx)**2 + 2*\sec(e + fx) + 1), x))/a**2$$

3.172.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. $2(66) = 132$.

Time = 0.23 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.06

$$\int \frac{\sec^2(e + fx)(c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx$$

$$= \frac{c \left(\frac{9 \sin(fx+e) + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3}}{a^2} - \frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^2} + \frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^2} \right) + \frac{c \left(\frac{3 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^2}}{6f}$$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output
$$\frac{1/6*(c*((9*\sin(f*x + e)/(\cos(f*x + e) + 1) + \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2 - 6*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/a^2 + 6*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/a^2) + c*(3*\sin(f*x + e)/(\cos(f*x + e) + 1) + \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2)/f}{6f}$$

3.172.
$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$$

3.172.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.16

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$$

$$= -\frac{\frac{3c\log(|\tan(\frac{1}{2}fx+\frac{1}{2}e)+1|)}{a^2} - \frac{3c\log(|\tan(\frac{1}{2}fx+\frac{1}{2}e)-1|)}{a^2} - \frac{a^4c\tan(\frac{1}{2}fx+\frac{1}{2}e)^3+6a^4c\tan(\frac{1}{2}fx+\frac{1}{2}e)}{a^6}}{3f}$$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output `-1/3*(3*c*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^2 - 3*c*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a^2 - (a^4*c*tan(1/2*f*x + 1/2*e)^3 + 6*a^4*c*tan(1/2*f*x + 1/2*e))/a^6)/f`

3.172.9 Mupad [B] (verification not implemented)

Time = 13.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.63

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{c\left(6\tan\left(\frac{e}{2}+\frac{fx}{2}\right)-6\operatorname{atanh}\left(\tan\left(\frac{e}{2}+\frac{fx}{2}\right)\right)+\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^3\right)}{3a^2f}$$

input `int((c - c/cos(e + f*x))/(cos(e + f*x)^2*(a + a/cos(e + f*x))^2),x)`

output `(c*(6*tan(e/2 + (f*x)/2) - 6*atanh(tan(e/2 + (f*x)/2)) + tan(e/2 + (f*x)/2)^3))/(3*a^2*f)`

$$3.173 \quad \int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

3.173.1 Optimal result	1183
3.173.2 Mathematica [A] (verified)	1183
3.173.3 Rubi [A] (verified)	1184
3.173.4 Maple [A] (verified)	1186
3.173.5 Fricas [A] (verification not implemented)	1186
3.173.6 Sympy [F]	1187
3.173.7 Maxima [A] (verification not implemented)	1187
3.173.8 Giac [A] (verification not implemented)	1188
3.173.9 Mupad [B] (verification not implemented)	1188

3.173.1 Optimal result

Integrand size = 32, antiderivative size = 86

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx = -\frac{2c\tan(e+fx)}{5f(a+a\sec(e+fx))^3} + \frac{11c\tan(e+fx)}{15af(a+a\sec(e+fx))^2} - \frac{4c\tan(e+fx)}{15f(a^3+a^3\sec(e+fx))}$$

output `-2/5*c*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+11/15*c*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2-4/15*c*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))`

3.173.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.55

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx = \frac{c(1+3\sec(e+fx)-4\sec^2(e+fx))\tan(e+fx)}{15a^3f(1+\sec(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]^2*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^3,x]`

output `(c*(1 + 3*Sec[e + f*x] - 4*Sec[e + f*x]^2)*Tan[e + f*x])/(15*a^3*f*(1 + Sec[e + f*x])^3)`

3.173. $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$

3.173.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4496, 25, 3042, 4488, 3042, 4281}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a\sec(e+fx)+a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})^2(c-c\csc(e+fx+\frac{\pi}{2}))}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\
 & \quad \downarrow \text{4496} \\
 & -\frac{\int -\frac{\sec(e+fx)(6ac-5ac\sec(e+fx))}{(\sec(e+fx)a+a)^2} dx}{5a^2} - \frac{2c\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{\sec(e+fx)(6ac-5ac\sec(e+fx))}{(\sec(e+fx)a+a)^2} dx}{5a^2} - \frac{2c\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(6ac-5ac\csc(e+fx+\frac{\pi}{2}))}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a^2} - \frac{2c\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\
 & \quad \downarrow \text{4488} \\
 & \frac{11ac\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} - \frac{4}{3}c \int \frac{\sec(e+fx)}{\sec(e+fx)a+a} dx - \frac{2c\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{11ac\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} - \frac{4}{3}c \int \frac{\csc(e+fx+\frac{\pi}{2})}{\csc(e+fx+\frac{\pi}{2})a+a} dx - \frac{2c\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\
 & \quad \downarrow \text{4281} \\
 & \frac{11ac\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} - \frac{4c\tan(e+fx)}{3f(a\sec(e+fx)+a)} - \frac{2c\tan(e+fx)}{5f(a\sec(e+fx)+a)^3}
 \end{aligned}$$

3.173. $\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$

input `Int[(Sec[e + f*x]^2*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^3,x]`

output `(-2*c*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) + ((11*a*c*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) - (4*c*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x]))) / (5*a^2)`

3.173.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4281 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[-Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4488 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(A*b - a*B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*B*m + A*b*(m + 1))/(a*b*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && NeQ[a*B*m + A*b*(m + 1), 0] && LtQ[m, -2^(-1)]`

rule 4496 `Int[csc[(e_.) + (f_.)*(x_)]^2*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-A*b - a*B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(b*f*(2*m + 1))), x] + Simp[1/(b^2*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[A*b*m - a*B*m + b*B*(2*m + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`

3.173.4 Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.42

method	result	size
parallelrisc	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 \left(3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 + 5\right)}{30a^3 f}$	36
derivativedivides	$c \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} \right) / (2f a^3)$	37
default	$c \left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} \right) / (2f a^3)$	37
risc	$\frac{2ic(15e^{3i(fx+e)} - 5e^{2i(fx+e)} + 5e^{i(fx+e)} + 1)}{15f a^3 (e^{i(fx+e)} + 1)^5}$	59
norman	$\frac{-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{6af} + \frac{7c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{30af} + \frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{30af} - \frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{10af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 a^2}$	101

```
input int(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output -1/30*c*tan(1/2*f*x+1/2*e)^3*(3*tan(1/2*f*x+1/2*e)^2+5)/a^3/f
```

3.173.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.91

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{(c \cos(fx+e))^2 + 3c \cos(fx+e) - 4c) \sin(fx+e)}{15(a^3 f \cos(fx+e))^3 + 3a^3 f \cos(fx+e)^2 + 3a^3 f \cos(fx+e) + a^3 f}$$

```
input integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="fracas")
```

```
output 1/15*(c*cos(f*x + e)^2 + 3*c*cos(f*x + e) - 4*c)*sin(f*x + e)/(a^3*f*cos(f*x + e)^3 + 3*a^3*f*cos(f*x + e)^2 + 3*a^3*f*cos(f*x + e) + a^3*f)
```

3.173.
$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

3.173.6 Sympy [F]

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{c \left(\int \left(-\frac{\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} \right) dx + \int \frac{\sec^3(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx \right)}{a^3}$$

input `integrate(sec(f*x+e)**2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**3,x)`

output `-c*(Integral(-sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(sec(e + f*x)**3/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3`

3.173.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.34

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= -\frac{c \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} + \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3} - \frac{3c \left(\frac{5 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3}$$

$60 f$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output `-1/60*(c*(15*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 3*c*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3)/f`

3.173.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.43

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx = -\frac{3c\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^5+5c\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3}{30a^3f}$$

input `integrate(sec(f*x+e)^2*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output `-1/30*(3*c*tan(1/2*f*x + 1/2*e)^5 + 5*c*tan(1/2*f*x + 1/2*e)^3)/(a^3*f)`

3.173.9 Mupad [B] (verification not implemented)

Time = 13.29 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.41

$$\int \frac{\sec^2(e+fx)(c-c\sec(e+fx))}{(a+a\sec(e+fx))^3} dx = -\frac{c\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^3\left(3\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^2+5\right)}{30a^3f}$$

input `int((c - c/cos(e + f*x))/(cos(e + f*x)^2*(a + a/cos(e + f*x))^3),x)`

output `-(c*tan(e/2 + (f*x)/2)^3*(3*tan(e/2 + (f*x)/2)^2 + 5))/(30*a^3*f)`

3.174 $\int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx$

3.174.1 Optimal result	1189
3.174.2 Mathematica [B] (verified)	1189
3.174.3 Rubi [A] (verified)	1190
3.174.4 Maple [F]	1191
3.174.5 Fricas [F]	1192
3.174.6 Sympy [F]	1192
3.174.7 Maxima [F]	1192
3.174.8 Giac [F]	1193
3.174.9 Mupad [F(-1)]	1193

3.174.1 Optimal result

Integrand size = 34, antiderivative size = 140

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx =$$

$$\frac{a^2 c \cos^2(e + fx)^{\frac{3+p}{2}} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{3+p}{2}, \frac{5}{2}, \sin^2(e + fx)\right) (g \sec(e + fx))^p \tan^3(e + fx)}{3f}$$

$$- \frac{a^2 c \cos^2(e + fx)^{\frac{4+p}{2}} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{4+p}{2}, \frac{5}{2}, \sin^2(e + fx)\right) (g \sec(e + fx))^{1+p} \tan^3(e + fx)}{3fg}$$

output `-1/3*a^2*c*(cos(f*x+e)^2)^(3/2+1/2*p)*hypergeom([3/2, 3/2+1/2*p],[5/2],sin(f*x+e)^2)*(g*sec(f*x+e))^p*tan(f*x+e)^3/f-1/3*a^2*c*(cos(f*x+e)^2)^(2+1/2*p)*hypergeom([3/2, 2+1/2*p],[5/2],sin(f*x+e)^2)*(g*sec(f*x+e))^(p+1)*tan(f*x+e)^3/f/g`

3.174.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 325 vs. 2(140) = 280.

Time = 3.02 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.32

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx$$

$$= \frac{a^2 \csc^2\left(\frac{1}{2}(e + fx)\right) \sec^4\left(\frac{1}{2}(e + fx)\right) (g \sec(e + fx))^p (1 + \sec(e + fx))^2 (c - c \sec(e + fx)) \left(-2 \cos^3(e + fx)\right)}{\dots}$$

input `Integrate[(g*Sec[e + f*x])^p*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x]),x]`

output `(a^2*Csc[(e + f*x)/2]^2*Sec[(e + f*x)/2]^4*(g*Sec[e + f*x])^p*(1 + Sec[e + f*x])^2*(c - c*Sec[e + f*x])*(-2*Cos[e + f*x]^3*(Cos[e + f*x]^2)^(p/2)*(2*Hypergeometric2F1[1/2, (2 + p)/2, 3/2, Sin[e + f*x]^2] - Hypergeometric2F1[1/2, (4 + p)/2, 3/2, Sin[e + f*x]^2])*Sin[e + f*x] - ((Sec[e + f*x]^2)^(-1 - p/2)*((4 + p)*Hypergeometric2F1[1/2, 1 - p/2, 3/2, -Tan[e + f*x]^2] - 3*(Sec[e + f*x]^2)^(p/2))*Sin[e + f*x])/(1 + p) - (Hypergeometric2F1[1/2, (2 + p)/2, (4 + p)/2, Sec[e + f*x]^2]*Sin[e + f*x])/((2 + p)*Sqrt[-Tan[e + f*x]^2]) + (2*Cot[e + f*x]*Hypergeometric2F1[1/2, (3 + p)/2, (5 + p)/2, Sec[e + f*x]^2]*Sqrt[-Tan[e + f*x]^2])/(3 + p))/(32*f)`

3.174.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.088$, Rules used = {3042, 4450, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec(e + fx) + a)^2 (c - c \sec(e + fx)) (g \sec(e + fx))^p dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a \right)^2 \left(c - c \csc\left(e + fx + \frac{\pi}{2}\right) \right) \left(g \csc\left(e + fx + \frac{\pi}{2}\right) \right)^p dx \\
 & \quad \downarrow \text{4450} \\
 & -ac \int (a \tan^2(e + fx) (g \sec(e + fx))^p + a \sec(e + fx) \tan^2(e + fx) (g \sec(e + fx))^p) dx \\
 & \quad \downarrow \text{2009} \\
 & -ac \left(\frac{a \tan^3(e + fx) \cos^2(e + fx)^{\frac{p+3}{2}} (g \sec(e + fx))^p \text{Hypergeometric2F1}\left(\frac{3}{2}, \frac{p+3}{2}, \frac{5}{2}, \sin^2(e + fx)\right)}{3f} + \frac{a \tan^3(e + fx)}{3f} \right)
 \end{aligned}$$

input `Int[(g*Sec[e + f*x])^p*(a + a*Sec[e + f*x])^2*(c - c*Sec[e + f*x]),x]`

output $-(a*c*((a*(\cos[e + f*x]^2)^{(3+p)/2})*\text{Hypergeometric2F1}[3/2, (3+p)/2, 5/2, \sin[e + f*x]^2]*(g*\sec[e + f*x])^p*\tan[e + f*x]^3)/(3*f) + (a*(\cos[e + f*x]^2)^{(4+p)/2})*\text{Hypergeometric2F1}[3/2, (4+p)/2, 5/2, \sin[e + f*x]^2]*(g*\sec[e + f*x])^{(1+p)}*\tan[e + f*x]^3)/(3*f*g))$

3.174.3.1 Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4450 $\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(g_.))^{(p_.)}*(\csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^{(m_.)}*(\csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[\text{ExpandTrig}[(g*\csc[e + f*x])^p*\cot[e + f*x]^{(2*m)}, (c + d*\csc[e + f*x])^{(n-m)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, n, p\}, x] \&\& \text{EqQ}[b*c + a*d, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{IntegersQ}[m, n] \&\& \text{GeQ}[n - m, 0] \&\& \text{GtQ}[m*n, 0]$

3.174.4 Maple [F]

$$\int (g \sec(fx + e))^p (a + a \sec(fx + e))^2 (c - c \sec(fx + e)) dx$$

input $\text{int}((g*\sec(f*x+e))^p*(a+a*\sec(f*x+e))^2*(c-c*\sec(f*x+e)),x)$

output $\text{int}((g*\sec(f*x+e))^p*(a+a*\sec(f*x+e))^2*(c-c*\sec(f*x+e)),x)$

3.174.5 Fracas [F]

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx$$

$$= \int -(a \sec(fx + e) + a)^2 (c \sec(fx + e) - c) (g \sec(fx + e))^p dx$$

input `integrate((g*sec(f*x+e))^p*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="fricas")`

output `integral(-(a^2*c*sec(f*x + e)^3 + a^2*c*sec(f*x + e)^2 - a^2*c*sec(f*x + e) - a^2*c)*(g*sec(f*x + e))^p, x)`

3.174.6 Sympy [F]

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx$$

$$= -a^2 c \left(\int -(g \sec(e + fx))^p dx + \int -(g \sec(e + fx))^p \sec(e + fx) dx \right.$$

$$\left. + \int (g \sec(e + fx))^p \sec^2(e + fx) dx + \int (g \sec(e + fx))^p \sec^3(e + fx) dx \right)$$

input `integrate((g*sec(f*x+e))**p*(a+a*sec(f*x+e))**2*(c-c*sec(f*x+e)),x)`

output `-a**2*c*(Integral(-(g*sec(e + f*x))**p, x) + Integral(-(g*sec(e + f*x))**p*sec(e + f*x), x) + Integral((g*sec(e + f*x))**p*sec(e + f*x)**2, x) + Integral((g*sec(e + f*x))**p*sec(e + f*x)**3, x))`

3.174.7 Maxima [F]

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx$$

$$= \int -(a \sec(fx + e) + a)^2 (c \sec(fx + e) - c) (g \sec(fx + e))^p dx$$

input `integrate((g*sec(f*x+e))^p*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-integrate((a*sec(f*x + e) + a)^2*(c*sec(f*x + e) - c)*(g*sec(f*x + e))^p, x)`

3.174.8 Giac [F]

$$\begin{aligned} & \int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx \\ &= \int -(a \sec(fx + e) + a)^2 (c \sec(fx + e) - c) (g \sec(fx + e))^p dx \end{aligned}$$

input `integrate((g*sec(f*x+e))^p*(a+a*sec(f*x+e))^2*(c-c*sec(f*x+e)),x, algorithm="giac")`

output `integrate(-(a*sec(f*x + e) + a)^2*(c*sec(f*x + e) - c)*(g*sec(f*x + e))^p, x)`

3.174.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (g \sec(e + fx))^p (a + a \sec(e + fx))^2 (c - c \sec(e + fx)) dx \\ &= \int \left(a + \frac{a}{\cos(e + fx)} \right)^2 \left(c - \frac{c}{\cos(e + fx)} \right) \left(\frac{g}{\cos(e + fx)} \right)^p dx \end{aligned}$$

input `int((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))*(g/cos(e + f*x))^p,x)`

output `int((a + a/cos(e + f*x))^2*(c - c/cos(e + f*x))*(g/cos(e + f*x))^p, x)`

3.175 $\int (g \sec(e + fx))^p (a + a \sec(e + fx))(c - c \sec(e + fx)) dx$

3.175.1 Optimal result	1194
3.175.2 Mathematica [A] (verified)	1194
3.175.3 Rubi [A] (verified)	1195
3.175.4 Maple [F]	1196
3.175.5 Fricas [F]	1196
3.175.6 Sympy [F]	1197
3.175.7 Maxima [F]	1197
3.175.8 Giac [F]	1198
3.175.9 Mupad [F(-1)]	1198

3.175.1 Optimal result

Integrand size = 32, antiderivative size = 65

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))(c - c \sec(e + fx)) dx = \frac{ac \cos^2(e + fx)^{\frac{3+p}{2}} \text{Hypergeometric2F1}\left(\frac{3}{2}, \frac{3+p}{2}, \frac{5}{2}, \sin^2(e + fx)\right) (g \sec(e + fx))^p \tan^3(e + fx)}{3f}$$

```
output -1/3*a*c*(cos(f*x+e)^2)^(3/2+1/2*p)*hypergeom([3/2, 3/2+1/2*p], [5/2], sin(f*x+e)^2)*(g*sec(f*x+e))^p*tan(f*x+e)^3/f
```

3.175.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))(c - c \sec(e + fx)) dx = - \frac{ac(g \sec(e + fx))^p \tan(e + fx) \left(p + \frac{\text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{p}{2}, \frac{2+p}{2}, \sec^2(e + fx)\right)}{\sqrt{-\tan^2(e + fx)}} \right)}{fp(1 + p)}$$

```
input Integrate[(g*Sec[e + f*x])^p*(a + a*Sec[e + f*x])*(c - c*Sec[e + f*x]),x]
```

output $-\left(\frac{a c (g \sec [e+f x])^p \tan [e+f x] \left(p+\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{p}{2}, 2+\frac{p}{2}, \sec [e+f x]^2\right] / \sqrt{-\tan [e+f x]^2}\right)}{f p(1+p)}\right)$

3.175.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4450, 3042, 3097}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \sec(e+fx) + a)(c - c \sec(e+fx))(g \sec(e+fx))^p dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a \csc\left(e+fx+\frac{\pi}{2}\right) + a \right) \left(c - c \csc\left(e+fx+\frac{\pi}{2}\right) \right) \left(g \csc\left(e+fx+\frac{\pi}{2}\right) \right)^p dx \\ & \quad \downarrow \text{4450} \\ & -ac \int (g \sec(e+fx))^p \tan^2(e+fx) dx \\ & \quad \downarrow \text{3042} \\ & -ac \int (g \sec(e+fx))^p \tan(e+fx)^2 dx \\ & \quad \downarrow \text{3097} \\ & \frac{a c \tan^3(e+fx) \cos^2(e+fx)^{\frac{p+3}{2}} (g \sec(e+fx))^p \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{p+3}{2}, \frac{5}{2}, \sin^2(e+fx)\right)}{3f} \end{aligned}$$

input $\text{Int}[(g \sec [e+f x])^p (a+a \sec [e+f x])(c-c \sec [e+f x]), x]$

output $-\frac{1}{3} a c (\cos [e+f x]^2)^{\frac{3+p}{2}} \operatorname{Hypergeometric2F1}\left[\frac{3}{2}, \frac{3+p}{2}, \frac{5}{2}, \sin [e+f x]^2\right] (g \sec [e+f x])^p \tan [e+f x]^3 / f$

3.175.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3097 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n + 1)*((Cos[e + f*x]^2)^(m + n + 1)/2)/(b*f*(n + 1))*Hypergeometric2F1[(n + 1)/2, (m + n + 1)/2, (n + 3)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[(n - 1)/2] && !IntegerQ[m/2]`

rule 4450 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[((-a)*c)^m Int[ExpandTrig[(g*csc[e + f*x])^p*cot[e + f*x]^(2*m), (c + d*csc[e + f*x])^(n - m), x], x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegersQ[m, n] && GeQ[n - m, 0] && GtQ[m*n, 0]`

3.175.4 Maple [F]

$$\int (g \sec(fx + e))^p (a + a \sec(fx + e)) (c - c \sec(fx + e)) dx$$

input `int((g*sec(f*x+e))^p*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x)`

output `int((g*sec(f*x+e))^p*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x)`

3.175.5 Fracas [F]

$$\begin{aligned} & \int (g \sec(e + fx))^p (a + a \sec(e + fx)) (c - c \sec(e + fx)) dx \\ &= \int -(a \sec(fx + e) + a) (c \sec(fx + e) - c) (g \sec(fx + e))^p dx \end{aligned}$$

input `integrate((g*sec(f*x+e))^p*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="fracas")`

output `integral(-(a*c*sec(f*x + e)^2 - a*c)*(g*sec(f*x + e))^p, x)`

3.175.6 Sympy [F]

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= -ac \left(\int -(g \sec(e + fx))^p dx + \int (g \sec(e + fx))^p \sec^2(e + fx) dx \right)$$

input `integrate((g*sec(f*x+e))**p*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x)`

output `-a*c*(Integral(-(g*sec(e + f*x))**p, x) + Integral((g*sec(e + f*x))**p*sec(e + f*x)**2, x))`

3.175.7 Maxima [F]

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \int -(a \sec(fx + e) + a)(c \sec(fx + e) - c)(g \sec(fx + e))^p dx$$

input `integrate((g*sec(f*x+e))^p*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="maxima")`

output `-integrate((a*sec(f*x + e) + a)*(c*sec(f*x + e) - c)*(g*sec(f*x + e))^p, x)`

3.175.8 Giac [F]

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \int -(a \sec(fx + e) + a)(c \sec(fx + e) - c)(g \sec(fx + e))^p dx$$

input `integrate((g*sec(f*x+e))^p*(a+a*sec(f*x+e))*(c-c*sec(f*x+e)),x, algorithm="giac")`

output `integrate(-(a*sec(f*x + e) + a)*(c*sec(f*x + e) - c)*(g*sec(f*x + e))^p, x)`

3.175.9 Mupad [F(-1)]

Timed out.

$$\int (g \sec(e + fx))^p (a + a \sec(e + fx))(c - c \sec(e + fx)) dx$$

$$= \int \left(a + \frac{a}{\cos(e + fx)} \right) \left(c - \frac{c}{\cos(e + fx)} \right) \left(\frac{g}{\cos(e + fx)} \right)^p dx$$

input `int((a + a/cos(e + f*x))*(c - c/cos(e + f*x))*(g/cos(e + f*x))^p,x)`

output `int((a + a/cos(e + f*x))*(c - c/cos(e + f*x))*(g/cos(e + f*x))^p, x)`

3.176 $\int \frac{(g \sec(e+fx))^p (c-c \sec(e+fx))}{a+a \sec(e+fx)} dx$

3.176.1 Optimal result 1199
 3.176.2 Mathematica [C] (warning: unable to verify) 1200
 3.176.3 Rubi [A] (verified) 1200
 3.176.4 Maple [F] 1203
 3.176.5 Fracas [F] 1203
 3.176.6 Sympy [F] 1203
 3.176.7 Maxima [F] 1204
 3.176.8 Giac [F(-2)] 1204
 3.176.9 Mupad [F(-1)] 1205

3.176.1 Optimal result

Integrand size = 34, antiderivative size = 180

$$\int \frac{(g \sec(e+fx))^p (c-c \sec(e+fx))}{a+a \sec(e+fx)} dx =$$

$$\frac{cg(1-2p) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-p}{2}, \frac{3-p}{2}, \cos^2(e+fx)\right) (g \sec(e+fx))^{-1+p} \sin(e+fx)}{af(1-p)\sqrt{\sin^2(e+fx)}} +$$

$$\frac{2c \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{p}{2}, \frac{2-p}{2}, \cos^2(e+fx)\right) (g \sec(e+fx))^p \sin(e+fx)}{af\sqrt{\sin^2(e+fx)}} -$$

$$\frac{2c(g \sec(e+fx))^p \tan(e+fx)}{f(a+a \sec(e+fx))}$$

output

```
-c*g*(1-2*p)*hypergeom([1/2, 1/2-1/2*p], [3/2-1/2*p], cos(f*x+e)^2)*(g*sec(f*x+e))^(1-p)*sin(f*x+e)/a/f/(1-p)/(sin(f*x+e)^2)^(1/2)+2*c*hypergeom([1/2, -1/2*p], [1-1/2*p], cos(f*x+e)^2)*(g*sec(f*x+e))^p*sin(f*x+e)/a/f/(sin(f*x+e)^2)^(1/2)-2*c*(g*sec(f*x+e))^p*tan(f*x+e)/f/(a+a*sec(f*x+e))
```


3.176.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 14.79 (sec) , antiderivative size = 3396, normalized size of antiderivative = 18.87

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{a + a \sec(e + fx)} dx = \text{Result too large to show}$$

input `Integrate[((g*Sec[e + f*x])^p*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output `(-6*c*Sec[e + f*x]^p*(g*Sec[e + f*x])^p*Tan[(e + f*x)/2]^3*(-((AppellF1[1/2, p, 1 - p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2)/(3*AppellF1[1/2, p, 1 - p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-1 + p)*AppellF1[3/2, p, 2 - p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + p*AppellF1[3/2, 1 + p, 1 - p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2) + AppellF1[1/2, p, -p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]/(3*AppellF1[1/2, p, -p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*p*(AppellF1[3/2, p, 1 - p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + p, -p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2)))/(a*f*(3*Sec[(e + f*x)/2]^2*Sec[e + f*x]^p*(-((AppellF1[1/2, p, 1 - p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2)/(3*AppellF1[1/2, p, 1 - p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-1 + p)*AppellF1[3/2, p, 2 - p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + p*AppellF1[3/2, 1 + p, 1 - p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2) + AppellF1[1/2, p, -p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]/(3*AppellF1[1/2, p, -p, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*p*(AppellF1[3/2, p, 1 - p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + p, -p, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2)) + 6*p*Sec[e + f*x]^(1 + p)*Sin[e + f*x]*Tan[(e + f*...`

3.176.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4508, 3042, 4274, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.176. $\int \frac{(g \sec(e+fx))^p (c-c \sec(e+fx))}{a+a \sec(e+fx)} dx$

$$\begin{aligned}
& \int \frac{(c - c \sec(e + fx))(g \sec(e + fx))^p}{a \sec(e + fx) + a} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(c - c \csc(e + fx + \frac{\pi}{2}))(g \csc(e + fx + \frac{\pi}{2}))^p}{a \csc(e + fx + \frac{\pi}{2}) + a} dx \\
& \quad \downarrow \text{4508} \\
& \frac{\int (g \sec(e + fx))^p (ac(1 - 2p) + 2acp \sec(e + fx)) dx}{a^2} - \frac{2c \tan(e + fx)(g \sec(e + fx))^p}{f(a \sec(e + fx) + a)} \\
& \quad \downarrow \text{3042} \\
& \frac{\int (g \csc(e + fx + \frac{\pi}{2}))^p (ac(1 - 2p) + 2acp \csc(e + fx + \frac{\pi}{2})) dx}{a^2} - \frac{2c \tan(e + fx)(g \sec(e + fx))^p}{f(a \sec(e + fx) + a)} \\
& \quad \downarrow \text{4274} \\
& \frac{ac(1 - 2p) \int (g \sec(e + fx))^p dx + \frac{2acp \int (g \sec(e + fx))^{p+1} dx}{g}}{a^2} - \frac{2c \tan(e + fx)(g \sec(e + fx))^p}{f(a \sec(e + fx) + a)} \\
& \quad \downarrow \text{3042} \\
& \frac{ac(1 - 2p) \int (g \csc(e + fx + \frac{\pi}{2}))^p dx + \frac{2acp \int (g \csc(e + fx + \frac{\pi}{2}))^{p+1} dx}{g}}{a^2} - \frac{2c \tan(e + fx)(g \sec(e + fx))^p}{f(a \sec(e + fx) + a)} \\
& \quad \downarrow \text{4259} \\
& \frac{\frac{2acp \left(\frac{\cos(e + fx)}{g}\right)^p (g \sec(e + fx))^p \int \left(\frac{\cos(e + fx)}{g}\right)^{-p-1} dx}{g} + ac(1 - 2p) \left(\frac{\cos(e + fx)}{g}\right)^p (g \sec(e + fx))^p \int \left(\frac{\cos(e + fx)}{g}\right)^{-p} dx}{a^2} - \frac{2c \tan(e + fx)(g \sec(e + fx))^p}{f(a \sec(e + fx) + a)} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{2acp \left(\frac{\cos(e + fx)}{g}\right)^p (g \sec(e + fx))^p \int \left(\frac{\sin(e + fx + \frac{\pi}{2})}{g}\right)^{-p-1} dx}{g} + ac(1 - 2p) \left(\frac{\cos(e + fx)}{g}\right)^p (g \sec(e + fx))^p \int \left(\frac{\sin(e + fx + \frac{\pi}{2})}{g}\right)^{-p} dx}{a^2} - \frac{2c \tan(e + fx)(g \sec(e + fx))^p}{f(a \sec(e + fx) + a)} \\
& \quad \downarrow \text{3122}
\end{aligned}$$

3.176. $\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{a + a \sec(e + fx)} dx$

$$\frac{2ac \sin(e+fx)(g \sec(e+fx))^p \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{p}{2}, \frac{2-p}{2}, \cos^2(e+fx)\right)}{f \sqrt{\sin^2(e+fx)}} - \frac{acg(1-2p) \sin(e+fx)(g \sec(e+fx))^{p-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-p}{2}, \frac{3-p}{2}, \cos^2(e+fx)\right)}{f(1-p) \sqrt{\sin^2(e+fx)}}$$

$$\frac{2c \tan(e+fx)(g \sec(e+fx))^p}{f(a \sec(e+fx) + a)}$$

input `Int[((g*Sec[e + f*x])^p*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output `((-(a*c*g*(1 - 2*p)*Hypergeometric2F1[1/2, (1 - p)/2, (3 - p)/2, Cos[e + f*x]^2]*(g*Sec[e + f*x])^(-1 + p)*Sin[e + f*x])/(f*(1 - p)*Sqrt[Sin[e + f*x]^2])) + (2*a*c*Hypergeometric2F1[1/2, -1/2*p, (2 - p)/2, Cos[e + f*x]^2]*(g*Sec[e + f*x])^p*Ssin[e + f*x])/(f*Sqrt[Sin[e + f*x]^2]))/a^2 - (2*c*(g*Sec[e + f*x])^p*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))`

3.176.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4508 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.176.4 Maple [F]

$$\int \frac{(g \sec(fx + e))^p (c - c \sec(fx + e))}{a + a \sec(fx + e)} dx$$

input `int((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x)`

output `int((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x)`

3.176.5 Fricas [F]

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{a + a \sec(e + fx)} dx = \int -\frac{(c \sec(fx + e) - c)(g \sec(fx + e))^p}{a \sec(fx + e) + a} dx$$

input `integrate((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="fricas")`

output `integral(-(c*sec(f*x + e) - c)*(g*sec(f*x + e))^p/(a*sec(f*x + e) + a), x)`

3.176.6 Sympy [F]

$$\begin{aligned} & \int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{a + a \sec(e + fx)} dx \\ &= -\frac{c \left(\int \left(-\frac{(g \sec(e + fx))^p}{\sec(e + fx) + 1} \right) dx + \int \frac{(g \sec(e + fx))^p \sec(e + fx)}{\sec(e + fx) + 1} dx \right)}{a} \end{aligned}$$

input `integrate((g*sec(f*x+e))**p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x)`

output `-c*(Integral(-(g*sec(e + f*x))**p/(sec(e + f*x) + 1), x) + Integral((g*sec(e + f*x))**p*sec(e + f*x)/(sec(e + f*x) + 1), x))/a`

3.176.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{a + a \sec(e + fx)} dx = \int -\frac{(c \sec(fx + e) - c)(g \sec(fx + e))^p}{a \sec(fx + e) + a} dx$$

input `integrate((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `-integrate((c*sec(f*x + e) - c)*(g*sec(f*x + e))^p/(a*sec(f*x + e) + a), x)`

3.176.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{a + a \sec(e + fx)} dx = \text{Exception raised: TypeError}$$

input `integrate((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0,1,2,0]%%}+%%{1,[0,1,0,0]%%} / %%{2,[0,0,0,1]%%}Error: Ba`

3.176.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{a + a \sec(e + fx)} dx = \int \frac{\left(c - \frac{c}{\cos(e + fx)}\right) \left(\frac{g}{\cos(e + fx)}\right)^p}{a + \frac{a}{\cos(e + fx)}} dx$$

input `int(((c - c/cos(e + f*x))*(g/cos(e + f*x))^p)/(a + a/cos(e + f*x)),x)`

output `int(((c - c/cos(e + f*x))*(g/cos(e + f*x))^p)/(a + a/cos(e + f*x)), x)`

$$3.177 \quad \int \frac{(g \sec(e+fx))^p (c - c \sec(e+fx))}{(a+a \sec(e+fx))^2} dx$$

3.177.1 Optimal result 1206
 3.177.2 Mathematica [A] (verified) 1207
 3.177.3 Rubi [A] (verified) 1207
 3.177.4 Maple [F] 1210
 3.177.5 Fricas [F] 1210
 3.177.6 Sympy [F] 1211
 3.177.7 Maxima [F] 1211
 3.177.8 Giac [F(-2)] 1211
 3.177.9 Mupad [F(-1)] 1212

3.177.1 Optimal result

Integrand size = 34, antiderivative size = 226

$$\int \frac{(g \sec(e+fx))^p (c - c \sec(e+fx))}{(a+a \sec(e+fx))^2} dx =$$

$$\frac{cg(3-4p) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-p}{2}, \frac{3-p}{2}, \cos^2(e+fx)\right) (g \sec(e+fx))^{-1+p} \sin(e+fx)}{3a^2 f \sqrt{\sin^2(e+fx)}} +$$

$$\frac{c(5-4p) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{p}{2}, \frac{2-p}{2}, \cos^2(e+fx)\right) (g \sec(e+fx))^p \sin(e+fx)}{3a^2 f \sqrt{\sin^2(e+fx)}} -$$

$$\frac{c(5-4p)(g \sec(e+fx))^p \tan(e+fx)}{3a^2 f (1+\sec(e+fx))} - \frac{2c(g \sec(e+fx))^p \tan(e+fx)}{3f(a+a \sec(e+fx))^2}$$

output

```
-1/3*c*g*(3-4*p)*hypergeom([1/2, 1/2-1/2*p], [3/2-1/2*p], cos(f*x+e)^2)*(g*sec(f*x+e))^(1-p)*sin(f*x+e)/a^2/f/(sin(f*x+e)^2)^(1/2)+1/3*c*(5-4*p)*hypergeom([1/2, -1/2*p], [1-1/2*p], cos(f*x+e)^2)*(g*sec(f*x+e))^p*sin(f*x+e)/a^2/f/(sin(f*x+e)^2)^(1/2)-1/3*c*(5-4*p)*(g*sec(f*x+e))^p*tan(f*x+e)/a^2/f/(1+sec(f*x+e))-2/3*c*(g*sec(f*x+e))^p*tan(f*x+e)/f/(a+a*sec(f*x+e))^2
```

3.177.2 Mathematica [A] (verified)

Time = 1.99 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.83

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx =$$

$$\frac{c(g \sec(e + fx))^p \left(2p(1 + p) \tan(e + fx) + (1 + \sec(e + fx)) \left(-p(1 + p)(-5 + 4p) \tan(e + fx) - \left((-1 + p)(1 + p)(-3 + 4p) \cot(e + fx) \right) \right) \right)}{(a + a \sec(e + fx))^2}$$

input `Integrate[((g*Sec[e + f*x])^p*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2, x]`

output `-1/3*(c*(g*Sec[e + f*x])^p*(2*p*(1 + p)*Tan[e + f*x] + (1 + Sec[e + f*x])*(-(p*(1 + p)*(-5 + 4*p)*Tan[e + f*x] - ((-1 + p)*(1 + p)*(-3 + 4*p)*Cot[e + f*x]*Hypergeometric2F1[1/2, p/2, (2 + p)/2, Sec[e + f*x]^2] + (5 - 4*p)*p^2*Csc[e + f*x]*Hypergeometric2F1[1/2, (1 + p)/2, (3 + p)/2, Sec[e + f*x]^2]))*(1 + Sec[e + f*x])*Sqrt[-Tan[e + f*x]^2]))/(a^2*f*p*(1 + p)*(1 + Sec[e + f*x])^2)`

3.177.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4508, 3042, 4508, 3042, 4274, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - c \sec(e + fx))(g \sec(e + fx))^p}{(a \sec(e + fx) + a)^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c - c \csc(e + fx + \frac{\pi}{2}))(g \csc(e + fx + \frac{\pi}{2}))^p}{(a \csc(e + fx + \frac{\pi}{2}) + a)^2} dx$$

$$\downarrow \text{4508}$$

$$\frac{\int \frac{(g \sec(e + fx))^p (ac(3 - 2p) - 2ac(1 - p) \sec(e + fx))}{\sec(e + fx)a + a} dx}{3a^2} - \frac{2c \tan(e + fx)(g \sec(e + fx))^p}{3f(a \sec(e + fx) + a)^2}$$

$$\downarrow \text{3042}$$

3.177. $\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx$

$$\frac{\int \frac{(g \csc(e+fx+\frac{\pi}{2}))^p (ac(3-2p)-2ac(1-p) \csc(e+fx+\frac{\pi}{2}))}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a^2} - \frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx)+a)^2}$$

↓ 4508

$$\frac{\int (g \sec(e+fx))^p (c(3-4p)(1-p)a^2+c(5-4p)p \sec(e+fx)a^2) dx}{a^2} - \frac{c(5-4p) \tan(e+fx)(g \sec(e+fx))^p}{f(\sec(e+fx)+1)}$$

$$\frac{3a^2}{3f(a \sec(e+fx)+a)^2} \frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx)+a)^2}$$

↓ 3042

$$\frac{\int (g \csc(e+fx+\frac{\pi}{2}))^p (c(3-4p)(1-p)a^2+c(5-4p)p \csc(e+fx+\frac{\pi}{2})a^2) dx}{a^2} - \frac{c(5-4p) \tan(e+fx)(g \sec(e+fx))^p}{f(\sec(e+fx)+1)}$$

$$\frac{3a^2}{3f(a \sec(e+fx)+a)^2} \frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx)+a)^2}$$

↓ 4274

$$\frac{a^2c(3-4p)(1-p) \int (g \sec(e+fx))^p dx + \frac{a^2c(5-4p)p}{g} \int (g \sec(e+fx))^{p+1} dx}{a^2} - \frac{c(5-4p) \tan(e+fx)(g \sec(e+fx))^p}{f(\sec(e+fx)+1)}$$

$$\frac{3a^2}{3f(a \sec(e+fx)+a)^2} \frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx)+a)^2}$$

↓ 3042

$$\frac{a^2c(3-4p)(1-p) \int (g \csc(e+fx+\frac{\pi}{2}))^p dx + \frac{a^2c(5-4p)p}{g} \int (g \csc(e+fx+\frac{\pi}{2}))^{p+1} dx}{a^2} - \frac{c(5-4p) \tan(e+fx)(g \sec(e+fx))^p}{f(\sec(e+fx)+1)}$$

$$\frac{3a^2}{3f(a \sec(e+fx)+a)^2} \frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx)+a)^2}$$

↓ 4259

$$\frac{a^2c(5-4p)p \left(\frac{\cos(e+fx)}{g}\right)^p (g \sec(e+fx))^p \int \left(\frac{\cos(e+fx)}{g}\right)^{-p-1} dx + a^2c(3-4p)(1-p) \left(\frac{\cos(e+fx)}{g}\right)^p (g \sec(e+fx))^p \int \left(\frac{\cos(e+fx)}{g}\right)^{-p} dx}{a^2} - \frac{c(5-4p) \tan(e+fx)(g \sec(e+fx))^p}{f(\sec(e+fx)+1)}$$

$$\frac{3a^2}{3f(a \sec(e+fx)+a)^2} \frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx)+a)^2}$$

↓ 3042

3.177. $\int \frac{(g \sec(e+fx))^p (c-c \sec(e+fx))}{(a+a \sec(e+fx))^2} dx$

$$\frac{a^2 c(5-4p) \left(\frac{\cos(e+fx)}{g}\right)^p (g \sec(e+fx))^p \int \left(\frac{\sin\left(e+fx+\frac{\pi}{2}\right)}{g}\right)^{-p-1} dx}{g} + \frac{a^2 c(3-4p)(1-p) \left(\frac{\cos(e+fx)}{g}\right)^p (g \sec(e+fx))^p \int \left(\frac{\sin\left(e+fx+\frac{\pi}{2}\right)}{g}\right)^{-p} dx}{a^2} - \frac{c(5-4p) \left(\frac{\cos(e+fx)}{g}\right)^p (g \sec(e+fx))^p \int \left(\frac{\sin\left(e+fx+\frac{\pi}{2}\right)}{g}\right)^{-p-1} dx}{a^2}$$

$$\frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx) + a)^2} \quad 3a^2$$

↓ 3122

$$\frac{a^2 c(5-4p) \sin(e+fx)(g \sec(e+fx))^p \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{p}{2}, \frac{2-p}{2}, \cos^2(e+fx)\right)}{f \sqrt{\sin^2(e+fx)}} - \frac{a^2 c g(3-4p) \sin(e+fx)(g \sec(e+fx))^{p-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-p}{2}, \frac{3-p}{2}, \cos^2(e+fx)\right)}{f \sqrt{\sin^2(e+fx)}} - \frac{c(5-4p) \sin(e+fx)(g \sec(e+fx))^p \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{p}{2}, \frac{2-p}{2}, \cos^2(e+fx)\right)}{a^2}$$

$$\frac{2c \tan(e+fx)(g \sec(e+fx))^p}{3f(a \sec(e+fx) + a)^2} \quad 3a^2$$

input `Int[((g*Sec[e + f*x])^p*(c - c*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2,x]`

output `(-2*c*(g*Sec[e + f*x])^p*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) + ((-(a^2*c*g*(3 - 4*p)*Hypergeometric2F1[1/2, (1 - p)/2, (3 - p)/2, Cos[e + f*x]^2]*(g*Sec[e + f*x])^(-1 + p)*Sin[e + f*x])/(f*Sqrt[Sin[e + f*x]^2])) + (a^2*c*(5 - 4*p)*Hypergeometric2F1[1/2, -1/2*p, (2 - p)/2, Cos[e + f*x]^2]*(g*Sec[e + f*x])^p*Ssin[e + f*x])/(f*Sqrt[Sin[e + f*x]^2]))/a^2 - (c*(5 - 4*p)*(g*Sec[e + f*x])^p*Tan[e + f*x])/(f*(1 + Sec[e + f*x]))/(3*a^2)`

3.177.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.177. $\int \frac{(g \sec(e+fx))^p (c - c \sec(e+fx))}{(a + a \sec(e+fx))^2} dx$

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4508 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.177.4 Maple [F]

$$\int \frac{(g \sec(fx + e))^p (c - c \sec(fx + e))}{(a + a \sec(fx + e))^2} dx$$

input `int((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x)`

output `int((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x)`

3.177.5 Fracas [F]

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx = \int -\frac{(c \sec(fx + e) - c)(g \sec(fx + e))^p}{(a \sec(fx + e) + a)^2} dx$$

input `integrate((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="fricas")`

output `integral(-(c*sec(f*x + e) - c)*(g*sec(f*x + e))^p/(a^2*sec(f*x + e)^2 + 2*a^2*sec(f*x + e) + a^2), x)`

3.177.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx$$

$$= \frac{c \left(\int \left(-\frac{(g \sec(e + fx))^p}{\sec^2(e + fx) + 2 \sec(e + fx) + 1} \right) dx + \int \frac{(g \sec(e + fx))^p \sec(e + fx)}{\sec^2(e + fx) + 2 \sec(e + fx) + 1} dx \right)}{a^2}$$

input `integrate((g*sec(f*x+e))**p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**2,x)`

output `-c*(Integral(-(g*sec(e + f*x))**p/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral((g*sec(e + f*x))**p*sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.177.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx = \int -\frac{(c \sec(fx + e) - c)(g \sec(fx + e))^p}{(a \sec(fx + e) + a)^2} dx$$

input `integrate((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output `-integrate((c*sec(f*x + e) - c)*(g*sec(f*x + e))^p/(a*sec(f*x + e) + a)^2, x)`

3.177.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate((g*sec(f*x+e))^p*(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN
 PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
 unding error%%{-1,[0,1,4,0]%%}+%%{1,[0,1,0,0]%%} / %%{4,[0,0,0,2]%%}
 Error: B

3.177.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^p (c - c \sec(e + fx))}{(a + a \sec(e + fx))^2} dx = \int \frac{\left(c - \frac{c}{\cos(e + fx)}\right) \left(\frac{g}{\cos(e + fx)}\right)^p}{\left(a + \frac{a}{\cos(e + fx)}\right)^2} dx$$

input `int(((c - c/cos(e + f*x))*(g/cos(e + f*x))^p)/(a + a/cos(e + f*x))^2,x)`

output `int(((c - c/cos(e + f*x))*(g/cos(e + f*x))^p)/(a + a/cos(e + f*x))^2, x)`

3.178
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c-c \sec(e+fx)} dx$$

3.178.1 Optimal result 1213
 3.178.2 Mathematica [A] (verified) 1213
 3.178.3 Rubi [A] (verified) 1214
 3.178.4 Maple [B] (verified) 1216
 3.178.5 Fricas [A] (verification not implemented) 1217
 3.178.6 Sympy [F] 1217
 3.178.7 Maxima [B] (verification not implemented) 1218
 3.178.8 Giac [F] 1218
 3.178.9 Mupad [F(-1)] 1219

3.178.1 Optimal result

Integrand size = 40, antiderivative size = 104

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c-c \sec(e+fx)} dx =$$

$$\frac{2\sqrt{a}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{g \sec(e+fx)} \sqrt{a+a \sec(e+fx)}}\right)}{cf}$$

$$+ \frac{2g \cot(e+fx) \sqrt{g \sec(e+fx)} \sqrt{a+a \sec(e+fx)}}{cf}$$

output `-2*g^(3/2)*arctanh(a^(1/2)*g^(1/2)*tan(f*x+e)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))*a^(1/2)/c/f+2*g*cot(f*x+e)*(g*sec(f*x+e))^(1/2)*(a+a*sec(f*x+e))^(1/2)/c/f`

3.178.2 Mathematica [A] (verified)

Time = 4.33 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.56

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c-c \sec(e+fx)} dx = \frac{2 \cot\left(\frac{1}{2}(e+fx)\right) (g \sec(e+fx))^{3/2} \sqrt{a(1+\sec(e+fx))}}{\sqrt{a+a \sec(e+fx)}} \left(\sqrt{\frac{a+a \sec(e+fx)}{a(1+\sec(e+fx))}}\right)$$

input `Integrate[((g*Sec[e + f*x])^(3/2)*Sqrt[a + a*Sec[e + f*x]])/(c - c*Sec[e + f*x]),x]`

output $(2*\text{Cot}[(e + f*x)/2]*(g*\text{Sec}[e + f*x])^{3/2}*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]*(\text{Sqrt}[\text{Sec}[e + f*x]]*\text{Sqrt}[1 + \text{Sec}[e + f*x]] + (\text{Log}[1 + \text{Sec}[e + f*x]] - \text{Log}[\text{Sqrt}[\text{Sec}[e + f*x]] + \text{Sec}[e + f*x]^{3/2} + \text{Sqrt}[1 + \text{Sec}[e + f*x]]*\text{Sqrt}[\text{Tan}[e + f*x]^2])*\text{Sqrt}[\text{Tan}[e + f*x]^2]))/(c*f*\text{Sec}[e + f*x]^{3/2}*(1 + \text{Sec}[e + f*x])^{3/2})$

3.178.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.22, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4452, 57, 65, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{a \sec(e + fx) + a}(g \sec(e + fx))^{3/2}}{c - c \sec(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a}(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{c - c \csc(e + fx + \frac{\pi}{2})} dx \\ & \quad \downarrow \text{4452} \\ & \frac{acg \tan(e + fx) \int \frac{\sqrt{g \sec(e + fx)}}{(c - c \sec(e + fx))^{3/2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}} \\ & \quad \downarrow \text{57} \\ & \frac{acg \tan(e + fx) \left(\frac{2\sqrt{g \sec(e + fx)}}{c\sqrt{c - c \sec(e + fx)}} - \frac{g \int \frac{1}{\sqrt{g \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} d \sec(e + fx)}{c} \right)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}} \\ & \quad \downarrow \text{65} \\ & \frac{acg \tan(e + fx) \left(\frac{2\sqrt{g \sec(e + fx)}}{c\sqrt{c - c \sec(e + fx)}} - \frac{2g \int \frac{1}{\frac{c \sec(e + fx)g}{c - c \sec(e + fx)} + g} d \frac{\sqrt{g \sec(e + fx)}}{\sqrt{c - c \sec(e + fx)}}}{c} \right)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}} \\ & \quad \downarrow \text{218} \end{aligned}$$

3.178. $\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx$

$$\frac{acg \tan(e + fx) \left(\frac{2\sqrt{g \sec(e+fx)}}{c\sqrt{c-c\sec(e+fx)}} - \frac{2\sqrt{g} \arctan\left(\frac{\sqrt{c}\sqrt{g \sec(e+fx)}}{\sqrt{g}\sqrt{c-c\sec(e+fx)}}\right)}{c^{3/2}} \right)}{f\sqrt{a \sec(e + fx) + a}\sqrt{c - c\sec(e + fx)}}$$

input `Int[((g*Sec[e + f*x])^(3/2)*Sqrt[a + a*Sec[e + f*x]])/(c - c*Sec[e + f*x]),x]`

output `-((a*c*g*((-2*Sqrt[g]*ArcTan[(Sqrt[c]*Sqrt[g*Sec[e + f*x]])/(Sqrt[g]*Sqrt[c - c*Sec[e + f*x]])])/c^(3/2) + (2*Sqrt[g*Sec[e + f*x]])/(c*Sqrt[c - c*Sec[e + f*x]]))*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]))`

3.178.3.1 Defintions of rubi rules used

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4452 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) +
(a_)^(m_)*(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a
*c*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]))
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Cs
c[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && EqQ[b*c + a*
d, 0] && EqQ[a^2 - b^2, 0]
```

3.178.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(88) = 176.

Time = 5.28 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.82

method	result
default	$\frac{g \left(\operatorname{arctanh} \left(\frac{\cos(fx+e) - \sin(fx+e) + 1}{2(\cos(fx+e)+1)\sqrt{\frac{1}{\cos(fx+e)+1}}} \right) \sin(fx+e) - \operatorname{arctanh} \left(\frac{\cos(fx+e) + \sin(fx+e) + 1}{2(\cos(fx+e)+1)\sqrt{\frac{1}{\cos(fx+e)+1}}} \right) \sin(fx+e) + 2\sqrt{\frac{1}{\cos(fx+e)+1}} \cos(fx+e) \right)}{cf(\cos(fx+e)+1)\sqrt{\frac{1}{\cos(fx+e)+1}}}$

```
input int((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e)),x,method=
_RETURNVERBOSE)
```

```
output g/c/f*(arctanh(1/2*(cos(f*x+e)-sin(f*x+e)+1)/(cos(f*x+e)+1)/(1/(cos(f*x+e)
+1))^(1/2))*sin(f*x+e)-arctanh(1/2*(cos(f*x+e)+sin(f*x+e)+1)/(cos(f*x+e)+1
)/(1/(cos(f*x+e)+1))^(1/2))*sin(f*x+e)+2*(1/(cos(f*x+e)+1))^(1/2)*cos(f*x+
e)+2*(1/(cos(f*x+e)+1))^(1/2))*(a*(sec(f*x+e)+1))^(1/2)*(g*sec(f*x+e))^(1/
2)/(cos(f*x+e)+1)/(1/(cos(f*x+e)+1))^(1/2)*cot(f*x+e)
```

3.178. $\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c-c \sec(e+fx)} dx$

3.178.5 Fricas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 340, normalized size of antiderivative = 3.27

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx = \frac{\sqrt{agg} \log \left(\frac{ag \cos(fx+e)^3 - 7 ag \cos(fx+e)^2 + 4 \sqrt{ag} (\cos(fx+e)^2 - 2 \cos(fx+e) + 1)}{\cos(fx+e)^3 + \cos(fx+e)} \right) + \sqrt{-agg} \arctan \left(\frac{2 \sqrt{-ag} \sqrt{\frac{a \cos(fx+e) + a}{\cos(fx+e)}} \sqrt{\frac{g}{\cos(fx+e)}} \cos(fx+e) \sin(fx+e)}{ag \cos(fx+e)^2 - ag \cos(fx+e) - 2 ag} \right) \sin(fx+e) - 2 g \sqrt{\frac{a \cos(fx+e) + a}{\cos(fx+e)}} \sqrt{\frac{g}{\cos(fx+e)}}}{cf \sin(fx+e)}$$

input `integrate((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e)),x,
algorithm="fricas")`

output `[1/2*(sqrt(a*g)*g*log((a*g*cos(f*x + e)^3 - 7*a*g*cos(f*x + e)^2 + 4*sqrt(a*g)*(cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) + 8*a*g)/(cos(f*x + e)^3 + cos(f*x + e)^2))*sin(f*x + e) + 4*g*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e))/(c*f*sin(f*x + e)), -(sqrt(-a*g)*g*arctan(2*sqrt(-a*g)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)/(a*g*cos(f*x + e)^2 - a*g*cos(f*x + e) - 2*a*g))*sin(f*x + e) - 2*g*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e))/(c*f*sin(f*x + e))]`

3.178.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx = - \frac{\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a \sec(e + fx) + a}}{\sec(e + fx) - 1} dx}{c}$$

input `integrate((g*sec(f*x+e))**(3/2)*(a+a*sec(f*x+e))**(1/2)/(c-c*sec(f*x+e)),x)`

output `-Integral((g*sec(e + f*x))**(3/2)*sqrt(a*sec(e + f*x) + a)/(sec(e + f*x) - 1), x)/c`

3.178. $\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx$

3.178.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 979 vs. 2(88) = 176.

Time = 0.45 (sec) , antiderivative size = 979, normalized size of antiderivative = 9.41

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx = \text{Too large to display}$$

```
input integrate((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e)),x,
algorithm="maxima")
```

```
output 1/2*(4*sqrt(2)*g*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*sin(
1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 4*sqrt(2)*g*cos(1/2*arc
tan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*sin(1/4*arctan2(sin(2*f*x + 2*e)
, cos(2*f*x + 2*e))) + 4*sqrt(2)*g*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2
*f*x + 2*e))) - (g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2
+ g*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*g*cos(1/2*a
rctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + g)*log(2*cos(1/4*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e)))^2 + 2*sqrt(2)*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e))) + 2*sqrt(2)*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)
)) + 2) + (g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + g*si
n(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*g*cos(1/2*arctan2
(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + g)*log(2*cos(1/4*arctan2(sin(2*f*x
+ 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*
f*x + 2*e)))^2 + 2*sqrt(2)*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2
*e))) - 2*sqrt(2)*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2
) - (g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + g*sin(1/2*
arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*g*cos(1/2*arctan2(sin(2
*f*x + 2*e), cos(2*f*x + 2*e))) + g)*log(2*cos(1/4*arctan2(sin(2*f*x + 2*e
), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x...
```

3.178.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx = \int -\frac{\sqrt{a \sec(fx + e) + a} (g \sec(fx + e))^{3/2}}{c \sec(fx + e) - c} dx$$

```
input integrate((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e)),x,
algorithm="giac")
```

3.178. $\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx$

output `sage0*x`

3.178.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c - c \sec(e + fx)} dx = \int \frac{\sqrt{a + \frac{a}{\cos(e + fx)}} \left(\frac{g}{\cos(e + fx)}\right)^{3/2}}{c - \frac{c}{\cos(e + fx)}} dx$$

input `int(((a + a/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c - c/cos(e + f*x))),x)`

output `int(((a + a/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c - c/cos(e + f*x))), x)`

3.179
$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$$

3.179.1 Optimal result 1220
 3.179.2 Mathematica [A] (verified) 1220
 3.179.3 Rubi [A] (verified) 1221
 3.179.4 Maple [A] (verified) 1223
 3.179.5 Fricas [A] (verification not implemented) 1223
 3.179.6 Sympy [F] 1224
 3.179.7 Maxima [F] 1224
 3.179.8 Giac [A] (verification not implemented) 1225
 3.179.9 Mupad [F(-1)] 1225

3.179.1 Optimal result

Integrand size = 36, antiderivative size = 81

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx = -\frac{\arctan\left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{2}\sqrt{acf}} + \frac{\cot(e+fx)\sqrt{a+a \sec(e+fx)}}{acf}$$

output `-1/2*arctan(1/2*a^(1/2)*tan(f*x+e)*2^(1/2)/(a+a*sec(f*x+e))^(1/2))/c/f*2^(1/2)/a^(1/2)+cot(f*x+e)*(a+a*sec(f*x+e))^(1/2)/a/c/f`

3.179.2 Mathematica [A] (verified)

Time = 1.68 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.90

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx = -\frac{\cot\left(\frac{1}{2}(e+fx)\right)\left(-2+\sqrt{2}\arctan\left(\frac{\sqrt{-1+\sec(e+fx)}}{\sqrt{2}}\right)\sqrt{-1+\sec(e+fx)}\right)}{2cf\sqrt{a(1+\sec(e+fx))}}$$

input `Integrate[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output
$$\frac{-1/2*(\text{Cot}[(e + f*x)/2]*(-2 + \text{Sqrt}[2]*\text{ArcTan}[\text{Sqrt}[-1 + \text{Sec}[e + f*x]]/\text{Sqrt}[2]]*\text{Sqrt}[-1 + \text{Sec}[e + f*x]]))/(c*f*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x]]))}{}$$

3.179.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.22, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4452, 27, 87, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^2(e + fx)}{\sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e + fx + \frac{\pi}{2})^2}{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a(c - c \csc(e + fx + \frac{\pi}{2}))}} dx \\ & \quad \downarrow \text{4452} \\ & \frac{a \tan(e + fx) \int \frac{\sec(e + fx)}{a(\sec(e + fx) + 1)(c - c \sec(e + fx))^{3/2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} \\ & \quad \downarrow \text{27} \\ & \frac{c \tan(e + fx) \int \frac{\sec(e + fx)}{(\sec(e + fx) + 1)(c - c \sec(e + fx))^{3/2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} \\ & \quad \downarrow \text{87} \\ & \frac{c \tan(e + fx) \left(\frac{1}{c \sqrt{c - c \sec(e + fx)}} - \frac{\int \frac{1}{(\sec(e + fx) + 1) \sqrt{c - c \sec(e + fx)}} d \sec(e + fx)}{2c} \right)}{f \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} \\ & \quad \downarrow \text{73} \\ & \frac{c \tan(e + fx) \left(\frac{\int \frac{1}{2 - \frac{c - c \sec(e + fx)}{c}} d \sqrt{c - c \sec(e + fx)}}{c^2} + \frac{1}{c \sqrt{c - c \sec(e + fx)}} \right)}{f \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} \\ & \quad \downarrow \text{219} \end{aligned}$$

3.179.
$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)(c - c \sec(e + fx))}} dx$$

$$\frac{c \tan(e + fx) \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c - c \sec(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} + \frac{1}{c\sqrt{c - c \sec(e + fx)}} \right)}{f\sqrt{a \sec(e + fx) + a}\sqrt{c - c \sec(e + fx)}}$$

input `Int[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output `-((c*(ArcTanh[Sqrt[c - c*Sec[e + f*x]]/(Sqrt[2]*Sqrt[c]])/(Sqrt[2]*c^(3/2)) + 1/(c*Sqrt[c - c*Sec[e + f*x]]))*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]))`

3.179.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.179. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$

```
rule 4452 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.)^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[a
*c*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]))
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Cs
c[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && EqQ[b*c + a*
d, 0] && EqQ[a^2 - b^2, 0]
```

3.179.4 Maple [A] (verified)

Time = 2.40 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.32

method	result
default	$-\frac{\sqrt{a(\sec(fx+e)+1)} \left(\sqrt{2} \sqrt{-\frac{\cos(fx+e)}{\cos(fx+e)+1}} \ln \left(\csc(fx+e) - \cot(fx+e) + \sqrt{\cot(fx+e)^2 - 2 \csc(fx+e) \cot(fx+e) + \csc(fx+e)^2 - 1} \right) - 1 \right)}{2cfa}$

```
input int(sec(f*x+e)^2/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/2/c/f/a*(a*(sec(f*x+e)+1))^(1/2)*(2^(1/2)*(-cos(f*x+e)/(cos(f*x+e)+1))^(
1/2)*ln(csc(f*x+e)-cot(f*x+e)+(cot(f*x+e)^2-2*csc(f*x+e)*cot(f*x+e)+csc(f
*x+e)^2-1)^(1/2))-2*cot(f*x+e))
```

3.179.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 260, normalized size of antiderivative = 3.21

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx$$

$$= \frac{\left[\sqrt{2a} \sqrt{-\frac{1}{a}} \log \left(\frac{2\sqrt{2} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}} \sqrt{-\frac{1}{a}} \cos(fx+e) \sin(fx+e) + 3 \cos(fx+e)^2 + 2 \cos(fx+e) - 1}{\cos(fx+e)^2 + 2 \cos(fx+e) + 1} \right) \sin(fx + e) + 4 \sqrt{\frac{a \cos(fx+e)}{\cos(fx+e)}} \right]}{4acf \sin(fx + e)}$$

```
input integrate(sec(f*x+e)^2/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorith
m="fricas")
```

3.179. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$

output `[1/4*(sqrt(2)*a*sqrt(-1/a)*log((2*sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(-1/a)*cos(f*x + e)*sin(f*x + e) + 3*cos(f*x + e)^2 + 2*cos(f*x + e) - 1)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1))*sin(f*x + e) + 4*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*cos(f*x + e))/(a*c*f*sin(f*x + e)), 1/2*(sqrt(2)*sqrt(a)*arctan(sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*cos(f*x + e)/(sqrt(a)*sin(f*x + e)))*sin(f*x + e) + 2*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*cos(f*x + e))/(a*c*f*sin(f*x + e))]`

3.179.6 Sympy [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = -\frac{\int \frac{\sec^2(e + fx)}{\sqrt{a \sec(e + fx) + a} \sec(e + fx) - \sqrt{a \sec(e + fx) + a}} dx}{c}$$

input `integrate(sec(f*x+e)**2/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x)`

output `-Integral(sec(e + f*x)**2/(sqrt(a*sec(e + f*x) + a)*sec(e + f*x) - sqrt(a*sec(e + f*x) + a)), x)/c`

3.179.7 Maxima [F]

$$\begin{aligned} & \int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx \\ &= \int -\frac{\sec^2(fx + e)}{\sqrt{a \sec(fx + e) + a}(c \sec(fx + e) - c)} dx \end{aligned}$$

input `integrate(sec(f*x+e)^2/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `-integrate(sec(f*x + e)^2/(sqrt(a*sec(f*x + e) + a)*(c*sec(f*x + e) - c)), x)`

3.179.8 Giac [A] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.63

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

$$= \frac{\sqrt{2} \log\left(\left(\sqrt{-a} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - \sqrt{-a \tan^2\left(\frac{1}{2}fx + \frac{1}{2}e\right) + a}\right)^2\right)}{\sqrt{-a} \operatorname{sgn}(\cos(fx+e))} - \frac{4\sqrt{2}\sqrt{-a}}{\left(\left(\sqrt{-a} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - \sqrt{-a \tan^2\left(\frac{1}{2}fx + \frac{1}{2}e\right) + a}\right)^2 - a\right) \operatorname{sgn}(\cos(fx+e))}$$

input `integrate(sec(f*x+e)^2/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm m="giac")`

output `1/4*(sqrt(2)*log((sqrt(-a)*tan(1/2*f*x + 1/2*e) - sqrt(-a*tan(1/2*f*x + 1/2*e)^2 + a))^2)/(sqrt(-a)*c*sgn(cos(f*x + e)))) - 4*sqrt(2)*sqrt(-a)/(((sqrt(-a)*tan(1/2*f*x + 1/2*e) - sqrt(-a*tan(1/2*f*x + 1/2*e)^2 + a))^2 - a)*c*sgn(cos(f*x + e))))/f`

3.179.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

$$= - \int \frac{1}{\sqrt{a + \frac{a}{\cos(e+fx)}} \left(c \cos(e+fx) - c \left(\frac{\cos(2e+2fx)}{2} + \frac{1}{2} \right) \right)} dx$$

input `int(1/(cos(e + f*x)^2*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))),x)`

output `-int(1/((a + a/cos(e + f*x))^(1/2)*(c*cos(e + f*x) - c*(cos(2*e + 2*f*x)/2 + 1/2))), x)`

3.180
$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$$

3.180.1 Optimal result 1226
 3.180.2 Mathematica [B] (verified) 1227
 3.180.3 Rubi [A] (verified) 1228
 3.180.4 Maple [B] (verified) 1231
 3.180.5 Fricas [A] (verification not implemented) 1232
 3.180.6 Sympy [F(-1)] 1233
 3.180.7 Maxima [B] (verification not implemented) 1233
 3.180.8 Giac [F] 1234
 3.180.9 Mupad [F(-1)] 1235

3.180.1 Optimal result

Integrand size = 38, antiderivative size = 140

$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx = -\frac{2\operatorname{arcsinh}\left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{ac}f} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{\sec(e+fx)} \sin(e+fx)}{\sqrt{2}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{2}\sqrt{ac}f} + \frac{\csc(e+fx)\sqrt{a+a \sec(e+fx)}}{acf\sqrt{\sec(e+fx)}}$$

output

```
-2*arcsinh(a^(1/2)*tan(f*x+e)/(a+a*sec(f*x+e))^(1/2))/c/f/a^(1/2)+1/2*arctanh(1/2*sin(f*x+e)*a^(1/2)*sec(f*x+e)^(1/2)*2^(1/2)/(a+a*sec(f*x+e))^(1/2))/c/f*2^(1/2)/a^(1/2)+csc(f*x+e)*(a+a*sec(f*x+e))^(1/2)/a/c/f/sec(f*x+e)^(1/2)
```

3.180.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 724 vs. $2(140) = 280$.

Time = 11.25 (sec) , antiderivative size = 724, normalized size of antiderivative = 5.17

$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

$$= \frac{\sec^{\frac{3}{2}}(e+fx)\sqrt{(1+\cos(e+fx))\sec(e+fx)}\sqrt{1+\sec(e+fx)}\left(-\frac{2\cot(e)}{f} + \frac{\csc(\frac{e}{2})\csc(\frac{e}{2}+\frac{fx}{2})\sin(\frac{fx}{2})}{f} + \frac{\sec(\frac{e}{2})}{f}\right)}{\sqrt{a(1+\sec(e+fx))}(c-c\sec(e+fx))} + \frac{\cos(e+fx)\left(\log\left(1-2\sec(e+fx)-3\sec^2(e+fx)-2\sqrt{2}\sqrt{\sec(e+fx)}\sqrt{1+\sec(e+fx)}\sqrt{-1+\sec(e+fx)}\right)\right)}{2f(1+\sec(e+fx))} + \frac{\cos(e+fx)\left(-8\log(1+\sec(e+fx))+8\log\left(\sqrt{\sec(e+fx)}+\sec^{\frac{3}{2}}(e+fx)+\sqrt{1+\sec(e+fx)}\sqrt{-1+\sec(e+fx)}\right)\right)}{2f(1+\sec(e+fx))}$$

input `Integrate[Sec[e + f*x]^(5/2)/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output `(Sec[e + f*x]^(3/2)*Sqrt[(1 + Cos[e + f*x])*Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*((-2*Cot[e])/f + (Csc[e/2]*Csc[e/2 + (f*x)/2]*Sin[(f*x)/2])/f + (Sec[e/2]*Sec[e/2 + (f*x)/2]*Sin[(f*x)/2])/f)*Sin[e/2 + (f*x)/2]^2/(Sqrt[a*(1 + Sec[e + f*x])]*(c - c*Sec[e + f*x])) + (Cos[e + f*x]*(Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 - 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[-1 + Sec[e + f*x]]*Sqrt[-1 + Sec[e + f*x]^2]] - Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 + 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[-1 + Sec[e + f*x]^2]])*(1 + Sec[e + f*x])^(3/2)*Sqrt[-1 + Sec[e + f*x]^2]*Sin[e/2 + (f*x)/2]^2*Sin[e + f*x])/(2*f*(1 + Cos[e + f*x])*Sqrt[2 - 2*Cos[e + f*x]^2]*Sqrt[1 - Cos[e + f*x]^2]*Sqrt[a*(1 + Sec[e + f*x])]*(c - c*Sec[e + f*x])) + (Cos[e + f*x]*(-8*Log[1 + Sec[e + f*x]] + 8*Log[Sqrt[Sec[e + f*x]] + Sec[e + f*x]^(3/2) + Sqrt[1 + Sec[e + f*x]]*Sqrt[-1 + Sec[e + f*x]^2]] + Sqrt[2]*(-Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 - 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[-1 + Sec[e + f*x]^2]] + Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 + 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[-1 + Sec[e + f*x]^2]]))*(1 + Sec[e + f*x])^(3/2)*Sqrt[-1 + Sec[e + f*x]^2]*Sin[e/2 + (f*x)/2]^2*Sin[e + f*x])/(2*f*(1 + Cos[e + f*x])*(1 - Cos[e + f*x]^2)*Sqrt[a*(1 + Sec[e + f*x])]*(c - c*Sec[e + f*x]))`

$$3.180. \quad \int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

3.180.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.12, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {3042, 4452, 27, 109, 27, 175, 64, 104, 216, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a \sec(e+fx) + a(c - c \sec(e+fx))}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(e+fx + \frac{\pi}{2})^{5/2}}{\sqrt{a \csc(e+fx + \frac{\pi}{2}) + a(c - c \csc(e+fx + \frac{\pi}{2}))}} dx$$

$$\downarrow \text{4452}$$

$$\frac{a \tan(e+fx) \int \frac{\sec^{\frac{3}{2}}(e+fx)}{a(\sec(e+fx)+1)(c - c \sec(e+fx))^{3/2}} d \sec(e+fx)}{f \sqrt{a \sec(e+fx) + a \sqrt{c - c \sec(e+fx)}}$$

$$\downarrow \text{27}$$

$$\frac{c \tan(e+fx) \int \frac{\sec^{\frac{3}{2}}(e+fx)}{(\sec(e+fx)+1)(c - c \sec(e+fx))^{3/2}} d \sec(e+fx)}{f \sqrt{a \sec(e+fx) + a \sqrt{c - c \sec(e+fx)}}$$

$$\downarrow \text{109}$$

$$\frac{c \tan(e+fx) \left(\frac{\sqrt{\sec(e+fx)}}{c \sqrt{c - c \sec(e+fx)}} - \frac{\int \frac{c(2 \sec(e+fx)+1)}{2 \sqrt{\sec(e+fx)}(\sec(e+fx)+1) \sqrt{c - c \sec(e+fx)}} d \sec(e+fx)}{c^2} \right)}{f \sqrt{a \sec(e+fx) + a \sqrt{c - c \sec(e+fx)}}$$

$$\downarrow \text{27}$$

$$\frac{c \tan(e+fx) \left(\frac{\sqrt{\sec(e+fx)}}{c \sqrt{c - c \sec(e+fx)}} - \frac{\int \frac{2 \sec(e+fx)+1}{\sqrt{\sec(e+fx)}(\sec(e+fx)+1) \sqrt{c - c \sec(e+fx)}} d \sec(e+fx)}{2c} \right)}{f \sqrt{a \sec(e+fx) + a \sqrt{c - c \sec(e+fx)}}$$

$$\downarrow \text{175}$$

$$\frac{c \tan(e+fx) \left(\frac{\sqrt{\sec(e+fx)}}{c \sqrt{c - c \sec(e+fx)}} - \frac{2 \int \frac{1}{\sqrt{\sec(e+fx)} \sqrt{c - c \sec(e+fx)}} d \sec(e+fx) - \int \frac{1}{\sqrt{\sec(e+fx)}(\sec(e+fx)+1) \sqrt{c - c \sec(e+fx)}} d \sec(e+fx)}{2c} \right)}{f \sqrt{a \sec(e+fx) + a \sqrt{c - c \sec(e+fx)}}$$

$$\downarrow \text{64}$$

$$3.180. \quad \int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))}} dx$$

$$c \tan(e + fx) \left(\frac{\frac{\sqrt{\sec(e+fx)}}{c\sqrt{c-c\sec(e+fx)}} - \frac{-\int \frac{1}{\sqrt{\sec(e+fx)(\sec(e+fx)+1)\sqrt{c-c\sec(e+fx)}} d\sec(e+fx)}{2c} - \frac{4 \int \frac{1}{\sqrt{1-\frac{c-c\sec(e+fx)}{c}}} d\sqrt{c-c\sec(e+fx)}}{c}}{\frac{f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}}{2c}} \right)$$

$$f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}$$

↓ 104

$$c \tan(e + fx) \left(\frac{\frac{\sqrt{\sec(e+fx)}}{c\sqrt{c-c\sec(e+fx)}} - \frac{-2 \int \frac{1}{\frac{2c\sec(e+fx)}{c-c\sec(e+fx)}+1} d\frac{\sqrt{\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}} - \frac{4 \int \frac{1}{\sqrt{1-\frac{c-c\sec(e+fx)}{c}}} d\sqrt{c-c\sec(e+fx)}}{c}}{\frac{f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}}{2c}} \right)$$

$$f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}$$

↓ 216

$$c \tan(e + fx) \left(\frac{\frac{\sqrt{\sec(e+fx)}}{c\sqrt{c-c\sec(e+fx)}} - \frac{4 \int \frac{1}{\sqrt{1-\frac{c-c\sec(e+fx)}{c}}} d\sqrt{c-c\sec(e+fx)}}{c} - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{c}}}{\frac{f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}}{2c}} \right)$$

$$f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}$$

↓ 223

$$c \tan(e + fx) \left(\frac{\frac{\sqrt{\sec(e+fx)}}{c\sqrt{c-c\sec(e+fx)}} - \frac{4 \arcsin\left(\frac{\sqrt{c-c\sec(e+fx)}}{\sqrt{c}}\right)}{\sqrt{c}} - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{\sec(e+fx)}}{\sqrt{c-c\sec(e+fx)}}\right)}{\sqrt{c}}}{\frac{f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}}{2c}} \right)$$

$$f\sqrt{a\sec(e+fx)+a\sqrt{c-c\sec(e+fx)}}$$

input `Int[Sec[e + f*x]^(5/2)/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output `-((c*(-1/2*((-4*ArcSin[Sqrt[c - c*Sec[e + f*x]]/Sqrt[c]])/Sqrt[c] - (Sqrt[2]*ArcTan[(Sqrt[2]*Sqrt[c]*Sqrt[Sec[e + f*x]])/Sqrt[c - c*Sec[e + f*x]]])/Sqrt[c])/c + Sqrt[Sec[e + f*x]]/(c*Sqrt[c - c*Sec[e + f*x]]))*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

$$3.180. \quad \int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

3.180.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 64 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2/b Subst[Int[1/Sqrt[c - a*(d/b) + d*(x^2/b)], x], x, Sqrt[a + b*x]], x] /; FreeQ[{a, b, c, d}, x] && GtQ[c - a*(d/b), 0] && (!GtQ[a - c*(b/d), 0] || PosQ[b])`
- rule 104 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 109 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 175 `Int[(((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)*((g_) + (h_)*(x_)))/((a_) + (b_)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.180.
$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)(c-c\sec(e+fx))}} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4452 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a*c*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.180.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 254 vs. $2(120) = 240$.

Time = 5.05 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.82

method	result
default	$\frac{\left(\sqrt{2} \arctan\left(\frac{\sin(fx+e)\sqrt{2}}{2(\cos(fx+e)+1)\sqrt{-\frac{1}{\cos(fx+e)+1}}}\right) \sin(fx+e) - 2\sqrt{-\frac{1}{\cos(fx+e)+1}} \cos(fx+e) - 2 \arctan\left(\frac{-\cos(fx+e)+\sin(fx+e)-1}{2(\cos(fx+e)+1)\sqrt{-\frac{1}{\cos(fx+e)+1}}}\right)}{2cfa(\cos(fx+e))}$

input `int(sec(f*x+e)^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2), x, method=_RETURNVERBOSE)`

output `-1/2/c/f/a*(2^(1/2)*arctan(1/2*sin(f*x+e)*2^(1/2)/(cos(f*x+e)+1)/(-1/(cos(f*x+e)+1))^(1/2))*sin(f*x+e)-2*(-1/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)-2*arctan(1/2*(-cos(f*x+e)+sin(f*x+e)-1)/(cos(f*x+e)+1)/(-1/(cos(f*x+e)+1))^(1/2))*sin(f*x+e)-2*arctan(1/2*(cos(f*x+e)+sin(f*x+e)+1)/(cos(f*x+e)+1)/(-1/(cos(f*x+e)+1))^(1/2))*sin(f*x+e)-2*(-1/(cos(f*x+e)+1))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)*sec(f*x+e)^(5/2)/(cos(f*x+e)+1)/(-1/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2*cot(f*x+e)`

3.180.
$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

3.180.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 462, normalized size of antiderivative = 3.30

$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

$$= \frac{\sqrt{2}\sqrt{a} \log\left(-\frac{\cos(fx+e)^2 - \frac{2\sqrt{2}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\cos(fx+e)}\sin(fx+e)}{\sqrt{a}} - 2\cos(fx+e) - 3}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right) \sin(fx+e) + 2\sqrt{a} \log\left(\frac{a\cos(fx+e)}{4acf \sin(fx+e)}\right)}{\sqrt{2}a\sqrt{-\frac{1}{a}} \arctan\left(\frac{\sqrt{2}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{-\frac{1}{a}}\sqrt{\cos(fx+e)}}{\sin(fx+e)}\right) \sin(fx+e) + 2\sqrt{-a} \arctan\left(\frac{2\sqrt{-a}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\cos(fx+e)}}{a\cos(fx+e)^2 - a\cos(fx+e)}\right)}{2acf \sin(fx+e)}$$

```
input integrate(sec(f*x+e)^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algo
rithm="fricas")
```

```
output [1/4*(sqrt(2)*sqrt(a)*log(-(cos(f*x + e)^2 - 2*sqrt(2)*sqrt((a*cos(f*x + e)
+ a)/cos(f*x + e))*sqrt(cos(f*x + e))*sin(f*x + e)/sqrt(a) - 2*cos(f*x +
e) - 3)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1))*sin(f*x + e) + 2*sqrt(a)*l
og((a*cos(f*x + e)^3 - 7*a*cos(f*x + e)^2 + 4*(cos(f*x + e)^2 - 2*cos(f*x
+ e))*sqrt(a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sin(f*x + e)/sqrt(co
s(f*x + e)) + 8*a)/(cos(f*x + e)^3 + cos(f*x + e)^2))*sin(f*x + e) + 4*sq
rt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(cos(f*x + e)))/(a*c*f*sin(f*x +
e)), -1/2*(sqrt(2)*a*sqrt(-1/a)*arctan(sqrt(2)*sqrt((a*cos(f*x + e) + a)/c
os(f*x + e))*sqrt(-1/a)*sqrt(cos(f*x + e))/sin(f*x + e))*sin(f*x + e) + 2*
sqrt(-a)*arctan(2*sqrt(-a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(co
s(f*x + e))*sin(f*x + e)/(a*cos(f*x + e)^2 - a*cos(f*x + e) - 2*a))*sin(f*
x + e) - 2*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(cos(f*x + e)))/(a*
c*f*sin(f*x + e))]
```

$$3.180. \quad \int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx$$

3.180.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)**(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x)`output `Timed out`**3.180.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1310 vs. 2(120) = 240.

Time = 0.45 (sec) , antiderivative size = 1310, normalized size of antiderivative = 9.36

$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output

```
-1/2*((sqrt(2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sqrt(2)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + sqrt(2))*log(2*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sqrt(2)*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*sqrt(2)*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 2) - (sqrt(2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sqrt(2)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + sqrt(2))*log(2*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sqrt(2)*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - 2*sqrt(2)*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 2) + (sqrt(2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sqrt(2)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + sqrt(2))*log(2*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*sqrt(2)*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 2) - (sqrt(2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sqrt(2)*sin(1/2*arc...
```

3.180.8 Giac [F]

$$\int \frac{\sec^{\frac{5}{2}}(e + fx)}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx$$

$$= \int -\frac{\sec^{\frac{5}{2}}(fx + e)}{\sqrt{a \sec(fx + e) + a}(c \sec(fx + e) - c)} dx$$

input

```
integrate(sec(f*x+e)^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
integrate(-sec(f*x + e)^(5/2)/(sqrt(a*sec(f*x + e) + a)*(c*sec(f*x + e) - c)), x)
```

3.180. $\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$

3.180.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{\frac{5}{2}}(e+fx)}{\sqrt{a+a\sec(e+fx)}(c-c\sec(e+fx))} dx = \int \frac{\left(\frac{1}{\cos(e+fx)}\right)^{5/2}}{\sqrt{a+\frac{a}{\cos(e+fx)}}\left(c-\frac{c}{\cos(e+fx)}\right)} dx$$

input `int((1/cos(e + f*x))^(5/2)/((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))),x)`

output `int((1/cos(e + f*x))^(5/2)/((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))), x)`

$$3.181 \quad \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$$

3.181.1 Optimal result	1236
3.181.2 Mathematica [B] (verified)	1236
3.181.3 Rubi [A] (verified)	1237
3.181.4 Maple [A] (verified)	1239
3.181.5 Fricas [A] (verification not implemented)	1240
3.181.6 Sympy [F]	1240
3.181.7 Maxima [B] (verification not implemented)	1241
3.181.8 Giac [F]	1241
3.181.9 Mupad [F(-1)]	1242

3.181.1 Optimal result

Integrand size = 40, antiderivative size = 116

$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx =$$

$$-\frac{g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{2}\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{2}\sqrt{acf}}$$

$$+ \frac{g \cot(e+fx) \sqrt{g \sec(e+fx)} \sqrt{a+a \sec(e+fx)}}{acf}$$

```
output -1/2*g^(3/2)*arctanh(1/2*a^(1/2)*g^(1/2)*tan(f*x+e)*2^(1/2)/(g*sec(f*x+e))
^(1/2)/(a+a*sec(f*x+e))^(1/2))/c/f*2^(1/2)/a^(1/2)+g*cot(f*x+e)*(g*sec(f*x
+e))^(1/2)*(a+a*sec(f*x+e))^(1/2)/a/c/f
```

3.181.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 236 vs. 2(116) = 232.

Time = 3.92 (sec) , antiderivative size = 236, normalized size of antiderivative = 2.03

$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx =$$

$$\frac{a \cos\left(\frac{1}{2}(e+fx)\right) (g \sec(e+fx))^{5/2} \sin^3\left(\frac{1}{2}(e+fx)\right) \left(-4 - 4 \sec(e+fx) + \frac{(\log(1-2 \sec(e+fx)-3 \sec^2(e+fx))-2)}{c f g(-1 + \sec(e+fx))}\right)}{c f g(-1 + \sec(e+fx))}$$

$$3.181. \quad \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$$

input `Integrate[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output `-((a*Cos[(e + f*x)/2]*(g*Sec[e + f*x])^(5/2)*Sin[(e + f*x)/2]^3*(-4 - 4*Sec[e + f*x] + ((Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 - 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[Tan[e + f*x]^2]] - Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 + 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[Tan[e + f*x]^2]))*Sqrt[Tan[e + f*x]^2])/Sqrt[Sec[(e + f*x)/2]^2]))/(c*f*g*(-1 + Sec[e + f*x])^2*(a*(1 + Sec[e + f*x]))^(3/2))`

3.181.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3042, 4452, 27, 105, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a \sec(e + fx) + a(c - c \sec(e + fx))}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a(c - c \csc(e + fx + \frac{\pi}{2}))}} dx \\
 & \quad \downarrow \text{4452} \\
 & -\frac{acg \tan(e + fx) \int \frac{\sqrt{g \sec(e + fx)}}{a(\sec(e + fx) + 1)(c - c \sec(e + fx))^{3/2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} \\
 & \quad \downarrow \text{27} \\
 & -\frac{cg \tan(e + fx) \int \frac{\sqrt{g \sec(e + fx)}}{(\sec(e + fx) + 1)(c - c \sec(e + fx))^{3/2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}} \\
 & \quad \downarrow \text{105} \\
 & -\frac{cg \tan(e + fx) \left(\frac{\sqrt{g \sec(e + fx)}}{c \sqrt{c - c \sec(e + fx)}} - \frac{g \int \frac{1}{\sqrt{g \sec(e + fx)}(\sec(e + fx) + 1)\sqrt{c - c \sec(e + fx)}} d \sec(e + fx)}{2c} \right)}{f \sqrt{a \sec(e + fx) + a \sqrt{c - c \sec(e + fx)}}}
 \end{aligned}$$

3.181. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx$

$$\begin{array}{c}
 \downarrow 104 \\
 \frac{cg \tan(e + fx) \left(\frac{\sqrt{g \sec(e + fx)}}{c\sqrt{c - c \sec(e + fx)}} - \frac{g \int \frac{1}{\frac{2c \sec(e + fx)g}{c - c \sec(e + fx)} + g} d \frac{\sqrt{g \sec(e + fx)}}{\sqrt{c - c \sec(e + fx)}}}{c} \right)}{f\sqrt{a \sec(e + fx) + a}\sqrt{c - c \sec(e + fx)}} \\
 \downarrow 218 \\
 \frac{cg \tan(e + fx) \left(\frac{\sqrt{g \sec(e + fx)}}{c\sqrt{c - c \sec(e + fx)}} - \frac{\sqrt{g} \arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{g \sec(e + fx)}}{\sqrt{g}\sqrt{c - c \sec(e + fx)}}\right)}{\sqrt{2}c^{3/2}} \right)}{f\sqrt{a \sec(e + fx) + a}\sqrt{c - c \sec(e + fx)}}
 \end{array}$$

input `Int[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output `-((c*g*(-((Sqrt[g]*ArcTan[(Sqrt[2]*Sqrt[c]*Sqrt[g*Sec[e + f*x]])/(Sqrt[g]*Sqrt[c - c*Sec[e + f*x]])]))/(Sqrt[2]*c^(3/2))) + Sqrt[g*Sec[e + f*x]]/(c*Sqrt[c - c*Sec[e + f*x]]))*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.181.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 105 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((m + 1)*(b*e - a*f))), x] - Simp[n*((d*e - c*f)/((m + 1)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`

3.181. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)(c - c \sec(e + fx))}} dx$

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4452 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a*c*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.181.4 Maple [A] (verified)

Time = 3.07 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.07

method	result
default	$\frac{g \left(\operatorname{arcsinh}(\cot(fx+e) - \csc(fx+e)) \sqrt{2} \sin(fx+e) + 2 \sqrt{\frac{1}{\cos(fx+e)+1}} \cos(fx+e) + 2 \sqrt{\frac{1}{\cos(fx+e)+1}} \right) \sqrt{g \sec(fx+e)} \sqrt{a(\sec(fx+e)+1)}}{2cfa(\cos(fx+e)+1) \sqrt{\frac{1}{\cos(fx+e)+1}}}$

input `int((g*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*g/c/f/a*(arcsinh(cot(f*x+e)-csc(f*x+e))*2^(1/2)*sin(f*x+e)+2*(1/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)+2*(1/(cos(f*x+e)+1))^(1/2))*(g*sec(f*x+e))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)/(cos(f*x+e)+1)/(1/(cos(f*x+e)+1))^(1/2)*cot(f*x+e)`

3.181.
$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$$

3.181.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 330, normalized size of antiderivative = 2.84

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \left[\frac{\sqrt{2}ag \sqrt{\frac{g}{a}} \log \left(-\frac{2\sqrt{2}\sqrt{\frac{g}{a}} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{g}{\cos(fx+e)}} \cos(fx+e) \sin(fx+e)}{\cos(fx+e)^2 + 2 \cos(fx+e)} \right)}{\dots} \right]$$

```
input integrate((g*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="fricas")
```

```
output [1/4*(sqrt(2)*a*g*sqrt(g/a)*log(-(2*sqrt(2)*sqrt(g/a)*sqrt((a*cos(f*x + e)
+ a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) + g*cos
(f*x + e)^2 - 2*g*cos(f*x + e) - 3*g)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1
))*sin(f*x + e) + 4*g*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f
*x + e))*cos(f*x + e))/(a*c*f*sin(f*x + e)), 1/2*(sqrt(2)*a*g*sqrt(-g/a)*a
rctan(sqrt(2)*sqrt(-g/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/co
s(f*x + e))*cos(f*x + e)/(g*sin(f*x + e))*sin(f*x + e) + 2*g*sqrt((a*cos(
f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e))/(a*c*f*sin(
f*x + e))]
```

3.181.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = -\frac{\int \frac{(g \sec(e+fx))^{\frac{3}{2}}}{\sqrt{a \sec(e+fx)+a \sec(e+fx)-\sqrt{a \sec(e+fx)+a}} dx}{c}$$

```
input integrate((g*sec(f*x+e))**(3/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x
)
```

```
output -Integral((g*sec(e + f*x))**(3/2)/(sqrt(a*sec(e + f*x) + a)*sec(e + f*x) -
sqrt(a*sec(e + f*x) + a)), x)/c
```

3.181.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 536 vs. $2(97) = 194$.

Time = 0.42 (sec) , antiderivative size = 536, normalized size of antiderivative = 4.62

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \frac{\left(4g \cos\left(\frac{1}{4} \arctan(\sin(2fx + 2e), \cos(2fx + 2e))\right) \sin\left(\frac{1}{2}\right)}{\right)}$$

input `integrate((g*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="maxima")`

output `1/2*(4*g*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 4*g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + g*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + g)*log(cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + (g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + g*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*g*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + g)*log(cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 4*g*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(g)/((sqrt(2)*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sqrt(2)*c*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + sqrt(2)*c)*sqrt(a)*f)`

3.181.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \int -\frac{(g \sec(fx + e))^{3/2}}{\sqrt{a \sec(fx + e) + a}(c \sec(fx + e) - c)} dx$$

input `integrate((g*sec(f*x+e))^(3/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="giac")`

3.181. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx$

output `integrate(-(g*sec(f*x + e))^(3/2)/(sqrt(a*sec(f*x + e) + a)*(c*sec(f*x + e) - c)), x)`

3.181.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{3/2}}{\sqrt{a + \frac{a}{\cos(e+fx)}} \left(c - \frac{c}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(3/2)/((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))),x)`

output `int((g/cos(e + f*x))^(3/2)/((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))), x)`

$$3.182 \quad \int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$$

3.182.1 Optimal result	1243
3.182.2 Mathematica [A] (verified)	1243
3.182.3 Rubi [A] (verified)	1244
3.182.4 Maple [A] (verified)	1247
3.182.5 Fricas [A] (verification not implemented)	1248
3.182.6 Sympy [F(-1)]	1249
3.182.7 Maxima [B] (verification not implemented)	1249
3.182.8 Giac [F]	1250
3.182.9 Mupad [F(-1)]	1251

3.182.1 Optimal result

Integrand size = 40, antiderivative size = 179

$$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx =$$

$$\frac{2g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{ac}f} + \frac{g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{2}\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{2}\sqrt{ac}f}$$

$$+ \frac{g^2 \cot(e+fx) \sqrt{g \sec(e+fx)} \sqrt{a+a \sec(e+fx)}}{acf}$$

```
output -2*g^(5/2)*arctanh(a^(1/2)*g^(1/2)*tan(f*x+e)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))/c/f/a^(1/2)+1/2*g^(5/2)*arctanh(1/2*a^(1/2)*g^(1/2)*tan(f*x+e)*2^(1/2)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))/c/f*2^(1/2)/a^(1/2)+g^2*cot(f*x+e)*(g*sec(f*x+e))^(1/2)*(a+a*sec(f*x+e))^(1/2)/a/c/f
```

3.182.2 Mathematica [A] (verified)

Time = 3.07 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.83

$$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx =$$

$$\frac{(g \sec(e+fx))^{5/2} \sqrt{1+\sec(e+fx)} \sin^3(e+fx) \left(8\sqrt{\sec(e+fx)}\sqrt{1+\sec(e+fx)} + (16 \log(1+\sec(e+fx)))\right)}{\dots}$$

3.182. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c-c \sec(e+fx))} dx$

input `Integrate[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output `-1/8*((g*Sec[e + f*x])^(5/2)*Sqrt[1 + Sec[e + f*x]]*Sin[e + f*x]^3*(8*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]] + (16*Log[1 + Sec[e + f*x]] - 16*Log[Sqrt[Sec[e + f*x]] + Sec[e + f*x]^(3/2) + Sqrt[1 + Sec[e + f*x]]*Sqrt[Tan[e + f*x]^2]] + Sqrt[2]*(Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 - 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[Tan[e + f*x]^2]] - Log[1 - 2*Sec[e + f*x] - 3*Sec[e + f*x]^2 + 2*Sqrt[2]*Sqrt[Sec[e + f*x]]*Sqrt[1 + Sec[e + f*x]]*Sqrt[Tan[e + f*x]^2]]))*Sqrt[Tan[e + f*x]^2))/(c*f*(-1 + Cos[e + f*x])*(1 + Cos[e + f*x])^2*(-1 + Sec[e + f*x])*Sec[e + f*x]^(5/2)*Sqrt[a*(1 + Sec[e + f*x])])`

3.182.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.11, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.225$, Rules used = {3042, 4452, 27, 109, 27, 175, 65, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a \sec(e + fx) + a}(c - c \sec(e + fx))} dx$$

↓ 3042

$$\int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{5/2}}{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a}(c - c \csc(e + fx + \frac{\pi}{2}))} dx$$

↓ 4452

$$\frac{acg \tan(e + fx) \int \frac{(g \sec(e + fx))^{3/2}}{a(\sec(e + fx) + 1)(c - c \sec(e + fx))^{3/2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}}$$

↓ 27

$$\frac{cg \tan(e + fx) \int \frac{(g \sec(e + fx))^{3/2}}{(\sec(e + fx) + 1)(c - c \sec(e + fx))^{3/2}} d \sec(e + fx)}{f \sqrt{a \sec(e + fx) + a} \sqrt{c - c \sec(e + fx)}}$$

↓ 109

3.182. $\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx$

$$\begin{aligned}
& \frac{cg \tan(e+fx) \left(\frac{g\sqrt{g \sec(e+fx)}}{c\sqrt{c-c \sec(e+fx)}} - \frac{\int \frac{cg^2(2 \sec(e+fx)+1)}{2\sqrt{g \sec(e+fx)(\sec(e+fx)+1)\sqrt{c-c \sec(e+fx)}} d \sec(e+fx)}{c^2} \right)}{f\sqrt{a \sec(e+fx) + a\sqrt{c-c \sec(e+fx)}}} \\
& \quad \downarrow 27 \\
& \frac{cg \tan(e+fx) \left(\frac{g\sqrt{g \sec(e+fx)}}{c\sqrt{c-c \sec(e+fx)}} - \frac{g^2 \int \frac{2 \sec(e+fx)+1}{\sqrt{g \sec(e+fx)(\sec(e+fx)+1)\sqrt{c-c \sec(e+fx)}} d \sec(e+fx)}{2c} \right)}{f\sqrt{a \sec(e+fx) + a\sqrt{c-c \sec(e+fx)}}} \\
& \quad \downarrow 175 \\
& \frac{cg \tan(e+fx) \left(\frac{g\sqrt{g \sec(e+fx)}}{c\sqrt{c-c \sec(e+fx)}} - \frac{g^2 \left(2 \int \frac{1}{\sqrt{g \sec(e+fx)\sqrt{c-c \sec(e+fx)}} d \sec(e+fx)} - \int \frac{1}{\sqrt{g \sec(e+fx)(\sec(e+fx)+1)\sqrt{c-c \sec(e+fx)}} d \sec(e+fx)} \right)}{2c} \right)}{f\sqrt{a \sec(e+fx) + a\sqrt{c-c \sec(e+fx)}}} \\
& \quad \downarrow 65 \\
& \frac{cg \tan(e+fx) \left(\frac{g\sqrt{g \sec(e+fx)}}{c\sqrt{c-c \sec(e+fx)}} - \frac{g^2 \left(4 \int \frac{1}{\frac{c \sec(e+fx)g}{c-c \sec(e+fx)} + g} d \frac{\sqrt{g \sec(e+fx)}}{\sqrt{c-c \sec(e+fx)}} - \int \frac{1}{\sqrt{g \sec(e+fx)(\sec(e+fx)+1)\sqrt{c-c \sec(e+fx)}} d \sec(e+fx)} \right)}{2c} \right)}{f\sqrt{a \sec(e+fx) + a\sqrt{c-c \sec(e+fx)}}} \\
& \quad \downarrow 104 \\
& \frac{cg \tan(e+fx) \left(\frac{g\sqrt{g \sec(e+fx)}}{c\sqrt{c-c \sec(e+fx)}} - \frac{g^2 \left(4 \int \frac{1}{\frac{c \sec(e+fx)g}{c-c \sec(e+fx)} + g} d \frac{\sqrt{g \sec(e+fx)}}{\sqrt{c-c \sec(e+fx)}} - 2 \int \frac{1}{\frac{2c \sec(e+fx)g}{c-c \sec(e+fx)} + g} d \frac{\sqrt{g \sec(e+fx)}}{\sqrt{c-c \sec(e+fx)}} \right)}{2c} \right)}{f\sqrt{a \sec(e+fx) + a\sqrt{c-c \sec(e+fx)}}} \\
& \quad \downarrow 218 \\
& \frac{cg \tan(e+fx) \left(\frac{g\sqrt{g \sec(e+fx)}}{c\sqrt{c-c \sec(e+fx)}} - \frac{g^2 \left(\frac{4 \arctan \left(\frac{\sqrt{c}\sqrt{g \sec(e+fx)}}{\sqrt{g}\sqrt{c-c \sec(e+fx)}} \right)}{\sqrt{c}\sqrt{g}} - \frac{\sqrt{2} \arctan \left(\frac{\sqrt{2}\sqrt{c}\sqrt{g \sec(e+fx)}}{\sqrt{g}\sqrt{c-c \sec(e+fx)}} \right)}{\sqrt{c}\sqrt{g}} \right)}{2c} \right)}{f\sqrt{a \sec(e+fx) + a\sqrt{c-c \sec(e+fx)}}}
\end{aligned}$$

3.182. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))}} dx$

input `Int[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + a*Sec[e + f*x]]*(c - c*Sec[e + f*x])),x]`

output `-((c*g*(-1/2*(g^2*((4*ArcTan[(Sqrt[c]*Sqrt[g*Sec[e + f*x]])/(Sqrt[g]*Sqrt[c - c*Sec[e + f*x]])))/(Sqrt[c]*Sqrt[g]) - (Sqrt[2]*ArcTan[(Sqrt[2]*Sqrt[c]*Sqrt[g*Sec[e + f*x]])/(Sqrt[g]*Sqrt[c - c*Sec[e + f*x]])])/(Sqrt[c]*Sqrt[g])))/c + (g*Sqrt[g*Sec[e + f*x]])/(c*Sqrt[c - c*Sec[e + f*x]]))*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]))`

3.182.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 104 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 109 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

rule 175 `Int[(((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_))*((g_) + (h_)*(x_))) / ((a_) + (b_)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4452 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a*c*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*(c + d*x)^(n - 1/2), x], x, Cs c[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

3.182.4 Maple [A] (verified)

Time = 5.20 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.24

method	result
default	$-\frac{g^2 \left(\operatorname{arcsinh}(\cot(fx+e) - \csc(fx+e))\sqrt{2} \sin(fx+e) - 2\sqrt{\frac{1}{\cos(fx+e)+1}} \cos(fx+e) + 2 \operatorname{arctanh}\left(\frac{-\cos(fx+e) + \sin(fx+e) - 1}{2(\cos(fx+e)+1)\sqrt{\frac{1}{\cos(fx+e)+1}}}\right) \right) \sin(fx+e)}{2cfa(\cos(fx+e)+1)}$

input `int((g*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2/c/f/a*g^2*(arcsinh(cot(f*x+e)-csc(f*x+e))*2^(1/2)*sin(f*x+e)-2*(1/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)+2*arctanh(1/2*(-cos(f*x+e)+sin(f*x+e)-1)/(cos(f*x+e)+1)/(1/(cos(f*x+e)+1))^(1/2))*sin(f*x+e)+2*arctanh(1/2*(cos(f*x+e)+sin(f*x+e)+1)/(cos(f*x+e)+1)/(1/(cos(f*x+e)+1))^(1/2))*sin(f*x+e)-2*(1/(cos(f*x+e)+1))^(1/2))*(a*(sec(f*x+e)+1))^(1/2)*(g*sec(f*x+e))^(1/2)/(cos(f*x+e)+1)/(1/(cos(f*x+e)+1))^(1/2)*cot(f*x+e)`

3.182.
$$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)(c-c \sec(e+fx))}} dx$$

3.182.5 Fracas [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 569, normalized size of antiderivative = 3.18

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \frac{\sqrt{2}ag^2 \sqrt{\frac{g}{a}} \log \left(\frac{2\sqrt{2}\sqrt{\frac{g}{a}} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{g}{\cos(fx+e)}} \cos(fx+e) \sin(fx+e)}{\cos(fx+e)^2 + 2 \cos(fx+e)} \right)}{\sqrt{2}ag^2 \sqrt{-\frac{g}{a}} \arctan \left(\frac{\sqrt{2}\sqrt{-\frac{g}{a}} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{g}{\cos(fx+e)}} \cos(fx+e)}{g \sin(fx+e)} \right) \sin(fx+e) + 2ag^2 \sqrt{-\frac{g}{a}} \arctan \left(\frac{2\sqrt{-\frac{g}{a}} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}}}{g} \right)}{2acf \sin(fx+e)}$$

input `integrate((g*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="fracas")`

output `[1/4*(sqrt(2)*a*g^2*sqrt(g/a)*log((2*sqrt(2)*sqrt(g/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) - g*cos(f*x + e)^2 + 2*g*cos(f*x + e) + 3*g)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1))*sin(f*x + e) + 2*a*g^2*sqrt(g/a)*log((g*cos(f*x + e)^3 + 4*(cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt(g/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) - 7*g*cos(f*x + e)^2 + 8*g)/(cos(f*x + e)^3 + cos(f*x + e)^2))*sin(f*x + e) + 4*g^2*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e))/(a*c*f*sin(f*x + e)), -1/2*(sqrt(2)*a*g^2*sqrt(-g/a)*arctan(sqrt(2)*sqrt(-g/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)/(g*sin(f*x + e)))*sin(f*x + e) + 2*a*g^2*sqrt(-g/a)*arctan(2*sqrt(-g/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)/(g*cos(f*x + e)^2 - g*cos(f*x + e) - 2*g))*sin(f*x + e) - 2*g^2*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)/(a*c*f*sin(f*x + e))]`

3.182.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \text{Timed out}$$

input `integrate((g*sec(f*x+e))**(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x)`

output `Timed out`

3.182.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1400 vs. 2(149) = 298.

Time = 0.45 (sec) , antiderivative size = 1400, normalized size of antiderivative = 7.82

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \text{Too large to display}$$

input `integrate((g*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,algorithm="maxima")`

output

```

1/2*(4*g^2*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))*sin(1/2*ar
ctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 4*g^2*cos(1/2*arctan2(sin(2*f
*x + 2*e), cos(2*f*x + 2*e)))*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e))) + 4*g^2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - (s
qrt(2)*g^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sqrt(2
)*g^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*g
^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + sqrt(2)*g^2*log
(2*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/4*arct
an2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sqrt(2)*cos(1/4*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*sqrt(2)*sin(1/4*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e))) + 2) + (sqrt(2)*g^2*cos(1/2*arctan2(sin(2*f*x + 2
*e), cos(2*f*x + 2*e)))^2 + sqrt(2)*g^2*sin(1/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*g^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos
(2*f*x + 2*e))) + sqrt(2)*g^2*log(2*cos(1/4*arctan2(sin(2*f*x + 2*e), cos
(2*f*x + 2*e)))^2 + 2*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))
^2 + 2*sqrt(2)*cos(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 2*sq
rt(2)*sin(1/4*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2) - (sqrt(2)
*g^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sqrt(2)*g^2*
sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sqrt(2)*g^2*cos
(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + sqrt(2)*g^2*log(2*...

```

3.182.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \int -\frac{(g \sec(fx + e))^{5/2}}{\sqrt{a \sec(fx + e) + a}(c \sec(fx + e) - c)} dx$$

input

```

integrate((g*sec(f*x+e))^(5/2)/(c-c*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="giac")

```

output

```

integrate(-(g*sec(f*x + e))^(5/2)/(sqrt(a*sec(f*x + e) + a)*(c*sec(f*x + e
) - c)), x)

```

3.182.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c - c \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{5/2}}{\sqrt{a + \frac{a}{\cos(e+fx)}} \left(c - \frac{c}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(5/2)/((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))),x)`

output `int((g/cos(e + f*x))^(5/2)/((a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))), x)`

3.183
$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx$$

3.183.1 Optimal result 1252
 3.183.2 Mathematica [C] (verified) 1252
 3.183.3 Rubi [A] (verified) 1253
 3.183.4 Maple [B] (verified) 1254
 3.183.5 Fricas [B] (verification not implemented) 1255
 3.183.6 Sympy [F] 1255
 3.183.7 Maxima [A] (verification not implemented) 1256
 3.183.8 Giac [F(-2)] 1256
 3.183.9 Mupad [F(-1)] 1257

3.183.1 Optimal result

Integrand size = 38, antiderivative size = 46

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx = \frac{\log(\tan(e+fx)) \tan(e+fx)}{f \sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}}$$

output `ln(tan(f*x+e))*tan(f*x+e)/f/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2)`

3.183.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.76 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.04

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx = \frac{4i(-1 + e^{i(e+fx)}) \operatorname{arctanh}(e^{2i(e+fx)}) \cos^2(\frac{1}{2}(e+fx)) \sec(e+fx)}{(1 + e^{i(e+fx)}) f \sqrt{a(1 + \sec(e+fx))} \sqrt{c - c \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]),x]`

output `((4*I)*(-1 + E^(I*(e + f*x)))*ArcTanh[E^((2*I)*(e + f*x))]*Cos[(e + f*x)/2]^2*Sec[e + f*x])/((1 + E^(I*(e + f*x)))*f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c - c*Sec[e + f*x]])`

3.183.
$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c-c \sec(e+fx)}} dx$$

3.183.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3042, 4451, 25, 3042, 3100, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(e+fx)}{\sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})^2}{\sqrt{a \csc(e+fx + \frac{\pi}{2}) + a} \sqrt{c - c \csc(e+fx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4451} \\
 & \frac{\tan(e+fx) \int -\csc(e+fx) \sec(e+fx) dx}{\sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\tan(e+fx) \int \csc(e+fx) \sec(e+fx) dx}{\sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tan(e+fx) \int \csc(e+fx) \sec(e+fx) dx}{\sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{3100} \\
 & \frac{\tan(e+fx) \int \cot(e+fx) d \tan(e+fx)}{f \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}} \\
 & \quad \downarrow \text{14} \\
 & \frac{\tan(e+fx) \log(\tan(e+fx))}{f \sqrt{a \sec(e+fx) + a} \sqrt{c - c \sec(e+fx)}}
 \end{aligned}$$

input `Int[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]]),x]`

output `(Log[Tan[e + f*x]]*Tan[e + f*x])/(f*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c - c*Sec[e + f*x]])`

3.183. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)} \sqrt{c-c \sec(e+fx)}} dx$

3.183.3.1 Defintions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3100 `Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[1/f Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]`
- rule 4451 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(m_.), x_Symbol] := Simp[((-a)*c)^(m + 1/2)*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])) Int[(g*Csc[e + f*x])^p*Cot[e + f*x]^(2*m), x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IntegerQ[m + 1/2]`

3.183.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 100 vs. $2(42) = 84$.

Time = 2.58 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.20

method	result
default	$\frac{\sqrt{a(\sec(fx+e)+1)}(\ln(-\cot(fx+e)+\csc(fx+e)+1)+\ln(-\cot(fx+e)+\csc(fx+e)-1)-\ln(-\cot(fx+e)+\csc(fx+e)))}{fa\sqrt{-c(\sec(fx+e)-1)}}(\cot(fx+e)-c)$
risch	$\frac{i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)\ln(1+e^{2i(fx+e)})}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}(1+e^{2i(fx+e)})}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f} - \frac{i(e^{i(fx+e)}+1)(e^{i(fx+e)}-1)\ln(e^{2i(fx+e)}-1)}{\sqrt{\frac{a(e^{i(fx+e)}+1)^2}{1+e^{2i(fx+e)}}(1+e^{2i(fx+e)})}\sqrt{\frac{c(e^{i(fx+e)}-1)^2}{1+e^{2i(fx+e)}}}f}$

input `int(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x,method=_R ETURNVERBOSE)`

$$3.183. \quad \int \frac{\sec^2(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} dx$$

output $1/f/a*(a*(\sec(f*x+e)+1))^{(1/2)}*(\ln(-\cot(f*x+e)+\csc(f*x+e)+1)+\ln(-\cot(f*x+e)+\csc(f*x+e)-1)-\ln(-\cot(f*x+e)+\csc(f*x+e)))/(-c*(\sec(f*x+e)-1))^{(1/2)}*(\cot(f*x+e)-\csc(f*x+e))$

3.183.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(42) = 84$.

Time = 0.34 (sec) , antiderivative size = 255, normalized size of antiderivative = 5.54

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} dx$$

$$= \left[\frac{\sqrt{-ac} \log \left(-\frac{8 \left((2 \cos(fx+e))^3 - \cos(fx+e) \right) \sqrt{-ac} \sqrt{\frac{a \cos(fx+e)+a}{\cos(fx+e)}} \sqrt{\frac{c \cos(fx+e)-c}{\cos(fx+e)}} + (2ac \cos(fx+e)^4 - 2ac \cos(fx+e)^2 + ac) \sin(fx+e)}{(\cos(fx+e)^4 - \cos(fx+e)^2) \sin(fx+e)} \right)}{2acf} \right]$$

input `integrate(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `[-1/2*sqrt(-a*c)*log(-8*((2*cos(f*x + e))^3 - cos(f*x + e))*sqrt(-a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e)) + (2*a*c*cos(f*x + e)^4 - 2*a*c*cos(f*x + e)^2 + a*c)*sin(f*x + e))/((cos(f*x + e)^4 - cos(f*x + e)^2)*sin(f*x + e)))/(a*c*f), sqrt(a*c)*arctan(sqrt(a*c)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) - c)/cos(f*x + e))*cos(f*x + e)/((2*a*c*cos(f*x + e)^2 - a*c)*sin(f*x + e)))/(a*c*f)]`

3.183.6 Sympy [F]

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} dx$$

$$= \int \frac{\sec^2(e+fx)}{\sqrt{a(\sec(e+fx)+1)}\sqrt{-c(\sec(e+fx)-1)}} dx$$

input `integrate(sec(f*x+e)**2/(a+a*sec(f*x+e))**(1/2)/(c-c*sec(f*x+e))**(1/2),x)`

3.183. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c-c\sec(e+fx)}} dx$

output `Integral(sec(e + f*x)**2/(sqrt(a*(sec(e + f*x) + 1))*sqrt(-c*(sec(e + f*x) - 1))), x)`

3.183.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.22

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx = \frac{\arctan(\sin(2fx + 2e), \cos(2fx + 2e) + 1) - \arctan(\sin(2fx + 2e), \cos(2fx + 2e) - 1)}{\sqrt{a}\sqrt{c}f}$$

input `integrate(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `-(arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) - 1))/(sqrt(a)*sqrt(c)*f)`

3.183.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c-c*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Limit: Max order reached or unable to make series expansion Error: Bad Argument Value`

3.183.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c - c \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx)^2 \sqrt{a + \frac{a}{\cos(e + fx)}} \sqrt{c - \frac{c}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x)^2*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(1/2)),x)`

output `int(1/(cos(e + f*x)^2*(a + a/cos(e + f*x))^(1/2)*(c - c/cos(e + f*x))^(1/2)), x)`

3.184 $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx$

3.184.1 Optimal result 1258
 3.184.2 Mathematica [A] (verified) 1258
 3.184.3 Rubi [A] (verified) 1259
 3.184.4 Maple [B] (warning: unable to verify) 1260
 3.184.5 Fracas [B] (verification not implemented) 1261
 3.184.6 Sympy [F] 1261
 3.184.7 Maxima [F] 1262
 3.184.8 Giac [F] 1262
 3.184.9 Mupad [F(-1)] 1262

3.184.1 Optimal result

Integrand size = 34, antiderivative size = 65

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx = \frac{2\sqrt{a}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{c-d}\sqrt{a+a\sec(e+fx)}}\right)}{\sqrt{c-d}\sqrt{d}f}$$

output `2*arctanh(a^(1/2)*d^(1/2)*tan(f*x+e)/(c-d)^(1/2)/(a+a*sec(f*x+e))^(1/2))*a^(1/2)/f/(c-d)^(1/2)/d^(1/2)`

3.184.2 Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx = \frac{\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{d}\sin(\frac{1}{2}(e+fx))}{\sqrt{c-d}\sqrt{\cos(e+fx)}}\right)\sqrt{\cos(e+fx)}\sec\left(\frac{1}{2}(e+fx)\right)\sqrt{a(1+\sec(e+fx))}}{\sqrt{c-d}\sqrt{d}f}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c - d*Sec[e + f*x]),x]`

output `(Sqrt[2]*ArcTanh[(Sqrt[2]*Sqrt[d]*Sin[(e + f*x)/2])/(Sqrt[c - d]*Sqrt[Cos[e + f*x]])]*Sqrt[Cos[e + f*x]]*Sec[(e + f*x)/2]*Sqrt[a*(1 + Sec[e + f*x])])/(Sqrt[c - d]*Sqrt[d]*f)`

3.184. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx$

3.184.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.088$, Rules used = {3042, 4455, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a\sec(e+fx)+a}}{c-d\sec(e+fx)} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}}{c-d\csc(e+fx+\frac{\pi}{2})} dx$$

↓ 4455

$$-\frac{2a \int \frac{1}{a(c-d) - \frac{a^2 d \tan^2(e+fx)}{\sec(e+fx)a+a}} d\left(-\frac{a \tan(e+fx)}{\sqrt{\sec(e+fx)a+a}}\right)}{f}$$

↓ 221

$$\frac{2\sqrt{a}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{c-d}\sqrt{a\sec(e+fx)+a}}\right)}{\sqrt{d}f\sqrt{c-d}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c - d*Sec[e + f*x]),x]`

output `(2*Sqrt[a]*ArcTanh[(Sqrt[a]*Sqrt[d]*Tan[e + f*x])/(Sqrt[c - d]*Sqrt[a + a*Sec[e + f*x]])])/(Sqrt[c - d]*Sqrt[d]*f)`

3.184.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4455 Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d + d*x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2,
0]
```

3.184.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 401 vs. 2(51) = 102.

Time = 18.88 (sec) , antiderivative size = 402, normalized size of antiderivative = 6.18

method	result
default	$\left(\ln \left(\frac{2 \left(\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \sqrt{-\frac{2d}{c+d}} c + \sqrt{-\frac{2d}{c+d}} \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} d + \sqrt{(c+d)(c-d)} (-\cot(fx+e) + \csc(fx+e)) - c}{-c(-\cot(fx+e) + \csc(fx+e)) - (-\cot(fx+e) + \csc(fx+e))d + \sqrt{(c+d)(c-d)}} \right) \right)$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-d*sec(f*x+e)),x,method=_RETURNVER
BOSE)
```

```
output 1/f/(-2*d/(c+d))^(1/2)/((c+d)*(c-d))^(1/2)*(ln(-2*((1-cos(f*x+e))^2*csc(f
*x+e)^2-1)^(1/2)*(-2*d/(c+d))^(1/2)*c+(-2*d/(c+d))^(1/2)*((1-cos(f*x+e))^2
*csc(f*x+e)^2-1)^(1/2)*d+((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))-c-d
/(-c*(-cot(f*x+e)+csc(f*x+e))-(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/
2)))-ln(-2*(-((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(-2*d/(c+d))^(1/2)*c-
(-2*d/(c+d))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*d+((c+d)*(c-d))
^(1/2)*(-cot(f*x+e)+csc(f*x+e))+c+d)/(c*(-cot(f*x+e)+csc(f*x+e))+(-cot(f*x
+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2))))*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)
^(1/2)*(-2*a/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)
```

3.184. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx$

3.184.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(51) = 102.

Time = 0.42 (sec) , antiderivative size = 357, normalized size of antiderivative = 5.49

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx$$

$$= \left[\frac{\sqrt{\frac{a}{cd-d^2}} \log\left(-\frac{(ac^2-8acd+8ad^2)\cos(fx+e)^3+ad^2+(ac^2-2acd)\cos(fx+e)^2+4((c^2d-3cd^2+2d^3)\cos(fx+e)^2+(cd^2-d^3)\cos(fx+e))}{c^2\cos(fx+e)^3+(c^2-2cd)\cos(fx+e)^2+d^2-(2cd-d^2)\cos(fx+e)}\right)}{2f} \right.$$

$$\left. - \frac{\sqrt{-\frac{a}{cd-d^2}} \arctan\left(\frac{2(cd-d^2)\sqrt{-\frac{a}{cd-d^2}}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\cos(fx+e)\sin(fx+e)}{(ac-2ad)\cos(fx+e)^2+ad+(ac-ad)\cos(fx+e)}\right)}{f} \right]$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-d*sec(f*x+e)),x, algorithm="fricas")`

output `[1/2*sqrt(a/(c*d - d^2))*log(-((a*c^2 - 8*a*c*d + 8*a*d^2)*cos(f*x + e)^3 + a*d^2 + (a*c^2 - 2*a*c*d)*cos(f*x + e)^2 + 4*((c^2*d - 3*c*d^2 + 2*d^3)*cos(f*x + e)^2 + (c*d^2 - d^3)*cos(f*x + e))*sqrt(a/(c*d - d^2))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sin(f*x + e) + (6*a*c*d - 7*a*d^2)*cos(f*x + e))/(c^2*cos(f*x + e)^3 + (c^2 - 2*c*d)*cos(f*x + e)^2 + d^2 - (2*c*d - d^2)*cos(f*x + e)))/f, -sqrt(-a/(c*d - d^2))*arctan(2*(c*d - d^2)*sqrt(-a/(c*d - d^2))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)/((a*c - 2*a*d)*cos(f*x + e)^2 + a*d + (a*c - a*d)*cos(f*x + e)))/f]`

3.184.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx = \int \frac{\sqrt{a(\sec(e+fx)+1)}\sec(e+fx)}{c-d\sec(e+fx)} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(1/2)/(c-d*sec(f*x+e)),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*sec(e + f*x)/(c - d*sec(e + f*x)), x)`

3.184. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx$

3.184.7 Maxima [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx = \int -\frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{d\sec(fx+e)-c} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-d*sec(f*x+e)),x, algorithm="maxima")`

output `-integrate(sqrt(a*sec(f*x + e) + a)*sec(f*x + e)/(d*sec(f*x + e) - c), x)`

3.184.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx = \int -\frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{d\sec(fx+e)-c} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c-d*sec(f*x+e)),x, algorithm="giac")`

output `sage0*x`

3.184.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c-d\sec(e+fx)} dx = -\int \frac{\sqrt{a+\frac{a}{\cos(e+fx)}}}{d-c\cos(e+fx)} dx$$

input `int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c - d/cos(e + f*x))),x)`

output `-int((a + a/cos(e + f*x))^(1/2)/(d - c*cos(e + f*x)), x)`

3.185 $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$

3.185.1 Optimal result	1263
3.185.2 Mathematica [A] (verified)	1264
3.185.3 Rubi [A] (verified)	1264
3.185.4 Maple [A] (verified)	1268
3.185.5 Fricas [A] (verification not implemented)	1269
3.185.6 Sympy [F]	1269
3.185.7 Maxima [A] (verification not implemented)	1270
3.185.8 Giac [B] (verification not implemented)	1271
3.185.9 Mupad [B] (verification not implemented)	1272

3.185.1 Optimal result

Integrand size = 29, antiderivative size = 236

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{a(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4) \operatorname{arctanh}(\sin(e + fx))}{8f}$$

$$+ \frac{a(12c^4 + 95c^3d + 112c^2d^2 + 80cd^3 + 16d^4) \tan(e + fx)}{30f}$$

$$+ \frac{ad(24c^3 + 130c^2d + 116cd^2 + 45d^3) \sec(e + fx) \tan(e + fx)}{120f}$$

$$+ \frac{a(12c^2 + 35cd + 16d^2) (c + d \sec(e + fx))^2 \tan(e + fx)}{60f}$$

$$+ \frac{a(4c + 5d)(c + d \sec(e + fx))^3 \tan(e + fx)}{20f} + \frac{a(c + d \sec(e + fx))^4 \tan(e + fx)}{5f}$$

output `1/8*a*(8*c^4+16*c^3*d+24*c^2*d^2+12*c*d^3+3*d^4)*arctanh(sin(f*x+e))/f+1/30*a*(12*c^4+95*c^3*d+112*c^2*d^2+80*c*d^3+16*d^4)*tan(f*x+e)/f+1/120*a*d*(24*c^3+130*c^2*d+116*c*d^2+45*d^3)*sec(f*x+e)*tan(f*x+e)/f+1/60*a*(12*c^2+35*c*d+16*d^2)*(c+d*sec(f*x+e))^2*tan(f*x+e)/f+1/20*a*(4*c+5*d)*(c+d*sec(f*x+e))^3*tan(f*x+e)/f+1/5*a*(c+d*sec(f*x+e))^4*tan(f*x+e)/f`

3.185.2 Mathematica [A] (verified)

Time = 1.80 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.65

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{a(15(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx)(120(c + d)^4 + 15d(16c^3 +$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^4,x]`

output `(a*(15*(8*c^4 + 16*c^3*d + 24*c^2*d^2 + 12*c*d^3 + 3*d^4)*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(120*(c + d)^4 + 15*d*(16*c^3 + 24*c^2*d + 12*c*d^2 + 3*d^3)*Sec[e + f*x] + 30*d^3*(4*c + d)*Sec[e + f*x]^3 + 80*d^2*(3*c^2 + 2*c*d + d^2)*Tan[e + f*x]^2 + 24*d^4*Tan[e + f*x]^4))/(120*f)`

3.185.3 Rubi [A] (verified)

Time = 1.40 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$, Rules used = {3042, 4490, 3042, 4490, 3042, 4490, 3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)(c + d \sec(e + fx))^4 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^4 dx$$

$$\downarrow \text{4490}$$

$$\frac{1}{5} \int \sec(e + fx)(c + d \sec(e + fx))^3(a(5c + 4d) + a(4c + 5d) \sec(e + fx)) dx +$$

$$\frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f}$$

$$\downarrow \text{3042}$$

3.185. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$

$$\frac{1}{5} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 \left(a(5c + 4d) + a(4c + 5d) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f}$$

↓ 4490

$$\frac{1}{5} \left(\frac{1}{4} \int \sec(e + fx)(c + d \sec(e + fx))^2 (a(20c^2 + 28dc + 15d^2) + a(12c^2 + 35dc + 16d^2) \sec(e + fx)) dx + \frac{a(4c + d \sec(e + fx))^4}{5f}\right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 (a(20c^2 + 28dc + 15d^2) + a(12c^2 + 35dc + 16d^2) \csc\left(e + fx + \frac{\pi}{2}\right)) dx + \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)$$

↓ 4490

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \sec(e + fx)(c + d \sec(e + fx)) (a(60c^3 + 108dc^2 + 115d^2c + 32d^3) + a(24c^3 + 130dc^2 + 116d^2c + 45d^3) \sec(e + fx)) dx + \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)\right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) (a(60c^3 + 108dc^2 + 115d^2c + 32d^3) + a(24c^3 + 130dc^2 + 116d^2c + 45d^3) \csc\left(e + fx + \frac{\pi}{2}\right)) dx + \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)\right)$$

↓ 4485

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \sec(e + fx) (15a(8c^4 + 16dc^3 + 24d^2c^2 + 12d^3c + 3d^4) + 4a(12c^4 + 95dc^3 + 112d^2c^2 + 80d^3c + 16d^4) \sec(e + fx)) dx + \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)\right)\right)$$

↓ 3042

3.185. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \csc \left(e + fx + \frac{\pi}{2} \right) \left(15a(8c^4 + 16dc^3 + 24d^2c^2 + 12d^3c + 3d^4) + 4a(12c^4 + 95dc^3 + 112d^2c^2 + 80d^3c + 3d^4) \right) dx \right. \right. \right. \\ \left. \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right) \right) \right) \downarrow 4274$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(4a(12c^4 + 95c^3d + 112c^2d^2 + 80cd^3 + 16d^4) \int \sec^2(e + fx) dx + 15a(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4) \right) \right. \right. \right. \\ \left. \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right) \right) \right) \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15a(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + 4a(12c^4 + 95c^3d + 112c^2d^2 + 80d^3c + 3d^4) \right) \right. \right. \right. \\ \left. \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right) \right) \right) \downarrow 4254$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15a(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx - \frac{4a(12c^4 + 95c^3d + 112c^2d^2 + 80d^3c + 3d^4)}{f} \right) \right. \right. \right. \\ \left. \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right) \right) \right) \downarrow 24$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15a(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{4a(12c^4 + 95c^3d + 112c^2d^2 + 80d^3c + 3d^4)}{f} \right) \right. \right. \right. \\ \left. \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right) \right) \right) \downarrow 4257$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\frac{15a(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4) \operatorname{arctanh}(\sin(e + fx))}{f} + \frac{4a(12c^4 + 95c^3d + 112c^2d^2 + 80d^3c + 3d^4)}{f} \right) \right. \right. \right. \\ \left. \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right) \right) \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^4,x]`

3.185. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$

output $(a*(c + d*\text{Sec}[e + f*x])^4*\text{Tan}[e + f*x])/(5*f) + ((a*(4*c + 5*d)*(c + d*\text{Sec}[e + f*x])^3*\text{Tan}[e + f*x])/(4*f) + ((a*(12*c^2 + 35*c*d + 16*d^2)*(c + d*\text{Sec}[e + f*x])^2*\text{Tan}[e + f*x])/(3*f) + ((a*d*(24*c^3 + 130*c^2*d + 116*c*d^2 + 45*d^3)*\text{Sec}[e + f*x]*\text{Tan}[e + f*x])/(2*f) + ((15*a*(8*c^4 + 16*c^3*d + 24*c^2*d^2 + 12*c*d^3 + 3*d^4)*\text{ArcTanh}[\text{Sin}[e + f*x]])/f + (4*a*(12*c^4 + 95*c^3*d + 112*c^2*d^2 + 80*c*d^3 + 16*d^4)*\text{Tan}[e + f*x])/f)/2)/3)/4)/5$

3.185.3.1 Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ /; FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 4254 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[-d^{(-1)} \text{ Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] \text{ /; FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

rule 4257 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] \text{ /; FreeQ}[\{c, d\}, x]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{ Int}[(d*\text{Csc}[e + f*x])^{(n + 1)}, x], x] \text{ /; FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4485 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))*(\text{csc}[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*((d*\text{Csc}[e + f*x])^n/(f*(n + 1))), x] + \text{Simp}[1/(n + 1) \text{ Int}[(d*\text{Csc}[e + f*x])^n*\text{Simp}[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*\text{Csc}[e + f*x], x], x], x] \text{ /; FreeQ}[\{a, b, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[A*b - a*B, 0] \ \&\& \ !\text{LeQ}[n, -1]$

```
rule 4490 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-B)*Cot[e + f*x]*((
a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[Csc[e + f*x]*
(a + b*Csc[e + f*x])^(m - 1)*Simp[b*B*m + a*A*(m + 1) + (a*B*m + A*b*(m +
1))*Csc[e + f*x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*
B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0]
```

3.185.4 Maple [A] (verified)

Time = 5.16 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.00

method	result
parts	$\frac{(4ac d^3 + a d^4) \left(- \left(- \frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f} - \frac{(6a^2 c^2 d^2 + 4ac d^3) \left(- \frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) + 4ac d^3}{f}$
derivativedivides	$\frac{a c^4 \tan(fx+e) + 4a c^3 d \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) - 6a c^2 d^2 \left(- \frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) + 4ac d^3}{f}$
default	$\frac{a c^4 \tan(fx+e) + 4a c^3 d \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) - 6a c^2 d^2 \left(- \frac{2}{3} - \frac{\sec(fx+e)^2}{3} \right) \tan(fx+e) + 4ac d^3}{f}$
norman	$- \frac{a(8c^4 + 16c^3 d + 24c^2 d^2 + 12c d^3 + 3d^4) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{4f} - \frac{a(8c^4 + 48c^3 d + 72c^2 d^2 + 52c d^3 + 13d^4) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f} - \frac{4a(45c^4 + 180c^3 d + 150c^2 d^2 + 60c d^3 + 10d^4)}{4f}$
parallelrisch	$2a \left(-5(2c^3 d + 3c^2 d^2 + \frac{3}{2} c d^3 + \frac{3}{8} d^4 + c^4) \left(\frac{\cos(5fx+5e)}{10} + \frac{\cos(3fx+3e)}{2} + \cos(fx+e) \right) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) + 5(2c^3 d + 3c^2 d^2 + \frac{3}{2} c d^3 + \frac{3}{8} d^4 + c^4) \right)$
risch	$\frac{ia(64d^4 + 120c^4 - 720c^2 d^2 e^{7i(fx+e)} + 960c d^3 e^{6i(fx+e)} + 2880c^3 d e^{4i(fx+e)} - 180c d^3 e^{9i(fx+e)} + 480c^3 d e^{8i(fx+e)} + 1440c^2 d^2 e^{7i(fx+e)} + 1440c^2 d^2 e^{6i(fx+e)} + 1440c^2 d^2 e^{5i(fx+e)} + 1440c^2 d^2 e^{4i(fx+e)} + 1440c^2 d^2 e^{3i(fx+e)} + 1440c^2 d^2 e^{2i(fx+e)} + 1440c^2 d^2 e^{i(fx+e)} + 1440c^2 d^2)}{f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^4,x,method=_RETURNVERBOSE)
)
```

```
output (4*a*c*d^3+a*d^4)/f*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln
(sec(f*x+e)+tan(f*x+e)))-(6*a*c^2*d^2+4*a*c*d^3)/f*(-2/3-1/3*sec(f*x+e)^2)
*tan(f*x+e)+(4*a*c^3*d+6*a*c^2*d^2)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(se
c(f*x+e)+tan(f*x+e)))+(a*c^4+4*a*c^3*d)/f*tan(f*x+e)+a*c^4/f*ln(sec(f*x+e)
+tan(f*x+e))-a*d^4/f*(-8/15-1/5*sec(f*x+e)^4-4/15*sec(f*x+e)^2)*tan(f*x+e)
```

3.185. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$

3.185.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.19

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{15(8ac^4 + 16ac^3d + 24ac^2d^2 + 12acd^3 + 3ad^4) \cos(fx + e)^5 \log(\sin(fx + e) + 1) - 15(8ac^4 + 16ac^3d + 24ac^2d^2 + 12acd^3 + 3ad^4) \cos(fx + e)^5 \log(-\sin(fx + e) + 1) + 2(24a^2d^4 + 8(15ac^4 + 60ac^3d + 60ac^2d^2 + 40acd^3 + 8ad^4) \cos(fx + e)^4 + 15(16ac^3d + 24ac^2d^2 + 12acd^3 + 3ad^4) \cos(fx + e)^3 + 16(15ac^2d^2 + 10acd^3 + 2ad^4) \cos(fx + e)^2 + 30(4ac^2d^3 + ad^4) \cos(fx + e)) \sin(fx + e)}{(f \cos(fx + e))^5}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^4,x, algorithm="fricas")`

output `1/240*(15*(8*a*c^4 + 16*a*c^3*d + 24*a*c^2*d^2 + 12*a*c*d^3 + 3*a*d^4)*cos(f*x + e)^5*log(sin(f*x + e) + 1) - 15*(8*a*c^4 + 16*a*c^3*d + 24*a*c^2*d^2 + 12*a*c*d^3 + 3*a*d^4)*cos(f*x + e)^5*log(-sin(f*x + e) + 1) + 2*(24*a*d^4 + 8*(15*a*c^4 + 60*a*c^3*d + 60*a*c^2*d^2 + 40*a*c*d^3 + 8*a*d^4)*cos(f*x + e)^4 + 15*(16*a*c^3*d + 24*a*c^2*d^2 + 12*a*c*d^3 + 3*a*d^4)*cos(f*x + e)^3 + 16*(15*a*c^2*d^2 + 10*a*c*d^3 + 2*a*d^4)*cos(f*x + e)^2 + 30*(4*a*c*d^3 + a*d^4)*cos(f*x + e))*sin(f*x + e)/(f*cos(f*x + e))^5)`

3.185.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= a \left(\int c^4 \sec(e + fx) dx + \int c^4 \sec^2(e + fx) dx + \int d^4 \sec^5(e + fx) dx \right.$$

$$+ \int d^4 \sec^6(e + fx) dx + \int 4cd^3 \sec^4(e + fx) dx + \int 4cd^3 \sec^5(e + fx) dx$$

$$+ \int 6c^2d^2 \sec^3(e + fx) dx + \int 6c^2d^2 \sec^4(e + fx) dx + \int 4c^3d \sec^2(e + fx) dx$$

$$\left. + \int 4c^3d \sec^3(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))**4,x)`

```
output a*(Integral(c**4*sec(e + f*x), x) + Integral(c**4*sec(e + f*x)**2, x) + In
tegral(d**4*sec(e + f*x)**5, x) + Integral(d**4*sec(e + f*x)**6, x) + Inte
gral(4*c*d**3*sec(e + f*x)**4, x) + Integral(4*c*d**3*sec(e + f*x)**5, x)
+ Integral(6*c**2*d**2*sec(e + f*x)**3, x) + Integral(6*c**2*d**2*sec(e +
f*x)**4, x) + Integral(4*c**3*d*sec(e + f*x)**2, x) + Integral(4*c**3*d*se
c(e + f*x)**3, x))
```

3.185.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.61

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{480 (\tan (fx + e)^3 + 3 \tan (fx + e)) ac^2 d^2 + 320 (\tan (fx + e)^3 + 3 \tan (fx + e)) acd^3 + 16 (3 \tan (fx + e)^3 + 3 \tan (fx + e)) a^2 c^2 d^2 + 16 (3 \tan (fx + e)^3 + 3 \tan (fx + e)) a^2 c d^3 + 16 (3 \tan (fx + e)^3 + 3 \tan (fx + e)) a^2 d^3}{1}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^4,x, algorithm="max
ima")
```

```
output 1/240*(480*(tan(f*x + e)^3 + 3*tan(f*x + e))*a*c^2*d^2 + 320*(tan(f*x + e)
^3 + 3*tan(f*x + e))*a*c*d^3 + 16*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 +
15*tan(f*x + e))*a*d^4 - 60*a*c*d^3*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))
/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log
(sin(f*x + e) - 1)) - 15*a*d^4*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin
(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(
f*x + e) - 1)) - 240*a*c^3*d*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(si
n(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 360*a*c^2*d^2*(2*sin(f*x + e)/(
sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 240
*a*c^4*log(sec(f*x + e) + tan(f*x + e)) + 240*a*c^4*tan(f*x + e) + 960*a*c
^3*d*tan(f*x + e))/f
```

3.185.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 566 vs. $2(224) = 448$.

Time = 0.39 (sec) , antiderivative size = 566, normalized size of antiderivative = 2.40

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{15(8ac^4 + 16ac^3d + 24ac^2d^2 + 12acd^3 + 3ad^4) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 15(8ac^4 + 16ac^3d + 24ac^2d^2 + 12acd^3 + 3ad^4) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right) + 2(120a^2c^4 + 240a^2cd^3 + 360a^2d^2c^2 + 180a^2d^2c^3 + 45a^2d^4) \tan^9\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 180a^2cd^3 \tan^9\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 45a^2d^4 \tan^9\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 480a^2c^4 \tan^7\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1440a^2c^3d \tan^7\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1160a^2c^3d^2 \tan^7\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 130a^2d^4 \tan^7\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 720a^2c^4 \tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 2880a^2c^3d \tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 2400a^2c^2d^2 \tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1600a^2cd^3 \tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 464a^2d^4 \tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 480a^2c^4 \tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 2400a^2c^3d \tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 2640a^2c^2d^2 \tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1400a^2cd^3 \tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 190a^2d^4 \tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 120a^2c^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 720a^2c^3d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1080a^2c^2d^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 780a^2cd^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 195a^2d^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)\right)^2 - 1} \frac{1}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^4,x, algorithm="giac")`

output `1/120*(15*(8*a*c^4 + 16*a*c^3*d + 24*a*c^2*d^2 + 12*a*c*d^3 + 3*a*d^4)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*(8*a*c^4 + 16*a*c^3*d + 24*a*c^2*d^2 + 12*a*c*d^3 + 3*a*d^4)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(120*a*c^4*tan(1/2*f*x + 1/2*e)^9 + 240*a*c^3*d*tan(1/2*f*x + 1/2*e)^9 + 360*a*c^2*d^2*tan(1/2*f*x + 1/2*e)^9 + 180*a*c*d^3*tan(1/2*f*x + 1/2*e)^9 + 45*a*d^4*tan(1/2*f*x + 1/2*e)^9 - 480*a*c^4*tan(1/2*f*x + 1/2*e)^7 - 1440*a*c^3*d*tan(1/2*f*x + 1/2*e)^7 - 1200*a*c^2*d^2*tan(1/2*f*x + 1/2*e)^7 - 1160*a*c*d^3*tan(1/2*f*x + 1/2*e)^7 - 130*a*d^4*tan(1/2*f*x + 1/2*e)^7 + 720*a*c^4*tan(1/2*f*x + 1/2*e)^5 + 2880*a*c^3*d*tan(1/2*f*x + 1/2*e)^5 + 2400*a*c^2*d^2*tan(1/2*f*x + 1/2*e)^5 + 1600*a*c*d^3*tan(1/2*f*x + 1/2*e)^5 + 464*a*d^4*tan(1/2*f*x + 1/2*e)^5 - 480*a*c^4*tan(1/2*f*x + 1/2*e)^3 - 2400*a*c^3*d*tan(1/2*f*x + 1/2*e)^3 - 2640*a*c^2*d^2*tan(1/2*f*x + 1/2*e)^3 - 1400*a*c*d^3*tan(1/2*f*x + 1/2*e)^3 - 190*a*d^4*tan(1/2*f*x + 1/2*e)^3 + 120*a*c^4*tan(1/2*f*x + 1/2*e) + 720*a*c^3*d*tan(1/2*f*x + 1/2*e) + 1080*a*c^2*d^2*tan(1/2*f*x + 1/2*e) + 780*a*c*d^3*tan(1/2*f*x + 1/2*e) + 195*a*d^4*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^5)/f`

3.185.9 Mupad [B] (verification not implemented)

Time = 17.12 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.53

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{a \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4)}{2(4c^4 + 8c^3d + 12c^2d^2 + 6cd^3 + \frac{3d^4}{2})}\right)(8c^4 + 16c^3d + 24c^2d^2 + 12cd^3 + 3d^4)}{4f} \\ + \left(2ac^4 + 4ac^3d + 6ac^2d^2 + 3acd^3 + \frac{3ad^4}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9 + \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7 + \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right) + \left(-8ac^4 - 24ac^3d - 20ac^2d^2 - \frac{58acd}{3} - 5d^3\right)$$

input `int(((a + a/cos(e + f*x))*(c + d/cos(e + f*x))^4)/cos(e + f*x),x)`output `(a*atanh((tan(e/2 + (f*x)/2)*(12*c*d^3 + 16*c^3*d + 8*c^4 + 3*d^4 + 24*c^2*d^2))/(2*(6*c*d^3 + 8*c^3*d + 4*c^4 + (3*d^4)/2 + 12*c^2*d^2)))*(12*c*d^3 + 16*c^3*d + 8*c^4 + 3*d^4 + 24*c^2*d^2))/(4*f) - (tan(e/2 + (f*x)/2)^9*(2*a*c^4 + (3*a*d^4)/4 + 6*a*c^2*d^2 + 3*a*c*d^3 + 4*a*c^3*d) - tan(e/2 + (f*x)/2)^7*(8*a*c^4 + (13*a*d^4)/6 + 20*a*c^2*d^2 + (58*a*c*d^3)/3 + 24*a*c^3*d) - tan(e/2 + (f*x)/2)^5*(8*a*c^4 + (19*a*d^4)/6 + 44*a*c^2*d^2 + (70*a*c*d^3)/3 + 40*a*c^3*d) + tan(e/2 + (f*x)/2)^3*(8*a*c^4 + (19*a*d^4)/6 + 44*a*c^2*d^2 + (70*a*c*d^3)/3 + 40*a*c^3*d) + tan(e/2 + (f*x)/2)^1*(8*a*c^4 + (19*a*d^4)/6 + 44*a*c^2*d^2 + (70*a*c*d^3)/3 + 40*a*c^3*d) + tan(e/2 + (f*x)/2)*(2*a*c^4 + (13*a*d^4)/4 + 18*a*c^2*d^2 + 13*a*c*d^3 + 12*a*c^3*d))/(f*(5*tan(e/2 + (f*x)/2)^2 - 10*tan(e/2 + (f*x)/2)^4 + 10*tan(e/2 + (f*x)/2)^6 - 5*tan(e/2 + (f*x)/2)^8 + tan(e/2 + (f*x)/2)^10 - 1))`

3.186 $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$

3.186.1 Optimal result	1273
3.186.2 Mathematica [A] (verified)	1274
3.186.3 Rubi [A] (verified)	1274
3.186.4 Maple [A] (verified)	1277
3.186.5 Fricas [A] (verification not implemented)	1278
3.186.6 Sympy [F]	1279
3.186.7 Maxima [A] (verification not implemented)	1279
3.186.8 Giac [B] (verification not implemented)	1280
3.186.9 Mupad [B] (verification not implemented)	1281

3.186.1 Optimal result

Integrand size = 29, antiderivative size = 171

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{a(8c^3 + 12c^2d + 12cd^2 + 3d^3) \operatorname{arctanh}(\sin(e + fx))}{8f}$$

$$+ \frac{a(3c^3 + 16c^2d + 12cd^2 + 4d^3) \tan(e + fx)}{6f}$$

$$+ \frac{ad(6c^2 + 20cd + 9d^2) \sec(e + fx) \tan(e + fx)}{24f}$$

$$+ \frac{a(3c + 4d)(c + d \sec(e + fx))^2 \tan(e + fx)}{12f} + \frac{a(c + d \sec(e + fx))^3 \tan(e + fx)}{4f}$$

output

```
1/8*a*(8*c^3+12*c^2*d+12*c*d^2+3*d^3)*arctanh(sin(f*x+e))/f+1/6*a*(3*c^3+1
6*c^2*d+12*c*d^2+4*d^3)*tan(f*x+e)/f+1/24*a*d*(6*c^2+20*c*d+9*d^2)*sec(f*x
+e)*tan(f*x+e)/f+1/12*a*(3*c+4*d)*(c+d*sec(f*x+e))^2*tan(f*x+e)/f+1/4*a*(c
+d*sec(f*x+e))^3*tan(f*x+e)/f
```

3.186.2 Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.60

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{a(3(8c^3 + 12c^2d + 12cd^2 + 3d^3) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx)(24(c + d)^3 + 9d(2c + d)^2 \sec(e + fx))}{24f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^3,x]`

output `(a*(3*(8*c^3 + 12*c^2*d + 12*c*d^2 + 3*d^3)*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(24*(c + d)^3 + 9*d*(2*c + d)^2*Sec[e + f*x] + 6*d^3*Sec[e + f*x]^3 + 8*d^2*(3*c + d)*Tan[e + f*x]^2))/(24*f)`

3.186.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 4490, 3042, 4490, 3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)(c + d \sec(e + fx))^3 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 dx$$

$$\downarrow \text{4490}$$

$$\frac{1}{4} \int \sec(e + fx)(c + d \sec(e + fx))^2 (a(4c + 3d) + a(3c + 4d) \sec(e + fx)) dx + \frac{a \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 \left(a(4c + 3d) + a(3c + 4d) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \frac{a \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

3.186. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$

↓ 4490

$$\frac{1}{4} \left(\frac{1}{3} \int \sec(e+fx)(c+d\sec(e+fx)) (a(12c^2+15dc+8d^2) + a(6c^2+20dc+9d^2)\sec(e+fx)) dx + \frac{a(3c+4d)}{4f} \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right) \left(a(12c^2+15dc+8d^2) + a(6c^2+20dc+9d^2)\csc\left(e+fx+\frac{\pi}{2}\right)\right) dx + \frac{a \tan(e+fx)(c+d\sec(e+fx))^3}{4f} \right)$$

↓ 4485

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \sec(e+fx) (3a(8c^3+12dc^2+12d^2c+3d^3) + 4a(3c^3+16dc^2+12d^2c+4d^3)\sec(e+fx)) dx + \frac{ad(6c^2+12cd+4d^2)}{4f} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(3a(8c^3+12dc^2+12d^2c+3d^3) + 4a(3c^3+16dc^2+12d^2c+4d^3)\csc\left(e+fx+\frac{\pi}{2}\right)\right) dx + \frac{a \tan(e+fx)(c+d\sec(e+fx))^3}{4f} \right) \right)$$

↓ 4274

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(4a(3c^3+16c^2d+12cd^2+4d^3) \int \sec^2(e+fx) dx + 3a(8c^3+12c^2d+12cd^2+3d^3) \int \sec(e+fx) dx \right) + \frac{a \tan(e+fx)(c+d\sec(e+fx))^3}{4f} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3a(8c^3+12c^2d+12cd^2+3d^3) \int \csc\left(e+fx+\frac{\pi}{2}\right) dx + 4a(3c^3+16c^2d+12cd^2+4d^3) \int \csc\left(e+fx+\frac{\pi}{2}\right) dx \right) + \frac{a \tan(e+fx)(c+d\sec(e+fx))^3}{4f} \right) \right)$$

↓ 4254

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3a(8c^3 + 12c^2d + 12cd^2 + 3d^3) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx - \frac{4a(3c^3 + 16c^2d + 12cd^2 + 4d^3) \int 1d(-\tan(e + fx))}{f} \right. \right. \right. \\ \left. \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^3}{4f} \right) \right) \right)$$

↓ 24

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3a(8c^3 + 12c^2d + 12cd^2 + 3d^3) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{4a(3c^3 + 16c^2d + 12cd^2 + 4d^3) \tan(e + fx)}{f} \right) \right. \right. \\ \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^3}{4f} \right) \right)$$

↓ 4257

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\frac{3a(8c^3 + 12c^2d + 12cd^2 + 3d^3) \operatorname{arctanh}(\sin(e + fx))}{f} + \frac{4a(3c^3 + 16c^2d + 12cd^2 + 4d^3) \tan(e + fx)}{f} \right) \right. \right. \\ \left. \left. \frac{a \tan(e + fx)(c + d \sec(e + fx))^3}{4f} \right) \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^3,x]`

output `(a*(c + d*Sec[e + f*x])^3*Tan[e + f*x])/(4*f) + ((a*(3*c + 4*d)*(c + d*Sec[e + f*x])^2*Tan[e + f*x])/(3*f) + ((a*d*(6*c^2 + 20*c*d + 9*d^2)*Sec[e + f*x]*Tan[e + f*x])/(2*f) + ((3*a*(8*c^3 + 12*c^2*d + 12*c*d^2 + 3*d^3)*ArcTanh[Sin[e + f*x]])/f + (4*a*(3*c^3 + 16*c^2*d + 12*c*d^2 + 4*d^3)*Tan[e + f*x])/f)/2)/3)/4`

3.186.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^(n_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4485 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^(n_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)*(csc[(e_.) + (f_.)*(x_)])*(B_.) + (A_.), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`

rule 4490 `Int[csc[(e_.) + (f_.)*(x_)])*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_.)*(csc[(e_.) + (f_.)*(x_)])*(B_.) + (A_.), x_Symbol] := Simp[(-B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*Simp[b*B*m + a*A*(m + 1) + (a*B*m + A*b*(m + 1))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0]`

3.186.4 Maple [A] (verified)

Time = 4.00 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.08

method	result
parts	$-\frac{(3ac^2d+ad^3)\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)}{f} + \frac{(3ac^2d+3acd^2)\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)}{f}$
derivatividevides	$\frac{ac^3\tan(fx+e)+3ac^2d\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)-3acd^2\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+ad^3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)}{f}$
default	$\frac{ac^3\tan(fx+e)+3ac^2d\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)-3acd^2\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+ad^3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)}{f}$
parallelrisch	$2\left(-2\left(\frac{3}{4} + \frac{\cos(4fx+4e)}{4} + \cos(2fx+2e)\right)\left(\frac{3}{2}c^2d + \frac{3}{2}cd^2 + \frac{3}{8}d^3 + c^3\right)\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 2\left(\frac{3}{4} + \frac{\cos(4fx+4e)}{4} + \cos(2fx+2e)\right)\left(\frac{3}{2}c^2d + \frac{3}{2}cd^2 + \frac{3}{8}d^3 + c^3\right)\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)\right)$
norman	$\frac{-\frac{a(8c^3+12c^2d+12cd^2+3d^3)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{4f} + \frac{a(8c^3+36c^2d+36cd^2+13d^3)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f} + \frac{a(72c^3+180c^2d+84cd^2+49d^3)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{12f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^4}$
risch	$\frac{ia(24c^3+48cd^2+16d^3+72c^2d+36cd^2e^{i(fx+e)}+192cd^2e^{2i(fx+e)}+36c^2de^{i(fx+e)}+72c^2de^{6i(fx+e)}-36c^2de^{5i(fx+e)}-36c^2de^{4i(fx+e)}-36c^2de^{3i(fx+e)}-36c^2de^{2i(fx+e)}-36c^2de^{i(fx+e)}-36c^2d-36c^2)}{f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output -(3*a*c*d^2+a*d^3)/f*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)+(3*a*c^2*d+3*a*c*d^2)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+(a*c^3+3*a*c^2*d)/f*tan(f*x+e)+a*c^3/f*ln(sec(f*x+e)+tan(f*x+e))+a*d^3/f*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln(sec(f*x+e)+tan(f*x+e)))
```

3.186.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.23

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{3(8ac^3 + 12ac^2d + 12acd^2 + 3ad^3) \cos(fx + e)^4 \log(\sin(fx + e) + 1) - 3(8ac^3 + 12ac^2d + 12acd^2 + 3ad^3) \cos(fx + e)^4 \log(\sin(fx + e) - 1)}{f}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^3,x, algorithm="fracas")
```

3.186. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$

output $\frac{1}{48}(3(8ac^3 + 12a^2cd + 12acd^2 + 3ad^3)\cos(fx + e)^4 \log(\sin(fx + e) + 1) - 3(8ac^3 + 12a^2cd + 12acd^2 + 3ad^3)\cos(fx + e)^4 \log(-\sin(fx + e) + 1) + 2(6ad^3 + 8(3ac^3 + 9a^2cd + 6acd^2 + 2ad^3)\cos(fx + e)^3 + 9(4a^2cd + 4acd^2 + ad^3)\cos(fx + e)^2 + 8(3acd^2 + ad^3)\cos(fx + e))\sin(fx + e))/(f\cos(fx + e)^4)$

3.186.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx \\ &= a \left(\int c^3 \sec(e + fx) dx + \int c^3 \sec^2(e + fx) dx + \int d^3 \sec^4(e + fx) dx \right. \\ & \quad \left. + \int d^3 \sec^5(e + fx) dx + \int 3cd^2 \sec^3(e + fx) dx + \int 3cd^2 \sec^4(e + fx) dx \right. \\ & \quad \left. + \int 3c^2d \sec^2(e + fx) dx + \int 3c^2d \sec^3(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))**3,x)`

output `a*(Integral(c**3*sec(e + f*x), x) + Integral(c**3*sec(e + f*x)**2, x) + Integral(d**3*sec(e + f*x)**4, x) + Integral(d**3*sec(e + f*x)**5, x) + Integral(3*c*d**2*sec(e + f*x)**3, x) + Integral(3*c*d**2*sec(e + f*x)**4, x) + Integral(3*c**2*d*sec(e + f*x)**2, x) + Integral(3*c**2*d*sec(e + f*x)**3, x))`

3.186.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.56

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx \\ &= \frac{48(\tan(fx + e)^3 + 3 \tan(fx + e))acd^2 + 16(\tan(fx + e)^3 + 3 \tan(fx + e))ad^3 - 3ad^3 \left(\frac{2(3 \sin(fx + e)^3 - \sin(fx + e)^4 - 2 \sin(fx + e))}{\sin(fx + e)^4 - 2 \sin(fx + e)^3} \right)}{\dots} \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output $\frac{1}{48}(48(\tan(fx + e))^3 + 3\tan(fx + e))a^2cd^2 + 16(\tan(fx + e))^3 + 3\tan(fx + e)a^2d^3 - 3a^2d^3(2(3\sin(fx + e)^3 - 5\sin(fx + e)))/(\sin(fx + e)^4 - 2\sin(fx + e)^2 + 1) - 3\log(\sin(fx + e) + 1) + 3\log(\sin(fx + e) - 1) - 36a^2cd^2(2\sin(fx + e))/(\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1) - 36a^2cd^2(2\sin(fx + e))/(\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1) + 48a^2c^3\log(\sec(fx + e) + \tan(fx + e)) + 48a^2c^3\tan(fx + e) + 144a^2cd^2\tan(fx + e))/f$

3.186.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 380 vs. $2(161) = 322$.

Time = 0.36 (sec) , antiderivative size = 380, normalized size of antiderivative = 2.22

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{3(8ac^3 + 12ac^2d + 12acd^2 + 3ad^3) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3(8ac^3 + 12ac^2d + 12acd^2 + 3ad^3) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right) + 2(24a^2c^3\tan^7\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 36a^2cd^2\tan^7\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 9a^2d^3\tan^7\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 72a^2c^3\tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 180a^2cd^2\tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 84a^2cd^2\tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 49a^2d^3\tan^5\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 72a^2c^3\tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 252a^2cd^2\tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 156a^2cd^2\tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 31a^2d^3\tan^3\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 24a^2c^3\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 108a^2cd^2\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 108a^2cd^2\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 39a^2d^3\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right))}{(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 1)^4}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output $\frac{1}{24}(3(8a^2c^3 + 12a^2c^2d + 12a^2cd^2 + 3a^2d^3)\log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1) - 3(8a^2c^3 + 12a^2c^2d + 12a^2cd^2 + 3a^2d^3)\log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1) - 2(24a^2c^3\tan^7(\frac{1}{2}fx + \frac{1}{2}e) + 36a^2cd^2\tan^7(\frac{1}{2}fx + \frac{1}{2}e) + 9a^2d^3\tan^7(\frac{1}{2}fx + \frac{1}{2}e) - 72a^2c^3\tan^5(\frac{1}{2}fx + \frac{1}{2}e) - 180a^2cd^2\tan^5(\frac{1}{2}fx + \frac{1}{2}e) - 84a^2cd^2\tan^5(\frac{1}{2}fx + \frac{1}{2}e) - 49a^2d^3\tan^5(\frac{1}{2}fx + \frac{1}{2}e) + 72a^2c^3\tan^3(\frac{1}{2}fx + \frac{1}{2}e) + 252a^2cd^2\tan^3(\frac{1}{2}fx + \frac{1}{2}e) + 156a^2cd^2\tan^3(\frac{1}{2}fx + \frac{1}{2}e) + 31a^2d^3\tan^3(\frac{1}{2}fx + \frac{1}{2}e) - 24a^2c^3\tan(\frac{1}{2}fx + \frac{1}{2}e) - 108a^2cd^2\tan(\frac{1}{2}fx + \frac{1}{2}e) - 108a^2cd^2\tan(\frac{1}{2}fx + \frac{1}{2}e) - 39a^2d^3\tan(\frac{1}{2}fx + \frac{1}{2}e))/(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)^4)/f$

3.186. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$

3.186.9 Mupad [B] (verification not implemented)

Time = 17.05 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.49

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{\left(-2ac^3 - 3ac^2d - 3acd^2 - \frac{3ad^3}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7 + \left(6ac^3 + 15ac^2d + 7acd^2 + \frac{49ad^3}{12}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6\right)}$$

$$+ \frac{a \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (8c^3 + 12c^2d + 12cd^2 + 3d^3)}{2(4c^3 + 6c^2d + 6cd^2 + \frac{3d^3}{2})}\right)}{4f} (8c^3 + 12c^2d + 12cd^2 + 3d^3)$$

input `int(((a + a/cos(e + f*x))*(c + d/cos(e + f*x))^3)/cos(e + f*x),x)`output `(tan(e/2 + (f*x)/2)*(2*a*c^3 + (13*a*d^3)/4 + 9*a*c*d^2 + 9*a*c^2*d) - tan(e/2 + (f*x)/2)^7*(2*a*c^3 + (3*a*d^3)/4 + 3*a*c*d^2 + 3*a*c^2*d) - tan(e/2 + (f*x)/2)^3*(6*a*c^3 + (31*a*d^3)/12 + 13*a*c*d^2 + 21*a*c^2*d) + tan(e/2 + (f*x)/2)^5*(6*a*c^3 + (49*a*d^3)/12 + 7*a*c*d^2 + 15*a*c^2*d))/(f*(6*tan(e/2 + (f*x)/2)^4 - 4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/2)^8 + 1)) + (a*atanh((tan(e/2 + (f*x)/2)*(12*c*d^2 + 12*c^2*d + 8*c^3 + 3*d^3))/(2*(6*c*d^2 + 6*c^2*d + 4*c^3 + (3*d^3)/2)))*(12*c*d^2 + 12*c^2*d + 8*c^3 + 3*d^3))/(4*f)`

3.187 $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$

3.187.1 Optimal result	1282
3.187.2 Mathematica [A] (verified)	1282
3.187.3 Rubi [A] (verified)	1283
3.187.4 Maple [A] (verified)	1286
3.187.5 Fricas [A] (verification not implemented)	1286
3.187.6 Sympy [F]	1287
3.187.7 Maxima [A] (verification not implemented)	1287
3.187.8 Giac [B] (verification not implemented)	1288
3.187.9 Mupad [B] (verification not implemented)	1288

3.187.1 Optimal result

Integrand size = 29, antiderivative size = 108

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{a(2c^2 + 2cd + d^2) \operatorname{arctanh}(\sin(e + fx))}{2f} + \frac{2a(c^2 + 3cd + d^2) \tan(e + fx)}{3f}$$

$$+ \frac{ad(2c + 3d) \sec(e + fx) \tan(e + fx)}{6f} + \frac{a(c + d \sec(e + fx))^2 \tan(e + fx)}{3f}$$

```
output 1/2*a*(2*c^2+2*c*d+d^2)*arctanh(sin(f*x+e))/f+2/3*a*(c^2+3*c*d+d^2)*tan(f*x+e)/f+1/6*a*d*(2*c+3*d)*sec(f*x+e)*tan(f*x+e)/f+1/3*a*(c+d*sec(f*x+e))^2*tan(f*x+e)/f
```

3.187.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.69

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{a(3(2c^2 + 2cd + d^2) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx) (3d(2c + d) \sec(e + fx) + 2(3(c + d)^2 + d^2 \tan^2(e + fx))))}{6f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^2,x]`

output `(a*(3*(2*c^2 + 2*c*d + d^2)*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(3*d*(2*c + d)*Sec[e + f*x] + 2*(3*(c + d)^2 + d^2*Tan[e + f*x]^2)))/(6*f)`

3.187.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4490, 3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)(c + d \sec(e + fx))^2 dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 dx \\
 & \quad \downarrow \text{4490} \\
 & \frac{1}{3} \int \sec(e + fx)(c + d \sec(e + fx))(a(3c + 2d) + a(2c + 3d) \sec(e + fx)) dx + \\
 & \quad \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(a(3c + 2d) + a(2c + 3d) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \\
 & \quad \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f} \\
 & \quad \downarrow \text{4485} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \sec(e + fx) (3a(2c^2 + 2dc + d^2) + 4a(c^2 + 3dc + d^2) \sec(e + fx)) dx + \frac{ad(2c + 3d) \tan(e + fx) \sec(e + fx)}{2f} \right) \\
 & \quad \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.187. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$

$$\frac{1}{3} \left(\frac{1}{2} \int \csc \left(e + fx + \frac{\pi}{2} \right) \left(3a(2c^2 + 2dc + d^2) + 4a(c^2 + 3dc + d^2) \csc \left(e + fx + \frac{\pi}{2} \right) \right) dx + \frac{ad(2c + 3d) \tan(e + fx)}{2f} \right) \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 4274

$$\frac{1}{3} \left(\frac{1}{2} \left(4a(c^2 + 3cd + d^2) \int \sec^2(e + fx) dx + 3a(2c^2 + 2cd + d^2) \int \sec(e + fx) dx \right) + \frac{ad(2c + 3d) \tan(e + fx) \sec(e + fx)}{2f} \right) \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3a(2c^2 + 2cd + d^2) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + 4a(c^2 + 3cd + d^2) \int \csc \left(e + fx + \frac{\pi}{2} \right)^2 dx \right) + \frac{ad(2c + 3d) \tan(e + fx)}{2f} \right) \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 4254

$$\frac{1}{3} \left(\frac{1}{2} \left(3a(2c^2 + 2cd + d^2) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx - \frac{4a(c^2 + 3cd + d^2) \int 1d(-\tan(e + fx))}{f} \right) + \frac{ad(2c + 3d) \tan(e + fx)}{2f} \right) \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 24

$$\frac{1}{3} \left(\frac{1}{2} \left(3a(2c^2 + 2cd + d^2) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{4a(c^2 + 3cd + d^2) \tan(e + fx)}{f} \right) + \frac{ad(2c + 3d) \tan(e + fx)}{2f} \right) \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{3a(2c^2 + 2cd + d^2) \operatorname{arctanh}(\sin(e + fx))}{f} + \frac{4a(c^2 + 3cd + d^2) \tan(e + fx)}{f} \right) + \frac{ad(2c + 3d) \tan(e + fx) \sec(e + fx)}{2f} \right) \frac{a \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^2,x]`

output $(a*(c + d*\text{Sec}[e + f*x])^2*\text{Tan}[e + f*x]/(3*f) + ((a*d*(2*c + 3*d)*\text{Sec}[e + f*x]*\text{Tan}[e + f*x])/(2*f) + ((3*a*(2*c^2 + 2*c*d + d^2)*\text{ArcTanh}[\text{Sin}[e + f*x]])/f + (4*a*(c^2 + 3*c*d + d^2)*\text{Tan}[e + f*x])/f)/2)/3$

3.187.3.1 Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ ; FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 4254 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[-d^{(-1)} \text{ Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] \text{ ; FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

rule 4257 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{ Int}[(d*\text{Csc}[e + f*x])^{(n + 1)}, x], x] \text{ ; FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4485 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))*(\text{csc}[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*((d*\text{Csc}[e + f*x])^n/(f*(n + 1))), x] + \text{Simp}[1/(n + 1) \text{ Int}[(d*\text{Csc}[e + f*x])^n*\text{Simp}[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*\text{Csc}[e + f*x], x], x], x] \text{ ; FreeQ}[\{a, b, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[A*b - a*B, 0] \ \&\& \ !\text{LeQ}[n, -1]$

rule 4490 $\text{Int}[\text{csc}[(e_.) + (f_.)*(x_.)]*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^{(m_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-B)*\text{Cot}[e + f*x]*((a + b*\text{Csc}[e + f*x])^m/(f*(m + 1))), x] + \text{Simp}[1/(m + 1) \text{ Int}[\text{Csc}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m - 1)}*\text{Simp}[b*B*m + a*A*(m + 1) + (a*B*m + A*b*(m + 1))*\text{Csc}[e + f*x], x], x], x] \text{ ; FreeQ}[\{a, b, A, B, e, f\}, x] \ \&\& \ \text{NeQ}[A*b - a*B, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$

3.187.4 Maple [A] (verified)

Time = 3.18 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.09

method	result
parts	$\frac{(2acd+ad^2)\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)}{f} + \frac{(ac^2+2acd)\tan(fx+e)}{f} + \frac{ac^2\ln(\sec(fx+e)+\tan(fx+e))}{f}$
derivativedivides	$\frac{ac^2\tan(fx+e)+2acd\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) - ad^2\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + ac^2\ln(\sec(fx+e)+\tan(fx+e))}{f}$
default	$\frac{ac^2\tan(fx+e)+2acd\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) - ad^2\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + ac^2\ln(\sec(fx+e)+\tan(fx+e))}{f}$
norman	$\frac{-\frac{a(2c^2+2cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{f} - \frac{a(2c^2+6cd+3d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{f} + \frac{4a(3c^2+6cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3f}}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2-1\right)^3} - \frac{a(2c^2+2cd+d^2)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)\right)}{2f}$
parallelrisc	$2\left(\frac{3\left(cd+\frac{1}{2}d^2+c^2\right)\left(\cos(fx+e)+\frac{\cos(3fx+3e)}{3}\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{2} + \frac{3\left(cd+\frac{1}{2}d^2+c^2\right)\left(\cos(fx+e)+\frac{\cos(3fx+3e)}{3}\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{2}\right)$
risc	$\frac{ia(6cde^{5i(fx+e)}+3d^2e^{5i(fx+e)}-6c^2e^{4i(fx+e)}-12cde^{4i(fx+e)}-12c^2e^{2i(fx+e)}-24cde^{2i(fx+e)}-12d^2e^{2i(fx+e)}-6de^{2i(fx+e)})}{3f(1+e^{2i(fx+e)})^3}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output (2*a*c*d+a*d^2)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+(a*c^2+2*a*c*d)/f*tan(f*x+e)+a*c^2/f*ln(sec(f*x+e)+tan(f*x+e))-a*d^2/f*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)
```

3.187.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.39

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{3(2ac^2 + 2acd + ad^2) \cos(fx + e)^3 \log(\sin(fx + e) + 1) - 3(2ac^2 + 2acd + ad^2) \cos(fx + e)^3 \log(-\sin(fx + e) + 1)}{12f \cos(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^2,x, algorithm="fricas")
```

output $1/12*(3*(2*a*c^2 + 2*a*c*d + a*d^2)*\cos(f*x + e)^3*\log(\sin(f*x + e) + 1) - 3*(2*a*c^2 + 2*a*c*d + a*d^2)*\cos(f*x + e)^3*\log(-\sin(f*x + e) + 1) + 2*(2*a*d^2 + 2*(3*a*c^2 + 6*a*c*d + 2*a*d^2)*\cos(f*x + e)^2 + 3*(2*a*c*d + a*d^2)*\cos(f*x + e))*\sin(f*x + e))/(f*\cos(f*x + e)^3)$

3.187.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx \\ &= a \left(\int c^2 \sec(e + fx) dx + \int c^2 \sec^2(e + fx) dx + \int d^2 \sec^3(e + fx) dx \right. \\ & \quad \left. + \int d^2 \sec^4(e + fx) dx + \int 2cd \sec^2(e + fx) dx + \int 2cd \sec^3(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))**2,x)`

output `a*(Integral(c**2*sec(e + f*x), x) + Integral(c**2*sec(e + f*x)**2, x) + Integral(d**2*sec(e + f*x)**3, x) + Integral(d**2*sec(e + f*x)**4, x) + Integral(2*c*d*sec(e + f*x)**2, x) + Integral(2*c*d*sec(e + f*x)**3, x))`

3.187.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.53

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx \\ &= \frac{4(\tan(fx + e)^3 + 3 \tan(fx + e))ad^2 - 6acd \left(\frac{2 \sin(fx + e)}{\sin(fx + e)^2 - 1} - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1) \right) - 3a^2d^2(2 \sin(fx + e)/(\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1)) + 12a^2c^2 \log(\sec(fx + e) + \tan(fx + e)) + 12a^2c^2 \tan(fx + e) + 24a^2c*d*\tan(fx + e))/f} \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output $1/12*(4*(\tan(f*x + e)^3 + 3*\tan(f*x + e))*a*d^2 - 6*a*c*d*(2*\sin(f*x + e)/(\sin(f*x + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) - 3*a*d^2*(2*\sin(f*x + e)/(\sin(f*x + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) + 12*a*c^2*\log(\sec(f*x + e) + \tan(f*x + e)) + 12*a*c^2*\tan(f*x + e) + 24*a*c*d*\tan(f*x + e))/f$

3.187. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$

3.187.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 232 vs. $2(100) = 200$.

Time = 0.33 (sec) , antiderivative size = 232, normalized size of antiderivative = 2.15

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{3(2ac^2 + 2acd + ad^2) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3(2ac^2 + 2acd + ad^2) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output `1/6*(3*(2*a*c^2 + 2*a*c*d + a*d^2)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 3*(2*a*c^2 + 2*a*c*d + a*d^2)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(6*a*c^2*tan(1/2*f*x + 1/2*e)^5 + 6*a*c*d*tan(1/2*f*x + 1/2*e)^5 + 3*a*d^2*tan(1/2*f*x + 1/2*e)^5 - 12*a*c^2*tan(1/2*f*x + 1/2*e)^3 - 24*a*c*d*tan(1/2*f*x + 1/2*e)^3 - 4*a*d^2*tan(1/2*f*x + 1/2*e)^3 + 6*a*c^2*tan(1/2*f*x + 1/2*e) + 18*a*c*d*tan(1/2*f*x + 1/2*e) + 9*a*d^2*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^3)/f`

3.187.9 Mupad [B] (verification not implemented)

Time = 16.37 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.81

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{a \operatorname{atanh}\left(\frac{2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) (2c^2 + 2cd + d^2)}{4c^2 + 4cd + 2d^2}\right) (2c^2 + 2cd + d^2)}{f}$$

$$- \frac{(2ac^2 + 2acd + ad^2) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-4ac^2 - 8acd - \frac{4ad^2}{3}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + (2ac^2 + 6acd + 3ad^2) \tan\left(\frac{e}{2} + \frac{fx}{2}\right) - (2ac^2 + 2acd + ad^2) \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)}{f}$$

input `int(((a + a/cos(e + f*x))*(c + d/cos(e + f*x))^2)/cos(e + f*x), x)`

output $(a \operatorname{atanh}((2 \tan(e/2 + (f*x)/2) * (2*c*d + 2*c^2 + d^2)) / (4*c*d + 4*c^2 + 2*d^2)) * (2*c*d + 2*c^2 + d^2)) / f - (\tan(e/2 + (f*x)/2) * (2*a*c^2 + 3*a*d^2 + 6*a*c*d) + \tan(e/2 + (f*x)/2)^5 * (2*a*c^2 + a*d^2 + 2*a*c*d) - \tan(e/2 + (f*x)/2)^3 * (4*a*c^2 + (4*a*d^2)/3 + 8*a*c*d)) / (f * (3*\tan(e/2 + (f*x)/2)^2 - 3*\tan(e/2 + (f*x)/2)^4 + \tan(e/2 + (f*x)/2)^6 - 1))$

3.188 $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$

3.188.1 Optimal result	1290
3.188.2 Mathematica [A] (verified)	1290
3.188.3 Rubi [A] (verified)	1291
3.188.4 Maple [A] (verified)	1293
3.188.5 Fricas [A] (verification not implemented)	1294
3.188.6 Sympy [F]	1294
3.188.7 Maxima [A] (verification not implemented)	1295
3.188.8 Giac [B] (verification not implemented)	1295
3.188.9 Mupad [B] (verification not implemented)	1296

3.188.1 Optimal result

Integrand size = 27, antiderivative size = 56

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{a(2c + d)\operatorname{arctanh}(\sin(e + fx))}{2f} + \frac{a(c + d)\tan(e + fx)}{f} + \frac{ad \sec(e + fx)\tan(e + fx)}{2f}$$

output `1/2*a*(2*c+d)*arctanh(sin(f*x+e))/f+a*(c+d)*tan(f*x+e)/f+1/2*a*d*sec(f*x+e)*tan(f*x+e)/f`

3.188.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.34

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{a\operatorname{arctanh}(\sin(e + fx))}{f} + \frac{ad\operatorname{arctanh}(\sin(e + fx))}{2f}$$

$$+ \frac{a c \tan(e + fx)}{f} + \frac{ad \tan(e + fx)}{f} + \frac{ad \sec(e + fx)\tan(e + fx)}{2f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x]),x]`

output `(a*c*ArcTanh[Sin[e + f*x]])/f + (a*d*ArcTanh[Sin[e + f*x]])/(2*f) + (a*c*Tan[e + f*x])/f + (a*d*Tan[e + f*x])/f + (a*d*Sec[e + f*x]*Tan[e + f*x])/(2*f)`

3.188.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)(c + d \sec(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4485} \\
 & \frac{1}{2} \int \sec(e + fx)(a(2c + d) + 2a(c + d) \sec(e + fx)) dx + \frac{ad \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a(2c + d) + 2a(c + d) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \frac{ad \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{4274} \\
 & \frac{1}{2} \left(2a(c + d) \int \sec^2(e + fx) dx + a(2c + d) \int \sec(e + fx) dx \right) + \frac{ad \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(a(2c + d) \int \csc\left(e + fx + \frac{\pi}{2}\right) dx + 2a(c + d) \int \csc\left(e + fx + \frac{\pi}{2}\right)^2 dx \right) + \frac{ad \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{4254} \\
 & \frac{1}{2} \left(a(2c + d) \int \csc\left(e + fx + \frac{\pi}{2}\right) dx - \frac{2a(c + d) \int 1d(-\tan(e + fx))}{f} \right) + \frac{ad \tan(e + fx) \sec(e + fx)}{2f}
 \end{aligned}$$

$$\frac{1}{2} \left(a(2c+d) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{2a(c+d) \tan(e+fx)}{f} \right) + \frac{ad \tan(e+fx) \sec(e+fx)}{2f}$$

↓ 24

$$\frac{1}{2} \left(\frac{a(2c+d) \operatorname{arctanh}(\sin(e+fx))}{f} + \frac{2a(c+d) \tan(e+fx)}{f} \right) + \frac{ad \tan(e+fx) \sec(e+fx)}{2f}$$

↓ 4257

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])*(c + d*Sec[e + f*x]),x]`

output `(a*d*Sec[e + f*x]*Tan[e + f*x])/(2*f) + ((a*(2*c + d)*ArcTanh[Sin[e + f*x]])/f + (2*a*(c + d)*Tan[e + f*x])/f)/2`

3.188.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[t[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

3.188.4 Maple [A] (verified)

Time = 2.20 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.34

method	result
derivativedivides	$\frac{ac \tan(fx+e) + ad \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) + ac \ln(\sec(fx+e) + \tan(fx+e)) + ad \tan(fx+e)}{f}$
default	$\frac{ac \tan(fx+e) + ad \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) + ac \ln(\sec(fx+e) + \tan(fx+e)) + ad \tan(fx+e)}{f}$
parts	$\frac{(ac+ad) \tan(fx+e)}{f} + \frac{ac \ln(\sec(fx+e) + \tan(fx+e))}{f} + \frac{ad \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right)}{f}$
parallelrisc	$-\frac{\left(\left(c + \frac{d}{2} \right) (1 + \cos(2fx+2e)) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) - \left(c + \frac{d}{2} \right) (1 + \cos(2fx+2e)) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right) + (-c-d) \sin(2fx+2e) \right)}{f(1 + \cos(2fx+2e))}$
norman	$\frac{a(2c+3d) \tan \left(\frac{fx}{2} + \frac{e}{2} \right) - \frac{a(2c+d) \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^3}{\left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right)^2 - 1 \right)^2} - \frac{a(2c+d) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right)}{2f} + \frac{a(2c+d) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) + 1 \right)}{2f}$
risc	$-\frac{ia(d e^{3i(fx+e)} - 2 e^{2i(fx+e)} c - 2 d e^{2i(fx+e)} - d e^{i(fx+e)} - 2c - 2d)}{f(1 + e^{2i(fx+e)})^2} - \frac{ac \ln(e^{i(fx+e)} - i)}{f} - \frac{a \ln(e^{i(fx+e)} - i) d}{2f} + \frac{ac}{f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f*(a*c*tan(f*x+e)+a*d*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f
*x+e)))+a*c*ln(sec(f*x+e)+tan(f*x+e))+a*d*tan(f*x+e))
```

3.188. $\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$

3.188.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.71

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{(2ac + ad) \cos(fx + e)^2 \log(\sin(fx + e) + 1) - (2ac + ad) \cos(fx + e)^2 \log(-\sin(fx + e) + 1) + 2(ad^2 + a^2c) \sin(fx + e)}{4f \cos(fx + e)^2}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `1/4*((2*a*c + a*d)*cos(f*x + e)^2*log(sin(f*x + e) + 1) - (2*a*c + a*d)*cos(f*x + e)^2*log(-sin(f*x + e) + 1) + 2*(a*d + 2*(a*c + a*d)*cos(f*x + e))*sin(f*x + e))/(f*cos(f*x + e)^2)`

3.188.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= a \left(\int c \sec(e + fx) dx + \int c \sec^2(e + fx) dx + \int d \sec^2(e + fx) dx + \int d \sec^3(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e)),x)`

output `a*(Integral(c*sec(e + f*x), x) + Integral(c*sec(e + f*x)**2, x) + Integral(d*sec(e + f*x)**2, x) + Integral(d*sec(e + f*x)**3, x))`

3.188.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.57

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx =$$

$$\frac{ad \left(\frac{2 \sin(fx+e)}{\sin(fx+e)^2-1} - \log(\sin(fx+e) + 1) + \log(\sin(fx+e) - 1) \right) - 4ac \log(\sec(fx+e) + \tan(fx+e)) - 4ad \tan(fx+e)}{4f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `-1/4*(a*d*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 4*a*c*log(sec(f*x + e) + tan(f*x + e)) - 4*a*d*tan(f*x + e) - 4*a*d*tan(f*x + e))/f`

3.188.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(52) = 104.

Time = 0.30 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.21

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{(2ac + ad) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|) - (2ac + ad) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|) - \frac{2(2ac \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + ad \tan(\frac{1}{2}fx + \frac{1}{2}e))}{2f}}{2f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))*(c+d*sec(f*x+e)),x, algorithm="giac")`

output `1/2*((2*a*c + a*d)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - (2*a*c + a*d)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(2*a*c*tan(1/2*f*x + 1/2*e)^3 + a*d*tan(1/2*f*x + 1/2*e)^3 - 2*a*c*tan(1/2*f*x + 1/2*e) - 3*a*d*tan(1/2*f*x + 1/2*e)))/(tan(1/2*f*x + 1/2*e)^2 - 1)^2)/f`

3.188.9 Mupad [B] (verification not implemented)

Time = 14.43 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.98

$$\int \sec(e + fx)(a + a \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{a \operatorname{atanh}\left(\frac{2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)(2c+d)}{4c+2d}\right) (2c+d)}{f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (2ac + ad) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right) (2ac + 3ad)}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1\right)}$$

input `int(((a + a/cos(e + f*x))*(c + d/cos(e + f*x)))/cos(e + f*x),x)`output `(a*atanh((2*tan(e/2 + (f*x)/2)*(2*c + d))/(4*c + 2*d))*(2*c + d))/f - (tan(e/2 + (f*x)/2)^3*(2*a*c + a*d) - tan(e/2 + (f*x)/2)*(2*a*c + 3*a*d))/(f*(tan(e/2 + (f*x)/2)^4 - 2*tan(e/2 + (f*x)/2)^2 + 1))`

3.189 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c+d \sec(e+fx)} dx$

3.189.1 Optimal result 1297
 3.189.2 Mathematica [A] (verified) 1297
 3.189.3 Rubi [A] (verified) 1298
 3.189.4 Maple [A] (verified) 1300
 3.189.5 Fricas [A] (verification not implemented) 1301
 3.189.6 Sympy [F] 1301
 3.189.7 Maxima [F(-2)] 1302
 3.189.8 Giac [B] (verification not implemented) 1302
 3.189.9 Mupad [B] (verification not implemented) 1303

3.189.1 Optimal result

Integrand size = 29, antiderivative size = 69

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c+d \sec(e+fx)} dx = \frac{a \operatorname{arctanh}(\sin(e+fx))}{df} - \frac{2a\sqrt{c-d} \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{d\sqrt{c+d}}$$

output `a*arctanh(sin(f*x+e))/d/f-2*a*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))*(c-d)^(1/2)/d/f/(c+d)^(1/2)`

3.189.2 Mathematica [A] (verified)

Time = 0.76 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.55

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c+d \sec(e+fx)} dx = \frac{a \left(\frac{2(c-d) \operatorname{arctanh}\left(\frac{(-c+d) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}} - \log\left(\cos\left(\frac{1}{2}(e+fx)\right) - \sin\left(\frac{1}{2}(e+fx)\right)\right) + \log\left(\cos\left(\frac{1}{2}(e+fx)\right) + \sin\left(\frac{1}{2}(e+fx)\right)\right) \right)}{df}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x]),x]`

3.189. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{c+d \sec(e+fx)} dx$

output $(a*((2*(c - d)*\text{ArcTanh}[((-c + d)*\text{Tan}[(e + f*x)/2])/\text{Sqrt}[c^2 - d^2]])/\text{Sqrt}[c^2 - d^2] - \text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] + \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]])/(d*f)$

3.189.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4486, 3042, 4257, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e + fx)(a \sec(e + fx) + a)}{c + d \sec(e + fx)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\csc(e + fx + \frac{\pi}{2})(a \csc(e + fx + \frac{\pi}{2}) + a)}{c + d \csc(e + fx + \frac{\pi}{2})} dx \\ & \quad \downarrow 4486 \\ & \frac{a \int \sec(e + fx) dx}{d} - \frac{a(c - d) \int \frac{\sec(e + fx)}{c + d \sec(e + fx)} dx}{d} \\ & \quad \downarrow 3042 \\ & \frac{a \int \csc(e + fx + \frac{\pi}{2}) dx}{d} - \frac{a(c - d) \int \frac{\csc(e + fx + \frac{\pi}{2})}{c + d \csc(e + fx + \frac{\pi}{2})} dx}{d} \\ & \quad \downarrow 4257 \\ & \frac{a \arctanh(\sin(e + fx))}{df} - \frac{a(c - d) \int \frac{\csc(e + fx + \frac{\pi}{2})}{c + d \csc(e + fx + \frac{\pi}{2})} dx}{d} \\ & \quad \downarrow 4318 \\ & \frac{a \arctanh(\sin(e + fx))}{df} - \frac{a(c - d) \int \frac{1}{\frac{c \cos(e + fx)}{d} + 1} dx}{d^2} \\ & \quad \downarrow 3042 \\ & \frac{a \arctanh(\sin(e + fx))}{df} - \frac{a(c - d) \int \frac{1}{\frac{c \sin(e + fx + \frac{\pi}{2})}{d} + 1} dx}{d^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 3138 \\ & \frac{\operatorname{arctanh}(\sin(e+fx))}{df} - \frac{2a(c-d) \int \frac{1}{(1-\frac{c}{d}) \tan^2(\frac{1}{2}(e+fx)) + \frac{c+d}{d}} d \tan(\frac{1}{2}(e+fx))}{d^2 f} \\ & \downarrow 221 \\ & \frac{\operatorname{arctanh}(\sin(e+fx))}{df} - \frac{2a\sqrt{c-d} \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{df\sqrt{c+d}} \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x]),x]`

output `(a*ArcTanh[Sin[e + f*x]])/(d*f) - (2*a*Sqrt[c - d]*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]])/(d*Sqrt[c + d]*f)`

3.189.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4318 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

```
rule 4486 Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[B/b Int[Csc[e + f*x], x], x] + Simp[(A*b - a*B)/b Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0]
```

3.189.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.30

method	result
derivativedivides	$4a \left(-\frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4d} - \frac{(c-d) \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2d\sqrt{(c+d)(c-d)}} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4d} \right)$
default	$4a \left(-\frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{4d} - \frac{(c-d) \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2d\sqrt{(c+d)(c-d)}} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{4d} \right)$
risch	$\frac{\sqrt{(c+d)(c-d)} a \ln\left(\frac{e^{i(fx+e)} - i\sqrt{(c+d)(c-d)-d}}{c}\right)}{(c+d)fd} - \frac{\sqrt{(c+d)(c-d)} a \ln\left(\frac{e^{i(fx+e)} + i\sqrt{(c+d)(c-d)+d}}{c}\right)}{(c+d)fd} + \frac{a \ln(e^{i(fx+e)})}{df}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 4/f*a*(-1/4/d*ln(tan(1/2*f*x+1/2*e)-1)-1/2*(c-d)/d/((c+d)*(c-d))^(1/2)*arc
tanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))+1/4/d*ln(tan(1/2*f*x+1/
2*e)+1))
```

3.189.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 255, normalized size of antiderivative = 3.70

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{c+d\sec(e+fx)} dx$$

$$= \frac{a\sqrt{\frac{c-d}{c+d}} \log\left(\frac{2cd\cos(fx+e)-(c^2-2d^2)\cos(fx+e)-2(c^2+cd+(cd+d^2)\cos(fx+e))\sqrt{\frac{c-d}{c+d}}\sin(fx+e)+2c^2-d^2}{c^2\cos(fx+e)^2+2cd\cos(fx+e)+d^2}\right) + a\log(\sin(fx+e))}{2df}$$

$$- \frac{2a\sqrt{-\frac{c-d}{c+d}} \arctan\left(-\frac{(d\cos(fx+e)+c)\sqrt{-\frac{c-d}{c+d}}}{(c-d)\sin(fx+e)}\right) - a\log(\sin(fx+e)+1) + a\log(-\sin(fx+e)+1)}{2df}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="fricas")
```

```
output [1/2*(a*sqrt((c - d)/(c + d))*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*(c^2 + c*d + (c*d + d^2)*cos(f*x + e))*sqrt((c - d)/(c + d))*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + a*log(sin(f*x + e) + 1) - a*log(-sin(f*x + e) + 1))/(d*f), -1/2*(2*a*sqrt(-(c - d)/(c + d))*arctan(-(d*cos(f*x + e) + c)*sqrt(-(c - d)/(c + d)))/((c - d)*sin(f*x + e))) - a*log(sin(f*x + e) + 1) + a*log(-sin(f*x + e) + 1))/(d*f)]
```

3.189.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{c+d\sec(e+fx)} dx$$

$$= a\left(\int \frac{\sec(e+fx)}{c+d\sec(e+fx)} dx + \int \frac{\sec^2(e+fx)}{c+d\sec(e+fx)} dx\right)$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x)
```

```
output a*(Integral(sec(e + f*x)/(c + d*sec(e + f*x)), x) + Integral(sec(e + f*x)*2/(c + d*sec(e + f*x)), x))
```

3.189. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{c+d\sec(e+fx)} dx$

3.189.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{c + d \sec(e + fx)} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f
or more de
```

3.189.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 127 vs. 2(60) = 120.

Time = 0.33 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.84

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{c + d \sec(e + fx)} dx$$

$$= \frac{\frac{a \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) + 1|)}{d} - \frac{a \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) - 1|)}{d} + \frac{2 \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e) - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-c^2+d^2}} \right) \right)}{\sqrt{-c^2+d^2}d}}{f} (ac-$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="giac")
```

```
output (a*log(abs(tan(1/2*f*x + 1/2*e) + 1))/d - a*log(abs(tan(1/2*f*x + 1/2*e) -
1))/d + 2*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan
(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*(a*c - a*d
))/(sqrt(-c^2 + d^2)*d)/f
```

3.189.9 Mupad [B] (verification not implemented)

Time = 13.83 (sec) , antiderivative size = 195, normalized size of antiderivative = 2.83

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{c+d\sec(e+fx)} dx = \frac{2a \operatorname{atanh}\left(\frac{\sin\left(\frac{e}{2}+\frac{fx}{2}\right)}{\cos\left(\frac{e}{2}+\frac{fx}{2}\right)}\right)}{f(c+d)} + \frac{2a \left(\operatorname{atanh}\left(\frac{d^3 \sin\left(\frac{e}{2}+\frac{fx}{2}\right) - c^3 \sin\left(\frac{e}{2}+\frac{fx}{2}\right) + cd^2 \sin\left(\frac{e}{2}+\frac{fx}{2}\right) - c^2 d \sin\left(\frac{e}{2}+\frac{fx}{2}\right) + c \sin\left(\frac{e}{2}+\frac{fx}{2}\right) (c^2-d^2)}{\cos\left(\frac{e}{2}+\frac{fx}{2}\right) \sqrt{c^2-d^2} (d^2+cd)}\right) \sqrt{c^2-d^2} + c \operatorname{atanh}\left(\frac{\sin\left(\frac{e}{2}+\frac{fx}{2}\right)}{\cos\left(\frac{e}{2}+\frac{fx}{2}\right)}\right)}{df(c+d)}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c + d/cos(e + f*x))),x)`output `(2*a*atanh(sin(e/2 + (f*x)/2)/cos(e/2 + (f*x)/2)))/(f*(c + d)) + (2*a*(atanh((d^3*sin(e/2 + (f*x)/2) - c^3*sin(e/2 + (f*x)/2) + c*d^2*sin(e/2 + (f*x)/2) - c^2*d*sin(e/2 + (f*x)/2) + c*sin(e/2 + (f*x)/2)*(c^2 - d^2))/(cos(e/2 + (f*x)/2)*(c^2 - d^2)^(1/2)*(c*d + d^2)))*(c^2 - d^2)^(1/2) + c*atanh(sin(e/2 + (f*x)/2)/cos(e/2 + (f*x)/2)))/(d*f*(c + d))`

3.190 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^2} dx$

3.190.1 Optimal result 1304
 3.190.2 Mathematica [A] (verified) 1304
 3.190.3 Rubi [A] (verified) 1305
 3.190.4 Maple [A] (verified) 1307
 3.190.5 Fricas [B] (verification not implemented) 1308
 3.190.6 Sympy [F] 1308
 3.190.7 Maxima [F(-2)] 1309
 3.190.8 Giac [A] (verification not implemented) 1309
 3.190.9 Mupad [B] (verification not implemented) 1310

3.190.1 Optimal result

Integrand size = 29, antiderivative size = 79

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^2} dx = \frac{2a \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{\sqrt{c-d}(c+d)^{3/2}f} + \frac{a \tan(e+fx)}{(c+d)f(c+d \sec(e+fx))}$$

output `2*a*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c+d)^(3/2)/f/(c-d)^(1/2)+a*tan(f*x+e)/(c+d)/f/(c+d*sec(f*x+e))`

3.190.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^2} dx = \frac{a \left(-\frac{2 \operatorname{arctanh}\left(\frac{(-c+d) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}} + \frac{\sin(e+fx)}{d+c \cos(e+fx)} \right)}{(c+d)f}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x])^2,x]`

output `(a*((-2*ArcTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]])/Sqrt[c^2 - d^2] + Sin[e + f*x]/(d + c*Cos[e + f*x]))/((c + d)*f)`

3.190. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^2} dx$

3.190.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.14, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 4491, 25, 27, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c+d\sec(e+fx))^2} dx$$

$$\downarrow 3042$$

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)}{(c+d\csc(e+fx+\frac{\pi}{2}))^2} dx$$

$$\downarrow 4491$$

$$\frac{a \tan(e+fx)}{f(c+d)(c+d\sec(e+fx))} - \frac{\int -\frac{a(c-d)\sec(e+fx)}{c+d\sec(e+fx)} dx}{c^2-d^2}$$

$$\downarrow 25$$

$$\frac{\int \frac{a(c-d)\sec(e+fx)}{c+d\sec(e+fx)} dx}{c^2-d^2} + \frac{a \tan(e+fx)}{f(c+d)(c+d\sec(e+fx))}$$

$$\downarrow 27$$

$$\frac{a(c-d) \int \frac{\sec(e+fx)}{c+d\sec(e+fx)} dx}{c^2-d^2} + \frac{a \tan(e+fx)}{f(c+d)(c+d\sec(e+fx))}$$

$$\downarrow 3042$$

$$\frac{a(c-d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{c+d\csc(e+fx+\frac{\pi}{2})} dx}{c^2-d^2} + \frac{a \tan(e+fx)}{f(c+d)(c+d\sec(e+fx))}$$

$$\downarrow 4318$$

$$\frac{a(c-d) \int \frac{1}{\frac{c \cos(e+fx)}{d} + 1} dx}{d(c^2-d^2)} + \frac{a \tan(e+fx)}{f(c+d)(c+d\sec(e+fx))}$$

$$\downarrow 3042$$

$$\frac{a(c-d) \int \frac{1}{\frac{c \sin(e+fx+\frac{\pi}{2})}{d} + 1} dx}{d(c^2-d^2)} + \frac{a \tan(e+fx)}{f(c+d)(c+d\sec(e+fx))}$$

$$\downarrow 3138$$

$$\frac{2a(c-d) \int \frac{1}{(1-\frac{c}{d}) \tan^2(\frac{1}{2}(e+fx)) + \frac{c+d}{d}} d \tan(\frac{1}{2}(e+fx))}{df(c^2-d^2)} + \frac{a \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))}$$

↓ 221

$$\frac{2a\sqrt{c-d} \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{f\sqrt{c+d}(c^2-d^2)} + \frac{a \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x])^2,x]`

output `(2*a*sqrt[c - d]*ArcTanh[(sqrt[c - d]*Tan[(e + f*x)/2]]/sqrt[c + d])/(sqrt[c + d]*(c^2 - d^2)*f) + (a*Tan[e + f*x])/((c + d)*f*(c + d*Sec[e + f*x]))`

3.190.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4318 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

```
rule 4491 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(cs
c[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e
+ f*x]*((a + b*Csc[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1
/((m + 1)*(a^2 - b^2)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp
[(a*A - b*B)*(m + 1) - (A*b - a*B)*(m + 2)*Csc[e + f*x], x], x] /; Free
Q[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m
, -1]
```

3.190.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.33

method	result
derivativedivides	$4a \frac{\left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}\right) + \frac{\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c+d)\sqrt{(c+d)(c-d)}} \right)}{f}$
default	$4a \frac{\left(-\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}\right) + \frac{\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c+d)\sqrt{(c+d)(c-d)}} \right)}{f}$
risch	$\frac{2ia(d e^{i(fx+e)} + c)}{cf(c+d)(e^{2i(fx+e)}c + 2d e^{i(fx+e)} + c)} + \frac{a \ln\left(e^{i(fx+e)} + \frac{ic^2 - id^2 + \sqrt{c^2 - d^2}d}{\sqrt{c^2 - d^2}c}\right)}{\sqrt{c^2 - d^2}(c+d)f} - \frac{a \ln\left(e^{i(fx+e)} + \frac{-ic^2 + id^2 + \sqrt{c^2 - d^2}d}{c\sqrt{c^2 - d^2}}\right)}{\sqrt{c^2 - d^2}(c+d)f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE
)
```

```
output 4/f*a*(-1/2*tan(1/2*f*x+1/2*e)/(c+d)/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1
/2*e)^2*d-c-d)+1/2/(c+d)/(((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2
*e)/((c+d)*(c-d))^(1/2)))
```

3.190.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 149 vs. 2(70) = 140.

Time = 0.29 (sec) , antiderivative size = 357, normalized size of antiderivative = 4.52

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^2} dx$$

$$= \left[\frac{(ac \cos(fx+e) + ad)\sqrt{c^2-d^2} \log\left(\frac{2cd\cos(fx+e) - (c^2-2d^2)\cos(fx+e)^2 + 2\sqrt{c^2-d^2}(d\cos(fx+e)+c)\sin(fx+e) + 2c^2-d^2}{c^2\cos(fx+e)^2 + 2cd\cos(fx+e) + d^2}\right) - 2((c^4 + c^3d - c^2d^2 - cd^3)f \cos(fx+e) + (c^3d + c^2d^2 - cd^3 - d^4)f)}{2((c^4 + c^3d - c^2d^2 - cd^3)f \cos(fx+e) + (c^3d + c^2d^2 - cd^3 - d^4)f)} \right]$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="fricas")`

output `[1/2*((a*c*cos(f*x + e) + a*d)*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(a*c^2 - a*d^2)*sin(f*x + e))/((c^4 + c^3*d - c^2*d^2 - c*d^3)*f*cos(f*x + e) + (c^3*d + c^2*d^2 - c*d^3 - d^4)*f), ((a*c*cos(f*x + e) + a*d)*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (a*c^2 - a*d^2)*sin(f*x + e))/((c^4 + c^3*d - c^2*d^2 - c*d^3)*f*cos(f*x + e) + (c^3*d + c^2*d^2 - c*d^3 - d^4)*f)]`

3.190.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^2} dx = a \left(\int \frac{\sec(e+fx)}{c^2 + 2cd\sec(e+fx) + d^2\sec^2(e+fx)} dx + \int \frac{\sec^2(e+fx)}{c^2 + 2cd\sec(e+fx) + d^2\sec^2(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))**2,x)`

output `a*(Integral(sec(e + f*x)/(c**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)**2), x) + Integral(sec(e + f*x)**2/(c**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)**2), x))`

3.190.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c + d \sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de
```

3.190.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.73

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c + d \sec(e + fx))^2} dx =$$

$$2 \left(\frac{\left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan\left(\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e) - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-c^2+d^2}}\right) \right) a}{\sqrt{-c^2+d^2}(c+d)} + \frac{a \tan(\frac{1}{2} fx + \frac{1}{2} e)}{(c \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - d \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - c - d)(c+d)} \right) / f$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="giac")
```

```
output -2*((pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*a/(sqrt(-c^2 + d^2)*(c + d)) + a*tan(1/2*f*x + 1/2*e)/((c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)*(c + d)))/f
```

3.190.9 Mupad [B] (verification not implemented)

Time = 13.47 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.08

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^2} dx = \frac{2a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f(c+d) \left((d-c) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + c+d \right)} + \frac{2a \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \sqrt{c-d}}{\sqrt{c+d}}\right)}{f(c+d)^{3/2} \sqrt{c-d}}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c + d/cos(e + f*x))^2),x)`output `(2*a*tan(e/2 + (f*x)/2))/(f*(c + d)*(c + d - tan(e/2 + (f*x)/2)^2*(c - d)) + (2*a*atanh((tan(e/2 + (f*x)/2)*(c - d)^(1/2))/(c + d)^(1/2)))/(f*(c + d)^(3/2)*(c - d)^(1/2))`

3.191 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^3} dx$

3.191.1 Optimal result 1311
 3.191.2 Mathematica [A] (verified) 1311
 3.191.3 Rubi [A] (verified) 1312
 3.191.4 Maple [A] (verified) 1315
 3.191.5 Fricas [B] (verification not implemented) 1315
 3.191.6 Sympy [F] 1316
 3.191.7 Maxima [F(-2)] 1317
 3.191.8 Giac [B] (verification not implemented) 1317
 3.191.9 Mupad [B] (verification not implemented) 1318

3.191.1 Optimal result

Integrand size = 29, antiderivative size = 131

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^3} dx = \frac{a(2c-d)\operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{(c-d)^{3/2}(c+d)^{5/2}f} + \frac{a \tan(e+fx)}{2(c+d)f(c+d \sec(e+fx))^2} + \frac{a(c-2d) \tan(e+fx)}{2(c-d)(c+d)^2f(c+d \sec(e+fx))}$$

```
output a*(2*c-d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c-d)^(3/2)/(c+d)^(5/2)/f+1/2*a*tan(f*x+e)/(c+d)/f/(c+d*sec(f*x+e))^2+1/2*a*(c-2*d)*tan(f*x+e)/(c-d)/(c+d)^2/f/(c+d*sec(f*x+e))
```

3.191.2 Mathematica [A] (verified)

Time = 1.54 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.27

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^3} dx = \frac{a(1+\cos(e+fx)) \sec^2\left(\frac{1}{2}(e+fx)\right) \left(-2(2c-d)\operatorname{arctanh}\left(\frac{(-c+d)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right) (d+c \cos(e+fx))^2 + \sqrt{c^2-d^2}\right)}{4(c-d)(c+d)^2\sqrt{c^2-d^2}f(d+c \cos(e+fx))}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x])^3,x]`

output `(a*(1 + Cos[e + f*x])*Sec[(e + f*x)/2]^2*(-2*(2*c - d)*ArcTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]]*(d + c*Cos[e + f*x])^2 + Sqrt[c^2 - d^2]*((c - 2*d)*d + (2*c^2 - 2*c*d - d^2)*Cos[e + f*x])*Sin[e + f*x))/(4*(c - d)*(c + d)^2*Sqrt[c^2 - d^2]*f*(d + c*Cos[e + f*x])^2)`

3.191.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.14, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 4491, 25, 3042, 4491, 25, 27, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a+\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a+\csc(e+fx+\frac{\pi}{2}))}{(c+d\csc(e+fx+\frac{\pi}{2}))^3} dx$$

↓ 4491

$$\frac{a \tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2} - \frac{\int -\frac{\sec(e+fx)(2a(c-d)+a\sec(e+fx)(c-d))}{(c+d\sec(e+fx))^2} dx}{2(c^2-d^2)}$$

↓ 25

$$\frac{\int \frac{\sec(e+fx)(2a(c-d)+a\sec(e+fx)(c-d))}{(c+d\sec(e+fx))^2} dx}{2(c^2-d^2)} + \frac{a \tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2}$$

↓ 3042

$$\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(2a(c-d)+a\csc(e+fx+\frac{\pi}{2})(c-d))}{(c+d\csc(e+fx+\frac{\pi}{2}))^2} dx}{2(c^2-d^2)} + \frac{a \tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2}$$

↓ 4491

$$\frac{a(c-2d) \tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))} - \frac{\int -\frac{a(c-d)(2c-d)\sec(e+fx)}{c+d\sec(e+fx)} dx}{c^2-d^2} + \frac{a \tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2}$$

3.191. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{\int \frac{a(c-d)(2c-d) \sec(e+fx)}{c+d \sec(e+fx)} dx}{c^2-d^2} + \frac{a(c-2d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))} + \frac{a \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} \\
& \downarrow 27 \\
& \frac{a(c-d)(2c-d) \int \frac{\sec(e+fx)}{c+d \sec(e+fx)} dx}{c^2-d^2} + \frac{a(c-2d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))} + \frac{a \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} \\
& \downarrow 3042 \\
& \frac{a(c-d)(2c-d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{c+d \csc(e+fx+\frac{\pi}{2})} dx}{c^2-d^2} + \frac{a(c-2d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))} + \frac{a \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} \\
& \downarrow 4318 \\
& \frac{a(c-d)(2c-d) \int \frac{\frac{1}{c \cos(e+fx)+1}}{d} dx}{d(c^2-d^2)} + \frac{a(c-2d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))} + \frac{a \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} \\
& \downarrow 3042 \\
& \frac{a(c-d)(2c-d) \int \frac{\frac{1}{c \sin(e+fx+\frac{\pi}{2})+1}}{d} dx}{d(c^2-d^2)} + \frac{a(c-2d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))} + \frac{a \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} \\
& \downarrow 3138 \\
& \frac{2a(c-d)(2c-d) \int \frac{\frac{1}{(1-\frac{c}{d}) \tan^2(\frac{1}{2}(e+fx)) + \frac{c+d}{d}} d \tan(\frac{1}{2}(e+fx))}{df(c^2-d^2)}}{2(c^2-d^2)} + \frac{a(c-2d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))} + \\
& \frac{a \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} \\
& \downarrow 221 \\
& \frac{2a\sqrt{c-d}(2c-d) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{f\sqrt{c+d}(c^2-d^2)} + \frac{a(c-2d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))} + \frac{a \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x])^3,x]`

```
output (a*Tan[e + f*x])/(2*(c + d)*f*(c + d*Sec[e + f*x])^2) + ((2*a*Sqrt[c - d]*
(2*c - d)*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]])/(Sqrt[c + d
]*(c^2 - d^2)*f) + (a*(c - 2*d)*Tan[e + f*x])/((c + d)*f*(c + d*Sec[e + f*
x]))) / (2*(c^2 - d^2))
```

3.191.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 221 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3138 Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

```
rule 4318 Int[csc[(e_) + (f_)*(x_)]/(csc[(e_) + (f_)*(x_)]*(b_) + (a_)), x_Symbo
l] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e,
f}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 4491 Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(cs
c[(e_) + (f_)*(x_)]*(B_) + (A_)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e
+ f*x]*((a + b*Csc[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1
/((m + 1)*(a^2 - b^2)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp
[(a*A - b*B)*(m + 1) - (A*b - a*B)*(m + 2)*Csc[e + f*x], x], x], x] /; Free
Q[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m
, -1]
```

3.191.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.36

method	result
derivativedivides	$4a \frac{\left(-\frac{(2c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{4(c^2+2cd+d^2)} + \frac{(2c-3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{4(c+d)(c-d)} + \frac{(2c-d)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{4(c^3+c^2d-cd^2-d^3)\sqrt{(c+d)(c-d)}} \right)}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 d - c - d \right)^2} + \frac{(2c-d)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{4(c^3+c^2d-cd^2-d^3)\sqrt{(c+d)(c-d)}}$
default	$4a \frac{\left(-\frac{(2c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{4(c^2+2cd+d^2)} + \frac{(2c-3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{4(c+d)(c-d)} + \frac{(2c-d)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{4(c^3+c^2d-cd^2-d^3)\sqrt{(c+d)(c-d)}} \right)}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 d - c - d \right)^2} + \frac{(2c-d)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{4(c^3+c^2d-cd^2-d^3)\sqrt{(c+d)(c-d)}}$
risch	$\frac{ia(-3c^3de^{3i(fx+e)}+2c^2d^2e^{3i(fx+e)}+2cd^3e^{3i(fx+e)}-2c^4e^{2i(fx+e)}+2c^3de^{2i(fx+e)}-3c^2d^2e^{2i(fx+e)}+4cd^3e^{2i(fx+e)}+2c^2d^2e^{i(fx+e)}+2cd^3e^{i(fx+e)}+2c^2de^{i(fx+e)}+2cde^{i(fx+e)}+2c^2e^{i(fx+e)}+2cde^{i(fx+e)}+2ce^{i(fx+e)}+2e^{i(fx+e)})}{c^2(-c^2+d^2)f(e^{2i(fx+e)}c+2de^{i(fx+e)}+c)^2(c^2+2cd+d^2)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 4/f*a*((-1/4*(2*c-d)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3+1/4*(2*c-3*d)/(c+d)/(c-d)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^2+1/4*(2*c-d)/(c^3+c^2*d-c*d^2-d^3)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))
```

3.191.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. 2(118) = 236.

Time = 0.31 (sec) , antiderivative size = 736, normalized size of antiderivative = 5.62

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$$

$$= \left[\frac{(2acd^2 - ad^3 + (2ac^3 - ac^2d)\cos(fx+e)^2 + 2(2ac^2d - acd^2)\cos(fx+e))\sqrt{c^2 - d^2} \log\left(\frac{2cd\cos(fx+e) - (c^2 - d^2)\sec(fx+e)}{2((c^7 + c^6d - 2c^5d^2 - 2c^4d^3 + c^3d^4 + c^2d^5)f\cos(fx+e) + c^2d^2 + cd^3 + c^2d + d^3)}\right)}{4((c^7 + c^6d - 2c^5d^2 - 2c^4d^3 + c^3d^4 + c^2d^5)f\cos(fx+e) + c^2d^2 + cd^3 + c^2d + d^3)} \right]$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="fracas")
```

output `[1/4*((2*a*c*d^2 - a*d^3 + (2*a*c^3 - a*c^2*d)*cos(f*x + e)^2 + 2*(2*a*c^2*d - a*c*d^2)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c))*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(a*c^3*d - 2*a*c^2*d^2 - a*c*d^3 + 2*a*d^4 + (2*a*c^4 - 2*a*c^3*d - 3*a*c^2*d^2 + 2*a*c*d^3 + a*d^4)*cos(f*x + e))*sin(f*x + e)/((c^7 + c^6*d - 2*c^5*d^2 - 2*c^4*d^3 + c^3*d^4 + c^2*d^5)*f*cos(f*x + e)^2 + 2*(c^6*d + c^5*d^2 - 2*c^4*d^3 - 2*c^3*d^4 + c^2*d^5 + c*d^6)*f*cos(f*x + e) + (c^5*d^2 + c^4*d^3 - 2*c^3*d^4 - 2*c^2*d^5 + c*d^6 + d^7)*f), 1/2*((2*a*c*d^2 - a*d^3 + (2*a*c^3 - a*c^2*d)*cos(f*x + e)^2 + 2*(2*a*c^2*d - a*c*d^2)*cos(f*x + e))*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (a*c^3*d - 2*a*c^2*d^2 - a*c*d^3 + 2*a*d^4 + (2*a*c^4 - 2*a*c^3*d - 3*a*c^2*d^2 + 2*a*c*d^3 + a*d^4)*cos(f*x + e))*sin(f*x + e)/((c^7 + c^6*d - 2*c^5*d^2 - 2*c^4*d^3 + c^3*d^4 + c^2*d^5)*f*cos(f*x + e)^2 + 2*(c^6*d + c^5*d^2 - 2*c^4*d^3 - 2*c^3*d^4 + c^2*d^5 + c*d^6)*f*cos(f*x + e) + (c^5*d^2 + c^4*d^3 - 2*c^3*d^4 - 2*c^2*d^5 + c*d^6 + d^7)*f)]`

3.191.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$$

$$= a \left(\int \frac{\sec(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx + \int \frac{\sec^2(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))**3,x)`

output `a*(Integral(sec(e + f*x)/(c**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**3), x) + Integral(sec(e + f*x)**2/(c**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**3), x))`

3.191.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c + d \sec(e + fx))^3} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de
```

3.191.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 263 vs. 2(118) = 236.

Time = 0.36 (sec) , antiderivative size = 263, normalized size of antiderivative = 2.01

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c + d \sec(e + fx))^3} dx$$

$$= \frac{\left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2c+2d) + \arctan\left(-\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right) (2ac-ad)}{(c^3+c^2d-cd^2-d^3)\sqrt{-c^2+d^2}} - \frac{2ac^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 3acd \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + ad^2}{(c^3+c^2d-cd^2-d^3)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="giac")
```

```
output ((pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*(2*a*c - a*d)/((c^3 + c^2*d - c*d^2 - d^3)*sqrt(-c^2 + d^2)) - (2*a*c^2*tan(1/2*f*x + 1/2*e)^3 - 3*a*c*d*tan(1/2*f*x + 1/2*e)^3 + a*d^2*tan(1/2*f*x + 1/2*e)^3 - 2*a*c^2*tan(1/2*f*x + 1/2*e) + a*c*d*tan(1/2*f*x + 1/2*e) + 3*a*d^2*tan(1/2*f*x + 1/2*e))/((c^3 + c^2*d - c*d^2 - d^3)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^2))/f
```

3.191. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$

3.191.9 Mupad [B] (verification not implemented)

Time = 15.42 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.31

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$$

$$= \frac{a \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2}+\frac{fx}{2}\right)\sqrt{c-d}}{\sqrt{c+d}}\right)(2c-d)}{f(c+d)^{5/2}(c-d)^{3/2}}$$

$$- \frac{\frac{\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^3(2ac-ad)}{(c+d)^2} - \frac{a \tan\left(\frac{e}{2}+\frac{fx}{2}\right)(2c-3d)}{(c+d)(c-d)}}{f\left(2cd - \tan\left(\frac{e}{2}+\frac{fx}{2}\right)^2(2c^2-2d^2) + \tan\left(\frac{e}{2}+\frac{fx}{2}\right)^4(c^2-2cd+d^2) + c^2+d^2\right)}$$

input `int((a + a/cos(e + f*x))/(cos(e + f*x)*(c + d/cos(e + f*x))^3),x)`output `(a*atanh((tan(e/2 + (f*x)/2)*(c - d)^(1/2))/(c + d)^(1/2))*(2*c - d))/(f*(c + d)^(5/2)*(c - d)^(3/2)) - ((tan(e/2 + (f*x)/2)^3*(2*a*c - a*d))/(c + d)^2 - (a*tan(e/2 + (f*x)/2)*(2*c - 3*d))/((c + d)*(c - d)))/(f*(2*c*d - tan(e/2 + (f*x)/2)^2*(2*c^2 - 2*d^2) + tan(e/2 + (f*x)/2)^4*(c^2 - 2*c*d + d^2) + c^2 + d^2))`

3.192 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^4} dx$

3.192.1 Optimal result 1319
 3.192.2 Mathematica [A] (verified) 1320
 3.192.3 Rubi [A] (verified) 1320
 3.192.4 Maple [A] (verified) 1324
 3.192.5 Fricas [B] (verification not implemented) 1324
 3.192.6 Sympy [F] 1325
 3.192.7 Maxima [F(-2)] 1326
 3.192.8 Giac [B] (verification not implemented) 1326
 3.192.9 Mupad [B] (verification not implemented) 1327

3.192.1 Optimal result

Integrand size = 29, antiderivative size = 189

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))}{(c+d \sec(e+fx))^4} dx = \frac{a(2c^2 - 2cd + d^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{(c-d)^{5/2}(c+d)^{7/2}f} + \frac{a \tan(e+fx)}{3(c+d)f(c+d \sec(e+fx))^3} + \frac{a(2c-3d) \tan(e+fx)}{6(c-d)(c+d)^2f(c+d \sec(e+fx))^2} + \frac{a(c-4d)(2c-d) \tan(e+fx)}{6(c-d)^2(c+d)^3f(c+d \sec(e+fx))}$$

```
output a*(2*c^2-2*c*d+d^2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c-d)^(5/2)/(c+d)^(7/2)/f+1/3*a*tan(f*x+e)/(c+d)/f/(c+d*sec(f*x+e))^3+1/6*a*(2*c-3*d)*tan(f*x+e)/(c-d)/(c+d)^2/f/(c+d*sec(f*x+e))^2+1/6*a*(c-4*d)*(2*c-d)*tan(f*x+e)/(c-d)^2/(c+d)^3/f/(c+d*sec(f*x+e))
```


3.192.2 Mathematica [A] (verified)

Time = 3.65 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.31

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^4} dx =$$

$$\frac{a(1+\cos(e+fx))\sec^2\left(\frac{1}{2}(e+fx)\right)\left(6(2c^2-2cd+d^2)\operatorname{arctanh}\left(\frac{(-c+d)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)\right)(d+c\cos(e+fx))}{\dots}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x])^4,x]`output `-1/12*(a*(1 + Cos[e + f*x])*Sec[(e + f*x)/2]^2*(6*(2*c^2 - 2*c*d + d^2)*Ar
cTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]]*(d + c*Cos[e + f*x])^3
- (Sqrt[c^2 - d^2]*(6*c^4 - 12*c^3*d + 2*c^2*d^2 - 15*c*d^3 + 10*d^4 + 6*d
*(2*c^3 - 7*c^2*d + 2*c*d^2 + d^3)*Cos[e + f*x] + (6*c^4 - 12*c^3*d - 2*c^
2*d^2 + 3*c*d^3 + 2*d^4)*Cos[2*(e + f*x)]*Sin[e + f*x])/2))/((c - d)^2*(c
+ d)^3*Sqrt[c^2 - d^2]*f*(d + c*Cos[e + f*x])^3)`**3.192.3 Rubi [A] (verified)**Time = 1.07 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.14, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$, Rules used = {3042, 4491, 25, 3042, 4491, 25, 3042, 4491, 27, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)}{(c+d\sec(e+fx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(a\csc\left(e+fx+\frac{\pi}{2}\right)+a)}{(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^4} dx$$

$$\downarrow \text{4491}$$

$$\frac{a\tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3} - \frac{\int -\frac{\sec(e+fx)(3a(c-d)+2a\sec(e+fx)(c-d))}{(c+d\sec(e+fx))^3} dx}{3(c^2-d^2)}$$

$$\downarrow \text{25}$$

3.192. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$

$$\begin{aligned}
& \frac{\int \frac{\sec(e+fx)(3a(c-d)+2a\sec(e+fx)(c-d))}{(c+d\sec(e+fx))^3} dx}{3(c^2-d^2)} + \frac{a \tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(3a(c-d)+2a\csc(e+fx+\frac{\pi}{2})(c-d))}{(c+d\csc(e+fx+\frac{\pi}{2}))^3} dx}{3(c^2-d^2)} + \frac{a \tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3} \\
& \quad \downarrow \text{4491} \\
& \frac{\frac{a(2c-3d)\tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2} - \frac{\int -\frac{\sec(e+fx)(2a(3c-2d)(c-d)+a(2c-3d)\sec(e+fx)(c-d))}{(c+d\sec(e+fx))^2} dx}{2(c^2-d^2)}}{3(c^2-d^2)} + \frac{a \tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3} \\
& \quad \downarrow \text{25} \\
& \frac{\frac{\int \frac{\sec(e+fx)(2a(3c-2d)(c-d)+a(2c-3d)\sec(e+fx)(c-d))}{(c+d\sec(e+fx))^2} dx}{2(c^2-d^2)} + \frac{a(2c-3d)\tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2} + \frac{a \tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3}}{3(c^2-d^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(2a(3c-2d)(c-d)+a(2c-3d)\csc(e+fx+\frac{\pi}{2})(c-d))}{(c+d\csc(e+fx+\frac{\pi}{2}))^2} dx}{2(c^2-d^2)} + \frac{a(2c-3d)\tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2} + \frac{a \tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3}}{3(c^2-d^2)} \\
& \quad \downarrow \text{4491} \\
& \frac{\frac{\frac{a(c-4d)(2c-d)\tan(e+fx)}{f(c+d)(c+d\sec(e+fx))} - \frac{\int -\frac{3a(c-d)(2c^2-2dc+d^2)\sec(e+fx)}{c+d\sec(e+fx)} dx}{c^2-d^2}}{2(c^2-d^2)} + \frac{a(2c-3d)\tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2} + \frac{a \tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3}}{3(c^2-d^2)} \\
& \quad \downarrow \text{27} \\
& \frac{\frac{3a(c-d)(2c^2-2cd+d^2) \int \frac{\sec(e+fx)}{c+d\sec(e+fx)} dx}{c^2-d^2} + \frac{a(c-4d)(2c-d)\tan(e+fx)}{f(c+d)(c+d\sec(e+fx))} + \frac{a(2c-3d)\tan(e+fx)}{2f(c+d)(c+d\sec(e+fx))^2}}{2(c^2-d^2)} + \frac{a \tan(e+fx)}{3f(c+d)(c+d\sec(e+fx))^3} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.192. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$

$$\frac{3a(c-d)(2c^2-2cd+d^2) \int \frac{\csc(e+fx+\frac{\pi}{2})}{c+d \csc(e+fx+\frac{\pi}{2})} dx + \frac{a(c-4d)(2c-d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))}}{c^2-d^2} + \frac{a(2c-3d) \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)}{a \tan(e+fx)}$$

$$\frac{3f(c+d)(c+d \sec(e+fx))^3}{}$$

↓ 4318

$$\frac{3a(c-d)(2c^2-2cd+d^2) \int \frac{1}{c \cos(e+fx)+1} dx + \frac{a(c-4d)(2c-d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))}}{d(c^2-d^2)} + \frac{a(2c-3d) \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)}{a \tan(e+fx)}$$

$$\frac{3f(c+d)(c+d \sec(e+fx))^3}{}$$

↓ 3042

$$\frac{3a(c-d)(2c^2-2cd+d^2) \int \frac{1}{c \sin(e+fx+\frac{\pi}{2})+1} dx + \frac{a(c-4d)(2c-d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))}}{d(c^2-d^2)} + \frac{a(2c-3d) \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)}{a \tan(e+fx)}$$

$$\frac{3f(c+d)(c+d \sec(e+fx))^3}{}$$

↓ 3138

$$\frac{6a(c-d)(2c^2-2cd+d^2) \int \frac{1}{(1-\frac{c}{d}) \tan^2(\frac{1}{2}(e+fx)) + \frac{c+d}{d}} d \tan(\frac{1}{2}(e+fx)) + \frac{a(c-4d)(2c-d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))}}{df(c^2-d^2)} + \frac{a(2c-3d) \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)}{a \tan(e+fx)}$$

$$\frac{3f(c+d)(c+d \sec(e+fx))^3}{}$$

↓ 221

$$\frac{6a\sqrt{c-d}(2c^2-2cd+d^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right) + \frac{a(c-4d)(2c-d) \tan(e+fx)}{f(c+d)(c+d \sec(e+fx))}}{f\sqrt{c+d}(c^2-d^2)} + \frac{a(2c-3d) \tan(e+fx)}{2f(c+d)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)}{a \tan(e+fx)}$$

$$\frac{3f(c+d)(c+d \sec(e+fx))^3}{}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x]))/(c + d*Sec[e + f*x])^4,x]`

```
output (a*Tan[e + f*x])/(3*(c + d)*f*(c + d*Sec[e + f*x])^3) + ((a*(2*c - 3*d)*Tan[e + f*x])/(2*(c + d)*f*(c + d*Sec[e + f*x])^2) + ((6*a*Sqrt[c - d]*(2*c^2 - 2*c*d + d^2)*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]])/(Sqrt[c + d]*(c^2 - d^2)*f) + (a*(c - 4*d)*(2*c - d)*Tan[e + f*x])/((c + d)*f*(c + d*Sec[e + f*x])))/(2*(c^2 - d^2))/(3*(c^2 - d^2))
```

3.192.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3138 Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 4318 Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 4491 Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[(a*A - b*B)*(m + 1) - (A*b - a*B)*(m + 2)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

3.192.4 Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.43

method	result
derivativedivides	$4a \left(\frac{-\frac{(2c^2-2cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{4(c^3+3c^2d+3cd^2+d^3)} + \frac{(3c^2-6cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3(c-d)(c^2+2cd+d^2)} - \frac{(2c^2-6cd+3d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{4(c+d)(c^2-2cd+d^2)} + \frac{(2c^2-2cd+d^2)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{c-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}\right)}{4(c^5+c^4d-2c^3d^2-2c^2d^3+d^4)} \right) \frac{1}{f}$
default	$4a \left(\frac{-\frac{(2c^2-2cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{4(c^3+3c^2d+3cd^2+d^3)} + \frac{(3c^2-6cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3(c-d)(c^2+2cd+d^2)} - \frac{(2c^2-6cd+3d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{4(c+d)(c^2-2cd+d^2)} + \frac{(2c^2-2cd+d^2)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{c-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}\right)}{4(c^5+c^4d-2c^3d^2-2c^2d^3+d^4)} \right) \frac{1}{f}$
risch	Expression too large to display

input `int(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^4,x,method=_RETURNVERBOSE)`

output `4/f*a*((-1/4*(2*c^2-2*c*d+d^2)/(c^3+3*c^2*d+3*c*d^2+d^3)*tan(1/2*f*x+1/2*e))^5+1/3*(3*c^2-6*c*d+d^2)/(c-d)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3-1/4*(2*c^2-6*c*d+3*d^2)/(c+d)/(c^2-2*c*d+d^2)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^3+1/4*(2*c^2-2*c*d+d^2)/(c^5+c^4*d-2*c^3*d^2-2*c^2*d^3+d^4)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))`

3.192.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 610 vs. 2(174) = 348.

Time = 0.36 (sec) , antiderivative size = 1278, normalized size of antiderivative = 6.76

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^4} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^4,x, algorithm="fricas")`

output

```
[1/12*(3*(2*a*c^2*d^3 - 2*a*c*d^4 + a*d^5 + (2*a*c^5 - 2*a*c^4*d + a*c^3*d^2)*cos(f*x + e)^3 + 3*(2*a*c^4*d - 2*a*c^3*d^2 + a*c^2*d^3)*cos(f*x + e)^2 + 3*(2*a*c^3*d^2 - 2*a*c^2*d^3 + a*c*d^4)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(2*a*c^4*d^2 - 9*a*c^3*d^3 + 2*a*c^2*d^4 + 9*a*c*d^5 - 4*a*d^6 + (6*a*c^6 - 12*a*c^5*d - 8*a*c^4*d^2 + 15*a*c^3*d^3 + 4*a*c^2*d^4 - 3*a*c*d^5 - 2*a*d^6)*cos(f*x + e)^2 + 3*(2*a*c^5*d - 7*a*c^4*d^2 + 8*a*c^3*d^4 - 2*a*c*d^5 - a*d^6)*cos(f*x + e))*sin(f*x + e))/((c^10 + c^9*d - 3*c^8*d^2 - 3*c^7*d^3 + 3*c^6*d^4 + 3*c^5*d^5 - c^4*d^6 - c^3*d^7)*f*cos(f*x + e)^3 + 3*(c^9*d + c^8*d^2 - 3*c^7*d^3 - 3*c^6*d^4 + 3*c^5*d^5 + 3*c^4*d^6 - c^3*d^7 - c^2*d^8)*f*cos(f*x + e)^2 + 3*(c^8*d^2 + c^7*d^3 - 3*c^6*d^4 - 3*c^5*d^5 + 3*c^4*d^6 + 3*c^3*d^7 - c^2*d^8 - c*d^9)*f*cos(f*x + e) + (c^7*d^3 + c^6*d^4 - 3*c^5*d^5 - 3*c^4*d^6 + 3*c^3*d^7 + 3*c^2*d^8 - c*d^9 - d^10)*f), 1/6*(3*(2*a*c^2*d^3 - 2*a*c*d^4 + a*d^5 + (2*a*c^5 - 2*a*c^4*d + a*c^3*d^2)*cos(f*x + e)^3 + 3*(2*a*c^4*d - 2*a*c^3*d^2 + a*c^2*d^3)*cos(f*x + e)^2 + 3*(2*a*c^3*d^2 - 2*a*c^2*d^3 + a*c*d^4)*cos(f*x + e))*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (2*a*c^4*d^2 - 9*a*c^3*d^3 + 2*a*c^2*d^4 + 9*a*c*d^5 - 4*a*d^6 + (6*a*c^6 - 12*a*c^5*d - 8*a*c^4*d^2 + 15*a*c^3*d^3 + 4...
```

3.192.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$$

$$= a \left(\int \frac{\sec(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx \right. \\ \left. + \int \frac{\sec^2(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))**4,x)`

output `a*(Integral(sec(e + f*x)/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x) + Integral(sec(e + f*x)**2/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x))`

3.192.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c + d \sec(e + fx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.192.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 449 vs. 2(174) = 348.

Time = 0.37 (sec) , antiderivative size = 449, normalized size of antiderivative = 2.38

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c + d \sec(e + fx))^4} dx = \frac{3(2ac^2 - 2acd + ad^2) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right)}{(c^5 + c^4d - 2c^3d^2 - 2c^2d^3 + cd^4 + d^5)\sqrt{-c^2+d^2}} + \frac{6ac^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 18ac^3d \tan(\frac{1}{2}fx + \frac{1}{2}e)^4 + \dots}{(c^5 + c^4d - 2c^3d^2 - 2c^2d^3 + cd^4 + d^5)\sqrt{-c^2+d^2}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^4,x, algorithm="giac")`

output
$$\frac{-1/3*(3*(2*a*c^2 - 2*a*c*d + a*d^2)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + \arctan((c*\tan(1/2*f*x + 1/2*e) - d*\tan(1/2*f*x + 1/2*e))/\sqrt{(-c^2 + d^2)})))/((c^5 + c^4*d - 2*c^3*d^2 - 2*c^2*d^3 + c*d^4 + d^5)*\sqrt{(-c^2 + d^2)}) + (6*a*c^4*\tan(1/2*f*x + 1/2*e)^5 - 18*a*c^3*d*\tan(1/2*f*x + 1/2*e)^5 + 21*a*c^2*d^2*\tan(1/2*f*x + 1/2*e)^5 - 12*a*c*d^3*\tan(1/2*f*x + 1/2*e)^5 + 3*a*d^4*\tan(1/2*f*x + 1/2*e)^5 - 12*a*c^4*\tan(1/2*f*x + 1/2*e)^3 + 24*a*c^3*d*\tan(1/2*f*x + 1/2*e)^3 + 8*a*c^2*d^2*\tan(1/2*f*x + 1/2*e)^3 - 24*a*c*d^3*\tan(1/2*f*x + 1/2*e)^3 + 4*a*d^4*\tan(1/2*f*x + 1/2*e)^3 + 6*a*c^4*\tan(1/2*f*x + 1/2*e) - 6*a*c^3*d*\tan(1/2*f*x + 1/2*e) - 21*a*c^2*d^2*\tan(1/2*f*x + 1/2*e) + 9*a*d^4*\tan(1/2*f*x + 1/2*e))/((c^5 + c^4*d - 2*c^3*d^2 - 2*c^2*d^3 + c*d^4 + d^5)*(c*\tan(1/2*f*x + 1/2*e)^2 - d*\tan(1/2*f*x + 1/2*e)^2 - c - d)^3))/f$$

3.192.9 Mupad [B] (verification not implemented)

Time = 16.76 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.70

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))}{(c + d \sec(e + fx))^4} dx$$

$$= \frac{\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (2ac^2 - 2acd + ad^2)}{(c+d)^3} + \frac{a \tan\left(\frac{e}{2} + \frac{fx}{2}\right) (2c^2 - 6cd + 3d^2)}{(c+d)(c^2 - 2cd + d^2)} - \frac{4a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{(c+d)(c^2 - 2cd + d^2)}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (-3c^3 - 3c^2d + 3cd^2 + 3d^3) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 (-3c^3 + 3c^2d + 3cd^2 - 3d^3) + 3cd^2 \right)} + \frac{a \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (2c - 2d) (c^2 - 2cd + d^2)}{2\sqrt{c+d}(c-d)^{5/2}}\right) (2c^2 - 2cd + d^2)}{f (c+d)^{7/2} (c-d)^{5/2}}$$

input $\text{int}((a + a/\cos(e + f*x))/(\cos(e + f*x)*(c + d/\cos(e + f*x))^4), x)$

output
$$\frac{((\tan(e/2 + (f*x)/2)^5*(2*a*c^2 + a*d^2 - 2*a*c*d))/(c + d)^3 + (a*\tan(e/2 + (f*x)/2)*(2*c^2 - 6*c*d + 3*d^2))/((c + d)*(c^2 - 2*c*d + d^2)) - (4*a*\tan(e/2 + (f*x)/2)^3*(3*c^2 - 6*c*d + d^2))/(3*(c + d)^2*(c - d)))/(f*(\tan(e/2 + (f*x)/2)^2*(3*c*d^2 - 3*c^2*d - 3*c^3 + 3*d^3) - \tan(e/2 + (f*x)/2)^4*(3*c*d^2 + 3*c^2*d - 3*c^3 - 3*d^3) + 3*c*d^2 + 3*c^2*d + c^3 + d^3 - \tan(e/2 + (f*x)/2)^6*(3*c*d^2 - 3*c^2*d + c^3 - d^3)) + (a*\operatorname{atanh}((\tan(e/2 + (f*x)/2)*(2*c - 2*d)*(c^2 - 2*c*d + d^2))/(2*(c + d)^{(1/2)}*(c - d)^{(5/2)})))*(2*c^2 - 2*c*d + d^2))/(f*(c + d)^{(7/2)}*(c - d)^{(5/2)})$$

3.193 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$

3.193.1 Optimal result	1328
3.193.2 Mathematica [A] (verified)	1329
3.193.3 Rubi [A] (verified)	1329
3.193.4 Maple [A] (verified)	1333
3.193.5 Fricas [A] (verification not implemented)	1334
3.193.6 Sympy [F]	1335
3.193.7 Maxima [B] (verification not implemented)	1336
3.193.8 Giac [B] (verification not implemented)	1337
3.193.9 Mupad [B] (verification not implemented)	1338

3.193.1 Optimal result

Integrand size = 31, antiderivative size = 327

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$$

$$= \frac{a^2(24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \operatorname{arctanh}(\sin(e + fx))}{16f}$$

$$- \frac{a^2(4c^5 - 48c^4d - 311c^3d^2 - 448c^2d^3 - 288cd^4 - 64d^5) \tan(e + fx)}{60df}$$

$$- \frac{a^2(8c^4 - 96c^3d - 438c^2d^2 - 464cd^3 - 165d^4) \sec(e + fx) \tan(e + fx)}{240f}$$

$$- \frac{a^2(4c^3 - 48c^2d - 123cd^2 - 64d^3) (c + d \sec(e + fx))^2 \tan(e + fx)}{120df}$$

$$- \frac{a^2(4c^2 - 48cd - 55d^2) (c + d \sec(e + fx))^3 \tan(e + fx)}{120df}$$

$$- \frac{a^2(c - 12d)(c + d \sec(e + fx))^4 \tan(e + fx)}{30df} + \frac{a^2(c + d \sec(e + fx))^5 \tan(e + fx)}{6df}$$

output

```
1/16*a^2*(24*c^4+64*c^3*d+84*c^2*d^2+48*c*d^3+11*d^4)*arctanh(sin(f*x+e))/
f-1/60*a^2*(4*c^5-48*c^4*d-311*c^3*d^2-448*c^2*d^3-288*c*d^4-64*d^5)*tan(f
*x+e)/d/f-1/240*a^2*(8*c^4-96*c^3*d-438*c^2*d^2-464*c*d^3-165*d^4)*sec(f*x
+e)*tan(f*x+e)/f-1/120*a^2*(4*c^3-48*c^2*d-123*c*d^2-64*d^3)*(c+d*sec(f*x+
e))^2*tan(f*x+e)/d/f-1/120*a^2*(4*c^2-48*c*d-55*d^2)*(c+d*sec(f*x+e))^3*ta
n(f*x+e)/d/f-1/30*a^2*(c-12*d)*(c+d*sec(f*x+e))^4*tan(f*x+e)/d/f+1/6*a^2*(
c+d*sec(f*x+e))^5*tan(f*x+e)/d/f
```

3.193.2 Mathematica [A] (verified)

Time = 6.19 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.55

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$$

$$= \frac{a^2(15(24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx)(480(c + d)^4 + 15(8c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \sec(e + fx) + 10d^2(36c^2 + 48cd + 11d^2) \sec^2(e + fx) + 40d^4 \sec^3(e + fx) + 320d(c + d)^3 \tan(e + fx)^2 + 96d^3(2c + d) \tan(e + fx)^4))}{240f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^4,x]`output `(a^2*(15*(24*c^4 + 64*c^3*d + 84*c^2*d^2 + 48*c*d^3 + 11*d^4)*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(480*(c + d)^4 + 15*(8*c^4 + 64*c^3*d + 84*c^2*d^2 + 48*c*d^3 + 11*d^4)*Sec[e + f*x] + 10*d^2*(36*c^2 + 48*c*d + 11*d^2)*Sec[e + f*x]^2 + 40*d^4*Sec[e + f*x]^3 + 320*d*(c + d)^3*Tan[e + f*x]^2 + 96*d^3*(2*c + d)*Tan[e + f*x]^4))/(240*f)`**3.193.3 Rubi [A] (verified)**Time = 0.55 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.19, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4475, 111, 25, 27, 170, 25, 27, 164, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^2(c + d \sec(e + fx))^4 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^4 dx$$

$$\downarrow 4475$$

$$\frac{a^2 \tan(e + fx) \int \frac{(\sec(e + fx)a + a)^{3/2} (c + d \sec(e + fx))^4 d \sec(e + fx)}{\sqrt{a - a \sec(e + fx)}}}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

$$\downarrow 111$$

$$a^2 \tan(e + fx) \left(-\frac{\int -\frac{a^2(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))^2(6c^2+2dc+3d^2+d(9c+2d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx)}{6a^2} - \frac{d\sqrt{a-a \sec(e+fx)}(a \sec(e+fx))}{6a^2} \right)$$

$$\frac{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{\downarrow 25}$$

$$a^2 \tan(e + fx) \left(\int \frac{a^2(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))^2(6c^2+2dc+3d^2+d(9c+2d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d\sqrt{a-a \sec(e+fx)}(a \sec(e+fx))}{6a^2} \right)$$

$$\frac{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{\downarrow 27}$$

$$a^2 \tan(e + fx) \left(\frac{1}{6} \int \frac{(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))^2(6c^2+2dc+3d^2+d(9c+2d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d\sqrt{a-a \sec(e+fx)}(a \sec(e+fx))}{6a^2} \right)$$

$$\frac{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{\downarrow 170}$$

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(-\int -\frac{a^2(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))(30c^3+28dc^2+37d^2c+4d^3+d(48c^2+32dc+19d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx)}{5a^2} - \frac{d(9c+2d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx))}{6a^2} \right) \right)$$

$$\frac{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{\downarrow 25}$$

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\int \frac{a^2(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))(30c^3+28dc^2+37d^2c+4d^3+d(48c^2+32dc+19d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d(9c+2d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx))}{6a^2} \right) \right)$$

$$\frac{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{\downarrow 27}$$

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{1}{5} \int \frac{(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))(30c^3+28dc^2+37d^2c+4d^3+d(48c^2+32dc+19d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d(9c+2d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx))}{6a^2} \right) \right)$$

$$\frac{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{\downarrow 164}$$

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{5}{4} (24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d\sqrt{a-a \sec(e+fx)}(a \sec(e+fx))}{6a^2} \right) \right) \right)$$

$$\downarrow 60$$

3.193. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{5}{4} (24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \left(\frac{3}{2} a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a\sec(e+fx)}} d \sec(e+fx) - \sqrt{a-a\sec(e+fx)} \right) \right) \right) \right)$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{5}{4} (24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{a-a\sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) \right) \right) \right) \right) \right)$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{5}{4} (24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \left(\frac{3}{2} a \left(2a \int \frac{1}{-\frac{(a-a\sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} \right) \right) \right) \right) \right)$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{5}{4} (24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4) \left(\frac{3}{2} a \left(-2 \arctan \left(\frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{a\sec(e+fx)+a}} \right) - \sqrt{a-a\sec(e+fx)} \right) \right) \right) \right) \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^4,x]`

output `-((a^2*(-1/6*(d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2)*(c + d*Sec[e + f*x])^3)/a^2 + (-1/5*(d*(9*c + 2*d)*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2)*(c + d*Sec[e + f*x])^2)/a^2 + (-1/4*(d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2)*(2*(52*c^3 + 56*c^2*d + 48*c*d^2 + 9*d^3) + d*(48*c^2 + 32*c*d + 19*d^2)*Sec[e + f*x]))/a^2 + (5*(24*c^4 + 64*c^3*d + 84*c^2*d^2 + 48*c*d^3 + 11*d^4)*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/a + (3*a*(-2*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/a))/2)/4)/5)/6)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]))`

3.193.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 111 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 164 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))*((g_) + (h_)*(x_)), x_] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`

```
rule 170 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])
```

3.193.4 Maple [A] (verified)

Time = 5.92 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.10

method	result
parts	$-\frac{(4a^2d^3c+2a^2d^4)\left(-\frac{8}{15}-\frac{\sec(fx+e)^4}{5}-\frac{4\sec(fx+e)^2}{15}\right)\tan(fx+e)}{f} + \frac{(2c^4a^2+4a^2c^3d)\tan(fx+e)}{f} + \frac{(6a^2c^2d^2+8a^2d^3c+2a^2d^4)}{f}$
norman	$\frac{17a^2(24c^4+64c^3d+84c^2d^2+48cd^3+11d^4)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^9}{24f} - \frac{a^2(24c^4+64c^3d+84c^2d^2+48cd^3+11d^4)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^{11}}{8f} + \frac{a^2(40c^4+192c^3d+24c^2d^2+16cd^3+5d^4)}{4f}$
parallelrisch	$2a^2\left(-\frac{45\left(c^4+\frac{8}{3}c^3d+\frac{7}{2}c^2d^2+2cd^3+\frac{11}{24}d^4\right)\left(\frac{2\cos(4fx+4e)}{5}+\frac{2}{3}+\cos(2fx+2e)+\frac{\cos(6fx+6e)}{15}\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{4} + \frac{45\left(c^4+\frac{8}{3}c^3d+\frac{7}{2}c^2d^2+2cd^3+\frac{11}{24}d^4\right)}{4f}\right)$
derivativedivides	$c^4a^2\left(\frac{\sec(fx+e)\tan(fx+e)}{2}+\frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)-4a^2c^3d\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+6a^2c^2d^2\left(-\left(-\frac{\sec(fx+e)}{4}\right)\right)$
default	$c^4a^2\left(\frac{\sec(fx+e)\tan(fx+e)}{2}+\frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)-4a^2c^3d\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+6a^2c^2d^2\left(-\left(-\frac{\sec(fx+e)}{4}\right)\right)$
risch	Expression too large to display

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^4,x,method=_RETURNVERBOSE)
```

```
output -(4*a^2*c*d^3+2*a^2*d^4)/f*(-8/15-1/5*sec(f*x+e)^4-4/15*sec(f*x+e)^2)*tan(f*x+e)+(2*a^2*c^4+4*a^2*c^3*d)/f*tan(f*x+e)+(6*a^2*c^2*d^2+8*a^2*c*d^3+a^2*d^4)/f*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln(sec(f*x+e)+tan(f*x+e)))-(4*a^2*c^3*d+12*a^2*c^2*d^2+4*a^2*c*d^3)/f*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)+(a^2*c^4+8*a^2*c^3*d+6*a^2*c^2*d^2)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+a^2*d^4/f*(-(-1/6*sec(f*x+e)^5-5/24*sec(f*x+e)^3-5/16*sec(f*x+e))*tan(f*x+e)+5/16*ln(sec(f*x+e)+tan(f*x+e)))+1/f*ln(sec(f*x+e)+tan(f*x+e))*a^2*c^4
```

3.193.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.18

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$$

$$= \frac{15(24a^2c^4 + 64a^2c^3d + 84a^2c^2d^2 + 48a^2cd^3 + 11a^2d^4) \cos(fx + e)^6 \log(\sin(fx + e) + 1) - 15(24a^2c^4 + 64a^2c^3d + 84a^2c^2d^2 + 48a^2cd^3 + 11a^2d^4) \cos(fx + e)^6}{15(24a^2c^4 + 64a^2c^3d + 84a^2c^2d^2 + 48a^2cd^3 + 11a^2d^4)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^4,x, algorithm="fracas")
```

output $1/480*(15*(24*a^2*c^4 + 64*a^2*c^3*d + 84*a^2*c^2*d^2 + 48*a^2*c*d^3 + 11*a^2*d^4)*\cos(f*x + e)^6*\log(\sin(f*x + e) + 1) - 15*(24*a^2*c^4 + 64*a^2*c^3*d + 84*a^2*c^2*d^2 + 48*a^2*c*d^3 + 11*a^2*d^4)*\cos(f*x + e)^6*\log(-\sin(f*x + e) + 1) + 2*(40*a^2*d^4 + 32*(15*a^2*c^4 + 50*a^2*c^3*d + 60*a^2*c^2*d^2 + 36*a^2*c*d^3 + 8*a^2*d^4)*\cos(f*x + e)^5 + 15*(8*a^2*c^4 + 64*a^2*c^3*d + 84*a^2*c^2*d^2 + 48*a^2*c*d^3 + 11*a^2*d^4)*\cos(f*x + e)^4 + 64*(5*a^2*c^3*d + 15*a^2*c^2*d^2 + 9*a^2*c*d^3 + 2*a^2*d^4)*\cos(f*x + e)^3 + 10*(36*a^2*c^2*d^2 + 48*a^2*c*d^3 + 11*a^2*d^4)*\cos(f*x + e)^2 + 96*(2*a^2*c*d^3 + a^2*d^4)*\cos(f*x + e))*\sin(f*x + e))/(f*\cos(f*x + e)^6)$

3.193.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx \\ &= a^2 \left(\int c^4 \sec(e + fx) dx + \int 2c^4 \sec^2(e + fx) dx + \int c^4 \sec^3(e + fx) dx \right. \\ & \quad + \int d^4 \sec^5(e + fx) dx + \int 2d^4 \sec^6(e + fx) dx + \int d^4 \sec^7(e + fx) dx \\ & \quad + \int 4cd^3 \sec^4(e + fx) dx + \int 8cd^3 \sec^5(e + fx) dx + \int 4cd^3 \sec^6(e + fx) dx \\ & \quad + \int 6c^2d^2 \sec^3(e + fx) dx + \int 12c^2d^2 \sec^4(e + fx) dx + \int 6c^2d^2 \sec^5(e + fx) dx \\ & \quad \left. + \int 4c^3d \sec^2(e + fx) dx + \int 8c^3d \sec^3(e + fx) dx + \int 4c^3d \sec^4(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c+d*sec(f*x+e))**4,x)`

output `a**2*(Integral(c**4*sec(e + f*x), x) + Integral(2*c**4*sec(e + f*x)**2, x) + Integral(c**4*sec(e + f*x)**3, x) + Integral(d**4*sec(e + f*x)**5, x) + Integral(2*d**4*sec(e + f*x)**6, x) + Integral(d**4*sec(e + f*x)**7, x) + Integral(4*c*d**3*sec(e + f*x)**4, x) + Integral(8*c*d**3*sec(e + f*x)**5, x) + Integral(4*c*d**3*sec(e + f*x)**6, x) + Integral(6*c**2*d**2*sec(e + f*x)**3, x) + Integral(12*c**2*d**2*sec(e + f*x)**4, x) + Integral(6*c**2*d**2*sec(e + f*x)**5, x) + Integral(4*c**3*d*sec(e + f*x)**2, x) + Integral(8*c**3*d*sec(e + f*x)**3, x) + Integral(4*c**3*d*sec(e + f*x)**4, x))`

3.193.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs. $2(313) = 626$.

Time = 0.22 (sec) , antiderivative size = 683, normalized size of antiderivative = 2.09

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$$

$$= \frac{640 (\tan (fx + e)^3 + 3 \tan (fx + e)) a^2 c^3 d + 1920 (\tan (fx + e)^3 + 3 \tan (fx + e)) a^2 c^2 d^2 + 128 (3 \tan (fx + e) + \tan (fx + e)^3) a^2 c d^3 + 64 (3 \tan (fx + e)^5 + 10 \tan (fx + e)^3 + 15 \tan (fx + e)) a^2 d^4 - 5 a^2 d^4 (2 (15 \sin (fx + e)^5 - 40 \sin (fx + e)^3 + 33 \sin (fx + e)) / (\sin (fx + e)^6 - 3 \sin (fx + e)^4 + 3 \sin (fx + e)^2 - 1) - 15 \log (\sin (fx + e) + 1) + 15 \log (\sin (fx + e) - 1)) - 180 a^2 c^2 d^2 (2 (3 \sin (fx + e)^3 - 5 \sin (fx + e)) / (\sin (fx + e)^4 - 2 \sin (fx + e)^2 + 1) - 3 \log (\sin (fx + e) + 1) + 3 \log (\sin (fx + e) - 1)) - 240 a^2 c d^3 (2 (3 \sin (fx + e)^3 - 5 \sin (fx + e)) / (\sin (fx + e)^4 - 2 \sin (fx + e)^2 + 1) - 3 \log (\sin (fx + e) + 1) + 3 \log (\sin (fx + e) - 1)) - 30 a^2 d^4 (2 (3 \sin (fx + e)^3 - 5 \sin (fx + e)) / (\sin (fx + e)^4 - 2 \sin (fx + e)^2 + 1) - 3 \log (\sin (fx + e) + 1) + 3 \log (\sin (fx + e) - 1)) - 120 a^2 c^4 (2 \sin (fx + e) / (\sin (fx + e)^2 - 1) - \log (\sin (fx + e) + 1) + \log (\sin (fx + e) - 1)) - 960 a^2 c^3 d (2 \sin (fx + e) / (\sin (fx + e)^2 - 1) - \log (\sin (fx + e) + 1) + \log (\sin (fx + e) - 1)) - 720 a^2 c^2 d^2 (2 \sin (fx + e) / (\sin (fx + e)^2 - 1) - \log (\sin (fx + e) + 1) + \log (\sin (fx + e) - 1)) + 480 a^2 c^4 \log (\sec (fx + e) + \tan (fx + e)) + 960 a^2 c^4 \tan (fx + e) + 1920 a^2 c^3 d \tan (fx + e)) / f$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^4,x, algorithm="maxima")
```

```
output 1/480*(640*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^2*c^3*d + 1920*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^2*c^2*d^2 + 128*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^2*c*d^3 + 64*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^2*c*d^3 + 64*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^2*d^4 - 5*a^2*d^4*(2*(15*sin(f*x + e)^5 - 40*sin(f*x + e)^3 + 33*sin(f*x + e))/(sin(f*x + e)^6 - 3*sin(f*x + e)^4 + 3*sin(f*x + e)^2 - 1) - 15*log(sin(f*x + e) + 1) + 15*log(sin(f*x + e) - 1)) - 180*a^2*c^2*d^2*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 240*a^2*c*d^3*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 30*a^2*d^4*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 120*a^2*c^4*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 960*a^2*c^3*d*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 720*a^2*c^2*d^2*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 480*a^2*c^4*log(sec(f*x + e) + tan(f*x + e)) + 960*a^2*c^4*tan(f*x + e) + 1920*a^2*c^3*d*tan(f*x + e))/f
```

3.193.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 736 vs. $2(313) = 626$.

Time = 0.41 (sec) , antiderivative size = 736, normalized size of antiderivative = 2.25

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^4,x, algorithm="giac")`

output

```
1/240*(15*(24*a^2*c^4 + 64*a^2*c^3*d + 84*a^2*c^2*d^2 + 48*a^2*c*d^3 + 11*
a^2*d^4)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*(24*a^2*c^4 + 64*a^2*c^3*
d + 84*a^2*c^2*d^2 + 48*a^2*c*d^3 + 11*a^2*d^4)*log(abs(tan(1/2*f*x + 1/2*
e) - 1)) - 2*(360*a^2*c^4*tan(1/2*f*x + 1/2*e)^11 + 960*a^2*c^3*d*tan(1/2*
f*x + 1/2*e)^11 + 1260*a^2*c^2*d^2*tan(1/2*f*x + 1/2*e)^11 + 720*a^2*c*d^3
*tan(1/2*f*x + 1/2*e)^11 + 165*a^2*d^4*tan(1/2*f*x + 1/2*e)^11 - 2040*a^2*
c^4*tan(1/2*f*x + 1/2*e)^9 - 5440*a^2*c^3*d*tan(1/2*f*x + 1/2*e)^9 - 7140*
a^2*c^2*d^2*tan(1/2*f*x + 1/2*e)^9 - 4080*a^2*c*d^3*tan(1/2*f*x + 1/2*e)^9
- 935*a^2*d^4*tan(1/2*f*x + 1/2*e)^9 + 4560*a^2*c^4*tan(1/2*f*x + 1/2*e)^
7 + 13440*a^2*c^3*d*tan(1/2*f*x + 1/2*e)^7 + 15480*a^2*c^2*d^2*tan(1/2*f*x
+ 1/2*e)^7 + 10272*a^2*c*d^3*tan(1/2*f*x + 1/2*e)^7 + 1986*a^2*d^4*tan(1/
2*f*x + 1/2*e)^7 - 5040*a^2*c^4*tan(1/2*f*x + 1/2*e)^5 - 17280*a^2*c^3*d*t
an(1/2*f*x + 1/2*e)^5 - 19080*a^2*c^2*d^2*tan(1/2*f*x + 1/2*e)^5 - 11232*a
^2*c*d^3*tan(1/2*f*x + 1/2*e)^5 - 3006*a^2*d^4*tan(1/2*f*x + 1/2*e)^5 + 27
60*a^2*c^4*tan(1/2*f*x + 1/2*e)^3 + 11200*a^2*c^3*d*tan(1/2*f*x + 1/2*e)^3
+ 13980*a^2*c^2*d^2*tan(1/2*f*x + 1/2*e)^3 + 7440*a^2*c*d^3*tan(1/2*f*x +
1/2*e)^3 + 1305*a^2*d^4*tan(1/2*f*x + 1/2*e)^3 - 600*a^2*c^4*tan(1/2*f*x
+ 1/2*e) - 2880*a^2*c^3*d*tan(1/2*f*x + 1/2*e) - 4500*a^2*c^2*d^2*tan(1/2*
f*x + 1/2*e) - 3120*a^2*c*d^3*tan(1/2*f*x + 1/2*e) - 795*a^2*d^4*tan(1/2*f
*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^6)/f
```

3.193.9 Mupad [B] (verification not implemented)

Time = 17.08 (sec) , antiderivative size = 484, normalized size of antiderivative = 1.48

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^4 dx$$

$$= \frac{\left(-3a^2c^4 - 8a^2c^3d - \frac{21a^2c^2d^2}{2} - 6a^2cd^3 - \frac{11a^2d^4}{8}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11} + \left(17a^2c^4 + \frac{136a^2c^3d}{3} + \frac{119a^2c^2d^2}{2} + \frac{a^2 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)(24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4)}{4(6c^4 + 16c^3d + 21c^2d^2 + 12cd^3 + \frac{11d^4}{4})}\right)}{8f} (24c^4 + 64c^3d + 84c^2d^2 + 48cd^3 + 11d^4)}{8f}$$

input `int(((a + a/cos(e + f*x))^2*(c + d/cos(e + f*x))^4)/cos(e + f*x),x)`

output

```
(tan(e/2 + (f*x)/2)*(5*a^2*c^4 + (53*a^2*d^4)/8 + 26*a^2*c*d^3 + 24*a^2*c^3*d + (75*a^2*c^2*d^2)/2) - tan(e/2 + (f*x)/2)^11*(3*a^2*c^4 + (11*a^2*d^4)/8 + 6*a^2*c*d^3 + 8*a^2*c^3*d + (21*a^2*c^2*d^2)/2) + tan(e/2 + (f*x)/2)^9*(17*a^2*c^4 + (187*a^2*d^4)/24 + 34*a^2*c*d^3 + (136*a^2*c^3*d)/3 + (119*a^2*c^2*d^2)/2) - tan(e/2 + (f*x)/2)^3*(23*a^2*c^4 + (87*a^2*d^4)/8 + 62*a^2*c*d^3 + (280*a^2*c^3*d)/3 + (233*a^2*c^2*d^2)/2) - tan(e/2 + (f*x)/2)^7*(38*a^2*c^4 + (331*a^2*d^4)/20 + (428*a^2*c*d^3)/5 + 112*a^2*c^3*d + 129*a^2*c^2*d^2) + tan(e/2 + (f*x)/2)^5*(42*a^2*c^4 + (501*a^2*d^4)/20 + (468*a^2*c*d^3)/5 + 144*a^2*c^3*d + 159*a^2*c^2*d^2)/(f*(15*tan(e/2 + (f*x)/2)^4 - 6*tan(e/2 + (f*x)/2)^2 - 20*tan(e/2 + (f*x)/2)^6 + 15*tan(e/2 + (f*x)/2)^8 - 6*tan(e/2 + (f*x)/2)^10 + tan(e/2 + (f*x)/2)^12 + 1)) + (a^2*atanh((tan(e/2 + (f*x)/2)*(48*c*d^3 + 64*c^3*d + 24*c^4 + 11*d^4 + 84*c^2*d^2))/(4*(12*c*d^3 + 16*c^3*d + 6*c^4 + (11*d^4)/4 + 21*c^2*d^2)))*(48*c*d^3 + 64*c^3*d + 24*c^4 + 11*d^4 + 84*c^2*d^2))/(8*f)
```

3.194 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx$

3.194.1 Optimal result	1339
3.194.2 Mathematica [A] (verified)	1340
3.194.3 Rubi [A] (verified)	1340
3.194.4 Maple [A] (verified)	1344
3.194.5 Fricas [A] (verification not implemented)	1344
3.194.6 Sympy [F]	1345
3.194.7 Maxima [B] (verification not implemented)	1346
3.194.8 Giac [B] (verification not implemented)	1346
3.194.9 Mupad [B] (verification not implemented)	1347

3.194.1 Optimal result

Integrand size = 31, antiderivative size = 242

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx$$

$$= \frac{3a^2(2c + d)(2c^2 + 3cd + 2d^2) \operatorname{arctanh}(\sin(e + fx))}{8f}$$

$$- \frac{a^2(c^4 - 10c^3d - 44c^2d^2 - 40cd^3 - 12d^4) \tan(e + fx)}{10df}$$

$$- \frac{a^2(2c^3 - 20c^2d - 57cd^2 - 30d^3) \sec(e + fx) \tan(e + fx)}{40f}$$

$$- \frac{a^2(c^2 - 10cd - 12d^2)(c + d \sec(e + fx))^2 \tan(e + fx)}{20df}$$

$$- \frac{a^2(c - 10d)(c + d \sec(e + fx))^3 \tan(e + fx)}{20df} + \frac{a^2(c + d \sec(e + fx))^4 \tan(e + fx)}{5df}$$

output

```
3/8*a^2*(2*c+d)*(2*c^2+3*c*d+2*d^2)*arctanh(sin(f*x+e))/f-1/10*a^2*(c^4-10*c^3*d-44*c^2*d^2-40*c*d^3-12*d^4)*tan(f*x+e)/d/f-1/40*a^2*(2*c^3-20*c^2*d-57*c*d^2-30*d^3)*sec(f*x+e)*tan(f*x+e)/f-1/20*a^2*(c^2-10*c*d-12*d^2)*(c+d*sec(f*x+e))^2*tan(f*x+e)/d/f-1/20*a^2*(c-10*d)*(c+d*sec(f*x+e))^3*tan(f*x+e)/d/f+1/5*a^2*(c+d*sec(f*x+e))^4*tan(f*x+e)/d/f
```

3.194.2 Mathematica [A] (verified)

Time = 3.45 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.58

$$\int \sec(e+fx)(a+a\sec(e+fx))^2(c+d\sec(e+fx))^3 dx$$

$$= \frac{a^2(15(4c^3+8c^2d+7cd^2+2d^3)\operatorname{arctanh}(\sin(e+fx)) + \tan(e+fx)(5(4c^3+24c^2d+21cd^2+6d^3)\sec(e+fx) + 10d^2(3c+2d)\sec(e+fx)^3 + 8(10(c+d)^3+5d(c+d)^2\tan(e+fx)^2+d^3\tan(e+fx)^4))}{40f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^3,x]`output `(a^2*(15*(4*c^3 + 8*c^2*d + 7*c*d^2 + 2*d^3)*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(5*(4*c^3 + 24*c^2*d + 21*c*d^2 + 6*d^3)*Sec[e + f*x] + 10*d^2*(3*c + 2*d)*Sec[e + f*x]^3 + 8*(10*(c + d)^3 + 5*d*(c + d)^2*Tan[e + f*x]^2 + d^3*Tan[e + f*x]^4)))/(40*f)`**3.194.3 Rubi [A] (verified)**Time = 0.46 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.25, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4475, 111, 25, 27, 164, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e+fx)(a\sec(e+fx)+a)^2(c+d\sec(e+fx))^3 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^2\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)^3 dx$$

$$\downarrow 4475$$

$$-\frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{3/2}(c+d\sec(e+fx))^3 d\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}}}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

$$\downarrow 111$$

$$a^2 \tan(e + fx) \left(- \frac{\int - \frac{a^2 (\sec(e+fx)a+a)^{3/2} (c+d \sec(e+fx)) (5c^2+2dc+2d^2+d(7c+2d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx)}{5a^2} - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{5a^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 25

$$a^2 \tan(e + fx) \left(\int \frac{a^2 (\sec(e+fx)a+a)^{3/2} (c+d \sec(e+fx)) (5c^2+2dc+2d^2+d(7c+2d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{5a^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{1}{5} \int \frac{(\sec(e+fx)a+a)^{3/2} (c+d \sec(e+fx)) (5c^2+2dc+2d^2+d(7c+2d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e + fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{5a^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 164

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{5}{4} (2c + d) (2c^2 + 3cd + 2d^2) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}} d \sec(e + fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{5a^2} \right) \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{5}{4} (2c + d) (2c^2 + 3cd + 2d^2) \left(\frac{3}{2} a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}} d \sec(e + fx) - \frac{\sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{2a} \right) \right) \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{5}{4} (2c + d) (2c^2 + 3cd + 2d^2) \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e + fx) - \frac{\sqrt{a-a \sec(e+fx)}}{a} \right) \right) \right) \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{5}{4} (2c + d) (2c^2 + 3cd + 2d^2) \left(\frac{3}{2} a \left(2a \int \frac{1}{-\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a}}{a} \right) \right) \right) \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 218

3.194. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx$

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{5}{4} (2c + d) (2c^2 + 3cd + 2d^2) \left(\frac{3}{2} a \left(-2 \arctan \left(\frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{a \sec(e + fx) + a}} \right) - \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{a} \right) \right) \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^3,x]`

output `-((a^2*(-1/5*(d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2)*(c + d*Sec[e + f*x])^2)/a^2 + (-1/4*(d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2)*(2*(8*c^2 + 5*c*d + 2*d^2) + d*(7*c + 2*d)*Sec[e + f*x]))/a^2 + (5*(2*c + d)*(2*c^2 + 3*c*d + 2*d^2)*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/a + (3*a*(-2*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/a))/2)/4)/5)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))`

3.194.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 111 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`

rule 164 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x))*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.194.4 Maple [A] (verified)

Time = 5.18 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.11

method	result
parts	$\frac{(3a^2cd^2+2a^2d^3)\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)}{f} + \frac{(2a^2c^3+3a^2c^2d)\tan(fx+e)}{f}$
norman	$\frac{7a^2(4c^3+8c^2d+7cd^2+2d^3)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{2f} - \frac{3a^2(4c^3+8c^2d+7cd^2+2d^3)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^9}{4f} - \frac{8a^2(15c^3+35c^2d+25cd^2+9d^3)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{5f} - \frac{1}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)^5}$
parallelrisch	$4\left(-\frac{15\left(c+\frac{d}{2}\right)\left(\frac{\cos(5fx+5e)}{10}+\frac{\cos(3fx+3e)}{2}+\cos(fx+e)\right)\left(c^2+\frac{3}{2}cd+d^2\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{4} + \frac{15\left(c+\frac{d}{2}\right)\left(\frac{\cos(5fx+5e)}{10}+\frac{\cos(3fx+3e)}{2}\right)}{4}\right)$
derivativedivides	$a^2c^3\left(\frac{\sec(fx+e)\tan(fx+e)}{2}+\frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)-3a^2c^2d\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+3a^2cd^2\left(-\left(-\frac{\sec(fx+e)}{4}\right)^5\right)$
default	$a^2c^3\left(\frac{\sec(fx+e)\tan(fx+e)}{2}+\frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)-3a^2c^2d\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+3a^2cd^2\left(-\left(-\frac{\sec(fx+e)}{4}\right)^5\right)$
risch	$-\frac{ia^2(-80c^3-160cd^2-48d^3-200c^2d-105cd^2e^{i(fx+e)}-800cd^2e^{2i(fx+e)}-330cd^2e^{3i(fx+e)}-480cd^2e^{6i(fx+e)}-120c^2d^3e^{9i(fx+e)})}{(1-\tan^2(fx+e))^5}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$(3a^2cd^2+2a^2d^3)/f*(-(-1/4*\sec(f*x+e)^3-3/8*\sec(f*x+e))*\tan(f*x+e)+3/8*\ln(\sec(f*x+e)+\tan(f*x+e)))+(2a^2c^3+3a^2c^2d)/f*\tan(f*x+e)-(3a^2c^2d+6a^2c*d^2+a^2d^3)/f*(-2/3-1/3*\sec(f*x+e)^2)*\tan(f*x+e)+(a^2c^3+6a^2c^2d+3a^2cd^2)/f*(1/2*\sec(f*x+e)*\tan(f*x+e)+1/2*\ln(\sec(f*x+e)+\tan(f*x+e)))+1/f*\ln(\sec(f*x+e)+\tan(f*x+e))*a^2c^3-a^2d^3/f*(-8/15-1/5*\sec(f*x+e)^4-4/15*\sec(f*x+e)^2)*\tan(f*x+e)$$

3.194.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.21

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx$$

$$= \frac{15(4a^2c^3 + 8a^2c^2d + 7a^2cd^2 + 2a^2d^3) \cos(fx + e)^5 \log(\sin(fx + e) + 1) - 15(4a^2c^3 + 8a^2c^2d + 7a^2cd^2 + 2a^2d^3) \cos(fx + e)^5 \log(\sin(fx + e) - 1) - 15(4a^2c^3 + 8a^2c^2d + 7a^2cd^2 + 2a^2d^3) \cos(fx + e)^5 \log(\sin(fx + e) + 1) - 15(4a^2c^3 + 8a^2c^2d + 7a^2cd^2 + 2a^2d^3) \cos(fx + e)^5 \log(\sin(fx + e) - 1)}{(1 - \tan^2(fx + e))^5}$$

3.194. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^3,x, algorithm="fricas")`

output `1/80*(15*(4*a^2*c^3 + 8*a^2*c^2*d + 7*a^2*c*d^2 + 2*a^2*d^3)*cos(f*x + e)^5*log(sin(f*x + e) + 1) - 15*(4*a^2*c^3 + 8*a^2*c^2*d + 7*a^2*c*d^2 + 2*a^2*d^3)*cos(f*x + e)^5*log(-sin(f*x + e) + 1) + 2*(8*a^2*d^3 + 8*(10*a^2*c^3 + 25*a^2*c^2*d + 20*a^2*c*d^2 + 6*a^2*d^3)*cos(f*x + e)^4 + 5*(4*a^2*c^3 + 24*a^2*c^2*d + 21*a^2*c*d^2 + 6*a^2*d^3)*cos(f*x + e)^3 + 8*(5*a^2*c^2*d + 10*a^2*c*d^2 + 3*a^2*d^3)*cos(f*x + e)^2 + 10*(3*a^2*c*d^2 + 2*a^2*d^3)*cos(f*x + e))*sin(f*x + e)/(f*cos(f*x + e)^5)`

3.194.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx \\ &= a^2 \left(\int c^3 \sec(e + fx) dx + \int 2c^3 \sec^2(e + fx) dx + \int c^3 \sec^3(e + fx) dx \right. \\ & \quad + \int d^3 \sec^4(e + fx) dx + \int 2d^3 \sec^5(e + fx) dx + \int d^3 \sec^6(e + fx) dx \\ & \quad + \int 3cd^2 \sec^3(e + fx) dx + \int 6cd^2 \sec^4(e + fx) dx + \int 3cd^2 \sec^5(e + fx) dx \\ & \quad \left. + \int 3c^2d \sec^2(e + fx) dx + \int 6c^2d \sec^3(e + fx) dx + \int 3c^2d \sec^4(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c+d*sec(f*x+e))**3,x)`

output `a**2*(Integral(c**3*sec(e + f*x), x) + Integral(2*c**3*sec(e + f*x)**2, x) + Integral(c**3*sec(e + f*x)**3, x) + Integral(d**3*sec(e + f*x)**4, x) + Integral(2*d**3*sec(e + f*x)**5, x) + Integral(d**3*sec(e + f*x)**6, x) + Integral(3*c*d**2*sec(e + f*x)**3, x) + Integral(6*c*d**2*sec(e + f*x)**4, x) + Integral(3*c*d**2*sec(e + f*x)**5, x) + Integral(3*c**2*d*sec(e + f*x)**2, x) + Integral(6*c**2*d*sec(e + f*x)**3, x) + Integral(3*c**2*d*sec(e + f*x)**4, x))`

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output `1/40*(15*(4*a^2*c^3 + 8*a^2*c^2*d + 7*a^2*c*d^2 + 2*a^2*d^3)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*(4*a^2*c^3 + 8*a^2*c^2*d + 7*a^2*c*d^2 + 2*a^2*d^3)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(60*a^2*c^3*tan(1/2*f*x + 1/2*e)^9 + 120*a^2*c^2*d*tan(1/2*f*x + 1/2*e)^9 + 105*a^2*c*d^2*tan(1/2*f*x + 1/2*e)^9 + 30*a^2*d^3*tan(1/2*f*x + 1/2*e)^9 - 280*a^2*c^3*tan(1/2*f*x + 1/2*e)^7 - 560*a^2*c^2*d*tan(1/2*f*x + 1/2*e)^7 - 490*a^2*c*d^2*tan(1/2*f*x + 1/2*e)^7 - 140*a^2*d^3*tan(1/2*f*x + 1/2*e)^7 + 480*a^2*c^3*tan(1/2*f*x + 1/2*e)^5 + 1120*a^2*c^2*d*tan(1/2*f*x + 1/2*e)^5 + 800*a^2*c*d^2*tan(1/2*f*x + 1/2*e)^5 + 288*a^2*d^3*tan(1/2*f*x + 1/2*e)^5 - 360*a^2*c^3*tan(1/2*f*x + 1/2*e)^3 - 1040*a^2*c^2*d*tan(1/2*f*x + 1/2*e)^3 - 790*a^2*c*d^2*tan(1/2*f*x + 1/2*e)^3 - 180*a^2*d^3*tan(1/2*f*x + 1/2*e)^3 + 100*a^2*c^3*tan(1/2*f*x + 1/2*e) + 360*a^2*c^2*d*tan(1/2*f*x + 1/2*e) + 375*a^2*c*d^2*tan(1/2*f*x + 1/2*e) + 130*a^2*d^3*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^5)/f`

3.194.9 Mupad [B] (verification not implemented)

Time = 17.06 (sec) , antiderivative size = 394, normalized size of antiderivative = 1.63

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 dx$$

$$= \frac{3a^2 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) (2c+d) (2c^2+3cd+2d^2)}{2(6c^3+12c^2d+\frac{21c^2d^2}{2}+3d^3)}\right) (2c+d) (2c^2+3cd+2d^2)}{4f} - \frac{\left(3a^2c^3 + 6a^2c^2d + \frac{21a^2cd^2}{4} + \frac{3a^2d^3}{2}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9 + \left(-14a^2c^3 - 28a^2c^2d - \frac{49a^2cd^2}{2} - 7a^2d^3\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)^2 - 1}$$

input `int(((a + a/cos(e + f*x))^2*(c + d/cos(e + f*x))^3)/cos(e + f*x),x)`

output $(3a^2 \operatorname{atanh}((3 \tan(e/2 + (f*x)/2) * (2c + d) * (3cd + 2c^2 + 2d^2)) / (2 * ((21cd^2)/2 + 12c^2d + 6c^3 + 3d^3))) * (2c + d) * (3cd + 2c^2 + 2d^2)) / (4f) - (\tan(e/2 + (f*x)/2)^9 * (3a^2c^3 + (3a^2d^3)/2 + (21a^2cd^2)/4 + 6a^2c^2d) - \tan(e/2 + (f*x)/2)^7 * (14a^2c^3 + 7a^2d^3 + (49a^2cd^2)/2 + 28a^2c^2d) - \tan(e/2 + (f*x)/2)^3 * (18a^2c^3 + 9a^2d^3 + (79a^2cd^2)/2 + 52a^2c^2d) + \tan(e/2 + (f*x)/2)^5 * (24a^2c^3 + (72a^2d^3)/5 + 40a^2cd^2 + 56a^2c^2d) + \tan(e/2 + (f*x)/2) * (5a^2c^3 + (13a^2d^3)/2 + (75a^2cd^2)/4 + 18a^2c^2d)) / (f * (5 \tan(e/2 + (f*x)/2)^2 - 10 \tan(e/2 + (f*x)/2)^4 + 10 \tan(e/2 + (f*x)/2)^6 - 5 \tan(e/2 + (f*x)/2)^8 + \tan(e/2 + (f*x)/2)^{10} - 1))$

3.195 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$

3.195.1 Optimal result	1349
3.195.2 Mathematica [A] (verified)	1350
3.195.3 Rubi [A] (verified)	1350
3.195.4 Maple [A] (verified)	1353
3.195.5 Fricas [A] (verification not implemented)	1354
3.195.6 Sympy [F]	1355
3.195.7 Maxima [A] (verification not implemented)	1355
3.195.8 Giac [A] (verification not implemented)	1356
3.195.9 Mupad [B] (verification not implemented)	1357

3.195.1 Optimal result

Integrand size = 31, antiderivative size = 176

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$$

$$= \frac{a^2(12c^2 + 16cd + 7d^2) \operatorname{arctanh}(\sin(e + fx))}{8f} - \frac{a^2(c^3 - 8c^2d - 20cd^2 - 8d^3) \tan(e + fx)}{6df}$$

$$- \frac{a^2(2c(c - 8d) - 21d^2) \sec(e + fx) \tan(e + fx)}{24f}$$

$$- \frac{a^2(c - 8d)(c + d \sec(e + fx))^2 \tan(e + fx)}{12df} + \frac{a^2(c + d \sec(e + fx))^3 \tan(e + fx)}{4df}$$

```
output 1/8*a^2*(12*c^2+16*c*d+7*d^2)*arctanh(sin(f*x+e))/f-1/6*a^2*(c^3-8*c^2*d-20*c*d^2-8*d^3)*tan(f*x+e)/d/f-1/24*a^2*(2*c*(c-8*d)-21*d^2)*sec(f*x+e)*tan(f*x+e)/f-1/12*a^2*(c-8*d)*(c+d*sec(f*x+e))^2*tan(f*x+e)/d/f+1/4*a^2*(c+d*sec(f*x+e))^3*tan(f*x+e)/d/f
```

3.195.2 Mathematica [A] (verified)

Time = 1.68 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.57

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$$

$$= \frac{a^2(3(12c^2 + 16cd + 7d^2) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx)(3(4c^2 + 16cd + 7d^2) \sec(e + fx) + 6d^2 \sec^3))}{24f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^2,x]`

output $(a^2*(3*(12*c^2 + 16*c*d + 7*d^2)*\operatorname{ArcTanh}[\operatorname{Sin}[e + f*x]] + \operatorname{Tan}[e + f*x]*(3*(4*c^2 + 16*c*d + 7*d^2)*\operatorname{Sec}[e + f*x] + 6*d^2*\operatorname{Sec}[e + f*x]^3 + 16*(c + d)*(3*(c + d) + d*\operatorname{Tan}[e + f*x]^2)))/(24*f)$

3.195.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.53, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4475, 101, 25, 27, 90, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^2(c + d \sec(e + fx))^2 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 dx$$

$$\downarrow 4475$$

$$\frac{a^2 \tan(e + fx) \int \frac{(\sec(e + fx)a + a)^{3/2}(c + d \sec(e + fx))^2}{\sqrt{a - a \sec(e + fx)}} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

$$\downarrow 101$$

$$\frac{a^2 \tan(e + fx) \left(-\frac{\int \frac{a^2 (\sec(e + fx)a + a)^{3/2} (4c^2 + 2dc + d^2 + d(5c + 2d) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)}} d \sec(e + fx)}{4a^2} - \frac{d \sqrt{a - a \sec(e + fx)} (a \sec(e + fx) + a)^{5/2} (c + d \sec(e + fx))}{4a^2} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

3.195. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$

↓ 25

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(\sec(e+fx)a+a)^{3/2}(4c^2+2dc+d^2+d(5c+2d)\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}} d\sec(e+fx)}{4a^2} - \frac{d\sqrt{a-a\sec(e+fx)}(a\sec(e+fx)+a)^{5/2}(c+d\sec(e+fx))}{4a^2} \right)$$

$$f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{\frac{1}{4} \int \frac{(\sec(e+fx)a+a)^{3/2}(4c^2+2dc+d^2+d(5c+2d)\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}} d\sec(e+fx) - \frac{d\sqrt{a-a\sec(e+fx)}(a\sec(e+fx)+a)^{5/2}(c+d\sec(e+fx))}{4a^2}}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \right)$$

↓ 90

$$a^2 \tan(e + fx) \left(\frac{\frac{1}{4} \left(\frac{1}{3} (12c^2 + 16cd + 7d^2) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a\sec(e+fx)}} d\sec(e+fx) - \frac{d(5c+2d)\sqrt{a-a\sec(e+fx)}(a\sec(e+fx)+a)^{5/2}}{3a^2} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \right)$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{\frac{1}{4} \left(\frac{1}{3} (12c^2 + 16cd + 7d^2) \left(\frac{3}{2} a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a\sec(e+fx)}} d\sec(e+fx) - \frac{\sqrt{a-a\sec(e+fx)}(a\sec(e+fx)+a)^{3/2}}{2a} \right) \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \right)$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{\frac{1}{4} \left(\frac{1}{3} (12c^2 + 16cd + 7d^2) \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d\sec(e+fx) - \frac{\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{a} \right) \right) \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \right)$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{\frac{1}{4} \left(\frac{1}{3} (12c^2 + 16cd + 7d^2) \left(\frac{3}{2} a \left(2a \int \frac{1}{-\frac{(a-a\sec(e+fx))a}{\sec(e+fx)a+a} - a} d\frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} - \frac{\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{a} \right) \right) \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \right)$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{\frac{1}{4} \left(\frac{1}{3} (12c^2 + 16cd + 7d^2) \left(\frac{3}{2} a \left(-2 \arctan \left(\frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{a\sec(e+fx)+a}} \right) - \frac{\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{a} \right) \right) \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^2,x]`

3.195. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$


```
output 
$$-\left(\frac{a^2(-1/4(d\sqrt{a - a\sec[e + fx]})(a + a\sec[e + fx])^{5/2}(c + d\sec[e + fx]))}{a^2} + \frac{(-1/3(d(5c + 2d)\sqrt{a - a\sec[e + fx]})(a + a\sec[e + fx])^{5/2})}{a^2} + \frac{((12c^2 + 16cd + 7d^2)(-1/2(\sqrt{a - a\sec[e + fx]})(a + a\sec[e + fx])^{3/2})}{a} + \frac{3a(-2\text{ArcTan}[\sqrt{a - a\sec[e + fx]}/\sqrt{a + a\sec[e + fx]}) - (\sqrt{a - a\sec[e + fx]}\sqrt{a + a\sec[e + fx]})/a)}{2}\right)/3/4 * \text{Tan}[e + fx] / (f\sqrt{a - a\sec[e + fx]} * \sqrt{a + a\sec[e + fx]})$$

```

3.195.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 45 Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

```
rule 90 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]
```

- rule 101 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)*(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)n(e + f*x)p*Simp[a2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 218 `Int[((a_) + (b_.)*(x_)2)(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))(p_.)(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))(m_)(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))(n_), x_Symbol] := Simp[a2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)(p - 1)(a + b*x)(m - 1/2)((c + d*x)n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a2 - b2, 0] && NeQ[c2 - d2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.195.4 Maple [A] (verified)

Time = 4.36 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.15

method	result
parts	$-\frac{(2a^2cd+2a^2d^2)\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)}{f} + \frac{(2a^2c^2+2a^2cd)\tan(fx+e)}{f} + \frac{(a^2c^2+4a^2cd+a^2d^2)\left(\frac{\sec(fx+e)}{2}\right)}{f(3+\cos(fx+e))}$
parallelrisch	$4a^2\left(-\frac{3\left(c^2+\frac{4}{3}cd+\frac{7}{12}d^2\right)\left(\frac{3}{4}+\frac{\cos(4fx+4e)}{4}+\cos(2fx+2e)\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{2} + \frac{3\left(c^2+\frac{4}{3}cd+\frac{7}{12}d^2\right)\left(\frac{3}{4}+\frac{\cos(4fx+4e)}{4}+\cos(2fx+2e)\right)}{2}\right)$
norman	$\frac{11a^2(12c^2+16cd+7d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{12f} - \frac{a^2(12c^2+16cd+7d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{4f} + \frac{a^2(20c^2+48cd+25d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{4f} - \frac{a^2(156c^2+272cd+125d^2)}{4f\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)^4}$
derivativedivides	$a^2c^2\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) - 2a^2cd\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + a^2d^2\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3}{8}\ln(\sec(fx+e)+\tan(fx+e))\right)\right)$
default	$a^2c^2\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) - 2a^2cd\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + a^2d^2\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3}{8}\ln(\sec(fx+e)+\tan(fx+e))\right)\right)$
risch	$-\frac{ia^2(12c^2e^{7i(fx+e)}+48cde^{7i(fx+e)}+21d^2e^{7i(fx+e)}-48c^2e^{6i(fx+e)}-48cde^{6i(fx+e)}+12c^2e^{5i(fx+e)}+48cde^{5i(fx+e)}+21d^2e^{5i(fx+e)}-48c^2e^{4i(fx+e)}-48cde^{4i(fx+e)}+12c^2e^{3i(fx+e)}+48cde^{3i(fx+e)}+21d^2e^{3i(fx+e)}-48c^2e^{2i(fx+e)}-48cde^{2i(fx+e)}+12c^2e^{i(fx+e)}+48cde^{i(fx+e)}+21d^2e^{i(fx+e)}-48c^2-48cd+12c^2+48cd+21d^2)}{12f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output -(2*a^2*c*d+2*a^2*d^2)/f*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)+(2*a^2*c^2+2*a^2*c*d)/f*tan(f*x+e)+(a^2*c^2+4*a^2*c*d+a^2*d^2)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+1/f*ln(sec(f*x+e)+tan(f*x+e))*a^2*c^2+a^2*d^2/f*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln(sec(f*x+e)+tan(f*x+e)))
```

3.195.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.19

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$$

$$= \frac{3(12a^2c^2 + 16a^2cd + 7a^2d^2) \cos(fx + e)^4 \log(\sin(fx + e) + 1) - 3(12a^2c^2 + 16a^2cd + 7a^2d^2) \cos(fx + e)^4}{1}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^2,x, algorithm="fracas")
```

output $1/48*(3*(12*a^2*c^2 + 16*a^2*c*d + 7*a^2*d^2)*\cos(f*x + e)^4*\log(\sin(f*x + e) + 1) - 3*(12*a^2*c^2 + 16*a^2*c*d + 7*a^2*d^2)*\cos(f*x + e)^4*\log(-\sin(f*x + e) + 1) + 2*(6*a^2*d^2 + 16*(3*a^2*c^2 + 5*a^2*c*d + 2*a^2*d^2)*\cos(f*x + e)^3 + 3*(4*a^2*c^2 + 16*a^2*c*d + 7*a^2*d^2)*\cos(f*x + e)^2 + 16*(a^2*c*d + a^2*d^2)*\cos(f*x + e))*\sin(f*x + e))/(f*\cos(f*x + e)^4)$

3.195.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx \\ &= a^2 \left(\int c^2 \sec(e + fx) dx + \int 2c^2 \sec^2(e + fx) dx + \int c^2 \sec^3(e + fx) dx \right. \\ & \quad + \int d^2 \sec^3(e + fx) dx + \int 2d^2 \sec^4(e + fx) dx + \int d^2 \sec^5(e + fx) dx \\ & \quad \left. + \int 2cd \sec^2(e + fx) dx + \int 4cd \sec^3(e + fx) dx + \int 2cd \sec^4(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c+d*sec(f*x+e))**2,x)`

output `a**2*(Integral(c**2*sec(e + f*x), x) + Integral(2*c**2*sec(e + f*x)**2, x) + Integral(c**2*sec(e + f*x)**3, x) + Integral(d**2*sec(e + f*x)**3, x) + Integral(2*d**2*sec(e + f*x)**4, x) + Integral(d**2*sec(e + f*x)**5, x) + Integral(2*c*d*sec(e + f*x)**2, x) + Integral(4*c*d*sec(e + f*x)**3, x) + Integral(2*c*d*sec(e + f*x)**4, x))`

3.195.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 324, normalized size of antiderivative = 1.84

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx \\ &= \frac{32 (\tan(fx + e)^3 + 3 \tan(fx + e)) a^2 cd + 32 (\tan(fx + e)^3 + 3 \tan(fx + e)) a^2 d^2 - 3 a^2 d^2 \left(\frac{2 (3 \sin(fx + e))}{\sin(fx + e)^4 - 2} \right)}{1} \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

3.195. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$

output $\frac{1}{48}(32(\tan(fx + e))^3 + 3\tan(fx + e))a^2cd + 32(\tan(fx + e))^3 + 3\tan(fx + e)a^2d^2 - 3a^2d^2(2(3\sin(fx + e))^3 - 5\sin(fx + e)) / (\sin(fx + e)^4 - 2\sin(fx + e)^2 + 1) - 3\log(\sin(fx + e) + 1) + 3\log(\sin(fx + e) - 1) - 12a^2c^2(2\sin(fx + e) / (\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1)) - 48a^2cd(2\sin(fx + e) / (\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1)) - 12a^2d^2(2\sin(fx + e) / (\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1)) + 48a^2c^2\log(\sec(fx + e) + \tan(fx + e)) + 96a^2c^2\tan(fx + e) + 96a^2cd\tan(fx + e))/f$

3.195.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.82

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$$

$$= \frac{3(12a^2c^2 + 16a^2cd + 7a^2d^2) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3(12a^2c^2 + 16a^2cd + 7a^2d^2) \log\left(\left|\tan\left(\frac{1}{2}fx\right)\right.\right)}{}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output $\frac{1}{24}(3(12a^2c^2 + 16a^2cd + 7a^2d^2)\log(\tan(1/2fx + 1/2e) + 1) - 3(12a^2c^2 + 16a^2cd + 7a^2d^2)\log(\tan(1/2fx + 1/2e) - 1) - 2(36a^2c^2\tan(1/2fx + 1/2e)^7 + 48a^2cd\tan(1/2fx + 1/2e)^7 + 21a^2d^2\tan(1/2fx + 1/2e)^7 - 132a^2c^2\tan(1/2fx + 1/2e)^5 - 176a^2cd\tan(1/2fx + 1/2e)^5 - 77a^2d^2\tan(1/2fx + 1/2e)^5 + 156a^2c^2\tan(1/2fx + 1/2e)^3 + 272a^2cd\tan(1/2fx + 1/2e)^3 + 83a^2d^2\tan(1/2fx + 1/2e)^3 - 60a^2c^2\tan(1/2fx + 1/2e) - 144a^2cd\tan(1/2fx + 1/2e) - 75a^2d^2\tan(1/2fx + 1/2e)) / (\tan(1/2fx + 1/2e)^2 - 1)^4)/f$

3.195.9 Mupad [B] (verification not implemented)

Time = 17.20 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.35

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 dx$$

$$= \frac{\left(-3a^2c^2 - 4a^2cd - \frac{7a^2d^2}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7 + \left(11a^2c^2 + \frac{44a^2cd}{3} + \frac{77a^2d^2}{12}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-13a^2c^2 - \frac{11a^2cd}{3} - \frac{7a^2d^2}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + \left(3a^2c^2 + \frac{4a^2cd}{3} + \frac{a^2d^2}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right) + \frac{a^2 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)(12c^2 + 16cd + 7d^2)}{2(6c^2 + 8cd + \frac{7d^2}{2})}\right)(12c^2 + 16cd + 7d^2)}{4f}$$

input `int(((a + a/cos(e + f*x))^2*(c + d/cos(e + f*x))^2)/cos(e + f*x),x)`

```
output (tan(e/2 + (f*x)/2)*(5*a^2*c^2 + (25*a^2*d^2)/4 + 12*a^2*c*d) - tan(e/2 +
(f*x)/2)^7*(3*a^2*c^2 + (7*a^2*d^2)/4 + 4*a^2*c*d) + tan(e/2 + (f*x)/2)^5*
(11*a^2*c^2 + (77*a^2*d^2)/12 + (44*a^2*c*d)/3) - tan(e/2 + (f*x)/2)^3*(13
*a^2*c^2 + (83*a^2*d^2)/12 + (68*a^2*c*d)/3))/(f*(6*tan(e/2 + (f*x)/2)^4 -
4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/2)^8 +
1)) + (a^2*atanh((tan(e/2 + (f*x)/2)*(16*c*d + 12*c^2 + 7*d^2))/(2*(8*c*d
+ 6*c^2 + (7*d^2)/2)))*(16*c*d + 12*c^2 + 7*d^2))/(4*f)
```

3.196 $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$

3.196.1 Optimal result	1358
3.196.2 Mathematica [A] (verified)	1358
3.196.3 Rubi [A] (verified)	1359
3.196.4 Maple [A] (verified)	1362
3.196.5 Fricas [A] (verification not implemented)	1362
3.196.6 Sympy [F]	1363
3.196.7 Maxima [A] (verification not implemented)	1363
3.196.8 Giac [A] (verification not implemented)	1364
3.196.9 Mupad [B] (verification not implemented)	1364

3.196.1 Optimal result

Integrand size = 29, antiderivative size = 103

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$$

$$= \frac{a^2(3c + 2d)\operatorname{arctanh}(\sin(e + fx))}{2f} + \frac{2a^2(3c + 2d)\tan(e + fx)}{3f}$$

$$+ \frac{a^2(3c + 2d)\sec(e + fx)\tan(e + fx)}{6f} + \frac{d(a + a \sec(e + fx))^2\tan(e + fx)}{3f}$$

```
output 1/2*a^2*(3*c+2*d)*arctanh(sin(f*x+e))/f+2/3*a^2*(3*c+2*d)*tan(f*x+e)/f+1/6
*a^2*(3*c+2*d)*sec(f*x+e)*tan(f*x+e)/f+1/3*d*(a+a*sec(f*x+e))^2*tan(f*x+e)
/f
```

3.196.2 Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.61

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$$

$$= \frac{a^2((9c + 6d)\operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx)(12(c + d) + 3(c + 2d)\sec(e + fx) + 2d\tan^2(e + fx)))}{6f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x]),x]`

output $(a^2*((9*c + 6*d)*\text{ArcTanh}[\text{Sin}[e + f*x]] + \text{Tan}[e + f*x]*(12*(c + d) + 3*(c + 2*d)*\text{Sec}[e + f*x] + 2*d*\text{Tan}[e + f*x]^2)))/(6*f)$

3.196.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.89, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4489, 3042, 4275, 3042, 4254, 24, 4534, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a \sec(e + fx) + a)^2(c + d \sec(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4489} \\
 & \frac{1}{3}(3c + 2d) \int \sec(e + fx)(\sec(e + fx)a + a)^2 dx + \frac{d \tan(e + fx)(a \sec(e + fx) + a)^2}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3}(3c + 2d) \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)a + a\right)^2 dx + \frac{d \tan(e + fx)(a \sec(e + fx) + a)^2}{3f} \\
 & \quad \downarrow \text{4275} \\
 & \frac{1}{3}(3c + 2d) \left(2a^2 \int \sec^2(e + fx) dx + \int \sec(e + fx) (\sec^2(e + fx)a^2 + a^2) dx\right) + \\
 & \quad \frac{d \tan(e + fx)(a \sec(e + fx) + a)^2}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3}(3c + 2d) \left(2a^2 \int \csc\left(e + fx + \frac{\pi}{2}\right)^2 dx + \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(\csc\left(e + fx + \frac{\pi}{2}\right)^2 a^2 + a^2\right) dx\right) + \\
 & \quad \frac{d \tan(e + fx)(a \sec(e + fx) + a)^2}{3f}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 4254 \\
& \frac{1}{3}(3c+2d) \left(\int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)^2 a^2 + a^2 \right) dx - \frac{2a^2 \int 1d(-\tan(e+fx))}{f} \right) + \\
& \quad \frac{d \tan(e+fx)(a \sec(e+fx) + a)^2}{3f} \\
& \downarrow 24 \\
& \frac{1}{3}(3c+2d) \left(\int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)^2 a^2 + a^2 \right) dx + \frac{2a^2 \tan(e+fx)}{f} \right) + \\
& \quad \frac{d \tan(e+fx)(a \sec(e+fx) + a)^2}{3f} \\
& \downarrow 4534 \\
& \frac{1}{3}(3c+2d) \left(\frac{3}{2}a^2 \int \sec(e+fx) dx + \frac{2a^2 \tan(e+fx)}{f} + \frac{a^2 \tan(e+fx) \sec(e+fx)}{2f} \right) + \\
& \quad \frac{d \tan(e+fx)(a \sec(e+fx) + a)^2}{3f} \\
& \downarrow 3042 \\
& \frac{1}{3}(3c+2d) \left(\frac{3}{2}a^2 \int \csc\left(e+fx+\frac{\pi}{2}\right) dx + \frac{2a^2 \tan(e+fx)}{f} + \frac{a^2 \tan(e+fx) \sec(e+fx)}{2f} \right) + \\
& \quad \frac{d \tan(e+fx)(a \sec(e+fx) + a)^2}{3f} \\
& \downarrow 4257 \\
& \frac{1}{3}(3c+2d) \left(\frac{3a^2 \operatorname{arctanh}(\sin(e+fx))}{2f} + \frac{2a^2 \tan(e+fx)}{f} + \frac{a^2 \tan(e+fx) \sec(e+fx)}{2f} \right) + \\
& \quad \frac{d \tan(e+fx)(a \sec(e+fx) + a)^2}{3f}
\end{aligned}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x]),x]`

output `(d*(a + a*Sec[e + f*x])^2*Tan[e + f*x])/(3*f) + ((3*c + 2*d)*((3*a^2*ArcTanh[Sin[e + f*x]])/(2*f) + (2*a^2*Tan[e + f*x])/f + (a^2*Sec[e + f*x]*Tan[e + f*x])/(2*f)))/3`

3.196.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4275 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(2), x_Symbol] := Simp[2*a*(b/d) Int[(d*Csc[e + f*x])^(n + 1), x], x] + Int[(d*Csc[e + f*x])^n*(a^2 + b^2*Csc[e + f*x]^2), x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4489 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*B*m + A*b*(m + 1))/(b*(m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x], x] /; FreeQ[{a, b, A, B, e, f, m}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && NeQ[a*B*m + A*b*(m + 1), 0] && !LtQ[m, -2^(-1)]`
- rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.196.4 Maple [A] (verified)

Time = 2.93 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.17

method	result
parts	$\frac{(a^2c+2a^2d)\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)}{f} + \frac{(2a^2c+a^2d)\tan(fx+e)}{f} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{f}$
derivativedivides	$\frac{a^2c\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) - a^2d\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + 2a^2c\tan(fx+e) + 2a^2d\left(\frac{\sec(fx+e)}{2}\right)}{f}$
default	$\frac{a^2c\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right) - a^2d\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + 2a^2c\tan(fx+e) + 2a^2d\left(\frac{\sec(fx+e)}{2}\right)}{f}$
parallelrisch	$\frac{\left(-\frac{9\left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3}\right)\left(c + \frac{2d}{3}\right)\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2} + \frac{9\left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3}\right)\left(c + \frac{2d}{3}\right)\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2}\right) + (c+2d)}{f(\cos(3fx+3e)+3\cos(fx+e))}$
norman	$\frac{\frac{8a^2(3c+2d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3f} - \frac{a^2(3c+2d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f} - \frac{a^2(5c+6d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^3} - \frac{a^2(3c+2d)\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2f} + \frac{a^2(3c+2d)\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2f}$
risch	$-\frac{ia^2(3ce^{5i(fx+e)} + 6de^{5i(fx+e)} - 12ce^{4i(fx+e)} - 6de^{4i(fx+e)} - 24e^{2i(fx+e)}c - 24de^{2i(fx+e)} - 3e^{i(fx+e)}c - 6de^{i(fx+e)}d)}{3f(1+e^{2i(fx+e)})^3}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output (a^2*c+2*a^2*d)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+(2*a^2*c+a^2*d)/f*tan(f*x+e)+1/f*ln(sec(f*x+e)+tan(f*x+e))*a^2*c-a^2*d/f*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)
```

3.196.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.34

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$$

$$= \frac{3(3a^2c + 2a^2d) \cos(fx + e)^3 \log(\sin(fx + e) + 1) - 3(3a^2c + 2a^2d) \cos(fx + e)^3 \log(-\sin(fx + e) + 1)}{12f \cos(fx + e)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e)),x, algorithm="fracas")
```

output $1/12*(3*(3*a^2*c + 2*a^2*d)*\cos(f*x + e)^3*\log(\sin(f*x + e) + 1) - 3*(3*a^2*c + 2*a^2*d)*\cos(f*x + e)^3*\log(-\sin(f*x + e) + 1) + 2*(2*a^2*d + 2*(6*a^2*c + 5*a^2*d)*\cos(f*x + e)^2 + 3*(a^2*c + 2*a^2*d)*\cos(f*x + e))*\sin(f*x + e))/(f*\cos(f*x + e)^3)$

3.196.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx \\ &= a^2 \left(\int c \sec(e + fx) dx + \int 2c \sec^2(e + fx) dx + \int c \sec^3(e + fx) dx \right. \\ & \quad \left. + \int d \sec^2(e + fx) dx + \int 2d \sec^3(e + fx) dx + \int d \sec^4(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2*(c+d*sec(f*x+e)),x)`

output `a**2*(Integral(c*sec(e + f*x), x) + Integral(2*c*sec(e + f*x)**2, x) + Integral(c*sec(e + f*x)**3, x) + Integral(d*sec(e + f*x)**2, x) + Integral(2*d*sec(e + f*x)**3, x) + Integral(d*sec(e + f*x)**4, x))`

3.196.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.62

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx \\ &= \frac{4(\tan(fx + e)^3 + 3 \tan(fx + e))a^2d - 3a^2c \left(\frac{2 \sin(fx + e)}{\sin(fx + e)^2 - 1} - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1) \right)}{f} \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e)),x, algorithm="maxima")`

output $1/12*(4*(\tan(f*x + e)^3 + 3*\tan(f*x + e))*a^2*d - 3*a^2*c*(2*\sin(f*x + e)/(\sin(f*x + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) - 6*a^2*d*(2*\sin(f*x + e)/(\sin(f*x + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) + 12*a^2*c*\log(\sec(f*x + e) + \tan(f*x + e)) + 24*a^2*c*\tan(f*x + e) + 12*a^2*d*\tan(f*x + e))/f$

3.196.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.73

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$$

$$= \frac{3(3a^2c + 2a^2d) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3(3a^2c + 2a^2d) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right) - \frac{2(9a^2c \tan(\frac{1}{2}fx + \frac{1}{2}e) + 6a^2d)}{6f}}{6f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e)),x, algorithm="giac")`

output `1/6*(3*(3*a^2*c + 2*a^2*d)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 3*(3*a^2*c + 2*a^2*d)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(9*a^2*c*tan(1/2*f*x + 1/2*e)^5 + 6*a^2*d*tan(1/2*f*x + 1/2*e)^5 - 24*a^2*c*tan(1/2*f*x + 1/2*e)^3 - 16*a^2*d*tan(1/2*f*x + 1/2*e)^3 + 15*a^2*c*tan(1/2*f*x + 1/2*e) + 18*a^2*d*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^3)/f`

3.196.9 Mupad [B] (verification not implemented)

Time = 16.21 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.56

$$\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$$

$$= \frac{2a^2 \operatorname{atanh}\left(\frac{4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{3c}{2} + d\right)}{6c + 4d}\right) \left(\frac{3c}{2} + d\right)}{f} - \frac{(3a^2c + 2a^2d) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-8a^2c - \frac{16a^2d}{3}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + (5a^2c + 6a^2d) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)}$$

input `int(((a + a/cos(e + f*x))^2*(c + d/cos(e + f*x)))/cos(e + f*x),x)`

output `(2*a^2*atanh((4*tan(e/2 + (f*x)/2)*((3*c)/2 + d))/(6*c + 4*d))*((3*c)/2 + d)/f - (tan(e/2 + (f*x)/2)*(5*a^2*c + 6*a^2*d) + tan(e/2 + (f*x)/2)^5*(3*a^2*c + 2*a^2*d) - tan(e/2 + (f*x)/2)^3*(8*a^2*c + (16*a^2*d)/3))/(f*(3*tan(e/2 + (f*x)/2)^2 - 3*tan(e/2 + (f*x)/2)^4 + tan(e/2 + (f*x)/2)^6 - 1))`

3.196. $\int \sec(e + fx)(a + a \sec(e + fx))^2(c + d \sec(e + fx)) dx$

3.197 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c+d \sec(e+fx)} dx$

3.197.1 Optimal result 1365
 3.197.2 Mathematica [C] (verified) 1365
 3.197.3 Rubi [B] (verified) 1366
 3.197.4 Maple [A] (verified) 1370
 3.197.5 Fricas [A] (verification not implemented) 1370
 3.197.6 Sympy [F] 1371
 3.197.7 Maxima [F(-2)] 1371
 3.197.8 Giac [B] (verification not implemented) 1372
 3.197.9 Mupad [B] (verification not implemented) 1373

3.197.1 Optimal result

Integrand size = 31, antiderivative size = 95

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c+d \sec(e+fx)} dx = -\frac{a^2(c-2d)\operatorname{arctanh}(\sin(e+fx))}{d^2 f} + \frac{2a^2(c-d)^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{d^2\sqrt{c+df}} + \frac{a^2 \tan(e+fx)}{df}$$

```
output -a^2*(c-2*d)*arctanh(sin(f*x+e))/d^2/f+2*a^2*(c-d)^(3/2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/d^2/f/(c+d)^(1/2)+a^2*tan(f*x+e)/d/f
```

3.197.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.85 (sec) , antiderivative size = 329, normalized size of antiderivative = 3.46

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{c+d \sec(e+fx)} dx = \frac{a^2 \cos(e+fx)(d+c \cos(e+fx)) \sec^4\left(\frac{1}{2}(e+fx)\right) (1+\sec(e+fx))^2 \left((c-2d) \log\left(\cos\left(\frac{1}{2}(e+fx)\right)\right) - \operatorname{si}\left(\frac{1}{2}(e+fx)\right) \right)}{d^2 f (c+d)}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x]),x]`

output `(a^2*Cos[e + f*x]*(d + c*Cos[e + f*x])*Sec[(e + f*x)/2]^4*(1 + Sec[e + f*x])^2*((c - 2*d)*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] - (c - 2*d)*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] - ((2*I)*(c - d)^2*ArcTan[((I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]))*(Cos[e] - I*Sin[e]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + (d*Sin[(f*x)/2])/((Cos[e/2] - Sin[e/2])*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])) + (d*Sin[(f*x)/2])/((Cos[e/2] + Sin[e/2])*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2]))) / (4*d^2*f*(c + d*Sec[e + f*x]))`

3.197.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 201 vs. $2(95) = 190$.

Time = 0.41 (sec) , antiderivative size = 201, normalized size of antiderivative = 2.12, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4475, 113, 25, 27, 175, 45, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{c+d\sec(e+fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{c+d\csc(e+fx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4475} \\
 & -\frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \\
 & \quad \downarrow \text{113} \\
 & -\frac{a^2 \tan(e+fx) \left(-\frac{\int -\frac{a^3(d-(c-2d)\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}(c+d\sec(e+fx))} d\sec(e+fx)}{ad} - \frac{\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{d} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

3.197. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c+d\sec(e+fx)} dx$

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^3(d-(c-2d)\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx) - \frac{\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{d} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e + fx) \left(\frac{a^2 \int \frac{d-(c-2d)\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx) - \frac{\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{d} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 175

$$\frac{a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{(c-d)^2 \int \frac{1}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx) - \frac{(c-2d) \int \frac{1}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d\sec(e+fx)}{d} \right)}{d} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 45

$$\frac{a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{(c-d)^2 \int \frac{1}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx) - \frac{2(c-2d) \int \frac{1}{\frac{(a-a\sec(e+fx))a}{\sec(e+fx)a+a} - a} d\frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}} d\sec(e+fx)}{d} \right)}{d} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 104

$$\frac{a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{2(c-d)^2 \int \frac{1}{\frac{a(c+d)(\sec(e+fx)a+a)}{a-a\sec(e+fx)}} d\frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a\sec(e+fx)}}} d\sec(e+fx) - \frac{2(c-2d) \int \frac{1}{\frac{(a-a\sec(e+fx))a}{\sec(e+fx)a+a} - a} d\frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}} d\sec(e+fx)}{d} \right)}{d} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 218

3.197. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c+d\sec(e+fx)} dx$

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{2(c-d)^{3/2} \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}\right)}{ad\sqrt{c+d}} + \frac{2(c-2d) \arctan\left(\frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}}\right)}{ad} \right)}{d} - \frac{\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{d} \right) - \frac{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}{d}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x]),x]`

output `-((a^2*((a^2*((2*(c - 2*d)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])])/(a*d) + (2*(c - d)^(3/2)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]/(a*d*Sqrt[c + d])))/d - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]/d)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])`

3.197.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 104 `Int[(((a_) + (b_)*(x_))^(m_))*((c_) + (d_)*(x_))^(n_)/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 113 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 175 `Int[(((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_))) / ((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.197.4 Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.58

method	result
derivativedivides	$8a^2 \left(-\frac{(-c^2+2cd-d^2) \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}\right)}{4d^2\sqrt{(c+d)(c-d)}} - \frac{1}{8d\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)} + \frac{(-c+2d)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{8d^2} - \frac{1}{8d\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)} \right) f$
default	$8a^2 \left(-\frac{(-c^2+2cd-d^2) \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}\right)}{4d^2\sqrt{(c+d)(c-d)}} - \frac{1}{8d\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)} + \frac{(-c+2d)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{8d^2} - \frac{1}{8d\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)} \right) f$
risch	$\frac{2ia^2}{fd(1+e^{2i(fx+e)})} + \frac{\sqrt{(c+d)(c-d)} a^2 \ln\left(e^{i(fx+e)} + \frac{i\sqrt{(c+d)(c-d)+d}}{c}\right)}{(c+d)fd^2} - \frac{\sqrt{(c+d)(c-d)} a^2 \ln\left(e^{i(fx+e)} + \frac{i\sqrt{(c+d)(c-d)-d}}{c}\right)}{(c+d)fd}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 8/f*a^2*(-1/4/d^2*(-c^2+2*c*d-d^2)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))-1/8/d/(tan(1/2*f*x+1/2*e)+1)+1/8/d^2*(-c+2*d)*ln(tan(1/2*f*x+1/2*e)+1)-1/8/d/(tan(1/2*f*x+1/2*e)-1)+1/8*(c-2*d)/d^2*ln(tan(1/2*f*x+1/2*e)-1))
```

3.197.5 Fracas [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 398, normalized size of antiderivative = 4.19

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c+d\sec(e+fx)} dx$$

$$= \left[\frac{2a^2d\sin(fx+e) - (a^2c - a^2d)\sqrt{\frac{c-d}{c+d}} \cos(fx+e) \log\left(\frac{2cd\cos(fx+e) - (c^2-2d^2)\cos(fx+e)^2 - 2(c^2+cd+(cd+d^2)\cos(fx+e))}{c^2\cos(fx+e)^2 + 2cd\cos(fx+e) + c^2}\right)}{\dots} \right]$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="fracas")
```

output `[1/2*(2*a^2*d*sin(f*x + e) - (a^2*c - a^2*d)*sqrt((c - d)/(c + d))*cos(f*x + e)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*(c^2 + c*d + (c*d + d^2)*cos(f*x + e))*sqrt((c - d)/(c + d))*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) - (a^2*c - 2*a^2*d)*cos(f*x + e)*log(sin(f*x + e) + 1) + (a^2*c - 2*a^2*d)*cos(f*x + e)*log(-sin(f*x + e) + 1))/(d^2*f*cos(f*x + e)), 1/2*(2*a^2*d*sin(f*x + e) + 2*(a^2*c - a^2*d)*sqrt(-(c - d)/(c + d))*arctan(-(d*cos(f*x + e) + c)*sqrt(-(c - d)/(c + d)))/((c - d)*sin(f*x + e)))*cos(f*x + e) - (a^2*c - 2*a^2*d)*cos(f*x + e)*log(sin(f*x + e) + 1) + (a^2*c - 2*a^2*d)*cos(f*x + e)*log(-sin(f*x + e) + 1))/(d^2*f*cos(f*x + e))]`

3.197.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{c + d \sec(e + fx)} dx = a^2 \left(\int \frac{\sec(e + fx)}{c + d \sec(e + fx)} dx + \int \frac{2 \sec^2(e + fx)}{c + d \sec(e + fx)} dx + \int \frac{\sec^3(e + fx)}{c + d \sec(e + fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e)),x)`

output `a**2*(Integral(sec(e + f*x)/(c + d*sec(e + f*x)), x) + Integral(2*sec(e + f*x)**2/(c + d*sec(e + f*x)), x) + Integral(sec(e + f*x)**3/(c + d*sec(e + f*x)), x))`

3.197.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{c + d \sec(e + fx)} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de

3.197.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. $2(86) = 172$.

Time = 0.35 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.06

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c+d\sec(e+fx)} dx =$$

$$\frac{2a^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)d} + \frac{(a^2c - 2a^2d) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{d^2} - \frac{(a^2c - 2a^2d) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{d^2} + \frac{2(a^2c^2 - 2a^2cd + a^2d^2)}{f} \left(\pi \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="giac")`

output `-(2*a^2*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1/2*e)^2 - 1)*d) + (a^2*c - 2*a^2*d)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/d^2 - (a^2*c - 2*a^2*d)*log(abs(tan(1/2*f*x + 1/2*e) - 1))/d^2 + 2*(a^2*c^2 - 2*a^2*c*d + a^2*d^2)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/sqrt(-c^2 + d^2))/f`

3.197.9 Mupad [B] (verification not implemented)

Time = 14.26 (sec) , antiderivative size = 529, normalized size of antiderivative = 5.57

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{c+d\sec(e+fx)} dx$$

$$= \frac{2a^2 \left(\frac{\sin(e+fx)}{2} + 2\cos(e+fx) \operatorname{atanh} \left(\frac{\sin\left(\frac{e}{2} + \frac{fx}{2}\right)}{\cos\left(\frac{e}{2} + \frac{fx}{2}\right)} \right) \right)}{f \cos(e+fx)(c+d)}$$

$$+ \frac{2a^2 \left(\frac{c \sin(e+fx)}{2} + c \cos(e+fx) \operatorname{atanh} \left(\frac{\sin\left(\frac{e}{2} + \frac{fx}{2}\right)}{\cos\left(\frac{e}{2} + \frac{fx}{2}\right)} \right) \right)}{df \cos(e+fx)(c+d)}$$

$$- \frac{2a^2 \left(c^2 \cos(e+fx) \operatorname{atanh} \left(\frac{\sin\left(\frac{e}{2} + \frac{fx}{2}\right)}{\cos\left(\frac{e}{2} + \frac{fx}{2}\right)} \right) + \cos(e+fx) \operatorname{atan} \left(\frac{(2c \sin\left(\frac{e}{2} + \frac{fx}{2}\right) (c^4 - 2c^3d + 2cd^3 - d^4))^{3/2} - 2c^5 \sin\left(\frac{e}{2} + \frac{fx}{2}\right)}{\dots} \right) \right)}{\dots}$$

```
input int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c + d/cos(e + f*x))),x)
```

```
output (2*a^2*(sin(e + f*x)/2 + 2*cos(e + f*x)*atanh(sin(e/2 + (f*x)/2)/cos(e/2 + (f*x)/2)))/(f*cos(e + f*x)*(c + d)) + (2*a^2*((c*sin(e + f*x))/2 + c*cos(e + f*x)*atanh(sin(e/2 + (f*x)/2)/cos(e/2 + (f*x)/2)))/(d*f*cos(e + f*x)*(c + d)) - (2*a^2*(c^2*cos(e + f*x)*atanh(sin(e/2 + (f*x)/2)/cos(e/2 + (f*x)/2)) + cos(e + f*x)*atan(((2*c*sin(e/2 + (f*x)/2)*(2*c*d^3 - 2*c^3*d + c^4 - d^4))^(3/2) - 2*c^5*sin(e/2 + (f*x)/2)*(2*c*d^3 - 2*c^3*d + c^4 - d^4))^(1/2) + 5*d^5*sin(e/2 + (f*x)/2)*(2*c*d^3 - 2*c^3*d + c^4 - d^4))^(1/2) - c*d^4*sin(e/2 + (f*x)/2)*(2*c*d^3 - 2*c^3*d + c^4 - d^4))^(1/2) + 4*c^4*d*sin(e/2 + (f*x)/2)*(2*c*d^3 - 2*c^3*d + c^4 - d^4))^(1/2) - 9*c^2*d^3*sin(e/2 + (f*x)/2)*(2*c*d^3 - 2*c^3*d + c^4 - d^4))^(1/2) + 3*c^3*d^2*sin(e/2 + (f*x)/2)*(2*c*d^3 - 2*c^3*d + c^4 - d^4))^(1/2))*1i)/(d*cos(e/2 + (f*x)/2)*(c + d)*(8*c*d^4 + 3*c^4*d - 5*d^5 + 2*c^2*d^3 - 8*c^3*d^2))*((c + d)*(c - d)^3)^(1/2)*1i))/(d^2*f*cos(e + f*x)*(c + d))
```

3.198
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^2} dx$$

3.198.1 Optimal result 1374
 3.198.2 Mathematica [C] (verified) 1374
 3.198.3 Rubi [A] (verified) 1375
 3.198.4 Maple [A] (verified) 1379
 3.198.5 Fricas [B] (verification not implemented) 1380
 3.198.6 Sympy [F] 1381
 3.198.7 Maxima [F(-2)] 1381
 3.198.8 Giac [B] (verification not implemented) 1382
 3.198.9 Mupad [B] (verification not implemented) 1382

3.198.1 Optimal result

Integrand size = 31, antiderivative size = 117

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^2} dx = \frac{a^2 \operatorname{arctanh}(\sin(e + fx))}{d^2 f} - \frac{2a^2 \sqrt{c - d}(c + 2d) \operatorname{arctanh}\left(\frac{\sqrt{c - d} \tan\left(\frac{1}{2}(e + fx)\right)}{\sqrt{c + d}}\right)}{d^2 (c + d)^{3/2} f} - \frac{a^2 (c - d) \tan(e + fx)}{d(c + d)f(c + d \sec(e + fx))}$$

output

```
a^2*arctanh(sin(f*x+e))/d^2/f-2*a^2*(c+2*d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))*(c-d)^(1/2)/d^2/(c+d)^(3/2)/f-a^2*(c-d)*tan(f*x+e)/d/(c+d)/f/(c+d*sec(f*x+e))
```

3.198.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.44 (sec) , antiderivative size = 312, normalized size of antiderivative = 2.67

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^2} dx = \frac{a^2(d + c \cos(e + fx)) \sec^4\left(\frac{1}{2}(e + fx)\right) (1 + \sec(e + fx))^2 \left(-((d + c \cos(e + fx)) \log(\cos\left(\frac{1}{2}(e + fx)\right)) - \right)}{\dots}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^2,x]`

output `(a^2*(d + c*Cos[e + f*x])*Sec[(e + f*x)/2]^4*(1 + Sec[e + f*x])^2*(-((d + c*Cos[e + f*x])*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]]) + (d + c*Cos[e + f*x])*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] + (2*(c^2 + c*d - 2*d^2)*ArcTan[((I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))]/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]))*(d + c*Cos[e + f*x])*(I*Cos[e] + Sin[e]))/((c + d)*Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + ((c - d)*d*(d*Sin[e] - c*Sin[f*x]))/(c*(c + d)*(Cos[e/2] - Sin[e/2])*(Cos[e/2] + Sin[e/2])))/(4*d^2*f*(c + d*Sec[e + f*x])^2)`

3.198.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.97, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4475, 109, 25, 27, 175, 45, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a \sec(e+fx) + a)^2}{(c + d \sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx + \frac{\pi}{2})(a \csc(e+fx + \frac{\pi}{2}) + a)^2}{(c + d \csc(e+fx + \frac{\pi}{2}))^2} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx) + a}}$$

↓ 109

$$\frac{a^2 \tan(e+fx) \left(\frac{(c-d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx) + a}}{d(c+d)(c+d \sec(e+fx))} - \frac{\int -\frac{a^3(2d+(c+d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{ad(c+d)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx) + a}}$$

↓ 25

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^3(2d+(c+d)\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx)}{ad(c+d)} + \frac{(c-d)\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{d(c+d)(c+d\sec(e+fx))} \right)$$

$$f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{a^2 \int \frac{2d+(c+d)\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx)}{d(c+d)} + \frac{(c-d)\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}{d(c+d)(c+d\sec(e+fx))} \right)$$

$$f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}$$

↓ 175

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{(c+d) \int \frac{1}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d\sec(e+fx)}{d} - \frac{(c-d)(c+2d) \int \frac{1}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx)}{d} \right)}{d(c+d)} \right)$$

$$f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{2(c+d) \int \frac{1}{\frac{(a-a\sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}}{d} - \frac{(c-d)(c+2d) \int \frac{1}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a(c+d\sec(e+fx))}} d\sec(e+fx)}{d} \right)}{d(c+d)} \right)$$

$$f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}$$

↓ 104

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{2(c+d) \int \frac{1}{\frac{(a-a\sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a\sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}}{d} - \frac{2(c-d)(c+2d) \int \frac{1}{\frac{a(c-d)+a(c+d)(\sec(e+fx)a+a)}{a-a\sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a\sec(e+fx)}}}{d} \right)}{d(c+d)} \right) +$$

$$f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}$$

↓ 218

3.198. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^2} dx$

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(-\frac{2(c+d) \arctan\left(\frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}}\right)}{ad} - \frac{2\sqrt{c-d}(c+2d) \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}\right)}{ad\sqrt{c+d}} \right)}{d(c+d)} \right) + \frac{(c-d)\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)}}{d(c+d)(c+d \sec(e+fx))}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^2,x]`

output `-((a^2*((a^2*(-2*(c + d)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])])/(a*d) - (2*Sqrt[c - d]*(c + 2*d)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(a*d*Sqrt[c + d])))/(d*(c + d)) + ((c - d)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(d*(c + d)*(c + d*Sec[e + f*x]))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]))`

3.198.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 104 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

- rule 109 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 175 `Int[(((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_))) / ((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.198.4 Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.34

method	result
derivativedivides	$8a^2 \frac{\left((c-d) \left(\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} \right) - \frac{(c+2d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c+d) \sqrt{(c+d)(c-d)}} \right)}{4d^2} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{8d^2} - \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{8d^2}$
default	$8a^2 \frac{\left((c-d) \left(\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} \right) - \frac{(c+2d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c+d) \sqrt{(c+d)(c-d)}} \right)}{4d^2} + \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{8d^2} - \frac{\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{8d^2}$
risch	$-\frac{2ia^2(c-d)(de^{i(fx+e)}+c)}{df(c+d)c(e^{2i(fx+e)}c+2de^{i(fx+e)}+c)} + \frac{\sqrt{(c+d)(c-d)}a^2 \ln\left(e^{i(fx+e)} - \frac{i\sqrt{(c+d)(c-d)}-d}{c}\right)c}{(c+d)^2 f d^2} + \frac{2\sqrt{(c+d)(c-d)}a^2}{8d^2}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `8/f*a^2*(1/4*(c-d)/d^2*(d/(c+d)*tan(1/2*f*x+1/2*e)/(tan(1/2*f*x+1/2*e)^2*c - tan(1/2*f*x+1/2*e)^2*d-c-d)-(c+2*d)/(c+d)/((c+d)*(c-d)^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d)^(1/2))))+1/8/d^2*ln(tan(1/2*f*x+1/2*e)+1)-1/8/d^2*ln(tan(1/2*f*x+1/2*e)-1))`

3.198.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 249 vs. $2(108) = 216$.

Time = 0.42 (sec) , antiderivative size = 567, normalized size of antiderivative = 4.85

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^2} dx$$

$$= \frac{(a^2cd + 2a^2d^2 + (a^2c^2 + 2a^2cd)\cos(fx+e))\sqrt{\frac{c-d}{c+d}} \log\left(\frac{2cd\cos(fx+e) - (c^2 - 2d^2)\cos(fx+e)^2 - 2(c^2 + cd + (cd+d^2)\cos(fx+e))}{c^2\cos(fx+e)^2 + 2cd\cos(fx+e)}\right)}{2(a^2cd + 2a^2d^2 + (a^2c^2 + 2a^2cd)\cos(fx+e))\sqrt{-\frac{c-d}{c+d}} \arctan\left(-\frac{(d\cos(fx+e)+c)\sqrt{-\frac{c-d}{c+d}}}{(c-d)\sin(fx+e)}\right) - (a^2cd + a^2d^2)}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x, algorithm="fracas")`

output `[1/2*((a^2*c*d + 2*a^2*d^2 + (a^2*c^2 + 2*a^2*c*d)*cos(f*x + e))*sqrt((c - d)/(c + d))*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*(c^2 + c*d + (c*d + d^2)*cos(f*x + e))*sqrt((c - d)/(c + d))*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + (a^2*c*d + a^2*d^2 + (a^2*c^2 + a^2*c*d)*cos(f*x + e))*log(sin(f*x + e) + 1) - (a^2*c*d + a^2*d^2 + (a^2*c^2 + a^2*c*d)*cos(f*x + e))*log(-sin(f*x + e) + 1) - 2*(a^2*c*d - a^2*d^2)*sin(f*x + e))/((c^2*d^2 + c*d^3)*f*cos(f*x + e) + (c*d^3 + d^4)*f), -1/2*(2*(a^2*c*d + 2*a^2*d^2 + (a^2*c^2 + 2*a^2*c*d)*cos(f*x + e))*sqrt(-(c - d)/(c + d))*arctan(-(d*cos(f*x + e) + c)*sqrt(-(c - d)/(c + d))/((c - d)*sin(f*x + e))) - (a^2*c*d + a^2*d^2 + (a^2*c^2 + a^2*c*d)*cos(f*x + e))*log(sin(f*x + e) + 1) + (a^2*c*d + a^2*d^2 + (a^2*c^2 + a^2*c*d)*cos(f*x + e))*log(-sin(f*x + e) + 1) + 2*(a^2*c*d - a^2*d^2)*sin(f*x + e))/((c^2*d^2 + c*d^3)*f*cos(f*x + e) + (c*d^3 + d^4)*f)]`

3.198.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^2} dx = a^2 \left(\int \frac{\sec(e+fx)}{c^2+2cd\sec(e+fx)+d^2\sec^2(e+fx)} dx + \int \frac{2\sec^2(e+fx)}{c^2+2cd\sec(e+fx)+d^2\sec^2(e+fx)} dx + \int \frac{\sec^3(e+fx)}{c^2+2cd\sec(e+fx)+d^2\sec^2(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e))**2,x)`

output `a**2*(Integral(sec(e + f*x)/(c**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)**2), x) + Integral(2*sec(e + f*x)**2/(c**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)**2), x) + Integral(sec(e + f*x)**3/(c**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)**2), x))`

3.198.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.198.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 230 vs. $2(108) = 216$.

Time = 0.36 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.97

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^2} dx$$

$$= \frac{\frac{a^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e)|) + a^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{d^2} + \frac{2(a^2c^2 + a^2cd - 2a^2d^2) \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan\left(\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d}{\sqrt{-c^2+d^2}}\right) \right)}{(cd^2+d^3)\sqrt{-c^2+d^2}}}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output `(a^2*log(abs(tan(1/2*f*x + 1/2*e) + 1))/d^2 - a^2*log(abs(tan(1/2*f*x + 1/2*e) - 1))/d^2 + 2*(a^2*c^2 + a^2*c*d - 2*a^2*d^2)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((c*d^2 + d^3)*sqrt(-c^2 + d^2)) + 2*(a^2*c*tan(1/2*f*x + 1/2*e) - a^2*d*tan(1/2*f*x + 1/2*e))/((c*tan(1/2*f*x + 1/2*e))^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)*(c*d + d^2))/f`

3.198.9 Mupad [B] (verification not implemented)

Time = 16.42 (sec) , antiderivative size = 2563, normalized size of antiderivative = 21.91

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^2} dx = \text{Too large to display}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c + d/cos(e + f*x))^2),x)`

output $(a^2 \operatorname{atan}(((a^2((a^2((32(3a^2d^8 - 2a^2cd^7 - 4a^2c^2d^6 + 2a^2c^3d^5 + a^2c^4d^4))/(2cd^4 + d^5 + c^2d^3) - (32a^2 \tan(e/2 + (f*x)/2)*(2cd^8 - 4c^3d^6 + 2c^5d^4))/(d^2(2cd^3 + d^4 + c^2d^2)))))/d^2 + (32 \tan(e/2 + (f*x)/2)*(2a^4c^5 - 5a^4d^5 + 9a^4cd^4 + a^4c^2d^3 - 7a^4c^3d^2))/(2cd^3 + d^4 + c^2d^2))*1i)/d^2 - (a^2((a^2((32(3a^2d^8 - 2a^2cd^7 - 4a^2c^2d^6 + 2a^2c^3d^5 + a^2c^4d^4))/(2cd^4 + d^5 + c^2d^3) + (32a^2 \tan(e/2 + (f*x)/2)*(2cd^8 - 4c^3d^6 + 2c^5d^4))/(d^2(2cd^3 + d^4 + c^2d^2)))))/d^2 - (32 \tan(e/2 + (f*x)/2)*(2a^4c^5 - 5a^4d^5 + 9a^4cd^4 + a^4c^2d^3 - 7a^4c^3d^2))/(2cd^3 + d^4 + c^2d^2))*1i)/d^2)/((64(2a^6d^4 - a^6c^4 - 5a^6cd^3 + a^6c^3d + 3a^6c^2d^2))/(2cd^4 + d^5 + c^2d^3) + (a^2((a^2((32(3a^2d^8 - 2a^2cd^7 - 4a^2c^2d^6 + 2a^2c^3d^5 + a^2c^4d^4))/(2cd^4 + d^5 + c^2d^3) - (32a^2 \tan(e/2 + (f*x)/2)*(2cd^8 - 4c^3d^6 + 2c^5d^4))/(d^2(2cd^3 + d^4 + c^2d^2)))))/d^2 + (32 \tan(e/2 + (f*x)/2)*(2a^4c^5 - 5a^4d^5 + 9a^4cd^4 + a^4c^2d^3 - 7a^4c^3d^2))/(2cd^3 + d^4 + c^2d^2))))/d^2 + (a^2((a^2((32(3a^2d^8 - 2a^2cd^7 - 4a^2c^2d^6 + 2a^2c^3d^5 + a^2c^4d^4))/(2cd^4 + d^5 + c^2d^3) + (32a^2 \tan(e/2 + (f*x)/2)*(2cd^8 - 4c^3d^6 + 2c^5d^4))/(d^2(2cd^3 + d^4 + c^2d^2)))))/d^2 - (32 \tan(e/2 + (f*x)/2)*(2a^4c^5 - 5a^4d^5 + 9a^4cd^4 + a^4c^2d^3 - 7a^4c^3d^2))/(2cd^3 + d^4 + \dots$

3.199 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^3} dx$

3.199.1 Optimal result 1384
 3.199.2 Mathematica [C] (verified) 1384
 3.199.3 Rubi [A] (verified) 1385
 3.199.4 Maple [A] (verified) 1388
 3.199.5 Fricas [B] (verification not implemented) 1388
 3.199.6 Sympy [F] 1389
 3.199.7 Maxima [F(-2)] 1390
 3.199.8 Giac [A] (verification not implemented) 1390
 3.199.9 Mupad [B] (verification not implemented) 1391

3.199.1 Optimal result

Integrand size = 31, antiderivative size = 130

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^3} dx = \frac{3a^2 \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{\sqrt{c-d}(c+d)^{5/2} f} + \frac{(a^2 + a^2 \sec(e+fx)) \tan(e+fx)}{2(c+d)f(c+d \sec(e+fx))^2} + \frac{3a^2 \tan(e+fx)}{2(c+d)^2 f(c+d \sec(e+fx))}$$

```
output 3*a^2*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c+d)^(5/2)/f/(c-d)^(1/2)+1/2*(a^2+a^2*sec(f*x+e))*tan(f*x+e)/(c+d)/f/(c+d*sec(f*x+e))^2+3/2*a^2*tan(f*x+e)/(c+d)^2/f/(c+d*sec(f*x+e))
```

3.199.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.85 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.92

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^3} dx = \frac{a^2(d+c \cos(e+fx)) \sec^4\left(\frac{1}{2}(e+fx)\right) \sec(e+fx)(1+\sec(e+fx))^2}{8(c+d)}$$

3.199. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^3} dx$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^3,x]`

output $(a^2*(d + c*\cos[e + f*x])*Sec[(e + f*x)/2]^4*Sec[e + f*x]*(1 + Sec[e + f*x])^2*((-6*I)*ArcTan[((I*\cos[e] + \sin[e])*(c*\sin[e] + (-d + c*\cos[e]))*\tan[(f*x)/2]))/(Sqrt[c^2 - d^2]*Sqrt[(\cos[e] - I*\sin[e])^2]))*(d + c*\cos[e + f*x])^2*(\cos[e] - I*\sin[e])/(Sqrt[c^2 - d^2]*Sqrt[(\cos[e] - I*\sin[e])^2]) + ((c - d)*(c + d)*Sec[e]*(-d*\sin[e]) + c*\sin[f*x])/c^2 + ((d + c*\cos[e + f*x])*Sec[e]*((c^2 - 4*c*d - 2*d^2)*\sin[e] + c*(4*c + d)*\sin[f*x])/c^2)/(8*(c + d)^2*f*(c + d*Sec[e + f*x])^3)$

3.199.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.72, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 4475, 105, 105, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c+d\sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c+d\csc(e+fx+\frac{\pi}{2}))^3} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^3} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 105

$$\frac{a^2 \tan(e+fx) \left(\frac{3a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^2} d\sec(e+fx)}{2(c+d)} - \frac{\sqrt{a-a\sec(e+fx)}(a\sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d\sec(e+fx))^2} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 105

$$\begin{aligned}
 & a^2 \tan(e + fx) \left(\frac{3a \left(\frac{a \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))} d \sec(e+fx)}{c+d} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)}{2a(c+d)(c+d \sec(e+fx))} \right) \\
 & \hline
 & \qquad \qquad \qquad f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} \\
 & \qquad \qquad \qquad \downarrow 104 \\
 & a^2 \tan(e + fx) \left(\frac{3a \left(\frac{2a \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)}{2a(c+d)(c+d \sec(e+fx))} \right) \\
 & \hline
 & \qquad \qquad \qquad f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} \\
 & \qquad \qquad \qquad \downarrow 218 \\
 & a^2 \tan(e + fx) \left(\frac{3a \left(\frac{2 \arctan \left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a \sec(e+fx)}} \right)}{\sqrt{c-d}(c+d)^{3/2}} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d \sec(e+fx))^2} \right) \\
 & \hline
 & \qquad \qquad \qquad f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}
 \end{aligned}$$

```
input Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^3,x]
```

```
output -((a^2*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2)))/(a*(c + d)*(c + d*Sec[e + f*x])^2) + (3*a*((2*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(Sqrt[c - d]*(c + d)^(3/2)) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a*(c + d)*(c + d*Sec[e + f*x]))))/(2*(c + d))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.199.3.1 Defintions of rubi rules used

- rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 105 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((m + 1)*(b*e - a*f))), x] - Simp[n*((d*e - c*f)/((m + 1)*(b*e - a*f))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_)^(m_)*(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_)^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.199.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.28

method	result
derivativedivides	$8a^2 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)^2} + \frac{-\frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{8(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)}{c+d} + \frac{3 \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}\right)}}{8(c+d)\sqrt{(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}\right)}} \right) f$
default	$8a^2 \left(\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)^2} + \frac{-\frac{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{8(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)}{c+d} + \frac{3 \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}\right)}}{8(c+d)\sqrt{(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}\right)}} \right) f$
risch	$\frac{ia^2(-c^3 e^{3i(fx+e)} + 4c^2 d e^{3i(fx+e)} + 2c d^2 e^{3i(fx+e)} + 4c^3 e^{2i(fx+e)} + c^2 d e^{2i(fx+e)} + 8c d^2 e^{2i(fx+e)} + 2d^3 e^{2i(fx+e)} + c^3 e^{i(fx+e)})}{c^2(c+d)^2 f(e^{2i(fx+e)} c + 2d e^{i(fx+e)} + c)^2}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `8/f*a^2*(1/4*tan(1/2*f*x+1/2*e)/(c+d)/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^2+3/4/(c+d)*(-1/2*tan(1/2*f*x+1/2*e)/(c+d)/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)+1/2/(c+d)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))))`

3.199.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 282 vs. 2(117) = 234.

Time = 0.31 (sec) , antiderivative size = 622, normalized size of antiderivative = 4.78

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^3} dx$$

$$= \left[\frac{3(a^2 c^2 \cos^2(fx+e) + 2a^2 cd \cos(fx+e) + a^2 d^2) \sqrt{c^2 - d^2} \log\left(\frac{2cd \cos(fx+e) - (c^2 - 2d^2) \cos(fx+e)^2 + 2\sqrt{c^2 - d^2} d \cos(fx+e)}{c^2 \cos^2(fx+e) + 2cd \cos(fx+e) + d^2}\right)}{4((c^6 + 2c^5 d - 2c^3 d^3 - c^2 d^4) f \cos(fx+e)^2 + 2(c^6 + 2c^5 d - 2c^3 d^3 - c^2 d^4) f \cos(fx+e) + 2(c^6 + 2c^5 d - 2c^3 d^3 - c^2 d^4))} \right]$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x, algorithm="fricas")`

output `[1/4*(3*(a^2*c^2*cos(f*x + e)^2 + 2*a^2*c*d*cos(f*x + e) + a^2*d^2)*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(a^2*c^3 + 4*a^2*c^2*d - a^2*c*d^2 - 4*a^2*d^3 + (4*a^2*c^3 + a^2*c^2*d - 4*a^2*c*d^2 - a^2*d^3)*cos(f*x + e))*sin(f*x + e))/((c^6 + 2*c^5*d - 2*c^3*d^3 - c^2*d^4)*f*cos(f*x + e)^2 + 2*(c^5*d + 2*c^4*d^2 - 2*c^2*d^4 - c*d^5)*f*cos(f*x + e) + (c^4*d^2 + 2*c^3*d^3 - 2*c*d^5 - d^6)*f), 1/2*(3*(a^2*c^2*cos(f*x + e)^2 + 2*a^2*c*d*cos(f*x + e) + a^2*d^2)*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (a^2*c^3 + 4*a^2*c^2*d - a^2*c*d^2 - 4*a^2*d^3 + (4*a^2*c^3 + a^2*c^2*d - 4*a^2*c*d^2 - a^2*d^3)*cos(f*x + e))*sin(f*x + e))/((c^6 + 2*c^5*d - 2*c^3*d^3 - c^2*d^4)*f*cos(f*x + e)^2 + 2*(c^5*d + 2*c^4*d^2 - 2*c^2*d^4 - c*d^5)*f*cos(f*x + e) + (c^4*d^2 + 2*c^3*d^3 - 2*c*d^5 - d^6)*f)]`

3.199.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^3} dx \\ &= a^2 \left(\int \frac{\sec(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \right. \\ & \quad + \int \frac{2\sec^2(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \\ & \quad \left. + \int \frac{\sec^3(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e))**3,x)`

output `a**2*(Integral(sec(e + f*x)/(c**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**3), x) + Integral(2*sec(e + f*x)**2/(c**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**3), x) + Integral(sec(e + f*x)**3/(c**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**3), x))`

3.199.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de

3.199.8 Giac [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.62

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^3} dx = \frac{3 \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right) a^2}{(c^2+2cd+d^2)\sqrt{-c^2+d^2}} + \frac{3a^2c \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 3a^2d \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 5a^2c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - d \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c}{(c \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - d \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - c - d)^2 - c} f$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output $-(3*(\pi*\text{floor}(1/2*(f*x + e)/\pi + 1/2)*\text{sgn}(2*c - 2*d) + \arctan((c*\tan(1/2*f*x + 1/2*e) - d*\tan(1/2*f*x + 1/2*e))/\text{sqrt}(-c^2 + d^2)))*a^2/((c^2 + 2*c*d + d^2)*\text{sqrt}(-c^2 + d^2)) + (3*a^2*c*\tan(1/2*f*x + 1/2*e)^3 - 3*a^2*d*\tan(1/2*f*x + 1/2*e)^3 - 5*a^2*c*\tan(1/2*f*x + 1/2*e)^2 - 5*a^2*d*\tan(1/2*f*x + 1/2*e)^2 - c - d)^2*(c^2 + 2*c*d + d^2))/f$

3.199.9 Mupad [B] (verification not implemented)

Time = 15.30 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.22

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^3} dx$$

$$= \frac{\frac{5a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{c+d} - \frac{3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (a^2 c - a^2 d)}{(c+d)^2}}{f \left(2cd - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (2c^2 - 2d^2) + \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 (c^2 - 2cd + d^2) + c^2 + d^2 \right)}$$

$$+ \frac{3a^2 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \sqrt{c-d}}{\sqrt{c+d}}\right)}{f(c+d)^{5/2} \sqrt{c-d}}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c + d/cos(e + f*x))^3),x)`output `((5*a^2*tan(e/2 + (f*x)/2))/(c + d) - (3*tan(e/2 + (f*x)/2)^3*(a^2*c - a^2*d))/(c + d)^2)/(f*(2*c*d - tan(e/2 + (f*x)/2)^2*(2*c^2 - 2*d^2) + tan(e/2 + (f*x)/2)^4*(c^2 - 2*c*d + d^2) + c^2 + d^2)) + (3*a^2*atanh((tan(e/2 + (f*x)/2)*(c - d)^(1/2))/(c + d)^(1/2)))/(f*(c + d)^(5/2)*(c - d)^(1/2))`

3.200 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^4} dx$

3.200.1 Optimal result 1392
 3.200.2 Mathematica [A] (verified) 1393
 3.200.3 Rubi [A] (verified) 1393
 3.200.4 Maple [A] (verified) 1396
 3.200.5 Fricas [B] (verification not implemented) 1397
 3.200.6 Sympy [F] 1398
 3.200.7 Maxima [F(-2)] 1399
 3.200.8 Giac [B] (verification not implemented) 1399
 3.200.9 Mupad [B] (verification not implemented) 1400

3.200.1 Optimal result

Integrand size = 31, antiderivative size = 213

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^4} dx = \frac{a^2(3c-2d)\operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{(c-d)^{3/2}(c+d)^{7/2}f} - \frac{d(a+a \sec(e+fx))^2 \tan(e+fx)}{3(c^2-d^2)f(c+d \sec(e+fx))^3} + \frac{(3c-2d)(a^2+a^2 \sec(e+fx)) \tan(e+fx)}{6(c-d)(c+d)^2 f(c+d \sec(e+fx))^2} + \frac{a^2(3c-2d) \tan(e+fx)}{2(c-d)(c+d)^3 f(c+d \sec(e+fx))}$$

```
output a^2*(3*c-2*d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c-d)^(3/2)/(c+d)^(7/2)/f-1/3*d*(a+a*sec(f*x+e))^2*tan(f*x+e)/(c^2-d^2)/f/(c+d*sec(f*x+e))^3+1/6*(3*c-2*d)*(a^2+a^2*sec(f*x+e))*tan(f*x+e)/(c-d)/(c+d)^2/f/(c+d*sec(f*x+e))^2+1/2*a^2*(3*c-2*d)*tan(f*x+e)/(c-d)/(c+d)^3/f/(c+d*sec(f*x+e))
```

3.200.2 Mathematica [A] (verified)

Time = 3.98 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.99

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^4} dx$$

$$= \frac{a^2(c-d)^2 \left(24(3c-2d)\operatorname{arctanh}\left(\frac{(-c+d)\tan(\frac{1}{2}(e+fx))}{\sqrt{c^2-d^2}}\right) (d+c\cos(e+fx))^3 - 2\sqrt{c^2-d^2}(12c^3-5c^2d+6cd^2) \right)}{24(-c+d)^3(c+d)^3\sqrt{c^2-d^2}}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^4,x]`output `(a^2*(c - d)^2*(24*(3*c - 2*d)*ArcTanh[(-c + d)*Tan[(e + f*x)/2]]/Sqrt[c^2 - d^2])*(d + c*Cos[e + f*x])^3 - 2*Sqrt[c^2 - d^2]*(12*c^3 - 5*c^2*d + 6*c*d^2 - 22*d^3 + 6*(c^3 + 6*c^2*d - 7*c*d^2 - 2*d^3)*Cos[e + f*x] + (12*c^3 - 7*c^2*d - 6*c*d^2 - 2*d^3)*Cos[2*(e + f*x)])*Sin[e + f*x])/(24*(-c + d)^3*(c + d)^3*Sqrt[c^2 - d^2]*f*(d + c*Cos[e + f*x])^3)`**3.200.3 Rubi [A] (verified)**Time = 0.43 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.44, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3042, 4475, 107, 105, 105, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c+d\sec(e+fx))^4} dx$$

$$\downarrow 3042$$

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(a\csc(e+fx+\frac{\pi}{2})+a)^2}{(c+d\csc(e+fx+\frac{\pi}{2}))^4} dx$$

$$\downarrow 4475$$

$$\frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^4} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

$$\downarrow 107$$

3.200. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^4} dx$

$$a^2 \tan(e + fx) \left(\frac{(3c-2d) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3} d \sec(e+fx)}{3(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a^2(c^2-d^2)(c+d \sec(e+fx))^3} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 105

$$a^2 \tan(e + fx) \left(\frac{(3c-2d) \left(\frac{3a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d \sec(e+fx))^2} \right)}{3(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a^2(c^2-d^2)(c+d \sec(e+fx))^3} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 105

$$a^2 \tan(e + fx) \left(\frac{(3c-2d) \left(\frac{3a \left(\frac{a \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{c+d} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d \sec(e+fx))^2} \right)}{3(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a^2(c^2-d^2)(c+d \sec(e+fx))^3} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 104

$$a^2 \tan(e + fx) \left(\frac{(3c-2d) \left(\frac{3a \left(\frac{2a \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{c+d} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d \sec(e+fx))^2} \right)}{3(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a^2(c^2-d^2)(c+d \sec(e+fx))^3} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 218

3.200. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^4} dx$

$$a^2 \tan(e + fx) \left(\frac{d\sqrt{a - a \sec(e + fx)}(a \sec(e + fx) + a)^{5/2}}{3a^2(c^2 - d^2)(c + d \sec(e + fx))^3} + \frac{(3c - 2d) \left(\frac{3a \left(\frac{2 \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e + fx) + a}}{\sqrt{c-d}\sqrt{a - a \sec(e + fx)}}\right)}{\sqrt{c-d}(c+d)^{3/2}} - \frac{\sqrt{a - a \sec(e + fx)}\sqrt{a \sec(e + fx) + a}}{a(c+d)(c+d \sec(e + fx))} \right)}{2(c+d)} \right)}{3(c^2 - d^2)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^4,x]`

output `-((a^2*((d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2))/(3*a^2*(c^2 - d^2)*(c + d*Sec[e + f*x])^3) + ((3*c - 2*d)*(-1/2*(Sqrt[a - a*Sec[e + f*x]])*(a + a*Sec[e + f*x])^(3/2))/(a*(c + d)*(c + d*Sec[e + f*x])^2) + (3*a*((2*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(Sqrt[c - d]*(c + d)^(3/2)) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a*(c + d)*(c + d*Sec[e + f*x])))/(2*(c + d)))/(3*(c^2 - d^2))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))`

3.200.3.1 Defintions of rubi rules used

rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 105 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((m + 1)*(b*e - a*f))), x] - Simp[n*((d*e - c*f)/((m + 1)*(b*e - a*f))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`

rule 107 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.200.4 Maple [A] (verified)

Time = 1.05 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.07

method	result
derivativedivides	$8a^2 \left(\frac{\frac{(3c-2d)(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{8c^3+24c^2d+24cd^2+8d^3} - \frac{(3c-2d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3(c^2+2cd+d^2)} + \frac{(5c-6d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{8(c+d)(c-d)} \frac{(3c-2d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{8(c^4+2c^3d-2cd^3-d^4)\sqrt{(c+d)(c-d)}} \right) \frac{f}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)^3}$
default	$8a^2 \left(\frac{\frac{(3c-2d)(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{8c^3+24c^2d+24cd^2+8d^3} - \frac{(3c-2d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3(c^2+2cd+d^2)} + \frac{(5c-6d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{8(c+d)(c-d)} \frac{(3c-2d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{8(c^4+2c^3d-2cd^3-d^4)\sqrt{(c+d)(c-d)}} \right) \frac{f}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)^3}$
risch	$\frac{ia^2(7c^5d+2c^3d^3+6c^4d^2-12c^6+12cd^5e^{2i(fx+e)}+24c^3d^3e^{4i(fx+e)}+36c^2d^4e^{4i(fx+e)}+12cd^5e^{4i(fx+e)}-72c^5de^{3i(fx+e)})}{\dots}$

3.200. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^4} dx$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^4,x,method=_RETURNVERBOSE)
```

```
output 8/f*a^2*(-(1/8*(3*c-2*d)*(c-d)/(c^3+3*c^2*d+3*c*d^2+d^3)*tan(1/2*f*x+1/2*e)^5-1/3*(3*c-2*d)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3+1/8*(5*c-6*d)/(c+d)/(c-d)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^3+1/8*(3*c-2*d)/(c^4+2*c^3*d-2*c*d^3-d^4)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))
```

3.200.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 588 vs. $2(198) = 396$.

Time = 0.34 (sec) , antiderivative size = 1234, normalized size of antiderivative = 5.79

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^4} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^4,x, algorithm="fracas")
```

output

```
[1/12*(3*(3*a^2*c*d^3 - 2*a^2*d^4 + (3*a^2*c^4 - 2*a^2*c^3*d)*cos(f*x + e)
^3 + 3*(3*a^2*c^3*d - 2*a^2*c^2*d^2)*cos(f*x + e)^2 + 3*(3*a^2*c^2*d^2 - 2
*a^2*c*d^3)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 -
2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x +
e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(a^
2*c^4*d + 6*a^2*c^3*d^2 - 11*a^2*c^2*d^3 - 6*a^2*c*d^4 + 10*a^2*d^5 + (12*
a^2*c^5 - 7*a^2*c^4*d - 18*a^2*c^3*d^2 + 5*a^2*c^2*d^3 + 6*a^2*c*d^4 + 2*a
^2*d^5)*cos(f*x + e)^2 + 3*(a^2*c^5 + 6*a^2*c^4*d - 8*a^2*c^3*d^2 - 8*a^2*
c^2*d^3 + 7*a^2*c*d^4 + 2*a^2*d^5)*cos(f*x + e))*sin(f*x + e))/((c^9 + 2*c
^8*d - c^7*d^2 - 4*c^6*d^3 - c^5*d^4 + 2*c^4*d^5 + c^3*d^6)*f*cos(f*x + e)
^3 + 3*(c^8*d + 2*c^7*d^2 - c^6*d^3 - 4*c^5*d^4 - c^4*d^5 + 2*c^3*d^6 + c^
2*d^7)*f*cos(f*x + e)^2 + 3*(c^7*d^2 + 2*c^6*d^3 - c^5*d^4 - 4*c^4*d^5 - c
^3*d^6 + 2*c^2*d^7 + c*d^8)*f*cos(f*x + e) + (c^6*d^3 + 2*c^5*d^4 - c^4*d^
5 - 4*c^3*d^6 - c^2*d^7 + 2*c*d^8 + d^9)*f), 1/6*(3*(3*a^2*c*d^3 - 2*a^2*d
^4 + (3*a^2*c^4 - 2*a^2*c^3*d)*cos(f*x + e)^3 + 3*(3*a^2*c^3*d - 2*a^2*c^2
*d^2)*cos(f*x + e)^2 + 3*(3*a^2*c^2*d^2 - 2*a^2*c*d^3)*cos(f*x + e))*sqrt(
-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin
(f*x + e))) + (a^2*c^4*d + 6*a^2*c^3*d^2 - 11*a^2*c^2*d^3 - 6*a^2*c*d^4 +
10*a^2*d^5 + (12*a^2*c^5 - 7*a^2*c^4*d - 18*a^2*c^3*d^2 + 5*a^2*c^2*d^3 +
6*a^2*c*d^4 + 2*a^2*d^5)*cos(f*x + e)^2 + 3*(a^2*c^5 + 6*a^2*c^4*d - 8*...
```

3.200.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^4} dx$$

$$= a^2 \left(\int \frac{\sec(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx \right.$$

$$+ \int \frac{2\sec^2(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx$$

$$\left. + \int \frac{\sec^3(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e))**4,x)`

```
output a**2*(Integral(sec(e + f*x)/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x) + Integral(2*sec(e + f*x)**2/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x) + Integral(sec(e + f*x)**3/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x))
```

3.200.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^4} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^4,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de
```

3.200.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 403 vs. $2(198) = 396$.

Time = 0.41 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.89

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^4} dx = \frac{3(3a^2c - 2a^2d) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right)}{(c^4 + 2c^3d - 2cd^3 - d^4)\sqrt{-c^2+d^2}} + \frac{9a^2c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 24a^2c^2d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\dots}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^4,x, algorithm="giac")
```



```
output -1/3*(3*(3*a^2*c - 2*a^2*d)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*
d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 +
d^2)))/((c^4 + 2*c^3*d - 2*c*d^3 - d^4)*sqrt(-c^2 + d^2)) + (9*a^2*c^3*tan
(1/2*f*x + 1/2*e)^5 - 24*a^2*c^2*d*tan(1/2*f*x + 1/2*e)^5 + 21*a^2*c*d^2*t
an(1/2*f*x + 1/2*e)^5 - 6*a^2*d^3*tan(1/2*f*x + 1/2*e)^5 - 24*a^2*c^3*tan(
1/2*f*x + 1/2*e)^3 + 16*a^2*c^2*d*tan(1/2*f*x + 1/2*e)^3 + 24*a^2*c*d^2*ta
n(1/2*f*x + 1/2*e)^3 - 16*a^2*d^3*tan(1/2*f*x + 1/2*e)^3 + 15*a^2*c^3*tan(
1/2*f*x + 1/2*e) + 12*a^2*c^2*d*tan(1/2*f*x + 1/2*e) - 21*a^2*c*d^2*tan(1/
2*f*x + 1/2*e) - 18*a^2*d^3*tan(1/2*f*x + 1/2*e))/((c^4 + 2*c^3*d - 2*c*d^
3 - d^4)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^3))
/f
```

3.200.9 Mupad [B] (verification not implemented)

Time = 16.58 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.34

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^4} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (3a^2c^2 - 5a^2cd + 2a^2d^2)}{(c+d)^3} - \frac{8 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (3a^2c - 2a^2d)}{3(c+d)^2} + \frac{a^2}{c+d}$$

$$+ \frac{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (-3c^3 - 3c^2d + 3cd^2 + 3d^3) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 (-3c^3 + 3c^2d + 3cd^2 - 3d^3) + 3cd^2 \right)}{f(c+d)^{7/2}(c-d)^{3/2}}$$

$$+ \frac{2a^2 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\sqrt{c-d}}{\sqrt{c+d}}\right) \left(\frac{3c}{2} - d\right)}{f(c+d)^{7/2}(c-d)^{3/2}}$$

```
input int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c + d/cos(e + f*x))^4),x)
```

```
output ((tan(e/2 + (f*x)/2)^5*(3*a^2*c^2 + 2*a^2*d^2 - 5*a^2*c*d))/(c + d)^3 - (8
*tan(e/2 + (f*x)/2)^3*(3*a^2*c - 2*a^2*d))/(3*(c + d)^2) + (a^2*tan(e/2 +
(f*x)/2)*(5*c - 6*d))/((c + d)*(c - d))/(f*(tan(e/2 + (f*x)/2)^2*(3*c*d^2
- 3*c^2*d - 3*c^3 + 3*d^3) - tan(e/2 + (f*x)/2)^4*(3*c*d^2 + 3*c^2*d - 3*
c^3 - 3*d^3) + 3*c*d^2 + 3*c^2*d + c^3 + d^3 - tan(e/2 + (f*x)/2)^6*(3*c*d
^2 - 3*c^2*d + c^3 - d^3))) + (2*a^2*atanh((tan(e/2 + (f*x)/2)*(c - d)^(1/
2)))/(c + d)^(1/2))*((3*c)/2 - d)/(f*(c + d)^(7/2)*(c - d)^(3/2))
```

3.201 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^5} dx$

3.201.1 Optimal result	1401
3.201.2 Mathematica [A] (verified)	1402
3.201.3 Rubi [A] (verified)	1402
3.201.4 Maple [A] (verified)	1408
3.201.5 Fricas [B] (verification not implemented)	1409
3.201.6 Sympy [F]	1410
3.201.7 Maxima [F(-2)]	1411
3.201.8 Giac [B] (verification not implemented)	1411
3.201.9 Mupad [B] (verification not implemented)	1412

3.201.1 Optimal result

Integrand size = 31, antiderivative size = 276

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^5} dx = \frac{a^2(12c^2 - 16cd + 7d^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{4(c-d)^{5/2}(c+d)^{9/2}f} - \frac{a^2(c-d) \tan(e+fx)}{4d(c+d)f(c+d \sec(e+fx))^4} + \frac{a^2(c+8d) \tan(e+fx)}{12d(c+d)^2 f(c+d \sec(e+fx))^3} + \frac{a^2(2c^2 + 16cd - 21d^2) \tan(e+fx)}{24(c-d)d(c+d)^3 f(c+d \sec(e+fx))^2} + \frac{a^2(2c^3 + 16c^2d - 59cd^2 + 32d^3) \tan(e+fx)}{24(c-d)^2 d(c+d)^4 f(c+d \sec(e+fx))}$$

```
output 1/4*a^2*(12*c^2-16*c*d+7*d^2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c-d)^(5/2)/(c+d)^(9/2)/f-1/4*a^2*(c-d)*tan(f*x+e)/d/(c+d)/f/(c+d*sec(f*x+e))^4+1/12*a^2*(c+8*d)*tan(f*x+e)/d/(c+d)^2/f/(c+d*sec(f*x+e))^3+1/24*a^2*(2*c^2+16*c*d-21*d^2)*tan(f*x+e)/(c-d)/d/(c+d)^3/f/(c+d*sec(f*x+e))^2+1/24*a^2*(2*c^3+16*c^2*d-59*c*d^2+32*d^3)*tan(f*x+e)/(c-d)^2/d/(c+d)^4/f/(c+d*sec(f*x+e))
```

3.201.2 Mathematica [A] (verified)

Time = 8.01 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.17

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^5} dx$$

$$= a^2 \left(-\frac{24(12c^2-16cd+7d^2)\operatorname{arctanh}\left(\frac{(-c+d)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}} + \frac{(24c^5+192c^4d-446c^3d^2+128c^2d^3-148cd^4+160d^5+(144c^5-172c^4d+208c^3d^2-785c^2d^3+368cd^4+102d^5)\cos[e+fx]+2(12c^5+96c^4d-227c^3d^2+32c^2d^3+44cd^4+16d^5)\cos[2(e+fx)]+48c^5\cos[3(e+fx)]-68c^4d\cos[3(e+fx)]-16c^3d^2\cos[3(e+fx)]+5c^2d^3\cos[3(e+fx)]+16cd^4\cos[3(e+fx)]+6d^5\cos[3(e+fx)])\sin[e+fx]}{(d+c\cos[e+fx])^4(96(c-d)^2(c+d)^4f)} \right)$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^5,x]`output `(a^2*((-24*(12*c^2 - 16*c*d + 7*d^2)*ArcTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]])/Sqrt[c^2 - d^2] + ((24*c^5 + 192*c^4*d - 446*c^3*d^2 + 128*c^2*d^3 - 148*c*d^4 + 160*d^5 + (144*c^5 - 172*c^4*d + 208*c^3*d^2 - 785*c^2*d^3 + 368*c*d^4 + 102*d^5)*Cos[e + f*x] + 2*(12*c^5 + 96*c^4*d - 227*c^3*d^2 + 32*c^2*d^3 + 44*c*d^4 + 16*d^5)*Cos[2*(e + f*x)] + 48*c^5*Cos[3*(e + f*x)] - 68*c^4*d*Cos[3*(e + f*x)] - 16*c^3*d^2*Cos[3*(e + f*x)] + 5*c^2*d^3*Cos[3*(e + f*x)] + 16*c*d^4*Cos[3*(e + f*x)] + 6*d^5*Cos[3*(e + f*x)]))*Sin[e + f*x])/(d + c*Cos[e + f*x])^4)/(96*(c - d)^2*(c + d)^4*f)`**3.201.3 Rubi [A] (verified)**Time = 0.60 (sec) , antiderivative size = 462, normalized size of antiderivative = 1.67, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4475, 109, 25, 27, 168, 27, 168, 27, 168, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^2}{(c+d\sec(e+fx))^5} dx$$

$$\downarrow 3042$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^2}{\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)^5} dx$$

$$\downarrow 4475$$

3.201. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^5} dx$

$$\frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^5} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 109

$$\frac{a^2 \tan(e+fx) \left(\frac{(c-d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{4d(c+d)(c+d \sec(e+fx))^4} - \frac{\int -\frac{a^3(8d+(c+7d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))^4} d \sec(e+fx)}{4ad(c+d)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^3(8d+(c+7d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))^4} d \sec(e+fx)}{4ad(c+d)} + \frac{(c-d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{4d(c+d)(c+d \sec(e+fx))^4} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{a^2 \int \frac{8d+(c+7d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))^4} d \sec(e+fx)}{4d(c+d)} + \frac{(c-d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{4d(c+d)(c+d \sec(e+fx))^4} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 168

$$\frac{a^2 \tan(e+fx) \left(\frac{a^2 \left(\frac{\int \frac{a^2(c-d)(21d+2(c+8d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))^3} d \sec(e+fx)}{3a^2(c^2-d^2)} - \frac{(c+8d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{3a^2(c+d)(c+d \sec(e+fx))^3} \right)}{4d(c+d)} + \frac{(c-d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{4d(c+d)(c+d \sec(e+fx))^4} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{a^2 \left(\frac{(c-d) \int \frac{21d+2(c+8d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))^3} d \sec(e+fx)}{3(c^2-d^2)} - \frac{(c+8d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{3a^2(c+d)(c+d \sec(e+fx))^3} \right)}{4d(c+d)} + \frac{(c-d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{4d(c+d)(c+d \sec(e+fx))^4} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 168

3.201. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^5} dx$

$$a^2 \tan(e + fx) \left(\frac{(c-d) \left(\frac{\int \frac{a^2(2(19c-16d)d - (21d^2 - 2c(c+8d)) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))^2} d \sec(e+fx)}{2a^2(c^2-d^2)} - \frac{(2c^2+16cd-21d^2) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{2a^2(c^2-d^2)(c+d \sec(e+fx))^2} \right)}{3(c^2-d^2)} \right)}{4d(c+d)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{(c-d) \left(\frac{\int \frac{2(19c-16d)d - (21d^2 - 2c(c+8d)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))^2} d \sec(e+fx)}{2(c^2-d^2)} - \frac{(2c^2+16cd-21d^2) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{2a^2(c^2-d^2)(c+d \sec(e+fx))^2} \right)}{3(c^2-d^2)} \right)}{4d(c+d)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 168

3.201. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^5} dx$

$$\left. \begin{array}{l} a^2 \tan(e+fx) \\ a^2 \end{array} \right\} \left((c-d) \left(\frac{\int \frac{3a^2 d(12c^2 - 16dc + 7d^2)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)} a + a(c+d \sec(e+fx))} d \sec(e+fx) - \frac{(2c^3 + 16c^2 d - 59cd^2 + 32d^3) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)}}{a^2(c^2-d^2)(c+d \sec(e+fx))}}{2(c^2-d^2)} \right) \right.$$

$f \sqrt{a - a \sec(e + fx)}$

↓ 27

$$\left. \begin{array}{l} a^2 \tan(e+fx) \\ a^2 \end{array} \right\} \left((c-d) \left(\frac{\int \frac{3d(12c^2 - 16cd + 7d^2)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)} a + a(c+d \sec(e+fx))} d \sec(e+fx) - \frac{(2c^3 + 16c^2 d - 59cd^2 + 32d^3) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)}}{a^2(c^2-d^2)(c+d \sec(e+fx))}}{2(c^2-d^2)} \right) \right.$$

$f \sqrt{a - a \sec(e + fx)}$

↓ 104

3.201. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^5} dx$

$$a^2 \tan(e + fx) \left((c-d) \frac{6d(12c^2 - 16cd + 7d^2) \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}} - \frac{(2c^3 + 16c^2d - 59cd^2 + 32d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(c+d \sec(e+fx))} \right) - \frac{2(c^2-d^2)}{3(c^2-d^2)} - \frac{4d(c+d)}{4d(c+d)}$$

$f \sqrt{a - a \sec(e + fx)}$

218

$$a^2 \tan(e + fx) \left((c-d) \frac{6d(12c^2 - 16cd + 7d^2) \arctan\left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a \sec(e+fx)}}\right) - \frac{(2c^3 + 16c^2d - 59cd^2 + 32d^3) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a^2(c^2-d^2)(c+d \sec(e+fx))}}{a \sqrt{c-d} \sqrt{c+d} (c^2-d^2)} \right) - \frac{2(c^2-d^2)}{3(c^2-d^2)} - \frac{4d(c+d)}{4d(c+d)}$$

$f \sqrt{a - a \sec(e + fx)}$

```
input Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^2)/(c + d*Sec[e + f*x])^5,x]
```

3.201. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^2}{(c+d \sec(e+fx))^5} dx$

```
output -((a^2*(((c - d)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(4*d*(c + d)*(c + d*Sec[e + f*x])^4) + (a^2*(-1/3*((c + 8*d)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a^2*(c + d)*(c + d*Sec[e + f*x])^3) + ((c - d)*(-1/2*((2*c^2 + 16*c*d - 21*d^2)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a^2*(c^2 - d^2)*(c + d*Sec[e + f*x])^2) + ((6*d*(12*c^2 - 16*c*d + 7*d^2)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(a*Sqrt[c - d]*Sqrt[c + d]*(c^2 - d^2)) - ((2*c^3 + 16*c^2*d - 59*c*d^2 + 32*d^3)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]/(a^2*(c^2 - d^2)*(c + d*Sec[e + f*x])))/(2*(c^2 - d^2)))/(3*(c^2 - d^2)))/(4*d*(c + d))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.201.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 109 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```



```
rule 168 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_)*((csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])
```

3.201.4 Maple [A] (verified)

Time = 1.44 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.28

method	result
derivativedivides	$8a^2 \left(\frac{\frac{(12c^2 - 16cd + 7d^2)(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{32c^4 + 128c^3d + 192c^2d^2 + 128cd^3 + 32d^4} - \frac{11(12c^2 - 16cd + 7d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{96(c^3 + 3c^2d + 3cd^2 + d^3)} + \frac{(156c^2 - 272cd + 83d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{96(c-d)(c^2 + 2cd + d^2)} - \frac{(20c^2 - 16cd + 7d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{96(c-d)(c^2 + 2cd + d^2)} \right) \frac{f}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)^4}$
default	$8a^2 \left(\frac{\frac{(12c^2 - 16cd + 7d^2)(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{32c^4 + 128c^3d + 192c^2d^2 + 128cd^3 + 32d^4} - \frac{11(12c^2 - 16cd + 7d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{96(c^3 + 3c^2d + 3cd^2 + d^3)} + \frac{(156c^2 - 272cd + 83d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{96(c-d)(c^2 + 2cd + d^2)} - \frac{(20c^2 - 16cd + 7d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{96(c-d)(c^2 + 2cd + d^2)} \right) \frac{f}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)^4}$
risch	Expression too large to display

3.201. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^5} dx$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^5,x,method=_RETURNVERBOSE)
```

```
output 8/f*a^2*(-(1/32*(12*c^2-16*c*d+7*d^2)*(c-d)/(c^4+4*c^3*d+6*c^2*d^2+4*c*d^3+d^4)*tan(1/2*f*x+1/2*e)^7-11/96*(12*c^2-16*c*d+7*d^2)/(c^3+3*c^2*d+3*c*d^2+d^3)*tan(1/2*f*x+1/2*e)^5+1/96*(156*c^2-272*c*d+83*d^2)/(c-d)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3-1/32*(20*c^2-48*c*d+25*d^2)/(c+d)/(c^2-2*c*d+d^2)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^4+1/32*(12*c^2-16*c*d+7*d^2)/(c^6+2*c^5*d-c^4*d^2-4*c^3*d^3-c^2*d^4+2*c*d^5+d^6)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))
```

3.201.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 925 vs. $2(257) = 514$.

Time = 0.40 (sec) , antiderivative size = 1908, normalized size of antiderivative = 6.91

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^5} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^5,x, algorithm="fracas")
```

output

```
[1/48*(3*(12*a^2*c^2*d^4 - 16*a^2*c*d^5 + 7*a^2*d^6 + (12*a^2*c^6 - 16*a^2*c^5*d + 7*a^2*c^4*d^2)*cos(f*x + e)^4 + 4*(12*a^2*c^5*d - 16*a^2*c^4*d^2 + 7*a^2*c^3*d^3)*cos(f*x + e)^3 + 6*(12*a^2*c^4*d^2 - 16*a^2*c^3*d^3 + 7*a^2*c^2*d^4)*cos(f*x + e)^2 + 4*(12*a^2*c^3*d^3 - 16*a^2*c^2*d^4 + 7*a^2*c*d^5)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(2*a^2*c^5*d^2 + 16*a^2*c^4*d^3 - 61*a^2*c^3*d^4 + 16*a^2*c^2*d^5 + 59*a^2*c*d^6 - 32*a^2*d^7 + (48*a^2*c^7 - 68*a^2*c^6*d - 64*a^2*c^5*d^2 + 73*a^2*c^4*d^3 + 32*a^2*c^3*d^4 + a^2*c^2*d^5 - 16*a^2*c*d^6 - 6*a^2*d^7)*cos(f*x + e)^3 + (12*a^2*c^7 + 96*a^2*c^6*d - 239*a^2*c^5*d^2 - 64*a^2*c^4*d^3 + 271*a^2*c^3*d^4 - 16*a^2*c^2*d^5 - 44*a^2*c*d^6 - 16*a^2*d^7)*cos(f*x + e)^2 + (8*a^2*c^6*d + 64*a^2*c^5*d^2 - 208*a^2*c^4*d^3 + 16*a^2*c^3*d^4 + 221*a^2*c^2*d^5 - 80*a^2*c*d^6 - 21*a^2*d^7)*cos(f*x + e))*sin(f*x + e))/((c^12 + 2*c^11*d - 2*c^10*d^2 - 6*c^9*d^3 + 6*c^7*d^5 + 2*c^6*d^6 - 2*c^5*d^7 - c^4*d^8)*f*cos(f*x + e)^4 + 4*(c^11*d + 2*c^10*d^2 - 2*c^9*d^3 - 6*c^8*d^4 + 6*c^6*d^6 + 2*c^5*d^7 - 2*c^4*d^8 - c^3*d^9)*f*cos(f*x + e)^3 + 6*(c^10*d^2 + 2*c^9*d^3 - 2*c^8*d^4 - 6*c^7*d^5 + 6*c^5*d^7 + 2*c^4*d^8 - 2*c^3*d^9 - c^2*d^10)*f*cos(f*x + e)^2 + 4*(c^9*d^3 + 2*c^8*d^4 - 2*c^7*d^5 - 6*c^6*d^6 + 6*c^4*d^8 + 2*c^3*d^9 - 2*c^2*d^10 - c*d^11)*f*cos(f*x + e) + (c^8*...
```

3.201.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^5} dx$$

$$= a^2 \left(\int \frac{\sec(e+fx)}{c^5 + 5c^4d\sec(e+fx) + 10c^3d^2\sec^2(e+fx) + 10c^2d^3\sec^3(e+fx) + 5cd^4\sec^4(e+fx) + d^5\sec^5(e+fx)} dx \right.$$

$$+ \int \frac{2\sec^2(e+fx)}{c^5 + 5c^4d\sec(e+fx) + 10c^3d^2\sec^2(e+fx) + 10c^2d^3\sec^3(e+fx) + 5cd^4\sec^4(e+fx) + d^5\sec^5(e+fx)} dx$$

$$\left. + \int \frac{\sec^3(e+fx)}{c^5 + 5c^4d\sec(e+fx) + 10c^3d^2\sec^2(e+fx) + 10c^2d^3\sec^3(e+fx) + 5cd^4\sec^4(e+fx) + d^5\sec^5(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e))**5,x)`

```
output a**2*(Integral(sec(e + f*x)/(c**5 + 5*c**4*d*sec(e + f*x) + 10*c**3*d**2*sec(e + f*x)**2 + 10*c**2*d**3*sec(e + f*x)**3 + 5*c*d**4*sec(e + f*x)**4 + d**5*sec(e + f*x)**5), x) + Integral(2*sec(e + f*x)**2/(c**5 + 5*c**4*d*sec(e + f*x) + 10*c**3*d**2*sec(e + f*x)**2 + 10*c**2*d**3*sec(e + f*x)**3 + 5*c*d**4*sec(e + f*x)**4 + d**5*sec(e + f*x)**5), x) + Integral(sec(e + f*x)**3/(c**5 + 5*c**4*d*sec(e + f*x) + 10*c**3*d**2*sec(e + f*x)**2 + 10*c**2*d**3*sec(e + f*x)**3 + 5*c*d**4*sec(e + f*x)**4 + d**5*sec(e + f*x)**5), x))
```

3.201.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^5} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^5,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` for more de
```

3.201.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 710 vs. $2(257) = 514$.

Time = 0.51 (sec) , antiderivative size = 710, normalized size of antiderivative = 2.57

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^5} dx$$

$$\frac{3(12a^2c^2 - 16a^2cd + 7a^2d^2) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2c+2d) + \arctan \left(-\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right)}{(c^6 + 2c^5d - c^4d^2 - 4c^3d^3 - c^2d^4 + 2cd^5 + d^6)\sqrt{-c^2+d^2}} - \frac{36a^2c^5 \tan(\frac{1}{2}fx + \frac{1}{2}e)^7 - 156a^2c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 156a^2c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 36a^2c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(c^6 + 2c^5d - c^4d^2 - 4c^3d^3 - c^2d^4 + 2cd^5 + d^6)\sqrt{-c^2+d^2}}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^5,x, algorithm="giac")
```

3.201. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^2}{(c+d\sec(e+fx))^5} dx$

output

```

1/12*(3*(12*a^2*c^2 - 16*a^2*c*d + 7*a^2*d^2)*(pi*floor(1/2*(f*x + e)/pi +
1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x +
1/2*e))/sqrt(-c^2 + d^2)))/((c^6 + 2*c^5*d - c^4*d^2 - 4*c^3*d^3 - c^2*d^4
+ 2*c*d^5 + d^6)*sqrt(-c^2 + d^2)) - (36*a^2*c^5*tan(1/2*f*x + 1/2*e)^7 -
156*a^2*c^4*d*tan(1/2*f*x + 1/2*e)^7 + 273*a^2*c^3*d^2*tan(1/2*f*x + 1/2*
e)^7 - 243*a^2*c^2*d^3*tan(1/2*f*x + 1/2*e)^7 + 111*a^2*c*d^4*tan(1/2*f*x
+ 1/2*e)^7 - 21*a^2*d^5*tan(1/2*f*x + 1/2*e)^7 - 132*a^2*c^5*tan(1/2*f*x +
1/2*e)^5 + 308*a^2*c^4*d*tan(1/2*f*x + 1/2*e)^5 - 121*a^2*c^3*d^2*tan(1/2
*f*x + 1/2*e)^5 - 231*a^2*c^2*d^3*tan(1/2*f*x + 1/2*e)^5 + 253*a^2*c*d^4*t
an(1/2*f*x + 1/2*e)^5 - 77*a^2*d^5*tan(1/2*f*x + 1/2*e)^5 + 156*a^2*c^5*ta
n(1/2*f*x + 1/2*e)^3 - 116*a^2*c^4*d*tan(1/2*f*x + 1/2*e)^3 - 345*a^2*c^3*
d^2*tan(1/2*f*x + 1/2*e)^3 + 199*a^2*c^2*d^3*tan(1/2*f*x + 1/2*e)^3 + 189*
a^2*c*d^4*tan(1/2*f*x + 1/2*e)^3 - 83*a^2*d^5*tan(1/2*f*x + 1/2*e)^3 - 60*
a^2*c^5*tan(1/2*f*x + 1/2*e) - 36*a^2*c^4*d*tan(1/2*f*x + 1/2*e) + 177*a^2
*c^3*d^2*tan(1/2*f*x + 1/2*e) + 147*a^2*c^2*d^3*tan(1/2*f*x + 1/2*e) - 81*
a^2*c*d^4*tan(1/2*f*x + 1/2*e) - 75*a^2*d^5*tan(1/2*f*x + 1/2*e))/((c^6 +
2*c^5*d - c^4*d^2 - 4*c^3*d^3 - c^2*d^4 + 2*c*d^5 + d^6)*(c*tan(1/2*f*x +
1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^4))/f

```

3.201.9 Mupad [B] (verification not implemented)

Time = 17.02 (sec) , antiderivative size = 438, normalized size of antiderivative = 1.59

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^2}{(c + d \sec(e + fx))^5} dx$$

$$= \frac{\frac{11 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (12a^2c^2 - 16a^2cd + 7a^2d^2)}{12(c+d)^3} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7 (12a^2c^3 - 28a^2c^2d + 28a^2cd^2 - 12a^2d^3)}{4(c+d)^4}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 (6c^4 - 12c^2d^2 + 6d^4) + \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (-4c^4 - 8c^3d + 8cd^3 + 4d^4) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 \right)} + \frac{a^2 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)(2c-2d)(c^2-2cd+d^2)}{2\sqrt{c+d}(c-d)^{5/2}}\right) (12c^2 - 16cd + 7d^2)}{4f(c+d)^{9/2}(c-d)^{5/2}}$$

input `int((a + a/cos(e + f*x))^2/(cos(e + f*x)*(c + d/cos(e + f*x))^5),x)`

output
$$\begin{aligned} & ((11*\tan(e/2 + (f*x)/2)^5*(12*a^2*c^2 + 7*a^2*d^2 - 16*a^2*c*d))/(12*(c + \\ & d)^3) - (\tan(e/2 + (f*x)/2)^7*(12*a^2*c^3 - 7*a^2*d^3 + 23*a^2*c*d^2 - 28* \\ & a^2*c^2*d))/(4*(c + d)^4) - (a^2*\tan(e/2 + (f*x)/2)^3*(156*c^2 - 272*c*d + \\ & 83*d^2))/(12*(c + d)^2*(c - d)) + (a^2*\tan(e/2 + (f*x)/2)*(20*c^2 - 48*c* \\ & d + 25*d^2))/(4*(c + d)*(c^2 - 2*c*d + d^2)))/(f*(\tan(e/2 + (f*x)/2)^4*(6* \\ & c^4 + 6*d^4 - 12*c^2*d^2) + \tan(e/2 + (f*x)/2)^2*(8*c*d^3 - 8*c^3*d - 4*c^4 \\ & + 4*d^4) - \tan(e/2 + (f*x)/2)^6*(8*c*d^3 - 8*c^3*d + 4*c^4 - 4*d^4) + \tan \\ & (e/2 + (f*x)/2)^8*(c^4 - 4*c^3*d - 4*c*d^3 + d^4 + 6*c^2*d^2) + 4*c*d^3 + \\ & 4*c^3*d + c^4 + d^4 + 6*c^2*d^2)) + (a^2*\operatorname{atanh}((\tan(e/2 + (f*x)/2)*(2*c - \\ & 2*d)*(c^2 - 2*c*d + d^2))/(2*(c + d)^{(1/2)}*(c - d)^{(5/2)})))*(12*c^2 - 16*c \\ & *d + 7*d^2))/(4*f*(c + d)^{(9/2)}*(c - d)^{(5/2)}) \end{aligned}$$

3.202 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$

3.202.1 Optimal result	1414
3.202.2 Mathematica [A] (verified)	1415
3.202.3 Rubi [A] (verified)	1415
3.202.4 Maple [A] (verified)	1419
3.202.5 Fricas [A] (verification not implemented)	1420
3.202.6 Sympy [F]	1420
3.202.7 Maxima [B] (verification not implemented)	1421
3.202.8 Giac [B] (verification not implemented)	1422
3.202.9 Mupad [B] (verification not implemented)	1423

3.202.1 Optimal result

Integrand size = 31, antiderivative size = 288

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$$

$$= \frac{a^3(40c^3 + 90c^2d + 78cd^2 + 23d^3) \operatorname{arctanh}(\sin(e + fx))}{16f}$$

$$+ \frac{a^3(40c^3 + 90c^2d + 78cd^2 + 23d^3) \tan(e + fx)}{16f}$$

$$+ \frac{(40c^3 + 90c^2d + 78cd^2 + 23d^3)(a^3 + a^3 \sec(e + fx)) \tan(e + fx)}{48f}$$

$$+ \frac{a(3c + 8d)(a + a \sec(e + fx))^2(c + d \sec(e + fx))^2 \tan(e + fx)}{30f}$$

$$+ \frac{a(a + a \sec(e + fx))^2(c + d \sec(e + fx))^3 \tan(e + fx)}{6f}$$

$$+ \frac{a(a + a \sec(e + fx))^2(2(4c^3 + 74c^2d + 66cd^2 + 21d^3) + d(6c^2 + 62cd + 31d^2) \sec(e + fx)) \tan(e + fx)}{120f}$$

output `1/16*a^3*(40*c^3+90*c^2*d+78*c*d^2+23*d^3)*arctanh(sin(f*x+e))/f+1/16*a^3*(40*c^3+90*c^2*d+78*c*d^2+23*d^3)*tan(f*x+e)/f+1/48*(40*c^3+90*c^2*d+78*c*d^2+23*d^3)*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/f+1/30*a*(3*c+8*d)*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^2*tan(f*x+e)/f+1/6*a*(a+a*sec(f*x+e))^2*(c+d*sec(f*x+e))^3*tan(f*x+e)/f+1/120*a*(a+a*sec(f*x+e))^2*(8*c^3+148*c^2*d+132*c*d^2+42*d^3+d*(6*c^2+62*c*d+31*d^2)*sec(f*x+e))*tan(f*x+e)/f`

3.202. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$

3.202.2 Mathematica [A] (verified)

Time = 6.09 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.59

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$$

$$= \frac{a^3(15(40c^3 + 90c^2d + 78cd^2 + 23d^3) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx)(15(24c^3 + 90c^2d + 78cd^2 + 23d^3) \sec(e + fx) + 10d(18c^2 + 54cd + 23d^2) \sec^3(e + fx) + 40d^3 \sec^5(e + fx) + 16(c + d)(60(c + d)^2 + 5(c^2 + 8cd + 7d^2) \tan^2(e + fx) + 9d^2 \tan^4(e + fx))))}{240f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^3,x]`output `(a^3*(15*(40*c^3 + 90*c^2*d + 78*c*d^2 + 23*d^3)*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(15*(24*c^3 + 90*c^2*d + 78*c*d^2 + 23*d^3)*Sec[e + f*x] + 10*d*(18*c^2 + 54*c*d + 23*d^2)*Sec[e + f*x]^3 + 40*d^3*Sec[e + f*x]^5 + 16*(c + d)*(60*(c + d)^2 + 5*(c^2 + 8*c*d + 7*d^2)*Tan[e + f*x]^2 + 9*d^2*Tan[e + f*x]^4)))/(240*f)`**3.202.3 Rubi [A] (verified)**Time = 0.48 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.20, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4475, 111, 25, 27, 164, 60, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3(c + d \sec(e + fx))^3 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 dx$$

$$\downarrow \text{4475}$$

$$\frac{a^2 \tan(e + fx) \int \frac{(\sec(e + fx)a + a)^{5/2} (c + d \sec(e + fx))^3 d \sec(e + fx)}{\sqrt{a - a \sec(e + fx)}}}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

$$\downarrow \text{111}$$

$$\frac{a^2 \tan(e + fx) \left(- \int \frac{a^2 (\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx)) (6c^2+3dc+2d^2+d(8c+3d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} 6a^2} d \sec(e+fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{6a^2} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e + fx) \left(\int \frac{a^2 (\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx)) (6c^2+3dc+2d^2+d(8c+3d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} 6a^2} d \sec(e+fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{6a^2} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e + fx) \left(\frac{1}{6} \int \frac{(\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx)) (6c^2+3dc+2d^2+d(8c+3d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{6a^2} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 164

$$\frac{a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{3}{20} (40c^3 + 90c^2d + 78cd^2 + 23d^3) \int \frac{(\sec(e+fx)a+a)^{5/2}}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{6a^2} \right) \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 60

$$\frac{a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{3}{20} (40c^3 + 90c^2d + 78cd^2 + 23d^3) \left(\frac{5}{3} a \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{\sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{3a} \right) \right) \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 60

$$\frac{a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{3}{20} (40c^3 + 90c^2d + 78cd^2 + 23d^3) \left(\frac{5}{3} a \left(\frac{3}{2} a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{\sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)}{2a} \right) \right) \right) \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 60

$$\frac{a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{3}{20} (40c^3 + 90c^2d + 78cd^2 + 23d^3) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) - \frac{1}{2a} \int \frac{1}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) \right) \right) \right) \right) \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{3}{20} (40c^3 + 90c^2d + 78cd^2 + 23d^3) \right) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(2a \int \frac{1}{-\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} - \frac{\sqrt{a-a \sec(e+fx)}}{a} \right) \right) \right)$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{1}{6} \left(\frac{3}{20} (40c^3 + 90c^2d + 78cd^2 + 23d^3) \right) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(-2 \arctan \left(\frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}} \right) - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a} \right) \right) \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^3,x]`

output `-((a^2*(-1/6*(d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(7/2)*(c + d*Sec[e + f*x])^2)/a^2 + (-1/20*(d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(7/2)*(70*c^2 + 54*c*d + 19*d^2 + 4*d*(8*c + 3*d)*Sec[e + f*x]))/a^2 + (3*(40*c^3 + 90*c^2*d + 78*c*d^2 + 23*d^3)*(-1/3*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2))/a + (5*a*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/a + (3*a*(-2*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/a))/2))/3))/20)/6)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))`

3.202.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 111 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_, x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1
))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)
^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m
- 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m
+ n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] &
& GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

```
rule 164 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_
))*((g_.) + (h_.)*(x_)), x_] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) -
b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x))*(a + b*x)^(m + 1)*((
c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h
*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n +
3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) +
d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(
a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x]
&& NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[
^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]),
], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[
b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || In
tegerQ[m - 1/2])
```

3.202.4 Maple [A] (verified)

Time = 6.19 (sec) , antiderivative size = 355, normalized size of antiderivative = 1.23

method	result
norman	$-\frac{33a^3(40c^3+90c^2d+78cd^2+23d^3)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{20f} + \frac{17a^3(40c^3+90c^2d+78cd^2+23d^3)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^9}{24f} - \frac{a^3(40c^3+90c^2d+78cd^2+23d^3)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{8f}$
parallelrisch	$6\left(-\frac{25\left(\frac{2\cos(4fx+4e)}{5}+\frac{2}{3}+\cos(2fx+2e)+\frac{\cos(6fx+6e)}{15}\right)\left(c^3+\frac{9}{4}c^2d+\frac{39}{20}cd^2+\frac{23}{40}d^3\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{4} + \frac{25\left(\frac{2\cos(4fx+4e)}{5}+\frac{2}{3}\right)}{4}\right)$
parts	$-\frac{(3a^3cd^2+3a^3d^3)\left(-\frac{8}{15}-\frac{\sec(fx+e)^4}{5}-\frac{4\sec(fx+e)^2}{15}\right)\tan(fx+e)}{f} + \frac{(3c^3a^3+3a^3c^2d)\tan(fx+e)}{f} + \frac{(3a^3c^2d+9a^3cd^2)\tan(fx+e)}{f}$
derivativedivides	$-c^3a^3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+3a^3c^2d\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)$
default	$-c^3a^3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+3a^3c^2d\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)$
risch	$-\frac{ia^3(-880c^3-1824cd^2-544d^3-2160c^2d-1170cd^2e^{i(fx+e)}-10944cd^2e^{2i(fx+e)}-5670cd^2e^{3i(fx+e)}-18240cd^2e^{6i(fx+e)})}{f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^3,x,method=_RETURNVERBO
SE)
```

```
output (-33/20*a^3*(40*c^3+90*c^2*d+78*c*d^2+23*d^3)/f*tan(1/2*f*x+1/2*e)^7+17/24
*a^3*(40*c^3+90*c^2*d+78*c*d^2+23*d^3)/f*tan(1/2*f*x+1/2*e)^9-1/8*a^3*(40*
c^3+90*c^2*d+78*c*d^2+23*d^3)/f*tan(1/2*f*x+1/2*e)^11+1/8*a^3*(88*c^3+294*
c^2*d+306*c*d^2+105*d^3)/f*tan(1/2*f*x+1/2*e)+3/20*a^3*(520*c^3+1250*c^2*d
+998*c*d^2+323*d^3)/f*tan(1/2*f*x+1/2*e)^5-1/24*a^3*(1112*c^3+3078*c^2*d+2
514*c*d^2+633*d^3)/f*tan(1/2*f*x+1/2*e)^3)/(tan(1/2*f*x+1/2*e)^2-1)^6-1/16
*a^3*(40*c^3+90*c^2*d+78*c*d^2+23*d^3)/f*ln(tan(1/2*f*x+1/2*e)-1)+1/16*a^3
*(40*c^3+90*c^2*d+78*c*d^2+23*d^3)/f*ln(tan(1/2*f*x+1/2*e)+1)
```

3.202. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$

3.202.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.17

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$$

$$= \frac{15(40a^3c^3 + 90a^3c^2d + 78a^3cd^2 + 23a^3d^3) \cos(fx + e)^6 \log(\sin(fx + e) + 1) - 15(40a^3c^3 + 90a^3c^2d + 78a^3cd^2 + 23a^3d^3) \cos(fx + e)^6 \log(-\sin(fx + e) + 1) + 2(40a^3d^3 + 16(55a^3c^3 + 135a^3c^2d + 114a^3cd^2 + 34a^3d^3) \cos(fx + e)^5 + 15(24a^3c^3 + 90a^3c^2d + 78a^3cd^2 + 23a^3d^3) \cos(fx + e)^4 + 16(5a^3c^3 + 45a^3c^2d + 57a^3cd^2 + 17a^3d^3) \cos(fx + e)^3 + 10(18a^3c^2d + 54a^3cd^2 + 23a^3d^3) \cos(fx + e)^2 + 144(a^3cd^2 + a^3d^3) \cos(fx + e)) \sin(fx + e)}{(f \cos(fx + e))^6}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^3,x, algorithm="fricas")
```

```
output 1/480*(15*(40*a^3*c^3 + 90*a^3*c^2*d + 78*a^3*c*d^2 + 23*a^3*d^3)*cos(f*x + e)^6*log(sin(f*x + e) + 1) - 15*(40*a^3*c^3 + 90*a^3*c^2*d + 78*a^3*c*d^2 + 23*a^3*d^3)*cos(f*x + e)^6*log(-sin(f*x + e) + 1) + 2*(40*a^3*d^3 + 16*(55*a^3*c^3 + 135*a^3*c^2*d + 114*a^3*c*d^2 + 34*a^3*d^3)*cos(f*x + e)^5 + 15*(24*a^3*c^3 + 90*a^3*c^2*d + 78*a^3*c*d^2 + 23*a^3*d^3)*cos(f*x + e)^4 + 16*(5*a^3*c^3 + 45*a^3*c^2*d + 57*a^3*c*d^2 + 17*a^3*d^3)*cos(f*x + e)^3 + 10*(18*a^3*c^2*d + 54*a^3*c*d^2 + 23*a^3*d^3)*cos(f*x + e)^2 + 144*(a^3*c*d^2 + a^3*d^3)*cos(f*x + e))*sin(f*x + e))/(f*cos(f*x + e)^6)
```

3.202.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$$

$$= a^3 \left(\int c^3 \sec(e + fx) dx + \int 3c^3 \sec^2(e + fx) dx + \int 3c^3 \sec^3(e + fx) dx \right.$$

$$+ \int c^3 \sec^4(e + fx) dx + \int d^3 \sec^4(e + fx) dx + \int 3d^3 \sec^5(e + fx) dx$$

$$+ \int 3d^3 \sec^6(e + fx) dx + \int d^3 \sec^7(e + fx) dx + \int 3cd^2 \sec^3(e + fx) dx$$

$$+ \int 9cd^2 \sec^4(e + fx) dx + \int 9cd^2 \sec^5(e + fx) dx + \int 3cd^2 \sec^6(e + fx) dx$$

$$+ \left. \int 3c^2d \sec^2(e + fx) dx + \int 9c^2d \sec^3(e + fx) dx + \int 9c^2d \sec^4(e + fx) dx \right.$$

$$\left. + \int 3c^2d \sec^5(e + fx) dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c+d*sec(f*x+e))**3,x)`

output `a**3*(Integral(c**3*sec(e + f*x), x) + Integral(3*c**3*sec(e + f*x)**2, x) + Integral(3*c**3*sec(e + f*x)**3, x) + Integral(c**3*sec(e + f*x)**4, x) + Integral(d**3*sec(e + f*x)**4, x) + Integral(3*d**3*sec(e + f*x)**5, x) + Integral(3*d**3*sec(e + f*x)**6, x) + Integral(d**3*sec(e + f*x)**7, x) + Integral(3*c*d**2*sec(e + f*x)**3, x) + Integral(9*c*d**2*sec(e + f*x)**4, x) + Integral(9*c*d**2*sec(e + f*x)**5, x) + Integral(3*c*d**2*sec(e + f*x)**6, x) + Integral(3*c**2*d*sec(e + f*x)**2, x) + Integral(9*c**2*d*sec(e + f*x)**3, x) + Integral(9*c**2*d*sec(e + f*x)**4, x) + Integral(3*c**2*d*sec(e + f*x)**5, x))`

3.202.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 701 vs. $2(273) = 546$.

Time = 0.24 (sec) , antiderivative size = 701, normalized size of antiderivative = 2.43

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$$

$$= \frac{160 (\tan (fx + e))^3 + 3 \tan (fx + e) a^3 c^3 + 1440 (\tan (fx + e))^3 + 3 \tan (fx + e) a^3 c^2 d + 96 (3 \tan (fx + e) - 1) a^3 c^2 d + 96 (3 \tan (fx + e) - 1) a^3 c d + 96 (3 \tan (fx + e) - 1) a^3 d}{1}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output

```

1/480*(160*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c^3 + 1440*(tan(f*x + e)^
3 + 3*tan(f*x + e))*a^3*c^2*d + 96*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 +
15*tan(f*x + e))*a^3*c*d^2 + 1440*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c
*d^2 + 96*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^3*d^3
+ 160*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*d^3 - 5*a^3*d^3*(2*(15*sin(f*
x + e)^5 - 40*sin(f*x + e)^3 + 33*sin(f*x + e))/(sin(f*x + e)^6 - 3*sin(f*
x + e)^4 + 3*sin(f*x + e)^2 - 1) - 15*log(sin(f*x + e) + 1) + 15*log(sin(f
*x + e) - 1)) - 90*a^3*c^2*d*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f
*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*
x + e) - 1)) - 270*a^3*c*d^2*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f
*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*
x + e) - 1)) - 90*a^3*d^3*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x
+ e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x +
e) - 1)) - 360*a^3*c^3*(2*sin(f*x + e))/(sin(f*x + e)^2 - 1) - log(sin(f*x
+ e) + 1) + log(sin(f*x + e) - 1)) - 1080*a^3*c^2*d*(2*sin(f*x + e))/(sin(
f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 360*a^3
*c*d^2*(2*sin(f*x + e))/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(
sin(f*x + e) - 1)) + 480*a^3*c^3*log(sec(f*x + e) + tan(f*x + e)) + 1440*a
^3*c^3*tan(f*x + e) + 1440*a^3*c^2*d*tan(f*x + e))/f

```

3.202.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 584 vs. $2(273) = 546$.

Time = 0.44 (sec) , antiderivative size = 584, normalized size of antiderivative = 2.03

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$$

$$= \frac{15(40a^3c^3 + 90a^3c^2d + 78a^3cd^2 + 23a^3d^3) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 15(40a^3c^3 + 90a^3c^2d + 78a^3cd^2 + 23a^3d^3)}{\dots}$$

input

```

integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^3,x, algorithm="g
iac")

```

```

output 1/240*(15*(40*a^3*c^3 + 90*a^3*c^2*d + 78*a^3*c*d^2 + 23*a^3*d^3)*log(abs(
tan(1/2*f*x + 1/2*e) + 1)) - 15*(40*a^3*c^3 + 90*a^3*c^2*d + 78*a^3*c*d^2
+ 23*a^3*d^3)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(600*a^3*c^3*tan(1/2*
f*x + 1/2*e)^11 + 1350*a^3*c^2*d*tan(1/2*f*x + 1/2*e)^11 + 1170*a^3*c*d^2*
tan(1/2*f*x + 1/2*e)^11 + 345*a^3*d^3*tan(1/2*f*x + 1/2*e)^11 - 3400*a^3*c
^3*tan(1/2*f*x + 1/2*e)^9 - 7650*a^3*c^2*d*tan(1/2*f*x + 1/2*e)^9 - 6630*a
^3*c*d^2*tan(1/2*f*x + 1/2*e)^9 - 1955*a^3*d^3*tan(1/2*f*x + 1/2*e)^9 + 79
20*a^3*c^3*tan(1/2*f*x + 1/2*e)^7 + 17820*a^3*c^2*d*tan(1/2*f*x + 1/2*e)^7
+ 15444*a^3*c*d^2*tan(1/2*f*x + 1/2*e)^7 + 4554*a^3*d^3*tan(1/2*f*x + 1/2
*e)^7 - 9360*a^3*c^3*tan(1/2*f*x + 1/2*e)^5 - 22500*a^3*c^2*d*tan(1/2*f*x
+ 1/2*e)^5 - 17964*a^3*c*d^2*tan(1/2*f*x + 1/2*e)^5 - 5814*a^3*d^3*tan(1/2
*f*x + 1/2*e)^5 + 5560*a^3*c^3*tan(1/2*f*x + 1/2*e)^3 + 15390*a^3*c^2*d*ta
n(1/2*f*x + 1/2*e)^3 + 12570*a^3*c*d^2*tan(1/2*f*x + 1/2*e)^3 + 3165*a^3*d
^3*tan(1/2*f*x + 1/2*e)^3 - 1320*a^3*c^3*tan(1/2*f*x + 1/2*e) - 4410*a^3*c
^2*d*tan(1/2*f*x + 1/2*e) - 4590*a^3*c*d^2*tan(1/2*f*x + 1/2*e) - 1575*a^3
*d^3*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^6)/f

```

3.202.9 Mupad [B] (verification not implemented)

Time = 17.05 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.43

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^3 dx$$

$$= \frac{a^3 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)(40c^3 + 90c^2d + 78cd^2 + 23d^3)}{4(10c^3 + \frac{45c^2d}{2} + \frac{39cd^2}{2} + \frac{23d^3}{4})}\right)(40c^3 + 90c^2d + 78cd^2 + 23d^3)}{8f}$$

$$+ \left(5a^3c^3 + \frac{45a^3c^2d}{4} + \frac{39a^3cd^2}{4} + \frac{23a^3d^3}{8}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11} + \left(-\frac{85a^3c^3}{3} - \frac{255a^3c^2d}{4} - \frac{221a^3cd^2}{4} - \frac{391a^3d^3}{24}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9$$

```

input int(((a + a/cos(e + f*x))^3*(c + d/cos(e + f*x))^3)/cos(e + f*x),x)

```


output $(a^3 \operatorname{atanh}(\tan(e/2 + (f*x)/2) * (78*c*d^2 + 90*c^2*d + 40*c^3 + 23*d^3)) / (4 * ((39*c*d^2)/2 + (45*c^2*d)/2 + 10*c^3 + (23*d^3)/4)) * (78*c*d^2 + 90*c^2*d + 40*c^3 + 23*d^3) / (8*f) - (\tan(e/2 + (f*x)/2)^{11} * (5*a^3*c^3 + (23*a^3*d^3)/8 + (39*a^3*c*d^2)/4 + (45*a^3*c^2*d)/4) - \tan(e/2 + (f*x)/2)^9 * ((85*a^3*c^3)/3 + (391*a^3*d^3)/24 + (221*a^3*c*d^2)/4 + (255*a^3*c^2*d)/4) + \tan(e/2 + (f*x)/2)^3 * ((139*a^3*c^3)/3 + (211*a^3*d^3)/8 + (419*a^3*c*d^2)/4 + (513*a^3*c^2*d)/4) + \tan(e/2 + (f*x)/2)^7 * (66*a^3*c^3 + (759*a^3*d^3)/20 + (1287*a^3*c*d^2)/10 + (297*a^3*c^2*d)/2) - \tan(e/2 + (f*x)/2)^5 * (78*a^3*c^3 + (969*a^3*d^3)/20 + (1497*a^3*c*d^2)/10 + (375*a^3*c^2*d)/2) - \tan(e/2 + (f*x)/2) * (11*a^3*c^3 + (105*a^3*d^3)/8 + (153*a^3*c*d^2)/4 + (147*a^3*c^2*d)/4) / (f * (15*\tan(e/2 + (f*x)/2)^4 - 6*\tan(e/2 + (f*x)/2)^2 - 20*\tan(e/2 + (f*x)/2)^6 + 15*\tan(e/2 + (f*x)/2)^8 - 6*\tan(e/2 + (f*x)/2)^{10} + \tan(e/2 + (f*x)/2)^{12} + 1))$

3.203 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$

3.203.1 Optimal result	1425
3.203.2 Mathematica [A] (verified)	1426
3.203.3 Rubi [A] (verified)	1426
3.203.4 Maple [A] (verified)	1430
3.203.5 Fricas [A] (verification not implemented)	1430
3.203.6 Sympy [F]	1431
3.203.7 Maxima [A] (verification not implemented)	1432
3.203.8 Giac [A] (verification not implemented)	1432
3.203.9 Mupad [B] (verification not implemented)	1433

3.203.1 Optimal result

Integrand size = 31, antiderivative size = 257

$$\begin{aligned}
 & \int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx \\
 &= \frac{a^3(20c^2 + 30cd + 13d^2) \operatorname{arctanh}(\sin(e + fx))}{8f} \\
 &+ \frac{a^3(2c^4 - 15c^3d + 72c^2d^2 + 180cd^3 + 76d^4) \tan(e + fx)}{30d^2f} \\
 &+ \frac{a^3(4c^3 - 30c^2d + 146cd^2 + 195d^3) \sec(e + fx) \tan(e + fx)}{120df} \\
 &+ \frac{a^3(2c^2 - 15cd + 76d^2)(c + d \sec(e + fx))^2 \tan(e + fx)}{60d^2f} \\
 &- \frac{a^3(2c - 11d)(c + d \sec(e + fx))^3 \tan(e + fx)}{20d^2f} \\
 &+ \frac{(a^3 + a^3 \sec(e + fx))(c + d \sec(e + fx))^3 \tan(e + fx)}{5df}
 \end{aligned}$$

output

```

1/8*a^3*(20*c^2+30*c*d+13*d^2)*arctanh(sin(f*x+e))/f+1/30*a^3*(2*c^4-15*c^
3*d+72*c^2*d^2+180*c*d^3+76*d^4)*tan(f*x+e)/d^2/f+1/120*a^3*(4*c^3-30*c^2*
d+146*c*d^2+195*d^3)*sec(f*x+e)*tan(f*x+e)/d/f+1/60*a^3*(2*c^2-15*c*d+76*d
^2)*(c+d*sec(f*x+e))^2*tan(f*x+e)/d^2/f-1/20*a^3*(2*c-11*d)*(c+d*sec(f*x+e
))^3*tan(f*x+e)/d^2/f+1/5*(a^3+a^3*sec(f*x+e))*(c+d*sec(f*x+e))^3*tan(f*x+
e)/d/f

```

3.203.2 Mathematica [A] (verified)

Time = 4.43 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.51

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$$

$$= \frac{a^3(15(20c^2 + 30cd + 13d^2) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx) (15(12c^2 + 30cd + 13d^2) \sec(e + fx) + 30cd + 13d^2))}{120f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^2,x]`output `(a^3*(15*(20*c^2 + 30*c*d + 13*d^2)*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(15*(12*c^2 + 30*c*d + 13*d^2)*Sec[e + f*x] + 30*d*(2*c + 3*d)*Sec[e + f*x]^3 + 8*(60*(c + d)^2 + 5*(c^2 + 6*c*d + 5*d^2)*Tan[e + f*x]^2 + 3*d^2*Tan[e + f*x]^4)))/(120*f)`**3.203.3 Rubi [A] (verified)**Time = 0.41 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.20, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4475, 101, 25, 27, 90, 60, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a \sec(e + fx) + a)^3(c + d \sec(e + fx))^2 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 dx$$

$$\downarrow 4475$$

$$- \frac{a^2 \tan(e + fx) \int \frac{(\sec(e + fx)a + a)^{5/2}(c + d \sec(e + fx))^2 d \sec(e + fx)}{\sqrt{a - a \sec(e + fx)}}}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

$$\downarrow 101$$

3.203. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$

$$a^2 \tan(e + fx) \left(- \frac{\int - \frac{a^2 (\sec(e+fx)a+a)^{5/2} (5c^2+3dc+d^2+3d(2c+d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx)}{5a^2} - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)^{7/2} (c+d \sec(e+fx))}{5a^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 25

$$a^2 \tan(e + fx) \left(\int \frac{a^2 (\sec(e+fx)a+a)^{5/2} (5c^2+3dc+d^2+3d(2c+d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e+fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)^{7/2} (c+d \sec(e+fx))}{5a^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{1}{5} \int \frac{(\sec(e+fx)a+a)^{5/2} (5c^2+3dc+d^2+3d(2c+d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}} d \sec(e + fx) - \frac{d \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)^{7/2}}{5a^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 90

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{1}{4} (20c^2 + 30cd + 13d^2) \int \frac{(\sec(e+fx)a+a)^{5/2}}{\sqrt{a-a \sec(e+fx)}} d \sec(e + fx) - \frac{3d(2c+d) \sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)^{7/2}}{4a^2} \right) \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{1}{4} (20c^2 + 30cd + 13d^2) \left(\frac{5}{3} a \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}} d \sec(e + fx) - \frac{\sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)^{5/2}}{3a} \right) \right) \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{1}{4} (20c^2 + 30cd + 13d^2) \left(\frac{5}{3} a \left(\frac{3}{2} a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}} d \sec(e + fx) - \frac{\sqrt{a-a \sec(e+fx)} (a \sec(e+fx)+a)^{3/2}}{2a} \right) \right) \right) \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 60

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{1}{4} (20c^2 + 30cd + 13d^2) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e + fx) - \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} \right) \right) \right) \right) \right)$$

↓ 45

3.203. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{1}{4} (20c^2 + 30cd + 13d^2) \right) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(2a \int \frac{1}{\frac{(a - a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a - a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} - \frac{\sqrt{a - a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a} \right) \right) \right)$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{1}{5} \left(\frac{1}{4} (20c^2 + 30cd + 13d^2) \right) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(-2 \arctan \left(\frac{\sqrt{a - a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}} \right) - \frac{\sqrt{a - a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a} \right) \right) \right)$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^2,x]`

output `-((a^2*(-1/5*(d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(7/2)*(c + d*Sec[e + f*x]))/a^2 + ((-3*d*(2*c + d)*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(7/2))/(4*a^2) + ((20*c^2 + 30*c*d + 13*d^2)*(-1/3*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2))/a + (5*a*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/a + (3*a*(-2*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/a))/2))/3))/4)/5)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))`

3.203.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

- rule 60 $\text{Int}[(a_.) + (b_.)(x_)^m)((c_.) + (d_.)(x_)^n], x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1}((c + d*x)^n/(b*(m+n+1))), x] + \text{Simp}[n*((b*c - a*d)/(b*(m+n+1))) \text{Int}[(a + b*x)^m(c + d*x)^{n-1}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m+n+1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m-n, 0]))) \ \&\& \ !\text{ILtQ}[m+n+2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 90 $\text{Int}[(a_.) + (b_.)(x_)((c_.) + (d_.)(x_)^n)((e_.) + (f_.)(x_)^p), x_] \rightarrow \text{Simp}[b*(c + d*x)^{n+1}((e + f*x)^{p+1}/(d*f*(n+p+2))), x] + \text{Simp}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(d*f*(n+p+2)) \text{Int}[(c + d*x)^n(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x \ \&\& \ \text{NeQ}[n+p+2, 0]$
- rule 101 $\text{Int}[(a_.) + (b_.)(x_)^2((c_.) + (d_.)(x_)^n)((e_.) + (f_.)(x_)^p), x_] \rightarrow \text{Simp}[b*(a + b*x)(c + d*x)^{n+1}((e + f*x)^{p+1}/(d*f*(n+p+3))), x] + \text{Simp}[1/(d*f*(n+p+3)) \text{Int}[(c + d*x)^n(e + f*x)^p \text{Simp}[a^2*d*f*(n+p+3) - b*(b*c*e + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(n+p+4) - b*(d*e*(n+2) + c*f*(p+2))]*x, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x \ \&\& \ \text{NeQ}[n+p+3, 0]$
- rule 218 $\text{Int}[(a_) + (b_)(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4475 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(g_.))^{p_.}(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.))^{m_.}(\text{csc}[(e_.) + (f_.)(x_)]*(d_.) + (c_.))^{n_.}], x_Symbol] \rightarrow \text{Simp}[a^2*g*(\text{Cot}[e + f*x]/(f*\text{Sqrt}[a + b*\text{Csc}[e + f*x]]*\text{Sqrt}[a - b*\text{Csc}[e + f*x]])) \text{Subst}[\text{Int}[(g*x)^{p-1}(a + b*x)^{m-1/2}((c + d*x)^n/\text{Sqrt}[a - b*x]), x], x, \text{Csc}[e + f*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, m, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ \text{IntegerQ}[m - 1/2])$

3.203.4 Maple [A] (verified)

Time = 4.54 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.00

method	result
norman	$-\frac{32a^3(20c^2+30cd+13d^2)\tan(\frac{fx}{2}+\frac{e}{2})^5}{15f} + \frac{7a^3(20c^2+30cd+13d^2)\tan(\frac{fx}{2}+\frac{e}{2})^7}{6f} - \frac{a^3(20c^2+30cd+13d^2)\tan(\frac{fx}{2}+\frac{e}{2})^9}{4f} - \frac{a^3(44c^2+30cd+13d^2)\tan(\frac{fx}{2}+\frac{e}{2})^{11}}{3f} - \frac{a^3(20c^2+30cd+13d^2)\tan(\frac{fx}{2}+\frac{e}{2})^{13}}{2f} + \frac{a^3(20c^2+30cd+13d^2)\tan(\frac{fx}{2}+\frac{e}{2})^{15}}{f}$
parts	$\frac{(2a^3cd+3a^3d^2)\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)}{f} + \frac{(3c^2a^3+2a^3cd)\tan(fx+e)}{f}$
parallelrisch	$26a^3\left(-\frac{75\left(\frac{\cos(5fx+5e)}{10}+\frac{\cos(3fx+3e)}{2}+\cos(fx+e)\right)\left(c^2+\frac{3}{2}cd+\frac{1}{20}d^2\right)\ln(\tan(\frac{fx}{2}+\frac{e}{2})-1)}{26} + \frac{75\left(\frac{\cos(5fx+5e)}{10}+\frac{\cos(3fx+3e)}{2}+\cos(fx+e)\right)\left(c^2+\frac{3}{2}cd+\frac{1}{20}d^2\right)\ln(\tan(\frac{fx}{2}+\frac{e}{2})+1)}{26}\right)$
derivativedivides	$-c^2a^3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+2a^3cd\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)$
default	$-c^2a^3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)+2a^3cd\left(-\left(-\frac{\sec(fx+e)^3}{4}-\frac{3\sec(fx+e)}{8}\right)\tan(fx+e)+\frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)$
risch	$-\frac{ia^3(-440c^2-4800cde^{4i(fx+e)}-3360cde^{2i(fx+e)}-450de^{i(fx+e)}c-720cd-304d^2-1520d^2e^{2i(fx+e)}-195d^2e^{i(fx+e)}-155d^2e^{-i(fx+e)}-155d^2e^{-2i(fx+e)})}{155f}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output (-32/15*a^3*(20*c^2+30*c*d+13*d^2)/f*tan(1/2*f*x+1/2*e)^5+7/6*a^3*(20*c^2+30*c*d+13*d^2)/f*tan(1/2*f*x+1/2*e)^7-1/4*a^3*(20*c^2+30*c*d+13*d^2)/f*tan(1/2*f*x+1/2*e)^9-1/4*a^3*(44*c^2+98*c*d+51*d^2)/f*tan(1/2*f*x+1/2*e)+1/6*a^3*(212*c^2+366*c*d+133*d^2)/f*tan(1/2*f*x+1/2*e)^3/(tan(1/2*f*x+1/2*e)^2-1)^5-1/8*a^3*(20*c^2+30*c*d+13*d^2)/f*ln(tan(1/2*f*x+1/2*e)-1)+1/8*a^3*(20*c^2+30*c*d+13*d^2)/f*ln(tan(1/2*f*x+1/2*e)+1)
```

3.203.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 245, normalized size of antiderivative = 0.95

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx = \frac{15(20a^3c^2 + 30a^3cd + 13a^3d^2)\cos(fx + e)^5 \log(\sin(fx + e) + 1) - 15(20a^3c^2 + 30a^3cd + 13a^3d^2)\cos(fx + e)^6 \log(\sin(fx + e) - 1)}{155f}$$

3.203. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^2,x, algorithm="f
ricas")`

output `1/240*(15*(20*a^3*c^2 + 30*a^3*c*d + 13*a^3*d^2)*cos(f*x + e)^5*log(sin(f*
x + e) + 1) - 15*(20*a^3*c^2 + 30*a^3*c*d + 13*a^3*d^2)*cos(f*x + e)^5*log
(-sin(f*x + e) + 1) + 2*(24*a^3*d^2 + 8*(55*a^3*c^2 + 90*a^3*c*d + 38*a^3*
d^2)*cos(f*x + e)^4 + 15*(12*a^3*c^2 + 30*a^3*c*d + 13*a^3*d^2)*cos(f*x +
e)^3 + 8*(5*a^3*c^2 + 30*a^3*c*d + 19*a^3*d^2)*cos(f*x + e)^2 + 30*(2*a^3*
c*d + 3*a^3*d^2)*cos(f*x + e))*sin(f*x + e))/(f*cos(f*x + e)^5)`

3.203.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx \\ &= a^3 \left(\int c^2 \sec(e + fx) dx + \int 3c^2 \sec^2(e + fx) dx + \int 3c^2 \sec^3(e + fx) dx \right. \\ & \quad + \int c^2 \sec^4(e + fx) dx + \int d^2 \sec^3(e + fx) dx + \int 3d^2 \sec^4(e + fx) dx \\ & \quad + \int 3d^2 \sec^5(e + fx) dx + \int d^2 \sec^6(e + fx) dx + \int 2cd \sec^2(e + fx) dx \\ & \quad \left. + \int 6cd \sec^3(e + fx) dx + \int 6cd \sec^4(e + fx) dx + \int 2cd \sec^5(e + fx) dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c+d*sec(f*x+e))**2,x)`

output `a**3*(Integral(c**2*sec(e + f*x), x) + Integral(3*c**2*sec(e + f*x)**2, x)
+ Integral(3*c**2*sec(e + f*x)**3, x) + Integral(c**2*sec(e + f*x)**4, x)
+ Integral(d**2*sec(e + f*x)**3, x) + Integral(3*d**2*sec(e + f*x)**4, x)
+ Integral(3*d**2*sec(e + f*x)**5, x) + Integral(d**2*sec(e + f*x)**6, x)
+ Integral(2*c*d*sec(e + f*x)**2, x) + Integral(6*c*d*sec(e + f*x)**3, x)
+ Integral(6*c*d*sec(e + f*x)**4, x) + Integral(2*c*d*sec(e + f*x)**5, x)
)`

3.203.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 459, normalized size of antiderivative = 1.79

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$$

$$= \frac{80 (\tan (fx + e)^3 + 3 \tan (fx + e)) a^3 c^2 + 480 (\tan (fx + e)^3 + 3 \tan (fx + e)) a^3 c d + 16 (3 \tan (fx + e))$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^2,x, algorithm="maxima")
```

```
output 1/240*(80*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c^2 + 480*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c*d + 16*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*a^3*d^2 + 240*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*d^2 - 30*a^3*c*d*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 45*a^3*d^2*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 180*a^3*c^2*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 360*a^3*c*d*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 60*a^3*d^2*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 240*a^3*c^2*log(sec(f*x + e) + tan(f*x + e)) + 720*a^3*c^2*tan(f*x + e) + 480*a^3*c*d*tan(f*x + e))/f
```

3.203.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.46

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$$

$$= \frac{15 (20 a^3 c^2 + 30 a^3 c d + 13 a^3 d^2) \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + 1 \right| \right) - 15 (20 a^3 c^2 + 30 a^3 c d + 13 a^3 d^2) \log \left(\left| \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) - 1 \right| \right)}{2}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e))^2,x, algorithm="giac")
```

```
output 1/120*(15*(20*a^3*c^2 + 30*a^3*c*d + 13*a^3*d^2)*log(abs(tan(1/2*f*x + 1/2
*e) + 1)) - 15*(20*a^3*c^2 + 30*a^3*c*d + 13*a^3*d^2)*log(abs(tan(1/2*f*x
+ 1/2*e) - 1)) - 2*(300*a^3*c^2*tan(1/2*f*x + 1/2*e)^9 + 450*a^3*c*d*tan(1
/2*f*x + 1/2*e)^9 + 195*a^3*d^2*tan(1/2*f*x + 1/2*e)^9 - 1400*a^3*c^2*tan(
1/2*f*x + 1/2*e)^7 - 2100*a^3*c*d*tan(1/2*f*x + 1/2*e)^7 - 910*a^3*d^2*tan
(1/2*f*x + 1/2*e)^7 + 2560*a^3*c^2*tan(1/2*f*x + 1/2*e)^5 + 3840*a^3*c*d*t
an(1/2*f*x + 1/2*e)^5 + 1664*a^3*d^2*tan(1/2*f*x + 1/2*e)^5 - 2120*a^3*c^2
*tan(1/2*f*x + 1/2*e)^3 - 3660*a^3*c*d*tan(1/2*f*x + 1/2*e)^3 - 1330*a^3*d
^2*tan(1/2*f*x + 1/2*e)^3 + 660*a^3*c^2*tan(1/2*f*x + 1/2*e) + 1470*a^3*c*
d*tan(1/2*f*x + 1/2*e) + 765*a^3*d^2*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x +
1/2*e)^2 - 1)^5)/f
```

3.203.9 Mupad [B] (verification not implemented)

Time = 17.10 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.12

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx))^2 dx$$

$$= \frac{a^3 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)(20c^2 + 30cd + 13d^2)}{2(10c^2 + 15cd + \frac{13d^2}{2})}\right)(20c^2 + 30cd + 13d^2)}{4f} - \frac{\left(5a^3c^2 + \frac{15a^3cd}{2} + \frac{13a^3d^2}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9 + \left(-\frac{70a^3c^2}{3} - 35a^3cd - \frac{91a^3d^2}{6}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7 + \left(\frac{128a^3c^2}{3} - \frac{128a^3cd}{3} - \frac{128a^3d^2}{3}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{10} - 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 + 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 10 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)}$$

```
input int(((a + a/cos(e + f*x))^3*(c + d/cos(e + f*x))^2)/cos(e + f*x),x)
```

```
output (a^3*atanh((tan(e/2 + (f*x)/2)*(30*c*d + 20*c^2 + 13*d^2))/(2*(15*c*d + 10
*c^2 + (13*d^2)/2)))*(30*c*d + 20*c^2 + 13*d^2))/(4*f) - (tan(e/2 + (f*x)/
2)*(11*a^3*c^2 + (51*a^3*d^2)/4 + (49*a^3*c*d)/2) + tan(e/2 + (f*x)/2)^9*(
5*a^3*c^2 + (13*a^3*d^2)/4 + (15*a^3*c*d)/2) - tan(e/2 + (f*x)/2)^7*((70*a
^3*c^2)/3 + (91*a^3*d^2)/6 + 35*a^3*c*d) - tan(e/2 + (f*x)/2)^3*((106*a^3*
c^2)/3 + (133*a^3*d^2)/6 + 61*a^3*c*d) + tan(e/2 + (f*x)/2)^5*((128*a^3*c
^2)/3 + (416*a^3*d^2)/15 + 64*a^3*c*d)/(f*(5*tan(e/2 + (f*x)/2)^2 - 10*tan
(e/2 + (f*x)/2)^4 + 10*tan(e/2 + (f*x)/2)^6 - 5*tan(e/2 + (f*x)/2)^8 + tan
(e/2 + (f*x)/2)^10 - 1))
```

3.204 $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx$

3.204.1 Optimal result	1434
3.204.2 Mathematica [A] (verified)	1435
3.204.3 Rubi [A] (verified)	1435
3.204.4 Maple [A] (verified)	1437
3.204.5 Fricas [A] (verification not implemented)	1438
3.204.6 Sympy [F]	1438
3.204.7 Maxima [B] (verification not implemented)	1439
3.204.8 Giac [A] (verification not implemented)	1439
3.204.9 Mupad [B] (verification not implemented)	1440

3.204.1 Optimal result

Integrand size = 29, antiderivative size = 125

$$\begin{aligned} & \int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx \\ &= \frac{5a^3(4c + 3d)\operatorname{arctanh}(\sin(e + fx))}{8f} + \frac{a^3(4c + 3d)\tan(e + fx)}{f} \\ & \quad + \frac{3a^3(4c + 3d)\sec(e + fx)\tan(e + fx)}{8f} \\ & \quad + \frac{d(a + a \sec(e + fx))^3 \tan(e + fx)}{4f} + \frac{a^3(4c + 3d)\tan^3(e + fx)}{12f} \end{aligned}$$

output `5/8*a^3*(4*c+3*d)*arctanh(sin(f*x+e))/f+a^3*(4*c+3*d)*tan(f*x+e)/f+3/8*a^3*(4*c+3*d)*sec(f*x+e)*tan(f*x+e)/f+1/4*d*(a+a*sec(f*x+e))^3*tan(f*x+e)/f+1/12*a^3*(4*c+3*d)*tan(f*x+e)^3/f`

3.204.2 Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.38

$$\int \sec(e+fx)(a+a\sec(e+fx))^3(c+d\sec(e+fx))dx$$

$$= \frac{5a^3c \operatorname{arctanh}(\sin(e+fx))}{2f} + \frac{15a^3d \operatorname{arctanh}(\sin(e+fx))}{8f} + \frac{4a^3c \tan(e+fx)}{f}$$

$$+ \frac{4a^3d \tan(e+fx)}{f} + \frac{3a^3c \sec(e+fx) \tan(e+fx)}{2f} + \frac{15a^3d \sec(e+fx) \tan(e+fx)}{8f}$$

$$+ \frac{a^3d \sec^3(e+fx) \tan(e+fx)}{4f} + \frac{a^3c \tan^3(e+fx)}{3f} + \frac{a^3d \tan^3(e+fx)}{f}$$

input `Integrate[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x]),x]`output `(5*a^3*c*ArcTanh[Sin[e + f*x]])/(2*f) + (15*a^3*d*ArcTanh[Sin[e + f*x]])/(8*f) + (4*a^3*c*Tan[e + f*x])/f + (4*a^3*d*Tan[e + f*x])/f + (3*a^3*c*Sec[e + f*x]*Tan[e + f*x])/(2*f) + (15*a^3*d*Sec[e + f*x]*Tan[e + f*x])/(8*f) + (a^3*d*Sec[e + f*x]^3*Tan[e + f*x])/(4*f) + (a^3*c*Tan[e + f*x]^3)/(3*f) + (a^3*d*Tan[e + f*x]^3)/f`**3.204.3 Rubi [A] (verified)**Time = 0.41 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.88, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.172$, Rules used = {3042, 4489, 3042, 4278, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e+fx)(a\sec(e+fx)+a)^3(c+d\sec(e+fx))dx$$

$$\downarrow 3042$$

$$\int \csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^3\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)dx$$

$$\downarrow 4489$$

$$\frac{1}{4}(4c+3d)\int \sec(e+fx)(\sec(e+fx)a+a)^3dx + \frac{d \tan(e+fx)(a \sec(e+fx)+a)^3}{4f}$$

$$\downarrow 3042$$

3.204. $\int \sec(e+fx)(a+a\sec(e+fx))^3(c+d\sec(e+fx))dx$

$$\frac{1}{4}(4c+3d) \int \csc\left(e+fx+\frac{\pi}{2}\right) \left(\csc\left(e+fx+\frac{\pi}{2}\right)a+a\right)^3 dx + \frac{d \tan(e+fx)(a \sec(e+fx)+a)^3}{4f}$$

↓ 4278

$$\frac{1}{4}(4c+3d) \int (a^3 \sec^4(e+fx) + 3a^3 \sec^3(e+fx) + 3a^3 \sec^2(e+fx) + a^3 \sec(e+fx)) dx + \frac{d \tan(e+fx)(a \sec(e+fx)+a)^3}{4f}$$

↓ 2009

$$3d) \left(\frac{5a^3 \operatorname{arctanh}(\sin(e+fx))}{2f} + \frac{a^3 \tan^3(e+fx)}{3f} + \frac{4a^3 \tan(e+fx)}{f} + \frac{3a^3 \tan(e+fx) \sec(e+fx)}{2f} \right) + \frac{d \tan(e+fx)(a \sec(e+fx)+a)^3}{4f}$$

input `Int[Sec[e + f*x]*(a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x]),x]`

output `(d*(a + a*Sec[e + f*x])^3*Tan[e + f*x])/(4*f) + ((4*c + 3*d)*((5*a^3*ArcTanh[Sin[e + f*x]])/(2*f) + (4*a^3*Tan[e + f*x])/f + (3*a^3*Sec[e + f*x]*Tan[e + f*x])/(2*f) + (a^3*Tan[e + f*x]^3)/(3*f)))/4`

3.204.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4278 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n_.*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m_.], x_Symbol] := Int[ExpandTrig[(a + b*csc[e + f*x])^m*(d*csc[e + f*x])^n, x], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && I GtQ[m, 0] && RationalQ[n]`

```
rule 4489 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*B*m + A*b*(m + 1))/(b*(m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^m, x], x] /; FreeQ[{a, b, A, B, e, f, m}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && NeQ[a*B*m + A*b*(m + 1), 0] && !LtQ[m, -2^(-1)]
```

3.204.4 Maple [A] (verified)

Time = 3.98 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.40

method	result
norman	$-\frac{73a^3(4c+3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{12f} + \frac{55a^3(4c+3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{12f} - \frac{5a^3(4c+3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{4f} + \frac{a^3(49d+44c)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{4f} - \frac{5a^3(4c+3d)}{12f}$
parallelrisc	$26 \left(-\frac{15\left(\frac{3}{4} + \frac{\cos(4fx+4e)}{4} + \cos(2fx+2e)\right)\left(c + \frac{3d}{4}\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)}{13} + \frac{15\left(\frac{3}{4} + \frac{\cos(4fx+4e)}{4} + \cos(2fx+2e)\right)\left(c + \frac{3d}{4}\right)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{13} \right)$
parts	$-\frac{(a^3c+3a^3d)\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right)\tan(fx+e)}{f} + \frac{(3a^3c+a^3d)\tan(fx+e)}{f} + \frac{(3a^3c+3a^3d)\left(\frac{\sec(fx+e)\tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{8}\right)}{f}$
derivativedivides	$-a^3c\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + a^3d\left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8}\right)\tan(fx+e) + \frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right) + 3a^3d$
default	$-a^3c\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right)\tan(fx+e) + a^3d\left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8}\right)\tan(fx+e) + \frac{3\ln(\sec(fx+e)+\tan(fx+e))}{8}\right) + 3a^3d$
risc	$-\frac{ia^3(36ce^{7i(fx+e)}+45de^{7i(fx+e)}-72ce^{6i(fx+e)}-24de^{6i(fx+e)}+36ce^{5i(fx+e)}+69de^{5i(fx+e)}-264ce^{4i(fx+e)}-216de^{4i(fx+e)}+12f(1+e^{2i(fx+e)}))}{12f(1+e^{2i(fx+e)})}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output (-73/12*a^3*(4*c+3*d)/f*tan(1/2*f*x+1/2*e)^3+55/12*a^3*(4*c+3*d)/f*tan(1/2*f*x+1/2*e)^5-5/4*a^3*(4*c+3*d)/f*tan(1/2*f*x+1/2*e)^7+1/4*a^3*(49*d+44*c)/f*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2-1)^4-5/8*a^3*(4*c+3*d)/f*ln(tan(1/2*f*x+1/2*e)-1)+5/8*a^3*(4*c+3*d)/f*ln(tan(1/2*f*x+1/2*e)+1)
```

3.204. $\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx$

3.204.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.29

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx$$

$$= \frac{15(4a^3c + 3a^3d) \cos(fx + e)^4 \log(\sin(fx + e) + 1) - 15(4a^3c + 3a^3d) \cos(fx + e)^4 \log(-\sin(fx + e))}{f \cos(fx + e)^4}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e)),x, algorithm="fricas")
```

```
output 1/48*(15*(4*a^3*c + 3*a^3*d)*cos(f*x + e)^4*log(sin(f*x + e) + 1) - 15*(4*a^3*c + 3*a^3*d)*cos(f*x + e)^4*log(-sin(f*x + e) + 1) + 2*(6*a^3*d + 8*(11*a^3*c + 9*a^3*d)*cos(f*x + e)^3 + 9*(4*a^3*c + 5*a^3*d)*cos(f*x + e)^2 + 8*(a^3*c + 3*a^3*d)*cos(f*x + e))*sin(f*x + e))/(f*cos(f*x + e)^4)
```

3.204.6 Sympy [F]

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx$$

$$= a^3 \left(\int c \sec(e + fx) dx + \int 3c \sec^2(e + fx) dx + \int 3c \sec^3(e + fx) dx \right.$$

$$\left. + \int c \sec^4(e + fx) dx + \int d \sec^2(e + fx) dx + \int 3d \sec^3(e + fx) dx \right.$$

$$\left. + \int 3d \sec^4(e + fx) dx + \int d \sec^5(e + fx) dx \right)$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3*(c+d*sec(f*x+e)),x)
```

```
output a**3*(Integral(c*sec(e + f*x), x) + Integral(3*c*sec(e + f*x)**2, x) + Integral(3*c*sec(e + f*x)**3, x) + Integral(c*sec(e + f*x)**4, x) + Integral(d*sec(e + f*x)**2, x) + Integral(3*d*sec(e + f*x)**3, x) + Integral(3*d*sec(e + f*x)**4, x) + Integral(d*sec(e + f*x)**5, x))
```

3.204.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 262 vs. $2(117) = 234$.

Time = 0.23 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.10

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx$$

$$= \frac{16 (\tan (fx + e))^3 + 3 \tan (fx + e) a^3 c + 48 (\tan (fx + e))^3 + 3 \tan (fx + e) a^3 d - 3 a^3 d \left(\frac{2 (3 \sin (fx + e))^3 - 5}{\sin (fx + e)^4 - 2 \sin} \right)}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `1/48*(16*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*c + 48*(tan(f*x + e)^3 + 3*tan(f*x + e))*a^3*d - 3*a^3*d*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 36*a^3*c*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 36*a^3*d*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 48*a^3*c*log(sec(f*x + e) + tan(f*x + e)) + 144*a^3*c*tan(f*x + e) + 48*a^3*d*tan(f*x + e))/f`

3.204.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.70

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx$$

$$= \frac{15 (4 a^3 c + 3 a^3 d) \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + 1 \right| \right) - 15 (4 a^3 c + 3 a^3 d) \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 (60 a^3 c \tan (fx + e) + 3 a^3 d \tan (fx + e))}{\sin (fx + e)^4 - 2 \sin (fx + e)^2 + 1}}{f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3*(c+d*sec(f*x+e)),x, algorithm="giac")`


```
output 1/24*(15*(4*a^3*c + 3*a^3*d)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*(4*a^
3*c + 3*a^3*d)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(60*a^3*c*tan(1/2*f*
x + 1/2*e)^7 + 45*a^3*d*tan(1/2*f*x + 1/2*e)^7 - 220*a^3*c*tan(1/2*f*x + 1
/2*e)^5 - 165*a^3*d*tan(1/2*f*x + 1/2*e)^5 + 292*a^3*c*tan(1/2*f*x + 1/2*e
)^3 + 219*a^3*d*tan(1/2*f*x + 1/2*e)^3 - 132*a^3*c*tan(1/2*f*x + 1/2*e) -
147*a^3*d*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^4)/f
```

3.204.9 Mupad [B] (verification not implemented)

Time = 16.88 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.62

$$\int \sec(e + fx)(a + a \sec(e + fx))^3(c + d \sec(e + fx)) dx$$

$$= \frac{\left(-5a^3c - \frac{15a^3d}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7 + \left(\frac{55a^3c}{3} + \frac{55a^3d}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-\frac{73a^3c}{3} - \frac{73a^3d}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + \left(\frac{11a^3c}{3} + \frac{11a^3d}{4}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right) + 1}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^8 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 1\right)} + \frac{5a^3 \operatorname{atanh}\left(\frac{5 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)(4c + 3d)}{2(10c + \frac{15d}{2})}\right)(4c + 3d)}{4f}$$

```
input int(((a + a/cos(e + f*x))^3*(c + d/cos(e + f*x)))/cos(e + f*x),x)
```

```
output (tan(e/2 + (f*x)/2)*(11*a^3*c + (49*a^3*d)/4) - tan(e/2 + (f*x)/2)^7*(5*a^
3*c + (15*a^3*d)/4) + tan(e/2 + (f*x)/2)^5*((55*a^3*c)/3 + (55*a^3*d)/4) -
tan(e/2 + (f*x)/2)^3*((73*a^3*c)/3 + (73*a^3*d)/4))/(f*(6*tan(e/2 + (f*x)
/2)^4 - 4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/
2)^8 + 1)) + (5*a^3*atanh((5*tan(e/2 + (f*x)/2)*(4*c + 3*d))/(2*(10*c + (1
5*d)/2)))*(4*c + 3*d))/(4*f)
```

3.205 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c+d \sec(e+fx)} dx$

3.205.1 Optimal result 1441
 3.205.2 Mathematica [C] (warning: unable to verify) 1442
 3.205.3 Rubi [A] (verified) 1442
 3.205.4 Maple [A] (verified) 1447
 3.205.5 Fricas [A] (verification not implemented) 1448
 3.205.6 Sympy [F] 1449
 3.205.7 Maxima [F(-2)] 1449
 3.205.8 Giac [B] (verification not implemented) 1450
 3.205.9 Mupad [B] (verification not implemented) 1450

3.205.1 Optimal result

Integrand size = 31, antiderivative size = 153

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c+d \sec(e+fx)} dx = \frac{a^3 \operatorname{arctanh}(\sin(e+fx))}{2df} + \frac{a^3(c^2-3cd+3d^2) \operatorname{arctanh}(\sin(e+fx))}{d^3 f} - \frac{2a^3(c-d)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{d^3 \sqrt{c+d} f} - \frac{a^3(c-3d) \tan(e+fx)}{d^2 f} + \frac{a^3 \sec(e+fx) \tan(e+fx)}{2df}$$

output

```
1/2*a^3*arctanh(sin(f*x+e))/d/f+a^3*(c^2-3*c*d+3*d^2)*arctanh(sin(f*x+e))/d^3/f-2*a^3*(c-d)^(5/2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/d^3/f/(c+d)^(1/2)-a^3*(c-3*d)*tan(f*x+e)/d^2/f+1/2*a^3*sec(f*x+e)*tan(f*x+e)/d/f
```

3.205.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 3.38 (sec) , antiderivative size = 419, normalized size of antiderivative = 2.74

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx$$

$$= \frac{a^3 \cos^2(e+fx)(d+c\cos(e+fx)) \sec^6\left(\frac{1}{2}(e+fx)\right) (1+\sec(e+fx))^3 \left(-2(2c^2-6cd+7d^2) \log\left(\cos\left(\frac{1}{2}(e+fx)\right)\right)\right)}{c+d\sec(e+fx)}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x]),x]`

output `(a^3*Cos[e + f*x]^2*(d + c*Cos[e + f*x])*Sec[(e + f*x)/2]^6*(1 + Sec[e + f*x])^3*(-2*(2*c^2 - 6*c*d + 7*d^2)*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + 2*(2*c^2 - 6*c*d + 7*d^2)*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] + (8*(c - d)^3*ArcTan[((I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))]/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]))*(I*Cos[e] + Sin[e]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + d^2/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2 - (4*(c - 3*d)*d*Sin[(f*x)/2])/((Cos[e/2] - Sin[e/2])*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])) - d^2/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])^2 - (4*(c - 3*d)*d*Sin[(f*x)/2])/((Cos[e/2] + Sin[e/2])*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])))/(32*d^3*f*(c + d*Sec[e + f*x]))`

3.205.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.73, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4475, 113, 25, 27, 171, 25, 27, 175, 45, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{c+d\sec(e+fx)} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^3}{c+d\csc\left(e+fx+\frac{\pi}{2}\right)} dx$$

3.205. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx$

$$\begin{aligned} & \downarrow 4475 \\ & \frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{5/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 113 \\ & \frac{a^2 \tan(e+fx) \left(-\frac{\int -\frac{a^3 \sqrt{\sec(e+fx)a+a}(c+2d-(2c-5d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))} d \sec(e+fx)}{2ad} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 25 \\ & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^3 \sqrt{\sec(e+fx)a+a}(c+2d-(2c-5d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))} d \sec(e+fx)}{2ad} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 27 \\ & \frac{a^2 \tan(e+fx) \left(\frac{a^2 \int \frac{\sqrt{\sec(e+fx)a+a}(c+2d-(2c-5d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))} d \sec(e+fx)}{2d} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 171 \\ & \frac{a^2 \tan(e+fx) \left(\frac{a^2 \left(\frac{(2c-5d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{ad} - \frac{\int -\frac{a^2 (d(c+2d)+(2c^2-6dc+7d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{ad} \right)}{2d} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 25 \\ & \frac{a^2 \tan(e+fx) \left(\frac{a^2 \left(\frac{\int \frac{a^2 (d(c+2d)+(2c^2-6dc+7d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{ad} + \frac{(2c-5d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{ad} \right)}{2d} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 27 \end{aligned}$$

3.205. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c+d \sec(e+fx)} dx$

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{d(c+2d) + (2c^2 - 6dc + 7d^2) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}} \frac{d \sec(e+fx)}{d} + \frac{(2c-5d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{ad} \right)}{2d} - \sqrt{a-a \sec(e+fx)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 175

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{(2c^2 - 6cd + 7d^2) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{d} - \frac{2(c-d)^3 \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}}}{d} \right)}{2d} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{a^2 \left(\frac{2(2c^2 - 6cd + 7d^2) \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}}{d} - \frac{2(c-d)^3 \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}}}{d} \right)}{2d} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$$

↓ 104

3.205. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{c+d \sec(e+fx)} dx$

$$a^2 \tan(e + fx) \left(\frac{a \left(\frac{2(2c^2 - 6cd + 7d^2) \int \frac{1}{\frac{(a - a \sec(e + fx))a}{\sec(e + fx)a + a} - a} \frac{d \sqrt{a - a \sec(e + fx)}}{\sqrt{\sec(e + fx)a + a}} - \frac{4(c - d)^3 \int \frac{1}{a(c - d) + \frac{a(c + d)(\sec(e + fx)a + a)}{a - a \sec(e + fx)}} \frac{d \sqrt{\sec(e + fx)a + a}}{\sqrt{a - a \sec(e + fx)}}}{d} \right)}{2d} \right)$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{a \left(\frac{2(2c^2 - 6cd + 7d^2) \arctan\left(\frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{a \sec(e + fx) + a}}\right)}{ad} - \frac{4(c - d)^{5/2} \arctan\left(\frac{\sqrt{c + d} \sqrt{a \sec(e + fx) + a}}{\sqrt{c - d} \sqrt{a - a \sec(e + fx)}}\right)}{ad \sqrt{c + d}} \right)}{2d} + \frac{(2c - 5d) \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{ad} \right)$$

```
input Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x]),x]
```

```
output -((a^2*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/d + (a^2*((a*((-2*(2*c^2 - 6*c*d + 7*d^2)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])/(a*d) - (4*(c - d)^(5/2)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]/(a*d*Sqrt[c + d])))/d + ((2*c - 5*d)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a*d)))/(2*d))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.205.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 113 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegersQ[2*m, 2*n, 2*p]`
- rule 171 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 2)), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegersQ[2*m, 2*n, 2*p]`

```
rule 175 Int[(((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)*((g_) + (h_)*(x_))) / ((a_) + (b_)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4475 Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])
```

3.205.4 Maple [A] (verified)

Time = 0.87 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.46

method	result
derivativedivides	$16a^3 \left(-\frac{(c^3 - 3c^2d + 3cd^2 - d^3) \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{8d^3\sqrt{(c+d)(c-d)}} + \frac{1}{32d\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^2} - \frac{5d-2c}{32d^2\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{(-2c^2 + 6cd - 7d^2)}{f} \right)$
default	$16a^3 \left(-\frac{(c^3 - 3c^2d + 3cd^2 - d^3) \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{8d^3\sqrt{(c+d)(c-d)}} + \frac{1}{32d\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^2} - \frac{5d-2c}{32d^2\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{(-2c^2 + 6cd - 7d^2)}{f} \right)$
risch	$-\frac{ia^3(d e^{3i(fx+e)} + 2e^{2i(fx+e)}c - 6de^{2i(fx+e)} - de^{i(fx+e)} + 2c - 6d)}{f d^2(1 + e^{2i(fx+e)})^2} + \frac{\sqrt{(c+d)(c-d)} a^3 \ln\left(\frac{e^{i(fx+e)} - i\sqrt{(c+d)(c-d)} - d}{c}\right)}{(c+d)f d^3}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)), x, method=_RETURNVERBOSE)
```

3.205. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx$


```
output 16/f*a^3*(-1/8*(c^3-3*c^2*d+3*c*d^2-d^3)/d^3/((c+d)*(c-d))^(1/2)*arctanh((
c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))+1/32/d/(tan(1/2*f*x+1/2*e)-1)
^2-1/32*(5*d-2*c)/d^2/(tan(1/2*f*x+1/2*e)-1)+1/32/d^3*(-2*c^2+6*c*d-7*d^2)
*ln(tan(1/2*f*x+1/2*e)-1)-1/32/d/(tan(1/2*f*x+1/2*e)+1)^2-1/32*(5*d-2*c)/d
^2/(tan(1/2*f*x+1/2*e)+1)+1/32*(2*c^2-6*c*d+7*d^2)/d^3*ln(tan(1/2*f*x+1/2*
e)+1))
```

3.205.5 Fracas [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 532, normalized size of antiderivative = 3.48

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx$$

$$= \frac{2(a^3c^2 - 2a^3cd + a^3d^2)\sqrt{\frac{c-d}{c+d}} \cos(fx+e)^2 \log\left(\frac{2cd\cos(fx+e) - (c^2-2d^2)\cos(fx+e)^2 - 2(c^2+cd+(cd+d^2)\cos(fx+e))\sqrt{\frac{c-d}{c+d}}}{c^2\cos(fx+e)^2 + 2cd\cos(fx+e) + d^2}\right)}{4(a^3c^2 - 2a^3cd + a^3d^2)\sqrt{\frac{c-d}{c+d}} \arctan\left(-\frac{(d\cos(fx+e)+c)\sqrt{\frac{c-d}{c+d}}}{(c-d)\sin(fx+e)}\right) \cos(fx+e)^2 - (2a^3c^2 - 6a^3cd + 7a^3d^2)}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)),x, algorithm="fri
cas")
```

```
output [1/4*(2*(a^3*c^2 - 2*a^3*c*d + a^3*d^2)*sqrt((c - d)/(c + d))*cos(f*x + e)
^2*log(((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*(c^2 + c*d +
(c*d + d^2)*cos(f*x + e))*sqrt((c - d)/(c + d))*sin(f*x + e) + 2*c^2 - d
^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + (2*a^3*c^2 - 6*a^3*c
*d + 7*a^3*d^2)*cos(f*x + e)^2*log(sin(f*x + e) + 1) - (2*a^3*c^2 - 6*a^3*
c*d + 7*a^3*d^2)*cos(f*x + e)^2*log(-sin(f*x + e) + 1) + 2*(a^3*d^2 - 2*(a
^3*c*d - 3*a^3*d^2)*cos(f*x + e))*sin(f*x + e))/(d^3*f*cos(f*x + e)^2), -1
/4*(4*(a^3*c^2 - 2*a^3*c*d + a^3*d^2)*sqrt(-(c - d)/(c + d))*arctan(-(d*co
s(f*x + e) + c)*sqrt(-(c - d)/(c + d))/((c - d)*sin(f*x + e)))*cos(f*x + e
)^2 - (2*a^3*c^2 - 6*a^3*c*d + 7*a^3*d^2)*cos(f*x + e)^2*log(sin(f*x + e)
+ 1) + (2*a^3*c^2 - 6*a^3*c*d + 7*a^3*d^2)*cos(f*x + e)^2*log(-sin(f*x + e)
+ 1) - 2*(a^3*d^2 - 2*(a^3*c*d - 3*a^3*d^2)*cos(f*x + e))*sin(f*x + e))/
(d^3*f*cos(f*x + e)^2)]
```

3.205.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx = a^3 \left(\int \frac{\sec(e+fx)}{c+d\sec(e+fx)} dx + \int \frac{3\sec^2(e+fx)}{c+d\sec(e+fx)} dx + \int \frac{3\sec^3(e+fx)}{c+d\sec(e+fx)} dx + \int \frac{\sec^4(e+fx)}{c+d\sec(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e)),x)`

output `a**3*(Integral(sec(e + f*x)/(c + d*sec(e + f*x)), x) + Integral(3*sec(e + f*x)**2/(c + d*sec(e + f*x)), x) + Integral(3*sec(e + f*x)**3/(c + d*sec(e + f*x)), x) + Integral(sec(e + f*x)**4/(c + d*sec(e + f*x)), x))`

3.205.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.205.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 285 vs. $2(140) = 280$.

Time = 0.37 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.86

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx$$

$$= \frac{(2a^3c^2-6a^3cd+7a^3d^2)\log(|\tan(\frac{1}{2}fx+\frac{1}{2}e)+1|)}{d^3} - \frac{(2a^3c^2-6a^3cd+7a^3d^2)\log(|\tan(\frac{1}{2}fx+\frac{1}{2}e)-1|)}{d^3} + \frac{4(a^3c^3-3a^3c^2d+3a^3cd^2-a^3d^3)}{d^3} \left(\frac{\arctan\left(\frac{c\tan(\frac{1}{2}fx+\frac{1}{2}e)-d}{\sqrt{-c^2+d^2}}\right)}{\sqrt{-c^2+d^2}} + \frac{\arctan\left(\frac{c\tan(\frac{1}{2}fx+\frac{1}{2}e)+d}{\sqrt{-c^2+d^2}}\right)}{\sqrt{-c^2+d^2}} \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)),x, algorithm="giac")`

output `1/2*((2*a^3*c^2 - 6*a^3*c*d + 7*a^3*d^2)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/d^3 - (2*a^3*c^2 - 6*a^3*c*d + 7*a^3*d^2)*log(abs(tan(1/2*f*x + 1/2*e) - 1))/d^3 + 4*(a^3*c^3 - 3*a^3*c^2*d + 3*a^3*c*d^2 - a^3*d^3)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/(sqrt(-c^2 + d^2)*d^3) + 2*(2*a^3*c*tan(1/2*f*x + 1/2*e)^3 - 5*a^3*d*tan(1/2*f*x + 1/2*e)^3 - 2*a^3*c*tan(1/2*f*x + 1/2*e) + 7*a^3*d*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1/2*e)^2 - 1)^2*d^2))/f`

3.205.9 Mupad [B] (verification not implemented)

Time = 14.36 (sec) , antiderivative size = 1902, normalized size of antiderivative = 12.43

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{c+d\sec(e+fx)} dx = \text{Too large to display}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c + d/cos(e + f*x))),x)`

output

$$\begin{aligned} & (\operatorname{atanh}((18824a^9c^2\tan(e/2 + (f*x)/2))/(18824a^9c^2 + 2968a^9d^2 - \\ & (16680a^9c^3)/d + (8608a^9c^4)/d^2 - (2480a^9c^5)/d^3 + (320a^9c^6)/d^4 - 11560a^9c*d) - \\ & (16680a^9c^3\tan(e/2 + (f*x)/2))/(2968a^9d^3 - 16680a^9c^3 - 11560a^9c*d^2 + 18824a^9c^2*d + (8608a^9c^4)/d - \\ & (2480a^9c^5)/d^2 + (320a^9c^6)/d^3) + (8608a^9c^4\tan(e/2 + (f*x)/2)) \\ & /((8608a^9c^4 + 2968a^9d^4 - 11560a^9c*d^3 - 16680a^9c^3*d + 18824a^9c^2*d^2 - \\ & (2480a^9c^5)/d + (320a^9c^6)/d^2) - (2480a^9c^5\tan(e/2 + (f*x)/2))/(2968a^9d^5 - \\ & 2480a^9c^5 - 11560a^9c*d^4 + 8608a^9c^4*d + 18824a^9c^2*d^3 - 16680a^9c^3*d^2 + \\ & (320a^9c^6)/d) + (320a^9c^6\tan(e/2 + (f*x)/2))/(320a^9c^6 + 2968a^9d^6 - 11560a^9c*d^5 - \\ & 2480a^9c^5*d + 18824a^9c^2*d^4 - 16680a^9c^3*d^3 + 8608a^9c^4*d^2) + \\ & (2968a^9d^2\tan(e/2 + (f*x)/2))/(18824a^9c^2 + 2968a^9d^2 - (16680a^9c^3)/d + \\ & (8608a^9c^4)/d^2 - (2480a^9c^5)/d^3 + (320a^9c^6)/d^4 - 11560a^9c*d) - \\ & (11560a^9c*d\tan(e/2 + (f*x)/2))/(18824a^9c^2 + 2968a^9d^2 - (16680a^9c^3)/d + \\ & (8608a^9c^4)/d^2 - (2480a^9c^5)/d^3 + (320a^9c^6)/d^4 - 11560a^9c*d)) * (2a^3c^2 + 7a^3d^2 - 6a^3c*d) / (d^3f - \\ & ((\tan(e/2 + (f*x)/2)*(2a^3c - 7a^3d))/d^2 - (a^3\tan(e/2 + (f*x)/2)^3*(2c - 5d))/d^2) / \\ & (f*(\tan(e/2 + (f*x)/2)^4 - 2\tan(e/2 + (f*x)/2)^2 + 1)) - (a^3*\operatorname{atan}(((a^3*((c + d)*(c - d)^5)^{(1/2)}*((8*\tan(e/2 + (f*x)/2) \\ & *(8a^6c^7 - 53a^6d^7 + 259a^6c*d^6 - 64a^6c^6*d - 547a^6c^2*d\dots \end{aligned}$$

3.206 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^2} dx$

3.206.1 Optimal result 1452
 3.206.2 Mathematica [C] (warning: unable to verify) 1453
 3.206.3 Rubi [A] (verified) 1453
 3.206.4 Maple [A] (verified) 1458
 3.206.5 Fricas [B] (verification not implemented) 1459
 3.206.6 Sympy [F] 1460
 3.206.7 Maxima [F(-2)] 1461
 3.206.8 Giac [B] (verification not implemented) 1461
 3.206.9 Mupad [B] (verification not implemented) 1462

3.206.1 Optimal result

Integrand size = 31, antiderivative size = 161

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^2} dx = -\frac{a^3(2c-3d)\operatorname{arctanh}(\sin(e+fx))}{d^3 f} + \frac{2a^3(c-d)^{3/2}(2c+3d)\operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{d^3(c+d)^{3/2}f} + \frac{2a^3c \tan(e+fx)}{d^2(c+d)f} - \frac{(c-d)(a^3+a^3 \sec(e+fx)) \tan(e+fx)}{d(c+d)f(c+d \sec(e+fx))}$$

```
output -a^3*(2*c-3*d)*arctanh(sin(f*x+e))/d^3/f+2*a^3*(c-d)^(3/2)*(2*c+3*d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/d^3/(c+d)^(3/2)/f+2*a^3*c*tan(f*x+e)/d^2/(c+d)/f-(c-d)*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/d/(c+d)/f/(c+d*sec(f*x+e))
```

3.206.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 5.18 (sec) , antiderivative size = 455, normalized size of antiderivative = 2.83

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^2} dx$$

$$= \frac{a^3 \cos(e + fx)(d + c \cos(e + fx)) \sec^6\left(\frac{1}{2}(e + fx)\right) (1 + \sec(e + fx))^3 \left((2c - 3d)(d + c \cos(e + fx)) \log \left(\frac{\cos\left(\frac{e + fx}{2}\right) - \sin\left(\frac{e + fx}{2}\right)}{\cos\left(\frac{e + fx}{2}\right) + \sin\left(\frac{e + fx}{2}\right)} \right) + (-2c + 3d)(d + c \cos(e + fx)) \log\left(\frac{\cos\left(\frac{e + fx}{2}\right) + \sin\left(\frac{e + fx}{2}\right)}{\cos\left(\frac{e + fx}{2}\right) - \sin\left(\frac{e + fx}{2}\right)}\right) - ((2I)(c - d)^2(2c + 3d) \operatorname{ArcTan}\left[\frac{(I \cos[e] + \sin[e])(c \sin[e] + (-d + c \cos[e]) \tan[(fx)/2])}{\sqrt{c^2 - d^2} \sqrt{(\cos[e] - I \sin[e])^2}}\right]) \right)}{(c + d) \sqrt{c^2 - d^2} \sqrt{(\cos[e] - I \sin[e])^2} + ((c - d)^2 d^* (-d \sin[e] + c \sin[fx])) / (c(c + d)(\cos[e/2] - \sin[e/2])(\cos[e/2] + \sin[e/2])) + (d(d + c \cos[e + fx]) \sin[(fx)/2]) / ((\cos[e/2] - \sin[e/2])(\cos[(e + fx)/2] - \sin[(e + fx)/2])) + (d(d + c \cos[e + fx]) \sin[(fx)/2]) / ((\cos[e/2] + \sin[e/2])(\cos[(e + fx)/2] + \sin[(e + fx)/2])))) / (8d^3 f (c + d \sec[e + fx])^2}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^2,x]`

output `(a^3*Cos[e + f*x]*(d + c*Cos[e + f*x])*Sec[(e + f*x)/2]^6*(1 + Sec[e + f*x])^3*((2*c - 3*d)*(d + c*Cos[e + f*x])*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + (-2*c + 3*d)*(d + c*Cos[e + f*x])*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] - ((2*I)*(c - d)^2*(2*c + 3*d)*ArcTan[((I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2])])*(d + c*Cos[e + f*x])*(Cos[e] - I*Sin[e]))/((c + d)*Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + ((c - d)^2*d*(-(d*Sin[e]) + c*Sin[fx]))/(c*(c + d)*(Cos[e/2] - Sin[e/2])*(Cos[e/2] + Sin[e/2])) + (d*(d + c*Cos[e + f*x])*Sin[(f*x)/2])/((Cos[e/2] - Sin[e/2])*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])) + (d*(d + c*Cos[e + f*x])*Sin[(f*x)/2])/((Cos[e/2] + Sin[e/2])*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])))))/(8*d^3*f*(c + d*Sec[e + f*x])^2)`

3.206.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.76, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4475, 109, 27, 171, 25, 27, 175, 45, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(a \sec(e + fx) + a)^3}{(c + d \sec(e + fx))^2} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right) \left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3}{\left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2} dx$$

↓ 4475

$$\frac{a^2 \tan(e + fx) \int \frac{(\sec(e + fx)a + a)^{5/2}}{\sqrt{a - a \sec(e + fx)}(c + d \sec(e + fx))^2} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 109

$$\frac{a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a - a \sec(e + fx)}(a \sec(e + fx) + a)^{3/2}}{d(c+d)(c + d \sec(e + fx))} - \frac{\int \frac{a^3 \sqrt{\sec(e + fx)a + a}(-2 \sec(e + fx)c + c - 3d)}{\sqrt{a - a \sec(e + fx)}(c + d \sec(e + fx))} d \sec(e + fx)}{ad(c+d)} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 27

$$\frac{a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a - a \sec(e + fx)}(a \sec(e + fx) + a)^{3/2}}{d(c+d)(c + d \sec(e + fx))} - \frac{a^2 \int \frac{\sqrt{\sec(e + fx)a + a}(-2 \sec(e + fx)c + c - 3d)}{\sqrt{a - a \sec(e + fx)}(c + d \sec(e + fx))} d \sec(e + fx)}{d(c+d)} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 171

$$\frac{a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a - a \sec(e + fx)}(a \sec(e + fx) + a)^{3/2}}{d(c+d)(c + d \sec(e + fx))} - \frac{a^2 \left(\frac{2c\sqrt{a - a \sec(e + fx)}\sqrt{a \sec(e + fx) + a}}{ad} - \frac{\int \frac{a^2((c-3d)d + (2c-3d)(c+d)\sec(e + fx))}{\sqrt{a - a \sec(e + fx)}\sqrt{\sec(e + fx)a + a}} d \sec(e + fx)}{ad} \right)}{d(c+d)} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 25

$$\frac{a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a - a \sec(e + fx)}(a \sec(e + fx) + a)^{3/2}}{d(c+d)(c + d \sec(e + fx))} - \frac{a^2 \left(\frac{\int \frac{a^2((c-3d)d + (2c-3d)(c+d)\sec(e + fx))}{\sqrt{a - a \sec(e + fx)}\sqrt{\sec(e + fx)a + a} + 2c\sqrt{a - a \sec(e + fx)}}{ad} d \sec(e + fx)}{d(c+d)} \right)}{d(c+d)} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 27

3.206. $\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^2} dx$

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{d(c+d)(c+d \sec(e+fx))} - \frac{a^2 \left(\frac{a \int \frac{(c-3d)d+(2c-3d)(c+d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{d} + 2c\sqrt{a-a} \right)}{d(c+d)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 175

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{d(c+d)(c+d \sec(e+fx))} - \frac{a^2 \left(\frac{(2c-3d)(c+d) \int \frac{1}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{d} - \frac{(c-d)^2(2c-d)}{d} \right)}{d} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{d(c+d)(c+d \sec(e+fx))} - \frac{a^2 \left(\frac{2(2c-3d)(c+d) \int \frac{1}{-\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \sqrt{a-a \sec(e+fx)}}{d} - \frac{(c-d)^2(2c-d)}{d} \right)}{d} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 104

3.206. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^2} dx$

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{d(c+d)(c+d \sec(e+fx))} - \frac{a \left(\frac{2(2c-3d)(c+d) f \frac{1}{\sec(e+fx)a+a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}}{d} - \frac{2(c-d)^2(2c-3d)}{d} \right)}{d} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

218

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{d(c+d)(c+d \sec(e+fx))} - \frac{a \left(\frac{2(2c+3d)(c-d)^{3/2} \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}\right)}{ad\sqrt{c+d}} - \frac{2(2c-3d)(c+d) \arctan\left(\frac{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}\right)}{d} \right)}{d(c+d)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

```
input Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^2,x]
```

```
output -((a^2*(((c - d)*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/(d*(c + d)*(c + d*Sec[e + f*x])) - (a^2*((a*((-2*(2*c - 3*d)*(c + d)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])]/(a*d) - (2*(c - d)^(3/2)*(2*c + 3*d)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]/(a*d*Sqrt[c + d])))/d + (2*c*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a*d)))/(d*(c + d))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.206.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 109 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 171 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegersQ[2*m, 2*n, 2*p]`

- rule 175 `Int[(((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_))) / ((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.206.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.34

method	result
derivativedivides	$16a^3 \frac{\left((c^2 - 2cd + d^2) \left(\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} \right) - \frac{(2c+3d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c+d) \sqrt{(c+d)(c-d)}} \right)}{4d^3} - \frac{f}{16d^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)}$
default	$16a^3 \frac{\left((c^2 - 2cd + d^2) \left(\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} \right) - \frac{(2c+3d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c+d) \sqrt{(c+d)(c-d)}} \right)}{4d^3} - \frac{f}{16d^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)}$
risch	$\frac{2ia^3 (c^2 d e^{3i(fx+e)} - 2c d^2 e^{3i(fx+e)} + d^3 e^{3i(fx+e)} + 2c^3 e^{2i(fx+e)} - c^2 d e^{2i(fx+e)} + c d^2 e^{2i(fx+e)} + 3c^2 d e^{i(fx+e)} + d^3 e^{i(fx+e)})}{f d^2 (1 + e^{2i(fx+e)}) (c+d) c (e^{2i(fx+e)} c + 2d e^{i(fx+e)} + c)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 16/f*a^3*(-1/4*(c^2-2*c*d+d^2)/d^3*(1/2*d/(c+d)*tan(1/2*f*x+1/2*e)/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)-1/2*(2*c+3*d)/(c+d)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))-1/16/d^2/(tan(1/2*f*x+1/2*e)-1)+1/16*(2*c-3*d)/d^3*ln(tan(1/2*f*x+1/2*e)-1)-1/16/d^2/(tan(1/2*f*x+1/2*e)+1)+1/16/d^3*(-2*c+3*d)*ln(tan(1/2*f*x+1/2*e)+1))
```

3.206.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 395 vs. 2(152) = 304.

Time = 0.56 (sec) , antiderivative size = 859, normalized size of antiderivative = 5.34

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^2} dx$$

$$= \left[\frac{((2a^3c^3 + a^3c^2d - 3a^3cd^2) \cos(fx+e))^2 + (2a^3c^2d + a^3cd^2 - 3a^3d^3) \cos(fx+e)}{f d^2 (1 + e^{2i(fx+e)}) (c+d) c (e^{2i(fx+e)} c + 2d e^{i(fx+e)} + c)} \sqrt{\frac{c-d}{c+d}} \log \left(\frac{2cd \cos(fx+e)}{c+d} \right) \right]$$

3.206. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^2} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x, algorithm="ricas")`

output `[-1/2*(((2*a^3*c^3 + a^3*c^2*d - 3*a^3*c*d^2)*cos(f*x + e)^2 + (2*a^3*c^2*d + a^3*c*d^2 - 3*a^3*d^3)*cos(f*x + e))*sqrt((c - d)/(c + d))*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*(c^2 + c*d + (c*d + d^2)*cos(f*x + e))*sqrt((c - d)/(c + d))*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + ((2*a^3*c^3 - a^3*c^2*d - 3*a^3*c*d^2)*cos(f*x + e)^2 + (2*a^3*c^2*d - a^3*c*d^2 - 3*a^3*d^3)*cos(f*x + e))*log(sin(f*x + e) + 1) - ((2*a^3*c^3 - a^3*c^2*d - 3*a^3*c*d^2)*cos(f*x + e)^2 + (2*a^3*c^2*d - a^3*c*d^2 - 3*a^3*d^3)*cos(f*x + e))*log(-sin(f*x + e) + 1) - 2*(a^3*c*d^2 + a^3*d^3 + (2*a^3*c^2*d - a^3*c*d^2 + a^3*d^3)*cos(f*x + e))*sin(f*x + e)/((c^2*d^3 + c*d^4)*f*cos(f*x + e)^2 + (c*d^4 + d^5)*f*cos(f*x + e)), 1/2*(2*(((2*a^3*c^3 + a^3*c^2*d - 3*a^3*c*d^2)*cos(f*x + e)^2 + (2*a^3*c^2*d + a^3*c*d^2 - 3*a^3*d^3)*cos(f*x + e))*sqrt(-(c - d)/(c + d))*arctan(-(d*cos(f*x + e) + c)*sqrt(-(c - d)/(c + d))/((c - d)*sin(f*x + e))) - ((2*a^3*c^3 - a^3*c^2*d - 3*a^3*c*d^2)*cos(f*x + e)^2 + (2*a^3*c^2*d - a^3*c*d^2 - 3*a^3*d^3)*cos(f*x + e))*log(sin(f*x + e) + 1) + ((2*a^3*c^3 - a^3*c^2*d - 3*a^3*c*d^2)*cos(f*x + e)^2 + (2*a^3*c^2*d - a^3*c*d^2 - 3*a^3*d^3)*cos(f*x + e))*log(-sin(f*x + e) + 1) + 2*(a^3*c*d^2 + a^3*d^3 + (2*a^3*c^2*d - a^3*c*d^2 + a^3*d^3)*cos(f*x + e))*sin(f*x + e)/((c^2*d^3 + c*d^4)*f*cos(f*x + e)^2 + (c*d^4 + d^5)*f*cos(f*x + e)))]`

3.206.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^2} dx = a^3 \left(\int \frac{\sec(e+fx)}{c^2 + 2cd\sec(e+fx) + d^2\sec^2(e+fx)} dx + \int \frac{3\sec^2(e+fx)}{c^2 + 2cd\sec(e+fx) + d^2\sec^2(e+fx)} dx + \int \frac{3\sec^3(e+fx)}{c^2 + 2cd\sec(e+fx) + d^2\sec^2(e+fx)} dx + \int \frac{\sec^4(e+fx)}{c^2 + 2cd\sec(e+fx) + d^2\sec^2(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e))**2,x)`

```
output a**3*(Integral(sec(e + f*x)/(c**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)
**2), x) + Integral(3*sec(e + f*x)**2/(c**2 + 2*c*d*sec(e + f*x) + d**2*se
c(e + f*x)**2), x) + Integral(3*sec(e + f*x)**3/(c**2 + 2*c*d*sec(e + f*x)
+ d**2*sec(e + f*x)**2), x) + Integral(sec(e + f*x)**4/(c**2 + 2*c*d*sec(
e + f*x) + d**2*sec(e + f*x)**2), x))
```

3.206.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x, algorithm="m
axima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f
or more de
```

3.206.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(152) = 304$.

Time = 0.37 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.97

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^2} dx =$$

$$\frac{2(2a^3c^3 - a^3c^2d - 4a^3cd^2 + 3a^3d^3) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e) - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-c^2+d^2}} \right) \right)}{(cd^3+d^4)\sqrt{-c^2+d^2}} + \frac{4(a^3c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 - c}{(c \tan(\frac{1}{2} fx + \frac{1}{2} e))^4}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x, algorithm="g
iac")
```

output $-(2*(2*a^3*c^3 - a^3*c^2*d - 4*a^3*c*d^2 + 3*a^3*d^3)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((c*d^3 + d^4)*sqrt(-c^2 + d^2)) + 4*(a^3*c^2*tan(1/2*f*x + 1/2*e)^3 - a^3*c*d*tan(1/2*f*x + 1/2*e)^3 - a^3*c^2*tan(1/2*f*x + 1/2*e) - a^3*d^2*tan(1/2*f*x + 1/2*e))/((c*tan(1/2*f*x + 1/2*e))^4 - d*tan(1/2*f*x + 1/2*e)^4 - 2*c*tan(1/2*f*x + 1/2*e)^2 + c + d)*(c*d^2 + d^3)) + (2*a^3*c - 3*a^3*d)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/d^3 - (2*a^3*c - 3*a^3*d)*log(abs(tan(1/2*f*x + 1/2*e) - 1))/d^3)/f$

3.206.9 Mupad [B] (verification not implemented)

Time = 16.91 (sec) , antiderivative size = 3135, normalized size of antiderivative = 19.47

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^2} dx = \text{Too large to display}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c + d/cos(e + f*x))^2),x)`

output $(a^3*atan(((a^3*((64*tan(e/2 + (f*x)/2)*(4*a^6*c^7 - 9*a^6*d^7 + 27*a^6*c*d^6 - 12*a^6*c^6*d - 16*a^6*c^2*d^5 - 24*a^6*c^3*d^4 + 29*a^6*c^4*d^3 + a^6*c^5*d^2)))/(2*c*d^5 + d^6 + c^2*d^4) + (a^3*((64*(3*a^3*d^11 - 3*a^3*c*d^10 - 4*a^3*c^2*d^9 + 4*a^3*c^3*d^8 + a^3*c^4*d^7 - a^3*c^5*d^6)))/(2*c*d^7 + d^8 + c^2*d^6) - (64*a^3*tan(e/2 + (f*x)/2)*(2*c - 3*d)*(c*d^10 - 2*c^3*d^8 + c^5*d^6)))/(d^3*(2*c*d^5 + d^6 + c^2*d^4)))*(2*c - 3*d))/d^3)*(2*c - 3*d)*1i)/d^3 + (a^3*((64*tan(e/2 + (f*x)/2)*(4*a^6*c^7 - 9*a^6*d^7 + 27*a^6*c*d^6 - 12*a^6*c^6*d - 16*a^6*c^2*d^5 - 24*a^6*c^3*d^4 + 29*a^6*c^4*d^3 + a^6*c^5*d^2)))/(2*c*d^5 + d^6 + c^2*d^4) - (a^3*((64*(3*a^3*d^11 - 3*a^3*c*d^10 - 4*a^3*c^2*d^9 + 4*a^3*c^3*d^8 + a^3*c^4*d^7 - a^3*c^5*d^6)))/(2*c*d^7 + d^8 + c^2*d^6) + (64*a^3*tan(e/2 + (f*x)/2)*(2*c - 3*d)*(c*d^10 - 2*c^3*d^8 + c^5*d^6)))/(d^3*(2*c*d^5 + d^6 + c^2*d^4)))*(2*c - 3*d))/d^3)*(2*c - 3*d)*1i)/d^3)/((128*(4*a^9*c^7 - 9*a^9*c*d^6 - 16*a^9*c^6*d + 36*a^9*c^2*d^5 - 50*a^9*c^3*d^4 + 20*a^9*c^4*d^3 + 15*a^9*c^5*d^2))/(2*c*d^7 + d^8 + c^2*d^6) + (a^3*((64*tan(e/2 + (f*x)/2)*(4*a^6*c^7 - 9*a^6*d^7 + 27*a^6*c*d^6 - 12*a^6*c^6*d - 16*a^6*c^2*d^5 - 24*a^6*c^3*d^4 + 29*a^6*c^4*d^3 + a^6*c^5*d^2)))/(2*c*d^5 + d^6 + c^2*d^4) + (a^3*((64*(3*a^3*d^11 - 3*a^3*c*d^10 - 4*a^3*c^2*d^9 + 4*a^3*c^3*d^8 + a^3*c^4*d^7 - a^3*c^5*d^6)))/(2*c*d^7 + d^8 + c^2*d^6) - (64*a^3*tan(e/2 + (f*x)/2)*(2*c - 3*d)*(c*d^10 - 2*c^3*d^8 + c^5*d^6)))/(d^3*(2*c*d^5 + d^6 + c^2*d^4)))*(2*c - 3*d))/d^3)*(...$

3.207 $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^3} dx$

3.207.1 Optimal result 1463
 3.207.2 Mathematica [C] (warning: unable to verify) 1463
 3.207.3 Rubi [A] (verified) 1464
 3.207.4 Maple [A] (verified) 1469
 3.207.5 Fricas [B] (verification not implemented) 1469
 3.207.6 Sympy [F] 1470
 3.207.7 Maxima [F(-2)] 1471
 3.207.8 Giac [B] (verification not implemented) 1471
 3.207.9 Mupad [B] (verification not implemented) 1472

3.207.1 Optimal result

Integrand size = 31, antiderivative size = 188

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^3} dx$$

$$= \frac{a^3 \operatorname{arctanh}(\sin(e+fx))}{d^3 f} - \frac{a^3 \sqrt{c-d}(2c^2+6cd+7d^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{d^3 (c+d)^{5/2} f}$$

$$- \frac{(c-d)(a^3+a^3 \sec(e+fx)) \tan(e+fx)}{2d(c+d)f(c+d \sec(e+fx))^2} - \frac{a^3(c-d)(2c+5d) \tan(e+fx)}{2d^2(c+d)^2 f(c+d \sec(e+fx))}$$

output `a^3*arctanh(sin(f*x+e))/d^3/f-a^3*(2*c^2+6*c*d+7*d^2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))*(c-d)^(1/2)/d^3/(c+d)^(5/2)/f-1/2*(c-d)*(a^3+a^3*sec(f*x+e))*tan(f*x+e)/d/(c+d)/f/(c+d*sec(f*x+e))^2-1/2*a^3*(c-d)*(2*c+5*d)*tan(f*x+e)/d^2/(c+d)^2/f/(c+d*sec(f*x+e))`

3.207.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 4.33 (sec) , antiderivative size = 393, normalized size of antiderivative = 2.09

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^3} dx$$

$$= \frac{a^3(d+c \cos(e+fx)) \sec^6\left(\frac{1}{2}(e+fx)\right) (1+\sec(e+fx))^3 \left(-4(d+c \cos(e+fx))^2 \log\left(\cos\left(\frac{1}{2}(e+fx)\right)\right) - \dots\right)}{\dots}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^3,x]`

output $(a^3(d + c\cos[e + fx])\sec[(e + fx)/2]^6(1 + \sec[e + fx])^3(-4(d + c\cos[e + fx])^2\log[\cos[(e + fx)/2] - \sin[(e + fx)/2]] + 4(d + c\cos[e + fx])^2\log[\cos[(e + fx)/2] + \sin[(e + fx)/2]] + (4(2c^3 + 4c^2d + c^2d^2 - 7d^3)\operatorname{ArcTan}[(I\cos[e] + \sin[e])(c\sin[e] + (-d + c\cos[e])\tan[(fx)/2]))/(\sqrt{c^2 - d^2}\sqrt{(\cos[e] - I\sin[e])^2}) + (d + c\cos[e + fx])^2(I\cos[e] + \sin[e]))/((c + d)^2\sqrt{c^2 - d^2}\sqrt{(\cos[e] - I\sin[e])^2}) + ((c - d)d\sec[e]((2c^4 + 6c^3d + 5c^2d^2 + 12cd^3 + 2d^4)\sin[e] - c(d(7c^2 + 18cd + 2d^2)\sin[fx] - d(c^2 + 6cd + 2d^2)\sin[2e + fx] + c(2c^2 + 6cd + d^2)\sin[e + 2fx])))/(c^2*(c + d)^2))/((32d^3f*(c + d\sec[e + fx])^3)$

3.207.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.76, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4475, 109, 27, 166, 25, 27, 175, 45, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(a \sec(e + fx) + a)^3}{(c + d \sec(e + fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})(a \csc(e + fx + \frac{\pi}{2}) + a)^3}{(c + d \csc(e + fx + \frac{\pi}{2}))^3} dx$$

↓ 4475

$$\frac{a^2 \tan(e + fx) \int \frac{(\sec(e + fx)a + a)^{5/2}}{\sqrt{a - a \sec(e + fx)}(c + d \sec(e + fx))^3} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 109

$$\frac{a^2 \tan(e + fx) \left(\frac{(c - d) \sqrt{a - a \sec(e + fx)} (a \sec(e + fx) + a)^{3/2}}{2d(c + d)(c + d \sec(e + fx))^2} - \frac{\int \frac{a^3 \sqrt{\sec(e + fx)a + a} (c - 5d - 2(c + d) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} (c + d \sec(e + fx))^2} d \sec(e + fx)}{2ad(c + d)} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 27

3.207. $\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a^2 \int \frac{\sqrt{\sec(e+fx)a+a}(c-5d-2(c+d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2d(c+d)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 166

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a^2 \left(\int \frac{a^2 (2 \sec(e+fx)(c+d)^2 + d(c+7d))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx) - \frac{(c-d)(2c+d)}{ad(c+d)} \right)}{2d(c+d)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 25

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a^2 \left(- \int \frac{a^2 (2 \sec(e+fx)(c+d)^2 + d(c+7d))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx) - \frac{(c-d)(2c+d)}{ad(c+d)} \right)}{2d(c+d)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a^2 \left(- \frac{a \int \frac{2 \sec(e+fx)(c+d)^2 + d(c+7d)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx) - \frac{(c-d)(2c+d)}{d(c+d)}}{2d(c+d)} \right)}{2d(c+d)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 175

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a^2 \left(- \frac{a \left(\frac{2(c+d)^2 \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) - \frac{(2c(c+d)^2 - (c+d))}{d} \right)}{2d(c+d)} \right)}{2d(c+d)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

3.207. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^3} dx$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a \left(\frac{4(c+d)^2 \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} (2c(c+d)^2 - d^2)}{d(c+d)} \right)}{d(c+d)} \right)$$

$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$

↓ 104

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a \left(\frac{4(c+d)^2 \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} 2(2c(c+d)^2 - d^2)}{d(c+d)} \right)}{d(c+d)} \right)$$

$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{(c-d)\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2d(c+d)(c+d \sec(e+fx))^2} - \frac{a \left(\frac{2(2c(c+d)^2 - d^2(c+7d)) \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}\right) - 4(c+d)^2}{ad\sqrt{c-d}\sqrt{c+d}} \right)}{d(c+d)} \right)$$

$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$

3.207. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^3} dx$

input `Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^3,x]`

output `-((a^2*(((c - d)*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/(2*d*(c + d)*(c + d*Sec[e + f*x])^2) - (a^2*(-((a*(-4*(c + d)^2*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]]))/(a*d) - (2*(2*c*(c + d)^2 - d^2*(c + 7*d))*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]))/(a*Sqrt[c - d]*d*Sqrt[c + d])))/(d*(c + d))) - ((c - d)*(2*c + 5*d)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a*d*(c + d)*(c + d*Sec[e + f*x])))/(2*d*(c + d))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]))`

3.207.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 104 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 109 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

rule 166 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] - Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[b*c*(f*g - e*h)*(m + 1) + (b*g - a*h)*(d*e*n + c*f*(p + 1)) + d*(b*(f*g - e*h)*(m + 1) + f*(b*g - a*h)*(n + p + 1))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, p}, x] && ILtQ[m, -1] && GtQ[n, 0]`

rule 175 `Int[(((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_))))/((a_.) + (b_.)*(x_)), x_] := Simp[h/b Int[(c + d*x)^n*(e + f*x)^p, x], x] + Simp[(b*g - a*h)/b Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.207.4 Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.21

method	result
derivativedivides	$16a^3 \left(\frac{\ln(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)}{16d^3} + \frac{(c-d) \left(\frac{d(2c^2+3cd-5d^2) \tan(\frac{fx}{2} + \frac{e}{2})^3}{2c^2+4cd+2d^2} - \frac{d(2c+7d) \tan(\frac{fx}{2} + \frac{e}{2})}{2(c+d)} \right)}{8d^3} - \frac{(2c^2+6cd+7d^2) \operatorname{arctanh}\left(\frac{(c-d) \tan(\frac{fx}{2} + \frac{e}{2})}{\sqrt{(c+d)}}\right)}{2(c^2+2cd+d^2)\sqrt{(c+d)}} \right)$
default	$16a^3 \left(\frac{\ln(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)}{16d^3} + \frac{(c-d) \left(\frac{d(2c^2+3cd-5d^2) \tan(\frac{fx}{2} + \frac{e}{2})^3}{2c^2+4cd+2d^2} - \frac{d(2c+7d) \tan(\frac{fx}{2} + \frac{e}{2})}{2(c+d)} \right)}{8d^3} - \frac{(2c^2+6cd+7d^2) \operatorname{arctanh}\left(\frac{(c-d) \tan(\frac{fx}{2} + \frac{e}{2})}{\sqrt{(c+d)}}\right)}{2(c^2+2cd+d^2)\sqrt{(c+d)}} \right)$
risch	$\frac{ia^3(-c^4 d e^{3i(fx+e)} - 5c^3 d^2 e^{3i(fx+e)} + 4c^2 d^3 e^{3i(fx+e)} + 2c d^4 e^{3i(fx+e)} - 2c^5 e^{2i(fx+e)} - 4c^4 d e^{2i(fx+e)} + c^3 d^2 e^{2i(fx+e)} - 7c^2 d^3 e^{2i(fx+e)} - 4c d^4 e^{2i(fx+e)} - 5d^5 e^{2i(fx+e)})}{c^2 d^2 (c+d)^2}$

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `16/f*a^3*(1/16/d^3*ln(tan(1/2*f*x+1/2*e)+1)+1/8*(c-d)/d^3*((1/2*d*(2*c^2+3*c*d-5*d^2)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3-1/2*d*(2*c+7*d)/(c+d)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^2-1/2*(2*c^2+6*c*d+7*d^2)/(c^2+2*c*d+d^2)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))-1/16/d^3*ln(tan(1/2*f*x+1/2*e)-1))`

3.207.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 552 vs. 2(175) = 350.

Time = 0.60 (sec) , antiderivative size = 1176, normalized size of antiderivative = 6.26

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^3} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x, algorithm="fricas")`

output `[1/4*((2*a^3*c^2*d^2 + 6*a^3*c*d^3 + 7*a^3*d^4 + (2*a^3*c^4 + 6*a^3*c^3*d + 7*a^3*c^2*d^2)*cos(f*x + e)^2 + 2*(2*a^3*c^3*d + 6*a^3*c^2*d^2 + 7*a^3*c*d^3)*cos(f*x + e))*sqrt((c - d)/(c + d))*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*(c^2 + c*d + (c*d + d^2)*cos(f*x + e))*sqrt((c - d)/(c + d))*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(a^3*c^2*d^2 + 2*a^3*c*d^3 + a^3*d^4 + (a^3*c^4 + 2*a^3*c^3*d + a^3*c^2*d^2)*cos(f*x + e)^2 + 2*(a^3*c^3*d + 2*a^3*c^2*d^2 + a^3*c*d^3)*cos(f*x + e))*log(sin(f*x + e) + 1) - 2*(a^3*c^2*d^2 + 2*a^3*c*d^3 + a^3*d^4 + (a^3*c^4 + 2*a^3*c^3*d + a^3*c^2*d^2)*cos(f*x + e)^2 + 2*(a^3*c^3*d + 2*a^3*c^2*d^2 + a^3*c*d^3)*cos(f*x + e))*log(-sin(f*x + e) + 1) - 2*(3*a^3*c^2*d^2 + 3*a^3*c*d^3 - 6*a^3*d^4 + (2*a^3*c^3*d + 4*a^3*c^2*d^2 - 5*a^3*c*d^3 - a^3*d^4)*cos(f*x + e))*sin(f*x + e))/((c^4*d^3 + 2*c^3*d^4 + c^2*d^5)*f*cos(f*x + e)^2 + 2*(c^3*d^4 + 2*c^2*d^5 + c*d^6)*f*cos(f*x + e) + (c^2*d^5 + 2*c*d^6 + d^7)*f), -1/2*((2*a^3*c^2*d^2 + 6*a^3*c*d^3 + 7*a^3*d^4 + (2*a^3*c^4 + 6*a^3*c^3*d + 7*a^3*c^2*d^2)*cos(f*x + e)^2 + 2*(2*a^3*c^3*d + 6*a^3*c^2*d^2 + 7*a^3*c*d^3)*cos(f*x + e))*sqrt(-(c - d)/(c + d))*arctan(-(d*cos(f*x + e) + c)*sqrt(-(c - d)/(c + d))/((c - d)*sin(f*x + e))) - (a^3*c^2*d^2 + 2*a^3*c*d^3 + a^3*d^4 + (a^3*c^4 + 2*a^3*c^3*d + a^3*c^2*d^2)*cos(f*x + e)^2 + 2*(a^3*c^3*d + 2*a^3*c^2*d^2 + a^3*c*d^3)*cos(f*x + e))*log(sin(f*x + e) + 1) + (a^3*c^2*d^2 + 2*a^3*c*d^3 + a^3*d...`

3.207.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^3} dx \\ &= a^3 \left(\int \frac{\sec(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \right. \\ & \quad + \int \frac{3\sec^2(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \\ & \quad + \int \frac{3\sec^3(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \\ & \quad \left. + \int \frac{\sec^4(e+fx)}{c^3 + 3c^2d\sec(e+fx) + 3cd^2\sec^2(e+fx) + d^3\sec^3(e+fx)} dx \right) \end{aligned}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e))**3,x)`

3.207. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^3} dx$

```
output a**3*(Integral(sec(e + f*x)/(c**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e
+ f*x)**2 + d**3*sec(e + f*x)**3), x) + Integral(3*sec(e + f*x)**2/(c**3
+ 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**3)
, x) + Integral(3*sec(e + f*x)**3/(c**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2
*sec(e + f*x)**2 + d**3*sec(e + f*x)**3), x) + Integral(sec(e + f*x)**4/(c
**3 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)
**3), x))
```

3.207.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^3} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x, algorithm="m
axima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f
or more de
```

3.207.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 376 vs. $2(175) = 350$.

Time = 0.42 (sec) , antiderivative size = 376, normalized size of antiderivative = 2.00

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^3} dx$$

$$= \frac{a^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{d^3} - \frac{a^3 \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{d^3} + \frac{(2a^3c^3 + 4a^3c^2d + a^3cd^2 - 7a^3d^3) \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan\left(\frac{c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{d}\right) \right)}{(c^2d^3 + 2cd^4 + d^5)\sqrt{-c^2+d^2}}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x, algorithm="g
iac")
```

3.207. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^3} dx$


```
output (a^3*log(abs(tan(1/2*f*x + 1/2*e) + 1))/d^3 - a^3*log(abs(tan(1/2*f*x + 1/2*e) - 1))/d^3 + (2*a^3*c^3 + 4*a^3*c^2*d + a^3*c*d^2 - 7*a^3*d^3)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((c^2*d^3 + 2*c*d^4 + d^5)*sqrt(-c^2 + d^2)) + (2*a^3*c^3*tan(1/2*f*x + 1/2*e)^3 + a^3*c^2*d*tan(1/2*f*x + 1/2*e)^3 - 8*a^3*c*d^2*tan(1/2*f*x + 1/2*e)^3 + 5*a^3*d^3*tan(1/2*f*x + 1/2*e)^3 - 2*a^3*c^3*tan(1/2*f*x + 1/2*e) - 7*a^3*c^2*d*tan(1/2*f*x + 1/2*e) + 2*a^3*c*d^2*tan(1/2*f*x + 1/2*e) + 7*a^3*d^3*tan(1/2*f*x + 1/2*e))/((c^2*d^2 + 2*c*d^3 + d^4)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^2))/f
```

3.207.9 Mupad [B] (verification not implemented)

Time = 20.18 (sec) , antiderivative size = 4131, normalized size of antiderivative = 21.97

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^3} dx = \text{Too large to display}$$

```
input int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c + d/cos(e + f*x))^3),x)
```

```
output - ((a^3*tan(e/2 + (f*x)/2)*(5*c*d + 2*c^2 - 7*d^2))/(d^2*(c + d)) - (a^3*tan(e/2 + (f*x)/2)^3*(c^2*d - 8*c*d^2 + 2*c^3 + 5*d^3))/(d^2*(c + d)^2)/(f*(2*c*d - tan(e/2 + (f*x)/2)^2*(2*c^2 - 2*d^2) + tan(e/2 + (f*x)/2)^4*(c^2 - 2*c*d + d^2) + c^2 + d^2)) - (a^3*atan(((a^3*((8*tan(e/2 + (f*x)/2)*(8*a^6*c^7 - 53*a^6*d^7 + 59*a^6*c*d^6 + 16*a^6*c^6*d + 53*a^6*c^2*d^5 - 23*a^6*c^3*d^4 - 52*a^6*c^4*d^3 - 8*a^6*c^5*d^2)))/(4*c*d^7 + d^8 + 6*c^2*d^6 + 4*c^3*d^5 + c^4*d^4) + (a^3*((8*(18*a^3*d^12 + 10*a^3*c*d^11 - 32*a^3*c^2*d^10 - 20*a^3*c^3*d^9 + 10*a^3*c^4*d^8 + 10*a^3*c^5*d^7 + 4*a^3*c^6*d^6)))/(4*c*d^9 + d^10 + 6*c^2*d^8 + 4*c^3*d^7 + c^4*d^6) - (8*a^3*tan(e/2 + (f*x)/2)*(8*c*d^12 + 16*c^2*d^11 - 8*c^3*d^10 - 32*c^4*d^9 - 8*c^5*d^8 + 16*c^6*d^7 + 8*c^7*d^6)))/(d^3*(4*c*d^7 + d^8 + 6*c^2*d^6 + 4*c^3*d^5 + c^4*d^4))))/d^3)*i)/d^3 + (a^3*((8*tan(e/2 + (f*x)/2)*(8*a^6*c^7 - 53*a^6*d^7 + 59*a^6*c*d^6 + 16*a^6*c^6*d + 53*a^6*c^2*d^5 - 23*a^6*c^3*d^4 - 52*a^6*c^4*d^3 - 8*a^6*c^5*d^2)))/(4*c*d^7 + d^8 + 6*c^2*d^6 + 4*c^3*d^5 + c^4*d^4) - (a^3*((8*(18*a^3*d^12 + 10*a^3*c*d^11 - 32*a^3*c^2*d^10 - 20*a^3*c^3*d^9 + 10*a^3*c^4*d^8 + 10*a^3*c^5*d^7 + 4*a^3*c^6*d^6)))/(4*c*d^9 + d^10 + 6*c^2*d^8 + 4*c^3*d^7 + c^4*d^6) + (8*a^3*tan(e/2 + (f*x)/2)*(8*c*d^12 + 16*c^2*d^11 - 8*c^3*d^10 - 32*c^4*d^9 - 8*c^5*d^8 + 16*c^6*d^7 + 8*c^7*d^6)))/(d^3*(4*c*d^7 + d^8 + 6*c^2*d^6 + 4*c^3*d^5 + c^4*d^4))))/d^3)*i)/d^3)/((16*(4*a^9*c^6 - 35*a^9*d^6 + 61*a^9*c*d^5 + 10*a^9*c^5*d + 5*a^9*c^2*d^4 ...
```

3.208
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^4} dx$$

3.208.1 Optimal result 1473
 3.208.2 Mathematica [C] (verified) 1474
 3.208.3 Rubi [A] (verified) 1474
 3.208.4 Maple [A] (verified) 1478
 3.208.5 Fracas [B] (verification not implemented) 1479
 3.208.6 Sympy [F] 1480
 3.208.7 Maxima [F(-2)] 1480
 3.208.8 Giac [A] (verification not implemented) 1481
 3.208.9 Mupad [B] (verification not implemented) 1481

3.208.1 Optimal result

Integrand size = 31, antiderivative size = 178

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^4} dx = \frac{5a^3 \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{\sqrt{c-d}(c+d)^{7/2} f} + \frac{a(a+a \sec(e+fx))^2 \tan(e+fx)}{3(c+d)f(c+d \sec(e+fx))^3} - \frac{5a^3(c-d) \tan(e+fx)}{6d(c+d)^2 f(c+d \sec(e+fx))^2} + \frac{5a^3(c+4d) \tan(e+fx)}{6d(c+d)^3 f(c+d \sec(e+fx))}$$

output `5*a^3*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c+d)^(7/2)/f/(c-d)^(1/2)+1/3*a*(a+a*sec(f*x+e))^2*tan(f*x+e)/(c+d)/f/(c+d*sec(f*x+e))^3-5/6*a^3*(c-d)*tan(f*x+e)/d/(c+d)^2/f/(c+d*sec(f*x+e))^2+5/6*a^3*(c+4*d)*tan(f*x+e)/d/(c+d)^3/f/(c+d*sec(f*x+e))`

3.208.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.88 (sec) , antiderivative size = 398, normalized size of antiderivative = 2.24

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^4} dx$$

$$= \frac{a^3(d + c \cos(e + fx)) \sec^6\left(\frac{1}{2}(e + fx)\right) \sec(e + fx)(1 + \sec(e + fx))^3 \left(-\frac{120i \arctan\left(\frac{(i \cos(e) + \sin(e))(c \sin(e) + (-d + c \cos(e) + \sin(e)))}{\sqrt{c^2 - d^2} \sqrt{(\cos(e) - i \sin(e))}}\right)}{\sqrt{c^2 - d^2}} \right)}{\dots}$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^4,x]`

output `(a^3*(d + c*Cos[e + f*x])*Sec[(e + f*x)/2]^6*Sec[e + f*x]*(1 + Sec[e + f*x])^3*(((120*I)*ArcTan[((I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2])]*(d + c*Cos[e + f*x])^3*(Cos[e] - I*Sin[e]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + (c*Sec[e]*(6*(8*c^4 + 6*c^3*d + 30*c^2*d^2 + 9*c*d^3 + 2*d^4)*Sin[f*x] - 3*(6*c^4 - 3*c^3*d + 30*c^2*d^2 + 18*c*d^3 + 4*d^4)*Sin[2*e + f*x] + c*(3*(3*c^3 + 38*c^2*d + 12*c*d^2 + 2*d^3)*Sin[e + 2*f*x] + 3*(3*c^3 - 6*c^2*d - 6*c*d^2 - 2*d^3)*Sin[3*e + 2*f*x] + c*(22*c^2 + 9*c*d + 2*d^2)*Sin[2*e + 3*f*x])) - 2*d*(66*c^4 + 27*c^3*d + 50*c^2*d^2 + 18*c*d^3 + 4*d^4)*Tan[e])/c^3))/(192*(c + d)^3*f*(c + d*Sec[e + f*x])^4)`

3.208.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.62, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3042, 4475, 105, 105, 105, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(a \sec(e + fx) + a)^3}{(c + d \sec(e + fx))^4} dx$$

↓ 3042

$$\begin{aligned}
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a \csc(e+fx+\frac{\pi}{2})+a)^3}{(c+d \csc(e+fx+\frac{\pi}{2}))^4} dx \\
 & \quad \downarrow 4475 \\
 & \frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{5/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 105 \\
 & \frac{a^2 \tan(e+fx) \left(\frac{5a \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3} d \sec(e+fx)}{3(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a(c+d)(c+d \sec(e+fx))^3} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 105 \\
 & \frac{a^2 \tan(e+fx) \left(\frac{5a \left(\frac{3a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d \sec(e+fx))^2} \right)}{3(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a(c+d)(c+d \sec(e+fx))^3} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 105 \\
 & \frac{a^2 \tan(e+fx) \left(\frac{5a \left(\frac{3a \left(\frac{a \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{c+d} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{2a(c+d)(c+d \sec(e+fx))^3} \right)}{3(c+d)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 104
 \end{aligned}$$

3.208. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^4} dx$

$$a^2 \tan(e + fx) \left(\frac{5a \left(\frac{3a \left(\frac{2af \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a\sec(e+fx)}}{c+d}}{d \sqrt{a-a\sec(e+fx)}} - \frac{\sqrt{a-a\sec(e+fx)} \sqrt{a\sec(e+fx)+a}}{a(c+d)(c+d\sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a\sec(e+fx)}(a\sec(e+fx)+a)}{2a(c+d)(c+d\sec(e+fx))} \right)}{3(c+d)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

218

$$a^2 \tan(e + fx) \left(\frac{5a \left(\frac{3a \left(\frac{2 \arctan \left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a\sec(e+fx)}} \right)}{\sqrt{c-d}(c+d)^{3/2}} - \frac{\sqrt{a-a\sec(e+fx)} \sqrt{a\sec(e+fx)+a}}{a(c+d)(c+d\sec(e+fx))} \right)}{2(c+d)} - \frac{\sqrt{a-a\sec(e+fx)}(a\sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d\sec(e+fx))^2} \right)}{3(c+d)} \right) - \sqrt{a - a \sec(e + fx)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

```
input Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^4,x]
```

```
output -((a^2*(-1/3*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2))/(a*(c + d)*(c + d*Sec[e + f*x])^3) + (5*a*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/(a*(c + d)*(c + d*Sec[e + f*x])^2) + (3*a*((2*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(Sqrt[c - d]*(c + d)^(3/2)) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a*(c + d)*(c + d*Sec[e + f*x])))/(2*(c + d)))/(3*(c + d))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.208.3.1 Defintions of rubi rules used

- rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 105 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((m + 1)*(b*e - a*f))), x] - Simp[n*((d*e - c*f)/((m + 1)*(b*e - a*f))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.208.4 Maple [A] (verified)

Time = 1.22 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.28

method	result
derivativedivides	$16a^3 \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{6(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d)}$
default	$16a^3 \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{6(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4(c+d)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d)}$
risch	$ia^3 (22c^5 + 132c^4 d e^{3i(fx+e)} + 54c^3 d^2 e^{3i(fx+e)} + 100c^2 d^3 e^{3i(fx+e)} + 36c d^4 e^{3i(fx+e)} + 36c^4 d e^{2i(fx+e)} + 180c^3 d^2 e^{2i(fx+e)})$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^4,x,method=_RETURNVERBOSE)
```

3.208. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^4} dx$

output
$$\frac{16/f a^3 (-1/6 \tan(1/2 f x + 1/2 e) / (c+d) / (\tan(1/2 f x + 1/2 e)^2 c - \tan(1/2 f x + 1/2 e)^2 d - c-d)^3 - 5/6 / (c+d) * (-1/4 \tan(1/2 f x + 1/2 e) / (c+d) / (\tan(1/2 f x + 1/2 e)^2 c - \tan(1/2 f x + 1/2 e)^2 d - c-d)^2 - 3/4 / (c+d) * (-1/2 \tan(1/2 f x + 1/2 e) / (c+d) / (\tan(1/2 f x + 1/2 e)^2 c - \tan(1/2 f x + 1/2 e)^2 d - c-d) + 1/2 / (c+d) / ((c+d) * (c-d))^{1/2} * \operatorname{arctanh}((c-d) * \tan(1/2 f x + 1/2 e) / ((c+d) * (c-d))^{1/2}))}{12 ((c^8 + 3 c^7 d + 2 c^6 d^2 - 2 c^5 d^3 - 3 c^4 d^4 - c^3 d^5) f \cos$$

3.208.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 477 vs. $2(163) = 326$.

Time = 0.34 (sec) , antiderivative size = 1012, normalized size of antiderivative = 5.69

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^4} dx$$

$$= \left[\frac{15 (a^3 c^3 \cos (fx+e)^3 + 3 a^3 c^2 d \cos (fx+e)^2 + 3 a^3 c d^2 \cos (fx+e) + a^3 d^3) \sqrt{c^2-d^2} \log \left(\frac{2 c d \cos (fx+e) - (c^2-d^2) \cos (fx+e)^2 + 2 \sqrt{c^2-d^2} (d \cos (fx+e) + c) \sin (fx+e) + 2 c^2-d^2}{(c^2 \cos (fx+e)^2 + 2 c d \cos (fx+e) + d^2)} \right) + 2 * (2 a^3 c^4 + 9 a^3 c^3 d + 20 a^3 c^2 d^2 - 9 a^3 c d^3 - 22 a^3 d^4 + (22 a^3 c^4 + 9 a^3 c^3 d - 20 a^3 c^2 d^2 - 9 a^3 c d^3 - 2 a^3 d^4) * \cos (fx+e)^2 + 3 * (3 a^3 c^4 + 16 a^3 c^3 d - 16 a^3 c d^3 - 3 a^3 d^4) * \cos (fx+e)) * \sin (fx+e)}{(c^8 + 3 c^7 d + 2 c^6 d^2 - 2 c^5 d^3 - 3 c^4 d^4 - c^3 d^5) * f \cos (fx+e)^3 + 3 * (c^7 d + 3 c^6 d^2 + 2 c^5 d^3 - 2 c^4 d^4 - 3 c^3 d^5 - c^2 d^6) * f \cos (fx+e)^2 + 3 * (c^6 d^2 + 3 c^5 d^3 + 2 c^4 d^4 - 2 c^3 d^5 - 3 c^2 d^6 - c d^7) * f \cos (fx+e) + (c^5 d^3 + 3 c^4 d^4 + 2 c^3 d^5 - 2 c^2 d^6 - 3 c d^7 - d^8) * f}{12 ((c^8 + 3 c^7 d + 2 c^6 d^2 - 2 c^5 d^3 - 3 c^4 d^4 - c^3 d^5) f \cos$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^4,x, algorithm="fracas")`

output
$$\left[\frac{1}{12} * \left(15 * (a^3 * c^3 * \cos (f * x + e)^3 + 3 * a^3 * c^2 * d * \cos (f * x + e)^2 + 3 * a^3 * c * d^2 * \cos (f * x + e) + a^3 * d^3) * \sqrt{c^2 - d^2} * \log \left(\frac{2 * c * d * \cos (f * x + e) - (c^2 - 2 * d^2) * \cos (f * x + e)^2 + 2 * \sqrt{c^2 - d^2} * (d * \cos (f * x + e) + c) * \sin (f * x + e) + 2 * c^2 - d^2}{(c^2 * \cos (f * x + e)^2 + 2 * c * d * \cos (f * x + e) + d^2)} \right) + 2 * (2 * a^3 * c^4 + 9 * a^3 * c^3 * d + 20 * a^3 * c^2 * d^2 - 9 * a^3 * c * d^3 - 22 * a^3 * d^4 + (22 * a^3 * c^4 + 9 * a^3 * c^3 * d - 20 * a^3 * c^2 * d^2 - 9 * a^3 * c * d^3 - 2 * a^3 * d^4) * \cos (f * x + e)^2 + 3 * (3 * a^3 * c^4 + 16 * a^3 * c^3 * d - 16 * a^3 * c * d^3 - 3 * a^3 * d^4) * \cos (f * x + e)) * \sin (f * x + e) \right) / \left((c^8 + 3 * c^7 * d + 2 * c^6 * d^2 - 2 * c^5 * d^3 - 3 * c^4 * d^4 - c^3 * d^5) * f * \cos (f * x + e)^3 + 3 * (c^7 * d + 3 * c^6 * d^2 + 2 * c^5 * d^3 - 2 * c^4 * d^4 - 3 * c^3 * d^5 - c^2 * d^6) * f * \cos (f * x + e)^2 + 3 * (c^6 * d^2 + 3 * c^5 * d^3 + 2 * c^4 * d^4 - 2 * c^3 * d^5 - 3 * c^2 * d^6 - c * d^7) * f * \cos (f * x + e) + (c^5 * d^3 + 3 * c^4 * d^4 + 2 * c^3 * d^5 - 2 * c^2 * d^6 - 3 * c * d^7 - d^8) * f \right), \frac{1}{6} * \left(15 * (a^3 * c^3 * \cos (f * x + e)^3 + 3 * a^3 * c^2 * d * \cos (f * x + e)^2 + 3 * a^3 * c * d^2 * \cos (f * x + e) + a^3 * d^3) * \sqrt{-c^2 + d^2} * \operatorname{arctan} \left(-\sqrt{-c^2 + d^2} * (d * \cos (f * x + e) + c) / ((c^2 - d^2) * \sin (f * x + e)) \right) + (2 * a^3 * c^4 + 9 * a^3 * c^3 * d + 20 * a^3 * c^2 * d^2 - 9 * a^3 * c * d^3 - 22 * a^3 * d^4 + (22 * a^3 * c^4 + 9 * a^3 * c^3 * d - 20 * a^3 * c^2 * d^2 - 9 * a^3 * c * d^3 - 2 * a^3 * d^4) * \cos (f * x + e)^2 + 3 * (3 * a^3 * c^4 + 16 * a^3 * c^3 * d - 16 * a^3 * c * d^3 - 3 * a^3 * d^4) * \cos (f * x + e)) * \sin (f * x + e) \right) / \left((c^8 + 3 * c^7 * d + 2 * c^6 * d^2 - 2 * c^5 * d^3 - 3 * c^4 * d^4 - c^3 * d^5) * f * \cos (f * x + e)^3 + 3 * (c^7 * d + 3 * c^6 * d^2 + 2 * c^5 * d^3 - 2 * c^4 * d^4 - 3 * c^3 * d^5 - c^2 * d^6) * f * \cos (f * x + e)^2 + 3 * (c^6 * d^2 + 3 * c^5 * d^3 + 2 * c^4 * d^4 - 2 * c^3 * d^5 - c^2 * d^6) * f * \cos (f * x + e) + (c^5 * d^3 + 3 * c^4 * d^4 + 2 * c^3 * d^5 - 2 * c^2 * d^6 - 3 * c * d^7 - d^8) * f \right)$$

3.208.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^4} dx$$

$$= a^3 \left(\int \frac{\sec(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx \right.$$

$$+ \int \frac{3\sec^2(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx$$

$$+ \int \frac{3\sec^3(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx$$

$$\left. + \int \frac{\sec^4(e+fx)}{c^4 + 4c^3d\sec(e+fx) + 6c^2d^2\sec^2(e+fx) + 4cd^3\sec^3(e+fx) + d^4\sec^4(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e))**4,x)`

output `a**3*(Integral(sec(e + f*x)/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x) + Integral(3*sec(e + f*x)**2/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x) + Integral(3*sec(e + f*x)**3/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x) + Integral(sec(e + f*x)**4/(c**4 + 4*c**3*d*sec(e + f*x) + 6*c**2*d**2*sec(e + f*x)**2 + 4*c*d**3*sec(e + f*x)**3 + d**4*sec(e + f*x)**4), x))`

3.208.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.208. $\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^4} dx$

3.208.8 Giac [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.72

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^4} dx =$$

$$\frac{15 \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right) a^3}{(c^3+3c^2d+3cd^2+d^3)\sqrt{-c^2+d^2}} + \frac{15a^3c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 30a^3cd \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 15a^3d^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5}{(c+d)^3} - \frac{40 \tan(\frac{e}{2} + \frac{fx}{2})}{c+d} + \frac{5 \tan(\frac{e}{2} + \frac{fx}{2})^5 (a^3c^2 - 2a^3cd + a^3d^2)}{(c+d)^3} + \frac{11a^3 \tan(\frac{e}{2} + \frac{fx}{2})}{c+d} - \frac{40 \tan(\frac{e}{2} + \frac{fx}{2})}{c+d} + \frac{5a^3 \operatorname{atanh} \left(\frac{\tan(\frac{e}{2} + \frac{fx}{2}) \sqrt{c-d}}{\sqrt{c+d}} \right)}{f(c+d)^{7/2} \sqrt{c-d}}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^4,x, algorithm="giac")`

output `-1/3*(15*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*a^3/((c^3 + 3*c^2*d + 3*c*d^2 + d^3)*sqrt(-c^2 + d^2)) + (15*a^3*c^2*tan(1/2*f*x + 1/2*e)^5 - 30*a^3*c*d*tan(1/2*f*x + 1/2*e)^5 + 15*a^3*d^2*tan(1/2*f*x + 1/2*e)^5 - 40*a^3*c^2*tan(1/2*f*x + 1/2*e)^3 + 40*a^3*d^2*tan(1/2*f*x + 1/2*e)^3 + 33*a^3*c^2*tan(1/2*f*x + 1/2*e) + 66*a^3*c*d*tan(1/2*f*x + 1/2*e) + 33*a^3*d^2*tan(1/2*f*x + 1/2*e))/(c^3 + 3*c^2*d + 3*c*d^2 + d^3)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^3)/f`

3.208.9 Mupad [B] (verification not implemented)

Time = 17.09 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.48

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^4} dx$$

$$= \frac{5 \tan(\frac{e}{2} + \frac{fx}{2})^5 (a^3c^2 - 2a^3cd + a^3d^2)}{(c+d)^3} + \frac{11a^3 \tan(\frac{e}{2} + \frac{fx}{2})}{c+d} - \frac{40 \tan(\frac{e}{2} + \frac{fx}{2})}{c+d} + \frac{5a^3 \operatorname{atanh} \left(\frac{\tan(\frac{e}{2} + \frac{fx}{2}) \sqrt{c-d}}{\sqrt{c+d}} \right)}{f(c+d)^{7/2} \sqrt{c-d}}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c + d/cos(e + f*x))^4),x)`

output $((5*\tan(e/2 + (f*x)/2)^5*(a^3*c^2 + a^3*d^2 - 2*a^3*c*d))/(c + d)^3 + (11*a^3*\tan(e/2 + (f*x)/2))/(c + d) - (40*\tan(e/2 + (f*x)/2)^3*(a^3*c - a^3*d))/(3*(c + d)^2))/(f*(\tan(e/2 + (f*x)/2)^2*(3*c*d^2 - 3*c^2*d - 3*c^3 + 3*d^3) - \tan(e/2 + (f*x)/2)^4*(3*c*d^2 + 3*c^2*d - 3*c^3 - 3*d^3) + 3*c*d^2 + 3*c^2*d + c^3 + d^3 - \tan(e/2 + (f*x)/2)^6*(3*c*d^2 - 3*c^2*d + c^3 - d^3))) + (5*a^3*atanh((\tan(e/2 + (f*x)/2)*(c - d)^{(1/2)})/(c + d)^{(1/2)}))/(f*(c + d)^{(7/2)*(c - d)^{(1/2)})}$

3.209
$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^5} dx$$

3.209.1 Optimal result 1483
 3.209.2 Mathematica [A] (verified) 1484
 3.209.3 Rubi [A] (verified) 1484
 3.209.4 Maple [A] (verified) 1489
 3.209.5 Fricas [B] (verification not implemented) 1489
 3.209.6 Sympy [F] 1490
 3.209.7 Maxima [F(-2)] 1491
 3.209.8 Giac [B] (verification not implemented) 1491
 3.209.9 Mupad [B] (verification not implemented) 1492

3.209.1 Optimal result

Integrand size = 31, antiderivative size = 266

$$\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^5} dx = \frac{5a^3(4c-3d) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{4(c-d)^{3/2}(c+d)^{9/2}f} - \frac{d(a+a \sec(e+fx))^3 \tan(e+fx)}{4(c^2-d^2)f(c+d \sec(e+fx))^4} + \frac{a(4c-3d)(a+a \sec(e+fx))^2 \tan(e+fx)}{12(c-d)(c+d)^2 f(c+d \sec(e+fx))^3} - \frac{5a^3(4c-3d) \tan(e+fx)}{24d(c+d)^3 f(c+d \sec(e+fx))^2} + \frac{5a^3(4c-3d)(c+4d) \tan(e+fx)}{24(c-d)d(c+d)^4 f(c+d \sec(e+fx))}$$

```
output 5/4*a^3*(4*c-3*d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c-d)^(3/2)/(c+d)^(9/2)/f-1/4*d*(a+a*sec(f*x+e))^3*tan(f*x+e)/(c^2-d^2)/f/(c+d*sec(f*x+e))^4+1/12*a*(4*c-3*d)*(a+a*sec(f*x+e))^2*tan(f*x+e)/(c-d)/(c+d)^2/f/(c+d*sec(f*x+e))^3-5/24*a^3*(4*c-3*d)*tan(f*x+e)/d/(c+d)^3/f/(c+d*sec(f*x+e))^2+5/24*a^3*(4*c-3*d)*(c+4*d)*tan(f*x+e)/(c-d)/d/(c+d)^4/f/(c+d*sec(f*x+e))
```

3.209.2 Mathematica [A] (verified)

Time = 6.45 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.03

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^5} dx$$

$$a^3 \left(-\frac{120(4c-3d)\operatorname{arctanh}\left(\frac{(-c+d)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}} - \frac{(-72c^4-478c^3d+336c^2d^2+28cd^3+336d^4+(-296c^4-84c^3d-577c^2d^2+984cd^3+198d^4)\cos[e+fx]+(-72c^4-470c^3d+384c^2d^2+200cd^3+48d^4)\cos[2(e+fx)]-88c^4\cos[3(e+fx)]+36c^3d\cos[3(e+fx)]+37c^2d^2\cos[3(e+fx)]+24cd^3\cos[3(e+fx)]+6d^4\cos[3(e+fx)])\sin[e+fx]}{(d+c\cos[e+fx])^4} \right) / (96(c-d)(c+d)^4 f)$$

input `Integrate[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^5,x]`output `(a^3*((-120*(4*c - 3*d)*ArcTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]])/Sqrt[c^2 - d^2] - ((-72*c^4 - 478*c^3*d + 336*c^2*d^2 + 28*c*d^3 + 336*d^4 + (-296*c^4 - 84*c^3*d - 577*c^2*d^2 + 984*c*d^3 + 198*d^4)*Cos[e + f*x] + (-72*c^4 - 470*c^3*d + 384*c^2*d^2 + 200*c*d^3 + 48*d^4)*Cos[2*(e + f*x)] - 88*c^4*Cos[3*(e + f*x)] + 36*c^3*d*Cos[3*(e + f*x)] + 37*c^2*d^2*Cos[3*(e + f*x)] + 24*c*d^3*Cos[3*(e + f*x)] + 6*d^4*Cos[3*(e + f*x)])*Sin[e + f*x])/(d + c*Cos[e + f*x])^4)/(96*(c - d)*(c + d)^4*f)`**3.209.3 Rubi [A] (verified)**Time = 0.46 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.39, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 4475, 107, 105, 105, 105, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a\sec(e+fx)+a)^3}{(c+d\sec(e+fx))^5} dx$$

$$\downarrow 3042$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^3}{\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)^5} dx$$

$$\downarrow 4475$$

$$\begin{aligned}
 & \frac{a^2 \tan(e+fx) \int \frac{(\sec(e+fx)a+a)^{5/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^5} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 107 \\
 & \frac{a^2 \tan(e+fx) \left(\frac{(4c-3d) \int \frac{(\sec(e+fx)a+a)^{5/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4} d \sec(e+fx)}{4(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{7/2}}{4a^2(c^2-d^2)(c+d \sec(e+fx))^4} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 105 \\
 & \frac{a^2 \tan(e+fx) \left(\frac{(4c-3d) \left(\frac{5a \int \frac{(\sec(e+fx)a+a)^{3/2}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3} d \sec(e+fx)}{3(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a(c+d)(c+d \sec(e+fx))^3} \right)}{4(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{7/2}}{4a^2(c^2-d^2)(c+d \sec(e+fx))^4} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 105 \\
 & \frac{a^2 \tan(e+fx) \left(\frac{(4c-3d) \left(\frac{5a \left(\frac{3a \int \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{3/2}}{2a(c+d)(c+d \sec(e+fx))^2} \right)}{3(c+d)} - \frac{\sqrt{a-a \sec(e+fx)}(a \sec(e+fx)+a)^{5/2}}{3a(c+d)(c+d \sec(e+fx))^3} \right)}{4(c^2-d^2)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 105
 \end{aligned}$$

3.209. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^5} dx$

$$\begin{array}{l}
 a^2 \tan(e + fx) \\
 \left. \begin{array}{l}
 (4c-3d) \\
 5a \\
 3a \\
 2(c+d) \\
 3(c+d) \\
 4(c^2-d^2)
 \end{array} \right\} \left(\frac{a \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}} d \sec(e+fx)}{c+d} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right) - \frac{\sqrt{a-a \sec(e+fx)}}{2a}
 \end{array}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

104

$$\begin{array}{l}
 a^2 \tan(e + fx) \\
 \left. \begin{array}{l}
 (4c-3d) \\
 5a \\
 3a \\
 2(c+d) \\
 3(c+d) \\
 4(c^2-d^2)
 \end{array} \right\} \left(\frac{2a \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{c+d} - \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a(c+d)(c+d \sec(e+fx))} \right) - \frac{\sqrt{a-a \sec(e+fx)}}{2a(c+d)(c-d)}
 \end{array}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

3.209. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^5} dx$

↓ 218

$$a^2 \tan(e + fx) \frac{d\sqrt{a - a \sec(e + fx)}(a \sec(e + fx) + a)^{7/2}}{4a^2(c^2 - d^2)(c + d \sec(e + fx))^4} + \frac{5a \left(\frac{3a \left(\frac{2 \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}\right) - \frac{\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)}}{a(c+d)(c+d \sec(e+fx))} \right)}{\sqrt{c-d}(c+d)^{3/2}} \right)}{2(c+d)} \right)}{(4c-3d) \frac{3(c+d)}{4(c+d)}}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

```
input Int[(Sec[e + f*x]*(a + a*Sec[e + f*x])^3)/(c + d*Sec[e + f*x])^5,x]
```

```
output -((a^2*((d*Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(7/2))/(4*a^2*(c^2 - d^2)*(c + d*Sec[e + f*x])^4) + ((4*c - 3*d)*(-1/3*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(5/2))/(a*(c + d)*(c + d*Sec[e + f*x])^3) + (5*a*(-1/2*(Sqrt[a - a*Sec[e + f*x]]*(a + a*Sec[e + f*x])^(3/2))/(a*(c + d)*(c + d*Sec[e + f*x])^2) + (3*a*((2*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(Sqrt[c - d]*(c + d)^(3/2)) - (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])/(a*(c + d)*(c + d*Sec[e + f*x])))/(2*(c + d)))/(3*(c + d)))/(4*(c^2 - d^2))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```


3.209.3.1 Defintions of rubi rules used

- rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 105 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((m + 1)*(b*e - a*f))), x] - Simp[n*((d*e - c*f)/((m + 1)*(b*e - a*f))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`
- rule 107 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Simp[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f))] Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.209.4 Maple [A] (verified)

Time = 1.70 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.14

method	result
derivativedivides	$16a^3 \frac{\left(-\frac{5(4c-3d)(c^2-2cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{64(c^4+4c^3d+6c^2d^2+4cd^3+d^4)} + \frac{55(c-d)(4c-3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{192(c^3+3c^2d+3cd^2+d^3)} - \frac{73(4c-3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{192(c^2+2cd+d^2)} + \frac{(44c-49d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{64(c+d)(c-d)} \right)}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 d - c - d \right)^4}$
default	$16a^3 \frac{\left(-\frac{5(4c-3d)(c^2-2cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{64(c^4+4c^3d+6c^2d^2+4cd^3+d^4)} + \frac{55(c-d)(4c-3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{192(c^3+3c^2d+3cd^2+d^3)} - \frac{73(4c-3d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{192(c^2+2cd+d^2)} + \frac{(44c-49d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{64(c+d)(c-d)} \right)}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 d - c - d \right)^4}$
risch	Expression too large to display

input `int(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^5,x,method=_RETURNVERBOSE)`

output `16/f*a^3*((-5/64*(4*c-3*d)*(c^2-2*c*d+d^2)/(c^4+4*c^3*d+6*c^2*d^2+4*c*d^3+d^4)*tan(1/2*f*x+1/2*e)^7+55/192*(c-d)*(4*c-3*d)/(c^3+3*c^2*d+3*c*d^2+d^3)*tan(1/2*f*x+1/2*e)^5-73/192*(4*c-3*d)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3+1/64*(44*c-49*d)/(c+d)/(c-d)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^4+5/64*(4*c-3*d)/(c^5+3*c^4*d+2*c^3*d^2-2*c^2*d^3-3*c*d^4-d^5)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))`

3.209.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 828 vs. 2(247) = 494.

Time = 0.36 (sec) , antiderivative size = 1714, normalized size of antiderivative = 6.44

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^5} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^5,x, algorithm="fricas")`

output

```
[1/48*(15*(4*a^3*c*d^4 - 3*a^3*d^5 + (4*a^3*c^5 - 3*a^3*c^4*d)*cos(f*x + e)
)^4 + 4*(4*a^3*c^4*d - 3*a^3*c^3*d^2)*cos(f*x + e)^3 + 6*(4*a^3*c^3*d^2 -
3*a^3*c^2*d^3)*cos(f*x + e)^2 + 4*(4*a^3*c^2*d^3 - 3*a^3*c*d^4)*cos(f*x +
e))*sqrt(c^2 - d^2)*log(((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2
+ 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2
*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(2*a^3*c^5*d + 12*a^3*c^4
*d^2 + 41*a^3*c^3*d^3 - 84*a^3*c^2*d^4 - 43*a^3*c*d^5 + 72*a^3*d^6 + (88*a
^3*c^6 - 36*a^3*c^5*d - 125*a^3*c^4*d^2 + 12*a^3*c^3*d^3 + 31*a^3*c^2*d^4
+ 24*a^3*c*d^5 + 6*a^3*d^6)*cos(f*x + e)^3 + (36*a^3*c^6 + 235*a^3*c^5*d -
228*a^3*c^4*d^2 - 335*a^3*c^3*d^3 + 168*a^3*c^2*d^4 + 100*a^3*c*d^5 + 24*
a^3*d^6)*cos(f*x + e)^2 + (8*a^3*c^6 + 48*a^3*c^5*d + 164*a^3*c^4*d^2 - 27
6*a^3*c^3*d^3 - 217*a^3*c^2*d^4 + 228*a^3*c*d^5 + 45*a^3*d^6)*cos(f*x + e)
)*sin(f*x + e))/((c^11 + 3*c^10*d + c^9*d^2 - 5*c^8*d^3 - 5*c^7*d^4 + c^6*
d^5 + 3*c^5*d^6 + c^4*d^7)*f*cos(f*x + e)^4 + 4*(c^10*d + 3*c^9*d^2 + c^8*
d^3 - 5*c^7*d^4 - 5*c^6*d^5 + c^5*d^6 + 3*c^4*d^7 + c^3*d^8)*f*cos(f*x + e)
)^3 + 6*(c^9*d^2 + 3*c^8*d^3 + c^7*d^4 - 5*c^6*d^5 - 5*c^5*d^6 + c^4*d^7 +
3*c^3*d^8 + c^2*d^9)*f*cos(f*x + e)^2 + 4*(c^8*d^3 + 3*c^7*d^4 + c^6*d^5
- 5*c^5*d^6 - 5*c^4*d^7 + c^3*d^8 + 3*c^2*d^9 + c*d^10)*f*cos(f*x + e) + (
c^7*d^4 + 3*c^6*d^5 + c^5*d^6 - 5*c^4*d^7 - 5*c^3*d^8 + c^2*d^9 + 3*c*d^10
+ d^11)*f), 1/24*(15*(4*a^3*c*d^4 - 3*a^3*d^5 + (4*a^3*c^5 - 3*a^3*c^4...
```

3.209.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^5} dx$$

$$= a^3 \left(\int \frac{\sec(e+fx)}{c^5 + 5c^4d\sec(e+fx) + 10c^3d^2\sec^2(e+fx) + 10c^2d^3\sec^3(e+fx) + 5cd^4\sec^4(e+fx) + d^5\sec^5(e+fx)} dx \right.$$

$$+ \int \frac{3\sec^2(e+fx)}{c^5 + 5c^4d\sec(e+fx) + 10c^3d^2\sec^2(e+fx) + 10c^2d^3\sec^3(e+fx) + 5cd^4\sec^4(e+fx) + d^5\sec^5(e+fx)} dx$$

$$+ \int \frac{3\sec^3(e+fx)}{c^5 + 5c^4d\sec(e+fx) + 10c^3d^2\sec^2(e+fx) + 10c^2d^3\sec^3(e+fx) + 5cd^4\sec^4(e+fx) + d^5\sec^5(e+fx)} dx$$

$$+ \left. \int \frac{\sec^4(e+fx)}{c^5 + 5c^4d\sec(e+fx) + 10c^3d^2\sec^2(e+fx) + 10c^2d^3\sec^3(e+fx) + 5cd^4\sec^4(e+fx) + d^5\sec^5(e+fx)} dx \right)$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e))**5,x)`

```
output a**3*(Integral(sec(e + f*x)/(c**5 + 5*c**4*d*sec(e + f*x) + 10*c**3*d**2*s
ec(e + f*x)**2 + 10*c**2*d**3*sec(e + f*x)**3 + 5*c*d**4*sec(e + f*x)**4 +
d**5*sec(e + f*x)**5), x) + Integral(3*sec(e + f*x)**2/(c**5 + 5*c**4*d*s
ec(e + f*x) + 10*c**3*d**2*sec(e + f*x)**2 + 10*c**2*d**3*sec(e + f*x)**3
+ 5*c*d**4*sec(e + f*x)**4 + d**5*sec(e + f*x)**5), x) + Integral(3*sec(e
+ f*x)**3/(c**5 + 5*c**4*d*sec(e + f*x) + 10*c**3*d**2*sec(e + f*x)**2 + 1
0*c**2*d**3*sec(e + f*x)**3 + 5*c*d**4*sec(e + f*x)**4 + d**5*sec(e + f*x)
**5), x) + Integral(sec(e + f*x)**4/(c**5 + 5*c**4*d*sec(e + f*x) + 10*c**
3*d**2*sec(e + f*x)**2 + 10*c**2*d**3*sec(e + f*x)**3 + 5*c*d**4*sec(e + f
*x)**4 + d**5*sec(e + f*x)**5), x))
```

3.209.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^5} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^5,x, algorithm="m
axima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f
or more de
```

3.209.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 601 vs. $2(247) = 494$.

Time = 0.48 (sec) , antiderivative size = 601, normalized size of antiderivative = 2.26

$$\int \frac{\sec(e + fx)(a + a \sec(e + fx))^3}{(c + d \sec(e + fx))^5} dx$$

$$= \frac{15(4a^3c - 3a^3d) \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(-2c+2d) + \arctan \left(-\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right)}{(c^5+3c^4d+2c^3d^2-2c^2d^3-3cd^4-d^5)\sqrt{-c^2+d^2}} - \frac{60a^3c^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^7 - 225a^3c^3d \tan(\frac{1}{2}fx + \frac{1}{2}e)^6 + \dots}{(c^5+3c^4d+2c^3d^2-2c^2d^3-3cd^4-d^5)\sqrt{-c^2+d^2}}$$

3.209. $\int \frac{\sec(e+fx)(a+a \sec(e+fx))^3}{(c+d \sec(e+fx))^5} dx$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^5,x, algorithm="giac")`

output `1/12*(15*(4*a^3*c - 3*a^3*d)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((c^5 + 3*c^4*d + 2*c^3*d^2 - 2*c^2*d^3 - 3*c*d^4 - d^5)*sqrt(-c^2 + d^2)) - (60*a^3*c^4*tan(1/2*f*x + 1/2*e)^7 - 225*a^3*c^3*d*tan(1/2*f*x + 1/2*e)^7 + 315*a^3*c^2*d^2*tan(1/2*f*x + 1/2*e)^7 - 195*a^3*c*d^3*tan(1/2*f*x + 1/2*e)^7 + 45*a^3*d^4*tan(1/2*f*x + 1/2*e)^7 - 220*a^3*c^4*tan(1/2*f*x + 1/2*e)^5 + 385*a^3*c^3*d*tan(1/2*f*x + 1/2*e)^5 + 55*a^3*c^2*d^2*tan(1/2*f*x + 1/2*e)^5 - 385*a^3*c*d^3*tan(1/2*f*x + 1/2*e)^5 + 165*a^3*d^4*tan(1/2*f*x + 1/2*e)^5 + 292*a^3*c^4*tan(1/2*f*x + 1/2*e)^3 + 73*a^3*c^3*d*tan(1/2*f*x + 1/2*e)^3 - 511*a^3*c^2*d^2*tan(1/2*f*x + 1/2*e)^3 - 73*a^3*c*d^3*tan(1/2*f*x + 1/2*e)^3 + 219*a^3*d^4*tan(1/2*f*x + 1/2*e)^3 - 132*a^3*c^4*tan(1/2*f*x + 1/2*e) - 249*a^3*c^3*d*tan(1/2*f*x + 1/2*e) + 45*a^3*c^2*d^2*tan(1/2*f*x + 1/2*e) + 309*a^3*c*d^3*tan(1/2*f*x + 1/2*e) + 147*a^3*d^4*tan(1/2*f*x + 1/2*e))/((c^5 + 3*c^4*d + 2*c^3*d^2 - 2*c^2*d^3 - 3*c*d^4 - d^5)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^4)/f`

3.209.9 Mupad [B] (verification not implemented)

Time = 17.09 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.45

$$\int \frac{\sec(e+fx)(a+a\sec(e+fx))^3}{(c+d\sec(e+fx))^5} dx$$

$$= \frac{\frac{55 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (4a^3c^2 - 7a^3cd + 3a^3d^2)}{12(c+d)^3} - \frac{73 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (4a^3c - 3a^3d)}{12(c+d)^2}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 (6c^4 - 12c^2d^2 + 6d^4) + \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (-4c^4 - 8c^3d + 8cd^3 + 4d^4) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right) \right)^6} + \frac{5a^3 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\sqrt{c-d}}{\sqrt{c+d}}\right) (4c - 3d)}{4f(c+d)^{9/2}(c-d)^{3/2}}$$

input `int((a + a/cos(e + f*x))^3/(cos(e + f*x)*(c + d/cos(e + f*x))^5),x)`

output $((55*\tan(e/2 + (f*x)/2)^5*(4*a^3*c^2 + 3*a^3*d^2 - 7*a^3*c*d))/(12*(c + d)^3) - (73*\tan(e/2 + (f*x)/2)^3*(4*a^3*c - 3*a^3*d))/(12*(c + d)^2) - (5*\tan(e/2 + (f*x)/2)^7*(4*a^3*c^3 - 3*a^3*d^3 + 10*a^3*c*d^2 - 11*a^3*c^2*d))/(4*(c + d)^4) + (a^3*\tan(e/2 + (f*x)/2)*(44*c - 49*d))/(4*(c + d)*(c - d)))/(f*(\tan(e/2 + (f*x)/2)^4*(6*c^4 + 6*d^4 - 12*c^2*d^2) + \tan(e/2 + (f*x)/2)^2*(8*c*d^3 - 8*c^3*d - 4*c^4 + 4*d^4) - \tan(e/2 + (f*x)/2)^6*(8*c*d^3 - 8*c^3*d + 4*c^4 - 4*d^4) + \tan(e/2 + (f*x)/2)^8*(c^4 - 4*c^3*d - 4*c*d^3 + d^4 + 6*c^2*d^2) + 4*c*d^3 + 4*c^3*d + c^4 + d^4 + 6*c^2*d^2)) + (5*a^3*\operatorname{atanh}((\tan(e/2 + (f*x)/2)*(c - d)^{1/2})/(c + d)^{1/2})*(4*c - 3*d))/(4*f*(c + d)^{9/2}*(c - d)^{3/2}))$

3.210 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+a \sec(e+fx)} dx$

3.210.1 Optimal result 1494
 3.210.2 Mathematica [B] (verified) 1495
 3.210.3 Rubi [A] (verified) 1495
 3.210.4 Maple [A] (verified) 1500
 3.210.5 Fricas [A] (verification not implemented) 1500
 3.210.6 Sympy [F] 1501
 3.210.7 Maxima [B] (verification not implemented) 1501
 3.210.8 Giac [A] (verification not implemented) 1502
 3.210.9 Mupad [B] (verification not implemented) 1503

3.210.1 Optimal result

Integrand size = 31, antiderivative size = 183

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+a \sec(e+fx)} dx$$

$$= \frac{d(8c^3 - 12c^2d + 12cd^2 - 3d^3) \operatorname{arctanh}(\sin(e+fx))}{2af}$$

$$- \frac{(3c-4d)d(c+d \sec(e+fx))^2 \tan(e+fx)}{3af} + \frac{(c-d)(c+d \sec(e+fx))^3 \tan(e+fx)}{f(a+a \sec(e+fx))}$$

$$- \frac{d(4(3c^3 - 16c^2d + 12cd^2 - 4d^3) + d(6c^2 - 20cd + 9d^2) \sec(e+fx)) \tan(e+fx)}{6af}$$

output

```
1/2*d*(8*c^3-12*c^2*d+12*c*d^2-3*d^3)*arctanh(sin(f*x+e))/a/f-1/3*(3*c-4*d)
)*d*(c+d*sec(f*x+e))^2*tan(f*x+e)/a/f+(c-d)*(c+d*sec(f*x+e))^3*tan(f*x+e)/
f/(a+a*sec(f*x+e))-1/6*d*(12*c^3-64*c^2*d+48*c*d^2-16*d^3+d*(6*c^2-20*c*d+
9*d^2)*sec(f*x+e))*tan(f*x+e)/a/f
```

3.210.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1243 vs. $2(183) = 366$.

Time = 7.85 (sec) , antiderivative size = 1243, normalized size of antiderivative = 6.79

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+a\sec(e+fx)} dx = \text{Too large to display}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + a*Sec[e + f*x]),x]`

output

```
((-8*c^3*d + 12*c^2*d^2 - 12*c*d^3 + 3*d^4)*Cos[e/2 + (f*x)/2]^2*Cos[e + f*x]^3*Log[Cos[e/2 + (f*x)/2] - Sin[e/2 + (f*x)/2]]*(c + d*Sec[e + f*x])^4)/(f*(d + c*Cos[e + f*x])^4*(a + a*Sec[e + f*x])) + ((8*c^3*d - 12*c^2*d^2 + 12*c*d^3 - 3*d^4)*Cos[e/2 + (f*x)/2]^2*Cos[e + f*x]^3*Log[Cos[e/2 + (f*x)/2] + Sin[e/2 + (f*x)/2]]*(c + d*Sec[e + f*x])^4)/(f*(d + c*Cos[e + f*x])^4*(a + a*Sec[e + f*x])) + (Cos[e/2 + (f*x)/2]*Sec[e/2]*Sec[e]*(c + d*Sec[e + f*x])^4*(-18*c^4*Sin[(f*x)/2] + 72*c^3*d*Sin[(f*x)/2] - 36*c^2*d^2*Sin[(f*x)/2] + 24*c*d^3*Sin[(f*x)/2] + 6*d^4*Sin[(f*x)/2] + 18*c^4*Sin[(3*f*x)/2] - 72*c^3*d*Sin[(3*f*x)/2] + 180*c^2*d^2*Sin[(3*f*x)/2] - 108*c*d^3*Sin[(3*f*x)/2] + 39*d^4*Sin[(3*f*x)/2] - 72*c^2*d^2*Sin[e - (f*x)/2] + 48*c*d^3*Sin[e - (f*x)/2] - 24*d^4*Sin[e - (f*x)/2] - 36*c^2*d^2*Sin[e + (f*x)/2] + 24*c*d^3*Sin[e + (f*x)/2] - 6*d^4*Sin[e + (f*x)/2] - 18*c^4*Sin[2*e + (f*x)/2] + 72*c^3*d*Sin[2*e + (f*x)/2] - 144*c^2*d^2*Sin[2*e + (f*x)/2] + 96*c*d^3*Sin[2*e + (f*x)/2] - 24*d^4*Sin[2*e + (f*x)/2] + 72*c^2*d^2*Sin[e + (3*f*x)/2] - 36*c*d^3*Sin[e + (3*f*x)/2] + 21*d^4*Sin[e + (3*f*x)/2] + 18*c^4*Sin[2*e + (3*f*x)/2] - 72*c^3*d*Sin[2*e + (3*f*x)/2] + 72*c^2*d^2*Sin[2*e + (3*f*x)/2] - 36*c*d^3*Sin[2*e + (3*f*x)/2] + 9*d^4*Sin[2*e + (3*f*x)/2] - 36*c^2*d^2*Sin[3*e + (3*f*x)/2] + 36*c*d^3*Sin[3*e + (3*f*x)/2] - 9*d^4*Sin[3*e + (3*f*x)/2] + 36*c^2*d^2*Sin[e + (5*f*x)/2] - 12*c*d^3*Sin[e + (5*f*x)/2] + 7*d^4*Sin[e + (5*f*x)/2] - 6*c^4*Sin[2*e + (5*f*x)/2]...
```

3.210.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.67, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4475, 109, 25, 27, 170, 25, 27, 164, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.210. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+a\sec(e+fx)} dx$

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)^4}{a\csc\left(e+fx+\frac{\pi}{2}\right)+a} dx \\
 & \quad \downarrow \text{4475} \\
 & \frac{a^2 \tan(e+fx) \int \frac{(c+d\sec(e+fx))^4}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \\
 & \quad \downarrow \text{109} \\
 & \frac{a^2 \tan(e+fx) \left(-\frac{\int -\frac{a^2 d(4c-3d-(3c-4d)\sec(e+fx))(c+d\sec(e+fx))^2}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d\sec(e+fx)}{a^3} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^3}{a^2\sqrt{a\sec(e+fx)+a}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \\
 & \quad \downarrow \text{25} \\
 & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2 d(4c-3d-(3c-4d)\sec(e+fx))(c+d\sec(e+fx))^2}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d\sec(e+fx)}{a^3} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^3}{a^2\sqrt{a\sec(e+fx)+a}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 \tan(e+fx) \left(\frac{d \int \frac{(4c-3d-(3c-4d)\sec(e+fx))(c+d\sec(e+fx))^2}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d\sec(e+fx)}{a} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^3}{a^2\sqrt{a\sec(e+fx)+a}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \\
 & \quad \downarrow \text{170} \\
 & \frac{a^2 \tan(e+fx) \left(\frac{d \left(\frac{(3c-4d)\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}(c+d\sec(e+fx))^2}{3a^2} - \frac{\int -\frac{a^2(c+d\sec(e+fx))(12c^2-15dc+8d^2-(6c^2-20dc+9d^2)\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}}}{3a^2} \right)}{a} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

3.210. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+a\sec(e+fx)} dx$

$$a^2 \tan(e + fx) \left(\frac{d \left(\int \frac{a^2(c+d \sec(e+fx))(12c^2-15dc+8d^2-(6c^2-20dc+9d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) + \frac{(3c-4d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}(c+d)}{3a^2} \right)}{a} \right) = f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}$$

27

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{1}{3} \int \frac{(c+d \sec(e+fx))(12c^2-15dc+8d^2-(6c^2-20dc+9d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) + \frac{(3c-4d) \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}(c+d)}{3a^2} \right)}{a} \right) = f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}$$

164

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{1}{3} \left(\frac{3}{2} (8c^3-12c^2d+12cd^2-3d^3) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) + \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a} (d(6c^2-20cd+9d^2))}{a} \right) \right)}{a} \right) = f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}$$

45

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{1}{3} \left(3(8c^3-12c^2d+12cd^2-3d^3) \int \frac{1}{-\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} + \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a} (d(6c^2-20cd+9d^2))}{2a^2} \right) \right)}{a} \right) = f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}$$

218

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{1}{3} \left(\frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a} (d(6c^2-20cd+9d^2)) \sec(e+fx) + 4(3c^3-16c^2d+12cd^2-4d^3)}{2a^2} \right) \right) - \frac{3(8c^3-12c^2d+12cd^2-3d^3)}{a}}{a} \right) = f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}$$

```
input Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + a*Sec[e + f*x]), x]
```

3.210. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+a \sec(e+fx)} dx$

```
output -((a^2*(-(((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^3)/(a^2*S
qrt[a + a*Sec[e + f*x]]))) + (d*(((3*c - 4*d)*Sqrt[a - a*Sec[e + f*x]]*Sqrt
[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(3*a^2) + ((-3*(8*c^3 - 12*c^
2*d + 12*c*d^2 - 3*d^3)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e +
f*x]]])/a + (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(4*(3*c^3
- 16*c^2*d + 12*c*d^2 - 4*d^3) + d*(6*c^2 - 20*c*d + 9*d^2)*Sec[e + f*x]))
/(2*a^2))/3))/a)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[
e + f*x]]))
```

3.210.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 109 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_
))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f
*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1)
+ c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p)
+ b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c,
d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] ||
IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

rule 164 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*(g_.) + (h_.)*(x_)), x_] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x))*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`

rule 170 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.210.4 Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.50

method	result
parallelrisch	$-12\left(c^3 - \frac{3}{2}c^2d + \frac{3}{2}cd^2 - \frac{3}{8}d^3\right)\left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3}\right)d\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 12\left(c^3 - \frac{3}{2}c^2d + \frac{3}{2}cd^2 - \frac{3}{8}d^3\right)\left(\cos(fx+e)\right)$
derivativedivides	$\tan\left(\frac{fx}{2} + \frac{e}{2}\right)c^4 - 4\tan\left(\frac{fx}{2} + \frac{e}{2}\right)c^3d + 6\tan\left(\frac{fx}{2} + \frac{e}{2}\right)c^2d^2 - 4\tan\left(\frac{fx}{2} + \frac{e}{2}\right)cd^3 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)d^4 - \frac{d^4}{3\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^3} - \frac{d(8c^3 - 3c^2d + 3cd^2 - 3d^3)}{3\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^3}$
default	$\tan\left(\frac{fx}{2} + \frac{e}{2}\right)c^4 - 4\tan\left(\frac{fx}{2} + \frac{e}{2}\right)c^3d + 6\tan\left(\frac{fx}{2} + \frac{e}{2}\right)c^2d^2 - 4\tan\left(\frac{fx}{2} + \frac{e}{2}\right)cd^3 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)d^4 - \frac{d^4}{3\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^3} - \frac{d(8c^3 - 3c^2d + 3cd^2 - 3d^3)}{3\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^3}$
norman	$\frac{(c^4 - 4c^3d + 6c^2d^2 - 4cd^3 + d^4)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{af} + \frac{(c^4 - 4c^3d + 18c^2d^2 - 8cd^3 + 4d^4)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} - \frac{(4c^4 - 16c^3d + 36c^2d^2 - 28cd^3 + 9d^4)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af}$
risch	$i(36c^2d^2e^{6i(fx+e)} + 144c^2d^2e^{4i(fx+e)} - 108cd^3e^{2i(fx+e)} - 72c^3de^{2i(fx+e)} - 24c^3de^{6i(fx+e)} - 36cd^3e^{5i(fx+e)} - 96cd^3e^{4i(fx+e)})$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$3*(-4*(c^3-3/2*c^2*d+3/2*c*d^2-3/8*d^3)*(\cos(f*x+e)+1/3*\cos(3*f*x+3*e))*d*\ln(\tan(1/2*f*x+1/2*e)-1)+4*(c^3-3/2*c^2*d+3/2*c*d^2-3/8*d^3)*(\cos(f*x+e)+1/3*\cos(3*f*x+3*e))*d*\ln(\tan(1/2*f*x+1/2*e)+1)+\tan(1/2*f*x+1/2*e)*((1/3*c^4-4/3*c^3*d+4*c^2*d^2-8/3*c*d^3+8/9*d^4)*\cos(3*f*x+3*e)+(4*c^2*d^2-4/3*c*d^3+7/9*d^4)*\cos(2*f*x+2*e)+(c^4-4*c^3*d+12*c^2*d^2-16/3*c*d^3+22/9*d^4)*\cos(f*x+e)+4*c^2*d^2-4/3*c*d^3+11/9*d^4))/a/f/(\cos(3*f*x+3*e)+3*\cos(f*x+e))$$

3.210.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.62

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+a\sec(e+fx)} dx$$

$$= \frac{3((8c^3d - 12c^2d^2 + 12cd^3 - 3d^4)\cos(fx+e)^4 + (8c^3d - 12c^2d^2 + 12cd^3 - 3d^4)\cos(fx+e)^3)\log(\sin(fx+e))}{a}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e)),x, algorithm="fricas")`

3.210.
$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+a\sec(e+fx)} dx$$

output
$$\frac{1}{12} * (3 * ((8 * c^3 * d - 12 * c^2 * d^2 + 12 * c * d^3 - 3 * d^4) * \cos(f * x + e)^4 + (8 * c^3 * d - 12 * c^2 * d^2 + 12 * c * d^3 - 3 * d^4) * \cos(f * x + e)^3) * \log(\sin(f * x + e) + 1) - 3 * ((8 * c^3 * d - 12 * c^2 * d^2 + 12 * c * d^3 - 3 * d^4) * \cos(f * x + e)^4 + (8 * c^3 * d - 12 * c^2 * d^2 + 12 * c * d^3 - 3 * d^4) * \cos(f * x + e)^3) * \log(-\sin(f * x + e) + 1) + 2 * (2 * d^4 + 2 * (3 * c^4 - 12 * c^3 * d + 36 * c^2 * d^2 - 24 * c * d^3 + 8 * d^4) * \cos(f * x + e))^3 + (36 * c^2 * d^2 - 12 * c * d^3 + 7 * d^4) * \cos(f * x + e)^2 + (12 * c * d^3 - d^4) * \cos(f * x + e) * \sin(f * x + e)) / (a * f * \cos(f * x + e)^4 + a * f * \cos(f * x + e)^3)$$

3.210.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{a + a \sec(e + fx)} dx$$

$$= \frac{\int \frac{c^4 \sec(e + fx)}{\sec(e + fx) + 1} dx + \int \frac{d^4 \sec^5(e + fx)}{\sec(e + fx) + 1} dx + \int \frac{4cd^3 \sec^4(e + fx)}{\sec(e + fx) + 1} dx + \int \frac{6c^2 d^2 \sec^3(e + fx)}{\sec(e + fx) + 1} dx + \int \frac{4c^3 d \sec^2(e + fx)}{\sec(e + fx) + 1} dx}{a}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**4/(a+a*sec(f*x+e)),x)`

output `(Integral(c**4*sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(d**4*sec(e + f*x)**5/(sec(e + f*x) + 1), x) + Integral(4*c*d**3*sec(e + f*x)**4/(sec(e + f*x) + 1), x) + Integral(6*c**2*d**2*sec(e + f*x)**3/(sec(e + f*x) + 1), x) + Integral(4*c**3*d*sec(e + f*x)**2/(sec(e + f*x) + 1), x))/a`

3.210.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 596 vs. $2(174) = 348$.

Time = 0.22 (sec) , antiderivative size = 596, normalized size of antiderivative = 3.26

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{a + a \sec(e + fx)} dx$$

$$= \frac{d^4 \left(2 \left(\frac{9 \sin(fx+e)}{\cos(fx+e)+1} - \frac{16 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{15 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right) - \frac{9 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a} + \frac{9 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a} + \frac{6 \sin(fx+e)}{a(\cos(fx+e)+1)} \right) - 1}{a}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e)),x, algorithm="maxima")`

3.210.
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+a \sec(e+fx)} dx$$

output $\frac{1}{6}(d^4(2(9\sin(fx + e)/(\cos(fx + e) + 1) - 16\sin(fx + e)^3/(\cos(fx + e) + 1)^3 + 15\sin(fx + e)^5/(\cos(fx + e) + 1)^5)/(a - 3a\sin(fx + e)^2/(\cos(fx + e) + 1)^2 + 3a\sin(fx + e)^4/(\cos(fx + e) + 1)^4 - a\sin(fx + e)^6/(\cos(fx + e) + 1)^6) - 9\log(\sin(fx + e)/(\cos(fx + e) + 1) + 1)/a + 9\log(\sin(fx + e)/(\cos(fx + e) + 1) - 1)/a + 6\sin(fx + e)/(a(\cos(fx + e) + 1))) - 12*c*d^3(2(\sin(fx + e)/(\cos(fx + e) + 1) - 3\sin(fx + e)^3/(\cos(fx + e) + 1)^3)/(a - 2a\sin(fx + e)^2/(\cos(fx + e) + 1)^2 + a\sin(fx + e)^4/(\cos(fx + e) + 1)^4) - 3\log(\sin(fx + e)/(\cos(fx + e) + 1) + 1)/a + 3\log(\sin(fx + e)/(\cos(fx + e) + 1) - 1)/a + 2\sin(fx + e)/(a(\cos(fx + e) + 1))) - 36*c^2*d^2(\log(\sin(fx + e)/(\cos(fx + e) + 1) + 1)/a - \log(\sin(fx + e)/(\cos(fx + e) + 1) - 1)/a - 2\sin(fx + e)/((a - a\sin(fx + e)^2/(\cos(fx + e) + 1)^2)*(\cos(fx + e) + 1)) - \sin(fx + e)/(a(\cos(fx + e) + 1))) + 24*c^3*d(\log(\sin(fx + e)/(\cos(fx + e) + 1) + 1)/a - \log(\sin(fx + e)/(\cos(fx + e) + 1) - 1)/a - \sin(fx + e)/(a(\cos(fx + e) + 1))) + 6*c^4*\sin(fx + e)/(a(\cos(fx + e) + 1)))/f$

3.210.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.88

$$\int \frac{\sec(e + fx)(c + d\sec(e + fx))^4}{a + a\sec(e + fx)} dx$$

$$= \frac{3(8c^3d - 12c^2d^2 + 12cd^3 - 3d^4) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{a} - \frac{3(8c^3d - 12c^2d^2 + 12cd^3 - 3d^4) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{a} + \frac{6(c^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right))}{f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e)),x, algorithm="giac")`

output $\frac{1}{6}(3*(8*c^3*d - 12*c^2*d^2 + 12*c*d^3 - 3*d^4)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a - 3*(8*c^3*d - 12*c^2*d^2 + 12*c*d^3 - 3*d^4)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a + 6*(c^4*\tan(1/2*f*x + 1/2*e) - 4*c^3*d*\tan(1/2*f*x + 1/2*e) + 6*c^2*d^2*\tan(1/2*f*x + 1/2*e) - 4*c*d^3*\tan(1/2*f*x + 1/2*e) + d^4*\tan(1/2*f*x + 1/2*e))/a - 2*(36*c^2*d^2*\tan(1/2*f*x + 1/2*e)^5 - 36*c*d^3*\tan(1/2*f*x + 1/2*e)^5 + 15*d^4*\tan(1/2*f*x + 1/2*e)^5 - 72*c^2*d^2*\tan(1/2*f*x + 1/2*e)^3 + 48*c*d^3*\tan(1/2*f*x + 1/2*e)^3 - 16*d^4*\tan(1/2*f*x + 1/2*e)^3 + 36*c^2*d^2*\tan(1/2*f*x + 1/2*e) - 12*c*d^3*\tan(1/2*f*x + 1/2*e) + 9*d^4*\tan(1/2*f*x + 1/2*e))/((\tan(1/2*f*x + 1/2*e)^2 - 1)^3*a))/f$

3.210.9 Mupad [B] (verification not implemented)

Time = 14.17 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.15

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+a\sec(e+fx)} dx$$

$$= \frac{(12c^2d^2 - 12cd^3 + 5d^4) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-24c^2d^2 + 16cd^3 - \frac{16d^4}{3}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + (12c^2d^2 - 4cd^3)}{f \left(-a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 + 3a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 3a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + a\right)}$$

$$+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (c-d)^4}{af} + \frac{d \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (8c^3 - 12c^2d + 12cd^2 - 3d^3)}{af}$$

input `int((c + d/cos(e + f*x))^4/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`output `(tan(e/2 + (f*x)/2)*(3*d^4 - 4*c*d^3 + 12*c^2*d^2) + tan(e/2 + (f*x)/2)^5*(5*d^4 - 12*c*d^3 + 12*c^2*d^2) - tan(e/2 + (f*x)/2)^3*((16*d^4)/3 - 16*c*d^3 + 24*c^2*d^2))/(f*(a - 3*a*tan(e/2 + (f*x)/2)^2 + 3*a*tan(e/2 + (f*x)/2)^4 - a*tan(e/2 + (f*x)/2)^6)) + (tan(e/2 + (f*x)/2)*(c - d)^4)/(a*f) + (d*atanh(tan(e/2 + (f*x)/2))*(12*c*d^2 - 12*c^2*d + 8*c^3 - 3*d^3))/(a*f)`

3.211
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+a \sec(e+fx)} dx$$

3.211.1 Optimal result 1504
 3.211.2 Mathematica [A] (verified) 1504
 3.211.3 Rubi [A] (verified) 1505
 3.211.4 Maple [A] (verified) 1508
 3.211.5 Fracas [A] (verification not implemented) 1509
 3.211.6 Sympy [F] 1509
 3.211.7 Maxima [B] (verification not implemented) 1510
 3.211.8 Giac [A] (verification not implemented) 1510
 3.211.9 Mupad [B] (verification not implemented) 1511

3.211.1 Optimal result

Integrand size = 31, antiderivative size = 117

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+a \sec(e+fx)} dx$$

$$= \frac{3d(2c^2 - 2cd + d^2) \operatorname{arctanh}(\sin(e+fx))}{2af} + \frac{(c-d)(c+d \sec(e+fx))^2 \tan(e+fx)}{f(a+a \sec(e+fx))}$$

$$- \frac{d(4(c^2 - 3cd + d^2) + (2c - 3d)d \sec(e+fx)) \tan(e+fx)}{2af}$$

output `3/2*d*(2*c^2-2*c*d+d^2)*arctanh(sin(f*x+e))/a/f+(c-d)*(c+d*sec(f*x+e))^2*tan(f*x+e)/f/(a+a*sec(f*x+e))-1/2*d*(4*c^2-12*c*d+4*d^2+(2*c-3*d)*d*sec(f*x+e))*tan(f*x+e)/a/f`

3.211.2 Mathematica [A] (verified)

Time = 1.60 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+a \sec(e+fx)} dx$$

$$= \frac{3d(2c^2 - 2cd + d^2) \operatorname{arctanh}(\sin(e+fx)) + 2(c-d)^3 \tan\left(\frac{1}{2}(e+fx)\right) + d^2(6c - 2d + d \sec(e+fx)) \tan(e+fx)}{2af}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + a*Sec[e + f*x]),x]`

output $(3*d*(2*c^2 - 2*c*d + d^2)*\text{ArcTanh}[\text{Sin}[e + f*x]] + 2*(c - d)^3*\text{Tan}[(e + f*x)/2] + d^2*(6*c - 2*d + d*\text{Sec}[e + f*x])* \text{Tan}[e + f*x])/(2*a*f)$

3.211.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.84, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 4475, 109, 25, 27, 164, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a\sec(e+fx)+a} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))^3}{a\csc(e+fx+\frac{\pi}{2})+a} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{(c+d\sec(e+fx))^3}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 109

$$\frac{a^2 \tan(e+fx) \left(-\frac{\int \frac{a^2 d(3c-2d-(2c-3d)\sec(e+fx))(c+d\sec(e+fx)) d\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}}}{a^3} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^2}{a^2\sqrt{a\sec(e+fx)+a}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2 d(3c-2d-(2c-3d)\sec(e+fx))(c+d\sec(e+fx)) d\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}}}{a^3} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^2}{a^2\sqrt{a\sec(e+fx)+a}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{d \int \frac{(3c-2d-(2c-3d)\sec(e+fx))(c+d\sec(e+fx)) d\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}}}{a} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^2}{a^2\sqrt{a\sec(e+fx)+a}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

3.211. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx$

↓ 164

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{3}{2}(2c^2 - 2cd + d^2) \int \frac{1}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a}} d \sec(e + fx) + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (4(c^2 - 3cd + d^2) + d(2c - 3d))}{2a^2} \right)}{a} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{d \left(3(2c^2 - 2cd + d^2) \int \frac{1}{\frac{(a - a \sec(e + fx))a}{\sec(e + fx)a + a} - a} d \frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{\sec(e + fx)a + a}} + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (4(c^2 - 3cd + d^2) + d(2c - 3d))}{2a^2} \right)}{a} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (4(c^2 - 3cd + d^2) + d(2c - 3d) \sec(e + fx))}{2a^2} - \frac{3(2c^2 - 2cd + d^2) \arctan\left(\frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{a \sec(e + fx) + a}}\right)}{a} \right)}{a} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

```
input Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + a*Sec[e + f*x]),x]
```

```
output -((a^2*(-(((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(a^2*Sqrt[a + a*Sec[e + f*x]])) + (d*((-3*(2*c^2 - 2*c*d + d^2)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])/a + (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(4*(c^2 - 3*c*d + d^2) + (2*c - 3*d)*d*Sec[e + f*x]))/(2*a^2)))/a)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.211. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+a \sec(e+fx)} dx$

3.211.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 109 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 164 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))*((g_) + (h_)*(x_)), x_] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a
^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]]))
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x
], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[
b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || In
tegerQ[m - 1/2])
```

3.211.4 Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.55

method	result
parallelrisch	$\frac{-3(c^2 - cd + \frac{1}{2}d^2)(1 + \cos(2fx + 2e))d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 3(c^2 - cd + \frac{1}{2}d^2)(1 + \cos(2fx + 2e))d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af(1 + \cos(2fx + 2e))}$
derivativedivides	$c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 3c^2d \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)cd^2 - d^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{d^3}{2\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{3d(2c^2 - 2cd + d^2) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)}{2}$
default	$c^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 3c^2d \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)cd^2 - d^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \frac{d^3}{2\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{3d(2c^2 - 2cd + d^2) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)}{2}$
norman	$\frac{(c^3 - 3c^2d + 3cd^2 - d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{af} + \frac{(3c^3 - 9c^2d + 21cd^2 - 7d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} - \frac{3(c^3 - 3c^2d + 5cd^2 - 2d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{af} - \frac{(c^3 - 3c^2d + 3cd^2 - d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af}$
risch	$\frac{i(2c^3e^{4i(fx+e)} - 6c^2de^{4i(fx+e)} + 6cd^2e^{4i(fx+e)} - 3d^3e^{4i(fx+e)} + 6cd^2e^{3i(fx+e)} - 3d^3e^{3i(fx+e)} + 4c^3e^{2i(fx+e)} - 12c^2de^{2i(fx+e)} + 12cd^2e^{2i(fx+e)} - 3d^3e^{2i(fx+e)})}{fa(e^{i(fx+e)} + 1)(1 + e^{2i(fx+e)})}$

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE
)
```

```
output (-3*(c^2-c*d+1/2*d^2)*(1+cos(2*f*x+2*e))*d*ln(tan(1/2*f*x+1/2*e)-1)+3*(c^2
-c*d+1/2*d^2)*(1+cos(2*f*x+2*e))*d*ln(tan(1/2*f*x+1/2*e)+1)+tan(1/2*f*x+1/
2*e)*((c^3-3*c^2*d+6*c*d^2-2*d^3)*cos(2*f*x+2*e)+(6*c*d^2-d^3)*cos(f*x+e)+
c^3-3*c^2*d+6*c*d^2-d^3)/a/f/(1+cos(2*f*x+2*e))
```

$$3.211. \int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx$$

3.211.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.85

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx$$

$$= \frac{3((2c^2d-2cd^2+d^3)\cos(fx+e)^3 + (2c^2d-2cd^2+d^3)\cos(fx+e)^2)\log(\sin(fx+e)+1) - 3((2c^2d-2cd^2+d^3)\cos(fx+e)^3 + (2c^2d-2cd^2+d^3)\cos(fx+e)^2)\log(-\sin(fx+e)+1) + 2(d^3+2(c^3-3c^2d+6cd^2-2d^3)\cos(fx+e)^2 + (6cd^2-d^3)\cos(fx+e))\sin(fx+e)}{(a^2\cos(fx+e)^3 + a^2\cos(fx+e)^2)}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e)),x, algorithm="fricas")
```

```
output 1/4*(3*((2*c^2*d - 2*c*d^2 + d^3)*cos(f*x + e)^3 + (2*c^2*d - 2*c*d^2 + d^3)*cos(f*x + e)^2)*log(sin(f*x + e) + 1) - 3*((2*c^2*d - 2*c*d^2 + d^3)*cos(f*x + e)^3 + (2*c^2*d - 2*c*d^2 + d^3)*cos(f*x + e)^2)*log(-sin(f*x + e) + 1) + 2*(d^3 + 2*(c^3 - 3*c^2*d + 6*c*d^2 - 2*d^3)*cos(f*x + e)^2 + (6*c*d^2 - d^3)*cos(f*x + e))*sin(f*x + e))/(a*f*cos(f*x + e)^3 + a*f*cos(f*x + e)^2)
```

3.211.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx$$

$$= \frac{\int \frac{c^3 \sec(e+fx)}{\sec(e+fx)+1} dx + \int \frac{d^3 \sec^4(e+fx)}{\sec(e+fx)+1} dx + \int \frac{3cd^2 \sec^3(e+fx)}{\sec(e+fx)+1} dx + \int \frac{3c^2d \sec^2(e+fx)}{\sec(e+fx)+1} dx}{a}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))**3/(a+a*sec(f*x+e)),x)
```

```
output (Integral(c**3*sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(d**3*sec(e + f*x)**4/(sec(e + f*x) + 1), x) + Integral(3*c*d**2*sec(e + f*x)**3/(sec(e + f*x) + 1), x) + Integral(3*c**2*d*sec(e + f*x)**2/(sec(e + f*x) + 1), x))/a
```

3.211.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 388 vs. $2(114) = 228$.

Time = 0.22 (sec) , antiderivative size = 388, normalized size of antiderivative = 3.32

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx =$$

$$d^3 \left(\frac{2 \left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - \frac{3 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a - \frac{2a \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{a \sin(fx+e)^4}{(\cos(fx+e)+1)^4}} - \frac{3 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a} + \frac{3 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a} + \frac{2 \sin(fx+e)}{a(\cos(fx+e)+1)} \right) + 6cd^2 \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a} \right)$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `-1/2*(d^3*(2*(sin(f*x + e)/(cos(f*x + e) + 1) - 3*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a - 2*a*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a*sin(f*x + e)^4/(cos(f*x + e) + 1)^4) - 3*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a + 3*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a + 2*sin(f*x + e)/(a*(cos(f*x + e) + 1))) + 6*c*d^2*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a - 2*sin(f*x + e)/((a - a*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1)) - sin(f*x + e)/(a*(cos(f*x + e) + 1))) - 6*c^2*d*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a - sin(f*x + e)/(a*(cos(f*x + e) + 1))) - 2*c^3*sin(f*x + e)/(a*(cos(f*x + e) + 1)))/f`

3.211.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.87

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx$$

$$\frac{3(2c^2d-2cd^2+d^3) \log\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+1\right|}{a} - \frac{3(2c^2d-2cd^2+d^3) \log\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-1\right|}{a} + \frac{2(c^3 \tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-3c^2d \tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+cd^2 \tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-d^3)}{a}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e)),x, algorithm="giac")`

3.211. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx$

output $1/2*(3*(2*c^2*d - 2*c*d^2 + d^3)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a - 3*(2*c^2*d - 2*c*d^2 + d^3)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a + 2*(c^3*\tan(1/2*f*x + 1/2*e) - 3*c^2*d*\tan(1/2*f*x + 1/2*e) + 3*c*d^2*\tan(1/2*f*x + 1/2*e) - d^3*\tan(1/2*f*x + 1/2*e))/a - 2*(6*c*d^2*\tan(1/2*f*x + 1/2*e)^3 - 3*d^3*\tan(1/2*f*x + 1/2*e)^3 - 6*c*d^2*\tan(1/2*f*x + 1/2*e) + d^3*\tan(1/2*f*x + 1/2*e))/((\tan(1/2*f*x + 1/2*e)^2 - 1)^2*a))/f$

3.211.9 Mupad [B] (verification not implemented)

Time = 13.68 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.19

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+a\sec(e+fx)} dx \\ &= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (6cd^2 - d^3) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (6cd^2 - 3d^3)}{f \left(a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2a \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + a \right)} \\ & \quad + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (c-d)^3}{af} + \frac{3d \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (2c^2 - 2cd + d^2)}{af} \end{aligned}$$

input `int((c + d/cos(e + f*x))^3/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output $(\tan(e/2 + (f*x)/2)*(6*c*d^2 - d^3) - \tan(e/2 + (f*x)/2)^3*(6*c*d^2 - 3*d^3))/(f*(a - 2*a*\tan(e/2 + (f*x)/2)^2 + a*\tan(e/2 + (f*x)/2)^4) + (\tan(e/2 + (f*x)/2)*(c - d)^3)/(a*f) + (3*d*\operatorname{atanh}(\tan(e/2 + (f*x)/2))*(2*c^2 - 2*c*d + d^2))/(a*f)$

3.212 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+a \sec(e+fx)} dx$

3.212.1 Optimal result 1512
 3.212.2 Mathematica [B] (verified) 1512
 3.212.3 Rubi [B] (verified) 1513
 3.212.4 Maple [A] (verified) 1516
 3.212.5 Fracas [B] (verification not implemented) 1516
 3.212.6 Sympy [F] 1517
 3.212.7 Maxima [B] (verification not implemented) 1517
 3.212.8 Giac [A] (verification not implemented) 1518
 3.212.9 Mupad [B] (verification not implemented) 1518

3.212.1 Optimal result

Integrand size = 31, antiderivative size = 68

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+a \sec(e+fx)} dx = \frac{(2c-d) \operatorname{darctanh}(\sin(e+fx))}{af} + \frac{d^2 \tan(e+fx)}{af} + \frac{(c-d)^2 \tan(e+fx)}{f(a+a \sec(e+fx))}$$

output `(2*c-d)*d*arctanh(sin(f*x+e))/a/f+d^2*tan(f*x+e)/a/f+(c-d)^2*tan(f*x+e)/f/(a+a*sec(f*x+e))`

3.212.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 237 vs. 2(68) = 136.

Time = 2.66 (sec) , antiderivative size = 237, normalized size of antiderivative = 3.49

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+a \sec(e+fx)} dx$$

$$= \frac{2 \cos\left(\frac{1}{2}(e+fx)\right) \cos(e+fx)(c+d \sec(e+fx))^2 \left((c-d)^2 \sec\left(\frac{e}{2}\right) \sin\left(\frac{fx}{2}\right) + d \cos\left(\frac{1}{2}(e+fx)\right) \left(-((2c-d) \operatorname{darctanh}(\sin(e+fx))) \right) \right)}{af^2}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + a*Sec[e + f*x]),x]`

output $(2*\text{Cos}[(e + f*x)/2]*\text{Cos}[e + f*x]*(c + d*\text{Sec}[e + f*x])^2*((c - d)^2*\text{Sec}[e/2]*\text{Sin}[(f*x)/2] + d*\text{Cos}[(e + f*x)/2]*(-(2*c - d)*(\text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] - \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]])) + (d*\text{Sin}[f*x])/((\text{Cos}[e/2] - \text{Sin}[e/2])*(\text{Cos}[e/2] + \text{Sin}[e/2])*(\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2])*(\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]))))/(a*f*(d + c*\text{Cos}[e + f*x])^2*(1 + \text{Sec}[e + f*x]))$

3.212.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 169 vs. $2(68) = 136$.

Time = 0.34 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.49, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3042, 4475, 100, 27, 90, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a \sec(e + fx) + a} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})(c + d \csc(e + fx + \frac{\pi}{2}))^2}{a \csc(e + fx + \frac{\pi}{2}) + a} dx$$

↓ 4475

$$\frac{a^2 \tan(e + fx) \int \frac{(c + d \sec(e + fx))^2}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{3/2}} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 100

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^3 d(2c - d + d \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a}} d \sec(e + fx)}{a^4} - \frac{(c - d)^2 \sqrt{a - a \sec(e + fx)}}{a^2 \sqrt{a \sec(e + fx) + a}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 27

$$\frac{a^2 \tan(e + fx) \left(\frac{d \int \frac{2c - d + d \sec(e + fx)}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a}} d \sec(e + fx)}{a} - \frac{(c - d)^2 \sqrt{a - a \sec(e + fx)}}{a^2 \sqrt{a \sec(e + fx) + a}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 90

3.212. $\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + a \sec(e + fx)} dx$

$$\begin{aligned}
 & \frac{a^2 \tan(e + fx) \left(\frac{d \left((2c-d) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) - \frac{d \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a^2} \right)}{a} - \frac{(c-d)^2 \sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 45 \\
 & \frac{a^2 \tan(e + fx) \left(\frac{d \left(2(2c-d) \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} - \frac{d \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a^2} \right)}{a} - \frac{(c-d)^2 \sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow 218 \\
 & \frac{a^2 \tan(e + fx) \left(\frac{d \left(-\frac{d \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}{a^2} - \frac{2(2c-d) \arctan\left(\frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}}\right)}{a} \right)}{a} - \frac{(c-d)^2 \sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + a*Sec[e + f*x]),x]`

output `-((a^2*(-(((c - d)^2*Sqrt[a - a*Sec[e + f*x]])/(a^2*Sqrt[a + a*Sec[e + f*x]]))) + (d*((-2*(2*c - d)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]])/a - (d*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]/a^2))/a)*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])`

3.212.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

- rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`
- rule 100 `Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d^2*(d*e - c*f)*(n + 1))), x] - Simp[1/(d^2*(d*e - c*f)*(n + 1)) Int[(c + d*x)^(n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n + 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.212.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.51

method	result
parallelrisc	$\frac{-2\left(c-\frac{d}{2}\right)\cos(fx+e)d\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)+2\left(c-\frac{d}{2}\right)\cos(fx+e)d\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)+\tan\left(\frac{fx}{2}+\frac{e}{2}\right)\left((c^2-2cd+2d^2)\cos(fx+e)-d\right)}{af\cos(fx+e)}$
derivativedivides	$\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)c^2-2\tan\left(\frac{fx}{2}+\frac{e}{2}\right)cd+\tan\left(\frac{fx}{2}+\frac{e}{2}\right)d^2-\frac{d^2}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1}-d(2c-d)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)-\frac{d^2}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1}+d(2c-d)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{fa}$
default	$\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)c^2-2\tan\left(\frac{fx}{2}+\frac{e}{2}\right)cd+\tan\left(\frac{fx}{2}+\frac{e}{2}\right)d^2-\frac{d^2}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1}-d(2c-d)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)-\frac{d^2}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1}+d(2c-d)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{fa}$
norman	$\frac{\frac{(c^2-2cd+d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{af}+\frac{(c^2-2cd+3d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{af}-\frac{2(c^2-2cd+2d^2)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{af}}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2-1\right)^2}+\frac{d(2c-d)\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{af}$
risc	$\frac{2i(c^2e^{2i(fx+e)}-2cde^{2i(fx+e)}+d^2e^{2i(fx+e)}+d^2e^{i(fx+e)}+c^2-2cd+2d^2)}{fa(1+e^{2i(fx+e)})(e^{i(fx+e)}+1)}+\frac{2d\ln(e^{i(fx+e)}+i)c}{af}-\frac{d^2\ln(e^{i(fx+e)}+i)}{af}$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `(-2*(c-1/2*d)*cos(f*x+e)*d*ln(tan(1/2*f*x+1/2*e)-1)+2*(c-1/2*d)*cos(f*x+e)*d*ln(tan(1/2*f*x+1/2*e)+1)+tan(1/2*f*x+1/2*e)*((c^2-2*c*d+2*d^2)*cos(f*x+e)+d^2))/a/f/cos(f*x+e)`

3.212.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 155 vs. 2(68) = 136.

Time = 0.27 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.28

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{a+a\sec(e+fx)} dx$$

$$= \frac{((2cd-d^2)\cos(fx+e))^2+(2cd-d^2)\cos(fx+e)\log(\sin(fx+e)+1)-((2cd-d^2)\cos(fx+e))^2}{2(af\cos(fx+e))^2}-$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e)),x, algorithm="fricas")`

3.212.
$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{a+a\sec(e+fx)} dx$$

output $1/2*((2*c*d - d^2)*\cos(f*x + e)^2 + (2*c*d - d^2)*\cos(f*x + e))*\log(\sin(f*x + e) + 1) - ((2*c*d - d^2)*\cos(f*x + e)^2 + (2*c*d - d^2)*\cos(f*x + e))*\log(-\sin(f*x + e) + 1) + 2*(d^2 + (c^2 - 2*c*d + 2*d^2)*\cos(f*x + e))*\sin(f*x + e)/(a*f*\cos(f*x + e)^2 + a*f*\cos(f*x + e))$

3.212.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + a \sec(e + fx)} dx$$

$$= \frac{\int \frac{c^2 \sec(e+fx)}{\sec(e+fx)+1} dx + \int \frac{d^2 \sec^3(e+fx)}{\sec(e+fx)+1} dx + \int \frac{2cd \sec^2(e+fx)}{\sec(e+fx)+1} dx}{a}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**2/(a+a*sec(f*x+e)),x)`

output `(Integral(c**2*sec(e + f*x)/(sec(e + f*x) + 1), x) + Integral(d**2*sec(e + f*x)**3/(sec(e + f*x) + 1), x) + Integral(2*c*d*sec(e + f*x)**2/(sec(e + f*x) + 1), x))/a`

3.212.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 223 vs. $2(68) = 136$.

Time = 0.22 (sec) , antiderivative size = 223, normalized size of antiderivative = 3.28

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + a \sec(e + fx)} dx =$$

$$\frac{d^2 \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a} - \frac{2 \sin(fx+e)}{\left(a - \frac{a \sin(fx+e)^2}{(\cos(fx+e)+1)^2}\right)(\cos(fx+e)+1)} - \frac{\sin(fx+e)}{a(\cos(fx+e)+1)} \right) - 2cd \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1}\right)}{\cos(fx+e)+1} \right)}{f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output $-(d^2*(\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/a - \log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/a - 2*\sin(f*x + e)/((a - a*\sin(f*x + e)^2/(\cos(f*x + e) + 1)^2)*(\cos(f*x + e) + 1)) - \sin(f*x + e)/(a*(\cos(f*x + e) + 1))) - 2*c*d*(\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/a - \log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/a - \sin(f*x + e)/(a*(\cos(f*x + e) + 1))) - c^2*\sin(f*x + e)/(a*(\cos(f*x + e) + 1)))/f$

3.212.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.00

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + a \sec(e + fx)} dx$$

$$= \frac{(2cd - d^2) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a} - \frac{(2cd - d^2) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a} - \frac{2d^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)a} + \frac{c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e) - 2cd \tan(\frac{1}{2}fx + \frac{1}{2}e)}{a}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e)),x, algorithm="giac")`

output $((2*c*d - d^2)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a - (2*c*d - d^2)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a - 2*d^2*\tan(1/2*f*x + 1/2*e)/((\tan(1/2*f*x + 1/2*e)^2 - 1)*a) + (c^2*\tan(1/2*f*x + 1/2*e) - 2*c*d*\tan(1/2*f*x + 1/2*e) + d^2*\tan(1/2*f*x + 1/2*e))/a)/f$

3.212.9 Mupad [B] (verification not implemented)

Time = 13.53 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.25

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + a \sec(e + fx)} dx = \frac{\tan(\frac{e}{2} + \frac{fx}{2}) (c - d)^2}{af} + \frac{2d^2 \tan(\frac{e}{2} + \frac{fx}{2})}{f (a - a \tan(\frac{e}{2} + \frac{fx}{2})^2)}$$

$$+ \frac{2d \operatorname{atanh}(\tan(\frac{e}{2} + \frac{fx}{2})) (2c - d)}{af}$$

input `int((c + d/cos(e + f*x))^2/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output $(\tan(e/2 + (f*x)/2)*(c - d)^2)/(a*f) + (2*d^2*\tan(e/2 + (f*x)/2))/(f*(a - a*\tan(e/2 + (f*x)/2)^2)) + (2*d*\operatorname{atanh}(\tan(e/2 + (f*x)/2))*(2*c - d))/(a*f)$

3.212. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{a+a\sec(e+fx)} dx$

3.213 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+a \sec(e+fx)} dx$

3.213.1 Optimal result 1519
 3.213.2 Mathematica [B] (verified) 1519
 3.213.3 Rubi [A] (verified) 1520
 3.213.4 Maple [A] (verified) 1521
 3.213.5 Fricas [A] (verification not implemented) 1522
 3.213.6 Sympy [F] 1522
 3.213.7 Maxima [B] (verification not implemented) 1523
 3.213.8 Giac [A] (verification not implemented) 1523
 3.213.9 Mupad [B] (verification not implemented) 1524

3.213.1 Optimal result

Integrand size = 29, antiderivative size = 43

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+a \sec(e+fx)} dx = \frac{\operatorname{darctanh}(\sin(e+fx))}{af} + \frac{(c-d) \tan(e+fx)}{f(a+a \sec(e+fx))}$$

output `d*arctanh(sin(f*x+e))/a/f+(c-d)*tan(f*x+e)/f/(a+a*sec(f*x+e))`

3.213.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 109 vs. 2(43) = 86.

Time = 0.90 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.53

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+a \sec(e+fx)} dx = \frac{2 \cos\left(\frac{1}{2}(e+fx)\right) \left(d \cos\left(\frac{1}{2}(e+fx)\right) \left(-\log\left(\cos\left(\frac{1}{2}(e+fx)\right) - \sin\left(\frac{1}{2}(e+fx)\right)\right) + \log\left(\cos\left(\frac{1}{2}(e+fx)\right) + \sin\left(\frac{1}{2}(e+fx)\right)\right)\right) + (c-d) \operatorname{Sec}[e/2] \operatorname{Sin}[(f*x)/2]}{af(1 + \cos(e+fx))}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output `(2*Cos[(e + f*x)/2]*(d*Cos[(e + f*x)/2]*(-Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]]) + (c - d)*Sec[e/2]*Sin[(f*x)/2))/(a*f*(1 + Cos[e + f*x]))`

3.213. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+a \sec(e+fx)} dx$

3.213.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.172$, Rules used = {3042, 4486, 3042, 4257, 4281}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c+d\sec(e+fx))}{a\sec(e+fx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))}{a\csc(e+fx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{4486} \\
 & (c-d) \int \frac{\sec(e+fx)}{\sec(e+fx)a+a} dx + \frac{d \int \sec(e+fx) dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & (c-d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\csc(e+fx+\frac{\pi}{2})a+a} dx + \frac{d \int \csc(e+fx+\frac{\pi}{2}) dx}{a} \\
 & \quad \downarrow \text{4257} \\
 & (c-d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\csc(e+fx+\frac{\pi}{2})a+a} dx + \frac{\operatorname{darctanh}(\sin(e+fx))}{af} \\
 & \quad \downarrow \text{4281} \\
 & \frac{\operatorname{darctanh}(\sin(e+fx))}{af} + \frac{(c-d)\tan(e+fx)}{f(a\sec(e+fx)+a)}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + a*Sec[e + f*x]),x]`

output `(d*ArcTanh[Sin[e + f*x]])/(a*f) + ((c - d)*Tan[e + f*x])/(f*(a + a*Sec[e + f*x]))`

3.213.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4281 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)*(b_.) + (a_.)], x_Symbol] := Simp[-Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4486 `Int[(csc[(e_.) + (f_.)*(x_)*(csc[(e_.) + (f_.)*(x_)*(B_.) + (A_.)])/(csc[(e_.) + (f_.)*(x_)*(b_.) + (a_.)], x_Symbol] := Simp[B/b Int[Csc[e + f*x], x] + Simp[(A*b - a*B)/b Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0]`

3.213.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.23

method	result	size
parallelrisch	$\frac{-\ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)d + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)d + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)(c-d)}{af}$	53
derivativedivides	$\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - d \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)d + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)d}{fa}$	61
default	$\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - d \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)d + \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)d}{fa}$	61
risch	$\frac{2ic}{fa(e^{i(fx+e)}+1)} - \frac{2id}{fa(e^{i(fx+e)}+1)} + \frac{d \ln(e^{i(fx+e)}+i)}{af} - \frac{d \ln(e^{i(fx+e)}-i)}{af}$	91
norman	$\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{af} - \frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{af} + \frac{d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{af} - \frac{d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{af}$	105

input `int(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output $(-\ln(\tan(1/2*f*x+1/2*e)-1)*d+\ln(\tan(1/2*f*x+1/2*e)+1)*d+\tan(1/2*f*x+1/2*e)*(c-d))/a/f$

3.213.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.72

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{a+a\sec(e+fx)} dx$$

$$= \frac{(d\cos(fx+e)+d)\log(\sin(fx+e)+1) - (d\cos(fx+e)+d)\log(-\sin(fx+e)+1) + 2(c-d)\sin(fx+e)}{2(af\cos(fx+e)+af)}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="fricas")`

output $1/2*((d*\cos(f*x + e) + d)*\log(\sin(f*x + e) + 1) - (d*\cos(f*x + e) + d)*\log(-\sin(f*x + e) + 1) + 2*(c - d)*\sin(f*x + e))/(a*f*\cos(f*x + e) + a*f)$

3.213.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{a+a\sec(e+fx)} dx = \frac{\int \frac{c\sec(e+fx)}{\sec(e+fx)+1} dx + \int \frac{d\sec^2(e+fx)}{\sec(e+fx)+1} dx}{a}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e)),x)`

output $(\text{Integral}(c*\sec(e + f*x)/(\sec(e + f*x) + 1), x) + \text{Integral}(d*\sec(e + f*x)*2/(\sec(e + f*x) + 1), x))/a$

3.213.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(43) = 86$.

Time = 0.20 (sec) , antiderivative size = 99, normalized size of antiderivative = 2.30

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{a + a \sec(e + fx)} dx$$

$$= \frac{d \left(\frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a} - \frac{\log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a} - \frac{\sin(fx+e)}{a(\cos(fx+e)+1)} \right) + \frac{c \sin(fx+e)}{a(\cos(fx+e)+1)}}{f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="maxima")`

output `(d*(log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a - log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a - sin(f*x + e)/(a*(cos(f*x + e) + 1))) + c*sin(f*x + e)/(a*(cos(f*x + e) + 1)))/f`

3.213.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.63

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{a + a \sec(e + fx)} dx$$

$$= \frac{\frac{d \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) + 1|)}{a} - \frac{d \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) - 1|)}{a} + \frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e) - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}{a}}{f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e)),x, algorithm="giac")`

output `(d*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a - d*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a + (c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/a)/f`

3.213.9 Mupad [B] (verification not implemented)

Time = 13.87 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{a + a \sec(e + fx)} dx = \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (c - d)}{a f} + \frac{2 d \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a f}$$

input `int((c + d/cos(e + f*x))/(cos(e + f*x)*(a + a/cos(e + f*x))),x)`

output `(tan(e/2 + (f*x)/2)*(c - d))/(a*f) + (2*d*atanh(tan(e/2 + (f*x)/2)))/(a*f)`

$$3.214 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx$$

3.214.1 Optimal result 1525
 3.214.2 Mathematica [C] (verified) 1525
 3.214.3 Rubi [A] (verified) 1526
 3.214.4 Maple [A] (verified) 1528
 3.214.5 Fricas [A] (verification not implemented) 1528
 3.214.6 Sympy [F] 1529
 3.214.7 Maxima [F(-2)] 1529
 3.214.8 Giac [A] (verification not implemented) 1530
 3.214.9 Mupad [B] (verification not implemented) 1530

3.214.1 Optimal result

Integrand size = 31, antiderivative size = 83

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx = -\frac{2d \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a(c-d)^{3/2} \sqrt{c+df}} + \frac{\tan(e+fx)}{(c-d)f(a+a \sec(e+fx))}$$

output

```
-2*d*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a/(c-d)^(3/2)/f/(c+d)^(1/2)+tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))
```

3.214.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.06 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.93

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx$$

$$= \frac{2 \cos\left(\frac{1}{2}(e+fx)\right) \left(\frac{2d \arctan\left(\frac{(i \cos(e)+\sin(e))(c \sin(e)+(-d+c \cos(e)) \tan\left(\frac{fx}{2}\right))}{\sqrt{c^2-d^2} \sqrt{(\cos(e)-i \sin(e))^2}}\right) \cos\left(\frac{1}{2}(e+fx)\right) (i \cos(e)+\sin(e))}{\sqrt{c^2-d^2} \sqrt{(\cos(e)-i \sin(e))^2}} + \sec\left(\frac{e}{2}\right) \sin\left(\frac{fx}{2}\right) \right)}{a(c-d)f(1+\cos(e+fx))}$$

3.214. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])),x]`

output `(2*Cos[(e + f*x)/2]*((2*d*ArcTan[(I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]))*Cos[(e + f*x)/2]*(I*Cos[e] + Sin[e]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + Sec[e/2]*Sin[(f*x)/2))/(a*(c - d)*f*(1 + Cos[e + f*x]))`

3.214.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.86, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4475, 107, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)(c + d \sec(e+fx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)(c + d \csc(e+fx + \frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{4475} \\
 & -\frac{a^2 \tan(e+fx) \int \frac{1}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx) + a}} \\
 & \quad \downarrow \text{107} \\
 & \frac{a^2 \tan(e+fx) \left(-\frac{d \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{a(c-d)} - \frac{\sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx) + a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx) + a}} \\
 & \quad \downarrow \text{104} \\
 & \frac{a^2 \tan(e+fx) \left(-\frac{2d \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{a(c-d)} - \frac{\sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx) + a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx) + a}} \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

3.214. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx$

$$\frac{a^2 \tan(e + fx) \left(-\frac{2d \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}\right)}{a^2(c-d)^{3/2}\sqrt{c+d}} - \frac{\sqrt{a-a \sec(e+fx)}}{a^2(c-d)\sqrt{a \sec(e+fx)+a}} \right)}{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])),x]`

output `-((a^2*((-2*d*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]))/(a^2*(c - d)^(3/2)*Sqrt[c + d]) - Sqrt[a - a*Sec[e + f*x]]/(a^2*(c - d)*Sqrt[a + a*Sec[e + f*x]]))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])`

3.214.3.1 Defintions of rubi rules used

rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 107 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[
^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x
], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[
b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || In
tegerQ[m - 1/2])
```

3.214.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{c-d} - \frac{2d \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c-d)\sqrt{(c+d)(c-d)}}$	74
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{c-d} - \frac{2d \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c-d)\sqrt{(c+d)(c-d)}}$	74
risch	$\frac{2i}{fa(c-d)(e^{i(fx+e)}+1)} + \frac{d \ln\left(e^{i(fx+e)} + \frac{-ic^2+id^2+\sqrt{c^2-d^2}d}{c\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}(c-d)fa} - \frac{d \ln\left(e^{i(fx+e)} + \frac{ic^2-id^2+\sqrt{c^2-d^2}d}{\sqrt{c^2-d^2}c}\right)}{\sqrt{c^2-d^2}(c-d)fa}$	188

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f/a*(tan(1/2*f*x+1/2*e)/(c-d)-2*d/(c-d)/((c+d)*(c-d))^(1/2)*arctanh((c-d)
)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))
```

3.214.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 353, normalized size of antiderivative = 4.25

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))} dx$$

$$= \left[\frac{\sqrt{c^2 - d^2}(d \cos(fx + e) + d) \log\left(\frac{2cd \cos(fx+e) - (c^2 - 2d^2) \cos(fx+e)^2 + 2\sqrt{c^2-d^2}(d \cos(fx+e) + c) \sin(fx+e) + 2c^2 - d^2}{c^2 \cos(fx+e)^2 + 2cd \cos(fx+e) + d^2}\right)}{2((ac^3 - ac^2d - acd^2 + ad^3)f \cos(fx + e) + (ac^3 - ac^2d - acd^2 + ad^3)f)} - \frac{\sqrt{-c^2 + d^2}(d \cos(fx + e) + d) \arctan\left(-\frac{\sqrt{-c^2+d^2}(d \cos(fx+e)+c)}{(c^2-d^2) \sin(fx+e)}\right) - (c^2 - d^2) \sin(fx + e)}{(ac^3 - ac^2d - acd^2 + ad^3)f \cos(fx + e) + (ac^3 - ac^2d - acd^2 + ad^3)f} \right]$$

3.214. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `[-1/2*(sqrt(c^2 - d^2)*(d*cos(f*x + e) + d)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) - 2*(c^2 - d^2)*sin(f*x + e))/((a*c^3 - a*c^2*d - a*c*d^2 + a*d^3)*f*cos(f*x + e) + (a*c^3 - a*c^2*d - a*c*d^2 + a*d^3)*f), -(sqrt(-c^2 + d^2)*(d*cos(f*x + e) + d)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) - (c^2 - d^2)*sin(f*x + e))/((a*c^3 - a*c^2*d - a*c*d^2 + a*d^3)*f*cos(f*x + e) + (a*c^3 - a*c^2*d - a*c*d^2 + a*d^3)*f)]`

3.214.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))} dx = \frac{\int \frac{\sec(e + fx)}{c \sec(e + fx) + c + d \sec^2(e + fx) + d \sec(e + fx)} dx}{a}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x)`

output `Integral(sec(e + f*x)/(c*sec(e + f*x) + c + d*sec(e + f*x)**2 + d*sec(e + f*x)), x)/a`

3.214.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.214. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))} dx$

3.214.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.33

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))} dx$$

$$= \frac{2 \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e) - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-c^2+d^2}} \right) \right) d}{(ac-ad)\sqrt{-c^2+d^2}} + \frac{\tan(\frac{1}{2} fx + \frac{1}{2} e)}{ac-ad}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="giac")
```

```
output (2*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*d/((a*c - a*d)*sqrt(-c^2 + d^2)) + tan(1/2*f*x + 1/2*e)/(a*c - a*d))/f
```

3.214.9 Mupad [B] (verification not implemented)

Time = 13.58 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.33

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))} dx$$

$$= \frac{\tan(\frac{e}{2} + \frac{fx}{2})}{af(c-d)} - \frac{2d \operatorname{atanh} \left(\frac{\sin(\frac{e}{2} + \frac{fx}{2}) c^2 - 2 \sin(\frac{e}{2} + \frac{fx}{2}) cd + \sin(\frac{e}{2} + \frac{fx}{2}) d^2}{\cos(\frac{e}{2} + \frac{fx}{2}) \sqrt{c+d} (c-d)^{3/2}} \right)}{af \sqrt{c+d} (c-d)^{3/2}}$$

```
input int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c + d/cos(e + f*x))),x)
```

```
output tan(e/2 + (f*x)/2)/(a*f*(c - d)) - (2*d*atanh((c^2*sin(e/2 + (f*x)/2) + d^2*sin(e/2 + (f*x)/2) - 2*c*d*sin(e/2 + (f*x)/2))/(cos(e/2 + (f*x)/2)*(c + d)^(1/2)*(c - d)^(3/2)))/(a*f*(c + d)^(1/2)*(c - d)^(3/2))
```

$$3.215 \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$$

3.215.1 Optimal result 1531
 3.215.2 Mathematica [C] (verified) 1531
 3.215.3 Rubi [A] (verified) 1532
 3.215.4 Maple [A] (verified) 1535
 3.215.5 Fracas [B] (verification not implemented) 1536
 3.215.6 Sympy [F] 1537
 3.215.7 Maxima [F(-2)] 1537
 3.215.8 Giac [A] (verification not implemented) 1537
 3.215.9 Mupad [B] (verification not implemented) 1538

3.215.1 Optimal result

Integrand size = 31, antiderivative size = 145

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$$

$$= -\frac{2d(2c+d) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a(c-d)^{5/2}(c+d)^{3/2}f} + \frac{(c+2d) \tan(e+fx)}{(c-d)^2(c+d)f(a+a \sec(e+fx))}$$

$$- \frac{d \tan(e+fx)}{(c^2-d^2)f(a+a \sec(e+fx))(c+d \sec(e+fx))}$$

```
output -2*d*(2*c+d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a/(c-d)^(
5/2)/(c+d)^(3/2)/f+(c+2*d)*tan(f*x+e)/(c-d)^2/(c+d)/f/(a+a*sec(f*x+e))-d*t
an(f*x+e)/(c^2-d^2)/f/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))
```

3.215.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.14 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.97

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$$

$$= \frac{2 \cos\left(\frac{1}{2}(e+fx)\right) (d+c \cos(e+fx)) \sec^3(e+fx) \left(\frac{2d(2c+d) \arctan\left(\frac{(i \cos(e)+\sin(e))(c \sin(e)+(-d+c \cos(e)) \tan\left(\frac{fx}{2}\right))}{\sqrt{c^2-d^2} \sqrt{(\cos(e)-i \sin(e))^2}}\right)}{(c+d) \sqrt{c^2-d^2} \sqrt{(\cos(e)-i \sin(e))}} \right) \cos\left(\frac{1}{2}(e+fx)\right)}{a(c-d)^2 f (1 + \sec(e+fx))}$$

3.215. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^2),x]`

output $(2*\text{Cos}[(e + f*x)/2]*(d + c*\text{Cos}[e + f*x])* \text{Sec}[e + f*x]^3*((2*d*(2*c + d)*\text{ArcTan}[(I*\text{Cos}[e] + \text{Sin}[e])*(c*\text{Sin}[e] + (-d + c*\text{Cos}[e])* \text{Tan}[(f*x)/2]))/(\text{Sqrt}[c^2 - d^2]*\text{Sqrt}[(\text{Cos}[e] - I*\text{Sin}[e])^2]))*\text{Cos}[(e + f*x)/2]*(d + c*\text{Cos}[e + f*x])*(I*\text{Cos}[e] + \text{Sin}[e]))/((c + d)*\text{Sqrt}[c^2 - d^2]*\text{Sqrt}[(\text{Cos}[e] - I*\text{Sin}[e])^2]) + (d + c*\text{Cos}[e + f*x])* \text{Sec}[e/2]*\text{Sin}[(f*x)/2] + (d^2*\text{Cos}[(e + f*x)/2]*(-d*\text{Sin}[e] + c*\text{Sin}[f*x]))/(c*(c + d)*(\text{Cos}[e/2] - \text{Sin}[e/2])*(\text{Cos}[e/2] + \text{Sin}[e/2]))))/(a*(c - d)^2*f*(1 + \text{Sec}[e + f*x])*(c + d*\text{Sec}[e + f*x])^2)$

3.215.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.61, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 4475, 114, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{(a \sec(e + fx) + a)(c + d \sec(e + fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{(a \csc(e + fx + \frac{\pi}{2}) + a)(c + d \csc(e + fx + \frac{\pi}{2}))^2} dx$$

↓ 4475

$$\frac{a^2 \tan(e + fx) \int \frac{1}{\sqrt{a - a \sec(e + fx)} (\sec(e + fx) a + a)^{3/2} (c + d \sec(e + fx))^2} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 114

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^2 (c + d - d \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} (\sec(e + fx) a + a)^{3/2} (c + d \sec(e + fx))} d \sec(e + fx)}{a^2 (c^2 - d^2)} + \frac{d \sqrt{a - a \sec(e + fx)}}{a^2 (c^2 - d^2) \sqrt{a \sec(e + fx) + a} (c + d \sec(e + fx))} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 27

3.215. $\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^2} dx$

$$\begin{aligned}
& \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{c+d-d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{c^2-d^2} + \frac{d\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)\sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))} \right)}{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}} \\
& \quad \downarrow 169 \\
& \frac{a^2 \tan(e+fx) \left(-\frac{\int \frac{a^2 d(2c+d)}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{a^3(c-d)} - \frac{(c+2d)\sqrt{a-a \sec(e+fx)}}{a^2(c-d)\sqrt{a \sec(e+fx)+a}} + \frac{d\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)\sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))} \right)}{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}} \\
& \quad \downarrow 27 \\
& \frac{a^2 \tan(e+fx) \left(-\frac{d(2c+d) \int \frac{1}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{a(c-d)} - \frac{(c+2d)\sqrt{a-a \sec(e+fx)}}{a^2(c-d)\sqrt{a \sec(e+fx)+a}} + \frac{d\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)\sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))} \right)}{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}} \\
& \quad \downarrow 104 \\
& \frac{a^2 \tan(e+fx) \left(-\frac{2d(2c+d) \int \frac{1}{a(c-d)+\frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{a(c-d)} - \frac{(c+2d)\sqrt{a-a \sec(e+fx)}}{a^2(c-d)\sqrt{a \sec(e+fx)+a}} + \frac{d\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)\sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))} \right)}{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}} \\
& \quad \downarrow 218 \\
& \frac{a^2 \tan(e+fx) \left(-\frac{2d(2c+d) \arctan\left(\frac{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}}{\sqrt{c-d}\sqrt{a-a \sec(e+fx)}}\right)}{a^2(c-d)^{3/2}\sqrt{c+d}} - \frac{(c+2d)\sqrt{a-a \sec(e+fx)}}{a^2(c-d)\sqrt{a \sec(e+fx)+a}} + \frac{d\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)\sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))} \right)}{f\sqrt{a-a \sec(e+fx)}\sqrt{a \sec(e+fx)+a}}
\end{aligned}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^2), x]`

```
output -((a^2*((d*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c^2 - d^2)*Sqrt[a + a*Sec[e + f
*x]]*(c + d*Sec[e + f*x])) + ((-2*d*(2*c + d)*ArcTan[(Sqrt[c + d]*Sqrt[a +
a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]/(a^2*(c - d)^(3
/2)*Sqrt[c + d]) - ((c + 2*d)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*Sqrt[
a + a*Sec[e + f*x]])))/(c^2 - d^2)*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x
]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.215.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 114 Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_
))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)
)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e
- a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1)
- b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] ||
IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

```
rule 169 Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_
))^(p_)*((g_) + (h_)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c +
d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + S
imp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n
*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a
h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[
2*m, 2*n, 2*p]
```

```
rule 218 Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

$$3.215. \int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c+d\sec(e+fx))^2} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*sqrt[a + b*Csc[e + f*x]]*sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.215.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.01

method	result
derivativedivides	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{c^2 - 2cd + d^2} + \frac{4d \left(-\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d \right)} - \frac{(2c+d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c+d) \sqrt{(c+d)(c-d)}} \right)}{fa} \frac{(c-d)^2}{(c-d)^2}$
default	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{c^2 - 2cd + d^2} + \frac{4d \left(-\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d \right)} - \frac{(2c+d) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c+d) \sqrt{(c+d)(c-d)}} \right)}{fa} \frac{(c-d)^2}{(c-d)^2}$
risch	$\frac{2i(c^3 e^{2i(fx+e)} + c^2 d e^{2i(fx+e)} + d^3 e^{2i(fx+e)} + 2c^2 d e^{i(fx+e)} + 3c d^2 e^{i(fx+e)} + d^3 e^{i(fx+e)} + c^3 + c^2 d + c d^2)}{(e^{i(fx+e)} + 1)(e^{2i(fx+e)} c + 2d e^{i(fx+e)} + c) f (c-d)^2 a c (c+d)} + \frac{2d \ln\left(e^{i(fx+e)}\right)}{\sqrt{c^2}}$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `1/f/a*(tan(1/2*f*x+1/2*e)/(c^2-2*c*d+d^2)+4*d/(c-d)^2*(-1/2*d/(c+d)*tan(1/2*f*x+1/2*e)/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)-1/2*(2*c+d)/(c+d)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))))`

$$3.215. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$$

3.215.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(136) = 272$.

Time = 0.32 (sec) , antiderivative size = 691, normalized size of antiderivative = 4.77

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))(c+d\sec(e+fx))^2} dx$$

$$= \left[\frac{(2cd^2+d^3+(2c^2d+cd^2)\cos(fx+e))^2+(2c^2d+3cd^2+d^3)\cos(fx+e)\sqrt{c^2-d^2}\log\left(\frac{2cd\cos(fx+e)-(c^2-d^2)\sin(fx+e)}{(ac^6-ac^5d-2ac^4d^2+2ac^3d^3+ac^2d^4-acd^5)f\cos(fx+e)^2+(ac^6-3ac^4d^2+3acd^5)f\sin(fx+e)}\right)}{2((ac^6-ac^5d-2ac^4d^2+2ac^3d^3+ac^2d^4-acd^5)f\cos(fx+e))^2+(ac^6-3ac^4d^2+3acd^5)f\sin(fx+e)} \right. \\ \left. - \frac{(2cd^2+d^3+(2c^2d+cd^2)\cos(fx+e))^2+(2c^2d+3cd^2+d^3)\cos(fx+e)\sqrt{-c^2+d^2}\arctan\left(-\frac{\sqrt{-c^2+d^2}\sin(fx+e)}{(c^2-d^2)\cos(fx+e)}\right)}{(ac^6-ac^5d-2ac^4d^2+2ac^3d^3+ac^2d^4-acd^5)f\cos(fx+e)^2+(ac^6-3ac^4d^2+3acd^5)f\sin(fx+e)} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="fracas")`

output `[1/2*((2*c*d^2 + d^3 + (2*c^2*d + c*d^2)*cos(f*x + e))^2 + (2*c^2*d + 3*c*d^2 + d^3)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e))^2 - 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(c^3*d + 2*c^2*d^2 - c*d^3 - 2*d^4 + (c^4 + c^3*d - c*d^3 - d^4)*cos(f*x + e))*sin(f*x + e)/((a*c^6 - a*c^5*d - 2*a*c^4*d^2 + 2*a*c^3*d^3 + a*c^2*d^4 - a*c*d^5)*f*cos(f*x + e)^2 + (a*c^6 - 3*a*c^4*d^2 + 3*a*c^2*d^4 - a*d^6)*f*cos(f*x + e) + (a*c^5*d - a*c^4*d^2 - 2*a*c^3*d^3 + 2*a*c^2*d^4 + a*c*d^5 - a*d^6)*f), -((2*c*d^2 + d^3 + (2*c^2*d + c*d^2)*cos(f*x + e))^2 + (2*c^2*d + 3*c*d^2 + d^3)*cos(f*x + e))*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) - (c^3*d + 2*c^2*d^2 - c*d^3 - 2*d^4 + (c^4 + c^3*d - c*d^3 - d^4)*cos(f*x + e))*sin(f*x + e)/((a*c^6 - a*c^5*d - 2*a*c^4*d^2 + 2*a*c^3*d^3 + a*c^2*d^4 - a*c*d^5)*f*cos(f*x + e)^2 + (a*c^6 - 3*a*c^4*d^2 + 3*a*c^2*d^4 - a*d^6)*f*cos(f*x + e) + (a*c^5*d - a*c^4*d^2 - 2*a*c^3*d^3 + 2*a*c^2*d^4 + a*c*d^5 - a*d^6)*f)]`

3.215.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^2} dx$$

$$= \frac{\int \frac{\sec(e + fx)}{c^2 \sec(e + fx) + c^2 + 2cd \sec^2(e + fx) + 2cd \sec(e + fx) + d^2 \sec^3(e + fx) + d^2 \sec^2(e + fx)} dx}{a}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))**2,x)`

output `Integral(sec(e + f*x)/(c**2*sec(e + f*x) + c**2 + 2*c*d*sec(e + f*x)**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)**3 + d**2*sec(e + f*x)**2), x)/a`

3.215.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.215.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.52

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^2} dx =$$

$$\frac{2d^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{(ac^3 - ac^2d - acd^2 + ad^3) \left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - c - d\right)} - \frac{2 \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan\left(\frac{c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{\sqrt{-c^2+d^2}}\right)\right)}{(ac^3 - ac^2d - acd^2 + ad^3) \sqrt{-c^2+d^2}}$$

f

3.215. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output `-(2*d^2*tan(1/2*f*x + 1/2*e)/((a*c^3 - a*c^2*d - a*c*d^2 + a*d^3)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)) - 2*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*(2*c*d + d^2)/((a*c^3 - a*c^2*d - a*c*d^2 + a*d^3)*sqrt(-c^2 + d^2)) - tan(1/2*f*x + 1/2*e)/(a*c^2 - 2*a*c*d + a*d^2))/f`

3.215.9 Mupad [B] (verification not implemented)

Time = 13.54 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.29

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^2} dx = \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a f (c - d)^2} - \frac{2 d^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f (c + d) \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (a c^3 - 3 a c^2 d + 3 a c d^2 - a d^3) - a d^3 - a c^3 + a c d^2 + a c^2 d \right)} - \frac{2 d \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (2 c - 2 d) (a c^2 - 2 a c d + a d^2)}{2 a \sqrt{c + d} (c - d)^{5/2}}\right) (2 c + d)}{a f (c + d)^{3/2} (c - d)^{5/2}}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c + d/cos(e + f*x))^2),x)`

output `tan(e/2 + (f*x)/2)/(a*f*(c - d)^2) - (2*d^2*tan(e/2 + (f*x)/2))/(f*(c + d)*(tan(e/2 + (f*x)/2)^2*(a*c^3 - a*d^3 + 3*a*c*d^2 - 3*a*c^2*d) - a*d^3 - a*c^3 + a*c*d^2 + a*c^2*d)) - (2*d*atanh((tan(e/2 + (f*x)/2)*(2*c - 2*d)*(a*c^2 + a*d^2 - 2*a*c*d))/(2*a*(c + d)^(1/2)*(c - d)^(5/2)))*(2*c + d))/(a*f*(c + d)^(3/2)*(c - d)^(5/2))`

3.215. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^2} dx$

3.216 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$

3.216.1 Optimal result 1539
 3.216.2 Mathematica [C] (warning: unable to verify) 1540
 3.216.3 Rubi [A] (verified) 1540
 3.216.4 Maple [A] (verified) 1544
 3.216.5 Fricas [B] (verification not implemented) 1545
 3.216.6 Sympy [F] 1546
 3.216.7 Maxima [F(-2)] 1547
 3.216.8 Giac [A] (verification not implemented) 1547
 3.216.9 Mupad [B] (verification not implemented) 1548

3.216.1 Optimal result

Integrand size = 31, antiderivative size = 207

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$$

$$= -\frac{3d(2c^2+2cd+d^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a(c-d)^{7/2}(c+d)^{5/2}f}$$

$$+ \frac{d(2c+3d) \tan(e+fx)}{2a(c-d)^2(c+d)f(c+d \sec(e+fx))^2}$$

$$+ \frac{\tan(e+fx)}{(c-d)f(a+a \sec(e+fx))(c+d \sec(e+fx))^2}$$

$$+ \frac{d(2c+d)(c+4d) \tan(e+fx)}{2a(c-d)^3(c+d)^2f(c+d \sec(e+fx))}$$

output

```
-3*d*(2*c^2+2*c*d+d^2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))
/a/(c-d)^(7/2)/(c+d)^(5/2)/f+1/2*d*(2*c+3*d)*tan(f*x+e)/a/(c-d)^2/(c+d)/f/
(c+d*sec(f*x+e))^2+tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^2+
1/2*d*(2*c+d)*(c+4*d)*tan(f*x+e)/a/(c-d)^3/(c+d)^2/f/(c+d*sec(f*x+e))
```

3.216.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.45 (sec) , antiderivative size = 1422, normalized size of antiderivative = 6.87

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^3} dx = \text{Too large to display}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x))*(c + d*Sec[e + f*x])^3),x]`

output

```
((2*c^2 + 2*c*d + d^2)*Cos[e/2 + (f*x)/2]^2*(d + c*Cos[e + f*x])^3*Sec[e +
f*x]^4*((-6*I)*d*ArcTan[Sec[(f*x)/2]*(Cos[e]/(Sqrt[c^2 - d^2])*Sqrt[Cos[2*
e] - I*Sin[2*e]]) - (I*Sin[e])/(Sqrt[c^2 - d^2])*Sqrt[Cos[2*e] - I*Sin[2*e
]]))*((-I)*d*Sin[(f*x)/2] + I*c*Sin[e + (f*x)/2])*Cos[e]/(Sqrt[c^2 - d^2
]*f*Sqrt[Cos[2*e] - I*Sin[2*e]]) - (6*d*ArcTan[Sec[(f*x)/2]*(Cos[e]/(Sqrt[
c^2 - d^2])*Sqrt[Cos[2*e] - I*Sin[2*e]]) - (I*Sin[e])/(Sqrt[c^2 - d^2])*Sqrt
[Cos[2*e] - I*Sin[2*e]])*((-I)*d*Sin[(f*x)/2] + I*c*Sin[e + (f*x)/2])*Si
n[e]/(Sqrt[c^2 - d^2]*f*Sqrt[Cos[2*e] - I*Sin[2*e]])))/((-c + d)^3*(c + d
)^2*(a + a*Sec[e + f*x))*(c + d*Sec[e + f*x])^3) + (Cos[e/2 + (f*x)/2]*(d
+ c*Cos[e + f*x])*Sec[e/2]*Sec[e]*Sec[e + f*x]^4*(8*c^5*d*Sin[(f*x)/2] + 1
0*c^4*d^2*Sin[(f*x)/2] - 11*c^3*d^3*Sin[(f*x)/2] - 17*c^2*d^4*Sin[(f*x)/2]
- 2*c*d^5*Sin[(f*x)/2] + 2*d^6*Sin[(f*x)/2] - 8*c^5*d*Sin[(3*f*x)/2] - 22
*c^4*d^2*Sin[(3*f*x)/2] - 27*c^3*d^3*Sin[(3*f*x)/2] - 5*c^2*d^4*Sin[(3*f*x
)/2] + 2*c*d^5*Sin[(3*f*x)/2] + 4*c^6*Sin[e - (f*x)/2] + 8*c^5*d*Sin[e - (
f*x)/2] + 18*c^4*d^2*Sin[e - (f*x)/2] + 35*c^3*d^3*Sin[e - (f*x)/2] + 25*c
^2*d^4*Sin[e - (f*x)/2] + 2*c*d^5*Sin[e - (f*x)/2] - 2*d^6*Sin[e - (f*x)/2
] - 4*c^6*Sin[e + (f*x)/2] - 8*c^5*d*Sin[e + (f*x)/2] - 6*c^4*d^2*Sin[e +
(f*x)/2] - 7*c^3*d^3*Sin[e + (f*x)/2] + 5*c^2*d^4*Sin[e + (f*x)/2] + 2*c*d
^5*Sin[e + (f*x)/2] - 2*d^6*Sin[e + (f*x)/2] + 8*c^5*d*Sin[2*e + (f*x)/2]
+ 22*c^4*d^2*Sin[2*e + (f*x)/2] + 17*c^3*d^3*Sin[2*e + (f*x)/2] + 13*c^...
```

3.216.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.58, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4475, 114, 27, 168, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.216. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)(c+d \sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)(c+d \csc(e+fx+\frac{\pi}{2}))^3} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{1}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))^3} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 114

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(2c+d-2d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2a^2(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}}{2a^2(c^2-d^2) \sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))^2} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{2c+d-2d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}}{2a^2(c^2-d^2) \sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))^2} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 168

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2((c+d)(2c+3d)-d(4c+d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{a^2(c^2-d^2)} + \frac{d(4c+d) \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2) \sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))} + \frac{1}{2a^2(c^2-d^2) \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{(c+d)(2c+3d)-d(4c+d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{c^2-d^2} + \frac{d(4c+d) \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2) \sqrt{a \sec(e+fx)+a}(c+d \sec(e+fx))} + \frac{1}{2a^2(c^2-d^2) \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 169

3.216. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{3a^2 d(2c^2 + 2cd + d^2)}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a(c + d \sec(e + fx))}} d \sec(e + fx)}{a^3(c - d)} - \frac{(2c + d)(c + 4d)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} + \frac{d(4c + d)\sqrt{a - a \sec(e + fx)}}{a^2(c^2 - d^2)\sqrt{a \sec(e + fx) + a(c + d \sec(e + fx))}} \right) \frac{c^2 - d^2}{2(c^2 - d^2)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

27

$$a^2 \tan(e + fx) \left(\frac{3d(2c^2 + 2cd + d^2) \int \frac{1}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a(c + d \sec(e + fx))}} d \sec(e + fx)}{a(c - d)} - \frac{(2c + d)(c + 4d)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} + \frac{d(4c + d)\sqrt{a - a \sec(e + fx)}}{a^2(c^2 - d^2)\sqrt{a \sec(e + fx) + a(c + d \sec(e + fx))}} \right) \frac{c^2 - d^2}{2(c^2 - d^2)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

104

$$a^2 \tan(e + fx) \left(\frac{6d(2c^2 + 2cd + d^2) \int \frac{1}{a(c - d) + \frac{a(c + d)(\sec(e + fx)a + a)}{a - a \sec(e + fx)}} d \frac{\sqrt{\sec(e + fx)a + a}}{\sqrt{a - a \sec(e + fx)}}}{a(c - d)} - \frac{(2c + d)(c + 4d)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} + \frac{d(4c + d)\sqrt{a - a \sec(e + fx)}}{a^2(c^2 - d^2)\sqrt{a \sec(e + fx) + a(c + d \sec(e + fx))}} \right) \frac{c^2 - d^2}{2(c^2 - d^2)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

218

$$a^2 \tan(e + fx) \left(\frac{6d(2c^2 + 2cd + d^2) \arctan\left(\frac{\sqrt{c + d}\sqrt{a \sec(e + fx) + a}}{\sqrt{c - d}\sqrt{a - a \sec(e + fx)}}\right)}{a^2(c - d)^{3/2}\sqrt{c + d}} - \frac{(2c + d)(c + 4d)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} + \frac{d(4c + d)\sqrt{a - a \sec(e + fx)}}{a^2(c^2 - d^2)\sqrt{a \sec(e + fx) + a(c + d \sec(e + fx))}} \right) \frac{c^2 - d^2}{2(c^2 - d^2)}$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

```
input Int[Sec[e + f*x]/((a + a*Sec[e + f*x])*(c + d*Sec[e + f*x])^3),x]
```

```
output -((a^2*((d*Sqrt[a - a*Sec[e + f*x]])/(2*a^2*(c^2 - d^2)*Sqrt[a + a*Sec[e +
f*x]]*(c + d*Sec[e + f*x])^2) + ((d*(4*c + d)*Sqrt[a - a*Sec[e + f*x]])/(
a^2*(c^2 - d^2)*Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])) + ((-6*d*(2
*c^2 + 2*c*d + d^2)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c
- d]*Sqrt[a - a*Sec[e + f*x]])])/(a^2*(c - d)^(3/2)*Sqrt[c + d]) - ((2*c +
d)*(c + 4*d)*Sqrt[a - a*Sec[e + f*x]]/(a^2*(c - d)*Sqrt[a + a*Sec[e + f*
x]])))/(c^2 - d^2)/(2*(c^2 - d^2))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*
x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.216.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 114 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)
)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e
- a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1)
- b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] ||
IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

```
rule 168 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c +
d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + S
imp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n
*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*
h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```


rule 169 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_))*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_))*((csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.)^(n_)), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*(c + d*x)^n/Sqrt[a - b*x], x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.216.4 Maple [A] (verified)

Time = 0.99 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.07

3.216.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$$

method	result
derivativedivides	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{c^3 - 3c^2d + 3cd^2 - d^3} + \frac{2d \left(\frac{-3d(2c^2 - cd - d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + \frac{d(6c+d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c^2 + 2cd + d^2)} \right)}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{3(2c^2 + 2cd + d^2) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c^2 + 2cd + d^2) \sqrt{(c+d)(c-d)}}}{(c-d)^3}$
default	$\frac{\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{c^3 - 3c^2d + 3cd^2 - d^3} + \frac{2d \left(\frac{-3d(2c^2 - cd - d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + \frac{d(6c+d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2(c^2 + 2cd + d^2)} \right)}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{3(2c^2 + 2cd + d^2) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{2(c^2 + 2cd + d^2) \sqrt{(c+d)(c-d)}}}{fa}$
risch	$i(2d^6 e^{2i(fx+e)} + c^2 d^4 - 4c^5 d - 2c^3 d^3 - 8c^4 d^2 - 2c^6 - 2c d^5 e^{2i(fx+e)} - 7c^3 d^3 e^{4i(fx+e)} - 2c^2 d^4 e^{4i(fx+e)} + 2c d^5 e^{4i(fx+e)} - 8c^5)$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/f/a*(tan(1/2*f*x+1/2*e)/(c^3-3*c^2*d+3*c*d^2-d^3)+2*d/(c-d)^3*((-3/2*d*(2*c^2-c*d-d^2)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3+1/2*d*(6*c+d)/(c+d)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^2-3/2*(2*c^2+2*c*d+d^2)/(c^2+2*c*d+d^2)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))))
```

3.216.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 637 vs. 2(194) = 388.

Time = 0.36 (sec) , antiderivative size = 1331, normalized size of antiderivative = 6.43

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^3} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="fricas")
```

output

```

[-1/4*(3*(2*c^2*d^3 + 2*c*d^4 + d^5 + (2*c^4*d + 2*c^3*d^2 + c^2*d^3)*cos(
f*x + e)^3 + (2*c^4*d + 6*c^3*d^2 + 5*c^2*d^3 + 2*c*d^4)*cos(f*x + e)^2 +
(4*c^3*d^2 + 6*c^2*d^3 + 4*c*d^4 + d^5)*cos(f*x + e))*sqrt(c^2 - d^2)*log(
(2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*
cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*
cos(f*x + e) + d^2)) - 2*(2*c^4*d^2 + 9*c^3*d^3 + 2*c^2*d^4 - 9*c*d^5 - 4*
d^6 + (2*c^6 + 4*c^5*d + 6*c^4*d^2 - 2*c^3*d^3 - 9*c^2*d^4 - 2*c*d^5 + d^6
)*cos(f*x + e)^2 + (4*c^5*d + 14*c^4*d^2 + 7*c^3*d^3 - 13*c^2*d^4 - 11*c*d
^5 - d^6)*cos(f*x + e))*sin(f*x + e))/((a*c^9 - a*c^8*d - 3*a*c^7*d^2 + 3*
a*c^6*d^3 + 3*a*c^5*d^4 - 3*a*c^4*d^5 - a*c^3*d^6 + a*c^2*d^7)*f*cos(f*x +
e)^3 + (a*c^9 + a*c^8*d - 5*a*c^7*d^2 - 3*a*c^6*d^3 + 9*a*c^5*d^4 + 3*a*c
^4*d^5 - 7*a*c^3*d^6 - a*c^2*d^7 + 2*a*c*d^8)*f*cos(f*x + e)^2 + (2*a*c^8*
d - a*c^7*d^2 - 7*a*c^6*d^3 + 3*a*c^5*d^4 + 9*a*c^4*d^5 - 3*a*c^3*d^6 - 5*
a*c^2*d^7 + a*c*d^8 + a*d^9)*f*cos(f*x + e) + (a*c^7*d^2 - a*c^6*d^3 - 3*a
*c^5*d^4 + 3*a*c^4*d^5 + 3*a*c^3*d^6 - 3*a*c^2*d^7 - a*c*d^8 + a*d^9)*f),
-1/2*(3*(2*c^2*d^3 + 2*c*d^4 + d^5 + (2*c^4*d + 2*c^3*d^2 + c^2*d^3)*cos(f
*x + e)^3 + (2*c^4*d + 6*c^3*d^2 + 5*c^2*d^3 + 2*c*d^4)*cos(f*x + e)^2 + (
4*c^3*d^2 + 6*c^2*d^3 + 4*c*d^4 + d^5)*cos(f*x + e))*sqrt(-c^2 + d^2)*arct
an(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) - (2
*c^4*d^2 + 9*c^3*d^3 + 2*c^2*d^4 - 9*c*d^5 - 4*d^6 + (2*c^6 + 4*c^5*d + ...

```

3.216.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^3} dx$$

$$= \frac{\int \frac{\sec(e+fx)}{c^3 \sec(e+fx) + c^3 + 3c^2 d \sec^2(e+fx) + 3c^2 d \sec(e+fx) + 3cd^2 \sec^3(e+fx) + 3cd^2 \sec^2(e+fx) + d^3 \sec^4(e+fx) + d^3 \sec^3(e+fx)}{a} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))**3,x)`

output `Integral(sec(e + f*x)/(c**3*sec(e + f*x) + c**3 + 3*c**2*d*sec(e + f*x)**2 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**3 + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**4 + d**3*sec(e + f*x)**3), x)/a`

3.216. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$

3.216.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see 'assume?' f or more de

3.216.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.75

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^3} dx = \frac{3(2c^2d + 2cd^2 + d^3) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2c+2d) + \arctan \left(-\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right)}{(ac^5 - ac^4d - 2ac^3d^2 + 2ac^2d^3 + acd^4 - ad^5)\sqrt{-c^2+d^2}} - \frac{\tan(\frac{1}{2}fx + \frac{1}{2}e)}{ac^3 - 3ac^2d + 3acd^2 - ad^3} + \frac{6c^2d^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output `-(3*(2*c^2*d + 2*c*d^2 + d^3)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((a*c^5 - a*c^4*d - 2*a*c^3*d^2 + 2*a*c^2*d^3 + a*c*d^4 - a*d^5)*sqrt(-c^2 + d^2)) - tan(1/2*f*x + 1/2*e)/(a*c^3 - 3*a*c^2*d + 3*a*c*d^2 - a*d^3) + (6*c^2*d^2*tan(1/2*f*x + 1/2*e)^3 - 3*c*d^3*tan(1/2*f*x + 1/2*e)^3 - 3*d^4*tan(1/2*f*x + 1/2*e)^3 - 6*c^2*d^2*tan(1/2*f*x + 1/2*e) - 7*c*d^3*tan(1/2*f*x + 1/2*e) - d^4*tan(1/2*f*x + 1/2*e))/((a*c^5 - a*c^4*d - 2*a*c^3*d^2 + 2*a*c^2*d^3 + a*c*d^4 - a*d^5)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^2)/f`

3.216. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))(c+d \sec(e+fx))^3} dx$

3.216.9 Mupad [B] (verification not implemented)

Time = 14.58 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.83

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))(c + d \sec(e + fx))^3} dx = \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a f (c - d)^3}$$

$$- \frac{\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)(d^3 + 6cd^2)}{c+d} + \frac{3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{c+d}}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (2ac^5 - 6ac^4d + 4ac^3d^2 + 4ac^2d^3 - 6acd^4 + 2ad^5) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 (ac^5 - 5acd^4 + d^5) \right)}$$

$$+ \frac{d \operatorname{atan}\left(\frac{\operatorname{li} \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^4 - 4i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^3 d + 6i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^2 d^2 - 4i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c d^3 + \operatorname{li} \tan\left(\frac{e}{2} + \frac{fx}{2}\right) d^4}{\sqrt{c+d}(c-d)^{7/2}}\right)}{a f (c + d)^{5/2} (c - d)^{7/2}} (2c^2 + 2cd + d^2)$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))*(c + d/cos(e + f*x))^3),x)`

output

```
tan(e/2 + (f*x)/2)/(a*f*(c - d)^3) - ((tan(e/2 + (f*x)/2)*(6*c*d^2 + d^3))
/(c + d) + (3*tan(e/2 + (f*x)/2)^3*(c*d^3 + d^4 - 2*c^2*d^2))/(c + d)^2)/(
f*(tan(e/2 + (f*x)/2)^2*(2*a*c^5 + 2*a*d^5 + 4*a*c^2*d^3 + 4*a*c^3*d^2 - 6
*a*c*d^4 - 6*a*c^4*d) - tan(e/2 + (f*x)/2)^4*(a*c^5 - a*d^5 - 10*a*c^2*d^3
+ 10*a*c^3*d^2 + 5*a*c*d^4 - 5*a*c^4*d) - a*c^5 + a*d^5 - 2*a*c^2*d^3 + 2
*a*c^3*d^2 - a*c*d^4 + a*c^4*d)) + (d*atan((c^4*tan(e/2 + (f*x)/2)*1i + d^
4*tan(e/2 + (f*x)/2)*1i - c*d^3*tan(e/2 + (f*x)/2)*4i - c^3*d*tan(e/2 + (f
*x)/2)*4i + c^2*d^2*tan(e/2 + (f*x)/2)*6i)/((c + d)^(1/2)*(c - d)^(7/2)))*
(2*c*d + 2*c^2 + d^2)*3i)/(a*f*(c + d)^(5/2)*(c - d)^(7/2))
```

3.217 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^2} dx$

3.217.1 Optimal result 1549
 3.217.2 Mathematica [A] (verified) 1550
 3.217.3 Rubi [A] (verified) 1550
 3.217.4 Maple [A] (verified) 1555
 3.217.5 Fracas [A] (verification not implemented) 1556
 3.217.6 Sympy [F] 1557
 3.217.7 Maxima [B] (verification not implemented) 1557
 3.217.8 Giac [B] (verification not implemented) 1558
 3.217.9 Mupad [B] (verification not implemented) 1559

3.217.1 Optimal result

Integrand size = 31, antiderivative size = 258

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^2} dx = \frac{5(2c-d)d^2(2c^2-3cd+2d^2) \operatorname{arctanh}(\sin(e+fx))}{2a^2 f} - \frac{d(c^2+10cd-12d^2)(c+d \sec(e+fx))^2 \tan(e+fx)}{3a^2 f} + \frac{(c-d)(c+10d)(c+d \sec(e+fx))^3 \tan(e+fx)}{3f(a^2+a^2 \sec(e+fx))} + \frac{(c-d)(c+d \sec(e+fx))^4 \tan(e+fx)}{3f(a+a \sec(e+fx))^2} - \frac{d(4(c^4+10c^3d-44c^2d^2+40cd^3-12d^4)+d(2c^3+20c^2d-57cd^2+30d^3) \sec(e+fx)) \tan(e+fx)}{6a^2 f}$$

output

```
5/2*(2*c-d)*d^2*(2*c^2-3*c*d+2*d^2)*arctanh(sin(f*x+e))/a^2/f-1/3*d*(c^2+10*c*d-12*d^2)*(c+d*sec(f*x+e))^2*tan(f*x+e)/a^2/f+1/3*(c-d)*(c+10*d)*(c+d*sec(f*x+e))^3*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))+1/3*(c-d)*(c+d*sec(f*x+e))^4*tan(f*x+e)/f/(a+a*sec(f*x+e))^2-1/6*d*(4*c^4+40*c^3*d-176*c^2*d^2+160*c*d^3-48*d^4+d*(2*c^3+20*c^2*d-57*c*d^2+30*d^3)*sec(f*x+e))*tan(f*x+e)/a^2/f
```

3.217.2 Mathematica [A] (verified)

Time = 7.42 (sec) , antiderivative size = 446, normalized size of antiderivative = 1.73

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{240d^2(-4c^3+8c^2d-7cd^2+2d^3)\cos^4\left(\frac{1}{2}(e+fx)\right)\left(\log\left(\cos\left(\frac{1}{2}(e+fx)\right)-\sin\left(\frac{1}{2}(e+fx)\right)\right)-\log\left(\cos\left(\frac{1}{2}(e+fx)\right)+\sin\left(\frac{1}{2}(e+fx)\right)\right)\right)}{(a+a\sec(e+fx))^2}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^2,x]`

output

$$\frac{(240d^2(-4c^3+8c^2d-7cd^2+2d^3)\cos^4\left(\frac{e+fx}{2}\right)\left(\log\left[\frac{\cos\left(\frac{e+fx}{2}\right)-\sin\left(\frac{e+fx}{2}\right)}{\cos\left(\frac{e+fx}{2}\right)+\sin\left(\frac{e+fx}{2}\right)}\right)+2\cos\left(\frac{e+fx}{2}\right)\left(6c^5+15c^4d-120c^3d^2+420c^2d^3-300cd^4+104d^5+(6c^5+60c^4d-300c^3d^2+840c^2d^3-585cd^4+190d^5)\cos[e+fx]+4(2c^5+5c^4d-40c^3d^2+130c^2d^3-95cd^4+30d^5)\cos[2(e+fx)]+2c^5\cos[3(e+fx)]+20c^4d\cos[3(e+fx)]-100c^3d^2\cos[3(e+fx)]+280c^2d^3\cos[3(e+fx)]-215cd^4\cos[3(e+fx)]+66d^5\cos[3(e+fx)]+2c^5\cos[4(e+fx)]+5c^4d\cos[4(e+fx)]-40c^3d^2\cos[4(e+fx)]+100c^2d^3\cos[4(e+fx)]-80cd^4\cos[4(e+fx)]+24d^5\cos[4(e+fx)]\right)\sec[e+fx]^3\sin\left(\frac{e+fx}{2}\right)}{(24a^2f(1+\cos[e+fx])^2)}$$
3.217.3 Rubi [A] (verified)Time = 0.59 (sec) , antiderivative size = 392, normalized size of antiderivative = 1.52, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4475, 109, 25, 27, 167, 27, 170, 25, 27, 164, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a\sec(e+fx)+a)^2} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)^5}{\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^2} dx$$

↓ 4475

3.217. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$

$$\frac{a^2 \tan(e+fx) \int \frac{(c+d \sec(e+fx))^5}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 109

$$\frac{a^2 \tan(e+fx) \left(-\frac{\int \frac{a^2(c+d \sec(e+fx))^3(c^2+6dc-4d^2-3(c-2d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))^3(c^2+6dc-4d^2-3(c-2d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{(c+d \sec(e+fx))^3(c^2+6dc-4d^2-3(c-2d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 167

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{3a^2 d(c+d \sec(e+fx))^2((11c-10d)d-(c^2+10dc-12d^2)\sec(e+fx))}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{3a} - \frac{(c-d)(c+10d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{3d \int \frac{(c+d \sec(e+fx))^2((11c-10d)d-(c^2+10dc-12d^2)\sec(e+fx))}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{a} - \frac{(c-d)(c+10d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 170

3.217. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^2} dx$

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{(c^2 + 10cd - 12d^2) \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (c + d \sec(e + fx))^2}{3a^2} \int - \frac{a^2 (c + d \sec(e + fx)) (d(31c^2 - 50dc + 24d^2) - (2c^3 + 20dc^2 - 57d^2c + 30d^3) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx) a + a}} \right)}{a} \right) \frac{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{3a}$$

25

$$a^2 \tan(e + fx) \left(\frac{3d \left(\int \frac{a^2 (c + d \sec(e + fx)) (d(31c^2 - 50dc + 24d^2) - (2c^3 + 20dc^2 - 57d^2c + 30d^3) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx) a + a}} d \sec(e + fx) + \frac{(c^2 + 10cd - 12d^2) \sqrt{a - a \sec(e + fx)}}{3a^2} \right)}{a} \right) \frac{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{3a}$$

27

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \int \frac{(c + d \sec(e + fx)) (d(31c^2 - 50dc + 24d^2) - (2c^3 + 20dc^2 - 57d^2c + 30d^3) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx) a + a}} d \sec(e + fx) + \frac{(c^2 + 10cd - 12d^2) \sqrt{a - a \sec(e + fx)}}{3a^2} \right)}{a} \right) \frac{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{3a}$$

164

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \left(\frac{15}{2} d(2c - d)(2c^2 - 3cd + 2d^2) \int \frac{1}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx) a + a}} d \sec(e + fx) + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(2c^3 + 20c^2d - 57cd^2 + 30d^3) \sec(e + fx))}{\sqrt{\sec(e + fx) a + a}} \right) \right)}{a} \right) \frac{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{3a}$$

45

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \left(15d(2c - d)(2c^2 - 3cd + 2d^2) \int \frac{1}{\frac{(a - a \sec(e + fx)) a}{\sec(e + fx) a + a} - a} d \frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{\sec(e + fx) a + a}} + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(2c^3 + 20c^2d - 57cd^2 + 30d^3) \sec(e + fx))}{\sqrt{\sec(e + fx) a + a}} \right) \right)}{a} \right) \frac{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{3a}$$

3.217. $\int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + a \sec(e + fx))^2} dx$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \left(\frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(2c^3 + 20c^2d - 57cd^2 + 30d^3) \sec(e + fx) + 4(c^4 + 10c^3d - 44c^2d^2 + 40cd^3 - 12d^4))}{2a^2} \right) - \frac{15d(2c - d)}{a} \right)}{\dots} \right)$$

```
input Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^2,x]
```

```
output -((a^2*(-1/3*((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^4)/(a^2*(a + a*Sec[e + f*x])^(3/2)) + (-(((c - d)*(c + 10*d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^3)/(a^2*Sqrt[a + a*Sec[e + f*x]])) + (3*d*((c^2 + 10*c*d - 12*d^2)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(3*a^2) + ((-15*(2*c - d)*d*(2*c^2 - 3*c*d + 2*d^2)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])/a + (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(4*(c^4 + 10*c^3*d - 44*c^2*d^2 + 40*c*d^3 - 12*d^4) + d*(2*c^3 + 20*c^2*d - 57*c*d^2 + 30*d^3)*Sec[e + f*x]))/(2*a^2))/3)/a)/(3*a))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.217.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 45 Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

rule 109 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

rule 164 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`

rule 167 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] - Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[b*c*(f*g - e*h)*(m + 1) + (b*g - a*h)*(d*e*n + c*f*(p + 1)) + d*(b*(f*g - e*h)*(m + 1) + f*(b*g - a*h)*(n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, p}, x] && LtQ[m, -1] && GtQ[n, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 170 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.217.4 Maple [A] (verified)

Time = 1.27 (sec) , antiderivative size = 382, normalized size of antiderivative = 1.48

method	result
parallelrisch	$-360\left(c-\frac{d}{2}\right)\left(c^2-\frac{3}{2}cd+d^2\right)\left(\cos(fx+e)+\frac{\cos(3fx+3e)}{3}\right)d^2\ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)+360\left(c-\frac{d}{2}\right)\left(c^2-\frac{3}{2}cd+d^2\right)\left(\cos(fx+e)+\frac{\cos(3fx+3e)}{3}\right)d^2$
derivativedivides	$-\frac{c^5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{5c^4 d \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} - \frac{10c^3 d^2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{10c^2 d^3 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} - \frac{5c d^4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{d^5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3}$
default	$-\frac{c^5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{5c^4 d \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} - \frac{10c^3 d^2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{10c^2 d^3 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} - \frac{5c d^4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{d^5 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3}$
norman	$-\frac{(c^5-5c^4d+10c^3d^2-10c^2d^3+5cd^4-d^5)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^{13}}{6af} - \frac{(c^5+5c^4d-30c^3d^2+90c^2d^3-65cd^4+21d^5)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{2af} + \frac{10(2c^5+5c^4d-10c^3d^2+10c^2d^3-5cd^4+d^5)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{af}$
risch	Expression too large to display

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

3.217.
$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$$

output $1/12*(-360*(c-1/2*d)*(c^2-3/2*c*d+d^2)*(\cos(f*x+e)+1/3*\cos(3*f*x+3*e))*d^2*\ln(\tan(1/2*f*x+1/2*e)-1)+360*(c-1/2*d)*(c^2-3/2*c*d+d^2)*(\cos(f*x+e)+1/3*\cos(3*f*x+3*e))*d^2*\ln(\tan(1/2*f*x+1/2*e)+1)+2*\sec(1/2*f*x+1/2*e)^2*((c^5+33*d^5+10*c^4*d-50*c^3*d^2+140*c^2*d^3-215/2*c*d^4)*\cos(3*f*x+3*e)+(4*c^5+10*c^4*d-80*c^3*d^2+260*c^2*d^3-190*c*d^4+60*d^5)*\cos(2*f*x+2*e)+(c^5+12*d^5+5/2*c^4*d-20*c^3*d^2+50*c^2*d^3-40*c*d^4)*\cos(4*f*x+4*e)+(95*d^5-150*c^3*d^2+3*c^5+30*c^4*d+420*c^2*d^3-585/2*c*d^4)*\cos(f*x+e)+52*d^5+3*c^5+15/2*c^4*d-60*c^3*d^2+210*c^2*d^3-150*c*d^4)*\tan(1/2*f*x+1/2*e))/f/a^2/(\cos(3*f*x+3*e)+3*\cos(f*x+e))$

3.217.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 456, normalized size of antiderivative = 1.77

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{15((4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5)\cos(fx+e)^5 + 2(4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5)\cos(fx+e)^4 + (4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5)\cos(fx+e)^3 \log(\sin(fx+e)+1) - 15((4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5)\cos(fx+e)^5 + 2(4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5)\cos(fx+e)^4 + (4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5)\cos(fx+e)^3 \log(-\sin(fx+e)+1) + 2*(2d^5 + 2*(2c^5 + 5c^4d - 40c^3d^2 + 100c^2d^3 - 80c*d^4 + 24d^5)*\cos(f*x+e)^4 + (2c^5 + 20c^4d - 100c^3d^2 + 280c^2d^3 - 215c*d^4 + 66d^5)*\cos(f*x+e)^3 + 6*(10c^2d^3 - 5c*d^4 + 2d^5)*\cos(f*x+e)^2 + (15c*d^4 - 2d^5)*\cos(f*x+e))*\sin(f*x+e))/(a^2*f*\cos(f*x+e)^5 + 2*a^2*f*\cos(f*x+e)^4 + a^2*f*\cos(f*x+e)^3)}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x, algorithm="fracas")`

output $1/12*(15*((4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\cos(f*x + e)^5 + 2*(4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\cos(f*x + e)^4 + (4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\cos(f*x + e)^3)*\log(\sin(f*x + e) + 1) - 15*((4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\cos(f*x + e)^5 + 2*(4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\cos(f*x + e)^4 + (4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\cos(f*x + e)^3)*\log(-\sin(f*x + e) + 1) + 2*(2*d^5 + 2*(2*c^5 + 5*c^4*d - 40*c^3*d^2 + 100*c^2*d^3 - 80*c*d^4 + 24*d^5)*\cos(f*x + e)^4 + (2*c^5 + 20*c^4*d - 100*c^3*d^2 + 280*c^2*d^3 - 215*c*d^4 + 66*d^5)*\cos(f*x + e)^3 + 6*(10*c^2*d^3 - 5*c*d^4 + 2*d^5)*\cos(f*x + e)^2 + (15*c*d^4 - 2*d^5)*\cos(f*x + e))*\sin(f*x + e))/(a^2*f*\cos(f*x + e)^5 + 2*a^2*f*\cos(f*x + e)^4 + a^2*f*\cos(f*x + e)^3)$

3.217.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{\int \frac{c^5 \sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{d^5 \sec^6(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{5cd^4 \sec^5(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{10c^2 d^3 \sec^4(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx}{a^2}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**5/(a+a*sec(f*x+e))**2,x)`

output `(Integral(c**5*sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(d**5*sec(e + f*x)**6/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(5*c*d**4*sec(e + f*x)**5/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(10*c**2*d**3*sec(e + f*x)**4/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(10*c**3*d**2*sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(5*c**4*d*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.217.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 772 vs. $2(247) = 494$.

Time = 0.24 (sec) , antiderivative size = 772, normalized size of antiderivative = 2.99

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

```
output 1/6*(d^5*(4*(9*sin(f*x + e)/(cos(f*x + e) + 1) - 20*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 15*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^2 - 3*a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 3*a^2*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - a^2*sin(f*x + e)^6/(cos(f*x + e) + 1)^6) + (27*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 30*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 30*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2) - 5*c*d^4*(6*(3*sin(f*x + e)/(cos(f*x + e) + 1) - 5*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^2 - 2*a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^2*sin(f*x + e)^4/(cos(f*x + e) + 1)^4) + (21*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 21*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 21*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2) + 10*c^2*d^3*((15*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 12*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 12*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 + 12*sin(f*x + e)/((a^2 - a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1))) - 10*c^3*d^2*((9*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 6*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 6*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2) + 5*c^4*d*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 + c^5*(3*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f
```

3.217.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 506 vs. $2(247) = 494$.

Time = 0.44 (sec) , antiderivative size = 506, normalized size of antiderivative = 1.96

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + a \sec(e + fx))^2} dx$$

$$= \frac{15(4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a^2} - \frac{15(4c^3d^2 - 8c^2d^3 + 7cd^4 - 2d^5) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a^2} - \frac{2(60c^2d^3 \tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a^2}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^2,x, algorithm="giac")
```

$$3.217. \int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^2} dx$$

output $\frac{1}{6}*(15*(4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a^2 - 15*(4*c^3*d^2 - 8*c^2*d^3 + 7*c*d^4 - 2*d^5)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a^2 - 2*(60*c^2*d^3*\tan(1/2*f*x + 1/2*e)^5 - 75*c*d^4*\tan(1/2*f*x + 1/2*e)^5 + 30*d^5*\tan(1/2*f*x + 1/2*e)^5 - 120*c^2*d^3*\tan(1/2*f*x + 1/2*e)^3 + 120*c*d^4*\tan(1/2*f*x + 1/2*e)^3 - 40*d^5*\tan(1/2*f*x + 1/2*e)^3 + 60*c^2*d^3*\tan(1/2*f*x + 1/2*e) - 45*c*d^4*\tan(1/2*f*x + 1/2*e) + 18*d^5*\tan(1/2*f*x + 1/2*e))/((\tan(1/2*f*x + 1/2*e)^2 - 1)^3*a^2) - (a^4*c^5*\tan(1/2*f*x + 1/2*e)^3 - 5*a^4*c^4*d*\tan(1/2*f*x + 1/2*e)^3 + 10*a^4*c^3*d^2*\tan(1/2*f*x + 1/2*e)^3 - 10*a^4*c^2*d^3*\tan(1/2*f*x + 1/2*e)^3 + 5*a^4*c*d^4*\tan(1/2*f*x + 1/2*e)^3 - a^4*d^5*\tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^5*\tan(1/2*f*x + 1/2*e) - 15*a^4*c^4*d*\tan(1/2*f*x + 1/2*e) + 90*a^4*c^3*d^2*\tan(1/2*f*x + 1/2*e) - 150*a^4*c^2*d^3*\tan(1/2*f*x + 1/2*e) + 105*a^4*c*d^4*\tan(1/2*f*x + 1/2*e) - 27*a^4*d^5*\tan(1/2*f*x + 1/2*e))/a^6)/f$

3.217.9 Mupad [B] (verification not implemented)

Time = 13.55 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.04

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + a \sec(e + fx))^2} dx$$

$$= \frac{5d^2 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (2c - d) (2c^2 - 3cd + 2d^2)}{a^2 f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{2(c-d)^5}{a^2} - \frac{5(c+d)(c-d)^4}{2a^2}\right)}{f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (c-d)^5}{6a^2 f}$$

$$- \frac{(20c^2d^3 - 25cd^4 + 10d^5) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-40c^2d^3 + 40cd^4 - \frac{40d^5}{3}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + (20c^2d^3 - 15cd^4 + 5d^5) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - a^2\right)}$$

input `int((c + d/cos(e + f*x))^5/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output $(5*d^2*\operatorname{atanh}(\tan(e/2 + (f*x)/2))*(2*c - d)*(2*c^2 - 3*c*d + 2*d^2))/(a^2*f) - (\tan(e/2 + (f*x)/2)*((2*(c - d)^5)/a^2 - (5*(c + d)*(c - d)^4)/(2*a^2)))/f - (\tan(e/2 + (f*x)/2)^3*(c - d)^5)/(6*a^2*f) - (\tan(e/2 + (f*x)/2)*(6*d^5 - 15*c*d^4 + 20*c^2*d^3) + \tan(e/2 + (f*x)/2)^5*(10*d^5 - 25*c*d^4 + 20*c^2*d^3) - \tan(e/2 + (f*x)/2)^3*((40*d^5)/3 - 40*c*d^4 + 40*c^2*d^3))/(f*(3*a^2*\tan(e/2 + (f*x)/2)^2 - 3*a^2*\tan(e/2 + (f*x)/2)^4 + a^2*\tan(e/2 + (f*x)/2)^6 - a^2))$

3.217. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^2} dx$

3.218
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^2} dx$$

3.218.1 Optimal result 1560
 3.218.2 Mathematica [A] (verified) 1561
 3.218.3 Rubi [A] (verified) 1561
 3.218.4 Maple [A] (verified) 1565
 3.218.5 Fricas [A] (verification not implemented) 1566
 3.218.6 Sympy [F] 1566
 3.218.7 Maxima [B] (verification not implemented) 1567
 3.218.8 Giac [A] (verification not implemented) 1568
 3.218.9 Mupad [B] (verification not implemented) 1568

3.218.1 Optimal result

Integrand size = 31, antiderivative size = 193

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^2} dx$$

$$= \frac{d^2(12c^2 - 16cd + 7d^2) \operatorname{arctanh}(\sin(e+fx))}{2a^2 f}$$

$$+ \frac{(c-d)(c+8d)(c+d \sec(e+fx))^2 \tan(e+fx)}{3f(a^2 + a^2 \sec(e+fx))}$$

$$+ \frac{(c-d)(c+d \sec(e+fx))^3 \tan(e+fx)}{3f(a+a \sec(e+fx))^2}$$

$$- \frac{d(4(c^3 + 8c^2d - 20cd^2 + 8d^3) + d(2c^2 + 16cd - 21d^2) \sec(e+fx)) \tan(e+fx)}{6a^2 f}$$

```
output 1/2*d^2*(12*c^2-16*c*d+7*d^2)*arctanh(sin(f*x+e))/a^2/f+1/3*(c-d)*(c+8*d)*
(c+d*sec(f*x+e))^2*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))+1/3*(c-d)*(c+d*sec(f*
x+e))^3*tan(f*x+e)/f/(a+a*sec(f*x+e))^2-1/6*d*(4*c^3+32*c^2*d-80*c*d^2+32*
d^3+d*(2*c^2+16*c*d-21*d^2)*sec(f*x+e))*tan(f*x+e)/a^2/f
```

3.218.2 Mathematica [A] (verified)

Time = 5.10 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.61

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{-24d^2(12c^2 - 16cd + 7d^2) \cos^4\left(\frac{1}{2}(e+fx)\right) \left(\log\left(\cos\left(\frac{1}{2}(e+fx)\right) - \sin\left(\frac{1}{2}(e+fx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(e+fx)\right) + \sin\left(\frac{1}{2}(e+fx)\right)\right)\right)}{(12a^2f(1+\cos(e+fx)))^2}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^2,x]`output `(-24*d^2*(12*c^2 - 16*c*d + 7*d^2)*Cos[(e + f*x)/2]^4*(Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] - Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]]) + 2*Cos[(e + f*x)/2]*(2*c^4 + 16*c^3*d - 60*c^2*d^2 + 112*c*d^3 - 37*d^4 + 6*(c^4 + 2*c^3*d - 12*c^2*d^2 + 28*c*d^3 - 10*d^4)*Cos[e + f*x] + (2*c^4 + 16*c^3*d - 60*c^2*d^2 + 112*c*d^3 - 43*d^4)*Cos[2*(e + f*x)] + 2*c^4*Cos[3*(e + f*x)] + 4*c^3*d*Cos[3*(e + f*x)] - 24*c^2*d^2*Cos[3*(e + f*x)] + 40*c*d^3*Cos[3*(e + f*x)] - 16*d^4*Cos[3*(e + f*x)])*Sec[e + f*x]^2*Sin[(e + f*x)/2]/(12*a^2*f*(1 + Cos[e + f*x])^2)`**3.218.3 Rubi [A] (verified)**Time = 0.48 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.56, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4475, 109, 25, 27, 167, 27, 164, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a\sec(e+fx)+a)^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^4}{(a\csc\left(e+fx+\frac{\pi}{2}\right)+a)^2} dx$$

$$\downarrow \text{4475}$$

$$-\frac{a^2 \tan(e+fx) \int \frac{(c+d\sec(e+fx))^4}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

3.218. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

↓ 109

$$\frac{a^2 \tan(e + fx) \left(-\frac{\int \frac{a^2(c+d \sec(e+fx))^2(c^2+5dc-3d^2-(2c-5d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))^2(c^2+5dc-3d^2-(2c-5d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d \sec(e+fx))^2(c^2+5dc-3d^2-(2c-5d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 167

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^2 d(c+d \sec(e+fx))((19c-16d)d-(2c^2+16dc-21d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{3a} - \frac{(c-d)(c+8d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e + fx) \left(\frac{d \int \frac{(c+d \sec(e+fx))((19c-16d)d-(2c^2+16dc-21d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{3a} - \frac{(c-d)(c+8d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 164

3.218. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^2} dx$

$$\begin{array}{l}
 a^2 \tan(e + fx) \left(\frac{d \left(\frac{3}{2} d(12c^2 - 16cd + 7d^2) \int \frac{1}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a}} d \sec(e + fx) + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(2c^2 + 16cd - 21d^2) \sec(e + fx))}{2a^2}}{a} \right)}{3a} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)} \\
 \downarrow 45 \\
 a^2 \tan(e + fx) \left(\frac{d \left(3d(12c^2 - 16cd + 7d^2) \int \frac{(a - a \sec(e + fx))a}{\sec(e + fx)a + a} - a \frac{d \sqrt{a - a \sec(e + fx)}}{\sqrt{\sec(e + fx)a + a}} + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(2c^2 + 16cd - 21d^2) \sec(e + fx))}{2a^2}}{a} \right)}{3a} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)} \\
 \downarrow 218 \\
 a^2 \tan(e + fx) \left(\frac{d \left(\frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(2c^2 + 16cd - 21d^2) \sec(e + fx) + 4(c^3 + 8c^2d - 20cd^2 + 8d^3))}{2a^2} - \frac{3d(12c^2 - 16cd + 7d^2) \arctan\left(\frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{a \sec(e + fx) + a}}\right)}{a} \right)}{a} \right)}{3a} \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}
 \end{array}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^2,x]`

output `-((a^2*(-1/3*((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^3)/(a^2*(a + a*Sec[e + f*x])^(3/2)) + (-(((c - d)*(c + 8*d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(a^2*Sqrt[a + a*Sec[e + f*x]])) + (d*((-3*d*(12*c^2 - 16*c*d + 7*d^2)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]])/a + (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(4*(c^3 + 8*c^2*d - 20*c*d^2 + 8*d^3) + d*(2*c^2 + 16*c*d - 21*d^2)*Sec[e + f*x]))/(2*a^2)))/a)/(3*a))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]))`

3.218. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^2} dx$

3.218.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 109 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 164 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`
- rule 167 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] - Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[b*c*(f*g - e*h)*(m + 1) + (b*g - a*h)*(d*e*n + c*f*(p + 1)) + d*(b*(f*g - e*h)*(m + 1) + f*(b*g - a*h)*(n + p + 1))*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, p}, x] && LtQ[m, -1] && GtQ[n, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]]) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.218.4 Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.42

method	result
parallelrisch	$\frac{-12(1+\cos(2fx+2e))(c^2-\frac{4}{3}cd+\frac{7}{12}d^2)d^2 \ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)-1\right)+12(1+\cos(2fx+2e))(c^2-\frac{4}{3}cd+\frac{7}{12}d^2)d^2 \ln\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^4 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^3 d}{3} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^2 d^2 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c d^3}{3} - \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^4}{3} + \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^4 + 4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^3 d}$
derivativedivides	$\frac{-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^4 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^3 d}{3} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^2 d^2 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c d^3}{3} - \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^4}{3} + \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^4 + 4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^3 d}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^4 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^3 d}{3} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^2 d^2 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c d^3}{3} - \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^4}{3} + \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^4 + 4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^3 d}$
default	$\frac{-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^4 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^3 d}{3} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^2 d^2 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c d^3}{3} - \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^4}{3} + \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^4 + 4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^3 d}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^4 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^3 d}{3} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^2 d^2 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c d^3}{3} - \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^4}{3} + \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^4 + 4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^3 d}$
norman	$\frac{(c^4-4c^3d+6c^2d^2-4cd^3+d^4) \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^{11}}{6af} + \frac{(c^4+4c^3d-18c^2d^2+36cd^3-13d^4) \tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{2af} - \frac{(3c^4+4c^3d-30c^2d^2+44cd^3-19d^4)}{af}$
risch	$\frac{i(-36c^2d^2e^{6i(fx+e)}-120c^2d^2e^{4i(fx+e)}+256cd^3e^{2i(fx+e)}+16c^3de^{2i(fx+e)}+144cd^3e^{5i(fx+e)}+224cd^3e^{4i(fx+e)}-108c^4d^4)}{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^4 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^3 d}{3} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c^2 d^2 + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 c d^3}{3} - \frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^4}{3} + \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^4 + 4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right) c^3 d}$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

$$3.218. \int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

```
output 1/2*(-12*(1+cos(2*f*x+2*e))*(c^2-4/3*c*d+7/12*d^2)*d^2*ln(tan(1/2*f*x+1/2*
e)-1)+12*(1+cos(2*f*x+2*e))*(c^2-4/3*c*d+7/12*d^2)*d^2*ln(tan(1/2*f*x+1/2*
e)+1)+sec(1/2*f*x+1/2*e)^2*tan(1/2*f*x+1/2*e)*((1/3*c^4+8/3*c^3*d-10*c^2*d
^2+56/3*c*d^3-43/6*d^4)*cos(2*f*x+2*e)+(1/3*c^4+2/3*c^3*d-4*c^2*d^2+20/3*c
*d^3-8/3*d^4)*cos(3*f*x+3*e)+(c^4+2*c^3*d-12*c^2*d^2+28*c*d^3-10*d^4)*cos(
f*x+e)+1/3*c^4+8/3*c^3*d-10*c^2*d^2+56/3*c*d^3-37/6*d^4))/f/a^2/(1+cos(2*f
*x+2*e))
```

3.218.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.87

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{3((12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e)^4 + 2(12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e)^3 + (12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e)^2 + 2(12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e) + 12c^2d^2 - 16cd^3 + 7d^4)}{a^2}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x, algorithm="f
ricas")
```

```
output 1/12*(3*((12*c^2*d^2 - 16*c*d^3 + 7*d^4)*cos(f*x + e)^4 + 2*(12*c^2*d^2 -
16*c*d^3 + 7*d^4)*cos(f*x + e)^3 + (12*c^2*d^2 - 16*c*d^3 + 7*d^4)*cos(f*x
+ e)^2)*log(sin(f*x + e) + 1) - 3*((12*c^2*d^2 - 16*c*d^3 + 7*d^4)*cos(f*
x + e)^4 + 2*(12*c^2*d^2 - 16*c*d^3 + 7*d^4)*cos(f*x + e)^3 + (12*c^2*d^2
- 16*c*d^3 + 7*d^4)*cos(f*x + e)^2)*log(-sin(f*x + e) + 1) + 2*(3*d^4 + 4*
(c^4 + 2*c^3*d - 12*c^2*d^2 + 20*c*d^3 - 8*d^4)*cos(f*x + e)^3 + (2*c^4 +
16*c^3*d - 60*c^2*d^2 + 112*c*d^3 - 43*d^4)*cos(f*x + e)^2 + 6*(4*c*d^3 -
d^4)*cos(f*x + e))*sin(f*x + e))/(a^2*f*cos(f*x + e)^4 + 2*a^2*f*cos(f*x +
e)^3 + a^2*f*cos(f*x + e)^2)
```

3.218.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \int \frac{c^4 \sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{d^4 \sec^5(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{4cd^3 \sec^4(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{6c^2d^2 \sec^3(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx$$

$$= \frac{3((12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e)^4 + 2(12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e)^3 + (12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e)^2 + 2(12c^2d^2 - 16cd^3 + 7d^4)\cos(fx+e) + 12c^2d^2 - 16cd^3 + 7d^4)}{a^2}$$

3.218. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**4/(a+a*sec(f*x+e))**2,x)`

output `(Integral(c**4*sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(d**4*sec(e + f*x)**5/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(4*c*d**3*sec(e + f*x)**4/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(6*c**2*d**2*sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(4*c**3*d*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.218.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 536 vs. $2(184) = 368$.

Time = 0.23 (sec) , antiderivative size = 536, normalized size of antiderivative = 2.78

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{(a + a \sec(e + fx))^2} dx =$$

$$d^4 \left(\frac{6 \left(\frac{3 \sin(fx+e)}{\cos(fx+e)+1} - \frac{5 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^2 - \frac{2 a^2 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{a^2 \sin(fx+e)^4}{(\cos(fx+e)+1)^4}} + \frac{21 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} - \frac{21 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^2} + \frac{21 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^2} \right) -$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output `-1/6*(d^4*(6*(3*sin(f*x + e)/(cos(f*x + e) + 1) - 5*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^2 - 2*a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^2*sin(f*x + e)^4/(cos(f*x + e) + 1)^4) + (21*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 21*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 21*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 - 4*c*d^3*((15*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 12*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 12*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 + 12*sin(f*x + e)/((a^2 - a^2*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1))) + 6*c^2*d^2*((9*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 6*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 6*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 - 4*c^3*d*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - c^4*(3*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f`

3.218.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.86

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{3(12c^2d^2-16cd^3+7d^4)\log\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+1\right|}{a^2} - \frac{3(12c^2d^2-16cd^3+7d^4)\log\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-1\right|}{a^2} - \frac{6\left(8cd^3\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3-5d^4\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3-8cd^3\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+3d^4\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)\right)}{a^2} + \frac{a^4c^4\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3-4a^4c^3d\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3+6a^4c^2d^2\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3-4a^4c^2d^3\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3+a^4d^4\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3-3a^4c^4\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-12a^4c^3d\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+54a^4c^2d^2\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-60a^4c^2d^3\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+21a^4d^4\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)}{a^6} / f$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^2,x, algorithm="giac")
```

```
output 1/6*(3*(12*c^2*d^2 - 16*c*d^3 + 7*d^4)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^2 - 3*(12*c^2*d^2 - 16*c*d^3 + 7*d^4)*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a^2 - 6*(8*c*d^3*tan(1/2*f*x + 1/2*e)^3 - 5*d^4*tan(1/2*f*x + 1/2*e)^3 - 8*c*d^3*tan(1/2*f*x + 1/2*e) + 3*d^4*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1/2*e)^2 - 1)^2*a^2) - (a^4*c^4*tan(1/2*f*x + 1/2*e)^3 - 4*a^4*c^3*d*tan(1/2*f*x + 1/2*e)^3 + 6*a^4*c^2*d^2*tan(1/2*f*x + 1/2*e)^3 - 4*a^4*c^2*d^3*tan(1/2*f*x + 1/2*e)^3 + a^4*d^4*tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^4*tan(1/2*f*x + 1/2*e) - 12*a^4*c^3*d*tan(1/2*f*x + 1/2*e) + 54*a^4*c^2*d^2*tan(1/2*f*x + 1/2*e) - 60*a^4*c^2*d^3*tan(1/2*f*x + 1/2*e) + 21*a^4*d^4*tan(1/2*f*x + 1/2*e))/a^6)/f
```

3.218.9 Mupad [B] (verification not implemented)

Time = 13.70 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (8cd^3 - 3d^4) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (8cd^3 - 5d^4)}{f \left(a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + a^2 \right)}$$

$$- \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{3(c-d)^4}{2a^2} - \frac{2(c+d)(c-d)^3}{a^2} \right)}{f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (c-d)^4}{6a^2 f}$$

$$+ \frac{d^2 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (12c^2 - 16cd + 7d^2)}{a^2 f}$$

input `int((c + d/cos(e + f*x))^4/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output `(tan(e/2 + (f*x)/2)*(8*c*d^3 - 3*d^4) - tan(e/2 + (f*x)/2)^3*(8*c*d^3 - 5*d^4))/(f*(a^2*tan(e/2 + (f*x)/2)^4 - 2*a^2*tan(e/2 + (f*x)/2)^2 + a^2)) - (tan(e/2 + (f*x)/2)*((3*(c - d)^4)/(2*a^2) - (2*(c + d)*(c - d)^3)/a^2))/f - (tan(e/2 + (f*x)/2)^3*(c - d)^4)/(6*a^2*f) + (d^2*atanh(tan(e/2 + (f*x)/2))*(12*c^2 - 16*c*d + 7*d^2))/(a^2*f)`

3.218. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^2} dx$

3.219
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^2} dx$$

3.219.1 Optimal result 1570
 3.219.2 Mathematica [B] (verified) 1570
 3.219.3 Rubi [A] (verified) 1571
 3.219.4 Maple [A] (verified) 1574
 3.219.5 Fricas [B] (verification not implemented) 1575
 3.219.6 Sympy [F] 1575
 3.219.7 Maxima [B] (verification not implemented) 1576
 3.219.8 Giac [A] (verification not implemented) 1577
 3.219.9 Mupad [B] (verification not implemented) 1577

3.219.1 Optimal result

Integrand size = 31, antiderivative size = 133

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^2} dx$$

$$= \frac{(3c-2d)d^2 \arctanh(\sin(e+fx))}{a^2 f} + \frac{(c-d)(c+d \sec(e+fx))^2 \tan(e+fx)}{3f(a+a \sec(e+fx))^2}$$

$$+ \frac{(c^3+4c^2d-12cd^2+10d^3-(c-4d)d^2 \sec(e+fx)) \tan(e+fx)}{3f(a^2+a^2 \sec(e+fx))}$$

output `(3*c-2*d)*d^2*arctanh(sin(f*x+e))/a^2/f+1/3*(c-d)*(c+d*sec(f*x+e))^2*tan(f*x+e)/f/(a+a*sec(f*x+e))^2+1/3*(c^3+4*c^2*d-12*c*d^2+10*d^3-(c-4*d)*d^2*sec(f*x+e))*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))`

3.219.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 294 vs. 2(133) = 266.

Time = 3.65 (sec) , antiderivative size = 294, normalized size of antiderivative = 2.21

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^2} dx$$

$$= \frac{2 \cos^6\left(\frac{1}{2}(e+fx)\right) \sec(e+fx) (6d^2(-3c+2d) (\log(\cos(\frac{1}{2}(e+fx))) - \sin(\frac{1}{2}(e+fx)))) - \log(\cos(\frac{1}{2}(e+fx)))}{(a+a \sec(e+fx))^2}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^2,x]`

output $(2*\text{Cos}[(e + f*x)/2]^6*\text{Sec}[e + f*x]*(6*d^2*(-3*c + 2*d)*(\text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] - \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]]) - 8*(c - d)^3*\text{Csc}[e + f*x]^3*\text{Sin}[(e + f*x)/2]^4 + 32*(c - d)^3*\text{Csc}[e + f*x]^5*\text{Sin}[(e + f*x)/2]^8 + 2*(2*c^3 + 3*c^2*d - 12*c*d^2 + 13*d^3)*\text{Tan}[(e + f*x)/2] + 6*(3*c - 2*d)*d^2*(\text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] - \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]])*\text{Tan}[(e + f*x)/2]^2 - 2*(c - d)^2*(2*c + 7*d)*\text{Tan}[(e + f*x)/2]^3))/(3*a^2*f*(1 + \text{Cos}[e + f*x])^2)$

3.219.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.67, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 4475, 109, 25, 27, 160, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a \sec(e + fx) + a)^2} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})(c + d \csc(e + fx + \frac{\pi}{2}))^3}{(a \csc(e + fx + \frac{\pi}{2}) + a)^2} dx$$

↓ 4475

$$-\frac{a^2 \tan(e + fx) \int \frac{(c + d \sec(e + fx))^3}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{5/2}} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 109

$$a^2 \tan(e + fx) \left(-\frac{\int -\frac{a^2(c + d \sec(e + fx))(c^2 + 4dc - 2d^2 - (c - 4d)d \sec(e + fx))}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{3/2}} d \sec(e + fx)}{3a^3} - \frac{(c - d)\sqrt{a - a \sec(e + fx)}(c + d \sec(e + fx))^2}{3a^2(a \sec(e + fx) + a)^{3/2}} \right)$$

$$\frac{a^2 \tan(e + fx) \left(-\frac{\int -\frac{a^2(c + d \sec(e + fx))(c^2 + 4dc - 2d^2 - (c - 4d)d \sec(e + fx))}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{3/2}} d \sec(e + fx)}{3a^3} - \frac{(c - d)\sqrt{a - a \sec(e + fx)}(c + d \sec(e + fx))^2}{3a^2(a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 25

3.219. $\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + a \sec(e + fx))^2} dx$

$$\begin{aligned}
 & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))(c^2+4dc-2d^2-(c-4d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{(c+d \sec(e+fx))(c^2+4dc-2d^2-(c-4d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow \text{160} \\
 & \frac{a^2 \tan(e+fx) \left(\frac{3d^2(3c-2d) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{a} - \frac{\sqrt{a-a \sec(e+fx)}(c^3+4c^2d-d^2(c-4d) \sec(e+fx)-12cd^2+10d^3)}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow \text{45} \\
 & \frac{a^2 \tan(e+fx) \left(\frac{6d^2(3c-2d) \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}}{a} - \frac{\sqrt{a-a \sec(e+fx)}(c^3+4c^2d-d^2(c-4d) \sec(e+fx)-12cd^2+10d^3)}{a^2 \sqrt{a \sec(e+fx)+a}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow \text{218} \\
 & \frac{a^2 \tan(e+fx) \left(-\frac{6d^2(3c-2d) \arctan\left(\frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}}\right)}{a^2} - \frac{\sqrt{a-a \sec(e+fx)}(c^3+4c^2d-d^2(c-4d) \sec(e+fx)-12cd^2+10d^3)}{a^2 \sqrt{a \sec(e+fx)+a}} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^2,x]`

3.219. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^2} dx$

```
output 
$$-\left(\frac{a^2(-1/3((c-d)\sqrt{a-a\sec[e+fx]})(c+d\sec[e+fx])^2)}{(a^2(a+a\sec[e+fx])^{3/2})} + \frac{((-6(3c-2d)d^2\text{ArcTan}[\sqrt{a-a\sec[e+fx]}/\sqrt{a+a\sec[e+fx]})]/a^2 - (\sqrt{a-a\sec[e+fx]})(c^3 + 4c^2d - 12cd^2 + 10d^3 - (c-4d)d^2\sec[e+fx]))}{(a^2\sqrt{a+a\sec[e+fx]})} \right) / (3a) \cdot \frac{\tan[e+fx]}{(f\sqrt{a-a\sec[e+fx]}\sqrt{a+a\sec[e+fx]})}$$

```

3.219.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 109 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

```
rule 160 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))*((g_) + (h_)*(x_)), x_] := Simp[(b^2*d*e*g - a^2*d*f*h*m - a*b*(d*(f*g + e*h) - c*f*h*(m + 1)) + b*f*h*(b*c - a*d)*(m + 1)*x*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d*(b*c - a*d)*(m + 1))), x] + Simp[(a*d*f*h*m + b*(d*(f*g + e*h) - c*f*h*(m + 2)))/(b^2*d) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])
```


output $\frac{1}{6}(-18(c-2/3d)\cos(fx+e)d^2\ln(\tan(1/2fx+1/2e)-1)+18(c-2/3d)\cos(fx+e)d^2\ln(\tan(1/2fx+1/2e)+1)+\tan(1/2fx+1/2e)\sec(1/2fx+1/2e)^2((c^3+3/2c^2d-6cd^2+5d^3)\cos(2fx+2e)+(c^3+6c^2d-15cd^2+14d^3)\cos(fx+e)+c^3+3/2c^2d-6cd^2+8d^3))/f/a^2/\cos(fx+e)$

3.219.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. $2(130) = 260$.

Time = 0.28 (sec) , antiderivative size = 268, normalized size of antiderivative = 2.02

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{3((3cd^2-2d^3)\cos(fx+e)^3+2(3cd^2-2d^3)\cos(fx+e)^2+(3cd^2-2d^3)\cos(fx+e))\log(\sin(fx+e))}{a^2}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e))^2,x, algorithm="fracas")`

output $\frac{1}{6}(3*((3c^2d^2-2d^3)\cos(fx+e)^3+2*(3c^2d^2-2d^3)\cos(fx+e)^2+(3c^2d^2-2d^3)\cos(fx+e))*\log(\sin(fx+e)+1)-3*((3c^2d^2-2d^3)\cos(fx+e)^3+2*(3c^2d^2-2d^3)\cos(fx+e)^2+(3c^2d^2-2d^3)\cos(fx+e))*\log(-\sin(fx+e)+1)+2*(3d^3+(2c^3+3c^2d-12cd^2+10d^3)\cos(fx+e)^2+(c^3+6c^2d-15cd^2+14d^3)\cos(fx+e))*\sin(fx+e))/(a^2f\cos(fx+e)^3+2a^2f\cos(fx+e)^2+a^2f\cos(fx+e))$

3.219.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{\int \frac{c^3\sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{d^3\sec^4(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{3cd^2\sec^3(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{3c^2d\sec^2(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx}{a^2}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**3/(a+a*sec(f*x+e))**2,x)`

3.219. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$


```
output (Integral(c**3*sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + I
ntegral(d**3*sec(e + f*x)**4/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) +
Integral(3*c*d**2*sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1),
x) + Integral(3*c**2*d*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) +
1), x))/a**2
```

3.219.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 342 vs. $2(130) = 260$.

Time = 0.25 (sec) , antiderivative size = 342, normalized size of antiderivative = 2.57

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + a \sec(e + fx))^2} dx$$

$$= d^3 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} - \frac{12 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^2} + \frac{12 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^2} + \frac{12 \sin(fx+e)}{\left(a^2 - \frac{a^2 \sin^2(fx+e)}{(\cos(fx+e)+1)^2}\right)(\cos(fx+e)+1)} \right) - 3$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e))^2,x, algorithm="m
axima")
```

```
output 1/6*(d^3*((15*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x +
e) + 1)^3)/a^2 - 12*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 12*log(
sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 + 12*sin(f*x + e)/((a^2 - a^2*sin
(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1))) - 3*c*d^2*((9*sin(f
*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 6*
log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 6*log(sin(f*x + e)/(cos(f*x
+ e) + 1) - 1)/a^2) + 3*c^2*d*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*
x + e)^3/(cos(f*x + e) + 1)^3)/a^2 + c^3*(3*sin(f*x + e)/(cos(f*x + e) + 1
) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f
```

3.219.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.88

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx =$$

$$\frac{12d^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)a^2} - \frac{6(3cd^2 - 2d^3) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a^2} + \frac{6(3cd^2 - 2d^3) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a^2} + \frac{a^4 c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3}{a^2}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output `-1/6*(12*d^3*tan(1/2*f*x + 1/2*e)/((tan(1/2*f*x + 1/2*e)^2 - 1)*a^2) - 6*(3*c*d^2 - 2*d^3)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^2 + 6*(3*c*d^2 - 2*d^3)*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a^2 + (a^4*c^3*tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^2*d*tan(1/2*f*x + 1/2*e)^3 + 3*a^4*c*d^2*tan(1/2*f*x + 1/2*e)^3 - a^4*d^3*tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^3*tan(1/2*f*x + 1/2*e) - 9*a^4*c^2*d*tan(1/2*f*x + 1/2*e) + 27*a^4*c*d^2*tan(1/2*f*x + 1/2*e) - 15*a^4*d^3*tan(1/2*f*x + 1/2*e))/a^6)/f`

3.219.9 Mupad [B] (verification not implemented)

Time = 13.52 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.02

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx = \frac{2d^2 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (3c - 2d)}{a^2 f}$$

$$- \frac{2d^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - a^2\right)}$$

$$- \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (c - d)^3}{6a^2 f}$$

$$- \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{(c-d)^3}{a^2} - \frac{3(c+d)(c-d)^2}{2a^2}\right)}{f}$$

input `int((c + d/cos(e + f*x))^3/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output $(2*d^2*atanh(\tan(e/2 + (f*x)/2))*(3*c - 2*d))/(a^2*f) - (2*d^3*\tan(e/2 + (f*x)/2))/(f*(a^2*\tan(e/2 + (f*x)/2)^2 - a^2)) - (\tan(e/2 + (f*x)/2)^3*(c - d)^3)/(6*a^2*f) - (\tan(e/2 + (f*x)/2)*((c - d)^3/a^2 - (3*(c + d)*(c - d)^2)/(2*a^2)))/f$

3.219. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^2} dx$

3.220 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^2} dx$

3.220.1 Optimal result 1579
 3.220.2 Mathematica [B] (verified) 1579
 3.220.3 Rubi [A] (verified) 1580
 3.220.4 Maple [A] (verified) 1583
 3.220.5 Fracas [A] (verification not implemented) 1583
 3.220.6 Sympy [F] 1584
 3.220.7 Maxima [B] (verification not implemented) 1584
 3.220.8 Giac [A] (verification not implemented) 1585
 3.220.9 Mupad [B] (verification not implemented) 1585

3.220.1 Optimal result

Integrand size = 31, antiderivative size = 89

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{(a + a \sec(e + fx))^2} dx = \frac{d^2 \operatorname{arctanh}(\sin(e + fx))}{a^2 f} + \frac{(c - d)^2 \tan(e + fx)}{3f(a + a \sec(e + fx))^2} + \frac{(c - d)(c + 5d) \tan(e + fx)}{3f(a^2 + a^2 \sec(e + fx))}$$

output `d^2*arctanh(sin(f*x+e))/a^2/f+1/3*(c-d)^2*tan(f*x+e)/f/(a+a*sec(f*x+e))^2+1/3*(c-d)*(c+5*d)*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))`

3.220.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 181 vs. 2(89) = 178.

Time = 2.41 (sec) , antiderivative size = 181, normalized size of antiderivative = 2.03

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{(a + a \sec(e + fx))^2} dx = \frac{2 \cos\left(\frac{1}{2}(e + fx)\right) (6d^2 \cos^3\left(\frac{1}{2}(e + fx)\right) (\log(\cos\left(\frac{1}{2}(e + fx)\right) - \sin\left(\frac{1}{2}(e + fx)\right))) - \log(\cos\left(\frac{1}{2}(e + fx)\right))}{(a + a \sec(e + fx))^2}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^2,x]`

output $(-2*\text{Cos}[(e + f*x)/2]*(6*d^2*\text{Cos}[(e + f*x)/2]^3*(\text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] - \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]]) + (c - d)^2*\text{Sec}[e/2]*\text{Sin}[(f*x)/2] - 4*(c^2 + c*d - 2*d^2)*\text{Cos}[(e + f*x)/2]^2*\text{Sec}[e/2]*\text{Sin}[(f*x)/2] + (c - d)^2*\text{Cos}[(e + f*x)/2]*\text{Tan}[e/2]))/(3*a^2*f*(1 + \text{Cos}[e + f*x])^2)$

3.220.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 178, normalized size of antiderivative = 2.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3042, 4475, 100, 27, 87, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{(a \sec(e + fx) + a)^2} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})(c + d \csc(e + fx + \frac{\pi}{2}))^2}{(a \csc(e + fx + \frac{\pi}{2}) + a)^2} dx$$

↓ 4475

$$\frac{a^2 \tan(e + fx) \int \frac{(c + d \sec(e + fx))^2}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{5/2}} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 100

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^3(c^2 + 4dc - 2d^2 + 3d^2 \sec(e + fx))}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{3/2}} d \sec(e + fx)}{3a^4} - \frac{(c - d)^2 \sqrt{a - a \sec(e + fx)}}{3a^2(a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 27

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{c^2 + 4dc - 2d^2 + 3d^2 \sec(e + fx)}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{3/2}} d \sec(e + fx)}{3a} - \frac{(c - d)^2 \sqrt{a - a \sec(e + fx)}}{3a^2(a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 87

$$\begin{aligned}
 & \frac{a^2 \tan(e + fx) \left(\frac{3a^2 \int \frac{1}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a}} d \sec(e + fx)}{a} - \frac{(c-d)(c+5d)\sqrt{a - a \sec(e + fx)}}{a^2 \sqrt{a \sec(e + fx) + a}} - \frac{(c-d)^2 \sqrt{a - a \sec(e + fx)}}{3a^2 (a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}} \\
 & \quad \downarrow 45 \\
 & \frac{a^2 \tan(e + fx) \left(\frac{6a^2 \int \frac{1}{\frac{(a - a \sec(e + fx))a}{\sec(e + fx)a + a} - a} d \frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{\sec(e + fx)a + a}}}{a} - \frac{(c-d)(c+5d)\sqrt{a - a \sec(e + fx)}}{a^2 \sqrt{a \sec(e + fx) + a}} - \frac{(c-d)^2 \sqrt{a - a \sec(e + fx)}}{3a^2 (a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}} \\
 & \quad \downarrow 218 \\
 & \frac{a^2 \tan(e + fx) \left(\frac{6a^2 \arctan\left(\frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{a \sec(e + fx) + a}}\right)}{a^2} - \frac{(c-d)(c+5d)\sqrt{a - a \sec(e + fx)}}{a^2 \sqrt{a \sec(e + fx) + a}} - \frac{(c-d)^2 \sqrt{a - a \sec(e + fx)}}{3a^2 (a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^2,x]`

output `-((a^2*(-1/3*((c - d)^2*Sqrt[a - a*Sec[e + f*x]])/(a^2*(a + a*Sec[e + f*x])^(3/2)) + ((-6*d^2*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]]])/a^2 - ((c - d)*(c + 5*d)*Sqrt[a - a*Sec[e + f*x]])/(a^2*Sqrt[a + a*Sec[e + f*x]]))/(3*a))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])`

3.220.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(- (b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`
- rule 100 `Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d^2*(d*e - c*f)*(n + 1))), x] - Simp[1/(d^2*(d*e - c*f)*(n + 1)) Int[(c + d*x)^(n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n + 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_)*(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.)^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.220.4 Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.94

method	result
parallelrisch	$\frac{-6 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) d^2 + 6 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) d^2 - (c-d) \left((c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 3c - 9d \right) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{6a^2 f}$
derivativedivides	$\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) cd + \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 cd}{3} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) c^2 - 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) d^2 - 2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) d^2 - \frac{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \tan\left(\frac{fx}{2} + \frac{e}{2}\right) d^2}{2f a^2}$
default	$\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) cd + \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 cd}{3} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right) c^2 - 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) d^2 - 2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) d^2 - \frac{c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \tan\left(\frac{fx}{2} + \frac{e}{2}\right) d^2}{2f a^2}$
risch	$\frac{2i(3c^2 e^{2i(fx+e)} - 3d^2 e^{2i(fx+e)} + 3c^2 e^{i(fx+e)} + 6d e^{i(fx+e)} c - 9d^2 e^{i(fx+e)} + 2c^2 + 2cd - 4d^2)}{3f a^2 (e^{i(fx+e)} + 1)^3} + \frac{d^2 \ln(e^{i(fx+e)} + i)}{a^2 f} - \frac{d^2}{a^2 f}$
norman	$\frac{-\frac{(c^2 - 2cd + d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{6af} + \frac{(c^2 + 2cd - 3d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2af} + \frac{(5c^2 + 2cd - 7d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{6af} - \frac{(7c^2 + 10cd - 17d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{6af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 a}$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `1/6*(-6*ln(tan(1/2*f*x+1/2*e)-1)*d^2+6*ln(tan(1/2*f*x+1/2*e)+1)*d^2-(c-d)*((c-d)*tan(1/2*f*x+1/2*e)^2-3*c-9*d)*tan(1/2*f*x+1/2*e))/a^2/f`

3.220.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.74

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{(a + a \sec(e + fx))^2} dx$$

$$= \frac{3(d^2 \cos(fx + e)^2 + 2d^2 \cos(fx + e) + d^2) \log(\sin(fx + e) + 1) - 3(d^2 \cos(fx + e)^2 + 2d^2 \cos(fx + e) + d^2) \log(-\sin(fx + e) + 1) + 2(c^2 + 4cd - 5d^2 + 2(c^2 + cd - 2d^2) \cos(fx + e)) \sin(fx + e)}{6(a^2 f \cos(fx + e)^2 + 2a^2 f)}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x, algorithm="fricas")`

output `1/6*(3*(d^2*cos(f*x + e)^2 + 2*d^2*cos(f*x + e) + d^2)*log(sin(f*x + e) + 1) - 3*(d^2*cos(f*x + e)^2 + 2*d^2*cos(f*x + e) + d^2)*log(-sin(f*x + e) + 1) + 2*(c^2 + 4*c*d - 5*d^2 + 2*(c^2 + c*d - 2*d^2)*cos(f*x + e))*sin(f*x + e))/(a^2*f*cos(f*x + e)^2 + 2*a^2*f*cos(f*x + e) + a^2*f)`

3.220. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^2} dx$

3.220.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{\int \frac{c^2 \sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{d^2 \sec^3(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{2cd \sec^2(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx}{a^2}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**2/(a+a*sec(f*x+e))**2,x)`

output `(Integral(c**2*sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(d**2*sec(e + f*x)**3/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(2*c*d*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.220.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 195 vs. $2(85) = 170$.

Time = 0.22 (sec) , antiderivative size = 195, normalized size of antiderivative = 2.19

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx =$$

$$\frac{d^2 \left(\frac{9 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} - \frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^2} + \frac{6 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^2} \right) - \frac{2cd \left(\frac{3 \sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^2} - c}{6f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output `-1/6*(d^2*((9*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - 6*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^2 + 6*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^2 - 2*c*d*(3*sin(f*x + e)/(cos(f*x + e) + 1) + sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2 - c^2*(3*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/a^2)/f`

3.220.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.78

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx$$

$$= \frac{\frac{6d^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a^2} - \frac{6d^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a^2} - \frac{a^4 c^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 2a^4 cd \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + a^4 d^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 3a^4 c}{a^6}}{6f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output `1/6*(6*d^2*log(abs(tan(1/2*f*x + 1/2*e) + 1))/a^2 - 6*d^2*log(abs(tan(1/2*f*x + 1/2*e) - 1))/a^2 - (a^4*c^2*tan(1/2*f*x + 1/2*e)^3 - 2*a^4*c*d*tan(1/2*f*x + 1/2*e)^3 + a^4*d^2*tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^2*tan(1/2*f*x + 1/2*e) - 6*a^4*c*d*tan(1/2*f*x + 1/2*e) + 9*a^4*d^2*tan(1/2*f*x + 1/2*e))/a^6)/f`

3.220.9 Mupad [B] (verification not implemented)

Time = 13.42 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^2} dx = \frac{2d^2 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a^2 f}$$

$$- \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{(c-d)^2}{2a^2} - \frac{c^2-d^2}{a^2}\right)}{f}$$

$$- \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (c-d)^2}{6a^2 f}$$

input `int((c + d/cos(e + f*x))^2/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output `(2*d^2*atanh(tan(e/2 + (f*x)/2)))/(a^2*f) - (tan(e/2 + (f*x)/2)*((c - d)^2/(2*a^2) - (c^2 - d^2)/a^2))/f - (tan(e/2 + (f*x)/2)^3*(c - d)^2)/(6*a^2*f)`

3.221 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^2} dx$

3.221.1 Optimal result 1586
 3.221.2 Mathematica [A] (verified) 1586
 3.221.3 Rubi [A] (verified) 1587
 3.221.4 Maple [A] (verified) 1588
 3.221.5 Fricas [A] (verification not implemented) 1589
 3.221.6 Sympy [F] 1589
 3.221.7 Maxima [A] (verification not implemented) 1589
 3.221.8 Giac [A] (verification not implemented) 1590
 3.221.9 Mupad [B] (verification not implemented) 1590

3.221.1 Optimal result

Integrand size = 29, antiderivative size = 65

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^2} dx = \frac{(c-d) \tan(e+fx)}{3f(a+a \sec(e+fx))^2} + \frac{(c+2d) \tan(e+fx)}{3f(a^2+a^2 \sec(e+fx))}$$

output `1/3*(c-d)*tan(f*x+e)/f/(a+a*sec(f*x+e))^2+1/3*(c+2*d)*tan(f*x+e)/f/(a^2+a^2*sec(f*x+e))`

3.221.2 Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.17

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^2} dx = \frac{\cos(\frac{1}{2}(e+fx)) \sec(\frac{e}{2}) (3(c+d) \sin(\frac{fx}{2}) - 3c \sin(e+\frac{fx}{2}) + (2c+d) \sin(e+\frac{3fx}{2}))}{3a^2 f(1+\cos(e+fx))^2}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2,x]`

output `(Cos[(e + f*x)/2]*Sec[e/2]*(3*(c + d)*Sin[(f*x)/2] - 3*c*Sin[e + (f*x)/2] + (2*c + d)*Sin[e + (3*f*x)/2]))/(3*a^2*f*(1 + Cos[e + f*x])^2)`

3.221.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$, Rules used = {3042, 4488, 3042, 4281}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a\sec(e+fx)+a)^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))}{(a\csc(e+fx+\frac{\pi}{2})+a)^2} dx$$

↓ 4488

$$\frac{(c+2d) \int \frac{\sec(e+fx)}{\sec(e+fx)a+a} dx}{3a} + \frac{(c-d)\tan(e+fx)}{3f(a\sec(e+fx)+a)^2}$$

↓ 3042

$$\frac{(c+2d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} + \frac{(c-d)\tan(e+fx)}{3f(a\sec(e+fx)+a)^2}$$

↓ 4281

$$\frac{(c+2d)\tan(e+fx)}{3af(a\sec(e+fx)+a)} + \frac{(c-d)\tan(e+fx)}{3f(a\sec(e+fx)+a)^2}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + a*Sec[e + f*x])^2,x]`

output `((c - d)*Tan[e + f*x])/(3*f*(a + a*Sec[e + f*x])^2) + ((c + 2*d)*Tan[e + f*x])/(3*a*f*(a + a*Sec[e + f*x]))`

3.221.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4281 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[-Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4488 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(A*b - a*B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*B*m + A*b*(m + 1))/(a*b*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && NeQ[a*B*m + A*b*(m + 1), 0] && LtQ[m, -2^(-1)]`

3.221.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

method	result	size
parallelrisch	$-\frac{\left(-3c-3d+(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)\right)^2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{6a^2f}$	42
derivativdivides	$-\frac{c \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{d \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + c \tan\left(\frac{fx}{2}+\frac{e}{2}\right) + d \tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{2fa^2}$	60
default	$-\frac{c \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + \frac{d \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3} + c \tan\left(\frac{fx}{2}+\frac{e}{2}\right) + d \tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{2fa^2}$	60
risch	$\frac{2i(3e^{2i(fx+e)}c+3e^{i(fx+e)}c+3de^{i(fx+e)}+2c+d)}{3fa^2(e^{i(fx+e)}+1)^3}$	64
norman	$-\frac{(c-d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{6af} - \frac{(c+d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{2af} + \frac{(2c+d)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{3af}$ $a\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2-1\right)$	89

input `int(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `-1/6*(-3*c-3*d+(c-d)*tan(1/2*f*x+1/2*e)^2)*tan(1/2*f*x+1/2*e)/a^2/f`

3.221.
$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^2} dx$$

3.221.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.89

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^2} dx = \frac{((2c+d)\cos(fx+e)+c+2d)\sin(fx+e)}{3(a^2f\cos(fx+e)^2+2a^2f\cos(fx+e)+a^2f)}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="fricas")`

output `1/3*((2*c + d)*cos(f*x + e) + c + 2*d)*sin(f*x + e)/(a^2*f*cos(f*x + e)^2 + 2*a^2*f*cos(f*x + e) + a^2*f)`

3.221.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^2} dx \\ &= \int \frac{c\sec(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx + \int \frac{d\sec^2(e+fx)}{\sec^2(e+fx)+2\sec(e+fx)+1} dx \\ & \qquad \qquad \qquad a^2 \end{aligned}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))**2,x)`

output `(Integral(c*sec(e + f*x)/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x) + Integral(d*sec(e + f*x)**2/(sec(e + f*x)**2 + 2*sec(e + f*x) + 1), x))/a**2`

3.221.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.43

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^2} dx = \frac{d\left(\frac{3\sin(fx+e)}{\cos(fx+e)+1} + \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3}\right)}{a^2} + \frac{c\left(\frac{3\sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^3}{(\cos(fx+e)+1)^3}\right)}{a^2}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="maxima")`

output $1/6*(d*(3*\sin(f*x + e)/(\cos(f*x + e) + 1) + \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2 + c*(3*\sin(f*x + e)/(\cos(f*x + e) + 1) - \sin(f*x + e)^3/(\cos(f*x + e) + 1)^3)/a^2)/f$

3.221.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.92

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{(a + a \sec(e + fx))^2} dx$$

$$= -\frac{c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 - d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 - 3c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 3d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{6a^2f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^2,x, algorithm="giac")`

output $-1/6*(c*\tan(1/2*f*x + 1/2*e)^3 - d*\tan(1/2*f*x + 1/2*e)^3 - 3*c*\tan(1/2*f*x + 1/2*e) - 3*d*\tan(1/2*f*x + 1/2*e))/(a^2*f)$

3.221.9 Mupad [B] (verification not implemented)

Time = 13.58 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.69

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{(a + a \sec(e + fx))^2} dx = \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (c + d)}{2a^2f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (c - d)}{6a^2f}$$

input `int((c + d/cos(e + f*x))/(cos(e + f*x)*(a + a/cos(e + f*x))^2),x)`

output $(\tan(e/2 + (f*x)/2)*(c + d))/(2*a^2*f) - (\tan(e/2 + (f*x)/2)^3*(c - d))/(6*a^2*f)$

3.222 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))} dx$

3.222.1 Optimal result 1591
 3.222.2 Mathematica [C] (verified) 1591
 3.222.3 Rubi [A] (verified) 1592
 3.222.4 Maple [A] (verified) 1595
 3.222.5 Fracas [B] (verification not implemented) 1596
 3.222.6 Sympy [F] 1596
 3.222.7 Maxima [F(-2)] 1597
 3.222.8 Giac [B] (verification not implemented) 1597
 3.222.9 Mupad [B] (verification not implemented) 1598

3.222.1 Optimal result

Integrand size = 31, antiderivative size = 129

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))} dx = \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a^2(c-d)^{5/2} \sqrt{c+df}} + \frac{\tan(e+fx)}{3(c-d)f(a+a \sec(e+fx))^2} + \frac{(c-4d) \tan(e+fx)}{3(c-d)^2 f(a^2+a^2 \sec(e+fx))}$$

```
output 2*d^2*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a^2/(c-d)^(5/2)/
f/(c+d)^(1/2)+1/3*tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))^2+1/3*(c-4*d)*tan(f*
x+e)/(c-d)^2/f/(a^2+a^2*sec(f*x+e))
```

3.222.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.05 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.62

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))} dx = \frac{\cos\left(\frac{1}{2}(e+fx)\right) \left(-\frac{24id^2 \arctan\left(\frac{(i \cos(e)+\sin(e))(c \sin(e)+(-d+c \cos(e)) \tan\left(\frac{fx}{2}\right))}{\sqrt{c^2-d^2} \sqrt{(\cos(e)-i \sin(e))^2}}\right) \cos^3\left(\frac{1}{2}(e+fx)\right) (\cos(e)-i \sin(e))}{\sqrt{c^2-d^2} \sqrt{(\cos(e)-i \sin(e))^2}} + \sec\left(\frac{e}{2}\right) (3(c-d)^2 f(1+\cos(e+fx)))^2 \right)}{3a^2(c-d)^2 f(1+\cos(e+fx))^2}$$

3.222. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))} dx$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])),x]`

output $(\text{Cos}[(e + f*x)/2]*(((-24*I)*d^2*\text{ArcTan}[\frac{(I*\text{Cos}[e] + \text{Sin}[e])*(c*\text{Sin}[e] + (-d + c*\text{Cos}[e])*\text{Tan}[(f*x)/2])}{\sqrt{c^2 - d^2}*\sqrt{(\text{Cos}[e] - I*\text{Sin}[e])^2}}] * \text{Cos}[(e + f*x)/2]^3 * (\text{Cos}[e] - I*\text{Sin}[e])) / (\sqrt{c^2 - d^2}*\sqrt{(\text{Cos}[e] - I*\text{Sin}[e])^2}) + \text{Sec}[e/2]*(3*(c - 3*d)*\text{Sin}[(f*x)/2] - 3*(c - 2*d)*\text{Sin}[e + (f*x)/2] + (2*c - 5*d)*\text{Sin}[e + (3*f*x)/2])) / (3*a^2*(c - d)^2*f*(1 + \text{Cos}[e + f*x])^2)$

3.222.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.70, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4475, 115, 25, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{(a \sec(e + fx) + a)^2 (c + d \sec(e + fx))} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{(a \csc(e + fx + \frac{\pi}{2}) + a)^2 (c + d \csc(e + fx + \frac{\pi}{2}))} dx$$

↓ 4475

$$\frac{a^2 \tan(e + fx) \int \frac{1}{\sqrt{a - a \sec(e + fx)} (\sec(e + fx) a + a)^{5/2} (c + d \sec(e + fx))} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 115

$$\frac{a^2 \tan(e + fx) \left(-\frac{\int -\frac{a^2 (c - 3d + d \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} (\sec(e + fx) a + a)^{3/2} (c + d \sec(e + fx))} d \sec(e + fx)}{3a^3 (c - d)} - \frac{\sqrt{a - a \sec(e + fx)}}{3a^2 (c - d) (a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 25

$$\frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^2 (c - 3d + d \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} (\sec(e + fx) a + a)^{3/2} (c + d \sec(e + fx))} d \sec(e + fx)}{3a^3 (c - d)} - \frac{\sqrt{a - a \sec(e + fx)}}{3a^2 (c - d) (a \sec(e + fx) + a)^{3/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

3.222. $\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))} dx$

$$\begin{array}{c}
\downarrow 27 \\
\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{c-3d+d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{3/2} (c+d \sec(e+fx))} d \sec(e+fx)}{3a(c-d)} - \frac{\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 169 \\
\frac{a^2 \tan(e+fx) \left(-\frac{\int -\frac{3a^2 d^2}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a} (c+d \sec(e+fx))} d \sec(e+fx)}{a^3(c-d)} - \frac{(c-4d) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 27 \\
\frac{a^2 \tan(e+fx) \left(\frac{3d^2 \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a} (c+d \sec(e+fx))} d \sec(e+fx)}{a(c-d)} - \frac{(c-4d) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 104 \\
\frac{a^2 \tan(e+fx) \left(\frac{6d^2 \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{a(c-d)} - \frac{(c-4d) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 218 \\
\frac{a^2 \tan(e+fx) \left(\frac{6d^2 \arctan\left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a \sec(e+fx)}}\right)}{a^2(c-d)^{3/2} \sqrt{c+d}} - \frac{(c-4d) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}
\end{array}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])),x]`

3.222. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))} dx$

```
output -((a^2*(-1/3*Sqrt[a - a*Sec[e + f*x]]/(a^2*(c - d)*(a + a*Sec[e + f*x])^(3/2)) + ((6*d^2*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]/(a^2*(c - d)^(3/2)*Sqrt[c + d]) - ((c - 4*d)*Sqrt[a - a*Sec[e + f*x]]/(a^2*(c - d)*Sqrt[a + a*Sec[e + f*x]]))/(3*a*(c - d))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.222.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 115 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]
```

```
rule 169 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]
```

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.222.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.95

method	result
derivativedivides	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{(c-d)^2} - \frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 3d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2fa^2} + \frac{4d^2 \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c-d)^2 \sqrt{(c+d)(c-d)}}$
default	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 - d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{(c-d)^2} - \frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 3d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{2fa^2} + \frac{4d^2 \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c-d)^2 \sqrt{(c+d)(c-d)}}$
risch	$\frac{2i(3e^{2i(fx+e)}c - 6de^{2i(fx+e)} + 3e^{i(fx+e)}c - 9de^{i(fx+e)} + 2c - 5d)}{3fa^2(c-d)^2(e^{i(fx+e)} + 1)^3} + \frac{d^2 \ln\left(\frac{e^{i(fx+e)} + \frac{ic^2 - id^2 + \sqrt{c^2 - d^2}d}{\sqrt{c^2 - d^2}c}}{\sqrt{c^2 - d^2}c}\right)}{\sqrt{c^2 - d^2}(c-d)^2fa^2} - \frac{d^2 \ln(e^{i(fx+e)})}{\sqrt{c^2 - d^2}(c-d)^2fa^2}$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/2/f/a^2*(-1/(c-d)^2*(1/3*c*tan(1/2*f*x+1/2*e)^3-1/3*d*tan(1/2*f*x+1/2*e)^3-c*tan(1/2*f*x+1/2*e)+3*d*tan(1/2*f*x+1/2*e))+4*d^2/(c-d)^2/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))`

$$3.222. \quad \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))} dx$$

3.222.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 270 vs. $2(116) = 232$.

Time = 0.29 (sec) , antiderivative size = 598, normalized size of antiderivative = 4.64

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))} dx$$

$$= \left[\frac{3(d^2 \cos^2(fx+e) + 2d^2 \cos(fx+e) + d^2) \sqrt{c^2-d^2} \log\left(\frac{2cd \cos(fx+e) - (c^2-2d^2) \cos(fx+e)^2 + 2\sqrt{c^2-d^2}(d \cos(fx+e) + c) \sin(fx+e) + c^2 \cos^2(fx+e) + 2cd \cos(fx+e) + d^2}{c^2 \cos^2(fx+e) + 2cd \cos(fx+e) + d^2}\right)}{6((a^2c^4 - 2a^2c^3d + 2a^2cd^3 - a^2d^4)f \cos(fx+e)^2 + 2(a^2c^4 - 2a^2c^3d + 2a^2cd^3 - a^2d^4)f \cos(fx+e) + (a^2c^4 - 2a^2c^3d + 2a^2cd^3 - a^2d^4))} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `[1/6*(3*(d^2*cos(f*x + e)^2 + 2*d^2*cos(f*x + e) + d^2)*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(c^3 - 4*c^2*d - c*d^2 + 4*d^3 + (2*c^3 - 5*c^2*d - 2*c*d^2 + 5*d^3)*cos(f*x + e))*sin(f*x + e))/((a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*f*cos(f*x + e)^2 + 2*(a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*f*cos(f*x + e) + (a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*f), 1/3*(3*(d^2*cos(f*x + e)^2 + 2*d^2*cos(f*x + e) + d^2)*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (c^3 - 4*c^2*d - c*d^2 + 4*d^3 + (2*c^3 - 5*c^2*d - 2*c*d^2 + 5*d^3)*cos(f*x + e))*sin(f*x + e))/((a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*f*cos(f*x + e)^2 + 2*(a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*f*cos(f*x + e) + (a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*f)]`

3.222.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))} dx$$

$$= \int \frac{\sec(e+fx)}{c \sec^2(e+fx) + 2c \sec(e+fx) + c + d \sec^3(e+fx) + 2d \sec^2(e+fx) + d \sec(e+fx)} dx$$

$$a^2$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e)),x)`

3.222. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))} dx$

output `Integral(sec(e + f*x)/(c*sec(e + f*x)**2 + 2*c*sec(e + f*x) + c + d*sec(e + f*x)**3 + 2*d*sec(e + f*x)**2 + d*sec(e + f*x)), x)/a**2`

3.222.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` for more de`

3.222.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 249 vs. $2(116) = 232$.

Time = 0.33 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.93

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))} dx = \frac{12 \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e) - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-c^2+d^2}} \right) \right) d^2}{(a^2 c^2 - 2 a^2 c d + a^2 d^2) \sqrt{-c^2+d^2}} + \frac{a^4 c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 - 2 a^4 c d \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + a^4 d^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3}{6 f}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="giac")`

output `-1/6*(12*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*d^2/((a^2*c^2 - 2*a^2*c*d + a^2*d^2)*sqrt(-c^2 + d^2)) + (a^4*c^2*tan(1/2*f*x + 1/2*e)^3 - 2*a^4*c*d*tan(1/2*f*x + 1/2*e)^3 + a^4*d^2*tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^2*tan(1/2*f*x + 1/2*e) + 12*a^4*c*d*tan(1/2*f*x + 1/2*e) - 9*a^4*d^2*tan(1/2*f*x + 1/2*e))/(a^6*c^3 - 3*a^6*c^2*d + 3*a^6*c*d^2 - a^6*d^3))/f`

3.222. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c+d \sec(e+fx))} dx$

3.222.9 Mupad [B] (verification not implemented)

Time = 13.88 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.30

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{1}{a^2(c-d)} - \frac{c+d}{2a^2(c-d)^2}\right)}{f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{6a^2 f (c-d)}$$

$$- \frac{d^2 \operatorname{atan}\left(\frac{\operatorname{li} \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^3 - 3i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^2 d + 3i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c d^2 - \operatorname{li} \tan\left(\frac{e}{2} + \frac{fx}{2}\right) d^3}{\sqrt{c+d}(c-d)^{5/2}}\right) 2i}{a^2 f \sqrt{c+d} (c-d)^{5/2}}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c + d/cos(e + f*x))),x)`output `(tan(e/2 + (f*x)/2)*(1/(a^2*(c - d)) - (c + d)/(2*a^2*(c - d)^2))/f - tan(e/2 + (f*x)/2)^3/(6*a^2*f*(c - d)) - (d^2*atan((c^3*tan(e/2 + (f*x)/2)*1i - d^3*tan(e/2 + (f*x)/2)*1i + c*d^2*tan(e/2 + (f*x)/2)*3i - c^2*d*tan(e/2 + (f*x)/2)*3i)/((c + d)^(1/2)*(c - d)^(5/2)))*2i)/(a^2*f*(c + d)^(1/2)*(c - d)^(5/2))`

3.223 $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2} dx$

3.223.1 Optimal result 1599
 3.223.2 Mathematica [C] (verified) 1600
 3.223.3 Rubi [A] (verified) 1600
 3.223.4 Maple [A] (verified) 1604
 3.223.5 Fricas [B] (verification not implemented) 1605
 3.223.6 Sympy [F] 1606
 3.223.7 Maxima [F(-2)] 1606
 3.223.8 Giac [B] (verification not implemented) 1606
 3.223.9 Mupad [B] (verification not implemented) 1607

3.223.1 Optimal result

Integrand size = 31, antiderivative size = 211

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2} dx$$

$$= \frac{2d^2(3c+2d)\operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a^2(c-d)^{7/2}(c+d)^{3/2}f} + \frac{d(c^2-6cd-10d^2)\tan(e+fx)}{3a^2(c-d)^3(c+d)f(c+d \sec(e+fx))}$$

$$+ \frac{(c-6d)\tan(e+fx)}{3a^2(c-d)^2f(1+\sec(e+fx))(c+d \sec(e+fx))}$$

$$+ \frac{\tan(e+fx)}{3(c-d)f(a+a \sec(e+fx))^2(c+d \sec(e+fx))}$$

output

```
2*d^2*(3*c+2*d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a^2/(c-d)^(7/2)/(c+d)^(3/2)/f+1/3*d*(c^2-6*c*d-10*d^2)*tan(f*x+e)/a^2/(c-d)^3/(c+d)/f/(c+d*sec(f*x+e))+1/3*(c-6*d)*tan(f*x+e)/a^2/(c-d)^2/f/(1+sec(f*x+e))/(c+d*sec(f*x+e))+1/3*tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))
```


3.223.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.96 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.78

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^2} dx$$

$$= \frac{2 \cos\left(\frac{1}{2}(e + fx)\right) (d + c \cos(e + fx)) \sec^4(e + fx) \left(\frac{12d^2(3c+2d) \arctan\left(\frac{(i \cos(e) + \sin(e))(c \sin(e) + (-d + c \cos(e)) \tan\left(\frac{fx}{2}\right))}{\sqrt{c^2 - d^2} \sqrt{(\cos(e) - i \sin(e))^2}}\right)}{(c+d)\sqrt{c^2 - d^2} \sqrt{(\cos(e) - i \sin(e))}} \right)}{\dots}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^2),x]`

output `(2*Cos[(e + f*x)/2]*(d + c*Cos[e + f*x])*Sec[e + f*x]^4*((12*d^2*(3*c + 2*d)*ArcTan[((I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))]/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]))*Cos[(e + f*x)/2]^3*(d + c*Cos[e + f*x])*(I*Cos[e] + Sin[e]))/((c + d)*Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + (c - d)*(d + c*Cos[e + f*x])*Sec[e/2]*Sin[(f*x)/2] - 4*(c - 4*d)*Cos[(e + f*x)/2]^2*(d + c*Cos[e + f*x])*Sec[e/2]*Sin[(f*x)/2] + (6*d^3*Cos[(e + f*x)/2]^3*(-(d*Sin[e] + c*Sin[f*x]))/(c*(c + d)*(Cos[e/2] - Sin[e/2])*(Cos[e/2] + Sin[e/2])) + (c - d)*Cos[(e + f*x)/2]*(d + c*Cos[e + f*x])*Tan[e/2]))/(3*a^2*(-c + d)^3*f*(1 + Sec[e + f*x])^2*(c + d*Sec[e + f*x])^2)`

3.223.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.46, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4475, 114, 27, 169, 25, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{(a \sec(e + fx) + a)^2 (c + d \sec(e + fx))^2} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right)}{\left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^2 \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2} dx$$

3.223. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2} dx$

$$\begin{aligned} & \downarrow 4475 \\ & \frac{a^2 \tan(e+fx) \int \frac{1}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 114 \\ & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(c+2d-2d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{a^2(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{3/2}(c+d \sec(e+fx))} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 27 \\ & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{c+2d-2d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{c^2-d^2} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{3/2}(c+d \sec(e+fx))} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 169 \\ & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2((c-6d)(c+d)+d(c+4d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(c+4d) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 25 \\ & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2((c-6d)(c+d)+d(c+4d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(c+4d) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 27 \\ & \frac{a^2 \tan(e+fx) \left(\frac{\int \frac{(c-6d)(c+d)+d(c+4d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a(c-d)} - \frac{(c+4d) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\ & \downarrow 169 \end{aligned}$$

3.223. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2} dx$

$$\begin{array}{c}
 a^2 \tan(e + fx) \left(\frac{\int - \frac{3a^2 d^2 (3c+2d)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}} d \sec(e+fx) - \frac{(c^2-6cd-10d^2) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(c+4d) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}}}{\frac{3a(c-d)}{c^2-d^2}} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} \\
 \downarrow 27 \\
 a^2 \tan(e + fx) \left(\frac{3d^2(3c+2d) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}} d \sec(e+fx) - \frac{(c^2-6cd-10d^2) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(c+4d) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}}}{\frac{3a(c-d)}{c^2-d^2}} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} \\
 \downarrow 104 \\
 a^2 \tan(e + fx) \left(\frac{6d^2(3c+2d) \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}} - \frac{(c^2-6cd-10d^2) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(c+4d) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}}}{\frac{3a(c-d)}{c^2-d^2}} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} \\
 \downarrow 218 \\
 a^2 \tan(e + fx) \left(\frac{6d^2(3c+2d) \arctan\left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a \sec(e+fx)}}\right) - \frac{(c^2-6cd-10d^2) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(c+4d) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}}}{\frac{3a(c-d)}{c^2-d^2}} + \frac{1}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{3/2}} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}
 \end{array}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^2),x]`

3.223. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2} dx$

```
output -((a^2*((d*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c^2 - d^2)*(a + a*Sec[e + f*x])
^(3/2)*(c + d*Sec[e + f*x])) + (-1/3*((c + 4*d)*Sqrt[a - a*Sec[e + f*x]])/
(a^2*(c - d)*(a + a*Sec[e + f*x])^(3/2)) + ((6*d^2*(3*c + 2*d)*ArcTan[(Sqr
t[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])]
)/(a^2*(c - d)^(3/2)*Sqrt[c + d]) - ((c^2 - 6*c*d - 10*d^2)*Sqrt[a - a*Sec
[e + f*x]])/(a^2*(c - d)*Sqrt[a + a*Sec[e + f*x]]))/(3*a*(c - d))/(c^2 -
d^2))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])
```

3.223.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 114 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)
)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e
- a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1)
- b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] ||
IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

```
rule 169 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c +
d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + S
imp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n
*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a
h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[
2*m, 2*n, 2*p]
```

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*sqrt[a + b*Csc[e + f*x]]*sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.223.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.96

method	result
derivativedivides	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 5d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^2 - 2cd + d^2)(c - d)} - \frac{4d^2 \left(-\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} \right)^2}{(c-d)^3} \right)}{2fa^2}$
default	$-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 5d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^2 - 2cd + d^2)(c - d)} - \frac{4d^2 \left(-\frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c+d) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} \right)^2}{(c-d)^3} \right)}{2fa^2}$
risch	$\frac{2i(-3c^4 e^{4i(fx+e)} + 6c^3 d e^{4i(fx+e)} + 9c^2 d^2 e^{4i(fx+e)} + 3d^4 e^{4i(fx+e)} - 3c^4 e^{3i(fx+e)} + 6c^3 d e^{3i(fx+e)} + 27c^2 d^2 e^{3i(fx+e)} + 21c d^3 e^{3i(fx+e)} - 3c^3 d^3 e^{2i(fx+e)} + 6c^2 d^3 e^{2i(fx+e)} + 3c d^4 e^{2i(fx+e)} - 3c^2 d^4 e^{i(fx+e)} + 6c d^4 e^{i(fx+e)} - 3d^5 e^{i(fx+e)})}{(c^2 - 2cd + d^2)^2 (c + d \sec(e + fx))^2}$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output $\frac{1}{2}f/a^2*(-1/(c^2-2*c*d+d^2)/(c-d)*(1/3*c*\tan(1/2*f*x+1/2*e)^3-1/3*d*\tan(1/2*f*x+1/2*e)^3-c*\tan(1/2*f*x+1/2*e)+5*d*\tan(1/2*f*x+1/2*e))-4*d^2/(c-d)^3*(-d/(c+d)*\tan(1/2*f*x+1/2*e)/(\tan(1/2*f*x+1/2*e)^2*c-\tan(1/2*f*x+1/2*e)^2*d-c-d)-(3*c+2*d)/(c+d)/((c+d)*(c-d))^{1/2}*\operatorname{arctanh}((c-d)*\tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^{1/2})))$

3.223.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 592 vs. $2(196) = 392$.

Time = 0.33 (sec) , antiderivative size = 1242, normalized size of antiderivative = 5.89

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x, algorithm="fracas")`

output $[-1/6*(3*(3*c*d^3 + 2*d^4 + (3*c^2*d^2 + 2*c*d^3)*\cos(f*x + e))^3 + (6*c^2*d^2 + 7*c*d^3 + 2*d^4)*\cos(f*x + e)^2 + (3*c^2*d^2 + 8*c*d^3 + 4*d^4)*\cos(f*x + e))*\sqrt{c^2 - d^2}*\log((2*c*d*\cos(f*x + e) - (c^2 - 2*d^2)*\cos(f*x + e)^2 - 2*\sqrt{c^2 - d^2}*(d*\cos(f*x + e) + c)*\sin(f*x + e) + 2*c^2 - d^2)/(c^2*\cos(f*x + e)^2 + 2*c*d*\cos(f*x + e) + d^2)) - 2*(c^4*d - 6*c^3*d^2 - 11*c^2*d^3 + 6*c*d^4 + 10*d^5 + (2*c^5 - 6*c^4*d - 10*c^3*d^2 + 3*c^2*d^3 + 8*c*d^4 + 3*d^5)*\cos(f*x + e)^2 + (c^5 - 4*c^4*d - 14*c^3*d^2 - 10*c^2*d^3 + 13*c*d^4 + 14*d^5)*\cos(f*x + e))*\sin(f*x + e))/((a^2*c^7 - 2*a^2*c^6*d - a^2*c^5*d^2 + 4*a^2*c^4*d^3 - a^2*c^3*d^4 - 2*a^2*c^2*d^5 + a^2*c*d^6)*f*\cos(f*x + e)^3 + (2*a^2*c^7 - 3*a^2*c^6*d - 4*a^2*c^5*d^2 + 7*a^2*c^4*d^3 + 2*a^2*c^3*d^4 - 5*a^2*c^2*d^5 + a^2*d^7)*f*\cos(f*x + e)^2 + (a^2*c^7 - 5*a^2*c^5*d^2 + 2*a^2*c^4*d^3 + 7*a^2*c^3*d^4 - 4*a^2*c^2*d^5 - 3*a^2*c*d^6 + 2*a^2*d^7)*f*\cos(f*x + e) + (a^2*c^6*d - 2*a^2*c^5*d^2 - a^2*c^4*d^3 + 4*a^2*c^3*d^4 - a^2*c^2*d^5 - 2*a^2*c*d^6 + a^2*d^7)*f), 1/3*(3*(3*c*d^3 + 2*d^4 + (3*c^2*d^2 + 2*c*d^3)*\cos(f*x + e))^3 + (6*c^2*d^2 + 7*c*d^3 + 2*d^4)*\cos(f*x + e)^2 + (3*c^2*d^2 + 8*c*d^3 + 4*d^4)*\cos(f*x + e))*\sqrt{-(c^2 + d^2)}*\operatorname{arctan}(-\sqrt{-(c^2 + d^2)}*(d*\cos(f*x + e) + c)/((c^2 - d^2)*\sin(f*x + e))) + (c^4*d - 6*c^3*d^2 - 11*c^2*d^3 + 6*c*d^4 + 10*d^5 + (2*c^5 - 6*c^4*d - 10*c^3*d^2 + 3*c^2*d^3 + 8*c*d^4 + 3*d^5)*\cos(f*x + e)^2 + (c^5 - 4*c^4*d - 14*c^3*d^2 - 10*c^2*d^3 + 13*c*d^4 + 14*d^5)*\cos(f*x + e) + e...]$

3.223.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^2} dx$$

$$= \frac{\int \frac{\sec(e+fx)}{c^2 \sec^2(e+fx) + 2c^2 \sec(e+fx) + c^2 + 2cd \sec^3(e+fx) + 4cd \sec^2(e+fx) + 2cd \sec(e+fx) + d^2 \sec^4(e+fx) + 2d^2 \sec^3(e+fx) + d^2 \sec^2(e+fx)}{a^2} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e))**2,x)`

output `Integral(sec(e + f*x)/(c**2*sec(e + f*x)**2 + 2*c**2*sec(e + f*x) + c**2 + 2*c*d*sec(e + f*x)**3 + 4*c*d*sec(e + f*x)**2 + 2*c*d*sec(e + f*x) + d**2 *sec(e + f*x)**4 + 2*d**2*sec(e + f*x)**3 + d**2*sec(e + f*x)**2), x)/a**2`

3.223.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.223.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 474 vs. $2(196) = 392$.

Time = 0.33 (sec) , antiderivative size = 474, normalized size of antiderivative = 2.25

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^2} dx$$

$$= \frac{12 d^3 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{(a^2c^4 - 2a^2c^3d + 2a^2cd^3 - a^2d^4) \left(c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - c - d\right)} + \frac{12(3cd^2 + 2d^3) \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(-2c+2d) + \arctan\left(-\frac{c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - c - d}\right)\right)}{(a^2c^4 - 2a^2c^3d + 2a^2cd^3 - a^2d^4) \sqrt{-c^2 - d^2}}$$

3.223. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^2} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output `1/6*(12*d^3*tan(1/2*f*x + 1/2*e)/((a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)) + 12*(3*c*d^2 + 2*d^3)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((a^2*c^4 - 2*a^2*c^3*d + 2*a^2*c*d^3 - a^2*d^4)*sqrt(-c^2 + d^2)) - (a^4*c^4*tan(1/2*f*x + 1/2*e)^3 - 4*a^4*c^3*d*tan(1/2*f*x + 1/2*e)^3 + 6*a^4*c^2*d^2*tan(1/2*f*x + 1/2*e)^3 - 4*a^4*c*d^3*tan(1/2*f*x + 1/2*e)^3 + a^4*d^4*tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^4*tan(1/2*f*x + 1/2*e) + 24*a^4*c^3*d*tan(1/2*f*x + 1/2*e) - 54*a^4*c^2*d^2*tan(1/2*f*x + 1/2*e) + 48*a^4*c*d^3*tan(1/2*f*x + 1/2*e) - 15*a^4*d^4*tan(1/2*f*x + 1/2*e))/(a^6*c^6 - 6*a^6*c^5*d + 15*a^6*c^4*d^2 - 20*a^6*c^3*d^3 + 15*a^6*c^2*d^4 - 6*a^6*c*d^5 + a^6*d^6))/f`

3.223.9 Mupad [B] (verification not implemented)

Time = 14.29 (sec) , antiderivative size = 314, normalized size of antiderivative = 1.49

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^2} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{3}{2a^2(c-d)^2} - \frac{c^2-d^2}{a^2(c-d)^4}\right)}{f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{6a^2 f (c-d)^2}$$

$$+ \frac{2d^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f(c+d) \left(a^2 d^4 - a^2 c^4 + \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (a^2 c^4 - 4a^2 c^3 d + 6a^2 c^2 d^2 - 4a^2 c d^3 + a^2 d^4) - 2a^2 c d^3 + 2d^2 \operatorname{atan}\left(\frac{1 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^4 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^3 d + 6 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^2 d^2 - 4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c d^3 + 1 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) d^4}{\sqrt{c+d}(c-d)^{7/2}}\right) (3c + 2d) 2i}$$

$$- \frac{a^2 f (c+d)^{3/2} (c-d)^{7/2}}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c + d/cos(e + f*x))^2),x)`

output $(\tan(e/2 + (f*x)/2)*(3/(2*a^2*(c - d)^2) - (c^2 - d^2)/(a^2*(c - d)^4)))/f$
 $- \tan(e/2 + (f*x)/2)^3/(6*a^2*f*(c - d)^2) + (2*d^3*\tan(e/2 + (f*x)/2))/($
 $f*(c + d)*(a^2*d^4 - a^2*c^4 + \tan(e/2 + (f*x)/2)^2*(a^2*c^4 + a^2*d^4 - 4$
 $*a^2*c*d^3 - 4*a^2*c^3*d + 6*a^2*c^2*d^2) - 2*a^2*c*d^3 + 2*a^2*c^3*d) -$
 $(d^2*atan((c^4*\tan(e/2 + (f*x)/2)*1i + d^4*\tan(e/2 + (f*x)/2)*1i - c*d^3*t$
 $an(e/2 + (f*x)/2)*4i - c^3*d*\tan(e/2 + (f*x)/2)*4i + c^2*d^2*\tan(e/2 + (f*$
 $x)/2)*6i)/((c + d)^(1/2)*(c - d)^(7/2)))*(3*c + 2*d)*2i)/(a^2*f*(c + d)^(3$
 $/2)*(c - d)^(7/2))$

3.223. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2} dx$

3.224
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^3} dx$$

3.224.1 Optimal result 1609
 3.224.2 Mathematica [C] (warning: unable to verify) 1610
 3.224.3 Rubi [A] (verified) 1610
 3.224.4 Maple [A] (verified) 1616
 3.224.5 Fricas [B] (verification not implemented) 1616
 3.224.6 Sympy [F] 1617
 3.224.7 Maxima [F(-2)] 1618
 3.224.8 Giac [B] (verification not implemented) 1618
 3.224.9 Mupad [B] (verification not implemented) 1619

3.224.1 Optimal result

Integrand size = 31, antiderivative size = 284

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^3} dx$$

$$= \frac{d^2(12c^2 + 16cd + 7d^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{a^2(c-d)^{9/2}(c+d)^{5/2}f}$$

$$+ \frac{d(2c^2 - 16cd - 21d^2) \tan(e+fx)}{6a^2(c-d)^3(c+d)f(c+d \sec(e+fx))^2}$$

$$+ \frac{(c-8d) \tan(e+fx)}{3a^2(c-d)^2f(1+\sec(e+fx))(c+d \sec(e+fx))^2}$$

$$+ \frac{\tan(e+fx)}{3(c-d)f(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2}$$

$$+ \frac{d(2c^3 - 16c^2d - 59cd^2 - 32d^3) \tan(e+fx)}{6a^2(c-d)^4(c+d)^2f(c+d \sec(e+fx))}$$

output `d^2*(12*c^2+16*c*d+7*d^2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a^2/(c-d)^(9/2)/(c+d)^(5/2)/f+1/6*d*(2*c^2-16*c*d-21*d^2)*tan(f*x+e)/a^2/(c-d)^3/(c+d)/f/(c+d*sec(f*x+e))^2+1/3*(c-8*d)*tan(f*x+e)/a^2/(c-d)^2/f/(1+sec(f*x+e))/(c+d*sec(f*x+e))^2+1/3*tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2+1/6*d*(2*c^3-16*c^2*d-59*c*d^2-32*d^3)*tan(f*x+e)/a^2/(c-d)^4/(c+d)^2/f/(c+d*sec(f*x+e))`

3.224.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 8.18 (sec) , antiderivative size = 2220, normalized size of antiderivative = 7.82

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^3} dx = \text{Result too large to show}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^3),x]`

output

```
((12*c^2 + 16*c*d + 7*d^2)*Cos[e/2 + (f*x)/2]^4*(d + c*Cos[e + f*x])^3*Sec[e + f*x]^5*((-4*I)*d^2*ArcTan[Sec[(f*x)/2]*(Cos[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]]) - (I*Sin[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]])))*((-I)*d*Sin[(f*x)/2] + I*c*Sin[e + (f*x)/2])*Cos[e]/(Sqrt[c^2 - d^2]*f*Sqrt[Cos[2*e] - I*Sin[2*e]]) - (4*d^2*ArcTan[Sec[(f*x)/2]*(Cos[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]]) - (I*Sin[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]])))*((-I)*d*Sin[(f*x)/2] + I*c*Sin[e + (f*x)/2])*Sin[e]/(Sqrt[c^2 - d^2]*f*Sqrt[Cos[2*e] - I*Sin[2*e]]))/((-c + d)^4*(c + d)^2*(a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^3 + (Cos[e/2 + (f*x)/2]*(d + c*Cos[e + f*x])*Sec[e/2]*Sec[e]*Sec[e + f*x]^5*(-16*c^7*Sin[(f*x)/2] + 14*c^6*d*Sin[(f*x)/2] + 220*c^5*d^2*Sin[(f*x)/2] + 334*c^4*d^3*Sin[(f*x)/2] + 54*c^3*d^4*Sin[(f*x)/2] - 156*c^2*d^5*Sin[(f*x)/2] - 48*c*d^6*Sin[(f*x)/2] + 18*d^7*Sin[(f*x)/2] + 14*c^7*Sin[(3*f*x)/2] - 16*c^6*d*Sin[(3*f*x)/2] - 226*c^5*d^2*Sin[(3*f*x)/2] - 532*c^4*d^3*Sin[(3*f*x)/2] - 583*c^3*d^4*Sin[(3*f*x)/2] - 232*c^2*d^5*Sin[(3*f*x)/2] - 6*c*d^6*Sin[(3*f*x)/2] + 6*d^7*Sin[(3*f*x)/2] - 12*c^7*Sin[e - (f*x)/2] + 20*c^6*d*Sin[e - (f*x)/2] + 236*c^5*d^2*Sin[e - (f*x)/2] + 628*c^4*d^3*Sin[e - (f*x)/2] + 778*c^3*d^4*Sin[e - (f*x)/2] + 420*c^2*d^5*Sin[e - (f*x)/2] + 48*c*d^6*Sin[e - (f*x)/2] - 18*d^7*Sin[e - (f*x)/2] + 12*c^7*Sin[e + (f*x)/2] - 20*c^6*d*Sin[e + (f*x)/2] - 236*c^5*d^2*Sin[e + (f*x)/2] - 460*c^4*d^3*Sin[e + ...
```

3.224.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.48, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4475, 114, 27, 168, 27, 169, 25, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.224. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^3} dx$

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)^2(c+d \sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)^2(c+d \csc(e+fx+\frac{\pi}{2}))^3} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{1}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))^3} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 114

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(2(c+d)-3d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2a^2(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}}{2a^2(c^2-d^2)(a \sec(e+fx)+a)^{3/2}(c+d \sec(e+fx))^2} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{2(c+d)-3d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}}{2a^2(c^2-d^2)(a \sec(e+fx)+a)^{3/2}(c+d \sec(e+fx))^2} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 168

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(2c^2+12dc+7d^2-2d(5c+2d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{a^2(c^2-d^2)} + \frac{d(5c+2d) \sqrt{a-a \sec(e+fx)}}{2(c^2-d^2)(a \sec(e+fx)+a)^{3/2}(c+d \sec(e+fx))} + \frac{1}{2a^2(c^2-d^2)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{2c^2+12dc+7d^2-2d(5c+2d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{c^2-d^2} + \frac{d(5c+2d) \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{3/2}(c+d \sec(e+fx))} + \frac{1}{2a^2(c^2-d^2)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 169

3.224. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2((c+d)(2c^2-16dc-21d^2)+d(2c^2+22dc+11d^2)\sec(e+fx))\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d\sec(e+fx))} d\sec(e+fx) - \frac{(2c^2+22cd+11d^2)\sqrt{a-a\sec(e+fx)}}{3a^2(c-d)(a\sec(e+fx)+a)^{3/2}}}{c^2-d^2} + \frac{d(5c+2d)\sqrt{a-a\sec(e+fx)}}{a^2(c^2-d^2)(a\sec(e+fx)+a)} \right) dx$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

25

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2((c+d)(2c^2-16dc-21d^2)+d(2c^2+22dc+11d^2)\sec(e+fx))\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d\sec(e+fx))} d\sec(e+fx) - \frac{(2c^2+22cd+11d^2)\sqrt{a-a\sec(e+fx)}}{3a^2(c-d)(a\sec(e+fx)+a)^{3/2}}}{c^2-d^2} + \frac{d(5c+2d)\sqrt{a-a\sec(e+fx)}}{a^2(c^2-d^2)(a\sec(e+fx)+a)} \right) dx$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d)(2c^2-16dc-21d^2)+d(2c^2+22dc+11d^2)\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d\sec(e+fx))} d\sec(e+fx) - \frac{(2c^2+22cd+11d^2)\sqrt{a-a\sec(e+fx)}}{3a^2(c-d)(a\sec(e+fx)+a)^{3/2}}}{c^2-d^2} + \frac{d(5c+2d)\sqrt{a-a\sec(e+fx)}}{a^2(c^2-d^2)(a\sec(e+fx)+a)} \right) dx$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

169

$$a^2 \tan(e + fx) \left(\frac{\int \frac{3a^2 d^2 (12c^2+16dc+7d^2)}{\sqrt{a-a\sec(e+fx)}\sqrt{\sec(e+fx)a+a}(c+d\sec(e+fx))} d\sec(e+fx) - \frac{(2c^3-16c^2d-59cd^2-32d^3)\sqrt{a-a\sec(e+fx)}}{a^2(c-d)\sqrt{a\sec(e+fx)+a}}}{3a(c-d)} - \frac{(2c^2+22cd+11d^2)\sqrt{a-a\sec(e+fx)}}{3a^2(c-d)(a\sec(e+fx)+a)^{3/2}}}{c^2-d^2} \right) dx$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

27

3.224. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^2(c+d\sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{3d^2(12c^2+16cd+7d^2) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}} d \sec(e+fx)}{a(c-d)} - \frac{(2c^3-16c^2d-59cd^2-32d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} \right) \frac{3a(c-d)}{c^2-d^2} \frac{1}{2(c^2-d^2)}$$

$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$

104

$$a^2 \tan(e + fx) \left(\frac{6d^2(12c^2+16cd+7d^2) \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{a(c-d)} - \frac{(2c^3-16c^2d-59cd^2-32d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} \right) \frac{3a(c-d)}{c^2-d^2} \frac{1}{2(c^2-d^2)}$$

$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$

218

$$a^2 \tan(e + fx) \left(\frac{6d^2(12c^2+16cd+7d^2) \arctan\left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a \sec(e+fx)}}\right)}{a^2(c-d)^{3/2} \sqrt{c+d}} - \frac{(2c^3-16c^2d-59cd^2-32d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} \right) \frac{3a(c-d)}{c^2-d^2} \frac{1}{2(c^2-d^2)}$$

$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$

```
input Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^2*(c + d*Sec[e + f*x])^3),x]
```

```
output -((a^2*((d*Sqrt[a - a*Sec[e + f*x]])/(2*a^2*(c^2 - d^2)*(a + a*Sec[e + f*x
])^(3/2)*(c + d*Sec[e + f*x])^2) + ((d*(5*c + 2*d)*Sqrt[a - a*Sec[e + f*x]
])/(a^2*(c^2 - d^2)*(a + a*Sec[e + f*x])^(3/2)*(c + d*Sec[e + f*x])) + (-1
/3*((2*c^2 + 22*c*d + 11*d^2)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*(a +
a*Sec[e + f*x])^(3/2)) + ((6*d^2*(12*c^2 + 16*c*d + 7*d^2)*ArcTan[(Sqrt[c
+ d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(a
^2*(c - d)^(3/2)*Sqrt[c + d]) - ((2*c^3 - 16*c^2*d - 59*c*d^2 - 32*d^3)*Sq
rt[a - a*Sec[e + f*x]])/(a^2*(c - d)*Sqrt[a + a*Sec[e + f*x]))/(3*a*(c -
d))/(c^2 - d^2)/(2*(c^2 - d^2))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x
]]*Sqrt[a + a*Sec[e + f*x]))
```

3.224.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 114 Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)
)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e
- a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1)
- b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] ||
IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

rule 168 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]`

rule 169 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.224.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 280, normalized size of antiderivative = 0.99

method	result
derivativedivides	$\frac{-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 7d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^3 - 3c^2d + 3cd^2 - d^3)(c-d)} - \frac{8d^2 \left(-\frac{d(8c^2 - 3cd - 5d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{4(c^2 + 2cd + d^2)} + \frac{d(8c + 3d) \tan\left(\frac{fx}{2}\right)}{4c + 4d} \right)}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}^2} {2fa^2}$
default	$\frac{-\frac{c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - \frac{d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} - c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 7d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^3 - 3c^2d + 3cd^2 - d^3)(c-d)} - \frac{8d^2 \left(-\frac{d(8c^2 - 3cd - 5d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{4(c^2 + 2cd + d^2)} + \frac{d(8c + 3d) \tan\left(\frac{fx}{2}\right)}{4c + 4d} \right)}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d}^2} {2fa^2}$
risch	Expression too large to display

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `1/2/f/a^2*(-1/(c^3-3*c^2*d+3*c*d^2-d^3)/(c-d)*(1/3*c*tan(1/2*f*x+1/2*e)^3-1/3*d*tan(1/2*f*x+1/2*e)^3-c*tan(1/2*f*x+1/2*e)+7*d*tan(1/2*f*x+1/2*e))-8*d^2/(c-d)^4*((-1/4*d*(8*c^2-3*c*d-5*d^2)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3+1/4*d*(8*c+3*d)/(c+d)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^2-1/4*(12*c^2+16*c*d+7*d^2)/(c^2+2*c*d+d^2)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))))`

3.224.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 986 vs. 2(267) = 534.

Time = 0.36 (sec) , antiderivative size = 2030, normalized size of antiderivative = 7.15

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^3} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x, algorithm="fracas")`

output

```
[1/12*(3*(12*c^2*d^4 + 16*c*d^5 + 7*d^6 + (12*c^4*d^2 + 16*c^3*d^3 + 7*c^2*d^4)*cos(f*x + e)^4 + 2*(12*c^4*d^2 + 28*c^3*d^3 + 23*c^2*d^4 + 7*c*d^5)*cos(f*x + e)^3 + (12*c^4*d^2 + 64*c^3*d^3 + 83*c^2*d^4 + 44*c*d^5 + 7*d^6)*cos(f*x + e)^2 + 2*(12*c^3*d^3 + 28*c^2*d^4 + 23*c*d^5 + 7*d^6)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(2*c^5*d^2 - 16*c^4*d^3 - 61*c^3*d^4 - 16*c^2*d^5 + 59*c*d^6 + 32*d^7 + (4*c^7 - 14*c^6*d - 44*c^5*d^2 - 32*c^4*d^3 + 28*c^3*d^4 + 49*c^2*d^5 + 12*c*d^6 - 3*d^7)*cos(f*x + e)^3 + (2*c^7 - 8*c^6*d - 68*c^5*d^2 - 140*c^4*d^3 - 23*c^3*d^4 + 142*c^2*d^5 + 89*c*d^6 + 6*d^7)*cos(f*x + e)^2 + (4*c^6*d - 28*c^5*d^2 - 118*c^4*d^3 - 106*c^3*d^4 + 71*c^2*d^5 + 134*c*d^6 + 43*d^7)*cos(f*x + e))*sin(f*x + e))/((a^2*c^10 - 2*a^2*c^9*d - 2*a^2*c^8*d^2 + 6*a^2*c^7*d^3 - 6*a^2*c^5*d^5 + 2*a^2*c^4*d^6 + 2*a^2*c^3*d^7 - a^2*c^2*d^8)*f*cos(f*x + e)^4 + 2*(a^2*c^10 - a^2*c^9*d - 4*a^2*c^8*d^2 + 4*a^2*c^7*d^3 + 6*a^2*c^6*d^4 - 6*a^2*c^5*d^5 - 4*a^2*c^4*d^6 + 4*a^2*c^3*d^7 + a^2*c^2*d^8 - a^2*c*d^9)*f*cos(f*x + e)^3 + (a^2*c^10 + 2*a^2*c^9*d - 9*a^2*c^8*d^2 - 4*a^2*c^7*d^3 + 22*a^2*c^6*d^4 - 22*a^2*c^4*d^6 + 4*a^2*c^3*d^7 + 9*a^2*c^2*d^8 - 2*a^2*c*d^9 - a^2*d^10)*f*cos(f*x + e)^2 + 2*(a^2*c^9*d - a^2*c^8*d^2 - 4*a^2*c^7*d^3 + 4*a^2*c^6*d^4 + 6*a^2*c^5*d^5 - 6*a^2*c^4*d^6 - 4*a^2*c^3*d^7 + 4*...
```

3.224.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^3} dx$$

$$= \frac{\int \frac{\sec(e+fx)}{c^3 \sec^2(e+fx) + 2c^3 \sec(e+fx) + c^3 + 3c^2 d \sec^3(e+fx) + 6c^2 d \sec^2(e+fx) + 3c^2 d \sec(e+fx) + 3cd^2 \sec^4(e+fx) + 6cd^2 \sec^3(e+fx) + 3cd^2 \sec^2(e+fx) + 3cd^2 \sec(e+fx) + d^3} dx}{a^2}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**2/(c+d*sec(f*x+e))**3,x)`

output

```
Integral(sec(e + f*x)/(c**3*sec(e + f*x)**2 + 2*c**3*sec(e + f*x) + c**3 + 3*c**2*d*sec(e + f*x)**3 + 6*c**2*d*sec(e + f*x)**2 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**4 + 6*c*d**2*sec(e + f*x)**3 + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**5 + 2*d**3*sec(e + f*x)**4 + d**3*sec(e + f*x)**3), x)/a**2
```

3.224. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2 (c+d \sec(e+fx))^3} dx$

3.224.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.224.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 751 vs. 2(267) = 534.

Time = 0.40 (sec) , antiderivative size = 751, normalized size of antiderivative = 2.64

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^3} dx$$

$$= \frac{6(12c^2d^2 + 16cd^3 + 7d^4) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2c+2d) + \arctan \left(-\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right)}{(a^2c^6 - 2a^2c^5d - a^2c^4d^2 + 4a^2c^3d^3 - a^2c^2d^4 - 2a^2cd^5 + a^2d^6)\sqrt{-c^2+d^2}} - \frac{a^4c^6 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 6a^4c^5d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output

```

1/6*(6*(12*c^2*d^2 + 16*c*d^3 + 7*d^4)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((a^2*c^6 - 2*a^2*c^5*d - a^2*c^4*d^2 + 4*a^2*c^3*d^3 - a^2*c^2*d^4 - 2*a^2*c*d^5 + a^2*d^6)*sqrt(-c^2 + d^2)) - (a^4*c^6*tan(1/2*f*x + 1/2*e)^3 - 6*a^4*c^5*d*tan(1/2*f*x + 1/2*e)^3 + 15*a^4*c^4*d^2*tan(1/2*f*x + 1/2*e)^3 - 20*a^4*c^3*d^3*tan(1/2*f*x + 1/2*e)^3 + 15*a^4*c^2*d^4*tan(1/2*f*x + 1/2*e)^3 - 6*a^4*c*d^5*tan(1/2*f*x + 1/2*e)^3 + a^4*d^6*tan(1/2*f*x + 1/2*e)^3 - 3*a^4*c^6*tan(1/2*f*x + 1/2*e) + 36*a^4*c^5*d*tan(1/2*f*x + 1/2*e) - 135*a^4*c^4*d^2*tan(1/2*f*x + 1/2*e) + 240*a^4*c^3*d^3*tan(1/2*f*x + 1/2*e) - 225*a^4*c^2*d^4*tan(1/2*f*x + 1/2*e) + 108*a^4*c*d^5*tan(1/2*f*x + 1/2*e) - 21*a^4*d^6*tan(1/2*f*x + 1/2*e))/(a^6*c^9 - 9*a^6*c^8*d + 36*a^6*c^7*d^2 - 84*a^6*c^6*d^3 + 126*a^6*c^5*d^4 - 126*a^6*c^4*d^5 + 84*a^6*c^3*d^6 - 36*a^6*c^2*d^7 + 9*a^6*c*d^8 - a^6*d^9) + 6*(8*c^2*d^3*tan(1/2*f*x + 1/2*e)^3 - 3*c*d^4*tan(1/2*f*x + 1/2*e)^3 - 5*d^5*tan(1/2*f*x + 1/2*e)^3 - 8*c^2*d^3*tan(1/2*f*x + 1/2*e) - 11*c*d^4*tan(1/2*f*x + 1/2*e) - 3*d^5*tan(1/2*f*x + 1/2*e))/((a^2*c^6 - 2*a^2*c^5*d - a^2*c^4*d^2 + 4*a^2*c^3*d^3 - a^2*c^2*d^4 - 2*a^2*c*d^5 + a^2*d^6)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^2))/f

```

3.224.9 Mupad [B] (verification not implemented)

Time = 14.20 (sec) , antiderivative size = 505, normalized size of antiderivative = 1.78

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^3} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)} \frac{\sec(e + fx)}{(a + a \sec(e + fx))^2 (c + d \sec(e + fx))^3} dx$$

$$= \frac{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (2a^2 c^6 - 8a^2 c^5 d + 10a^2 c^4 d^2 - 10a^2 c^2 d^4 + 8a^2 c d^5 - 2a^2 d^6) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 (a^2 c^6 + 6a^2 c^5 d + 15a^2 c^4 d^2 + 20a^2 c^3 d^3 + 15a^2 c^2 d^4 + 6a^2 c d^5 + d^6) \right)}{a^2 f (c + d)^3} + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{2}{a^2 (c-d)^3} - \frac{3(c+d)}{2a^2 (c-d)^4} \right)}{f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3}{6a^2 f (c-d)^3}$$

$$- \frac{d^2 \operatorname{atan}\left(\frac{1i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^5 - 5i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^4 d + 10i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^3 d^2 - 10i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^2 d^3 + 5i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c d^4 - 1i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) d^5}{\sqrt{c+d}(c-d)^{9/2}}\right)}{a^2 f (c+d)^{5/2} (c-d)^{9/2}}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^2*(c + d/cos(e + f*x))^3),x)`

$$3.224. \int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^3} dx$$

output

$$\begin{aligned} & ((\tan(e/2 + (f*x)/2)^3*(3*c*d^4 + 5*d^5 - 8*c^2*d^3))/(c + d)^2 + (\tan(e/2 \\ & + (f*x)/2)*(8*c*d^3 + 3*d^4))/(c + d))/(f*(\tan(e/2 + (f*x)/2)^2*(2*a^2*c^6 \\ & - 2*a^2*d^6 + 8*a^2*c*d^5 - 8*a^2*c^5*d - 10*a^2*c^2*d^4 + 10*a^2*c^4*d^2) \\ & - \tan(e/2 + (f*x)/2)^4*(a^2*c^6 + a^2*d^6 - 6*a^2*c*d^5 - 6*a^2*c^5*d + \\ & 15*a^2*c^2*d^4 - 20*a^2*c^3*d^3 + 15*a^2*c^4*d^2) - a^2*c^6 - a^2*d^6 + 2 \\ & *a^2*c*d^5 + 2*a^2*c^5*d + a^2*c^2*d^4 - 4*a^2*c^3*d^3 + a^2*c^4*d^2)) + (\\ & \tan(e/2 + (f*x)/2)*(2/(a^2*(c - d)^3) - (3*(c + d))/(2*a^2*(c - d)^4))/f \\ & - \tan(e/2 + (f*x)/2)^3/(6*a^2*f*(c - d)^3) - (d^2*atan((c^5*\tan(e/2 + (f*x) \\ &)/2)*1i - d^5*\tan(e/2 + (f*x)/2)*1i + c*d^4*\tan(e/2 + (f*x)/2)*5i - c^4*d* \\ & \tan(e/2 + (f*x)/2)*5i - c^2*d^3*\tan(e/2 + (f*x)/2)*10i + c^3*d^2*\tan(e/2 + \\ & (f*x)/2)*10i)/((c + d)^(1/2)*(c - d)^(9/2)))*(16*c*d + 12*c^2 + 7*d^2)*1i \\ &)/(a^2*f*(c + d)^(5/2)*(c - d)^(9/2)) \end{aligned}$$

3.224. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^2(c+d \sec(e+fx))^3} dx$

3.225 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$

3.225.1 Optimal result 1621
 3.225.2 Mathematica [B] (verified) 1622
 3.225.3 Rubi [A] (verified) 1623
 3.225.4 Maple [A] (verified) 1628
 3.225.5 Fricas [A] (verification not implemented) 1629
 3.225.6 Sympy [F] 1630
 3.225.7 Maxima [B] (verification not implemented) 1631
 3.225.8 Giac [A] (verification not implemented) 1631
 3.225.9 Mupad [B] (verification not implemented) 1632

3.225.1 Optimal result

Integrand size = 31, antiderivative size = 363

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx \\ &= \frac{d^3(40c^3 - 90c^2d + 78cd^2 - 23d^3) \operatorname{arctanh}(\sin(e+fx))}{2a^3 f} \\ & \quad - \frac{2d(2c^5 + 18c^4d + 107c^3d^2 - 472c^2d^3 + 456cd^4 - 136d^5) \tan(e+fx)}{15a^3 f} \\ & \quad - \frac{d^2(4c^4 + 36c^3d + 216c^2d^2 - 626cd^3 + 345d^4) \sec(e+fx) \tan(e+fx)}{30a^3 f} \\ & \quad - \frac{d(2c^3 + 18c^2d + 111cd^2 - 136d^3) (c+d \sec(e+fx))^2 \tan(e+fx)}{15a^3 f} \\ & \quad + \frac{(c-d)(2c^2 + 18cd + 115d^2) (c+d \sec(e+fx))^3 \tan(e+fx)}{15f(a^3 + a^3 \sec(e+fx))} \\ & \quad + \frac{(c-d)(2c + 13d)(c+d \sec(e+fx))^4 \tan(e+fx)}{15af(a+a \sec(e+fx))^2} \\ & \quad + \frac{(c-d)(c+d \sec(e+fx))^5 \tan(e+fx)}{5f(a+a \sec(e+fx))^3} \end{aligned}$$

output $\frac{1}{2}d^3(40c^3-90c^2d+78cd^2-23d^3)\operatorname{arctanh}(\sin(fx+e))/a^3/f-2/15d$
 $\cdot(2c^5+18c^4d+107c^3d^2-472c^2d^3+456cd^4-136d^5)\tan(fx+e)/a^3$
 $/f-1/30d^2(4c^4+36c^3d+216c^2d^2-626cd^3+345d^4)\sec(fx+e)\tan$
 $(fx+e)/a^3/f-1/15d(2c^3+18c^2d+111cd^2-136d^3)(c+d\sec(fx+e))^2$
 $\tan(fx+e)/a^3/f+1/15(c-d)(2c^2+18cd+115d^2)(c+d\sec(fx+e))^3\tan$
 $(fx+e)/f/(a^3+a^3\sec(fx+e))+1/15(c-d)(2c+13d)(c+d\sec(fx+e))^4\tan$
 $(fx+e)/a/f/(a+a\sec(fx+e))^2+1/5(c-d)(c+d\sec(fx+e))^5\tan(fx+e)/f/$
 $(a+a\sec(fx+e))^3$

3.225.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1338 vs. $2(363) = 726$.

Time = 9.81 (sec) , antiderivative size = 1338, normalized size of antiderivative = 3.69

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx = \text{Too large to display}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^6)/(a + a*Sec[e + f*x])^3,x]`

output $(4*(-40c^3d^3 + 90c^2d^4 - 78cd^5 + 23d^6)\operatorname{Cos}[e/2 + (fx)/2]^6\operatorname{Cos}$
 $[e + fx]^3\operatorname{Log}[\operatorname{Cos}[e/2 + (fx)/2] - \operatorname{Sin}[e/2 + (fx)/2]](c + d\sec[e + f$
 $x])^6)/(f*(d + c\operatorname{Cos}[e + fx])^6(a + a\sec[e + f*x])^3) - (4*(-40c^3d^3$
 $+ 90c^2d^4 - 78cd^5 + 23d^6)\operatorname{Cos}[e/2 + (fx)/2]^6\operatorname{Cos}[e + fx]^3\operatorname{Log}$
 $[\operatorname{Cos}[e/2 + (fx)/2] + \operatorname{Sin}[e/2 + (fx)/2]](c + d\sec[e + f*x])^6)/(f*(d +$
 $c\operatorname{Cos}[e + fx])^6(a + a\sec[e + f*x])^3) + (2\operatorname{Cos}[e/2 + (fx)/2]^2\operatorname{Cos}[e$
 $+ fx]^3\sec[e/2](c + d\sec[e + f*x])^6(c^6\operatorname{Sin}[e/2] - 6c^5d\operatorname{Sin}[e/2]$
 $+ 15c^4d^2\operatorname{Sin}[e/2] - 20c^3d^3\operatorname{Sin}[e/2] + 15c^2d^4\operatorname{Sin}[e/2] - 6cd^5$
 $5\operatorname{Sin}[e/2] + d^6\operatorname{Sin}[e/2]))/(5f*(d + c\operatorname{Cos}[e + fx])^6(a + a\sec[e + f*x]$
 $)^3) + (8\operatorname{Cos}[e/2 + (fx)/2]^4\operatorname{Cos}[e + fx]^3\sec[e/2](c + d\sec[e + f*x]$
 $)^6(-4c^6\operatorname{Sin}[e/2] + 9c^5d\operatorname{Sin}[e/2] + 15c^4d^2\operatorname{Sin}[e/2] - 70c^3d^3$
 $3\operatorname{Sin}[e/2] + 90c^2d^4\operatorname{Sin}[e/2] - 51cd^5\operatorname{Sin}[e/2] + 11d^6\operatorname{Sin}[e/2]))/($
 $15f*(d + c\operatorname{Cos}[e + fx])^6(a + a\sec[e + f*x])^3) + (2\operatorname{Cos}[e/2 + (fx)/2]$
 $]\operatorname{Cos}[e + fx]^3\sec[e/2](c + d\sec[e + f*x])^6(c^6\operatorname{Sin}[(fx)/2] - 6c^5$
 $d\operatorname{Sin}[(fx)/2] + 15c^4d^2\operatorname{Sin}[(fx)/2] - 20c^3d^3\operatorname{Sin}[(fx)/2] + 15c$
 $^2d^4\operatorname{Sin}[(fx)/2] - 6cd^5\operatorname{Sin}[(fx)/2] + d^6\operatorname{Sin}[(fx)/2]))/(5f*(d +$
 $c\operatorname{Cos}[e + fx])^6(a + a\sec[e + f*x])^3) + (8\operatorname{Cos}[e/2 + (fx)/2]^3\operatorname{Cos}[e$
 $+ fx]^3\sec[e/2](c + d\sec[e + f*x])^6(-4c^6\operatorname{Sin}[(fx)/2] + 9c^5d\operatorname{Si}$
 $n[(fx)/2] + 15c^4d^2\operatorname{Sin}[(fx)/2] - 70c^3d^3\operatorname{Sin}[(fx)/2] + 90c^2d^$
 $4\operatorname{Sin}[(fx)/2] - 51cd^5\operatorname{Sin}[(fx)/2] + 11d^6\operatorname{Sin}[(fx)/2]))/(15f*(d...$

3.225.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 501, normalized size of antiderivative = 1.38, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4475, 109, 25, 27, 167, 27, 167, 27, 170, 25, 27, 164, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a\sec(e+fx)+a)^3} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)^6}{\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^3} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{(c+d\sec(e+fx))^6}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{7/2}} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 109

$$\frac{a^2 \tan(e+fx) \left(-\frac{\int \frac{a^2(c+d\sec(e+fx))^4(2c^2+8dc-5d^2-(3c-8d)d\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d\sec(e+fx)}{5a^3} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^5}{5a^2(a\sec(e+fx)+a)^{5/2}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(c+d\sec(e+fx))^4(2c^2+8dc-5d^2-(3c-8d)d\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d\sec(e+fx)}{5a^3} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^5}{5a^2(a\sec(e+fx)+a)^{5/2}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{(c+d\sec(e+fx))^4(2c^2+8dc-5d^2-(3c-8d)d\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d\sec(e+fx)}{5a} - \frac{(c-d)\sqrt{a-a\sec(e+fx)}(c+d\sec(e+fx))^5}{5a^2(a\sec(e+fx)+a)^{5/2}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 167

3.225. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))^3 (2c^3+10dc^2+55d^2c-52d^3-3d(2c^2+14dc-21d^2) \sec(e+fx)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)(2c+13d)\sqrt{a-a \sec(e+fx)}(c+d)}{3a^2(a \sec(e+fx)+a)^{3/2}}}{5a} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d \sec(e+fx))^3 (2c^3+10dc^2+55d^2c-52d^3-3d(2c^2+14dc-21d^2) \sec(e+fx)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a} - \frac{(c-d)(2c+13d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))}{3a^2(a \sec(e+fx)+a)^{3/2}}}{5a} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 167

$$a^2 \tan(e + fx) \left(\frac{\int \frac{3a^2 d(c+d \sec(e+fx))^2 (d(2c^2+118dc-115d^2) - (2c^3+18dc^2+111d^2c-136d^3) \sec(e+fx)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{a^3} - \frac{(c-d)(2c^2+18cd+115d^2)\sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)+a}}}{3a} - \frac{(c-d)(2c^2+18cd+115d^2)\sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)+a}}}{5a} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{3d \int \frac{(c+d \sec(e+fx))^2 (d(2c^2+118dc-115d^2) - (2c^3+18dc^2+111d^2c-136d^3) \sec(e+fx)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{a} - \frac{(c-d)(2c^2+18cd+115d^2)\sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)+a}}}{3a} - \frac{(c-d)(2c^2+18cd+115d^2)\sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)+a}}}{5a} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 170

3.225. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{(2c^3 + 18c^2d + 111cd^2 - 136d^3) \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (c + d \sec(e + fx))^2}{3a^2} \int - \frac{a^2 (c + d \sec(e + fx)) (d(2c^3 + 318dc^2 - 567d^2c + 272d^3) - (4c^4 + 36dc^3 + 216d^2c^2 - 626d^3c + 345d^4) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}} \right)}{a} \right)$$

↓ 25

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{\int \frac{a^2 (c + d \sec(e + fx)) (d(2c^3 + 318dc^2 - 567d^2c + 272d^3) - (4c^4 + 36dc^3 + 216d^2c^2 - 626d^3c + 345d^4) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}} d \sec(e + fx) + \frac{(2c^3 + 18c^2d + 111cd^2 - 136d^3) \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (c + d \sec(e + fx))^2}{3a^2}}{a} \right)$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \int \frac{(c + d \sec(e + fx)) (d(2c^3 + 318dc^2 - 567d^2c + 272d^3) - (4c^4 + 36dc^3 + 216d^2c^2 - 626d^3c + 345d^4) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}} d \sec(e + fx) + \frac{(2c^3 + 18c^2d + 111cd^2 - 136d^3) \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (c + d \sec(e + fx))^2}{3a^2}}{a} \right)$$

↓ 164

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \left(\frac{15}{2} d^2 (40c^3 - 90c^2d + 78cd^2 - 23d^3) \int \frac{1}{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}} d \sec(e + fx) + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(4c^4 + 36dc^3 + 216d^2c^2 - 626d^3c + 345d^4) \sec(e + fx))}{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}} \right)}{a} \right)$$

↓ 45

3.225. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \left(15d^2 (40c^3 - 90c^2d + 78cd^2 - 23d^3) \int \frac{1}{\frac{(a - a \sec(e + fx))a}{\sec(e + fx)a + a} - a} d \frac{\sqrt{a - a \sec(e + fx)}}{\sqrt{\sec(e + fx)a + a}} + \frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}{d(4c^4 + 36c^3d + 216c^2d^2 - 626cd^3 + 345d^4)} \right) \right)}{\dots} \right)$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{3d \left(\frac{1}{3} \left(\frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(4c^4 + 36c^3d + 216c^2d^2 - 626cd^3 + 345d^4) \sec(e + fx) + 4(2c^5 + 18c^4d + 107c^3d^2 - 472c^2d^3 + 456cd^4 - 136d^5) + d(4c^4 + 36c^3d + 216c^2d^2 - 626cd^3 + 345d^4))}{2a^2} \right) \right)}{\dots} \right)$$

```
input Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^6)/(a + a*Sec[e + f*x])^3,x]
```

```
output -((a^2*(-1/5*((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^5)/(a^2*(a + a*Sec[e + f*x])^(5/2)) + (-1/3*((c - d)*(2*c + 13*d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^4)/(a^2*(a + a*Sec[e + f*x])^(3/2)) + (-((c - d)*(2*c^2 + 18*c*d + 115*d^2)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^3)/(a^2*Sqrt[a + a*Sec[e + f*x]])) + (3*d*((2*c^3 + 18*c^2*d + 111*c*d^2 - 136*d^3)*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(3*a^2) + ((-15*d^2*(40*c^3 - 90*c^2*d + 78*c*d^2 - 23*d^3)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])]/a + (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(4*(2*c^5 + 18*c^4*d + 107*c^3*d^2 - 472*c^2*d^3 + 456*c*d^4 - 136*d^5) + d*(4*c^4 + 36*c^3*d + 216*c^2*d^2 - 626*c*d^3 + 345*d^4)*Sec[e + f*x]))/(2*a^2))/3)/a)/(3*a))/(5*a))*Tan[e + f*x))/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.225.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 109 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 164 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`
- rule 167 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] - Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[b*c*(f*g - e*h)*(m + 1) + (b*g - a*h)*(d*e*n + c*f*(p + 1)) + d*(b*(f*g - e*h)*(m + 1) + f*(b*g - a*h)*(n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, p}, x] && LtQ[m, -1] && GtQ[n, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 170 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_))*((g_.) + (h_.)*(x_)), x_] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Simp[1/(d*f*(m + n + p + 2)) Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_))*((csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.)^(n_)), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.225.4 Maple [A] (verified)

Time = 1.47 (sec) , antiderivative size = 491, normalized size of antiderivative = 1.35

method	result
parallelrisch	$-14400(c^3 - \frac{9}{4}c^2d + \frac{39}{20}cd^2 - \frac{23}{40}d^3) \left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3} \right) d^3 \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) + 14400(c^3 - \frac{9}{4}c^2d + \frac{39}{20}cd^2 - \frac{23}{40}d^3)$
derivativedivides	$-\frac{4d^6}{3(\tan(\frac{fx}{2} + \frac{e}{2}) - 1)^3} + c^6 \tan(\frac{fx}{2} + \frac{e}{2}) + 49d^6 \tan(\frac{fx}{2} + \frac{e}{2}) - \frac{4d^6}{3(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)^3} + \frac{d^6 \tan(\frac{fx}{2} + \frac{e}{2})^5}{5} - \frac{2c^6 \tan(\frac{fx}{2} + \frac{e}{2})^3}{3} + 10$
default	$-\frac{4d^6}{3(\tan(\frac{fx}{2} + \frac{e}{2}) - 1)^3} + c^6 \tan(\frac{fx}{2} + \frac{e}{2}) + 49d^6 \tan(\frac{fx}{2} + \frac{e}{2}) - \frac{4d^6}{3(\tan(\frac{fx}{2} + \frac{e}{2}) + 1)^3} + \frac{d^6 \tan(\frac{fx}{2} + \frac{e}{2})^5}{5} - \frac{2c^6 \tan(\frac{fx}{2} + \frac{e}{2})^3}{3} + 10$
risch	Expression too large to display

3.225.
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & \frac{1}{240}(-14400(c^3-9/4c^2d+39/20cd^2-23/40d^3)(\cos(fx+e)+1/3\cos(3fx+3e))*d^3\ln(\tan(1/2fx+1/2e))-1)+14400(c^3-9/4c^2d+39/20cd^2-23/40d^3)(\cos(fx+e)+1/3\cos(3fx+3e))*d^3\ln(\tan(1/2fx+1/2e)+1)+12\tan(1/2fx+1/2e)*((43/12c^6+1549/6d^6-859cd^5+95/2c^4d^2-1190/3c^3d^3+1035c^2d^4+27/2c^5d)\cos(3fx+3e)+(36c^5d+4c^6+1382/3d^6-1524cd^5+60c^4d^2-680c^3d^3+1860c^2d^4)\cos(2fx+2e)+(c^6+429/4d^6+9c^5d+15c^4d^2-170c^3d^3+855/2c^2d^4-717/2cd^5)\cos(4fx+4e)+(7/12c^6+68/3d^6-76cd^5+3/2c^5d+5/2c^4d^2-110/3c^3d^3+90c^2d^4)\cos(5fx+5e)+(3907/6d^6+33c^5d+47/6c^6-3020/3c^3d^3+2655c^2d^4-2137cd^5+130c^4d^2)\cos(fx+e)+4321/12d^6+3c^6+27c^5d+45c^4d^2-510c^3d^3+2865/2c^2d^4-2331/2cd^5)*\sec(1/2fx+1/2e)^4)/f/a^3/(\cos(3fx+3e)+3\cos(fx+e)) \end{aligned}$$

3.225.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 620, normalized size of antiderivative = 1.71

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{15((40c^3d^3-90c^2d^4+78cd^5-23d^6)\cos(fx+e)^6+3(40c^3d^3-90c^2d^4+78cd^5-23d^6)\cos(fx+e))}{f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x, algorithm="fracas")`

output $\frac{1}{60} \cdot (15 \cdot ((40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^6 + 3 \cdot (40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^5 + 3 \cdot (40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^4 + (40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^3) \cdot \log(\sin(fx + e) + 1) - 15 \cdot ((40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^6 + 3 \cdot (40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^5 + 3 \cdot (40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^4 + (40c^3d^3 - 90c^2d^4 + 78cd^5 - 23d^6) \cos(fx + e)^3) \cdot \log(-\sin(fx + e) + 1) + 2 \cdot (10d^6 + 2 \cdot (7c^6 + 18c^5d + 30c^4d^2 - 440c^3d^3 + 1080c^2d^4 - 912cd^5 + 272d^6) \cos(fx + e)^5 + 3 \cdot (4c^6 + 36c^5d + 60c^4d^2 - 680c^3d^3 + 1710c^2d^4 - 1434cd^5 + 429d^6) \cos(fx + e)^4 + (4c^6 + 36c^5d + 210c^4d^2 - 1280c^3d^3 + 3510c^2d^4 - 2874cd^5 + 869d^6) \cos(fx + e)^3 + 5 \cdot (90c^2d^4 - 54cd^5 + 19d^6) \cos(fx + e)^2 + 15 \cdot (6cd^5 - d^6) \cos(fx + e) \cdot \sin(fx + e)) / (a^3 f \cos(fx + e)^6 + 3a^3 f \cos(fx + e)^5 + 3a^3 f \cos(fx + e)^4 + a^3 f \cos(fx + e)^3)$

3.225.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^6}{(a + a \sec(e + fx))^3} dx$$

$$= \int \frac{c^6 \sec(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx + \int \frac{d^6 \sec^7(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx + \int \frac{6cd^5 \sec^6(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**6/(a+a*sec(f*x+e))**3,x)`

output `(Integral(c**6*sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(d**6*sec(e + f*x)**7/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(6*c*d**5*sec(e + f*x)**6/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(15*c**2*d**4*sec(e + f*x)**5/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(20*c**3*d**3*sec(e + f*x)**4/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(15*c**4*d**2*sec(e + f*x)**3/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(6*c**5*d*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3`

3.225. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^6}{(a+a \sec(e+fx))^3} dx$

3.225.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 946 vs. $2(349) = 698$.

Time = 0.25 (sec) , antiderivative size = 946, normalized size of antiderivative = 2.61

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/60*(d^6*(20*(33*sin(f*x + e)/(cos(f*x + e) + 1) - 76*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 51*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/(a^3 - 3*a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 3*a^3*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - a^3*sin(f*x + e)^6/(cos(f*x + e) + 1)^6) + (735*sin(f*x + e)/(cos(f*x + e) + 1) + 50*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 690*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 690*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3 - 6*c*d^5*(60*(5*sin(f*x + e)/(cos(f*x + e) + 1) - 7*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^3 - 2*a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^3*sin(f*x + e)^4/(cos(f*x + e) + 1)^4) + (465*sin(f*x + e)/(cos(f*x + e) + 1) + 40*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 390*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 390*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3 + 45*c^2*d^4*(40*sin(f*x + e)/((a^3 - a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1)) + (85*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3 - 20*c^3*d^3*((105*sin(f*x + e)/(cos(f*x + e) + 1) + 20*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) + 15*c^4*d...`

3.225.8 Giac [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 672, normalized size of antiderivative = 1.85

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$$

$$\frac{30(40c^3d^3-90c^2d^4+78cd^5-23d^6)\log\left(\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)+1\right|\right)}{a^3} - \frac{30(40c^3d^3-90c^2d^4+78cd^5-23d^6)\log\left(\left|\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-1\right|\right)}{a^3} - \frac{20(90c^2d^4}{a^3}$$

=

3.225. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^6/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output `1/60*(30*(40*c^3*d^3 - 90*c^2*d^4 + 78*c*d^5 - 23*d^6)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) / a^3 - 30*(40*c^3*d^3 - 90*c^2*d^4 + 78*c*d^5 - 23*d^6)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) / a^3 - 20*(90*c^2*d^4*tan(1/2*f*x + 1/2*e)^5 - 126*c*d^5*tan(1/2*f*x + 1/2*e)^5 + 51*d^6*tan(1/2*f*x + 1/2*e)^5 - 180*c^2*d^4*tan(1/2*f*x + 1/2*e)^3 + 216*c*d^5*tan(1/2*f*x + 1/2*e)^3 - 76*d^6*tan(1/2*f*x + 1/2*e)^3 + 90*c^2*d^4*tan(1/2*f*x + 1/2*e) - 90*c*d^5*tan(1/2*f*x + 1/2*e) + 33*d^6*tan(1/2*f*x + 1/2*e)) / ((tan(1/2*f*x + 1/2*e)^2 - 1)^3*a^3) + (3*a^12*c^6*tan(1/2*f*x + 1/2*e)^5 - 18*a^12*c^5*d*tan(1/2*f*x + 1/2*e)^5 + 45*a^12*c^4*d^2*tan(1/2*f*x + 1/2*e)^5 - 60*a^12*c^3*d^3*tan(1/2*f*x + 1/2*e)^5 + 45*a^12*c^2*d^4*tan(1/2*f*x + 1/2*e)^5 - 18*a^12*c*d^5*tan(1/2*f*x + 1/2*e)^5 + 3*a^12*d^6*tan(1/2*f*x + 1/2*e)^5 - 10*a^12*c^6*tan(1/2*f*x + 1/2*e)^3 + 150*a^12*c^4*d^2*tan(1/2*f*x + 1/2*e)^3 - 400*a^12*c^3*d^3*tan(1/2*f*x + 1/2*e)^3 + 450*a^12*c^2*d^4*tan(1/2*f*x + 1/2*e)^3 - 240*a^12*c*d^5*tan(1/2*f*x + 1/2*e)^3 + 50*a^12*d^6*tan(1/2*f*x + 1/2*e)^3 + 15*a^12*c^6*tan(1/2*f*x + 1/2*e) + 90*a^12*c^5*d*tan(1/2*f*x + 1/2*e) + 225*a^12*c^4*d^2*tan(1/2*f*x + 1/2*e) - 2100*a^12*c^3*d^3*tan(1/2*f*x + 1/2*e) + 3825*a^12*c^2*d^4*tan(1/2*f*x + 1/2*e) - 2790*a^12*c*d^5*tan(1/2*f*x + 1/2*e) + 735*a^12*d^6*tan(1/2*f*x + 1/2*e)) / a^15) / f`

3.225.9 Mupad [B] (verification not implemented)

Time = 13.62 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.90

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{5(c-d)^6}{2a^3} - \frac{6(c+d)(c-d)^5}{a^3} + \frac{15(c+d)^2(c-d)^4}{4a^3} \right)}{f}$$

$$- \frac{(30c^2d^4 - 42cd^5 + 17d^6) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-60c^2d^4 + 72cd^5 - \frac{76d^6}{3}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + (30c^2d^4 - 30cd^5 + 17d^6) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - a^3 \right)}$$

$$+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\frac{(c-d)^6}{3a^3} - \frac{(c+d)(c-d)^5}{2a^3} \right)}{f} + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (c-d)^6}{20a^3 f}$$

$$+ \frac{d^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (40c^3 - 90c^2d + 78cd^2 - 23d^3)}{a^3 f}$$

3.225. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^6}{(a+a\sec(e+fx))^3} dx$

input `int((c + d/cos(e + f*x))^6/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`

output `(tan(e/2 + (f*x)/2)*((5*(c - d)^6)/(2*a^3) - (6*(c + d)*(c - d)^5)/a^3 + (15*(c + d)^2*(c - d)^4)/(4*a^3)))/f - (tan(e/2 + (f*x)/2)*(11*d^6 - 30*c*d^5 + 30*c^2*d^4) + tan(e/2 + (f*x)/2)^5*(17*d^6 - 42*c*d^5 + 30*c^2*d^4) - tan(e/2 + (f*x)/2)^3*((76*d^6)/3 - 72*c*d^5 + 60*c^2*d^4))/(f*(3*a^3*tan(e/2 + (f*x)/2)^2 - 3*a^3*tan(e/2 + (f*x)/2)^4 + a^3*tan(e/2 + (f*x)/2)^6 - a^3)) + (tan(e/2 + (f*x)/2)^3*((c - d)^6/(3*a^3) - ((c + d)*(c - d)^5)/(2*a^3)))/f + (tan(e/2 + (f*x)/2)^5*(c - d)^6)/(20*a^3*f) + (d^3*atanh(tan(e/2 + (f*x)/2)))*(78*c*d^2 - 90*c^2*d + 40*c^3 - 23*d^3))/(a^3*f)`

3.226 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$

3.226.1 Optimal result 1634
 3.226.2 Mathematica [A] (verified) 1635
 3.226.3 Rubi [A] (verified) 1635
 3.226.4 Maple [A] (verified) 1640
 3.226.5 Fricas [A] (verification not implemented) 1641
 3.226.6 Sympy [F] 1641
 3.226.7 Maxima [B] (verification not implemented) 1642
 3.226.8 Giac [A] (verification not implemented) 1643
 3.226.9 Mupad [B] (verification not implemented) 1644

3.226.1 Optimal result

Integrand size = 31, antiderivative size = 287

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$$

$$= \frac{d^3(20c^2 - 30cd + 13d^2) \operatorname{arctanh}(\sin(e+fx))}{2a^3 f}$$

$$- \frac{2d(2c^4 + 15c^3d + 72c^2d^2 - 180cd^3 + 76d^4) \tan(e+fx)}{15a^3 f}$$

$$- \frac{d^2(4c^3 + 30c^2d + 146cd^2 - 195d^3) \sec(e+fx) \tan(e+fx)}{30a^3 f}$$

$$+ \frac{(c-d)(2c^2 + 15cd + 76d^2)(c+d \sec(e+fx))^2 \tan(e+fx)}{15f(a^3 + a^3 \sec(e+fx))}$$

$$+ \frac{(c-d)(2c+11d)(c+d \sec(e+fx))^3 \tan(e+fx)}{15af(a+a \sec(e+fx))^2}$$

$$+ \frac{(c-d)(c+d \sec(e+fx))^4 \tan(e+fx)}{5f(a+a \sec(e+fx))^3}$$

```
output 1/2*d^3*(20*c^2-30*c*d+13*d^2)*arctanh(sin(f*x+e))/a^3/f-2/15*d*(2*c^4+15*
c^3*d+72*c^2*d^2-180*c*d^3+76*d^4)*tan(f*x+e)/a^3/f-1/30*d^2*(4*c^3+30*c^2
*d+146*c*d^2-195*d^3)*sec(f*x+e)*tan(f*x+e)/a^3/f+1/15*(c-d)*(2*c^2+15*c*d
+76*d^2)*(c+d*sec(f*x+e))^2*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))+1/15*(c-d)*
(2*c+11*d)*(c+d*sec(f*x+e))^3*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2+1/5*(c-d)*
(c+d*sec(f*x+e))^4*tan(f*x+e)/f/(a+a*sec(f*x+e))^3
```

3.226.2 Mathematica [A] (verified)

Time = 5.51 (sec) , antiderivative size = 439, normalized size of antiderivative = 1.53

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{-480d^3(20c^2 - 30cd + 13d^2) \cos^6\left(\frac{1}{2}(e+fx)\right) \left(\log\left(\cos\left(\frac{1}{2}(e+fx)\right) - \sin\left(\frac{1}{2}(e+fx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(e+fx)\right) + \sin\left(\frac{1}{2}(e+fx)\right)\right)\right)}{(120a^3f(1+\cos(e+fx)))^3}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^3,x]`output `(-480*d^3*(20*c^2 - 30*c*d + 13*d^2)*Cos[(e + f*x)/2]^6*(Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] - Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]]) + 2*Cos[(e + f*x)/2]*(29*c^5 + 105*c^4*d + 340*c^3*d^2 - 1940*c^2*d^3 + 3420*c*d^4 - 1354*d^5 + 3*(12*c^5 + 90*c^4*d + 120*c^3*d^2 - 1020*c^2*d^3 + 1910*c*d^4 - 777*d^5)*Cos[e + f*x] + 6*(6*c^5 + 20*c^4*d + 60*c^3*d^2 - 360*c^2*d^3 + 630*c*d^4 - 261*d^5)*Cos[2*(e + f*x)] + 12*c^5*Cos[3*(e + f*x)] + 90*c^4*d*Cos[3*(e + f*x)] + 120*c^3*d^2*Cos[3*(e + f*x)] - 1020*c^2*d^3*Cos[3*(e + f*x)] + 1710*c*d^4*Cos[3*(e + f*x)] - 717*d^5*Cos[3*(e + f*x)] + 7*c^5*Cos[4*(e + f*x)] + 15*c^4*d*Cos[4*(e + f*x)] + 20*c^3*d^2*Cos[4*(e + f*x)] - 220*c^2*d^3*Cos[4*(e + f*x)] + 360*c*d^4*Cos[4*(e + f*x)] - 152*d^5*Cos[4*(e + f*x)])*Sec[e + f*x]^2*Sin[(e + f*x)/2])/(120*a^3*f*(1 + Cos[e + f*x])^3)`**3.226.3 Rubi [A] (verified)**Time = 0.60 (sec) , antiderivative size = 400, normalized size of antiderivative = 1.39, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4475, 109, 25, 27, 167, 27, 167, 27, 164, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a\sec(e+fx)+a)^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^5}{(a\csc\left(e+fx+\frac{\pi}{2}\right)+a)^3} dx$$

$$\downarrow \text{4475}$$

3.226. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$

$$\frac{a^2 \tan(e+fx) \int \frac{(c+d \sec(e+fx))^5}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{7/2}} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 109

$$\frac{a^2 \tan(e+fx) \left(-\frac{\int -\frac{a^2(c+d \sec(e+fx))^3((2c-d)(c+4d)-(2c-7d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))^3((2c-d)(c+4d)-(2c-7d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{(c+d \sec(e+fx))^3((2c-d)(c+4d)-(2c-7d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 167

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))^2(2c^3+9dc^2+37d^2c-33d^3-d(4c^2+24dc-43d^2)\sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)(2c+11d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{(c+d \sec(e+fx))^2(2c^3+9dc^2+37d^2c-33d^3-d(4c^2+24dc-43d^2)\sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a} - \frac{(c-d)(2c+11d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^4}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 167

3.226. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2 d(c+d \sec(e+fx)) (d(2c^2+165dc-152d^2) - (4c^3+30dc^2+146d^2c-195d^3) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) - \frac{(c-d)(2c^2+15cd+76d^2) \sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)}}}{\frac{3a}{5a}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{d \int \frac{(c+d \sec(e+fx)) (d(2c^2+165dc-152d^2) - (4c^3+30dc^2+146d^2c-195d^3) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) - \frac{(c-d)(2c^2+15cd+76d^2) \sqrt{a-a \sec(e+fx)}}{a^2 \sqrt{a \sec(e+fx)}}}{\frac{3a}{5a}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$$

↓ 164

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{15}{2} d^2 (20c^2 - 30cd + 13d^2) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx) + \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a} (d(4c^3+30c^2d+146cd^2-195d^3) \sec(e+fx))}{a} \right)}{\frac{3a}{5a}} \right)$$

↓ 45

$$a^2 \tan(e + fx) \left(\frac{d \left(15d^2 (20c^2 - 30cd + 13d^2) \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} \frac{d \sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} + \frac{\sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a} (d(4c^3+30c^2d+146cd^2-195d^3) \sec(e+fx))}{2a} \right)}{\frac{3a}{5a}} \right)$$

↓ 218

3.226. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+a \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{d \left(\frac{\sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} (d(4c^3 + 30c^2d + 146cd^2 - 195d^3) \sec(e + fx) + 4(2c^4 + 15c^3d + 72c^2d^2 - 180cd^3 + 76d^4))}{2a^2} \right)}{\frac{a}{3a}} \right)$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^5)/(a + a*Sec[e + f*x])^3,x]`

output `-(a^2*(-1/5*((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^4)/(a^2*(a + a*Sec[e + f*x])^(5/2)) + (-1/3*((c - d)*(2*c + 11*d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^3)/(a^2*(a + a*Sec[e + f*x])^(3/2)) + (-((c - d)*(2*c^2 + 15*c*d + 76*d^2)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(a^2*Sqrt[a + a*Sec[e + f*x]])) + (d*((-15*d^2*(20*c^2 - 30*c*d + 13*d^2)*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])]/a + (Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]*(4*(2*c^4 + 15*c^3*d + 72*c^2*d^2 - 180*c*d^3 + 76*d^4) + d*(4*c^3 + 30*c^2*d + 146*c*d^2 - 195*d^3)*Sec[e + f*x]))/(2*a^2)))/a)/(3*a))/(5*a))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])`

3.226.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 109 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

rule 164 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_] := Simp[-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`

rule 167 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] - Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[b*c*(f*g - e*h)*(m + 1) + (b*g - a*h)*(d*e*n + c*f*(p + 1)) + d*(b*(f*g - e*h)*(m + 1) + f*(b*g - a*h)*(n + p + 1))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, p}, x] && LtQ[m, -1] && GtQ[n, 0] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[
^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x
], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[
b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || In
tegerQ[m - 1/2])
```

3.226.4 Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.25

method	result
parallelrisch	$-2400(c^2 - \frac{3}{2}cd + \frac{13}{20}d^2)(1 + \cos(2fx + 2e))d^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 2400(c^2 - \frac{3}{2}cd + \frac{13}{20}d^2)(1 + \cos(2fx + 2e))d^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)$
derivativedivides	$\frac{c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - c^4 d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 + 2c^3 d^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 2c^2 d^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 + c d^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - \frac{d^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5}$
default	$\frac{c^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - c^4 d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 + 2c^3 d^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - 2c^2 d^3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 + c d^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - \frac{d^5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5}$
norman	$\frac{(c^5 - 5c^4 d + 10c^3 d^2 - 10c^2 d^3 + 5c d^4 - d^5) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{15}}{20af} - \frac{5(c^5 - 3c^4 d + 2c^3 d^2 + 2c^2 d^3 - 3c d^4 + d^5) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{13}}{12af} - \frac{(c^5 + 5c^4 d + 10c^3 d^2 - 10c^2 d^3 + 5c d^4 - d^5) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{20af}$
risch	Expression too large to display

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/240*(-2400*(c^2-3/2*c*d+13/20*d^2)*(1+cos(2*f*x+2*e))*d^3*ln(tan(1/2*f*x
+1/2*e)-1)+2400*(c^2-3/2*c*d+13/20*d^2)*(1+cos(2*f*x+2*e))*d^3*ln(tan(1/2*
f*x+1/2*e)+1)+7*sec(1/2*f*x+1/2*e)^4*tan(1/2*f*x+1/2*e)*(6*(20/7*c^4*d+6/7
*c^5-261/7*d^5+60/7*c^3*d^2-360/7*c^2*d^3+90*c*d^4)*cos(2*f*x+2*e)+3/7*(4*
c^5+30*c^4*d+40*c^3*d^2-340*c^2*d^3+570*c*d^4-239*d^5)*cos(3*f*x+3*e)+(c^5
-152/7*d^5+15/7*c^4*d+20/7*c^3*d^2-220/7*c^2*d^3+360/7*c*d^4)*cos(4*f*x+4*
e)+3*(-111*d^5+120/7*c^3*d^2+12/7*c^5+90/7*c^4*d-1020/7*c^2*d^3+1910/7*c*d
^4)*cos(f*x+e)-1354/7*d^5+29/7*c^5+15*c^4*d+340/7*c^3*d^2-1940/7*c^2*d^3+3
420/7*c*d^4)/f/a^3/(1+cos(2*f*x+2*e))
```

3.226. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$

3.226.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 502, normalized size of antiderivative = 1.75

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{15((20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^5 + 3(20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^4 + 3(20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^3 + (20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^2 \log(\sin(fx+e)+1) - 15((20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^5 + 3(20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^4 + 3(20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^3 + (20c^2d^3 - 30cd^4 + 13d^5)\cos(fx+e)^2) \log(-\sin(fx+e)+1) + 2(15d^5 + 2(7c^5 + 15c^4d + 20c^3d^2 - 220c^2d^3 + 360cd^4 - 152d^5)\cos(fx+e)^4 + 3(4c^5 + 30c^4d + 40c^3d^2 - 340c^2d^3 + 570cd^4 - 239d^5)\cos(fx+e)^3 + (4c^5 + 30c^4d + 140c^3d^2 - 640c^2d^3 + 1170cd^4 - 479d^5)\cos(fx+e)^2 + 15(10cd^4 - 3d^5)\cos(fx+e))\sin(fx+e))/(a^3f\cos(fx+e)^5 + 3a^3f\cos(fx+e)^4 + 3a^3f\cos(fx+e)^3 + a^3f\cos(fx+e)^2)}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x, algorithm="fricas")
```

```
output 1/60*(15*((20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^5 + 3*(20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^4 + 3*(20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^3 + (20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^2)*log(sin(f*x + e) + 1) - 15*((20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^5 + 3*(20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^4 + 3*(20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^3 + (20*c^2*d^3 - 30*c*d^4 + 13*d^5)*cos(f*x + e)^2)*log(-sin(f*x + e) + 1) + 2*(15*d^5 + 2*(7*c^5 + 15*c^4*d + 20*c^3*d^2 - 220*c^2*d^3 + 360*c*d^4 - 152*d^5)*cos(f*x + e)^4 + 3*(4*c^5 + 30*c^4*d + 40*c^3*d^2 - 340*c^2*d^3 + 570*c*d^4 - 239*d^5)*cos(f*x + e)^3 + (4*c^5 + 30*c^4*d + 140*c^3*d^2 - 640*c^2*d^3 + 1170*c*d^4 - 479*d^5)*cos(f*x + e)^2 + 15*(10*c*d^4 - 3*d^5)*cos(f*x + e))*sin(f*x + e))/(a^3*f*cos(f*x + e)^5 + 3*a^3*f*cos(f*x + e)^4 + 3*a^3*f*cos(f*x + e)^3 + a^3*f*cos(f*x + e)^2)
```

3.226.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$$

$$= \int \frac{c^5 \sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \frac{d^5 \sec^6(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \frac{5cd^4 \sec^5(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))**5/(a+a*sec(f*x+e))**3,x)
```

```
output (Integral(c**5*sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e
+ f*x) + 1), x) + Integral(d**5*sec(e + f*x)**6/(sec(e + f*x)**3 + 3*sec(
e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(5*c*d**4*sec(e + f*x)**5/
(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(
10*c**2*d**3*sec(e + f*x)**4/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(
e + f*x) + 1), x) + Integral(10*c**3*d**2*sec(e + f*x)**3/(sec(e + f*x)**3
+ 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(5*c**4*d*sec(e +
f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a
**3
```

3.226.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 689 vs. $2(275) = 550$.

Time = 0.24 (sec) , antiderivative size = 689, normalized size of antiderivative = 2.40

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + a \sec(e + fx))^3} dx =$$

$$d^5 \left(\frac{60 \left(\frac{5 \sin(fx+e)}{\cos(fx+e)+1} - \frac{7 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right)}{a^3 - \frac{2a^3 \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{a^3 \sin(fx+e)^4}{(\cos(fx+e)+1)^4}} + \frac{\frac{465 \sin(fx+e)}{\cos(fx+e)+1} + \frac{40 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5}}{a^3} - \frac{390 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^3} + \frac{390 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^3} \right)$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x, algorithm="m
axima")
```

output

```
-1/60*(d^5*(60*(5*sin(f*x + e)/(cos(f*x + e) + 1) - 7*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)/(a^3 - 2*a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^3*sin(f*x + e)^4/(cos(f*x + e) + 1)^4) + (465*sin(f*x + e)/(cos(f*x + e) + 1) + 40*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 390*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 390*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) - 15*c*d^4*(40*sin(f*x + e)/((a^3 - a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1)) + (85*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) + 10*c^2*d^3*((105*sin(f*x + e)/(cos(f*x + e) + 1) + 20*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3) - 10*c^3*d^2*(15*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - c^5*(15*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 15*c^4*d*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3)/f
```

3.226.8 Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 504, normalized size of antiderivative = 1.76

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + a \sec(e + fx))^3} dx$$

$$\frac{30(20c^2d^3 - 30cd^4 + 13d^5) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a^3} - \frac{30(20c^2d^3 - 30cd^4 + 13d^5) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a^3} - \frac{60(10cd^4 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 7cd^5 \tan(\frac{1}{2}fx + \frac{1}{2}e))}{a^3}$$

=

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

```

output 1/60*(30*(20*c^2*d^3 - 30*c*d^4 + 13*d^5)*log(abs(tan(1/2*f*x + 1/2*e) + 1
)))/a^3 - 30*(20*c^2*d^3 - 30*c*d^4 + 13*d^5)*log(abs(tan(1/2*f*x + 1/2*e)
- 1))/a^3 - 60*(10*c*d^4*tan(1/2*f*x + 1/2*e)^3 - 7*d^5*tan(1/2*f*x + 1/2*
e)^3 - 10*c*d^4*tan(1/2*f*x + 1/2*e) + 5*d^5*tan(1/2*f*x + 1/2*e))/((tan(1
/2*f*x + 1/2*e)^2 - 1)^2*a^3) + (3*a^12*c^5*tan(1/2*f*x + 1/2*e)^5 - 15*a^
12*c^4*d*tan(1/2*f*x + 1/2*e)^5 + 30*a^12*c^3*d^2*tan(1/2*f*x + 1/2*e)^5 -
30*a^12*c^2*d^3*tan(1/2*f*x + 1/2*e)^5 + 15*a^12*c*d^4*tan(1/2*f*x + 1/2*
e)^5 - 3*a^12*d^5*tan(1/2*f*x + 1/2*e)^5 - 10*a^12*c^5*tan(1/2*f*x + 1/2*e
)^3 + 100*a^12*c^3*d^2*tan(1/2*f*x + 1/2*e)^3 - 200*a^12*c^2*d^3*tan(1/2*f
*x + 1/2*e)^3 + 150*a^12*c*d^4*tan(1/2*f*x + 1/2*e)^3 - 40*a^12*d^5*tan(1/
2*f*x + 1/2*e)^3 + 15*a^12*c^5*tan(1/2*f*x + 1/2*e) + 75*a^12*c^4*d*tan(1/
2*f*x + 1/2*e) + 150*a^12*c^3*d^2*tan(1/2*f*x + 1/2*e) - 1050*a^12*c^2*d^3
*tan(1/2*f*x + 1/2*e) + 1275*a^12*c*d^4*tan(1/2*f*x + 1/2*e) - 465*a^12*d^
5*tan(1/2*f*x + 1/2*e))/a^15)/f

```

3.226.9 Mupad [B] (verification not implemented)

Time = 13.86 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.88

$$\begin{aligned}
 & \int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + a \sec(e + fx))^3} dx \\
 &= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{3(c-d)^5}{2a^3} - \frac{15(c+d)(c-d)^4}{4a^3} + \frac{5(c+d)^2(c-d)^3}{2a^3}\right)}{f} \\
 &+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (10cd^4 - 5d^5) - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (10cd^4 - 7d^5)}{f \left(a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 2a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + a^3\right)} \\
 &+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\frac{(c-d)^5}{4a^3} - \frac{5(c+d)(c-d)^4}{12a^3}\right)}{f} + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (c-d)^5}{20a^3 f} \\
 &+ \frac{d^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (20c^2 - 30cd + 13d^2)}{a^3 f}
 \end{aligned}$$

```

input int((c + d/cos(e + f*x))^5/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)

```

output $(\tan(e/2 + (f*x)/2)*((3*(c - d)^5)/(2*a^3) - (15*(c + d)*(c - d)^4)/(4*a^3) + (5*(c + d)^2*(c - d)^3)/(2*a^3)))/f + (\tan(e/2 + (f*x)/2)*(10*c*d^4 - 5*d^5) - \tan(e/2 + (f*x)/2)^3*(10*c*d^4 - 7*d^5))/(f*(a^3*\tan(e/2 + (f*x)/2)^4 - 2*a^3*\tan(e/2 + (f*x)/2)^2 + a^3)) + (\tan(e/2 + (f*x)/2)^3*((c - d)^5/(4*a^3) - (5*(c + d)*(c - d)^4)/(12*a^3)))/f + (\tan(e/2 + (f*x)/2)^5*(c - d)^5)/(20*a^3*f) + (d^3*atanh(\tan(e/2 + (f*x)/2))*(20*c^2 - 30*c*d + 13*d^2))/(a^3*f)$

3.226. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+a\sec(e+fx))^3} dx$

$$3.227 \quad \int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^3} dx$$

3.227.1 Optimal result	1646
3.227.2 Mathematica [A] (verified)	1647
3.227.3 Rubi [A] (verified)	1647
3.227.4 Maple [A] (verified)	1651
3.227.5 Fricas [A] (verification not implemented)	1652
3.227.6 Sympy [F]	1652
3.227.7 Maxima [B] (verification not implemented)	1653
3.227.8 Giac [A] (verification not implemented)	1653
3.227.9 Mupad [B] (verification not implemented)	1654

3.227.1 Optimal result

Integrand size = 31, antiderivative size = 205

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^3} dx \\ &= \frac{(4c-3d)d^3 \operatorname{arctanh}(\sin(e+fx))}{a^3 f} + \frac{(c-d)(2c+9d)(c+d \sec(e+fx))^2 \tan(e+fx)}{15af(a+a \sec(e+fx))^2} \\ & \quad + \frac{(c-d)(c+d \sec(e+fx))^3 \tan(e+fx)}{5f(a+a \sec(e+fx))^3} \\ & \quad + \frac{(2c^4+8c^3d+21c^2d^2-88cd^3+72d^4-d^2(2c^2+10cd-27d^2) \sec(e+fx)) \tan(e+fx)}{15f(a^3+a^3 \sec(e+fx))} \end{aligned}$$

output $(4*c-3*d)*d^3*\operatorname{arctanh}(\sin(f*x+e))/a^3/f+1/15*(c-d)*(2*c+9*d)*(c+d*\sec(f*x+e))^2*\tan(f*x+e)/a/f/(a+a*\sec(f*x+e))^2+1/5*(c-d)*(c+d*\sec(f*x+e))^3*\tan(f*x+e)/f/(a+a*\sec(f*x+e))^3+1/15*(2*c^4+8*c^3*d+21*c^2*d^2-88*c*d^3+72*d^4-d^2*(2*c^2+10*c*d-27*d^2)*\sec(f*x+e))*\tan(f*x+e)/f/(a^3+a^3*\sec(f*x+e))$

3.227.2 Mathematica [A] (verified)

Time = 3.86 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.42

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{2 \cos\left(\frac{1}{2}(e+fx)\right) \left(3(c-d)^4 \sec\left(\frac{e}{2}\right) \sin\left(\frac{fx}{2}\right) - 8(c-d)^3(2c+3d) \cos^2\left(\frac{1}{2}(e+fx)\right) \sec\left(\frac{e}{2}\right) \sin\left(\frac{fx}{2}\right) + 4(c-d)^2(2c+3d)^2 \cos\left(\frac{1}{2}(e+fx)\right) \sec\left(\frac{e}{2}\right) \sin\left(\frac{fx}{2}\right) - 4(c-d)(2c+3d)^3 \cos^2\left(\frac{1}{2}(e+fx)\right) \sec\left(\frac{e}{2}\right) \sin\left(\frac{fx}{2}\right) + (2c+3d)^4 \sec\left(\frac{e}{2}\right) \sin\left(\frac{fx}{2}\right)\right)}{(a+a\sec(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^3,x]`output $(2*\cos[(e + f*x)/2]*(3*(c - d)^4*\sec[e/2]*\sin[(f*x)/2] - 8*(c - d)^3*(2*c + 3*d)*\cos[(e + f*x)/2]^2*\sec[e/2]*\sin[(f*x)/2] + 4*(c - d)^2*(7*c^2 + 26*c*d + 57*d^2)*\cos[(e + f*x)/2]^4*\sec[e/2]*\sin[(f*x)/2] - 60*d^3*\cos[(e + f*x)/2]^5*((4*c - 3*d)*(\log[\cos[(e + f*x)/2] - \sin[(e + f*x)/2]] - \log[\cos[(e + f*x)/2] + \sin[(e + f*x)/2]]) - d*\sec[e]*\sec[e + f*x]*\sin[f*x]) + 3*(c - d)^4*\cos[(e + f*x)/2]*\tan[e/2] - 8*(c - d)^3*(2*c + 3*d)*\cos[(e + f*x)/2]^3*\tan[e/2))/(15*a^3*f*(1 + \cos[e + f*x])^3)$ **3.227.3 Rubi [A] (verified)**Time = 0.50 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.51, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4475, 109, 25, 27, 167, 27, 160, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a\sec(e+fx)+a)^3} dx$$

$$\downarrow 3042$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\left(c+d\csc\left(e+fx+\frac{\pi}{2}\right)\right)^4}{\left(a\csc\left(e+fx+\frac{\pi}{2}\right)+a\right)^3} dx$$

$$\downarrow 4475$$

$$-\frac{a^2 \tan(e+fx) \int \frac{(c+d\sec(e+fx))^4}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{7/2}} d\sec(e+fx)}{f \sqrt{a-a\sec(e+fx)} \sqrt{a\sec(e+fx)+a}}$$

$$\downarrow 109$$

3.227. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{\int -\frac{a^2(c+d \sec(e+fx))^2(2c^2+6dc-3d^2-(c-6d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 25

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))^2(2c^2+6dc-3d^2-(c-6d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d \sec(e+fx))^2(2c^2+6dc-3d^2-(c-6d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^3}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 167

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))(2c^3+8dc^2+23d^2c-18a^3-d(2c^2+10dc-27d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a^3} - \frac{(c-d)(2c+9d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d \sec(e+fx))(2c^3+8dc^2+23d^2c-18a^3-d(2c^2+10dc-27d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d \sec(e+fx)}{3a} - \frac{(c-d)(2c+9d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 160

$$a^2 \tan(e + fx) \left(\frac{15d^3(4c-3d) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{a} - \frac{\sqrt{a-a \sec(e+fx)}(2c^4+8c^3d-d^2(2c^2+10cd-27d^2) \sec(e+fx)+21c^2d^2-d^3)}{a^2 \sqrt{a \sec(e+fx)+a}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

3.227. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+a \sec(e+fx))^3} dx$

$$\begin{array}{c}
 \downarrow 45 \\
 a^2 \tan(e + fx) \left(\frac{30d^3(4c-3d) \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} - \frac{\sqrt{a-a \sec(e+fx)}(2c^4+8c^3d-d^2(2c^2+10cd-27d^2)) \sec(e+fx)+21c^2d^2-88c^3d+72d^4}{a^2 \sqrt{a \sec(e+fx)+a}}}{3a} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a} \\
 \downarrow 218 \\
 a^2 \tan(e + fx) \left(-\frac{30d^3(4c-3d) \arctan\left(\frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}}\right) - \frac{\sqrt{a-a \sec(e+fx)}(2c^4+8c^3d-d^2(2c^2+10cd-27d^2)) \sec(e+fx)+21c^2d^2-88c^3d+72d^4}{a^2 \sqrt{a \sec(e+fx)+a}}}{a^2} \right) \\
 \hline
 f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}
 \end{array}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + a*Sec[e + f*x])^3,x]`

output `-(a^2*(-1/5*((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^3)/(a^2*(a + a*Sec[e + f*x])^(5/2)) + (-1/3*((c - d)*(2*c + 9*d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(a^2*(a + a*Sec[e + f*x])^(3/2)) + ((-30*(4*c - 3*d)*d^3*ArcTan[Sqrt[a - a*Sec[e + f*x]]/Sqrt[a + a*Sec[e + f*x]])/a^2 - (Sqrt[a - a*Sec[e + f*x]]*(2*c^4 + 8*c^3*d + 21*c^2*d^2 - 88*c*d^3 + 72*d^4 - d^2*(2*c^2 + 10*c*d - 27*d^2))*Sec[e + f*x]))/(a^2*Sqrt[a + a*Sec[e + f*x]]))/(3*a))/(5*a))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x])))`

3.227.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 109 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 160 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))*((g_) + (h_)*(x_)), x_] := Simp[(b^2*d*e*g - a^2*d*f*h*m - a*b*(d*(f*g + e*h) - c*f*h*(m + 1)) + b*f*h*(b*c - a*d)*(m + 1)*x*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d*(b*c - a*d)*(m + 1))), x] + Simp[(a*d*f*h*m + b*(d*(f*g + e*h) - c*f*h*(m + 2)))/(b^2*d) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`
- rule 167 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)*((g_) + (h_)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] - Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[b*c*(f*g - e*h)*(m + 1) + (b*g - a*h)*(d*e*n + c*f*(p + 1)) + d*(b*(f*g - e*h)*(m + 1) + f*(b*g - a*h)*(n + p + 1))*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, p}, x] && LtQ[m, -1] && GtQ[n, 0] && IntegersQ[2*m, 2*n, 2*p]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[
^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x
], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[
b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || In
tegerQ[m - 1/2])
```

3.227.4 Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.20

method	result
parallelrisch	$-960\left(c - \frac{3d}{4}\right) \cos(fx+e)d^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 960\left(c - \frac{3d}{4}\right) \cos(fx+e)d^3 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) + 29 \sec\left(\frac{fx}{2} + \frac{e}{2}\right)^4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)$
derivativedivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^4}{5} - \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^3 d}{5} + \frac{6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^2 d^2}{5} - \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c d^3}{5} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 d^4}{5} - \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 c^4}{3} + 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)$
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^4}{5} - \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^3 d}{5} + \frac{6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^2 d^2}{5} - \frac{4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c d^3}{5} + \frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 d^4}{5} - \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 c^4}{3} + 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)$
norman	$\frac{(c^4 - 4c^3d + 6c^2d^2 - 4cd^3 + d^4) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{13}}{20af} + \frac{(c^4 + 4c^3d + 6c^2d^2 - 28cd^3 + 25d^4) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4af} - \frac{(7c^4 + 24c^3d + 30c^2d^2 - 160cd^3 + 13d^4)}{6af}$
risch	$\frac{2i(120c^2d^2e^{4i(fx+e)} - 668cd^3e^{2i(fx+e)} + 72c^3de^{2i(fx+e)} - 300cd^3e^{5i(fx+e)} - 640cd^3e^{4i(fx+e)} + 60c^2d^2e^{i(fx+e)} - 380cd^3e^{i(fx+e)})}{1}$

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/240*(-960*(c-3/4*d)*cos(f*x+e)*d^3*ln(tan(1/2*f*x+1/2*e)-1)+960*(c-3/4*d
)*cos(f*x+e)*d^3*ln(tan(1/2*f*x+1/2*e)+1)+29*sec(1/2*f*x+1/2*e)^4*tan(1/2*
f*x+1/2*e)*(6/29*(2*c^4+12*c^3*d+12*c^2*d^2-68*c*d^3+57*d^4)*cos(2*f*x+2*e
)+1/29*(7*c^4+12*c^3*d+12*c^2*d^2-88*c*d^3+72*d^4)*cos(3*f*x+3*e)+(c^4+84/
29*c^3*d+204/29*c^2*d^2-776/29*c*d^3+684/29*d^4)*cos(f*x+e)+12/29*c^4+72/2
9*c^3*d+72/29*c^2*d^2-408/29*c*d^3+402/29*d^4))/f/a^3/cos(f*x+e)
```

$$3.227. \int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

3.227.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.88

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{15((4cd^3-3d^4)\cos(fx+e)^4 + 3(4cd^3-3d^4)\cos(fx+e)^3 + 3(4cd^3-3d^4)\cos(fx+e)^2 + (4cd^3-3d^4)\cos(fx+e))\log(\sin(fx+e)+1) - 15((4cd^3-3d^4)\cos(fx+e)^4 + 3(4cd^3-3d^4)\cos(fx+e)^3 + 3(4cd^3-3d^4)\cos(fx+e)^2 + (4cd^3-3d^4)\cos(fx+e))\log(-\sin(fx+e)+1) + 2(15d^4 + (7c^4 + 12c^3d + 12c^2d^2 - 88cd^3 + 72d^4)\cos(fx+e)^3 + 3(2c^4 + 12c^3d + 42c^2d^2 - 68cd^3 + 57d^4)\cos(fx+e)^2 + (2c^4 + 12c^3d + 42c^2d^2 - 128cd^3 + 117d^4)\cos(fx+e))\sin(fx+e)}{a^3 f \cos(fx+e)^4 + 3a^3 f \cos(fx+e)^3 + 3a^3 f \cos(fx+e)^2 + a^3 f \cos(fx+e)}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x, algorithm="fracas")
```

```
output 1/30*(15*((4*c*d^3 - 3*d^4)*cos(f*x + e)^4 + 3*(4*c*d^3 - 3*d^4)*cos(f*x + e)^3 + 3*(4*c*d^3 - 3*d^4)*cos(f*x + e)^2 + (4*c*d^3 - 3*d^4)*cos(f*x + e))*log(sin(f*x + e) + 1) - 15*((4*c*d^3 - 3*d^4)*cos(f*x + e)^4 + 3*(4*c*d^3 - 3*d^4)*cos(f*x + e)^3 + 3*(4*c*d^3 - 3*d^4)*cos(f*x + e)^2 + (4*c*d^3 - 3*d^4)*cos(f*x + e))*log(-sin(f*x + e) + 1) + 2*(15*d^4 + (7*c^4 + 12*c^3*d + 12*c^2*d^2 - 88*c*d^3 + 72*d^4)*cos(f*x + e)^3 + 3*(2*c^4 + 12*c^3*d + 42*c^2*d^2 - 68*c*d^3 + 57*d^4)*cos(f*x + e)^2 + (2*c^4 + 12*c^3*d + 42*c^2*d^2 - 128*c*d^3 + 117*d^4)*cos(f*x + e))*sin(f*x + e))/(a^3*f*cos(f*x + e)^4 + 3*a^3*f*cos(f*x + e)^3 + 3*a^3*f*cos(f*x + e)^2 + a^3*f*cos(f*x + e))
```

3.227.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

$$= \int \frac{c^4 \sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \frac{d^4 \sec^5(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \frac{4cd^3 \sec^4(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))**4/(a+a*sec(f*x+e))**3,x)
```

```
output (Integral(c**4*sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(d**4*sec(e + f*x)**5/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(4*c*d**3*sec(e + f*x)**4/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(6*c**2*d**2*sec(e + f*x)**3/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(4*c**3*d*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3
```

3.227. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$

3.227.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 475 vs. $2(199) = 398$.

Time = 0.23 (sec) , antiderivative size = 475, normalized size of antiderivative = 2.32

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

$$= 3d^4 \left(\frac{40 \sin(fx+e)}{\left(a^3 - \frac{a^3 \sin(fx+e)^2}{(\cos(fx+e)+1)^2}\right) (\cos(fx+e)+1)} + \frac{85 \sin(fx+e)}{\cos(fx+e)+1} + \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5} - \frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} + 1\right)}{a^3} + \frac{60 \log\left(\frac{\sin(fx+e)}{\cos(fx+e)+1} - 1\right)}{a^3} \right)$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x, algorithm="maxima")
```

```
output 1/60*(3*d^4*(40*sin(f*x + e)/((a^3 - a^3*sin(f*x + e)^2/(cos(f*x + e) + 1)^2)*(cos(f*x + e) + 1)) + (85*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3 - 4*c*d^3*((105*sin(f*x + e)/(cos(f*x + e) + 1) + 20*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 - 60*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)/a^3 + 60*log(sin(f*x + e)/(cos(f*x + e) + 1) - 1)/a^3 + 6*c^2*d^2*(15*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 + c^4*(15*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 + 12*c^3*d*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3)/f
```

3.227.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.82

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx =$$

$$\frac{120d^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{\left(\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right)a^3} - \frac{60(4cd^3 - 3d^4) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{a^3} + \frac{60(4cd^3 - 3d^4) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{a^3} - \frac{3a^{12}c^4 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{a^3}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+a*sec(f*x+e))^3,x, algorithm="giac")
```

$$3.227. \int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+a\sec(e+fx))^3} dx$$

output
$$\begin{aligned} & -1/60*(120*d^4*\tan(1/2*f*x + 1/2*e)/((\tan(1/2*f*x + 1/2*e)^2 - 1)*a^3) - 6 \\ & 0*(4*c*d^3 - 3*d^4)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a^3 + 60*(4*c*d^3 - \\ & 3*d^4)*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a^3 - (3*a^{12}*c^4*\tan(1/2*f*x + \\ & 1/2*e)^5 - 12*a^{12}*c^3*d*\tan(1/2*f*x + 1/2*e)^5 + 18*a^{12}*c^2*d^2*\tan(1/2 \\ & *f*x + 1/2*e)^5 - 12*a^{12}*c*d^3*\tan(1/2*f*x + 1/2*e)^5 + 3*a^{12}*d^4*\tan(1/ \\ & 2*f*x + 1/2*e)^5 - 10*a^{12}*c^4*\tan(1/2*f*x + 1/2*e)^3 + 60*a^{12}*c^2*d^2*\tan \\ & (1/2*f*x + 1/2*e)^3 - 80*a^{12}*c*d^3*\tan(1/2*f*x + 1/2*e)^3 + 30*a^{12}*d^4* \\ & \tan(1/2*f*x + 1/2*e)^3 + 15*a^{12}*c^4*\tan(1/2*f*x + 1/2*e) + 60*a^{12}*c^3*d* \\ & \tan(1/2*f*x + 1/2*e) + 90*a^{12}*c^2*d^2*\tan(1/2*f*x + 1/2*e) - 420*a^{12}*c*d \\ & ^3*\tan(1/2*f*x + 1/2*e) + 255*a^{12}*d^4*\tan(1/2*f*x + 1/2*e))/a^{15}/f \end{aligned}$$

3.227.9 Mupad [B] (verification not implemented)

Time = 13.54 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.95

$$\begin{aligned} \int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{(a + a \sec(e + fx))^3} dx &= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{3(c-d)^4}{4a^3} + \frac{3(c^2-d^2)^2}{2a^3} - \frac{2(c+d)(c-d)^3}{a^3}\right)}{f} \\ &+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\frac{(c-d)^4}{6a^3} - \frac{(c+d)(c-d)^3}{3a^3}\right)}{f} \\ &- \frac{2d^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(a^3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - a^3\right)} \\ &+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (c-d)^4}{20a^3 f} \\ &+ \frac{2d^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right) (4c - 3d)}{a^3 f} \end{aligned}$$

input $\text{int}((c + d/\cos(e + f*x))^4/(\cos(e + f*x)*(a + a/\cos(e + f*x))^3),x)$

output
$$\begin{aligned} & (\tan(e/2 + (f*x)/2)*((3*(c - d)^4)/(4*a^3) + (3*(c^2 - d^2)^2)/(2*a^3) - (\\ & 2*(c + d)*(c - d)^3/a^3))/f + (\tan(e/2 + (f*x)/2)^3*((c - d)^4/(6*a^3) - \\ & ((c + d)*(c - d)^3)/(3*a^3)))/f - (2*d^4*\tan(e/2 + (f*x)/2))/(f*(a^3*\tan(e \\ & /2 + (f*x)/2)^2 - a^3)) + (\tan(e/2 + (f*x)/2)^5*(c - d)^4)/(20*a^3*f) + (2 \\ & *d^3*\operatorname{atanh}(\tan(e/2 + (f*x)/2))*(4*c - 3*d))/(a^3*f) \end{aligned}$$

3.228 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^3} dx$

3.228.1 Optimal result 1655
 3.228.2 Mathematica [B] (verified) 1655
 3.228.3 Rubi [A] (verified) 1656
 3.228.4 Maple [A] (verified) 1659
 3.228.5 Fracas [A] (verification not implemented) 1660
 3.228.6 Sympy [F] 1661
 3.228.7 Maxima [B] (verification not implemented) 1661
 3.228.8 Giac [B] (verification not implemented) 1662
 3.228.9 Mupad [B] (verification not implemented) 1663

3.228.1 Optimal result

Integrand size = 31, antiderivative size = 133

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^3} dx$$

$$= \frac{d^3 \arctanh(\sin(e+fx))}{a^3 f} + \frac{(c-d)(c+d \sec(e+fx))^2 \tan(e+fx)}{5f(a+a \sec(e+fx))^3}$$

$$+ \frac{(c-d)(2(2c^2+8cd+11d^2)+(2c^2+11cd+29d^2)\sec(e+fx))\tan(e+fx)}{15af(a+a \sec(e+fx))^2}$$

```
output d^3*arctanh(sin(f*x+e))/a^3/f+1/5*(c-d)*(c+d*sec(f*x+e))^2*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+1/15*(c-d)*(4*c^2+16*c*d+22*d^2+(2*c^2+11*c*d+29*d^2)*sec(f*x+e))*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2
```

3.228.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 295 vs. 2(133) = 266.

Time = 2.59 (sec) , antiderivative size = 295, normalized size of antiderivative = 2.22

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^3} dx$$

$$= \frac{-240d^3 \cos^6\left(\frac{1}{2}(e+fx)\right) \left(\log\left(\cos\left(\frac{1}{2}(e+fx)\right) - \sin\left(\frac{1}{2}(e+fx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(e+fx)\right) + \sin\left(\frac{1}{2}(e+fx)\right)\right)\right)}{\dots}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^3,x]`

output $(-240*d^3*\text{Cos}[(e + f*x)/2]^6*(\text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] - \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]]) + (c - d)*\text{Cos}[(e + f*x)/2]*\text{Sec}[e/2]*(5*(8*c^2 + 17*c*d + 29*d^2)*\text{Sin}[(f*x)/2] - 15*(2*c^2 + 5*c*d + 5*d^2)*\text{Sin}[e + (f*x)/2] + 20*c^2*\text{Sin}[e + (3*f*x)/2] + 65*c*d*\text{Sin}[e + (3*f*x)/2] + 95*d^2*\text{Sin}[e + (3*f*x)/2] - 15*c^2*\text{Sin}[2*e + (3*f*x)/2] - 15*c*d*\text{Sin}[2*e + (3*f*x)/2] - 15*d^2*\text{Sin}[2*e + (3*f*x)/2] + 7*c^2*\text{Sin}[2*e + (5*f*x)/2] + 16*c*d*\text{Sin}[2*e + (5*f*x)/2] + 22*d^2*\text{Sin}[2*e + (5*f*x)/2]))/(30*a^3*f*(1 + \text{Cos}[e + f*x])^3)$

3.228.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.69, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 4475, 109, 25, 27, 162, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a \sec(e + fx) + a)^3} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})(c + d \csc(e + fx + \frac{\pi}{2}))^3}{(a \csc(e + fx + \frac{\pi}{2}) + a)^3} dx$$

↓ 4475

$$\frac{a^2 \tan(e + fx) \int \frac{(c + d \sec(e + fx))^3}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{7/2}} d \sec(e + fx)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 109

$$\frac{a^2 \tan(e + fx) \left(-\frac{\int \frac{a^2(c + d \sec(e + fx))(2c^2 + 5dc - 2d^2 + 5d^2 \sec(e + fx))}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{5/2}} d \sec(e + fx)}{5a^3} - \frac{(c - d)\sqrt{a - a \sec(e + fx)}(c + d \sec(e + fx))^2}{5a^2(a \sec(e + fx) + a)^{5/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

↓ 25

3.228. $\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + a \sec(e + fx))^3} dx$

$$\begin{aligned}
 & \frac{a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(c+d \sec(e+fx))(2c^2+5dc-2d^2+5d^2 \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d \sec(e+fx))(2c^2+5dc-2d^2+5d^2 \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d \sec(e+fx)}{5a} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}(c+d \sec(e+fx))^2}{5a^2(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
 & \quad \downarrow \text{162} \\
 & \frac{a^2 \tan(e + fx) \left(\frac{5d^3 \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a}} d \sec(e+fx)}{a^2} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}((2c^2+11cd+29d^2) \sec(e+fx)+2(2c^2+8cd+11d^2))}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{5a} \\
 & \quad \downarrow \text{45} \\
 & \frac{a^2 \tan(e + fx) \left(\frac{10d^3 \int \frac{1}{\frac{(a-a \sec(e+fx))a}{\sec(e+fx)a+a} - a} d \frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}}}{a^2} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}((2c^2+11cd+29d^2) \sec(e+fx)+2(2c^2+8cd+11d^2))}{3a^2(a \sec(e+fx)+a)^{3/2}} \right)}{5a} \\
 & \quad \downarrow \text{218} \\
 & \frac{a^2 \tan(e + fx) \left(-\frac{10d^3 \arctan\left(\frac{\sqrt{a-a \sec(e+fx)}}{\sqrt{a \sec(e+fx)+a}}\right)}{a^3} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}((2c^2+11cd+29d^2) \sec(e+fx)+2(2c^2+8cd+11d^2))}{3a^2(a \sec(e+fx)+a)^{3/2}} - \frac{(c-d)\sqrt{a-a \sec(e+fx)}}{5a^2(a \sec(e+fx)+a)^{3/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + a*Sec[e + f*x])^3,x]`

3.228. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+a \sec(e+fx))^3} dx$

```
output -((a^2*(-1/5*((c - d)*Sqrt[a - a*Sec[e + f*x]]*(c + d*Sec[e + f*x])^2)/(a^
2*(a + a*Sec[e + f*x])^(5/2)) + ((-10*d^3*ArcTan[Sqrt[a - a*Sec[e + f*x]]/
Sqrt[a + a*Sec[e + f*x]]])/a^3 - ((c - d)*Sqrt[a - a*Sec[e + f*x]]*(2*(2*c
^2 + 8*c*d + 11*d^2) + (2*c^2 + 11*c*d + 29*d^2)*Sec[e + f*x]))/(3*a^2*(a
+ a*Sec[e + f*x])^(3/2)))/(5*a))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]
*Sqrt[a + a*Sec[e + f*x]]))
```

3.228.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 109 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_
)^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f
*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1)
+ c*f*(p + 1) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p)
+ b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c,
d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] ||
IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

```
rule 162 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))
*((g_.) + (h_.)*(x_)), x_] := Simp[((b^3*c*e*g*(m + 2) - a^3*d*f*h*(n + 2)
- a^2*b*(c*f*h*m - d*(f*g + e*h)*(m + n + 3)) - a*b^2*(c*(f*g + e*h) + d*e
*g*(2*m + n + 4)) + b*(a^2*d*f*h*(m - n) - a*b*(2*c*f*h*(m + 1) - d*(f*g +
e*h)*(n + 1)) + b^2*(c*(f*g + e*h)*(m + 1) - d*e*g*(m + n + 2)))*x]/(b^2*(b
*c - a*d)^2*(m + 1)*(m + 2))*((a + b*x)^(m + 1)*(c + d*x)^(n + 1), x] + Sim
p[(f*(h/b^2) - (d*(m + n + 3)*(a^2*d*f*h*(m - n) - a*b*(2*c*f*h*(m + 1) - d
*(f*g + e*h)*(n + 1)) + b^2*(c*(f*g + e*h)*(m + 1) - d*e*g*(m + n + 2)))]/(
b^2*(b*c - a*d)^2*(m + 1)*(m + 2)) Int[(a + b*x)^(m + 2)*(c + d*x)^n, x]
, x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && (LtQ[m, -2] || (EqQ[m +
n + 3, 0] && !LtQ[n, -2]))
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))^(n_), x_Symbol] := Simp[a
^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]]))
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x
], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[
b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || In
tegerQ[m - 1/2])
```

3.228.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.88

method	result
parallelrisch	$\frac{-60 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) d^3 + 60 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) d^3 + 3(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \left((c-d)^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4 - \frac{10(c+2d)(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{3} \right)}{60a^3 f}$
derivativedivides	$-\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 c^3}{3} + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 c d^2 + 3c^2 d \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) c d^2 - \frac{3c^2 d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{3c d^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5}$
default	$-\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 c^3}{3} + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 c d^2 + 3c^2 d \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) c d^2 - \frac{3c^2 d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{3c d^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5}$
risch	$\frac{2i(15c^3 e^{4i(fx+e)} - 15d^3 e^{4i(fx+e)} + 30c^3 e^{3i(fx+e)} + 45c^2 d e^{3i(fx+e)} - 75d^3 e^{3i(fx+e)} + 40c^3 e^{2i(fx+e)} + 45c^2 d e^{2i(fx+e)} + 60d^3 e^{2i(fx+e)})}{15f a^3 (e^{i(fx+e)} - 1)}$
norman	$\frac{(c^3 - 3c^2 d + 3c d^2 - d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{20af} - \frac{(c^3 + 3c^2 d + 3c d^2 - 7d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4af} + \frac{3(3c^3 + c^2 d - c d^2 - 3d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{10af} + \frac{(11c^3 + 27c^2 d - 27c d^2 - 11d^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{10af} + \frac{1}{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}$

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBO
SE)
```

```
output 1/60*(-60*ln(tan(1/2*f*x+1/2*e)-1)*d^3+60*ln(tan(1/2*f*x+1/2*e)+1)*d^3+3*(
c-d)*tan(1/2*f*x+1/2*e)*((c-d)^2*tan(1/2*f*x+1/2*e)^4-10/3*(c+2*d)*(c-d)*t
an(1/2*f*x+1/2*e)^2+5*c^2+20*c*d+35*d^2))/a^3/f
```

3.228.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.86

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{15(d^3 \cos(fx+e)^3 + 3d^3 \cos(fx+e)^2 + 3d^3 \cos(fx+e) + d^3) \log(\sin(fx+e) + 1) - 15(d^3 \cos(fx+e)^3 + 3d^3 \cos(fx+e)^2 + 3d^3 \cos(fx+e) + d^3)}{a^3}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x, algorithm="f
ricas")
```

output
$$\frac{1}{30} \cdot (15 \cdot (d^3 \cos(fx + e))^3 + 3 \cdot d^3 \cos(fx + e)^2 + 3 \cdot d^3 \cos(fx + e) + d^3) \cdot \log(\sin(fx + e) + 1) - 15 \cdot (d^3 \cos(fx + e))^3 + 3 \cdot d^3 \cos(fx + e)^2 + 3 \cdot d^3 \cos(fx + e) + d^3) \cdot \log(-\sin(fx + e) + 1) + 2 \cdot (2 \cdot c^3 + 9 \cdot c^2 \cdot d + 21 \cdot c \cdot d^2 - 32 \cdot d^3 + (7 \cdot c^3 + 9 \cdot c^2 \cdot d + 6 \cdot c \cdot d^2 - 22 \cdot d^3) \cdot \cos(fx + e)^2 + 3 \cdot (2 \cdot c^3 + 9 \cdot c^2 \cdot d + 6 \cdot c \cdot d^2 - 17 \cdot d^3) \cdot \cos(fx + e)) \cdot \sin(fx + e) / (a^3 \cdot f \cdot \cos(fx + e)^3 + 3 \cdot a^3 \cdot f \cdot \cos(fx + e)^2 + 3 \cdot a^3 \cdot f \cdot \cos(fx + e) + a^3 \cdot f)$$

3.228.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + a \sec(e + fx))^3} dx$$

$$= \frac{\int \frac{c^3 \sec(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx + \int \frac{d^3 \sec^4(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx + \int \frac{3cd^2 \sec^3(e + fx)}{\sec^3(e + fx) + 3 \sec^2(e + fx) + 3 \sec(e + fx) + 1} dx}{a^3}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**3/(a+a*sec(f*x+e))**3,x)`

output
$$\left(\text{Integral}(c^3 \sec(e + fx) / (\sec(e + fx)^3 + 3 \sec(e + fx)^2 + 3 \sec(e + fx) + 1), x) + \text{Integral}(d^3 \sec(e + fx)^4 / (\sec(e + fx)^3 + 3 \sec(e + fx)^2 + 3 \sec(e + fx) + 1), x) + \text{Integral}(3 \cdot c \cdot d^2 \sec(e + fx)^3 / (\sec(e + fx)^3 + 3 \sec(e + fx)^2 + 3 \sec(e + fx) + 1), x) + \text{Integral}(3 \cdot c^2 \cdot d \sec(e + fx)^2 / (\sec(e + fx)^3 + 3 \sec(e + fx)^2 + 3 \sec(e + fx) + 1), x) \right) / a^3$$

3.228.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 307 vs. $2(126) = 252$.

Time = 0.23 (sec) , antiderivative size = 307, normalized size of antiderivative = 2.31

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + a \sec(e + fx))^3} dx =$$

$$d^3 \left(\frac{105 \sin(fx + e) + 20 \sin(fx + e)^3 + 3 \sin(fx + e)^5}{a^3 \cos(fx + e) + 1} - \frac{60 \log\left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} + 1\right)}{a^3} + \frac{60 \log\left(\frac{\sin(fx + e)}{\cos(fx + e) + 1} - 1\right)}{a^3} \right) - \frac{3cd^2 \left(\frac{15 \sin(fx + e)}{\cos(fx + e) + 1} + \dots \right)}{60f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

3.228.
$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + a \sec(e + fx))^3} dx$$

output
$$\begin{aligned} & -1/60*(d^3*((105*\sin(f*x + e)/(\cos(f*x + e) + 1) + 20*\sin(f*x + e)^3/(\cos(f*x + e) + 1)^3 + 3*\sin(f*x + e)^5/(\cos(f*x + e) + 1)^5)/a^3 - 60*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) + 1)/a^3 + 60*\log(\sin(f*x + e)/(\cos(f*x + e) + 1) - 1)/a^3) - 3*c*d^2*(15*\sin(f*x + e)/(\cos(f*x + e) + 1) + 10*\sin(f*x + e)^3/(\cos(f*x + e) + 1)^3 + 3*\sin(f*x + e)^5/(\cos(f*x + e) + 1)^5)/a^3 - c^3*(15*\sin(f*x + e)/(\cos(f*x + e) + 1) - 10*\sin(f*x + e)^3/(\cos(f*x + e) + 1)^3 + 3*\sin(f*x + e)^5/(\cos(f*x + e) + 1)^5)/a^3 - 9*c^2*d*(5*\sin(f*x + e)/(\cos(f*x + e) + 1) - \sin(f*x + e)^5/(\cos(f*x + e) + 1)^5)/a^3)/f \end{aligned}$$

3.228.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. $2(126) = 252$.

Time = 0.37 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.95

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{60d^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{a^3} - \frac{60d^3 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{a^3} + \frac{3a^{12}c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 9a^{12}c^2d \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 9a^{12}cd^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 3a^{12}d^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5}{a^{15}}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output
$$\begin{aligned} & 1/60*(60*d^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) + 1))/a^3 - 60*d^3*\log(\text{abs}(\tan(1/2*f*x + 1/2*e) - 1))/a^3 + (3*a^{12}*c^3*\tan(1/2*f*x + 1/2*e)^5 - 9*a^{12}*c^2*d*\tan(1/2*f*x + 1/2*e)^5 + 9*a^{12}*c*d^2*\tan(1/2*f*x + 1/2*e)^5 - 3*a^{12}*d^3*\tan(1/2*f*x + 1/2*e)^5 - 10*a^{12}*c^3*\tan(1/2*f*x + 1/2*e)^3 + 30*a^{12}*c*d^2*\tan(1/2*f*x + 1/2*e)^3 - 20*a^{12}*d^3*\tan(1/2*f*x + 1/2*e)^3 + 15*a^{12}*c^3*\tan(1/2*f*x + 1/2*e) + 45*a^{12}*c^2*d*\tan(1/2*f*x + 1/2*e) + 45*a^{12}*c*d^2*\tan(1/2*f*x + 1/2*e) - 105*a^{12}*d^3*\tan(1/2*f*x + 1/2*e))/a^{15})/f \end{aligned}$$

3.228.9 Mupad [B] (verification not implemented)

Time = 13.52 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.11

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+a\sec(e+fx))^3} dx = \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{(c-d)^3}{4a^3} - \frac{3(c+d)(c-d)^2}{4a^3} + \frac{3(c+d)^2(c-d)}{4a^3} \right)}{f}$$

$$+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\frac{(c-d)^3}{12a^3} - \frac{(c+d)(c-d)^2}{4a^3} \right)}{f}$$

$$+ \frac{2d^3 \operatorname{atanh}\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{a^3 f}$$

$$+ \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (c-d)^3}{20a^3 f}$$

input `int((c + d/cos(e + f*x))^3/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`output `(tan(e/2 + (f*x)/2)*((c - d)^3/(4*a^3) - (3*(c + d)*(c - d)^2)/(4*a^3) + (3*(c + d)^2*(c - d))/(4*a^3)))/f + (tan(e/2 + (f*x)/2)^3*((c - d)^3/(12*a^3) - ((c + d)*(c - d)^2)/(4*a^3)))/f + (2*d^3*atanh(tan(e/2 + (f*x)/2)))/(a^3*f) + (tan(e/2 + (f*x)/2)^5*(c - d)^3)/(20*a^3*f)`

3.229 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^3} dx$

3.229.1 Optimal result 1664
 3.229.2 Mathematica [A] (verified) 1664
 3.229.3 Rubi [A] (verified) 1665
 3.229.4 Maple [A] (verified) 1667
 3.229.5 Fricas [A] (verification not implemented) 1668
 3.229.6 Sympy [F] 1668
 3.229.7 Maxima [A] (verification not implemented) 1669
 3.229.8 Giac [A] (verification not implemented) 1669
 3.229.9 Mupad [B] (verification not implemented) 1670

3.229.1 Optimal result

Integrand size = 31, antiderivative size = 115

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^3} dx = \frac{(c-d)^2 \tan(e+fx)}{5f(a+a \sec(e+fx))^3} + \frac{2(c-d)(c+4d) \tan(e+fx)}{15af(a+a \sec(e+fx))^2} + \frac{(2c^2+6cd+7d^2) \tan(e+fx)}{15f(a^3+a^3 \sec(e+fx))}$$

output `1/5*(c-d)^2*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+2/15*(c-d)*(c+4*d)*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2+1/15*(2*c^2+6*c*d+7*d^2)*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))`

3.229.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.73

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^3} dx = \frac{(2c^2+6cd+7d^2+6(c^2+3cd+d^2) \cos(e+fx) + (7c^2+6cd+2d^2) \cos^2(e+fx)) \sin(e+fx)}{15a^3 f(1+\cos(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^3,x]`

output $((2*c^2 + 6*c*d + 7*d^2 + 6*(c^2 + 3*c*d + d^2)*\text{Cos}[e + f*x] + (7*c^2 + 6*c*d + 2*d^2)*\text{Cos}[e + f*x]^2)*\text{Sin}[e + f*x])/(15*a^3*f*(1 + \text{Cos}[e + f*x])^3)$

3.229.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.67, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 4475, 100, 27, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a\sec(e+fx)+a)^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))^2}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx$$

↓ 4475

$$-\frac{a^2 \tan(e+fx) \int \frac{(c+d\sec(e+fx))^2}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{7/2}} d\sec(e+fx)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

↓ 100

$$a^2 \tan(e+fx) \left(\frac{\int \frac{a^3(2c^2+6dc-3d^2+5d^2\sec(e+fx))}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d\sec(e+fx)}{5a^4} - \frac{(c-d)^2\sqrt{a-a\sec(e+fx)}}{5a^2(a\sec(e+fx)+a)^{5/2}} \right)$$

↓ 27

$$a^2 \tan(e+fx) \left(\frac{\int \frac{2c^2+6dc-3d^2+5d^2\sec(e+fx)}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{5/2}} d\sec(e+fx)}{5a} - \frac{(c-d)^2\sqrt{a-a\sec(e+fx)}}{5a^2(a\sec(e+fx)+a)^{5/2}} \right)$$

↓ 87

$$a^2 \tan(e+fx) \left(\frac{(2c^2+6cd+7d^2) \int \frac{1}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d\sec(e+fx)}{3a} - \frac{2(c-d)(c+4d)\sqrt{a-a\sec(e+fx)}}{3a^2(a\sec(e+fx)+a)^{3/2}} - \frac{(c-d)^2\sqrt{a-a\sec(e+fx)}}{5a^2(a\sec(e+fx)+a)^{5/2}} \right)$$

↓

$$-\frac{a^2 \tan(e+fx) \left(\frac{(2c^2+6cd+7d^2) \int \frac{1}{\sqrt{a-a\sec(e+fx)}(\sec(e+fx)a+a)^{3/2}} d\sec(e+fx)}{3a} - \frac{2(c-d)(c+4d)\sqrt{a-a\sec(e+fx)}}{3a^2(a\sec(e+fx)+a)^{3/2}} - \frac{(c-d)^2\sqrt{a-a\sec(e+fx)}}{5a^2(a\sec(e+fx)+a)^{5/2}} \right)}{f\sqrt{a-a\sec(e+fx)}\sqrt{a\sec(e+fx)+a}}$$

3.229. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$

$$\frac{a^2 \tan(e + fx) \left(\frac{-(2c^2 + 6cd + 7d^2)\sqrt{a - a \sec(e + fx)}}{3a^3 \sqrt{a \sec(e + fx) + a}} - \frac{2(c-d)(c+4d)\sqrt{a - a \sec(e + fx)}}{3a^2 (a \sec(e + fx) + a)^{3/2}} - \frac{(c-d)^2 \sqrt{a - a \sec(e + fx)}}{5a^2 (a \sec(e + fx) + a)^{5/2}} \right)}{f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + a*Sec[e + f*x])^3,x]`

output `-((a^2*(-1/5*((c - d)^2*Sqrt[a - a*Sec[e + f*x]])/(a^2*(a + a*Sec[e + f*x])^(5/2)) + ((-2*(c - d)*(c + 4*d)*Sqrt[a - a*Sec[e + f*x]])/(3*a^2*(a + a*Sec[e + f*x])^(3/2)) - ((2*c^2 + 6*c*d + 7*d^2)*Sqrt[a - a*Sec[e + f*x]])/(3*a^3*Sqrt[a + a*Sec[e + f*x]]))/(5*a))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]))`

3.229.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

```
rule 100 Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(
p_), x_] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d^2*(d
*e - c*f)*(n + 1))), x] - Simp[1/(d^2*(d*e - c*f)*(n + 1)) Int[(c + d*x)^(
n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1) + c*f*(
p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n + 1)*x
, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n
+ p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4475 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[a
^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]]))
Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x
], x, Csc[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[
b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || In
tegerQ[m - 1/2])
```

3.229.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.58

method	result
parallelrisch	$\frac{\left((c-d)^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4 + \frac{10(-c^2+d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2}{3} + 5(c+d)^2 \right) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{20a^3 f}$
derivativedivides	$\frac{(c-d)^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{2(-c-d)(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)(-c-d)^2}{4f a^3}$
default	$\frac{(c-d)^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} + \frac{2(-c-d)(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)(-c-d)^2}{4f a^3}$
risch	$\frac{2i(15c^2 e^{4i(fx+e)} + 30c^2 e^{3i(fx+e)} + 30cd e^{3i(fx+e)} + 40c^2 e^{2i(fx+e)} + 30cd e^{2i(fx+e)} + 20d^2 e^{2i(fx+e)} + 20c^2 e^{i(fx+e)} + 30de^{i(fx+e)})}{15f a^3 (e^{i(fx+e)} + 1)^5}$
norman	$\frac{\frac{(c^2 - 2cd + d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{20af} + \frac{(c^2 + 2cd + d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4af} - \frac{(2c^2 + 3cd + d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3af} - \frac{(4c^2 - 3cd - d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{15af} + \frac{(19c^2 - 19cd + 19d^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{15af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2 a^2}$

3.229. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+a \sec(e+fx))^3} dx$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output $\frac{1}{20}((c-d)^2 \tan(1/2*f*x+1/2*e)^4 + 10/3*(-c^2+d^2)*\tan(1/2*f*x+1/2*e)^2 + 5*(c+d)^2*\tan(1/2*f*x+1/2*e))/a^3/f$

3.229.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.98

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{((7c^2+6cd+2d^2)\cos(fx+e)^2+2c^2+6cd+7d^2+6(c^2+3cd+d^2)\cos(fx+e))\sin(fx+e)}{15(a^3f\cos(fx+e)^3+3a^3f\cos(fx+e)^2+3a^3f\cos(fx+e)+a^3f)}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x, algorithm="fricas")`

output $\frac{1}{15}((7c^2+6cd+2d^2)\cos(f*x+e)^2+2c^2+6cd+7d^2+6(c^2+3cd+d^2)\cos(f*x+e))*\sin(f*x+e)/(a^3f*\cos(f*x+e)^3+3a^3f*\cos(f*x+e)^2+3a^3f*\cos(f*x+e)+a^3f)$

3.229.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

$$= \int \frac{c^2\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \frac{d^2\sec^3(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \frac{2cd\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**2/(a+a*sec(f*x+e))**3,x)`

output $(\text{Integral}(c**2*\sec(e+f*x)/(\sec(e+f*x)**3+3*\sec(e+f*x)**2+3*\sec(e+f*x)+1),x)+\text{Integral}(d**2*\sec(e+f*x)**3/(\sec(e+f*x)**3+3*\sec(e+f*x)**2+3*\sec(e+f*x)+1),x)+\text{Integral}(2*c*d*\sec(e+f*x)**2/(\sec(e+f*x)**3+3*\sec(e+f*x)**2+3*\sec(e+f*x)+1),x))/a**3$

3.229. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$

3.229.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.60

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{d^2 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} + \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3} + \frac{c^2 \left(\frac{15 \sin(fx+e)}{\cos(fx+e)+1} - \frac{10 \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3 \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{a^3} + \frac{6cd \left(\frac{5 \sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)}{\cos(fx+e)} \right)}{a^3}$$

$60 f$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x, algorithm="maxima")
```

```
output 1/60*(d^2*(15*sin(f*x + e)/(cos(f*x + e) + 1) + 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 + c^2*(15*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 + 6*c*d*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3/f
```

3.229.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.12

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{3c^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 6cd \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 + 3d^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 10c^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + 10d^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{60a^3f}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+a*sec(f*x+e))^3,x, algorithm="giac")
```

```
output 1/60*(3*c^2*tan(1/2*f*x + 1/2*e)^5 - 6*c*d*tan(1/2*f*x + 1/2*e)^5 + 3*d^2*tan(1/2*f*x + 1/2*e)^5 - 10*c^2*tan(1/2*f*x + 1/2*e)^3 + 10*d^2*tan(1/2*f*x + 1/2*e)^3 + 15*c^2*tan(1/2*f*x + 1/2*e) + 30*c*d*tan(1/2*f*x + 1/2*e) + 15*d^2*tan(1/2*f*x + 1/2*e))/(a^3*f)
```

3.229.9 Mupad [B] (verification not implemented)

Time = 13.75 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.69

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+a\sec(e+fx))^3} dx = \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (c+d)^2}{4a^3 f} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 (2c^2 - 2d^2)}{12a^3 f} + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 (c-d)^2}{20a^3 f}$$

input `int((c + d/cos(e + f*x))^2/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`output `(tan(e/2 + (f*x)/2)*(c + d)^2)/(4*a^3*f) - (tan(e/2 + (f*x)/2)^3*(2*c^2 - 2*d^2))/(12*a^3*f) + (tan(e/2 + (f*x)/2)^5*(c - d)^2)/(20*a^3*f)`

3.230 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^3} dx$

3.230.1 Optimal result 1671
 3.230.2 Mathematica [A] (verified) 1671
 3.230.3 Rubi [A] (verified) 1672
 3.230.4 Maple [A] (verified) 1674
 3.230.5 Fricas [A] (verification not implemented) 1674
 3.230.6 Sympy [F] 1675
 3.230.7 Maxima [A] (verification not implemented) 1675
 3.230.8 Giac [A] (verification not implemented) 1676
 3.230.9 Mupad [B] (verification not implemented) 1676

3.230.1 Optimal result

Integrand size = 29, antiderivative size = 102

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^3} dx = \frac{(c-d) \tan(e+fx)}{5f(a+a \sec(e+fx))^3} + \frac{(2c+3d) \tan(e+fx)}{15af(a+a \sec(e+fx))^2} + \frac{(2c+3d) \tan(e+fx)}{15f(a^3+a^3 \sec(e+fx))}$$

output `1/5*(c-d)*tan(f*x+e)/f/(a+a*sec(f*x+e))^3+1/15*(2*c+3*d)*tan(f*x+e)/a/f/(a+a*sec(f*x+e))^2+1/15*(2*c+3*d)*tan(f*x+e)/f/(a^3+a^3*sec(f*x+e))`

3.230.2 Mathematica [A] (verified)

Time = 0.95 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.32

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^3} dx = \frac{\cos(\frac{1}{2}(e+fx)) \sec(\frac{e}{2}) (5(8c+3d) \sin(\frac{fx}{2}) - 15(2c+d) \sin(e+\frac{fx}{2}) + 20c \sin(e+\frac{3fx}{2}) + 15d \sin(e+\frac{5fx}{2}))}{30a^3 f(1+\cos(e+fx))^3}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + a*Sec[e + f*x])^3,x]`

output $(\text{Cos}[(e + f*x)/2]*\text{Sec}[e/2]*(5*(8*c + 3*d)*\text{Sin}[(f*x)/2] - 15*(2*c + d)*\text{Sin}[e + (f*x)/2] + 20*c*\text{Sin}[e + (3*f*x)/2] + 15*d*\text{Sin}[e + (3*f*x)/2] - 15*c*\text{Sin}[2*e + (3*f*x)/2] + 7*c*\text{Sin}[2*e + (5*f*x)/2] + 3*d*\text{Sin}[2*e + (5*f*x)/2]))/(30*a^3*f*(1 + \text{Cos}[e + f*x])^3)$

3.230.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.207$, Rules used = {3042, 4488, 3042, 4283, 3042, 4281}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a\sec(e+fx)+a)^3} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))}{(a\csc(e+fx+\frac{\pi}{2})+a)^3} dx \\ & \quad \downarrow 4488 \\ & \frac{(2c+3d) \int \frac{\sec(e+fx)}{(\sec(e+fx)a+a)^2} dx}{5a} + \frac{(c-d)\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\ & \quad \downarrow 3042 \\ & \frac{(2c+3d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{(\csc(e+fx+\frac{\pi}{2})a+a)^2} dx}{5a} + \frac{(c-d)\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\ & \quad \downarrow 4283 \\ & \frac{(2c+3d) \left(\frac{\int \frac{\sec(e+fx)}{\sec(e+fx)a+a} dx}{3a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \right)}{5a} + \frac{(c-d)\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\ & \quad \downarrow 3042 \\ & \frac{(2c+3d) \left(\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\csc(e+fx+\frac{\pi}{2})a+a} dx}{3a} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2} \right)}{5a} + \frac{(c-d)\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} \\ & \quad \downarrow 4281 \end{aligned}$$

3.230. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$

$$\frac{(c-d)\tan(e+fx)}{5f(a\sec(e+fx)+a)^3} + \frac{(2c+3d)\left(\frac{\tan(e+fx)}{3af(a\sec(e+fx)+a)} + \frac{\tan(e+fx)}{3f(a\sec(e+fx)+a)^2}\right)}{5a}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + a*Sec[e + f*x])^3,x]`

output `((c - d)*Tan[e + f*x])/(5*f*(a + a*Sec[e + f*x])^3) + ((2*c + 3*d)*(Tan[e + f*x]/(3*f*(a + a*Sec[e + f*x])^2) + Tan[e + f*x]/(3*a*f*(a + a*Sec[e + f*x]))))/(5*a)`

3.230.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4281 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[-Cot[e + f*x]/(f*(b + a*Csc[e + f*x])), x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4283 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[b*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(m + 1)/(a*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && IntegerQ[2*m]`

rule 4488 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(A*b - a*B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*B*m + A*b*(m + 1))/(a*b*(2*m + 1)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && NeQ[a*B*m + A*b*(m + 1), 0] && LtQ[m, -2^(-1)]`

3.230.4 Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.55

method	result
parallelrisch	$\frac{\left((c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^4 - \frac{10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c}{3} + 5c + 5d \right) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{20a^3 f}$
derivativedivides	$\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f a^3}$
default	$\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{5} - \frac{2c \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3} + c \tan\left(\frac{fx}{2} + \frac{e}{2}\right) + d \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4f a^3}$
risch	$\frac{2i(15ce^{4i(fx+e)} + 30ce^{3i(fx+e)} + 15de^{3i(fx+e)} + 40e^{2i(fx+e)}c + 15de^{2i(fx+e)} + 20e^{i(fx+e)}c + 15de^{i(fx+e)} + 7c + 3d)}{15f a^3 (e^{i(fx+e)} + 1)^5}$
norman	$\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{20af} - \frac{(c+d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{4af} + \frac{(5c+3d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{12af} - \frac{(13c-3d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{60af}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1 \right) a^2}$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `1/20*((c-d)*tan(1/2*f*x+1/2*e)^4-10/3*tan(1/2*f*x+1/2*e)^2*c+5*c+5*d)*tan(1/2*f*x+1/2*e)/a^3/f`

3.230.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{(a + a \sec(e + fx))^3} dx$$

$$= \frac{((7c + 3d) \cos(fx + e)^2 + 3(2c + 3d) \cos(fx + e) + 2c + 3d) \sin(fx + e)}{15(a^3 f \cos(fx + e)^3 + 3a^3 f \cos(fx + e)^2 + 3a^3 f \cos(fx + e) + a^3 f)}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="fricas")`

output `1/15*((7*c + 3*d)*cos(f*x + e)^2 + 3*(2*c + 3*d)*cos(f*x + e) + 2*c + 3*d)*sin(f*x + e)/(a^3*f*cos(f*x + e)^3 + 3*a^3*f*cos(f*x + e)^2 + 3*a^3*f*cos(f*x + e) + a^3*f)`

3.230.
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+a \sec(e+fx))^3} dx$$

3.230.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{\int \frac{c\sec(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx + \int \frac{d\sec^2(e+fx)}{\sec^3(e+fx)+3\sec^2(e+fx)+3\sec(e+fx)+1} dx}{a^3}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))**3,x)`

output `(Integral(c*sec(e + f*x)/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x) + Integral(d*sec(e + f*x)**2/(sec(e + f*x)**3 + 3*sec(e + f*x)**2 + 3*sec(e + f*x) + 1), x))/a**3`

3.230.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.13

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{c\left(\frac{15\sin(fx+e)}{\cos(fx+e)+1} - \frac{10\sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{3\sin(fx+e)^5}{(\cos(fx+e)+1)^5}\right) + \frac{3d\left(\frac{5\sin(fx+e)}{\cos(fx+e)+1} - \frac{\sin(fx+e)^5}{(\cos(fx+e)+1)^5}\right)}{60f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="maxima")`

output `1/60*(c*(15*sin(f*x + e)/(cos(f*x + e) + 1) - 10*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 3*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3 + 3*d*(5*sin(f*x + e)/(cos(f*x + e) + 1) - sin(f*x + e)^5/(cos(f*x + e) + 1)^5)/a^3)/f`

3.230.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.74

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{3c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 3d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 - 10c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + 15c \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 15d \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{60a^3f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^3,x, algorithm="giac")`

output `1/60*(3*c*tan(1/2*f*x + 1/2*e)^5 - 3*d*tan(1/2*f*x + 1/2*e)^5 - 10*c*tan(1/2*f*x + 1/2*e)^3 + 15*c*tan(1/2*f*x + 1/2*e) + 15*d*tan(1/2*f*x + 1/2*e))/(a^3*f)`

3.230.9 Mupad [B] (verification not implemented)

Time = 13.51 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.65

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+a\sec(e+fx))^3} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(15c + 15d - 10c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 3c \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 3d \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4\right)}{60a^3f}$$

input `int((c + d/cos(e + f*x))/(cos(e + f*x)*(a + a/cos(e + f*x))^3),x)`

output `(tan(e/2 + (f*x)/2)*(15*c + 15*d - 10*c*tan(e/2 + (f*x)/2)^2 + 3*c*tan(e/2 + (f*x)/2)^4 - 3*d*tan(e/2 + (f*x)/2)^4)/(60*a^3*f)`

3.231
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))} dx$$

3.231.1 Optimal result 1677
 3.231.2 Mathematica [C] (verified) 1678
 3.231.3 Rubi [A] (verified) 1678
 3.231.4 Maple [A] (verified) 1682
 3.231.5 Fricas [B] (verification not implemented) 1683
 3.231.6 Sympy [F] 1684
 3.231.7 Maxima [F(-2)] 1685
 3.231.8 Giac [B] (verification not implemented) 1685
 3.231.9 Mupad [B] (verification not implemented) 1686

3.231.1 Optimal result

Integrand size = 31, antiderivative size = 181

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))} dx = -\frac{2d^3 \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a^3(c-d)^{7/2} \sqrt{c+d} f} + \frac{\tan(e+fx)}{5(c-d)f(a+a \sec(e+fx))^3} + \frac{(2c-7d) \tan(e+fx)}{15a(c-d)^2 f(a+a \sec(e+fx))^2} + \frac{(2c^2-9cd+22d^2) \tan(e+fx)}{15(c-d)^3 f(a^3+a^3 \sec(e+fx))}$$

output

```
-2*d^3*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a^3/(c-d)^(7/2)
/f/(c+d)^(1/2)+1/5*tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))^3+1/15*(2*c-7*d)*ta
n(f*x+e)/a/(c-d)^2/f/(a+a*sec(f*x+e))^2+1/15*(2*c^2-9*c*d+22*d^2)*tan(f*x+
e)/(c-d)^3/f/(a^3+a^3*sec(f*x+e))
```

3.231.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.16 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.91

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))} dx$$

$$\cos\left(\frac{1}{2}(e + fx)\right) \left(\frac{480d^3 \arctan\left(\frac{(i \cos(e) + \sin(e))(c \sin(e) + (-d + c \cos(e)) \tan\left(\frac{fx}{2}\right))}{\sqrt{c^2 - d^2} \sqrt{(\cos(e) - i \sin(e))^2}}\right)}{\sqrt{c^2 - d^2} \sqrt{(\cos(e) - i \sin(e))^2}} \right) \cos^5\left(\frac{1}{2}(e + fx)\right) (i \cos(e) + \sin(e)) + \sec\left(\frac{e}{2}\right) (5(8c^2$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])),x]`

output `(Cos[(e + f*x)/2]*((480*d^3*ArcTan[((I*Cos[e] + Sin[e])*(c*Sin[e] + (-d + c*Cos[e])*Tan[(f*x)/2]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]))*Cos[(e + f*x)/2]^5*(I*Cos[e] + Sin[e]))/(Sqrt[c^2 - d^2]*Sqrt[(Cos[e] - I*Sin[e])^2]) + Sec[e/2]*(5*(8*c^2 - 27*c*d + 37*d^2)*Sin[(f*x)/2] - 15*(2*c^2 - 7*c*d + 9*d^2)*Sin[e + (f*x)/2] + 20*c^2*Sin[e + (3*f*x)/2] - 75*c*d*Sin[e + (3*f*x)/2] + 115*d^2*Sin[e + (3*f*x)/2] - 15*c^2*Sin[2*e + (3*f*x)/2] + 45*c*d*Sin[2*e + (3*f*x)/2] - 45*d^2*Sin[2*e + (3*f*x)/2] + 7*c^2*Sin[2*e + (5*f*x)/2] - 24*c*d*Sin[2*e + (5*f*x)/2] + 32*d^2*Sin[2*e + (5*f*x)/2]))/(30*a^3*(c - d)^3*f*(1 + Cos[e + f*x])^3)`

3.231.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.62, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4475, 115, 25, 27, 169, 25, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{(a \sec(e + fx) + a)^3 (c + d \sec(e + fx))} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right)}{\left(a \csc\left(e + fx + \frac{\pi}{2}\right) + a\right)^3 (c + d \csc\left(e + fx + \frac{\pi}{2}\right))} dx$$

3.231. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))} dx$

$$\begin{array}{c}
\downarrow 4475 \\
\frac{a^2 \tan(e+fx) \int \frac{1}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{7/2} (c+d \sec(e+fx))} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 115 \\
\frac{a^2 \tan(e+fx) \left(-\frac{\int \frac{a^2(2c-5d+2d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx))} d \sec(e+fx)}{5a^3(c-d)} - \frac{\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 25 \\
\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(2c-5d+2d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx))} d \sec(e+fx)}{5a^3(c-d)} - \frac{\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 27 \\
\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{2c-5d+2d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx))} d \sec(e+fx)}{5a(c-d)} - \frac{\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 169 \\
\frac{a^2 \tan(e+fx) \left(-\frac{\int \frac{a^2(2c^2-7dc+15d^2+(2c-7d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{3/2} (c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(2c-7d)\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} - \frac{\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 25 \\
\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(2c^2-7dc+15d^2+(2c-7d)d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{3/2} (c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(2c-7d)\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} - \frac{\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}} \\
\downarrow 27
\end{array}$$

3.231. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 (c+d \sec(e+fx))} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{2c^2 - 7dc + 15d^2 + (2c - 7d)d \sec(e + fx)}{\sqrt{a - a \sec(e + fx)}(\sec(e + fx)a + a)^{3/2}(c + d \sec(e + fx))} d \sec(e + fx)}{3a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}} - \frac{\sqrt{a - a \sec(e + fx)}}{5a^2(c - d)(a \sec(e + fx) + a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 169

$$a^2 \tan(e + fx) \left(\frac{-\frac{\int \frac{15a^2 d^3}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a}(c + d \sec(e + fx))} d \sec(e + fx)}{a^3(c - d)} - \frac{(2c^2 - 9cd + 22d^2)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}}}{3a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{5a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{-\frac{15d^3 \int \frac{1}{\sqrt{a - a \sec(e + fx)} \sqrt{\sec(e + fx)a + a}(c + d \sec(e + fx))} d \sec(e + fx)}{a(c - d)} - \frac{(2c^2 - 9cd + 22d^2)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}}}{3a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{5a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 104

$$a^2 \tan(e + fx) \left(\frac{-\frac{30d^3 \int \frac{1}{a(c - d) + \frac{a(c + d)(\sec(e + fx)a + a)}{a - a \sec(e + fx)}} d \frac{\sqrt{\sec(e + fx)a + a}}{\sqrt{a - a \sec(e + fx)}}}{a(c - d)} - \frac{(2c^2 - 9cd + 22d^2)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}}}{3a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{5a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 218

$$a^2 \tan(e + fx) \left(\frac{-\frac{30d^3 \arctan\left(\frac{\sqrt{c + d}\sqrt{a \sec(e + fx) + a}}{\sqrt{c - d}\sqrt{a - a \sec(e + fx)}}\right)}{a^2(c - d)^{3/2}\sqrt{c + d}} - \frac{(2c^2 - 9cd + 22d^2)\sqrt{a - a \sec(e + fx)}}{a^2(c - d)\sqrt{a \sec(e + fx) + a}} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{3a^2(c - d)(a \sec(e + fx) + a)^{3/2}}}{3a(c - d)} - \frac{(2c - 7d)\sqrt{a - a \sec(e + fx)}}{5a(c - d)} - \frac{\sqrt{a - a \sec(e + fx)}}{5a^2(c - d)(a \sec(e + fx) + a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

input `Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])),x]`

output `-((a^2*(-1/5*Sqrt[a - a*Sec[e + f*x]]/(a^2*(c - d)*(a + a*Sec[e + f*x])^(5/2)) + (-1/3*((2*c - 7*d)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*(a + a*Sec[e + f*x])^(3/2)) + ((-30*d^3*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(a^2*(c - d)^(3/2)*Sqrt[c + d]) - ((2*c^2 - 9*c*d + 22*d^2)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*Sqrt[a + a*Sec[e + f*x]]))/(3*a*(c - d))/(5*a*(c - d))*Tan[e + f*x]/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))`

3.231.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 115 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 169 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_))*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_))*((csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.)^(n_)), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.231.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.12

method	result
derivativedivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^2 - 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 cd + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 d^2 - 2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 cd - 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{(c-d)^3} \frac{4fa^3}{4fa^3}$
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^2 - 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 cd + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 d^2 - 2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 cd - 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{(c-d)^3} \frac{4fa^3}{4fa^3}$
risch	$\frac{2i(15c^2e^{4i(fx+e)} - 45cde^{4i(fx+e)} + 45d^2e^{4i(fx+e)} + 30c^2e^{3i(fx+e)} - 105cde^{3i(fx+e)} + 135d^2e^{3i(fx+e)} + 40c^2e^{2i(fx+e)} - 120cde^{2i(fx+e)} + 120d^2e^{2i(fx+e)} + 15c^2e^{i(fx+e)} - 30cde^{i(fx+e)} + 15d^2e^{i(fx+e)})}{15fa^3(c-d)^3} (e^{i(fx+e)} + 1)$

3.231. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))} dx$

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/4/f/a^3*(1/(c-d)^3*(1/5*tan(1/2*f*x+1/2*e)^5*c^2-2/5*tan(1/2*f*x+1/2*e)^5*c*d+1/5*tan(1/2*f*x+1/2*e)^5*d^2-2/3*c^2*tan(1/2*f*x+1/2*e)^3+2*tan(1/2*f*x+1/2*e)^3*c*d-4/3*tan(1/2*f*x+1/2*e)^3*d^2+tan(1/2*f*x+1/2*e)*c^2-4*tan(1/2*f*x+1/2*e)*c*d+7*tan(1/2*f*x+1/2*e)*d^2)-8*d^3/(c-d)^3/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))`

3.231.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. $2(166) = 332$.

Time = 0.30 (sec) , antiderivative size = 1001, normalized size of antiderivative = 5.53

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c+d\sec(e+fx))} dx$$

$$= \left[\frac{15(d^3 \cos^3(fx+e) + 3d^3 \cos^2(fx+e) + 3d^3 \cos(fx+e) + d^3) \sqrt{c^2-d^2} \log\left(\frac{2cd \cos(fx+e) - (c^2-2d^2) \cos^2(fx+e) + d^2}{c^2-d^2}\right)}{30((a^3c^5 - 3a^3c^4d + 2a^3c^3d^2 + 2a^3c^2d^3 - 3a^3cd^4 + a^3d^5)f \cos^3(fx+e) + 3(a^3c^5 - 3a^3c^4d + 2a^3c^3d^2 + 2a^3c^2d^3 - 3a^3cd^4 + a^3d^5)f \cos^2(fx+e) + 3(a^3c^5 - 3a^3c^4d + 2a^3c^3d^2 + 2a^3c^2d^3 - 3a^3cd^4 + a^3d^5)f \cos(fx+e) + 3(a^3c^5 - 3a^3c^4d + 2a^3c^3d^2 + 2a^3c^2d^3 - 3a^3cd^4 + a^3d^5))} \right. \\ \left. - \frac{15(d^3 \cos^3(fx+e) + 3d^3 \cos^2(fx+e) + 3d^3 \cos(fx+e) + d^3) \sqrt{-c^2+d^2} \arctan\left(-\frac{\sqrt{-c^2+d^2}(d \cos(fx+e) + c)}{(c^2-d^2) \sin(fx+e)}\right)}{15((a^3c^5 - 3a^3c^4d + 2a^3c^3d^2 + 2a^3c^2d^3 - 3a^3cd^4 + a^3d^5)f \cos^3(fx+e) + 3(a^3c^5 - 3a^3c^4d + 2a^3c^3d^2 + 2a^3c^2d^3 - 3a^3cd^4 + a^3d^5))} \right]$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output

```

[-1/30*(15*(d^3*cos(f*x + e)^3 + 3*d^3*cos(f*x + e)^2 + 3*d^3*cos(f*x + e)
+ d^3)*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x +
e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/
(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) - 2*(2*c^4 - 9*c^3*d + 20
*c^2*d^2 + 9*c*d^3 - 22*d^4 + (7*c^4 - 24*c^3*d + 25*c^2*d^2 + 24*c*d^3 -
32*d^4)*cos(f*x + e)^2 + 3*(2*c^4 - 9*c^3*d + 15*c^2*d^2 + 9*c*d^3 - 17*d^
4)*cos(f*x + e))*sin(f*x + e))/((a^3*c^5 - 3*a^3*c^4*d + 2*a^3*c^3*d^2 + 2
*a^3*c^2*d^3 - 3*a^3*c*d^4 + a^3*d^5)*f*cos(f*x + e)^3 + 3*(a^3*c^5 - 3*a^
3*c^4*d + 2*a^3*c^3*d^2 + 2*a^3*c^2*d^3 - 3*a^3*c*d^4 + a^3*d^5)*f*cos(f*x
+ e)^2 + 3*(a^3*c^5 - 3*a^3*c^4*d + 2*a^3*c^3*d^2 + 2*a^3*c^2*d^3 - 3*a^3
*c*d^4 + a^3*d^5)*f*cos(f*x + e) + (a^3*c^5 - 3*a^3*c^4*d + 2*a^3*c^3*d^2
+ 2*a^3*c^2*d^3 - 3*a^3*c*d^4 + a^3*d^5)*f), -1/15*(15*(d^3*cos(f*x + e)^3
+ 3*d^3*cos(f*x + e)^2 + 3*d^3*cos(f*x + e) + d^3)*sqrt(-c^2 + d^2)*arcta
n(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) - (2*
c^4 - 9*c^3*d + 20*c^2*d^2 + 9*c*d^3 - 22*d^4 + (7*c^4 - 24*c^3*d + 25*c^2
*d^2 + 24*c*d^3 - 32*d^4)*cos(f*x + e)^2 + 3*(2*c^4 - 9*c^3*d + 15*c^2*d^2
+ 9*c*d^3 - 17*d^4)*cos(f*x + e))*sin(f*x + e))/((a^3*c^5 - 3*a^3*c^4*d +
2*a^3*c^3*d^2 + 2*a^3*c^2*d^3 - 3*a^3*c*d^4 + a^3*d^5)*f*cos(f*x + e)^3 +
3*(a^3*c^5 - 3*a^3*c^4*d + 2*a^3*c^3*d^2 + 2*a^3*c^2*d^3 - 3*a^3*c*d^4 +
a^3*d^5)*f*cos(f*x + e)^2 + 3*(a^3*c^5 - 3*a^3*c^4*d + 2*a^3*c^3*d^2 + ...

```

3.231.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))} dx$$

$$= \frac{\int \frac{\sec(e + fx)}{c \sec^3(e + fx) + 3c \sec^2(e + fx) + 3c \sec(e + fx) + c + d \sec^4(e + fx) + 3d \sec^3(e + fx) + 3d \sec^2(e + fx) + d \sec(e + fx)} dx}{a^3}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e)),x)`

output `Integral(sec(e + f*x)/(c*sec(e + f*x)**3 + 3*c*sec(e + f*x)**2 + 3*c*sec(e + f*x) + c + d*sec(e + f*x)**4 + 3*d*sec(e + f*x)**3 + 3*d*sec(e + f*x)**2 + d*sec(e + f*x)), x)/a**3`

3.231.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de
```

3.231.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 471 vs. 2(166) = 332.

Time = 0.36 (sec) , antiderivative size = 471, normalized size of antiderivative = 2.60

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))} dx = \frac{120 \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2c+2d) + \arctan \left(-\frac{c \tan(\frac{1}{2} fx + \frac{1}{2} e) - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-c^2+d^2}} \right) \right) d^3}{(a^3 c^3 - 3 a^3 c^2 d + 3 a^3 c d^2 - a^3 d^3) \sqrt{-c^2+d^2}} - \frac{3 a^{12} c^4 \tan(\frac{1}{2} fx + \frac{1}{2} e)^5 - 12 a^{12} c^3 d \tan(\frac{1}{2} fx + \frac{1}{2} e)^4}{(a^3 c^3 - 3 a^3 c^2 d + 3 a^3 c d^2 - a^3 d^3) \sqrt{-c^2+d^2}}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e)),x, algorithm="giac")
```

output
$$\begin{aligned} & -1/60*(120*(\pi*\text{floor}(1/2*(f*x + e)/\pi + 1/2)*\text{sgn}(-2*c + 2*d) + \arctan(-(c*\tan(1/2*f*x + 1/2*e) - d*\tan(1/2*f*x + 1/2*e))/\sqrt{-c^2 + d^2}))*d^3/((a^3*c^3 - 3*a^3*c^2*d + 3*a^3*c*d^2 - a^3*d^3)*\sqrt{-c^2 + d^2}) - (3*a^{12}*c^4*\tan(1/2*f*x + 1/2*e)^5 - 12*a^{12}*c^3*d*\tan(1/2*f*x + 1/2*e)^5 + 18*a^{12}*c^2*d^2*\tan(1/2*f*x + 1/2*e)^5 - 12*a^{12}*c*d^3*\tan(1/2*f*x + 1/2*e)^5 + 3*a^{12}*d^4*\tan(1/2*f*x + 1/2*e)^5 - 10*a^{12}*c^4*\tan(1/2*f*x + 1/2*e)^3 + 50*a^{12}*c^3*d*\tan(1/2*f*x + 1/2*e)^3 - 90*a^{12}*c^2*d^2*\tan(1/2*f*x + 1/2*e)^3 + 70*a^{12}*c*d^3*\tan(1/2*f*x + 1/2*e)^3 - 20*a^{12}*d^4*\tan(1/2*f*x + 1/2*e)^3 + 15*a^{12}*c^4*\tan(1/2*f*x + 1/2*e) - 90*a^{12}*c^3*d*\tan(1/2*f*x + 1/2*e) + 240*a^{12}*c^2*d^2*\tan(1/2*f*x + 1/2*e) - 270*a^{12}*c*d^3*\tan(1/2*f*x + 1/2*e) + 105*a^{12}*d^4*\tan(1/2*f*x + 1/2*e))/(a^{15}*c^5 - 5*a^{15}*c^4*d + 10*a^{15}*c^3*d^2 - 10*a^{15}*c^2*d^3 + 5*a^{15}*c*d^4 - a^{15}*d^5))/f \end{aligned}$$

3.231.9 Mupad [B] (verification not implemented)

Time = 13.80 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.26

$$\begin{aligned} & \int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))} dx \\ & = \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{3}{4a^3(c-d)} - \frac{(c+d) \left(\frac{3}{4a^3(c-d)} - \frac{c+d}{4a^3(c-d)^2} \right)}{c-d} \right)}{f} \\ & \quad - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\frac{1}{4a^3(c-d)} - \frac{c+d}{12a^3(c-d)^2} \right)}{f} + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{20a^3 f (c-d)} \\ & \quad - \frac{2d^3 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) (2c-2d) (a^3 c^3 - 3a^3 c^2 d + 3a^3 c d^2 - a^3 d^3)}{2a^3 \sqrt{c+d} (c-d)^{7/2}} \right)}{a^3 f \sqrt{c+d} (c-d)^{7/2}} \end{aligned}$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c + d/cos(e + f*x))),x)`

output
$$\begin{aligned} & (\tan(e/2 + (f*x)/2)*(3/(4*a^3*(c - d)) - ((c + d)*(3/(4*a^3*(c - d)) - (c + d)/(4*a^3*(c - d)^2)))/(c - d))/f - (\tan(e/2 + (f*x)/2)^3*(1/(4*a^3*(c - d)) - (c + d)/(12*a^3*(c - d)^2))/f + \tan(e/2 + (f*x)/2)^5/(20*a^3*f*(c - d)) - (2*d^3*\operatorname{atanh}((\tan(e/2 + (f*x)/2)*(2*c - 2*d)*(a^3*c^3 - a^3*d^3 + 3*a^3*c*d^2 - 3*a^3*c^2*d))/(2*a^3*(c + d)^{(1/2)*(c - d)^{(7/2)}})))/(a^3*f*(c + d)^{(1/2)*(c - d)^{(7/2)}}) \end{aligned}$$

3.232
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^2} dx$$

3.232.1 Optimal result 1687
 3.232.2 Mathematica [C] (warning: unable to verify) 1688
 3.232.3 Rubi [A] (verified) 1688
 3.232.4 Maple [A] (verified) 1693
 3.232.5 Fricas [B] (verification not implemented) 1694
 3.232.6 Sympy [F] 1695
 3.232.7 Maxima [F(-2)] 1696
 3.232.8 Giac [B] (verification not implemented) 1696
 3.232.9 Mupad [B] (verification not implemented) 1697

3.232.1 Optimal result

Integrand size = 31, antiderivative size = 288

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^2} dx$$

$$= -\frac{2d^3(4c+3d)\operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a^3(c-d)^{9/2}(c+d)^{3/2}f} + \frac{d(2c^3-12c^2d+43cd^2+72d^3)\tan(e+fx)}{15a^3(c-d)^4(c+d)f(c+d \sec(e+fx))}$$

$$+ \frac{\tan(e+fx)}{5(c-d)f(a+a \sec(e+fx))^3(c+d \sec(e+fx))}$$

$$+ \frac{(2c-9d)\tan(e+fx)}{15a(c-d)^2f(a+a \sec(e+fx))^2(c+d \sec(e+fx))}$$

$$+ \frac{(2c^2-12cd+45d^2)\tan(e+fx)}{15(c-d)^3f(a^3+a^3 \sec(e+fx))(c+d \sec(e+fx))}$$

output

```
-2*d^3*(4*c+3*d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a^3/(c-d)^(9/2)/(c+d)^(3/2)/f+1/15*d*(2*c^3-12*c^2*d+43*c*d^2+72*d^3)*tan(f*x+e)/a^3/(c-d)^4/(c+d)/f/(c+d*sec(f*x+e))+1/5*tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))+1/15*(2*c-9*d)*tan(f*x+e)/a/(c-d)^2/f/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))+1/15*(2*c^2-12*c*d+45*d^2)*tan(f*x+e)/(c-d)^3/f/(a^3+a^3*sec(f*x+e))/(c+d*sec(f*x+e))
```


3.232.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.96 (sec) , antiderivative size = 1772, normalized size of antiderivative = 6.15

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c+d\sec(e+fx))^2} dx = \text{Too large to display}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^2),x]`

output

```
((4*c + 3*d)*Cos[e/2 + (f*x)/2]^6*(d + c*Cos[e + f*x])^2*Sec[e + f*x]^5*((
(16*I)*d^3*ArcTan[Sec[(f*x)/2]*(Cos[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*
Sin[2*e]]) - (I*Sin[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]])))*((-
I)*d*Sin[(f*x)/2] + I*c*Sin[e + (f*x)/2]))*Cos[e]/(Sqrt[c^2 - d^2]*f*Sqrt
[Cos[2*e] - I*Sin[2*e]]) + (16*d^3*ArcTan[Sec[(f*x)/2]*(Cos[e]/(Sqrt[c^2 -
d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]]) - (I*Sin[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[
2*e] - I*Sin[2*e]])))*((-I)*d*Sin[(f*x)/2] + I*c*Sin[e + (f*x)/2]))*Sin[e]
/(Sqrt[c^2 - d^2]*f*Sqrt[Cos[2*e] - I*Sin[2*e]])))/((-c + d)^4*(c + d)*(a
+ a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^2) + (Cos[e/2 + (f*x)/2]*(d + c*C
os[e + f*x])*Sec[e/2]*Sec[e]*Sec[e + f*x]^5*(-55*c^5*Sin[(f*x)/2] + 135*c^
4*d*Sin[(f*x)/2] - 20*c^3*d^2*Sin[(f*x)/2] - 810*c^2*d^3*Sin[(f*x)/2] - 45
0*c*d^4*Sin[(f*x)/2] + 150*d^5*Sin[(f*x)/2] + 47*c^5*Sin[(3*f*x)/2] - 137*
c^4*d*Sin[(3*f*x)/2] + 88*c^3*d^2*Sin[(3*f*x)/2] + 812*c^2*d^3*Sin[(3*f*x)
/2] + 690*c*d^4*Sin[(3*f*x)/2] + 75*d^5*Sin[(3*f*x)/2] - 50*c^5*Sin[e - (f
*x)/2] + 130*c^4*d*Sin[e - (f*x)/2] - 10*c^3*d^2*Sin[e - (f*x)/2] - 1030*c
^2*d^3*Sin[e - (f*x)/2] - 990*c*d^4*Sin[e - (f*x)/2] - 150*d^5*Sin[e - (f*
x)/2] + 50*c^5*Sin[e + (f*x)/2] - 130*c^4*d*Sin[e + (f*x)/2] + 10*c^3*d^2*
Sin[e + (f*x)/2] + 1030*c^2*d^3*Sin[e + (f*x)/2] + 765*c*d^4*Sin[e + (f*x)
/2] - 150*d^5*Sin[e + (f*x)/2] - 55*c^5*Sin[2*e + (f*x)/2] + 135*c^4*d*Sin
[2*e + (f*x)/2] - 20*c^3*d^2*Sin[2*e + (f*x)/2] - 810*c^2*d^3*Sin[2*e + ...
```

3.232.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 392, normalized size of antiderivative = 1.36, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 4475, 114, 27, 169, 25, 27, 169, 25, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.232. $\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c+d\sec(e+fx))^2} dx$

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx)+a)^3 (c+d \sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})}{(a \csc(e+fx+\frac{\pi}{2})+a)^3 (c+d \csc(e+fx+\frac{\pi}{2}))^2} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{1}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{7/2} (c+d \sec(e+fx))^2} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 114

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(c+3d-3d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{7/2} (c+d \sec(e+fx))} d \sec(e+fx)}{a^2(c^2-d^2)} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{5/2} (c+d \sec(e+fx))} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{c+3d-3d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{7/2} (c+d \sec(e+fx))} d \sec(e+fx)}{c^2-d^2} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{5/2} (c+d \sec(e+fx))} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 169

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(2c^2-8dc-15d^2+2d(c+6d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx))} d \sec(e+fx)}{5a^3(c-d)} - \frac{(c+6d) \sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 25

$$\frac{a^2 \tan(e+fx) \left(\frac{\int \frac{a^2(2c^2-8dc-15d^2+2d(c+6d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{5/2} (c+d \sec(e+fx))} d \sec(e+fx)}{5a^3(c-d)} - \frac{(c+6d) \sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} + \frac{d \sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)} \right)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}}$$

↓ 27

3.232. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 (c+d \sec(e+fx))^2} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{2c^2 - 8dc - 15d^2 + 2d(c+6d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{5a(c-d)} - \frac{(c+6d)\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} + \frac{d\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 169

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2((c+d)(2c^2-12dc+45d^2)+d(2c^2-10dc-27d^2)\sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(2c^2-10cd-27d^2)\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} - \frac{(c+6d)\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 25

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2((c+d)(2c^2-12dc+45d^2)+d(2c^2-10dc-27d^2)\sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(2c^2-10cd-27d^2)\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} - \frac{(c+6d)\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d)(2c^2-12dc+45d^2)+d(2c^2-10dc-27d^2)\sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a(c-d)} - \frac{(2c^2-10cd-27d^2)\sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} - \frac{(c+6d)\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 169

3.232. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^2} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{15a^2 d^3 (4c+3d)}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}} d \sec(e+fx)}{a^3(c-d)} - \frac{(2c^3-12c^2d+43cd^2+72d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(2c^2-10cd-27d^2)}{3a^2(c-d)(a \sec(e+fx))} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$$

27

$$a^2 \tan(e + fx) \left(\frac{15d^3(4c+3d) \int \frac{1}{\sqrt{a-a \sec(e+fx)} \sqrt{\sec(e+fx)a+a(c+d \sec(e+fx))}} d \sec(e+fx)}{a(c-d)} - \frac{(2c^3-12c^2d+43cd^2+72d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(2c^2-10cd-27d^2)}{3a^2(c-d)(a \sec(e+fx))} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$$

104

$$a^2 \tan(e + fx) \left(\frac{30d^3(4c+3d) \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{a(c-d)} - \frac{(2c^3-12c^2d+43cd^2+72d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(2c^2-10cd-27d^2)}{3a^2(c-d)(a \sec(e+fx))} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx)}$$

218

$$a^2 \tan(e + fx) \left(\frac{30d^3(4c+3d) \arctan\left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a \sec(e+fx)}}\right)}{a^2(c-d)^{3/2} \sqrt{c+d}} - \frac{(2c^3-12c^2d+43cd^2+72d^3) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} - \frac{(2c^2-10cd-27d^2) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

```
input Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^2),x]
```

3.232. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^2} dx$

```
output -((a^2*((d*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c^2 - d^2)*(a + a*Sec[e + f*x])
^(5/2)*(c + d*Sec[e + f*x]))) + (-1/5*((c + 6*d)*Sqrt[a - a*Sec[e + f*x]])/
(a^2*(c - d)*(a + a*Sec[e + f*x])^(5/2)) + (-1/3*((2*c^2 - 10*c*d - 27*d^2
)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*(a + a*Sec[e + f*x])^(3/2)) + ((-
30*d^3*(4*c + 3*d)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c -
d]*Sqrt[a - a*Sec[e + f*x]])])/(a^2*(c - d)^(3/2)*Sqrt[c + d]) - ((2*c^3
- 12*c^2*d + 43*c*d^2 + 72*d^3)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*Sqr
t[a + a*Sec[e + f*x]]))/(3*a*(c - d))/(5*a*(c - d))/(c^2 - d^2))*Tan[e +
f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])
```

3.232.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 114 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)
)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e
- a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1)
- b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x],
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] ||
IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

rule 169 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_))*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_))*((csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.)^(n_)), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.232.4 Maple [A] (verified)

Time = 0.89 (sec) , antiderivative size = 284, normalized size of antiderivative = 0.99

method	result
derivativedivides	$\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 c^2 - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 cd + \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 d^2 - 2c^2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 + \frac{8 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 cd}{(c^2-2cd+d^2)(c-d)^2} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^2}{(c^2-2cd+d^2)(c-d)^2}$
default	$\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 c^2 - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 cd + \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5 d^2 - 2c^2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 + \frac{8 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 cd}{(c^2-2cd+d^2)(c-d)^2} - 2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3 d^2}{(c^2-2cd+d^2)(c-d)^2}$
risch	$2i(7c^5+10c^3d^2e^{3i(fx+e)}-137c^4de^{2i(fx+e)}+106c^3d^2e^{i(fx+e)}-76c^4de^{i(fx+e)}+195cd^4e^{5i(fx+e)}+990cd^4e^{3i(fx+e)}+60d^5e^{i(fx+e)})$

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 1/4/f/a^3*(1/(c^2-2*c*d+d^2)/(c-d)^2*(1/5*tan(1/2*f*x+1/2*e)^5*c^2-2/5*tan(1/2*f*x+1/2*e)^5*c*d+1/5*tan(1/2*f*x+1/2*e)^5*d^2-2/3*c^2*tan(1/2*f*x+1/2*e)^3+8/3*tan(1/2*f*x+1/2*e)^3*c*d-2*tan(1/2*f*x+1/2*e)^3*d^2+tan(1/2*f*x+1/2*e)*c^2-6*tan(1/2*f*x+1/2*e)*c*d+17*tan(1/2*f*x+1/2*e)*d^2)+16*d^3/(c-d)^4*(-1/2*d/(c+d)*tan(1/2*f*x+1/2*e)/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)-1/2*(4*c+3*d)/(c+d)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))))
```

3.232.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 818 vs. 2(271) = 542.

Time = 0.34 (sec) , antiderivative size = 1693, normalized size of antiderivative = 5.88

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^2} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x, algorithm="fracas")
```

```

output [1/30*(15*(4*c*d^4 + 3*d^5 + (4*c^2*d^3 + 3*c*d^4)*cos(f*x + e)^4 + (12*c^
2*d^3 + 13*c*d^4 + 3*d^5)*cos(f*x + e)^3 + 3*(4*c^2*d^3 + 7*c*d^4 + 3*d^5)
*cos(f*x + e)^2 + (4*c^2*d^3 + 15*c*d^4 + 9*d^5)*cos(f*x + e))*sqrt(c^2 -
d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*sqrt(c^2 -
d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2
+ 2*c*d*cos(f*x + e) + d^2)) + 2*(2*c^5*d - 12*c^4*d^2 + 41*c^3*d^3 + 84*
c^2*d^4 - 43*c*d^5 - 72*d^6 + (7*c^6 - 27*c^5*d + 31*c^4*d^2 + 99*c^3*d^3
- 23*c^2*d^4 - 72*c*d^5 - 15*d^6)*cos(f*x + e)^3 + (6*c^6 - 29*c^5*d + 51*
c^4*d^2 + 193*c^3*d^3 + 60*c^2*d^4 - 164*c*d^5 - 117*d^6)*cos(f*x + e)^2 +
(2*c^6 - 6*c^5*d + 5*c^4*d^2 + 147*c^3*d^3 + 164*c^2*d^4 - 141*c*d^5 - 17
1*d^6)*cos(f*x + e))*sin(f*x + e))/((a^3*c^8 - 3*a^3*c^7*d + a^3*c^6*d^2 +
5*a^3*c^5*d^3 - 5*a^3*c^4*d^4 - a^3*c^3*d^5 + 3*a^3*c^2*d^6 - a^3*c*d^7)*
f*cos(f*x + e)^4 + (3*a^3*c^8 - 8*a^3*c^7*d + 16*a^3*c^5*d^3 - 10*a^3*c^4*
d^4 - 8*a^3*c^3*d^5 + 8*a^3*c^2*d^6 - a^3*d^8)*f*cos(f*x + e)^3 + 3*(a^3*c
^8 - 2*a^3*c^7*d - 2*a^3*c^6*d^2 + 6*a^3*c^5*d^3 - 6*a^3*c^3*d^5 + 2*a^3*c
^2*d^6 + 2*a^3*c*d^7 - a^3*d^8)*f*cos(f*x + e)^2 + (a^3*c^8 - 8*a^3*c^6*d
^2 + 8*a^3*c^5*d^3 + 10*a^3*c^4*d^4 - 16*a^3*c^3*d^5 + 8*a^3*c*d^7 - 3*a^3*
d^8)*f*cos(f*x + e) + (a^3*c^7*d - 3*a^3*c^6*d^2 + a^3*c^5*d^3 + 5*a^3*c^4
*d^4 - 5*a^3*c^3*d^5 - a^3*c^2*d^6 + 3*a^3*c*d^7 - a^3*d^8)*f), -1/15*(15*
(4*c*d^4 + 3*d^5 + (4*c^2*d^3 + 3*c*d^4)*cos(f*x + e)^4 + (12*c^2*d^3 + ...

```

3.232.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^2} dx$$

$$= \frac{\int \frac{\sec(e + fx)}{c^2 \sec^3(e + fx) + 3c^2 \sec^2(e + fx) + 3c^2 \sec(e + fx) + c^2 + 2cd \sec^4(e + fx) + 6cd \sec^3(e + fx) + 6cd \sec^2(e + fx) + 2cd \sec(e + fx) + d^2 \sec^5(e + fx) + 3cd \sec^4(e + fx) + 3d^2 \sec^3(e + fx) + 3d^2 \sec^2(e + fx) + 3d^2 \sec(e + fx) + d^2} dx}{a^3}$$

```

input integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e))**2,x)

```

```

output Integral(sec(e + f*x)/(c**2*sec(e + f*x)**3 + 3*c**2*sec(e + f*x)**2 + 3*c
**2*sec(e + f*x) + c**2 + 2*c*d*sec(e + f*x)**4 + 6*c*d*sec(e + f*x)**3 +
6*c*d*sec(e + f*x)**2 + 2*c*d*sec(e + f*x) + d**2*sec(e + f*x)**5 + 3*d**2
*sec(e + f*x)**4 + 3*d**2*sec(e + f*x)**3 + d**2*sec(e + f*x)**2), x)/a**3

```


3.232.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.232.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 918 vs. 2(271) = 542.

Time = 0.39 (sec) , antiderivative size = 918, normalized size of antiderivative = 3.19

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output

```
-1/60*(120*d^4*tan(1/2*f*x + 1/2*e)/((a^3*c^5 - 3*a^3*c^4*d + 2*a^3*c^3*d^2 + 2*a^3*c^2*d^3 - 3*a^3*c*d^4 + a^3*d^5)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)) + 120*(4*c*d^3 + 3*d^4)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((a^3*c^5 - 3*a^3*c^4*d + 2*a^3*c^3*d^2 + 2*a^3*c^2*d^3 - 3*a^3*c*d^4 + a^3*d^5)*sqrt(-c^2 + d^2)) - (3*a^12*c^8*tan(1/2*f*x + 1/2*e)^5 - 24*a^12*c^7*d*tan(1/2*f*x + 1/2*e)^5 + 84*a^12*c^6*d^2*tan(1/2*f*x + 1/2*e)^5 - 168*a^12*c^5*d^3*tan(1/2*f*x + 1/2*e)^5 + 210*a^12*c^4*d^4*tan(1/2*f*x + 1/2*e)^5 - 168*a^12*c^3*d^5*tan(1/2*f*x + 1/2*e)^5 + 84*a^12*c^2*d^6*tan(1/2*f*x + 1/2*e)^5 - 24*a^12*c*d^7*tan(1/2*f*x + 1/2*e)^5 + 3*a^12*d^8*tan(1/2*f*x + 1/2*e)^5 - 10*a^12*c^8*tan(1/2*f*x + 1/2*e)^3 + 100*a^12*c^7*d*tan(1/2*f*x + 1/2*e)^3 - 420*a^12*c^6*d^2*tan(1/2*f*x + 1/2*e)^3 + 980*a^12*c^5*d^3*tan(1/2*f*x + 1/2*e)^3 - 1400*a^12*c^4*d^4*tan(1/2*f*x + 1/2*e)^3 + 1260*a^12*c^3*d^5*tan(1/2*f*x + 1/2*e)^3 - 700*a^12*c^2*d^6*tan(1/2*f*x + 1/2*e)^3 + 220*a^12*c*d^7*tan(1/2*f*x + 1/2*e)^3 - 30*a^12*d^8*tan(1/2*f*x + 1/2*e)^3 + 15*a^12*c^8*tan(1/2*f*x + 1/2*e) - 180*a^12*c^7*d*tan(1/2*f*x + 1/2*e) + 1020*a^12*c^6*d^2*tan(1/2*f*x + 1/2*e) - 3180*a^12*c^5*d^3*tan(1/2*f*x + 1/2*e) + 5850*a^12*c^4*d^4*tan(1/2*f*x + 1/2*e) - 6540*a^12*c^3*d^5*tan(1/2*f*x + 1/2*e) + 4380*a^12*c^2*d^6*tan(1/2*f*x + 1/2*e) - 1620*a^12*c*d^7*tan(1/2*f*x + 1/2*e) + 2...
```

3.232.9 Mupad [B] (verification not implemented)

Time = 13.88 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.61

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^2} dx$$

$$= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{20 a^3 f (c - d)^2} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{2(c^2 - d^2) \left(\frac{1}{a^3 (c-d)^2} - \frac{c^2 - d^2}{2 a^3 (c-d)^4} \right)}{(c-d)^2} - \frac{3}{2 a^3 (c-d)^2} + \frac{(c+d)^2}{4 a^3 (c-d)^4} \right)}{f}$$

$$- \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\frac{1}{3 a^3 (c-d)^2} - \frac{c^2 - d^2}{6 a^3 (c-d)^4} \right)}{f}$$

$$+ \frac{2 d^4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f (c + d) \left(a^3 c^5 - \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (a^3 c^5 - 5 a^3 c^4 d + 10 a^3 c^3 d^2 - 10 a^3 c^2 d^3 + 5 a^3 c d^4 - a^3 d^5) + a^3 d^5 \right)}$$

$$+ \frac{d^3 \operatorname{atan}\left(\frac{11 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^5 - 5i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^4 d + 10i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^3 d^2 - 10i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^2 d^3 + 5i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c d^4 - 11 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) d^5}{\sqrt{c+d} (c-d)^{9/2}} \right)}{a^3 f (c + d)^{3/2} (c - d)^{9/2}}$$

3.232. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 (c+d \sec(e+fx))^2} dx$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c + d/cos(e + f*x))^2),x)`

output `tan(e/2 + (f*x)/2)^5/(20*a^3*f*(c - d)^2) - (tan(e/2 + (f*x)/2)*((2*(c^2 - d^2)*(1/(a^3*(c - d)^2) - (c^2 - d^2)/(2*a^3*(c - d)^4)))/(c - d)^2 - 3/(2*a^3*(c - d)^2) + (c + d)^2/(4*a^3*(c - d)^4))/f - (tan(e/2 + (f*x)/2)^3*(1/(3*a^3*(c - d)^2) - (c^2 - d^2)/(6*a^3*(c - d)^4))/f + (2*d^4*tan(e/2 + (f*x)/2))/(f*(c + d)*(a^3*c^5 - tan(e/2 + (f*x)/2)^2*(a^3*c^5 - a^3*d^5 + 5*a^3*c*d^4 - 5*a^3*c^4*d - 10*a^3*c^2*d^3 + 10*a^3*c^3*d^2) + a^3*d^5 - 3*a^3*c*d^4 - 3*a^3*c^4*d + 2*a^3*c^2*d^3 + 2*a^3*c^3*d^2)) + (d^3*atan((c^5*tan(e/2 + (f*x)/2)*1i - d^5*tan(e/2 + (f*x)/2)*1i + c*d^4*tan(e/2 + (f*x)/2)*5i - c^4*d*tan(e/2 + (f*x)/2)*5i - c^2*d^3*tan(e/2 + (f*x)/2)*10i + c^3*d^2*tan(e/2 + (f*x)/2)*10i)/((c + d)^(1/2)*(c - d)^(9/2)))*(4*c + 3*d)*2i)/(a^3*f*(c + d)^(3/2)*(c - d)^(9/2))`

3.233
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$$

3.233.1 Optimal result 1699
 3.233.2 Mathematica [C] (warning: unable to verify) 1700
 3.233.3 Rubi [A] (verified) 1701
 3.233.4 Maple [A] (verified) 1707
 3.233.5 Fricas [B] (verification not implemented) 1708
 3.233.6 Sympy [F] 1709
 3.233.7 Maxima [F(-2)] 1710
 3.233.8 Giac [B] (verification not implemented) 1710
 3.233.9 Mupad [B] (verification not implemented) 1711

3.233.1 Optimal result

Integrand size = 31, antiderivative size = 368

$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$$

$$= -\frac{d^3(20c^2+30cd+13d^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{a^3(c-d)^{11/2}(c+d)^{5/2}f}$$

$$+ \frac{d(4c^3-30c^2d+146cd^2+195d^3) \tan(e+fx)}{30a^3(c-d)^4(c+d)f(c+d \sec(e+fx))^2}$$

$$+ \frac{\tan(e+fx)}{5(c-d)f(a+a \sec(e+fx))^3(c+d \sec(e+fx))^2}$$

$$+ \frac{(2c-11d) \tan(e+fx)}{15a(c-d)^2f(a+a \sec(e+fx))^2(c+d \sec(e+fx))^2}$$

$$+ \frac{(2c^2-15cd+76d^2) \tan(e+fx)}{15(c-d)^3f(a^3+a^3 \sec(e+fx))(c+d \sec(e+fx))^2}$$

$$+ \frac{d(4c^4-30c^3d+142c^2d^2+525cd^3+304d^4) \tan(e+fx)}{30a^3(c-d)^5(c+d)^2f(c+d \sec(e+fx))}$$

output

```
-d^3*(20*c^2+30*c*d+13*d^2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/a^3/(c-d)^(11/2)/(c+d)^(5/2)/f+1/30*d*(4*c^3-30*c^2*d+146*c*d^2+195*d^3)*tan(f*x+e)/a^3/(c-d)^4/(c+d)/f/(c+d*sec(f*x+e))^2+1/5*tan(f*x+e)/(c-d)/f/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^2+1/5*(2*c-11*d)*tan(f*x+e)/a/(c-d)^2/f/(a+a*sec(f*x+e))^2/(c+d*sec(f*x+e))^2+1/15*(2*c^2-15*c*d+76*d^2)*tan(f*x+e)/(c-d)^3/f/(a^3+a^3*sec(f*x+e))/(c+d*sec(f*x+e))^2+1/30*d*(4*c^4-30*c^3*d+142*c^2*d^2+525*c*d^3+304*d^4)*tan(f*x+e)/a^3/(c-d)^5/(c+d)^2/f/(c+d*sec(f*x+e))
```

3.233.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 9.05 (sec) , antiderivative size = 1096, normalized size of antiderivative = 2.98

$$\int \frac{\sec(e+fx)}{(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3} dx$$

$$= \frac{4\cos^4\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))^3\sec\left(\frac{e}{2}\right)\sec^6(e+fx)(-8c\sin\left(\frac{e}{2}\right)+23d\sin\left(\frac{e}{2}\right))}{15(-c+d)^4f(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3}$$

$$+ \frac{(20c^2+30cd+13d^2)\cos^6\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))^3\sec^6(e+fx)\left(-\frac{8id^3\arctan\left(\sec\left(\frac{fx}{2}\right)\right)\left(\frac{\cos(e)}{\sqrt{c^2-d^2}\sqrt{\cos(2e)}}\right)}{(-c+d)}\right)}{(-c+d)}$$

$$- \frac{2\cos\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))^3\sec\left(\frac{e}{2}\right)\sec^6(e+fx)\sin\left(\frac{fx}{2}\right)}{5(-c+d)^3f(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3}$$

$$+ \frac{4\cos^3\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))^3\sec\left(\frac{e}{2}\right)\sec^6(e+fx)(-8c\sin\left(\frac{fx}{2}\right)+23d\sin\left(\frac{fx}{2}\right))}{15(-c+d)^4f(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3}$$

$$- \frac{8\cos^5\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))^3\sec\left(\frac{e}{2}\right)\sec^6(e+fx)(7c^2\sin\left(\frac{fx}{2}\right)-44cd\sin\left(\frac{fx}{2}\right)+127d^2\sin\left(\frac{fx}{2}\right))}{15(-c+d)^5f(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3}$$

$$+ \frac{4\cos^6\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))\sec(e)\sec^6(e+fx)(d^6\sin(e)-cd^5\sin(fx))}{c^2(-c+d)^4(c+d)f(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3}$$

$$- \frac{4\cos^6\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))^2\sec(e)\sec^6(e+fx)(-11c^2d^5\sin(e)-6cd^6\sin(e)+2d^7\sin(e)+10c^2d^5\sin(fx))}{c^2(-c+d)^5(c+d)^2f(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3}$$

$$- \frac{2\cos^2\left(\frac{e}{2}+\frac{fx}{2}\right)(d+c\cos(e+fx))^3\sec^6(e+fx)\tan\left(\frac{e}{2}\right)}{5(-c+d)^3f(a+a\sec(e+fx))^3(c+d\sec(e+fx))^3}$$

input `Integrate[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^3),x]`

output

```
(4*cos[e/2 + (f*x)/2]^4*(d + c*cos[e + f*x])^3*sec[e/2]*sec[e + f*x]^6*(-8
*c*sin[e/2] + 23*d*sin[e/2]))/(15*(-c + d)^4*f*(a + a*sec[e + f*x])^3*(c +
d*sec[e + f*x])^3) + ((20*c^2 + 30*c*d + 13*d^2)*cos[e/2 + (f*x)/2]^6*(d
+ c*cos[e + f*x])^3*sec[e + f*x]^6*((-8*I)*d^3*ArcTan[Sec[(f*x)/2]*(Cos[e
]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]])] - (I*Sin[e]/(Sqrt[c^2 - d
^2]*Sqrt[Cos[2*e] - I*Sin[2*e]])))*((-I)*d*sin[(f*x)/2] + I*c*sin[e + (f*x)
/2]))*cos[e]/(Sqrt[c^2 - d^2]*f*Sqrt[Cos[2*e] - I*Sin[2*e]]) - (8*d^3*Arc
Tan[Sec[(f*x)/2]*(Cos[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]])] - (
I*Sin[e]/(Sqrt[c^2 - d^2]*Sqrt[Cos[2*e] - I*Sin[2*e]])))*((-I)*d*sin[(f*x)
/2] + I*c*sin[e + (f*x)/2]))*sin[e]/(Sqrt[c^2 - d^2]*f*Sqrt[Cos[2*e] - I*
Sin[2*e]])))/((-c + d)^5*(c + d)^2*(a + a*sec[e + f*x])^3*(c + d*sec[e + f
*x])^3) - (2*cos[e/2 + (f*x)/2]*(d + c*cos[e + f*x])^3*sec[e/2]*sec[e + f*
x]^6*sin[(f*x)/2])/(5*(-c + d)^3*f*(a + a*sec[e + f*x])^3*(c + d*sec[e + f
*x])^3) + (4*cos[e/2 + (f*x)/2]^3*(d + c*cos[e + f*x])^3*sec[e/2]*sec[e +
f*x]^6*(-8*c*sin[(f*x)/2] + 23*d*sin[(f*x)/2]))/(15*(-c + d)^4*f*(a + a*se
c[e + f*x])^3*(c + d*sec[e + f*x])^3) - (8*cos[e/2 + (f*x)/2]^5*(d + c*cos
[e + f*x])^3*sec[e/2]*sec[e + f*x]^6*(7*c^2*sin[(f*x)/2] - 44*c*d*sin[(f*x)
]/2) + 127*d^2*sin[(f*x)/2))/(15*(-c + d)^5*f*(a + a*sec[e + f*x])^3*(c +
d*sec[e + f*x])^3) + (4*cos[e/2 + (f*x)/2]^6*(d + c*cos[e + f*x])*sec[e]*
sec[e + f*x]^6*(d^6*sin[e] - c*d^5*sin[f*x]))/(c^2*(-c + d)^4*(c + d)*f...
```

3.233.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 508, normalized size of antiderivative = 1.38, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 4475, 114, 27, 168, 27, 169, 25, 27, 169, 25, 27, 169, 27, 104, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(a \sec(e+fx) + a)^3 (c + d \sec(e+fx))^3} dx$$

↓ 3042

$$\int \frac{\csc(e+fx + \frac{\pi}{2})}{(a \csc(e+fx + \frac{\pi}{2}) + a)^3 (c + d \csc(e+fx + \frac{\pi}{2}))^3} dx$$

↓ 4475

$$\frac{a^2 \tan(e+fx) \int \frac{1}{\sqrt{a-a \sec(e+fx)} (\sec(e+fx)a+a)^{7/2} (c+d \sec(e+fx))^3} d \sec(e+fx)}{f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx) + a}}$$

3.233. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 (c+d \sec(e+fx))^3} dx$

↓ 114

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(2c+3d-4d \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{7/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2a^2(c^2-d^2)} + \frac{d\sqrt{a-a \sec(e+fx)}}{2a^2(c^2-d^2)(a \sec(e+fx)+a)^{5/2}(c+d \sec(e+fx))^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{2c+3d-4d \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{7/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{2(c^2-d^2)} + \frac{d\sqrt{a-a \sec(e+fx)}}{2a^2(c^2-d^2)(a \sec(e+fx)+a)^{5/2}(c+d \sec(e+fx))^2} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 168

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2(2c^2+21dc+13d^2-9d(2c+d) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{7/2}(c+d \sec(e+fx))^2} d \sec(e+fx)}{a^2(c^2-d^2)} + \frac{3d(2c+d)\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{5/2}(c+d \sec(e+fx))} + \frac{1}{2a^2(c^2-d^2)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{2c^2+21dc+13d^2-9d(2c+d) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{7/2}(c+d \sec(e+fx))} d \sec(e+fx)}{c^2-d^2} + \frac{3d(2c+d)\sqrt{a-a \sec(e+fx)}}{a^2(c^2-d^2)(a \sec(e+fx)+a)^{5/2}(c+d \sec(e+fx))} + \frac{1}{2a^2(c^2-d^2)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 169

$$a^2 \tan(e + fx) \left(\frac{\int -\frac{a^2((2c+5d)(2c^2-16dc-13d^2)+2d(2c^2+39dc+22d^2) \sec(e+fx))}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{5a^3(c-d)} - \frac{(2c^2+39cd+22d^2)\sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} + \frac{1}{a^2(c^2-d^2)} \right)$$

$$f \sqrt{a - a \sec(e + fx)} \sqrt{a \sec(e + fx) + a}$$

↓ 25

3.233. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2((2c+5d)(2c^2-16dc-13d^2)+2d(2c^2+39dc+22d^2)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{5a^3(c-d)} - \frac{(2c^2+39cd+22d^2) \sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} + \frac{3d}{a^2(c^2-d^2)} \right) \frac{c^2-d^2}{2(c^2-d^2)} + f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}$$

27

$$a^2 \tan(e + fx) \left(\frac{\int \frac{(2c+5d)(2c^2-16dc-13d^2)+2d(2c^2+39dc+22d^2) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{5/2}(c+d \sec(e+fx))} d \sec(e+fx)}{5a(c-d)} - \frac{(2c^2+39cd+22d^2) \sqrt{a-a \sec(e+fx)}}{5a^2(c-d)(a \sec(e+fx)+a)^{5/2}} + \frac{3d(2c+d)}{a^2(c^2-d^2)} \right) \frac{c^2-d^2}{2(c^2-d^2)} + f \sqrt{a-a \sec(e+fx)} \sqrt{a \sec(e+fx)+a}$$

169

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2((c+d)(4c^3-30dc^2+146d^2c+195d^3)+d(4c^3-26dc^2-184d^2c-109d^3)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(4c^3-26c^2d-184cd^2-109d^3) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} + \frac{3d(2c+d)}{a^2(c^2-d^2)} \right) \frac{c^2-d^2}{2(c^2-d^2)} + f \sqrt{a-a \sec(e+fx)}$$

25

$$a^2 \tan(e + fx) \left(\frac{\int \frac{a^2((c+d)(4c^3-30dc^2+146d^2c+195d^3)+d(4c^3-26dc^2-184d^2c-109d^3)) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a^3(c-d)} - \frac{(4c^3-26c^2d-184cd^2-109d^3) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} + \frac{3d(2c+d)}{a^2(c^2-d^2)} \right) \frac{c^2-d^2}{2(c^2-d^2)} + f \sqrt{a-a \sec(e+fx)}$$

27

3.233. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{\int \frac{(c+d)(4c^3 - 30dc^2 + 146d^2c + 195d^3) + d(4c^3 - 26dc^2 - 184d^2c - 109d^3) \sec(e+fx)}{\sqrt{a-a \sec(e+fx)}(\sec(e+fx)a+a)^{3/2}(c+d \sec(e+fx))} d \sec(e+fx)}{3a(c-d)} - \frac{(4c^3 - 26c^2d - 184cd^2 - 109d^3) \sqrt{a-a \sec(e+fx)}}{3a^2(c-d)(a \sec(e+fx)+a)^{3/2}} \right) \frac{d \sec(e+fx)}{5a(c-d)} - \frac{c^2 - d^2}{2(c^2 - d^2)}$$

$f \sqrt{a - a \sec(e + fx)}$

↓ 169

$$a^2 \tan(e + fx) \left(- \frac{\int \frac{15a^2d^3(20c^2 + 30dc + 13d^2)}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{a^3(c-d)} - \frac{(4c^4 - 30c^3d + 142c^2d^2 + 525cd^3 + 304d^4) \sqrt{a-a \sec(e+fx)}}{a^2(c-d)\sqrt{a \sec(e+fx)+a}} - (4c^3 - 30c^2d - 184cd^2 - 109d^3) \sqrt{a-a \sec(e+fx)}}{3a(c-d)} - \frac{c^2 - d^2}{5a(c-d)} - \frac{c^2 - d^2}{2(c^2 - d^2)} \right)$$

$f \sqrt{a - a \sec(e + fx)}$

↓ 27

$$a^2 \tan(e + fx) \left(- \frac{15d^3(20c^2 + 30cd + 13d^2) \int \frac{1}{\sqrt{a-a \sec(e+fx)}\sqrt{\sec(e+fx)a+a}(c+d \sec(e+fx))} d \sec(e+fx)}{a(c-d)} - \frac{(4c^4 - 30c^3d + 142c^2d^2 + 525cd^3 + 304d^4) \sqrt{a-a \sec(e+fx)}}{a^2(c-d)\sqrt{a \sec(e+fx)+a}} - (4c^3 - 30c^2d - 184cd^2 - 109d^3) \sqrt{a-a \sec(e+fx)}}{3a(c-d)} - \frac{c^2 - d^2}{5a(c-d)} - \frac{c^2 - d^2}{2(c^2 - d^2)} \right)$$

↓ 104

3.233. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$

$$a^2 \tan(e + fx) \left(\frac{30d^3(20c^2 + 30cd + 13d^2) \int \frac{1}{a(c-d) + \frac{a(c+d)(\sec(e+fx)a+a)}{a-a \sec(e+fx)}} d \frac{\sqrt{\sec(e+fx)a+a}}{\sqrt{a-a \sec(e+fx)}}}{a(c-d)} - \frac{(4c^4 - 30c^3d + 142c^2d^2 + 525cd^3 + 304d^4) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}} \right)$$

218

$$a^2 \tan(e + fx) \left(\frac{30d^3(20c^2 + 30cd + 13d^2) \arctan\left(\frac{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}}{\sqrt{c-d} \sqrt{a-a \sec(e+fx)}}\right) - \frac{(4c^4 - 30c^3d + 142c^2d^2 + 525cd^3 + 304d^4) \sqrt{a-a \sec(e+fx)}}{a^2(c-d) \sqrt{a \sec(e+fx)+a}}}{a^2(c-d)^{3/2} \sqrt{c+d}} - \frac{(4c^3 - 26c^2d + 184cd^2 - 109d^3) \sqrt{a-a \sec(e+fx)}}{3a(c-d)} \right)$$

$f \sqrt{a - a \sec(e+fx)}$

```
input Int[Sec[e + f*x]/((a + a*Sec[e + f*x])^3*(c + d*Sec[e + f*x])^3),x]
```

```
output -((a^2*((d*Sqrt[a - a*Sec[e + f*x]])/(2*a^2*(c^2 - d^2)*(a + a*Sec[e + f*x])^(5/2)*(c + d*Sec[e + f*x])^2) + ((3*d*(2*c + d)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c^2 - d^2)*(a + a*Sec[e + f*x])^(5/2)*(c + d*Sec[e + f*x]))) + (-1/5*((2*c^2 + 39*c*d + 22*d^2)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*(a + a*Sec[e + f*x])^(5/2)) + (-1/3*((4*c^3 - 26*c^2*d - 184*c*d^2 - 109*d^3)*Sqrt[a - a*Sec[e + f*x]])/(a^2*(c - d)*(a + a*Sec[e + f*x])^(3/2)) + ((-30*d^3*(20*c^2 + 30*c*d + 13*d^2)*ArcTan[(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])/(Sqrt[c - d]*Sqrt[a - a*Sec[e + f*x]])])/(a^2*(c - d)^(3/2)*Sqrt[c + d]) - ((4*c^4 - 30*c^3*d + 142*c^2*d^2 + 525*c*d^3 + 304*d^4)*Sqrt[a - a*Sec[e + f*x]]/(a^2*(c - d)*Sqrt[a + a*Sec[e + f*x]]))/(3*a*(c - d)))/(5*a*(c - d)))/(c^2 - d^2)/(2*(c^2 - d^2))*Tan[e + f*x])/(f*Sqrt[a - a*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]]))
```

3.233. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$

3.233.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 114 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`
- rule 168 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]`
- rule 169 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4475 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(d_) + (c_))^(n_), x_Symbol] := Simp[a^2*g*(Cot[e + f*x]/(f*Sqrt[a + b*Csc[e + f*x]]*Sqrt[a - b*Csc[e + f*x]])) Subst[Int[(g*x)^(p - 1)*(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Csc[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && (EqQ[p, 1] || IntegerQ[m - 1/2])`

3.233.4 Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 365, normalized size of antiderivative = 0.99

method	result
derivativedivides	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^2 - 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 cd + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 d^2 - 2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 cd - 8 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^3 - 3c^2d + 3cd^2 - d^3)(c^2 - 2cd + d^2)}$
default	$\frac{\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 c^2 - 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 cd + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 d^2 - 2c^2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 cd - 8 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 d^2 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^3 - 3c^2d + 3cd^2 - d^3)(c^2 - 2cd + d^2)}$
risch	Expression too large to display

input `int(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

3.233.
$$\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3(c+d \sec(e+fx))^3} dx$$

output $\frac{1}{4} f/a^3 \left(\frac{1}{(c^3 - 3c^2d + 3cd^2 - d^3)} \frac{1}{(c^2 - 2cd + d^2)} \left(\frac{1}{5} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 c^2 - \frac{2}{5} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 cd + \frac{1}{5} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 d^2 - \frac{2}{3} c^2 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + \frac{10}{3} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 cd - \frac{8}{3} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 d^2 + \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) c^2 - 8 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) cd + 31 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) d^2 \right) + 16d^3 / (c-d)^5 \left(\frac{-1/4 d (10c^2 - 3cd - 7d^2)}{(c^2 + 2cd + d^2)} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^3 + \frac{5/4 d (2c+d)}{(c+d)} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) \right) / \left(\frac{\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 c - \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 d - c - d}{(c+d)(c-d)} \right)^2 - \frac{1}{4} \frac{(20c^2 + 30cd + 13d^2)}{(c^2 + 2cd + d^2)} \frac{1}{((c+d)(c-d))^{1/2}} \operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{((c+d)(c-d))^{1/2}}\right) \right)$

3.233.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1310 vs. 2(349) = 698.

Time = 0.39 (sec) , antiderivative size = 2677, normalized size of antiderivative = 7.27

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^3} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x, algorithm="fricas")`

output `[-1/60*(15*(20*c^2*d^5 + 30*c*d^6 + 13*d^7 + (20*c^4*d^3 + 30*c^3*d^4 + 13*c^2*d^5)*cos(f*x + e)^5 + (60*c^4*d^3 + 130*c^3*d^4 + 99*c^2*d^5 + 26*c*d^6)*cos(f*x + e)^4 + (60*c^4*d^3 + 210*c^3*d^4 + 239*c^2*d^5 + 108*c*d^6 + 13*d^7)*cos(f*x + e)^3 + (20*c^4*d^3 + 150*c^3*d^4 + 253*c^2*d^5 + 168*c*d^6 + 39*d^7)*cos(f*x + e)^2 + (40*c^3*d^4 + 120*c^2*d^5 + 116*c*d^6 + 39*d^7)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) - 2*(4*c^6*d^2 - 30*c^5*d^3 + 138*c^4*d^4 + 555*c^3*d^5 + 162*c^2*d^6 - 525*c*d^7 - 304*d^8 + (14*c^8 - 60*c^7*d + 78*c^6*d^2 + 480*c^5*d^3 + 312*c^4*d^4 - 330*c^3*d^5 - 419*c^2*d^6 - 90*c*d^7 + 15*d^8)*cos(f*x + e)^4 + (12*c^8 - 62*c^7*d + 114*c^6*d^2 + 1056*c^5*d^3 + 1626*c^4*d^4 - 81*c^3*d^5 - 1707*c^2*d^6 - 913*c*d^7 - 45*d^8)*cos(f*x + e)^3 + (4*c^8 - 6*c^7*d - 28*c^6*d^2 + 828*c^5*d^3 + 2400*c^4*d^4 + 1197*c^3*d^5 - 1897*c^2*d^6 - 2019*c*d^7 - 479*d^8)*cos(f*x + e)^2 + (8*c^7*d - 48*c^6*d^2 + 186*c^5*d^3 + 1224*c^4*d^4 + 1539*c^3*d^5 - 459*c^2*d^6 - 1733*c*d^7 - 717*d^8)*cos(f*x + e))*sin(f*x + e))/((a^3*c^11 - 3*a^3*c^10*d + 8*a^3*c^8*d^3 - 6*a^3*c^7*d^4 - 6*a^3*c^6*d^5 + 8*a^3*c^5*d^6 - 3*a^3*c^3*d^8 + a^3*c^2*d^9)*f*cos(f*x + e)^5 + (3*a^3*c^11 - 7*a^3*c^10*d - 6*a^3*c^9*d^2 + 24*a^3*c^8*d^3 - 2*a^3*c^7*d^4 - 30*a^3*c^6*d^5 + 12*a^3*c^5*d^6 + 16*a^3*c^4*d^7 - 9*a^3*c^3*d^8 - 3*...`

3.233.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^3} dx$$

$$= \int \frac{\sec(e+fx)}{c^3 \sec^3(e+fx) + 3c^3 \sec^2(e+fx) + 3c^3 \sec(e+fx) + c^3 + 3c^2 d \sec^4(e+fx) + 9c^2 d \sec^3(e+fx) + 9c^2 d \sec^2(e+fx) + 3c^2 d \sec(e+fx) + 3cd^2 \sec^5(e+fx) + 3cd^2 \sec^4(e+fx) + 3cd^2 \sec^3(e+fx) + 3cd^2 \sec^2(e+fx) + 3cd^2 \sec(e+fx) + d^3} a^3$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**3/(c+d*sec(f*x+e))**3,x)`

output `Integral(sec(e + f*x)/(c**3*sec(e + f*x)**3 + 3*c**3*sec(e + f*x)**2 + 3*c**3*sec(e + f*x) + c**3 + 3*c**2*d*sec(e + f*x)**4 + 9*c**2*d*sec(e + f*x)**3 + 9*c**2*d*sec(e + f*x)**2 + 3*c**2*d*sec(e + f*x) + 3*c*d**2*sec(e + f*x)**5 + 9*c*d**2*sec(e + f*x)**4 + 9*c*d**2*sec(e + f*x)**3 + 3*c*d**2*sec(e + f*x)**2 + d**3*sec(e + f*x)**6 + 3*d**3*sec(e + f*x)**5 + 3*d**3*sec(e + f*x)**4 + d**3*sec(e + f*x)**3), x)/a**3`

3.233.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.233.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1369 vs. 2(349) = 698.

Time = 0.47 (sec) , antiderivative size = 1369, normalized size of antiderivative = 3.72

$$\int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^3} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^3/(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output

```

-1/60*(60*(20*c^2*d^3 + 30*c*d^4 + 13*d^5)*(pi*floor(1/2*(f*x + e)/pi + 1/
2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2
*e))/sqrt(-c^2 + d^2)))/((a^3*c^7 - 3*a^3*c^6*d + a^3*c^5*d^2 + 5*a^3*c^4*
d^3 - 5*a^3*c^3*d^4 - a^3*c^2*d^5 + 3*a^3*c*d^6 - a^3*d^7)*sqrt(-c^2 + d^2
)) - (3*a^12*c^12*tan(1/2*f*x + 1/2*e)^5 - 36*a^12*c^11*d*tan(1/2*f*x + 1/
2*e)^5 + 198*a^12*c^10*d^2*tan(1/2*f*x + 1/2*e)^5 - 660*a^12*c^9*d^3*tan(1
/2*f*x + 1/2*e)^5 + 1485*a^12*c^8*d^4*tan(1/2*f*x + 1/2*e)^5 - 2376*a^12*c
^7*d^5*tan(1/2*f*x + 1/2*e)^5 + 2772*a^12*c^6*d^6*tan(1/2*f*x + 1/2*e)^5 -
2376*a^12*c^5*d^7*tan(1/2*f*x + 1/2*e)^5 + 1485*a^12*c^4*d^8*tan(1/2*f*x
+ 1/2*e)^5 - 660*a^12*c^3*d^9*tan(1/2*f*x + 1/2*e)^5 + 198*a^12*c^2*d^10*t
an(1/2*f*x + 1/2*e)^5 - 36*a^12*c*d^11*tan(1/2*f*x + 1/2*e)^5 + 3*a^12*d^1
2*tan(1/2*f*x + 1/2*e)^5 - 10*a^12*c^12*tan(1/2*f*x + 1/2*e)^3 + 150*a^12*
c^11*d*tan(1/2*f*x + 1/2*e)^3 - 990*a^12*c^10*d^2*tan(1/2*f*x + 1/2*e)^3 +
3850*a^12*c^9*d^3*tan(1/2*f*x + 1/2*e)^3 - 9900*a^12*c^8*d^4*tan(1/2*f*x
+ 1/2*e)^3 + 17820*a^12*c^7*d^5*tan(1/2*f*x + 1/2*e)^3 - 23100*a^12*c^6*d^
6*tan(1/2*f*x + 1/2*e)^3 + 21780*a^12*c^5*d^7*tan(1/2*f*x + 1/2*e)^3 - 148
50*a^12*c^4*d^8*tan(1/2*f*x + 1/2*e)^3 + 7150*a^12*c^3*d^9*tan(1/2*f*x + 1
/2*e)^3 - 2310*a^12*c^2*d^10*tan(1/2*f*x + 1/2*e)^3 + 450*a^12*c*d^11*tan(
1/2*f*x + 1/2*e)^3 - 40*a^12*d^12*tan(1/2*f*x + 1/2*e)^3 + 15*a^12*c^12*ta
n(1/2*f*x + 1/2*e) - 270*a^12*c^11*d*tan(1/2*f*x + 1/2*e) + 2340*a^12*c...

```

3.233.9 Mupad [B] (verification not implemented)

Time = 14.09 (sec) , antiderivative size = 655, normalized size of antiderivative = 1.78

$$\begin{aligned}
& \int \frac{\sec(e + fx)}{(a + a \sec(e + fx))^3 (c + d \sec(e + fx))^3} dx \\
&= \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5}{20 a^3 f (c - d)^3} - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(\frac{3(c+d)^2}{4 a^3 (c-d)^5} - \frac{5}{2 a^3 (c-d)^3} + \frac{3(c+d) \left(\frac{5}{4 a^3 (c-d)^3} - \frac{3(c+d)}{4 a^3 (c-d)^4} \right)}{c-d} \right)}{f} \\
&\quad - \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 \left(\frac{5}{12 a^3 (c-d)^3} - \frac{c+d}{4 a^3 (c-d)^4} \right)}{f} \\
&\quad - \frac{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 (2 a^3 c^7 - 10 a^3 c^6 d + 18 a^3 c^5 d^2 - 10 a^3 c^4 d^3 - 10 a^3 c^3 d^4 + 18 a^3 c^2 d^5 - 10 a^3 c d^6 + \right.}{a^3 f (c + d)^{5/2} (c - d)^{11/2}} \\
&\quad \left. d^3 \operatorname{atan}\left(\frac{11 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^6 - 6i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^5 d + 15i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^4 d^2 - 20i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^3 d^3 + 15i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c^2 d^4 - 6i \tan\left(\frac{e}{2} + \frac{fx}{2}\right) c}{\sqrt{c+d} (c-d)^{11/2}} \right)}{a^3 f (c + d)^{5/2} (c - d)^{11/2}} \\
&\quad + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{a^3 f (c + d)^{5/2} (c - d)^{11/2}}
\end{aligned}$$

3.233. $\int \frac{\sec(e+fx)}{(a+a \sec(e+fx))^3 (c+d \sec(e+fx))^3} dx$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^3*(c + d/cos(e + f*x))^3),x)`

output `tan(e/2 + (f*x)/2)^5/(20*a^3*f*(c - d)^3) - (tan(e/2 + (f*x)/2)*((3*(c + d)^2)/(4*a^3*(c - d)^5) - 5/(2*a^3*(c - d)^3) + (3*(c + d)*(5/(4*a^3*(c - d)^3) - (3*(c + d))/(4*a^3*(c - d)^4)))/(c - d))/f - (tan(e/2 + (f*x)/2)^3*(5/(12*a^3*(c - d)^3) - (c + d)/(4*a^3*(c - d)^4))/f - ((tan(e/2 + (f*x)/2)^3*(3*c*d^5 + 7*d^6 - 10*c^2*d^4))/(c + d)^2 + (5*tan(e/2 + (f*x)/2)*(2*c*d^4 + d^5))/(c + d))/(f*(tan(e/2 + (f*x)/2)^2*(2*a^3*c^7 + 2*a^3*d^7 - 10*a^3*c*d^6 - 10*a^3*c^6*d + 18*a^3*c^2*d^5 - 10*a^3*c^3*d^4 - 10*a^3*c^4*d^3 + 18*a^3*c^5*d^2) - tan(e/2 + (f*x)/2)^4*(a^3*c^7 - a^3*d^7 + 7*a^3*c*d^6 - 7*a^3*c^6*d - 21*a^3*c^2*d^5 + 35*a^3*c^3*d^4 - 35*a^3*c^4*d^3 + 21*a^3*c^5*d^2) - a^3*c^7 + a^3*d^7 - 3*a^3*c*d^6 + 3*a^3*c^6*d + a^3*c^2*d^5 + 5*a^3*c^3*d^4 - 5*a^3*c^4*d^3 - a^3*c^5*d^2)) + (d^3*atan((c^6*tan(e/2 + (f*x)/2)*1i + d^6*tan(e/2 + (f*x)/2)*1i - c*d^5*tan(e/2 + (f*x)/2)*6i - c^5*d*tan(e/2 + (f*x)/2)*6i + c^2*d^4*tan(e/2 + (f*x)/2)*15i - c^3*d^3*tan(e/2 + (f*x)/2)*20i + c^4*d^2*tan(e/2 + (f*x)/2)*15i)/((c + d)^(1/2)*(c - d)^(11/2)))*(30*c*d + 20*c^2 + 13*d^2)*1i)/(a^3*f*(c + d)^(5/2)*(c - d)^(11/2))`

3.234
$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$$

3.234.1 Optimal result 1713
 3.234.2 Mathematica [A] (verified) 1713
 3.234.3 Rubi [A] (verified) 1714
 3.234.4 Maple [B] (warning: unable to verify) 1715
 3.234.5 Fricas [B] (verification not implemented) 1716
 3.234.6 Sympy [F] 1716
 3.234.7 Maxima [F] 1717
 3.234.8 Giac [F] 1717
 3.234.9 Mupad [F(-1)] 1717

3.234.1 Optimal result

Integrand size = 35, antiderivative size = 61

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \frac{2\sqrt{a}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{d}f}$$

output `2*arctanh(a^(1/2)*d^(1/2)*tan(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2))*a^(1/2)/f/d^(1/2)`

3.234.2 Mathematica [A] (verified)

Time = 0.72 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.67

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \frac{\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{d}\sin(\frac{1}{2}(e+fx))}{\sqrt{d+c\cos(e+fx)}}\right)\sqrt{d+c\cos(e+fx)}\sec\left(\frac{1}{2}(e+fx)\right)\sqrt{a(1+\sec(e+fx))}}{\sqrt{d}f\sqrt{c+d\sec(e+fx)}}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/Sqrt[c + d*Sec[e + f*x]],x]`

output $(\text{Sqrt}[2]*\text{ArcTanh}[(\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sin}[e + f*x])/2])/(\text{Sqrt}[d + c*\text{Cos}[e + f*x]])*\text{Sqrt}[d + c*\text{Cos}[e + f*x]]*\text{Sec}[(e + f*x)/2]*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])]/(\text{Sqrt}[d]*f*\text{Sqrt}[c + d*\text{Sec}[e + f*x]])$

3.234.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.086$, Rules used = {3042, 4468, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a\sec(e+fx)+a}}{\sqrt{c+d\sec(e+fx)}} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}}{\sqrt{c+d\csc(e+fx+\frac{\pi}{2})}} dx$$

↓ 4468

$$\frac{2a \int \frac{1}{1-\frac{ad \tan^2(e+fx)}{(\sec(e+fx)a+a)(c+d\sec(e+fx))}} d\left(-\frac{\tan(e+fx)}{\sqrt{\sec(e+fx)a+a}\sqrt{c+d\sec(e+fx)}}\right)}{f}$$

↓ 219

$$\frac{2\sqrt{a}\text{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{d}f}$$

input $\text{Int}[(\text{Sec}[e + f*x]*\text{Sqrt}[a + a*\text{Sec}[e + f*x]])/(\text{Sqrt}[c + d*\text{Sec}[e + f*x]]),x]$

output $(2*\text{Sqrt}[a]*\text{ArcTanh}[(\text{Sqrt}[a]*\text{Sqrt}[d]*\text{Tan}[e + f*x])/(\text{Sqrt}[a + a*\text{Sec}[e + f*x]])*\text{Sqrt}[c + d*\text{Sec}[e + f*x]])/(\text{Sqrt}[d]*f)$

3.234.3.1 Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4468 Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/Sq
rt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)], x_Symbol] :> Simp[-2*(b/f) Subs
t[Int[1/(1 - b*d*x^2), x], x, Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c
+ d*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.234.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 277 vs. 2(49) = 98.

Time = 5.51 (sec) , antiderivative size = 278, normalized size of antiderivative = 4.56

method	result
default	$\frac{\sqrt{2} \sqrt{a(\sec(fx+e)+1)} \sqrt{c+d\sec(fx+e)} \left(\ln \left(\frac{2\sqrt{2} \sqrt{-d} \sqrt{-\frac{2(d+c \cos(fx+e))}{\cos(fx+e)+1}} \sin(fx+e) - 2 \sin(fx+e)c - 2 \sin(fx+e)d + 2c \cos(fx+e) - 2d \cos(fx+e)}}{-\cos(fx+e)+1+\sin(fx+e)} \right) \right)}{f\sqrt{-d}(\cos(fx+e)+1)}$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 1/f*2^(1/2)/(-d)^(1/2)*(a*(sec(f*x+e)+1))^(1/2)*(c+d*sec(f*x+e))^(1/2)*(ln
(2*(2^(1/2)*(-d)^(1/2)*(-2*(d+c*cos(f*x+e)))/(cos(f*x+e)+1))^(1/2)*sin(f*x+
e)-sin(f*x+e)*c-sin(f*x+e)*d+c*cos(f*x+e)-d*cos(f*x+e)-c+d)/(-cos(f*x+e)+1
+sin(f*x+e)))-ln(-2*(2^(1/2)*(-d)^(1/2)*(-2*(d+c*cos(f*x+e)))/(cos(f*x+e)+1
))^(1/2)*sin(f*x+e)-sin(f*x+e)*c-sin(f*x+e)*d-c*cos(f*x+e)+d*cos(f*x+e)+c-
d)/(cos(f*x+e)-1+sin(f*x+e))))*cos(f*x+e)/(cos(f*x+e)+1)/(-2*(d+c*cos(f*x+
e))/(cos(f*x+e)+1))^(1/2)
```

3.234.
$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$$

3.234.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. $2(49) = 98$.

Time = 0.41 (sec) , antiderivative size = 307, normalized size of antiderivative = 5.03

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$$

$$= \frac{\sqrt{\frac{a}{d}} \log \left(-\frac{8acd\cos(fx+e)+(ac^2-6acd+ad^2)\cos(fx+e)^3+4(2d^2\cos(fx+e)+(cd-d^2)\cos(fx+e)^2)\sqrt{\frac{a}{d}}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)+d}{\cos(fx+e)}}}{\cos(fx+e)^3+\cos(fx+e)^2} \right)}{2f}$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `[1/2*sqrt(a/d)*log(-(8*a*c*d*cos(f*x + e) + (a*c^2 - 6*a*c*d + a*d^2)*cos(f*x + e)^3 + 4*(2*d^2*cos(f*x + e) + (c*d - d^2)*cos(f*x + e)^2)*sqrt(a/d)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*sin(f*x + e) + 8*a*d^2 + (a*c^2 + 2*a*c*d - 7*a*d^2)*cos(f*x + e)^2)/(cos(f*x + e)^3 + cos(f*x + e)^2))/f, sqrt(-a/d)*arctan(-2*d*sqrt(-a/d)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)/((a*c - a*d)*cos(f*x + e)^2 + 2*a*d + (a*c + a*d)*cos(f*x + e)))/f]`

3.234.6 SymPy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \int \frac{\sqrt{a(\sec(e+fx)+1)}\sec(e+fx)}{\sqrt{c+d\sec(e+fx)}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e))**(1/2),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*sec(e + f*x)/sqrt(c + d*sec(e + f*x)), x)`

3.234.7 Maxima [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \int \frac{\sqrt{a\sec(fx+e)+a}\sec(fx+e)}{\sqrt{d\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="maxima")`

output `integrate(sqrt(a*sec(f*x + e) + a)*sec(f*x + e)/sqrt(d*sec(f*x + e) + c),
x)`

3.234.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \int \frac{\sqrt{a\sec(fx+e)+a}\sec(fx+e)}{\sqrt{d\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="giac")`

output `sage0*x`

3.234.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \int \frac{\sqrt{a+\frac{a}{\cos(e+fx)}}}{\cos(e+fx)\sqrt{c+\frac{d}{\cos(e+fx)}}} dx$$

input `int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))^(1/2)),x
)`

output `int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))^(1/2)),
x)`

3.234. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$

3.235
$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx$$

3.235.1 Optimal result 1718
 3.235.2 Mathematica [A] (verified) 1718
 3.235.3 Rubi [A] (verified) 1719
 3.235.4 Maple [B] (warning: unable to verify) 1721
 3.235.5 Fricas [A] (verification not implemented) 1722
 3.235.6 Sympy [F] 1723
 3.235.7 Maxima [F] 1724
 3.235.8 Giac [F] 1724
 3.235.9 Mupad [F(-1)] 1724

3.235.1 Optimal result

Integrand size = 35, antiderivative size = 140

$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \frac{\sqrt{2}\sqrt{c-d}\arctan\left(\frac{\sqrt{a}\sqrt{c-d}\tan(e+fx)}{\sqrt{2}\sqrt{a+a\sec(e+fx)}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{a}f} + \frac{2\sqrt{d}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{a}f}$$

output `arctan(1/2*a^(1/2)*(c-d)^(1/2)*tan(f*x+e)*2^(1/2)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2))*2^(1/2)*(c-d)^(1/2)/f/a^(1/2)+2*arctanh(a^(1/2)*d^(1/2)*tan(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2))*d^(1/2)/f/a^(1/2)`

3.235.2 Mathematica [A] (verified)

Time = 14.28 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.34

$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \frac{\sqrt{c}\left(-\sqrt{2}\sqrt{c-d}\arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{d+c\cos(e+fx)}}{\sqrt{c-d}\sqrt{c-c\cos(e+fx)}}\right) + 2\sqrt{d}\operatorname{arctanh}\left(\frac{\sqrt{c}\sqrt{d+c\cos(e+fx)}}{\sqrt{d}\sqrt{c-c\cos(e+fx)}}\right)\right)\sqrt{c+d\sec(e+fx)}\sin(e+fx)}{f\sqrt{c-c\cos(e+fx)}\sqrt{d+c\cos(e+fx)}\sqrt{a(1+\sec(e+fx))}}$$

3.235.
$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx$$

input `Integrate[(Sec[e + f*x]*Sqrt[c + d*Sec[e + f*x]])/Sqrt[a + a*Sec[e + f*x]],x]`

output `(Sqrt[c]*(-(Sqrt[2]*Sqrt[c - d]*ArcTan[(Sqrt[2]*Sqrt[c]*Sqrt[d + c*Cos[e + f*x]])/(Sqrt[c - d]*Sqrt[c - c*Cos[e + f*x]])]) + 2*Sqrt[d]*ArcTanh[(Sqrt[c]*Sqrt[d + c*Cos[e + f*x]])/(Sqrt[d]*Sqrt[c - c*Cos[e + f*x]])])*Sqrt[c + d*Sec[e + f*x]]*Sin[e + f*x])/(f*Sqrt[c - c*Cos[e + f*x]]*Sqrt[d + c*Cos[e + f*x]]*Sqrt[a*(1 + Sec[e + f*x])])`

3.235.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4469, 3042, 4468, 219, 4471, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a\sec(e+fx)+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{c+d\csc(e+fx+\frac{\pi}{2})}}{\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}} dx \\
 & \quad \downarrow \text{4469} \\
 & (c-d) \int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a+a}\sqrt{c+d\sec(e+fx)}} dx + \frac{d \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{\sqrt{c+d\sec(e+fx)}} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & (c-d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}\sqrt{c+d\csc(e+fx+\frac{\pi}{2})}} dx + \frac{d \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{\sqrt{c+d\csc(e+fx+\frac{\pi}{2})}} dx}{a} \\
 & \quad \downarrow \text{4468}
 \end{aligned}$$

3.235. $\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx$

$$\begin{aligned}
& (c-d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a\sqrt{c+d\csc(e+fx+\frac{\pi}{2})}}} dx - \\
& \frac{2d \int \frac{1}{1-\frac{ad\tan^2(e+fx)}{(\sec(e+fx)a+a)(c+d\sec(e+fx))}}} d\left(-\frac{\tan(e+fx)}{\sqrt{\sec(e+fx)a+a\sqrt{c+d\sec(e+fx)}}}\right)}{f} \\
& \quad \downarrow \text{219} \\
& (c-d) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a\sqrt{c+d\csc(e+fx+\frac{\pi}{2})}}} dx + \\
& \frac{2\sqrt{d}\operatorname{darctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a\sqrt{c+d\sec(e+fx)}}}\right)}{\sqrt{a}f} \\
& \quad \downarrow \text{4471} \\
& \frac{2\sqrt{d}\operatorname{darctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a\sqrt{c+d\sec(e+fx)}}}\right)}{\sqrt{a}f} - \\
& \frac{2(c-d) \int \frac{1}{\frac{a(c-d)\tan^2(e+fx)}{(\sec(e+fx)a+a)(c+d\sec(e+fx))}+2}} d\left(-\frac{\tan(e+fx)}{\sqrt{\sec(e+fx)a+a\sqrt{c+d\sec(e+fx)}}}\right)}{f} \\
& \quad \downarrow \text{216} \\
& \frac{\sqrt{2}\sqrt{c-d}\operatorname{arctan}\left(\frac{\sqrt{a}\sqrt{c-d}\tan(e+fx)}{\sqrt{2}\sqrt{a\sec(e+fx)+a\sqrt{c+d\sec(e+fx)}}}\right)}{\sqrt{a}f} + \frac{2\sqrt{d}\operatorname{darctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a\sqrt{c+d\sec(e+fx)}}}\right)}{\sqrt{a}f}
\end{aligned}$$

input `Int[(Sec[e + f*x]*Sqrt[c + d*Sec[e + f*x]])/Sqrt[a + a*Sec[e + f*x]],x]`

output `(Sqrt[2]*Sqrt[c - d]*ArcTan[(Sqrt[a]*Sqrt[c - d]*Tan[e + f*x])/(Sqrt[2]*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])]/(Sqrt[a]*f) + (2*Sqrt[d]*ArcTanh[(Sqrt[a]*Sqrt[d]*Tan[e + f*x])/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])]/(Sqrt[a]*f))`

3.235.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4468 Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/Sq
rt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)], x_Symbol] := Simp[-2*(b/f) Subs
t[Int[1/(1 - b*d*x^2), x], x, Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c
+ d*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 4469 Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/Sq
rt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)], x_Symbol] := Simp[-(b*c - a*d)/d
Int[Csc[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])], x],
x] + Simp[b/d Int[Csc[e + f*x]*(Sqrt[c + d*Csc[e + f*x]]/Sqrt[a + b*Csc[
e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 - b^2, 0] && EqQ[c^2 - d^2, 0]
```

```
rule 4471 Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*Sqr
t[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)])], x_Symbol] := Simp[-2*(a/(b*f))
Subst[Int[1/(2 + (a*c - b*d)*x^2), x], x, Cot[e + f*x]/(Sqrt[a + b*Csc[e +
f*x]]*Sqrt[c + d*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.235.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 477 vs. 2(113) = 226.

Time = 5.30 (sec) , antiderivative size = 478, normalized size of antiderivative = 3.41

method	result
default	$-\frac{\sqrt{2} \sqrt{c+d \sec (f x+e)} \sqrt{a(\sec (f x+e)+1)} \left(\ln \left(\frac{\sqrt{-\frac{2(d+c \cos (f x+e))}{\cos (f x+e)+1}} \sqrt{c-d}-c \cot (f x+e)+d \cot (f x+e)+c \csc (f x+e)-d \csc (f x+e)}}{\sqrt{c-d}} \right) \right) \sqrt{2} \sqrt{c+d \sec (f x+e)}}{2 a \sqrt{a+d \sec (f x+e)}}$

3.235.
$$\int \frac{\sec(e+fx)\sqrt{c+d \sec(e+fx)}}{\sqrt{a+a \sec(e+fx)}} dx$$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/f/a/(c-d)^(1/2)*2^(1/2)/(-d)^(1/2)*(c+d*sec(f*x+e))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)*(ln(1/(c-d)^(1/2)*((-2*(d+c*cos(f*x+e)))/(cos(f*x+e)+1))^(1/2)*(c-d)^(1/2)-c*cot(f*x+e)+d*cot(f*x+e)+c*csc(f*x+e)-d*csc(f*x+e)))*2^(1/2)*(-d)^(1/2)*c-ln(1/(c-d)^(1/2)*((-2*(d+c*cos(f*x+e)))/(cos(f*x+e)+1))^(1/2)*(c-d)^(1/2)-c*cot(f*x+e)+d*cot(f*x+e)+c*csc(f*x+e)-d*csc(f*x+e)))*2^(1/2)*(-d)^(1/2)*d+d*ln(2*(-2^(1/2)*(-d)^(1/2)*(-2*(d+c*cos(f*x+e)))/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)+c*cos(f*x+e)-d*cos(f*x+e)+sin(f*x+e)*c+sin(f*x+e)*d-c+d)/(cos(f*x+e)-1+sin(f*x+e)))*(c-d)^(1/2)-d*ln(-2*(2^(1/2)*(-d)^(1/2)*(-2*(d+c*cos(f*x+e)))/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-sin(f*x+e)*c-sin(f*x+e)*d+c*cos(f*x+e)-d*cos(f*x+e)-c+d)/(cos(f*x+e)-1-sin(f*x+e)))*(c-d)^(1/2)*cos(f*x+e)/(cos(f*x+e)+1)/(-2*(d+c*cos(f*x+e)))/(cos(f*x+e)+1))^(1/2)`

3.235.5 Fracas [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 1048, normalized size of antiderivative = 7.49

$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="fracas")`

output `[1/2*(sqrt(2)*sqrt(-(c - d)/a)*log(-(2*sqrt(2)*sqrt(-(c - d)/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) - (3*c - d)*cos(f*x + e)^2 - 2*(c + d)*cos(f*x + e) + c - 3*d)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + sqrt(d/a)*log(-((c^2 - 6*c*d + d^2)*cos(f*x + e)^3 + 4*((c - d)*cos(f*x + e)^2 + 2*d*cos(f*x + e)))*sqrt(d/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*sin(f*x + e) + 8*c*d*cos(f*x + e) + (c^2 + 2*c*d - 7*d^2)*cos(f*x + e)^2 + 8*d^2)/(cos(f*x + e)^3 + cos(f*x + e)^2))/f, 1/2*(2*sqrt(2)*sqrt((c - d)/a)*arctan(-sqrt(2)*sqrt((c - d)/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*cos(f*x + e)/((c - d)*sin(f*x + e))) + sqrt(d/a)*log(-((c^2 - 6*c*d + d^2)*cos(f*x + e)^3 + 4*((c - d)*cos(f*x + e)^2 + 2*d*cos(f*x + e))*sqrt(d/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*sin(f*x + e) + 8*c*d*cos(f*x + e) + (c^2 + 2*c*d - 7*d^2)*cos(f*x + e)^2 + 8*d^2)/(cos(f*x + e)^3 + cos(f*x + e)^2))/f, 1/2*(sqrt(2)*sqrt(-(c - d)/a)*log(-(2*sqrt(2)*sqrt(-(c - d)/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) - (3*c - d)*cos(f*x + e)^2 - 2*(c + d)*cos(f*x + e) + c - 3*d)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + 2*sqrt(-d/a)*arctan(-2*sqrt(-d/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*cos(f*x + e)*s...`

3.235.6 Sympy [F]

$$\int \frac{\sec(e + fx)\sqrt{c + d\sec(e + fx)}}{\sqrt{a + a\sec(e + fx)}} dx = \int \frac{\sqrt{c + d\sec(e + fx)}\sec(e + fx)}{\sqrt{a(\sec(e + fx) + 1)}} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**(1/2)/(a+a*sec(f*x+e))**(1/2),x)`

output `Integral(sqrt(c + d*sec(e + f*x))*sec(e + f*x)/sqrt(a*(sec(e + f*x) + 1)), x)`

3.235.7 Maxima [F]

$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{\sqrt{d\sec(fx+e)+c}\sec(fx+e)}{\sqrt{a\sec(fx+e)+a}} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(d*sec(f*x + e) + c)*sec(f*x + e)/sqrt(a*sec(f*x + e) + a), x)`

3.235.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{\sqrt{d\sec(fx+e)+c}\sec(fx+e)}{\sqrt{a\sec(fx+e)+a}} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(d*sec(f*x + e) + c)*sec(f*x + e)/sqrt(a*sec(f*x + e) + a), x)`

3.235.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx = \int \frac{\sqrt{c+\frac{d}{\cos(e+fx)}}}{\cos(e+fx)\sqrt{a+\frac{a}{\cos(e+fx)}}} dx$$

input `int((c + d/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

output `int((c + d/cos(e + f*x))^(1/2)/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)),x)`

3.235. $\int \frac{\sec(e+fx)\sqrt{c+d\sec(e+fx)}}{\sqrt{a+a\sec(e+fx)}} dx$

3.236
$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$$

3.236.1 Optimal result 1725
 3.236.2 Mathematica [A] (verified) 1725
 3.236.3 Rubi [A] (verified) 1726
 3.236.4 Maple [B] (verified) 1727
 3.236.5 Fricas [A] (verification not implemented) 1727
 3.236.6 Sympy [F] 1728
 3.236.7 Maxima [F] 1728
 3.236.8 Giac [F] 1729
 3.236.9 Mupad [F(-1)] 1729

3.236.1 Optimal result

Integrand size = 35, antiderivative size = 78

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a}\sqrt{c-d} \tan(e+fx)}{\sqrt{2}\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}}\right)}{\sqrt{a}\sqrt{c-d}f}$$

output `arctan(1/2*a^(1/2)*(c-d)^(1/2)*tan(f*x+e)*2^(1/2)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2))*2^(1/2)/f/a^(1/2)/(c-d)^(1/2)`

3.236.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.37

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx = \frac{2 \arctan\left(\frac{\sqrt{c-d} \sin(\frac{1}{2}(e+fx))}{\sqrt{d+c \cos(e+fx)}}\right) \cos\left(\frac{1}{2}(e+fx)\right) \sqrt{d+c \cos(e+fx)} \sec(e+fx)}{\sqrt{c-d}f\sqrt{a(1+\sec(e+fx))}\sqrt{c+d \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `(2*ArcTan[(Sqrt[c - d]*Sin[(e + f*x)/2])/Sqrt[d + c*Cos[e + f*x]]]*Cos[(e + f*x)/2]*Sqrt[d + c*Cos[e + f*x]]*Sec[e + f*x])/((Sqrt[c - d]*f*Sqrt[a*(1 + Sec[e + f*x])])*Sqrt[c + d*Sec[e + f*x]])`

3.236.
$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$$

3.236.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.086$, Rules used = {3042, 4471, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{\sqrt{a \sec(e+fx) + a} \sqrt{c+d \sec(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)}{\sqrt{a \csc\left(e+fx+\frac{\pi}{2}\right) + a} \sqrt{c+d \csc\left(e+fx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{4471} \\
 & -\frac{2 \int \frac{1}{\frac{a(c-d) \tan^2(e+fx)}{(\sec(e+fx)a+a)(c+d \sec(e+fx))} + 2}}{f} d\left(-\frac{\tan(e+fx)}{\sqrt{\sec(e+fx)a+a} \sqrt{c+d \sec(e+fx)}}\right) \\
 & \quad \downarrow \text{216} \\
 & \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a} \sqrt{c-d} \tan(e+fx)}{\sqrt{2} \sqrt{a \sec(e+fx) + a} \sqrt{c+d \sec(e+fx)}}\right)}{\sqrt{a} f \sqrt{c-d}}
 \end{aligned}$$

input `Int[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `(Sqrt[2]*ArcTan[(Sqrt[a]*Sqrt[c - d]*Tan[e + f*x])/(Sqrt[2]*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])]/(Sqrt[a]*Sqrt[c - d]*f))`

3.236.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.236. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx$

```
rule 4471 Int[csc[(e_.) + (f_.)*(x_.)]/(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)]*Sqr
t[csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)]), x_Symbol] :> Simp[-2*(a/(b*f))
Subst[Int[1/(2 + (a*c - b*d)*x^2), x], x, Cot[e + f*x]/(Sqrt[a + b*Csc[e +
f*x]]*Sqrt[c + d*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.236.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(63) = 126.

Time = 2.86 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.73

method	result	size
default	$-\frac{2\sqrt{c+d\sec(fx+e)}\sqrt{a(\sec(fx+e)+1)}\ln\left(\sqrt{\frac{-2(d+c\cos(fx+e))}{\cos(fx+e)+1}}-\sqrt{c-d}\cot(fx+e)+\sqrt{c-d}\csc(fx+e)\right)\cos(fx+e)}{fa\sqrt{c-d}(\cos(fx+e)+1)\sqrt{\frac{-2(d+c\cos(fx+e))}{\cos(fx+e)+1}}}$	135

```
input int(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -2/f/a/(c-d)^(1/2)*(c+d*sec(f*x+e))^(1/2)*(a*(sec(f*x+e)+1))^(1/2)*ln((-2*
(d+c*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)-(c-d)^(1/2)*cot(f*x+e)+(c-d)^(1/2)*
csc(f*x+e))*cos(f*x+e)/(cos(f*x+e)+1)/(-2*(d+c*cos(f*x+e))/(cos(f*x+e)+1))
^(1/2)
```

3.236.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 246, normalized size of antiderivative = 3.15

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}\sqrt{c + d \sec(e + fx)}} dx$$

$$= \left[\frac{\sqrt{2}\sqrt{-\frac{1}{ac-ad}} \log\left(-\frac{2\sqrt{2}(c-d)\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)+d}{\cos(fx+e)}}\sqrt{-\frac{1}{ac-ad}}\cos(fx+e)\sin(fx+e)-(3c-d)\cos(fx+e)^2-2(c+d)\cos(fx+e)}}{\cos(fx+e)^2+2\cos(fx+e)+1}\right)}{2f} \right.$$

$$\left. - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{\frac{a\cos(fx+e)+a}{\cos(fx+e)}}\sqrt{\frac{c\cos(fx+e)+d}{\cos(fx+e)}}\cos(fx+e)}{\sqrt{ac-ad}\sin(fx+e)}\right)}{\sqrt{ac-ad}f} \right]$$

3.236. $\int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="fricas")`

output `[1/2*sqrt(2)*sqrt(-1/(a*c - a*d))*log(-(2*sqrt(2)*(c - d)*sqrt((a*cos(f*x
+ e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*sqrt(-1/(a
*c - a*d))*cos(f*x + e)*sin(f*x + e) - (3*c - d)*cos(f*x + e)^2 - 2*(c + d
) *cos(f*x + e) + c - 3*d)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1))/f, -sqrt(
2)*arctan(sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x
+ e) + d)/cos(f*x + e))*cos(f*x + e)/(sqrt(a*c - a*d)*sin(f*x + e)))/(sqrt
(a*c - a*d)*f)]`

3.236.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec(e + fx)}{\sqrt{a (\sec(e + fx) + 1)} \sqrt{c + d \sec(e + fx)}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(a*(sec(e + f*x) + 1))*sqrt(c + d*sec(e + f*x))
, x)`

3.236.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{a \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="maxima")`

output `integrate(sec(f*x + e)/(sqrt(a*sec(f*x + e) + a))*sqrt(d*sec(f*x + e) + c)
, x)`

3.236. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx$

3.236.8 Giac [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{a \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorith="giac")`

output `integrate(sec(f*x + e)/(sqrt(a*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)), x)`

3.236.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx) \sqrt{a + \frac{a}{\cos(e + fx)}} \sqrt{c + \frac{d}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2)),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2)), x)`

3.237 $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$

3.237.1 Optimal result	1730
3.237.2 Mathematica [A] (verified)	1730
3.237.3 Rubi [A] (verified)	1731
3.237.4 Maple [B] (warning: unable to verify)	1734
3.237.5 Fricas [A] (verification not implemented)	1734
3.237.6 Sympy [F]	1735
3.237.7 Maxima [F]	1736
3.237.8 Giac [F]	1736
3.237.9 Mupad [F(-1)]	1736

3.237.1 Optimal result

Integrand size = 37, antiderivative size = 141

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx = -\frac{\sqrt{2} \arctan\left(\frac{\sqrt{a}\sqrt{c-d} \tan(e+fx)}{\sqrt{2}\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}}\right)}{\sqrt{a}\sqrt{c-d}f} + \frac{2\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d} \tan(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}}\right)}{\sqrt{a}\sqrt{d}f}$$

output

```
-arctan(1/2*a^(1/2)*(c-d)^(1/2)*tan(f*x+e)*2^(1/2)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2))*2^(1/2)/f/a^(1/2)/(c-d)^(1/2)+2*arctanh(a^(1/2)*d^(1/2)*tan(f*x+e)/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2))/f/a^(1/2)/d^(1/2)
```

3.237.2 Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.21

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx = \frac{2\left(-\sqrt{d} \arctan\left(\frac{\sqrt{c-d} \sin(\frac{1}{2}(e+fx))}{\sqrt{d+c \cos(e+fx)}}\right) + \sqrt{2}\sqrt{c-d} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{d} \sin(\frac{1}{2}(e+fx))}{\sqrt{d+c \cos(e+fx)}}\right)\right) \cos\left(\frac{1}{2}(e+fx)\right) \sqrt{d+c \cos(e+fx)}}{\sqrt{c-d}\sqrt{d}f\sqrt{a(1+\sec(e+fx))}\sqrt{c+d \sec(e+fx)}}$$

3.237. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$

input `Integrate[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `(2*(-(Sqrt[d]*ArcTan[(Sqrt[c - d]*Sin[(e + f*x)/2])/Sqrt[d + c*Cos[e + f*x]]]) + Sqrt[2]*Sqrt[c - d]*ArcTanh[(Sqrt[2]*Sqrt[d]*Sin[(e + f*x)/2])/Sqrt[d + c*Cos[e + f*x]]])*Cos[(e + f*x)/2]*Sqrt[d + c*Cos[e + f*x]]*Sec[e + f*x])/(Sqrt[c - d]*Sqrt[d]*f*Sqrt[a*(1 + Sec[e + f*x])]*Sqrt[c + d*Sec[e + f*x]])`

3.237.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.189$, Rules used = {3042, 4473, 3042, 4468, 219, 4471, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a \sec(e + fx) + a} \sqrt{c + d \sec(e + fx)}} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})^2}{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a} \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx$$

↓ 4473

$$\frac{\int \frac{\sec(e + fx) \sqrt{\sec(e + fx) a + a}}{\sqrt{c + d \sec(e + fx)}} dx}{a} - \int \frac{\sec(e + fx)}{\sqrt{\sec(e + fx) a + a} \sqrt{c + d \sec(e + fx)}} dx$$

↓ 3042

$$\frac{\int \frac{\csc(e + fx + \frac{\pi}{2}) \sqrt{\csc(e + fx + \frac{\pi}{2}) a + a}}{\sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx}{a} - \int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{\csc(e + fx + \frac{\pi}{2}) a + a} \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx$$

↓ 4468

3.237. $\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$

$$\begin{aligned}
 & - \int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{\csc(e + fx + \frac{\pi}{2})a + a\sqrt{c + d\csc(e + fx + \frac{\pi}{2})}}} dx - \\
 & \frac{2 \int \frac{1}{1 - \frac{ad \tan^2(e+fx)}{(\sec(e+fx)a+a)(c+d\sec(e+fx))}} d\left(-\frac{\tan(e+fx)}{\sqrt{\sec(e+fx)a+a}\sqrt{c+d\sec(e+fx)}}\right)}{f} \\
 & \quad \downarrow \text{219} \\
 & \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{a}\sqrt{d}f} - \int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{\csc(e + fx + \frac{\pi}{2})a + a\sqrt{c + d\csc(e + fx + \frac{\pi}{2})}}} dx \\
 & \quad \downarrow \text{4471} \\
 & \frac{2 \int \frac{1}{\frac{a(c-d)\tan^2(e+fx)}{(\sec(e+fx)a+a)(c+d\sec(e+fx))} + 2} d\left(-\frac{\tan(e+fx)}{\sqrt{\sec(e+fx)a+a}\sqrt{c+d\sec(e+fx)}}\right)}{f} + \\
 & \quad \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{a}\sqrt{d}f} \\
 & \quad \downarrow \text{216} \\
 & \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{a}\sqrt{d}f} - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a}\sqrt{c-d}\tan(e+fx)}{\sqrt{2}\sqrt{a\sec(e+fx)+a}\sqrt{c+d\sec(e+fx)}}\right)}{\sqrt{a}f\sqrt{c-d}}
 \end{aligned}$$

input `Int[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `-((Sqrt[2]*ArcTan[(Sqrt[a]*Sqrt[c - d]*Tan[e + f*x])/(Sqrt[2]*Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])])/(Sqrt[a]*Sqrt[c - d]*f)) + (2*ArcTanh[(Sqrt[a]*Sqrt[d]*Tan[e + f*x])/(Sqrt[a + a*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])])/(Sqrt[a]*Sqrt[d]*f)`

3.237.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4468 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)], x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(1 - b*d*x^2), x], x, Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4471 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)]), x_Symbol] := Simp[-2*(a/(b*f)) Subst[Int[1/(2 + (a*c - b*d)*x^2), x], x, Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4473 `Int[csc[(e_.) + (f_.)*(x_)]^2/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)]), x_Symbol] := Simp[-a/b Int[Csc[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]]), x], x] + Simp[1/b Int[Csc[e + f*x]*(Sqrt[a + b*Csc[e + f*x]]/Sqrt[c + d*Csc[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

3.237.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 365 vs. $2(114) = 228$.

Time = 5.16 (sec) , antiderivative size = 366, normalized size of antiderivative = 2.60

method	result
default	$\frac{\sqrt{2} \sqrt{c+d \sec(fx+e)} \sqrt{a(\sec(fx+e)+1)} \left(\ln \left(\sqrt{-\frac{2(d+c \cos(fx+e))}{\cos(fx+e)+1}} - \sqrt{c-d} \cot(fx+e) + \sqrt{c-d} \csc(fx+e) \right) \sqrt{2} \sqrt{-d} - \ln \left(\frac{-2\sqrt{2}\sqrt{-d}}{\dots} \right) \right)}{\dots}$

input `int(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{f a (c-d)^{1/2} 2^{1/2} (-d)^{1/2} (c+d \sec(fx+e))^{1/2} (a(\sec(fx+e)+1))^{1/2} \left(\ln \left(\frac{-2(d+c \cos(fx+e))}{\cos(fx+e)+1} \right)^{1/2} - (c-d)^{1/2} \cot(fx+e) + (c-d)^{1/2} \csc(fx+e) \right) 2^{1/2} (-d)^{1/2} - \ln \left(2(-2)^{1/2} (-d)^{1/2} (-2(d+c \cos(fx+e)) / (\cos(fx+e)+1))^{1/2} \sin(fx+e) + c \cos(fx+e) - d \cos(fx+e) + \sin(fx+e) * c + \sin(fx+e) * d - c + d / (\cos(fx+e) - 1 + \sin(fx+e)) \right) * (c-d)^{1/2} + \ln \left(-2(2)^{1/2} (-d)^{1/2} (-2(d+c \cos(fx+e)) / (\cos(fx+e)+1))^{1/2} \sin(fx+e) - \sin(fx+e) * c - \sin(fx+e) * d + c \cos(fx+e) - d \cos(fx+e) - c + d / (\cos(fx+e) - 1 - \sin(fx+e)) \right) * (c-d)^{1/2} \right) \cos(fx+e) / (\cos(fx+e)+1) / (-2(d+c \cos(fx+e)) / (\cos(fx+e)+1))^{1/2}}$$

3.237.5 Fracas [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 1100, normalized size of antiderivative = 7.80

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorithm="fracas")`

```
output [1/2*(sqrt(2)*a*d*sqrt(-1/(a*c - a*d))*log((2*sqrt(2)*(c - d)*sqrt((a*cos(
f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*sqrt(-
1/(a*c - a*d))*cos(f*x + e)*sin(f*x + e) + (3*c - d)*cos(f*x + e)^2 + 2*(c
+ d)*cos(f*x + e) - c + 3*d)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + sqr
t(a*d)*log(-(8*a*c*d*cos(f*x + e) + (a*c^2 - 6*a*c*d + a*d^2)*cos(f*x + e)
^3 + 4*((c - d)*cos(f*x + e)^2 + 2*d*cos(f*x + e))*sqrt(a*d)*sqrt((a*cos(f
*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*sin(f*x
+ e) + 8*a*d^2 + (a*c^2 + 2*a*c*d - 7*a*d^2)*cos(f*x + e)^2)/(cos(f*x + e)
^3 + cos(f*x + e)^2)))/(a*d*f), 1/2*(2*sqrt(2)*a*d*arctan(sqrt(2)*sqrt((a
*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*c
os(f*x + e)/(sqrt(a*c - a*d)*sin(f*x + e)))/sqrt(a*c - a*d) + sqrt(a*d)*lo
g(-(8*a*c*d*cos(f*x + e) + (a*c^2 - 6*a*c*d + a*d^2)*cos(f*x + e)^3 + 4*((
c - d)*cos(f*x + e)^2 + 2*d*cos(f*x + e))*sqrt(a*d)*sqrt((a*cos(f*x + e) +
a)/cos(f*x + e))*sqrt((c*cos(f*x + e) + d)/cos(f*x + e))*sin(f*x + e) + 8
*a*d^2 + (a*c^2 + 2*a*c*d - 7*a*d^2)*cos(f*x + e)^2)/(cos(f*x + e)^3 + cos
(f*x + e)^2)))/(a*d*f), 1/2*(sqrt(2)*a*d*sqrt(-1/(a*c - a*d))*log((2*sqrt(
2)*(c - d)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt((c*cos(f*x + e) +
d)/cos(f*x + e))*sqrt(-1/(a*c - a*d))*cos(f*x + e)*sin(f*x + e) + (3*c - d)
)*cos(f*x + e)^2 + 2*(c + d)*cos(f*x + e) - c + 3*d)/(cos(f*x + e)^2 + 2*c
os(f*x + e) + 1)) + 2*sqrt(-a*d)*arctan(-2*sqrt(-a*d)*sqrt((a*cos(f*x + ...
```

3.237.6 Sympy [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec^2(e + fx)}{\sqrt{a(\sec(e + fx) + 1)} \sqrt{c + d \sec(e + fx)}} dx$$

```
input integrate(sec(f*x+e)**2/(a+a*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e))**(1/2),x)
```

```
output Integral(sec(e + f*x)**2/(sqrt(a*(sec(e + f*x) + 1))*sqrt(c + d*sec(e + f*
x))), x)
```


3.237.7 Maxima [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec^2(fx + e)}{\sqrt{a \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)^2/(sqrt(a*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)), x)`

3.237.8 Giac [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec^2(fx + e)}{\sqrt{a \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)^2/(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate(sec(f*x + e)^2/(sqrt(a*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)), x)`

3.237.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx)^2 \sqrt{a + \frac{a}{\cos(e + fx)}} \sqrt{c + \frac{d}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x)^2*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2)),x)`

output `int(1/(cos(e + f*x)^2*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2)), x)`

3.237. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$

3.238
$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

3.238.1 Optimal result	1738
3.238.2 Mathematica [A] (verified)	1738
3.238.3 Rubi [A] (verified)	1739
3.238.4 Maple [B] (warning: unable to verify)	1740
3.238.5 Fracas [B] (verification not implemented)	1741
3.238.6 Sympy [F]	1741
3.238.7 Maxima [F]	1742
3.238.8 Giac [F]	1742
3.238.9 Mupad [F(-1)]	1742

3.238.1 Optimal result

Integrand size = 33, antiderivative size = 61

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx = \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{c+d}\sqrt{a+a\sec(e+fx)}}\right)}{\sqrt{d}\sqrt{c+df}}$$

output `2*arctan(a^(1/2)*d^(1/2)*tan(f*x+e)/(c+d)^(1/2)/(a+a*sec(f*x+e))^(1/2))*a^(1/2)/f/d^(1/2)/(c+d)^(1/2)`

3.238.2 Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.54

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{d}\sin(\frac{1}{2}(e+fx))}{\sqrt{c+d}\sqrt{\cos(e+fx)}}\right) \sqrt{\cos(e+fx)} \sec\left(\frac{1}{2}(e+fx)\right) \sqrt{a(1+\sec(e+fx))}}{\sqrt{d}\sqrt{c+df}}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`

output `(Sqrt[2]*ArcTan[(Sqrt[2]*Sqrt[d]*Sin[(e + f*x)/2])/(Sqrt[c + d]*Sqrt[Cos[e + f*x]])]*Sqrt[Cos[e + f*x]]*Sec[(e + f*x)/2]*Sqrt[a*(1 + Sec[e + f*x])])/(Sqrt[d]*Sqrt[c + d]*f)`

3.238.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {3042, 4455, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a\sec(e+fx)+a}}{c+d\sec(e+fx)} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a\csc(e+fx+\frac{\pi}{2})+a}}{c+d\csc(e+fx+\frac{\pi}{2})} dx$$

↓ 4455

$$-\frac{2a \int \frac{1}{\frac{a^2 d \tan^2(e+fx)}{\sec(e+fx)a+a} + a(c+d)} d\left(-\frac{a \tan(e+fx)}{\sqrt{\sec(e+fx)a+a}}\right)}{f}$$

↓ 218

$$\frac{2\sqrt{a} \arctan\left(\frac{\sqrt{a}\sqrt{d}\tan(e+fx)}{\sqrt{c+d}\sqrt{a\sec(e+fx)+a}}\right)}{\sqrt{d}f\sqrt{c+d}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + a*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`

output `(2*Sqrt[a]*ArcTan[(Sqrt[a]*Sqrt[d]*Tan[e + f*x])/(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])])/(Sqrt[d]*Sqrt[c + d]*f)`

3.238.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.238. $\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx$

```
rule 4455 Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d + d*x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2,
0]
```

3.238.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 423 vs. 2(47) = 94.

Time = 19.38 (sec) , antiderivative size = 424, normalized size of antiderivative = 6.95

method	result
default	$\sqrt{2} \left(\ln \left(-\frac{2 \left(\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} \sqrt{2} \sqrt{\frac{d}{c-d}} c - \sqrt{2} \sqrt{\frac{d}{c-d}} \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1} d + \sqrt{(c+d)(c-d)} (-\cot(fx+e) + \csc(fx+e)) \right)}{-c(-\cot(fx+e) + \csc(fx+e)) + (-\cot(fx+e) + \csc(fx+e))d + \sqrt{(c+d)(c-d)}} \right) \right)$

```
input int(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,method=_RETURNVER
BOSE)
```

```
output 1/2/f*2^(1/2)/(d/(c-d))^(1/2)/((c+d)*(c-d))^(1/2)*(ln(-2*(((1-cos(f*x+e))^
2*csc(f*x+e)^2-1)^(1/2)*2^(1/2)*(d/(c-d))^(1/2)*c-2^(1/2)*(d/(c-d))^(1/2)*
((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*d+((c+d)*(c-d))^(1/2)*(-cot(f*x+e)
+csc(f*x+e))-c+d)/(-c*(-cot(f*x+e)+csc(f*x+e))+(-cot(f*x+e)+csc(f*x+e))*d+
((c+d)*(c-d))^(1/2)))-ln(2*(((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*2^(1/2)
)*(d/(c-d))^(1/2)*c-2^(1/2)*(d/(c-d))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2
-1)^(1/2)*d-((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))-c+d)/(c*(-cot(f*x
+e)+csc(f*x+e))-(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2))))*((1-cos(
f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(-2*a/(((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(
1/2)
```

$$3.238. \int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

3.238.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(47) = 94$.

Time = 0.44 (sec) , antiderivative size = 343, normalized size of antiderivative = 5.62

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

$$= \left[\frac{\sqrt{-\frac{a}{cd+d^2}} \log\left(-\frac{(ac^2+8acd+8ad^2)\cos(fx+e)^3+ad^2+(ac^2+2acd)\cos(fx+e)^2-4((c^2d+3cd^2+2d^3)\cos(fx+e)^2-(cd^2+d^3)\cos(fx+e))}{c^2\cos(fx+e)^3+(c^2+2cd)\cos(fx+e)^2+d^2+(2cd+d^2)\cos(fx+e)}\right)}{2f} \right]$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `[1/2*sqrt(-a/(c*d + d^2))*log(-((a*c^2 + 8*a*c*d + 8*a*d^2)*cos(f*x + e)^3 + a*d^2 + (a*c^2 + 2*a*c*d)*cos(f*x + e)^2 - 4*((c^2*d + 3*c*d^2 + 2*d^3)*cos(f*x + e)^2 - (c*d^2 + d^3)*cos(f*x + e))*sqrt(-a/(c*d + d^2))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sin(f*x + e) - (6*a*c*d + 7*a*d^2)*cos(f*x + e))/(c^2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 + d^2 + (2*c*d + d^2)*cos(f*x + e)))/f, sqrt(a/(c*d + d^2))*arctan(2*(c*d + d^2)*sqrt(a/(c*d + d^2))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)/((a*c + 2*a*d)*cos(f*x + e)^2 - a*d + (a*c + a*d)*cos(f*x + e)))/f]`

3.238.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{a(\sec(e+fx)+1)}\sec(e+fx)}{c+d\sec(e+fx)} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e)),x)`

output `Integral(sqrt(a*(sec(e + f*x) + 1))*sec(e + f*x)/(c + d*sec(e + f*x)), x)`

3.238.7 Maxima [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{d\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `integrate(sqrt(a*sec(f*x + e) + a)*sec(f*x + e)/(d*sec(f*x + e) + c), x)`

3.238.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{a\sec(fx+e)+a\sec(fx+e)}}{d\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="giac")`

output `sage0*x`

3.238.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+a\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{a + \frac{a}{\cos(e+fx)}}}{\cos(e+fx) \left(c + \frac{d}{\cos(e+fx)}\right)} dx$$

input `int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))),x)`

output `int((a + a/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))), x)`

3.239
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx$$

3.239.1 Optimal result	1743
3.239.2 Mathematica [A] (verified)	1743
3.239.3 Rubi [A] (verified)	1744
3.239.4 Maple [B] (warning: unable to verify)	1746
3.239.5 Fracas [A] (verification not implemented)	1747
3.239.6 Sympy [F]	1748
3.239.7 Maxima [F(-2)]	1749
3.239.8 Giac [F]	1749
3.239.9 Mupad [F(-1)]	1749

3.239.1 Optimal result

Integrand size = 39, antiderivative size = 149

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx = \frac{2\sqrt{a}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{g \sec(e+fx)} \sqrt{a+a \sec(e+fx)}}\right)}{df} - \frac{2\sqrt{a}\sqrt{c}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{g \sec(e+fx)} \sqrt{a+a \sec(e+fx)}}\right)}{d\sqrt{c+d}f}$$

output

```

2*g^(3/2)*arctanh(a^(1/2)*g^(1/2)*tan(f*x+e)/(g*sec(f*x+e))^(1/2)/(a+a*sec
(f*x+e))^(1/2))*a^(1/2)/d/f-2*g^(3/2)*arctanh(a^(1/2)*c^(1/2)*g^(1/2)*tan(
f*x+e)/(c+d)^(1/2)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))*a^(1/2)*c^
(1/2)/d/f/(c+d)^(1/2)
    
```

3.239.2 Mathematica [A] (verified)

Time = 2.03 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.26

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx = \frac{g^2(\sqrt{c+d} \log(\sqrt{2}-2 \sin(\frac{1}{2}(e+fx))) - \sqrt{c+d} \log(\sqrt{2}+2 \sin(\frac{1}{2}(e+fx))) + \sqrt{c}(-\log(\sqrt{2}\sqrt{c+d} - \sqrt{2d\sqrt{c+d}f\sqrt{c+d}}))}{\sqrt{2d\sqrt{c+d}f\sqrt{c+d}}}$$

input `Integrate[((g*Sec[e + f*x])^(3/2)*Sqrt[a + a*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`

output `-((g^2*(Sqrt[c + d]*Log[Sqrt[2] - 2*Sin[(e + f*x)/2]] - Sqrt[c + d]*Log[Sqrt[2] + 2*Sin[(e + f*x)/2]] + Sqrt[c]*(-Log[Sqrt[2]*Sqrt[c + d] - 2*Sqrt[c]*Sin[(e + f*x)/2]] + Log[Sqrt[2]*Sqrt[c + d] + 2*Sqrt[c]*Sin[(e + f*x)/2]])*Sec[(e + f*x)/2]*Sqrt[a*(1 + Sec[e + f*x])])/(Sqrt[2]*d*Sqrt[c + d]*Sqrt[g*Sec[e + f*x]])`

3.239.3 Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {3042, 4458, 3042, 4289, 221, 4453, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a \sec(e+fx) + a} (g \sec(e+fx))^{3/2}}{c + d \sec(e+fx)} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\sqrt{a \csc(e+fx + \frac{\pi}{2}) + a} (g \csc(e+fx + \frac{\pi}{2}))^{3/2}}{c + d \csc(e+fx + \frac{\pi}{2})} dx \\
 & \quad \downarrow 4458 \\
 & \frac{g \int \sqrt{g \sec(e+fx)} \sqrt{\sec(e+fx)a + a} dx}{d} - \frac{cg \int \frac{\sqrt{g \sec(e+fx)} \sqrt{\sec(e+fx)a + a}}{c + d \sec(e+fx)} dx}{d} \\
 & \quad \downarrow 3042 \\
 & \frac{g \int \sqrt{g \csc(e+fx + \frac{\pi}{2})} \sqrt{\csc(e+fx + \frac{\pi}{2})a + a} dx}{d} - \frac{cg \int \frac{\sqrt{g \csc(e+fx + \frac{\pi}{2})} \sqrt{\csc(e+fx + \frac{\pi}{2})a + a}}{c + d \csc(e+fx + \frac{\pi}{2})} dx}{d} \\
 & \quad \downarrow 4289 \\
 & - \frac{2ag^2 \int \frac{1}{a - \frac{a^2 \sin(e+fx) \tan(e+fx)}{\sec(e+fx)a + a}} d\left(-\frac{a \tan(e+fx)}{\sqrt{g \sec(e+fx)} \sqrt{\sec(e+fx)a + a}}\right)}{d} \\
 & \quad \downarrow \\
 & \frac{cg \int \frac{\sqrt{g \csc(e+fx + \frac{\pi}{2})} \sqrt{\csc(e+fx + \frac{\pi}{2})a + a}}{c + d \csc(e+fx + \frac{\pi}{2})} dx}{d}
 \end{aligned}$$

3.239. $\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a + a \sec(e+fx)}}{c + d \sec(e+fx)} dx$

$$\begin{aligned}
 & \downarrow 221 \\
 & \frac{2\sqrt{a}g^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{g\sec(e+fx)}}\right)}{df} - \frac{cg \int \frac{\sqrt{g\csc(e+fx+\frac{\pi}{2})}\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{c+d\csc(e+fx+\frac{\pi}{2})} dx}{d} \\
 & \downarrow 4453 \\
 & \frac{2acg^2 \int \frac{1}{a(c+d) - \frac{a^2 c \sin(e+fx)\tan(e+fx)}{\sec(e+fx)a+a}} d\left(-\frac{a\tan(e+fx)}{\sqrt{g\sec(e+fx)}\sqrt{\sec(e+fx)a+a}}\right)}{df} + \\
 & \frac{2\sqrt{a}g^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{g\sec(e+fx)}}\right)}{df} \\
 & \downarrow 221 \\
 & \frac{2\sqrt{a}g^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g}\tan(e+fx)}{\sqrt{a\sec(e+fx)+a}\sqrt{g\sec(e+fx)}}\right)}{df} - \frac{2\sqrt{a}\sqrt{c}g^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g}\tan(e+fx)}{\sqrt{c+d}\sqrt{a\sec(e+fx)+a}\sqrt{g\sec(e+fx)}}\right)}{df\sqrt{c+d}}
 \end{aligned}$$

input `Int[((g*Sec[e + f*x])^(3/2)*Sqrt[a + a*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`

output `(2*Sqrt[a]*g^(3/2)*ArcTanh[(Sqrt[a]*Sqrt[g]*Tan[e + f*x])/(Sqrt[g*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])]/(d*f) - (2*Sqrt[a]*Sqrt[c]*g^(3/2)*ArcTanh[(Sqrt[a]*Sqrt[c]*Sqrt[g]*Tan[e + f*x])/(Sqrt[c + d]*Sqrt[g*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])]/(d*Sqrt[c + d]*f))`

3.239.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4289 `Int[Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*b*(d/f) Subst[Int[1/(b - d*x^2), x], x, b*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[d*Csc[e + f*x]])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && !GtQ[a*(d/b), 0]`

3.239. $\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx$

rule 4453 `Int[(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(g_.)]*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)])/(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)), x_Symbol] := Simp[-2*b*(g/f) Subst[Int[1/(b*c + a*d - c*g*x^2), x], x, b*(Cot[e + f*x]/(Sqrt[g*Csc[e + f*x]]*Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4458 `Int[((csc[(e_.) + (f_.)*(x_.)]*(g_.))^(3/2)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)])/(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)), x_Symbol] := Simp[g/d Int[Sqrt[g*Csc[e + f*x]]*Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[c*(g/d) Int[Sqrt[g*Csc[e + f*x]]*(Sqrt[a + b*Csc[e + f*x]]/(c + d*Csc[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0]`

3.239.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 684 vs. 2(117) = 234.

Time = 22.71 (sec) , antiderivative size = 685, normalized size of antiderivative = 4.60

method	result
default	$-\frac{g\sqrt{2}(c-d)\sqrt{-\frac{g((1-\cos(fx+e))^2 \csc(fx+e)^2+1)}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}}}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1} \sqrt{-\frac{2a}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}} \left(\sqrt{(c+d)(c-d)} \right)$

input `int((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)`

3.239.
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx$$

output

```
-g/f*2^(1/2)*(c-d)/((c+d)*(c-d))^(1/2)/(c-d+((c+d)*(c-d))^(1/2))/(-c+d+((c+d)*(c-d))^(1/2))/(c/(c-d))^(1/2)*(-g*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*(-2*a/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)*((c+d)*(c-d))^(1/2)*arctanh(1/2*(-cot(f*x+e)+csc(f*x+e)+1)*2^(1/2)/((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2))*(c/(c-d))^(1/2)+((c+d)*(c-d))^(1/2)*arctanh(1/2*(-cot(f*x+e)+csc(f*x+e)-1)*2^(1/2)/((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2))*(c/(c-d))^(1/2)-c*ln(-2*(2^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*c-2^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*d+((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))+c-d)/(-c*(-cot(f*x+e)+csc(f*x+e))+(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2)))+c*ln(2*(2^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*c-2^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*d-((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))+c-d)/(c*(-cot(f*x+e)+csc(f*x+e))-(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2)))/((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)
```

3.239.5 Fracas [A] (verification not implemented)

Time = 4.81 (sec) , antiderivative size = 1126, normalized size of antiderivative = 7.56

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c + d \sec(e + fx)} dx = \text{Too large to display}$$

input

```
integrate((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,
algorithm="fricas")
```

output

```
[1/2*(sqrt(a*c*g/(c + d))*g*log((a*c^2*g*cos(f*x + e)^3 - (7*a*c^2 + 6*a*c*d)*g*cos(f*x + e)^2 + 4*((c^2 + c*d)*cos(f*x + e)^2 - (2*c^2 + 3*c*d + d^2)*cos(f*x + e))*sqrt(a*c*g/(c + d))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) + (2*a*c*d + a*d^2)*g*cos(f*x + e) + (8*a*c^2 + 8*a*c*d + a*d^2)*g)/(c^2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 + d^2 + (2*c*d + d^2)*cos(f*x + e))) + sqrt(a*g)*g*log((a*g*cos(f*x + e)^3 - 7*a*g*cos(f*x + e)^2 - 4*sqrt(a*g)*(cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) + 8*a*g)/(cos(f*x + e)^3 + cos(f*x + e)^2)))/(d*f), -1/2*(2*sqrt(-a*c*g/(c + d))*g*arctan(1/2*(c*cos(f*x + e)^2 - (2*c + d)*cos(f*x + e))*sqrt(-a*c*g/(c + d))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e)))/(a*c*g*sin(f*x + e))) - sqrt(a*g)*g*log((a*g*cos(f*x + e)^3 - 7*a*g*cos(f*x + e)^2 - 4*sqrt(a*g)*(cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) + 8*a*g)/(cos(f*x + e)^3 + cos(f*x + e)^2)))/(d*f), 1/2*(2*sqrt(-a*g)*g*arctan(2*sqrt(-a*g)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)/(a*g*cos(f*x + e)^2 - a*g*cos(f*x + e) - 2*a*g)) + sqrt(a*c*g/(c + d))*g*log((a*c^2*g*cos(f*x + e)^3 - (7*a*c^2 + 6*a*c*d)*g*cos(f*x + e)^2 + 4*((c^2 + c*d)*cos(f*x + e)^2 - (2*c^2 + 3*c*d + d^2)*cos(f*x + e))*sqrt(a*c*g/(c + d))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(...
```

3.239.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c + d \sec(e + fx)} dx = \int \frac{\sqrt{a (\sec(e + fx) + 1)} (g \sec(e + fx))^{3/2}}{c + d \sec(e + fx)} dx$$

input

```
integrate((g*sec(f*x+e))**(3/2)*(a+a*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e)),x)
```

output

```
Integral(sqrt(a*(sec(e + f*x) + 1))*(g*sec(e + f*x))**(3/2)/(c + d*sec(e + f*x)), x)
```

3.239.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c + d \sec(e + fx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,
algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: sign: argument cannot be imagi
nary; found %i
```

3.239.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c + d \sec(e + fx)} dx = \int \frac{\sqrt{a \sec(fx + e) + a} (g \sec(fx + e))^{3/2}}{d \sec(fx + e) + c} dx$$

```
input integrate((g*sec(f*x+e))^(3/2)*(a+a*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,
algorithm="giac")
```

```
output sage0*x
```

3.239.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + a \sec(e + fx)}}{c + d \sec(e + fx)} dx = \int \frac{\sqrt{a + \frac{a}{\cos(e+fx)}} \left(\frac{g}{\cos(e+fx)} \right)^{3/2}}{c + \frac{d}{\cos(e+fx)}} dx$$

```
input int(((a + a/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c + d/cos(e + f*x
)),x)
```

```
output int(((a + a/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c + d/cos(e + f*x
)), x)
```

3.239. $\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+a \sec(e+fx)}}{c+d \sec(e+fx)} dx$

$$3.240 \quad \int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.240.1 Optimal result	1750
3.240.2 Mathematica [A] (verified)	1750
3.240.3 Rubi [A] (verified)	1751
3.240.4 Maple [B] (warning: unable to verify)	1753
3.240.5 Fracas [A] (verification not implemented)	1754
3.240.6 Sympy [F]	1754
3.240.7 Maxima [F]	1755
3.240.8 Giac [F(-2)]	1755
3.240.9 Mupad [F(-1)]	1755

3.240.1 Optimal result

Integrand size = 33, antiderivative size = 122

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)f} - \frac{2\sqrt{d} \arctan\left(\frac{\sqrt{a}\sqrt{d} \tan(e+fx)}{\sqrt{c+d}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)\sqrt{c+d}f}$$

output `arctan(1/2*a^(1/2)*tan(f*x+e)*2^(1/2)/(a+a*sec(f*x+e))^(1/2))*2^(1/2)/(c-d)/f/a^(1/2)-2*arctan(a^(1/2)*d^(1/2)*tan(f*x+e)/(c+d)^(1/2)/(a+a*sec(f*x+e))^(1/2))*d^(1/2)/(c-d)/f/a^(1/2)/(c+d)^(1/2)`

3.240.2 Mathematica [A] (verified)

Time = 1.43 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.18

$$\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{2\left(\sqrt{-c-d} \arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right)\right) + \sqrt{2}\sqrt{d} \operatorname{arctanh}\left(\frac{\sqrt{d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{-c-d}\sqrt{\frac{\cos(e+fx)}{1+\cos(e+fx)}}}\right)}{\sqrt{-c-d}(c-d)f\sqrt{\cos(e+fx) \sec^2\left(\frac{1}{2}(e+fx)\right)}\sqrt{a(1+\sec(e+fx))}}$$

input `Integrate[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(2*(Sqrt[-c - d]*ArcSin[Tan[(e + f*x)/2]] + Sqrt[2]*Sqrt[d]*ArcTanh[(Sqrt[d]*Tan[(e + f*x)/2])/(Sqrt[-c - d]*Sqrt[Cos[e + f*x]/(1 + Cos[e + f*x])])])/(Sqrt[-c - d]*(c - d)*f*Sqrt[Cos[e + f*x]*Sec[(e + f*x)/2]^2]*Sqrt[a*(1 + Sec[e + f*x])])`

3.240.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3042, 4460, 3042, 4282, 216, 4455, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)}{\sqrt{a \sec(e+fx) + a(c+d \sec(e+fx))}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx + \frac{\pi}{2})}{\sqrt{a \csc(e+fx + \frac{\pi}{2}) + a(c+d \csc(e+fx + \frac{\pi}{2}))}} dx \\
 & \quad \downarrow \text{4460} \\
 & \frac{\int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a+a}} dx}{c-d} - \frac{d \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{c+d \sec(e+fx)} dx}{a(c-d)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx + \frac{\pi}{2})}{\sqrt{\csc(e+fx + \frac{\pi}{2})a+a}} dx}{c-d} - \frac{d \int \frac{\csc(e+fx + \frac{\pi}{2})\sqrt{\csc(e+fx + \frac{\pi}{2})a+a}}{c+d \csc(e+fx + \frac{\pi}{2})} dx}{a(c-d)} \\
 & \quad \downarrow \text{4282} \\
 & - \frac{2 \int \frac{1}{\frac{a^2 \tan^2(e+fx)}{\sec(e+fx)a+a} + 2a} d\left(-\frac{a \tan(e+fx)}{\sqrt{\sec(e+fx)a+a}}\right)}{f(c-d)} - \frac{d \int \frac{\csc(e+fx + \frac{\pi}{2})\sqrt{\csc(e+fx + \frac{\pi}{2})a+a}}{c+d \csc(e+fx + \frac{\pi}{2})} dx}{a(c-d)} \\
 & \quad \downarrow \text{216} \\
 & \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2}\sqrt{a \sec(e+fx)+a}}\right)}{\sqrt{a}f(c-d)} - \frac{d \int \frac{\csc(e+fx + \frac{\pi}{2})\sqrt{\csc(e+fx + \frac{\pi}{2})a+a}}{c+d \csc(e+fx + \frac{\pi}{2})} dx}{a(c-d)}
 \end{aligned}$$

3.240. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

$$\begin{aligned}
& \downarrow 4455 \\
& \frac{2d \int \frac{1}{\frac{a^2 d \tan^2(e+fx)}{\sec(e+fx)a+a} + a(c+d)} d \left(-\frac{a \tan(e+fx)}{\sqrt{\sec(e+fx)a+a}} \right)}{f(c-d)} + \frac{\sqrt{2} \arctan \left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2} \sqrt{a \sec(e+fx)+a}} \right)}{\sqrt{a} f(c-d)} \\
& \downarrow 218 \\
& \frac{\sqrt{2} \arctan \left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2} \sqrt{a \sec(e+fx)+a}} \right)}{\sqrt{a} f(c-d)} - \frac{2\sqrt{d} \arctan \left(\frac{\sqrt{a} \sqrt{d} \tan(e+fx)}{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}} \right)}{\sqrt{a} f(c-d) \sqrt{c+d}}
\end{aligned}$$

input `Int[Sec[e + f*x]/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(Sqrt[2]*ArcTan[(Sqrt[a]*Tan[e + f*x])/(Sqrt[2]*Sqrt[a + a*Sec[e + f*x]])])/(Sqrt[a]*(c - d)*f) - (2*Sqrt[d]*ArcTan[(Sqrt[a]*Sqrt[d]*Tan[e + f*x])/(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])])/(Sqrt[a]*(c - d)*Sqrt[c + d]*f)`

3.240.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x])/Sqrt[a + b*Csc[e + f*x]]], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

```
rule 4455 Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(c
sc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d + d*x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x])]],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2,
0]
```

```
rule 4460 Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])*(cs
c[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[b/(b*c - a*d) Int
[Csc[e + f*x]/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[d/(b*c - a*d) Int[C
sc[e + f*x]*(Sqrt[a + b*Csc[e + f*x])/(c + d*Csc[e + f*x]), x], x] /; Free
Q[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && (EqQ[a^2 - b^2, 0] || EqQ[
c^2 - d^2, 0])
```

3.240.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 503 vs. 2(99) = 198.

Time = 19.78 (sec) , antiderivative size = 504, normalized size of antiderivative = 4.13

method	result
default	$\left(2\sqrt{(c+d)(c-d)} \ln\left(\frac{\csc(fx+e) - \cot(fx+e) + \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}}{2\sqrt{(c+d)(c-d)}}\right) + d\sqrt{2} \ln\left(-\frac{2\left(-\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2 - 1}\right)}{c-d}\right)\right)$

```
input int(sec(f*x+e)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVER
BOSE)
```

```
output 1/2/f/(d/(c-d))^(1/2)/(c-d)/((c+d)*(c-d))^(1/2)/a*(2*((c+d)*(c-d))^(1/2)*l
n(csc(f*x+e)-cot(f*x+e)+((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2))*(d/(c-d))
^(1/2)+d*2^(1/2)*ln(-2*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*2^(1/2)*(
d/(c-d))^(1/2)*c+2^(1/2)*(d/(c-d))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)
^(1/2)*d+((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))+c-d)/(c*(-cot(f*x+e)
+csc(f*x+e))-(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2))-d*2^(1/2)*ln
(-2*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*2^(1/2)*(d/(c-d))^(1/2)*c-2^(
1/2)*(d/(c-d))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*d+((c+d)*(c-d)
)^(1/2)*(-cot(f*x+e)+csc(f*x+e))-c+d)/(-c*(-cot(f*x+e)+csc(f*x+e))+(-cot(
f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2))))*((1-cos(f*x+e))^2*csc(f*x+e)^2
-1)^(1/2)*(-2*a/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)
```

$$3.240. \int \frac{\sec(e+fx)}{\sqrt{a+a\sec(e+fx)(c+d\sec(e+fx))}} dx$$

3.240.5 Fricas [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 963, normalized size of antiderivative = 7.89

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm=
"fricas")
```

```
output [-1/2*(sqrt(2)*sqrt(-1/a)*log((2*sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x
+ e))*sqrt(-1/a)*cos(f*x + e)*sin(f*x + e) + 3*cos(f*x + e)^2 + 2*cos(f*x
+ e) - 1)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + sqrt(-d/(a*c + a*d))*l
og(-((c^2 + 8*c*d + 8*d^2)*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 -
4*((c^2 + 3*c*d + 2*d^2)*cos(f*x + e)^2 - (c*d + d^2)*cos(f*x + e))*sqrt(
-d/(a*c + a*d))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sin(f*x + e) + d^2
- (6*c*d + 7*d^2)*cos(f*x + e))/(c^2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f
*x + e)^2 + d^2 + (2*c*d + d^2)*cos(f*x + e)))/((c - d)*f), -1/2*(sqrt(2)
*sqrt(-1/a)*log((2*sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(-1
/a)*cos(f*x + e)*sin(f*x + e) + 3*cos(f*x + e)^2 + 2*cos(f*x + e) - 1)/(co
s(f*x + e)^2 + 2*cos(f*x + e) + 1)) + 2*sqrt(d/(a*c + a*d))*arctan(2*(c +
d)*sqrt(d/(a*c + a*d))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*cos(f*x + e
)*sin(f*x + e)/((c + 2*d)*cos(f*x + e)^2 + (c + d)*cos(f*x + e) - d)))/((c
- d)*f), -1/2*(sqrt(-d/(a*c + a*d))*log(-((c^2 + 8*c*d + 8*d^2)*cos(f*x +
e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 - 4*((c^2 + 3*c*d + 2*d^2)*cos(f*x +
e)^2 - (c*d + d^2)*cos(f*x + e))*sqrt(-d/(a*c + a*d))*sqrt((a*cos(f*x + e)
+ a)/cos(f*x + e))*sin(f*x + e) + d^2 - (6*c*d + 7*d^2)*cos(f*x + e))/(c^
2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 + d^2 + (2*c*d + d^2)*cos(
f*x + e)) + 2*sqrt(2)*arctan(sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x +
e))*cos(f*x + e)/(sqrt(a)*sin(f*x + e)))/sqrt(a))/((c - d)*f), -(sqrt(d...
```

3.240.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx$$

$$= \int \frac{\sec(e + fx)}{\sqrt{a(\sec(e + fx) + 1)}(c + d \sec(e + fx))} dx$$

```
input integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x)
```

3.240. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

output `Integral(sec(e + f*x)/(sqrt(a*(sec(e + f*x) + 1))*(c + d*sec(e + f*x))), x)`

3.240.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\sec(fx + e)}{\sqrt{a \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/(sqrt(a*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.240.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Exception raised: TypeError}$$

input `integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:index.cc index_m i_lex_is_greater E rror: Bad Argument Value`

3.240.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sec(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx \\ &= \int \frac{1}{\cos(e + fx) \sqrt{a + \frac{a}{\cos(e + fx)}} \left(c + \frac{d}{\cos(e + fx)} \right)} dx \end{aligned}$$

3.240. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

input `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))),x)`

output `int(1/(cos(e + f*x)*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))), x)`

3.240. $\int \frac{\sec(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

$$3.241 \quad \int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.241.1 Optimal result	1757
3.241.2 Mathematica [A] (verified)	1757
3.241.3 Rubi [A] (verified)	1758
3.241.4 Maple [B] (warning: unable to verify)	1760
3.241.5 Fricas [A] (verification not implemented)	1761
3.241.6 Sympy [F]	1761
3.241.7 Maxima [F]	1762
3.241.8 Giac [F(-2)]	1762
3.241.9 Mupad [F(-1)]	1763

3.241.1 Optimal result

Integrand size = 35, antiderivative size = 124

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx = -\frac{\sqrt{2} \arctan\left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)f} + \frac{2c \arctan\left(\frac{\sqrt{a}\sqrt{d} \tan(e+fx)}{\sqrt{c+d}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)\sqrt{d}\sqrt{c+d}f}$$

output `-arctan(1/2*a^(1/2)*tan(f*x+e)*2^(1/2)/(a+a*sec(f*x+e))^(1/2))*2^(1/2)/(c-d)/f/a^(1/2)+2*c*arctan(a^(1/2)*d^(1/2)*tan(f*x+e)/(c+d)^(1/2)/(a+a*sec(f*x+e))^(1/2))/(c-d)/f/a^(1/2)/d^(1/2)/(c+d)^(1/2)`

3.241.2 Mathematica [A] (verified)

Time = 0.72 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.14

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx = -\frac{2\left(\sqrt{d}\sqrt{c+d} \arctan\left(\frac{\sin(\frac{1}{2}(e+fx))}{\sqrt{\cos(e+fx)}}\right) - \sqrt{2}c \arctan\left(\frac{\sqrt{2}\sqrt{d} \sin(\frac{1}{2}(e+fx))}{\sqrt{c+d}\sqrt{\cos(e+fx)}}\right)\right) \cos\left(\frac{1}{2}(e+fx)\right)}{(c-d)\sqrt{d}\sqrt{c+d}f\sqrt{\cos(e+fx)}\sqrt{a(1+\sec(e+fx))}}$$

input `Integrate[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

3.241. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

output $(-2*(\text{Sqrt}[d]*\text{Sqrt}[c + d]*\text{ArcTan}[\text{Sin}[(e + f*x)/2]/\text{Sqrt}[\text{Cos}[e + f*x]]] - \text{Sqrt}[2]*c*\text{ArcTan}[(\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sin}[(e + f*x)/2])/(\text{Sqrt}[c + d]*\text{Sqrt}[\text{Cos}[e + f*x]])])*\text{Cos}[(e + f*x)/2])/((c - d)*\text{Sqrt}[d]*\text{Sqrt}[c + d]*f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[a*(1 + \text{Sec}[e + f*x])])$

3.241.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4464, 3042, 4282, 216, 4455, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(e+fx)}{\sqrt{a \sec(e+fx) + a}(c+d \sec(e+fx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})^2}{\sqrt{a \csc(e+fx+\frac{\pi}{2}) + a}(c+d \csc(e+fx+\frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{4464} \\
 & \frac{c \int \frac{\sec(e+fx)\sqrt{\sec(e+fx)a+a}}{c+d \sec(e+fx)} dx}{a(c-d)} - \frac{\int \frac{\sec(e+fx)}{\sqrt{\sec(e+fx)a+a}} dx}{c-d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{c+d \csc(e+fx+\frac{\pi}{2})} dx}{a(c-d)} - \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx}{c-d} \\
 & \quad \downarrow \text{4282} \\
 & \frac{2 \int \frac{1}{\frac{a^2 \tan^2(e+fx)}{\sec(e+fx)a+a} + 2a} d\left(-\frac{a \tan(e+fx)}{\sqrt{\sec(e+fx)a+a}}\right)}{f(c-d)} + \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{c+d \csc(e+fx+\frac{\pi}{2})} dx}{a(c-d)} \\
 & \quad \downarrow \text{216} \\
 & \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{c+d \csc(e+fx+\frac{\pi}{2})} dx}{a(c-d)} - \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2}\sqrt{a \sec(e+fx)+a}}\right)}{\sqrt{a}f(c-d)} \\
 & \quad \downarrow \text{4455}
 \end{aligned}$$

3.241. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

$$\frac{2c \int \frac{1}{\frac{a^2 d \tan^2(e+fx)}{\sec(e+fx)a+a} + a(c+d)} d \left(-\frac{a \tan(e+fx)}{\sqrt{\sec(e+fx)a+a}} \right)}{f(c-d)} - \frac{\sqrt{2} \arctan \left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2} \sqrt{a \sec(e+fx)+a}} \right)}{\sqrt{a} f(c-d)}$$

↓ 218

$$\frac{2c \arctan \left(\frac{\sqrt{a} \sqrt{d} \tan(e+fx)}{\sqrt{c+d} \sqrt{a \sec(e+fx)+a}} \right)}{\sqrt{a} \sqrt{d} f(c-d) \sqrt{c+d}} - \frac{\sqrt{2} \arctan \left(\frac{\sqrt{a} \tan(e+fx)}{\sqrt{2} \sqrt{a \sec(e+fx)+a}} \right)}{\sqrt{a} f(c-d)}$$

input `Int[Sec[e + f*x]^2/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `-((Sqrt[2]*ArcTan[(Sqrt[a]*Tan[e + f*x])/(Sqrt[2]*Sqrt[a + a*Sec[e + f*x]])])/(Sqrt[a]*(c - d)*f)) + (2*c*ArcTan[(Sqrt[a]*Sqrt[d]*Tan[e + f*x])/(Sqrt[c + d]*Sqrt[a + a*Sec[e + f*x]])])/(Sqrt[a]*(c - d)*Sqrt[d]*Sqrt[c + d]*f)`

3.241.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4282 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2/f Subst[Int[1/(2*a + x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, e, f}, x] && EqQ[a^2 - b^2, 0]`

rule 4455 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d + d*x^2), x], x, b*(Cot[e + f*x]/Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0]`


```
rule 4464 Int[csc[(e_.) + (f_.)*(x_.)]^2/(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)]*(
csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))), x_Symbol] := Simp[-a/(b*c - a*d)
Int[Csc[e + f*x]/Sqrt[a + b*Csc[e + f*x]], x], x] + Simp[c/(b*c - a*d) In
t[Csc[e + f*x]*(Sqrt[a + b*Csc[e + f*x]]/(c + d*Csc[e + f*x])), x], x] /; F
reeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && (EqQ[a^2 - b^2, 0] || E
qQ[c^2 - d^2, 0])
```

3.241.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 503 vs. 2(101) = 202.

Time = 18.95 (sec) , antiderivative size = 504, normalized size of antiderivative = 4.06

method	result
default	$-\frac{\sqrt{-\frac{2a}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}} \sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2-1} \left(2\sqrt{(c+d)(c-d)} \ln\left(\csc(fx+e)-\cot(fx+e)+\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}\right) \right)}{1}$

```
input int(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/2/f/a/((c+d)*(c-d))^(1/2)/(c-d)/(d/(c-d))^(1/2)*(-2*a/((1-cos(f*x+e))^2
*csc(f*x+e)^2-1))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*(2*((c+d)*
(c-d))^(1/2)*ln(csc(f*x+e)-cot(f*x+e)+((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1
/2))*(d/(c-d))^(1/2)-c*2^(1/2)*ln(-2*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1
/2)*2^(1/2)*(d/(c-d))^(1/2)*c-2^(1/2)*(d/(c-d))^(1/2)*((1-cos(f*x+e))^2*cs
c(f*x+e)^2-1)^(1/2)*d+((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))-c+d)/(-
c*(-cot(f*x+e)+csc(f*x+e))+(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2)
)+c*2^(1/2)*ln(-2*(-((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2)*2^(1/2)*(d/(c-
d))^(1/2)*c+2^(1/2)*(d/(c-d))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)^(1/2
)*d+((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))+c-d)/(c*(-cot(f*x+e)+csc(
f*x+e))-(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2))))
```

3.241.
$$\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)(c+d \sec(e+fx))}} dx$$

3.241.5 Fricas [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 1041, normalized size of antiderivative = 8.40

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
output [-1/2*(sqrt(2)*(a*c*d + a*d^2)*sqrt(-1/a)*log(-(2*sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(-1/a)*cos(f*x + e)*sin(f*x + e) - 3*cos(f*x + e)^2 - 2*cos(f*x + e) + 1)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) - sqrt(-a*c*d - a*d^2)*c*log(-((a*c^2 + 8*a*c*d + 8*a*d^2)*cos(f*x + e)^3 + a*d^2 + (a*c^2 + 2*a*c*d)*cos(f*x + e)^2 - 4*sqrt(-a*c*d - a*d^2)*((c + 2*d)*cos(f*x + e)^2 - d*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sin(f*x + e) - (6*a*c*d + 7*a*d^2)*cos(f*x + e))/(c^2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 + d^2 + (2*c*d + d^2)*cos(f*x + e)))/((a*c^2*d - a*d^3)*f), -1/2*(sqrt(2)*(a*c*d + a*d^2)*sqrt(-1/a)*log(-(2*sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(-1/a)*cos(f*x + e)*sin(f*x + e) - 3*cos(f*x + e)^2 - 2*cos(f*x + e) + 1)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) - 2*sqrt(a*c*d + a*d^2)*c*arctan(2*sqrt(a*c*d + a*d^2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)/((a*c + 2*a*d)*cos(f*x + e)^2 - a*d + (a*c + a*d)*cos(f*x + e)))/((a*c^2*d - a*d^3)*f), 1/2*(sqrt(-a*c*d - a*d^2)*c*log(-((a*c^2 + 8*a*c*d + 8*a*d^2)*cos(f*x + e)^3 + a*d^2 + (a*c^2 + 2*a*c*d)*cos(f*x + e)^2 - 4*sqrt(-a*c*d - a*d^2)*((c + 2*d)*cos(f*x + e)^2 - d*cos(f*x + e))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sin(f*x + e) - (6*a*c*d + 7*a*d^2)*cos(f*x + e))/(c^2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 + d^2 + (2*c*d + d^2)*cos(f*x + e))) + 2*sqrt(2)*(a*c*d + a*d^2)*arctan(sqrt(2)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*co...
```

3.241.6 Sympy [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx$$

$$= \int \frac{\sec^2(e + fx)}{\sqrt{a(\sec(e + fx) + 1)}(c + d \sec(e + fx))} dx$$

```
input integrate(sec(f*x+e)**2/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x)
```

3.241. $\int \frac{\sec^2(e+fx)}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

output `Integral(sec(e + f*x)**2/(sqrt(a*(sec(e + f*x) + 1))*(c + d*sec(e + f*x))), x)`

3.241.7 Maxima [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\sec^2(fx + e)}{\sqrt{a \sec(fx + e) + a(d \sec(fx + e) + c)}} dx$$

input `integrate(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)^2/(sqrt(a*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.241.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Exception raised: TypeError}$$

input `integrate(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);;OUTPUT:index.cc index_m i_lex_is_greater Error: Bad Argument Value`

3.241.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx$$

$$= \int \frac{1}{\cos(e + fx)^2 \sqrt{a + \frac{a}{\cos(e + fx)}} \left(c + \frac{d}{\cos(e + fx)} \right)} dx$$

input `int(1/(cos(e + f*x)^2*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))),x)`output `int(1/(cos(e + f*x)^2*(a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))), x)`

$$3.242 \quad \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.242.1 Optimal result	1764
3.242.2 Mathematica [A] (verified)	1764
3.242.3 Rubi [A] (verified)	1765
3.242.4 Maple [B] (warning: unable to verify)	1767
3.242.5 Fracas [A] (verification not implemented)	1768
3.242.6 Sympy [F]	1768
3.242.7 Maxima [F]	1769
3.242.8 Giac [F(-2)]	1770
3.242.9 Mupad [F(-1)]	1770

3.242.1 Optimal result

Integrand size = 39, antiderivative size = 167

$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx =$$

$$\frac{\sqrt{2}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{2}\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)f}$$

$$+ \frac{2\sqrt{c}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)\sqrt{c+d}f}$$

output `-g^(3/2)*arctanh(1/2*a^(1/2)*g^(1/2)*tan(f*x+e)*2^(1/2)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))*2^(1/2)/(c-d)/f/a^(1/2)+2*g^(3/2)*arctanh(a^(1/2)*c^(1/2)*g^(1/2)*tan(f*x+e)/(c+d)^(1/2)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))*c^(1/2)/(c-d)/f/a^(1/2)/(c+d)^(1/2)`

3.242.2 Mathematica [A] (verified)

Time = 1.76 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.19

$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{g \cos\left(\frac{1}{2}(e+fx)\right) (2\sqrt{c+d} \log\left(\cos\left(\frac{1}{4}(e+fx)\right) - \sin\left(\frac{1}{4}(e+fx)\right)\right))}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))}$$

3.242. $\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

input `Integrate[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(g*Cos[(e + f*x)/2]*(2*Sqrt[c + d]*Log[Cos[(e + f*x)/4] - Sin[(e + f*x)/4]] - 2*Sqrt[c + d]*Log[Cos[(e + f*x)/4] + Sin[(e + f*x)/4]] + Sqrt[2]*Sqrt[c]*(-Log[Sqrt[2]*Sqrt[c + d] - 2*Sqrt[c]*Sin[(e + f*x)/2]] + Log[Sqrt[2]*Sqrt[c + d] + 2*Sqrt[c]*Sin[(e + f*x)/2]]))*Sqrt[g*Sec[e + f*x]]/((c - d)*Sqrt[c + d]*f*Sqrt[a*(1 + Sec[e + f*x])])`

3.242.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {3042, 4462, 3042, 4295, 221, 4453, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a \sec(e + fx) + a}(c + d \sec(e + fx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a}(c + d \csc(e + fx + \frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{4462} \\
 & \frac{cg \int \frac{\sqrt{g \sec(e + fx)} \sqrt{\sec(e + fx)a + a}}{c + d \sec(e + fx)} dx}{a(c - d)} - \frac{g \int \frac{\sqrt{g \sec(e + fx)}}{\sqrt{\sec(e + fx)a + a}} dx}{c - d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{cg \int \frac{\sqrt{g \csc(e + fx + \frac{\pi}{2})} \sqrt{\csc(e + fx + \frac{\pi}{2})a + a}}{c + d \csc(e + fx + \frac{\pi}{2})} dx}{a(c - d)} - \frac{g \int \frac{\sqrt{g \csc(e + fx + \frac{\pi}{2})}}{\sqrt{\csc(e + fx + \frac{\pi}{2})a + a}} dx}{c - d} \\
 & \quad \downarrow \text{4295} \\
 & \frac{2g^2 \int \frac{1}{2a - \frac{a^2 \sin(e + fx) \tan(e + fx)}{\sec(e + fx)a + a}} d\left(-\frac{a \tan(e + fx)}{\sqrt{g \sec(e + fx)} \sqrt{\sec(e + fx)a + a}}\right)}{f(c - d)} + \frac{cg \int \frac{\sqrt{g \csc(e + fx + \frac{\pi}{2})} \sqrt{\csc(e + fx + \frac{\pi}{2})a + a}}{c + d \csc(e + fx + \frac{\pi}{2})} dx}{a(c - d)} \\
 & \quad \downarrow \text{221}
 \end{aligned}$$

3.242. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx$

$$\begin{aligned}
 & \frac{cg \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})} \sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{c+d \csc(e+fx+\frac{\pi}{2})} dx}{a(c-d)} - \frac{\sqrt{2}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{2}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{a}f(c-d)} \\
 & \quad \downarrow \text{4453} \\
 & - \frac{2cg^2 \int \frac{1}{a(c+d) - \frac{a^2 \csc(e+fx) \tan(e+fx)}{\sec(e+fx)a+a}} d\left(-\frac{a \tan(e+fx)}{\sqrt{g \sec(e+fx)}\sqrt{\sec(e+fx)a+a}}\right)}{f(c-d)} - \frac{\sqrt{2}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{2}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{a}f(c-d)} \\
 & \quad \downarrow \text{221} \\
 & \frac{2\sqrt{c}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{a}f(c-d)\sqrt{c+d}} - \frac{\sqrt{2}g^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{2}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{a}f(c-d)}
 \end{aligned}$$

input `Int[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `-((Sqrt[2]*g^(3/2)*ArcTanh[(Sqrt[a]*Sqrt[g]*Tan[e + f*x])/(Sqrt[2]*Sqrt[g*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])]/(Sqrt[a]*(c - d)*f)) + (2*Sqrt[c]*g^(3/2)*ArcTanh[(Sqrt[a]*Sqrt[c]*Sqrt[g]*Tan[e + f*x])/(Sqrt[c + d]*Sqrt[g*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])]/(Sqrt[a]*(c - d)*Sqrt[c + d]*f))`

3.242.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4295 `Int[Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_. + (a_.))], x_Symbol] := Simp[-2*b*(d/(a*f)) Subst[Int[1/(2*b - d*x^2), x], x, b*(Cot[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[d*Csc[e + f*x]])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0]`

3.242. $\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

rule 4453 `Int[(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(g_.)]*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)])/(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)), x_Symbol] := Simp[-2*b*(g/f) Subst[Int[1/(b*c + a*d - c*g*x^2), x], x, b*(Cot[e + f*x]/(Sqrt[g*Csc[e + f*x]]*Sqrt[a + b*Csc[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0]`

rule 4462 `Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(3/2)/(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))), x_Symbol] := Simp[(-a)*(g/(b*c - a*d)) Int[Sqrt[g*Csc[e + f*x]]/Sqrt[a + b*Csc[e + f*x]], x], x] + Simp[c*(g/(b*c - a*d)) Int[Sqrt[g*Csc[e + f*x]]*(Sqrt[a + b*Csc[e + f*x]])/(c + d*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0]`

3.242.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 556 vs. 2(132) = 264.

Time = 20.66 (sec) , antiderivative size = 557, normalized size of antiderivative = 3.34

method	result
default	$g \sqrt{-\frac{g((1-\cos(fx+e))^2 \csc(fx+e)^2+1)}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}} \left((1-\cos(fx+e))^2 \csc(fx+e)^2-1 \right) \sqrt{-\frac{2a}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1}} \left(c\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{(1-\cos(fx+e))^2 \csc(fx+e)^2+1}}{(1-\cos(fx+e))^2 \csc(fx+e)^2-1} \right) \right)$

input `int((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*g/f/a/(c/(c-d))^(1/2)/(c-d)/((c+d)*(c-d))^(1/2)*(-g*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)*((1-cos(f*x+e))^2*csc(f*x+e)^2-1)*(-2*a/((1-cos(f*x+e))^2*csc(f*x+e)^2-1))^(1/2)*(c^2^(1/2)*ln(2*(2^(1/2))*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*c-2^(1/2))*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*d-((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))+c-d)/(c*(-cot(f*x+e)+csc(f*x+e))-(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2))-c^2^(1/2)*ln(-2*(2^(1/2))*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*c-2^(1/2))*((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)*(c/(c-d))^(1/2)*d+((c+d)*(c-d))^(1/2)*(-cot(f*x+e)+csc(f*x+e))+c-d)/(-c*(-cot(f*x+e)+csc(f*x+e))+(-cot(f*x+e)+csc(f*x+e))*d+((c+d)*(c-d))^(1/2))-2*((c+d)*(c-d))^(1/2)*arcsinh(cot(f*x+e)-csc(f*x+e))*(c/(c-d))^(1/2))/((1-cos(f*x+e))^2*csc(f*x+e)^2+1)^(1/2)`

$$3.242. \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.242.5 Fricas [A] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 1103, normalized size of antiderivative = 6.60

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Too large to display}$$

```
input integrate((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="fricas")
```

```
output [-1/2*(sqrt(2)*g*sqrt(g/a)*log((2*sqrt(2)*sqrt(g/a)*sqrt((a*cos(f*x + e) +
a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) - g*cos(f
*x + e)^2 + 2*g*cos(f*x + e) + 3*g)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1))
+ sqrt(c*g/(a*c + a*d))*g*log((c^2*g*cos(f*x + e)^3 - (7*c^2 + 6*c*d)*g*cos
os(f*x + e)^2 + 4*((c^2 + c*d)*cos(f*x + e)^2 - (2*c^2 + 3*c*d + d^2)*cos(
f*x + e))*sqrt(c*g/(a*c + a*d))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sq
rt(g/cos(f*x + e))*sin(f*x + e) + (2*c*d + d^2)*g*cos(f*x + e) + (8*c^2 +
8*c*d + d^2)*g)/(c^2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 + d^2 +
(2*c*d + d^2)*cos(f*x + e)))/((c - d)*f), 1/2*(2*sqrt(2)*g*sqrt(-g/a)*ar
ctan(sqrt(2)*sqrt(-g/a)*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos
(f*x + e))*cos(f*x + e)/(g*sin(f*x + e))) - sqrt(c*g/(a*c + a*d))*g*log((c
^2*g*cos(f*x + e)^3 - (7*c^2 + 6*c*d)*g*cos(f*x + e)^2 + 4*((c^2 + c*d)*co
s(f*x + e)^2 - (2*c^2 + 3*c*d + d^2)*cos(f*x + e))*sqrt(c*g/(a*c + a*d))*s
qrt((a*cos(f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) +
(2*c*d + d^2)*g*cos(f*x + e) + (8*c^2 + 8*c*d + d^2)*g)/(c^2*cos(f*x + e)
^3 + (c^2 + 2*c*d)*cos(f*x + e)^2 + d^2 + (2*c*d + d^2)*cos(f*x + e)))/((
c - d)*f), -1/2*(sqrt(2)*g*sqrt(g/a)*log((2*sqrt(2)*sqrt(g/a)*sqrt((a*cos(
f*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e)
- g*cos(f*x + e)^2 + 2*g*cos(f*x + e) + 3*g)/(cos(f*x + e)^2 + 2*cos(f*x
+ e) + 1)) - 2*sqrt(-c*g/(a*c + a*d))*g*arctan(1/2*(c*cos(f*x + e)^2 - ...
```

3.242.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(e + fx))^{\frac{3}{2}}}{\sqrt{a (\sec(e + fx) + 1)}(c + d \sec(e + fx))} dx$$

```
input integrate((g*sec(f*x+e))**(3/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x
)
```

3.242. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx$

output `Integral((g*sec(e + f*x))**(3/2)/(sqrt(a*(sec(e + f*x) + 1))*(c + d*sec(e + f*x))), x)`

3.242.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{\frac{3}{2}}}{\sqrt{a \sec(fx + e) + a(d \sec(fx + e) + c)}} dx$$

input `integrate((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `1/2*(sqrt(2)*c*f*g*integrate(((c^2*cos(2*f*x + 2*e))^2 + c^2*sin(2*f*x + 2*e))^2 - 2*(c*d - 2*d^2)*cos(f*x + e)^2 - (c^2 - 4*c*d)*sin(2*f*x + 2*e)*sin(f*x + e) - 2*(c*d - 2*d^2)*sin(f*x + e)^2 + (c^2 - (c^2 - 4*c*d)*cos(f*x + e))*cos(2*f*x + 2*e) - (c^2 - 2*c*d)*cos(f*x + e))*cos(1/2*arctan2(sin(f*x + e), cos(f*x + e))) - (c^2*cos(2*f*x + 2*e)*sin(f*x + e) - (c^2*cos(f*x + e) + c^2)*sin(2*f*x + 2*e) + (c^2 - 2*c*d)*sin(f*x + e))*sin(1/2*arctan2(sin(f*x + e), cos(f*x + e))))/((c^2*cos(2*f*x + 2*e))^2 + 4*d^2*cos(f*x + e)^2 + c^2*sin(2*f*x + 2*e)^2 + 4*c*d*sin(2*f*x + 2*e)*sin(f*x + e) + 4*d^2*sin(f*x + e)^2 + 4*c*d*cos(f*x + e) + c^2 + 2*(2*c*d*cos(f*x + e) + c^2)*cos(2*f*x + 2*e))*cos(1/2*arctan2(sin(f*x + e), cos(f*x + e)))^2 + (c^2*cos(2*f*x + 2*e))^2 + 4*d^2*cos(f*x + e)^2 + c^2*sin(2*f*x + 2*e)^2 + 4*c*d*sin(2*f*x + 2*e)*sin(f*x + e) + 4*d^2*sin(f*x + e)^2 + 4*c*d*cos(f*x + e) + c^2 + 2*(2*c*d*cos(f*x + e) + c^2)*cos(2*f*x + 2*e))*sin(1/2*arctan2(sin(f*x + e), cos(f*x + e)))^2), x) + sqrt(2)*c*f*g*integrate(((2*c*d*cos(f*x + e))^2 + 2*c*d*sin(f*x + e)^2 - (c^2 - 2*c*d)*cos(2*f*x + 2*e))^2 + c^2*cos(f*x + e) - (c^2 - 2*c*d)*sin(2*f*x + 2*e))^2 + (c^2 - 2*c*d + 4*d^2)*sin(2*f*x + 2*e)*sin(f*x + e) - (c^2 - 2*c*d - (c^2 - 2*c*d + 4*d^2)*cos(f*x + e))*cos(2*f*x + 2*e))*cos(1/2*arctan2(sin(f*x + e), cos(f*x + e))) + (c^2*sin(f*x + e) + (c^2 + 2*c*d - 4*d^2)*cos(2*f*x + 2*e)*sin(f*x + e) - (c^2 - 2*c*d + (c^2 + 2*c*d - 4*d^2)*cos(f*x + e))*sin(2*f*x + 2*e))*sin(...`

3.242.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Exception raised: TypeError}$$

input `integrate((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:index.cc index_m i_lex_is_greater E
rror: Bad Argument Value`

3.242.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{3/2}}{\sqrt{a + \frac{a}{\cos(e+fx)}} \left(c + \frac{d}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(3/2)/((a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x)
)),x)`

output `int((g/cos(e + f*x))^(3/2)/((a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x)
)), x)`

$$3.243 \quad \int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.243.1 Optimal result	1771
3.243.2 Mathematica [A] (verified)	1772
3.243.3 Rubi [A] (verified)	1772
3.243.4 Maple [B] (warning: unable to verify)	1776
3.243.5 Fricas [A] (verification not implemented)	1777
3.243.6 Sympy [F(-1)]	1777
3.243.7 Maxima [F]	1778
3.243.8 Giac [F(-2)]	1779
3.243.9 Mupad [F(-1)]	1779

3.243.1 Optimal result

Integrand size = 39, antiderivative size = 231

$$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{2g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}df}$$

$$+ \frac{\sqrt{2}g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g} \tan(e+fx)}{\sqrt{2}\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)f}$$

$$- \frac{2c^{3/2}g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{g \sec(e+fx)}\sqrt{a+a \sec(e+fx)}}\right)}{\sqrt{a}(c-d)d\sqrt{c+df}}$$

```
output 2*g^(5/2)*arctanh(a^(1/2)*g^(1/2)*tan(f*x+e)/(g*sec(f*x+e))^(1/2)/(a+a*sec
(f*x+e))^(1/2))/d/f/a^(1/2)+g^(5/2)*arctanh(1/2*a^(1/2)*g^(1/2)*tan(f*x+e)
*2^(1/2)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))*2^(1/2)/(c-d)/f/a^(1
/2)-2*c^(3/2)*g^(5/2)*arctanh(a^(1/2)*c^(1/2)*g^(1/2)*tan(f*x+e)/(c+d)^(1/
2)/(g*sec(f*x+e))^(1/2)/(a+a*sec(f*x+e))^(1/2))/(c-d)/d/f/a^(1/2)/(c+d)^(1
/2)
```

3.243.2 Mathematica [A] (verified)

Time = 1.49 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.67

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \frac{2g^2 \left(d\sqrt{c+d} \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(e+fx)\right)\right) + \sqrt{2}\left((c-d)\sqrt{c+d}\right) \right)}{\dots}$$

input `Integrate[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(2*g^2*(d*Sqrt[c + d]*ArcTanh[Sin[(e + f*x)/2]] + Sqrt[2]*((c - d)*Sqrt[c + d]*ArcTanh[Sqrt[2]*Sin[(e + f*x)/2]] - c^(3/2)*ArcTanh[(Sqrt[2]*Sqrt[c]*Sin[(e + f*x)/2])/Sqrt[c + d]]))*Cos[(e + f*x)/2]*Sqrt[g*Sec[e + f*x]]/((c - d)*d*Sqrt[c + d]*f*Sqrt[a*(1 + Sec[e + f*x])])`

3.243.3 Rubi [A] (verified)Time = 1.41 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$, Rules used = {3042, 4466, 3042, 4453, 221, 4511, 3042, 4289, 221, 4295, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a \sec(e + fx) + a}(c + d \sec(e + fx))} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{5/2}}{\sqrt{a \csc(e + fx + \frac{\pi}{2}) + a}(c + d \csc(e + fx + \frac{\pi}{2}))} dx \\ & \quad \downarrow \text{4466} \\ & \frac{g^2 \int \frac{\sqrt{g \sec(e+fx)}(ac+a(c-d) \sec(e+fx))}{\sqrt{\sec(e+fx)a+a}} dx}{ad(c-d)} - \frac{c^2 g^2 \int \frac{\sqrt{g \sec(e+fx)} \sqrt{\sec(e+fx)a+a}}{c+d \sec(e+fx)} dx}{ad(c-d)} \\ & \quad \downarrow \text{3042} \\ & \frac{g^2 \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})}(ac+a(c-d) \csc(e+fx+\frac{\pi}{2}))}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx}{ad(c-d)} - \frac{c^2 g^2 \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})} \sqrt{\csc(e+fx+\frac{\pi}{2})a+a}}{c+d \csc(e+fx+\frac{\pi}{2})} dx}{ad(c-d)} \end{aligned}$$

3.243. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

$$\begin{aligned}
& \downarrow 4453 \\
& \frac{2c^2 g^3 \int \frac{1}{a(c+d) - \frac{a^2 c \sin(e+fx) \tan(e+fx)}{\sec(e+fx)a+a}} d\left(-\frac{a \tan(e+fx)}{\sqrt{g \sec(e+fx)} \sqrt{\sec(e+fx)a+a}}\right)}{df(c-d)} + \\
& \frac{g^2 \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})} (ac+a(c-d) \csc(e+fx+\frac{\pi}{2}))}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx}{ad(c-d)} \\
& \downarrow 221 \\
& \frac{g^2 \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})} (ac+a(c-d) \csc(e+fx+\frac{\pi}{2}))}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx}{ad(c-d)} - \frac{2c^{3/2} g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{adf}(c-d)\sqrt{c+d}} \\
& \downarrow 4511 \\
& \frac{g^2 \left((c-d) \int \sqrt{g \sec(e+fx)} \sqrt{\sec(e+fx)a+a} dx + ad \int \frac{\sqrt{g \sec(e+fx)}}{\sqrt{\sec(e+fx)a+a}} dx \right)}{ad(c-d)} - \\
& \frac{2c^{3/2} g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{adf}(c-d)\sqrt{c+d}} \\
& \downarrow 3042 \\
& \frac{g^2 \left((c-d) \int \sqrt{g \csc(e+fx+\frac{\pi}{2})} \sqrt{\csc(e+fx+\frac{\pi}{2})a+a} dx + ad \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})}}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx \right)}{ad(c-d)} - \\
& \frac{2c^{3/2} g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{adf}(c-d)\sqrt{c+d}} \\
& \downarrow 4289 \\
& \frac{g^2 \left(ad \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})}}{\sqrt{\csc(e+fx+\frac{\pi}{2})a+a}} dx - \frac{2ag(c-d) \int \frac{1}{a - \frac{a^2 \sin(e+fx) \tan(e+fx)}{\sec(e+fx)a+a}} d\left(-\frac{a \tan(e+fx)}{\sqrt{g \sec(e+fx)} \sqrt{\sec(e+fx)a+a}}\right)}{f} \right)}{ad(c-d)} - \\
& \frac{2c^{3/2} g^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g} \tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{adf}(c-d)\sqrt{c+d}} \\
& \downarrow 221
\end{aligned}$$

3.243. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$

$$\begin{aligned}
& g^2 \left(ad \int \frac{\sqrt{g \csc(e+fx+\frac{\pi}{2})}}{\sqrt{\csc(e+fx+\frac{\pi}{2})} a+a} dx + \frac{2\sqrt{a}\sqrt{g}(c-d)\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g}\tan(e+fx)}{\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{f} \right) \\
& \frac{2c^{3/2}g^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g}\tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{a}df(c-d)\sqrt{c+d}} \\
& \quad \downarrow 4295 \\
& g^2 \left(\frac{2\sqrt{a}\sqrt{g}(c-d)\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g}\tan(e+fx)}{\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{f} - \frac{2adg \int \frac{1}{2a-\frac{a^2 \sin(e+fx)\tan(e+fx)}{\sec(e+fx)a+a}} d\left(-\frac{a \tan(e+fx)}{\sqrt{g \sec(e+fx)}\sqrt{\sec(e+fx)a+a}}\right)}{f} \right) \\
& \frac{2c^{3/2}g^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g}\tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{a}df(c-d)\sqrt{c+d}} \\
& \quad \downarrow 221 \\
& g^2 \left(\frac{2\sqrt{a}\sqrt{g}(c-d)\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g}\tan(e+fx)}{\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{f} + \frac{\sqrt{2}\sqrt{ad}\sqrt{g}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{g}\tan(e+fx)}{\sqrt{2}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{f} \right) \\
& \frac{2c^{3/2}g^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{a}\sqrt{c}\sqrt{g}\tan(e+fx)}{\sqrt{c+d}\sqrt{a \sec(e+fx)+a}\sqrt{g \sec(e+fx)}}\right)}{\sqrt{a}df(c-d)\sqrt{c+d}}
\end{aligned}$$

input `Int[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + a*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(g^2*((2*Sqrt[a]*(c - d)*Sqrt[g]*ArcTanh[(Sqrt[a]*Sqrt[g]*Tan[e + f*x])/(Sqrt[g*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])])/f + (Sqrt[2]*Sqrt[a]*d*Sqrt[g]*ArcTanh[(Sqrt[a]*Sqrt[g]*Tan[e + f*x])/(Sqrt[2]*Sqrt[g*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])])/f)/(a*(c - d)*d) - (2*c^(3/2)*g^(5/2)*ArcTanh[(Sqrt[a]*Sqrt[c]*Sqrt[g]*Tan[e + f*x])/(Sqrt[c + d]*Sqrt[g*Sec[e + f*x]]*Sqrt[a + a*Sec[e + f*x]])]/(Sqrt[a]*(c - d)*d*Sqrt[c + d]*f)`

3.243.3.1 Defintions of rubi rules used

- rule 221 $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) * \text{ArcTanh}[x / \text{Rt}[-a/b, 2]], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b]$
- rule 3042 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4289 $\text{Int}[\text{Sqrt}[\text{csc}[e_+] + (f_+)(x_+)] * (d_+) * \text{Sqrt}[\text{csc}[e_+] + (f_+)(x_+) * (b_+ + a_+)], x_Symbol] \rightarrow \text{Simp}[-2 * b * (d/f) \ \text{Subst}[\text{Int}[1/(b - d * x^2), x], x, b * (\text{Cot}[e + f * x] / (\text{Sqrt}[a + b * \text{Csc}[e + f * x]] * \text{Sqrt}[d * \text{Csc}[e + f * x]]))], x] /;$ $\text{FreeQ}\{a, b, d, e, f, x\} \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ !\text{GtQ}[a * (d/b), 0]$
- rule 4295 $\text{Int}[\text{Sqrt}[\text{csc}[e_+] + (f_+)(x_+)] * (d_+) / \text{Sqrt}[\text{csc}[e_+] + (f_+)(x_+) * (b_+ + a_+)], x_Symbol] \rightarrow \text{Simp}[-2 * b * (d/(a * f)) \ \text{Subst}[\text{Int}[1/(2 * b - d * x^2), x], x, b * (\text{Cot}[e + f * x] / (\text{Sqrt}[a + b * \text{Csc}[e + f * x]] * \text{Sqrt}[d * \text{Csc}[e + f * x]]))], x] /;$ $\text{FreeQ}\{a, b, d, e, f, x\} \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$
- rule 4453 $\text{Int}[(\text{Sqrt}[\text{csc}[e_+] + (f_+)(x_+)] * (g_+) * \text{Sqrt}[\text{csc}[e_+] + (f_+)(x_+) * (b_+ + a_+)]) / (\text{csc}[e_+] + (f_+)(x_+) * (d_+) + (c_+)), x_Symbol] \rightarrow \text{Simp}[-2 * b * (g/f) \ \text{Subst}[\text{Int}[1/(b * c + a * d - c * g * x^2), x], x, b * (\text{Cot}[e + f * x] / (\text{Sqrt}[g * \text{Csc}[e + f * x]] * \text{Sqrt}[a + b * \text{Csc}[e + f * x]]))], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, g, x\} \ \&\& \ \text{NeQ}[b * c - a * d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$
- rule 4466 $\text{Int}[(\text{csc}[e_+] + (f_+)(x_+) * (g_+))^{5/2} / (\text{Sqrt}[\text{csc}[e_+] + (f_+)(x_+) * (b_+ + a_+)] * (\text{csc}[e_+] + (f_+)(x_+) * (d_+) + (c_+))), x_Symbol] \rightarrow \text{Simp}[(-c^2) * (g^2 / (d * (b * c - a * d))) \ \text{Int}[\text{Sqrt}[g * \text{Csc}[e + f * x]] * (\text{Sqrt}[a + b * \text{Csc}[e + f * x]] / (c + d * \text{Csc}[e + f * x])), x], x] + \text{Simp}[g^2 / (d * (b * c - a * d)) \ \text{Int}[\text{Sqrt}[g * \text{Csc}[e + f * x]] * ((a * c + (b * c - a * d) * \text{Csc}[e + f * x]) / \text{Sqrt}[a + b * \text{Csc}[e + f * x]]), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, g, x\} \ \&\& \ \text{NeQ}[b * c - a * d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$
- rule 4511 $\text{Int}[(\text{csc}[e_+] + (f_+)(x_+) * (d_+))^{(n_+)} * (\text{csc}[e_+] + (f_+)(x_+) * (b_+ + a_+))^{(m_+)} * (\text{csc}[e_+] + (f_+)(x_+) * (B_+) + (A_+)), x_Symbol] \rightarrow \text{Simp}[(A * b - a * B) / b \ \text{Int}[(a + b * \text{Csc}[e + f * x])^{(m_+)} * (d * \text{Csc}[e + f * x])^{(n_+)}, x], x] + \text{Simp}[B / b \ \text{Int}[(a + b * \text{Csc}[e + f * x])^{(m_+ + 1)} * (d * \text{Csc}[e + f * x])^{(n_+)}, x], x] /;$ $\text{FreeQ}\{a, b, d, e, f, A, B, m\}, x\} \ \&\& \ \text{NeQ}[A * b - a * B, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

3.243.
$$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)(c+d \sec(e+fx))}} dx$$

3.243.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 795 vs. $2(184) = 368$.

Time = 22.87 (sec) , antiderivative size = 796, normalized size of antiderivative = 3.45

method	result
default	$-\frac{2\left(\sqrt{(c+d)(c-d)} \operatorname{arcsinh}(\cot(fx+e)-\csc(fx+e))\sqrt{\frac{c}{c-d}}d\sqrt{2}-\operatorname{arctanh}\left(\frac{\cos(fx+e)+\sin(fx+e)+1}{2(\cos(fx+e)+1)\sqrt{\frac{1}{\cos(fx+e)+1}}}\right)\right)\sqrt{\frac{c}{c-d}}\sqrt{(c+d)(c-d)}c+}{}$

```
input int((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,method=
_RETURNVERBOSE)
```

```
output -2/f/a/(c/(c-d))^(1/2)/((c+d)*(c-d))^(1/2)/(-c+d+((c+d)*(c-d))^(1/2))/(c-d
+((c+d)*(c-d))^(1/2))*(((c+d)*(c-d))^(1/2)*arcsinh(cot(f*x+e)-csc(f*x+e))*
(c/(c-d))^(1/2)*d*2^(1/2)-arctanh(1/2*(cos(f*x+e)+sin(f*x+e)+1)/(cos(f*x+e
)+1)/(1/(cos(f*x+e)+1))^(1/2))*(c/(c-d))^(1/2)*((c+d)*(c-d))^(1/2)*c+arcta
nh(1/2*(cos(f*x+e)+sin(f*x+e)+1)/(cos(f*x+e)+1)/(1/(cos(f*x+e)+1))^(1/2))*
(c/(c-d))^(1/2)*((c+d)*(c-d))^(1/2)*d-arctanh(1/2*(-cos(f*x+e)+sin(f*x+e)-
1)/(cos(f*x+e)+1)/(1/(cos(f*x+e)+1))^(1/2))*(c/(c-d))^(1/2)*((c+d)*(c-d))^(
1/2)*c+arctanh(1/2*(-cos(f*x+e)+sin(f*x+e)-1)/(cos(f*x+e)+1)/(1/(cos(f*x+
e)+1))^(1/2))*(c/(c-d))^(1/2)*((c+d)*(c-d))^(1/2)*d+ln(-2*(2*(1/(cos(f*x+e
)+1))^(1/2)*(c/(c-d))^(1/2)*c*sin(f*x+e)-2*(1/(cos(f*x+e)+1))^(1/2)*(c/(c-
d))^(1/2)*d*sin(f*x+e)+sin(f*x+e)*c-sin(f*x+e)*d-((c+d)*(c-d))^(1/2)*cos(f
*x+e)+((c+d)*(c-d))^(1/2))/(((c+d)*(c-d))^(1/2)*sin(f*x+e)+c*cos(f*x+e)-d*
cos(f*x+e)-c+d))*c^2-ln(-2*(-2*(c/(c-d))^(1/2)*(1/(cos(f*x+e)+1))^(1/2)*c*
cos(f*x+e)+2*(c/(c-d))^(1/2)*(1/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)*d-2*(1/(c
os(f*x+e)+1))^(1/2)*(c/(c-d))^(1/2)*c+2*(1/(cos(f*x+e)+1))^(1/2)*(c/(c-d))
^(1/2)*d+((c+d)*(c-d))^(1/2)*sin(f*x+e)-c*cos(f*x+e)+d*cos(f*x+e)-c+d)/(((
c+d)*(c-d))^(1/2)*cos(f*x+e)+sin(f*x+e)*c-sin(f*x+e)*d+((c+d)*(c-d))^(1/2)
))*c^2)*(a*(sec(f*x+e)+1))^(1/2)*(g*sec(f*x+e))^(1/2)*g^2/(cos(f*x+e)+1)/(
1/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)
```

$$3.243. \int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+a \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.243.5 Fricas [A] (verification not implemented)

Time = 52.75 (sec) , antiderivative size = 1597, normalized size of antiderivative = 6.91

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Too large to display}$$

```
input integrate((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="fricas")
```

```
output [-1/2*(sqrt(2)*d*g^2*sqrt(g/a)*log(-(2*sqrt(2)*sqrt(g/a)*sqrt((a*cos(f*x +
e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) + g*
cos(f*x + e)^2 - 2*g*cos(f*x + e) - 3*g)/(cos(f*x + e)^2 + 2*cos(f*x + e)
+ 1)) + c*sqrt(c*g/(a*c + a*d))*g^2*log((c^2*g*cos(f*x + e)^3 - (7*c^2 + 6
*c*d)*g*cos(f*x + e)^2 - 4*((c^2 + c*d)*cos(f*x + e)^2 - (2*c^2 + 3*c*d +
d^2)*cos(f*x + e))*sqrt(c*g/(a*c + a*d))*sqrt((a*cos(f*x + e) + a)/cos(f*x
+ e))*sqrt(g/cos(f*x + e))*sin(f*x + e) + (2*c*d + d^2)*g*cos(f*x + e) +
(8*c^2 + 8*c*d + d^2)*g)/(c^2*cos(f*x + e)^3 + (c^2 + 2*c*d)*cos(f*x + e)^
2 + d^2 + (2*c*d + d^2)*cos(f*x + e))) - (c - d)*g^2*sqrt(g/a)*log((g*cos(
f*x + e)^3 - 4*(cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt(g/a)*sqrt((a*cos(f*x
+ e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) - 7*g*cos(f*x +
e)^2 + 8*g)/(cos(f*x + e)^3 + cos(f*x + e)^2)))/((c*d - d^2)*f), -1/2*(sq
rt(2)*d*g^2*sqrt(g/a)*log(-(2*sqrt(2)*sqrt(g/a)*sqrt((a*cos(f*x + e) + a)/
cos(f*x + e))*sqrt(g/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) + g*cos(f*x +
e)^2 - 2*g*cos(f*x + e) - 3*g)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + 2
*c*sqrt(-c*g/(a*c + a*d))*g^2*arctan(1/2*(c*cos(f*x + e)^2 - (2*c + d)*cos
(f*x + e))*sqrt(-c*g/(a*c + a*d))*sqrt((a*cos(f*x + e) + a)/cos(f*x + e))*
sqrt(g/cos(f*x + e))/(c*g*sin(f*x + e))) - (c - d)*g^2*sqrt(g/a)*log((g*co
s(f*x + e)^3 - 4*(cos(f*x + e)^2 - 2*cos(f*x + e))*sqrt(g/a)*sqrt((a*cos(f
*x + e) + a)/cos(f*x + e))*sqrt(g/cos(f*x + e))*sin(f*x + e) - 7*g*cos(...
```

3.243.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Timed out}$$

```
input integrate((g*sec(f*x+e))**(5/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))**(1/2),x
)
```

3.243. $\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx$

output Timed out

3.243.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{5/2}}{\sqrt{a \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="maxima")`

output

```
-1/2*(sqrt(2)*c^2*f*g^2*integrate(((c^2*cos(2*f*x + 2*e))^2 + c^2*sin(2*f*x
+ 2*e)^2 - 2*(c*d - 2*d^2)*cos(f*x + e)^2 - (c^2 - 4*c*d)*sin(2*f*x + 2*e
)*sin(f*x + e) - 2*(c*d - 2*d^2)*sin(f*x + e)^2 + (c^2 - (c^2 - 4*c*d)*cos
(f*x + e))*cos(2*f*x + 2*e) - (c^2 - 2*c*d)*cos(f*x + e))*cos(1/2*arctan2(
sin(f*x + e), cos(f*x + e))) - (c^2*cos(2*f*x + 2*e)*sin(f*x + e) - (c^2*c
os(f*x + e) + c^2)*sin(2*f*x + 2*e) + (c^2 - 2*c*d)*sin(f*x + e))*sin(1/2*
arctan2(sin(f*x + e), cos(f*x + e))))/((c^2*cos(2*f*x + 2*e))^2 + 4*d^2*cos
(f*x + e)^2 + c^2*sin(2*f*x + 2*e)^2 + 4*c*d*sin(2*f*x + 2*e)*sin(f*x + e)
+ 4*d^2*sin(f*x + e)^2 + 4*c*d*cos(f*x + e) + c^2 + 2*(2*c*d*cos(f*x + e)
+ c^2)*cos(2*f*x + 2*e))*cos(1/2*arctan2(sin(f*x + e), cos(f*x + e)))^2 +
(c^2*cos(2*f*x + 2*e))^2 + 4*d^2*cos(f*x + e)^2 + c^2*sin(2*f*x + 2*e)^2 +
4*c*d*sin(2*f*x + 2*e)*sin(f*x + e) + 4*d^2*sin(f*x + e)^2 + 4*c*d*cos(f*
x + e) + c^2 + 2*(2*c*d*cos(f*x + e) + c^2)*cos(2*f*x + 2*e))*sin(1/2*arct
an2(sin(f*x + e), cos(f*x + e)))^2), x) + sqrt(2)*c^2*f*g^2*integrate(((2*
c*d*cos(f*x + e)^2 + 2*c*d*sin(f*x + e)^2 - (c^2 - 2*c*d)*cos(2*f*x + 2*e)
^2 + c^2*cos(f*x + e) - (c^2 - 2*c*d)*sin(2*f*x + 2*e)^2 + (c^2 - 2*c*d +
4*d^2)*sin(2*f*x + 2*e)*sin(f*x + e) - (c^2 - 2*c*d - (c^2 - 2*c*d + 4*d^2
)*cos(f*x + e))*cos(2*f*x + 2*e))*cos(1/2*arctan2(sin(f*x + e), cos(f*x +
e))) + (c^2*sin(f*x + e) + (c^2 + 2*c*d - 4*d^2)*cos(2*f*x + 2*e)*sin(f*x
+ e) - (c^2 - 2*c*d + (c^2 + 2*c*d - 4*d^2)*cos(f*x + e))*sin(2*f*x + 2...
```

3.243.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+a*sec(f*x+e))^(1/2),x,
algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:index.cc index_m i_lex_is_greater Error: Bad Argument V
alue`

3.243.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + a \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{5/2}}{\sqrt{a + \frac{a}{\cos(e+fx)}} \left(c + \frac{d}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(5/2)/((a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))
)),x)`

output `int((g/cos(e + f*x))^(5/2)/((a + a/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))
)), x)`

3.244 $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$

3.244.1 Optimal result	1780
3.244.2 Mathematica [A] (verified)	1781
3.244.3 Rubi [A] (verified)	1781
3.244.4 Maple [A] (verified)	1785
3.244.5 Fricas [A] (verification not implemented)	1786
3.244.6 Sympy [F]	1786
3.244.7 Maxima [A] (verification not implemented)	1787
3.244.8 Giac [B] (verification not implemented)	1787
3.244.9 Mupad [B] (verification not implemented)	1788

3.244.1 Optimal result

Integrand size = 29, antiderivative size = 250

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{(8ac^4 + 16bc^3d + 24ac^2d^2 + 12bcd^3 + 3ad^4) \operatorname{arctanh}(\sin(e + fx))}{8f}$$

$$+ \frac{(12bc^4 + 95ac^3d + 112bc^2d^2 + 80acd^3 + 16bd^4) \tan(e + fx)}{30f}$$

$$+ \frac{d(24bc^3 + 130ac^2d + 116bcd^2 + 45ad^3) \sec(e + fx) \tan(e + fx)}{120f}$$

$$+ \frac{(12bc^2 + 35acd + 16bd^2) (c + d \sec(e + fx))^2 \tan(e + fx)}{60f}$$

$$+ \frac{(4bc + 5ad)(c + d \sec(e + fx))^3 \tan(e + fx)}{20f} + \frac{b(c + d \sec(e + fx))^4 \tan(e + fx)}{5f}$$

output `1/8*(8*a*c^4+24*a*c^2*d^2+3*a*d^4+16*b*c^3*d+12*b*c*d^3)*arctanh(sin(f*x+e))/f+1/30*(95*a*c^3*d+80*a*c*d^3+12*b*c^4+112*b*c^2*d^2+16*b*d^4)*tan(f*x+e)/f+1/120*d*(130*a*c^2*d+45*a*d^3+24*b*c^3+116*b*c*d^2)*sec(f*x+e)*tan(f*x+e)/f+1/60*(35*a*c*d+12*b*c^2+16*b*d^2)*(c+d*sec(f*x+e))^2*tan(f*x+e)/f+1/20*(5*a*d+4*b*c)*(c+d*sec(f*x+e))^3*tan(f*x+e)/f+1/5*b*(c+d*sec(f*x+e))^4*tan(f*x+e)/f`

3.244.2 Mathematica [A] (verified)

Time = 2.84 (sec) , antiderivative size = 201, normalized size of antiderivative = 0.80

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{15(4bcd(4c^2 + 3d^2) + a(8c^4 + 24c^2d^2 + 3d^4)) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx) (15d(3ad(8c^2 + d^2) + 4$$

input `Integrate[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^4,x]`

output $(15*(4*b*c*d*(4*c^2 + 3*d^2) + a*(8*c^4 + 24*c^2*d^2 + 3*d^4))*\operatorname{ArcTanh}[\operatorname{Sin}[e + f*x]] + \operatorname{Tan}[e + f*x]*(15*d*(3*a*d*(8*c^2 + d^2) + 4*b*(4*c^3 + 3*c*d^2))*\operatorname{Sec}[e + f*x] + 30*d^3*(4*b*c + a*d)*\operatorname{Sec}[e + f*x]^3 + 8*(15*(4*a*c*d*(c^2 + d^2) + b*(c^4 + 6*c^2*d^2 + d^4)) + 10*d^2*(2*a*c*d + b*(3*c^2 + d^2))*\operatorname{Tan}[e + f*x]^2 + 3*b*d^4*\operatorname{Tan}[e + f*x]^4))/(120*f)$

3.244.3 Rubi [A] (verified)

Time = 1.42 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$, Rules used = {3042, 4490, 3042, 4490, 3042, 4490, 3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$\downarrow 3042$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a + b \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^4 dx$$

$$\downarrow 4490$$

$$\frac{1}{5} \int \sec(e + fx)(c + d \sec(e + fx))^3(5ac + 4bd + (4bc + 5ad) \sec(e + fx))dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f}$$

$$\downarrow 3042$$

$$\frac{1}{5} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 \left(5ac + 4bd + (4bc + 5ad) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f}$$

↓ 4490

$$\frac{1}{5} \left(\frac{1}{4} \int \sec(e + fx)(c + d \sec(e + fx))^2 (20ac^2 + 28bdc + 15ad^2 + (12bc^2 + 35adc + 16bd^2) \sec(e + fx)) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 (20ac^2 + 28bdc + 15ad^2 + (12bc^2 + 35adc + 16bd^2) \csc\left(e + fx + \frac{\pi}{2}\right)) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)$$

↓ 4490

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \sec(e + fx)(c + d \sec(e + fx)) (60ac^3 + 108bdc^2 + 115ad^2c + 32bd^3 + (24bc^3 + 130adc^2 + 116bd^2c + 15ad^3) \sec(e + fx)) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)\right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) (60ac^3 + 108bdc^2 + 115ad^2c + 32bd^3 + (24bc^3 + 130adc^2 + 116bd^2c + 15ad^3) \csc\left(e + fx + \frac{\pi}{2}\right)) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)\right)$$

↓ 4485

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \sec(e + fx) (15(8ac^4 + 16bdc^3 + 24ad^2c^2 + 12bd^3c + 3ad^4) + 4(12bc^4 + 95adc^3 + 112bd^2c^2 + 80ad^3c + 15ad^4) \sec(e + fx)) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f}\right)\right)\right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \csc \left(e + fx + \frac{\pi}{2} \right) \left(15(8ac^4 + 16bdc^3 + 24ad^2c^2 + 12bd^3c + 3ad^4) + 4(12bc^4 + 95adc^3 + 112bd^2c^2) \right. \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right. \right. \right. \\ \left. \left. \downarrow 4274 \right. \right. \right.$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(4(95ac^3d + 80acd^3 + 12bc^4 + 112bc^2d^2 + 16bd^4) \int \sec^2(e + fx) dx + 15(8ac^4 + 24ac^2d^2 + 3ad^4 + 112bd^3c + 3ad^4) \right. \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right. \right. \right. \\ \left. \left. \downarrow 3042 \right. \right. \right.$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15(8ac^4 + 24ac^2d^2 + 3ad^4 + 16bc^3d + 12bcd^3) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + 4(95ac^3d + 80acd^3 + 12bc^4 + 112bd^2c^2) \right. \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right. \right. \right. \\ \left. \left. \downarrow 4254 \right. \right. \right.$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15(8ac^4 + 24ac^2d^2 + 3ad^4 + 16bc^3d + 12bcd^3) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx - \frac{4(95ac^3d + 80acd^3 + 12bc^4 + 112bd^2c^2)}{5f} \right. \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right. \right. \right. \\ \left. \left. \downarrow 24 \right. \right. \right.$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15(8ac^4 + 24ac^2d^2 + 3ad^4 + 16bc^3d + 12bcd^3) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{4(95ac^3d + 80acd^3 + 12bc^4 + 112bd^2c^2)}{5f} \right. \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right. \right. \right. \\ \left. \left. \downarrow 4257 \right. \right. \right.$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\frac{15(8ac^4 + 24ac^2d^2 + 3ad^4 + 16bc^3d + 12bcd^3) \operatorname{arctanh}(\sin(e + fx))}{f} + \frac{4(95ac^3d + 80acd^3 + 12bc^4 + 112bd^2c^2)}{5f} \right. \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^4}{5f} \right. \right. \right. \right.$$

input `Int[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^4,x]`

3.244. $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$


```
output (b*(c + d*Sec[e + f*x])^4*Tan[e + f*x])/(5*f) + (((4*b*c + 5*a*d)*(c + d*Sec[e + f*x])^3*Tan[e + f*x])/(4*f) + (((12*b*c^2 + 35*a*c*d + 16*b*d^2)*(c + d*Sec[e + f*x])^2*Tan[e + f*x])/(3*f) + ((d*(24*b*c^3 + 130*a*c^2*d + 16*b*c*d^2 + 45*a*d^3)*Sec[e + f*x]*Tan[e + f*x])/(2*f) + ((15*(8*a*c^4 + 16*b*c^3*d + 24*a*c^2*d^2 + 12*b*c*d^3 + 3*a*d^4)*ArcTanh[Sin[e + f*x]])/f + (4*(12*b*c^4 + 95*a*c^3*d + 112*b*c^2*d^2 + 80*a*c*d^3 + 16*b*d^4)*Tan[e + f*x])/f)/2)/3)/4)/5
```

3.244.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4274 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]
```

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]
```

```
rule 4490 Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*Simp[b*B*m + a*A*(m + 1) + (a*B*m + A*b*(m + 1))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0]
```

3.244.4 Maple [A] (verified)

Time = 5.14 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.95

method	result
parts	$\frac{(a d^4 + 4bc d^3) \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3\sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e) + \tan(fx+e))}{8} \right)}{f} - \frac{(4ac d^3 + 6b c^2 d^2) \left(-\frac{2}{3} - \frac{\sec(fx+e)}{f} \right)}{f}$
derivativedivides	$\frac{a c^4 \ln(\sec(fx+e) + \tan(fx+e)) + 4a c^3 d \tan(fx+e) + 6a c^2 d^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) - 4ac d^3 \left(-\frac{2}{3} - \frac{\sec(fx+e)}{f} \right)}{f}$
default	$\frac{a c^4 \ln(\sec(fx+e) + \tan(fx+e)) + 4a c^3 d \tan(fx+e) + 6a c^2 d^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) - 4ac d^3 \left(-\frac{2}{3} - \frac{\sec(fx+e)}{f} \right)}{f}$
parallelrisch	$\frac{-120(\cos(5fx+5e) + 5 \cos(3fx+3e) + 10 \cos(fx+e)) (a^4 + 3a^2 c^2 d^2 + \frac{3}{8} a d^4 + 2b c^3 d + \frac{3}{2} bc d^3) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + 120(c^4 d^2 + 4ac d^3 + 4b c^2 d^2 + 4a^2 c d^4)}{f}$
norman	$\frac{-4(180a^3 c^3 d + 100ac d^3 + 45b c^4 + 150b c^2 d^2 + 29b d^4) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5 - (32a^3 c^3 d - 24a^2 c^2 d^2 + 32ac d^3 - 5a d^4 + 8b c^4 - 16b c^3 d + 48b c^2 d^2 - 16b^2 c d^3 + 16b^2 d^4)}{15f}$
risch	$\frac{i(480b c^2 d^2 + 320ac d^3 + 480a c^3 d + 64b d^4 + 120b c^4 - 210a d^4 e^{7i(fx+e)} + 480b c^4 e^{6i(fx+e)} + 320b d^4 e^{2i(fx+e)} + 720b c^4 e^{4i(fx+e)})}{f}$

```
input int(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^4,x,method=_RETURNVERBOSE)
```

```
output (a*d^4+4*b*c*d^3)/f*(-(-1/4*sec(f*x+e)^3-3/8*sec(f*x+e))*tan(f*x+e)+3/8*ln(sec(f*x+e)+tan(f*x+e)))-(4*a*c*d^3+6*b*c^2*d^2)/f*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)+(6*a*c^2*d^2+4*b*c^3*d)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+(4*a*c^3*d+b*c^4)/f*tan(f*x+e)+a*c^4/f*ln(sec(f*x+e)+tan(f*x+e))-b*d^4/f*(-8/15-1/5*sec(f*x+e)^4-4/15*sec(f*x+e)^2)*tan(f*x+e)
```

3.244. $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$

3.244.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.12

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{15(8ac^4 + 16bc^3d + 24ac^2d^2 + 12bcd^3 + 3ad^4) \cos(fx + e)^5 \log(\sin(fx + e) + 1) - 15(8ac^4 + 16bc^3d$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^4,x, algorithm="fricas")`

output `1/240*(15*(8*a*c^4 + 16*b*c^3*d + 24*a*c^2*d^2 + 12*b*c*d^3 + 3*a*d^4)*cos(f*x + e)^5*log(sin(f*x + e) + 1) - 15*(8*a*c^4 + 16*b*c^3*d + 24*a*c^2*d^2 + 12*b*c*d^3 + 3*a*d^4)*cos(f*x + e)^5*log(-sin(f*x + e) + 1) + 2*(24*b*d^4 + 8*(15*b*c^4 + 60*a*c^3*d + 60*b*c^2*d^2 + 40*a*c*d^3 + 8*b*d^4)*cos(f*x + e)^4 + 15*(16*b*c^3*d + 24*a*c^2*d^2 + 12*b*c*d^3 + 3*a*d^4)*cos(f*x + e)^3 + 16*(15*b*c^2*d^2 + 10*a*c*d^3 + 2*b*d^4)*cos(f*x + e)^2 + 30*(4*b*c*d^3 + a*d^4)*cos(f*x + e))*sin(f*x + e))/(f*cos(f*x + e)^5)`

3.244.6 Sympy [F]

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \int (a + b \sec(e + fx))(c + d \sec(e + fx))^4 \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))**4,x)`

output `Integral((a + b*sec(e + f*x))*(c + d*sec(e + f*x))**4*sec(e + f*x), x)`

3.244.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.52

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{480 (\tan (fx + e)^3 + 3 \tan (fx + e))bc^2d^2 + 320 (\tan (fx + e)^3 + 3 \tan (fx + e))acd^3 + 16 (3 \tan (fx + e) + \dots}{\dots}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^4,x, algorithm="maxima")`

output `1/240*(480*(tan(f*x + e)^3 + 3*tan(f*x + e))*b*c^2*d^2 + 320*(tan(f*x + e)^3 + 3*tan(f*x + e))*a*c*d^3 + 16*(3*tan(f*x + e)^5 + 10*tan(f*x + e)^3 + 15*tan(f*x + e))*b*d^4 - 60*b*c*d^3*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e)))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 15*a*d^4*(2*(3*sin(f*x + e)^3 - 5*sin(f*x + e)))/(sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1) - 3*log(sin(f*x + e) + 1) + 3*log(sin(f*x + e) - 1)) - 240*b*c^3*d*(2*sin(f*x + e))/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 360*a*c^2*d^2*(2*sin(f*x + e))/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) + 240*a*c^4*log(sec(f*x + e) + tan(f*x + e)) + 240*b*c^4*tan(f*x + e) + 960*a*c^3*d*tan(f*x + e))/f`

3.244.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 850 vs. 2(238) = 476.

Time = 0.37 (sec) , antiderivative size = 850, normalized size of antiderivative = 3.40

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^4,x, algorithm="giac")`

```

output 1/120*(15*(8*a*c^4 + 16*b*c^3*d + 24*a*c^2*d^2 + 12*b*c*d^3 + 3*a*d^4)*log
(abs(tan(1/2*f*x + 1/2*e) + 1)) - 15*(8*a*c^4 + 16*b*c^3*d + 24*a*c^2*d^2
+ 12*b*c*d^3 + 3*a*d^4)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(120*b*c^4*
tan(1/2*f*x + 1/2*e)^9 + 480*a*c^3*d*tan(1/2*f*x + 1/2*e)^9 - 240*b*c^3*d*
tan(1/2*f*x + 1/2*e)^9 - 360*a*c^2*d^2*tan(1/2*f*x + 1/2*e)^9 + 720*b*c^2*
d^2*tan(1/2*f*x + 1/2*e)^9 + 480*a*c*d^3*tan(1/2*f*x + 1/2*e)^9 - 300*b*c*
d^3*tan(1/2*f*x + 1/2*e)^9 - 75*a*d^4*tan(1/2*f*x + 1/2*e)^9 + 120*b*d^4*t
an(1/2*f*x + 1/2*e)^9 - 480*b*c^4*tan(1/2*f*x + 1/2*e)^7 - 1920*a*c^3*d*ta
n(1/2*f*x + 1/2*e)^7 + 480*b*c^3*d*tan(1/2*f*x + 1/2*e)^7 + 720*a*c^2*d^2*
tan(1/2*f*x + 1/2*e)^7 - 1920*b*c^2*d^2*tan(1/2*f*x + 1/2*e)^7 - 1280*a*c*
d^3*tan(1/2*f*x + 1/2*e)^7 + 120*b*c*d^3*tan(1/2*f*x + 1/2*e)^7 + 30*a*d^4
*tan(1/2*f*x + 1/2*e)^7 - 160*b*d^4*tan(1/2*f*x + 1/2*e)^7 + 720*b*c^4*tan
(1/2*f*x + 1/2*e)^5 + 2880*a*c^3*d*tan(1/2*f*x + 1/2*e)^5 + 2400*b*c^2*d^2
*tan(1/2*f*x + 1/2*e)^5 + 1600*a*c*d^3*tan(1/2*f*x + 1/2*e)^5 + 464*b*d^4*
tan(1/2*f*x + 1/2*e)^5 - 480*b*c^4*tan(1/2*f*x + 1/2*e)^3 - 1920*a*c^3*d*t
an(1/2*f*x + 1/2*e)^3 - 480*b*c^3*d*tan(1/2*f*x + 1/2*e)^3 - 720*a*c^2*d^2
*tan(1/2*f*x + 1/2*e)^3 - 1920*b*c^2*d^2*tan(1/2*f*x + 1/2*e)^3 - 1280*a*c
*d^3*tan(1/2*f*x + 1/2*e)^3 - 120*b*c*d^3*tan(1/2*f*x + 1/2*e)^3 - 30*a*d^
4*tan(1/2*f*x + 1/2*e)^3 - 160*b*d^4*tan(1/2*f*x + 1/2*e)^3 + 120*b*c^4*ta
n(1/2*f*x + 1/2*e) + 480*a*c^3*d*tan(1/2*f*x + 1/2*e) + 240*b*c^3*d*tan...

```

3.244.9 Mupad [B] (verification not implemented)

Time = 17.35 (sec) , antiderivative size = 555, normalized size of antiderivative = 2.22

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^4 dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(ac^4 + 2bc^3d + 3ac^2d^2 + \frac{3bcd^3}{2} + \frac{3ad^4}{8}\right)}{4ac^4 + 8bc^3d + 12ac^2d^2 + 6bcd^3 + \frac{3ad^4}{2}}\right) \left(2ac^4 + 4bc^3d + 6ac^2d^2 + 3bcd^3 + \frac{3ad^4}{4}\right)}{f}$$

$$- \frac{\left(2bc^4 - \frac{5ad^4}{4} + 2bd^4 - 6ac^2d^2 + 12bc^2d^2 + 8acd^3 + 8ac^3d - 5bcd^3 - 4bc^3d\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}{f} + \dots$$

```

input int(((a + b/cos(e + f*x))*(c + d/cos(e + f*x))^4)/cos(e + f*x),x)

```

output

$$\begin{aligned} & (\operatorname{atanh}((4*\tan(e/2 + (f*x)/2)*(a*c^4 + (3*a*d^4)/8 + 3*a*c^2*d^2 + (3*b*c*d^3)/2 + 2*b*c^3*d))/(4*a*c^4 + (3*a*d^4)/2 + 12*a*c^2*d^2 + 6*b*c*d^3 + 8*b*c^3*d))*(2*a*c^4 + (3*a*d^4)/4 + 6*a*c^2*d^2 + 3*b*c*d^3 + 4*b*c^3*d))/f \\ & - (\tan(e/2 + (f*x)/2)^5*(12*b*c^4 + (116*b*d^4)/15 + 40*b*c^2*d^2 + (80*a*c*d^3)/3 + 48*a*c^3*d) + \tan(e/2 + (f*x)/2)*((5*a*d^4)/4 + 2*b*c^4 + 2*b*d^4 + 6*a*c^2*d^2 + 12*b*c^2*d^2 + 8*a*c*d^3 + 8*a*c^3*d + 5*b*c*d^3 + 4*b*c^3*d) + \tan(e/2 + (f*x)/2)^9*(2*b*c^4 - (5*a*d^4)/4 + 2*b*d^4 - 6*a*c^2*d^2 + 12*b*c^2*d^2 + 8*a*c*d^3 + 8*a*c^3*d - 5*b*c*d^3 - 4*b*c^3*d) - \tan(e/2 + (f*x)/2)^3*((a*d^4)/2 + 8*b*c^4 + (8*b*d^4)/3 + 12*a*c^2*d^2 + 32*b*c^2*d^2 + (64*a*c*d^3)/3 + 32*a*c^3*d + 2*b*c*d^3 + 8*b*c^3*d) - \tan(e/2 + (f*x)/2)^7*(8*b*c^4 - (a*d^4)/2 + (8*b*d^4)/3 - 12*a*c^2*d^2 + 32*b*c^2*d^2 + (64*a*c*d^3)/3 + 32*a*c^3*d - 2*b*c*d^3 - 8*b*c^3*d))/(f*(5*\tan(e/2 + (f*x)/2)^2 - 10*\tan(e/2 + (f*x)/2)^4 + 10*\tan(e/2 + (f*x)/2)^6 - 5*\tan(e/2 + (f*x)/2)^8 + \tan(e/2 + (f*x)/2)^10 - 1)) \end{aligned}$$

3.245 $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^3 dx$

3.245.1 Optimal result	1790
3.245.2 Mathematica [A] (verified)	1791
3.245.3 Rubi [A] (verified)	1791
3.245.4 Maple [A] (verified)	1795
3.245.5 Fricas [A] (verification not implemented)	1795
3.245.6 Sympy [F]	1796
3.245.7 Maxima [A] (verification not implemented)	1796
3.245.8 Giac [B] (verification not implemented)	1797
3.245.9 Mupad [B] (verification not implemented)	1798

3.245.1 Optimal result

Integrand size = 29, antiderivative size = 180

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{(8ac^3 + 12bc^2d + 12acd^2 + 3bd^3) \operatorname{arctanh}(\sin(e + fx))}{8f}$$

$$+ \frac{(4ad(4c^2 + d^2) + 3b(c^3 + 4cd^2)) \tan(e + fx)}{6f}$$

$$+ \frac{d(6bc^2 + 20acd + 9bd^2) \sec(e + fx) \tan(e + fx)}{24f}$$

$$+ \frac{(3bc + 4ad)(c + d \sec(e + fx))^2 \tan(e + fx)}{12f} + \frac{b(c + d \sec(e + fx))^3 \tan(e + fx)}{4f}$$

```
output 1/8*(8*a*c^3+12*a*c*d^2+12*b*c^2*d+3*b*d^3)*arctanh(sin(f*x+e))/f+1/6*(4*a
*d*(4*c^2+d^2)+3*b*(c^3+4*c*d^2))*tan(f*x+e)/f+1/24*d*(20*a*c*d+6*b*c^2+9*
b*d^2)*sec(f*x+e)*tan(f*x+e)/f+1/12*(4*a*d+3*b*c)*(c+d*sec(f*x+e))^2*tan(f
*x+e)/f+1/4*b*(c+d*sec(f*x+e))^3*tan(f*x+e)/f
```

3.245.2 Mathematica [A] (verified)

Time = 0.78 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.79

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{3(3bd(4c^2 + d^2) + 4a(2c^3 + 3cd^2)) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx) (9d(4acd + b(4c^2 + d^2)) \sec(e + fx) + 6bd^2 \tan(e + fx))}{24f}$$

input `Integrate[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^3,x]`output `(3*(3*b*d*(4*c^2 + d^2) + 4*a*(2*c^3 + 3*c*d^2))*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(9*d*(4*a*c*d + b*(4*c^2 + d^2))*Sec[e + f*x] + 6*b*d^3*Sec[e + f*x]^3 + 8*(3*a*d*(3*c^2 + d^2) + 3*b*(c^3 + 3*c*d^2) + d^2*(3*b*c + a*d))*Tan[e + f*x]^2))/(24*f)`**3.245.3 Rubi [A] (verified)**Time = 1.05 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 4490, 3042, 4490, 3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a + b \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^3 dx$$

$$\downarrow \text{4490}$$

$$\frac{1}{4} \int \sec(e + fx)(c + d \sec(e + fx))^2(4ac + 3bd + (3bc + 4ad) \sec(e + fx)) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

$$\downarrow \text{3042}$$

3.245. $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^3 dx$

$$\frac{1}{4} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 \left(4ac + 3bd + (3bc + 4ad) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

↓ 4490

$$\frac{1}{4} \left(\frac{1}{3} \int \sec(e + fx)(c + d \sec(e + fx)) (12ac^2 + 15bdc + 8ad^2 + (6bc^2 + 20adc + 9bd^2) \sec(e + fx)) dx + \frac{(4ad + 3bd^2) \sec(e + fx)}{3}\right) \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(12ac^2 + 15bdc + 8ad^2 + (6bc^2 + 20adc + 9bd^2) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \frac{(4ad + 3bd^2) \csc\left(e + fx + \frac{\pi}{2}\right)}{3}\right) \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

↓ 4485

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \sec(e + fx) (3(3bd(4c^2 + d^2) + 4a(2c^3 + 3d^2c)) + 4(4ad(4c^2 + d^2) + 3b(c^3 + 4d^2c))) \sec(e + fx) dx + \frac{(4ad + 3bd^2) \sec(e + fx)}{3}\right)\right) \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(3(3bd(4c^2 + d^2) + 4a(2c^3 + 3d^2c)) + 4(4ad(4c^2 + d^2) + 3b(c^3 + 4d^2c))\right) \csc\left(e + fx + \frac{\pi}{2}\right) dx + \frac{(4ad + 3bd^2) \csc\left(e + fx + \frac{\pi}{2}\right)}{3}\right)\right) \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

↓ 4274

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(4(4ad(4c^2 + d^2) + 3b(c^3 + 4cd^2)) \int \sec^2(e + fx) dx + 3(8ac^3 + 12acd^2 + 12bc^2d + 3bd^3) \int \sec(e + fx) dx + \frac{(4ad + 3bd^2) \sec(e + fx)}{3}\right)\right)\right) \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(4(4ad(4c^2 + d^2) + 3b(c^3 + 4cd^2)) \int \csc \left(e + fx + \frac{\pi}{2} \right)^2 dx + 3(8ac^3 + 12acd^2 + 12bc^2d + 3bd^3) \int \csc \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f} \right) \right) \right. \\ \left. \downarrow 4254 \right.$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3(8ac^3 + 12acd^2 + 12bc^2d + 3bd^3) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx - \frac{4(4ad(4c^2 + d^2) + 3b(c^3 + 4cd^2)) \int 1d(-}{f} \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f} \right) \right) \right. \\ \left. \downarrow 24 \right.$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3(8ac^3 + 12acd^2 + 12bc^2d + 3bd^3) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{4(4ad(4c^2 + d^2) + 3b(c^3 + 4cd^2)) \tan(e + }{f} \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f} \right) \right) \right. \\ \left. \downarrow 4257 \right.$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\frac{3(8ac^3 + 12acd^2 + 12bc^2d + 3bd^3) \operatorname{arctanh}(\sin(e + fx))}{f} + \frac{4(4ad(4c^2 + d^2) + 3b(c^3 + 4cd^2)) \tan(e + }{f} \right. \right. \right. \\ \left. \left. \left. \frac{b \tan(e + fx)(c + d \sec(e + fx))^3}{4f} \right) \right) \right.$$

input `Int[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^3,x]`

output `(b*(c + d*Sec[e + f*x])^3*Tan[e + f*x])/(4*f) + (((3*b*c + 4*a*d)*(c + d*Sec[e + f*x])^2*Tan[e + f*x])/(3*f) + ((d*(6*b*c^2 + 20*a*c*d + 9*b*d^2)*Sec[e + f*x]*Tan[e + f*x])/(2*f) + ((3*(8*a*c^3 + 12*b*c^2*d + 12*a*c*d^2 + 3*b*d^3)*ArcTanh[Sin[e + f*x]])/f + (4*(4*a*d*(4*c^2 + d^2) + 3*b*(c^3 + 4*c*d^2))*Tan[e + f*x])/f)/2)/3)/4`

3.245.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4485 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`
- rule 4490 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*Simp[b*B*m + a*A*(m + 1) + (a*B*m + A*b*(m + 1))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0]`

3.245.4 Maple [A] (verified)

Time = 3.81 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.03

method	result
parts	$-\frac{(a d^3+3 b c d^2)\left(-\frac{2}{3}-\frac{\sec(f x+e)^2}{3}\right) \tan(f x+e)}{f}+\frac{(3 a c d^2+3 b c^2 d)\left(\frac{\sec(f x+e) \tan(f x+e)}{2}+\frac{\ln(\sec(f x+e)+\tan(f x+e))}{2}\right)}{f}$
derivativedivides	$\frac{a c^3 \ln(\sec(f x+e)+\tan(f x+e))+3 a c^2 d \tan(f x+e)+3 a c d^2\left(\frac{\sec(f x+e) \tan(f x+e)}{2}+\frac{\ln(\sec(f x+e)+\tan(f x+e))}{2}\right)-a d^3\left(-\frac{2}{3}-\frac{\sec(f x+e)^2}{3}\right) \tan(f x+e)}{f}$
default	$\frac{a c^3 \ln(\sec(f x+e)+\tan(f x+e))+3 a c^2 d \tan(f x+e)+3 a c d^2\left(\frac{\sec(f x+e) \tan(f x+e)}{2}+\frac{\ln(\sec(f x+e)+\tan(f x+e))}{2}\right)-a d^3\left(-\frac{2}{3}-\frac{\sec(f x+e)^2}{3}\right) \tan(f x+e)}{f}$
parallelrisch	$-96\left(\frac{3}{4}+\frac{\cos(4 f x+4 e)}{4}+\cos(2 f x+2 e)\right)\left(a c^3+\frac{3}{2} a c d^2+\frac{3}{2} b c^2 d+\frac{3}{8} b d^3\right) \ln\left(\tan\left(\frac{f x}{2}+\frac{e}{2}\right)-1\right)+96\left(\frac{3}{4}+\frac{\cos(4 f x+4 e)}{4}+\cos(2 f x+2 e)\right)$
norman	$-\frac{(24 a c^2 d-12 a c d^2+8 a d^3+8 b c^3-12 b c^2 d+24 b c d^2-5 b d^3) \tan\left(\frac{f x}{2}+\frac{e}{2}\right)^7}{4 f}+\frac{(24 a c^2 d+12 a c d^2+8 a d^3+8 b c^3+12 b c^2 d+24 b c d^2+5 b d^3) \tan\left(\frac{f x}{2}+\frac{e}{2}\right)}{4 f}$
risch	$-\frac{i(-48 b c d^2-72 a c^2 d-16 a d^3-24 b c^3-33 b d^3 e^{3 i(f x+e)}-9 d^3 b e^{i(f x+e)}-72 b c^3 e^{4 i(f x+e)}+33 b d^3 e^{5 i(f x+e)}-48 a d^3 e^{4 i(f x+e)})}{4 f}$

input `int(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$-(a d^3+3 b c d^2) / f * (-2 / 3-1 / 3 * \sec(f x+e)^2) * \tan(f x+e)+(3 a c d^2+3 b c^2 d) / f * (1 / 2 * \sec(f x+e) * \tan(f x+e)+1 / 2 * \ln(\sec(f x+e)+\tan(f x+e)))+(3 a c^2 d+b c^3) / f * \tan(f x+e)+a c^3 / f * \ln(\sec(f x+e)+\tan(f x+e))+b d^3 / f * (-(-1 / 4 * \sec(f x+e)^3-3 / 8 * \sec(f x+e)) * \tan(f x+e)+3 / 8 * \ln(\sec(f x+e)+\tan(f x+e)))$$

3.245.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.17

$$\int \sec(e+fx)(a+b \sec(e+fx))(c+d \sec(e+fx))^3 dx$$

$$= \frac{3(8 a c^3+12 b c^2 d+12 a c d^2+3 b d^3) \cos(f x+e)^4 \log(\sin(f x+e)+1)-3(8 a c^3+12 b c^2 d+12 a c d^2+3 b d^3) \sec(f x+e)^3}{4}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^3,x, algorithm="fricas")`

3.245.
$$\int \sec(e+fx)(a+b \sec(e+fx))(c+d \sec(e+fx))^3 dx$$

output $\frac{1}{48}(3(8a^3c^3 + 12b^2cd + 12acd^2 + 3b^3d^3)\cos(fx + e)^4 \log(\sin(fx + e) + 1) - 3(8a^3c^3 + 12b^2cd + 12acd^2 + 3b^3d^3)\cos(fx + e)^4 \log(-\sin(fx + e) + 1) + 2(6b^3d^3 + 8(3b^2c^3 + 9a^2c^2d + 6b^2cd^2 + 2ad^3)\cos(fx + e)^3 + 9(4b^2c^2d + 4acd^2 + bd^3)\cos(fx + e)^2 + 8(3b^2cd^2 + ad^3)\cos(fx + e))\sin(fx + e))/(f\cos(fx + e)^4)$

3.245.6 Sympy [F]

$$\begin{aligned} & \int \sec(e + fx)(a + b\sec(e + fx))(c + d\sec(e + fx))^3 dx \\ &= \int (a + b\sec(e + fx))(c + d\sec(e + fx))^3 \sec(e + fx) dx \end{aligned}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))**3,x)`

output `Integral((a + b*sec(e + f*x))*(c + d*sec(e + f*x))**3*sec(e + f*x), x)`

3.245.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.48

$$\begin{aligned} & \int \sec(e + fx)(a + b\sec(e + fx))(c + d\sec(e + fx))^3 dx \\ &= \frac{48(\tan(fx + e)^3 + 3\tan(fx + e))bcd^2 + 16(\tan(fx + e)^3 + 3\tan(fx + e))ad^3 - 3bd^3 \left(\frac{2(3\sin(fx+e)^3 - \sin(fx+e)^4 - 2\sin(fx+e))}{\sin(fx+e)^4 - 2\sin(fx+e)} \right)}{\sin(fx+e)^4 - 2\sin(fx+e)} \end{aligned}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output $\frac{1}{48}(48(\tan(fx + e))^3 + 3\tan(fx + e))*b*c*d^2 + 16(\tan(fx + e))^3 + 3\tan(fx + e)*a*d^3 - 3*b*d^3*(2*(3*\sin(fx + e))^3 - 5*\sin(fx + e))/(\sin(fx + e)^4 - 2*\sin(fx + e)^2 + 1) - 3*\log(\sin(fx + e) + 1) + 3*\log(\sin(fx + e) - 1) - 36*b*c^2*d*(2*\sin(fx + e))/(\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1) - 36*a*c*d^2*(2*\sin(fx + e))/(\sin(fx + e)^2 - 1) - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1) + 48*a*c^3*\log(\sec(fx + e) + \tan(fx + e)) + 48*b*c^3*\tan(fx + e) + 144*a*c^2*d*\tan(fx + e))/f$

3.245.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 586 vs. $2(170) = 340$.

Time = 0.36 (sec) , antiderivative size = 586, normalized size of antiderivative = 3.26

$$\int \sec(e + fx)(a + b\sec(e + fx))(c + d\sec(e + fx))^3 dx$$

$$= \frac{3(8ac^3 + 12bc^2d + 12acd^2 + 3bd^3) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3(8ac^3 + 12bc^2d + 12acd^2 + 3bd^3) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{f}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output $\frac{1}{24}(3*(8*a*c^3 + 12*b*c^2*d + 12*a*c*d^2 + 3*b*d^3)*\log(\tan(1/2*f*x + 1/2*e) + 1) - 3*(8*a*c^3 + 12*b*c^2*d + 12*a*c*d^2 + 3*b*d^3)*\log(\tan(1/2*f*x + 1/2*e) - 1) - 2*(24*b*c^3*\tan(1/2*f*x + 1/2*e)^7 + 72*a*c^2*d*\tan(1/2*f*x + 1/2*e)^7 - 36*b*c^2*d*\tan(1/2*f*x + 1/2*e)^7 - 36*a*c*d^2*\tan(1/2*f*x + 1/2*e)^7 + 72*b*c*d^2*\tan(1/2*f*x + 1/2*e)^7 + 24*a*d^3*\tan(1/2*f*x + 1/2*e)^7 - 15*b*d^3*\tan(1/2*f*x + 1/2*e)^7 - 72*b*c^3*\tan(1/2*f*x + 1/2*e)^5 - 216*a*c^2*d*\tan(1/2*f*x + 1/2*e)^5 + 36*b*c^2*d*\tan(1/2*f*x + 1/2*e)^5 + 36*a*c*d^2*\tan(1/2*f*x + 1/2*e)^5 - 120*b*c*d^2*\tan(1/2*f*x + 1/2*e)^5 - 40*a*d^3*\tan(1/2*f*x + 1/2*e)^5 - 9*b*d^3*\tan(1/2*f*x + 1/2*e)^5 + 72*b*c^3*\tan(1/2*f*x + 1/2*e)^3 + 216*a*c^2*d*\tan(1/2*f*x + 1/2*e)^3 + 36*b*c^2*d*\tan(1/2*f*x + 1/2*e)^3 + 36*a*c*d^2*\tan(1/2*f*x + 1/2*e)^3 + 120*b*c*d^2*\tan(1/2*f*x + 1/2*e)^3 + 40*a*d^3*\tan(1/2*f*x + 1/2*e)^3 - 9*b*d^3*\tan(1/2*f*x + 1/2*e)^3 - 24*b*c^3*\tan(1/2*f*x + 1/2*e) - 72*a*c^2*d*\tan(1/2*f*x + 1/2*e) - 36*b*c^2*d*\tan(1/2*f*x + 1/2*e) - 36*a*c*d^2*\tan(1/2*f*x + 1/2*e) - 72*b*c*d^2*\tan(1/2*f*x + 1/2*e) - 24*a*d^3*\tan(1/2*f*x + 1/2*e) - 15*b*d^3*\tan(1/2*f*x + 1/2*e))/(\tan(1/2*f*x + 1/2*e)^2 - 1)^4)/f$

3.245. $\int \sec(e + fx)(a + b\sec(e + fx))(c + d\sec(e + fx))^3 dx$

3.245.9 Mupad [B] (verification not implemented)

Time = 17.20 (sec) , antiderivative size = 395, normalized size of antiderivative = 2.19

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^3 dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(ac^3 + \frac{3bc^2d}{2} + \frac{3acd^2}{2} + \frac{3bd^3}{8}\right)}{4ac^3 + 6bc^2d + 6acd^2 + \frac{3bd^3}{2}}\right) \left(2ac^3 + 3bc^2d + 3acd^2 + \frac{3bd^3}{4}\right)}{f} \\ - \frac{\left(2ad^3 + 2bc^3 - \frac{5bd^3}{4} - 3acd^2 + 6ac^2d + 6bcd^2 - 3b^2c^2d\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^7 + \left(3acd^2 - 6bc^3 - \frac{3bd^3}{4}\right)}{f}$$

input `int(((a + b/cos(e + f*x))*(c + d/cos(e + f*x))^3)/cos(e + f*x),x)`

output

```
(atanh((4*tan(e/2 + (f*x)/2)*(a*c^3 + (3*b*d^3)/8 + (3*a*c*d^2)/2 + (3*b*c^2*d)/2)))/(4*a*c^3 + (3*b*d^3)/2 + 6*a*c*d^2 + 6*b*c^2*d))*(2*a*c^3 + (3*b*d^3)/4 + 3*a*c*d^2 + 3*b*c^2*d))/f - (tan(e/2 + (f*x)/2)^7*(2*a*d^3 + 2*b*c^3 - (5*b*d^3)/4 - 3*a*c*d^2 + 6*a*c^2*d + 6*b*c*d^2 - 3*b*c^2*d) + tan(e/2 + (f*x)/2)^3*((10*a*d^3)/3 + 6*b*c^3 - (3*b*d^3)/4 + 3*a*c*d^2 + 18*a*c^2*d + 10*b*c*d^2 + 3*b*c^2*d) - tan(e/2 + (f*x)/2)^5*((10*a*d^3)/3 + 6*b*c^3 + (3*b*d^3)/4 - 3*a*c*d^2 + 18*a*c^2*d + 10*b*c*d^2 - 3*b*c^2*d) - tan(e/2 + (f*x)/2)*(2*a*d^3 + 2*b*c^3 + (5*b*d^3)/4 + 3*a*c*d^2 + 6*a*c^2*d + 6*b*c*d^2 + 3*b*c^2*d))/(f*(6*tan(e/2 + (f*x)/2)^4 - 4*tan(e/2 + (f*x)/2)^2 - 4*tan(e/2 + (f*x)/2)^6 + tan(e/2 + (f*x)/2)^8 + 1))
```

3.246 $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$

3.246.1 Optimal result	1799
3.246.2 Mathematica [A] (verified)	1799
3.246.3 Rubi [A] (verified)	1800
3.246.4 Maple [A] (verified)	1803
3.246.5 Fricas [A] (verification not implemented)	1803
3.246.6 Sympy [F]	1804
3.246.7 Maxima [A] (verification not implemented)	1804
3.246.8 Giac [B] (verification not implemented)	1805
3.246.9 Mupad [B] (verification not implemented)	1805

3.246.1 Optimal result

Integrand size = 29, antiderivative size = 115

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{(2bcd + a(2c^2 + d^2)) \operatorname{arctanh}(\sin(e + fx))}{2f} + \frac{2(3acd + b(c^2 + d^2)) \tan(e + fx)}{3f}$$

$$+ \frac{d(2bc + 3ad) \sec(e + fx) \tan(e + fx)}{6f} + \frac{b(c + d \sec(e + fx))^2 \tan(e + fx)}{3f}$$

```
output 1/2*(2*b*c*d+a*(2*c^2+d^2))*arctanh(sin(f*x+e))/f+2/3*(3*a*c*d+b*(c^2+d^2)
)*tan(f*x+e)/f+1/6*d*(3*a*d+2*b*c)*sec(f*x+e)*tan(f*x+e)/f+1/3*b*(c+d*sec(
f*x+e))^2*tan(f*x+e)/f
```

3.246.2 Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.77

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{3(2bcd + a(2c^2 + d^2)) \operatorname{arctanh}(\sin(e + fx)) + \tan(e + fx) (12acd + 6b(c^2 + d^2) + 3d(2bc + ad) \sec(e + fx))}{6f}$$

input `Integrate[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^2,x]`

output `(3*(2*b*c*d + a*(2*c^2 + d^2))*ArcTanh[Sin[e + f*x]] + Tan[e + f*x]*(12*a*c*d + 6*b*(c^2 + d^2) + 3*d*(2*b*c + a*d)*Sec[e + f*x] + 2*b*d^2*Tan[e + f*x]^2))/(6*f)`

3.246.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4490, 3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a + b \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2 dx \\
 & \quad \downarrow \text{4490} \\
 & \frac{1}{3} \int \sec(e + fx)(c + d \sec(e + fx))(3ac + 2bd + (2bc + 3ad) \sec(e + fx)) dx + \\
 & \quad \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(3ac + 2bd + (2bc + 3ad) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \\
 & \quad \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f} \\
 & \quad \downarrow \text{4485} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \sec(e + fx) (3(2bcd + a(2c^2 + d^2)) + 4(3acd + b(c^2 + d^2)) \sec(e + fx)) dx + \frac{d(3ad + 2bc) \tan(e + fx) \sec(e + fx)}{2f} \right) \\
 & \quad \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.246. $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$

$$\frac{1}{3} \left(\frac{1}{2} \int \csc \left(e + fx + \frac{\pi}{2} \right) \left(3(2bcd + a(2c^2 + d^2)) + 4(3acd + b(c^2 + d^2)) \csc \left(e + fx + \frac{\pi}{2} \right) \right) dx + \frac{d(3ad + 2bc) \tan(e + fx)}{2f} \right) + \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 4274

$$\frac{1}{3} \left(\frac{1}{2} \left(4(3acd + b(c^2 + d^2)) \int \sec^2(e + fx) dx + 3(a(2c^2 + d^2) + 2bcd) \int \sec(e + fx) dx \right) + \frac{d(3ad + 2bc) \tan(e + fx)}{2f} \right) + \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3(a(2c^2 + d^2) + 2bcd) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + 4(3acd + b(c^2 + d^2)) \int \csc \left(e + fx + \frac{\pi}{2} \right)^2 dx \right) + \frac{d(3ad + 2bc) \tan(e + fx)}{2f} \right) + \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 4254

$$\frac{1}{3} \left(\frac{1}{2} \left(3(a(2c^2 + d^2) + 2bcd) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx - \frac{4(3acd + b(c^2 + d^2)) \int 1d(-\tan(e + fx))}{f} \right) + \frac{d(3ad + 2bc) \tan(e + fx)}{2f} \right) + \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 24

$$\frac{1}{3} \left(\frac{1}{2} \left(3(a(2c^2 + d^2) + 2bcd) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{4(3acd + b(c^2 + d^2)) \tan(e + fx)}{f} \right) + \frac{d(3ad + 2bc) \tan(e + fx)}{2f} \right) + \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{3(a(2c^2 + d^2) + 2bcd) \operatorname{arctanh}(\sin(e + fx))}{f} + \frac{4(3acd + b(c^2 + d^2)) \tan(e + fx)}{f} \right) + \frac{d(3ad + 2bc) \tan(e + fx)}{2f} \right) + \frac{b \tan(e + fx)(c + d \sec(e + fx))^2}{3f}$$

input `Int[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^2,x]`

output $(b*(c + d*\text{Sec}[e + f*x])^2*\text{Tan}[e + f*x])/(3*f) + ((d*(2*b*c + 3*a*d)*\text{Sec}[e + f*x]*\text{Tan}[e + f*x])/(2*f) + ((3*(2*b*c*d + a*(2*c^2 + d^2))*\text{ArcTanh}[\text{Sin}[e + f*x]])/f + (4*(3*a*c*d + b*(c^2 + d^2))*\text{Tan}[e + f*x])/f)/2)/3$

3.246.3.1 Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ ; FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 4254 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[-d^{(-1)} \text{ Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] \text{ ; FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

rule 4257 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{ Int}[(d*\text{Csc}[e + f*x])^{(n + 1)}, x], x] \text{ ; FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4485 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))*(\text{csc}[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*((d*\text{Csc}[e + f*x])^n/(f*(n + 1))), x] + \text{Simp}[1/(n + 1) \text{ Int}[(d*\text{Csc}[e + f*x])^n*\text{Simp}[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*\text{Csc}[e + f*x], x], x], x] \text{ ; FreeQ}[\{a, b, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[A*b - a*B, 0] \ \&\& \ !\text{LeQ}[n, -1]$

rule 4490 $\text{Int}[\text{csc}[(e_.) + (f_.)*(x_.)]*(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^{(m_.)}*(\text{csc}[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-B)*\text{Cot}[e + f*x]*((a + b*\text{Csc}[e + f*x])^m/(f*(m + 1))), x] + \text{Simp}[1/(m + 1) \text{ Int}[\text{Csc}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m - 1)}*\text{Simp}[b*B*m + a*A*(m + 1) + (a*B*m + A*b*(m + 1))*\text{Csc}[e + f*x], x], x], x] \text{ ; FreeQ}[\{a, b, A, B, e, f\}, x] \ \&\& \ \text{NeQ}[A*b - a*B, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$

3.246.4 Maple [A] (verified)

Time = 3.40 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.03

method	result
parts	$\frac{(a d^2 + 2bcd) \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right)}{f} + \frac{(2acd + b c^2) \tan(fx+e)}{f} + \frac{a c^2 \ln(\sec(fx+e) + \tan(fx+e))}{f}$
derivativedivides	$\frac{a c^2 \ln(\sec(fx+e) + \tan(fx+e)) + 2acd \tan(fx+e) + a d^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) + b c^2 \tan(fx+e)}{f}$
default	$\frac{a c^2 \ln(\sec(fx+e) + \tan(fx+e)) + 2acd \tan(fx+e) + a d^2 \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e) + \tan(fx+e))}{2} \right) + b c^2 \tan(fx+e)}{f}$
parallelrisch	$\frac{-9 \left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3} \right) \left(a c^2 + \frac{1}{2} a d^2 + bcd \right) \ln \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right) - 1 \right) + 9 \left(\cos(fx+e) + \frac{\cos(3fx+3e)}{3} \right) \left(a c^2 + \frac{1}{2} a d^2 + bcd \right)}{3f \cos(3fx+3e)}$
norman	$\frac{4 \left(6acd + 3b c^2 + b d^2 \right) \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^3 - \left(4acd - a d^2 + 2b c^2 - 2bcd + 2b d^2 \right) \tan \left(\frac{fx}{2} + \frac{e}{2} \right)^5 - \left(4acd + a d^2 + 2b c^2 + 2bcd + 2b d^2 \right) \tan \left(\frac{fx}{2} + \frac{e}{2} \right)}{3f \left(\tan \left(\frac{fx}{2} + \frac{e}{2} \right)^2 - 1 \right)^3}$
risch	$-\frac{i \left(3a d^2 e^{5i(fx+e)} + 6bcd e^{5i(fx+e)} - 12acd e^{4i(fx+e)} - 6b c^2 e^{4i(fx+e)} - 24acd e^{2i(fx+e)} - 12b c^2 e^{2i(fx+e)} - 12b d^2 e^{2i(fx+e)} \right)}{3f \left(1 + e^{2i(fx+e)} \right)^3}$

input `int(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `(a*d^2+2*b*c*d)/f*(1/2*sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e)))+(2*a*c*d+b*c^2)/f*tan(f*x+e)+a*c^2/f*ln(sec(f*x+e)+tan(f*x+e))-b*d^2/f*(-2/3-1/3*sec(f*x+e)^2)*tan(f*x+e)`

3.246.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.30

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{3(2ac^2 + 2bcd + ad^2) \cos(fx + e)^3 \log(\sin(fx + e) + 1) - 3(2ac^2 + 2bcd + ad^2) \cos(fx + e)^3 \log(-\sin(fx + e) + 1)}{12f \cos(fx + e)^3}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^2,x, algorithm="fricas")`

3.246. $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$

output $1/12*(3*(2*a*c^2 + 2*b*c*d + a*d^2)*\cos(f*x + e)^3*\log(\sin(f*x + e) + 1) - 3*(2*a*c^2 + 2*b*c*d + a*d^2)*\cos(f*x + e)^3*\log(-\sin(f*x + e) + 1) + 2*(2*b*d^2 + 2*(3*b*c^2 + 6*a*c*d + 2*b*d^2)*\cos(f*x + e)^2 + 3*(2*b*c*d + a*d^2)*\cos(f*x + e))*\sin(f*x + e))/(f*\cos(f*x + e)^3)$

3.246.6 Sympy [F]

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \int (a + b \sec(e + fx))(c + d \sec(e + fx))^2 \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))**2,x)`

output `Integral((a + b*sec(e + f*x))*(c + d*sec(e + f*x))**2*sec(e + f*x), x)`

3.246.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.43

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{4(\tan(fx + e))^3 + 3 \tan(fx + e)bd^2 - 6bcd\left(\frac{2 \sin(fx+e)}{\sin(fx+e)^2-1} - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1)\right) - 3ad^2\left(\frac{2 \sin(fx+e)}{\sin(fx+e)^2-1} - \log(\sin(fx + e) + 1) + \log(\sin(fx + e) - 1)\right) + 12a*c^2*\log(\sec(f*x + e) + \tan(f*x + e)) + 12*b*c^2*tan(f*x + e) + 24*a*c*d*tan(f*x + e))/f}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output $1/12*(4*(\tan(f*x + e)^3 + 3*\tan(f*x + e))*b*d^2 - 6*b*c*d*(2*\sin(f*x + e)/(\sin(f*x + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) - 3*a*d^2*(2*\sin(f*x + e)/(\sin(f*x + e)^2 - 1) - \log(\sin(f*x + e) + 1) + \log(\sin(f*x + e) - 1)) + 12*a*c^2*\log(\sec(f*x + e) + \tan(f*x + e)) + 12*b*c^2*tan(f*x + e) + 24*a*c*d*tan(f*x + e))/f$

3.246.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. $2(107) = 214$.

Time = 0.34 (sec) , antiderivative size = 294, normalized size of antiderivative = 2.56

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{3(2ac^2 + 2bcd + ad^2) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right) - 3(2ac^2 + 2bcd + ad^2) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{f}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output `1/6*(3*(2*a*c^2 + 2*b*c*d + a*d^2)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - 3*(2*a*c^2 + 2*b*c*d + a*d^2)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(6*b*c^2*tan(1/2*f*x + 1/2*e)^5 + 12*a*c*d*tan(1/2*f*x + 1/2*e)^5 - 6*b*c*d*tan(1/2*f*x + 1/2*e)^5 - 3*a*d^2*tan(1/2*f*x + 1/2*e)^5 + 6*b*d^2*tan(1/2*f*x + 1/2*e)^5 - 12*b*c^2*tan(1/2*f*x + 1/2*e)^3 - 24*a*c*d*tan(1/2*f*x + 1/2*e)^3 - 4*b*d^2*tan(1/2*f*x + 1/2*e)^3 + 6*b*c^2*tan(1/2*f*x + 1/2*e) + 12*a*c*d*tan(1/2*f*x + 1/2*e) + 6*b*c*d*tan(1/2*f*x + 1/2*e) + 3*a*d^2*tan(1/2*f*x + 1/2*e) + 6*b*d^2*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^3)/f`

3.246.9 Mupad [B] (verification not implemented)

Time = 17.02 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.97

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx))^2 dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) \left(ac^2 + bcd + \frac{ad^2}{2}\right)}{4ac^2 + 4bcd + 2ad^2}\right) (2ac^2 + 2bcd + ad^2)}{f}$$

$$- \frac{(2bc^2 - ad^2 + 2bd^2 + 4acd - 2bcd) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^5 + \left(-4bc^2 - 8acd - \frac{4bd^2}{3}\right) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^3 + (ac^2 + bcd + \frac{ad^2}{2}) \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}{f \left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6 - 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 + 3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)}$$

input `int(((a + b/cos(e + f*x))*(c + d/cos(e + f*x))^2)/cos(e + f*x),x)`

output $(\operatorname{atanh}((4*\tan(e/2 + (f*x)/2)*(a*c^2 + (a*d^2)/2 + b*c*d))/(4*a*c^2 + 2*a*d^2 + 4*b*c*d))*(2*a*c^2 + a*d^2 + 2*b*c*d))/f - (\tan(e/2 + (f*x)/2)*(a*d^2 + 2*b*c^2 + 2*b*d^2 + 4*a*c*d + 2*b*c*d) - \tan(e/2 + (f*x)/2)^3*(4*b*c^2 + (4*b*d^2)/3 + 8*a*c*d) + \tan(e/2 + (f*x)/2)^5*(2*b*c^2 - a*d^2 + 2*b*d^2 + 4*a*c*d - 2*b*c*d))/(f*(3*\tan(e/2 + (f*x)/2)^2 - 3*\tan(e/2 + (f*x)/2)^4 + \tan(e/2 + (f*x)/2)^6 - 1))$

3.247 $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$

3.247.1 Optimal result	1807
3.247.2 Mathematica [A] (verified)	1807
3.247.3 Rubi [A] (verified)	1808
3.247.4 Maple [A] (verified)	1810
3.247.5 Fricas [A] (verification not implemented)	1811
3.247.6 Sympy [F]	1811
3.247.7 Maxima [A] (verification not implemented)	1811
3.247.8 Giac [B] (verification not implemented)	1812
3.247.9 Mupad [B] (verification not implemented)	1812

3.247.1 Optimal result

Integrand size = 27, antiderivative size = 61

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{(2ac + bd)\operatorname{arctanh}(\sin(e + fx))}{2f} + \frac{(bc + ad)\tan(e + fx)}{f} + \frac{bd \sec(e + fx)\tan(e + fx)}{2f}$$

output `1/2*(2*a*c+b*d)*arctanh(sin(f*x+e))/f+(a*d+b*c)*tan(f*x+e)/f+1/2*b*d*sec(f*x+e)*tan(f*x+e)/f`

3.247.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.23

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{a\operatorname{arctanh}(\sin(e + fx))}{f} + \frac{b\operatorname{arctanh}(\sin(e + fx))}{2f}$$

$$+ \frac{bc \tan(e + fx)}{f} + \frac{ad \tan(e + fx)}{f} + \frac{bd \sec(e + fx)\tan(e + fx)}{2f}$$

input `Integrate[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x]),x]`

output `(a*c*ArcTanh[Sin[e + f*x]])/f + (b*d*ArcTanh[Sin[e + f*x]])/(2*f) + (b*c*Tan[e + f*x])/f + (a*d*Tan[e + f*x])/f + (b*d*Sec[e + f*x]*Tan[e + f*x])/(2*f)`

3.247.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.05, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {3042, 4485, 3042, 4274, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(a + b \csc\left(e + fx + \frac{\pi}{2}\right)\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4485} \\
 & \frac{1}{2} \int \sec(e + fx)(2ac + bd + 2(bc + ad) \sec(e + fx)) dx + \frac{bd \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \csc\left(e + fx + \frac{\pi}{2}\right) \left(2ac + bd + 2(bc + ad) \csc\left(e + fx + \frac{\pi}{2}\right)\right) dx + \frac{bd \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{4274} \\
 & \frac{1}{2} \left(2(ad + bc) \int \sec^2(e + fx) dx + (2ac + bd) \int \sec(e + fx) dx \right) + \frac{bd \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left((2ac + bd) \int \csc\left(e + fx + \frac{\pi}{2}\right) dx + 2(ad + bc) \int \csc\left(e + fx + \frac{\pi}{2}\right)^2 dx \right) + \frac{bd \tan(e + fx) \sec(e + fx)}{2f} \\
 & \quad \downarrow \text{4254} \\
 & \frac{1}{2} \left((2ac + bd) \int \csc\left(e + fx + \frac{\pi}{2}\right) dx - \frac{2(ad + bc) \int 1d(-\tan(e + fx))}{f} \right) + \frac{bd \tan(e + fx) \sec(e + fx)}{2f}
 \end{aligned}$$

3.247. $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$

$$\frac{1}{2} \left((2ac + bd) \int \csc \left(e + fx + \frac{\pi}{2} \right) dx + \frac{2(ad + bc) \tan(e + fx)}{f} \right) + \frac{bd \tan(e + fx) \sec(e + fx)}{2f}$$

$$\frac{1}{2} \left(\frac{(2ac + bd) \operatorname{arctanh}(\sin(e + fx))}{f} + \frac{2(ad + bc) \tan(e + fx)}{f} \right) + \frac{bd \tan(e + fx) \sec(e + fx)}{2f}$$

input `Int[Sec[e + f*x]*(a + b*Sec[e + f*x])*(c + d*Sec[e + f*x]),x]`

output `(b*d*Sec[e + f*x]*Tan[e + f*x])/(2*f) + (((2*a*c + b*d)*ArcTanh[Sin[e + f*x]])/f + (2*(b*c + a*d)*Tan[e + f*x])/f)/2`

3.247.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

3.247.4 Maple [A] (verified)

Time = 2.28 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.23

method	result
derivativedivides	$\frac{ac \ln(\sec(fx+e)+\tan(fx+e))+ad \tan(fx+e)+bc \tan(fx+e)+bd \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)}{f}$
default	$\frac{ac \ln(\sec(fx+e)+\tan(fx+e))+ad \tan(fx+e)+bc \tan(fx+e)+bd \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)}{f}$
parts	$\frac{(ad+bc) \tan(fx+e)}{f} + \frac{ac \ln(\sec(fx+e)+\tan(fx+e))}{f} + \frac{bd \left(\frac{\sec(fx+e) \tan(fx+e)}{2} + \frac{\ln(\sec(fx+e)+\tan(fx+e))}{2}\right)}{f}$
parallelrisc	$\frac{-\left(ac + \frac{bd}{2}\right)(1+\cos(2fx+2e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) + \left(ac + \frac{bd}{2}\right)(1+\cos(2fx+2e)) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) + (ad+bc) \sin(2fx)}{f(1+\cos(2fx+2e))}$
norman	$\frac{(2ad+2bc+bd) \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - (2ad+2bc-bd) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{f \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^2} - \frac{(2ac+bd) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{2f} + \frac{(2ac+bd) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2f}$
risc	$-\frac{i(bde^{3i(fx+e)} - 2ade^{2i(fx+e)} - 2bce^{2i(fx+e)} - bde^{i(fx+e)} - 2ad - 2bc)}{f(1+e^{2i(fx+e)})^2} - \frac{ac \ln(e^{i(fx+e)} - i)}{f} - \frac{\ln(e^{i(fx+e)} - i)bd}{2f} +$

```
input int(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f*(a*c*ln(sec(f*x+e)+tan(f*x+e))+a*d*tan(f*x+e)+b*c*tan(f*x+e)+b*d*(1/2*
sec(f*x+e)*tan(f*x+e)+1/2*ln(sec(f*x+e)+tan(f*x+e))))
```

3.247.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.57

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{(2ac + bd) \cos(fx + e)^2 \log(\sin(fx + e) + 1) - (2ac + bd) \cos(fx + e)^2 \log(-\sin(fx + e) + 1) + 2(bd \sin(fx + e))}{4f \cos(fx + e)^2}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `1/4*((2*a*c + b*d)*cos(f*x + e)^2*log(sin(f*x + e) + 1) - (2*a*c + b*d)*cos(f*x + e)^2*log(-sin(f*x + e) + 1) + 2*(b*d + 2*(b*c + a*d)*cos(f*x + e))*sin(f*x + e))/(f*cos(f*x + e)^2)`

3.247.6 Sympy [F]

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \int (a + b \sec(e + fx))(c + d \sec(e + fx)) \sec(e + fx) dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e)),x)`

output `Integral((a + b*sec(e + f*x))*(c + d*sec(e + f*x))*sec(e + f*x), x)`

3.247.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.44

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx =$$

$$\frac{bd \left(\frac{2 \sin(fx+e)}{\sin(fx+e)^2-1} - \log(\sin(fx+e)+1) + \log(\sin(fx+e)-1) \right) - 4ac \log(\sec(fx+e) + \tan(fx+e))}{4f}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `-1/4*(b*d*(2*sin(f*x + e)/(sin(f*x + e)^2 - 1) - log(sin(f*x + e) + 1) + log(sin(f*x + e) - 1)) - 4*a*c*log(sec(f*x + e) + tan(f*x + e)) - 4*b*c*tan(f*x + e) - 4*a*d*tan(f*x + e))/f`

3.247.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(57) = 114.

Time = 0.31 (sec) , antiderivative size = 153, normalized size of antiderivative = 2.51

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{(2ac + bd) \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + 1 \right| \right) - (2ac + bd) \log \left(\left| \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - 1 \right| \right) - \frac{2 \left(2bc \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + 2ad \right)}{2f}}{2f}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))*(c+d*sec(f*x+e)),x, algorithm="giac")`

output `1/2*((2*a*c + b*d)*log(abs(tan(1/2*f*x + 1/2*e) + 1)) - (2*a*c + b*d)*log(abs(tan(1/2*f*x + 1/2*e) - 1)) - 2*(2*b*c*tan(1/2*f*x + 1/2*e)^3 + 2*a*d*tan(1/2*f*x + 1/2*e)^3 - b*d*tan(1/2*f*x + 1/2*e)^3 - 2*b*c*tan(1/2*f*x + 1/2*e) - 2*a*d*tan(1/2*f*x + 1/2*e) - b*d*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^2 - 1)^2)/f`

3.247.9 Mupad [B] (verification not implemented)

Time = 14.56 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.70

$$\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$$

$$= \frac{\operatorname{atanh} \left(\tan \left(\frac{e}{2} + \frac{fx}{2} \right) \right) (2ac + bd)}{f} + \frac{\tan \left(\frac{e}{2} + \frac{fx}{2} \right) (2ad + 2bc + bd) - \tan \left(\frac{e}{2} + \frac{fx}{2} \right)^3 (2ad + 2bc - bd)}{f \left(\tan \left(\frac{e}{2} + \frac{fx}{2} \right)^4 - 2 \tan \left(\frac{e}{2} + \frac{fx}{2} \right)^2 + 1 \right)}$$

3.247. $\int \sec(e + fx)(a + b \sec(e + fx))(c + d \sec(e + fx)) dx$

input `int(((a + b/cos(e + f*x))*(c + d/cos(e + f*x)))/cos(e + f*x),x)`

output `(atanh(tan(e/2 + (f*x)/2))*(2*a*c + b*d))/f + (tan(e/2 + (f*x)/2)*(2*a*d + 2*b*c + b*d) - tan(e/2 + (f*x)/2)^3*(2*a*d + 2*b*c - b*d))/(f*(tan(e/2 + (f*x)/2)^4 - 2*tan(e/2 + (f*x)/2)^2 + 1))`

output $((2*(b*c - a*d)*\text{ArcTanh}[((-c + d)*\text{Tan}[(e + f*x)/2])/\text{Sqrt}[c^2 - d^2]])/\text{Sqrt}[c^2 - d^2] + b*(-\text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] + \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]]))/ (d*f)$

3.248.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4486, 3042, 4257, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(a+b\sec(e+fx))}{c+d\sec(e+fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(a+b\csc(e+fx+\frac{\pi}{2}))}{c+d\csc(e+fx+\frac{\pi}{2})} dx \\ & \quad \downarrow \text{4486} \\ & \frac{b \int \sec(e+fx) dx}{d} - \frac{(bc-ad) \int \frac{\sec(e+fx)}{c+d\sec(e+fx)} dx}{d} \\ & \quad \downarrow \text{3042} \\ & \frac{b \int \csc(e+fx+\frac{\pi}{2}) dx}{d} - \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{c+d\csc(e+fx+\frac{\pi}{2})} dx}{d} \\ & \quad \downarrow \text{4257} \\ & \frac{\text{barctanh}(\sin(e+fx))}{df} - \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{c+d\csc(e+fx+\frac{\pi}{2})} dx}{d} \\ & \quad \downarrow \text{4318} \\ & \frac{\text{barctanh}(\sin(e+fx))}{df} - \frac{(bc-ad) \int \frac{1}{\frac{c \cos(e+fx)}{d} + 1} dx}{d^2} \\ & \quad \downarrow \text{3042} \\ & \frac{\text{barctanh}(\sin(e+fx))}{df} - \frac{(bc-ad) \int \frac{1}{\frac{c \sin(e+fx+\frac{\pi}{2})}{d} + 1} dx}{d^2} \end{aligned}$$

3.248. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{c+d\sec(e+fx)} dx$

$$\begin{aligned} & \downarrow 3138 \\ & \frac{\operatorname{barctanh}(\sin(e+fx))}{df} - \frac{2(bc-ad) \int \frac{1}{(1-\frac{c}{d}) \tan^2(\frac{1}{2}(e+fx)) + \frac{c+d}{d}} d \tan(\frac{1}{2}(e+fx))}{d^2 f} \\ & \downarrow 221 \\ & \frac{\operatorname{barctanh}(\sin(e+fx))}{df} - \frac{2(bc-ad) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{df \sqrt{c-d} \sqrt{c+d}} \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + b*Sec[e + f*x]))/(c + d*Sec[e + f*x]),x]`

output `(b*ArcTanh[Sin[e + f*x]])/(d*f) - (2*(b*c - a*d)*ArcTanh[(Sqrt[c - d]*Tan[e + f*x]/2])/Sqrt[c + d])/Sqrt[c - d]*d*Sqrt[c + d]*f)`

3.248.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4318 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

```
rule 4486 Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Simp[B/b Int[Csc[e + f*x], x], x] + Simp[(A*b - a*B)/b Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0]
```

3.248.4 Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{b \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) - b \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \frac{2(-ad+bc) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{d\sqrt{(c+d)(c-d)}}}{f}$
default	$\frac{b \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right) - b \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right) - \frac{2(-ad+bc) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{d\sqrt{(c+d)(c-d)}}}{f}$
risch	$\frac{\ln\left(\frac{e^{i(fx+e)} + ic^2 - id^2 + \sqrt{c^2 - d^2} d}{\sqrt{c^2 - d^2} c}\right) a}{\sqrt{c^2 - d^2} f} - \frac{\ln\left(\frac{e^{i(fx+e)} + ic^2 - id^2 + \sqrt{c^2 - d^2} d}{\sqrt{c^2 - d^2} c}\right) bc}{\sqrt{c^2 - d^2} fd} - \frac{\ln\left(\frac{e^{i(fx+e)} - ic^2 - id^2 - \sqrt{c^2 - d^2} d}{\sqrt{c^2 - d^2} c}\right) a}{\sqrt{c^2 - d^2} f}$

```
input int(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f*(b/d*ln(tan(1/2*f*x+1/2*e)+1)-b/d*ln(tan(1/2*f*x+1/2*e)-1)-2/d*(-a*d+b*c)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))
```

3.248.5 Fricas [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 316, normalized size of antiderivative = 4.16

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{c + d \sec(e + fx)} dx$$

$$= \left[\frac{(bc - ad)\sqrt{c^2 - d^2} \log\left(\frac{2cd \cos(fx+e) - (c^2 - 2d^2) \cos(fx+e)^2 + 2\sqrt{c^2 - d^2}(d \cos(fx+e) + c) \sin(fx+e) + 2c^2 - d^2}{c^2 \cos(fx+e)^2 + 2cd \cos(fx+e) + d^2}\right) - (bc^2 - bd^2)}{2(c^2d - d^3)f} \right. \\ \left. - \frac{2(bc - ad)\sqrt{-c^2 + d^2} \arctan\left(-\frac{\sqrt{-c^2 + d^2}(d \cos(fx+e) + c)}{(c^2 - d^2) \sin(fx+e)}\right) - (bc^2 - bd^2) \log(\sin(fx + e) + 1) + (bc^2 - bd^2)}{2(c^2d - d^3)f} \right]$$

3.248. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{c+d\sec(e+fx)} dx$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `[-1/2*((b*c - a*d)*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) - (b*c^2 - b*d^2)*log(sin(f*x + e) + 1) + (b*c^2 - b*d^2)*log(-sin(f*x + e) + 1))/((c^2*d - d^3)*f), -1/2*(2*(b*c - a*d)*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) - (b*c^2 - b*d^2)*log(sin(f*x + e) + 1) + (b*c^2 - b*d^2)*log(-sin(f*x + e) + 1))/((c^2*d - d^3)*f)]`

3.248.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + b\sec(e + fx))}{c + d\sec(e + fx)} dx = \int \frac{(a + b\sec(e + fx))\sec(e + fx)}{c + d\sec(e + fx)} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x)`

output `Integral((a + b*sec(e + f*x))*sec(e + f*x)/(c + d*sec(e + f*x)), x)`

3.248.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + b\sec(e + fx))}{c + d\sec(e + fx)} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` for more de`

3.248.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.67

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{c+d\sec(e+fx)} dx$$

$$= \frac{\frac{b \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{d} - \frac{b \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{d} + \frac{2 \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan\left(\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}}\right) \right) (bc-a)}{\sqrt{-c^2+d^2}d}}{f}$$

```
input integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="giac")
```

```
output (b*log(abs(tan(1/2*f*x + 1/2*e) + 1))/d - b*log(abs(tan(1/2*f*x + 1/2*e) - 1))/d + 2*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*(b*c - a*d)/(sqrt(-c^2 + d^2)*d))/f
```

3.248.9 Mupad [B] (verification not implemented)

Time = 14.54 (sec) , antiderivative size = 573, normalized size of antiderivative = 7.54

$$\begin{aligned}
& \int \frac{\sec(e+fx)(a+b\sec(e+fx))}{c+d\sec(e+fx)} dx \\
&= \frac{a c^2 \ln\left(\frac{c \sin\left(\frac{e+fx}{2}\right) - d \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{c^2-d^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f (c^2-d^2)^{3/2}} \\
&\quad - \frac{a d^2 \ln\left(\frac{c \sin\left(\frac{e+fx}{2}\right) - d \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{c^2-d^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f (c^2-d^2)^{3/2}} - \frac{2 b d \operatorname{atanh}\left(\frac{\sin\left(\frac{e+fx}{2}\right)}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f (c^2-d^2)} \\
&\quad - \frac{a \ln\left(\frac{c \cos\left(\frac{e+fx}{2}\right) + d \cos\left(\frac{e+fx}{2}\right) - \sin\left(\frac{e+fx}{2}\right) \sqrt{c^2-d^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f (c^2-d^2)} \sqrt{(c+d)(c-d)} \\
&\quad + \frac{b c d \ln\left(\frac{c \sin\left(\frac{e+fx}{2}\right) - d \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{c^2-d^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f (c^2-d^2)^{3/2}} \\
&\quad + \frac{2 b c^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{e+fx}{2}\right)}{\cos\left(\frac{e+fx}{2}\right)}\right)}{d f (c^2-d^2)} - \frac{b c^3 \ln\left(\frac{c \sin\left(\frac{e+fx}{2}\right) - d \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{c^2-d^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{d f (c^2-d^2)^{3/2}} \\
&\quad + \frac{b c \ln\left(\frac{c \cos\left(\frac{e+fx}{2}\right) + d \cos\left(\frac{e+fx}{2}\right) - \sin\left(\frac{e+fx}{2}\right) \sqrt{c^2-d^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{d f (c^2-d^2)} \sqrt{(c+d)(c-d)}
\end{aligned}$$

input `int((a + b/cos(e + f*x))/(cos(e + f*x)*(c + d/cos(e + f*x))),x)`

output

```
(a*c^2*log((c*sin(e/2 + (f*x)/2) - d*sin(e/2 + (f*x)/2) + cos(e/2 + (f*x)/2)*(c^2 - d^2)^(1/2))/cos(e/2 + (f*x)/2)))/(f*(c^2 - d^2)^(3/2)) - (a*d^2*log((c*sin(e/2 + (f*x)/2) - d*sin(e/2 + (f*x)/2) + cos(e/2 + (f*x)/2)*(c^2 - d^2)^(1/2))/cos(e/2 + (f*x)/2)))/(f*(c^2 - d^2)^(3/2)) - (2*b*d*atanh(sin(e/2 + (f*x)/2)/cos(e/2 + (f*x)/2)))/(f*(c^2 - d^2)) - (a*log((c*cos(e/2 + (f*x)/2) + d*cos(e/2 + (f*x)/2) - sin(e/2 + (f*x)/2)*(c^2 - d^2)^(1/2))/cos(e/2 + (f*x)/2))*((c + d)*(c - d))^(1/2))/(f*(c^2 - d^2)) + (b*c*d*log((c*sin(e/2 + (f*x)/2) - d*sin(e/2 + (f*x)/2) + cos(e/2 + (f*x)/2)*(c^2 - d^2)^(1/2))/cos(e/2 + (f*x)/2)))/(f*(c^2 - d^2)^(3/2)) + (2*b*c^2*atanh(sin(e/2 + (f*x)/2)/cos(e/2 + (f*x)/2)))/(d*f*(c^2 - d^2)) - (b*c^3*log((c*sin(e/2 + (f*x)/2) - d*sin(e/2 + (f*x)/2) + cos(e/2 + (f*x)/2)*(c^2 - d^2)^(1/2))/cos(e/2 + (f*x)/2)))/(d*f*(c^2 - d^2)^(3/2)) + (b*c*log((c*cos(e/2 + (f*x)/2) + d*cos(e/2 + (f*x)/2) - sin(e/2 + (f*x)/2)*(c^2 - d^2)^(1/2))/cos(e/2 + (f*x)/2))*((c + d)*(c - d))^(1/2))/(d*f*(c^2 - d^2))
```

3.249 $\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^2} dx$

3.249.1 Optimal result 1822
 3.249.2 Mathematica [A] (verified) 1822
 3.249.3 Rubi [A] (verified) 1823
 3.249.4 Maple [A] (verified) 1825
 3.249.5 Fricas [A] (verification not implemented) 1826
 3.249.6 Sympy [F] 1826
 3.249.7 Maxima [F(-2)] 1827
 3.249.8 Giac [A] (verification not implemented) 1827
 3.249.9 Mupad [B] (verification not implemented) 1828

3.249.1 Optimal result

Integrand size = 29, antiderivative size = 99

$$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^2} dx = \frac{2(ac-bd) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{(c-d)^{3/2}(c+d)^{3/2} f} + \frac{(bc-ad) \tan(e+fx)}{(c^2-d^2) f(c+d \sec(e+fx))}$$

output `2*(a*c-b*d)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c-d)^(3/2)/(c+d)^(3/2)/f+(-a*d+b*c)*tan(f*x+e)/(c^2-d^2)/f/(c+d*sec(f*x+e))`

3.249.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.98

$$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^2} dx = \frac{2(ac-bd) \operatorname{arctanh}\left(\frac{(-c+d) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{(c^2-d^2)^{3/2}} + \frac{(bc-ad) \sin(e+fx)}{(c-d)(c+d)(d+c \cos(e+fx))} f$$

input `Integrate[(Sec[e + f*x]*(a + b*Sec[e + f*x]))/(c + d*Sec[e + f*x])^2,x]`

output $((-2*(a*c - b*d)*\text{ArcTanh}[((-c + d)*\text{Tan}[(e + f*x)/2])/\text{Sqrt}[c^2 - d^2]])/(c^2 - d^2)^{(3/2)} + ((b*c - a*d)*\text{Sin}[e + f*x])/((c - d)*(c + d)*(d + c*\text{Cos}[e + f*x]))) / f$

3.249.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.11, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 4491, 25, 27, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{(c + d \sec(e + fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})(a + b \csc(e + fx + \frac{\pi}{2}))}{(c + d \csc(e + fx + \frac{\pi}{2}))^2} dx$$

↓ 4491

$$\frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))} - \frac{\int -\frac{(ac - bd) \sec(e + fx)}{c + d \sec(e + fx)} dx}{c^2 - d^2}$$

↓ 25

$$\frac{\int \frac{(ac - bd) \sec(e + fx)}{c + d \sec(e + fx)} dx}{c^2 - d^2} + \frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))}$$

↓ 27

$$\frac{(ac - bd) \int \frac{\sec(e + fx)}{c + d \sec(e + fx)} dx}{c^2 - d^2} + \frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))}$$

↓ 3042

$$\frac{(ac - bd) \int \frac{\csc(e + fx + \frac{\pi}{2})}{c + d \csc(e + fx + \frac{\pi}{2})} dx}{c^2 - d^2} + \frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))}$$

↓ 4318

$$\frac{(ac - bd) \int \frac{1}{d \cos(e + fx) + 1} dx}{d(c^2 - d^2)} + \frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))}$$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \frac{(ac - bd) \int \frac{1}{\frac{c \sin(e+fx+\frac{\pi}{2})}{d} + 1} dx}{d(c^2 - d^2)} + \frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))} \\
 & \downarrow \text{3138} \\
 & \frac{2(ac - bd) \int \frac{1}{(1 - \frac{c}{d}) \tan^2(\frac{1}{2}(e+fx)) + \frac{c+d}{d}} d \tan(\frac{1}{2}(e + fx))}{df(c^2 - d^2)} + \frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))} \\
 & \downarrow \text{221} \\
 & \frac{2(ac - bd) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{f\sqrt{c-d}\sqrt{c+d}(c^2 - d^2)} + \frac{(bc - ad) \tan(e + fx)}{f(c^2 - d^2)(c + d \sec(e + fx))}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(a + b*Sec[e + f*x]))/(c + d*Sec[e + f*x])^2,x]`

output `(2*(a*c - b*d)*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]]/(Sqrt[c - d]*Sqrt[c + d]*(c^2 - d^2)*f) + ((b*c - a*d)*Tan[e + f*x])/((c^2 - d^2)*f*(c + d*Sec[e + f*x]))`

3.249.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4318 `Int[csc[(e_) + (f_)*(x_)]/(csc[(e_) + (f_)*(x_)]*(b_) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4491 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(B_) + (A_)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[(a*A - b*B)*(m + 1) - (A*b - a*B)*(m + 2)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.249.4 Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.33

method	result
derivativedivides	$\frac{2(ad-bc) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^2-d^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)} + \frac{2(ac-bd) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c+d)(c-d)\sqrt{(c+d)(c-d)}}$
default	$\frac{2(ad-bc) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^2-d^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d\right)} + \frac{2(ac-bd) \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(c+d)(c-d)\sqrt{(c+d)(c-d)}}$
risch	$\frac{2i(-ad+bc)(de^{i(fx+e)}+c)}{c(c^2-d^2)f(e^{2i(fx+e)}c+2de^{i(fx+e)}+c)} + \frac{\ln\left(e^{i(fx+e)} + \frac{ic^2-id^2+\sqrt{c^2-d^2}d}{\sqrt{c^2-d^2}c}\right)ac}{\sqrt{c^2-d^2}(c+d)(c-d)f} - \frac{\ln\left(e^{i(fx+e)} + \frac{ic^2-id^2+\sqrt{c^2-d^2}d}{\sqrt{c^2-d^2}c}\right)}{\sqrt{c^2-d^2}(c+d)(c-d)f}$

input `int(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

3.249.
$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^2} dx$$

output $\frac{1}{f} \cdot \frac{2(a*d-b*c)}{(c^2-d^2)} \cdot \frac{\tan(1/2*f*x+1/2*e)}{(\tan(1/2*f*x+1/2*e)^2*c-\tan(1/2*f*x+1/2*e)^2*d-c-d)} + \frac{2(a*c-b*d)}{(c+d)} \cdot \frac{1}{(c-d)} \cdot \frac{1}{((c+d)*(c-d))^{1/2}} \cdot \operatorname{arctan} \left(\frac{h((c-d)*\tan(1/2*f*x+1/2*e))}{((c+d)*(c-d))^{1/2}} \right)$

3.249.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 389, normalized size of antiderivative = 3.93

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^2} dx$$

$$= \left[\frac{(acd - bd^2 + (ac^2 - bcd) \cos(fx + e)) \sqrt{c^2 - d^2} \log \left(\frac{2cd \cos(fx+e) - (c^2 - 2d^2) \cos(fx+e)^2 + 2\sqrt{c^2 - d^2} (d \cos(fx+e) + c) \sin(fx+e)}{c^2 \cos(fx+e)^2 + 2cd \cos(fx+e) + d^2} \right)}{2((c^5 - 2c^3d^2 + cd^4)f \cos(fx + e) + (c^4d - 2c^2d^3 + d^5)f)} \right]$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="fricas")`

output $\left[\frac{1}{2} \cdot \frac{(a*c*d - b*d^2 + (a*c^2 - b*c*d)*\cos(f*x + e))*\sqrt{c^2 - d^2}*\log((2*c*d*\cos(f*x + e) - (c^2 - 2*d^2)*\cos(f*x + e)^2 + 2*\sqrt{c^2 - d^2}*(d*\cos(f*x + e) + c)*\sin(f*x + e) + 2*c^2 - d^2)/(c^2*\cos(f*x + e)^2 + 2*c*d*\cos(f*x + e) + d^2)) + 2*(b*c^3 - a*c^2*d - b*c*d^2 + a*d^3)*\sin(f*x + e)}{((c^5 - 2*c^3*d^2 + c*d^4)*f*\cos(f*x + e) + (c^4*d - 2*c^2*d^3 + d^5)*f)}, \frac{(a*c*d - b*d^2 + (a*c^2 - b*c*d)*\cos(f*x + e))*\sqrt{-c^2 + d^2}*\operatorname{arctan}(\sqrt{-c^2 + d^2}*(d*\cos(f*x + e) + c)/((c^2 - d^2)*\sin(f*x + e))) + (b*c^3 - a*c^2*d - b*c*d^2 + a*d^3)*\sin(f*x + e)}{((c^5 - 2*c^3*d^2 + c*d^4)*f*\cos(f*x + e) + (c^4*d - 2*c^2*d^3 + d^5)*f)} \right]$

3.249.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^2} dx = \int \frac{(a+b\sec(e+fx))\sec(e+fx)}{(c+d\sec(e+fx))^2} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))**2,x)`

output `Integral((a + b*sec(e + f*x))*sec(e + f*x)/(c + d*sec(e + f*x))**2, x)`

3.249.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{(c + d \sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de

3.249.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.74

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{(c + d \sec(e + fx))^2} dx =$$

$$2 \left(\frac{\left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2c-2d) + \arctan\left(\frac{c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) - d \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)}{\sqrt{-c^2+d^2}}\right) \right) (ac-bd)}{(c^2-d^2)\sqrt{-c^2+d^2}} \right) + \frac{bc \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) - ad \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)}{\left(c \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - d \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - c - d\right) (c^2 - d^2)}$$

f

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="giac")`

output `-2*((pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))*(a*c - b*d)/((c^2 - d^2)*sqrt(-c^2 + d^2)) + (b*c*tan(1/2*f*x + 1/2*e) - a*d*tan(1/2*f*x + 1/2*e))/((c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)*(c^2 - d^2)))/f`

3.249.9 Mupad [B] (verification not implemented)

Time = 13.88 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.07

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^2} dx = \frac{2 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \sqrt{c-d}}{\sqrt{c+d}}\right) (ac-bd)}{f(c+d)^{3/2}(c-d)^{3/2}} - \frac{2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) (ad-bc)}{f(c+d)(c-d)\left((d-c)\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + c+d\right)}$$

input `int((a + b/cos(e + f*x))/(cos(e + f*x)*(c + d/cos(e + f*x))^2),x)`output `(2*atanh((tan(e/2 + (f*x)/2)*(c - d)^(1/2))/(c + d)^(1/2))*(a*c - b*d))/(f*(c + d)^(3/2)*(c - d)^(3/2)) - (2*tan(e/2 + (f*x)/2)*(a*d - b*c))/(f*(c + d)*(c - d)*(c + d - tan(e/2 + (f*x)/2)^2*(c - d)))`

3.250
$$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^3} dx$$

3.250.1 Optimal result	1829
3.250.2 Mathematica [A] (verified)	1829
3.250.3 Rubi [A] (verified)	1830
3.250.4 Maple [A] (verified)	1833
3.250.5 Fricas [B] (verification not implemented)	1833
3.250.6 Sympy [F]	1834
3.250.7 Maxima [F(-2)]	1834
3.250.8 Giac [B] (verification not implemented)	1835
3.250.9 Mupad [B] (verification not implemented)	1836

3.250.1 Optimal result

Integrand size = 29, antiderivative size = 166

$$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^3} dx = -\frac{(3bcd - a(2c^2 + d^2)) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{(c-d)^{5/2}(c+d)^{5/2}f} + \frac{(bc - ad) \tan(e+fx)}{2(c^2 - d^2) f(c+d \sec(e+fx))^2} - \frac{(3acd - b(c^2 + 2d^2)) \tan(e+fx)}{2(c^2 - d^2)^2 f(c+d \sec(e+fx))}$$

```
output - (3*b*c*d-a*(2*c^2+d^2))*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2)))/(c-d)^(5/2)/(c+d)^(5/2)/f+1/2*(-a*d+b*c)*tan(f*x+e)/(c^2-d^2)/f/(c+d*sec(c(f*x+e))^2-1/2*(3*a*c*d-b*(c^2+2*d^2))*tan(f*x+e)/(c^2-d^2)^2/f/(c+d*sec(f*x+e)))
```

3.250.2 Mathematica [A] (verified)

Time = 1.14 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.04

$$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^3} dx = \frac{2(-3bcd+a(2c^2+d^2)) \operatorname{arctanh}\left(\frac{(-c+d) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{(c^2-d^2)^{5/2}} + \frac{d(-bc+ad) \sin(e+fx)}{c(c-d)(c+d)(d+c \cos(e+fx))^2} + \frac{(ad(-4c^2+d^2)+bc(2c^2+d^2)) \sin(e+fx)}{c(c-d)^2(c+d)^2(d+c \cos(e+fx))}$$

$2f$

3.250.
$$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^3} dx$$

input `Integrate[(Sec[e + f*x]*(a + b*Sec[e + f*x]))/(c + d*Sec[e + f*x])^3,x]`

output `((-2*(-3*b*c*d + a*(2*c^2 + d^2))*ArcTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]])/(c^2 - d^2)^(5/2) + (d*(-(b*c) + a*d)*Sin[e + f*x])/(c*(c - d)*(c + d)*(d + c*Cos[e + f*x])^2) + ((a*d*(-4*c^2 + d^2) + b*c*(2*c^2 + d^2))*Sin[e + f*x])/(c*(c - d)^2*(c + d)^2*(d + c*Cos[e + f*x]))/(2*f)`

3.250.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.15, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 4491, 25, 3042, 4491, 27, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(a+b\csc(e+fx+\frac{\pi}{2}))}{(c+d\csc(e+fx+\frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{4491} \\
 & \frac{(bc-ad)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} - \int \frac{\sec(e+fx)(2(ac-bd)+(bc-ad)\sec(e+fx))}{2(c^2-d^2)(c+d\sec(e+fx))^2} dx \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{\sec(e+fx)(2(ac-bd)+(bc-ad)\sec(e+fx))}{(c+d\sec(e+fx))^2} dx}{2(c^2-d^2)} + \frac{(bc-ad)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(2(ac-bd)+(bc-ad)\csc(e+fx+\frac{\pi}{2}))}{(c+d\csc(e+fx+\frac{\pi}{2}))^2} dx}{2(c^2-d^2)} + \frac{(bc-ad)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} \\
 & \quad \downarrow \text{4491} \\
 & \frac{-\int \frac{(3bcd-a(2c^2+d^2))\sec(e+fx)}{c^2-d^2} dx - \frac{(3acd-b(c^2+2d^2))\tan(e+fx)}{f(c^2-d^2)(c+d\sec(e+fx))}}{2(c^2-d^2)} + \frac{(bc-ad)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2}
 \end{aligned}$$

3.250. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{(3bcd-a(2c^2+d^2)) \int \frac{\sec(e+fx)}{c+d \sec(e+fx)} dx}{c^2-d^2} - \frac{(3acd-b(c^2+2d^2)) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))} + \frac{(bc-ad) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} \\
& \downarrow 3042 \\
& \frac{(3bcd-a(2c^2+d^2)) \int \frac{\csc(e+fx+\frac{\pi}{2})}{c+d \csc(e+fx+\frac{\pi}{2})} dx}{c^2-d^2} - \frac{(3acd-b(c^2+2d^2)) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))} + \frac{(bc-ad) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} \\
& \downarrow 4318 \\
& \frac{(3bcd-a(2c^2+d^2)) \int \frac{1}{\frac{c \cos(e+fx)}{d} + 1} dx}{d(c^2-d^2)} - \frac{(3acd-b(c^2+2d^2)) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))} + \frac{(bc-ad) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} \\
& \downarrow 3042 \\
& \frac{(3bcd-a(2c^2+d^2)) \int \frac{1}{\frac{c \sin(e+fx+\frac{\pi}{2})}{d} + 1} dx}{d(c^2-d^2)} - \frac{(3acd-b(c^2+2d^2)) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))} + \frac{(bc-ad) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} \\
& \downarrow 3138 \\
& \frac{2(3bcd-a(2c^2+d^2)) \int \frac{1}{(1-\frac{c}{d}) \tan^2(\frac{1}{2}(e+fx)) + \frac{c+d}{d}} d \tan(\frac{1}{2}(e+fx))}{df(c^2-d^2)} - \frac{(3acd-b(c^2+2d^2)) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))} + \\
& \quad \frac{(bc-ad) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} \\
& \downarrow 221 \\
& \frac{2(3bcd-a(2c^2+d^2)) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{f\sqrt{c-d}\sqrt{c+d}(c^2-d^2)} - \frac{(3acd-b(c^2+2d^2)) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))} + \\
& \quad \frac{(bc-ad) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(a + b*Sec[e + f*x]))/(c + d*Sec[e + f*x])^3,x]`

output `((b*c - a*d)*Tan[e + f*x])/(2*(c^2 - d^2)*f*(c + d*Sec[e + f*x])^2) + ((-2 * (3*b*c*d - a*(2*c^2 + d^2))*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]])/(Sqrt[c - d]*Sqrt[c + d]*(c^2 - d^2)*f) - ((3*a*c*d - b*(c^2 + 2*d^2))*Tan[e + f*x])/((c^2 - d^2)*f*(c + d*Sec[e + f*x]))/(2*(c^2 - d^2))`

3.250. $\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^3} dx$

3.250.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 4318 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`
- rule 4491 `Int[csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[(a*A - b*B)*(m + 1) - (A*b - a*B)*(m + 2)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.250.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.42

method	result
derivativedivides	$2 \left(-\frac{(4acd+ad^2-2bc^2-bcd-2bd^2) \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{2(c-d)(c^2+2cd+d^2)} + \frac{(4acd-ad^2-2bc^2+bcd-2bd^2) \tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{2(c+d)(c^2-2cd+d^2)} \right) \frac{(2ac^2+ad^2-3bcd) \arctan\left(\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{c-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}\right)}{(c^4-2c^2d^2+d^4)} + \frac{f}{(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 d - c - d)^2}$
default	$2 \left(-\frac{(4acd+ad^2-2bc^2-bcd-2bd^2) \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{2(c-d)(c^2+2cd+d^2)} + \frac{(4acd-ad^2-2bc^2+bcd-2bd^2) \tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{2(c+d)(c^2-2cd+d^2)} \right) \frac{(2ac^2+ad^2-3bcd) \arctan\left(\frac{\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{c-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}\right)}{(c^4-2c^2d^2+d^4)} + \frac{f}{(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 d - c - d)^2}$
risch	$\frac{i(-5ac^3d^2e^{3i(fx+e)}+2ac^4e^{3i(fx+e)}+3bc^4de^{3i(fx+e)}-4ac^4de^{2i(fx+e)}-7ac^2d^3e^{2i(fx+e)}+2ad^5e^{2i(fx+e)}+2bc^5e^{2i(fx+e)}+2bd^5e^{2i(fx+e)})}{c^2(-c^2+d^2)}$

```
input int(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 1/f*(-2*(-1/2*(4*a*c*d+a*d^2-2*b*c^2-b*c*d-2*b*d^2)/(c-d)/(c^2+2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3+1/2*(4*a*c*d-a*d^2-2*b*c^2+b*c*d-2*b*d^2)/(c+d)/(c^2-2*c*d+d^2)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^2+(2*a*c^2+a*d^2-3*b*c*d)/(c^4-2*c^2*d^2+d^4)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))
```

3.250.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 347 vs. 2(153) = 306.

Time = 0.32 (sec) , antiderivative size = 752, normalized size of antiderivative = 4.53

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$$

$$= \frac{\left[(2ac^2d^2 - 3bcd^3 + ad^4 + (2ac^4 - 3bc^3d + ac^2d^2) \cos^2(fx+e) + 2(2ac^3d - 3bc^2d^2 + acd^3) \cos(fx+e) + 2ad^5 \sin^2(fx+e) \right]}{4((c^8 - 3c^6d^2 + 3c^4d^4 - 3c^2d^6 + d^8))}$$

```
input integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="fracas")
```

3.250. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$

output `[1/4*((2*a*c^2*d^2 - 3*b*c*d^3 + a*d^4 + (2*a*c^4 - 3*b*c^3*d + a*c^2*d^2)*cos(f*x + e)^2 + 2*(2*a*c^3*d - 3*b*c^2*d^2 + a*c*d^3)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(b*c^4*d - 3*a*c^3*d^2 + b*c^2*d^3 + 3*a*c*d^4 - 2*b*d^5 + (2*b*c^5 - 4*a*c^4*d - b*c^3*d^2 + 5*a*c^2*d^3 - b*c*d^4 - a*d^5)*cos(f*x + e))*sin(f*x + e))/((c^8 - 3*c^6*d^2 + 3*c^4*d^4 - c^2*d^6)*f*cos(f*x + e)^2 + 2*(c^7*d - 3*c^5*d^3 + 3*c^3*d^5 - c*d^7)*f*cos(f*x + e) + (c^6*d^2 - 3*c^4*d^4 + 3*c^2*d^6 - d^8)*f), 1/2*((2*a*c^2*d^2 - 3*b*c*d^3 + a*d^4 + (2*a*c^4 - 3*b*c^3*d + a*c^2*d^2)*cos(f*x + e)^2 + 2*(2*a*c^3*d - 3*b*c^2*d^2 + a*c*d^3)*cos(f*x + e))*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (b*c^4*d - 3*a*c^3*d^2 + b*c^2*d^3 + 3*a*c*d^4 - 2*b*d^5 + (2*b*c^5 - 4*a*c^4*d - b*c^3*d^2 + 5*a*c^2*d^3 - b*c*d^4 - a*d^5)*cos(f*x + e))*sin(f*x + e))/((c^8 - 3*c^6*d^2 + 3*c^4*d^4 - c^2*d^6)*f*cos(f*x + e)^2 + 2*(c^7*d - 3*c^5*d^3 + 3*c^3*d^5 - c*d^7)*f*cos(f*x + e) + (c^6*d^2 - 3*c^4*d^4 + 3*c^2*d^6 - d^8)*f)]`

3.250.6 Sympy [F]

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{(c + d \sec(e + fx))^3} dx = \int \frac{(a + b \sec(e + fx)) \sec(e + fx)}{(c + d \sec(e + fx))^3} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))**3,x)`

output `Integral((a + b*sec(e + f*x))*sec(e + f*x)/(c + d*sec(e + f*x))**3, x)`

3.250.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{(c + d \sec(e + fx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de

3.250.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 399 vs. $2(153) = 306$.

Time = 0.38 (sec) , antiderivative size = 399, normalized size of antiderivative = 2.40

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$$

$$\frac{(2ac^2-3bcd+ad^2)\left(\pi\left[\frac{fx+e}{2\pi}+\frac{1}{2}\right]\operatorname{sgn}(-2c+2d)+\arctan\left(\frac{-c\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)-d\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)}{\sqrt{-c^2+d^2}}\right)\right)}{(c^4-2c^2d^2+d^4)\sqrt{-c^2+d^2}} - \frac{2bc^3\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)^3-4ac^2d\tan\left(\frac{1}{2}fx+\frac{1}{2}e\right)}{(c^4-2c^2d^2+d^4)\sqrt{-c^2+d^2}}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^3,x, algorithm="giac")`

output $((2*a*c^2 - 3*b*c*d + a*d^2)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(-2*c + 2*d) + arctan(-(c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x + 1/2*e))/sqrt(-c^2 + d^2)))/((c^4 - 2*c^2*d^2 + d^4)*sqrt(-c^2 + d^2)) - (2*b*c^3*tan(1/2*f*x + 1/2*e)^3 - 4*a*c^2*d*tan(1/2*f*x + 1/2*e)^3 - b*c^2*d*tan(1/2*f*x + 1/2*e)^3 + 3*a*c*d^2*tan(1/2*f*x + 1/2*e)^3 + b*c*d^2*tan(1/2*f*x + 1/2*e)^3 + a*d^3*tan(1/2*f*x + 1/2*e)^3 - 2*b*d^3*tan(1/2*f*x + 1/2*e)^3 - 2*b*c^3*tan(1/2*f*x + 1/2*e) + 4*a*c^2*d*tan(1/2*f*x + 1/2*e) - b*c^2*d*tan(1/2*f*x + 1/2*e) + 3*a*c*d^2*tan(1/2*f*x + 1/2*e) - b*c*d^2*tan(1/2*f*x + 1/2*e) - a*d^3*tan(1/2*f*x + 1/2*e) - 2*b*d^3*tan(1/2*f*x + 1/2*e))/((c^4 - 2*c^2*d^2 + d^4)*(c*tan(1/2*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^2))/f$

3.250.9 Mupad [B] (verification not implemented)

Time = 16.70 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.51

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^3} dx$$

$$= \frac{\frac{\tan\left(\frac{e}{2}+\frac{fx}{2}\right)(ad^2+2bc^2+2bd^2-4acd-bcd)}{(c+d)(c^2-2cd+d^2)} - \frac{\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^3(2bc^2-ad^2+2bd^2-4acd+bcd)}{(c+d)^2(c-d)}}{f\left(2cd - \tan\left(\frac{e}{2}+\frac{fx}{2}\right)^2(2c^2-2d^2) + \tan\left(\frac{e}{2}+\frac{fx}{2}\right)^4(c^2-2cd+d^2) + c^2+d^2\right)}$$

$$+ \frac{\operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2}+\frac{fx}{2}\right)(2c-2d)(c^2-2cd+d^2)}{2\sqrt{c+d}(c-d)^{5/2}}\right)(2ac^2-3bcd+ad^2)}{f(c+d)^{5/2}(c-d)^{5/2}}$$

input `int((a + b/cos(e + f*x))/(cos(e + f*x)*(c + d/cos(e + f*x))^3),x)`output `((tan(e/2 + (f*x)/2)*(a*d^2 + 2*b*c^2 + 2*b*d^2 - 4*a*c*d - b*c*d))/((c + d)*(c^2 - 2*c*d + d^2)) - (tan(e/2 + (f*x)/2)^3*(2*b*c^2 - a*d^2 + 2*b*d^2 - 4*a*c*d + b*c*d))/((c + d)^2*(c - d)))/(f*(2*c*d - tan(e/2 + (f*x)/2)^2*(2*c^2 - 2*d^2) + tan(e/2 + (f*x)/2)^4*(c^2 - 2*c*d + d^2) + c^2 + d^2)) + (atanh((tan(e/2 + (f*x)/2)*(2*c - 2*d)*(c^2 - 2*c*d + d^2))/(2*(c + d)^(1/2)*(c - d)^(5/2)))*(2*a*c^2 + a*d^2 - 3*b*c*d))/(f*(c + d)^(5/2)*(c - d)^(5/2))`

3.251 $\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^4} dx$

3.251.1 Optimal result 1837
 3.251.2 Mathematica [A] (verified) 1838
 3.251.3 Rubi [A] (verified) 1838
 3.251.4 Maple [A] (verified) 1842
 3.251.5 Fricas [B] (verification not implemented) 1843
 3.251.6 Sympy [F] 1843
 3.251.7 Maxima [F(-2)] 1844
 3.251.8 Giac [B] (verification not implemented) 1844
 3.251.9 Mupad [B] (verification not implemented) 1845

3.251.1 Optimal result

Integrand size = 29, antiderivative size = 237

$$\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^4} dx$$

$$= \frac{(2ac^3 - 4bc^2d + 3acd^2 - bd^3) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{(c-d)^{7/2}(c+d)^{7/2}f}$$

$$+ \frac{(bc-ad) \tan(e+fx)}{3(c^2-d^2)f(c+d \sec(e+fx))^3} + \frac{(2bc^2 - 5acd + 3bd^2) \tan(e+fx)}{6(c^2-d^2)^2 f(c+d \sec(e+fx))^2}$$

$$+ \frac{(2bc^3 - 11ac^2d + 13bcd^2 - 4ad^3) \tan(e+fx)}{6(c^2-d^2)^3 f(c+d \sec(e+fx))}$$

```
output (2*a*c^3+3*a*c*d^2-4*b*c^2*d-b*d^3)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)
/(c+d)^(1/2))/(c-d)^(7/2)/(c+d)^(7/2)/f+1/3*(-a*d+b*c)*tan(f*x+e)/(c^2-d^2)
)/f/(c+d*sec(f*x+e))^3+1/6*(-5*a*c*d+2*b*c^2+3*b*d^2)*tan(f*x+e)/(c^2-d^2)
^2/f/(c+d*sec(f*x+e))^2+1/6*(-11*a*c^2*d-4*a*d^3+2*b*c^3+13*b*c*d^2)*tan(f
*x+e)/(c^2-d^2)^3/f/(c+d*sec(f*x+e))
```

3.251.2 Mathematica [A] (verified)

Time = 2.01 (sec) , antiderivative size = 405, normalized size of antiderivative = 1.71

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$$

$$= \frac{(d+c\cos(e+fx))\sec^3(e+fx)(a+b\sec(e+fx)) \left(\frac{24(-bd(4c^2+d^2)+a(2c^3+3cd^2))\operatorname{arctanh}\left(\frac{(-c+d)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}} \right)}{\dots}$$

input `Integrate[(Sec[e + f*x]*(a + b*Sec[e + f*x]))/(c + d*Sec[e + f*x])^4,x]`

output

```
((d + c*Cos[e + f*x])*Sec[e + f*x]^3*(a + b*Sec[e + f*x])*((24*(-(b*d*(4*c^2 + d^2)) + a*(2*c^3 + 3*c*d^2))*ArcTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]])*(d + c*Cos[e + f*x])^3)/Sqrt[c^2 - d^2] - 6*b*c^5*Sin[e + f*x] + 18*a*c^4*d*Sin[e + f*x] - 18*b*c^3*d^2*Sin[e + f*x] + 39*a*c^2*d^3*Sin[e + f*x] - 51*b*c*d^4*Sin[e + f*x] + 18*a*d^5*Sin[e + f*x] - 12*b*c^4*d*Sin[2*(e + f*x)] + 54*a*c^3*d^2*Sin[2*(e + f*x)] - 54*b*c^2*d^3*Sin[2*(e + f*x)] + 6*a*c*d^4*Sin[2*(e + f*x)] + 6*b*d^5*Sin[2*(e + f*x)] - 6*b*c^5*Sin[3*(e + f*x)] + 18*a*c^4*d*Sin[3*(e + f*x)] - 10*b*c^3*d^2*Sin[3*(e + f*x)] - 5*a*c^2*d^3*Sin[3*(e + f*x)] + b*c*d^4*Sin[3*(e + f*x)] + 2*a*d^5*Sin[3*(e + f*x)]))/(24*(-c^2 + d^2)^3*f*(b + a*Cos[e + f*x])*(c + d*Sec[e + f*x])^4)
```

3.251.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.17, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$, Rules used = {3042, 4491, 25, 3042, 4491, 25, 3042, 4491, 27, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(a+b\csc\left(e+fx+\frac{\pi}{2}\right))}{(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^4} dx$$

3.251. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$

$$\begin{aligned}
& \downarrow 4491 \\
& \frac{(bc-ad)\tan(e+fx)}{3f(c^2-d^2)(c+d\sec(e+fx))^3} - \frac{\int -\frac{\sec(e+fx)(3(ac-bd)+2(bc-ad)\sec(e+fx))}{(c+d\sec(e+fx))^3} dx}{3(c^2-d^2)} \\
& \downarrow 25 \\
& \frac{\int \frac{\sec(e+fx)(3(ac-bd)+2(bc-ad)\sec(e+fx))}{(c+d\sec(e+fx))^3} dx}{3(c^2-d^2)} + \frac{(bc-ad)\tan(e+fx)}{3f(c^2-d^2)(c+d\sec(e+fx))^3} \\
& \downarrow 3042 \\
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(3(ac-bd)+2(bc-ad)\csc(e+fx+\frac{\pi}{2}))}{(c+d\csc(e+fx+\frac{\pi}{2}))^3} dx}{3(c^2-d^2)} + \frac{(bc-ad)\tan(e+fx)}{3f(c^2-d^2)(c+d\sec(e+fx))^3} \\
& \downarrow 4491 \\
& \frac{(-5acd+2bc^2+3bd^2)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} - \frac{\int -\frac{\sec(e+fx)(2(3ac^2-5bdc+2ad^2)+(2bc^2-5adc+3bd^2)\sec(e+fx))}{(c+d\sec(e+fx))^2} dx}{2(c^2-d^2)} + \\
& \frac{3(c^2-d^2)}{3f(c^2-d^2)(c+d\sec(e+fx))^3} \\
& \downarrow 25 \\
& \frac{\int \frac{\sec(e+fx)(2(3ac^2-5bdc+2ad^2)+(2bc^2-5adc+3bd^2)\sec(e+fx))}{(c+d\sec(e+fx))^2} dx}{2(c^2-d^2)} + \frac{(-5acd+2bc^2+3bd^2)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} + \\
& \frac{3(c^2-d^2)}{3f(c^2-d^2)(c+d\sec(e+fx))^3} \\
& \downarrow 3042 \\
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})(2(3ac^2-5bdc+2ad^2)+(2bc^2-5adc+3bd^2)\csc(e+fx+\frac{\pi}{2}))}{(c+d\csc(e+fx+\frac{\pi}{2}))^2} dx}{2(c^2-d^2)} + \frac{(-5acd+2bc^2+3bd^2)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} + \\
& \frac{3(c^2-d^2)}{3f(c^2-d^2)(c+d\sec(e+fx))^3} \\
& \downarrow 4491 \\
& \frac{(-11ac^2d-4ad^3+2bc^3+13bcd^2)\tan(e+fx)}{f(c^2-d^2)(c+d\sec(e+fx))} - \frac{\int -\frac{3(2ac^3-4bdc^2+3ad^2c-bd^3)\sec(e+fx)}{c+d\sec(e+fx)} dx}{c^2-d^2} + \frac{(-5acd+2bc^2+3bd^2)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} + \\
& \frac{3(c^2-d^2)}{3f(c^2-d^2)(c+d\sec(e+fx))^3} \\
& \downarrow 27
\end{aligned}$$

3.251. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$

$$\frac{\frac{3(2ac^3+3acd^2-4bc^2d-bd^3) \int \frac{\sec(e+fx)}{c+d \sec(e+fx)} dx + \frac{(-11ac^2d-4ad^3+2bc^3+13bcd^2) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))}}{c^2-d^2}}{2(c^2-d^2)} + \frac{(-5acd+2bc^2+3bd^2) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)(bc-ad) \tan(e+fx)}{3f(c^2-d^2)(c+d \sec(e+fx))^3}$$

↓ 3042

$$\frac{\frac{3(2ac^3+3acd^2-4bc^2d-bd^3) \int \frac{\csc(e+fx+\frac{\pi}{2})}{c+d \csc(e+fx+\frac{\pi}{2})} dx + \frac{(-11ac^2d-4ad^3+2bc^3+13bcd^2) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))}}{c^2-d^2}}{2(c^2-d^2)} + \frac{(-5acd+2bc^2+3bd^2) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)(bc-ad) \tan(e+fx)}{3f(c^2-d^2)(c+d \sec(e+fx))^3}$$

↓ 4318

$$\frac{\frac{3(2ac^3+3acd^2-4bc^2d-bd^3) \int \frac{1}{\frac{c \cos(e+fx)}{d} + 1} dx + \frac{(-11ac^2d-4ad^3+2bc^3+13bcd^2) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))}}{d(c^2-d^2)}}{2(c^2-d^2)} + \frac{(-5acd+2bc^2+3bd^2) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)(bc-ad) \tan(e+fx)}{3f(c^2-d^2)(c+d \sec(e+fx))^3}$$

↓ 3042

$$\frac{\frac{3(2ac^3+3acd^2-4bc^2d-bd^3) \int \frac{1}{\frac{c \sin(e+fx+\frac{\pi}{2})}{d} + 1} dx + \frac{(-11ac^2d-4ad^3+2bc^3+13bcd^2) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))}}{d(c^2-d^2)}}{2(c^2-d^2)} + \frac{(-5acd+2bc^2+3bd^2) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))^2} +$$

$$\frac{3(c^2-d^2)(bc-ad) \tan(e+fx)}{3f(c^2-d^2)(c+d \sec(e+fx))^3}$$

↓ 3138

$$\frac{\frac{6(2ac^3+3acd^2-4bc^2d-bd^3) \int \frac{1}{\left(1-\frac{c}{d}\right) \tan^2\left(\frac{1}{2}(e+fx)\right) + \frac{c+d}{d}} d \tan\left(\frac{1}{2}(e+fx)\right) + \frac{(-11ac^2d-4ad^3+2bc^3+13bcd^2) \tan(e+fx)}{f(c^2-d^2)(c+d \sec(e+fx))}}{df(c^2-d^2)}}{2(c^2-d^2)} + \frac{(-5acd+2bc^2+3bd^2) \tan(e+fx)}{2f(c^2-d^2)(c+d \sec(e+fx))} +$$

$$\frac{3(c^2-d^2)(bc-ad) \tan(e+fx)}{3f(c^2-d^2)(c+d \sec(e+fx))^3}$$

↓ 221

3.251. $\int \frac{\sec(e+fx)(a+b \sec(e+fx))}{(c+d \sec(e+fx))^4} dx$

$$\frac{6(2ac^3+3acd^2-4bc^2d-bd^3)\operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{f\sqrt{c-d}\sqrt{c+d}(c^2-d^2)} + \frac{(-11ac^2d-4ad^3+2bc^3+13bcd^2)\tan(e+fx)}{f(c^2-d^2)(c+d\sec(e+fx))} + \frac{(-5acd+2bc^2+3bd^2)\tan(e+fx)}{2f(c^2-d^2)(c+d\sec(e+fx))^2} + \frac{3(c^2-d^2)(bc-ad)\tan(e+fx)}{3f(c^2-d^2)(c+d\sec(e+fx))^3}$$

input `Int[(Sec[e + f*x]*(a + b*Sec[e + f*x]))/(c + d*Sec[e + f*x])^4,x]`

output `((b*c - a*d)*Tan[e + f*x])/(3*(c^2 - d^2)*f*(c + d*Sec[e + f*x])^3) + (((2 *b*c^2 - 5*a*c*d + 3*b*d^2)*Tan[e + f*x])/(2*(c^2 - d^2)*f*(c + d*Sec[e + f*x])^2) + ((6*(2*a*c^3 - 4*b*c^2*d + 3*a*c*d^2 - b*d^3)*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]])/(Sqrt[c - d]*Sqrt[c + d]*(c^2 - d^2)*f) + ((2*b*c^3 - 11*a*c^2*d + 13*b*c*d^2 - 4*a*d^3)*Tan[e + f*x])/((c^2 - d^2)*f*(c + d*Sec[e + f*x]))/(2*(c^2 - d^2)))/(3*(c^2 - d^2))`

3.251.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4318 `Int[csc[(e_.) + (f_.)*(x_.)]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4491 `Int[csc[(e_.) + (f_.)*(x_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-A*b - a*B)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[(a*A - b*B)*(m + 1) - (A*b - a*B)*(m + 2)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.251.4 Maple [A] (verified)

Time = 1.28 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.59

method	result
derivativedivides	$2 \left(-\frac{(6ac^2d + 3acd^2 + 2ad^3 - 2bc^3 - 2bc^2d - 6bcd^2 - bd^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{2(c-d)(c^3 + 3c^2d + 3cd^2 + d^3)} + \frac{2(9ac^2d + ad^3 - 3bc^3 - 7bcd^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3(c^2 + 2cd + d^2)(c^2 - 2cd + d^2)} - \frac{(6ac^2d - b^2d^2)}{(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d)^3} \right) \frac{f}{f}$
default	$2 \left(-\frac{(6ac^2d + 3acd^2 + 2ad^3 - 2bc^3 - 2bc^2d - 6bcd^2 - bd^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{2(c-d)(c^3 + 3c^2d + 3cd^2 + d^3)} + \frac{2(9ac^2d + ad^3 - 3bc^3 - 7bcd^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{3(c^2 + 2cd + d^2)(c^2 - 2cd + d^2)} - \frac{(6ac^2d - b^2d^2)}{(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d)^3} \right) \frac{f}{f}$
risch	Expression too large to display

input `int(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^4,x,method=_RETURNVERBOSE)`

output `1/f*(-2*(-1/2*(6*a*c^2*d+3*a*c*d^2+2*a*d^3-2*b*c^3-2*b*c^2*d-6*b*c*d^2-b*d^3)/(c-d)/(c^3+3*c^2*d+3*c*d^2+d^3)*tan(1/2*f*x+1/2*e)^5+2/3*(9*a*c^2*d+a*d^3-3*b*c^3-7*b*c*d^2)/(c^2+2*c*d+d^2)/(c^2-2*c*d+d^2)*tan(1/2*f*x+1/2*e)^3-1/2*(6*a*c^2*d-3*a*c*d^2+2*a*d^3-2*b*c^3+2*b*c^2*d-6*b*c*d^2+b*d^3)/(c+d)/(c^3-3*c^2*d+3*c*d^2-d^3)*tan(1/2*f*x+1/2*e))/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)^3+(2*a*c^3+3*a*c*d^2-4*b*c^2*d-b*d^3)/(c^6-3*c^4*d^2+3*c^2*d^4-d^6)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2)))`

3.251. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$

3.251.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 590 vs. $2(222) = 444$.

Time = 0.36 (sec) , antiderivative size = 1238, normalized size of antiderivative = 5.22

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^4,x, algorithm="fricas")`

output `[1/12*(3*(2*a*c^3*d^3 - 4*b*c^2*d^4 + 3*a*c*d^5 - b*d^6 + (2*a*c^6 - 4*b*c^5*d + 3*a*c^4*d^2 - b*c^3*d^3)*cos(f*x + e)^3 + 3*(2*a*c^5*d - 4*b*c^4*d^2 + 3*a*c^3*d^3 - b*c^2*d^4)*cos(f*x + e)^2 + 3*(2*a*c^4*d^2 - 4*b*c^3*d^3 + 3*a*c^2*d^4 - b*c*d^5)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*(2*b*c^5*d^2 - 11*a*c^4*d^3 + 11*b*c^3*d^4 + 7*a*c^2*d^5 - 13*b*c*d^6 + 4*a*d^7 + (6*b*c^7 - 18*a*c^6*d + 4*b*c^5*d^2 + 23*a*c^4*d^3 - 11*b*c^3*d^4 - 7*a*c^2*d^5 + b*c*d^6 + 2*a*d^7)*cos(f*x + e)^2 + 3*(2*b*c^6*d - 9*a*c^5*d^2 + 7*b*c^4*d^3 + 8*a*c^3*d^4 - 10*b*c^2*d^5 + a*c*d^6 + b*d^7)*cos(f*x + e))*sin(f*x + e))/((c^11 - 4*c^9*d^2 + 6*c^7*d^4 - 4*c^5*d^6 + c^3*d^8)*f*cos(f*x + e)^3 + 3*(c^10*d - 4*c^8*d^3 + 6*c^6*d^5 - 4*c^4*d^7 + c^2*d^9)*f*cos(f*x + e)^2 + 3*(c^9*d^2 - 4*c^7*d^4 + 6*c^5*d^6 - 4*c^3*d^8 + c*d^10)*f*cos(f*x + e) + (c^8*d^3 - 4*c^6*d^5 + 6*c^4*d^7 - 4*c^2*d^9 + d^11)*f), 1/6*(3*(2*a*c^3*d^3 - 4*b*c^2*d^4 + 3*a*c*d^5 - b*d^6 + (2*a*c^6 - 4*b*c^5*d + 3*a*c^4*d^2 - b*c^3*d^3)*cos(f*x + e)^3 + 3*(2*a*c^5*d - 4*b*c^4*d^2 + 3*a*c^3*d^3 - b*c^2*d^4)*cos(f*x + e)^2 + 3*(2*a*c^4*d^2 - 4*b*c^3*d^3 + 3*a*c^2*d^4 - b*c*d^5)*cos(f*x + e))*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (2*b*c^5*d^2 - 11*a*c^4*d^3 + 11*b*c^3*d^4 + 7*a*c^2*d^5 - 13*b*c*d^6 + ...`

3.251.6 Sympy [F]

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx = \int \frac{(a+b\sec(e+fx))\sec(e+fx)}{(c+d\sec(e+fx))^4} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))**4,x)`

3.251. $\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$

output `Integral((a + b*sec(e + f*x))*sec(e + f*x)/(c + d*sec(e + f*x))**4, x)`

3.251.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{(c + d \sec(e + fx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see `assume?` f or more de`

3.251.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 693 vs. 2(222) = 444.

Time = 0.40 (sec) , antiderivative size = 693, normalized size of antiderivative = 2.92

$$\int \frac{\sec(e + fx)(a + b \sec(e + fx))}{(c + d \sec(e + fx))^4} dx = \frac{3(2ac^3 - 4bc^2d + 3acd^2 - bd^3) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2c-2d) + \arctan \left(\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right)}{(c^6 - 3c^4d^2 + 3c^2d^4 - d^6)\sqrt{-c^2+d^2}} + \frac{6bc^5 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 18ac^4d}{\dots}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^4,x, algorithm="giac")`

output

```

-1/3*(3*(2*a*c^3 - 4*b*c^2*d + 3*a*c*d^2 - b*d^3)*(pi*floor(1/2*(f*x + e)/
pi + 1/2)*sgn(2*c - 2*d) + arctan((c*tan(1/2*f*x + 1/2*e) - d*tan(1/2*f*x
+ 1/2*e))/sqrt(-c^2 + d^2)))/((c^6 - 3*c^4*d^2 + 3*c^2*d^4 - d^6)*sqrt(-c^
2 + d^2)) + (6*b*c^5*tan(1/2*f*x + 1/2*e)^5 - 18*a*c^4*d*tan(1/2*f*x + 1/2
*e)^5 - 6*b*c^4*d*tan(1/2*f*x + 1/2*e)^5 + 27*a*c^3*d^2*tan(1/2*f*x + 1/2*
e)^5 + 12*b*c^3*d^2*tan(1/2*f*x + 1/2*e)^5 - 6*a*c^2*d^3*tan(1/2*f*x + 1/2
*e)^5 - 27*b*c^2*d^3*tan(1/2*f*x + 1/2*e)^5 + 3*a*c*d^4*tan(1/2*f*x + 1/2*
e)^5 + 12*b*c*d^4*tan(1/2*f*x + 1/2*e)^5 - 6*a*d^5*tan(1/2*f*x + 1/2*e)^5
+ 3*b*d^5*tan(1/2*f*x + 1/2*e)^5 - 12*b*c^5*tan(1/2*f*x + 1/2*e)^3 + 36*a*
c^4*d*tan(1/2*f*x + 1/2*e)^3 - 16*b*c^3*d^2*tan(1/2*f*x + 1/2*e)^3 - 32*a*
c^2*d^3*tan(1/2*f*x + 1/2*e)^3 + 28*b*c*d^4*tan(1/2*f*x + 1/2*e)^3 - 4*a*d
^5*tan(1/2*f*x + 1/2*e)^3 + 6*b*c^5*tan(1/2*f*x + 1/2*e) - 18*a*c^4*d*tan(
1/2*f*x + 1/2*e) + 6*b*c^4*d*tan(1/2*f*x + 1/2*e) - 27*a*c^3*d^2*tan(1/2*f
*x + 1/2*e) + 12*b*c^3*d^2*tan(1/2*f*x + 1/2*e) - 6*a*c^2*d^3*tan(1/2*f*x
+ 1/2*e) + 27*b*c^2*d^3*tan(1/2*f*x + 1/2*e) - 3*a*c*d^4*tan(1/2*f*x + 1/2
*e) + 12*b*c*d^4*tan(1/2*f*x + 1/2*e) - 6*a*d^5*tan(1/2*f*x + 1/2*e) - 3*b
*d^5*tan(1/2*f*x + 1/2*e))/((c^6 - 3*c^4*d^2 + 3*c^2*d^4 - d^6)*(c*tan(1/2
*f*x + 1/2*e)^2 - d*tan(1/2*f*x + 1/2*e)^2 - c - d)^3))/f

```

3.251.9 Mupad [B] (verification not implemented)

Time = 18.53 (sec) , antiderivative size = 439, normalized size of antiderivative = 1.85

$$\int \frac{\sec(e+fx)(a+b\sec(e+fx))}{(c+d\sec(e+fx))^4} dx$$

$$= \frac{\frac{\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^5 (2bc^3-2ad^3+bd^3-3acd^2-6ac^2d+6bcd^2+2bc^2d)}{(c+d)^3(c-d)} + \frac{4\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^3 (-3bc^3+9ac^2d-7bcd^2+ad^3)}{3(c+d)^2(c^2-2cd+d^2)}}{f \left(\tan\left(\frac{e}{2}+\frac{fx}{2}\right)^2 (-3c^3-3c^2d+3cd^2+3d^3) - \tan\left(\frac{e}{2}+\frac{fx}{2}\right)^4 (-3c^3+3c^2d+3cd^2-3d^3) + 3cd^2 \right)} + \frac{\operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2}+\frac{fx}{2}\right)(2c-2d)(c^3-3c^2d+3cd^2-d^3)}{2\sqrt{c+d}(c-d)^{7/2}}\right) (2ac^3-4bc^2d+3acd^2-bd^3)}{f(c+d)^{7/2}(c-d)^{7/2}}$$

input `int((a + b/cos(e + f*x))/(cos(e + f*x)*(c + d/cos(e + f*x))^4),x)`

output

$$\begin{aligned} & ((\tan(e/2 + (f*x)/2)^5*(2*b*c^3 - 2*a*d^3 + b*d^3 - 3*a*c*d^2 - 6*a*c^2*d \\ & + 6*b*c*d^2 + 2*b*c^2*d))/((c + d)^3*(c - d)) + (4*\tan(e/2 + (f*x)/2)^3*(a \\ & *d^3 - 3*b*c^3 + 9*a*c^2*d - 7*b*c*d^2))/(3*(c + d)^2*(c^2 - 2*c*d + d^2)) \\ & - (\tan(e/2 + (f*x)/2)*(2*a*d^3 - 2*b*c^3 + b*d^3 - 3*a*c*d^2 + 6*a*c^2*d \\ & - 6*b*c*d^2 + 2*b*c^2*d))/((c + d)*(3*c*d^2 - 3*c^2*d + c^3 - d^3)))/(f*(\tan \\ & (e/2 + (f*x)/2)^2*(3*c*d^2 - 3*c^2*d - 3*c^3 + 3*d^3) - \tan(e/2 + (f*x)/ \\ & 2)^4*(3*c*d^2 + 3*c^2*d - 3*c^3 - 3*d^3) + 3*c*d^2 + 3*c^2*d + c^3 + d^3 - \\ & \tan(e/2 + (f*x)/2)^6*(3*c*d^2 - 3*c^2*d + c^3 - d^3))) + (\operatorname{atanh}((\tan(e/2 \\ & + (f*x)/2)*(2*c - 2*d)*(3*c*d^2 - 3*c^2*d + c^3 - d^3)))/(2*(c + d)^{(1/2)}*(\\ & c - d)^{(7/2)})))*(2*a*c^3 - b*d^3 + 3*a*c*d^2 - 4*b*c^2*d))/(f*(c + d)^{(7/2)} \\ & *(c - d)^{(7/2)}) \end{aligned}$$

3.252
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+b \sec(e+fx)} dx$$

3.252.1 Optimal result 1847
 3.252.2 Mathematica [B] (verified) 1848
 3.252.3 Rubi [A] (verified) 1848
 3.252.4 Maple [B] (verified) 1850
 3.252.5 Fricas [B] (verification not implemented) 1851
 3.252.6 Sympy [F] 1852
 3.252.7 Maxima [F(-2)] 1853
 3.252.8 Giac [B] (verification not implemented) 1853
 3.252.9 Mupad [B] (verification not implemented) 1854

3.252.1 Optimal result

Integrand size = 31, antiderivative size = 247

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{a+b \sec(e+fx)} dx \\ &= \frac{d^3(4bc-ad)\operatorname{arctanh}(\sin(e+fx))}{2b^2f} \\ & \quad + \frac{d(2bc-ad)(2b^2c^2-2abcd+a^2d^2)\operatorname{arctanh}(\sin(e+fx))}{b^4f} \\ & \quad + \frac{2(bc-ad)^4\operatorname{arctanh}\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b}b^4\sqrt{a+bf}} \\ & \quad + \frac{d^4 \tan(e+fx)}{bf} + \frac{d^2(6b^2c^2-4abcd+a^2d^2)\tan(e+fx)}{b^3f} \\ & \quad + \frac{d^3(4bc-ad)\sec(e+fx)\tan(e+fx)}{2b^2f} + \frac{d^4 \tan^3(e+fx)}{3bf} \end{aligned}$$

```
output 1/2*d^3*(-a*d+4*b*c)*arctanh(sin(f*x+e))/b^2/f+d*(-a*d+2*b*c)*(a^2*d^2-2*a
*b*c*d+2*b^2*c^2)*arctanh(sin(f*x+e))/b^4/f+2*(-a*d+b*c)^4*arctanh((a-b)^(
1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/b^4/f/(a-b)^(1/2)/(a+b)^(1/2)+d^4*tan
(f*x+e)/b/f+d^2*(a^2*d^2-4*a*b*c*d+6*b^2*c^2)*tan(f*x+e)/b^3/f+1/2*d^3*(-a
*d+4*b*c)*sec(f*x+e)*tan(f*x+e)/b^2/f+1/3*d^4*tan(f*x+e)^3/b/f
```


3.252.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 580 vs. $2(247) = 494$.

Time = 5.68 (sec) , antiderivative size = 580, normalized size of antiderivative = 2.35

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+b\sec(e+fx)} dx$$

$$= \frac{\cos^3(e+fx)(b+a\cos(e+fx))(c+d\sec(e+fx))^4 \left(-\frac{24(bc-ad)^4 \operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}} \right) - 6d(8a^2bcd^2}{\dots}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + b*Sec[e + f*x]),x]`

output `(Cos[e + f*x]^3*(b + a*Cos[e + f*x])*(c + d*Sec[e + f*x])^4*((-24*(b*c - a*d)^4*ArcTanh[(-a + b)*Tan[(e + f*x)/2]]/Sqrt[a^2 - b^2])/Sqrt[a^2 - b^2] - 6*d*(8*a^2*b*c*d^2 - 2*a^3*d^3 + 4*b^3*c*(2*c^2 + d^2) - a*b^2*d*(12*c^2 + d^2))*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] - 6*d*(-8*a^2*b*c*d^2 + 2*a^3*d^3 - 4*b^3*c*(2*c^2 + d^2) + a*b^2*d*(12*c^2 + d^2))*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] + (b^2*d^3*(-3*a*d + b*(12*c + d)))/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2 + (2*b^3*d^4*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^3 + (4*b*d^2*(-12*a*b*c*d + 3*a^2*d^2 + 2*b^2*(9*c^2 + d^2))*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]) + (2*b^3*d^4*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])^3 - (b^2*d^3*(-3*a*d + b*(12*c + d)))/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])^2 + (4*b*d^2*(-12*a*b*c*d + 3*a^2*d^2 + 2*b^2*(9*c^2 + d^2))*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2]))/(12*b^4*f*(d + c*Cos[e + f*x])^4*(a + b*Sec[e + f*x]))`

3.252.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4476, 3042, 3431, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.252. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+b\sec(e+fx)} dx$

$$\begin{aligned}
& \int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+b\sec(e+fx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))^4}{a+b\csc(e+fx+\frac{\pi}{2})} dx \\
& \quad \downarrow \text{4476} \\
& \int \frac{\sec^4(e+fx)(c\cos(e+fx)+d)^4}{a\cos(e+fx)+b} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(c\sin(e+fx+\frac{\pi}{2})+d)^4}{\sin(e+fx+\frac{\pi}{2})^4(a\sin(e+fx+\frac{\pi}{2})+b)} dx \\
& \quad \downarrow \text{3431} \\
& \int \left(\frac{d(2bc-ad)(a^2d^2-2abcd+2b^2c^2)\sec(e+fx)}{b^4} + \frac{d^2(a^2d^2-4abcd+6b^2c^2)\sec^2(e+fx)}{b^3} + \frac{(bc-ad)^4}{b^4(a\cos(e+fx))} \right) dx \\
& \quad \downarrow \text{2009} \\
& \frac{d(2bc-ad)(a^2d^2-2abcd+2b^2c^2)\operatorname{arctanh}(\sin(e+fx))}{b^4f} + \frac{d^2(a^2d^2-4abcd+6b^2c^2)\tan(e+fx)}{b^3f} + \\
& \quad \frac{2(bc-ad)^4\operatorname{arctanh}\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{b^4f\sqrt{a-b}\sqrt{a+b}} + \frac{d^3(4bc-ad)\operatorname{arctanh}(\sin(e+fx))}{2b^2f} + \\
& \quad \frac{d^3(4bc-ad)\tan(e+fx)\sec(e+fx)}{2b^2f} + \frac{d^4\tan^3(e+fx)}{3bf} + \frac{d^4\tan(e+fx)}{bf}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + b*Sec[e + f*x]),x]`

output `(d^3*(4*b*c - a*d)*ArcTanh[Sin[e + f*x]]/(2*b^2*f) + (d*(2*b*c - a*d)*(2*b^2*c^2 - 2*a*b*c*d + a^2*d^2)*ArcTanh[Sin[e + f*x]])/(b^4*f) + (2*(b*c - a*d)^4*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b^4*Sqrt[a + b]*f) + (d^4*Tan[e + f*x])/(b*f) + (d^2*(6*b^2*c^2 - 4*a*b*c*d + a^2*d^2)*Tan[e + f*x])/(b^3*f) + (d^3*(4*b*c - a*d)*Sec[e + f*x]*Tan[e + f*x])/(2*b^2*f) + (d^4*Tan[e + f*x]^3)/(3*b*f)`

3.252.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3431 Int[((g_.)*sin[(e_.) + (f_.)*(x_)])^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(g*sin[e + f*x])^p*(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && NeQ[b*c - a*d, 0] && (IntegersQ[m, n] || IntegersQ[m, p] || IntegersQ[n, p]) && NeQ[p, 2]
```

```
rule 4476 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_)*(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.)^(n_), x_Symbol] := Simp[1/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.252.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 479 vs. 2(232) = 464.

Time = 1.44 (sec) , antiderivative size = 480, normalized size of antiderivative = 1.94

method	result
derivativedivides	$-\frac{2(-a^4d^4+4a^3bcd^3-6a^2b^2c^2d^2+4ab^3c^3d-b^4c^4) \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^4\sqrt{(a-b)(a+b)}} - \frac{d^4}{3b\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)^3} - \frac{d(2a^3d^3-8a^2bcd^2+...)}{...}$
default	$-\frac{2(-a^4d^4+4a^3bcd^3-6a^2b^2c^2d^2+4ab^3c^3d-b^4c^4) \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^4\sqrt{(a-b)(a+b)}} - \frac{d^4}{3b\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)+1\right)^3} - \frac{d(2a^3d^3-8a^2bcd^2+...)}{...}$
risch	Expression too large to display

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

3.252.
$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+b\sec(e+fx)} dx$$

output
$$\frac{1}{f} \frac{(-2/b^4(-a^4d^4+4a^3b^3cd^3-6a^2b^2c^2d^2+4ab^3c^3d-b^4c^4)/((a-b)(a+b))^{1/2} \operatorname{arctanh}((a-b)\tan(1/2fx+1/2e))/((a-b)(a+b))^{1/2}) - 1/3d^4/b/(\tan(1/2fx+1/2e)+1)^3 - 1/2d(2a^3d^3-8a^2b^3cd^2+12ab^2c^2d+a^2b^2d^3-8b^3c^3-4b^3cd^2)/b^4 \ln(\tan(1/2fx+1/2e)+1) - 1/2d^2(2a^2d^2-8ab^2cd+a^2bd^2+12b^2c^2-4b^2cd+2b^2d^2)/b^3/(\tan(1/2fx+1/2e)+1) + 1/2d^3(ad-4bc+bd)/b^2/(\tan(1/2fx+1/2e)+1)^2 - 1/3d^4/b/(\tan(1/2fx+1/2e)-1)^3 + 1/2d(2a^3d^3-8a^2b^3cd^2+12ab^2c^2d+a^2b^2d^3-8b^3c^3-4b^3cd^2)/b^4 \ln(\tan(1/2fx+1/2e)-1) - 1/2d^2(2a^2d^2-8ab^2cd+a^2bd^2+12b^2c^2-4b^2cd+2b^2d^2)/b^3/(\tan(1/2fx+1/2e)-1) - 1/2d^3(ad-4bc+bd)/b^2/(\tan(1/2fx+1/2e)-1)^2}$$

3.252.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 518 vs. $2(232) = 464$.

Time = 102.85 (sec) , antiderivative size = 1093, normalized size of antiderivative = 4.43

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+b\sec(e+fx)} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e)),x, algorithm="fricas")`

output

```
[1/12*(6*(b^4*c^4 - 4*a*b^3*c^3*d + 6*a^2*b^2*c^2*d^2 - 4*a^3*b*c*d^3 + a^4*d^4)*sqrt(a^2 - b^2)*cos(f*x + e)^3*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 + 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)) + 3*(8*(a^2*b^3 - b^5)*c^3*d - 12*(a^3*b^2 - a*b^4)*c^2*d^2 + 4*(2*a^4*b - a^2*b^3 - b^5)*c*d^3 - (2*a^5 - a^3*b^2 - a*b^4)*d^4)*cos(f*x + e)^3*log(sin(f*x + e) + 1) - 3*(8*(a^2*b^3 - b^5)*c^3*d - 12*(a^3*b^2 - a*b^4)*c^2*d^2 + 4*(2*a^4*b - a^2*b^3 - b^5)*c*d^3 - (2*a^5 - a^3*b^2 - a*b^4)*d^4)*cos(f*x + e)^3*log(-sin(f*x + e) + 1) + 2*(2*(a^2*b^3 - b^5)*d^4 + 2*(18*(a^2*b^3 - b^5)*c^2*d^2 - 12*(a^3*b^2 - a*b^4)*c*d^3 + (3*a^4*b - a^2*b^3 - 2*b^5)*d^4)*cos(f*x + e)^2 + 3*(4*(a^2*b^3 - b^5)*c*d^3 - (a^3*b^2 - a*b^4)*d^4)*cos(f*x + e))*sin(f*x + e))/((a^2*b^4 - b^6)*f*cos(f*x + e)^3), 1/12*(12*(b^4*c^4 - 4*a*b^3*c^3*d + 6*a^2*b^2*c^2*d^2 - 4*a^3*b*c*d^3 + a^4*d^4)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e)))*cos(f*x + e)^3 + 3*(8*(a^2*b^3 - b^5)*c^3*d - 12*(a^3*b^2 - a*b^4)*c^2*d^2 + 4*(2*a^4*b - a^2*b^3 - b^5)*c*d^3 - (2*a^5 - a^3*b^2 - a*b^4)*d^4)*cos(f*x + e)^3*log(sin(f*x + e) + 1) - 3*(8*(a^2*b^3 - b^5)*c^3*d - 12*(a^3*b^2 - a*b^4)*c^2*d^2 + 4*(2*a^4*b - a^2*b^3 - b^5)*c*d^3 - (2*a^5 - a^3*b^2 - a*b^4)*d^4)*cos(f*x + e)^3*log(-sin(f*x + e) + 1) + 2*(2*(a^2*b^3 - b^5)*d^4 + 2*(18*(a^2*b^3 - b^5)*c^2*d^2 - 12*(a^3*b^2 - a*b^4)*c...
```

3.252.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+b\sec(e+fx)} dx = \int \frac{(c+d\sec(e+fx))^4 \sec(e+fx)}{a+b\sec(e+fx)} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**4/(a+b*sec(f*x+e)),x)`

output `Integral((c + d*sec(e + f*x))**4*sec(e + f*x)/(a + b*sec(e + f*x)), x)`

3.252.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{a + b \sec(e + fx)} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e)),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de
```

3.252.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 606 vs. 2(232) = 464.

Time = 0.40 (sec) , antiderivative size = 606, normalized size of antiderivative = 2.45

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{a + b \sec(e + fx)} dx$$

$$= \frac{3(8b^3c^3d - 12ab^2c^2d^2 + 8a^2bcd^3 + 4b^3cd^3 - 2a^3d^4 - ab^2d^4) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{b^4} - \frac{3(8b^3c^3d - 12ab^2c^2d^2 + 8a^2bcd^3 + 4b^3cd^3 - 2a^3d^4 - ab^2d^4)}{b^4}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e)),x, algorithm="giac")
```

output $\frac{1}{6} \cdot (3 \cdot (8b^3c^3d - 12ab^2c^2d^2 + 8a^2b^2cd^3 + 4b^3cd^3 - 2a^3d^4 - ab^2d^4) \cdot \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)) / b^4 - 3 \cdot (8b^3c^3d - 12ab^2c^2d^2 + 8a^2b^2cd^3 + 4b^3cd^3 - 2a^3d^4 - ab^2d^4) \cdot \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)) / b^4 - 12 \cdot (b^4c^4 - 4ab^3c^3d + 6a^2b^2c^2d^2 - 4a^3b^2cd^3 + a^4d^4) \cdot (\pi \cdot \text{floor}(\frac{1}{2}(fx + e)) / \pi + \frac{1}{2}) \cdot \text{sgn}(2a - 2b) + \arctan((a \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e) - b \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)) / \sqrt{-a^2 + b^2})) / (\sqrt{-a^2 + b^2} \cdot b^4) - 2 \cdot (36b^2c^2d^2 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 24ab^2cd^3 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 12b^2c^2d^3 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 6a^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 3ab^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 + 6b^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 72b^2c^2d^2 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + 48ab^2cd^3 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 12a^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 4b^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + 36b^2c^2d^2 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e) - 24ab^2cd^3 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e) + 12b^2c^2d^3 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e) + 6a^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e) - 3ab^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e) + 6b^2d^4 \cdot \tan(\frac{1}{2}fx + \frac{1}{2}e)) / ((\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)^3 \cdot b^3)) / f$

3.252.9 Mupad [B] (verification not implemented)

Time = 23.13 (sec) , antiderivative size = 9987, normalized size of antiderivative = 40.43

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{a+b\sec(e+fx)} dx = \text{Too large to display}$$

input `int((c + d/cos(e + f*x))^4/(cos(e + f*x)*(a + b/cos(e + f*x))),x)`

output $(\operatorname{atan}(\frac{(((((8(4b^{13}c^4 - 8ab^{12}c^4 - 2a^2b^{12}d^4 + 8b^{13}c^3d + 16b^{13}c^3d + 4a^2b^{11}c^4 + 2a^2b^{11}d^4 - 2a^3b^{10}d^4 + 6a^4b^9d^4 - 4a^5b^8d^4 - 24a^2b^{12}c^2d^2 + 8a^2b^{11}c^3d + 16a^2b^{11}c^3d - 24a^3b^{10}c^2d^3 + 16a^4b^9c^2d^3 + 48a^2b^{11}c^2d^2 - 24a^3b^{10}c^2d^2 - 8a^2b^{12}c^3d - 32a^2b^{12}c^3d)))/b^9 - (8\tan(e/2 + (fx)/2)*(8a^2b^{10} - 16a^2b^9 + 8a^3b^8)*(b^2((a^4d^4)/2 + 6a^2c^2d^2) - b^3(2c^2d^3 + 4c^3d) + a^3d^4 - 4a^2b^2c^3d))/b^{10}*(b^2((a^4d^4)/2 + 6a^2c^2d^2) - b^3(2c^2d^3 + 4c^3d) + a^3d^4 - 4a^2b^2c^3d))/b^4 + (8\tan(e/2 + (fx)/2)*(8a^9d^8 - 4b^9c^8 + 4ab^8c^8 - 16a^8b^8d^8 - a^2b^7d^8 + 3a^3b^6d^8 - 7a^4b^5d^8 + 13a^5b^4d^8 - 16a^6b^3d^8 + 16a^7b^2d^8 - 16b^9c^2d^6 - 64b^9c^4d^4 - 64b^9c^6d^2 + 48ab^8c^2d^6 + 112ab^8c^3d^5 + 192ab^8c^4d^4 + 192ab^8c^5d^3 + 192ab^8c^6d^2 - 24a^2b^7c^2d^7 - 32a^2b^7c^7d + 56a^3b^6c^2d^7 - 104a^4b^5c^2d^7 + 128a^5b^4c^2d^7 - 128a^6b^3c^2d^7 + 128a^7b^2c^2d^7 - 136a^2b^7c^2d^6 - 336a^2b^7c^3d^5 - 464a^2b^7c^4d^4 - 576a^2b^7c^5d^3 - 304a^2b^7c^6d^2 + 280a^3b^6c^2d^6 + 560a^3b^6c^3d^5 + 880a^3b^6c^4d^4 + 800a^3b^6c^5d^3 + 176a^3b^6c^6d^2 - 376a^4b^5c^2d^6 - 784a^4b^5c^3d^5 - 1096a^4b^5c^4d^4 - 416a^4b^5c^5d^3 + 424a^5b^4c^2d^6 + 896a^5b^4c^3d^5 + 552a^5b^4c^4d^4 - 448a^6b^3c^2d^6 - 448a^6b^3c^3d^5 + 224\dots$

3.253 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+b \sec(e+fx)} dx$

3.253.1 Optimal result 1856
 3.253.2 Mathematica [B] (verified) 1857
 3.253.3 Rubi [A] (verified) 1857
 3.253.4 Maple [A] (verified) 1859
 3.253.5 Fricas [B] (verification not implemented) 1860
 3.253.6 Sympy [F] 1860
 3.253.7 Maxima [F(-2)] 1861
 3.253.8 Giac [B] (verification not implemented) 1861
 3.253.9 Mupad [B] (verification not implemented) 1862

3.253.1 Optimal result

Integrand size = 31, antiderivative size = 170

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{a+b \sec(e+fx)} dx = \frac{d^3 \operatorname{arctanh}(\sin(e+fx))}{2bf} + \frac{d(3b^2c^2 - 3abcd + a^2d^2) \operatorname{arctanh}(\sin(e+fx))}{b^3f} + \frac{2(bc - ad)^3 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{\sqrt{a-b} b^3 \sqrt{a+b} f} + \frac{d^2(3bc - ad) \tan(e+fx)}{b^2f} + \frac{d^3 \sec(e+fx) \tan(e+fx)}{2bf}$$

```
output 1/2*d^3*arctanh(sin(f*x+e))/b/f+d*(a^2*d^2-3*a*b*c*d+3*b^2*c^2)*arctanh(sin(f*x+e))/b^3/f+2*(-a*d+b*c)^3*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/b^3/f/(a-b)^(1/2)/(a+b)^(1/2)+d^2*(-a*d+3*b*c)*tan(f*x+e)/b^2/f+1/2*d^3*sec(f*x+e)*tan(f*x+e)/b/f
```

3.253.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 389 vs. $2(170) = 340$.

Time = 3.11 (sec) , antiderivative size = 389, normalized size of antiderivative = 2.29

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+b\sec(e+fx)} dx$$

$$= \frac{\cos^2(e+fx)(b+a\cos(e+fx))(c+d\sec(e+fx))^3 \left(\frac{8(-bc+ad)^3 \operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}} - 2d(-6abcd + \dots) \right)}{\dots}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + b*Sec[e + f*x]),x]`

output `(Cos[e + f*x]^2*(b + a*Cos[e + f*x])*(c + d*Sec[e + f*x])^3*((8*(-(b*c) + a*d)^3*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]])/Sqrt[a^2 - b^2] - 2*d*(-6*a*b*c*d + 2*a^2*d^2 + b^2*(6*c^2 + d^2))*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + 2*d*(-6*a*b*c*d + 2*a^2*d^2 + b^2*(6*c^2 + d^2))*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] + (b^2*d^3)/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2 + (4*b*d^2*(3*b*c - a*d)*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]) - (b^2*d^3)/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])^2 + (4*b*d^2*(3*b*c - a*d)*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2]))/(4*b^3*f*(d + c*Cos[e + f*x])^3*(a + b*Sec[e + f*x]))`

3.253.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4476, 3042, 3431, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+b\sec(e+fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^3}{a+b\csc\left(e+fx+\frac{\pi}{2}\right)} dx$$

$$\begin{aligned}
& \downarrow 4476 \\
& \int \frac{\sec^3(e+fx)(c \cos(e+fx) + d)^3}{a \cos(e+fx) + b} dx \\
& \downarrow 3042 \\
& \int \frac{(c \sin(e+fx + \frac{\pi}{2}) + d)^3}{\sin(e+fx + \frac{\pi}{2})^3 (a \sin(e+fx + \frac{\pi}{2}) + b)} dx \\
& \downarrow 3431 \\
& \int \left(\frac{d(a^2 d^2 - 3abcd + 3b^2 c^2) \sec(e+fx)}{b^3} + \frac{(bc - ad)^3}{b^3 (a \cos(e+fx) + b)} + \frac{d^2 (3bc - ad) \sec^2(e+fx)}{b^2} + \frac{d^3 \sec^3(e+fx)}{b} \right) dx \\
& \downarrow 2009 \\
& \frac{d(a^2 d^2 - 3abcd + 3b^2 c^2) \operatorname{arctanh}(\sin(e+fx))}{b^3 f} + \frac{2(bc - ad)^3 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{b^3 f \sqrt{a-b} \sqrt{a+b}} + \\
& \frac{d^2 (3bc - ad) \tan(e+fx)}{b^2 f} + \frac{d^3 \operatorname{arctanh}(\sin(e+fx))}{2bf} + \frac{d^3 \tan(e+fx) \sec(e+fx)}{2bf}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + b*Sec[e + f*x]),x]`

output `(d^3*ArcTanh[Sin[e + f*x]])/(2*b*f) + (d*(3*b^2*c^2 - 3*a*b*c*d + a^2*d^2)*ArcTanh[Sin[e + f*x]])/(b^3*f) + (2*(b*c - a*d)^3*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b^3*Sqrt[a + b]*f) + (d^2*(3*b*c - a*d)*Tan[e + f*x])/(b^2*f) + (d^3*Sec[e + f*x]*Tan[e + f*x])/(2*b*f)`

3.253.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3431 Int[((g_.)*sin[(e_.) + (f_.)*(x_)])^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(g*sin[e + f*x])^p*(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && NeQ[b*c - a*d, 0] && (IntegersQ[m, n] || IntegersQ[m, p] || IntegersQ[n, p]) && NeQ[p, 2]
```

```
rule 4476 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_)^(m_)*(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_)^(n_), x_Symbol] := Simp[1/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.253.4 Maple [A] (verified)

Time = 1.04 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.70

method	result
derivativedivides	$-\frac{d^3}{2b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{d(2a^2d^2 - 6abcd + 6b^2c^2 + b^2d^2) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2b^3} + \frac{d^2(2ad - 6bc + bd)}{2b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} + \frac{d^3}{2b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^2} - \frac{d^2(2ad - 6bc + bd)}{2b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}$
default	$-\frac{d^3}{2b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{d(2a^2d^2 - 6abcd + 6b^2c^2 + b^2d^2) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2b^3} + \frac{d^2(2ad - 6bc + bd)}{2b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} + \frac{d^3}{2b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)^2} - \frac{d^2(2ad - 6bc + bd)}{2b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}$
risch	$-\frac{id^2(bde^{3i(fx+e)} + 2ade^{2i(fx+e)} - 6bce^{2i(fx+e)} - bde^{i(fx+e)} + 2ad - 6bc)}{fb^2(1+e^{2i(fx+e)})^2} + \frac{\ln\left(e^{i(fx+e)} - \frac{ia^2 - ib^2 - b\sqrt{a^2 - b^2}}{\sqrt{a^2 - b^2}a}\right)a^3d^3}{\sqrt{a^2 - b^2}fb^3}$

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f*(-1/2*d^3/b/(tan(1/2*f*x+1/2*e)+1)^2+1/2*d*(2*a^2*d^2-6*a*b*c*d+6*b^2*c^2+b^2*d^2)/b^3*ln(tan(1/2*f*x+1/2*e)+1)+1/2*d^2*(2*a*d-6*b*c+b*d)/b^2/(tan(1/2*f*x+1/2*e)+1)+1/2*d^3/b/(tan(1/2*f*x+1/2*e)-1)^2-1/2*d*(2*a^2*d^2-6*a*b*c*d+6*b^2*c^2+b^2*d^2)/b^3*ln(tan(1/2*f*x+1/2*e)-1)+1/2*d^2*(2*a*d-6*b*c+b*d)/b^2/(tan(1/2*f*x+1/2*e)-1)-2*(a^3*d^3-3*a^2*b*c*d^2+3*a*b^2*c^2*d-b^3*c^3)/b^3/((a-b)*(a+b))^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e)/((a-b)*(a+b))^(1/2)))
```

$$3.253. \int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+b\sec(e+fx)} dx$$

3.253.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 361 vs. $2(157) = 314$.

Time = 22.06 (sec) , antiderivative size = 779, normalized size of antiderivative = 4.58

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+b\sec(e+fx)} dx$$

$$= \left[-\frac{2(b^3c^3 - 3ab^2c^2d + 3a^2bcd^2 - a^3d^3)\sqrt{a^2 - b^2} \cos(fx+e)^2 \log\left(\frac{2ab\cos(fx+e) - (a^2 - 2b^2)\cos(fx+e)^2 - 2\sqrt{a^2 - b^2}}{a^2\cos(fx+e)^2 + 2ab\cos(fx+e)}\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e)),x, algorithm="fricas")`

output `[-1/4*(2*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(a^2 - b^2)*cos(f*x + e)^2*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 - 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)) - (6*(a^2*b^2 - b^4)*c^2*d - 6*(a^3*b - a*b^3)*c*d^2 + (2*a^4 - a^2*b^2 - b^4)*d^3)*cos(f*x + e)^2*log(sin(f*x + e) + 1) + (6*(a^2*b^2 - b^4)*c^2*d - 6*(a^3*b - a*b^3)*c*d^2 + (2*a^4 - a^2*b^2 - b^4)*d^3)*cos(f*x + e)^2*log(-sin(f*x + e) + 1) - 2*((a^2*b^2 - b^4)*d^3 + 2*(3*(a^2*b^2 - b^4)*c*d^2 - (a^3*b - a*b^3)*d^3)*cos(f*x + e))*sin(f*x + e))/((a^2*b^3 - b^5)*f*cos(f*x + e)^2), 1/4*(4*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e)))*cos(f*x + e)^2 + (6*(a^2*b^2 - b^4)*c^2*d - 6*(a^3*b - a*b^3)*c*d^2 + (2*a^4 - a^2*b^2 - b^4)*d^3)*cos(f*x + e)^2*log(sin(f*x + e) + 1) - (6*(a^2*b^2 - b^4)*c^2*d - 6*(a^3*b - a*b^3)*c*d^2 + (2*a^4 - a^2*b^2 - b^4)*d^3)*cos(f*x + e)^2*log(-sin(f*x + e) + 1) + 2*((a^2*b^2 - b^4)*d^3 + 2*(3*(a^2*b^2 - b^4)*c*d^2 - (a^3*b - a*b^3)*d^3)*cos(f*x + e))*sin(f*x + e))/((a^2*b^3 - b^5)*f*cos(f*x + e)^2)]`

3.253.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+b\sec(e+fx)} dx = \int \frac{(c+d\sec(e+fx))^3 \sec(e+fx)}{a+b\sec(e+fx)} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**3/(a+b*sec(f*x+e)),x)`

3.253. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+b\sec(e+fx)} dx$

output `Integral((c + d*sec(e + f*x))^3*sec(e + f*x)/(a + b*sec(e + f*x)), x)`

3.253.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(c + d\sec(e + fx))^3}{a + b\sec(e + fx)} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de`

3.253.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. $2(157) = 314$.

Time = 0.40 (sec) , antiderivative size = 339, normalized size of antiderivative = 1.99

$$\int \frac{\sec(e + fx)(c + d\sec(e + fx))^3}{a + b\sec(e + fx)} dx$$

$$= \frac{(6b^2c^2d - 6abcd^2 + 2a^2d^3 + b^2d^3) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) + 1\right|\right)}{b^3} - \frac{(6b^2c^2d - 6abcd^2 + 2a^2d^3 + b^2d^3) \log\left(\left|\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right) - 1\right|\right)}{b^3} - \frac{4(b^3c^3 - 3ab^2c^2d + \dots)}{b^3}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e)),x, algorithm="giac")`

```
output 1/2*((6*b^2*c^2*d - 6*a*b*c*d^2 + 2*a^2*d^3 + b^2*d^3)*log(abs(tan(1/2*f*x
+ 1/2*e) + 1)))/b^3 - (6*b^2*c^2*d - 6*a*b*c*d^2 + 2*a^2*d^3 + b^2*d^3)*lo
g(abs(tan(1/2*f*x + 1/2*e) - 1))/b^3 - 4*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*
b*c*d^2 - a^3*d^3)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*a - 2*b) + arct
an((a*tan(1/2*f*x + 1/2*e) - b*tan(1/2*f*x + 1/2*e))/sqrt(-a^2 + b^2)))/(s
qrt(-a^2 + b^2)*b^3) - 2*(6*b*c*d^2*tan(1/2*f*x + 1/2*e)^3 - 2*a*d^3*tan(1
/2*f*x + 1/2*e)^3 - b*d^3*tan(1/2*f*x + 1/2*e)^3 - 6*b*c*d^2*tan(1/2*f*x +
1/2*e) + 2*a*d^3*tan(1/2*f*x + 1/2*e) - b*d^3*tan(1/2*f*x + 1/2*e))/((tan
(1/2*f*x + 1/2*e)^2 - 1)^2*b^2))/f
```

3.253.9 Mupad [B] (verification not implemented)

Time = 21.04 (sec) , antiderivative size = 6730, normalized size of antiderivative = 39.59

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{a+b\sec(e+fx)} dx = \text{Too large to display}$$

```
input int((c + d/cos(e + f*x))^3/(cos(e + f*x)*(a + b/cos(e + f*x))),x)
```

```
output ((tan(e/2 + (f*x)/2)*(b*d^3 - 2*a*d^3 + 6*b*c*d^2))/b^2 + (tan(e/2 + (f*x)
/2)^3*(2*a*d^3 + b*d^3 - 6*b*c*d^2))/b^2)/(f*(tan(e/2 + (f*x)/2)^4 - 2*tan
(e/2 + (f*x)/2)^2 + 1)) - (atan((((8*tan(e/2 + (f*x)/2)*(8*a^7*d^6 - 4*b^
7*c^6 - b^7*d^6 + 4*a*b^6*c^6 + 3*a*b^6*d^6 - 16*a^6*b*d^6 - 7*a^2*b^5*d^6
+ 13*a^3*b^4*d^6 - 16*a^4*b^3*d^6 + 16*a^5*b^2*d^6 - 12*b^7*c^2*d^4 - 36*
b^7*c^4*d^2 + 36*a*b^6*c^2*d^4 + 72*a*b^6*c^3*d^3 + 108*a*b^6*c^4*d^2 - 36
*a^2*b^5*c*d^5 - 24*a^2*b^5*c^5*d + 60*a^3*b^4*c*d^5 - 84*a^4*b^3*c*d^5 +
96*a^5*b^2*c*d^5 - 96*a^2*b^5*c^2*d^4 - 216*a^2*b^5*c^3*d^3 - 168*a^2*b^5*
c^4*d^2 + 192*a^3*b^4*c^2*d^4 + 296*a^3*b^4*c^3*d^3 + 96*a^3*b^4*c^4*d^2 -
240*a^4*b^3*c^2*d^4 - 152*a^4*b^3*c^3*d^3 + 120*a^5*b^2*c^2*d^4 + 12*a*b^
6*c*d^5 + 24*a*b^6*c^5*d - 48*a^6*b*c*d^5))/b^4 + (((8*(4*b^10*c^3 + 2*b^1
0*d^3 - 8*a*b^9*c^3 - 2*a*b^9*d^3 + 12*b^10*c^2*d + 4*a^2*b^8*c^3 + 2*a^2*
b^8*d^3 - 6*a^3*b^7*d^3 + 4*a^4*b^6*d^3 + 24*a^2*b^8*c*d^2 + 12*a^2*b^8*c^
2*d - 12*a^3*b^7*c*d^2 - 12*a*b^9*c*d^2 - 24*a*b^9*c^2*d))/b^6 - (8*tan(e/
2 + (f*x)/2)*(8*a*b^8 - 16*a^2*b^7 + 8*a^3*b^6)*(b^2*(3*c^2*d + d^3/2) + a
^2*d^3 - 3*a*b*c*d^2))/b^7)*(b^2*(3*c^2*d + d^3/2) + a^2*d^3 - 3*a*b*c*d^2
))/b^3)*(b^2*(3*c^2*d + d^3/2) + a^2*d^3 - 3*a*b*c*d^2)*1i)/b^3 + (((8*tan
(e/2 + (f*x)/2)*(8*a^7*d^6 - 4*b^7*c^6 - b^7*d^6 + 4*a*b^6*c^6 + 3*a*b^6*d
^6 - 16*a^6*b*d^6 - 7*a^2*b^5*d^6 + 13*a^3*b^4*d^6 - 16*a^4*b^3*d^6 + 16*a
^5*b^2*d^6 - 12*b^7*c^2*d^4 - 36*b^7*c^4*d^2 + 36*a*b^6*c^2*d^4 + 72*a*...
```

3.254 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+b \sec(e+fx)} dx$

3.254.1 Optimal result 1863
 3.254.2 Mathematica [A] (verified) 1863
 3.254.3 Rubi [A] (verified) 1864
 3.254.4 Maple [A] (verified) 1866
 3.254.5 Fricas [B] (verification not implemented) 1866
 3.254.6 Sympy [F] 1867
 3.254.7 Maxima [F(-2)] 1867
 3.254.8 Giac [B] (verification not implemented) 1868
 3.254.9 Mupad [B] (verification not implemented) 1868

3.254.1 Optimal result

Integrand size = 31, antiderivative size = 103

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+b \sec(e+fx)} dx = \frac{d(2bc-ad)\operatorname{arctanh}(\sin(e+fx))}{b^2 f} + \frac{2(bc-ad)^2 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b} b^2 \sqrt{a+b} f} + \frac{d^2 \tan(e+fx)}{b f}$$

```
output d*(-a*d+2*b*c)*arctanh(sin(f*x+e))/b^2/f+2*(-a*d+b*c)^2*arctanh((a-b)^(1/2)
)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/b^2/f/(a-b)^(1/2)/(a+b)^(1/2)+d^2*tan(f*
x+e)/b/f
```

3.254.2 Mathematica [A] (verified)

Time = 2.51 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.31

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{a+b \sec(e+fx)} dx = \frac{2(bc-ad)^2 \operatorname{arctanh}\left(\frac{(-a+b) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}} + d \frac{-((2bc-ad) (\log(\cos(\frac{1}{2}(e+fx))) - \sin(\frac{1}{2}(e+fx)))) - \log(\cos(\frac{1}{2}(e+fx)))}{b^2 f}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + b*Sec[e + f*x]),x]`

output `((-2*(b*c - a*d)^2*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]])/Sqrt[a^2 - b^2] + d*(-((2*b*c - a*d)*(Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] - Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]])) + b*d*Tan[e + f*x]))/(b^2*f)`

3.254.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4476, 3042, 3431, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{a+b\sec(e+fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))^2}{a+b\csc(e+fx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4476} \\
 & \int \frac{\sec^2(e+fx)(c\cos(e+fx)+d)^2}{a\cos(e+fx)+b} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c\sin(e+fx+\frac{\pi}{2})+d)^2}{\sin(e+fx+\frac{\pi}{2})^2(a\sin(e+fx+\frac{\pi}{2})+b)} dx \\
 & \quad \downarrow \text{3431} \\
 & \int \left(\frac{(bc-ad)^2}{b^2(a\cos(e+fx)+b)} + \frac{d(2bc-ad)\sec(e+fx)}{b^2} + \frac{d^2\sec^2(e+fx)}{b} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{d(2bc-ad)\operatorname{arctanh}(\sin(e+fx))}{b^2f} + \frac{2(bc-ad)^2\operatorname{arctanh}\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{b^2f\sqrt{a-b}\sqrt{a+b}} + \frac{d^2\tan(e+fx)}{bf}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + b*Sec[e + f*x]),x]`

output `(d*(2*b*c - a*d)*ArcTanh[Sin[e + f*x]]/(b^2*f) + (2*(b*c - a*d)^2*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b^2*Sqrt[a + b]*f) + (d^2*Tan[e + f*x])/(b*f)`

3.254.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3431 `Int[((g_.)*sin[(e_.) + (f_.)*(x_)])^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(g*sin[e + f*x])^p*(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && NeQ[b*c - a*d, 0] && (IntegersQ[m, n] || IntegersQ[m, p] || IntegersQ[n, p]) && NeQ[p, 2]`

rule 4476 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_)^(m_)*(csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_)^(n_), x_Symbol] := Simp[1/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && IntegerQ[m] && IntegerQ[n]`

3.254.4 Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.60

method	result
derivativedivides	$\frac{\frac{d^2}{b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} - \frac{d(ad-2bc) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{b^2} - \frac{2(-a^2d^2+2abcd-b^2c^2) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^2 \sqrt{(a-b)(a+b)}}}{f} - \frac{d^2}{b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}$
default	$\frac{\frac{d^2}{b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} - \frac{d(ad-2bc) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{b^2} - \frac{2(-a^2d^2+2abcd-b^2c^2) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^2 \sqrt{(a-b)(a+b)}}}{f} - \frac{d^2}{b \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}$
risch	$\frac{2id^2}{fb(1+e^{2i(fx+e)})} + \frac{d^2 \ln(e^{i(fx+e)} - i)a}{b^2 f} - \frac{2d \ln(e^{i(fx+e)} - i)c}{bf} + \frac{\ln\left(e^{i(fx+e)} + \frac{ia^2 - ib^2 + b\sqrt{a^2 - b^2}}{\sqrt{a^2 - b^2} a}\right) a^2 d^2}{\sqrt{a^2 - b^2} f b^2} - \frac{2 \ln(e^{i(fx+e)} - i)c}{bf}$

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f*(-d^2/b/(tan(1/2*f*x+1/2*e)+1)-d*(a*d-2*b*c)/b^2*ln(tan(1/2*f*x+1/2*e)+1)-2/b^2*(-a^2*d^2+2*a*b*c*d-b^2*c^2)/((a-b)*(a+b))^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e)/((a-b)*(a+b))^(1/2))-d^2/b/(tan(1/2*f*x+1/2*e)-1)+d*(a*d-2*b*c)/b^2*ln(tan(1/2*f*x+1/2*e)-1))
```

3.254.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 231 vs. 2(94) = 188.

Time = 3.30 (sec) , antiderivative size = 518, normalized size of antiderivative = 5.03

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{a+b\sec(e+fx)} dx = \left[\frac{2(a^2b-b^3)d^2 \sin(fx+e) + (b^2c^2-2abcd+a^2d^2)\sqrt{a^2-b^2} \cos(fx+e) \log\left(\frac{2ab \cos(fx+e) - (a^2-2b^2) \cos(fx+e)}{a^2 \cos(fx+e)}\right)}{\dots} \right]$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e)),x, algorithm="fricas")
```

```
output [1/2*(2*(a^2*b - b^3)*d^2*sin(f*x + e) + (b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(a^2 - b^2)*cos(f*x + e)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 + 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)) + (2*(a^2*b - b^3)*c*d - (a^3 - a*b^2)*d^2)*cos(f*x + e)*log(sin(f*x + e) + 1) - (2*(a^2*b - b^3)*c*d - (a^3 - a*b^2)*d^2)*cos(f*x + e)*log(-sin(f*x + e) + 1))/((a^2*b^2 - b^4)*f*cos(f*x + e)), 1/2*(2*(a^2*b - b^3)*d^2*sin(f*x + e) + 2*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e)))*cos(f*x + e) + (2*(a^2*b - b^3)*c*d - (a^3 - a*b^2)*d^2)*cos(f*x + e)*log(sin(f*x + e) + 1) - (2*(a^2*b - b^3)*c*d - (a^3 - a*b^2)*d^2)*cos(f*x + e)*log(-sin(f*x + e) + 1))/((a^2*b^2 - b^4)*f*cos(f*x + e))]
```

3.254.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + b \sec(e + fx)} dx = \int \frac{(c + d \sec(e + fx))^2 \sec(e + fx)}{a + b \sec(e + fx)} dx$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))**2/(a+b*sec(f*x+e)),x)
```

```
output Integral((c + d*sec(e + f*x))**2*sec(e + f*x)/(a + b*sec(e + f*x)), x)
```

3.254.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + b \sec(e + fx)} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e)),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de
```

3.254. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{a+b\sec(e+fx)} dx$

3.254.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 195 vs. 2(94) = 188.

Time = 0.35 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.89

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + b \sec(e + fx)} dx =$$

$$\frac{\frac{2d^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(\tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 1)b} - \frac{(2bcd - ad^2) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{b^2} + \frac{(2bcd - ad^2) \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{b^2} + \frac{2(b^2c^2 - 2abcd + a^2d^2)}{f} \left(\pi \left[\dots \right] \right)}{f}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e)),x, algorithm="giac")`

output `-(2*d^2*tan(1/2*f*x + 1/2*e)/((tan(1/2*f*x + 1/2*e)^2 - 1)*b) - (2*b*c*d - a*d^2)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/b^2 + (2*b*c*d - a*d^2)*log(abs(tan(1/2*f*x + 1/2*e) - 1))/b^2 + 2*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*f*x + 1/2*e) - b*tan(1/2*f*x + 1/2*e))/sqrt(-a^2 + b^2)))/sqrt(-a^2 + b^2)*b^2)/f`

3.254.9 Mupad [B] (verification not implemented)

Time = 18.71 (sec) , antiderivative size = 3559, normalized size of antiderivative = 34.55

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^2}{a + b \sec(e + fx)} dx = \text{Too large to display}$$

input `int((c + d/cos(e + f*x))^2/(cos(e + f*x)*(a + b/cos(e + f*x))),x)`

output

```

- (2*d^2*tan(e/2 + (f*x)/2))/(b*f*(tan(e/2 + (f*x)/2)^2 - 1)) - (atan((((
a + b)*(a - b))^(1/2)*((32*tan(e/2 + (f*x)/2)*(2*a^5*d^4 - b^5*c^4 + a*b^4
*c^4 - 4*a^4*b*d^4 - a^2*b^3*d^4 + 3*a^3*b^2*d^4 - 4*b^5*c^2*d^2 + 12*a*b^
4*c^2*d^2 - 12*a^2*b^3*c*d^3 - 4*a^2*b^3*c^3*d + 16*a^3*b^2*c*d^3 - 18*a^2
*b^3*c^2*d^2 + 10*a^3*b^2*c^2*d^2 + 4*a*b^4*c*d^3 + 4*a*b^4*c^3*d - 8*a^4
*b*c*d^3))/b^2 + (((a + b)*(a - b))^(1/2)*((32*(b^7*c^2 - 2*a*b^6*c^2 - a*b
^6*d^2 + a^2*b^5*c^2 + 2*a^2*b^5*d^2 - a^3*b^4*d^2 + 2*b^7*c*d - 4*a*b^6*c
*d + 2*a^2*b^5*c*d))/b^3 - (32*tan(e/2 + (f*x)/2)*((a + b)*(a - b))^(1/2)*
(a*d - b*c)^2*(2*a*b^6 - 4*a^2*b^5 + 2*a^3*b^4))/(b^2*(b^4 - a^2*b^2)))*(a
*d - b*c)^2)/(b^4 - a^2*b^2))*(a*d - b*c)^2*1i)/(b^4 - a^2*b^2) + (((a + b
)*(a - b))^(1/2)*((32*tan(e/2 + (f*x)/2)*(2*a^5*d^4 - b^5*c^4 + a*b^4*c^4
- 4*a^4*b*d^4 - a^2*b^3*d^4 + 3*a^3*b^2*d^4 - 4*b^5*c^2*d^2 + 12*a*b^4*c^2
*d^2 - 12*a^2*b^3*c*d^3 - 4*a^2*b^3*c^3*d + 16*a^3*b^2*c*d^3 - 18*a^2*b^3
*c^2*d^2 + 10*a^3*b^2*c^2*d^2 + 4*a*b^4*c*d^3 + 4*a*b^4*c^3*d - 8*a^4*b*c*d
^3))/b^2 - (((a + b)*(a - b))^(1/2)*((32*(b^7*c^2 - 2*a*b^6*c^2 - a*b^6*d
^2 + a^2*b^5*c^2 + 2*a^2*b^5*d^2 - a^3*b^4*d^2 + 2*b^7*c*d - 4*a*b^6*c*d +
2*a^2*b^5*c*d))/b^3 + (32*tan(e/2 + (f*x)/2)*((a + b)*(a - b))^(1/2)*(a*d
- b*c)^2*(2*a*b^6 - 4*a^2*b^5 + 2*a^3*b^4))/(b^2*(b^4 - a^2*b^2)))*(a*d -
b*c)^2)/(b^4 - a^2*b^2))*(a*d - b*c)^2*1i)/(b^4 - a^2*b^2))/((64*(a^4*b*d^
6 - a^5*d^6 - 2*b^5*c^5*d + 4*b^5*c^4*d^2 - 12*a*b^4*c^3*d^3 + a*b^4*c^...

```

3.255 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+b \sec(e+fx)} dx$

3.255.1 Optimal result 1870
 3.255.2 Mathematica [A] (verified) 1870
 3.255.3 Rubi [A] (verified) 1871
 3.255.4 Maple [A] (verified) 1873
 3.255.5 Fricas [A] (verification not implemented) 1873
 3.255.6 Sympy [F] 1874
 3.255.7 Maxima [F(-2)] 1874
 3.255.8 Giac [A] (verification not implemented) 1875
 3.255.9 Mupad [B] (verification not implemented) 1876

3.255.1 Optimal result

Integrand size = 29, antiderivative size = 76

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+b \sec(e+fx)} dx = \frac{d \operatorname{arctanh}(\sin(e+fx))}{bf} + \frac{2(bc-ad) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b} b \sqrt{a+bf}}$$

output `d*arctanh(sin(f*x+e))/b/f+2*(-a*d+b*c)*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/b/f/(a-b)^(1/2)/(a+b)^(1/2)`

3.255.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.47

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{a+b \sec(e+fx)} dx = \frac{2(-bc+ad) \operatorname{arctanh}\left(\frac{(-a+b) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}} + \frac{d(-\log(\cos(\frac{1}{2}(e+fx))) - \sin(\frac{1}{2}(e+fx))) + \log(\cos(\frac{1}{2}(e+fx)))}{bf}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + b*Sec[e + f*x]),x]`

output $((2*(-(b*c) + a*d)*\text{ArcTanh}[((-a + b)*\text{Tan}[(e + f*x)/2])/ \text{Sqrt}[a^2 - b^2]])/ \text{Sqrt}[a^2 - b^2] + d*(-\text{Log}[\text{Cos}[(e + f*x)/2] - \text{Sin}[(e + f*x)/2]] + \text{Log}[\text{Cos}[(e + f*x)/2] + \text{Sin}[(e + f*x)/2]]))/(b*f)$

3.255.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4486, 3042, 4257, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d\sec(e+fx))}{a+b\sec(e+fx)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))}{a+b\csc(e+fx+\frac{\pi}{2})} dx \\ & \quad \downarrow 4486 \\ & \frac{(bc-ad) \int \frac{\sec(e+fx)}{a+b\sec(e+fx)} dx}{b} + \frac{d \int \sec(e+fx) dx}{b} \\ & \quad \downarrow 3042 \\ & \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{a+b\csc(e+fx+\frac{\pi}{2})} dx}{b} + \frac{d \int \csc(e+fx+\frac{\pi}{2}) dx}{b} \\ & \quad \downarrow 4257 \\ & \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{a+b\csc(e+fx+\frac{\pi}{2})} dx}{b} + \frac{\text{darctanh}(\sin(e+fx))}{bf} \\ & \quad \downarrow 4318 \\ & \frac{(bc-ad) \int \frac{1}{\frac{a \cos(e+fx)}{b} + 1} dx}{b^2} + \frac{\text{darctanh}(\sin(e+fx))}{bf} \\ & \quad \downarrow 3042 \\ & \frac{(bc-ad) \int \frac{1}{\frac{a \sin(e+fx+\frac{\pi}{2})}{b} + 1} dx}{b^2} + \frac{\text{darctanh}(\sin(e+fx))}{bf} \end{aligned}$$

3.255. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{a+b\sec(e+fx)} dx$

$$\begin{aligned} & \downarrow 3138 \\ & \frac{2(bc - ad) \int \frac{1}{(1 - \frac{a}{b}) \tan^2(\frac{1}{2}(e+fx)) + \frac{a+b}{b}} d \tan(\frac{1}{2}(e+fx))}{b^2 f} + \frac{\operatorname{darctanh}(\sin(e+fx))}{bf} \\ & \downarrow 221 \\ & \frac{2(bc - ad) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{bf \sqrt{a-b} \sqrt{a+b}} + \frac{\operatorname{darctanh}(\sin(e+fx))}{bf} \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + b*Sec[e + f*x]),x]`

output `(d*ArcTanh[Sin[e + f*x]])/(b*f) + (2*(b*c - a*d)*ArcTanh[(Sqrt[a - b]*Tan[e + f*x]/2])/Sqrt[a + b]]/(Sqrt[a - b]*b*Sqrt[a + b]*f)`

3.255.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4318 `Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

```
rule 4486 Int[(csc[(e_.) + (f_.)*(x_)]*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Simp[B/b Int[Csc[e + f*x], x], x] + Simp[(A*b - a*B)/b Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f, A, B}, x] && NeQ[A*b - a*B, 0]
```

3.255.4 Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.21

method	result
derivativedivides	$-\frac{2(ad-bc) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b\sqrt{(a-b)(a+b)}} + \frac{d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{b} - \frac{d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{b}$
default	$-\frac{2(ad-bc) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b\sqrt{(a-b)(a+b)}} + \frac{d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{b} - \frac{d \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{b}$
risch	$\frac{\ln\left(e^{i(fx+e)} + \frac{-ia^2+ib^2+b\sqrt{a^2-b^2}}{a\sqrt{a^2-b^2}}\right)ad}{\sqrt{a^2-b^2}fb} - \frac{\ln\left(e^{i(fx+e)} + \frac{-ia^2+ib^2+b\sqrt{a^2-b^2}}{a\sqrt{a^2-b^2}}\right)c}{\sqrt{a^2-b^2}f} - \frac{\ln\left(e^{i(fx+e)} + \frac{ia^2-ib^2+b\sqrt{a^2-b^2}}{\sqrt{a^2-b^2}a}\right)a}{\sqrt{a^2-b^2}fb}$

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f*(-2*(a*d-b*c)/b/((a-b)*(a+b))^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e)/((a-b)*(a+b))^(1/2))+d/b*ln(tan(1/2*f*x+1/2*e)+1)-d/b*ln(tan(1/2*f*x+1/2*e)-1))
```

3.255.5 Fracas [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 309, normalized size of antiderivative = 4.07

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{a + b \sec(e + fx)} dx$$

$$= \frac{\left[(a^2 - b^2)d \log(\sin(fx + e) + 1) - (a^2 - b^2)d \log(-\sin(fx + e) + 1) - \sqrt{a^2 - b^2}(bc - ad) \log\left(\frac{2ab \cos(fx + e) + a^2 - b^2}{\sqrt{a^2 - b^2}(a + b \sec(e + fx))}\right) \right]}{2(a^2b - b^3)f}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e)),x, algorithm="fraca s")
```

```
output [1/2*((a^2 - b^2)*d*log(sin(f*x + e) + 1) - (a^2 - b^2)*d*log(-sin(f*x + e) + 1) - sqrt(a^2 - b^2)*(b*c - a*d)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 - 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)))/((a^2*b - b^3)*f), 1/2*((a^2 - b^2)*d*log(sin(f*x + e) + 1) - (a^2 - b^2)*d*log(-sin(f*x + e) + 1) + 2*sqrt(-a^2 + b^2)*(b*c - a*d)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e))))/((a^2*b - b^3)*f)]
```

3.255.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{a + b \sec(e + fx)} dx = \int \frac{(c + d \sec(e + fx)) \sec(e + fx)}{a + b \sec(e + fx)} dx$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e)),x)
```

```
output Integral((c + d*sec(e + f*x))*sec(e + f*x)/(a + b*sec(e + f*x)), x)
```

3.255.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{a + b \sec(e + fx)} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e)),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de
```

3.255.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.67

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{a + b \sec(e + fx)} dx$$

$$= \frac{\frac{d \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) + 1|)}{b} - \frac{d \log(|\tan(\frac{1}{2} fx + \frac{1}{2} e) - 1|)}{b} - \frac{2 \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan\left(\frac{a \tan(\frac{1}{2} fx + \frac{1}{2} e) - b \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-a^2+b^2}}\right) \right)}{\sqrt{-a^2+b^2}}}{f} (bc - a^2)$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e)),x, algorithm="giac")
```

```
output (d*log(abs(tan(1/2*f*x + 1/2*e) + 1))/b - d*log(abs(tan(1/2*f*x + 1/2*e) - 1))/b - 2*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*f*x + 1/2*e) - b*tan(1/2*f*x + 1/2*e))/sqrt(-a^2 + b^2)))*(b*c - a*d)/(sqrt(-a^2 + b^2)*b))/f
```

3.255.9 Mupad [B] (verification not implemented)

Time = 14.62 (sec) , antiderivative size = 571, normalized size of antiderivative = 7.51

$$\begin{aligned}
& \int \frac{\sec(e+fx)(c+d\sec(e+fx))}{a+b\sec(e+fx)} dx \\
&= \frac{b^2 c \ln\left(\frac{b \sin\left(\frac{e+fx}{2}\right) - a \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{a^2-b^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f(a^2-b^2)^{3/2}} \\
&\quad - \frac{a^2 c \ln\left(\frac{b \sin\left(\frac{e+fx}{2}\right) - a \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{a^2-b^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f(a^2-b^2)^{3/2}} - \frac{2bd \operatorname{atanh}\left(\frac{\sin\left(\frac{e+fx}{2}\right)}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f(a^2-b^2)} \\
&\quad + \frac{c \ln\left(\frac{a \cos\left(\frac{e+fx}{2}\right) + b \cos\left(\frac{e+fx}{2}\right) + \sin\left(\frac{e+fx}{2}\right) \sqrt{a^2-b^2}}{\cos\left(\frac{e+fx}{2}\right)}\right) \sqrt{(a+b)(a-b)}}{f(a^2-b^2)} \\
&\quad - \frac{abd \ln\left(\frac{b \sin\left(\frac{e+fx}{2}\right) - a \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{a^2-b^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{f(a^2-b^2)^{3/2}} \\
&\quad + \frac{2a^2 d \operatorname{atanh}\left(\frac{\sin\left(\frac{e+fx}{2}\right)}{\cos\left(\frac{e+fx}{2}\right)}\right)}{bf(a^2-b^2)} + \frac{a^3 d \ln\left(\frac{b \sin\left(\frac{e+fx}{2}\right) - a \sin\left(\frac{e+fx}{2}\right) + \cos\left(\frac{e+fx}{2}\right) \sqrt{a^2-b^2}}{\cos\left(\frac{e+fx}{2}\right)}\right)}{bf(a^2-b^2)^{3/2}} \\
&\quad - \frac{ad \ln\left(\frac{a \cos\left(\frac{e+fx}{2}\right) + b \cos\left(\frac{e+fx}{2}\right) + \sin\left(\frac{e+fx}{2}\right) \sqrt{a^2-b^2}}{\cos\left(\frac{e+fx}{2}\right)}\right) \sqrt{(a+b)(a-b)}}{bf(a^2-b^2)}
\end{aligned}$$

input `int((c + d/cos(e + f*x))/(cos(e + f*x)*(a + b/cos(e + f*x))),x)`

output

$$\begin{aligned} & (b^2*c*\log((b*\sin(e/2 + (f*x)/2) - a*\sin(e/2 + (f*x)/2) + \cos(e/2 + (f*x)/2) \\ & /2)*(a^2 - b^2)^{(1/2)}/\cos(e/2 + (f*x)/2)))/(f*(a^2 - b^2)^{(3/2)}) - (a^2*c* \\ & \log((b*\sin(e/2 + (f*x)/2) - a*\sin(e/2 + (f*x)/2) + \cos(e/2 + (f*x)/2)*(a^2 \\ & - b^2)^{(1/2)}/\cos(e/2 + (f*x)/2)))/(f*(a^2 - b^2)^{(3/2)}) - (2*b*d*atanh(s \\ & \sin(e/2 + (f*x)/2)/\cos(e/2 + (f*x)/2)))/(f*(a^2 - b^2)) + (c*\log((a*\cos(e/2 \\ & + (f*x)/2) + b*\cos(e/2 + (f*x)/2) + \sin(e/2 + (f*x)/2)*(a^2 - b^2)^{(1/2)}) \\ & / \cos(e/2 + (f*x)/2))*((a + b)*(a - b))^{(1/2)})/(f*(a^2 - b^2)) - (a*b*d*\log \\ & ((b*\sin(e/2 + (f*x)/2) - a*\sin(e/2 + (f*x)/2) + \cos(e/2 + (f*x)/2)*(a^2 - \\ & b^2)^{(1/2)})/\cos(e/2 + (f*x)/2)))/(f*(a^2 - b^2)^{(3/2)}) + (2*a^2*d*atanh(si \\ & n(e/2 + (f*x)/2)/\cos(e/2 + (f*x)/2)))/(b*f*(a^2 - b^2)) + (a^3*d*\log((b*si \\ & n(e/2 + (f*x)/2) - a*\sin(e/2 + (f*x)/2) + \cos(e/2 + (f*x)/2)*(a^2 - b^2)^{(\\ & 1/2)})/\cos(e/2 + (f*x)/2)))/(b*f*(a^2 - b^2)^{(3/2)}) - (a*d*\log((a*\cos(e/2 + \\ & (f*x)/2) + b*\cos(e/2 + (f*x)/2) + \sin(e/2 + (f*x)/2)*(a^2 - b^2)^{(1/2)})/c \\ & \cos(e/2 + (f*x)/2))*((a + b)*(a - b))^{(1/2)})/(b*f*(a^2 - b^2)) \end{aligned}$$

3.256
$$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))} dx$$

3.256.1 Optimal result	1878
3.256.2 Mathematica [A] (verified)	1878
3.256.3 Rubi [A] (verified)	1879
3.256.4 Maple [A] (verified)	1881
3.256.5 Fricas [A] (verification not implemented)	1881
3.256.6 Sympy [F]	1882
3.256.7 Maxima [F(-2)]	1883
3.256.8 Giac [B] (verification not implemented)	1883
3.256.9 Mupad [B] (verification not implemented)	1884

3.256.1 Optimal result

Integrand size = 31, antiderivative size = 121

$$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))} dx = \frac{2b \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b} \sqrt{a+b} (bc-ad) f} - \frac{2d \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{\sqrt{c-d} \sqrt{c+d} (bc-ad) f}$$

output `2*b*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/(-a*d+b*c)/f/(a-b)^(1/2)/(a+b)^(1/2)-2*d*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(-a*d+b*c)/f/(c-d)^(1/2)/(c+d)^(1/2)`

3.256.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.98

$$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))} dx = -\frac{2b \operatorname{arctanh}\left(\frac{(-a+b) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2} (bc-ad) f} - \frac{2d \operatorname{arctanh}\left(\frac{(-c+d) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{(-bc+ad) \sqrt{c^2-d^2} f}$$

input `Integrate[Sec[e + f*x]/((a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])),x]`

3.256.
$$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))} dx$$

output
$$\frac{(-2*b*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]]/(Sqrt[a^2 - b^2]*(b*c - a*d)*f) - (2*d*ArcTanh[((-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2]]/((-(b*c) + a*d)*Sqrt[c^2 - d^2]*f)}$$

3.256.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3042, 4476, 3042, 3480, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e + fx)}{(a + b \sec(e + fx))(c + d \sec(e + fx))} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\csc(e + fx + \frac{\pi}{2})}{(a + b \csc(e + fx + \frac{\pi}{2}))(c + d \csc(e + fx + \frac{\pi}{2}))} dx \\ & \quad \downarrow 4476 \\ & \int \frac{\cos(e + fx)}{(a \cos(e + fx) + b)(c \cos(e + fx) + d)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\sin(e + fx + \frac{\pi}{2})}{(a \sin(e + fx + \frac{\pi}{2}) + b)(c \sin(e + fx + \frac{\pi}{2}) + d)} dx \\ & \quad \downarrow 3480 \\ & \frac{b \int \frac{1}{b+a \cos(e+fx)} dx}{bc - ad} - \frac{d \int \frac{1}{d+c \cos(e+fx)} dx}{bc - ad} \\ & \quad \downarrow 3042 \\ & \frac{b \int \frac{1}{b+a \sin(e+fx+\frac{\pi}{2})} dx}{bc - ad} - \frac{d \int \frac{1}{d+c \sin(e+fx+\frac{\pi}{2})} dx}{bc - ad} \\ & \quad \downarrow 3138 \\ & \frac{2b \int \frac{1}{-((a-b) \tan^2(\frac{1}{2}(e+fx)))+a+b} d \tan(\frac{1}{2}(e+fx))}{f(bc - ad)} - \frac{2d \int \frac{1}{-((c-d) \tan^2(\frac{1}{2}(e+fx)))+c+d} d \tan(\frac{1}{2}(e+fx))}{f(bc - ad)} \\ & \quad \downarrow 221 \end{aligned}$$

3.256.
$$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))} dx$$

$$\frac{2b \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{f\sqrt{a-b}\sqrt{a+b}(bc-ad)} - \frac{2d \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{f\sqrt{c-d}\sqrt{c+d}(bc-ad)}$$

input `Int[Sec[e + f*x]/((a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])),x]`

output `(2*b*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]]/(Sqrt[a - b]*Sqrt[a + b]*(b*c - a*d)*f) - (2*d*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]]/(Sqrt[c - d]*Sqrt[c + d]*(b*c - a*d)*f)`

3.256.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4476 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*(csc[(e_) + (f_)*(x_)])*(b_) + (a_)^(m_)*(csc[(e_) + (f_)*(x_)])*(d_) + (c_)^(n_), x_Symbol] := Simp[1/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && IntegerQ[m] && IntegerQ[n]`

3.256.4 Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{2d \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(ad-bc)\sqrt{(c+d)(c-d)}} - \frac{2b \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(ad-bc)\sqrt{(a-b)(a+b)}}$
default	$\frac{2d \operatorname{arctanh}\left(\frac{(c-d) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c+d)(c-d)}}\right)}{(ad-bc)\sqrt{(c+d)(c-d)}} - \frac{2b \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(ad-bc)\sqrt{(a-b)(a+b)}}$
risch	$\frac{b \ln\left(\frac{e^{i(fx+e)} - \frac{ia^2-ib^2-b\sqrt{a^2-b^2}}{\sqrt{a^2-b^2}a}}{\sqrt{a^2-b^2}(ad-bc)f}\right)}{\sqrt{a^2-b^2}(ad-bc)f} - \frac{b \ln\left(\frac{e^{i(fx+e)} + \frac{ia^2-ib^2+b\sqrt{a^2-b^2}}{\sqrt{a^2-b^2}a}}{\sqrt{a^2-b^2}(ad-bc)f}\right)}{\sqrt{a^2-b^2}(ad-bc)f} + \frac{d \ln\left(\frac{e^{i(fx+e)} + \frac{ic^2-id^2+\sqrt{c^2-d^2}d}{\sqrt{c^2-d^2}c}}{\sqrt{c^2-d^2}(ad-bc)f}\right)}{\sqrt{c^2-d^2}(ad-bc)f}$

input `int(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/f*(2*d/(a*d-b*c)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/((c+d)*(c-d))^(1/2))-2*b/(a*d-b*c)/((a-b)*(a+b))^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e)/((a-b)*(a+b))^(1/2)))`

3.256.5 Fracas [A] (verification not implemented)

Time = 1.86 (sec) , antiderivative size = 1040, normalized size of antiderivative = 8.60

$$\int \frac{\sec(e + fx)}{(a + b \sec(e + fx))(c + d \sec(e + fx))} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="fracas")`

output

```

[-1/2*((a^2 - b^2)*sqrt(c^2 - d^2)*d*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + (b*c^2 - b*d^2)*sqrt(a^2 - b^2)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 - 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)))/(((a^2*b - b^3)*c^3 - (a^3 - a*b^2)*c^2*d - (a^2*b - b^3)*c*d^2 + (a^3 - a*b^2)*d^3)*f), -1/2*((a^2 - b^2)*sqrt(c^2 - d^2)*d*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) - 2*(b*c^2 - b*d^2)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e)))/(((a^2*b - b^3)*c^3 - (a^3 - a*b^2)*c^2*d - (a^2*b - b^3)*c*d^2 + (a^3 - a*b^2)*d^3)*f), -1/2*(2*(a^2 - b^2)*sqrt(-c^2 + d^2)*d*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) + (b*c^2 - b*d^2)*sqrt(a^2 - b^2)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 - 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)))/(((a^2*b - b^3)*c^3 - (a^3 - a*b^2)*c^2*d - (a^2*b - b^3)*c*d^2 + (a^3 - a*b^2)*d^3)*f), -((a^2 - b^2)*sqrt(-c^2 + d^2)*d*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) - (b*c^2 - b*d^2)*sqrt(-a^2 + b^2)*arctan(-...

```

3.256.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{(a + b\sec(e + fx))(c + d\sec(e + fx))} dx = \int \frac{\sec(e + fx)}{(a + b\sec(e + fx))(c + d\sec(e + fx))} dx$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x)`

output `Integral(sec(e + f*x)/((a + b*sec(e + f*x))*(c + d*sec(e + f*x))), x)`

3.256.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + b \sec(e + fx))(c + d \sec(e + fx))} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see 'assume?' f or more de

3.256.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 522 vs. 2(103) = 206.

Time = 0.40 (sec) , antiderivative size = 522, normalized size of antiderivative = 4.31

$$\int \frac{\sec(e + fx)}{(a + b \sec(e + fx))(c + d \sec(e + fx))} dx$$

$$= \frac{(\sqrt{-c^2+d^2}b(c-2d)|c-d|+\sqrt{-c^2+d^2}ad|c-d|+\sqrt{-c^2+d^2}|-bc+ad||c-d|) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] + \arctan \left(\frac{2\sqrt{\frac{1}{2}} \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)}{\sqrt{-2ac-2bd+\sqrt{-4(ac+bc+ad+bd)(ac-bc-ad+bd)}}} \right)}{(bc-ad)^2(c^2-2cd+d^2)+(c^3-2c^2d+cd^2)a|-bc+ad|-(c^2d-2cd^2+d^3)b|-bc+ad|} \right)}{(bc-ad)^2(c^2-2cd+d^2)+(c^3-2c^2d+cd^2)a|-bc+ad|-(c^2d-2cd^2+d^3)b|-bc+ad|}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e)),x, algorithm="giac")`

output $((\sqrt{-c^2 + d^2} * b * (c - 2*d) * \text{abs}(c - d) + \sqrt{-c^2 + d^2} * a * d * \text{abs}(c - d) + \sqrt{-c^2 + d^2} * \text{abs}(-b*c + a*d) * \text{abs}(c - d)) * (\pi * \text{floor}(1/2 * (f*x + e)) / \pi + 1/2) + \arctan(2 * \sqrt{1/2} * \tan(1/2 * f*x + 1/2 * e) / \sqrt{-(2*a*c - 2*b*d + \sqrt{-4*(a*c + b*c + a*d + b*d)*(a*c - b*c - a*d + b*d) + 4*(a*c - b*d)^2}) / (a*c - b*c - a*d + b*d))) / ((b*c - a*d)^2 * (c^2 - 2*c*d + d^2) + (c^3 - 2*c^2*d + c*d^2) * a * \text{abs}(-b*c + a*d) - (c^2*d - 2*c*d^2 + d^3) * b * \text{abs}(-b*c + a*d)) + (\sqrt{-a^2 + b^2} * b * c * \text{abs}(a - b) + \sqrt{-a^2 + b^2} * (a - 2*b) * d * \text{abs}(a - b) - \sqrt{-a^2 + b^2} * \text{abs}(-b*c + a*d) * \text{abs}(a - b)) * (\pi * \text{floor}(1/2 * (f*x + e)) / \pi + 1/2) + \arctan(2 * \sqrt{1/2} * \tan(1/2 * f*x + 1/2 * e) / \sqrt{-(2*a*c - 2*b*d - \sqrt{-4*(a*c + b*c + a*d + b*d)*(a*c - b*c - a*d + b*d) + 4*(a*c - b*d)^2}) / (a*c - b*c - a*d + b*d))) / ((a^2 - 2*a*b + b^2) * (b*c - a*d)^2 - (a^3 - 2*a^2*b + a*b^2) * c * \text{abs}(-b*c + a*d) + (a^2*b - 2*a*b^2 + b^3) * d * \text{abs}(-b*c + a*d))) / f$

3.256.9 Mupad [B] (verification not implemented)

Time = 15.93 (sec) , antiderivative size = 2665, normalized size of antiderivative = 22.02

$$\int \frac{\sec(e + fx)}{(a + b \sec(e + fx))(c + d \sec(e + fx))} dx = \text{Too large to display}$$

input `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))*(c + d/cos(e + f*x))),x)`

output

$$\begin{aligned}
 & (b^5 c^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a^5 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i + b^3 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{3/2} i \\
 & + b^5 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a^2 b^3 c^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a^3 b^2 c^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i \\
 & - a^2 b^3 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i + a^3 b^2 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - b^3 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{3/2} i \\
 & - b^5 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i + a b^2 c^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{3/2} i + a b^4 c^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i \\
 & + a^4 b d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i + a^2 b^3 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i + a^2 b^3 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i \\
 & + a^3 b^2 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a b^2 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{3/2} i - a b^4 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i \\
 & + a^4 b^2 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a b^2 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{3/2} i - a b^4 c d \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i \\
 &) / (a^6 d^2 - b^6 c^2 + 2 a^2 b^4 c^2 - a^4 b^2 c^2 + a^2 b^4 d^2 - 2 a^4 b^2 d^2) (a^2 - b^2)^{1/2} i / (f (a^3 d^3 - b^3 c^3 + a^2 b c^3 - a b^2 d^3 - a^3 c^2 d + b^3 c d^2 + a b^2 c^2 d - a^2 b c d^2)) - (b d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i + b^3 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{3/2} i + b^5 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a^2 b^3 c^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a^3 b^2 c^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i - a^2 b^3 d^2 \tan(e/2 + (f*x)/2) (a^2 - b^2)^{1/2} i + a \dots
 \end{aligned}$$

3.257
$$\int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx$$

3.257.1 Optimal result	1886
3.257.2 Mathematica [A] (verified)	1886
3.257.3 Rubi [A] (verified)	1887
3.257.4 Maple [A] (verified)	1890
3.257.5 Fricas [B] (verification not implemented)	1891
3.257.6 Sympy [F]	1891
3.257.7 Maxima [F(-2)]	1892
3.257.8 Giac [A] (verification not implemented)	1892
3.257.9 Mupad [B] (verification not implemented)	1893

3.257.1 Optimal result

Integrand size = 31, antiderivative size = 187

$$\int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx$$

$$= \frac{2b^2 \operatorname{arctanh}\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b}\sqrt{a+b}(bc-ad)^2 f} - \frac{2d(2bc^2 - acd - bd^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{(c-d)^{3/2}(c+d)^{3/2}(bc-ad)^2 f}$$

$$+ \frac{d^2 \sin(e+fx)}{(bc-ad)(c^2-d^2)f(d+c\cos(e+fx))}$$

```
output -2*d*(-a*c*d+2*b*c^2-b*d^2)*arctanh((c-d)^(1/2)*tan(1/2*f*x+1/2*e)/(c+d)^(1/2))/(c-d)^(3/2)/(c+d)^(3/2)/(-a*d+b*c)^2/f+d^2*sin(f*x+e)/(-a*d+b*c)/(c^2-d^2)/f/(d+c*cos(f*x+e))+2*b^2*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/(-a*d+b*c)^2/f/(a-b)^(1/2)/(a+b)^(1/2)
```

3.257.2 Mathematica [A] (verified)

Time = 1.34 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.22

$$\int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx$$

$$= \frac{-2b^2(c^2-d^2)^{3/2} \operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right) (d+c\cos(e+fx)) - \sqrt{a^2-b^2}d \left(-2(2bc^2 - acd - bd^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)\right)}{\sqrt{a^2-b^2}(c-d)(c+d)(bc-ad)^2\sqrt{c^2-d^2}f}$$

3.257.
$$\int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx$$

input `Integrate[Sec[e + f*x]/((a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^2),x]`

output $(-2*b^2*(c^2 - d^2)^{3/2}*ArcTanh[(-a + b)*Tan[(e + f*x)/2]]/Sqrt[a^2 - b^2])*(d + c*Cos[e + f*x]) - Sqrt[a^2 - b^2]*d*(-2*(2*b*c^2 - a*c*d - b*d^2)*ArcTanh[(-c + d)*Tan[(e + f*x)/2]]/Sqrt[c^2 - d^2])*(d + c*Cos[e + f*x]) + d*(-(b*c) + a*d)*Sqrt[c^2 - d^2]*Sin[e + f*x])/(Sqrt[a^2 - b^2]*(c - d)*(c + d)*(b*c - a*d)^2*Sqrt[c^2 - d^2]*f*(d + c*Cos[e + f*x]))$

3.257.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.18, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4476, 3042, 3535, 25, 3042, 3480, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(e + fx)}{(a + b \sec(e + fx))(c + d \sec(e + fx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e + fx + \frac{\pi}{2})}{(a + b \csc(e + fx + \frac{\pi}{2}))(c + d \csc(e + fx + \frac{\pi}{2}))^2} dx \\ & \quad \downarrow \text{4476} \\ & \int \frac{\cos^2(e + fx)}{(a \cos(e + fx) + b)(c \cos(e + fx) + d)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(e + fx + \frac{\pi}{2})^2}{(a \sin(e + fx + \frac{\pi}{2}) + b)(c \sin(e + fx + \frac{\pi}{2}) + d)^2} dx \\ & \quad \downarrow \text{3535} \\ & \frac{\int -\frac{bcd + (acd - b(c^2 - d^2)) \cos(e + fx)}{(b + a \cos(e + fx))(d + c \cos(e + fx))} dx}{(c^2 - d^2)(bc - ad)} + \frac{d^2 \sin(e + fx)}{f(c^2 - d^2)(bc - ad)(c \cos(e + fx) + d)} \\ & \quad \downarrow \text{25} \\ & \frac{d^2 \sin(e + fx)}{f(c^2 - d^2)(bc - ad)(c \cos(e + fx) + d)} - \frac{\int \frac{bcd + (acd - b(c^2 - d^2)) \cos(e + fx)}{(b + a \cos(e + fx))(d + c \cos(e + fx))} dx}{(c^2 - d^2)(bc - ad)} \end{aligned}$$

3.257. $\int \frac{\sec(e + fx)}{(a + b \sec(e + fx))(c + d \sec(e + fx))^2} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{d^2 \sin(e+fx)}{f(c^2-d^2)(bc-ad)(c \cos(e+fx)+d)} - \frac{\int \frac{bcd+(acd-b(c^2-d^2)) \sin(e+fx+\frac{\pi}{2})}{(b+a \sin(e+fx+\frac{\pi}{2}))(d+c \sin(e+fx+\frac{\pi}{2}))} dx}{(c^2-d^2)(bc-ad)} \\
& \downarrow \text{3480} \\
& \frac{d^2 \sin(e+fx)}{f(c^2-d^2)(bc-ad)(c \cos(e+fx)+d)} - \frac{b^2(c^2-d^2) \int \frac{1}{b+a \cos(e+fx)} dx}{bc-ad} - \frac{d(acd-b(2c^2-d^2)) \int \frac{1}{d+c \cos(e+fx)} dx}{bc-ad} \\
& \downarrow \text{3042} \\
& \frac{d^2 \sin(e+fx)}{f(c^2-d^2)(bc-ad)(c \cos(e+fx)+d)} - \frac{b^2(c^2-d^2) \int \frac{1}{b+a \sin(e+fx+\frac{\pi}{2})} dx}{bc-ad} - \frac{d(acd-b(2c^2-d^2)) \int \frac{1}{d+c \sin(e+fx+\frac{\pi}{2})} dx}{bc-ad} \\
& \downarrow \text{3138} \\
& \frac{d^2 \sin(e+fx)}{f(c^2-d^2)(bc-ad)(c \cos(e+fx)+d)} - \frac{2b^2(c^2-d^2) \int \frac{1}{-(a-b) \tan^2(\frac{1}{2}(e+fx))+a+b} d \tan(\frac{1}{2}(e+fx))}{f(bc-ad)} - \frac{2d(acd-b(2c^2-d^2)) \int \frac{1}{-(c-d) \tan^2(\frac{1}{2}(e+fx))+c+d} d \tan(\frac{1}{2}(e+fx))}{f(bc-ad)} \\
& \downarrow \text{221} \\
& \frac{d^2 \sin(e+fx)}{f \sqrt{a-b} \sqrt{a+b} (bc-ad)} - \frac{2d(acd-b(2c^2-d^2)) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{f \sqrt{c-d} \sqrt{c+d} (bc-ad)} \\
& \frac{2b^2(c^2-d^2) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{f \sqrt{a-b} \sqrt{a+b} (bc-ad)} - \frac{2d(acd-b(2c^2-d^2)) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{f \sqrt{c-d} \sqrt{c+d} (bc-ad)} \\
& \frac{d^2 \sin(e+fx)}{f(c^2-d^2)(bc-ad)(c \cos(e+fx)+d)} - \frac{2b^2(c^2-d^2) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{f \sqrt{a-b} \sqrt{a+b} (bc-ad)} - \frac{2d(acd-b(2c^2-d^2)) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e+fx))}{\sqrt{c+d}}\right)}{f \sqrt{c-d} \sqrt{c+d} (bc-ad)}
\end{aligned}$$

input `Int[Sec[e + f*x]/((a + b*Sec[e + f*x])*(c + d*Sec[e + f*x])^2),x]`

output `-(((-2*b^2*(c^2 - d^2)*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*(b*c - a*d)*f) - (2*d*(a*c*d - b*(2*c^2 - d^2))*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]])/(Sqrt[c - d]*Sqrt[c + d]*(b*c - a*d)*f))/((b*c - a*d)*(c^2 - d^2))) + (d^2*Sin[e + f*x])/((b*c - a*d)*(c^2 - d^2)*f*(d + c*Cos[e + f*x]))`

$$3.257. \int \frac{\sec(e+fx)}{(a+b \sec(e+fx))(c+d \sec(e+fx))^2} dx$$

3.257.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3535 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[a*(m + 1)*(b*c - a*d)*(A + C) + d*(A*b^2 + a^2*C)*(m + n + 2) - (c*(A*b^2 + a^2*C) + b*(m + 1)*(b*c - a*d)*(A + C))*Sin[e + f*x] - d*(A*b^2 + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 4476 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.)^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[1
/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c
*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c -
a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.257.4 Maple [A] (verified)

Time = 1.37 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{2b^2 \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(ad-bc)^2\sqrt{(a-b)(a+b)}} - \frac{2d \left(-\frac{d(ad-bc)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^2-d^2)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{(acd-2bc^2+bd^2)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c-d)(c+d)}}\right)}{(c+d)(c-d)\sqrt{(c+d)(c-d)}} \right)}{(ad-bc)^2}$
default	$\frac{2b^2 \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(ad-bc)^2\sqrt{(a-b)(a+b)}} - \frac{2d \left(-\frac{d(ad-bc)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(c^2-d^2)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 c - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 d - c - d} - \frac{(acd-2bc^2+bd^2)\operatorname{arctanh}\left(\frac{(c-d)\tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(c-d)(c+d)}}\right)}{(c+d)(c-d)\sqrt{(c+d)(c-d)}} \right)}{(ad-bc)^2}$
risch	$\frac{2id^2(d e^{i(fx+e)} + c)}{c(c^2-d^2)(-ad+bc)f(e^{2i(fx+e)}c+2de^{i(fx+e)}+c)} + \frac{d^2 \ln\left(e^{i(fx+e)} + \frac{ic^2-id^2+\sqrt{c^2-d^2}d}{\sqrt{c^2-d^2}c}\right)ac}{\sqrt{c^2-d^2}(ad-bc)^2(c+d)(c-d)f} - \frac{2d \ln\left(e^{i(fx+e)} + \frac{ic^2-d^2}{\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}(ad-bc)}$

```
input int(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x,method=_RETURNVERBOSE
)
```

```
output 1/f*(2*b^2/(a*d-b*c)^2/((a-b)*(a+b))^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e
))/(a-b)*(a+b))^(1/2))-2*d/(a*d-b*c)^2*(-d*(a*d-b*c)/(c^2-d^2)*tan(1/2*f*x
+1/2*e)/(tan(1/2*f*x+1/2*e)^2*c-tan(1/2*f*x+1/2*e)^2*d-c-d)-(a*c*d-2*b*c^2
+b*d^2)/(c+d)/(c-d)/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/(
(c+d)*(c-d))^(1/2))))
```

$$3.257. \int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx$$

3.257.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 658 vs. $2(169) = 338$.

Time = 96.83 (sec) , antiderivative size = 2863, normalized size of antiderivative = 15.31

$$\int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="fricas")`

output `[1/2*((b^2*c^4*d - 2*b^2*c^2*d^3 + b^2*d^5 + (b^2*c^5 - 2*b^2*c^3*d^2 + b^2*c*d^4)*cos(f*x + e))*sqrt(a^2 - b^2)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 + 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)) + (2*(a^2*b - b^3)*c^2*d^2 - (a^3 - a*b^2)*c*d^3 - (a^2*b - b^3)*d^4 + (2*(a^2*b - b^3)*c^3*d - (a^3 - a*b^2)*c^2*d^2 - (a^2*b - b^3)*c*d^3)*cos(f*x + e))*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 - 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*((a^2*b - b^3)*c^3*d^2 - (a^3 - a*b^2)*c^2*d^3 - (a^2*b - b^3)*c*d^4 + (a^3 - a*b^2)*d^5)*sin(f*x + e))/((a^2*b^2 - b^4)*c^7 - 2*(a^3*b - a*b^3)*c^6*d + (a^4 - 3*a^2*b^2 + 2*b^4)*c^5*d^2 + 4*(a^3*b - a*b^3)*c^4*d^3 - (2*a^4 - 3*a^2*b^2 + b^4)*c^3*d^4 - 2*(a^3*b - a*b^3)*c^2*d^5 + (a^4 - a^2*b^2)*c*d^6)*f*cos(f*x + e) + ((a^2*b^2 - b^4)*c^6*d - 2*(a^3*b - a*b^3)*c^5*d^2 + (a^4 - 3*a^2*b^2 + 2*b^4)*c^4*d^3 + 4*(a^3*b - a*b^3)*c^3*d^4 - (2*a^4 - 3*a^2*b^2 + b^4)*c^2*d^5 - 2*(a^3*b - a*b^3)*c*d^6 + (a^4 - a^2*b^2)*d^7)*f), 1/2*(2*(b^2*c^4*d - 2*b^2*c^2*d^3 + b^2*d^5 + (b^2*c^5 - 2*b^2*c^3*d^2 + b^2*c*d^4)*cos(f*x + e))*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e))) + (2*(a^2*b - b^3)*c^2*d^2 - (a^3 - a*b^2)*c*d^3 - (a^2*b - b^3)*d^4 + (2*(a^2*b - b^3)*c^3*d - (a^3 - a*b^2)*c^2*d^2 - (a^2*b - ...`

3.257.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx \\ &= \int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx \end{aligned}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))**2,x)`

output `Integral(sec(e + f*x)/((a + b*sec(e + f*x))*(c + d*sec(e + f*x))**2), x)`

3.257.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + b\sec(e + fx))(c + d\sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see 'assume?' f or more de`

3.257.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 331, normalized size of antiderivative = 1.77

$$\int \frac{\sec(e + fx)}{(a + b\sec(e + fx))(c + d\sec(e + fx))^2} dx =$$

$$2 \left(\frac{\left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} fx + \frac{1}{2} e) - b \tan(\frac{1}{2} fx + \frac{1}{2} e)}{\sqrt{-a^2+b^2}} \right) \right) b^2}{(b^2c^2 - 2abcd + a^2d^2)\sqrt{-a^2+b^2}} \right) + \frac{d^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)}{(bc^3 - ac^2d - bcd^2 + ad^3) \left(c \tan(\frac{1}{2} fx + \frac{1}{2} e) \right)^2 - d \tan(\frac{1}{2} fx + \frac{1}{2} e)}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^2,x, algorithm="giac")`

3.257. $\int \frac{\sec(e+fx)}{(a+b\sec(e+fx))(c+d\sec(e+fx))^2} dx$

output
$$-2*((\pi*\text{floor}(1/2*(f*x + e)/\pi + 1/2)*\text{sgn}(2*a - 2*b) + \arctan((a*\tan(1/2*f*x + 1/2*e) - b*\tan(1/2*f*x + 1/2*e))/\sqrt{-a^2 + b^2}))*b^2/((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*\sqrt{-a^2 + b^2}) + d^2*\tan(1/2*f*x + 1/2*e)/((b*c^3 - a*c^2*d - b*c*d^2 + a*d^3)*(c*\tan(1/2*f*x + 1/2*e)^2 - d*\tan(1/2*f*x + 1/2*e)^2 - c - d)) - (2*b*c^2*d - a*c*d^2 - b*d^3)*(\pi*\text{floor}(1/2*(f*x + e)/\pi + 1/2)*\text{sgn}(2*c - 2*d) + \arctan((c*\tan(1/2*f*x + 1/2*e) - d*\tan(1/2*f*x + 1/2*e))/\sqrt{-c^2 + d^2}))/((b^2*c^4 - 2*a*b*c^3*d + a^2*c^2*d^2 - b^2*c^2*d^2 + 2*a*b*c*d^3 - a^2*d^4)*\sqrt{-c^2 + d^2}))/f$$

3.257.9 Mupad [B] (verification not implemented)

Time = 27.24 (sec) , antiderivative size = 20827, normalized size of antiderivative = 111.37

$$\int \frac{\sec(e + fx)}{(a + b\sec(e + fx))(c + d\sec(e + fx))^2} dx = \text{Too large to display}$$

input `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))*(c + d/cos(e + f*x))^2),x)`

output
$$(2*d^2*\tan(e/2 + (f*x)/2))/((f*(c + d)*(c + d - \tan(e/2 + (f*x)/2))^2*(c - d)))*(a*d^2 + b*c^2 - a*c*d - b*c*d) - (d*\text{atan}(((d*((32*\tan(e/2 + (f*x)/2)*(b^5*c^6 + 2*b^5*d^6 - a*b^4*c^6 - 4*a*b^4*d^6 - 2*b^5*c*d^5 - 2*b^5*c^5*d + 3*a^2*b^3*d^6 - a^3*b^2*d^6 - a^5*c^2*d^4 - 5*b^5*c^2*d^4 + 4*b^5*c^3*d^3 + 3*b^5*c^4*d^2 + 13*a*b^4*c^2*d^4 - 8*a*b^4*c^3*d^3 - 11*a*b^4*c^4*d^2 - 6*a^2*b^3*c*d^5 + 6*a^3*b^2*c*d^5 + 3*a^4*b*c^2*d^4 + 4*a^4*b*c^3*d^3 - 11*a^2*b^3*c^2*d^4 + 12*a^2*b^3*c^3*d^3 + 12*a^2*b^3*c^4*d^2 + a^3*b^2*c^2*d^4 - 12*a^3*b^2*c^3*d^3 - 4*a^3*b^2*c^4*d^2 + 4*a*b^4*c*d^5 + 2*a*b^4*c^5*d - 2*a^4*b*c*d^5)))/(a^2*d^5 - b^2*c^5 + a^2*c*d^4 - b^2*c^4*d - a^2*c^2*d^3 - a^2*c^3*d^2 + b^2*c^2*d^3 + b^2*c^3*d^2 - 2*a*b*c*d^4 + 2*a*b*c^4*d - 2*a*b*c^2*d^3 + 2*a*b*c^3*d^2) + (d*((32*(2*a*b^6*c^9 - b^7*c^9 + a^6*b*d^9 + a^7*c*d^8 + 2*b^7*c^8*d - a^2*b^5*c^9 + a^4*b^3*d^9 - 2*a^5*b^2*d^9 - a^7*c^2*d^7 - a^7*c^3*d^6 + a^7*c^4*d^5 + b^7*c^4*d^5 - 3*b^7*c^6*d^3 + b^7*c^7*d^2 - 4*a*b^6*c^3*d^6 - 2*a*b^6*c^4*d^5 + 13*a*b^6*c^5*d^4 + a*b^6*c^6*d^3 - 11*a*b^6*c^7*d^2 - 8*a^2*b^5*c^8*d - 4*a^3*b^4*c*d^8 + 5*a^3*b^4*c^8*d + 8*a^4*b^3*c*d^8 - 3*a^5*b^2*c*d^8 - 5*a^6*b*c^2*d^7 + 7*a^6*b*c^3*d^6 + 4*a^6*b*c^4*d^5 - 5*a^6*b*c^5*d^4 + 6*a^2*b^5*c^2*d^7 + 8*a^2*b^5*c^3*d^6 - 21*a^2*b^5*c^4*d^5 - 16*a^2*b^5*c^5*d^4 + 23*a^2*b^5*c^6*d^3 + 9*a^2*b^5*c^7*d^2 - 12*a^3*b^4*c^2*d^7 + 14*a^3*b^4*c^3*d^6 + 34*a^3*b^4*c^4*d^5 - 21*a^3*b^4*c^5*d^4 - 27*a^3*b^4*c^6*d^3 + 11*a^3*b^4*c^7*d^2 ...$$

3.258 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+b \sec(e+fx))^2} dx$

3.258.1 Optimal result 1894
 3.258.2 Mathematica [B] (verified) 1895
 3.258.3 Rubi [A] (verified) 1896
 3.258.4 Maple [A] (verified) 1898
 3.258.5 Fracas [F(-1)] 1899
 3.258.6 Sympy [F] 1899
 3.258.7 Maxima [F(-2)] 1900
 3.258.8 Giac [B] (verification not implemented) 1900
 3.258.9 Mupad [B] (verification not implemented) 1901

3.258.1 Optimal result

Integrand size = 31, antiderivative size = 379

$$\begin{aligned} & \int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+b \sec(e+fx))^2} dx \\ &= \frac{d^4(5bc-2ad)\operatorname{arctanh}(\sin(e+fx))}{2b^3f} \\ &+ \frac{d^2(10b^3c^3-20ab^2c^2d+15a^2bcd^2-4a^3d^3)\operatorname{arctanh}(\sin(e+fx))}{b^5f} \\ &+ \frac{2(bc-ad)^5\operatorname{arctanh}\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{a(a-b)^{3/2}b^3(a+b)^{3/2}f} \\ &+ \frac{2(bc-ad)^4(bc+4ad)\operatorname{arctanh}\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{a\sqrt{a-b}b^5\sqrt{a+b}f} - \frac{(bc-ad)^5\sin(e+fx)}{b^4(a^2-b^2)f(b+a\cos(e+fx))} \\ &+ \frac{d^5\tan(e+fx)}{b^2f} + \frac{d^3(10b^2c^2-10abcd+3a^2d^2)\tan(e+fx)}{b^4f} \\ &+ \frac{d^4(5bc-2ad)\sec(e+fx)\tan(e+fx)}{2b^3f} + \frac{d^5\tan^3(e+fx)}{3b^2f} \end{aligned}$$

output $\frac{1}{2}d^4(-2ad+5bc)\operatorname{arctanh}(\sin(fx+e))/b^3/f+d^2(-4a^3d^3+15a^2bc^2d^2-20ab^2c^2d+10b^3c^3)\operatorname{arctanh}(\sin(fx+e))/b^5/f+2(-ad+bc)^5\operatorname{arctanh}((a-b)^{1/2}\tan(1/2fx+1/2e)/(a+b)^{1/2})/a(a-b)^{3/2}/b^3/(a+b)^{3/2}/f-(-ad+bc)^5\sin(fx+e)/b^4/(a^2-b^2)/f/(b+a\cos(fx+e))+2(-ad+bc)^4(4ad+bc)\operatorname{arctanh}((a-b)^{1/2}\tan(1/2fx+1/2e)/(a+b)^{1/2})/a/b^5/f/(a-b)^{1/2}/(a+b)^{1/2}+d^5\tan(fx+e)/b^2/f+d^3(3a^2d^2-10ab^2c^2d+10b^2c^2)\tan(fx+e)/b^4/f+1/2d^4(-2ad+5bc)\sec(fx+e)\tan(fx+e)/b^3/f+1/3d^5\tan(fx+e)^3/b^2/f$

3.258.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 784 vs. $2(379) = 758$.

Time = 10.09 (sec) , antiderivative size = 784, normalized size of antiderivative = 2.07

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+b\sec(e+fx))^2} dx$$

$$= \frac{(b+a\cos(e+fx))(c+d\sec(e+fx))^5 \left(-\frac{24(bc-ad)^4(abc+4a^2d-5b^2d)\operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} \right) \cos^3(e+fx)(b+a\cos(e+fx))}{(a+b\sec(e+fx))^2}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^5)/(a + b*Sec[e + f*x])^2,x]`

output

```
((b + a*cos[e + f*x])*(c + d*sec[e + f*x])^5*((-24*(b*c - a*d)^4*(a*b*c +
4*a^2*d - 5*b^2*d)*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]]*Co
s[e + f*x]^3*(b + a*cos[e + f*x]))/(a^2 - b^2)^(3/2) + 6*d^2*(-30*a^2*b*c*
d^2 + 8*a^3*d^3 - 5*b^3*c*(4*c^2 + d^2) + 2*a*b^2*d*(20*c^2 + d^2))*Cos[e
+ f*x]^3*(b + a*cos[e + f*x])*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + 6
*d^2*(30*a^2*b*c*d^2 - 8*a^3*d^3 + 5*b^3*c*(4*c^2 + d^2) - 2*a*b^2*d*(20*c
^2 + d^2))*Cos[e + f*x]^3*(b + a*cos[e + f*x])*Log[Cos[(e + f*x)/2] + Sin[
(e + f*x)/2]] + (b*(-60*a^2*b^3*c^2*d^3 + 60*b^5*c^2*d^3 + 45*a^3*b^2*c*d^
4 - 45*a*b^4*c*d^4 - 12*a^4*b*d^5 + 4*a^2*b^3*d^5 + 8*b^5*d^5 + (135*a^4*b
*c*d^4 - 36*a^5*d^5 + 30*a^2*b^3*c*d^2*(3*c^2 - 4*d^2) + a^3*b^2*d^3*(-180
*c^2 + 29*d^2) + a*b^4*d*(-45*c^4 + 90*c^2*d^2 - 2*d^4) + b^5*(9*c^5 + 30*
c*d^4))*Cos[e + f*x] + b*(-a^2 + b^2)*d^3*(-45*a*b*c*d + 12*a^2*d^2 + 4*b^
2*(15*c^2 + d^2))*Cos[2*(e + f*x)] + 3*b^5*c^5*Cos[3*(e + f*x)] - 15*a*b^4
*c^4*d*Cos[3*(e + f*x)] + 30*a^2*b^3*c^3*d^2*Cos[3*(e + f*x)] - 60*a^3*b^2
*c^2*d^3*Cos[3*(e + f*x)] + 30*a*b^4*c^2*d^3*Cos[3*(e + f*x)] + 45*a^4*b*c
*d^4*Cos[3*(e + f*x)] - 30*a^2*b^3*c*d^4*Cos[3*(e + f*x)] - 12*a^5*d^5*Cos
[3*(e + f*x)] + 7*a^3*b^2*d^5*Cos[3*(e + f*x)] + 2*a*b^4*d^5*Cos[3*(e + f
x)])*Sin[e + f*x])/(-a^2 + b^2)))/(12*b^5*f*(d + c*cos[e + f*x])^5*(a + b*
Sec[e + f*x])^2)
```

3.258.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4476, 3042, 3431, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+b\sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^5}{(a+b\csc\left(e+fx+\frac{\pi}{2}\right))^2} dx$$

↓ 4476

$$\int \frac{\sec^4(e+fx)(c\cos(e+fx)+d)^5}{(a\cos(e+fx)+b)^2} dx$$

↓ 3042

3.258. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+b\sec(e+fx))^2} dx$

$$\int \frac{(c \sin(e + fx + \frac{\pi}{2}) + d)^5}{\sin(e + fx + \frac{\pi}{2})^4 (a \sin(e + fx + \frac{\pi}{2}) + b)^2} dx$$

↓ 3431

$$\int \left(\frac{d^3(3a^2d^2 - 10abcd + 10b^2c^2) \sec^2(e + fx)}{b^4} + \frac{d^2(-4a^3d^3 + 15a^2bcd^2 - 20ab^2c^2d + 10b^3c^3) \sec(e + fx)}{b^5} + \frac{(a}{ab^5}$$

↓ 2009

$$\begin{aligned} & \frac{d^3(3a^2d^2 - 10abcd + 10b^2c^2) \tan(e + fx)}{b^4 f} - \frac{(bc - ad)^5 \sin(e + fx)}{b^4 f (a^2 - b^2) (a \cos(e + fx) + b)} + \\ & \frac{d^2(-4a^3d^3 + 15a^2bcd^2 - 20ab^2c^2d + 10b^3c^3) \operatorname{arctanh}(\sin(e + fx))}{b^5 f} + \\ & \frac{2(bc - ad)^4(4ad + bc) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{ab^5 f \sqrt{a-b} \sqrt{a+b}} + \frac{d^4(5bc - 2ad) \operatorname{arctanh}(\sin(e + fx))}{2b^3 f} + \\ & \frac{2(bc - ad)^5 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{ab^3 f (a-b)^{3/2} (a+b)^{3/2}} + \frac{d^4(5bc - 2ad) \tan(e + fx) \sec(e + fx)}{2b^3 f} + \\ & \frac{d^5 \tan^3(e + fx)}{3b^2 f} + \frac{d^5 \tan(e + fx)}{b^2 f} \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^5)/(a + b*Sec[e + f*x])^2,x]`

output `(d^4*(5*b*c - 2*a*d)*ArcTanh[Sin[e + f*x]]/(2*b^3*f) + (d^2*(10*b^3*c^3 - 20*a*b^2*c^2*d + 15*a^2*b*c*d^2 - 4*a^3*d^3)*ArcTanh[Sin[e + f*x]]/(b^5*f) + (2*(b*c - a*d)^5*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(a*(a - b)^(3/2)*b^3*(a + b)^(3/2)*f) + (2*(b*c - a*d)^4*(b*c + 4*a*d)*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*b^5*Sqrt[a + b]*f) - ((b*c - a*d)^5*Sin[e + f*x])/(b^4*(a^2 - b^2)*f*(b + a*Cos[e + f*x])) + (d^5*Tan[e + f*x])/(b^2*f) + (d^3*(10*b^2*c^2 - 10*a*b*c*d + 3*a^2*d^2)*Tan[e + f*x])/(b^4*f) + (d^4*(5*b*c - 2*a*d)*Sec[e + f*x]*Tan[e + f*x])/(2*b^3*f) + (d^5*Tan[e + f*x]^3)/(3*b^2*f)`

3.258.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3431 Int[((g_.)*sin[(e_.) + (f_.)*(x_)])^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(g*sin[e + f*x])^p*(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && NeQ[b*c - a*d, 0] && (IntegersQ[m, n] || IntegersQ[m, p] || IntegersQ[n, p]) && NeQ[p, 2]
```

```
rule 4476 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_)*((csc[(e_.) + (f_.)*(x_)])*(d_.) + (c_.))^(n_), x_Symbol] := Simp[1/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.258.4 Maple [A] (verified)

Time = 2.51 (sec) , antiderivative size = 689, normalized size of antiderivative = 1.82

method	result
derivativedivides	$\frac{\frac{d^5}{3b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^3} - \frac{d^2(8a^3d^3 - 30a^2bcd^2 + 40ab^2c^2d + 2ab^2d^3 - 20b^3c^3 - 5cd^2b^3) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2b^5}}{2b^4}$
default	$\frac{\frac{d^5}{3b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^3} - \frac{d^2(8a^3d^3 - 30a^2bcd^2 + 40ab^2c^2d + 2ab^2d^3 - 20b^3c^3 - 5cd^2b^3) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2b^5}}{2b^4}$
risch	Expression too large to display

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+b*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

3.258.
$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+b\sec(e+fx))^2} dx$$

output
$$\frac{1}{f} \left(-\frac{1}{3} d^5 / b^2 / (\tan(1/2 f x + 1/2 e) + 1)^3 - \frac{1}{2} d^2 (8 a^3 d^3 - 30 a^2 b c d^2 + 40 a b^2 c^2 d + 2 a^2 b^2 d^3 - 20 b^3 c^3 - 5 b^3 c d^2) / b^5 \ln(\tan(1/2 f x + 1/2 e) + 1) - \frac{1}{2} d^3 (6 a^2 d^2 - 20 a b c d + 2 a b d^2 + 20 b^2 c^2 - 5 b^2 c d + 2 b^2 d^2) / b^4 / (\tan(1/2 f x + 1/2 e) + 1) + \frac{1}{2} d^4 (2 a d - 5 b c + b d) / b^3 / (\tan(1/2 f x + 1/2 e) + 1)^2 - \frac{1}{3} d^5 / b^2 / (\tan(1/2 f x + 1/2 e) - 1)^3 + \frac{1}{2} d^2 (8 a^3 d^3 - 30 a^2 b c d^2 + 40 a b^2 c^2 d + 2 a^2 b^2 d^3 - 20 b^3 c^3 - 5 b^3 c d^2) / b^5 \ln(\tan(1/2 f x + 1/2 e) - 1) - \frac{1}{2} d^3 (6 a^2 d^2 - 20 a b c d + 2 a b d^2 + 20 b^2 c^2 - 5 b^2 c d + 2 b^2 d^2) / b^4 / (\tan(1/2 f x + 1/2 e) - 1) - \frac{1}{2} d^4 (2 a d - 5 b c + b d) / b^3 / (\tan(1/2 f x + 1/2 e) - 1)^2 - \frac{2}{b^5} (b (a^5 d^5 - 5 a^4 b c d^4 + 10 a^3 b^2 c^2 d^3 - 10 a^2 b^3 c^3 d^2 + 5 a b^4 c^4 d - b^5 c^5) / (a^2 - b^2) \tan(1/2 f x + 1/2 e) / (\tan(1/2 f x + 1/2 e)^2 a - \tan(1/2 f x + 1/2 e)^2 b - a - b) - (4 a^6 d^5 - 15 a^5 b c d^4 + 20 a^4 b^2 c^2 d^3 - 5 a^4 b^2 d^5 - 10 a^3 b^3 c^3 d^2 + 20 a^3 b^3 c d^4 - 30 a^2 b^4 c^2 d^3 + a b^5 c^5 + 20 a b^5 c^3 d^2 - 5 b^6 c^4 d) / (a - b) / (a + b) / ((a - b) * (a + b))^{1/2} \operatorname{arctanh}((a - b) \tan(1/2 f x + 1/2 e) / ((a - b) * (a + b))^{1/2}) \right)$$

3.258.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + b \sec(e + fx))^2} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+b*sec(f*x+e))^2,x, algorithm="fricas")`

output Timed out

3.258.6 Sympy [F]

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^5}{(a + b \sec(e + fx))^2} dx = \int \frac{(c + d \sec(e + fx))^5 \sec(e + fx)}{(a + b \sec(e + fx))^2} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**5/(a+b*sec(f*x+e))**2,x)`

output `Integral((c + d*sec(e + f*x))**5*sec(e + f*x)/(a + b*sec(e + f*x))**2, x)`

3.258.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+b\sec(e+fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+b*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de`

3.258.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 857 vs. 2(355) = 710.

Time = 0.45 (sec) , antiderivative size = 857, normalized size of antiderivative = 2.26

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+b\sec(e+fx))^2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^5/(a+b*sec(f*x+e))^2,x, algorithm="giac")`

```

output -1/6*(12*(a*b^5*c^5 - 5*b^6*c^4*d - 10*a^3*b^3*c^3*d^2 + 20*a*b^5*c^3*d^2
+ 20*a^4*b^2*c^2*d^3 - 30*a^2*b^4*c^2*d^3 - 15*a^5*b*c*d^4 + 20*a^3*b^3*c*
d^4 + 4*a^6*d^5 - 5*a^4*b^2*d^5)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*a
- 2*b) + arctan((a*tan(1/2*f*x + 1/2*e) - b*tan(1/2*f*x + 1/2*e))/sqrt(-a
^2 + b^2)))/((a^2*b^5 - b^7)*sqrt(-a^2 + b^2)) - 12*(b^5*c^5*tan(1/2*f*x +
1/2*e) - 5*a*b^4*c^4*d*tan(1/2*f*x + 1/2*e) + 10*a^2*b^3*c^3*d^2*tan(1/2*
f*x + 1/2*e) - 10*a^3*b^2*c^2*d^3*tan(1/2*f*x + 1/2*e) + 5*a^4*b*c*d^4*tan
(1/2*f*x + 1/2*e) - a^5*d^5*tan(1/2*f*x + 1/2*e))/((a^2*b^4 - b^6)*(a*tan(
1/2*f*x + 1/2*e)^2 - b*tan(1/2*f*x + 1/2*e)^2 - a - b)) - 3*(20*b^3*c^3*d^
2 - 40*a*b^2*c^2*d^3 + 30*a^2*b*c*d^4 + 5*b^3*c*d^4 - 8*a^3*d^5 - 2*a*b^2*
d^5)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/b^5 + 3*(20*b^3*c^3*d^2 - 40*a*b^2
*c^2*d^3 + 30*a^2*b*c*d^4 + 5*b^3*c*d^4 - 8*a^3*d^5 - 2*a*b^2*d^5)*log(abs
(tan(1/2*f*x + 1/2*e) - 1))/b^5 + 2*(60*b^2*c^2*d^3*tan(1/2*f*x + 1/2*e)^5
- 60*a*b*c*d^4*tan(1/2*f*x + 1/2*e)^5 - 15*b^2*c*d^4*tan(1/2*f*x + 1/2*e)
^5 + 18*a^2*d^5*tan(1/2*f*x + 1/2*e)^5 + 6*a*b*d^5*tan(1/2*f*x + 1/2*e)^5
+ 6*b^2*d^5*tan(1/2*f*x + 1/2*e)^5 - 120*b^2*c^2*d^3*tan(1/2*f*x + 1/2*e)^
3 + 120*a*b*c*d^4*tan(1/2*f*x + 1/2*e)^3 - 36*a^2*d^5*tan(1/2*f*x + 1/2*e)
^3 - 4*b^2*d^5*tan(1/2*f*x + 1/2*e)^3 + 60*b^2*c^2*d^3*tan(1/2*f*x + 1/2*e)
) - 60*a*b*c*d^4*tan(1/2*f*x + 1/2*e) + 15*b^2*c*d^4*tan(1/2*f*x + 1/2*e)
+ 18*a^2*d^5*tan(1/2*f*x + 1/2*e) - 6*a*b*d^5*tan(1/2*f*x + 1/2*e) + 6*...

```

3.258.9 Mupad [B] (verification not implemented)

Time = 28.86 (sec) , antiderivative size = 17256, normalized size of antiderivative = 45.53

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^5}{(a+b\sec(e+fx))^2} dx = \text{Too large to display}$$

```

input int((c + d/cos(e + f*x))^5/(cos(e + f*x)*(a + b/cos(e + f*x))^2),x)

```

output $(\operatorname{atan}(\frac{((8 \tan(e/2 + (f*x)/2) * (128*a^{12}*d^{10} - 128*a^{11}*b*d^{10} + 4*a^2*b^{10}*c^{10} + 4*a^2*b^{10}*d^{10} - 8*a^3*b^9*d^{10} + 28*a^4*b^8*d^{10} - 48*a^5*b^7*d^{10} + 28*a^6*b^6*d^{10} - 8*a^7*b^5*d^{10} + 8*a^8*b^4*d^{10} + 192*a^9*b^3*d^{10} - 192*a^{10}*b^2*d^{10} + 25*b^{12}*c^2*d^8 + 200*b^{12}*c^4*d^6 + 400*b^{12}*c^6*d^4 + 100*b^{12}*c^8*d^2 - 50*a*b^{11}*c^2*d^8 - 480*a*b^{11}*c^3*d^7 - 400*a*b^{11}*c^4*d^6 - 1600*a*b^{11}*c^5*d^5 - 800*a*b^{11}*c^6*d^4 - 800*a*b^{11}*c^7*d^3 + 40*a^2*b^{10}*c*d^9 - 180*a^3*b^9*c*d^9 + 320*a^4*b^8*c*d^9 - 260*a^5*b^7*c*d^9 + 200*a^6*b^6*c*d^9 - 140*a^7*b^5*c*d^9 - 1520*a^8*b^4*c*d^9 + 1520*a^9*b^3*c*d^9 + 960*a^{10}*b^2*c*d^9 + 435*a^2*b^{10}*c^2*d^8 + 960*a^2*b^{10}*c^3*d^7 + 2600*a^2*b^{10}*c^4*d^6 + 3200*a^2*b^{10}*c^5*d^5 + 2400*a^2*b^{10}*c^6*d^4 + 160*a^2*b^{10}*c^8*d^2 - 820*a^3*b^9*c^2*d^8 - 2240*a^3*b^9*c^3*d^7 - 4800*a^3*b^9*c^4*d^6 - 4000*a^3*b^9*c^5*d^5 + 1600*a^3*b^9*c^6*d^4 + 160*a^3*b^9*c^7*d^3 + 1055*a^4*b^8*c^2*d^8 + 3520*a^4*b^8*c^3*d^7 + 4000*a^4*b^8*c^4*d^6 - 6400*a^4*b^8*c^5*d^5 - 2640*a^4*b^8*c^6*d^4 - 80*a^4*b^8*c^8*d^2 - 1290*a^5*b^7*c^2*d^8 - 2400*a^5*b^7*c^3*d^7 + 10800*a^5*b^7*c^4*d^6 + 7760*a^5*b^7*c^5*d^5 - 800*a^5*b^7*c^6*d^4 + 160*a^5*b^7*c^7*d^3 + 825*a^6*b^6*c^2*d^8 - 9920*a^6*b^6*c^3*d^7 - 11560*a^6*b^6*c^4*d^6 + 3200*a^6*b^6*c^5*d^5 + 680*a^6*b^6*c^6*d^4 + 5240*a^7*b^5*c^2*d^8 + 10080*a^7*b^5*c^3*d^7 - 5600*a^7*b^5*c^4*d^6 - 3168*a^7*b^5*c^5*d^5 - 5240*a^8*b^4*c^2*d^8 + 5440*a^8*b^4*c^3*d^7 + 5600*a^8*b^4*c^4*d^6 - 3080*a^9*b^3*c^2*d^8 \dots$

3.258. $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^5}{(a+b \sec(e+fx))^2} dx$

3.259
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+b \sec(e+fx))^2} dx$$

3.259.1 Optimal result 1903
 3.259.2 Mathematica [A] (verified) 1904
 3.259.3 Rubi [A] (verified) 1904
 3.259.4 Maple [A] (verified) 1906
 3.259.5 Fricas [B] (verification not implemented) 1907
 3.259.6 Sympy [F] 1908
 3.259.7 Maxima [F(-2)] 1909
 3.259.8 Giac [B] (verification not implemented) 1909
 3.259.9 Mupad [B] (verification not implemented) 1910

3.259.1 Optimal result

Integrand size = 31, antiderivative size = 297

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^4}{(a+b \sec(e+fx))^2} dx = \frac{d^4 \operatorname{arctanh}(\sin(e+fx))}{2b^2 f} + \frac{d^2(6b^2c^2 - 8abcd + 3a^2d^2) \operatorname{arctanh}(\sin(e+fx))}{b^4 f} + \frac{2(bc - ad)^4 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{a(a-b)^{3/2}b^2(a+b)^{3/2}f} + \frac{2(bc - ad)^3(bc + 3ad) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{a\sqrt{a-b}b^4\sqrt{a+b}f} - \frac{(bc - ad)^4 \sin(e+fx)}{b^3(a^2 - b^2)f(b + a \cos(e+fx))} + \frac{2d^3(2bc - ad) \tan(e+fx)}{b^3 f} + \frac{d^4 \sec(e+fx) \tan(e+fx)}{2b^2 f}$$

output

```
1/2*d^4*arctanh(sin(f*x+e))/b^2/f+d^2*(3*a^2*d^2-8*a*b*c*d+6*b^2*c^2)*arctanh(sin(f*x+e))/b^4/f+2*(-a*d+b*c)^4*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e))/(a+b)^(1/2)/a/(a-b)^(3/2)/b^2/(a+b)^(3/2)/f-(-a*d+b*c)^4*sin(f*x+e)/b^3/(a^2-b^2)/f/(b+a*cos(f*x+e))+2*(-a*d+b*c)^3*(3*a*d+b*c)*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e))/(a+b)^(1/2))/a/b^4/f/(a-b)^(1/2)/(a+b)^(1/2)+2*d^3*(-a*d+2*b*c)*tan(f*x+e)/b^3/f+1/2*d^4*sec(f*x+e)*tan(f*x+e)/b^2/f
```


3.259.2 Mathematica [A] (verified)

Time = 5.40 (sec) , antiderivative size = 511, normalized size of antiderivative = 1.72

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+b\sec(e+fx))^2} dx$$

$$= \frac{\cos^2(e+fx)(b+a\cos(e+fx))(c+d\sec(e+fx))^4 \left(\frac{8(-bc+ad)^3(abc+3a^2d-4b^2d)\operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} \right) (b+...)}{...}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + b*Sec[e + f*x])^2,x]`

output `(Cos[e + f*x]^2*(b + a*Cos[e + f*x])*(c + d*Sec[e + f*x])^4*((8*(-(b*c) + a*d)^3*(a*b*c + 3*a^2*d - 4*b^2*d)*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]]*(b + a*Cos[e + f*x]))/(a^2 - b^2)^(3/2) - 2*d^2*(-16*a*b*c*d + 6*a^2*d^2 + b^2*(12*c^2 + d^2))*(b + a*Cos[e + f*x])*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + 2*d^2*(-16*a*b*c*d + 6*a^2*d^2 + b^2*(12*c^2 + d^2))*(b + a*Cos[e + f*x])*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] + (b^2*d^4*(b + a*Cos[e + f*x]))/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2 + (8*b*d^3*(2*b*c - a*d)*(b + a*Cos[e + f*x])*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]) - (b^2*d^4*(b + a*Cos[e + f*x]))/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])^2 + (8*b*d^3*(2*b*c - a*d)*(b + a*Cos[e + f*x])*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2]) + (4*b*(b*c - a*d)^4*Sin[e + f*x])/((-a + b)*(a + b)))/(4*b^4*f*(d + c*Cos[e + f*x])^4*(a + b*Sec[e + f*x])^2)`

3.259.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4476, 3042, 3431, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+b\sec(e+fx))^2} dx$$

↓ 3042

3.259. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+b\sec(e+fx))^2} dx$

$$\begin{aligned}
& \int \frac{\csc\left(e + fx + \frac{\pi}{2}\right) \left(c + d \csc\left(e + fx + \frac{\pi}{2}\right)\right)^4}{\left(a + b \csc\left(e + fx + \frac{\pi}{2}\right)\right)^2} dx \\
& \quad \downarrow 4476 \\
& \int \frac{\sec^3(e + fx)(c \cos(e + fx) + d)^4}{(a \cos(e + fx) + b)^2} dx \\
& \quad \downarrow 3042 \\
& \int \frac{\left(c \sin\left(e + fx + \frac{\pi}{2}\right) + d\right)^4}{\sin\left(e + fx + \frac{\pi}{2}\right)^3 \left(a \sin\left(e + fx + \frac{\pi}{2}\right) + b\right)^2} dx \\
& \quad \downarrow 3431 \\
& \int \left(\frac{d^2(3a^2d^2 - 8abcd + 6b^2c^2) \sec(e + fx)}{b^4} - \frac{(ad - bc)^3(3ad + bc)}{ab^4(a \cos(e + fx) + b)} + \frac{2d^3(2bc - ad) \sec^2(e + fx)}{b^3} - \frac{(ad - bc)^3}{ab^3(a \cos(e + fx) + b)} \right) dx \\
& \quad \downarrow 2009 \\
& \frac{d^2(3a^2d^2 - 8abcd + 6b^2c^2) \operatorname{arctanh}(\sin(e + fx))}{b^4 f} - \frac{(bc - ad)^4 \sin(e + fx)}{b^3 f (a^2 - b^2) (a \cos(e + fx) + b)} + \\
& \frac{2(bc - ad)^3(3ad + bc) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e + fx)\right)}{\sqrt{a+b}}\right)}{ab^4 f \sqrt{a-b} \sqrt{a+b}} + \frac{2(bc - ad)^4 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e + fx)\right)}{\sqrt{a+b}}\right)}{ab^2 f (a - b)^{3/2} (a + b)^{3/2}} + \\
& \frac{2d^3(2bc - ad) \tan(e + fx)}{b^3 f} + \frac{d^4 \operatorname{arctanh}(\sin(e + fx))}{2b^2 f} + \frac{d^4 \tan(e + fx) \sec(e + fx)}{2b^2 f}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^4)/(a + b*Sec[e + f*x])^2,x]`

output `(d^4*ArcTanh[Sin[e + f*x]])/(2*b^2*f) + (d^2*(6*b^2*c^2 - 8*a*b*c*d + 3*a^2*d^2)*ArcTanh[Sin[e + f*x]])/(b^4*f) + (2*(b*c - a*d)^4*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(a*(a - b)^(3/2)*b^2*(a + b)^(3/2)*f) + (2*(b*c - a*d)^3*(b*c + 3*a*d)*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*b^4*Sqrt[a + b]*f) - ((b*c - a*d)^4*Sin[e + f*x])/(b^3*(a^2 - b^2)*f*(b + a*Cos[e + f*x])) + (2*d^3*(2*b*c - a*d)*Tan[e + f*x])/(b^3*f) + (d^4*Sec[e + f*x]*Tan[e + f*x])/(2*b^2*f)`

3.259.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3431 Int[((g_)*sin[(e_)+(f_)*(x_)])^(p_)*((a_)+(b_)*sin[(e_)+(f_)*(x_)])^(m_)*((c_)+(d_)*sin[(e_)+(f_)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(g*sin[e+f*x])^p*(a+b*sin[e+f*x])^m*(c+d*sin[e+f*x])^n, x], x] /; FreeQ[{a,b,c,d,e,f,g,n,p}, x] && NeQ[b*c-a*d, 0] && (IntegersQ[m,n] || IntegersQ[m,p] || IntegersQ[n,p]) && NeQ[p, 2]
```

```
rule 4476 Int[(csc[(e_)+(f_)*(x_)])*(g_)^(p_)*(csc[(e_)+(f_)*(x_)])*(b_)+(a_)^(m_)*(csc[(e_)+(f_)*(x_)])*(d_)+(c_)^(n_), x_Symbol] := Simp[1/g^(m+n) Int[(g*Csc[e+f*x])^(m+n+p)*(b+a*Sin[e+f*x])^m*(d+c*Sin[e+f*x])^n, x], x] /; FreeQ[{a,b,c,d,e,f,g,p}, x] && NeQ[b*c-a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.259.4 Maple [A] (verified)

Time = 1.72 (sec) , antiderivative size = 465, normalized size of antiderivative = 1.57

method	result
derivativedivides	$\frac{\frac{d^4}{2b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{d^2(6a^2d^2 - 16abcd + 12b^2c^2 + b^2d^2) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2b^4} + \frac{d^3(4ad - 8bc + bd)}{2b^3 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} + \frac{2b(a^4d^4 - 4a^3bcd^3 + 6a^2b^2cd^2 - 4ab^3cd + b^4d^2)}{(a^2 - b^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2}$
default	$\frac{\frac{d^4}{2b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2} + \frac{d^2(6a^2d^2 - 16abcd + 12b^2c^2 + b^2d^2) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{2b^4} + \frac{d^3(4ad - 8bc + bd)}{2b^3 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)} + \frac{2b(a^4d^4 - 4a^3bcd^3 + 6a^2b^2cd^2 - 4ab^3cd + b^4d^2)}{(a^2 - b^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)^2}$
risch	Expression too large to display

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

3.259.
$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+b\sec(e+fx))^2} dx$$

output
$$\frac{1}{f} \left(-\frac{1}{2} d^4 / b^2 / (\tan(1/2 f x + 1/2 e) + 1)^2 + \frac{1}{2} d^2 (6 a^2 d^2 - 16 a b c d + 12 b^2 c^2 + b^2 d^2) / b^4 \ln(\tan(1/2 f x + 1/2 e) + 1) + \frac{1}{2} d^3 (4 a d - 8 b c + b d) / b^3 / (\tan(1/2 f x + 1/2 e) + 1) + \frac{2}{b^4} (b (a^4 d^4 - 4 a^3 b c d^3 + 6 a^2 b^2 c^2 d^2 - 4 a a b^3 c^3 d + b^4 c^4) / (a^2 - b^2) \tan(1/2 f x + 1/2 e) / (\tan(1/2 f x + 1/2 e)^2 a - \tan(1/2 f x + 1/2 e)^2 b - a - b) - (3 a^5 d^4 - 8 a^4 b c d^3 + 6 a^3 b^2 c^2 d^2 - 4 a^3 b^2 d^4 + 12 a^2 b^3 c d^3 - a b^4 c^4 - 12 a b^4 c^2 d^2 + 4 b^5 c^3 d) / ((a - b) / (a + b) / ((a - b) * (a + b))^{1/2} * \operatorname{arctanh}((a - b) * \tan(1/2 f x + 1/2 e) / ((a - b) * (a + b))^{1/2})) + \frac{1}{2} d^4 / b^2 / (\tan(1/2 f x + 1/2 e) - 1)^2 - \frac{1}{2} d^2 (6 a^2 d^2 - 16 a b c d + 12 b^2 c^2 + b^2 d^2) / b^4 \ln(\tan(1/2 f x + 1/2 e) - 1) + \frac{1}{2} d^3 (4 a d - 8 b c + b d) / b^3 / (\tan(1/2 f x + 1/2 e) - 1) \right)$$

3.259.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 934 vs. $2(275) = 550$.

Time = 170.55 (sec) , antiderivative size = 1925, normalized size of antiderivative = 6.48

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{(a + b \sec(e + fx))^2} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e))^2,x, algorithm="fracas")`

output

```

[-1/4*(2*((a^2*b^4*c^4 - 4*a*b^5*c^3*d - 6*(a^4*b^2 - 2*a^2*b^4)*c^2*d^2 +
4*(2*a^5*b - 3*a^3*b^3)*c*d^3 - (3*a^6 - 4*a^4*b^2)*d^4)*cos(f*x + e)^3 +
(a*b^5*c^4 - 4*b^6*c^3*d - 6*(a^3*b^3 - 2*a*b^5)*c^2*d^2 + 4*(2*a^4*b^2 -
3*a^2*b^4)*c*d^3 - (3*a^5*b - 4*a^3*b^3)*d^4)*cos(f*x + e)^2)*sqrt(a^2 -
b^2)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 - 2*sqrt(a^2 -
b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2
+ 2*a*b*cos(f*x + e) + b^2)) - ((12*(a^5*b^2 - 2*a^3*b^4 + a*b^6)*c^2*d^2
- 16*(a^6*b - 2*a^4*b^3 + a^2*b^5)*c*d^3 + (6*a^7 - 11*a^5*b^2 + 4*a^3*b^
4 + a*b^6)*d^4)*cos(f*x + e)^3 + (12*(a^4*b^3 - 2*a^2*b^5 + b^7)*c^2*d^2 -
16*(a^5*b^2 - 2*a^3*b^4 + a*b^6)*c*d^3 + (6*a^6*b - 11*a^4*b^3 + 4*a^2*b^
5 + b^7)*d^4)*cos(f*x + e)^2)*log(sin(f*x + e) + 1) + ((12*(a^5*b^2 - 2*a^
3*b^4 + a*b^6)*c^2*d^2 - 16*(a^6*b - 2*a^4*b^3 + a^2*b^5)*c*d^3 + (6*a^7 -
11*a^5*b^2 + 4*a^3*b^4 + a*b^6)*d^4)*cos(f*x + e)^3 + (12*(a^4*b^3 - 2*a^
2*b^5 + b^7)*c^2*d^2 - 16*(a^5*b^2 - 2*a^3*b^4 + a*b^6)*c*d^3 + (6*a^6*b -
11*a^4*b^3 + 4*a^2*b^5 + b^7)*d^4)*cos(f*x + e)^2)*log(-sin(f*x + e) + 1)
- 2*((a^4*b^3 - 2*a^2*b^5 + b^7)*d^4 - 2*((a^2*b^5 - b^7)*c^4 - 4*(a^3*b^
4 - a*b^6)*c^3*d + 6*(a^4*b^3 - a^2*b^5)*c^2*d^2 - 4*(2*a^5*b^2 - 3*a^3*b^
4 + a*b^6)*c*d^3 + (3*a^6*b - 5*a^4*b^3 + 2*a^2*b^5)*d^4)*cos(f*x + e)^2 +
(8*(a^4*b^3 - 2*a^2*b^5 + b^7)*c*d^3 - 3*(a^5*b^2 - 2*a^3*b^4 + a*b^6)*d^
4)*cos(f*x + e))*sin(f*x + e))/((a^5*b^4 - 2*a^3*b^6 + a*b^8)*f*cos(f*x...

```

3.259.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+b\sec(e+fx))^2} dx = \int \frac{(c+d\sec(e+fx))^4 \sec(e+fx)}{(a+b\sec(e+fx))^2} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**4/(a+b*sec(f*x+e))**2,x)`

output `Integral((c + d*sec(e + f*x))**4*sec(e + f*x)/(a + b*sec(e + f*x))**2, x)`

3.259.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+b\sec(e+fx))^2} dx = \text{Exception raised: ValueError}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e))^2,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de
```

3.259.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 551 vs. 2(275) = 550.

Time = 0.39 (sec) , antiderivative size = 551, normalized size of antiderivative = 1.86

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^4}{(a+b\sec(e+fx))^2} dx = \frac{4(ab^4c^4 - 4b^5c^3d - 6a^3b^2c^2d^2 + 12ab^4c^2d^2 + 8a^4bcd^3 - 12a^2b^3cd^3 - 3a^5d^4 + 4a^3b^2d^4) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) - b}{\sqrt{-a^2 + b^2}} \right) \right)}{(a^2b^4 - b^6)\sqrt{-a^2 + b^2}}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^4/(a+b*sec(f*x+e))^2,x, algorithm="giac")
```

output

```
-1/2*(4*(a*b^4*c^4 - 4*b^5*c^3*d - 6*a^3*b^2*c^2*d^2 + 12*a*b^4*c^2*d^2 +
8*a^4*b*c*d^3 - 12*a^2*b^3*c*d^3 - 3*a^5*d^4 + 4*a^3*b^2*d^4)*(pi*floor(1/
2*(f*x + e)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*f*x + 1/2*e) - b*
tan(1/2*f*x + 1/2*e))/sqrt(-a^2 + b^2)))/((a^2*b^4 - b^6)*sqrt(-a^2 + b^2)
) - 4*(b^4*c^4*tan(1/2*f*x + 1/2*e) - 4*a*b^3*c^3*d*tan(1/2*f*x + 1/2*e) +
6*a^2*b^2*c^2*d^2*tan(1/2*f*x + 1/2*e) - 4*a^3*b*c*d^3*tan(1/2*f*x + 1/2*
e) + a^4*d^4*tan(1/2*f*x + 1/2*e))/((a^2*b^3 - b^5)*(a*tan(1/2*f*x + 1/2*e
)^2 - b*tan(1/2*f*x + 1/2*e)^2 - a - b)) - (12*b^2*c^2*d^2 - 16*a*b*c*d^3
+ 6*a^2*d^4 + b^2*d^4)*log(abs(tan(1/2*f*x + 1/2*e) + 1))/b^4 + (12*b^2*c^
2*d^2 - 16*a*b*c*d^3 + 6*a^2*d^4 + b^2*d^4)*log(abs(tan(1/2*f*x + 1/2*e) -
1))/b^4 + 2*(8*b*c*d^3*tan(1/2*f*x + 1/2*e)^3 - 4*a*d^4*tan(1/2*f*x + 1/2
*e)^3 - b*d^4*tan(1/2*f*x + 1/2*e)^3 - 8*b*c*d^3*tan(1/2*f*x + 1/2*e) + 4*
a*d^4*tan(1/2*f*x + 1/2*e) - b*d^4*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1
/2*e)^2 - 1)^2*b^3))/f
```

3.259.9 Mupad [B] (verification not implemented)

Time = 26.19 (sec) , antiderivative size = 12483, normalized size of antiderivative = 42.03

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^4}{(a + b \sec(e + fx))^2} dx = \text{Too large to display}$$

input `int((c + d/cos(e + f*x))^4/(cos(e + f*x)*(a + b/cos(e + f*x))^2),x)`

output $(\text{atan}(\frac{(8(2b^{15}d^4 - 4a^2b^{13}c^4 + 16b^{15}c^3d + 4a^2b^{13}c^4 + 4a^3b^{12}c^4 - 4a^4b^{11}c^4 + 6a^2b^{13}d^4 - 16a^3b^{12}d^4 - 14a^4b^{11}d^4 + 28a^5b^{10}d^4 + 6a^6b^9d^4 - 12a^7b^8d^4 + 24b^{15}c^2d^2 - 48a^2b^{14}c^2d^2 + 48a^2b^{13}cd^3 - 16a^2b^{13}c^3d + 48a^3b^{12}cd^3 + 16a^3b^{12}c^3d - 80a^4b^{11}cd^3 - 16a^5b^{10}cd^3 + 32a^6b^9cd^3 - 24a^2b^{13}c^2d^2 + 72a^3b^{12}c^2d^2 - 24a^5b^{10}c^2d^2 - 32a^2b^{14}cd^3 - 16a^2b^{14}c^3d))}{(ab^{11} + b^{12} - a^2b^{10} - a^3b^9) - (8\tan(e/2 + (f*x)/2)*(b^2(d^4/2 + 6c^2d^2) + 3a^2d^4 - 8a^2b^9cd^3)*(8a^2b^{13} - 8a^2b^{12} - 16a^3b^{11} + 16a^4b^{10} + 8a^5b^9 - 8a^6b^8))}{(b^4(a^2b^8 + b^9 - a^2b^7 - a^3b^6))}*(b^2(d^4/2 + 6c^2d^2) + 3a^2d^4 - 8a^2b^9cd^3))/b^4 - (8\tan(e/2 + (f*x)/2)*(72a^{10}d^8 + b^{10}d^8 - 2a^2b^9d^8 - 72a^9b^8d^8 + 4a^2b^8c^8 + 11a^2b^8d^8 - 20a^3b^7d^8 + 23a^4b^6d^8 - 26a^5b^5d^8 + 17a^6b^4d^8 + 120a^7b^3d^8 - 120a^8b^2d^8 + 24b^{10}c^2d^6 + 144b^{10}c^4d^4 + 64b^{10}c^6d^2 - 48a^2b^9c^2d^6 - 384a^2b^9c^3d^5 - 288a^2b^9c^4d^4 - 384a^2b^9c^5d^3 + 64a^2b^8c^2d^7 - 160a^3b^7cd^7 + 256a^4b^6cd^7 - 160a^5b^5cd^7 - 704a^6b^4cd^7 + 704a^7b^3cd^7 + 384a^8b^2cd^7 + 376a^2b^8c^2d^6 + 768a^2b^8c^3d^5 + 816a^2b^8c^4d^4 + 96a^2b^8c^6d^2 - 704a^3b^7c^2d^6 - 896a^3b^7c^3d^5 + 576a^3b^7c^4d^4 + 96a^3b^7c^5d^3 + 536a^4b^6c^2d^6 - 1536a^4b^6c^3d^5 + 1536a^4b^6c^4d^4 + 96a^4b^6c^5d^3 - 536a^4b^6c^6d^2 - 1536a^4b^6c^7d^1 + 1536a^4b^6c^8d^0))}{(a+b^2c^2d^2)^2}) dx$

3.260
$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+b \sec(e+fx))^2} dx$$

3.260.1 Optimal result 1912
 3.260.2 Mathematica [A] (verified) 1913
 3.260.3 Rubi [A] (verified) 1913
 3.260.4 Maple [A] (verified) 1915
 3.260.5 Fracas [B] (verification not implemented) 1916
 3.260.6 Sympy [F] 1916
 3.260.7 Maxima [F(-2)] 1917
 3.260.8 Giac [B] (verification not implemented) 1917
 3.260.9 Mupad [B] (verification not implemented) 1918

3.260.1 Optimal result

Integrand size = 31, antiderivative size = 228

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^3}{(a+b \sec(e+fx))^2} dx = \frac{d^2(3bc-2ad)\operatorname{arctanh}(\sin(e+fx))}{b^3 f} + \frac{2(bc-ad)^3 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{a(a-b)^{3/2} b(a+b)^{3/2} f} + \frac{2(bc-ad)^2(bc+2ad)\operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{a\sqrt{a-b} b^3 \sqrt{a+b} f} - \frac{(bc-ad)^3 \sin(e+fx)}{b^2(a^2-b^2) f(b+a \cos(e+fx))} + \frac{d^3 \tan(e+fx)}{b^2 f}$$

```
output d^2*(-2*a*d+3*b*c)*arctanh(sin(f*x+e))/b^3/f+2*(-a*d+b*c)^3*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/a/(a-b)^(3/2)/b/(a+b)^(3/2)/f-(-a*d+b*c)^3*sin(f*x+e)/b^2/(a^2-b^2)/f/(b+a*cos(f*x+e))+2*(-a*d+b*c)^2*(2*a*d+b*c)*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/a/b^3/f/(a-b)^(1/2)/(a+b)^(1/2)+d^3*tan(f*x+e)/b^2/f
```

3.260.2 Mathematica [A] (verified)

Time = 3.41 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.59

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+b\sec(e+fx))^2} dx$$

$$= \frac{\cos(e+fx)(b+a\cos(e+fx))(c+d\sec(e+fx))^3 \left(-\frac{2(bc-ad)^2(abc+2a^2d-3b^2d)\operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} \right)}{(a^2-b^2)^{3/2}}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + b*Sec[e + f*x])^2,x]`

output `(Cos[e + f*x]*(b + a*Cos[e + f*x])*(c + d*Sec[e + f*x])^3*((-2*(b*c - a*d)^2*(a*b*c + 2*a^2*d - 3*b^2*d)*ArcTanh[(-a + b)*Tan[(e + f*x)/2]]/Sqrt[a^2 - b^2])*(b + a*Cos[e + f*x]))/(a^2 - b^2)^(3/2) + d^2*(-3*b*c + 2*a*d)*(b + a*Cos[e + f*x])*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + d^2*(3*b*c - 2*a*d)*(b + a*Cos[e + f*x])*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] + (b*d^3*(b + a*Cos[e + f*x])*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]) + (b*d^3*(b + a*Cos[e + f*x])*Sin[(e + f*x)/2])/(Cos[(e + f*x)/2] + Sin[(e + f*x)/2]) + (b*(b*c - a*d)^3*Sin[e + f*x])/((-a + b)*(a + b)))/(b^3*f*(d + c*Cos[e + f*x])^3*(a + b*Sec[e + f*x])^2)`

3.260.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4476, 3042, 3431, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+b\sec(e+fx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^3}{(a+b\csc\left(e+fx+\frac{\pi}{2}\right))^2} dx$$

$$\downarrow \text{4476}$$

3.260. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+b\sec(e+fx))^2} dx$

$$\begin{aligned}
& \int \frac{\sec^2(e+fx)(c \cos(e+fx)+d)^3}{(a \cos(e+fx)+b)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(c \sin(e+fx+\frac{\pi}{2})+d)^3}{\sin(e+fx+\frac{\pi}{2})^2 (a \sin(e+fx+\frac{\pi}{2})+b)^2} dx \\
& \quad \downarrow \text{3431} \\
& \int \left(\frac{d^2(3bc-2ad) \sec(e+fx)}{b^3} + \frac{(ad-bc)^2(2ad+bc)}{ab^3(a \cos(e+fx)+b)} + \frac{(ad-bc)^3}{ab^2(a \cos(e+fx)+b)^2} + \frac{d^3 \sec^2(e+fx)}{b^2} \right) dx \\
& \quad \downarrow \text{2009} \\
& -\frac{(bc-ad)^3 \sin(e+fx)}{b^2 f (a^2-b^2) (a \cos(e+fx)+b)} + \frac{d^2(3bc-2ad) \operatorname{arctanh}(\sin(e+fx))}{b^3 f} + \\
& \frac{2(bc-ad)^2(2ad+bc) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{ab^3 f \sqrt{a-b} \sqrt{a+b}} + \frac{2(bc-ad)^3 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{ab f (a-b)^{3/2} (a+b)^{3/2}} + \\
& \frac{d^3 \tan(e+fx)}{b^2 f}
\end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^3)/(a + b*Sec[e + f*x])^2,x]`

output `(d^2*(3*b*c - 2*a*d)*ArcTanh[Sin[e + f*x]]/(b^3*f) + (2*(b*c - a*d)^3*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(a*(a - b)^(3/2)*b*(a + b)^(3/2)*f) + (2*(b*c - a*d)^2*(b*c + 2*a*d)*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*b^3*Sqrt[a + b]*f) - ((b*c - a*d)^3*Sin[e + f*x])/(b^2*(a^2 - b^2)*f*(b + a*Cos[e + f*x])) + (d^3*Tan[e + f*x])/b^2*f)`

3.260.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3431 Int[((g_.)*sin[(e_.) + (f_.)*(x_)])^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(g*sin[e + f*x])^p*(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && NeQ[b*c - a*d, 0] && (IntegersQ[m, n] || IntegersQ[m, p] || IntegersQ[n, p]) && NeQ[p, 2]
```

```
rule 4476 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[1/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.260.4 Maple [A] (verified)

Time = 1.47 (sec) , antiderivative size = 314, normalized size of antiderivative = 1.38

method	result
derivativedivides	$-\frac{a^3}{b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{d^2(2ad - 3bc) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{b^3} - \frac{\left(\frac{b(a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(a^2 - b^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 a - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 b - a - b}\right)^2}{(2a^4d^3 - 3a^3bd^2)}$
default	$-\frac{a^3}{b^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)} + \frac{d^2(2ad - 3bc) \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{b^3} - \frac{\left(\frac{b(a^3d^3 - 3a^2bcd^2 + 3ab^2c^2d - b^3c^3) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(a^2 - b^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2 a - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 b - a - b}\right)^2}{(2a^4d^3 - 3a^3bd^2)}$
risch	Expression too large to display

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 1/f*(-d^3/b^2/(tan(1/2*f*x+1/2*e)-1)+d^2*(2*a*d-3*b*c)/b^3*ln(tan(1/2*f*x+1/2*e)-1)-2/b^3*(b*(a^3*d^3-3*a^2*b*c*d^2+3*a*b^2*c^2*d-b^3*c^3)/(a^2-b^2)*tan(1/2*f*x+1/2*e)/(tan(1/2*f*x+1/2*e)^2*a-tan(1/2*f*x+1/2*e)^2*b-a-b)-(2*a^4*d^3-3*a^3*b*c*d^2-3*a^2*b^2*d^3+a*b^3*c^3+6*a*b^3*c*d^2-3*b^4*c^2*d)/(a-b)/(a+b)/((a-b)*(a+b))^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e)/((a-b)*(a+b))^(1/2)))-d^3/b^2/(tan(1/2*f*x+1/2*e)+1)-d^2*(2*a*d-3*b*c)/b^3*ln(tan(1/2*f*x+1/2*e)+1))
```

$$3.260. \int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+b\sec(e+fx))^2} dx$$

3.260.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 635 vs. $2(210) = 420$.

Time = 39.31 (sec) , antiderivative size = 1326, normalized size of antiderivative = 5.82

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+b\sec(e+fx))^2} dx = \text{Too large to display}$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e))^2,x, algorithm="fracas")
```

```
output [1/2*((a^2*b^3*c^3 - 3*a*b^4*c^2*d - 3*(a^4*b - 2*a^2*b^3)*c*d^2 + (2*a^5 - 3*a^3*b^2)*d^3)*cos(f*x + e)^2 + (a*b^4*c^3 - 3*b^5*c^2*d - 3*(a^3*b^2 - 2*a*b^4)*c*d^2 + (2*a^4*b - 3*a^2*b^3)*d^3)*cos(f*x + e)*sqrt(a^2 - b^2)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 + 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)) + ((3*(a^5*b - 2*a^3*b^3 + a*b^5)*c*d^2 - 2*(a^6 - 2*a^4*b^2 + a^2*b^4)*d^3)*cos(f*x + e)^2 + (3*(a^4*b^2 - 2*a^2*b^4 + b^6)*c*d^2 - 2*(a^5*b - 2*a^3*b^3 + a*b^5)*d^3)*cos(f*x + e))*log(sin(f*x + e) + 1) - ((3*(a^5*b - 2*a^3*b^3 + a*b^5)*c*d^2 - 2*(a^6 - 2*a^4*b^2 + a^2*b^4)*d^3)*cos(f*x + e)^2 + (3*(a^4*b^2 - 2*a^2*b^4 + b^6)*c*d^2 - 2*(a^5*b - 2*a^3*b^3 + a*b^5)*d^3)*cos(f*x + e))*log(-sin(f*x + e) + 1) + 2*((a^4*b^2 - 2*a^2*b^4 + b^6)*d^3 - ((a^2*b^4 - b^6)*c^3 - 3*(a^3*b^3 - a*b^5)*c^2*d + 3*(a^4*b^2 - a^2*b^4)*c*d^2 - (2*a^5*b - 3*a^3*b^3 + a*b^5)*d^3)*cos(f*x + e)*sin(f*x + e))/((a^5*b^3 - 2*a^3*b^5 + a*b^7)*f*cos(f*x + e)^2 + (a^4*b^4 - 2*a^2*b^6 + b^8)*f*cos(f*x + e)), 1/2*(2*((a^2*b^3*c^3 - 3*a*b^4*c^2*d - 3*(a^4*b - 2*a^2*b^3)*c*d^2 + (2*a^5 - 3*a^3*b^2)*d^3)*cos(f*x + e)^2 + (a*b^4*c^3 - 3*b^5*c^2*d - 3*(a^3*b^2 - 2*a*b^4)*c*d^2 + (2*a^4*b - 3*a^2*b^3)*d^3)*cos(f*x + e))*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e))) + ((3*(a^5*b - 2*a^3*b^3 + a*b^5)*c*d^2 - 2*(a^6 - 2*a^4*b^2 + a^2*b^4)*d^3)*cos(f*x + e)^2 + ...
```

3.260.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+b\sec(e+fx))^2} dx = \int \frac{(c+d\sec(e+fx))^3 \sec(e+fx)}{(a+b\sec(e+fx))^2} dx$$

```
input integrate(sec(f*x+e)*(c+d*sec(f*x+e))**3/(a+b*sec(f*x+e))**2,x)
```

output `Integral((c + d*sec(e + f*x))**3*sec(e + f*x)/(a + b*sec(e + f*x))**2, x)`

3.260.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + b \sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de`

3.260.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 539 vs. $2(210) = 420$.

Time = 0.38 (sec) , antiderivative size = 539, normalized size of antiderivative = 2.36

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))^3}{(a + b \sec(e + fx))^2} dx = \frac{2(ab^3c^3 - 3b^4c^2d - 3a^3bcd^2 + 6ab^3cd^2 + 2a^4d^3 - 3a^2b^2d^3) \left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2}fx + \frac{1}{2}e) - b \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-a^2+b^2}} \right) \right)}{(a^2b^3 - b^5)\sqrt{-a^2+b^2}} - \frac{2(b^2c^3 - 3b^3cd^2 + 2a^2b^2d^3)}{(a^2b^3 - b^5)\sqrt{-a^2+b^2}}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^3/(a+b*sec(f*x+e))^2,x, algorithm="giac")`

output

```

-(2*(a*b^3*c^3 - 3*b^4*c^2*d - 3*a^3*b*c*d^2 + 6*a*b^3*c*d^2 + 2*a^4*d^3 -
  3*a^2*b^2*d^3)*(pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*a - 2*b) + arctan(
  (a*tan(1/2*f*x + 1/2*e) - b*tan(1/2*f*x + 1/2*e))/sqrt(-a^2 + b^2)))/((a^2
  *b^3 - b^5)*sqrt(-a^2 + b^2)) - 2*(b^3*c^3*tan(1/2*f*x + 1/2*e)^3 - 3*a*b^
  2*c^2*d*tan(1/2*f*x + 1/2*e)^3 + 3*a^2*b*c*d^2*tan(1/2*f*x + 1/2*e)^3 - 2*
  a^3*d^3*tan(1/2*f*x + 1/2*e)^3 + a^2*b*d^3*tan(1/2*f*x + 1/2*e)^3 + a*b^2*
  d^3*tan(1/2*f*x + 1/2*e)^3 - b^3*d^3*tan(1/2*f*x + 1/2*e)^3 - b^3*c^3*tan(
  1/2*f*x + 1/2*e) + 3*a*b^2*c^2*d*tan(1/2*f*x + 1/2*e) - 3*a^2*b*c*d^2*tan(
  1/2*f*x + 1/2*e) + 2*a^3*d^3*tan(1/2*f*x + 1/2*e) + a^2*b*d^3*tan(1/2*f*x
  + 1/2*e) - a*b^2*d^3*tan(1/2*f*x + 1/2*e) - b^3*d^3*tan(1/2*f*x + 1/2*e))/
  ((a*tan(1/2*f*x + 1/2*e)^4 - b*tan(1/2*f*x + 1/2*e)^4 - 2*a*tan(1/2*f*x +
  1/2*e)^2 + a + b)*(a^2*b^2 - b^4)) - (3*b*c*d^2 - 2*a*d^3)*log(abs(tan(1/2
  *f*x + 1/2*e) + 1))/b^3 + (3*b*c*d^2 - 2*a*d^3)*log(abs(tan(1/2*f*x + 1/2*
  e) - 1))/b^3)/f

```

3.260.9 Mupad [B] (verification not implemented)

Time = 22.92 (sec) , antiderivative size = 7958, normalized size of antiderivative = 34.90

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^3}{(a+b\sec(e+fx))^2} dx = \text{Too large to display}$$

input `int((c + d/cos(e + f*x))^3/(cos(e + f*x)*(a + b/cos(e + f*x))^2),x)`

output

$$\begin{aligned}
& (d^2 \operatorname{atan}\left(\left(d^2 \left((32 \tan(e/2 + (f*x)/2) * (8*a^8*d^6 - 8*a^7*b*d^6 + a^2*b^6*c^6 + 4*a^2*b^6*d^6 - 8*a^3*b^5*d^6 + 5*a^4*b^4*d^6 + 16*a^5*b^3*d^6 - 16*a^6*b^2*d^6 + 9*b^8*c^2*d^4 + 9*b^8*c^4*d^2 - 18*a*b^7*c^2*d^4 - 36*a*b^7*c^3*d^3 + 24*a^2*b^6*c*d^5 - 24*a^3*b^5*c*d^5 - 48*a^4*b^4*c*d^5 + 54*a^5*b^3*c*d^5 + 24*a^6*b^2*c*d^5 + 45*a^2*b^6*c^2*d^4 + 12*a^2*b^6*c^4*d^2 + 36*a^3*b^5*c^2*d^4 + 12*a^3*b^5*c^3*d^3 - 57*a^4*b^4*c^2*d^4 - 6*a^4*b^4*c^4*d^2 - 18*a^5*b^3*c^2*d^4 + 4*a^5*b^3*c^3*d^3 + 18*a^6*b^2*c^2*d^4 - 12*a*b^7*c*d^5 - 6*a*b^7*c^5*d - 24*a^7*b*c*d^5) \right) / (a*b^6 + b^7 - a^2*b^5 - a^3*b^4) \right) + (d^2 \left((32 * (a*b^{11}*c^3 + 2*a*b^{11}*d^3 - 3*b^{12}*c*d^2 - 3*b^{12}*c^2*d - a^2*b^{10}*c^3 - a^3*b^9*c^3 + a^4*b^8*c^3 - 3*a^2*b^{10}*d^3 - 3*a^3*b^9*d^3 + 5*a^4*b^8*d^3 + a^5*b^7*d^3 - 2*a^6*b^6*d^3 + 3*a^2*b^{10}*c*d^2 + 3*a^2*b^{10}*c^2*d - 9*a^3*b^9*c*d^2 - 3*a^3*b^9*c^2*d + 3*a^5*b^7*c*d^2 + 6*a*b^{11}*c*d^2 + 3*a*b^{11}*c^2*d) \right) / (a*b^8 + b^9 - a^2*b^7 - a^3*b^6) + (32*d^2 * \tan(e/2 + (f*x)/2) * (2*a*d - 3*b*c) * (2*a*b^{11} - 2*a^2*b^{10} - 4*a^3*b^9 + 4*a^4*b^8 + 2*a^5*b^7 - 2*a^6*b^6) \right) / (b^3 * (a*b^6 + b^7 - a^2*b^5 - a^3*b^4))) * (2*a*d - 3*b*c) / b^3 * (2*a*d - 3*b*c) * i) / b^3 + (d^2 \left((32 * \tan(e/2 + (f*x)/2) * (8*a^8*d^6 - 8*a^7*b*d^6 + a^2*b^6*c^6 + 4*a^2*b^6*d^6 - 8*a^3*b^5*d^6 + 5*a^4*b^4*d^6 + 16*a^5*b^3*d^6 - 16*a^6*b^2*d^6 + 9*b^8*c^2*d^4 + 9*b^8*c^4*d^2 - 18*a*b^7*c^2*d^4 - 36*a*b^7*c^3*d^3 + 24*a^2*b^6*c*d^5 - 24*a^3*b^5*c*d^5 - 48*a^4*b^4*c*d^5 + 54*a^5*b^3*c*d^5 + 24*a^6*b^2*c*d^5 + 4... \right)
\end{aligned}$$

3.261 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+b \sec(e+fx))^2} dx$

3.261.1 Optimal result 1920
 3.261.2 Mathematica [A] (verified) 1921
 3.261.3 Rubi [A] (verified) 1921
 3.261.4 Maple [A] (verified) 1923
 3.261.5 Fricas [B] (verification not implemented) 1924
 3.261.6 Sympy [F] 1925
 3.261.7 Maxima [F(-2)] 1925
 3.261.8 Giac [A] (verification not implemented) 1925
 3.261.9 Mupad [B] (verification not implemented) 1926

3.261.1 Optimal result

Integrand size = 31, antiderivative size = 198

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))^2}{(a+b \sec(e+fx))^2} dx = \frac{d^2 \operatorname{arctanh}(\sin(e+fx))}{b^2 f} + \frac{2(bc-ad)^2 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{a(a-b)^{3/2}(a+b)^{3/2} f} + \frac{2(b^2 c^2 - a^2 d^2) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{a\sqrt{a-b} b^2 \sqrt{a+b} f} - \frac{(bc-ad)^2 \sin(e+fx)}{b(a^2-b^2) f(b+a \cos(e+fx))}$$

```
output d^2*arctanh(sin(f*x+e))/b^2/f+2*(-a*d+b*c)^2*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/a/(a-b)^(3/2)/(a+b)^(3/2)/f-(-a*d+b*c)^2*sin(f*x+e)/b/(a^2-b^2)/f/(b+a*cos(f*x+e))+2*(-a^2*d^2+b^2*c^2)*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/a/b^2/f/(a-b)^(1/2)/(a+b)^(1/2)
```

3.261.2 Mathematica [A] (verified)

Time = 1.24 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.91

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx$$

$$= \frac{2(2b^3cd+a^3d^2-ab^2(c^2+2d^2))\operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right) - d^2 \log\left(\cos\left(\frac{1}{2}(e+fx)\right) - \sin\left(\frac{1}{2}(e+fx)\right)\right) + d^2 \log\left(\cos\left(\frac{1}{2}(e+fx)\right) + \sin\left(\frac{1}{2}(e+fx)\right)\right)}{(a^2-b^2)^{3/2} b^2 f}$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + b*Sec[e + f*x])^2,x]`output `((2*(2*b^3*c*d + a^3*d^2 - a*b^2*(c^2 + 2*d^2))*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]])/(a^2 - b^2)^(3/2) - d^2*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]] + d^2*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]] + (b*(b*c - a*d)^2*Sin[e + f*x])/((-a + b)*(a + b)*(b + a*Cos[e + f*x]))/(b^2*f)`**3.261.3 Rubi [A] (verified)**Time = 0.63 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 4476, 3042, 3431, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)(c+d\csc\left(e+fx+\frac{\pi}{2}\right))^2}{(a+b\csc\left(e+fx+\frac{\pi}{2}\right))^2} dx$$

$$\downarrow \text{4476}$$

$$\int \frac{\sec(e+fx)(c\cos(e+fx)+d)^2}{(a\cos(e+fx)+b)^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c\sin\left(e+fx+\frac{\pi}{2}\right)+d)^2}{\sin\left(e+fx+\frac{\pi}{2}\right)(a\sin\left(e+fx+\frac{\pi}{2}\right)+b)^2} dx$$

3.261. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx$

$$\begin{aligned}
 & \int \left(\frac{b^2 c^2 - a^2 d^2}{ab^2(a \cos(e + fx) + b)} - \frac{(ad - bc)^2}{ab(a \cos(e + fx) + b)^2} + \frac{d^2 \sec(e + fx)}{b^2} \right) dx \\
 & \quad \downarrow \text{3431} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2(b^2 c^2 - a^2 d^2) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{ab^2 f \sqrt{a-b} \sqrt{a+b}} - \frac{(bc - ad)^2 \sin(e + fx)}{bf(a^2 - b^2)(a \cos(e + fx) + b)} + \\
 & \frac{2(bc - ad)^2 \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{af(a-b)^{3/2}(a+b)^{3/2}} + \frac{d^2 \operatorname{arctanh}(\sin(e + fx))}{b^2 f}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x])^2)/(a + b*Sec[e + f*x])^2,x]`

output `(d^2*ArcTanh[Sin[e + f*x]])/(b^2*f) + (2*(b*c - a*d)^2*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]]/(a*(a - b)^(3/2)*(a + b)^(3/2)*f) + (2*(b^2*c^2 - a^2*d^2)*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]]/(a*Sqrt[a - b]*b^2*Sqrt[a + b]*f) - ((b*c - a*d)^2*Sin[e + f*x])/(b*(a^2 - b^2)*f*(b + a*Cos[e + f*x]))`

3.261.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3431 `Int[((g_.)*sin[(e_.) + (f_.)*(x_)])^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(g*sin[e + f*x])^p*(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && NeQ[b*c - a*d, 0] && (IntegersQ[m, n] || IntegersQ[m, p] || IntegersQ[n, p]) && NeQ[p, 2]`

```
rule 4476 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))^(n_), x_Symbol] := Simp[1
/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c
*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c -
a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.261.4 Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.09

method	result
derivativedivides	$-\frac{d^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{b^2} + \frac{d^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{b^2} + \frac{2b(a^2d^2 - 2abcd + b^2c^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(a^2 - b^2)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 a - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 b - a - b\right)} - \frac{2(a^3d^2 - b^2c^2a - 2b^2d^2a + (a-b))}{b^2}$
default	$-\frac{d^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 1\right)}{b^2} + \frac{d^2 \ln\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right) + 1\right)}{b^2} + \frac{2b(a^2d^2 - 2abcd + b^2c^2) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(a^2 - b^2)\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 a - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 b - a - b\right)} - \frac{2(a^3d^2 - b^2c^2a - 2b^2d^2a + (a-b))}{b^2}$
risch	$-\frac{2i(a^2d^2 - 2abcd + b^2c^2)(e^{i(fx+e)}b+a)}{(a^2 - b^2)fa(e^{2i(fx+e)}a + 2e^{i(fx+e)}b+a)} + \frac{\ln\left(\frac{e^{i(fx+e)} - ia^2 - ib^2 - b\sqrt{a^2 - b^2}}{\sqrt{a^2 - b^2}a}\right)a^3d^2}{\sqrt{a^2 - b^2}(a+b)(a-b)fb^2} - \frac{\ln\left(\frac{e^{i(fx+e)} - ia^2 - ib^2 - b\sqrt{a^2 - b^2}}{\sqrt{a^2 - b^2}a}\right)}{\sqrt{a^2 - b^2}(a+b)(a-b)}$

```
input int(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e))^2,x,method=_RETURNVERBO
SE)
```

```
output 1/f*(-d^2/b^2*ln(tan(1/2*f*x+1/2*e)-1)+d^2/b^2*ln(tan(1/2*f*x+1/2*e)+1)+2/
b^2*(b*(a^2*d^2-2*a*b*c*d+b^2*c^2)/(a^2-b^2)*tan(1/2*f*x+1/2*e)/(tan(1/2*f
*x+1/2*e)^2*a-tan(1/2*f*x+1/2*e)^2*b-a-b)-(a^3*d^2-a*b^2*c^2-2*a*b^2*d^2+2
*b^3*c*d)/(a-b)/(a+b)/((a-b)*(a+b))^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e)
/((a-b)*(a+b))^(1/2))))
```

3.261.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 371 vs. $2(180) = 360$.

Time = 5.70 (sec) , antiderivative size = 798, normalized size of antiderivative = 4.03

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx$$

$$= \left[\frac{(ab^3c^2 - 2b^4cd - (a^3b - 2ab^3)d^2 + (a^2b^2c^2 - 2ab^3cd - (a^4 - 2a^2b^2)d^2) \cos(fx+e))\sqrt{a^2-b^2} \log\left(\frac{2a}{\dots}\right)}{\dots} \right]$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e))^2,x, algorithm="fracas")`

output `[-1/2*((a*b^3*c^2 - 2*b^4*c*d - (a^3*b - 2*a*b^3)*d^2 + (a^2*b^2*c^2 - 2*a*b^3*c*d - (a^4 - 2*a^2*b^2)*d^2)*cos(f*x + e))*sqrt(a^2 - b^2)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 - 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)) - ((a^5 - 2*a^3*b^2 + a*b^4)*d^2*cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*d^2)*log(sin(f*x + e) + 1) + ((a^5 - 2*a^3*b^2 + a*b^4)*d^2*cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*d^2)*log(-sin(f*x + e) + 1) + 2*((a^2*b^3 - b^5)*c^2 - 2*(a^3*b^2 - a*b^4)*c*d + (a^4*b - a^2*b^3)*d^2)*sin(f*x + e)/((a^5*b^2 - 2*a^3*b^4 + a*b^6)*f*cos(f*x + e) + (a^4*b^3 - 2*a^2*b^5 + b^7)*f), 1/2*(2*(a*b^3*c^2 - 2*b^4*c*d - (a^3*b - 2*a*b^3)*d^2 + (a^2*b^2*c^2 - 2*a*b^3*c*d - (a^4 - 2*a^2*b^2)*d^2)*cos(f*x + e))*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*cos(f*x + e) + a)/((a^2 - b^2)*sin(f*x + e))) + ((a^5 - 2*a^3*b^2 + a*b^4)*d^2*cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*d^2)*log(sin(f*x + e) + 1) - ((a^5 - 2*a^3*b^2 + a*b^4)*d^2*cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*d^2)*log(-sin(f*x + e) + 1) - 2*((a^2*b^3 - b^5)*c^2 - 2*(a^3*b^2 - a*b^4)*c*d + (a^4*b - a^2*b^3)*d^2)*sin(f*x + e)/((a^5*b^2 - 2*a^3*b^4 + a*b^6)*f*cos(f*x + e) + (a^4*b^3 - 2*a^2*b^5 + b^7)*f)]`

3.261.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx = \int \frac{(c+d\sec(e+fx))^2 \sec(e+fx)}{(a+b\sec(e+fx))^2} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))**2/(a+b*sec(f*x+e))**2,x)`

output `Integral((c + d*sec(e + f*x))**2*sec(e + f*x)/(a + b*sec(e + f*x))**2, x)`

3.261.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de`

3.261.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.35

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx$$

$$= \frac{d^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1|)}{b^2} - \frac{d^2 \log(|\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1|)}{b^2} - \frac{2(ab^2c^2 - 2b^3cd - a^3d^2 + 2ab^2d^2) \left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2}fx + \frac{1}{2}e)}{b} \right) \right)}{(a^2b^2 - b^4)\sqrt{-a^2 + b^2}}$$

3.261. $\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))^2/(a+b*sec(f*x+e))^2,x, algorithm="giac")`

output $(d^2 \log(\abs{\tan(1/2fx + 1/2e) + 1})/b^2 - d^2 \log(\abs{\tan(1/2fx + 1/2e) - 1})/b^2 - 2(a^2b^2c^2 - 2b^3cd - a^3d^2 + 2ab^2d^2)(\pi \operatorname{floor}(1/2(fx + e)/\pi + 1/2) \operatorname{sgn}(2a - 2b) + \arctan((a \tan(1/2fx + 1/2e) - b \tan(1/2fx + 1/2e))/\sqrt{-a^2 + b^2}))/((a^2b^2 - b^4)\sqrt{-a^2 + b^2}) + 2(b^2c^2 \tan(1/2fx + 1/2e) - 2ab^2cd \tan(1/2fx + 1/2e) + a^2d^2 \tan(1/2fx + 1/2e))/((a^2b - b^3)(a \tan(1/2fx + 1/2e)^2 - b \tan(1/2fx + 1/2e)^2 - a - b))/f$

3.261.9 Mupad [B] (verification not implemented)

Time = 21.29 (sec) , antiderivative size = 4926, normalized size of antiderivative = 24.88

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))^2}{(a+b\sec(e+fx))^2} dx = \text{Too large to display}$$

input `int((c + d/cos(e + f*x))^2/(cos(e + f*x)*(a + b/cos(e + f*x))^2),x)`

output $- (d^2 \operatorname{atan}(((d^2((32 \tan(e/2 + (fx)/2)(2a^6d^4 + b^6d^4 - 2ab^5d^4 - 2a^5bd^4 + a^2b^4c^4 + 3a^2b^4d^4 + 4a^3b^3d^4 - 5a^4b^2d^4 + 4b^6c^2d^2 + 4a^3b^3cd^3 + 4a^2b^4c^2d^2 - 2a^4b^2c^2d^2 - 8ab^5cd^3 - 4ab^5c^3d)))/(ab^4 + b^5 - a^2b^3 - a^3b^2) + (d^2((32(ab^8c^2 - b^9d^2 + 2ab^8d^2 - a^2b^7c^2 - a^3b^6c^2 + a^4b^5c^2 + a^2b^7d^2 - 3a^3b^6d^2 + a^5b^4d^2 - 2b^9cd + 2ab^8cd + 2a^2b^7cd - 2a^3b^6cd)))/(ab^5 + b^6 - a^2b^4 - a^3b^3) + (32d^2 \tan(e/2 + (fx)/2)(2ab^9 - 2a^2b^8 - 4a^3b^7 + 4a^4b^6 + 2a^5b^5 - 2a^6b^4))/(b^2(ab^4 + b^5 - a^2b^3 - a^3b^2)))))/b^2) * i) / b^2 + (d^2((32 \tan(e/2 + (fx)/2)(2a^6d^4 + b^6d^4 - 2ab^5d^4 - 2a^5bd^4 + a^2b^4c^4 + 3a^2b^4d^4 + 4a^3b^3d^4 - 5a^4b^2d^4 + 4b^6c^2d^2 + 4a^3b^3cd^3 + 4a^2b^4c^2d^2 - 2a^4b^2c^2d^2 - 8ab^5cd^3 - 4ab^5c^3d)))/(ab^4 + b^5 - a^2b^3 - a^3b^2) - (d^2((32(ab^8c^2 - b^9d^2 + 2ab^8d^2 - a^2b^7c^2 - a^3b^6c^2 + a^4b^5c^2 + a^2b^7d^2 - 3a^3b^6d^2 + a^5b^4d^2 - 2b^9cd + 2ab^8cd + 2a^2b^7cd - 2a^3b^6cd)))/(ab^5 + b^6 - a^2b^4 - a^3b^3) - (32d^2 \tan(e/2 + (fx)/2)(2ab^9 - 2a^2b^8 - 4a^3b^7 + 4a^4b^6 + 2a^5b^5 - 2a^6b^4))/(b^2(ab^4 + b^5 - a^2b^3 - a^3b^2)))))/b^2) * i) / b^2) / ((64(a^5d^6 + 2ab^4d^6 - a^4bd^6 - 2b^5cd^5 + 2a^2b^3d^6 - 3a^3b^2d^6 + 4b^5c^2d^4 + ab^4c^2d^4 - 4ab^4c^3d...$

3.262 $\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+b \sec(e+fx))^2} dx$

3.262.1 Optimal result 1927
 3.262.2 Mathematica [A] (verified) 1927
 3.262.3 Rubi [A] (verified) 1928
 3.262.4 Maple [A] (verified) 1930
 3.262.5 Fricas [A] (verification not implemented) 1931
 3.262.6 Sympy [F] 1931
 3.262.7 Maxima [F(-2)] 1932
 3.262.8 Giac [A] (verification not implemented) 1932
 3.262.9 Mupad [B] (verification not implemented) 1933

3.262.1 Optimal result

Integrand size = 29, antiderivative size = 100

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+b \sec(e+fx))^2} dx = \frac{2(ac-bd) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{(a-b)^{3/2}(a+b)^{3/2} f} - \frac{(bc-ad) \tan(e+fx)}{(a^2-b^2) f(a+b \sec(e+fx))}$$

output `2*(a*c-b*d)*arctanh((a-b)^(1/2)*tan(1/2*f*x+1/2*e)/(a+b)^(1/2))/(a-b)^(3/2)/(a+b)^(3/2)/f-(-a*d+b*c)*tan(f*x+e)/(a^2-b^2)/f/(a+b*sec(f*x+e))`

3.262.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.97

$$\int \frac{\sec(e+fx)(c+d \sec(e+fx))}{(a+b \sec(e+fx))^2} dx = \frac{2(ac-bd) \operatorname{arctanh}\left(\frac{(-a+b) \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} + \frac{(-bc+ad) \sin(e+fx)}{(a-b)(a+b)(b+a \cos(e+fx))} f$$

input `Integrate[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + b*Sec[e + f*x])^2,x]`


```
output ((-2*(a*c - b*d)*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]])/(a^
2 - b^2)^(3/2) + (((-b*c) + a*d)*Sin[e + f*x])/((a - b)*(a + b)*(b + a*Cos
[e + f*x])))/f
```

3.262.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.11, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 4491, 25, 27, 3042, 4318, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+b\sec(e+fx))^2} dx$$

↓ 3042

$$\int \frac{\csc(e+fx+\frac{\pi}{2})(c+d\csc(e+fx+\frac{\pi}{2}))}{(a+b\csc(e+fx+\frac{\pi}{2}))^2} dx$$

↓ 4491

$$-\frac{\int -\frac{(ac-bd)\sec(e+fx)}{a+b\sec(e+fx)} dx}{a^2-b^2} - \frac{(bc-ad)\tan(e+fx)}{f(a^2-b^2)(a+b\sec(e+fx))}$$

↓ 25

$$\frac{\int \frac{(ac-bd)\sec(e+fx)}{a+b\sec(e+fx)} dx}{a^2-b^2} - \frac{(bc-ad)\tan(e+fx)}{f(a^2-b^2)(a+b\sec(e+fx))}$$

↓ 27

$$\frac{(ac-bd)\int \frac{\sec(e+fx)}{a+b\sec(e+fx)} dx}{a^2-b^2} - \frac{(bc-ad)\tan(e+fx)}{f(a^2-b^2)(a+b\sec(e+fx))}$$

↓ 3042

$$\frac{(ac-bd)\int \frac{\csc(e+fx+\frac{\pi}{2})}{a+b\csc(e+fx+\frac{\pi}{2})} dx}{a^2-b^2} - \frac{(bc-ad)\tan(e+fx)}{f(a^2-b^2)(a+b\sec(e+fx))}$$

↓ 4318

$$\frac{(ac-bd)\int \frac{1}{a\cos(\frac{e+fx}{b})+1} dx}{b(a^2-b^2)} - \frac{(bc-ad)\tan(e+fx)}{f(a^2-b^2)(a+b\sec(e+fx))}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{(ac - bd) \int \frac{1}{\frac{a \sin(e+fx+\frac{\pi}{2})}{b} + 1} dx}{b(a^2 - b^2)} - \frac{(bc - ad) \tan(e + fx)}{f(a^2 - b^2)(a + b \sec(e + fx))} \\
 & \downarrow 3138 \\
 & \frac{2(ac - bd) \int \frac{1}{(1 - \frac{a}{b}) \tan^2(\frac{1}{2}(e+fx)) + \frac{a+b}{b}} d \tan(\frac{1}{2}(e + fx))}{bf(a^2 - b^2)} - \frac{(bc - ad) \tan(e + fx)}{f(a^2 - b^2)(a + b \sec(e + fx))} \\
 & \downarrow 221 \\
 & \frac{2(ac - bd) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e+fx))}{\sqrt{a+b}}\right)}{f\sqrt{a-b}\sqrt{a+b}(a^2 - b^2)} - \frac{(bc - ad) \tan(e + fx)}{f(a^2 - b^2)(a + b \sec(e + fx))}
 \end{aligned}$$

input `Int[(Sec[e + f*x]*(c + d*Sec[e + f*x]))/(a + b*Sec[e + f*x])^2,x]`

output `(2*(a*c - b*d)*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]]/(Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*f) - ((b*c - a*d)*Tan[e + f*x])/((a^2 - b^2)*f*(a + b*Sec[e + f*x]))`

3.262.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 4318 `Int[csc[(e_) + (f_)*(x_)]/(csc[(e_) + (f_)*(x_)]*(b_) + (a_)), x_Symbol] := Simp[1/b Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4491 `Int[csc[(e_) + (f_)*(x_)]*(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(m_)*(csc[(e_) + (f_)*(x_)]*(B_) + (A_)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[Csc[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*Simp[(a*A - b*B)*(m + 1) - (A*b - a*B)*(m + 2)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, A, B, e, f}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.262.4 Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.32

method	result
derivativedivides	$-\frac{2(ad-bc)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{(a^2-b^2)\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 a-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 b-a-b\right)}+\frac{2(ac-bd)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}}$
default	$-\frac{2(ad-bc)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{(a^2-b^2)\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 a-\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2 b-a-b\right)}+\frac{2(ac-bd)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}}$
risch	$\frac{2i(ad-bc)(e^{i(fx+e)}b+a)}{a(a^2-b^2)f(e^{2i(fx+e)}a+2e^{i(fx+e)}b+a)}+\frac{\ln\left(e^{i(fx+e)}+\frac{ia^2-ib^2+b\sqrt{a^2-b^2}}{\sqrt{a^2-b^2}a}\right)ac}{\sqrt{a^2-b^2}(a+b)(a-b)f}-\frac{\ln\left(e^{i(fx+e)}+\frac{ia^2-ib^2+b\sqrt{a^2-b^2}}{\sqrt{a^2-b^2}a}\right)}{\sqrt{a^2-b^2}(a+b)(a-b)f}$

input `int(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^2,x,method=_RETURNVERBOSE)`

output $1/f*(-2*(a*d-b*c)/(a^2-b^2)*\tan(1/2*f*x+1/2*e)/(\tan(1/2*f*x+1/2*e)^2*a-\tan(1/2*f*x+1/2*e)^2*b-a-b)+2*(a*c-b*d)/(a-b)/(a+b)/((a-b)*(a+b))^{(1/2)*\arctan((a-b)*\tan(1/2*f*x+1/2*e)/((a-b)*(a+b))^{(1/2)})}$

3.262.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 394, normalized size of antiderivative = 3.94

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+b\sec(e+fx))^2} dx$$

$$= \left[\frac{(abc - b^2d + (a^2c - abd) \cos(fx + e))\sqrt{a^2 - b^2} \log\left(\frac{2ab \cos(fx+e) - (a^2 - 2b^2) \cos(fx+e)^2 + 2\sqrt{a^2 - b^2}(b \cos(fx+e) + a) \sin(fx+e)}{a^2 \cos(fx+e)^2 + 2ab \cos(fx+e) + b^2}\right)}{2((a^5 - 2a^3b^2 + ab^4)f \cos(fx + e) + (a^4b - 2a^2b^3 + b^5)f)}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^2,x, algorithm="fricas")`

output $[1/2*((a*b*c - b^2*d + (a^2*c - a*b*d)*\cos(f*x + e))*\sqrt{a^2 - b^2}*\log((2*a*b*\cos(f*x + e) - (a^2 - 2*b^2)*\cos(f*x + e)^2 + 2*\sqrt{a^2 - b^2}*(b*\cos(f*x + e) + a)*\sin(f*x + e) + 2*a^2 - b^2)/(a^2*\cos(f*x + e)^2 + 2*a*b*\cos(f*x + e) + b^2)) - 2*((a^2*b - b^3)*c - (a^3 - a*b^2)*d)*\sin(f*x + e)]/((a^5 - 2*a^3*b^2 + a*b^4)*f*\cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*f), ((a*b*c - b^2*d + (a^2*c - a*b*d)*\cos(f*x + e))*\sqrt{-a^2 + b^2}*\arctan(\sqrt{-a^2 + b^2}*(b*\cos(f*x + e) + a)/((a^2 - b^2)*\sin(f*x + e))) - ((a^2*b - b^3)*c - (a^3 - a*b^2)*d)*\sin(f*x + e)]/((a^5 - 2*a^3*b^2 + a*b^4)*f*\cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*f)]$

3.262.6 Sympy [F]

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+b\sec(e+fx))^2} dx = \int \frac{(c+d\sec(e+fx))\sec(e+fx)}{(a+b\sec(e+fx))^2} dx$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e))**2,x)`

output `Integral((c + d*sec(e + f*x))*sec(e + f*x)/(a + b*sec(e + f*x))**2, x)`

3.262.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{(a + b \sec(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` f or more de`

3.262.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.73

$$\int \frac{\sec(e + fx)(c + d \sec(e + fx))}{(a + b \sec(e + fx))^2} dx =$$

$$2 \left(\frac{\left(\pi \left\lfloor \frac{fx+e}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan\left(\frac{a \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) - b \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)}{\sqrt{-a^2+b^2}} \right) \right) (ac-bd)}{(a^2-b^2)\sqrt{-a^2+b^2}} - \frac{bc \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) - ad \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)}{(a \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - b \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - a - b)} \right) / f$$

input `integrate(sec(f*x+e)*(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^2,x, algorithm="giac")`

output `-2*((pi*floor(1/2*(f*x + e)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*f*x + 1/2*e) - b*tan(1/2*f*x + 1/2*e))/sqrt(-a^2 + b^2)))*(a*c - b*d)/((a^2 - b^2)*sqrt(-a^2 + b^2)) - (b*c*tan(1/2*f*x + 1/2*e) - a*d*tan(1/2*f*x + 1/2*e))/((a*tan(1/2*f*x + 1/2*e)^2 - b*tan(1/2*f*x + 1/2*e)^2 - a - b)*(a^2 - b^2)))/f`

3.262.9 Mupad [B] (verification not implemented)

Time = 13.75 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.06

$$\int \frac{\sec(e+fx)(c+d\sec(e+fx))}{(a+b\sec(e+fx))^2} dx = \frac{2 \operatorname{atanh}\left(\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right) \sqrt{a-b}}{\sqrt{a+b}}\right) (ac - bd)}{f(a+b)^{3/2}(a-b)^{3/2}} + \frac{2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) (ad - bc)}{f(a+b)(a-b)\left((b-a)\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + a+b\right)}$$

input `int((c + d/cos(e + f*x))/(cos(e + f*x)*(a + b/cos(e + f*x))^2),x)`output `(2*atanh((tan(e/2 + (f*x)/2)*(a - b)^(1/2))/(a + b)^(1/2))*(a*c - b*d))/(f*(a + b)^(3/2)*(a - b)^(3/2)) + (2*tan(e/2 + (f*x)/2)*(a*d - b*c))/(f*(a + b)*(a - b)*(a + b - tan(e/2 + (f*x)/2)^2*(a - b)))`

3.263 $\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))^2(c+d \sec(e+fx))} dx$

3.263.1 Optimal result	1934
3.263.2 Mathematica [A] (verified)	1934
3.263.3 Rubi [A] (verified)	1935
3.263.4 Maple [A] (verified)	1938
3.263.5 Fricas [B] (verification not implemented)	1939
3.263.6 Sympy [F]	1939
3.263.7 Maxima [F(-2)]	1940
3.263.8 Giac [A] (verification not implemented)	1940
3.263.9 Mupad [B] (verification not implemented)	1941

3.263.1 Optimal result

Integrand size = 31, antiderivative size = 186

$$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))^2(c+d \sec(e+fx))} dx$$

$$= \frac{2b(abc - 2a^2d + b^2d) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a+b}}\right)}{(a-b)^{3/2}(a+b)^{3/2}(bc-ad)^2 f}$$

$$+ \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c+d}}\right)}{\sqrt{c-d}\sqrt{c+d}(bc-ad)^2 f} - \frac{b^2 \sin(e+fx)}{(a^2-b^2)(bc-ad)f(b+a \cos(e+fx))}$$

output $2*b*(-2*a^2*d+a*b*c+b^2*d)*\operatorname{arctanh}\left(\frac{(a-b)^{1/2}*\tan(1/2*f*x+1/2*e)}{(a+b)^{1/2}}\right)/(a-b)^{3/2}/(a+b)^{3/2}/(-a*d+b*c)^2/f-b^2*\sin(f*x+e)/(a^2-b^2)/(-a*d+b*c)/f/(b+a*\cos(f*x+e))+2*d^2*\operatorname{arctanh}\left(\frac{(c-d)^{1/2}*\tan(1/2*f*x+1/2*e)}{(c+d)^{1/2}}\right)/(-a*d+b*c)^2/f/(c-d)^{1/2}/(c+d)^{1/2}$

3.263.2 Mathematica [A] (verified)

Time = 1.59 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.95

$$\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))^2(c+d \sec(e+fx))} dx$$

$$= \frac{2b(abc-2a^2d+b^2d)\operatorname{arctanh}\left(\frac{(-a+b)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}} + \frac{2(a^2-b^2)d^2\operatorname{arctanh}\left(\frac{(-c+d)\tan\left(\frac{1}{2}(e+fx)\right)}{\sqrt{c^2-d^2}}\right)}{\sqrt{c^2-d^2}} + \frac{b^2(bc-ad)\sin(e+fx)}{b+a \cos(e+fx)}$$

$$(-a+b)(a+b)(bc-ad)^2 f$$

3.263. $\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))^2(c+d \sec(e+fx))} dx$

input `Integrate[Sec[e + f*x]/((a + b*Sec[e + f*x])^2*(c + d*Sec[e + f*x])),x]`

output `((2*b*(a*b*c - 2*a^2*d + b^2*d)*ArcTanh[((-a + b)*Tan[(e + f*x)/2])/Sqrt[a^2 - b^2]])/Sqrt[a^2 - b^2] + (2*(a^2 - b^2)*d^2*ArcTanh[(-c + d)*Tan[(e + f*x)/2])/Sqrt[c^2 - d^2])/Sqrt[c^2 - d^2] + (b^2*(b*c - a*d)*Sin[e + f*x])/(b + a*cos[e + f*x])/((-a + b)*(a + b)*(b*c - a*d)^2*f)`

3.263.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.17, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4476, 3042, 3535, 25, 3042, 3480, 3042, 3138, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(e + fx)}{(a + b \sec(e + fx))^2 (c + d \sec(e + fx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(e + fx + \frac{\pi}{2})}{(a + b \csc(e + fx + \frac{\pi}{2}))^2 (c + d \csc(e + fx + \frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{4476} \\
 & \int \frac{\cos^2(e + fx)}{(a \cos(e + fx) + b)^2 (c \cos(e + fx) + d)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(e + fx + \frac{\pi}{2})^2}{(a \sin(e + fx + \frac{\pi}{2}) + b)^2 (c \sin(e + fx + \frac{\pi}{2}) + d)} dx \\
 & \quad \downarrow \text{3535} \\
 & \int \frac{-\frac{abd + (-da^2 + bca + b^2d) \cos(e + fx)}{(b + a \cos(e + fx))(d + c \cos(e + fx))} dx}{(a^2 - b^2)(bc - ad)} - \frac{b^2 \sin(e + fx)}{f(a^2 - b^2)(bc - ad)(a \cos(e + fx) + b)} \\
 & \quad \downarrow \text{25} \\
 & \int \frac{abd + (-da^2 + bca + b^2d) \cos(e + fx)}{(b + a \cos(e + fx))(d + c \cos(e + fx))} dx - \frac{b^2 \sin(e + fx)}{f(a^2 - b^2)(bc - ad)(a \cos(e + fx) + b)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.263. $\int \frac{\sec(e + fx)}{(a + b \sec(e + fx))^2 (c + d \sec(e + fx))} dx$

$$\begin{aligned}
 & \int \frac{abd + (-da^2 + bca + b^2d) \sin(e + fx + \frac{\pi}{2})}{(b + a \sin(e + fx + \frac{\pi}{2}))(d + c \sin(e + fx + \frac{\pi}{2}))} dx - \frac{b^2 \sin(e + fx)}{f(a^2 - b^2)(bc - ad)(a \cos(e + fx) + b)} \\
 & \quad \downarrow \text{3480} \\
 & \frac{d^2(a^2 - b^2) \int \frac{1}{d + c \cos(e + fx)} dx}{bc - ad} + \frac{b(-2a^2d + abc + b^2d) \int \frac{1}{b + a \cos(e + fx)} dx}{bc - ad} - \frac{b^2 \sin(e + fx)}{f(a^2 - b^2)(bc - ad)(a \cos(e + fx) + b)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{d^2(a^2 - b^2) \int \frac{1}{d + c \sin(e + fx + \frac{\pi}{2})} dx}{bc - ad} + \frac{b(-2a^2d + abc + b^2d) \int \frac{1}{b + a \sin(e + fx + \frac{\pi}{2})} dx}{bc - ad} - \\
 & \quad \frac{b^2 \sin(e + fx)}{f(a^2 - b^2)(bc - ad)(a \cos(e + fx) + b)} \\
 & \quad \downarrow \text{3138} \\
 & \frac{2d^2(a^2 - b^2) \int \frac{1}{-(c - d) \tan^2(\frac{1}{2}(e + fx)) + c + d} d \tan(\frac{1}{2}(e + fx))}{f(bc - ad)} + \frac{2b(-2a^2d + abc + b^2d) \int \frac{1}{-(a - b) \tan^2(\frac{1}{2}(e + fx)) + a + b} d \tan(\frac{1}{2}(e + fx))}{f(bc - ad)} - \\
 & \quad \frac{b^2 \sin(e + fx)}{f(a^2 - b^2)(bc - ad)(a \cos(e + fx) + b)} \\
 & \quad \downarrow \text{221} \\
 & \frac{2d^2(a^2 - b^2) \operatorname{arctanh}\left(\frac{\sqrt{c-d} \tan(\frac{1}{2}(e + fx))}{\sqrt{c+d}}\right)}{f\sqrt{c-d}\sqrt{c+d}(bc - ad)} + \frac{2b(-2a^2d + abc + b^2d) \operatorname{arctanh}\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(e + fx))}{\sqrt{a+b}}\right)}{f\sqrt{a-b}\sqrt{a+b}(bc - ad)} - \\
 & \quad \frac{b^2 \sin(e + fx)}{f(a^2 - b^2)(bc - ad)(a \cos(e + fx) + b)}
 \end{aligned}$$

input `Int[Sec[e + f*x]/((a + b*Sec[e + f*x])^2*(c + d*Sec[e + f*x])),x]`

output `((2*b*(a*b*c - 2*a^2*d + b^2*d)*ArcTanh[(Sqrt[a - b]*Tan[(e + f*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*(b*c - a*d)*f) + (2*(a^2 - b^2)*d^2*ArcTanh[(Sqrt[c - d]*Tan[(e + f*x)/2])/Sqrt[c + d]])/(Sqrt[c - d]*Sqrt[c + d]*(b*c - a*d)*f)/((a^2 - b^2)*(b*c - a*d)) - (b^2*Sin[e + f*x])/((a^2 - b^2)*(b*c - a*d)*f*(b + a*Cos[e + f*x]))`

3.263.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3535 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[a*(m + 1)*(b*c - a*d)*(A + C) + d*(A*b^2 + a^2*C)*(m + n + 2) - (c*(A*b^2 + a^2*C) + b*(m + 1)*(b*c - a*d)*(A + C))*Sin[e + f*x] - d*(A*b^2 + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 4476 Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))^(n_.), x_Symbol] := Simp[1
/g^(m + n) Int[(g*Csc[e + f*x])^(m + n + p)*(b + a*Sin[e + f*x])^m*(d + c
*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c -
a*d, 0] && IntegerQ[m] && IntegerQ[n]
```

3.263.4 Maple [A] (verified)

Time = 1.45 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.13

method	result
derivativedivides	$\frac{2b \left(\frac{b(ad-bc) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(a^2-b^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 a - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 b - a - b} \right) - \frac{(2a^2d-abc-db^2) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}} \right)}{(ad-bc)^2} + \frac{2d^2 \operatorname{arctanh}\left(\frac{(c-d)}{\sqrt{(a-b)(a+b)}}\right)}{(ad-bc)^2 \sqrt{(a-b)(a+b)}}$
default	$\frac{2b \left(\frac{b(ad-bc) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{(a^2-b^2) \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 a - \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 b - a - b} \right) - \frac{(2a^2d-abc-db^2) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}} \right)}{(ad-bc)^2} + \frac{2d^2 \operatorname{arctanh}\left(\frac{(c-d)}{\sqrt{(a-b)(a+b)}}\right)}{(ad-bc)^2 \sqrt{(a-b)(a+b)}}$
risch	$\frac{2ib^2(e^{i(fx+e)}b+a)}{a(a^2-b^2)(ad-bc)f(e^{2i(fx+e)}a+2e^{i(fx+e)}b+a)} + \frac{2b \ln\left(\frac{e^{i(fx+e)} - ia^2 - ib^2 - b\sqrt{a^2-b^2}}{\sqrt{a^2-b^2}a}\right) a^2 d}{\sqrt{a^2-b^2} (ad-bc)^2 (a+b)(a-b)f} - \frac{b^2 \ln\left(\frac{e^{i(fx+e)} - ia^2 - ib^2 - b\sqrt{a^2-b^2}}{\sqrt{a^2-b^2}a}\right)}{\sqrt{a^2-b^2} (ad-bc)^2}$

```
input int(sec(f*x+e)/(a+b*sec(f*x+e))^2/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE
)
```

```
output 1/f*(2*b/(a*d-b*c)^2*(-b*(a*d-b*c)/(a^2-b^2)*tan(1/2*f*x+1/2*e)/(tan(1/2*f
*x+1/2*e)^2*a-tan(1/2*f*x+1/2*e)^2*b-a-b)-(2*a^2*d-a*b*c-b^2*d)/(a-b)/(a+b
))/(a-b)*(a+b)^(1/2)*arctanh((a-b)*tan(1/2*f*x+1/2*e)/((a-b)*(a+b))^(1/2)
))+2*d^2/(a*d-b*c)^2/((c+d)*(c-d))^(1/2)*arctanh((c-d)*tan(1/2*f*x+1/2*e)/
((c+d)*(c-d))^(1/2)))
```

3.263. $\int \frac{\sec(e+fx)}{(a+b \sec(e+fx))^2(c+d \sec(e+fx))} dx$

3.263.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 655 vs. $2(168) = 336$.

Time = 60.79 (sec) , antiderivative size = 2852, normalized size of antiderivative = 15.33

$$\int \frac{\sec(e+fx)}{(a+b\sec(e+fx))^2(c+d\sec(e+fx))} dx = \text{Too large to display}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `[-1/2*((a*b^3*c^3 - a*b^3*c*d^2 - (2*a^2*b^2 - b^4)*c^2*d + (2*a^2*b^2 - b^4)*d^3 + (a^2*b^2*c^3 - a^2*b^2*c*d^2 - (2*a^3*b - a*b^3)*c^2*d + (2*a^3*b - a*b^3)*d^3)*cos(f*x + e))*sqrt(a^2 - b^2)*log((2*a*b*cos(f*x + e) - (a^2 - 2*b^2)*cos(f*x + e)^2 - 2*sqrt(a^2 - b^2)*(b*cos(f*x + e) + a)*sin(f*x + e) + 2*a^2 - b^2)/(a^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + b^2)) - ((a^5 - 2*a^3*b^2 + a*b^4)*d^2*cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*d^2)*sqrt(c^2 - d^2)*log((2*c*d*cos(f*x + e) - (c^2 - 2*d^2)*cos(f*x + e)^2 + 2*sqrt(c^2 - d^2)*(d*cos(f*x + e) + c)*sin(f*x + e) + 2*c^2 - d^2)/(c^2*cos(f*x + e)^2 + 2*c*d*cos(f*x + e) + d^2)) + 2*((a^2*b^3 - b^5)*c^3 - (a^3*b^2 - a*b^4)*c^2*d - (a^2*b^3 - b^5)*c*d^2 + (a^3*b^2 - a*b^4)*d^3)*sin(f*x + e)/(((a^5*b^2 - 2*a^3*b^4 + a*b^6)*c^4 - 2*(a^6*b - 2*a^4*b^3 + a^2*b^5)*c^3*d + (a^7 - 3*a^5*b^2 + 3*a^3*b^4 - a*b^6)*c^2*d^2 + 2*(a^6*b - 2*a^4*b^3 + a^2*b^5)*c*d^3 - (a^7 - 2*a^5*b^2 + a^3*b^4)*d^4)*f*cos(f*x + e) + ((a^4*b^3 - 2*a^2*b^5 + b^7)*c^4 - 2*(a^5*b^2 - 2*a^3*b^4 + a*b^6)*c^3*d + (a^6*b - 3*a^4*b^3 + 3*a^2*b^5 - b^7)*c^2*d^2 + 2*(a^5*b^2 - 2*a^3*b^4 + a*b^6)*c*d^3 - (a^6*b - 2*a^4*b^3 + a^2*b^5)*d^4)*f), 1/2*(2*((a^5 - 2*a^3*b^2 + a*b^4)*d^2*cos(f*x + e) + (a^4*b - 2*a^2*b^3 + b^5)*d^2)*sqrt(-c^2 + d^2)*arctan(-sqrt(-c^2 + d^2)*(d*cos(f*x + e) + c)/((c^2 - d^2)*sin(f*x + e))) - (a*b^3*c^3 - a*b^3*c*d^2 - (2*a^2*b^2 - b^4)*c^2*d + (2*a^2*b^2 - b^4)*d^3 + (a^2*b^2*c^3 - a^2*b^2*c*d^2 - (2*a^3*b - a*b^3)*c^2*d + ...`

3.263.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sec(e+fx)}{(a+b\sec(e+fx))^2(c+d\sec(e+fx))} dx \\ &= \int \frac{\sec(e+fx)}{(a+b\sec(e+fx))^2(c+d\sec(e+fx))} dx \end{aligned}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))**2/(c+d*sec(f*x+e)),x)`

output `Integral(sec(e + f*x)/((a + b*sec(e + f*x))**2*(c + d*sec(e + f*x))), x)`

3.263.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sec(e + fx)}{(a + b\sec(e + fx))^2(c + d\sec(e + fx))} dx = \text{Exception raised: ValueError}$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c^2-4*d^2>0)', see 'assume?' f or more de`

3.263.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.77

$$\int \frac{\sec(e + fx)}{(a + b\sec(e + fx))^2(c + d\sec(e + fx))} dx$$

$$= 2 \left(\frac{\left(\pi \left[\frac{fx+e}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2c+2d) + \arctan \left(-\frac{c \tan(\frac{1}{2}fx + \frac{1}{2}e) - d \tan(\frac{1}{2}fx + \frac{1}{2}e)}{\sqrt{-c^2+d^2}} \right) \right) d^2}{(b^2c^2 - 2abcd + a^2d^2)\sqrt{-c^2+d^2}} + \frac{b^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{(a^2bc - b^3c - a^3d + ab^2d) \left(a \tan(\frac{1}{2}fx + \frac{1}{2}e) \right)^2 - b \tan(\frac{1}{2}fx + \frac{1}{2}e)} \right)$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))^2/(c+d*sec(f*x+e)),x, algorithm="giac")`

output $2*((\pi*\text{floor}(1/2*(f*x + e)/\pi + 1/2)*\text{sgn}(-2*c + 2*d) + \arctan(-(c*\tan(1/2*f*x + 1/2*e) - d*\tan(1/2*f*x + 1/2*e))/\sqrt{-c^2 + d^2}))*d^2/((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*\sqrt{-c^2 + d^2}) + b^2*\tan(1/2*f*x + 1/2*e)/((a^2*b*c - b^3*c - a^3*d + a*b^2*d)*(a*\tan(1/2*f*x + 1/2*e)^2 - b*\tan(1/2*f*x + 1/2*e)^2 - a - b)) - (a*b^2*c - 2*a^2*b*d + b^3*d)*(\pi*\text{floor}(1/2*(f*x + e)/\pi + 1/2)*\text{sgn}(2*a - 2*b) + \arctan((a*\tan(1/2*f*x + 1/2*e) - b*\tan(1/2*f*x + 1/2*e))/\sqrt{-a^2 + b^2}))/((a^2*b^2*c^2 - b^4*c^2 - 2*a^3*b*c*d + 2*a*b^3*c*d + a^4*d^2 - a^2*b^2*d^2)*\sqrt{-a^2 + b^2}))/f$

3.263.9 Mupad [B] (verification not implemented)

Time = 27.48 (sec) , antiderivative size = 20827, normalized size of antiderivative = 111.97

$$\int \frac{\sec(e + fx)}{(a + b\sec(e + fx))^2(c + d\sec(e + fx))} dx = \text{Too large to display}$$

input `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))^2*(c + d/cos(e + f*x))),x)`

output $(d^2*\text{atan}(((d^2*(c^2 - d^2)^{(1/2)}*((32*\tan(e/2 + (f*x)/2)*(a^6*d^5 + 2*b^6*d^5 - 2*a*b^5*d^5 - 2*a^5*b*d^5 - a^6*c*d^4 - 4*b^6*c*d^4 - a^2*b^4*c^5 - 5*a^2*b^4*d^5 + 4*a^3*b^3*d^5 + 3*a^4*b^2*d^5 + 3*b^6*c^2*d^3 - b^6*c^3*d^2 - 6*a*b^5*c^2*d^3 + 6*a*b^5*c^3*d^2 + 13*a^2*b^4*c*d^4 + 3*a^2*b^4*c^4*d - 8*a^3*b^3*c*d^4 + 4*a^3*b^3*c^4*d - 11*a^4*b^2*c*d^4 - 11*a^2*b^4*c^2*d^3 + a^2*b^4*c^3*d^2 + 12*a^3*b^3*c^2*d^3 - 12*a^3*b^3*c^3*d^2 + 12*a^4*b^2*c^2*d^3 - 4*a^4*b^2*c^3*d^2 + 4*a*b^5*c*d^4 - 2*a*b^5*c^4*d + 2*a^5*b*c*d^4))/(a^5*d^2 - b^5*c^2 - a*b^4*c^2 + a^4*b*d^2 + a^2*b^3*c^2 + a^3*b^2*c^2 - a^2*b^3*d^2 - a^3*b^2*d^2 + 2*a*b^4*c*d - 2*a^4*b*c*d + 2*a^2*b^3*c*d - 2*a^3*b^2*c*d) + (d^2*(c^2 - d^2)^{(1/2)}*((32*(a*b^8*c^7 - a^9*d^7 + 2*a^8*b*d^7 + 2*a^9*c*d^6 + b^9*c^6*d - a^2*b^7*c^7 - a^3*b^6*c^7 + a^4*b^5*c^7 + a^4*b^5*d^7 - 3*a^6*b^3*d^7 + a^7*b^2*d^7 - a^9*c^2*d^5 + b^9*c^4*d^3 - 2*b^9*c^5*d^2 - 4*a*b^8*c^3*d^4 + 8*a*b^8*c^4*d^3 - 3*a*b^8*c^5*d^2 - 5*a^2*b^7*c^6*d - 4*a^3*b^6*c*d^6 + 7*a^3*b^6*c^6*d - 2*a^4*b^5*c*d^6 + 4*a^4*b^5*c^6*d + 13*a^5*b^4*c*d^6 - 5*a^5*b^4*c^6*d + a^6*b^3*c*d^6 - 11*a^7*b^2*c*d^6 - 8*a^8*b*c^2*d^5 + 5*a^8*b*c^3*d^4 + 6*a^2*b^7*c^2*d^5 - 12*a^2*b^7*c^3*d^4 - a^2*b^7*c^4*d^3 + 13*a^2*b^7*c^5*d^2 + 8*a^3*b^6*c^2*d^5 + 14*a^3*b^6*c^3*d^4 - 31*a^3*b^6*c^4*d^3 + 7*a^3*b^6*c^5*d^2 - 21*a^4*b^5*c^2*d^5 + 34*a^4*b^5*c^3*d^4 + 4*a^4*b^5*c^4*d^3 - 21*a^4*b^5*c^5*d^2 - 16*a^5*b^4*c^2*d^5 - 21*a^5*b^4*c^3*d^4 + 33*a^5*b^4*c^4*d^3 - 4*a^5*b^4...$

3.264 $\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$

3.264.1 Optimal result 1942
 3.264.2 Mathematica [A] (verified) 1943
 3.264.3 Rubi [A] (verified) 1943
 3.264.4 Maple [A] (verified) 1945
 3.264.5 Fracas [F(-1)] 1946
 3.264.6 Sympy [F] 1946
 3.264.7 Maxima [F] 1946
 3.264.8 Giac [F] 1947
 3.264.9 Mupad [F(-1)] 1947

3.264.1 Optimal result

Integrand size = 33, antiderivative size = 213

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

$$= \frac{2\sqrt{a+b}\cot(e+fx)\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b\sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)\sqrt{\frac{b(1-\sec(e+fx))}{a+b}}\sqrt{-\frac{b(1+\sec(e+fx))}{a-b}}}{df} - \frac{2(bc-ad)\operatorname{EllipticPi}\left(\frac{2d}{c+d}, \arcsin\left(\frac{\sqrt{1-\sec(e+fx)}}{\sqrt{2}}\right), \frac{2b}{a+b}\right)\sqrt{\frac{a+b\sec(e+fx)}{a+b}}\tan(e+fx)}{d(c+d)f\sqrt{a+b\sec(e+fx)}\sqrt{-\tan^2(e+fx)}}$$

```
output 2*cot(f*x+e)*EllipticF((a+b*sec(f*x+e))^(1/2)/(a+b)^(1/2),((a+b)/(a-b))^(1/2))*(a+b)^(1/2)*(b*(1-sec(f*x+e))/(a+b))^(1/2)*(-b*(1+sec(f*x+e))/(a-b))^(1/2)/d/f-2*(-a*d+b*c)*EllipticPi(1/2*(1-sec(f*x+e))^(1/2)*2^(1/2),2*d/(c+d),2^(1/2)*(b/(a+b))^(1/2))*((a+b*sec(f*x+e))/(a+b))^(1/2)*tan(f*x+e)/d/(c+d)/f/(a+b*sec(f*x+e))^(1/2)/(-tan(f*x+e)^2)^(1/2)
```

3.264.2 Mathematica [A] (verified)

Time = 4.10 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.86

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

$$= \frac{4 \cos^2\left(\frac{1}{2}(e+fx)\right) \sqrt{\frac{\cos(e+fx)}{1+\cos(e+fx)}} \sqrt{\frac{b+a\cos(e+fx)}{(a+b)(1+\cos(e+fx))}} \left((a-b)(c+d) \operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right), \frac{a-b}{a+b}\right) \right)}{(c-d)(c+d)f(b+a\cos(e+fx))}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`output `(4*Cos[(e + f*x)/2]^2*Sqrt[Cos[e + f*x]/(1 + Cos[e + f*x])] * Sqrt[(b + a*Cos[e + f*x])/((a + b)*(1 + Cos[e + f*x]))] * ((a - b)*(c + d)*EllipticF[ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)] + 2*(b*c - a*d)*EllipticPi[(c - d)/(c + d), ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)]) * Sqrt[a + b*Sec[e + f*x]])/((c - d)*(c + d)*f*(b + a*Cos[e + f*x]))`**3.264.3 Rubi [A] (verified)**Time = 0.71 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {3042, 4457, 3042, 4319, 4461}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\sqrt{a+b\csc\left(e+fx+\frac{\pi}{2}\right)}}{c+d\csc\left(e+fx+\frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4457}$$

$$\frac{b \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}} dx}{d} - \frac{(bc-ad) \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx}{d}$$

$$\downarrow \text{3042}$$

$$\frac{b \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})}} dx}{d} - \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d}$$

↓ 4319

$$\frac{2\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{df} - \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d}$$

↓ 4461

$$\frac{2\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{df} - \frac{2(bc-ad) \tan(e+fx) \sqrt{\frac{a+b \sec(e+fx)}{a+b}} \text{EllipticPi}\left(\frac{2d}{c+d}, \arcsin\left(\frac{\sqrt{1-\sec(e+fx)}}{\sqrt{2}}\right), \frac{2b}{a+b}\right)}{df(c+d) \sqrt{-\tan^2(e+fx)} \sqrt{a+b \sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`

output `(2*Sqrt[a + b]*Cot[e + f*x]*EllipticF[ArcSin[Sqrt[a + b*Sec[e + f*x]]/Sqrt[a + b]], (a + b)/(a - b)]*Sqrt[(b*(1 - Sec[e + f*x]))/(a + b)]*Sqrt[-((b*(1 + Sec[e + f*x]))/(a - b))])/(d*f) - (2*(b*c - a*d)*EllipticPi[(2*d)/(c + d), ArcSin[Sqrt[1 - Sec[e + f*x]]/Sqrt[2]], (2*b)/(a + b)]*Sqrt[(a + b*Sec[e + f*x])/(a + b)]*Tan[e + f*x])/(d*(c + d)*f*Sqrt[a + b*Sec[e + f*x]]*Sqrt[-Tan[e + f*x]^2])`

3.264.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4319 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*(Rt[a + b, 2]/(b*f*Cot[e + f*x]))*Sqrt[(b*(1 - Csc[e + f*x]))/(a + b)]*Sqrt[(-b)*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Csc[e + f*x]]/Rt[a + b, 2]], (a + b)/(a - b)], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4457 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[b/d Int[Csc[e + f*x]/Sqrt[a + b*Csc[e + f*x]], x, x] - Simp[(b*c - a*d)/d Int[Csc[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x])), x, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4461 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))), x_Symbol] := Simp[-2*(Cot[e + f*x]/(f*(c + d)*Sqrt[a + b*Csc[e + f*x]]*Sqrt[-Cot[e + f*x]^2]))*Sqrt[(a + b*Csc[e + f*x])]/(a + b)*EllipticPi[2*(d/(c + d)), ArcSin[Sqrt[1 - Csc[e + f*x]]]/Sqrt[2]], 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.264.4 Maple [A] (verified)

Time = 9.15 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.48

method	result
default	$-\frac{2(\cos(fx+e)+1)\left(\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)ac+\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)ad-\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)bc-\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)bd-2\text{EllipticPi}\left(\cot(fx+e)-\csc(fx+e),\frac{c-d}{c+d},\sqrt{\frac{a-b}{a+b}}\right)a*d+2\text{EllipticPi}\left(\cot(fx+e)-\csc(fx+e),\frac{c-d}{c+d},\sqrt{\frac{a-b}{a+b}}\right)b*c\right)}{(a+b)\sqrt{a+b}\sqrt{\cos(fx+e)+1}}\frac{1}{(a+b)\sqrt{a+b}\sqrt{\cos(fx+e)+1}}\frac{1}{(a+b)\sqrt{a+b}\sqrt{\cos(fx+e)+1}}$

input `int(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `-2/f/(c+d)/(c-d)*(cos(f*x+e)+1)*(EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*a*c+EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*a*d-EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*b*c-EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*b*d-2*EllipticPi(cot(f*x+e)-csc(f*x+e),(c-d)/(c+d),((a-b)/(a+b))^(1/2))*a*d+2*EllipticPi(cot(f*x+e)-csc(f*x+e),(c-d)/(c+d),((a-b)/(a+b))^(1/2))*b*c)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*(a+b*sec(f*x+e))^(1/2)/(b+a*cos(f*x+e))`

3.264.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `Timed out`

3.264.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{a+b\sec(e+fx)}\sec(e+fx)}{c+d\sec(e+fx)} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e)),x)`

output `Integral(sqrt(a + b*sec(e + f*x))*sec(e + f*x)/(c + d*sec(e + f*x)), x)`

3.264.7 Maxima [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{b\sec(fx+e)+a}\sec(fx+e)}{d\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(d*sec(f*x + e) + c), x)`

3.264.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{b\sec(fx+e)+a}\sec(fx+e)}{d\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="giac")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(d*sec(f*x + e) + c), x)`

3.264.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{a + \frac{b}{\cos(e+fx)}}}{\cos(e+fx) \left(c + \frac{d}{\cos(e+fx)}\right)} dx$$

input `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))),x)`

output `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))), x)`

3.265
$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$$

3.265.1 Optimal result 1948
 3.265.2 Mathematica [C] (warning: unable to verify) 1948
 3.265.3 Rubi [A] (verified) 1949
 3.265.4 Maple [A] (verified) 1950
 3.265.5 Fricas [F(-1)] 1951
 3.265.6 Sympy [F] 1951
 3.265.7 Maxima [F] 1951
 3.265.8 Giac [F] 1952
 3.265.9 Mupad [F(-1)] 1952

3.265.1 Optimal result

Integrand size = 35, antiderivative size = 196

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$$

$$= \frac{2 \cot(e+fx) \operatorname{EllipticPi}\left(\frac{b(c+d)}{(a+b)d}, \arcsin\left(\frac{\sqrt{\frac{a+b}{c+d}}\sqrt{c+d\sec(e+fx)}}{\sqrt{a+b\sec(e+fx)}}\right), \frac{(a-b)(c+d)}{(a+b)(c-d)}\right) \sqrt{-\frac{(bc-ad)(1-\sec(e+fx))}{(c+d)(a+b\sec(e+fx))}} \sqrt{\frac{(bc-ad)(1-\sec(e+fx))}{(c-d)(a+b\sec(e+fx))}}}{d\sqrt{\frac{a+b}{c+d}}f}$$

```
output 2*cot(f*x+e)*EllipticPi(((a+b)/(c+d))^(1/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*sec(f*x+e))^(1/2),b*(c+d)/(a+b)/d,((a-b)*(c+d)/(a+b)/(c-d))^(1/2))*(a+b*sec(f*x+e))*(-(-a*d+b*c)*(1-sec(f*x+e))/(c+d)/(a+b*sec(f*x+e)))^(1/2)*((-a*d+b*c)*(1+sec(f*x+e))/(c-d)/(a+b*sec(f*x+e)))^(1/2)/d/f/((a+b)/(c+d))^(1/2)
```

3.265.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 37.86 (sec) , antiderivative size = 44664, normalized size of antiderivative = 227.88

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \text{Result too large to show}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/Sqrt[c + d*Sec[e + f*x]],x]`

output `Result too large to show`

3.265.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.057$, Rules used = {3042, 4470}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)\sqrt{a + b\sec(e + fx)}}{\sqrt{c + d\sec(e + fx)}} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right)\sqrt{a + b\csc\left(e + fx + \frac{\pi}{2}\right)}}{\sqrt{c + d\csc\left(e + fx + \frac{\pi}{2}\right)}} dx$$

↓ 4470

$$\frac{2 \cot(e + fx)(a + b\sec(e + fx))\sqrt{-\frac{(bc-ad)(1-\sec(e+fx))}{(c+d)(a+b\sec(e+fx))}}\sqrt{\frac{(bc-ad)(\sec(e+fx)+1)}{(c-d)(a+b\sec(e+fx))}} \text{EllipticPi}\left(\frac{b(c+d)}{(a+b)d}, \arcsin\left(\frac{\sqrt{\frac{a+b}{c+d}}\sqrt{c+d}}{\sqrt{a+b\sec(e+fx)}}\right)\right)}{df\sqrt{\frac{a+b}{c+d}}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/Sqrt[c + d*Sec[e + f*x]],x]`

output `(2*Cot[e + f*x]*EllipticPi[(b*(c + d))/((a + b)*d), ArcSin[(Sqrt[(a + b)/(c + d)]*Sqrt[c + d*Sec[e + f*x]])/Sqrt[a + b*Sec[e + f*x]]], ((a - b)*(c + d))/((a + b)*(c - d)))*Sqrt[-((b*c - a*d)*(1 - Sec[e + f*x]))/((c + d)*(a + b*Sec[e + f*x]))]*Sqrt[((b*c - a*d)*(1 + Sec[e + f*x]))/((c - d)*(a + b*Sec[e + f*x]))]*(a + b*Sec[e + f*x])]/(d*Sqrt[(a + b)/(c + d)]*f)`

3.265. $\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$

3.265.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4470 Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)], x_Symbol] :> Simp[-2*((a + b*Csc[e + f*x])/(d*f*Sqrt[(a + b)/(c + d)]*Cot[e + f*x]))*Sqrt[(-(b*c - a*d))*((1 - Csc[e + f*x])/((c + d)*(a + b*Csc[e + f*x])))]*Sqrt[(b*c - a*d)*((1 + Csc[e + f*x])/((c - d)*(a + b*Csc[e + f*x])))]*EllipticPi[b*((c + d)/(d*(a + b))), ArcSin[Sqrt[(a + b)/(c + d)]*(Sqrt[c + d*Csc[e + f*x]]/Sqrt[a + b*Csc[e + f*x]])], (a - b)*((c + d)/((a + b)*(c - d))), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.265.4 Maple [A] (verified)

Time = 12.84 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.64

method	result
default	$2\sqrt{\frac{d+c\cos(fx+e)}{(c+d)(\cos(fx+e)+1)}}\sqrt{a+b\sec(fx+e)}\sqrt{c+d\sec(fx+e)}\sqrt{\frac{b+a\cos(fx+e)}{(a+b)(\cos(fx+e)+1)}}\left(\text{EllipticF}\left(\sqrt{\frac{a-b}{a+b}}(-\cot(fx+e)+\csc(fx+e)),\sqrt{\frac{a}{a+b}}\right)\right)$

```
input int(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 2/f/((a-b)/(a+b))^(1/2)*(1/(c+d)*(d+c*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(a+b*sec(f*x+e))^(1/2)*(c+d*sec(f*x+e))^(1/2)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(EllipticF(((a-b)/(a+b))^(1/2)*(-cot(f*x+e)+csc(f*x+e)), ((a+b)*(c-d)/(a-b)/(c+d))^(1/2))*a-EllipticF(((a-b)/(a+b))^(1/2)*(-cot(f*x+e)+csc(f*x+e)), ((a+b)*(c-d)/(a-b)/(c+d))^(1/2))*b+2*EllipticPi(((a-b)/(a+b))^(1/2)*(-cot(f*x+e)+csc(f*x+e)), (a+b)/(a-b), ((c-d)/(c+d))^(1/2)/((a-b)/(a+b))^(1/2))*b)/(d+c*cos(f*x+e))/(b+a*cos(f*x+e))*(cos(f*x+e)^2+cos(f*x+e))
```

3.265. $\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx$

3.265.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)\sqrt{a + b\sec(e + fx)}}{\sqrt{c + d\sec(e + fx)}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="fricas")`

output `Timed out`

3.265.6 Sympy [F]

$$\int \frac{\sec(e + fx)\sqrt{a + b\sec(e + fx)}}{\sqrt{c + d\sec(e + fx)}} dx = \int \frac{\sqrt{a + b\sec(e + fx)}\sec(e + fx)}{\sqrt{c + d\sec(e + fx)}} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e))**(1/2),x)`

output `Integral(sqrt(a + b*sec(e + f*x))*sec(e + f*x)/sqrt(c + d*sec(e + f*x)), x
)`

3.265.7 Maxima [F]

$$\int \frac{\sec(e + fx)\sqrt{a + b\sec(e + fx)}}{\sqrt{c + d\sec(e + fx)}} dx = \int \frac{\sqrt{b\sec(fx + e) + a}\sec(fx + e)}{\sqrt{d\sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="maxima")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/sqrt(d*sec(f*x + e) + c),
x)`

3.265.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \int \frac{\sqrt{b\sec(fx+e)+a}\sec(fx+e)}{\sqrt{d\sec(fx+e)+c}} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorith="giac")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/sqrt(d*sec(f*x + e) + c), x)`

3.265.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sqrt{c+d\sec(e+fx)}} dx = \int \frac{\sqrt{a + \frac{b}{\cos(e+fx)}}}{\cos(e+fx) \sqrt{c + \frac{d}{\cos(e+fx)}}} dx$$

input `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))^(1/2)),x)`

output `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))^(1/2)),x)`

3.266 $\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx$

3.266.1 Optimal result	1953
3.266.2 Mathematica [A] (verified)	1953
3.266.3 Rubi [A] (verified)	1954
3.266.4 Maple [A] (verified)	1955
3.266.5 Fracas [F]	1956
3.266.6 Sympy [F]	1956
3.266.7 Maxima [F]	1956
3.266.8 Giac [F]	1957
3.266.9 Mupad [F(-1)]	1957

3.266.1 Optimal result

Integrand size = 35, antiderivative size = 192

$$\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx = \frac{2\sqrt{a+b} \cot(e+fx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{c+d}\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}\sqrt{c+d \sec(e+fx)}}\right), \frac{(a+b)(c-d)}{(a-b)(c+d)}\right) \sqrt{\frac{(bc-ad)(1-\sec(e+fx))}{(a+b)(c+d \sec(e+fx))}} \sqrt{-\frac{(bc-ad)(1+\sec(e+fx))}{(a-b)(c+d \sec(e+fx))}}}{\sqrt{c+d}(bc-ad)f}$$

output

```
2*cot(f*x+e)*EllipticF((c+d)^(1/2)*(a+b*sec(f*x+e))^(1/2)/(a+b)^(1/2)/(c+d*sec(f*x+e))^(1/2),((a+b)*(c-d)/(a-b)/(c+d))^(1/2)*(c+d*sec(f*x+e))*(a+b)^(1/2)*((-a*d+b*c)*(1-sec(f*x+e))/(a+b)/(c+d*sec(f*x+e)))^(1/2)*(-(-a*d+b*c)*(1+sec(f*x+e))/(a-b)/(c+d*sec(f*x+e)))^(1/2)/(-a*d+b*c)/f/(c+d)^(1/2)
```

3.266.2 Mathematica [A] (verified)

Time = 3.20 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.21

$$\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx = \frac{4\sqrt{\frac{(c+d) \cot^2(\frac{1}{2}(e+fx))}{c-d}} \sqrt{\frac{(a+b)(d+c \cos(e+fx)) \csc^2(\frac{1}{2}(e+fx))}{-bc+ad}} \csc(e+fx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{\frac{(a+b)(d+c \cos(e+fx)) \csc^2(\frac{1}{2}(e+fx))}{-bc+ad}}}{\sqrt{2}}\right)\right)}{(a+b)f \sqrt{\frac{(c+d)(b+a \cos(e+fx)) \csc^2(\frac{1}{2}(e+fx))}{bc-ad}} \sqrt{c+d \sec(e+fx)}}$$

3.266. $\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx$

input `Integrate[Sec[e + f*x]/(Sqrt[a + b*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `(4*Sqrt[((c + d)*Cot[(e + f*x)/2]^2)/(c - d])*Sqrt[((a + b)*(d + c*Cos[e + f*x])*Csc[(e + f*x)/2]^2)/(-b*c) + a*d])*Csc[e + f*x]*EllipticF[ArcSin[Sqrt[((a + b)*(d + c*Cos[e + f*x])*Csc[(e + f*x)/2]^2)/(-b*c) + a*d)]/Sqrt[2]], (2*(b*c - a*d))/((a + b)*(c - d))]*Sqrt[a + b*Sec[e + f*x]]*Sin[(e + f*x)/2]^2)/((a + b)*f*Sqrt[((c + d)*(b + a*Cos[e + f*x])*Csc[(e + f*x)/2]^2)/(b*c - a*d)]*Sqrt[c + d*Sec[e + f*x]])]`

3.266.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.057$, Rules used = {3042, 4472}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

↓ 3042

$$\int \frac{\csc\left(e + fx + \frac{\pi}{2}\right)}{\sqrt{a + b \csc\left(e + fx + \frac{\pi}{2}\right)} \sqrt{c + d \csc\left(e + fx + \frac{\pi}{2}\right)}} dx$$

↓ 4472

$$\frac{2\sqrt{a+b} \cot(e+fx)(c+d \sec(e+fx)) \sqrt{\frac{(bc-ad)(1-\sec(e+fx))}{(a+b)(c+d \sec(e+fx))}} \sqrt{-\frac{(bc-ad)(\sec(e+fx)+1)}{(a-b)(c+d \sec(e+fx))}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{c+d} \sqrt{a+b \sec(e+fx)}}{\sqrt{a+b} \sqrt{c+d \sec(e+fx)}}\right)\right)}{f \sqrt{c+d} (bc-ad)}$$

input `Int[Sec[e + f*x]/(Sqrt[a + b*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `(2*Sqrt[a + b]*Cot[e + f*x]*EllipticF[ArcSin[(Sqrt[c + d]*Sqrt[a + b*Sec[e + f*x]])/(Sqrt[a + b]*Sqrt[c + d*Sec[e + f*x]])], ((a + b)*(c - d))/((a - b)*(c + d))]*Sqrt[((b*c - a*d)*(1 - Sec[e + f*x]))/((a + b)*(c + d*Sec[e + f*x]))]*Sqrt[-(((b*c - a*d)*(1 + Sec[e + f*x]))/((a - b)*(c + d*Sec[e + f*x])))]*(c + d*Sec[e + f*x]))/(Sqrt[c + d]*(b*c - a*d)*f)`

3.266. $\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx$

3.266.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4472 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]), x_Symbol] := Simp[-2*((c + d*Csc[e + f*x])/(f*(b*c - a*d)*Rt[(c + d)/(a + b), 2]*Cot[e + f*x]))*Sqrt[(b*c - a*d)*((1 - Csc[e + f*x])/((a + b)*(c + d*Csc[e + f*x])))]*Sqrt[(-(b*c - a*d))*((1 + Csc[e + f*x])/((a - b)*(c + d*Csc[e + f*x])))]*EllipticF[ArcSin[Rt[(c + d)/(a + b), 2]*(Sqrt[a + b*Csc[e + f*x]]/Sqrt[c + d*Csc[e + f*x]])], (a + b)*((c - d)/((a - b)*(c + d))), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.266.4 Maple [A] (verified)

Time = 10.71 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.99

method	result
default	$\frac{2\sqrt{c+d\sec(fx+e)}\sqrt{a+b\sec(fx+e)}\sqrt{\frac{b+a\cos(fx+e)}{(a+b)(\cos(fx+e)+1)}}\sqrt{\frac{d+c\cos(fx+e)}{(c+d)(\cos(fx+e)+1)}}\text{EllipticF}\left(\sqrt{\frac{a-b}{a+b}}(-\cot(fx+e)+\csc(fx+e)),\sqrt{\frac{a-b}{a-b}}\right)}{f\sqrt{\frac{a-b}{a+b}}(d+c\cos(fx+e))(b+a\cos(fx+e))}$

input `int(sec(f*x+e)/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{2/f/((a-b)/(a+b))^{1/2}*(c+d*\sec(f*x+e))^{1/2}*(a+b*\sec(f*x+e))^{1/2}*(1/(a+b)*(b+a*\cos(f*x+e))/(\cos(f*x+e)+1))^{1/2}*(1/(c+d)*(d+c*\cos(f*x+e))/(\cos(f*x+e)+1))^{1/2}*\text{EllipticF}(((a-b)/(a+b))^{1/2}*(-\cot(f*x+e)+\csc(f*x+e)),(a+b)*(c-d)/(a-b)/(c+d))^{1/2}/(d+c*\cos(f*x+e))/(b+a*\cos(f*x+e))*(\cos(f*x+e)^2+\cos(f*x+e))$$

3.266.
$$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx$$

3.266.5 Fricas [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{b \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="fricas")`

output `integral(sqrt(b*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)*sec(f*x + e)/(b
*d*sec(f*x + e)^2 + a*c + (b*c + a*d)*sec(f*x + e)), x)`

3.266.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(a + b*sec(e + f*x))*sqrt(c + d*sec(e + f*x))),
x)`

3.266.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{b \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="maxima")`

output `integrate(sec(f*x + e)/(sqrt(b*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c))
, x)`

3.266.8 Giac [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{b \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algo
rithm="giac")`

output `integrate(sec(f*x + e)/(sqrt(b*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c))
, x)`

3.266.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx) \sqrt{a + \frac{b}{\cos(e + fx)}} \sqrt{c + \frac{d}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2))
,x)`

output `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2))
, x)`

$$3.267 \quad \int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx$$

3.267.1 Optimal result	1958
3.267.2 Mathematica [A] (verified)	1958
3.267.3 Rubi [A] (verified)	1959
3.267.4 Maple [C] (verified)	1960
3.267.5 Fricas [F]	1961
3.267.6 Sympy [F]	1961
3.267.7 Maxima [F]	1961
3.267.8 Giac [F]	1962
3.267.9 Mupad [F(-1)]	1962

3.267.1 Optimal result

Integrand size = 35, antiderivative size = 110

$$\int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx = \frac{2 \cot(e+fx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{2+3\sec(e+fx)}}{\sqrt{5}\sqrt{-4+5\sec(e+fx)}}\right), 45\right) (4-5\sec(e+fx)) \sqrt{\frac{1-\sec(e+fx)}{4-5\sec(e+fx)}} \sqrt{\frac{1+\sec(e+fx)}{4-5\sec(e+fx)}}}{f}$$

```
output 2*cot(f*x+e)*EllipticF(1/5*(2+3*sec(f*x+e))^(1/2)*5^(1/2)/(-4+5*sec(f*x+e))^(1/2),3*5^(1/2))*(4-5*sec(f*x+e))*((1-sec(f*x+e))/(4-5*sec(f*x+e)))^(1/2)*((1+sec(f*x+e))/(4-5*sec(f*x+e)))^(1/2)/f
```

3.267.2 Mathematica [A] (verified)

Time = 2.65 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.60

$$\int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx = \frac{4\sqrt{-\cot^2\left(\frac{1}{2}(e+fx)\right)}\sqrt{-\left((3+2\cos(e+fx))\csc^2\left(\frac{1}{2}(e+fx)\right)\right)}\sqrt{-\left((-5+4\cos(e+fx))\csc^2\left(\frac{1}{2}(e+fx)\right)\right)}}{3\sqrt{5}f\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}}$$

```
input Integrate[Sec[e + f*x]/(Sqrt[2 + 3*Sec[e + f*x]]*Sqrt[-4 + 5*Sec[e + f*x]]),x]
```

$$3.267. \quad \int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx$$

output $(-4*\text{Sqrt}[-\text{Cot}[(e + f*x)/2]^2]*\text{Sqrt}[-((3 + 2*\text{Cos}[e + f*x])* \text{Csc}[(e + f*x)/2]^2)]*\text{Sqrt}[-((-5 + 4*\text{Cos}[e + f*x])* \text{Csc}[(e + f*x)/2]^2)]*\text{Csc}[e + f*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[5/22]*\text{Sqrt}[(-5 + 4*\text{Cos}[e + f*x])/(-1 + \text{Cos}[e + f*x])]], 44/45]*\text{Sec}[e + f*x]*\text{Sin}[(e + f*x)/2]^4)/(3*\text{Sqrt}[5]*f*\text{Sqrt}[2 + 3*\text{Sec}[e + f*x]]*\text{Sqrt}[-4 + 5*\text{Sec}[e + f*x]])$

3.267.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.057$, Rules used = {3042, 4472}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{\sqrt{3 \sec(e + fx) + 2} \sqrt{5 \sec(e + fx) - 4}} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{3 \csc(e + fx + \frac{\pi}{2}) + 2} \sqrt{5 \csc(e + fx + \frac{\pi}{2}) - 4}} dx$$

↓ 4472

$$\frac{2 \cot(e + fx)(4 - 5 \sec(e + fx)) \sqrt{\frac{1 - \sec(e + fx)}{4 - 5 \sec(e + fx)}} \sqrt{\frac{\sec(e + fx) + 1}{4 - 5 \sec(e + fx)}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{3 \sec(e + fx) + 2}}{\sqrt{5} \sqrt{5 \sec(e + fx) - 4}}\right), 45\right)}{f}$$

input $\text{Int}[\text{Sec}[e + f*x]/(\text{Sqrt}[2 + 3*\text{Sec}[e + f*x]]*\text{Sqrt}[-4 + 5*\text{Sec}[e + f*x]]),x]$

output $(2*\text{Cot}[e + f*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[2 + 3*\text{Sec}[e + f*x]]/(\text{Sqrt}[5]*\text{Sqrt}[-4 + 5*\text{Sec}[e + f*x]])], 45]*(4 - 5*\text{Sec}[e + f*x])* \text{Sqrt}[(1 - \text{Sec}[e + f*x])/(4 - 5*\text{Sec}[e + f*x])]*\text{Sqrt}[(1 + \text{Sec}[e + f*x])/(4 - 5*\text{Sec}[e + f*x])])/f$

3.267.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4472 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]), x_Symbol] := Simp[-2*((c + d*Csc[e + f*x])/(f*(b*c - a*d)*Rt[(c + d)/(a + b), 2]*Cot[e + f*x]))*Sqrt[(b*c - a*d)*((1 - Csc[e + f*x])/((a + b)*(c + d*Csc[e + f*x])))]*Sqrt[(-(b*c - a*d))*((1 + Csc[e + f*x])/((a - b)*(c + d*Csc[e + f*x])))]*EllipticF[ArcSin[Rt[(c + d)/(a + b), 2]*(Sqrt[a + b*Csc[e + f*x]]/Sqrt[c + d*Csc[e + f*x]])], (a + b)*((c - d)/((a - b)*(c + d)))]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.267.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 7.54 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.35

method	result
default	$-\frac{i\sqrt{5} \operatorname{EllipticF}\left(\frac{i\sqrt{5}(\cot(fx+e)-\csc(fx+e))}{5}, 3\sqrt{5}\right) \sqrt{2+3\sec(fx+e)} \sqrt{-4+5\sec(fx+e)} \sqrt{-\frac{2(4\cos(fx+e)-5)}{\cos(fx+e)+1}} \sqrt{10} \sqrt{\frac{2\cos(fx+e)+3}{\cos(fx+e)+1}}}{5f(8\cos(fx+e)^2+2\cos(fx+e)-15)}$

input `int(sec(f*x+e)/(2+3*sec(f*x+e))^(1/2)/(-4+5*sec(f*x+e))^(1/2), x, method=_RE
TURNVERBOSE)`

output `-1/5*I/f*5^(1/2)*EllipticF(1/5*I*5^(1/2)*(cot(f*x+e)-csc(f*x+e)), 3*5^(1/2)
)*(2+3*sec(f*x+e))^(1/2)*(-4+5*sec(f*x+e))^(1/2)*(-2*(4*cos(f*x+e)-5)/(cos
(f*x+e)+1))^(1/2)*10^(1/2)*((2*cos(f*x+e)+3)/(cos(f*x+e)+1))^(1/2)/(8*cos
(f*x+e)^2+2*cos(f*x+e)-15)*(cos(f*x+e)^2+cos(f*x+e))`

$$3.267. \quad \int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx$$

3.267.5 Fricas [F]

$$\int \frac{\sec(e + fx)}{\sqrt{2 + 3 \sec(e + fx)} \sqrt{-4 + 5 \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{5 \sec(fx + e) - 4} \sqrt{3 \sec(fx + e) + 2}} dx$$

input `integrate(sec(f*x+e)/(2+3*sec(f*x+e))^(1/2)/(-4+5*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral(sqrt(5*sec(f*x + e) - 4)*sqrt(3*sec(f*x + e) + 2)*sec(f*x + e)/(15*sec(f*x + e)^2 - 2*sec(f*x + e) - 8), x)`

3.267.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{2 + 3 \sec(e + fx)} \sqrt{-4 + 5 \sec(e + fx)}} dx$$

$$= \int \frac{\sec(e + fx)}{\sqrt{3 \sec(e + fx) + 2} \sqrt{5 \sec(e + fx) - 4}} dx$$

input `integrate(sec(f*x+e)/(2+3*sec(f*x+e))**(1/2)/(-4+5*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(3*sec(e + f*x) + 2)*sqrt(5*sec(e + f*x) - 4)), x)`

3.267.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{\sqrt{2 + 3 \sec(e + fx)} \sqrt{-4 + 5 \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{5 \sec(fx + e) - 4} \sqrt{3 \sec(fx + e) + 2}} dx$$

input `integrate(sec(f*x+e)/(2+3*sec(f*x+e))^(1/2)/(-4+5*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/(sqrt(5*sec(f*x + e) - 4)*sqrt(3*sec(f*x + e) + 2)), x)`

3.267.8 Giac [F]

$$\begin{aligned} & \int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx \\ &= \int \frac{\sec(fx+e)}{\sqrt{5\sec(fx+e)-4}\sqrt{3\sec(fx+e)+2}} dx \end{aligned}$$

input `integrate(sec(f*x+e)/(2+3*sec(f*x+e))^(1/2)/(-4+5*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate(sec(f*x + e)/(sqrt(5*sec(f*x + e) - 4)*sqrt(3*sec(f*x + e) + 2)), x)`

3.267.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sec(e+fx)}{\sqrt{2+3\sec(e+fx)}\sqrt{-4+5\sec(e+fx)}} dx \\ &= \int \frac{1}{\cos(e+fx)\sqrt{\frac{3}{\cos(e+fx)}+2}\sqrt{\frac{5}{\cos(e+fx)}-4}} dx \end{aligned}$$

input `int(1/(cos(e + f*x)*(3/cos(e + f*x) + 2)^(1/2)*(5/cos(e + f*x) - 4)^(1/2)),x)`

output `int(1/(cos(e + f*x)*(3/cos(e + f*x) + 2)^(1/2)*(5/cos(e + f*x) - 4)^(1/2)), x)`

3.268 $\int \frac{\sec(e+fx)}{\sqrt{4-5\sec(e+fx)}\sqrt{2+3\sec(e+fx)}} dx$

3.268.1 Optimal result	1963
3.268.2 Mathematica [A] (verified)	1963
3.268.3 Rubi [A] (verified)	1964
3.268.4 Maple [A] (verified)	1965
3.268.5 Fricas [F]	1966
3.268.6 Sympy [F]	1966
3.268.7 Maxima [F]	1966
3.268.8 Giac [F]	1967
3.268.9 Mupad [F(-1)]	1967

3.268.1 Optimal result

Integrand size = 35, antiderivative size = 125

$$\int \frac{\sec(e+fx)}{\sqrt{4-5\sec(e+fx)}\sqrt{2+3\sec(e+fx)}} dx = \frac{2i \cot(e+fx) \operatorname{EllipticF}\left(\operatorname{iarcsinh}\left(\frac{\sqrt{5}\sqrt{4-5\sec(e+fx)}}{\sqrt{2+3\sec(e+fx)}}\right), \frac{1}{45}\right) \sqrt{\frac{1-\sec(e+fx)}{2+3\sec(e+fx)}} \sqrt{\frac{1+\sec(e+fx)}{2+3\sec(e+fx)}} (2+3\sec(e+fx))}{3\sqrt{5}f}$$

output

```
2/15*I*cot(f*x+e)*EllipticF(I*5^(1/2)*(4-5*sec(f*x+e))^(1/2)/(2+3*sec(f*x+e))^(1/2),1/15*5^(1/2))*(2+3*sec(f*x+e))*((1-sec(f*x+e))/(2+3*sec(f*x+e)))^(1/2)*((1+sec(f*x+e))/(2+3*sec(f*x+e)))^(1/2)/f*5^(1/2)
```

3.268.2 Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.41

$$\int \frac{\sec(e+fx)}{\sqrt{4-5\sec(e+fx)}\sqrt{2+3\sec(e+fx)}} dx = \frac{4\sqrt{-\cot^2\left(\frac{1}{2}(e+fx)\right)}\sqrt{-((3+2\cos(e+fx))\operatorname{csc}^2\left(\frac{1}{2}(e+fx)\right))}\sqrt{-((-5+4\cos(e+fx))\operatorname{csc}^2\left(\frac{1}{2}(e+fx)\right))}}{3\sqrt{5}f\sqrt{4-5\sec(e+fx)}}$$

input

```
Integrate[Sec[e + f*x]/(Sqrt[4 - 5*Sec[e + f*x]]*Sqrt[2 + 3*Sec[e + f*x]]),x]
```

3.268. $\int \frac{\sec(e+fx)}{\sqrt{4-5\sec(e+fx)}\sqrt{2+3\sec(e+fx)}} dx$

output $(-4*\text{Sqrt}[-\text{Cot}[(e + f*x)/2]^2]*\text{Sqrt}[-((3 + 2*\text{Cos}[e + f*x])* \text{Csc}[(e + f*x)/2]^2)]*\text{Sqrt}[-((-5 + 4*\text{Cos}[e + f*x])* \text{Csc}[(e + f*x)/2]^2)]*\text{Csc}[e + f*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[5/22]*\text{Sqrt}[-(5 + 4*\text{Cos}[e + f*x])/(-1 + \text{Cos}[e + f*x])]], 44/45]*\text{Sec}[e + f*x]*\text{Sin}[(e + f*x)/2]^4)/(3*\text{Sqrt}[5]*f*\text{Sqrt}[4 - 5*\text{Sec}[e + f*x]]*\text{Sqrt}[2 + 3*\text{Sec}[e + f*x]])$

3.268.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.057$, Rules used = {3042, 4472}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{\sqrt{4 - 5 \sec(e + fx)} \sqrt{3 \sec(e + fx) + 2}} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{4 - 5 \csc(e + fx + \frac{\pi}{2})} \sqrt{3 \csc(e + fx + \frac{\pi}{2}) + 2}} dx$$

↓ 4472

$$\frac{2i \cot(e + fx) \sqrt{\frac{1 - \sec(e + fx)}{3 \sec(e + fx) + 2}} \sqrt{\frac{\sec(e + fx) + 1}{3 \sec(e + fx) + 2}} (3 \sec(e + fx) + 2) \text{EllipticF}\left(i \operatorname{arcsinh}\left(\frac{\sqrt{5} \sqrt{4 - 5 \sec(e + fx)}}{\sqrt{3 \sec(e + fx) + 2}}\right), \frac{1}{45}\right)}{3\sqrt{5}f}$$

input $\text{Int}[\text{Sec}[e + f*x]/(\text{Sqrt}[4 - 5*\text{Sec}[e + f*x]]*\text{Sqrt}[2 + 3*\text{Sec}[e + f*x]]),x]$

output $((2*I)/3)*\text{Cot}[e + f*x]*\text{EllipticF}[I*\text{ArcSinh}[(\text{Sqrt}[5]*\text{Sqrt}[4 - 5*\text{Sec}[e + f*x]])/\text{Sqrt}[2 + 3*\text{Sec}[e + f*x]]], 1/45]*\text{Sqrt}[(1 - \text{Sec}[e + f*x])/ (2 + 3*\text{Sec}[e + f*x])] * \text{Sqrt}[(1 + \text{Sec}[e + f*x])/ (2 + 3*\text{Sec}[e + f*x])] * (2 + 3*\text{Sec}[e + f*x])]/(\text{Sqrt}[5]*f)$

3.268.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4472 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.)]), x_Symbol] := Simp[-2*((c + d*Csc[e + f*x])/(f*(b*c - a*d)*Rt[(c + d)/(a + b), 2]*Cot[e + f*x]))*Sqrt[(b*c - a*d)*((1 - Csc[e + f*x])/((a + b)*(c + d*Csc[e + f*x])))]*Sqrt[(-(b*c - a*d))*((1 + Csc[e + f*x])/((a - b)*(c + d*Csc[e + f*x])))]*EllipticF[ArcSin[Rt[(c + d)/(a + b), 2]*(Sqrt[a + b*Csc[e + f*x]]/Sqrt[c + d*Csc[e + f*x]])], (a + b)*((c - d)/((a - b)*(c + d)))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.268.4 Maple [A] (verified)

Time = 7.36 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.14

method	result
default	$-\frac{i\sqrt{2+3\sec(fx+e)}\sqrt{4-5\sec(fx+e)}\sqrt{-\frac{2(4\cos(fx+e)-5)}{\cos(fx+e)+1}}\sqrt{10}\sqrt{\frac{2\cos(fx+e)+3}{\cos(fx+e)+1}}\text{EllipticF}\left(3i(-\cot(fx+e)+\csc(fx+e)),\frac{\sqrt{5}}{15}\right)(\cos(fx+e)-1)}{15f(8\cos(fx+e)^2+2\cos(fx+e)-15)}$

input `int(sec(f*x+e)/(4-5*sec(f*x+e))^(1/2)/(2+3*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-1/15*I/f*(2+3*sec(f*x+e))^(1/2)*(4-5*sec(f*x+e))^(1/2)*(-2*(4*cos(f*x+e)-5)/(cos(f*x+e)+1))^(1/2)*10^(1/2)*((2*cos(f*x+e)+3)/(cos(f*x+e)+1))^(1/2)*EllipticF(3*I*(-cot(f*x+e)+csc(f*x+e)),1/15*5^(1/2))/(8*cos(f*x+e)^2+2*cos(f*x+e)-15)*(cos(f*x+e)^2+cos(f*x+e))`

$$3.268. \int \frac{\sec(e+fx)}{\sqrt{4-5\sec(e+fx)}\sqrt{2+3\sec(e+fx)}} dx$$

3.268.5 Fricas [F]

$$\int \frac{\sec(e + fx)}{\sqrt{4 - 5 \sec(e + fx)} \sqrt{2 + 3 \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{3 \sec(fx + e) + 2} \sqrt{-5 \sec(fx + e) + 4}} dx$$

input `integrate(sec(f*x+e)/(4-5*sec(f*x+e))^(1/2)/(2+3*sec(f*x+e))^(1/2),x, algo
rithm="fricas")`

output `integral(-sqrt(3*sec(f*x + e) + 2)*sqrt(-5*sec(f*x + e) + 4)*sec(f*x + e)/
(15*sec(f*x + e)^2 - 2*sec(f*x + e) - 8), x)`

3.268.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{4 - 5 \sec(e + fx)} \sqrt{2 + 3 \sec(e + fx)}} dx$$

$$= \int \frac{\sec(e + fx)}{\sqrt{4 - 5 \sec(e + fx)} \sqrt{3 \sec(e + fx) + 2}} dx$$

input `integrate(sec(f*x+e)/(4-5*sec(f*x+e))**(1/2)/(2+3*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(4 - 5*sec(e + f*x))*sqrt(3*sec(e + f*x) + 2)),
x)`

3.268.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{\sqrt{4 - 5 \sec(e + fx)} \sqrt{2 + 3 \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{3 \sec(fx + e) + 2} \sqrt{-5 \sec(fx + e) + 4}} dx$$

input `integrate(sec(f*x+e)/(4-5*sec(f*x+e))^(1/2)/(2+3*sec(f*x+e))^(1/2),x, algo
rithm="maxima")`

output `integrate(sec(f*x + e)/(sqrt(3*sec(f*x + e) + 2)*sqrt(-5*sec(f*x + e) + 4)
, x)`

3.268.8 Giac [F]

$$\int \frac{\sec(e + fx)}{\sqrt{4 - 5 \sec(e + fx)} \sqrt{2 + 3 \sec(e + fx)}} dx$$

$$= \int \frac{\sec(fx + e)}{\sqrt{3 \sec(fx + e) + 2} \sqrt{-5 \sec(fx + e) + 4}} dx$$

input `integrate(sec(f*x+e)/(4-5*sec(f*x+e))^(1/2)/(2+3*sec(f*x+e))^(1/2),x, algo
rithm="giac")`

output `integrate(sec(f*x + e)/(sqrt(3*sec(f*x + e) + 2)*sqrt(-5*sec(f*x + e) + 4)
, x)`

3.268.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{\sqrt{4 - 5 \sec(e + fx)} \sqrt{2 + 3 \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx) \sqrt{\frac{3}{\cos(e+fx)} + 2} \sqrt{4 - \frac{5}{\cos(e+fx)}}} dx$$

input `int(1/(cos(e + f*x)*(3/cos(e + f*x) + 2)^(1/2)*(4 - 5/cos(e + f*x))^(1/2))
,x)`

output `int(1/(cos(e + f*x)*(3/cos(e + f*x) + 2)^(1/2)*(4 - 5/cos(e + f*x))^(1/2))
, x)`

3.269
$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$$

3.269.1 Optimal result	1968
3.269.2 Mathematica [C] (warning: unable to verify)	1969
3.269.3 Rubi [A] (verified)	1969
3.269.4 Maple [A] (verified)	1971
3.269.5 Fracas [F(-1)]	1972
3.269.6 Sympy [F]	1972
3.269.7 Maxima [F]	1973
3.269.8 Giac [F]	1973
3.269.9 Mupad [F(-1)]	1973

3.269.1 Optimal result

Integrand size = 37, antiderivative size = 396

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}\sqrt{c+d \sec(e+fx)}} dx$$

$$= \frac{2 \cot(e+fx) \operatorname{EllipticPi}\left(\frac{b(c+d)}{(a+b)d}, \arcsin\left(\frac{\sqrt{\frac{a+b}{c+d}}\sqrt{c+d \sec(e+fx)}}{\sqrt{a+b \sec(e+fx)}}\right), \frac{(a-b)(c+d)}{(a+b)(c-d)}\right) \sqrt{-\frac{(bc-ad)(1-\sec(e+fx))}{(c+d)(a+b \sec(e+fx))}} \sqrt{\frac{(bc-ad)(1+\sec(e+fx))}{(c-d)(a+b \sec(e+fx))}}}{bd \sqrt{\frac{a+b}{c+d}} f}$$

$$= \frac{2a\sqrt{a+b} \cot(e+fx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{c+d}\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}\sqrt{c+d \sec(e+fx)}}\right), \frac{(a+b)(c-d)}{(a-b)(c+d)}\right) \sqrt{\frac{(bc-ad)(1-\sec(e+fx))}{(a+b)(c+d \sec(e+fx))}} \sqrt{-\frac{(bc-ad)(1+\sec(e+fx))}{(a-b)(c+d \sec(e+fx))}}}{b\sqrt{c+d}(bc-ad)f}$$

```
output 2*cot(f*x+e)*EllipticPi(((a+b)/(c+d))^(1/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*sec(f*x+e))^(1/2),b*(c+d)/(a+b)/d,((a-b)*(c+d)/(a+b)/(c-d))^(1/2))*(a+b*sec(f*x+e))*(-a*d+b*c)*(1-sec(f*x+e))/(c+d)/(a+b*sec(f*x+e))^(1/2)*((-a*d+b*c)*(1+sec(f*x+e))/(c-d)/(a+b*sec(f*x+e)))^(1/2)/b/d/f/((a+b)/(c+d))^(1/2)
-2*a*cot(f*x+e)*EllipticF((c+d)^(1/2)*(a+b*sec(f*x+e))^(1/2)/(a+b)^(1/2)/(c+d*sec(f*x+e))^(1/2),((a+b)*(c-d)/(a-b)/(c+d))^(1/2)*(c+d*sec(f*x+e))*(a+b)^(1/2)*((-a*d+b*c)*(1-sec(f*x+e))/(a+b)/(c+d*sec(f*x+e)))^(1/2)*((-a*d+b*c)*(1+sec(f*x+e))/(a-b)/(c+d*sec(f*x+e)))^(1/2)/b/(-a*d+b*c)/f/(c+d)^(1/2)
```

3.269.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 36.53 (sec) , antiderivative size = 39359, normalized size of antiderivative = 99.39

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx = \text{Result too large to show}$$

input `Integrate[Sec[e + f*x]^2/(Sqrt[a + b*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `Result too large to show`

3.269.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.135$, Rules used = {3042, 4473, 3042, 4470, 4472}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(e + fx + \frac{\pi}{2})^2}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})} \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{4473} \\ & \frac{\int \frac{\sec(e + fx) \sqrt{a + b \sec(e + fx)}}{\sqrt{c + d \sec(e + fx)}} dx}{b} - \frac{a \int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx}{b} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{\csc(e + fx + \frac{\pi}{2}) \sqrt{a + b \csc(e + fx + \frac{\pi}{2})}}{\sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx}{b} - \frac{a \int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})} \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx}{b} \\ & \quad \downarrow \text{4470} \end{aligned}$$

3.269. $\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$

$$\frac{2 \cot(e + fx)(a + b \sec(e + fx)) \sqrt{-\frac{(bc-ad)(1-\sec(e+fx))}{(c+d)(a+b \sec(e+fx))}} \sqrt{\frac{(bc-ad)(\sec(e+fx)+1)}{(c-d)(a+b \sec(e+fx))}} \operatorname{EllipticPi}\left(\frac{b(c+d)}{(a+b)d}, \arcsin\left(\frac{\sqrt{\frac{a+b}{c+d}} \sqrt{c+d}}{\sqrt{a+b \sec(e+fx)}}\right)\right)}{a \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} \sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx}$$

$bdf \sqrt{\frac{a+b}{c+d}}$
 \downarrow 4472

$$\frac{2 \cot(e + fx)(a + b \sec(e + fx)) \sqrt{-\frac{(bc-ad)(1-\sec(e+fx))}{(c+d)(a+b \sec(e+fx))}} \sqrt{\frac{(bc-ad)(\sec(e+fx)+1)}{(c-d)(a+b \sec(e+fx))}} \operatorname{EllipticPi}\left(\frac{b(c+d)}{(a+b)d}, \arcsin\left(\frac{\sqrt{\frac{a+b}{c+d}} \sqrt{c+d}}{\sqrt{a+b \sec(e+fx)}}\right)\right)}{2a\sqrt{a+b} \cot(e + fx)(c + d \sec(e + fx)) \sqrt{\frac{(bc-ad)(1-\sec(e+fx))}{(a+b)(c+d \sec(e+fx))}} \sqrt{-\frac{(bc-ad)(\sec(e+fx)+1)}{(a-b)(c+d \sec(e+fx))}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{c+d} \sqrt{a+b}}{\sqrt{a+b} \sqrt{c+d}}\right)\right)}{bf\sqrt{c+d}(bc - ad)}$$

input `Int[Sec[e + f*x]^2/(Sqrt[a + b*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]]),x]`

output `(2*Cot[e + f*x]*EllipticPi[(b*(c + d))/((a + b)*d), ArcSin[(Sqrt[(a + b)/(c + d)]*Sqrt[c + d*Sec[e + f*x]])/Sqrt[a + b*Sec[e + f*x]]], ((a - b)*(c + d))/((a + b)*(c - d))*Sqrt[-((b*c - a*d)*(1 - Sec[e + f*x]))/((c + d)*(a + b*Sec[e + f*x]))])*Sqrt[((b*c - a*d)*(1 + Sec[e + f*x]))/((c - d)*(a + b*Sec[e + f*x]))]*(a + b*Sec[e + f*x])/(b*d*Sqrt[(a + b)/(c + d)]*f) - (2*a*Sqrt[a + b]*Cot[e + f*x]*EllipticF[ArcSin[(Sqrt[c + d]*Sqrt[a + b*Sec[e + f*x]])/(Sqrt[a + b]*Sqrt[c + d*Sec[e + f*x]])], ((a + b)*(c - d))/((a - b)*(c + d))*Sqrt[((b*c - a*d)*(1 - Sec[e + f*x]))/((a + b)*(c + d*Sec[e + f*x]))])*Sqrt[-((b*c - a*d)*(1 + Sec[e + f*x]))/((a - b)*(c + d*Sec[e + f*x]))]*(c + d*Sec[e + f*x])/(b*Sqrt[c + d]*(b*c - a*d)*f)`

3.269.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4470 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)], x_Symbol] := Simp[-2*((a + b*Csc[e + f*x])/(d*f*Sqrt[(a + b)/(c + d)]*Cot[e + f*x]))*Sqrt[(-(b*c - a*d))*((1 - Csc[e + f*x])/((c + d)*(a + b*Csc[e + f*x])))]*Sqrt[(b*c - a*d)*((1 + Csc[e + f*x])/((c - d)*(a + b*Csc[e + f*x])))]*EllipticPi[b*((c + d)/(d*(a + b))), ArcSin[Sqrt[(a + b)/(c + d)]*(Sqrt[c + d*Csc[e + f*x]]/Sqrt[a + b*Csc[e + f*x]])], (a - b)*((c + d)/((a + b)*(c - d))), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4472 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)], x_Symbol] := Simp[-2*((c + d*Csc[e + f*x])/(f*(b*c - a*d)*Rt[(c + d)/(a + b), 2]*Cot[e + f*x]))*Sqrt[(b*c - a*d)*((1 - Csc[e + f*x])/((a + b)*(c + d*Csc[e + f*x])))]*Sqrt[(-(b*c - a*d))*((1 + Csc[e + f*x])/((a - b)*(c + d*Csc[e + f*x])))]*EllipticF[ArcSin[Rt[(c + d)/(a + b), 2]*(Sqrt[a + b*Csc[e + f*x]]/Sqrt[c + d*Csc[e + f*x]])], (a + b)*((c - d)/((a - b)*(c + d))), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4473 `Int[csc[(e_.) + (f_.)*(x_)]^2/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)], x_Symbol] := Simp[-a/b Int[Csc[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*Sqrt[c + d*Csc[e + f*x]]), x], x] + Simp[1/b Int[Csc[e + f*x]*(Sqrt[a + b*Csc[e + f*x]]/Sqrt[c + d*Csc[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

3.269.4 Maple [A] (verified)

Time = 15.93 (sec) , antiderivative size = 262, normalized size of antiderivative = 0.66

method	result
default	$-\frac{2\sqrt{c+d}\sec(fx+e)\sqrt{a+b}\sec(fx+e)\left(\text{EllipticF}\left(\sqrt{\frac{a-b}{a+b}}(-\cot(fx+e)+\csc(fx+e)),\sqrt{\frac{(a+b)(c-d)}{(a-b)(c+d)}}\right)-2\text{EllipticPi}\left(\sqrt{\frac{a-b}{a+b}}(-\cot(fx+e)+\csc(fx+e)),\sqrt{\frac{(a+b)(c-d)}{(a-b)(c+d)}}\right)\right)}{f\sqrt{\frac{a-b}{a+b}}(d+c\cos(fx+e))(b+a\cos(fx+e))}$

3.269. $\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx$

input `int(sec(f*x+e)^2/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x,method=_RETURVERBOSE)`

output `-2/f/((a-b)/(a+b))^(1/2)*(c+d*sec(f*x+e))^(1/2)*(a+b*sec(f*x+e))^(1/2)*(EllipticF(((a-b)/(a+b))^(1/2)*(-cot(f*x+e)+csc(f*x+e)),((a+b)*(c-d)/(a-b)/(c+d))^(1/2))-2*EllipticPi(((a-b)/(a+b))^(1/2)*(-cot(f*x+e)+csc(f*x+e)),(a+b)/(a-b),((c-d)/(c+d))^(1/2)/((a-b)/(a+b))^(1/2)))*(1/(c+d)*(d+c*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)/(d+c*cos(f*x+e))/(b+a*cos(f*x+e))*(cos(f*x+e)^2+cos(f*x+e))`

3.269.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)^2/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x,algorithm="fricas")`

output `Timed out`

3.269.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx \\ &= \int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}\sqrt{c+d\sec(e+fx)}} dx \end{aligned}$$

input `integrate(sec(f*x+e)**2/(a+b*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)**2/(sqrt(a + b*sec(e + f*x))*sqrt(c + d*sec(e + f*x))), x)`

3.269.7 Maxima [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec^2(fx + e)}{\sqrt{b \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)^2/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)^2/(sqrt(b*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)), x)`

3.269.8 Giac [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{\sec^2(fx + e)}{\sqrt{b \sec(fx + e) + a} \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate(sec(f*x+e)^2/(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate(sec(f*x + e)^2/(sqrt(b*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)), x)`

3.269.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)} \sqrt{c + d \sec(e + fx)}} dx$$

$$= \int \frac{1}{\cos(e + fx)^2 \sqrt{a + \frac{b}{\cos(e + fx)}} \sqrt{c + \frac{d}{\cos(e + fx)}}} dx$$

input `int(1/(cos(e + f*x)^2*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2)),x)`

output `int(1/(cos(e + f*x)^2*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))^(1/2)), x)`

3.269. $\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)} \sqrt{c+d \sec(e+fx)}} dx$

3.270
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{c+d \sec(e+fx)}}{a+b \sec(e+fx)} dx$$

3.270.1 Optimal result	1975
3.270.2 Mathematica [C] (verified)	1975
3.270.3 Rubi [A] (verified)	1976
3.270.4 Maple [C] (verified)	1980
3.270.5 Fricas [F(-1)]	1980
3.270.6 Sympy [F]	1981
3.270.7 Maxima [F]	1981
3.270.8 Giac [F]	1981
3.270.9 Mupad [F(-1)]	1982

3.270.1 Optimal result

Integrand size = 39, antiderivative size = 170

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{c+d \sec(e+fx)}}{a+b \sec(e+fx)} dx = \frac{2dg \sqrt{\frac{d+c \cos(e+fx)}{c+d}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right) \sqrt{g \sec(e+fx)}}{bf \sqrt{c+d \sec(e+fx)}} + \frac{2(bc-ad)g \sqrt{\frac{d+c \cos(e+fx)}{c+d}} \text{EllipticPi}\left(\frac{2a}{a+b}, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right) \sqrt{g \sec(e+fx)}}{b(a+b)f \sqrt{c+d \sec(e+fx)}}$$

output

```
2*d*g*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e), 2, 2^(1/2)*(c/(c+d))^(1/2))*((d+c*cos(f*x+e))/(c+d))^(1/2)*(g*sec(f*x+e))^(1/2)/b/f/(c+d*sec(f*x+e))^(1/2)+2*(-a*d+b*c)*g*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e), 2*a/(a+b), 2^(1/2)*(c/(c+d))^(1/2))*((d+c*cos(f*x+e))/(c+d))^(1/2)*(g*sec(f*x+e))^(1/2)/b/(a+b)/f/(c+d*sec(f*x+e))^(1/2)
```

3.270.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 25.51 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.31

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{c+d \sec(e+fx)}}{a+b \sec(e+fx)} dx = \frac{2ig \sqrt{-\frac{c(-1+\cos(e+fx))}{c+d}} \sqrt{\frac{c(1+\cos(e+fx))}{c-d}} \cot(e+fx) \left(\text{EllipticPi}\left(1 - \frac{c}{d}, i \operatorname{arcsinh}\left(\sqrt{\frac{1}{c-d}} \sqrt{d+c \cos(e+fx)}\right)\right) \right)}{b \sqrt{\frac{1}{c-d}} f \sqrt{c+d \sec(e+fx)}}$$

3.270.
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{c+d \sec(e+fx)}}{a+b \sec(e+fx)} dx$$

input `Integrate[((g*Sec[e + f*x])^(3/2)*Sqrt[c + d*Sec[e + f*x]])/(a + b*Sec[e + f*x]),x]`

output `((-2*I)*g*Sqrt[-((c*(-1 + Cos[e + f*x]))/(c + d))]*Sqrt[(c*(1 + Cos[e + f*x]))/(c - d)]*Cot[e + f*x]*(EllipticPi[1 - c/d, I*ArcSinh[Sqrt[(c - d)^(-1)]*Sqrt[d + c*Cos[e + f*x]]], (-c + d)/(c + d)] - EllipticPi[(a*(-c + d))/(-b*c) + a*d, I*ArcSinh[Sqrt[(c - d)^(-1)]*Sqrt[d + c*Cos[e + f*x]]], (-c + d)/(c + d)]*Sqrt[g*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])/(b*Sqrt[(c - d)^(-1)]*f*Sqrt[d + c*Cos[e + f*x]])`

3.270.3 Rubi [A] (verified)

Time = 2.05 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4459, 3042, 4346, 3042, 3286, 3042, 3284, 4463, 3042, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2} \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}}{a + b \csc(e + fx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4459} \\
 & \frac{(bc - ad) \int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx}{b} + \frac{d \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{c + d \sec(e + fx)}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(bc - ad) \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{(a + b \csc(e + fx + \frac{\pi}{2})) \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx}{b} + \frac{d \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx}{b} \\
 & \quad \downarrow \text{4346}
 \end{aligned}$$

3.270. $\int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx$

$$\begin{aligned}
& \frac{(bc - ad) \int \frac{(g \csc(e+fx + \frac{\pi}{2}))^{3/2}}{(a+b \csc(e+fx + \frac{\pi}{2})) \sqrt{c+d \csc(e+fx + \frac{\pi}{2})}} dx}{b} + \\
& \frac{dg \sqrt{g \sec(e+fx)} \sqrt{c \cos(e+fx) + d} \int \frac{\sec(e+fx)}{\sqrt{d+c \cos(e+fx)}} dx}{b \sqrt{c+d \sec(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(bc - ad) \int \frac{(g \csc(e+fx + \frac{\pi}{2}))^{3/2}}{(a+b \csc(e+fx + \frac{\pi}{2})) \sqrt{c+d \csc(e+fx + \frac{\pi}{2})}} dx}{b} + \\
& \frac{dg \sqrt{g \sec(e+fx)} \sqrt{c \cos(e+fx) + d} \int \frac{1}{\sin(e+fx + \frac{\pi}{2}) \sqrt{d+c \sin(e+fx + \frac{\pi}{2})}} dx}{b \sqrt{c+d \sec(e+fx)}} \\
& \quad \downarrow \text{3286} \\
& \frac{(bc - ad) \int \frac{(g \csc(e+fx + \frac{\pi}{2}))^{3/2}}{(a+b \csc(e+fx + \frac{\pi}{2})) \sqrt{c+d \csc(e+fx + \frac{\pi}{2})}} dx}{b} + \\
& \frac{dg \sqrt{g \sec(e+fx)} \sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{\sec(e+fx)}{\sqrt{\frac{d}{c+d} + \frac{c \cos(e+fx)}{c+d}}} dx}{b \sqrt{c+d \sec(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(bc - ad) \int \frac{(g \csc(e+fx + \frac{\pi}{2}))^{3/2}}{(a+b \csc(e+fx + \frac{\pi}{2})) \sqrt{c+d \csc(e+fx + \frac{\pi}{2})}} dx}{b} + \\
& \frac{2dg \sqrt{g \sec(e+fx)} \sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{1}{\sin(e+fx + \frac{\pi}{2}) \sqrt{\frac{d}{c+d} + \frac{c \sin(e+fx + \frac{\pi}{2})}{c+d}}} dx}{b \sqrt{c+d \sec(e+fx)}} \\
& \quad \downarrow \text{3284} \\
& \frac{(bc - ad) \int \frac{(g \csc(e+fx + \frac{\pi}{2}))^{3/2}}{(a+b \csc(e+fx + \frac{\pi}{2})) \sqrt{c+d \csc(e+fx + \frac{\pi}{2})}} dx}{b} + \\
& \frac{2dg \sqrt{g \sec(e+fx)} \sqrt{\frac{c \cos(e+fx)+d}{c+d}} \text{EllipticPi} \left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d} \right)}{bf \sqrt{c+d \sec(e+fx)}} \\
& \quad \downarrow \text{4463} \\
& \frac{g(bc - ad) \sqrt{g \sec(e+fx)} \sqrt{c \cos(e+fx) + d} \int \frac{1}{(b+a \cos(e+fx)) \sqrt{d+c \cos(e+fx)}} dx}{b \sqrt{c+d \sec(e+fx)}} + \\
& \frac{2dg \sqrt{g \sec(e+fx)} \sqrt{\frac{c \cos(e+fx)+d}{c+d}} \text{EllipticPi} \left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d} \right)}{bf \sqrt{c+d \sec(e+fx)}}
\end{aligned}$$

3.270. $\int \frac{(g \sec(e+fx))^{3/2} \sqrt{c+d \sec(e+fx)}}{a+b \sec(e+fx)} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{g(bc - ad)\sqrt{g \sec(e + fx)}\sqrt{c \cos(e + fx) + d} \int \frac{1}{(b+a \sin(e+fx+\frac{\pi}{2}))\sqrt{d+c \sin(e+fx+\frac{\pi}{2})}} dx}{b\sqrt{c + d \sec(e + fx)}} + \\
 & \frac{2dg\sqrt{g \sec(e + fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \text{EllipticPi}\left(2, \frac{1}{2}(e + fx), \frac{2c}{c+d}\right)}{bf\sqrt{c + d \sec(e + fx)}} \\
 & \downarrow 3286 \\
 & \frac{g(bc - ad)\sqrt{g \sec(e + fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{1}{(b+a \cos(e+fx))\sqrt{\frac{d}{c+d} + \frac{c \cos(e+fx)}{c+d}}} dx}{b\sqrt{c + d \sec(e + fx)}} + \\
 & \frac{2dg\sqrt{g \sec(e + fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \text{EllipticPi}\left(2, \frac{1}{2}(e + fx), \frac{2c}{c+d}\right)}{bf\sqrt{c + d \sec(e + fx)}} \\
 & \downarrow 3042 \\
 & \frac{g(bc - ad)\sqrt{g \sec(e + fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{1}{(b+a \sin(e+fx+\frac{\pi}{2}))\sqrt{\frac{d}{c+d} + \frac{c \sin(e+fx+\frac{\pi}{2})}{c+d}}} dx}{b\sqrt{c + d \sec(e + fx)}} + \\
 & \frac{2dg\sqrt{g \sec(e + fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \text{EllipticPi}\left(2, \frac{1}{2}(e + fx), \frac{2c}{c+d}\right)}{bf\sqrt{c + d \sec(e + fx)}} \\
 & \downarrow 3284 \\
 & \frac{2g(bc - ad)\sqrt{g \sec(e + fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \text{EllipticPi}\left(\frac{2a}{a+b}, \frac{1}{2}(e + fx), \frac{2c}{c+d}\right)}{bf(a + b)\sqrt{c + d \sec(e + fx)}} + \\
 & \frac{2dg\sqrt{g \sec(e + fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \text{EllipticPi}\left(2, \frac{1}{2}(e + fx), \frac{2c}{c+d}\right)}{bf\sqrt{c + d \sec(e + fx)}}
 \end{aligned}$$

input `Int[((g*Sec[e + f*x])^(3/2)*Sqrt[c + d*Sec[e + f*x]])/(a + b*Sec[e + f*x]),x]`

output `(2*d*g*Sqrt[(d + c*Cos[e + f*x])/(c + d)]*EllipticPi[2, (e + f*x)/2, (2*c)/(c + d)]*Sqrt[g*Sec[e + f*x]])/(b*f*Sqrt[c + d*Sec[e + f*x]]) + (2*(b*c - a*d)*g*Sqrt[(d + c*Cos[e + f*x])/(c + d)]*EllipticPi[(2*a)/(a + b), (e + f*x)/2, (2*c)/(c + d)]*Sqrt[g*Sec[e + f*x]])/(b*(a + b)*f*Sqrt[c + d*Sec[e + f*x]])`

3.270.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 4346 `Int[(csc[(e_) + (f_)*(x_)]*(d_))^(3/2)/Sqrt[csc[(e_) + (f_)*(x_)]*(b_) + (a_)], x_Symbol] := Simp[d*Sqrt[d*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sin[e + f*x]*Sqrt[b + a*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4459 `Int[((csc[(e_) + (f_)*(x_)]*(g_))^(3/2)*Sqrt[csc[(e_) + (f_)*(x_)]*(b_) + (a_)])/(csc[(e_) + (f_)*(x_)]*(d_) + (c_)), x_Symbol] := Simp[b/d Int[(g*Csc[e + f*x])^(3/2)/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[(b*c - a*d)/d Int[(g*Csc[e + f*x])^(3/2)/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 4463 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(3/2)/(Sqrt[csc[(e_) + (f_)*(x_)]*(b_) + (a_)])*(csc[(e_) + (f_)*(x_)]*(d_) + (c_)), x_Symbol] := Simp[g*Sqrt[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

3.270.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 8.49 (sec) , antiderivative size = 445, normalized size of antiderivative = 2.62

method	result
default	$-\frac{2ig\sqrt{g\sec(fx+e)}\cos(fx+e)\sqrt{c+d\sec(fx+e)}\sqrt{\frac{d+c\cos(fx+e)}{(c+d)(\cos(fx+e)+1)}}}{(c+d)(\cos(fx+e)+1)}\left(\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{c-d}{c+d}}\right)abc-\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{c-d}{c+d}}\right)abc-\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{c-d}{c+d}}\right)abc-\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{c-d}{c+d}}\right)abc\right)$

```
input int((g*sec(f*x+e))^(3/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*sec(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output -2*I*g/f/b/(a-b)/(a+b)*(g*sec(f*x+e))^(1/2)*cos(f*x+e)*(c+d*sec(f*x+e))^(1/2)*(1/(c+d)*(d+c*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-c-d)/(c+d))^(1/2))*a*b*c-EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-c-d)/(c+d))^(1/2))*a*b*d+EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-c-d)/(c+d))^(1/2))*b^2*c-EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-c-d)/(c+d))^(1/2))*b^2*d-2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-1,I*((c-d)/(c+d))^(1/2))*a^2*d+2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-1,I*((c-d)/(c+d))^(1/2))*b^2*d+2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-(a-b)/(a+b),I*((c-d)/(c+d))^(1/2))*a^2*d-2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-(a-b)/(a+b),I*((c-d)/(c+d))^(1/2))*a*b*c)/(d+c*cos(f*x+e))/(1/(cos(f*x+e)+1))^(1/2)
```

3.270.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx = \text{Timed out}$$

```
input integrate((g*sec(f*x+e))^(3/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*sec(f*x+e)),x,algorithm="fricas")
```

```
output Timed out
```

3.270.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx = \int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx$$

input `integrate((g*sec(f*x+e))**(3/2)*(c+d*sec(f*x+e))**(1/2)/(a+b*sec(f*x+e)),x)`

output `Integral((g*sec(e + f*x))**(3/2)*sqrt(c + d*sec(e + f*x))/(a + b*sec(e + f*x)), x)`

3.270.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx = \int \frac{\sqrt{d \sec(fx + e) + c} (g \sec(fx + e))^{3/2}}{b \sec(fx + e) + a} dx$$

input `integrate((g*sec(f*x+e))^(3/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*sec(f*x+e)),x, algorithm="maxima")`

output `integrate(sqrt(d*sec(f*x + e) + c)*(g*sec(f*x + e))^(3/2)/(b*sec(f*x + e) + a), x)`

3.270.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx = \int \frac{\sqrt{d \sec(fx + e) + c} (g \sec(fx + e))^{3/2}}{b \sec(fx + e) + a} dx$$

input `integrate((g*sec(f*x+e))^(3/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*sec(f*x+e)),x, algorithm="giac")`

output `integrate(sqrt(d*sec(f*x + e) + c)*(g*sec(f*x + e))^(3/2)/(b*sec(f*x + e) + a), x)`

3.270.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{c + d \sec(e + fx)}}{a + b \sec(e + fx)} dx = \int \frac{\sqrt{c + \frac{d}{\cos(e + fx)}} \left(\frac{g}{\cos(e + fx)} \right)^{3/2}}{a + \frac{b}{\cos(e + fx)}} dx$$

input `int(((c + d/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(a + b/cos(e + f*x))),x)`

output `int(((c + d/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(a + b/cos(e + f*x))), x)`

3.271
$$\int \frac{(g \sec(e+fx))^{3/2}}{(a+b \sec(e+fx))\sqrt{c+d \sec(e+fx)}} dx$$

3.271.1 Optimal result	1983
3.271.2 Mathematica [A] (verified)	1983
3.271.3 Rubi [A] (verified)	1984
3.271.4 Maple [C] (verified)	1986
3.271.5 Fracas [F(-1)]	1986
3.271.6 Sympy [F]	1987
3.271.7 Maxima [F]	1987
3.271.8 Giac [F]	1987
3.271.9 Mupad [F(-1)]	1988

3.271.1 Optimal result

Integrand size = 39, antiderivative size = 83

$$\int \frac{(g \sec(e+fx))^{3/2}}{(a+b \sec(e+fx))\sqrt{c+d \sec(e+fx)}} dx = \frac{2g\sqrt{\frac{d+c \cos(e+fx)}{c+d}} \text{EllipticPi}\left(\frac{2a}{a+b}, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right) \sqrt{g \sec(e+fx)}}{(a+b)f\sqrt{c+d \sec(e+fx)}}$$

output

```
2*g*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e),2*a/(a+b),2^(1/2)*(c/(c+d))^(1/2))*((d+c*cos(f*x+e))/(c+d))^(1/2)*(g*sec(f*x+e))^(1/2)/(a+b)/f/(c+d*sec(f*x+e))^(1/2)
```

3.271.2 Mathematica [A] (verified)

Time = 1.35 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00

$$\int \frac{(g \sec(e+fx))^{3/2}}{(a+b \sec(e+fx))\sqrt{c+d \sec(e+fx)}} dx = \frac{2g\sqrt{\frac{d+c \cos(e+fx)}{c+d}} \text{EllipticPi}\left(\frac{2a}{a+b}, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right) \sqrt{g \sec(e+fx)}}{(a+b)f\sqrt{c+d \sec(e+fx)}}$$

input

```
Integrate[(g*Sec[e + f*x])^(3/2)/((a + b*Sec[e + f*x])*Sqrt[c + d*Sec[e + f*x]]),x]
```

output

```
(2*g*Sqrt[(d + c*Cos[e + f*x])/(c + d)]*EllipticPi[(2*a)/(a + b), (e + f*x)/2, (2*c)/(c + d)]*Sqrt[g*Sec[e + f*x]]/((a + b)*f*Sqrt[c + d*Sec[e + f*x]])
```

3.271.
$$\int \frac{(g \sec(e+fx))^{3/2}}{(a+b \sec(e+fx))\sqrt{c+d \sec(e+fx)}} dx$$

3.271.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3042, 4463, 3042, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{(a + b \csc(e + fx + \frac{\pi}{2})) \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4463} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{c \cos(e + fx) + d} \int \frac{1}{(b + a \cos(e + fx)) \sqrt{d + c \cos(e + fx)}} dx}{\sqrt{c + d \sec(e + fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{c \cos(e + fx) + d} \int \frac{1}{(b + a \sin(e + fx + \frac{\pi}{2})) \sqrt{d + c \sin(e + fx + \frac{\pi}{2})}} dx}{\sqrt{c + d \sec(e + fx)}} \\
 & \quad \downarrow \text{3286} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{\frac{c \cos(e + fx) + d}{c + d}} \int \frac{1}{(b + a \cos(e + fx)) \sqrt{\frac{d}{c + d} + \frac{c \cos(e + fx)}{c + d}}} dx}{\sqrt{c + d \sec(e + fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{\frac{c \cos(e + fx) + d}{c + d}} \int \frac{1}{(b + a \sin(e + fx + \frac{\pi}{2})) \sqrt{\frac{d}{c + d} + \frac{c \sin(e + fx + \frac{\pi}{2})}{c + d}}} dx}{\sqrt{c + d \sec(e + fx)}} \\
 & \quad \downarrow \text{3284} \\
 & \frac{2g \sqrt{g \sec(e + fx)} \sqrt{\frac{c \cos(e + fx) + d}{c + d}} \text{EllipticPi}\left(\frac{2a}{a + b}, \frac{1}{2}(e + fx), \frac{2c}{c + d}\right)}{f(a + b) \sqrt{c + d \sec(e + fx)}}
 \end{aligned}$$

3.271. $\int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx$

input `Int[(g*Sec[e + f*x])^(3/2)/((a + b*Sec[e + f*x])*Sqrt[c + d*Sec[e + f*x]]),x]`

output `(2*g*Sqrt[(d + c*Cos[e + f*x])/(c + d)]*EllipticPi[(2*a)/(a + b), (e + f*x)/2, (2*c)/(c + d)]*Sqrt[g*Sec[e + f*x]])/((a + b)*f*Sqrt[c + d*Sec[e + f*x]])`

3.271.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 4463 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(3/2)/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))), x_Symbol] := Simp[g*Sqrt[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

3.271.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 7.34 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.69

method	result
default	$-\frac{2ig \cos(fx+e)\sqrt{c+d \sec(fx+e)} \left(2a \operatorname{EllipticPi}\left(i(-\cot(fx+e)+\csc(fx+e)), -\frac{a-b}{a+b}, i\sqrt{\frac{c-d}{c+d}}\right) - a \operatorname{EllipticF}\left(i(-\cot(fx+e)+\csc(fx+e)), \frac{c-d}{c+d}\right) - b \operatorname{EllipticF}\left(i(-\cot(fx+e)+\csc(fx+e)), \frac{c-d}{c+d}\right) \right)}{f(a-b)(a+b)(d+c \cos(fx+e))\sqrt{\frac{c-d}{c+d}}}$

input `int((g*sec(f*x+e))^(3/2)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output
$$-2*I*g/f/(a-b)/(a+b)*\cos(f*x+e)*(c+d*\sec(f*x+e))^{1/2}*(2*a*\operatorname{EllipticPi}(I*(-\cot(f*x+e)+\csc(f*x+e)), -(a-b)/(a+b), I*((c-d)/(c+d))^{1/2})-a*\operatorname{EllipticF}(I*(-\cot(f*x+e)+\csc(f*x+e)), (-(c-d)/(c+d))^{1/2})-b*\operatorname{EllipticF}(I*(-\cot(f*x+e)+\csc(f*x+e)), (-(c-d)/(c+d))^{1/2}))* (g*\sec(f*x+e))^{1/2}*(1/(c+d))*(d+c*\cos(f*x+e))/(\cos(f*x+e)+1))^{1/2}/(d+c*\cos(f*x+e))/(1/(\cos(f*x+e)+1))^{1/2}$$

3.271.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx))\sqrt{c + d \sec(e + fx)}} dx = \text{Timed out}$$

input `integrate((g*sec(f*x+e))^(3/2)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^(1/2),x,algorithm="fricas")`

output `Timed out`

3.271.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx = \int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx$$

input `integrate((g*sec(f*x+e))**(3/2)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))**(1/2),x)`

output `Integral((g*sec(e + f*x))**(3/2)/((a + b*sec(e + f*x))*sqrt(c + d*sec(e + f*x))), x)`

3.271.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx = \int \frac{(g \sec(fx + e))^{3/2}}{(b \sec(fx + e) + a) \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate((g*sec(f*x+e))^(3/2)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^(1/2),x,algorithm="maxima")`

output `integrate((g*sec(f*x + e))^(3/2)/((b*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)), x)`

3.271.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx = \int \frac{(g \sec(fx + e))^{3/2}}{(b \sec(fx + e) + a) \sqrt{d \sec(fx + e) + c}} dx$$

input `integrate((g*sec(f*x+e))^(3/2)/(a+b*sec(f*x+e))/(c+d*sec(f*x+e))^(1/2),x,algorithm="giac")`

output `integrate((g*sec(f*x + e))^(3/2)/((b*sec(f*x + e) + a)*sqrt(d*sec(f*x + e) + c)), x)`

3.271. $\int \frac{(g \sec(e+fx))^{3/2}}{(a+b \sec(e+fx)) \sqrt{c+d \sec(e+fx)}} dx$

3.271.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2}}{(a + b \sec(e + fx)) \sqrt{c + d \sec(e + fx)}} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{3/2}}{\left(a + \frac{b}{\cos(e+fx)}\right) \sqrt{c + \frac{d}{\cos(e+fx)}}} dx$$

input `int((g/cos(e + f*x))^(3/2)/((a + b/cos(e + f*x))*(c + d/cos(e + f*x))^(1/2)),x)`

output `int((g/cos(e + f*x))^(3/2)/((a + b/cos(e + f*x))*(c + d/cos(e + f*x))^(1/2)), x)`

3.272
$$\int \frac{\sqrt{g \sec(e+fx)} \sqrt{c+d \sec(e+fx)}}{a+b \cos(e+fx)} dx$$

3.272.1 Optimal result	1989
3.272.2 Mathematica [C] (verified)	1990
3.272.3 Rubi [A] (verified)	1990
3.272.4 Maple [C] (verified)	1994
3.272.5 Fricas [F(-1)]	1995
3.272.6 Sympy [F]	1995
3.272.7 Maxima [F]	1995
3.272.8 Giac [F]	1996
3.272.9 Mupad [F(-1)]	1996

3.272.1 Optimal result

Integrand size = 39, antiderivative size = 168

$$\int \frac{\sqrt{g \sec(e+fx)} \sqrt{c+d \sec(e+fx)}}{a+b \cos(e+fx)} dx$$

$$= \frac{2d \sqrt{\frac{d+c \cos(e+fx)}{c+d}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right) \sqrt{g \sec(e+fx)}}{af \sqrt{c+d \sec(e+fx)}} + \frac{2(ac-bd) \sqrt{\frac{d+c \cos(e+fx)}{c+d}} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right) \sqrt{g \sec(e+fx)}}{a(a+b)f \sqrt{c+d \sec(e+fx)}}$$

```
output 2*d*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e),2,2^(1/2)*(c/(c+d))^(1/2))*((d+c*cos(f*x+e))/(c+d))^(1/2)*(g*sec(f*x+e))^(1/2)/a/f/(c+d*sec(f*x+e))^(1/2)+2*(a*c-b*d)*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e),2*b/(a+b),2^(1/2)*(c/(c+d))^(1/2))*((d+c*cos(f*x+e))/(c+d))^(1/2)*(g*sec(f*x+e))^(1/2)/a/(a+b)/f/(c+d*sec(f*x+e))^(1/2)
```

3.272.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 25.12 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.32

$$\int \frac{\sqrt{g \sec(e + fx)} \sqrt{c + d \sec(e + fx)}}{a + b \cos(e + fx)} dx =$$

$$\frac{2i \sqrt{-\frac{c(-1+\cos(e+fx))}{c+d}} \sqrt{\frac{c(1+\cos(e+fx))}{c-d}} \cot(e + fx) \left(\text{EllipticPi} \left(1 - \frac{c}{d}, \text{I} \text{arcsinh} \left(\sqrt{\frac{1}{c-d}} \sqrt{d + c \cos(e + fx)} \right) \right) \right)}{a \sqrt{\frac{1}{c-d}} f}$$

input `Integrate[(Sqrt[g*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])/(a + b*Cos[e + f*x]),x]`

output `((-2*I)*Sqrt[-((c*(-1 + Cos[e + f*x]))/(c + d))]*Sqrt[(c*(1 + Cos[e + f*x]))/(c - d)]*Cot[e + f*x]*(EllipticPi[1 - c/d, I*ArcSinh[Sqrt[(c - d)^(-1)]*Sqrt[d + c*Cos[e + f*x]]], (-c + d)/(c + d)] - EllipticPi[(b*(-c + d))/(-(a*c) + b*d), I*ArcSinh[Sqrt[(c - d)^(-1)]*Sqrt[d + c*Cos[e + f*x]]], (-c + d)/(c + d)]*Sqrt[g*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])/(a*Sqrt[(c - d)^(-1)]*f*Sqrt[d + c*Cos[e + f*x]])`

3.272.3 Rubi [A] (verified)

Time = 2.33 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 3441, 3042, 4459, 3042, 4346, 3042, 3286, 3042, 3284, 4463, 3042, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{g \sec(e + fx)} \sqrt{c + d \sec(e + fx)}}{a + b \cos(e + fx)} dx$$

↓ 3042

$$\int \frac{\sqrt{g \csc(e + fx + \frac{\pi}{2})} \sqrt{c + d \csc(e + fx + \frac{\pi}{2})}}{a + b \sin(e + fx + \frac{\pi}{2})} dx$$

↓ 3441

3.272. $\int \frac{\sqrt{g \sec(e+fx)} \sqrt{c+d \sec(e+fx)}}{a+b \cos(e+fx)} dx$

$$\begin{aligned}
 & \int \frac{(g \sec(e+fx))^{3/2} \sqrt{c+d \sec(e+fx)}}{b+a \sec(e+fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2} \sqrt{c+d \csc(e+fx+\frac{\pi}{2})}}{b+a \csc(e+fx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4459} \\
 & \frac{(ac-bd) \int \frac{(g \sec(e+fx))^{3/2}}{(b+a \sec(e+fx)) \sqrt{c+d \sec(e+fx)}} dx}{a} + \frac{d \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{c+d \sec(e+fx)}} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(ac-bd) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{(b+a \csc(e+fx+\frac{\pi}{2})) \sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{d \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx}{a} \\
 & \quad \downarrow \text{4346} \\
 & \frac{(ac-bd) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{(b+a \csc(e+fx+\frac{\pi}{2})) \sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{dg \sqrt{g \sec(e+fx)} \sqrt{c \cos(e+fx)+d} \int \frac{\sec(e+fx)}{\sqrt{d+c \cos(e+fx)}} dx}{a \sqrt{c+d \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(ac-bd) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{(b+a \csc(e+fx+\frac{\pi}{2})) \sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{dg \sqrt{g \sec(e+fx)} \sqrt{c \cos(e+fx)+d} \int \frac{1}{\sin(e+fx+\frac{\pi}{2}) \sqrt{d+c \sin(e+fx+\frac{\pi}{2})}} dx}{a \sqrt{c+d \sec(e+fx)}} \\
 & \quad \downarrow \text{3286} \\
 & \frac{(ac-bd) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{(b+a \csc(e+fx+\frac{\pi}{2})) \sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{dg \sqrt{g \sec(e+fx)} \sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{\sec(e+fx)}{\sqrt{\frac{d}{c+d} + \frac{c \cos(e+fx)}{c+d}}} dx}{a \sqrt{c+d \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(ac-bd) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{(b+a \csc(e+fx+\frac{\pi}{2})) \sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{dg \sqrt{g \sec(e+fx)} \sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{1}{\sin(e+fx+\frac{\pi}{2}) \sqrt{\frac{d}{c+d} + \frac{c \sin(e+fx+\frac{\pi}{2})}{c+d}}} dx}{a \sqrt{c+d \sec(e+fx)}} \\
 & \quad \downarrow \text{g}
 \end{aligned}$$

3.272. $\int \frac{\sqrt{g \sec(e+fx)} \sqrt{c+d \sec(e+fx)}}{a+b \cos(e+fx)} dx$

$$\begin{aligned}
 & \downarrow 3284 \\
 & \frac{(ac-bd) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{(b+a \csc(e+fx+\frac{\pi}{2}))\sqrt{c+d \csc(e+fx+\frac{\pi}{2})}} dx + \frac{2dg\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right)}{af\sqrt{c+d \sec(e+fx)}}}{g} \\
 & \downarrow 4463 \\
 & \frac{g(ac-bd)\sqrt{g \sec(e+fx)}\sqrt{c \cos(e+fx)+d} \int \frac{1}{(a+b \cos(e+fx))\sqrt{d+c \cos(e+fx)}} dx + \frac{2dg\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right)}{af\sqrt{c+d \sec(e+fx)}}}{g} \\
 & \downarrow 3042 \\
 & \frac{g(ac-bd)\sqrt{g \sec(e+fx)}\sqrt{c \cos(e+fx)+d} \int \frac{1}{(a+b \sin(e+fx+\frac{\pi}{2}))\sqrt{d+c \sin(e+fx+\frac{\pi}{2})}} dx + \frac{2dg\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right)}{af\sqrt{c+d \sec(e+fx)}}}{g} \\
 & \downarrow 3286 \\
 & \frac{g(ac-bd)\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{1}{(a+b \cos(e+fx))\sqrt{\frac{d}{c+d} + \frac{c \cos(e+fx)}{c+d}}} dx + \frac{2dg\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right)}{af\sqrt{c+d \sec(e+fx)}}}{g} \\
 & \downarrow 3042 \\
 & \frac{g(ac-bd)\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \int \frac{1}{(a+b \sin(e+fx+\frac{\pi}{2}))\sqrt{\frac{d}{c+d} + \frac{c \sin(e+fx+\frac{\pi}{2})}{c+d}}} dx + \frac{2dg\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right)}{af\sqrt{c+d \sec(e+fx)}}}{g} \\
 & \downarrow 3284 \\
 & \frac{2g(ac-bd)\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right) + \frac{2dg\sqrt{g \sec(e+fx)}\sqrt{\frac{c \cos(e+fx)+d}{c+d}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2c}{c+d}\right)}{af\sqrt{c+d \sec(e+fx)}}}{g}
 \end{aligned}$$

input `Int[(Sqrt[g*Sec[e + f*x]]*Sqrt[c + d*Sec[e + f*x]])/(a + b*Cos[e + f*x]),x]`

3.272. $\int \frac{\sqrt{g \sec(e+fx)}\sqrt{c+d \sec(e+fx)}}{a+b \cos(e+fx)} dx$

```
output ((2*d*g*Sqrt[(d + c*Cos[e + f*x])/(c + d)]*EllipticPi[2, (e + f*x)/2, (2*c
)/(c + d)]*Sqrt[g*Sec[e + f*x]])/(a*f*Sqrt[c + d*Sec[e + f*x]]) + (2*(a*c
- b*d)*g*Sqrt[(d + c*Cos[e + f*x])/(c + d)]*EllipticPi[(2*b)/(a + b), (e +
f*x)/2, (2*c)/(c + d)]*Sqrt[g*Sec[e + f*x]])/(a*(a + b)*f*Sqrt[c + d*Sec[
e + f*x]]))/g
```

3.272.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

```
rule 3286 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt
[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c +
d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]
```

```
rule 3441 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(d_.) +
(c_.))^(n_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp
[g^m Int[(g*Csc[e + f*x])^(p - m)*(b + a*Csc[e + f*x])^m*(c + d*Csc[e + f
*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && IntegerQ[m]
```

```
rule 4346 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.
) + (a_.)], x_Symbol] := Simp[d*Sqrt[d*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x
]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sin[e + f*x]*Sqrt[b + a*Sin[e + f*x]]
), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
```

rule 4459 `Int[((csc[(e_.) + (f_.)*(x_.)]*(g_.))^(3/2)*Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)]/(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.)), x_Symbol] := Simp[b/d Int[(g*Csc[e + f*x])^(3/2)/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[(b*c - a*d)/d Int[(g*Csc[e + f*x])^(3/2)/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 4463 `Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(3/2)/(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))), x_Symbol] := Simp[g*Sqrt[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

3.272.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 8.24 (sec) , antiderivative size = 442, normalized size of antiderivative = 2.63

method	result
default	$\frac{2i\sqrt{g\sec(fx+e)}\cos(fx+e)\sqrt{c+d\sec(fx+e)}\left(\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{c-d}{c+d}}\right)a^2c-\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{c-d}{c+d}}\right)a^2d\right)}{(a+b)\cos(fx+e)}$

input `int((g*sec(f*x+e))^(1/2)*(c+d*sec(f*x+e))^(1/2)/(a+cos(f*x+e)*b),x,method=_RETURNVERBOSE)`

output
$$2*I/f/a/(a-b)/(a+b)*(g*\sec(f*x+e))^{1/2}*\cos(f*x+e)*(c+d*\sec(f*x+e))^{1/2}*(\text{EllipticF}(I*(\cot(f*x+e)-\csc(f*x+e)),(-\frac{c-d}{c+d})^{1/2})*a^2*c-\text{EllipticF}(I*(\cot(f*x+e)-\csc(f*x+e)),(-\frac{c-d}{c+d})^{1/2})*a^2*d+\text{EllipticF}(I*(\cot(f*x+e)-\csc(f*x+e)),(-\frac{c-d}{c+d})^{1/2})*a*b*c-\text{EllipticF}(I*(\cot(f*x+e)-\csc(f*x+e)),(-\frac{c-d}{c+d})^{1/2})*a*b*d+2*\text{EllipticPi}(I*(\cot(f*x+e)-\csc(f*x+e)),-1,I*((\frac{c-d}{c+d})^{1/2})*a^2*d-2*\text{EllipticPi}(I*(\cot(f*x+e)-\csc(f*x+e)),-1,I*((\frac{c-d}{c+d})^{1/2})*b^2*d-2*\text{EllipticPi}(I*(\cot(f*x+e)-\csc(f*x+e)),(a-b)/(a+b),I*((\frac{c-d}{c+d})^{1/2})*a*b*c+2*\text{EllipticPi}(I*(\cot(f*x+e)-\csc(f*x+e)),(a-b)/(a+b),I*((\frac{c-d}{c+d})^{1/2})*b^2*d)*(1/(c+d)*(d+c*\cos(f*x+e)))/(\cos(f*x+e)+1))^{1/2}/(d+c*\cos(f*x+e))/(1/(\cos(f*x+e)+1))^{1/2}$$

3.272.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{g \sec(e + fx)} \sqrt{c + d \sec(e + fx)}}{a + b \cos(e + fx)} dx = \text{Timed out}$$

```
input integrate((g*sec(f*x+e))^(1/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*cos(f*x+e)),x,
algorithm="fricas")
```

```
output Timed out
```

3.272.6 Sympy [F]

$$\int \frac{\sqrt{g \sec(e + fx)} \sqrt{c + d \sec(e + fx)}}{a + b \cos(e + fx)} dx = \int \frac{\sqrt{g \sec(e + fx)} \sqrt{c + d \sec(e + fx)}}{a + b \cos(e + fx)} dx$$

```
input integrate((g*sec(f*x+e))**(1/2)*(c+d*sec(f*x+e))**(1/2)/(a+b*cos(f*x+e)),x
)
```

```
output Integral(sqrt(g*sec(e + f*x))*sqrt(c + d*sec(e + f*x))/(a + b*cos(e + f*x)
), x)
```

3.272.7 Maxima [F]

$$\int \frac{\sqrt{g \sec(e + fx)} \sqrt{c + d \sec(e + fx)}}{a + b \cos(e + fx)} dx = \int \frac{\sqrt{d \sec(fx + e) + c} \sqrt{g \sec(fx + e)}}{b \cos(fx + e) + a} dx$$

```
input integrate((g*sec(f*x+e))^(1/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*cos(f*x+e)),x,
algorithm="maxima")
```

```
output integrate(sqrt(d*sec(f*x + e) + c)*sqrt(g*sec(f*x + e))/(b*cos(f*x + e) +
a), x)
```

3.272.8 Giac [F]

$$\int \frac{\sqrt{g \sec(e+fx)} \sqrt{c+d \sec(e+fx)}}{a+b \cos(e+fx)} dx = \int \frac{\sqrt{d \sec(fx+e)+c} \sqrt{g \sec(fx+e)}}{b \cos(fx+e)+a} dx$$

input `integrate((g*sec(f*x+e))^(1/2)*(c+d*sec(f*x+e))^(1/2)/(a+b*cos(f*x+e)),x,
algorithm="giac")`

output `integrate(sqrt(d*sec(f*x + e) + c)*sqrt(g*sec(f*x + e))/(b*cos(f*x + e) +
a), x)`

3.272.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{g \sec(e+fx)} \sqrt{c+d \sec(e+fx)}}{a+b \cos(e+fx)} dx = \int \frac{\sqrt{c + \frac{d}{\cos(e+fx)}} \sqrt{\frac{g}{\cos(e+fx)}}}{a+b \cos(e+fx)} dx$$

input `int(((c + d/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(1/2))/(a + b*cos(e + f*x
)),x)`

output `int(((c + d/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(1/2))/(a + b*cos(e + f*x
)), x)`

3.273
$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx$$

3.273.1 Optimal result	1997
3.273.2 Mathematica [B] (verified)	1997
3.273.3 Rubi [A] (verified)	1998
3.273.4 Maple [A] (verified)	1999
3.273.5 Fricas [F]	1999
3.273.6 Sympy [F]	2000
3.273.7 Maxima [F]	2000
3.273.8 Giac [F]	2000
3.273.9 Mupad [F(-1)]	2001

3.273.1 Optimal result

Integrand size = 33, antiderivative size = 95

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx$$

$$= \frac{E\left(\arcsin\left(\frac{\tan(e+fx)}{1+\sec(e+fx)}\right) \middle| \frac{a-b}{a+b}\right) \sqrt{\frac{1}{1+\sec(e+fx)}} \sqrt{a+b\sec(e+fx)}}{cf \sqrt{\frac{a+b\sec(e+fx)}{(a+b)(1+\sec(e+fx))}}}$$

```
output EllipticE(tan(f*x+e)/(1+sec(f*x+e)), ((a-b)/(a+b))^(1/2))*(1/(1+sec(f*x+e)))^(1/2)*(a+b*sec(f*x+e))^(1/2)/c/f/((a+b*sec(f*x+e))/(a+b)/(1+sec(f*x+e)))^(1/2)
```

3.273.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 264 vs. 2(95) = 190.

Time = 6.44 (sec) , antiderivative size = 264, normalized size of antiderivative = 2.78

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx$$

$$= \frac{\cos^2\left(\frac{1}{2}(e+fx)\right) \sqrt{\sec(e+fx)} \sqrt{a+b\sec(e+fx)} \left(\frac{2\sqrt{\frac{\cos(e+fx)}{1+\cos(e+fx)}} E\left(\arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right) \middle| \frac{a-b}{a+b}\right) \sec^4\left(\frac{1}{2}(e+fx)\right) \sqrt{1+\sec(e+fx)}}{\left(\frac{1}{1+\cos(e+fx)}\right)^{3/2} \sqrt{\frac{b+a\cos(e+fx)}{(a+b)(1+\cos(e+fx))}}}\right)}{4cf(1 + \dots)}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/(c + c*Sec[e + f*x]),x]`

output `(Cos[(e + f*x)/2]^2*Sqrt[Sec[e + f*x]]*Sqrt[a + b*Sec[e + f*x]]*((2*Sqrt[Cos[e + f*x]/(1 + Cos[e + f*x]])*EllipticE[ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)]*Sec[(e + f*x)/2]^4*Sqrt[1 + Sec[e + f*x]])/(((1 + Cos[e + f*x])^(-1))^3/2)*Sqrt[(b + a*Cos[e + f*x])/((a + b)*(1 + Cos[e + f*x]))]) + (Sec[(e + f*x)/2]^5*Sqrt[1 + Sec[e + f*x]]*(-Sin[(e + f*x)/2] + Sin[(3*(e + f*x))/2]))/((1 + Cos[e + f*x])^(-1))^3/2 - 8*Sqrt[Sec[e + f*x]]*(Sin[e + f*x] - Tan[(e + f*x)/2]))/(4*c*f*(1 + Sec[e + f*x]))`

3.273.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {3042, 4456}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c\sec(e+fx)+c} dx$$

↓ 3042

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\sqrt{a+b\csc\left(e+fx+\frac{\pi}{2}\right)}}{c\csc\left(e+fx+\frac{\pi}{2}\right)+c} dx$$

↓ 4456

$$\frac{\sqrt{\frac{1}{\sec(e+fx)+1}}\sqrt{a+b\sec(e+fx)}E\left(\arcsin\left(\frac{\tan(e+fx)}{\sec(e+fx)+1}\right)\middle|\frac{a-b}{a+b}\right)}{cf\sqrt{\frac{a+b\sec(e+fx)}{(a+b)(\sec(e+fx)+1)}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/(c + c*Sec[e + f*x]),x]`

output `(EllipticE[ArcSin[Tan[e + f*x]/(1 + Sec[e + f*x])], (a - b)/(a + b)]*Sqrt[(1 + Sec[e + f*x])^(-1)]*Sqrt[a + b*Sec[e + f*x]])/(c*f*Sqrt[(a + b*Sec[e + f*x])/((a + b)*(1 + Sec[e + f*x]))])`

3.273.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4456 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[(-Sqrt[a + b*Csc[e + f*x]])*(Sqrt[c/(c + d*Csc[e + f*x])]/(d*f*Sqrt[c*d*((a + b*Csc[e + f*x])/((b*c + a*d)*(c + d*Csc[e + f*x]))])))*EllipticE[ArcSin[c*(Cot[e + f*x]/(c + d*Csc[e + f*x])], -(b*c - a*d)/(b*c + a*d)], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && EqQ[c^2 - d^2, 0]`

3.273.4 Maple [A] (verified)

Time = 7.30 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.29

method	result	size
default	$\frac{(-a-b)(\cos(fx+e)+1)\sqrt{\frac{b+a\cos(fx+e)}{(a+b)(\cos(fx+e)+1)}}\sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}}\text{EllipticE}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)\sqrt{a+b\sec(fx+e)}}{cf(b+a\cos(fx+e))}$	123

input `int(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/c/f*(-a-b)*(cos(f*x+e)+1)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*(a+b*sec(f*x+e))^(1/2)/(b+a*cos(f*x+e))`

3.273.5 Fracas [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx = \int \frac{\sqrt{b\sec(fx+e)+a\sec(fx+e)}}{c\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x, algorithm="fracas")`

output `integral(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(c*sec(f*x + e) + c), x)`

3.273.
$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx$$

3.273.6 Sympy [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx = \frac{\int \frac{\sqrt{a+b\sec(e+fx)}\sec(e+fx)}{\sec(e+fx)+1} dx}{c}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))**(1/2)/(c+c*sec(f*x+e)),x)`

output `Integral(sqrt(a + b*sec(e + f*x))*sec(e + f*x)/(sec(e + f*x) + 1), x)/c`

3.273.7 Maxima [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx = \int \frac{\sqrt{b\sec(fx+e)+a}\sec(fx+e)}{c\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(c*sec(f*x + e) + c), x)`

3.273.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx = \int \frac{\sqrt{b\sec(fx+e)+a}\sec(fx+e)}{c\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x, algorithm="giac")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(c*sec(f*x + e) + c), x)`

3.273.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx) \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx = \int \frac{\sqrt{a + \frac{b}{\cos(e + fx)}}}{\cos(e + fx) \left(c + \frac{c}{\cos(e + fx)}\right)} dx$$

input `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + c/cos(e + f*x))),x)`

output `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + c/cos(e + f*x))), x)`

3.274
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+c \sec(e+fx)} dx$$

3.274.1 Optimal result	2002
3.274.2 Mathematica [F]	2003
3.274.3 Rubi [A] (verified)	2003
3.274.4 Maple [C] (verified)	2010
3.274.5 Fracas [F(-1)]	2011
3.274.6 Sympy [F]	2011
3.274.7 Maxima [F]	2012
3.274.8 Giac [F]	2012
3.274.9 Mupad [F(-1)]	2012

3.274.1 Optimal result

Integrand size = 39, antiderivative size = 295

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+c \sec(e+fx)} dx = \frac{g(b+a \cos(e+fx))E(\frac{1}{2}(e+fx)|\frac{2a}{a+b}) \sqrt{g \sec(e+fx)}}{cf \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \sqrt{a+b \sec(e+fx)}} + \frac{(a-b)g \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticF}(\frac{1}{2}(e+fx), \frac{2a}{a+b}) \sqrt{g \sec(e+fx)}}{cf \sqrt{a+b \sec(e+fx)}} + \frac{2bg \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticPi}(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}) \sqrt{g \sec(e+fx)}}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g(b+a \cos(e+fx)) \sqrt{g \sec(e+fx)} \sin(e+fx)}{f(c+c \cos(e+fx)) \sqrt{a+b \sec(e+fx)}}$$

output

```
-g*(b+a*cos(f*x+e))*sin(f*x+e)*(g*sec(f*x+e))^(1/2)/f/(c+c*cos(f*x+e))/(a+b*sec(f*x+e))^(1/2)+g*(b+a*cos(f*x+e))*(cos(1/2*f*x+1/2*e))^2^(1/2)/cos(1/2*f*x+1/2*e)*EllipticE(sin(1/2*f*x+1/2*e),2^(1/2)*(a/(a+b))^(1/2))*(g*sec(f*x+e))^(1/2)/c/f/((b+a*cos(f*x+e))/(a+b))^(1/2)/(a+b*sec(f*x+e))^(1/2)+(a-b)*g*(cos(1/2*f*x+1/2*e))^2^(1/2)/cos(1/2*f*x+1/2*e)*EllipticF(sin(1/2*f*x+1/2*e),2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/c/f/(a+b*sec(f*x+e))^(1/2)+2*b*g*(cos(1/2*f*x+1/2*e))^2^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e),2,2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/c/f/(a+b*sec(f*x+e))^(1/2)
```

3.274.2 Mathematica [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx = \int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx$$

input `Integrate[((g*Sec[e + f*x])^(3/2)*Sqrt[a + b*Sec[e + f*x]])/(c + c*Sec[e + f*x]),x]`

output `Integrate[((g*Sec[e + f*x])^(3/2)*Sqrt[a + b*Sec[e + f*x]])/(c + c*Sec[e + f*x]), x]`

3.274.3 Rubi [A] (verified)

Time = 2.75 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.05, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3042, 4459, 3042, 4346, 3042, 3286, 3042, 3284, 4463, 3042, 3247, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c \sec(e + fx) + c} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2} \sqrt{a + b \csc(e + fx + \frac{\pi}{2})}}{c \csc(e + fx + \frac{\pi}{2}) + c} dx \\ & \quad \downarrow \text{4459} \\ & \frac{b \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}} dx}{c} + (a - b) \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)} (\sec(e + fx) c + c)} dx \\ & \quad \downarrow \text{3042} \\ & \frac{b \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}} dx}{c} + (a - b) \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})} (\csc(e + fx + \frac{\pi}{2}) c + c)} dx \\ & \quad \downarrow \text{4346} \end{aligned}$$

3.274. $\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx$

$$\begin{aligned}
& \frac{(a-b) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (\csc(e+fx+\frac{\pi}{2})c+c)} dx +}{bg \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{\sec(e+fx)}{\sqrt{b+a \cos(e+fx)}} dx} \\
& \qquad \qquad \qquad \frac{dx}{c \sqrt{a+b \sec(e+fx)}} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{(a-b) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (\csc(e+fx+\frac{\pi}{2})c+c)} dx +}{bg \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{1}{\sin(e+fx+\frac{\pi}{2}) \sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx} \\
& \qquad \qquad \qquad \frac{dx}{c \sqrt{a+b \sec(e+fx)}} \\
& \qquad \qquad \qquad \downarrow \text{3286} \\
& \frac{(a-b) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (\csc(e+fx+\frac{\pi}{2})c+c)} dx +}{bg \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{\sec(e+fx)}{\sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b}}} dx} \\
& \qquad \qquad \qquad \frac{dx}{c \sqrt{a+b \sec(e+fx)}} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{(a-b) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (\csc(e+fx+\frac{\pi}{2})c+c)} dx +}{bg \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sin(e+fx+\frac{\pi}{2}) \sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b}}} dx} \\
& \qquad \qquad \qquad \frac{dx}{c \sqrt{a+b \sec(e+fx)}} \\
& \qquad \qquad \qquad \downarrow \text{3284} \\
& \frac{(a-b) \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (\csc(e+fx+\frac{\pi}{2})c+c)} dx +}{2bg \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)} \\
& \qquad \qquad \qquad \frac{dx}{cf \sqrt{a+b \sec(e+fx)}} \\
& \qquad \qquad \qquad \downarrow \text{4463} \\
& \frac{g(a-b) \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{1}{\sqrt{b+a \cos(e+fx)} (\cos(e+fx)c+c)} dx +}{2bg \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)} \\
& \qquad \qquad \qquad \frac{dx}{cf \sqrt{a+b \sec(e+fx)}} +
\end{aligned}$$

3.274. $\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+c \sec(e+fx)} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}(\sin(e+fx+\frac{\pi}{2})c+c)} dx}{\sqrt{a+b \sec(e+fx)}} + \\
& \frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3247} \\
& \frac{g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(-\frac{a \int -\frac{\cos(e+fx)c+c}{2\sqrt{b+a \cos(e+fx)}} dx}{c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} + \\
& \frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{27} \\
& \frac{g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \int \frac{\cos(e+fx)c+c}{\sqrt{b+a \cos(e+fx)}} dx}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} + \\
& \frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3042} \\
& \frac{g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \int \frac{\sin(e+fx+\frac{\pi}{2})c+c}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} + \\
& \frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3231} \\
& \frac{g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \cos(e+fx)}} dx}{a} + \frac{c \int \sqrt{b+a \cos(e+fx)} dx}{a} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} + \\
& \frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b \sec(e+fx)}}
\end{aligned}$$

3.274. $\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+c \sec(e+fx)} dx$

$$g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \downarrow \text{3042} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c \int \sqrt{b+a \sin(e+fx+\frac{\pi}{2})} dx}{a} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+b)} \right)$$

$$\frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi} \left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b} \right)}{cf\sqrt{a+b \sec(e+fx)}} \sqrt{a+b \sec(e+fx)}$$

$$g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \downarrow \text{3134} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c\sqrt{a \cos(e+fx)+b} \int \sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b}} dx}{a\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+b)} \right)$$

$$\frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi} \left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b} \right)}{cf\sqrt{a+b \sec(e+fx)}} \sqrt{a+b \sec(e+fx)}$$

$$g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \downarrow \text{3042} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c\sqrt{a \cos(e+fx)+b} \int \sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b}} dx}{a\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+b)} \right)$$

$$\frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi} \left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b} \right)}{cf\sqrt{a+b \sec(e+fx)}} \sqrt{a+b \sec(e+fx)}$$

$$\downarrow \text{3132}$$

$$g(a-b)\sqrt{g\sec(e+fx)}\sqrt{a\cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a\sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{2c\sqrt{a\cos(e+fx)+b}E\left(\frac{1}{2}(e+fx)\mid\frac{2a}{a+b}\right)}{af\sqrt{\frac{a\cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} \right) - \frac{\sin(e+fx)}{f(a-b)}$$

$$\frac{\sqrt{a+b\sec(e+fx)} \cdot 2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b\sec(e+fx)}}$$

↓ 3142

$$g(a-b)\sqrt{g\sec(e+fx)}\sqrt{a\cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b)\sqrt{\frac{a\cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a\cos(e+fx)}{a+b}}} dx}{a\sqrt{a\cos(e+fx)+b}} + \frac{2c\sqrt{a\cos(e+fx)+b}E\left(\frac{1}{2}(e+fx)\mid\frac{2a}{a+b}\right)}{af\sqrt{\frac{a\cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} \right)$$

$$\frac{\sqrt{a+b\sec(e+fx)} \cdot 2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b\sec(e+fx)}}$$

↓ 3042

$$g(a-b)\sqrt{g\sec(e+fx)}\sqrt{a\cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b)\sqrt{\frac{a\cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a\sin(e+fx+\frac{\pi}{2})}{a+b}}} dx}{a\sqrt{a\cos(e+fx)+b}} + \frac{2c\sqrt{a\cos(e+fx)+b}E\left(\frac{1}{2}(e+fx)\mid\frac{2a}{a+b}\right)}{af\sqrt{\frac{a\cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} \right)$$

$$\frac{\sqrt{a+b\sec(e+fx)} \cdot 2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b\sec(e+fx)}}$$

↓ 3140

3.274. $\int \frac{(g\sec(e+fx))^{3/2}\sqrt{a+b\sec(e+fx)}}{c+c\sec(e+fx)} dx$

$$g(a-b)\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{2c(a-b)\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{af\sqrt{a \cos(e+fx)+b}} + \frac{2c\sqrt{a \cos(e+fx)+b} E\left(\frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{af\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} \right)$$

$$\frac{2bg\sqrt{g \sec(e+fx)}\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf\sqrt{a+b \sec(e+fx)}}$$

input `Int[((g*Sec[e + f*x])^(3/2)*Sqrt[a + b*Sec[e + f*x]])/(c + c*Sec[e + f*x]),x]`

output `(2*b*g*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[2, (e + f*x)/2, (2*a)/(a + b)]*Sqrt[g*Sec[e + f*x]])/(c*f*Sqrt[a + b*Sec[e + f*x]]) + ((a - b)*g*Sqrt[b + a*Cos[e + f*x]]*Sqrt[g*Sec[e + f*x]]*((a*((2*c*Sqrt[b + a*Cos[e + f*x]]*EllipticE[(e + f*x)/2, (2*a)/(a + b)])/(a*f*Sqrt[(b + a*Cos[e + f*x])/(a + b)])) + (2*(a - b)*c*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticF[(e + f*x)/2, (2*a)/(a + b)]/(a*f*Sqrt[b + a*Cos[e + f*x]])))/(2*(a - b)*c^2) - (Sqrt[b + a*Cos[e + f*x]]*Sin[e + f*x])/((a - b)*f*(c + c*Cos[e + f*x])))/Sqrt[a + b*Sec[e + f*x]]`

3.274.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3247 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])^n/((a_) + (b_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(b*c - a*d)*(a + b*Sin[e + f*x]))], x] + Simp[d/(a*(b*c - a*d)) Int[(c + d*Sin[e + f*x])^n*(a*n - b*(n + 1)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, 0] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

```
rule 3286 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt
[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c +
d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]
```

```
rule 4346 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.
) + (a_)], x_Symbol] := Simp[d*Sqrt[d*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x
]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sin[e + f*x]*Sqrt[b + a*Sin[e + f*x]
]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 4459 Int[((csc[(e_.) + (f_.)*(x_)]*(g_.))^(3/2)*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.
) + (a_)])/(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[b/d
Int[(g*Csc[e + f*x])^(3/2)/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[(b*c -
a*d)/d Int[(g*Csc[e + f*x])^(3/2)/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e
+ f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 - b^2, 0]
```

```
rule 4463 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(3/2)/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.
) + (a_)])*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[g*Sqr
t[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int
[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b,
c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
```

3.274.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 6.87 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.93

method	result
default	$\frac{ig\sqrt{a+b\sec(fx+e)}\sqrt{g\sec(fx+e)}\cos(fx+e)\left(2a\operatorname{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{a-b}{a+b}}\right)-2b\operatorname{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{a-b}{a+b}}\right)\right)}{c+c\sec(e+fx)}$

```
input int((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x,method=
_RETURNVERBOSE)
```

$$3.274. \int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+c \sec(e+fx)} dx$$

output `I*g/c/f*(a+b*sec(f*x+e))^(1/2)*(g*sec(f*x+e))^(1/2)*cos(f*x+e)*(2*a*EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^(1/2))-2*b*EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^(1/2))-a*EllipticE(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^(1/2))-b*EllipticE(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^(1/2))+4*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-1,I*((a-b)/(a+b))^(1/2))*b*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)/(b+a*cos(f*x+e))/(1/(cos(f*x+e)+1))^(1/2)`

3.274.5 Fricas [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx = \text{Timed out}$$

input `integrate((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x,algorithm="fricas")`

output Timed out

3.274.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx = \frac{\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{\sec(e + fx) + 1} dx}{c}$$

input `integrate((g*sec(f*x+e))**(3/2)*(a+b*sec(f*x+e))**(1/2)/(c+c*sec(f*x+e)),x)`

output `Integral((g*sec(e + f*x))**(3/2)*sqrt(a + b*sec(e + f*x))/(sec(e + f*x) + 1), x)/c`

3.274.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx = \int \frac{\sqrt{b \sec(fx + e) + a} (g \sec(fx + e))^{3/2}}{c \sec(fx + e) + c} dx$$

input `integrate((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x,
algorithm="maxima")`

output `integrate(sqrt(b*sec(f*x + e) + a)*(g*sec(f*x + e))^(3/2)/(c*sec(f*x + e)
+ c), x)`

3.274.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx = \int \frac{\sqrt{b \sec(fx + e) + a} (g \sec(fx + e))^{3/2}}{c \sec(fx + e) + c} dx$$

input `integrate((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+c*sec(f*x+e)),x,
algorithm="giac")`

output `integrate(sqrt(b*sec(f*x + e) + a)*(g*sec(f*x + e))^(3/2)/(c*sec(f*x + e)
+ c), x)`

3.274.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + c \sec(e + fx)} dx = \int \frac{\sqrt{a + \frac{b}{\cos(e+fx)}} \left(\frac{g}{\cos(e+fx)}\right)^{3/2}}{c + \frac{c}{\cos(e+fx)}} dx$$

input `int(((a + b/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c + c/cos(e + f*x
)),x)`

output `int(((a + b/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c + c/cos(e + f*x
)), x)`

$$3.275 \quad \int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$$

3.275.1 Optimal result	2013
3.275.2 Mathematica [A] (warning: unable to verify)	2014
3.275.3 Rubi [A] (verified)	2014
3.275.4 Maple [A] (verified)	2016
3.275.5 Fracas [F]	2017
3.275.6 Sympy [F]	2017
3.275.7 Maxima [F]	2018
3.275.8 Giac [F]	2018
3.275.9 Mupad [F(-1)]	2018

3.275.1 Optimal result

Integrand size = 33, antiderivative size = 209

$$\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx =$$

$$\frac{2\sqrt{a+b} \cot(e+fx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(1+\sec(e+fx))}{a-b}}}{(a-b)cf}$$

$$+ \frac{E\left(\arcsin\left(\frac{\tan(e+fx)}{1+\sec(e+fx)}\right) \middle| \frac{a-b}{a+b}\right) \sqrt{\frac{1}{1+\sec(e+fx)}} \sqrt{a+b \sec(e+fx)}}{(a-b)cf \sqrt{\frac{a+b \sec(e+fx)}{(a+b)(1+\sec(e+fx))}}}$$

output `-2*cot(f*x+e)*EllipticF((a+b*sec(f*x+e))^(1/2)/(a+b)^(1/2),((a+b)/(a-b))^(1/2))*(a+b)^(1/2)*(b*(1-sec(f*x+e))/(a+b))^(1/2)*(-b*(1+sec(f*x+e))/(a-b))^(1/2)/(a-b)/c/f+EllipticE(tan(f*x+e)/(1+sec(f*x+e)),((a-b)/(a+b))^(1/2))*(1/(1+sec(f*x+e)))^(1/2)*(a+b*sec(f*x+e))^(1/2)/(a-b)/c/f/((a+b*sec(f*x+e))/(a+b)/(1+sec(f*x+e)))^(1/2)`

3.275.2 Mathematica [A] (warning: unable to verify)

Time = 13.34 (sec) , antiderivative size = 375, normalized size of antiderivative = 1.79

$$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$$

$$= \frac{\cos^2\left(\frac{e}{2} + \frac{fx}{2}\right) (b+a\cos(e+fx)) \sec^2(e+fx) \left(\frac{2\sin(e+fx)}{-a+b} - \frac{2\tan\left(\frac{1}{2}(e+fx)\right)}{-a+b}\right)}{f\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))}$$

$$= \frac{2\cos^2\left(\frac{e}{2} + \frac{fx}{2}\right) \sec^{\frac{3}{2}}(e+fx) \sqrt{\cos^2\left(\frac{1}{2}(e+fx)\right) \sec(e+fx)} \left((a-b)E\left(\arcsin\left(\sqrt{\frac{a-b}{a+b}} \tan\left(\frac{1}{2}(e+fx)\right)\right)\right)\right)}{\left(\frac{a-b}{a+b}\right)^{\frac{3}{2}} (a+b)f\sqrt{\cos(e+fx) \sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])),x]`output `(Cos[e/2 + (f*x)/2]^2*(b + a*Cos[e + f*x])*Sec[e + f*x]^2*((2*Sin[e + f*x])/(-a + b) - (2*Tan[(e + f*x)/2])/(-a + b)))/(f*Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])) - (2*Cos[e/2 + (f*x)/2]^2*Sec[e + f*x]^(3/2)*Sqrt[Cos[(e + f*x)/2]^2*Sec[e + f*x]]*((a - b)*EllipticE[ArcSin[Sqrt[(a - b)/(a + b)]]*Tan[(e + f*x)/2]], (a + b)/(a - b))*Sqrt[((b + a*Cos[e + f*x])*Sec[(e + f*x)/2]^2)/(a + b)] + Sqrt[2]*Sqrt[(a - b)/(a + b)]*Sqrt[Cos[e + f*x]/(1 + Cos[e + f*x])]*(b + a*Cos[e + f*x])*Tan[(e + f*x)/2]*(-1 + Tan[(e + f*x)/2]^2))/(((a - b)/(a + b))^(3/2)*(a + b)*f*Sqrt[Cos[e + f*x]*Sec[(e + f*x)/2]^4]*Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x]))`**3.275.3 Rubi [A] (verified)**Time = 0.73 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {3042, 4460, 3042, 4319, 4456}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)}{(c\sec(e+fx)+c)\sqrt{a+b\sec(e+fx)}} dx$$

↓ 3042

3.275. $\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$

$$\begin{aligned}
& \int \frac{\csc\left(e + fx + \frac{\pi}{2}\right)}{\left(c \csc\left(e + fx + \frac{\pi}{2}\right) + c\right) \sqrt{a + b \csc\left(e + fx + \frac{\pi}{2}\right)}} dx \\
& \quad \downarrow \text{4460} \\
& \frac{\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sec(e+fx)c+c} dx}{a-b} - \frac{b \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}} dx}{c(a-b)} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}}{\csc(e+fx+\frac{\pi}{2})c+c} dx}{a-b} - \frac{b \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}} dx}{c(a-b)} \\
& \quad \downarrow \text{4319} \\
& \frac{\int \frac{\csc(e+fx+\frac{\pi}{2})\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}}{\csc(e+fx+\frac{\pi}{2})c+c} dx}{a-b} - \\
& \frac{2\sqrt{a+b}\cot(e+fx)\sqrt{\frac{b(1-\sec(e+fx))}{a+b}}\sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b\sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{cf(a-b)} \\
& \quad \downarrow \text{4456} \\
& \frac{\sqrt{\frac{1}{\sec(e+fx)+1}}\sqrt{a+b\sec(e+fx)}E\left(\arcsin\left(\frac{\tan(e+fx)}{\sec(e+fx)+1}\right) \middle| \frac{a-b}{a+b}\right)}{cf(a-b)\sqrt{\frac{a+b\sec(e+fx)}{(a+b)(\sec(e+fx)+1)}}} - \\
& \frac{2\sqrt{a+b}\cot(e+fx)\sqrt{\frac{b(1-\sec(e+fx))}{a+b}}\sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b\sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{cf(a-b)}
\end{aligned}$$

input `Int[Sec[e + f*x]/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])),x]`

output `(-2*Sqrt[a + b]*Cot[e + f*x]*EllipticF[ArcSin[Sqrt[a + b*Sec[e + f*x]]/Sqrt[a + b]], (a + b)/(a - b)]*Sqrt[(b*(1 - Sec[e + f*x]))/(a + b)]*Sqrt[-((b*(1 + Sec[e + f*x]))/(a - b))]/((a - b)*c*f) + (EllipticE[ArcSin[Tan[e + f*x]/(1 + Sec[e + f*x])], (a - b)/(a + b)]*Sqrt[(1 + Sec[e + f*x])^(-1)]*Sqrt[a + b*Sec[e + f*x]]/((a - b)*c*f*Sqrt[(a + b*Sec[e + f*x])]/((a + b)*(1 + Sec[e + f*x]))))]`

3.275.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4319 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*(Rt[a + b, 2]/(b*f*Cot[e + f*x]))*Sqrt[(b*(1 - Csc[e + f*x]))/(a + b)]*Sqrt[(-b)*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Csc[e + f*x]]/Rt[a + b, 2]], (a + b)/(a - b)], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4456 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[(-Sqrt[a + b*Csc[e + f*x]])*(Sqrt[c/(c + d*Csc[e + f*x])]/(d*f*Sqrt[c*d*((a + b*Csc[e + f*x])/((b*c + a*d)*(c + d*Csc[e + f*x]))])))*EllipticE[ArcSin[c*(Cot[e + f*x]/(c + d*Csc[e + f*x]))], -(b*c - a*d)/(b*c + a*d)], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && EqQ[c^2 - d^2, 0]`

rule 4460 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[b/(b*c - a*d) Int[Csc[e + f*x]/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[d/(b*c - a*d) Int[Csc[e + f*x]*(Sqrt[a + b*Csc[e + f*x])/(c + d*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && (EqQ[a^2 - b^2, 0] || EqQ[c^2 - d^2, 0])`

3.275.4 Maple [A] (verified)

Time = 5.33 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.91

method	result
default	$\frac{(\cos(fx+e)+1)\left(2 \operatorname{EllipticF}\left(\cot(fx+e)-\csc(fx+e), \sqrt{\frac{a-b}{a+b}}\right) b - a \operatorname{EllipticE}\left(\cot(fx+e)-\csc(fx+e), \sqrt{\frac{a-b}{a+b}}\right) - b \operatorname{EllipticE}\left(\cot(fx+e), \sqrt{\frac{a-b}{a+b}}\right)\right)}{cf(a-b)(b+a \cos(fx+e))}$

input `int(sec(f*x+e)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2), x, method=_RETURNVERBOSE)`

$$3.275. \int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)(c+c \sec(e+fx))}} dx$$

output $1/c/f/(a-b)*(\cos(f*x+e)+1)*(2*EllipticF(\cot(f*x+e)-\csc(f*x+e),((a-b)/(a+b))^{(1/2)})*b-a*EllipticE(\cot(f*x+e)-\csc(f*x+e),((a-b)/(a+b))^{(1/2)})-b*EllipticE(\cot(f*x+e)-\csc(f*x+e),((a-b)/(a+b))^{(1/2)}))*(\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*(1/(a+b)*(b+a*\cos(f*x+e))/(\cos(f*x+e)+1))^{(1/2)}*(a+b*\sec(f*x+e))^{(1/2)}/(b+a*\cos(f*x+e))$

3.275.5 Fricas [F]

$$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx = \int \frac{\sec(fx+e)}{\sqrt{b\sec(fx+e)+a}(c\sec(fx+e)+c)} dx$$

input `integrate(sec(f*x+e)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(b*c*sec(f*x + e)^2 + (a + b)*c*sec(f*x + e) + a*c), x)`

3.275.6 Sympy [F]

$$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx = \frac{\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}\sec(e+fx)+\sqrt{a+b\sec(e+fx)}} dx}{c}$$

input `integrate(sec(f*x+e)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(a + b*sec(e + f*x))*sec(e + f*x) + sqrt(a + b*sec(e + f*x))), x)/c`

3.275.7 Maxima [F]

$$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx = \int \frac{\sec(fx+e)}{\sqrt{b\sec(fx+e)+a}(c\sec(fx+e)+c)} dx$$

input `integrate(sec(f*x+e)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.275.8 Giac [F]

$$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx = \int \frac{\sec(fx+e)}{\sqrt{b\sec(fx+e)+a}(c\sec(fx+e)+c)} dx$$

input `integrate(sec(f*x+e)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate(sec(f*x + e)/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.275.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx \\ &= \int \frac{1}{\cos(e+fx) \sqrt{a + \frac{b}{\cos(e+fx)}} \left(c + \frac{c}{\cos(e+fx)} \right)} dx \end{aligned}$$

input `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))),x)`

output `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))), x)`

$$3.276 \quad \int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$$

3.276.1 Optimal result	2019
3.276.2 Mathematica [A] (verified)	2020
3.276.3 Rubi [A] (verified)	2020
3.276.4 Maple [A] (verified)	2022
3.276.5 Fricas [F]	2023
3.276.6 Sympy [F]	2023
3.276.7 Maxima [F]	2024
3.276.8 Giac [F]	2024
3.276.9 Mupad [F(-1)]	2024

3.276.1 Optimal result

Integrand size = 35, antiderivative size = 214

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$$

$$= \frac{2a\sqrt{a+b} \cot(e+fx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(1+\sec(e+fx))}{a-b}}}{(a-b)bcf}$$

$$- \frac{E\left(\arcsin\left(\frac{\tan(e+fx)}{1+\sec(e+fx)}\right) \middle| \frac{a-b}{a+b}\right) \sqrt{\frac{1}{1+\sec(e+fx)}} \sqrt{a+b \sec(e+fx)}}{(a-b)cf \sqrt{\frac{a+b \sec(e+fx)}{(a+b)(1+\sec(e+fx))}}}$$

output `2*a*cot(f*x+e)*EllipticF((a+b*sec(f*x+e))^(1/2)/(a+b)^(1/2),((a+b)/(a-b))^(1/2))*(a+b)^(1/2)*(b*(1-sec(f*x+e))/(a+b))^(1/2)*(-b*(1+sec(f*x+e))/(a-b))^(1/2)/(a-b)/b/c/f-EllipticE(tan(f*x+e)/(1+sec(f*x+e)),((a-b)/(a+b))^(1/2))*(1/(1+sec(f*x+e)))^(1/2)*(a+b*sec(f*x+e))^(1/2)/(a-b)/c/f/((a+b*sec(f*x+e))/(a+b)/(1+sec(f*x+e)))^(1/2)`

3.276.2 Mathematica [A] (verified)

Time = 5.48 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.73

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$$

$$= \frac{4 \cos^4\left(\frac{1}{2}(e+fx)\right) \sqrt{\frac{b+a\cos(e+fx)}{(a+b)(1+\cos(e+fx))}} \left((a+b)E\left(\arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right) \middle| \frac{a-b}{a+b}\right) - 2a \operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right)\right) \right)}{(-a+b)cf \sqrt{\frac{\cos(e+fx)}{1+\cos(e+fx)}} (1+\cos(e+fx))^2 \sqrt{a+b\sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]^2/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])),x]`

output `(4*Cos[(e + f*x)/2]^4*Sqrt[(b + a*Cos[e + f*x])/((a + b)*(1 + Cos[e + f*x]))])*((a + b)*EllipticE[ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)] - 2*a*EllipticF[ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)]))/((-a + b)*c*f*Sqrt[Cos[e + f*x]/(1 + Cos[e + f*x])]*(1 + Cos[e + f*x])^2*Sqrt[a + b*Sec[e + f*x]])`

3.276.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4464, 3042, 4319, 4456}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(e+fx)}{(c\sec(e+fx)+c)\sqrt{a+b\sec(e+fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)^2}{\left(c\csc\left(e+fx+\frac{\pi}{2}\right)+c\right)\sqrt{a+b\csc\left(e+fx+\frac{\pi}{2}\right)}} dx$$

$$\downarrow \text{4464}$$

$$\frac{a \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}} dx}{c(a-b)} - \frac{\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{\sec(e+fx)c+c} dx}{a-b}$$

3.276. $\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx$

$$\begin{aligned}
& \int \frac{a \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})}} dx}{c(a-b)} \quad \downarrow \text{3042} \quad \int \frac{\csc(e+fx+\frac{\pi}{2}) \sqrt{a+b \csc(e+fx+\frac{\pi}{2})}}{\csc(e+fx+\frac{\pi}{2})c+c} dx \\
& \frac{2a\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{bcf(a-b)} \\
& \quad \downarrow \text{4319} \quad \frac{\int \frac{\csc(e+fx+\frac{\pi}{2}) \sqrt{a+b \csc(e+fx+\frac{\pi}{2})}}{\csc(e+fx+\frac{\pi}{2})c+c} dx}{a-b} \\
& \quad \downarrow \text{4456} \quad \frac{2a\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{bcf(a-b)} \\
& \quad \frac{\sqrt{\frac{1}{\sec(e+fx)+1}} \sqrt{a+b \sec(e+fx)} E\left(\arcsin\left(\frac{\tan(e+fx)}{\sec(e+fx)+1}\right) \middle| \frac{a-b}{a+b}\right)}{cf(a-b) \sqrt{\frac{a+b \sec(e+fx)}{(a+b)(\sec(e+fx)+1)}}}
\end{aligned}$$

input `Int[Sec[e + f*x]^2/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])),x]`

output `(2*a*Sqrt[a + b]*Cot[e + f*x]*EllipticF[ArcSin[Sqrt[a + b*Sec[e + f*x]]/Sqrt[a + b]], (a + b)/(a - b)]*Sqrt[(b*(1 - Sec[e + f*x]))/(a + b)]*Sqrt[-((b*(1 + Sec[e + f*x]))/(a - b))]/((a - b)*b*c*f) - (EllipticE[ArcSin[Tan[e + f*x]/(1 + Sec[e + f*x])], (a - b)/(a + b)]*Sqrt[(1 + Sec[e + f*x])^(-1)]*Sqrt[a + b*Sec[e + f*x]]/((a - b)*c*f*Sqrt[(a + b*Sec[e + f*x])/(a + b)]*(1 + Sec[e + f*x])))`

3.276.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4319 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*(Rt[a + b, 2]/(b*f*Cot[e + f*x]))*Sqrt[(b*(1 - Csc[e + f*x]))/(a + b)]*Sqrt[(-b)*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Csc[e + f*x]]/Rt[a + b, 2]], (a + b)/(a - b)], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4456 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[(-Sqrt[a + b*Csc[e + f*x]])*(Sqrt[c/(c + d*Csc[e + f*x])]/(d*f*Sqrt[c*d*((a + b*Csc[e + f*x])/((b*c + a*d)*(c + d*Csc[e + f*x]))])))*EllipticE[ArcSin[c*(Cot[e + f*x]/(c + d*Csc[e + f*x]))], -(b*c - a*d)/(b*c + a*d)], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && EqQ[c^2 - d^2, 0]`

rule 4464 `Int[csc[(e_.) + (f_.)*(x_)]^2/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))), x_Symbol] := Simp[-a/(b*c - a*d) Int[Csc[e + f*x]/Sqrt[a + b*Csc[e + f*x]], x], x] + Simp[c/(b*c - a*d) Int[Csc[e + f*x]*(Sqrt[a + b*Csc[e + f*x])/(c + d*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && (EqQ[a^2 - b^2, 0] || EqQ[c^2 - d^2, 0])`

3.276.4 Maple [A] (verified)

Time = 8.77 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.90

method	result
default	$-\frac{(\cos(fx+e)+1)\left(-a \operatorname{EllipticE}\left(\cot(fx+e)-\operatorname{csc}(fx+e), \sqrt{\frac{a-b}{a+b}}\right)-b \operatorname{EllipticE}\left(\cot(fx+e)-\operatorname{csc}(fx+e), \sqrt{\frac{a-b}{a+b}}\right)+2 \operatorname{EllipticF}\left(\cot(fx+e), \sqrt{\frac{a-b}{a+b}}\right)\right)}{cf(a-b)(b+a \cos(fx+e))}$

input `int(sec(f*x+e)^2/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2), x, method=_RETURNV ERBOSE)`

$$3.276. \int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)(c+c \sec(e+fx))}} dx$$

output `-1/c/f/(a-b)*(cos(f*x+e)+1)*(-a*EllipticE(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))-b*EllipticE(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))+2*EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*a)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(a+b*sec(f*x+e))^(1/2)/(b+a*cos(f*x+e))`

3.276.5 Fricas [F]

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx = \int \frac{\sec^2(fx+e)}{\sqrt{b\sec(fx+e)+a}(c\sec(fx+e)+c)} dx$$

input `integrate(sec(f*x+e)^2/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)^2/(b*c*sec(f*x + e)^2 + (a + b)*c*sec(f*x + e) + a*c), x)`

3.276.6 Sympy [F]

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+c\sec(e+fx))} dx = \frac{\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}\sec(e+fx)+\sqrt{a+b\sec(e+fx)}} dx}{c}$$

input `integrate(sec(f*x+e)**2/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)**2/(sqrt(a + b*sec(e + f*x))*sec(e + f*x) + sqrt(a + b*sec(e + f*x))), x)/c`

3.276.7 Maxima [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{\sec(fx + e)^2}{\sqrt{b \sec(fx + e) + a}(c \sec(fx + e) + c)} dx$$

input `integrate(sec(f*x+e)^2/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm m="maxima")`

output `integrate(sec(f*x + e)^2/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.276.8 Giac [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{\sec(fx + e)^2}{\sqrt{b \sec(fx + e) + a}(c \sec(fx + e) + c)} dx$$

input `integrate(sec(f*x+e)^2/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm m="giac")`

output `integrate(sec(f*x + e)^2/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.276.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx \\ &= \int \frac{1}{\cos(e + fx)^2 \sqrt{a + \frac{b}{\cos(e + fx)}} \left(c + \frac{c}{\cos(e + fx)} \right)} dx \end{aligned}$$

input `int(1/(cos(e + f*x)^2*(a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))),x)`

output `int(1/(cos(e + f*x)^2*(a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))), x)`

3.276. $\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$

$$3.277 \quad \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$$

3.277.1 Optimal result	2025
3.277.2 Mathematica [C] (warning: unable to verify)	2026
3.277.3 Rubi [A] (verified)	2027
3.277.4 Maple [C] (verified)	2032
3.277.5 Fricas [C] (verification not implemented)	2032
3.277.6 Sympy [F]	2033
3.277.7 Maxima [F]	2033
3.277.8 Giac [F]	2034
3.277.9 Mupad [F(-1)]	2034

3.277.1 Optimal result

Integrand size = 39, antiderivative size = 229

$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx = \frac{g(b+a \cos(e+fx))E(\frac{1}{2}(e+fx)|\frac{2a}{a+b}) \sqrt{g \sec(e+fx)}}{(a-b)cf \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \sqrt{a+b \sec(e+fx)}} + \frac{g \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \operatorname{EllipticF}(\frac{1}{2}(e+fx), \frac{2a}{a+b}) \sqrt{g \sec(e+fx)}}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g(b+a \cos(e+fx)) \sqrt{g \sec(e+fx)} \sin(e+fx)}{(a-b)f(c+c \cos(e+fx)) \sqrt{a+b \sec(e+fx)}}$$

output

```
-g*(b+a*cos(f*x+e))*sin(f*x+e)*(g*sec(f*x+e))^(1/2)/(a-b)/f/(c+c*cos(f*x+e)))/(a+b*sec(f*x+e))^(1/2)+g*(b+a*cos(f*x+e))*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticE(sin(1/2*f*x+1/2*e),2^(1/2)*(a/(a+b))^(1/2))*(g*sec(f*x+e))^(1/2)/(a-b)/c/f/((b+a*cos(f*x+e))/(a+b))^(1/2)/(a+b*sec(f*x+e))^(1/2)+g*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticF(sin(1/2*f*x+1/2*e),2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/c/f/(a+b*sec(f*x+e))^(1/2)
```

3.277.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 4 in optimal.

Time = 8.39 (sec) , antiderivative size = 1019, normalized size of antiderivative = 4.45

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \frac{\cos^2\left(\frac{e}{2} + \frac{fx}{2}\right) (b + a \cos(e + fx))(g \sec(e + fx))^{3/2} \left(\frac{2 \csc(e)}{(-a+b)f}\right)}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} +$$

$$\frac{\text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{\csc(e)(b - a\sqrt{1 + \cot^2(e)}) \sin(e) \sin(fx - \arctan(\cot(e)))}{a\sqrt{1 + \cot^2(e)}\left(1 + \frac{b \csc(e)}{a\sqrt{1 + \cot^2(e)}}\right)}, \frac{\csc(e)(b - a\sqrt{1 + \cot^2(e)}) \sin(e) \sin(fx - \arctan(\cot(e)))}{a\sqrt{1 + \cot^2(e)}\left(-1 + \frac{b \csc(e)}{a\sqrt{1 + \cot^2(e)}}\right)}\right)}{+}$$

$$a \cos^2\left(\frac{e}{2} + \frac{fx}{2}\right) \sqrt{b + a \cos(e + fx)} \csc\left(\frac{e}{2}\right) \sec\left(\frac{e}{2}\right) (g \sec(e + fx))^{3/2} \left(\frac{\text{AppellF1}\left(-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{\sec(e)(b + a \cos(e) \cos(fx + \arctan(\tan(e) \sqrt{1 + \tan^2(e)}))}{a\sqrt{1 + \tan^2(e)}}\right)}{\sqrt{1 + \tan^2(e)} \sqrt{\frac{a\sqrt{1 + \tan^2(e)} - a \cos(fx + \arctan(\tan(e) \sqrt{1 + \tan^2(e)})}{b \sec(e) + a\sqrt{1 + \tan^2(e)}}}\right)}{2(-}$$

```
input Integrate[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])),x]
```

output $(\cos[e/2 + (f*x)/2]^2*(b + a*\cos[e + f*x])*(g*\sec[e + f*x])^{3/2}*((2*\csc[e])/((-a + b)*f) + (2*\sec[e/2]*\sec[e/2 + (f*x)/2]*\sin[(f*x)/2])/((-a + b)*f))/(\sqrt{a + b*\sec[e + f*x]}*(c + c*\sec[e + f*x])) + (\text{AppellF1}[1/2, 1/2, 1/2, 3/2, (\csc[e]*(b - a*\sqrt{1 + \cot[e]^2}*\sin[e]*\sin[f*x - \text{ArcTan}[\cot[e]]])/(a*\sqrt{1 + \cot[e]^2}*(1 + (b*\csc[e])/(a*\sqrt{1 + \cot[e]^2}))), (\csc[e]*(b - a*\sqrt{1 + \cot[e]^2}*\sin[e]*\sin[f*x - \text{ArcTan}[\cot[e]]])/(a*\sqrt{1 + \cot[e]^2}*(-1 + (b*\csc[e])/(a*\sqrt{1 + \cot[e]^2})))])*\cos[e/2 + (f*x)/2]^2*\sqrt{b + a*\cos[e + f*x]}*csc[e/2]*\sec[e/2]*(g*\sec[e + f*x])^{3/2}*\sec[f*x - \text{ArcTan}[\cot[e]]]*\sqrt{(a*\sqrt{1 + \cot[e]^2} - a*\sqrt{1 + \cot[e]^2}*\sin[f*x - \text{ArcTan}[\cot[e]]])/(a*\sqrt{1 + \cot[e]^2} - b*\csc[e])}*\sqrt{(a*\sqrt{1 + \cot[e]^2} + a*\sqrt{1 + \cot[e]^2}*\sin[f*x - \text{ArcTan}[\cot[e]]])/(a*\sqrt{1 + \cot[e]^2} + b*\csc[e])}*\sqrt{b - a*\sqrt{1 + \cot[e]^2}*\sin[e]*\sin[f*x - \text{ArcTan}[\cot[e]]]})/((-a + b)*f*\sqrt{1 + \cot[e]^2}*\sqrt{a + b*\sec[e + f*x]}*(c + c*\sec[e + f*x])) + (a*\cos[e/2 + (f*x)/2]^2*\sqrt{b + a*\cos[e + f*x]}*csc[e/2]*\sec[e/2]*(g*\sec[e + f*x])^{3/2}*((\text{AppellF1}[-1/2, -1/2, -1/2, 1/2, -((\sec[e]*(b + a*\cos[e]*\cos[f*x + \text{ArcTan}[\tan[e]])*\sqrt{1 + \tan[e]^2}))/ (a*\sqrt{1 + \tan[e]^2}*(1 - (b*\sec[e])/(a*\sqrt{1 + \tan[e]^2}))) , -((\sec[e]*(b + a*\cos[e]*\cos[f*x + \text{ArcTan}[\tan[e]])*\sqrt{1 + \tan[e]^2}))/ (a*\sqrt{1 + \tan[e]^2})*(-1 - (b*\sec[e])/(a*\sqrt{1 + \tan[e]^2})))])*\sin[f*x + \text{ArcTan}[\tan[e]]]*\tan[e])/(\sqrt{1 + \tan[e]^2}*\sqrt{(a*\sqrt{1 + \tan[e]^2} - a*\cos[f*x + \text{ArcTan}[\tan[e]]])})$

3.277.3 Rubi [A] (verified)

Time = 1.27 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.359$, Rules used = {3042, 4463, 3042, 3247, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(g \sec(e + fx))^{3/2}}{(c \sec(e + fx) + c) \sqrt{a + b \sec(e + fx)}} dx$$

↓ 3042

$$\int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{(c \csc(e + fx + \frac{\pi}{2}) + c) \sqrt{a + b \csc(e + fx + \frac{\pi}{2})}} dx$$

↓ 4463

$$\frac{g \sqrt{g \sec(e + fx)} \sqrt{a \cos(e + fx) + b} \int \frac{1}{\sqrt{b + a \cos(e + fx)} (\cos(e + fx) c + c)}}{\sqrt{a + b \sec(e + fx)}} dx$$

3.277. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)} (c + c \sec(e + fx))} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}(\sin(e+fx+\frac{\pi}{2})c+c)} dx}{\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3247} \\
& \frac{g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(-\frac{a \int -\frac{\cos(e+fx)c+c}{2\sqrt{b+a \cos(e+fx)}} dx}{c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{27} \\
& \frac{g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \int \frac{\cos(e+fx)c+c}{\sqrt{b+a \cos(e+fx)}} dx}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3042} \\
& \frac{g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(a \int \frac{\sin(e+fx+\frac{\pi}{2})c+c}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3231} \\
& \frac{g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \cos(e+fx)}} dx}{a} + \frac{c \int \sqrt{b+a \cos(e+fx)} dx}{a} \right) - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3042} \\
& \frac{g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c \int \sqrt{b+a \sin(e+fx+\frac{\pi}{2})} dx}{a} \right) - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
& \downarrow \text{3134}
\end{aligned}$$

3.277. $\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$

$$g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c\sqrt{a \cos(e+fx)+b} \int \frac{\sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b}} dx}{a\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a}}{f(a-b)(c \cos(e+fx))} \right)$$

$$\sqrt{a+b \sec(e+fx)}$$

↓ 3042

$$g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c\sqrt{a \cos(e+fx)+b} \int \frac{\sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b}} dx}{a\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)}{f(a-b)(c \cos(e+fx))} \right)$$

$$\sqrt{a+b \sec(e+fx)}$$

↓ 3132

$$g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{2c\sqrt{a \cos(e+fx)+b} E\left(\frac{1}{2}(e+fx) \middle| \frac{2a}{a+b}\right)}{af\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)}}{f(a-b)(c \cos(e+fx))} \right)$$

$$\sqrt{a+b \sec(e+fx)}$$

↓ 3142

$$g\sqrt{g \sec(e+fx)}\sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b)\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b}}} dx}{a\sqrt{a \cos(e+fx)+b}} + \frac{2c\sqrt{a \cos(e+fx)+b} E\left(\frac{1}{2}(e+fx) \middle| \frac{2a}{a+b}\right)}{af\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right)}{2c^2(a-b)} - \frac{\sin(e+fx)\sqrt{a \cos(e+fx)}}{f(a-b)(c \cos(e+fx))} \right)$$

$$\sqrt{a+b \sec(e+fx)}$$

↓ 3042

3.277. $\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$

$$g\sqrt{g \sec(e + fx)}\sqrt{a \cos(e + fx) + b} \left(\frac{c(a-b)\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b}}} dx}{a\sqrt{a \cos(e+fx)+b}} + \frac{2c\sqrt{a \cos(e+fx)+b}E\left(\frac{1}{2}(e+fx) \mid \frac{2a}{a+b}\right)}{af\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) \frac{1}{2c^2(a-b)}$$

$$\sqrt{a + b \sec(e + fx)}$$

3140

$$g\sqrt{g \sec(e + fx)}\sqrt{a \cos(e + fx) + b} \left(\frac{2c(a-b)\sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{af\sqrt{a \cos(e+fx)+b}} + \frac{2c\sqrt{a \cos(e+fx)+b}E\left(\frac{1}{2}(e+fx) \mid \frac{2a}{a+b}\right)}{af\sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) \frac{1}{2c^2(a-b)} - \frac{\sin}{f}$$

$$\sqrt{a + b \sec(e + fx)}$$

input `Int[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])),x]`

output `(g*Sqrt[b + a*Cos[e + f*x]]*Sqrt[g*Sec[e + f*x]]*((a*((2*c*Sqrt[b + a*Cos[e + f*x]]*EllipticE[(e + f*x)/2, (2*a)/(a + b)])/(a*f*Sqrt[(b + a*Cos[e + f*x])/(a + b)]) + (2*(a - b)*c*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticF[(e + f*x)/2, (2*a)/(a + b)]/(a*f*Sqrt[b + a*Cos[e + f*x]])))/(2*(a - b)*c^2 - (Sqrt[b + a*Cos[e + f*x]]*Sin[e + f*x])/((a - b)*f*(c + c*Cos[e + f*x]))))/Sqrt[a + b*Sec[e + f*x]]`

3.277.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.277. \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$$

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3247 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)/((a_) + (b_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(b*c - a*d)*(a + b*Sin[e + f*x]))), x] + Simp[d/(a*(b*c - a*d)) Int[(c + d*Sin[e + f*x])^n*(a^n - b*(n + 1)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, 0] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 4463 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(3/2)/(Sqrt[csc[(e_) + (f_)*(x_)]*(b_ + (a_)*csc[(e_) + (f_)*(x_)]*(d_) + (c_))), x_Symbol] := Simp[g*Sqrt[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

3.277.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.70 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.90

method	result
default	$-\frac{ig \cos(fx+e) \sqrt{g \sec(fx+e)} \sqrt{a+b \sec(fx+e)} \left(2 \operatorname{EllipticF}\left(i(-\cot(fx+e)+\csc(fx+e)), \sqrt{-\frac{a-b}{a+b}}\right) a - a \operatorname{EllipticE}\left(i(-\cot(fx+e)+\csc(fx+e)), \sqrt{-\frac{a-b}{a+b}}\right) \right)}{cf(a-b)(b+a \cos(fx+e)) \sqrt{\frac{1}{\cos(fx+e)}}}$

input `int((g*sec(f*x+e))^(3/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output
$$-I*g/c/f/(a-b)*\cos(f*x+e)*(g*\sec(f*x+e))^{(1/2)}*(a+b*\sec(f*x+e))^{(1/2)}*(2*\operatorname{EllipticF}(I*(-\cot(f*x+e)+\csc(f*x+e)),(-a-b)/(a+b))^{(1/2)})*a-a*\operatorname{EllipticE}(I*(-\cot(f*x+e)+\csc(f*x+e)),(-a-b)/(a+b))^{(1/2)}-b*\operatorname{EllipticE}(I*(-\cot(f*x+e)+\csc(f*x+e)),(-a-b)/(a+b))^{(1/2)})*(1/(a+b)*(b+a*\cos(f*x+e))/(\cos(f*x+e)+1))^{(1/2)}/(b+a*\cos(f*x+e))/(1/(\cos(f*x+e)+1))^{(1/2)}$$

3.277.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 517, normalized size of antiderivative = 2.26

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx =$$

$$\frac{6 ag \sqrt{\frac{a \cos(fx+e)+b}{\cos(fx+e)}} \sqrt{\frac{g}{\cos(fx+e)}} \cos(fx+e) \sin(fx+e) + \sqrt{2}(i(3a-2b)g \cos(fx+e) + i(3a-2b)g) \sqrt{a}}{\dots}$$

input `integrate((g*sec(f*x+e))^(3/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="fricas")`

3.277.
$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$$

output

```
-1/6*(6*a*g*sqrt((a*cos(f*x + e) + b)/cos(f*x + e))*sqrt(g/cos(f*x + e))*c
os(f*x + e)*sin(f*x + e) + sqrt(2)*(I*(3*a - 2*b)*g*cos(f*x + e) + I*(3*a
- 2*b)*g)*sqrt(a*g)*weierstrassPInverse(-4/3*(3*a^2 - 4*b^2)/a^2, 8/27*(9*
a^2*b - 8*b^3)/a^3, 1/3*(3*a*cos(f*x + e) + 3*I*a*sin(f*x + e) + 2*b)/a) +
sqrt(2)*(-I*(3*a - 2*b)*g*cos(f*x + e) - I*(3*a - 2*b)*g)*sqrt(a*g)*weier
strassPInverse(-4/3*(3*a^2 - 4*b^2)/a^2, 8/27*(9*a^2*b - 8*b^3)/a^3, 1/3*(
3*a*cos(f*x + e) - 3*I*a*sin(f*x + e) + 2*b)/a) - 3*sqrt(2)*(I*a*g*cos(f*x
+ e) + I*a*g)*sqrt(a*g)*weierstrassZeta(-4/3*(3*a^2 - 4*b^2)/a^2, 8/27*(9
*a^2*b - 8*b^3)/a^3, weierstrassPInverse(-4/3*(3*a^2 - 4*b^2)/a^2, 8/27*(9
*a^2*b - 8*b^3)/a^3, 1/3*(3*a*cos(f*x + e) + 3*I*a*sin(f*x + e) + 2*b)/a))
- 3*sqrt(2)*(-I*a*g*cos(f*x + e) - I*a*g)*sqrt(a*g)*weierstrassZeta(-4/3*
(3*a^2 - 4*b^2)/a^2, 8/27*(9*a^2*b - 8*b^3)/a^3, weierstrassPInverse(-4/3*
(3*a^2 - 4*b^2)/a^2, 8/27*(9*a^2*b - 8*b^3)/a^3, 1/3*(3*a*cos(f*x + e) - 3
*I*a*sin(f*x + e) + 2*b)/a)))/((a^2 - a*b)*c*f*cos(f*x + e) + (a^2 - a*b)*
c*f)
```

3.277.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \frac{\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)} \sec(e + fx) + \sqrt{a + b \sec(e + fx)}} dx}{c}$$

input

```
integrate((g*sec(f*x+e))**(3/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x
)
```

output

```
Integral((g*sec(e + f*x))**(3/2)/(sqrt(a + b*sec(e + f*x))*sec(e + f*x) +
sqrt(a + b*sec(e + f*x))), x)/c
```

3.277.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{3/2}}{\sqrt{b \sec(fx + e) + a}(c \sec(fx + e) + c)} dx$$

input

```
integrate((g*sec(f*x+e))^(3/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,
algorithm="maxima")
```

3.277. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx$

output `integrate((g*sec(f*x + e))^(3/2)/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.277.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{3/2}}{\sqrt{b \sec(fx + e) + a}(c \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(3/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((g*sec(f*x + e))^(3/2)/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.277.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{3/2}}{\sqrt{a + \frac{b}{\cos(e+fx)}} \left(c + \frac{c}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(3/2)/((a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))),x)`

output `int((g/cos(e + f*x))^(3/2)/((a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))), x)`

$$3.278 \quad \int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$$

3.278.1 Optimal result	2035
3.278.2 Mathematica [F]	2036
3.278.3 Rubi [A] (verified)	2036
3.278.4 Maple [C] (verified)	2044
3.278.5 Fricas [F(-1)]	2044
3.278.6 Sympy [F(-1)]	2045
3.278.7 Maxima [F]	2045
3.278.8 Giac [F]	2045
3.278.9 Mupad [F(-1)]	2046

3.278.1 Optimal result

Integrand size = 39, antiderivative size = 312

$$\begin{aligned} & \int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx = \\ & \frac{g^2(b+a \cos(e+fx))E(\frac{1}{2}(e+fx)|\frac{2a}{a+b})\sqrt{g \sec(e+fx)}}{(a-b)cf\sqrt{\frac{b+a \cos(e+fx)}{a+b}}\sqrt{a+b \sec(e+fx)}} \\ & - \frac{g^2\sqrt{\frac{b+a \cos(e+fx)}{a+b}}\text{EllipticF}(\frac{1}{2}(e+fx),\frac{2a}{a+b})\sqrt{g \sec(e+fx)}}{cf\sqrt{a+b \sec(e+fx)}} \\ & + \frac{2g^2\sqrt{\frac{b+a \cos(e+fx)}{a+b}}\text{EllipticPi}(2,\frac{1}{2}(e+fx),\frac{2a}{a+b})\sqrt{g \sec(e+fx)}}{cf\sqrt{a+b \sec(e+fx)}} \\ & + \frac{g^2(b+a \cos(e+fx))\sqrt{g \sec(e+fx)}\sin(e+fx)}{(a-b)f(c+c \cos(e+fx))\sqrt{a+b \sec(e+fx)}} \end{aligned}$$

output
$$g^2(b+a\cos(fx+e))\sin(fx+e)(g\sec(fx+e))^{1/2}/(a-b)/f/(c+c\cos(fx+e))/(a+b\sec(fx+e))^{1/2}-g^2(b+a\cos(fx+e))(\cos(1/2fx+1/2e))^2)^{1/2}/\cos(1/2fx+1/2e)*\text{EllipticE}(\sin(1/2fx+1/2e),2^{1/2}*(a/(a+b))^{1/2})*(g\sec(fx+e))^{1/2}/(a-b)/c/f/((b+a\cos(fx+e))/(a+b))^{1/2}/(a+b\sec(fx+e))^{1/2}-g^2(\cos(1/2fx+1/2e))^2)^{1/2}/\cos(1/2fx+1/2e)*\text{EllipticF}(\sin(1/2fx+1/2e),2^{1/2}*(a/(a+b))^{1/2})*((b+a\cos(fx+e))/(a+b))^{1/2})*(g\sec(fx+e))^{1/2}/c/f/(a+b\sec(fx+e))^{1/2}+2g^2(\cos(1/2fx+1/2e))^2)^{1/2}/\cos(1/2fx+1/2e)*\text{EllipticPi}(\sin(1/2fx+1/2e),2^{1/2}*(a/(a+b))^{1/2})*((b+a\cos(fx+e))/(a+b))^{1/2})*(g\sec(fx+e))^{1/2}/c/f/(a+b\sec(fx+e))^{1/2}$$

3.278.2 Mathematica [F]

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx$$

input `Integrate[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])),x]`

output `Integrate[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])), x]`

3.278.3 Rubi [A] (verified)

Time = 2.65 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.99, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3042, 4467, 3042, 4346, 3042, 3286, 3042, 3284, 4463, 3042, 3247, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(g \sec(e + fx))^{5/2}}{(c \sec(e + fx) + c)\sqrt{a + b \sec(e + fx)}} dx$$

↓ 3042

3.278. $\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx$

$$\begin{aligned}
& \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{5/2}}{(c \csc(e + fx + \frac{\pi}{2}) + c) \sqrt{a + b \csc(e + fx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4467} \\
& \frac{g \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}} dx}{c} - g \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(\sec(e + fx)c + c)} dx \\
& \quad \downarrow \text{3042} \\
& \frac{g \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}} dx}{c} - g \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(\csc(e + fx + \frac{\pi}{2})c + c)} dx \\
& \quad \downarrow \text{4346} \\
& \frac{g^2 \sqrt{g \sec(e + fx)} \sqrt{a \cos(e + fx) + b} \int \frac{\sec(e + fx)}{\sqrt{b + a \cos(e + fx)}} dx}{c \sqrt{a + b \sec(e + fx)}} - \\
& \quad g \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(\csc(e + fx + \frac{\pi}{2})c + c)} dx \\
& \quad \downarrow \text{3042} \\
& \frac{g^2 \sqrt{g \sec(e + fx)} \sqrt{a \cos(e + fx) + b} \int \frac{1}{\sin(e + fx + \frac{\pi}{2}) \sqrt{b + a \sin(e + fx + \frac{\pi}{2})}} dx}{c \sqrt{a + b \sec(e + fx)}} - \\
& \quad g \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(\csc(e + fx + \frac{\pi}{2})c + c)} dx \\
& \quad \downarrow \text{3286} \\
& \frac{g^2 \sqrt{g \sec(e + fx)} \sqrt{\frac{a \cos(e + fx) + b}{a + b}} \int \frac{\sec(e + fx)}{\sqrt{\frac{b}{a + b} + \frac{a \cos(e + fx)}{a + b}}} dx}{c \sqrt{a + b \sec(e + fx)}} - \\
& \quad g \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(\csc(e + fx + \frac{\pi}{2})c + c)} dx \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.278. $\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx$

$$\begin{aligned}
& \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sin(e+fx+\frac{\pi}{2}) \sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b}}} dx}{c \sqrt{a+b \sec(e+fx)}} \\
& \frac{g \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (\csc(e+fx+\frac{\pi}{2}) c+c)} dx}{\sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{3284} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} \\
& \frac{g \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (\csc(e+fx+\frac{\pi}{2}) c+c)} dx}{\sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{4463} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} \\
& \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{1}{\sqrt{b+a \cos(e+fx)} (\cos(e+fx)c+c)} dx}{\sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} \\
& \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})} (\sin(e+fx+\frac{\pi}{2})c+c)} dx}{\sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{3247} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} \\
& \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \left(-\frac{a \int -\frac{\cos(e+fx)c+c}{2\sqrt{b+a \cos(e+fx)}} dx}{c^2(a-b)} - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.278. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$

$$\begin{aligned}
 & \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \\
 & \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \left(\frac{a \int \frac{\cos(e+fx)c+c}{\sqrt{b+a \cos(e+fx)}} dx}{2c^2(a-b)} - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \\
 & \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \left(\frac{a \int \frac{\sin(e+fx+\frac{\pi}{2})c+c}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{2c^2(a-b)} - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
 & \quad \downarrow \text{3231} \\
 & \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \\
 & \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \cos(e+fx)}} dx}{a} + \frac{c \int \sqrt{b+a \cos(e+fx)} dx}{a} \right)}{2c^2(a-b)} - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \\
 & \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \left(\frac{a \left(\frac{c(a-b) \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c \int \sqrt{b+a \sin(e+fx+\frac{\pi}{2})} dx}{a} \right)}{2c^2(a-b)} - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx)+c)} \right)}{\sqrt{a+b \sec(e+fx)}} \\
 & \quad \downarrow \text{3134}
 \end{aligned}$$

3.278. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$

$$\frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b}}{2c^2(a-b)} \left(a \left(\frac{c^{(a-b)} \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c \sqrt{a \cos(e+fx)+b} \int \sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b}} dx}{a \sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx))} \right)$$

$$\sqrt{a+b \sec(e+fx)}$$

3042

$$\frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b}}{2c^2(a-b)} \left(a \left(\frac{c^{(a-b)} \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{c \sqrt{a \cos(e+fx)+b} \int \sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b}} dx}{a \sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx))} \right)$$

$$\sqrt{a+b \sec(e+fx)}$$

3132

$$\frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b}}{2c^2(a-b)} \left(a \left(\frac{c^{(a-b)} \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{a} + \frac{2c \sqrt{a \cos(e+fx)+b} E\left(\frac{1}{2}(e+fx) \mid \frac{2a}{a+b}\right)}{af \sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) - \frac{\sin(e+fx) \sqrt{a \cos(e+fx)+b}}{f(a-b)(c \cos(e+fx))} \right)$$

$$\sqrt{a+b \sec(e+fx)}$$

3142

3.278. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$

$$\frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b}}{2c^2(a-b)} \left(a \frac{c(a-b) \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b}}} dx}{a \sqrt{a \cos(e+fx)+b}} + \frac{2c \sqrt{a \cos(e+fx)+b} E\left(\frac{1}{2}(e+fx) \mid \frac{2a}{a+b}\right)}{af \sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) - \frac{\sin}{j}$$

$$\sqrt{a+b \sec(e+fx)}$$

↓ 3042

$$\frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b}}{2c^2(a-b)} \left(a \frac{c(a-b) \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a \sin\left(e+fx+\frac{\pi}{2}\right)}{a+b}}} dx}{a \sqrt{a \cos(e+fx)+b}} + \frac{2c \sqrt{a \cos(e+fx)+b} E\left(\frac{1}{2}(e+fx) \mid \frac{2a}{a+b}\right)}{af \sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) - \frac{\sin}{j}$$

$$\sqrt{a+b \sec(e+fx)}$$

↓ 3140

$$\frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{cf \sqrt{a+b \sec(e+fx)}} - \frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b}}{2c^2(a-b)} \left(a \frac{2c(a-b) \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{af \sqrt{a \cos(e+fx)+b}} + \frac{2c \sqrt{a \cos(e+fx)+b} E\left(\frac{1}{2}(e+fx) \mid \frac{2a}{a+b}\right)}{af \sqrt{\frac{a \cos(e+fx)+b}{a+b}}} \right) - \frac{\sin}{j}$$

$$\sqrt{a+b \sec(e+fx)}$$

input `Int[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + c*Sec[e + f*x])) ,x]`

3.278. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+c \sec(e+fx))} dx$

```
output (2*g^2*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[2, (e + f*x)/2, (2*a)
/(a + b)]*Sqrt[g*Sec[e + f*x]]/(c*f*Sqrt[a + b*Sec[e + f*x]]) - (g^2*Sqrt
[b + a*Cos[e + f*x]]*Sqrt[g*Sec[e + f*x]]*((a*((2*c*Sqrt[b + a*Cos[e + f*x
]]*EllipticE[(e + f*x)/2, (2*a)/(a + b)])/(a*f*Sqrt[(b + a*Cos[e + f*x])/(
a + b)]) + (2*(a - b)*c*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticF[(e +
f*x)/2, (2*a)/(a + b)]/(a*f*Sqrt[b + a*Cos[e + f*x]])))/(2*(a - b)*c^2) -
(Sqrt[b + a*Cos[e + f*x]]*Sin[e + f*x])/((a - b)*f*(c + c*Cos[e + f*x]))
)/Sqrt[a + b*Sec[e + f*x]]
```

3.278.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3142 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a
+ b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b)
+ (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && !GtQ[a + b, 0]
```

rule 3231 `Int[((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])/Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3247 `Int[((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)/((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(b*c - a*d)*(a + b*Sin[e + f*x]))], x] + Simp[d/(a*(b*c - a*d)) Int[(c + d*Sin[e + f*x])^n*(a*n - b*(n + 1)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, 0] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 4346 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[d*Sqrt[d*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sin[e + f*x]*Sqrt[b + a*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4463 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(3/2)/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))], x_Symbol] := Simp[g*Sqrt[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

```
rule 4467 Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(5/2)/(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))), x_Symbol] := Simp[g/d Int[(g*Csc[e + f*x])^(3/2)/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[c*(g/d Int[(g*Csc[e + f*x])^(3/2)/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
```

3.278.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 7.97 (sec) , antiderivative size = 323, normalized size of antiderivative = 1.04

method	result
default	$i \left(4 \operatorname{EllipticF} \left(i(-\cot(fx+e)+\csc(fx+e)), \sqrt{-\frac{a-b}{a+b}} \right) a - 2b \operatorname{EllipticF} \left(i(-\cot(fx+e)+\csc(fx+e)), \sqrt{-\frac{a-b}{a+b}} \right) - a \operatorname{EllipticE} \left(i(-\cot(fx+e)+\csc(fx+e)), \sqrt{-\frac{a-b}{a+b}} \right) \right)$

```
input int((g*sec(f*x+e))^(5/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
output I/c/f/(a-b)*(4*EllipticF(I*(-cot(f*x+e)+csc(f*x+e)),(-a-b)/(a+b))^(1/2))*a-2*b*EllipticF(I*(-cot(f*x+e)+csc(f*x+e)),(-a-b)/(a+b))^(1/2))-a*EllipticE(I*(-cot(f*x+e)+csc(f*x+e)),(-a-b)/(a+b))^(1/2))-b*EllipticE(I*(-cot(f*x+e)+csc(f*x+e)),(-a-b)/(a+b))^(1/2))-4*EllipticPi(I*(-cot(f*x+e)+csc(f*x+e)),-1,I*((a-b)/(a+b))^(1/2))*a+4*b*EllipticPi(I*(-cot(f*x+e)+csc(f*x+e)),-1,I*((a-b)/(a+b))^(1/2)))*(a+b*sec(f*x+e))^(1/2)*(g*sec(f*x+e))^(1/2)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*g^2/(1/(cos(f*x+e)+1))^(1/2)/(b+a*cos(f*x+e))*cos(f*x+e)
```

3.278.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \text{Timed out}$$

```
input integrate((g*sec(f*x+e))^(5/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="fracas")
```

3.278. $\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx$

output Timed out

3.278.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \text{Timed out}$$

input `integrate((g*sec(f*x+e))**(5/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x)`

output Timed out

3.278.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{5/2}}{\sqrt{b \sec(fx + e) + a}(c \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(5/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="maxima")`

output `integrate((g*sec(f*x + e))^(5/2)/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.278.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{5/2}}{\sqrt{b \sec(fx + e) + a}(c \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(5/2)/(c+c*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="giac")`

output `integrate((g*sec(f*x + e))^(5/2)/(sqrt(b*sec(f*x + e) + a)*(c*sec(f*x + e) + c)), x)`

3.278. $\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx$

3.278.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + c \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{5/2}}{\sqrt{a + \frac{b}{\cos(e+fx)}} \left(c + \frac{c}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(5/2)/((a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))),x)`

output `int((g/cos(e + f*x))^(5/2)/((a + b/cos(e + f*x))^(1/2)*(c + c/cos(e + f*x))), x)`

3.279
$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

3.279.1 Optimal result 2047
 3.279.2 Mathematica [A] (verified) 2048
 3.279.3 Rubi [A] (verified) 2048
 3.279.4 Maple [A] (verified) 2050
 3.279.5 Fracas [F(-1)] 2051
 3.279.6 Sympy [F] 2051
 3.279.7 Maxima [F] 2051
 3.279.8 Giac [F] 2052
 3.279.9 Mupad [F(-1)] 2052

3.279.1 Optimal result

Integrand size = 33, antiderivative size = 213

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

$$= \frac{2\sqrt{a+b}\cot(e+fx)\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b\sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)\sqrt{\frac{b(1-\sec(e+fx))}{a+b}}\sqrt{-\frac{b(1+\sec(e+fx))}{a-b}}}{df} - \frac{2(bc-ad)\operatorname{EllipticPi}\left(\frac{2d}{c+d}, \arcsin\left(\frac{\sqrt{1-\sec(e+fx)}}{\sqrt{2}}\right), \frac{2b}{a+b}\right)\sqrt{\frac{a+b\sec(e+fx)}{a+b}}\tan(e+fx)}{d(c+d)f\sqrt{a+b\sec(e+fx)}\sqrt{-\tan^2(e+fx)}}$$

output

```
2*cot(f*x+e)*EllipticF((a+b*sec(f*x+e))^(1/2)/(a+b)^(1/2),((a+b)/(a-b))^(1/2))*
(a+b)^(1/2)*(b*(1-sec(f*x+e))/(a+b))^(1/2)*(-b*(1+sec(f*x+e))/(a-b))^(1/2)/d/f-
2*(-a*d+b*c)*EllipticPi(1/2*(1-sec(f*x+e))^(1/2)*2^(1/2),2*d/(c+d),2^(1/2)*
(b/(a+b))^(1/2))*((a+b*sec(f*x+e))/(a+b))^(1/2)*tan(f*x+e)/d/(c+d)/f/
(a+b*sec(f*x+e))^(1/2)/(-tan(f*x+e)^2)^(1/2)
```


3.279.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.86

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

$$= \frac{4 \cos^2\left(\frac{1}{2}(e+fx)\right) \sqrt{\frac{\cos(e+fx)}{1+\cos(e+fx)}} \sqrt{\frac{b+a\cos(e+fx)}{(a+b)(1+\cos(e+fx))}} \left((a-b)(c+d) \operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right), \frac{a-b}{a+b}\right) \right)}{(c-d)(c+d)f(b+a\cos(e+fx))}$$

input `Integrate[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`output `(4*Cos[(e + f*x)/2]^2*Sqrt[Cos[e + f*x]/(1 + Cos[e + f*x])] * Sqrt[(b + a*Cos[e + f*x])/((a + b)*(1 + Cos[e + f*x]))] * ((a - b)*(c + d)*EllipticF[ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)] + 2*(b*c - a*d)*EllipticPi[(c - d)/(c + d), ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)]) * Sqrt[a + b*Sec[e + f*x]])/((c - d)*(c + d)*f*(b + a*Cos[e + f*x]))`**3.279.3 Rubi [A] (verified)**Time = 0.70 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {3042, 4457, 3042, 4319, 4461}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)\sqrt{a+b\csc\left(e+fx+\frac{\pi}{2}\right)}}{c+d\csc\left(e+fx+\frac{\pi}{2}\right)} dx$$

$$\downarrow \text{4457}$$

$$\frac{b \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}} dx}{d} - \frac{(bc-ad) \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx}{d}$$

$$\downarrow \text{3042}$$

$$\frac{b \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})}} dx}{d} - \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d}$$

↓ 4319

$$\frac{2\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{df} - \frac{(bc-ad) \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d}$$

↓ 4461

$$\frac{2\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{df} - \frac{2(bc-ad) \tan(e+fx) \sqrt{\frac{a+b \sec(e+fx)}{a+b}} \text{EllipticPi}\left(\frac{2d}{c+d}, \arcsin\left(\frac{\sqrt{1-\sec(e+fx)}}{\sqrt{2}}\right), \frac{2b}{a+b}\right)}{df(c+d) \sqrt{-\tan^2(e+fx)} \sqrt{a+b \sec(e+fx)}}$$

input `Int[(Sec[e + f*x]*Sqrt[a + b*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`

output `(2*Sqrt[a + b]*Cot[e + f*x]*EllipticF[ArcSin[Sqrt[a + b*Sec[e + f*x]]/Sqrt[a + b]], (a + b)/(a - b)]*Sqrt[(b*(1 - Sec[e + f*x]))/(a + b)]*Sqrt[-((b*(1 + Sec[e + f*x]))/(a - b))])/(d*f) - (2*(b*c - a*d)*EllipticPi[(2*d)/(c + d), ArcSin[Sqrt[1 - Sec[e + f*x]]/Sqrt[2]], (2*b)/(a + b)]*Sqrt[(a + b*Sec[e + f*x])/(a + b)]*Tan[e + f*x])/(d*(c + d)*f*Sqrt[a + b*Sec[e + f*x]]*Sqrt[-Tan[e + f*x]^2])`

3.279.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4319 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*(Rt[a + b, 2]/(b*f*Cot[e + f*x]))*Sqrt[(b*(1 - Csc[e + f*x]))/(a + b)]*Sqrt[(-b)*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Csc[e + f*x]]/Rt[a + b, 2]], (a + b)/(a - b)], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

3.279. $\int \frac{\sec(e+fx)\sqrt{a+b \sec(e+fx)}}{c+d \sec(e+fx)} dx$

rule 4457 `Int[(csc[(e_.) + (f_.)*(x_)]*Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)])/(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[b/d Int[Csc[e + f*x]/Sqrt[a + b*Csc[e + f*x]], x, x] - Simp[(b*c - a*d)/d Int[Csc[e + f*x]/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x])), x, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4461 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))), x_Symbol] := Simp[-2*(Cot[e + f*x]/(f*(c + d)*Sqrt[a + b*Csc[e + f*x]]*Sqrt[-Cot[e + f*x]^2]))*Sqrt[(a + b*Csc[e + f*x])]/(a + b)*EllipticPi[2*(d/(c + d)), ArcSin[Sqrt[1 - Csc[e + f*x]]]/Sqrt[2]], 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.279.4 Maple [A] (verified)

Time = 8.32 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.48

method	result
default	$-\frac{2(\cos(fx+e)+1)\left(\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)ac+\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)ad-\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)bc-\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)bd-2\text{EllipticPi}\left(\cot(fx+e)-\csc(fx+e),\frac{c-d}{c+d},\sqrt{\frac{a-b}{a+b}}\right)a*d+2\text{EllipticPi}\left(\cot(fx+e)-\csc(fx+e),\frac{c-d}{c+d},\sqrt{\frac{a-b}{a+b}}\right)b*c\right)}{(a+b)\sqrt{a+b}\sqrt{\cos(fx+e)+1}}\frac{1}{(a+b)\sqrt{a+b}\sqrt{\cos(fx+e)+1}}\frac{1}{(a+b)\sqrt{a+b}\sqrt{\cos(fx+e)+1}}$

input `int(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output `-2/f/(c+d)/(c-d)*(cos(f*x+e)+1)*(EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*a*c+EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*a*d-EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*b*c-EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*b*d-2*EllipticPi(cot(f*x+e)-csc(f*x+e),(c-d)/(c+d),((a-b)/(a+b))^(1/2))*a*d+2*EllipticPi(cot(f*x+e)-csc(f*x+e),(c-d)/(c+d),((a-b)/(a+b))^(1/2))*b*c)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*(a+b*sec(f*x+e))^(1/2)/(b+a*cos(f*x+e))`

3.279.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)\sqrt{a + b\sec(e + fx)}}{c + d\sec(e + fx)} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="fricas")`

output `Timed out`

3.279.6 Sympy [F]

$$\int \frac{\sec(e + fx)\sqrt{a + b\sec(e + fx)}}{c + d\sec(e + fx)} dx = \int \frac{\sqrt{a + b\sec(e + fx)}\sec(e + fx)}{c + d\sec(e + fx)} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e)),x)`

output `Integral(sqrt(a + b*sec(e + f*x))*sec(e + f*x)/(c + d*sec(e + f*x)), x)`

3.279.7 Maxima [F]

$$\int \frac{\sec(e + fx)\sqrt{a + b\sec(e + fx)}}{c + d\sec(e + fx)} dx = \int \frac{\sqrt{b\sec(fx + e) + a}\sec(fx + e)}{d\sec(fx + e) + c} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="maxima")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(d*sec(f*x + e) + c), x)`

3.279.8 Giac [F]

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{b\sec(fx+e)+a}\sec(fx+e)}{d\sec(fx+e)+c} dx$$

input `integrate(sec(f*x+e)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x, algorithm="giac")`

output `integrate(sqrt(b*sec(f*x + e) + a)*sec(f*x + e)/(d*sec(f*x + e) + c), x)`

3.279.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(e+fx)\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx = \int \frac{\sqrt{a + \frac{b}{\cos(e+fx)}}}{\cos(e+fx) \left(c + \frac{d}{\cos(e+fx)}\right)} dx$$

input `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))),x)`

output `int((a + b/cos(e + f*x))^(1/2)/(cos(e + f*x)*(c + d/cos(e + f*x))), x)`

3.280
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+d \sec(e+fx)} dx$$

3.280.1 Optimal result 2053
 3.280.2 Mathematica [C] (verified) 2053
 3.280.3 Rubi [A] (verified) 2054
 3.280.4 Maple [C] (verified) 2058
 3.280.5 Fracas [F(-1)] 2058
 3.280.6 Sympy [F] 2059
 3.280.7 Maxima [F] 2059
 3.280.8 Giac [F] 2059
 3.280.9 Mupad [F(-1)] 2060

3.280.1 Optimal result

Integrand size = 39, antiderivative size = 170

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+d \sec(e+fx)} dx = \frac{2bg \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right) \sqrt{g \sec(e+fx)}}{df \sqrt{a+b \sec(e+fx)}} - \frac{2(bc-ad)g \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticPi}\left(\frac{2c}{c+d}, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right) \sqrt{g \sec(e+fx)}}{d(c+d)f \sqrt{a+b \sec(e+fx)}}$$

output

```
2*b*g*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e), 2, 2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/d/f/(a+b*sec(f*x+e))^(1/2)-2*(-a*d+b*c)*g*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e), 2*c/(c+d), 2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/d/(c+d)/f/(a+b*sec(f*x+e))^(1/2)
```

3.280.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 25.56 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.31

$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+d \sec(e+fx)} dx = \frac{2ig \sqrt{-\frac{a(-1+\cos(e+fx))}{a+b}} \sqrt{\frac{a(1+\cos(e+fx))}{a-b}} \cot(e+fx) \left(\text{EllipticPi}\left(1 - \frac{a}{b}, \text{iarcsinh}\left(\sqrt{\frac{1}{a-b}} \sqrt{b+a \cos(e+fx)}\right)\right) \sqrt{\frac{1}{a-b}} df \sqrt{b+a \cos(e+fx)} \right)}{\sqrt{\frac{1}{a-b}} df \sqrt{b+a \cos(e+fx)}}$$

3.280.
$$\int \frac{(g \sec(e+fx))^{3/2} \sqrt{a+b \sec(e+fx)}}{c+d \sec(e+fx)} dx$$

input `Integrate[((g*Sec[e + f*x])^(3/2)*Sqrt[a + b*Sec[e + f*x]])/(c + d*Sec[e + f*x]),x]`

output `((-2*I)*g*Sqrt[-((a*(-1 + Cos[e + f*x]))/(a + b))]*Sqrt[(a*(1 + Cos[e + f*x]))/(a - b)]*Cot[e + f*x]*(EllipticPi[1 - a/b, I*ArcSinh[Sqrt[(a - b)^(-1)]*Sqrt[b + a*Cos[e + f*x]]], (-a + b)/(a + b)] - EllipticPi[((a - b)*c)/(-(b*c) + a*d), I*ArcSinh[Sqrt[(a - b)^(-1)]*Sqrt[b + a*Cos[e + f*x]]], (-a + b)/(a + b)]*Sqrt[g*Sec[e + f*x]]*Sqrt[a + b*Sec[e + f*x]])/(Sqrt[(a - b)^(-1)]*d*f*Sqrt[b + a*Cos[e + f*x]])`

3.280.3 Rubi [A] (verified)

Time = 2.06 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4459, 3042, 4346, 3042, 3286, 3042, 3284, 4463, 3042, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2} \sqrt{a + b \csc(e + fx + \frac{\pi}{2})}}{c + d \csc(e + fx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4459} \\
 & \frac{b \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}} dx}{d} - \frac{(bc - ad) \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}} dx}{d} - \frac{(bc - ad) \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(c + d \csc(e + fx + \frac{\pi}{2}))} dx}{d} \\
 & \quad \downarrow \text{4346}
 \end{aligned}$$

3.280. $\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx$

$$\begin{aligned}
& \frac{bg\sqrt{g\sec(e+fx)}\sqrt{a\cos(e+fx)+b}\int\frac{\sec(e+fx)}{\sqrt{b+a\cos(e+fx)}}dx}{d\sqrt{a+b\sec(e+fx)}} \\
& \frac{(bc-ad)\int\frac{(g\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}(c+d\csc(e+fx+\frac{\pi}{2}))}dx}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{bg\sqrt{g\sec(e+fx)}\sqrt{a\cos(e+fx)+b}\int\frac{1}{\sin(e+fx+\frac{\pi}{2})\sqrt{b+a\sin(e+fx+\frac{\pi}{2})}}dx}{d\sqrt{a+b\sec(e+fx)}} \\
& \frac{(bc-ad)\int\frac{(g\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}(c+d\csc(e+fx+\frac{\pi}{2}))}dx}{d} \\
& \quad \downarrow \text{3286} \\
& \frac{bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\int\frac{\sec(e+fx)}{\sqrt{\frac{b}{a+b}+\frac{a\cos(e+fx)}{a+b}}}dx}{d\sqrt{a+b\sec(e+fx)}} \\
& \frac{(bc-ad)\int\frac{(g\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}(c+d\csc(e+fx+\frac{\pi}{2}))}dx}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\int\frac{1}{\sin(e+fx+\frac{\pi}{2})\sqrt{\frac{b}{a+b}+\frac{a\sin(e+fx+\frac{\pi}{2})}{a+b}}}dx}{d\sqrt{a+b\sec(e+fx)}} \\
& \frac{(bc-ad)\int\frac{(g\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}(c+d\csc(e+fx+\frac{\pi}{2}))}dx}{d} \\
& \quad \downarrow \text{3284} \\
& \frac{2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\text{EllipticPi}\left(2,\frac{1}{2}(e+fx),\frac{2a}{a+b}\right)}{df\sqrt{a+b\sec(e+fx)}} \\
& \frac{(bc-ad)\int\frac{(g\csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b\csc(e+fx+\frac{\pi}{2})}(c+d\csc(e+fx+\frac{\pi}{2}))}dx}{d} \\
& \quad \downarrow \text{4463} \\
& \frac{2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\text{EllipticPi}\left(2,\frac{1}{2}(e+fx),\frac{2a}{a+b}\right)}{df\sqrt{a+b\sec(e+fx)}} \\
& \frac{g(bc-ad)\sqrt{g\sec(e+fx)}\sqrt{a\cos(e+fx)+b}\int\frac{1}{\sqrt{b+a\cos(e+fx)}(d+c\cos(e+fx))}dx}{d\sqrt{a+b\sec(e+fx)}}
\end{aligned}$$

3.280. $\int \frac{(g\sec(e+fx))^{3/2}\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df\sqrt{a+b\sec(e+fx)}} - \\
& \frac{g(bc-ad)\sqrt{g\sec(e+fx)}\sqrt{a\cos(e+fx)+b}\int\frac{1}{\sqrt{b+a\sin(e+fx+\frac{\pi}{2})}(d+c\sin(e+fx+\frac{\pi}{2}))}dx}{d\sqrt{a+b\sec(e+fx)}} \\
& \downarrow 3286 \\
& \frac{2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df\sqrt{a+b\sec(e+fx)}} - \\
& \frac{g(bc-ad)\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\int\frac{1}{\sqrt{\frac{b}{a+b}+\frac{a\cos(e+fx)}{a+b}}(d+c\cos(e+fx))}dx}{d\sqrt{a+b\sec(e+fx)}} \\
& \downarrow 3042 \\
& \frac{2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df\sqrt{a+b\sec(e+fx)}} - \\
& \frac{g(bc-ad)\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\int\frac{1}{\sqrt{\frac{b}{a+b}+\frac{a\sin(e+fx+\frac{\pi}{2})}{a+b}}(d+c\sin(e+fx+\frac{\pi}{2}))}dx}{d\sqrt{a+b\sec(e+fx)}} \\
& \downarrow 3284 \\
& \frac{2bg\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df\sqrt{a+b\sec(e+fx)}} - \\
& \frac{2g(bc-ad)\sqrt{g\sec(e+fx)}\sqrt{\frac{a\cos(e+fx)+b}{a+b}}\operatorname{EllipticPi}\left(\frac{2c}{c+d}, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df(c+d)\sqrt{a+b\sec(e+fx)}}
\end{aligned}$$

input `Int[((g*Sec[e + f*x])^(3/2)*Sqrt[a + b*Sec[e + f*x]])/(c + d*Sec[e + f*x]), x]`

output `(2*b*g*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[2, (e + f*x)/2, (2*a)/(a + b)]*Sqrt[g*Sec[e + f*x]])/(d*f*Sqrt[a + b*Sec[e + f*x]]) - (2*(b*c - a*d)*g*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[(2*c)/(c + d), (e + f*x)/2, (2*a)/(a + b)]*Sqrt[g*Sec[e + f*x]])/(d*(c + d)*f*Sqrt[a + b*Sec[e + f*x]])`

3.280. $\int \frac{(g\sec(e+fx))^{3/2}\sqrt{a+b\sec(e+fx)}}{c+d\sec(e+fx)} dx$

3.280.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 4346 `Int[(csc[(e_) + (f_)*(x_)]*(d_))^(3/2)/Sqrt[csc[(e_) + (f_)*(x_)]*(b_) + (a_)], x_Symbol] := Simp[d*Sqrt[d*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sin[e + f*x]*Sqrt[b + a*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4459 `Int[((csc[(e_) + (f_)*(x_)]*(g_))^(3/2)*Sqrt[csc[(e_) + (f_)*(x_)]*(b_) + (a_)])/(csc[(e_) + (f_)*(x_)]*(d_) + (c_)), x_Symbol] := Simp[b/d Int[(g*Csc[e + f*x])^(3/2)/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[(b*c - a*d)/d Int[(g*Csc[e + f*x])^(3/2)/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 4463 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(3/2)/(Sqrt[csc[(e_) + (f_)*(x_)]*(b_) + (a_)])*(csc[(e_) + (f_)*(x_)]*(d_) + (c_)), x_Symbol] := Simp[g*Sqrt[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

3.280.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 8.22 (sec) , antiderivative size = 445, normalized size of antiderivative = 2.62

method	result
default	$-\frac{2ig\left(\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{a-b}{a+b}}\right)acd+\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{a-b}{a+b}}\right)ad^2-\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{a-b}{a+b}}\right)ad-\text{EllipticF}\left(i(\cot(fx+e)-\csc(fx+e)),\sqrt{-\frac{a-b}{a+b}}\right)ad\right)}{\dots}$

input `int((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -2*I*g/f/d/(c+d)/(c-d)*(EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b)) \\ & ^{(1/2)})*a*c*d+EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^{(1/2)}*a* \\ & d^2-EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^{(1/2)}*b*c*d-EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^{(1/2)}*b*d^2-2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-1,I*((a-b)/(a+b))^{(1/2)})*b*c^2+2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-1,I*((a-b)/(a+b))^{(1/2)})*b*d^2-2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-(c-d)/(c+d),I*((a-b)/(a+b))^{(1/2)})*a*c*d+2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)),-(c-d)/(c+d),I*((a-b)/(a+b))^{(1/2)})*b*c^2)*cos(f*x+e)*(a+b*sec(f*x+e))^{(1/2)}*(g*sec(f*x+e))^{(1/2)}*(1/(a+b)*(b+a*cos(f*x+e)))/(cos(f*x+e)+1)^{(1/2)}/(b+a*cos(f*x+e))/(1/(cos(f*x+e)+1))^{(1/2)} \end{aligned}$$

3.280.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx = \text{Timed out}$$

input `integrate((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,algorithm="fricas")`

output `Timed out`

3.280.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx = \int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx$$

input `integrate((g*sec(f*x+e))**(3/2)*(a+b*sec(f*x+e))**(1/2)/(c+d*sec(f*x+e)),x)`

output `Integral((g*sec(e + f*x))**(3/2)*sqrt(a + b*sec(e + f*x))/(c + d*sec(e + f*x)), x)`

3.280.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx = \int \frac{\sqrt{b \sec(fx + e) + a} (g \sec(fx + e))^{3/2}}{d \sec(fx + e) + c} dx$$

input `integrate((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,algorithm="maxima")`

output `integrate(sqrt(b*sec(f*x + e) + a)*(g*sec(f*x + e))^(3/2)/(d*sec(f*x + e) + c), x)`

3.280.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx = \int \frac{\sqrt{b \sec(fx + e) + a} (g \sec(fx + e))^{3/2}}{d \sec(fx + e) + c} dx$$

input `integrate((g*sec(f*x+e))^(3/2)*(a+b*sec(f*x+e))^(1/2)/(c+d*sec(f*x+e)),x,algorithm="giac")`

output `integrate(sqrt(b*sec(f*x + e) + a)*(g*sec(f*x + e))^(3/2)/(d*sec(f*x + e) + c), x)`

3.280.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2} \sqrt{a + b \sec(e + fx)}}{c + d \sec(e + fx)} dx = \int \frac{\sqrt{a + \frac{b}{\cos(e+fx)}} \left(\frac{g}{\cos(e+fx)}\right)^{3/2}}{c + \frac{d}{\cos(e+fx)}} dx$$

input `int(((a + b/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c + d/cos(e + f*x)),x)`

output `int(((a + b/cos(e + f*x))^(1/2)*(g/cos(e + f*x))^(3/2))/(c + d/cos(e + f*x)), x)`

3.281
$$\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.281.1 Optimal result 2061
 3.281.2 Mathematica [A] (verified) 2061
 3.281.3 Rubi [A] (verified) 2062
 3.281.4 Maple [B] (verified) 2063
 3.281.5 Fricas [F(-1)] 2064
 3.281.6 Sympy [F] 2064
 3.281.7 Maxima [F] 2064
 3.281.8 Giac [F] 2065
 3.281.9 Mupad [F(-1)] 2065

3.281.1 Optimal result

Integrand size = 33, antiderivative size = 102

$$\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{2 \operatorname{EllipticPi}\left(\frac{2d}{c+d}, \arcsin\left(\frac{\sqrt{1-\sec(e+fx)}}{\sqrt{2}}\right), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sec(e+fx)}{a+b}} \tan(e+fx)}{(c+d)f \sqrt{a+b \sec(e+fx)} \sqrt{-\tan^2(e+fx)}}$$

output `2*EllipticPi(1/2*(1-sec(f*x+e))^(1/2)*2^(1/2), 2*d/(c+d), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*sec(f*x+e))/(a+b))^(1/2)*tan(f*x+e)/(c+d)/f/(a+b*sec(f*x+e))^(1/2)/(-tan(f*x+e)^2)^(1/2)`

3.281.2 Mathematica [A] (verified)

Time = 8.83 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.83

$$\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{2 \sqrt{\frac{b+a \cos(e+fx)}{(a+b)(1+\cos(e+fx))}} \left((c+d) \operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right), \frac{a-b}{a+b}\right) - 2d \operatorname{EllipticPi}\left(\frac{c-d}{c+d}, \arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right)\right) \right)}{(c-d)(c+d)f \sqrt{\sec^2\left(\frac{1}{2}(e+fx)\right)} \sqrt{a}}$$

input `Integrate[Sec[e + f*x]/(Sqrt[a + b*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

3.281.
$$\int \frac{\sec(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$$

output $(2\sqrt{(b + a\cos(e + fx))/((a + b)(1 + \cos(e + fx)))}) * ((c + d) * \text{EllipticF}[\text{ArcSin}[\text{Tan}[(e + fx)/2]], (a - b)/(a + b)] - 2d * \text{EllipticPi}[(c - d)/(c + d), \text{ArcSin}[\text{Tan}[(e + fx)/2]], (a - b)/(a + b)]) * \sqrt{\cos(e + fx) * \text{Sec}[(e + fx)/2]^2} * \sqrt{\text{Sec}[e + fx]} * \sqrt{1 + \text{Sec}[e + fx]}) / ((c - d)(c + d) * f * \sqrt{\text{Sec}[(e + fx)/2]^2} * \sqrt{a + b * \text{Sec}[e + fx]})$

3.281.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {3042, 4461}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx$$

↓ 3042

$$\int \frac{\csc(e + fx + \frac{\pi}{2})}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(c + d \csc(e + fx + \frac{\pi}{2}))} dx$$

↓ 4461

$$\frac{2 \tan(e + fx) \sqrt{\frac{a + b \sec(e + fx)}{a + b}} \text{EllipticPi}\left(\frac{2d}{c + d}, \arcsin\left(\frac{\sqrt{1 - \sec(e + fx)}}{\sqrt{2}}\right), \frac{2b}{a + b}\right)}{f(c + d) \sqrt{-\tan^2(e + fx)} \sqrt{a + b \sec(e + fx)}}$$

input $\text{Int}[\text{Sec}[e + fx]/(\sqrt{a + b * \text{Sec}[e + fx]} * (c + d * \text{Sec}[e + fx])), x]$

output $(2 * \text{EllipticPi}[(2 * d)/(c + d), \text{ArcSin}[\sqrt{1 - \text{Sec}[e + fx]}/\sqrt{2}], (2 * b)/(a + b)] * \sqrt{(a + b * \text{Sec}[e + fx])/(a + b)} * \text{Tan}[e + fx]) / ((c + d) * f * \sqrt{a + b * \text{Sec}[e + fx]} * \sqrt{-\text{Tan}[e + fx]^2})$

3.281.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4461 `Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))), x_Symbol] := Simp[-2*(Cot[e + f*x]/(f*(c + d)*Sqrt[a + b*Csc[e + f*x]]*Sqrt[-Cot[e + f*x]^2]))*Sqrt[(a + b*Csc[e + f*x])]/(a + b)*EllipticPi[2*(d/(c + d)), ArcSin[Sqrt[1 - Csc[e + f*x]]]/Sqrt[2]], 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.281.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 202 vs. $2(97) = 194$.

Time = 7.64 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.99

method	result
default	$-\frac{2(\cos(fx+e)+1)\left(\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)c+\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)d-2\text{EllipticPi}\left(\cot(fx+e)-\csc(fx+e),\frac{c-d}{c+d}\right)\right)}{f(c-d)(c+d)(b+a\cos(fx+e))}$

input `int(sec(f*x+e)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/f/(c-d)/(c+d)*(cos(f*x+e)+1)*(EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*c+EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*d-2*EllipticPi(cot(f*x+e)-csc(f*x+e),(c-d)/(c+d),((a-b)/(a+b))^(1/2))*d)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(a+b*sec(f*x+e))^(1/2)/(b+a*cos(f*x+e))`

3.281.
$$\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)(c+d\sec(e+fx))}} dx$$

3.281.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

3.281.6 Sympy [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx$$

input `integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)/(sqrt(a + b*sec(e + f*x))*(c + d*sec(e + f*x))), x)`

3.281.7 Maxima [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\sec(fx + e)}{\sqrt{b \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.281.8 Giac [F]

$$\int \frac{\sec(e + fx)}{\sqrt{a + b\sec(e + fx)}(c + d\sec(e + fx))} dx = \int \frac{\sec(fx + e)}{\sqrt{b\sec(fx + e) + a}(d\sec(fx + e) + c)} dx$$

input `integrate(sec(f*x+e)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate(sec(f*x + e)/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.281.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sec(e + fx)}{\sqrt{a + b\sec(e + fx)}(c + d\sec(e + fx))} dx \\ &= \int \frac{1}{\cos(e + fx) \sqrt{a + \frac{b}{\cos(e+fx)}} \left(c + \frac{d}{\cos(e+fx)}\right)} dx \end{aligned}$$

input `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))),x)`

output `int(1/(cos(e + f*x)*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))), x)`

3.282
$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.282.1 Optimal result 2066
 3.282.2 Mathematica [A] (verified) 2067
 3.282.3 Rubi [A] (verified) 2067
 3.282.4 Maple [A] (verified) 2069
 3.282.5 Fracas [F(-1)] 2070
 3.282.6 Sympy [F] 2070
 3.282.7 Maxima [F] 2070
 3.282.8 Giac [F] 2071
 3.282.9 Mupad [F(-1)] 2071

3.282.1 Optimal result

Integrand size = 35, antiderivative size = 209

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$$

$$= \frac{2\sqrt{a+b} \cot(e+fx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(1+\sec(e+fx))}{a-b}}}{bdf}$$

$$- \frac{2c \operatorname{EllipticPi}\left(\frac{2d}{c+d}, \arcsin\left(\frac{\sqrt{1-\sec(e+fx)}}{\sqrt{2}}\right), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sec(e+fx)}{a+b}} \tan(e+fx)}{d(c+d)f \sqrt{a+b \sec(e+fx)} \sqrt{-\tan^2(e+fx)}}$$

```
output 2*cot(f*x+e)*EllipticF((a+b*sec(f*x+e))^(1/2)/(a+b)^(1/2),((a+b)/(a-b))^(1/2))*
(a+b)^(1/2)*(b*(1-sec(f*x+e))/(a+b))^(1/2)*(-b*(1+sec(f*x+e))/(a-b))^(1/2)/
b/d/f-2*c*EllipticPi(1/2*(1-sec(f*x+e))^(1/2)*2^(1/2),2*d/(c+d),2^(1/2)*
(b/(a+b))^(1/2))*((a+b*sec(f*x+e))/(a+b))^(1/2)*tan(f*x+e)/d/(c+d)/f/
(a+b*sec(f*x+e))^(1/2)/(-tan(f*x+e)^2)^(1/2)
```

3.282.2 Mathematica [A] (verified)

Time = 4.29 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.79

$$\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx = \frac{4 \cos^2\left(\frac{1}{2}(e+fx)\right) \sqrt{\frac{\cos(e+fx)}{1+\cos(e+fx)}} \sqrt{\frac{b+a\cos(e+fx)}{(a+b)(1+\cos(e+fx))}} \left((c+d) \operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(e+fx)\right)\right), \frac{a-b}{a+b}\right) - (c-d)(c+d)f\sqrt{a+b\sec(e+fx)}\right)}{(c-d)(c+d)f\sqrt{a+b\sec(e+fx)}}$$

input `Integrate[Sec[e + f*x]^2/(Sqrt[a + b*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(-4*Cos[(e + f*x)/2]^2*Sqrt[Cos[e + f*x]/(1 + Cos[e + f*x])] * Sqrt[(b + a*Cos[e + f*x])/((a + b)*(1 + Cos[e + f*x]))] * ((c + d)*EllipticF[ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)] - 2*c*EllipticPi[(c - d)/(c + d), ArcSin[Tan[(e + f*x)/2]], (a - b)/(a + b)]) * Sec[e + f*x])/((c - d)*(c + d)*f*Sqrt[a + b*Sec[e + f*x]])`

3.282.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4465, 3042, 4319, 4461}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc\left(e+fx+\frac{\pi}{2}\right)^2}{\sqrt{a+b\csc\left(e+fx+\frac{\pi}{2}\right)}(c+d\csc\left(e+fx+\frac{\pi}{2}\right))} dx \\ & \quad \downarrow \text{4465} \\ & \frac{\int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}} dx}{d} - \frac{c \int \frac{\sec(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx}{d} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.282. $\int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx$

$$\frac{\int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})}} dx}{d} - \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d}$$

↓ 4319

$$\frac{2\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{bdf} - \frac{c \int \frac{\csc(e+fx+\frac{\pi}{2})}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d}$$

↓ 4461

$$\frac{2\sqrt{a+b} \cot(e+fx) \sqrt{\frac{b(1-\sec(e+fx))}{a+b}} \sqrt{-\frac{b(\sec(e+fx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \sec(e+fx)}}{\sqrt{a+b}}\right), \frac{a+b}{a-b}\right)}{bdf} - \frac{2c \tan(e+fx) \sqrt{\frac{a+b \sec(e+fx)}{a+b}} \text{EllipticPi}\left(\frac{2d}{c+d}, \arcsin\left(\frac{\sqrt{1-\sec(e+fx)}}{\sqrt{2}}\right), \frac{2b}{a+b}\right)}{df(c+d) \sqrt{-\tan^2(e+fx)} \sqrt{a+b \sec(e+fx)}}$$

input `Int[Sec[e + f*x]^2/(Sqrt[a + b*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(2*Sqrt[a + b]*Cot[e + f*x]*EllipticF[ArcSin[Sqrt[a + b*Sec[e + f*x]]/Sqrt[a + b]], (a + b)/(a - b)]*Sqrt[(b*(1 - Sec[e + f*x]))/(a + b)]*Sqrt[-((b*(1 + Sec[e + f*x]))/(a - b))]/(b*d*f) - (2*c*EllipticPi[(2*d)/(c + d), ArcSin[Sqrt[1 - Sec[e + f*x]]/Sqrt[2]], (2*b)/(a + b)]*Sqrt[(a + b*Sec[e + f*x])/(a + b)]*Tan[e + f*x])/(d*(c + d)*f*Sqrt[a + b*Sec[e + f*x]]*Sqrt[-Tan[e + f*x]^2])`

3.282.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4319 `Int[csc[(e_.) + (f_.)*(x_)]/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)], x_Symbol] := Simp[-2*(Rt[a + b, 2]/(b*f*Cot[e + f*x]))*Sqrt[(b*(1 - Csc[e + f*x]))/(a + b)]*Sqrt[(-b)*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Csc[e + f*x]]/Rt[a + b, 2]], (a + b)/(a - b)], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]`

3.282. $\int \frac{\sec^2(e+fx)}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$

```
rule 4461 Int[csc[(e_.) + (f_.)*(x_)]/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*(csc
c[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))), x_Symbol] := Simp[-2*(Cot[e + f*x]/(f
*(c + d)*Sqrt[a + b*Csc[e + f*x]]*Sqrt[-Cot[e + f*x]^2]))*Sqrt[(a + b*Csc[e
+ f*x])/(a + b)]*EllipticPi[2*(d/(c + d)), ArcSin[Sqrt[1 - Csc[e + f*x]]/S
qrt[2]], 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 4465 Int[csc[(e_.) + (f_.)*(x_)]^2/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*(
csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))), x_Symbol] := Simp[1/d Int[Csc[e +
f*x]/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[c/d Int[Csc[e + f*x]/(Sqrt[
a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.282.4 Maple [A] (verified)

Time = 8.54 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.97

method	result
default	$\frac{2(\cos(fx+e)+1)\left(\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)c+\text{EllipticF}\left(\cot(fx+e)-\csc(fx+e),\sqrt{\frac{a-b}{a+b}}\right)d-2c\text{EllipticPi}\left(\cot(fx+e),\frac{f(c+d)(c-d)(b+a\cos(fx+e))}{f(c+d)(c-d)(b+a\cos(fx+e))}\right)\right)}{f(c+d)(c-d)(b+a\cos(fx+e))}$

```
input int(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output 2/f/(c+d)/(c-d)*(cos(f*x+e)+1)*(EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+
b))^(1/2))*c+EllipticF(cot(f*x+e)-csc(f*x+e),((a-b)/(a+b))^(1/2))*d-2*c*El
lipticPi(cot(f*x+e)-csc(f*x+e),(c-d)/(c+d),((a-b)/(a+b))^(1/2)))*(cos(f*x+
e)/(cos(f*x+e)+1))^(1/2)*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)*(
a+b*sec(f*x+e))^(1/2)/(b+a*cos(f*x+e))
```

$$3.282. \int \frac{\sec^2(e+fx)}{\sqrt{a+b\sec(e+fx)}(c+d\sec(e+fx))} dx$$

3.282.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Timed out}$$

input `integrate(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

3.282.6 Sympy [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx$$

input `integrate(sec(f*x+e)**2/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x)`

output `Integral(sec(e + f*x)**2/(sqrt(a + b*sec(e + f*x))*(c + d*sec(e + f*x))), x)`

3.282.7 Maxima [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\sec^2(fx + e)}{\sqrt{b \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate(sec(f*x + e)^2/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.282.8 Giac [F]

$$\int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\sec^2(fx + e)}{\sqrt{b \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate(sec(f*x+e)^2/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x, algorithm m="giac")`

output `integrate(sec(f*x + e)^2/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.282.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sec^2(e + fx)}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx \\ &= \int \frac{1}{\cos(e + fx)^2 \sqrt{a + \frac{b}{\cos(e + fx)}} \left(c + \frac{d}{\cos(e + fx)} \right)} dx \end{aligned}$$

input `int(1/(cos(e + f*x)^2*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))),x)`

output `int(1/(cos(e + f*x)^2*(a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))), x)`

3.283
$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.283.1 Optimal result 2072
 3.283.2 Mathematica [A] (verified) 2072
 3.283.3 Rubi [A] (verified) 2073
 3.283.4 Maple [C] (verified) 2075
 3.283.5 Fricas [F(-1)] 2075
 3.283.6 Sympy [F] 2076
 3.283.7 Maxima [F] 2076
 3.283.8 Giac [F] 2076
 3.283.9 Mupad [F(-1)] 2077

3.283.1 Optimal result

Integrand size = 39, antiderivative size = 83

$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{2g \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticPi}\left(\frac{2c}{c+d}, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right) \sqrt{g \sec(e+fx)}}{(c+d)f \sqrt{a+b \sec(e+fx)}}$$

output `2*g*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e), 2*c/(c+d), 2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/(c+d)/f/(a+b*sec(f*x+e))^(1/2)`

3.283.2 Mathematica [A] (verified)

Time = 1.38 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00

$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{2g \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticPi}\left(\frac{2c}{c+d}, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right) \sqrt{g \sec(e+fx)}}{(c+d)f \sqrt{a+b \sec(e+fx)}}$$

input `Integrate[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(2*g*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[(2*c)/(c + d), (e + f*x)/2, (2*a)/(a + b)]*Sqrt[g*Sec[e + f*x]]/((c + d)*f*Sqrt[a + b*Sec[e + f*x]])`

3.283.
$$\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$$

3.283.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3042, 4463, 3042, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{3/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(c + d \csc(e + fx + \frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{4463} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{a \cos(e + fx) + b} \int \frac{1}{\sqrt{b + a \cos(e + fx)}(d + c \cos(e + fx))} dx}{\sqrt{a + b \sec(e + fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{a \cos(e + fx) + b} \int \frac{1}{\sqrt{b + a \sin(e + fx + \frac{\pi}{2})}(d + c \sin(e + fx + \frac{\pi}{2}))} dx}{\sqrt{a + b \sec(e + fx)}} \\
 & \quad \downarrow \text{3286} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{\frac{a \cos(e + fx) + b}{a + b}} \int \frac{1}{\sqrt{\frac{b}{a + b} + \frac{a \cos(e + fx)}{a + b}}(d + c \cos(e + fx))} dx}{\sqrt{a + b \sec(e + fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{g \sqrt{g \sec(e + fx)} \sqrt{\frac{a \cos(e + fx) + b}{a + b}} \int \frac{1}{\sqrt{\frac{b}{a + b} + \frac{a \sin(e + fx + \frac{\pi}{2})}{a + b}}(d + c \sin(e + fx + \frac{\pi}{2}))} dx}{\sqrt{a + b \sec(e + fx)}} \\
 & \quad \downarrow \text{3284} \\
 & \frac{2g \sqrt{g \sec(e + fx)} \sqrt{\frac{a \cos(e + fx) + b}{a + b}} \text{EllipticPi}\left(\frac{2c}{c + d}, \frac{1}{2}(e + fx), \frac{2a}{a + b}\right)}{f(c + d) \sqrt{a + b \sec(e + fx)}}
 \end{aligned}$$

3.283. $\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx$

input `Int[(g*Sec[e + f*x])^(3/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `(2*g*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[(2*c)/(c + d), (e + f*x)/2, (2*a)/(a + b)]*Sqrt[g*Sec[e + f*x]])/((c + d)*f*Sqrt[a + b*Sec[e + f*x]])`

3.283.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 4463 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(3/2)/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_.))), x_Symbol] := Simp[g*Sqrt[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

3.283.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 7.46 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.69

method	result
default	$-\frac{2ig\sqrt{a+b\sec(fx+e)}\cos(fx+e)\sqrt{g\sec(fx+e)}\left(2c\operatorname{EllipticPi}\left(i(-\cot(fx+e)+\csc(fx+e)),-\frac{c-d}{c+d},i\sqrt{\frac{a-b}{a+b}}\right)-c\operatorname{EllipticF}\left(i(-\cot(fx+e)+\csc(fx+e)),\frac{c-d}{c+d}\right)\right)}{f(c-d)(c+d)(b+a\cos(fx+e))\sqrt{\frac{a-b}{a+b}}}$

input `int((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `-2*I*g/f/(c-d)/(c+d)*(a+b*sec(f*x+e))^(1/2)*cos(f*x+e)*(g*sec(f*x+e))^(1/2)*(2*c*EllipticPi(I*(-cot(f*x+e)+csc(f*x+e)),-(c-d)/(c+d),I*((a-b)/(a+b))^(1/2))-c*EllipticF(I*(-cot(f*x+e)+csc(f*x+e)),(-(a-b)/(a+b))^(1/2))-d*EllipticF(I*(-cot(f*x+e)+csc(f*x+e)),(-(a-b)/(a+b))^(1/2)))*(1/(a+b)*(b+a*cos(f*x+e))/(cos(f*x+e)+1))^(1/2)/(b+a*cos(f*x+e))/(1/(cos(f*x+e)+1))^(1/2)`

3.283.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Timed out}$$

input `integrate((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="fracas")`

output `Timed out`

3.283.6 Sympy [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx$$

input `integrate((g*sec(f*x+e))**(3/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x)`

output `Integral((g*sec(e + f*x))**(3/2)/(sqrt(a + b*sec(e + f*x))*(c + d*sec(e + f*x))), x)`

3.283.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{3/2}}{\sqrt{b \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="maxima")`

output `integrate((g*sec(f*x + e))^(3/2)/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.283.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{3/2}}{\sqrt{b \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(3/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="giac")`

output `integrate((g*sec(f*x + e))^(3/2)/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.283. $\int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$

3.283.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{3/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{3/2}}{\sqrt{a + \frac{b}{\cos(e+fx)}} \left(c + \frac{d}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(3/2)/((a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))),x)`

output `int((g/cos(e + f*x))^(3/2)/((a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))), x)`

3.284 $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$

3.284.1 Optimal result 2078
 3.284.2 Mathematica [C] (verified) 2078
 3.284.3 Rubi [A] (verified) 2079
 3.284.4 Maple [C] (verified) 2083
 3.284.5 Fricas [F(-1)] 2083
 3.284.6 Sympy [F(-1)] 2084
 3.284.7 Maxima [F] 2084
 3.284.8 Giac [F] 2084
 3.284.9 Mupad [F(-1)] 2085

3.284.1 Optimal result

Integrand size = 39, antiderivative size = 166

$$\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx = \frac{2g^2 \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right) \sqrt{g \sec(e+fx)}}{df \sqrt{a+b \sec(e+fx)}} - \frac{2cg^2 \sqrt{\frac{b+a \cos(e+fx)}{a+b}} \text{EllipticPi}\left(\frac{2c}{c+d}, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right) \sqrt{g \sec(e+fx)}}{d(c+d)f \sqrt{a+b \sec(e+fx)}}$$

output

```
2*g^2*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e),2,2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/d/f/(a+b*sec(f*x+e))^(1/2)-2*c*g^2*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticPi(sin(1/2*f*x+1/2*e),2*c/(c+d),2^(1/2)*(a/(a+b))^(1/2))*((b+a*cos(f*x+e))/(a+b))^(1/2)*(g*sec(f*x+e))^(1/2)/d/(c+d)/f/(a+b*sec(f*x+e))^(1/2)
```

3.284.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

3.284. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$

Time = 25.39 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.48

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \frac{2ig \sqrt{-\frac{a(-1+\cos(e+fx))}{a+b}} \sqrt{\frac{a(1+\cos(e+fx))}{a-b}} \sqrt{b + a \cos(e + fx)} \cot(e + fx) \left((-bc + ad) \operatorname{EllipticPi} \left(1 - \frac{a}{b}, i \operatorname{arcsin} \left(\sqrt{\frac{1}{a-b} bd} \right) \right) \right)}{\dots}$$

input `Integrate[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

output `((-2*I)*g*Sqrt[-((a*(-1 + Cos[e + f*x]))/(a + b))]*Sqrt[(a*(1 + Cos[e + f*x]))/(a - b)]*Sqrt[b + a*Cos[e + f*x]]*Cot[e + f*x]*((-b*c) + a*d)*EllipticPi[1 - a/b, I*ArcSinh[Sqrt[(a - b)^(-1)]*Sqrt[b + a*Cos[e + f*x]]], (-a + b)/(a + b)] + b*c*EllipticPi[((a - b)*c)/(-b*c) + a*d, I*ArcSinh[Sqrt[(a - b)^(-1)]*Sqrt[b + a*Cos[e + f*x]]], (-a + b)/(a + b)]*(g*Sec[e + f*x])^(3/2))/(Sqrt[(a - b)^(-1)]*b*d*(-b*c) + a*d)*f*Sqrt[a + b*Sec[e + f*x]]]`

3.284.3 Rubi [A] (verified)

Time = 2.07 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4467, 3042, 4346, 3042, 3286, 3042, 3284, 4463, 3042, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx \xrightarrow{3042} \int \frac{(g \csc(e + fx + \frac{\pi}{2}))^{5/2}}{\sqrt{a + b \csc(e + fx + \frac{\pi}{2})}(c + d \csc(e + fx + \frac{\pi}{2}))} dx \xrightarrow{4467} \frac{g \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}} dx}{d} - \frac{cg \int \frac{(g \sec(e+fx))^{3/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx}{d}$$

3.284. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$

$$\begin{array}{c}
\downarrow 3042 \\
\frac{g \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})}} dx}{d} - \frac{cg \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d} \\
\downarrow 4346 \\
\frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{\sec(e+fx)}{\sqrt{b+a \cos(e+fx)}} dx}{d \sqrt{a+b \sec(e+fx)}} - \frac{cg \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d} \\
\downarrow 3042 \\
\frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{1}{\sin(e+fx+\frac{\pi}{2}) \sqrt{b+a \sin(e+fx+\frac{\pi}{2})}} dx}{d \sqrt{a+b \sec(e+fx)}} - \frac{cg \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d} \\
\downarrow 3286 \\
\frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{\sec(e+fx)}{\sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b}}} dx}{d \sqrt{a+b \sec(e+fx)}} - \frac{cg \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d} \\
\downarrow 3042 \\
\frac{g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sin(e+fx+\frac{\pi}{2}) \sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b}}} dx}{d \sqrt{a+b \sec(e+fx)}} - \frac{cg \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d} \\
\downarrow 3284 \\
\frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df \sqrt{a+b \sec(e+fx)}} - \frac{cg \int \frac{(g \csc(e+fx+\frac{\pi}{2}))^{3/2}}{\sqrt{a+b \csc(e+fx+\frac{\pi}{2})} (c+d \csc(e+fx+\frac{\pi}{2}))} dx}{d} \\
\downarrow 4463
\end{array}$$

3.284. $\int \frac{(g \sec(e+fx))^{5/2}}{\sqrt{a+b \sec(e+fx)}(c+d \sec(e+fx))} dx$

$$\begin{aligned}
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df \sqrt{a+b \sec(e+fx)}} - \\
& \frac{cg^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{1}{\sqrt{b+a \cos(e+fx)(d+c \cos(e+fx))}} dx}{d \sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df \sqrt{a+b \sec(e+fx)}} - \\
& \frac{cg^2 \sqrt{g \sec(e+fx)} \sqrt{a \cos(e+fx)+b} \int \frac{1}{\sqrt{b+a \sin(e+fx+\frac{\pi}{2})(d+c \sin(e+fx+\frac{\pi}{2}))}} dx}{d \sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{3286} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df \sqrt{a+b \sec(e+fx)}} - \\
& \frac{cg^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a \cos(e+fx)}{a+b} (d+c \cos(e+fx))}} dx}{d \sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df \sqrt{a+b \sec(e+fx)}} - \\
& \frac{cg^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \int \frac{1}{\sqrt{\frac{b}{a+b} + \frac{a \sin(e+fx+\frac{\pi}{2})}{a+b} (d+c \sin(e+fx+\frac{\pi}{2}))}} dx}{d \sqrt{a+b \sec(e+fx)}} \\
& \quad \downarrow \text{3284} \\
& \frac{2g^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df \sqrt{a+b \sec(e+fx)}} - \\
& \frac{2cg^2 \sqrt{g \sec(e+fx)} \sqrt{\frac{a \cos(e+fx)+b}{a+b}} \operatorname{EllipticPi}\left(\frac{2c}{c+d}, \frac{1}{2}(e+fx), \frac{2a}{a+b}\right)}{df(c+d) \sqrt{a+b \sec(e+fx)}}
\end{aligned}$$

input `Int[(g*Sec[e + f*x])^(5/2)/(Sqrt[a + b*Sec[e + f*x]]*(c + d*Sec[e + f*x])),x]`

```
output (2*g^2*Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[2, (e + f*x)/2, (2*a)
/(a + b)]*Sqrt[g*Sec[e + f*x]]/(d*f*Sqrt[a + b*Sec[e + f*x]]) - (2*c*g^2*
Sqrt[(b + a*Cos[e + f*x])/(a + b)]*EllipticPi[(2*c)/(c + d), (e + f*x)/2,
(2*a)/(a + b)]*Sqrt[g*Sec[e + f*x]]/(d*(c + d)*f*Sqrt[a + b*Sec[e + f*x]]
)
```

3.284.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

```
rule 3286 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt
[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c +
d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]
```

```
rule 4346 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.
) + (a_)], x_Symbol] := Simp[d*Sqrt[d*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]
])/Sqrt[a + b*Csc[e + f*x]] Int[1/(Sin[e + f*x]*Sqrt[b + a*Sin[e + f*x]]
), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 4463 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(3/2)/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(b_.
) + (a_)])*(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_)), x_Symbol] := Simp[g*Sqr
t[g*Csc[e + f*x]]*(Sqrt[b + a*Sin[e + f*x]]/Sqrt[a + b*Csc[e + f*x]]) Int
[1/(Sqrt[b + a*Sin[e + f*x]]*(d + c*Sin[e + f*x])), x], x] /; FreeQ[{a, b,
c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
```

```
rule 4467 Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(5/2)/(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)]*(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))), x_Symbol] := Simp[g/d
  Int[(g*Csc[e + f*x])^(3/2)/Sqrt[a + b*Csc[e + f*x]], x], x] - Simp[c*(g/d
  Int[(g*Csc[e + f*x])^(3/2)/(Sqrt[a + b*Csc[e + f*x]]*(c + d*Csc[e + f*x]
  )), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b*c - a*d, 0] && NeQ[a
  ^2 - b^2, 0]
```

3.284.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 9.42 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.89

method	result
default	$\frac{2i \left(\text{EllipticF} \left(i(\cot(fx+e) - \csc(fx+e)), \sqrt{-\frac{a-b}{a+b}} \right) dc + \text{EllipticF} \left(i(\cot(fx+e) - \csc(fx+e)), \sqrt{-\frac{a-b}{a+b}} \right) d^2 + 2c^2 \text{EllipticPi} \left(i(\cot(fx+e) \right. \right. \right.$

```
input int((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,method=
_RETURNVERBOSE)
```

```
output 2*I/f/d/(c+d)/(c-d)*(EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^(1
/2))*d*c+EllipticF(I*(cot(f*x+e)-csc(f*x+e)),(-a-b)/(a+b))^(1/2))*d^2+2*c
^2*EllipticPi(I*(cot(f*x+e)-csc(f*x+e)), -1, I*((a-b)/(a+b))^(1/2))-2*Ellipt
icPi(I*(cot(f*x+e)-csc(f*x+e)), -1, I*((a-b)/(a+b))^(1/2))*d^2-2*c^2*Ellipti
cPi(I*(cot(f*x+e)-csc(f*x+e)), -(c-d)/(c+d), I*((a-b)/(a+b))^(1/2))*1/(a+b
)*(b+a*cos(f*x+e))/(cos(f*x+e)+1)^(1/2)*(a+b*sec(f*x+e))^(1/2)*(g*sec(f*x
+e))^(1/2)*g^2/(1/(cos(f*x+e)+1))^(1/2)/(b+a*cos(f*x+e))*cos(f*x+e)
```

3.284.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Timed out}$$

```
input integrate((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,
algorithm="fricas")
```

```
output Timed out
```

3.284. $\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx$

3.284.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \text{Timed out}$$

input `integrate((g*sec(f*x+e))**(5/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))**(1/2),x)`

output `Timed out`

3.284.7 Maxima [F]

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{5/2}}{\sqrt{b \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="maxima")`

output `integrate((g*sec(f*x + e))^(5/2)/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.284.8 Giac [F]

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{(g \sec(fx + e))^{5/2}}{\sqrt{b \sec(fx + e) + a}(d \sec(fx + e) + c)} dx$$

input `integrate((g*sec(f*x+e))^(5/2)/(c+d*sec(f*x+e))/(a+b*sec(f*x+e))^(1/2),x,algorithm="giac")`

output `integrate((g*sec(f*x + e))^(5/2)/(sqrt(b*sec(f*x + e) + a)*(d*sec(f*x + e) + c)), x)`

3.284.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(g \sec(e + fx))^{5/2}}{\sqrt{a + b \sec(e + fx)}(c + d \sec(e + fx))} dx = \int \frac{\left(\frac{g}{\cos(e+fx)}\right)^{5/2}}{\sqrt{a + \frac{b}{\cos(e+fx)}} \left(c + \frac{d}{\cos(e+fx)}\right)} dx$$

input `int((g/cos(e + f*x))^(5/2)/((a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))),x)`

output `int((g/cos(e + f*x))^(5/2)/((a + b/cos(e + f*x))^(1/2)*(c + d/cos(e + f*x))), x)`

3.285 $\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx$

3.285.1 Optimal result 2086
 3.285.2 Mathematica [B] (verified) 2086
 3.285.3 Rubi [A] (verified) 2087
 3.285.4 Maple [A] (verified) 2089
 3.285.5 Fricas [B] (verification not implemented) 2089
 3.285.6 Sympy [F] 2090
 3.285.7 Maxima [A] (verification not implemented) 2090
 3.285.8 Giac [A] (verification not implemented) 2090
 3.285.9 Mupad [B] (verification not implemented) 2091

3.285.1 Optimal result

Integrand size = 28, antiderivative size = 67

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx = \frac{\cot^5(\frac{1}{2}(e+fx))}{20c^7f} - \frac{\cot^7(\frac{1}{2}(e+fx))}{14c^7f} + \frac{\cot^9(\frac{1}{2}(e+fx))}{36c^7f}$$

output `1/20*cot(1/2*f*x+1/2*e)^5/c^7/f-1/14*cot(1/2*f*x+1/2*e)^7/c^7/f+1/36*cot(1/2*f*x+1/2*e)^9/c^7/f`

3.285.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 151 vs. 2(67) = 134.

Time = 4.03 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.25

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx = \frac{\csc(\frac{e}{2}) \csc^9(\frac{1}{2}(e+fx)) (-971082 \sin(\frac{fx}{2}) - 718830 \sin(e + \frac{fx}{2}) + 467208 \sin(e + \frac{3fx}{2}) + 659400 \sin(2e + \frac{fx}{2}))}{(c-c\sec(e+fx))^7}$$

input `Integrate[(Sec[e + f*x]*Tan[e + f*x]^4)/(c - c*Sec[e + f*x])^7,x]`

output $(\text{Csc}[e/2]*\text{Csc}[(e + f*x)/2]^9*(-971082*\text{Sin}[(f*x)/2] - 718830*\text{Sin}[e + (f*x)/2] + 467208*\text{Sin}[e + (3*f*x)/2] + 659400*\text{Sin}[2*e + (3*f*x)/2] - 303192*\text{Sin}[2*e + (5*f*x)/2] - 179640*\text{Sin}[3*e + (5*f*x)/2] + 30753*\text{Sin}[3*e + (7*f*x)/2] + 89955*\text{Sin}[4*e + (7*f*x)/2] - 13427*\text{Sin}[4*e + (9*f*x)/2] + 15*\text{Sin}[5*e + (9*f*x)/2]))/(23063040*c^7*f)$

3.285.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.88, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {3042, 4902, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^4(e + fx) \sec(e + fx)}{(c - c \sec(e + fx))^7} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(e + fx)^4 \sec(e + fx)}{(c - c \sec(e + fx))^7} dx \\ & \quad \downarrow \text{4902} \\ & \frac{2 \int -\frac{\cot^{10}(\frac{1}{2}(e + fx))(1 - \tan^2(\frac{1}{2}(e + fx)))^2}{8c^7} d \tan(\frac{1}{2}(e + fx))}{f} \\ & \quad \downarrow \text{27} \\ & -\frac{\int \cot^{10}(\frac{1}{2}(e + fx))(1 - \tan^2(\frac{1}{2}(e + fx)))^2 d \tan(\frac{1}{2}(e + fx))}{4c^7 f} \\ & \quad \downarrow \text{244} \\ & -\frac{\int (\cot^{10}(\frac{1}{2}(e + fx)) - 2 \cot^8(\frac{1}{2}(e + fx)) + \cot^6(\frac{1}{2}(e + fx))) d \tan(\frac{1}{2}(e + fx))}{4c^7 f} \\ & \quad \downarrow \text{2009} \\ & -\frac{\frac{1}{9} \cot^9(\frac{1}{2}(e + fx)) + \frac{2}{7} \cot^7(\frac{1}{2}(e + fx)) - \frac{1}{5} \cot^5(\frac{1}{2}(e + fx))}{4c^7 f} \end{aligned}$$

input $\text{Int}[(\text{Sec}[e + f*x]*\text{Tan}[e + f*x]^4)/(c - c*\text{Sec}[e + f*x])^7, x]$

output
$$\frac{-1/4*(-1/5*\cot[(e + f*x)/2]^5 + (2*\cot[(e + f*x)/2]^7)/7 - \cot[(e + f*x)/2]^9/9)}{(c^7*f)}$$

3.285.3.1 Defintions of rubi rules used

rule 27
$$\text{Int}[(a_*)(Fx_), x_Symbol] \text{ :> Simp}[a \text{ Int}[Fx, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_) \text{ /; FreeQ}[b, x]]$$

rule 244
$$\text{Int}[(c_*)(x_)^{(m_)*((a_)+(b_)*(x_)^2)^{(p_)}}, x_Symbol] \text{ :> Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^2)^p, x], x] \text{ /; FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \text{IGtQ}[p, 0]$$

rule 2009
$$\text{Int}[u_, x_Symbol] \text{ :> Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 3042
$$\text{Int}[u_, x_Symbol] \text{ :> Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4902
$$\text{Int}[u_, x_Symbol] \text{ :> With}[\{w = \text{Block}[\{\$ShowSteps = \text{False}, \$StepCounter = \text{Null}\}, \text{Int}[\text{SubstFor}[1/(1 + \text{FreeFactors}[\text{Tan}[\text{FunctionOfTrig}[u, x]/2], x]^2*x^2), \text{Tan}[\text{FunctionOfTrig}[u, x]/2]/\text{FreeFactors}[\text{Tan}[\text{FunctionOfTrig}[u, x]/2], x], u, x], x]\}], \text{Module}[\{v = \text{FunctionOfTrig}[u, x], d\}, \text{Simp}[d = \text{FreeFactors}[\text{Tan}[v/2], x]; 2*(d/\text{Coefficient}[v, x, 1]) \text{ Subst}[\text{Int}[\text{SubstFor}[1/(1 + d^2*x^2), \text{Tan}[v/2]/d, u, x], x], x, \text{Tan}[v/2]/d, x]] \text{ /; CalculusFreeQ}[w, x]] \text{ /; InverseFunctionFreeQ}[u, x] \ \&\& \ !\text{FalseQ}[\text{FunctionOfTrig}[u, x]]$$

3.285.4 Maple [A] (verified)

Time = 11.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.73

method	result
derivativedivides	$\frac{\frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{2}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}}{4f c^7}$
default	$\frac{\frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} - \frac{2}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} + \frac{1}{9 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}}{4f c^7}$
risch	$\frac{2i(315 e^{8i(fx+e)} - 630 e^{7i(fx+e)} + 2310 e^{6i(fx+e)} - 2520 e^{5i(fx+e)} + 3402 e^{4i(fx+e)} - 1638 e^{3i(fx+e)} + 1062 e^{2i(fx+e)} - 108)}{315 f c^7 (e^{i(fx+e)} - 1)^9}$

```
input int(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^7,x,method=_RETURNVERBOSE)
```

```
output 1/4/f/c^7*(1/5/tan(1/2*f*x+1/2*e)^5-2/7/tan(1/2*f*x+1/2*e)^7+1/9/tan(1/2*f*x+1/2*e)^9)
```

3.285.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. 2(58) = 116.

Time = 0.31 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.81

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx$$

$$= \frac{47 \cos^5(fx+e) + 127 \cos^4(fx+e) + 101 \cos^3(fx+e) + 11 \cos^2(fx+e) - 8 \cos(fx+e) + 2}{315 (c^7 f \cos^4(fx+e) - 4 c^7 f \cos^3(fx+e) + 6 c^7 f \cos^2(fx+e) - 4 c^7 f \cos(fx+e) + c^7 f) \sin(fx+e)}$$

```
input integrate(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^7,x, algorithm="fracas")
```

```
output 1/315*(47*cos(f*x + e)^5 + 127*cos(f*x + e)^4 + 101*cos(f*x + e)^3 + 11*cos(f*x + e)^2 - 8*cos(f*x + e) + 2)/((c^7*f*cos(f*x + e)^4 - 4*c^7*f*cos(f*x + e)^3 + 6*c^7*f*cos(f*x + e)^2 - 4*c^7*f*cos(f*x + e) + c^7*f)*sin(f*x + e))
```

3.285.6 Sympy [F]

$$\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx$$

$$= -\frac{\int \frac{\tan^4(e+fx)\sec(e+fx)}{\sec^7(e+fx)-7\sec^6(e+fx)+21\sec^5(e+fx)-35\sec^4(e+fx)+35\sec^3(e+fx)-21\sec^2(e+fx)+7\sec(e+fx)-1} dx}{c^7}$$

input `integrate(sec(f*x+e)*tan(f*x+e)**4/(c-c*sec(f*x+e))**7,x)`

output `-Integral(tan(e + f*x)**4*sec(e + f*x)/(sec(e + f*x)**7 - 7*sec(e + f*x)**6 + 21*sec(e + f*x)**5 - 35*sec(e + f*x)**4 + 35*sec(e + f*x)**3 - 21*sec(e + f*x)**2 + 7*sec(e + f*x) - 1), x)/c**7`

3.285.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.01

$$\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx = -\frac{\left(\frac{90\sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{63\sin^4(fx+e)}{(\cos(fx+e)+1)^4} - 35\right)(\cos(fx+e)+1)^9}{1260c^7f\sin(fx+e)^9}$$

input `integrate(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^7,x, algorithm="maxima")`

output `-1/1260*(90*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 63*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 35)*(cos(f*x + e) + 1)^9/(c^7*f*sin(f*x + e)^9)`

3.285.8 Giac [A] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.70

$$\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx = \frac{63\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^4 - 90\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 + 35}{1260c^7f\tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^9}$$

input `integrate(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^7,x, algorithm="giac")`

output `1/1260*(63*tan(1/2*f*x + 1/2*e)^4 - 90*tan(1/2*f*x + 1/2*e)^2 + 35)/(c^7*f*tan(1/2*f*x + 1/2*e)^9)`

3.285. $\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^7} dx$

3.285.9 Mupad [B] (verification not implemented)

Time = 13.52 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.70

$$\int \frac{\sec(e + fx) \tan^4(e + fx)}{(c - c \sec(e + fx))^7} dx = \frac{63 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4 - 90 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 + 35}{1260 c^7 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^9}$$

input `int(tan(e + f*x)^4/(cos(e + f*x)*(c - c/cos(e + f*x))^7),x)`output `(63*tan(e/2 + (f*x)/2)^4 - 90*tan(e/2 + (f*x)/2)^2 + 35)/(1260*c^7*f*tan(e/2 + (f*x)/2)^9)`

3.286 $\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^8} dx$

3.286.1 Optimal result 2092
 3.286.2 Mathematica [A] (verified) 2092
 3.286.3 Rubi [A] (verified) 2093
 3.286.4 Maple [A] (verified) 2095
 3.286.5 Fricas [A] (verification not implemented) 2095
 3.286.6 Sympy [F] 2096
 3.286.7 Maxima [A] (verification not implemented) 2096
 3.286.8 Giac [A] (verification not implemented) 2097
 3.286.9 Mupad [B] (verification not implemented) 2097

3.286.1 Optimal result

Integrand size = 28, antiderivative size = 89

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^8} dx = \frac{\cot^5\left(\frac{1}{2}(e+fx)\right)}{40c^8f} - \frac{3\cot^7\left(\frac{1}{2}(e+fx)\right)}{56c^8f} + \frac{\cot^9\left(\frac{1}{2}(e+fx)\right)}{24c^8f} - \frac{\cot^{11}\left(\frac{1}{2}(e+fx)\right)}{88c^8f}$$

output `1/40*cot(1/2*f*x+1/2*e)^5/c^8/f-3/56*cot(1/2*f*x+1/2*e)^7/c^8/f+1/24*cot(1/2*f*x+1/2*e)^9/c^8/f-1/88*cot(1/2*f*x+1/2*e)^11/c^8/f`

3.286.2 Mathematica [A] (verified)

Time = 5.02 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.97

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^8} dx = \frac{\csc\left(\frac{e}{2}\right) \csc^{11}\left(\frac{1}{2}(e+fx)\right) \left(425964 \sin\left(\frac{fx}{2}\right) + 486024 \sin\left(e + \frac{fx}{2}\right) - 351450 \sin\left(e + \frac{3fx}{2}\right) - 299970 \sin\left(e + 2fx\right) + 199985 \sin\left(e + \frac{5fx}{2}\right) - 49996 \sin\left(e + 3fx\right) + 4999 \sin\left(e + \frac{7fx}{2}\right) - 499 \sin\left(e + 4fx\right) + 49 \sin\left(e + \frac{9fx}{2}\right) - 4 \sin\left(e + 5fx\right) + \sin\left(e + \frac{11fx}{2}\right)\right)}{88c^8f}$$

input `Integrate[(Sec[e + f*x]*Tan[e + f*x]^4)/(c - c*Sec[e + f*x])^8,x]`

output $-1/15375360*(\text{Csc}[e/2]*\text{Csc}[(e + f*x)/2]^11*(425964*\text{Sin}[(f*x)/2] + 486024*\text{Sin}[e + (f*x)/2] - 351450*\text{Sin}[e + (3*f*x)/2] - 299970*\text{Sin}[2*e + (3*f*x)/2] + 145695*\text{Sin}[2*e + (5*f*x)/2] + 180015*\text{Sin}[3*e + (5*f*x)/2] - 63580*\text{Sin}[3*e + (7*f*x)/2] - 44990*\text{Sin}[4*e + (7*f*x)/2] + 6710*\text{Sin}[4*e + (9*f*x)/2] + 15004*\text{Sin}[5*e + (9*f*x)/2] - 1975*\text{Sin}[5*e + (11*f*x)/2] + \text{Sin}[6*e + (11*f*x)/2]))/(c^8*f)$

3.286.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.84, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {3042, 4902, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^4(e + fx) \sec(e + fx)}{(c - c \sec(e + fx))^8} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(e + fx)^4 \sec(e + fx)}{(c - c \sec(e + fx))^8} dx \\ & \quad \downarrow \text{4902} \\ & \frac{2 \int \frac{\cot^{12}(\frac{1}{2}(e + fx)) (1 - \tan^2(\frac{1}{2}(e + fx)))^3}{16c^8} d \tan(\frac{1}{2}(e + fx))}{f} \\ & \quad \downarrow \text{27} \\ & \frac{\int \cot^{12}(\frac{1}{2}(e + fx)) (1 - \tan^2(\frac{1}{2}(e + fx)))^3 d \tan(\frac{1}{2}(e + fx))}{8c^8 f} \\ & \quad \downarrow \text{244} \\ & \frac{\int (\cot^{12}(\frac{1}{2}(e + fx)) - 3 \cot^{10}(\frac{1}{2}(e + fx)) + 3 \cot^8(\frac{1}{2}(e + fx)) - \cot^6(\frac{1}{2}(e + fx))) d \tan(\frac{1}{2}(e + fx))}{8c^8 f} \\ & \quad \downarrow \text{2009} \\ & \frac{-\frac{1}{11} \cot^{11}(\frac{1}{2}(e + fx)) + \frac{1}{3} \cot^9(\frac{1}{2}(e + fx)) - \frac{3}{7} \cot^7(\frac{1}{2}(e + fx)) + \frac{1}{5} \cot^5(\frac{1}{2}(e + fx))}{8c^8 f} \end{aligned}$$

input $\text{Int}[(\text{Sec}[e + f*x]*\text{Tan}[e + f*x]^4)/(c - c*\text{Sec}[e + f*x])^8, x]$

3.286. $\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c - c \sec(e+fx))^8} dx$

output $(\text{Cot}[(e + f*x)/2]^5/5 - (3*\text{Cot}[(e + f*x)/2]^7)/7 + \text{Cot}[(e + f*x)/2]^9/3 - \text{Cot}[(e + f*x)/2]^{11}/(8*c^8*f))$

3.286.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 244 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4902 `Int[u_, x_Symbol] := With[{w = Block[{$ShowSteps = False, $StepCounter = Null}, Int[SubstFor[1/(1 + FreeFactors[Tan[FunctionOfTrig[u, x]/2], x]^2*x^2), Tan[FunctionOfTrig[u, x]/2]/FreeFactors[Tan[FunctionOfTrig[u, x]/2], x], u, x], x]}], Module[{v = FunctionOfTrig[u, x], d}, Simp[d = FreeFactors[Tan[v/2], x]; 2*(d/Coefficient[v, x, 1]) Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v/2]/d, u, x], x], x, Tan[v/2]/d, x]] /; CalculusFreeQ[w, x]] /; InverseFunctionFreeQ[u, x] && !FalseQ[FunctionOfTrig[u, x]]`

3.286.4 Maple [A] (verified)

Time = 17.28 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.70

method	result
derivativedivides	$-\frac{3}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} - \frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} + \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$
default	$-\frac{3}{7 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7} - \frac{1}{11 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}} + \frac{1}{5 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5} + \frac{1}{3 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}$
risch	$\frac{2i(1155 e^{10i(fx+e)} - 3465 e^{9i(fx+e)} + 13860 e^{8i(fx+e)} - 23100 e^{7i(fx+e)} + 37422 e^{6i(fx+e)} - 32802 e^{5i(fx+e)} + 27060 e^{4i(fx+e)} - 15510 e^{3i(fx+e)} + 5511 e^{2i(fx+e)} - 1155 e^{i(fx+e)} + 1155)}{1155 f c^8 (e^{i(fx+e)} - 1)^{11}}$

```
input int(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^8,x,method=_RETURNVERBOSE)
```

```
output 1/8/f/c^8*(-3/7/tan(1/2*f*x+1/2*e)^7-1/11/tan(1/2*f*x+1/2*e)^11+1/5/tan(1/2*f*x+1/2*e)^5+1/3/tan(1/2*f*x+1/2*e)^9)
```

3.286.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.64

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c-c\sec(e+fx))^8} dx$$

$$= \frac{152 \cos^6(fx+e) + 395 \cos^5(fx+e) + 289 \cos^4(fx+e) + 15 \cos^3(fx+e) - 19 \cos^2(fx+e) + 10 \cos(fx+e) - 2}{1155 (c^8 f \cos^5(fx+e) - 5 c^8 f \cos^4(fx+e) + 10 c^8 f \cos^3(fx+e) - 10 c^8 f \cos^2(fx+e) + 5 c^8 f \cos(fx+e) - c^8 f) \sin(fx+e)}$$

```
input integrate(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^8,x, algorithm="fracas")
```

```
output 1/1155*(152*cos(f*x + e)^6 + 395*cos(f*x + e)^5 + 289*cos(f*x + e)^4 + 15*cos(f*x + e)^3 - 19*cos(f*x + e)^2 + 10*cos(f*x + e) - 2)/((c^8*f*cos(f*x + e)^5 - 5*c^8*f*cos(f*x + e)^4 + 10*c^8*f*cos(f*x + e)^3 - 10*c^8*f*cos(f*x + e)^2 + 5*c^8*f*cos(f*x + e) - c^8*f)*sin(f*x + e))
```


3.286.6 Sympy [F]

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c - c \sec(e+fx))^8} dx$$

$$= \frac{\int \frac{\tan^4(e+fx) \sec(e+fx)}{\sec^8(e+fx) - 8 \sec^7(e+fx) + 28 \sec^6(e+fx) - 56 \sec^5(e+fx) + 70 \sec^4(e+fx) - 56 \sec^3(e+fx) + 28 \sec^2(e+fx) - 8 \sec(e+fx) + 1} dx}{c^8}$$

input `integrate(sec(f*x+e)*tan(f*x+e)**4/(c-c*sec(f*x+e))**8,x)`

output `Integral(tan(e + f*x)**4*sec(e + f*x)/(sec(e + f*x)**8 - 8*sec(e + f*x)**7 + 28*sec(e + f*x)**6 - 56*sec(e + f*x)**5 + 70*sec(e + f*x)**4 - 56*sec(e + f*x)**3 + 28*sec(e + f*x)**2 - 8*sec(e + f*x) + 1), x)/c**8`

3.286.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.99

$$\int \frac{\sec(e+fx) \tan^4(e+fx)}{(c - c \sec(e+fx))^8} dx$$

$$= \frac{\left(\frac{385 \sin^2(fx+e)}{(\cos(fx+e)+1)^2} - \frac{495 \sin^4(fx+e)}{(\cos(fx+e)+1)^4} + \frac{231 \sin^6(fx+e)}{(\cos(fx+e)+1)^6} - 105 \right) (\cos(fx+e) + 1)^{11}}{9240 c^8 f \sin(fx+e)^{11}}$$

input `integrate(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^8,x, algorithm="maxima")`

output `1/9240*(385*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - 495*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 231*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 - 105)*(cos(f*x + e) + 1)^11/(c^8*f*sin(f*x + e)^11)`

3.286.8 Giac [A] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^8} dx$$

$$= \frac{231 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^6 - 495 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^4 + 385 \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^2 - 105}{9240 c^8 f \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^{11}}$$

input `integrate(sec(f*x+e)*tan(f*x+e)^4/(c-c*sec(f*x+e))^8,x, algorithm="giac")`output `1/9240*(231*tan(1/2*f*x + 1/2*e)^6 - 495*tan(1/2*f*x + 1/2*e)^4 + 385*tan(1/2*f*x + 1/2*e)^2 - 105)/(c^8*f*tan(1/2*f*x + 1/2*e)^11)`**3.286.9 Mupad [B] (verification not implemented)**

Time = 14.18 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.67

$$\int \frac{\sec(e+fx)\tan^4(e+fx)}{(c-c\sec(e+fx))^8} dx = \frac{\frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^6}{5} - \frac{3 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^4}{7} + \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2}{3} - \frac{1}{11}}{8 c^8 f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)^{11}}$$

input `int(tan(e + f*x)^4/(cos(e + f*x)*(c - c/cos(e + f*x))^8),x)`output `(tan(e/2 + (f*x)/2)^2/3 - (3*tan(e/2 + (f*x)/2)^4)/7 + tan(e/2 + (f*x)/2)^6/5 - 1/11)/(8*c^8*f*tan(e/2 + (f*x)/2)^11)`

APPENDIX

4.1 Listing of Grading functions	2098
--	------

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(" ,
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```



```
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if
```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

```



```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```