

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.5-Secant/124-4.5.4.1-a+b-sec^m-A+B-sec+C-
sec²-

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 1:40am

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	45
4	Appendix	465

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	15
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [70]. This is test number [124].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (70)	0.00 (0)
Mathematica	100.00 (70)	0.00 (0)
Maple	100.00 (70)	0.00 (0)
Fricas	100.00 (70)	0.00 (0)
Mupad	70.00 (49)	30.00 (21)
Maxima	68.57 (48)	31.43 (22)
Giac	65.71 (46)	34.29 (24)
Sympy	4.29 (3)	95.71 (67)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	78.571	2.857	18.571	0.000
Maple	65.714	1.429	32.857	0.000
Fricas	64.286	5.714	30.000	0.000
Maxima	61.429	7.143	0.000	31.429
Giac	35.714	30.000	0.000	34.286
Sympy	4.286	0.000	0.000	95.714
Mupad	0.000	70.000	0.000	30.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	0	0.00	0.00	0.00
Mupad	21	0.00	100.00	0.00
Maxima	22	100.00	0.00	0.00
Giac	24	100.00	0.00	0.00
Sympy	67	92.54	7.46	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.21
Maxima	0.23
Giac	0.29
Rubi	0.39
Mathematica	0.48
Maple	1.44
Sympy	2.07
Mupad	13.21

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Sympy	38.00	1.93	49.00	1.65
Mupad	60.37	1.37	42.00	1.10
Maxima	68.92	1.90	44.50	1.18
Rubi	69.96	1.00	61.50	1.00
Mathematica	75.79	1.27	60.50	1.00
Fricas	80.67	1.21	69.50	1.16
Giac	84.67	1.66	60.00	1.62
Maple	213.20	2.38	59.00	1.33

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v0.6

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	24
2.3	Detailed conclusion table specific for Rubi results	42

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	21
2.1.3	Maple	22
2.1.4	Fricas	22
2.1.5	Maxima	22
2.1.6	Giac	23
2.1.7	Mupad	23
2.1.8	Sympy	23

2.1.1 Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 16, 18, 20, 22, 24, 26, 27, 28, 29, 30, 31, 33, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64 }

B grade { 32, 34 }

C grade { 14, 15, 17, 19, 21, 23, 25, 65, 66, 67, 68, 69, 70 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.3 Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 25, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64 }

B grade { 24 }

C grade { 16, 17, 18, 19, 20, 21, 22, 23, 26, 27, 47, 48, 49, 50, 51, 52, 53, 65, 66, 67, 68, 69, 70 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.4 Fricas

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 42, 43, 44, 45, 46, 54, 55, 56, 57, 59, 60, 61, 62, 63, 64 }

B grade { 7, 40, 41, 58 }

C grade { 16, 17, 18, 19, 20, 21, 22, 23, 47, 48, 49, 50, 51, 52, 53, 65, 66, 67, 68, 69, 70 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.5 Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64 }

B grade { 14, 15, 25, 26, 41 }

C grade { }

F normal fail { 16, 17, 18, 19, 20, 21, 22, 23, 24, 47, 48, 49, 50, 51, 52, 53, 65, 66, 67, 68, 69, 70 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.6 Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 46 }

B grade { 24, 37, 38, 39, 40, 41, 42, 43, 44, 45, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64 }

C grade { }

F normal fail { 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 47, 48, 49, 50, 51, 52, 53, 65, 66, 67, 68, 69, 70 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64 }

C grade { }

F normal fail { }

F(-1) timeout fail { 16, 17, 18, 19, 20, 21, 22, 23, 47, 48, 49, 50, 51, 52, 53, 65, 66, 67, 68, 69, 70 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { 9, 32, 33 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 47, 48, 49, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70 }

F(-1) timeout fail { 13, 23, 46, 53, 64 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	70	81	78	60	74	0	79	56
N.S.	1	0.80	0.93	0.90	0.69	0.85	0.00	0.91	0.64
time (sec)	N/A	0.302	0.259	4.590	0.214	0.291	0.000	0.289	15.170

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	94	137	108	126	114	0	121	102
N.S.	1	0.96	1.40	1.10	1.29	1.16	0.00	1.23	1.04
time (sec)	N/A	0.449	0.022	0.490	0.231	0.304	0.000	0.295	0.238

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	58	61	58	43	56	0	57	42
N.S.	1	0.89	0.94	0.89	0.66	0.86	0.00	0.88	0.65
time (sec)	N/A	0.306	0.161	0.245	0.278	0.268	0.000	0.306	14.966

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	68	93	85	97	95	0	98	77
N.S.	1	0.97	1.33	1.21	1.39	1.36	0.00	1.40	1.10
time (sec)	N/A	0.357	0.009	0.312	0.227	0.291	0.000	0.306	15.548

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	36	35	34	37	0	34	28
N.S.	1	1.00	0.84	0.81	0.79	0.86	0.00	0.79	0.65
time (sec)	N/A	0.274	0.068	0.191	0.205	0.289	0.000	0.300	15.745

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	48	55	58	72	0	60	41
N.S.	1	1.00	1.20	1.38	1.45	1.80	0.00	1.50	1.02
time (sec)	N/A	0.254	0.009	0.118	0.208	0.267	0.000	0.299	16.080

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	16	15	31	0	15	17
N.S.	1	1.00	1.00	1.07	1.00	2.07	0.00	1.00	1.13
time (sec)	N/A	0.149	0.002	0.045	0.192	0.246	0.000	0.268	16.025

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	35	30	38	40	0	40	22
N.S.	1	1.00	1.46	1.25	1.58	1.67	0.00	1.67	0.92
time (sec)	N/A	0.242	0.013	0.099	0.200	0.252	0.000	0.270	0.056

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	33	24	37	28	51	37	25
N.S.	1	1.00	1.06	0.77	1.19	0.90	1.65	1.19	0.81
time (sec)	N/A	0.203	0.027	0.080	0.286	0.248	2.523	0.274	15.911

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	50	31	27	28	0	34	28
N.S.	1	1.00	1.67	1.03	0.90	0.93	0.00	1.13	0.93
time (sec)	N/A	0.270	0.009	0.154	0.197	0.254	0.000	0.273	0.046

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	59	45	44	73	49	0	73	67
N.S.	1	0.97	0.74	0.72	1.20	0.80	0.00	1.20	1.10
time (sec)	N/A	0.272	0.075	0.219	0.290	0.272	0.000	0.279	16.433

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	47	71	49	43	45	0	57	43
N.S.	1	0.94	1.42	0.98	0.86	0.90	0.00	1.14	0.86
time (sec)	N/A	0.296	0.008	0.228	0.197	0.265	0.000	0.276	15.990

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	85	68	61	103	68	0	96	91
N.S.	1	0.96	0.76	0.69	1.16	0.76	0.00	1.08	1.02
time (sec)	N/A	0.341	0.079	0.256	0.279	0.252	0.000	0.280	16.137

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	113	46	304	33	0	0	42
N.S.	1	1.00	4.35	1.77	11.69	1.27	0.00	0.00	1.62
time (sec)	N/A	0.213	0.442	0.724	0.429	0.273	0.000	0.000	15.610

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	111	44	296	33	0	0	41
N.S.	1	1.00	4.44	1.76	11.84	1.32	0.00	0.00	1.64
time (sec)	N/A	0.218	0.299	0.340	0.428	0.257	0.000	0.000	15.212

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	108	84	291	0	142	0	0	0
N.S.	1	0.98	0.76	2.65	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	0.468	1.113	22.693	0.000	0.099	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	104	184	798	0	143	0	0	0
N.S.	1	0.95	1.67	7.25	0.00	1.30	0.00	0.00	0.00
time (sec)	N/A	0.464	1.201	5.330	0.000	0.098	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	58	220	0	111	0	0	0
N.S.	1	1.00	0.81	3.06	0.00	1.54	0.00	0.00	0.00
time (sec)	N/A	0.360	0.458	2.141	0.000	0.094	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	126	763	0	99	0	0	0
N.S.	1	1.00	1.85	11.22	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.355	0.795	3.781	0.000	0.088	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	66	218	0	100	0	0	0
N.S.	1	1.00	0.88	2.91	0.00	1.33	0.00	0.00	0.00
time (sec)	N/A	0.361	0.443	2.913	0.000	0.095	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	77	133	798	0	108	0	0	0
N.S.	1	1.00	1.73	10.36	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	0.362	1.218	5.274	0.000	0.091	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	113	79	291	0	119	0	0	0
N.S.	1	1.01	0.71	2.60	0.00	1.06	0.00	0.00	0.00
time (sec)	N/A	0.484	0.925	4.658	0.000	0.092	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	113	143	866	0	129	0	0	0
N.S.	1	1.01	1.28	7.73	0.00	1.15	0.00	0.00	0.00
time (sec)	N/A	0.488	1.325	7.760	0.000	0.101	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	41	0	19	0	46	21
N.S.	1	1.00	1.00	1.95	0.00	0.90	0.00	2.19	1.00
time (sec)	N/A	0.193	0.217	1.561	0.000	0.247	0.000	0.323	0.215

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	107	40	283	29	0	0	37
N.S.	1	1.00	5.10	1.90	13.48	1.38	0.00	0.00	1.76
time (sec)	N/A	0.204	0.386	1.148	0.428	0.249	0.000	0.000	15.392

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	41	42	19	0	22	42
N.S.	1	1.00	1.00	2.16	2.21	1.00	0.00	1.16	2.21
time (sec)	N/A	0.190	0.031	0.319	0.195	0.235	0.000	0.312	0.126

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	41	30	19	0	30	19
N.S.	1	1.00	1.00	2.16	1.58	1.00	0.00	1.58	1.00
time (sec)	N/A	0.192	0.035	0.240	0.203	0.251	0.000	0.296	15.172

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	36	33	19	0	23	23
N.S.	1	1.00	1.00	1.89	1.74	1.00	0.00	1.21	1.21
time (sec)	N/A	0.186	0.028	0.194	0.199	0.245	0.000	0.299	0.059

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	32	20	19	0	20	21
N.S.	1	1.00	1.00	1.68	1.05	1.00	0.00	1.05	1.11
time (sec)	N/A	0.189	0.032	0.138	0.202	0.241	0.000	0.285	15.139

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	18	22	19	0	24	22
N.S.	1	1.00	1.00	1.06	1.29	1.12	0.00	1.41	1.29
time (sec)	N/A	0.182	0.015	0.109	0.194	0.232	0.000	0.290	15.104

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	19	0	11	11
N.S.	1	1.00	1.00	1.09	1.00	1.73	0.00	1.00	1.00
time (sec)	N/A	0.170	0.002	0.062	0.197	0.236	0.000	0.269	15.275

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	23	12	11	11	14	11	11
N.S.	1	1.00	2.09	1.09	1.00	1.00	1.27	1.00	1.00
time (sec)	N/A	0.154	0.014	0.078	0.199	0.251	0.059	0.266	0.022

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	33	15	23	17	49	14	14
N.S.	1	1.00	1.94	0.88	1.35	1.00	2.88	0.82	0.82
time (sec)	N/A	0.188	0.008	0.122	0.207	0.249	3.627	0.269	15.225

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	46	26	21	19	0	21	22
N.S.	1	1.00	2.42	1.37	1.11	1.00	0.00	1.11	1.16
time (sec)	N/A	0.191	0.035	0.150	0.232	0.250	0.000	0.276	0.046

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	31	29	33	19	0	23	19
N.S.	1	1.00	1.63	1.53	1.74	1.00	0.00	1.21	1.00
time (sec)	N/A	0.189	0.086	0.165	0.191	0.261	0.000	0.282	15.435

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	38	37	30	19	0	30	23
N.S.	1	1.00	2.00	1.95	1.58	1.00	0.00	1.58	1.21
time (sec)	N/A	0.196	0.030	0.204	0.207	0.251	0.000	0.287	15.531

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	90	59	73	95	99	0	164	153
N.S.	1	1.06	0.69	0.86	1.12	1.16	0.00	1.93	1.80
time (sec)	N/A	0.487	0.175	0.345	0.201	0.268	0.000	0.314	18.027

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	64	60	60	70	88	0	122	109
N.S.	1	1.02	0.95	0.95	1.11	1.40	0.00	1.94	1.73
time (sec)	N/A	0.404	0.121	0.256	0.208	0.275	0.000	0.298	17.120

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	48	47	47	58	74	0	107	85
N.S.	1	1.02	1.00	1.00	1.23	1.57	0.00	2.28	1.81
time (sec)	N/A	0.360	0.009	0.182	0.198	0.270	0.000	0.326	15.824

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	30	31	60	0	57	47
N.S.	1	1.00	1.00	1.25	1.29	2.50	0.00	2.38	1.96
time (sec)	N/A	0.153	0.007	0.100	0.194	0.264	0.000	0.278	15.308

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	29	37	36	0	43	57
N.S.	1	1.00	1.00	1.81	2.31	2.25	0.00	2.69	3.56
time (sec)	N/A	0.235	0.002	0.115	0.196	0.260	0.000	0.284	15.211

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	26	16	20	17	0	39	17
N.S.	1	1.00	1.73	1.07	1.33	1.13	0.00	2.60	1.13
time (sec)	N/A	0.228	0.004	0.121	0.195	0.260	0.000	0.285	15.416

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	39	35	32	34	29	0	82	31
N.S.	1	1.03	0.92	0.84	0.89	0.76	0.00	2.16	0.82
time (sec)	N/A	0.281	0.049	0.150	0.197	0.250	0.000	0.290	15.318

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	55	57	44	46	42	0	98	55
N.S.	1	1.02	1.06	0.81	0.85	0.78	0.00	1.81	1.02
time (sec)	N/A	0.310	0.046	0.184	0.214	0.256	0.000	0.292	15.615

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	81	73	57	57	53	0	140	75
N.S.	1	1.07	0.96	0.75	0.75	0.70	0.00	1.84	0.99
time (sec)	N/A	0.381	0.087	0.227	0.205	0.259	0.000	0.293	15.253

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	93	89	69	69	64	0	154	113
N.S.	1	1.01	0.97	0.75	0.75	0.70	0.00	1.67	1.23
time (sec)	N/A	0.391	0.076	0.275	0.216	0.255	0.000	0.287	19.245

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	181	102	555	0	207	0	0	0
N.S.	1	1.07	0.60	3.28	0.00	1.22	0.00	0.00	0.00
time (sec)	N/A	0.756	0.821	1.923	0.000	0.099	0.000	0.000	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	147	90	534	0	183	0	0	0
N.S.	1	1.09	0.67	3.96	0.00	1.36	0.00	0.00	0.00
time (sec)	N/A	0.593	0.527	2.878	0.000	0.094	0.000	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	114	73	460	0	143	0	0	0
N.S.	1	1.05	0.67	4.22	0.00	1.31	0.00	0.00	0.00
time (sec)	N/A	0.541	0.792	1.839	0.000	0.096	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	57	414	0	122	0	0	0
N.S.	1	1.00	0.67	4.87	0.00	1.44	0.00	0.00	0.00
time (sec)	N/A	0.435	0.350	2.864	0.000	0.094	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	120	81	536	0	150	0	0	0
N.S.	1	1.03	0.70	4.62	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	0.550	0.511	1.727	0.000	0.100	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	155	91	566	0	164	0	0	0
N.S.	1	1.05	0.62	3.85	0.00	1.12	0.00	0.00	0.00
time (sec)	N/A	0.606	1.237	2.811	0.000	0.105	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	176	191	104	582	0	175	0	0	0
N.S.	1	1.09	0.59	3.31	0.00	0.99	0.00	0.00	0.00
time (sec)	N/A	0.775	1.324	1.725	0.000	0.108	0.000	0.000	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	120	80	104	127	122	0	246	197
N.S.	1	0.98	0.66	0.85	1.04	1.00	0.00	2.02	1.61
time (sec)	N/A	0.671	0.485	0.474	0.215	0.271	0.000	0.309	17.674

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	95	71	106	139	117	0	230	160
N.S.	1	0.98	0.73	1.09	1.43	1.21	0.00	2.37	1.65
time (sec)	N/A	0.564	0.198	0.394	0.219	0.279	0.000	0.306	17.938

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	79	51	68	79	94	0	162	123
N.S.	1	1.01	0.65	0.87	1.01	1.21	0.00	2.08	1.58
time (sec)	N/A	0.538	0.165	0.261	0.205	0.265	0.000	0.299	17.097

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	59	63	75	82	0	115	89
N.S.	1	1.00	1.16	1.24	1.47	1.61	0.00	2.25	1.75
time (sec)	N/A	0.394	0.005	0.206	0.206	0.268	0.000	0.314	15.737

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	27	35	34	71	0	60	161
N.S.	1	1.00	1.00	1.30	1.26	2.63	0.00	2.22	5.96
time (sec)	N/A	0.161	0.001	0.026	0.205	0.264	0.000	0.294	14.946

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	38	37	46	45	0	70	68
N.S.	1	1.00	1.41	1.37	1.70	1.67	0.00	2.59	2.52
time (sec)	N/A	0.329	0.014	0.157	0.209	0.270	0.000	0.306	14.916

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	55	35	42	33	0	86	34
N.S.	1	1.00	1.31	0.83	1.00	0.79	0.00	2.05	0.81
time (sec)	N/A	0.326	0.037	0.151	0.195	0.246	0.000	0.288	14.848

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	58	53	49	55	45	0	138	66
N.S.	1	1.04	0.95	0.88	0.98	0.80	0.00	2.46	1.18
time (sec)	N/A	0.430	0.077	0.184	0.199	0.272	0.000	0.297	14.711

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	86	70	63	77	65	0	200	81
N.S.	1	0.98	0.80	0.72	0.88	0.74	0.00	2.27	0.92
time (sec)	N/A	0.441	0.124	0.239	0.206	0.254	0.000	0.286	15.058

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	101	87	79	89	73	0	222	104
N.S.	1	1.03	0.89	0.81	0.91	0.74	0.00	2.27	1.06
time (sec)	N/A	0.571	0.128	0.282	0.196	0.272	0.000	0.308	14.919

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	124	102	95	115	93	0	284	126
N.S.	1	0.94	0.77	0.72	0.87	0.70	0.00	2.15	0.95
time (sec)	N/A	0.533	0.217	0.342	0.219	0.279	0.000	0.297	15.013

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	178	179	308	949	0	225	0	0	0
N.S.	1	1.01	1.73	5.33	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.974	5.708	2.950	0.000	0.105	0.000	0.000	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	143	302	611	0	193	0	0	0
N.S.	1	1.05	2.22	4.49	0.00	1.42	0.00	0.00	0.00
time (sec)	N/A	0.822	2.201	1.528	0.000	0.093	0.000	0.000	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	156	847	0	153	0	0	0
N.S.	1	1.00	1.42	7.70	0.00	1.39	0.00	0.00	0.00
time (sec)	N/A	0.676	1.394	2.734	0.000	0.094	0.000	0.000	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	143	608	0	160	0	0	0
N.S.	1	1.00	1.22	5.20	0.00	1.37	0.00	0.00	0.00
time (sec)	N/A	0.682	1.721	1.594	0.000	0.098	0.000	0.000	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	154	169	956	0	176	0	0	0
N.S.	1	1.03	1.13	6.37	0.00	1.17	0.00	0.00	0.00
time (sec)	N/A	0.839	2.198	2.692	0.000	0.102	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	190	177	726	0	193	0	0	0
N.S.	1	1.03	0.96	3.92	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.972	2.600	1.543	0.000	0.106	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [65] had the largest ratio of [.454544999999999977]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	5	0.80	21	0.238
2	A	8	8	0.96	21	0.381
3	A	6	5	0.89	21	0.238
4	A	6	6	0.97	21	0.286
5	A	6	5	1.00	21	0.238
6	A	4	4	1.00	19	0.211
7	A	1	1	1.00	12	0.083
8	A	4	4	1.00	19	0.211
9	A	3	3	1.00	21	0.143
10	A	6	5	1.00	21	0.238
11	A	5	5	0.97	21	0.238
12	A	7	6	0.94	21	0.286
13	A	7	7	0.96	21	0.333
14	A	2	2	1.00	29	0.069
15	A	2	2	1.00	28	0.071
16	A	8	8	0.98	25	0.320
17	A	8	8	0.95	25	0.320
18	A	6	6	1.00	25	0.240
19	A	6	6	1.00	25	0.240
20	A	6	6	1.00	25	0.240
21	A	6	6	1.00	25	0.240
22	A	8	8	1.01	25	0.320

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	8	8	1.01	25	0.320
24	A	2	2	1.00	23	0.087
25	A	2	2	1.00	24	0.083
26	A	2	2	1.00	21	0.095
27	A	2	2	1.00	21	0.095
28	A	2	2	1.00	21	0.095
29	A	2	2	1.00	21	0.095
30	A	2	2	1.00	19	0.105
31	A	5	4	1.00	10	0.400
32	A	3	3	1.00	8	0.375
33	A	2	2	1.00	19	0.105
34	A	2	2	1.00	21	0.095
35	A	2	2	1.00	21	0.095
36	A	2	2	1.00	21	0.095
37	A	12	11	1.06	28	0.393
38	A	10	9	1.02	28	0.321
39	A	10	9	1.02	26	0.346
40	A	1	1	1.00	19	0.053
41	A	6	6	1.00	26	0.231
42	A	5	5	1.00	28	0.179
43	A	7	7	1.03	28	0.250
44	A	9	8	1.02	28	0.286
45	A	11	10	1.07	28	0.357
46	A	11	10	1.01	28	0.357
47	A	13	13	1.07	32	0.406
48	A	11	11	1.09	32	0.344
49	A	11	11	1.05	32	0.344
50	A	9	9	1.00	32	0.281
51	A	11	11	1.03	32	0.344
52	A	11	11	1.05	32	0.344
53	A	13	13	1.09	32	0.406
54	A	13	12	0.98	29	0.414
55	A	11	10	0.98	29	0.345
56	A	11	10	1.01	29	0.345

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	9	8	1.00	27	0.296
58	A	1	1	1.00	20	0.050
59	A	7	7	1.00	27	0.259
60	A	6	6	1.00	29	0.207
61	A	10	9	1.04	29	0.310
62	A	10	9	0.98	29	0.310
63	A	13	12	1.03	29	0.414
64	A	12	11	0.94	29	0.379
65	A	15	15	1.01	33	0.455
66	A	13	13	1.05	33	0.394
67	A	11	11	1.00	33	0.333
68	A	11	11	1.00	33	0.333
69	A	13	13	1.03	33	0.394
70	A	15	15	1.03	33	0.455

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx$	48
3.2	$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx$	53
3.3	$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx$	59
3.4	$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx$	64
3.5	$\int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx$	70
3.6	$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx$	75
3.7	$\int (A + C \sec^2(c + dx)) dx$	80
3.8	$\int \cos(c + dx) (A + C \sec^2(c + dx)) dx$	84
3.9	$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx$	89
3.10	$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx$	94
3.11	$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx$	99
3.12	$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx$	104
3.13	$\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx$	110
3.14	$\int \sec^m(c + dx) \left(-\frac{Cm}{1+m} + C \sec^2(c + dx)\right) dx$	116
3.15	$\int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m}\right) dx$	121
3.16	$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx$	126
3.17	$\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx$	132
3.18	$\int \sqrt{b \sec(c + dx)} (A + C \sec^2(c + dx)) dx$	139
3.19	$\int \frac{A+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	144
3.20	$\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	150
3.21	$\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	155
3.22	$\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$	161
3.23	$\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx$	167
3.24	$\int \frac{3+3 \sec^2(c+dx)}{\sqrt{\sec(c+dx)}} dx$	173
3.25	$\int \sec^m(e + fx) (m - (1 + m) \sec^2(e + fx)) dx$	177
3.26	$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx$	182

3.27	$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx$	187
3.28	$\int \sec^3(e + fx) (3 - 4 \sec^2(e + fx)) dx$	192
3.29	$\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx$	197
3.30	$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx$	202
3.31	$\int -\sec^2(e + fx) dx$	207
3.32	$\int -\cos(e + fx) dx$	212
3.33	$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx$	217
3.34	$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx$	221
3.35	$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx$	226
3.36	$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx$	231
3.37	$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	236
3.38	$\int \sec^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	243
3.39	$\int \sec(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	249
3.40	$\int (B \sec(c + dx) + C \sec^2(c + dx)) dx$	255
3.41	$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	260
3.42	$\int \cos^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	265
3.43	$\int \cos^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	270
3.44	$\int \cos^4(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	276
3.45	$\int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	282
3.46	$\int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$	289
3.47	$\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx$	296
3.48	$\int \sqrt{b \sec(c + dx)} (B \sec(c + dx) + C \sec^2(c + dx)) dx$	304
3.49	$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	312
3.50	$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	319
3.51	$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	325
3.52	$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$	332
3.53	$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx$	339
3.54	$\int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	347
3.55	$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	355
3.56	$\int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	362
3.57	$\int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	369
3.58	$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	375
3.59	$\int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	380
3.60	$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	386
3.61	$\int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	391
3.62	$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	397
3.63	$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	404
3.64	$\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	411
3.65	$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$	418

3.66	$\int \sqrt{b \sec(c+dx)}(A + B \sec(c+dx) + C \sec^2(c+dx)) dx$	427
3.67	$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	435
3.68	$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	442
3.69	$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	449
3.70	$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$	457

3.1 $\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx$

3.1.1	Optimal result	48
3.1.2	Mathematica [A] (verified)	48
3.1.3	Rubi [A] (verified)	49
3.1.4	Maple [A] (verified)	50
3.1.5	Fricas [A] (verification not implemented)	51
3.1.6	Sympy [F]	51
3.1.7	Maxima [A] (verification not implemented)	51
3.1.8	Giac [A] (verification not implemented)	52
3.1.9	Mupad [B] (verification not implemented)	52

3.1.1 Optimal result

Integrand size = 21, antiderivative size = 87

$$\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(7A + 6C) \tan(c + dx)}{7d} + \frac{C \sec^6(c + dx) \tan(c + dx)}{7d} + \frac{2(7A + 6C) \tan^3(c + dx)}{21d} + \frac{(7A + 6C) \tan^5(c + dx)}{35d}$$

output `1/7*(7*A+6*C)*tan(d*x+c)/d+1/7*C*sec(d*x+c)^6*tan(d*x+c)/d+2/21*(7*A+6*C)*tan(d*x+c)^3/d+1/35*(7*A+6*C)*tan(d*x+c)^5/d`

3.1.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.93

$$\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx = \frac{A(\tan(c + dx) + \frac{2}{3} \tan^3(c + dx) + \frac{1}{5} \tan^5(c + dx))}{d} + \frac{C(\tan(c + dx) + \tan^3(c + dx) + \frac{3}{5} \tan^5(c + dx) + \frac{1}{7} \tan^7(c + dx))}{d}$$

input `Integrate[Sec[c + d*x]^6*(A + C*Sec[c + d*x]^2),x]`

output `(A*(Tan[c + d*x] + (2*Tan[c + d*x]^3)/3 + Tan[c + d*x]^5/5))/d + (C*(Tan[c + d*x] + Tan[c + d*x]^3 + (3*Tan[c + d*x]^5)/5 + Tan[c + d*x]^7/7))/d`

3.1.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 4534, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^6 \left(A + C \csc\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{7}(7A + 6C) \int \sec^6(c + dx) dx + \frac{C \tan(c + dx) \sec^6(c + dx)}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{7}(7A + 6C) \int \csc\left(c + dx + \frac{\pi}{2}\right)^6 dx + \frac{C \tan(c + dx) \sec^6(c + dx)}{7d} \\
 & \quad \downarrow \text{4254} \\
 & \frac{C \tan(c + dx) \sec^6(c + dx)}{7d} - \frac{(7A + 6C) \int (\tan^4(c + dx) + 2 \tan^2(c + dx) + 1) d(-\tan(c + dx))}{7d} \\
 & \quad \downarrow \text{2009} \\
 & \frac{C \tan(c + dx) \sec^6(c + dx)}{7d} - \frac{(7A + 6C) \left(-\frac{1}{5} \tan^5(c + dx) - \frac{2}{3} \tan^3(c + dx) - \tan(c + dx)\right)}{7d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^6*(A + C*Sec[c + d*x]^2),x]`

output `(C*Sec[c + d*x]^6*Tan[c + d*x])/(7*d) - ((7*A + 6*C)*(-Tan[c + d*x] - (2*Tan[c + d*x]^3)/3 - Tan[c + d*x]^5/5))/(7*d)`

3.1. $\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx$

3.1.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.1.4 Maple [A] (verified)

Time = 4.59 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.90

method	result
derivativedivides	$\frac{-A\left(-\frac{8}{15}-\frac{\sec(dx+c)^4}{5}-\frac{4\sec(dx+c)^2}{15}\right)\tan(dx+c)-C\left(-\frac{16}{35}-\frac{\sec(dx+c)^6}{7}-\frac{6\sec(dx+c)^4}{35}-\frac{8\sec(dx+c)^2}{35}\right)\tan(dx+c)}{d}$
default	$\frac{-A\left(-\frac{8}{15}-\frac{\sec(dx+c)^4}{5}-\frac{4\sec(dx+c)^2}{15}\right)\tan(dx+c)-C\left(-\frac{16}{35}-\frac{\sec(dx+c)^6}{7}-\frac{6\sec(dx+c)^4}{35}-\frac{8\sec(dx+c)^2}{35}\right)\tan(dx+c)}{d}$
parts	$\frac{A\left(-\frac{8}{15}-\frac{\sec(dx+c)^4}{5}-\frac{4\sec(dx+c)^2}{15}\right)\tan(dx+c)}{d}-\frac{C\left(-\frac{16}{35}-\frac{\sec(dx+c)^6}{7}-\frac{6\sec(dx+c)^4}{35}-\frac{8\sec(dx+c)^2}{35}\right)\tan(dx+c)}{d}$
risch	$\frac{16i(70Ae^{8i(dx+c)}+175Ae^{6i(dx+c)}+210Ce^{6i(dx+c)}+147Ae^{4i(dx+c)}+126Ce^{4i(dx+c)}+49Ae^{2i(dx+c)}+42Ce^{2i(dx+c)}+7)}{105d(e^{2i(dx+c)}+1)^7}$
parallelrisch	$\frac{(1176A+1008C)\sin(3dx+3c)+(392A+336C)\sin(5dx+5c)+(56A+48C)\sin(7dx+7c)+840\sin(dx+c)(A+2C)}{105d(\cos(7dx+7c)+7\cos(5dx+5c)+21\cos(3dx+3c)+35\cos(dx+c))}$
norman	$\frac{2(A+C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d}-\frac{2(A+C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^{13}}{d}+\frac{4(5A+3C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{3d}+\frac{4(5A+3C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^{11}}{3d}+\frac{8(91A+53C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{35d}-\frac{1}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-1\right)^7}$

input `int(sec(d*x+c)^6*(A+C*sec(d*x+c)^2), x, method=_RETURNVERBOSE)`

output $1/d*(-A*(-8/15-1/5*\sec(d*x+c)^4-4/15*\sec(d*x+c)^2)*\tan(d*x+c)-C*(-16/35-1/7*\sec(d*x+c)^6-6/35*\sec(d*x+c)^4-8/35*\sec(d*x+c)^2)*\tan(d*x+c))$

3.1.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.85

$$\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(8(7A + 6C) \cos(dx + c)^6 + 4(7A + 6C) \cos(dx + c)^4 + 3(7A + 6C) \cos(dx + c)^2 + 15C) \sin(dx + c)}{105 d \cos(dx + c)^7}$$

input `integrate(sec(d*x+c)^6*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output $1/105*(8*(7*A + 6*C)*\cos(d*x + c)^6 + 4*(7*A + 6*C)*\cos(d*x + c)^4 + 3*(7*A + 6*C)*\cos(d*x + c)^2 + 15*C)*\sin(d*x + c)/(d*\cos(d*x + c)^7)$

3.1.6 Sympy [F]

$$\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \sec^6(c + dx) dx$$

input `integrate(sec(d*x+c)**6*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*sec(c + d*x)**6, x)`

3.1.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.69

$$\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{15 C \tan(dx + c)^7 + 21 (A + 3 C) \tan(dx + c)^5 + 35 (2 A + 3 C) \tan(dx + c)^3 + 105 (A + C) \tan(dx + c)}{105 d}$$

3.1. $\int \sec^6(c + dx) (A + C \sec^2(c + dx)) dx$

input `integrate(sec(d*x+c)^6*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output $\frac{1}{105} \cdot (15C \tan(dx+c)^7 + 21(A+3C) \tan(dx+c)^5 + 35(2A+3C) \tan(dx+c)^3 + 105(A+C) \tan(dx+c)) / d$

3.1.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.91

$$\int \sec^6(c+dx) (A+C \sec^2(c+dx)) dx$$

$$= \frac{15C \tan(dx+c)^7 + 21A \tan(dx+c)^5 + 63C \tan(dx+c)^5 + 70A \tan(dx+c)^3 + 105C \tan(dx+c)^3 + 105(A+C) \tan(dx+c)}{105d}$$

input `integrate(sec(d*x+c)^6*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output $\frac{1}{105} \cdot (15C \tan(dx+c)^7 + 21A \tan(dx+c)^5 + 63C \tan(dx+c)^5 + 70A \tan(dx+c)^3 + 105C \tan(dx+c)^3 + 105A \tan(dx+c) + 105C \tan(dx+c)) / d$

3.1.9 Mupad [B] (verification not implemented)

Time = 15.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.64

$$\int \sec^6(c+dx) (A+C \sec^2(c+dx)) dx$$

$$= \frac{\frac{C \tan(c+dx)^7}{7} + \left(\frac{A}{5} + \frac{3C}{5}\right) \tan(c+dx)^5 + \left(\frac{2A}{3} + C\right) \tan(c+dx)^3 + (A+C) \tan(c+dx)}{d}$$

input `int((A + C/cos(c + d*x)^2)/cos(c + d*x)^6,x)`

output $\frac{(\tan(c+d*x))^3 \cdot ((2A)/3 + C) + (C \cdot (\tan(c+d*x))^7)/7 + \tan(c+d*x) \cdot (A + C) + \tan(c+d*x)^5 \cdot (A/5 + (3C)/5)}{d}$

3.1. $\int \sec^6(c+dx) (A+C \sec^2(c+dx)) dx$

3.2 $\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx$

3.2.1	Optimal result	53
3.2.2	Mathematica [A] (verified)	53
3.2.3	Rubi [A] (verified)	54
3.2.4	Maple [A] (verified)	56
3.2.5	Fricas [A] (verification not implemented)	56
3.2.6	Sympy [F]	57
3.2.7	Maxima [A] (verification not implemented)	57
3.2.8	Giac [A] (verification not implemented)	58
3.2.9	Mupad [B] (verification not implemented)	58

3.2.1 Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(6A + 5C)\operatorname{arctanh}(\sin(c + dx))}{16d} + \frac{(6A + 5C)\sec(c + dx)\tan(c + dx)}{16d} + \frac{(6A + 5C)\sec^3(c + dx)\tan(c + dx)}{24d} + \frac{C\sec^5(c + dx)\tan(c + dx)}{6d}$$

output `1/16*(6*A+5*C)*arctanh(sin(d*x+c))/d+1/16*(6*A+5*C)*sec(d*x+c)*tan(d*x+c)/d+1/24*(6*A+5*C)*sec(d*x+c)^3*tan(d*x+c)/d+1/6*C*sec(d*x+c)^5*tan(d*x+c)/d`

3.2.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.40

$$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx = \frac{3A\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{5C\operatorname{arctanh}(\sin(c + dx))}{16d} + \frac{3A\sec(c + dx)\tan(c + dx)}{8d} + \frac{5C\sec(c + dx)\tan(c + dx)}{16d} + \frac{A\sec^3(c + dx)\tan(c + dx)}{4d} + \frac{5C\sec^3(c + dx)\tan(c + dx)}{24d} + \frac{C\sec^5(c + dx)\tan(c + dx)}{6d}$$

input `Integrate[Sec[c + d*x]^5*(A + C*Sec[c + d*x]^2),x]`

output `(3*A*ArcTanh[Sin[c + d*x]])/(8*d) + (5*C*ArcTanh[Sin[c + d*x]])/(16*d) + (3*A*Sec[c + d*x]*Tan[c + d*x])/(8*d) + (5*C*Sec[c + d*x]*Tan[c + d*x])/(16*d) + (A*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (5*C*Sec[c + d*x]^3*Tan[c + d*x])/(24*d) + (C*Sec[c + d*x]^5*Tan[c + d*x])/(6*d)`

3.2.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {3042, 4534, 3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^5 \left(A + C \csc\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{6}(6A + 5C) \int \sec^5(c + dx) dx + \frac{C \tan(c + dx) \sec^5(c + dx)}{6d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6}(6A + 5C) \int \csc\left(c + dx + \frac{\pi}{2}\right)^5 dx + \frac{C \tan(c + dx) \sec^5(c + dx)}{6d} \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{6}(6A + 5C) \left(\frac{3}{4} \int \sec^3(c + dx) dx + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right) + \frac{C \tan(c + dx) \sec^5(c + dx)}{6d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6}(6A + 5C) \left(\frac{3}{4} \int \csc\left(c + dx + \frac{\pi}{2}\right)^3 dx + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right) + \frac{C \tan(c + dx) \sec^5(c + dx)}{6d} \\
 & \quad \downarrow \text{4255}
 \end{aligned}$$

$$\frac{1}{6}(6A + 5C) \left(\frac{3}{4} \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right) + \frac{C \tan(c + dx) \sec^5(c + dx)}{6d}$$

↓ 3042

$$5C \left(\frac{3}{4} \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right) + \frac{C \tan(c + dx) \sec^5(c + dx)}{6d}$$

↓ 4257

$$\frac{1}{6}(6A + 5C) \left(\frac{3}{4} \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right) + \frac{C \tan(c + dx) \sec^5(c + dx)}{6d}$$

input `Int[Sec[c + d*x]^5*(A + C*Sec[c + d*x]^2), x]`

output `(C*Sec[c + d*x]^5*Tan[c + d*x])/(6*d) + ((6*A + 5*C)*((Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (3*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d))))/4)/6`

3.2.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`


```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.)
+ (A_.)), x_Symbol] :> Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1)
)), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

3.2.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{A\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)+C\left(-\left(-\frac{\sec(dx+c)^5}{6}-\frac{5\sec(dx+c)^3}{24}-\frac{5\sec(dx+c)}{16}\right)\right)}{d}$
default	$\frac{A\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)+C\left(-\left(-\frac{\sec(dx+c)^5}{6}-\frac{5\sec(dx+c)^3}{24}-\frac{5\sec(dx+c)}{16}\right)\right)}{d}$
parts	$\frac{A\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d} + \frac{C\left(-\left(-\frac{\sec(dx+c)^5}{6}-\frac{5\sec(dx+c)^3}{24}-\frac{5\sec(dx+c)}{16}\right)\right)}{d}$
parallelrisch	$\frac{-270\left(\frac{\cos(6dx+6c)}{15}+\frac{2\cos(4dx+4c)}{5}+\cos(2dx+2c)+\frac{2}{3}\right)\left(A+\frac{5C}{6}\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+270\left(\frac{\cos(6dx+6c)}{15}+\frac{2\cos(4dx+4c)}{5}+\cos(2dx+2c)+\frac{2}{3}\right)}{48d(\cos(6dx+6c)+6\cos(2dx+2c)+3)}$
norman	$\frac{(2A+15C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^5}{4d}+\frac{(2A+15C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^7}{4d}+\frac{(10A+11C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{8d}+\frac{(10A+11C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^{11}}{8d}-\frac{(42A-5C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{24d}}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^6}$
risch	$-\frac{ie^{i(dx+c)}(18Ae^{10i(dx+c)}+15Ce^{10i(dx+c)}+102Ae^{8i(dx+c)}+85Ce^{8i(dx+c)}+84Ae^{6i(dx+c)}+198Ce^{6i(dx+c)}-84Ae^{4i(dx+c)}-15Ce^{4i(dx+c)}-102Ae^{2i(dx+c)}-85Ce^{2i(dx+c)}-18Ae^{2i(dx+c)}-15Ce^{2i(dx+c)}-102A-85C-18A-15C)}{24d(e^{2i(dx+c)}+1)^6}$

```
input int(sec(d*x+c)^5*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)
```

```
output 1/d*(A*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))+C*(-(-1/6*sec(d*x+c)^5-5/24*sec(d*x+c)^3-5/16*sec(d*x+c))*tan(d*x+c)+5/16*ln(sec(d*x+c)+tan(d*x+c))))
```

3.2.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.16

$$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3(6A + 5C) \cos(dx + c)^6 \log(\sin(dx + c) + 1) - 3(6A + 5C) \cos(dx + c)^6 \log(-\sin(dx + c) + 1) + 24A \cos(dx + c)^6 \log(\cos(dx + c) + 1) - 24C \cos(dx + c)^6 \log(\cos(dx + c) - 1)}{96d \cos(dx + c)^6}$$

3.2. $\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx$

input `integrate(sec(d*x+c)^5*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/96*(3*(6*A + 5*C)*cos(d*x + c)^6*log(sin(d*x + c) + 1) - 3*(6*A + 5*C)*cos(d*x + c)^6*log(-sin(d*x + c) + 1) + 2*(3*(6*A + 5*C)*cos(d*x + c)^4 + 2*(6*A + 5*C)*cos(d*x + c)^2 + 8*C)*sin(d*x + c))/(d*cos(d*x + c)^6)`

3.2.6 Sympy [F]

$$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \sec^5(c + dx) dx$$

input `integrate(sec(d*x+c)**5*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*sec(c + d*x)**5, x)`

3.2.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.29

$$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3(6A + 5C) \log(\sin(dx + c) + 1) - 3(6A + 5C) \log(\sin(dx + c) - 1) - \frac{2(3(6A + 5C) \sin(dx + c)^5 - 8(6A + 5C) \sin(dx + c)^3 + 3(10A + 11C) \sin(dx + c))}{\sin(dx + c)^6 - 3 \sin(dx + c)^4 + 3 \sin(dx + c)^2 - 1}}{96d}$$

input `integrate(sec(d*x+c)^5*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/96*(3*(6*A + 5*C)*log(sin(d*x + c) + 1) - 3*(6*A + 5*C)*log(sin(d*x + c) - 1) - 2*(3*(6*A + 5*C)*sin(d*x + c)^5 - 8*(6*A + 5*C)*sin(d*x + c)^3 + 3*(10*A + 11*C)*sin(d*x + c))/(sin(d*x + c)^6 - 3*sin(d*x + c)^4 + 3*sin(d*x + c)^2 - 1))/d`

3.2.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.23

$$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3(6A + 5C) \log(|\sin(dx + c) + 1|) - 3(6A + 5C) \log(|\sin(dx + c) - 1|) - \frac{2(18A \sin(dx+c)^5 + 15C \sin(dx+c)^5)}{96d}}{96d}$$

input `integrate(sec(d*x+c)^5*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/96*(3*(6*A + 5*C)*log(abs(sin(d*x + c) + 1)) - 3*(6*A + 5*C)*log(abs(sin(d*x + c) - 1)) - 2*(18*A*sin(d*x + c)^5 + 15*C*sin(d*x + c)^5 - 48*A*sin(d*x + c)^3 - 40*C*sin(d*x + c)^3 + 30*A*sin(d*x + c) + 33*C*sin(d*x + c)))/(sin(d*x + c)^2 - 1)^3/d`

3.2.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.04

$$\int \sec^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{\operatorname{atanh}(\sin(c + dx)) \left(\frac{3A}{8} + \frac{5C}{16} \right)}{d} - \frac{\left(\frac{3A}{8} + \frac{5C}{16} \right) \sin(c + dx)^5 + \left(-A - \frac{5C}{6} \right) \sin(c + dx)^3 + \left(\frac{5A}{8} + \frac{11C}{16} \right) \sin(c + dx)}{d \left(\sin(c + dx)^6 - 3 \sin(c + dx)^4 + 3 \sin(c + dx)^2 - 1 \right)}$$

input `int((A + C/cos(c + d*x)^2)/cos(c + d*x)^5,x)`

output `(atanh(sin(c + d*x))*((3*A)/8 + (5*C)/16))/d - (sin(c + d*x)*((5*A)/8 + (11*C)/16) - sin(c + d*x)^3*(A + (5*C)/6) + sin(c + d*x)^5*((3*A)/8 + (5*C)/16))/(d*(3*sin(c + d*x)^2 - 3*sin(c + d*x)^4 + sin(c + d*x)^6 - 1))`

3.3 $\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx$

3.3.1	Optimal result	59
3.3.2	Mathematica [A] (verified)	59
3.3.3	Rubi [A] (verified)	60
3.3.4	Maple [A] (verified)	61
3.3.5	Fricas [A] (verification not implemented)	62
3.3.6	Sympy [F]	62
3.3.7	Maxima [A] (verification not implemented)	62
3.3.8	Giac [A] (verification not implemented)	63
3.3.9	Mupad [B] (verification not implemented)	63

3.3.1 Optimal result

Integrand size = 21, antiderivative size = 65

$$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(5A + 4C) \tan(c + dx)}{5d} + \frac{C \sec^4(c + dx) \tan(c + dx)}{5d} + \frac{(5A + 4C) \tan^3(c + dx)}{15d}$$

output `1/5*(5*A+4*C)*tan(d*x+c)/d+1/5*C*sec(d*x+c)^4*tan(d*x+c)/d+1/15*(5*A+4*C)*tan(d*x+c)^3/d`

3.3.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.94

$$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx = \frac{A(\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d} + \frac{C(\tan(c + dx) + \frac{2}{3} \tan^3(c + dx) + \frac{1}{5} \tan^5(c + dx))}{d}$$

input `Integrate[Sec[c + d*x]^4*(A + C*Sec[c + d*x]^2),x]`

output `(A*(Tan[c + d*x] + Tan[c + d*x]^3/3))/d + (C*(Tan[c + d*x] + (2*Tan[c + d*x]^3)/3 + Tan[c + d*x]^5/5))/d`

3.3.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.89, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 4534, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c+dx) (A + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 \left(A + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{5}(5A+4C) \int \sec^4(c+dx) dx + \frac{C \tan(c+dx) \sec^4(c+dx)}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5}(5A+4C) \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 dx + \frac{C \tan(c+dx) \sec^4(c+dx)}{5d} \\
 & \quad \downarrow \text{4254} \\
 & \frac{C \tan(c+dx) \sec^4(c+dx)}{5d} - \frac{(5A+4C) \int (\tan^2(c+dx)+1) d(-\tan(c+dx))}{5d} \\
 & \quad \downarrow \text{2009} \\
 & \frac{C \tan(c+dx) \sec^4(c+dx)}{5d} - \frac{(5A+4C) \left(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx)\right)}{5d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^4*(A + C*Sec[c + d*x]^2),x]`

output `(C*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) - ((5*A + 4*C)*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/(5*d)`

3.3.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.3.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{-A\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)-C\left(-\frac{8}{15}-\frac{\sec(dx+c)^4}{5}-\frac{4\sec(dx+c)^2}{15}\right)\tan(dx+c)}{d}$
default	$\frac{-A\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)-C\left(-\frac{8}{15}-\frac{\sec(dx+c)^4}{5}-\frac{4\sec(dx+c)^2}{15}\right)\tan(dx+c)}{d}$
parts	$\frac{A\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)}{d}-\frac{C\left(-\frac{8}{15}-\frac{\sec(dx+c)^4}{5}-\frac{4\sec(dx+c)^2}{15}\right)\tan(dx+c)}{d}$
parallelrisch	$\frac{(50A+40C)\sin(3dx+3c)+(10A+8C)\sin(5dx+5c)+40\sin(dx+c)(A+2C)}{15d(\cos(5dx+5c)+5\cos(3dx+3c)+10\cos(dx+c))}$
risch	$\frac{4i(15Ae^{6i(dx+c)}+35Ae^{4i(dx+c)}+40Ce^{4i(dx+c)}+25Ae^{2i(dx+c)}+20Ce^{2i(dx+c)}+5A+4C)}{15d(e^{2i(dx+c)}+1)^5}$
norman	$\frac{-\frac{2(A+C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d}-\frac{2(A+C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^9}{d}+\frac{8(2A+C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{3d}+\frac{8(2A+C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^7}{3d}-\frac{4(25A+29C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^5}{15d}}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-1\right)^5}$

input `int(sec(d*x+c)^4*(A+C*sec(d*x+c)^2), x, method=_RETURNVERBOSE)`

output `1/d*(-A*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)-C*(-8/15-1/5*sec(d*x+c)^4-4/15*sec(d*x+c)^2)*tan(d*x+c))`

3.3.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.86

$$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(2(5A + 4C) \cos(dx + c)^4 + (5A + 4C) \cos(dx + c)^2 + 3C) \sin(dx + c)}{15d \cos(dx + c)^5}$$

input `integrate(sec(d*x+c)^4*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/15*(2*(5*A + 4*C)*cos(d*x + c)^4 + (5*A + 4*C)*cos(d*x + c)^2 + 3*C)*sin(d*x + c)/(d*cos(d*x + c)^5)`

3.3.6 Sympy [F]

$$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \sec^4(c + dx) dx$$

input `integrate(sec(d*x+c)**4*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*sec(c + d*x)**4, x)`

3.3.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.66

$$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3C \tan(dx + c)^5 + 5(A + 2C) \tan(dx + c)^3 + 15(A + C) \tan(dx + c)}{15d}$$

input `integrate(sec(d*x+c)^4*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/15*(3*C*tan(d*x + c)^5 + 5*(A + 2*C)*tan(d*x + c)^3 + 15*(A + C)*tan(d*x + c))/d`

3.3.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.88

$$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3 C \tan(dx + c)^5 + 5 A \tan(dx + c)^3 + 10 C \tan(dx + c)^3 + 15 A \tan(dx + c) + 15 C \tan(dx + c)}{15 d}$$

input `integrate(sec(d*x+c)^4*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/15*(3*C*tan(d*x + c)^5 + 5*A*tan(d*x + c)^3 + 10*C*tan(d*x + c)^3 + 15*A*tan(d*x + c) + 15*C*tan(d*x + c))/d`

3.3.9 Mupad [B] (verification not implemented)

Time = 14.97 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

$$\int \sec^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{\frac{C \tan(c+dx)^5}{5} + \left(\frac{A}{3} + \frac{2C}{3}\right) \tan(c + dx)^3 + (A + C) \tan(c + dx)}{d}$$

input `int((A + C/cos(c + d*x)^2)/cos(c + d*x)^4,x)`

output `((C*tan(c + d*x)^5)/5 + tan(c + d*x)*(A + C) + tan(c + d*x)^3*(A/3 + (2*C)/3))/d`

3.4 $\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx$

3.4.1	Optimal result	64
3.4.2	Mathematica [A] (verified)	64
3.4.3	Rubi [A] (verified)	65
3.4.4	Maple [A] (verified)	66
3.4.5	Fricas [A] (verification not implemented)	67
3.4.6	Sympy [F]	68
3.4.7	Maxima [A] (verification not implemented)	68
3.4.8	Giac [A] (verification not implemented)	68
3.4.9	Mupad [B] (verification not implemented)	69

3.4.1 Optimal result

Integrand size = 21, antiderivative size = 70

$$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(4A + 3C)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{(4A + 3C) \sec(c + dx) \tan(c + dx)}{8d} + \frac{C \sec^3(c + dx) \tan(c + dx)}{4d}$$

output `1/8*(4*A+3*C)*arctanh(sin(d*x+c))/d+1/8*(4*A+3*C)*sec(d*x+c)*tan(d*x+c)/d+1/4*C*sec(d*x+c)^3*tan(d*x+c)/d`

3.4.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.33

$$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx = \frac{A\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{3C\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{A \sec(c + dx) \tan(c + dx)}{2d} + \frac{3C \sec(c + dx) \tan(c + dx)}{8d} + \frac{C \sec^3(c + dx) \tan(c + dx)}{4d}$$

input `Integrate[Sec[c + d*x]^3*(A + C*Sec[c + d*x]^2),x]`

output `(A*ArcTanh[Sin[c + d*x]])/(2*d) + (3*C*ArcTanh[Sin[c + d*x]])/(8*d) + (A*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (3*C*Sec[c + d*x]*Tan[c + d*x])/(8*d) + (C*Sec[c + d*x]^3*Tan[c + d*x])/(4*d)`

3.4.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4534, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^3 \left(A + C \csc\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{4}(4A + 3C) \int \sec^3(c + dx) dx + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4}(4A + 3C) \int \csc\left(c + dx + \frac{\pi}{2}\right)^3 dx + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d} \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{4}(4A + 3C) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4}(4A + 3C) \left(\frac{1}{2} \int \csc\left(c + dx + \frac{\pi}{2}\right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d} \\
 & \quad \downarrow \text{4257} \\
 & \frac{1}{4}(4A + 3C) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^3*(A + C*Sec[c + d*x]^2),x]`

output `(C*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((4*A + 3*C)*(ArcTanh[Sin[c + d*x]])/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d))/4`

3.4.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.4.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{A\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) + C\left(-\left(-\frac{\sec(dx+c)^3}{4} - \frac{3\sec(dx+c)}{8}\right)\tan(dx+c) + \frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d}$
default	$\frac{A\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) + C\left(-\left(-\frac{\sec(dx+c)^3}{4} - \frac{3\sec(dx+c)}{8}\right)\tan(dx+c) + \frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d}$
parts	$\frac{A\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)}{d} + \frac{C\left(-\left(-\frac{\sec(dx+c)^3}{4} - \frac{3\sec(dx+c)}{8}\right)\tan(dx+c) + \frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d}$
parallelrisch	$\frac{-8\left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4}\right) + \cos(2dx+2c)\left(A + \frac{3C}{4}\right)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 8\left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4}\right) + \cos(2dx+2c)\left(A + \frac{3C}{4}\right)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4d(\cos(4dx+4c) + 4\cos(2dx+2c) + 3)}$
norman	$\frac{-\frac{(4A-3C)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{4d} - \frac{(4A-3C)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{4d} + \frac{(4A+5C)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4d} + \frac{(4A+5C)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^7}{4d}}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^4} - \frac{(4A+3C)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{8d}$
risch	$-\frac{ie^{i(dx+c)}(4Ae^{6i(dx+c)} + 3Ce^{6i(dx+c)} + 4Ae^{4i(dx+c)} + 11Ce^{4i(dx+c)} - 4Ae^{2i(dx+c)} - 11Ce^{2i(dx+c)} - 4A - 3C)}{4d(e^{2i(dx+c)} + 1)^4} + \frac{\ln(e^{i(dx+c)} + 1)}{4d}$

input `int(sec(d*x+c)^3*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(A*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+C*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c))))`

3.4.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.36

$$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(4A + 3C) \cos(dx + c)^4 \log(\sin(dx + c) + 1) - (4A + 3C) \cos(dx + c)^4 \log(-\sin(dx + c) + 1) + 2((4A + 3C) \cos(dx + c)^2 \sin(dx + c) \log(\sin(dx + c) + 1) - (4A + 3C) \cos(dx + c)^2 \sin(dx + c) \log(-\sin(dx + c) + 1))}{16d \cos(dx + c)^4}$$

input `integrate(sec(d*x+c)^3*(A+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/16*((4*A + 3*C)*cos(d*x + c)^4*log(sin(d*x + c) + 1) - (4*A + 3*C)*cos(d*x + c)^4*log(-sin(d*x + c) + 1) + 2*((4*A + 3*C)*cos(d*x + c)^2 + 2*C)*sin(d*x + c))/d*cos(d*x + c)^4`

3.4.6 Sympy [F]

$$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \sec^3(c + dx) dx$$

input `integrate(sec(d*x+c)**3*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*sec(c + d*x)**3, x)`

3.4.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.39

$$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(4A + 3C) \log(\sin(dx + c) + 1) - (4A + 3C) \log(\sin(dx + c) - 1) - \frac{2((4A + 3C) \sin(dx + c)^3 - (4A + 5C) \sin(dx + c))}{\sin(dx + c)^4 - 2 \sin(dx + c)^2 + 1}}{16d}$$

input `integrate(sec(d*x+c)^3*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/16*((4*A + 3*C)*log(sin(d*x + c) + 1) - (4*A + 3*C)*log(sin(d*x + c) - 1) - 2*((4*A + 3*C)*sin(d*x + c)^3 - (4*A + 5*C)*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1))/d`

3.4.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.40

$$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(4A + 3C) \log(|\sin(dx + c) + 1|) - (4A + 3C) \log(|\sin(dx + c) - 1|) - \frac{2(4A \sin(dx + c)^3 + 3C \sin(dx + c)^3 - 4A \sin(dx + c))}{(\sin(dx + c)^2 - 1)}}{16d}$$

input `integrate(sec(d*x+c)^3*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output $1/16*((4*A + 3*C)*\log(\text{abs}(\sin(d*x + c) + 1)) - (4*A + 3*C)*\log(\text{abs}(\sin(d*x + c) - 1)) - 2*(4*A*\sin(d*x + c)^3 + 3*C*\sin(d*x + c)^3 - 4*A*\sin(d*x + c) - 5*C*\sin(d*x + c)))/(\sin(d*x + c)^2 - 1)^2)/d$

3.4.9 Mupad [B] (verification not implemented)

Time = 15.55 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.10

$$\int \sec^3(c + dx) (A + C \sec^2(c + dx)) dx = \frac{\sin(c + dx) \left(\frac{A}{2} + \frac{5C}{8}\right) - \sin(c + dx)^3 \left(\frac{A}{2} + \frac{3C}{8}\right)}{d (\sin(c + dx)^4 - 2 \sin(c + dx)^2 + 1)} + \frac{\operatorname{atanh}(\sin(c + dx)) \left(\frac{A}{2} + \frac{3C}{8}\right)}{d}$$

input $\text{int}((A + C/\cos(c + d*x)^2)/\cos(c + d*x)^3,x)$

output $(\sin(c + d*x)*(A/2 + (5*C)/8) - \sin(c + d*x)^3*(A/2 + (3*C)/8))/(d*(\sin(c + d*x)^4 - 2*\sin(c + d*x)^2 + 1)) + (\operatorname{atanh}(\sin(c + d*x))*(A/2 + (3*C)/8))/d$

3.5 $\int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx$

3.5.1	Optimal result	70
3.5.2	Mathematica [A] (verified)	70
3.5.3	Rubi [A] (verified)	71
3.5.4	Maple [A] (verified)	72
3.5.5	Fricas [A] (verification not implemented)	73
3.5.6	Sympy [F]	73
3.5.7	Maxima [A] (verification not implemented)	73
3.5.8	Giac [A] (verification not implemented)	74
3.5.9	Mupad [B] (verification not implemented)	74

3.5.1 Optimal result

Integrand size = 21, antiderivative size = 43

$$\int \sec^2(c+dx) (A+C \sec^2(c+dx)) dx = \frac{(3A + 2C) \tan(c + dx)}{3d} + \frac{C \sec^2(c + dx) \tan(c + dx)}{3d}$$

output `1/3*(3*A+2*C)*tan(d*x+c)/d+1/3*C*sec(d*x+c)^2*tan(d*x+c)/d`

3.5.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

$$\int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{A \tan(c + dx)}{d} + \frac{C(\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d}$$

input `Integrate[Sec[c + d*x]^2*(A + C*Sec[c + d*x]^2),x]`

output `(A*Tan[c + d*x])/d + (C*(Tan[c + d*x] + Tan[c + d*x]^3/3))/d`

3.5.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 4534, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx) (A + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 \left(A + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{3}(3A+2C) \int \sec^2(c+dx) dx + \frac{C \tan(c+dx) \sec^2(c+dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3}(3A+2C) \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx + \frac{C \tan(c+dx) \sec^2(c+dx)}{3d} \\
 & \quad \downarrow \text{4254} \\
 & \frac{C \tan(c+dx) \sec^2(c+dx)}{3d} - \frac{(3A+2C) \int 1d(-\tan(c+dx))}{3d} \\
 & \quad \downarrow \text{24} \\
 & \frac{(3A+2C) \tan(c+dx)}{3d} + \frac{C \tan(c+dx) \sec^2(c+dx)}{3d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^2*(A + C*Sec[c + d*x]^2), x]`

output `((3*A + 2*C)*Tan[c + d*x])/(3*d) + (C*Sec[c + d*x]^2*Tan[c + d*x])/(3*d)`

3.5.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_. + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.5.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{A \tan(dx+c) - C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$	35
default	$\frac{A \tan(dx+c) - C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$	35
parts	$\frac{A \tan(dx+c)}{d} - \frac{C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$	37
parallelrisc	$\frac{(3A+2C) \sin(3dx+3c) + 3 \sin(dx+c)(A+2C)}{3d(\cos(3dx+3c) + 3 \cos(dx+c))}$	57
risc	$\frac{2i(3A e^{4i(dx+c)} + 6A e^{2i(dx+c)} + 6C e^{2i(dx+c)} + 3A + 2C)}{3d(e^{2i(dx+c)} + 1)^3}$	63
norman	$\frac{-\frac{2(A+C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} - \frac{2(A+C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{d} + \frac{4(3A+C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{3d}}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)^3}$	75

input `int(sec(d*x+c)^2*(A+C*sec(d*x+c)^2), x, method=_RETURNVERBOSE)`

output `1/d*(A*tan(d*x+c)-C*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c))`

3.5.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.86

$$\int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{((3A + 2C) \cos(dx + c)^2 + C) \sin(dx + c)}{3d \cos(dx + c)^3}$$

input `integrate(sec(d*x+c)^2*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/3*((3*A + 2*C)*cos(d*x + c)^2 + C)*sin(d*x + c)/(d*cos(d*x + c)^3)`

3.5.6 Sympy [F]

$$\int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*sec(c + d*x)**2, x)`

3.5.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\begin{aligned} \int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx \\ = \frac{(\tan(dx + c)^3 + 3 \tan(dx + c))C + 3A \tan(dx + c)}{3d} \end{aligned}$$

input `integrate(sec(d*x+c)^2*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/3*((tan(d*x + c)^3 + 3*tan(d*x + c))*C + 3*A*tan(d*x + c))/d`

3.5.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{C \tan(dx + c)^3 + 3A \tan(dx + c) + 3C \tan(dx + c)}{3d}$$

input `integrate(sec(d*x+c)^2*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/3*(C*tan(d*x + c)^3 + 3*A*tan(d*x + c) + 3*C*tan(d*x + c))/d`

3.5.9 Mupad [B] (verification not implemented)

Time = 15.74 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.65

$$\int \sec^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{C \tan(c + dx)^3}{3d} + \frac{\tan(c + dx) (A + C)}{d}$$

input `int((A + C/cos(c + d*x)^2)/cos(c + d*x)^2,x)`

output `(C*tan(c + d*x)^3)/(3*d) + (tan(c + d*x)*(A + C))/d`

3.6 $\int \sec(c + dx) (A + C \sec^2(c + dx)) dx$

3.6.1	Optimal result	75
3.6.2	Mathematica [A] (verified)	75
3.6.3	Rubi [A] (verified)	76
3.6.4	Maple [A] (verified)	77
3.6.5	Fricas [A] (verification not implemented)	77
3.6.6	Sympy [F]	78
3.6.7	Maxima [A] (verification not implemented)	78
3.6.8	Giac [A] (verification not implemented)	78
3.6.9	Mupad [B] (verification not implemented)	79

3.6.1 Optimal result

Integrand size = 19, antiderivative size = 40

$$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(2A + C)\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{C \sec(c + dx) \tan(c + dx)}{2d}$$

```
output 1/2*(2*A+C)*arctanh(sin(d*x+c))/d+1/2*C*sec(d*x+c)*tan(d*x+c)/d
```

3.6.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.20

$$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx = \frac{A\operatorname{arctanh}(\sin(c + dx))}{d} + \frac{C\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{C \sec(c + dx) \tan(c + dx)}{2d}$$

```
input Integrate[Sec[c + d*x]*(A + C*Sec[c + d*x]^2),x]
```

```
output (A*ArcTanh[Sin[c + d*x]])/d + (C*ArcTanh[Sin[c + d*x]])/(2*d) + (C*Sec[c + d*x]*Tan[c + d*x])/(2*d)
```

3.6.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {3042, 4534, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c + dx) (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right) \left(A + C \csc\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{2}(2A + C) \int \sec(c + dx) dx + \frac{C \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(2A + C) \int \csc\left(c + dx + \frac{\pi}{2}\right) dx + \frac{C \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow \text{4257} \\
 & \frac{(2A + C) \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{C \tan(c + dx) \sec(c + dx)}{2d}
 \end{aligned}$$

input `Int[Sec[c + d*x]*(A + C*Sec[c + d*x]^2),x]`

output `((2*A + C)*ArcTanh[Sin[c + d*x]]/(2*d) + (C*Sec[c + d*x]*Tan[c + d*x])/(2*d)`

3.6.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.)
+ (A_.)), x_Symbol] :> Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1)
)), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

3.6.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.38

method	result	size
derivativedivides	$\frac{A \ln(\sec(dx+c)+\tan(dx+c))+C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$	55
default	$\frac{A \ln(\sec(dx+c)+\tan(dx+c))+C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$	55
parts	$\frac{A \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$	57
parallelrisc	$\frac{-\left(A+\frac{C}{2}\right)(1+\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+\left(A+\frac{C}{2}\right)(1+\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)+C \sin(dx+c)}{d(1+\cos(2dx+2c))}$	86
norman	$\frac{\frac{C \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d} + \frac{C \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{d^2}}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^2} - \frac{(2A+C) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{2d} + \frac{(2A+C) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{2d}$	93
risc	$-\frac{iC(e^{3i(dx+c)}-e^{i(dx+c)})}{d(e^{2i(dx+c)}+1)^2} - \frac{\ln(e^{i(dx+c)}-i)A}{d} - \frac{\ln(e^{i(dx+c)}-i)C}{2d} + \frac{\ln(e^{i(dx+c)}+i)A}{d} + \frac{\ln(e^{i(dx+c)}+i)C}{2d}$	111

```
input int(sec(d*x+c)*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)
```

```
output 1/d*(A*ln(sec(d*x+c)+tan(d*x+c))+C*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d
*x+c)+tan(d*x+c))))
```

3.6.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.80

$$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(2A + C) \cos(dx + c)^2 \log(\sin(dx + c) + 1) - (2A + C) \cos(dx + c)^2 \log(-\sin(dx + c) + 1) + 2C \sin(dx + c)}{4d \cos(dx + c)^2}$$

```
input integrate(sec(d*x+c)*(A+C*sec(d*x+c)^2),x, algorithm="fricas")
```

output $1/4*((2*A + C)*\cos(d*x + c)^2*\log(\sin(d*x + c) + 1) - (2*A + C)*\cos(d*x + c)^2*\log(-\sin(d*x + c) + 1) + 2*C*\sin(d*x + c))/(d*\cos(d*x + c)^2)$

3.6.6 Sympy [F]

$$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \sec(c + dx) dx$$

input `integrate(sec(d*x+c)*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*sec(c + d*x), x)`

3.6.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.45

$$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(2A + C) \log(\sin(dx + c) + 1) - (2A + C) \log(\sin(dx + c) - 1) - \frac{2C \sin(dx+c)}{\sin(dx+c)^2 - 1}}{4d}$$

input `integrate(sec(d*x+c)*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output $1/4*((2*A + C)*\log(\sin(d*x + c) + 1) - (2*A + C)*\log(\sin(d*x + c) - 1) - 2*C*\sin(d*x + c)/(\sin(d*x + c)^2 - 1))/d$

3.6.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.50

$$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(2A + C) \log(|\sin(dx + c) + 1|) - (2A + C) \log(|\sin(dx + c) - 1|) - \frac{2C \sin(dx+c)}{\sin(dx+c)^2 - 1}}{4d}$$

3.6. $\int \sec(c + dx) (A + C \sec^2(c + dx)) dx$

input `integrate(sec(d*x+c)*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/4*((2*A + C)*log(abs(sin(d*x + c) + 1)) - (2*A + C)*log(abs(sin(d*x + c) - 1)) - 2*C*sin(d*x + c)/(sin(d*x + c)^2 - 1))/d`

3.6.9 Mupad [B] (verification not implemented)

Time = 16.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02

$$\int \sec(c + dx) (A + C \sec^2(c + dx)) dx = \frac{\operatorname{atanh}(\sin(c + dx)) (A + \frac{C}{2})}{d} - \frac{C \sin(c + dx)}{2d (\sin(c + dx)^2 - 1)}$$

input `int((A + C/cos(c + d*x)^2)/cos(c + d*x),x)`

output `(atanh(sin(c + d*x))*(A + C/2))/d - (C*sin(c + d*x))/(2*d*(sin(c + d*x)^2 - 1))`

3.7 $\int (A + C \sec^2(c + dx)) dx$

3.7.1	Optimal result	80
3.7.2	Mathematica [A] (verified)	80
3.7.3	Rubi [A] (verified)	81
3.7.4	Maple [A] (verified)	81
3.7.5	Fricas [B] (verification not implemented)	82
3.7.6	Sympy [F]	82
3.7.7	Maxima [A] (verification not implemented)	82
3.7.8	Giac [A] (verification not implemented)	83
3.7.9	Mupad [B] (verification not implemented)	83

3.7.1 Optimal result

Integrand size = 12, antiderivative size = 15

$$\int (A + C \sec^2(c + dx)) dx = Ax + \frac{C \tan(c + dx)}{d}$$

output `A*x+C*tan(d*x+c)/d`

3.7.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (A + C \sec^2(c + dx)) dx = Ax + \frac{C \tan(c + dx)}{d}$$

input `Integrate[A + C*Sec[c + d*x]^2,x]`

output `A*x + (C*Tan[c + d*x])/d`

3.7.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (A + C \sec^2(c + dx)) dx$$

$$\downarrow \text{2009}$$

$$Ax + \frac{C \tan(c + dx)}{d}$$

input `Int[A + C*Sec[c + d*x]^2,x]`

output `A*x + (C*Tan[c + d*x])/d`

3.7.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.7.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$Ax + \frac{C \tan(dx+c)}{d}$	16
parts	$Ax + \frac{C \tan(dx+c)}{d}$	16
derivativedivides	$\frac{(dx+c)A+C \tan(dx+c)}{d}$	21
risch	$Ax + \frac{2iC}{d(e^{2i(dx+c)}+1)}$	25
parallelrisk	$-\frac{2C \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)} + Ax$	35
norman	$\frac{Ax \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - Ax - \frac{2C \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1}$	51

input `int(A+C*sec(d*x+c)^2,x,method=_RETURNVERBOSE)`

output `A*x+C*tan(d*x+c)/d`

3.7.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. $2(15) = 30$.

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.07

$$\int (A + C \sec^2(c + dx)) dx = \frac{A dx \cos(dx + c) + C \sin(dx + c)}{d \cos(dx + c)}$$

input `integrate(A+C*sec(d*x+c)^2,x, algorithm="fricas")`

output `(A*d*x*cos(d*x + c) + C*sin(d*x + c))/(d*cos(d*x + c))`

3.7.6 Sympy [F]

$$\int (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) dx$$

input `integrate(A+C*sec(d*x+c)**2,x)`

output `Integral(A + C*sec(c + d*x)**2, x)`

3.7.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (A + C \sec^2(c + dx)) dx = Ax + \frac{C \tan(dx + c)}{d}$$

input `integrate(A+C*sec(d*x+c)^2,x, algorithm="maxima")`

output `A*x + C*tan(d*x + c)/d`

3.7.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (A + C \sec^2(c + dx)) dx = Ax + \frac{C \tan(dx + c)}{d}$$

input `integrate(A+C*sec(d*x+c)^2,x, algorithm="giac")`

output `A*x + C*tan(d*x + c)/d`

3.7.9 Mupad [B] (verification not implemented)

Time = 16.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int (A + C \sec^2(c + dx)) dx = \frac{C \tan(c + dx) + A dx}{d}$$

input `int(A + C/cos(c + d*x)^2,x)`

output `(C*tan(c + d*x) + A*d*x)/d`

3.8 $\int \cos(c + dx) (A + C \sec^2(c + dx)) dx$

3.8.1	Optimal result	84
3.8.2	Mathematica [A] (verified)	84
3.8.3	Rubi [A] (verified)	85
3.8.4	Maple [A] (verified)	86
3.8.5	Fricas [A] (verification not implemented)	86
3.8.6	Sympy [F]	87
3.8.7	Maxima [A] (verification not implemented)	87
3.8.8	Giac [A] (verification not implemented)	87
3.8.9	Mupad [B] (verification not implemented)	88

3.8.1 Optimal result

Integrand size = 19, antiderivative size = 24

$$\int \cos(c + dx) (A + C \sec^2(c + dx)) dx = \frac{C \arctanh(\sin(c + dx))}{d} + \frac{A \sin(c + dx)}{d}$$

output `C*arctanh(sin(d*x+c))/d+A*sin(d*x+c)/d`

3.8.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.46

$$\int \cos(c + dx) (A + C \sec^2(c + dx)) dx = \frac{C \arctanh(\sin(c + dx))}{d} + \frac{A \cos(dx) \sin(c)}{d} + \frac{A \cos(c) \sin(dx)}{d}$$

input `Integrate[Cos[c + d*x]*(A + C*Sec[c + d*x]^2),x]`

output `(C*ArcTanh[Sin[c + d*x]])/d + (A*Cos[d*x]*Sin[c])/d + (A*Cos[c]*Sin[d*x])/d`

3.8.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {3042, 4533, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx) (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4533} \\
 & C \int \sec(c + dx) dx + \frac{A \sin(c + dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & C \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{A \sin(c + dx)}{d} \\
 & \quad \downarrow \text{4257} \\
 & \frac{A \sin(c + dx)}{d} + \frac{C \operatorname{arctanh}(\sin(c + dx))}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]*(A + C*Sec[c + d*x]^2),x]`

output `(C*ArcTanh[Sin[c + d*x]])/d + (A*Sin[c + d*x])/d`

3.8.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4533 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.)
+ (A_.)), x_Symbol] :> Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] +
Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; Fr
eeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]
```

3.8.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.25

method	result	size
derivativedivides	$\frac{A \sin(dx+c) + C \ln(\sec(dx+c) + \tan(dx+c))}{d}$	30
default	$\frac{A \sin(dx+c) + C \ln(\sec(dx+c) + \tan(dx+c))}{d}$	30
parallelrisch	$\frac{A \sin(dx+c) - C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{d}$	43
risch	$-\frac{iA e^{i(dx+c)}}{2d} + \frac{iA e^{-i(dx+c)}}{2d} + \frac{\ln(e^{i(dx+c)} + i)C}{d} - \frac{\ln(e^{i(dx+c)} - i)C}{d}$	71
norman	$\frac{-\frac{2A \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{2A \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{d}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)} + \frac{C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{d} - \frac{C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{d}$	101

```
input int(cos(d*x+c)*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)
```

```
output 1/d*(A*sin(d*x+c)+C*ln(sec(d*x+c)+tan(d*x+c)))
```

3.8.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.67

$$\int \cos(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{C \log(\sin(dx + c) + 1) - C \log(-\sin(dx + c) + 1) + 2A \sin(dx + c)}{2d}$$

```
input integrate(cos(d*x+c)*(A+C*sec(d*x+c)^2),x, algorithm="fracas")
```

```
output 1/2*(C*log(sin(d*x + c) + 1) - C*log(-sin(d*x + c) + 1) + 2*A*sin(d*x + c)
)/d
```

3.8.6 Sympy [F]

$$\int \cos(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \cos(c + dx) dx$$

input `integrate(cos(d*x+c)*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*cos(c + d*x), x)`

3.8.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.58

$$\begin{aligned} & \int \cos(c + dx) (A + C \sec^2(c + dx)) dx \\ &= \frac{C(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 2A \sin(dx + c)}{2d} \end{aligned}$$

input `integrate(cos(d*x+c)*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/2*(C*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 2*A*sin(d*x + c)) /d`

3.8.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.67

$$\begin{aligned} & \int \cos(c + dx) (A + C \sec^2(c + dx)) dx \\ &= \frac{C \log(|\sin(dx + c) + 1|) - C \log(|\sin(dx + c) - 1|) + 2A \sin(dx + c)}{2d} \end{aligned}$$

input `integrate(cos(d*x+c)*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/2*(C*log(abs(sin(d*x + c) + 1)) - C*log(abs(sin(d*x + c) - 1)) + 2*A*sin(d*x + c))/d`

3.8.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \cos(c + dx) (A + C \sec^2(c + dx)) dx = \frac{A \sin(c + dx) + C \operatorname{atanh}(\sin(c + dx))}{d}$$

input `int(cos(c + d*x)*(A + C/cos(c + d*x)^2),x)`

output `(A*sin(c + d*x) + C*atanh(sin(c + d*x)))/d`

3.9 $\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx$

3.9.1	Optimal result	89
3.9.2	Mathematica [A] (verified)	89
3.9.3	Rubi [A] (verified)	90
3.9.4	Maple [A] (verified)	91
3.9.5	Fricas [A] (verification not implemented)	91
3.9.6	Sympy [A] (verification not implemented)	92
3.9.7	Maxima [A] (verification not implemented)	92
3.9.8	Giac [A] (verification not implemented)	92
3.9.9	Mupad [B] (verification not implemented)	93

3.9.1 Optimal result

Integrand size = 21, antiderivative size = 31

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{1}{2}(A + 2C)x + \frac{A \cos(c + dx) \sin(c + dx)}{2d}$$

output `1/2*(A+2*C)*x+1/2*A*cos(d*x+c)*sin(d*x+c)/d`

3.9.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx = Cx + \frac{A(c + dx)}{2d} + \frac{A \sin(2(c + dx))}{4d}$$

input `Integrate[Cos[c + d*x]^2*(A + C*Sec[c + d*x]^2),x]`

output `C*x + (A*(c + d*x))/(2*d) + (A*Sin[2*(c + d*x)])/(4*d)`

3.9.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4533, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})^2} dx$$

$$\downarrow \text{4533}$$

$$\frac{1}{2}(A + 2C) \int 1 dx + \frac{A \sin(c + dx) \cos(c + dx)}{2d}$$

$$\downarrow \text{24}$$

$$\frac{A \sin(c + dx) \cos(c + dx)}{2d} + \frac{1}{2}x(A + 2C)$$

input `Int[Cos[c + d*x]^2*(A + C*Sec[c + d*x]^2),x]`

output `((A + 2*C)*x)/2 + (A*Cos[c + d*x]*Sin[c + d*x])/(2*d)`

3.9.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

3.9.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

method	result
risch	$\frac{Ax}{2} + Cx + \frac{A \sin(2dx+2c)}{4d}$
parallelrisch	$\frac{A \sin(2dx+2c)+2(A+2C)xd}{4d}$
derivativedivides	$\frac{A \left(\frac{\sin(dx+c)}{2} \cos(dx+c) + \frac{dx}{2} + \frac{c}{2} \right) + C(dx+c)}{d}$
default	$\frac{A \left(\frac{\sin(dx+c)}{2} \cos(dx+c) + \frac{dx}{2} + \frac{c}{2} \right) + C(dx+c)}{d}$
norman	$\frac{\left(-\frac{A}{2}-C\right)x + \left(-\frac{A}{2}-C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \left(\frac{A}{2}+C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + \left(\frac{A}{2}+C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 - \frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{2A \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$

input `int(cos(d*x+c)^2*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/2*A*x+C*x+1/4*A/d*sin(2*d*x+2*c)`

3.9.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(A + 2C)dx + A \cos(dx + c) \sin(dx + c)}{2d}$$

input `integrate(cos(d*x+c)^2*(A+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/2*((A + 2*C)*d*x + A*cos(d*x + c)*sin(d*x + c))/d`

3.9.6 Sympy [A] (verification not implemented)

Time = 2.52 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.65

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= A \left(\begin{cases} \frac{x \sin^2(c+dx)}{2} + \frac{x \cos^2(c+dx)}{2} + \frac{\sin(c+dx)\cos(c+dx)}{2d} & \text{for } d \neq 0 \\ x \cos^2(c) & \text{otherwise} \end{cases} \right) + Cx$$

input `integrate(cos(d*x+c)**2*(A+C*sec(d*x+c)**2),x)`output `A*Piecewise((x*sin(c + d*x)**2/2 + x*cos(c + d*x)**2/2 + sin(c + d*x)*cos(c + d*x)/(2*d), Ne(d, 0)), (x*cos(c)**2, True)) + C*x`**3.9.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(dx + c)(A + 2C) + \frac{A \tan(dx+c)}{\tan(dx+c)^2+1}}{2d}$$

input `integrate(cos(d*x+c)^2*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`output `1/2*((d*x + c)*(A + 2*C) + A*tan(d*x + c)/(tan(d*x + c)^2 + 1))/d`**3.9.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(dx + c)(A + 2C) + \frac{A \tan(dx+c)}{\tan(dx+c)^2+1}}{2d}$$

input `integrate(cos(d*x+c)^2*(A+C*sec(d*x+c)^2),x, algorithm="giac")`output `1/2*((d*x + c)*(A + 2*C) + A*tan(d*x + c)/(tan(d*x + c)^2 + 1))/d`

3.9.9 Mupad [B] (verification not implemented)

Time = 15.91 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \cos^2(c + dx) (A + C \sec^2(c + dx)) dx = \frac{\frac{A \sin(2c+2dx)}{4} + dx \left(\frac{A}{2} + C\right)}{d}$$

input `int(cos(c + d*x)^2*(A + C/cos(c + d*x)^2),x)`

output `((A*sin(2*c + 2*d*x))/4 + d*x*(A/2 + C))/d`

3.10 $\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx$

3.10.1	Optimal result	94
3.10.2	Mathematica [A] (verified)	94
3.10.3	Rubi [A] (verified)	95
3.10.4	Maple [A] (verified)	96
3.10.5	Fricas [A] (verification not implemented)	97
3.10.6	Sympy [F]	97
3.10.7	Maxima [A] (verification not implemented)	97
3.10.8	Giac [A] (verification not implemented)	98
3.10.9	Mupad [B] (verification not implemented)	98

3.10.1 Optimal result

Integrand size = 21, antiderivative size = 30

$$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(A + C) \sin(c + dx)}{d} - \frac{A \sin^3(c + dx)}{3d}$$

output `(A+C)*sin(d*x+c)/d-1/3*A*sin(d*x+c)^3/d`

3.10.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.67

$$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx = \frac{C \cos(dx) \sin(c)}{d} + \frac{C \cos(c) \sin(dx)}{d} + \frac{A \sin(c + dx)}{d} - \frac{A \sin^3(c + dx)}{3d}$$

input `Integrate[Cos[c + d*x]^3*(A + C*Sec[c + d*x]^2),x]`

output `(C*Cos[d*x]*Sin[c])/d + (C*Cos[c]*Sin[d*x])/d + (A*SIn[c + d*x])/d - (A*Sin[c + d*x]^3)/(3*d)`

3.10.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 4532, 3042, 3492, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c+dx) (A + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c+dx + \frac{\pi}{2})^2}{\csc(c+dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{4532} \\
 & \int \cos(c+dx) (A \cos^2(c+dx) + C) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(c+dx + \frac{\pi}{2}) \left(A \sin(c+dx + \frac{\pi}{2})^2 + C \right) dx \\
 & \quad \downarrow \text{3492} \\
 & - \frac{\int (-A \sin^2(c+dx) + A + C) d(-\sin(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & - \frac{\frac{1}{3} A \sin^3(c+dx) - (A + C) \sin(c+dx)}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^3*(A + C*Sec[c + d*x]^2),x]`

output `-((-((A + C)*Sin[c + d*x]) + (A*SIN[c + d*x]^3)/3)/d)`

3.10.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3492 Int[sin[(e_.) + (f_.)*(x_.)]^(m_.)*((A_) + (C_.)*sin[(e_.) + (f_.)*(x_.)]^2), x_Symbol] := Simp[-f^(-1) Subst[Int[(1 - x^2)^((m - 1)/2)*(A + C - C*x^2), x], x, Cos[e + f*x]], x] /; FreeQ[{e, f, A, C}, x] && IGtQ[(m + 1)/2, 0]
```

```
rule 4532 Int[csc[(e_.) + (f_.)*(x_.)]^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_)), x_Symbol] := Int[(C + A*Sin[e + f*x]^2)/Sin[e + f*x]^(m + 2), x] /; FreeQ[{e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && ILtQ[(m + 1)/2, 0]
```

3.10.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

method	result	size
parallelrisch	$\frac{A \sin(3dx+3c)+9\left(A+\frac{4C}{3}\right) \sin(dx+c)}{12d}$	31
derivativedivides	$\frac{A(2+\cos(dx+c)^2) \sin(dx+c)}{3d} + C \sin(dx+c)$	33
default	$\frac{A(2+\cos(dx+c)^2) \sin(dx+c)}{3d} + C \sin(dx+c)$	33
risch	$\frac{3A \sin(dx+c)}{4d} + \frac{C \sin(dx+c)}{d} + \frac{A \sin(3dx+3c)}{12d}$	40
norman	$\frac{2(A-3C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{3d} - \frac{2(A-3C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^5}{3d} - \frac{2(A+C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d} + \frac{2(A+C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^7}{d}$ $\frac{\left(1+\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2\right)^3 \left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-1\right)}{\left(1+\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2\right)^3 \left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-1\right)}$	111

```
input int(cos(d*x+c)^3*(A+C*sec(d*x+c)^2), x, method=_RETURNVERBOSE)
```

```
output 1/12*(A*sin(3*d*x+3*c)+9*(A+4/3*C)*sin(d*x+c))/d
```

3.10. $\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx$

3.10.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(A \cos(dx + c)^2 + 2A + 3C) \sin(dx + c)}{3d}$$

input `integrate(cos(d*x+c)^3*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`output `1/3*(A*cos(d*x + c)^2 + 2*A + 3*C)*sin(d*x + c)/d`**3.10.6 Sympy [F]**

$$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \cos^3(c + dx) dx$$

input `integrate(cos(d*x+c)**3*(A+C*sec(d*x+c)**2),x)`output `Integral((A + C*sec(c + d*x)**2)*cos(c + d*x)**3, x)`**3.10.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.90

$$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx = -\frac{A \sin(dx + c)^3 - 3(A + C) \sin(dx + c)}{3d}$$

input `integrate(cos(d*x+c)^3*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`output `-1/3*(A*sin(d*x + c)^3 - 3*(A + C)*sin(d*x + c))/d`

3.10.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

$$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= -\frac{A \sin(dx + c)^3 - 3A \sin(dx + c) - 3C \sin(dx + c)}{3d}$$

input `integrate(cos(d*x+c)^3*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `-1/3*(A*sin(d*x + c)^3 - 3*A*sin(d*x + c) - 3*C*sin(d*x + c))/d`

3.10.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \cos^3(c + dx) (A + C \sec^2(c + dx)) dx = -\frac{\frac{A \sin(c+dx)^3}{3} - \sin(c + dx) (A + C)}{d}$$

input `int(cos(c + d*x)^3*(A + C/cos(c + d*x)^2),x)`

output `-((A*sin(c + d*x)^3)/3 - sin(c + d*x)*(A + C))/d`

3.11 $\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx$

3.11.1	Optimal result	99
3.11.2	Mathematica [A] (verified)	99
3.11.3	Rubi [A] (verified)	100
3.11.4	Maple [A] (verified)	101
3.11.5	Fricas [A] (verification not implemented)	102
3.11.6	Sympy [F]	102
3.11.7	Maxima [A] (verification not implemented)	102
3.11.8	Giac [A] (verification not implemented)	103
3.11.9	Mupad [B] (verification not implemented)	103

3.11.1 Optimal result

Integrand size = 21, antiderivative size = 61

$$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx = \frac{1}{8}(3A + 4C)x + \frac{(3A + 4C) \cos(c + dx) \sin(c + dx)}{8d} + \frac{A \cos^3(c + dx) \sin(c + dx)}{4d}$$

output `1/8*(3*A+4*C)*x+1/8*(3*A+4*C)*cos(d*x+c)*sin(d*x+c)/d+1/4*A*cos(d*x+c)^3*sin(d*x+c)/d`

3.11.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.74

$$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx = \frac{4(3A + 4C)(c + dx) + 8(A + C) \sin(2(c + dx)) + A \sin(4(c + dx))}{32d}$$

input `Integrate[Cos[c + d*x]^4*(A + C*Sec[c + d*x]^2),x]`

output `(4*(3*A + 4*C)*(c + d*x) + 8*(A + C)*Sin[2*(c + d*x)] + A*Sine[4*(c + d*x)])/(32*d)`

3.11.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.97, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 4533, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(c+dx) (A + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c+dx + \frac{\pi}{2})^2}{\csc(c+dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{4533} \\
 & \frac{1}{4}(3A + 4C) \int \cos^2(c+dx) dx + \frac{A \sin(c+dx) \cos^3(c+dx)}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4}(3A + 4C) \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{A \sin(c+dx) \cos^3(c+dx)}{4d} \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{4}(3A + 4C) \left(\int \frac{1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{A \sin(c+dx) \cos^3(c+dx)}{4d} \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{4}(3A + 4C) \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) + \frac{A \sin(c+dx) \cos^3(c+dx)}{4d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^4*(A + C*Sec[c + d*x]^2), x]`

output `(A*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + ((3*A + 4*C)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/4`

3.11.3.1 Defintions of rubi rules used

rule 244 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.)^(m_.)*(csc[(e_.) + (f_.)*(x_)^2*(C_.) + (A_.)], x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

3.11.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.72

method	result
parallelrisch	$\frac{(8A+8C) \sin(2dx+2c)+A \sin(4dx+4c)+12d \left(A+\frac{4C}{3}\right) x}{32d}$
risch	$\frac{3Ax}{8} + \frac{Cx}{2} + \frac{A \sin(4dx+4c)}{32d} + \frac{A \sin(2dx+2c)}{4d} + \frac{\sin(2dx+2c)C}{4d}$
derivativedivides	$A \left(\frac{\left(\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + C \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
default	$A \left(\frac{\left(\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + C \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
norman	$\frac{\left(-\frac{3A}{8}-\frac{C}{2}\right)x + \left(-\frac{9A}{8}-\frac{3C}{2}\right)x \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2 + \left(-\frac{3A}{4}-C\right)x \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^4 + \left(\frac{3A}{4}+C\right)x \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^6 + \left(\frac{3A}{8}+\frac{C}{2}\right)x \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^8}{(1+\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2)^4}$

input `int(cos(d*x+c)^4*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

3.11. $\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx$

output $1/32*((8*A+8*C)*\sin(2*d*x+2*c)+A*\sin(4*d*x+4*c)+12*d*(A+4/3*C)*x)/d$

3.11.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.80

$$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(3A + 4C)dx + (2A \cos(dx + c))^3 + (3A + 4C) \cos(dx + c) \sin(dx + c)}{8d}$$

input `integrate(cos(d*x+c)^4*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output $1/8*((3*A + 4*C)*d*x + (2*A*\cos(d*x + c))^3 + (3*A + 4*C)*\cos(d*x + c))*\sin(d*x + c)/d$

3.11.6 Sympy [F]

$$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \cos^4(c + dx) dx$$

input `integrate(cos(d*x+c)**4*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*cos(c + d*x)**4, x)`

3.11.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.20

$$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(dx + c)(3A + 4C) + \frac{(3A+4C) \tan(dx+c)^3 + (5A+4C) \tan(dx+c)}{\tan(dx+c)^4 + 2 \tan(dx+c)^2 + 1}}{8d}$$

input `integrate(cos(d*x+c)^4*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/8*((d*x + c)*(3*A + 4*C) + ((3*A + 4*C)*tan(d*x + c)^3 + (5*A + 4*C)*tan(d*x + c)))/(tan(d*x + c)^4 + 2*tan(d*x + c)^2 + 1))/d`

3.11.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.20

$$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(dx + c)(3A + 4C) + \frac{3A \tan(dx+c)^3 + 4C \tan(dx+c)^3 + 5A \tan(dx+c) + 4C \tan(dx+c)}{(\tan(dx+c)^2 + 1)^2}}{8d}$$

input `integrate(cos(d*x+c)^4*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/8*((d*x + c)*(3*A + 4*C) + (3*A*tan(d*x + c)^3 + 4*C*tan(d*x + c)^3 + 5*A*tan(d*x + c) + 4*C*tan(d*x + c)))/(tan(d*x + c)^2 + 1)^2)/d`

3.11.9 Mupad [B] (verification not implemented)

Time = 16.43 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.10

$$\int \cos^4(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= x \left(\frac{3A}{8} + \frac{C}{2} \right) + \frac{\left(\frac{3A}{8} + \frac{C}{2} \right) \tan(c + dx)^3 + \left(\frac{5A}{8} + \frac{C}{2} \right) \tan(c + dx)}{d (\tan(c + dx)^4 + 2 \tan(c + dx)^2 + 1)}$$

input `int(cos(c + d*x)^4*(A + C/cos(c + d*x)^2),x)`

output `x*((3*A)/8 + C/2) + (tan(c + d*x)*((5*A)/8 + C/2) + tan(c + d*x)^3*((3*A)/8 + C/2))/(d*(2*tan(c + d*x)^2 + tan(c + d*x)^4 + 1))`

3.12 $\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx$

3.12.1	Optimal result	104
3.12.2	Mathematica [A] (verified)	104
3.12.3	Rubi [A] (verified)	105
3.12.4	Maple [A] (verified)	106
3.12.5	Fricas [A] (verification not implemented)	107
3.12.6	Sympy [F]	107
3.12.7	Maxima [A] (verification not implemented)	108
3.12.8	Giac [A] (verification not implemented)	108
3.12.9	Mupad [B] (verification not implemented)	108

3.12.1 Optimal result

Integrand size = 21, antiderivative size = 50

$$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx = \frac{(A + C) \sin(c + dx)}{d} - \frac{(2A + C) \sin^3(c + dx)}{3d} + \frac{A \sin^5(c + dx)}{5d}$$

output $(A+C)*\sin(d*x+c)/d-1/3*(2*A+C)*\sin(d*x+c)^3/d+1/5*A*\sin(d*x+c)^5/d$

3.12.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.42

$$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx = \frac{A \sin(c + dx)}{d} + \frac{C \sin(c + dx)}{d} - \frac{2A \sin^3(c + dx)}{3d} - \frac{C \sin^3(c + dx)}{3d} + \frac{A \sin^5(c + dx)}{5d}$$

input $\text{Integrate}[\text{Cos}[c + d*x]^5*(A + C*\text{Sec}[c + d*x]^2),x]$

output $(A*\text{Sin}[c + d*x])/d + (C*\text{Sin}[c + d*x])/d - (2*A*\text{Sin}[c + d*x]^3)/(3*d) - (C*\text{Sin}[c + d*x]^3)/(3*d) + (A*\text{Sin}[c + d*x]^5)/(5*d)$

3.12.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4532, 3042, 3492, 290, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^5(c+dx) (A + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c+dx + \frac{\pi}{2})^2}{\csc(c+dx + \frac{\pi}{2})^5} dx \\
 & \quad \downarrow \text{4532} \\
 & \int \cos^3(c+dx) (A \cos^2(c+dx) + C) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(c+dx + \frac{\pi}{2})^3 \left(A \sin(c+dx + \frac{\pi}{2})^2 + C \right) dx \\
 & \quad \downarrow \text{3492} \\
 & \frac{\int (1 - \sin^2(c+dx)) (-A \sin^2(c+dx) + A + C) d(-\sin(c+dx))}{d} \\
 & \quad \downarrow \text{290} \\
 & \frac{\int (A \sin^4(c+dx) - (2A + C) \sin^2(c+dx) + A(\frac{C}{A} + 1)) d(-\sin(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\frac{1}{3}(2A + C) \sin^3(c+dx) - (A + C) \sin(c+dx) - \frac{1}{5} A \sin^5(c+dx)}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^5*(A + C*Sec[c + d*x]^2),x]`

output `-((-((A + C)*Sin[c + d*x]) + ((2*A + C)*Sin[c + d*x]^3)/3 - (A*Ssin[c + d*x]^5)/5)/d)`

3.12.3.1 Defintions of rubi rules used

- rule 290 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]`

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 3492 `Int[sin[(e_) + (f_)*(x_)]^(m_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[-f^(-1) Subst[Int[(1 - x^2)^((m - 1)/2)*(A + C - C*x^2), x], x, Cos[e + f*x]], x] /; FreeQ[{e, f, A, C}, x] && IGtQ[(m + 1)/2, 0]`

- rule 4532 `Int[csc[(e_) + (f_)*(x_)]^(m_)*(csc[(e_) + (f_)*(x_)]^2*(C_) + (A_)), x_Symbol] := Int[(C + A*Sin[e + f*x]^2)/Sin[e + f*x]^(m + 2), x] /; FreeQ[{e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && ILtQ[(m + 1)/2, 0]`

3.12.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.98

method	result
parallelrisc	$\frac{(25A+20C) \sin(3dx+3c)+3A \sin(5dx+5c)+150 \sin(dx+c) \left(A+\frac{6C}{5}\right)}{240d}$
derivativedivides	$\frac{A \left(\frac{8}{3}+\cos(dx+c)\right)^4+\frac{4 \cos(dx+c)^2}{3}}{5} \frac{\sin(dx+c)}{d} + \frac{C(2+\cos(dx+c)^2) \sin(dx+c)}{3}$
default	$\frac{A \left(\frac{8}{3}+\cos(dx+c)\right)^4+\frac{4 \cos(dx+c)^2}{3}}{5} \frac{\sin(dx+c)}{d} + \frac{C(2+\cos(dx+c)^2) \sin(dx+c)}{3}$
risc	$\frac{5A \sin(dx+c)}{8d} + \frac{3C \sin(dx+c)}{4d} + \frac{A \sin(5dx+5c)}{80d} + \frac{5A \sin(3dx+3c)}{48d} + \frac{\sin(3dx+3c)C}{12d}$
norman	$\frac{-\frac{2(A+C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d} + \frac{2(A+C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^{11}}{d} - \frac{2(A+5C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{3d} + \frac{2(A+5C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^9}{3d} - \frac{4(19A+5C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^5}{15d}}{\left(1+\tan\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^5 \left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-1\right)}$

3.12. $\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx$

input `int(cos(d*x+c)^5*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/240*((25*A+20*C)*sin(3*d*x+3*c)+3*A*sin(5*d*x+5*c)+150*sin(d*x+c)*(A+6/5*C))/d`

3.12.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.90

$$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{(3 A \cos(dx + c)^4 + (4 A + 5 C) \cos(dx + c)^2 + 8 A + 10 C) \sin(dx + c)}{15 d}$$

input `integrate(cos(d*x+c)^5*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/15*(3*A*cos(d*x + c)^4 + (4*A + 5*C)*cos(d*x + c)^2 + 8*A + 10*C)*sin(d*x + c)/d`

3.12.6 Sympy [F]

$$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx = \int (A + C \sec^2(c + dx)) \cos^5(c + dx) dx$$

input `integrate(cos(d*x+c)**5*(A+C*sec(d*x+c)**2),x)`

output `Integral((A + C*sec(c + d*x)**2)*cos(c + d*x)**5, x)`

3.12.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.86

$$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3 A \sin(dx + c)^5 - 5(2 A + C) \sin(dx + c)^3 + 15(A + C) \sin(dx + c)}{15 d}$$

input `integrate(cos(d*x+c)^5*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/15*(3*A*sin(d*x + c)^5 - 5*(2*A + C)*sin(d*x + c)^3 + 15*(A + C)*sin(d*x + c))/d`

3.12.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.14

$$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3 A \sin(dx + c)^5 - 10 A \sin(dx + c)^3 - 5 C \sin(dx + c)^3 + 15 A \sin(dx + c) + 15 C \sin(dx + c)}{15 d}$$

input `integrate(cos(d*x+c)^5*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/15*(3*A*sin(d*x + c)^5 - 10*A*sin(d*x + c)^3 - 5*C*sin(d*x + c)^3 + 15*A*sin(d*x + c) + 15*C*sin(d*x + c))/d`

3.12.9 Mupad [B] (verification not implemented)

Time = 15.99 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.86

$$\int \cos^5(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{\frac{A \sin(c+dx)^5}{5} + \left(-\frac{2A}{3} - \frac{C}{3}\right) \sin(c + dx)^3 + (A + C) \sin(c + dx)}{d}$$

input `int(cos(c + d*x)^5*(A + C/cos(c + d*x)^2),x)`

output `((A*sin(c + d*x)^5)/5 + sin(c + d*x)*(A + C) - sin(c + d*x)^3*((2*A)/3 + C/3))/d`

3.13 $\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx$

3.13.1	Optimal result	110
3.13.2	Mathematica [A] (verified)	110
3.13.3	Rubi [A] (verified)	111
3.13.4	Maple [A] (verified)	112
3.13.5	Fricas [A] (verification not implemented)	113
3.13.6	Sympy [F(-1)]	114
3.13.7	Maxima [A] (verification not implemented)	114
3.13.8	Giac [A] (verification not implemented)	114
3.13.9	Mupad [B] (verification not implemented)	115

3.13.1 Optimal result

Integrand size = 21, antiderivative size = 89

$$\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx = \frac{1}{16}(5A + 6C)x + \frac{(5A + 6C) \cos(c + dx) \sin(c + dx)}{16d} + \frac{(5A + 6C) \cos^3(c + dx) \sin(c + dx)}{24d} + \frac{A \cos^5(c + dx) \sin(c + dx)}{6d}$$

```
output 1/16*(5*A+6*C)*x+1/16*(5*A+6*C)*cos(d*x+c)*sin(d*x+c)/d+1/24*(5*A+6*C)*cos
(d*x+c)^3*sin(d*x+c)/d+1/6*A*cos(d*x+c)^5*sin(d*x+c)/d
```

3.13.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.76

$$\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx = \frac{60Ac + 72cC + 60A dx + 72C dx + (45A + 48C) \sin(2(c + dx)) + (9A + 6C) \sin(4(c + dx)) + A \sin(6(c + dx))}{192d}$$

```
input Integrate[Cos[c + d*x]^6*(A + C*Sec[c + d*x]^2),x]
```

```
output (60*A*c + 72*c*C + 60*A*d*x + 72*C*d*x + (45*A + 48*C)*Sin[2*(c + d*x)] +
(9*A + 6*C)*Sin[4*(c + d*x)] + A*SIN[6*(c + d*x)]/(192*d)
```

3.13.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4533, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^6(c+dx) (A + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c+dx + \frac{\pi}{2})^2}{\csc(c+dx + \frac{\pi}{2})^6} dx \\
 & \quad \downarrow \text{4533} \\
 & \frac{1}{6}(5A + 6C) \int \cos^4(c+dx) dx + \frac{A \sin(c+dx) \cos^5(c+dx)}{6d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6}(5A + 6C) \int \sin(c+dx + \frac{\pi}{2})^4 dx + \frac{A \sin(c+dx) \cos^5(c+dx)}{6d} \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{6}(5A + 6C) \left(\frac{3}{4} \int \cos^2(c+dx) dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) + \frac{A \sin(c+dx) \cos^5(c+dx)}{6d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6}(5A + 6C) \left(\frac{3}{4} \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) + \frac{A \sin(c+dx) \cos^5(c+dx)}{6d} \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{6}(5A + 6C) \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) + \frac{A \sin(c+dx) \cos^5(c+dx)}{6d} \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{6}(5A + 6C) \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right) + \frac{A \sin(c+dx) \cos^5(c+dx)}{6d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^6*(A + C*Sec[c + d*x]^2),x]`

output `(A*Cos[c + d*x]^5*Sin[c + d*x])/(6*d) + ((5*A + 6*C)*((Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (3*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d))/4))/6`

3.13.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.)^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

3.13.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.69

method	result
parallelrisch	$\frac{(45A+48C)\sin(2dx+2c)+(9A+6C)\sin(4dx+4c)+A\sin(6dx+6c)+60dx\left(A+\frac{6C}{5}\right)}{192d}$
risch	$\frac{5Ax}{16} + \frac{3Cx}{8} + \frac{A\sin(6dx+6c)}{192d} + \frac{3A\sin(4dx+4c)}{64d} + \frac{\sin(4dx+4c)C}{32d} + \frac{15A\sin(2dx+2c)}{64d} + \frac{\sin(2dx+2c)C}{4d}$
derivativedivides	$A\left(\frac{\left(\cos(dx+c)^5 + \frac{5\cos(dx+c)^3}{4} + \frac{15\cos(dx+c)}{8}\right)\sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16}\right) + C\left(\frac{\left(\cos(dx+c)^3 + \frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right)$
default	$A\left(\frac{\left(\cos(dx+c)^5 + \frac{5\cos(dx+c)^3}{4} + \frac{15\cos(dx+c)}{8}\right)\sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16}\right) + C\left(\frac{\left(\cos(dx+c)^3 + \frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right)$
norman	$\left(-\frac{5A}{16} - \frac{3C}{8}\right)x + \left(-\frac{45A}{16} - \frac{27C}{8}\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + \left(-\frac{25A}{16} - \frac{15C}{8}\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \left(-\frac{25A}{16} - \frac{15C}{8}\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + \left(\frac{5A}{16} + \frac{3C}{8}\right)x$

input `int(cos(d*x+c)^6*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/192*((45*A+48*C)*sin(2*d*x+2*c)+(9*A+6*C)*sin(4*d*x+4*c)+A*sin(6*d*x+6*c)+60*d*x*(A+6/5*C))/d`

3.13.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.76

$$\int \cos^6(c+dx)(A+C\sec^2(c+dx))dx$$

$$= \frac{3(5A+6C)dx + (8A\cos(dx+c)^5 + 2(5A+6C)\cos(dx+c)^3 + 3(5A+6C)\cos(dx+c))\sin(dx+c)}{48d}$$

input `integrate(cos(d*x+c)^6*(A+C*sec(d*x+c)^2),x,algorithm="fricas")`

output `1/48*(3*(5*A+6*C)*d*x+(8*A*cos(d*x+c)^5+2*(5*A+6*C)*cos(d*x+c)^3+3*(5*A+6*C)*cos(d*x+c))*sin(d*x+c)/d`

3.13.6 Sympy [F(-1)]

Timed out.

$$\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**6*(A+C*sec(d*x+c)**2),x)`output `Timed out`**3.13.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.16

$$\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3(dx+c)(5A+6C) + \frac{3(5A+6C)\tan(dx+c)^5 + 8(5A+6C)\tan(dx+c)^3 + 3(11A+10C)\tan(dx+c)}{\tan(dx+c)^6 + 3\tan(dx+c)^4 + 3\tan(dx+c)^2 + 1}}{48d}$$

input `integrate(cos(d*x+c)^6*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`output `1/48*(3*(d*x + c)*(5*A + 6*C) + (3*(5*A + 6*C)*tan(d*x + c)^5 + 8*(5*A + 6*C)*tan(d*x + c)^3 + 3*(11*A + 10*C)*tan(d*x + c)))/(tan(d*x + c)^6 + 3*tan(d*x + c)^4 + 3*tan(d*x + c)^2 + 1))/d`**3.13.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.08

$$\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= \frac{3(dx+c)(5A+6C) + \frac{15A\tan(dx+c)^5 + 18C\tan(dx+c)^5 + 40A\tan(dx+c)^3 + 48C\tan(dx+c)^3 + 33A\tan(dx+c) + 30C\tan(dx+c)}{(\tan(dx+c)^2 + 1)^3}}{48d}$$

input `integrate(cos(d*x+c)^6*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output $\frac{1}{48}(3(dx + c)(5A + 6C) + (15A \tan(dx + c)^5 + 18C \tan(dx + c)^5 + 40A \tan(dx + c)^3 + 48C \tan(dx + c)^3 + 33A \tan(dx + c) + 30C \tan(dx + c)) / (\tan(dx + c)^2 + 1)^3) / d$

3.13.9 Mupad [B] (verification not implemented)

Time = 16.14 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.02

$$\int \cos^6(c + dx) (A + C \sec^2(c + dx)) dx$$

$$= x \left(\frac{5A}{16} + \frac{3C}{8} \right) + \frac{\left(\frac{5A}{16} + \frac{3C}{8} \right) \tan(c + dx)^5 + \left(\frac{5A}{6} + C \right) \tan(c + dx)^3 + \left(\frac{11A}{16} + \frac{5C}{8} \right) \tan(c + dx)}{d (\tan(c + dx)^6 + 3 \tan(c + dx)^4 + 3 \tan(c + dx)^2 + 1)}$$

input `int(cos(c + d*x)^6*(A + C/cos(c + d*x)^2),x)`

output `x*((5*A)/16 + (3*C)/8) + (tan(c + d*x)*((11*A)/16 + (5*C)/8) + tan(c + d*x)^3*((5*A)/6 + C) + tan(c + d*x)^5*((5*A)/16 + (3*C)/8))/(d*(3*tan(c + d*x)^2 + 3*tan(c + d*x)^4 + tan(c + d*x)^6 + 1))`

3.14 $\int \sec^m(c + dx) \left(-\frac{Cm}{1+m} + C \sec^2(c + dx)\right) dx$

3.14.1	Optimal result	116
3.14.2	Mathematica [C] (verified)	116
3.14.3	Rubi [A] (verified)	117
3.14.4	Maple [A] (verified)	118
3.14.5	Fricas [A] (verification not implemented)	118
3.14.6	Sympy [F]	118
3.14.7	Maxima [B] (verification not implemented)	119
3.14.8	Giac [F]	119
3.14.9	Mupad [B] (verification not implemented)	120

3.14.1 Optimal result

Integrand size = 29, antiderivative size = 26

$$\int \sec^m(c + dx) \left(-\frac{Cm}{1+m} + C \sec^2(c + dx)\right) dx = \frac{C \sec^{1+m}(c + dx) \sin(c + dx)}{d(1+m)}$$

output `C*sec(d*x+c)^(1+m)*sin(d*x+c)/d/(1+m)`

3.14.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.44 (sec) , antiderivative size = 113, normalized size of antiderivative = 4.35

$$\int \sec^m(c + dx) \left(-\frac{Cm}{1+m} + C \sec^2(c + dx)\right) dx = \frac{C \csc(c + dx) \sec^{-1+m}(c + dx) \left(-((2+m) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, \sec^2(c + dx)\right)) + (1+m) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, \sec^2(c + dx)\right)\right)}{d(1+m)(2+m)}$$

input `Integrate[Sec[c + d*x]^m*(-((C*m)/(1+m)) + C*Sec[c + d*x]^2),x]`

output `(C*Csc[c + d*x]*Sec[c + d*x]^(-1+m)*(-((2+m)*Hypergeometric2F1[1/2, m/2, (2+m)/2, Sec[c + d*x]^2]) + (1+m)*Hypergeometric2F1[1/2, (2+m)/2, (4+m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^2)*Sqrt[-Tan[c + d*x]^2])/(d*(1+m)*(2+m))`

3.14.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^m(c + dx) \left(C \sec^2(c + dx) - \frac{Cm}{m+1} \right) dx$$

↓ 3042

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^m \left(C \csc\left(c + dx + \frac{\pi}{2}\right)^2 - \frac{Cm}{m+1} \right) dx$$

↓ 4531

$$\frac{C \sin(c + dx) \sec^{m+1}(c + dx)}{d(m+1)}$$

input `Int[Sec[c + d*x]^m*(-((C*m)/(1 + m)) + C*Sec[c + d*x]^2),x]`

output `(C*Sec[c + d*x]^(1 + m)*Sin[c + d*x])/(d*(1 + m))`

3.14.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m]*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] :> Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.14. $\int \sec^m(c + dx) \left(-\frac{Cm}{1+m} + C \sec^2(c + dx) \right) dx$

3.14.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.77

method	result
parallelrisch	$-\frac{2C \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\frac{1}{\cos(dx+c)}\right)^m}{(1+m)d \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$
risch	$-\frac{iC(e^{i(dx+c)})^m (e^{2i(dx+c)}+1)^{-m} 2^m \left(e^{\frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2i(dx+c)}+1}\right) \operatorname{csgn}\left(\frac{ie^{i(dx+c)}}{e^{2i(dx+c)}+1}\right)^2}{2}} e^{-\frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2i(dx+c)}+1}\right) \operatorname{csgn}\left(\frac{ie^{i(dx+c)}}{e^{2i(dx+c)}+1}\right)}{2}} \right)}{1}$

input `int(sec(d*x+c)^m*(-C*m/(1+m)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `-2*C/(1+m)/d*tan(1/2*d*x+1/2*c)*(1/cos(d*x+c))^m/(tan(1/2*d*x+1/2*c)^2-1)`

3.14.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.27

$$\int \sec^m(c+dx) \left(-\frac{Cm}{1+m} + C \sec^2(c+dx) \right) dx = \frac{C \frac{1}{\cos(dx+c)} \sin(dx+c)^m}{(dm+d) \cos(dx+c)}$$

input `integrate(sec(d*x+c)^m*(-C*m/(1+m)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `C*(1/cos(d*x + c))^m*sin(d*x + c)/((d*m + d)*cos(d*x + c))`

3.14.6 Sympy [F]

$$\int \sec^m(c+dx) \left(-\frac{Cm}{1+m} + C \sec^2(c+dx) \right) dx$$

$$= \frac{C \left(\int (-m \sec^m(c+dx)) dx + \int \sec^2(c+dx) \sec^m(c+dx) dx + \int m \sec^2(c+dx) \sec^m(c+dx) dx \right)}{m+1}$$

input `integrate(sec(d*x+c)**m*(-C*m/(1+m)+C*sec(d*x+c)**2),x)`

3.14. $\int \sec^m(c+dx) \left(-\frac{Cm}{1+m} + C \sec^2(c+dx) \right) dx$

3.14.9 Mupad [B] (verification not implemented)

Time = 15.61 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.62

$$\int \sec^m(c + dx) \left(-\frac{Cm}{1+m} + C \sec^2(c + dx) \right) dx = \frac{C \sin(2c + 2dx) \left(\frac{1}{\cos(c+dx)} \right)^m}{d (\cos(2c + 2dx) + 1) (m + 1)}$$

input `int((1/cos(c + d*x))^m*(C/cos(c + d*x)^2 - (C*m)/(m + 1)),x)`

output `(C*sin(2*c + 2*d*x)*(1/cos(c + d*x))^m)/(d*(cos(2*c + 2*d*x) + 1)*(m + 1))`

$$3.15 \quad \int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m} \right) dx$$

3.15.1	Optimal result	121
3.15.2	Mathematica [C] (verified)	121
3.15.3	Rubi [A] (verified)	122
3.15.4	Maple [A] (verified)	123
3.15.5	Fricas [A] (verification not implemented)	123
3.15.6	Sympy [F]	123
3.15.7	Maxima [B] (verification not implemented)	124
3.15.8	Giac [F]	124
3.15.9	Mupad [B] (verification not implemented)	125

3.15.1 Optimal result

Integrand size = 28, antiderivative size = 25

$$\int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c + dx)}{m} \right) dx = -\frac{A \sec^{1+m}(c + dx) \sin(c + dx)}{dm}$$

output `-A*sec(d*x+c)^(1+m)*sin(d*x+c)/d/m`

3.15.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.30 (sec) , antiderivative size = 111, normalized size of antiderivative = 4.44

$$\int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c + dx)}{m} \right) dx = \frac{A \csc(c + dx) \sec^{-1+m}(c + dx) ((2 + m) \text{Hypergeometric2F1}(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, \sec^2(c + dx)) - (1 + m) \text{Hypergeometric2F1}(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, \sec^2(c + dx)))}{dm(2 + m)}$$

input `Integrate[Sec[c + d*x]^m*(A - (A*(1 + m)*Sec[c + d*x]^2)/m), x]`

output `(A*Csc[c + d*x]*Sec[c + d*x]^(-1 + m)*((2 + m)*Hypergeometric2F1[1/2, m/2, (2 + m)/2, Sec[c + d*x]^2] - (1 + m)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^2)*Sqrt[-Tan[c + d*x]^2])/(d*m*(2 + m))`

$$3.15. \quad \int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m} \right) dx$$

3.15.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^m(c+dx) \left(A - \frac{A(m+1)\sec^2(c+dx)}{m} \right) dx$$

↓ 3042

$$\int \csc\left(c+dx+\frac{\pi}{2}\right)^m \left(A - \frac{A(m+1)\csc\left(c+dx+\frac{\pi}{2}\right)^2}{m} \right) dx$$

↓ 4531

$$-\frac{A \sin(c+dx) \sec^{m+1}(c+dx)}{dm}$$

input `Int[Sec[c + d*x]^m*(A - (A*(1 + m)*Sec[c + d*x]^2)/m), x]`

output `-((A*Sec[c + d*x]^(1 + m)*Sin[c + d*x])/(d*m))`

3.15.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m]*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.15. $\int \sec^m(c+dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m} \right) dx$

3.15.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.76

method	result
parallelrisch	$\frac{2A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\frac{1}{\cos(dx+c)}\right)^m}{md \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$
risch	$iA(e^{i(dx+c)})^m (e^{2i(dx+c)}+1)^{-m} 2^m \left(e^{\frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2i(dx+c)}+1}\right) \operatorname{csgn}\left(\frac{ie^{i(dx+c)}}{e^{2i(dx+c)}+1}\right)^2}{2} m} e^{-\frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2i(dx+c)}+1}\right) \operatorname{csgn}\left(\frac{ie^{i(dx+c)}}{e^{2i(dx+c)}+1}\right)}{2}}$

input `int(sec(d*x+c)^m*(A-A*(1+m)*sec(d*x+c)^2/m),x,method=_RETURNVERBOSE)`

output `2*A/m/d*tan(1/2*d*x+1/2*c)*(1/cos(d*x+c))^m/(tan(1/2*d*x+1/2*c)^2-1)`

3.15.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.32

$$\int \sec^m(c+dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m} \right) dx = -\frac{A \frac{1}{\cos(dx+c)} \sin(dx+c)}{dm \cos(dx+c)}$$

input `integrate(sec(d*x+c)^m*(A-A*(1+m)*sec(d*x+c)^2/m),x, algorithm="fricas")`

output `-A*(1/cos(d*x + c))^m*sin(d*x + c)/(d*m*cos(d*x + c))`

3.15.6 Sympy [F]

$$\int \sec^m(c+dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m} \right) dx = \frac{A \left(\int (-m \sec^m(c+dx)) dx + \int \sec^2(c+dx) \sec^m(c+dx) dx + \int m \sec^2(c+dx) \sec^m(c+dx) dx \right)}{m}$$

input `integrate(sec(d*x+c)**m*(A-A*(1+m)*sec(d*x+c)**2/m),x)`

3.15. $\int \sec^m(c+dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m} \right) dx$

output `-A*(Integral(-m*sec(c + d*x)**m, x) + Integral(sec(c + d*x)**2*sec(c + d*x)**m, x) + Integral(m*sec(c + d*x)**2*sec(c + d*x)**m, x))/m`

3.15.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 296 vs. $2(25) = 50$.

Time = 0.43 (sec) , antiderivative size = 296, normalized size of antiderivative = 11.84

$$\int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c + dx)}{m} \right) dx$$

$$= \frac{2^m A \cos(-(dx + c)(m + 2) + m \arctan(\sin(2dx + 2c), \cos(2dx + 2c) + 1)) \sin(2dx + 2c) - 2^m A \cos$$

input `integrate(sec(d*x+c)^m*(A-A*(1+m)*sec(d*x+c)^2/m),x, algorithm="maxima")`

output `(2^m*A*cos(-(d*x + c)*(m + 2) + m*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(2*d*x + 2*c) - 2^m*A*cos(-(d*x + c)*m + m*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(2*d*x + 2*c) + (2^m*A*cos(2*d*x + 2*c) + 2^m*A)*sin(-(d*x + c)*(m + 2) + m*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - (2^m*A*cos(2*d*x + 2*c) + 2^m*A)*sin(-(d*x + c)*m + m*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))/((m*cos(2*d*x + 2*c)^2 + m*sin(2*d*x + 2*c)^2 + 2*m*cos(2*d*x + 2*c) + m)*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/2*m)*d)`

3.15.8 Giac [F]

$$\int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c + dx)}{m} \right) dx$$

$$= \int - \left(\frac{A(m+1)\sec(dx+c)^2}{m} - A \right) \sec(dx+c)^m dx$$

input `integrate(sec(d*x+c)^m*(A-A*(1+m)*sec(d*x+c)^2/m),x, algorithm="giac")`

output `integrate(-(A*(m + 1)*sec(d*x + c)^2/m - A)*sec(d*x + c)^m, x)`

3.15. $\int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c+dx)}{m} \right) dx$

3.15.9 Mupad [B] (verification not implemented)

Time = 15.21 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int \sec^m(c + dx) \left(A - \frac{A(1+m)\sec^2(c + dx)}{m} \right) dx = -\frac{A \sin(2c + 2dx) \left(\frac{1}{\cos(c+dx)} \right)^m}{dm (\cos(2c + 2dx) + 1)}$$

input `int((A - (A*(m + 1))/(m*cos(c + d*x)^2))*(1/cos(c + d*x))^m,x)`

output `-(A*sin(2*c + 2*d*x)*(1/cos(c + d*x))^m)/(d*m*(cos(2*c + 2*d*x) + 1))`

3.16 $\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx$

3.16.1	Optimal result	126
3.16.2	Mathematica [A] (verified)	126
3.16.3	Rubi [A] (verified)	127
3.16.4	Maple [C] (verified)	129
3.16.5	Fricas [C] (verification not implemented)	130
3.16.6	Sympy [F]	130
3.16.7	Maxima [F]	130
3.16.8	Giac [F]	131
3.16.9	Mupad [F(-1)]	131

3.16.1 Optimal result

Integrand size = 25, antiderivative size = 110

$$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx = \frac{2b^2(7A + 5C)\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{2b(7A + 5C)(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} + \frac{2C(b \sec(c + dx))^{5/2} \tan(c + dx)}{7d}$$

output

```
2/21*b*(7*A+5*C)*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/d+2/21*b^2*(7*A+5*C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d+2/7*C*(b*sec(d*x+c))^(5/2)*tan(d*x+c)/d
```

3.16.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.76

$$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx = \frac{(b \sec(c + dx))^{7/2} \left(4(7A + 5C) \cos^{\frac{7}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 2(7A + 11C) \right)}{42bd}$$

input `Integrate[(b*Sec[c + d*x])^(5/2)*(A + C*Sec[c + d*x]^2),x]`

output `((b*Sec[c + d*x])^(7/2)*(4*(7*A + 5*C)*Cos[c + d*x]^(7/2)*EllipticF[(c + d*x)/2, 2] + 2*(7*A + 11*C + (7*A + 5*C)*Cos[2*(c + d*x)])*Sin[c + d*x]))/(42*b*d)`

3.16.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {3042, 4534, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^{5/2} \left(A + C \csc\left(c + dx + \frac{\pi}{2}\right)^2 \right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{7}(7A + 5C) \int (b \sec(c + dx))^{5/2} dx + \frac{2C \tan(c + dx)(b \sec(c + dx))^{5/2}}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{7}(7A + 5C) \int \left(b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^{5/2} dx + \frac{2C \tan(c + dx)(b \sec(c + dx))^{5/2}}{7d} \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{7}(7A + 5C) \left(\frac{1}{3}b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} \right) + \\
 & \quad \frac{2C \tan(c + dx)(b \sec(c + dx))^{5/2}}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{7}(7A + 5C) \left(\frac{1}{3}b^2 \int \sqrt{b \csc\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} \right) + \\
 & \quad \frac{2C \tan(c + dx)(b \sec(c + dx))^{5/2}}{7d}
 \end{aligned}$$

$$\begin{array}{c}
\downarrow 4258 \\
5C) \left(\frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right) + \\
\frac{2C \tan(c+dx) (b \sec(c+dx))^{5/2}}{7d} \\
\downarrow 3042 \\
5C) \left(\frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right) + \\
\frac{2C \tan(c+dx) (b \sec(c+dx))^{5/2}}{7d} \\
\downarrow 3120 \\
5C) \left(\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right) + \\
\frac{2C \tan(c+dx) (b \sec(c+dx))^{5/2}}{7d}
\end{array}$$

input `Int[(b*Sec[c + d*x])^(5/2)*(A + C*Sec[c + d*x]^2),x]`

output `((7*A + 5*C)*((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d))/7 + (2*C*(b*Sec[c + d*x])^(5/2)*Tan[c + d*x])/(7*d)`

3.16.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_. + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.16.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 22.69 (sec) , antiderivative size = 291, normalized size of antiderivative = 2.65

method	result
default	$\frac{2b^2 \sqrt{b \sec(dx+c)} \left(7iA \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \sqrt{\frac{1}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(\cot(dx+c)-\csc(dx+c)), i) \cos(dx+c) + 5iC \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \sqrt{\frac{1}{\cos(dx+c)+1}} \right)}{3d}$
parts	$\frac{2A \sqrt{b \sec(dx+c)} b^2 \left(i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(\cot(dx+c)-\csc(dx+c)), i) \cos(dx+c) + i \operatorname{EllipticF}(i(\cot(dx+c)-\csc(dx+c)), i) \cos(dx+c) \right)}{3d}$

input `int((b*sec(d*x+c))^(5/2)*(A+C*sec(d*x+c)^2), x, method=_RETURNVERBOSE)`

output `2/21*b^2/d*(b*sec(d*x+c))^(1/2)*(7*I*A*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)), I)*cos(d*x+c)+5*I*C*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)), I)*cos(d*x+c)+7*I*A*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)), I)+5*I*C*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)), I)+7*A*tan(d*x+c)+5*C*tan(d*x+c)+3*C*tan(d*x+c)*sec(d*x+c)^2)`

3.16.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.29

$$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx = \frac{-i \sqrt{2} (7A + 5C) b^{5/2} \cos(dx + c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} (7A + 5C) b^{5/2} \cos(dx + c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2((7A + 5C) b^2 \cos(dx + c)^2 + 3C b^2) \sqrt{b/\cos(dx + c)} \sin(dx + c)}{(d \cos(dx + c))^3}$$

input `integrate((b*sec(d*x+c))^(5/2)*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/21*(-I*sqrt(2)*(7*A + 5*C)*b^(5/2)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*(7*A + 5*C)*b^(5/2)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*((7*A + 5*C)*b^2*cos(d*x + c)^2 + 3*C*b^2)*sqrt(b/cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.16.6 Sympy [F]

$$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx = \int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx$$

input `integrate((b*sec(d*x+c))**(5/2)*(A+C*sec(d*x+c)**2),x)`

output `Integral((b*sec(c + d*x))**(5/2)*(A + C*sec(c + d*x)**2), x)`

3.16.7 Maxima [F]

$$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + A) (b \sec(dx + c))^{5/2} dx$$

input `integrate((b*sec(d*x+c))^(5/2)*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)*(b*sec(d*x + c))^(5/2), x)`

3.16.8 Giac [F]

$$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + A)(b \sec(dx + c))^{5/2} dx$$

input `integrate((b*sec(d*x+c))^(5/2)*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)*(b*sec(d*x + c))^(5/2), x)`

3.16.9 Mupad [F(-1)]

Timed out.

$$\int (b \sec(c + dx))^{5/2} (A + C \sec^2(c + dx)) dx = \int \left(A + \frac{C}{\cos(c + dx)^2} \right) \left(\frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

input `int((A + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(5/2),x)`

output `int((A + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(5/2), x)`

3.17 $\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx$

3.17.1	Optimal result	132
3.17.2	Mathematica [C] (verified)	132
3.17.3	Rubi [A] (verified)	133
3.17.4	Maple [C] (verified)	135
3.17.5	Fricas [C] (verification not implemented)	136
3.17.6	Sympy [F]	137
3.17.7	Maxima [F]	137
3.17.8	Giac [F]	137
3.17.9	Mupad [F(-1)]	138

3.17.1 Optimal result

Integrand size = 25, antiderivative size = 110

$$\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx = -\frac{2b^2(5A + 3C)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2b(5A + 3C)\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2C(b \sec(c + dx))^{3/2} \tan(c + dx)}{5d}$$

output

```
-2/5*b^2*(5*A+3*C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/5*b*(5*A+3*C)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+2/5*C*(b*sec(d*x+c))^(3/2)*tan(d*x+c)/d
```

3.17.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.20 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.67

$$\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx = \frac{4ie^{i(c+dx)} \cos^3(c + dx) \left(-3 \left(5A(1 + e^{2i(c+dx)})^2 + C(1 + 8e^{2i(c+dx)} + 3e^{4i(c+dx)}) \right) + (5A + C) \right)}{15d(1 + e^{2i(c+dx)})}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)*(A + C*Sec[c + d*x]^2),x]`

output `((((4*I)/15)*E^(I*(c + d*x))*Cos[c + d*x]^3*(-3*(5*A*(1 + E^((2*I)*(c + d*x))))^2 + C*(1 + 8*E^((2*I)*(c + d*x)) + 3*E^((4*I)*(c + d*x)))) + (5*A + 3*C)*(1 + E^((2*I)*(c + d*x)))^(5/2)*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])*(b*Sec[c + d*x])^(3/2)*(A + C*Sec[c + d*x]^2))/(d*(1 + E^((2*I)*(c + d*x)))^2*(A + 2*C + A*Cos[2*(c + d*x)]))`

3.17.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.95, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {3042, 4534, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^{3/2} \left(A + C \csc\left(c + dx + \frac{\pi}{2}\right)^2 \right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{5}(5A + 3C) \int (b \sec(c + dx))^{3/2} dx + \frac{2C \tan(c + dx)(b \sec(c + dx))^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5}(5A + 3C) \int \left(b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^{3/2} dx + \frac{2C \tan(c + dx)(b \sec(c + dx))^{3/2}}{5d} \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \right) + \\
 & \quad \frac{2C \tan(c + dx)(b \sec(c + dx))^{3/2}}{5d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2C \tan(c + dx) (b \sec(c + dx))^{3/2}}{5d}$$

↓ 4258

$$\frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) + \frac{2C \tan(c + dx) (b \sec(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) + \frac{2C \tan(c + dx) (b \sec(c + dx))^{3/2}}{5d}$$

↓ 3119

$$\frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) + \frac{2C \tan(c + dx) (b \sec(c + dx))^{3/2}}{5d}$$

input `Int[(b*Sec[c + d*x])^(3/2)*(A + C*Sec[c + d*x]^2),x]`

output `((5*A + 3*C)*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5 + (2*C*(b*Sec[c + d*x])^(3/2)*Tan[c + d*x])/(5*d)`

3.17.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 4255 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
  && IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
  EqQ[n^2, 1/4]
```

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.)
  + (A_)), x_Symbol] :> Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1)
  )), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
  FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

3.17.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 5.33 (sec) , antiderivative size = 798, normalized size of antiderivative = 7.25

method	result	size
default	Expression too large to display	798
parts	Expression too large to display	809

```
input int((b*sec(d*x+c))^(3/2)*(A+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)
```


output

```

-2/5*b/d*(b*sec(d*x+c))^(1/2)/(cos(d*x+c)+1)*(-3*I*C*EllipticE(I*(cot(d*x+
c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/
2)*cos(d*x+c)^2+6*I*C*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c
)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+5*I*A*(1/(cos(d*x
+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-cs
c(d*x+c)),I)+5*I*A*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1
))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2+3*I*C*EllipticF(I*
(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c
)+1))^(1/2)*cos(d*x+c)^2-10*I*A*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/
(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+10*I*A*
EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c
)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+3*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x
+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)-5*I*A*Ell
ipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(
cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2-5*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+
c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-6*I*C*Ell
ipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(c
os(d*x+c)+1))^(1/2)*cos(d*x+c)-3*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/
(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-5*A*sin(d*x+c
)-3*C*sin(d*x+c)-C*tan(d*x+c)-C*sec(d*x+c)*tan(d*x+c)

```

3.17.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.30

$$\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx = \frac{-i \sqrt{2} (5A + 3C) b^{3/2} \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + I \sin(dx + c))) + I \sqrt{2} (5A + 3C) b^{3/2} \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - I \sin(dx + c))) + 2((5A + 3C) b \cos(dx + c)^2 + C b) \sqrt{b/\cos(dx + c)} \sin(dx + c)}{(d \cos(dx + c))^2}$$

input `integrate((b*sec(d*x+c))^(3/2)*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output

```

1/5*(-I*sqrt(2)*(5*A + 3*C)*b^(3/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0,
weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*(5*
A + 3*C)*b^(3/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse
(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*((5*A + 3*C)*b*cos(d*x + c)^2
+ C*b)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^2)

```

3.17. $\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx$

3.17.6 Sympy [F]

$$\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx = \int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx$$

input `integrate((b*sec(d*x+c))**(3/2)*(A+C*sec(d*x+c)**2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*(A + C*sec(c + d*x)**2), x)`

3.17.7 Maxima [F]

$$\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + A)(b \sec(dx + c))^{3/2} dx$$

input `integrate((b*sec(d*x+c))^(3/2)*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)*(b*sec(d*x + c))^(3/2), x)`

3.17.8 Giac [F]

$$\int (b \sec(c + dx))^{3/2} (A + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + A)(b \sec(dx + c))^{3/2} dx$$

input `integrate((b*sec(d*x+c))^(3/2)*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)*(b*sec(d*x + c))^(3/2), x)`

3.17.9 Mupad [F(-1)]

Timed out.

$$\int (b \sec(c+dx))^{3/2} (A+C \sec^2(c+dx)) dx = \int \left(A + \frac{C}{\cos(c+dx)^2} \right) \left(\frac{b}{\cos(c+dx)} \right)^{3/2} dx$$

input `int((A + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(3/2),x)`output `int((A + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(3/2), x)`

3.18 $\int \sqrt{b \sec(c + dx)}(A + C \sec^2(c + dx)) dx$

3.18.1	Optimal result	139
3.18.2	Mathematica [A] (verified)	139
3.18.3	Rubi [A] (verified)	140
3.18.4	Maple [C] (verified)	141
3.18.5	Fricas [C] (verification not implemented)	142
3.18.6	Sympy [F]	142
3.18.7	Maxima [F]	143
3.18.8	Giac [F]	143
3.18.9	Mupad [F(-1)]	143

3.18.1 Optimal result

Integrand size = 25, antiderivative size = 72

$$\begin{aligned} & \int \sqrt{b \sec(c + dx)}(A + C \sec^2(c + dx)) dx \\ &= \frac{2(3A + C) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} \\ & \quad + \frac{2C \sqrt{b \sec(c + dx)} \tan(c + dx)}{3d} \end{aligned}$$

output `2/3*(3*A+C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d+2/3*C*(b*sec(d*x+c))^(1/2)*tan(d*x+c)/d`

3.18.2 Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int \sqrt{b \sec(c + dx)}(A + C \sec^2(c + dx)) dx \\ &= \frac{2(b \sec(c + dx))^{3/2} \left((3A + C) \cos^{\frac{3}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + C \sin(c + dx) \right)}{3bd} \end{aligned}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]*(A + C*Sec[c + d*x]^2),x]`

output $(2*(b*\text{Sec}[c + d*x])^{(3/2)}*((3*A + C)*\text{Cos}[c + d*x]^{(3/2)}*\text{EllipticF}[(c + d*x)/2, 2] + C*\text{Sin}[c + d*x]))/(3*b*d)$

3.18.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {3042, 4534, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{b \sec(c + dx)} (A + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{b \csc\left(c + dx + \frac{\pi}{2}\right)} \left(A + C \csc\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{3}(3A + C) \int \sqrt{b \sec(c + dx)} dx + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3}(3A + C) \int \sqrt{b \csc\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
 & \quad \downarrow \text{4258} \\
 & \frac{1}{3}(3A + C) \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3}(3A + C) \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2(3A + C) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d}
 \end{aligned}$$

input $\text{Int}[\text{Sqrt}[b*\text{Sec}[c + d*x]]*(A + C*\text{Sec}[c + d*x]^2), x]$

3.18. $\int \sqrt{b \sec(c + dx)} (A + C \sec^2(c + dx)) dx$

output $(2*(3*A + C)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*C*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Tan}[c + d*x])/(3*d)$

3.18.3.1 Defintions of rubi rules used

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\text{sin}[c_.] + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] \text{ ; FreeQ}\{c, d\}, x]$

rule 4258 $\text{Int}[(\text{csc}[c_.] + (d_.)*(x_.))*(b_.))^n, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{ Int}[1/\text{Sin}[c + d*x]^n, x], x] \text{ ; FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

rule 4534 $\text{Int}[(\text{csc}[e_.] + (f_.)*(x_.))*(b_.))^m*(\text{csc}[e_.] + (f_.)*(x_.))^2*(C_. + A_.), x_Symbol] \rightarrow \text{Simp}[(-C)*\text{Cot}[e + f*x]*((b*\text{Csc}[e + f*x])^m/(f*(m + 1))), x] + \text{Simp}[(C*m + A*(m + 1))/(m + 1) \text{ Int}[(b*\text{Csc}[e + f*x])^m, x], x] \text{ ; FreeQ}\{b, e, f, A, C, m\}, x \ \&\& \ \text{NeQ}[C*m + A*(m + 1), 0] \ \&\& \ \text{!LeQ}[m, -1]$

3.18.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.14 (sec) , antiderivative size = 220, normalized size of antiderivative = 3.06

method	result
parts	$-\frac{2iA(\cos(dx+c)+1)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}{d}\text{EllipticF}(i(-\cot(dx+c)+\text{csc}(dx+c)),i)\sqrt{b\sec(dx+c)} - \frac{2C\sqrt{b\sec(dx+c)}}{d}\left(i\sqrt{\frac{1}{\cos(dx+c)+1}}\right)$
default	$\frac{2\sqrt{b\sec(dx+c)}}{d}\left(3iA\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\sqrt{\frac{1}{\cos(dx+c)+1}}\text{EllipticF}(i(\cot(dx+c)-\text{csc}(dx+c)),i)\cos(dx+c)+iC\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\sqrt{\frac{1}{\cos(dx+c)+1}}\right)$

input $\text{int}((b*\text{sec}(d*x+c))^(1/2)*(A+C*\text{sec}(d*x+c)^2), x, \text{method}=_RETURNVERBOSE)$

output $-2IA/d(\cos(dx+c)+1)*(1/(\cos(dx+c)+1))^{1/2}*(\cos(dx+c)/(\cos(dx+c)+1))^{1/2}*EllipticF(I*(-\cot(dx+c)+\csc(dx+c)),I)*(b*\sec(dx+c))^{1/2}-2/3*C/d*(b*\sec(dx+c))^{1/2}*(I*(1/(\cos(dx+c)+1))^{1/2}*(\cos(dx+c)/(\cos(dx+c)+1))^{1/2}*EllipticF(I*(-\cot(dx+c)+\csc(dx+c)),I)*\cos(dx+c)+I*(1/(\cos(dx+c)+1))^{1/2}*(\cos(dx+c)/(\cos(dx+c)+1))^{1/2}*EllipticF(I*(-\cot(dx+c)+\csc(dx+c)),I)-\tan(dx+c))$

3.18.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.54

$$\int \sqrt{b \sec(c+dx)} (A + C \sec^2(c+dx)) dx$$

$$= \frac{\sqrt{2}(-3iA - iC)\sqrt{b} \cos(dx+c) \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c)) + \sqrt{2}(3iA + iC)\sqrt{b} \cos(dx+c) \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i \sin(dx+c))}{3d \cos(dx+c)}$$

input `integrate((b*sec(d*x+c))^(1/2)*(A+C*sec(d*x+c)^2),x, algorithm="fricas")`

output $1/3*(\sqrt{2})*(-3IA - IC)*\sqrt{b}*\cos(dx+c)*\text{weierstrassPInverse}(-4, 0, \cos(dx+c) + I*\sin(dx+c)) + \sqrt{2}*(3IA + IC)*\sqrt{b}*\cos(dx+c)*\text{weierstrassPInverse}(-4, 0, \cos(dx+c) - I*\sin(dx+c)) + 2C*\sqrt{b}/(\cos(dx+c))*\sin(dx+c)/(d*\cos(dx+c))$

3.18.6 Sympy [F]

$$\int \sqrt{b \sec(c+dx)} (A + C \sec^2(c+dx)) dx = \int \sqrt{b \sec(c+dx)} (A + C \sec^2(c+dx)) dx$$

input `integrate((b*sec(d*x+c))**(1/2)*(A+C*sec(d*x+c)**2),x)`

output `Integral(sqrt(b*sec(c+d*x))*(A+C*sec(c+d*x)**2),x)`

3.18.7 Maxima [F]

$$\int \sqrt{b \sec(c + dx)} (A + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + A) \sqrt{b \sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2)*(A+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)*sqrt(b*sec(d*x + c)), x)`

3.18.8 Giac [F]

$$\int \sqrt{b \sec(c + dx)} (A + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + A) \sqrt{b \sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2)*(A+C*sec(d*x+c)^2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)*sqrt(b*sec(d*x + c)), x)`

3.18.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \sec(c + dx)} (A + C \sec^2(c + dx)) dx = \int \left(A + \frac{C}{\cos(c + dx)^2} \right) \sqrt{\frac{b}{\cos(c + dx)}} dx$$

input `int((A + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(1/2),x)`

output `int((A + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(1/2), x)`

3.19 $\int \frac{A+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.19.1 Optimal result 144
 3.19.2 Mathematica [C] (verified) 144
 3.19.3 Rubi [A] (verified) 145
 3.19.4 Maple [C] (verified) 146
 3.19.5 Fracas [C] (verification not implemented) 147
 3.19.6 Sympy [F] 148
 3.19.7 Maxima [F] 148
 3.19.8 Giac [F] 148
 3.19.9 Mupad [F(-1)] 149

3.19.1 Optimal result

Integrand size = 25, antiderivative size = 68

$$\int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{2(A - C)E(\frac{1}{2}(c + dx)|2)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2C \tan(c + dx)}{d\sqrt{b \sec(c + dx)}}$$

output `2*(A-C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*C*tan(d*x+c)/d/(b*sec(d*x+c))^(1/2)`

3.19.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.79 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.85

$$\int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \frac{2i(-3(A + Ae^{2i(c+dx)} - 2Ce^{2i(c+dx)}) + 2(A - C)e^{2i(c+dx)}\sqrt{1 + e^{2i(c+dx)}} \text{Hypergeometric2F1}(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -))}{3d(1 + e^{2i(c+dx)})\sqrt{b \sec(c + dx)}}$$

input `Integrate[(A + C*Sec[c + d*x]^2)/Sqrt[b*Sec[c + d*x]],x]`

output $(((-2*I)/3)*(-3*(A + A*E^{((2*I)*(c + d*x))} - 2*C*E^{((2*I)*(c + d*x))}) + 2*(A - C)*E^{((2*I)*(c + d*x))}*Sqrt[1 + E^{((2*I)*(c + d*x))}]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^{((2*I)*(c + d*x))}])/(d*(1 + E^{((2*I)*(c + d*x))})*Sqrt[b*Sec[c + d*x]))$

3.19.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {3042, 4534, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + C \csc(c + dx + \frac{\pi}{2})^2}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{4534} \\ & (A - C) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx + \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \\ & \quad \downarrow \text{3042} \\ & (A - C) \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx + \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \\ & \quad \downarrow \text{4258} \\ & \frac{(A - C) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{(A - C) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \\ & \quad \downarrow \text{3119} \\ & \frac{2(A - C)E(\frac{1}{2}(c + dx) | 2)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \end{aligned}$$

input `Int[(A + C*Sec[c + d*x]^2)/Sqrt[b*Sec[c + d*x]],x]`

output `(2*(A - C)*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*C*Tan[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])`

3.19.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.19.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.78 (sec) , antiderivative size = 763, normalized size of antiderivative = 11.22

method	result	size
default	Expression too large to display	763
parts	Expression too large to display	778

input `int((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `2/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(I*A*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*A*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*C*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+2*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)-2*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-2*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)+2*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)+I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)-I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)-I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)+I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)+A*sin(d*x+c)+C*tan(d*x+c)`

3.19.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.46

$$\int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{\sqrt{2}(i A - i C)\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + \sqrt{2}(-i A + i C)\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{2} + 2C\sqrt{b/\cos(dx + c)}\sin(dx + c)/(b*d)$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `(sqrt(2)*(I*A - I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + sqrt(2)*(-I*A + I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*C*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b*d)`

3.19. $\int \frac{A+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.19.6 Sympy [F]

$$\int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate((A+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(1/2),x)`

output `Integral((A + C*sec(c + d*x)**2)/sqrt(b*sec(c + d*x)), x)`

3.19.7 Maxima [F]

$$\int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{C \sec(dx + c)^2 + A}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)/sqrt(b*sec(d*x + c)), x)`

3.19.8 Giac [F]

$$\int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{C \sec(dx + c)^2 + A}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)/sqrt(b*sec(d*x + c)), x)`

3.19.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{A + \frac{C}{\cos(c+dx)^2}}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(1/2), x)`output `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(1/2), x)`

3.20 $\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.20.1	Optimal result	150
3.20.2	Mathematica [A] (verified)	150
3.20.3	Rubi [A] (verified)	151
3.20.4	Maple [C] (verified)	152
3.20.5	Fricas [C] (verification not implemented)	153
3.20.6	Sympy [F]	153
3.20.7	Maxima [F]	154
3.20.8	Giac [F]	154
3.20.9	Mupad [F(-1)]	154

3.20.1 Optimal result

Integrand size = 25, antiderivative size = 75

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{2(A + 3C) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3b^2d} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}}$$

output `2/3*(A+3*C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d+2/3*A*tan(d*x+c)/d/(b*sec(d*x+c))^(3/2)`

3.20.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.88

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{\sec^2(c + dx) \left(2(A + 3C) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + A \sin(2(c + dx)) \right)}{3d(b \sec(c + dx))^{3/2}}$$

input `Integrate[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(3/2), x]`

output `(Sec[c + d*x]^2*(2*(A + 3*C)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + A*Sin[2*(c + d*x)])/(3*d*(b*Sec[c + d*x])^(3/2))`

3.20.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {3042, 4533, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4533} \\
 & \frac{(A + 3C) \int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A + 3C) \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{(A + 3C) \sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A + 3C) \sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2(A + 3C) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3b^2 d} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}}
 \end{aligned}$$

input `Int[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(3/2),x]`

output `(2*(A + 3*C)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*A*Tan[c + d*x])/(3*d*(b*Sec[c + d*x])^(3/2))`

3.20. $\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx$

3.20.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^n_, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.))^m_*(csc[(e_.) + (f_.)*(x_)]^2*(C_. + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

3.20.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.91 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.91

method	result
parts	$-\frac{2A \left(i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)), i\right) + i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)), i\right) \right)}{3d\sqrt{b \sec(dx+c)} b}$
default	$\frac{2iA \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \sqrt{\frac{1}{\cos(dx+c)+1}} \operatorname{EllipticF}\left(i(\cot(dx+c)-\csc(dx+c)), i\right)}{3} + 2iC \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \sqrt{\frac{1}{\cos(dx+c)+1}} \operatorname{EllipticF}\left(i(\cot(dx+c)-\csc(dx+c)), i\right)$

input `int((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2), x, method=_RETURNVERBOSE)`

output
$$-2/3*A/d/(b*\sec(d*x+c))^{1/2}/b*(I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\text{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)+I*(1/(\cos(d*x+c)+1))^{1/2}*(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}*\text{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)*\sec(d*x+c)-\sin(d*x+c))-2*I*C/d*(1/(\cos(d*x+c)+1))^{1/2}*\text{EllipticF}(I*(-\cot(d*x+c)+\csc(d*x+c)),I)/b/(b*\sec(d*x+c))^{1/2}/(\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2}$$

3.20.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.33

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{2A \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) + \sqrt{2}(-iA - 3iC) \sqrt{b} \text{weierstrassPInverse}}{}$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")`

output
$$1/3*(2*A*\sqrt{b/\cos(d*x+c)}*\cos(d*x+c)*\sin(d*x+c) + \sqrt{2}*(-I*A - 3*I*C)*\sqrt{b}*\text{weierstrassPInverse}(-4, 0, \cos(d*x+c) + I*\sin(d*x+c)) + \sqrt{2}*(I*A + 3*I*C)*\sqrt{b}*\text{weierstrassPInverse}(-4, 0, \cos(d*x+c) - I*\sin(d*x+c)))/(b^2*d)$$

3.20.6 Sympy [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(3/2),x)`

output `Integral((A + C*sec(c + d*x)**2)/(b*sec(c + d*x))**(3/2), x)`

3.20.7 Maxima [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(3/2), x)`

3.20.8 Giac [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(3/2), x)`

3.20.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{A + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(3/2),x)`

output `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(3/2), x)`

3.21 $\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.21.1	Optimal result	155
3.21.2	Mathematica [C] (verified)	155
3.21.3	Rubi [A] (verified)	156
3.21.4	Maple [C] (verified)	157
3.21.5	Fricas [C] (verification not implemented)	158
3.21.6	Sympy [F]	159
3.21.7	Maxima [F]	159
3.21.8	Giac [F]	159
3.21.9	Mupad [F(-1)]	160

3.21.1 Optimal result

Integrand size = 25, antiderivative size = 77

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{2(3A + 5C)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5b^2d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}}$$

```
output 2/5*(3*A+5*C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/5*A*tan(d*x+c)/d/(b*sec(d*x+c))^(5/2)
```

3.21.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.22 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.73

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{e^{-idx} \sec^2(c + dx)(\cos(dx) + i \sin(dx)) \left(12i(3A + 5C) - \frac{8i(3A+5C)e^{2i(c+dx)} \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{\left(\frac{2i}{b} (c + dx)\right)}\right]}{\sqrt{1+e^{2i(c+dx)}}}\right)}{30d(b \sec(c + dx))^{5/2}}$$

```
input Integrate[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(5/2),x]
```

```
output (Sec[c + d*x]^2*(Cos[d*x] + I*Sin[d*x])*((12*I)*(3*A + 5*C) - ((8*I)*(3*A + 5*C)*E^((2*I)*(c + d*x))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/Sqrt[1 + E^((2*I)*(c + d*x))] + 6*A*Sin[2*(c + d*x)]))/(30*d*E^(I*d*x)*(b*Sec[c + d*x])^(5/2))
```

3.21.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {3042, 4533, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4533} \\
 & \frac{(3A + 5C) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{5b^2} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(3A + 5C) \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx}{5b^2} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{(3A + 5C) \int \frac{\sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}}{5b^2} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(3A + 5C) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}}{5b^2} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2(3A + 5C)E(\frac{1}{2}(c + dx) | 2)}{5b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}}
 \end{aligned}$$

input `Int[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(5/2),x]`

output `(2*(3*A + 5*C)*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*A*Tan[c + d*x])/(5*d*(b*Sec[c + d*x])^(5/2))`

3.21. $\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx$

3.21.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

3.21.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 5.27 (sec) , antiderivative size = 798, normalized size of antiderivative = 10.36

method	result	size
default	Expression too large to display	798
parts	Expression too large to display	812

input `int((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `2/5/b^2/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(-3*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)-3*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)+6*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)-10*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)+5*I*C*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)+10*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)+3*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)-6*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-5*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)-5*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)+A*cos(d*x+c)^2*sin(d*x+c)+5*I*C*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)+3*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)+A*cos(d*x+c)*sin(d*x+c)+3*A*sin(d*x+c)+5*C*sin(d*x+c))`

3.21.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.40

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{2A \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^2 \sin(dx+c) + \sqrt{2}(3iA + 5iC) \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + I \sin(dx+c))) + \sqrt{2}(-3iA - 5iC) \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - I \sin(dx+c)))}{(b^3 d)}$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/5*(2*A*sqrt(b/cos(d*x + c))*cos(d*x + c)^2*sin(d*x + c) + sqrt(2)*(3*I*A + 5*I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + sqrt(2)*(-3*I*A - 5*I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b^3*d)`

3.21.6 Sympy [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(5/2),x)`

output `Integral((A + C*sec(c + d*x)**2)/(b*sec(c + d*x))**(5/2), x)`

3.21.7 Maxima [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(5/2), x)`

3.21.8 Giac [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(5/2), x)`

3.21.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{A + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(5/2),x)`output `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(5/2), x)`

3.22 $\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$

3.22.1	Optimal result	161
3.22.2	Mathematica [A] (verified)	161
3.22.3	Rubi [A] (verified)	162
3.22.4	Maple [C] (verified)	164
3.22.5	Fricas [C] (verification not implemented)	164
3.22.6	Sympy [F]	165
3.22.7	Maxima [F]	165
3.22.8	Giac [F]	165
3.22.9	Mupad [F(-1)]	166

3.22.1 Optimal result

Integrand size = 25, antiderivative size = 112

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{2(5A + 7C) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{21b^4d} + \frac{2(5A + 7C) \sin(c + dx)}{21b^3d \sqrt{b \sec(c + dx)}} + \frac{2A \tan(c + dx)}{7d(b \sec(c + dx))^{7/2}}$$

output `2/21*(5*A+7*C)*sin(d*x+c)/b^3/d/(b*sec(d*x+c))^(1/2)+2/21*(5*A+7*C)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^4/d+2/7*A*tan(d*x+c)/d/(b*sec(d*x+c))^(7/2)`

3.22.2 Mathematica [A] (verified)

Time = 0.92 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.71

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{\frac{4(5A+7C) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{\sqrt{\cos(c+dx)}} + 2(13A + 14C + 3A \cos(2(c + dx))) \sin(c + dx)}{42b^3d \sqrt{b \sec(c + dx)}}$$

input `Integrate[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(7/2),x]`

output `((4*(5*A + 7*C)*EllipticF[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*(13*A + 14*C + 3*A*Cos[2*(c + d*x)])*Sin[c + d*x])/(42*b^3*d*Sqrt[b*Sec[c + d*x]])`

3.22.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {3042, 4533, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4533} \\
 & \frac{(5A + 7C) \int \frac{1}{(b \sec(c + dx))^{3/2}} dx}{7b^2} + \frac{2A \tan(c + dx)}{7d(b \sec(c + dx))^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(5A + 7C) \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2A \tan(c + dx)}{7d(b \sec(c + dx))^{7/2}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{(5A + 7C) \left(\frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right)}{7b^2} + \frac{2A \tan(c + dx)}{7d(b \sec(c + dx))^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(5A + 7C) \left(\frac{\int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right)}{7b^2} + \frac{2A \tan(c + dx)}{7d(b \sec(c + dx))^{7/2}} \\
 & \quad \downarrow \text{4258} \\
 & \frac{(5A + 7C) \left(\frac{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right)}{7b^2} + \frac{2A \tan(c + dx)}{7d(b \sec(c + dx))^{7/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{(5A + 7C) \left(\frac{\int \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}}$$

↓ 3120

$$\frac{(5A + 7C) \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2, \sqrt{b \sec(c+dx)}\right)}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}}$$

input `Int[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(7/2), x]`

output `((5*A + 7*C)*((2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*sqrt[b*Sec[c + d*x]])))/(7*b^2) + (2*A*Tan[c + d*x])/(7*d*(b*Sec[c + d*x])^(7/2))`

3.2.2.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_. + (A_.)), x_Symbol] :> Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

3.22.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.66 (sec) , antiderivative size = 291, normalized size of antiderivative = 2.60

method	result
default	$\frac{10iA\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\sqrt{\frac{1}{\cos(dx+c)+1}}\text{EllipticF}(i(\cot(dx+c)-\csc(dx+c)),i)}{21} + \frac{2iC\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\sqrt{\frac{1}{\cos(dx+c)+1}}\text{EllipticF}(i(\cot(dx+c)-\csc(dx+c)))}{3}$
parts	$-\frac{2A\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)))\right)}{21d\sqrt{b\sec(dx+c)}b^3}$

input `int((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output
$$\frac{2}{21} \frac{1}{b^3} \frac{d}{(b \sec(d*x+c))^{1/2}} * (5*I*A*(1/(\cos(d*x+c)+1))^{1/2} * (\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2} * \text{EllipticF}(I*(\cot(d*x+c)-\csc(d*x+c)), I) + 7*I*C*(1/(\cos(d*x+c)+1))^{1/2} * (\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2} * \text{EllipticF}(I*(\cot(d*x+c)-\csc(d*x+c)), I) + 5*I*A*(1/(\cos(d*x+c)+1))^{1/2} * (\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2} * \text{EllipticF}(I*(\cot(d*x+c)-\csc(d*x+c)), I) * \sec(d*x+c) + 3*A*\cos(d*x+c)^2 * \sin(d*x+c) + 7*I*C*(1/(\cos(d*x+c)+1))^{1/2} * (\cos(d*x+c)/(\cos(d*x+c)+1))^{1/2} * \text{EllipticF}(I*(\cot(d*x+c)-\csc(d*x+c)), I) * \sec(d*x+c) + 5*A*\sin(d*x+c) + 7*C*\sin(d*x+c))$$

3.22.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.06

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{\sqrt{2}(-5iA - 7iC)\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{21d\sqrt{b\sec(dx+c)}b^3} + \dots$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm="fracas")`

3.22.
$$\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$$

output `1/21*(sqrt(2)*(-5*I*A - 7*I*C)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(5*I*A + 7*I*C)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(3*A*cos(d*x + c)^3 + (5*A + 7*C)*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b^4*d)`

3.22.6 Sympy [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx$$

input `integrate((A+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(7/2),x)`

output `Integral((A + C*sec(c + d*x)**2)/(b*sec(c + d*x))**(7/2), x)`

3.22.7 Maxima [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{7/2}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(7/2), x)`

3.22.8 Giac [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{7/2}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(7/2), x)`

3.22.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{A + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{7/2}} dx$$

input `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(7/2),x)`output `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(7/2), x)`

3.23 $\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx$

3.23.1	Optimal result	167
3.23.2	Mathematica [C] (verified)	167
3.23.3	Rubi [A] (verified)	168
3.23.4	Maple [C] (verified)	170
3.23.5	Fricas [C] (verification not implemented)	171
3.23.6	Sympy [F(-1)]	171
3.23.7	Maxima [F]	171
3.23.8	Giac [F]	172
3.23.9	Mupad [F(-1)]	172

3.23.1 Optimal result

Integrand size = 25, antiderivative size = 112

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \frac{2(7A + 9C)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15b^4d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2(7A + 9C) \sin(c + dx)}{45b^3d(b \sec(c + dx))^{3/2}} + \frac{2A \tan(c + dx)}{9d(b \sec(c + dx))^{9/2}}$$

output `2/45*(7*A+9*C)*sin(d*x+c)/b^3/d/(b*sec(d*x+c))^(3/2)+2/15*(7*A+9*C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^4/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/9*A*tan(d*x+c)/d/(b*sec(d*x+c))^(9/2)`

3.23.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.32 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.28

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \frac{e^{-idx}(\cos(dx) + i \sin(dx)) \left(336iA + 432iC - \frac{32i(7A+9C)e^{2i(c+dx)} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\right)}{\sqrt{1+e^{2i(c+dx)}}} \right)}{360b^4d\sqrt{b \sec(c + dx)}}$$

input `Integrate[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(9/2),x]`


```
output ((Cos[d*x] + I*Sin[d*x])*((336*I)*A + (432*I)*C - ((32*I)*(7*A + 9*C)*E^((2*I)*(c + d*x))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/Sqrt[1 + E^((2*I)*(c + d*x))] + (76*A + 72*C)*Sin[2*(c + d*x)] + 10*A*Sin[4*(c + d*x)]))/(360*b^4*d*E^(I*d*x)*Sqrt[b*Sec[c + d*x]])
```

3.23.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {3042, 4533, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{9/2}} dx \\
 & \quad \downarrow \text{4533} \\
 & \frac{(7A + 9C) \int \frac{1}{(b \sec(c + dx))^{5/2}} dx}{9b^2} + \frac{2A \tan(c + dx)}{9d(b \sec(c + dx))^{9/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(7A + 9C) \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx}{9b^2} + \frac{2A \tan(c + dx)}{9d(b \sec(c + dx))^{9/2}} \\
 & \quad \downarrow \text{4256} \\
 & \frac{(7A + 9C) \left(\frac{3 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{5b^2} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} \right)}{9b^2} + \frac{2A \tan(c + dx)}{9d(b \sec(c + dx))^{9/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(7A + 9C) \left(\frac{3 \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} \right)}{9b^2} + \frac{2A \tan(c + dx)}{9d(b \sec(c + dx))^{9/2}} \\
 & \quad \downarrow \text{4258}
 \end{aligned}$$

$$\frac{(7A + 9C) \left(\frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2A \tan(c+dx)}{9d(b \sec(c+dx))^{9/2}}$$

↓ 3042

$$\frac{(7A + 9C) \left(\frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2A \tan(c+dx)}{9d(b \sec(c+dx))^{9/2}}$$

↓ 3119

$$\frac{(7A + 9C) \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{9b^2} + \frac{2A \tan(c+dx)}{9d(b \sec(c+dx))^{9/2}}$$

input `Int[(A + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(9/2),x]`

output `((7*A + 9*C)*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])) + (2*Sin[c + d*x]/(5*b*d*(b*Sec[c + d*x])^(3/2))))/(9*b^2) + (2*A*Tan[c + d*x])/(9*d*(b*Sec[c + d*x])^(9/2))`

3.23.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

```
rule 4533 Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.
+ (A_.)), x_Symbol] :> Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] +
Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; Fr
eeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]
```

3.23.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 7.76 (sec) , antiderivative size = 866, normalized size of antiderivative = 7.73

method	result	size
default	Expression too large to display	866
parts	Expression too large to display	876

```
input int((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x,method=_RETURNVERBOSE)
```

```
output -2/45/b^4/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(-27*I*C*(1/(cos(d*x+c)+1)
)^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+
c)),I)*sec(d*x+c)+42*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+
1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-5*A*cos(d*x+c)^4*sin(d*x+
c)-21*I*A*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Ellip
ticF(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)+21*I*A*(1/(cos(d*x+c)+1))^(1/
2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I
)*sec(d*x+c)+54*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(
1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-42*I*A*(1/(cos(d*x+c)+1))^(1/2
)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)
-5*A*cos(d*x+c)^3*sin(d*x+c)+27*I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(
cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*sec(d*x+c)-54*
I*C*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I
*(cot(d*x+c)-csc(d*x+c)),I)-21*I*A*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*
(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-27*I
*C*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*
x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-7*A*cos(d*x+c)^2*sin(d*x+c)+21*I*A*E
llipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)
/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+27*I*C*EllipticE(I*(cot(d*x+c)-csc(d*x+c
)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*...
```

3.23.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.15

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx =$$

$$3\sqrt{2}(-7iA - 9iC)\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) +$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x, algorithm="fracas")`

output `-1/45*(3*sqrt(2)*(-7*I*A - 9*I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(7*I*A + 9*I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(5*A*cos(d*x + c)^4 + (7*A + 9*C)*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^5*d)`

3.23.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \text{Timed out}$$

input `integrate((A+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(9/2),x)`

output `Timed out`

3.23.7 Maxima [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{\frac{9}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(9/2), x)`

3.23. $\int \frac{A+C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx$

3.23.8 Giac [F]

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \int \frac{C \sec(dx + c)^2 + A}{(b \sec(dx + c))^{\frac{9}{2}}} dx$$

input `integrate((A+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + A)/(b*sec(d*x + c))^(9/2), x)`

3.23.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \int \frac{A + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{9/2}} dx$$

input `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(9/2),x)`

output `int((A + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(9/2), x)`

$$3.24 \quad \int \frac{3+3 \sec^2(c+dx)}{\sqrt{\sec(c+dx)}} dx$$

3.24.1	Optimal result	173
3.24.2	Mathematica [A] (verified)	173
3.24.3	Rubi [A] (verified)	174
3.24.4	Maple [B] (verified)	175
3.24.5	Fricas [A] (verification not implemented)	175
3.24.6	Sympy [F]	175
3.24.7	Maxima [F]	176
3.24.8	Giac [B] (verification not implemented)	176
3.24.9	Mupad [B] (verification not implemented)	176

3.24.1 Optimal result

Integrand size = 23, antiderivative size = 21

$$\int \frac{3 + 3 \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx = \frac{6\sqrt{\sec(c + dx)} \sin(c + dx)}{d}$$

output `6*sin(d*x+c)*sec(d*x+c)^(1/2)/d`

3.24.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{3 + 3 \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx = \frac{6\sqrt{\sec(c + dx)} \sin(c + dx)}{d}$$

input `Integrate[(3 + 3*Sec[c + d*x]^2)/Sqrt[Sec[c + d*x]],x]`

output `(6*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d`

3.24.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3 \sec^2(c + dx) + 3}{\sqrt{\sec(c + dx)}} dx$$

↓ 3042

$$\int \frac{3 \csc(c + dx + \frac{\pi}{2})^2 + 3}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

↓ 4531

$$\frac{6 \sin(c + dx) \sqrt{\sec(c + dx)}}{d}$$

input `Int[(3 + 3*Sec[c + d*x]^2)/Sqrt[Sec[c + d*x]],x]`

output `(6*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d`

3.24.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.24.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. $2(19) = 38$.

Time = 1.56 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.95

method	result
default	$\frac{12 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1} d}$
parts	$\frac{6 \sqrt{\left(2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right)^2} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1} \operatorname{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - 6 \left(-2 \sqrt{-2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1}\right)}{\sqrt{-2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + \sin\left(\frac{dx}{2} + \frac{c}{2}\right)^2} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1} d}$

input `int((3+3*sec(d*x+c)^2)/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `12*sin(1/2*d*x+1/2*c)*cos(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.24.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{3 + 3 \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx = \frac{6 \sin(dx + c)}{d \sqrt{\cos(dx + c)}}$$

input `integrate((3+3*sec(d*x+c)^2)/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `6*sin(d*x + c)/(d*sqrt(cos(d*x + c)))`

3.24.6 Sympy [F]

$$\int \frac{3 + 3 \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx = 3 \left(\int \frac{1}{\sqrt{\sec(c + dx)}} dx + \int \sec^{\frac{3}{2}}(c + dx) dx \right)$$

input `integrate((3+3*sec(d*x+c)**2)/sec(d*x+c)**(1/2),x)`

output `3*(Integral(1/sqrt(sec(c + d*x)), x) + Integral(sec(c + d*x)**(3/2), x))`

3.24. $\int \frac{3+3\sec^2(c+dx)}{\sqrt{\sec(c+dx)}} dx$

3.24.7 Maxima [F]

$$\int \frac{3 + 3 \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx = \int \frac{3 (\sec(dx + c)^2 + 1)}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((3+3*sec(d*x+c)^2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `3*integrate((sec(d*x + c)^2 + 1)/sqrt(sec(d*x + c)), x)`

3.24.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(19) = 38$.

Time = 0.32 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.19

$$\int \frac{3 + 3 \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx = -\frac{12 \sqrt{-\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 1} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 1\right) d}$$

input `integrate((3+3*sec(d*x+c)^2)/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `-12*sqrt(-tan(1/2*d*x + 1/2*c)^4 + 1)*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^4 - 1)*d)`

3.24.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{3 + 3 \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx = \frac{6 \sin(c + dx) \sqrt{\frac{1}{\cos(c + dx)}}}{d}$$

input `int((3/cos(c + d*x)^2 + 3)/(1/cos(c + d*x))^(1/2),x)`

output `(6*sin(c + d*x)*(1/cos(c + d*x))^(1/2))/d`

3.25 $\int \sec^m(e+fx) (m - (1 + m) \sec^2(e + fx)) dx$

3.25.1	Optimal result	177
3.25.2	Mathematica [C] (verified)	177
3.25.3	Rubi [A] (verified)	178
3.25.4	Maple [A] (verified)	179
3.25.5	Fricas [A] (verification not implemented)	179
3.25.6	Sympy [F]	179
3.25.7	Maxima [B] (verification not implemented)	180
3.25.8	Giac [F]	180
3.25.9	Mupad [B] (verification not implemented)	181

3.25.1 Optimal result

Integrand size = 24, antiderivative size = 21

$$\int \sec^m(e + fx) (m - (1 + m) \sec^2(e + fx)) dx = -\frac{\sec^{1+m}(e + fx) \sin(e + fx)}{f}$$

output `-sec(f*x+e)^(1+m)*sin(f*x+e)/f`

3.25.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.39 (sec) , antiderivative size = 107, normalized size of antiderivative = 5.10

$$\int \sec^m(e + fx) (m - (1 + m) \sec^2(e + fx)) dx = \frac{\csc(e + fx) \sec^{-1+m}(e + fx) ((2 + m) \text{Hypergeometric2F1}(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, \sec^2(e + fx)) - (1 + m) \text{Hypergeometric2F1}(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, \sec^2(e + fx)))}{f(2 + m)}$$

input `Integrate[Sec[e + f*x]^m*(m - (1 + m)*Sec[e + f*x]^2),x]`

output `(Csc[e + f*x]*Sec[e + f*x]^(-1 + m)*((2 + m)*Hypergeometric2F1[1/2, m/2, (2 + m)/2, Sec[e + f*x]^2] - (1 + m)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Sec[e + f*x]^2]*Sec[e + f*x]^2)*Sqrt[-Tan[e + f*x]^2])/(f*(2 + m))`

3.25.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^m(e + fx) (m - (m + 1) \sec^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right)^m \left((-m - 1) \csc\left(e + fx + \frac{\pi}{2}\right)^2 + m\right) dx$$

$$\downarrow \text{4531}$$

$$-\frac{\sin(e + fx) \sec^{m+1}(e + fx)}{f}$$

input `Int[Sec[e + f*x]^m*(m - (1 + m)*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^(1 + m)*Sin[e + f*x])/f)`

3.25.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m]*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.25.4 Maple [A] (verified)

Time = 1.15 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.90

method	result
parallelrisch	$\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) \left(\frac{1}{\cos(fx+e)}\right)^m}{f \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)}$
risch	$i(e^{i(fx+e)})^m (e^{2i(fx+e)+1})^{-m} 2^m \left(e^{\frac{i \operatorname{csgn}\left(\frac{i}{e^{2i(fx+e)+1}}\right) \operatorname{csgn}\left(\frac{ie^{i(fx+e)}}{e^{2i(fx+e)+1}}\right)^2 \pi m}{2}} e^{-\frac{i \operatorname{csgn}\left(\frac{i}{e^{2i(fx+e)+1}}\right) \operatorname{csgn}\left(\frac{ie^{i(fx+e)}}{e^{2i(fx+e)+1}}\right)}{2}} $

input `int(sec(f*x+e)^m*(m-(1+m)*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `2/f*tan(1/2*f*x+1/2*e)*(1/cos(f*x+e))^m/(tan(1/2*f*x+1/2*e)^2-1)`

3.25.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.38

$$\int \sec^m(e+fx) (m - (1+m) \sec^2(e+fx)) dx = -\frac{\frac{1}{\cos(fx+e)}^m \sin(fx+e)}{f \cos(fx+e)}$$

input `integrate(sec(f*x+e)^m*(m-(1+m)*sec(f*x+e)^2),x, algorithm="fracas")`

output `-(1/cos(f*x + e))^m*sin(f*x + e)/(f*cos(f*x + e))`

3.25.6 Sympy [F]

$$\begin{aligned} & \int \sec^m(e+fx) (m - (1+m) \sec^2(e+fx)) dx \\ &= - \int (-m \sec^m(e+fx)) dx - \int \sec^2(e+fx) \sec^m(e+fx) dx \\ & \quad - \int m \sec^2(e+fx) \sec^m(e+fx) dx \end{aligned}$$

input `integrate(sec(f*x+e)**m*(m-(1+m)*sec(f*x+e)**2),x)`

output `-Integral(-m*sec(e + f*x)**m, x) - Integral(sec(e + f*x)**2*sec(e + f*x)**m, x) - Integral(m*sec(e + f*x)**2*sec(e + f*x)**m, x)`

3.25.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 283 vs. $2(21) = 42$.

Time = 0.43 (sec) , antiderivative size = 283, normalized size of antiderivative = 13.48

$$\int \sec^m(e + fx) (m - (1 + m) \sec^2(e + fx)) dx$$

$$= \frac{2^m \cos(-(fx + e)(m + 2) + m \arctan(\sin(2fx + 2e), \cos(2fx + 2e) + 1)) \sin(2fx + 2e) - 2^m \cos(-$$

input `integrate(sec(f*x+e)^m*(m-(1+m)*sec(f*x+e)^2),x, algorithm="maxima")`

output `(2^m*cos(-(f*x + e)*(m + 2) + m*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1))*sin(2*f*x + 2*e) - 2^m*cos(-(f*x + e)*m + m*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1))*sin(2*f*x + 2*e) + (2^m*cos(2*f*x + 2*e) + 2^m)*sin(-(f*x + e)*(m + 2) + m*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - (2^m*cos(2*f*x + 2*e) + 2^m)*sin(-(f*x + e)*m + m*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)))/((cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/2*m)*f)`

3.25.8 Giac [F]

$$\int \sec^m(e + fx) (m - (1 + m) \sec^2(e + fx)) dx$$

$$= \int -((m + 1) \sec^2(fx + e) - m) \sec^m(fx + e) dx$$

input `integrate(sec(f*x+e)^m*(m-(1+m)*sec(f*x+e)^2),x, algorithm="giac")`

output `integrate(-((m + 1)*sec(f*x + e)^2 - m)*sec(f*x + e)^m, x)`

3.25. $\int \sec^m(e + fx) (m - (1 + m) \sec^2(e + fx)) dx$

3.25.9 Mupad [B] (verification not implemented)

Time = 15.39 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \sec^m(e + fx) (m - (1 + m) \sec^2(e + fx)) dx = -\frac{\sin(2e + 2fx) \left(\frac{1}{\cos(e+fx)}\right)^m}{f (\cos(2e + 2fx) + 1)}$$

input `int((m - (m + 1)/cos(e + f*x)^2)*(1/cos(e + f*x))^m,x)`

output `-(sin(2*e + 2*f*x)*(1/cos(e + f*x))^m)/(f*(cos(2*e + 2*f*x) + 1))`

3.26 $\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx$

3.26.1	Optimal result	182
3.26.2	Mathematica [A] (verified)	182
3.26.3	Rubi [A] (verified)	183
3.26.4	Maple [C] (verified)	184
3.26.5	Fricas [A] (verification not implemented)	184
3.26.6	Sympy [F]	185
3.26.7	Maxima [B] (verification not implemented)	185
3.26.8	Giac [A] (verification not implemented)	185
3.26.9	Mupad [B] (verification not implemented)	186

3.26.1 Optimal result

Integrand size = 21, antiderivative size = 19

$$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx = -\frac{\sec^5(e + fx) \tan(e + fx)}{f}$$

output `-sec(f*x+e)^5*tan(f*x+e)/f`

3.26.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx = -\frac{\sec^5(e + fx) \tan(e + fx)}{f}$$

input `Integrate[Sec[e + f*x]^5*(5 - 6*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^5*Tan[e + f*x])/f)`

3.26.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right)^5 \left(5 - 6 \csc\left(e + fx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{4531}$$

$$-\frac{\tan(e + fx) \sec^5(e + fx)}{f}$$

input `Int[Sec[e + f*x]^5*(5 - 6*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^5*Tan[e + f*x])/f)`

3.26.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.26.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.16

method	result
risch	$\frac{32i(e^{7i(fx+e)} - e^{5i(fx+e)})}{f(e^{2i(fx+e)} + 1)^6}$
parallelrisch	$-\frac{32 \sin(fx+e)}{f(\cos(6fx+6e) + 6 \cos(4fx+4e) + 15 \cos(2fx+2e) + 10)}$
derivativedivides	$-\frac{5 \left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + 6 \left(-\frac{\sec(fx+e)^5}{6} - \frac{5 \sec(fx+e)^3}{24} - \frac{5 \sec(fx+e)}{16} \right) \tan(fx+e)}{f}$
default	$-\frac{5 \left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + 6 \left(-\frac{\sec(fx+e)^5}{6} - \frac{5 \sec(fx+e)^3}{24} - \frac{5 \sec(fx+e)}{16} \right) \tan(fx+e)}{f}$
parts	$-\frac{5 \left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{15 \ln(\sec(fx+e) + \tan(fx+e))}{8}}{f} - \frac{6 \left(-\frac{\sec(fx+e)^5}{6} - \frac{5 \sec(fx+e)^3}{24} - \frac{5 \sec(fx+e)}{16} \right) \tan(fx+e)}{f}$
norman	$-\frac{\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f} - \frac{10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{f} - \frac{20 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f} - \frac{20 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{f} - \frac{10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{f} - \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^6}$

input `int(sec(f*x+e)^5*(5-6*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `32*I/f/(exp(2*I*(f*x+e))+1)^6*(exp(7*I*(f*x+e))-exp(5*I*(f*x+e)))`

3.26.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx = -\frac{\sin(fx + e)}{f \cos(fx + e)^6}$$

input `integrate(sec(f*x+e)^5*(5-6*sec(f*x+e)^2),x, algorithm="fricas")`

output `-sin(f*x + e)/(f*cos(f*x + e)^6)`

3.26.6 Sympy [F]

$$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx = - \int (-5 \sec^5(e + fx)) dx - \int 6 \sec^7(e + fx) dx$$

input `integrate(sec(f*x+e)**5*(5-6*sec(f*x+e)**2),x)`

output `-Integral(-5*sec(e + f*x)**5, x) - Integral(6*sec(e + f*x)**7, x)`

3.26.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(19) = 38.

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.21

$$\begin{aligned} & \int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx \\ &= \frac{\sin(fx + e)}{(\sin(fx + e))^6 - 3 \sin(fx + e)^4 + 3 \sin(fx + e)^2 - 1} f \end{aligned}$$

input `integrate(sec(f*x+e)^5*(5-6*sec(f*x+e)^2),x, algorithm="maxima")`

output `sin(f*x + e)/((sin(f*x + e)^6 - 3*sin(f*x + e)^4 + 3*sin(f*x + e)^2 - 1)*f)`

3.26.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx = \frac{\sin(fx + e)}{(\sin(fx + e))^2 - 1}^3 f$$

input `integrate(sec(f*x+e)^5*(5-6*sec(f*x+e)^2),x, algorithm="giac")`

output `sin(f*x + e)/((sin(f*x + e)^2 - 1)^3*f)`

3.26.9 Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.21

$$\int \sec^5(e + fx) (5 - 6 \sec^2(e + fx)) dx$$

$$= \frac{\sin(e + fx)}{f (\sin(e + fx)^6 - 3 \sin(e + fx)^4 + 3 \sin(e + fx)^2 - 1)}$$

input `int(-(6/cos(e + f*x)^2 - 5)/cos(e + f*x)^5,x)`

output `sin(e + f*x)/(f*(3*sin(e + f*x)^2 - 3*sin(e + f*x)^4 + sin(e + f*x)^6 - 1))`

3.27 $\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx$

3.27.1	Optimal result	187
3.27.2	Mathematica [A] (verified)	187
3.27.3	Rubi [A] (verified)	188
3.27.4	Maple [C] (verified)	189
3.27.5	Fricas [A] (verification not implemented)	189
3.27.6	Sympy [F]	190
3.27.7	Maxima [A] (verification not implemented)	190
3.27.8	Giac [A] (verification not implemented)	190
3.27.9	Mupad [B] (verification not implemented)	191

3.27.1 Optimal result

Integrand size = 21, antiderivative size = 19

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx = -\frac{\sec^4(e + fx) \tan(e + fx)}{f}$$

output `-sec(f*x+e)^4*tan(f*x+e)/f`

3.27.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx = -\frac{\sec^4(e + fx) \tan(e + fx)}{f}$$

input `Integrate[Sec[e + f*x]^4*(4 - 5*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^4*Tan[e + f*x])/f)`

3.27.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right)^4 \left(4 - 5 \csc\left(e + fx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{4531}$$

$$-\frac{\tan(e + fx) \sec^4(e + fx)}{f}$$

input `Int[Sec[e + f*x]^4*(4 - 5*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^4*Tan[e + f*x])/f)`

3.27.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.27.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.16

method	result	size
risch	$\frac{16i(e^{6i(fx+e)} - e^{4i(fx+e)})}{f(e^{2i(fx+e)} + 1)^5}$	41
parallelrisch	$-\frac{16 \sin(fx+e)}{f(\cos(5fx+5e) + 5 \cos(3fx+3e) + 10 \cos(fx+e))}$	43
derivativedivides	$-\frac{4\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e) + 5\left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15}\right) \tan(fx+e)}{f}$	56
default	$-\frac{4\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e) + 5\left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15}\right) \tan(fx+e)}{f}$	56
parts	$-\frac{4\left(-\frac{2}{3} - \frac{\sec(fx+e)^2}{3}\right) \tan(fx+e)}{f} + \frac{5\left(-\frac{8}{15} - \frac{\sec(fx+e)^4}{5} - \frac{4 \sec(fx+e)^2}{15}\right) \tan(fx+e)}{f}$	58
norman	$\frac{\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f} + \frac{8 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{f} + \frac{12 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f} + \frac{8 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{f} + \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^5}$	96

input `int(sec(f*x+e)^4*(4-5*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`

output $16*I/f/(exp(2*I*(f*x+e))+1)^5*(exp(6*I*(f*x+e))-exp(4*I*(f*x+e)))$

3.27.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx = -\frac{\sin(fx + e)}{f \cos(fx + e)^5}$$

input `integrate(sec(f*x+e)^4*(4-5*sec(f*x+e)^2),x, algorithm="fracas")`

output `-sin(f*x + e)/(f*cos(f*x + e)^5)`

3.27.6 Sympy [F]

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx = - \int (-4 \sec^4(e + fx)) dx - \int 5 \sec^6(e + fx) dx$$

input `integrate(sec(f*x+e)**4*(4-5*sec(f*x+e)**2),x)`

output `-Integral(-4*sec(e + f*x)**4, x) - Integral(5*sec(e + f*x)**6, x)`

3.27.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.58

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx = - \frac{\tan(fx + e)^5 + 2 \tan(fx + e)^3 + \tan(fx + e)}{f}$$

input `integrate(sec(f*x+e)^4*(4-5*sec(f*x+e)^2),x, algorithm="maxima")`

output `-(tan(f*x + e)^5 + 2*tan(f*x + e)^3 + tan(f*x + e))/f`

3.27.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.58

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx = - \frac{\tan(fx + e)^5 + 2 \tan(fx + e)^3 + \tan(fx + e)}{f}$$

input `integrate(sec(f*x+e)^4*(4-5*sec(f*x+e)^2),x, algorithm="giac")`

output `-(tan(f*x + e)^5 + 2*tan(f*x + e)^3 + tan(f*x + e))/f`

3.27.9 Mupad [B] (verification not implemented)

Time = 15.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^4(e + fx) (4 - 5 \sec^2(e + fx)) dx = -\frac{\sin(e + fx)}{f \cos(e + fx)^5}$$

input `int(-(5/cos(e + f*x)^2 - 4)/cos(e + f*x)^4,x)`

output `-sin(e + f*x)/(f*cos(e + f*x)^5)`

3.28 $\int \sec^3(e + fx) (3 - 4 \sec^2(e + fx)) dx$

3.28.1	Optimal result	192
3.28.2	Mathematica [A] (verified)	192
3.28.3	Rubi [A] (verified)	193
3.28.4	Maple [A] (verified)	194
3.28.5	Fricas [A] (verification not implemented)	194
3.28.6	Sympy [F]	195
3.28.7	Maxima [A] (verification not implemented)	195
3.28.8	Giac [A] (verification not implemented)	195
3.28.9	Mupad [B] (verification not implemented)	196

3.28.1 Optimal result

Integrand size = 21, antiderivative size = 19

$$\int \sec^3(e + fx) (3 - 4 \sec^2(e + fx)) dx = -\frac{\sec^3(e + fx) \tan(e + fx)}{f}$$

output `-sec(f*x+e)^3*tan(f*x+e)/f`

3.28.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^3(e + fx) (3 - 4 \sec^2(e + fx)) dx = -\frac{\sec^3(e + fx) \tan(e + fx)}{f}$$

input `Integrate[Sec[e + f*x]^3*(3 - 4*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^3*Tan[e + f*x])/f)`

3.28.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^3(e + fx) (3 - 4 \sec^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right)^3 \left(3 - 4 \csc\left(e + fx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{4531}$$

$$-\frac{\tan(e + fx) \sec^3(e + fx)}{f}$$

input `Int[Sec[e + f*x]^3*(3 - 4*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^3*Tan[e + f*x])/f)`

3.28.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.28.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.89

method	result
parallelrisc	$-\frac{8 \sin(fx+e)}{f(\cos(4fx+4e)+4 \cos(2fx+2e)+3)}$
risc	$\frac{8i(e^{5i(fx+e)}-e^{3i(fx+e)})}{f(e^{2i(fx+e)}+1)^4}$
derivativedivides	$\frac{\frac{3 \sec(fx+e) \tan(fx+e)}{2} + 4 \left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e)}{f}$
default	$\frac{\frac{3 \sec(fx+e) \tan(fx+e)}{2} + 4 \left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e)}{f}$
norman	$\frac{\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f} - \frac{6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{f} - \frac{6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f} - \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{f}}{\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)^4}$
parts	$\frac{3 \sec(fx+e) \tan(fx+e)}{2f} + \frac{3 \ln(\sec(fx+e)+\tan(fx+e))}{2f} - \frac{4 \left(-\left(-\frac{\sec(fx+e)^3}{4} - \frac{3 \sec(fx+e)}{8} \right) \tan(fx+e) + \frac{3 \ln(\sec(fx+e)+\tan(fx+e))}{2f} \right)}{f}$

input `int(sec(f*x+e)^3*(3-4*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `-8/f*sin(f*x+e)/(cos(4*f*x+4*e)+4*cos(2*f*x+2*e)+3)`

3.28.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^3(e + fx) (3 - 4 \sec^2(e + fx)) dx = -\frac{\sin(fx + e)}{f \cos(fx + e)^4}$$

input `integrate(sec(f*x+e)^3*(3-4*sec(f*x+e)^2),x, algorithm="fricas")`

output `-sin(f*x + e)/(f*cos(f*x + e)^4)`

3.28.6 Sympy [F]

$$\int \sec^3(e + fx) (3 - 4\sec^2(e + fx)) dx = - \int (-3\sec^3(e + fx)) dx - \int 4\sec^5(e + fx) dx$$

input `integrate(sec(f*x+e)**3*(3-4*sec(f*x+e)**2),x)`

output `-Integral(-3*sec(e + f*x)**3, x) - Integral(4*sec(e + f*x)**5, x)`

3.28.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.74

$$\int \sec^3(e + fx) (3 - 4\sec^2(e + fx)) dx = -\frac{\sin(fx + e)}{(\sin(fx + e)^4 - 2\sin(fx + e)^2 + 1)f}$$

input `integrate(sec(f*x+e)^3*(3-4*sec(f*x+e)^2),x, algorithm="maxima")`

output `-sin(f*x + e)/((sin(f*x + e)^4 - 2*sin(f*x + e)^2 + 1)*f)`

3.28.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.21

$$\int \sec^3(e + fx) (3 - 4\sec^2(e + fx)) dx = -\frac{\sin(fx + e)}{(\sin(fx + e)^2 - 1)^2 f}$$

input `integrate(sec(f*x+e)^3*(3-4*sec(f*x+e)^2),x, algorithm="giac")`

output `-sin(f*x + e)/((sin(f*x + e)^2 - 1)^2*f)`

3.28.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.21

$$\int \sec^3(e + fx) (3 - 4 \sec^2(e + fx)) dx = -\frac{\sin(e + fx)}{f (\sin(e + fx)^2 - 1)^2}$$

input `int(-(4/cos(e + f*x)^2 - 3)/cos(e + f*x)^3,x)`

output `-sin(e + f*x)/(f*(sin(e + f*x)^2 - 1)^2)`

3.29 $\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx$

3.29.1	Optimal result	197
3.29.2	Mathematica [A] (verified)	197
3.29.3	Rubi [A] (verified)	198
3.29.4	Maple [A] (verified)	199
3.29.5	Fricas [A] (verification not implemented)	199
3.29.6	Sympy [F]	200
3.29.7	Maxima [A] (verification not implemented)	200
3.29.8	Giac [A] (verification not implemented)	200
3.29.9	Mupad [B] (verification not implemented)	201

3.29.1 Optimal result

Integrand size = 21, antiderivative size = 19

$$\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx = -\frac{\sec^2(e + fx) \tan(e + fx)}{f}$$

output `-sec(f*x+e)^2*tan(f*x+e)/f`

3.29.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx = -\frac{\sec^2(e + fx) \tan(e + fx)}{f}$$

input `Integrate[Sec[e + f*x]^2*(2 - 3*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^2*Tan[e + f*x])/f)`

3.29.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right)^2 \left(2 - 3 \csc\left(e + fx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{4531}$$

$$-\frac{\tan(e + fx) \sec^2(e + fx)}{f}$$

input `Int[Sec[e + f*x]^2*(2 - 3*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]^2*Tan[e + f*x])/f)`

3.29.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.29.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.68

method	result	size
parallelrisc	$-\frac{4 \sin(fx+e)}{f(\cos(3fx+3e)+3 \cos(fx+e))}$	32
derivativedivides	$\frac{2 \tan(fx+e)+3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right) \tan(fx+e)}{f}$	34
default	$\frac{2 \tan(fx+e)+3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right) \tan(fx+e)}{f}$	34
parts	$\frac{2 \tan(fx+e)}{f} + \frac{3\left(-\frac{2}{3}-\frac{\sec(fx+e)^2}{3}\right) \tan(fx+e)}{f}$	36
risch	$\frac{4i(e^{4i(fx+e)}-e^{2i(fx+e)})}{f(e^{2i(fx+e)}+1)^3}$	41
norman	$\frac{\frac{2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{f} + \frac{4 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{f} + \frac{2 \tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{f}}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2-1\right)^3}$	64

input `int(sec(f*x+e)^2*(2-3*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`output `-4/f*sin(f*x+e)/(cos(3*f*x+3*e)+3*cos(f*x+e))`**3.29.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^2(e+fx)(2-3\sec^2(e+fx)) dx = -\frac{\sin(fx+e)}{f \cos(fx+e)^3}$$

input `integrate(sec(f*x+e)^2*(2-3*sec(f*x+e)^2),x, algorithm="fricas")`output `-sin(f*x + e)/(f*cos(f*x + e)^3)`

3.29.6 Sympy [F]

$$\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx = - \int (-2 \sec^2(e + fx)) dx - \int 3 \sec^4(e + fx) dx$$

input `integrate(sec(f*x+e)**2*(2-3*sec(f*x+e)**2),x)`

output `-Integral(-2*sec(e + f*x)**2, x) - Integral(3*sec(e + f*x)**4, x)`

3.29.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx = -\frac{\tan(fx + e)^3 + \tan(fx + e)}{f}$$

input `integrate(sec(f*x+e)^2*(2-3*sec(f*x+e)^2),x, algorithm="maxima")`

output `-(tan(f*x + e)^3 + tan(f*x + e))/f`

3.29.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \sec^2(e + fx) (2 - 3 \sec^2(e + fx)) dx = -\frac{\tan(fx + e)^3 + \tan(fx + e)}{f}$$

input `integrate(sec(f*x+e)^2*(2-3*sec(f*x+e)^2),x, algorithm="giac")`

output `-(tan(f*x + e)^3 + tan(f*x + e))/f`

3.29.9 Mupad [B] (verification not implemented)

Time = 15.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \sec^2(e + fx) (2 - 3\sec^2(e + fx)) dx = -\frac{\tan(e + fx) (\tan(e + fx)^2 + 1)}{f}$$

input `int(-(3/cos(e + f*x)^2 - 2)/cos(e + f*x)^2,x)`

output `-(tan(e + f*x)*(tan(e + f*x)^2 + 1))/f`

3.30 $\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx$

3.30.1	Optimal result	202
3.30.2	Mathematica [A] (verified)	202
3.30.3	Rubi [A] (verified)	203
3.30.4	Maple [A] (verified)	204
3.30.5	Fricas [A] (verification not implemented)	204
3.30.6	Sympy [F]	205
3.30.7	Maxima [A] (verification not implemented)	205
3.30.8	Giac [A] (verification not implemented)	205
3.30.9	Mupad [B] (verification not implemented)	206

3.30.1 Optimal result

Integrand size = 19, antiderivative size = 17

$$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx = -\frac{\sec(e + fx) \tan(e + fx)}{f}$$

output `-sec(f*x+e)*tan(f*x+e)/f`

3.30.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx = -\frac{\sec(e + fx) \tan(e + fx)}{f}$$

input `Integrate[Sec[e + f*x]*(1 - 2*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]*Tan[e + f*x])/f)`

3.30.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(e + fx + \frac{\pi}{2}\right) \left(1 - 2 \csc\left(e + fx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{4531}$$

$$-\frac{\tan(e + fx) \sec(e + fx)}{f}$$

input `Int[Sec[e + f*x]*(1 - 2*Sec[e + f*x]^2),x]`

output `-((Sec[e + f*x]*Tan[e + f*x])/f)`

3.30.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.30.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

method	result	size
derivativedivides	$-\frac{\sec(fx+e)\tan(fx+e)}{f}$	18
default	$-\frac{\sec(fx+e)\tan(fx+e)}{f}$	18
parts	$-\frac{\sec(fx+e)\tan(fx+e)}{f}$	18
parallelrisch	$-\frac{2\sin(fx+e)}{f(1+\cos(2fx+2e))}$	25
risch	$\frac{2i(e^{3i(fx+e)}-e^{i(fx+e)})}{f(e^{2i(fx+e)}+1)^2}$	41
norman	$-\frac{2\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{f}-\frac{2\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{\left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2-1\right)^2}$	48

input `int(sec(f*x+e)*(1-2*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`output `-sec(f*x+e)*tan(f*x+e)/f`**3.30.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx = -\frac{\sin(fx + e)}{f \cos(fx + e)^2}$$

input `integrate(sec(f*x+e)*(1-2*sec(f*x+e)^2),x, algorithm="fracas")`output `-sin(f*x + e)/(f*cos(f*x + e)^2)`

3.30.6 Sympy [F]

$$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx = - \int (-\sec(e + fx)) dx - \int 2 \sec^3(e + fx) dx$$

input `integrate(sec(f*x+e)*(1-2*sec(f*x+e)**2),x)`

output `-Integral(-sec(e + f*x), x) - Integral(2*sec(e + f*x)**3, x)`

3.30.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx = \frac{\sin(fx + e)}{(\sin(fx + e))^2 - 1} f$$

input `integrate(sec(f*x+e)*(1-2*sec(f*x+e)^2),x, algorithm="maxima")`

output `sin(f*x + e)/((sin(f*x + e)^2 - 1)*f)`

3.30.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.41

$$\int \sec(e + fx) (1 - 2 \sec^2(e + fx)) dx = - \frac{1}{f \left(\frac{1}{\sin(fx+e)} - \sin(fx + e) \right)}$$

input `integrate(sec(f*x+e)*(1-2*sec(f*x+e)^2),x, algorithm="giac")`

output `-1/(f*(1/sin(f*x + e) - sin(f*x + e)))`

3.30.9 Mupad [B] (verification not implemented)

Time = 15.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \sec(e + fx) (1 - 2\sec^2(e + fx)) dx = \frac{\sin(e + fx)}{f (\sin(e + fx)^2 - 1)}$$

input `int(-(2/cos(e + f*x)^2 - 1)/cos(e + f*x),x)`

output `sin(e + f*x)/(f*(sin(e + f*x)^2 - 1))`

3.31 $\int -\sec^2(e + fx) dx$

3.31.1	Optimal result	207
3.31.2	Mathematica [A] (verified)	207
3.31.3	Rubi [A] (verified)	208
3.31.4	Maple [A] (verified)	209
3.31.5	Fricas [A] (verification not implemented)	209
3.31.6	Sympy [F]	210
3.31.7	Maxima [A] (verification not implemented)	210
3.31.8	Giac [A] (verification not implemented)	210
3.31.9	Mupad [B] (verification not implemented)	211

3.31.1 Optimal result

Integrand size = 10, antiderivative size = 11

$$\int -\sec^2(e + fx) dx = -\frac{\tan(e + fx)}{f}$$

output `-tan(f*x+e)/f`

3.31.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\sec^2(e + fx) dx = -\frac{\tan(e + fx)}{f}$$

input `Integrate[-Sec[e + f*x]^2,x]`

output `-(Tan[e + f*x]/f)`

3.31.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {25, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int -\sec^2(e + fx) dx \\
 \downarrow 25 \\
 -\int \sec^2(e + fx) dx \\
 \downarrow 3042 \\
 -\int \csc\left(e + fx + \frac{\pi}{2}\right)^2 dx \\
 \downarrow 4254 \\
 \frac{\int 1d(-\tan(e + fx))}{f} \\
 \downarrow 24 \\
 -\frac{\tan(e + fx)}{f}
 \end{array}$$

input `Int[-Sec[e + f*x]^2,x]`

output `-(Tan[e + f*x]/f)`

3.31.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

3.31.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$-\frac{\tan(fx+e)}{f}$	12
default	$-\frac{\tan(fx+e)}{f}$	12
risch	$-\frac{2i}{f(e^{2i(fx+e)}+1)}$	20
norman	$\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)}$	30
parallelrisc	$\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f\left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)}$	30

```
input int(-sec(f*x+e)^2,x,method=_RETURNVERBOSE)
```

```
output -tan(f*x+e)/f
```

3.31.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int -\sec^2(e + fx) dx = -\frac{\sin(fx + e)}{f \cos(fx + e)}$$

```
input integrate(-sec(f*x+e)^2,x, algorithm="fricas")
```

```
output -sin(f*x + e)/(f*cos(f*x + e))
```

3.31.6 Sympy [F]

$$\int -\sec^2(e + fx) dx = -\int \sec^2(e + fx) dx$$

input `integrate(-sec(f*x+e)**2,x)`

output `-Integral(sec(e + f*x)**2, x)`

3.31.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\sec^2(e + fx) dx = -\frac{\tan(fx + e)}{f}$$

input `integrate(-sec(f*x+e)^2,x, algorithm="maxima")`

output `-tan(f*x + e)/f`

3.31.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\sec^2(e + fx) dx = -\frac{\tan(fx + e)}{f}$$

input `integrate(-sec(f*x+e)^2,x, algorithm="giac")`

output `-tan(f*x + e)/f`

3.31.9 Mupad [B] (verification not implemented)

Time = 15.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\sec^2(e + fx) dx = -\frac{\tan(e + fx)}{f}$$

input `int(-1/cos(e + f*x)^2,x)`

output `-tan(e + f*x)/f`

3.32 $\int -\cos(e + fx) dx$

3.32.1	Optimal result	212
3.32.2	Mathematica [B] (verified)	212
3.32.3	Rubi [A] (verified)	213
3.32.4	Maple [A] (verified)	214
3.32.5	Fricas [A] (verification not implemented)	214
3.32.6	Sympy [A] (verification not implemented)	215
3.32.7	Maxima [A] (verification not implemented)	215
3.32.8	Giac [A] (verification not implemented)	215
3.32.9	Mupad [B] (verification not implemented)	216

3.32.1 Optimal result

Integrand size = 8, antiderivative size = 11

$$\int -\cos(e + fx) dx = -\frac{\sin(e + fx)}{f}$$

output `-sin(f*x+e)/f`

3.32.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. 2(11) = 22.

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.09

$$\int -\cos(e + fx) dx = -\frac{\cos(fx) \sin(e)}{f} - \frac{\cos(e) \sin(fx)}{f}$$

input `Integrate[-Cos[e + f*x],x]`

output `-((Cos[f*x]*Sin[e])/f) - (Cos[e]*Sin[f*x])/f`

3.32.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {25, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int -\cos(e + fx) dx \\ & \quad \downarrow \text{25} \\ & - \int \cos(e + fx) dx \\ & \quad \downarrow \text{3042} \\ & - \int \sin\left(e + fx + \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{3117} \\ & - \frac{\sin(e + fx)}{f} \end{aligned}$$

input `Int[-Cos[e + f*x],x]`

output `-(Sin[e + f*x]/f)`

3.32.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

3.32.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$-\frac{\sin(fx+e)}{f}$	12
default	$-\frac{\sin(fx+e)}{f}$	12
risch	$-\frac{\sin(fx+e)}{f}$	12
parallelrisch	$-\frac{\sin(fx+e)}{f}$	12
parts	$-\frac{\sin(fx+e)}{f}$	12
norman	$-\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f \left(1 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)\right)^2}$	30
meijerg	$-\frac{\cos(e) \sin(fx)}{f} + \frac{\sin(e) \sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(fx)}{\sqrt{\pi}}\right)}{f}$	35

input `int(-cos(f*x+e),x,method=_RETURNVERBOSE)`output `-sin(f*x+e)/f`**3.32.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\cos(e + fx) dx = -\frac{\sin(fx + e)}{f}$$

input `integrate(-cos(f*x+e),x, algorithm="fricas")`output `-sin(f*x + e)/f`

3.32.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.27

$$\int -\cos(e + fx) dx = -\begin{cases} \frac{\sin(e+fx)}{f} & \text{for } f \neq 0 \\ x \cos(e) & \text{otherwise} \end{cases}$$

input `integrate(-cos(f*x+e),x)`output `-Piecewise((sin(e + f*x)/f, Ne(f, 0)), (x*cos(e), True))`**3.32.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\cos(e + fx) dx = -\frac{\sin(fx + e)}{f}$$

input `integrate(-cos(f*x+e),x, algorithm="maxima")`output `-sin(f*x + e)/f`**3.32.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\cos(e + fx) dx = -\frac{\sin(fx + e)}{f}$$

input `integrate(-cos(f*x+e),x, algorithm="giac")`output `-sin(f*x + e)/f`

3.32.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int -\cos(e + fx) dx = -\frac{\sin(e + fx)}{f}$$

input `int(-cos(e + f*x),x)`

output `-sin(e + f*x)/f`

3.33 $\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx$

3.33.1	Optimal result	217
3.33.2	Mathematica [A] (verified)	217
3.33.3	Rubi [A] (verified)	218
3.33.4	Maple [A] (verified)	219
3.33.5	Fricas [A] (verification not implemented)	219
3.33.6	Sympy [A] (verification not implemented)	219
3.33.7	Maxima [A] (verification not implemented)	220
3.33.8	Giac [A] (verification not implemented)	220
3.33.9	Mupad [B] (verification not implemented)	220

3.33.1 Optimal result

Integrand size = 19, antiderivative size = 17

$$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx = -\frac{\cos(e + fx) \sin(e + fx)}{f}$$

output `-cos(f*x+e)*sin(f*x+e)/f`

3.33.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.94

$$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx = -\frac{\cos(2fx) \sin(2e)}{2f} - \frac{\cos(2e) \sin(2fx)}{2f}$$

input `Integrate[Cos[e + f*x]^2*(-2 + Sec[e + f*x]^2),x]`

output `-1/2*(Cos[2*f*x]*Sin[2*e])/f - (Cos[2*e]*Sin[2*f*x])/(2*f)`

3.33.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(e + fx) (\sec^2(e + fx) - 2) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(e + fx + \frac{\pi}{2})^2 - 2}{\csc(e + fx + \frac{\pi}{2})^2} dx$$

$$\downarrow \text{4531}$$

$$-\frac{\sin(e + fx) \cos(e + fx)}{f}$$

input `Int[Cos[e + f*x]^2*(-2 + Sec[e + f*x]^2), x]`

output `-((Cos[e + f*x]*Sin[e + f*x])/f)`

3.33.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.33.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

method	result	size
risch	$-\frac{\sin(2fx+2e)}{2f}$	15
parallelrisch	$-\frac{\sin(2fx+2e)}{2f}$	15
derivativdivides	$-\frac{\cos(fx+e)\sin(fx+e)}{f}$	18
default	$-\frac{\cos(fx+e)\sin(fx+e)}{f}$	18
norman	$\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right) - 4 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3 + 2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{\left(1 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2\right)^2 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)}$	79

input `int(cos(f*x+e)^2*(-2+sec(f*x+e)^2),x,method=_RETURNVERBOSE)`output `-1/2/f*sin(2*f*x+2*e)`**3.33.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx = -\frac{\cos(fx + e)\sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^2*(-2+sec(f*x+e)^2),x, algorithm="fricas")`output `-cos(f*x + e)*sin(f*x + e)/f`**3.33.6 Sympy [A] (verification not implemented)**

Time = 3.63 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.88

$$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx$$

$$= x - 2 \left(\begin{cases} \frac{x \sin^2(e+fx)}{2} + \frac{x \cos^2(e+fx)}{2} + \frac{\sin(e+fx)\cos(e+fx)}{2f} & \text{for } f \neq 0 \\ x \cos^2(e) & \text{otherwise} \end{cases} \right)$$

input `integrate(cos(f*x+e)**2*(-2+sec(f*x+e)**2),x)`

output `x - 2*Piecewise((x*sin(e + f*x)**2/2 + x*cos(e + f*x)**2/2 + sin(e + f*x)*cos(e + f*x)/(2*f), Ne(f, 0)), (x*cos(e)**2, True))`

3.33.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx = -\frac{\tan(fx + e)}{(\tan(fx + e)^2 + 1)f}$$

input `integrate(cos(f*x+e)^2*(-2+sec(f*x+e)^2),x, algorithm="maxima")`

output `-tan(f*x + e)/((tan(f*x + e)^2 + 1)*f)`

3.33.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx = -\frac{\sin(2fx + 2e)}{2f}$$

input `integrate(cos(f*x+e)^2*(-2+sec(f*x+e)^2),x, algorithm="giac")`

output `-1/2*sin(2*f*x + 2*e)/f`

3.33.9 Mupad [B] (verification not implemented)

Time = 15.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cos^2(e + fx) (-2 + \sec^2(e + fx)) dx = -\frac{\sin(2e + 2fx)}{2f}$$

input `int(cos(e + f*x)^2*(1/cos(e + f*x)^2 - 2),x)`

output `-sin(2*e + 2*f*x)/(2*f)`

3.34 $\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx$

3.34.1	Optimal result	221
3.34.2	Mathematica [B] (verified)	221
3.34.3	Rubi [A] (verified)	222
3.34.4	Maple [A] (verified)	223
3.34.5	Fricas [A] (verification not implemented)	223
3.34.6	Sympy [F]	224
3.34.7	Maxima [A] (verification not implemented)	224
3.34.8	Giac [A] (verification not implemented)	224
3.34.9	Mupad [B] (verification not implemented)	225

3.34.1 Optimal result

Integrand size = 21, antiderivative size = 19

$$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx = -\frac{\cos^2(e + fx) \sin(e + fx)}{f}$$

output `-cos(f*x+e)^2*sin(f*x+e)/f`

3.34.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 46 vs. 2(19) = 38.

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.42

$$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx = \frac{2 \cos(fx) \sin(e)}{f} + \frac{2 \cos(e) \sin(fx)}{f} - \frac{3 \sin(e + fx)}{f} + \frac{\sin^3(e + fx)}{f}$$

input `Integrate[Cos[e + f*x]^3*(-3 + 2*Sec[e + f*x]^2),x]`

output `(2*Cos[f*x]*Sin[e])/f + (2*Cos[e]*Sin[f*x])/f - (3*Sine + f*x])/f + Sin[e + f*x]^3/f`

3.34.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(e + fx) (2 \sec^2(e + fx) - 3) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{2 \csc(e + fx + \frac{\pi}{2})^2 - 3}{\csc(e + fx + \frac{\pi}{2})^3} dx$$

$$\downarrow \text{4531}$$

$$\frac{\sin(e + fx) \cos^2(e + fx)}{f}$$

input `Int[Cos[e + f*x]^3*(-3 + 2*Sec[e + f*x]^2),x]`

output `-((Cos[e + f*x]^2*Sin[e + f*x])/f)`

3.34.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.34.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.37

method	result	size
parallelrisch	$\frac{-\sin(fx+e)-\sin(3fx+3e)}{4f}$	26
risch	$\frac{-\sin(fx+e)}{4f} - \frac{\sin(3fx+3e)}{4f}$	27
derivativedivides	$\frac{-(2+\cos(fx+e)^2)\sin(fx+e)+2\sin(fx+e)}{f}$	32
default	$\frac{-(2+\cos(fx+e)^2)\sin(fx+e)+2\sin(fx+e)}{f}$	32
norman	$\frac{\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f} - \frac{6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{f} + \frac{6 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f} - \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{f}}{\left(1 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2\right)^3 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)}$	95

input `int(cos(f*x+e)^3*(-3+2*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`output `1/4*(-sin(f*x+e)-sin(3*f*x+3*e))/f`**3.34.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx = -\frac{\cos(fx + e)^2 \sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^3*(-3+2*sec(f*x+e)^2),x, algorithm="fricas")`output `-cos(f*x + e)^2*sin(f*x + e)/f`

3.34.6 Sympy [F]

$$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx = \int (2 \sec^2(e + fx) - 3) \cos^3(e + fx) dx$$

input `integrate(cos(f*x+e)**3*(-3+2*sec(f*x+e)**2),x)`

output `Integral((2*sec(e + f*x)**2 - 3)*cos(e + f*x)**3, x)`

3.34.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx = \frac{\sin(fx + e)^3 - \sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^3*(-3+2*sec(f*x+e)^2),x, algorithm="maxima")`

output `(sin(f*x + e)^3 - sin(f*x + e))/f`

3.34.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx = \frac{\sin(fx + e)^3 - \sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^3*(-3+2*sec(f*x+e)^2),x, algorithm="giac")`

output `(sin(f*x + e)^3 - sin(f*x + e))/f`

3.34.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \cos^3(e + fx) (-3 + 2 \sec^2(e + fx)) dx = -\frac{\sin(e + fx) - \sin(e + fx)^3}{f}$$

input `int(cos(e + f*x)^3*(2/cos(e + f*x)^2 - 3),x)`

output `-(sin(e + f*x) - sin(e + f*x)^3)/f`

3.35 $\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx$

3.35.1	Optimal result	226
3.35.2	Mathematica [A] (verified)	226
3.35.3	Rubi [A] (verified)	227
3.35.4	Maple [A] (verified)	228
3.35.5	Fricas [A] (verification not implemented)	228
3.35.6	Sympy [F]	229
3.35.7	Maxima [A] (verification not implemented)	229
3.35.8	Giac [A] (verification not implemented)	229
3.35.9	Mupad [B] (verification not implemented)	230

3.35.1 Optimal result

Integrand size = 21, antiderivative size = 19

$$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx = -\frac{\cos^3(e + fx) \sin(e + fx)}{f}$$

output `-cos(f*x+e)^3*sin(f*x+e)/f`

3.35.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.63

$$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx = -\frac{\sin(2(e + fx))}{4f} - \frac{\sin(4(e + fx))}{8f}$$

input `Integrate[Cos[e + f*x]^4*(-4 + 3*Sec[e + f*x]^2),x]`

output `-1/4*Sin[2*(e + f*x)]/f - Sin[4*(e + f*x)]/(8*f)`

3.35.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^4(e + fx) (3 \sec^2(e + fx) - 4) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{3 \csc(e + fx + \frac{\pi}{2})^2 - 4}{\csc(e + fx + \frac{\pi}{2})^4} dx$$

$$\downarrow \text{4531}$$

$$\frac{\sin(e + fx) \cos^3(e + fx)}{f}$$

input `Int[Cos[e + f*x]^4*(-4 + 3*Sec[e + f*x]^2),x]`

output `-((Cos[e + f*x]^3*Sin[e + f*x])/f)`

3.35.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.35.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.53

method	result	size
parallelrisc	$\frac{-\sin(4fx+4e)-2\sin(2fx+2e)}{8f}$	29
risc	$-\frac{\sin(4fx+4e)}{8f} - \frac{\sin(2fx+2e)}{4f}$	30
derivativedivides	$\frac{-\left(\cos(fx+e)^3 + \frac{3\cos(fx+e)}{2}\right)\sin(fx+e) + \frac{3\cos(fx+e)\sin(fx+e)}{2}}{f}$	45
default	$\frac{-\left(\cos(fx+e)^3 + \frac{3\cos(fx+e)}{2}\right)\sin(fx+e) + \frac{3\cos(fx+e)\sin(fx+e)}{2}}{f}$	45
norman	$\frac{\frac{2\tan\left(\frac{fx}{2}+\frac{e}{2}\right)}{f} - \frac{8\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^3}{f} + \frac{12\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^5}{f} - \frac{8\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^7}{f} + \frac{2\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^9}{f}}{\left(1+\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2\right)^4 \left(\tan\left(\frac{fx}{2}+\frac{e}{2}\right)^2-1\right)}$	111

input `int(cos(f*x+e)^4*(-4+3*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `1/8*(-sin(4*f*x+4*e)-2*sin(2*f*x+2*e))/f`

3.35.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx = -\frac{\cos(fx + e)^3 \sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^4*(-4+3*sec(f*x+e)^2),x, algorithm="fricas")`

output `-cos(f*x + e)^3*sin(f*x + e)/f`

3.35.6 Sympy [F]

$$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx = \int (3 \sec^2(e + fx) - 4) \cos^4(e + fx) dx$$

input `integrate(cos(f*x+e)**4*(-4+3*sec(f*x+e)**2),x)`

output `Integral((3*sec(e + f*x)**2 - 4)*cos(e + f*x)**4, x)`

3.35.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.74

$$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx = -\frac{\tan(fx + e)}{(\tan(fx + e)^4 + 2 \tan(fx + e)^2 + 1)f}$$

input `integrate(cos(f*x+e)^4*(-4+3*sec(f*x+e)^2),x, algorithm="maxima")`

output `-tan(f*x + e)/((tan(f*x + e)^4 + 2*tan(f*x + e)^2 + 1)*f)`

3.35.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.21

$$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx = -\frac{\tan(fx + e)}{(\tan(fx + e)^2 + 1)^2 f}$$

input `integrate(cos(f*x+e)^4*(-4+3*sec(f*x+e)^2),x, algorithm="giac")`

output `-tan(f*x + e)/((tan(f*x + e)^2 + 1)^2*f)`

3.35.9 Mupad [B] (verification not implemented)

Time = 15.43 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \cos^4(e + fx) (-4 + 3 \sec^2(e + fx)) dx = -\frac{\cos(e + fx)^3 \sin(e + fx)}{f}$$

input `int(cos(e + f*x)^4*(3/cos(e + f*x)^2 - 4),x)`

output `-(cos(e + f*x)^3*sin(e + f*x))/f`

3.36 $\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx$

3.36.1	Optimal result	231
3.36.2	Mathematica [A] (verified)	231
3.36.3	Rubi [A] (verified)	232
3.36.4	Maple [A] (verified)	233
3.36.5	Fricas [A] (verification not implemented)	233
3.36.6	Sympy [F]	234
3.36.7	Maxima [A] (verification not implemented)	234
3.36.8	Giac [A] (verification not implemented)	234
3.36.9	Mupad [B] (verification not implemented)	235

3.36.1 Optimal result

Integrand size = 21, antiderivative size = 19

$$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx = -\frac{\cos^4(e + fx) \sin(e + fx)}{f}$$

output `-cos(f*x+e)^4*sin(f*x+e)/f`

3.36.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.00

$$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx = -\frac{\sin(e + fx)}{f} + \frac{2 \sin^3(e + fx)}{f} - \frac{\sin^5(e + fx)}{f}$$

input `Integrate[Cos[e + f*x]^5*(-5 + 4*Sec[e + f*x]^2),x]`

output `-(Sin[e + f*x]/f) + (2*SIN[e + f*x]^3)/f - Sin[e + f*x]^5/f`

3.36.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 4531}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^5(e + fx) (4 \sec^2(e + fx) - 5) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{4 \csc(e + fx + \frac{\pi}{2})^2 - 5}{\csc(e + fx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{4531}$$

$$\frac{\sin(e + fx) \cos^4(e + fx)}{f}$$

input `Int[Cos[e + f*x]^5*(-5 + 4*Sec[e + f*x]^2),x]`

output `-((Cos[e + f*x]^4*Sin[e + f*x])/f)`

3.36.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4531 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^m_.*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] /; FreeQ[{b, e, f, A, C, m}, x] && EqQ[C*m + A*(m + 1), 0]`

3.36.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.95

method	result	size
parallelrisc	$\frac{-2 \sin(fx+e) - \sin(5fx+5e) - 3 \sin(3fx+3e)}{16f}$	37
risc	$-\frac{\sin(fx+e)}{8f} - \frac{\sin(5fx+5e)}{16f} - \frac{3 \sin(3fx+3e)}{16f}$	41
derivativedivides	$-\frac{\left(\frac{8}{3} + \cos(fx+e)^4 + \frac{4 \cos(fx+e)^2}{3}\right) \sin(fx+e) + \frac{4(2 + \cos(fx+e)^2) \sin(fx+e)}{3}}{f}$	52
default	$-\frac{\left(\frac{8}{3} + \cos(fx+e)^4 + \frac{4 \cos(fx+e)^2}{3}\right) \sin(fx+e) + \frac{4(2 + \cos(fx+e)^2) \sin(fx+e)}{3}}{f}$	52
norman	$\frac{\frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)}{f} - \frac{10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^3}{f} + \frac{20 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^5}{f} - \frac{20 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^7}{f} + \frac{10 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^9}{f} - \frac{2 \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^{11}}{f}}{\left(1 + \tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2\right)^5 \left(\tan\left(\frac{fx}{2} + \frac{e}{2}\right)^2 - 1\right)}$	127

input `int(cos(f*x+e)^5*(-5+4*sec(f*x+e)^2),x,method=_RETURNVERBOSE)`output `1/16*(-2*sin(f*x+e)-sin(5*f*x+5*e)-3*sin(3*f*x+3*e))/f`**3.36.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx = -\frac{\cos(fx + e)^4 \sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^5*(-5+4*sec(f*x+e)^2),x, algorithm="fricas")`output `-cos(f*x + e)^4*sin(f*x + e)/f`

3.36.6 Sympy [F]

$$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx = \int (4 \sec^2(e + fx) - 5) \cos^5(e + fx) dx$$

input `integrate(cos(f*x+e)**5*(-5+4*sec(f*x+e)**2),x)`

output `Integral((4*sec(e + f*x)**2 - 5)*cos(e + f*x)**5, x)`

3.36.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.58

$$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx = -\frac{\sin(fx + e)^5 - 2 \sin(fx + e)^3 + \sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^5*(-5+4*sec(f*x+e)^2),x, algorithm="maxima")`

output `-(sin(f*x + e)^5 - 2*sin(f*x + e)^3 + sin(f*x + e))/f`

3.36.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.58

$$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx = -\frac{\sin(fx + e)^5 - 2 \sin(fx + e)^3 + \sin(fx + e)}{f}$$

input `integrate(cos(f*x+e)^5*(-5+4*sec(f*x+e)^2),x, algorithm="giac")`

output `-(sin(f*x + e)^5 - 2*sin(f*x + e)^3 + sin(f*x + e))/f`

3.36.9 Mupad [B] (verification not implemented)

Time = 15.53 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.21

$$\int \cos^5(e + fx) (-5 + 4 \sec^2(e + fx)) dx = -\frac{\sin(e + fx) (\sin(e + fx)^2 - 1)^2}{f}$$

input `int(cos(e + f*x)^5*(4/cos(e + f*x)^2 - 5),x)`

output `-(sin(e + f*x)*(sin(e + f*x)^2 - 1)^2)/f`

3.37 $\int \sec^3(c+dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.37.1	Optimal result	236
3.37.2	Mathematica [A] (verified)	236
3.37.3	Rubi [A] (verified)	237
3.37.4	Maple [A] (verified)	239
3.37.5	Fricas [A] (verification not implemented)	240
3.37.6	Sympy [F]	240
3.37.7	Maxima [A] (verification not implemented)	240
3.37.8	Giac [B] (verification not implemented)	241
3.37.9	Mupad [B] (verification not implemented)	241

3.37.1 Optimal result

Integrand size = 28, antiderivative size = 85

$$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3C \arctanh(\sin(c + dx))}{8d} + \frac{B \tan(c + dx)}{d} + \frac{3C \sec(c + dx) \tan(c + dx)}{8d}$$

$$+ \frac{C \sec^3(c + dx) \tan(c + dx)}{4d} + \frac{B \tan^3(c + dx)}{3d}$$

output `3/8*C*arctanh(sin(d*x+c))/d+B*tan(d*x+c)/d+3/8*C*sec(d*x+c)*tan(d*x+c)/d+1/4*C*sec(d*x+c)^3*tan(d*x+c)/d+1/3*B*tan(d*x+c)^3/d`

3.37.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.69

$$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{9C \arctanh(\sin(c + dx)) + \tan(c + dx) (9C \sec(c + dx) + 6C \sec^3(c + dx) + 8B(3 + \tan^2(c + dx)))}{24d}$$

input `Integrate[Sec[c + d*x]^3*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(9*C*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(9*C*Sec[c + d*x] + 6*C*Sec[c + d*x]^3 + 8*B*(3 + Tan[c + d*x]^2)))/(24*d)`

3.37.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.393$, Rules used = {3042, 4535, 27, 3042, 4254, 2009, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 \left(B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \sec^4(c+dx) dx + \int C \sec^5(c+dx) dx \\
 & \quad \downarrow \text{27} \\
 & B \int \sec^4(c+dx) dx + C \int \sec^5(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 dx + C \int \csc\left(c+dx+\frac{\pi}{2}\right)^5 dx \\
 & \quad \downarrow \text{4254} \\
 & C \int \csc\left(c+dx+\frac{\pi}{2}\right)^5 dx - \frac{B \int (\tan^2(c+dx) + 1) d(-\tan(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & C \int \csc\left(c+dx+\frac{\pi}{2}\right)^5 dx - \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
 & \quad \downarrow \text{4255} \\
 & C \left(\frac{3}{4} \int \sec^3(c+dx) dx + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right) - \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & C \left(\frac{3}{4} \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right) - \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
 & \quad \downarrow \text{4255}
 \end{aligned}$$

$$\begin{aligned}
& C \left(\frac{3}{4} \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right) - \\
& \quad \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
& \quad \downarrow \text{3042} \\
& C \left(\frac{3}{4} \left(\frac{1}{2} \int \csc\left(c+dx + \frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right) - \\
& \quad \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
& \quad \downarrow \text{4257} \\
& C \left(\frac{3}{4} \left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d} \right) - \\
& \quad \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d}
\end{aligned}$$

input `Int[Sec[c + d*x]^3*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `-((B*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/d) + C*((Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (3*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/4)`

3.37.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.37.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{-B\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)+C\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d}$
default	$\frac{-B\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)+C\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d}$
parts	$-\frac{B\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)}{d} + \frac{C\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d}$
risch	$-\frac{i(9C e^{7i(dx+c)}+33C e^{5i(dx+c)}-48B e^{4i(dx+c)}-33C e^{3i(dx+c)}-64B e^{2i(dx+c)}-9C e^{i(dx+c)}-16B)}{12d(e^{2i(dx+c)}+1)^4} + \frac{3\ln(e^{i(dx+c)}+1)}{8d}$
norman	$-\frac{(8B-5C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^7}{4d} + \frac{(8B+5C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{4d} - \frac{(40B-9C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{12d} + \frac{(40B+9C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^5}{12d} - \frac{3C\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{8d}$
parallelrisc	$\frac{-18\left(\frac{3}{4}+\frac{\cos(4dx+4c)}{4}\right)+\cos(2dx+2c)}{12d(\cos(4dx+4c)+4\cos(2dx+2c)+3)}C\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+18\left(\frac{3}{4}+\frac{\cos(4dx+4c)}{4}\right)+\cos(2dx+2c)}{12d(\cos(4dx+4c)+4\cos(2dx+2c)+3)}C\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)$

input `int(sec(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(-B*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+C*(-(1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c))))`

$$3.37. \quad \int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

3.37.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.16

$$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{9C \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 9C \cos(dx + c)^4 \log(-\sin(dx + c) + 1) + 2(16B \cos(dx + c)^3 + 9C \cos(dx + c)^2 + 8B \cos(dx + c) + 6C) \sin(dx + c)}{48d \cos(dx + c)^4}$$

```
input integrate(sec(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")
```

```
output 1/48*(9*C*cos(d*x + c)^4*log(sin(d*x + c) + 1) - 9*C*cos(d*x + c)^4*log(-sin(d*x + c) + 1) + 2*(16*B*cos(d*x + c)^3 + 9*C*cos(d*x + c)^2 + 8*B*cos(d*x + c) + 6*C)*sin(d*x + c))/(d*cos(d*x + c)^4)
```

3.37.6 Sympy [F]

$$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (B + C \sec(c + dx)) \sec^4(c + dx) dx$$

```
input integrate(sec(d*x+c)**3*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)
```

```
output Integral((B + C*sec(c + d*x))*sec(c + d*x)**4, x)
```

3.37.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.12

$$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{16(\tan(dx + c)^3 + 3 \tan(dx + c))B - 3C \left(\frac{2(3 \sin(dx + c)^3 - 5 \sin(dx + c))}{\sin(dx + c)^4 - 2 \sin(dx + c)^2 + 1} - 3 \log(\sin(dx + c) + 1) + 3 \log(\sin(dx + c) - 1) \right)}{48d}$$

input `integrate(sec(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output $\frac{1}{48}*(16*(\tan(dx + c))^3 + 3*\tan(dx + c))*B - 3*C*(2*(3*\sin(dx + c))^3 - 5*\sin(dx + c))/(\sin(dx + c)^4 - 2*\sin(dx + c)^2 + 1) - 3*\log(\sin(dx + c) + 1) + 3*\log(\sin(dx + c) - 1))/d$

3.37.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 164 vs. $2(77) = 154$.

Time = 0.31 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.93

$$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{9C \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 9C \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) - \frac{2(24B \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 15C \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 4)}{d}}{24d}$$

input `integrate(sec(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output $\frac{1}{24}*(9*C*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 9*C*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(24*B*\tan(1/2*d*x + 1/2*c)^7 - 15*C*\tan(1/2*d*x + 1/2*c)^7 - 40*B*\tan(1/2*d*x + 1/2*c)^5 - 9*C*\tan(1/2*d*x + 1/2*c)^5 + 40*B*\tan(1/2*d*x + 1/2*c)^3 - 9*C*\tan(1/2*d*x + 1/2*c)^3 - 24*B*\tan(1/2*d*x + 1/2*c) - 15*C*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^4/d$

3.37.9 Mupad [B] (verification not implemented)

Time = 18.03 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.80

$$\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{3C \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{4d}$$

$$- \frac{\left(2B - \frac{5C}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(-\frac{10B}{3} - \frac{3C}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{10B}{3} - \frac{3C}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(-2B - \frac{5C}{4}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)}$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/cos(c + d*x)^3,x)`

3.37. $\int \sec^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

output $(3*C*\operatorname{atanh}(\tan(c/2 + (d*x)/2)))/(4*d) - (\tan(c/2 + (d*x)/2)^7*(2*B - (5*C)/4) + \tan(c/2 + (d*x)/2)^3*((10*B)/3 - (3*C)/4) - \tan(c/2 + (d*x)/2)^5*((10*B)/3 + (3*C)/4) - \tan(c/2 + (d*x)/2)*(2*B + (5*C)/4))/(d*(6*\tan(c/2 + (d*x)/2)^4 - 4*\tan(c/2 + (d*x)/2)^2 - 4*\tan(c/2 + (d*x)/2)^6 + \tan(c/2 + (d*x)/2)^8 + 1))$

3.38 $\int \sec^2(c+dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.38.1	Optimal result	243
3.38.2	Mathematica [A] (verified)	243
3.38.3	Rubi [A] (verified)	244
3.38.4	Maple [A] (verified)	246
3.38.5	Fricas [A] (verification not implemented)	246
3.38.6	Sympy [F]	247
3.38.7	Maxima [A] (verification not implemented)	247
3.38.8	Giac [B] (verification not implemented)	247
3.38.9	Mupad [B] (verification not implemented)	248

3.38.1 Optimal result

Integrand size = 28, antiderivative size = 63

$$\int \sec^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{B \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{C \tan(c + dx)}{d} + \frac{B \sec(c + dx) \tan(c + dx)}{2d} + \frac{C \tan^3(c + dx)}{3d}$$

```
output 1/2*B*arctanh(sin(d*x+c))/d+C*tan(d*x+c)/d+1/2*B*sec(d*x+c)*tan(d*x+c)/d+1/3*C*tan(d*x+c)^3/d
```

3.38.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.95

$$\int \sec^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{B \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{B \sec(c + dx) \tan(c + dx)}{2d} + \frac{C(\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d}$$

```
input Integrate[Sec[c + d*x]^2*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]
```

```
output (B*ArcTanh[Sin[c + d*x]])/(2*d) + (B*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (C*(Tan[c + d*x] + Tan[c + d*x]^3/3))/d
```

3.38.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.321$, Rules used = {3042, 4535, 27, 3042, 4254, 2009, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 \left(B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \sec^3(c+dx) dx + \int C \sec^4(c+dx) dx \\
 & \quad \downarrow \text{27} \\
 & B \int \sec^3(c+dx) dx + C \int \sec^4(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx + C \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{4254} \\
 & B \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx - \frac{C \int (\tan^2(c+dx) + 1) d(-\tan(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & B \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx - \frac{C(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
 & \quad \downarrow \text{4255} \\
 & B \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{C(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{C(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

3.38. $\int \sec^2(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$

$$B \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{C(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d}$$

input `Int[Sec[c + d*x]^2*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `B*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)) - (C*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/d`

3.38.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4535 `Int[(csc[(e_) + (f_)*(x_)]*(b_))^(m_)*((A_) + csc[(e_) + (f_)*(x_)]*(B_) + csc[(e_) + (f_)*(x_)]^2*(C_)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.38.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.95

method	result
derivativedivides	$\frac{B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) - C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$
default	$\frac{B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) - C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$
parts	$\frac{B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d} - \frac{C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$
risch	$-\frac{i(3B e^{5i(dx+c)} - 12C e^{2i(dx+c)} - 3B e^{i(dx+c)} - 4C)}{3d(e^{2i(dx+c)} + 1)^3} + \frac{B \ln(e^{i(dx+c)} + i)}{2d} - \frac{B \ln(e^{i(dx+c)} - i)}{2d}$
norman	$\frac{(B-2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5 + \frac{4C \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{3d} - \frac{(B+2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d}}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)^3} - \frac{B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2d} + \frac{B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2d}$
parallelrisch	$\frac{-9 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 9 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 6B \sin(2dx+c)}{6d(\cos(3dx+3c) + 3 \cos(dx+c))}$

input `int(sec(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(B*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))-C*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c))`

3.38.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.40

$$\int \sec^2(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \frac{3B \cos(dx+c)^3 \log(\sin(dx+c)+1) - 3B \cos(dx+c)^3 \log(-\sin(dx+c)+1) + 2(4C \cos(dx+c)^2 - 3B \cos(dx+c) + 2C) \sin(dx+c)}{12d \cos(dx+c)^3}$$

input `integrate(sec(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/12*(3*B*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*B*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(4*C*cos(d*x + c)^2 + 3*B*cos(d*x + c) + 2*C)*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.38. $\int \sec^2(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$

3.38.6 Sympy [F]

$$\int \sec^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (B + C \sec(c + dx)) \sec^3(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((B + C*sec(c + d*x))*sec(c + d*x)**3, x)`

3.38.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11

$$\int \sec^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{4 (\tan(dx + c)^3 + 3 \tan(dx + c))C - 3B \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2 - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right)}{12d}$$

input `integrate(sec(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/12*(4*(tan(d*x + c)^3 + 3*tan(d*x + c))*C - 3*B*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)))/d`

3.38.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 122 vs. 2(57) = 114.

Time = 0.30 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.94

$$\int \sec^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3B \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 3B \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + \frac{2 \left(3B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^5 - 6C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^5 + 4C \right)}{\left(\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right)^2}{6d}$$

input `integrate(sec(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/6*(3*B*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*B*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + 2*(3*B*tan(1/2*d*x + 1/2*c)^5 - 6*C*tan(1/2*d*x + 1/2*c)^5 + 4*C*tan(1/2*d*x + 1/2*c)^3 - 3*B*tan(1/2*d*x + 1/2*c) - 6*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^3)/d`

3.38.9 Mupad [B] (verification not implemented)

Time = 17.12 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.73

$$\int \sec^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{B \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d}$$

$$+ \frac{(B - 2C) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \frac{4C \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{3} + (-B - 2C) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1 \right)}$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/cos(c + d*x)^2,x)`

output `(B*atanh(tan(c/2 + (d*x)/2)))/d + ((4*C*tan(c/2 + (d*x)/2)^3)/3 - tan(c/2 + (d*x)/2)*(B + 2*C) + tan(c/2 + (d*x)/2)^5*(B - 2*C))/(d*(3*tan(c/2 + (d*x)/2)^2 - 3*tan(c/2 + (d*x)/2)^4 + tan(c/2 + (d*x)/2)^6 - 1))`

3.39 $\int \sec(c+dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.39.1	Optimal result	249
3.39.2	Mathematica [A] (verified)	249
3.39.3	Rubi [A] (verified)	250
3.39.4	Maple [A] (verified)	252
3.39.5	Fricas [A] (verification not implemented)	252
3.39.6	Sympy [F]	253
3.39.7	Maxima [A] (verification not implemented)	253
3.39.8	Giac [B] (verification not implemented)	253
3.39.9	Mupad [B] (verification not implemented)	254

3.39.1 Optimal result

Integrand size = 26, antiderivative size = 47

$$\int \sec(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{C \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{B \tan(c + dx)}{d} + \frac{C \sec(c + dx) \tan(c + dx)}{2d}$$

output `1/2*C*arctanh(sin(d*x+c))/d+B*tan(d*x+c)/d+1/2*C*sec(d*x+c)*tan(d*x+c)/d`

3.39.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int \sec(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{C \operatorname{ArcTanh}[\sin(c + dx)]}{2d} + \frac{B \tan(c + dx)}{d} + \frac{C \sec(c + dx) \tan(c + dx)}{2d}$$

input `Integrate[Sec[c + d*x]*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(C*ArcTanh[Sin[c + d*x]])/(2*d) + (B*Tan[c + d*x])/d + (C*Sec[c + d*x]*Tan[c + d*x])/(2*d)`

3.39.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.346$, Rules used = {3042, 4535, 27, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right) \left(B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \sec^2(c+dx) dx + \int C \sec^3(c+dx) dx \\
 & \quad \downarrow \text{27} \\
 & B \int \sec^2(c+dx) dx + C \int \sec^3(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx + C \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{4254} \\
 & C \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx - \frac{B \int 1d(-\tan(c+dx))}{d} \\
 & \quad \downarrow \text{24} \\
 & C \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx + \frac{B \tan(c+dx)}{d} \\
 & \quad \downarrow \text{4255} \\
 & C \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{B \tan(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & C \left(\frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{B \tan(c+dx)}{d} \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

$$C \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{B \tan(c + dx)}{d}$$

input `Int[Sec[c + d*x]*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(B*Tan[c + d*x])/d + C*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d))`

3.39.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4535 `Int[(csc[(e_) + (f_)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_) + (f_)*(x_)]*(B_.) + csc[(e_) + (f_)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.39.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{B \tan(dx+c) + C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d}$
default	$\frac{B \tan(dx+c) + C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d}$
parts	$\frac{B \tan(dx+c)}{d} + \frac{C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d}$
parallelrisch	$\frac{-C(1+\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + C(1+\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 2B \sin(2dx+2c) + 2C \sin(dx+c)}{2d(1+\cos(2dx+2c))}$
norman	$\frac{(2B+C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - (2B-C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2d} + \frac{C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2d}$
risch	$-\frac{i(C e^{3i(dx+c)} - 2B e^{2i(dx+c)} - C e^{i(dx+c)} - 2B)}{d(e^{2i(dx+c)} + 1)^2} + \frac{\ln(e^{i(dx+c)} + i)C}{2d} - \frac{\ln(e^{i(dx+c)} - i)C}{2d}$

input `int(sec(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`output `1/d*(B*tan(d*x+c)+C*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c))))`**3.39.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.57

$$\int \sec(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \frac{C \cos(dx+c)^2 \log(\sin(dx+c)+1) - C \cos(dx+c)^2 \log(-\sin(dx+c)+1) + 2(2B \cos(dx+c) + C)}{4d \cos(dx+c)^2}$$

input `integrate(sec(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`output `1/4*(C*cos(d*x + c)^2*log(sin(d*x + c) + 1) - C*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(2*B*cos(d*x + c) + C)*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.39.6 Sympy [F]

$$\int \sec(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (B + C \sec(c + dx)) \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((B + C*sec(c + d*x))*sec(c + d*x)**2, x)`

3.39.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.23

$$\int \sec(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= -\frac{C \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx+c)+1) + \log(\sin(dx+c)-1) \right) - 4B \tan(dx+c)}{4d}$$

input `integrate(sec(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `-1/4*(C*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 4*B*tan(d*x + c))/d`

3.39.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. $2(43) = 86$.

Time = 0.33 (sec) , antiderivative size = 107, normalized size of antiderivative = 2.28

$$\int \sec(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{C \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - C \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) - \frac{2 \left(2B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 - C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 - 2B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right)}{\left(\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 - 1 \right)^2}}{2d}$$

input `integrate(sec(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output $\frac{1}{2}*(C*\log(\tan(\frac{1}{2}*d*x + \frac{1}{2}*c) + 1)) - C*\log(\tan(\frac{1}{2}*d*x + \frac{1}{2}*c) - 1)) - 2*(2*B*\tan(\frac{1}{2}*d*x + \frac{1}{2}*c)^3 - C*\tan(\frac{1}{2}*d*x + \frac{1}{2}*c)^3 - 2*B*\tan(\frac{1}{2}*d*x + \frac{1}{2}*c) - C*\tan(\frac{1}{2}*d*x + \frac{1}{2}*c))/(\tan(\frac{1}{2}*d*x + \frac{1}{2}*c)^2 - 1)^2)/d$

3.39.9 Mupad [B] (verification not implemented)

Time = 15.82 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.81

$$\int \sec(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{C \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (2B - C) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (2B + C)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)}$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/cos(c + d*x),x)`

output $(C*\operatorname{atanh}(\tan(c/2 + (d*x)/2)))/d - (\tan(c/2 + (d*x)/2)^3*(2*B - C) - \tan(c/2 + (d*x)/2)*(2*B + C))/(d*(\tan(c/2 + (d*x)/2)^4 - 2*\tan(c/2 + (d*x)/2)^2 + 1))$

3.40 $\int (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.40.1	Optimal result	255
3.40.2	Mathematica [A] (verified)	255
3.40.3	Rubi [A] (verified)	256
3.40.4	Maple [A] (verified)	256
3.40.5	Fricas [B] (verification not implemented)	257
3.40.6	Sympy [F]	257
3.40.7	Maxima [A] (verification not implemented)	258
3.40.8	Giac [B] (verification not implemented)	258
3.40.9	Mupad [B] (verification not implemented)	258

3.40.1 Optimal result

Integrand size = 19, antiderivative size = 24

$$\int (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{C \tan(c + dx)}{d}$$

output `B*arctanh(sin(d*x+c))/d+C*tan(d*x+c)/d`

3.40.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{C \tan(c + dx)}{d}$$

input `Integrate[B*Sec[c + d*x] + C*Sec[c + d*x]^2,x]`

output `(B*ArcTanh[Sin[c + d*x]])/d + (C*Tan[c + d*x])/d`

3.40.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

↓ 2009

$$\frac{B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{C \tan(c + dx)}{d}$$

input `Int[B*Sec[c + d*x] + C*Sec[c + d*x]^2,x]`

output `(B*ArcTanh[Sin[c + d*x]])/d + (C*Tan[c + d*x])/d`

3.40.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.40.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.25

method	result	size
derivativedivides	$\frac{B \ln(\sec(dx+c)+\tan(dx+c))+C \tan(dx+c)}{d}$	30
default	$\frac{B \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{C \tan(dx+c)}{d}$	32
parts	$\frac{B \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{C \tan(dx+c)}{d}$	32
risch	$-\frac{B \ln(e^{i(dx+c)}-i)}{d} + \frac{B \ln(e^{i(dx+c)}+i)}{d} + \frac{2iC}{d(e^{2i(dx+c)}+1)}$	59
parallelrisc	$\frac{-B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right) \cos(dx+c)+B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right) \cos(dx+c)+C \sin(dx+c)}{d \cos(dx+c)}$	63
norman	$-\frac{2C \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-1\right)} + \frac{B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{d} - \frac{B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{d}$	67

input `int(B*sec(d*x+c)+C*sec(d*x+c)^2,x,method=_RETURNVERBOSE)`

output `1/d*(B*ln(sec(d*x+c)+tan(d*x+c))+C*tan(d*x+c))`

3.40.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(24) = 48$.

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.50

$$\int (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{B \cos(dx + c) \log(\sin(dx + c) + 1) - B \cos(dx + c) \log(-\sin(dx + c) + 1) + 2C \sin(dx + c)}{2d \cos(dx + c)}$$

input `integrate(B*sec(d*x+c)+C*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/2*(B*cos(d*x + c)*log(sin(d*x + c) + 1) - B*cos(d*x + c)*log(-sin(d*x + c) + 1) + 2*C*sin(d*x + c))/(d*cos(d*x + c))`

3.40.6 Sympy [F]

$$\int (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (B + C \sec(c + dx)) \sec(c + dx) dx$$

input `integrate(B*sec(d*x+c)+C*sec(d*x+c)**2,x)`

output `Integral((B + C*sec(c + d*x))*sec(c + d*x), x)`

3.40.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.29

$$\int (B \sec(c+dx) + C \sec^2(c+dx)) dx = \frac{B \log(\sec(dx+c) + \tan(dx+c))}{d} + \frac{C \tan(dx+c)}{d}$$

input `integrate(B*sec(d*x+c)+C*sec(d*x+c)^2,x, algorithm="maxima")`

output `B*log(sec(d*x + c) + tan(d*x + c))/d + C*tan(d*x + c)/d`

3.40.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs. 2(24) = 48.

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.38

$$\begin{aligned} & \int (B \sec(c+dx) + C \sec^2(c+dx)) dx \\ &= \frac{B \left(\log \left(\left| \frac{1}{\sin(dx+c)} + \sin(dx+c) + 2 \right| \right) - \log \left(\left| \frac{1}{\sin(dx+c)} + \sin(dx+c) - 2 \right| \right) \right)}{4d} \\ & \quad + \frac{C \tan(dx+c)}{d} \end{aligned}$$

input `integrate(B*sec(d*x+c)+C*sec(d*x+c)^2,x, algorithm="giac")`

output `1/4*B*(log(abs(1/sin(d*x + c) + sin(d*x + c) + 2)) - log(abs(1/sin(d*x + c) + sin(d*x + c) - 2)))/d + C*tan(d*x + c)/d`

3.40.9 Mupad [B] (verification not implemented)

Time = 15.31 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.96

$$\int (B \sec(c+dx) + C \sec^2(c+dx)) dx = \frac{2B \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d} - \frac{2C \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1\right)}$$

input `int(B/cos(c + d*x) + C/cos(c + d*x)^2,x)`

output `(2*B*atanh(tan(c/2 + (d*x)/2)))/d - (2*C*tan(c/2 + (d*x)/2))/(d*(tan(c/2 + (d*x)/2)^2 - 1))`

3.41 $\int \cos(c+dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.41.1	Optimal result	260
3.41.2	Mathematica [A] (verified)	260
3.41.3	Rubi [A] (verified)	261
3.41.4	Maple [A] (verified)	262
3.41.5	Fricas [B] (verification not implemented)	263
3.41.6	Sympy [F]	263
3.41.7	Maxima [B] (verification not implemented)	263
3.41.8	Giac [B] (verification not implemented)	264
3.41.9	Mupad [B] (verification not implemented)	264

3.41.1 Optimal result

Integrand size = 26, antiderivative size = 16

$$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = Bx + \frac{C \arctanh(\sin(c + dx))}{d}$$

output `B*x+C*arctanh(sin(d*x+c))/d`

3.41.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = Bx + \frac{C \arctanh(\sin(c + dx))}{d}$$

input `Integrate[Cos[c + d*x]*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `B*x + (C*ArcTanh[Sin[c + d*x]])/d`

3.41.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 4535, 24, 27, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4535} \\
 & B \int 1 dx + \int C \sec(c + dx) dx \\
 & \quad \downarrow \text{24} \\
 & \int C \sec(c + dx) dx + Bx \\
 & \quad \downarrow \text{27} \\
 & C \int \sec(c + dx) dx + Bx \\
 & \quad \downarrow \text{3042} \\
 & C \int \csc(c + dx + \frac{\pi}{2}) dx + Bx \\
 & \quad \downarrow \text{4257} \\
 & \frac{C \operatorname{arctanh}(\sin(c + dx))}{d} + Bx
 \end{aligned}$$

input `Int[Cos[c + d*x]*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `B*x + (C*ArcTanh[Sin[c + d*x]])/d`

3.41.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4535 `Int[(csc[(e_) + (f_)*(x_)]*(b_))^(m_)*((A_) + csc[(e_) + (f_)*(x_)]*(B_) + csc[(e_) + (f_)*(x_)]^2*(C_)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.41.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

method	result	size
derivativedivides	$\frac{C \ln(\sec(dx+c)+\tan(dx+c))+B(dx+c)}{d}$	29
default	$\frac{C \ln(\sec(dx+c)+\tan(dx+c))+B(dx+c)}{d}$	29
parallelrisch	$\frac{Bxd - C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{d}$	39
risch	$Bx + \frac{\ln(e^{i(dx+c)}+i)C}{d} - \frac{\ln(e^{i(dx+c)}-i)C}{d}$	42
norman	$\frac{Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 - Bx}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)} + \frac{C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{d} - \frac{C \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{d}$	87

input `int(cos(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(C*ln(sec(d*x+c)+tan(d*x+c))+B*(d*x+c))`

3.41. $\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.41.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(16) = 32$.

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.25

$$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2 B dx + C \log(\sin(dx + c) + 1) - C \log(-\sin(dx + c) + 1)}{2d}$$

input `integrate(cos(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/2*(2*B*d*x + C*log(sin(d*x + c) + 1) - C*log(-sin(d*x + c) + 1))/d`

3.41.6 Sympy [F]

$$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (B + C \sec(c + dx)) \cos(c + dx) \sec(c + dx) dx$$

input `integrate(cos(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((B + C*sec(c + d*x))*cos(c + d*x)*sec(c + d*x), x)`

3.41.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(16) = 32$.

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.31

$$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2(dx + c)B + C(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1))}{2d}$$

input `integrate(cos(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*B + C*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)))/d`

3.41. $\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.41.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. $2(16) = 32$.

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.69

$$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(dx + c)B + C \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - C \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right)}{d}$$

input `integrate(cos(d*x+c)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `((d*x + c)*B + C*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - C*log(abs(tan(1/2*d*x + 1/2*c) - 1)))/d`

3.41.9 Mupad [B] (verification not implemented)

Time = 15.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 3.56

$$\int \cos(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2 B \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d} + \frac{2 C \operatorname{atanh} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d}$$

input `int(cos(c + d*x)*(B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(2*B*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*C*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d`

3.42 $\int \cos^2(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$

3.42.1	Optimal result	265
3.42.2	Mathematica [A] (verified)	265
3.42.3	Rubi [A] (verified)	266
3.42.4	Maple [A] (verified)	267
3.42.5	Fricas [A] (verification not implemented)	268
3.42.6	Sympy [F]	268
3.42.7	Maxima [A] (verification not implemented)	268
3.42.8	Giac [B] (verification not implemented)	269
3.42.9	Mupad [B] (verification not implemented)	269

3.42.1 Optimal result

Integrand size = 28, antiderivative size = 15

$$\int \cos^2(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx = Cx + \frac{B \sin(c+dx)}{d}$$

output `C*x+B*sin(d*x+c)/d`

3.42.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\begin{aligned} & \int \cos^2(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\ &= Cx + \frac{B \cos(dx) \sin(c)}{d} + \frac{B \cos(c) \sin(dx)}{d} \end{aligned}$$

input `Integrate[Cos[c + d*x]^2*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `C*x + (B*Cos[d*x]*Sin[c])/d + (B*Cos[c]*Sin[d*x])/d`

3.42.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {3042, 4535, 24, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \cos(c + dx) dx + \int C dx \\
 & \quad \downarrow \text{24} \\
 & B \int \cos(c + dx) dx + Cx \\
 & \quad \downarrow \text{3042} \\
 & B \int \sin(c + dx + \frac{\pi}{2}) dx + Cx \\
 & \quad \downarrow \text{3117} \\
 & \frac{B \sin(c + dx)}{d} + Cx
 \end{aligned}$$

input `Int[Cos[c + d*x]^2*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `C*x + (B*Sin[c + d*x])/d`

3.42.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)])*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.), x_Symbol] := Simp[B/b Int[(b*Cs c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.42.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
risch	$Cx + \frac{B \sin(dx+c)}{d}$	16
parallelrisc	$\frac{dx C + B \sin(dx+c)}{d}$	18
derivativedivides	$\frac{B \sin(dx+c) + C(dx+c)}{d}$	21
default	$\frac{B \sin(dx+c) + C(dx+c)}{d}$	21
norman	$\frac{Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 - Cx - \frac{2B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{2B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{d} - Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$	112

input `int(cos(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `C*x+B*sin(d*x+c)/d`

3.42.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \cos^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{Cdx + B \sin(dx + c)}{d}$$

input `integrate(cos(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `(C*d*x + B*sin(d*x + c))/d`

3.42.6 Sympy [F]

$$\begin{aligned} & \int \cos^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int (B + C \sec(c + dx)) \cos^2(c + dx) \sec(c + dx) dx \end{aligned}$$

input `integrate(cos(d*x+c)**2*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((B + C*sec(c + d*x))*cos(c + d*x)**2*sec(c + d*x), x)`

3.42.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.33

$$\int \cos^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{(dx + c)C + B \sin(dx + c)}{d}$$

input `integrate(cos(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `((d*x + c)*C + B*sin(d*x + c))/d`

3.42.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(15) = 30$.

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.60

$$\int \cos^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{(dx + c)C + \frac{2B \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1}}{d}$$

input `integrate(cos(d*x+c)^2*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `((d*x + c)*C + 2*B*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 + 1))/d`

3.42.9 Mupad [B] (verification not implemented)

Time = 15.42 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \cos^2(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{B \sin(c + dx) + C dx}{d}$$

input `int(cos(c + d*x)^2*(B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(B*sin(c + d*x) + C*d*x)/d`

3.43 $\int \cos^3(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$

3.43.1	Optimal result	270
3.43.2	Mathematica [A] (verified)	270
3.43.3	Rubi [A] (verified)	271
3.43.4	Maple [A] (verified)	272
3.43.5	Fricas [A] (verification not implemented)	273
3.43.6	Sympy [F]	273
3.43.7	Maxima [A] (verification not implemented)	274
3.43.8	Giac [B] (verification not implemented)	274
3.43.9	Mupad [B] (verification not implemented)	275

3.43.1 Optimal result

Integrand size = 28, antiderivative size = 38

$$\begin{aligned} & \int \cos^3(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\ &= \frac{Bx}{2} + \frac{C \sin(c+dx)}{d} + \frac{B \cos(c+dx) \sin(c+dx)}{2d} \end{aligned}$$

output `1/2*B*x+C*sin(d*x+c)/d+1/2*B*cos(d*x+c)*sin(d*x+c)/d`

3.43.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

$$\begin{aligned} & \int \cos^3(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\ &= \frac{4C \sin(c+dx) + B(2(c+dx) + \sin(2(c+dx)))}{4d} \end{aligned}$$

input `Integrate[Cos[c + d*x]^3*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(4*C*Sin[c + d*x] + B*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d)`

3.43.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4535, 27, 3042, 3115, 24, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \cos^2(c + dx) dx + \int C \cos(c + dx) dx \\
 & \quad \downarrow \text{27} \\
 & B \int \cos^2(c + dx) dx + C \int \cos(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \sin(c + dx + \frac{\pi}{2})^2 dx + C \int \sin(c + dx + \frac{\pi}{2}) dx \\
 & \quad \downarrow \text{3115} \\
 & B \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) + C \int \sin(c + dx + \frac{\pi}{2}) dx \\
 & \quad \downarrow \text{24} \\
 & C \int \sin(c + dx + \frac{\pi}{2}) dx + B \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) \\
 & \quad \downarrow \text{3117} \\
 & B \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) + \frac{C \sin(c + dx)}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^3*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(C*Sin[c + d*x])/d + B*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d))`

3.43.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.)^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)])*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.43.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

method	result
risch	$\frac{Bx}{2} + \frac{C \sin(dx+c)}{d} + \frac{B \sin(2dx+2c)}{4d}$
parallelrisch	$\frac{2Bxd+B \sin(2dx+2c)+4C \sin(dx+c)}{4d}$
derivativedivides	$\frac{B \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + C \sin(dx+c)}{d}$
default	$\frac{B \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + C \sin(dx+c)}{d}$
norman	$\frac{Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + \frac{(B-2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{d} + \frac{(B+2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{d} - \frac{Bx}{2} - Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^8}{2} - \frac{(B-2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$

input `int(cos(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/2*B*x+C*sin(d*x+c)/d+1/4*B/d*sin(2*d*x+2*c)`

3.43.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

$$\int \cos^3(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \frac{Bdx + (B \cos(dx+c) + 2C) \sin(dx+c)}{2d}$$

input `integrate(cos(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/2*(B*d*x + (B*cos(d*x + c) + 2*C)*sin(d*x + c))/d`

3.43.6 Sympy [F]

$$\int \cos^3(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \int (B + C \sec(c+dx)) \cos^3(c+dx) \sec(c+dx) dx$$

input `integrate(cos(d*x+c)**3*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((B + C*sec(c + d*x))*cos(c + d*x)**3*sec(c + d*x), x)`

3.43.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \cos^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(2 dx + 2 c + \sin(2 dx + 2 c))B + 4 C \sin(dx + c)}{4 d}$$

input `integrate(cos(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/4*((2*d*x + 2*c + sin(2*d*x + 2*c))*B + 4*C*sin(d*x + c))/d`

3.43.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(34) = 68$.

Time = 0.29 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.16

$$\int \cos^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(dx + c)B - \frac{2(B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 2C \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - B \tan(\frac{1}{2} dx + \frac{1}{2} c) - 2C \tan(\frac{1}{2} dx + \frac{1}{2} c))}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^2}}{2 d}$$

input `integrate(cos(d*x+c)^3*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/2*((d*x + c)*B - 2*(B*tan(1/2*d*x + 1/2*c)^3 - 2*C*tan(1/2*d*x + 1/2*c)^3 - B*tan(1/2*d*x + 1/2*c) - 2*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^2)/d`

3.43.9 Mupad [B] (verification not implemented)

Time = 15.32 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \cos^3(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{Bx}{2} + \frac{B \sin(2c + 2dx)}{4d} + \frac{C \sin(c + dx)}{d}$$

input `int(cos(c + d*x)^3*(B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(B*x)/2 + (B*sin(2*c + 2*d*x))/(4*d) + (C*sin(c + d*x))/d`

3.44 $\int \cos^4(c+dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.44.1	Optimal result	276
3.44.2	Mathematica [A] (verified)	276
3.44.3	Rubi [A] (verified)	277
3.44.4	Maple [A] (verified)	279
3.44.5	Fricas [A] (verification not implemented)	279
3.44.6	Sympy [F]	280
3.44.7	Maxima [A] (verification not implemented)	280
3.44.8	Giac [B] (verification not implemented)	280
3.44.9	Mupad [B] (verification not implemented)	281

3.44.1 Optimal result

Integrand size = 28, antiderivative size = 54

$$\int \cos^4(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{Cx}{2} + \frac{B \sin(c + dx)}{d} + \frac{C \cos(c + dx) \sin(c + dx)}{2d} - \frac{B \sin^3(c + dx)}{3d}$$

```
output 1/2*C*x+B*sin(d*x+c)/d+1/2*C*cos(d*x+c)*sin(d*x+c)/d-1/3*B*sin(d*x+c)^3/d
```

3.44.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06

$$\int \cos^4(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{C(c + dx)}{2d} + \frac{B \sin(c + dx)}{d} - \frac{B \sin^3(c + dx)}{3d} + \frac{C \sin(2(c + dx))}{4d}$$

```
input Integrate[Cos[c + d*x]^4*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]
```

```
output (C*(c + d*x))/(2*d) + (B*Sin[c + d*x])/d - (B*Sin[c + d*x]^3)/(3*d) + (C*S
in[2*(c + d*x)]/(4*d)
```

3.44.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.02, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4535, 27, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c+dx+\frac{\pi}{2}) + C \csc(c+dx+\frac{\pi}{2})^2}{\csc(c+dx+\frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \cos^3(c+dx) dx + \int C \cos^2(c+dx) dx \\
 & \quad \downarrow \text{27} \\
 & B \int \cos^3(c+dx) dx + C \int \cos^2(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \sin(c+dx+\frac{\pi}{2})^3 dx + C \int \sin(c+dx+\frac{\pi}{2})^2 dx \\
 & \quad \downarrow \text{3113} \\
 & C \int \sin(c+dx+\frac{\pi}{2})^2 dx - \frac{B \int (1 - \sin^2(c+dx)) d(-\sin(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & C \int \sin(c+dx+\frac{\pi}{2})^2 dx - \frac{B(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} \\
 & \quad \downarrow \text{3115} \\
 & C \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) - \frac{B(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} \\
 & \quad \downarrow \text{24} \\
 & C \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) - \frac{B(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^4*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `C*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (B*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d`

3.44.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.44.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

method	result
parallelrisch	$\frac{6dxC+9B\sin(dx+c)+B\sin(3dx+3c)+3\sin(2dx+2c)C}{12d}$
risch	$\frac{Cx}{2} + \frac{3B\sin(dx+c)}{4d} + \frac{B\sin(3dx+3c)}{12d} + \frac{\sin(2dx+2c)C}{4d}$
derivativedivides	$\frac{B(2+\cos(dx+c)^2)\sin(dx+c)}{3} + C\left(\frac{\sin(dx+c)\cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)$
default	$\frac{B(2+\cos(dx+c)^2)\sin(dx+c)}{3} + C\left(\frac{\sin(dx+c)\cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)$
norman	$\frac{Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + \frac{(2B-C)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^9}{d} - \frac{Cx}{2} - \frac{4B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{3d} + \frac{4B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^7}{3d} + \frac{2C \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{d} - \frac{3Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{2}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^4 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$

input `int(cos(d*x+c)^4*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/12*(6*d*x*C+9*B*sin(d*x+c)+B*sin(3*d*x+3*c)+3*sin(2*d*x+2*c)*C)/d`

3.44.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.78

$$\int \cos^4(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \frac{3Cdx + (2B \cos(dx+c))^2 + 3C \cos(dx+c) + 4B \sin(dx+c)}{6d}$$

input `integrate(cos(d*x+c)^4*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/6*(3*C*d*x + (2*B*cos(d*x + c))^2 + 3*C*cos(d*x + c) + 4*B)*sin(d*x + c)/d`

3.44.6 Sympy [F]

$$\begin{aligned} & \int \cos^4(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int (B + C \sec(c + dx)) \cos^4(c + dx) \sec(c + dx) dx \end{aligned}$$

input `integrate(cos(d*x+c)**4*(B*sec(d*x+c)+C*sec(d*x+c)**2), x)`

output `Integral((B + C*sec(c + d*x))*cos(c + d*x)**4*sec(c + d*x), x)`

3.44.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.85

$$\begin{aligned} & \int \cos^4(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= -\frac{4(\sin(dx + c)^3 - 3\sin(dx + c))B - 3(2dx + 2c + \sin(2dx + 2c))C}{12d} \end{aligned}$$

input `integrate(cos(d*x+c)^4*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `-1/12*(4*(sin(d*x + c)^3 - 3*sin(d*x + c))*B - 3*(2*d*x + 2*c + sin(2*d*x + 2*c))*C)/d`

3.44.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 98 vs. $2(48) = 96$.

Time = 0.29 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.81

$$\begin{aligned} & \int \cos^4(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{3(dx + c)C + \frac{2(6B \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 3C \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 + 4B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 6B \tan(\frac{1}{2} dx + \frac{1}{2} c) + 3C \tan(\frac{1}{2} dx + \frac{1}{2} c))}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^3}}{6d} \end{aligned}$$

input `integrate(cos(d*x+c)^4*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output $\frac{1}{6}*(3*(d*x + c)*C + 2*(6*B*\tan(1/2*d*x + 1/2*c)^5 - 3*C*\tan(1/2*d*x + 1/2*c)^5 + 4*B*\tan(1/2*d*x + 1/2*c)^3 + 6*B*\tan(1/2*d*x + 1/2*c) + 3*C*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 + 1)^3/d$

3.44.9 Mupad [B] (verification not implemented)

Time = 15.62 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.02

$$\int \cos^4(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{C x}{2} + \frac{2 B \sin(c + dx)}{3 d} + \frac{C \cos(c + dx) \sin(c + dx)}{2 d} + \frac{B \cos(c + dx)^2 \sin(c + dx)}{3 d}$$

input `int(cos(c + d*x)^4*(B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output $(C*x)/2 + (2*B*\sin(c + d*x))/(3*d) + (C*\cos(c + d*x)*\sin(c + d*x))/(2*d) + (B*\cos(c + d*x)^2*\sin(c + d*x))/(3*d)$

3.45 $\int \cos^5(c+dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.45.1	Optimal result	282
3.45.2	Mathematica [A] (verified)	282
3.45.3	Rubi [A] (verified)	283
3.45.4	Maple [A] (verified)	285
3.45.5	Fricas [A] (verification not implemented)	286
3.45.6	Sympy [F]	286
3.45.7	Maxima [A] (verification not implemented)	286
3.45.8	Giac [B] (verification not implemented)	287
3.45.9	Mupad [B] (verification not implemented)	287

3.45.1 Optimal result

Integrand size = 28, antiderivative size = 76

$$\begin{aligned} & \int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{3Bx}{8} + \frac{C \sin(c + dx)}{d} + \frac{3B \cos(c + dx) \sin(c + dx)}{8d} \\ & \quad + \frac{B \cos^3(c + dx) \sin(c + dx)}{4d} - \frac{C \sin^3(c + dx)}{3d} \end{aligned}$$

output `3/8*B*x+C*sin(d*x+c)/d+3/8*B*cos(d*x+c)*sin(d*x+c)/d+1/4*B*cos(d*x+c)^3*sin(d*x+c)/d-1/3*C*sin(d*x+c)^3/d`

3.45.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{3B(c + dx)}{8d} + \frac{C \sin(c + dx)}{d} - \frac{C \sin^3(c + dx)}{3d} + \frac{B \sin(2(c + dx))}{4d} + \frac{B \sin(4(c + dx))}{32d} \end{aligned}$$

input `Integrate[Cos[c + d*x]^5*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(3*B*(c + d*x))/(8*d) + (C*Sin[c + d*x])/d - (C*Sin[c + d*x]^3)/(3*d) + (B*Sin[2*(c + d*x)])/(4*d) + (B*Sin[4*(c + d*x)])/(32*d)`

3.45.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3042, 4535, 27, 3042, 3113, 2009, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^5(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c+dx+\frac{\pi}{2}) + C \csc(c+dx+\frac{\pi}{2})^2}{\csc(c+dx+\frac{\pi}{2})^5} dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \cos^4(c+dx) dx + \int C \cos^3(c+dx) dx \\
 & \quad \downarrow \text{27} \\
 & B \int \cos^4(c+dx) dx + C \int \cos^3(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \sin(c+dx+\frac{\pi}{2})^4 dx + C \int \sin(c+dx+\frac{\pi}{2})^3 dx \\
 & \quad \downarrow \text{3113} \\
 & B \int \sin(c+dx+\frac{\pi}{2})^4 dx - \frac{C \int (1 - \sin^2(c+dx)) d(-\sin(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & B \int \sin(c+dx+\frac{\pi}{2})^4 dx - \frac{C(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} \\
 & \quad \downarrow \text{3115} \\
 & B \left(\frac{3}{4} \int \cos^2(c+dx) dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) - \frac{C(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{3}{4} \int \sin(c+dx+\frac{\pi}{2})^2 dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) - \frac{C(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3115} \\
 B\left(\frac{3}{4}\left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx)\cos(c+dx)}{2d}\right) + \frac{\sin(c+dx)\cos^3(c+dx)}{4d}\right) - \\
 \frac{C\left(\frac{1}{3}\sin^3(c+dx) - \sin(c+dx)\right)}{d} \\
 \downarrow \text{24} \\
 B\left(\frac{\sin(c+dx)\cos^3(c+dx)}{4d} + \frac{3}{4}\left(\frac{\sin(c+dx)\cos(c+dx)}{2d} + \frac{x}{2}\right)\right) - \\
 \frac{C\left(\frac{1}{3}\sin^3(c+dx) - \sin(c+dx)\right)}{d}
 \end{array}$$

input `Int[Cos[c + d*x]^5*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `-((C*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d) + B*((Cos[c + d*x]^3*Sin[c + d*x])/((4*d) + (3*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/((2*d))))/4)`

3.45.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.45.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.75

method	result
parallelrisch	$\frac{36Bxd+3B\sin(4dx+4c)+24B\sin(2dx+2c)+72C\sin(dx+c)+8C\sin(3dx+3c)}{96d}$
derivativedivides	$B \left(\frac{\left(\cos(dx+c)^3 + \frac{3\cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{C(2+\cos(dx+c)^2)\sin(dx+c)}{3}$
default	$B \left(\frac{\left(\cos(dx+c)^3 + \frac{3\cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{C(2+\cos(dx+c)^2)\sin(dx+c)}{3}$
risch	$\frac{3Bx}{8} + \frac{3C\sin(dx+c)}{4d} + \frac{B\sin(4dx+4c)}{32d} + \frac{\sin(3dx+3c)C}{12d} + \frac{B\sin(2dx+2c)}{4d}$
norman	$\frac{-\frac{3Bx}{8} - \frac{3Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{2} - \frac{15Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{8} + \frac{15Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^8}{8} + \frac{3Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^{10}}{2} + \frac{3Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^{12}}{8} + \frac{(3B-8C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^{14}}{6}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^{14}}$

input `int(cos(d*x+c)^5*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/96*(36*B*x*d+3*B*sin(4*d*x+4*c)+24*B*sin(2*d*x+2*c)+72*C*sin(d*x+c)+8*C*sin(3*d*x+3*c))/d`

3.45.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.70

$$\int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{9 B dx + (6 B \cos(dx + c)^3 + 8 C \cos(dx + c)^2 + 9 B \cos(dx + c) + 16 C) \sin(dx + c)}{24 d}$$

input `integrate(cos(d*x+c)^5*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/24*(9*B*d*x + (6*B*cos(d*x + c)^3 + 8*C*cos(d*x + c)^2 + 9*B*cos(d*x + c) + 16*C)*sin(d*x + c))/d`

3.45.6 Sympy [F]

$$\int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (B + C \sec(c + dx)) \cos^5(c + dx) \sec(c + dx) dx$$

input `integrate(cos(d*x+c)**5*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((B + C*sec(c + d*x))*cos(c + d*x)**5*sec(c + d*x), x)`

3.45.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.75

$$\int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(2 dx + 2 c))B - 32(\sin(dx + c)^3 - 3 \sin(dx + c))C}{96 d}$$

input `integrate(cos(d*x+c)^5*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/96*(3*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B - 32*(sin(d*x + c)^3 - 3*sin(d*x + c))*C)/d`

3.45.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 140 vs. $2(68) = 136$.

Time = 0.29 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.84

$$\int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{9(dx + c)B - \frac{2(15B \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 - 24C \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 - 9B \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 40C \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 + 9B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 40C \tan(\frac{1}{2} dx + \frac{1}{2} c)^3)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^4}}{24d}$$

input `integrate(cos(d*x+c)^5*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/24*(9*(d*x + c)*B - 2*(15*B*tan(1/2*d*x + 1/2*c)^7 - 24*C*tan(1/2*d*x + 1/2*c)^7 - 9*B*tan(1/2*d*x + 1/2*c)^5 - 40*C*tan(1/2*d*x + 1/2*c)^5 + 9*B*tan(1/2*d*x + 1/2*c)^3 - 40*C*tan(1/2*d*x + 1/2*c)^3 - 15*B*tan(1/2*d*x + 1/2*c) - 24*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^4/d`

3.45.9 Mupad [B] (verification not implemented)

Time = 15.25 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.99

$$\int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3Bx}{8} + \frac{2C \sin(c + dx)}{3d} + \frac{3B \cos(c + dx) \sin(c + dx)}{8d}$$

$$+ \frac{B \cos(c + dx)^3 \sin(c + dx)}{4d} + \frac{C \cos(c + dx)^2 \sin(c + dx)}{3d}$$

input `int(cos(c + d*x)^5*(B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

3.45. $\int \cos^5(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

output $(3Bx)/8 + (2C\sin(c + dx))/(3d) + (3B\cos(c + dx)\sin(c + dx))/(8d) + (B\cos(c + dx)^3\sin(c + dx))/(4d) + (C\cos(c + dx)^2\sin(c + dx))/(3d)$

3.46 $\int \cos^6(c+dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.46.1	Optimal result	289
3.46.2	Mathematica [A] (verified)	289
3.46.3	Rubi [A] (verified)	290
3.46.4	Maple [A] (verified)	292
3.46.5	Fricas [A] (verification not implemented)	293
3.46.6	Sympy [F(-1)]	293
3.46.7	Maxima [A] (verification not implemented)	294
3.46.8	Giac [A] (verification not implemented)	294
3.46.9	Mupad [B] (verification not implemented)	295

3.46.1 Optimal result

Integrand size = 28, antiderivative size = 92

$$\int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3Cx}{8} + \frac{B \sin(c + dx)}{d} + \frac{3C \cos(c + dx) \sin(c + dx)}{8d}$$

$$+ \frac{C \cos^3(c + dx) \sin(c + dx)}{4d} - \frac{2B \sin^3(c + dx)}{3d} + \frac{B \sin^5(c + dx)}{5d}$$

output `3/8*C*x+B*sin(d*x+c)/d+3/8*C*cos(d*x+c)*sin(d*x+c)/d+1/4*C*cos(d*x+c)^3*sin(d*x+c)/d-2/3*B*sin(d*x+c)^3/d+1/5*B*sin(d*x+c)^5/d`

3.46.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.97

$$\int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3C(c + dx)}{8d} + \frac{B \sin(c + dx)}{d} - \frac{2B \sin^3(c + dx)}{3d}$$

$$+ \frac{B \sin^5(c + dx)}{5d} + \frac{C \sin(2(c + dx))}{4d} + \frac{C \sin(4(c + dx))}{32d}$$

input `Integrate[Cos[c + d*x]^6*(B*Sec[c + d*x] + C*Sec[c + d*x]^2), x]`

output $(3*C*(c + d*x))/(8*d) + (B*\text{Sin}[c + d*x])/d - (2*B*\text{Sin}[c + d*x]^3)/(3*d) + (B*\text{Sin}[c + d*x]^5)/(5*d) + (C*\text{Sin}[2*(c + d*x)])/(4*d) + (C*\text{Sin}[4*(c + d*x)])/(32*d)$

3.46.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3042, 4535, 27, 3042, 3113, 2009, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})^6} dx \\
 & \quad \downarrow \text{4535} \\
 & B \int \cos^5(c + dx) dx + \int C \cos^4(c + dx) dx \\
 & \quad \downarrow \text{27} \\
 & B \int \cos^5(c + dx) dx + C \int \cos^4(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \sin(c + dx + \frac{\pi}{2})^5 dx + C \int \sin(c + dx + \frac{\pi}{2})^4 dx \\
 & \quad \downarrow \text{3113} \\
 & C \int \sin(c + dx + \frac{\pi}{2})^4 dx - \frac{B \int (\sin^4(c + dx) - 2 \sin^2(c + dx) + 1) d(-\sin(c + dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & C \int \sin(c + dx + \frac{\pi}{2})^4 dx - \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d} \\
 & \quad \downarrow \text{3115}
 \end{aligned}$$

$$\begin{aligned}
& \frac{C\left(\frac{3}{4} \int \cos^2(c+dx) dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d}\right) - B\left(-\frac{1}{5} \sin^5(c+dx) + \frac{2}{3} \sin^3(c+dx) - \sin(c+dx)\right)}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{C\left(\frac{3}{4} \int \sin\left(c+dx + \frac{\pi}{2}\right)^2 dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d}\right) - B\left(-\frac{1}{5} \sin^5(c+dx) + \frac{2}{3} \sin^3(c+dx) - \sin(c+dx)\right)}{d} \\
& \quad \downarrow \text{3115} \\
& \frac{C\left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d}\right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d}\right) - B\left(-\frac{1}{5} \sin^5(c+dx) + \frac{2}{3} \sin^3(c+dx) - \sin(c+dx)\right)}{d} \\
& \quad \downarrow \text{24} \\
& \frac{C\left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2}\right)\right) - B\left(-\frac{1}{5} \sin^5(c+dx) + \frac{2}{3} \sin^3(c+dx) - \sin(c+dx)\right)}{d}
\end{aligned}$$

input `Int[Cos[c + d*x]^6*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `-((B*(-Sin[c + d*x] + (2*Sin[c + d*x]^3)/3 - Sin[c + d*x]^5/5))/d) + C*((Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (3*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/4)`

3.46.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*(n - 1)/n Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.46.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.75

method	result
parallelrisch	$\frac{180dxC + 300B \sin(dx+c) + 6B \sin(5dx+5c) + 50B \sin(3dx+3c) + 15 \sin(4dx+4c)C + 120 \sin(2dx+2c)C}{480d}$
derivativedivides	$\frac{B \left(\frac{8}{3} + \cos(dx+c)^4 + \frac{4 \cos(dx+c)^2}{3} \right) \sin(dx+c)}{5} + C \left(\frac{\left(\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right)$
default	$\frac{B \left(\frac{8}{3} + \cos(dx+c)^4 + \frac{4 \cos(dx+c)^2}{3} \right) \sin(dx+c)}{5} + C \left(\frac{\left(\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right)$
risch	$\frac{3Cx}{8} + \frac{5B \sin(dx+c)}{8d} + \frac{B \sin(5dx+5c)}{80d} + \frac{\sin(4dx+4c)C}{32d} + \frac{5B \sin(3dx+3c)}{48d} + \frac{\sin(2dx+2c)C}{4d}$
norman	$\frac{C \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^7}{d} - \frac{3Cx}{8} - \frac{15Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{8} - \frac{27Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{8} - \frac{15Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6}{8} + \frac{15Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^8}{8} + \frac{27Cx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8}$

input `int(cos(d*x+c)^6*(B*sec(d*x+c)+C*sec(d*x+c)^2), x, method=_RETURNVERBOSE)`

3.46. $\int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$

output $\frac{1}{480} \cdot (180 \cdot d \cdot x \cdot C + 300 \cdot B \cdot \sin(dx+c) + 6 \cdot B \cdot \sin(5 \cdot dx+5 \cdot c) + 50 \cdot B \cdot \sin(3 \cdot dx+3 \cdot c) + 15 \cdot \sin(4 \cdot dx+4 \cdot c) \cdot C + 120 \cdot \sin(2 \cdot dx+2 \cdot c) \cdot C) / d$

3.46.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.70

$$\int \cos^6(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \frac{45 C dx + (24 B \cos(dx+c)^4 + 30 C \cos(dx+c)^3 + 32 B \cos(dx+c)^2 + 45 C \cos(dx+c) + 64 B) \sin(dx+c)}{120 d}$$

input `integrate(cos(d*x+c)^6*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output $\frac{1}{120} \cdot (45 \cdot C \cdot d \cdot x + (24 \cdot B \cdot \cos(dx+c)^4 + 30 \cdot C \cdot \cos(dx+c)^3 + 32 \cdot B \cdot \cos(dx+c)^2 + 45 \cdot C \cdot \cos(dx+c) + 64 \cdot B) \cdot \sin(dx+c)) / d$

3.46.6 Sympy [F(-1)]

Timed out.

$$\int \cos^6(c+dx) (B \sec(c+dx) + C \sec^2(c+dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**6*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Timed out`

3.46.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.75

$$\int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{32 (3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c)) B + 15 (12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(2 dx + 2 c)) C}{480 d}$$

```
input integrate(cos(d*x+c)^6*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")
```

```
output 1/480*(32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B + 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*C)/d
```

3.46.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.67

$$\int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{45(dx + c)C + \frac{2(120B \tan(\frac{1}{2} dx + \frac{1}{2} c)^9 - 75C \tan(\frac{1}{2} dx + \frac{1}{2} c)^9 + 160B \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 - 30C \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 + 464B \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 + 120B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 30C \tan(\frac{1}{2} dx + \frac{1}{2} c))}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^5}}{120 d}$$

```
input integrate(cos(d*x+c)^6*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")
```

```
output 1/120*(45*(d*x + c)*C + 2*(120*B*tan(1/2*d*x + 1/2*c)^9 - 75*C*tan(1/2*d*x + 1/2*c)^9 + 160*B*tan(1/2*d*x + 1/2*c)^7 - 30*C*tan(1/2*d*x + 1/2*c)^7 + 464*B*tan(1/2*d*x + 1/2*c)^5 + 160*B*tan(1/2*d*x + 1/2*c)^3 + 30*C*tan(1/2*d*x + 1/2*c)^3 + 120*B*tan(1/2*d*x + 1/2*c) + 75*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^5)/d
```

3.46.9 Mupad [B] (verification not implemented)

Time = 19.25 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.23

$$\int \cos^6(c + dx) (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{3Cx}{8} + \frac{(2B - \frac{5C}{4}) \tan(\frac{c}{2} + \frac{dx}{2})^9 + (\frac{8B}{3} - \frac{C}{2}) \tan(\frac{c}{2} + \frac{dx}{2})^7 + \frac{116B \tan(\frac{c}{2} + \frac{dx}{2})^5}{15} + (\frac{8B}{3} + \frac{C}{2}) \tan(\frac{c}{2} + \frac{dx}{2})^3 + (\frac{2B}{3} - \frac{C}{2}) \tan(\frac{c}{2} + \frac{dx}{2})}{d \left(\tan(\frac{c}{2} + \frac{dx}{2})^2 + 1 \right)^5}$$

input `int(cos(c + d*x)^6*(B/cos(c + d*x) + C/cos(c + d*x)^2),x)`output `(3*C*x)/8 + (tan(c/2 + (d*x)/2)^3*((8*B)/3 + C/2) + tan(c/2 + (d*x)/2)^9*(2*B - (5*C)/4) + tan(c/2 + (d*x)/2)^7*((8*B)/3 - C/2) + (116*B*tan(c/2 + (d*x)/2)^5)/15 + tan(c/2 + (d*x)/2)*(2*B + (5*C)/4))/(d*(tan(c/2 + (d*x)/2)^2 + 1)^5)`

3.47 $\int (b \sec(c+dx))^{3/2} (B \sec(c+dx) + C \sec^2(c+dx)) dx$

3.47.1	Optimal result	296
3.47.2	Mathematica [A] (verified)	297
3.47.3	Rubi [A] (verified)	297
3.47.4	Maple [C] (verified)	300
3.47.5	Fricas [C] (verification not implemented)	301
3.47.6	Sympy [F]	302
3.47.7	Maxima [F]	302
3.47.8	Giac [F]	302
3.47.9	Mupad [F(-1)]	303

3.47.1 Optimal result

Integrand size = 32, antiderivative size = 169

$$\int (b \sec(c+dx))^{3/2} (B \sec(c+dx) + C \sec^2(c+dx)) dx =$$

$$-\frac{6b^2CE\left(\frac{1}{2}(c+dx) \mid 2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{2bB\sqrt{\cos(c+dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)\sqrt{b\sec(c+dx)}}{3d}$$

$$+ \frac{6bC\sqrt{b\sec(c+dx)}\sin(c+dx)}{5d} + \frac{2B(b\sec(c+dx))^{3/2}\sin(c+dx)}{3d}$$

$$+ \frac{2C(b\sec(c+dx))^{5/2}\sin(c+dx)}{5bd}$$

output

```
2/3*B*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/d+2/5*C*(b*sec(d*x+c))^(5/2)*sin(d*x+c)/b/d-6/5*b^2*C*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+6/5*b*C*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+2/3*b*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d
```

3.47.2 Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.60

$$\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{(b \sec(c + dx))^{5/2} \left(-36C \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 20B \cos^{\frac{5}{2}}(c + dx) \operatorname{EllipticF}\right)}{30bd}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `((b*Sec[c + d*x])^(5/2)*(-36*C*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 20*B*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 21*C*Sin[c + d*x] + 10*B*Sin[2*(c + d*x)] + 9*C*Sin[3*(c + d*x)])/(30*b*d)`

3.47.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.406$, Rules used = {3042, 4535, 27, 2030, 3042, 4255, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \left(b \csc\left(c + dx + \frac{\pi}{2}\right) \right)^{3/2} \left(B \csc\left(c + dx + \frac{\pi}{2}\right) + C \csc\left(c + dx + \frac{\pi}{2}\right)^2 \right) dx \\ & \quad \downarrow \text{4535} \\ & \frac{B \int (b \sec(c + dx))^{5/2} dx}{b} + \int C \sec^2(c + dx) (b \sec(c + dx))^{3/2} dx \\ & \quad \downarrow \text{27} \\ & \frac{B \int (b \sec(c + dx))^{5/2} dx}{b} + C \int \sec^2(c + dx) (b \sec(c + dx))^{3/2} dx \\ & \quad \downarrow \text{2030} \end{aligned}$$

$$\begin{aligned}
& \frac{C \int (b \sec(c + dx))^{7/2} dx}{b^2} + \frac{B \int (b \sec(c + dx))^{5/2} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{C \int (b \csc(c + dx + \frac{\pi}{2}))^{7/2} dx}{b^2} + \frac{B \int (b \csc(c + dx + \frac{\pi}{2}))^{5/2} dx}{b} \\
& \quad \downarrow \text{4255} \\
& \frac{B \left(\frac{1}{3} b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} \right)}{b} + \\
& \frac{C \left(\frac{3}{5} b^2 \int (b \sec(c + dx))^{3/2} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \right)}{b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{B \left(\frac{1}{3} b^2 \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} \right)}{b} + \\
& \frac{C \left(\frac{3}{5} b^2 \int (b \csc(c + dx + \frac{\pi}{2}))^{3/2} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \right)}{b^2} \\
& \quad \downarrow \text{4255} \\
& \frac{B \left(\frac{1}{3} b^2 \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} \right)}{b} + \\
& \frac{C \left(\frac{3}{5} b^2 \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \right) + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \right)}{b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{B \left(\frac{1}{3} b^2 \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} \right)}{b} + \\
& \frac{C \left(\frac{3}{5} b^2 \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \right)}{b^2} \\
& \quad \downarrow \text{4258} \\
& \frac{B \left(\frac{1}{3} b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2b \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} \right)}{b} + \\
& \frac{C \left(\frac{3}{5} b^2 \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) + \frac{2b \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d} \right)}{b^2} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.47. $\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx$

$$\begin{aligned}
& \frac{B\left(\frac{1}{3}b^2\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2b\sin(c+dx)(b\sec(c+dx))^{3/2}}{3d}\right)}{b} + \\
& \frac{C\left(\frac{3}{5}b^2\left(\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d}-\frac{b^2\int\sqrt{\sin(c+dx+\frac{\pi}{2})}dx}{\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}\right)+\frac{2b\sin(c+dx)(b\sec(c+dx))^{5/2}}{5d}\right)}{b^2} \\
& \quad \downarrow \text{3119} \\
& \frac{B\left(\frac{1}{3}b^2\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2b\sin(c+dx)(b\sec(c+dx))^{3/2}}{3d}\right)}{b} + \\
& \frac{C\left(\frac{3}{5}b^2\left(\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d}-\frac{2b^2E\left(\frac{1}{2}(c+dx)|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}\right)+\frac{2b\sin(c+dx)(b\sec(c+dx))^{5/2}}{5d}\right)}{b^2} \\
& \quad \downarrow \text{3120} \\
& \frac{B\left(\frac{2b^2\sqrt{\cos(c+dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx),2\right)\sqrt{b\sec(c+dx)}}{3d}+\frac{2b\sin(c+dx)(b\sec(c+dx))^{3/2}}{3d}\right)}{b} + \\
& \frac{C\left(\frac{3}{5}b^2\left(\frac{2b\sin(c+dx)\sqrt{b\sec(c+dx)}}{d}-\frac{2b^2E\left(\frac{1}{2}(c+dx)|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}\right)+\frac{2b\sin(c+dx)(b\sec(c+dx))^{5/2}}{5d}\right)}{b^2}
\end{aligned}$$

input `Int[(b*Sec[c + d*x])^(3/2)*(B*Sec[c + d*x] + C*Sec[c + d*x]^2), x]`

output `(B*((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d))/b + (C*((2*b*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*d) + (3*b^2*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/5))/b^2`

3.47.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2030 `Int[(F_x_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F_x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.47.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.92 (sec) , antiderivative size = 555, normalized size of antiderivative = 3.28

method	result
parts	$\frac{2B\sqrt{b\sec(dx+c)}b\left(-i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\cos(dx+c)-i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\right)}{3d}$
default	$-\frac{2b\sqrt{b\sec(dx+c)}\left(5iB\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)^2-9iC\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\right)}{3d}$

3.47. $\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx$

```
input int((b*sec(d*x+c))^(3/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVER
BOSE)
```

```
output 2/3*B/d*(b*sec(d*x+c))^(1/2)*b*(-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)
*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*(
1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-co
t(d*x+c)+csc(d*x+c)),I)+tan(d*x+c))+2/5*C/d*(b*sec(d*x+c))^(1/2)*b/(cos(d*
x+c)+1)*(3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*El
lipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)^2-3*I*(1/(cos(d*x+c)+1))^(
1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)
),I)*cos(d*x+c)^2+6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(
1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)-6*I*(1/(cos(d*x+c)
+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d
*x+c)),I)*cos(d*x+c)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+
1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-3*I*EllipticF(I*(cot(d*x+
c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/
2)+3*sin(d*x+c)+tan(d*x+c)+sec(d*x+c)*tan(d*x+c))
```

3.47.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.22

$$\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{-5i \sqrt{2} B b^{3/2} \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} B b^{3/2} \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - 9i \sqrt{2} C b^{3/2} \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 9i \sqrt{2} C b^{3/2} \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 * (9 C b \cos(dx + c)^2 + 5 B b \cos(dx + c) + 3 C b) \sqrt{b / \cos(dx + c)} \sin(dx + c)}{(d \cos(dx + c))^2}$$

```
input integrate((b*sec(d*x+c))^(3/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm=
"fricas")
```

```
output 1/15*(-5*I*sqrt(2)*B*b^(3/2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos
(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*B*b^(3/2)*cos(d*x + c)^2*weierst
rassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 9*I*sqrt(2)*C*b^(3/2)
*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x
+ c) + I*sin(d*x + c))) + 9*I*sqrt(2)*C*b^(3/2)*cos(d*x + c)^2*weierstrass
Zeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2
*(9*C*b*cos(d*x + c)^2 + 5*B*b*cos(d*x + c) + 3*C*b)*sqrt(b/cos(d*x + c))*
sin(d*x + c))/(d*cos(d*x + c)^2)
```

3.47. $\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.47.6 Sympy [F]

$$\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (b \sec(c + dx))^{3/2} (B + C \sec(c + dx)) \sec(c + dx) dx$$

input `integrate((b*sec(d*x+c))**(3/2)*(B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*(B + C*sec(c + d*x))*sec(c + d*x), x)`

3.47.7 Maxima [F]

$$\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + B \sec(dx + c))(b \sec(dx + c))^{3/2} dx$$

input `integrate((b*sec(d*x+c))^(3/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))*(b*sec(d*x + c))^(3/2), x)`

3.47.8 Giac [F]

$$\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + B \sec(dx + c))(b \sec(dx + c))^{3/2} dx$$

input `integrate((b*sec(d*x+c))^(3/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))*(b*sec(d*x + c))^(3/2), x)`

3.47.9 Mupad [F(-1)]

Timed out.

$$\int (b \sec(c + dx))^{3/2} (B \sec(c + dx) + C \sec^2(c + dx)) dx = \int \left(\frac{B}{\cos(c + dx)} + \frac{C}{\cos(c + dx)^2} \right) \left(\frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(3/2), x)`output `int((B/cos(c + d*x) + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(3/2), x)`

3.48 $\int \sqrt{b \sec(c + dx)}(B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.48.1	Optimal result	304
3.48.2	Mathematica [A] (verified)	305
3.48.3	Rubi [A] (verified)	305
3.48.4	Maple [C] (verified)	308
3.48.5	Fricas [C] (verification not implemented)	309
3.48.6	Sympy [F]	310
3.48.7	Maxima [F]	310
3.48.8	Giac [F]	310
3.48.9	Mupad [F(-1)]	311

3.48.1 Optimal result

Integrand size = 32, antiderivative size = 135

$$\int \sqrt{b \sec(c + dx)}(B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= -\frac{2bBE\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2C\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2B\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} + \frac{2C(b \sec(c + dx))^{3/2} \sin(c + dx)}{3bd}$$

output $2/3*C*(b*\sec(d*x+c))^(3/2)*\sin(d*x+c)/b/d-2*b*B*(\cos(1/2*d*x+1/2*c))^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticE}(\sin(1/2*d*x+1/2*c), 2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)+2*B*\sin(d*x+c)*(b*\sec(d*x+c))^(1/2)/d+2/3*C*(\cos(1/2*d*x+1/2*c))^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)*(b*\sec(d*x+c))^(1/2)/d$

3.48.2 Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.67

$$\int \sqrt{b \sec(c+dx)} (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \frac{(b \sec(c+dx))^{3/2} \left(-6B \cos^{3/2}(c+dx) E\left(\frac{1}{2}(c+dx) \mid 2\right) + 2C \cos^{3/2}(c+dx) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 2(C - \dots) \right)}{3bd}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]*(B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `((b*Sec[c + d*x])^(3/2)*(-6*B*Cos[c + d*x]^(3/2)*EllipticE[(c + d*x)/2, 2] + 2*C*Cos[c + d*x]^(3/2)*EllipticF[(c + d*x)/2, 2] + 2*(C + 3*B*Cos[c + d*x])*Sin[c + d*x]))/(3*b*d)`

3.48.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.09, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.344$, Rules used = {3042, 4535, 27, 2030, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{b \sec(c+dx)} (B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)} \left(B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2 \right) dx$$

$$\downarrow 4535$$

$$\frac{B \int (b \sec(c+dx))^{3/2} dx}{b} + \int C \sec^2(c+dx) \sqrt{b \sec(c+dx)} dx$$

$$\downarrow 27$$

$$\frac{B \int (b \sec(c+dx))^{3/2} dx}{b} + C \int \sec^2(c+dx) \sqrt{b \sec(c+dx)} dx$$

$$\downarrow 2030$$

$$\begin{aligned}
& \frac{C \int (b \sec(c+dx))^{5/2} dx}{b^2} + \frac{B \int (b \sec(c+dx))^{3/2} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{C \int (b \csc(c+dx+\frac{\pi}{2}))^{5/2} dx}{b^2} + \frac{B \int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx}{b} \\
& \quad \downarrow \text{4255} \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right)}{b} + \\
& \frac{C \left(\frac{1}{3} b^2 \int \sqrt{b \sec(c+dx)} dx + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right)}{b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right)}{b} + \\
& \frac{C \left(\frac{1}{3} b^2 \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right)}{b^2} \\
& \quad \downarrow \text{4258} \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} + \\
& \frac{C \left(\frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right)}{b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} + \\
& \frac{C \left(\frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right)}{b^2} \\
& \quad \downarrow \text{3119} \\
& \frac{C \left(\frac{1}{3} b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sin(c+dx) (b \sec(c+dx))^{3/2}}{3d} \right)}{b^2} + \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b}
\end{aligned}$$

3.48. $\int \sqrt{b \sec(c+dx)} (B \sec(c+dx) + C \sec^2(c+dx)) dx$

$$\begin{array}{c}
 \downarrow \text{3120} \\
 \frac{B\left(\frac{2b \sin(c+dx)\sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c+dx)|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}\right) + C\left(\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d}\right)}{b^2}
 \end{array}$$

input `Int[Sqrt[b*Sec[c + d*x]]*(B*Sec[c + d*x] + C*Sec[c + d*x]^2), x]`

output `(B*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d)/b + (C*((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d)))/b^2`

3.48.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2030 `Int[(Fx_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 4255 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
  && IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
  EqQ[n^2, 1/4]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[c + d*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.48.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.88 (sec) , antiderivative size = 534, normalized size of antiderivative = 3.96

method	result
parts	$\frac{2B \left(i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)), i\right) \cos(dx+c)^2 - i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticE}\left(i(-\cot(dx+c)+\csc(dx+c)), i\right) \cos(dx+c) \right)}{2\sqrt{b \sec(dx+c)} \left(3iB \operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)), i\right) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)^2 - 3iB \operatorname{EllipticE}\left(i(-\cot(dx+c)+\csc(dx+c)), i\right) \cos(dx+c) \right)}$
default	

```
input int((b*sec(d*x+c))^(1/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2), x, method=_RETURNVER
BOSE)
```

```
output 2*B/d*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)^2-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)^2+2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+sin(d*x+c)*(b*sec(d*x+c))^(1/2)/(cos(d*x+c)+1)-2/3*C/d*(b*sec(d*x+c))^(1/2)*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-tan(d*x+c))
```

3.48.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.36

$$\int \sqrt{b \sec(c + dx)} (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{-i \sqrt{2} C \sqrt{b} \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} C \sqrt{b} \cos(dx + c)}{1}$$

```
input integrate((b*sec(d*x+c))^(1/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")
```

```
output 1/3*(-I*sqrt(2)*C*sqrt(b)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*C*sqrt(b)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*B*sqrt(b)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*B*sqrt(b)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*B*cos(d*x + c) + C)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c))
```

3.48.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{b \sec(c + dx)} (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int \sqrt{b \sec(c + dx)} (B + C \sec(c + dx)) \sec(c + dx) dx \end{aligned}$$

input `integrate((b*sec(d*x+c))**(1/2)*(B*sec(d*x+c)+C*sec(d*x+c)**2), x)`

output `Integral(sqrt(b*sec(c + d*x))*(B + C*sec(c + d*x))*sec(c + d*x), x)`

3.48.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{b \sec(c + dx)} (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int (C \sec(dx + c)^2 + B \sec(dx + c)) \sqrt{b \sec(dx + c)} dx \end{aligned}$$

input `integrate((b*sec(d*x+c))^(1/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2), x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))*sqrt(b*sec(d*x + c)), x)`

3.48.8 Giac [F]

$$\begin{aligned} & \int \sqrt{b \sec(c + dx)} (B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int (C \sec(dx + c)^2 + B \sec(dx + c)) \sqrt{b \sec(dx + c)} dx \end{aligned}$$

input `integrate((b*sec(d*x+c))^(1/2)*(B*sec(d*x+c)+C*sec(d*x+c)^2), x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))*sqrt(b*sec(d*x + c)), x)`

3.48.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \sec(c + dx)} (B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int \left(\frac{B}{\cos(c + dx)} + \frac{C}{\cos(c + dx)^2} \right) \sqrt{\frac{b}{\cos(c + dx)}} dx$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(1/2), x)`output `int((B/cos(c + d*x) + C/cos(c + d*x)^2)*(b/cos(c + d*x))^(1/2), x)`

3.49 $\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.49.1 Optimal result 312
 3.49.2 Mathematica [A] (verified) 313
 3.49.3 Rubi [A] (verified) 313
 3.49.4 Maple [C] (verified) 316
 3.49.5 Fracas [C] (verification not implemented) 317
 3.49.6 Sympy [F] 317
 3.49.7 Maxima [F] 318
 3.49.8 Giac [F] 318
 3.49.9 Mupad [F(-1)] 318

3.49.1 Optimal result

Integrand size = 32, antiderivative size = 109

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= -\frac{2CE\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2B\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{bd} + \frac{2C\sqrt{b \sec(c + dx)} \sin(c + dx)}{bd}$$

output

```
-2*C*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*C*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/b/d+2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b/d
```

3.49.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.67

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{2 \left(-CE\left(\frac{1}{2}(c + dx) \mid 2\right) + B \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \frac{C \sin(c + dx)}{\sqrt{\cos(c + dx)}} \right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

input `Integrate[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/Sqrt[b*Sec[c + d*x]],x]`

output `(2*(-(C*EllipticE[(c + d*x)/2, 2]) + B*EllipticF[(c + d*x)/2, 2] + (C*Sin[c + d*x])/Sqrt[Cos[c + d*x]]))/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

3.49.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.344$, Rules used = {3042, 4535, 27, 2030, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{B \csc\left(c + dx + \frac{\pi}{2}\right) + C \csc\left(c + dx + \frac{\pi}{2}\right)^2}{\sqrt{b \csc\left(c + dx + \frac{\pi}{2}\right)}} dx$$

$$\downarrow \text{4535}$$

$$\frac{B \int \sqrt{b \sec(c + dx)} dx}{b} + \int \frac{C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$\downarrow \text{27}$$

$$\frac{B \int \sqrt{b \sec(c + dx)} dx}{b} + C \int \frac{\sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$\downarrow \text{2030}$$

$$\begin{aligned}
& \frac{C \int (b \sec(c+dx))^{3/2} dx}{b^2} + \frac{B \int \sqrt{b \sec(c+dx)} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{C \int (b \csc(c+dx+\frac{\pi}{2}))^{3/2} dx}{b^2} + \frac{B \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{b} \\
& \quad \downarrow \text{4255} \\
& \frac{C \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right)}{b^2} + \frac{B \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{C \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx \right)}{b^2} + \frac{B \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{b} \\
& \quad \downarrow \text{4258} \\
& \frac{C \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b^2} + \frac{B \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{C \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b^2} + \\
& \quad \frac{B \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
& \quad \downarrow \text{3119} \\
& \frac{B \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} + \\
& \quad \frac{C \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b^2} \\
& \quad \downarrow \text{3120} \\
& \frac{C \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b^2} + \\
& \quad \frac{2B \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{bd}
\end{aligned}$$

3.49. $\int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

input `Int[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/Sqrt[b*Sec[c + d*x]],x]`

output `(2*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b*d) + (C*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/b^2`

3.49.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2030 `Int[(F_x_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Simp[b^2*((n-2)/(n-1)) Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.49.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.84 (sec) , antiderivative size = 460, normalized size of antiderivative = 4.22

method	result
parts	$-\frac{2iB\sqrt{\frac{1}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)}{d\sqrt{b\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}} - \frac{2C(i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}})}{d\sqrt{b\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}$
default	$-\frac{2(iB \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - iC \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i))}{d\sqrt{b\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}}$

```
input int((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2), x, method=_RETURNVERBOSE)
```

```
output -2*I*B/d*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)/(b*sec(d*x+c))^(1/2)/(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)-2*C/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)), I)*sec(d*x+c)-tan(d*x+c))
```

$$3.49. \int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

3.49.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.31

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{-i \sqrt{2} B \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} B \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - i \sqrt{2} C \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \sqrt{2} C \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 C \sqrt{b} \sin(dx + c)}{b \sqrt{b \sec(c + dx)}}$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - I*sqrt(2)*C*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*C*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*C*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b*d)`

3.49.6 Sympy [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{(B + C \sec(c + dx)) \sec(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(1/2),x)`

output `Integral((B + C*sec(c + d*x))*sec(c + d*x)/sqrt(b*sec(c + d*x)), x)`

3.49.7 Maxima [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/sqrt(b*sec(d*x + c)), x)`

3.49.8 Giac [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/sqrt(b*sec(d*x + c)), x)`

3.49.9 Mupad [F(-1)]

Timed out.

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{\frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(1/2),x)`

output `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(1/2), x)`

3.50
$$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

3.50.1	Optimal result	319
3.50.2	Mathematica [A] (verified)	319
3.50.3	Rubi [A] (verified)	320
3.50.4	Maple [C] (verified)	322
3.50.5	Fricas [C] (verification not implemented)	323
3.50.6	Sympy [F]	323
3.50.7	Maxima [F]	323
3.50.8	Giac [F]	324
3.50.9	Mupad [F(-1)]	324

3.50.1 Optimal result

Integrand size = 32, antiderivative size = 85

$$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2BE(\frac{1}{2}(c+dx)|2)}{bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{2C\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)\sqrt{b \sec(c+dx)}}{b^2d}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+
1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*C*(cos(1/2*d*x
+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*
cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d
```

3.50.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

$$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx = \frac{2(BE(\frac{1}{2}(c+dx)|2)+C \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2))}{bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

```
input Integrate[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(3/2),x]
```

```
output (2*(B*EllipticE[(c + d*x)/2, 2] + C*EllipticF[(c + d*x)/2, 2]))/(b*d*Sqrt[
Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])
```


3.50.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.281$, Rules used = {3042, 4535, 27, 2030, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c+dx+\frac{\pi}{2}) + C \csc(c+dx+\frac{\pi}{2})^2}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4535} \\
 & \frac{B \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b} + \int \frac{C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{B \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b} + C \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{C \int \sqrt{b \sec(c+dx)} dx}{b^2} + \frac{B \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{C \int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{b^2} + \frac{B \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \quad \downarrow \text{4258} \\
 & \frac{C \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^2} + \frac{B \int \sqrt{\cos(c+dx)} dx}{b \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{C \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b^2} + \frac{B \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
 & \quad \downarrow \text{3119}
 \end{aligned}$$

3.50. $\int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

$$\frac{C\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx}{b^2} + \frac{2BE(\frac{1}{2}(c+dx)|2)}{bd\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}$$

↓ 3120

$$\frac{2C\sqrt{\cos(c+dx)}\operatorname{EllipticF}(\frac{1}{2}(c+dx),2)\sqrt{b\sec(c+dx)}}{b^2d} + \frac{2BE(\frac{1}{2}(c+dx)|2)}{bd\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}$$

input `Int[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(3/2), x]`

output `(2*B*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*C*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b^2*d)`

3.50.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2030 `Int[(F_x_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*F_x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.50.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.86 (sec) , antiderivative size = 414, normalized size of antiderivative = 4.87

method	result
risch	$-\frac{iB\sqrt{2}}{db\sqrt{\frac{be^{i(dx+c)}}{e^{2i(dx+c)}+1}}}-i\left(\frac{iC\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}\operatorname{EllipticF}\left(\sqrt{-i(e^{i(dx+c)}+i)},\frac{\sqrt{2}}{2}\right)}{\sqrt{be^{3i(dx+c)}+be^{i(dx+c)}}}\right)+B\left(-\frac{2(b e^{2i(dx+c)}}{b\sqrt{e^{i(dx+c)}}(be^{2i(dx+c)}+1)}\right)$
parts	$2B\left(i\operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\cos(dx+c)-i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\right)$
default	$\frac{2iB\sqrt{\left((1-\cos(dx+c))^2\csc(dx+c)^2+1\right)\left((1-\cos(dx+c))^2\csc(dx+c)^2-1\right)}\sqrt{(1-\cos(dx+c))^2\csc(dx+c)^2+1}\sqrt{-(1-\cos(dx+c))^2\csc(dx+c)^2-1}}{\left((1-\cos(dx+c))^2\csc(dx+c)^2+1\right)\left((1-\cos(dx+c))^2\csc(dx+c)^2-1\right)}$

```
input int((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -I*B/d*2^(1/2)/b/(b*exp(I*(d*x+c))/(exp(I*(d*x+c))^2+1))^(1/2)-I/d*(I*C*(-I*(exp(I*(d*x+c))+I))^(1/2)*2^(1/2)*(I*(exp(I*(d*x+c))-I))^(1/2)*(I*exp(I*(d*x+c)))^(1/2)/(b*exp(I*(d*x+c))^3+b*exp(I*(d*x+c)))^(1/2)*EllipticF((-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))+B*(-2*(b*exp(I*(d*x+c))^2+b)/b/(exp(I*(d*x+c))*(b*exp(I*(d*x+c))^2+b))^(1/2)+I*(-I*(exp(I*(d*x+c))+I))^(1/2)*2^(1/2)*(I*(exp(I*(d*x+c))-I))^(1/2)*(I*exp(I*(d*x+c)))^(1/2)/(b*exp(I*(d*x+c))^3+b*exp(I*(d*x+c)))^(1/2)*(-2*I*EllipticE((-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))+I*EllipticF((-I*(exp(I*(d*x+c))+I))^(1/2),1/2*2^(1/2))))*2^(1/2)/b/(exp(I*(d*x+c))^2+1)/(b*exp(I*(d*x+c))/(exp(I*(d*x+c))^2+1))^(1/2)*(b*exp(I*(d*x+c))*(exp(I*(d*x+c))^2+1))^(1/2)
```

$$3.50. \int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

3.50.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.44

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{-i \sqrt{2} C \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{(b \sec(c + dx))^{3/2}}$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm="fracas")`

output `(-I*sqrt(2)*C*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*C*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^2*d)`

3.50.6 Sympy [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{(B + C \sec(c + dx)) \sec(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(3/2),x)`

output `Integral((B + C*sec(c + d*x))*sec(c + d*x)/(b*sec(c + d*x))**(3/2), x)`

3.50.7 Maxima [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(3/2), x)`

3.50. $\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.50.8 Giac [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(3/2), x)`

3.50.9 Mupad [F(-1)]

Timed out.

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{\frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(3/2),x)`

output `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(3/2), x)`

3.51
$$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

3.51.1	Optimal result	325
3.51.2	Mathematica [A] (verified)	325
3.51.3	Rubi [A] (verified)	326
3.51.4	Maple [C] (verified)	329
3.51.5	Fricas [C] (verification not implemented)	329
3.51.6	Sympy [F]	330
3.51.7	Maxima [F]	330
3.51.8	Giac [F]	331
3.51.9	Mupad [F(-1)]	331

3.51.1 Optimal result

Integrand size = 32, antiderivative size = 116

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{2CE(\frac{1}{2}(c + dx)|2)}{b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2B \sqrt{\cos(c + dx)} \text{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{3b^3 d} + \frac{2B \sin(c + dx)}{3b^2 d \sqrt{b \sec(c + dx)}}$$

```
output 2/3*B*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(1/2)+2*C*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/3*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d
```

3.51.2 Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.70

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{6CE(\frac{1}{2}(c + dx)|2) + 2B \text{EllipticF}(\frac{1}{2}(c + dx), 2) + \frac{B \sin(2(c+dx))}{\sqrt{\cos(c+dx)}}}{3b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

```
input Integrate[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(5/2),x]
```

```
output (6*C*EllipticE[(c + d*x)/2, 2] + 2*B*EllipticF[(c + d*x)/2, 2] + (B*Sin[2*
(c + d*x)]/Sqrt[Cos[c + d*x]])/(3*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c +
d*x]])
```

3.51.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.344$, Rules used = {3042, 4535, 27, 2030, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4535} \\
 & \frac{B \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{b} + \int \frac{C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{B \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{b} + C \int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{C \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b^2} + \frac{B \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{C \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{b^2} + \frac{B \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} \\
 & \quad \downarrow \text{4256} \\
 & \frac{B \left(\frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{C \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{b^2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{B \left(\frac{\int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{C \int \frac{1}{\sqrt{b \csc(c+dx + \frac{\pi}{2})}} dx}{b^2} \\
& \downarrow \text{4258} \\
& \frac{B \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{C \int \sqrt{\cos(c+dx)} dx}{b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
& \downarrow \text{3042} \\
& \frac{B \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{C \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx}{b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
& \downarrow \text{3119} \\
& \frac{B \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{2CE(\frac{1}{2}(c+dx)|2)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
& \downarrow \text{3120} \\
& \frac{B \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{2CE(\frac{1}{2}(c+dx)|2)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}
\end{aligned}$$

input `Int[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(5/2), x]`

output `(2*C*EllipticE[(c + d*x)/2, 2])/(b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (B*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/b`

3.51.3.1 Defintions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2030 $\text{Int}[(Fx_*)(v_)^{(m_)*((b_*)(v_))^{(n_)}}, x_Symbol] \rightarrow \text{Simp}[1/b^m \text{ Int}[(b*v)^{(m+n)*Fx}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4256 $\text{Int}[(\text{csc}[(c_.) + (d_*)(x_)]*(b_.)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n+1)}/(b*d^n)), x] + \text{Simp}[(n+1)/(b^2*n) \text{ Int}[(b*\text{Csc}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_*)(x_)]*(b_.)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{ Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{EqQ}[n^2, 1/4]$
- rule 4535 $\text{Int}[(\text{csc}[(e_.) + (f_*)(x_)]*(b_.)^{(m_)*((A_.) + \text{csc}[(e_.) + (f_*)(x_)]*(B_.) + \text{csc}[(e_.) + (f_*)(x_)]^2*(C_.)}), x_Symbol] \rightarrow \text{Simp}[B/b \text{ Int}[(b*\text{Csc}[e + f*x])^{(m+1)}, x], x] + \text{Int}[(b*\text{Csc}[e + f*x])^m*(A + C*\text{Csc}[e + f*x]^2), x] /; \text{FreeQ}[\{b, e, f, A, B, C, m\}, x]$

3.51.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.73 (sec) , antiderivative size = 536, normalized size of antiderivative = 4.62

method	result
parts	$-\frac{2B \left(i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) + i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \right)}{3d\sqrt{b\sec(dx+c)}b^2}$
default	$-\frac{2 \left(iB \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) + 3iC \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \right)}{b^2 d \sqrt{b \sec(dx+c)}}$

input `int((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2), x, method=_RETURNVERBOSE)`

output
$$-\frac{2}{3} \frac{B}{d} \frac{1}{(b \sec(dx+c))^{1/2}} \frac{1}{b^2} \left(I \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \operatorname{EllipticF}(I(-\cot(dx+c)+\csc(dx+c)), I) + I \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \operatorname{EllipticF}(I(-\cot(dx+c)+\csc(dx+c)), I) \right) \sec(dx+c) - \sin(dx+c) + 2 \frac{C}{b^2} \frac{1}{d} \frac{1}{(\cos(dx+c)+1)} \frac{1}{(b \sec(dx+c))^{1/2}} \left(I \operatorname{EllipticE}(I(-\cot(dx+c)+\csc(dx+c)), I) \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \operatorname{EllipticF}(I(-\cot(dx+c)+\csc(dx+c)), I) \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cos(dx+c) - I \operatorname{EllipticF}(I(-\cot(dx+c)+\csc(dx+c)), I) \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \cos(dx+c) + 2 I \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \operatorname{EllipticE}(I(-\cot(dx+c)+\csc(dx+c)), I) - 2 I \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \operatorname{EllipticF}(I(-\cot(dx+c)+\csc(dx+c)), I) + I \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \operatorname{EllipticE}(I(-\cot(dx+c)+\csc(dx+c)), I) \sec(dx+c) - I \frac{1}{(\cos(dx+c)+1)^{1/2}} \frac{\cos(dx+c)}{(\cos(dx+c)+1)^{1/2}} \operatorname{EllipticF}(I(-\cot(dx+c)+\csc(dx+c)), I) \sec(dx+c) + \sin(dx+c) \right)$$

3.51.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.29

$$\int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx = \frac{2B \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) - i \sqrt{2} B \sqrt{b} \operatorname{weierstrassPInve}}{b^2 d \sqrt{b \sec(dx+c)}}$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")`

3.51.
$$\int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

output `1/3*(2*B*sqrt(b/cos(d*x + c))*cos(d*x + c)*sin(d*x + c) - I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*C*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*C*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/((b^3*d)`

3.51.6 Sympy [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{(B + C \sec(c + dx)) \sec(c + dx)}{(b \sec(c + dx))^{5/2}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(5/2), x)`

output `Integral((B + C*sec(c + d*x))*sec(c + d*x)/(b*sec(c + d*x))**(5/2), x)`

3.51.7 Maxima [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{5/2}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(5/2), x)`

3.51.8 Giac [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(5/2), x)`

3.51.9 Mupad [F(-1)]

Timed out.

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{\frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(5/2),x)`

output `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(5/2), x)`

3.52
$$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$$

3.52.1	Optimal result	332
3.52.2	Mathematica [A] (verified)	332
3.52.3	Rubi [A] (verified)	333
3.52.4	Maple [C] (verified)	336
3.52.5	Fricas [C] (verification not implemented)	337
3.52.6	Sympy [F]	337
3.52.7	Maxima [F]	337
3.52.8	Giac [F]	338
3.52.9	Mupad [F(-1)]	338

3.52.1 Optimal result

Integrand size = 32, antiderivative size = 147

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{6BE(\frac{1}{2}(c + dx) | 2)}{5b^3d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2C\sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{3b^4d} + \frac{2B \sin(c + dx)}{5b^2d(b \sec(c + dx))^{3/2}} + \frac{2C \sin(c + dx)}{3b^3d\sqrt{b \sec(c + dx)}}$$

```
output 2/5*B*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(3/2)+2/3*C*sin(d*x+c)/b^3/d/(b*sec(d*x+c))^(1/2)+6/5*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^3/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/3*C*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^4/d
```

3.52.2 Mathematica [A] (verified)

Time = 1.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.62

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{2\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}\left(9BE(\frac{1}{2}(c + dx) | 2) + 5C \operatorname{EllipticF}\right)}{15b^3d}$$

input `Integrate[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(7/2),x]`

output `(2*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]*(9*B*EllipticE[(c + d*x)/2, 2] + 5*C*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(5*C + 3*B*Cos[c + d*x])*Sin[c + d*x]))/(15*b^4*d)`

3.52.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.344$, Rules used = {3042, 4535, 27, 2030, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4535} \\
 & \frac{B \int \frac{1}{(b \sec(c + dx))^{5/2}} dx}{b} + \int \frac{C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{B \int \frac{1}{(b \sec(c + dx))^{5/2}} dx}{b} + C \int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx \\
 & \quad \downarrow \text{2030} \\
 & \frac{C \int \frac{1}{(b \sec(c + dx))^{3/2}} dx}{b^2} + \frac{B \int \frac{1}{(b \sec(c + dx))^{5/2}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{C \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx}{b^2} + \frac{B \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx}{b} \\
 & \quad \downarrow \text{4256}
 \end{aligned}$$

$$\begin{aligned}
& \frac{B\left(\frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}}\right)}{b} + \frac{C\left(\frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}\right)}{b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{B\left(\frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}}\right)}{b} + \frac{C\left(\frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}\right)}{b^2} \\
& \quad \downarrow \text{4258} \\
& \frac{B\left(\frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}}\right)}{b} + \\
& \frac{C\left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}\right)}{b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{B\left(\frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}}\right)}{b} + \\
& \frac{C\left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}\right)}{b^2} \\
& \quad \downarrow \text{3119} \\
& \frac{C\left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}\right)}{b^2} + \\
& \frac{B\left(\frac{6E\left(\frac{1}{2}(c+dx)\right)|2}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}}\right)}{b} \\
& \quad \downarrow \text{3120} \\
& \frac{B\left(\frac{6E\left(\frac{1}{2}(c+dx)\right)|2}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}}\right)}{b} + \\
& \frac{C\left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}\right)}{b^2}
\end{aligned}$$

input `Int[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(7/2), x]`

```
output (B*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c
+ d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))))/b + (C*((2*S
qrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d
) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]]))))/b^2
```

3.52.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 2030 Int[(F_x_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)
^(m + n)*F_x, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 4256 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Csc[c + d*x])^(n + 1)/(b*d*n), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c
+ d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```



```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*
(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2)
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.52.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.81 (sec) , antiderivative size = 566, normalized size of antiderivative = 3.85

method	result
parts	$\frac{2B \left(3i \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) - 3i \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c) \right)}{5}$
default	$-\frac{6iB \operatorname{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{5} + \frac{6iB \operatorname{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)), i) \sqrt{\frac{1}{\cos(dx+c)+1}} \sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}} \cos(dx+c)}{5}$

```
input int((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2), x, method=_RETURNVER
BOSE)
```

```
output 2/5*B/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^3*(3*I*(1/(cos(d*x+c)+1))^(1
/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c))
,I)*cos(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1
/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+
1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d
*x+c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*E
llipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d
*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*
x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Ellipt
icF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+cos(d*x+c)^2*sin(d*x+c)+sin(d
*x+c)*cos(d*x+c)+3*sin(d*x+c)-2/3*C/d/(b*sec(d*x+c))^(1/2)/b^3*(I*(1/(cos
(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+
c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(
1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-sin(d*x+c))
```

3.52.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.12

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{-5i \sqrt{2} C \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{(b \sec(c + dx))^{7/2}}$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm="fricas")`

output `1/15*(-5*I*sqrt(2)*C*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*C*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 9*I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 9*I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*B*cos(d*x + c)^2 + 5*C*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^4*d)`

3.52.6 Sympy [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{(B + C \sec(c + dx)) \sec(c + dx)}{(b \sec(c + dx))^{7/2}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(7/2),x)`

output `Integral((B + C*sec(c + d*x))*sec(c + d*x)/(b*sec(c + d*x))**(7/2), x)`

3.52.7 Maxima [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{7/2}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(7/2), x)`

3.52.8 Giac [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{7/2}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(7/2), x)`

3.52.9 Mupad [F(-1)]

Timed out.

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{\frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{7/2}} dx$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(7/2),x)`

output `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(7/2), x)`

3.53
$$\int \frac{B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx$$

3.53.1	Optimal result	339
3.53.2	Mathematica [A] (verified)	340
3.53.3	Rubi [A] (verified)	340
3.53.4	Maple [C] (verified)	344
3.53.5	Fricas [C] (verification not implemented)	344
3.53.6	Sympy [F(-1)]	345
3.53.7	Maxima [F]	345
3.53.8	Giac [F]	346
3.53.9	Mupad [F(-1)]	346

3.53.1 Optimal result

Integrand size = 32, antiderivative size = 176

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \frac{6CE(\frac{1}{2}(c + dx)|2)}{5b^4d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{10B\sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{21b^5d} + \frac{2B \sin(c + dx)}{7b^2d(b \sec(c + dx))^{5/2}} + \frac{2C \sin(c + dx)}{5b^3d(b \sec(c + dx))^{3/2}} + \frac{10B \sin(c + dx)}{21b^4d\sqrt{b \sec(c + dx)}}$$

```
output 2/7*B*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(5/2)+2/5*C*sin(d*x+c)/b^3/d/(b*sec(d*x+c))^(3/2)+10/21*B*sin(d*x+c)/b^4/d/(b*sec(d*x+c))^(1/2)+6/5*C*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^4/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+10/21*B*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^5/d
```

3.53.2 Mathematica [A] (verified)

Time = 1.32 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.59

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \frac{\sqrt{b \sec(c + dx)} \left(252C \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 100B \sqrt{\cos(c + dx)} \right)}{(b \sec(c + dx))^{9/2}}$$

input `Integrate[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(9/2),x]`

output `(Sqrt[b*Sec[c + d*x]]*(252*C*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 100*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (65*B + 42*C*Cos[c + d*x] + 15*B*Cos[2*(c + d*x)])*Sin[2*(c + d*x)])/(210*b^5*d)`

3.53.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.09, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.406$, Rules used = {3042, 4535, 27, 2030, 3042, 4256, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{B \csc\left(c + dx + \frac{\pi}{2}\right) + C \csc\left(c + dx + \frac{\pi}{2}\right)^2}{(b \csc\left(c + dx + \frac{\pi}{2}\right))^{9/2}} dx \\ & \quad \downarrow \text{4535} \\ & \frac{B \int \frac{1}{(b \sec(c + dx))^{7/2}} dx}{b} + \int \frac{C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx \\ & \quad \downarrow \text{27} \\ & \frac{B \int \frac{1}{(b \sec(c + dx))^{7/2}} dx}{b} + C \int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx \\ & \quad \downarrow \text{2030} \\ & \frac{C \int \frac{1}{(b \sec(c + dx))^{5/2}} dx}{b^2} + \frac{B \int \frac{1}{(b \sec(c + dx))^{7/2}} dx}{b} \end{aligned}$$

3.53. $\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{C \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{b^2} + \frac{B \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx}{b} \\
 & \downarrow 4256 \\
 & \frac{B \left(\frac{5 \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)}{b} + \frac{C \left(\frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b^2} \\
 & \downarrow 3042 \\
 & \frac{B \left(\frac{5 \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)}{b} + \frac{C \left(\frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b^2} \\
 & \downarrow 4256 \\
 & \frac{B \left(\frac{5 \left(\frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)}{b} + \\
 & \frac{C \left(\frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b^2} \\
 & \downarrow 3042 \\
 & \frac{B \left(\frac{5 \left(\frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)}{b} + \\
 & \frac{C \left(\frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b^2} \\
 & \downarrow 4258 \\
 & \frac{B \left(\frac{5 \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right)}{b} + \\
 & \frac{C \left(\frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b^2}
 \end{aligned}$$

3.53. $\int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx$

$$\begin{array}{c}
 \downarrow 3042 \\
 B \left(\frac{5 \left(\frac{\int \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}{3b^2} dx + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 \hline
 C \left(\frac{3 \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})}}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} dx + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}}}{b^2} \right) \\
 \downarrow 3119 \\
 B \left(\frac{5 \left(\frac{\int \frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}{3b^2} dx + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 \hline
 C \left(\frac{6E\left(\frac{1}{2}(c+dx)|2\right)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 \downarrow 3120 \\
 B \left(\frac{5 \left(\frac{2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}} \right) \\
 \hline
 C \left(\frac{6E\left(\frac{1}{2}(c+dx)|2\right)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right) \\
 \downarrow
 \end{array}$$

input `Int[(B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(9/2), x]`

output `(C*((6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2)))/b^2 + (B*((2*Sin[c + d*x])/(7*b*d*(b*Sec[c + d*x])^(5/2)) + (5*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/(7*b^2))/b`

3.53.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 2030 `Int[(Fx_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[1/b^m Int[(b*v)^(m+n)*Fx, x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 4256 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d*n)), x] + Simp[(n+1)/(b^2*n) Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4535 `Int[(csc[(e_.) + (f_)*(x_)]*(b_))^(m_)*((A_.) + csc[(e_.) + (f_)*(x_)]*(B_.) + csc[(e_.) + (f_)*(x_)]^2*(C_)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m+1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.53.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.72 (sec) , antiderivative size = 582, normalized size of antiderivative = 3.31

method	result
parts	$-\frac{2B\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i)\right)}{21d\sqrt{b\sec(dx+c)}b^4}$
default	$-\frac{2(-63iC\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i)\sec(dx+c)+50iB\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticE}(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i))}{21d\sqrt{b\sec(dx+c)}b^4}$

input `int((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x,method=_RETURNVERBOSE)`

output

```
-2/21*B/d/(b*sec(d*x+c))^(1/2)/b^4*(5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*cos(d*x+c)^2*sin(d*x+c)-5*sin(d*x+c))+2/5*C/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^4*(3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+cos(d*x+c)^2*sin(d*x+c)+sin(d*x+c)*cos(d*x+c)+3*sin(d*x+c))
```

3.53.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.99

$$\int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx = \frac{-25i\sqrt{2}B\sqrt{b}\operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c))}{(b \sec(c+dx))^{9/2}}$$

3.53. $\int \frac{B \sec(c+dx) + C \sec^2(c+dx)}{(b \sec(c+dx))^{9/2}} dx$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x, algorithm="fricas")`

output `1/105*(-25*I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 25*I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 63*I*sqrt(2)*C*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 63*I*sqrt(2)*C*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(15*B*cos(d*x + c)^3 + 21*C*cos(d*x + c)^2 + 25*B*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b^5*d)`

3.53.6 Sympy [F(-1)]

Timed out.

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \text{Timed out}$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(9/2),x)`

output Timed out

3.53.7 Maxima [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{9/2}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(9/2), x)`

3.53.8 Giac [F]

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c)}{(b \sec(dx + c))^{\frac{9}{2}}} dx$$

input `integrate((B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(9/2),x, algorithm="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c))/(b*sec(d*x + c))^(9/2), x)`

3.53.9 Mupad [F(-1)]

Timed out.

$$\int \frac{B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{9/2}} dx = \int \frac{\frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{9/2}} dx$$

input `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(9/2),x)`

output `int((B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(9/2), x)`

3.54 $\int \sec^4(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.54.1	Optimal result	347
3.54.2	Mathematica [A] (verified)	347
3.54.3	Rubi [A] (verified)	348
3.54.4	Maple [A] (verified)	351
3.54.5	Fricas [A] (verification not implemented)	351
3.54.6	Sympy [F]	352
3.54.7	Maxima [A] (verification not implemented)	352
3.54.8	Giac [B] (verification not implemented)	353
3.54.9	Mupad [B] (verification not implemented)	353

3.54.1 Optimal result

Integrand size = 29, antiderivative size = 122

$$\int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3B \operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{(5A + 4C) \tan(c + dx)}{5d} + \frac{3B \sec(c + dx) \tan(c + dx)}{8d}$$

$$+ \frac{B \sec^3(c + dx) \tan(c + dx)}{4d} + \frac{C \sec^4(c + dx) \tan(c + dx)}{5d} + \frac{(5A + 4C) \tan^3(c + dx)}{15d}$$

output `3/8*B*arctanh(sin(d*x+c))/d+1/5*(5*A+4*C)*tan(d*x+c)/d+3/8*B*sec(d*x+c)*tan(d*x+c)/d+1/4*B*sec(d*x+c)^3*tan(d*x+c)/d+1/5*C*sec(d*x+c)^4*tan(d*x+c)/d+1/15*(5*A+4*C)*tan(d*x+c)^3/d`

3.54.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.66

$$\int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{45B \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (45B \sec(c + dx) + 30B \sec^3(c + dx) + 8(15(A + C) + 5(A + 2C)))}{120d}$$

input `Integrate[Sec[c + d*x]^4*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output $(45*B*ArcTanh[\sin[c + dx]] + \tan[c + dx]*(45*B*\sec[c + dx] + 30*B*\sec[c + dx]^3 + 8*(15*(A + C) + 5*(A + 2*C))*\tan[c + dx]^2 + 3*C*\tan[c + dx]^4))/(120*d)$

3.54.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.98, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 4535, 3042, 4255, 3042, 4255, 3042, 4257, 4534, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^4 \left(A + B \csc\left(c + dx + \frac{\pi}{2}\right) + C \csc\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4535} \\
 & \int \sec^4(c + dx) (C \sec^2(c + dx) + A) dx + B \int \sec^5(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^4 \left(C \csc\left(c + dx + \frac{\pi}{2}\right)^2 + A\right) dx + B \int \csc\left(c + dx + \frac{\pi}{2}\right)^5 dx \\
 & \quad \downarrow \text{4255} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^4 \left(C \csc\left(c + dx + \frac{\pi}{2}\right)^2 + A\right) dx + \\
 & \quad B \left(\frac{3}{4} \int \sec^3(c + dx) dx + \frac{\tan(c + dx) \sec^3(c + dx)}{4d}\right) \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^4 \left(C \csc\left(c + dx + \frac{\pi}{2}\right)^2 + A\right) dx + \\
 & \quad B \left(\frac{3}{4} \int \csc\left(c + dx + \frac{\pi}{2}\right)^3 dx + \frac{\tan(c + dx) \sec^3(c + dx)}{4d}\right) \\
 & \quad \downarrow \text{4255}
 \end{aligned}$$

$$\begin{aligned}
& \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + \\
& B\left(\frac{3}{4}\left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d}\right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d}\right) \\
& \quad \downarrow \text{3042} \\
& \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + \\
& B\left(\frac{3}{4}\left(\frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d}\right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d}\right) \\
& \quad \downarrow \text{4257} \\
& \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + \\
& B\left(\frac{3}{4}\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d}\right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d}\right) \\
& \quad \downarrow \text{4534} \\
& \frac{1}{5}(5A+4C) \int \sec^4(c+dx) dx + \\
& B\left(\frac{3}{4}\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d}\right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d}\right) + \\
& \quad \frac{C \tan(c+dx) \sec^4(c+dx)}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5}(5A+4C) \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 dx + \\
& B\left(\frac{3}{4}\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d}\right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d}\right) + \\
& \quad \frac{C \tan(c+dx) \sec^4(c+dx)}{5d} \\
& \quad \downarrow \text{4254} \\
& -\frac{(5A+4C) \int (\tan^2(c+dx)+1) d(-\tan(c+dx))}{5d} + \\
& B\left(\frac{3}{4}\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d}\right) + \frac{\tan(c+dx) \sec^3(c+dx)}{4d}\right) + \\
& \quad \frac{C \tan(c+dx) \sec^4(c+dx)}{5d} \\
& \quad \downarrow \text{2009}
\end{aligned}$$

$$\begin{aligned}
 & - \frac{(5A + 4C) \left(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx) \right)}{5d} + \\
 & B \left(\frac{3}{4} \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{\tan(c + dx) \sec^3(c + dx)}{4d} \right) + \\
 & \frac{C \tan(c + dx) \sec^4(c + dx)}{5d}
 \end{aligned}$$

input `Int[Sec[c + d*x]^4*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2), x]`

output `(C*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) - ((5*A + 4*C)*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/(5*d) + B*((Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (3*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/4)`

3.54.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_. + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

3.54. $\int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*
(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.54.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.85

method	result
derivativedivides	$-A\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)+B\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)-C\left(-\frac{8}{15}\right)$
default	$-A\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)+B\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)-C\left(-\frac{8}{15}\right)$
parts	$\frac{A\left(-\frac{2}{3}-\frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)}{d} + \frac{B\left(-\left(-\frac{\sec(dx+c)^3}{4}-\frac{3\sec(dx+c)}{8}\right)\tan(dx+c)+\frac{3\ln(\sec(dx+c)+\tan(dx+c))}{8}\right)}{d}$
risch	$\frac{i(45B e^{9i(dx+c)}+210B e^{7i(dx+c)}-240A e^{6i(dx+c)}-560A e^{4i(dx+c)}-640C e^{4i(dx+c)}-210B e^{3i(dx+c)}-400A e^{2i(dx+c)})}{60d(e^{2i(dx+c)}+1)^5}$
norman	$\frac{4(25A+29C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^5}{15d}-\frac{(8A-5B+8C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^9}{4d}-\frac{(8A+5B+8C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{4d}+\frac{(32A-3B+16C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^7}{6d}+\frac{(32A+16C)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{6d}$
parallelrisch	$\frac{-450B\left(\frac{\cos(5dx+5c)}{10}+\frac{\cos(3dx+3c)}{2}+\cos(dx+c)\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+450B\left(\frac{\cos(5dx+5c)}{10}+\frac{\cos(3dx+3c)}{2}+\cos(dx+c)\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{120d(\cos(5dx+5c)+1)}$

```
input int(sec(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)
```

```
output 1/d*(-A*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+B*(-(-1/4*sec(d*x+c)^3-3/8*sec(
d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))-C*(-8/15-1/5*sec(d*x+c)^
4-4/15*sec(d*x+c)^2)*tan(d*x+c))
```

3.54.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00

$$\int \sec^4(c+dx) (A+B\sec(c+dx)+C\sec^2(c+dx)) dx$$

$$= \frac{45B\cos(dx+c)^5\log(\sin(dx+c)+1)-45B\cos(dx+c)^5\log(-\sin(dx+c)+1)+2(16(5A+4C)\cos(dx+c)+240d\cos(dx+c))\tan(dx+c)}{240d\cos(dx+c)}$$

3.54. $\int \sec^4(c+dx) (A+B\sec(c+dx)+C\sec^2(c+dx)) dx$

input `integrate(sec(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/240*(45*B*cos(d*x + c)^5*log(sin(d*x + c) + 1) - 45*B*cos(d*x + c)^5*log(-sin(d*x + c) + 1) + 2*(16*(5*A + 4*C)*cos(d*x + c)^4 + 45*B*cos(d*x + c)^3 + 8*(5*A + 4*C)*cos(d*x + c)^2 + 30*B*cos(d*x + c) + 24*C)*sin(d*x + c))/(d*cos(d*x + c)^5)`

3.54.6 Sympy [F]

$$\begin{aligned} & \int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \sec^4(c + dx) dx \end{aligned}$$

input `integrate(sec(d*x+c)**4*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*sec(c + d*x)**4, x)`

3.54.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.04

$$\begin{aligned} & \int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{80 (\tan(dx + c)^3 + 3 \tan(dx + c))A + 16 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 + 15 \tan(dx + c))C - 15B}{240 d} \end{aligned}$$

input `integrate(sec(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/240*(80*(tan(d*x + c)^3 + 3*tan(d*x + c))*A + 16*(3*tan(d*x + c)^5 + 10*tan(d*x + c)^3 + 15*tan(d*x + c))*C - 15*B*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)))/d`

3.54. $\int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.54.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. $2(110) = 220$.

Time = 0.31 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.02

$$\int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{45 B \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 45 B \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) - \frac{2 \left(120 A \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^9 - 75 B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^9 + 120 C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^9 - 320 A \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^7 + 30 B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^7 - 160 C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^7 + 400 A \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^5 + 464 C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^5 - 320 A \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 - 30 B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 - 160 C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 + 120 A \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 75 B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 120 C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 - 1}}{d}$$

input `integrate(sec(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/120*(45*B*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 45*B*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(120*A*tan(1/2*d*x + 1/2*c)^9 - 75*B*tan(1/2*d*x + 1/2*c)^9 + 120*C*tan(1/2*d*x + 1/2*c)^9 - 320*A*tan(1/2*d*x + 1/2*c)^7 + 30*B*tan(1/2*d*x + 1/2*c)^7 - 160*C*tan(1/2*d*x + 1/2*c)^7 + 400*A*tan(1/2*d*x + 1/2*c)^5 + 464*C*tan(1/2*d*x + 1/2*c)^5 - 320*A*tan(1/2*d*x + 1/2*c)^3 - 30*B*tan(1/2*d*x + 1/2*c)^3 - 160*C*tan(1/2*d*x + 1/2*c)^3 + 120*A*tan(1/2*d*x + 1/2*c) + 75*B*tan(1/2*d*x + 1/2*c) + 120*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)/d`

3.54.9 Mupad [B] (verification not implemented)

Time = 17.67 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.61

$$\int \sec^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{3 B \operatorname{atanh} \left(\tan \left(\frac{c}{2} + \frac{dx}{2} \right) \right)}{4 d}$$

$$- \frac{\left(2 A - \frac{5 B}{4} + 2 C \right) \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^9 + \left(\frac{B}{2} - \frac{16 A}{3} - \frac{8 C}{3} \right) \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^7 + \left(\frac{20 A}{3} + \frac{116 C}{15} \right) \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^5 + \left(- \frac{2 A}{3} - \frac{5 B}{15} - \frac{2 C}{3} \right) \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^3 + \left(\frac{A}{3} + \frac{B}{15} + \frac{C}{3} \right) \tan \left(\frac{c}{2} + \frac{dx}{2} \right) + \frac{d \left(\tan \left(\frac{c}{2} + \frac{dx}{2} \right)^{10} - 5 \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^8 + 10 \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^6 - 10 \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^4 + 5 \tan \left(\frac{c}{2} + \frac{dx}{2} \right)^2 - 1 \right)}{d \left(\tan \left(\frac{c}{2} + \frac{dx}{2} \right)^2 - 1 \right)^5}$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/cos(c + d*x)^4,x)`

output $(3*B*atanh(\tan(c/2 + (d*x)/2)))/(4*d) - (\tan(c/2 + (d*x)/2)^5*((20*A)/3 + (116*C)/15) + \tan(c/2 + (d*x)/2)*(2*A + (5*B)/4 + 2*C) + \tan(c/2 + (d*x)/2)^9*(2*A - (5*B)/4 + 2*C) - \tan(c/2 + (d*x)/2)^3*((16*A)/3 + B/2 + (8*C)/3) - \tan(c/2 + (d*x)/2)^7*((16*A)/3 - B/2 + (8*C)/3)/(d*(5*\tan(c/2 + (d*x)/2)^2 - 10*\tan(c/2 + (d*x)/2)^4 + 10*\tan(c/2 + (d*x)/2)^6 - 5*\tan(c/2 + (d*x)/2)^8 + \tan(c/2 + (d*x)/2)^{10} - 1))$

3.55 $\int \sec^3(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.55.1	Optimal result	355
3.55.2	Mathematica [A] (verified)	355
3.55.3	Rubi [A] (verified)	356
3.55.4	Maple [A] (verified)	358
3.55.5	Fricas [A] (verification not implemented)	359
3.55.6	Sympy [F]	359
3.55.7	Maxima [A] (verification not implemented)	359
3.55.8	Giac [B] (verification not implemented)	360
3.55.9	Mupad [B] (verification not implemented)	361

3.55.1 Optimal result

Integrand size = 29, antiderivative size = 97

$$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(4A + 3C)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{B \tan(c + dx)}{d}$$

$$+ \frac{(4A + 3C)\sec(c + dx)\tan(c + dx)}{8d} + \frac{C \sec^3(c + dx)\tan(c + dx)}{4d} + \frac{B \tan^3(c + dx)}{3d}$$

output `1/8*(4*A+3*C)*arctanh(sin(d*x+c))/d+B*tan(d*x+c)/d+1/8*(4*A+3*C)*sec(d*x+c)*tan(d*x+c)/d+1/4*C*sec(d*x+c)^3*tan(d*x+c)/d+1/3*B*tan(d*x+c)^3/d`

3.55.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.73

$$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(4A + 3C)\operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (3(4A + 3C)\sec(c + dx) + 6C \sec^3(c + dx) + 8B(3 + \tan^2(c + dx)))}{24d}$$

input `Integrate[Sec[c + d*x]^3*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(3*(4*A + 3*C)*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(3*(4*A + 3*C)*Sec[c + d*x] + 6*C*Sec[c + d*x]^3 + 8*B*(3 + Tan[c + d*x]^2)))/(24*d)`

3.55. $\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.55.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4535, 3042, 4254, 2009, 4534, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c+dx) (A + B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 \left(A + B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4535} \\
 & \int \sec^3(c+dx) (C \sec^2(c+dx) + A) dx + B \int \sec^4(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + B \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{4254} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx - \frac{B \int (\tan^2(c+dx) + 1) d(-\tan(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx - \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{4}(4A + 3C) \int \sec^3(c+dx) dx - \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} + \frac{C \tan(c+dx) \sec^3(c+dx)}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4}(4A + 3C) \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx - \frac{B(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d} + \\
 & \quad \frac{C \tan(c+dx) \sec^3(c+dx)}{4d} \\
 & \quad \downarrow \text{4255}
 \end{aligned}$$

$$\frac{\frac{1}{4}(4A + 3C) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{B(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d} + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d}}{\downarrow 3042}$$

$$\frac{\frac{1}{4}(4A + 3C) \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{B(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d} + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d}}{\downarrow 4257}$$

$$\frac{\frac{1}{4}(4A + 3C) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{B(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d} + \frac{C \tan(c + dx) \sec^3(c + dx)}{4d}}$$

input `Int[Sec[c + d*x]^3*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2), x]`

output `(C*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((4*A + 3*C)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/4 - (B*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/d`

3.55.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
;/; FreeQ[{c, d}, x]
```

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]^2*(C_.)
+ (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1)
)), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_.)]*
(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2)
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.55.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{A\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) - B\left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3}\right)\tan(dx+c) + C\left(-\left(-\frac{\sec(dx+c)^3}{4} - \frac{3\sec(dx+c)}{8}\right)\right)}{d}$
default	$\frac{A\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) - B\left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3}\right)\tan(dx+c) + C\left(-\left(-\frac{\sec(dx+c)^3}{4} - \frac{3\sec(dx+c)}{8}\right)\right)}{d}$
parts	$\frac{A\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)}{d} - \frac{B\left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3}\right)\tan(dx+c)}{d} + \frac{C\left(-\left(-\frac{\sec(dx+c)^3}{4} - \frac{3\sec(dx+c)}{8}\right)\right)}{d}$
parallelrisch	$\frac{-48\left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4}\right) + \cos(2dx+2c)\left(A + \frac{3C}{4}\right)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 48\left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4}\right) + \cos(2dx+2c)\left(A + \frac{3C}{4}\right)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{24d(\cos(4dx+4c) + 4\cos(2dx+2c))}$
norman	$\frac{(4A+5C-8B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^7}{4d} + \frac{(4A+5C+8B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4d} - \frac{(12A-9C-40B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{12d} - \frac{(12A-9C+40B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{12d} - \frac{A}{d} - \frac{B}{d} + \frac{C}{d}$
risch	$\frac{i(12Ae^{7i(dx+c)} + 9Ce^{7i(dx+c)} + 12Ae^{5i(dx+c)} + 33Ce^{5i(dx+c)} - 48Be^{4i(dx+c)} - 12Ae^{3i(dx+c)} - 33Ce^{3i(dx+c)} - 64Be^{2i(dx+c)} - 12Ae^{2i(dx+c)} - 9Ce^{2i(dx+c)} - 12Ae^{i(dx+c)} - 9Ce^{i(dx+c)} - 12A - 9C)}{12d(e^{2i(dx+c)} + 1)^4}$

```
input int(sec(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2), x, method=_RETURNVERBOSE)
```

```
output 1/d*(A*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))-B*(-2/3-1
/3*sec(d*x+c)^2)*tan(d*x+c)+C*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x
+c)+3/8*ln(sec(d*x+c)+tan(d*x+c))))
```

$$3.55. \int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

3.55.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.21

$$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(4A + 3C) \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 3(4A + 3C) \cos(dx + c)^4 \log(-\sin(dx + c) + 1) + 2(16B \cos(dx + c)^3 + 3(4A + 3C) \cos(dx + c)^2 + 8B \cos(dx + c) + 6C) \sin(dx + c)}{48 d \cos(dx + c)^4}$$

input `integrate(sec(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/48*(3*(4*A + 3*C)*cos(d*x + c)^4*log(sin(d*x + c) + 1) - 3*(4*A + 3*C)*cos(d*x + c)^4*log(-sin(d*x + c) + 1) + 2*(16*B*cos(d*x + c)^3 + 3*(4*A + 3*C)*cos(d*x + c)^2 + 8*B*cos(d*x + c) + 6*C)*sin(d*x + c))/(d*cos(d*x + c)^4)`

3.55.6 Sympy [F]

$$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \sec^3(c + dx) dx$$

input `integrate(sec(d*x+c)**3*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*sec(c + d*x)**3, x)`

3.55.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.43

$$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{16(\tan(dx + c)^3 + 3 \tan(dx + c))B - 3C \left(\frac{2(3 \sin(dx+c)^3 - 5 \sin(dx+c))}{\sin(dx+c)^4 - 2 \sin(dx+c)^2 + 1} - 3 \log(\sin(dx + c) + 1) + 3 \log(\sin(dx + c) - 1) \right)}{48 d}$$

3.55. $\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

input `integrate(sec(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output $\frac{1}{48}*(16*(\tan(dx + c)^3 + 3*\tan(dx + c))*B - 3*C*(2*(3*\sin(dx + c)^3 - 5*\sin(dx + c))/(\sin(dx + c)^4 - 2*\sin(dx + c)^2 + 1) - 3*\log(\sin(dx + c) + 1) + 3*\log(\sin(dx + c) - 1)) - 12*A*(2*\sin(dx + c)/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)))/d$

3.55.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 230 vs. $2(89) = 178$.

Time = 0.31 (sec) , antiderivative size = 230, normalized size of antiderivative = 2.37

$$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(4A + 3C) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - 3(4A + 3C) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right) + \frac{2(12A \tan(\frac{1}{2} dx + \frac{1}{2} c))^7}{\dots}}{\dots}$$

input `integrate(sec(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output $\frac{1}{24}*(3*(4*A + 3*C)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(4*A + 3*C)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) + 2*(12*A*\tan(1/2*d*x + 1/2*c)^7 - 24*B*\tan(1/2*d*x + 1/2*c)^7 + 15*C*\tan(1/2*d*x + 1/2*c)^7 - 12*A*\tan(1/2*d*x + 1/2*c)^5 + 40*B*\tan(1/2*d*x + 1/2*c)^5 + 9*C*\tan(1/2*d*x + 1/2*c)^5 - 12*A*\tan(1/2*d*x + 1/2*c)^3 - 40*B*\tan(1/2*d*x + 1/2*c)^3 + 9*C*\tan(1/2*d*x + 1/2*c)^3 + 12*A*\tan(1/2*d*x + 1/2*c) + 24*B*\tan(1/2*d*x + 1/2*c) + 15*C*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^4/d$

3.55.9 Mupad [B] (verification not implemented)

Time = 17.94 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.65

$$\int \sec^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) \left(A + \frac{3C}{4}\right)}{d} + \frac{\left(A - 2B + \frac{5C}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{10B}{3} - A + \frac{3C}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{3C}{4} - \frac{10B}{3} - A\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{10B}{3} - A + \frac{3C}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)}$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/cos(c + d*x)^3,x)`output `(atanh(tan(c/2 + (d*x)/2))*(A + (3*C)/4))/d + (tan(c/2 + (d*x)/2)*(A + 2*B + (5*C)/4) + tan(c/2 + (d*x)/2)^7*(A - 2*B + (5*C)/4) - tan(c/2 + (d*x)/2)^3*(A + (10*B)/3 - (3*C)/4) + tan(c/2 + (d*x)/2)^5*((10*B)/3 - A + (3*C)/4))/(d*(6*tan(c/2 + (d*x)/2)^4 - 4*tan(c/2 + (d*x)/2)^2 - 4*tan(c/2 + (d*x)/2)/2)^6 + tan(c/2 + (d*x)/2)^8 + 1))`

3.56 $\int \sec^2(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.56.1	Optimal result	362
3.56.2	Mathematica [A] (verified)	362
3.56.3	Rubi [A] (verified)	363
3.56.4	Maple [A] (verified)	365
3.56.5	Fricas [A] (verification not implemented)	366
3.56.6	Sympy [F]	366
3.56.7	Maxima [A] (verification not implemented)	367
3.56.8	Giac [B] (verification not implemented)	367
3.56.9	Mupad [B] (verification not implemented)	368

3.56.1 Optimal result

Integrand size = 29, antiderivative size = 78

$$\begin{aligned} & \int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{\text{Barctanh}(\sin(c + dx))}{2d} + \frac{(3A + 2C) \tan(c + dx)}{3d} \\ & \quad + \frac{B \sec(c + dx) \tan(c + dx)}{2d} + \frac{C \sec^2(c + dx) \tan(c + dx)}{3d} \end{aligned}$$

output `1/2*B*arctanh(sin(d*x+c))/d+1/3*(3*A+2*C)*tan(d*x+c)/d+1/2*B*sec(d*x+c)*tan(d*x+c)/d+1/3*C*sec(d*x+c)^2*tan(d*x+c)/d`

3.56.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.65

$$\begin{aligned} & \int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{3\text{Barctanh}(\sin(c + dx)) + \tan(c + dx) (6(A + C) + 3B \sec(c + dx) + 2C \tan^2(c + dx))}{6d} \end{aligned}$$

input `Integrate[Sec[c + d*x]^2*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(3*B*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(6*(A + C) + 3*B*Sec[c + d*x] + 2*C*Tan[c + d*x]^2))/(6*d)`

3.56.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4535, 3042, 4255, 3042, 4257, 4534, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx) (A + B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 \left(A + B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4535} \\
 & \int \sec^2(c+dx) (C \sec^2(c+dx) + A) dx + B \int \sec^3(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + B \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{4255} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + \\
 & \quad B \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + \\
 & \quad B \left(\frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) \\
 & \quad \downarrow \text{4257} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + \\
 & \quad B \left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) \\
 & \quad \downarrow \text{4534}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{3}(3A + 2C) \int \sec^2(c + dx) dx + B \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \\
& \quad \frac{C \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3}(3A + 2C) \int \csc \left(c + dx + \frac{\pi}{2} \right)^2 dx + B \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \\
& \quad \frac{C \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{4254} \\
& -\frac{(3A + 2C) \int 1d(-\tan(c + dx))}{3d} + B \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \\
& \quad \frac{C \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{24} \\
& \frac{(3A + 2C) \tan(c + dx)}{3d} + B \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \\
& \quad \frac{C \tan(c + dx) \sec^2(c + dx)}{3d}
\end{aligned}$$

input `Int[Sec[c + d*x]^2*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2), x]`

output `((3*A + 2*C)*Tan[c + d*x])/(3*d) + (C*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + B*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d))`

3.56.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.56.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{A \tan(dx+c) + B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) - C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$
default	$\frac{A \tan(dx+c) + B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) - C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$
parts	$\frac{A \tan(dx+c)}{d} + \frac{B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d} - \frac{C \left(-\frac{2}{3} - \frac{\sec(dx+c)^2}{3} \right) \tan(dx+c)}{d}$
norman	$\frac{4(3A+C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{3d} - \frac{(2A-B+2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{d} - \frac{(2A+B+2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} - \frac{B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2d} + \frac{B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2d}$
risch	$-\frac{i(3B e^{5i(dx+c)} - 6A e^{4i(dx+c)} - 12A e^{2i(dx+c)} - 12C e^{2i(dx+c)} - 3B e^{i(dx+c)} - 6A - 4C)}{3d(e^{2i(dx+c)} + 1)^3} + \frac{B \ln(e^{i(dx+c)} + i)}{2d} - \frac{B \ln(e^{i(dx+c)} - i)}{2d}$
parallelrisc	$-\frac{9 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 9 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) B \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + (6A+4C) \sin(dx+c)}{6d(\cos(3dx+3c) + 3 \cos(dx+c))}$

3.56. $\int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

input `int(sec(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(A*tan(d*x+c)+B*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c))) - C*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c))`

3.56.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.21

$$\int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3 B \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3 B \cos(dx + c)^3 \log(-\sin(dx + c) + 1) + 2(2(3A + 2C) \cos(dx + c) + 2C) \sin(dx + c)}{12 d \cos(dx + c)^3}$$

input `integrate(sec(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/12*(3*B*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*B*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(2*(3*A + 2*C)*cos(d*x + c)^2 + 3*B*cos(d*x + c) + 2*C)*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.56.6 Sympy [F]

$$\int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \sec^2(c + dx) dx$$

input `integrate(sec(d*x+c)**2*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*sec(c + d*x)**2, x)`

3.56.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.01

$$\int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{4 (\tan(dx + c))^3 + 3 \tan(dx + c) C - 3 B \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right)}{12 d}$$

input `integrate(sec(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/12*(4*(tan(d*x + c)^3 + 3*tan(d*x + c))*C - 3*B*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 12*A*tan(d*x + c))/d`

3.56.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(70) = 140$.

Time = 0.30 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.08

$$\int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3 B \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 3 B \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) - \frac{2 \left(6 A \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^5 - 3 B \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^5 + 6 C \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 \right)}{6 d}$$

input `integrate(sec(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/6*(3*B*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*B*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(6*A*tan(1/2*d*x + 1/2*c)^5 - 3*B*tan(1/2*d*x + 1/2*c)^5 + 6*C*tan(1/2*d*x + 1/2*c)^3 - 12*A*tan(1/2*d*x + 1/2*c)^3 - 4*C*tan(1/2*d*x + 1/2*c)^3 + 6*A*tan(1/2*d*x + 1/2*c) + 3*B*tan(1/2*d*x + 1/2*c) + 6*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^3/d`

3.56.9 Mupad [B] (verification not implemented)

Time = 17.10 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.58

$$\int \sec^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{B \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d} - \frac{(2A - B + 2C) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-4A - \frac{4C}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (2A + B + 2C) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1 \right)}$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/cos(c + d*x)^2,x)`

output `(B*atanh(tan(c/2 + (d*x)/2)))/d - (tan(c/2 + (d*x)/2)*(2*A + B + 2*C) - tan(c/2 + (d*x)/2)^3*(4*A + (4*C)/3) + tan(c/2 + (d*x)/2)^5*(2*A - B + 2*C))/(d*(3*tan(c/2 + (d*x)/2)^2 - 3*tan(c/2 + (d*x)/2)^4 + tan(c/2 + (d*x)/2)^6 - 1))`

3.57 $\int \sec(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.57.1	Optimal result	369
3.57.2	Mathematica [A] (verified)	369
3.57.3	Rubi [A] (verified)	370
3.57.4	Maple [A] (verified)	372
3.57.5	Fricas [A] (verification not implemented)	372
3.57.6	Sympy [F]	373
3.57.7	Maxima [A] (verification not implemented)	373
3.57.8	Giac [B] (verification not implemented)	373
3.57.9	Mupad [B] (verification not implemented)	374

3.57.1 Optimal result

Integrand size = 27, antiderivative size = 51

$$\int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(2A + C) \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{B \tan(c + dx)}{d} + \frac{C \sec(c + dx) \tan(c + dx)}{2d}$$

output `1/2*(2*A+C)*arctanh(sin(d*x+c))/d+B*tan(d*x+c)/d+1/2*C*sec(d*x+c)*tan(d*x+c)/d`

3.57.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.16

$$\int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{A \operatorname{ArcTanh}(\sin(c + dx))}{d} + \frac{C \operatorname{ArcTanh}(\sin(c + dx))}{2d}$$

$$+ \frac{B \tan(c + dx)}{d} + \frac{C \sec(c + dx) \tan(c + dx)}{2d}$$

input `Integrate[Sec[c + d*x]*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(A*ArcTanh[Sin[c + d*x]])/d + (C*ArcTanh[Sin[c + d*x]])/(2*d) + (B*Tan[c + d*x])/d + (C*Sec[c + d*x]*Tan[c + d*x])/(2*d)`

3.57.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {3042, 4535, 3042, 4254, 24, 4534, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c+dx) (A + B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right) \left(A + B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{4535} \\
 & \int \sec(c+dx) (C \sec^2(c+dx) + A) dx + B \int \sec^2(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right) \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + B \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{4254} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right) \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx - \frac{B \int 1d(-\tan(c+dx))}{d} \\
 & \quad \downarrow \text{24} \\
 & \int \csc\left(c+dx+\frac{\pi}{2}\right) \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A\right) dx + \frac{B \tan(c+dx)}{d} \\
 & \quad \downarrow \text{4534} \\
 & \frac{1}{2}(2A+C) \int \sec(c+dx) dx + \frac{B \tan(c+dx)}{d} + \frac{C \tan(c+dx) \sec(c+dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(2A+C) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{B \tan(c+dx)}{d} + \frac{C \tan(c+dx) \sec(c+dx)}{2d} \\
 & \quad \downarrow \text{4257} \\
 & \frac{(2A+C) \operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{B \tan(c+dx)}{d} + \frac{C \tan(c+dx) \sec(c+dx)}{2d}
 \end{aligned}$$

input `Int[Sec[c + d*x]*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `((2*A + C)*ArcTanh[Sin[c + d*x]]/(2*d) + (B*Tan[c + d*x])/d + (C*Sec[c + d*x]*Tan[c + d*x])/(2*d)`

3.57.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.57.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.24

method	result
derivativedivides	$\frac{A \ln(\sec(dx+c)+\tan(dx+c))+B \tan(dx+c)+C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$
default	$\frac{A \ln(\sec(dx+c)+\tan(dx+c))+B \tan(dx+c)+C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$
parts	$\frac{A \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{B \tan(dx+c)}{d} + \frac{C \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$
parallelrisch	$\frac{-\left(A+\frac{C}{2}\right)(1+\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+\left(A+\frac{C}{2}\right)(1+\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)+B \sin(2dx+2c)+C \sin(2dx+2c)}{d(1+\cos(2dx+2c))}$
norman	$\frac{\frac{(2B+C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d} - \frac{(2B-C) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{d}}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^2} - \frac{(2A+C) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{2d} + \frac{(2A+C) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{2d}$
risch	$-\frac{i(C e^{3i(dx+c)}-2B e^{2i(dx+c)}-C e^{i(dx+c)}-2B)}{d(e^{2i(dx+c)}+1)^2} - \frac{\ln(e^{i(dx+c)}-i)A}{d} - \frac{\ln(e^{i(dx+c)}-i)C}{2d} + \frac{\ln(e^{i(dx+c)}+i)A}{d} + \frac{\ln(e^{i(dx+c)}+i)C}{2d}$

input `int(sec(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(A*ln(sec(d*x+c)+tan(d*x+c))+B*tan(d*x+c)+C*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c))))`

3.57.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.61

$$\int \sec(c+dx) (A+B \sec(c+dx)+C \sec^2(c+dx)) dx$$

$$= \frac{(2A+C) \cos(dx+c)^2 \log(\sin(dx+c)+1) - (2A+C) \cos(dx+c)^2 \log(-\sin(dx+c)+1) + 2(2B \cos(dx+c)+C) \sin(dx+c)}{4d \cos(dx+c)^2}$$

input `integrate(sec(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/4*((2*A + C)*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (2*A + C)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(2*B*cos(d*x + c) + C)*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.57.6 Sympy [F]

$$\begin{aligned} & \int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \sec(c + dx) dx \end{aligned}$$

input `integrate(sec(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*sec(c + d*x), x)`

3.57.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.47

$$\begin{aligned} & \int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \\ & \frac{C \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx+c)+1) + \log(\sin(dx+c)-1) \right) - 4A \log(\sec(dx+c) + \tan(dx+c))}{4d} \end{aligned}$$

input `integrate(sec(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `-1/4*(C*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 4*A*log(sec(d*x + c) + tan(d*x + c)) - 4*B*tan(d*x + c))/d`

3.57.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(47) = 94.

Time = 0.31 (sec) , antiderivative size = 115, normalized size of antiderivative = 2.25

$$\begin{aligned} & \int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{(2A + C) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) - (2A + C) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|) - \frac{2(2B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - C \tan(\frac{1}{2} dx + \frac{1}{2} c))}{2d}}{2d} \end{aligned}$$

3.57. $\int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

input `integrate(sec(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/2*((2*A + C)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (2*A + C)*log(abs(tan(1/2*d*x + 1/2*c) - 1))) - 2*(2*B*tan(1/2*d*x + 1/2*c)^3 - C*tan(1/2*d*x + 1/2*c)^3 - 2*B*tan(1/2*d*x + 1/2*c) - C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^2/d`

3.57.9 Mupad [B] (verification not implemented)

Time = 15.74 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.75

$$\int \sec(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (2A + C)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (2B - C) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (2B + C)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)}$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/cos(c + d*x),x)`

output `(atanh(tan(c/2 + (d*x)/2))*(2*A + C))/d - (tan(c/2 + (d*x)/2)^3*(2*B - C) - tan(c/2 + (d*x)/2)*(2*B + C))/(d*(tan(c/2 + (d*x)/2)^4 - 2*tan(c/2 + (d*x)/2)^2 + 1))`

3.58 $\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.58.1	Optimal result	375
3.58.2	Mathematica [A] (verified)	375
3.58.3	Rubi [A] (verified)	376
3.58.4	Maple [A] (verified)	376
3.58.5	Fricas [B] (verification not implemented)	377
3.58.6	Sympy [F]	377
3.58.7	Maxima [A] (verification not implemented)	378
3.58.8	Giac [B] (verification not implemented)	378
3.58.9	Mupad [B] (verification not implemented)	379

3.58.1 Optimal result

Integrand size = 20, antiderivative size = 27

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = Ax + \frac{B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{C \tan(c + dx)}{d}$$

output `A*x+B*arctanh(sin(d*x+c))/d+C*tan(d*x+c)/d`

3.58.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = Ax + \frac{B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{C \tan(c + dx)}{d}$$

input `Integrate[A + B*Sec[c + d*x] + C*Sec[c + d*x]^2,x]`

output `A*x + (B*ArcTanh[Sin[c + d*x]])/d + (C*Tan[c + d*x])/d`

3.58.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

↓ 2009

$$Ax + \frac{B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{C \tan(c + dx)}{d}$$

input `Int[A + B*Sec[c + d*x] + C*Sec[c + d*x]^2,x]`

output `A*x + (B*ArcTanh[Sin[c + d*x]])/d + (C*Tan[c + d*x])/d`

3.58.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.58.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.30

method	result	size
default	$Ax + \frac{B \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{C \tan(dx+c)}{d}$	35
parts	$Ax + \frac{B \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{C \tan(dx+c)}{d}$	35
derivativedivides	$\frac{(dx+c)A+B \ln(\sec(dx+c)+\tan(dx+c))+C \tan(dx+c)}{d}$	37
risch	$Ax - \frac{B \ln(e^{i(dx+c)}-i)}{d} + \frac{B \ln(e^{i(dx+c)}+i)}{d} + \frac{2iC}{d(e^{2i(dx+c)}+1)}$	62
parallelrisch	$\frac{-B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right) \cos(dx+c)+B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right) \cos(dx+c)+C \sin(dx+c)}{d \cos(dx+c)} + Ax$	67
norman	$\frac{Ax \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2 - Ax - \frac{2C \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d}}{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2 - 1} + \frac{B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{d} - \frac{B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{d}$	87

input `int(A+B*sec(d*x+c)+C*sec(d*x+c)^2,x,method=_RETURNVERBOSE)`

output `A*x+B/d*ln(sec(d*x+c)+tan(d*x+c))+C*tan(d*x+c)/d`

3.58.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71 vs. $2(27) = 54$.

Time = 0.26 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.63

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2 A dx \cos(dx + c) + B \cos(dx + c) \log(\sin(dx + c) + 1) - B \cos(dx + c) \log(-\sin(dx + c) + 1) + 2 C \sin(dx + c)}{2 d \cos(dx + c)}$$

input `integrate(A+B*sec(d*x+c)+C*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/2*(2*A*d*x*cos(d*x + c) + B*cos(d*x + c)*log(sin(d*x + c) + 1) - B*cos(d*x + c)*log(-sin(d*x + c) + 1) + 2*C*sin(d*x + c))/(d*cos(d*x + c))`

3.58.6 Sympy [F]

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

input `integrate(A+B*sec(d*x+c)+C*sec(d*x+c)**2,x)`

output `Integral(A + B*sec(c + d*x) + C*sec(c + d*x)**2, x)`

3.58.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.26

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = Ax + \frac{B \log(\sec(dx + c) + \tan(dx + c))}{d} + \frac{C \tan(dx + c)}{d}$$

input `integrate(A+B*sec(d*x+c)+C*sec(d*x+c)^2,x, algorithm="maxima")`

output `A*x + B*log(sec(d*x + c) + tan(d*x + c))/d + C*tan(d*x + c)/d`

3.58.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. 2(27) = 54.

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.22

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = Ax + \frac{B \left(\log \left(\left| \frac{1}{\sin(dx+c)} + \sin(dx+c) + 2 \right| \right) - \log \left(\left| \frac{1}{\sin(dx+c)} + \sin(dx+c) - 2 \right| \right) \right)}{4d} + \frac{C \tan(dx+c)}{d}$$

input `integrate(A+B*sec(d*x+c)+C*sec(d*x+c)^2,x, algorithm="giac")`

output `A*x + 1/4*B*(log(abs(1/sin(d*x + c) + sin(d*x + c) + 2)) - log(abs(1/sin(d*x + c) + sin(d*x + c) - 2)))/d + C*tan(d*x + c)/d`

3.58.9 Mupad [B] (verification not implemented)

Time = 14.95 (sec) , antiderivative size = 161, normalized size of antiderivative = 5.96

$$\int (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2 A \operatorname{atan}\left(\frac{64 A^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{64 A^3 + 64 A B^2} + \frac{64 A B^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{64 A^3 + 64 A B^2}\right)}{d}$$

$$+ \frac{2 B \operatorname{atanh}\left(\frac{64 B^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{64 A^2 B + 64 B^3} + \frac{64 A^2 B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{64 A^2 B + 64 B^3}\right)}{d} - \frac{2 C \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1\right)}$$

input `int(A + B/cos(c + d*x) + C/cos(c + d*x)^2,x)`output `(2*A*atan((64*A^3*tan(c/2 + (d*x)/2))/(64*A*B^2 + 64*A^3) + (64*A*B^2*tan(c/2 + (d*x)/2))/(64*A*B^2 + 64*A^3))/d + (2*B*atanh((64*B^3*tan(c/2 + (d*x)/2))/(64*A^2*B + 64*B^3) + (64*A^2*B*tan(c/2 + (d*x)/2))/(64*A^2*B + 64*B^3))/d - (2*C*tan(c/2 + (d*x)/2))/(d*(tan(c/2 + (d*x)/2)^2 - 1))`

3.59 $\int \cos(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.59.1	Optimal result	380
3.59.2	Mathematica [A] (verified)	380
3.59.3	Rubi [A] (verified)	381
3.59.4	Maple [A] (verified)	382
3.59.5	Fricas [A] (verification not implemented)	383
3.59.6	Sympy [F]	383
3.59.7	Maxima [A] (verification not implemented)	384
3.59.8	Giac [B] (verification not implemented)	384
3.59.9	Mupad [B] (verification not implemented)	385

3.59.1 Optimal result

Integrand size = 27, antiderivative size = 27

$$\begin{aligned} & \int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= Bx + \frac{C \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{A \sin(c + dx)}{d} \end{aligned}$$

output `B*x+C*arctanh(sin(d*x+c))/d+A*sin(d*x+c)/d`

3.59.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.41

$$\begin{aligned} & \int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= Bx + \frac{C \operatorname{ArcTanh}(\sin(c + dx))}{d} + \frac{A \cos(dx) \sin(c)}{d} + \frac{A \cos(c) \sin(dx)}{d} \end{aligned}$$

input `Integrate[Cos[c + d*x]*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `B*x + (C*ArcTanh[Sin[c + d*x]])/d + (A*Cos[d*x]*Sin[c])/d + (A*Cos[c]*Sin[d*x])/d`

3.59.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$, Rules used = {3042, 4535, 24, 3042, 4533, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{4535} \\
 & \int \cos(c + dx) (C \sec^2(c + dx) + A) dx + B \int 1 dx \\
 & \quad \downarrow \text{24} \\
 & \int \cos(c + dx) (C \sec^2(c + dx) + A) dx + Bx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\csc(c + dx + \frac{\pi}{2})} dx + Bx \\
 & \quad \downarrow \text{4533} \\
 & C \int \sec(c + dx) dx + \frac{A \sin(c + dx)}{d} + Bx \\
 & \quad \downarrow \text{3042} \\
 & C \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{A \sin(c + dx)}{d} + Bx \\
 & \quad \downarrow \text{4257} \\
 & \frac{A \sin(c + dx)}{d} + \frac{C \operatorname{arctanh}(\sin(c + dx))}{d} + Bx
 \end{aligned}$$

input `Int[Cos[c + d*x]*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `B*x + (C*ArcTanh[Sin[c + d*x]])/d + (A*Sin[c + d*x])/d`

3.59.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)*(b_.)]^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`
- rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)*(b_.)]^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.59.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.37

method	result	size
derivativedivides	$\frac{A \sin(dx+c)+B(dx+c)+C \ln(\sec(dx+c)+\tan(dx+c))}{d}$	37
default	$\frac{A \sin(dx+c)+B(dx+c)+C \ln(\sec(dx+c)+\tan(dx+c))}{d}$	37
parallelrisch	$\frac{Bxd+A \sin(dx+c)-C \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+C \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{d}$	47
risch	$Bx - \frac{iAe^{i(dx+c)}}{2d} + \frac{iAe^{-i(dx+c)}}{2d} + \frac{\ln(e^{i(dx+c)}+i)C}{d} - \frac{\ln(e^{i(dx+c)}-i)C}{d}$	74
norman	$\frac{Bx \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^4 - Bx - \frac{2A \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d} + \frac{2A \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{d}}{\left(1+\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2\right)\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-1\right)} + \frac{C \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{d} - \frac{C \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{d}$	11

input `int(cos(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

3.59. $\int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

output `1/d*(A*sin(d*x+c)+B*(d*x+c)+C*ln(sec(d*x+c)+tan(d*x+c)))`

3.59.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.67

$$\int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2 B dx + C \log(\sin(dx + c) + 1) - C \log(-\sin(dx + c) + 1) + 2 A \sin(dx + c)}{2 d}$$

input `integrate(cos(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/2*(2*B*d*x + C*log(sin(d*x + c) + 1) - C*log(-sin(d*x + c) + 1) + 2*A*sin(d*x + c))/d`

3.59.6 Sympy [F]

$$\int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \cos(c + dx) dx$$

input `integrate(cos(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*cos(c + d*x), x)`

3.59.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.70

$$\int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2(dx + c)B + C(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 2A \sin(dx + c)}{2d}$$

input `integrate(cos(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*B + C*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 2*A*sin(d*x + c))/d`

3.59.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(27) = 54$.

Time = 0.31 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.59

$$\int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(dx + c)B + C \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - C \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) + \frac{2A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1}}{d}$$

input `integrate(cos(d*x+c)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `((d*x + c)*B + C*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - C*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + 2*A*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 + 1))/d`

3.59.9 Mupad [B] (verification not implemented)

Time = 14.92 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.52

$$\int \cos(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{2 B \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2 C \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{A \sin(c + dx)}{d}$$

input `int(cos(c + d*x)*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`output `(2*B*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*C*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (A*sin(c + d*x))/d`

3.60 $\int \cos^2(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.60.1	Optimal result	386
3.60.2	Mathematica [A] (verified)	386
3.60.3	Rubi [A] (verified)	387
3.60.4	Maple [A] (verified)	388
3.60.5	Fricas [A] (verification not implemented)	389
3.60.6	Sympy [F]	389
3.60.7	Maxima [A] (verification not implemented)	389
3.60.8	Giac [B] (verification not implemented)	390
3.60.9	Mupad [B] (verification not implemented)	390

3.60.1 Optimal result

Integrand size = 29, antiderivative size = 42

$$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{1}{2}(A + 2C)x + \frac{B \sin(c + dx)}{d} + \frac{A \cos(c + dx) \sin(c + dx)}{2d}$$

```
output 1/2*(A+2*C)*x+B*sin(d*x+c)/d+1/2*A*cos(d*x+c)*sin(d*x+c)/d
```

3.60.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.31

$$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= Cx + \frac{A(c + dx)}{2d} + \frac{B \cos(dx) \sin(c)}{d} + \frac{B \cos(c) \sin(dx)}{d} + \frac{A \sin(2(c + dx))}{4d}$$

```
input Integrate[Cos[c + d*x]^2*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]
```

```
output C*x + (A*(c + d*x))/(2*d) + (B*Cos[d*x]*Sin[c])/d + (B*Cos[c]*Sin[d*x])/d + (A*Sine[2*(c + d*x)])/(4*d)
```

3.60.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.207$, Rules used = {3042, 4535, 3042, 3117, 4533, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{4535} \\
 & \int \cos^2(c + dx) (C \sec^2(c + dx) + A) dx + B \int \cos(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\csc(c + dx + \frac{\pi}{2})^2} dx + B \int \sin(c + dx + \frac{\pi}{2}) dx \\
 & \quad \downarrow \text{3117} \\
 & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\csc(c + dx + \frac{\pi}{2})^2} dx + \frac{B \sin(c + dx)}{d} \\
 & \quad \downarrow \text{4533} \\
 & \frac{1}{2}(A + 2C) \int 1 dx + \frac{A \sin(c + dx) \cos(c + dx)}{2d} + \frac{B \sin(c + dx)}{d} \\
 & \quad \downarrow \text{24} \\
 & \frac{A \sin(c + dx) \cos(c + dx)}{2d} + \frac{1}{2}x(A + 2C) + \frac{B \sin(c + dx)}{d}
 \end{aligned}$$

input `Int[Cos[c + d*x]^2*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2), x]`

output `((A + 2*C)*x)/2 + (B*Sin[c + d*x])/d + (A*Cos[c + d*x]*Sin[c + d*x])/(2*d)`

3.60.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`
- rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.60.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.83

method	result
risch	$\frac{Ax}{2} + Cx + \frac{B \sin(dx+c)}{d} + \frac{A \sin(2dx+2c)}{4d}$
parallelrisch	$\frac{A \sin(2dx+2c)+4B \sin(dx+c)+2(A+2C)xd}{4d}$
derivativedivides	$\frac{A \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + B \sin(dx+c) + C(dx+c)}{d}$
default	$\frac{A \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + B \sin(dx+c) + C(dx+c)}{d}$
norman	$\frac{\left(-\frac{A}{2}-C\right)x + \left(-\frac{A}{2}-C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \left(\frac{A}{2}+C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + \left(\frac{A}{2}+C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + \frac{2A \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{d} - \frac{(A-2B)}{d}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$

```
input int(cos(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)
```

$$3.60. \int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

output $1/2*A*x+C*x+B*\sin(d*x+c)/d+1/4*A/d*\sin(2*d*x+2*c)$

3.60.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.79

$$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(A + 2C)dx + (A \cos(dx + c) + 2B) \sin(dx + c)}{2d}$$

input `integrate(cos(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output $1/2*((A + 2*C)*d*x + (A*\cos(d*x + c) + 2*B)*\sin(d*x + c))/d$

3.60.6 Sympy [F]

$$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \cos^2(c + dx) dx$$

input `integrate(cos(d*x+c)**2*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*cos(c + d*x)**2, x)`

3.60.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(2 dx + 2 c + \sin(2 dx + 2 c))A + 4 (dx + c)C + 4 B \sin(dx + c)}{4 d}$$

input `integrate(cos(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/4*((2*d*x + 2*c + sin(2*d*x + 2*c))*A + 4*(d*x + c)*C + 4*B*sin(d*x + c))/d`

3.60.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 86 vs. $2(38) = 76$.

Time = 0.29 (sec) , antiderivative size = 86, normalized size of antiderivative = 2.05

$$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{(dx + c)(A + 2C) - \frac{2(A \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 2B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - A \tan(\frac{1}{2} dx + \frac{1}{2} c) - 2B \tan(\frac{1}{2} dx + \frac{1}{2} c))}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^2}}{2d}$$

input `integrate(cos(d*x+c)^2*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/2*((d*x + c)*(A + 2*C) - 2*(A*tan(1/2*d*x + 1/2*c)^3 - 2*B*tan(1/2*d*x + 1/2*c)^3 - A*tan(1/2*d*x + 1/2*c) - 2*B*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^2)/d`

3.60.9 Mupad [B] (verification not implemented)

Time = 14.85 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.81

$$\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{Ax}{2} + Cx + \frac{A \sin(2c + 2dx)}{4d} + \frac{B \sin(c + dx)}{d}$$

input `int(cos(c + d*x)^2*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(A*x)/2 + C*x + (A*sin(2*c + 2*d*x))/(4*d) + (B*sin(c + d*x))/d`

3.60. $\int \cos^2(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.61 $\int \cos^3(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.61.1	Optimal result	391
3.61.2	Mathematica [A] (verified)	391
3.61.3	Rubi [A] (verified)	392
3.61.4	Maple [A] (verified)	394
3.61.5	Fricas [A] (verification not implemented)	394
3.61.6	Sympy [F]	395
3.61.7	Maxima [A] (verification not implemented)	395
3.61.8	Giac [B] (verification not implemented)	396
3.61.9	Mupad [B] (verification not implemented)	396

3.61.1 Optimal result

Integrand size = 29, antiderivative size = 56

$$\int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{Bx}{2} + \frac{(A + C) \sin(c + dx)}{d} + \frac{B \cos(c + dx) \sin(c + dx)}{2d} - \frac{A \sin^3(c + dx)}{3d}$$

```
output 1/2*B*x+(A+C)*sin(d*x+c)/d+1/2*B*cos(d*x+c)*sin(d*x+c)/d-1/3*A*sin(d*x+c)^3/d
```

3.61.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

$$\int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{6Bc + 6Bdx + 3(3A + 4C) \sin(c + dx) + 3B \sin(2(c + dx)) + A \sin(3(c + dx))}{12d}$$

```
input Integrate[Cos[c + d*x]^3*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]
```

```
output (6*B*c + 6*B*d*x + 3*(3*A + 4*C)*Sin[c + d*x] + 3*B*Sine[2*(c + d*x)] + A*Sin[3*(c + d*x)]/(12*d)
```


3.61.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 4535, 3042, 3115, 24, 4532, 3042, 3492, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c+dx) (A + B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \csc(c+dx + \frac{\pi}{2}) + C \csc(c+dx + \frac{\pi}{2})^2}{\csc(c+dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{4535} \\
 & \int \cos^3(c+dx) (C \sec^2(c+dx) + A) dx + B \int \cos^2(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^3} dx + B \int \sin(c+dx + \frac{\pi}{2})^2 dx \\
 & \quad \downarrow \text{3115} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^3} dx + B \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) \\
 & \quad \downarrow \text{24} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^3} dx + B \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \\
 & \quad \downarrow \text{4532} \\
 & \int \cos(c+dx) (A \cos^2(c+dx) + C) dx + B \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(c+dx + \frac{\pi}{2}) \left(A \sin(c+dx + \frac{\pi}{2})^2 + C \right) dx + B \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \\
 & \quad \downarrow \text{3492} \\
 & B \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) - \frac{\int (-A \sin^2(c+dx) + A + C) d(-\sin(c+dx))}{d}
 \end{aligned}$$

3.61. $\int \cos^3(c+dx) (A + B \sec(c+dx) + C \sec^2(c+dx)) dx$

$$B \left(\frac{\sin(c+dx)\cos(c+dx)}{2d} + \frac{x}{2} \right) - \frac{\overset{\downarrow 2009}{\frac{1}{3}A \sin^3(c+dx) - (A+C)\sin(c+dx)}}{d}$$

input `Int[Cos[c + d*x]^3*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `B*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (-((A + C)*Sin[c + d*x]) + (A*Sin[c + d*x]^3)/3)/d`

3.61.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3492 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[-f^(-1) Subst[Int[(1 - x^2)^((m - 1)/2)*(A + C - C*x^2)], x], x, Cos[e + f*x]], x] /; FreeQ[{e, f, A, C}, x] && IGtQ[(m + 1)/2, 0]`

rule 4532 `Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_)), x_Symbol] := Int[(C + A*Sin[e + f*x]^2)/Sin[e + f*x]^(m + 2), x] /; FreeQ[{e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && ILtQ[(m + 1)/2, 0]`

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.61.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.88

method	result
parallelrisch	$\frac{3B \sin(2dx+2c)+A \sin(3dx+3c)+(9A+12C) \sin(dx+c)+6Bxd}{12d}$
derivativedivides	$\frac{A(2+\cos(dx+c)^2) \sin(dx+c)}{3} + B \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + C \sin(dx+c)$
default	$\frac{A(2+\cos(dx+c)^2) \sin(dx+c)}{3} + B \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + C \sin(dx+c)$
risch	$\frac{Bx}{2} + \frac{3A \sin(dx+c)}{4d} + \frac{C \sin(dx+c)}{d} + \frac{A \sin(3dx+3c)}{12d} + \frac{B \sin(2dx+2c)}{4d}$
norman	$\frac{Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + \frac{(2A-B+2C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^7}{d} - \frac{Bx}{2} - Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^8}{2} - \frac{(2A-3B-6C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{3d} - \frac{(2A-3B-6C) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{3d}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1\right)}$

```
input int(cos(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)
```

```
output 1/12*(3*B*sin(2*d*x+2*c)+A*sin(3*d*x+3*c)+(9*A+12*C)*sin(d*x+c)+6*B*x*d)/d
```

3.61.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.80

$$\int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3 B dx + (2 A \cos(dx + c)^2 + 3 B \cos(dx + c) + 4 A + 6 C) \sin(dx + c)}{6 d}$$

```
input integrate(cos(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")
```

3.61. $\int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

output $1/6*(3*B*d*x + (2*A*\cos(d*x + c)^2 + 3*B*\cos(d*x + c) + 4*A + 6*C)*\sin(d*x + c))/d$

3.61.6 Sympy [F]

$$\begin{aligned} & \int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \cos^3(c + dx) dx \end{aligned}$$

input `integrate(cos(d*x+c)**3*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*cos(c + d*x)**3, x)`

3.61.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98

$$\begin{aligned} & \int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \\ & \frac{4 (\sin(dx + c)^3 - 3 \sin(dx + c))A - 3(2dx + 2c + \sin(2dx + 2c))B - 12C \sin(dx + c)}{12d} \end{aligned}$$

input `integrate(cos(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `-1/12*(4*(sin(d*x + c)^3 - 3*sin(d*x + c))*A - 3*(2*d*x + 2*c + sin(2*d*x + 2*c))*B - 12*C*sin(d*x + c))/d`

3.61.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 138 vs. $2(50) = 100$.

Time = 0.30 (sec) , antiderivative size = 138, normalized size of antiderivative = 2.46

$$\int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(dx + c)B + \frac{2(6A \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 3B \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 + 6C \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 + 4A \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 12C \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 6A \tan(\frac{1}{2} dx + \frac{1}{2} c))}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^3}}{6d}$$

input `integrate(cos(d*x+c)^3*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/6*(3*(d*x + c)*B + 2*(6*A*tan(1/2*d*x + 1/2*c)^5 - 3*B*tan(1/2*d*x + 1/2*c)^5 + 6*C*tan(1/2*d*x + 1/2*c)^5 + 4*A*tan(1/2*d*x + 1/2*c)^3 + 12*C*tan(1/2*d*x + 1/2*c)^3 + 6*A*tan(1/2*d*x + 1/2*c) + 3*B*tan(1/2*d*x + 1/2*c) + 6*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^3)/d`

3.61.9 Mupad [B] (verification not implemented)

Time = 14.71 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.18

$$\int \cos^3(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{Bx}{2} + \frac{2A \sin(c + dx)}{3d} + \frac{C \sin(c + dx)}{d}$$

$$+ \frac{B \cos(c + dx) \sin(c + dx)}{2d} + \frac{A \cos(c + dx)^2 \sin(c + dx)}{3d}$$

input `int(cos(c + d*x)^3*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(B*x)/2 + (2*A*sin(c + d*x))/(3*d) + (C*sin(c + d*x))/d + (B*cos(c + d*x)*sin(c + d*x))/(2*d) + (A*cos(c + d*x)^2*sin(c + d*x))/(3*d)`

3.62 $\int \cos^4(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.62.1	Optimal result	397
3.62.2	Mathematica [A] (verified)	397
3.62.3	Rubi [A] (verified)	398
3.62.4	Maple [A] (verified)	400
3.62.5	Fricas [A] (verification not implemented)	401
3.62.6	Sympy [F]	401
3.62.7	Maxima [A] (verification not implemented)	401
3.62.8	Giac [B] (verification not implemented)	402
3.62.9	Mupad [B] (verification not implemented)	402

3.62.1 Optimal result

Integrand size = 29, antiderivative size = 88

$$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{1}{8}(3A + 4C)x + \frac{B \sin(c + dx)}{d} + \frac{(3A + 4C) \cos(c + dx) \sin(c + dx)}{8d}$$

$$+ \frac{A \cos^3(c + dx) \sin(c + dx)}{4d} - \frac{B \sin^3(c + dx)}{3d}$$

output `1/8*(3*A+4*C)*x+B*sin(d*x+c)/d+1/8*(3*A+4*C)*cos(d*x+c)*sin(d*x+c)/d+1/4*A*cos(d*x+c)^3*sin(d*x+c)/d-1/3*B*sin(d*x+c)^3/d`

3.62.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.80

$$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{36Ac + 48cC + 36Adx + 48Cdx + 96B \sin(c + dx) - 32B \sin^3(c + dx) + 24(A + C) \sin(2(c + dx)) + 3A \sin^4(c + dx)}{96d}$$

input `Integrate[Cos[c + d*x]^4*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(36*A*c + 48*c*C + 36*A*d*x + 48*C*d*x + 96*B*Sin[c + d*x] - 32*B*Sin[c + d*x]^3 + 24*(A + C)*Sin[2*(c + d*x)] + 3*A*Sin[4*(c + d*x)])/(96*d)`

3.62. $\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.62.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.98, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 4535, 3042, 3113, 2009, 4533, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(c+dx) (A + B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \csc(c+dx + \frac{\pi}{2}) + C \csc(c+dx + \frac{\pi}{2})^2}{\csc(c+dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{4535} \\
 & \int \cos^4(c+dx) (C \sec^2(c+dx) + A) dx + B \int \cos^3(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^4} dx + B \int \sin(c+dx + \frac{\pi}{2})^3 dx \\
 & \quad \downarrow \text{3113} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^4} dx - \frac{B \int (1 - \sin^2(c+dx)) d(-\sin(c+dx))}{d} \\
 & \quad \downarrow \text{2009} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^4} dx - \frac{B(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} \\
 & \quad \downarrow \text{4533} \\
 & \frac{1}{4}(3A+4C) \int \cos^2(c+dx) dx + \frac{A \sin(c+dx) \cos^3(c+dx)}{4d} - \frac{B(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4}(3A+4C) \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{A \sin(c+dx) \cos^3(c+dx)}{4d} - \frac{B(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} \\
 & \quad \downarrow \text{3115}
 \end{aligned}$$

$$\frac{1}{4}(3A + 4C) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) + \frac{A \sin(c + dx) \cos^3(c + dx)}{4d} - \frac{B(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx))}{d}$$

↓ 24

$$\frac{1}{4}(3A + 4C) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) + \frac{A \sin(c + dx) \cos^3(c + dx)}{4d} - \frac{B(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx))}{d}$$

input `Int[Cos[c + d*x]^4*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(A*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + ((3*A + 4*C)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/4 - (B*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d`

3.62.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] :> Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.62.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.72

method	result
parallelrisch	$\frac{24(A+C) \sin(2dx+2c)+8B \sin(3dx+3c)+3A \sin(4dx+4c)+72B \sin(dx+c)+36d \left(A+\frac{4C}{3}\right)x}{96d}$
risch	$\frac{3Ax}{8} + \frac{Cx}{2} + \frac{3B \sin(dx+c)}{4d} + \frac{A \sin(4dx+4c)}{32d} + \frac{B \sin(3dx+3c)}{12d} + \frac{A \sin(2dx+2c)}{4d} + \frac{\sin(2dx+2c)C}{4d}$
derivativedivides	$A \left(\frac{\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{B(2+\cos(dx+c)^2) \sin(dx+c)}{3} + C \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
default	$A \left(\frac{\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{B(2+\cos(dx+c)^2) \sin(dx+c)}{3} + C \left(\frac{\sin(dx+c) \cos(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
norman	$\left(-\frac{3A}{8} - \frac{C}{2}\right)x + \left(-\frac{9A}{8} - \frac{3C}{2}\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \left(-\frac{3A}{4} - C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + \left(\frac{3A}{4} + C\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + \left(\frac{3A}{8} + \frac{C}{2}\right)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^8$

input `int(cos(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/96*(24*(A+C)*sin(2*d*x+2*c)+8*B*sin(3*d*x+3*c)+3*A*sin(4*d*x+4*c)+72*B*sin(dx+c)+36*d*(A+4/3*C)*x)/d`

3.62.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.74

$$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(3A + 4C)dx + (6A \cos(dx + c)^3 + 8B \cos(dx + c)^2 + 3(3A + 4C) \cos(dx + c) + 16B) \sin(dx + c)}{24d}$$

input `integrate(cos(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/24*(3*(3*A + 4*C)*d*x + (6*A*cos(d*x + c)^3 + 8*B*cos(d*x + c)^2 + 3*(3*A + 4*C)*cos(d*x + c) + 16*B)*sin(d*x + c))/d`

3.62.6 Sympy [F]

$$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \cos^4(c + dx) dx$$

input `integrate(cos(d*x+c)**4*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*cos(c + d*x)**4, x)`

3.62.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.88

$$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(12dx + 12c + \sin(4dx + 4c) + 8 \sin(2dx + 2c))A - 32(\sin(dx + c)^3 - 3 \sin(dx + c))B + 24(2dx + 2c)C}{96d}$$

input `integrate(cos(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/96*(3*(12*d*x + 12*c + sin(4*d*x + 4*c)) + 8*sin(2*d*x + 2*c))*A - 32*(sin(d*x + c)^3 - 3*sin(d*x + c))*B + 24*(2*d*x + 2*c + sin(2*d*x + 2*c))*C)/d`

3.62.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(80) = 160$.

Time = 0.29 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.27

$$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3(dx + c)(3A + 4C) - \frac{2}{9} \left(15A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^7 - 24B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^7 + 12C \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^7 - 9A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 40B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 12C \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 9A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 40B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 12C \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 15A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 24B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 12C \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 1} dx$$

input `integrate(cos(d*x+c)^4*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/24*(3*(d*x + c)*(3*A + 4*C) - 2*(15*A*tan(1/2*d*x + 1/2*c)^7 - 24*B*tan(1/2*d*x + 1/2*c)^7 + 12*C*tan(1/2*d*x + 1/2*c)^7 - 9*A*tan(1/2*d*x + 1/2*c)^5 - 40*B*tan(1/2*d*x + 1/2*c)^5 + 12*C*tan(1/2*d*x + 1/2*c)^5 + 9*A*tan(1/2*d*x + 1/2*c)^3 - 40*B*tan(1/2*d*x + 1/2*c)^3 - 12*C*tan(1/2*d*x + 1/2*c)^3 - 15*A*tan(1/2*d*x + 1/2*c) - 24*B*tan(1/2*d*x + 1/2*c) - 12*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^4/d`

3.62.9 Mupad [B] (verification not implemented)

Time = 15.06 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.92

$$\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3Ax}{8} + \frac{Cx}{2} + \frac{A \sin(2c + 2dx)}{4d} + \frac{A \sin(4c + 4dx)}{32d} + \frac{B \sin(3c + 3dx)}{12d} + \frac{C \sin(2c + 2dx)}{4d} + \frac{3B \sin(c + dx)}{4d}$$

3.62. $\int \cos^4(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

input `int(cos(c + d*x)^4*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(3*A*x)/8 + (C*x)/2 + (A*sin(2*c + 2*d*x))/(4*d) + (A*sin(4*c + 4*d*x))/(3
2*d) + (B*sin(3*c + 3*d*x))/(12*d) + (C*sin(2*c + 2*d*x))/(4*d) + (3*B*sin
(c + d*x))/(4*d)`

3.63 $\int \cos^5(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.63.1	Optimal result	404
3.63.2	Mathematica [A] (verified)	404
3.63.3	Rubi [A] (verified)	405
3.63.4	Maple [A] (verified)	408
3.63.5	Fricas [A] (verification not implemented)	408
3.63.6	Sympy [F]	409
3.63.7	Maxima [A] (verification not implemented)	409
3.63.8	Giac [B] (verification not implemented)	409
3.63.9	Mupad [B] (verification not implemented)	410

3.63.1 Optimal result

Integrand size = 29, antiderivative size = 98

$$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3Bx}{8} + \frac{(A + C) \sin(c + dx)}{d} + \frac{3B \cos(c + dx) \sin(c + dx)}{8d}$$

$$+ \frac{B \cos^3(c + dx) \sin(c + dx)}{4d} - \frac{(2A + C) \sin^3(c + dx)}{3d} + \frac{A \sin^5(c + dx)}{5d}$$

output `3/8*B*x+(A+C)*sin(d*x+c)/d+3/8*B*cos(d*x+c)*sin(d*x+c)/d+1/4*B*cos(d*x+c)^3*sin(d*x+c)/d-1/3*(2*A+C)*sin(d*x+c)^3/d+1/5*A*sin(d*x+c)^5/d`

3.63.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.89

$$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{180Bc + 180Bdx + 60(5A + 6C) \sin(c + dx) + 120B \sin(2(c + dx)) + 50A \sin(3(c + dx)) + 40C \sin(3(c + dx)) + 15B \sin(4(c + dx)) + 6A \sin(5(c + dx))}{480d}$$

input `Integrate[Cos[c + d*x]^5*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(180*B*c + 180*B*d*x + 60*(5*A + 6*C)*Sin[c + d*x] + 120*B*Sine[2*(c + d*x)] + 50*A*Sine[3*(c + d*x)] + 40*C*Sine[3*(c + d*x)] + 15*B*Sine[4*(c + d*x)] + 6*A*Sine[5*(c + d*x)])/(480*d)`

3.63.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 4535, 3042, 3115, 3042, 3115, 24, 4532, 3042, 3492, 290, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^5(c+dx) (A + B \sec(c+dx) + C \sec^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \csc(c+dx + \frac{\pi}{2}) + C \csc(c+dx + \frac{\pi}{2})^2}{\csc(c+dx + \frac{\pi}{2})^5} dx \\
 & \quad \downarrow \text{4535} \\
 & \int \cos^5(c+dx) (C \sec^2(c+dx) + A) dx + B \int \cos^4(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^5} dx + B \int \sin(c+dx + \frac{\pi}{2})^4 dx \\
 & \quad \downarrow \text{3115} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^5} dx + B \left(\frac{3}{4} \int \cos^2(c+dx) dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^5} dx + B \left(\frac{3}{4} \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) \\
 & \quad \downarrow \text{3115} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^5} dx + \\
 & \quad B \left(\frac{3}{4} \left(\int \frac{1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) \\
 & \quad \downarrow \text{24} \\
 & \int \frac{C \csc(c+dx + \frac{\pi}{2})^2 + A}{\csc(c+dx + \frac{\pi}{2})^5} dx + B \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 4532 \\
& \int \cos^3(c+dx) (A \cos^2(c+dx) + C) dx + \\
& B \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right) \\
& \downarrow 3042 \\
& \int \sin \left(c+dx + \frac{\pi}{2} \right)^3 \left(A \sin \left(c+dx + \frac{\pi}{2} \right)^2 + C \right) dx + \\
& B \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right) \\
& \downarrow 3492 \\
& \frac{B \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right) - \int (1 - \sin^2(c+dx)) (-A \sin^2(c+dx) + A + C) d(-\sin(c+dx))}{d} \\
& \downarrow 290 \\
& \frac{B \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right) - \int (A \sin^4(c+dx) - (2A + C) \sin^2(c+dx) + A \left(\frac{C}{A} + 1 \right)) d(-\sin(c+dx))}{d} \\
& \downarrow 2009 \\
& \frac{B \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right) - \frac{1}{3}(2A + C) \sin^3(c+dx) - (A + C) \sin(c+dx) - \frac{1}{5} A \sin^5(c+dx)}{d}
\end{aligned}$$

input `Int[Cos[c + d*x]^5*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `-((-((A + C)*Sin[c + d*x]) + ((2*A + C)*Sin[c + d*x]^3)/3 - (A*Sin[c + d*x]^5)/5)/d) + B*((Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (3*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d))))/4)`

3.63.3.1 Defintions of rubi rules used

- rule 294 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 290 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3492 `Int[sin[(e_) + (f_)*(x_)]^(m_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[-f^(-1) Subst[Int[(1 - x^2)^(m - 1)/2*(A + C - C*x^2), x], x, Cos[e + f*x]], x] /; FreeQ[{e, f, A, C}, x] && IGtQ[(m + 1)/2, 0]`
- rule 4532 `Int[csc[(e_) + (f_)*(x_)]^(m_)*(csc[(e_) + (f_)*(x_)]^2*(C_) + (A_)), x_Symbol] := Int[(C + A*Sin[e + f*x]^2)/Sin[e + f*x]^(m + 2), x] /; FreeQ[{e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && ILtQ[(m + 1)/2, 0]`
- rule 4535 `Int[(csc[(e_) + (f_)*(x_)]*(b_))^(m_)*((A_) + csc[(e_) + (f_)*(x_)]*(B_) + csc[(e_) + (f_)*(x_)]^2*(C_)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.63.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.81

method	result
parallelrisch	$\frac{(50A+40C) \sin(3dx+3c)+120B \sin(2dx+2c)+15B \sin(4dx+4c)+6A \sin(5dx+5c)+(300A+360C) \sin(dx+c)+180Bxd}{480d}$
derivativedivides	$\frac{A \left(\frac{8}{3} + \cos(dx+c)^4 + \frac{4 \cos(dx+c)^2}{3} \right) \sin(dx+c)}{5} + B \left(\frac{\left(\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{C(2+\cos(dx+c)^2) \sin(dx+c)}{3}$
default	$\frac{A \left(\frac{8}{3} + \cos(dx+c)^4 + \frac{4 \cos(dx+c)^2}{3} \right) \sin(dx+c)}{5} + B \left(\frac{\left(\cos(dx+c)^3 + \frac{3 \cos(dx+c)}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{C(2+\cos(dx+c)^2) \sin(dx+c)}{3}$
risch	$\frac{3Bx}{8} + \frac{5A \sin(dx+c)}{8d} + \frac{3C \sin(dx+c)}{4d} + \frac{A \sin(5dx+5c)}{80d} + \frac{B \sin(4dx+4c)}{32d} + \frac{5A \sin(3dx+3c)}{48d} + \frac{\sin(3dx+3c)C}{12d}$
norman	$\frac{-\frac{3Bx}{8} - \frac{3Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{2} - \frac{15Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{8} + \frac{15Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^8}{8} + \frac{3Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^{10}}{2} + \frac{3Bx \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^{12}}{8} - (8A-9B+40C) \sin(dx+c)}{120d}$

input `int(cos(d*x+c)^5*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/480*((50*A+40*C)*sin(3*d*x+3*c)+120*B*sin(2*d*x+2*c)+15*B*sin(4*d*x+4*c)+6*A*sin(5*d*x+5*c)+(300*A+360*C)*sin(d*x+c)+180*B*x*d)/d`

3.63.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.74

$$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{45 B dx + (24 A \cos(dx + c)^4 + 30 B \cos(dx + c)^3 + 8(4 A + 5 C) \cos(dx + c)^2 + 45 B \cos(dx + c) + 64 A + 80 C) \sin(dx + c)}{120 d}$$

input `integrate(cos(d*x+c)^5*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/120*(45*B*d*x + (24*A*cos(d*x + c)^4 + 30*B*cos(d*x + c)^3 + 8*(4*A + 5*C)*cos(d*x + c)^2 + 45*B*cos(d*x + c) + 64*A + 80*C)*sin(d*x + c))/d`

3.63.6 Sympy [F]

$$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (A + B \sec(c + dx) + C \sec^2(c + dx)) \cos^5(c + dx) dx$$

input `integrate(cos(d*x+c)**5*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)*cos(c + d*x)**5, x)`

3.63.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.91

$$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{32 (3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c))A + 15 (12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(2 dx + 2 c))B - 160 (\sin(dx + c)^3 - 3 \sin(dx + c))C}{480 d}$$

input `integrate(cos(d*x+c)^5*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `1/480*(32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*A + 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B - 160*(sin(d*x + c)^3 - 3*sin(d*x + c))*C)/d`

3.63.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 222 vs. 2(88) = 176.

Time = 0.31 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.27

$$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$45 (dx + c)B + \frac{2 (120 A \tan(\frac{1}{2} dx + \frac{1}{2} c)^9 - 75 B \tan(\frac{1}{2} dx + \frac{1}{2} c)^9 + 120 C \tan(\frac{1}{2} dx + \frac{1}{2} c)^9 + 160 A \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 - 30 B \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 + 30 C \tan(\frac{1}{2} dx + \frac{1}{2} c)^7)}{480 d}$$

3.63. $\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

input `integrate(cos(d*x+c)^5*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/120*(45*(d*x + c)*B + 2*(120*A*tan(1/2*d*x + 1/2*c)^9 - 75*B*tan(1/2*d*x + 1/2*c)^9 + 120*C*tan(1/2*d*x + 1/2*c)^9 + 160*A*tan(1/2*d*x + 1/2*c)^7 - 30*B*tan(1/2*d*x + 1/2*c)^7 + 320*C*tan(1/2*d*x + 1/2*c)^7 + 464*A*tan(1/2*d*x + 1/2*c)^5 + 400*C*tan(1/2*d*x + 1/2*c)^5 + 160*A*tan(1/2*d*x + 1/2*c)^3 + 30*B*tan(1/2*d*x + 1/2*c)^3 + 320*C*tan(1/2*d*x + 1/2*c)^3 + 120*A*tan(1/2*d*x + 1/2*c) + 75*B*tan(1/2*d*x + 1/2*c) + 120*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^5/d`

3.63.9 Mupad [B] (verification not implemented)

Time = 14.92 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.06

$$\int \cos^5(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{3Bx}{8} + \frac{5A \sin(3c + 3dx)}{48d} + \frac{A \sin(5c + 5dx)}{80d} + \frac{B \sin(2c + 2dx)}{4d}$$

$$+ \frac{B \sin(4c + 4dx)}{32d} + \frac{C \sin(3c + 3dx)}{12d} + \frac{5A \sin(c + dx)}{8d} + \frac{3C \sin(c + dx)}{4d}$$

input `int(cos(c + d*x)^5*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(3*B*x)/8 + (5*A*sin(3*c + 3*d*x))/(48*d) + (A*sin(5*c + 5*d*x))/(80*d) + (B*sin(2*c + 2*d*x))/(4*d) + (B*sin(4*c + 4*d*x))/(32*d) + (C*sin(3*c + 3*d*x))/(12*d) + (5*A*sin(c + d*x))/(8*d) + (3*C*sin(c + d*x))/(4*d)`

3.64 $\int \cos^6(c+dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.64.1	Optimal result	411
3.64.2	Mathematica [A] (verified)	411
3.64.3	Rubi [A] (verified)	412
3.64.4	Maple [A] (verified)	415
3.64.5	Fricas [A] (verification not implemented)	415
3.64.6	Sympy [F(-1)]	416
3.64.7	Maxima [A] (verification not implemented)	416
3.64.8	Giac [B] (verification not implemented)	416
3.64.9	Mupad [B] (verification not implemented)	417

3.64.1 Optimal result

Integrand size = 29, antiderivative size = 132

$$\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{1}{16}(5A + 6C)x + \frac{B \sin(c + dx)}{d} + \frac{(5A + 6C) \cos(c + dx) \sin(c + dx)}{16d}$$

$$+ \frac{(5A + 6C) \cos^3(c + dx) \sin(c + dx)}{24d}$$

$$+ \frac{A \cos^5(c + dx) \sin(c + dx)}{6d} - \frac{2B \sin^3(c + dx)}{3d} + \frac{B \sin^5(c + dx)}{5d}$$

```
output 1/16*(5*A+6*C)*x+B*sin(d*x+c)/d+1/16*(5*A+6*C)*cos(d*x+c)*sin(d*x+c)/d+1/2
4*(5*A+6*C)*cos(d*x+c)^3*sin(d*x+c)/d+1/6*A*cos(d*x+c)^5*sin(d*x+c)/d-2/3*
B*sin(d*x+c)^3/d+1/5*B*sin(d*x+c)^5/d
```

3.64.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.77

$$\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{960B \sin(c + dx) - 640B \sin^3(c + dx) + 192B \sin^5(c + dx) + 5(60Ac + 72cC + 60Adx + 72Cdx + (45A$$

960d

input `Integrate[Cos[c + d*x]^6*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output $(960*B*\sin[c + d*x] - 640*B*\sin[c + d*x]^3 + 192*B*\sin[c + d*x]^5 + 5*(60*A*c + 72*c*C + 60*A*d*x + 72*C*d*x + (45*A + 48*C)*\sin[2*(c + d*x)] + (9*A + 6*C)*\sin[4*(c + d*x)] + A*\sin[6*(c + d*x)])/(960*d)$

3.64.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.94, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 4535, 3042, 3113, 2009, 4533, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{\csc(c + dx + \frac{\pi}{2})^6} dx \\ & \quad \downarrow \text{4535} \\ & \int \cos^6(c + dx) (C \sec^2(c + dx) + A) dx + B \int \cos^5(c + dx) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\csc(c + dx + \frac{\pi}{2})^6} dx + B \int \sin(c + dx + \frac{\pi}{2})^5 dx \\ & \quad \downarrow \text{3113} \\ & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\csc(c + dx + \frac{\pi}{2})^6} dx - \frac{B \int (\sin^4(c + dx) - 2 \sin^2(c + dx) + 1) d(-\sin(c + dx))}{d} \\ & \quad \downarrow \text{2009} \\ & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\csc(c + dx + \frac{\pi}{2})^6} dx - \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d} \\ & \quad \downarrow \text{4533} \end{aligned}$$

3.64. $\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

$$\begin{aligned}
& \frac{1}{6}(5A + 6C) \int \cos^4(c + dx) dx + \frac{A \sin(c + dx) \cos^5(c + dx)}{6d} - \\
& \quad \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{6}(5A + 6C) \int \sin\left(c + dx + \frac{\pi}{2}\right)^4 dx + \frac{A \sin(c + dx) \cos^5(c + dx)}{6d} - \\
& \quad \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d} \\
& \quad \downarrow \text{3115} \\
& \frac{1}{6}(5A + 6C) \left(\frac{3}{4} \int \cos^2(c + dx) dx + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) + \frac{A \sin(c + dx) \cos^5(c + dx)}{6d} - \\
& \quad \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{6}(5A + 6C) \left(\frac{3}{4} \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 dx + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) + \\
& \quad \frac{A \sin(c + dx) \cos^5(c + dx)}{6d} - \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d} \\
& \quad \downarrow \text{3115} \\
& \frac{1}{6}(5A + 6C) \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) + \\
& \quad \frac{A \sin(c + dx) \cos^5(c + dx)}{6d} - \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d} \\
& \quad \downarrow \text{24} \\
& \frac{1}{6}(5A + 6C) \left(\frac{\sin(c + dx) \cos^3(c + dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) \right) + \\
& \quad \frac{A \sin(c + dx) \cos^5(c + dx)}{6d} - \frac{B(-\frac{1}{5} \sin^5(c + dx) + \frac{2}{3} \sin^3(c + dx) - \sin(c + dx))}{d}
\end{aligned}$$

input `Int[Cos[c + d*x]^6*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(A*Cos[c + d*x]^5*Sin[c + d*x])/(6*d) - (B*(-Sin[c + d*x] + (2*Sin[c + d*x]^3)/3 - Sin[c + d*x]^5/5))/d + ((5*A + 6*C)*((Cos[c + d*x]^3*Sin[c + d*x])/4d) + (3*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)))/4)/6`

3.64.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`
- rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)])*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.64.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.72

method	result
parallelrisch	$(225A+240C) \sin(2dx+2c)+(45A+30C) \sin(4dx+4c)+100B \sin(3dx+3c)+12B \sin(5dx+5c)+5A \sin(6dx+6c)+600B \sin(dx+c)$
derivativedivides	$A \left(\frac{\left(\cos(dx+c)^5 + \frac{5 \cos(dx+c)^3}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{B \left(\frac{8}{3} + \cos(dx+c)^4 + \frac{4 \cos(dx+c)^2}{3} \right) \sin(dx+c)}{5} + C \left(\frac{\cos(dx+c)}{d} \right)$
default	$A \left(\frac{\left(\cos(dx+c)^5 + \frac{5 \cos(dx+c)^3}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{B \left(\frac{8}{3} + \cos(dx+c)^4 + \frac{4 \cos(dx+c)^2}{3} \right) \sin(dx+c)}{5} + C \left(\frac{\cos(dx+c)}{d} \right)$
risch	$\frac{5Ax}{16} + \frac{3Cx}{8} + \frac{5B \sin(dx+c)}{8d} + \frac{A \sin(6dx+6c)}{192d} + \frac{B \sin(5dx+5c)}{80d} + \frac{3A \sin(4dx+4c)}{64d} + \frac{\sin(4dx+4c)C}{32d} + \frac{5B \sin(dx+c)}{d}$
norman	$\left(-\frac{5A}{16} - \frac{3C}{8} \right) x + \left(-\frac{45A}{16} - \frac{27C}{8} \right) x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + \left(-\frac{25A}{16} - \frac{15C}{8} \right) x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \left(-\frac{25A}{16} - \frac{15C}{8} \right) x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + \left(\frac{5A}{16} + \frac{3C}{8} \right) x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^8$

input `int(cos(d*x+c)^6*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/960*((225*A+240*C)*sin(2*d*x+2*c)+(45*A+30*C)*sin(4*d*x+4*c)+100*B*sin(3*d*x+3*c)+12*B*sin(5*d*x+5*c)+5*A*sin(6*d*x+6*c)+600*B*sin(d*x+c)+300*d*x*(A+6/5*C))/d`

3.64.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.70

$$\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{15(5A + 6C)dx + (40A \cos(dx + c)^5 + 48B \cos(dx + c)^4 + 10(5A + 6C) \cos(dx + c)^3 + 64B \cos(dx + c)^2 + 15(5A + 6C) \cos(dx + c) + 128B) \sin(dx + c)}{240d}$$

input `integrate(cos(d*x+c)^6*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fracas")`

output `1/240*(15*(5*A + 6*C)*d*x + (40*A*cos(d*x + c)^5 + 48*B*cos(d*x + c)^4 + 10*(5*A + 6*C)*cos(d*x + c)^3 + 64*B*cos(d*x + c)^2 + 15*(5*A + 6*C)*cos(d*x + c) + 128*B)*sin(d*x + c))/d`

3.64. $\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.64.6 Sympy [F(-1)]

Timed out.

$$\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**6*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`output `Timed out`**3.64.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.87

$$\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{5(4 \sin(2dx + 2c)^3 - 60dx - 60c - 9 \sin(4dx + 4c) - 48 \sin(2dx + 2c))A - 64(3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c))B - 30(12dx + 12c + \sin(4dx + 4c) + 8 \sin(2dx + 2c))C}{960d}$$

input `integrate(cos(d*x+c)^6*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`output `-1/960*(5*(4*sin(2*d*x + 2*c)^3 - 60*d*x - 60*c - 9*sin(4*d*x + 4*c) - 48*sin(2*d*x + 2*c))*A - 64*(3*sin(dx + c)^5 - 10*sin(dx + c)^3 + 15*sin(dx + c))*B - 30*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*C)/d`**3.64.8 Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 284 vs. 2(120) = 240.

Time = 0.30 (sec) , antiderivative size = 284, normalized size of antiderivative = 2.15

$$\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{15(dx + c)(5A + 6C) - 2(165A \tan(\frac{1}{2}dx + \frac{1}{2}c)^{11} - 240B \tan(\frac{1}{2}dx + \frac{1}{2}c)^{11} + 150C \tan(\frac{1}{2}dx + \frac{1}{2}c)^{11} - 25A \tan(\frac{1}{2}dx + \frac{1}{2}c)^9 - 560B \tan(\frac{1}{2}dx + \frac{1}{2}c)^9 + 150C \tan(\frac{1}{2}dx + \frac{1}{2}c)^9)}{150d}$$

3.64. $\int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

input `integrate(cos(d*x+c)^6*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="giac")`

output `1/240*(15*(d*x + c)*(5*A + 6*C) - 2*(165*A*tan(1/2*d*x + 1/2*c)^11 - 240*B*tan(1/2*d*x + 1/2*c)^11 + 150*C*tan(1/2*d*x + 1/2*c)^11 - 25*A*tan(1/2*d*x + 1/2*c)^9 - 560*B*tan(1/2*d*x + 1/2*c)^9 + 210*C*tan(1/2*d*x + 1/2*c)^9 + 450*A*tan(1/2*d*x + 1/2*c)^7 - 1248*B*tan(1/2*d*x + 1/2*c)^7 + 60*C*tan(1/2*d*x + 1/2*c)^7 - 450*A*tan(1/2*d*x + 1/2*c)^5 - 1248*B*tan(1/2*d*x + 1/2*c)^5 - 60*C*tan(1/2*d*x + 1/2*c)^5 + 25*A*tan(1/2*d*x + 1/2*c)^3 - 560*B*tan(1/2*d*x + 1/2*c)^3 - 210*C*tan(1/2*d*x + 1/2*c)^3 - 165*A*tan(1/2*d*x + 1/2*c) - 240*B*tan(1/2*d*x + 1/2*c) - 150*C*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^6)/d`

3.64.9 Mupad [B] (verification not implemented)

Time = 15.01 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.95

$$\begin{aligned} & \int \cos^6(c + dx) (A + B \sec(c + dx) + C \sec^2(c + dx)) dx \\ &= \frac{5Ax}{16} + \frac{3Cx}{8} + \frac{15A \sin(2c + 2dx)}{64d} + \frac{3A \sin(4c + 4dx)}{64d} \\ &+ \frac{A \sin(6c + 6dx)}{192d} + \frac{5B \sin(3c + 3dx)}{48d} + \frac{B \sin(5c + 5dx)}{80d} \\ &+ \frac{C \sin(2c + 2dx)}{4d} + \frac{C \sin(4c + 4dx)}{32d} + \frac{5B \sin(c + dx)}{8d} \end{aligned}$$

input `int(cos(c + d*x)^6*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `(5*A*x)/16 + (3*C*x)/8 + (15*A*sin(2*c + 2*d*x))/(64*d) + (3*A*sin(4*c + 4*d*x))/(64*d) + (A*sin(6*c + 6*d*x))/(192*d) + (5*B*sin(3*c + 3*d*x))/(48*d) + (B*sin(5*c + 5*d*x))/(80*d) + (C*sin(2*c + 2*d*x))/(4*d) + (C*sin(4*c + 4*d*x))/(32*d) + (5*B*sin(c + d*x))/(8*d)`

3.65 $\int (b \sec(c+dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)$

3.65.1	Optimal result	418
3.65.2	Mathematica [C] (verified)	419
3.65.3	Rubi [A] (verified)	419
3.65.4	Maple [C] (verified)	423
3.65.5	Fricas [C] (verification not implemented)	424
3.65.6	Sympy [F]	425
3.65.7	Maxima [F]	425
3.65.8	Giac [F]	426
3.65.9	Mupad [F(-1)]	426

3.65.1 Optimal result

Integrand size = 33, antiderivative size = 178

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx =$$

$$\frac{2b^2(5A + 3C)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2bB\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

$$+ \frac{2b(5A + 3C)\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2B(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d} + \frac{2C(b \sec(c + dx))^{3/2} \tan(c + dx)}{5d}$$

output

```
2/3*B*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/d-2/5*b^2*(5*A+3*C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/5*b*(5*A+3*C)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+2/3*b*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d+2/5*C*(b*sec(d*x+c))^(3/2)*tan(d*x+c)/d
```

3.65.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.71 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.73

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{2e^{-ic}(-1 + e^{2ic}) \cos^3(c + dx) \csc(c) \left(5B - 15Ae^{i(c+dx)} - 3Ce^{i(c+dx)} - 30Ae^{3i(c+dx)} - \dots\right)}{\dots}$$

input `Integrate[(b*Sec[c + d*x])^(3/2)*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(2*(-1 + E^((2*I)*c))*Cos[c + d*x]^3*Csc[c]*(5*B - 15*A*E^(I*(c + d*x)) - 3*C*E^(I*(c + d*x)) - 30*A*E^((3*I)*(c + d*x)) - 24*C*E^((3*I)*(c + d*x)) - 5*B*E^((4*I)*(c + d*x)) - 15*A*E^((5*I)*(c + d*x)) - 9*C*E^((5*I)*(c + d*x)) - (5*I)*B*(1 + E^((2*I)*(c + d*x)))^2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (5*A + 3*C)*E^(I*(c + d*x))*(1 + E^((2*I)*(c + d*x)))^(5/2))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])*(b*Sec[c + d*x])^(3/2)*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2))/(15*d*E^(I*c)*(1 + E^((2*I)*(c + d*x)))^2*(A + 2*C + 2*B*cos[c + d*x] + A*cos[2*(c + d*x)]))`

3.65.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4535, 3042, 4255, 3042, 4258, 3042, 3120, 4534, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(b \csc\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \csc\left(c + dx + \frac{\pi}{2}\right) + C \csc\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{4535}$$

3.65. $\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

$$\begin{aligned}
& \int (b \sec(c + dx))^{3/2} (C \sec^2(c + dx) + A) dx + \frac{B \int (b \sec(c + dx))^{5/2} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \int \left(b \csc \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(C \csc \left(c + dx + \frac{\pi}{2} \right)^2 + A \right) dx + \frac{B \int (b \csc (c + dx + \frac{\pi}{2}))^{5/2} dx}{b} \\
& \quad \downarrow \text{4255} \\
& \frac{\int \left(b \csc \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(C \csc \left(c + dx + \frac{\pi}{2} \right)^2 + A \right) dx + B \left(\frac{1}{3} b^2 \int \sqrt{b \sec(c + dx)} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \right)}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \left(b \csc \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(C \csc \left(c + dx + \frac{\pi}{2} \right)^2 + A \right) dx + B \left(\frac{1}{3} b^2 \int \sqrt{b \csc (c + dx + \frac{\pi}{2})} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \right)}{b} \\
& \quad \downarrow \text{4258} \\
& \frac{\int \left(b \csc \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(C \csc \left(c + dx + \frac{\pi}{2} \right)^2 + A \right) dx + B \left(\frac{1}{3} b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \right)}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \left(b \csc \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(C \csc \left(c + dx + \frac{\pi}{2} \right)^2 + A \right) dx + B \left(\frac{1}{3} b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \right)}{b} \\
& \quad \downarrow \text{3120} \\
& \frac{\int \left(b \csc \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(C \csc \left(c + dx + \frac{\pi}{2} \right)^2 + A \right) dx + B \left(\frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF} \left(\frac{1}{2} (c + dx), 2 \right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \right)}{b} \\
& \quad \downarrow \text{4534}
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{5}(5A + 3C) \int (b \sec(c + dx))^{3/2} dx + \\
& \frac{B \left(\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right)}{2C \tan(c + dx)(b \sec(c + dx))^{3/2}} + \\
& \frac{b}{5d} \downarrow \text{3042} \\
& \frac{1}{5}(5A + 3C) \int \left(b \csc \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} dx + \\
& \frac{B \left(\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right)}{2C \tan(c + dx)(b \sec(c + dx))^{3/2}} + \\
& \frac{b}{5d} \downarrow \text{4255} \\
& \frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \right) + \\
& \frac{B \left(\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right)}{2C \tan(c + dx)(b \sec(c + dx))^{3/2}} + \\
& \frac{b}{5d} \downarrow \text{3042} \\
& \frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \frac{B \left(\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right)}{2C \tan(c + dx)(b \sec(c + dx))^{3/2}} + \\
& \frac{b}{5d} \downarrow \text{4258} \\
& \frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) + \\
& \frac{B \left(\frac{2b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3d} + \frac{2b \sin(c+dx)(b \sec(c+dx))^{3/2}}{3d} \right)}{2C \tan(c + dx)(b \sec(c + dx))^{3/2}} + \\
& \frac{b}{5d} \downarrow \text{3042}
\end{aligned}$$

$$\frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{b^2 \int \sqrt{\sin(c + dx + \frac{\pi}{2}) dx}}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) +$$

$$\frac{B \left(\frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \right)}{b} +$$

$$\frac{2C \tan(c + dx) (b \sec(c + dx))^{3/2}}{5d}$$

↓ 3119

$$\frac{1}{5}(5A + 3C) \left(\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c + dx) | 2)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \right) +$$

$$\frac{B \left(\frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d} \right)}{b} +$$

$$\frac{2C \tan(c + dx) (b \sec(c + dx))^{3/2}}{5d}$$

input `Int[(b*Sec[c + d*x])^(3/2)*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `((5*A + 3*C)*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d)/5 + (B*((2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (2*b*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(3*d)))/b + (2*C*(b*Sec[c + d*x])^(3/2)*Tan[c + d*x])/(5*d)`

3.65.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 4255 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
  EqQ[n^2, 1/4]
```

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.)
  + (A_)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
  FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.65.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.95 (sec) , antiderivative size = 949, normalized size of antiderivative = 5.33

method	result	size
parts	Expression too large to display	949
default	Expression too large to display	998

```
input int((b*sec(d*x+c))^(3/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNV
ERBOSE)
```


output

```

-2*A/d*(I*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)^2-I*EllipticE(I*(cot(d*x+c)-c
sc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*c
os(d*x+c)^2+2*I*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(
1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-2*I*EllipticE(I*(cot(d*
x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(
1/2)*cos(d*x+c)+I*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*(1/(cos(d*x+c)+1)
)^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(
d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+c)),I)-sin(d*
x+c))*(b*sec(d*x+c))^(1/2)*b/(cos(d*x+c)+1)+2/3*B/d*(b*sec(d*x+c))^(1/2)*b
*(-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos
(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d
*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+tan(d*
x+c))+2/5*C/d*(b*sec(d*x+c))^(1/2)*b/(cos(d*x+c)+1)*(3*I*(1/(cos(d*x+c)+1)
)^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-csc(d*x+
c)),I)*cos(d*x+c)^2-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1
))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)^2+6*I*(1/(cos(d
*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(cot(d*x+c)-
csc(d*x+c)),I)*cos(d*x+c)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*
x+c)+1))^(1/2)*EllipticF(I*(cot(d*x+c)-csc(d*x+c)),I)*cos(d*x+c)+3*I*(1...

```

3.65.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.26

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \frac{-5i \sqrt{2} B b^{3/2} \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5 \dots}{\dots}$$

input

```

integrate((b*sec(d*x+c))^(3/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm
m="fricas")

```

output `1/15*(-5*I*sqrt(2)*B*b^(3/2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*B*b^(3/2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*(5*A + 3*C)*b^(3/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*(5*A + 3*C)*b^(3/2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*(5*A + 3*C)*b*cos(d*x + c)^2 + 5*B*b*cos(d*x + c) + 3*C*b)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^2)`

3.65.6 Sympy [F]

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

input `integrate((b*sec(d*x+c))**(3/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral((b*sec(c + d*x))**(3/2)*(A + B*sec(c + d*x) + C*sec(c + d*x)**2), x)`

3.65.7 Maxima [F]

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + B \sec(dx + c) + A)(b \sec(dx + c))^{3/2} dx$$

input `integrate((b*sec(d*x+c))^(3/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)*(b*sec(d*x + c))^(3/2), x)`

3.65.8 Giac [F]

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \int (C \sec(dx + c)^2 + B \sec(dx + c) + A)(b \sec(dx + c))^{\frac{3}{2}} dx$$

input `integrate((b*sec(d*x+c))^(3/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm m="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)*(b*sec(d*x + c))^(3/2), x)`

3.65.9 Mupad [F(-1)]

Timed out.

$$\int (b \sec(c + dx))^{3/2} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx = \int \left(\frac{b}{\cos(c + dx)} \right)^{3/2} \left(A + \frac{B}{\cos(c + dx)} + \frac{C}{\cos(c + dx)^2} \right) dx$$

input `int((b/cos(c + d*x))^(3/2)*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `int((b/cos(c + d*x))^(3/2)*(A + B/cos(c + d*x) + C/cos(c + d*x)^2), x)`

3.66 $\int \sqrt{b \sec(c + dx)}(A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

3.66.1	Optimal result	427
3.66.2	Mathematica [C] (verified)	428
3.66.3	Rubi [A] (verified)	428
3.66.4	Maple [C] (verified)	432
3.66.5	Fricas [C] (verification not implemented)	432
3.66.6	Sympy [F]	433
3.66.7	Maxima [F]	433
3.66.8	Giac [F]	434
3.66.9	Mupad [F(-1)]	434

3.66.1 Optimal result

Integrand size = 33, antiderivative size = 136

$$\int \sqrt{b \sec(c + dx)}(A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= -\frac{2bBE\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2(3A + C)\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2B\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} + \frac{2C\sqrt{b \sec(c + dx)} \tan(c + dx)}{3d}$$

```
output -2*b*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*B*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+2/3*(3*A+C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d+2/3*C*(b*sec(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.66.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.20 (sec) , antiderivative size = 302, normalized size of antiderivative = 2.22

$$\int \sqrt{b \sec(c+dx)}(A + B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$= \frac{4\sqrt{b \sec(c+dx)}(A + B \sec(c+dx) + C \sec^2(c+dx)) \left(-\frac{i\sqrt{2}e^{-i(c+dx)}\sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}}(3B(1+e^{2i(c+dx)})+3B(-1+e^{2ic}))}{3d(A + B \sec(c+dx) + C \sec^2(c+dx))} \right)}{3d(A + B \sec(c+dx) + C \sec^2(c+dx))}$$

input `Integrate[Sqrt[b*Sec[c + d*x]]*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(4*Sqrt[b*Sec[c + d*x]]*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)*((-I)*Sqrt[2]*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*(3*B*(1 + E^((2*I)*(c + d*x))) + 3*B*(-1 + E^((2*I)*c))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[-1/4, 1/2, 3/4, -E^((2*I)*(c + d*x))]] + (3*A + C)*E^(I*(c + d*x))*(-1 + E^((2*I)*c))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/4, 1/2, 5/4, -E^((2*I)*(c + d*x))]))/(E^(I*(c + d*x))*(-1 + E^((2*I)*c))) + Sqrt[Sec[c + d*x]]*(3*B*Cos[d*x]*Csc[c] + C*Tan[c + d*x]))/(3*d*(A + 2*C + 2*B*Cos[c + d*x] + A*Cos[2*(c + d*x)])*Sec[c + d*x]^(5/2))`

3.66.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 4535, 3042, 4255, 3042, 4258, 3042, 3119, 4534, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{b \sec(c+dx)}(A + B \sec(c+dx) + C \sec^2(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)}\left(A + B \csc\left(c+dx+\frac{\pi}{2}\right) + C \csc\left(c+dx+\frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{4535}$$

$$\begin{aligned}
& \int \sqrt{b \sec(c+dx)} (C \sec^2(c+dx) + A) dx + \frac{B \int (b \sec(c+dx))^{3/2} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)} \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A \right) dx + \frac{B \int (b \csc\left(c+dx+\frac{\pi}{2}\right))^{3/2} dx}{b} \\
& \quad \downarrow \text{4255} \\
& \frac{\int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)} \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A \right) dx +}{b} \\
& \quad \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \right)}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)} \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A \right) dx +}{b} \\
& \quad \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - b^2 \int \frac{1}{\sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)}} dx \right)}{b} \\
& \quad \downarrow \text{4258} \\
& \frac{\int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)} \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A \right) dx +}{b} \\
& \quad \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)} \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A \right) dx +}{b} \\
& \quad \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{b^2 \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow \text{3119} \\
& \frac{\int \sqrt{b \csc\left(c+dx+\frac{\pi}{2}\right)} \left(C \csc\left(c+dx+\frac{\pi}{2}\right)^2 + A \right) dx +}{b} \\
& \quad \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c+dx)|2\right)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow \text{4534}
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{3}(3A + C) \int \sqrt{b \sec(c + dx)} dx + \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} + \\
& \qquad \qquad \qquad \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{1}{3}(3A + C) \int \sqrt{b \csc \left(c + dx + \frac{\pi}{2} \right)} dx + \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} + \\
& \qquad \qquad \qquad \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
& \qquad \qquad \qquad \downarrow \text{4258} \\
& \frac{1}{3}(3A + C) \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{1}{3}(3A + C) \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)}} dx + \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d} \\
& \qquad \qquad \qquad \downarrow \text{3120} \\
& \frac{2(3A + C) \sqrt{\cos(c + dx)} \operatorname{EllipticF} \left(\frac{1}{2}(c + dx), 2 \right) \sqrt{b \sec(c + dx)}}{3d} + \\
& \frac{B \left(\frac{2b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d} - \frac{2b^2 E(\frac{1}{2}(c+dx)|2)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \right)}{b} + \frac{2C \tan(c + dx) \sqrt{b \sec(c + dx)}}{3d}
\end{aligned}$$

input `Int[Sqrt[b*Sec[c + d*x]]*(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2),x]`

output `(2*(3*A + C)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*d) + (B*((-2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d))/b + (2*C*Sqrt[b*Sec[c + d*x]]*Tan[c + d*x])/(3*d)`

3.66.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.66.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.53 (sec) , antiderivative size = 611, normalized size of antiderivative = 4.49

method	result
parts	$-\frac{2iA(\cos(dx+c)+1)\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),i)\sqrt{b\sec(dx+c)}}{d} + \frac{2B(i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\text{EllipticE}(i(-\cot(dx+c)+\csc(dx+c)),i)\cos(dx+c)-2i(1/\cos(dx+c)+1)^{1/2}(\cos(dx+c)/(\cos(dx+c)+1))^{1/2}\text{EllipticF}(i(-\cot(dx+c)+\csc(dx+c)),I)-\tan(dx+c))}{d}$
default	Expression too large to display

```
input int((b*sec(d*x+c))^(1/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x,method=_RETURNV
ERBOSE)
```

```
output -2*I*A/d*(cos(d*x+c)+1)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(b*sec(d*x+c))^(1/2)+2*B/d*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)^2-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)^2+2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)+sin(d*x+c)*(b*sec(d*x+c))^(1/2)/(cos(d*x+c)+1)-2/3*C/d*(b*sec(d*x+c))^(1/2)*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)-tan(d*x+c))
```

3.66.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.42

$$\int \sqrt{b \sec(c + dx)} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \frac{\sqrt{2}(-3iA - iC)\sqrt{b} \cos(dx + c) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \sqrt{2}(3iA + iC)\sqrt{b} \cos(dx + c) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\cos(dx + c)}$$

3.66. $\int \sqrt{b \sec(c + dx)} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$

input `integrate((b*sec(d*x+c))^(1/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="fricas")`

output `1/3*(sqrt(2)*(-3*I*A - I*C)*sqrt(b)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(3*I*A + I*C)*sqrt(b)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*B*sqrt(b)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*B*sqrt(b)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*B*cos(d*x + c) + C)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c))`

3.66.6 Sympy [F]

$$\int \sqrt{b \sec(c + dx)} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int \sqrt{b \sec(c + dx)} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

input `integrate((b*sec(d*x+c))**(1/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)**2),x)`

output `Integral(sqrt(b*sec(c + d*x))*(A + B*sec(c + d*x) + C*sec(c + d*x)**2), x)`

3.66.7 Maxima [F]

$$\int \sqrt{b \sec(c + dx)} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (C \sec(dx + c)^2 + B \sec(dx + c) + A) \sqrt{b \sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)*sqrt(b*sec(d*x + c)), x)`

3.66.8 Giac [F]

$$\int \sqrt{b \sec(c + dx)} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int (C \sec(dx + c)^2 + B \sec(dx + c) + A) \sqrt{b \sec(dx + c)} dx$$

input `integrate((b*sec(d*x+c))^(1/2)*(A+B*sec(d*x+c)+C*sec(d*x+c)^2),x, algorithm m="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)*sqrt(b*sec(d*x + c)), x)`

3.66.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \sec(c + dx)} (A + B \sec(c + dx) + C \sec^2(c + dx)) dx$$

$$= \int \sqrt{\frac{b}{\cos(c + dx)}} \left(A + \frac{B}{\cos(c + dx)} + \frac{C}{\cos(c + dx)^2} \right) dx$$

input `int((b/cos(c + d*x))^(1/2)*(A + B/cos(c + d*x) + C/cos(c + d*x)^2),x)`

output `int((b/cos(c + d*x))^(1/2)*(A + B/cos(c + d*x) + C/cos(c + d*x)^2), x)`

3.67 $\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$

3.67.1 Optimal result 435
 3.67.2 Mathematica [C] (verified) 436
 3.67.3 Rubi [A] (verified) 436
 3.67.4 Maple [C] (verified) 439
 3.67.5 Fricas [C] (verification not implemented) 440
 3.67.6 Sympy [F] 440
 3.67.7 Maxima [F] 441
 3.67.8 Giac [F] 441
 3.67.9 Mupad [F(-1)] 441

3.67.1 Optimal result

Integrand size = 33, antiderivative size = 110

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{2(A - C)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2B\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{b \sec(c + dx)}}{bd} + \frac{2C \tan(c + dx)}{d\sqrt{b \sec(c + dx)}}$$

```
output 2*(A-C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*
d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2*B*(cos(1/2*d
*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)
)*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b/d+2*C*tan(d*x+c)/d/(b*sec(d*x+c)
)^(1/2)
```

3.67.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.39 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.42

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{2e^{-idx} \sqrt{b \sec(c + dx)} (-i \cos(dx) + \sin(dx)) \left(-3A \cos(c + dx) + 3C \cos(c + dx) + 3iB \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left[\frac{c + dx}{2}, 2\right] + (A - C) E^{I(c + dx)} \sqrt{1 + E^{(2I)(c + dx)}} \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{(2I)(c + dx)}\right] + (3I)C \sin[c + dx] \right)}{(3bdE^{I dx})}$$

input `Integrate[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/Sqrt[b*Sec[c + d*x]],x]`

output `(2*Sqrt[b*Sec[c + d*x]]*((-I)*Cos[d*x] + Sin[d*x])*(-3*A*Cos[c + d*x] + 3*C*Cos[c + d*x] + (3*I)*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (A - C)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + (3*I)*C*Sin[c + d*x]))/(3*b*d*E^(I*d*x))`

3.67.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4535, 3042, 4258, 3042, 3120, 4534, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \csc\left(c + dx + \frac{\pi}{2}\right) + C \csc\left(c + dx + \frac{\pi}{2}\right)^2}{\sqrt{b \csc\left(c + dx + \frac{\pi}{2}\right)}} dx$$

$$\downarrow \text{4535}$$

$$\int \frac{C \sec^2(c + dx) + A}{\sqrt{b \sec(c + dx)}} dx + \frac{B \int \sqrt{b \sec(c + dx)} dx}{b}$$

$$\downarrow \text{3042}$$

3.67. $\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$

$$\begin{aligned}
& \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx + \frac{B \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx}{b} \\
& \quad \downarrow 4258 \\
& \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx + \frac{B \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{b} \\
& \quad \downarrow 3042 \\
& \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx + \frac{B \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{b} \\
& \quad \downarrow 3120 \\
& \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx + \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{bd} \\
& \quad \downarrow 4534 \\
& (A - C) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx + \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{bd} + \\
& \quad \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow 3042 \\
& (A - C) \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx + \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{bd} + \\
& \quad \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow 4258 \\
& \frac{(A - C) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{bd} + \\
& \quad \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow 3042 \\
& \frac{(A - C) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{bd} + \\
& \quad \frac{2C \tan(c + dx)}{d \sqrt{b \sec(c + dx)}}
\end{aligned}$$

3.67. $\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$

$$\begin{aligned} & \downarrow 3119 \\ & \frac{2(A-C)E\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{2B\sqrt{\cos(c+dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx),2\right)\sqrt{b\sec(c+dx)}}{bd} + \\ & \frac{2C\tan(c+dx)}{d\sqrt{b\sec(c+dx)}} \end{aligned}$$

input `Int[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/Sqrt[b*Sec[c + d*x]],x]`

output `(2*(A - C)*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b*d) + (2*C*Tan[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])`

3.67.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m*(csc[(e_.) + (f_.)*(x_)]^2*(C_. + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.67.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.73 (sec) , antiderivative size = 847, normalized size of antiderivative = 7.70

method	result	size
parts	Expression too large to display	847
default	Expression too large to display	945

```
input int((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output 2*A/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(I*EllipticE(I*(-cot(d*x+c)+csc(
d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(
d*x+c)-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+2*I*(1/(cos(d*x+c)+1))^(1/2)*
(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-
2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I
*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d
*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-I*(1/(c
os(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*
x+c)+csc(d*x+c)),I)*sec(d*x+c)+sin(d*x+c))-2*I*B/d*(1/(cos(d*x+c)+1))^(1/2
)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)/(b*sec(d*x+c))^(1/2)/(cos(d*x+c)
/(cos(d*x+c)+1))^(1/2)-2*C/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(I*Ellipt
icE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(co
s(d*x+c)+1))^(1/2)*cos(d*x+c)-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1
/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+2*I*(1
/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot
(d*x+c)+csc(d*x+c)),I)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c
)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(
1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)
),I)*sec(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(...
```


3.67.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.39

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

$$= \frac{-i \sqrt{2} B \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} B \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + \sqrt{2} (I A - I C) \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + \sqrt{2} (-I A + I C) \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 C \sqrt{b / \cos(dx + c)} \sin(dx + c)}{b d}$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + sqrt(2)*(I*A - I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + sqrt(2)*(-I*A + I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*C*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b*d)`

3.67.6 Sympy [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(1/2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)/sqrt(b*sec(c + d*x)), x)`

3.67.7 Maxima [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm m="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/sqrt(b*sec(d*x + c)), x)`

3.67.8 Giac [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{\sqrt{b \sec(dx + c)}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(1/2),x, algorithm m="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/sqrt(b*sec(d*x + c)), x)`

3.67.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx = \int \frac{A + \frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(1/2),x)`

output `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(1/2), x)`

3.68
$$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

3.68.1	Optimal result	442
3.68.2	Mathematica [C] (verified)	442
3.68.3	Rubi [A] (verified)	443
3.68.4	Maple [C] (verified)	445
3.68.5	Fricas [C] (verification not implemented)	446
3.68.6	Sympy [F]	447
3.68.7	Maxima [F]	447
3.68.8	Giac [F]	447
3.68.9	Mupad [F(-1)]	448

3.68.1 Optimal result

Integrand size = 33, antiderivative size = 117

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{2BE(\frac{1}{2}(c + dx) | 2)}{bd\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2(A + 3C)\sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{3b^2d} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/3*(A+3*C)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d+2/3*A*tan(d*x+c)/d/(b*sec(d*x+c))^(3/2)
```

3.68.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.72 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.22

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{2\left(6iB \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1}{2}, \frac{3}{4}, -e^{2i(c+dx)}\right) - 2i(A + 3C)\right)}{3b}$$

input `Integrate[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(3/2),x
]`

output `(2*((6*I)*B*Hypergeometric2F1[-1/4, 1/2, 3/4, -E^((2*I)*(c + d*x))] - (2*I)
)*(A + 3*C)*E^(I*(c + d*x))*Hypergeometric2F1[1/4, 1/2, 5/4, -E^((2*I)*(c
+ d*x))] + Sqrt[1 + E^((2*I)*(c + d*x))]*((-3*I)*B + A*Sin[c + d*x]))/(3*
b*d*Sqrt[1 + E^((2*I)*(c + d*x))]*Sqrt[b*Sec[c + d*x]])`

3.68.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4535, 3042, 4258, 3042, 3119, 4533, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx \\ & \quad \downarrow \text{4535} \\ & \int \frac{C \sec^2(c + dx) + A}{(b \sec(c + dx))^{3/2}} dx + \frac{B \int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{b} \\ & \quad \downarrow \text{3042} \\ & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{B \int \frac{1}{\sqrt{b \csc(c + dx + \frac{\pi}{2})}} dx}{b} \\ & \quad \downarrow \text{4258} \\ & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{B \int \sqrt{\cos(c + dx)} dx}{b \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\begin{aligned}
& \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{B \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{b \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow \text{3119} \\
& \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{2BE(\frac{1}{2}(c + dx) | 2)}{bd \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow \text{4533} \\
& \frac{(A + 3C) \int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} + \frac{2BE(\frac{1}{2}(c + dx) | 2)}{bd \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(A + 3C) \int \sqrt{b \csc(c + dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} + \frac{2BE(\frac{1}{2}(c + dx) | 2)}{bd \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow \text{4258} \\
& \frac{(A + 3C) \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} + \\
& \quad \frac{2BE(\frac{1}{2}(c + dx) | 2)}{bd \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(A + 3C) \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} + \\
& \quad \frac{2BE(\frac{1}{2}(c + dx) | 2)}{bd \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
& \quad \downarrow \text{3120} \\
& \frac{2(A + 3C) \sqrt{\cos(c + dx)} \text{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{3b^2 d} + \frac{2A \tan(c + dx)}{3d(b \sec(c + dx))^{3/2}} + \\
& \quad \frac{2BE(\frac{1}{2}(c + dx) | 2)}{bd \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}
\end{aligned}$$

input `Int[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(3/2), x]`

output `(2*B*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*(A + 3*C)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*A*Tan[c + d*x])/(3*d*(b*Sec[c + d*x])^(3/2))`

3.68. $\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$

3.68.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^m*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.68.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.59 (sec) , antiderivative size = 608, normalized size of antiderivative = 5.20

method	result
parts	$-\frac{2A\left(i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)+i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\csc(dx+c)),i\right)\right)}{3d\sqrt{b\sec(dx+c)}b}$
default	Expression too large to display

3.68.
$$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

input `int((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x,method=_RETURNV
ERBOSE)`

output `-2/3*A/d/(b*sec(d*x+c))^(1/2)/b*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-sin(d*x+c))+2*B/b/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+sin(d*x+c))-2*I*C/d*(1/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)/b/(b*sec(d*x+c))^(1/2)/(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)`

3.68.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.37

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \frac{2A \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c) \sin(dx+c) + \sqrt{2}(-iA - 3iC) \sqrt{b}}{\dots}$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm
m="fricas")`

output `1/3*(2*A*sqrt(b/cos(d*x + c))*cos(d*x + c)*sin(d*x + c) + sqrt(2)*(-I*A - 3*I*C)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(I*A + 3*I*C)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b^2*d)`

3.68.6 Sympy [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(3/2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)/(b*sec(c + d*x))**(3/2), x)`

3.68.7 Maxima [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm m="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/(b*sec(d*x + c))^(3/2), x)`

3.68.8 Giac [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(3/2),x, algorithm m="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/(b*sec(d*x + c))^(3/2), x)`

3.68.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx = \int \frac{A + \frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(3/2), x)`output `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(3/2), x)`

3.69
$$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

3.69.1	Optimal result	449
3.69.2	Mathematica [C] (verified)	449
3.69.3	Rubi [A] (verified)	450
3.69.4	Maple [C] (verified)	453
3.69.5	Fricas [C] (verification not implemented)	454
3.69.6	Sympy [F]	455
3.69.7	Maxima [F]	455
3.69.8	Giac [F]	455
3.69.9	Mupad [F(-1)]	456

3.69.1 Optimal result

Integrand size = 33, antiderivative size = 150

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{2(3A + 5C)E(\frac{1}{2}(c + dx)|2)}{5b^2d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2B\sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3b^3d\sqrt{b \sec(c + dx)}} + \frac{2B \sin(c + dx)}{3b^2d\sqrt{b \sec(c + dx)}} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}}$$

output $2/3*B*\sin(d*x+c)/b^2/d/(b*\sec(d*x+c))^{(1/2)}+2/5*(3*A+5*C)*(\cos(1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+2/3*B*(\cos(1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b^3/d+2/5*A*\tan(d*x+c)/d/(b*\sec(d*x+c))^{(5/2)}$

3.69.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.20 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.13

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{e^{-idx} \sqrt{b \sec(c + dx)} (\cos(dx) + i \sin(dx)) \left(10B \sqrt{\cos(c + dx)} E \right)}{\dots}$$

input `Integrate[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(5/2), x]`

output `(Sqrt[b*Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*(10*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (2*I)*(3*A + 5*C)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((6*I)*(3*A + 5*C) + 10*B*Sin[c + d*x] + 3*A*Sin[2*(c + d*x)])))/(15*b^3*d*E^(I*d*x))`

3.69.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 4535, 3042, 4256, 3042, 4258, 3042, 3120, 4533, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \csc(c + dx + \frac{\pi}{2}) + C \csc(c + dx + \frac{\pi}{2})^2}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4535} \\
 & \int \frac{C \sec^2(c + dx) + A}{(b \sec(c + dx))^{5/2}} dx + \frac{B \int \frac{1}{(b \sec(c + dx))^{3/2}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx + \frac{B \int \frac{1}{(b \csc(c + dx + \frac{\pi}{2}))^{3/2}} dx}{b} \\
 & \quad \downarrow \text{4256} \\
 & \int \frac{C \csc(c + dx + \frac{\pi}{2})^2 + A}{(b \csc(c + dx + \frac{\pi}{2}))^{5/2}} dx + \frac{B \left(\frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} + \frac{2 \sin(c + dx)}{3bd \sqrt{b \sec(c + dx)}} \right)}{b} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx + \frac{B \left(\frac{\int \sqrt{b \csc(c+dx+\frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow 4258 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx + \frac{B \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow 3042 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx + \frac{B \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow 3120 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx + \frac{B \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow 4533 \\
& \frac{(3A+5C) \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2A \tan(c+dx)}{5d(b \sec(c+dx))^{5/2}} + \\
& \frac{B \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow 3042 \\
& \frac{(3A+5C) \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2A \tan(c+dx)}{5d(b \sec(c+dx))^{5/2}} + \\
& \frac{B \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow 4258 \\
& \frac{(3A+5C) \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2A \tan(c+dx)}{5d(b \sec(c+dx))^{5/2}} + \\
& \frac{B \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}} \right)}{b} \\
& \quad \downarrow 3042
\end{aligned}$$

3.69. $\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

$$\frac{(3A + 5C) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}} +$$

$$B \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)$$

b

↓ 3119

$$\frac{2(3A + 5C)E(\frac{1}{2}(c + dx) | 2)}{5b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2A \tan(c + dx)}{5d(b \sec(c + dx))^{5/2}} +$$

$$B \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)$$

b

input `Int[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(5/2), x]`

output `(2*(3*A + 5*C)*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (B*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])))/b + (2*A*Tan[c + d*x])/(5*d*(b*Sec[c + d*x])^(5/2))`

3.69.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n)*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

```
rule 4533 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.)
+ (A_)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] +
Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; Fr
eeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*
(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Cs
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2)
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.69.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.69 (sec) , antiderivative size = 956, normalized size of antiderivative = 6.37

method	result	size
parts	Expression too large to display	956
default	Expression too large to display	1004

```
input int((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 2/5*A/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^2*(3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+cos(d*x+c)^2*sin(d*x+c)+sin(d*x+c)*cos(d*x+c)+3*sin(d*x+c))-2/3*B/d/(b*sec(d*x+c))^(1/2)/b^2*(I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-sin(d*x+c))+2*C/b^2/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)*(I*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)-I*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*cos(d*x+c)+2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)-2*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(...
```

3.69.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.17

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \frac{-5i \sqrt{2} B \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{(b \sec(c + dx))^{5/2}}$$

```
input integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x, algorithm="fracas")
```

```
output 1/15*(-5*I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*B*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*sqrt(2)*(-3*I*A - 5*I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(3*I*A + 5*I*C)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*A*cos(d*x + c)^2 + 5*B*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c))/(b^3*d)
```

3.69. $\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$

3.69.6 Sympy [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(5/2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)/(b*sec(c + d*x))**(5/2), x)`

3.69.7 Maxima [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x, algorithm m="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/(b*sec(d*x + c))^(5/2), x)`

3.69.8 Giac [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(5/2),x, algorithm m="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/(b*sec(d*x + c))^(5/2), x)`

3.69.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx = \int \frac{A + \frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(5/2), x)`output `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(5/2), x)`

3.70
$$\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$$

3.70.1 Optimal result 457
 3.70.2 Mathematica [C] (verified) 458
 3.70.3 Rubi [A] (verified) 458
 3.70.4 Maple [C] (verified) 462
 3.70.5 Fracas [C] (verification not implemented) 463
 3.70.6 Sympy [F] 463
 3.70.7 Maxima [F] 464
 3.70.8 Giac [F] 464
 3.70.9 Mupad [F(-1)] 464

3.70.1 Optimal result

Integrand size = 33, antiderivative size = 185

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{6BE(\frac{1}{2}(c + dx) | 2)}{5b^3d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2(5A + 7C)\sqrt{\cos(c + dx)} \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2) \sqrt{b \sec(c + dx)}}{21b^4d} + \frac{2B \sin(c + dx)}{5b^2d(b \sec(c + dx))^{3/2}} + \frac{2(5A + 7C) \sin(c + dx)}{21b^3d\sqrt{b \sec(c + dx)}} + \frac{2A \tan(c + dx)}{7d(b \sec(c + dx))^{7/2}}$$

```
output 2/5*B*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(3/2)+2/21*(5*A+7*C)*sin(d*x+c)/b^3/d/(b*sec(d*x+c))^(1/2)+6/5*B*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^3/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+2/21*(5*A+7*C)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^4/d+2/7*A*tan(d*x+c)/d/(b*sec(d*x+c))^(7/2)
```

3.70.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.60 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.96

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \frac{504iB \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1}{2}, \frac{3}{4}, -e^{2i(c+dx)}\right) - 40i(5A + 7C)}{(b \sec(c + dx))^{7/2}}$$

input `Integrate[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(7/2), x]`

output `((504*I)*B*Hypergeometric2F1[-1/4, 1/2, 3/4, -E^((2*I)*(c + d*x))] - (40*I)*(5*A + 7*C)*E^(I*(c + d*x))*Hypergeometric2F1[1/4, 1/2, 5/4, -E^((2*I)*(c + d*x))] + Sqrt[1 + E^((2*I)*(c + d*x))]*(5*(23*A + 28*C)*Sin[c + d*x] + 3*((-84*I)*B + 14*B*Sin[2*(c + d*x)] + 5*A*Sin[3*(c + d*x)])))/(210*b^3*d*Sqrt[1 + E^((2*I)*(c + d*x))]*Sqrt[b*Sec[c + d*x]])`

3.70.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4535, 3042, 4256, 3042, 4258, 3042, 3119, 4533, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \csc\left(c + dx + \frac{\pi}{2}\right) + C \csc\left(c + dx + \frac{\pi}{2}\right)^2}{(b \csc\left(c + dx + \frac{\pi}{2}\right))^{7/2}} dx \\ & \quad \downarrow \text{4535} \\ & \int \frac{C \sec^2(c + dx) + A}{(b \sec(c + dx))^{7/2}} dx + \frac{B \int \frac{1}{(b \sec(c + dx))^{5/2}} dx}{b} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.70. $\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx$

$$\begin{aligned}
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx + \frac{B \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{5/2}} dx}{b} \\
& \quad \downarrow 4256 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx + \frac{B \left(\frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
& \quad \downarrow 3042 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx + \frac{B \left(\frac{3 \int \frac{1}{\sqrt{b \csc(c+dx+\frac{\pi}{2})}} dx}{5b^2} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
& \quad \downarrow 4258 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx + \frac{B \left(\frac{3 \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
& \quad \downarrow 3042 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx + \frac{B \left(\frac{3 \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{5b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
& \quad \downarrow 3119 \\
& \int \frac{C \csc(c+dx+\frac{\pi}{2})^2 + A}{(b \csc(c+dx+\frac{\pi}{2}))^{7/2}} dx + \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
& \quad \downarrow 4533 \\
& \frac{(5A+7C) \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}} + \\
& \quad \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
& \quad \downarrow 3042 \\
& \frac{(5A+7C) \int \frac{1}{(b \csc(c+dx+\frac{\pi}{2}))^{3/2}} dx}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}} + \\
& \quad \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b}
\end{aligned}$$

3.70. $\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$

$$\begin{aligned}
 & \downarrow 4256 \\
 & \frac{(5A + 7C) \left(\frac{\int \sqrt{b \sec(c+dx)} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}} + \\
 & \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
 & \downarrow 3042 \\
 & \frac{(5A + 7C) \left(\frac{\int \sqrt{b \csc(c+dx + \frac{\pi}{2})} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}} + \\
 & \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
 & \downarrow 4258 \\
 & \frac{(5A + 7C) \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}} + \\
 & \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
 & \downarrow 3042 \\
 & \frac{(5A + 7C) \left(\frac{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{3b^2} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}} + \\
 & \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b} \\
 & \downarrow 3120 \\
 & \frac{(5A + 7C) \left(\frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2) \sqrt{b \sec(c+dx)}}{3b^2 d} + \frac{2 \sin(c+dx)}{3bd \sqrt{b \sec(c+dx)}} \right)}{7b^2} + \frac{2A \tan(c+dx)}{7d(b \sec(c+dx))^{7/2}} + \\
 & \frac{B \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \sec(c+dx))^{3/2}} \right)}{b}
 \end{aligned}$$

input `Int[(A + B*Sec[c + d*x] + C*Sec[c + d*x]^2)/(b*Sec[c + d*x])^(7/2), x]`

3.70. $\int \frac{A+B \sec(c+dx)+C \sec^2(c+dx)}{(b \sec(c+dx))^{7/2}} dx$

output $(B*((6*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*\text{Sin}[c + d*x])/(5*b*d*(b*\text{Sec}[c + d*x])^{(3/2)})))/b + ((5*A + 7*C)*((2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*b^2*d) + (2*\text{Sin}[c + d*x])/(3*b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])))/(7*b^2) + (2*A*\text{Tan}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(7/2)})$

3.70.3.1 Defintions of rubi rules used

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3119 $\text{Int}[\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

rule 4256 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n + 1)}/(b*d^n)), x] + \text{Simp}[(n + 1)/(b^2*n) \text{Int}[(b*\text{Csc}[c + d*x])^{(n + 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

rule 4533 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.))^{(m_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]^{2*(C_.) + (A_.)}), x_Symbol] \rightarrow \text{Simp}[A*\text{Cot}[e + f*x]*((b*\text{Csc}[e + f*x])^m/(f*m)), x] + \text{Simp}[(C*m + A*(m + 1))/(b^2*m) \text{Int}[(b*\text{Csc}[e + f*x])^{(m + 2)}, x], x] /; \text{FreeQ}\{b, e, f, A, C\}, x \ \&\& \ \text{NeQ}[C*m + A*(m + 1), 0] \ \&\& \ \text{LeQ}[m, -1]$

rule 4535 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.))^{(m_.)}*((A_.) + \text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + \text{csc}[(e_.) + (f_.)*(x_)]^{2*(C_.)}), x_Symbol] \rightarrow \text{Simp}[B/b \text{Int}[(b*\text{Csc}[e + f*x])^{(m + 1)}, x], x] + \text{Int}[(b*\text{Csc}[e + f*x])^m*(A + C*\text{Csc}[e + f*x]^2), x] /; \text{FreeQ}\{b, e, f, A, B, C, m\}, x]$

3.70.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.54 (sec) , antiderivative size = 726, normalized size of antiderivative = 3.92

method	result
parts	$-\frac{2A\left(5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)+5i\sqrt{\frac{1}{\cos(dx+c)+1}}\sqrt{\frac{\cos(dx+c)}{\cos(dx+c)+1}}\operatorname{EllipticF}\left(i(-\cot(dx+c)+\operatorname{csc}(dx+c)),i\right)\right)}{21d\sqrt{b\sec(dx+c)}b^3}$
default	Expression too large to display

```
input int((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/21*A/d/(b*sec(d*x+c))^(1/2)/b^3*(5*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+
c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+5*I*(1/(c
os(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*
x+c)+csc(d*x+c)),I)*sec(d*x+c)-3*cos(d*x+c)^2*sin(d*x+c)-5*sin(d*x+c))+2/5
*B/d/(cos(d*x+c)+1)/(b*sec(d*x+c))^(1/2)/b^3*(3*I*(1/(cos(d*x+c)+1))^(1/2)
*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)
*cos(d*x+c)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)
*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*cos(d*x+c)+6*I*(1/(cos(d*x+c)+1))
^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+
c)),I)-6*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*Elli
pticF(I*(-cot(d*x+c)+csc(d*x+c)),I)+3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+
c)/(cos(d*x+c)+1))^(1/2)*EllipticE(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c
)-3*I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF
(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)+cos(d*x+c)^2*sin(d*x+c)+sin(d*x+
c)*cos(d*x+c)+3*sin(d*x+c))-2/3*C/d/(b*sec(d*x+c))^(1/2)/b^3*(I*(1/(cos(d*
x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)*EllipticF(I*(-cot(d*x+c)+
csc(d*x+c)),I)+I*(1/(cos(d*x+c)+1))^(1/2)*(cos(d*x+c)/(cos(d*x+c)+1))^(1/2)
)*EllipticF(I*(-cot(d*x+c)+csc(d*x+c)),I)*sec(d*x+c)-sin(d*x+c))
```

3.70.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.04

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx =$$

$$5 \sqrt{2}(5i A + 7i C) \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5 \sqrt{2}(-5i A - 7i C) \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - 63i \sqrt{2} B \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 63i \sqrt{2} B \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - 2(15A \cos(dx + c)^3 + 21B \cos(dx + c)^2 + 5(5A + 7C) \cos(dx + c)) \sqrt{b/\cos(dx + c)} \sin(dx + c) / (b^4 d)$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm="fricas")`

output `-1/105*(5*sqrt(2)*(5*I*A + 7*I*C)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-5*I*A - 7*I*C)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 63*I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 63*I*sqrt(2)*B*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(15*A*cos(d*x + c)^3 + 21*B*cos(d*x + c)^2 + 5*(5*A + 7*C)*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^4*d)`

3.70.6 Sympy [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{\frac{7}{2}}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)**2)/(b*sec(d*x+c))**(7/2),x)`

output `Integral((A + B*sec(c + d*x) + C*sec(c + d*x)**2)/(b*sec(c + d*x))**(7/2), x)`

3.70.7 Maxima [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{(b \sec(dx + c))^{7/2}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm m="maxima")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/(b*sec(d*x + c))^(7/2), x)`

3.70.8 Giac [F]

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{C \sec(dx + c)^2 + B \sec(dx + c) + A}{(b \sec(dx + c))^{7/2}} dx$$

input `integrate((A+B*sec(d*x+c)+C*sec(d*x+c)^2)/(b*sec(d*x+c))^(7/2),x, algorithm m="giac")`

output `integrate((C*sec(d*x + c)^2 + B*sec(d*x + c) + A)/(b*sec(d*x + c))^(7/2), x)`

3.70.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \sec(c + dx) + C \sec^2(c + dx)}{(b \sec(c + dx))^{7/2}} dx = \int \frac{A + \frac{B}{\cos(c+dx)} + \frac{C}{\cos(c+dx)^2}}{\left(\frac{b}{\cos(c+dx)}\right)^{7/2}} dx$$

input `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(7/2),x)`

output `int((A + B/cos(c + d*x) + C/cos(c + d*x)^2)/(b/cos(c + d*x))^(7/2), x)`

APPENDIX

4.1 Listing of Grading functions	465
--	-----

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(" ,
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end proc:

```



```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```

if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):

```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)+"/"+str(ExpnType_optimal)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```



```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf_
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```