

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

6-Hyperbolic-functions/6.4-Hyperbolic-cotangent/176-6.4.7-d-
hyper- \hat{m} -a+b-c-coth- \hat{n} - \hat{p}

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 7:22am

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	40
4	Appendix	398

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	15
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [53]. This is test number [176].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (53)	0.00 (0)
Mathematica	100.00 (53)	0.00 (0)
Fricas	100.00 (53)	0.00 (0)
Maple	81.13 (43)	18.87 (10)
Giac	66.04 (35)	33.96 (18)
Mupad	60.38 (32)	39.62 (21)
Maxima	30.19 (16)	69.81 (37)
Sympy	13.21 (7)	86.79 (46)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	66.038	15.094	18.868	0.000
Maple	33.962	45.283	1.887	18.868
Giac	7.547	47.170	11.321	33.962
Maxima	7.547	16.981	5.660	69.811
Fricas	3.774	90.566	5.660	0.000
Sympy	1.887	11.321	0.000	86.792
Mupad	0.000	60.377	0.000	39.623

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	10	100.00	0.00	0.00
Giac	18	22.22	0.00	77.78
Mupad	21	0.00	100.00	0.00
Maxima	37	100.00	0.00	0.00
Sympy	46	93.48	6.52	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maple	0.19
Maxima	0.29
Rubi	0.29
Fricas	0.40
Giac	0.47
Mathematica	0.80
Mupad	2.44
Sympy	3.84

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	69.38	1.07	60.00	1.07
Mathematica	110.08	1.72	65.00	1.05
Maple	186.91	3.01	137.00	2.74
Maxima	209.25	2.63	64.50	2.18
Mupad	249.41	2.21	44.00	1.00
Sympy	292.71	3.96	253.00	4.35
Giac	316.83	4.81	241.00	4.06
Fricas	3314.85	40.97	2111.00	37.57

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {37, 42, 44, 47, 49}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2013
Design v0.01

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	24
2.3	Detailed conclusion table specific for Rubi results	38

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	21
2.1.3	Maple	22
2.1.4	Fricas	22
2.1.5	Maxima	22
2.1.6	Giac	23
2.1.7	Mupad	23
2.1.8	Sympy	23

2.1.1 Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 13, 14, 15, 16, 18, 19, 20, 22, 23, 24, 25, 26, 28, 29, 31, 32, 34, 36, 38, 48, 49, 50, 51, 52, 53 }

B grade { 10, 12, 17, 27, 30, 33, 35, 39 }

C grade { 21, 37, 40, 41, 42, 43, 44, 45, 46, 47 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.3 Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 50, 51, 52 }

B grade { 16, 17, 18, 19, 22, 23, 24, 25, 28, 29, 30, 31, 32, 33, 34, 35, 38, 39, 40, 43, 44, 45, 48, 49 }

C grade { 53 }

F normal fail { 20, 21, 26, 27, 36, 37, 41, 42, 46, 47 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.4 Fricas

A grade { 11, 15 }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 50, 51, 52, 53 }

C grade { 29, 31, 49 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.5 Maxima

A grade { 5, 11, 13, 15 }

B grade { 1, 2, 3, 4, 6, 7, 8, 9, 50 }

C grade { 10, 12, 14 }

F normal fail { 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 51, 52, 53 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.6 Giac

A grade { 5, 11, 15, 50 }

B grade { 1, 2, 3, 4, 6, 7, 8, 9, 13, 19, 20, 26, 28, 30, 34, 35, 38, 39, 40, 42, 43, 44, 45, 47, 48 }

C grade { 10, 12, 14, 29, 31, 49 }

F normal fail { 27, 51, 52, 53 }

F(-1) timeout fail { }

F(-2) exception fail { 16, 17, 18, 21, 22, 23, 24, 25, 32, 33, 36, 37, 41, 46 }

2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 22, 24, 28, 29, 30, 32, 34, 35, 38, 40, 43, 45, 48, 49, 50 }

C grade { }

F normal fail { }

F(-1) timeout fail { 17, 19, 20, 21, 23, 25, 26, 27, 31, 33, 36, 37, 39, 41, 42, 44, 46, 47, 51, 52, 53 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { 9 }

B grade { 1, 2, 3, 4, 5, 50 }

C grade { }

F normal fail { 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 51, 52, 53 }

F(-1) timeout fail { 6, 7, 8 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	155	231	153	624	2111	639	721	158
N.S.	1	0.97	1.44	0.96	3.90	13.19	3.99	4.51	0.99
time (sec)	N/A	0.329	3.219	0.343	0.216	0.270	14.402	0.330	2.114

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	108	127	112	410	1164	478	447	111
N.S.	1	0.98	1.15	1.02	3.73	10.58	4.35	4.06	1.01
time (sec)	N/A	0.288	2.254	0.245	0.206	0.254	5.832	0.316	2.045

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	75	100	75	239	557	332	241	72
N.S.	1	1.01	1.35	1.01	3.23	7.53	4.49	3.26	0.97
time (sec)	N/A	0.259	1.508	0.177	0.203	0.267	2.329	0.291	2.006

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	47	65	46	114	197	211	103	41
N.S.	1	1.09	1.51	1.07	2.65	4.58	4.91	2.40	0.95
time (sec)	N/A	0.234	0.494	0.119	0.194	0.262	1.086	0.277	0.125

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	45	47	71	56	488	253	65	37
N.S.	1	0.98	1.02	1.54	1.22	10.61	5.50	1.41	0.80
time (sec)	N/A	0.339	0.080	0.130	0.291	0.280	2.489	0.276	0.122

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	106	97	103	207	1952	0	198	110
N.S.	1	1.19	1.09	1.16	2.33	21.93	0.00	2.22	1.24
time (sec)	N/A	0.277	0.626	0.211	0.320	0.327	0.000	0.293	2.313

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	170	147	156	509	7508	0	409	2719
N.S.	1	1.20	1.04	1.10	3.58	52.87	0.00	2.88	19.15
time (sec)	N/A	0.337	0.349	0.306	0.386	0.382	0.000	0.325	2.712

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	239	203	219	925	20031	0	750	3685
N.S.	1	1.19	1.01	1.09	4.60	99.66	0.00	3.73	18.33
time (sec)	N/A	0.437	0.598	0.485	0.506	0.521	0.000	0.357	2.810

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	27	38	70	34	38	15
N.S.	1	1.00	1.00	1.42	2.00	3.68	1.79	2.00	0.79
time (sec)	N/A	0.279	0.115	0.088	0.283	0.265	0.274	0.268	0.114

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	30	4	19	45	0	26	3
N.S.	1	1.00	10.00	1.33	6.33	15.00	0.00	8.67	1.00
time (sec)	N/A	0.223	0.053	0.188	0.277	0.253	0.000	0.284	1.905

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	16	28	15	17	17	0	23	14
N.S.	1	1.14	2.00	1.07	1.21	1.21	0.00	1.64	1.00
time (sec)	N/A	0.237	0.048	0.109	0.279	0.255	0.000	0.291	2.089

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	26	51	21	49	279	0	60	20
N.S.	1	1.08	2.12	0.88	2.04	11.62	0.00	2.50	0.83
time (sec)	N/A	0.238	0.102	0.095	0.302	0.282	0.000	0.274	1.946

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	35	49	28	46	211	0	52	27
N.S.	1	1.13	1.58	0.90	1.48	6.81	0.00	1.68	0.87
time (sec)	N/A	0.241	0.087	0.143	0.301	0.259	0.000	0.268	1.917

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	15	13	14	11	40	0	24	18
N.S.	1	1.15	1.00	1.08	0.85	3.08	0.00	1.85	1.38
time (sec)	N/A	0.233	0.042	0.127	0.280	0.261	0.000	0.271	2.092

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	13	11	12	11	2	0	18	15
N.S.	1	1.18	1.00	1.09	1.00	0.18	0.00	1.64	1.36
time (sec)	N/A	0.228	0.040	0.059	0.276	0.255	0.000	0.283	1.969

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	68	60	253	0	2367	0	0	66
N.S.	1	1.08	0.95	4.02	0.00	37.57	0.00	0.00	1.05
time (sec)	N/A	0.302	0.200	0.211	0.000	0.363	0.000	0.000	3.902

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	89	191	276	0	4877	0	0	0
N.S.	1	1.05	2.25	3.25	0.00	57.38	0.00	0.00	0.00
time (sec)	N/A	0.325	0.836	0.140	0.000	0.434	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	49	44	238	0	1551	0	0	51
N.S.	1	1.11	1.00	5.41	0.00	35.25	0.00	0.00	1.16
time (sec)	N/A	0.265	0.035	0.094	0.000	0.313	0.000	0.000	2.335

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	82	238	0	3455	0	262	0
N.S.	1	1.00	1.37	3.97	0.00	57.58	0.00	4.37	0.00
time (sec)	N/A	0.237	0.329	0.139	0.000	0.369	0.000	0.477	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	61	56	0	0	3479	0	259	0
N.S.	1	1.09	1.00	0.00	0.00	62.12	0.00	4.62	0.00
time (sec)	N/A	0.289	0.038	0.000	0.000	0.380	0.000	0.512	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	48	48	42	0	0	1539	0	0	0
N.S.	1	1.00	0.88	0.00	0.00	32.06	0.00	0.00	0.00
time (sec)	N/A	0.278	0.101	0.000	0.000	0.324	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	89	86	488	0	4940	0	0	112
N.S.	1	1.09	1.05	5.95	0.00	60.24	0.00	0.00	1.37
time (sec)	N/A	0.317	0.437	0.093	0.000	0.470	0.000	0.000	9.988

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	132	219	529	0	10286	0	0	0
N.S.	1	1.07	1.78	4.30	0.00	83.63	0.00	0.00	0.00
time (sec)	N/A	0.431	0.951	0.098	0.000	0.597	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	70	59	473	0	2362	0	0	64
N.S.	1	1.11	0.94	7.51	0.00	37.49	0.00	0.00	1.02
time (sec)	N/A	0.276	0.164	0.074	0.000	0.373	0.000	0.000	4.154

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	92	111	473	0	5037	0	0	0
N.S.	1	1.05	1.26	5.38	0.00	57.24	0.00	0.00	0.00
time (sec)	N/A	0.284	0.578	0.125	0.000	0.429	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	71	76	71	0	0	3949	0	470	0
N.S.	1	1.07	1.00	0.00	0.00	55.62	0.00	6.62	0.00
time (sec)	N/A	0.324	0.076	0.000	0.000	0.426	0.000	3.809	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	77	77	180	0	0	4025	0	0	0
N.S.	1	1.00	2.34	0.00	0.00	52.27	0.00	0.00	0.00
time (sec)	N/A	0.342	0.468	0.000	0.000	0.417	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	60	97	0	683	0	119	68
N.S.	1	1.00	1.94	3.13	0.00	22.03	0.00	3.84	2.19
time (sec)	N/A	0.209	0.120	0.229	0.000	0.261	0.000	0.282	2.038

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	62	142	0	226	0	124	43
N.S.	1	1.00	1.38	3.16	0.00	5.02	0.00	2.76	0.96
time (sec)	N/A	0.227	0.082	0.203	0.000	0.255	0.000	0.302	2.014

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	53	116	158	0	1043	0	265	78
N.S.	1	1.06	2.32	3.16	0.00	20.86	0.00	5.30	1.56
time (sec)	N/A	0.230	0.415	0.103	0.000	0.276	0.000	0.287	2.120

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	70	118	211	0	361	0	285	0
N.S.	1	1.04	1.76	3.15	0.00	5.39	0.00	4.25	0.00
time (sec)	N/A	0.248	0.264	0.107	0.000	0.262	0.000	0.309	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	52	47	129	0	1576	0	0	39
N.S.	1	1.11	1.00	2.74	0.00	33.53	0.00	0.00	0.83
time (sec)	N/A	0.280	0.099	0.117	0.000	0.315	0.000	0.000	2.388

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	134	137	0	3513	0	0	0
N.S.	1	1.00	2.23	2.28	0.00	58.55	0.00	0.00	0.00
time (sec)	N/A	0.297	0.238	0.118	0.000	0.394	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	114	0	1298	0	196	23
N.S.	1	1.00	1.00	3.93	0.00	44.76	0.00	6.76	0.79
time (sec)	N/A	0.255	0.021	0.155	0.000	0.296	0.000	0.419	2.306

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	77	114	0	1357	0	196	25
N.S.	1	1.00	2.48	3.68	0.00	43.77	0.00	6.32	0.81
time (sec)	N/A	0.202	0.137	0.137	0.000	0.298	0.000	0.388	2.223

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	61	56	0	0	3397	0	0	0
N.S.	1	1.09	1.00	0.00	0.00	60.66	0.00	0.00	0.00
time (sec)	N/A	0.292	0.045	0.000	0.000	0.386	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	51	51	127	0	0	1621	0	0	0
N.S.	1	1.00	2.49	0.00	0.00	31.78	0.00	0.00	0.00
time (sec)	N/A	0.287	0.774	0.000	0.000	0.325	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	58	52	287	0	2541	0	359	45
N.S.	1	1.12	1.00	5.52	0.00	48.87	0.00	6.90	0.87
time (sec)	N/A	0.301	0.164	0.097	0.000	0.358	0.000	0.459	2.769

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	109	289	0	2279	0	363	0
N.S.	1	1.00	2.06	5.45	0.00	43.00	0.00	6.85	0.00
time (sec)	N/A	0.294	1.107	0.100	0.000	0.351	0.000	0.459	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	54	41	273	0	2299	0	364	41
N.S.	1	1.10	0.84	5.57	0.00	46.92	0.00	7.43	0.84
time (sec)	N/A	0.278	0.047	0.075	0.000	0.355	0.000	0.471	2.613

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	78	98	70	0	0	6991	0	0	0
N.S.	1	1.26	0.90	0.00	0.00	89.63	0.00	0.00	0.00
time (sec)	N/A	0.345	0.063	0.000	0.000	0.545	0.000	0.000	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	B	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	85	91	260	0	0	3931	0	540	0
N.S.	1	1.07	3.06	0.00	0.00	46.25	0.00	6.35	0.00
time (sec)	N/A	0.369	7.926	0.000	0.000	0.506	0.000	0.599	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	86	63	435	0	6560	0	951	82
N.S.	1	1.16	0.85	5.88	0.00	88.65	0.00	12.85	1.11
time (sec)	N/A	0.322	0.119	0.105	0.000	0.689	0.000	0.596	4.250

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	88	97	215	454	0	6591	0	952	0
N.S.	1	1.10	2.44	5.16	0.00	74.90	0.00	10.82	0.00
time (sec)	N/A	0.334	7.067	0.104	0.000	0.682	0.000	0.551	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	82	43	420	0	5736	0	909	76
N.S.	1	1.17	0.61	6.00	0.00	81.94	0.00	12.99	1.09
time (sec)	N/A	0.292	0.048	0.081	0.000	0.630	0.000	0.575	4.332

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	108	142	73	0	0	19199	0	0	0
N.S.	1	1.31	0.68	0.00	0.00	177.77	0.00	0.00	0.00
time (sec)	N/A	0.392	0.072	0.000	0.000	1.242	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	B	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	131	148	1350	0	0	10729	0	1133	0
N.S.	1	1.13	10.31	0.00	0.00	81.90	0.00	8.65	0.00
time (sec)	N/A	0.454	8.544	0.000	0.000	1.296	0.000	0.769	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	62	0	547	0	69	63
N.S.	1	1.00	1.00	2.48	0.00	21.88	0.00	2.76	2.52
time (sec)	N/A	0.199	0.071	0.200	0.000	0.253	0.000	0.265	2.096

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	C	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	49	66	0	175	0	73	22
N.S.	1	1.00	1.81	2.44	0.00	6.48	0.00	2.70	0.81
time (sec)	N/A	0.202	0.079	0.198	0.000	0.263	0.000	0.267	2.189

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	40	40	41	73	95	102	25	38
N.S.	1	1.05	1.05	1.08	1.92	2.50	2.68	0.66	1.00
time (sec)	N/A	0.317	0.093	0.099	0.289	0.273	0.500	0.263	0.106

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	90	86	116	0	5172	0	0	0
N.S.	1	1.01	0.97	1.30	0.00	58.11	0.00	0.00	0.00
time (sec)	N/A	0.343	0.170	0.815	0.000	0.402	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	43	40	37	0	1290	0	0	0
N.S.	1	1.08	1.00	0.92	0.00	32.25	0.00	0.00	0.00
time (sec)	N/A	0.275	0.021	0.577	0.000	0.406	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	77	73	431	0	3938	0	0	0
N.S.	1	1.04	0.99	5.82	0.00	53.22	0.00	0.00	0.00
time (sec)	N/A	0.316	0.528	0.670	0.000	0.495	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [30] had the largest ratio of [.800000000000000044]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	5	4	0.97	14	0.286
2	A	5	4	0.98	14	0.286
3	A	5	4	1.01	14	0.286
4	A	5	4	1.09	14	0.286
5	A	8	7	0.98	14	0.500
6	A	7	6	1.19	14	0.429
7	A	8	7	1.20	14	0.500
8	A	11	10	1.19	14	0.714
9	A	8	7	1.00	10	0.700
10	A	6	5	1.00	12	0.417
11	A	7	6	1.14	10	0.600
12	A	7	6	1.08	12	0.500
13	A	8	7	1.13	10	0.700
14	A	6	5	1.15	12	0.417
15	A	6	5	1.18	10	0.500
16	A	10	9	1.08	17	0.529
17	A	11	10	1.05	17	0.588
18	A	9	8	1.11	15	0.533
19	A	8	7	1.00	12	0.583
20	A	9	8	1.09	15	0.533
21	A	9	8	1.00	17	0.471
22	A	11	10	1.09	17	0.588

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	14	13	1.07	17	0.765
24	A	10	9	1.11	15	0.600
25	A	10	9	1.05	12	0.750
26	A	11	10	1.07	15	0.667
27	A	11	10	1.00	17	0.588
28	A	7	6	1.00	10	0.600
29	A	8	7	1.00	12	0.583
30	A	9	8	1.06	10	0.800
31	A	10	9	1.04	12	0.750
32	A	9	8	1.11	17	0.471
33	A	10	9	1.00	17	0.529
34	A	8	7	1.00	15	0.467
35	A	5	4	1.00	12	0.333
36	A	9	8	1.09	15	0.533
37	A	9	8	1.00	17	0.471
38	A	9	8	1.12	17	0.471
39	A	8	7	1.00	17	0.412
40	A	9	8	1.10	15	0.533
41	A	11	10	1.26	15	0.667
42	A	12	11	1.07	17	0.647
43	A	10	9	1.16	17	0.529
44	A	10	9	1.10	17	0.529
45	A	10	9	1.17	15	0.600
46	A	13	12	1.31	15	0.800
47	A	13	12	1.13	17	0.706
48	A	5	4	1.00	10	0.400
49	A	5	4	1.00	12	0.333
50	A	5	4	1.05	8	0.500
51	A	13	12	1.01	15	0.800
52	A	8	7	1.08	15	0.467
53	A	11	10	1.04	15	0.667

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (a + b \coth^2(c + dx))^5 dx$	43
3.2	$\int (a + b \coth^2(c + dx))^4 dx$	51
3.3	$\int (a + b \coth^2(c + dx))^3 dx$	58
3.4	$\int (a + b \coth^2(c + dx))^2 dx$	64
3.5	$\int \frac{1}{a + b \coth^2(c + dx)} dx$	70
3.6	$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx$	77
3.7	$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx$	84
3.8	$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx$	92
3.9	$\int \frac{1}{1 - 2 \coth^2(x)} dx$	102
3.10	$\int \sqrt{1 - \coth^2(x)} dx$	107
3.11	$\int \sqrt{-1 + \coth^2(x)} dx$	112
3.12	$\int (1 - \coth^2(x))^{3/2} dx$	117
3.13	$\int (-1 + \coth^2(x))^{3/2} dx$	123
3.14	$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx$	129
3.15	$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx$	134
3.16	$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx$	139
3.17	$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx$	146
3.18	$\int \coth(x) \sqrt{a + b \coth^2(x)} dx$	153
3.19	$\int \sqrt{a + b \coth^2(x)} dx$	159
3.20	$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx$	166
3.21	$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx$	173
3.22	$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx$	179

3.23	$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx$	186
3.24	$\int \coth(x) (a + b \coth^2(x))^{3/2} dx$	194
3.25	$\int (a + b \coth^2(x))^{3/2} dx$	201
3.26	$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx$	208
3.27	$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx$	216
3.28	$\int \sqrt{1 + \coth^2(x)} dx$	223
3.29	$\int \sqrt{-1 - \coth^2(x)} dx$	230
3.30	$\int (1 + \coth^2(x))^{3/2} dx$	236
3.31	$\int (-1 - \coth^2(x))^{3/2} dx$	243
3.32	$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx$	250
3.33	$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx$	256
3.34	$\int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx$	263
3.35	$\int \frac{1}{\sqrt{a+b \coth^2(x)}} dx$	270
3.36	$\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx$	276
3.37	$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx$	282
3.38	$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx$	288
3.39	$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx$	295
3.40	$\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx$	302
3.41	$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx$	309
3.42	$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx$	316
3.43	$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx$	324
3.44	$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx$	332
3.45	$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx$	340
3.46	$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx$	347
3.47	$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx$	354
3.48	$\int \frac{1}{\sqrt{1+\coth^2(x)}} dx$	362
3.49	$\int \frac{1}{\sqrt{-1-\coth^2(x)}} dx$	368
3.50	$\int \frac{1}{1+\coth^3(x)} dx$	373
3.51	$\int \coth(x) \sqrt{a + b \coth^4(x)} dx$	378

3.52	$\int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx$	385
3.53	$\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx$	391

3.1 $\int (a + b \coth^2(c + dx))^5 dx$

3.1.1	Optimal result	43
3.1.2	Mathematica [A] (verified)	43
3.1.3	Rubi [A] (verified)	44
3.1.4	Maple [A] (verified)	46
3.1.5	Fricas [B] (verification not implemented)	46
3.1.6	Sympy [B] (verification not implemented)	47
3.1.7	Maxima [B] (verification not implemented)	48
3.1.8	Giac [B] (verification not implemented)	49
3.1.9	Mupad [B] (verification not implemented)	50

3.1.1 Optimal result

Integrand size = 14, antiderivative size = 160

$$\int (a + b \coth^2(c + dx))^5 dx = (a + b)^5 x - \frac{b(5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4) \coth(c + dx)}{d} - \frac{b^2(10a^3 + 10a^2b + 5ab^2 + b^3) \coth^3(c + dx)}{3d} - \frac{b^3(10a^2 + 5ab + b^2) \coth^5(c + dx)}{5d} - \frac{b^4(5a + b) \coth^7(c + dx)}{7d} - \frac{b^5 \coth^9(c + dx)}{9d}$$

```
output (a+b)^5*x-b*(5*a^4+10*a^3*b+10*a^2*b^2+5*a*b^3+b^4)*coth(d*x+c)/d-1/3*b^2*(10*a^3+10*a^2*b+5*a*b^2+b^3)*coth(d*x+c)^3/d-1/5*b^3*(10*a^2+5*a*b+b^2)*coth(d*x+c)^5/d-1/7*b^4*(5*a+b)*coth(d*x+c)^7/d-1/9*b^5*coth(d*x+c)^9/d
```

3.1.2 Mathematica [A] (verified)

Time = 3.22 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.44

$$\int (a + b \coth^2(c + dx))^5 dx = \frac{b^5 \coth^9(c + dx) \left(35 + 45 \tanh^2(c + dx) + 63 \tanh^4(c + dx) + 105 \tanh^6(c + dx) + 315 \tanh^8(c + dx) \right)}{d}$$

input `Integrate[(a + b*Coth[c + d*x]^2)^5,x]`

output
$$\begin{aligned} & -1/315*(b^5*Coth[c + d*x]^9*(35 + 45*Tanh[c + d*x]^2 + 63*Tanh[c + d*x]^4 \\ & + 105*Tanh[c + d*x]^6 + 315*Tanh[c + d*x]^8 + (1575*a^4*Tanh[c + d*x]^8)/b \\ & ^4 - (315*(a + b)^5*ArcTanh[Sqrt[Tanh[c + d*x]^2]]*Tanh[c + d*x]^10)/(b^5* \\ & Sqrt[Tanh[c + d*x]^2]) + (1050*a^3*Tanh[c + d*x]^6*(1 + 3*Tanh[c + d*x]^2) \\ &)/b^3 + (210*a^2*Tanh[c + d*x]^4*(3 + 5*Tanh[c + d*x]^2 + 15*Tanh[c + d*x] \\ & ^4))/b^2 + (15*a*Tanh[c + d*x]^2*(15 + 21*Tanh[c + d*x]^2 + 35*Tanh[c + d* \\ & x]^4 + 105*Tanh[c + d*x]^6))/b)/d \end{aligned}$$

3.1.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.97, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4144, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \coth^2(c + dx))^5 dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a - b \tan \left(ic + idx + \frac{\pi}{2} \right)^2 \right)^5 dx \\ & \quad \downarrow \text{4144} \\ & \int \frac{(b \coth^2(c+dx)+a)^5}{1-\coth^2(c+dx)} d \coth(c + dx) \\ & \quad \downarrow \text{300} \\ & \frac{\int \left(-b^5 \coth^8(c + dx) - b^4(5a + b) \coth^6(c + dx) - b^3(10a^2 + 5ba + b^2) \coth^4(c + dx) - b^2(10a^3 + 10ba^2 + 5b^2a - \right.}{d} \\ & \quad \downarrow \text{2009} \\ & \left. -\frac{1}{5}b^3(10a^2 + 5ab + b^2) \coth^5(c + dx) - \frac{1}{3}b^2(10a^3 + 10a^2b + 5ab^2 + b^3) \coth^3(c + dx) - b(5a^4 + 10a^3b + 10a^2b^2 + \right.}{d} \end{aligned}$$

3.1. $\int (a + b \coth^2(c + dx))^5 dx$

input `Int[(a + b*Coth[c + d*x]^2)^5,x]`

output `((a + b)^5*ArcTanh[Coth[c + d*x]] - b*(5*a^4 + 10*a^3*b + 10*a^2*b^2 + 5*a*b^3 + b^4)*Coth[c + d*x] - (b^2*(10*a^3 + 10*a^2*b + 5*a*b^2 + b^3)*Coth[c + d*x]^3)/3 - (b^3*(10*a^2 + 5*a*b + b^2)*Coth[c + d*x]^5)/5 - (b^4*(5*a + b)*Coth[c + d*x]^7)/7 - (b^5*Coth[c + d*x]^9)/9)/d`

3.1.3.1 Defintions of rubi rules used

rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.1.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.96

method	result
parallelrisch	$\frac{-35b^5 \coth(dx+c)^9 + (-225ab^4 - 45b^5) \coth(dx+c)^7 + (-630a^2b^3 - 315ab^4 - 63b^5) \coth(dx+c)^5 + (-1050a^3b^2 - 1050a^2b^3 - 525ab^4 - 105b^5) \coth(dx+c)^3 - 1575(a^4 + 2a^3b + 2a^2b^2 + ab^3 + 1/5b^4) b \coth(dx+c) + 315d^2 x^2 (a+b)^5}{315d}$
derivativedivides	$\frac{-5a^4b \coth(dx+c) - 10a^3b^2 \coth(dx+c) - 10a^2b^3 \coth(dx+c) - 5ab^4 \coth(dx+c) - \frac{5ab^4 \coth(dx+c)^7}{7} - 2a^2b^3 \coth(dx+c)^5}{d}$
default	$\frac{-5a^4b \coth(dx+c) - 10a^3b^2 \coth(dx+c) - 10a^2b^3 \coth(dx+c) - 5ab^4 \coth(dx+c) - \frac{5ab^4 \coth(dx+c)^7}{7} - 2a^2b^3 \coth(dx+c)^5}{d}$
parts	$a^5x + \frac{b^5 \left(-\frac{\coth(dx+c)^9}{9} - \frac{\coth(dx+c)^7}{7} - \frac{\coth(dx+c)^5}{5} - \frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} + \frac{\ln(\coth(dx+c)+1)}{2} \right)}{d}$
risch	$a^5x + 5ba^4x + 10a^3b^2x + 10b^3a^2x + 5ab^4x + b^5x - \frac{2b(-31500a^3be^{2dx+2c} - 34020a^2b^2e^{2dx+2c} - 17010ab^3e^{2dx+2c} - 1050b^4e^{2dx+2c} - 1050a^4e^{2dx+2c})}{d}$

input `int((a+coth(d*x+c)^2*b)^5,x,method=_RETURNVERBOSE)`

output `1/315*(-35*b^5*coth(d*x+c)^9+(-225*a*b^4-45*b^5)*coth(d*x+c)^7+(-630*a^2*b^3-315*a*b^4-63*b^5)*coth(d*x+c)^5+(-1050*a^3*b^2-1050*a^2*b^3-525*a*b^4-105*b^5)*coth(d*x+c)^3-1575*(a^4+2*a^3*b+2*a^2*b^2+a*b^3+1/5*b^4)*b*coth(d*x+c)+315*d*x*(a+b)^5)/d`

3.1.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2111 vs. $2(152) = 304$.

Time = 0.27 (sec) , antiderivative size = 2111, normalized size of antiderivative = 13.19

$$\int (a + b \coth^2(c + dx))^5 dx = \text{Too large to display}$$

input `integrate((a+b*coth(d*x+c)^2)^5,x, algorithm="fricas")`

output

```
-1/315*((1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5)*
cosh(d*x + c)^9 + 9*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4
+ 563*b^5)*cosh(d*x + c)*sinh(d*x + c)^8 - (1575*a^4*b + 4200*a^3*b^2 + 4
830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*
a^2*b^3 + 5*a*b^4 + b^5)*d*x)*sinh(d*x + c)^9 - 9*(1225*a^4*b + 2800*a^3*b
^2 + 2730*a^2*b^3 + 1240*a*b^4 + 213*b^5)*cosh(d*x + c)^7 + 9*(1575*a^4*b
+ 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b
+ 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x - 4*(1575*a^4*b + 4200*a^3*
b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b
^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)^7 + 21
*(4*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5)*cosh
(d*x + c)^3 - 3*(1225*a^4*b + 2800*a^3*b^2 + 2730*a^2*b^3 + 1240*a*b^4 + 2
13*b^5)*cosh(d*x + c))*sinh(d*x + c)^6 + 36*(875*a^4*b + 1750*a^3*b^2 + 16
80*a^2*b^3 + 890*a*b^4 + 213*b^5)*cosh(d*x + c)^5 - 9*(6300*a^4*b + 16800*
a^3*b^2 + 19320*a^2*b^3 + 10560*a*b^4 + 2252*b^5 + 14*(1575*a^4*b + 4200*a
^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3
*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^4 + 1260*(a^5 + 5*a
^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x - 21*(1575*a^4*b + 4200
*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a
^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)...
```

3.1.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 639 vs. $2(148) = 296$.

Time = 14.40 (sec) , antiderivative size = 639, normalized size of antiderivative = 3.99

$$\int (a + b \coth^2(c + dx))^5 dx$$

$$= \begin{cases} x(a + b \coth^2(c))^5 \\ -\frac{a^5 \log(-e^{-dx})}{d} - \frac{5a^4 b \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{10a^3 b^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} - \frac{10a^2 b^3 \log(-e^{-dx}) \coth^6(dx + \log(-e^{-dx}))}{d} \\ a^5 x + 5a^4 b x \coth^2(dx + \log(e^{-dx})) + 10a^3 b^2 x \coth^4(dx + \log(e^{-dx})) + 10a^2 b^3 x \coth^6(dx + \log(e^{-dx})) \\ a^5 x + 5a^4 b x - \frac{5a^4 b}{d \tanh(c+dx)} + 10a^3 b^2 x - \frac{10a^3 b^2}{d \tanh(c+dx)} - \frac{10a^3 b^2}{3d \tanh^3(c+dx)} + 10a^2 b^3 x - \frac{10a^2 b^3}{d \tanh(c+dx)} - \frac{10a^2 b^3}{3d \tanh^3(c+dx)} \end{cases}$$

input `integrate((a+b*coth(d*x+c)**2)**5,x)`

3.1. $\int (a + b \coth^2(c + dx))^5 dx$


```
output Piecewise((x*(a + b*coth(c)**2)**5, Eq(d, 0)), (-a**5*log(-exp(-d*x))/d -
5*a**4*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**2/d - 10*a**3*b**2*1
og(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**4/d - 10*a**2*b**3*log(-exp(-d
*x))*coth(d*x + log(-exp(-d*x)))**6/d - 5*a*b**4*log(-exp(-d*x))*coth(d*x
+ log(-exp(-d*x)))**8/d - b**5*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))
**10/d, Eq(c, log(-exp(-d*x)))), (a**5*x + 5*a**4*b*x*coth(d*x + log(exp(-
d*x)))**2 + 10*a**3*b**2*x*coth(d*x + log(exp(-d*x)))**4 + 10*a**2*b**3*x*
coth(d*x + log(exp(-d*x)))**6 + 5*a*b**4*x*coth(d*x + log(exp(-d*x)))**8 +
b**5*x*coth(d*x + log(exp(-d*x)))**10, Eq(c, log(exp(-d*x)))), (a**5*x +
5*a**4*b*x - 5*a**4*b/(d*tanh(c + d*x)) + 10*a**3*b**2*x - 10*a**3*b**2/(d
*tanh(c + d*x)) - 10*a**3*b**2/(3*d*tanh(c + d*x)**3) + 10*a**2*b**3*x - 1
0*a**2*b**3/(d*tanh(c + d*x)) - 10*a**2*b**3/(3*d*tanh(c + d*x)**3) - 2*a*
*2*b**3/(d*tanh(c + d*x)**5) + 5*a*b**4*x - 5*a*b**4/(d*tanh(c + d*x)) - 5
*a*b**4/(3*d*tanh(c + d*x)**3) - a*b**4/(d*tanh(c + d*x)**5) - 5*a*b**4/(7
*d*tanh(c + d*x)**7) + b**5*x - b**5/(d*tanh(c + d*x)) - b**5/(3*d*tanh(c
+ d*x)**3) - b**5/(5*d*tanh(c + d*x)**5) - b**5/(7*d*tanh(c + d*x)**7) - b
**5/(9*d*tanh(c + d*x)**9), True))
```

3.1.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 624 vs. $2(152) = 304$.

Time = 0.22 (sec) , antiderivative size = 624, normalized size of antiderivative = 3.90

$$\int (a + b \coth^2(c + dx))^5 dx$$

$$= \frac{1}{315} b^5 \left(315x + \frac{315c}{d} - \frac{2(3492e^{(-2dx-2c)} - 13968e^{(-4dx-4c)} + 26292e^{(-6dx-6c)} - 39438e^{(-8dx-8c)} + 315e^{(-10dx-10c)})}{d(9e^{(-2dx-2c)} - 36e^{(-4dx-4c)} + 84e^{(-6dx-6c)} - 126e^{(-8dx-8c)} + 126e^{(-10dx-10c)})} \right)$$

$$+ \frac{1}{21} ab^4 \left(105x + \frac{105c}{d} - \frac{8(203e^{(-2dx-2c)} - 609e^{(-4dx-4c)} + 770e^{(-6dx-6c)} - 770e^{(-8dx-8c)} + 315e^{(-10dx-10c)})}{d(7e^{(-2dx-2c)} - 21e^{(-4dx-4c)} + 35e^{(-6dx-6c)} - 35e^{(-8dx-8c)} + 21e^{(-10dx-10c)})} \right)$$

$$+ \frac{2}{3} a^2 b^3 \left(15x + \frac{15c}{d} - \frac{2(70e^{(-2dx-2c)} - 140e^{(-4dx-4c)} + 90e^{(-6dx-6c)} - 45e^{(-8dx-8c)} - 23)}{d(5e^{(-2dx-2c)} - 10e^{(-4dx-4c)} + 10e^{(-6dx-6c)} - 5e^{(-8dx-8c)} + e^{(-10dx-10c)} - 1)} \right)$$

$$+ \frac{10}{3} a^3 b^2 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)}{d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1)} \right)$$

$$+ 5a^4 b \left(x + \frac{c}{d} + \frac{2}{d(e^{(-2dx-2c)} - 1)} \right) + a^5 x$$

```
input integrate((a+b*coth(d*x+c)^2)^5,x, algorithm="maxima")
```

3.1. $\int (a + b \coth^2(c + dx))^5 dx$

output

$$\frac{1}{315}b^5(315x + 315c/d - 2(3492e^{(-2dx - 2c)} - 13968e^{(-4dx - 4c)} + 26292e^{(-6dx - 6c)} - 39438e^{(-8dx - 8c)} + 31500e^{(-10dx - 10c)} - 21000e^{(-12dx - 12c)} + 6300e^{(-14dx - 14c)} - 1575e^{(-16dx - 16c)} - 563)/(d(9e^{(-2dx - 2c)} - 36e^{(-4dx - 4c)} + 84e^{(-6dx - 6c)} - 126e^{(-8dx - 8c)} + 126e^{(-10dx - 10c)} - 84e^{(-12dx - 12c)} + 36e^{(-14dx - 14c)} - 9e^{(-16dx - 16c)} + e^{(-18dx - 18c)} - 1))) + \frac{1}{21}ab^4(105x + 105c/d - 8(203e^{(-2dx - 2c)} - 609e^{(-4dx - 4c)} + 770e^{(-6dx - 6c)} - 770e^{(-8dx - 8c)} + 315e^{(-10dx - 10c)} - 105e^{(-12dx - 12c)} - 44)/(d(7e^{(-2dx - 2c)} - 21e^{(-4dx - 4c)} + 35e^{(-6dx - 6c)} - 35e^{(-8dx - 8c)} + 21e^{(-10dx - 10c)} - 7e^{(-12dx - 12c)} + e^{(-14dx - 14c)} - 1))) + \frac{2}{3}a^2b^3(15x + 15c/d - 2(70e^{(-2dx - 2c)} - 140e^{(-4dx - 4c)} + 90e^{(-6dx - 6c)} - 45e^{(-8dx - 8c)} - 23)/(d(5e^{(-2dx - 2c)} - 10e^{(-4dx - 4c)} + 10e^{(-6dx - 6c)} - 5e^{(-8dx - 8c)} + e^{(-10dx - 10c)} - 1))) + \frac{10}{3}a^3b^2(3x + 3c/d - 4(3e^{(-2dx - 2c)} - 3e^{(-4dx - 4c)} - 2)/(d(3e^{(-2dx - 2c)} - 3e^{(-4dx - 4c)} + e^{(-6dx - 6c)} - 1))) + 5a^4b(x + c/d + 2/(d(e^{(-2dx - 2c)} - 1))) + a^5x$$

3.1.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 721 vs. $2(152) = 304$.

Time = 0.33 (sec) , antiderivative size = 721, normalized size of antiderivative = 4.51

$$\int (a + b \coth^2(c + dx))^5 dx$$

$$= \frac{315(a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5)(dx + c) - \frac{2(1575a^4be^{(16dx+16c)} + 6300a^3b^2e^{(16dx+16c)} + 9450a^2b^3e^{(16dx+16c)} + 2700ab^4e^{(16dx+16c)} + 270b^5e^{(16dx+16c)})}{d^2}}{d^2}$$

input `integrate((a+b*coth(d*x+c)^2)^5,x, algorithm="giac")`

```
output 1/315*(315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*(d*x
+ c) - 2*(1575*a^4*b*e^(16*d*x + 16*c) + 6300*a^3*b^2*e^(16*d*x + 16*c) +
9450*a^2*b^3*e^(16*d*x + 16*c) + 6300*a*b^4*e^(16*d*x + 16*c) + 1575*b^5*e
^(16*d*x + 16*c) - 12600*a^4*b*e^(14*d*x + 14*c) - 44100*a^3*b^2*e^(14*d*x
+ 14*c) - 56700*a^2*b^3*e^(14*d*x + 14*c) - 31500*a*b^4*e^(14*d*x + 14*c)
- 6300*b^5*e^(14*d*x + 14*c) + 44100*a^4*b*e^(12*d*x + 12*c) + 136500*a^3
*b^2*e^(12*d*x + 12*c) + 161700*a^2*b^3*e^(12*d*x + 12*c) + 90300*a*b^4*e
^(12*d*x + 12*c) + 21000*b^5*e^(12*d*x + 12*c) - 88200*a^4*b*e^(10*d*x + 10
*c) - 245700*a^3*b^2*e^(10*d*x + 10*c) - 283500*a^2*b^3*e^(10*d*x + 10*c)
- 157500*a*b^4*e^(10*d*x + 10*c) - 31500*b^5*e^(10*d*x + 10*c) + 110250*a^
4*b*e^(8*d*x + 8*c) + 283500*a^3*b^2*e^(8*d*x + 8*c) + 325080*a^2*b^3*e^(8
*d*x + 8*c) + 175140*a*b^4*e^(8*d*x + 8*c) + 39438*b^5*e^(8*d*x + 8*c) - 8
8200*a^4*b*e^(6*d*x + 6*c) - 216300*a^3*b^2*e^(6*d*x + 6*c) - 244020*a^2*b
^3*e^(6*d*x + 6*c) - 131460*a*b^4*e^(6*d*x + 6*c) - 26292*b^5*e^(6*d*x + 6
*c) + 44100*a^4*b*e^(4*d*x + 4*c) + 107100*a^3*b^2*e^(4*d*x + 4*c) + 11718
0*a^2*b^3*e^(4*d*x + 4*c) + 63540*a*b^4*e^(4*d*x + 4*c) + 13968*b^5*e^(4*d
*x + 4*c) - 12600*a^4*b*e^(2*d*x + 2*c) - 31500*a^3*b^2*e^(2*d*x + 2*c) -
34020*a^2*b^3*e^(2*d*x + 2*c) - 17460*a*b^4*e^(2*d*x + 2*c) - 3492*b^5*e^(
2*d*x + 2*c) + 1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563
*b^5)/(e^(2*d*x + 2*c) - 1)^9/d
```

3.1.9 Mupad [B] (verification not implemented)

Time = 2.11 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.99

$$\int (a + b \coth^2(c + dx))^5 dx = x(a + b)^5 - \frac{\coth(c + dx)^3 (10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5)}{3d} - \frac{\coth(c + dx)^5 (10a^2b^3 + 5ab^4 + b^5)}{5d} - \frac{\coth(c + dx)^7 (b^5 + 5ab^4)}{7d} - \frac{b^5 \coth(c + dx)^9}{9d} - \frac{b \coth(c + dx) (5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4)}{d}$$

```
input int((a + b*coth(c + d*x)^2)^5,x)
```

```
output x*(a + b)^5 - (coth(c + d*x)^3*(5*a*b^4 + b^5 + 10*a^2*b^3 + 10*a^3*b^2))/
(3*d) - (coth(c + d*x)^5*(5*a*b^4 + b^5 + 10*a^2*b^3))/(5*d) - (coth(c + d
*x)^7*(5*a*b^4 + b^5))/(7*d) - (b^5*coth(c + d*x)^9)/(9*d) - (b*coth(c + d
*x)*(5*a*b^3 + 10*a^3*b + 5*a^4 + b^4 + 10*a^2*b^2))/d
```

3.1. $\int (a + b \coth^2(c + dx))^5 dx$

3.2 $\int (a + b \coth^2(c + dx))^4 dx$

3.2.1	Optimal result	51
3.2.2	Mathematica [A] (verified)	51
3.2.3	Rubi [A] (verified)	52
3.2.4	Maple [A] (verified)	53
3.2.5	Fricas [B] (verification not implemented)	54
3.2.6	Sympy [B] (verification not implemented)	55
3.2.7	Maxima [B] (verification not implemented)	56
3.2.8	Giac [B] (verification not implemented)	56
3.2.9	Mupad [B] (verification not implemented)	57

3.2.1 Optimal result

Integrand size = 14, antiderivative size = 110

$$\int (a + b \coth^2(c + dx))^4 dx = (a + b)^4 x - \frac{b(2a + b)(2a^2 + 2ab + b^2) \coth(c + dx)}{d} - \frac{b^2(6a^2 + 4ab + b^2) \coth^3(c + dx)}{3d} - \frac{b^3(4a + b) \coth^5(c + dx)}{5d} - \frac{b^4 \coth^7(c + dx)}{7d}$$

```
output (a+b)^4*x-b*(2*a+b)*(2*a^2+2*a*b+b^2)*coth(d*x+c)/d-1/3*b^2*(6*a^2+4*a*b+b^2)*coth(d*x+c)^3/d-1/5*b^3*(4*a+b)*coth(d*x+c)^5/d-1/7*b^4*coth(d*x+c)^7/d
```

3.2.2 Mathematica [A] (verified)

Time = 2.25 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.15

$$\int (a + b \coth^2(c + dx))^4 dx = \frac{\coth(c + dx) \left(b(105(4a^3 + 6a^2b + 4ab^2 + b^3) + 35b(6a^2 + 4ab + b^2) \coth^2(c + dx) + 21b^2(4a + b) \coth^4(c + dx) \right)}{105d}$$

```
input Integrate[(a + b*Coth[c + d*x]^2)^4,x]
```

output
$$\frac{-1/105*(\text{Coth}[c + d*x]*(b*(105*(4*a^3 + 6*a^2*b + 4*a*b^2 + b^3) + 35*b*(6*a^2 + 4*a*b + b^2))*\text{Coth}[c + d*x]^2 + 21*b^2*(4*a + b)*\text{Coth}[c + d*x]^4 + 15*b^3*\text{Coth}[c + d*x]^6) - 105*(a + b)^4*\text{ArcTanh}[\text{Sqrt}[\text{Tanh}[c + d*x]^2]]*\text{Sqrt}[\text{Tanh}[c + d*x]^2])}{d}$$

3.2.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4144, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \coth^2(c + dx))^4 dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a - b \tan \left(ic + idx + \frac{\pi}{2} \right)^2 \right)^4 dx \\ & \quad \downarrow \text{4144} \\ & \int \frac{(b \coth^2(c+dx)+a)^4}{1-\coth^2(c+dx)} d \coth(c + dx) \\ & \quad \downarrow \text{300} \\ & \frac{\int \left(-b^4 \coth^6(c + dx) - b^3(4a + b) \coth^4(c + dx) - b^2(6a^2 + 4ba + b^2) \coth^2(c + dx) - b(2a + b)(2a^2 + 2ba + b^2) \right) dx}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{-\frac{1}{3}b^2(6a^2 + 4ab + b^2) \coth^3(c + dx) - b(2a + b)(2a^2 + 2ab + b^2) \coth(c + dx) + (a + b)^4 \text{arctanh}(\coth(c + dx))}{d} \end{aligned}$$

input $\text{Int}[(a + b*\text{Coth}[c + d*x]^2)^4, x]$

output
$$\frac{((a + b)^4*\text{ArcTanh}[\text{Coth}[c + d*x]] - b*(2*a + b)*(2*a^2 + 2*a*b + b^2)*\text{Coth}[c + d*x] - (b^2*(6*a^2 + 4*a*b + b^2)*\text{Coth}[c + d*x]^3)/3 - (b^3*(4*a + b)*\text{Coth}[c + d*x]^5)/5 - (b^4*\text{Coth}[c + d*x]^7)/7)}{d}$$

3.2. $\int (a + b \coth^2(c + dx))^4 dx$

3.2.3.1 Defintions of rubi rules used

```
rule 300 Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int
[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c
, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

rule 4144 Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*
(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] ||
EqQ[n^2, 16])
```

3.2.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.02

method	result
parallelrisch	$\frac{-15b^4 \coth(dx+c)^7 + (-84ab^3 - 21b^4) \coth(dx+c)^5 + (-210a^2b^2 - 140ab^3 - 35b^4) \coth(dx+c)^3 + (-420a^3b - 630a^2b^2 - 420ab^3 - 35b^4) \coth(dx+c) - 420a^4}{105d}$
derivativedivides	$\frac{-4 \coth(dx+c)a^3b - 6a^2b^2 \coth(dx+c) - 4ab^3 \coth(dx+c) - \frac{4ab^3 \coth(dx+c)^5}{5} - 2a^2b^2 \coth(dx+c)^3 - \frac{4 \coth(dx+c)^3 a b^3}{3} - b^4}{d}$
default	$\frac{-4 \coth(dx+c)a^3b - 6a^2b^2 \coth(dx+c) - 4ab^3 \coth(dx+c) - \frac{4ab^3 \coth(dx+c)^5}{5} - 2a^2b^2 \coth(dx+c)^3 - \frac{4 \coth(dx+c)^3 a b^3}{3} - b^4}{d}$
parts	$x a^4 + \frac{b^4 \left(-\frac{\coth(dx+c)^7}{7} - \frac{\coth(dx+c)^5}{5} - \frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} + \frac{\ln(\coth(dx+c)+1)}{2} \right)}{d} + \frac{4ab^3}{d}$
risch	$x a^4 + 4b a^3 x + 6a^2 b^2 x + 4a b^3 x + b^4 x - \frac{8b(161a^2b^2 + 105b^3 e^{12dx+12c} + 105a^3 + 609 e^{4dx+4c} b^3 + 315a^2 b e^{12dx+12c})}{d}$

```
input int((a+coth(d*x+c)^2*b)^4,x,method=_RETURNVERBOSE)
```

output $1/105*(-15*b^4*\coth(d*x+c)^7+(-84*a*b^3-21*b^4)*\coth(d*x+c)^5+(-210*a^2*b^2-140*a*b^3-35*b^4)*\coth(d*x+c)^3+(-420*a^3*b-630*a^2*b^2-420*a*b^3-105*b^4)*\coth(d*x+c)+105*d*x*(a+b)^4)/d$

3.2.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1164 vs. $2(104) = 208$.

Time = 0.25 (sec) , antiderivative size = 1164, normalized size of antiderivative = 10.58

$$\int (a + b \coth^2(c + dx))^4 dx = \text{Too large to display}$$

input `integrate((a+b*coth(d*x+c)^2)^4,x, algorithm="fracas")`

output $-1/105*(4*(105*a^3*b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)^7 + 28*(105*a^3*b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)*\sinh(d*x + c)^6 - (420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x)*\sinh(d*x + c)^7 - 28*(75*a^3*b + 120*a^2*b^2 + 71*a*b^3 + 14*b^4)*\cosh(d*x + c)^5 + 7*(420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x - 3*(420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^2)*\sinh(d*x + c)^5 + 140*((105*a^3*b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)^3 - (75*a^3*b + 120*a^2*b^2 + 71*a*b^3 + 14*b^4)*\cosh(d*x + c))*\sinh(d*x + c)^4 + 84*(45*a^3*b + 60*a^2*b^2 + 41*a*b^3 + 14*b^4)*\cosh(d*x + c)^3 - 7*(5*(420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^4 + 1260*a^3*b + 2520*a^2*b^2 + 1932*a*b^3 + 528*b^4 + 315*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x - 10*(420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^2)*\sinh(d*x + c)^3 + 28*(3*(105*a^3*b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)^5 - 10*(75*a^3*b + 120*a^2*b^2 + 71*a*b^3 + 14*b^4)*\cosh(d*x + c)^3 + 9*(45*a^3*b + 60*a^2*b^2 + 41*a*b^3 + 14*b^4)*\cosh(d*x + c))*\sinh(d*x + c)^2 - 420*(5*a^3*b + 6*a^2*b^2 + 5*a*b^3)*\cosh(d*x + c) - 7*((420*a^3*b + 840*a^2*b^2 + 644*a*b^3...$

3.2.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 478 vs. $2(99) = 198$.

Time = 5.83 (sec) , antiderivative size = 478, normalized size of antiderivative = 4.35

$$\int (a + b \coth^2(c + dx))^4 dx$$

$$= \begin{cases} x(a + b \coth^2(c))^4 \\ -\frac{a^4 \log(-e^{-dx})}{d} - \frac{4a^3 b \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{6a^2 b^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} - \frac{4ab^3 \log(-e^{-dx}) \coth^6(dx + \log(-e^{-dx}))}{d} \\ a^4 x + 4a^3 b x \coth^2(dx + \log(e^{-dx})) + 6a^2 b^2 x \coth^4(dx + \log(e^{-dx})) + 4ab^3 x \coth^6(dx + \log(e^{-dx})) \\ a^4 x + 4a^3 b x - \frac{4a^3 b}{d \tanh(c+dx)} + 6a^2 b^2 x - \frac{6a^2 b^2}{d \tanh(c+dx)} - \frac{2a^2 b^2}{d \tanh^3(c+dx)} + 4ab^3 x - \frac{4ab^3}{d \tanh(c+dx)} - \frac{4ab^3}{3d \tanh^3(c+dx)} - \end{cases}$$

input `integrate((a+b*coth(d*x+c)**2)**4,x)`

output `Piecewise((x*(a + b*coth(c)**2)**4, Eq(d, 0)), (-a**4*log(-exp(-d*x))/d - 4*a**3*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**2/d - 6*a**2*b**2*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**4/d - 4*a*b**3*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**6/d - b**4*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**8/d, Eq(c, log(-exp(-d*x)))), (a**4*x + 4*a**3*b*x*coth(d*x + log(exp(-d*x)))**2 + 6*a**2*b**2*x*coth(d*x + log(exp(-d*x)))**4 + 4*a*b**3*x*coth(d*x + log(exp(-d*x)))**6 + b**4*x*coth(d*x + log(exp(-d*x)))**8, Eq(c, log(exp(-d*x)))), (a**4*x + 4*a**3*b*x - 4*a**3*b/(d*tanh(c + d*x)) + 6*a**2*b**2*x - 6*a**2*b**2/(d*tanh(c + d*x)) - 2*a**2*b**2/(d*tanh(c + d*x)**3) + 4*a*b**3*x - 4*a*b**3/(d*tanh(c + d*x)) - 4*a*b**3/(3*d*tanh(c + d*x)**3) - 4*a*b**3/(5*d*tanh(c + d*x)**5) + b**4*x - b**4/(d*tanh(c + d*x)) - b**4/(3*d*tanh(c + d*x)**3) - b**4/(5*d*tanh(c + d*x)**5) - b**4/(7*d*tanh(c + d*x)**7), True))`

3.2.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 410 vs. $2(104) = 208$.

Time = 0.21 (sec) , antiderivative size = 410, normalized size of antiderivative = 3.73

$$\int (a + b \coth^2(c + dx))^4 dx$$

$$= \frac{1}{105} b^4 \left(105x + \frac{105c}{d} - \frac{8(203e^{(-2dx-2c)} - 609e^{(-4dx-4c)} + 770e^{(-6dx-6c)} - 770e^{(-8dx-8c)} + 315e^{(-10dx-10c)} - 105e^{(-12dx-12c)} - 44)}{d(7e^{(-2dx-2c)} - 21e^{(-4dx-4c)} + 35e^{(-6dx-6c)} - 35e^{(-8dx-8c)} + 21e^{(-10dx-10c)} - 7e^{(-12dx-12c)} + e^{(-14dx-14c)} - 1)} \right) + \frac{4}{15} ab^3 \left(15x + \frac{15c}{d} - \frac{2(70e^{(-2dx-2c)} - 140e^{(-4dx-4c)} + 90e^{(-6dx-6c)} - 45e^{(-8dx-8c)} - 23)}{d(5e^{(-2dx-2c)} - 10e^{(-4dx-4c)} + 10e^{(-6dx-6c)} - 5e^{(-8dx-8c)} + e^{(-10dx-10c)} - 1)} \right) + 2a^2b^2 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)}{d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1)} \right) + 4a^3b \left(x + \frac{c}{d} + \frac{2}{d(e^{(-2dx-2c)} - 1)} \right) + a^4x$$

input `integrate((a+b*coth(d*x+c)^2)^4,x, algorithm="maxima")`

output `1/105*b^4*(105*x + 105*c/d - 8*(203*e^(-2*d*x - 2*c) - 609*e^(-4*d*x - 4*c) + 770*e^(-6*d*x - 6*c) - 770*e^(-8*d*x - 8*c) + 315*e^(-10*d*x - 10*c) - 105*e^(-12*d*x - 12*c) - 44)/(d*(7*e^(-2*d*x - 2*c) - 21*e^(-4*d*x - 4*c) + 35*e^(-6*d*x - 6*c) - 35*e^(-8*d*x - 8*c) + 21*e^(-10*d*x - 10*c) - 7*e^(-12*d*x - 12*c) + e^(-14*d*x - 14*c) - 1))) + 4/15*a*b^3*(15*x + 15*c/d - 2*(70*e^(-2*d*x - 2*c) - 140*e^(-4*d*x - 4*c) + 90*e^(-6*d*x - 6*c) - 45*e^(-8*d*x - 8*c) - 23)/(d*(5*e^(-2*d*x - 2*c) - 10*e^(-4*d*x - 4*c) + 10*e^(-6*d*x - 6*c) - 5*e^(-8*d*x - 8*c) + e^(-10*d*x - 10*c) - 1))) + 2*a^2*b^2*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) - 2)/(d*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) - 1))) + 4*a^3*b*(x + c/d + 2/(d*(e^(-2*d*x - 2*c) - 1))) + a^4*x`

3.2.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 447 vs. $2(104) = 208$.

Time = 0.32 (sec) , antiderivative size = 447, normalized size of antiderivative = 4.06

$$\int (a + b \coth^2(c + dx))^4 dx$$

$$= \frac{105(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4)(dx + c) - \frac{8(105a^3be^{(12dx+12c)} + 315a^2b^2e^{(12dx+12c)} + 315ab^3e^{(12dx+12c)} + 105b^4e^{(12dx+12c)})}{d(7e^{(-2dx-2c)} - 21e^{(-4dx-4c)} + 35e^{(-6dx-6c)} - 35e^{(-8dx-8c)} + 21e^{(-10dx-10c)} - 7e^{(-12dx-12c)} + e^{(-14dx-14c)} - 1)}}{d}$$

3.2. $\int (a + b \coth^2(c + dx))^4 dx$

input `integrate((a+b*coth(d*x+c))^2)^4,x, algorithm="giac")`

output
$$\frac{1}{105} \cdot (105 \cdot (a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4) \cdot (dx + c) - 8 \cdot (105a^3be^{12dx+12c} + 315a^2b^2e^{12dx+12c} + 315ab^3e^{12dx+12c} + 105b^4e^{12dx+12c}) - 630a^3be^{10dx+10c} - 1575a^2b^2e^{10dx+10c} - 1260ab^3e^{10dx+10c} - 315b^4e^{10dx+10c} + 1575a^3be^{8dx+8c} + 3360a^2b^2e^{8dx+8c} + 2555ab^3e^{8dx+8c} + 770b^4e^{8dx+8c} - 2100a^3be^{6dx+6c} - 3990a^2b^2e^{6dx+6c} - 3080ab^3e^{6dx+6c} - 770b^4e^{6dx+6c} + 1575a^3be^{4dx+4c} + 2835a^2b^2e^{4dx+4c} + 2121ab^3e^{4dx+4c} + 609b^4e^{4dx+4c} - 630a^3be^{2dx+2c} - 1155a^2b^2e^{2dx+2c} - 812ab^3e^{2dx+2c} - 203b^4e^{2dx+2c} + 105a^3b + 210a^2b^2 + 161ab^3 + 44b^4) / (e^{2dx+2c} - 1)^7 / d$$

3.2.9 Mupad [B] (verification not implemented)

Time = 2.05 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.01

$$\int (a + b \coth^2(c + dx))^4 dx = x(a + b)^4 - \frac{\coth(c + dx)^3 (6a^2b^2 + 4ab^3 + b^4)}{3d} - \frac{\coth(c + dx)^5 (b^4 + 4ab^3)}{5d} - \frac{b^4 \coth(c + dx)^7}{7d} - \frac{b \coth(c + dx) (4a^3 + 6a^2b + 4ab^2 + b^3)}{d}$$

input `int((a + b*coth(c + d*x))^2)^4,x)`

output
$$x \cdot (a + b)^4 - (\coth(c + d \cdot x))^3 \cdot (4 \cdot a \cdot b^3 + b^4 + 6 \cdot a^2 \cdot b^2) / (3 \cdot d) - (\coth(c + d \cdot x))^5 \cdot (4 \cdot a \cdot b^3 + b^4) / (5 \cdot d) - (b^4 \cdot \coth(c + d \cdot x)^7) / (7 \cdot d) - (b \cdot \coth(c + d \cdot x) \cdot (4 \cdot a \cdot b^2 + 6 \cdot a^2 \cdot b + 4 \cdot a^3 + b^3)) / d$$

3.3 $\int (a + b \coth^2(c + dx))^3 dx$

3.3.1	Optimal result	58
3.3.2	Mathematica [A] (verified)	58
3.3.3	Rubi [A] (verified)	59
3.3.4	Maple [A] (verified)	60
3.3.5	Fricas [B] (verification not implemented)	61
3.3.6	Sympy [B] (verification not implemented)	61
3.3.7	Maxima [B] (verification not implemented)	62
3.3.8	Giac [B] (verification not implemented)	63
3.3.9	Mupad [B] (verification not implemented)	63

3.3.1 Optimal result

Integrand size = 14, antiderivative size = 74

$$\int (a + b \coth^2(c + dx))^3 dx = (a + b)^3 x - \frac{b(3a^2 + 3ab + b^2) \coth(c + dx)}{d} - \frac{b^2(3a + b) \coth^3(c + dx)}{3d} - \frac{b^3 \coth^5(c + dx)}{5d}$$

output $(a+b)^3x - b(3a^2+3ab+b^2)*\coth(d*x+c)/d - 1/3*b^2*(3a+b)*\coth(d*x+c)^3/d - 1/5*b^3*\coth(d*x+c)^5/d$

3.3.2 Mathematica [A] (verified)

Time = 1.51 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.35

$$\int (a + b \coth^2(c + dx))^3 dx = -\frac{b \coth(c + dx) (15(3a^2 + 3ab + b^2) + 5b(3a + b) \coth^2(c + dx) + 3b^2 \coth^4(c + dx))}{15d} + \frac{(a + b)^3 \operatorname{arctanh}\left(\sqrt{\tanh^2(c + dx)}\right) \tanh(c + dx)}{d \sqrt{\tanh^2(c + dx)}}$$

input `Integrate[(a + b*Coth[c + d*x]^2)^3,x]`

output
$$\frac{-1/15*(b*\text{Coth}[c + d*x]*(15*(3*a^2 + 3*a*b + b^2) + 5*b*(3*a + b)*\text{Coth}[c + d*x]^2 + 3*b^2*\text{Coth}[c + d*x]^4))/d + ((a + b)^3*\text{ArcTanh}[\text{Sqrt}[\text{Tanh}[c + d*x]^2]]*\text{Tanh}[c + d*x])/(d*\text{Sqrt}[\text{Tanh}[c + d*x]^2])}{1}$$

3.3.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4144, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \coth^2(c + dx))^3 dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a - b \tan \left(ic + idx + \frac{\pi}{2} \right)^2 \right)^3 dx \\ & \quad \downarrow \text{4144} \\ & \frac{\int \frac{(b \coth^2(c+dx)+a)^3}{1-\coth^2(c+dx)} d \coth(c + dx)}{d} \\ & \quad \downarrow \text{300} \\ & \frac{\int \left(-b^3 \coth^4(c + dx) - b^2(3a + b) \coth^2(c + dx) - b(3a^2 + 3ba + b^2) + \frac{(a+b)^3}{1-\coth^2(c+dx)} \right) d \coth(c + dx)}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{-b(3a^2 + 3ab + b^2) \coth(c + dx) + (a + b)^3 \text{arctanh}(\coth(c + dx)) - \frac{1}{3}b^2(3a + b) \coth^3(c + dx) - \frac{1}{5}b^3 \coth^5(c + dx)}{d} \end{aligned}$$

input $\text{Int}[(a + b*\text{Coth}[c + d*x]^2)^3, x]$

output
$$\frac{((a + b)^3*\text{ArcTanh}[\text{Coth}[c + d*x]] - b*(3*a^2 + 3*a*b + b^2)*\text{Coth}[c + d*x] - (b^2*(3*a + b)*\text{Coth}[c + d*x]^3)/3 - (b^3*\text{Coth}[c + d*x]^5)/5)/d}{1}$$

3.3.3.1 Defintions of rubi rules used

rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int [PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.3.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.01

method	result
parallelrisch	$\frac{-3b^3 \coth(dx+c)^5 + (-15ab^2 - 5b^3) \coth(dx+c)^3 + (-45a^2b - 45ab^2 - 15b^3) \coth(dx+c) + 15dx(a+b)^3}{15d}$
derivativedivides	$\frac{-3a^2b \coth(dx+c) - 3 \coth(dx+c)ab^2 - ab^2 \coth(dx+c)^3 - \frac{b^3 \coth(dx+c)^3}{3} - b^3 \coth(dx+c) - \frac{b^3 \coth(dx+c)^5}{5} + \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \coth(dx+c)}{d}}{d}$
default	$\frac{-3a^2b \coth(dx+c) - 3 \coth(dx+c)ab^2 - ab^2 \coth(dx+c)^3 - \frac{b^3 \coth(dx+c)^3}{3} - b^3 \coth(dx+c) - \frac{b^3 \coth(dx+c)^5}{5} + \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \coth(dx+c)}{d}}{d}$
parts	$a^3x + \frac{b^3 \left(-\frac{\coth(dx+c)^5}{5} - \frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} + \frac{\ln(\coth(dx+c)+1)}{2} \right)}{d} + \frac{3ab^2 \left(-\frac{\coth(dx+c)^3}{3} - \coth(dx+c) \right)}{d}$
risch	$a^3x + 3ba^2x + 3ab^2x + b^3x - \frac{2b(45a^2e^{8dx+8c} + 90abe^{8dx+8c} + 45b^2e^{8dx+8c} - 180a^2e^{6dx+6c} - 270abe^{6dx+6c} - 90b^3e^{6dx+6c})}{d}$

input `int((a+coth(d*x+c)^2*b)^3,x,method=_RETURNVERBOSE)`

output `1/15*(-3*b^3*coth(d*x+c)^5+(-15*a*b^2-5*b^3)*coth(d*x+c)^3+(-45*a^2*b-45*a*b^2-15*b^3)*coth(d*x+c)+15*d*x*(a+b)^3)/d`

3.3. $\int (a + b \coth^2(c + dx))^3 dx$

3.3.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 557 vs. $2(70) = 140$.

Time = 0.27 (sec) , antiderivative size = 557, normalized size of antiderivative = 7.53

$$\int (a + b \coth^2(c + dx))^3 dx = \frac{(45 a^2 b + 60 a b^2 + 23 b^3) \cosh(dx + c)^5 + 5(45 a^2 b + 60 a b^2 + 23 b^3) \cosh(dx + c) \sinh(dx + c)^4 - (45 a^3 + 3 a^2 b + 3 a b^2 + b^3) d \cosh(dx + c)^5 - 5(27 a^2 b + 24 a b^2 + 5 b^3) \cosh(dx + c)^3 + 5(45 a^2 b + 60 a b^2 + 23 b^3 + 15(a^3 + 3 a^2 b + 3 a b^2 + b^3) d) \sinh(dx + c)^5 - 5(27 a^2 b + 24 a b^2 + 5 b^3) \cosh(dx + c)^3 + 5(45 a^2 b + 60 a b^2 + 23 b^3 + 15(a^3 + 3 a^2 b + 3 a b^2 + b^3) d) \sinh(dx + c)^3 + 5(2(45 a^2 b + 60 a b^2 + 23 b^3) \cosh(dx + c)^3 - 3(27 a^2 b + 24 a b^2 + 5 b^3) \cosh(dx + c)) \sinh(dx + c)^2 + 10(9 a^2 b + 6 a b^2 + 5 b^3) \cosh(dx + c) - 5((45 a^2 b + 60 a b^2 + 23 b^3 + 15(a^3 + 3 a^2 b + 3 a b^2 + b^3) d) \cosh(dx + c)^4 + 90 a^2 b + 120 a b^2 + 46 b^3 + 30(a^3 + 3 a^2 b + 3 a b^2 + b^3) d) \sinh(dx + c) - 3(45 a^2 b + 60 a b^2 + 23 b^3 + 15(a^3 + 3 a^2 b + 3 a b^2 + b^3) d) \cosh(dx + c)^2 \sinh(dx + c)}{(d \sinh(dx + c))^5 + 5(2 d \cosh(dx + c)^2 - d) \sinh(dx + c)^3 + 5(d \cosh(dx + c))^4 - 3 d \cosh(dx + c)^2 + 2 d) \sinh(dx + c)}$$

input `integrate((a+b*coth(d*x+c))^3,x, algorithm="fracas")`

output

```
-1/15*((45*a^2*b + 60*a*b^2 + 23*b^3)*cosh(d*x + c)^5 + 5*(45*a^2*b + 60*a*b^2 + 23*b^3)*cosh(d*x + c)*sinh(d*x + c)^4 - (45*a^2*b + 60*a*b^2 + 23*b^3 + 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x)*sinh(d*x + c)^5 - 5*(27*a^2*b + 24*a*b^2 + 5*b^3)*cosh(d*x + c)^3 + 5*(45*a^2*b + 60*a*b^2 + 23*b^3 + 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x - 2*(45*a^2*b + 60*a*b^2 + 23*b^3 + 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)^3 + 5*(2*(45*a^2*b + 60*a*b^2 + 23*b^3)*cosh(d*x + c)^3 - 3*(27*a^2*b + 24*a*b^2 + 5*b^3)*cosh(d*x + c))*sinh(d*x + c)^2 + 10*(9*a^2*b + 6*a*b^2 + 5*b^3)*cosh(d*x + c) - 5*((45*a^2*b + 60*a*b^2 + 23*b^3 + 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x)*cosh(d*x + c)^4 + 90*a^2*b + 120*a*b^2 + 46*b^3 + 30*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x - 3*(45*a^2*b + 60*a*b^2 + 23*b^3 + 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c))/(d*sinh(d*x + c))^5 + 5*(2*d*cosh(d*x + c)^2 - d)*sinh(d*x + c)^3 + 5*(d*cosh(d*x + c))^4 - 3*d*cosh(d*x + c)^2 + 2*d)*sinh(d*x + c))
```

3.3.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(65) = 130$.

Time = 2.33 (sec) , antiderivative size = 332, normalized size of antiderivative = 4.49

$$\int (a + b \coth^2(c + dx))^3 dx = \begin{cases} x(a + b \coth^2(c))^3 \\ -\frac{a^3 \log(-e^{-dx})}{d} - \frac{3a^2 b \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{3ab^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} - \frac{b^3 \log(-e^{-dx}) \coth^6(dx + \log(-e^{-dx}))}{d} \\ a^3 x + 3a^2 b x \coth^2(dx + \log(e^{-dx})) + 3ab^2 x \coth^4(dx + \log(e^{-dx})) + b^3 x \coth^6(dx + \log(e^{-dx})) \\ a^3 x + 3a^2 b x - \frac{3a^2 b}{d \tanh(c+dx)} + 3ab^2 x - \frac{3ab^2}{d \tanh(c+dx)} - \frac{ab^2}{d \tanh^3(c+dx)} + b^3 x - \frac{b^3}{d \tanh(c+dx)} - \frac{b^3}{3d \tanh^3(c+dx)} - \frac{b^3}{5d \tanh^5(c+dx)} \end{cases}$$

3.3. $\int (a + b \coth^2(c + dx))^3 dx$

input `integrate((a+b*coth(d*x+c)**2)**3,x)`

output `Piecewise((x*(a + b*coth(c)**2)**3, Eq(d, 0)), (-a**3*log(-exp(-d*x))/d - 3*a**2*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))*2/d - 3*a*b**2*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))*4/d - b**3*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))*6/d, Eq(c, log(-exp(-d*x)))), (a**3*x + 3*a**2*b*x*coth(d*x + log(exp(-d*x)))*2 + 3*a*b**2*x*coth(d*x + log(exp(-d*x)))*4 + b**3*x*coth(d*x + log(exp(-d*x)))*6, Eq(c, log(exp(-d*x)))), (a**3*x + 3*a**2*b*x - 3*a**2*b/(d*tanh(c + d*x)) + 3*a*b**2*x - 3*a*b**2/(d*tanh(c + d*x)) - a*b**2/(d*tanh(c + d*x)**3) + b**3*x - b**3/(d*tanh(c + d*x)) - b**3/(3*d*tanh(c + d*x)**3) - b**3/(5*d*tanh(c + d*x)**5), True))`

3.3.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(70) = 140$.

Time = 0.20 (sec) , antiderivative size = 239, normalized size of antiderivative = 3.23

$$\int (a + b \coth^2(c + dx))^3 dx$$

$$= \frac{1}{15} b^3 \left(15x + \frac{15c}{d} - \frac{2(70e^{(-2dx-2c)} - 140e^{(-4dx-4c)} + 90e^{(-6dx-6c)} - 45e^{(-8dx-8c)} - 23)}{d(5e^{(-2dx-2c)} - 10e^{(-4dx-4c)} + 10e^{(-6dx-6c)} - 5e^{(-8dx-8c)} + e^{(-10dx-10c)} - 1)} \right)$$

$$+ ab^2 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)}{d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1)} \right)$$

$$+ 3a^2b \left(x + \frac{c}{d} + \frac{2}{d(e^{(-2dx-2c)} - 1)} \right) + a^3x$$

input `integrate((a+b*coth(d*x+c)^2)^3,x, algorithm="maxima")`

output `1/15*b^3*(15*x + 15*c/d - 2*(70*e^(-2*d*x - 2*c) - 140*e^(-4*d*x - 4*c) + 90*e^(-6*d*x - 6*c) - 45*e^(-8*d*x - 8*c) - 23)/(d*(5*e^(-2*d*x - 2*c) - 10*e^(-4*d*x - 4*c) + 10*e^(-6*d*x - 6*c) - 5*e^(-8*d*x - 8*c) + e^(-10*d*x - 10*c) - 1))) + a*b^2*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) - 2)/(d*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) - 1))) + 3*a^2*b*(x + c/d + 2/(d*(e^(-2*d*x - 2*c) - 1))) + a^3*x`

3.3.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 241 vs. $2(70) = 140$.

Time = 0.29 (sec) , antiderivative size = 241, normalized size of antiderivative = 3.26

$$\int (a + b \coth^2(c + dx))^3 dx$$

$$= \frac{15(a^3 + 3a^2b + 3ab^2 + b^3)(dx + c) - \frac{2(45a^2be^{(8dx+8c)} + 90ab^2e^{(8dx+8c)} + 45b^3e^{(8dx+8c)} - 180a^2be^{(6dx+6c)} - 270ab^2e^{(6dx+6c)} - 90a^2be^{(4dx+4c)} - 270ab^2e^{(4dx+4c)} - 45a^2be^{(2dx+2c)} - 135ab^2e^{(2dx+2c)} - 45b^3e^{(2dx+2c)} - 180a^2be^{(2dx+2c)} - 270ab^2e^{(2dx+2c)} - 45b^3e^{(2dx+2c)})}{e^{(2dx+2c)} - 1}}{d}$$

input `integrate((a+b*coth(d*x+c)^2)^3,x, algorithm="giac")`

output `1/15*(15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*(d*x + c) - 2*(45*a^2*b*e^(8*d*x + 8*c) + 90*a*b^2*e^(8*d*x + 8*c) + 45*b^3*e^(8*d*x + 8*c) - 180*a^2*b*e^(6*d*x + 6*c) - 270*a*b^2*e^(6*d*x + 6*c) - 90*b^3*e^(6*d*x + 6*c) + 270*a^2*b*e^(4*d*x + 4*c) + 330*a*b^2*e^(4*d*x + 4*c) + 140*b^3*e^(4*d*x + 4*c) - 180*a^2*b*e^(2*d*x + 2*c) - 210*a*b^2*e^(2*d*x + 2*c) - 70*b^3*e^(2*d*x + 2*c) + 45*a^2*b + 60*a*b^2 + 23*b^3)/(e^(2*d*x + 2*c) - 1)^5)/d`

3.3.9 Mupad [B] (verification not implemented)

Time = 2.01 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.97

$$\int (a + b \coth^2(c + dx))^3 dx = x(a + b)^3 - \frac{\coth(c + dx)^3 (b^3 + 3ab^2)}{3d}$$

$$- \frac{b^3 \coth(c + dx)^5}{5d} - \frac{b \coth(c + dx) (3a^2 + 3ab + b^2)}{d}$$

input `int((a + b*coth(c + d*x)^2)^3,x)`

output `x*(a + b)^3 - (coth(c + d*x)^3*(3*a*b^2 + b^3))/(3*d) - (b^3*coth(c + d*x)^5)/(5*d) - (b*coth(c + d*x)*(3*a^2 + 3*a*b + b^2))/d`

3.4 $\int (a + b \coth^2(c + dx))^2 dx$

3.4.1	Optimal result	64
3.4.2	Mathematica [A] (verified)	64
3.4.3	Rubi [A] (verified)	65
3.4.4	Maple [A] (verified)	66
3.4.5	Fricas [B] (verification not implemented)	67
3.4.6	Sympy [B] (verification not implemented)	67
3.4.7	Maxima [B] (verification not implemented)	68
3.4.8	Giac [B] (verification not implemented)	68
3.4.9	Mupad [B] (verification not implemented)	69

3.4.1 Optimal result

Integrand size = 14, antiderivative size = 43

$$\int (a + b \coth^2(c + dx))^2 dx = (a + b)^2 x - \frac{b(2a + b) \coth(c + dx)}{d} - \frac{b^2 \coth^3(c + dx)}{3d}$$

output `(a+b)^2*x-b*(2*a+b)*coth(d*x+c)/d-1/3*b^2*coth(d*x+c)^3/d`

3.4.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.51

$$\int (a + b \coth^2(c + dx))^2 dx = \frac{\coth(c + dx) \left(b(6a + 3b + b \coth^2(c + dx)) - 3(a + b)^2 \operatorname{arctanh} \left(\sqrt{\tanh^2(c + dx)} \right) \sqrt{\tanh^2(c + dx)} \right)}{3d}$$

input `Integrate[(a + b*Coth[c + d*x]^2)^2,x]`

output `-1/3*(Coth[c + d*x]*(b*(6*a + 3*b + b*Coth[c + d*x]^2) - 3*(a + b)^2*ArcTanh[Sqrt[Tanh[c + d*x]^2]]*Sqrt[Tanh[c + d*x]^2]))/d`

3.4.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4144, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \coth^2(c + dx))^2 dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a - b \tan \left(ic + idx + \frac{\pi}{2} \right)^2 \right)^2 dx \\
 & \quad \downarrow \text{4144} \\
 & \frac{\int \frac{(b \coth^2(c+dx)+a)^2}{1-\coth^2(c+dx)} d \coth(c + dx)}{d} \\
 & \quad \downarrow \text{300} \\
 & \frac{\int \left(\frac{(a+b)^2}{1-\coth^2(c+dx)} - b^2 \coth^2(c + dx) - b(2a + b) \right) d \coth(c + dx)}{d} \\
 & \quad \downarrow \text{2009} \\
 & \frac{(a + b)^2 \operatorname{arctanh}(\coth(c + dx)) - b(2a + b) \coth(c + dx) - \frac{1}{3} b^2 \coth^3(c + dx)}{d}
 \end{aligned}$$

input `Int[(a + b*Coth[c + d*x]^2)^2,x]`

output `((a + b)^2*ArcTanh[Coth[c + d*x]] - b*(2*a + b)*Coth[c + d*x] - (b^2*Coth[c + d*x]^3)/3)/d`

3.4.3.1 Defintions of rubi rules used

- rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int [PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`
- rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.4.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.07

method	result
parallelrisch	$\frac{-b^2 \coth(dx+c)^3 + (-6ab-3b^2) \coth(dx+c) + 3dx(a+b)^2}{3d}$
derivativedivides	$\frac{-\frac{b^2 \coth(dx+c)^3}{3} - 2 \coth(dx+c)ab - \coth(dx+c)b^2 - \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)-1)}{2} + \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)+1)}{2}}{d}$
default	$\frac{-\frac{b^2 \coth(dx+c)^3}{3} - 2 \coth(dx+c)ab - \coth(dx+c)b^2 - \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)-1)}{2} + \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)+1)}{2}}{d}$
risch	$a^2x + 2abx + b^2x - \frac{4b(3ae^{4dx+4c} + 3be^{4dx+4c} - 6e^{2dx+2c}a - 3be^{2dx+2c} + 3a+2b)}{3d(e^{2dx+2c}-1)^3}$
parts	$a^2x + \frac{b^2 \left(-\frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} + \frac{\ln(\coth(dx+c)+1)}{2} \right)}{d} + \frac{2ab \left(-\coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} \right)}{d}$

input `int((a+coth(d*x+c)^2*b)^2,x,method=_RETURNVERBOSE)`

output `1/3*(-b^2*coth(d*x+c)^3+(-6*a*b-3*b^2)*coth(d*x+c)+3*d*x*(a+b)^2)/d`

3.4. $\int (a + b \coth^2(c + dx))^2 dx$

3.4.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. $2(41) = 82$.

Time = 0.26 (sec) , antiderivative size = 197, normalized size of antiderivative = 4.58

$$\int (a + b \coth^2(c + dx))^2 dx = \frac{2(3ab + 2b^2) \cosh(dx + c)^3 + 6(3ab + 2b^2) \cosh(dx + c) \sinh(dx + c)^2 - (3(a^2 + 2ab + b^2)dx + 6ab)}{3(d \sinh(dx + c))^3 + 3(d \cosh(dx + c))^2 - d \sinh(dx + c)}$$

input `integrate((a+b*coth(d*x+c))^2,x, algorithm="fracas")`

output `-1/3*(2*(3*a*b + 2*b^2)*cosh(d*x + c)^3 + 6*(3*a*b + 2*b^2)*cosh(d*x + c)*sinh(d*x + c)^2 - (3*(a^2 + 2*a*b + b^2)*d*x + 6*a*b + 4*b^2)*sinh(d*x + c)^3 - 6*a*b*cosh(d*x + c) + 3*(3*(a^2 + 2*a*b + b^2)*d*x - (3*(a^2 + 2*a*b + b^2)*d*x + 6*a*b + 4*b^2)*cosh(d*x + c)^2 + 6*a*b + 4*b^2)*sinh(d*x + c))/((d*sinh(d*x + c))^3 + 3*(d*cosh(d*x + c))^2 - d*sinh(d*x + c))`

3.4.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. $2(36) = 72$.

Time = 1.09 (sec) , antiderivative size = 211, normalized size of antiderivative = 4.91

$$\int (a + b \coth^2(c + dx))^2 dx = \begin{cases} x(a + b \coth^2(c))^2 & \text{for } d = 0 \\ -\frac{a^2 \log(-e^{-dx})}{d} - \frac{2ab \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{b^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} & \text{for } c = \log(-e^{-dx}) \\ a^2x + 2abx \coth^2(dx + \log(e^{-dx})) + b^2x \coth^4(dx + \log(e^{-dx})) & \text{for } c = \log(e^{-dx}) \\ a^2x + 2abx - \frac{2ab}{d \tanh(c+dx)} + b^2x - \frac{b^2}{d \tanh(c+dx)} - \frac{b^2}{3d \tanh^3(c+dx)} & \text{otherwise} \end{cases}$$

input `integrate((a+b*coth(d*x+c)**2)**2,x)`

output `Piecewise((x*(a + b*coth(c)**2)**2, Eq(d, 0)), (-a**2*log(-exp(-d*x))/d - 2*a*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**2/d - b**2*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**4/d, Eq(c, log(-exp(-d*x)))), (a**2*x + 2*a*b*x*coth(d*x + log(exp(-d*x)))**2 + b**2*x*coth(d*x + log(exp(-d*x)))**4, Eq(c, log(exp(-d*x)))), (a**2*x + 2*a*b*x - 2*a*b/(d*tanh(c + d*x)) + b**2*x - b**2/(d*tanh(c + d*x)) - b**2/(3*d*tanh(c + d*x)**3), True))`

3.4. $\int (a + b \coth^2(c + dx))^2 dx$

3.4.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(41) = 82$.

Time = 0.19 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.65

$$\int (a + b \coth^2(c + dx))^2 dx$$

$$= \frac{1}{3} b^2 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)}{d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1)} \right)$$

$$+ 2ab \left(x + \frac{c}{d} + \frac{2}{d(e^{(-2dx-2c)} - 1)} \right) + a^2 x$$

input `integrate((a+b*coth(d*x+c)^2)^2,x, algorithm="maxima")`

output `1/3*b^2*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) - 2)/(d*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) - 1))) + 2*a*b*(x + c/d + 2/(d*(e^(-2*d*x - 2*c) - 1))) + a^2*x`

3.4.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(41) = 82$.

Time = 0.28 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.40

$$\int (a + b \coth^2(c + dx))^2 dx$$

$$= \frac{3(a^2 + 2ab + b^2)(dx + c) - \frac{4(3abe^{(4dx+4c)} + 3b^2e^{(4dx+4c)} - 6abe^{(2dx+2c)} - 3b^2e^{(2dx+2c)} + 3ab + 2b^2)}{(e^{(2dx+2c)} - 1)^3}}{3d}$$

input `integrate((a+b*coth(d*x+c)^2)^2,x, algorithm="giac")`

output `1/3*(3*(a^2 + 2*a*b + b^2)*(d*x + c) - 4*(3*a*b*e^(4*d*x + 4*c) + 3*b^2*e^(4*d*x + 4*c) - 6*a*b*e^(2*d*x + 2*c) - 3*b^2*e^(2*d*x + 2*c) + 3*a*b + 2*b^2)/(e^(2*d*x + 2*c) - 1)^3)/d`

3.4.9 Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int (a + b \coth^2(c + dx))^2 dx = x(a + b)^2 - \frac{b^2 \coth(c + dx)^3}{3d} - \frac{b \coth(c + dx)(2a + b)}{d}$$

input `int((a + b*coth(c + d*x)^2)^2,x)`

output `x*(a + b)^2 - (b^2*coth(c + d*x)^3)/(3*d) - (b*coth(c + d*x)*(2*a + b))/d`

3.5 $\int \frac{1}{a+b \coth^2(c+dx)} dx$

3.5.1	Optimal result	70
3.5.2	Mathematica [A] (verified)	70
3.5.3	Rubi [A] (verified)	71
3.5.4	Maple [A] (verified)	73
3.5.5	Fricas [B] (verification not implemented)	73
3.5.6	Sympy [B] (verification not implemented)	74
3.5.7	Maxima [A] (verification not implemented)	75
3.5.8	Giac [A] (verification not implemented)	75
3.5.9	Mupad [B] (verification not implemented)	76

3.5.1 Optimal result

Integrand size = 14, antiderivative size = 46

$$\int \frac{1}{a+b \coth^2(c+dx)} dx = \frac{x}{a+b} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{\sqrt{a}(a+b)d}$$

output `x/(a+b)-arctan(a^(1/2)*tanh(d*x+c)/b^(1/2))*b^(1/2)/(a+b)/d/a^(1/2)`

3.5.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02

$$\int \frac{1}{a+b \coth^2(c+dx)} dx = \frac{-\frac{\sqrt{b} \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{\sqrt{a}} + \operatorname{arctanh}(\tanh(c+dx))}{(a+b)d}$$

input `Integrate[(a + b*Coth[c + d*x]^2)^(-1),x]`

output `(-((Sqrt[b]*ArcTan[(Sqrt[a]*Tanh[c + d*x])/Sqrt[b]])/Sqrt[a]) + ArcTanh[Tanh[c + d*x]])/((a + b)*d)`

3.5.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4143, 25, 3042, 25, 4158, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + b \coth^2(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{a - b \tan\left(ic + idx + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{4143} \\
 & \frac{b \int -\frac{\operatorname{csch}^2(c+dx)}{b \coth^2(c+dx)+a} dx}{a+b} + \frac{x}{a+b} \\
 & \quad \downarrow \text{25} \\
 & \frac{x}{a+b} - \frac{b \int \frac{\operatorname{csch}^2(c+dx)}{b \coth^2(c+dx)+a} dx}{a+b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x}{a+b} - \frac{b \int -\frac{\sec\left(ic+idx+\frac{\pi}{2}\right)^2}{a-b \tan\left(ic+idx+\frac{\pi}{2}\right)^2} dx}{a+b} \\
 & \quad \downarrow \text{25} \\
 & \frac{x}{a+b} + \frac{b \int \frac{\sec\left(\frac{1}{2}(2ic+\pi)+idx\right)^2}{a-b \tan\left(\frac{1}{2}(2ic+\pi)+idx\right)^2} dx}{a+b} \\
 & \quad \downarrow \text{4158} \\
 & \frac{x}{a+b} - \frac{ib \int \frac{1}{b \coth^2(c+dx)+a} d(i \coth(c + dx))}{d(a+b)} \\
 & \quad \downarrow \text{221} \\
 & \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{\sqrt{ad}(a+b)} + \frac{x}{a+b}
 \end{aligned}$$

input `Int[(a + b*Coth[c + d*x]^2)^(-1),x]`

output `x/(a + b) + (Sqrt[b]*ArcTan[(Sqrt[b]*Coth[c + d*x])/Sqrt[a]])/(Sqrt[a]*(a + b)*d)`

3.5.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4143 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := Simp[x/(a - b), x] - Simp[b/(a - b) Int[Sec[e + f*x]^2/(a + b*Tan[e + f*x]^2), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a, b]`

rule 4158 `Int[sec[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/(c^(m - 1)*f) Subst[Int[(c^2 + ff^2*x^2)^(m/2 - 1)*(a + b*(ff*x)^n)^p, x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[m/2] && (IntegersQ[n, p] || IGtQ[m, 0] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.5.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.54

method	result	size
derivativedivides	$\frac{b \arctan\left(\frac{b \coth(dx+c)}{\sqrt{ab}}\right) - \frac{\ln(\coth(dx+c)-1)}{2a+2b} + \frac{\ln(\coth(dx+c)+1)}{2a+2b}}{d}$	71
default	$\frac{b \arctan\left(\frac{b \coth(dx+c)}{\sqrt{ab}}\right) - \frac{\ln(\coth(dx+c)-1)}{2a+2b} + \frac{\ln(\coth(dx+c)+1)}{2a+2b}}{d}$	71
risch	$\frac{x}{a+b} + \frac{\sqrt{-ab} \ln\left(e^{2dx+2c} - \frac{2\sqrt{-ab}+a-b}{a+b}\right)}{2a(a+b)d} - \frac{\sqrt{-ab} \ln\left(e^{2dx+2c} + \frac{2\sqrt{-ab}-a+b}{a+b}\right)}{2a(a+b)d}$	108

input `int(1/(a+coth(d*x+c)^2*b),x,method=_RETURNVERBOSE)`

output `1/d*(b/(a+b)/(a*b)^(1/2)*arctan(b*coth(d*x+c)/(a*b)^(1/2))-1/(2*a+2*b)*ln(coth(d*x+c)-1)+1/(2*a+2*b)*ln(coth(d*x+c)+1))`

3.5.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 81 vs. 2(38) = 76.

Time = 0.28 (sec) , antiderivative size = 488, normalized size of antiderivative = 10.61

$$\int \frac{1}{a + b \coth^2(c + dx)} dx$$

$$= \left[2 dx + \sqrt{-\frac{b}{a}} \log \left(\frac{(a^2 + 2ab + b^2) \cosh(dx+c)^4 + 4(a^2 + 2ab + b^2) \cosh(dx+c) \sinh(dx+c)^3 + (a^2 + 2ab + b^2) \sinh(dx+c)^4 - 2(a^2 - b^2) \cosh(dx+c) \sinh(dx+c)^2}{(a+b) \cosh(dx+c)^4 + 4(a+b) \cosh(dx+c) \sinh(dx+c)^3 + (a+b) \sinh(dx+c)^4} \right) \right]$$

input `integrate(1/(a+b*coth(d*x+c)^2),x, algorithm="fricas")`

output `[1/2*(2*d*x + sqrt(-b/a)*log(((a^2 + 2*a*b + b^2)*cosh(d*x + c)^4 + 4*(a^2 + 2*a*b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (a^2 + 2*a*b + b^2)*sinh(d*x + c)^4 - 2*(a^2 - b^2)*cosh(d*x + c)^2 + 2*(3*(a^2 + 2*a*b + b^2)*cosh(d*x + c)^2 - a^2 + b^2)*sinh(d*x + c)^2 + a^2 - 6*a*b + b^2 + 4*((a^2 + 2*a*b + b^2)*cosh(d*x + c)^3 - (a^2 - b^2)*cosh(d*x + c))*sinh(d*x + c) - 4*((a^2 + a*b)*cosh(d*x + c)^2 + 2*(a^2 + a*b)*cosh(d*x + c)*sinh(d*x + c) + (a^2 + a*b)*sinh(d*x + c)^2 - a^2 + a*b)*sqrt(-b/a))/((a + b)*cosh(d*x + c)^4 + 4*(a + b)*cosh(d*x + c)*sinh(d*x + c)^3 + (a + b)*sinh(d*x + c)^4 - 2*(a - b)*cosh(d*x + c)^2 + 2*(3*(a + b)*cosh(d*x + c)^2 - a + b)*sinh(d*x + c)^2 + 4*((a + b)*cosh(d*x + c)^3 - (a - b)*cosh(d*x + c))*sinh(d*x + c) + a + b)))/((a + b)*d), (d*x - sqrt(b/a)*arctan(1/2*((a + b)*cosh(d*x + c)^2 + 2*(a + b)*cosh(d*x + c)*sinh(d*x + c) + (a + b)*sinh(d*x + c)^2 - a + b)*sqrt(b/a/b)))/((a + b)*d)]`

3.5.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(37) = 74.

Time = 2.49 (sec) , antiderivative size = 253, normalized size of antiderivative = 5.50

$$\int \frac{1}{a + b \coth^2(c + dx)} dx$$

$$= \begin{cases} \frac{\infty x}{\coth^2(c)} & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ \frac{x - \frac{\tanh(c+dx)}{d}}{b} & \text{for } a = 0 \\ -\frac{dx \tanh^2(c+dx)}{2bd \tanh^2(c+dx) - 2bd} + \frac{dx}{2bd \tanh^2(c+dx) - 2bd} - \frac{\tanh(c+dx)}{2bd \tanh^2(c+dx) - 2bd} & \text{for } a = -b \\ \frac{x}{a} & \text{for } b = 0 \\ \frac{x}{a + b \coth^2(c)} & \text{for } d = 0 \\ \frac{2adx\sqrt{-\frac{b}{a}}}{2a^2d\sqrt{-\frac{b}{a}} + 2abd\sqrt{-\frac{b}{a}}} - \frac{b \log\left(-\sqrt{-\frac{b}{a}} + \tanh(c+dx)\right)}{2a^2d\sqrt{-\frac{b}{a}} + 2abd\sqrt{-\frac{b}{a}}} + \frac{b \log\left(\sqrt{-\frac{b}{a}} + \tanh(c+dx)\right)}{2a^2d\sqrt{-\frac{b}{a}} + 2abd\sqrt{-\frac{b}{a}}} & \text{otherwise} \end{cases}$$

input `integrate(1/(a+b*coth(d*x+c)**2), x)`

```
output Piecewise((zoo*x/coth(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((x - tanh(c
+ d*x)/d)/b, Eq(a, 0)), (-d*x*tanh(c + d*x)**2/(2*b*d*tanh(c + d*x)**2 -
2*b*d) + d*x/(2*b*d*tanh(c + d*x)**2 - 2*b*d) - tanh(c + d*x)/(2*b*d*tanh(
c + d*x)**2 - 2*b*d), Eq(a, -b)), (x/a, Eq(b, 0)), (x/(a + b*coth(c)**2),
Eq(d, 0)), (2*a*d*x*sqrt(-b/a)/(2*a**2*d*sqrt(-b/a) + 2*a*b*d*sqrt(-b/a))
- b*log(-sqrt(-b/a) + tanh(c + d*x))/(2*a**2*d*sqrt(-b/a) + 2*a*b*d*sqrt(-
b/a)) + b*log(sqrt(-b/a) + tanh(c + d*x))/(2*a**2*d*sqrt(-b/a) + 2*a*b*d*s
qrt(-b/a)), True))
```

3.5.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.22

$$\int \frac{1}{a + b \coth^2(c + dx)} dx = \frac{b \arctan\left(\frac{(a+b)e^{(-2dx-2c)} - a + b}{2\sqrt{ab}}\right)}{\sqrt{ab}(a+b)d} + \frac{dx + c}{(a+b)d}$$

```
input integrate(1/(a+b*coth(d*x+c)^2),x, algorithm="maxima")
```

```
output b*arctan(1/2*((a + b)*e^(-2*d*x - 2*c) - a + b)/sqrt(a*b))/(sqrt(a*b)*(a +
b)*d) + (d*x + c)/((a + b)*d)
```

3.5.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.41

$$\int \frac{1}{a + b \coth^2(c + dx)} dx = -\frac{b \arctan\left(\frac{ae^{(2dx+2c)} + be^{(2dx+2c)} - a + b}{2\sqrt{ab}}\right)}{\sqrt{ab}(a+b)} - \frac{dx+c}{a+b}$$

```
input integrate(1/(a+b*coth(d*x+c)^2),x, algorithm="giac")
```

```
output -(b*arctan(1/2*(a*e^(2*d*x + 2*c) + b*e^(2*d*x + 2*c) - a + b)/sqrt(a*b))/
(sqrt(a*b)*(a + b)) - (d*x + c)/(a + b))/d
```

3.5.9 Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \frac{1}{a + b \coth^2(c + dx)} dx = \frac{x}{a + b} + \frac{b \operatorname{atan}\left(\frac{b \coth(c + dx)}{\sqrt{ab}}\right)}{d \sqrt{ab} (a + b)}$$

input `int(1/(a + b*coth(c + d*x)^2),x)`

output `x/(a + b) + (b*atan((b*coth(c + d*x))/(a*b)^(1/2)))/(d*(a*b)^(1/2)*(a + b))`

3.6 $\int \frac{1}{(a+b \coth^2(c+dx))^2} dx$

3.6.1	Optimal result	77
3.6.2	Mathematica [A] (verified)	77
3.6.3	Rubi [A] (verified)	78
3.6.4	Maple [A] (verified)	80
3.6.5	Fricas [B] (verification not implemented)	80
3.6.6	Sympy [F(-1)]	81
3.6.7	Maxima [B] (verification not implemented)	82
3.6.8	Giac [B] (verification not implemented)	82
3.6.9	Mupad [B] (verification not implemented)	83

3.6.1 Optimal result

Integrand size = 14, antiderivative size = 89

$$\int \frac{1}{(a+b \coth^2(c+dx))^2} dx = \frac{x}{(a+b)^2} - \frac{\sqrt{b}(3a+b) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{2a^{3/2}(a+b)^2d} + \frac{b \coth(c+dx)}{2a(a+b)d(a+b \coth^2(c+dx))}$$

output `x/(a+b)^2+1/2*b*coth(d*x+c)/a/(a+b)/d/(a+b*coth(d*x+c)^2)-1/2*(3*a+b)*arctan(a^(1/2)*tanh(d*x+c)/b^(1/2))*b^(1/2)/a^(3/2)/(a+b)^2/d`

3.6.2 Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.09

$$\int \frac{1}{(a+b \coth^2(c+dx))^2} dx = \frac{\sqrt{b}(3a+b) \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{a^{3/2}} + \frac{b(a+b) \coth(c+dx)}{a(a+b \coth^2(c+dx))} - \log(1 - \coth(c+dx)) + \log(1 + \coth(c+dx))$$

$$= \frac{\sqrt{b}(3a+b) \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right) + \frac{b(a+b) \coth(c+dx)}{a(a+b \coth^2(c+dx))} - \log(1 - \coth(c+dx)) + \log(1 + \coth(c+dx))}{2(a+b)^2d}$$

input `Integrate[(a + b*Coth[c + d*x]^2)^(-2),x]`

output $((\text{Sqrt}[b]*(3*a + b)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Coth}[c + d*x])/\text{Sqrt}[a]])/a^{(3/2)} + (b*(a + b)*\text{Coth}[c + d*x])/(a*(a + b*\text{Coth}[c + d*x]^2)) - \text{Log}[1 - \text{Coth}[c + d*x]] + \text{Log}[1 + \text{Coth}[c + d*x]])/(2*(a + b)^{2*d})$

3.6.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.19, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4144, 316, 397, 218, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + b \coth^2(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\left(a - b \tan\left(ic + idx + \frac{\pi}{2}\right)\right)^2} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{1}{(1 - \coth^2(c + dx))(b \coth^2(c + dx) + a)^2} d \coth(c + dx) \\
 & \quad \downarrow \text{316} \\
 & \frac{b \coth(c + dx)}{2a(a + b)(a + b \coth^2(c + dx))} - \frac{\int \frac{b \coth^2(c + dx) + b - 2(a + b)}{(1 - \coth^2(c + dx))(b \coth^2(c + dx) + a)} d \coth(c + dx)}{2a(a + b)} \\
 & \quad \downarrow \text{397} \\
 & \frac{b \coth(c + dx)}{2a(a + b)(a + b \coth^2(c + dx))} - \frac{2a \int \frac{1}{1 - \coth^2(c + dx)} d \coth(c + dx)}{a + b} - \frac{b(3a + b) \int \frac{1}{b \coth^2(c + dx) + a} d \coth(c + dx)}{a + b} \\
 & \quad \downarrow \text{218} \\
 & \frac{b \coth(c + dx)}{2a(a + b)(a + b \coth^2(c + dx))} - \frac{2a \int \frac{1}{1 - \coth^2(c + dx)} d \coth(c + dx)}{a + b} - \frac{\sqrt{b(3a + b)} \arctan\left(\frac{\sqrt{b} \coth(c + dx)}{\sqrt{a}}\right)}{\sqrt{a}(a + b)} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

3.6. $\int \frac{1}{(a + b \coth^2(c + dx))^2} dx$

$$\frac{\frac{b \coth(c+dx)}{2a(a+b)(a+b \coth^2(c+dx))} - \frac{\sqrt{b(3a+b)} \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)} - \frac{2a \operatorname{arctanh}(\coth(c+dx))}{a+b}}{2a(a+b)}$$

d

input `Int[(a + b*Coth[c + d*x]^2)^(-2), x]`

output `(-1/2*(-((Sqrt[b]*(3*a + b)*ArcTan[(Sqrt[b]*Coth[c + d*x])/Sqrt[a]])/(Sqrt[a]*(a + b))) - (2*a*ArcTanh[Coth[c + d*x]]/(a + b))/(a*(a + b)) + (b*Cot h[c + d*x])/(2*a*(a + b)*(a + b*Coth[c + d*x]^2)))/d`

3.6.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))* ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt Q[a, 0] || LtQ[b, 0])`

rule 316 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Sim p[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))), x] + Simp[1/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x ^2)^q*Simp[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && ! (!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, 2, p, q, x]`

rule 397 `Int[((e_) + (f_.)*(x_)^2)/((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e , f}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`


```
rule 4144 Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*
(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] ||
EqQ[n^2, 16])
```

3.6.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.16

method	result
derivativedivides	$\frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^2} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^2} + \frac{b \left(\frac{(a+b) \coth(dx+c)}{2a(a+\coth(dx+c)^2b)} + \frac{(3a+b) \arctan\left(\frac{b \coth(dx+c)}{\sqrt{ab}}\right)}{2a\sqrt{ab}} \right)}{(a+b)^2}}{d}$
default	$\frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^2} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^2} + \frac{b \left(\frac{(a+b) \coth(dx+c)}{2a(a+\coth(dx+c)^2b)} + \frac{(3a+b) \arctan\left(\frac{b \coth(dx+c)}{\sqrt{ab}}\right)}{2a\sqrt{ab}} \right)}{(a+b)^2}}{d}$
risch	$\frac{x}{a^2+2ab+b^2} + \frac{b(e^{2dx+2c}a-b e^{2dx+2c}-a-b)}{da(a+b)^2(ae^{4dx+4c}+be^{4dx+4c}-2e^{2dx+2c}a+2be^{2dx+2c}+a+b)} + \frac{3\sqrt{-ab} \ln\left(\frac{e^{2dx+2c}-2\sqrt{-ab}+a-b}{a+b}\right)}{4a(a+b)^2d}$

```
input int(1/(a+coth(d*x+c)^2*b)^2,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-1/2/(a+b)^2*ln(coth(d*x+c)-1)+1/2/(a+b)^2*ln(coth(d*x+c)+1)+b/(a+b)^
2*(1/2*(a+b)/a*coth(d*x+c)/(a+coth(d*x+c)^2*b)+1/2*(3*a+b)/a/(a*b)^(1/2)*a
rctan(b*coth(d*x+c)/(a*b)^(1/2))))
```

3.6.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 814 vs. 2(77) = 154.

Time = 0.33 (sec) , antiderivative size = 1952, normalized size of antiderivative = 21.93

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \text{Too large to display}$$

```
input integrate(1/(a+b*coth(d*x+c)^2)^2,x, algorithm="fricas")
```

3.6. $\int \frac{1}{(a+b \coth^2(c+dx))^2} dx$

output

```
[1/4*(4*(a^2 + a*b)*d*x*cosh(d*x + c)^4 + 16*(a^2 + a*b)*d*x*cosh(d*x + c)
*sinh(d*x + c)^3 + 4*(a^2 + a*b)*d*x*sinh(d*x + c)^4 + 4*(a^2 + a*b)*d*x -
4*(2*(a^2 - a*b)*d*x - a*b + b^2)*cosh(d*x + c)^2 + 4*(6*(a^2 + a*b)*d*x*
cosh(d*x + c)^2 - 2*(a^2 - a*b)*d*x + a*b - b^2)*sinh(d*x + c)^2 + ((3*a^2
+ 4*a*b + b^2)*cosh(d*x + c)^4 + 4*(3*a^2 + 4*a*b + b^2)*cosh(d*x + c)*si
nh(d*x + c)^3 + (3*a^2 + 4*a*b + b^2)*sinh(d*x + c)^4 - 2*(3*a^2 - 2*a*b -
b^2)*cosh(d*x + c)^2 + 2*(3*(3*a^2 + 4*a*b + b^2)*cosh(d*x + c)^2 - 3*a^2
+ 2*a*b + b^2)*sinh(d*x + c)^2 + 3*a^2 + 4*a*b + b^2 + 4*((3*a^2 + 4*a*b
+ b^2)*cosh(d*x + c)^3 - (3*a^2 - 2*a*b - b^2)*cosh(d*x + c))*sinh(d*x + c
))*sqrt(-b/a)*log(((a^2 + 2*a*b + b^2)*cosh(d*x + c)^4 + 4*(a^2 + 2*a*b +
b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (a^2 + 2*a*b + b^2)*sinh(d*x + c)^4 -
2*(a^2 - b^2)*cosh(d*x + c)^2 + 2*(3*(a^2 + 2*a*b + b^2)*cosh(d*x + c)^2
- a^2 + b^2)*sinh(d*x + c)^2 + a^2 - 6*a*b + b^2 + 4*((a^2 + 2*a*b + b^2)*
cosh(d*x + c)^3 - (a^2 - b^2)*cosh(d*x + c))*sinh(d*x + c) - 4*((a^2 + a*b
)*cosh(d*x + c)^2 + 2*(a^2 + a*b)*cosh(d*x + c)*sinh(d*x + c) + (a^2 + a*b
)*sinh(d*x + c)^2 - a^2 + a*b)*sqrt(-b/a))/((a + b)*cosh(d*x + c)^4 + 4*(a
+ b)*cosh(d*x + c)*sinh(d*x + c)^3 + (a + b)*sinh(d*x + c)^4 - 2*(a - b)*
cosh(d*x + c)^2 + 2*(3*(a + b)*cosh(d*x + c)^2 - a + b)*sinh(d*x + c)^2 +
4*((a + b)*cosh(d*x + c)^3 - (a - b)*cosh(d*x + c))*sinh(d*x + c) + a + b)
) - 4*a*b - 4*b^2 + 8*(2*(a^2 + a*b)*d*x*cosh(d*x + c)^3 - (2*(a^2 - a*...
```

3.6.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \text{Timed out}$$

input `integrate(1/(a+b*coth(d*x+c)**2)**2,x)`

output `Timed out`

3.6.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 207 vs. 2(77) = 154.

Time = 0.32 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.33

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \frac{(3ab + b^2) \arctan\left(\frac{(a+b)e^{(-2dx-2c)} - a + b}{2\sqrt{ab}}\right)}{2(a^3 + 2a^2b + ab^2)\sqrt{abd}} + \frac{ab + b^2 - (ab - b^2)e^{(-2dx-2c)}}{(a^4 + 3a^3b + 3a^2b^2 + ab^3 - 2(a^4 + a^3b - a^2b^2 - ab^3))e^{(-2dx-2c)} + (a^4 + 3a^3b + 3a^2b^2 + ab^3)e^{(-4dx-4c)}} + \frac{dx + c}{(a^2 + 2ab + b^2)d}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^2,x, algorithm="maxima")`

output `1/2*(3*a*b + b^2)*arctan(1/2*((a + b)*e^(-2*d*x - 2*c) - a + b)/sqrt(a*b)) /((a^3 + 2*a^2*b + a*b^2)*sqrt(a*b)*d) + (a*b + b^2 - (a*b - b^2)*e^(-2*d*x - 2*c))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 - 2*(a^4 + a^3*b - a^2*b^2 - a*b^3))*e^(-2*d*x - 2*c) + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*e^(-4*d*x - 4*c))*d) + (d*x + c)/((a^2 + 2*a*b + b^2)*d)`

3.6.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. 2(77) = 154.

Time = 0.29 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.22

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \frac{(3ab + b^2) \arctan\left(\frac{ae^{(2dx+2c)} + be^{(2dx+2c)} - a + b}{2\sqrt{ab}}\right)}{(a^3 + 2a^2b + ab^2)\sqrt{ab}} - \frac{2(dx+c)}{a^2 + 2ab + b^2} - \frac{2(ab e^{(2dx+2c)} - b^2 e^{(2dx+2c)} - ab - b^2)}{(a^3 + 2a^2b + ab^2)(ae^{(4dx+4c)} + be^{(4dx+4c)} - 2ae^{(2dx+2c)} + 2be^{(2dx+2c)})}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^2,x, algorithm="giac")`

output `-1/2*((3*a*b + b^2)*arctan(1/2*(a*e^(2*d*x + 2*c) + b*e^(2*d*x + 2*c) - a + b)/sqrt(a*b))/((a^3 + 2*a^2*b + a*b^2)*sqrt(a*b)) - 2*(d*x + c)/(a^2 + 2*a*b + b^2) - 2*(a*b*e^(2*d*x + 2*c) - b^2*e^(2*d*x + 2*c) - a*b - b^2)/((a^3 + 2*a^2*b + a*b^2)*(a*e^(4*d*x + 4*c) + b*e^(4*d*x + 4*c) - 2*a*e^(2*d*x + 2*c) + 2*b*e^(2*d*x + 2*c) + a + b)))/d`

3.6. $\int \frac{1}{(a + b \coth^2(c + dx))^2} dx$

3.6.9 Mupad [B] (verification not implemented)

Time = 2.31 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.24

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \frac{\frac{ax}{(a+b)^2} + \frac{bx \coth(c+dx)^2}{(a+b)^2} + \frac{b \coth(c+dx)}{2ad(a+b)}}{b \coth(c + dx)^2 + a} + \frac{\operatorname{atan}\left(\frac{b \coth(c+dx)}{\sqrt{ab}}\right) (b^2 + 3ab)}{\sqrt{ab} (2a^3d + ab(4ad + 2bd))}$$

input `int(1/(a + b*coth(c + d*x)^2)^2,x)`output `((a*x)/(a + b)^2 + (b*x*coth(c + d*x)^2)/(a + b)^2 + (b*coth(c + d*x))/(2*a*d*(a + b)))/(a + b*coth(c + d*x)^2) + (atan((b*coth(c + d*x))/(a*b)^(1/2)))*(3*a*b + b^2))/((a*b)^(1/2)*(2*a^3*d + a*b*(4*a*d + 2*b*d)))`

3.7 $\int \frac{1}{(a+b \coth^2(c+dx))^3} dx$

3.7.1	Optimal result	84
3.7.2	Mathematica [A] (verified)	84
3.7.3	Rubi [A] (verified)	85
3.7.4	Maple [A] (verified)	88
3.7.5	Fricas [B] (verification not implemented)	88
3.7.6	Sympy [F(-1)]	89
3.7.7	Maxima [B] (verification not implemented)	89
3.7.8	Giac [B] (verification not implemented)	90
3.7.9	Mupad [B] (verification not implemented)	90

3.7.1 Optimal result

Integrand size = 14, antiderivative size = 142

$$\int \frac{1}{(a+b \coth^2(c+dx))^3} dx = \frac{x}{(a+b)^3} - \frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{8a^{5/2}(a+b)^3d} + \frac{b \coth(c+dx)}{4a(a+b)d(a+b \coth^2(c+dx))^2} + \frac{b(7a+3b) \coth(c+dx)}{8a^2(a+b)^2d(a+b \coth^2(c+dx))}$$

output

```
x/(a+b)^3+1/4*b*coth(d*x+c)/a/(a+b)/d/(a+b*coth(d*x+c)^2)^2+1/8*b*(7*a+3*b)*coth(d*x+c)/a^2/(a+b)^2/d/(a+b*coth(d*x+c)^2)-1/8*(15*a^2+10*a*b+3*b^2)*arctan(a^(1/2)*tanh(d*x+c)/b^(1/2))*b^(1/2)/a^(5/2)/(a+b)^3/d
```

3.7.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.04

$$\int \frac{1}{(a+b \coth^2(c+dx))^3} dx = \frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{a^{5/2}} + \frac{2b(a+b)^2 \coth(c+dx)}{a(a+b \coth^2(c+dx))^2} + \frac{b(a+b)(7a+3b) \coth(c+dx)}{a^2(a+b \coth^2(c+dx))} - 4 \log(1 - \coth(c+dx)) + \dots$$

input `Integrate[(a + b*Coth[c + d*x]^2)^(-3), x]`

output $((\sqrt{b}*(15*a^2 + 10*a*b + 3*b^2)*\text{ArcTan}[(\sqrt{b}*\text{Coth}[c + d*x])/ \sqrt{a}]) / a^{5/2} + (2*b*(a + b)^2*\text{Coth}[c + d*x]) / (a*(a + b*\text{Coth}[c + d*x]^2)^2) + (b*(a + b)*(7*a + 3*b)*\text{Coth}[c + d*x]) / (a^2*(a + b*\text{Coth}[c + d*x]^2)) - 4*\text{Log}[1 - \text{Coth}[c + d*x]] + 4*\text{Log}[1 + \text{Coth}[c + d*x]]) / (8*(a + b)^3*d)$

3.7.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.20, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4144, 316, 402, 397, 218, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx$$

↓ 3042

$$\int \frac{1}{\left(a - b \tan\left(ic + idx + \frac{\pi}{2}\right)\right)^3} dx$$

↓ 4144

$$\frac{\int \frac{1}{(1 - \coth^2(c + dx))(b \coth^2(c + dx) + a)^3} d \coth(c + dx)}{d}$$

↓ 316

$$\frac{\frac{b \coth(c + dx)}{4a(a + b)(a + b \coth^2(c + dx))^2} - \int \frac{3b \coth^2(c + dx) + b - 4(a + b)}{(1 - \coth^2(c + dx))(b \coth^2(c + dx) + a)^2} d \coth(c + dx)}{4a(a + b)}}$$

↓ 402

$$\frac{\frac{b \coth(c + dx)}{4a(a + b)(a + b \coth^2(c + dx))^2} - \frac{\int \frac{8a^2 + 7ba + 3b^2 - b(7a + 3b) \coth^2(c + dx)}{(1 - \coth^2(c + dx))(b \coth^2(c + dx) + a)} d \coth(c + dx)}{2a(a + b)} - \frac{b(7a + 3b) \coth(c + dx)}{2a(a + b)(a + b \coth^2(c + dx))}}{4a(a + b)}}$$

↓ 397

3.7. $\int \frac{1}{(a + b \coth^2(c + dx))^3} dx$

$$\begin{aligned}
& \frac{b \operatorname{coth}(c+dx)}{4a(a+b)(a+b \operatorname{coth}^2(c+dx))^2} - \frac{\frac{b(15a^2+10ab+3b^2) \int \frac{1}{b \operatorname{coth}^2(c+dx)+a} d \operatorname{coth}(c+dx)}{a+b} + \frac{8a^2 \int \frac{1}{1-\operatorname{coth}^2(c+dx)} d \operatorname{coth}(c+dx)}{a+b}}{2a(a+b)} - \frac{b(7a+3b) \operatorname{coth}(c+dx)}{2a(a+b)(a+b \operatorname{coth}^2(c+dx))} \\
& \frac{b \operatorname{coth}(c+dx)}{4a(a+b)(a+b \operatorname{coth}^2(c+dx))^2} - \frac{d}{4a(a+b)} \\
& \quad \downarrow \text{218} \\
& \frac{b \operatorname{coth}(c+dx)}{4a(a+b)(a+b \operatorname{coth}^2(c+dx))^2} - \frac{\frac{8a^2 \int \frac{1}{1-\operatorname{coth}^2(c+dx)} d \operatorname{coth}(c+dx)}{a+b} + \frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{b} \operatorname{coth}(c+dx)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)}}{2a(a+b)} - \frac{b(7a+3b) \operatorname{coth}(c+dx)}{2a(a+b)(a+b \operatorname{coth}^2(c+dx))}}{4a(a+b)} \\
& \frac{b \operatorname{coth}(c+dx)}{4a(a+b)(a+b \operatorname{coth}^2(c+dx))^2} - \frac{d}{4a(a+b)} \\
& \quad \downarrow \text{219} \\
& \frac{b \operatorname{coth}(c+dx)}{4a(a+b)(a+b \operatorname{coth}^2(c+dx))^2} - \frac{\frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{b} \operatorname{coth}(c+dx)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)} + \frac{8a^2 \operatorname{arctanh}(\operatorname{coth}(c+dx))}{a+b}}{2a(a+b)} - \frac{b(7a+3b) \operatorname{coth}(c+dx)}{2a(a+b)(a+b \operatorname{coth}^2(c+dx))}}{4a(a+b)} \\
& \frac{b \operatorname{coth}(c+dx)}{4a(a+b)(a+b \operatorname{coth}^2(c+dx))^2} - \frac{d}{4a(a+b)}
\end{aligned}$$

input `Int[(a + b*Coth[c + d*x]^2)^(-3), x]`

output `((b*Coth[c + d*x])/(4*a*(a + b)*(a + b*Coth[c + d*x]^2)^2) - (-1/2*((Sqrt[b]*(15*a^2 + 10*a*b + 3*b^2)*ArcTan[(Sqrt[b]*Coth[c + d*x])/Sqrt[a]])/(Sqrt[a]*(a + b)) + (8*a^2*ArcTanh[Coth[c + d*x]])/(a + b))/(a*(a + b)) - (b*(7*a + 3*b)*Coth[c + d*x])/(2*a*(a + b)*(a + b*Coth[c + d*x]^2)))/(4*a*(a + b)))/d`

3.7.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

- rule 316 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))], x] + Simp[1/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && (!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, 2, p, q, x]`
- rule 397 `Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`
- rule 402 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a^2*(b*c - a*d)*(p + 1))), x] + Simp[1/(a^2*(b*c - a*d)*(p + 1)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(b*e - a*f) + e*2*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && LtQ[p, -1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.7.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{\frac{\ln(\coth(dx+c)+1)}{2(a+b)^3} - \frac{\ln(\coth(dx+c)-1)}{2(a+b)^3} + \frac{b \left(\frac{b(7a^2+10ab+3b^2)\coth(dx+c)^3}{8a^2} + \frac{(9a^2+14ab+5b^2)\coth(dx+c)}{8a} + \frac{(15a^2+10ab+3b^2)\arctan\left(\frac{b\coth(dx+c)}{a+b}\right)}{8a^2} \right)}{(a+\coth(dx+c)^2b)^2}}{(a+b)^3} + \frac{d}{(a+b)^3}$
default	$\frac{\frac{\ln(\coth(dx+c)+1)}{2(a+b)^3} - \frac{\ln(\coth(dx+c)-1)}{2(a+b)^3} + \frac{b \left(\frac{b(7a^2+10ab+3b^2)\coth(dx+c)^3}{8a^2} + \frac{(9a^2+14ab+5b^2)\coth(dx+c)}{8a} + \frac{(15a^2+10ab+3b^2)\arctan\left(\frac{b\coth(dx+c)}{a+b}\right)}{8a^2} \right)}{(a+\coth(dx+c)^2b)^2}}{(a+b)^3} + \frac{d}{(a+b)^3}$
risch	$\frac{x}{a^3+3a^2b+3ab^2+b^3} + \frac{b(9a^3e^{6dx+6c} - a^2be^{6dx+6c} - 13ab^2e^{6dx+6c} - 3e^{6dx+6c}b^3 - 27a^3e^{4dx+4c} + 9a^2be^{4dx+4c} - 21ab^2e^{4dx+4c} + 3b^3e^{4dx+4c})}{4(ae^{4dx+4c} + be^{4dx+4c} - 2e^{2dx+2c})}$

input `int(1/(a+coth(d*x+c)^2*b)^3,x,method=_RETURNVERBOSE)`

output `1/d*(1/2/(a+b)^3*ln(coth(d*x+c)+1)-1/2/(a+b)^3*ln(coth(d*x+c)-1)+b/(a+b)^3*((1/8*b*(7*a^2+10*a*b+3*b^2)/a^2*coth(d*x+c)^3+1/8*(9*a^2+14*a*b+5*b^2)/a*coth(d*x+c))/(a+coth(d*x+c)^2*b)^2+1/8*(15*a^2+10*a*b+3*b^2)/a^2/(a*b)^(1/2)*arctan(b*coth(d*x+c)/(a*b)^(1/2))))`

3.7.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3592 vs. 2(128) = 256.

Time = 0.38 (sec) , antiderivative size = 7508, normalized size of antiderivative = 52.87

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \text{Too large to display}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^3,x, algorithm="fracas")`

output `Too large to include`

3.7.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \text{Timed out}$$

input `integrate(1/(a+b*coth(d*x+c)**2)**3,x)`

output `Timed out`

3.7.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 509 vs. $2(128) = 256$.

Time = 0.39 (sec) , antiderivative size = 509, normalized size of antiderivative = 3.58

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \frac{(15 a^2 b + 10 a b^2 + 3 b^3) \arctan\left(\frac{(a+b)e^{(-2 dx - 2c)} - a + b}{2\sqrt{ab}}\right)}{8(a^5 + 3 a^4 b + 3 a^3 b^2 + a^2 b^3)\sqrt{abd}}$$

$$+ \frac{9 a^3 b + 21 a^2 b^2 + 15 a b^3 + 3 b^4 - (27 a^5 + 13 a^4 b + 9 a^3 b^2 + 7 a^2 b^3 + 3 a b^4 + b^5)e^{(-2 dx - 2c)}}{4(a^7 + 5 a^6 b + 10 a^5 b^2 + 10 a^4 b^3 + 5 a^3 b^4 + a^2 b^5 - 4(a^7 + 3 a^6 b + 2 a^5 b^2 - 2 a^4 b^3 - 3 a^3 b^4 - a^2 b^5)e^{(-2 dx - 2c)} + 3(9 a^3 b - 3 a^2 b^2 + 7 a b^3 + 3 b^4)e^{(-4 dx - 4c)} - (9 a^3 b - a^2 b^2 - 13 a b^3 - 3 b^4)e^{(-6 dx - 6c)})}$$

$$+ \frac{dx + c}{(a^3 + 3 a^2 b + 3 a b^2 + b^3)d}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^3,x, algorithm="maxima")`

output `1/8*(15*a^2*b + 10*a*b^2 + 3*b^3)*arctan(1/2*((a + b)*e^(-2*d*x - 2*c) - a + b)/sqrt(a*b))/((a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*sqrt(a*b)*d) + 1/4*(9*a^3*b + 21*a^2*b^2 + 15*a*b^3 + 3*b^4 - (27*a^5 + 13*a^4*b + 9*a^3*b^2 + 7*a^2*b^3 + 3*a*b^4 + b^5)*e^(-2*d*x - 2*c) + 3*(9*a^3*b - 3*a^2*b^2 + 7*a*b^3 + 3*b^4)*e^(-4*d*x - 4*c) - (9*a^3*b - a^2*b^2 - 13*a*b^3 - 3*b^4)*e^(-6*d*x - 6*c))/((a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5 - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*e^(-2*d*x - 2*c) + 2*(3*a^7 + 7*a^6*b + 6*a^5*b^2 + 6*a^4*b^3 + 7*a^3*b^4 + 3*a^2*b^5)*e^(-4*d*x - 4*c) - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*e^(-6*d*x - 6*c) + (a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5)*e^(-8*d*x - 8*c))*d) + (d*x + c)/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d)`

3.7.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 409 vs. $2(128) = 256$.

Time = 0.32 (sec) , antiderivative size = 409, normalized size of antiderivative = 2.88

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \frac{(15a^2b + 10ab^2 + 3b^3) \arctan\left(\frac{ae^{2dx+2c} + be^{2dx+2c} - a + b}{2\sqrt{ab}}\right)}{(a^5 + 3a^4b + 3a^3b^2 + a^2b^3)\sqrt{ab}} - \frac{8(dx+c)}{a^3 + 3a^2b + 3ab^2 + b^3} - \frac{2(9a^3be^{6dx+6c} - a^2b^2e^{6dx+6c} - 13ab^3e^{6dx+6c} - 27a^3b^2e^{4dx+4c} + 9a^2b^2e^{4dx+4c} - 21a^2b^3e^{4dx+4c} - 9b^4e^{4dx+4c} + 27a^3be^{2dx+2c} + 13a^2b^2e^{2dx+2c} - 23a^2b^3e^{2dx+2c} - 9b^4e^{2dx+2c} - 9a^3b - 21a^2b^2 - 15a^2b^3 - 3b^4)}{(a^5 + 3a^4b + 3a^3b^2 + a^2b^3)(ae^{4dx+4c} + be^{4dx+4c} - 2ae^{2dx+2c} + 2be^{2dx+2c} + a + b)^2}}{d}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^3,x, algorithm="giac")`

output `-1/8*((15*a^2*b + 10*a*b^2 + 3*b^3)*arctan(1/2*(a*e^(2*d*x + 2*c) + b*e^(2*d*x + 2*c) - a + b)/sqrt(a*b))/((a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*sqrt(a*b)) - 8*(d*x + c)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3) - 2*(9*a^3*b*e^(6*d*x + 6*c) - a^2*b^2*e^(6*d*x + 6*c) - 13*a*b^3*e^(6*d*x + 6*c) - 3*b^4*e^(6*d*x + 6*c) - 27*a^3*b^2*e^(4*d*x + 4*c) + 9*a^2*b^2*e^(4*d*x + 4*c) - 21*a*b^3*e^(4*d*x + 4*c) - 9*b^4*e^(4*d*x + 4*c) + 27*a^3*b*e^(2*d*x + 2*c) + 13*a^2*b^2*e^(2*d*x + 2*c) - 23*a^2*b^3*e^(2*d*x + 2*c) - 9*b^4*e^(2*d*x + 2*c) - 9*a^3*b - 21*a^2*b^2 - 15*a^2*b^3 - 3*b^4)/((a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*(a*e^(4*d*x + 4*c) + b*e^(4*d*x + 4*c) - 2*a*e^(2*d*x + 2*c) + 2*b*e^(2*d*x + 2*c) + a + b)^2))/d`

3.7.9 Mupad [B] (verification not implemented)

Time = 2.71 (sec) , antiderivative size = 2719, normalized size of antiderivative = 19.15

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \text{Too large to display}$$

input `int(1/(a + b*coth(c + d*x)^2)^3,x)`

output

```

log(coth(c + d*x) + 1)/(2*a^3*d + 2*b^3*d + 6*a*b^2*d + 6*a^2*b*d) + ((cot
h(c + d*x)*(9*a*b + 5*b^2))/(8*a*(2*a*b + a^2 + b^2)) + (b*coth(c + d*x)^3
*(7*a*b + 3*b^2))/(8*a*(a*b^2 + 2*a^2*b + a^3)))/(a^2*d + b^2*d*coth(c + d
*x)^4 + 2*a*b*d*coth(c + d*x)^2) - log(coth(c + d*x) - 1)/(2*d*(a + b)^3)
- (atan((((-a^5*b)^(1/2))*((coth(c + d*x)*(60*a*b^6 + 9*b^7 + 190*a^2*b^5 +
300*a^3*b^4 + 289*a^4*b^3)))/(32*(a^8*d^2 + 4*a^7*b*d^2 + a^4*b^4*d^2 + 4*
a^5*b^3*d^2 + 6*a^6*b^2*d^2)) + (((96*a^2*b^10*d^2 + 800*a^3*b^9*d^2 + 304
0*a^4*b^8*d^2 + 6816*a^5*b^7*d^2 + 9760*a^6*b^6*d^2 + 9056*a^7*b^5*d^2 + 5
280*a^8*b^4*d^2 + 1760*a^9*b^3*d^2 + 256*a^10*b^2*d^2)/(64*(a^10*d^3 + 6*a
^9*b*d^3 + a^4*b^6*d^3 + 6*a^5*b^5*d^3 + 15*a^6*b^4*d^3 + 20*a^7*b^3*d^3 +
15*a^8*b^2*d^3)) - (coth(c + d*x)*(-a^5*b)^(1/2)*(10*a*b + 15*a^2 + 3*b^2
)*(256*a^4*b^9*d^2 + 1280*a^5*b^8*d^2 + 2304*a^6*b^7*d^2 + 1280*a^7*b^6*d^
2 - 1280*a^8*b^5*d^2 - 2304*a^9*b^4*d^2 - 1280*a^10*b^3*d^2 - 256*a^11*b^2
*d^2))/(512*(a^8*d + a^5*b^3*d + 3*a^6*b^2*d + 3*a^7*b*d))*(a^8*d^2 + 4*a^7
*b*d^2 + a^4*b^4*d^2 + 4*a^5*b^3*d^2 + 6*a^6*b^2*d^2)))*(-a^5*b)^(1/2)*(10
*a*b + 15*a^2 + 3*b^2))/(16*(a^8*d + a^5*b^3*d + 3*a^6*b^2*d + 3*a^7*b*d))
)*(10*a*b + 15*a^2 + 3*b^2)*i)/(16*(a^8*d + a^5*b^3*d + 3*a^6*b^2*d + 3*a
^7*b*d)) + ((-a^5*b)^(1/2))*((coth(c + d*x)*(60*a*b^6 + 9*b^7 + 190*a^2*b^5
+ 300*a^3*b^4 + 289*a^4*b^3)))/(32*(a^8*d^2 + 4*a^7*b*d^2 + a^4*b^4*d^2 +
4*a^5*b^3*d^2 + 6*a^6*b^2*d^2)) - (((96*a^2*b^10*d^2 + 800*a^3*b^9*d^2 ...

```

3.8 $\int \frac{1}{(a+b \coth^2(c+dx))^4} dx$

3.8.1	Optimal result	92
3.8.2	Mathematica [A] (verified)	93
3.8.3	Rubi [A] (verified)	93
3.8.4	Maple [A] (verified)	97
3.8.5	Fricas [B] (verification not implemented)	97
3.8.6	Sympy [F(-1)]	98
3.8.7	Maxima [B] (verification not implemented)	98
3.8.8	Giac [B] (verification not implemented)	99
3.8.9	Mupad [B] (verification not implemented)	100

3.8.1 Optimal result

Integrand size = 14, antiderivative size = 201

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \frac{x}{(a + b)^4} - \frac{\sqrt{b}(35a^3 + 35a^2b + 21ab^2 + 5b^3) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{16a^{7/2}(a + b)^4d} + \frac{b \coth(c + dx)}{6a(a + b)d (a + b \coth^2(c + dx))^3} + \frac{b(11a + 5b) \coth(c + dx)}{24a^2(a + b)^2d (a + b \coth^2(c + dx))^2} + \frac{b(19a^2 + 16ab + 5b^2) \coth(c + dx)}{16a^3(a + b)^3d (a + b \coth^2(c + dx))}$$

```
output x/(a+b)^4+1/6*b*coth(d*x+c)/a/(a+b)/d/(a+b*coth(d*x+c)^2)^3+1/24*b*(11*a+5
*b)*coth(d*x+c)/a^2/(a+b)^2/d/(a+b*coth(d*x+c)^2)^2+1/16*b*(19*a^2+16*a*b+
5*b^2)*coth(d*x+c)/a^3/(a+b)^3/d/(a+b*coth(d*x+c)^2)-1/16*(35*a^3+35*a^2*b
+21*a*b^2+5*b^3)*arctan(a^(1/2)*tanh(d*x+c)/b^(1/2))*b^(1/2)/a^(7/2)/(a+b)
^4/d
```

3.8.2 Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.01

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx$$

$$= \frac{3\sqrt{b}(35a^3 + 35a^2b + 21ab^2 + 5b^3) \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{a^{7/2}} + \frac{8b(a+b)^3 \coth(c+dx)}{a(a+b \coth^2(c+dx))^3} + \frac{2b(a+b)^2(11a+5b) \coth(c+dx)}{a^2(a+b \coth^2(c+dx))^2} + \frac{3b(a+b)(19a^2+16ab+b^2)}{a^3(a+b \coth^2(c+dx))} + \frac{3b(a+b)(19a^2+16ab+b^2)}{48(a+b)^4 d}$$

input `Integrate[(a + b*Coth[c + d*x]^2)^(-4), x]`

output `((3*sqrt[b]*(35*a^3 + 35*a^2*b + 21*a*b^2 + 5*b^3)*ArcTan[(sqrt[b]*Coth[c + d*x])/sqrt[a]])/a^(7/2) + (8*b*(a + b)^3*Coth[c + d*x])/(a*(a + b*Coth[c + d*x]^2)^3) + (2*b*(a + b)^2*(11*a + 5*b)*Coth[c + d*x])/(a^2*(a + b*Coth[c + d*x]^2)^2) + (3*b*(a + b)*(19*a^2 + 16*a*b + 5*b^2)*Coth[c + d*x])/(a^3*(a + b*Coth[c + d*x]^2)) - 24*Log[1 - Coth[c + d*x]] + 24*Log[1 + Coth[c + d*x]])/(48*(a + b)^4*d)`

3.8.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.19, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 4144, 316, 402, 27, 402, 25, 397, 218, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx$$

$$\downarrow 3042$$

$$\int \frac{1}{\left(a - b \tan\left(ic + idx + \frac{\pi}{2}\right)\right)^4} dx$$

$$\downarrow 4144$$

$$\int \frac{1}{(1 - \coth^2(c+dx))(b \coth^2(c+dx) + a)^4} d \coth(c + dx)$$

$$\downarrow 316$$

3.8. $\int \frac{1}{(a + b \coth^2(c + dx))^4} dx$

$$\begin{aligned}
 & \frac{\frac{b \coth(c+dx)}{6a(a+b)(a+b \coth^2(c+dx))^3} - \frac{\int \frac{5b \coth^2(c+dx)+b-6(a+b)}{(1-\coth^2(c+dx))(b \coth^2(c+dx)+a)^3} d \coth(c+dx)}{6a(a+b)}}{d} \\
 & \quad \downarrow 402 \\
 & \frac{\frac{b \coth(c+dx)}{6a(a+b)(a+b \coth^2(c+dx))^3} - \frac{\int \frac{3(8a^2+11ba+5b^2-b(11a+5b) \coth^2(c+dx))}{(1-\coth^2(c+dx))(b \coth^2(c+dx)+a)^2} d \coth(c+dx)}{4a(a+b)} - \frac{b(11a+5b) \coth(c+dx)}{4a(a+b)(a+b \coth^2(c+dx))^2}}{6a(a+b)} \\
 & \quad \downarrow 27 \\
 & \frac{\frac{b \coth(c+dx)}{6a(a+b)(a+b \coth^2(c+dx))^3} - \frac{3 \int \frac{8a^2+11ba+5b^2-b(11a+5b) \coth^2(c+dx)}{(1-\coth^2(c+dx))(b \coth^2(c+dx)+a)^2} d \coth(c+dx)}{4a(a+b)} - \frac{b(11a+5b) \coth(c+dx)}{4a(a+b)(a+b \coth^2(c+dx))^2}}{6a(a+b)} \\
 & \quad \downarrow 402 \\
 & \frac{\frac{b \coth(c+dx)}{6a(a+b)(a+b \coth^2(c+dx))^3} - \frac{3 \left(\frac{b(19a^2+16ab+5b^2) \coth(c+dx)}{2a(a+b)(a+b \coth^2(c+dx))} - \frac{\int \frac{16a^3+19ba^2+16b^2a+5b^3-b(19a^2+16ba+5b^2) \coth^2(c+dx)}{(1-\coth^2(c+dx))(b \coth^2(c+dx)+a)} d \coth(c+dx)}{2a(a+b)} \right)}{4a(a+b)}}{6a(a+b)} \\
 & \quad \downarrow 25 \\
 & \frac{\frac{b \coth(c+dx)}{6a(a+b)(a+b \coth^2(c+dx))^3} - \frac{3 \left(\frac{\int \frac{16a^3+19ba^2+16b^2a+5b^3-b(19a^2+16ba+5b^2) \coth^2(c+dx)}{(1-\coth^2(c+dx))(b \coth^2(c+dx)+a)} d \coth(c+dx)}{2a(a+b)} + \frac{b(19a^2+16ab+5b^2) \coth(c+dx)}{2a(a+b)(a+b \coth^2(c+dx))} \right)}{4a(a+b)}}{6a(a+b)} \\
 & \quad \downarrow 397 \\
 & \frac{\frac{b \coth(c+dx)}{6a(a+b)(a+b \coth^2(c+dx))^3} - \frac{3 \left(\frac{16a^3 \int \frac{1}{1-\coth^2(c+dx)} d \coth(c+dx)}{a+b} + \frac{b(35a^3+35a^2b+21ab^2+5b^3) \int \frac{1}{b \coth^2(c+dx)+a} d \coth(c+dx)}{2a(a+b)} + \frac{b(19a^2+16ab+5b^2) \coth(c+dx)}{2a(a+b)(a+b \coth^2(c+dx))} \right)}{4a(a+b)}}{6a(a+b)} \\
 & \quad \downarrow 218
 \end{aligned}$$

3.8. $\int \frac{1}{(a+b \coth^2(c+dx))^4} dx$

$$\frac{\frac{b \operatorname{coth}(c+dx)}{6a(a+b)(a+b \operatorname{coth}^2(c+dx))^3} - \frac{\frac{16a^3 \int \frac{1}{1-\operatorname{coth}^2(c+dx)} d \operatorname{coth}(c+dx) + \frac{\sqrt{b}(35a^3+35a^2b+21ab^2+5b^3) \arctan\left(\frac{\sqrt{b} \operatorname{coth}(c+dx)}{\sqrt{a}}\right)}{2a(a+b)\sqrt{a(a+b)}} + \frac{b(19a^2+16ab+5b^2) \operatorname{coth}(c+dx)}{2a(a+b)(a+b \operatorname{coth}^2(c+dx))}}{4a(a+b)}}{6a(a+b)}}{d}$$

↓ 219

$$\frac{\frac{b \operatorname{coth}(c+dx)}{6a(a+b)(a+b \operatorname{coth}^2(c+dx))^3} - \frac{\frac{b(19a^2+16ab+5b^2) \operatorname{coth}(c+dx)}{2a(a+b)(a+b \operatorname{coth}^2(c+dx))} + \frac{16a^3 \operatorname{arctanh}(\operatorname{coth}(c+dx))}{a+b} + \frac{\sqrt{b}(35a^3+35a^2b+21ab^2+5b^3) \arctan\left(\frac{\sqrt{b} \operatorname{coth}(c+dx)}{\sqrt{a}}\right)}{2a(a+b)\sqrt{a(a+b)}}}{4a(a+b)}}{6a(a+b)}}{d}$$

input `Int[(a + b*Coth[c + d*x]^2)^(-4), x]`

output `((b*Coth[c + d*x])/(6*a*(a + b)*(a + b*Coth[c + d*x]^2)^3) - (-1/4*(b*(11*a + 5*b)*Coth[c + d*x])/(a*(a + b)*(a + b*Coth[c + d*x]^2)^2) - (3*(((Sqrt[b]*(35*a^3 + 35*a^2*b + 21*a*b^2 + 5*b^3)*ArcTan[(Sqrt[b]*Coth[c + d*x])/Sqrt[a]])/(Sqrt[a]*(a + b)) + (16*a^3*ArcTanh[Coth[c + d*x]])/(a + b))/(2*a*(a + b)) + (b*(19*a^2 + 16*a*b + 5*b^2)*Coth[c + d*x])/(2*a*(a + b)*(a + b*Coth[c + d*x]^2))))/(4*a*(a + b)))/(6*a*(a + b))/d`

3.8.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.8. $\int \frac{1}{(a+b \operatorname{coth}^2(c+dx))^4} dx$

- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 316 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d)), x] + Simp[1/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && (!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, 2, p, q, x]`
- rule 397 `Int[((e_) + (f_.)*(x_)^2)/((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`
- rule 402 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a^2*(b*c - a*d)*(p + 1))), x] + Simp[1/(a^2*(b*c - a*d)*(p + 1)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(b*e - a*f) + e*2*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && LtQ[p, -1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.8.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^4} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^4} + \frac{b \left(\frac{b^2(19a^3+35a^2b+21ab^2+5b^3)}{16a^3} \coth(dx+c)^5 + \frac{b(17a^3+33a^2b+21ab^2+5b^3)}{6a^2} \coth(dx+c) \right)}{(a+\coth(dx+c)^2b)^3}}{d}$
default	$\frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^4} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^4} + \frac{b \left(\frac{b^2(19a^3+35a^2b+21ab^2+5b^3)}{16a^3} \coth(dx+c)^5 + \frac{b(17a^3+33a^2b+21ab^2+5b^3)}{6a^2} \coth(dx+c) \right)}{(a+\coth(dx+c)^2b)^3}}{d}$
risch	Expression too large to display

input `int(1/(a+coth(d*x+c)^2*b)^4,x,method=_RETURNVERBOSE)`

output `1/d*(-1/2/(a+b)^4*ln(coth(d*x+c)-1)+1/2/(a+b)^4*ln(coth(d*x+c)+1)+1/(a+b)^4*b*((1/16*b^2*(19*a^3+35*a^2*b+21*a*b^2+5*b^3)/a^3*coth(d*x+c)^5+1/6*b*(17*a^3+33*a^2*b+21*a*b^2+5*b^3)/a^2*coth(d*x+c)^3+1/16*(29*a^3+61*a^2*b+43*a*b^2+11*b^3)/a*coth(d*x+c))/(a+coth(d*x+c)^2*b)^3+1/16*(35*a^3+35*a^2*b+21*a*b^2+5*b^3)/a^3/(a*b)^(1/2)*arctan(b*coth(d*x+c)/(a*b)^(1/2))))`

3.8.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9853 vs. 2(185) = 370.

Time = 0.52 (sec) , antiderivative size = 20031, normalized size of antiderivative = 99.66

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^4,x, algorithm="fricas")`

output Too large to include

3.8.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \text{Timed out}$$

input `integrate(1/(a+b*coth(d*x+c)**2)**4,x)`

output `Timed out`

3.8.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 925 vs. 2(185) = 370.

Time = 0.51 (sec) , antiderivative size = 925, normalized size of antiderivative = 4.60

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \frac{(35 a^3 b + 35 a^2 b^2 + 21 a b^3 + 5 b^4) \arctan\left(\frac{(a+b)e^{(-2 dx - 2c)} - a + b}{2\sqrt{ab}}\right)}{16 (a^7 + 4 a^6 b + 6 a^5 b^2 + 4 a^4 b^3 + a^3 b^4) \sqrt{abd}}$$

$$+ \frac{87 a^5 b + 319 a^4 b^2 + 450}{24 (a^{10} + 7 a^9 b + 21 a^8 b^2 + 35 a^7 b^3 + 35 a^6 b^4 + 21 a^5 b^5 + 7 a^4 b^6 + a^3 b^7 - 6 (a^{10} + 5 a^9 b + 9 a^8 b^2 + 5 a^7 b^3$$

$$+ \frac{dx + c}{(a^4 + 4 a^3 b + 6 a^2 b^2 + 4 a b^3 + b^4) d}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^4,x, algorithm="maxima")`

output $1/16*(35*a^3*b + 35*a^2*b^2 + 21*a*b^3 + 5*b^4)*\arctan(1/2*((a + b)*e^{(-2*d*x - 2*c)} - a + b)/\sqrt{a*b})/((a^7 + 4*a^6*b + 6*a^5*b^2 + 4*a^4*b^3 + a^3*b^4)*\sqrt{a*b}*d) + 1/24*(87*a^5*b + 319*a^4*b^2 + 450*a^3*b^3 + 306*a^2*b^4 + 103*a*b^5 + 15*b^6 - 3*(145*a^5*b + 267*a^4*b^2 + 34*a^3*b^3 - 178*a^2*b^4 - 115*a*b^5 - 25*b^6))*e^{(-2*d*x - 2*c)} + 6*(145*a^5*b + 93*a^4*b^2 - 6*a^3*b^3 + 106*a^2*b^4 + 85*a*b^5 + 25*b^6))*e^{(-4*d*x - 4*c)} - 2*(435*a^5*b + 29*a^4*b^2 + 162*a^3*b^3 - 306*a^2*b^4 - 245*a*b^5 - 75*b^6))*e^{(-6*d*x - 6*c)} + 3*(145*a^5*b + 17*a^4*b^2 - 58*a^3*b^3 + 150*a^2*b^4 + 105*a*b^5 + 25*b^6))*e^{(-8*d*x - 8*c)} - 3*(29*a^5*b + 23*a^4*b^2 - 62*a^3*b^3 - 82*a^2*b^4 - 31*a*b^5 - 5*b^6))*e^{(-10*d*x - 10*c)})/((a^10 + 7*a^9*b + 21*a^8*b^2 + 35*a^7*b^3 + 35*a^6*b^4 + 21*a^5*b^5 + 7*a^4*b^6 + a^3*b^7 - 6*(a^10 + 5*a^9*b + 9*a^8*b^2 + 5*a^7*b^3 - 5*a^6*b^4 - 9*a^5*b^5 - 5*a^4*b^6 - a^3*b^7))*e^{(-2*d*x - 2*c)} + 3*(5*a^10 + 19*a^9*b + 25*a^8*b^2 + 15*a^7*b^3 + 15*a^6*b^4 + 25*a^5*b^5 + 19*a^4*b^6 + 5*a^3*b^7))*e^{(-4*d*x - 4*c)} - 4*(5*a^10 + 17*a^9*b + 21*a^8*b^2 + 9*a^7*b^3 - 9*a^6*b^4 - 21*a^5*b^5 - 17*a^4*b^6 - 5*a^3*b^7))*e^{(-6*d*x - 6*c)} + 3*(5*a^10 + 19*a^9*b + 25*a^8*b^2 + 15*a^7*b^3 + 15*a^6*b^4 + 25*a^5*b^5 + 19*a^4*b^6 + 5*a^3*b^7))*e^{(-8*d*x - 8*c)} - 6*(a^10 + 5*a^9*b + 9*a^8*b^2 + 5*a^7*b^3 - 5*a^6*b^4 - 9*a^5*b^5 - 5*a^4*b^6 - a^3*b^7))*e^{(-10*d*x - 10*c)} + (a^10 + 7*a^9*b + 21*a^8*b^2 + 35*a^7*b^3 + 35*a^6*b^4 + 21*a^5*b^5 + 7*a^4*b^6 + a^3*b^7))*e^{(-1...$

3.8.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 750 vs. $2(185) = 370$.

Time = 0.36 (sec) , antiderivative size = 750, normalized size of antiderivative = 3.73

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \frac{3(35a^3b + 35a^2b^2 + 21ab^3 + 5b^4) \arctan\left(\frac{ae^{(2dx+2c)} + be^{(2dx+2c)} - a + b}{2\sqrt{ab}}\right)}{(a^7 + 4a^6b + 6a^5b^2 + 4a^4b^3 + a^3b^4)\sqrt{ab}} - \frac{48(dx+c)}{a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4} - \frac{2(87a^5be^{(10dx+10c)} + 69a^4b^2e^{(10dx+10c)})}{(a^7 + 4a^6b + 6a^5b^2 + 4a^4b^3 + a^3b^4)\sqrt{ab}}$$

input `integrate(1/(a+b*coth(d*x+c)^2)^4,x, algorithm="giac")`

3.8. $\int \frac{1}{(a+b \coth^2(c+dx))^4} dx$

output

```
-1/48*(3*(35*a^3*b + 35*a^2*b^2 + 21*a*b^3 + 5*b^4)*arctan(1/2*(a*e^(2*d*x
+ 2*c) + b*e^(2*d*x + 2*c) - a + b)/sqrt(a*b))/((a^7 + 4*a^6*b + 6*a^5*b^
2 + 4*a^4*b^3 + a^3*b^4)*sqrt(a*b)) - 48*(d*x + c)/(a^4 + 4*a^3*b + 6*a^2*
b^2 + 4*a*b^3 + b^4) - 2*(87*a^5*b*e^(10*d*x + 10*c) + 69*a^4*b^2*e^(10*d*
x + 10*c) - 186*a^3*b^3*e^(10*d*x + 10*c) - 246*a^2*b^4*e^(10*d*x + 10*c)
- 93*a*b^5*e^(10*d*x + 10*c) - 15*b^6*e^(10*d*x + 10*c) - 435*a^5*b*e^(8*d
*x + 8*c) - 51*a^4*b^2*e^(8*d*x + 8*c) + 174*a^3*b^3*e^(8*d*x + 8*c) - 450
*a^2*b^4*e^(8*d*x + 8*c) - 315*a*b^5*e^(8*d*x + 8*c) - 75*b^6*e^(8*d*x + 8
*c) + 870*a^5*b*e^(6*d*x + 6*c) + 58*a^4*b^2*e^(6*d*x + 6*c) + 324*a^3*b^3
*e^(6*d*x + 6*c) - 612*a^2*b^4*e^(6*d*x + 6*c) - 490*a*b^5*e^(6*d*x + 6*c)
- 150*b^6*e^(6*d*x + 6*c) - 870*a^5*b*e^(4*d*x + 4*c) - 558*a^4*b^2*e^(4*
d*x + 4*c) + 36*a^3*b^3*e^(4*d*x + 4*c) - 636*a^2*b^4*e^(4*d*x + 4*c) - 51
0*a*b^5*e^(4*d*x + 4*c) - 150*b^6*e^(4*d*x + 4*c) + 435*a^5*b*e^(2*d*x + 2
*c) + 801*a^4*b^2*e^(2*d*x + 2*c) + 102*a^3*b^3*e^(2*d*x + 2*c) - 534*a^2*
b^4*e^(2*d*x + 2*c) - 345*a*b^5*e^(2*d*x + 2*c) - 75*b^6*e^(2*d*x + 2*c) -
87*a^5*b - 319*a^4*b^2 - 450*a^3*b^3 - 306*a^2*b^4 - 103*a*b^5 - 15*b^6)/
((a^7 + 4*a^6*b + 6*a^5*b^2 + 4*a^4*b^3 + a^3*b^4)*(a*e^(4*d*x + 4*c) + b*
e^(4*d*x + 4*c) - 2*a*e^(2*d*x + 2*c) + 2*b*e^(2*d*x + 2*c) + a + b)^3))/d
```

3.8.9 Mupad [B] (verification not implemented)

Time = 2.81 (sec) , antiderivative size = 3685, normalized size of antiderivative = 18.33

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \text{Too large to display}$$

input `int(1/(a + b*coth(c + d*x)^2)^4,x)`

output

$$\begin{aligned} & \log(\coth(c + dx) + 1)/(2a^4d + 2b^4d + 12a^2b^2d + 8ab^3d + 8a^3bd) + ((\coth(c + dx))^3(16ab^3 + 5b^4 + 17a^2b^2))/(6a^2(3ab^2 + 3a^2b + a^3 + b^3)) + (\coth(c + dx)(32ab^2 + 29a^2b + 11b^3))/(16a(3ab^2 + 3a^2b + a^3 + b^3)) + (b^2\coth(c + dx)^5(16ab^2 + 19a^2b + 5b^3))/(16a^2(ab^3 + 3a^3b + a^4 + 3a^2b^2))/(a^3d + b^3d\coth(c + dx)^6 + 3a^2b^2d\coth(c + dx)^2 + 3ab^2d\coth(c + dx)^4) - \log(\coth(c + dx) - 1)/(2d(a + b)^4) - (\operatorname{atan}(((-a^7b)^{1/2})(\coth(c + dx)(210ab^8 + 25b^9 + 791a^2b^7 + 1820a^3b^6 + 2695a^4b^5 + 2450a^5b^4 + 1481a^6b^3)))/(128(a^{12}d^2 + 6a^{11}bd^2 + a^6b^6d^2 + 6a^7b^5d^2 + 15a^8b^4d^2 + 20a^9b^3d^2 + 15a^{10}b^2d^2))) + (((5a^3b^{13}d^2)/4 + 14a^4b^{12}d^2 + (287a^5b^{11}d^2)/4 + 224a^6b^{10}d^2 + (953a^7b^9d^2)/2 + 728a^8b^8d^2 + (1631a^9b^7d^2)/2 + 668a^{10}b^6d^2 + (1561a^{11}b^5d^2)/4 + 154a^{12}b^4d^2 + (147a^{13}b^3d^2)/4 + 4a^{14}b^2d^2)/(a^{15}d^3 + 9a^{14}bd^3 + a^6b^9d^3 + 9a^7b^8d^3 + 36a^8b^7d^3 + 84a^9b^6d^3 + 126a^{10}b^5d^3 + 126a^{11}b^4d^3 + 84a^{12}b^3d^3 + 36a^{13}b^2d^3) - (\coth(c + dx)(-a^7b)^{(1/2})(21ab^2 + 35a^2b + 35a^3 + 5b^3)(1024a^6b^{11}d^2 + 7168a^7b^{10}d^2 + 20480a^8b^9d^2 + 28672a^9b^8d^2 + 14336a^{10}b^7d^2 - 14336a^{11}b^6d^2 - 28672a^{12}b^5d^2 - 20480a^{13}b^4d^2 - 7168a^{14}b^3d^2 - 1024a^{15}b^2d^2))/(4096(a^{11}d + a^7b^4d + 4a^8b^3d + 6\dots \end{aligned}$$

3.9 $\int \frac{1}{1-2 \coth^2(x)} dx$

3.9.1	Optimal result	102
3.9.2	Mathematica [A] (verified)	102
3.9.3	Rubi [A] (verified)	103
3.9.4	Maple [A] (verified)	104
3.9.5	Fricas [B] (verification not implemented)	105
3.9.6	Sympy [A] (verification not implemented)	105
3.9.7	Maxima [B] (verification not implemented)	106
3.9.8	Giac [B] (verification not implemented)	106
3.9.9	Mupad [B] (verification not implemented)	106

3.9.1 Optimal result

Integrand size = 10, antiderivative size = 19

$$\int \frac{1}{1-2 \coth^2(x)} dx = -x + \sqrt{2} \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)$$

output `-x+arctanh(1/2*2^(1/2)*tanh(x))*2^(1/2)`

3.9.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-2 \coth^2(x)} dx = -x + \sqrt{2} \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)$$

input `Integrate[(1 - 2*Coth[x]^2)^(-1),x]`

output `-x + Sqrt[2]*ArcTanh[Tanh[x]/Sqrt[2]]`

3.9.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3042, 4143, 25, 3042, 25, 4158, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{1 - 2 \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{1 + 2 \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{4143} \\
 & 2 \int -\frac{\operatorname{csch}^2(x)}{1 - 2 \coth^2(x)} dx - x \\
 & \quad \downarrow \text{25} \\
 & -2 \int \frac{\operatorname{csch}^2(x)}{1 - 2 \coth^2(x)} dx - x \\
 & \quad \downarrow \text{3042} \\
 & -x - 2 \int -\frac{\sec\left(ix + \frac{\pi}{2}\right)^2}{2 \tan\left(ix + \frac{\pi}{2}\right)^2 + 1} dx \\
 & \quad \downarrow \text{25} \\
 & -x + 2 \int \frac{\sec\left(ix + \frac{\pi}{2}\right)^2}{2 \tan\left(ix + \frac{\pi}{2}\right)^2 + 1} dx \\
 & \quad \downarrow \text{4158} \\
 & 2 \int \frac{1}{1 - 2 \coth^2(x)} d \coth(x) - x \\
 & \quad \downarrow \text{219} \\
 & \sqrt{2} \operatorname{arctanh}\left(\sqrt{2} \coth(x)\right) - x
 \end{aligned}$$

input `Int[(1 - 2*Coth[x]^2)^(-1),x]`

output $-x + \text{Sqrt}[2] \cdot \text{ArcTanh}[\text{Sqrt}[2] \cdot \text{Coth}[x]]$

3.9.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 219 $\text{Int}[(\text{a}_) + (\text{b}_) \cdot (\text{x}_)^2]^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] \cdot \text{Rt}[-\text{b}, 2])) \cdot \text{ArcTanh}[\text{Rt}[-\text{b}, 2] \cdot (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /;$ $\text{FreeQ}\{\{\text{a}, \text{b}\}, \text{x}\} \ \&\& \ \text{NegQ}[\text{a}/\text{b}] \ \&\& \ (\text{GtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$

rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /;$ $\text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 4143 $\text{Int}[(\text{a}_) + (\text{b}_) \cdot \tan[(\text{e}_) + (\text{f}_) \cdot (\text{x}_)]^2]^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{x}/(\text{a} - \text{b}), \text{x}] - \text{Simp}[\text{b}/(\text{a} - \text{b}) \text{Int}[\text{Sec}[\text{e} + \text{f} \cdot \text{x}]^2/(\text{a} + \text{b} \cdot \text{Tan}[\text{e} + \text{f} \cdot \text{x}]^2), \text{x}], \text{x}] /;$ $\text{FreeQ}\{\{\text{a}, \text{b}, \text{e}, \text{f}\}, \text{x}\} \ \&\& \ \text{NeQ}[\text{a}, \text{b}]$

rule 4158 $\text{Int}[\text{sec}[(\text{e}_) + (\text{f}_) \cdot (\text{x}_)]^m] \cdot ((\text{a}_) + (\text{b}_) \cdot ((\text{c}_) \cdot \tan[(\text{e}_) + (\text{f}_) \cdot (\text{x}_)]^n))^p, \text{x_Symbol}] \rightarrow \text{With}\{\{\text{ff} = \text{FreeFactors}[\text{Tan}[\text{e} + \text{f} \cdot \text{x}], \text{x}]\}, \text{Simp}[\text{ff}/(\text{c}^{m-1} \cdot \text{f}) \text{Subst}[\text{Int}[(\text{c}^2 + \text{ff}^2 \cdot \text{x}^2)^{m/2-1} \cdot (\text{a} + \text{b} \cdot (\text{ff} \cdot \text{x})^n)^p, \text{x}], \text{x}, \text{c} \cdot (\text{Tan}[\text{e} + \text{f} \cdot \text{x}]/\text{ff})], \text{x}]\} /;$ $\text{FreeQ}\{\{\text{a}, \text{b}, \text{c}, \text{e}, \text{f}, \text{n}, \text{p}\}, \text{x}\} \ \&\& \ \text{IntegerQ}[m/2] \ \&\& \ (\text{IntegersQ}[\text{n}, \text{p}] \ || \ \text{IGtQ}[m, 0] \ || \ \text{IGtQ}[p, 0] \ || \ \text{EqQ}[\text{n}^2, 4] \ || \ \text{EqQ}[\text{n}^2, 16])$

3.9.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

method	result	size
derivativedivides	$-\frac{\ln(1+\coth(x))}{2} + \frac{\ln(\coth(x)-1)}{2} + \sqrt{2} \operatorname{arctanh}(\coth(x)\sqrt{2})$	27
default	$-\frac{\ln(1+\coth(x))}{2} + \frac{\ln(\coth(x)-1)}{2} + \sqrt{2} \operatorname{arctanh}(\coth(x)\sqrt{2})$	27
risch	$-x + \frac{\sqrt{2} \ln(e^{2x}+3-2\sqrt{2})}{2} - \frac{\sqrt{2} \ln(e^{2x}+3+2\sqrt{2})}{2}$	39

input `int(1/(1-2*coth(x)^2),x,method=_RETURNVERBOSE)`

output `-1/2*ln(1+coth(x))+1/2*ln(coth(x)-1)+2^(1/2)*arctanh(coth(x)*2^(1/2))`

3.9.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(16) = 32$.

Time = 0.27 (sec) , antiderivative size = 70, normalized size of antiderivative = 3.68

$$\int \frac{1}{1-2\coth^2(x)} dx$$

$$= \frac{1}{2} \sqrt{2} \log \left(-\frac{3(2\sqrt{2}-3)\cosh(x)^2 - 4(3\sqrt{2}-4)\cosh(x)\sinh(x) + 3(2\sqrt{2}-3)\sinh(x)^2 + 2\sqrt{2}-3}{\cosh(x)^2 + \sinh(x)^2 + 3} \right) - x$$

input `integrate(1/(1-2*coth(x)^2),x, algorithm="fricas")`

output `1/2*sqrt(2)*log(-(3*(2*sqrt(2) - 3)*cosh(x)^2 - 4*(3*sqrt(2) - 4)*cosh(x)*sinh(x) + 3*(2*sqrt(2) - 3)*sinh(x)^2 + 2*sqrt(2) - 3)/(cosh(x)^2 + sinh(x)^2 + 3)) - x`

3.9.6 Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.79

$$\int \frac{1}{1-2\coth^2(x)} dx = -x - \frac{\sqrt{2} \log(\tanh(x) - \sqrt{2})}{2} + \frac{\sqrt{2} \log(\tanh(x) + \sqrt{2})}{2}$$

input `integrate(1/(1-2*coth(x)**2),x)`

output `-x - sqrt(2)*log(tanh(x) - sqrt(2))/2 + sqrt(2)*log(tanh(x) + sqrt(2))/2`

3.9.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(16) = 32$.

Time = 0.28 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.00

$$\int \frac{1}{1 - 2 \coth^2(x)} dx = -\frac{1}{2} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(-2x)} - 3}{2\sqrt{2} + e^{(-2x)} + 3} \right) - x$$

input `integrate(1/(1-2*coth(x)^2),x, algorithm="maxima")`

output `-1/2*sqrt(2)*log(-(2*sqrt(2) - e^(-2*x) - 3)/(2*sqrt(2) + e^(-2*x) + 3)) - x`

3.9.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(16) = 32$.

Time = 0.27 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.00

$$\int \frac{1}{1 - 2 \coth^2(x)} dx = \frac{1}{2} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(2x)} - 3}{2\sqrt{2} + e^{(2x)} + 3} \right) - x$$

input `integrate(1/(1-2*coth(x)^2),x, algorithm="giac")`

output `1/2*sqrt(2)*log(-(2*sqrt(2) - e^(2*x) - 3)/(2*sqrt(2) + e^(2*x) + 3)) - x`

3.9.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{1 - 2 \coth^2(x)} dx = \sqrt{2} \operatorname{atanh}(\sqrt{2} \coth(x)) - x$$

input `int(-1/(2*coth(x)^2 - 1),x)`

output `2^(1/2)*atanh(2^(1/2)*coth(x)) - x`

3.10 $\int \sqrt{1 - \coth^2(x)} dx$

3.10.1	Optimal result	107
3.10.2	Mathematica [B] (verified)	107
3.10.3	Rubi [A] (verified)	108
3.10.4	Maple [A] (verified)	109
3.10.5	Fricas [B] (verification not implemented)	110
3.10.6	Sympy [F]	110
3.10.7	Maxima [C] (verification not implemented)	110
3.10.8	Giac [C] (verification not implemented)	111
3.10.9	Mupad [B] (verification not implemented)	111

3.10.1 Optimal result

Integrand size = 12, antiderivative size = 3

$$\int \sqrt{1 - \coth^2(x)} dx = \arcsin(\coth(x))$$

output `arcsin(coth(x))`

3.10.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 30 vs. $2(3) = 6$.

Time = 0.05 (sec) , antiderivative size = 30, normalized size of antiderivative = 10.00

$$\int \sqrt{1 - \coth^2(x)} dx = \sqrt{-\operatorname{csch}^2(x)} \left(-\log \left(\cosh \left(\frac{x}{2} \right) \right) + \log \left(\sinh \left(\frac{x}{2} \right) \right) \right) \sinh(x)$$

input `Integrate[Sqrt[1 - Coth[x]^2], x]`

output `Sqrt[-Csch[x]^2]*(-Log[Cosh[x/2]] + Log[Sinh[x/2]])*Sinh[x]`

3.10.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 4140, 3042, 4610, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{1 - \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{1 + \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \sqrt{-\operatorname{csch}^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\sec\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{4610} \\
 & \int \frac{1}{\sqrt{1 - \coth^2(x)}} d \coth(x) \\
 & \quad \downarrow \text{223} \\
 & \arcsin(\coth(x))
 \end{aligned}$$

input `Int[Sqrt[1 - Coth[x]^2],x]`

output `ArcSin[Coth[x]]`

3.10.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

3.10.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	$\arcsin(\coth(x))$	4
default	$\arcsin(\coth(x))$	4
risch	$\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1) \ln(e^x-1) - \sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1) \ln(e^x+1)$	67

input `int((1-coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsin(coth(x))`

3.10.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(3) = 6$.

Time = 0.25 (sec) , antiderivative size = 45, normalized size of antiderivative = 15.00

$$\int \sqrt{1 - \coth^2(x)} dx = -2 \arctan \left(\frac{\sqrt{-\frac{e^{(2x)}}{e^{(4x)} - 2e^{(2x)} + 1}} (e^{(2x)} - 1)}{\cosh(x) e^x + e^x \sinh(x)} \right)$$

input `integrate((1-coth(x)^2)^(1/2),x, algorithm="fracas")`

output `-2*arctan(sqrt(-e^(2*x)/(e^(4*x) - 2*e^(2*x) + 1))*(e^(2*x) - 1)/(cosh(x)*e^x + e^x*sinh(x)))`

3.10.6 Sympy [F]

$$\int \sqrt{1 - \coth^2(x)} dx = \int \sqrt{1 - \coth^2(x)} dx$$

input `integrate((1-coth(x)**2)**(1/2),x)`

output `Integral(sqrt(1 - coth(x)**2), x)`

3.10.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 6.33

$$\int \sqrt{1 - \coth^2(x)} dx = i \log(e^{-x} + 1) - i \log(e^{-x} - 1)$$

input `integrate((1-coth(x)^2)^(1/2),x, algorithm="maxima")`

output `I*log(e^(-x) + 1) - I*log(e^(-x) - 1)`

3.10. $\int \sqrt{1 - \coth^2(x)} dx$

3.10.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 8.67

$$\int \sqrt{1 - \coth^2(x)} dx = (i \log(e^x + 1) - i \log(|e^x - 1|)) \operatorname{sgn}(-e^{(2x)} + 1)$$

input `integrate((1-coth(x)^2)^(1/2),x, algorithm="giac")`

output `(I*log(e^x + 1) - I*log(abs(e^x - 1)))*sgn(-e^(2*x) + 1)`

3.10.9 Mupad [B] (verification not implemented)

Time = 1.90 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \sqrt{1 - \coth^2(x)} dx = \operatorname{asin}(\coth(x))$$

input `int((1 - coth(x)^2)^(1/2),x)`

output `asin(coth(x))`

3.11 $\int \sqrt{-1 + \coth^2(x)} dx$

3.11.1	Optimal result	112
3.11.2	Mathematica [A] (verified)	112
3.11.3	Rubi [A] (verified)	113
3.11.4	Maple [A] (verified)	114
3.11.5	Fricas [A] (verification not implemented)	115
3.11.6	Sympy [F]	115
3.11.7	Maxima [A] (verification not implemented)	115
3.11.8	Giac [A] (verification not implemented)	116
3.11.9	Mupad [B] (verification not implemented)	116

3.11.1 Optimal result

Integrand size = 10, antiderivative size = 14

$$\int \sqrt{-1 + \coth^2(x)} dx = -\operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}\right)$$

output `-arctanh(coth(x)/(csch(x)^2)^(1/2))`

3.11.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.00

$$\int \sqrt{-1 + \coth^2(x)} dx = \sqrt{\operatorname{csch}^2(x)} \left(-\log\left(\cosh\left(\frac{x}{2}\right)\right) + \log\left(\sinh\left(\frac{x}{2}\right)\right) \right) \sinh(x)$$

input `Integrate[Sqrt[-1 + Coth[x]^2], x]`

output `Sqrt[Csch[x]^2]*(-Log[Cosh[x/2]] + Log[Sinh[x/2]])*Sinh[x]`

3.11.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 4140, 3042, 4610, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\coth^2(x) - 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-1 - \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \sqrt{\operatorname{csch}^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-\sec\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{4610} \\
 & - \int \frac{1}{\sqrt{\coth^2(x) - 1}} d \coth(x) \\
 & \quad \downarrow \text{224} \\
 & - \int \frac{1}{1 - \frac{\coth^2(x)}{\coth^2(x) - 1}} d \frac{\coth(x)}{\sqrt{\coth^2(x) - 1}} \\
 & \quad \downarrow \text{219} \\
 & -\operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\coth^2(x) - 1}}\right)
 \end{aligned}$$

input `Int[Sqrt[-1 + Coth[x]^2], x]`

output `-ArcTanh[Coth[x]/Sqrt[-1 + Coth[x]^2]]`

3.11. $\int \sqrt{-1 + \coth^2(x)} dx$

3.11.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

3.11.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
derivativedivides	$-\ln\left(\coth(x) + \sqrt{\coth(x)^2 - 1}\right)$	15
default	$-\ln\left(\coth(x) + \sqrt{\coth(x)^2 - 1}\right)$	15
risch	$\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1) \ln(e^x-1) - \sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1) \ln(e^x+1)$	65

input `int((coth(x)^2-1)^(1/2),x,method=_RETURNVERBOSE)`

output `-ln(coth(x)+(coth(x)^2-1)^(1/2))`

3.11. $\int \sqrt{-1 + \coth^2(x)} dx$

3.11.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \sqrt{-1 + \coth^2(x)} dx = -\log(\cosh(x) + \sinh(x) + 1) + \log(\cosh(x) + \sinh(x) - 1)$$

input `integrate((-1+coth(x)^2)^(1/2),x, algorithm="fricas")`

output `-log(cosh(x) + sinh(x) + 1) + log(cosh(x) + sinh(x) - 1)`

3.11.6 Sympy [F]

$$\int \sqrt{-1 + \coth^2(x)} dx = \int \sqrt{\coth^2(x) - 1} dx$$

input `integrate((-1+coth(x)**2)**(1/2),x)`

output `Integral(sqrt(coth(x)**2 - 1), x)`

3.11.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \sqrt{-1 + \coth^2(x)} dx = \log(e^{-x} + 1) - \log(e^{-x} - 1)$$

input `integrate((-1+coth(x)^2)^(1/2),x, algorithm="maxima")`

output `log(e^(-x) + 1) - log(e^(-x) - 1)`

3.11.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.64

$$\int \sqrt{-1 + \coth^2(x)} dx = -(\log(e^x + 1) - \log(|e^x - 1|))\operatorname{sgn}(e^{2x} - 1)$$

input `integrate((-1+coth(x)^2)^(1/2),x, algorithm="giac")`

output `-(log(e^x + 1) - log(abs(e^x - 1)))*sgn(e^(2*x) - 1)`

3.11.9 Mupad [B] (verification not implemented)

Time = 2.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sqrt{-1 + \coth^2(x)} dx = -\ln\left(\coth(x) + \sqrt{\coth(x)^2 - 1}\right)$$

input `int((coth(x)^2 - 1)^(1/2),x)`

output `-log(coth(x) + (coth(x)^2 - 1)^(1/2))`

3.12 $\int (1 - \coth^2(x))^{3/2} dx$

3.12.1 Optimal result	117
3.12.2 Mathematica [B] (verified)	117
3.12.3 Rubi [A] (verified)	118
3.12.4 Maple [A] (verified)	119
3.12.5 Fricas [B] (verification not implemented)	120
3.12.6 Sympy [F]	120
3.12.7 Maxima [C] (verification not implemented)	121
3.12.8 Giac [C] (verification not implemented)	121
3.12.9 Mupad [B] (verification not implemented)	122

3.12.1 Optimal result

Integrand size = 12, antiderivative size = 24

$$\int (1 - \coth^2(x))^{3/2} dx = \frac{1}{2} \arcsin(\coth(x)) + \frac{1}{2} \coth(x) \sqrt{-\operatorname{csch}^2(x)}$$

output `1/2*arcsin(coth(x))+1/2*coth(x)*(-csch(x)^2)^(1/2)`

3.12.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs. $2(24) = 48$.

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.12

$$\int (1 - \coth^2(x))^{3/2} dx = \frac{1}{8} \sqrt{-\operatorname{csch}^2(x)} \left(\operatorname{csch}^2\left(\frac{x}{2}\right) - 4 \log\left(\cosh\left(\frac{x}{2}\right)\right) + 4 \log\left(\sinh\left(\frac{x}{2}\right)\right) + \operatorname{sech}^2\left(\frac{x}{2}\right) \right) \sinh(x)$$

input `Integrate[(1 - Coth[x]^2)^(3/2), x]`

output `(Sqrt[-Csch[x]^2]*(Csch[x/2]^2 - 4*Log[Cosh[x/2]] + 4*Log[Sinh[x/2]] + Sec h[x/2]^2)*Sinh[x])/8`

3.12.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4140, 3042, 4610, 211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (1 - \coth^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(1 + \tan\left(\frac{\pi}{2} + ix\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4140} \\
 & \int (-\operatorname{csch}^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(\sec\left(\frac{\pi}{2} + ix\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4610} \\
 & \int \sqrt{1 - \coth^2(x)} d\coth(x) \\
 & \quad \downarrow \text{211} \\
 & \frac{1}{2} \int \frac{1}{\sqrt{1 - \coth^2(x)}} d\coth(x) + \frac{1}{2} \sqrt{1 - \coth^2(x)} \coth(x) \\
 & \quad \downarrow \text{223} \\
 & \frac{1}{2} \arcsin(\coth(x)) + \frac{1}{2} \coth(x) \sqrt{1 - \coth^2(x)}
 \end{aligned}$$

input `Int[(1 - Coth[x]^2)^(3/2), x]`

output `ArcSin[Coth[x]]/2 + (Coth[x]*Sqrt[1 - Coth[x]^2])/2`

3.12.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

3.12.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\coth(x)\sqrt{1-\coth(x)^2}}{2} + \frac{\arcsin(\coth(x))}{2}$	21
default	$\frac{\coth(x)\sqrt{1-\coth(x)^2}}{2} + \frac{\arcsin(\coth(x))}{2}$	21
risch	$\frac{\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}(1+e^{2x})}{e^{2x}-1} - \frac{\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}e^{-x}(e^{2x}-1)\ln(e^x+1)}{2} + \frac{\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}e^{-x}(e^{2x}-1)\ln(e^x-1)}{2}$	99

input `int((1-coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `1/2*coth(x)*(1-coth(x)^2)^(1/2)+1/2*arcsin(coth(x))`

3.12. $\int (1 - \coth^2(x))^{3/2} dx$

3.12.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 279 vs. $2(18) = 36$.

Time = 0.28 (sec) , antiderivative size = 279, normalized size of antiderivative = 11.62

$$\int (1 - \coth^2(x))^{3/2} dx =$$

$$(4 \cosh(x) e^x \sinh(x)^3 + e^x \sinh(x)^4 + 2(3 \cosh(x)^2 - 1)e^x \sinh(x)^2 + 4(\cosh(x)^3 - \cosh(x))e^x \sinh(x) - \dots)$$

input `integrate((1-coth(x)^2)^(3/2),x, algorithm="fricas")`

output `-((4*cosh(x)*e^x*sinh(x)^3 + e^x*sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*e^x*sinh(x)^2 + 4*(cosh(x)^3 - cosh(x))*e^x*sinh(x) + (cosh(x)^4 - 2*cosh(x)^2 + 1)*e^x)*arctan(sqrt(-e^(2*x))/(e^(4*x) - 2*e^(2*x) + 1))*(e^(2*x) - 1)/(cosh(x)*e^x + e^x*sinh(x))) - ((e^(2*x) - 1)*sinh(x)^3 - cosh(x)^3 + 3*(cosh(x)*e^(2*x) - cosh(x))*sinh(x)^2 + (cosh(x)^3 + cosh(x))*e^(2*x) - (3*cosh(x)^2 - (3*cosh(x)^2 + 1)*e^(2*x) + 1)*sinh(x) - cosh(x))*sqrt(-e^(2*x)/(e^(4*x) - 2*e^(2*x) + 1)))/(4*cosh(x)*e^x*sinh(x)^3 + e^x*sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*e^x*sinh(x)^2 + 4*(cosh(x)^3 - cosh(x))*e^x*sinh(x) + (cosh(x)^4 - 2*cosh(x)^2 + 1)*e^x)`

3.12.6 Sympy [F]

$$\int (1 - \coth^2(x))^{3/2} dx = \int (1 - \coth^2(x))^{\frac{3}{2}} dx$$

input `integrate((1-coth(x)**2)**(3/2),x)`

output `Integral((1 - coth(x)**2)**(3/2), x)`

3.12.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.04

$$\int (1 - \coth^2(x))^{3/2} dx = \frac{i e^{(-x)} + i e^{(-3x)}}{2 e^{(-2x)} - e^{(-4x)} - 1} + \frac{1}{2} i \log(e^{(-x)} + 1) - \frac{1}{2} i \log(e^{(-x)} - 1)$$

input `integrate((1-coth(x)^2)^(3/2),x, algorithm="maxima")`

output `(I*e^(-x) + I*e^(-3*x))/(2*e^(-2*x) - e^(-4*x) - 1) + 1/2*I*log(e^(-x) + 1) - 1/2*I*log(e^(-x) - 1)`

3.12.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.50

$$\int (1 - \coth^2(x))^{3/2} dx = -\frac{1}{4} \left(\frac{4(i e^{(-x)} + i e^x)}{(e^{(-x)} + e^x)^2 - 4} - i \log(e^{(-x)} + e^x + 2) + i \log(e^{(-x)} + e^x - 2) \right) \operatorname{sgn}(-e^{(2x)} + 1)$$

input `integrate((1-coth(x)^2)^(3/2),x, algorithm="giac")`

output `-1/4*(4*(I*e^(-x) + I*e^x)/((e^(-x) + e^x)^2 - 4) - I*log(e^(-x) + e^x + 2) + I*log(e^(-x) + e^x - 2))*sgn(-e^(2*x) + 1)`

3.12.9 Mupad [B] (verification not implemented)

Time = 1.95 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int (1 - \operatorname{coth}^2(x))^{3/2} dx = \frac{\operatorname{asin}(\operatorname{coth}(x))}{2} + \frac{\operatorname{coth}(x) \sqrt{1 - \operatorname{coth}(x)^2}}{2}$$

input `int((1 - coth(x)^2)^(3/2),x)`

output `asin(coth(x))/2 + (coth(x)*(1 - coth(x)^2)^(1/2))/2`

3.13 $\int (-1 + \coth^2(x))^{3/2} dx$

3.13.1 Optimal result	123
3.13.2 Mathematica [A] (verified)	123
3.13.3 Rubi [A] (verified)	124
3.13.4 Maple [A] (verified)	126
3.13.5 Fricas [B] (verification not implemented)	126
3.13.6 Sympy [F]	127
3.13.7 Maxima [A] (verification not implemented)	127
3.13.8 Giac [B] (verification not implemented)	127
3.13.9 Mupad [B] (verification not implemented)	128

3.13.1 Optimal result

Integrand size = 10, antiderivative size = 31

$$\int (-1 + \coth^2(x))^{3/2} dx = \frac{1}{2} \operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{\operatorname{csch}^2(x)}$$

output `1/2*arctanh(coth(x)/(csch(x)^2)^(1/2))-1/2*coth(x)*(csch(x)^2)^(1/2)`

3.13.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int (-1 + \coth^2(x))^{3/2} dx = -\frac{1}{8} \sqrt{\operatorname{csch}^2(x)} \left(\operatorname{csch}^2\left(\frac{x}{2}\right) - 4 \log\left(\cosh\left(\frac{x}{2}\right)\right) + 4 \log\left(\sinh\left(\frac{x}{2}\right)\right) + \operatorname{sech}^2\left(\frac{x}{2}\right) \right) \sinh(x)$$

input `Integrate[(-1 + Coth[x]^2)^(3/2), x]`

output `-1/8*(Sqrt[Csch[x]^2]*(Csch[x/2]^2 - 4*Log[Cosh[x/2]] + 4*Log[Sinh[x/2]] + Sech[x/2]^2)*Sinh[x])`

3.13.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3042, 4140, 3042, 4610, 211, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (\coth^2(x) - 1)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(-1 - \tan\left(\frac{\pi}{2} + ix\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \operatorname{csch}^2(x)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(-\sec\left(\frac{\pi}{2} + ix\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4610} \\
 & - \int \sqrt{\coth^2(x) - 1} d\coth(x) \\
 & \quad \downarrow \text{211} \\
 & \frac{1}{2} \int \frac{1}{\sqrt{\coth^2(x) - 1}} d\coth(x) - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) - 1} \\
 & \quad \downarrow \text{224} \\
 & \frac{1}{2} \int \frac{1}{1 - \frac{\coth^2(x)}{\coth^2(x) - 1}} d\frac{\coth(x)}{\sqrt{\coth^2(x) - 1}} - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) - 1} \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{2} \operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\coth^2(x) - 1}}\right) - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) - 1}
 \end{aligned}$$

input `Int[(-1 + Coth[x]^2)^(3/2), x]`

3.13. $\int (-1 + \coth^2(x))^{3/2} dx$

output `ArcTanh[Coth[x]/Sqrt[-1 + Coth[x]^2]]/2 - (Coth[x]*Sqrt[-1 + Coth[x]^2])/2`

3.13.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

3.13.6 Sympy [F]

$$\int (-1 + \coth^2(x))^{3/2} dx = \int (\coth^2(x) - 1)^{\frac{3}{2}} dx$$

input `integrate((-1+coth(x)**2)**(3/2),x)`

output `Integral((coth(x)**2 - 1)**(3/2), x)`

3.13.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.48

$$\int (-1 + \coth^2(x))^{3/2} dx = -\frac{e^{-x} + e^{-3x}}{2e^{-2x} - e^{-4x} - 1} - \frac{1}{2} \log(e^{-x} + 1) + \frac{1}{2} \log(e^{-x} - 1)$$

input `integrate((-1+coth(x)^2)^(3/2),x, algorithm="maxima")`

output `-(e^(-x) + e^(-3*x))/(2*e^(-2*x) - e^(-4*x) - 1) - 1/2*log(e^(-x) + 1) + 1/2*log(e^(-x) - 1)`

3.13.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(23) = 46$.

Time = 0.27 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.68

$$\int (-1 + \coth^2(x))^{3/2} dx = -\frac{1}{4} \left(\frac{4(e^{-x} + e^x)}{(e^{-x} + e^x)^2 - 4} - \log(e^{-x} + e^x + 2) + \log(e^{-x} + e^x - 2) \right) \operatorname{sgn}(e^{2x} - 1)$$

input `integrate((-1+coth(x)^2)^(3/2),x, algorithm="giac")`

output `-1/4*(4*(e^(-x) + e^x)/((e^(-x) + e^x)^2 - 4) - log(e^(-x) + e^x + 2) + log(e^(-x) + e^x - 2))*sgn(e^(2*x) - 1)`

3.13. $\int (-1 + \coth^2(x))^{3/2} dx$

3.13.9 Mupad [B] (verification not implemented)

Time = 1.92 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int (-1 + \operatorname{coth}^2(x))^{3/2} dx = \frac{\ln\left(\operatorname{coth}(x) + \sqrt{\operatorname{coth}(x)^2 - 1}\right)}{2} - \frac{\operatorname{coth}(x) \sqrt{\operatorname{coth}(x)^2 - 1}}{2}$$

input `int((coth(x)^2 - 1)^(3/2),x)`

output `log(coth(x) + (coth(x)^2 - 1)^(1/2))/2 - (coth(x)*(coth(x)^2 - 1)^(1/2))/2`

3.14 $\int \frac{1}{\sqrt{1-\coth^2(x)}} dx$

3.14.1 Optimal result 129
 3.14.2 Mathematica [A] (verified) 129
 3.14.3 Rubi [A] (verified) 130
 3.14.4 Maple [A] (verified) 131
 3.14.5 Fricas [B] (verification not implemented) 132
 3.14.6 Sympy [F] 132
 3.14.7 Maxima [C] (verification not implemented) 132
 3.14.8 Giac [C] (verification not implemented) 133
 3.14.9 Mupad [B] (verification not implemented) 133

3.14.1 Optimal result

Integrand size = 12, antiderivative size = 13

$$\int \frac{1}{\sqrt{1-\coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{-\operatorname{csch}^2(x)}}$$

output `coth(x)/(-csch(x)^2)^(1/2)`

3.14.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-\coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{-\operatorname{csch}^2(x)}}$$

input `Integrate[1/Sqrt[1 - Coth[x]^2],x]`

output `Coth[x]/Sqrt[-Csch[x]^2]`

3.14.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 4140, 3042, 4610, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{1 - \coth^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{1 + \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \frac{1}{\sqrt{-\operatorname{csch}^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{\sec\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{4610} \\
 & \int \frac{1}{(1 - \coth^2(x))^{3/2}} d\coth(x) \\
 & \quad \downarrow \text{208} \\
 & \frac{\coth(x)}{\sqrt{1 - \coth^2(x)}}
 \end{aligned}$$

input `Int[1/Sqrt[1 - Coth[x]^2],x]`

output `Coth[x]/Sqrt[1 - Coth[x]^2]`

3.14.3.1 Defintions of rubi rules used

- rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`
- rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

3.14.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$\frac{\coth(x)}{\sqrt{1-\coth(x)^2}}$	14
default	$\frac{\coth(x)}{\sqrt{1-\coth(x)^2}}$	14
risch	$\frac{e^{2x}}{2\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}(e^{2x}-1)} + \frac{1}{2(e^{2x}-1)\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}}$	58

input `int(1/(1-coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `1/(1-coth(x)^2)^(1/2)*coth(x)`

3.14.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. $2(11) = 22$.

Time = 0.26 (sec) , antiderivative size = 40, normalized size of antiderivative = 3.08

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = -(\cosh(x) e^{(2x)} - \cosh(x)) \sqrt{-\frac{e^{(2x)}}{e^{(4x)} - 2e^{(2x)} + 1}} e^{(-x)}$$

input `integrate(1/(1-coth(x)^2)^(1/2),x, algorithm="fracas")`

output `-(cosh(x)*e^(2*x) - cosh(x))*sqrt(-e^(2*x)/(e^(4*x) - 2*e^(2*x) + 1))*e^(-x)`

3.14.6 Sympy [F]

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = \int \frac{1}{\sqrt{1 - \coth^2(x)}} dx$$

input `integrate(1/(1-coth(x)**2)**(1/2),x)`

output `Integral(1/sqrt(1 - coth(x)**2), x)`

3.14.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = \frac{1}{2}i e^{(-x)} + \frac{1}{2}i e^x$$

input `integrate(1/(1-coth(x)^2)^(1/2),x, algorithm="maxima")`

output `1/2*I*e^(-x) + 1/2*I*e^x`

3.14. $\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx$

3.14.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.85

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = -\frac{-i e^{(-x)} - i e^x}{2 \operatorname{sgn}(-e^{(2x)} + 1)}$$

input `integrate(1/(1-coth(x)^2)^(1/2),x, algorithm="giac")`

output `-1/2*(-I*e^(-x) - I*e^x)/sgn(-e^(2*x) + 1)`

3.14.9 Mupad [B] (verification not implemented)

Time = 2.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.38

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = -\cosh(x) \sinh(x) \sqrt{-\frac{1}{\cosh(x)^2 - 1}}$$

input `int(1/(1 - coth(x)^2)^(1/2),x)`

output `-cosh(x)*sinh(x)*(-1/(cosh(x)^2 - 1))^(1/2)`

$$3.15 \quad \int \frac{1}{\sqrt{-1+\coth^2(x)}} dx$$

3.15.1	Optimal result	134
3.15.2	Mathematica [A] (verified)	134
3.15.3	Rubi [A] (verified)	135
3.15.4	Maple [A] (verified)	136
3.15.5	Fricas [A] (verification not implemented)	137
3.15.6	Sympy [F]	137
3.15.7	Maxima [A] (verification not implemented)	137
3.15.8	Giac [A] (verification not implemented)	138
3.15.9	Mupad [B] (verification not implemented)	138

3.15.1 Optimal result

Integrand size = 10, antiderivative size = 11

$$\int \frac{1}{\sqrt{-1+\coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}$$

output `coth(x)/(csch(x)^2)^(1/2)`

3.15.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1+\coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}$$

input `Integrate[1/Sqrt[-1 + Coth[x]^2], x]`

output `Coth[x]/Sqrt[Csch[x]^2]`

3.15.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4140, 3042, 4610, 208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\coth^2(x) - 1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{-1 - \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \frac{1}{\sqrt{\operatorname{csch}^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{-\sec\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{4610} \\
 & - \int \frac{1}{(\coth^2(x) - 1)^{3/2}} d\coth(x) \\
 & \quad \downarrow \text{208} \\
 & \frac{\coth(x)}{\sqrt{\coth^2(x) - 1}}
 \end{aligned}$$

input `Int[1/Sqrt[-1 + Coth[x]^2],x]`

output `Coth[x]/Sqrt[-1 + Coth[x]^2]`

3.15.3.1 Defintions of rubi rules used

- rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`
- rule 4610 `Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[b*(ff/f) Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

3.15.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{\coth(x)}{\sqrt{\coth(x)^2 - 1}}$	12
default	$\frac{\coth(x)}{\sqrt{\coth(x)^2 - 1}}$	12
risch	$\frac{e^{2x}}{2\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}}(e^{2x}-1)} + \frac{1}{2(e^{2x}-1)\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}}}$	56

input `int(1/(coth(x)^2-1)^(1/2),x,method=_RETURNVERBOSE)`

output `coth(x)/(coth(x)^2-1)^(1/2)`

3.15.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.18

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \cosh(x)$$

input `integrate(1/(-1+coth(x)^2)^(1/2),x, algorithm="fracas")`output `cosh(x)`**3.15.6 Sympy [F]**

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \int \frac{1}{\sqrt{\coth^2(x) - 1}} dx$$

input `integrate(1/(-1+coth(x)**2)**(1/2),x)`output `Integral(1/sqrt(coth(x)**2 - 1), x)`**3.15.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = -\frac{1}{2} e^{(-x)} - \frac{1}{2} e^x$$

input `integrate(1/(-1+coth(x)^2)^(1/2),x, algorithm="maxima")`output `-1/2*e^(-x) - 1/2*e^x`

3.15.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.64

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \frac{e^{(-x)} + e^x}{2 \operatorname{sgn}(e^{(2x)} - 1)}$$

input `integrate(1/(-1+coth(x)^2)^(1/2),x, algorithm="giac")`

output `1/2*(e^(-x) + e^x)/sgn(e^(2*x) - 1)`

3.15.9 Mupad [B] (verification not implemented)

Time = 1.97 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \cosh(x) \sinh(x) \sqrt{\frac{1}{\cosh(x)^2 - 1}}$$

input `int(1/(coth(x)^2 - 1)^(1/2),x)`

output `cosh(x)*sinh(x)*(1/(cosh(x)^2 - 1))^(1/2)`

3.16 $\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx$

3.16.1	Optimal result	139
3.16.2	Mathematica [A] (verified)	139
3.16.3	Rubi [A] (verified)	140
3.16.4	Maple [B] (verified)	142
3.16.5	Fricas [B] (verification not implemented)	143
3.16.6	Sympy [F]	144
3.16.7	Maxima [F]	145
3.16.8	Giac [F(-2)]	145
3.16.9	Mupad [B] (verification not implemented)	145

3.16.1 Optimal result

Integrand size = 17, antiderivative size = 63

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a+b}} \right) - \sqrt{a + b \coth^2(x)} - \frac{(a + b \coth^2(x))^{3/2}}{3b}$$

output `-1/3*(a+b*coth(x)^2)^(3/2)/b+arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))*(a+b)^(1/2)-(a+b*coth(x)^2)^(1/2)`

3.16.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.95

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a+b}} \right) - \frac{\sqrt{a + b \coth^2(x)}(a + 3b + b \coth^2(x))}{3b}$$

input `Integrate[Coth[x]^3*Sqrt[a + b*Coth[x]^2],x]`

output `Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - (Sqrt[a + b*Coth[x]^2]*(a + 3*b + b*Coth[x]^2))/(3*b)`

3.16.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.08, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.529$, Rules used = {3042, 26, 4153, 26, 354, 90, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^3(x) \sqrt{a + b \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int i \tan\left(\frac{\pi}{2} + ix\right)^3 \sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \tan\left(ix + \frac{\pi}{2}\right)^3 \sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \coth^3(x) \sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth^3(x) \sqrt{a + b \coth^2(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\coth^2(x) \sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth^2(x) \\
 & \quad \downarrow \text{90} \\
 & \frac{1}{2} \left(\int \frac{\sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth^2(x) - \frac{2(a + b \coth^2(x))^{3/2}}{3b} \right) \\
 & \quad \downarrow \text{60}
 \end{aligned}$$

3.16. $\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx$

$$\frac{1}{2} \left((a+b) \int \frac{1}{(1-\coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - \frac{2(a+b \coth^2(x))^{3/2}}{3b} - 2\sqrt{a+b \coth^2(x)} \right)$$

↓ 73

$$\frac{1}{2} \left(\frac{2(a+b) \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d \sqrt{b \coth^2(x) + a}}{b} - \frac{2(a+b \coth^2(x))^{3/2}}{3b} - 2\sqrt{a+b \coth^2(x)} \right)$$

↓ 221

$$\frac{1}{2} \left(2\sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right) - \frac{2(a+b \coth^2(x))^{3/2}}{3b} - 2\sqrt{a+b \coth^2(x)} \right)$$

input `Int[Coth[x]^3*Sqrt[a + b*Coth[x]^2],x]`

output `(2*Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - 2*Sqrt[a + b*Coth[x]^2] - (2*(a + b*Coth[x]^2)^(3/2))/(3*b))/2`

3.16.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

- rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

- rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 4153 `Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_.))^p, x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.16.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(51) = 102.

Time = 0.21 (sec) , antiderivative size = 253, normalized size of antiderivative = 4.02

method	result
derivativedivides	$-\frac{(a+b \coth(x)^2)^{\frac{3}{2}}}{3b} - \frac{\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(\coth(x)-1)+b}{\sqrt{b}} + \sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}\right)}{2}$
default	$-\frac{(a+b \coth(x)^2)^{\frac{3}{2}}}{3b} - \frac{\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(\coth(x)-1)+b}{\sqrt{b}} + \sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}\right)}{2}$

3.16. $\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx$

```
input int(coth(x)^3*(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/3*(a+b*coth(x)^2)^(3/2)/b-1/2*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)-1/2*b^(1/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))+1/2*(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))-1/2*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)+1/2*b^(1/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))+1/2*(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))
```

3.16.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 903 vs. $2(51) = 102$.

Time = 0.36 (sec) , antiderivative size = 2367, normalized size of antiderivative = 37.57

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \text{Too large to display}$$

```
input integrate(coth(x)^3*(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")
```


output

```
[1/12*(3*(b*cosh(x)^6 + 6*b*cosh(x)*sinh(x)^5 + b*sinh(x)^6 - 3*b*cosh(x)^4 + 3*(5*b*cosh(x)^2 - b)*sinh(x)^4 + 4*(5*b*cosh(x)^3 - 3*b*cosh(x))*sinh(x)^3 + 3*b*cosh(x)^2 + 3*(5*b*cosh(x)^4 - 6*b*cosh(x)^2 + b)*sinh(x)^2 + 6*(b*cosh(x)^5 - 2*b*cosh(x)^3 + b*cosh(x))*sinh(x) - b)*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - ...
```

3.16.6 Sympy [F]

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} \coth^3(x) dx$$

input `integrate(coth(x)**3*(a+b*coth(x)**2)**(1/2),x)`

output `Integral(sqrt(a + b*coth(x)**2)*coth(x)**3, x)`

3.16.7 Maxima [F]

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth^2(x)^2 + a} \coth(x)^3 dx$$

input `integrate(coth(x)^3*(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*coth(x)^2 + a)*coth(x)^3, x)`

3.16.8 Giac [F(-2)]

Exception generated.

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)^3*(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Error: Bad Argument Type`

3.16.9 Mupad [B] (verification not implemented)

Time = 3.90 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.05

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = -\sqrt{b \coth^2(x)^2 + a} - \frac{(b \coth^2(x)^2 + a)^{3/2}}{3b} - 2 \operatorname{atan} \left(\frac{2 \sqrt{b \coth^2(x)^2 + a} \sqrt{-\frac{a}{4} - \frac{b}{4}}}{a + b} \right) \sqrt{-\frac{a}{4} - \frac{b}{4}}$$

input `int(coth(x)^3*(a + b*coth(x)^2)^(1/2),x)`

output `-(a + b*coth(x)^2)^(1/2) - (a + b*coth(x)^2)^(3/2)/(3*b) - 2*atan((2*(a + b*coth(x)^2)^(1/2)*(- a/4 - b/4)^(1/2))/(a + b))*(- a/4 - b/4)^(1/2)`

3.16. $\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx$

3.17 $\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx$

3.17.1	Optimal result	146
3.17.2	Mathematica [B] (verified)	146
3.17.3	Rubi [A] (verified)	147
3.17.4	Maple [B] (verified)	150
3.17.5	Fricas [B] (verification not implemented)	151
3.17.6	Sympy [F]	151
3.17.7	Maxima [F]	152
3.17.8	Giac [F(-2)]	152
3.17.9	Mupad [F(-1)]	152

3.17.1 Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = -\frac{(a + 2b) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{2\sqrt{b}} + \sqrt{a + b} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

```
output -1/2*(a+2*b)*arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))/b^(1/2)+arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))*(a+b)^(1/2)-1/2*coth(x)*(a+b*coth(x)^2)^(1/2)
```

3.17.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 191 vs. 2(85) = 170.

Time = 0.84 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.25

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \frac{\sqrt{(-a + b + (a + b) \cosh(2x)) \operatorname{csch}^2(x)} \left(\sqrt{2} \sqrt{a + b} (a + 2b) \operatorname{arctanh}\left(\frac{\sqrt{2} \sqrt{b} \cosh(x)}{\sqrt{-a + b + (a + b) \cosh(2x)}}\right) + \sqrt{b} (-2\sqrt{2} \sqrt{a + b} \sqrt{-a + b} \sqrt{a + b \coth^2(x)}) \right)}{2\sqrt{2} \sqrt{b} \sqrt{a + b} \sqrt{-a + b}}$$

3.17. $\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx$

input `Integrate[Coth[x]^2*Sqrt[a + b*Coth[x]^2],x]`

output `-1/2*(Sqrt[(-a + b + (a + b)*Cosh[2*x])*Csch[x]^2]*(Sqrt[2]*Sqrt[a + b]*(a + 2*b)*ArcTanh[(Sqrt[2]*Sqrt[b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]] + Sqrt[b]*(-2*Sqrt[2]*(a + b)*ArcTanh[(Sqrt[2]*Sqrt[a + b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]]) + Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])*Sinh[x])/(Sqrt[2]*Sqrt[b]*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])`

3.17.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.588$, Rules used = {3042, 25, 4153, 25, 380, 398, 224, 219, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^2(x) \sqrt{a + b \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(\frac{\pi}{2} + ix\right)^2 \sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(ix + \frac{\pi}{2}\right)^2 \sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\coth^2(x) \sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\coth^2(x) \sqrt{a + b \coth^2(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{380} \\
 & \frac{1}{2} \int \frac{(a + 2b) \coth^2(x) + a}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}
 \end{aligned}$$

3.17. $\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx$

↓ 398

$$\frac{1}{2} \left(2(a+b) \int \frac{1}{(1-\coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - (a+2b) \int \frac{1}{\sqrt{b \coth^2(x) + a}} d \coth(x) \right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

↓ 224

$$\frac{1}{2} \left(2(a+b) \int \frac{1}{(1-\coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - (a+2b) \int \frac{1}{1 - \frac{b \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} \right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

↓ 219

$$\frac{1}{2} \left(2(a+b) \int \frac{1}{(1-\coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{(a+2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)}{\sqrt{b}} \right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

↓ 291

$$\frac{1}{2} \left(2(a+b) \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} - \frac{(a+2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)}{\sqrt{b}} \right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

↓ 219

$$\frac{1}{2} \left(2\sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \frac{(a+2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)}{\sqrt{b}} \right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

input `Int[Coth[x]^2*Sqrt[a + b*Coth[x]^2], x]`

```
output (-((a + 2*b)*ArcTanh[(Sqrt[b]*Coth[x])/Sqrt[a + b*Coth[x]^2]])/Sqrt[b]) +
  2*Sqrt[a + b]*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]])/2 - (
  Coth[x]*Sqrt[a + b*Coth[x]^2])/2
```

3.17.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
  ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
  Q[a, 0] || LtQ[b, 0])
```

```
rule 224 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
  x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

```
rule 291 Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst
  [Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c,
  d}, x] && NeQ[b*c - a*d, 0]
```

```
rule 380 Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_
  ), x_Symbol] := Simp[e*(e*x)^(m - 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(b*
  (m + 2*(p + q) + 1))), x] - Simp[e^2/(b*(m + 2*(p + q) + 1)) Int[(e*x)^(m
  - 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[a*c*(m - 1) + (a*d*(m - 1) - 2
  *q*(b*c - a*d))*x^2, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c
  - a*d, 0] && GtQ[q, 0] && GtQ[m, 1] && IntBinomialQ[a, b, c, d, e, m, 2, p,
  q, x]
```

```
rule 398 Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*Sqrt[(c_) + (d_.)*(x_)^2])
  , x_Symbol] := Simp[f/b Int[1/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/
  b Int[1/((a + b*x^2)*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}
  , x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
  Q[u, x]
```

```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)]^(n_))^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.17.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. $2(67) = 134$.

Time = 0.14 (sec) , antiderivative size = 276, normalized size of antiderivative = 3.25

method	result
derivativedivides	$-\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} - \frac{a \ln\left(\sqrt{b} \coth(x) + \sqrt{a+b\coth(x)^2}\right)}{2\sqrt{b}} - \frac{\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{2} - \frac{\sqrt{b}}{2}$
default	$-\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} - \frac{a \ln\left(\sqrt{b} \coth(x) + \sqrt{a+b\coth(x)^2}\right)}{2\sqrt{b}} - \frac{\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{2} - \frac{\sqrt{b}}{2}$

```
input int(coth(x)^2*(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/2*coth(x)*(a+b*coth(x)^2)^(1/2)-1/2*a/b^(1/2)*ln(b^(1/2)*coth(x)+(a+b*c
oth(x)^2)^(1/2))-1/2*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)-1/2*b^(1/
2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2
))+1/2*(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1
)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))+1/2*(b*(1+coth(x))^2-2*b*(1+c
oth(x))+a+b)^(1/2)-1/2*b^(1/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))
^2-2*b*(1+coth(x))+a+b)^(1/2))-1/2*(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))
+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))
```

3.17. $\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx$

3.17.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 907 vs. 2(67) = 134.

Time = 0.43 (sec) , antiderivative size = 4877, normalized size of antiderivative = 57.38

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \text{Too large to display}$$

input `integrate(coth(x)^2*(a+b*coth(x)^2)^(1/2),x, algorithm="fracas")`

output

```
[1/4*((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*b*cosh(x)^2 +
2*(3*b*cosh(x)^2 - b)*sinh(x)^2 + 4*(b*cosh(x)^3 - b*cosh(x))*sinh(x) + b
)*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(
x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 +
2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)
)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^
3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3
+ 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5
+ 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))
*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cos
h(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^
3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)
^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*
b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 +
3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cos
h(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b +
b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)
)*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b
)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)
)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cos...
```

3.17.6 Sympy [F]

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} \coth^2(x) dx$$

input `integrate(coth(x)**2*(a+b*coth(x)**2)**(1/2),x)`

output `Integral(sqrt(a + b*coth(x)**2)*coth(x)**2, x)`

3.17. $\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx$

3.17.7 Maxima [F]

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth(x)^2 + a} \coth(x)^2 dx$$

input `integrate(coth(x)^2*(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*coth(x)^2 + a)*coth(x)^2, x)`

3.17.8 Giac [F(-2)]

Exception generated.

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)^2*(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:Error: Bad Argument Type`

3.17.9 Mupad [F(-1)]

Timed out.

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \int \coth(x)^2 \sqrt{b \coth(x)^2 + a} dx$$

input `int(coth(x)^2*(a + b*coth(x)^2)^(1/2),x)`

output `int(coth(x)^2*(a + b*coth(x)^2)^(1/2), x)`

3.18 $\int \coth(x) \sqrt{a + b \coth^2(x)} dx$

3.18.1	Optimal result	153
3.18.2	Mathematica [A] (verified)	153
3.18.3	Rubi [A] (verified)	154
3.18.4	Maple [B] (verified)	156
3.18.5	Fricas [B] (verification not implemented)	157
3.18.6	Sympy [F]	157
3.18.7	Maxima [F]	158
3.18.8	Giac [F(-2)]	158
3.18.9	Mupad [B] (verification not implemented)	158

3.18.1 Optimal result

Integrand size = 15, antiderivative size = 44

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - \sqrt{a + b \coth^2(x)}$$

output `arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))*(a+b)^(1/2)-(a+b*coth(x)^2)^(1/2)`

3.18.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - \sqrt{a + b \coth^2(x)}$$

input `Integrate[Coth[x]*Sqrt[a + b*Coth[x]^2],x]`

output `Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - Sqrt[a + b*Coth[x]^2]`

3.18.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3042, 26, 4153, 26, 353, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth(x) \sqrt{a + b \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i \tan\left(\frac{\pi}{2} + ix\right) \sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \tan\left(ix + \frac{\pi}{2}\right) \sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{4153} \\
 & -i \int \frac{i \coth(x) \sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x) \sqrt{a + b \coth^2(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{353} \\
 & \frac{1}{2} \int \frac{\sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth^2(x) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left((a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - 2 \sqrt{a + b \coth^2(x)} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(\frac{2(a + b) \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d \sqrt{b \coth^2(x) + a}}{b} - 2 \sqrt{a + b \coth^2(x)} \right)
 \end{aligned}$$

$$\frac{1}{2} \left(2\sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \operatorname{coth}^2(x)}}{\sqrt{a+b}} \right) - 2\sqrt{a+b \operatorname{coth}^2(x)} \right)$$

input `Int[Coth[x]*Sqrt[a + b*Coth[x]^2], x]`

output `(2*Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - 2*Sqrt[a + b*Coth[x]^2])/2`

3.18.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 353 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4153 `Int[((d_)*tan[(e_) + (f_)*(x_)]^(m_))*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.18.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(36) = 72.

Time = 0.09 (sec) , antiderivative size = 238, normalized size of antiderivative = 5.41

method	result
derivativedivides	$-\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} + \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} + \dots$
default	$-\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} + \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} + \dots$

input `int(coth(x)*(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)+1/2*b^(1/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))+1/2*(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))-1/2*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)-1/2*b^(1/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))+1/2*(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))`

3.18. $\int \coth(x) \sqrt{a + b \coth^2(x)} dx$

3.18.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 495 vs. $2(36) = 72$.

Time = 0.31 (sec) , antiderivative size = 1551, normalized size of antiderivative = 35.25

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \text{Too large to display}$$

```
input integrate(coth(x)*(a+b*coth(x)^2)^(1/2),x, algorithm="fracas")
```

```
output [1/4*((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(a + b)*log(-(a
^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*
sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2
*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*
b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a
^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)
*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)
*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*
a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 +
a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 +
3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cos
h(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*
a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(
x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)
)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)
)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)
*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh
(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*
b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*
b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)...
```

3.18.6 Sympy [F]

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} \coth(x) dx$$

```
input integrate(coth(x)*(a+b*coth(x)**2)**(1/2),x)
```

```
output Integral(sqrt(a + b*coth(x)**2)*coth(x), x)
```

3.18. $\int \coth(x) \sqrt{a + b \coth^2(x)} dx$

3.18.7 Maxima [F]

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth(x)^2 + a} \coth(x) dx$$

input `integrate(coth(x)*(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*coth(x)^2 + a)*coth(x), x)`

3.18.8 Giac [F(-2)]

Exception generated.

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)*(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Error: Bad Argument Type`

3.18.9 Mupad [B] (verification not implemented)

Time = 2.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.16

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = -\sqrt{b \coth(x)^2 + a} - 2 \operatorname{atan} \left(\frac{2 \sqrt{b \coth(x)^2 + a} \sqrt{-\frac{a}{4} - \frac{b}{4}}}{a + b} \right) \sqrt{-\frac{a}{4} - \frac{b}{4}}$$

input `int(coth(x)*(a + b*coth(x)^2)^(1/2),x)`

output `-(a + b*coth(x)^2)^(1/2) - 2*atan((2*(a + b*coth(x)^2)^(1/2)*(- a/4 - b/4)
)^(1/2))/(a + b)*(- a/4 - b/4)^(1/2)`

3.18. $\int \coth(x) \sqrt{a + b \coth^2(x)} dx$

3.19 $\int \sqrt{a + b \coth^2(x)} dx$

3.19.1	Optimal result	159
3.19.2	Mathematica [A] (verified)	159
3.19.3	Rubi [A] (verified)	160
3.19.4	Maple [B] (verified)	162
3.19.5	Fricas [B] (verification not implemented)	162
3.19.6	Sympy [F]	163
3.19.7	Maxima [F]	164
3.19.8	Giac [B] (verification not implemented)	164
3.19.9	Mupad [F(-1)]	165

3.19.1 Optimal result

Integrand size = 12, antiderivative size = 60

$$\int \sqrt{a + b \coth^2(x)} dx = -\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) + \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)$$

output `-arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))*b^(1/2)+arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))*(a+b)^(1/2)`

3.19.2 Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.37

$$\int \sqrt{a + b \coth^2(x)} dx = \sqrt{-a - b} \operatorname{arctan} \left(\frac{\coth(x) \sqrt{a + b \coth^2(x)} - \sqrt{b} \operatorname{csch}^2(x)}{\sqrt{-a - b}} \right) + \sqrt{b} \log \left(-\sqrt{b} \coth(x) + \sqrt{a + b \coth^2(x)} \right)$$

input `Integrate[Sqrt[a + b*Coth[x]^2], x]`

output `Sqrt[-a - b]*ArcTan[(Coth[x]*Sqrt[a + b*Coth[x]^2] - Sqrt[b]*Csch[x]^2)/Sqrt[-a - b]] + Sqrt[b]*Log[-(Sqrt[b]*Coth[x]) + Sqrt[a + b*Coth[x]^2]]`

3.19.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 4144, 301, 224, 219, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + b \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{\sqrt{a + b \coth^2(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{301} \\
 & (a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - b \int \frac{1}{\sqrt{b \coth^2(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{224} \\
 & (a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - b \int \frac{1}{1 - \frac{b \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} \\
 & \quad \downarrow \text{219} \\
 & (a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) \\
 & \quad \downarrow \text{291} \\
 & (a + b) \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} - \sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)
 \end{aligned}$$

$$\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a+b} \operatorname{coth}(x)}{\sqrt{a+b \operatorname{coth}^2(x)}}\right) - \sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \operatorname{coth}(x)}{\sqrt{a+b \operatorname{coth}^2(x)}}\right)$$

input `Int[Sqrt[a + b*Coth[x]^2], x]`

output `-(Sqrt[b]*ArcTanh[(Sqrt[b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]) + Sqrt[a + b]*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]`

3.19.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 301 `Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[b/d Int[(a + b*x^2)^(p - 1), x], x] - Simp[(b*c - a*d)/d Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4] || (EqQ[p, 2/3] && EqQ[b*c + 3*a*d, 0]))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4144 Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*
(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] ||
EqQ[n^2, 16])
```

3.19.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(48) = 96.

Time = 0.14 (sec) , antiderivative size = 238, normalized size of antiderivative = 3.97

method	result
derivativedivides	$\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} - \frac{\sqrt{a+b}}{2}$
default	$\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} - \frac{\sqrt{a+b}}{2}$

```
input int((a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/2*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-1/2*b^(1/2)*ln((b*(1+coth(
x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))-1/2*(a+b)^(1/2
)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x
))+a+b)^(1/2))/(1+coth(x)))-1/2*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2
)-1/2*b^(1/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1
)+a+b)^(1/2))+1/2*(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b
*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))
```

3.19.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 552 vs. 2(48) = 96.

Time = 0.37 (sec) , antiderivative size = 3455, normalized size of antiderivative = 57.58

$$\int \sqrt{a + b \coth^2(x)} dx = \text{Too large to display}$$

```
input integrate((a+b*coth(x)^2)^(1/2),x, algorithm="fracas")
```

3.19. $\int \sqrt{a + b \coth^2(x)} dx$

```
output [1/4*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x))^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)...
```

3.19.6 Sympy [F]

$$\int \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} dx$$

```
input integrate((a+b*coth(x)**2)**(1/2), x)
```

```
output Integral(sqrt(a + b*coth(x)**2), x)
```

3.19.7 Maxima [F]

$$\int \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth(x)^2 + a} dx$$

input `integrate((a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*coth(x)^2 + a), x)`

3.19.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(48) = 96.

Time = 0.48 (sec) , antiderivative size = 262, normalized size of antiderivative = 4.37

$$\int \sqrt{a + b \coth^2(x)} dx = \frac{1}{2} \left(\frac{4 b \arctan \left(-\frac{\sqrt{a+b}e^{(2x)} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a+b}}{2\sqrt{-b}} \right)}{\sqrt{-b}} - \sqrt{a+b} \log \left(\left| \left(\sqrt{a+b}e^{(2x)} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} \right) - 1 \right| \right) \right)$$

input `integrate((a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `1/2*(4*b*arctan(-1/2*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b))/sqrt(-b))/sqrt(-b) - sqrt(a + b)*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) - sqrt(a + b)*(a - b))) + sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))) - sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b))))*sgn(e^(2*x) - 1)`

3.19.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth(x)^2 + a} dx$$

input `int((a + b*coth(x)^2)^(1/2),x)`output `int((a + b*coth(x)^2)^(1/2), x)`

3.20 $\int \sqrt{a + b \coth^2(x)} \tanh(x) dx$

3.20.1	Optimal result	166
3.20.2	Mathematica [A] (verified)	166
3.20.3	Rubi [A] (verified)	167
3.20.4	Maple [F]	169
3.20.5	Fricas [B] (verification not implemented)	169
3.20.6	Sympy [F]	170
3.20.7	Maxima [F]	171
3.20.8	Giac [B] (verification not implemented)	171
3.20.9	Mupad [F(-1)]	172

3.20.1 Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = -\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}} \right) + \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right)$$

output `-arctanh((a+b*coth(x)^2)^(1/2)/a^(1/2))*a^(1/2)+arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))*(a+b)^(1/2)`

3.20.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = -\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}} \right) + \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right)$$

input `Integrate[Sqrt[a + b*Coth[x]^2]*Tanh[x],x]`

output `-(Sqrt[a]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]]) + Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]`

3.20.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3042, 26, 4153, 26, 354, 94, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(x) \sqrt{a + b \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \sqrt{a - b \tan^2\left(\frac{\pi}{2} + ix\right)}}{\tan\left(\frac{\pi}{2} + ix\right)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\sqrt{a - b \tan^2\left(ix + \frac{\pi}{2}\right)}}{\tan\left(ix + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \sqrt{b \coth^2(x) + a \tanh(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\tanh(x) \sqrt{a + b \coth^2(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\sqrt{b \coth^2(x) + a \tanh(x)}}{1 - \coth^2(x)} d \coth^2(x) \\
 & \quad \downarrow \text{94}
 \end{aligned}$$

$$\frac{1}{2} \left((a+b) \int \frac{1}{(1-\coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) + a \int \frac{\tanh(x)}{\sqrt{b \coth^2(x) + a}} d \coth^2(x) \right)$$

↓ 73

$$\frac{1}{2} \left(\frac{2(a+b) \int \frac{1}{\frac{a+b-\coth^4(x)}{b}} d \sqrt{b \coth^2(x) + a}}{b} + \frac{2a \int \frac{1}{\frac{\coth^4(x)-a}{b}} d \sqrt{b \coth^2(x) + a}}{b} \right)$$

↓ 221

$$\frac{1}{2} \left(2\sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right) - 2\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}} \right) \right)$$

input `Int[Sqrt[a + b*Coth[x]^2]*Tanh[x], x]`

output `(-2*Sqrt[a]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]] + 2*Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/2`

3.20.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 94 `Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] := Simp[(b*e - a*f)/(b*c - a*d) Int[(e + f*x)^(p - 1)/(a + b*x), x], x] - Simp[(d*e - c*f)/(b*c - a*d) Int[(e + f*x)^(p - 1)/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[0, p, 1]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 354 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4153 `Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.20.4 Maple [F]

$$\int \sqrt{a + b \coth(x)^2} \tanh(x) dx$$

input `int((a+b*coth(x)^2)^(1/2)*tanh(x),x)`

output `int((a+b*coth(x)^2)^(1/2)*tanh(x),x)`

3.20.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 557 vs. $2(44) = 88$.

Time = 0.38 (sec) , antiderivative size = 3479, normalized size of antiderivative = 62.12

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \text{Too large to display}$$

input `integrate((a+b*coth(x)^2)^(1/2)*tanh(x),x, algorithm="fracas")`

3.20. $\int \sqrt{a + b \coth^2(x)} \tanh(x) dx$

output `[1/4*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(...`

3.20.6 Sympy [F]

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \int \sqrt{a + b \coth^2(x)} \tanh(x) dx$$

input `integrate((a+b*coth(x)**2)**(1/2)*tanh(x),x)`

output `Integral(sqrt(a + b*coth(x)**2)*tanh(x), x)`

3.20.7 Maxima [F]

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \int \sqrt{b \coth^2(x) + a} \tanh(x) dx$$

input `integrate((a+b*coth(x)^2)^(1/2)*tanh(x),x, algorithm="maxima")`

output `integrate(sqrt(b*coth(x)^2 + a)*tanh(x), x)`

3.20.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. 2(44) = 88.

Time = 0.51 (sec) , antiderivative size = 259, normalized size of antiderivative = 4.62

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx =$$

$$-\frac{1}{2} \left(\frac{4a \arctan \left(-\frac{\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a+b}}}{2\sqrt{-a}} \right)}{\sqrt{-a}} + \sqrt{a+b} \log \left(\left| \left(\sqrt{a+be^{2x}} - \sqrt{a+b} \right) \right. \right. \right.$$

- 1)

input `integrate((a+b*coth(x)^2)^(1/2)*tanh(x),x, algorithm="giac")`

output `-1/2*(4*a*arctan(-1/2*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))/sqrt(-a))/sqrt(-a) + sqrt(a + b)*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) - sqrt(a + b)*(a - b))) + sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))) - sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b))))*sgn(e^(2*x) - 1)`

3.20.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \int \tanh(x) \sqrt{b \coth(x)^2 + a} dx$$

input `int(tanh(x)*(a + b*coth(x)^2)^(1/2), x)`output `int(tanh(x)*(a + b*coth(x)^2)^(1/2), x)`

3.21 $\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx$

3.21.1	Optimal result	173
3.21.2	Mathematica [C] (verified)	173
3.21.3	Rubi [A] (verified)	174
3.21.4	Maple [F]	176
3.21.5	Fricas [B] (verification not implemented)	176
3.21.6	Sympy [F]	177
3.21.7	Maxima [F]	178
3.21.8	Giac [F(-2)]	178
3.21.9	Mupad [F(-1)]	178

3.21.1 Optimal result

Integrand size = 17, antiderivative size = 48

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \sqrt{a + b \coth^2(x)} \tanh(x)$$

output `arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))*(a+b)^(1/2)-(a+b*coth(x)^2)^(1/2)*tanh(x)`

3.21.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.10 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.88

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = -\sqrt{a + b \coth^2(x)} \operatorname{Hypergeometric2F1} \left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{(a + b) \coth^2(x)}{a + b \coth^2(x)} \right) \tanh(x)$$

input `Integrate[Sqrt[a + b*Coth[x]^2]*Tanh[x]^2,x]`

output `-(Sqrt[a + b*Coth[x]^2]*Hypergeometric2F1[-1/2, 1, 1/2, ((a + b)*Coth[x]^2)/(a + b*Coth[x]^2)]*Tanh[x])`

3.21. $\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx$

3.21.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 25, 4153, 25, 377, 27, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^2(x) \sqrt{a + b \coth^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2}}{\tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2}}{\tan\left(ix + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\sqrt{b \coth^2(x) + a \tanh^2(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\tanh^2(x) \sqrt{a + b \coth^2(x)}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{377} \\
 & \int \frac{a + b}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \tanh(x) \sqrt{a + b \coth^2(x)} \\
 & \quad \downarrow \text{27} \\
 & (a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \tanh(x) \sqrt{a + b \coth^2(x)} \\
 & \quad \downarrow \text{291} \\
 & (a + b) \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} - \tanh(x) \sqrt{a + b \coth^2(x)}
 \end{aligned}$$

3.21. $\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx$

$$\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a+b} \operatorname{coth}(x)}{\sqrt{a+b \operatorname{coth}^2(x)}}\right) - \tanh(x) \sqrt{a+b \operatorname{coth}^2(x)}$$

input `Int[Sqrt[a + b*Coth[x]^2]*Tanh[x]^2,x]`

output `Sqrt[a + b]*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]] - Sqrt[a + b*Coth[x]^2]*Tanh[x]`

3.21.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 377 `Int[((e_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[(e*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(a*e*(m + 1))), x] - Simp[1/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[b*c*(m + 1) + 2*(b*c*(p + 1) + a*d*q) + d*(b*(m + 1) + 2*b*(p + q + 1))*x^2, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && LtQ[0, q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.21. $\int \sqrt{a + b \operatorname{coth}^2(x)} \tanh^2(x) dx$


```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)]^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.21.4 Maple [F]

$$\int \sqrt{a + b \coth(x)^2} \tanh(x)^2 dx$$

```
input int((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x)
```

```
output int((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x)
```

3.21.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 490 vs. $2(40) = 80$.

Time = 0.32 (sec) , antiderivative size = 1539, normalized size of antiderivative = 32.06

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \text{Too large to display}$$

```
input integrate((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x, algorithm="fricas")
```

output `[1/4*((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x))^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)...`

3.21.6 Sympy [F]

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx$$

input `integrate((a+b*coth(x)**2)**(1/2)*tanh(x)**2,x)`

output `Integral(sqrt(a + b*coth(x)**2)*tanh(x)**2, x)`

3.21.7 Maxima [F]

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \int \sqrt{b \coth(x)^2 + a} \tanh(x)^2 dx$$

input `integrate((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x, algorithm="maxima")`

output `integrate(sqrt(b*coth(x)^2 + a)*tanh(x)^2, x)`

3.21.8 Giac [F(-2)]

Exception generated.

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:Error: Bad Argument Type`

3.21.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \int \tanh(x)^2 \sqrt{b \coth(x)^2 + a} dx$$

input `int(tanh(x)^2*(a + b*coth(x)^2)^(1/2),x)`

output `int(tanh(x)^2*(a + b*coth(x)^2)^(1/2), x)`

3.22 $\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx$

3.22.1	Optimal result	179
3.22.2	Mathematica [A] (verified)	179
3.22.3	Rubi [A] (verified)	180
3.22.4	Maple [B] (verified)	183
3.22.5	Fricas [B] (verification not implemented)	183
3.22.6	Sympy [F]	184
3.22.7	Maxima [F]	185
3.22.8	Giac [F(-2)]	185
3.22.9	Mupad [B] (verification not implemented)	185

3.22.1 Optimal result

Integrand size = 17, antiderivative size = 82

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - (a + b) \sqrt{a + b \coth^2(x)} - \frac{1}{3} (a + b \coth^2(x))^{3/2} - \frac{(a + b \coth^2(x))^{5/2}}{5b}$$

output $(a+b)^{(3/2)}*\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2))/(a+b)^{(1/2)})-1/3*(a+b*\coth(x)^2)^{(3/2)}-1/5*(a+b*\coth(x)^2)^{(5/2)}/b-(a+b)*(a+b*\coth(x)^2)^{(1/2)}$

3.22.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.05

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - \frac{\sqrt{a + b \coth^2(x)} (3a^2 + 20ab + 15b^2 + b(6a + 5b) \coth^2(x) + 3b^2 \coth^4(x))}{15b}$$

input `Integrate[Coth[x]^3*(a + b*Coth[x]^2)^(3/2),x]`

output $(a + b)^{3/2} \text{ArcTanh}[\text{Sqrt}[a + b \text{Coth}[x]^2] / \text{Sqrt}[a + b]] - (\text{Sqrt}[a + b \text{Coth}[x]^2] * (3a^2 + 20ab + 15b^2 + b(6a + 5b) \text{Coth}[x]^2 + 3b^2 \text{Coth}[x]^4)) / (15b)$

3.22.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.09, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.588$, Rules used = {3042, 26, 4153, 26, 354, 90, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^3(x) (a + b \coth^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int i \tan\left(\frac{\pi}{2} + ix\right)^3 \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{3/2} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \tan\left(ix + \frac{\pi}{2}\right)^3 \left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{3/2} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \coth^3(x) (b \coth^2(x) + a)^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth^3(x) (a + b \coth^2(x))^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\coth^2(x) (b \coth^2(x) + a)^{3/2}}{1 - \coth^2(x)} d \coth^2(x) \\
 & \quad \downarrow \text{90} \\
 & \frac{1}{2} \left(\int \frac{(b \coth^2(x) + a)^{3/2}}{1 - \coth^2(x)} d \coth^2(x) - \frac{2(a + b \coth^2(x))^{5/2}}{5b} \right) \\
 & \quad \downarrow \text{60}
 \end{aligned}$$

$$\frac{1}{2} \left((a+b) \int \frac{\sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth^2(x) - \frac{2(a + b \coth^2(x))^{5/2}}{5b} - \frac{2}{3} (a + b \coth^2(x))^{3/2} \right)$$

↓ 60

$$\frac{1}{2} \left((a+b) \left((a+b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - 2\sqrt{a + b \coth^2(x)} \right) - \frac{2(a + b \coth^2(x))^{5/2}}{5b} \right)$$

↓ 73

$$\frac{1}{2} \left((a+b) \left(\frac{2(a+b) \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d \sqrt{b \coth^2(x) + a}}{b} - 2\sqrt{a + b \coth^2(x)} \right) - \frac{2(a + b \coth^2(x))^{5/2}}{5b} - \frac{2}{3} (a + b \coth^2(x))^{3/2} \right)$$

↓ 221

$$\frac{1}{2} \left((a+b) \left(2\sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a+b}} \right) - 2\sqrt{a + b \coth^2(x)} \right) - \frac{2(a + b \coth^2(x))^{5/2}}{5b} - \frac{2}{3} (a + b \coth^2(x))^{3/2} \right)$$

input `Int[Coth[x]^3*(a + b*Coth[x]^2)^(3/2),x]`

output `((-2*(a + b*Coth[x]^2)^(3/2))/3 - (2*(a + b*Coth[x]^2)^(5/2))/(5*b) + (a + b)*(2*Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - 2*Sqrt[a + b*Coth[x]^2]))/2`

3.22.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

3.22. $\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx$

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
 .), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
 x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
 + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
 p}, x] && NeQ[n + p + 2, 0]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_S
 ymbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x
 , x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ
 [(m - 1)/2]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4153 `Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
 (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
 x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
 f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m,
 n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
 nalQ[n]))`

3.22.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 487 vs. $2(66) = 132$.

Time = 0.09 (sec) , antiderivative size = 488, normalized size of antiderivative = 5.95

method	result
derivativedivides	$-\frac{(a+b \operatorname{coth}(x)^2)^{5/2}}{5b} - \frac{(b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b)^{3/2}}{6} - b \left(\frac{(2b(\operatorname{coth}(x)-1)+2b)\sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}}{4b} \right)$
default	$-\frac{(a+b \operatorname{coth}(x)^2)^{5/2}}{5b} - \frac{(b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b)^{3/2}}{6} - b \left(\frac{(2b(\operatorname{coth}(x)-1)+2b)\sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}}{4b} \right)$

input `int(coth(x)^3*(a+b*coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -1/5*(a+b*\operatorname{coth}(x)^2)^{(5/2)}/b-1/6*(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(3/2)} \\ & -1/2*b*(1/4*(2*b*(\operatorname{coth}(x)-1)+2*b)/b*(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b) \\ &)^{(1/2)}+1/8*(4*(a+b)*b-4*b^2)/b^{(3/2)}*\ln((b*(\operatorname{coth}(x)-1)+b)/b^{(1/2)}+(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)})) \\ & -1/2*(a+b)*((b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)}+b^{(1/2)}*\ln((b*(\operatorname{coth}(x)-1)+b)/b^{(1/2)}+(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)})) \\ & -(a+b)^{(1/2)}*\ln((2*a+2*b+2*b*(\operatorname{coth}(x)-1)+2*(a+b))^{(1/2)}*(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)})/(b*(\operatorname{coth}(x)-1))) \\ & -1/6*(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(3/2)}+1/2*b*(1/4*(2*b*(1+\operatorname{coth}(x))-2*b)/b*(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)}+1/8*(4*(a+b)*b-4*b^2)/b^{(3/2)} \\ &)*\ln((b*(1+\operatorname{coth}(x))-b)/b^{(1/2)}+(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)})) \\ & -1/2*(a+b)*((b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)}-b^{(1/2)}*\ln((b*(1+\operatorname{coth}(x))-b)/b^{(1/2)}+(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)})) \\ & -(a+b)^{(1/2)}*\ln((2*a+2*b-2*b*(1+\operatorname{coth}(x))+2*(a+b))^{(1/2)}*(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)})/(1+\operatorname{coth}(x)))) \end{aligned}$$

3.22.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2154 vs. $2(66) = 132$.

Time = 0.47 (sec) , antiderivative size = 4940, normalized size of antiderivative = 60.24

$$\int \operatorname{coth}^3(x) (a + b \operatorname{coth}^2(x))^{3/2} dx = \text{Too large to display}$$

input `integrate(coth(x)^3*(a+b*coth(x)^2)^(3/2),x, algorithm="fracas")`

3.22. $\int \operatorname{coth}^3(x) (a + b \operatorname{coth}^2(x))^{3/2} dx$


```
output [1/60*(15*((a*b + b^2)*cosh(x)^10 + 10*(a*b + b^2)*cosh(x)*sinh(x)^9 + (a*
b + b^2)*sinh(x)^10 - 5*(a*b + b^2)*cosh(x)^8 + 5*(9*(a*b + b^2)*cosh(x)^2
- a*b - b^2)*sinh(x)^8 + 40*(3*(a*b + b^2)*cosh(x)^3 - (a*b + b^2)*cosh(x
))*sinh(x)^7 + 10*(a*b + b^2)*cosh(x)^6 + 10*(21*(a*b + b^2)*cosh(x)^4 - 1
4*(a*b + b^2)*cosh(x)^2 + a*b + b^2)*sinh(x)^6 + 4*(63*(a*b + b^2)*cosh(x)
^5 - 70*(a*b + b^2)*cosh(x)^3 + 15*(a*b + b^2)*cosh(x))*sinh(x)^5 - 10*(a*
b + b^2)*cosh(x)^4 + 10*(21*(a*b + b^2)*cosh(x)^6 - 35*(a*b + b^2)*cosh(x)
^4 + 15*(a*b + b^2)*cosh(x)^2 - a*b - b^2)*sinh(x)^4 + 40*(3*(a*b + b^2)*c
osh(x)^7 - 7*(a*b + b^2)*cosh(x)^5 + 5*(a*b + b^2)*cosh(x)^3 - (a*b + b^2)
*cosh(x))*sinh(x)^3 + 5*(a*b + b^2)*cosh(x)^2 + 5*(9*(a*b + b^2)*cosh(x)^8
- 28*(a*b + b^2)*cosh(x)^6 + 30*(a*b + b^2)*cosh(x)^4 - 12*(a*b + b^2)*co
sh(x)^2 + a*b + b^2)*sinh(x)^2 - a*b - b^2 + 10*((a*b + b^2)*cosh(x)^9 - 4
*(a*b + b^2)*cosh(x)^7 + 6*(a*b + b^2)*cosh(x)^5 - 4*(a*b + b^2)*cosh(x)^3
+ (a*b + b^2)*cosh(x))*sinh(x))*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8
+ 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3
+ a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)
)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5
+ (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4
+ 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4
+ 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^...
```

3.22.6 Sympy [F]

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \coth^3(x) dx$$

```
input integrate(coth(x)**3*(a+b*coth(x)**2)**(3/2),x)
```

```
output Integral((a + b*coth(x)**2)**(3/2)*coth(x)**3, x)
```

3.22.7 Maxima [F]

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \coth(x)^3 dx$$

input `integrate(coth(x)^3*(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate((b*coth(x)^2 + a)^(3/2)*coth(x)^3, x)`

3.22.8 Giac [F(-2)]

Exception generated.

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)^3*(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

3.22.9 Mupad [B] (verification not implemented)

Time = 9.99 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.37

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = -\frac{(b \coth(x)^2 + a)^{5/2}}{5b} - \left(\frac{a+b}{3b} - \frac{a}{3b}\right) (b \coth(x)^2 + a)^{3/2} \\ - (a+b) \left(\frac{a+b}{b} - \frac{a}{b}\right) \sqrt{b \coth(x)^2 + a} - \operatorname{atan}\left(\frac{(a+b)^{3/2} \sqrt{b \coth(x)^2 + a} \operatorname{li}}{a^2 + 2ab + b^2}\right) (a+b)^{3/2} \operatorname{li}$$

input `int(coth(x)^3*(a + b*coth(x)^2)^(3/2),x)`

output `- (a + b*coth(x)^2)^(5/2)/(5*b) - ((a + b)/(3*b) - a/(3*b))*(a + b*coth(x)
^2)^(3/2) - atan(((a + b)^(3/2)*(a + b*coth(x)^2)^(1/2)*li)/(2*a*b + a^2 +
b^2))*(a + b)^(3/2)*li - (a + b)*((a + b)/b - a/b)*(a + b*coth(x)^2)^(1/2
)`

3.23 $\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx$

3.23.1	Optimal result	186
3.23.2	Mathematica [A] (verified)	186
3.23.3	Rubi [A] (verified)	187
3.23.4	Maple [B] (verified)	191
3.23.5	Fricas [B] (verification not implemented)	192
3.23.6	Sympy [F]	192
3.23.7	Maxima [F]	192
3.23.8	Giac [F(-2)]	193
3.23.9	Mupad [F(-1)]	193

3.23.1 Optimal result

Integrand size = 17, antiderivative size = 123

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = -\frac{(3a^2 + 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{8\sqrt{b}} + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right) - \frac{1}{8}(5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} - \frac{1}{4}b \coth^3(x) \sqrt{a + b \coth^2(x)}$$

```
output (a+b)^(3/2)*arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))-1/8*(3*a^2+
12*a*b+8*b^2)*arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))/b^(1/2)-1/8*(
5*a+4*b)*coth(x)*(a+b*coth(x)^2)^(1/2)-1/4*b*coth(x)^3*(a+b*coth(x)^2)^(1/
2)
```

3.23.2 Mathematica [A] (verified)

Time = 0.95 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.78

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \frac{\sqrt{(-a + b + (a + b) \cosh(2x)) \operatorname{csch}^2(x)} \left(-\sqrt{2} \sqrt{a + b} (3a^2 + 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right) \right)}{\dots}$$

input `Integrate[Coth[x]^2*(a + b*Coth[x]^2)^(3/2),x]`

output `(Sqrt[(-a + b + (a + b)*Cosh[2*x])*Csch[x]^2]*(-(Sqrt[2]*Sqrt[a + b]*(3*a^2 + 12*a*b + 8*b^2)*ArcTanh[(Sqrt[2]*Sqrt[b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]) + Sqrt[b]*(8*Sqrt[2]*(a + b)^2*ArcTanh[(Sqrt[2]*Sqrt[a + b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]) - Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x])*Coth[x]*Csch[x]*(5*a + 6*b + 2*b*Csch[x]^2)))*Sinh[x])/(8*Sqrt[2]*Sqrt[b]*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])`

3.23.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.765$, Rules used = {3042, 25, 4153, 25, 379, 25, 444, 27, 398, 224, 219, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^2(x) (a + b \coth^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(\frac{\pi}{2} + ix\right)^2 \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{3/2} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(ix + \frac{\pi}{2}\right)^2 \left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{3/2} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\coth^2(x) (b \coth^2(x) + a)^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\coth^2(x) (a + b \coth^2(x))^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{379} \\
 & -\frac{1}{4} \int -\frac{\coth^2(x) (b(5a + 4b) \coth^2(x) + a(4a + 3b))}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{1}{4} \int \frac{\coth^2(x) (b(5a + 4b) \coth^2(x) + a(4a + 3b))}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)} \\
& \downarrow 444 \\
& \frac{1}{4} \left(\frac{\int \frac{b((3a^2 + 12ba + 8b^2) \coth^2(x) + a(5a + 4b))}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x)}{2b} - \frac{1}{2} (5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} \right) - \\
& \qquad \qquad \qquad \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)} \\
& \downarrow 27 \\
& \frac{1}{4} \left(\frac{1}{2} \int \frac{(3a^2 + 12ba + 8b^2) \coth^2(x) + a(5a + 4b)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{1}{2} (5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} \right) - \\
& \qquad \qquad \qquad \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)} \\
& \downarrow 398 \\
& \frac{1}{4} \left(\frac{1}{2} \left(8(a + b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - (3a^2 + 12ab + 8b^2) \int \frac{1}{\sqrt{b \coth^2(x) + a}} d \coth(x) \right) - \right. \\
& \qquad \qquad \qquad \left. \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)} \right) \\
& \downarrow 224 \\
& \frac{1}{4} \left(\frac{1}{2} \left(8(a + b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - (3a^2 + 12ab + 8b^2) \int \frac{1}{1 - \frac{b \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} \right) - \right. \\
& \qquad \qquad \qquad \left. \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)} \right) \\
& \downarrow 219 \\
& \frac{1}{4} \left(\frac{1}{2} \left(8(a + b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{(3a^2 + 12ab + 8b^2) \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)}{\sqrt{b}} \right) - \right. \\
& \qquad \qquad \qquad \left. \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)} \right)
\end{aligned}$$

$$\begin{array}{c}
 \downarrow 291 \\
 \frac{1}{4} \left(\frac{1}{2} \left(8(a+b)^2 \int \frac{1}{1 - \frac{(a+b)\coth^2(x)}{b\coth^2(x)+a}} d \frac{\coth(x)}{\sqrt{b\coth^2(x)+a}} - \frac{(3a^2 + 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{\sqrt{b}} \right) - \frac{1}{2}(5a + \right. \\
 \left. \frac{1}{4}b\coth^3(x)\sqrt{a+b\coth^2(x)} \right) \\
 \downarrow 219 \\
 \frac{1}{4} \left(\frac{1}{2} \left(8(a+b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right) - \frac{(3a^2 + 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{\sqrt{b}} \right) - \frac{1}{2}(5a + 4b) \operatorname{coth}(x) \right. \\
 \left. \frac{1}{4}b\coth^3(x)\sqrt{a+b\coth^2(x)} \right)
 \end{array}$$

input `Int[Coth[x]^2*(a + b*Coth[x]^2)^(3/2), x]`

output `-1/4*(b*Coth[x]^3*Sqrt[a + b*Coth[x]^2]) + (((-(((3*a^2 + 12*a*b + 8*b^2)*ArcTanh[(Sqrt[b]*Coth[x])/Sqrt[a + b*Coth[x]^2]])/Sqrt[b]) + 8*(a + b)^(3/2))*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]])/2 - ((5*a + 4*b)*Coth[x]*Sqrt[a + b*Coth[x]^2])/2)/4`

3.23.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

- rule 224 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 291 `Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*((c_) + (d_)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 379 `Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1)/(b*e*(m + 2*(p + q) + 1))), x] + Simp[1/(b*(m + 2*(p + q) + 1)) Int[(e*x)^m*(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*((b*c - a*d)*(m + 1) + b*c*2*(p + q)) + (d*(b*c - a*d)*(m + 1) + d*2*(q - 1)*(b*c - a*d) + b*c*d*2*(p + q))*x^2, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`
- rule 398 `Int[((e_) + (f_)*(x_)^2)/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[f/b Int[1/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/((a + b*x^2)*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`
- rule 444 `Int[((g_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_) * ((e_) + (f_)*(x_)^2), x_Symbol] := Simp[f*g*(g*x)^(m - 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(b*d*(m + 2*(p + q) + 1))), x] - Simp[g^2/(b*d*(m + 2*(p + q) + 1)) Int[(g*x)^(m - 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m - 1) + (a*f*d*(m + 2*q + 1) + b*(f*c*(m + 2*p + 1) - e*d*(m + 2*(p + q) + 1)))*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && GtQ[m, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)]^(n_))^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.23.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 528 vs. 2(101) = 202.

Time = 0.10 (sec) , antiderivative size = 529, normalized size of antiderivative = 4.30

method	result
derivativedivides	$\frac{\coth(x)(a+b\coth(x)^2)^{\frac{3}{2}}}{4} - \frac{3a\left(\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} + \frac{a\ln(\sqrt{b}\coth(x)+\sqrt{a+b\coth(x)^2})}{2\sqrt{b}}\right)}{4} - \frac{(b(\coth(x)-1)^2+2b)}{4}$
default	$\frac{\coth(x)(a+b\coth(x)^2)^{\frac{3}{2}}}{4} - \frac{3a\left(\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} + \frac{a\ln(\sqrt{b}\coth(x)+\sqrt{a+b\coth(x)^2})}{2\sqrt{b}}\right)}{4} - \frac{(b(\coth(x)-1)^2+2b)}{4}$

```
input int(coth(x)^2*(a+b*coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -1/4*coth(x)*(a+b*coth(x)^2)^(3/2)-3/4*a*(1/2*coth(x)*(a+b*coth(x)^2)^(1/2)
)+1/2*a/b^(1/2)*ln(b^(1/2)*coth(x)+(a+b*coth(x)^2)^(1/2))-1/6*(b*(coth(x)
-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)-1/2*b*(1/4*(2*b*(coth(x)-1)+2*b)/b*(b*(co
th(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1/8*(4*(a+b)*b-4*b^2)/b^(3/2)*ln((b*
(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))-1/2*(
a+b)*((b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+b^(1/2)*ln((b*(coth(x)-1
)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))-(a+b)^(1/2)*ln((
2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b
)^(1/2))/(coth(x)-1))+1/6*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)-1/2
*b*(1/4*(2*b*(1+coth(x))-2*b)/b*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2
)+1/8*(4*(a+b)*b-4*b^2)/b^(3/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x)
)^2-2*b*(1+coth(x))+a+b)^(1/2))+1/2*(a+b)*((b*(1+coth(x))^2-2*b*(1+coth(x)
))+a+b)^(1/2)-b^(1/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1
+coth(x))+a+b)^(1/2))-(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2
)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x))))
```

3.23. $\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx$

3.23.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2260 vs. $2(101) = 202$.

Time = 0.60 (sec) , antiderivative size = 10286, normalized size of antiderivative = 83.63

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \text{Too large to display}$$

input `integrate(coth(x)^2*(a+b*coth(x)^2)^(3/2),x, algorithm="fracas")`

output `Too large to include`

3.23.6 Sympy [F]

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \coth^2(x) dx$$

input `integrate(coth(x)**2*(a+b*coth(x)**2)**(3/2),x)`

output `Integral((a + b*coth(x)**2)**(3/2)*coth(x)**2, x)`

3.23.7 Maxima [F]

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \coth(x)^2 dx$$

input `integrate(coth(x)^2*(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate((b*coth(x)^2 + a)^(3/2)*coth(x)^2, x)`

3.23.8 Giac [F(-2)]

Exception generated.

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)^2*(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

3.23.9 Mupad [F(-1)]

Timed out.

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \int \coth(x)^2 (b \coth(x)^2 + a)^{3/2} dx$$

input `int(coth(x)^2*(a + b*coth(x)^2)^(3/2),x)`

output `int(coth(x)^2*(a + b*coth(x)^2)^(3/2), x)`

3.24 $\int \coth(x) (a + b \coth^2(x))^{3/2} dx$

3.24.1	Optimal result	194
3.24.2	Mathematica [A] (verified)	194
3.24.3	Rubi [A] (verified)	195
3.24.4	Maple [B] (verified)	197
3.24.5	Fricas [B] (verification not implemented)	198
3.24.6	Sympy [F]	199
3.24.7	Maxima [F]	200
3.24.8	Giac [F(-2)]	200
3.24.9	Mupad [B] (verification not implemented)	200

3.24.1 Optimal result

Integrand size = 15, antiderivative size = 63

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - (a + b) \sqrt{a + b \coth^2(x)} - \frac{1}{3} (a + b \coth^2(x))^{3/2}$$

output `(a+b)^(3/2)*arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))-1/3*(a+b*coth(x)^2)^(3/2)-(a+b)*(a+b*coth(x)^2)^(1/2)`

3.24.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.94

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - \frac{1}{3} \sqrt{a + b \coth^2(x)} (4a + 3b + b \coth^2(x))$$

input `Integrate[Coth[x]*(a + b*Coth[x]^2)^(3/2),x]`

output `(a + b)^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - (Sqrt[a + b*Coth[x]^2]*(4*a + 3*b + b*Coth[x]^2))/3`

3.24.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 26, 4153, 26, 353, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth(x) (a + b \coth^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i \tan\left(\frac{\pi}{2} + ix\right) \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{3/2} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \tan\left(ix + \frac{\pi}{2}\right) \left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{3/2} dx \\
 & \quad \downarrow \text{4153} \\
 & -i \int \frac{i \coth(x) (b \coth^2(x) + a)^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x) (a + b \coth^2(x))^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{353} \\
 & \frac{1}{2} \int \frac{(b \coth^2(x) + a)^{3/2}}{1 - \coth^2(x)} d \coth^2(x) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left((a + b) \int \frac{\sqrt{b \coth^2(x) + a}}{1 - \coth^2(x)} d \coth^2(x) - \frac{2}{3} (a + b \coth^2(x))^{3/2} \right) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left((a + b) \left((a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - 2 \sqrt{a + b \coth^2(x)} \right) - \frac{2}{3} (a + b \coth^2(x))^{3/2} \right)
 \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & \frac{1}{2} \left((a+b) \left(\frac{2(a+b) \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} dx \sqrt{b \coth^2(x) + a}}{b} - 2\sqrt{a+b \coth^2(x)} \right) - \frac{2}{3} (a+b \coth^2(x))^{3/2} \right) \\ & \downarrow 221 \\ & \frac{1}{2} \left((a+b) \left(2\sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right) - 2\sqrt{a+b \coth^2(x)} \right) - \frac{2}{3} (a+b \coth^2(x))^{3/2} \right) \end{aligned}$$

input `Int[Coth[x]*(a + b*Coth[x]^2)^(3/2),x]`

output `((-2*(a + b*Coth[x]^2)^(3/2))/3 + (a + b)*(2*Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - 2*Sqrt[a + b*Coth[x]^2]))/2`

3.24.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

```
rule 353 Int[(x_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol]
  := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[
  {a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
  Q[u, x]
```

```
rule 4153 Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
  (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
  x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
  f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m,
  n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
  nalQ[n]))
```

3.24.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. 2(51) = 102.

Time = 0.07 (sec) , antiderivative size = 473, normalized size of antiderivative = 7.51

method	result
derivativedivides	$-\frac{(b(1+\coth(x))^2-2b(1+\coth(x))+a+b)^{\frac{3}{2}}}{6} + \frac{b \left(\frac{(2b(1+\coth(x))-2b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2} \right)}{2}$
default	$-\frac{(b(1+\coth(x))^2-2b(1+\coth(x))+a+b)^{\frac{3}{2}}}{6} + \frac{b \left(\frac{(2b(1+\coth(x))-2b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2} \right)}{2}$

```
input int(coth(x)*(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)
```

output

```

-1/6*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)+1/2*b*(1/4*(2*b*(1+coth(x))
)-2*b)/b*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)+1/8*(4*(a+b)*b-4*b^2
)/b^(3/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+
b)^(1/2))-1/2*(a+b)*((b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-b^(1/2)*
ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))-
(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b
*(1+coth(x))+a+b)^(1/2))/(1+coth(x))))-1/6*(b*(coth(x)-1)^2+2*b*(coth(x)-1
)+a+b)^(3/2)-1/2*b*(1/4*(2*b*(coth(x)-1)+2*b)/b*(b*(coth(x)-1)^2+2*b*(coth
(x)-1)+a+b)^(1/2)+1/8*(4*(a+b)*b-4*b^2)/b^(3/2)*ln((b*(coth(x)-1)+b)/b^(1/
2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))-1/2*(a+b)*((b*(coth(x)-1
)^2+2*b*(coth(x)-1)+a+b)^(1/2)+b^(1/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(cot
h(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))-(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)
-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1
))

```

3.24.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 865 vs. $2(51) = 102$.

Time = 0.37 (sec) , antiderivative size = 2362, normalized size of antiderivative = 37.49

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \text{Too large to display}$$

input `integrate(coth(x)*(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")`

output `[1/12*(3*((a + b)*cosh(x)^6 + 6*(a + b)*cosh(x)*sinh(x)^5 + (a + b)*sinh(x)^6 - 3*(a + b)*cosh(x)^4 + 3*(5*(a + b)*cosh(x)^2 - a - b)*sinh(x)^4 + 4*(5*(a + b)*cosh(x)^3 - 3*(a + b)*cosh(x))*sinh(x)^3 + 3*(a + b)*cosh(x)^2 + 3*(5*(a + b)*cosh(x)^4 - 6*(a + b)*cosh(x)^2 + a + b)*sinh(x)^2 + 6*((a + b)*cosh(x)^5 - 2*(a + b)*cosh(x)^3 + (a + b)*cosh(x))*sinh(x) - a - b)*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)...`

3.24.6 Sympy [F]

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \coth(x) dx$$

input `integrate(coth(x)*(a+b*coth(x)**2)**(3/2),x)`

output `Integral((a + b*coth(x)**2)**(3/2)*coth(x), x)`

3.24.7 Maxima [F]

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \coth(x) dx$$

input `integrate(coth(x)*(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate((b*coth(x)^2 + a)^(3/2)*coth(x), x)`

3.24.8 Giac [F(-2)]

Exception generated.

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)*(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:Error: Bad Argument Type`

3.24.9 Mupad [B] (verification not implemented)

Time = 4.15 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \operatorname{atanh}\left(\frac{(a+b)^{3/2} \sqrt{b \coth(x)^2 + a}}{a^2 + 2ab + b^2}\right) (a+b)^{3/2} \\ - (a+b) \sqrt{b \coth(x)^2 + a} - \frac{(b \coth(x)^2 + a)^{3/2}}{3}$$

input `int(coth(x)*(a + b*coth(x)^2)^(3/2),x)`

output `atanh(((a + b)^(3/2)*(a + b*coth(x)^2)^(1/2))/(2*a*b + a^2 + b^2))*(a + b)^(3/2) - (a + b)*(a + b*coth(x)^2)^(1/2) - (a + b*coth(x)^2)^(3/2)/3`

3.25 $\int (a + b \operatorname{coth}^2(x))^{3/2} dx$

3.25.1	Optimal result	201
3.25.2	Mathematica [A] (verified)	201
3.25.3	Rubi [A] (verified)	202
3.25.4	Maple [B] (verified)	205
3.25.5	Fricas [B] (verification not implemented)	205
3.25.6	Sympy [F]	206
3.25.7	Maxima [F]	206
3.25.8	Giac [F(-2)]	206
3.25.9	Mupad [F(-1)]	207

3.25.1 Optimal result

Integrand size = 12, antiderivative size = 88

$$\int (a + b \operatorname{coth}^2(x))^{3/2} dx = -\frac{1}{2}\sqrt{b}(3a + 2b)\operatorname{arctanh}\left(\frac{\sqrt{b} \operatorname{coth}(x)}{\sqrt{a + b \operatorname{coth}^2(x)}}\right) + (a + b)^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a + b} \operatorname{coth}(x)}{\sqrt{a + b \operatorname{coth}^2(x)}}\right) - \frac{1}{2}b \operatorname{coth}(x)\sqrt{a + b \operatorname{coth}^2(x)}$$

output

```
(a+b)^(3/2)*arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))-1/2*(3*a+2*b)*arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))*b^(1/2)-1/2*b*coth(x)*(a+b*coth(x)^2)^(1/2)
```

3.25.2 Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.26

$$\int (a + b \operatorname{coth}^2(x))^{3/2} dx = \frac{1}{2} \left(-2(-a-b)^{3/2} \arctan\left(\frac{\operatorname{coth}(x)\sqrt{a + b \operatorname{coth}^2(x)} - \sqrt{b}\operatorname{sch}^2(x)}{\sqrt{-a-b}}\right) - b \operatorname{coth}(x)\sqrt{a + b \operatorname{coth}^2(x)} \right)$$

input `Integrate[(a + b*Coth[x]^2)^(3/2), x]`

output `(-2*(-a - b)^(3/2)*ArcTan[(Coth[x]*Sqrt[a + b*Coth[x]^2] - Sqrt[b]*Csch[x]^2)/Sqrt[-a - b]] - b*Coth[x]*Sqrt[a + b*Coth[x]^2] + Sqrt[b]*(3*a + 2*b)*Log[-(Sqrt[b]*Coth[x]) + Sqrt[a + b*Coth[x]^2]])/2`

3.25.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4144, 318, 25, 398, 224, 219, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \coth^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2 \right)^{3/2} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{(a + b \coth^2(x))^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{318} \\
 & -\frac{1}{2} \int \frac{b(3a + 2b) \coth^2(x) + a(2a + b)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{1}{2} b \coth(x) \sqrt{a + b \coth^2(x)} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \int \frac{b(3a + 2b) \coth^2(x) + a(2a + b)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{1}{2} b \coth(x) \sqrt{a + b \coth^2(x)} \\
 & \quad \downarrow \text{398} \\
 & \frac{1}{2} \left(2(a + b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - b(3a + 2b) \int \frac{1}{\sqrt{b \coth^2(x) + a}} d \coth(x) \right) - \\
 & \quad \frac{1}{2} b \coth(x) \sqrt{a + b \coth^2(x)}
 \end{aligned}$$

↓ 224

$$\frac{1}{2} \left(2(a+b)^2 \int \frac{1}{(1-\coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - b(3a+2b) \int \frac{1}{1 - \frac{b \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} \right) - \frac{1}{2} b \coth(x) \sqrt{a + b \coth^2(x)}$$

↓ 219

$$\frac{1}{2} \left(2(a+b)^2 \int \frac{1}{(1-\coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \sqrt{b}(3a+2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \right) - \frac{1}{2} b \coth(x) \sqrt{a + b \coth^2(x)}$$

↓ 291

$$\frac{1}{2} \left(2(a+b)^2 \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} - \sqrt{b}(3a+2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \right) - \frac{1}{2} b \coth(x) \sqrt{a + b \coth^2(x)}$$

↓ 219

$$\frac{1}{2} \left(2(a+b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \sqrt{b}(3a+2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \right) - \frac{1}{2} b \coth(x) \sqrt{a + b \coth^2(x)}$$

input `Int[(a + b*Coth[x]^2)^(3/2), x]`

output `(-(Sqrt[b]*(3*a + 2*b)*ArcTanh[(Sqrt[b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]) + 2*(a + b)^(3/2)*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]])/2 - (b*Coth[x]*Sqrt[a + b*Coth[x]^2])/2`

3.25.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 224 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 291 `Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*((c_) + (d_)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 318 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[d*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1)/(b*(2*(p + q) + 1))), x] + Simp[1/(b*(2*(p + q) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*(b*c*(2*(p + q) + 1) - a*d) + d*(b*c*(2*(p + 2*q - 1) + 1) - a*d*(2*(q - 1) + 1))*x^2, x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[2*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, 2, p, q, x]`
- rule 398 `Int[((e_) + (f_)*(x_)^2)/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[f/b Int[1/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/((a + b*x^2)*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4144 `Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.25.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. $2(70) = 140$.

Time = 0.12 (sec) , antiderivative size = 473, normalized size of antiderivative = 5.38

method	result
derivativedivides	$-\frac{(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}}{6} - b \left(\frac{(2b(\coth(x)-1)+2b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2} \right)$
default	$-\frac{(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}}{6} - b \left(\frac{(2b(\coth(x)-1)+2b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2} \right)$

input `int((a+b*coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output

```
-1/6*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)-1/2*b*(1/4*(2*b*(coth(x)-1)+2*b)/b*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1/8*(4*(a+b)*b-4*b^2)/b^(3/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)))-1/2*(a+b)*((b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+b^(1/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)))-(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1)))+1/6*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)-1/2*b*(1/4*(2*b*(1+coth(x))-2*b)/b*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)+1/8*(4*(a+b)*b-4*b^2)/b^(3/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)))+1/2*(a+b)*((b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-b^(1/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)))-(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x))))
```

3.25.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 947 vs. $2(70) = 140$.

Time = 0.43 (sec) , antiderivative size = 5037, normalized size of antiderivative = 57.24

$$\int (a + b \coth^2(x))^{3/2} dx = \text{Too large to display}$$

input `integrate((a+b*coth(x)^2)^(3/2),x, algorithm="fricas")`

output Too large to include

3.25.6 Sympy [F]

$$\int (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} dx$$

input `integrate((a+b*coth(x)**2)**(3/2), x)`

output `Integral((a + b*coth(x)**2)**(3/2), x)`

3.25.7 Maxima [F]

$$\int (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} dx$$

input `integrate((a+b*coth(x)^2)^(3/2), x, algorithm="maxima")`

output `integrate((b*coth(x)^2 + a)^(3/2), x)`

3.25.8 Giac [F(-2)]

Exception generated.

$$\int (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*coth(x)^2)^(3/2), x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Error: Bad Argument Type`

3.25.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{3/2} dx$$

input `int((a + b*coth(x)^2)^(3/2),x)`output `int((a + b*coth(x)^2)^(3/2), x)`

3.26 $\int (a + b \coth^2(x))^{3/2} \tanh(x) dx$

3.26.1 Optimal result	208
3.26.2 Mathematica [A] (verified)	208
3.26.3 Rubi [A] (verified)	209
3.26.4 Maple [F]	212
3.26.5 Fricas [B] (verification not implemented)	212
3.26.6 Sympy [F]	213
3.26.7 Maxima [F]	213
3.26.8 Giac [B] (verification not implemented)	214
3.26.9 Mupad [F(-1)]	215

3.26.1 Optimal result

Integrand size = 15, antiderivative size = 71

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = -a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}}\right) + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}}\right) - b\sqrt{a + b \coth^2(x)}$$

output

```
-a^(3/2)*arctanh((a+b*coth(x)^2)^(1/2)/a^(1/2))+(a+b)^(3/2)*arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))-b*(a+b*coth(x)^2)^(1/2)
```

3.26.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = -a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}}\right) + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}}\right) - b\sqrt{a + b \coth^2(x)}$$

input `Integrate[(a + b*Coth[x]^2)^(3/2)*Tanh[x],x]`

output `-(a^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]]) + (a + b)^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - b*Sqrt[a + b*Coth[x]^2]`

3.26.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 26, 4153, 26, 354, 95, 25, 174, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(x) (a + b \coth^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \left(a - b \tan \left(\frac{\pi}{2} + ix \right)^2 \right)^{3/2}}{\tan \left(\frac{\pi}{2} + ix \right)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\left(a - b \tan \left(ix + \frac{\pi}{2} \right)^2 \right)^{3/2}}{\tan \left(ix + \frac{\pi}{2} \right)} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int - \frac{i (b \coth^2(x) + a)^{3/2} \tanh(x)}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\tanh(x) (a + b \coth^2(x))^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{(b \coth^2(x) + a)^{3/2} \tanh(x)}{1 - \coth^2(x)} d \coth^2(x) \\
 & \quad \downarrow \text{95}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left(- \int - \frac{(a^2 + b(2a + b) \coth^2(x)) \tanh(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - 2b \sqrt{a + b \coth^2(x)} \right) \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left(\int \frac{(a^2 + b(2a + b) \coth^2(x)) \tanh(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - 2b \sqrt{a + b \coth^2(x)} \right) \\
& \quad \downarrow \text{174} \\
& \frac{1}{2} \left(a^2 \int \frac{\tanh(x)}{\sqrt{b \coth^2(x) + a}} d \coth^2(x) + (a + b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - 2b \sqrt{a + b \coth^2(x)} \right) \\
& \quad \downarrow \text{73} \\
& \frac{1}{2} \left(\frac{2a^2 \int \frac{1}{\frac{\coth^4(x) - a}{b} - \frac{a}{b}} d \sqrt{b \coth^2(x) + a}}{b} + \frac{2(a + b)^2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d \sqrt{b \coth^2(x) + a}}{b} - 2b \sqrt{a + b \coth^2(x)} \right) \\
& \quad \downarrow \text{221} \\
& \frac{1}{2} \left(-2a^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}} \right) + 2(a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - 2b \sqrt{a + b \coth^2(x)} \right)
\end{aligned}$$

input `Int[(a + b*Coth[x]^2)^(3/2)*Tanh[x], x]`

output `(-2*a^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]] + 2*(a + b)^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - 2*b*Sqrt[a + b*Coth[x]^2])/2`

3.26.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 95 `Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
 x_] := Simp[f*(e + f*x)^(p - 1)/(b*d*(p - 1)), x] + Simp[1/(b*d) Int[(b
 *d*e^2 - a*c*f^2 + f*(2*b*d*e - b*c*f - a*d*f)*x)*((e + f*x)^(p - 2)/((a +
 b*x)*(c + d*x))], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 1]`
- rule 174 `Int[((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_))/(((a_.) + (b_.)*(x_))*
 ((c_.) + (d_.)*(x_))), x_] := Simp[(b*g - a*h)/(b*c - a*d) Int[(e + f*x)^(p
 / (a + b*x), x], x] - Simp[(d*g - c*h)/(b*c - a*d) Int[(e + f*x)^p/(c + d
 *x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_S
 ymbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x
 , x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ
 [(m - 1)/2]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4153 `Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
 (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
 x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
 f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m,
 n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
 nalQ[n]))`

3.26.4 Maple [F]

$$\int (a + b \coth(x)^2)^{\frac{3}{2}} \tanh(x) dx$$

input `int((a+b*coth(x)^2)^(3/2)*tanh(x),x)`

output `int((a+b*coth(x)^2)^(3/2)*tanh(x),x)`

3.26.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 641 vs. 2(57) = 114.

Time = 0.43 (sec) , antiderivative size = 3949, normalized size of antiderivative = 55.62

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \text{Too large to display}$$

input `integrate((a+b*coth(x)^2)^(3/2)*tanh(x),x, algorithm="fracas")`

output `[1/4*(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x)]/(cosh(x)^6 + 6*c...`

3.26.6 Sympy [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \tanh(x) dx$$

input `integrate((a+b*coth(x)**2)**(3/2)*tanh(x),x)`

output `Integral((a + b*coth(x)**2)**(3/2)*tanh(x), x)`

3.26.7 Maxima [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \tanh(x) dx$$

input `integrate((a+b*coth(x)^2)^(3/2)*tanh(x),x, algorithm="maxima")`

output `integrate((b*coth(x)^2 + a)^(3/2)*tanh(x), x)`

3.26.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 470 vs. 2(57) = 114.

Time = 3.81 (sec) , antiderivative size = 470, normalized size of antiderivative = 6.62

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx =$$

$$\frac{2a^2 \arctan\left(-\frac{\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a+b}}}{2\sqrt{-a}}\right) \operatorname{sgn}(e^{2x} - 1)}{\sqrt{-a}}$$

$$+ \frac{1}{2}(a+b)^{3/2} \log\left(\left|\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a+b}}\right|\right) \operatorname{sgn}(e^{2x} - 1)$$

$$- \frac{1}{2}(a+b)^{3/2} \log\left(\left|-\sqrt{a+be^{2x}} + \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a+b}}\right|\right) \operatorname{sgn}(e^{2x} - 1)$$

$$- \frac{(a^2 + 2ab + b^2) \log\left(\left|-\left(\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)(a+b) + \sqrt{a+b}(a+b)\right|\right)}{2\sqrt{a+b}}$$

$$+ \frac{4\left(\left(\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)b^2 \operatorname{sgn}(e^{2x} - 1) + \sqrt{a+b}\right)}{\left(\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)^2 - 2\left(\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)\sqrt{a+b}}$$

input `integrate((a+b*coth(x)^2)^(3/2)*tanh(x),x, algorithm="giac")`

output

```
-2*a^2*arctan(-1/2*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a
*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))/sqrt(-a))*sgn(e^(2*x) - 1)/
sqrt(-a) + 1/2*(a + b)^(3/2)*log(abs(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x)
+ b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b)))*sgn(e^(2*
x) - 1) - 1/2*(a + b)^(3/2)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x)
+ b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b)))*sgn(e^(2*
x) - 1) - 1/2*(a^2 + 2*a*b + b^2)*log(abs(-(sqrt(a + b)*e^(2*x) - sqrt(a*e
^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) + sqrt(a
+ b)*(a - b)))*sgn(e^(2*x) - 1)/sqrt(a + b) + 4*((sqrt(a + b)*e^(2*x) - sq
rt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*b^2*sgn(e^(
2*x) - 1) + sqrt(a + b)*b^2*sgn(e^(2*x) - 1))/((sqrt(a + b)*e^(2*x) - sqrt
(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))^2 - 2*(sqrt(a
+ b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a
+ b))*sqrt(a + b) + a - 3*b)
```

3.26.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \int \tanh(x) (b \coth(x)^2 + a)^{3/2} dx$$

input `int(tanh(x)*(a + b*coth(x)^2)^(3/2),x)`output `int(tanh(x)*(a + b*coth(x)^2)^(3/2), x)`

3.27 $\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx$

3.27.1	Optimal result	216
3.27.2	Mathematica [B] (verified)	216
3.27.3	Rubi [A] (verified)	217
3.27.4	Maple [F]	220
3.27.5	Fricas [B] (verification not implemented)	220
3.27.6	Sympy [F]	221
3.27.7	Maxima [F]	222
3.27.8	Giac [F]	222
3.27.9	Mupad [F(-1)]	222

3.27.1 Optimal result

Integrand size = 17, antiderivative size = 77

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = -b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) - a \sqrt{a + b \coth^2(x)} \tanh(x)$$

output

```
-b^(3/2)*arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))+(a+b)^(3/2)*arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))-a*(a+b*coth(x)^2)^(1/2)*tanh(x)
```

3.27.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 180 vs. 2(77) = 154.

Time = 0.47 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.34

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \frac{\left(-\sqrt{2}b^{3/2}\sqrt{a+b}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{b}\cosh(x)}{\sqrt{-a+b+(a+b)\cosh(2x)}}\right)\cosh(x) + \sqrt{2}(a+b)^2\operatorname{arctanh}\right)}{\dots}$$

input `Integrate[(a + b*Coth[x]^2)^(3/2)*Tanh[x]^2,x]`

output `((-(Sqrt[2]*b^(3/2)*Sqrt[a + b]*ArcTanh[(Sqrt[2]*Sqrt[b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]])*Cosh[x]) + Sqrt[2]*(a + b)^2*ArcTanh[(Sqrt[2]*Sqrt[a + b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]])*Cosh[x] - a*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])*Sqrt[(-a + b + (a + b)*Cosh[2*x])*Cosh[x]^2*Tanh[x])/(Sqrt[2]*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])`

3.27.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.588$, Rules used = {3042, 25, 4153, 25, 376, 398, 224, 219, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^2(x) (a + b \coth^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{(a - b \tan(\frac{\pi}{2} + ix))^2)^{3/2}}{\tan(\frac{\pi}{2} + ix)^2} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{(a - b \tan(ix + \frac{\pi}{2}))^2)^{3/2}}{\tan(ix + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{4153} \\
 & - \int -\frac{(b \coth^2(x) + a)^{3/2} \tanh^2(x)}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\tanh^2(x) (a + b \coth^2(x))^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{376} \\
 & \int \frac{b^2 \coth^2(x) + a(a + 2b)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - a \tanh(x) \sqrt{a + b \coth^2(x)}
 \end{aligned}$$

3.27. $\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx$

$$\begin{aligned}
& \downarrow \text{398} \\
& b^2 \left(- \int \frac{1}{\sqrt{b \coth^2(x) + a}} d \coth(x) \right) + (a + b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \\
& \qquad \qquad \qquad a \tanh(x) \sqrt{a + b \coth^2(x)} \\
& \downarrow \text{224} \\
& b^2 \left(- \int \frac{1}{1 - \frac{b \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} \right) + (a + \\
& b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - a \tanh(x) \sqrt{a + b \coth^2(x)} \\
& \downarrow \text{219} \\
& (a + b)^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - b^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \\
& \qquad \qquad \qquad a \tanh(x) \sqrt{a + b \coth^2(x)} \\
& \downarrow \text{291} \\
& (a + b)^2 \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} - b^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \\
& \qquad \qquad \qquad a \tanh(x) \sqrt{a + b \coth^2(x)} \\
& \downarrow \text{219} \\
& b^{3/2} \left(- \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \right) + (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \\
& \qquad \qquad \qquad a \tanh(x) \sqrt{a + b \coth^2(x)}
\end{aligned}$$

input `Int[(a + b*Coth[x]^2)^(3/2)*Tanh[x]^2, x]`

output `-(b^(3/2)*ArcTanh[(Sqrt[b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]) + (a + b)^(3/2)*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]] - a*Sqrt[a + b*Coth[x]^2]*Tanh[x]`

3.27.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 376 `Int[((e_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[c*(e*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1))/(a*e*(m + 1)), x] - Simp[1/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*(b*c - a*d)*(m + 1) + 2*c*(b*c*(p + 1) + a*d*(q - 1)) + d*((b*c - a*d)*(m + 1) + 2*b*c*(p + q))*x^2, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`
- rule 398 `Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[f/b Int[1/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/((a + b*x^2)*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)]^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.27.4 Maple [F]

$$\int (a + b \coth(x)^2)^{\frac{3}{2}} \tanh(x)^2 dx$$

```
input int((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x)
```

```
output int((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x)
```

3.27.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 695 vs. $2(63) = 126$.

Time = 0.42 (sec) , antiderivative size = 4025, normalized size of antiderivative = 52.27

$$\int (a + b \coth^2(x))^{\frac{3}{2}} \tanh^2(x) dx = \text{Too large to display}$$

```
input integrate((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x, algorithm="fricas")
```

output `[1/4*((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*co...`

3.27.6 Sympy [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \tanh^2(x) dx$$

input `integrate((a+b*coth(x)**2)**(3/2)*tanh(x)**2,x)`

output `Integral((a + b*coth(x)**2)**(3/2)*tanh(x)**2, x)`

3.27.7 Maxima [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int (b \coth(x)^2 + a)^{3/2} \tanh(x)^2 dx$$

input `integrate((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x, algorithm="maxima")`

output `integrate((b*coth(x)^2 + a)^(3/2)*tanh(x)^2, x)`

3.27.8 Giac [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int (b \coth(x)^2 + a)^{3/2} \tanh(x)^2 dx$$

input `integrate((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x, algorithm="giac")`

output `sage0*x`

3.27.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int \tanh(x)^2 (b \coth(x)^2 + a)^{3/2} dx$$

input `int(tanh(x)^2*(a + b*coth(x)^2)^(3/2), x)`

output `int(tanh(x)^2*(a + b*coth(x)^2)^(3/2), x)`

3.28 $\int \sqrt{1 + \coth^2(x)} dx$

3.28.1	Optimal result	223
3.28.2	Mathematica [A] (verified)	223
3.28.3	Rubi [A] (verified)	224
3.28.4	Maple [B] (verified)	225
3.28.5	Fricas [B] (verification not implemented)	226
3.28.6	Sympy [F]	227
3.28.7	Maxima [F]	228
3.28.8	Giac [B] (verification not implemented)	228
3.28.9	Mupad [B] (verification not implemented)	228

3.28.1 Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \sqrt{1 + \coth^2(x)} dx = -\operatorname{arcsinh}(\coth(x)) + \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{1 + \coth^2(x)}}\right)$$

output `-arcsinh(coth(x))+arctanh(coth(x)*2^(1/2)/(1+coth(x)^2)^(1/2))*2^(1/2)`

3.28.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.94

$$\begin{aligned} & \int \sqrt{1 + \coth^2(x)} dx \\ &= \frac{\sqrt{1 + \coth^2(x)} \left(-\operatorname{arctanh}\left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}}\right) + \sqrt{2} \log\left(\sqrt{2} \cosh(x) + \sqrt{\cosh(2x)}\right) \right) \sinh(x)}{\sqrt{\cosh(2x)}} \end{aligned}$$

input `Integrate[Sqrt[1 + Coth[x]^2], x]`

output `(Sqrt[1 + Coth[x]^2]*(-ArcTanh[Cosh[x]/Sqrt[Cosh[2*x]]] + Sqrt[2]*Log[Sqrt[2]*Cosh[x] + Sqrt[Cosh[2*x]]])*Sinh[x])/Sqrt[Cosh[2*x]]`

3.28.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 4144, 301, 222, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\coth^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{1 - \tan\left(\frac{\pi}{2} + ix\right)^2} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{\sqrt{\coth^2(x) + 1}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{301} \\
 & 2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{\coth^2(x) + 1}} d \coth(x) - \int \frac{1}{\sqrt{\coth^2(x) + 1}} d \coth(x) \\
 & \quad \downarrow \text{222} \\
 & 2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{\coth^2(x) + 1}} d \coth(x) - \operatorname{arcsinh}(\coth(x)) \\
 & \quad \downarrow \text{291} \\
 & 2 \int \frac{1}{1 - \frac{2 \coth^2(x)}{\coth^2(x) + 1}} d \frac{\coth(x)}{\sqrt{\coth^2(x) + 1}} - \operatorname{arcsinh}(\coth(x)) \\
 & \quad \downarrow \text{219} \\
 & \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{\coth^2(x) + 1}}\right) - \operatorname{arcsinh}(\coth(x))
 \end{aligned}$$

input `Int[Sqrt[1 + Coth[x]^2], x]`

output `-ArcSinh[Coth[x]] + Sqrt[2]*ArcTanh[(Sqrt[2]*Coth[x])/Sqrt[1 + Coth[x]^2]]`

3.28. $\int \sqrt{1 + \coth^2(x)} dx$

3.28.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 301 `Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[b/d Int[(a + b*x^2)^(p - 1), x], x] - Simp[(b*c - a*d)/d Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4] || (EqQ[p, 2/3] && EqQ[b*c + 3*a*d, 0]))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.28.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 96 vs. $2(25) = 50$.

Time = 0.23 (sec) , antiderivative size = 97, normalized size of antiderivative = 3.13

method	result
derivativedivides	$-\frac{\sqrt{(\coth(x)-1)^2+2\coth(x)}}{2} - \operatorname{arcsinh}(\coth(x)) + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2\coth(x))\sqrt{2}}{4\sqrt{(\coth(x)-1)^2+2\coth(x)}}\right)}{2} + \sqrt{1+\coth(x)}$
default	$-\frac{\sqrt{(\coth(x)-1)^2+2\coth(x)}}{2} - \operatorname{arcsinh}(\coth(x)) + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2\coth(x))\sqrt{2}}{4\sqrt{(\coth(x)-1)^2+2\coth(x)}}\right)}{2} + \sqrt{1+\coth(x)}$

input `int((1+coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*((coth(x)-1)^2+2*coth(x))^(1/2)-arcsinh(coth(x))+1/2*2^(1/2)*arctanh(1/4*(2+2*coth(x))*2^(1/2)/((coth(x)-1)^2+2*coth(x))^(1/2))+1/2*((1+coth(x))^2-2*coth(x))^(1/2)-1/2*2^(1/2)*arctanh(1/4*(2-2*coth(x))*2^(1/2)/((1+coth(x))^2-2*coth(x))^(1/2))`

3.28.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs. $2(25) = 50$.

Time = 0.26 (sec) , antiderivative size = 683, normalized size of antiderivative = 22.03

$$\int \sqrt{1 + \coth^2(x)} dx = \text{Too large to display}$$

input `integrate((1+coth(x)^2)^(1/2),x, algorithm="fricas")`

output `1/4*sqrt(2)*log(2*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 + 3)*sinh(x)^6 + 3*cosh(x)^6 + 2*(28*cosh(x)^3 + 9*cosh(x))*sinh(x)^5 + 5*(14*cosh(x)^4 + 9*cosh(x)^2 + 1)*sinh(x)^4 + 5*cosh(x)^4 + 4*(14*cosh(x)^5 + 15*cosh(x)^3 + 5*cosh(x))*sinh(x)^3 + (28*cosh(x)^6 + 45*cosh(x)^4 + 30*cosh(x)^2 + 4)*sinh(x)^2 + 4*cosh(x)^2 + 2*(4*cosh(x)^7 + 9*cosh(x)^5 + 10*cosh(x)^3 + 4*cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^6 + 6*sqrt(2)*cosh(x)*sinh(x)^5 + sqrt(2)*sinh(x)^6 + 3*(5*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^4 + 3*sqrt(2)*cosh(x)^4 + 4*(5*sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x))*sinh(x)^3 + (15*sqrt(2)*cosh(x)^4 + 18*sqrt(2)*cosh(x)^2 + 4*sqrt(2))*sinh(x)^2 + 4*sqrt(2)*cosh(x)^2 + 2*(3*sqrt(2)*cosh(x)^5 + 6*sqrt(2)*cosh(x)^3 + 4*sqrt(2)*cosh(x))*sinh(x) + 4*sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4)/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + 1/4*sqrt(2)*log(-2*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 - sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)) - 1/2*log((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 2*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sin...`

3.28.6 Sympy [F]

$$\int \sqrt{1 + \coth^2(x)} dx = \int \sqrt{\coth^2(x) + 1} dx$$

input `integrate((1+coth(x)**2)**(1/2), x)`

output `Integral(sqrt(coth(x)**2 + 1), x)`

3.28.7 Maxima [F]

$$\int \sqrt{1 + \coth^2(x)} dx = \int \sqrt{\coth(x)^2 + 1} dx$$

input `integrate((1+coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(coth(x)^2 + 1), x)`

3.28.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(25) = 50$.

Time = 0.28 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.84

$$\int \sqrt{1 + \coth^2(x)} dx$$

$$= \frac{1}{2} \sqrt{2} \left(\sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{4x} + 1} - 2e^{2x} + 2|}{2(\sqrt{2} + \sqrt{e^{4x} + 1} - e^{2x} + 1)} \right) + \log(\sqrt{e^{4x} + 1} - e^{2x} + 1) - \log(\sqrt{e^{4x} + 1} + e^{2x} + 1) \right) - 1$$

input `integrate((1+coth(x)^2)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(2)*(sqrt(2)*log(1/2*abs(-2*sqrt(2) + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) + 2)/(sqrt(2) + sqrt(e^(4*x) + 1) - e^(2*x) + 1)) + log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) - log(sqrt(e^(4*x) + 1) + e^(2*x) + 1))*sgn(e^(2*x) - 1)`

3.28.9 Mupad [B] (verification not implemented)

Time = 2.04 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.19

$$\int \sqrt{1 + \coth^2(x)} dx = \frac{\sqrt{2} \left(\ln \left(\coth(x) + \sqrt{2} \sqrt{\coth(x)^2 + 1} + 1 \right) - \ln(\coth(x) - 1) \right)}{2} - \operatorname{asinh}(\coth(x)) + \frac{\sqrt{2} \left(\ln(\coth(x) + 1) - \ln \left(\sqrt{2} \sqrt{\coth(x)^2 + 1} - \coth(x) + 1 \right) \right)}{2}$$

3.28. $\int \sqrt{1 + \coth^2(x)} dx$

input `int((coth(x)^2 + 1)^(1/2),x)`

output `(2^(1/2)*(log(coth(x) + 2^(1/2)*(coth(x)^2 + 1)^(1/2) + 1) - log(coth(x) - 1)))/2 - asinh(coth(x)) + (2^(1/2)*(log(coth(x) + 1) - log(2^(1/2)*(coth(x)^2 + 1)^(1/2) - coth(x) + 1)))/2`

3.29 $\int \sqrt{-1 - \coth^2(x)} dx$

3.29.1	Optimal result	230
3.29.2	Mathematica [A] (verified)	230
3.29.3	Rubi [A] (verified)	231
3.29.4	Maple [B] (verified)	233
3.29.5	Fricas [C] (verification not implemented)	233
3.29.6	Sympy [F]	234
3.29.7	Maxima [F]	234
3.29.8	Giac [C] (verification not implemented)	235
3.29.9	Mupad [B] (verification not implemented)	235

3.29.1 Optimal result

Integrand size = 12, antiderivative size = 45

$$\int \sqrt{-1 - \coth^2(x)} dx = \arctan\left(\frac{\coth(x)}{\sqrt{-1 - \coth^2(x)}}\right) - \sqrt{2} \arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-1 - \coth^2(x)}}\right)$$

output `arctan(coth(x)/(-1-coth(x)^2)^(1/2))-arctan(coth(x)*2^(1/2)/(-1-coth(x)^2)^(1/2))*2^(1/2)`

3.29.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.38

$$\begin{aligned} & \int \sqrt{-1 - \coth^2(x)} dx \\ &= \frac{\sqrt{-1 - \coth^2(x)} \left(-\operatorname{arctanh}\left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}}\right) + \sqrt{2} \log\left(\sqrt{2} \cosh(x) + \sqrt{\cosh(2x)}\right) \right) \sinh(x)}{\sqrt{\cosh(2x)}} \end{aligned}$$

input `Integrate[Sqrt[-1 - Coth[x]^2], x]`

output `(Sqrt[-1 - Coth[x]^2]*(-ArcTanh[Cosh[x]/Sqrt[Cosh[2*x]]] + Sqrt[2]*Log[Sqrt[2]*Cosh[x] + Sqrt[Cosh[2*x]]])*Sinh[x])/Sqrt[Cosh[2*x]]`

3.29.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 4144, 301, 224, 216, 291, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{-\coth^2(x) - 1} \, dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-1 + \tan\left(\frac{\pi}{2} + ix\right)^2} \, dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{\sqrt{-\coth^2(x) - 1}}{1 - \coth^2(x)} \, d\coth(x) \\
 & \quad \downarrow \text{301} \\
 & \int \frac{1}{\sqrt{-\coth^2(x) - 1}} \, d\coth(x) - 2 \int \frac{1}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} \, d\coth(x) \\
 & \quad \downarrow \text{224} \\
 & \int \frac{1}{\frac{\coth^2(x)}{-\coth^2(x)-1} + 1} \, d\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} - 2 \int \frac{1}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} \, d\coth(x) \\
 & \quad \downarrow \text{216} \\
 & \arctan\left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}}\right) - 2 \int \frac{1}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} \, d\coth(x) \\
 & \quad \downarrow \text{291} \\
 & \arctan\left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}}\right) - 2 \int \frac{1}{\frac{2\coth^2(x)}{-\coth^2(x)-1} + 1} \, d\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} \\
 & \quad \downarrow \text{216} \\
 & \arctan\left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}}\right) - \sqrt{2} \arctan\left(\frac{\sqrt{2}\coth(x)}{\sqrt{-\coth^2(x) - 1}}\right)
 \end{aligned}$$

input `Int[Sqrt[-1 - Coth[x]^2], x]`

output `ArcTan[Coth[x]/Sqrt[-1 - Coth[x]^2]] - Sqrt[2]*ArcTan[(Sqrt[2]*Coth[x])/Sqrt[-1 - Coth[x]^2]]`

3.29.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 301 `Int[((a_) + (b_.)*(x_)^2)^(p_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[b/d Int[(a + b*x^2)^(p - 1), x], x] - Simp[(b*c - a*d)/d Int[(a + b*x^2)^(p - 1)/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[p, 0] && (EqQ[p, 1/2] || EqQ[Denominator[p], 4] || (EqQ[p, 2/3] && EqQ[b*c + 3*a*d, 0]))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.29.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(37) = 74.

Time = 0.20 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.16

method	result
derivativedivides	$\frac{\sqrt{-(1+\coth(x))^2+2\coth(x)}}{2} + \frac{\arctan\left(\frac{\coth(x)}{\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2} - \frac{\sqrt{2}\arctan\left(\frac{(-2+2\coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2}$
default	$\frac{\sqrt{-(1+\coth(x))^2+2\coth(x)}}{2} + \frac{\arctan\left(\frac{\coth(x)}{\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2} - \frac{\sqrt{2}\arctan\left(\frac{(-2+2\coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2}$

input `int((-1-coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/2*(-(1+\coth(x))^2+2*\coth(x))^(1/2)+1/2*\arctan(\coth(x)/(-(1+\coth(x))^2+2* \\ & \coth(x))^(1/2))-1/2*2^(1/2)*\arctan(1/4*(-2+2*\coth(x))*2^(1/2)/(-(1+\coth(x)) \\ &)^2+2*\coth(x))^(1/2))-1/2*(-(\coth(x)-1)^2-2*\coth(x))^(1/2)+1/2*\arctan(\coth \\ & (x)/(-(\coth(x)-1)^2-2*\coth(x))^(1/2))+1/2*2^(1/2)*\arctan(1/4*(-2-2*\coth(x)) \\ &)*2^(1/2)/(-(\coth(x)-1)^2-2*\coth(x))^(1/2)) \end{aligned}$$

3.29.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 226, normalized size of antiderivative = 5.02

$$\begin{aligned} & \int \sqrt{-1 - \coth^2(x)} dx \\ & = -\frac{1}{4} \sqrt{-2} \log \left(-\left(\sqrt{-2} \sqrt{-2e^{4x} - 2} + 2e^{2x} - 2 \right) e^{(-2x)} \right) \\ & \quad + \frac{1}{4} \sqrt{-2} \log \left(\left(\sqrt{-2} \sqrt{-2e^{4x} - 2} - 2e^{2x} + 2 \right) e^{(-2x)} \right) \\ & \quad + \frac{1}{4} \sqrt{-2} \log \left(-2 \left(\sqrt{-2e^{4x} - 2} (e^{2x} + 2) + \sqrt{-2}e^{4x} + \sqrt{-2}e^{2x} + 2\sqrt{-2} \right) e^{(-4x)} \right) \\ & \quad - \frac{1}{4} \sqrt{-2} \log \left(-2 \left(\sqrt{-2e^{4x} - 2} (e^{2x} + 2) - \sqrt{-2}e^{4x} - \sqrt{-2}e^{2x} - 2\sqrt{-2} \right) e^{(-4x)} \right) \\ & \quad + \frac{1}{2} i \log \left(-4 \left(i \sqrt{-2e^{4x} - 2} + e^{2x} + 1 \right) e^{(-2x)} \right) \\ & \quad - \frac{1}{2} i \log \left(-4 \left(-i \sqrt{-2e^{4x} - 2} + e^{2x} + 1 \right) e^{(-2x)} \right) \end{aligned}$$

input `integrate((-1-coth(x)^2)^(1/2),x, algorithm="fricas")`

output `-1/4*sqrt(-2)*log(-(sqrt(-2)*sqrt(-2*e^(4*x) - 2) + 2*e^(2*x) - 2)*e^(-2*x)) + 1/4*sqrt(-2)*log((sqrt(-2)*sqrt(-2*e^(4*x) - 2) - 2*e^(2*x) + 2)*e^(-2*x)) + 1/4*sqrt(-2)*log(-2*(sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) + sqrt(-2)*e^(4*x) + sqrt(-2)*e^(2*x) + 2*sqrt(-2))*e^(-4*x)) - 1/4*sqrt(-2)*log(-2*(sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) - sqrt(-2)*e^(4*x) - sqrt(-2)*e^(2*x) - 2*sqrt(-2))*e^(-4*x)) + 1/2*I*log(-4*(I*sqrt(-2*e^(4*x) - 2) + e^(2*x) + 1)*e^(-2*x)) - 1/2*I*log(-4*(-I*sqrt(-2*e^(4*x) - 2) + e^(2*x) + 1)*e^(-2*x))`

3.29.6 Sympy [F]

$$\int \sqrt{-1 - \coth^2(x)} dx = \int \sqrt{-\coth^2(x) - 1} dx$$

input `integrate((-1-coth(x)**2)**(1/2),x)`

output `Integral(sqrt(-coth(x)**2 - 1), x)`

3.29.7 Maxima [F]

$$\int \sqrt{-1 - \coth^2(x)} dx = \int \sqrt{-\coth(x)^2 - 1} dx$$

input `integrate((-1-coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-coth(x)^2 - 1), x)`

3.29.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.76

$$\int \sqrt{-1 - \coth^2(x)} dx = -\frac{1}{2} \sqrt{2} \left(i \sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{4x} + 1} - 2e^{2x} + 2|}{2(\sqrt{2} + \sqrt{e^{4x} + 1} - e^{2x} + 1)} \right) + i \log(\sqrt{e^{4x} + 1} - e^{2x} + 1) - i \log(\sqrt{e^{4x} + 1} + e^{2x} + 1) \right)$$

input `integrate((-1-coth(x)^2)^(1/2),x, algorithm="giac")`

output `-1/2*sqrt(2)*(I*sqrt(2)*log(1/2*abs(-2*sqrt(2) + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) + 2)/(sqrt(2) + sqrt(e^(4*x) + 1) - e^(2*x) + 1)) + I*log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) - I*log(sqrt(e^(4*x) + 1) + e^(2*x) + 1))*sgn(-e^(2*x) + 1)`

3.29.9 Mupad [B] (verification not implemented)

Time = 2.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96

$$\int \sqrt{-1 - \coth^2(x)} dx = -\sqrt{2} \operatorname{atan} \left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth(x)^2 - 1}} \right) - \ln \left(\coth(x) - \sqrt{-\coth(x)^2 - 1} \operatorname{li} \right) \operatorname{li}$$

input `int((- coth(x)^2 - 1)^(1/2),x)`

output `- log(coth(x) - (- coth(x)^2 - 1)^(1/2)*1i)*1i - 2^(1/2)*atan((2^(1/2)*coth(x))/(- coth(x)^2 - 1)^(1/2))`

3.30 $\int (1 + \coth^2(x))^{3/2} dx$

3.30.1 Optimal result	236
3.30.2 Mathematica [B] (verified)	236
3.30.3 Rubi [A] (verified)	237
3.30.4 Maple [B] (verified)	239
3.30.5 Fricas [B] (verification not implemented)	240
3.30.6 Sympy [F]	241
3.30.7 Maxima [F]	241
3.30.8 Giac [B] (verification not implemented)	241
3.30.9 Mupad [B] (verification not implemented)	242

3.30.1 Optimal result

Integrand size = 10, antiderivative size = 50

$$\int (1 + \coth^2(x))^{3/2} dx = -\frac{5}{2} \operatorname{arcsinh}(\coth(x)) + 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{1 + \coth^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{1 + \coth^2(x)}$$

```
output -5/2*arcsinh(coth(x))+2*arctanh(coth(x)*2^(1/2)/(1+coth(x)^2)^(1/2))*2^(1/2)-1/2*coth(x)*(1+coth(x)^2)^(1/2)
```

3.30.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 116 vs. 2(50) = 100.

Time = 0.42 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.32

$$\int (1 + \coth^2(x))^{3/2} dx = -\frac{1}{8} (1 + \coth^2(x))^{3/2} \operatorname{sech}^2(2x) \left(16 \operatorname{arctanh}\left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}}\right) \sqrt{\cosh(2x)} \sinh^3(x) + 4 \left(\operatorname{arctan}\left(\frac{\cosh(x)}{\sqrt{-\cosh(2x)}}\right) \sqrt{-\cosh(2x)} \right) \right)$$

```
input Integrate[(1 + Coth[x]^2)^(3/2), x]
```

output $-1/8*((1 + \text{Coth}[x]^2)^{(3/2)}*\text{Sech}[2*x]^2*(16*\text{ArcTanh}[\text{Cosh}[x]/\text{Sqrt}[\text{Cosh}[2*x]]]*\text{Sqrt}[\text{Cosh}[2*x]]*\text{Sinh}[x]^3 + 4*(\text{ArcTan}[\text{Cosh}[x]/\text{Sqrt}[-\text{Cosh}[2*x]])*\text{Sqrt}[-\text{Cosh}[2*x]] - 4*\text{Sqrt}[2]*\text{Sqrt}[\text{Cosh}[2*x]]*\text{Log}[\text{Sqrt}[2]*\text{Cosh}[x] + \text{Sqrt}[\text{Cosh}[2*x]])*\text{Sinh}[x]^3 + \text{Sinh}[4*x]))$

3.30.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {3042, 4144, 318, 25, 398, 222, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (\coth^2(x) + 1)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(1 - \tan\left(\frac{\pi}{2} + ix\right)\right)^2)^{3/2} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{(\coth^2(x) + 1)^{3/2}}{1 - \coth^2(x)} d \coth(x) \\
 & \quad \downarrow \text{318} \\
 & -\frac{1}{2} \int -\frac{5 \coth^2(x) + 3}{(1 - \coth^2(x)) \sqrt{\coth^2(x) + 1}} d \coth(x) - \frac{1}{2} \sqrt{\coth^2(x) + 1} \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \int \frac{5 \coth^2(x) + 3}{(1 - \coth^2(x)) \sqrt{\coth^2(x) + 1}} d \coth(x) - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) + 1} \\
 & \quad \downarrow \text{398} \\
 & \frac{1}{2} \left(8 \int \frac{1}{(1 - \coth^2(x)) \sqrt{\coth^2(x) + 1}} d \coth(x) - 5 \int \frac{1}{\sqrt{\coth^2(x) + 1}} d \coth(x) \right) - \\
 & \quad \frac{1}{2} \coth(x) \sqrt{\coth^2(x) + 1} \\
 & \quad \downarrow \text{222}
 \end{aligned}$$

$$\frac{1}{2} \left(8 \int \frac{1}{(1 - \coth^2(x)) \sqrt{\coth^2(x) + 1}} d \coth(x) - 5 \operatorname{arcsinh}(\coth(x)) \right) - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) + 1}$$

↓ 291

$$\frac{1}{2} \left(8 \int \frac{1}{1 - \frac{2 \coth^2(x)}{\coth^2(x) + 1}} d \frac{\coth(x)}{\sqrt{\coth^2(x) + 1}} - 5 \operatorname{arcsinh}(\coth(x)) \right) - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) + 1}$$

↓ 219

$$\frac{1}{2} \left(4\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{2} \coth(x)}{\sqrt{\coth^2(x) + 1}} \right) - 5 \operatorname{arcsinh}(\coth(x)) \right) - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) + 1}$$

input `Int[(1 + Coth[x]^2)^(3/2), x]`

output `(-5*ArcSinh[Coth[x]] + 4*Sqrt[2]*ArcTanh[(Sqrt[2]*Coth[x])/Sqrt[1 + Coth[x]^2]])/2 - (Coth[x]*Sqrt[1 + Coth[x]^2])/2`

3.30.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

```
rule 318 Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Sim
p[d*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1)/(b*(2*(p + q) + 1))), x] + S
imp[1/(b*(2*(p + q) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*(b
*c*(2*(p + q) + 1) - a*d) + d*(b*c*(2*(p + 2*q - 1) + 1) - a*d*(2*(q - 1) +
1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && G
tQ[q, 1] && NeQ[2*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c,
d, 2, p, q, x]
```

```
rule 398 Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*Sqrt[(c_) + (d_.)*(x_)^2])
, x_Symbol] := Simp[f/b Int[1/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/
b Int[1/((a + b*x^2)*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}
, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4144 Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*
(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] ||
EqQ[n^2, 16])
```

3.30.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. 2(38) = 76.

Time = 0.10 (sec) , antiderivative size = 158, normalized size of antiderivative = 3.16

method	result
derivativedivides	$\frac{\left((1 + \coth(x))^2 - 2 \coth(x) \right)^{\frac{3}{2}}}{6} - \frac{\coth(x) \sqrt{(1 + \coth(x))^2 - 2 \coth(x)}}{4} - \frac{5 \operatorname{arcsinh}(\coth(x))}{2} + \sqrt{(1 + \coth(x))}$
default	$\frac{\left((1 + \coth(x))^2 - 2 \coth(x) \right)^{\frac{3}{2}}}{6} - \frac{\coth(x) \sqrt{(1 + \coth(x))^2 - 2 \coth(x)}}{4} - \frac{5 \operatorname{arcsinh}(\coth(x))}{2} + \sqrt{(1 + \coth(x))}$

```
input int((1+coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)
```



```
output 1/6*((1+coth(x))^2-2*coth(x))^(3/2)-1/4*coth(x)*((1+coth(x))^2-2*coth(x))^(1/2)-5/2*arcsinh(coth(x))+((1+coth(x))^2-2*coth(x))^(1/2)-2^(1/2)*arctanh(1/4*(2-2*coth(x))*2^(1/2)/((1+coth(x))^2-2*coth(x))^(1/2))-1/6*((coth(x)-1)^2+2*coth(x))^(3/2)-1/4*coth(x)*((coth(x)-1)^2+2*coth(x))^(1/2)-((coth(x)-1)^2+2*coth(x))^(1/2)+2^(1/2)*arctanh(1/4*(2+2*coth(x))*2^(1/2)/((coth(x)-1)^2+2*coth(x))^(1/2)))
```

3.30.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1043 vs. $2(38) = 76$.

Time = 0.28 (sec) , antiderivative size = 1043, normalized size of antiderivative = 20.86

$$\int (1 + \coth^2(x))^{3/2} dx = \text{Too large to display}$$

```
input integrate((1+coth(x)^2)^(3/2),x, algorithm="fracas")
```

```
output 1/4*(2*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 - sqrt(2))*sinh(x)^2 - 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 - sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(2*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 + 3)*sinh(x)^6 + 3*cosh(x)^6 + 2*(28*cosh(x)^3 + 9*cosh(x))*sinh(x)^5 + 5*(14*cosh(x)^4 + 9*cosh(x)^2 + 1)*sinh(x)^4 + 5*cosh(x)^4 + 4*(14*cosh(x)^5 + 15*cosh(x)^3 + 5*cosh(x))*sinh(x)^3 + (28*cosh(x)^6 + 45*cosh(x)^4 + 30*cosh(x)^2 + 4)*sinh(x)^2 + 4*cosh(x)^2 + 2*(4*cosh(x)^7 + 9*cosh(x)^5 + 10*cosh(x)^3 + 4*cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^6 + 6*sqrt(2)*cosh(x)*sinh(x)^5 + sqrt(2)*sinh(x)^6 + 3*(5*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^4 + 3*sqrt(2)*cosh(x)^4 + 4*(5*sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x))*sinh(x)^3 + (15*sqrt(2)*cosh(x)^4 + 18*sqrt(2)*cosh(x)^2 + 4*sqrt(2))*sinh(x)^2 + 4*sqrt(2)*cosh(x)^2 + 2*(3*sqrt(2)*cosh(x)^5 + 6*sqrt(2)*cosh(x)^3 + 4*sqrt(2)*cosh(x))*sinh(x) + 4*sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4)/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + 2*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 - sqrt(2))*sinh(x)^2 - 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 - sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-2*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sin...
```

3.30.6 Sympy [F]

$$\int (1 + \coth^2(x))^{3/2} dx = \int (\coth^2(x) + 1)^{\frac{3}{2}} dx$$

input `integrate((1+coth(x)**2)**(3/2),x)`

output `Integral((coth(x)**2 + 1)**(3/2), x)`

3.30.7 Maxima [F]

$$\int (1 + \coth^2(x))^{3/2} dx = \int (\coth(x)^2 + 1)^{\frac{3}{2}} dx$$

input `integrate((1+coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate((coth(x)^2 + 1)^(3/2), x)`

3.30.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 265 vs. $2(38) = 76$.

Time = 0.29 (sec) , antiderivative size = 265, normalized size of antiderivative = 5.30

$$\int (1 + \coth^2(x))^{3/2} dx = \frac{1}{4} \sqrt{2} \left(5 \sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{4x} + 1} - 2e^{2x} + 2|}{2(\sqrt{2} + \sqrt{e^{4x} + 1} - e^{2x} + 1)} \right) \operatorname{sgn}(e^{2x} - 1) + 4 \log(\sqrt{e^{4x} + 1}) \right)$$

input `integrate((1+coth(x)^2)^(3/2),x, algorithm="giac")`

output $\frac{1}{4}\sqrt{2}\left(5\sqrt{2}\log\left(\frac{1}{2}\left|\sqrt{2}-2\sqrt{e^{4x}+1}-2e^{2x}\right|\right)+2\sqrt{e^{4x}+1}-2e^{2x}\right)+2\sqrt{e^{4x}+1}-2e^{2x}\right)/\left(\sqrt{2}+\sqrt{e^{4x}+1}-e^{2x}\right)+\operatorname{sgn}\left(e^{2x}-1\right)+4\log\left(\sqrt{e^{4x}+1}-e^{2x}\right)+\operatorname{sgn}\left(e^{2x}-1\right)-4\log\left(\sqrt{e^{4x}+1}-e^{2x}\right)+\operatorname{sgn}\left(e^{2x}-1\right)-4\log\left(-\sqrt{e^{4x}+1}+e^{2x}\right)+\operatorname{sgn}\left(e^{2x}-1\right)-4\left(3\left(\sqrt{e^{4x}+1}-e^{2x}\right)^3\operatorname{sgn}\left(e^{2x}-1\right)+\left(\sqrt{e^{4x}+1}-e^{2x}\right)^2\operatorname{sgn}\left(e^{2x}-1\right)-\left(\sqrt{e^{4x}+1}-e^{2x}\right)\operatorname{sgn}\left(e^{2x}-1\right)+\operatorname{sgn}\left(e^{2x}-1\right)\right)/\left(\left(\sqrt{e^{4x}+1}-e^{2x}\right)^2+2\sqrt{e^{4x}+1}-2e^{2x}-1\right)^2$

3.30.9 Mupad [B] (verification not implemented)

Time = 2.12 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.56

$$\int (1 + \coth^2(x))^{3/2} dx = \sqrt{2} \left(\ln \left(\coth(x) + \sqrt{2} \sqrt{\coth(x)^2 + 1} + 1 \right) - \ln(\coth(x) - 1) \right) - \frac{\coth(x) \sqrt{\coth(x)^2 + 1}}{2} - \frac{5 \operatorname{asinh}(\coth(x))}{2} + \sqrt{2} \left(\ln(\coth(x) + 1) - \ln \left(\sqrt{2} \sqrt{\coth(x)^2 + 1} - \coth(x) + 1 \right) \right)$$

input `int((coth(x)^2 + 1)^(3/2),x)`

output $2^{1/2}\left(\log(\coth(x)+2^{1/2}\left(\coth(x)^2+1\right)^{1/2}+1)-\log(\coth(x)-1)\right)-\left(\coth(x)\left(\coth(x)^2+1\right)^{1/2}\right)/2-\left(5\operatorname{asinh}(\coth(x))\right)/2+2^{1/2}\left(\log(\coth(x)+1)-\log\left(2^{1/2}\left(\coth(x)^2+1\right)^{1/2}-\coth(x)+1\right)\right)$

3.31 $\int (-1 - \coth^2(x))^{3/2} dx$

3.31.1	Optimal result	243
3.31.2	Mathematica [A] (verified)	243
3.31.3	Rubi [A] (verified)	244
3.31.4	Maple [B] (verified)	247
3.31.5	Fricas [C] (verification not implemented)	247
3.31.6	Sympy [F]	248
3.31.7	Maxima [F]	248
3.31.8	Giac [C] (verification not implemented)	249
3.31.9	Mupad [F(-1)]	249

3.31.1 Optimal result

Integrand size = 12, antiderivative size = 67

$$\int (-1 - \coth^2(x))^{3/2} dx = -\frac{5}{2} \arctan\left(\frac{\coth(x)}{\sqrt{-1 - \coth^2(x)}}\right) + 2\sqrt{2} \arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-1 - \coth^2(x)}}\right) + \frac{1}{2} \coth(x) \sqrt{-1 - \coth^2(x)}$$

output `-5/2*arctan(coth(x)/(-1-coth(x)^2)^(1/2))+2*arctan(coth(x)*2^(1/2)/(-1-coth(x)^2)^(1/2))*2^(1/2)+1/2*coth(x)*(-1-coth(x)^2)^(1/2)`

3.31.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.76

$$\int (-1 - \coth^2(x))^{3/2} dx = -\frac{1}{8}(-1 - \coth^2(x))^{3/2} \operatorname{sech}^2(2x) \left(16 \operatorname{arctanh}\left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}}\right) \sqrt{\cosh(2x)} \sinh^3(x) + 4 \left(\arctan\left(\frac{\cosh(x)}{\sqrt{-\cosh(2x)}}\right) \sqrt{-\cosh(2x)} \right) \right)$$

input `Integrate[(-1 - Coth[x]^2)^(3/2), x]`

output `-1/8*((-1 - Coth[x]^2)^(3/2)*Sech[2*x]^2*(16*ArcTanh[Cosh[x]/Sqrt[Cosh[2*x]]]*Sqrt[Cosh[2*x]]*Sinh[x]^3 + 4*(ArcTan[Cosh[x]/Sqrt[-Cosh[2*x]]]*Sqrt[-Cosh[2*x]] - 4*Sqrt[2]*Sqrt[Cosh[2*x]]*Log[Sqrt[2]*Cosh[x] + Sqrt[Cosh[2*x]]])*Sinh[x]^3 + Sinh[4*x])`

3.31.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4144, 318, 25, 398, 224, 216, 291, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (-\coth^2(x) - 1)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(-1 + \tan\left(\frac{\pi}{2} + ix\right)\right)^{3/2} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{(-\coth^2(x) - 1)^{3/2}}{1 - \coth^2(x)} d\coth(x) \\
 & \quad \downarrow \text{318} \\
 & \frac{1}{2} \coth(x) \sqrt{-\coth^2(x) - 1} - \frac{1}{2} \int -\frac{5\coth^2(x) + 3}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} d\coth(x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \int \frac{5\coth^2(x) + 3}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} d\coth(x) + \frac{1}{2} \sqrt{-\coth^2(x) - 1} \coth(x) \\
 & \quad \downarrow \text{398} \\
 & \frac{1}{2} \left(8 \int \frac{1}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} d\coth(x) - 5 \int \frac{1}{\sqrt{-\coth^2(x) - 1}} d\coth(x) \right) + \\
 & \quad \frac{1}{2} \sqrt{-\coth^2(x) - 1} \coth(x) \\
 & \quad \downarrow \text{224}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left(8 \int \frac{1}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} d\coth(x) - 5 \int \frac{1}{\frac{\coth^2(x)}{-\coth^2(x)-1} + 1} d \frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} \right) + \\
& \qquad \qquad \qquad \frac{1}{2} \sqrt{-\coth^2(x) - 1} \coth(x) \\
& \qquad \qquad \qquad \downarrow \text{216} \\
& \frac{1}{2} \left(8 \int \frac{1}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} d\coth(x) - 5 \arctan \left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} \right) \right) + \\
& \qquad \qquad \qquad \frac{1}{2} \sqrt{-\coth^2(x) - 1} \coth(x) \\
& \qquad \qquad \qquad \downarrow \text{291} \\
& \frac{1}{2} \left(8 \int \frac{1}{\frac{2\coth^2(x)}{-\coth^2(x)-1} + 1} d \frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} - 5 \arctan \left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} \right) \right) + \\
& \qquad \qquad \qquad \frac{1}{2} \sqrt{-\coth^2(x) - 1} \coth(x) \\
& \qquad \qquad \qquad \downarrow \text{216} \\
& \frac{1}{2} \left(4\sqrt{2} \arctan \left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth^2(x) - 1}} \right) - 5 \arctan \left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} \right) \right) + \\
& \qquad \qquad \qquad \frac{1}{2} \coth(x) \sqrt{-\coth^2(x) - 1}
\end{aligned}$$

input `Int[(-1 - Coth[x]^2)^(3/2), x]`

output `(-5*ArcTan[Coth[x]/Sqrt[-1 - Coth[x]^2]] + 4*Sqrt[2]*ArcTan[(Sqrt[2]*Coth[x])/Sqrt[-1 - Coth[x]^2]])/2 + (Coth[x]*Sqrt[-1 - Coth[x]^2])/2`

3.31.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

- rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 318 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[d*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1)/(b*(2*(p + q) + 1))), x] + Simp[1/(b*(2*(p + q) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*(b*c*(2*(p + q) + 1) - a*d) + d*(b*c*(2*(p + 2*q - 1) + 1) - a*d*(2*(q - 1) + 1))*x^2, x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[2*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, 2, p, q, x]`
- rule 398 `Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := Simp[f/b Int[1/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/((a + b*x^2)*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.31.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(53) = 106.

Time = 0.11 (sec) , antiderivative size = 211, normalized size of antiderivative = 3.15

method	result
derivativedivides	$-\frac{(-\coth(x)-1)^2-2\coth(x)}{6}^{\frac{3}{2}} + \frac{\coth(x)\sqrt{-(\coth(x)-1)^2-2\coth(x)}}{4} - \frac{5\arctan\left(\frac{\coth(x)}{\sqrt{-(\coth(x)-1)^2-2\coth(x)}}\right)}{4}$
default	$-\frac{(-\coth(x)-1)^2-2\coth(x)}{6}^{\frac{3}{2}} + \frac{\coth(x)\sqrt{-(\coth(x)-1)^2-2\coth(x)}}{4} - \frac{5\arctan\left(\frac{\coth(x)}{\sqrt{-(\coth(x)-1)^2-2\coth(x)}}\right)}{4}$

input `int((-1-coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/6*(-(\coth(x)-1)^2-2*\coth(x))^(3/2)+1/4*\coth(x)*(-(\coth(x)-1)^2-2*\coth(x)) \\ &)^(1/2)-5/4*\arctan(\coth(x)/(-(\coth(x)-1)^2-2*\coth(x))^(1/2))+(-(\coth(x)-1) \\ &)^2-2*\coth(x))^(1/2)-2^(1/2)*\arctan(1/4*(-2-2*\coth(x))*2^(1/2)/(-(\coth(x)-1) \\ &)^2-2*\coth(x))^(1/2))+1/6*(-(1+\coth(x))^2+2*\coth(x))^(3/2)+1/4*\coth(x)*(- \\ & (1+\coth(x))^2+2*\coth(x))^(1/2)-5/4*\arctan(\coth(x)/(-(1+\coth(x))^2+2*\coth(x) \\ &))^(1/2))-(-(1+\coth(x))^2+2*\coth(x))^(1/2)+2^(1/2)*\arctan(1/4*(-2+2*\coth(x) \\ &))*2^(1/2)/(-(1+\coth(x))^2+2*\coth(x))^(1/2)) \end{aligned}$$

3.31.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 361, normalized size of antiderivative = 5.39

$$\int (-1 - \coth^2(x))^{3/2} dx = \frac{2(\sqrt{-2}e^{4x} - 2\sqrt{-2}e^{2x} + \sqrt{-2}) \log\left(2\left(\sqrt{-2}\sqrt{-2}e^{4x} - 2 + 2e^{2x} - 2\right)e^{-2x}\right)}{2}$$

input `integrate((-1-coth(x)^2)^(3/2),x, algorithm="fracas")`

output `1/4*(2*(sqrt(-2)*e^(4*x) - 2*sqrt(-2)*e^(2*x) + sqrt(-2))*log(2*(sqrt(-2)*sqrt(-2*e^(4*x) - 2) + 2*e^(2*x) - 2)*e^(-2*x)) - 2*(sqrt(-2)*e^(4*x) - 2*sqrt(-2)*e^(2*x) + sqrt(-2))*log(-2*(sqrt(-2)*sqrt(-2*e^(4*x) - 2) - 2*e^(2*x) + 2)*e^(-2*x)) - 5*(I*e^(4*x) - 2*I*e^(2*x) + I)*log(-4*(I*sqrt(-2*e^(4*x) - 2) + e^(2*x) + 1)*e^(-2*x)) - 5*(-I*e^(4*x) + 2*I*e^(2*x) - I)*log(-4*(-I*sqrt(-2*e^(4*x) - 2) + e^(2*x) + 1)*e^(-2*x)) - 2*(sqrt(-2)*e^(4*x) - 2*sqrt(-2)*e^(2*x) + sqrt(-2))*log(4*(sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) + sqrt(-2)*e^(4*x) + sqrt(-2)*e^(2*x) + 2*sqrt(-2))*e^(-4*x)) + 2*(sqrt(-2)*e^(4*x) - 2*sqrt(-2)*e^(2*x) + sqrt(-2))*log(4*(sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) - sqrt(-2)*e^(4*x) - sqrt(-2)*e^(2*x) - 2*sqrt(-2))*e^(-4*x)) + 2*sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 1))/(e^(4*x) - 2*e^(2*x) + 1)`

3.31.6 Sympy [F]

$$\int (-1 - \coth^2(x))^{3/2} dx = \int (-\coth^2(x) - 1)^{\frac{3}{2}} dx$$

input `integrate((-1-coth(x)**2)**(3/2), x)`

output `Integral((-coth(x)**2 - 1)**(3/2), x)`

3.31.7 Maxima [F]

$$\int (-1 - \coth^2(x))^{3/2} dx = \int (-\coth(x)^2 - 1)^{\frac{3}{2}} dx$$

input `integrate((-1-coth(x)^2)^(3/2), x, algorithm="maxima")`

output `integrate((-coth(x)^2 - 1)^(3/2), x)`

3.31.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 285, normalized size of antiderivative = 4.25

$$\int (-1 - \coth^2(x))^{3/2} dx =$$

$$-\frac{1}{4}\sqrt{2} \left(-5i\sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{4x} + 1} - 2e^{2x} + 2|}{2(\sqrt{2} + \sqrt{e^{4x} + 1} - e^{2x} + 1)} \right) \operatorname{sgn}(-e^{2x} + 1) - 4i \log(\sqrt{e^{4x} + 1} - e^{2x}) \right)$$

input `integrate((-1-coth(x)^2)^(3/2),x, algorithm="giac")`

output `-1/4*sqrt(2)*(-5*I*sqrt(2)*log(1/2*abs(-2*sqrt(2) + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) + 2)/(sqrt(2) + sqrt(e^(4*x) + 1) - e^(2*x) + 1))*sgn(-e^(2*x) + 1) - 4*I*log(sqrt(e^(4*x) + 1) - e^(2*x))*sgn(-e^(2*x) + 1) + 4*I*log(sqrt(e^(4*x) + 1) - e^(2*x))*sgn(-e^(2*x) + 1) + 4*I*log(-sqrt(e^(4*x) + 1) + e^(2*x) + 1)*sgn(-e^(2*x) + 1) + 4*(3*I*(sqrt(e^(4*x) + 1) - e^(2*x))^3*sgn(-e^(2*x) + 1) + I*(sqrt(e^(4*x) + 1) - e^(2*x))^2*sgn(-e^(2*x) + 1) + (-I*sqrt(e^(4*x) + 1) + I*e^(2*x))*sgn(-e^(2*x) + 1) + I*sgn(-e^(2*x) + 1))/((sqrt(e^(4*x) + 1) - e^(2*x))^2 + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) - 1)^2)`

3.31.9 Mupad [F(-1)]

Timed out.

$$\int (-1 - \coth^2(x))^{3/2} dx = \int (-\coth(x)^2 - 1)^{3/2} dx$$

input `int((- coth(x)^2 - 1)^(3/2),x)`

output `int((- coth(x)^2 - 1)^(3/2), x)`

3.32
$$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx$$

3.32.1	Optimal result	250
3.32.2	Mathematica [A] (verified)	250
3.32.3	Rubi [A] (verified)	251
3.32.4	Maple [B] (verified)	253
3.32.5	Fricas [B] (verification not implemented)	254
3.32.6	Sympy [F]	254
3.32.7	Maxima [F]	255
3.32.8	Giac [F(-2)]	255
3.32.9	Mupad [B] (verification not implemented)	255

3.32.1 Optimal result

Integrand size = 17, antiderivative size = 47

$$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b \coth^2(x)}}{b}$$

output `arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(1/2)-(a+b*coth(x)^2)^(1/2)/b`

3.32.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b \coth^2(x)}}{b}$$

input `Integrate[Coth[x]^3/Sqrt[a + b*Coth[x]^2],x]`

output `ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/Sqrt[a + b] - Sqrt[a + b*Coth[x]^2]/b`

3.32.
$$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx$$

3.32.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 26, 4153, 26, 354, 90, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \tan\left(\frac{\pi}{2} + ix\right)^3}{\sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)^3}{\sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2}} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \coth^3(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth^3(x)}{(1 - \coth^2(x)) \sqrt{a + b \coth^2(x)}} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\coth^2(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) \\
 & \quad \downarrow \text{90} \\
 & \frac{1}{2} \left(\int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) - \frac{2\sqrt{a + b \coth^2(x)}}{b} \right) \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d\sqrt{b \coth^2(x) + a}}{b} - \frac{2\sqrt{a + b \coth^2(x)}}{b} \right)$$

↓ 221

$$\frac{1}{2} \left(\frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right)}{\sqrt{a + b}} - \frac{2\sqrt{a + b \coth^2(x)}}{b} \right)$$

input `Int[Coth[x]^3/Sqrt[a + b*Coth[x]^2], x]`

output `((2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/Sqrt[a + b] - (2*Sqrt[a + b*Coth[x]^2])/b)/2`

3.32.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

```
rule 354 Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol]
:= Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x]
/; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4153 Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol]
:= With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]
/; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))
```

3.32.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 128 vs. 2(39) = 78.

Time = 0.12 (sec) , antiderivative size = 129, normalized size of antiderivative = 2.74

method	result
derivativedivides	$-\frac{\sqrt{a+b \operatorname{coth}(x)^2}}{b} + \frac{\ln\left(\frac{2a+2b+2b(\operatorname{coth}(x)-1)+2\sqrt{a+b}\sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}}{\operatorname{coth}(x)-1}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b-2b(1+\operatorname{coth}(x))}{\operatorname{coth}(x)-1}\right)}{2\sqrt{a+b}}$
default	$-\frac{\sqrt{a+b \operatorname{coth}(x)^2}}{b} + \frac{\ln\left(\frac{2a+2b+2b(\operatorname{coth}(x)-1)+2\sqrt{a+b}\sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}}{\operatorname{coth}(x)-1}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b-2b(1+\operatorname{coth}(x))}{\operatorname{coth}(x)-1}\right)}{2\sqrt{a+b}}$

```
input int(coth(x)^3/(a+b*coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)
```

```
output -(a+b*coth(x)^2)^(1/2)/b+1/2/(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))+1/2/(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))
```

3.32. $\int \frac{\operatorname{coth}^3(x)}{\sqrt{a+b \operatorname{coth}^2(x)}} dx$

3.32.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. 2(39) = 78.

Time = 0.31 (sec) , antiderivative size = 1576, normalized size of antiderivative = 33.53

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

```
input integrate(coth(x)^3/(a+b*coth(x)^2)^(1/2),x, algorithm="fracas")
```

```
output [1/4*((b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*...
```

3.32.6 Sympy [F]

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx$$

```
input integrate(coth(x)**3/(a+b*coth(x)**2)**(1/2),x)
```

3.32. $\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx$

output `Integral(coth(x)**3/sqrt(a + b*coth(x)**2), x)`

3.32.7 Maxima [F]

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)^3}{\sqrt{b \coth(x)^2 + a}} dx$$

input `integrate(coth(x)^3/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(coth(x)^3/sqrt(b*coth(x)^2 + a), x)`

3.32.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)^3/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:Error: Bad Argument Type`

3.32.9 Mupad [B] (verification not implemented)

Time = 2.39 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{b \coth(x)^2 + a}}{b}$$

input `int(coth(x)^3/(a + b*coth(x)^2)^(1/2),x)`

output `atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(1/2) - (a + b*coth(x)
)^2)^(1/2)/b`

3.33
$$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx$$

3.33.1	Optimal result	256
3.33.2	Mathematica [B] (verified)	256
3.33.3	Rubi [A] (verified)	257
3.33.4	Maple [B] (verified)	259
3.33.5	Fricas [B] (verification not implemented)	260
3.33.6	Sympy [F]	261
3.33.7	Maxima [F]	261
3.33.8	Giac [F(-2)]	261
3.33.9	Mupad [F(-1)]	262

3.33.1 Optimal result

Integrand size = 17, antiderivative size = 60

$$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}}$$

output `-arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))/b^(1/2)+arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))/(a+b)^(1/2)`

3.33.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 134 vs. 2(60) = 120.

Time = 0.24 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.23

$$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\left(-\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{b} \cosh(x)}{\sqrt{-a+b+(a+b) \cosh(2x)}}\right) + \sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{a+b} \cosh(x)}{\sqrt{-a+b+(a+b) \cosh(2x)}}\right)\right) \sqrt{(-a+b+(a+b) \cosh(2x))}}{\sqrt{b}\sqrt{a+b}\sqrt{-a+b+(a+b) \cosh(2x)}}$$

input `Integrate[Coth[x]^2/Sqrt[a + b*Coth[x]^2], x]`

3.33.
$$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx$$

output $((-\text{Sqrt}[a + b] \cdot \text{ArcTanh}[(\text{Sqrt}[2] \cdot \text{Sqrt}[b] \cdot \text{Cosh}[x]) / \text{Sqrt}[-a + b + (a + b) \cdot \text{Cosh}[2x]])] + \text{Sqrt}[b] \cdot \text{ArcTanh}[(\text{Sqrt}[2] \cdot \text{Sqrt}[a + b] \cdot \text{Cosh}[x]) / \text{Sqrt}[-a + b + (a + b) \cdot \text{Cosh}[2x]])] \cdot \text{Sqrt}[-a + b + (a + b) \cdot \text{Cosh}[2x]) \cdot \text{Csch}[x]^2 \cdot \text{Sinh}[x]) / (\text{Sqrt}[b] \cdot \text{Sqrt}[a + b] \cdot \text{Sqrt}[-a + b + (a + b) \cdot \text{Cosh}[2x]])$

3.33.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.529$, Rules used = {3042, 25, 4153, 25, 385, 224, 219, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan\left(\frac{\pi}{2} + ix\right)^2}{\sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\tan\left(ix + \frac{\pi}{2}\right)^2}{\sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\coth^2(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\coth^2(x)}{(1 - \coth^2(x)) \sqrt{a + b \coth^2(x)}} d \coth(x) \\
 & \quad \downarrow \text{385} \\
 & \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \int \frac{1}{\sqrt{b \coth^2(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{224}
 \end{aligned}$$

$$\begin{aligned}
& \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \int \frac{1}{1 - \frac{b \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} \\
& \quad \downarrow 219 \\
& \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{\sqrt{b}} \\
& \quad \downarrow 291 \\
& \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{\sqrt{b}} \\
& \quad \downarrow 219 \\
& \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{\sqrt{a + b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{\sqrt{b}}
\end{aligned}$$

input `Int[Coth[x]^2/Sqrt[a + b*Coth[x]^2], x]`

output `-(ArcTanh[(Sqrt[b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/Sqrt[b]) + ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/Sqrt[a + b]`

3.33.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 291 `Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*((c_) + (d_)*(x_)^2)), x_Symbol] := Subst [Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 385 `Int[(((e_)*(x_))^(m_)*((c_) + (d_)*(x_)^2)^(q_))/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[e^2/b Int[(e*x)^(m - 2)*(c + d*x^2)^q, x], x] - Simp[a*(e^2/b Int[(e*x)^(m - 2)*((c + d*x^2)^q/(a + b*x^2)), x], x] /; FreeQ[{a, b, c, d, e, m, q}, x] && NeQ[b*c - a*d, 0] && LeQ[2, m, 3] && IntBinomialQ[a, b, c, d, e, m, 2, -1, q, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4153 `Int[(((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f f^2*x^2)], x], x, c*(Tan[e + f*x]/ff), x]] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.33.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs. 2(48) = 96.

Time = 0.12 (sec) , antiderivative size = 137, normalized size of antiderivative = 2.28

method	result
derivativedivides	$-\frac{\ln\left(\sqrt{b} \coth(x) + \sqrt{a+b \coth(x)^2}\right)}{\sqrt{b}} - \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b} \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\dots\right)}{2\sqrt{a+b}}$
default	$-\frac{\ln\left(\sqrt{b} \coth(x) + \sqrt{a+b \coth(x)^2}\right)}{\sqrt{b}} - \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b} \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\dots\right)}{2\sqrt{a+b}}$

input `int(coth(x)^2/(a+b*coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)`

3.33. $\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx$

output $-\ln(b^{1/2} \coth(x) + (a+b \coth(x)^2)^{1/2})/b^{1/2} - 1/2/(a+b)^{1/2} \ln((2*a + 2*b - 2*b*(1+\coth(x)) + 2*(a+b)^{1/2}*(b*(1+\coth(x))^2 - 2*b*(1+\coth(x)) + a+b)^{1/2})/(1+\coth(x))) + 1/2/(a+b)^{1/2} \ln((2*a + 2*b + 2*b*(\coth(x)-1) + 2*(a+b)^{1/2}*(b*(\coth(x)-1)^2 + 2*b*(\coth(x)-1) + a+b)^{1/2})/(\coth(x)-1))$

3.33.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 567 vs. $2(48) = 96$.

Time = 0.39 (sec) , antiderivative size = 3513, normalized size of antiderivative = 58.55

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")`

output $[1/4*(\sqrt{a + b}*b*\log(((a*b^2 + b^3)*\cosh(x)^8 + 8*(a*b^2 + b^3)*\cosh(x)*\sinh(x)^7 + (a*b^2 + b^3)*\sinh(x)^8 + 2*(a*b^2 + 2*b^3)*\cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*\cosh(x)^2)*\sinh(x)^6 + 4*(14*(a*b^2 + b^3)*\cosh(x)^3 + 3*(a*b^2 + 2*b^3)*\cosh(x))*\sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^4 + (70*(a*b^2 + b^3)*\cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*\cosh(x)^2)*\sinh(x)^4 + 4*(14*(a*b^2 + b^3)*\cosh(x)^5 + 10*(a*b^2 + 2*b^3)*\cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x))*\sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*\cosh(x)^2 + 2*(14*(a*b^2 + b^3)*\cosh(x)^6 + 15*(a*b^2 + 2*b^3)*\cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^2)*\sinh(x)^2 + \sqrt{2}*(b^2*\cosh(x)^6 + 6*b^2*\cosh(x)*\sinh(x)^5 + b^2*\sinh(x)^6 + 3*b^2*\cosh(x)^4 + 3*(5*b^2*\cosh(x)^2 + b^2)*\sinh(x)^4 + 4*(5*b^2*\cosh(x)^3 + 3*b^2*\cosh(x))*\sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x)^2 + (15*b^2*\cosh(x)^4 + 18*b^2*\cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*\cosh(x)^5 + 6*b^2*\cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x))*\sinh(x))*\sqrt{a + b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)} + 4*(2*(a*b^2 + b^3)*\cosh(x)^7 + 3*(a*b^2 + 2*b^3)*\cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*\cosh(x))*\sinh(x)]/(\cosh(x)^6 + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4*\sinh(x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 15*\cos...$

3.33.6 Sympy [F]

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx$$

input `integrate(coth(x)**2/(a+b*coth(x)**2)**(1/2),x)`

output `Integral(coth(x)**2/sqrt(a + b*coth(x)**2), x)`

3.33.7 Maxima [F]

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(coth(x)^2/sqrt(b*coth(x)^2 + a), x)`

3.33.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

3.33.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

input `int(coth(x)^2/(a + b*coth(x)^2)^(1/2), x)`output `int(coth(x)^2/(a + b*coth(x)^2)^(1/2), x)`

$$3.34 \quad \int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx$$

3.34.1	Optimal result	263
3.34.2	Mathematica [A] (verified)	263
3.34.3	Rubi [A] (verified)	264
3.34.4	Maple [B] (verified)	266
3.34.5	Fricas [B] (verification not implemented)	266
3.34.6	Sympy [F]	267
3.34.7	Maxima [F]	268
3.34.8	Giac [B] (verification not implemented)	268
3.34.9	Mupad [B] (verification not implemented)	269

3.34.1 Optimal result

Integrand size = 15, antiderivative size = 29

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

output `arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(1/2)`

3.34.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

input `Integrate[Coth[x]/Sqrt[a + b*Coth[x]^2],x]`

output `ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/Sqrt[a + b]`

3.34. $\int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx$

3.34.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 26, 4153, 26, 353, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan\left(\frac{\pi}{2} + ix\right)}{\sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)}{\sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -i \int \frac{i \coth(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x)}{(1 - \coth^2(x)) \sqrt{a + b \coth^2(x)}} d \coth(x) \\
 & \quad \downarrow \text{353} \\
 & \frac{1}{2} \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) \\
 & \quad \downarrow \text{73} \\
 & \frac{\int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d \sqrt{b \coth^2(x) + a}}{b} \\
 & \quad \downarrow \text{221} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}}\right)}{\sqrt{a + b}}
 \end{aligned}$$

3.34. $\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx$

input `Int[Coth[x]/Sqrt[a + b*Coth[x]^2], x]`

output `ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/Sqrt[a + b]`

3.34.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 353 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4153 `Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.34.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. $2(23) = 46$.

Time = 0.16 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.93

method	result
derivativedivides	$\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)^2-2b(\coth(x)-1)+a+b}}{\coth(x)-1}\right)}{2\sqrt{a+b}}$
default	$\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)^2-2b(\coth(x)-1)+a+b}}{\coth(x)-1}\right)}{2\sqrt{a+b}}$

input `int(coth(x)/(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2}(a+b)^{-1/2} \ln\left(\frac{(2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b})}{(1+\coth(x))}\right) + \frac{1}{2}(a+b)^{-1/2} \ln\left(\frac{(2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)^2-2b(\coth(x)-1)+a+b})}{(\coth(x)-1)}\right)$$

3.34.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 333 vs. $2(23) = 46$.

Time = 0.30 (sec) , antiderivative size = 1298, normalized size of antiderivative = 44.76

$$\int \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}} dx = \text{Too large to display}$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="fracas")`

output `[1/4*(sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x))^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x)/(cosh(x)^6 + 6*cosh(x))^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh...`

3.34.6 Sympy [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx$$

input `integrate(coth(x)/(a+b*coth(x)**2)**(1/2),x)`

output `Integral(coth(x)/sqrt(a + b*coth(x)**2), x)`

3.34.7 Maxima [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth(x)^2 + a}} dx$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(coth(x)/sqrt(b*coth(x)^2 + a), x)`

3.34.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. $2(23) = 46$.

Time = 0.42 (sec) , antiderivative size = 196, normalized size of antiderivative = 6.76

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \frac{\log\left(\left|-\left(\sqrt{a+be^{2x}}-\sqrt{ae^{4x}+be^{4x}-2ae^{2x}+2be^{2x}+a+b}\right)(a+b)+\sqrt{a+b}(a-b)\right|\right)}{\sqrt{a+b}} + \frac{\log\left(\left|-\sqrt{a+be^{2x}}+\sqrt{ae^{4x}+be^{4x}-2ae^{2x}+2be^{2x}+a+b}\right|\right)}{\sqrt{a+b}}}{2 \operatorname{sgn}(e^{2x} - 1)}$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `-1/2*(log(abs(-(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) + sqrt(a + b)*(a - b)))/sqrt(a + b) + log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b)))/sqrt(a + b) - log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b)))/sqrt(a + b))/sgn(e^(2*x) - 1)`

3.34.9 Mupad [B] (verification not implemented)

Time = 2.31 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

input `int(coth(x)/(a + b*coth(x)^2)^(1/2),x)`output `atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(1/2)`

$$3.35 \quad \int \frac{1}{\sqrt{a+b \coth^2(x)}} dx$$

3.35.1	Optimal result	270
3.35.2	Mathematica [B] (verified)	270
3.35.3	Rubi [A] (verified)	271
3.35.4	Maple [B] (verified)	272
3.35.5	Fricas [B] (verification not implemented)	273
3.35.6	Sympy [F]	274
3.35.7	Maxima [F]	274
3.35.8	Giac [B] (verification not implemented)	274
3.35.9	Mupad [B] (verification not implemented)	275

3.35.1 Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}}$$

output `arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))/(a+b)^(1/2)`

3.35.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 77 vs. $2(31) = 62$.

Time = 0.14 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.48

$$\int \frac{1}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{\frac{(a+b) \coth^2(x)}{a}}}{\sqrt{1+\frac{b \coth^2(x)}{a}}}\right) \coth(x) \sqrt{1+\frac{b \coth^2(x)}{a}}}{\sqrt{\frac{(a+b) \coth^2(x)}{a}} \sqrt{a+b \coth^2(x)}}$$

input `Integrate[1/Sqrt[a + b*Coth[x]^2], x]`

output `(ArcTanh[Sqrt[((a + b)*Coth[x]^2)/a]/Sqrt[1 + (b*Coth[x]^2)/a]]*Coth[x]*Sqrt[1 + (b*Coth[x]^2)/a])/(Sqrt[((a + b)*Coth[x]^2)/a]*Sqrt[a + b*Coth[x]^2])`

3.35. $\int \frac{1}{\sqrt{a+b \coth^2(x)}} dx$

3.35.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4144, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a + b \coth^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{1}{(1 - \coth^2(x)) \sqrt{a + b \coth^2(x)}} d \coth(x) \\
 & \quad \downarrow \text{291} \\
 & \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{a+b \coth^2(x)}} d \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \\
 & \quad \downarrow \text{219} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}}
 \end{aligned}$$

input `Int[1/Sqrt[a + b*Coth[x]^2],x]`

output `ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/Sqrt[a + b]`

3.35.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.35.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. $2(25) = 50$.

Time = 0.14 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.68

method	result
derivativedivides	$-\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)}}{\coth(x)-1}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}}$
default	$-\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)}}{\coth(x)-1}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}}$

input `int(1/(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

```
output -1/2/(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^
2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))+1/2/(a+b)^(1/2)*ln((2*a+2*b+2*b
*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(c
oth(x)-1))
```

3.35.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 399 vs. $2(25) = 50$.

Time = 0.30 (sec) , antiderivative size = 1357, normalized size of antiderivative = 43.77

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

```
input integrate(1/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")
```

```
output [1/4*(sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*s
inh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^
2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*co
sh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 +
6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6
*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(
x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh
(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)
*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4
- a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sin
h(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6
+ 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)
^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*
cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*
b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cos
h(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a
+ b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*co
sh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*co
sh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)
^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(...
```

3.35.6 Sympy [F]

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{1}{\sqrt{a + b \coth^2(x)}} dx$$

input `integrate(1/(a+b*coth(x)**2)**(1/2),x)`

output `Integral(1/sqrt(a + b*coth(x)**2), x)`

3.35.7 Maxima [F]

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{1}{\sqrt{b \coth(x)^2 + a}} dx$$

input `integrate(1/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*coth(x)^2 + a), x)`

3.35.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(25) = 50.

Time = 0.39 (sec) , antiderivative size = 196, normalized size of antiderivative = 6.32

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \frac{\log\left(\frac{-(\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b})(a+b) + \sqrt{a+b}(a-b)}{\sqrt{a+b}}\right) - \frac{\log\left(\frac{-\sqrt{a+be^{2x}} + \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}}{\sqrt{a+b}}\right)}{2 \operatorname{sgn}(e^{2x} - 1)}}{2 \operatorname{sgn}(e^{2x} - 1)}$$

input `integrate(1/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output
$$-1/2*(\log(\text{abs}(-\sqrt{a+b}*e^{2*x} - \sqrt{a*e^{4*x} + b*e^{4*x} - 2*a*e^{2*x} + 2*b*e^{2*x} + a + b}))*\sqrt{a+b} + \sqrt{a+b}*(a-b)))/\sqrt{a+b} - \log(\text{abs}(-\sqrt{a+b}*e^{2*x} + \sqrt{a*e^{4*x} + b*e^{4*x} - 2*a*e^{2*x} + 2*b*e^{2*x} + a + b} + \sqrt{a+b}))/\sqrt{a+b} + \log(\text{abs}(-\sqrt{a+b}*e^{2*x} + \sqrt{a*e^{4*x} + b*e^{4*x} - 2*a*e^{2*x} + 2*b*e^{2*x} + a + b} - \sqrt{a+b}))/\sqrt{a+b})/\text{sgn}(e^{2*x} - 1)$$

3.35.9 Mupad [B] (verification not implemented)

Time = 2.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \frac{\text{atanh}\left(\frac{\coth(x) \sqrt{a+b}}{\sqrt{b \coth(x)^2 + a}}\right)}{\sqrt{a+b}}$$

input `int(1/(a + b*coth(x)^2)^(1/2),x)`

output `atanh((coth(x)*(a + b)^(1/2))/(a + b*coth(x)^2)^(1/2))/(a + b)^(1/2)`

3.36 $\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx$

3.36.1	Optimal result	276
3.36.2	Mathematica [A] (verified)	276
3.36.3	Rubi [A] (verified)	277
3.36.4	Maple [F]	279
3.36.5	Fricas [B] (verification not implemented)	279
3.36.6	Sympy [F]	280
3.36.7	Maxima [F]	281
3.36.8	Giac [F(-2)]	281
3.36.9	Mupad [F(-1)]	281

3.36.1 Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

output `-arctanh((a+b*coth(x)^2)^(1/2)/a^(1/2))/a^(1/2)+arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(1/2)`

3.36.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

input `Integrate[Tanh[x]/Sqrt[a + b*Coth[x]^2],x]`

output `-(ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]]/Sqrt[a]) + ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/Sqrt[a + b]`

3.36. $\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx$

3.36.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3042, 26, 4153, 26, 354, 97, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\tan\left(\frac{\pi}{2} + ix\right) \sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{\tan\left(ix + \frac{\pi}{2}\right) \sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2}} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \tanh(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\tanh(x)}{(1 - \coth^2(x)) \sqrt{a + b \coth^2(x)}} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\tanh(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) \\
 & \quad \downarrow \text{97} \\
 & \frac{1}{2} \left(\int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) + \int \frac{\tanh(x)}{\sqrt{b \coth^2(x) + a}} d \coth^2(x) \right) \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d\sqrt{b \coth^2(x) + a}}{b} + \frac{2 \int \frac{1}{\frac{\coth^4(x)}{b} - \frac{a}{b}} d\sqrt{b \coth^2(x) + a}}{b} \right)$$

↓ 221

$$\frac{1}{2} \left(\frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right)}{\sqrt{a+b}} - \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}} \right)}{\sqrt{a}} \right)$$

input `Int [Tanh[x]/Sqrt[a + b*Coth[x]^2], x]`

output `((-2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]])/Sqrt[a] + (2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/Sqrt[a + b])/2`

3.36.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 97 `Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] := Simp[b/(b*c - a*d) Int[(e + f*x)^p/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[(e + f*x)^p/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && !IntegerQ[p]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

```
rule 354 Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol]
:= Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x]
;/; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4153 Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol]
:= With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]
;/; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))
```

3.36.4 Maple [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth(x)^2}} dx$$

```
input int(tanh(x)/(a+b*coth(x)^2)^(1/2),x)
```

```
output int(tanh(x)/(a+b*coth(x)^2)^(1/2),x)
```

3.36.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 503 vs. $2(44) = 88$.

Time = 0.39 (sec) , antiderivative size = 3397, normalized size of antiderivative = 60.66

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

```
input integrate(tanh(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")
```


output `[1/4*(sqrt(a + b)*a*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x)/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*co...`

3.36.6 Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx$$

input `integrate(tanh(x)/(a+b*coth(x)**2)**(1/2),x)`

output `Integral(tanh(x)/sqrt(a + b*coth(x)**2), x)`

3.36.7 Maxima [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \coth(x)^2 + a}} dx$$

input `integrate(tanh(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(tanh(x)/sqrt(b*coth(x)^2 + a), x)`

3.36.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

input `integrate(tanh(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Error: Bad Argument Type`

3.36.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \coth(x)^2 + a}} dx$$

input `int(tanh(x)/(a + b*coth(x)^2)^(1/2),x)`

output `int(tanh(x)/(a + b*coth(x)^2)^(1/2), x)`

3.37
$$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx$$

3.37.1	Optimal result	282
3.37.2	Mathematica [C] (warning: unable to verify)	282
3.37.3	Rubi [A] (verified)	283
3.37.4	Maple [F]	285
3.37.5	Fricas [B] (verification not implemented)	285
3.37.6	Sympy [F]	286
3.37.7	Maxima [F]	287
3.37.8	Giac [F(-2)]	287
3.37.9	Mupad [F(-1)]	287

3.37.1 Optimal result

Integrand size = 17, antiderivative size = 51

$$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b \coth^2(x)} \tanh(x)}{a}$$

output `arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))/(a+b)^(1/2)-(a+b*coth(x)^2)^(1/2)*tanh(x)/a`

3.37.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.77 (sec) , antiderivative size = 127, normalized size of antiderivative = 2.49

$$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\left(1 + \frac{b \coth^2(x)}{a}\right) \sinh^2(x) \left(\frac{4(a+b) \cosh^2(x)(a+b \coth^2(x)) \operatorname{Hypergeometric2F1}\left(2, 2, \frac{5}{2}, \frac{(a+b) \cosh^2(x)}{a}\right)}{3a^2} + \frac{\arcsin\left(\sqrt{\frac{(a+b) \cosh^2(x)}{a}}\right)}{a\sqrt{-\frac{(a+b) \cosh^2(x)(a+b \coth^2(x))}{a^2}}}\right)}{\sqrt{a+b \coth^2(x)}}$$

3.37.
$$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx$$

input `Integrate[Tanh[x]^2/Sqrt[a + b*Coth[x]^2],x]`

output `((1 + (b*Coth[x]^2)/a)*Sinh[x]^2*((4*(a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Hypergeometric2F1[2, 2, 5/2, ((a + b)*Cosh[x]^2)/a])/(3*a^2) + (ArcSin[Sqrt[((a + b)*Cosh[x]^2)/a]]*(a + 2*b*Coth[x]^2))/(a*Sqrt[-((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2]))*Tanh[x])/Sqrt[a + b*Coth[x]^2]`

3.37.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 25, 4153, 25, 382, 27, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{\tan\left(\frac{\pi}{2} + ix\right)^2 \sqrt{a - b \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{\tan\left(ix + \frac{\pi}{2}\right)^2 \sqrt{a - b \tan\left(ix + \frac{\pi}{2}\right)^2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\tanh^2(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\tanh^2(x)}{(1 - \coth^2(x)) \sqrt{a + b \coth^2(x)}} d \coth(x) \\
 & \quad \downarrow \text{382} \\
 & \frac{\int \frac{a}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x)}{a} - \frac{\tanh(x) \sqrt{a + b \coth^2(x)}}{a}
 \end{aligned}$$

3.37. $\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx$

$$\begin{aligned}
 & \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x) - \frac{\tanh(x) \sqrt{a + b \coth^2(x)}}{a} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{1}{1 - \frac{(a+b) \coth^2(x)}{b \coth^2(x) + a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} - \frac{\tanh(x) \sqrt{a + b \coth^2(x)}}{a} \\
 & \quad \downarrow \text{291} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}} - \frac{\tanh(x) \sqrt{a + b \coth^2(x)}}{a} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

input `Int[Tanh[x]^2/Sqrt[a + b*Coth[x]^2], x]`

output `ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/Sqrt[a + b] - (Sqrt[a + b*Coth[x]^2]*Tanh[x])/a`

3.37.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

```
rule 382 Int[((e._)*(x._))^(m_)*((a_) + (b._)*(x_)^2)^(p_)*((c_) + (d._)*(x_)^2)^(q_)
, x_Symbol] := Simp[(e*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/
(a*c*e*(m + 1))), x] - Simp[1/(a*c*e^2*(m + 1)) Int[(e*x)^(m + 2)*(a + b*
x^2)^p*(c + d*x^2)^q*Simp[(b*c + a*d)*(m + 3) + 2*(b*c*p + a*d*q) + b*d*(m
+ 2*p + 2*q + 5)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && NeQ[
b*c - a*d, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4153 Int[((d._)*tan[(e._) + (f._)*(x_)]^(m_))*((a_) + (b._)*((c._)*tan[(e._) +
(f._)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff), x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.37.4 Maple [F]

$$\int \frac{\tanh(x)^2}{\sqrt{a + b \coth(x)^2}} dx$$

```
input int(tanh(x)^2/(a+b*coth(x)^2)^(1/2),x)
```

```
output int(tanh(x)^2/(a+b*coth(x)^2)^(1/2),x)
```

3.37.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 531 vs. $2(43) = 86$.

Time = 0.33 (sec) , antiderivative size = 1621, normalized size of antiderivative = 31.78

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

```
input integrate(tanh(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")
```

3.37. $\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx$

```
output [1/4*((a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 + a)*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x)*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*c...
```

3.37.6 Sympy [F]

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx$$

```
input integrate(tanh(x)**2/(a+b*coth(x)**2)**(1/2),x)
```

```
output Integral(tanh(x)**2/sqrt(a + b*coth(x)**2), x)
```

3.37.7 Maxima [F]

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

input `integrate(tanh(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(tanh(x)^2/sqrt(b*coth(x)^2 + a), x)`

3.37.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

input `integrate(tanh(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

3.37.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

input `int(tanh(x)^2/(a + b*coth(x)^2)^(1/2),x)`

output `int(tanh(x)^2/(a + b*coth(x)^2)^(1/2), x)`

3.38 $\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.38.1	Optimal result	288
3.38.2	Mathematica [A] (verified)	288
3.38.3	Rubi [A] (verified)	289
3.38.4	Maple [B] (verified)	291
3.38.5	Fricas [B] (verification not implemented)	292
3.38.6	Sympy [F]	293
3.38.7	Maxima [F]	293
3.38.8	Giac [B] (verification not implemented)	293
3.38.9	Mupad [B] (verification not implemented)	294

3.38.1 Optimal result

Integrand size = 17, antiderivative size = 52

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{a}{b(a+b)\sqrt{a+b \coth^2(x)}}$$

output `arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(3/2)+a/b/(a+b)/(a+b*coth(x)^2)^(1/2)`

3.38.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{a}{b(a+b)\sqrt{a+b \coth^2(x)}}$$

input `Integrate[Coth[x]^3/(a + b*Coth[x]^2)^(3/2),x]`

output `ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/(a + b)^(3/2) + a/(b*(a + b)*Sqrt[a + b*Coth[x]^2])`

3.38. $\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.38.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 26, 4153, 26, 354, 87, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \tan\left(\frac{\pi}{2} + ix\right)^3}{\left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)^3}{\left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \coth^3(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth^3(x)}{(1 - \coth^2(x)) (a + b \coth^2(x))^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\coth^2(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth^2(x) \\
 & \quad \downarrow \text{87} \\
 & \frac{1}{2} \left(\frac{\int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x)}{a + b} + \frac{2a}{b(a + b) \sqrt{a + b \coth^2(x)}} \right) \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d\sqrt{b \coth^2(x) + a}}{b(a+b)} + \frac{2a}{b(a+b)\sqrt{a+b \coth^2(x)}} \right)$$

↓ 221

$$\frac{1}{2} \left(\frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right)}{(a+b)^{3/2}} + \frac{2a}{b(a+b)\sqrt{a+b \coth^2(x)}} \right)$$

input `Int[Coth[x]^3/(a + b*Coth[x]^2)^(3/2), x]`

output `((2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/(a + b)^(3/2) + (2*a)/(b*(a + b)*Sqrt[a + b*Coth[x]^2]))/2`

3.38.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

```
rule 354 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol]
:= Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x]
&& NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_.))^p_.), x_Symbol]
:= With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]
&& (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))
```

3.38.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 286 vs. 2(44) = 88.

Time = 0.10 (sec) , antiderivative size = 287, normalized size of antiderivative = 5.52

method	result
derivativedivides	$\frac{1}{b\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}} - \frac{b(2b(1+\coth(x))-2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}$
default	$\frac{1}{b\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}} - \frac{b(2b(1+\coth(x))-2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}$

```
input int(coth(x)^3/(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)
```

```
output 1/b/(a+b*coth(x)^2)^(1/2)-1/2/(a+b)/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(
(1/2)-b/(a+b)*(2*b*(1+coth(x))-2*b)/(4*(a+b)*b-4*b^2)/(b*(1+coth(x))^2-2*b
*(1+coth(x))+a+b)^(1/2)+1/2/(a+b)^(3/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b
)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))-1/2/(a+b
)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+b/(a+b)*(2*b*(coth(x)-1)+2*b
)/(4*(a+b)*b-4*b^2)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1/2/(a+b)^(
(3/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth
(x)-1)+a+b)^(1/2))/(coth(x)-1))
```

3.38.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 990 vs. $2(44) = 88$.

Time = 0.36 (sec) , antiderivative size = 2541, normalized size of antiderivative = 48.87

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(coth(x)^3/(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")
```

```
output [1/4*(((a*b + b^2)*cosh(x)^4 + 4*(a*b + b^2)*cosh(x)*sinh(x)^3 + (a*b + b^
2)*sinh(x)^4 - 2*(a*b - b^2)*cosh(x)^2 + 2*(3*(a*b + b^2)*cosh(x)^2 - a*b
+ b^2)*sinh(x)^2 + a*b + b^2 + 4*((a*b + b^2)*cosh(x)^3 - (a*b - b^2)*cosh
(x))*sinh(x))*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*
cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6
- 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 +
a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b
- a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b
- a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2
*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 +
b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2
*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*c
osh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(
x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*s
inh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^
2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 +
(15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a
^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b -
b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh
(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3...
```

3.38. $\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.38.6 Sympy [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth^3(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

input `integrate(coth(x)**3/(a+b*coth(x)**2)**(3/2),x)`

output `Integral(coth(x)**3/(a + b*coth(x)**2)**(3/2), x)`

3.38.7 Maxima [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)^3}{(b \coth(x)^2 + a)^{\frac{3}{2}}} dx$$

input `integrate(coth(x)^3/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate(coth(x)^3/(b*coth(x)^2 + a)^(3/2), x)`

3.38.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 359 vs. $2(44) = 88$.

Time = 0.46 (sec) , antiderivative size = 359, normalized size of antiderivative = 6.90

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{(a^3 + a^2b)e^{(2x)}}{a^3 b \operatorname{sgn}(e^{(2x)} - 1) + 2a^2 b^2 \operatorname{sgn}(e^{(2x)} - 1) + ab^3 \operatorname{sgn}(e^{(2x)} - 1)} - \frac{a^3 + a^2b}{a^3 b \operatorname{sgn}(e^{(2x)} - 1) + 2a^2 b^2 \operatorname{sgn}(e^{(2x)} - 1) + ab^3 \operatorname{sgn}(e^{(2x)} - 1)}$$

$$\frac{\log \left(\left| \left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} \right) \sqrt{a + b} - a + b \right| \right)}{2(a + b)^{\frac{3}{2}} \operatorname{sgn}(e^{(2x)} - 1)}$$

$$- \frac{\log \left(\left| \left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} \right) \sqrt{a + b} - a - b \right| \right)}{2(a + b)^{\frac{3}{2}} \operatorname{sgn}(e^{(2x)} - 1)}$$

$$+ \frac{\log \left(\left| -\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b} \right| \right)}{2(a + b)^{\frac{3}{2}} \operatorname{sgn}(e^{(2x)} - 1)}$$

3.38. $\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx$

input `integrate(coth(x)^3/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output
$$\begin{aligned} & ((a^3 + a^2b)*e^{(2*x)}/(a^3*b*\operatorname{sgn}(e^{(2*x)} - 1) + 2*a^2*b^2*\operatorname{sgn}(e^{(2*x)} - 1) \\ & + a*b^3*\operatorname{sgn}(e^{(2*x)} - 1)) - (a^3 + a^2b)/(a^3*b*\operatorname{sgn}(e^{(2*x)} - 1) + 2*a^2*b^2*\operatorname{sgn}(e^{(2*x)} - 1) \\ & + a*b^3*\operatorname{sgn}(e^{(2*x)} - 1)))/\operatorname{sqrt}(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b) \\ & - 1/2*\log(\operatorname{abs}(\operatorname{sqrt}(a + b)*e^{(2*x)} - \operatorname{sqrt}(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b))*\operatorname{sqrt}(a + b) \\ & - a + b))/((a + b)^{(3/2)}*\operatorname{sgn}(e^{(2*x)} - 1)) - 1/2*\log(\operatorname{abs}(\operatorname{sqrt}(a + b)*e^{(2*x)} - \operatorname{sqrt}(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b) \\ &)*\operatorname{sqrt}(a + b) - a - b))/((a + b)^{(3/2)}*\operatorname{sgn}(e^{(2*x)} - 1)) + 1/2*\log(\operatorname{abs}(-\operatorname{sqrt}(a + b)*e^{(2*x)} + \operatorname{sqrt}(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b) \\ & - \operatorname{sqrt}(a + b)))/((a + b)^{(3/2)}*\operatorname{sgn}(e^{(2*x)} - 1)) \end{aligned}$$

3.38.9 Mupad [B] (verification not implemented)

Time = 2.77 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{coth}^3(x)}{(a + b \operatorname{coth}^2(x))^{3/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \operatorname{coth}(x)^2 + a}}{\sqrt{a+b}}\right)}{(a + b)^{3/2}} + \frac{a}{(b^2 + a b) \sqrt{b \operatorname{coth}(x)^2 + a}}$$

input `int(coth(x)^3/(a + b*coth(x)^2)^(3/2),x)`

output `atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(3/2) + a/((a*b + b^2)*(a + b*coth(x)^2)^(1/2))`

3.39 $\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.39.1	Optimal result	295
3.39.2	Mathematica [B] (verified)	295
3.39.3	Rubi [A] (verified)	296
3.39.4	Maple [B] (verified)	298
3.39.5	Fricas [B] (verification not implemented)	299
3.39.6	Sympy [F]	299
3.39.7	Maxima [F]	300
3.39.8	Giac [B] (verification not implemented)	300
3.39.9	Mupad [F(-1)]	301

3.39.1 Optimal result

Integrand size = 17, antiderivative size = 53

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a + b)^{3/2}} - \frac{\coth(x)}{(a + b)\sqrt{a + b \coth^2(x)}}$$

output `arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))/(a+b)^(3/2)-coth(x)/(a+b)/(a+b*coth(x)^2)^(1/2)`

3.39.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 109 vs. 2(53) = 106.

Time = 1.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.06

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{-2(a + b) \coth(x) + \frac{\operatorname{arctanh}\left(\frac{\sqrt{\frac{(a+b) \coth^2(x)}{a}}}{\sqrt{1 + \frac{b \coth^2(x)}{a}}}\right) (-a + b + (a+b) \cosh(2x)) \sqrt{\frac{(a+b) \coth^2(x)}{a}} \operatorname{csch}(2x)}{\sqrt{1 + \frac{b \coth^2(x)}{a}}}}{2(a + b)^2 \sqrt{a + b \coth^2(x)}}$$

input `Integrate[Coth[x]^2/(a + b*Coth[x]^2)^(3/2),x]`

3.39. $\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx$

output $(-2*(a + b)*\text{Coth}[x] + (\text{ArcTanh}[\text{Sqrt}[(a + b)*\text{Coth}[x]^2/a]/\text{Sqrt}[1 + (b*\text{Coth}[x]^2/a)])*(-a + b + (a + b)*\text{Cosh}[2*x])*\text{Sqrt}[(a + b)*\text{Coth}[x]^2/a]*\text{Csch}[x]*\text{Sech}[x])/\text{Sqrt}[1 + (b*\text{Coth}[x]^2/a)]/(2*(a + b)^2*\text{Sqrt}[a + b*\text{Coth}[x]^2])$

3.39.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3042, 25, 4153, 25, 373, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan(\frac{\pi}{2} + ix)^2}{(a - b \tan(\frac{\pi}{2} + ix)^2)^{3/2}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\tan(ix + \frac{\pi}{2})^2}{(a - b \tan(ix + \frac{\pi}{2})^2)^{3/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\coth^2(x)}{(1 - \coth^2(x))(b \coth^2(x) + a)^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\coth^2(x)}{(1 - \coth^2(x))(a + b \coth^2(x))^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{373} \\
 & \frac{\int \frac{1}{(1 - \coth^2(x))\sqrt{b \coth^2(x) + a}} d \coth(x)}{a + b} - \frac{\coth(x)}{(a + b)\sqrt{a + b \coth^2(x)}} \\
 & \quad \downarrow \text{291}
 \end{aligned}$$

$$\frac{\int \frac{1}{1 - \frac{(a+b)\coth^2(x)}{b\coth^2(x)+a}} d \frac{\coth(x)}{\sqrt{b\coth^2(x)+a}}}{a+b} - \frac{\coth(x)}{(a+b)\sqrt{a+b\coth^2(x)}}$$

↓ 219

$$\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{(a+b)^{3/2}} - \frac{\coth(x)}{(a+b)\sqrt{a+b\coth^2(x)}}$$

input `Int[Coth[x]^2/(a + b*Coth[x]^2)^(3/2), x]`

output `ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/(a + b)^(3/2) - Coth[x]/((a + b)*Sqrt[a + b*Coth[x]^2])`

3.39.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 373 `Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[e*(e*x)^(m - 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*(b*c - a*d)*(p + 1))), x] - Simp[e^2/(2*(b*c - a*d)*(p + 1)) Int[(e*x)^(m - 2)*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(m - 1) + d*(m + 2*p + 2*q + 3)*x^2, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[m, 1] && LeQ[m, 3] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4153 `Int[((d_)*tan[(e_) + (f_)*(x_)]^(m_))*((a_) + (b_))*((c_)*tan[(e_) + (f_)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.39.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 288 vs. $2(45) = 90$.

Time = 0.10 (sec) , antiderivative size = 289, normalized size of antiderivative = 5.45

method	result
derivativedivides	$-\frac{\coth(x)}{a\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}$
default	$-\frac{\coth(x)}{a\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}$

input `int(coth(x)^2/(a+b*coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -\coth(x)/a/(a+b\coth(x)^2)^{(1/2)} - 1/2/(a+b)/(b(\coth(x)-1)^2+2b(\coth(x)-1) \\ & +a+b)^{(1/2)} + b/(a+b)*(2b(\coth(x)-1)+2b)/(4(a+b)b-4b^2)/(b(\coth(x)-1) \\ & ^2+2b(\coth(x)-1)+a+b)^{(1/2)} + 1/2/(a+b)^{(3/2)}*\ln((2*a+2*b+2*b*(\coth(x)-1) \\ & +2*(a+b)^{(1/2)}*(b(\coth(x)-1)^2+2*b(\coth(x)-1)+a+b)^{(1/2)})/(\coth(x)-1)) + 1 \\ & /2/(a+b)/(b(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)} + b/(a+b)*(2*b*(1+\coth(\\ & x))-2*b)/(4(a+b)b-4*b^2)/(b(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)} - 1/2 \\ & / (a+b)^{(3/2)}*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^{(1/2)}*(b(1+\coth(x))^2-2* \\ & b*(1+\coth(x))+a+b)^{(1/2)})/(1+\coth(x))) \end{aligned}$$

3.39.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 860 vs. $2(45) = 90$.

Time = 0.35 (sec) , antiderivative size = 2279, normalized size of antiderivative = 43.00

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="fracas")`

output `[1/4*(((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x))^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^...`

3.39.6 SymPy [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth^2(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

input `integrate(coth(x)**2/(a+b*coth(x)**2)**(3/2),x)`

3.39. $\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx$

output `Integral(coth(x)**2/(a + b*coth(x)**2)**(3/2), x)`

3.39.7 Maxima [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{\frac{3}{2}}} dx$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate(coth(x)^2/(b*coth(x)^2 + a)^(3/2), x)`

3.39.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 363 vs. $2(45) = 90$.

Time = 0.46 (sec) , antiderivative size = 363, normalized size of antiderivative = 6.85

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx =$$

$$\frac{\frac{(a^2b+ab^2)e^{(2x)}}{a^3b\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)} + \frac{a^2b+ab^2}{a^3b\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)}}{\sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}}$$

$$\frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a + b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$+ \frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a - b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$\frac{\log\left(\left|-\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b}\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output $-\frac{(a^2b + ab^2)e^{2x}}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1} + \frac{a^2b + ab^2}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1} + \frac{a^2b^3 \operatorname{sgn}(e^{2x}) - 1}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1} + \frac{a^2b^3 \operatorname{sgn}(e^{2x}) - 1}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1}}{\sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}} - \frac{1}{2} \log(\operatorname{abs}(\sqrt{a+b}e^{2x} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}) \sqrt{a+b} - a + b)) / ((a+b)^{3/2} \operatorname{sgn}(e^{2x}) - 1)) + \frac{1}{2} \log(\operatorname{abs}(\sqrt{a+b}e^{2x} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}) \sqrt{a+b} - a - b)) / ((a+b)^{3/2} \operatorname{sgn}(e^{2x}) - 1)) - \frac{1}{2} \log(\operatorname{abs}(-\sqrt{a+b}e^{2x} + \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b} - \sqrt{a+b})) / ((a+b)^{3/2} \operatorname{sgn}(e^{2x}) - 1))$

3.39.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{3/2}} dx$$

input `int(coth(x)^2/(a + b*coth(x)^2)^(3/2), x)`

output `int(coth(x)^2/(a + b*coth(x)^2)^(3/2), x)`

3.40 $\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.40.1	Optimal result	302
3.40.2	Mathematica [C] (verified)	302
3.40.3	Rubi [A] (verified)	303
3.40.4	Maple [B] (verified)	305
3.40.5	Fricas [B] (verification not implemented)	306
3.40.6	Sympy [F]	306
3.40.7	Maxima [F]	307
3.40.8	Giac [B] (verification not implemented)	307
3.40.9	Mupad [B] (verification not implemented)	308

3.40.1 Optimal result

Integrand size = 15, antiderivative size = 49

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{1}{(a+b)\sqrt{a+b \coth^2(x)}}$$

output `arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(3/2)-1/(a+b)/(a+b*coth(x)^2)^(1/2)`

3.40.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.84

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = -\frac{\operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \coth^2(x)}{a+b}\right)}{(a+b)\sqrt{a+b \coth^2(x)}}$$

input `Integrate[Coth[x]/(a + b*Coth[x]^2)^(3/2),x]`

output `-(Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Coth[x]^2)/(a + b)]/((a + b)*Sqrt[a + b*Coth[x]^2]))`

3.40. $\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.40.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.10, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3042, 26, 4153, 26, 353, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan\left(\frac{\pi}{2} + ix\right)}{\left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)}{\left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -i \int \frac{i \coth(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x)}{(1 - \coth^2(x)) (a + b \coth^2(x))^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{353} \\
 & \frac{1}{2} \int \frac{1}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth^2(x) \\
 & \quad \downarrow \text{61} \\
 & \frac{1}{2} \left(\frac{\int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x)}{a + b} - \frac{2}{(a + b) \sqrt{a + b \coth^2(x)}} \right) \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d\sqrt{b \coth^2(x) + a}}{b(a+b)} - \frac{2}{(a+b)\sqrt{a+b \coth^2(x)}} \right)$$

↓ 221

$$\frac{1}{2} \left(\frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right)}{(a+b)^{3/2}} - \frac{2}{(a+b)\sqrt{a+b \coth^2(x)}} \right)$$

input `Int[Coth[x]/(a + b*Coth[x]^2)^(3/2), x]`

output `((2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/(a + b)^(3/2) - 2/((a + b)*Sqrt[a + b*Coth[x]^2]))/2`

3.40.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 353 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol]
 := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[
 {a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`

rule 4153 `Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
 (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
 x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
 f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m,
 n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
 nalQ[n]))`

3.40.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. 2(41) = 82.

Time = 0.08 (sec) , antiderivative size = 273, normalized size of antiderivative = 5.57

method	result
derivativedivides	$-\frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} +$
default	$-\frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} +$

input `int(coth(x)/(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)`

output `-1/2/(a+b)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+b/(a+b)*(2*b*(coth(
 x)-1)+2*b)/(4*(a+b)*b-4*b^2)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1
 /2/(a+b)^(3/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+
 2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))-1/2/(a+b)/(b*(1+coth(x))^2-2*b*(1
 +coth(x))+a+b)^(1/2)-b/(a+b)*(2*b*(1+coth(x))-2*b)/(4*(a+b)*b-4*b^2)/(b*(1
 +coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)+1/2/(a+b)^(3/2)*ln((2*a+2*b-2*b*(1+
 coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+cot
 h(x)))`

3.40.
$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx$$

3.40.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 869 vs. $2(41) = 82$.

Time = 0.35 (sec) , antiderivative size = 2299, normalized size of antiderivative = 46.92

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(coth(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="fracas")
```

```
output [1/4*(((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4
- 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a
+ b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)*sqrt(a + b)*log(-((a^3
+ a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*si
nh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b
)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)
*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3
+ a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*c
osh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*c
osh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^
2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a
^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3
*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(
x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^
2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)
^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^
2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^
5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*
sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)
)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2...
```

3.40.6 Sympy [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

```
input integrate(coth(x)/(a+b*coth(x)**2)**(3/2),x)
```

```
output Integral(coth(x)/(a + b*coth(x)**2)**(3/2), x)
```

3.40. $\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.40.7 Maxima [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth(x)^2 + a)^{3/2}} dx$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate(coth(x)/(b*coth(x)^2 + a)^(3/2), x)`

3.40.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 364 vs. 2(41) = 82.

Time = 0.47 (sec) , antiderivative size = 364, normalized size of antiderivative = 7.43

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx =$$

$$\frac{\frac{(a^2b+ab^2)e^{(2x)}}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)} - \frac{a^2b+ab^2}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)}}{\sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}}$$

$$\frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a + b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$\frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a - b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$+ \frac{\log\left(\left|-\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b}\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output $-\left(\frac{(a^2b + ab^2)e^{2x}}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1} + \frac{ab^3 \operatorname{sgn}(e^{2x})}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1}\right) - \left(\frac{(a^2b + ab^2)}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1} + \frac{ab^3 \operatorname{sgn}(e^{2x})}{(a^3b \operatorname{sgn}(e^{2x}) - 1) + 2a^2b^2 \operatorname{sgn}(e^{2x}) - 1}\right) / \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b} - \frac{1}{2} \log\left(\frac{\left(\sqrt{a+b}e^{2x} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right) \sqrt{a+b} - a + b}{(a+b)^{3/2} \operatorname{sgn}(e^{2x}) - 1}\right) - \frac{1}{2} \log\left(\frac{\left(\sqrt{a+b}e^{2x} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right) \sqrt{a+b} - a - b}{(a+b)^{3/2} \operatorname{sgn}(e^{2x}) - 1}\right) + \frac{1}{2} \log\left(\frac{a - \sqrt{a+b}e^{2x} + \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}}{(a+b)^{3/2} \operatorname{sgn}(e^{2x}) - 1}\right)$

3.40.9 Mupad [B] (verification not implemented)

Time = 2.61 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.84

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{1}{(a+b) \sqrt{b \coth(x)^2 + a}}$$

input `int(coth(x)/(a + b*coth(x)^2)^(3/2),x)`

output `atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(3/2) - 1/((a + b)*(a + b*coth(x)^2)^(1/2))`

3.41
$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx$$

3.41.1	Optimal result	309
3.41.2	Mathematica [C] (verified)	309
3.41.3	Rubi [A] (verified)	310
3.41.4	Maple [F]	313
3.41.5	Fricas [B] (verification not implemented)	313
3.41.6	Sympy [F]	314
3.41.7	Maxima [F]	314
3.41.8	Giac [F(-2)]	314
3.41.9	Mupad [F(-1)]	315

3.41.1 Optimal result

Integrand size = 15, antiderivative size = 78

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{a^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{b}{a(a+b)\sqrt{a+b \coth^2(x)}}$$

output

```
-arctanh((a+b*coth(x)^2)^(1/2)/a^(1/2))/a^(3/2)+arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(3/2)+b/a/(a+b)/(a+b*coth(x)^2)^(1/2)
```

3.41.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.90

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{-a \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \coth^2(x)}{a+b}\right) + (a+b) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \coth^2(x)}{a+b}\right)}{a(a+b)\sqrt{a+b \coth^2(x)}}$$

3.41.
$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx$$

input `Integrate[Tanh[x]/(a + b*Coth[x]^2)^(3/2),x]`

output `(-a*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Coth[x]^2)/(a + b)]) + (a + b)*Hypergeometric2F1[-1/2, 1, 1/2, 1 + (b*Coth[x]^2)/a]/(a*(a + b)*Sqrt[a + b*Coth[x]^2])`

3.41.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.26, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 26, 4153, 26, 354, 96, 25, 174, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\tan\left(\frac{\pi}{2} + ix\right) \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{\tan\left(ix + \frac{\pi}{2}\right) \left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \tanh(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\tanh(x)}{(1 - \coth^2(x)) (a + b \coth^2(x))^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\tanh(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth^2(x) \\
 & \quad \downarrow \text{96}
 \end{aligned}$$

3.41. $\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx$

$$\begin{aligned}
& \frac{1}{2} \left(\frac{2b}{a(a+b)\sqrt{a+b\coth^2(x)}} - \frac{\int -\frac{(-b\coth^2(x)+a+b)\tanh(x)}{(1-\coth^2(x))\sqrt{b\coth^2(x)+a}} d\coth^2(x)}{a(a+b)} \right) \\
& \quad \downarrow 25 \\
& \frac{1}{2} \left(\frac{\int \frac{(-b\coth^2(x)+a+b)\tanh(x)}{(1-\coth^2(x))\sqrt{b\coth^2(x)+a}} d\coth^2(x)}{a(a+b)} + \frac{2b}{a(a+b)\sqrt{a+b\coth^2(x)}} \right) \\
& \quad \downarrow 174 \\
& \frac{1}{2} \left(\frac{a \int \frac{1}{(1-\coth^2(x))\sqrt{b\coth^2(x)+a}} d\coth^2(x) + (a+b) \int \frac{\tanh(x)}{\sqrt{b\coth^2(x)+a}} d\coth^2(x)}{a(a+b)} + \frac{2b}{a(a+b)\sqrt{a+b\coth^2(x)}} \right) \\
& \quad \downarrow 73 \\
& \frac{1}{2} \left(\frac{2a \int \frac{\frac{a+b}{b} - \frac{\coth^4(x)}{b}}{b} d\sqrt{b\coth^2(x)+a} + \frac{2(a+b) \int \frac{\frac{1}{\coth^4(x)} - \frac{a}{b}}{b} d\sqrt{b\coth^2(x)+a}}{a(a+b)} + \frac{2b}{a(a+b)\sqrt{a+b\coth^2(x)}} \right) \\
& \quad \downarrow 221 \\
& \frac{1}{2} \left(\frac{2a \operatorname{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}} \right)}{\sqrt{a+b}} - \frac{2(a+b) \operatorname{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a}} \right)}{\sqrt{a}} + \frac{2b}{a(a+b)\sqrt{a+b\coth^2(x)}} \right)
\end{aligned}$$

input `Int[Tanh[x]/(a + b*Coth[x]^2)^(3/2), x]`

output `(((-2*(a + b)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]])/Sqrt[a] + (2*a*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/Sqrt[a + b])/(a*(a + b)) + (2*b)/(a*(a + b)*Sqrt[a + b*Coth[x]^2]))/2`

3.41.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 96 `Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] := Simp[f*(e + f*x)^(p + 1)/((p + 1)*(b*e - a*f)*(d*e - c*f)), x] + Simp[1/((b*e - a*f)*(d*e - c*f)) Int[(b*d*e - b*c*f - a*d*f - b*d*f*x)*((e + f*x)^(p + 1)/((a + b*x)*(c + d*x))), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[p, -1]`
- rule 174 `Int[(((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] := Simp[(b*g - a*h)/(b*c - a*d) Int[(e + f*x)^p/(a + b*x), x], x] - Simp[(d*g - c*h)/(b*c - a*d) Int[(e + f*x)^p/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4153 Int[((d_)*tan[(e_) + (f_)*(x_)]^(m_))*((a_) + (b_)*((c_)*tan[(e_) +
(f_)*(x_)]^(n_))^(p_)), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.41.4 Maple [F]

$$\int \frac{\tanh(x)}{(a + b \coth(x)^2)^{\frac{3}{2}}} dx$$

```
input int(tanh(x)/(a+b*coth(x)^2)^(3/2),x)
```

```
output int(tanh(x)/(a+b*coth(x)^2)^(3/2),x)
```

3.41.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1437 vs. $2(64) = 128$.

Time = 0.55 (sec) , antiderivative size = 6991, normalized size of antiderivative = 89.63

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(tanh(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")
```

```
output Too large to include
```

3.41.6 Sympy [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

input `integrate(tanh(x)/(a+b*coth(x)**2)**(3/2),x)`

output `Integral(tanh(x)/(a + b*coth(x)**2)**(3/2), x)`

3.41.7 Maxima [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{\frac{3}{2}}} dx$$

input `integrate(tanh(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate(tanh(x)/(b*coth(x)^2 + a)^(3/2), x)`

3.41.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(tanh(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type`

3.41.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{3/2}} dx$$

input `int(tanh(x)/(a + b*coth(x)^2)^(3/2), x)`output `int(tanh(x)/(a + b*coth(x)^2)^(3/2), x)`

3.42 $\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx$

3.42.1 Optimal result 316
 3.42.2 Mathematica [C] (warning: unable to verify) 316
 3.42.3 Rubi [A] (verified) 317
 3.42.4 Maple [F] 320
 3.42.5 Fricas [B] (verification not implemented) 320
 3.42.6 Sympy [F] 321
 3.42.7 Maxima [F] 322
 3.42.8 Giac [B] (verification not implemented) 322
 3.42.9 Mupad [F(-1)] 323

3.42.1 Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{3/2}} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{(a+2b)\sqrt{a+b \coth^2(x)} \tanh(x)}{a^2(a+b)}$$

```
output arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))/(a+b)^(3/2)+b*tanh(x)/a
/(a+b)/(a+b*coth(x)^2)^(1/2)-(a+2*b)*(a+b*coth(x)^2)^(1/2)*tanh(x)/a^2/(a+
b)
```

3.42.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.93 (sec) , antiderivative size = 260, normalized size of antiderivative = 3.06

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\sinh^2(x) \tanh(x) \left(\frac{8(a+b) \cosh^2(x) (2a^2+5ab \coth^2(x)+3b^2 \coth^4(x)) \operatorname{Hypergeometric2F1}\left(2, 2, \frac{7}{2}, \frac{(a+b) \cosh^2(x)}{a+b \coth^2(x)}\right)}{15a^3} \right)}{(a+b \coth^2(x))^{3/2}}$$

input `Integrate[Tanh[x]^2/(a + b*Coth[x]^2)^(3/2),x]`

output `(Sinh[x]^2*Tanh[x]*((8*(a + b)*Cosh[x]^2*(2*a^2 + 5*a*b*Coth[x]^2 + 3*b^2*Coth[x]^4)*Hypergeometric2F1[2, 2, 7/2, ((a + b)*Cosh[x]^2)/a])/(15*a^3) - (8*(a + b)*((-I)*a*Coth[x] - I*b*Coth[x]^3)^2*HypergeometricPFQ[{2, 2, 2}, {1, 7/2}, ((a + b)*Cosh[x]^2)/a]*Sinh[x]^2)/(15*a^3) - ((3*a^2 + 12*a*b*Coth[x]^2 + 8*b^2*Coth[x]^4)*(ArcSin[Sqrt[((a + b)*Cosh[x]^2)/a]]*(-a - b*Coth[x]^2) - a*Csch[x]^2*Sqrt[-(((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2]))*Tanh[x]^2)/(a^2*(a + b)*Sqrt[-(((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2)])))/(a*Sqrt[a + b*Coth[x]^2])`

3.42.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.07, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.647$, Rules used = {3042, 25, 4153, 25, 374, 25, 445, 25, 27, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{\tan\left(\frac{\pi}{2} + ix\right)^2 \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{\tan\left(ix + \frac{\pi}{2}\right)^2 \left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{3/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\tanh^2(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\tanh^2(x)}{(1 - \coth^2(x)) (a + b \coth^2(x))^{3/2}} d \coth(x)
 \end{aligned}$$

$$\begin{aligned}
& \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{\int -\frac{(-2b \coth^2(x)+a+2b) \tanh^2(x)}{(1-\coth^2(x))\sqrt{b \coth^2(x)+a}} d \coth(x)}{a(a+b)} \\
& \quad \downarrow 374 \\
& \frac{\int \frac{(-2b \coth^2(x)+a+2b) \tanh^2(x)}{(1-\coth^2(x))\sqrt{b \coth^2(x)+a}} d \coth(x)}{a(a+b)} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} \\
& \quad \downarrow 25 \\
& \frac{\int -\frac{a^2}{(1-\coth^2(x))\sqrt{b \coth^2(x)+a}} d \coth(x)}{a(a+b)} - \frac{(a+2b) \tanh(x)\sqrt{a+b \coth^2(x)}}{a} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} \\
& \quad \downarrow 445 \\
& \frac{\int \frac{a^2}{(1-\coth^2(x))\sqrt{b \coth^2(x)+a}} d \coth(x)}{a(a+b)} - \frac{(a+2b) \tanh(x)\sqrt{a+b \coth^2(x)}}{a} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} \\
& \quad \downarrow 25 \\
& \frac{a \int \frac{1}{(1-\coth^2(x))\sqrt{b \coth^2(x)+a}} d \coth(x)}{a(a+b)} - \frac{(a+2b) \tanh(x)\sqrt{a+b \coth^2(x)}}{a} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} \\
& \quad \downarrow 27 \\
& \frac{a \int \frac{1}{1-\frac{(a+b) \coth^2(x)}{b \coth^2(x)+a}} d \frac{\coth(x)}{\sqrt{b \coth^2(x)+a}} - \frac{(a+2b) \tanh(x)\sqrt{a+b \coth^2(x)}}{a}}{a(a+b)} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} \\
& \quad \downarrow 291 \\
& \frac{a \operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right) - \frac{(a+2b) \tanh(x)\sqrt{a+b \coth^2(x)}}{a}}{\sqrt{a+b}} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} \\
& \quad \downarrow 219
\end{aligned}$$

input `Int [Tanh [x]^2/(a + b*Coth [x]^2)^(3/2), x]`

$$3.42. \quad \int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx$$

output $(b \tanh[x]) / (a(a + b) \sqrt{a + b \coth[x]^2}) + ((a \operatorname{ArcTanh}[\sqrt{a + b} \coth[x]) / \sqrt{a + b \coth[x]^2}] / \sqrt{a + b} - ((a + 2b) \sqrt{a + b \coth[x]^2} \tanh[x]) / a) / (a(a + b))$

3.42.3.1 Defintions of rubi rules used

rule 25 $\operatorname{Int}[-(F_x), x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$

rule 27 $\operatorname{Int}[(a_*)(F_x), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \&\& \operatorname{!MatchQ}[F_x, (b_*)(G_x)] /; \operatorname{FreeQ}[b, x]$

rule 219 $\operatorname{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) * \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] * (x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \operatorname{||} \operatorname{LtQ}[b, 0])$

rule 291 $\operatorname{Int}[1/(\sqrt{(a_*) + (b_*)(x_)^2} * ((c_*) + (d_*)(x_)^2)), x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(c - (b*c - a*d)*x^2), x], x, x/\sqrt{a + b*x^2}] /; \operatorname{FreeQ}\{a, b, c, d\}, x \&\& \operatorname{NeQ}[b*c - a*d, 0]$

rule 374 $\operatorname{Int}[(e_*)(x_)^{(m_*)} * ((a_*) + (b_*)(x_)^2)^{(p_*)} * ((c_*) + (d_*)(x_)^2)^{(q_*)}, x_Symbol] \rightarrow \operatorname{Simp}[(-b) * (e*x)^{(m+1)} * (a + b*x^2)^{(p+1)} * (c + d*x^2)^{(q+1)} / (a*e*2*(b*c - a*d)*(p+1)), x] + \operatorname{Simp}[1/(a*2*(b*c - a*d)*(p+1)) \operatorname{Int}[(e*x)^m * (a + b*x^2)^{(p+1)} * (c + d*x^2)^q * \operatorname{Simp}[b*c*(m+1) + 2*(b*c - a*d)*(p+1) + d*b*(m+2*(p+q+2)+1)*x^2, x], x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, m, q\}, x \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[p, -1] \&\& \operatorname{IntBinomialQ}[a, b, c, d, e, m, 2, p, q, x]$

rule 445 $\operatorname{Int}[(g_*)(x_)^{(m_*)} * ((a_*) + (b_*)(x_)^2)^{(p_*)} * ((c_*) + (d_*)(x_)^2)^{(q_*)} * ((e_*) + (f_*)(x_)^2), x_Symbol] \rightarrow \operatorname{Simp}[e * (g*x)^{(m+1)} * (a + b*x^2)^{(p+1)} * (c + d*x^2)^{(q+1)} / (a*c*g*(m+1)), x] + \operatorname{Simp}[1/(a*c*g^2*(m+1)) \operatorname{Int}[(g*x)^{(m+2)} * (a + b*x^2)^p * (c + d*x^2)^q * \operatorname{Simp}[a*f*c*(m+1) - e*(b*c + a*d)*(m+2+1) - e*2*(b*c*p + a*d*q) - b*e*d*(m+2*(p+q+2)+1)*x^2, x], x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, g, p, q\}, x \&\& \operatorname{LtQ}[m, -1]$


```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)]^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.42.4 Maple [F]

$$\int \frac{\tanh(x)^2}{(a + b \coth(x)^2)^{\frac{3}{2}}} dx$$

```
input int(tanh(x)^2/(a+b*coth(x)^2)^(3/2),x)
```

```
output int(tanh(x)^2/(a+b*coth(x)^2)^(3/2),x)
```

3.42.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1686 vs. 2(75) = 150.

Time = 0.51 (sec) , antiderivative size = 3931, normalized size of antiderivative = 46.25

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(tanh(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")
```

output

```
[1/4*((a^3 + a^2*b)*cosh(x)^6 + 6*(a^3 + a^2*b)*cosh(x)*sinh(x)^5 + (a^3
+ a^2*b)*sinh(x)^6 - (a^3 - 3*a^2*b)*cosh(x)^4 - (a^3 - 3*a^2*b - 15*(a^3
+ a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(5*(a^3 + a^2*b)*cosh(x)^3 - (a^3 - 3*a^
2*b)*cosh(x))*sinh(x)^3 + a^3 + a^2*b - (a^3 - 3*a^2*b)*cosh(x)^2 + (15*(a
^3 + a^2*b)*cosh(x)^4 - a^3 + 3*a^2*b - 6*(a^3 - 3*a^2*b)*cosh(x)^2)*sinh(
x)^2 + 2*(3*(a^3 + a^2*b)*cosh(x)^5 - 2*(a^3 - 3*a^2*b)*cosh(x)^3 - (a^3 -
3*a^2*b)*cosh(x))*sinh(x))*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(
a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^
3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 +
4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^
3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3
- a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(
14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b +
4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(
a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b
^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 +
6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh
(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x
)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2
)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^...
```

3.42.6 Sympy [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

input `integrate(tanh(x)**2/(a+b*coth(x)**2)**(3/2), x)`

output `Integral(tanh(x)**2/(a + b*coth(x)**2)**(3/2), x)`

3.42.7 Maxima [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{\frac{3}{2}}} dx$$

input `integrate(tanh(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

output `integrate(tanh(x)^2/(b*coth(x)^2 + a)^(3/2), x)`

3.42.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 540 vs. $2(75) = 150$.

Time = 0.60 (sec) , antiderivative size = 540, normalized size of antiderivative = 6.35

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx =$$

$$\frac{\frac{(a^2b^3+ab^4)e^{(2x)}}{a^5b\operatorname{sgn}(e^{(2x)}-1)+2a^4b^2\operatorname{sgn}(e^{(2x)}-1)+a^3b^3\operatorname{sgn}(e^{(2x)}-1)} + \frac{a^2b^3+ab^4}{a^5b\operatorname{sgn}(e^{(2x)}-1)+2a^4b^2\operatorname{sgn}(e^{(2x)}-1)+a^3b^3\operatorname{sgn}(e^{(2x)}-1)}}{\sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}}$$

$$\frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a + b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$+ \frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a - b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$- \frac{\log\left(\left|-\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b}\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$- \frac{4\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)^2}{\left(\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)^2 + 2\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a + b\right)}$$

input `integrate(tanh(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

output $-\left(\frac{(a^2b^3 + ab^4)e^{2x}}{(a^5b \operatorname{sgn}(e^{2x}) - 1) + 2a^4b^2 \operatorname{sgn}(e^{2x}) - 1} + a^3b^3 \operatorname{sgn}(e^{2x} - 1)\right) + \left(\frac{a^2b^3 + ab^4}{(a^5b \operatorname{sgn}(e^{2x}) - 1) + 2a^4b^2 \operatorname{sgn}(e^{2x}) - 1} + a^3b^3 \operatorname{sgn}(e^{2x} - 1)\right) / \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b) - 1/2 \log(\operatorname{abs}((\sqrt{a + b})e^{2x} - \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b)}) \sqrt{a + b} - a + b)) / ((a + b)^{3/2} \operatorname{sgn}(e^{2x} - 1)) + 1/2 \log(\operatorname{abs}((\sqrt{a + b})e^{2x} - \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b)}) \sqrt{a + b} - a - b)) / ((a + b)^{3/2} \operatorname{sgn}(e^{2x} - 1)) - 1/2 \log(\operatorname{abs}(-\sqrt{a + b})e^{2x} + \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b) - \sqrt{a + b}})) / ((a + b)^{3/2} \operatorname{sgn}(e^{2x} - 1)) - 4(\sqrt{a + b})e^{2x} - \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b) - \sqrt{a + b}}) / (((\sqrt{a + b})e^{2x} - \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b) - \sqrt{a + b}})^2 + 2(\sqrt{a + b})e^{2x} - \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b) - \sqrt{a + b}}) \sqrt{(a^4e^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b) - \sqrt{a + b}}) \sqrt{(a + b) - 3a + b} a \operatorname{sgn}(e^{2x} - 1))$

3.42.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{3/2}} dx$$

input `int(tanh(x)^2/(a + b*coth(x)^2)^(3/2), x)`

output `int(tanh(x)^2/(a + b*coth(x)^2)^(3/2), x)`

3.43 $\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx$

3.43.1 Optimal result 324
 3.43.2 Mathematica [C] (verified) 324
 3.43.3 Rubi [A] (verified) 325
 3.43.4 Maple [B] (verified) 328
 3.43.5 Fricas [B] (verification not implemented) 329
 3.43.6 Sympy [F] 329
 3.43.7 Maxima [F] 329
 3.43.8 Giac [B] (verification not implemented) 330
 3.43.9 Mupad [B] (verification not implemented) 330

3.43.1 Optimal result

Integrand size = 17, antiderivative size = 74

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} + \frac{a}{3b(a+b)(a+b \coth^2(x))^{3/2}} - \frac{1}{(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

output `arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(5/2)+1/3*a/b/(a+b)/(a+b*coth(x)^2)^(3/2)-1/(a+b)^2/(a+b*coth(x)^2)^(1/2)`

3.43.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.12 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.85

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{a(a+b) - 3b(a+b \coth^2(x)) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \coth^2(x)}{a+b}\right)}{3b(a+b)^2 (a+b \coth^2(x))^{3/2}}$$

input `Integrate[Coth[x]^3/(a + b*Coth[x]^2)^(5/2), x]`

output $(a*(a + b) - 3*b*(a + b*\text{Coth}[x]^2)*\text{Hypergeometric2F1}[-1/2, 1, 1/2, (a + b*\text{Coth}[x]^2)/(a + b))]/(3*b*(a + b)^2*(a + b*\text{Coth}[x]^2)^{(3/2)})$

3.43.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.16, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.529$, Rules used = {3042, 26, 4153, 26, 354, 87, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \tan\left(\frac{\pi}{2} + ix\right)^3}{\left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)^3}{\left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \coth^3(x)}{(1 - \coth^2(x))(b \coth^2(x) + a)^{5/2}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth^3(x)}{(1 - \coth^2(x))(a + b \coth^2(x))^{5/2}} d \coth(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\coth^2(x)}{(1 - \coth^2(x))(b \coth^2(x) + a)^{5/2}} d \coth^2(x) \\
 & \quad \downarrow \text{87} \\
 & \frac{1}{2} \left(\frac{\int \frac{1}{(1 - \coth^2(x))(b \coth^2(x) + a)^{3/2}} d \coth^2(x)}{a + b} + \frac{2a}{3b(a + b)(a + b \coth^2(x))^{3/2}} \right)
 \end{aligned}$$

3.43. $\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx$

$$\begin{array}{c}
 \downarrow 61 \\
 \frac{1}{2} \left(\frac{\int \frac{1}{(1-\coth^2(x))\sqrt{b\coth^2(x)+a}} d\coth^2(x)}{a+b} - \frac{2}{(a+b)\sqrt{a+b\coth^2(x)}} + \frac{2a}{3b(a+b)(a+b\coth^2(x))^{3/2}} \right) \\
 \downarrow 73 \\
 \frac{1}{2} \left(\frac{2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d\sqrt{b\coth^2(x)+a}}{b(a+b)} - \frac{2}{(a+b)\sqrt{a+b\coth^2(x)}} + \frac{2a}{3b(a+b)(a+b\coth^2(x))^{3/2}} \right) \\
 \downarrow 221 \\
 \frac{1}{2} \left(\frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{2}{(a+b)\sqrt{a+b\coth^2(x)}} + \frac{2a}{3b(a+b)(a+b\coth^2(x))^{3/2}} \right)
 \end{array}$$

input `Int[Coth[x]^3/(a + b*Coth[x]^2)^(5/2), x]`

output `((2*a)/(3*b*(a + b)*(a + b*Coth[x]^2)^(3/2)) + ((2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/(a + b)^(3/2) - 2/((a + b)*Sqrt[a + b*Coth[x]^2]))/(a + b))/2`

3.43.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_.)]^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_.)]^(n_.))^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.43.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 434 vs. $2(62) = 124$.

Time = 0.10 (sec) , antiderivative size = 435, normalized size of antiderivative = 5.88

method	result
derivativedivides	$\frac{1}{3b(a+b \coth(x)^2)^{\frac{3}{2}}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}} + \frac{b \coth(x)}{6(a+b)a(b(\coth(x)-1)^2+2b(\coth(x)-1))}$
default	$\frac{1}{3b(a+b \coth(x)^2)^{\frac{3}{2}}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}} + \frac{b \coth(x)}{6(a+b)a(b(\coth(x)-1)^2+2b(\coth(x)-1))}$

```
input int(coth(x)^3/(a+b*coth(x)^2)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/3/b/(a+b*coth(x)^2)^(3/2)-1/6/(a+b)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b
)^(3/2)+1/6*b/(a+b)/a/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)*coth(x)+
1/3*b/(a+b)/a^2/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)*coth(x)-1/2/(a
+b)^2/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1/2/(a+b)^2/a/(b*(coth(x
)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)*b*coth(x)+1/2/(a+b)^(5/2)*ln((2*a+2*b+2*
b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(
coth(x)-1))-1/6/(a+b)/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)-1/6*b/(a
+b)/a/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)*coth(x)-1/3*b/(a+b)/a^2/
(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)*coth(x)-1/2/(a+b)^2/(b*(1+coth
(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-1/2/(a+b)^2/a/(b*(1+coth(x))^2-2*b*(1+co
th(x))+a+b)^(1/2)*b*coth(x)+1/2/(a+b)^(5/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*
(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))
```

3.43.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2964 vs. $2(62) = 124$.

Time = 0.69 (sec) , antiderivative size = 6560, normalized size of antiderivative = 88.65

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)^3/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")`

output Too large to include

3.43.6 Sympy [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth^3(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

input `integrate(coth(x)**3/(a+b*coth(x)**2)**(5/2),x)`

output `Integral(coth(x)**3/(a + b*coth(x)**2)**(5/2), x)`

3.43.7 Maxima [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)^3}{(b \coth(x)^2 + a)^{\frac{5}{2}}} dx$$

input `integrate(coth(x)^3/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")`

output `integrate(coth(x)^3/(b*coth(x)^2 + a)^(5/2), x)`

3.43.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 951 vs. $2(62) = 124$.

Time = 0.60 (sec) , antiderivative size = 951, normalized size of antiderivative = 12.85

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)^3/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")`

output

```

1/3*(((a^8*b*sgn(e^(2*x)) - 1) + 2*a^7*b^2*sgn(e^(2*x)) - 1) - 5*a^6*b^3*sgn(e^(2*x)) - 1) - 20*a^5*b^4*sgn(e^(2*x)) - 1) - 25*a^4*b^5*sgn(e^(2*x)) - 1) - 14*a^3*b^6*sgn(e^(2*x)) - 1) - 3*a^2*b^7*sgn(e^(2*x)) - 1))*e^(2*x)/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8) - 3*(a^8*b*sgn(e^(2*x)) - 1) + 2*a^7*b^2*sgn(e^(2*x)) - 1) - a^6*b^3*sgn(e^(2*x)) - 1) - 4*a^5*b^4*sgn(e^(2*x)) - 1) - a^4*b^5*sgn(e^(2*x)) - 1) + 2*a^3*b^6*sgn(e^(2*x)) - 1) + a^2*b^7*sgn(e^(2*x)) - 1))/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))*e^(2*x) + 3*(a^8*b*sgn(e^(2*x)) - 1) + 2*a^7*b^2*sgn(e^(2*x)) - 1) - a^6*b^3*sgn(e^(2*x)) - 1) - 4*a^5*b^4*sgn(e^(2*x)) - 1) - a^4*b^5*sgn(e^(2*x)) - 1) + 2*a^3*b^6*sgn(e^(2*x)) - 1) + a^2*b^7*sgn(e^(2*x)) - 1))/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))*e^(2*x) - (a^8*b*sgn(e^(2*x)) - 1) + 2*a^7*b^2*sgn(e^(2*x)) - 1) - 5*a^6*b^3*sgn(e^(2*x)) - 1) - 20*a^5*b^4*sgn(e^(2*x)) - 1) - 25*a^4*b^5*sgn(e^(2*x)) - 1) - 14*a^3*b^6*sgn(e^(2*x)) - 1) - 3*a^2*b^7*sgn(e^(2*x)) - 1))/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))/(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b)^(3/2) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a + b))/((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^(2*x) - 1)) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*...

```

3.43.9 Mupad [B] (verification not implemented)

Time = 4.25 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.11

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a(2a^2 + 4ab + 2b^2)}}{2(a+b)^{5/2}}\right)}{(a+b)^{5/2}} + \frac{\frac{a}{3(a+b)} - \frac{b(b \coth(x)^2 + a)}{(a+b)^2}}{b(b \coth(x)^2 + a)^{3/2}}$$

input `int(coth(x)^3/(a + b*coth(x)^2)^(5/2),x)`

output `atanh(((a + b*coth(x)^2)^(1/2)*(4*a*b + 2*a^2 + 2*b^2))/(2*(a + b)^(5/2)))/
(a + b)^(5/2) + (a/(3*(a + b)) - (b*(a + b*coth(x)^2))/(a + b)^2)/(b*(a +
b*coth(x)^2)^(3/2))`

3.44
$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx$$

3.44.1	Optimal result	332
3.44.2	Mathematica [C] (warning: unable to verify)	332
3.44.3	Rubi [A] (verified)	333
3.44.4	Maple [B] (verified)	336
3.44.5	Fricas [B] (verification not implemented)	337
3.44.6	Sympy [F]	337
3.44.7	Maxima [F]	337
3.44.8	Giac [B] (verification not implemented)	338
3.44.9	Mupad [F(-1)]	338

3.44.1 Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{5/2}} - \frac{\coth(x)}{3(a+b)(a+b \coth^2(x))^{3/2}} - \frac{(2a-b) \coth(x)}{3a(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

```
output arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))/(a+b)^(5/2)-1/3*coth(x)
/(a+b)/(a+b*coth(x)^2)^(3/2)-1/3*(2*a-b)*coth(x)/a/(a+b)^2/(a+b*coth(x)^2)^(1/2)
```

3.44.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.07 (sec) , antiderivative size = 215, normalized size of antiderivative = 2.44

$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\cosh^2(x) \coth(x) \left(\frac{4(a+b) \cosh^2(x) (a+b \coth^2(x)) \operatorname{Hypergeometric2F1}\left(2, 2, \frac{9}{2}, \frac{(a+b) \cosh^2(x)}{a}\right)}{35a^2} - \frac{(-5a-2b \coth^2(x)) \left(3 \arcsin\left(\sqrt{\frac{a+b \coth^2(x)}{a+b}}\right)\right)}{3a^2 \sqrt{a+b \coth^2(x)}} \right)}{3a^2 \sqrt{a+b \coth^2(x)}} \left(1 \right)$$

3.44.
$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx$$

input `Integrate[Coth[x]^2/(a + b*Coth[x]^2)^(5/2),x]`

output `-1/3*(Cosh[x]^2*Coth[x]*((4*(a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Hypergeometric2F1[2, 2, 9/2, ((a + b)*Cosh[x]^2)/a]/(35*a^2) - ((-5*a - 2*b*Coth[x]^2)*(3*ArcSin[Sqrt[((a + b)*Cosh[x]^2)/a]]*(a + b*Coth[x]^2)^2 - a*(-4*b*Coth[x]^2 + a*(-3 - Coth[x]^2))*Csch[x]^2*Sqrt[-(((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2]])*Tanh[x]^4)/(3*a*(a + b)^2*Sqrt[-(((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2]])))/(a^2*Sqrt[a + b*Coth[x]^2]*(1 + (b*Coth[x]^2)/a))`

3.44.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.529$, Rules used = {3042, 25, 4153, 25, 373, 402, 27, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan\left(\frac{\pi}{2} + ix\right)^2}{\left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\tan\left(ix + \frac{\pi}{2}\right)^2}{\left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -\int -\frac{\coth^2(x)}{(1 - \coth^2(x))(b \coth^2(x) + a)^{5/2}} d \coth(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\coth^2(x)}{(1 - \coth^2(x))(a + b \coth^2(x))^{5/2}} d \coth(x) \\
 & \quad \downarrow \text{373}
 \end{aligned}$$

3.44. $\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{2 \operatorname{coth}^2(x)+1}{(1-\operatorname{coth}^2(x))(b \operatorname{coth}^2(x)+a)^{3/2}} d \operatorname{coth}(x)}{3(a+b)} - \frac{\operatorname{coth}(x)}{3(a+b)(a+b \operatorname{coth}^2(x))^{3/2}} \\
 & \quad \downarrow 402 \\
 & \frac{\int -\frac{3a}{(1-\operatorname{coth}^2(x))\sqrt{b \operatorname{coth}^2(x)+a}} d \operatorname{coth}(x)}{3(a+b)} - \frac{(2a-b) \operatorname{coth}(x)}{a(a+b)\sqrt{a+b \operatorname{coth}^2(x)}} - \frac{\operatorname{coth}(x)}{3(a+b)(a+b \operatorname{coth}^2(x))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{3 \int \frac{1}{(1-\operatorname{coth}^2(x))\sqrt{b \operatorname{coth}^2(x)+a}} d \operatorname{coth}(x)}{3(a+b)} - \frac{(2a-b) \operatorname{coth}(x)}{a(a+b)\sqrt{a+b \operatorname{coth}^2(x)}} - \frac{\operatorname{coth}(x)}{3(a+b)(a+b \operatorname{coth}^2(x))^{3/2}} \\
 & \quad \downarrow 291 \\
 & \frac{3 \int \frac{1}{1-\frac{(a+b) \operatorname{coth}^2(x)}{b \operatorname{coth}^2(x)+a}} d \frac{\operatorname{coth}(x)}{\sqrt{b \operatorname{coth}^2(x)+a}}}{3(a+b)} - \frac{(2a-b) \operatorname{coth}(x)}{a(a+b)\sqrt{a+b \operatorname{coth}^2(x)}} - \frac{\operatorname{coth}(x)}{3(a+b)(a+b \operatorname{coth}^2(x))^{3/2}} \\
 & \quad \downarrow 219 \\
 & \frac{3 \operatorname{arctanh}\left(\frac{\sqrt{a+b} \operatorname{coth}(x)}{\sqrt{a+b \operatorname{coth}^2(x)}}\right)}{(a+b)^{3/2}} - \frac{(2a-b) \operatorname{coth}(x)}{a(a+b)\sqrt{a+b \operatorname{coth}^2(x)}} - \frac{\operatorname{coth}(x)}{3(a+b)(a+b \operatorname{coth}^2(x))^{3/2}}
 \end{aligned}$$

input `Int [Coth[x]^2/(a + b*Coth[x]^2)^(5/2),x]`

output `-1/3*Coth[x]/((a + b)*(a + b*Coth[x]^2)^(3/2)) + ((3*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]])/(a + b)^(3/2) - ((2*a - b)*Coth[x])/(a*(a + b)*Sqrt[a + b*Coth[x]^2]))/(3*(a + b))`

3.44.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 373 `Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[e*(e*x)^(m - 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*(b*c - a*d)*(p + 1))), x] - Simp[e^2/(2*(b*c - a*d)*(p + 1)) Int[(e*x)^(m - 2)*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(m - 1) + d*(m + 2*p + 2*q + 3)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[m, 1] && LeQ[m, 3] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`
- rule 402 `Int[((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*2*(b*c - a*d)*(p + 1))), x] + Simp[1/(a*2*(b*c - a*d)*(p + 1)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(b*e - a*f) + e*2*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(2*(p + q + 2) + 1)*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && LtQ[p, -1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4153 Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.44.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 453 vs. $2(74) = 148$.

Time = 0.10 (sec) , antiderivative size = 454, normalized size of antiderivative = 5.16

method	result
derivativedivides	$-\frac{\coth(x)}{3a(a+b\coth(x)^2)^{3/2}} - \frac{2\coth(x)}{3a^2\sqrt{a+b\coth(x)^2}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}} + \frac{1}{6(a+b)a(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}}$
default	$-\frac{\coth(x)}{3a(a+b\coth(x)^2)^{3/2}} - \frac{2\coth(x)}{3a^2\sqrt{a+b\coth(x)^2}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}} + \frac{1}{6(a+b)a(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}}$

```
input int(coth(x)^2/(a+b*coth(x)^2)^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/3*coth(x)/a/(a+b*coth(x)^2)^(3/2)-2/3/a^2*coth(x)/(a+b*coth(x)^2)^(1/2)
-1/6/(a+b)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)+1/6*b/(a+b)/a/(b*(c
oth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)*coth(x)+1/3*b/(a+b)/a^2/(b*(coth(x)
-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)*coth(x)-1/2/(a+b)^2/(b*(coth(x)-1)^2+2*b*
(coth(x)-1)+a+b)^(1/2)+1/2/(a+b)^2/a/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)
^(1/2)*b*coth(x)+1/2/(a+b)^(5/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)
*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))+1/6/(a+b)/(b*(1
+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)+1/6*b/(a+b)/a/(b*(1+coth(x))^2-2*b*
(1+coth(x))+a+b)^(3/2)*coth(x)+1/3*b/(a+b)/a^2/(b*(1+coth(x))^2-2*b*(1+cot
h(x))+a+b)^(1/2)*coth(x)+1/2/(a+b)^2/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)
^(1/2)+1/2/(a+b)^2/a/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)*b*coth(x)
-1/2/(a+b)^(5/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^
2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))
```

3.44.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3016 vs. $2(74) = 148$.

Time = 0.68 (sec) , antiderivative size = 6591, normalized size of antiderivative = 74.90

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")`

output Too large to include

3.44.6 Sympy [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth^2(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

input `integrate(coth(x)**2/(a+b*coth(x)**2)**(5/2),x)`

output `Integral(coth(x)**2/(a + b*coth(x)**2)**(5/2), x)`

3.44.7 Maxima [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{\frac{5}{2}}} dx$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")`

output `integrate(coth(x)^2/(b*coth(x)^2 + a)^(5/2), x)`

3.44.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 952 vs. 2(74) = 148.

Time = 0.55 (sec) , antiderivative size = 952, normalized size of antiderivative = 10.82

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")`

output

```
-1/3*(((3*a^7*b^2*sgn(e^(2*x)) - 1) + 14*a^6*b^3*sgn(e^(2*x)) - 1) + 25*a^5
*b^4*sgn(e^(2*x)) - 1) + 20*a^4*b^5*sgn(e^(2*x)) - 1) + 5*a^3*b^6*sgn(e^(2*x)
) - 1) - 2*a^2*b^7*sgn(e^(2*x)) - 1) - a*b^8*sgn(e^(2*x)) - 1))*e^(2*x)/(a^8
*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*
b^8) - 3*(a^7*b^2*sgn(e^(2*x)) - 1) + 2*a^6*b^3*sgn(e^(2*x)) - 1) - a^5*b^4*
sgn(e^(2*x)) - 1) - 4*a^4*b^5*sgn(e^(2*x)) - 1) - a^3*b^6*sgn(e^(2*x)) - 1) +
2*a^2*b^7*sgn(e^(2*x)) - 1) + a*b^8*sgn(e^(2*x)) - 1))/(a^8*b^2 + 6*a^7*b^3
+ 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))*e^(2*x) -
3*(a^7*b^2*sgn(e^(2*x)) - 1) + 2*a^6*b^3*sgn(e^(2*x)) - 1) - a^5*b^4*sgn(e(
2*x)) - 1) - 4*a^4*b^5*sgn(e^(2*x)) - 1) - a^3*b^6*sgn(e^(2*x)) - 1) + 2*a^2*
b^7*sgn(e^(2*x)) - 1) + a*b^8*sgn(e^(2*x)) - 1))/(a^8*b^2 + 6*a^7*b^3 + 15*a
^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))*e^(2*x) + (3*a^7*
b^2*sgn(e^(2*x)) - 1) + 14*a^6*b^3*sgn(e^(2*x)) - 1) + 25*a^5*b^4*sgn(e^(2*x)
) - 1) + 20*a^4*b^5*sgn(e^(2*x)) - 1) + 5*a^3*b^6*sgn(e^(2*x)) - 1) - 2*a^2*
b^7*sgn(e^(2*x)) - 1) - a*b^8*sgn(e^(2*x)) - 1))/(a^8*b^2 + 6*a^7*b^3 + 15*a
^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))/(a*e^(4*x) + b*e
(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b)^(3/2) - 1/2*log(abs((sqrt(a + b)
)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b
))*sqrt(a + b) - a + b))/((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^(2*x)) - 1)
) + 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a...
```

3.44.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{5/2}} dx$$

input `int(coth(x)^2/(a + b*coth(x)^2)^(5/2),x)`

output `int(coth(x)^2/(a + b*coth(x)^2)^(5/2), x)`

3.45
$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx$$

3.45.1	Optimal result	340
3.45.2	Mathematica [C] (verified)	340
3.45.3	Rubi [A] (verified)	341
3.45.4	Maple [B] (verified)	344
3.45.5	Fricas [B] (verification not implemented)	344
3.45.6	Sympy [F]	345
3.45.7	Maxima [F]	345
3.45.8	Giac [B] (verification not implemented)	345
3.45.9	Mupad [B] (verification not implemented)	346

3.45.1 Optimal result

Integrand size = 15, antiderivative size = 70

$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} - \frac{1}{3(a+b)(a+b \coth^2(x))^{3/2}} - \frac{1}{(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

output `arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(5/2)-1/3/(a+b)/(a+b*coth(x)^2)^(3/2)-1/(a+b)^2/(a+b*coth(x)^2)^(1/2)`

3.45.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.61

$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx = -\frac{\operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \coth^2(x)}{a+b}\right)}{3(a+b)(a+b \coth^2(x))^{3/2}}$$

input `Integrate[Coth[x]/(a + b*Coth[x]^2)^(5/2),x]`

output $-1/3*\text{Hypergeometric2F1}[-3/2, 1, -1/2, (a + b*\text{Coth}[x]^2)/(a + b)]/((a + b)*(a + b*\text{Coth}[x]^2)^{(3/2)})$

3.45.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.17, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 26, 4153, 26, 353, 61, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan\left(\frac{\pi}{2} + ix\right)}{\left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)}{\left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -i \int \frac{i \coth(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{5/2}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x)}{(1 - \coth^2(x)) (a + b \coth^2(x))^{5/2}} d \coth(x) \\
 & \quad \downarrow \text{353} \\
 & \frac{1}{2} \int \frac{1}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{5/2}} d \coth^2(x) \\
 & \quad \downarrow \text{61} \\
 & \frac{1}{2} \left(\frac{\int \frac{1}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth^2(x)}{a + b} - \frac{2}{3(a + b) (a + b \coth^2(x))^{3/2}} \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 61 \\
 \frac{1}{2} \left(\frac{\int \frac{1}{(1-\coth^2(x))\sqrt{b\coth^2(x)+a}} d\coth^2(x)}{a+b} - \frac{2}{(a+b)\sqrt{a+b\coth^2(x)}} - \frac{2}{3(a+b)(a+b\coth^2(x))^{3/2}} \right) \\
 \downarrow 73 \\
 \frac{1}{2} \left(\frac{2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d\sqrt{b\coth^2(x)+a}}{b(a+b)} - \frac{2}{(a+b)\sqrt{a+b\coth^2(x)}} - \frac{2}{3(a+b)(a+b\coth^2(x))^{3/2}} \right) \\
 \downarrow 221 \\
 \frac{1}{2} \left(\frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{2}{(a+b)\sqrt{a+b\coth^2(x)}} - \frac{2}{3(a+b)(a+b\coth^2(x))^{3/2}} \right)
 \end{array}$$

input `Int[Coth[x]/(a + b*Coth[x]^2)^(5/2),x]`

output `(-2/(3*(a + b)*(a + b*Coth[x]^2)^(3/2)) + ((2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/(a + b)^(3/2) - 2/((a + b)*Sqrt[a + b*Coth[x]^2]))/(a + b)/2`

3.45.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

- rule 61 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 353 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4153 `Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f*f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.45.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 419 vs. 2(58) = 116.

Time = 0.08 (sec) , antiderivative size = 420, normalized size of antiderivative = 6.00

method	result
derivativedivides	$-\frac{1}{6(a+b)\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{3/2}} + \frac{b \coth(x)}{6(a+b)a\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{3/2}} + \frac{1}{3(a+b)a}$
default	$-\frac{1}{6(a+b)\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{3/2}} + \frac{b \coth(x)}{6(a+b)a\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{3/2}} + \frac{1}{3(a+b)a}$

input `int(coth(x)/(a+b*coth(x)^2)^(5/2),x,method=_RETURNVERBOSE)`

output

```
-1/6/(a+b)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)+1/6*b/(a+b)/a/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)*coth(x)+1/3*b/(a+b)/a^2/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)*coth(x)-1/2/(a+b)^2/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1/2/(a+b)^2/a/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)*b*coth(x)+1/2/(a+b)^(5/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2))*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))-1/6/(a+b)/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)-1/6*b/(a+b)/a/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)*coth(x)-1/3*b/(a+b)/a^2/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)*coth(x)-1/2/(a+b)^2/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-1/2/(a+b)^2/a/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)*b*coth(x)+1/2/(a+b)^(5/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2))*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))
```

3.45.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2552 vs. 2(58) = 116.

Time = 0.63 (sec) , antiderivative size = 5736, normalized size of antiderivative = 81.94

$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")`

output Too large to include

3.45.6 Sympy [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

input `integrate(coth(x)/(a+b*coth(x)**2)**(5/2),x)`

output `Integral(coth(x)/(a + b*coth(x)**2)**(5/2), x)`

3.45.7 Maxima [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)}{(b \coth(x)^2 + a)^{\frac{5}{2}}} dx$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")`

output `integrate(coth(x)/(b*coth(x)^2 + a)^(5/2), x)`

3.45.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 909 vs. 2(58) = 116.

Time = 0.58 (sec) , antiderivative size = 909, normalized size of antiderivative = 12.99

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")`

output

```

-4/3*(((a^7*b^2 + 5*a^6*b^3 + 10*a^5*b^4 + 10*a^4*b^5 + 5*a^3*b^6 + a^2*b^7)*e^(2*x))/(a^8*b^2*sgn(e^(2*x) - 1) + 6*a^7*b^3*sgn(e^(2*x) - 1) + 15*a^6*b^4*sgn(e^(2*x) - 1) + 20*a^5*b^5*sgn(e^(2*x) - 1) + 15*a^4*b^6*sgn(e^(2*x) - 1) + 6*a^3*b^7*sgn(e^(2*x) - 1) + a^2*b^8*sgn(e^(2*x) - 1)) - 3*(a^7*b^2 + 4*a^6*b^3 + 6*a^5*b^4 + 4*a^4*b^5 + a^3*b^6)/(a^8*b^2*sgn(e^(2*x) - 1) + 6*a^7*b^3*sgn(e^(2*x) - 1) + 15*a^6*b^4*sgn(e^(2*x) - 1) + 20*a^5*b^5*sgn(e^(2*x) - 1) + 15*a^4*b^6*sgn(e^(2*x) - 1) + 6*a^3*b^7*sgn(e^(2*x) - 1) + a^2*b^8*sgn(e^(2*x) - 1))) * e^(2*x) + 3*(a^7*b^2 + 4*a^6*b^3 + 6*a^5*b^4 + 4*a^4*b^5 + a^3*b^6)/(a^8*b^2*sgn(e^(2*x) - 1) + 6*a^7*b^3*sgn(e^(2*x) - 1) + 15*a^6*b^4*sgn(e^(2*x) - 1) + 20*a^5*b^5*sgn(e^(2*x) - 1) + 15*a^4*b^6*sgn(e^(2*x) - 1) + 6*a^3*b^7*sgn(e^(2*x) - 1) + a^2*b^8*sgn(e^(2*x) - 1))) * e^(2*x) - (a^7*b^2 + 5*a^6*b^3 + 10*a^5*b^4 + 10*a^4*b^5 + 5*a^3*b^6 + a^2*b^7)/(a^8*b^2*sgn(e^(2*x) - 1) + 6*a^7*b^3*sgn(e^(2*x) - 1) + 15*a^6*b^4*sgn(e^(2*x) - 1) + 20*a^5*b^5*sgn(e^(2*x) - 1) + 15*a^4*b^6*sgn(e^(2*x) - 1) + 6*a^3*b^7*sgn(e^(2*x) - 1) + a^2*b^8*sgn(e^(2*x) - 1)))/(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b)^(3/2) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a + b))/((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^(2*x) - 1)) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a - b))/((a^2 + ...

```

3.45.9 Mupad [B] (verification not implemented)

Time = 4.33 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.09

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a} (2a^2 + 4ab + 2b^2)}{2(a+b)^{5/2}}\right)}{(a+b)^{5/2}} - \frac{\frac{1}{3(a+b)} + \frac{b \coth(x)^2 + a}{(a+b)^2}}{(b \coth(x)^2 + a)^{3/2}}$$

input `int(coth(x)/(a + b*coth(x)^2)^(5/2), x)`

output `atanh(((a + b*coth(x)^2)^(1/2)*(4*a*b + 2*a^2 + 2*b^2))/(2*(a + b)^(5/2)))/(a + b)^(5/2) - (1/(3*(a + b)) + (a + b*coth(x)^2)/(a + b)^2)/(a + b*coth(x)^2)^(3/2)`

$$3.46 \quad \int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx$$

3.46.1	Optimal result	347
3.46.2	Mathematica [C] (verified)	347
3.46.3	Rubi [A] (verified)	348
3.46.4	Maple [F]	352
3.46.5	Fricas [B] (verification not implemented)	352
3.46.6	Sympy [F]	352
3.46.7	Maxima [F]	353
3.46.8	Giac [F(-2)]	353
3.46.9	Mupad [F(-1)]	353

3.46.1 Optimal result

Integrand size = 15, antiderivative size = 108

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{a^{5/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} + \frac{b}{3a(a+b)(a+b \coth^2(x))^{3/2}} + \frac{b(2a+b)}{a^2(a+b)^2\sqrt{a+b \coth^2(x)}}$$

output `-arctanh((a+b*coth(x)^2)^(1/2)/a^(1/2))/a^(5/2)+arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))/(a+b)^(5/2)+1/3*b/a/(a+b)/(a+b*coth(x)^2)^(3/2)+b*(2*a+b)/a^2/(a+b)^2/(a+b*coth(x)^2)^(1/2)`

3.46.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.68

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{-a \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \coth^2(x)}{a+b}\right) + (a+b) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \coth^2(x)}{a+b}\right)}{3a(a+b)(a+b \coth^2(x))^{3/2}}$$

input `Integrate[Tanh[x]/(a + b*Coth[x]^2)^(5/2), x]`

3.46. $\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx$

output $(-a \operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, (a + b \operatorname{Coth}[x]^2)/(a + b)]) + (a + b) \operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, 1 + (b \operatorname{Coth}[x]^2)/a] / (3a(a + b)(a + b \operatorname{Coth}[x]^2)^{3/2})$

3.46.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.31, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {3042, 26, 4153, 26, 354, 96, 25, 169, 27, 174, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{(a + b \operatorname{coth}^2(x))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\tan\left(\frac{\pi}{2} + ix\right) \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{\tan\left(ix + \frac{\pi}{2}\right) \left(a - b \tan\left(ix + \frac{\pi}{2}\right)^2\right)^{5/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & i \int -\frac{i \tanh(x)}{(1 - \operatorname{coth}^2(x)) (b \operatorname{coth}^2(x) + a)^{5/2}} d \operatorname{coth}(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\tanh(x)}{(1 - \operatorname{coth}^2(x)) (a + b \operatorname{coth}^2(x))^{5/2}} d \operatorname{coth}(x) \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int \frac{\tanh(x)}{(1 - \operatorname{coth}^2(x)) (b \operatorname{coth}^2(x) + a)^{5/2}} d \operatorname{coth}^2(x) \\
 & \quad \downarrow \text{96} \\
 & \frac{1}{2} \left(\frac{2b}{3a(a + b) (a + b \operatorname{coth}^2(x))^{3/2}} - \frac{\int -\frac{(-b \operatorname{coth}^2(x) + a + b) \tanh(x)}{(1 - \operatorname{coth}^2(x)) (b \operatorname{coth}^2(x) + a)^{3/2}} d \operatorname{coth}^2(x)}{a(a + b)} \right)
 \end{aligned}$$

3.46. $\int \frac{\tanh(x)}{(a + b \operatorname{coth}^2(x))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{1}{2} \left(\frac{\int \frac{(-b \coth^2(x) + a + b) \tanh(x)}{(1 - \coth^2(x))(b \coth^2(x) + a)^{3/2}} d \coth^2(x)}{a(a+b)} + \frac{2b}{3a(a+b)(a+b \coth^2(x))^{3/2}} \right) \\
 & \downarrow 169 \\
 & \frac{1}{2} \left(\frac{2 \int \frac{((a+b)^2 - b(2a+b) \coth^2(x)) \tanh(x)}{2(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x)}{a(a+b)} + \frac{2b(2a+b)}{a(a+b) \sqrt{a+b \coth^2(x)}} + \frac{2b}{3a(a+b)(a+b \coth^2(x))^{3/2}} \right) \\
 & \downarrow 27 \\
 & \frac{1}{2} \left(\frac{\int \frac{((a+b)^2 - b(2a+b) \coth^2(x)) \tanh(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x)}{a(a+b)} + \frac{2b(2a+b)}{a(a+b) \sqrt{a+b \coth^2(x)}} + \frac{2b}{3a(a+b)(a+b \coth^2(x))^{3/2}} \right) \\
 & \downarrow 174 \\
 & \frac{1}{2} \left(\frac{a^2 \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth^2(x) + (a+b)^2 \int \frac{\tanh(x)}{\sqrt{b \coth^2(x) + a}} d \coth^2(x)}{a(a+b)} + \frac{2b(2a+b)}{a(a+b) \sqrt{a+b \coth^2(x)}} + \frac{2b}{3a(a+b)(a+b \coth^2(x))^{3/2}} \right) \\
 & \downarrow 73 \\
 & \frac{1}{2} \left(\frac{2a^2 \int \frac{1}{\frac{a+b}{b} - \frac{\coth^4(x)}{b}} d \sqrt{b \coth^2(x) + a} + 2(a+b)^2 \int \frac{1}{\frac{\coth^4(x)}{b} - \frac{a}{b}} d \sqrt{b \coth^2(x) + a}}{a(a+b)} + \frac{2b(2a+b)}{a(a+b) \sqrt{a+b \coth^2(x)}} + \frac{2b}{3a(a+b)(a+b \coth^2(x))^{3/2}} \right) \\
 & \downarrow 221
 \end{aligned}$$

3.46. $\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx$

$$\frac{1}{2} \left(\frac{\frac{2a^2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{2(a+b)^2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}}}{a(a+b)} + \frac{2b(2a+b)}{a(a+b)\sqrt{a+b \coth^2(x)}} + \frac{2b}{3a(a+b)(a+b \coth^2(x))^{3/2}} \right)$$

input `Int[Tanh[x]/(a + b*Coth[x]^2)^(5/2), x]`

output `((2*b)/(3*a*(a + b)*(a + b*Coth[x]^2)^(3/2)) + (((-2*(a + b)^2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]])/Sqrt[a] + (2*a^2*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]])/Sqrt[a + b]))/(a*(a + b)) + (2*b*(2*a + b))/(a*(a + b)*Sqrt[a + b*Coth[x]^2]))/(a*(a + b))/2`

3.46.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 96 `Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] := Simp[f*((e + f*x)^(p + 1)/((p + 1)*(b*e - a*f)*(d*e - c*f))), x] + Simp[1/((b*e - a*f)*(d*e - c*f)) Int[(b*d*e - b*c*f - a*d*f - b*d*f*x)*((e + f*x)^(p + 1)/((a + b*x)*(c + d*x))), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[p, -1]`

rule 169 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 174 `Int((((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] := Simp[(b*g - a*h)/(b*c - a*d) Int[(e + f*x)^p/(a + b*x), x], x] - Simp[(d*g - c*h)/(b*c - a*d) Int[(e + f*x)^p/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4153 `Int((((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.46.4 Maple [F]

$$\int \frac{\tanh(x)}{(a + b \coth(x)^2)^{\frac{5}{2}}} dx$$

input `int(tanh(x)/(a+b*coth(x)^2)^(5/2),x)`

output `int(tanh(x)/(a+b*coth(x)^2)^(5/2),x)`

3.46.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4452 vs. 2(90) = 180.

Time = 1.24 (sec) , antiderivative size = 19199, normalized size of antiderivative = 177.77

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx = \text{Too large to display}$$

input `integrate(tanh(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")`

output `Too large to include`

3.46.6 Sympy [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx = \int \frac{\tanh(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

input `integrate(tanh(x)/(a+b*coth(x)**2)**(5/2),x)`

output `Integral(tanh(x)/(a + b*coth(x)**2)**(5/2), x)`

3.46.7 Maxima [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{5/2}} dx$$

input `integrate(tanh(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")`

output `integrate(tanh(x)/(b*coth(x)^2 + a)^(5/2), x)`

3.46.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(tanh(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT>Error: Bad Argument Type`

3.46.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{5/2}} dx$$

input `int(tanh(x)/(a + b*coth(x)^2)^(5/2),x)`

output `int(tanh(x)/(a + b*coth(x)^2)^(5/2), x)`

3.47 $\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx$

3.47.1 Optimal result 354
 3.47.2 Mathematica [C] (warning: unable to verify) 354
 3.47.3 Rubi [A] (verified) 355
 3.47.4 Maple [F] 359
 3.47.5 Fricas [B] (verification not implemented) 359
 3.47.6 Sympy [F] 360
 3.47.7 Maxima [F] 360
 3.47.8 Giac [B] (verification not implemented) 360
 3.47.9 Mupad [F(-1)] 361

3.47.1 Optimal result

Integrand size = 17, antiderivative size = 131

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{5/2}} + \frac{b \tanh(x)}{3a(a+b)(a+b \coth^2(x))^{3/2}}$$

$$+ \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} - \frac{(3a+2b)(a+4b) \sqrt{a+b \coth^2(x)} \tanh(x)}{3a^3(a+b)^2}$$

```
output arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))/(a+b)^(5/2)+1/3*b*tanh(x)/a/(a+b)/(a+b*coth(x)^2)^(3/2)+1/3*b*(7*a+4*b)*tanh(x)/a^2/(a+b)^2/(a+b*coth(x)^2)^(1/2)-1/3*(3*a+2*b)*(a+4*b)*(a+b*coth(x)^2)^(1/2)*tanh(x)/a^3/(a+b)^2
```

3.47.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.54 (sec) , antiderivative size = 1350, normalized size of antiderivative = 10.31

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[Tanh[x]^2/(a + b*Coth[x]^2)^(5/2),x]`

output $(\text{Sinh}[x]^2*((16*b^3*((-1)*\text{Coth}[x] + 1*\text{Coth}[x]^3)^2)/(a*(a + b)^2) + (40*b*\text{Csch}[x]^2)/(a + b) + (160*b^2*\text{Coth}[x]^2*\text{Csch}[x]^2)/(3*a*(a + b)) + (64*b^3*\text{Coth}[x]^4*\text{Csch}[x]^2)/(3*a^2*(a + b)) - (40*b^2*\text{Csch}[x]^4)/(a + b)^2 + (92*(a + b)*\text{Cosh}[x]^2*\text{Hypergeometric2F1}[2, 2, 9/2, ((a + b)*\text{Cosh}[x]^2)/a])/(105*a) + (124*b*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^2*\text{Hypergeometric2F1}[2, 2, 9/2, ((a + b)*\text{Cosh}[x]^2)/a])/(35*a^2) + (152*b^2*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^4*\text{Hypergeometric2F1}[2, 2, 9/2, ((a + b)*\text{Cosh}[x]^2)/a])/(35*a^3) + (176*b^3*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^6*\text{Hypergeometric2F1}[2, 2, 9/2, ((a + b)*\text{Cosh}[x]^2)/a])/(105*a^4) + (24*(a + b)*\text{Cosh}[x]^2*\text{HypergeometricPFQ}[{2, 2, 2}, \{1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(35*a) + (16*b*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^2*\text{HypergeometricPFQ}[{2, 2, 2}, \{1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(7*a^2) + (88*b^2*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^4*\text{HypergeometricPFQ}[{2, 2, 2}, \{1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(35*a^3) + (32*b^3*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^6*\text{HypergeometricPFQ}[{2, 2, 2}, \{1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(35*a^4) + (16*(a + b)*\text{Cosh}[x]^2*\text{HypergeometricPFQ}[{2, 2, 2, 2}, \{1, 1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(105*a) + (16*b*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^2*\text{HypergeometricPFQ}[{2, 2, 2, 2}, \{1, 1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(35*a^2) + (16*b^2*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^4*\text{HypergeometricPFQ}[{2, 2, 2, 2}, \{1, 1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(35*a^3) + (16*b^3*(a + b)*\text{Cosh}[x]^2*\text{Coth}[x]^6*\text{HypergeometricPFQ}[{2, 2, 2, 2}, \{1, 1, 9/2\}, ((a + b)*\text{Cosh}[x]^2)/a])/(105*a^4) + (2...$

3.47.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.13, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.706$, Rules used = {3042, 25, 4153, 25, 374, 25, 441, 25, 445, 27, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx$$

↓ 3042

$$\int -\frac{1}{\tan\left(\frac{\pi}{2} + ix\right)^2 \left(a - b \tan\left(\frac{\pi}{2} + ix\right)^2\right)^{5/2}} dx$$

↓ 25

3.47. $\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx$

$$\begin{aligned}
& - \int \frac{1}{\tan\left(ix + \frac{\pi}{2}\right)^2 \left(a - b \tan\left(ix + \frac{\pi}{2}\right)\right)^{5/2}} dx \\
& \quad \downarrow \text{4153} \\
& - \int \frac{\tanh^2(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{5/2}} d \coth(x) \\
& \quad \downarrow \text{25} \\
& \int \frac{\tanh^2(x)}{(1 - \coth^2(x)) (a + b \coth^2(x))^{5/2}} d \coth(x) \\
& \quad \downarrow \text{374} \\
& \frac{b \tanh(x)}{3a(a+b) (a + b \coth^2(x))^{3/2}} - \frac{\int \frac{(-4b \coth^2(x) + 3a + 4b) \tanh^2(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth(x)}{3a(a+b)} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{(-4b \coth^2(x) + 3a + 4b) \tanh^2(x)}{(1 - \coth^2(x)) (b \coth^2(x) + a)^{3/2}} d \coth(x)}{3a(a+b)} + \frac{b \tanh(x)}{3a(a+b) (a + b \coth^2(x))^{3/2}} \\
& \quad \downarrow \text{441} \\
& \frac{b(7a+4b) \tanh(x)}{a(a+b) \sqrt{a + b \coth^2(x)}} - \frac{\int \frac{((3a+2b)(a+4b) - 2b(7a+4b) \coth^2(x)) \tanh^2(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x)}{a(a+b)} + \frac{b \tanh(x)}{3a(a+b) (a + b \coth^2(x))^{3/2}} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{((3a+2b)(a+4b) - 2b(7a+4b) \coth^2(x)) \tanh^2(x)}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x)}{a(a+b)} + \frac{b(7a+4b) \tanh(x)}{a(a+b) \sqrt{a + b \coth^2(x)}} + \frac{b \tanh(x)}{3a(a+b) (a + b \coth^2(x))^{3/2}} \\
& \quad \downarrow \text{445} \\
& \frac{\int \frac{3a^3}{(1 - \coth^2(x)) \sqrt{b \coth^2(x) + a}} d \coth(x)}{a} - \frac{(3a+2b)(a+4b) \tanh(x) \sqrt{a + b \coth^2(x)}}{a} + \frac{b(7a+4b) \tanh(x)}{a(a+b) \sqrt{a + b \coth^2(x)}} + \\
& \quad \frac{3a(a+b)}{b \tanh(x)} \\
& \quad \frac{3a(a+b) (a + b \coth^2(x))^{3/2}}{b \tanh(x)} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.47. $\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx$

$$\begin{aligned}
& \frac{3a^2 \int \frac{1}{(1-\coth^2(x))\sqrt{b\coth^2(x)+a}} d\coth(x) - \frac{(3a+2b)(a+4b)\tanh(x)\sqrt{a+b\coth^2(x)}}{a}}{a(a+b)} + \frac{b(7a+4b)\tanh(x)}{a(a+b)\sqrt{a+b\coth^2(x)}} + \\
& \frac{3a(a+b)}{b\tanh(x)} \\
& \frac{3a(a+b)(a+b\coth^2(x))^{3/2}}{3a(a+b)(a+b\coth^2(x))^{3/2}} \\
& \quad \downarrow \text{291} \\
& \frac{3a^2 \int \frac{1}{1-\frac{(a+b)\coth^2(x)}{b\coth^2(x)+a}} d\frac{\coth(x)}{\sqrt{b\coth^2(x)+a}} - \frac{(3a+2b)(a+4b)\tanh(x)\sqrt{a+b\coth^2(x)}}{a}}{a(a+b)} + \frac{b(7a+4b)\tanh(x)}{a(a+b)\sqrt{a+b\coth^2(x)}} + \\
& \frac{3a(a+b)}{b\tanh(x)} \\
& \frac{3a(a+b)(a+b\coth^2(x))^{3/2}}{3a(a+b)(a+b\coth^2(x))^{3/2}} \\
& \quad \downarrow \text{219} \\
& \frac{3a^2 \operatorname{arctanh}\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{\sqrt{a+b}} - \frac{(3a+2b)(a+4b)\tanh(x)\sqrt{a+b\coth^2(x)}}{a} + \frac{b(7a+4b)\tanh(x)}{a(a+b)\sqrt{a+b\coth^2(x)}} + \\
& \frac{3a(a+b)}{b\tanh(x)} \\
& \frac{3a(a+b)(a+b\coth^2(x))^{3/2}}{3a(a+b)(a+b\coth^2(x))^{3/2}}
\end{aligned}$$

input `Int [Tanh[x]^2/(a + b*Coth[x]^2)^(5/2), x]`

output `(b*Tanh[x])/(3*a*(a + b)*(a + b*Coth[x]^2)^(3/2)) + ((b*(7*a + 4*b)*Tanh[x])/ (a*(a + b)*Sqrt[a + b*Coth[x]^2]) + ((3*a^2*ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]])/Sqrt[a + b] - ((3*a + 2*b)*(a + 4*b)*Sqrt[a + b*Coth[x]^2]*Tanh[x])/a)/(a*(a + b)))/(3*a*(a + b))`

3.47.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

3.47. $\int \frac{\tanh^2(x)}{(a+b\coth^2(x))^{5/2}} dx$

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 291 `Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*((c_) + (d_)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 374 `Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[(-b)*(e*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*e*2*(b*c - a*d)*(p + 1))), x] + Simp[1/(a*2*(b*c - a*d)*(p + 1)) Int[(e*x)^m*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[b*c*(m + 1) + 2*(b*c - a*d)*(p + 1) + d*b*(m + 2*(p + q + 2) + 1)*x^2, x], x] /; FreeQ[{a, b, c, d, e, m, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`

rule 441 `Int[((g_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[(-b*e - a*f)*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*g*2*(b*c - a*d)*(p + 1))), x] + Simp[1/(a*2*(b*c - a*d)*(p + 1)) Int[(g*x)^m*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(b*e - a*f)*(m + 1) + e*2*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(m + 2*(p + q + 2) + 1)*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, q}, x] && LtQ[p, -1]`

rule 445 `Int[((g_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(a*c*g*(m + 1))), x] + Simp[1/(a*c*g^2*(m + 1)) Int[(g*x)^(m + 2)*(a + b*x^2)^p*(c + d*x^2)^q*Simp[a*f*c*(m + 1) - e*(b*c + a*d)*(m + 2 + 1) - e*2*(b*c*p + a*d*q) - b*e*d*(m + 2*(p + q + 2) + 1)*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && LtQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4153 Int[((d_.)*tan[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)]^(n_))^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.47.4 Maple [F]

$$\int \frac{\tanh(x)^2}{(a + b \coth(x)^2)^{5/2}} dx$$

```
input int(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x)
```

```
output int(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x)
```

3.47.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5085 vs. 2(113) = 226.

Time = 1.30 (sec) , antiderivative size = 10729, normalized size of antiderivative = 81.90

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

```
input integrate(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")
```

```
output Too large to include
```


3.47.6 Sympy [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

input `integrate(tanh(x)**2/(a+b*coth(x)**2)**(5/2),x)`

output `Integral(tanh(x)**2/(a + b*coth(x)**2)**(5/2), x)`

3.47.7 Maxima [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{\frac{5}{2}}} dx$$

input `integrate(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")`

output `integrate(tanh(x)^2/(b*coth(x)^2 + a)^(5/2), x)`

3.47.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1133 vs. $2(113) = 226$.

Time = 0.77 (sec) , antiderivative size = 1133, normalized size of antiderivative = 8.65

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

input `integrate(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")`

output

$$\begin{aligned}
& -1/3 * (((9*a^{13}*b^4 + 50*a^{12}*b^5 + 115*a^{11}*b^6 + 140*a^{10}*b^7 + 95*a^9*b^8 + 34*a^8*b^9 + 5*a^7*b^{10}) * e^{(2*x)} / (a^{16}*b^2 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{15}*b^3 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{14}*b^4 * \operatorname{sgn}(e^{(2*x)} - 1) + 20*a^{13}*b^5 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{12}*b^6 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{11}*b^7 * \operatorname{sgn}(e^{(2*x)} - 1) + a^{10}*b^8 * \operatorname{sgn}(e^{(2*x)} - 1)) - 3*(3*a^{13}*b^4 + 6*a^{12}*b^5 - 11*a^{11}*b^6 - 44*a^{10}*b^7 - 51*a^9*b^8 - 26*a^8*b^9 - 5*a^7*b^{10}) / (a^{16}*b^2 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{15}*b^3 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{14}*b^4 * \operatorname{sgn}(e^{(2*x)} - 1) + 20*a^{13}*b^5 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{12}*b^6 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{11}*b^7 * \operatorname{sgn}(e^{(2*x)} - 1) + a^{10}*b^8 * \operatorname{sgn}(e^{(2*x)} - 1))) * e^{(2*x)} - 3*(3*a^{13}*b^4 + 6*a^{12}*b^5 - 11*a^{11}*b^6 - 44*a^{10}*b^7 - 51*a^9*b^8 - 26*a^8*b^9 - 5*a^7*b^{10}) / (a^{16}*b^2 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{15}*b^3 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{14}*b^4 * \operatorname{sgn}(e^{(2*x)} - 1) + 20*a^{13}*b^5 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{12}*b^6 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{11}*b^7 * \operatorname{sgn}(e^{(2*x)} - 1) + a^{10}*b^8 * \operatorname{sgn}(e^{(2*x)} - 1))) * e^{(2*x)} + (9*a^{13}*b^4 + 50*a^{12}*b^5 + 115*a^{11}*b^6 + 140*a^{10}*b^7 + 95*a^9*b^8 + 34*a^8*b^9 + 5*a^7*b^{10}) / (a^{16}*b^2 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{15}*b^3 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{14}*b^4 * \operatorname{sgn}(e^{(2*x)} - 1) + 20*a^{13}*b^5 * \operatorname{sgn}(e^{(2*x)} - 1) + 15*a^{12}*b^6 * \operatorname{sgn}(e^{(2*x)} - 1) + 6*a^{11}*b^7 * \operatorname{sgn}(e^{(2*x)} - 1) + a^{10}*b^8 * \operatorname{sgn}(e^{(2*x)} - 1))) / (a * e^{(4*x)} + b * e^{(4*x)} - 2*a * e^{(2*x)} + 2*b * e^{(2*x)} + a + b)^{(3/2)} - 1/2 * \log(\operatorname{abs}((\operatorname{sqrt}(a + b) * e^{(2*x)} - \operatorname{sqrt}(a * e^{(4*x)} + b * e^{(4*x)} - 2*a * e^{(2*x)} + 2*b * e^{(2*x)} + a + b)) * \operatorname{sqrt}(a + b) - a + b)) / ((a^2 + 2*a*b + b^2) * \operatorname{sqrt}(a + b)))
\end{aligned}$$

3.47.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{5/2}} dx$$

input `int(tanh(x)^2/(a + b*coth(x)^2)^(5/2), x)`

output `int(tanh(x)^2/(a + b*coth(x)^2)^(5/2), x)`

3.48 $\int \frac{1}{\sqrt{1+\coth^2(x)}} dx$

3.48.1	Optimal result	362
3.48.2	Mathematica [A] (verified)	362
3.48.3	Rubi [A] (verified)	363
3.48.4	Maple [B] (verified)	364
3.48.5	Fricas [B] (verification not implemented)	365
3.48.6	Sympy [F]	365
3.48.7	Maxima [F]	366
3.48.8	Giac [B] (verification not implemented)	366
3.48.9	Mupad [B] (verification not implemented)	366

3.48.1 Optimal result

Integrand size = 10, antiderivative size = 25

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{1+\coth^2(x)}}\right)}{\sqrt{2}}$$

output `1/2*arctanh(coth(x)*2^(1/2)/(1+coth(x)^2)^(1/2))*2^(1/2)`

3.48.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{1+\coth^2(x)}}\right)}{\sqrt{2}}$$

input `Integrate[1/Sqrt[1 + Coth[x]^2],x]`

output `ArcTanh[(Sqrt[2]*Coth[x])/Sqrt[1 + Coth[x]^2]]/Sqrt[2]`

3.48.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 4144, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\coth^2(x) + 1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{1 - \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{1}{(1 - \coth^2(x)) \sqrt{\coth^2(x) + 1}} d \coth(x) \\
 & \quad \downarrow \text{291} \\
 & \int \frac{1}{1 - \frac{2 \coth^2(x)}{\coth^2(x)+1}} d \frac{\coth(x)}{\sqrt{\coth^2(x) + 1}} \\
 & \quad \downarrow \text{219} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{\coth^2(x)+1}}\right)}{\sqrt{2}}
 \end{aligned}$$

input `Int[1/Sqrt[1 + Coth[x]^2], x]`

output `ArcTanh[(Sqrt[2]*Coth[x])/Sqrt[1 + Coth[x]^2]]/Sqrt[2]`

3.48.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

3.48.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(20) = 40$.

Time = 0.20 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.48

method	result	size
derivativedivides	$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2-2 \operatorname{coth}(x))\sqrt{2}}{4\sqrt{(1+\operatorname{coth}(x))^2-2 \operatorname{coth}(x)}}\right)}{4} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2 \operatorname{coth}(x))\sqrt{2}}{4\sqrt{(\operatorname{coth}(x)-1)^2+2 \operatorname{coth}(x)}}\right)}{4}$	62
default	$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2-2 \operatorname{coth}(x))\sqrt{2}}{4\sqrt{(1+\operatorname{coth}(x))^2-2 \operatorname{coth}(x)}}\right)}{4} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2 \operatorname{coth}(x))\sqrt{2}}{4\sqrt{(\operatorname{coth}(x)-1)^2+2 \operatorname{coth}(x)}}\right)}{4}$	62

input `int(1/(1+coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4*2^(1/2)*arctanh(1/4*(2-2*coth(x))*2^(1/2)/((1+coth(x))^2-2*coth(x))^(1/2))+1/4*2^(1/2)*arctanh(1/4*(2+2*coth(x))*2^(1/2)/((coth(x)-1)^2+2*coth(x))^(1/2))`

3.48.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 547 vs. $2(20) = 40$.

Time = 0.25 (sec) , antiderivative size = 547, normalized size of antiderivative = 21.88

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \text{Too large to display}$$

```
input integrate(1/(1+coth(x)^2)^(1/2),x, algorithm="fracas")
```

```
output 1/8*sqrt(2)*log(2*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 + 3)*sinh(x)^6 + 3*cosh(x)^6 + 2*(28*cosh(x)^3 + 9*cosh(x))*sinh(x)^5 + 5*(14*cosh(x)^4 + 9*cosh(x)^2 + 1)*sinh(x)^4 + 5*cosh(x)^4 + 4*(14*cosh(x)^5 + 15*cosh(x)^3 + 5*cosh(x))*sinh(x)^3 + (28*cosh(x)^6 + 45*cosh(x)^4 + 30*cosh(x)^2 + 4)*sinh(x)^2 + 4*cosh(x)^2 + 2*(4*cosh(x)^7 + 9*cosh(x)^5 + 10*cosh(x)^3 + 4*cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^6 + 6*sqrt(2)*cosh(x)*sinh(x)^5 + sqrt(2)*sinh(x)^6 + 3*(5*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^4 + 3*sqrt(2)*cosh(x)^4 + 4*(5*sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x))*sinh(x)^3 + (15*sqrt(2)*cosh(x)^4 + 18*sqrt(2)*cosh(x)^2 + 4*sqrt(2))*sinh(x)^2 + 4*sqrt(2)*cosh(x)^2 + 2*(3*sqrt(2)*cosh(x)^5 + 6*sqrt(2)*cosh(x)^3 + 4*sqrt(2)*cosh(x))*sinh(x) + 4*sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4)/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + 1/8*sqrt(2)*log(-2*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 - sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2))
```

3.48.6 Sympy [F]

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \int \frac{1}{\sqrt{\coth^2(x) + 1}} dx$$

```
input integrate(1/(1+coth(x)**2)**(1/2),x)
```

```
output Integral(1/sqrt(coth(x)**2 + 1), x)
```

3.48.7 Maxima [F]

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \int \frac{1}{\sqrt{\coth(x)^2 + 1}} dx$$

input `integrate(1/(1+coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(coth(x)^2 + 1), x)`

3.48.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(20) = 40$.

Time = 0.26 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.76

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \frac{\sqrt{2} \left(\log \left(\sqrt{e^{(4x)} + 1} - e^{(2x)} + 1 \right) - \log \left(\sqrt{e^{(4x)} + 1} - e^{(2x)} \right) - \log \left(-\sqrt{e^{(4x)} + 1} + e^{(2x)} + 1 \right) \right)}{4 \operatorname{sgn}(e^{(2x)} - 1)}$$

input `integrate(1/(1+coth(x)^2)^(1/2),x, algorithm="giac")`

output `1/4*sqrt(2)*(log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) - log(sqrt(e^(4*x) + 1) - e^(2*x)) - log(-sqrt(e^(4*x) + 1) + e^(2*x) + 1))/sgn(e^(2*x) - 1)`

3.48.9 Mupad [B] (verification not implemented)

Time = 2.10 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.52

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \frac{\sqrt{2} \left(\ln \left(\coth(x) + \sqrt{2} \sqrt{\coth(x)^2 + 1} + 1 \right) - \ln(\coth(x) - 1) \right)}{4} + \frac{\sqrt{2} \left(\ln(\coth(x) + 1) - \ln \left(\sqrt{2} \sqrt{\coth(x)^2 + 1} - \coth(x) + 1 \right) \right)}{4}$$

input `int(1/(coth(x)^2 + 1)^(1/2),x)`

output $(2^{1/2} * (\log(\coth(x) + 2^{1/2} * (\coth(x)^2 + 1)^{1/2} + 1) - \log(\coth(x) - 1))) / 4 + (2^{1/2} * (\log(\coth(x) + 1) - \log(2^{1/2} * (\coth(x)^2 + 1)^{1/2} - \coth(x) + 1))) / 4$

$$3.49 \quad \int \frac{1}{\sqrt{-1-\coth^2(x)}} dx$$

3.49.1	Optimal result	368
3.49.2	Mathematica [A] (warning: unable to verify)	368
3.49.3	Rubi [A] (verified)	369
3.49.4	Maple [B] (verified)	370
3.49.5	Fricas [C] (verification not implemented)	371
3.49.6	Sympy [F]	371
3.49.7	Maxima [F]	372
3.49.8	Giac [C] (verification not implemented)	372
3.49.9	Mupad [B] (verification not implemented)	372

3.49.1 Optimal result

Integrand size = 12, antiderivative size = 27

$$\int \frac{1}{\sqrt{-1-\coth^2(x)}} dx = \frac{\arctan\left(\frac{\sqrt{2}\coth(x)}{\sqrt{-1-\coth^2(x)}}\right)}{\sqrt{2}}$$

output `1/2*arctan(coth(x)*2^(1/2)/(-1-coth(x)^2)^(1/2))*2^(1/2)`

3.49.2 Mathematica [A] (warning: unable to verify)

Time = 0.08 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.81

$$\int \frac{1}{\sqrt{-1-\coth^2(x)}} dx = \frac{\operatorname{arcsinh}\left(\frac{\sqrt{2}\coth(x)}{\sqrt{1-\coth^2(x)}}\right) \sqrt{1+\coth^2(x)}}{\sqrt{2}\sqrt{-1-\coth^2(x)}}$$

input `Integrate[1/Sqrt[-1 - Coth[x]^2], x]`

output `(ArcSinh[(Sqrt[2]*Coth[x])/Sqrt[1 - Coth[x]^2]]*Sqrt[1 + Coth[x]^2])/(Sqrt[2]*Sqrt[-1 - Coth[x]^2])`

3.49. $\int \frac{1}{\sqrt{-1-\coth^2(x)}} dx$

3.49.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4144, 291, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{-\coth^2(x) - 1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{-1 + \tan\left(\frac{\pi}{2} + ix\right)^2}} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{1}{\sqrt{-\coth^2(x) - 1} (1 - \coth^2(x))} d\coth(x) \\
 & \quad \downarrow \text{291} \\
 & \int \frac{1}{\frac{2\coth^2(x)}{-\coth^2(x)-1} + 1} d\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}} \\
 & \quad \downarrow \text{216} \\
 & \frac{\arctan\left(\frac{\sqrt{2}\coth(x)}{\sqrt{-\coth^2(x)-1}}\right)}{\sqrt{2}}
 \end{aligned}$$

input `Int[1/Sqrt[-1 - Coth[x]^2], x]`

output `ArcTan[(Sqrt[2]*Coth[x])/Sqrt[-1 - Coth[x]^2]]/Sqrt[2]`

3.49.3.1 Defintions of rubi rules used

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 291 Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4144 Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

3.49.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(22) = 44.

Time = 0.20 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.44

method	result	size
derivativedivides	$-\frac{\sqrt{2} \arctan\left(\frac{(-2-2 \coth(x))\sqrt{2}}{4\sqrt{-(\coth(x)-1)^2-2 \coth(x)}}\right)}{4} + \frac{\sqrt{2} \arctan\left(\frac{(-2+2 \coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2 \coth(x)}}\right)}{4}$	66
default	$-\frac{\sqrt{2} \arctan\left(\frac{(-2-2 \coth(x))\sqrt{2}}{4\sqrt{-(\coth(x)-1)^2-2 \coth(x)}}\right)}{4} + \frac{\sqrt{2} \arctan\left(\frac{(-2+2 \coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2 \coth(x)}}\right)}{4}$	66

```
input int(1/(-1-coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/4*2^(1/2)*arctan(1/4*(-2-2*coth(x))*2^(1/2)/(-(coth(x)-1)^2-2*coth(x))^(1/2))+1/4*2^(1/2)*arctan(1/4*(-2+2*coth(x))*2^(1/2)/(-(1+coth(x))^2+2*coth(x))^(1/2))
```

3.49. $\int \frac{1}{\sqrt{-1-\coth^2(x)}} dx$

3.49.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 175, normalized size of antiderivative = 6.48

$$\begin{aligned} & \int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx \\ &= \frac{1}{8} i \sqrt{2} \log \left(\frac{1}{2} \left(i \sqrt{2} \sqrt{-2 e^{4x} - 2} + 2 e^{2x} - 2 \right) e^{(-2x)} \right) \\ & \quad - \frac{1}{8} i \sqrt{2} \log \left(\frac{1}{2} \left(-i \sqrt{2} \sqrt{-2 e^{4x} - 2} + 2 e^{2x} - 2 \right) e^{(-2x)} \right) \\ & \quad - \frac{1}{8} i \sqrt{2} \log \left(\left(\sqrt{-2 e^{4x} - 2} (e^{2x} + 2) + i \sqrt{2} e^{4x} + i \sqrt{2} e^{2x} + 2i \sqrt{2} \right) e^{(-4x)} \right) \\ & \quad + \frac{1}{8} i \sqrt{2} \log \left(\left(\sqrt{-2 e^{4x} - 2} (e^{2x} + 2) - i \sqrt{2} e^{4x} - i \sqrt{2} e^{2x} - 2i \sqrt{2} \right) e^{(-4x)} \right) \end{aligned}$$

```
input integrate(1/(-1-coth(x)^2)^(1/2),x, algorithm="fricas")
```

```
output 1/8*I*sqrt(2)*log(1/2*(I*sqrt(2)*sqrt(-2*e^(4*x) - 2) + 2*e^(2*x) - 2)*e^(-2*x)) - 1/8*I*sqrt(2)*log(1/2*(-I*sqrt(2)*sqrt(-2*e^(4*x) - 2) + 2*e^(2*x) - 2)*e^(-2*x)) - 1/8*I*sqrt(2)*log((sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) + I*sqrt(2)*e^(4*x) + I*sqrt(2)*e^(2*x) + 2*I*sqrt(2))*e^(-4*x)) + 1/8*I*sqrt(2)*log((sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) - I*sqrt(2)*e^(4*x) - I*sqrt(2)*e^(2*x) - 2*I*sqrt(2))*e^(-4*x))
```

3.49.6 Sympy [F]

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \int \frac{1}{\sqrt{-\coth^2(x) - 1}} dx$$

```
input integrate(1/(-1-coth(x)**2)**(1/2),x)
```

```
output Integral(1/sqrt(-coth(x)**2 - 1), x)
```

3.49.7 Maxima [F]

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \int \frac{1}{\sqrt{-\coth(x)^2 - 1}} dx$$

input `integrate(1/(-1-coth(x)^2)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(-coth(x)^2 - 1), x)`

3.49.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.70

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \frac{\sqrt{2} \left(-i \log \left(\sqrt{e^{4x} + 1} - e^{2x} + 1 \right) + i \log \left(\sqrt{e^{4x} + 1} - e^{2x} \right) + i \log \left(-\sqrt{e^{4x} + 1} + e^{2x} + 1 \right) \right)}{4 \operatorname{sgn}(-e^{2x} + 1)}$$

input `integrate(1/(-1-coth(x)^2)^(1/2),x, algorithm="giac")`

output `-1/4*sqrt(2)*(-I*log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) + I*log(sqrt(e^(4*x) + 1) - e^(2*x)) + I*log(-sqrt(e^(4*x) + 1) + e^(2*x) + 1))/sgn(-e^(2*x) + 1)`

3.49.9 Mupad [B] (verification not implemented)

Time = 2.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \frac{\sqrt{2} \operatorname{atan} \left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth(x)^2 - 1}} \right)}{2}$$

input `int(1/(-coth(x)^2 - 1)^(1/2),x)`

output `(2^(1/2)*atan((2^(1/2)*coth(x))/(-coth(x)^2 - 1)^(1/2)))/2`

3.50 $\int \frac{1}{1+\coth^3(x)} dx$

3.50.1	Optimal result	373
3.50.2	Mathematica [A] (verified)	373
3.50.3	Rubi [A] (verified)	374
3.50.4	Maple [A] (verified)	375
3.50.5	Fricas [B] (verification not implemented)	376
3.50.6	Sympy [B] (verification not implemented)	376
3.50.7	Maxima [B] (verification not implemented)	377
3.50.8	Giac [A] (verification not implemented)	377
3.50.9	Mupad [B] (verification not implemented)	377

3.50.1 Optimal result

Integrand size = 8, antiderivative size = 38

$$\int \frac{1}{1+\coth^3(x)} dx = \frac{x}{2} - \frac{2 \arctan\left(\frac{1-2\coth(x)}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{1}{6(1+\coth(x))}$$

output `1/2*x-1/6/(1+coth(x))-2/9*arctan(1/3*(1-2*coth(x))*3^(1/2))*3^(1/2)`

3.50.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05

$$\int \frac{1}{1+\coth^3(x)} dx = \frac{2 \arctan\left(\frac{1-2\tanh(x)}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{1}{2} \operatorname{arctanh}(\tanh(x)) + \frac{1}{6(1+\tanh(x))}$$

input `Integrate[(1 + Coth[x]^3)^(-1), x]`

output `(2*ArcTan[(1 - 2*Tanh[x])/Sqrt[3]])/(3*Sqrt[3]) + ArcTanh[Tanh[x]]/2 + 1/(6*(1 + Tanh[x]))`

3.50.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4144, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\coth^3(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{1 + i \tan\left(\frac{\pi}{2} + ix\right)^3} dx \\
 & \quad \downarrow \text{4144} \\
 & \int \frac{1}{(1 - \coth^2(x)) (\coth^3(x) + 1)} d \coth(x) \\
 & \quad \downarrow \text{7276} \\
 & \int \left(\frac{1}{3 (\coth^2(x) - \coth(x) + 1)} - \frac{1}{2 (\coth^2(x) - 1)} + \frac{1}{6 (\coth(x) + 1)^2} \right) d \coth(x) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{2 \arctan\left(\frac{1-2\coth(x)}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{1}{2} \operatorname{arctanh}(\coth(x)) - \frac{1}{6(\coth(x) + 1)}
 \end{aligned}$$

input `Int[(1 + Coth[x]^3)^(-1),x]`

output `(-2*ArcTan[(1 - 2*Coth[x])/Sqrt[3]])/(3*Sqrt[3]) + ArcTanh[Coth[x]]/2 - 1/(6*(1 + Coth[x]))`

3.50.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4144 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

3.50.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$-\frac{1}{6(1+\coth(x))} + \frac{\ln(1+\coth(x))}{4} - \frac{\ln(\coth(x)-1)}{4} + \frac{2\sqrt{3} \arctan\left(\frac{(2\coth(x)-1)\sqrt{3}}{3}\right)}{9}$	41
default	$-\frac{1}{6(1+\coth(x))} + \frac{\ln(1+\coth(x))}{4} - \frac{\ln(\coth(x)-1)}{4} + \frac{2\sqrt{3} \arctan\left(\frac{(2\coth(x)-1)\sqrt{3}}{3}\right)}{9}$	41
risch	$\frac{x}{2} + \frac{e^{-2x}}{12} + \frac{i\sqrt{3} \ln(e^{2x}-i\sqrt{3})}{9} - \frac{i\sqrt{3} \ln(e^{2x}+i\sqrt{3})}{9}$	47

input `int(1/(1+coth(x)^3),x,method=_RETURNVERBOSE)`

output `-1/6/(1+coth(x))+1/4*ln(1+coth(x))-1/4*ln(coth(x)-1)+2/9*3^(1/2)*arctan(1/3*(2*coth(x)-1)*3^(1/2))`

3.50.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(29) = 58$.

Time = 0.27 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.50

$$\int \frac{1}{1 + \coth^3(x)} dx = \frac{18x \cosh(x)^2 + 36x \cosh(x) \sinh(x) + 18x \sinh(x)^2 + 8(\sqrt{3} \cosh(x)^2 + 2\sqrt{3} \cosh(x) \sinh(x) + \sqrt{3} \sinh(x)^2)}{36(\cosh(x)^2 + 2\cosh(x) \sinh(x) + \sinh(x)^2)}$$

input `integrate(1/(1+coth(x)^3),x, algorithm="fricas")`

output `1/36*(18*x*cosh(x)^2 + 36*x*cosh(x)*sinh(x) + 18*x*sinh(x)^2 + 8*(sqrt(3)*cosh(x)^2 + 2*sqrt(3)*cosh(x)*sinh(x) + sqrt(3)*sinh(x)^2)*arctan(-1/3*(sqrt(3)*cosh(x) + sqrt(3)*sinh(x))/(cosh(x) - sinh(x))) + 3/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)`

3.50.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. $2(36) = 72$.

Time = 0.50 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.68

$$\int \frac{1}{1 + \coth^3(x)} dx = \frac{9x \tanh(x)}{18 \tanh(x) + 18} + \frac{9x}{18 \tanh(x) + 18} - \frac{4\sqrt{3} \tanh(x) \operatorname{atan}\left(\frac{2\sqrt{3} \tanh(x)}{3} - \frac{\sqrt{3}}{3}\right)}{18 \tanh(x) + 18} - \frac{4\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3} \tanh(x)}{3} - \frac{\sqrt{3}}{3}\right)}{18 \tanh(x) + 18} + \frac{3}{18 \tanh(x) + 18}$$

input `integrate(1/(1+coth(x)**3),x)`

output `9*x*tanh(x)/(18*tanh(x) + 18) + 9*x/(18*tanh(x) + 18) - 4*sqrt(3)*tanh(x)*atan(2*sqrt(3)*tanh(x)/3 - sqrt(3)/3)/(18*tanh(x) + 18) - 4*sqrt(3)*atan(2*sqrt(3)*tanh(x)/3 - sqrt(3)/3)/(18*tanh(x) + 18) + 3/(18*tanh(x) + 18)`

3.50.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(29) = 58$.

Time = 0.29 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.92

$$\int \frac{1}{1 + \coth^3(x)} dx = -\frac{2}{9} \sqrt{3} \arctan \left(\frac{1}{6} \cdot 3^{\frac{3}{4}} \sqrt{2} \left(2 \sqrt{3} e^{-x} + 3^{\frac{1}{4}} \sqrt{2} \right) \right) \\ + \frac{2}{9} \sqrt{3} \arctan \left(\frac{1}{6} \cdot 3^{\frac{3}{4}} \sqrt{2} \left(2 \sqrt{3} e^{-x} - 3^{\frac{1}{4}} \sqrt{2} \right) \right) + \frac{1}{2} x + \frac{1}{12} e^{(-2x)}$$

input `integrate(1/(1+coth(x)^3),x, algorithm="maxima")`

output `-2/9*sqrt(3)*arctan(1/6*3^(3/4)*sqrt(2)*(2*sqrt(3)*e^(-x) + 3^(1/4)*sqrt(2))) + 2/9*sqrt(3)*arctan(1/6*3^(3/4)*sqrt(2)*(2*sqrt(3)*e^(-x) - 3^(1/4)*sqrt(2))) + 1/2*x + 1/12*e^(-2*x)`

3.50.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.66

$$\int \frac{1}{1 + \coth^3(x)} dx = -\frac{2}{9} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} e^{(2x)} \right) + \frac{1}{2} x + \frac{1}{12} e^{(-2x)}$$

input `integrate(1/(1+coth(x)^3),x, algorithm="giac")`

output `-2/9*sqrt(3)*arctan(1/3*sqrt(3)*e^(2*x)) + 1/2*x + 1/12*e^(-2*x)`

3.50.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \coth^3(x)} dx = \frac{\frac{x}{2} + \frac{\coth(x)}{6} + \frac{x \coth(x)}{2}}{\coth(x) + 1} + \frac{2 \sqrt{3} \operatorname{atan} \left(\frac{\sqrt{3} (2 \coth(x) - 1)}{3} \right)}{9}$$

input `int(1/(coth(x)^3 + 1),x)`

output `(x/2 + coth(x)/6 + (x*coth(x))/2)/(coth(x) + 1) + (2*3^(1/2)*atan((3^(1/2)*(2*coth(x) - 1))/3))/9`

3.51 $\int \coth(x) \sqrt{a + b \coth^4(x)} dx$

3.51.1	Optimal result	378
3.51.2	Mathematica [A] (verified)	379
3.51.3	Rubi [A] (verified)	379
3.51.4	Maple [A] (verified)	382
3.51.5	Fricas [B] (verification not implemented)	383
3.51.6	Sympy [F]	383
3.51.7	Maxima [F]	384
3.51.8	Giac [F]	384
3.51.9	Mupad [F(-1)]	384

3.51.1 Optimal result

Integrand size = 15, antiderivative size = 89

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = -\frac{1}{2} \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) + \frac{1}{2} \sqrt{a + b} \operatorname{arctanh} \left(\frac{a + b \coth^2(x)}{\sqrt{a + b} \sqrt{a + b \coth^4(x)}} \right) - \frac{1}{2} \sqrt{a + b \coth^4(x)}$$

output

```
-1/2*arctanh(coth(x)^2*b^(1/2)/(a+b*coth(x)^4)^(1/2))*b^(1/2)+1/2*arctanh(
(a+b*coth(x)^2)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))*(a+b)^(1/2)-1/2*(a+b*co
th(x)^4)^(1/2)
```

3.51.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.97

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \frac{1}{2} \left(-\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) \right. \\ \left. + \sqrt{a + b} \operatorname{arctanh} \left(\frac{a + b \coth^2(x)}{\sqrt{a + b} \sqrt{a + b \coth^4(x)}} \right) \right. \\ \left. - \sqrt{a + b \coth^4(x)} \right)$$

input `Integrate[Coth[x]*Sqrt[a + b*Coth[x]^4],x]`

output `(-(Sqrt[b]*ArcTanh[(Sqrt[b]*Coth[x]^2)/Sqrt[a + b*Coth[x]^4]]) + Sqrt[a + b]*ArcTanh[(a + b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4]]) - Sqrt[a + b*Coth[x]^4])/2`

3.51.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.01, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {3042, 26, 4153, 26, 1577, 493, 25, 719, 224, 219, 488, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx \\ \downarrow \text{3042} \\ \int -i \tan \left(\frac{\pi}{2} + ix \right) \sqrt{a + b \tan \left(\frac{\pi}{2} + ix \right)^4} dx \\ \downarrow \text{26} \\ -i \int \tan \left(ix + \frac{\pi}{2} \right) \sqrt{b \tan \left(ix + \frac{\pi}{2} \right)^4 + a} dx \\ \downarrow \text{4153}$$

$$\begin{aligned}
& -i \int \frac{i \coth(x) \sqrt{b \coth^4(x) + a}}{1 - \coth^2(x)} d \coth(x) \\
& \quad \downarrow \text{26} \\
& \int \frac{\coth(x) \sqrt{a + b \coth^4(x)}}{1 - \coth^2(x)} d \coth(x) \\
& \quad \downarrow \text{1577} \\
& \frac{1}{2} \int \frac{\sqrt{b \coth^4(x) + a}}{1 - \coth^2(x)} d \coth^2(x) \\
& \quad \downarrow \text{493} \\
& \frac{1}{2} \left(- \int \frac{b \coth^2(x) + a}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) - \sqrt{a + b \coth^4(x)} \right) \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left(\int \frac{b \coth^2(x) + a}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) - \sqrt{a + b \coth^4(x)} \right) \\
& \quad \downarrow \text{719} \\
& \frac{1}{2} \left(-b \int \frac{1}{\sqrt{b \coth^4(x) + a}} d \coth^2(x) + (a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) - \sqrt{a + b \coth^4(x)} \right) \\
& \quad \downarrow \text{224} \\
& \frac{1}{2} \left(-b \int \frac{1}{1 - b \coth^4(x)} d \frac{\coth^2(x)}{\sqrt{b \coth^4(x) + a}} + (a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) - \sqrt{a + b \coth^4(x)} \right) \\
& \quad \downarrow \text{219} \\
& \frac{1}{2} \left((a + b) \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) - \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) - \sqrt{a + b \coth^4(x)} \right) \\
& \quad \downarrow \text{488}
\end{aligned}$$

$$\frac{1}{2} \left(-(a+b) \int \frac{1}{-\coth^4(x) + a + b} dx \frac{-b \coth^2(x) - a}{\sqrt{b \coth^4(x) + a}} - \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) - \sqrt{a + b \coth^4(x)} \right)$$

↓ 219

$$\frac{1}{2} \left(-\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) - \sqrt{a + b} \operatorname{arctanh} \left(\frac{-a - b \coth^2(x)}{\sqrt{a + b} \sqrt{a + b \coth^4(x)}} \right) - \sqrt{a + b \coth^4(x)} \right)$$

input `Int[Coth[x]*Sqrt[a + b*Coth[x]^4], x]`

output `(-(Sqrt[b]*ArcTanh[(Sqrt[b]*Coth[x]^2)/Sqrt[a + b*Coth[x]^4]]) - Sqrt[a + b]*ArcTanh[(-a - b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4]]) - Sqrt[a + b*Coth[x]^4])/2`

3.51.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 488 `Int[1/(((c_) + (d_.)*(x_))*Sqrt[(a_) + (b_.)*(x_)^2]), x_Symbol] := -Subst[Int[1/(b*c^2 + a*d^2 - x^2), x], x, (a*d - b*c*x)/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x]`

```
rule 493 Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
(c + d*x)^(n + 1)*((a + b*x^2)^p/(d*(n + 2*p + 1))), x] + Simp[2*(p/(d*(n +
2*p + 1))) Int[(c + d*x)^n*(a + b*x^2)^(p - 1)*(a*d - b*c*x), x], x] /;
FreeQ[{a, b, c, d, n}, x] && GtQ[p, 0] && NeQ[n + 2*p + 1, 0] && (!Rationa
lQ[n] || LtQ[n, 1]) && !ILtQ[n + 2*p, 0] && IntQuadraticQ[a, 0, b, c, d, n
, p, x]
```

```
rule 719 Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^(p
_), x_Symbol] := Simp[g/e Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] +
Simp[(e*f - d*g)/e Int[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c,
d, e, f, g, m, p}, x] && !IGtQ[m, 0]
```

```
rule 1577 Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Simp[1/2 Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; Free
Q[{a, c, d, e, p, q}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4153 Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))
```

3.51.4 Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.30

method	result
derivativedivides	$-\frac{\sqrt{a+b \coth(x)^4}}{2} - \frac{\sqrt{b} \ln\left(2\sqrt{b} \coth(x)^2 + 2\sqrt{a+b \coth(x)^4}\right)}{2} + \frac{b \operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}} + \frac{a \operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}}$
default	$-\frac{\sqrt{a+b \coth(x)^4}}{2} - \frac{\sqrt{b} \ln\left(2\sqrt{b} \coth(x)^2 + 2\sqrt{a+b \coth(x)^4}\right)}{2} + \frac{b \operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}} + \frac{a \operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}}$

3.51. $\int \coth(x) \sqrt{a + b \coth^4(x)} dx$

input `int(coth(x)*(a+b*coth(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*(a+b*coth(x)^4)^(1/2)-1/2*b^(1/2)*ln(2*b^(1/2)*coth(x)^2+2*(a+b*coth(x)^4)^(1/2))+1/2*b/(a+b)^(1/2)*arctanh(1/2*(2*b*coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))+1/2*a/(a+b)^(1/2)*arctanh(1/2*(2*b*coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))`

3.51.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1053 vs. 2(69) = 138.

Time = 0.40 (sec) , antiderivative size = 5172, normalized size of antiderivative = 58.11

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \text{Too large to display}$$

input `integrate(coth(x)*(a+b*coth(x)^4)^(1/2),x, algorithm="fricas")`

output `Too large to include`

3.51.6 Sympy [F]

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \sqrt{a + b \coth^4(x)} \coth(x) dx$$

input `integrate(coth(x)*(a+b*coth(x)**4)**(1/2),x)`

output `Integral(sqrt(a + b*coth(x)**4)*coth(x), x)`

3.51.7 Maxima [F]

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \sqrt{b \coth(x)^4 + a} \coth(x) dx$$

input `integrate(coth(x)*(a+b*coth(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*coth(x)^4 + a)*coth(x), x)`

3.51.8 Giac [F]

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \sqrt{b \coth(x)^4 + a} \coth(x) dx$$

input `integrate(coth(x)*(a+b*coth(x)^4)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*coth(x)^4 + a)*coth(x), x)`

3.51.9 Mupad [F(-1)]

Timed out.

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \coth(x) \sqrt{b \coth(x)^4 + a} dx$$

input `int(coth(x)*(a + b*coth(x)^4)^(1/2), x)`

output `int(coth(x)*(a + b*coth(x)^4)^(1/2), x)`

3.52 $\int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx$

3.52.1	Optimal result	385
3.52.2	Mathematica [A] (verified)	385
3.52.3	Rubi [A] (verified)	386
3.52.4	Maple [A] (verified)	388
3.52.5	Fricas [B] (verification not implemented)	388
3.52.6	Sympy [F]	389
3.52.7	Maxima [F]	390
3.52.8	Giac [F]	390
3.52.9	Mupad [F(-1)]	390

3.52.1 Optimal result

Integrand size = 15, antiderivative size = 40

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{a+b \coth^2(x)}{\sqrt{a+b} \sqrt{a+b \coth^4(x)}}\right)}{2\sqrt{a+b}}$$

output `1/2*arctanh((a+b*coth(x)^2)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))/(a+b)^(1/2)`

3.52.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{a+b \coth^2(x)}{\sqrt{a+b} \sqrt{a+b \coth^4(x)}}\right)}{2\sqrt{a+b}}$$

input `Integrate[Coth[x]/Sqrt[a + b*Coth[x]^4],x]`

output `ArcTanh[(a + b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4])]/(2*Sqrt[a + b])`

3.52.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 26, 4153, 26, 1577, 488, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan\left(\frac{\pi}{2} + ix\right)}{\sqrt{a + b \tan\left(\frac{\pi}{2} + ix\right)^4}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)}{\sqrt{b \tan\left(ix + \frac{\pi}{2}\right)^4 + a}} dx \\
 & \quad \downarrow \text{4153} \\
 & -i \int \frac{i \coth(x)}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x)}{(1 - \coth^2(x)) \sqrt{a + b \coth^4(x)}} d \coth(x) \\
 & \quad \downarrow \text{1577} \\
 & \frac{1}{2} \int \frac{1}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) \\
 & \quad \downarrow \text{488} \\
 & -\frac{1}{2} \int \frac{1}{-\coth^4(x) + a + b} d \frac{-b \coth^2(x) - a}{\sqrt{b \coth^4(x) + a}} \\
 & \quad \downarrow \text{219} \\
 & -\frac{\operatorname{arctanh}\left(\frac{-a - b \coth^2(x)}{\sqrt{a+b} \sqrt{a + b \coth^4(x)}}\right)}{2\sqrt{a+b}}
 \end{aligned}$$

input `Int[Coth[x]/Sqrt[a + b*Coth[x]^4], x]`

output `-1/2*ArcTanh[(-a - b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4])]/Sqrt[a + b]`

3.52.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 488 `Int[1/(((c_) + (d_)*(x_))*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := -Subst[Int[1/(b*c^2 + a*d^2 - x^2), x], x, (a*d - b*c*x)/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x]`

rule 1577 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4153 `Int[((d_)*tan[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)]^(n_)))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

3.52.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}}$	37
default	$\frac{\operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}}$	37

input `int(coth(x)/(a+b*coth(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2/(a+b)^(1/2)*arctanh(1/2*(2*b*coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))`

3.52.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 626 vs. $2(32) = 64$.

Time = 0.41 (sec) , antiderivative size = 1290, normalized size of antiderivative = 32.25

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \text{Too large to display}$$

input `integrate(coth(x)/(a+b*coth(x)^4)^(1/2),x, algorithm="fricas")`

```
output [1/4*log(((a^2 + 2*a*b + b^2)*cosh(x)^8 + 8*(a^2 + 2*a*b + b^2)*cosh(x)*sinh(x)^7 + (a^2 + 2*a*b + b^2)*sinh(x)^8 - 4*(a^2 - b^2)*cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)^6 + 8*(7*(a^2 + 2*a*b + b^2)*cosh(x)^3 - 3*(a^2 - b^2)*cosh(x))*sinh(x)^5 + 2*(3*a^2 + 2*a*b + 3*b^2)*cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2)*cosh(x)^4 - 30*(a^2 - b^2)*cosh(x)^2 + 3*a^2 + 2*a*b + 3*b^2)*sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2)*cosh(x)^5 - 10*(a^2 - b^2)*cosh(x)^3 + (3*a^2 + 2*a*b + 3*b^2)*cosh(x))*sinh(x)^3 - 4*(a^2 - b^2)*cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^6 - 15*(a^2 - b^2)*cosh(x)^4 + 3*(3*a^2 + 2*a*b + 3*b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)^2 + sqrt(2)*((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^4 + (a + b)*sinh(x)^4 - 4*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - 2*a + 2*b)*sinh(x)^2 + 3*a + 3*b)/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sinh(x)^3 + sinh(x)^4)) + a^2 + 2*a*b + b^2 + 8*((a^2 + 2*a*b + b^2)*cosh(x)^7 - 3*(a^2 - b^2)*cosh(x)^5 + (3*a^2 + 2*a*b + 3*b^2)*cosh(x)^3 - (a^2 - b^2)*cosh(x))*sinh(x))/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4))/sqrt(a + b), -1/2*sqrt(-a - b)*arctan(sqrt(2)*((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*co...
```

3.52.6 Sympy [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx$$

```
input integrate(coth(x)/(a+b*coth(x)**4)**(1/2),x)
```

```
output Integral(coth(x)/sqrt(a + b*coth(x)**4), x)
```

3.52.7 Maxima [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth(x)^4 + a}} dx$$

input `integrate(coth(x)/(a+b*coth(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(coth(x)/sqrt(b*coth(x)^4 + a), x)`

3.52.8 Giac [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth(x)^4 + a}} dx$$

input `integrate(coth(x)/(a+b*coth(x)^4)^(1/2),x, algorithm="giac")`

output `integrate(coth(x)/sqrt(b*coth(x)^4 + a), x)`

3.52.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth(x)^4 + a}} dx$$

input `int(coth(x)/(a + b*coth(x)^4)^(1/2),x)`

output `int(coth(x)/(a + b*coth(x)^4)^(1/2), x)`

3.53 $\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx$

3.53.1	Optimal result	391
3.53.2	Mathematica [A] (verified)	391
3.53.3	Rubi [A] (verified)	392
3.53.4	Maple [C] (verified)	394
3.53.5	Fricas [B] (verification not implemented)	395
3.53.6	Sympy [F]	396
3.53.7	Maxima [F]	397
3.53.8	Giac [F]	397
3.53.9	Mupad [F(-1)]	397

3.53.1 Optimal result

Integrand size = 15, antiderivative size = 74

$$\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{a+b \coth^2(x)}{\sqrt{a+b}\sqrt{a+b \coth^4(x)}}\right)}{2(a+b)^{3/2}} - \frac{a-b \coth^2(x)}{2a(a+b)\sqrt{a+b \coth^4(x)}}$$

output `1/2*arctanh((a+b*coth(x)^2)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))/(a+b)^(3/2)
+1/2*(-a+b*coth(x)^2)/a/(a+b)/(a+b*coth(x)^4)^(1/2)`

3.53.2 Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.99

$$\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx = \frac{1}{2} \left(\frac{\operatorname{arctanh}\left(\frac{a+b \coth^2(x)}{\sqrt{a+b}\sqrt{a+b \coth^4(x)}}\right)}{(a+b)^{3/2}} - \frac{a-b \coth^2(x)}{a(a+b)\sqrt{a+b \coth^4(x)}} \right)$$

input `Integrate[Coth[x]/(a + b*Coth[x]^4)^(3/2),x]`

output `(ArcTanh[(a + b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4]])/(a + b)^(3/2) - (a - b*Coth[x]^2)/(a*(a + b)*Sqrt[a + b*Coth[x]^4]))/2`

3.53. $\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx$

3.53.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 26, 4153, 26, 1577, 496, 25, 27, 488, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan\left(\frac{\pi}{2} + ix\right)}{\left(a + b \tan\left(\frac{\pi}{2} + ix\right)^4\right)^{3/2}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan\left(ix + \frac{\pi}{2}\right)}{\left(b \tan\left(ix + \frac{\pi}{2}\right)^4 + a\right)^{3/2}} dx \\
 & \quad \downarrow \text{4153} \\
 & -i \int \frac{i \coth(x)}{(1 - \coth^2(x)) (b \coth^4(x) + a)^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x)}{(1 - \coth^2(x)) (a + b \coth^4(x))^{3/2}} d \coth(x) \\
 & \quad \downarrow \text{1577} \\
 & \frac{1}{2} \int \frac{1}{(1 - \coth^2(x)) (b \coth^4(x) + a)^{3/2}} d \coth^2(x) \\
 & \quad \downarrow \text{496} \\
 & \frac{1}{2} \left(\int -\frac{a}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) - \frac{a - b \coth^2(x)}{a(a + b) \sqrt{a + b \coth^4(x)}} \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \left(\int \frac{a}{(1 - \coth^2(x)) \sqrt{b \coth^4(x) + a}} d \coth^2(x) - \frac{a - b \coth^2(x)}{a(a + b) \sqrt{a + b \coth^4(x)}} \right)
 \end{aligned}$$

3.53. $\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{1}{2} \left(\frac{\int \frac{1}{(1-\coth^2(x))\sqrt{b\coth^4(x)+a}} d\coth^2(x)}{a+b} - \frac{a-b\coth^2(x)}{a(a+b)\sqrt{a+b\coth^4(x)}} \right) \\
 & \downarrow 488 \\
 & \frac{1}{2} \left(-\frac{\int \frac{1}{-\coth^4(x)+a+b} d\frac{-b\coth^2(x)-a}{\sqrt{b\coth^4(x)+a}}}{a+b} - \frac{a-b\coth^2(x)}{a(a+b)\sqrt{a+b\coth^4(x)}} \right) \\
 & \downarrow 219 \\
 & \frac{1}{2} \left(-\frac{\operatorname{arctanh}\left(\frac{-a-b\coth^2(x)}{\sqrt{a+b}\sqrt{a+b\coth^4(x)}}\right)}{(a+b)^{3/2}} - \frac{a-b\coth^2(x)}{a(a+b)\sqrt{a+b\coth^4(x)}} \right)
 \end{aligned}$$

input `Int[Coth[x]/(a + b*Coth[x]^4)^(3/2), x]`

output `(-(ArcTanh[(-a - b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4]])/(a + b)^(3/2)) - (a - b*Coth[x]^2)/(a*(a + b)*Sqrt[a + b*Coth[x]^4]))/2`

3.53.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 488 `Int[1/(((c_) + (d_)*(x_))*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := -Subst[
Int[1/(b*c^2 + a*d^2 - x^2), x], x, (a*d - b*c*x)/Sqrt[a + b*x^2]] /; FreeQ
[{a, b, c, d}, x]`

rule 496 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
(-(a*d + b*c*x)*(c + d*x)^(n + 1)*((a + b*x^2)^(p + 1)/(2*a*(p + 1)*(b*c^2
+ a*d^2))), x] + Simp[1/(2*a*(p + 1)*(b*c^2 + a*d^2)) Int[(c + d*x)^n*(a
+ b*x^2)^(p + 1)*Simp[b*c^2*(2*p + 3) + a*d^2*(n + 2*p + 3) + b*c*d*(n + 2
*p + 4)*x, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[p, -1] && IntQuad
raticQ[a, 0, b, c, d, n, p, x]`

rule 1577 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Simp[1/2 Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; Free
Q[{a, c, d, e, p, q}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4153 `Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Simp[c*(ff/f) Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + f
f^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m,
n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ratio
nalQ[n]))`

3.53.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.67 (sec) , antiderivative size = 431, normalized size of antiderivative = 5.82

3.53.
$$\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx$$

method	result
derivativedivides	$\frac{b \left(-\frac{\coth(x)^3}{4a(a+b)} + \frac{\coth(x)^2}{4a(a+b)} - \frac{\coth(x)}{4a(a+b)} - \frac{1}{4(a+b)b} \right)}{\sqrt{\left(\coth(x)^4 + \frac{a}{b} \right) b}} - \frac{\operatorname{arctanh} \left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}} \right) + \frac{\sqrt{1 - \frac{i\sqrt{b} \coth(x)^2}{\sqrt{a}}}}{\sqrt{a}} \sqrt{1 + \frac{i\sqrt{b} \coth(x)^2}{\sqrt{a}}}}{2(a+b) \sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}}$
default	$\frac{b \left(-\frac{\coth(x)^3}{4a(a+b)} + \frac{\coth(x)^2}{4a(a+b)} - \frac{\coth(x)}{4a(a+b)} - \frac{1}{4(a+b)b} \right)}{\sqrt{\left(\coth(x)^4 + \frac{a}{b} \right) b}} - \frac{\operatorname{arctanh} \left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}} \right) + \frac{\sqrt{1 - \frac{i\sqrt{b} \coth(x)^2}{\sqrt{a}}}}{\sqrt{a}} \sqrt{1 + \frac{i\sqrt{b} \coth(x)^2}{\sqrt{a}}}}{2(a+b) \sqrt{\frac{i\sqrt{b}}{\sqrt{a}}}}$

input `int(coth(x)/(a+b*coth(x)^4)^(3/2),x,method=_RETURNVERBOSE)`

output `b*(-1/4/a/(a+b)*coth(x)^3+1/4/a/(a+b)*coth(x)^2-1/4/a/(a+b)*coth(x)-1/4/(a+b)/b)/((coth(x)^4+a/b)*b)^(1/2)-1/2/(a+b)*(-1/2/(a+b)^(1/2)*arctanh(1/2*(2*b*coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))+1/(I/a^(1/2)*b^(1/2))^(1/2)*(1-I/a^(1/2)*b^(1/2)*coth(x)^2)^(1/2)*(1+I/a^(1/2)*b^(1/2)*coth(x)^2)^(1/2)/(a+b*coth(x)^4)^(1/2)*EllipticPi(coth(x)*(I/a^(1/2)*b^(1/2))^(1/2),-I*a^(1/2)/b^(1/2),(-I/a^(1/2)*b^(1/2))^(1/2)/(I/a^(1/2)*b^(1/2))^(1/2))+b*(1/4/a/(a+b)*coth(x)^3+1/4/a/(a+b)*coth(x)^2+1/4/a/(a+b)*coth(x)-1/4/(a+b)/b)/((coth(x)^4+a/b)*b)^(1/2)-1/2/(a+b)*(-1/2/(a+b)^(1/2)*arctanh(1/2*(2*b*coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))-1/(I/a^(1/2)*b^(1/2))^(1/2)*(1-I/a^(1/2)*b^(1/2)*coth(x)^2)^(1/2)*(1+I/a^(1/2)*b^(1/2)*coth(x)^2)^(1/2)/(a+b*coth(x)^4)^(1/2)*EllipticPi(coth(x)*(I/a^(1/2)*b^(1/2))^(1/2),-I*a^(1/2)/b^(1/2),(-I/a^(1/2)*b^(1/2))^(1/2)/(I/a^(1/2)*b^(1/2))^(1/2))`

3.53.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1947 vs. $2(63) = 126$.

Time = 0.50 (sec) , antiderivative size = 3938, normalized size of antiderivative = 53.22

$$\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx = \text{Too large to display}$$

input `integrate(coth(x)/(a+b*coth(x)^4)^(3/2),x, algorithm="fricas")`

output

```
[1/4*((a^2 + a*b)*cosh(x)^8 + 8*(a^2 + a*b)*cosh(x)*sinh(x)^7 + (a^2 + a*
b)*sinh(x)^8 - 4*(a^2 - a*b)*cosh(x)^6 + 4*(7*(a^2 + a*b)*cosh(x)^2 - a^2
+ a*b)*sinh(x)^6 + 8*(7*(a^2 + a*b)*cosh(x)^3 - 3*(a^2 - a*b)*cosh(x))*sin
h(x)^5 + 6*(a^2 + a*b)*cosh(x)^4 + 2*(35*(a^2 + a*b)*cosh(x)^4 - 30*(a^2 -
a*b)*cosh(x)^2 + 3*a^2 + 3*a*b)*sinh(x)^4 + 8*(7*(a^2 + a*b)*cosh(x)^5 -
10*(a^2 - a*b)*cosh(x)^3 + 3*(a^2 + a*b)*cosh(x))*sinh(x)^3 - 4*(a^2 - a*b
)*cosh(x)^2 + 4*(7*(a^2 + a*b)*cosh(x)^6 - 15*(a^2 - a*b)*cosh(x)^4 + 9*(a
^2 + a*b)*cosh(x)^2 - a^2 + a*b + 8*((a^2 + a*b)*co
sh(x)^7 - 3*(a^2 - a*b)*cosh(x)^5 + 3*(a^2 + a*b)*cosh(x)^3 - (a^2 - a*b)*
cosh(x))*sinh(x))*sqrt(a + b)*log(((a^2 + 2*a*b + b^2)*cosh(x)^8 + 8*(a^2
+ 2*a*b + b^2)*cosh(x)*sinh(x)^7 + (a^2 + 2*a*b + b^2)*sinh(x)^8 - 4*(a^2
- b^2)*cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)
^6 + 8*(7*(a^2 + 2*a*b + b^2)*cosh(x)^3 - 3*(a^2 - b^2)*cosh(x))*sinh(x)^5
+ 2*(3*a^2 + 2*a*b + 3*b^2)*cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2)*cosh(x)
^4 - 30*(a^2 - b^2)*cosh(x)^2 + 3*a^2 + 2*a*b + 3*b^2)*sinh(x)^4 + 8*(7*(a
^2 + 2*a*b + b^2)*cosh(x)^5 - 10*(a^2 - b^2)*cosh(x)^3 + (3*a^2 + 2*a*b +
3*b^2)*cosh(x))*sinh(x)^3 - 4*(a^2 - b^2)*cosh(x)^2 + 4*(7*(a^2 + 2*a*b +
b^2)*cosh(x)^6 - 15*(a^2 - b^2)*cosh(x)^4 + 3*(3*a^2 + 2*a*b + 3*b^2)*cosh
(x)^2 - a^2 + b^2)*sinh(x)^2 + sqrt(2)*((a + b)*cosh(x)^4 + 4*(a + b)*cosh
(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*...
```

3.53.6 Sympy [F]

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(a + b \coth^4(x))^{\frac{3}{2}}} dx$$

input `integrate(coth(x)/(a+b*coth(x)**4)**(3/2),x)`

output `Integral(coth(x)/(a + b*coth(x)**4)**(3/2), x)`

3.53.7 Maxima [F]

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth(x)^4 + a)^{\frac{3}{2}}} dx$$

input `integrate(coth(x)/(a+b*coth(x)^4)^(3/2),x, algorithm="maxima")`

output `integrate(coth(x)/(b*coth(x)^4 + a)^(3/2), x)`

3.53.8 Giac [F]

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth(x)^4 + a)^{\frac{3}{2}}} dx$$

input `integrate(coth(x)/(a+b*coth(x)^4)^(3/2),x, algorithm="giac")`

output `integrate(coth(x)/(b*coth(x)^4 + a)^(3/2), x)`

3.53.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth(x)^4 + a)^{3/2}} dx$$

input `int(coth(x)/(a + b*coth(x)^4)^(3/2),x)`

output `int(coth(x)/(a + b*coth(x)^4)^(3/2), x)`

APPENDIX

4.1 Listing of Grading functions	398
--	-----

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```



```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, CsCh,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsCh
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```

if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):

```



```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #instance(expn,Pow)
    if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #instance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #instance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #instance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```