

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

6-Hyperbolic-functions/6.6-Hyperbolic-cosecant/181-6.6.1-c+d-x-
 $\int \frac{1}{(c+dx)^m(a+b\operatorname{csch}^n(x))} dx$

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 6:24am

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	34
4	Appendix	279

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	15
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [29]. This is test number [181].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	100.00 (29)	0.00 (0)
Rubi	93.10 (27)	6.90 (2)
Fricas	86.21 (25)	13.79 (4)
Maple	65.52 (19)	34.48 (10)
Maxima	44.83 (13)	55.17 (16)
Mupad	27.59 (8)	72.41 (21)
Giac	27.59 (8)	72.41 (21)
Sympy	13.79 (4)	86.21 (25)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Mathematica	55.172	31.034	0.000	13.793
Rubi	27.586	0.000	51.724	20.690
Maple	13.793	37.931	0.000	48.276
Giac	10.345	3.448	0.000	86.207
Maxima	3.448	27.586	0.000	68.966
Fricas	0.000	72.414	0.000	27.586
Mupad	0.000	13.793	0.000	86.207
Sympy	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	0	0.00	0.00	0.00
Rubi	2	100.00	0.00	0.00
Fricas	4	0.00	0.00	100.00
Maple	10	100.00	0.00	0.00
Maxima	16	100.00	0.00	0.00
Mupad	21	0.00	100.00	0.00
Giac	21	100.00	0.00	0.00
Sympy	25	96.00	4.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.29
Maxima	0.30
Giac	0.36
Rubi	0.89
Sympy	1.80
Mupad	2.23
Maple	2.82
Mathematica	7.76

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Sympy	12.50	0.83	10.00	0.83
Giac	63.50	1.56	44.00	1.42
Mupad	68.00	1.48	30.50	1.28
Rubi	165.44	1.06	107.00	1.00
Maxima	213.62	3.72	168.00	2.44
Maple	333.63	2.36	240.00	2.28
Mathematica	675.97	1.98	137.00	1.72
Fricas	1522.68	5.76	692.00	4.69

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{4, 8, 12, 21}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {25}

Mathematica {17, 26, 27}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2013
Design v0.01

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	24
2.3	Detailed conclusion table specific for Rubi results	32

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	21
2.1.3	Maple	22
2.1.4	Fricas	22
2.1.5	Maxima	22
2.1.6	Giac	23
2.1.7	Mupad	23
2.1.8	Sympy	23

2.1.1 Rubi

A grade { 13, 14, 15, 16, 19, 20, 28, 29 }

B grade { }

C grade { 1, 2, 3, 5, 6, 7, 9, 10, 11, 17, 18, 22, 23, 24, 25 }

F normal fail { 26, 27 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 5, 6, 7, 9, 13, 14, 15, 16, 19, 20, 24, 25, 28, 29 }

B grade { 3, 10, 11, 17, 18, 22, 23, 26, 27 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.3 Maple

A grade { 3, 7, 20, 25 }

B grade { 1, 2, 5, 6, 9, 10, 11, 19, 24, 28, 29 }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 22, 23, 26, 27 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.4 Fricas

A grade { }

B grade { 1, 2, 3, 5, 6, 7, 9, 10, 11, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { 13, 14, 15, 16 }

2.1.5 Maxima

A grade { 25 }

B grade { 1, 2, 5, 7, 9, 10, 20, 29 }

C grade { }

F normal fail { 3, 6, 11, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 26, 27, 28 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.6 Giac

A grade { 20, 25, 29 }

B grade { 7 }

C grade { }

F normal fail { 1, 2, 3, 5, 6, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 26, 27, 28 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.7 Mupad

A grade { }

B grade { 7, 20, 25, 29 }

C grade { }

F normal fail { }

F(-1) timeout fail { 1, 2, 3, 5, 6, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 26, 27, 28 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 27, 28, 29 }

F(-1) timeout fail { 26 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column N.S. means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	149	173	168	541	333	396	0	0	0
N.S.	1	1.16	1.13	3.63	2.23	2.66	0.00	0.00	0.00
time (sec)	N/A	0.672	0.043	0.986	0.264	0.261	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	115	118	306	195	242	0	0	0
N.S.	1	1.16	1.19	3.09	1.97	2.44	0.00	0.00	0.00
time (sec)	N/A	0.455	0.042	0.945	0.282	0.265	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	61	116	89	0	119	0	0	0
N.S.	1	1.22	2.32	1.78	0.00	2.38	0.00	0.00	0.00
time (sec)	N/A	0.277	0.034	0.248	0.000	0.268	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	8	12	14
N.S.	1	1.00	1.20	1.00	1.20	1.20	0.80	1.20	1.40
time (sec)	N/A	0.189	20.670	0.047	0.432	0.260	0.296	0.320	2.141

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	146	185	473	320	1159	0	0	0
N.S.	1	1.42	1.80	4.59	3.11	11.25	0.00	0.00	0.00
time (sec)	N/A	0.709	0.485	0.802	0.347	0.271	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	106	137	240	0	623	0	0	0
N.S.	1	1.43	1.85	3.24	0.00	8.42	0.00	0.00	0.00
time (sec)	N/A	0.496	0.342	0.855	0.000	0.270	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	33	52	56	91	166	0	80	49
N.S.	1	1.14	1.79	1.93	3.14	5.72	0.00	2.76	1.69
time (sec)	N/A	0.262	0.048	0.805	0.198	0.274	0.000	0.295	2.113

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	73	14	10	14	14
N.S.	1	1.00	1.17	1.00	6.08	1.17	0.83	1.17	1.17
time (sec)	N/A	0.197	6.404	0.090	0.299	0.267	0.323	0.291	2.115

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	256	294	440	876	605	4008	0	0	0
N.S.	1	1.15	1.72	3.42	2.36	15.66	0.00	0.00	0.00
time (sec)	N/A	1.203	2.102	1.012	0.369	0.310	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	184	420	444	393	2218	0	0	0
N.S.	1	1.19	2.73	2.88	2.55	14.40	0.00	0.00	0.00
time (sec)	N/A	0.740	6.237	0.940	0.342	0.289	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	108	271	197	0	1026	0	0	0
N.S.	1	1.17	2.95	2.14	0.00	11.15	0.00	0.00	0.00
time (sec)	N/A	0.416	0.080	0.767	0.000	0.260	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	157	14	10	14	14
N.S.	1	1.00	1.17	1.00	13.08	1.17	0.83	1.17	1.17
time (sec)	N/A	0.202	42.663	0.097	0.332	0.244	0.326	0.643	2.138

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	17	0	0	0	0	0	0
N.S.	1	1.00	0.71	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.240	0.035	0.000	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	17	0	0	0	0	0	0
N.S.	1	1.00	0.71	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.241	0.051	0.000	0.000	0.000	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	47	47	45	0	0	0	0	0	0
N.S.	1	1.00	0.96	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.272	0.072	0.000	0.000	0.000	0.000	0.000	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	76	76	67	0	0	0	0	0	0
N.S.	1	1.00	0.88	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.343	0.188	0.000	0.000	0.000	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	F	F	B	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	448	452	1792	0	0	1976	0	0	0
N.S.	1	1.01	4.00	0.00	0.00	4.41	0.00	0.00	0.00
time (sec)	N/A	2.445	20.202	0.000	0.000	0.303	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	F	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	330	334	937	0	0	1265	0	0	0
N.S.	1	1.01	2.84	0.00	0.00	3.83	0.00	0.00	0.00
time (sec)	N/A	1.740	11.570	0.000	0.000	0.290	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	212	209	266	483	0	692	0	0	0
N.S.	1	0.99	1.25	2.28	0.00	3.26	0.00	0.00	0.00
time (sec)	N/A	1.038	0.392	1.170	0.000	0.289	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	30	30	49	83	132	0	60	31
N.S.	1	0.88	0.88	1.44	2.44	3.88	0.00	1.76	0.91
time (sec)	N/A	0.327	0.014	0.665	0.198	0.263	0.000	0.301	2.116

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	28	26	164	34	22	28	30
N.S.	1	1.00	1.08	1.00	6.31	1.31	0.85	1.08	1.15
time (sec)	N/A	0.336	45.950	0.201	0.393	0.267	6.271	0.405	2.318

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	F	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	696	641	2218	0	0	3847	0	0	0
N.S.	1	0.92	3.19	0.00	0.00	5.53	0.00	0.00	0.00
time (sec)	N/A	3.555	9.937	0.000	0.000	0.348	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	F	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	510	484	1216	0	0	2410	0	0	0
N.S.	1	0.95	2.38	0.00	0.00	4.73	0.00	0.00	0.00
time (sec)	N/A	2.829	4.935	0.000	0.000	0.314	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	327	324	610	1012	0	1284	0	0	0
N.S.	1	0.99	1.87	3.09	0.00	3.93	0.00	0.00	0.00
time (sec)	N/A	1.592	1.968	4.514	0.000	0.302	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	A	B	F	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	107	109	167	168	446	0	155	212
N.S.	1	1.13	1.15	1.76	1.77	4.69	0.00	1.63	2.23
time (sec)	N/A	0.611	0.343	4.001	0.283	0.279	0.000	0.304	2.440

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	B	F	F	B	F(-1)	F	F(-1)
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD
size	864	0	7404	0	0	7980	0	0	0
N.S.	1	0.00	8.57	0.00	0.00	9.24	0.00	0.00	0.00
time (sec)	N/A	0.000	32.297	0.000	0.000	0.377	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	B	F	F	B	F	F	F(-1)
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD
size	636	0	2129	0	0	4887	0	0	0
N.S.	1	0.00	3.35	0.00	0.00	7.68	0.00	0.00	0.00
time (sec)	N/A	0.000	16.795	0.000	0.000	0.336	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	400	360	696	1102	0	2465	0	0	0
N.S.	1	0.90	1.74	2.76	0.00	6.16	0.00	0.00	0.00
time (sec)	N/A	2.296	1.179	17.620	0.000	0.307	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	75	75	244	183	652	0	145	180
N.S.	1	0.88	0.88	2.87	2.15	7.67	0.00	1.71	2.12
time (sec)	N/A	0.394	0.101	17.864	0.220	0.268	0.000	0.317	2.472

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [22] had the largest ratio of [1.0714300000000010]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	C	8	7	1.16	14	0.500
2	C	7	6	1.16	14	0.429
3	C	6	5	1.22	12	0.417
4	N/A	3	0	1.00	10	0.000
5	C	12	11	1.42	16	0.688
6	C	11	10	1.43	16	0.625
7	C	7	7	1.14	14	0.500
8	N/A	3	0	1.00	12	0.000
9	C	14	13	1.15	16	0.812
10	C	12	11	1.19	16	0.688
11	C	10	9	1.17	14	0.643
12	N/A	3	0	1.00	12	0.000
13	A	1	1	1.00	20	0.050
14	A	1	1	1.00	20	0.050
15	A	1	1	1.00	20	0.050
16	A	1	1	1.00	24	0.042
17	C	21	20	1.01	26	0.769
18	C	16	15	1.01	26	0.577
19	A	13	12	0.99	24	0.500
20	A	12	11	0.88	19	0.579
21	N/A	2	0	1.00	26	0.000
22	C	31	30	0.92	28	1.071

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	C	28	27	0.95	28	0.964
24	C	20	19	0.99	26	0.731
25	C	15	14	1.13	21	0.667
26	F	0	0	N/A	0.000	N/A
27	F	0	0	N/A	0.000	N/A
28	A	27	26	0.90	26	1.000
29	A	12	11	0.88	21	0.524

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (c + dx)^3 \operatorname{csch}(a + bx) dx$	36
3.2	$\int (c + dx)^2 \operatorname{csch}(a + bx) dx$	44
3.3	$\int (c + dx) \operatorname{csch}(a + bx) dx$	51
3.4	$\int \frac{\operatorname{csch}(a+bx)}{x} dx$	57
3.5	$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx$	62
3.6	$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx$	70
3.7	$\int (c + dx) \operatorname{csch}^2(a + bx) dx$	76
3.8	$\int \frac{\operatorname{csch}^2(a+bx)}{x} dx$	81
3.9	$\int (c + dx)^3 \operatorname{csch}^3(a + bx) dx$	86
3.10	$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx$	96
3.11	$\int (c + dx) \operatorname{csch}^3(a + bx) dx$	105
3.12	$\int \frac{\operatorname{csch}^3(a+bx)}{x} dx$	112
3.13	$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx$	117
3.14	$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$	121
3.15	$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21}x\sqrt{\operatorname{csch}(x)} \right) dx$	125
3.16	$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} \right) dx$	129
3.17	$\int \frac{(e+fx)^3 \cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$	133
3.18	$\int \frac{(e+fx)^2 \cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$	144
3.19	$\int \frac{(e+fx) \cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$	154
3.20	$\int \frac{\cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$	162
3.21	$\int \frac{\cosh(c+dx)}{(e+fx)(a+b\operatorname{csch}(c+dx))} dx$	168
3.22	$\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$	172

3.23	$\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$	194
3.24	$\int \frac{(e+fx) \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$	210
3.25	$\int \frac{\cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$	221
3.26	$\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$	229
3.27	$\int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$	243
3.28	$\int \frac{(e+fx) \cosh^3(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$	259
3.29	$\int \frac{\cosh^3(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$	272

3.1 $\int (c + dx)^3 \operatorname{csch}(a + bx) dx$

3.1.1	Optimal result	36
3.1.2	Mathematica [A] (verified)	37
3.1.3	Rubi [C] (verified)	37
3.1.4	Maple [B] (verified)	40
3.1.5	Fricas [B] (verification not implemented)	41
3.1.6	Sympy [F]	41
3.1.7	Maxima [B] (verification not implemented)	42
3.1.8	Giac [F]	43
3.1.9	Mupad [F(-1)]	43

3.1.1 Optimal result

Integrand size = 14, antiderivative size = 149

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx = -\frac{2(c + dx)^3 \operatorname{arctanh}(e^{a+bx})}{b} - \frac{3d(c + dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b^2} + \frac{3d(c + dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b^2} + \frac{6d^2(c + dx) \operatorname{PolyLog}(3, -e^{a+bx})}{b^3} - \frac{6d^2(c + dx) \operatorname{PolyLog}(3, e^{a+bx})}{b^3} - \frac{6d^3 \operatorname{PolyLog}(4, -e^{a+bx})}{b^4} + \frac{6d^3 \operatorname{PolyLog}(4, e^{a+bx})}{b^4}$$

```
output -2*(d*x+c)^3*arctanh(exp(b*x+a))/b-3*d*(d*x+c)^2*polylog(2,-exp(b*x+a))/b^2+3*d*(d*x+c)^2*polylog(2,exp(b*x+a))/b^2+6*d^2*(d*x+c)*polylog(3,-exp(b*x+a))/b^3-6*d^2*(d*x+c)*polylog(3,exp(b*x+a))/b^3-6*d^3*polylog(4,-exp(b*x+a))/b^4+6*d^3*polylog(4,exp(b*x+a))/b^4
```

3.1.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.13

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx$$

$$= \frac{(c + dx)^3 \log(1 - e^{a+bx}) - (c + dx)^3 \log(1 + e^{a+bx}) - \frac{3d(b^2(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx}) - 2bd(c+dx) \operatorname{PolyLog}(3, -e^{a+bx}) - d^2 \operatorname{PolyLog}(4, -e^{a+bx}))}{b^3} + (c + dx)^3 \log(1 - e^{a+bx}) - (c + dx)^3 \log(1 + e^{a+bx}) - \frac{3d(b^2(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx}) - 2bd(c+dx) \operatorname{PolyLog}(3, e^{a+bx}) - d^2 \operatorname{PolyLog}(4, e^{a+bx}))}{b^3}}{b}$$

input `Integrate[(c + d*x)^3*Csch[a + b*x],x]`

output `((c + d*x)^3*Log[1 - E^(a + b*x)] - (c + d*x)^3*Log[1 + E^(a + b*x)] - (3*d*(b^2*(c + d*x)^2*PolyLog[2, -E^(a + b*x)] - 2*b*d*(c + d*x)*PolyLog[3, -E^(a + b*x)] + 2*d^2*PolyLog[4, -E^(a + b*x)]))/b^3 + (3*d*(b^2*(c + d*x)^2*PolyLog[2, E^(a + b*x)] - 2*b*d*(c + d*x)*PolyLog[3, E^(a + b*x)] + 2*d^2*PolyLog[4, E^(a + b*x)]))/b^3)/b`

3.1.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.67 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.16, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 26, 4670, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx$$

$$\downarrow 3042$$

$$\int i(c + dx)^3 \operatorname{csc}(ia + ibx) dx$$

$$\downarrow 26$$

$$i \int (c + dx)^3 \operatorname{csc}(ia + ibx) dx$$

$$\downarrow 4670$$

$$i \left(\frac{3id \int (c + dx)^2 \log(1 - e^{a+bx}) dx}{b} - \frac{3id \int (c + dx)^2 \log(1 + e^{a+bx}) dx}{b} + \frac{2i(c + dx)^3 \operatorname{arctanh}(e^{a+bx})}{b} \right)$$

↓ 3011

$$i \left(-\frac{3id \left(\frac{2d \int (c+dx) \operatorname{PolyLog}(2, -e^{a+bx}) dx}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{3id \left(\frac{2d \int (c+dx) \operatorname{PolyLog}(2, e^{a+bx}) dx}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right)$$

↓ 7163

$$i \left(-\frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{d \int \operatorname{PolyLog}(3, -e^{a+bx}) dx}{b} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{d \int \operatorname{PolyLog}(3, e^{a+bx}) dx}{b} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right)$$

↓ 2720

$$i \left(-\frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{d \int e^{-a-bx} \operatorname{PolyLog}(3, -e^{a+bx}) de^{a+bx}}{b^2} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{d \int e^{a+bx} \operatorname{PolyLog}(3, e^{a+bx}) de^{a+bx}}{b^2} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right)$$

↓ 7143

$$i \left(\frac{2i(c + dx)^3 \operatorname{arctanh}(e^{a+bx})}{b} - \frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{d \operatorname{PolyLog}(4, -e^{a+bx})}{b^2} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{d \operatorname{PolyLog}(4, e^{a+bx})}{b^2} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right)$$

input `Int[(c + d*x)^3*Csch[a + b*x],x]`

```
output I*(((2*I)*(c + d*x)^3*ArcTanh[E^(a + b*x)]/b - ((3*I)*d*(-(((c + d*x)^2*PolyLog[2, -E^(a + b*x)]/b) + (2*d*(((c + d*x)*PolyLog[3, -E^(a + b*x)]/b - (d*PolyLog[4, -E^(a + b*x)]/b^2))/b))/b) + ((3*I)*d*(-(((c + d*x)^2*PolyLog[2, E^(a + b*x)]/b) + (2*d*(((c + d*x)*PolyLog[3, E^(a + b*x)]/b - (d*PolyLog[4, E^(a + b*x)]/b^2))/b))/b))
```

3.1.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 3011 Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4670 Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```



```
rule 7163 Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

3.1.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 540 vs. $2(142) = 284$.

Time = 0.99 (sec) , antiderivative size = 541, normalized size of antiderivative = 3.63

method	result
risch	$-\frac{2c^3 \operatorname{arctanh}(e^{bx+a})}{b} - \frac{3d^3 \operatorname{polylog}(2, -e^{bx+a})x^2}{b^2} + \frac{6d^3 \operatorname{polylog}(3, -e^{bx+a})x}{b^3} + \frac{d^3 \ln(1 - e^{bx+a})x^3}{b} + \frac{d^3 \ln(1 - e^{bx+a})a^3}{b^4} +$

```
input int((d*x+c)^3*cscsch(b*x+a),x,method=_RETURNVERBOSE)
```

```
output -2/b*c^3*arctanh(exp(b*x+a))-3/b^2*d^3*polylog(2,-exp(b*x+a))*x^2+6/b^3*d^
3*polylog(3,-exp(b*x+a))*x+1/b*d^3*ln(1-exp(b*x+a))*x^3+1/b^4*d^3*ln(1-exp
(b*x+a))*a^3+3/b^2*d^3*polylog(2,exp(b*x+a))*x^2-6/b^3*d^3*polylog(3,exp(b
*x+a))*x-1/b*d^3*ln(exp(b*x+a)+1)*x^3-1/b^4*d^3*ln(exp(b*x+a)+1)*a^3+2/b^4
*d^3*a^3*arctanh(exp(b*x+a))+6/b^3*c*d^2*polylog(3,-exp(b*x+a))+3/b^2*d*c^
2*polylog(2,exp(b*x+a))-3/b^2*d*c^2*polylog(2,-exp(b*x+a))-6/b^3*c*d^2*pol
ylog(3,exp(b*x+a))+6/b^2*d*a*c^2*arctanh(exp(b*x+a))-3/b*d*c^2*ln(exp(b*x+
a)+1)*x-3/b^2*d*c^2*ln(exp(b*x+a)+1)*a-6/b^3*d^2*a^2*c*arctanh(exp(b*x+a))
+3/b*c*d^2*ln(1-exp(b*x+a))*x^2-3/b^3*c*d^2*ln(1-exp(b*x+a))*a^2+6/b^2*c*d
^2*polylog(2,exp(b*x+a))*x-3/b*c*d^2*ln(exp(b*x+a)+1)*x^2+3/b^3*c*d^2*ln(e
xp(b*x+a)+1)*a^2-6/b^2*c*d^2*polylog(2,-exp(b*x+a))*x+3/b*d*c^2*ln(1-exp(b
*x+a))*x+3/b^2*d*c^2*ln(1-exp(b*x+a))*a+6*d^3*polylog(4,exp(b*x+a))/b^4-6*
d^3*polylog(4,-exp(b*x+a))/b^4
```

3.1.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 396 vs. $2(140) = 280$.

Time = 0.26 (sec) , antiderivative size = 396, normalized size of antiderivative = 2.66

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx$$

$$= \frac{6d^3 \operatorname{polylog}(4, \cosh(bx + a) + \sinh(bx + a)) - 6d^3 \operatorname{polylog}(4, -\cosh(bx + a) - \sinh(bx + a)) + 3(b^2 d^3 x^3 + b^2 c^2 d^2 x^2 + 3b^3 c^2 d^2 x + b^3 c^3) \log(\cosh(bx + a) + \sinh(bx + a)) + 3(b^2 d^3 x^3 + 3b^3 c^2 d^2 x^2 + 3b^3 c^2 d^2 x + b^3 c^3) \log(-\cosh(bx + a) - \sinh(bx + a)) + 3(b^2 d^3 x^3 + 3b^3 c^2 d^2 x^2 + 3b^3 c^2 d^2 x + b^3 c^3) \log(\cosh(bx + a) + \sinh(bx + a) + 1) + 3(b^2 d^3 x^3 + 3b^3 c^2 d^2 x^2 + 3b^3 c^2 d^2 x + b^3 c^3) \log(-\cosh(bx + a) - \sinh(bx + a) + 1) - 6(b^2 d^3 x^3 + b^2 c^2 d^2 x^2) \operatorname{polylog}(3, \cosh(bx + a) + \sinh(bx + a)) + 6(b^2 d^3 x^3 + b^2 c^2 d^2 x^2) \operatorname{polylog}(3, -\cosh(bx + a) - \sinh(bx + a))}{b^4}$$

input `integrate((d*x+c)^3*csch(b*x+a),x, algorithm="fracas")`

output `(6*d^3*polylog(4, cosh(b*x + a) + sinh(b*x + a)) - 6*d^3*polylog(4, -cosh(b*x + a) - sinh(b*x + a)) + 3*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d)*dilog(cosh(b*x + a) + sinh(b*x + a)) - 3*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d)*dilog(-cosh(b*x + a) - sinh(b*x + a)) - (b^3*d^3*x^3 + 3*b^3*c*d^2*x^2 + 3*b^3*c^2*d*x + b^3*c^3)*log(cosh(b*x + a) + sinh(b*x + a) + 1) + (b^3*d^3*x^3 + 3*b^3*c*d^2*x^2 + 3*b^3*c^2*d*x + b^3*c^3)*log(-cosh(b*x + a) - sinh(b*x + a) + 1) - 6*(b*d^3*x + b*c*d^2)*polylog(3, cosh(b*x + a) + sinh(b*x + a)) + 6*(b*d^3*x + b*c*d^2)*polylog(3, -cosh(b*x + a) - sinh(b*x + a)))/b^4`

3.1.6 Sympy [F]

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx = \int (c + dx)^3 \operatorname{csch}(a + bx) dx$$

input `integrate((d*x+c)**3*csch(b*x+a),x)`

output `Integral((c + d*x)**3*csch(a + b*x), x)`

3.1.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 333 vs. $2(140) = 280$.

Time = 0.26 (sec) , antiderivative size = 333, normalized size of antiderivative = 2.23

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx$$

$$= -c^3 \left(\frac{\log(e^{-bx-a}) + 1}{b} - \frac{\log(e^{-bx-a}) - 1}{b} \right)$$

$$- \frac{3 (bx \log(e^{bx+a}) + 1) + \operatorname{Li}_2(-e^{bx+a})}{b^2} c^2 d$$

$$+ \frac{3 (bx \log(-e^{bx+a}) + 1) + \operatorname{Li}_2(e^{bx+a})}{b^2} c^2 d$$

$$- \frac{3 (b^2 x^2 \log(e^{bx+a}) + 1) + 2 bx \operatorname{Li}_2(-e^{bx+a}) - 2 \operatorname{Li}_3(-e^{bx+a})}{b^3} c d^2$$

$$+ \frac{3 (b^2 x^2 \log(-e^{bx+a}) + 1) + 2 bx \operatorname{Li}_2(e^{bx+a}) - 2 \operatorname{Li}_3(e^{bx+a})}{b^3} c d^2$$

$$- \frac{(b^3 x^3 \log(e^{bx+a}) + 1) + 3 b^2 x^2 \operatorname{Li}_2(-e^{bx+a}) - 6 bx \operatorname{Li}_3(-e^{bx+a}) + 6 \operatorname{Li}_4(-e^{bx+a})}{b^4} d^3$$

$$+ \frac{(b^3 x^3 \log(-e^{bx+a}) + 1) + 3 b^2 x^2 \operatorname{Li}_2(e^{bx+a}) - 6 bx \operatorname{Li}_3(e^{bx+a}) + 6 \operatorname{Li}_4(e^{bx+a})}{b^4} d^3$$

input `integrate((d*x+c)^3*csh(b*x+a),x, algorithm="maxima")`

output

```
-c^3*(log(e^(-b*x - a) + 1)/b - log(e^(-b*x - a) - 1)/b) - 3*(b*x*log(e^(b*x + a) + 1) + dilog(-e^(b*x + a)))*c^2*d/b^2 + 3*(b*x*log(-e^(b*x + a) + 1) + dilog(e^(b*x + a)))*c^2*d/b^2 - 3*(b^2*x^2*log(e^(b*x + a) + 1) + 2*b*x*dilog(-e^(b*x + a)) - 2*polylog(3, -e^(b*x + a)))*c*d^2/b^3 + 3*(b^2*x^2*log(-e^(b*x + a) + 1) + 2*b*x*dilog(e^(b*x + a)) - 2*polylog(3, e^(b*x + a)))*c*d^2/b^3 - (b^3*x^3*log(e^(b*x + a) + 1) + 3*b^2*x^2*dilog(-e^(b*x + a)) - 6*b*x*polylog(3, -e^(b*x + a)) + 6*polylog(4, -e^(b*x + a)))*d^3/b^4 + (b^3*x^3*log(-e^(b*x + a) + 1) + 3*b^2*x^2*dilog(e^(b*x + a)) - 6*b*x*polylog(3, e^(b*x + a)) + 6*polylog(4, e^(b*x + a)))*d^3/b^4
```

3.1.8 Giac [F]

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx = \int (dx + c)^3 \operatorname{csch}(bx + a) dx$$

input `integrate((d*x+c)^3*csch(b*x+a),x, algorithm="giac")`

output `integrate((d*x + c)^3*csch(b*x + a), x)`

3.1.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 \operatorname{csch}(a + bx) dx = \int \frac{(c + dx)^3}{\sinh(a + bx)} dx$$

input `int((c + d*x)^3/sinh(a + b*x),x)`

output `int((c + d*x)^3/sinh(a + b*x), x)`

3.2 $\int (c + dx)^2 \operatorname{csch}(a + bx) dx$

3.2.1	Optimal result	44
3.2.2	Mathematica [A] (verified)	44
3.2.3	Rubi [C] (verified)	45
3.2.4	Maple [B] (verified)	47
3.2.5	Fricas [B] (verification not implemented)	48
3.2.6	Sympy [F]	48
3.2.7	Maxima [B] (verification not implemented)	48
3.2.8	Giac [F]	49
3.2.9	Mupad [F(-1)]	50

3.2.1 Optimal result

Integrand size = 14, antiderivative size = 99

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx = -\frac{2(c + dx)^2 \operatorname{arctanh}(e^{a+bx})}{b} - \frac{2d(c + dx) \operatorname{PolyLog}(2, -e^{a+bx})}{b^2} + \frac{2d(c + dx) \operatorname{PolyLog}(2, e^{a+bx})}{b^2} + \frac{2d^2 \operatorname{PolyLog}(3, -e^{a+bx})}{b^3} - \frac{2d^2 \operatorname{PolyLog}(3, e^{a+bx})}{b^3}$$

```
output -2*(d*x+c)^2*arctanh(exp(b*x+a))/b-2*d*(d*x+c)*polylog(2,-exp(b*x+a))/b^2+
2*d*(d*x+c)*polylog(2,exp(b*x+a))/b^2+2*d^2*polylog(3,-exp(b*x+a))/b^3-2*d
^2*polylog(3,exp(b*x+a))/b^3
```

3.2.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.19

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx = \frac{(c + dx)^2 \log(1 - e^{a+bx}) - (c + dx)^2 \log(1 + e^{a+bx}) - \frac{2d(b(c+dx) \operatorname{PolyLog}(2, -e^{a+bx}) - d \operatorname{PolyLog}(3, -e^{a+bx}))}{b^2} + \frac{2d(b(c+dx) \operatorname{PolyLog}(2, e^{a+bx}) - d \operatorname{PolyLog}(3, e^{a+bx}))}{b^2}}{b}$$

```
input Integrate[(c + d*x)^2*Csch[a + b*x], x]
```

output $((c + dx)^2 \text{Log}[1 - E^{(a + b*x)}] - (c + dx)^2 \text{Log}[1 + E^{(a + b*x)}] - (2*d*(b*(c + d*x)*\text{PolyLog}[2, -E^{(a + b*x)}] - d*\text{PolyLog}[3, -E^{(a + b*x)}]))/b^2 + (2*d*(b*(c + d*x)*\text{PolyLog}[2, E^{(a + b*x)}] - d*\text{PolyLog}[3, E^{(a + b*x)}]))/b^2)/b$

3.2.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 26, 4670, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx$$

$$\downarrow 3042$$

$$\int i(c + dx)^2 \operatorname{csc}(ia + ibx) dx$$

$$\downarrow 26$$

$$i \int (c + dx)^2 \operatorname{csc}(ia + ibx) dx$$

$$\downarrow 4670$$

$$i \left(\frac{2id \int (c + dx) \log(1 - e^{a+bx}) dx}{b} - \frac{2id \int (c + dx) \log(1 + e^{a+bx}) dx}{b} + \frac{2i(c + dx)^2 \operatorname{arctanh}(e^{a+bx})}{b} \right)$$

$$\downarrow 3011$$

$$i \left(-\frac{2id \left(\frac{d \int \operatorname{PolyLog}(2, -e^{a+bx}) dx}{b} - \frac{(c+dx) \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{2id \left(\frac{d \int \operatorname{PolyLog}(2, e^{a+bx}) dx}{b} - \frac{(c+dx) \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} + \dots \right)$$

$$\downarrow 2720$$

$$i \left(-\frac{2id \left(\frac{d \int e^{-a-bx} \operatorname{PolyLog}(2, -e^{a+bx}) de^{a+bx}}{b^2} - \frac{(c+dx) \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{2id \left(\frac{d \int e^{-a-bx} \operatorname{PolyLog}(2, e^{a+bx}) de^{a+bx}}{b^2} - \frac{(c+dx) \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} + \dots \right)$$

↓ 7143

$$i \left(\frac{2i(c+dx)^2 \operatorname{arctanh}(e^{a+bx})}{b} - \frac{2id \left(\frac{d \operatorname{PolyLog}(3, -e^{a+bx})}{b^2} - \frac{(c+dx) \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{2id \left(\frac{d \operatorname{PolyLog}(3, e^{a+bx})}{b^2} - \frac{(c+dx) \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right)$$

input `Int[(c + d*x)^2*Csch[a + b*x], x]`

output `I*(((2*I)*(c + d*x)^2*ArcTanh[E^(a + b*x)])/b - ((2*I)*d*(-(((c + d*x)*PolyLog[2, -E^(a + b*x)])/b) + (d*PolyLog[3, -E^(a + b*x)]/b^2))/b + ((2*I)*d*(-(((c + d*x)*PolyLog[2, E^(a + b*x)])/b) + (d*PolyLog[3, E^(a + b*x)]/b^2))/b)`

3.2.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x_, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*x)))]^(n_)*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4670 Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

3.2.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 305 vs. $2(94) = 188$.

Time = 0.94 (sec) , antiderivative size = 306, normalized size of antiderivative = 3.09

method	result
risch	$\frac{2cd \ln(1 - e^{bx+a})x}{b} + \frac{2cd \operatorname{polylog}(2, e^{bx+a})}{b^2} - \frac{2cd \ln(e^{bx+a} + 1)x}{b} - \frac{2cd \operatorname{polylog}(2, -e^{bx+a})}{b^2} + \frac{2d^2 \operatorname{polylog}(2, e^{bx+a})x}{b^2} + \frac{d^2 \ln(e^{bx+a} + 1)}{b}$

```
input int((d*x+c)^2*csc(b*x+a),x,method=_RETURNVERBOSE)
```

```
output 2/b*c*d*ln(1-exp(b*x+a))*x+2/b^2*c*d*polylog(2,exp(b*x+a))-2/b*c*d*ln(exp(
b*x+a)+1)*x-2/b^2*c*d*polylog(2,-exp(b*x+a))+2/b^2*d^2*polylog(2,exp(b*x+a
))*x+1/b^3*d^2*ln(exp(b*x+a)+1)*a^2-2/b^2*d^2*polylog(2,-exp(b*x+a))*x-1/b
^3*d^2*ln(1-exp(b*x+a))*a^2-2/b^3*d^2*a^2*arctanh(exp(b*x+a))+1/b*d^2*ln(1
-exp(b*x+a))*x^2-2*d^2*polylog(3,exp(b*x+a))/b^3-1/b*d^2*ln(exp(b*x+a)+1)*
x^2+2*d^2*polylog(3,-exp(b*x+a))/b^3+2/b^2*c*d*ln(1-exp(b*x+a))*a-2/b^2*c*
d*ln(exp(b*x+a)+1)*a+4/b^2*d*a*c*arctanh(exp(b*x+a))-2/b*c^2*arctanh(exp(b
*x+a))
```


3.2.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 242 vs. 2(92) = 184.

Time = 0.27 (sec) , antiderivative size = 242, normalized size of antiderivative = 2.44

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx = \frac{-2d^2 \operatorname{polylog}(3, \cosh(bx + a) + \sinh(bx + a)) - 2d^2 \operatorname{polylog}(3, -\cosh(bx + a) - \sinh(bx + a)) - 2(bd^2 \operatorname{dilog}(\cosh(bx + a) + \sinh(bx + a)) + 2(bd^2 \operatorname{dilog}(-\cosh(bx + a) - \sinh(bx + a)) + (b^2 d^2 x^2 + 2b^2 c dx + b^2 c^2) \log(\cosh(bx + a) + \sinh(bx + a) + 1) - (b^2 c^2 - 2a b c d + a^2 d^2) \log(\cosh(bx + a) + \sinh(bx + a) - 1) - (b^2 d^2 x^2 + 2b^2 c dx + 2a b c d - a^2 d^2) \log(-\cosh(bx + a) - \sinh(bx + a) + 1))}{b^3}$$

input `integrate((d*x+c)^2*csch(b*x+a),x, algorithm="fricas")`

output `-(2*d^2*polylog(3, cosh(b*x + a) + sinh(b*x + a)) - 2*d^2*polylog(3, -cosh(b*x + a) - sinh(b*x + a)) - 2*(b*d^2*x + b*c*d)*dilog(cosh(b*x + a) + sinh(b*x + a)) + 2*(b*d^2*x + b*c*d)*dilog(-cosh(b*x + a) - sinh(b*x + a)) + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*log(cosh(b*x + a) + sinh(b*x + a) + 1) - (b^2*c^2 - 2*a*b*c*d + a^2*d^2)*log(cosh(b*x + a) + sinh(b*x + a) - 1) - (b^2*d^2*x^2 + 2*b^2*c*d*x + 2*a*b*c*d - a^2*d^2)*log(-cosh(b*x + a) - sinh(b*x + a) + 1))/b^3`

3.2.6 Sympy [F]

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx = \int (c + dx)^2 \operatorname{csch}(a + bx) dx$$

input `integrate((d*x+c)**2*csch(b*x+a),x)`

output `Integral((c + d*x)**2*csch(a + b*x), x)`

3.2.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 195 vs. 2(92) = 184.

Time = 0.28 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.97

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx$$

$$= -c^2 \left(\frac{\log(e^{-bx-a} + 1)}{b} - \frac{\log(e^{-bx-a} - 1)}{b} \right)$$

$$- \frac{2(bx \log(e^{(bx+a)} + 1) + \operatorname{Li}_2(-e^{(bx+a)}))cd}{b^2} + \frac{2(bx \log(-e^{(bx+a)} + 1) + \operatorname{Li}_2(e^{(bx+a)}))cd}{b^2}$$

$$- \frac{(b^2 x^2 \log(e^{(bx+a)} + 1) + 2bx \operatorname{Li}_2(-e^{(bx+a)}) - 2 \operatorname{Li}_3(-e^{(bx+a)}))d^2}{b^3}$$

$$+ \frac{(b^2 x^2 \log(-e^{(bx+a)} + 1) + 2bx \operatorname{Li}_2(e^{(bx+a)}) - 2 \operatorname{Li}_3(e^{(bx+a)}))d^2}{b^3}$$

input `integrate((d*x+c)^2*csch(b*x+a),x, algorithm="maxima")`

output `-c^2*(log(e^(-b*x - a) + 1)/b - log(e^(-b*x - a) - 1)/b) - 2*(b*x*log(e^(b*x + a) + 1) + dilog(-e^(b*x + a)))*c*d/b^2 + 2*(b*x*log(-e^(b*x + a) + 1) + dilog(e^(b*x + a)))*c*d/b^2 - (b^2*x^2*log(e^(b*x + a) + 1) + 2*b*x*dilog(-e^(b*x + a)) - 2*polylog(3, -e^(b*x + a)))*d^2/b^3 + (b^2*x^2*log(-e^(b*x + a) + 1) + 2*b*x*dilog(e^(b*x + a)) - 2*polylog(3, e^(b*x + a)))*d^2/b^3`

3.2.8 Giac [F]

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx = \int (dx + c)^2 \operatorname{csch}(bx + a) dx$$

input `integrate((d*x+c)^2*csch(b*x+a),x, algorithm="giac")`

output `integrate((d*x + c)^2*csch(b*x + a), x)`

3.2.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 \operatorname{csch}(a + bx) dx = \int \frac{(c + dx)^2}{\sinh(a + bx)} dx$$

input `int((c + d*x)^2/sinh(a + b*x),x)`output `int((c + d*x)^2/sinh(a + b*x), x)`

3.3 $\int (c + dx)\operatorname{csch}(a + bx) dx$

3.3.1	Optimal result	51
3.3.2	Mathematica [B] (verified)	51
3.3.3	Rubi [C] (verified)	52
3.3.4	Maple [A] (verified)	54
3.3.5	Fricas [B] (verification not implemented)	54
3.3.6	Sympy [F]	55
3.3.7	Maxima [F]	55
3.3.8	Giac [F]	55
3.3.9	Mupad [F(-1)]	56

3.3.1 Optimal result

Integrand size = 12, antiderivative size = 50

$$\int (c + dx)\operatorname{csch}(a + bx) dx = -\frac{2(c + dx)\operatorname{arctanh}(e^{a+bx})}{b} - \frac{d \operatorname{PolyLog}(2, -e^{a+bx})}{b^2} + \frac{d \operatorname{PolyLog}(2, e^{a+bx})}{b^2}$$

output `-2*(d*x+c)*arctanh(exp(b*x+a))/b-d*polylog(2,-exp(b*x+a))/b^2+d*polylog(2,exp(b*x+a))/b^2`

3.3.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 116 vs. 2(50) = 100.

Time = 0.03 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.32

$$\int (c + dx)\operatorname{csch}(a + bx) dx = -\frac{c \log(\cosh(\frac{a}{2} + \frac{bx}{2}))}{b} + \frac{c \log(\sinh(\frac{a}{2} + \frac{bx}{2}))}{b} + 2d \left(\frac{x \log(1 - e^{a+bx})}{2b} - \frac{x \log(1 + e^{a+bx})}{2b} - \frac{\operatorname{PolyLog}(2, -e^{a+bx})}{2b^2} + \frac{\operatorname{PolyLog}(2, e^{a+bx})}{2b^2} \right)$$

input `Integrate[(c + d*x)*Csch[a + b*x],x]`

output $-\frac{(c \operatorname{Log}[\operatorname{Cosh}[a/2 + (b*x)/2]])}{b} + \frac{(c \operatorname{Log}[\operatorname{Sinh}[a/2 + (b*x)/2]])}{b} + 2*d*(x \operatorname{Log}[1 - E^{(a + b*x)}])/(2*b) - (x \operatorname{Log}[1 + E^{(a + b*x)}])/(2*b) - \operatorname{PolyLog}[2, -E^{(a + b*x)}]/(2*b^2) + \operatorname{PolyLog}[2, E^{(a + b*x)}]/(2*b^2)$

3.3.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.22, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx) \operatorname{csch}(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int i(c + dx) \operatorname{csc}(ia + ibx) dx \\
 & \quad \downarrow \text{26} \\
 & i \int (c + dx) \operatorname{csc}(ia + ibx) dx \\
 & \quad \downarrow \text{4670} \\
 & i \left(\frac{id \int \log(1 - e^{a+bx}) dx}{b} - \frac{id \int \log(1 + e^{a+bx}) dx}{b} + \frac{2i(c + dx) \operatorname{arctanh}(e^{a+bx})}{b} \right) \\
 & \quad \downarrow \text{2715} \\
 & i \left(\frac{id \int e^{-a-bx} \log(1 - e^{a+bx}) de^{a+bx}}{b^2} - \frac{id \int e^{-a-bx} \log(1 + e^{a+bx}) de^{a+bx}}{b^2} + \frac{2i(c + dx) \operatorname{arctanh}(e^{a+bx})}{b} \right) \\
 & \quad \downarrow \text{2838} \\
 & i \left(\frac{2i(c + dx) \operatorname{arctanh}(e^{a+bx})}{b} + \frac{id \operatorname{PolyLog}(2, -e^{a+bx})}{b^2} - \frac{id \operatorname{PolyLog}(2, e^{a+bx})}{b^2} \right)
 \end{aligned}$$

input `Int[(c + d*x)*Csch[a + b*x],x]`

output `I*(((2*I)*(c + d*x)*ArcTanh[E^(a + b*x)])/b + (I*d*PolyLog[2, -E^(a + b*x)])/b^2 - (I*d*PolyLog[2, E^(a + b*x)])/b^2)`

3.3.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

3.3.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.78

method	result
derivativedivides	$\frac{d((bx+a)\ln(1-e^{bx+a})+\text{polylog}(2,e^{bx+a})-(bx+a)\ln(e^{bx+a}+1)-\text{polylog}(2,-e^{bx+a}))}{b} + \frac{2da \operatorname{arctanh}(e^{bx+a})}{b} - 2c \operatorname{arctanh}(e^{bx+a})$
default	$\frac{d((bx+a)\ln(1-e^{bx+a})+\text{polylog}(2,e^{bx+a})-(bx+a)\ln(e^{bx+a}+1)-\text{polylog}(2,-e^{bx+a}))}{b} + \frac{2da \operatorname{arctanh}(e^{bx+a})}{b} - 2c \operatorname{arctanh}(e^{bx+a})$
parts	$\frac{\ln\left(\tanh\left(\frac{bx}{2}+\frac{a}{2}\right)\right)dx}{b} + \frac{\ln\left(\tanh\left(\frac{bx}{2}+\frac{a}{2}\right)\right)c}{b} + \frac{2d\left(-\frac{\operatorname{dilog}\left(1+\tanh\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{2} - \frac{\ln\left(\tanh\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\ln\left(1+\tanh\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{2}\right)}{b^2}$
risch	$-\frac{2c \operatorname{arctanh}(e^{bx+a})}{b} + \frac{d\ln(1-e^{bx+a})x}{b} + \frac{d\ln(1-e^{bx+a})a}{b^2} + \frac{d\text{polylog}(2,e^{bx+a})}{b^2} - \frac{d\ln(e^{bx+a}+1)x}{b} - \frac{d\ln(e^{bx+a})}{b}$

input `int((d*x+c)*csch(b*x+a),x,method=_RETURNVERBOSE)`

output `1/b*(d/b*((b*x+a)*ln(1-exp(b*x+a))+polylog(2,exp(b*x+a))-(b*x+a)*ln(exp(b*x+a)+1)-polylog(2,-exp(b*x+a)))+2*d/b*a*arctanh(exp(b*x+a))-2*c*arctanh(exp(b*x+a)))`

3.3.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(45) = 90$.

Time = 0.27 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.38

$$\int (c + dx)\operatorname{csch}(a + bx) dx$$

$$= \frac{d\operatorname{Li}_2(\cosh(bx + a) + \sinh(bx + a)) - d\operatorname{Li}_2(-\cosh(bx + a) - \sinh(bx + a)) - (bdx + bc)\log(\cosh(bx + a) + \sinh(bx + a))}{b^2}$$

input `integrate((d*x+c)*csch(b*x+a),x, algorithm="fracas")`

output `(d*dilog(cosh(b*x + a) + sinh(b*x + a)) - d*dilog(-cosh(b*x + a) - sinh(b*x + a)) - (b*d*x + b*c)*log(cosh(b*x + a) + sinh(b*x + a) + 1) + (b*c - a*d)*log(cosh(b*x + a) + sinh(b*x + a) - 1) + (b*d*x + a*d)*log(-cosh(b*x + a) - sinh(b*x + a) + 1))/b^2`

3.3.6 Sympy [F]

$$\int (c + dx) \operatorname{csch}(a + bx) dx = \int (c + dx) \operatorname{csch}(a + bx) dx$$

input `integrate((d*x+c)*csch(b*x+a),x)`

output `Integral((c + d*x)*csch(a + b*x), x)`

3.3.7 Maxima [F]

$$\int (c + dx) \operatorname{csch}(a + bx) dx = \int (dx + c) \operatorname{csch}(bx + a) dx$$

input `integrate((d*x+c)*csch(b*x+a),x, algorithm="maxima")`

output `-c*(log(e^(-b*x - a) + 1)/b - log(e^(-b*x - a) - 1)/b) + 2*d*(integrate(1/2*x/(e^(b*x + a) + 1), x) + integrate(1/2*x/(e^(b*x + a) - 1), x))`

3.3.8 Giac [F]

$$\int (c + dx) \operatorname{csch}(a + bx) dx = \int (dx + c) \operatorname{csch}(bx + a) dx$$

input `integrate((d*x+c)*csch(b*x+a),x, algorithm="giac")`

output `integrate((d*x + c)*csch(b*x + a), x)`

3.3.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx) \operatorname{csch}(a + bx) dx = \int \frac{c + dx}{\sinh(a + bx)} dx$$

input `int((c + d*x)/sinh(a + b*x),x)`output `int((c + d*x)/sinh(a + b*x), x)`

3.4 $\int \frac{\operatorname{csch}(a+bx)}{x} dx$

3.4.1	Optimal result	57
3.4.2	Mathematica [N/A]	57
3.4.3	Rubi [N/A]	58
3.4.4	Maple [N/A] (verified)	59
3.4.5	Fricas [N/A]	59
3.4.6	Sympy [N/A]	60
3.4.7	Maxima [N/A]	60
3.4.8	Giac [N/A]	60
3.4.9	Mupad [N/A]	61

3.4.1 Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\operatorname{csch}(a + bx)}{x} dx = \operatorname{Int}\left(\frac{\operatorname{csch}(a + bx)}{x}, x\right)$$

output `Unintegrable(csch(b*x+a)/x,x)`

3.4.2 Mathematica [N/A]

Not integrable

Time = 20.67 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{csch}(a + bx)}{x} dx = \int \frac{\operatorname{csch}(a + bx)}{x} dx$$

input `Integrate[Csch[a + b*x]/x,x]`

output `Integrate[Csch[a + b*x]/x, x]`

3.4.3 Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 26, 4680}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\operatorname{csch}(a+bx)}{x} dx \\ \downarrow 3042 \\ \int \frac{i \operatorname{csc}(ia+ibx)}{x} dx \\ \downarrow 26 \\ i \int \frac{\operatorname{csc}(ia+ibx)}{x} dx \\ \downarrow 4680 \\ \int \frac{\operatorname{csch}(a+bx)}{x} dx \end{array}$$

input `Int[Csch[a + b*x]/x,x]`

output `$Aborted`

3.4.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

```
rule 4680 Int[csc[(e_.) + (f_.)*(x_.)]^(n_.)*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] :>
Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], Un
integrable[(c + d*x)^m*Sech[I*(e - Pi/2) + I*f*x]^n, x], (-I)^n*Unintegrabl
e[(c + d*x)^m*Csch[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], Uni
ntegrable[(c + d*x)^m*Sec[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*C
sc[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]
```

3.4.4 Maple [N/A] (verified)

Not integrable

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{csch}(bx+a)}{x} dx$$

input `int(csch(b*x+a)/x,x)`

output `int(csch(b*x+a)/x,x)`

3.4.5 Fracas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{csch}(a+bx)}{x} dx = \int \frac{\operatorname{csch}(bx+a)}{x} dx$$

input `integrate(csch(b*x+a)/x,x, algorithm="fricas")`

output `integral(csch(b*x + a)/x, x)`

3.4.6 Sympy [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\operatorname{csch}(a + bx)}{x} dx = \int \frac{\operatorname{csch}(a + bx)}{x} dx$$

input `integrate(csch(b*x+a)/x,x)`output `Integral(csch(a + b*x)/x, x)`**3.4.7 Maxima [N/A]**

Not integrable

Time = 0.43 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{csch}(a + bx)}{x} dx = \int \frac{\operatorname{csch}(bx + a)}{x} dx$$

input `integrate(csch(b*x+a)/x,x, algorithm="maxima")`output `integrate(csch(b*x + a)/x, x)`**3.4.8 Giac [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{csch}(a + bx)}{x} dx = \int \frac{\operatorname{csch}(bx + a)}{x} dx$$

input `integrate(csch(b*x+a)/x,x, algorithm="giac")`output `integrate(csch(b*x + a)/x, x)`

3.4.9 Mupad [N/A]

Not integrable

Time = 2.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40

$$\int \frac{\operatorname{csch}(a + bx)}{x} dx = \int \frac{1}{x \sinh(a + bx)} dx$$

input `int(1/(x*sinh(a + b*x)),x)`

output `int(1/(x*sinh(a + b*x)), x)`

3.5 $\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx$

3.5.1	Optimal result	62
3.5.2	Mathematica [A] (verified)	62
3.5.3	Rubi [C] (verified)	63
3.5.4	Maple [B] (verified)	66
3.5.5	Fricas [B] (verification not implemented)	67
3.5.6	Sympy [F]	67
3.5.7	Maxima [B] (verification not implemented)	68
3.5.8	Giac [F]	69
3.5.9	Mupad [F(-1)]	69

3.5.1 Optimal result

Integrand size = 16, antiderivative size = 103

$$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx = -\frac{(c + dx)^3}{b} - \frac{(c + dx)^3 \operatorname{coth}(a + bx)}{b} + \frac{3d(c + dx)^2 \log(1 - e^{2(a+bx)})}{b^2} + \frac{3d^2(c + dx) \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^3} - \frac{3d^3 \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^4}$$

```
output -(d*x+c)^3/b-(d*x+c)^3*coth(b*x+a)/b+3*d*(d*x+c)^2*ln(1-exp(2*b*x+2*a))/b^2+3*d^2*(d*x+c)*polylog(2,exp(2*b*x+2*a))/b^3-3/2*d^3*polylog(3,exp(2*b*x+2*a))/b^4
```

3.5.2 Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.80

$$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx = \frac{-\frac{2(c+dx)^3}{-1+e^{2a}} + \frac{3d(c+dx)^2 \log(1-e^{-a-bx})}{b} + \frac{3d(c+dx)^2 \log(1+e^{-a-bx})}{b} - \frac{6d^2(b(c+dx) \operatorname{PolyLog}(2, -e^{-a-bx}) + d \operatorname{PolyLog}(3, -e^{-a-bx}))}{b^3}}{b}$$

input `Integrate[(c + d*x)^3*Csch[a + b*x]^2,x]`

output `((-2*(c + d*x)^3)/(-1 + E^(2*a)) + (3*d*(c + d*x)^2*Log[1 - E^(-a - b*x)])
/b + (3*d*(c + d*x)^2*Log[1 + E^(-a - b*x)])/b - (6*d^2*(b*(c + d*x)*PolyL
og[2, -E^(-a - b*x)] + d*PolyLog[3, -E^(-a - b*x)]))/b^3 - (6*d^2*(b*(c +
d*x)*PolyLog[2, E^(-a - b*x)] + d*PolyLog[3, E^(-a - b*x)]))/b^3 + (c + d*
x)^3*Csch[a]*Csch[a + b*x]*Sinh[b*x])/b`

3.5.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.71 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.42, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {3042, 25, 4672, 26, 3042, 26, 4201, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^3 \operatorname{csch}^2(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -(c + dx)^3 \operatorname{csc}(ia + ibx)^2 dx \\
 & \quad \downarrow \text{25} \\
 & - \int (c + dx)^3 \operatorname{csc}(ia + ibx)^2 dx \\
 & \quad \downarrow \text{4672} \\
 & - \frac{(c + dx)^3 \operatorname{coth}(a + bx)}{b} + \frac{3id \int -i(c + dx)^2 \operatorname{coth}(a + bx) dx}{b} \\
 & \quad \downarrow \text{26} \\
 & \frac{3d \int (c + dx)^2 \operatorname{coth}(a + bx) dx}{b} - \frac{(c + dx)^3 \operatorname{coth}(a + bx)}{b} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{(c + dx)^3 \operatorname{coth}(a + bx)}{b} + \frac{3d \int -i(c + dx)^2 \tan\left(ia + ibx + \frac{\pi}{2}\right) dx}{b} \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{(c+dx)^3 \operatorname{coth}(a+bx)}{b} - \frac{3id \int (c+dx)^2 \tan\left(\frac{1}{2}(2ia+\pi)+ibx\right) dx}{b} \\
 & \quad \downarrow \text{4201} \\
 & -\frac{(c+dx)^3 \operatorname{coth}(a+bx)}{b} - \frac{3id \left(2i \int \frac{e^{2a+2bx-i\pi} (c+dx)^2 dx}{1+e^{2a+2bx-i\pi}} - \frac{i(c+dx)^3}{3d} \right)}{b} \\
 & \quad \downarrow \text{2620} \\
 & -\frac{(c+dx)^3 \operatorname{coth}(a+bx)}{b} - \frac{3id \left(2i \left(\frac{(c+dx)^2 \log(1+e^{2a+2bx-i\pi})}{2b} - \frac{d \int (c+dx) \log(1+e^{2a+2bx-i\pi}) dx}{b} \right) - \frac{i(c+dx)^3}{3d} \right)}{b} \\
 & \quad \downarrow \text{3011} \\
 & -\frac{(c+dx)^3 \operatorname{coth}(a+bx)}{b} - \frac{3id \left(2i \left(\frac{(c+dx)^2 \log(1+e^{2a+2bx-i\pi})}{2b} - \frac{d \left(\frac{\int \operatorname{PolyLog}(2, -e^{2a+2bx-i\pi}) dx}{2b} - \frac{(c+dx) \operatorname{PolyLog}(2, -e^{2a+2bx-i\pi})}{2b} \right)}{b} \right) - \frac{i(c+dx)^3}{3d} \right)}{b} \\
 & \quad \downarrow \text{2720} \\
 & -\frac{(c+dx)^3 \operatorname{coth}(a+bx)}{b} - \frac{3id \left(2i \left(\frac{(c+dx)^2 \log(1+e^{2a+2bx-i\pi})}{2b} - \frac{d \left(\frac{\int e^{-2a-2bx+i\pi} \operatorname{PolyLog}(2, -e^{2a+2bx-i\pi}) dx}{4b^2} - \frac{(c+dx) \operatorname{PolyLog}(2, -e^{2a+2bx-i\pi})}{2b} \right)}{b} \right) - \frac{i(c+dx)^3}{3d} \right)}{b} \\
 & \quad \downarrow \text{7143} \\
 & -\frac{(c+dx)^3 \operatorname{coth}(a+bx)}{b} - \frac{3id \left(2i \left(\frac{(c+dx)^2 \log(1+e^{2a+2bx-i\pi})}{2b} - \frac{d \left(\frac{\int \operatorname{PolyLog}(3, -e^{2a+2bx-i\pi}) dx}{4b^2} - \frac{(c+dx) \operatorname{PolyLog}(2, -e^{2a+2bx-i\pi})}{2b} \right)}{b} \right) - \frac{i(c+dx)^3}{3d} \right)}{b}
 \end{aligned}$$

input `Int[(c + d*x)^3*Csch[a + b*x]^2,x]`

```
output -(((c + d*x)^3*Coth[a + b*x])/b) - ((3*I)*d*(((1/3*I)*(c + d*x)^3)/d + (2
*I)*(((c + d*x)^2*Log[1 + E^(2*a - I*Pi + 2*b*x)])/(2*b) - (d*(-1/2*((c +
d*x)*PolyLog[2, -E^(2*a - I*Pi + 2*b*x)])/b + (d*PolyLog[3, -E^(2*a - I*Pi
+ 2*b*x)])/(4*b^2)))/b)))/b
```

3.5.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 2620 Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 3011 Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4201 Int[((c_.) + (d_.)*(x_.))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_.)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

```
rule 4672 Int[csc[(e_.) + (f_.)*(x_.)]^2*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] := Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)
*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

3.5.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. $2(101) = 202$.

Time = 0.80 (sec) , antiderivative size = 473, normalized size of antiderivative = 4.59

method	result
risch	$-\frac{2d^3x^3}{b} - \frac{6d^3 \operatorname{polylog}(3, e^{bx+a})}{b^4} - \frac{6d^3 \operatorname{polylog}(3, -e^{bx+a})}{b^4} + \frac{4d^3a^3}{b^4} - \frac{2(d^3x^3 + 3cd^2x^2 + 3c^2dx + c^3)}{(e^{2bx+2a}-1)b} - \frac{12d^2acx}{b^2} - \frac{6d^2ca}{b^2}$

```
input int((d*x+c)^3*csch(b*x+a)^2,x,method=_RETURNVERBOSE)
```

```
output -2/b*d^3*x^3-6/b^4*d^3*polylog(3,exp(b*x+a))-6/b^4*d^3*polylog(3,-exp(b*x+
a))+4/b^4*d^3*a^3-2*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)/(exp(2*b*x+2*a)-1)
/b-12/b^2*d^2*a*c*x-6/b^3*d^2*c*a*ln(exp(b*x+a)-1)+12/b^3*d^2*c*a*ln(exp(b
*x+a))+6/b^2*d^2*c*ln(exp(b*x+a)+1)*x+6/b^2*d^2*c*ln(1-exp(b*x+a))*x+6/b^3
*d^2*c*ln(1-exp(b*x+a))*a-6/b*d^2*c*x^2+6/b^3*d^3*a^2*x-6/b^3*d^2*c*a^2+3/
b^2*d*c^2*ln(exp(b*x+a)-1)+3/b^2*d*c^2*ln(exp(b*x+a)+1)-6/b^2*d*c^2*ln(exp
(b*x+a))+3/b^4*d^3*a^2*ln(exp(b*x+a)-1)-6/b^4*d^3*a^2*ln(exp(b*x+a))+3/b^2
*d^3*ln(exp(b*x+a)+1)*x^2+6/b^3*d^3*polylog(2,-exp(b*x+a))*x+3/b^2*d^3*ln(
1-exp(b*x+a))*x^2-3/b^4*d^3*ln(1-exp(b*x+a))*a^2+6/b^3*d^3*polylog(2,exp(b
*x+a))*x+6/b^3*d^2*c*polylog(2,-exp(b*x+a))+6/b^3*d^2*c*polylog(2,exp(b*x+
a))
```

3.5.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1159 vs. $2(100) = 200$.

Time = 0.27 (sec) , antiderivative size = 1159, normalized size of antiderivative = 11.25

$$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx = \text{Too large to display}$$

```
input integrate((d*x+c)^3*csh(b*x+a)^2,x, algorithm="fricas")
```

```
output -(2*b^3*c^3 - 6*a*b^2*c^2*d + 6*a^2*b*c*d^2 - 2*a^3*d^3 + 2*(b^3*d^3*x^3 +
3*b^3*c*d^2*x^2 + 3*b^3*c^2*d*x + 3*a*b^2*c^2*d - 3*a^2*b*c*d^2 + a^3*d^3
)*cosh(b*x + a)^2 + 4*(b^3*d^3*x^3 + 3*b^3*c*d^2*x^2 + 3*b^3*c^2*d*x + 3*a
*b^2*c^2*d - 3*a^2*b*c*d^2 + a^3*d^3)*cosh(b*x + a)*sinh(b*x + a) + 2*(b^3
*d^3*x^3 + 3*b^3*c*d^2*x^2 + 3*b^3*c^2*d*x + 3*a*b^2*c^2*d - 3*a^2*b*c*d^2
+ a^3*d^3)*sinh(b*x + a)^2 + 6*(b*d^3*x + b*c*d^2 - (b*d^3*x + b*c*d^2)*c
osh(b*x + a)^2 - 2*(b*d^3*x + b*c*d^2)*cosh(b*x + a)*sinh(b*x + a) - (b*d^
3*x + b*c*d^2)*sinh(b*x + a)^2)*dilog(cosh(b*x + a) + sinh(b*x + a)) + 6*(
b*d^3*x + b*c*d^2 - (b*d^3*x + b*c*d^2)*cosh(b*x + a)^2 - 2*(b*d^3*x + b*c
*d^2)*cosh(b*x + a)*sinh(b*x + a) - (b*d^3*x + b*c*d^2)*sinh(b*x + a)^2)*d
ilog(-cosh(b*x + a) - sinh(b*x + a)) + 3*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^
2*c^2*d - (b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d)*cosh(b*x + a)^2 - 2*(b
^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d)*cosh(b*x + a)*sinh(b*x + a) - (b^2
*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d)*sinh(b*x + a)^2)*log(cosh(b*x + a) +
sinh(b*x + a) + 1) + 3*(b^2*c^2*d - 2*a*b*c*d^2 + a^2*d^3 - (b^2*c^2*d -
2*a*b*c*d^2 + a^2*d^3)*cosh(b*x + a)^2 - 2*(b^2*c^2*d - 2*a*b*c*d^2 + a^2*
d^3)*cosh(b*x + a)*sinh(b*x + a) - (b^2*c^2*d - 2*a*b*c*d^2 + a^2*d^3)*sin
h(b*x + a)^2)*log(cosh(b*x + a) + sinh(b*x + a) - 1) + 3*(b^2*d^3*x^2 + 2*
b^2*c*d^2*x + 2*a*b*c*d^2 - a^2*d^3 - (b^2*d^3*x^2 + 2*b^2*c*d^2*x + 2*a*b
*c*d^2 - a^2*d^3)*cosh(b*x + a)^2 - 2*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + 2*...
```

3.5.6 Sympy [F]

$$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx = \int (c + dx)^3 \operatorname{csch}^2(a + bx) dx$$

```
input integrate((d*x+c)**3*csh(b*x+a)**2,x)
```

```
output Integral((c + d*x)**3*csh(a + b*x)**2, x)
```

3.5.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. $2(100) = 200$.

Time = 0.35 (sec) , antiderivative size = 320, normalized size of antiderivative = 3.11

$$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx$$

$$= -3c^2d \left(\frac{2xe^{(2bx+2a)}}{be^{(2bx+2a)} - b} - \frac{\log((e^{(bx+a)} + 1)e^{(-a)})}{b^2} - \frac{\log((e^{(bx+a)} - 1)e^{(-a)})}{b^2} \right)$$

$$+ \frac{6(bx \log(e^{(bx+a)} + 1) + \operatorname{Li}_2(-e^{(bx+a)}))cd^2}{b^3}$$

$$+ \frac{6(bx \log(-e^{(bx+a)} + 1) + \operatorname{Li}_2(e^{(bx+a)}))cd^2}{b^3} + \frac{2c^3}{b(e^{(-2bx-2a)} - 1)} - \frac{2(d^3x^3 + 3cd^2x^2)}{be^{(2bx+2a)} - b}$$

$$+ \frac{3(b^2x^2 \log(e^{(bx+a)} + 1) + 2bx \operatorname{Li}_2(-e^{(bx+a)}) - 2 \operatorname{Li}_3(-e^{(bx+a)}))d^3}{b^4}$$

$$+ \frac{3(b^2x^2 \log(-e^{(bx+a)} + 1) + 2bx \operatorname{Li}_2(e^{(bx+a)}) - 2 \operatorname{Li}_3(e^{(bx+a)}))d^3}{b^4}$$

$$- \frac{2(b^3d^3x^3 + 3b^3cd^2x^2)}{b^4}$$

input `integrate((d*x+c)^3*cshch(b*x+a)^2,x, algorithm="maxima")`

output `-3*c^2*d*(2*x*e^(2*b*x + 2*a)/(b*e^(2*b*x + 2*a) - b) - log((e^(b*x + a) + 1)*e^(-a))/b^2 - log((e^(b*x + a) - 1)*e^(-a))/b^2) + 6*(b*x*log(e^(b*x + a) + 1) + dilog(-e^(b*x + a)))*c*d^2/b^3 + 6*(b*x*log(-e^(b*x + a) + 1) + dilog(e^(b*x + a)))*c*d^2/b^3 + 2*c^3/(b*(e^(-2*b*x - 2*a) - 1)) - 2*(d^3*x^3 + 3*c*d^2*x^2)/(b*e^(2*b*x + 2*a) - b) + 3*(b^2*x^2*log(e^(b*x + a) + 1) + 2*b*x*dilog(-e^(b*x + a)) - 2*polylog(3, -e^(b*x + a)))*d^3/b^4 + 3*(b^2*x^2*log(-e^(b*x + a) + 1) + 2*b*x*dilog(e^(b*x + a)) - 2*polylog(3, e^(b*x + a)))*d^3/b^4 - 2*(b^3*d^3*x^3 + 3*b^3*c*d^2*x^2)/b^4`

3.5.8 Giac [F]

$$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx = \int (dx + c)^3 \operatorname{csch}(bx + a)^2 dx$$

input `integrate((d*x+c)^3*csh(b*x+a)^2,x, algorithm="giac")`

output `integrate((d*x + c)^3*csh(b*x + a)^2, x)`

3.5.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 \operatorname{csch}^2(a + bx) dx = \int \frac{(c + dx)^3}{\sinh(a + bx)^2} dx$$

input `int((c + d*x)^3/sinh(a + b*x)^2,x)`

output `int((c + d*x)^3/sinh(a + b*x)^2, x)`

3.6 $\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx$

3.6.1	Optimal result	70
3.6.2	Mathematica [A] (verified)	70
3.6.3	Rubi [C] (verified)	71
3.6.4	Maple [B] (verified)	73
3.6.5	Fricas [B] (verification not implemented)	74
3.6.6	Sympy [F]	74
3.6.7	Maxima [F]	75
3.6.8	Giac [F]	75
3.6.9	Mupad [F(-1)]	75

3.6.1 Optimal result

Integrand size = 16, antiderivative size = 74

$$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx = -\frac{(c + dx)^2}{b} - \frac{(c + dx)^2 \operatorname{coth}(a + bx)}{b} + \frac{2d(c + dx) \log(1 - e^{2(a+bx)})}{b^2} + \frac{d^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^3}$$

output $-(d*x+c)^2/b-(d*x+c)^2*\operatorname{coth}(b*x+a)/b+2*d*(d*x+c)*\ln(1-\exp(2*b*x+2*a))/b^2+d^2*\operatorname{polylog}(2,\exp(2*b*x+2*a))/b^3$

3.6.2 Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.85

$$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx = \frac{-\frac{2b(c+dx)(b(c+dx)-d(-1+e^{2a})\log(1-e^{-a-bx})-d(-1+e^{2a})\log(1+e^{-a-bx}))}{-1+e^{2a}} - 2d^2 \operatorname{PolyLog}(2, -e^{-a-bx}) - 2d^2 \operatorname{PolyLog}(2, e^{-a-bx})}{b^3}$$

input `Integrate[(c + d*x)^2*Csch[a + b*x]^2,x]`

output $((-2*b*(c + d*x)*(b*(c + d*x) - d*(-1 + E^{2*a}))*\operatorname{Log}[1 - E^{-a - b*x}] - d*(-1 + E^{2*a}))*\operatorname{Log}[1 + E^{-a - b*x}]))/(-1 + E^{2*a}) - 2*d^2*\operatorname{PolyLog}[2, -E^{-a - b*x}] - 2*d^2*\operatorname{PolyLog}[2, E^{-a - b*x}] + b^2*(c + d*x)^2*\operatorname{Csch}[a]*\operatorname{Csch}[a + b*x]*\operatorname{Sinh}[b*x])/b^3$

3.6.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.43, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3042, 25, 4672, 26, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c+dx)^2 \operatorname{csch}^2(a+bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -(c+dx)^2 \operatorname{csc}(ia+ibx)^2 dx \\
 & \quad \downarrow \text{25} \\
 & - \int (c+dx)^2 \operatorname{csc}(ia+ibx)^2 dx \\
 & \quad \downarrow \text{4672} \\
 & - \frac{(c+dx)^2 \operatorname{coth}(a+bx)}{b} + \frac{2id \int -i(c+dx) \operatorname{coth}(a+bx) dx}{b} \\
 & \quad \downarrow \text{26} \\
 & \frac{2d \int (c+dx) \operatorname{coth}(a+bx) dx}{b} - \frac{(c+dx)^2 \operatorname{coth}(a+bx)}{b} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{(c+dx)^2 \operatorname{coth}(a+bx)}{b} + \frac{2d \int -i(c+dx) \tan\left(ia+ibx+\frac{\pi}{2}\right) dx}{b} \\
 & \quad \downarrow \text{26} \\
 & - \frac{(c+dx)^2 \operatorname{coth}(a+bx)}{b} - \frac{2id \int (c+dx) \tan\left(\frac{1}{2}(2ia+\pi)+ibx\right) dx}{b} \\
 & \quad \downarrow \text{4201} \\
 & - \frac{(c+dx)^2 \operatorname{coth}(a+bx)}{b} - \frac{2id \left(2i \int \frac{e^{2a+2bx-i\pi}(c+dx)}{1+e^{2a+2bx-i\pi}} dx - \frac{i(c+dx)^2}{2d} \right)}{b} \\
 & \quad \downarrow \text{2620} \\
 & - \frac{(c+dx)^2 \operatorname{coth}(a+bx)}{b} - \frac{2id \left(2i \left(\frac{(c+dx) \log(1+e^{2a+2bx-i\pi})}{2b} - \frac{d \int \log(1+e^{2a+2bx-i\pi}) dx}{2b} \right) - \frac{i(c+dx)^2}{2d} \right)}{b}
 \end{aligned}$$

3.6. $\int (c+dx)^2 \operatorname{csch}^2(a+bx) dx$

$$\begin{array}{c}
 \downarrow 2715 \\
 \frac{(c+dx)^2 \coth(a+bx)}{b} - \frac{2id \left(2i \left(\frac{(c+dx) \log(1+e^{2a+2bx-i\pi})}{2b} - \frac{d \int e^{-2a-2bx+i\pi} \log(1+e^{2a+2bx-i\pi}) de^{2a+2bx-i\pi}}{4b^2} \right) - \frac{i(c+dx)^2}{2d} \right)}{b} \\
 \downarrow 2838 \\
 \frac{(c+dx)^2 \coth(a+bx)}{b} - \frac{2id \left(2i \left(\frac{d \text{PolyLog}(2, -e^{2a+2bx-i\pi})}{4b^2} + \frac{(c+dx) \log(1+e^{2a+2bx-i\pi})}{2b} \right) - \frac{i(c+dx)^2}{2d} \right)}{b}
 \end{array}$$

input `Int[(c + d*x)^2*Csch[a + b*x]^2,x]`

output `-(((c + d*x)^2*Coth[a + b*x])/b) - ((2*I)*d*(((1/2*I)*(c + d*x)^2)/d + (2*I)*(((c + d*x)*Log[1 + E^(2*a - I*Pi + 2*b*x)])/(2*b) + (d*PolyLog[2, -E^(2*a - I*Pi + 2*b*x)])/(4*b^2))))/b`

3.6.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 4672 `Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.6.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(74) = 148$.

Time = 0.86 (sec) , antiderivative size = 240, normalized size of antiderivative = 3.24

method	result
risch	$-\frac{2(x^2 d^2 + 2cdx + c^2)}{(e^{2bx+2a}-1)b} + \frac{2dc \ln(e^{bx+a}-1)}{b^2} + \frac{2dc \ln(e^{bx+a}+1)}{b^2} - \frac{4dc \ln(e^{bx+a})}{b^2} - \frac{2d^2 x^2}{b} - \frac{4d^2 ax}{b^2} - \frac{2d^2 a^2}{b^3} + \frac{2d^2 \ln(e^{bx+a})}{b^2}$

input `int((d*x+c)^2*csc(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `-2*(d^2*x^2+2*c*d*x+c^2)/(exp(2*b*x+2*a)-1)/b+2/b^2*d*c*ln(exp(b*x+a)-1)+2/b^2*d*c*ln(exp(b*x+a)+1)-4/b^2*d*c*ln(exp(b*x+a))-2/b*d^2*x^2-4/b^2*d^2*a*x-2/b^3*d^2*a^2+2/b^2*d^2*ln(exp(b*x+a)+1)*x+2/b^3*d^2*polylog(2,-exp(b*x+a))+2/b^2*d^2*ln(1-exp(b*x+a))*x+2/b^3*d^2*ln(1-exp(b*x+a))*a+2/b^3*d^2*polylog(2,exp(b*x+a))-2/b^3*d^2*a*ln(exp(b*x+a)-1)+4/b^3*d^2*a*ln(exp(b*x+a))`

3.6.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 623 vs. 2(73) = 146.

Time = 0.27 (sec) , antiderivative size = 623, normalized size of antiderivative = 8.42

$$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx = \frac{2(b^2c^2 - 2abcd + a^2d^2 + (b^2d^2x^2 + 2b^2cdx + 2abcd - a^2d^2) \cosh(bx + a)^2 + 2(b^2d^2x^2 + 2b^2cdx + 2abcd - a^2d^2) \sinh(bx + a)^2 + 2(b^2d^2x^2 + 2b^2cdx + 2abcd - a^2d^2) \operatorname{dilog}(\cosh(bx + a) + \sinh(bx + a)) + 2(b^2d^2x^2 + 2b^2cdx + 2abcd - a^2d^2) \operatorname{dilog}(-\cosh(bx + a) - \sinh(bx + a)) + 2(b^2d^2x^2 + 2b^2cdx + 2abcd - a^2d^2) \log(\cosh(bx + a) + \sinh(bx + a)) + 2(b^2d^2x^2 + 2b^2cdx + 2abcd - a^2d^2) \log(-\cosh(bx + a) - \sinh(bx + a))}{(b^3 \cosh(bx + a)^2 + 2b^3 \cosh(bx + a) \sinh(bx + a) + b^3 \sinh(bx + a)^2 - b^3)}$$

input `integrate((d*x+c)^2*csh(b*x+a)^2,x, algorithm="fricas")`

output

```
-2*(b^2*c^2 - 2*a*b*c*d + a^2*d^2 + (b^2*d^2*x^2 + 2*b^2*c*d*x + 2*a*b*c*d - a^2*d^2)*cosh(b*x + a)^2 + 2*(b^2*d^2*x^2 + 2*b^2*c*d*x + 2*a*b*c*d - a^2*d^2)*cosh(b*x + a)*sinh(b*x + a) + (b^2*d^2*x^2 + 2*b^2*c*d*x + 2*a*b*c*d - a^2*d^2)*sinh(b*x + a)^2 - (d^2*cosh(b*x + a)^2 + 2*d^2*cosh(b*x + a)*sinh(b*x + a) + d^2*sinh(b*x + a)^2 - d^2)*dilog(cosh(b*x + a) + sinh(b*x + a)) - (d^2*cosh(b*x + a)^2 + 2*d^2*cosh(b*x + a)*sinh(b*x + a) + d^2*sinh(b*x + a)^2 - d^2)*dilog(-cosh(b*x + a) - sinh(b*x + a)) + (b*d^2*x + b*c*d - (b*d^2*x + b*c*d)*cosh(b*x + a)^2 - 2*(b*d^2*x + b*c*d)*cosh(b*x + a)*sinh(b*x + a) - (b*d^2*x + b*c*d)*sinh(b*x + a)^2)*log(cosh(b*x + a) + sinh(b*x + a) + 1) + (b*c*d - a*d^2 - (b*c*d - a*d^2)*cosh(b*x + a)^2 - 2*(b*c*d - a*d^2)*cosh(b*x + a)*sinh(b*x + a) - (b*c*d - a*d^2)*sinh(b*x + a)^2)*log(cosh(b*x + a) + sinh(b*x + a) - 1) + (b*d^2*x + a*d^2 - (b*d^2*x + a*d^2)*cosh(b*x + a)^2 - 2*(b*d^2*x + a*d^2)*cosh(b*x + a)*sinh(b*x + a) - (b*d^2*x + a*d^2)*sinh(b*x + a)^2)*log(-cosh(b*x + a) - sinh(b*x + a) + 1))/(b^3*cosh(b*x + a)^2 + 2*b^3*cosh(b*x + a)*sinh(b*x + a) + b^3*sinh(b*x + a)^2 - b^3)
```

3.6.6 Sympy [F]

$$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx = \int (c + dx)^2 \operatorname{csch}^2(a + bx) dx$$

input `integrate((d*x+c)**2*csh(b*x+a)**2,x)`

output `Integral((c + d*x)**2*csh(a + b*x)**2, x)`

3.6.7 Maxima [F]

$$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx = \int (dx + c)^2 \operatorname{csch}(bx + a)^2 dx$$

input `integrate((d*x+c)^2*csh(b*x+a)^2,x, algorithm="maxima")`

output `-2*d^2*(x^2/(b*e^(2*b*x + 2*a) - b) + 2*integrate(1/2*x/(b*e^(b*x + a) + b), x) - 2*integrate(1/2*x/(b*e^(b*x + a) - b), x)) - 2*c*d*(2*x*e^(2*b*x + 2*a)/(b*e^(2*b*x + 2*a) - b) - log((e^(b*x + a) + 1)*e^(-a))/b^2 - log((e^(b*x + a) - 1)*e^(-a))/b^2) + 2*c^2/(b*(e^(-2*b*x - 2*a) - 1))`

3.6.8 Giac [F]

$$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx = \int (dx + c)^2 \operatorname{csch}(bx + a)^2 dx$$

input `integrate((d*x+c)^2*csh(b*x+a)^2,x, algorithm="giac")`

output `integrate((d*x + c)^2*csh(b*x + a)^2, x)`

3.6.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 \operatorname{csch}^2(a + bx) dx = \int \frac{(c + dx)^2}{\sinh(a + bx)^2} dx$$

input `int((c + d*x)^2/sinh(a + b*x)^2,x)`

output `int((c + d*x)^2/sinh(a + b*x)^2, x)`

3.7 $\int (c + dx) \operatorname{csch}^2(a + bx) dx$

3.7.1	Optimal result	76
3.7.2	Mathematica [A] (verified)	76
3.7.3	Rubi [C] (verified)	77
3.7.4	Maple [A] (verified)	78
3.7.5	Fricas [B] (verification not implemented)	79
3.7.6	Sympy [F]	79
3.7.7	Maxima [B] (verification not implemented)	79
3.7.8	Giac [B] (verification not implemented)	80
3.7.9	Mupad [B] (verification not implemented)	80

3.7.1 Optimal result

Integrand size = 14, antiderivative size = 29

$$\int (c + dx) \operatorname{csch}^2(a + bx) dx = -\frac{(c + dx) \operatorname{coth}(a + bx)}{b} + \frac{d \log(\sinh(a + bx))}{b^2}$$

output `-(d*x+c)*coth(b*x+a)/b+d*ln(sinh(b*x+a))/b^2`

3.7.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.79

$$\int (c + dx) \operatorname{csch}^2(a + bx) dx = -\frac{dx \operatorname{coth}(a)}{b} - \frac{c \operatorname{coth}(a + bx)}{b} + \frac{d \log(\sinh(a + bx))}{b^2} + \frac{dx \operatorname{csch}(a) \operatorname{csch}(a + bx) \sinh(bx)}{b}$$

input `Integrate[(c + d*x)*Csch[a + b*x]^2,x]`

output `-((d*x*Coth[a])/b) - (c*Coth[a + b*x])/b + (d*Log[Sinh[a + b*x]])/b^2 + (d*x*Csch[a]*Csch[a + b*x]*Sinh[b*x])/b`

3.7.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 25, 4672, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx) \operatorname{csch}^2(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -((c + dx) \operatorname{csc}(ia + ibx)^2) dx \\
 & \quad \downarrow \text{25} \\
 & - \int (c + dx) \operatorname{csc}(ia + ibx)^2 dx \\
 & \quad \downarrow \text{4672} \\
 & - \frac{(c + dx) \operatorname{coth}(a + bx)}{b} + \frac{id \int -i \operatorname{coth}(a + bx) dx}{b} \\
 & \quad \downarrow \text{26} \\
 & \frac{d \int \operatorname{coth}(a + bx) dx}{b} - \frac{(c + dx) \operatorname{coth}(a + bx)}{b} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{(c + dx) \operatorname{coth}(a + bx)}{b} + \frac{d \int -i \tan\left(ia + ibx + \frac{\pi}{2}\right) dx}{b} \\
 & \quad \downarrow \text{26} \\
 & - \frac{(c + dx) \operatorname{coth}(a + bx)}{b} - \frac{id \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right) dx}{b} \\
 & \quad \downarrow \text{3956} \\
 & - \frac{(c + dx) \operatorname{coth}(a + bx)}{b} + \frac{d \log(-i \sinh(a + bx))}{b^2}
 \end{aligned}$$

input `Int[(c + d*x)*Csch[a + b*x]^2,x]`

output $-\left(\frac{(c + dx)\operatorname{Coth}[a + bx]}{b} + \frac{d\operatorname{Log}[-1]\operatorname{Sinh}[a + bx]}{b^2}\right)$

3.7.3.1 Defintions of rubi rules used

rule 25 $\operatorname{Int}[-(F_x), x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$

rule 26 $\operatorname{Int}[(\operatorname{Complex}[0, a])*(F_x), x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Complex}[\operatorname{Identity}[0], a]) \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ \operatorname{EqQ}[a^2, 1]$

rule 3042 $\operatorname{Int}[u, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\operatorname{Int}[\tan[(c.) + (d.)*(x.)], x_Symbol] \rightarrow \operatorname{Simp}[-\operatorname{Log}[\operatorname{RemoveContent}[\operatorname{Cos}[c + dx], x]]/d, x] /; \operatorname{FreeQ}[\{c, d\}, x]$

rule 4672 $\operatorname{Int}[\operatorname{csc}[(e.) + (f.)*(x.)]^2*((c.) + (d.)*(x.))^{(m.)}, x_Symbol] \rightarrow \operatorname{Simp}[(-(c + dx)^m)*(\operatorname{Cot}[e + fx]/f), x] + \operatorname{Simp}[d*(m/f) \operatorname{Int}[(c + dx)^{(m - 1)} * \operatorname{Cot}[e + fx], x], x] /; \operatorname{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \operatorname{GtQ}[m, 0]$

3.7.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.93

method	result	size
risch	$-\frac{2dx}{b} - \frac{2da}{b^2} - \frac{2(dx+c)}{(e^{2bx+2a}-1)b} + \frac{d \ln(e^{2bx+2a}-1)}{b^2}$	56
parallelrisch	$\frac{-4 \ln\left(1 - \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + 2 \ln\left(\tanh\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d - b \left(\operatorname{coth}\left(\frac{bx}{2} + \frac{a}{2}\right)(dx+c) + (dx+c) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) + 2dx\right)}{2b^2}$	75

input $\operatorname{int}((dx+c)*\operatorname{csch}(bx+a)^2, x, \operatorname{method}=_RETURNVERBOSE)$

output $-2*d/b*x - 2*d/b^2*a - 2*(dx+c)/(exp(2*b*x+2*a)-1)/b + d/b^2*\ln(exp(2*b*x+2*a)-1)$

3.7. $\int (c + dx)\operatorname{csch}^2(a + bx) dx$

3.7.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 166 vs. 2(29) = 58.

Time = 0.27 (sec) , antiderivative size = 166, normalized size of antiderivative = 5.72

$$\int (c + dx) \operatorname{csch}^2(a + bx) dx = \frac{2 b d x \cosh (b x + a)^2 + 4 b d x \cosh (b x + a) \sinh (b x + a) + 2 b d x \sinh (b x + a)^2 + 2 b c - (d \cosh (b x + a) \sinh (b x + a))}{b^2 \cosh (b x + a)^2 + 2 b^2 \cosh (b x + a) \sinh (b x + a) + b^2 \sinh (b x + a)^2 - b^2}$$

input `integrate((d*x+c)*csch(b*x+a)^2,x, algorithm="fricas")`

output `-(2*b*d*x*cosh(b*x + a)^2 + 4*b*d*x*cosh(b*x + a)*sinh(b*x + a) + 2*b*d*x*sinh(b*x + a)^2 + 2*b*c - (d*cosh(b*x + a)^2 + 2*d*cosh(b*x + a)*sinh(b*x + a) + d*sinh(b*x + a)^2 - d)*log(2*sinh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a)))/(b^2*cosh(b*x + a)^2 + 2*b^2*cosh(b*x + a)*sinh(b*x + a) + b^2*sinh(b*x + a)^2 - b^2)`

3.7.6 Sympy [F]

$$\int (c + dx) \operatorname{csch}^2(a + bx) dx = \int (c + dx) \operatorname{csch}^2(a + bx) dx$$

input `integrate((d*x+c)*csch(b*x+a)**2,x)`

output `Integral((c + d*x)*csch(a + b*x)**2, x)`

3.7.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 91 vs. 2(29) = 58.

Time = 0.20 (sec) , antiderivative size = 91, normalized size of antiderivative = 3.14

$$\int (c + dx) \operatorname{csch}^2(a + bx) dx = -d \left(\frac{2 x e^{(2 b x + 2 a)}}{b e^{(2 b x + 2 a)} - b} - \frac{\log \left((e^{(b x + a)} + 1) e^{(-a)} \right)}{b^2} - \frac{\log \left((e^{(b x + a)} - 1) e^{(-a)} \right)}{b^2} \right) + \frac{2 c}{b (e^{(-2 b x - 2 a)} - 1)}$$

input `integrate((d*x+c)*csch(b*x+a)^2,x, algorithm="maxima")`

output
$$-d*(2*x*e^{(2*b*x + 2*a)}/(b*e^{(2*b*x + 2*a)} - b) - \log((e^{(b*x + a)} + 1)*e^{(-a)})/b^2 - \log((e^{(b*x + a)} - 1)*e^{(-a)})/b^2) + 2*c/(b*(e^{(-2*b*x - 2*a)} - 1))$$

3.7.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(29) = 58$.

Time = 0.30 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.76

$$\int (c + dx) \operatorname{csch}^2(a + bx) dx = -\frac{2bdxe^{(2bx+2a)} - de^{(2bx+2a)} \log(e^{(2bx+2a)} - 1) + 2bc + d \log(e^{(2bx+2a)} - 1)}{b^2e^{(2bx+2a)} - b^2}$$

input `integrate((d*x+c)*csch(b*x+a)^2,x, algorithm="giac")`

output
$$-(2*b*d*x*e^{(2*b*x + 2*a)} - d*e^{(2*b*x + 2*a)}*\log(e^{(2*b*x + 2*a)} - 1) + 2*b*c + d*\log(e^{(2*b*x + 2*a)} - 1))/(b^2*e^{(2*b*x + 2*a)} - b^2)$$

3.7.9 Mupad [B] (verification not implemented)

Time = 2.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.69

$$\int (c + dx) \operatorname{csch}^2(a + bx) dx = \frac{d \ln(e^{2a} e^{2bx} - 1)}{b^2} - \frac{2(c + dx)}{b(e^{2a+2bx} - 1)} - \frac{2dx}{b}$$

input `int((c + d*x)/sinh(a + b*x)^2,x)`

output
$$(d*\log(\exp(2*a)*\exp(2*b*x) - 1))/b^2 - (2*(c + d*x))/(b*(\exp(2*a + 2*b*x) - 1)) - (2*d*x)/b$$

3.8 $\int \frac{\operatorname{csch}^2(a+bx)}{x} dx$

3.8.1	Optimal result	81
3.8.2	Mathematica [N/A]	81
3.8.3	Rubi [N/A]	82
3.8.4	Maple [N/A] (verified)	83
3.8.5	Fricas [N/A]	83
3.8.6	Sympy [N/A]	84
3.8.7	Maxima [N/A]	84
3.8.8	Giac [N/A]	84
3.8.9	Mupad [N/A]	85

3.8.1 Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\operatorname{csch}^2(a+bx)}{x} dx = \operatorname{Int}\left(\frac{\operatorname{csch}^2(a+bx)}{x}, x\right)$$

output `Unintegrable(csch(b*x+a)^2/x,x)`

3.8.2 Mathematica [N/A]

Not integrable

Time = 6.40 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^2(a+bx)}{x} dx = \int \frac{\operatorname{csch}^2(a+bx)}{x} dx$$

input `Integrate[Csch[a + b*x]^2/x,x]`

output `Integrate[Csch[a + b*x]^2/x, x]`

3.8.3 Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 25, 4680}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\operatorname{csch}^2(a+bx)}{x} dx \\ \downarrow 3042 \\ \int -\frac{\csc(ia+ibx)^2}{x} dx \\ \downarrow 25 \\ -\int \frac{\csc(ia+ibx)^2}{x} dx \\ \downarrow 4680 \\ \int \frac{\operatorname{csch}^2(a+bx)}{x} dx \end{array}$$

input `Int[Csch[a + b*x]^2/x,x]`

output `$Aborted`

3.8.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

```
rule 4680 Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :>
Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], Un
integrable[(c + d*x)^m*Sech[I*(e - Pi/2) + I*f*x]^n, x], (-I)^n*Unintegrabl
e[(c + d*x)^m*Csch[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], Uni
ntegrable[(c + d*x)^m*Sec[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*C
sc[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]
```

3.8.4 Maple [N/A] (verified)

Not integrable

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{csch}(bx+a)^2}{x} dx$$

input `int(csch(b*x+a)^2/x,x)`

output `int(csch(b*x+a)^2/x,x)`

3.8.5 Fracas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^2(a+bx)}{x} dx = \int \frac{\operatorname{csch}(bx+a)^2}{x} dx$$

input `integrate(csch(b*x+a)^2/x,x, algorithm="fracas")`

output `integral(csch(b*x + a)^2/x, x)`

3.8.6 Sympy [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{csch}^2(a + bx)}{x} dx = \int \frac{\operatorname{csch}^2(a + bx)}{x} dx$$

input `integrate(csch(b*x+a)**2/x,x)`output `Integral(csch(a + b*x)**2/x, x)`**3.8.7 Maxima [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 73, normalized size of antiderivative = 6.08

$$\int \frac{\operatorname{csch}^2(a + bx)}{x} dx = \int \frac{\operatorname{csch}(bx + a)^2}{x} dx$$

input `integrate(csch(b*x+a)^2/x,x, algorithm="maxima")`output `-2/(b*x*e^(2*b*x + 2*a) - b*x) + 4*integrate(1/4/(b*x^2*e^(b*x + a) + b*x^2), x) - 4*integrate(1/4/(b*x^2*e^(b*x + a) - b*x^2), x)`**3.8.8 Giac [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^2(a + bx)}{x} dx = \int \frac{\operatorname{csch}(bx + a)^2}{x} dx$$

input `integrate(csch(b*x+a)^2/x,x, algorithm="giac")`output `integrate(csch(b*x + a)^2/x, x)`

3.8. $\int \frac{\operatorname{csch}^2(a+bx)}{x} dx$

3.8.9 Mupad [N/A]

Not integrable

Time = 2.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^2(a + bx)}{x} dx = \int \frac{1}{x \sinh(a + bx)^2} dx$$

input `int(1/(x*sinh(a + b*x)^2),x)`

output `int(1/(x*sinh(a + b*x)^2), x)`

3.9 $\int (c + dx)^3 \operatorname{csch}^3(a + bx) dx$

3.9.1	Optimal result	86
3.9.2	Mathematica [A] (verified)	87
3.9.3	Rubi [C] (verified)	88
3.9.4	Maple [B] (verified)	92
3.9.5	Fricas [B] (verification not implemented)	92
3.9.6	Sympy [F]	93
3.9.7	Maxima [B] (verification not implemented)	94
3.9.8	Giac [F]	95
3.9.9	Mupad [F(-1)]	95

3.9.1 Optimal result

Integrand size = 16, antiderivative size = 256

$$\begin{aligned}
 \int (c + dx)^3 \operatorname{csch}^3(a + bx) dx = & -\frac{6d^2(c + dx)\operatorname{arctanh}(e^{a+bx})}{b^3} \\
 & + \frac{(c + dx)^3 \operatorname{arctanh}(e^{a+bx})}{b} - \frac{3d(c + dx)^2 \operatorname{csch}(a + bx)}{2b^2} \\
 & - \frac{(c + dx)^3 \operatorname{coth}(a + bx)\operatorname{csch}(a + bx)}{2b} \\
 & - \frac{3d^3 \operatorname{PolyLog}(2, -e^{a+bx})}{b^4} \\
 & + \frac{3d(c + dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{2b^2} \\
 & + \frac{3d^3 \operatorname{PolyLog}(2, e^{a+bx})}{b^4} - \frac{3d(c + dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{2b^2} \\
 & - \frac{3d^2(c + dx) \operatorname{PolyLog}(3, -e^{a+bx})}{b^3} \\
 & + \frac{3d^2(c + dx) \operatorname{PolyLog}(3, e^{a+bx})}{b^3} \\
 & + \frac{3d^3 \operatorname{PolyLog}(4, -e^{a+bx})}{b^4} - \frac{3d^3 \operatorname{PolyLog}(4, e^{a+bx})}{b^4}
 \end{aligned}$$

output
$$\begin{aligned} & -6d^2(d*x+c)*\operatorname{arctanh}(\exp(b*x+a))/b^3+(d*x+c)^3*\operatorname{arctanh}(\exp(b*x+a))/b-3/2 \\ & *d*(d*x+c)^2*\operatorname{csch}(b*x+a)/b^2-1/2*(d*x+c)^3*\operatorname{coth}(b*x+a)*\operatorname{csch}(b*x+a)/b-3*d^3 \\ & *polylog(2,-\exp(b*x+a))/b^4+3/2*d*(d*x+c)^2*polylog(2,-\exp(b*x+a))/b^2+3*d \\ & ^3*polylog(2,\exp(b*x+a))/b^4-3/2*d*(d*x+c)^2*polylog(2,\exp(b*x+a))/b^2-3*d \\ & ^2*(d*x+c)*polylog(3,-\exp(b*x+a))/b^3+3*d^2*(d*x+c)*polylog(3,\exp(b*x+a))/ \\ & b^3+3*d^3*polylog(4,-\exp(b*x+a))/b^4-3*d^3*polylog(4,\exp(b*x+a))/b^4 \end{aligned}$$

3.9.2 Mathematica [A] (verified)

Time = 2.10 (sec) , antiderivative size = 440, normalized size of antiderivative = 1.72

$$\int (c + dx)^3 \operatorname{csch}^3(a + bx) dx = \frac{b^2(c + dx)^2(3d + b(c + dx) \operatorname{coth}(a + bx)) \operatorname{csch}(a + bx) + b^3 c^3 \log(1 - e^{a+bx}) - 6bcd^2 \log(1 - e^{a+bx}) + \dots}{\dots}$$

input `Integrate[(c + d*x)^3*Csch[a + b*x]^3,x]`

output
$$\begin{aligned} & -1/2*(b^2*(c + d*x)^2*(3*d + b*(c + d*x)*\operatorname{Coth}[a + b*x])* \operatorname{Csch}[a + b*x] + b^3 \\ & *c^3*\operatorname{Log}[1 - E^{(a + b*x)}] - 6*b*c*d^2*\operatorname{Log}[1 - E^{(a + b*x)}] + 3*b^3*c^2*d*x \\ & *\operatorname{Log}[1 - E^{(a + b*x)}] - 6*b*d^3*x*\operatorname{Log}[1 - E^{(a + b*x)}] + 3*b^3*c*d^2*x^2* \\ & \operatorname{Log}[1 - E^{(a + b*x)}] + b^3*d^3*x^3*\operatorname{Log}[1 - E^{(a + b*x)}] - b^3*c^3*\operatorname{Log}[1 + \\ & E^{(a + b*x)}] + 6*b*c*d^2*\operatorname{Log}[1 + E^{(a + b*x)}] - 3*b^3*c^2*d*x*\operatorname{Log}[1 + E^{(a \\ & + b*x)}] + 6*b*d^3*x*\operatorname{Log}[1 + E^{(a + b*x)}] - 3*b^3*c*d^2*x^2*\operatorname{Log}[1 + E^{(a + \\ & b*x)}] - b^3*d^3*x^3*\operatorname{Log}[1 + E^{(a + b*x)}] - 3*d*(-2*d^2 + b^2*(c + d*x)^2) \\ & *PolyLog[2, -E^{(a + b*x)}] + 3*d*(-2*d^2 + b^2*(c + d*x)^2)*PolyLog[2, E^{(a \\ & + b*x)}] + 6*b*c*d^2*PolyLog[3, -E^{(a + b*x)}] + 6*b*d^3*x*PolyLog[3, -E^{(a \\ & + b*x)}] - 6*b*c*d^2*PolyLog[3, E^{(a + b*x)}] - 6*b*d^3*x*PolyLog[3, E^{(a + \\ & b*x)}] - 6*d^3*PolyLog[4, -E^{(a + b*x)}] + 6*d^3*PolyLog[4, E^{(a + b*x)}])/b \\ & ^4 \end{aligned}$$

3.9.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.20 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.15, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.812$, Rules used = {3042, 26, 4674, 26, 3042, 26, 4670, 2715, 2838, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^3 \operatorname{csch}^3(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i(c + dx)^3 \operatorname{csc}(ia + ibx)^3 dx \\
 & \quad \downarrow \text{26} \\
 & -i \int (c + dx)^3 \operatorname{csc}(ia + ibx)^3 dx \\
 & \quad \downarrow \text{4674} \\
 & -i \left(-\frac{3d^2 \int -i(c + dx) \operatorname{csch}(a + bx) dx}{b^2} + \frac{1}{2} \int -i(c + dx)^3 \operatorname{csch}(a + bx) dx - \frac{3id(c + dx)^2 \operatorname{csch}(a + bx)}{2b^2} - \frac{i(c + dx)}{b} \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left(\frac{3id^2 \int (c + dx) \operatorname{csch}(a + bx) dx}{b^2} - \frac{1}{2} i \int (c + dx)^3 \operatorname{csch}(a + bx) dx - \frac{3id(c + dx)^2 \operatorname{csch}(a + bx)}{2b^2} - \frac{i(c + dx)^3 \operatorname{coth}(a + bx)}{b} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left(\frac{3id^2 \int i(c + dx) \operatorname{csc}(ia + ibx) dx}{b^2} - \frac{1}{2} i \int i(c + dx)^3 \operatorname{csc}(ia + ibx) dx - \frac{3id(c + dx)^2 \operatorname{csch}(a + bx)}{2b^2} - \frac{i(c + dx)^3 \operatorname{coth}(a + bx)}{b} \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left(-\frac{3d^2 \int (c + dx) \operatorname{csc}(ia + ibx) dx}{b^2} + \frac{1}{2} \int (c + dx)^3 \operatorname{csc}(ia + ibx) dx - \frac{3id(c + dx)^2 \operatorname{csch}(a + bx)}{2b^2} - \frac{i(c + dx)^3 \operatorname{coth}(a + bx)}{b} \right) \\
 & \quad \downarrow \text{4670}
 \end{aligned}$$

$$-i \left(-\frac{3d^2 \left(\frac{id \int \log(1-e^{a+bx}) dx}{b} - \frac{id \int \log(1+e^{a+bx}) dx}{b} + \frac{2i(c+dx) \operatorname{arctanh}(e^{a+bx})}{b} \right)}{b^2} + \frac{1}{2} \left(\frac{3id \int (c+dx)^2 \log(1-e^{a+bx}) dx}{b} \right) \right)$$

↓ 2715

$$-i \left(-\frac{3d^2 \left(\frac{id \int e^{-a-bx} \log(1-e^{a+bx}) de^{a+bx}}{b^2} - \frac{id \int e^{-a-bx} \log(1+e^{a+bx}) de^{a+bx}}{b^2} + \frac{2i(c+dx) \operatorname{arctanh}(e^{a+bx})}{b} \right)}{b^2} + \frac{1}{2} \left(\frac{3id \int (c+dx)^2 \log(1-e^{a+bx}) dx}{b} \right) \right)$$

↓ 2838

$$-i \left(\frac{1}{2} \left(\frac{3id \int (c+dx)^2 \log(1-e^{a+bx}) dx}{b} - \frac{3id \int (c+dx)^2 \log(1+e^{a+bx}) dx}{b} + \frac{2i(c+dx)^3 \operatorname{arctanh}(e^{a+bx})}{b} \right) \right)$$

↓ 3011

$$-i \left(\frac{1}{2} \left(-\frac{3id \left(\frac{2d \int (c+dx) \operatorname{PolyLog}(2, -e^{a+bx}) dx}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{3id \left(\frac{2d \int (c+dx) \operatorname{PolyLog}(2, e^{a+bx}) dx}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right) \right)$$

↓ 7163

$$-i \left(\frac{1}{2} \left(\frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{d \int \operatorname{PolyLog}(3, -e^{a+bx}) dx}{b} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{d \int \operatorname{PolyLog}(3, e^{a+bx}) dx}{b} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right) \right)$$

↓ 2720

$$-i \left(\frac{1}{2} \left(\frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{d \int e^{-a-bx} \operatorname{PolyLog}(3, -e^{a+bx}) de^{a+bx}}{b^2} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{3id \left(\frac{2d \left(\frac{(c+dx) \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{d \int e^{-a-bx} \operatorname{PolyLog}(3, e^{a+bx}) de^{a+bx}}{b^2} \right)}{b} - \frac{(c+dx)^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right) \right)$$

↓ 7143

$$-i \left(-\frac{3d^2 \left(\frac{2i(c+dx)\operatorname{arctanh}(e^{a+bx})}{b} + \frac{id \operatorname{PolyLog}(2, -e^{a+bx})}{b^2} - \frac{id \operatorname{PolyLog}(2, e^{a+bx})}{b^2} \right)}{b^2} \right) + \frac{1}{2} \left(\frac{2i(c+dx)^3 \operatorname{arctanh}(e^{a+bx})}{b} \right)$$

input `Int[(c + d*x)^3*Csch[a + b*x]^3,x]`

output `(-I)*((((-3*I)/2)*d*(c + d*x)^2*Csch[a + b*x])/b^2 - ((I/2)*(c + d*x)^3*Coth[a + b*x]*Csch[a + b*x])/b - (3*d^2*(((2*I)*(c + d*x)*ArcTanh[E^(a + b*x)]))/b + (I*d*PolyLog[2, -E^(a + b*x)]/b^2 - (I*d*PolyLog[2, E^(a + b*x)]/b^2))/b^2 + (((2*I)*(c + d*x)^3*ArcTanh[E^(a + b*x)])/b - ((3*I)*d*(-((c + d*x)^2*PolyLog[2, -E^(a + b*x)]/b) + (2*d*(((c + d*x)*PolyLog[3, -E^(a + b*x)])/b - (d*PolyLog[4, -E^(a + b*x)]/b^2))/b))/b + ((3*I)*d*(-((c + d*x)^2*PolyLog[2, E^(a + b*x)]/b) + (2*d*(((c + d*x)*PolyLog[3, E^(a + b*x)])/b - (d*PolyLog[4, E^(a + b*x)]/b^2))/b))/b)/2`

3.9.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^m] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 4674 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_)^(m_), x_Symbol] := Simp[(-b^2)*(c + d*x)^m*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (-Simp[b^2*d*m*(c + d*x)^(m - 1)*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + Simp[b^2*d^2*m*(m - 1)/(f^2*(n - 1)*(n - 2)) Int[(c + d*x)^(m - 2)*(b*Csc[e + f*x])^(n - 2), x], x] + Simp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)^m*(b*Csc[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_)^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

3.9.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 875 vs. $2(238) = 476$.

Time = 1.01 (sec) , antiderivative size = 876, normalized size of antiderivative = 3.42

method	result	size
risch	Expression too large to display	876

```
input int((d*x+c)^3*csch(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
output 1/b*c^3*arctanh(exp(b*x+a))+6/b^4*d^3*a*arctanh(exp(b*x+a))+3/b^3*d^3*ln(1
-exp(b*x+a))*x+3/b^4*d^3*ln(1-exp(b*x+a))*a-3/b^3*d^3*ln(exp(b*x+a)+1)*x-3
/b^4*d^3*ln(exp(b*x+a)+1)*a-6/b^3*d^2*c*arctanh(exp(b*x+a))+3/2/b^2*d^3*po
lylog(2,-exp(b*x+a))*x^2-3/b^3*d^3*polylog(3,-exp(b*x+a))*x-1/2/b*d^3*ln(1
-exp(b*x+a))*x^3-1/2/b^4*d^3*ln(1-exp(b*x+a))*a^3-3/2/b^2*d^3*polylog(2,ex
p(b*x+a))*x^2+3/b^3*d^3*polylog(3,exp(b*x+a))*x+1/2/b*d^3*ln(exp(b*x+a)+1)
*x^3+1/2/b^4*d^3*ln(exp(b*x+a)+1)*a^3-1/b^4*d^3*a^3*arctanh(exp(b*x+a))-3/
b^3*c*d^2*polylog(3,-exp(b*x+a))-3/2/b^2*d*c^2*polylog(2,exp(b*x+a))+3/2/b
^2*d*c^2*polylog(2,-exp(b*x+a))+3/b^3*c*d^2*polylog(3,exp(b*x+a))-3/b^2*d*
a*c^2*arctanh(exp(b*x+a))+3/2/b*d*c^2*ln(exp(b*x+a)+1)*x+3/2/b^2*d*c^2*ln(
exp(b*x+a)+1)*a+3/b^3*d^2*a^2*c*arctanh(exp(b*x+a))-3/2/b*c*d^2*ln(1-exp(b
*x+a))*x^2+3/2/b^3*c*d^2*ln(1-exp(b*x+a))*a^2-3/b^2*c*d^2*polylog(2,exp(b*
x+a))*x+3/2/b*c*d^2*ln(exp(b*x+a)+1)*x^2-3/2/b^3*c*d^2*ln(exp(b*x+a)+1)*a^
2+3/b^2*c*d^2*polylog(2,-exp(b*x+a))*x-3/2/b*d*c^2*ln(1-exp(b*x+a))*x-3/2/
b^2*d*c^2*ln(1-exp(b*x+a))*a-exp(b*x+a)*(exp(2*b*x+2*a)*b*d^3*x^3+3*exp(2*
b*x+2*a)*b*c*d^2*x^2+3*exp(2*b*x+2*a)*b*c^2*d*x+b*d^3*x^3+3*exp(2*b*x+2*a)
*d^3*x^2+exp(2*b*x+2*a)*b*c^3+3*b*c*d^2*x^2+6*exp(2*b*x+2*a)*c*d^2*x+3*b*c
^2*d*x+3*exp(2*b*x+2*a)*c^2*d-3*d^3*x^2+b*c^3-6*c*d^2*x-3*d*c^2)/b^2/(exp(
2*b*x+2*a)-1)^2-3*d^3*polylog(4,exp(b*x+a))/b^4+3*d^3*polylog(4,-exp(b*x+a)
)/b^4-3*d^3*polylog(2,-exp(b*x+a))/b^4+3*d^3*polylog(2,exp(b*x+a))/b^4
```

3.9.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4008 vs. $2(234) = 468$.

Time = 0.31 (sec) , antiderivative size = 4008, normalized size of antiderivative = 15.66

$$\int (c + dx)^3 \operatorname{csch}^3(a + bx) dx = \text{Too large to display}$$

```
input integrate((d*x+c)^3*csch(b*x+a)^3,x, algorithm="fricas")
```

```

output -1/2*(2*(b^3*d^3*x^3 + b^3*c^3 + 3*b^2*c^2*d + 3*(b^3*c*d^2 + b^2*d^3)*x^2
+ 3*(b^3*c^2*d + 2*b^2*c*d^2)*x)*cosh(b*x + a)^3 + 6*(b^3*d^3*x^3 + b^3*c
^3 + 3*b^2*c^2*d + 3*(b^3*c*d^2 + b^2*d^3)*x^2 + 3*(b^3*c^2*d + 2*b^2*c*d
^2)*x)*cosh(b*x + a)*sinh(b*x + a)^2 + 2*(b^3*d^3*x^3 + b^3*c^3 + 3*b^2*c^2
*d + 3*(b^3*c*d^2 + b^2*d^3)*x^2 + 3*(b^3*c^2*d + 2*b^2*c*d^2)*x)*sinh(b*x
+ a)^3 + 2*(b^3*d^3*x^3 + b^3*c^3 - 3*b^2*c^2*d + 3*(b^3*c*d^2 - b^2*d^3)
*x^2 + 3*(b^3*c^2*d - 2*b^2*c*d^2)*x)*cosh(b*x + a) + 3*(b^2*d^3*x^2 + 2*b
^2*c*d^2*x + b^2*c^2*d + (b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*d^3)
*cosh(b*x + a)^4 + 4*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*d^3)*cos
h(b*x + a)*sinh(b*x + a)^3 + (b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*
d^3)*sinh(b*x + a)^4 - 2*d^3 - 2*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d
- 2*d^3)*cosh(b*x + a)^2 - 2*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*
d^3 - 3*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*d^3)*cosh(b*x + a)^2)
*sinh(b*x + a)^2 + 4*((b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*d^3)*co
sh(b*x + a)^3 - (b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*d^3)*cosh(b*x
+ a))*sinh(b*x + a))*dilog(cosh(b*x + a) + sinh(b*x + a)) - 3*(b^2*d^3*x^
2 + 2*b^2*c*d^2*x + b^2*c^2*d + (b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d -
2*d^3)*cosh(b*x + a)^4 + 4*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2*d - 2*d
^3)*cosh(b*x + a)*sinh(b*x + a)^3 + (b^2*d^3*x^2 + 2*b^2*c*d^2*x + b^2*c^2
*d - 2*d^3)*sinh(b*x + a)^4 - 2*d^3 - 2*(b^2*d^3*x^2 + 2*b^2*c*d^2*x + ...

```

3.9.6 Sympy [F]

$$\int (c + dx)^3 \operatorname{csch}^3(a + bx) dx = \int (c + dx)^3 \operatorname{csch}^3(a + bx) dx$$

```
input integrate((d*x+c)**3*csch(b*x+a)**3,x)
```

```
output Integral((c + d*x)**3*csch(a + b*x)**3, x)
```

3.9.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 605 vs. $2(234) = 468$.

Time = 0.37 (sec) , antiderivative size = 605, normalized size of antiderivative = 2.36

$$\begin{aligned}
& \int (c + dx)^3 \operatorname{csch}^3(a + bx) dx \\
&= \frac{1}{2} c^3 \left(\frac{\log(e^{-bx-a}) + 1}{b} - \frac{\log(e^{-bx-a}) - 1}{b} + \frac{2(e^{-bx-a} + e^{-3bx-3a})}{b(2e^{-2bx-2a} - e^{-4bx-4a} - 1)} \right) \\
&+ \frac{3(b^2 x^2 \log(e^{(bx+a)} + 1) + 2bx \operatorname{Li}_2(-e^{(bx+a)}) - 2 \operatorname{Li}_3(-e^{(bx+a)})) cd^2}{2b^3} \\
&- \frac{3(b^2 x^2 \log(-e^{(bx+a)} + 1) + 2bx \operatorname{Li}_2(e^{(bx+a)}) - 2 \operatorname{Li}_3(e^{(bx+a)})) cd^2}{2b^3} \\
&- \frac{3cd^2 \log(e^{(bx+a)} + 1)}{b^3} + \frac{3cd^2 \log(e^{(bx+a)} - 1)}{b^3} \\
&- \frac{(bd^3 x^3 e^{(3a)} + 3c^2 d e^{(3a)} + 3(bcd^2 + d^3)x^2 e^{(3a)} + 3(bc^2 d + 2cd^2)xe^{(3a)})e^{(3bx)} + (bd^3 x^3 e^a - 3c^2 d e^a + 3c^2 d e^a)}{b^2 e^{(4bx+4a)} - 2b^2 e^{(2bx+2a)} + b^2} \\
&+ \frac{(b^3 x^3 \log(e^{(bx+a)} + 1) + 3b^2 x^2 \operatorname{Li}_2(-e^{(bx+a)}) - 6bx \operatorname{Li}_3(-e^{(bx+a)}) + 6 \operatorname{Li}_4(-e^{(bx+a)})) d^3}{2b^4} \\
&- \frac{(b^3 x^3 \log(-e^{(bx+a)} + 1) + 3b^2 x^2 \operatorname{Li}_2(e^{(bx+a)}) - 6bx \operatorname{Li}_3(e^{(bx+a)}) + 6 \operatorname{Li}_4(e^{(bx+a)})) d^3}{2b^4} \\
&+ \frac{3(b^2 c^2 d - 2d^3)(bx \log(e^{(bx+a)} + 1) + \operatorname{Li}_2(-e^{(bx+a)}))}{2b^4} \\
&- \frac{3(b^2 c^2 d - 2d^3)(bx \log(-e^{(bx+a)} + 1) + \operatorname{Li}_2(e^{(bx+a)}))}{2b^4}
\end{aligned}$$

input `integrate((d*x+c)^3*csch(b*x+a)^3,x, algorithm="maxima")`

output $\frac{1}{2}c^3(\log(e^{-bx-a})+1)/b - \log(e^{-bx-a}-1)/b + 2(e^{-bx-a} + e^{-3bx-3a})/(b(2e^{-2bx-2a} - e^{-4bx-4a} - 1)) + 3/2(b^2x^2\log(e^{bx+a})+1) + 2bx\operatorname{dilog}(-e^{bx+a}) - 2\operatorname{polylog}(3, -e^{bx+a}))cd^2/b^3 - 3/2(b^2x^2\log(-e^{bx+a})+1) + 2bx\operatorname{dilog}(e^{bx+a}) - 2\operatorname{polylog}(3, e^{bx+a}))cd^2/b^3 - 3cd^2\log(e^{bx+a})/b^3 + 3cd^2\log(e^{bx+a}-1)/b^3 - ((bd^3x^3e^{3a} + 3c^2de^{3a} + 3(bcd^2+d^3)x^2e^{3a} + 3(bc^2d+2cd^2)xe^{3a})e^{3bx} + (bd^3x^3e^a - 3c^2de^a + 3(bcd^2-d^3)x^2e^a + 3(bc^2d-2cd^2)xe^a)e^{bx})/(b^2e^{4bx+4a} - 2b^2e^{2bx+2a} + b^2) + 1/2(b^3x^3\log(e^{bx+a})+1) + 3b^2x^2\operatorname{dilog}(-e^{bx+a}) - 6bx\operatorname{polylog}(3, -e^{bx+a}) + 6\operatorname{polylog}(4, -e^{bx+a}))d^3/b^4 - 1/2(b^3x^3\log(-e^{bx+a})+1) + 3b^2x^2\operatorname{dilog}(e^{bx+a}) - 6bx\operatorname{polylog}(3, e^{bx+a}) + 6\operatorname{polylog}(4, e^{bx+a}))d^3/b^4 + 3/2(b^2c^2d - 2d^3)(bx\log(e^{bx+a})+1) + \operatorname{dilog}(-e^{bx+a})/b^4 - 3/2(b^2c^2d - 2d^3)(bx\log(-e^{bx+a})+1) + \operatorname{dilog}(e^{bx+a})/b^4$

3.9.8 Giac [F]

$$\int (c+dx)^3 \operatorname{csch}^3(a+bx) dx = \int (dx+c)^3 \operatorname{csch}(bx+a)^3 dx$$

input `integrate((d*x+c)^3*csch(b*x+a)^3,x, algorithm="giac")`

output `integrate((d*x + c)^3*csch(b*x + a)^3, x)`

3.9.9 Mupad [F(-1)]

Timed out.

$$\int (c+dx)^3 \operatorname{csch}^3(a+bx) dx = \int \frac{(c+dx)^3}{\sinh(a+bx)^3} dx$$

input `int((c + d*x)^3/sinh(a + b*x)^3,x)`

output `int((c + d*x)^3/sinh(a + b*x)^3, x)`

3.10 $\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx$

3.10.1	Optimal result	96
3.10.2	Mathematica [B] (verified)	97
3.10.3	Rubi [C] (verified)	97
3.10.4	Maple [B] (verified)	101
3.10.5	Fricas [B] (verification not implemented)	101
3.10.6	Sympy [F]	102
3.10.7	Maxima [B] (verification not implemented)	103
3.10.8	Giac [F]	104
3.10.9	Mupad [F(-1)]	104

3.10.1 Optimal result

Integrand size = 16, antiderivative size = 154

$$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx = \frac{(c + dx)^2 \operatorname{arctanh}(e^{a+bx})}{b} - \frac{d^2 \operatorname{arctanh}(\cosh(a + bx))}{b^3} - \frac{d(c + dx) \operatorname{csch}(a + bx)}{b^2} - \frac{(c + dx)^2 \operatorname{coth}(a + bx) \operatorname{csch}(a + bx)}{2b} + \frac{d(c + dx) \operatorname{PolyLog}(2, -e^{a+bx})}{b^2} - \frac{d(c + dx) \operatorname{PolyLog}(2, e^{a+bx})}{b^2} - \frac{d^2 \operatorname{PolyLog}(3, -e^{a+bx})}{b^3} + \frac{d^2 \operatorname{PolyLog}(3, e^{a+bx})}{b^3}$$

output $(d*x+c)^2*\operatorname{arctanh}(\exp(b*x+a))/b-d^2*\operatorname{arctanh}(\cosh(b*x+a))/b^3-d*(d*x+c)*\operatorname{csch}(b*x+a)/b^2-1/2*(d*x+c)^2*\operatorname{coth}(b*x+a)*\operatorname{csch}(b*x+a)/b+d*(d*x+c)*\operatorname{polylog}(2,-\exp(b*x+a))/b^2-d*(d*x+c)*\operatorname{polylog}(2,\exp(b*x+a))/b^2-d^2*\operatorname{polylog}(3,-\exp(b*x+a))/b^3+d^2*\operatorname{polylog}(3,\exp(b*x+a))/b^3$

3.10.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 420 vs. $2(154) = 308$.

Time = 6.24 (sec) , antiderivative size = 420, normalized size of antiderivative = 2.73

$$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx = -\frac{d(c + dx)\operatorname{csch}(a)}{b^2} + \frac{(-c^2 - 2cdx - d^2x^2)\operatorname{csch}^2\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b}$$

$$+ \frac{-b^2c^2 \log(1 - e^{a+bx}) + 2d^2 \log(1 - e^{a+bx}) - 2b^2cdx \log(1 - e^{a+bx}) - b^2d^2x^2 \log(1 - e^{a+bx}) + b^2c^2 \log(1 + e^{a+bx})}{8b}$$

$$+ \frac{(-c^2 - 2cdx - d^2x^2)\operatorname{sech}^2\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b}$$

$$+ \frac{\operatorname{csch}\left(\frac{a}{2}\right)\operatorname{csch}\left(\frac{a}{2} + \frac{bx}{2}\right)(cd \sinh\left(\frac{bx}{2}\right) + d^2x \sinh\left(\frac{bx}{2}\right))}{2b^2}$$

$$+ \frac{\operatorname{sech}\left(\frac{a}{2}\right)\operatorname{sech}\left(\frac{a}{2} + \frac{bx}{2}\right)(cd \sinh\left(\frac{bx}{2}\right) + d^2x \sinh\left(\frac{bx}{2}\right))}{2b^2}$$

input `Integrate[(c + d*x)^2*Csch[a + b*x]^3,x]`

output

```

-((d*(c + d*x)*Csch[a])/b^2) + ((-c^2 - 2*c*d*x - d^2*x^2)*Csch[a/2 + (b*x)/2]^2)/(8*b) + (-b^2*c^2*Log[1 - E^(a + b*x)]) + 2*d^2*Log[1 - E^(a + b*x)] - 2*b^2*c*d*x*Log[1 - E^(a + b*x)] - b^2*d^2*x^2*Log[1 - E^(a + b*x)] + b^2*c^2*Log[1 + E^(a + b*x)] - 2*d^2*Log[1 + E^(a + b*x)] + 2*b^2*c*d*x*Log[1 + E^(a + b*x)] + b^2*d^2*x^2*Log[1 + E^(a + b*x)] + 2*b*d*(c + d*x)*PolyLog[2, -E^(a + b*x)] - 2*b*d*(c + d*x)*PolyLog[2, E^(a + b*x)] - 2*d^2*PolyLog[3, -E^(a + b*x)] + 2*d^2*PolyLog[3, E^(a + b*x)]/(2*b^3) + ((-c^2 - 2*c*d*x - d^2*x^2)*Sech[a/2 + (b*x)/2]^2)/(8*b) + (Csch[a/2]*Csch[a/2 + (b*x)/2]*(c*d*Sinh[(b*x)/2] + d^2*x*Sinh[(b*x)/2]))/(2*b^2) + (Sech[a/2]*Sech[a/2 + (b*x)/2]*(c*d*Sinh[(b*x)/2] + d^2*x*Sinh[(b*x)/2]))/(2*b^2)

```

3.10.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.19, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {3042, 26, 4674, 26, 3042, 26, 4257, 4670, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.10. $\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx$

$$\begin{aligned}
& \int (c + dx)^2 \operatorname{csch}^3(a + bx) dx \\
& \quad \downarrow \text{3042} \\
& \int -i(c + dx)^2 \operatorname{csc}(ia + ibx)^3 dx \\
& \quad \downarrow \text{26} \\
& -i \int (c + dx)^2 \operatorname{csc}(ia + ibx)^3 dx \\
& \quad \downarrow \text{4674} \\
& -i \left(-\frac{d^2 \int -i \operatorname{csch}(a + bx) dx}{b^2} + \frac{1}{2} \int -i(c + dx)^2 \operatorname{csch}(a + bx) dx - \frac{id(c + dx) \operatorname{csch}(a + bx)}{b^2} - \frac{i(c + dx)^2 \operatorname{coth}(a + bx)}{2b} \right) \\
& \quad \downarrow \text{26} \\
& -i \left(\frac{id^2 \int \operatorname{csch}(a + bx) dx}{b^2} - \frac{1}{2} i \int (c + dx)^2 \operatorname{csch}(a + bx) dx - \frac{id(c + dx) \operatorname{csch}(a + bx)}{b^2} - \frac{i(c + dx)^2 \operatorname{coth}(a + bx) \operatorname{csch}(a + bx)}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& -i \left(\frac{id^2 \int i \operatorname{csc}(ia + ibx) dx}{b^2} - \frac{1}{2} i \int i(c + dx)^2 \operatorname{csc}(ia + ibx) dx - \frac{id(c + dx) \operatorname{csch}(a + bx)}{b^2} - \frac{i(c + dx)^2 \operatorname{coth}(a + bx)}{2b} \right) \\
& \quad \downarrow \text{26} \\
& -i \left(-\frac{d^2 \int \operatorname{csc}(ia + ibx) dx}{b^2} + \frac{1}{2} \int (c + dx)^2 \operatorname{csc}(ia + ibx) dx - \frac{id(c + dx) \operatorname{csch}(a + bx)}{b^2} - \frac{i(c + dx)^2 \operatorname{coth}(a + bx) \operatorname{csch}(a + bx)}{2b} \right) \\
& \quad \downarrow \text{4257} \\
& -i \left(\frac{1}{2} \int (c + dx)^2 \operatorname{csc}(ia + ibx) dx - \frac{id^2 \operatorname{arctanh}(\operatorname{cosh}(a + bx))}{b^3} - \frac{id(c + dx) \operatorname{csch}(a + bx)}{b^2} - \frac{i(c + dx)^2 \operatorname{coth}(a + bx)}{2b} \right) \\
& \quad \downarrow \text{4670} \\
& -i \left(\frac{1}{2} \left(\frac{2id \int (c + dx) \log(1 - e^{a+bx}) dx}{b} - \frac{2id \int (c + dx) \log(1 + e^{a+bx}) dx}{b} + \frac{2i(c + dx)^2 \operatorname{arctanh}(e^{a+bx})}{b} \right) - \frac{id^2 \operatorname{arctanh}(\operatorname{cosh}(a + bx))}{b^3} - \frac{id(c + dx) \operatorname{csch}(a + bx)}{b^2} - \frac{i(c + dx)^2 \operatorname{coth}(a + bx)}{2b} \right) \\
& \quad \downarrow \text{3011}
\end{aligned}$$

$$\begin{aligned}
& -i \left(\frac{1}{2} \left(-\frac{2id \left(\frac{d \int \text{PolyLog}(2, -e^{a+bx}) dx}{b} - \frac{(c+dx) \text{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{2id \left(\frac{d \int \text{PolyLog}(2, e^{a+bx}) dx}{b} - \frac{(c+dx) \text{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right) \right) \\
& \quad \downarrow \text{2720} \\
& -i \left(\frac{1}{2} \left(-\frac{2id \left(\frac{d \int e^{-a-bx} \text{PolyLog}(2, -e^{a+bx}) de^{a+bx}}{b^2} - \frac{(c+dx) \text{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} + \frac{2id \left(\frac{d \int e^{-a-bx} \text{PolyLog}(2, e^{a+bx}) de^{a+bx}}{b^2} - \frac{(c+dx) \text{PolyLog}(2, e^{a+bx})}{b} \right)}{b} \right) \right) \\
& \quad \downarrow \text{7143} \\
& -i \left(-\frac{id^2 \operatorname{arctanh}(\cosh(a+bx))}{b^3} + \frac{1}{2} \left(\frac{2i(c+dx)^2 \operatorname{arctanh}(e^{a+bx})}{b} - \frac{2id \left(\frac{d \text{PolyLog}(3, -e^{a+bx})}{b^2} - \frac{(c+dx) \text{PolyLog}(2, -e^{a+bx})}{b} \right)}{b} \right) \right)
\end{aligned}$$

input `Int[(c + d*x)^2*Csch[a + b*x]^3,x]`

output `(-I)*(((-I)*d^2*ArcTanh[Cosh[a + b*x]])/b^3 - (I*d*(c + d*x)*Csch[a + b*x])/b^2 - ((I/2)*(c + d*x)^2*Coth[a + b*x]*Csch[a + b*x])/b + (((2*I)*(c + d*x)^2*ArcTanh[E^(a + b*x)])/b - ((2*I)*d*(-(((c + d*x)*PolyLog[2, -E^(a + b*x)])/b) + (d*PolyLog[3, -E^(a + b*x)]/b^2))/b + ((2*I)*d*(-(((c + d*x)*PolyLog[2, E^(a + b*x)])/b) + (d*PolyLog[3, E^(a + b*x)]/b^2))/b)/2`

3.10.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 4674 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_))^(m_), x_Symbol] := Simp[(-b^2)*(c + d*x)^m*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (-Simp[b^2*d*m*(c + d*x)^(m - 1)*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + Simp[b^2*d^2*m*(m - 1)/(f^2*(n - 1)*(n - 2)) Int[(c + d*x)^(m - 2)*(b*Csc[e + f*x])^(n - 2), x], x] + Simp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)^m*(b*Csc[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

3.10.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 443 vs. $2(147) = 294$.

Time = 0.94 (sec) , antiderivative size = 444, normalized size of antiderivative = 2.88

method	result
risch	$-\frac{e^{bx+a}(e^{2bx+2a}bd^2x^2+2e^{2bx+2a}bcdx+e^{2bx+2a}b^2c^2+bd^2x^2+2e^{2bx+2a}d^2x+2bcdx+2e^{2bx+2a}cd+c^2b-2xd^2-2cd)}{b^2(e^{2bx+2a}-1)^2} + \frac{d^2 \ln(1-e^{bx+a})}{2b^3}$

input `int((d*x+c)^2*cscsch(b*x+a)^3,x,method=_RETURNVERBOSE)`

output

```
-exp(b*x+a)*(exp(2*b*x+2*a)*b*d^2*x^2+2*exp(2*b*x+2*a)*b*c*d*x+exp(2*b*x+2*a)*b*c^2+b*d^2*x^2+2*exp(2*b*x+2*a)*d^2*x+2*b*c*d*x+2*exp(2*b*x+2*a)*c*d+c^2*b-2*x*d^2-2*c*d)/b^2/(exp(2*b*x+2*a)-1)^2+1/2/b^3*d^2*ln(1-exp(b*x+a))*a^2-1/2/b^3*d^2*ln(exp(b*x+a)+1)*a^2+1/b^3*a^2*d^2*arctanh(exp(b*x+a))+1/b*c^2*arctanh(exp(b*x+a))-2/b^3*d^2*arctanh(exp(b*x+a))-2/b^2*a*c*d*arctanh(exp(b*x+a))-1/b*c*d*ln(1-exp(b*x+a))*x+d^2*polylog(3,exp(b*x+a))/b^3-d^2*polylog(3,-exp(b*x+a))/b^3-1/b^2*c*d*ln(1-exp(b*x+a))*a+1/b*c*d*ln(exp(b*x+a)+1)*x+1/b^2*c*d*ln(exp(b*x+a)+1)*a-1/2/b*d^2*ln(1-exp(b*x+a))*x^2-1/b^2*d^2*polylog(2,exp(b*x+a))*x+1/2/b*d^2*ln(exp(b*x+a)+1)*x^2+1/b^2*d^2*polylog(2,-exp(b*x+a))*x-1/b^2*c*d*polylog(2,exp(b*x+a))+1/b^2*c*d*polylog(2,-exp(b*x+a))
```

3.10.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2218 vs. $2(145) = 290$.

Time = 0.29 (sec) , antiderivative size = 2218, normalized size of antiderivative = 14.40

$$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx = \text{Too large to display}$$

input `integrate((d*x+c)^2*cscsch(b*x+a)^3,x, algorithm="fracas")`

output

```
-1/2*(2*(b^2*d^2*x^2 + b^2*c^2 + 2*b*c*d + 2*(b^2*c*d + b*d^2)*x)*cosh(b*x
+ a)^3 + 6*(b^2*d^2*x^2 + b^2*c^2 + 2*b*c*d + 2*(b^2*c*d + b*d^2)*x)*cosh
(b*x + a)*sinh(b*x + a)^2 + 2*(b^2*d^2*x^2 + b^2*c^2 + 2*b*c*d + 2*(b^2*c*
d + b*d^2)*x)*sinh(b*x + a)^3 + 2*(b^2*d^2*x^2 + b^2*c^2 - 2*b*c*d + 2*(b^
2*c*d - b*d^2)*x)*cosh(b*x + a) + 2*((b*d^2*x + b*c*d)*cosh(b*x + a)^4 + 4
*(b*d^2*x + b*c*d)*cosh(b*x + a)*sinh(b*x + a)^3 + (b*d^2*x + b*c*d)*sinh(
b*x + a)^4 + b*d^2*x + b*c*d - 2*(b*d^2*x + b*c*d)*cosh(b*x + a)^2 - 2*(b*
d^2*x + b*c*d - 3*(b*d^2*x + b*c*d)*cosh(b*x + a)^2)*sinh(b*x + a)^2 + 4*(
(b*d^2*x + b*c*d)*cosh(b*x + a)^3 - (b*d^2*x + b*c*d)*cosh(b*x + a))*sinh(
b*x + a))*dilog(cosh(b*x + a) + sinh(b*x + a)) - 2*((b*d^2*x + b*c*d)*cosh
(b*x + a)^4 + 4*(b*d^2*x + b*c*d)*cosh(b*x + a)*sinh(b*x + a)^3 + (b*d^2*x
+ b*c*d)*sinh(b*x + a)^4 + b*d^2*x + b*c*d - 2*(b*d^2*x + b*c*d)*cosh(b*x
+ a)^2 - 2*(b*d^2*x + b*c*d - 3*(b*d^2*x + b*c*d)*cosh(b*x + a)^2)*sinh(b
*x + a)^2 + 4*((b*d^2*x + b*c*d)*cosh(b*x + a)^3 - (b*d^2*x + b*c*d)*cosh(
b*x + a))*sinh(b*x + a))*dilog(-cosh(b*x + a) - sinh(b*x + a)) - (b^2*d^2*
x^2 + 2*b^2*c*d*x + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2 - 2*d^2)*cosh(b*x
+ a)^4 + 4*(b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2 - 2*d^2)*cosh(b*x + a)*si
nh(b*x + a)^3 + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2 - 2*d^2)*sinh(b*x + a
)^4 + b^2*c^2 - 2*(b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2 - 2*d^2)*cosh(b*x +
a)^2 - 2*(b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2 - 3*(b^2*d^2*x^2 + 2*b^2...
```

3.10.6 Sympy [F]

$$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx = \int (c + dx)^2 \operatorname{csch}^3(a + bx) dx$$

input `integrate((d*x+c)**2*csch(b*x+a)**3,x)`

output `Integral((c + d*x)**2*csch(a + b*x)**3, x)`

3.10.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 393 vs. $2(145) = 290$.

Time = 0.34 (sec) , antiderivative size = 393, normalized size of antiderivative = 2.55

$$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx$$

$$= \frac{1}{2} c^2 \left(\frac{\log(e^{-bx-a} + 1)}{b} - \frac{\log(e^{-bx-a} - 1)}{b} + \frac{2(e^{-bx-a} + e^{-3bx-3a})}{b(2e^{-2bx-2a} - e^{-4bx-4a} - 1)} \right)$$

$$+ \frac{(bx \log(e^{bx+a} + 1) + \operatorname{Li}_2(-e^{bx+a})) cd}{b^2} - \frac{(bx \log(-e^{bx+a} + 1) + \operatorname{Li}_2(e^{bx+a})) cd}{b^2}$$

$$- \frac{(bd^2 x^2 e^{3a} + 2cde^{3a} + 2(bcd + d^2)xe^{3a})e^{3bx} + (bd^2 x^2 e^a - 2cde^a + 2(bcd - d^2)xe^a)e^{bx}}{b^2 e^{4bx+4a} - 2b^2 e^{2bx+2a} + b^2}$$

$$+ \frac{(b^2 x^2 \log(e^{bx+a} + 1) + 2bx \operatorname{Li}_2(-e^{bx+a}) - 2\operatorname{Li}_3(-e^{bx+a})) d^2}{2b^3}$$

$$- \frac{(b^2 x^2 \log(-e^{bx+a} + 1) + 2bx \operatorname{Li}_2(e^{bx+a}) - 2\operatorname{Li}_3(e^{bx+a})) d^2}{2b^3}$$

$$- \frac{d^2 \log(e^{bx+a} + 1)}{b^3} + \frac{d^2 \log(e^{bx+a} - 1)}{b^3}$$

input `integrate((d*x+c)^2*csch(b*x+a)^3,x, algorithm="maxima")`

output `1/2*c^2*(log(e^(-b*x - a) + 1)/b - log(e^(-b*x - a) - 1)/b + 2*(e^(-b*x - a) + e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x - 2*a) - e^(-4*b*x - 4*a) - 1))) + (b*x*log(e^(b*x + a) + 1) + dilog(-e^(b*x + a)))*c*d/b^2 - (b*x*log(-e^(b*x + a) + 1) + dilog(e^(b*x + a)))*c*d/b^2 - ((b*d^2*x^2*e^(3*a) + 2*c*d*e^(3*a) + 2*(b*c*d + d^2)*x*e^(3*a))*e^(3*b*x) + (b*d^2*x^2*e^a - 2*c*d*e^a + 2*(b*c*d - d^2)*x*e^a)*e^(b*x))/(b^2*e^(4*b*x + 4*a) - 2*b^2*e^(2*b*x + 2*a) + b^2) + 1/2*(b^2*x^2*log(e^(b*x + a) + 1) + 2*b*x*dilog(-e^(b*x + a)) - 2*polylog(3, -e^(b*x + a)))*d^2/b^3 - 1/2*(b^2*x^2*log(-e^(b*x + a) + 1) + 2*b*x*dilog(e^(b*x + a)) - 2*polylog(3, e^(b*x + a)))*d^2/b^3 - d^2*log(e^(b*x + a) + 1)/b^3 + d^2*log(e^(b*x + a) - 1)/b^3`

3.10.8 Giac [F]

$$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx = \int (dx + c)^2 \operatorname{csch}(bx + a)^3 dx$$

input `integrate((d*x+c)^2*csh(b*x+a)^3,x, algorithm="giac")`

output `integrate((d*x + c)^2*csh(b*x + a)^3, x)`

3.10.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 \operatorname{csch}^3(a + bx) dx = \int \frac{(c + dx)^2}{\sinh(a + bx)^3} dx$$

input `int((c + d*x)^2/sinh(a + b*x)^3,x)`

output `int((c + d*x)^2/sinh(a + b*x)^3, x)`

3.11 $\int (c + dx)\operatorname{csch}^3(a + bx) dx$

3.11.1	Optimal result	105
3.11.2	Mathematica [B] (verified)	106
3.11.3	Rubi [C] (verified)	106
3.11.4	Maple [B] (verified)	109
3.11.5	Fricas [B] (verification not implemented)	109
3.11.6	Sympy [F]	110
3.11.7	Maxima [F]	111
3.11.8	Giac [F]	111
3.11.9	Mupad [F(-1)]	111

3.11.1 Optimal result

Integrand size = 14, antiderivative size = 92

$$\int (c + dx)\operatorname{csch}^3(a + bx) dx = \frac{(c + dx)\operatorname{arctanh}(e^{a+bx})}{b} - \frac{d\operatorname{csch}(a + bx)}{2b^2} - \frac{(c + dx)\operatorname{coth}(a + bx)\operatorname{csch}(a + bx)}{2b} + \frac{d\operatorname{PolyLog}(2, -e^{a+bx})}{2b^2} - \frac{d\operatorname{PolyLog}(2, e^{a+bx})}{2b^2}$$

```
output (d*x+c)*arctanh(exp(b*x+a))/b-1/2*d*csch(b*x+a)/b^2-1/2*(d*x+c)*coth(b*x+a)
)*csch(b*x+a)/b+1/2*d*polylog(2,-exp(b*x+a))/b^2-1/2*d*polylog(2,exp(b*x+a)
))/b^2
```

3.11.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 271 vs. $2(92) = 184$.

Time = 0.08 (sec) , antiderivative size = 271, normalized size of antiderivative = 2.95

$$\int (c + dx)\operatorname{csch}^3(a + bx) dx = -\frac{dx\operatorname{csch}^2\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b} - \frac{c\operatorname{csch}^2\left(\frac{1}{2}(a + bx)\right)}{8b} - \frac{dx \log(1 - e^{a+bx})}{2b}$$

$$+ \frac{dx \log(1 + e^{a+bx})}{2b} + \frac{c \log(\cosh\left(\frac{1}{2}(a + bx)\right))}{2b}$$

$$- \frac{c \log(\sinh\left(\frac{1}{2}(a + bx)\right))}{2b} + \frac{d \operatorname{PolyLog}(2, -e^{a+bx})}{2b^2}$$

$$- \frac{d \operatorname{PolyLog}(2, e^{a+bx})}{2b^2} - \frac{dx \operatorname{sech}^2\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b}$$

$$- \frac{c \operatorname{sech}^2\left(\frac{1}{2}(a + bx)\right)}{8b} + \frac{d \operatorname{csch}\left(\frac{a}{2}\right) \operatorname{csch}\left(\frac{a}{2} + \frac{bx}{2}\right) \sinh\left(\frac{bx}{2}\right)}{4b^2}$$

$$+ \frac{d \operatorname{sech}\left(\frac{a}{2}\right) \operatorname{sech}\left(\frac{a}{2} + \frac{bx}{2}\right) \sinh\left(\frac{bx}{2}\right)}{4b^2}$$

input `Integrate[(c + d*x)*Csch[a + b*x]^3,x]`

output `-1/8*(d*x*Csch[a/2 + (b*x)/2]^2)/b - (c*Csch[(a + b*x)/2]^2)/(8*b) - (d*x*Log[1 - E^(a + b*x)])/(2*b) + (d*x*Log[1 + E^(a + b*x)])/(2*b) + (c*Log[Cosh[(a + b*x)/2]])/(2*b) - (c*Log[Sinh[(a + b*x)/2]])/(2*b) + (d*PolyLog[2, -E^(a + b*x)])/(2*b^2) - (d*PolyLog[2, E^(a + b*x)])/(2*b^2) - (d*x*Sech[a/2 + (b*x)/2]^2)/(8*b) - (c*Sech[(a + b*x)/2]^2)/(8*b) + (d*Csch[a/2]*Csch[a/2 + (b*x)/2]*Sinh[(b*x)/2])/(4*b^2) + (d*Sech[a/2]*Sech[a/2 + (b*x)/2]*Sinh[(b*x)/2])/(4*b^2)`

3.11.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.17, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {3042, 26, 4673, 26, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)\operatorname{csch}^3(a + bx) dx$$

$$\begin{aligned}
& \int -i(c+dx) \csc(ia+ibx)^3 dx \\
& \quad \downarrow \text{3042} \\
& \int -i(c+dx) \csc(ia+ibx)^3 dx \\
& \quad \downarrow \text{26} \\
& -i \int (c+dx) \csc(ia+ibx)^3 dx \\
& \quad \downarrow \text{4673} \\
& -i \left(\frac{1}{2} \int -i(c+dx) \operatorname{csch}(a+bx) dx - \frac{id \operatorname{csch}(a+bx)}{2b^2} - \frac{i(c+dx) \operatorname{coth}(a+bx) \operatorname{csch}(a+bx)}{2b} \right) \\
& \quad \downarrow \text{26} \\
& -i \left(-\frac{1}{2} \int (c+dx) \operatorname{csch}(a+bx) dx - \frac{id \operatorname{csch}(a+bx)}{2b^2} - \frac{i(c+dx) \operatorname{coth}(a+bx) \operatorname{csch}(a+bx)}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& -i \left(-\frac{1}{2} i \int i(c+dx) \csc(ia+ibx) dx - \frac{id \operatorname{csch}(a+bx)}{2b^2} - \frac{i(c+dx) \operatorname{coth}(a+bx) \operatorname{csch}(a+bx)}{2b} \right) \\
& \quad \downarrow \text{26} \\
& -i \left(\frac{1}{2} \int (c+dx) \csc(ia+ibx) dx - \frac{id \operatorname{csch}(a+bx)}{2b^2} - \frac{i(c+dx) \operatorname{coth}(a+bx) \operatorname{csch}(a+bx)}{2b} \right) \\
& \quad \downarrow \text{4670} \\
& -i \left(\frac{1}{2} \left(\frac{id \int \log(1-e^{a+bx}) dx}{b} - \frac{id \int \log(1+e^{a+bx}) dx}{b} + \frac{2i(c+dx) \operatorname{arctanh}(e^{a+bx})}{b} \right) - \frac{id \operatorname{csch}(a+bx)}{2b^2} - \frac{i(c+dx) \operatorname{coth}(a+bx) \operatorname{csch}(a+bx)}{2b} \right) \\
& \quad \downarrow \text{2715} \\
& -i \left(\frac{1}{2} \left(\frac{id \int e^{-a-bx} \log(1-e^{a+bx}) de^{a+bx}}{b^2} - \frac{id \int e^{-a-bx} \log(1+e^{a+bx}) de^{a+bx}}{b^2} + \frac{2i(c+dx) \operatorname{arctanh}(e^{a+bx})}{b} \right) - \frac{id \operatorname{csch}(a+bx)}{2b^2} - \frac{i(c+dx) \operatorname{coth}(a+bx) \operatorname{csch}(a+bx)}{2b} \right) \\
& \quad \downarrow \text{2838} \\
& -i \left(\frac{1}{2} \left(\frac{2i(c+dx) \operatorname{arctanh}(e^{a+bx})}{b} + \frac{id \operatorname{PolyLog}(2, -e^{a+bx})}{b^2} - \frac{id \operatorname{PolyLog}(2, e^{a+bx})}{b^2} \right) - \frac{id \operatorname{csch}(a+bx)}{2b^2} - \frac{i(c+dx) \operatorname{coth}(a+bx) \operatorname{csch}(a+bx)}{2b} \right)
\end{aligned}$$

input `Int[(c + d*x)*Csch[a + b*x]^3, x]`

```
output (-I)*(((1/2*I)*d*Csch[a + b*x])/b^2 - ((I/2)*(c + d*x)*Coth[a + b*x]*Csch
[a + b*x])/b + (((2*I)*(c + d*x)*ArcTanh[E^(a + b*x)])/b + (I*d*PolyLog[2,
-E^(a + b*x)])/b^2 - (I*d*PolyLog[2, E^(a + b*x)])/b^2)/2)
```

3.11.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[F_x_, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 2715 Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4670 Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

```
rule 4673 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_)), x_Symbol] :=
Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))),
x] + (-Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x] + S
imp[b^2*((n - 2)/(n - 1)) Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x])
/; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]
```

3.11.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(81) = 162.

Time = 0.77 (sec) , antiderivative size = 197, normalized size of antiderivative = 2.14

method	result
risch	$-\frac{e^{bx+a}(e^{2bx+2a}bdx+e^{2bx+2a}bc+dxbe^{2bx+2a}d+cb-d)}{b^2(e^{2bx+2a}-1)^2} + \frac{c \operatorname{arctanh}(e^{bx+a})}{b} - \frac{d \ln(1-e^{bx+a})x}{2b} - \frac{d \ln(1-e^{bx+a})a}{2b^2} - \frac{d \operatorname{polylog}(2, \exp(bx+a))}{b^2}$

input `int((d*x+c)*csch(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `-exp(b*x+a)*(exp(2*b*x+2*a)*b*d*x+exp(2*b*x+2*a)*b*c+d*x*b+exp(2*b*x+2*a)*d+c*b-d)/b^2/(exp(2*b*x+2*a)-1)^2+1/b*c*arctanh(exp(b*x+a))-1/2/b*d*ln(1-exp(b*x+a))*x-1/2/b^2*d*ln(1-exp(b*x+a))*a-1/2*d*polylog(2,exp(b*x+a))/b^2+1/2/b*d*ln(exp(b*x+a)+1)*x+1/2/b^2*d*ln(exp(b*x+a)+1)*a+1/2*d*polylog(2,-exp(b*x+a))/b^2-1/b^2*d*a*arctanh(exp(b*x+a))`

3.11.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1026 vs. 2(79) = 158.

Time = 0.26 (sec) , antiderivative size = 1026, normalized size of antiderivative = 11.15

$$\int (c + dx) \operatorname{csch}^3(a + bx) dx = \text{Too large to display}$$

input `integrate((d*x+c)*csch(b*x+a)^3,x, algorithm="fracas")`

output

```

-1/2*(2*(b*d*x + b*c + d)*cosh(b*x + a)^3 + 6*(b*d*x + b*c + d)*cosh(b*x +
a)*sinh(b*x + a)^2 + 2*(b*d*x + b*c + d)*sinh(b*x + a)^3 + 2*(b*d*x + b*c
- d)*cosh(b*x + a) + (d*cosh(b*x + a)^4 + 4*d*cosh(b*x + a)*sinh(b*x + a)
^3 + d*sinh(b*x + a)^4 - 2*d*cosh(b*x + a)^2 + 2*(3*d*cosh(b*x + a)^2 - d)
*sinh(b*x + a)^2 + 4*(d*cosh(b*x + a)^3 - d*cosh(b*x + a))*sinh(b*x + a) +
d)*dilog(cosh(b*x + a) + sinh(b*x + a)) - (d*cosh(b*x + a)^4 + 4*d*cosh(b
*x + a)*sinh(b*x + a)^3 + d*sinh(b*x + a)^4 - 2*d*cosh(b*x + a)^2 + 2*(3*d
*cosh(b*x + a)^2 - d)*sinh(b*x + a)^2 + 4*(d*cosh(b*x + a)^3 - d*cosh(b*x
+ a))*sinh(b*x + a) + d)*dilog(-cosh(b*x + a) - sinh(b*x + a)) - ((b*d*x +
b*c)*cosh(b*x + a)^4 + 4*(b*d*x + b*c)*cosh(b*x + a)*sinh(b*x + a)^3 + (b
*d*x + b*c)*sinh(b*x + a)^4 + b*d*x - 2*(b*d*x + b*c)*cosh(b*x + a)^2 - 2*
(b*d*x - 3*(b*d*x + b*c)*cosh(b*x + a)^2 + b*c)*sinh(b*x + a)^2 + b*c + 4*
((b*d*x + b*c)*cosh(b*x + a)^3 - (b*d*x + b*c)*cosh(b*x + a))*sinh(b*x + a
))*log(cosh(b*x + a) + sinh(b*x + a) + 1) + ((b*c - a*d)*cosh(b*x + a)^4 +
4*(b*c - a*d)*cosh(b*x + a)*sinh(b*x + a)^3 + (b*c - a*d)*sinh(b*x + a)^4
- 2*(b*c - a*d)*cosh(b*x + a)^2 + 2*(3*(b*c - a*d)*cosh(b*x + a)^2 - b*c
+ a*d)*sinh(b*x + a)^2 + b*c - a*d + 4*((b*c - a*d)*cosh(b*x + a)^3 - (b*c
- a*d)*cosh(b*x + a))*sinh(b*x + a))*log(cosh(b*x + a) + sinh(b*x + a) -
1) + ((b*d*x + a*d)*cosh(b*x + a)^4 + 4*(b*d*x + a*d)*cosh(b*x + a)*sinh(b
*x + a)^3 + (b*d*x + a*d)*sinh(b*x + a)^4 + b*d*x - 2*(b*d*x + a*d)*cos...

```

3.11.6 Sympy [F]

$$\int (c + dx) \operatorname{csch}^3(a + bx) dx = \int (c + dx) \operatorname{csch}^3(a + bx) dx$$

input `integrate((d*x+c)*csch(b*x+a)**3,x)`

output `Integral((c + d*x)*csch(a + b*x)**3, x)`

3.11.7 Maxima [F]

$$\int (c + dx) \operatorname{csch}^3(a + bx) dx = \int (dx + c) \operatorname{csch}(bx + a)^3 dx$$

input `integrate((d*x+c)*csch(b*x+a)^3,x, algorithm="maxima")`

output `-d*((b*x*e^(3*a) + e^(3*a))*e^(3*b*x) + (b*x*e^a - e^a)*e^(b*x))/(b^2*e^(4*b*x + 4*a) - 2*b^2*e^(2*b*x + 2*a) + b^2) + 8*integrate(1/16*x/(e^(b*x + a) + 1), x) + 8*integrate(1/16*x/(e^(b*x + a) - 1), x) + 1/2*c*(log(e^(-b*x - a) + 1)/b - log(e^(-b*x - a) - 1)/b + 2*(e^(-b*x - a) + e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x - 2*a) - e^(-4*b*x - 4*a) - 1)))`

3.11.8 Giac [F]

$$\int (c + dx) \operatorname{csch}^3(a + bx) dx = \int (dx + c) \operatorname{csch}(bx + a)^3 dx$$

input `integrate((d*x+c)*csch(b*x+a)^3,x, algorithm="giac")`

output `integrate((d*x + c)*csch(b*x + a)^3, x)`

3.11.9 Mupad [F(-1)]

Timed out.

$$\int (c + dx) \operatorname{csch}^3(a + bx) dx = \int \frac{c + dx}{\sinh(a + bx)^3} dx$$

input `int((c + d*x)/sinh(a + b*x)^3,x)`

output `int((c + d*x)/sinh(a + b*x)^3, x)`

3.12 $\int \frac{\operatorname{csch}^3(a+bx)}{x} dx$

3.12.1	Optimal result	112
3.12.2	Mathematica [N/A]	112
3.12.3	Rubi [N/A]	113
3.12.4	Maple [N/A] (verified)	114
3.12.5	Fricas [N/A]	114
3.12.6	Sympy [N/A]	115
3.12.7	Maxima [N/A]	115
3.12.8	Giac [N/A]	115
3.12.9	Mupad [N/A]	116

3.12.1 Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\operatorname{csch}^3(a+bx)}{x} dx = \operatorname{Int}\left(\frac{\operatorname{csch}^3(a+bx)}{x}, x\right)$$

output `Unintegrable(csch(b*x+a)^3/x,x)`

3.12.2 Mathematica [N/A]

Not integrable

Time = 42.66 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^3(a+bx)}{x} dx = \int \frac{\operatorname{csch}^3(a+bx)}{x} dx$$

input `Integrate[Csch[a + b*x]^3/x,x]`

output `Integrate[Csch[a + b*x]^3/x, x]`

3.12.3 Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 26, 4680}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\operatorname{csch}^3(a+bx)}{x} dx \\ \downarrow 3042 \\ \int -\frac{i \csc(ia+ibx)^3}{x} dx \\ \downarrow 26 \\ -i \int \frac{\csc(ia+ibx)^3}{x} dx \\ \downarrow 4680 \\ \int \frac{\operatorname{csch}^3(a+bx)}{x} dx \end{array}$$

input `Int[Csch[a + b*x]^3/x,x]`

output `$Aborted`

3.12.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4680 Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :>
Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], Un
integrable[(c + d*x)^m*Sech[I*(e - Pi/2) + I*f*x]^n, x], (-I)^n*Unintegrabl
e[(c + d*x)^m*Csch[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], Uni
ntegrable[(c + d*x)^m*Sec[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*C
sc[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]
```

3.12.4 Maple [N/A] (verified)

Not integrable

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{csch}(bx+a)^3}{x} dx$$

input `int(csch(b*x+a)^3/x,x)`

output `int(csch(b*x+a)^3/x,x)`

3.12.5 Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^3(a+bx)}{x} dx = \int \frac{\operatorname{csch}(bx+a)^3}{x} dx$$

input `integrate(csch(b*x+a)^3/x,x, algorithm="fricas")`

output `integral(csch(b*x + a)^3/x, x)`

3.12.6 Sympy [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{csch}^3(a + bx)}{x} dx = \int \frac{\operatorname{csch}^3(a + bx)}{x} dx$$

input `integrate(csch(b*x+a)**3/x,x)`output `Integral(csch(a + b*x)**3/x, x)`**3.12.7 Maxima [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 157, normalized size of antiderivative = 13.08

$$\int \frac{\operatorname{csch}^3(a + bx)}{x} dx = \int \frac{\operatorname{csch}(bx + a)^3}{x} dx$$

input `integrate(csch(b*x+a)^3/x,x, algorithm="maxima")`output `-((b*x*e^(3*a) - e^(3*a))*e^(3*b*x) + (b*x*e^a + e^a)*e^(b*x))/(b^2*x^2*e^(4*b*x + 4*a) - 2*b^2*x^2*e^(2*b*x + 2*a) + b^2*x^2) - 8*integrate(1/16*(b^2*x^2 - 2)/(b^2*x^3*e^(b*x + a) + b^2*x^3), x) - 8*integrate(1/16*(b^2*x^2 - 2)/(b^2*x^3*e^(b*x + a) - b^2*x^3), x)`**3.12.8 Giac [N/A]**

Not integrable

Time = 0.64 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^3(a + bx)}{x} dx = \int \frac{\operatorname{csch}(bx + a)^3}{x} dx$$

input `integrate(csch(b*x+a)^3/x,x, algorithm="giac")`output `integrate(csch(b*x + a)^3/x, x)`

3.12.9 Mupad [N/A]

Not integrable

Time = 2.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{csch}^3(a + bx)}{x} dx = \int \frac{1}{x \sinh(a + bx)^3} dx$$

input `int(1/(x*sinh(a + b*x)^3),x)`output `int(1/(x*sinh(a + b*x)^3), x)`

3.13
$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx$$

3.13.1	Optimal result	117
3.13.2	Mathematica [A] (verified)	117
3.13.3	Rubi [A] (verified)	118
3.13.4	Maple [F]	118
3.13.5	Fricas [F(-2)]	119
3.13.6	Sympy [F]	119
3.13.7	Maxima [F]	119
3.13.8	Giac [F]	120
3.13.9	Mupad [F(-1)]	120

3.13.1 Optimal result

Integrand size = 20, antiderivative size = 24

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx = -\frac{4}{9\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{2x \cosh(x)}{3\sqrt{\operatorname{csch}(x)}}$$

output `-4/9/csch(x)^(3/2)+2/3*x*cosh(x)/csch(x)^(1/2)`

3.13.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx = \frac{2(-2 + 3x \operatorname{coth}(x))}{9\operatorname{csch}^{\frac{3}{2}}(x)}$$

input `Integrate[x/Csch[x]^(3/2) + (x*Sqrt[Csch[x]])/3,x]`

output `(2*(-2 + 3*x*Coth[x]))/(9*Csch[x]^(3/2))`

3.13.
$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx$$

3.13.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx$$

↓ 2009

$$\frac{2x \cosh(x)}{3\sqrt{\operatorname{csch}(x)}} - \frac{4}{9\operatorname{csch}^{\frac{3}{2}}(x)}$$

input `Int[x/Csch[x]^(3/2) + (x*Sqrt[Csch[x]])/3,x]`

output `-4/(9*Csch[x]^(3/2)) + (2*x*Cosh[x])/(3*Sqrt[Csch[x]])`

3.13.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.13.4 Maple [F]

$$\int \left(\frac{x}{\operatorname{csch}(x)^{\frac{3}{2}}} + \frac{x\sqrt{\operatorname{csch}(x)}}{3} \right) dx$$

input `int(x/csch(x)^(3/2)+1/3*x*csch(x)^(1/2),x)`

output `int(x/csch(x)^(3/2)+1/3*x*csch(x)^(1/2),x)`

3.13. $\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx$

3.13.5 Fricas [F(-2)]

Exception generated.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx = \text{Exception raised: TypeError}$$

input `integrate(x/csch(x)^(3/2)+1/3*x*csch(x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)`

3.13.6 Sympy [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx = \frac{\int \frac{3x}{\operatorname{csch}^{\frac{3}{2}}(x)} dx + \int x\sqrt{\operatorname{csch}(x)} dx}{3}$$

input `integrate(x/csch(x)**(3/2)+1/3*x*csch(x)**(1/2),x)`

output `(Integral(3*x/csch(x)**(3/2), x) + Integral(x*sqrt(csch(x)), x))/3`

3.13.7 Maxima [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx = \int \frac{1}{3}x\sqrt{\operatorname{csch}(x)} + \frac{x}{\operatorname{csch}(x)^{\frac{3}{2}}} dx$$

input `integrate(x/csch(x)^(3/2)+1/3*x*csch(x)^(1/2),x, algorithm="maxima")`

output `integrate(1/3*x*sqrt(csch(x)) + x/csch(x)^(3/2), x)`

3.13. $\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx$

3.13.8 Giac [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx = \int \frac{1}{3}x\sqrt{\operatorname{csch}(x)} + \frac{x}{\operatorname{csch}(x)^{\frac{3}{2}}} dx$$

input `integrate(x/csch(x)^(3/2)+1/3*x*csch(x)^(1/2),x, algorithm="giac")`

output `integrate(1/3*x*sqrt(csch(x)) + x/csch(x)^(3/2), x)`

3.13.9 Mupad [F(-1)]

Timed out.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx = \int \frac{x\sqrt{\frac{1}{\sinh(x)}}}{3} + \frac{x}{\left(\frac{1}{\sinh(x)}\right)^{3/2}} dx$$

input `int((x*(1/sinh(x))^(1/2))/3 + x/(1/sinh(x))^(3/2),x)`

output `int((x*(1/sinh(x))^(1/2))/3 + x/(1/sinh(x))^(3/2), x)`

3.13. $\int \left(\frac{x}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x\sqrt{\operatorname{csch}(x)} \right) dx$

$$3.14 \quad \int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$$

3.14.1	Optimal result	121
3.14.2	Mathematica [A] (verified)	121
3.14.3	Rubi [A] (verified)	122
3.14.4	Maple [F]	122
3.14.5	Fricas [F(-2)]	123
3.14.6	Sympy [F]	123
3.14.7	Maxima [F]	123
3.14.8	Giac [F]	124
3.14.9	Mupad [F(-1)]	124

3.14.1 Optimal result

Integrand size = 20, antiderivative size = 24

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx = -\frac{4}{25\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{2x \cosh(x)}{5\operatorname{csch}^{\frac{3}{2}}(x)}$$

output `-4/25/csch(x)^(5/2)+2/5*x*cosh(x)/csch(x)^(3/2)`

3.14.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx = \frac{2(-2 + 5x \operatorname{coth}(x))}{25\operatorname{csch}^{\frac{5}{2}}(x)}$$

input `Integrate[x/Csch[x]^(5/2) + (3*x)/(5*Sqrt[Csch[x]]),x]`

output `(2*(-2 + 5*x*Coth[x]))/(25*Csch[x]^(5/2))`

$$3.14. \quad \int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$$

3.14.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$$

↓ 2009

$$\frac{2x \cosh(x)}{5\operatorname{csch}^{\frac{3}{2}}(x)} - \frac{4}{25\operatorname{csch}^{\frac{5}{2}}(x)}$$

input `Int[x/Csch[x]^(5/2) + (3*x)/(5*Sqrt[Csch[x]]),x]`

output `-4/(25*Csch[x]^(5/2)) + (2*x*Cosh[x])/(5*Csch[x]^(3/2))`

3.14.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.14.4 Maple [F]

$$\int \left(\frac{x}{\operatorname{csch}(x)^{\frac{5}{2}}} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$$

input `int(x/csch(x)^(5/2)+3/5*x/csch(x)^(1/2),x)`

output `int(x/csch(x)^(5/2)+3/5*x/csch(x)^(1/2),x)`

3.14. $\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$

3.14.5 Fracas [F(-2)]

Exception generated.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx = \text{Exception raised: TypeError}$$

input `integrate(x/csch(x)^(5/2)+3/5*x/csch(x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)`

3.14.6 Sympy [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx = \frac{\int \frac{5x}{\operatorname{csch}^{\frac{5}{2}}(x)} dx + \int \frac{3x}{\sqrt{\operatorname{csch}(x)}} dx}{5}$$

input `integrate(x/csch(x)**(5/2)+3/5*x/csch(x)**(1/2),x)`

output `(Integral(5*x/csch(x)**(5/2), x) + Integral(3*x/sqrt(csch(x)), x))/5`

3.14.7 Maxima [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx = \int \frac{3x}{5\sqrt{\operatorname{csch}(x)}} + \frac{x}{\operatorname{csch}(x)^{\frac{5}{2}}} dx$$

input `integrate(x/csch(x)^(5/2)+3/5*x/csch(x)^(1/2),x, algorithm="maxima")`

output `integrate(3/5*x/sqrt(csch(x)) + x/csch(x)^(5/2), x)`

3.14. $\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$

3.14.8 Giac [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx = \int \frac{3x}{5\sqrt{\operatorname{csch}(x)}} + \frac{x}{\operatorname{csch}(x)^{\frac{5}{2}}} dx$$

input `integrate(x/csch(x)^(5/2)+3/5*x/csch(x)^(1/2),x, algorithm="giac")`

output `integrate(3/5*x/sqrt(csch(x)) + x/csch(x)^(5/2), x)`

3.14.9 Mupad [F(-1)]

Timed out.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx = \int \frac{3x}{5\sqrt{\frac{1}{\sinh(x)}}} + \frac{x}{\left(\frac{1}{\sinh(x)}\right)^{\frac{5}{2}}} dx$$

input `int((3*x)/(5*(1/sinh(x))^(1/2)) + x/(1/sinh(x))^(5/2),x)`

output `int((3*x)/(5*(1/sinh(x))^(1/2)) + x/(1/sinh(x))^(5/2), x)`

3.14. $\int \left(\frac{x}{\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{3x}{5\sqrt{\operatorname{csch}(x)}} \right) dx$

3.15
$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21}x\sqrt{\operatorname{csch}(x)} \right) dx$$

3.15.1	Optimal result	125
3.15.2	Mathematica [A] (verified)	125
3.15.3	Rubi [A] (verified)	126
3.15.4	Maple [F]	126
3.15.5	Fricas [F(-2)]	127
3.15.6	Sympy [F]	127
3.15.7	Maxima [F]	127
3.15.8	Giac [F]	128
3.15.9	Mupad [F(-1)]	128

3.15.1 Optimal result

Integrand size = 20, antiderivative size = 47

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21}x\sqrt{\operatorname{csch}(x)} \right) dx = -\frac{4}{49\operatorname{csch}^{\frac{7}{2}}(x)} + \frac{2x \cosh(x)}{7\operatorname{csch}^{\frac{5}{2}}(x)} + \frac{20}{63\operatorname{csch}^{\frac{3}{2}}(x)} - \frac{10x \cosh(x)}{21\sqrt{\operatorname{csch}(x)}}$$

output `-4/49/csch(x)^(7/2)+2/7*x*cosh(x)/csch(x)^(5/2)+20/63/csch(x)^(3/2)-10/21*x*cosh(x)/csch(x)^(1/2)`

3.15.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21}x\sqrt{\operatorname{csch}(x)} \right) dx = \sqrt{\operatorname{csch}(x)} \left(-\frac{167}{882} + \frac{88}{441} \cosh(2x) - \frac{1}{98} \cosh(4x) - \frac{13}{42}x \sinh(2x) + \frac{1}{28}x \sinh(4x) \right)$$

input `Integrate[x/Csch[x]^(7/2) - (5*x*Sqrt[Csch[x]])/21,x]`

output `Sqrt[Csch[x]]*(-167/882 + (88*Cosh[2*x])/441 - Cosh[4*x]/98 - (13*x*Sinh[2*x])/42 + (x*Sinh[4*x])/28)`

3.15.
$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21}x\sqrt{\operatorname{csch}(x)} \right) dx$$

3.15.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx$$

↓ 2009

$$\frac{20}{63 \operatorname{csch}^{\frac{3}{2}}(x)} - \frac{4}{49 \operatorname{csch}^{\frac{7}{2}}(x)} + \frac{2x \cosh(x)}{7 \operatorname{csch}^{\frac{5}{2}}(x)} - \frac{10x \cosh(x)}{21 \sqrt{\operatorname{csch}(x)}}$$

input `Int[x/Csch[x]^(7/2) - (5*x*Sqrt[Csch[x]])/21,x]`

output `-4/(49*Csch[x]^(7/2)) + (2*x*Cosh[x])/(7*Csch[x]^(5/2)) + 20/(63*Csch[x]^(3/2)) - (10*x*Cosh[x])/(21*Sqrt[Csch[x]])`

3.15.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.15.4 Maple [F]

$$\int \left(\frac{x}{\operatorname{csch}(x)^{\frac{7}{2}}} - \frac{5x \sqrt{\operatorname{csch}(x)}}{21} \right) dx$$

input `int(x/csch(x)^(7/2)-5/21*x*csch(x)^(1/2),x)`

output `int(x/csch(x)^(7/2)-5/21*x*csch(x)^(1/2),x)`

3.15. $\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx$

3.15.5 Fricas [F(-2)]

Exception generated.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx = \text{Exception raised: TypeError}$$

input `integrate(x/csch(x)^(7/2)-5/21*x*csch(x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)`

3.15.6 Sympy [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx = -\frac{\int \left(-\frac{21x}{\operatorname{csch}^{\frac{7}{2}}(x)} \right) dx + \int 5x \sqrt{\operatorname{csch}(x)} dx}{21}$$

input `integrate(x/csch(x)**(7/2)-5/21*x*csch(x)**(1/2),x)`

output `-(Integral(-21*x/csch(x)**(7/2), x) + Integral(5*x*sqrt(csch(x)), x))/21`

3.15.7 Maxima [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx = \int -\frac{5}{21} x \sqrt{\operatorname{csch}(x)} + \frac{x}{\operatorname{csch}(x)^{\frac{7}{2}}} dx$$

input `integrate(x/csch(x)^(7/2)-5/21*x*csch(x)^(1/2),x, algorithm="maxima")`

output `integrate(-5/21*x*sqrt(csch(x)) + x/csch(x)^(7/2), x)`

3.15. $\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx$

3.15.8 Giac [F]

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx = \int -\frac{5}{21} x \sqrt{\operatorname{csch}(x)} + \frac{x}{\operatorname{csch}(x)^{\frac{7}{2}}} dx$$

input `integrate(x/csch(x)^(7/2)-5/21*x*csch(x)^(1/2),x, algorithm="giac")`

output `integrate(-5/21*x*sqrt(csch(x)) + x/csch(x)^(7/2), x)`

3.15.9 Mupad [F(-1)]

Timed out.

$$\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx = - \int \frac{5x \sqrt{\frac{1}{\sinh(x)}}}{21} - \frac{x}{\left(\frac{1}{\sinh(x)}\right)^{7/2}} dx$$

input `int(x/(1/sinh(x))^(7/2) - (5*x*(1/sinh(x))^(1/2))/21,x)`

output `-int((5*x*(1/sinh(x))^(1/2))/21 - x/(1/sinh(x))^(7/2), x)`

3.15. $\int \left(\frac{x}{\operatorname{csch}^{\frac{7}{2}}(x)} - \frac{5}{21} x \sqrt{\operatorname{csch}(x)} \right) dx$

3.16
$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2 \sqrt{\operatorname{csch}(x)} \right) dx$$

3.16.1	Optimal result	129
3.16.2	Mathematica [A] (verified)	129
3.16.3	Rubi [A] (verified)	130
3.16.4	Maple [F]	131
3.16.5	Fricas [F(-2)]	131
3.16.6	Sympy [F]	131
3.16.7	Maxima [F]	132
3.16.8	Giac [F]	132
3.16.9	Mupad [F(-1)]	132

3.16.1 Optimal result

Integrand size = 24, antiderivative size = 76

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2 \sqrt{\operatorname{csch}(x)} \right) dx = -\frac{8x}{9\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{16 \cosh(x)}{27\sqrt{\operatorname{csch}(x)}} + \frac{2x^2 \cosh(x)}{3\sqrt{\operatorname{csch}(x)}} - \frac{16}{27}i\sqrt{\operatorname{csch}(x)} \operatorname{EllipticF}\left(\frac{\pi}{4} - \frac{ix}{2}, 2\right) \sqrt{i \sinh(x)}$$

output `-8/9*x/csch(x)^(3/2)+16/27*cosh(x)/csch(x)^(1/2)+2/3*x^2*cosh(x)/csch(x)^(1/2)-16/27*I*(sin(1/4*Pi+1/2*I*x)^2)^(1/2)/sin(1/4*Pi+1/2*I*x)*EllipticF(cos(1/4*Pi+1/2*I*x),2^(1/2))*csch(x)^(1/2)*(I*sinh(x))^(1/2)`

3.16.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.88

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2 \sqrt{\operatorname{csch}(x)} \right) dx = \frac{4(3 + \operatorname{csch}^2(x)) \left(-12x + (8 + 9x^2) \operatorname{coth}(x) + \frac{8\operatorname{csch}(x) \operatorname{EllipticF}\left(\frac{1}{4}(\pi - 2ix), 2\right)}{\sqrt{i \sinh(x)}} \right)}{27(-1 + 3 \cosh(2x))\operatorname{csch}^{\frac{7}{2}}(x)}$$

3.16.
$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2 \sqrt{\operatorname{csch}(x)} \right) dx$$

input `Integrate[x^2/Csch[x]^(3/2) + (x^2*Sqrt[Csch[x]])/3,x]`

output `(4*(3 + Csch[x]^2)*(-12*x + (8 + 9*x^2)*Coth[x] + (8*Csch[x]*EllipticF[(Pi - (2*I)*x)/4, 2])/Sqrt[I*Sinh[x]])/(27*(-1 + 3*Cosh[2*x])*Csch[x]^(7/2))`

3.16.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} \right) dx$$

↓ 2009

$$\frac{2x^2 \cosh(x)}{3\sqrt{\operatorname{csch}(x)}} - \frac{8x}{9\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{16 \cosh(x)}{27\sqrt{\operatorname{csch}(x)}} - \frac{16}{27}i\sqrt{i \sinh(x)}\sqrt{\operatorname{csch}(x)} \operatorname{EllipticF}\left(\frac{\pi}{4} - \frac{ix}{2}, 2\right)$$

input `Int[x^2/Csch[x]^(3/2) + (x^2*Sqrt[Csch[x]])/3,x]`

output `(-8*x)/(9*Csch[x]^(3/2)) + (16*Cosh[x])/(27*Sqrt[Csch[x]]) + (2*x^2*Cosh[x])/ (3*Sqrt[Csch[x]]) - ((16*I)/27)*Sqrt[Csch[x]]*EllipticF[Pi/4 - (I/2)*x, 2]*Sqrt[I*Sinh[x]]`

3.16. $\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} \right) dx$

3.16.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.16.4 Maple [F]

$$\int \left(\frac{x^2}{\operatorname{csch}(x)^{\frac{3}{2}}} + \frac{x^2 \sqrt{\operatorname{csch}(x)}}{3} \right) dx$$

input `int(x^2/csch(x)^(3/2)+1/3*x^2*csch(x)^(1/2),x)`

output `int(x^2/csch(x)^(3/2)+1/3*x^2*csch(x)^(1/2),x)`

3.16.5 Fricas [F(-2)]

Exception generated.

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2 \sqrt{\operatorname{csch}(x)} \right) dx = \text{Exception raised: TypeError}$$

input `integrate(x^2/csch(x)^(3/2)+1/3*x^2*csch(x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)`

3.16.6 Sympy [F]

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2 \sqrt{\operatorname{csch}(x)} \right) dx = \frac{\int \frac{3x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} dx + \int x^2 \sqrt{\operatorname{csch}(x)} dx}{3}$$

input `integrate(x**2/csch(x)**(3/2)+1/3*x**2*csch(x)**(1/2),x)`

output `(Integral(3*x**2/csch(x)**(3/2), x) + Integral(x**2*sqrt(csch(x)), x))/3`

3.16. $\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2 \sqrt{\operatorname{csch}(x)} \right) dx$

3.16.7 Maxima [F]

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} \right) dx = \int \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} + \frac{x^2}{\operatorname{csch}(x)^{\frac{3}{2}}} dx$$

input `integrate(x^2/csch(x)^(3/2)+1/3*x^2*csch(x)^(1/2),x, algorithm="maxima")`

output `integrate(1/3*x^2*sqrt(csch(x)) + x^2/csch(x)^(3/2), x)`

3.16.8 Giac [F]

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} \right) dx = \int \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} + \frac{x^2}{\operatorname{csch}(x)^{\frac{3}{2}}} dx$$

input `integrate(x^2/csch(x)^(3/2)+1/3*x^2*csch(x)^(1/2),x, algorithm="giac")`

output `integrate(1/3*x^2*sqrt(csch(x)) + x^2/csch(x)^(3/2), x)`

3.16.9 Mupad [F(-1)]

Timed out.

$$\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} \right) dx = \int \frac{x^2\sqrt{\frac{1}{\sinh(x)}}}{3} + \frac{x^2}{\left(\frac{1}{\sinh(x)}\right)^{3/2}} dx$$

input `int((x^2*(1/sinh(x))^(1/2))/3 + x^2/(1/sinh(x))^(3/2),x)`

output `int((x^2*(1/sinh(x))^(1/2))/3 + x^2/(1/sinh(x))^(3/2), x)`

3.16. $\int \left(\frac{x^2}{\operatorname{csch}^{\frac{3}{2}}(x)} + \frac{1}{3}x^2\sqrt{\operatorname{csch}(x)} \right) dx$

3.17 $\int \frac{(e+fx)^3 \cosh(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$

3.17.1	Optimal result	133
3.17.2	Mathematica [B] (warning: unable to verify)	134
3.17.3	Rubi [C] (verified)	135
3.17.4	Maple [F]	141
3.17.5	Fricas [B] (verification not implemented)	141
3.17.6	Sympy [F]	142
3.17.7	Maxima [F]	143
3.17.8	Giac [F]	143
3.17.9	Mupad [F(-1)]	143

3.17.1 Optimal result

Integrand size = 26, antiderivative size = 448

$$\int \frac{(e+fx)^3 \cosh(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{b(e+fx)^4}{4a^2f} - \frac{6f^3 \cosh(c+dx)}{ad^4} - \frac{3f(e+fx)^2 \cosh(c+dx)}{ad^2}$$

$$- \frac{b(e+fx)^3 \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d}$$

$$- \frac{b(e+fx)^3 \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d}$$

$$- \frac{3bf(e+fx)^2 \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d^2}$$

$$- \frac{3bf(e+fx)^2 \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d^2}$$

$$+ \frac{6bf^2(e+fx) \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d^3}$$

$$+ \frac{6bf^2(e+fx) \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d^3}$$

$$- \frac{6bf^3 \text{PolyLog}\left(4, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d^4}$$

$$- \frac{6bf^3 \text{PolyLog}\left(4, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d^4}$$

$$+ \frac{6f^2(e+fx) \sinh(c+dx)}{ad^3} + \frac{(e+fx)^3 \sinh(c+dx)}{ad}$$

output $\frac{1}{4}b*(f*x+e)^4/a^2/f-6*f^3*\cosh(d*x+c)/a/d^4-3*f*(f*x+e)^2*\cosh(d*x+c)/a/d^2-b*(f*x+e)^3*\ln(1+a*\exp(d*x+c)/(b-(a^2+b^2)^{(1/2)}))/a^2/d-b*(f*x+e)^3*\ln(1+a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2)}))/a^2/d-3*b*f*(f*x+e)^2*\text{polylog}(2,-a*\exp(d*x+c)/(b-(a^2+b^2)^{(1/2)}))/a^2/d^2-3*b*f*(f*x+e)^2*\text{polylog}(2,-a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2)}))/a^2/d^2+6*b*f^2*(f*x+e)*\text{polylog}(3,-a*\exp(d*x+c)/(b-(a^2+b^2)^{(1/2)}))/a^2/d^3+6*b*f^2*(f*x+e)*\text{polylog}(3,-a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2)}))/a^2/d^3-6*b*f^3*\text{polylog}(4,-a*\exp(d*x+c)/(b-(a^2+b^2)^{(1/2)}))/a^2/d^4+6*b*f^3*\text{polylog}(4,-a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2)}))/a^2/d^4+6*f^2*(f*x+e)*\sinh(d*x+c)/a/d^3+(f*x+e)^3*\sinh(d*x+c)/a/d$

3.17.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 1792 vs. $2(448) = 896$.

Time = 20.20 (sec) , antiderivative size = 1792, normalized size of antiderivative = 4.00

$$\int \frac{(e + fx)^3 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((e + f*x)^3*Cosh[c + d*x])/(a + b*CsSch[c + d*x]),x]`

output $(e*f^2*CsSch[c + d*x]*((4*b*x^3)/(-1 + E^{(2*c)}) - 2*b*x^3*Coth[c] - (6*a^2*b*(d^2*x^2*Log[1 + ((b - Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a] - 2*d*x*PolyLog[2, ((-b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a] - 2*PolyLog[3, ((-b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a]))/(Sqrt[a^2 + b^2]*(-b + Sqrt[a^2 + b^2])*d^3) - (6*a^2*b*(d^2*x^2*Log[1 + ((b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a] - 2*d*x*PolyLog[2, -(((b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a] - 2*PolyLog[3, -(((b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a)])))/(Sqrt[a^2 + b^2]*(b + Sqrt[a^2 + b^2])*d^3) + (6*b^2*(d^2*x^2*Log[1 + (a*E^{(c + d*x)})/(b - Sqrt[a^2 + b^2])]) + 2*d*x*PolyLog[2, (a*E^{(c + d*x)})/(-b + Sqrt[a^2 + b^2])]) - 2*PolyLog[3, (a*E^{(c + d*x)})/(-b + Sqrt[a^2 + b^2])]))/(Sqrt[a^2 + b^2]*d^3) - (6*b^2*(d^2*x^2*Log[1 + (a*E^{(c + d*x)})/(b + Sqrt[a^2 + b^2])]) + 2*d*x*PolyLog[2, -((a*E^{(c + d*x)})/(b + Sqrt[a^2 + b^2]))]) - 2*PolyLog[3, -((a*E^{(c + d*x)})/(b + Sqrt[a^2 + b^2]))]))/(Sqrt[a^2 + b^2]*d^3) + (6*a*Cosh[d*x]*(-2*d*x*Cosh[c] + (2 + d^2*x^2)*Sinh[c]))/d^3 + (6*a*((2 + d^2*x^2)*Cosh[c] - 2*d*x*Sinh[c])*Sinh[d*x])/d^3*(b + a*Sinh[c + d*x]))/(2*a^2*(a + b*CsSch[c + d*x])) + (f^3*CsSch[c + d*x]*((b*(x^4 - (2*a^2*(-1 + E^{(2*c)}))*(d^3*x^3*Log[1 + ((b - Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a] - 3*d^2*x^2*PolyLog[2, ((-b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a] - 6*d*x*PolyLog[3, ((-b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a] - 6*PolyLog[4, ((-b + Sqrt[a^2 + b^2])*E^{(-c - d*x)})/a]))/(Sqrt[a^2 + b^2]*(-b + Sqrt[a^2 + b^2])*d^4) - (2*a^2*(-1 + E^{(2*...$

3.17.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 2.45 (sec) , antiderivative size = 452, normalized size of antiderivative = 1.01, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.769$, Rules used = {6128, 6113, 3042, 3777, 26, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3118, 6095, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(e+fx)^3 \cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx \\
 & \quad \downarrow \text{6128} \\
 & \int \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{a \sinh(c+dx) + b} dx \\
 & \quad \downarrow \text{6113} \\
 & \frac{\int (e+fx)^3 \cosh(c+dx) dx}{a} - \frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx)^3 \sin\left(ic+idx+\frac{\pi}{2}\right) dx}{a} \\
 & \quad \downarrow \text{3777} \\
 & -\frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3if \int -i(e+fx)^2 \sinh(c+dx) dx}{d}}{a} \\
 & \quad \downarrow \text{26} \\
 & \frac{\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3f \int (e+fx)^2 \sinh(c+dx) dx}{d}}{a} - \frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3f \int -i(e+fx)^2 \sin(ic+idx) dx}{d}}{a} \\
 & \quad \downarrow \text{26} \\
 & -\frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \int (e+fx)^2 \sin(ic+idx) dx}{d}}{a} \\
 & \quad \downarrow \text{3777}
 \end{aligned}$$

3.17. $\int \frac{(e+fx)^3 \cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

$$\begin{aligned}
& -\frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \int (e+fx) \cosh(c+dx) dx}{d} \right)}{a} \\
& \quad \downarrow \text{3042} \\
& -\frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \int (e+fx) \sin\left(\frac{ic+idx+\frac{\pi}{2}}{d}\right) dx}{d} \right)}{a} \\
& \quad \downarrow \text{3777} \\
& \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{if \int -i \sinh(c+dx) dx}{d} \right)}{d} \right)}{a} \\
& \quad \downarrow \text{26} \\
& \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sinh(c+dx) dx}{d} \right)}{d} \right)}{a} \\
& \quad \downarrow \text{3042} \\
& \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int -i \sin\left(\frac{ic+idx}{d}\right) dx}{d} \right)}{d} \right)}{a} \\
& \quad \downarrow \text{26} \\
& \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} + \frac{if \int \sin\left(\frac{ic+idx}{d}\right) dx}{d} \right)}{d} \right)}{a} \\
& \quad \downarrow \text{3118} \\
& -\frac{b \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{d} \right)}{a}
\end{aligned}$$

3.17. $\int \frac{(e+fx)^3 \cosh(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow \text{6095} \\
 & \frac{b \left(\int \frac{e^{c+dx} (e+fx)^3}{e^{c+dx} a + b - \sqrt{a^2 + b^2}} dx + \int \frac{e^{c+dx} (e+fx)^3}{e^{c+dx} a + b + \sqrt{a^2 + b^2}} dx - \frac{(e+fx)^4}{4af} \right) + \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{d} \right)}{a}}{a} \\
 & \downarrow \text{2620} \\
 & b \left(-\frac{3f \int (e+fx)^2 \log \left(\frac{e^{c+dx} a}{b - \sqrt{a^2 + b^2}} + 1 \right) dx}{ad} - \frac{3f \int (e+fx)^2 \log \left(\frac{e^{c+dx} a}{b + \sqrt{a^2 + b^2}} + 1 \right) dx}{ad} + \frac{(e+fx)^3 \log \left(\frac{ae^{c+dx}}{b - \sqrt{a^2 + b^2}} + 1 \right)}{ad} + \frac{(e+fx)^3 \log \left(\frac{ae^{c+dx}}{b + \sqrt{a^2 + b^2}} + 1 \right)}{ad} \right) \\
 & \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{d} \right)}{a} \\
 & \downarrow \text{3011} \\
 & b \left(-\frac{3f \left(\frac{2f \int (e+fx) \operatorname{PolyLog} \left(2, -\frac{ae^{c+dx}}{b - \sqrt{a^2 + b^2}} \right) dx}{d} - \frac{(e+fx)^2 \operatorname{PolyLog} \left(2, -\frac{ae^{c+dx}}{b - \sqrt{a^2 + b^2}} \right)}{d} \right)}{ad} - \frac{3f \left(\frac{2f \int (e+fx) \operatorname{PolyLog} \left(2, -\frac{ae^{c+dx}}{b + \sqrt{a^2 + b^2}} \right) dx}{d} - \frac{(e+fx)^2 \operatorname{PolyLog} \left(2, -\frac{ae^{c+dx}}{b + \sqrt{a^2 + b^2}} \right)}{d} \right)}{ad} \right) \\
 & \frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{d} \right)}{a} \\
 & \downarrow \text{7163}
 \end{aligned}$$

3.17. $\int \frac{(e+fx)^3 \cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

$$b \left(\frac{3f \left(\frac{2f \left(\frac{(e+fx) \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{d} - \frac{f \int \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right) dx}{d} \right)}{d} \right) - (e+fx)^2 \operatorname{PolyLog} \left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{ad} \right) - \frac{3f \left(\frac{(e+fx) \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{d} \right)}{ad} \right)$$

$$\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{d} \right)}{a}$$

↓ 2720

$$b \left(\frac{3f \left(\frac{2f \left(\frac{(e+fx) \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{d} - \frac{f \int e^{-c-dx} \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right) de^{c+dx}}{d^2} \right)}{d} \right) - (e+fx)^2 \operatorname{PolyLog} \left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{ad} \right) - \frac{3f \left(\frac{(e+fx) \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{d} \right)}{ad} \right)$$

$$\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{d} \right)}{a}$$

↓ 7143

$$b \left(\frac{3f \left(\frac{2f \left(\frac{(e+fx) \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{d} - \frac{f \operatorname{PolyLog} \left(4, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{d^2} \right)}{d} \right) - (e+fx)^2 \operatorname{PolyLog} \left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{ad} \right) - \frac{3f \left(\frac{(e+fx) \operatorname{PolyLog} \left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} \right)}{d} \right)}{ad} \right)$$

$$\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{d} \right)}{a}$$

3.17. $\int \frac{(e+fx)^3 \cosh(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

input `Int[((e + f*x)^3*Cosh[c + d*x])/(a + b*Csch[c + d*x]),x]`

output `-((b*(-1/4*(e + f*x)^4/(a*f) + ((e + f*x)^3*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/(a*d) + ((e + f*x)^3*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]])))/(a*d) - (3*f*(-((e + f*x)^2*PolyLog[2, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d) + (2*f*(((e + f*x)*PolyLog[3, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d - (f*PolyLog[4, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d^2))/d)/(a*d) - (3*f*(-((e + f*x)^2*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]])))/d) + (2*f*(((e + f*x)*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]])))/d - (f*PolyLog[4, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]])))/d^2))/d)/(a*d))/a + (((e + f*x)^3*Sinh[c + d*x])/d + ((3*I)*f*((I*(e + f*x)^2*Cosh[c + d*x])/d - ((2*I)*f*(-((f*Cosh[c + d*x])/d^2) + ((e + f*x)*Sinh[c + d*x])/d))/d))/d)/a`

3.17.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))] * ((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 6095 `Int[(Cosh[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)), x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 + b^2, 2] + b*E^(c + d*x))), x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 + b^2, 2] + b*E^(c + d*x))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 + b^2, 0]`

rule 6113 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(p_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)]^(n_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^p*Sinh[c + d*x]^(n - 1), x], x] - Simp[a/b Int[(e + f*x)^m*Cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*Sinh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6128 `Int[((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)/(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && IntegersQ[m, n]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

```
rule 7163 Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

3.17.4 Maple [F]

$$\int \frac{(fx + e)^3 \cosh(dx + c)}{a + b \operatorname{csch}(dx + c)} dx$$

```
input int((f*x+e)^3*cosh(d*x+c)/(a+b*csch(d*x+c)),x)
```

```
output int((f*x+e)^3*cosh(d*x+c)/(a+b*csch(d*x+c)),x)
```

3.17.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1976 vs. $2(420) = 840$.

Time = 0.30 (sec) , antiderivative size = 1976, normalized size of antiderivative = 4.41

$$\int \frac{(e + fx)^3 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((f*x+e)^3*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="fracas")
```

```

output -1/4*(2*a*d^3*f^3*x^3 + 2*a*d^3*e^3 + 6*a*d^2*e^2*f + 12*a*d*e*f^2 + 12*a*
f^3 + 6*(a*d^3*e*f^2 + a*d^2*f^3)*x^2 - 2*(a*d^3*f^3*x^3 + a*d^3*e^3 - 3*a
*d^2*e^2*f + 6*a*d*e*f^2 - 6*a*f^3 + 3*(a*d^3*e*f^2 - a*d^2*f^3)*x^2 + 3*(
a*d^3*e^2*f - 2*a*d^2*e*f^2 + 2*a*d*f^3)*x)*cosh(d*x + c)^2 - 2*(a*d^3*f^3
*x^3 + a*d^3*e^3 - 3*a*d^2*e^2*f + 6*a*d*e*f^2 - 6*a*f^3 + 3*(a*d^3*e*f^2
- a*d^2*f^3)*x^2 + 3*(a*d^3*e^2*f - 2*a*d^2*e*f^2 + 2*a*d*f^3)*x)*sinh(d*x
+ c)^2 + 6*(a*d^3*e^2*f + 2*a*d^2*e*f^2 + 2*a*d*f^3)*x - (b*d^4*f^3*x^4 +
4*b*d^4*e*f^2*x^3 + 6*b*d^4*e^2*f*x^2 + 4*b*d^4*e^3*x + 8*b*c*d^3*e^3 - 1
2*b*c^2*d^2*e^2*f + 8*b*c^3*d*e*f^2 - 2*b*c^4*f^3)*cosh(d*x + c) + 12*((b*
d^2*f^3*x^2 + 2*b*d^2*e*f^2*x + b*d^2*e^2*f)*cosh(d*x + c) + (b*d^2*f^3*x^
2 + 2*b*d^2*e*f^2*x + b*d^2*e^2*f)*sinh(d*x + c))*dilog((b*cosh(d*x + c) +
b*sinh(d*x + c) + (a*cosh(d*x + c) + a*sinh(d*x + c))*sqrt((a^2 + b^2)/a^
2) - a)/a + 1) + 12*((b*d^2*f^3*x^2 + 2*b*d^2*e*f^2*x + b*d^2*e^2*f)*cosh(
d*x + c) + (b*d^2*f^3*x^2 + 2*b*d^2*e*f^2*x + b*d^2*e^2*f)*sinh(d*x + c))*
dilog((b*cosh(d*x + c) + b*sinh(d*x + c) - (a*cosh(d*x + c) + a*sinh(d*x +
c))*sqrt((a^2 + b^2)/a^2) - a)/a + 1) + 4*((b*d^3*e^3 - 3*b*c*d^2*e^2*f +
3*b*c^2*d*e*f^2 - b*c^3*f^3)*cosh(d*x + c) + (b*d^3*e^3 - 3*b*c*d^2*e^2*f
+ 3*b*c^2*d*e*f^2 - b*c^3*f^3)*sinh(d*x + c))*log(2*a*cosh(d*x + c) + 2*a
*sinh(d*x + c) + 2*a*sqrt((a^2 + b^2)/a^2) + 2*b) + 4*((b*d^3*e^3 - 3*b*c*
d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*cosh(d*x + c) + (b*d^3*e^3 - 3...

```

3.17.6 Sympy [F]

$$\int \frac{(e + fx)^3 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx)^3 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

```
input integrate((f*x+e)**3*cosh(d*x+c)/(a+b*csch(d*x+c)),x)
```

```
output Integral((e + f*x)**3*cosh(c + d*x)/(a + b*csch(c + d*x)), x)
```

3.17.7 Maxima [F]

$$\int \frac{(e + fx)^3 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^3 \cosh(dx + c)}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^3*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/2*e^3*(2*(d*x + c)*b/(a^2*d) - e^(d*x + c)/(a*d) + e^(-d*x - c)/(a*d) + 2*b*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a^2*d)) - 1/4*(b*d^4*f^3*x^4*e^c + 4*b*d^4*e*f^2*x^3*e^c + 6*b*d^4*e^2*f*x^2*e^c - 2*(a*d^3*f^3*x^3*e^(2*c) + 3*(d^3*e*f^2 - d^2*f^3)*a*x^2*e^(2*c) + 3*(d^3*e^2*f - 2*d^2*e*f^2 + 2*d*f^3)*a*x*e^(2*c) - 3*(d^2*e^2*f - 2*d*e*f^2 + 2*f^3)*a*e^(2*c))*e^(d*x) + 2*(a*d^3*f^3*x^3 + 3*(d^3*e*f^2 + d^2*f^3)*a*x^2 + 3*(d^3*e^2*f + 2*d^2*e*f^2 + 2*d*f^3)*a*x + 3*(d^2*e^2*f + 2*d*e*f^2 + 2*f^3)*a)*e^(-d*x)*e^(-c)/(a^2*d^4) + integrate(-2*(a*b*f^3*x^3 + 3*a*b*e*f^2*x^2 + 3*a*b*e^2*f*x - (b^2*f^3*x^3*e^c + 3*b^2*e*f^2*x^2*e^c + 3*b^2*e^2*f*x*e^c)*e^(d*x))/(a^3*e^(2*d*x + 2*c) + 2*a^2*b*e^(d*x + c) - a^3), x)`

3.17.8 Giac [F]

$$\int \frac{(e + fx)^3 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^3 \cosh(dx + c)}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^3*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)^3*cosh(d*x + c)/(b*csch(d*x + c) + a), x)`

3.17.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx)^3 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx) (e + fx)^3}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)*(e + f*x)^3)/(a + b/sinh(c + d*x)),x)`

output `int((cosh(c + d*x)*(e + f*x)^3)/(a + b/sinh(c + d*x)), x)`

3.18 $\int \frac{(e+fx)^2 \cosh(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$

3.18.1	Optimal result	144
3.18.2	Mathematica [B] (verified)	145
3.18.3	Rubi [C] (verified)	146
3.18.4	Maple [F]	151
3.18.5	Fricas [B] (verification not implemented)	151
3.18.6	Sympy [F]	152
3.18.7	Maxima [F]	152
3.18.8	Giac [F]	152
3.18.9	Mupad [F(-1)]	153

3.18.1 Optimal result

Integrand size = 26, antiderivative size = 330

$$\int \frac{(e+fx)^2 \cosh(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{b(e+fx)^3}{3a^2f} - \frac{2f(e+fx) \cosh(c+dx)}{ad^2} - \frac{b(e+fx)^2 \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d} - \frac{b(e+fx)^2 \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d} - \frac{2bf(e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d^2} - \frac{2bf(e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d^2} + \frac{2bf^2 \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d^3} + \frac{2bf^2 \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d^3} + \frac{2f^2 \sinh(c+dx)}{ad^3} + \frac{(e+fx)^2 \sinh(c+dx)}{ad}$$

output $\frac{1}{3}b(fx+e)^3/a^2/f-2f*(fx+e)*\cosh(dx+c)/a/d^2-b*(fx+e)^2*\ln(1+a*\exp(dx+c)/(b-(a^2+b^2)^{(1/2)}))/a^2/d-b*(fx+e)^2*\ln(1+a*\exp(dx+c)/(b+(a^2+b^2)^{(1/2)}))/a^2/d-2*b*f*(fx+e)*\text{polylog}(2,-a*\exp(dx+c)/(b-(a^2+b^2)^{(1/2)}))/a^2/d^2-2*b*f*(fx+e)*\text{polylog}(2,-a*\exp(dx+c)/(b+(a^2+b^2)^{(1/2)}))/a^2/d^2+2*b*f^2*\text{polylog}(3,-a*\exp(dx+c)/(b-(a^2+b^2)^{(1/2)}))/a^2/d^3+2*b*f^2*\text{polylog}(3,-a*\exp(dx+c)/(b+(a^2+b^2)^{(1/2)}))/a^2/d^3+2*f^2*\sinh(dx+c)/a/d^3+(fx+e)^2*\sinh(dx+c)/a/d$

3.18.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 937 vs. $2(330) = 660$.

Time = 11.57 (sec) , antiderivative size = 937, normalized size of antiderivative = 2.84

$$\int \frac{(e+fx)^2 \cosh(c+dx)}{a+b\text{csch}(c+dx)} dx$$

$$f^2 \text{csch}(c+dx) \left(\frac{4bx^3}{-1+e^{2c}} - 2bx^3 \coth(c) - \frac{6a^2b \left(d^2 x^2 \log \left(1 + \frac{(b-\sqrt{a^2+b^2})e^{-c-dx}}{a} \right) - 2dx \text{PolyLog} \left(2, \frac{(b+\sqrt{a^2+b^2})e^{-c-dx}}{a} \right) \right)}{\sqrt{a^2+b^2}(-b+\sqrt{a^2+b^2})d^3} \right)$$

$$- \frac{e^2 \text{csch}(c+dx) \left(\frac{b \log(b+a \sinh(c+dx))}{a^2} - \frac{\sinh(c+dx)}{a} \right) (b+a \sinh(c+dx))}{d(a+b\text{csch}(c+dx))}$$

$$+ \frac{ef \text{csch}(c+dx)(b+a \sinh(c+dx)) \left(-2a \cosh(c+dx) - b(2c(c+dx) - (c+dx)^2 + 2(c+dx) \log(1 + \dots) \right)}{d(a+b\text{csch}(c+dx))}$$

input `Integrate[((e + f*x)^2*Cosh[c + d*x])/(a + b*Csch[c + d*x]),x]`

output

```
(f^2*Csch[c + d*x]*((4*b*x^3)/(-1 + E^(2*c)) - 2*b*x^3*Coth[c] - (6*a^2*b*(d^2*x^2*Log[1 + ((b - Sqrt[a^2 + b^2])*E^(-c - d*x))/a] - 2*d*x*PolyLog[2, ((-b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a] - 2*PolyLog[3, ((-b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a]))/(Sqrt[a^2 + b^2]*(-b + Sqrt[a^2 + b^2])*d^3) - (6*a^2*b*(d^2*x^2*Log[1 + ((b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a] - 2*d*x*PolyLog[2, -(((b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a)] - 2*PolyLog[3, -(((b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a)]))/(Sqrt[a^2 + b^2]*(b + Sqrt[a^2 + b^2])*d^3) + (6*b^2*(d^2*x^2*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) + 2*d*x*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - 2*PolyLog[3, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2])]))/(Sqrt[a^2 + b^2]*d^3) - (6*b^2*(d^2*x^2*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) + 2*d*x*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))] - 2*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]))/(Sqrt[a^2 + b^2]*d^3) + (6*a*Cosh[d*x]*(-2*d*x*Cosh[c] + (2 + d^2*x^2)*Sinh[c]))/d^3 + (6*a*((2 + d^2*x^2)*Cosh[c] - 2*d*x*Sinh[c])*Sinh[d*x])/d^3*(b + a*Sinh[c + d*x]))/(6*a^2*(a + b*Csch[c + d*x])) - (e^2*Csch[c + d*x]*((b*Log[b + a*Sinh[c + d*x]])/a^2 - Sinh[c + d*x]/a)*(b + a*Sinh[c + d*x]))/(d*(a + b*Csch[c + d*x])) + (e*f*Csch[c + d*x]*(b + a*Sinh[c + d*x])*(-2*a*Cosh[c + d*x] - b*(2*c*(c + d*x) - (c + d*x)^2 + 2*(c + d*x)*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) + 2*(c + d*x)*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) - 2*c*Log[a - 2*b*E...
```

3.18.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.74 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.577$, Rules used = {6128, 6113, 3042, 3777, 26, 3042, 26, 3777, 3042, 3117, 6095, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

$$\downarrow 6128$$

$$\int \frac{(e + fx)^2 \sinh(c + dx) \cosh(c + dx)}{a \sinh(c + dx) + b} dx$$

$$\downarrow 6113$$

$$\frac{\int (e + fx)^2 \cosh(c + dx) dx}{a} - \frac{b \int \frac{(e + fx)^2 \cosh(c + dx)}{b + a \sinh(c + dx)} dx}{a}$$

3.18. $\int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{CSch}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx)^2 \sin\left(ic+idx+\frac{\pi}{2}\right) dx}{a} \\
 & \downarrow 3777 \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2if \int -i(e+fx) \sinh(c+dx) dx}{a}}{a} \\
 & \downarrow 26 \\
 & \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2f \int (e+fx) \sinh(c+dx) dx}{a}}{a} - \frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \downarrow 3042 \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2f \int -i(e+fx) \sin(ic+idx) dx}{a}}{a} \\
 & \downarrow 26 \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \int (e+fx) \sin(ic+idx) dx}{a}}{a} \\
 & \downarrow 3777 \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \cosh(c+dx) dx}{d} \right)}{a}}{a} \\
 & \downarrow 3042 \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \sin\left(ic+idx+\frac{\pi}{2}\right) dx}{d} \right)}{a}}{a} \\
 & \downarrow 3117 \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a}}{a} \\
 & \downarrow 6095 \\
 & -\frac{b \left(\int \frac{e^{c+dx} (e+fx)^2}{e^{c+dx} a + b - \sqrt{a^2 + b^2}} dx + \int \frac{e^{c+dx} (e+fx)^2}{e^{c+dx} a + b + \sqrt{a^2 + b^2}} dx - \frac{(e+fx)^3}{3af} \right)}{a} + \\
 & \frac{\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a}}{a} \\
 & \downarrow 2620
 \end{aligned}$$

3.18. $\int \frac{(e+fx)^2 \cosh(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & b \left(-\frac{2f \int (e+fx) \log\left(\frac{e^{c+dx} a}{b-\sqrt{a^2+b^2}}+1\right) dx}{ad} - \frac{2f \int (e+fx) \log\left(\frac{e^{c+dx} a}{b+\sqrt{a^2+b^2}}+1\right) dx}{ad} + \frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}+1\right)}{ad} + \frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}+1\right)}{ad} \right) \\
 & \frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \\
 & \quad \downarrow \text{3011} \\
 & b \left(-\frac{2f \left(\frac{f \int \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right) dx}{d} - (e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right) \right)}{ad} - \frac{2f \left(\frac{f \int \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) dx}{d} - (e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) \right)}{ad} \right) \\
 & \frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \\
 & \quad \downarrow \text{2720} \\
 & b \left(-\frac{2f \left(\frac{f \int e^{-c-dx} \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right) de^{c+dx}}{d^2} - (e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right) \right)}{ad} - \frac{2f \left(\frac{f \int e^{-c-dx} \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) de^{c+dx}}{d^2} - (e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) \right)}{ad} \right) \\
 & \frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \\
 & \quad \downarrow \text{7143} \\
 & b \left(-\frac{2f \left(\frac{f \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{d^2} - (e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right) \right)}{ad} - \frac{2f \left(\frac{f \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d^2} - (e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) \right)}{ad} \right) \\
 & \frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a}
 \end{aligned}$$

input `Int[((e + f*x)^2*Cosh[c + d*x])/(a + b*CsCh[c + d*x]),x]`

3.18. $\int \frac{(e+fx)^2 \cosh(c+dx)}{a+b\text{CSch}(c+dx)} dx$

```
output -((b*(-1/3*(e + f*x)^3/(a*f) + ((e + f*x)^2*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]))/(a*d) + ((e + f*x)^2*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]))/(a*d) - (2*f*(-((e + f*x)*PolyLog[2, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d) + (f*PolyLog[3, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]))/d^2))/(a*d) - (2*f*(-((e + f*x)*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]])))/d) + (f*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]))/d^2))/(a*d)))/a + (((e + f*x)^2*Sinh[c + d*x])/d + ((2*I)*f*((I*(e + f*x)*Cosh[c + d*x])/d - (I*f*Sinh[c + d*x])/d^2))/d)/a
```

3.18.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 2620 Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 3011 Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(`
`-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*C`
`os[e + f*x], x], x] /;` `FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 6095 `Int[(Cosh[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_.)*Sin`
`h[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)),`
`x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 + b^2, 2] + b*E^(c + d*x)))`
`, x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 + b^2, 2] + b*E^(c + d*x)))`
`, x]) /;` `FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 + b^2, 0]`

rule 6113 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(p_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) +`
`(d_.)*(x_)]^(n_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := S`
`imp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^p*Sinh[c + d*x]^(n - 1), x], x] - S`
`imp[a/b Int[(e + f*x)^m*Cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*Sinh`
`[c + d*x])), x], x] /;` `FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[`
`n, 0] && IGtQ[p, 0]`

rule 6128 `Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.))/(Csch[(c_.)`
`+ (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F`
`[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /;` `FreeQ[{a, b, c, d, e, f}, x] && H`
`yperbolicQ[F] && IntegersQ[m, n]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S`
`ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /;` `FreeQ[{a, b, c, d`
`, e, n, p}, x] && EqQ[b*d, a*e]`

3.18.4 Maple [F]

$$\int \frac{(fx + e)^2 \cosh(dx + c)}{a + b \operatorname{csch}(dx + c)} dx$$

input `int((f*x+e)^2*cosh(d*x+c)/(a+b*csch(d*x+c)),x)`

output `int((f*x+e)^2*cosh(d*x+c)/(a+b*csch(d*x+c)),x)`

3.18.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1265 vs. $2(308) = 616$.

Time = 0.29 (sec) , antiderivative size = 1265, normalized size of antiderivative = 3.83

$$\int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

input `integrate((f*x+e)^2*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="fricas")`

output

```
-1/6*(3*a*d^2*f^2*x^2 + 3*a*d^2*e^2 + 6*a*d*e*f + 6*a*f^2 - 3*(a*d^2*f^2*x^2 + a*d^2*e^2 - 2*a*d*e*f + 2*a*f^2 + 2*(a*d^2*e*f - a*d*f^2)*x)*cosh(d*x + c)^2 - 3*(a*d^2*f^2*x^2 + a*d^2*e^2 - 2*a*d*e*f + 2*a*f^2 + 2*(a*d^2*e*f - a*d*f^2)*x)*sinh(d*x + c)^2 + 6*(a*d^2*e*f + a*d*f^2)*x - 2*(b*d^3*f^2*x^3 + 3*b*d^3*e*f*x^2 + 3*b*d^3*e^2*x + 6*b*c*d^2*e^2 - 6*b*c^2*d*e*f + 2*b*c^3*f^2)*cosh(d*x + c) + 12*((b*d*f^2*x + b*d*e*f)*cosh(d*x + c) + (b*d*f^2*x + b*d*e*f)*sinh(d*x + c))*dilog((b*cosh(d*x + c) + b*sinh(d*x + c) + (a*cosh(d*x + c) + a*sinh(d*x + c))*sqrt((a^2 + b^2)/a^2) - a)/a + 1) + 12*((b*d*f^2*x + b*d*e*f)*cosh(d*x + c) + (b*d*f^2*x + b*d*e*f)*sinh(d*x + c))*dilog((b*cosh(d*x + c) + b*sinh(d*x + c) - (a*cosh(d*x + c) + a*sinh(d*x + c))*sqrt((a^2 + b^2)/a^2) - a)/a + 1) + 6*((b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*cosh(d*x + c) + (b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*sinh(d*x + c))*log(2*a*cosh(d*x + c) + 2*a*sinh(d*x + c) + 2*a*sqrt((a^2 + b^2)/a^2) + 2*b) + 6*((b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*cosh(d*x + c) + (b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*sinh(d*x + c))*log(2*a*cosh(d*x + c) + 2*a*sinh(d*x + c) - 2*a*sqrt((a^2 + b^2)/a^2) + 2*b) + 6*((b*d^2*f^2*x^2 + 2*b*d^2*e*f*x + 2*b*c*d*e*f - b*c^2*f^2)*cosh(d*x + c) + (b*d^2*f^2*x^2 + 2*b*d^2*e*f*x + 2*b*c*d*e*f - b*c^2*f^2)*sinh(d*x + c))*log(-(b*cosh(d*x + c) + b*sinh(d*x + c) + (a*cosh(d*x + c) + a*sinh(d*x + c))*sqrt((a^2 + b^2)/a^2) - a)/a) + 6*((b*d^2*f^2*x^2 + 2*b*d^2*e*f*x + 2*b*c*d*e*f - b*c...
```


3.18.6 Sympy [F]

$$\int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

input `integrate((f*x+e)**2*cosh(d*x+c)/(a+b*csch(d*x+c)),x)`

output `Integral((e + f*x)**2*cosh(c + d*x)/(a + b*csch(c + d*x)), x)`

3.18.7 Maxima [F]

$$\int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^2 \cosh(dx + c)}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^2*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/2*e^2*(2*(d*x + c)*b/(a^2*d) - e^(d*x + c)/(a*d) + e^(-d*x - c)/(a*d) + 2*b*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a^2*d)) - 1/6*(2*b*d^3*f^2*x^3*e^c + 6*b*d^3*e*f*x^2*e^c - 3*(a*d^2*f^2*x^2*e^(2*c) + 2*(d^2*e*f - d*f^2)*a*x*e^(2*c) - 2*(d*e*f - f^2)*a*e^(2*c))*e^(d*x) + 3*(a*d^2*f^2*x^2 + 2*(d^2*e*f + d*f^2)*a*x + 2*(d*e*f + f^2)*a)*e^(-d*x))*e^(-c)/(a^2*d^3) + integrate(-2*(a*b*f^2*x^2 + 2*a*b*e*f*x - (b^2*f^2*x^2*e^c + 2*b^2*e*f*x*e^c))*e^(d*x))/(a^3*e^(2*d*x + 2*c) + 2*a^2*b*e^(d*x + c) - a^3), x)`

3.18.8 Giac [F]

$$\int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^2 \cosh(dx + c)}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^2*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)^2*cosh(d*x + c)/(b*csch(d*x + c) + a), x)`

3.18.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx)^2 \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx) (e + fx)^2}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)*(e + f*x)^2)/(a + b/sinh(c + d*x)),x)`output `int((cosh(c + d*x)*(e + f*x)^2)/(a + b/sinh(c + d*x)), x)`

3.19 $\int \frac{(e+fx) \cosh(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$

3.19.1	Optimal result	154
3.19.2	Mathematica [A] (verified)	154
3.19.3	Rubi [A] (verified)	155
3.19.4	Maple [B] (verified)	158
3.19.5	Fricas [B] (verification not implemented)	159
3.19.6	Sympy [F]	160
3.19.7	Maxima [F]	160
3.19.8	Giac [F]	161
3.19.9	Mupad [F(-1)]	161

3.19.1 Optimal result

Integrand size = 24, antiderivative size = 212

$$\int \frac{(e+fx) \cosh(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{b(e+fx)^2}{2a^2f} - \frac{f \cosh(c+dx)}{ad^2} - \frac{b(e+fx) \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d}$$

$$- \frac{b(e+fx) \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d} - \frac{bf \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^2d^2}$$

$$- \frac{bf \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^2d^2} + \frac{(e+fx) \sinh(c+dx)}{ad}$$

output $1/2*b*(f*x+e)^2/a^2/f-f*\cosh(d*x+c)/a/d^2-b*(f*x+e)*\ln(1+a*\exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^2/d-b*(f*x+e)*\ln(1+a*\exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^2/d-b*f*\operatorname{polylog}(2,-a*\exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^2/d^2-b*f*\operatorname{polylog}(2,-a*\exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^2/d^2+(f*x+e)*\sinh(d*x+c)/a/d$

3.19.2 Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.25

$$\int \frac{(e+fx) \cosh(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{\mathbf{csch}(c+dx)(b+a \sinh(c+dx)) \left(2de(b \log(b+a \sinh(c+dx)) - a \sinh(c+dx)) + f(2a \cosh(c+dx))\right)}{a^2d^2}$$

input `Integrate[((e + f*x)*Cosh[c + d*x])/(a + b*Csch[c + d*x]),x]`

output `-1/2*(Csch[c + d*x]*(b + a*Sinh[c + d*x])*(2*d*(b*Log[b + a*Sinh[c + d*x]] - a*Sinh[c + d*x]) + f*(2*a*Cosh[c + d*x] + b*(2*c*(c + d*x) - (c + d*x)^2 + 2*(c + d*x)*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) + 2*(c + d*x)*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) - 2*c*Log[a - 2*b*E^(c + d*x) - a*E^(2*(c + d*x))] + 2*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) + 2*PolyLog[2, -(a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])]) - 2*a*d*x*Sinh[c + d*x]))/(a^2*d^2*(a + b*Csch[c + d*x]))`

3.19.3 Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.99, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6128, 6113, 3042, 3777, 26, 3042, 26, 3118, 6095, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(e + fx) \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx \\
 & \quad \downarrow \text{6128} \\
 & \int \frac{(e + fx) \sinh(c + dx) \cosh(c + dx)}{a \sinh(c + dx) + b} dx \\
 & \quad \downarrow \text{6113} \\
 & \frac{\int (e + fx) \cosh(c + dx) dx}{a} - \frac{b \int \frac{(e + fx) \cosh(c + dx)}{b + a \sinh(c + dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b \int \frac{(e + fx) \cosh(c + dx)}{b + a \sinh(c + dx)} dx}{a} + \frac{\int (e + fx) \sin\left(ic + idx + \frac{\pi}{2}\right) dx}{a} \\
 & \quad \downarrow \text{3777} \\
 & -\frac{b \int \frac{(e + fx) \cosh(c + dx)}{b + a \sinh(c + dx)} dx}{a} + \frac{(e + fx) \sinh(c + dx)}{d} - \frac{if \int -i \sinh(c + dx) dx}{d} \\
 & \quad \downarrow \text{26} \\
 & \frac{(e + fx) \sinh(c + dx)}{d} - \frac{f \int \sinh(c + dx) dx}{d} - \frac{b \int \frac{(e + fx) \cosh(c + dx)}{b + a \sinh(c + dx)} dx}{a}
 \end{aligned}$$

3.19. $\int \frac{(e + fx) \cosh(c + dx)}{a + b \operatorname{CSch}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & -\frac{b \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int -i \sin(ic+idx) dx}{a} \\
 & \downarrow 26 \\
 & -\frac{b \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{(e+fx) \sinh(c+dx)}{d} + \frac{if \int \sin(ic+idx) dx}{a} \\
 & \downarrow 3118 \\
 & \frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} - \frac{b \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \downarrow 6095 \\
 & \frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} - \frac{b \left(\int \frac{e^{c+dx} (e+fx)}{e^{c+dx} a + b - \sqrt{a^2+b^2}} dx + \int \frac{e^{c+dx} (e+fx)}{e^{c+dx} a + b + \sqrt{a^2+b^2}} dx - \frac{(e+fx)^2}{2af} \right)}{a} \\
 & \downarrow 2620 \\
 & \frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} - \\
 & b \left(-\frac{f \int \log\left(\frac{e^{c+dx} a}{b - \sqrt{a^2+b^2}} + 1\right) dx}{ad} - \frac{f \int \log\left(\frac{e^{c+dx} a}{b + \sqrt{a^2+b^2}} + 1\right) dx}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b - \sqrt{a^2+b^2}} + 1\right)}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2} + b} + 1\right)}{ad} - \frac{(e+fx)^2}{2af} \right) \\
 & \downarrow 2715 \\
 & \frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} - \\
 & b \left(-\frac{f \int e^{-c-dx} \log\left(\frac{e^{c+dx} a}{b - \sqrt{a^2+b^2}} + 1\right) dx}{ad^2} - \frac{f \int e^{-c-dx} \log\left(\frac{e^{c+dx} a}{b + \sqrt{a^2+b^2}} + 1\right) dx}{ad^2} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b - \sqrt{a^2+b^2}} + 1\right)}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2} + b} + 1\right)}{ad} \right) \\
 & \downarrow 2838 \\
 & \frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} - \\
 & b \left(\frac{f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b - \sqrt{a^2+b^2}}\right)}{ad^2} + \frac{f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b + \sqrt{a^2+b^2}}\right)}{ad^2} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b - \sqrt{a^2+b^2}} + 1\right)}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2} + b} + 1\right)}{ad} - \frac{(e+fx)^2}{2af} \right)
 \end{aligned}$$

input `Int[((e + f*x)*Cosh[c + d*x])/(a + b*CsCh[c + d*x]),x]`

3.19. $\int \frac{(e+fx) \cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

```
output -((b*(-1/2*(e + f*x)^2/(a*f) + ((e + f*x)*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]))/(a*d) + ((e + f*x)*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]))/(a*d) + (f*PolyLog[2, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]))]/(a*d^2) + (f*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]/(a*d^2)))/a + (-((f*Cosh[c + d*x])/d^2) + ((e + f*x)*Sinh[c + d*x])/d)/a
```

3.19.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x_, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 2620 Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3118 Int[sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 3777 Int[(((c_) + (d_)*(x_))^(m_))*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

```
rule 6095 Int[(Cosh[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))^(m_))/((a_) + (b_)*Sin
h[(c_) + (d_)*(x_)]), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)),
x] + (Int[(e + f*x)^m*(E^(c + d*x))/(a - Rt[a^2 + b^2, 2] + b*E^(c + d*x))
, x] + Int[(e + f*x)^m*(E^(c + d*x))/(a + Rt[a^2 + b^2, 2] + b*E^(c + d*x))
, x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 + b^2, 0]
```

```
rule 6113 Int[(Cosh[(c_) + (d_)*(x_)]^(p_)*((e_) + (f_)*(x_))^(m_)*Sinh[(c_) +
(d_)*(x_)]^(n_))/((a_) + (b_)*Sinh[(c_) + (d_)*(x_)]), x_Symbol] := S
imp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^p*Sinh[c + d*x]^(n - 1), x], x] - S
imp[a/b Int[(e + f*x)^m*Cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*Sinh
[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[
n, 0] && IGtQ[p, 0]
```

```
rule 6128 Int[(((e_) + (f_)*(x_))^(m_)*(F_)[(c_) + (d_)*(x_)]^(n_))/(Csch[(c_)
+ (d_)*(x_)]*(b_ + (a_))), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F
[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && H
yperbolicQ[F] && IntegersQ[m, n]
```

3.19.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 482 vs. $2(198) = 396$.

Time = 1.17 (sec) , antiderivative size = 483, normalized size of antiderivative = 2.28

method	result
risch	$\frac{bf x^2}{2a^2} - \frac{bex}{a^2} + \frac{(dx f + de - f)e^{dx+c}}{2a d^2} - \frac{(dx f + de + f)e^{-dx-c}}{2a d^2} - \frac{bf \ln\left(\frac{-a e^{dx+c} + \sqrt{a^2+b^2}-b}{-b + \sqrt{a^2+b^2}}\right)x}{d a^2} - \frac{bf \ln\left(\frac{a e^{dx+c} + \sqrt{a^2+b^2}+b}{b + \sqrt{a^2+b^2}}\right)x}{d a^2}$

```
input int((f*x+e)*cosh(d*x+c)/(a+b*csch(d*x+c)),x,method=_RETURNVERBOSE)
```

output $\frac{1}{2} \frac{1}{a^2 b f x^2} - \frac{1}{a^2 b e x} + \frac{1}{2} \frac{(d f x + d e - f)}{a d^2 \exp(d x + c)} - \frac{1}{2} \frac{(d f x + d e + f)}{a d^2 \exp(-d x - c)} - \frac{1}{d} \frac{1}{a^2 b f} \ln\left(\frac{-a \exp(d x + c) + (a^2 + b^2)^{1/2} - b}{-b + (a^2 + b^2)^{1/2}}\right) x - \frac{1}{d} \frac{1}{a^2 b f} \ln\left(\frac{a \exp(d x + c) + (a^2 + b^2)^{1/2} + b}{b + (a^2 + b^2)^{1/2}}\right) x - \frac{1}{d^2} \frac{1}{a^2 b f} \operatorname{dilog}\left(\frac{-a \exp(d x + c) + (a^2 + b^2)^{1/2} - b}{-b + (a^2 + b^2)^{1/2}}\right) - \frac{1}{d^2} \frac{1}{a^2 b f} \operatorname{dilog}\left(\frac{a \exp(d x + c) + (a^2 + b^2)^{1/2} + b}{b + (a^2 + b^2)^{1/2}}\right) - \frac{1}{d} \frac{1}{a^2 b e} \ln(\exp(2 d x + 2 c) * a + 2 \exp(d x + c) * b - a) + \frac{2}{d} \frac{1}{a^2 b f c x} + \frac{1}{d^2} \frac{1}{a^2 b f c^2} - \frac{1}{d^2} \frac{1}{a^2 b f} \ln\left(\frac{-a \exp(d x + c) + (a^2 + b^2)^{1/2} - b}{-b + (a^2 + b^2)^{1/2}}\right) * c - \frac{1}{d^2} \frac{1}{a^2 b f} \ln\left(\frac{a \exp(d x + c) + (a^2 + b^2)^{1/2} + b}{b + (a^2 + b^2)^{1/2}}\right) * c + \frac{1}{d^2} \frac{1}{a^2 b c f} \ln(\exp(2 d x + 2 c) * a + 2 \exp(d x + c) * b - a) - \frac{2}{d^2} \frac{1}{a^2 b c f} \ln(\exp(d x + c)) + \frac{2}{d} \frac{1}{a^2 b e} \ln(\exp(d x + c))$

3.19.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 692 vs. $2(196) = 392$.

Time = 0.29 (sec) , antiderivative size = 692, normalized size of antiderivative = 3.26

$$\int \frac{(e + f x) \cosh(c + d x)}{a + b \operatorname{csch}(c + d x)} dx =$$

$$\frac{a d f x + a d e - (a d f x + a d e - a f) \cosh(dx + c)^2 - (a d f x + a d e - a f) \sinh(dx + c)^2 + a f - (b d^2 f x^2 +$$

input `integrate((f*x+e)*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="fricas")`

output

```
-1/2*(a*d*f*x + a*d*e - (a*d*f*x + a*d*e - a*f)*cosh(d*x + c)^2 - (a*d*f*x
+ a*d*e - a*f)*sinh(d*x + c)^2 + a*f - (b*d^2*f*x^2 + 2*b*d^2*e*x + 4*b*c
*d*e - 2*b*c^2*f)*cosh(d*x + c) + 2*(b*f*cosh(d*x + c) + b*f*sinh(d*x + c)
)*dilog((b*cosh(d*x + c) + b*sinh(d*x + c) + (a*cosh(d*x + c) + a*sinh(d*x
+ c))*sqrt((a^2 + b^2)/a^2) - a)/a + 1) + 2*(b*f*cosh(d*x + c) + b*f*sinh
(d*x + c))*dilog((b*cosh(d*x + c) + b*sinh(d*x + c) - (a*cosh(d*x + c) + a
*sinh(d*x + c))*sqrt((a^2 + b^2)/a^2) - a)/a + 1) + 2*((b*d*e - b*c*f)*cos
h(d*x + c) + (b*d*e - b*c*f)*sinh(d*x + c))*log(2*a*cosh(d*x + c) + 2*a*si
nh(d*x + c) + 2*a*sqrt((a^2 + b^2)/a^2) + 2*b) + 2*((b*d*e - b*c*f)*cosh(d
*x + c) + (b*d*e - b*c*f)*sinh(d*x + c))*log(2*a*cosh(d*x + c) + 2*a*sinh(
d*x + c) - 2*a*sqrt((a^2 + b^2)/a^2) + 2*b) + 2*((b*d*f*x + b*c*f)*cosh(d*
x + c) + (b*d*f*x + b*c*f)*sinh(d*x + c))*log(-(b*cosh(d*x + c) + b*sinh(d
*x + c) + (a*cosh(d*x + c) + a*sinh(d*x + c))*sqrt((a^2 + b^2)/a^2) - a)/a
) + 2*((b*d*f*x + b*c*f)*cosh(d*x + c) + (b*d*f*x + b*c*f)*sinh(d*x + c))*
log(-(b*cosh(d*x + c) + b*sinh(d*x + c) - (a*cosh(d*x + c) + a*sinh(d*x +
c))*sqrt((a^2 + b^2)/a^2) - a)/a - (b*d^2*f*x^2 + 2*b*d^2*e*x + 4*b*c*d*e
- 2*b*c^2*f + 2*(a*d*f*x + a*d*e - a*f)*cosh(d*x + c))*sinh(d*x + c))/(a^
2*d^2*cosh(d*x + c) + a^2*d^2*sinh(d*x + c))
```

3.19.6 Sympy [F]

$$\int \frac{(e + fx) \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx) \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

input `integrate((f*x+e)*cosh(d*x+c)/(a+b*csch(d*x+c)),x)`

output `Integral((e + f*x)*cosh(c + d*x)/(a + b*csch(c + d*x)), x)`

3.19.7 Maxima [F]

$$\int \frac{(e + fx) \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e) \cosh(dx + c)}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/2*e*(2*(d*x + c)*b/(a^2*d) - e^(d*x + c)/(a*d) + e^(-d*x - c)/(a*d) + 2*b*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a^2*d)) - 1/2*f*((b*d^2*x^2*e^c - (a*d*x*e^(2*c) - a*e^(2*c))*e^(d*x) + (a*d*x + a)*e^(-d*x))*e^(-c)/(a^2*d^2) - integrate(4*(b^2*x*e^(d*x + c) - a*b*x)/(a^3*e^(2*d*x + 2*c) + 2*a^2*b*e^(d*x + c) - a^3), x))`

3.19.8 Giac [F]

$$\int \frac{(e + fx) \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e) \cosh(dx + c)}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)*cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)*cosh(d*x + c)/(b*csch(d*x + c) + a), x)`

3.19.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx) \cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx) (e + fx)}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)*(e + f*x))/(a + b/sinh(c + d*x)),x)`

output `int((cosh(c + d*x)*(e + f*x))/(a + b/sinh(c + d*x)), x)`

3.20 $\int \frac{\cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

3.20.1	Optimal result	162
3.20.2	Mathematica [A] (verified)	162
3.20.3	Rubi [A] (verified)	163
3.20.4	Maple [A] (verified)	165
3.20.5	Fricas [B] (verification not implemented)	165
3.20.6	Sympy [F]	166
3.20.7	Maxima [B] (verification not implemented)	166
3.20.8	Giac [A] (verification not implemented)	167
3.20.9	Mupad [B] (verification not implemented)	167

3.20.1 Optimal result

Integrand size = 19, antiderivative size = 34

$$\int \frac{\cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = -\frac{b \log(b+a \sinh(c+dx))}{a^2 d} + \frac{\sinh(c+dx)}{ad}$$

output `-b*ln(b+a*sinh(d*x+c))/a^2/d+sinh(d*x+c)/a/d`

3.20.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \frac{\cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = \frac{-b \log(b+a \sinh(c+dx)) + a \sinh(c+dx)}{a^2 d}$$

input `Integrate[Cosh[c + d*x]/(a + b*Csch[c + d*x]),x]`

output `(-(b*Log[b + a*Sinh[c + d*x]]) + a*Sinh[c + d*x])/(a^2*d)`

3.20.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.579$, Rules used = {3042, 4360, 26, 26, 3042, 26, 3312, 26, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(ic+idx)}{a+ib\operatorname{csc}(ic+idx)} dx \\
 & \quad \downarrow \text{4360} \\
 & \int \frac{i\sinh(c+dx)\cosh(c+dx)}{ia\sinh(c+dx)+ib} dx \\
 & \quad \downarrow \text{26} \\
 & i \int -\frac{i\cosh(c+dx)\sinh(c+dx)}{b+a\sinh(c+dx)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\sinh(c+dx)\cosh(c+dx)}{a\sinh(c+dx)+b} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i\sin(ic+idx)\cos(ic+idx)}{b-ia\sin(ic+idx)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\cos(ic+idx)\sin(ic+idx)}{b-ia\sin(ic+idx)} dx \\
 & \quad \downarrow \text{3312} \\
 & \frac{i \int \frac{i\sinh(c+dx)}{b+a\sinh(c+dx)} d(a\sinh(c+dx))}{ad} \\
 & \quad \downarrow \text{26} \\
 & \frac{\int \frac{\sinh(c+dx)}{b+a\sinh(c+dx)} d(a\sinh(c+dx))}{ad}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 27 \\
 \frac{\int \frac{a \sinh(c+dx)}{b+a \sinh(c+dx)} d(a \sinh(c+dx))}{a^2 d} \\
 \downarrow 49 \\
 \frac{\int \left(1 - \frac{b}{b+a \sinh(c+dx)}\right) d(a \sinh(c+dx))}{a^2 d} \\
 \downarrow 2009 \\
 \frac{a \sinh(c+dx) - b \log(a \sinh(c+dx) + b)}{a^2 d}
 \end{array}$$

input `Int[Cosh[c + d*x]/(a + b*CSch[c + d*x]),x]`

output `(-(b*Log[b + a*Sinh[c + d*x]]) + a*Sinh[c + d*x])/(a^2*d)`

3.20.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3312 `Int[cos[(e_.) + (f_.)*(x_.)]*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^m*(c + (d/b)*x)^n, x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

rule 4360 `Int[(cos[(e_.) + (f_.)*(x_.)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Int[(g*Cos[e + f*x])^p*((b + a*Sin[e + f*x])^m/Sin[e + f*x]^m), x] /; FreeQ[{a, b, e, f, g, p}, x] && IntegerQ[m]`

3.20.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.44

method	result	size
derivativedivides	$-\frac{\frac{b \ln(a+b \operatorname{csch}(dx+c))}{a^2} - \frac{1}{a \operatorname{csch}(dx+c)} - \frac{b \ln(\operatorname{csch}(dx+c))}{a^2}}{d}$	49
default	$-\frac{\frac{b \ln(a+b \operatorname{csch}(dx+c))}{a^2} - \frac{1}{a \operatorname{csch}(dx+c)} - \frac{b \ln(\operatorname{csch}(dx+c))}{a^2}}{d}$	49
risch	$\frac{bx}{a^2} + \frac{e^{dx+c}}{2ad} - \frac{e^{-dx-c}}{2ad} + \frac{2bc}{da^2} - \frac{b \ln\left(e^{2dx+2c} + \frac{2b e^{dx+c}}{a} - 1\right)}{da^2}$	82

input `int(cosh(d*x+c)/(a+b*csh(d*x+c)),x,method=_RETURNVERBOSE)`

output `-1/d*(1/a^2*b*ln(a+b*csh(d*x+c))-1/a/csh(d*x+c)-1/a^2*b*ln(csh(d*x+c)))`

3.20.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(34) = 68$.

Time = 0.26 (sec) , antiderivative size = 132, normalized size of antiderivative = 3.88

$$\int \frac{\cosh(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$$

$$= \frac{2bdx \cosh(dx+c) + a \cosh(dx+c)^2 + a \sinh(dx+c)^2 - 2(b \cosh(dx+c) + b \sinh(dx+c)) \log\left(\frac{2(a \cosh(dx+c) + b \sinh(dx+c))}{\cosh(dx+c)}\right)}{2(a^2d \cosh(dx+c) + a^2d \sinh(dx+c))}$$

input `integrate(cosh(d*x+c)/(a+b*csh(d*x+c)),x, algorithm="fricas")`

3.20.
$$\int \frac{\cosh(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$$

output `1/2*(2*b*d*x*cosh(d*x + c) + a*cosh(d*x + c)^2 + a*sinh(d*x + c)^2 - 2*(b*cosh(d*x + c) + b*sinh(d*x + c))*log(2*(a*sinh(d*x + c) + b)/(cosh(d*x + c) - sinh(d*x + c))) + 2*(b*d*x + a*cosh(d*x + c))*sinh(d*x + c) - a)/(a^2*d*cosh(d*x + c) + a^2*d*sinh(d*x + c))`

3.20.6 Sympy [F]

$$\int \frac{\cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

input `integrate(cosh(d*x+c)/(a+b*csch(d*x+c)),x)`

output `Integral(cosh(c + d*x)/(a + b*csch(c + d*x)), x)`

3.20.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(34) = 68$.

Time = 0.20 (sec) , antiderivative size = 83, normalized size of antiderivative = 2.44

$$\int \frac{\cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = -\frac{(dx + c)b}{a^2 d} + \frac{e^{(dx+c)}}{2ad} - \frac{e^{(-dx-c)}}{2ad} - \frac{b \log(-2be^{(-dx-c)} + ae^{(-2dx-2c)} - a)}{a^2 d}$$

input `integrate(cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-(d*x + c)*b/(a^2*d) + 1/2*e^(d*x + c)/(a*d) - 1/2*e^(-d*x - c)/(a*d) - b*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a^2*d)`

3.20.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.76

$$\int \frac{\cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \frac{e^{(dx+c)} - e^{(-dx-c)}}{a} - \frac{2b \log(|a(e^{(dx+c)} - e^{(-dx-c)}) + 2b|)}{a^2} \quad 2d$$

input `integrate(cosh(d*x+c)/(a+b*csch(d*x+c)),x, algorithm="giac")`output `1/2*((e^(d*x + c) - e^(-d*x - c))/a - 2*b*log(abs(a*(e^(d*x + c) - e^(-d*x - c)) + 2*b))/a^2)/d`**3.20.9 Mupad [B] (verification not implemented)**

Time = 2.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

$$\int \frac{\cosh(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = -\frac{b \ln(b + a \sinh(c + dx)) - a \sinh(c + dx)}{a^2 d}$$

input `int(cosh(c + d*x)/(a + b/sinh(c + d*x)),x)`output `-(b*log(b + a*sinh(c + d*x)) - a*sinh(c + d*x))/(a^2*d)`

3.21 $\int \frac{\cosh(c+dx)}{(e+fx)(a+b\mathbf{csch}(c+dx))} dx$

3.21.1	Optimal result	168
3.21.2	Mathematica [N/A]	168
3.21.3	Rubi [N/A]	169
3.21.4	Maple [N/A] (verified)	170
3.21.5	Fricas [N/A]	170
3.21.6	Sympy [N/A]	170
3.21.7	Maxima [N/A]	171
3.21.8	Giac [N/A]	171
3.21.9	Mupad [N/A]	171

3.21.1 Optimal result

Integrand size = 26, antiderivative size = 26

$$\int \frac{\cosh(c+dx)}{(e+fx)(a+b\mathbf{csch}(c+dx))} dx = \text{Int}\left(\frac{\cosh(c+dx)\sinh(c+dx)}{(e+fx)(b+a\sinh(c+dx))}, x\right)$$

output `Unintegrable(cosh(d*x+c)*sinh(d*x+c)/(f*x+e)/(b+a*sinh(d*x+c)),x)`

3.21.2 Mathematica [N/A]

Not integrable

Time = 45.95 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{\cosh(c+dx)}{(e+fx)(a+b\mathbf{csch}(c+dx))} dx = \int \frac{\cosh(c+dx)}{(e+fx)(a+b\mathbf{csch}(c+dx))} dx$$

input `Integrate[Cosh[c + d*x]/((e + f*x)*(a + b*Csch[c + d*x])),x]`

output `Integrate[Cosh[c + d*x]/((e + f*x)*(a + b*Csch[c + d*x])), x]`

3.21.3 Rubi [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {6128, 6125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cosh(c+dx)}{(e+fx)(a+b\operatorname{csch}(c+dx))} dx$$

↓ 6128

$$\int \frac{\sinh(c+dx)\cosh(c+dx)}{(e+fx)(a\sinh(c+dx)+b)} dx$$

↓ 6125

$$\int \frac{\sinh(c+dx)\cosh(c+dx)}{(e+fx)(a\sinh(c+dx)+b)} dx$$

input `Int[Cosh[c + d*x]/((e + f*x)*(a + b*Csch[c + d*x])),x]`

output `$Aborted`

3.21.3.1 Defintions of rubi rules used

rule 6125 `Int[(((e_.) + (f_.)*(x_.))^(m_.)*(F_)[(c_.) + (d_.)*(x_.)]^(n_.)*(G_)[(c_.) + (d_.)*(x_.)]^(p_.))/((a_) + (b_.)*Sinh[(c_.) + (d_.)*(x_.)]), x_Symbol] :> Unintegrable[((e + f*x)^m*F[c + d*x]^n*G[c + d*x]^p)/(a + b*Sinh[c + d*x]), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && HyperbolicQ[F] && HyperbolicQ[G]`

rule 6128 `Int[(((e_.) + (f_.)*(x_.))^(m_.)*(F_)[(c_.) + (d_.)*(x_.)]^(n_.))/(Csch[(c_.) + (d_.)*(x_.)]*(b_.) + (a_)), x_Symbol] :> Int[(e + f*x)^m*Sinh[c + d*x]*(F[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && IntegersQ[m, n]`

3.21. $\int \frac{\cosh(c+dx)}{(e+fx)(a+b\operatorname{CSch}(c+dx))} dx$

3.21.4 Maple [N/A] (verified)

Not integrable

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(dx + c)}{(fx + e)(a + b \operatorname{csch}(dx + c))} dx$$

input `int(cosh(d*x+c)/(f*x+e)/(a+b*csch(d*x+c)),x)`output `int(cosh(d*x+c)/(f*x+e)/(a+b*csch(d*x+c)),x)`**3.21.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.31

$$\int \frac{\cosh(c + dx)}{(e + fx)(a + b \operatorname{csch}(c + dx))} dx = \int \frac{\cosh(dx + c)}{(fx + e)(b \operatorname{csch}(dx + c) + a)} dx$$

input `integrate(cosh(d*x+c)/(f*x+e)/(a+b*csch(d*x+c)),x, algorithm="fricas")`output `integral(cosh(d*x + c)/(a*f*x + a*e + (b*f*x + b*e)*csch(d*x + c)), x)`**3.21.6 Sympy [N/A]**

Not integrable

Time = 6.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{\cosh(c + dx)}{(e + fx)(a + b \operatorname{csch}(c + dx))} dx = \int \frac{\cosh(c + dx)}{(a + b \operatorname{csch}(c + dx))(e + fx)} dx$$

input `integrate(cosh(d*x+c)/(f*x+e)/(a+b*csch(d*x+c)),x)`output `Integral(cosh(c + d*x)/((a + b*csch(c + d*x))*(e + f*x)), x)`

3.21.7 Maxima [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 164, normalized size of antiderivative = 6.31

$$\int \frac{\cosh(c + dx)}{(e + fx)(a + b\operatorname{csch}(c + dx))} dx = \int \frac{\cosh(dx + c)}{(fx + e)(b\operatorname{csch}(dx + c) + a)} dx$$

input `integrate(cosh(d*x+c)/(f*x+e)/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/2*e^(-c + d*e/f)*exp_integral_e(1, (f*x + e)*d/f)/(a*f) - 1/2*e^(c - d*e/f)*exp_integral_e(1, -(f*x + e)*d/f)/(a*f) - b*log(f*x + e)/(a^2*f) + 1/2*integrate(-4*(b^2*e^(d*x + c) - a*b)/(a^3*f*x + a^3*e - (a^3*f*x*e^(2*c) + a^3*e*e^(2*c)))*e^(2*d*x) - 2*(a^2*b*f*x*e^c + a^2*b*e*e^c)*e^(d*x)), x)`

3.21.8 Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{\cosh(c + dx)}{(e + fx)(a + b\operatorname{csch}(c + dx))} dx = \int \frac{\cosh(dx + c)}{(fx + e)(b\operatorname{csch}(dx + c) + a)} dx$$

input `integrate(cosh(d*x+c)/(f*x+e)/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate(cosh(d*x + c)/((f*x + e)*(b*csch(d*x + c) + a)), x)`

3.21.9 Mupad [N/A]

Not integrable

Time = 2.32 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.15

$$\int \frac{\cosh(c + dx)}{(e + fx)(a + b\operatorname{csch}(c + dx))} dx = \int \frac{\cosh(c + dx)}{(e + fx) \left(a + \frac{b}{\sinh(c + dx)} \right)} dx$$

input `int(cosh(c + d*x)/((e + f*x)*(a + b/sinh(c + d*x))),x)`

output `int(cosh(c + d*x)/((e + f*x)*(a + b/sinh(c + d*x))), x)`

3.21. $\int \frac{\cosh(c+dx)}{(e+fx)(a+b\operatorname{CSch}(c+dx))} dx$

$$3.22 \quad \int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$$

3.22.1	Optimal result	173
3.22.2	Mathematica [B] (verified)	174
3.22.3	Rubi [C] (verified)	175
3.22.4	Maple [F]	190
3.22.5	Fricas [B] (verification not implemented)	190
3.22.6	Sympy [F]	191
3.22.7	Maxima [F]	192
3.22.8	Giac [F]	192
3.22.9	Mupad [F(-1)]	193

3.22.1 Optimal result

Integrand size = 28, antiderivative size = 696

$$\begin{aligned}
\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = & \frac{3ef^2x}{4ad^2} + \frac{3f^3x^2}{8ad^2} + \frac{(e+fx)^4}{8af} + \frac{b^2(e+fx)^4}{4a^3f} \\
& - \frac{6bf^2(e+fx) \cosh(c+dx)}{a^2d^3} - \frac{b(e+fx)^3 \cosh(c+dx)}{a^2d} \\
& - \frac{3f^3 \cosh^2(c+dx)}{8ad^4} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4ad^2} \\
& - \frac{b\sqrt{a^2+b^2}(e+fx)^3 \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d} \\
& + \frac{b\sqrt{a^2+b^2}(e+fx)^3 \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d} \\
& - \frac{3b\sqrt{a^2+b^2}f(e+fx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d^2} \\
& + \frac{3b\sqrt{a^2+b^2}f(e+fx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d^2} \\
& + \frac{6b\sqrt{a^2+b^2}f^2(e+fx) \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d^3} \\
& - \frac{6b\sqrt{a^2+b^2}f^2(e+fx) \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d^3} \\
& - \frac{6b\sqrt{a^2+b^2}f^3 \operatorname{PolyLog}\left(4, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d^4} \\
& + \frac{6b\sqrt{a^2+b^2}f^3 \operatorname{PolyLog}\left(4, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d^4} \\
& + \frac{6bf^3 \sinh(c+dx)}{a^2d^4} + \frac{3bf(e+fx)^2 \sinh(c+dx)}{a^2d^2} \\
& + \frac{3f^2(e+fx) \cosh(c+dx) \sinh(c+dx)}{4ad^3} \\
& + \frac{(e+fx)^3 \cosh(c+dx) \sinh(c+dx)}{2ad}
\end{aligned}$$

output $\frac{3}{4}ef^2x/af^2+3/8f^3x^2/af^2+1/8(fxe)^4/af+1/4b^2(fxe)^4/a^3/f-6bf^2(fxe)\cosh(dx+c)/a^2/d^3-b(fxe)^3\cosh(dx+c)/a^2/d-3/8f^3\cosh(dx+c)^2/a/d^4-3/4f(fxe)^2\cosh(dx+c)^2/a/d^2+6bf^3\sinh(dx+c)/a^2/d^4+3bf(fxe)^2\sinh(dx+c)/a^2/d^2+3/4f^2(fxe)\cosh(dx+c)\sinh(dx+c)/a/d^3+1/2(fxe)^3\cosh(dx+c)\sinh(dx+c)/a/d-b(fxe)^3\ln(1+a\exp(dx+c)/(b-(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d+b(fxe)^3\ln(1+a\exp(dx+c)/(b+(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d-3bf(fxe)^2\operatorname{polylog}(2,-a\exp(dx+c)/(b-(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d^2+3bf(fxe)^2\operatorname{polylog}(2,-a\exp(dx+c)/(b+(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d^2+6bf^2(fxe)\operatorname{polylog}(3,-a\exp(dx+c)/(b-(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d^3-6bf^2(fxe)\operatorname{polylog}(3,-a\exp(dx+c)/(b+(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d^3-6bf^3\operatorname{polylog}(4,-a\exp(dx+c)/(b-(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d^4+6bf^3\operatorname{polylog}(4,-a\exp(dx+c)/(b+(a^2+b^2)^{1/2}))\cdot(a^2+b^2)^{1/2}/a^3/d^4$

3.22.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 2218 vs. $2(696) = 1392$.

Time = 9.94 (sec) , antiderivative size = 2218, normalized size of antiderivative = 3.19

$$\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = \text{Result too large to show}$$

input `Integrate[((e + f*x)^3*Cosh[c + d*x]^2)/(a + b*CsCh[c + d*x]),x]`

output $(e^3(c/d + x - (2*b*ArcTan[(a - b*Tanh[(c + d*x)/2])/Sqrt[-a^2 - b^2]])/Sqrt[-a^2 - b^2]*d)*Csch[c + d*x]*(b + a*Sinh[c + d*x])/(4*a*(a + b*Csch[c + d*x])) + (3*e^2*f*Csch[c + d*x]*(x^2 - (2*b*(d*x*(Log[1 + (a*E^(c + d*x)))/(b - Sqrt[a^2 + b^2]]) - Log[1 + (a*E^(c + d*x)))/(b + Sqrt[a^2 + b^2]]) + PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))))/(Sqrt[a^2 + b^2]*d^2)*(b + a*Sinh[c + d*x])/(8*a*(a + b*Csch[c + d*x])) + (e*f^2*Csch[c + d*x]*(x^3 - (3*b*(d^2*x^2*Log[1 + (a*E^(c + d*x)))/(b - Sqrt[a^2 + b^2]]) - d^2*x^2*Log[1 + (a*E^(c + d*x)))/(b + Sqrt[a^2 + b^2]]) + 2*d*x*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - 2*d*x*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]) - 2*PolyLog[3, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) + 2*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))))/(Sqrt[a^2 + b^2]*d^3)*(b + a*Sinh[c + d*x])/(4*a*(a + b*Csch[c + d*x])) + (f^3*Csch[c + d*x]*(x^4 - (4*b*(d^3*x^3*Log[1 + (a*E^(c + d*x)))/(b - Sqrt[a^2 + b^2]]) - d^3*x^3*Log[1 + (a*E^(c + d*x)))/(b + Sqrt[a^2 + b^2]]) + 3*d^2*x^2*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - 3*d^2*x^2*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))] - 6*d*x*PolyLog[3, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) + 6*d*x*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]) + 6*PolyLog[4, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - 6*PolyLog[4, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))))/(Sqrt[a^2 + b^2]*d^4)*(b + a...$

3.22.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 3.56 (sec) , antiderivative size = 641, normalized size of antiderivative = 0.92, number of steps used = 31, number of rules used = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$, Rules used = {6128, 6113, 3042, 3792, 17, 3042, 3791, 17, 6099, 17, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3777, 3042, 3117, 3803, 25, 2694, 27, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)^3 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

↓ 6128

$$\int \frac{(e + fx)^3 \sinh(c + dx) \cosh^2(c + dx)}{a \sinh(c + dx) + b} dx$$

↓ 6113

$$\begin{aligned}
 & \frac{\int (e+fx)^3 \cosh^2(c+dx) dx}{a} - \frac{b \int \frac{(e+fx)^3 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{b \int \frac{(e+fx)^3 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx)^3 \sin\left(ic+idx+\frac{\pi}{2}\right)^2 dx}{a} \\
 & \quad \downarrow \text{3792} \\
 & \frac{3f^2 \int (e+fx) \cosh^2(c+dx) dx}{2d^2} + \frac{1}{2} \int (e+fx)^3 dx - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} \\
 & \quad \frac{a}{b \int \frac{(e+fx)^3 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
 & \quad \downarrow \text{17} \\
 & \frac{3f^2 \int (e+fx) \cosh^2(c+dx) dx}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f} \\
 & \quad \frac{a}{b \int \frac{(e+fx)^3 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{b \int \frac{(e+fx)^3 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
 & \frac{3f^2 \int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right)^2 dx}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f} \\
 & \quad \downarrow \text{3791} \\
 & \frac{3f^2 \left(\frac{1}{2} \int (e+fx) dx - \frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f} \\
 & \quad \frac{a}{b \int \frac{(e+fx)^3 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
 & \quad \downarrow \text{17} \\
 & \frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f} \\
 & \quad \frac{a}{b \int \frac{(e+fx)^3 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
 & \quad \downarrow \text{6099}
 \end{aligned}$$

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \int (e+fx)^3 dx}{a^2} + \frac{\int (e+fx)^3 \sinh(c+dx) dx}{a} \right)$$

a
↓ 17

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b+a \sinh(c+dx)} dx}{a^2} + \frac{\int (e+fx)^3 \sinh(c+dx) dx}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 3042

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} + \frac{\int -i(e+fx)^3 \sin(ic+idx) dx}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 26

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \int (e+fx)^3 \sin(ic+idx) dx}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 3777

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - i \left(\frac{(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \int (e+fx)^2 \cosh(c+dx) dx}{d} \right) - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

↓ 3042

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{i(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \int (e+fx)^2 \sin(ic+idx + \frac{\pi}{2}) dx}{d} \right)}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 3777

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{i(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2if \int -i(e+fx) \sinh(c+dx) dx}{d} \right)}{d} \right)}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 26

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{i(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2f \int (e+fx) \sinh(c+dx) dx}{d} \right)}{d} \right)}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 3042

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{i(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2f \int -i(e+fx) \sin(ic+idx) dx}{d} \right)}{d} \right)}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b \operatorname{csch}(c+dx)} dx$

↓ 26

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{i(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \int (e+fx) \sin(ic+idx) dx}{d} \right)}{d} \right)}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 3777

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{i(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \cosh(c+dx) dx}{d} \right)}{d} \right)}{d} \right)}{a} - \frac{b(e+fx)^4}{4a^2 f} \right)$$

a
↓ 3042

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \sin\left(\frac{ic+idx+\frac{\pi}{2}}{d}\right) dx}{d} \right)}{d} \right)}{d} \right)}{a} \right) - b(\dots)$$

a
 \downarrow 3117

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{b(e+fx)^4}{4a^2 f} - \frac{i \left(\frac{(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh\left(\frac{c+dx}{d}\right)}{d^2} \right)}{d} \right)}{d} \right)}{a} \right)$$

a
 \downarrow 3803

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{2(a^2+b^2) \int -\frac{e^{c+dx} (e+fx)^3}{-e^{2(c+dx)} a+a-2be^{c+dx}} dx}{a^2} - \frac{b(e+fx)^4}{4a^2 f} - \frac{i \left(\frac{(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{d} \right)}{d} \right)}{a} \right)$$

a

↓ 25

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{2(a^2+b^2) \int \frac{e^{c+dx} (e+fx)^3}{-e^{2(c+dx)} a+a-2be^{c+dx}} dx}{a^2} - \frac{b(e+fx)^4}{4a^2 f} - \frac{i \left(\frac{(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{d} \right)}{d} \right)}{a} \right)$$

a

↓ 2694

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{2(a^2+b^2) \left(\frac{a \int -\frac{e^{c+dx} (e+fx)^3}{2(e^{c+dx} a+b-\sqrt{a^2+b^2})} dx}{\sqrt{a^2+b^2}} - \frac{a \int -\frac{e^{c+dx} (e+fx)^3}{2(e^{c+dx} a+b+\sqrt{a^2+b^2})} dx}{\sqrt{a^2+b^2}} \right)}{a^2} - \frac{b(e+fx)^4}{4a^2 f} - \frac{i \left(\frac{(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{d} \right)}{d} \right)}{a} \right)$$

a

↓ 27

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{2(a^2+b^2)}{a^2} \left(\frac{a \int \frac{e^{c+dx} (e+fx)^3}{e^{c+dx} a + b + \sqrt{a^2+b^2}} dx}{2\sqrt{a^2+b^2}} - \frac{a \int \frac{e^{c+dx} (e+fx)^3}{e^{c+dx} a + b - \sqrt{a^2+b^2}} dx}{2\sqrt{a^2+b^2}} \right) - \frac{b(e+fx)^4}{4a^2 f} - \frac{i \left(\frac{(e+fx)^3 \cosh(c+dx)}{d} - \frac{3if \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\dots \right)}{d} \right)}{a} \right)}{a} \right)$$

↓ 2620

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$b \left(\frac{2(a^2+b^2)}{a^2} \left(\frac{a \left(\frac{(e+fx)^3 \log \left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b} + 1 \right) - 3f \int (e+fx)^2 \log \left(\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}} + 1 \right) dx}{2\sqrt{a^2+b^2}} \right) - \frac{a \left(\frac{(e+fx)^3 \log \left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} + 1 \right) - 3f \int (e+fx)^2 \log \left(\frac{e^{c+dx} a}{b-\sqrt{a^2+b^2}} + 1 \right) dx}{2\sqrt{a^2+b^2}} \right)}{a^2} \right)$$

↓ 3011

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$\frac{2(a^2+b^2)}{2\sqrt{a^2+b^2}} \left(\frac{a}{ad} \left(\frac{(e+fx)^3 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}} + 1\right)}{ad} - \frac{3f \left(\frac{2f \int (e+fx) \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) dx}{a} - \frac{(e+fx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a} \right)}{ad} \right) \right) - \frac{a}{a} \left(\frac{(e+fx)^3 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}} + 1\right)}{ad} \right)$$

$$\frac{b}{a^2}$$

↓ 7163

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$\frac{2(a^2+b^2)}{2\sqrt{a^2+b^2}} \left(\frac{a}{ad} \left(\frac{(e+fx)^3 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}} + 1\right)}{ad} - \frac{2f \left(\frac{(e+fx) \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d} - \frac{f \int \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) dx}{d} \right)}{d} - \frac{(e+fx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d} \right) \right)$$

↓ 2720

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right)}{2d^2} - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}$$

$$\left(\frac{(e+fx)^3 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b} + 1\right)}{ad} - \frac{2f \left(\frac{(e+fx) \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d} - \frac{f \int e^{-c-dx} \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) de^{c+dx}}{d^2} \right)}{d} \right) (e+fx)^2$$

$$\frac{2(a^2+b^2)}{2\sqrt{a^2+b^2}}$$

$$b$$

↓ 7143

3.22. $\int \frac{(e+fx)^3 \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\frac{3f^2 \left(-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \right) - \frac{3f(e+fx)^2 \cosh^2(c+dx)}{4d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^4}{8f}}{2d^2}$$

$$\frac{2(a^2+b^2) \left(\frac{(e+fx)^3 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b}+1\right)}{ad} - \frac{2f \left(\frac{(e+fx) \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d} - \frac{f \operatorname{PolyLog}\left(4, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d^2} \right)}{ad} - \frac{(e+fx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d} \right)}{2\sqrt{a^2+b^2}}$$

```
input Int[((e + f*x)^3*Cosh[c + d*x]^2)/(a + b*CSch[c + d*x]),x]
```

```
output ((e + f*x)^4/(8*f) - (3*f*(e + f*x)^2*Cosh[c + d*x]^2)/(4*d^2) + ((e + f*x)
)^3*Cosh[c + d*x]*Sinh[c + d*x])/(2*d) + (3*f^2*((e + f*x)^2/(4*f) - (f*Co
sh[c + d*x]^2)/(4*d^2) + ((e + f*x)*Cosh[c + d*x]*Sinh[c + d*x])/(2*d)))/(
2*d^2))/a - (b*(-1/4*(b*(e + f*x)^4)/(a^2*f) - (2*(a^2 + b^2)*(-1/2*(a*((
e + f*x)^3*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/(a*d) - (3*f*(-
((e + f*x)^2*PolyLog[2, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d) + (
2*f*((e + f*x)*PolyLog[3, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d -
(f*PolyLog[4, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d^2))/d)/(a*d))
/Sqrt[a^2 + b^2] + (a*((e + f*x)^3*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2
+ b^2]])))/(a*d) - (3*f*(-((e + f*x)^2*PolyLog[2, -((a*E^(c + d*x))/(b + S
qrt[a^2 + b^2]])))/d) + (2*f*((e + f*x)*PolyLog[3, -((a*E^(c + d*x))/(b +
Sqrt[a^2 + b^2]])))/d - (f*PolyLog[4, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b
^2]])))/d^2))/d)/(a*d)))/(2*Sqrt[a^2 + b^2]))/a^2 - (I*(I*(e + f*x)^3*C
osh[c + d*x])/d - ((3*I)*f*((e + f*x)^2*Sinh[c + d*x])/d + ((2*I)*f*((I*(
e + f*x)*Cosh[c + d*x])/d - (I*f*Sinh[c + d*x])/d^2))/d))/a
```

3.2.2.3.1 Defintions of rubi rules used

```
rule 17 Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1
))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 2620 Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/
((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.))), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
))))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2694 `Int[((F_)^(u_)*((f_) + (g_)*(x_)^(m_)))/((a_) + (b_)*(F_)^(u_) + (c_) * (F_)^(v_)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[2*(c/q) Int [(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) * (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^(c_)*((a_) + (b_)*(x_)))^(n_)]*((f_) + (g_) * (x_)^(m_)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_) + (d_)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_) + (d_)*(x_)^(m_))*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[(-c + d*x)^m*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3791 `Int[((c_) + (d_)*(x_))*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x] * ((b*Sin[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)*(b*Sin[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

rule 3792 `Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*m*(c + d*x)^(m - 1)*((b*Sine + f*x)^(n)/(f^2*n^2)), x] + (-Simp[b*(c + d*x)^m*cos[e + f*x]*((b*Sine + f*x)^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)^m*(b*Sine + f*x)^(n - 2), x], x] - Simp[d^2*m*((m - 1)/(f^2*n^2)) Int[(c + d*x)^(m - 2)*(b*Sine + f*x)^(n), x], x] /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]`

rule 3803 `Int[((c_.) + (d_.)*(x_))^(m_)/((a_) + (b_.)*sin[(e_.) + (Complex[0, fz_]*) (f_.)*(x_)]), x_Symbol] := Simp[2 Int[(c + d*x)^m*(E^((-I)*e + f*fz*x)/((-I)*b + 2*a*E^((-I)*e + f*fz*x) + I*b*E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]`

rule 6099 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(n_)*((e_.) + (f_.)*(x_))^(m_))/((a_) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-a/b^2 Int[(e + f*x)^m*Cosh[c + d*x]^(n - 2), x], x] + (Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^(n - 2)*Sinh[c + d*x], x], x] + Simp[(a^2 + b^2)/b^2 Int[(e + f*x)^m*(Cosh[c + d*x]^(n - 2)/(a + b*Sinh[c + d*x]))], x], x) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[n, 1] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

rule 6113 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(p_)*((e_.) + (f_.)*(x_))^(m_)*Sinh[(c_.) + (d_.)*(x_)]^(n_))/((a_) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^p*Sinh[c + d*x]^(n - 1), x], x] - Simp[a/b Int[(e + f*x)^m*Cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*Sinh[c + d*x]))], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6128 `Int[(((e_.) + (f_.)*(x_))^(m_)*(F_)[(c_.) + (d_.)*(x_)]^(n_))/(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && IntegersQ[m, n]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

3.22.4 Maple [F]

$$\int \frac{(fx + e)^3 \cosh(dx + c)^2}{a + b \operatorname{csch}(dx + c)} dx$$

input `int((f*x+e)^3*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x)`

output `int((f*x+e)^3*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x)`

3.22.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3847 vs. $2(638) = 1276$.

Time = 0.35 (sec) , antiderivative size = 3847, normalized size of antiderivative = 5.53

$$\int \frac{(e + fx)^3 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

input `integrate((f*x+e)^3*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="fracas")`

```

output -1/32*(4*a^2*d^3*f^3*x^3 + 4*a^2*d^3*e^3 + 6*a^2*d^2*e^2*f + 6*a^2*d*e*f^2
+ 3*a^2*f^3 - (4*a^2*d^3*f^3*x^3 + 4*a^2*d^3*e^3 - 6*a^2*d^2*e^2*f + 6*a^
2*d*e*f^2 - 3*a^2*f^3 + 6*(2*a^2*d^3*e*f^2 - a^2*d^2*f^3)*x^2 + 6*(2*a^2*d
^3*e^2*f - 2*a^2*d^2*e*f^2 + a^2*d*f^3)*x)*cosh(d*x + c)^4 - (4*a^2*d^3*f^
3*x^3 + 4*a^2*d^3*e^3 - 6*a^2*d^2*e^2*f + 6*a^2*d*e*f^2 - 3*a^2*f^3 + 6*(2
*a^2*d^3*e*f^2 - a^2*d^2*f^3)*x^2 + 6*(2*a^2*d^3*e^2*f - 2*a^2*d^2*e*f^2 +
a^2*d*f^3)*x)*sinh(d*x + c)^4 + 16*(a*b*d^3*f^3*x^3 + a*b*d^3*e^3 - 3*a*b
*d^2*e^2*f + 6*a*b*d*e*f^2 - 6*a*b*f^3 + 3*(a*b*d^3*e*f^2 - a*b*d^2*f^3)*x
^2 + 3*(a*b*d^3*e^2*f - 2*a*b*d^2*e*f^2 + 2*a*b*d*f^3)*x)*cosh(d*x + c)^3
+ 4*(4*a*b*d^3*f^3*x^3 + 4*a*b*d^3*e^3 - 12*a*b*d^2*e^2*f + 24*a*b*d*e*f^2
- 24*a*b*f^3 + 12*(a*b*d^3*e*f^2 - a*b*d^2*f^3)*x^2 + 12*(a*b*d^3*e^2*f -
2*a*b*d^2*e*f^2 + 2*a*b*d*f^3)*x - (4*a^2*d^3*f^3*x^3 + 4*a^2*d^3*e^3 - 6
*a^2*d^2*e^2*f + 6*a^2*d*e*f^2 - 3*a^2*f^3 + 6*(2*a^2*d^3*e*f^2 - a^2*d^2*
f^3)*x^2 + 6*(2*a^2*d^3*e^2*f - 2*a^2*d^2*e*f^2 + a^2*d*f^3)*x)*cosh(d*x +
c))*sinh(d*x + c)^3 + 6*(2*a^2*d^3*e*f^2 + a^2*d^2*f^3)*x^2 - 4*((a^2 + 2
*b^2)*d^4*f^3*x^4 + 4*(a^2 + 2*b^2)*d^4*e*f^2*x^3 + 6*(a^2 + 2*b^2)*d^4*e^
2*f*x^2 + 4*(a^2 + 2*b^2)*d^4*e^3*x)*cosh(d*x + c)^2 - 2*(2*(a^2 + 2*b^2)*
d^4*f^3*x^4 + 8*(a^2 + 2*b^2)*d^4*e*f^2*x^3 + 12*(a^2 + 2*b^2)*d^4*e^2*f*x
^2 + 8*(a^2 + 2*b^2)*d^4*e^3*x + 3*(4*a^2*d^3*f^3*x^3 + 4*a^2*d^3*e^3 - 6*
a^2*d^2*e^2*f + 6*a^2*d*e*f^2 - 3*a^2*f^3 + 6*(2*a^2*d^3*e*f^2 - a^2*d^...

```

3.22.6 Sympy [F]

$$\int \frac{(e + fx)^3 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx)^3 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

```
input integrate((f*x+e)**3*cosh(d*x+c)**2/(a+b*csch(d*x+c)),x)
```

```
output Integral((e + f*x)**3*cosh(c + d*x)**2/(a + b*csch(c + d*x)), x)
```


3.22.7 Maxima [F]

$$\int \frac{(e + fx)^3 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^3 \cosh(dx + c)^2}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^3*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output

```
-1/8*e^3*((4*b*e^(-d*x - c) - a)*e^(2*d*x + 2*c)/(a^2*d) - 4*(a^2 + 2*b^2)
*(d*x + c)/(a^3*d) + (4*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c))/(a^2*d) + 8*(
a^2*b + b^3)*log((a*e^(-d*x - c) - b - sqrt(a^2 + b^2))/(a*e^(-d*x - c) -
b + sqrt(a^2 + b^2)))/(sqrt(a^2 + b^2)*a^3*d) + 1/32*(4*(a^2*d^4*f^3*e^(2
*c) + 2*b^2*d^4*f^3*e^(2*c))*x^4 + 16*(a^2*d^4*e*f^2*e^(2*c) + 2*b^2*d^4*e
*f^2*e^(2*c))*x^3 + 24*(a^2*d^4*e^2*f*e^(2*c) + 2*b^2*d^4*e^2*f*e^(2*c))*x
^2 + (4*a^2*d^3*f^3*x^3*e^(4*c) + 6*(2*d^3*e*f^2 - d^2*f^3)*a^2*x^2*e^(4*c
) + 6*(2*d^3*e^2*f - 2*d^2*e*f^2 + d*f^3)*a^2*x*e^(4*c) - 3*(2*d^2*e^2*f -
2*d*e*f^2 + f^3)*a^2*e^(4*c))*e^(2*d*x) - 16*(a*b*d^3*f^3*x^3*e^(3*c) + 3
*(d^3*e*f^2 - d^2*f^3)*a*b*x^2*e^(3*c) + 3*(d^3*e^2*f - 2*d^2*e*f^2 + 2*d*
f^3)*a*b*x*e^(3*c) - 3*(d^2*e^2*f - 2*d*e*f^2 + 2*f^3)*a*b*e^(3*c))*e^(d*x
) - 16*(a*b*d^3*f^3*x^3*e^c + 3*(d^3*e*f^2 + d^2*f^3)*a*b*x^2*e^c + 3*(d^3
*e^2*f + 2*d^2*e*f^2 + 2*d*f^3)*a*b*x*e^c + 3*(d^2*e^2*f + 2*d*e*f^2 + 2*f
^3)*a*b*e^c)*e^(-d*x) - (4*a^2*d^3*f^3*x^3 + 6*(2*d^3*e*f^2 + d^2*f^3)*a^2
*x^2 + 6*(2*d^3*e^2*f + 2*d^2*e*f^2 + d*f^3)*a^2*x + 3*(2*d^2*e^2*f + 2*d*
e*f^2 + f^3)*a^2)*e^(-2*d*x))*e^(-2*c)/(a^3*d^4) - integrate(2*((a^2*b*f^3
*e^c + b^3*f^3*e^c)*x^3 + 3*(a^2*b*e*f^2*e^c + b^3*e*f^2*e^c)*x^2 + 3*(a^2
*b*e^2*f*e^c + b^3*e^2*f*e^c)*x)*e^(d*x)/(a^4*e^(2*d*x + 2*c) + 2*a^3*b*e^
(d*x + c) - a^4), x)
```

3.22.8 Giac [F]

$$\int \frac{(e + fx)^3 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^3 \cosh(dx + c)^2}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^3*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)^3*cosh(d*x + c)^2/(b*csch(d*x + c) + a), x)`

3.22.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx)^3 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx)^2 (e + fx)^3}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)^2*(e + f*x)^3)/(a + b/sinh(c + d*x)),x)`output `int((cosh(c + d*x)^2*(e + f*x)^3)/(a + b/sinh(c + d*x)), x)`

3.23 $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$

3.23.1	Optimal result	194
3.23.2	Mathematica [B] (verified)	195
3.23.3	Rubi [C] (verified)	196
3.23.4	Maple [F]	206
3.23.5	Fricas [B] (verification not implemented)	207
3.23.6	Sympy [F]	207
3.23.7	Maxima [F]	208
3.23.8	Giac [F]	208
3.23.9	Mupad [F(-1)]	209

3.23.1 Optimal result

Integrand size = 28, antiderivative size = 510

$$\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{f^2x}{4ad^2} + \frac{(e+fx)^3}{6af} + \frac{b^2(e+fx)^3}{3a^3f} - \frac{2bf^2 \cosh(c+dx)}{a^2d^3}$$

$$- \frac{b(e+fx)^2 \cosh(c+dx)}{a^2d} - \frac{f(e+fx) \cosh^2(c+dx)}{2ad^2}$$

$$- \frac{b\sqrt{a^2+b^2}(e+fx)^2 \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d}$$

$$+ \frac{b\sqrt{a^2+b^2}(e+fx)^2 \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d}$$

$$- \frac{2b\sqrt{a^2+b^2}f(e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d^2}$$

$$+ \frac{2b\sqrt{a^2+b^2}f(e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d^2}$$

$$+ \frac{2b\sqrt{a^2+b^2}f^2 \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d^3}$$

$$- \frac{2b\sqrt{a^2+b^2}f^2 \text{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d^3}$$

$$+ \frac{2bf(e+fx) \sinh(c+dx)}{a^2d^2} + \frac{f^2 \cosh(c+dx) \sinh(c+dx)}{4ad^3}$$

$$+ \frac{(e+fx)^2 \cosh(c+dx) \sinh(c+dx)}{2ad}$$

output $\frac{1}{4}f^2x/a/d^2+1/6*(f*x+e)^3/a/f+1/3*b^2*(f*x+e)^3/a^3/f-2*b*f^2*\cosh(d*x+c)/a^2/d^3-b*(f*x+e)^2*\cosh(d*x+c)/a^2/d-1/2*f*(f*x+e)*\cosh(d*x+c)^2/a/d^2+2*b*f*(f*x+e)*\sinh(d*x+c)/a^2/d^2+1/4*f^2*\cosh(d*x+c)*\sinh(d*x+c)/a/d^3+1/2*(f*x+e)^2*\cosh(d*x+c)*\sinh(d*x+c)/a/d-b*(f*x+e)^2*\ln(1+a*\exp(d*x+c)/(b-(a^2+b^2)^{(1/2)}))*(a^2+b^2)^{(1/2)}/a^3/d+b*(f*x+e)^2*\ln(1+a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2)}))*(a^2+b^2)^{(1/2)}/a^3/d-2*b*f*(f*x+e)*\text{polylog}(2,-a*\exp(d*x+c)/(b-(a^2+b^2)^{(1/2)}))*(a^2+b^2)^{(1/2)}/a^3/d^2+2*b*f*(f*x+e)*\text{polylog}(2,-a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2)}))*(a^2+b^2)^{(1/2)}/a^3/d^2+2*b*f^2*\text{polylog}(3,-a*\exp(d*x+c)/(b-(a^2+b^2)^{(1/2)}))*(a^2+b^2)^{(1/2)}/a^3/d^3-2*b*f^2*\text{polylog}(3,-a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2)}))*(a^2+b^2)^{(1/2)}/a^3/d^3$

3.23.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1216 vs. $2(510) = 1020$.

Time = 4.93 (sec) , antiderivative size = 1216, normalized size of antiderivative = 2.38

$$\int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

$$\operatorname{csch}(c + dx)(b + a \sinh(c + dx)) \left(6a^2 e^2 \left(\frac{c}{d} + x - \frac{2b \arctan\left(\frac{a - b \tanh\left(\frac{1}{2}(c + dx)\right)}{\sqrt{-a^2 - b^2}}\right)}{\sqrt{-a^2 - b^2}d} \right) + 6a^2 e f \left(x^2 - \frac{2b(dx \log(1 + \dots))}{\dots} \right) \right)$$

input `Integrate[((e + f*x)^2*Cosh[c + d*x]^2)/(a + b*Csch[c + d*x]),x]`

output

```
(Csch[c + d*x]*(b + a*Sinh[c + d*x])*(6*a^2*e^2*(c/d + x - (2*b*ArcTan[(a - b*Tanh[(c + d*x)/2])/Sqrt[-a^2 - b^2]])/Sqrt[-a^2 - b^2]*d)) + 6*a^2*e*f*(x^2 - (2*b*(d*x*(Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) - Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) + PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]))/Sqrt[a^2 + b^2]*d^2) + 2*a^2*f^2*(x^3 - (3*b*(d^2*x^2*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) - d^2*x^2*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) + 2*d*x*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - 2*d*x*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))] - 2*PolyLog[3, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) + 2*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]))/Sqrt[a^2 + b^2]*d^3) + f^2*(2*(a^2 + 4*b^2)*x^3 - (6*b*(3*a^2 + 4*b^2)*(d^2*x^2*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) - d^2*x^2*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) + 2*d*x*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - 2*d*x*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))] - 2*PolyLog[3, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) + 2*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]))/Sqrt[a^2 + b^2]*d^3 - (24*a*b*Cosh[d*x]*((2 + d^2*x^2)*Cosh[c] - 2*d*x*Sinh[c]))/d^3 + (3*a^2*Cosh[2*d*x]*(-2*d*x*Cosh[2*c] + (1 + 2*d^2*x^2)*Sinh[2*c]))/d^3 - (24*a*b*(-2*d*x*Cosh[c] + (2 + d^2*x^2)*Sinh[c])*Sinh[d*x])/d^3 + (3*a^2*((1 + 2*d^2*x^2)*Cosh[2*c] - 2*d*x*Sinh[2*c])*Sinh[2*...
```

3.23.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 2.83 (sec) , antiderivative size = 484, normalized size of antiderivative = 0.95, number of steps used = 28, number of rules used = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.964$, Rules used = {6128, 6113, 3042, 3792, 17, 3042, 3115, 24, 6099, 17, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3118, 3803, 25, 2694, 27, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

↓ 6128

$$\int \frac{(e + fx)^2 \sinh(c + dx) \cosh^2(c + dx)}{a \sinh(c + dx) + b} dx$$

↓ 6113

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
& \frac{\int (e+fx)^2 \cosh^2(c+dx) dx}{a} - \frac{b \int \frac{(e+fx)^2 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
& \quad \downarrow \text{3042} \\
& - \frac{b \int \frac{(e+fx)^2 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx)^2 \sin\left(ic+idx+\frac{\pi}{2}\right)^2 dx}{a} \\
& \quad \downarrow \text{3792} \\
& \frac{f^2 \int \cosh^2(c+dx) dx}{2d^2} + \frac{1}{2} \int (e+fx)^2 dx - \frac{f(e+fx) \cosh^2(c+dx)}{2d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh(c+dx)}{2d} \\
& \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
& \quad \downarrow \text{17} \\
& \frac{f^2 \int \cosh^2(c+dx) dx}{2d^2} - \frac{f(e+fx) \cosh^2(c+dx)}{2d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f} \\
& \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
& \quad \downarrow \text{3042} \\
& - \frac{b \int \frac{(e+fx)^2 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
& \frac{f^2 \int \sin\left(ic+idx+\frac{\pi}{2}\right)^2 dx}{2d^2} - \frac{f(e+fx) \cosh^2(c+dx)}{2d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f} \\
& \quad \downarrow \text{3115} \\
& \frac{f^2 \left(\frac{\int 1 dx}{2} + \frac{\sinh(c+dx) \cosh(c+dx)}{2d} \right)}{2d^2} - \frac{f(e+fx) \cosh^2(c+dx)}{2d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f} \\
& \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
& \quad \downarrow \text{24} \\
& - \frac{f(e+fx) \cosh^2(c+dx)}{2d^2} + \frac{f^2 \left(\frac{\sinh(c+dx) \cosh(c+dx)}{2d} + \frac{x}{2} \right)}{2d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f} \\
& \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx} \\
& \quad \downarrow \text{6099}
\end{aligned}$$

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{a} \\
 & \quad b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\int(e+fx)^2dx}{a^2} + \frac{\int(e+fx)^2\sinh(c+dx)dx}{a}\right) \\
 & \quad \downarrow 17 \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{a} \\
 & \quad b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b+a\sinh(c+dx)}dx}{a^2} + \frac{\int(e+fx)^2\sinh(c+dx)dx}{a} - \frac{b(e+fx)^3}{3a^2f}\right) \\
 & \quad \downarrow 3042 \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{a} \\
 & \quad b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} + \frac{\int-i(e+fx)^2\sin(ic+idx)dx}{a} - \frac{b(e+fx)^3}{3a^2f}\right) \\
 & \quad \downarrow 26 \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{a} \\
 & \quad b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{i\int(e+fx)^2\sin(ic+idx)dx}{a} - \frac{b(e+fx)^3}{3a^2f}\right) \\
 & \quad \downarrow 3777 \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{a} \\
 & \quad b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{i\left(\frac{(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\int(e+fx)\cosh(c+dx)dx}{d}\right)}{a} - \frac{b(e+fx)^3}{3a^2f}\right) \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\int(e+fx)\sin\left(ic+idx+\frac{\pi}{2}\right)dx}{d}\right)}{a} - \frac{b(e+fx)^3}{3a^2f}\right)} \\
 & \quad \downarrow \text{3777} \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d} - \frac{if\int-i\sinh(c+dx)dx}{d}\right)}{a}\right)}{a} - \frac{b(e+fx)^3}{3a^2f}\right)} \\
 & \quad \downarrow \text{26} \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d} - \frac{f\int\sinh(c+dx)dx}{d}\right)}{a}\right)}{a} - \frac{b(e+fx)^3}{3a^2f}\right)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d} - \frac{f\int-i\sin(ic+idx)dx}{d}\right)}{a}\right)}{a} - \frac{b(e+fx)^3}{3a^2f}\right)} \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d} + \frac{if\int\sin(ic+idx)dx}{d}\right)}{d}\right)}{a} - \frac{b(e+fx)^3}{3a^2f}\right)} \\
 & \qquad \qquad \qquad \downarrow \mathbf{3118} \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2}{b-ia\sin(ic+idx)}dx}{a^2} - \frac{b(e+fx)^3}{3a^2f} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d} - \frac{f\cosh(c+dx)}{d^2}\right)}{d}\right)}{a}\right)} \\
 & \qquad \qquad \qquad \downarrow \mathbf{3803} \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(\frac{2(a^2+b^2)\int-\frac{e^{c+dx}(e+fx)^2}{-e^{2(c+dx)}a+a-2be^{c+dx}}dx}{a^2} - \frac{b(e+fx)^3}{3a^2f} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d} - \frac{f\cosh(c+dx)}{d^2}\right)}{d}\right)}{a}\right)} \\
 & \qquad \qquad \qquad \downarrow \mathbf{25} \\
 & \frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b\left(-\frac{2(a^2+b^2)\int\frac{e^{c+dx}(e+fx)^2}{-e^{2(c+dx)}a+a-2be^{c+dx}}dx}{a^2} - \frac{b(e+fx)^3}{3a^2f} - \frac{i\left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d} - \frac{f\cosh(c+dx)}{d^2}\right)}{d}\right)}{a}\right)} \\
 & \qquad \qquad \qquad \downarrow \mathbf{2694}
 \end{aligned}$$

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b \left(\frac{2(a^2+b^2) \left(\frac{a \int -\frac{e^{c+dx}(e+fx)^2}{2(e^{c+dx}a+b-\sqrt{a^2+b^2})} dx}{\sqrt{a^2+b^2}} - \frac{a \int -\frac{e^{c+dx}(e+fx)^2}{2(e^{c+dx}a+b+\sqrt{a^2+b^2})} dx}{\sqrt{a^2+b^2}} \right)}{a^2} - \frac{b(e+fx)^3}{3a^2f} - \frac{i \left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d}\right)}{a} \right)}{a} \right)}$$

27

$$\frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b \left(\frac{2(a^2+b^2) \left(\frac{a \int \frac{e^{c+dx}(e+fx)^2}{e^{c+dx}a+b+\sqrt{a^2+b^2}} dx}{2\sqrt{a^2+b^2}} - \frac{a \int \frac{e^{c+dx}(e+fx)^2}{e^{c+dx}a+b-\sqrt{a^2+b^2}} dx}{2\sqrt{a^2+b^2}} \right)}{a^2} - \frac{b(e+fx)^3}{3a^2f} - \frac{i \left(\frac{i(e+fx)^2\cosh(c+dx)}{d} - \frac{2if\left(\frac{(e+fx)\sinh(c+dx)}{d}\right) - \frac{f\cosh(c+dx)}{d}}{a} \right)}{a} \right)}$$

2620

$$\frac{-\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f}}{b \left(\frac{2(a^2+b^2) \left(\frac{a \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b} + 1\right)}{ad} - \frac{2f \int (e+fx) \log\left(\frac{e^{c+dx}a}{b+\sqrt{a^2+b^2}} + 1\right) dx}{ad} \right)}{2\sqrt{a^2+b^2}} - \frac{a \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} + 1\right)}{ad} - \frac{2f \int (e+fx) \log\left(\frac{e^{c+dx}a}{b-\sqrt{a^2+b^2}} + 1\right) dx}{ad} \right)}{2\sqrt{a^2+b^2}} \right)}{a^2} \right)}$$

3011

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f} \\
 & \left(\frac{a}{2(a^2+b^2)} \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}} + 1\right)}{ad} - \frac{2f\left(\frac{f \int \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) dx}{d} - \frac{(e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d}\right)}{ad} \right) \right) - \frac{a}{2\sqrt{a^2+b^2}} \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{ad} \right) \right) \\
 & \frac{b}{a^2}
 \end{aligned}$$

↓ 2720

$$\begin{aligned}
 & -\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f} \\
 & \left(\frac{a}{2(a^2+b^2)} \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}} + 1\right)}{ad} - \frac{2f\left(\frac{f \int e^{-c-dx} \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right) de^{c+dx}}{a^2} - \frac{(e+fx) \text{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d}\right)}{ad} \right) \right) - \frac{a}{2\sqrt{a^2+b^2}} \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{ad} \right) \right) \\
 & \frac{b}{a^2}
 \end{aligned}$$

↓ 7143

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{f(e+fx)\cosh^2(c+dx)}{2d^2} + \frac{f^2\left(\frac{\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{x}{2}\right)}{2d^2} + \frac{(e+fx)^2\sinh(c+dx)\cosh(c+dx)}{2d} + \frac{(e+fx)^3}{6f} \\
 & \left(\frac{a}{2(a^2+b^2)} \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b}+1\right)}{ad} - \frac{2f\left(\frac{f \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d^2} - \frac{(e+fx) \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{d}\right)}{ad} \right) \right) - \frac{a}{2\sqrt{a^2+b^2}} \left(\frac{(e+fx)^2 \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{ad} \right) \right) \\
 & \frac{b}{a^2}
 \end{aligned}$$

input `Int[((e + f*x)^2*Cosh[c + d*x]^2)/(a + b*Csch[c + d*x]),x]`

output `((e + f*x)^3/(6*f) - (f*(e + f*x)*Cosh[c + d*x]^2)/(2*d^2) + ((e + f*x)^2*Cosh[c + d*x]*Sinh[c + d*x])/(2*d) + (f^2*(x/2 + (Cosh[c + d*x]*Sinh[c + d*x])/(2*d)))/(2*d^2))/a - (b*(-1/3*(b*(e + f*x)^3)/(a^2*f) - (2*(a^2 + b^2)*(-1/2*(a*((e + f*x)^2*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/(a*d) - (2*f*(-((e + f*x)*PolyLog[2, -(a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/d) + (f*PolyLog[3, -(a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]))/d^2))/(a*d))/Sqrt[a^2 + b^2] + (a*((e + f*x)^2*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]))/(a*d) - (2*f*(-((e + f*x)*PolyLog[2, -(a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]])))/d) + (f*PolyLog[3, -(a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]))/d^2))/(a*d))/(2*Sqrt[a^2 + b^2]))/a^2 - (I*(I*(e + f*x)^2*Cosh[c + d*x])/d - ((2*I)*f*(-((f*Cosh[c + d*x])/d^2) + ((e + f*x)*Sinh[c + d*x])/d))/d)/a`

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

3.23.3.1 Defintions of rubi rules used

- rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`
- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2694 `Int[((F_)^(u_)*((f_.) + (g_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^(u_) + (c_.)*(F_)^(v_)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3792 `Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*m*(c + d*x)^(m - 1)*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)^m*Cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)^m*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[d^2*m*((m - 1)/(f^2*n^2)) Int[(c + d*x)^(m - 2)*(b*Sin[e + f*x])^n, x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]`

rule 3803 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] := Simp[2 Int[(c + d*x)^m*(E^((-I)*e + f*fz*x)/((-I)*b + 2*a*E^((-I)*e + f*fz*x) + I*b*E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]`

rule 6099 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(n_)*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-a/b^2 Int[(e + f*x)^m*Cosh[c + d*x]^(n - 2), x], x] + (Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^(n - 2)*Sinh[c + d*x], x], x] + Simp[(a^2 + b^2)/b^2 Int[(e + f*x)^m*(Cosh[c + d*x]^(n - 2)/(a + b*Sinh[c + d*x])), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[n, 1] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

rule 6113 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(p_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)]^(n_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^p*Sinh[c + d*x]^(n - 1), x], x] - Simp[a/b Int[(e + f*x)^m*Cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*Sinh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6128 `Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.))/((Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && IntegersQ[m, n]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

3.23.4 Maple [F]

$$\int \frac{(fx + e)^2 \cosh(dx + c)^2}{a + b \operatorname{csch}(dx + c)} dx$$

input `int((f*x+e)^2*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x)`

output `int((f*x+e)^2*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x)`

3.23.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2410 vs. $2(466) = 932$.

Time = 0.31 (sec) , antiderivative size = 2410, normalized size of antiderivative = 4.73

$$\int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((f*x+e)^2*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="fracas")
```

```
output -1/48*(6*a^2*d^2*f^2*x^2 + 6*a^2*d^2*e^2 + 6*a^2*d*e*f - 3*(2*a^2*d^2*f^2*x^2 + 2*a^2*d^2*e^2 - 2*a^2*d*e*f + a^2*f^2 + 2*(2*a^2*d^2*e*f - a^2*d*f^2)*x)*cosh(d*x + c)^4 - 3*(2*a^2*d^2*f^2*x^2 + 2*a^2*d^2*e^2 - 2*a^2*d*e*f + a^2*f^2 + 2*(2*a^2*d^2*e*f - a^2*d*f^2)*x)*sinh(d*x + c)^4 + 3*a^2*f^2 + 24*(a*b*d^2*f^2*x^2 + a*b*d^2*e^2 - 2*a*b*d*e*f + 2*a*b*f^2 + 2*(a*b*d^2*e*f - a*b*d*f^2)*x)*cosh(d*x + c)^3 + 12*(2*a*b*d^2*f^2*x^2 + 2*a*b*d^2*e^2 - 4*a*b*d*e*f + 4*a*b*f^2 + 4*(a*b*d^2*e*f - a*b*d*f^2)*x - (2*a^2*d^2*f^2*x^2 + 2*a^2*d^2*e^2 - 2*a^2*d*e*f + a^2*f^2 + 2*(2*a^2*d^2*e*f - a^2*d*f^2)*x)*cosh(d*x + c))*sinh(d*x + c)^3 - 8*((a^2 + 2*b^2)*d^3*f^2*x^3 + 3*(a^2 + 2*b^2)*d^3*e*f*x^2 + 3*(a^2 + 2*b^2)*d^3*e^2*x)*cosh(d*x + c)^2 - 2*(4*(a^2 + 2*b^2)*d^3*f^2*x^3 + 12*(a^2 + 2*b^2)*d^3*e*f*x^2 + 12*(a^2 + 2*b^2)*d^3*e^2*x + 9*(2*a^2*d^2*f^2*x^2 + 2*a^2*d^2*e^2 - 2*a^2*d*e*f + a^2*f^2 + 2*(2*a^2*d^2*e*f - a^2*d*f^2)*x)*cosh(d*x + c)^2 - 36*(a*b*d^2*f^2*x^2 + a*b*d^2*e^2 - 2*a*b*d*e*f + 2*a*b*f^2 + 2*(a*b*d^2*e*f - a*b*d*f^2)*x)*cosh(d*x + c)*sinh(d*x + c)^2 + 96*((a*b*d*f^2*x + a*b*d*e*f)*cosh(d*x + c)^2 + 2*(a*b*d*f^2*x + a*b*d*e*f)*cosh(d*x + c)*sinh(d*x + c) + (a*b*d*f^2*x + a*b*d*e*f)*sinh(d*x + c)^2)*sqrt((a^2 + b^2)/a^2)*dilog((b*cosh(d*x + c) + b*sinh(d*x + c) + (a*cosh(d*x + c) + a*sinh(d*x + c))*sqrt((a^2 + b^2)/a^2) - a)/a + 1) - 96*((a*b*d*f^2*x + a*b*d*e*f)*cosh(d*x + c)^2 + 2*(a*b*d*f^2*x + a*b*d*e*f)*cosh(d*x + c)*sinh(d*x + c) + (a*b*d*f^2*x ...
```

3.23.6 Sympy [F]

$$\int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

```
input integrate((f*x+e)**2*cosh(d*x+c)**2/(a+b*csch(d*x+c)),x)
```

```
output Integral((e + f*x)**2*cosh(c + d*x)**2/(a + b*csch(c + d*x)), x)
```

3.23. $\int \frac{(e+fx)^2 \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

3.23.7 Maxima [F]

$$\int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^2 \cosh(dx + c)^2}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^2*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/8*e^2*((4*b*e^(-d*x - c) - a)*e^(2*d*x + 2*c)/(a^2*d) - 4*(a^2 + 2*b^2)*(d*x + c)/(a^3*d) + (4*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c))/(a^2*d) + 8*(a^2*b + b^3)*log((a*e^(-d*x - c) - b - sqrt(a^2 + b^2))/(a*e^(-d*x - c) - b + sqrt(a^2 + b^2)))/(sqrt(a^2 + b^2)*a^3*d) + 1/48*(8*(a^2*d^3*f^2*e^(2*c) + 2*b^2*d^3*f^2*e^(2*c))*x^3 + 24*(a^2*d^3*e*f*e^(2*c) + 2*b^2*d^3*e*f*e^(2*c))*x^2 + 3*(2*a^2*d^2*f^2*x^2*e^(4*c) + 2*(2*d^2*e*f - d*f^2)*a^2*x*e^(4*c) - (2*d*e*f - f^2)*a^2*e^(4*c))*e^(2*d*x) - 24*(a*b*d^2*f^2*x^2*e^(3*c) + 2*(d^2*e*f - d*f^2)*a*b*x*e^(3*c))*e^(d*x) - 24*(a*b*d^2*f^2*x^2*e^c + 2*(d^2*e*f + d*f^2)*a*b*x*e^c + 2*(d*e*f + f^2)*a*b*e^c)*e^(-d*x) - 3*(2*a^2*d^2*f^2*x^2 + 2*(2*d^2*e*f + d*f^2)*a^2*x + (2*d*e*f + f^2)*a^2)*e^(-2*d*x))*e^(-2*c)/(a^3*d^3) - integrate(2*((a^2*b*f^2*e^c + b^3*f^2*e^c)*x^2 + 2*(a^2*b*e*f*e^c + b^3*e*f*e^c)*x)*e^(d*x)/(a^4*e^(2*d*x + 2*c) + 2*a^3*b*e^(d*x + c) - a^4), x)`

3.23.8 Giac [F]

$$\int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^2 \cosh(dx + c)^2}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^2*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)^2*cosh(d*x + c)^2/(b*csch(d*x + c) + a), x)`

3.23.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx)^2 \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx)^2 (e + fx)^2}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)^2*(e + f*x)^2)/(a + b/sinh(c + d*x)),x)`output `int((cosh(c + d*x)^2*(e + f*x)^2)/(a + b/sinh(c + d*x)), x)`

3.24 $\int \frac{(e+fx) \cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

3.24.1	Optimal result	210
3.24.2	Mathematica [A] (verified)	211
3.24.3	Rubi [C] (verified)	211
3.24.4	Maple [B] (verified)	218
3.24.5	Fricas [B] (verification not implemented)	218
3.24.6	Sympy [F]	219
3.24.7	Maxima [F]	220
3.24.8	Giac [F]	220
3.24.9	Mupad [F(-1)]	220

3.24.1 Optimal result

Integrand size = 26, antiderivative size = 327

$$\int \frac{(e+fx) \cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = \frac{ex}{2a} + \frac{b^2ex}{a^3} + \frac{fx^2}{4a} + \frac{b^2fx^2}{2a^3} - \frac{b(e+fx) \cosh(c+dx)}{a^2d}$$

$$- \frac{f \cosh^2(c+dx)}{4ad^2} - \frac{b\sqrt{a^2+b^2}(e+fx) \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d}$$

$$+ \frac{b\sqrt{a^2+b^2}(e+fx) \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d}$$

$$- \frac{b\sqrt{a^2+b^2}f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^3d^2}$$

$$+ \frac{b\sqrt{a^2+b^2}f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^3d^2}$$

$$+ \frac{bf \sinh(c+dx)}{a^2d^2} + \frac{(e+fx) \cosh(c+dx) \sinh(c+dx)}{2ad}$$

output

```
1/2*e*x/a+b^2*e*x/a^3+1/4*f*x^2/a+1/2*b^2*f*x^2/a^3-b*(f*x+e)*cosh(d*x+c)/
a^2/d-1/4*f*cosh(d*x+c)^2/a/d^2+b*f*sinh(d*x+c)/a^2/d^2+1/2*(f*x+e)*cosh(d
*x+c)*sinh(d*x+c)/a/d-b*(f*x+e)*ln(1+a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))*(a^
2+b^2)^(1/2)/a^3/d+b*(f*x+e)*ln(1+a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))*(a^2+b
^2)^(1/2)/a^3/d-b*f*polylog(2,-a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))*(a^2+b^2)
^(1/2)/a^3/d^2+b*f*polylog(2,-a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))*(a^2+b^2)
^(1/2)/a^3/d^2
```

3.24.2 Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 610, normalized size of antiderivative = 1.87

$$\int \frac{(e + fx) \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

$$= \operatorname{csch}(c + dx)(b + a \sinh(c + dx)) \left(2a^2 e \left(\frac{c}{d} + x - \frac{2b \arctan\left(\frac{a - b \tanh\left(\frac{1}{2}(c + dx)\right)}{\sqrt{-a^2 - b^2}}\right)}{\sqrt{-a^2 - b^2}d} \right) + a^2 f \left(x^2 - \frac{2b}{d} \left(\log\left(1 + \frac{ae^c}{b - \sqrt{-a^2 - b^2}}\right) \right) \right) \right)$$

input `Integrate[((e + f*x)*Cosh[c + d*x]^2)/(a + b*Csch[c + d*x]),x]`

output `(Csch[c + d*x]*(b + a*Sinh[c + d*x])*(2*a^2*e*(c/d + x - (2*b*ArcTan[(a - b*Tanh[(c + d*x)/2])/Sqrt[-a^2 - b^2]])/(Sqrt[-a^2 - b^2]*d) + a^2*f*(x^2 - (2*b*(d*x*(Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) - Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) + PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - PolyLog[2, -(a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])]))/(Sqrt[a^2 + b^2]*d^2) + (2*e*((a^2 + 4*b^2)*(c + d*x) - (2*b*(3*a^2 + 4*b^2)*ArcTan[(a - b*Tanh[(c + d*x)/2])/Sqrt[-a^2 - b^2]])/Sqrt[-a^2 - b^2] - 4*a*b*Cosh[c + d*x] + a^2*Sinh[2*(c + d*x)]))/d + (f*((a^2 + 4*b^2)*(-c + d*x)*(c + d*x) - 8*a*b*d*x*Cosh[c + d*x] - a^2*Cosh[2*(c + d*x)] - (2*b*(3*a^2 + 4*b^2)*(2*c*ArcTanh[(b + a*E^(c + d*x))/Sqrt[a^2 + b^2]] + (c + d*x)*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) - (c + d*x)*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) + PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - PolyLog[2, -(a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])]))/Sqrt[a^2 + b^2] + 8*a*b*Sinh[c + d*x] + 2*a^2*d*x*Sinh[2*(c + d*x)]))/d^2)/(8*a^3*(a + b*Csch[c + d*x]))`

3.24.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.59 (sec) , antiderivative size = 324, normalized size of antiderivative = 0.99, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.731$, Rules used = {6128, 6113, 3042, 3791, 17, 6099, 17, 3042, 26, 3777, 3042, 3117, 3803, 25, 2694, 27, 2620, 2715, 2838}

3.24. $\int \frac{(e+fx) \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(e+fx) \cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx \\
 & \quad \downarrow \text{6128} \\
 & \int \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{a \sinh(c+dx) + b} dx \\
 & \quad \downarrow \text{6113} \\
 & \frac{\int (e+fx) \cosh^2(c+dx) dx}{a} - \frac{b \int \frac{(e+fx) \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b \int \frac{(e+fx) \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right)^2 dx}{a} \\
 & \quad \downarrow \text{3791} \\
 & \frac{\frac{1}{2} \int (e+fx) dx - \frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d}}{a} - \frac{b \int \frac{(e+fx) \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{17} \\
 & -\frac{\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} - \frac{b \int \frac{(e+fx) \cosh^2(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{6099} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} - \\
 & \frac{b \left(\frac{(a^2+b^2) \int \frac{e+fx}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \int (e+fx) dx}{a^2} + \frac{\int (e+fx) \sinh(c+dx) dx}{a} \right)}{a} \\
 & \quad \downarrow \text{17} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} - \\
 & \frac{b \left(\frac{(a^2+b^2) \int \frac{e+fx}{b+a \sinh(c+dx)} dx}{a^2} + \frac{\int (e+fx) \sinh(c+dx) dx}{a} - \frac{b(e+fx)^2}{2a^2 f} \right)}{a} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.24. $\int \frac{(e+fx) \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{e+fx}{b-ia \sin(ic+idx)} dx}{a^2} + \frac{\int -i(e+fx) \sin(ic+idx) dx}{a} - \frac{b(e+fx)^2}{2a^2 f} \right) \\
 & \quad \downarrow \mathbf{26} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{e+fx}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \int (e+fx) \sin(ic+idx) dx}{a} - \frac{b(e+fx)^2}{2a^2 f} \right) \\
 & \quad \downarrow \mathbf{3777} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{e+fx}{b-ia \sin(ic+idx)} dx}{a^2} - i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \cosh(c+dx) dx}{d} \right) - \frac{b(e+fx)^2}{2a^2 f} \right) \\
 & \quad \downarrow \mathbf{3042} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{e+fx}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \sin(ic+idx + \frac{\pi}{2}) dx}{d} \right)}{a} - \frac{b(e+fx)^2}{2a^2 f} \right) \\
 & \quad \downarrow \mathbf{3117} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{e+fx}{b-ia \sin(ic+idx)} dx}{a^2} - \frac{b(e+fx)^2}{2a^2 f} - \frac{i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \right) \\
 & \quad \downarrow \mathbf{3803} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{2(a^2+b^2) \int -\frac{e^{c+dx}(e+fx)}{-e^{2(c+dx)}a+a-2be^{c+dx}} dx}{a^2} - \frac{b(e+fx)^2}{2a^2 f} - \frac{i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \right) \\
 & \quad \downarrow \mathbf{25}
 \end{aligned}$$

3.24. $\int \frac{(e+fx) \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{2(a^2+b^2) \int \frac{e^{c+dx}(e+fx)}{-e^{2(c+dx)}a+a-2be^{c+dx}} dx}{a^2} - \frac{b(e+fx)^2}{2a^2f} - \frac{i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \right) \\
 & \quad \downarrow \text{2694} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{2(a^2+b^2) \left(\frac{a \int \frac{e^{c+dx}(e+fx)}{2(e^{c+dx}a+b-\sqrt{a^2+b^2})} dx}{\sqrt{a^2+b^2}} - \frac{a \int \frac{e^{c+dx}(e+fx)}{2(e^{c+dx}a+b+\sqrt{a^2+b^2})} dx}{\sqrt{a^2+b^2}} \right)}{a^2} - \frac{b(e+fx)^2}{2a^2f} - \frac{i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \right) \\
 & \quad \downarrow \text{27} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{2(a^2+b^2) \left(\frac{a \int \frac{e^{c+dx}(e+fx)}{e^{c+dx}a+b+\sqrt{a^2+b^2}} dx}{2\sqrt{a^2+b^2}} - \frac{a \int \frac{e^{c+dx}(e+fx)}{e^{c+dx}a+b-\sqrt{a^2+b^2}} dx}{2\sqrt{a^2+b^2}} \right)}{a^2} - \frac{b(e+fx)^2}{2a^2f} - \frac{i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \right) \\
 & \quad \downarrow \text{2620} \\
 & \frac{-\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f}}{a} \\
 & b \left(\frac{2(a^2+b^2) \left(\frac{a \left(\frac{(e+fx) \log \left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b} + 1 \right)}{ad} - \frac{f \int \log \left(\frac{e^{c+dx}a}{b+\sqrt{a^2+b^2}} + 1 \right) dx}{ad} \right)}{2\sqrt{a^2+b^2}} - \frac{a \left(\frac{(e+fx) \log \left(\frac{-ae^{c+dx}}{b-\sqrt{a^2+b^2}} + 1 \right)}{ad} - \frac{f \int \log \left(\frac{-e^{c+dx}a}{b-\sqrt{a^2+b^2}} + 1 \right) dx}{ad} \right)}{2\sqrt{a^2+b^2}} \right)}{a^2} - \frac{b(e+fx)^2}{2a^2f} - \frac{i \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{a} \right) \\
 & \quad \downarrow \text{2715}
 \end{aligned}$$

3.24. $\int \frac{(e+fx) \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \\
 & \left(\frac{2(a^2+b^2)}{a^2} \left(\frac{a \left(\frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b} + 1\right)}{ad} - \frac{f \int e^{-c-dx} \log\left(\frac{e^{c+dx} a}{b+\sqrt{a^2+b^2}} + 1\right) de^{c+dx}}{ad^2} \right)}{2\sqrt{a^2+b^2}} - \frac{a \left(\frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} + 1\right)}{ad} - \frac{f \int e^{-c-dx} \log\left(\frac{e^{c+dx} a}{b-\sqrt{a^2+b^2}} + 1\right)}{ad^2} \right)}{2\sqrt{a^2+b^2}} \right) \right) \\
 & \hspace{15em} \downarrow \text{2838} \\
 & -\frac{f \cosh^2(c+dx)}{4d^2} + \frac{(e+fx) \sinh(c+dx) \cosh(c+dx)}{2d} + \frac{(e+fx)^2}{4f} \\
 & \left(\frac{2(a^2+b^2)}{a^2} \left(\frac{a \left(\frac{f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{ad^2} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b} + 1\right)}{ad} \right)}{2\sqrt{a^2+b^2}} - \frac{a \left(\frac{f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{ad^2} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}} + 1\right)}{ad} \right)}{2\sqrt{a^2+b^2}} \right) \right)
 \end{aligned}$$

```
input Int[((e + f*x)*Cosh[c + d*x]^2)/(a + b*CsCh[c + d*x]),x]
```

```
output ((e + f*x)^2/(4*f) - (f*Cosh[c + d*x]^2)/(4*d^2) + ((e + f*x)*Cosh[c + d*x]
]*Sinh[c + d*x])/(2*d))/a - (b*(-1/2*(b*(e + f*x)^2)/(a^2*f) - (2*(a^2 + b
^2)*(-1/2*(a*(((e + f*x)*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]])))/(
a*d) + (f*PolyLog[2, -((a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]))])/(a*d^2)))/
Sqrt[a^2 + b^2] + (a*(((e + f*x)*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b
^2]])))/(a*d) + (f*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))])/(a
*d^2)))/(2*Sqrt[a^2 + b^2]))/a^2 - (I*((I*(e + f*x)*Cosh[c + d*x])/d - (I
*f*Sinh[c + d*x])/d^2))/a)/a
```

3.24. $\int \frac{(e+fx) \cosh^2(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

3.24.3.1 Defintions of rubi rules used

- rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1)/(b*(m + 1))), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2694 `Int[((F_)^(u_)*((f_.) + (g_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^(u_) + (c_.)*(F_)^(v_)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-`
`(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*C`
`os[e + f*x], x], x] /;` `FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3791 `Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :=`
`Simp[d*((b*Ssin[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x`
`]*)((b*Ssin[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*`
`x)*(b*Ssin[e + f*x])^(n - 2), x], x] /;` `FreeQ[{b, c, d, e, f}, x] && GtQ[n,`
`1]`

rule 3803 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*sin[(e_.) + (Complex[0, fz_]*)`
`(f_.)*(x_)])], x_Symbol] := Simp[2 Int[(c + d*x)^m*(E^((-I)*e + f*fz*x)/((`
`-I)*b + 2*a*E^((-I)*e + f*fz*x) + I*b*E^(2*((-I)*e + f*fz*x)))]], x], x] /;`
`FreeQ[{a, b, c, d, e, f, fz}, x] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]`

rule 6099 `Int[(Cosh[(c_.) + (d_.)*(x_)])^(n_)*((e_.) + (f_.)*(x_))^(m_.)/((a_) + (b_.`
`)*Sinh[(c_.) + (d_.)*(x_)])], x_Symbol] := Simp[-a/b^2 Int[(e + f*x)^m*Cos`
`h[c + d*x]^(n - 2), x], x] + (Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^(n -`
`2)*Sinh[c + d*x], x], x] + Simp[(a^2 + b^2)/b^2 Int[(e + f*x)^m*(Cosh[c`
`+ d*x]^(n - 2)/(a + b*Sinh[c + d*x])], x], x]) /;` `FreeQ[{a, b, c, d, e, f},`
`x] && IGtQ[n, 1] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

rule 6113 `Int[(Cosh[(c_.) + (d_.)*(x_)])^(p_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) +`
`(d_.)*(x_)]^(n_.)/((a_) + (b_.)*Sinh[(c_.) + (d_.)*(x_)])], x_Symbol] := S`
`imp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^p*Sinh[c + d*x]^(n - 1), x], x] - S`
`imp[a/b Int[(e + f*x)^m*Cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*Sin`
`h[c + d*x])], x], x] /;` `FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[`
`n, 0] && IGtQ[p, 0]`

rule 6128 `Int[((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)/(Csch[(c_.)`
`+ (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F`
`[c + d*x]^n/(b + a*Sinh[c + d*x]), x] /;` `FreeQ[{a, b, c, d, e, f}, x] && H`
`yperbolicQ[F] && IntegersQ[m, n]`

3.24.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1011 vs. $2(297) = 594$.

Time = 4.51 (sec) , antiderivative size = 1012, normalized size of antiderivative = 3.09

method	result	size
risch	Expression too large to display	1012

```
input int((f*x+e)*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/4*f*x^2/a+1/2*e*x/a+1/2*b^2*f*x^2/a^3+b^2*e*x/a^3+1/16*(2*d*f*x+2*d*e-f)
/a/d^2*exp(2*d*x+2*c)-1/2*b*(d*f*x+d*e-f)/a^2/d^2*exp(d*x+c)-1/2*b*(d*f*x+
d*e+f)/a^2/d^2*exp(-d*x-c)-1/16*(2*d*f*x+2*d*e+f)/a/d^2*exp(-2*d*x-2*c)+2/
d/a*b*e/(a^2+b^2)^(1/2)*arctanh(1/2*(2*a*exp(d*x+c)+2*b)/(a^2+b^2)^(1/2))+
2/d/a^3*b^3*e/(a^2+b^2)^(1/2)*arctanh(1/2*(2*a*exp(d*x+c)+2*b)/(a^2+b^2)^(
1/2))-1/d/a*b*f/(a^2+b^2)^(1/2)*ln((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(
a^2+b^2)^(1/2)))*x+1/d/a*b*f/(a^2+b^2)^(1/2)*ln((a*exp(d*x+c)+(a^2+b^2)^(1
/2)+b)/(b+(a^2+b^2)^(1/2)))*x-1/d^2/a*b*f/(a^2+b^2)^(1/2)*ln((-a*exp(d*x+c
)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))*c+1/d^2/a*b*f/(a^2+b^2)^(1/2)*l
n((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))*c-1/d^2/a*b*f/(a^2
+b^2)^(1/2)*dilog((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))+
1/d^2/a*b*f/(a^2+b^2)^(1/2)*dilog((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2
+b^2)^(1/2)))-1/d/a^3*b^3*f/(a^2+b^2)^(1/2)*ln((-a*exp(d*x+c)+(a^2+b^2)^(1
/2)-b)/(-b+(a^2+b^2)^(1/2)))*x+1/d/a^3*b^3*f/(a^2+b^2)^(1/2)*ln((a*exp(d*x
+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))*x-1/d^2/a^3*b^3*f/(a^2+b^2)^(1
/2)*ln((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))*c+1/d^2/a^3
*b^3*f/(a^2+b^2)^(1/2)*ln((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1
/2)))*c-1/d^2/a^3*b^3*f/(a^2+b^2)^(1/2)*dilog((-a*exp(d*x+c)+(a^2+b^2)^(1/
2)-b)/(-b+(a^2+b^2)^(1/2)))+1/d^2/a^3*b^3*f/(a^2+b^2)^(1/2)*dilog((a*exp(d
*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))-2/d^2/a*b*f*c/(a^2+b^2)^(...
```

3.24.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1284 vs. $2(295) = 590$.

Time = 0.30 (sec) , antiderivative size = 1284, normalized size of antiderivative = 3.93

$$\int \frac{(e + fx) \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((f*x+e)*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="fricas")
```

3.24. $\int \frac{(e+fx) \cosh^2(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

output

```
-1/16*(2*a^2*d*f*x - (2*a^2*d*f*x + 2*a^2*d*e - a^2*f)*cosh(d*x + c)^4 - (
2*a^2*d*f*x + 2*a^2*d*e - a^2*f)*sinh(d*x + c)^4 + 2*a^2*d*e + 8*(a*b*d*f*
x + a*b*d*e - a*b*f)*cosh(d*x + c)^3 + 4*(2*a*b*d*f*x + 2*a*b*d*e - 2*a*b*
f - (2*a^2*d*f*x + 2*a^2*d*e - a^2*f)*cosh(d*x + c))*sinh(d*x + c)^3 + a^2
*f - 4*((a^2 + 2*b^2)*d^2*f*x^2 + 2*(a^2 + 2*b^2)*d^2*e*x)*cosh(d*x + c)^2
- 2*(2*(a^2 + 2*b^2)*d^2*f*x^2 + 4*(a^2 + 2*b^2)*d^2*e*x + 3*(2*a^2*d*f*x
+ 2*a^2*d*e - a^2*f)*cosh(d*x + c)^2 - 12*(a*b*d*f*x + a*b*d*e - a*b*f)*c
osh(d*x + c))*sinh(d*x + c)^2 + 16*(a*b*f*cosh(d*x + c)^2 + 2*a*b*f*cosh(d
*x + c)*sinh(d*x + c) + a*b*f*sinh(d*x + c)^2)*sqrt((a^2 + b^2)/a^2)*dilog
((b*cosh(d*x + c) + b*sinh(d*x + c) + (a*cosh(d*x + c) + a*sinh(d*x + c))*
sqrt((a^2 + b^2)/a^2) - a)/a + 1) - 16*(a*b*f*cosh(d*x + c)^2 + 2*a*b*f*co
sh(d*x + c)*sinh(d*x + c) + a*b*f*sinh(d*x + c)^2)*sqrt((a^2 + b^2)/a^2)*d
ilog((b*cosh(d*x + c) + b*sinh(d*x + c) - (a*cosh(d*x + c) + a*sinh(d*x +
c))*sqrt((a^2 + b^2)/a^2) - a)/a + 1) - 16*((a*b*d*e - a*b*c*f)*cosh(d*x +
c)^2 + 2*(a*b*d*e - a*b*c*f)*cosh(d*x + c)*sinh(d*x + c) + (a*b*d*e - a*b
*c*f)*sinh(d*x + c)^2)*sqrt((a^2 + b^2)/a^2)*log(2*a*cosh(d*x + c) + 2*a*s
inh(d*x + c) + 2*a*sqrt((a^2 + b^2)/a^2) + 2*b) + 16*((a*b*d*e - a*b*c*f)*
cosh(d*x + c)^2 + 2*(a*b*d*e - a*b*c*f)*cosh(d*x + c)*sinh(d*x + c) + (a*b
*d*e - a*b*c*f)*sinh(d*x + c)^2)*sqrt((a^2 + b^2)/a^2)*log(2*a*cosh(d*x +
c) + 2*a*sinh(d*x + c) - 2*a*sqrt((a^2 + b^2)/a^2) + 2*b) + 16*((a*b*d*...
```

3.24.6 Sympy [F]

$$\int \frac{(e + fx) \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx) \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

input `integrate((f*x+e)*cosh(d*x+c)**2/(a+b*csch(d*x+c)),x)`

output `Integral((e + f*x)*cosh(c + d*x)**2/(a + b*csch(c + d*x)), x)`

3.24.7 Maxima [F]

$$\int \frac{(e + fx) \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e) \cosh(dx + c)^2}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/16*(32*(a^2*b*e^c + b^3*e^c)*integrate(x*e^(d*x)/(a^4*e^(2*d*x + 2*c) + 2*a^3*b*e^(d*x + c) - a^4), x) - (4*(a^2*d^2*e^(2*c) + 2*b^2*d^2*e^(2*c)) *x^2 + (2*a^2*d*x*e^(4*c) - a^2*e^(4*c))*e^(2*d*x) - 8*(a*b*d*x*e^(3*c) - a*b*e^(3*c))*e^(d*x) - 8*(a*b*d*x*e^c + a*b*e^c)*e^(-d*x) - (2*a^2*d*x + a^2)*e^(-2*d*x))*e^(-2*c)/(a^3*d^2))*f - 1/8*e*((4*b*e^(-d*x - c) - a)*e^(2*d*x + 2*c)/(a^2*d) - 4*(a^2 + 2*b^2)*(d*x + c)/(a^3*d) + (4*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c))/(a^2*d) + 8*(a^2*b + b^3)*log((a*e^(-d*x - c) - b - sqrt(a^2 + b^2))/(a*e^(-d*x - c) - b + sqrt(a^2 + b^2)))/(sqrt(a^2 + b^2))*a^3*d)`

3.24.8 Giac [F]

$$\int \frac{(e + fx) \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e) \cosh(dx + c)^2}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)*cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)*cosh(d*x + c)^2/(b*csch(d*x + c) + a), x)`

3.24.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx) \cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx)^2 (e + fx)}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)^2*(e + f*x))/(a + b/sinh(c + d*x)),x)`

output `int((cosh(c + d*x)^2*(e + f*x))/(a + b/sinh(c + d*x)), x)`

3.25 $\int \frac{\cosh^2(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$

3.25.1	Optimal result	221
3.25.2	Mathematica [A] (verified)	221
3.25.3	Rubi [C] (warning: unable to verify)	222
3.25.4	Maple [A] (verified)	225
3.25.5	Fricas [B] (verification not implemented)	226
3.25.6	Sympy [F]	227
3.25.7	Maxima [A] (verification not implemented)	227
3.25.8	Giac [A] (verification not implemented)	227
3.25.9	Mupad [B] (verification not implemented)	228

3.25.1 Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \frac{\cosh^2(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{(a^2+2b^2)x}{2a^3} + \frac{2b\sqrt{a^2+b^2}\operatorname{arctanh}\left(\frac{a-b\tanh\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a^2+b^2}}\right)}{a^3d} - \frac{\cosh(c+dx)(2b-a\sinh(c+dx))}{2a^2d}$$

output `1/2*(a^2+2*b^2)*x/a^3-1/2*cosh(d*x+c)*(2*b-a*sinh(d*x+c))/a^2/d+2*b*arctan
h((a-b*tanh(1/2*d*x+1/2*c))/(a^2+b^2)^(1/2))*(a^2+b^2)^(1/2)/a^3/d`

3.25.2 Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.15

$$\int \frac{\cosh^2(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{2a^2c + 4b^2c + 2a^2dx + 4b^2dx + 8b\sqrt{-a^2-b^2}\operatorname{arctan}\left(\frac{a-b\tanh\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2-b^2}}\right) - 4ab\cosh(c+dx) + a^2\sinh(2(c+dx))}{4a^3d}$$

input `Integrate[Cosh[c + d*x]^2/(a + b*CsCh[c + d*x]),x]`

output $(2a^2c + 4b^2c + 2a^2dx + 4b^2dx + 8b\sqrt{-a^2 - b^2}\text{ArcTan}[(a - b\tanh[(c + dx)/2])/\sqrt{-a^2 - b^2}] - 4ab\cosh[c + dx] + a^2\sinh[2(c + dx)])/(4a^3d)$

3.25.3 Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.13, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4360, 26, 26, 3042, 26, 3344, 26, 3042, 3214, 3042, 3139, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cosh^2(c + dx)}{a + b\text{csch}(c + dx)} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\cos(ic + idx)^2}{a + ib\csc(ic + idx)} dx \\
 & \quad \downarrow 4360 \\
 & \int \frac{i \sinh(c + dx) \cosh^2(c + dx)}{ia \sinh(c + dx) + ib} dx \\
 & \quad \downarrow 26 \\
 & i \int -\frac{i \cosh^2(c + dx) \sinh(c + dx)}{b + a \sinh(c + dx)} dx \\
 & \quad \downarrow 26 \\
 & \int \frac{\sinh(c + dx) \cosh^2(c + dx)}{a \sinh(c + dx) + b} dx \\
 & \quad \downarrow 3042 \\
 & \int -\frac{i \sin(ic + idx) \cos(ic + idx)^2}{b - ia \sin(ic + idx)} dx \\
 & \quad \downarrow 26 \\
 & -i \int \frac{\cos(ic + idx)^2 \sin(ic + idx)}{b - ia \sin(ic + idx)} dx \\
 & \quad \downarrow 3344
 \end{aligned}$$

3.25. $\int \frac{\cosh^2(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\begin{aligned}
& -i \left(-\frac{\int \frac{i(ab - (a^2 + 2b^2) \sinh(c+dx))}{b+a \sinh(c+dx)} dx}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2 d} \right) \\
& \quad \downarrow \text{26} \\
& -i \left(-\frac{i \int \frac{ab - (a^2 + 2b^2) \sinh(c+dx)}{b+a \sinh(c+dx)} dx}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2 d} \right) \\
& \quad \downarrow \text{3042} \\
& -i \left(-\frac{i \int \frac{ab + i(a^2 + 2b^2) \sin(ic+idx)}{b-ia \sin(ic+idx)} dx}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2 d} \right) \\
& \quad \downarrow \text{3214} \\
& -i \left(-\frac{i \left(\frac{2b(a^2 + b^2) \int \frac{1}{b+a \sinh(c+dx)} dx}{a} - \frac{x(a^2 + 2b^2)}{a} \right)}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2 d} \right) \\
& \quad \downarrow \text{3042} \\
& -i \left(-\frac{i \left(-\frac{x(a^2 + 2b^2)}{a} + \frac{2b(a^2 + b^2) \int \frac{1}{b-ia \sin(ic+idx)} dx}{a} \right)}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2 d} \right) \\
& \quad \downarrow \text{3139} \\
& -i \left(-\frac{i \left(-\frac{x(a^2 + 2b^2)}{a} - \frac{4ib(a^2 + b^2) \int \frac{1}{-b \tanh^2(\frac{1}{2}(c+dx)) + 2a \tanh(\frac{1}{2}(c+dx)) + b} d(i \tanh(\frac{1}{2}(c+dx)))}{ad} \right)}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2 d} \right) \\
& \quad \downarrow \text{1083} \\
& -i \left(-\frac{i \left(-\frac{x(a^2 + 2b^2)}{a} + \frac{8ib(a^2 + b^2) \int \frac{1}{\tanh^2(\frac{1}{2}(c+dx)) - 4(a^2 + b^2)} d(2ib \tanh(\frac{1}{2}(c+dx)) - 2ia)}{ad} \right)}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2 d} \right) \\
& \quad \downarrow \text{217}
\end{aligned}$$

$$-i \left(\frac{i \left(\frac{4b\sqrt{a^2+b^2} \operatorname{arctanh}\left(\frac{\tanh\left(\frac{1}{2}(c+dx)\right)}{2\sqrt{a^2+b^2}}\right)}{ad} - \frac{x(a^2+2b^2)}{a} \right)}{2a^2} - \frac{i \cosh(c+dx)(2b - a \sinh(c+dx))}{2a^2d} \right)$$

input `Int[Cosh[c + d*x]^2/(a + b*Csch[c + d*x]),x]`

output `(-I)*(((-1/2*I)*(-((a^2 + 2*b^2)*x)/a) + (4*b*Sqrt[a^2 + b^2]*ArcTanh[Tanh[(c + d*x)/2]/(2*Sqrt[a^2 + b^2])])/(a*d)))/a^2 - ((I/2)*Cosh[c + d*x]*(2*b - a*Sinh[c + d*x]))/(a^2*d)`

3.25.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3344 `Int[(cos[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[g*(g*Cos[e + f*x])^(p - 1)*(a + b*Sin[e + f*x])^(m + 1)*((b*c*(m + p + 1) - a*d*p + b*d*(m + p)*Sin[e + f*x])/(b^2*f*(m + p)*(m + p + 1))), x] + Simp[g^2*(p - 1)/(b^2*(m + p)*(m + p + 1)) Int[(g*Cos[e + f*x])^(p - 2)*(a + b*Sin[e + f*x])^m*Simp[b*(a*d*m + b*c*(m + p + 1)) + (a*b*c*(m + p + 1) - d*(a^2*p - b^2*(m + p)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[a^2 - b^2, 0] && GtQ[p, 1] && NeQ[m + p, 0] && NeQ[m + p + 1, 0] && IntegerQ[2*m]`

rule 4360 `Int[(cos[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_.), x_Symbol] := Int[(g*Cos[e + f*x])^p*((b + a*Sin[e + f*x])^m/Sin[e + f*x]^m), x] /; FreeQ[{a, b, e, f, g, p}, x] && IntegerQ[m]`

3.25.4 Maple [A] (verified)

Time = 4.00 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.76

method	result
risch	$\frac{x}{2a} + \frac{xb^2}{a^3} + \frac{e^{2dx+2c}}{8da} - \frac{be^{dx+c}}{2da^2} - \frac{be^{-dx-c}}{2da^2} - \frac{e^{-2dx-2c}}{8da} + \frac{\sqrt{a^2+b^2} b \ln\left(\frac{e^{dx+c} + b + \sqrt{a^2+b^2}}{a}\right)}{da^3} - \frac{\sqrt{a^2+b^2} b \ln\left(\frac{e^{-dx-c} - b + \sqrt{a^2+b^2}}{a}\right)}{da^3}$
derivativedivides	$-\frac{1}{2a\left(1+\tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2} - \frac{-a+2b}{2a^2\left(1+\tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + \frac{(a^2+2b^2) \ln\left(1+\tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2a^3} + \frac{2b\sqrt{a^2+b^2} \operatorname{arctanh}\left(\frac{-2b \tanh\left(\frac{dx}{2} + \frac{c}{2}\right) + 2a}{2\sqrt{a^2+b^2}}\right)}{a^3}$
default	$-\frac{1}{2a\left(1+\tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2} - \frac{-a+2b}{2a^2\left(1+\tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + \frac{(a^2+2b^2) \ln\left(1+\tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2a^3} + \frac{2b\sqrt{a^2+b^2} \operatorname{arctanh}\left(\frac{-2b \tanh\left(\frac{dx}{2} + \frac{c}{2}\right) + 2a}{2\sqrt{a^2+b^2}}\right)}{a^3}$

input `int(cosh(d*x+c)^2/(a+b*cSch(d*x+c)),x,method=_RETURNVERBOSE)`

output $\frac{1}{2}x/a + x/a^3 b^2 + 1/8/d/a \exp(2dx+2c) - 1/2b/d/a^2 \exp(dx+c) - 1/2b/d/a^2 \exp(-dx-c) - 1/8/d/a \exp(-2dx-2c) + (a^2+b^2)^{1/2} b/d/a^3 \ln(\exp(dx+c)) + (b+(a^2+b^2)^{1/2})/a - (a^2+b^2)^{1/2} b/d/a^3 \ln(\exp(dx+c) - (-b+(a^2+b^2)^{1/2})/a)$

3.25.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 446 vs. $2(87) = 174$.

Time = 0.28 (sec) , antiderivative size = 446, normalized size of antiderivative = 4.69

$$\int \frac{\cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$$

$$= \frac{a^2 \cosh(dx+c)^4 + a^2 \sinh(dx+c)^4 + 4(a^2+2b^2)dx \cosh(dx+c)^2 - 4ab \cosh(dx+c)^3 + 4(a^2 \cosh(dx+c)^2 + a^2 \sinh(dx+c)^2 + 2ab \cosh(dx+c) + a^2 + 2b^2 + 2(a^2 \cosh(dx+c) + ab) \sinh(dx+c) + 2\sqrt{a^2+b^2}(a \cosh(dx+c) + a \sinh(dx+c) + b))}{(a \cosh(dx+c)^2 + a \sinh(dx+c)^2 + 2b \cosh(dx+c) + 2(a \cosh(dx+c) + b) \sinh(dx+c) - a)}$$

input `integrate(cosh(dx+c)^2/(a+b*csch(dx+c)),x, algorithm="fricas")`

output $\frac{1}{8}(a^2 \cosh(dx+c)^4 + a^2 \sinh(dx+c)^4 + 4(a^2 + 2b^2)dx \cosh(dx+c)^2 - 4ab \cosh(dx+c)^3 + 4(a^2 \cosh(dx+c) - ab) \sinh(dx+c)^3 - 4ab \cosh(dx+c) + 2(3a^2 \cosh(dx+c)^2 + 2(a^2 + 2b^2)dx - 6ab \cosh(dx+c)) \sinh(dx+c)^2 + 8(b \cosh(dx+c)^2 + 2b \cosh(dx+c) \sinh(dx+c) + b \sinh(dx+c)^2) \sqrt{a^2 + b^2} \log((a^2 \cosh(dx+c)^2 + a^2 \sinh(dx+c)^2 + 2ab \cosh(dx+c) + a^2 + 2b^2 + 2(a^2 \cosh(dx+c) + ab) \sinh(dx+c) + 2\sqrt{a^2 + b^2}(a \cosh(dx+c) + a \sinh(dx+c) + b)) / (a \cosh(dx+c)^2 + a \sinh(dx+c)^2 + 2b \cosh(dx+c) + 2(a \cosh(dx+c) + b) \sinh(dx+c) - a)) - a^2 + 4(a^2 \cosh(dx+c)^3 + 2(a^2 + 2b^2)dx \cosh(dx+c) - 3ab \cosh(dx+c)^2 - ab) \sinh(dx+c)) / (a^3 d \cosh(dx+c)^2 + 2a^3 d \cosh(dx+c) \sinh(dx+c) + a^3 d \sinh(dx+c)^2)$

3.25.6 Sympy [F]

$$\int \frac{\cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

input `integrate(cosh(d*x+c)**2/(a+b*csch(d*x+c)),x)`

output `Integral(cosh(c + d*x)**2/(a + b*csch(c + d*x)), x)`

3.25.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.77

$$\int \frac{\cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = -\frac{(4be^{(-dx-c)} - a)e^{(2dx+2c)}}{8a^2d} + \frac{(a^2 + 2b^2)(dx + c)}{2a^3d} - \frac{4be^{(-dx-c)} + ae^{(-2dx-2c)}}{8a^2d} - \frac{(a^2b + b^3) \log\left(\frac{ae^{(-dx-c)} - b - \sqrt{a^2 + b^2}}{ae^{(-dx-c)} - b + \sqrt{a^2 + b^2}}\right)}{\sqrt{a^2 + b^2}a^3d}$$

input `integrate(cosh(d*x+c)^2/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/8*(4*b*e^(-d*x - c) - a)*e^(2*d*x + 2*c)/(a^2*d) + 1/2*(a^2 + 2*b^2)*(d*x + c)/(a^3*d) - 1/8*(4*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c))/(a^2*d) - (a^2*b + b^3)*log((a*e^(-d*x - c) - b - sqrt(a^2 + b^2))/(a*e^(-d*x - c) - b + sqrt(a^2 + b^2)))/(sqrt(a^2 + b^2)*a^3*d)`

3.25.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.63

$$\int \frac{\cosh^2(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \frac{4(a^2 + 2b^2)(dx + c)}{a^3} + \frac{ae^{(2dx+2c)} - 4be^{(dx+c)}}{a^2} - \frac{(4abe^{(dx+c)} + a^2)e^{(-2dx-2c)}}{a^3} - \frac{8(a^2b + b^3) \log\left(\frac{2ae^{(dx+c)} + 2b - 2\sqrt{a^2 + b^2}}{2ae^{(dx+c)} + 2b + 2\sqrt{a^2 + b^2}}\right)}{\sqrt{a^2 + b^2}a^3}$$

input `integrate(cosh(d*x+c)^2/(a+b*cSch(d*x+c)),x, algorithm="giac")`

output `1/8*(4*(a^2 + 2*b^2)*(d*x + c)/a^3 + (a*e^(2*d*x + 2*c) - 4*b*e^(d*x + c))
/a^2 - (4*a*b*e^(d*x + c) + a^2)*e^(-2*d*x - 2*c)/a^3 - 8*(a^2*b + b^3)*lo
g(abs(2*a*e^(d*x + c) + 2*b - 2*sqrt(a^2 + b^2))/abs(2*a*e^(d*x + c) + 2*b
+ 2*sqrt(a^2 + b^2)))/(sqrt(a^2 + b^2)*a^3))/d`

3.25.9 Mupad [B] (verification not implemented)

Time = 2.44 (sec) , antiderivative size = 212, normalized size of antiderivative = 2.23

$$\int \frac{\cosh^2(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = \frac{e^{2c+2dx}}{8ad} - \frac{e^{-2c-2dx}}{8ad} + \frac{x(a^2+2b^2)}{2a^3} - \frac{be^{-c-dx}}{2a^2d} - \frac{be^{c+dx}}{2a^2d}$$

$$- \frac{b \ln\left(\frac{2be^{c+dx}(a^2+b^2)}{a^4} - \frac{2b\sqrt{a^2+b^2}(a-be^{c+dx})}{a^4}\right) \sqrt{a^2+b^2}}{a^3d}$$

$$+ \frac{b \ln\left(\frac{2b\sqrt{a^2+b^2}(a-be^{c+dx})}{a^4} + \frac{2be^{c+dx}(a^2+b^2)}{a^4}\right) \sqrt{a^2+b^2}}{a^3d}$$

input `int(cosh(c + d*x)^2/(a + b/sinh(c + d*x)),x)`

output `exp(2*c + 2*d*x)/(8*a*d) - exp(- 2*c - 2*d*x)/(8*a*d) + (x*(a^2 + 2*b^2))/
(2*a^3) - (b*exp(- c - d*x))/(2*a^2*d) - (b*exp(c + d*x))/(2*a^2*d) - (b*1
og((2*b*exp(c + d*x)*(a^2 + b^2))/a^4 - (2*b*(a^2 + b^2)^(1/2)*(a - b*exp(
c + d*x)))/a^4)*(a^2 + b^2)^(1/2)))/(a^3*d) + (b*log((2*b*(a^2 + b^2)^(1/2)
*(a - b*exp(c + d*x)))/a^4 + (2*b*exp(c + d*x)*(a^2 + b^2))/a^4)*(a^2 + b^
2)^(1/2)))/(a^3*d)`

$$3.26 \quad \int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$$

3.26.1	Optimal result	230
3.26.2	Mathematica [B] (warning: unable to verify)	231
3.26.3	Rubi [F]	231
3.26.4	Maple [F]	239
3.26.5	Fricas [B] (verification not implemented)	240
3.26.6	Sympy [F(-1)]	240
3.26.7	Maxima [F]	240
3.26.8	Giac [F]	241
3.26.9	Mupad [F(-1)]	242

3.26.1 Optimal result

Integrand size = 28, antiderivative size = 864

$$\begin{aligned}
 \int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = & -\frac{3bf^3x}{8a^2d^3} - \frac{b(e+fx)^3}{4a^2d} + \frac{b(a^2+b^2)(e+fx)^4}{4a^4f} \\
 & - \frac{40f^3 \cosh(c+dx)}{9ad^4} - \frac{6b^2f^3 \cosh(c+dx)}{a^3d^4} \\
 & - \frac{2f(e+fx)^2 \cosh(c+dx)}{ad^2} - \frac{3b^2f(e+fx)^2 \cosh(c+dx)}{a^3d^2} \\
 & - \frac{2f^3 \cosh^3(c+dx)}{27ad^4} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3ad^2} \\
 & - \frac{b(a^2+b^2)(e+fx)^3 \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d} \\
 & - \frac{b(a^2+b^2)(e+fx)^3 \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d} \\
 & - \frac{3b(a^2+b^2)f(e+fx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d^2} \\
 & - \frac{3b(a^2+b^2)f(e+fx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d^2} \\
 & + \frac{6b(a^2+b^2)f^2(e+fx) \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d^3} \\
 & + \frac{6b(a^2+b^2)f^2(e+fx) \operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d^3} \\
 & - \frac{6b(a^2+b^2)f^3 \operatorname{PolyLog}\left(4, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d^4} \\
 & - \frac{6b(a^2+b^2)f^3 \operatorname{PolyLog}\left(4, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d^4} \\
 & + \frac{40f^2(e+fx) \sinh(c+dx)}{9ad^3} + \frac{6b^2f^2(e+fx) \sinh(c+dx)}{a^3d^3} \\
 & + \frac{2(e+fx)^3 \sinh(c+dx)}{3ad} + \frac{b^2(e+fx)^3 \sinh(c+dx)}{a^3d} \\
 & + \frac{3bf^3 \cosh(c+dx) \sinh(c+dx)}{8a^2d^4} \\
 & + \frac{3bf(e+fx)^2 \cosh(c+dx) \sinh(c+dx)}{4a^2d^2} \\
 & + \frac{2f^2(e+fx) \cosh^2(c+dx) \sinh(c+dx)}{9ad^3} \\
 & + \frac{(e+fx)^3 \cosh^2(c+dx) \sinh(c+dx)}{3ad} \\
 & - \frac{3bf^2(e+fx) \sinh^2(c+dx)}{4a^2d^3} - \frac{b(e+fx)^3 \sinh^2(c+dx)}{2a^2d}
 \end{aligned}$$

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

output

```
-b*(a^2+b^2)*(f*x+e)^3*ln(1+a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^4/d-3*b^2*
f*(f*x+e)^2*cosh(d*x+c)/a^3/d^2+6*b^2*f^2*(f*x+e)*sinh(d*x+c)/a^3/d^3+3/8*
b*f^3*cosh(d*x+c)*sinh(d*x+c)/a^2/d^4+2/9*f^2*(f*x+e)*cosh(d*x+c)^2*sinh(d
*x+c)/a/d^3-3/4*b*f^2*(f*x+e)*sinh(d*x+c)^2/a^2/d^3-6*b*(a^2+b^2)*f^3*poly
log(4,-a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^4/d^4-6*b*(a^2+b^2)*f^3*polylog
(4,-a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^4/d^4-2*f*(f*x+e)^2*cosh(d*x+c)/a/
d^2+40/9*f^2*(f*x+e)*sinh(d*x+c)/a/d^3+1/3*(f*x+e)^3*cosh(d*x+c)^2*sinh(d*
x+c)/a/d-1/2*b*(f*x+e)^3*sinh(d*x+c)^2/a^2/d-1/3*f*(f*x+e)^2*cosh(d*x+c)^3
/a/d^2-6*b^2*f^3*cosh(d*x+c)/a^3/d^4+1/4*b*(a^2+b^2)*(f*x+e)^4/a^4/f-3/8*b
*f^3*x/a^2/d^3+2/3*(f*x+e)^3*sinh(d*x+c)/a/d-3*b*(a^2+b^2)*f*(f*x+e)^2*pol
ylog(2,-a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^4/d^2-3*b*(a^2+b^2)*f*(f*x+e)^
2*polylog(2,-a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^4/d^2+6*b*(a^2+b^2)*f^2*(
f*x+e)*polylog(3,-a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^4/d^3+6*b*(a^2+b^2)*
f^2*(f*x+e)*polylog(3,-a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^4/d^3+3/4*b*f*(
f*x+e)^2*cosh(d*x+c)*sinh(d*x+c)/a^2/d^2+b^2*(f*x+e)^3*sinh(d*x+c)/a^3/d-b
*(a^2+b^2)*(f*x+e)^3*ln(1+a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^4/d-40/9*f^3
*cosh(d*x+c)/a/d^4-1/4*b*(f*x+e)^3/a^2/d-2/27*f^3*cosh(d*x+c)^3/a/d^4
```

3.26.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 7404 vs. $2(864) = 1728$.

Time = 32.30 (sec) , antiderivative size = 7404, normalized size of antiderivative = 8.57

$$\int \frac{(e + fx)^3 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Result too large to show}$$

input `Integrate[((e + f*x)^3*Cosh[c + d*x]^3)/(a + b*Csch[c + d*x]),x]`

output `Result too large to show`

3.26.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e + fx)^3 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

↓ 6128

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \int \frac{(e+fx)^3 \sinh(c+dx) \cosh^3(c+dx)}{a \sinh(c+dx) + b} dx \\
 & \quad \downarrow \text{6113} \\
 & \frac{\int (e+fx)^3 \cosh^3(c+dx) dx}{a} - \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx)^3 \sin\left(ic+idx+\frac{\pi}{2}\right)^3 dx}{a} \\
 & \quad \downarrow \text{3792} \\
 & \frac{\frac{2f^2 \int (e+fx) \cosh^3(c+dx) dx}{3d^2} + \frac{2}{3} \int (e+fx)^3 \cosh(c+dx) dx - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & \quad - \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
 & \frac{\frac{2f^2 \int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right)^3 dx}{3d^2} + \frac{2}{3} \int (e+fx)^3 \sin\left(ic+idx+\frac{\pi}{2}\right) dx - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{(e+fx)^3 \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & \quad \downarrow \text{3777} \\
 & - \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
 & \frac{\frac{2f^2 \int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right)^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3if \int -i(e+fx)^2 \sinh(c+dx) dx}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{(e+fx)^3 \sinh(c+dx)}{3d}}{a} \\
 & \quad \downarrow \text{26} \\
 & - \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
 & \frac{\frac{2f^2 \int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right)^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3f \int (e+fx)^2 \sinh(c+dx) dx}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{(e+fx)^3 \sinh(c+dx)}{3d}}{a} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
 & \frac{\frac{2f^2 \int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right)^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3f \int -i(e+fx)^2 \sin(ic+idx) dx}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{(e+fx)^3 \sinh(c+dx)}{3d}}{a}
 \end{aligned}$$

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\begin{aligned} & \downarrow 26 \\ & \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{\frac{2f^2 \int (e+fx) \sin(ic+idx+\frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \int (e+fx)^2 \sin(ic+idx) dx}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{(e+fx)^3 \sinh(c+dx)}{3a}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow 3777 \\ & \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{\frac{2f^2 \int (e+fx) \sin(ic+idx+\frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \int (e+fx) \cosh(c+dx) dx}{d} \right)}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{\frac{2f^2 \int (e+fx) \sin(ic+idx+\frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \int (e+fx) \sin(ic+idx+\frac{\pi}{2}) dx}{d} \right)}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow 3777 \\ & \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{\frac{2f^2 \int (e+fx) \sin(ic+idx+\frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{if \int -i \sinh(c+dx) dx}{d} \right)}{d} \right)}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow 26 \\ & \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{\frac{2f^2 \int (e+fx) \sin(ic+idx+\frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sinh(c+dx) dx}{d} \right)}{d} \right)}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{\frac{2f^2 \int (e+fx) \sin(ic+idx+\frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sinh(c+dx) dx}{d} \right)}{d} \right)}{d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2}}{a} \end{aligned}$$

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{2f^2 \int (e+fx) \sin(ic+idx + \frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int -i \sin(ic+idx) dx}{d} \right)}{d} \right)}{d} \right)$$

26

$$\frac{2f^2 \int (e+fx) \sin(ic+idx + \frac{\pi}{2})^3 dx}{3d^2} + \frac{2}{3} \left(\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} + \frac{if \int \sin(ic+idx) dx}{d} \right)}{d} \right)}{d} \right)$$

3118

$$\frac{2f^2 \int (e+fx) \sin(ic+idx + \frac{\pi}{2})^3 dx}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sin(ic+idx) dx}{d} \right)}{d} \right)}{d} \right)$$

3791

$$\frac{2f^2 \left(\frac{2}{3} \int (e+fx) \cosh(c+dx) dx - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sin(ic+idx) dx}{d} \right)}{d} \right)}{d} \right)}{3d^2}$$

3042

$$\frac{2f^2 \left(\frac{2}{3} \int (e+fx) \sin(ic+idx + \frac{\pi}{2}) dx - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sin(ic+idx) dx}{d} \right)}{d} \right)}{d} \right)}{3d^2}$$

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned} & \downarrow \text{3777} \\ & -\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{if \int -i \sinh(c+dx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right)}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{26} \\ & -\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sinh(c+dx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right)}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3042} \\ & -\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int -i \sin(ic+idx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right)}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{26} \\ & -\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\ & \frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} + \frac{if \int \sin(ic+idx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right)}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a} \end{aligned}$$

$$\downarrow \text{3118}$$

$$3.26. \quad \int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$$

$$\frac{b \int \frac{(e+fx)^3 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{3d^2}$$

↓ 6099

$$\frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \int (e+fx)^3 \cosh(c+dx) dx}{a^2} + \frac{\int (e+fx)^3 \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a} + \frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{3d^2}$$

↓ 3042

$$\frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{3d^2}$$

$$\frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \int (e+fx)^3 \sin\left(ic+idx+\frac{\pi}{2}\right) dx}{a^2} + \frac{\int (e+fx)^3 \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a}$$

↓ 3777

$$\frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{3d^2}$$

$$\frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3if \int -i(e+fx)^2 \sinh(c+dx) dx}{d} \right)}{a^2} + \frac{\int (e+fx)^3 \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a}$$

↓ 26

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3f \int (e+fx)^2 \sinh(c+dx) dx}{d} \right)}{a^2} + \frac{\int (e+fx)^3 \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a} + \frac{2f^2 \left(\frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right)}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a}$$

↓ 3042

$$\frac{2f^2 \left(\frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right)}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a}$$

$$\frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} - \frac{3f \int -i(e+fx)^2 \sin(ic+idx) dx}{d} \right)}{a^2} + \frac{\int (e+fx)^3 \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a}$$

↓ 26

$$\frac{2f^2 \left(\frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right)}{3d^2} - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a}$$

$$\frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \int (e+fx)^2 \sin(ic+idx) dx}{d} \right)}{a^2} + \frac{\int (e+fx)^3 \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a}$$

↓ 3777

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{2f^2 \left(\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d} \right) - \frac{f(e+fx)^2 \cosh^3(c+dx)}{3d^2} + \frac{2}{3} \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} \right)}{a} \\ b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^3 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx)^3 \sinh(c+dx)}{d} + \frac{3if \left(\frac{i(e+fx)^2 \cosh(c+dx)}{d} - \frac{2if \int (e+fx) \cosh(c+dx) dx}{d} \right)}{d} \right)}{a^2} \right) + \frac{f(e+fx)^3 \cosh(c+dx)}{a}$$

input `Int[((e + f*x)^3*Cosh[c + d*x]^3)/(a + b*CsCh[c + d*x]),x]`

output `$Aborted`

3.26.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3118 `Int[sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3791 `Int[((c_) + (d_)*(x_))*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x] * ((b*Sin[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)*(b*Sin[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

3.26. $\int \frac{(e+fx)^3 \cosh^3(c+dx)}{a+b\text{CSch}(c+dx)} dx$

```
rule 3792 Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
  := Simp[d*m*(c + d*x)^(m - 1)*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (-Simp[
  b*(c + d*x)^m*cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^
  2*((n - 1)/n) Int[(c + d*x)^m*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[d^2
  *m*(m - 1)/(f^2*n^2) Int[(c + d*x)^(m - 2)*(b*Sin[e + f*x])^n, x], x]
  /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]
```

```
rule 6099 Int[(Cosh[(c_.) + (d_.)*(x_)]^(n_)*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_.)
  *Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-a/b^2 Int[(e + f*x)^m*cos
  h[c + d*x]^(n - 2), x], x] + (Simp[1/b Int[(e + f*x)^m*cosh[c + d*x]^(n -
  2)*sinh[c + d*x], x], x] + Simp[(a^2 + b^2)/b^2 Int[(e + f*x)^m*(Cosh[c
  + d*x]^(n - 2)/(a + b*Sinh[c + d*x])), x], x]) /; FreeQ[{a, b, c, d, e, f},
  x] && IGtQ[n, 1] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]
```

```
rule 6113 Int[(Cosh[(c_.) + (d_.)*(x_)]^(p_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) +
  (d_.)*(x_)]^(n_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := S
  imp[1/b Int[(e + f*x)^m*cosh[c + d*x]^p*sinh[c + d*x]^(n - 1), x], x] - S
  imp[a/b Int[(e + f*x)^m*cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*S
  inh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[
  n, 0] && IGtQ[p, 0]
```

```
rule 6128 Int[((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)/(Csch[(c_.)
  + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F
  [c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && H
  yperbolicQ[F] && IntegersQ[m, n]
```

3.26.4 Maple [F]

$$\int \frac{(fx + e)^3 \cosh(dx + c)^3}{a + b \operatorname{csch}(dx + c)} dx$$

```
input int((f*x+e)^3*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x)
```

```
output int((f*x+e)^3*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x)
```


3.26.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7980 vs. $2(810) = 1620$.

Time = 0.38 (sec) , antiderivative size = 7980, normalized size of antiderivative = 9.24

$$\int \frac{(e + fx)^3 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

input `integrate((f*x+e)^3*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.26.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(e + fx)^3 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Timed out}$$

input `integrate((f*x+e)**3*cosh(d*x+c)**3/(a+b*csch(d*x+c)),x)`

output Timed out

3.26.7 Maxima [F]

$$\int \frac{(e + fx)^3 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^3 \cosh(dx + c)^3}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^3*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output

```

-1/24*e^3*((3*a*b*e^(-d*x - c) - a^2 - 3*(3*a^2 + 4*b^2)*e^(-2*d*x - 2*c))
*e^(3*d*x + 3*c)/(a^3*d) + 24*(a^2*b + b^3)*(d*x + c)/(a^4*d) + (3*a*b*e^(
-2*d*x - 2*c) + a^2*e^(-3*d*x - 3*c) + 3*(3*a^2 + 4*b^2)*e^(-d*x - c))/(a^
3*d) + 24*(a^2*b + b^3)*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a
^4*d) - 1/864*(216*(a^2*b*d^4*f^3*e^(3*c) + b^3*d^4*f^3*e^(3*c))*x^4 + 86
4*(a^2*b*d^4*e*f^2*e^(3*c) + b^3*d^4*e*f^2*e^(3*c))*x^3 + 1296*(a^2*b*d^4*
e^2*f*e^(3*c) + b^3*d^4*e^2*f*e^(3*c))*x^2 - 4*(9*a^3*d^3*f^3*x^3*e^(6*c)
+ 9*(3*d^3*e*f^2 - d^2*f^3)*a^3*x^2*e^(6*c) + 3*(9*d^3*e^2*f - 6*d^2*e*f^2
+ 2*d*f^3)*a^3*x*e^(6*c) - (9*d^2*e^2*f - 6*d*e*f^2 + 2*f^3)*a^3*e^(6*c))
*e^(3*d*x) + 27*(4*a^2*b*d^3*f^3*x^3*e^(5*c) + 6*(2*d^3*e*f^2 - d^2*f^3)*a
^2*b*x^2*e^(5*c) + 6*(2*d^3*e^2*f - 2*d^2*e*f^2 + d*f^3)*a^2*b*x*e^(5*c) -
3*(2*d^2*e^2*f - 2*d*e*f^2 + f^3)*a^2*b*e^(5*c))*e^(2*d*x) + 108*(9*(d^2*
e^2*f - 2*d*e*f^2 + 2*f^3)*a^3*e^(4*c) + 12*(d^2*e^2*f - 2*d*e*f^2 + 2*f^3
)*a*b^2*e^(4*c) - (3*a^3*d^3*f^3*e^(4*c) + 4*a*b^2*d^3*f^3*e^(4*c))*x^3 -
3*(3*(d^3*e*f^2 - d^2*f^3)*a^3*e^(4*c) + 4*(d^3*e*f^2 - d^2*f^3)*a*b^2*e^(
4*c))*x^2 - 3*(3*(d^3*e^2*f - 2*d^2*e*f^2 + 2*d*f^3)*a^3*e^(4*c) + 4*(d^3*
e^2*f - 2*d^2*e*f^2 + 2*d*f^3)*a*b^2*e^(4*c))*x)*e^(d*x) + 108*(9*(d^2*e^2
*f + 2*d*e*f^2 + 2*f^3)*a^3*e^(2*c) + 12*(d^2*e^2*f + 2*d*e*f^2 + 2*f^3)*a
*b^2*e^(2*c) + (3*a^3*d^3*f^3*e^(2*c) + 4*a*b^2*d^3*f^3*e^(2*c))*x^3 + 3*(
3*(d^3*e*f^2 + d^2*f^3)*a^3*e^(2*c) + 4*(d^3*e*f^2 + d^2*f^3)*a*b^2*e^(...

```

3.26.8 Giac [F]

$$\int \frac{(e + fx)^3 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^3 \cosh(dx + c)^3}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^3*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)^3*cosh(d*x + c)^3/(b*csch(d*x + c) + a), x)`

3.26.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx)^3 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx)^3 (e + fx)^3}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)^3*(e + f*x)^3)/(a + b/sinh(c + d*x)),x)`output `int((cosh(c + d*x)^3*(e + f*x)^3)/(a + b/sinh(c + d*x)), x)`

$$3.27 \quad \int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$$

3.27.1	Optimal result	244
3.27.2	Mathematica [B] (warning: unable to verify)	245
3.27.3	Rubi [F]	246
3.27.4	Maple [F]	255
3.27.5	Fricas [B] (verification not implemented)	255
3.27.6	Sympy [F]	256
3.27.7	Maxima [F]	257
3.27.8	Giac [F]	257
3.27.9	Mupad [F(-1)]	258

3.27.1 Optimal result

Integrand size = 28, antiderivative size = 636

$$\begin{aligned}
\int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = & -\frac{befx}{2a^2d} - \frac{bf^2x^2}{4a^2d} + \frac{b(a^2+b^2)(e+fx)^3}{3a^4f} \\
& - \frac{4f(e+fx)\cosh(c+dx)}{3ad^2} - \frac{2b^2f(e+fx)\cosh(c+dx)}{a^3d^2} \\
& - \frac{2f(e+fx)\cosh^3(c+dx)}{9ad^2} \\
& - \frac{b(a^2+b^2)(e+fx)^2 \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d} \\
& - \frac{b(a^2+b^2)(e+fx)^2 \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d} \\
& - \frac{2b(a^2+b^2)f(e+fx)\operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d^2} \\
& - \frac{2b(a^2+b^2)f(e+fx)\operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d^2} \\
& + \frac{2b(a^2+b^2)f^2\operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d^3} \\
& + \frac{2b(a^2+b^2)f^2\operatorname{PolyLog}\left(3, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d^3} \\
& + \frac{14f^2\sinh(c+dx)}{9ad^3} + \frac{2b^2f^2\sinh(c+dx)}{a^3d^3} \\
& + \frac{2(e+fx)^2\sinh(c+dx)}{3ad} + \frac{b^2(e+fx)^2\sinh(c+dx)}{a^3d} \\
& + \frac{bf(e+fx)\cosh(c+dx)\sinh(c+dx)}{2a^2d^2} \\
& + \frac{(e+fx)^2\cosh^2(c+dx)\sinh(c+dx)}{3ad} - \frac{bf^2\sinh^2(c+dx)}{4a^2d^3} \\
& - \frac{b(e+fx)^2\sinh^2(c+dx)}{2a^2d} + \frac{2f^2\sinh^3(c+dx)}{27ad^3}
\end{aligned}$$

output
$$\begin{aligned}
& -1/2*b*e*f*x/a^2/d-1/4*b*f^2*x^2/a^2/d+1/3*b*(a^2+b^2)*(f*x+e)^3/a^4/f-4/3 \\
& *f*(f*x+e)*\cosh(d*x+c)/a/d^2-2*b^2*f*(f*x+e)*\cosh(d*x+c)/a^3/d^2-2/9*f*(f* \\
& x+e)*\cosh(d*x+c)^3/a/d^2-b*(a^2+b^2)*(f*x+e)^2*\ln(1+a*\exp(d*x+c)/(b-(a^2+b \\
& ^2)^{(1/2)}))/a^4/d-b*(a^2+b^2)*(f*x+e)^2*\ln(1+a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/ \\
& 2)}))/a^4/d-2*b*(a^2+b^2)*f*(f*x+e)*\text{polylog}(2,-a*\exp(d*x+c)/(b-(a^2+b^2)^{(1 \\
& /2)}))/a^4/d^2-2*b*(a^2+b^2)*f*(f*x+e)*\text{polylog}(2,-a*\exp(d*x+c)/(b+(a^2+b^2) \\
& ^{(1/2)}))/a^4/d^2+2*b*(a^2+b^2)*f^2*\text{polylog}(3,-a*\exp(d*x+c)/(b-(a^2+b^2)^{(1 \\
& /2)}))/a^4/d^3+2*b*(a^2+b^2)*f^2*\text{polylog}(3,-a*\exp(d*x+c)/(b+(a^2+b^2)^{(1/2) \\
&))/a^4/d^3+14/9*f^2*\sinh(d*x+c)/a/d^3+2*b^2*f^2*\sinh(d*x+c)/a^3/d^3+2/3*(f \\
& *x+e)^2*\sinh(d*x+c)/a/d+b^2*(f*x+e)^2*\sinh(d*x+c)/a^3/d+1/2*b*f*(f*x+e)*\text{co} \\
& \text{sh}(d*x+c)*\sinh(d*x+c)/a^2/d^2+1/3*(f*x+e)^2*\cosh(d*x+c)^2*\sinh(d*x+c)/a/d- \\
& 1/4*b*f^2*\sinh(d*x+c)^2/a^2/d^3-1/2*b*(f*x+e)^2*\sinh(d*x+c)^2/a^2/d+2/27*f \\
& ^2*\sinh(d*x+c)^3/a/d^3
\end{aligned}$$

3.27.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 2129 vs. $2(636) = 1272$.

Time = 16.80 (sec) , antiderivative size = 2129, normalized size of antiderivative = 3.35

$$\int \frac{(e + fx)^2 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Result too large to show}$$

input `Integrate[((e + f*x)^2*Cosh[c + d*x]^3)/(a + b*CsCh[c + d*x]),x]`

output

```
(f^2*Csch[c + d*x]*((4*b*x^3)/(-1 + E^(2*c)) - 2*b*x^3*Coth[c] - (6*a^2*b*(d^2*x^2*Log[1 + ((b - Sqrt[a^2 + b^2])*E^(-c - d*x))/a] - 2*d*x*PolyLog[2, ((-b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a] - 2*PolyLog[3, ((-b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a]))/(Sqrt[a^2 + b^2]*(-b + Sqrt[a^2 + b^2])*d^3) - (6*a^2*b*(d^2*x^2*Log[1 + ((b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a] - 2*d*x*PolyLog[2, -(((b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a)] - 2*PolyLog[3, -(((b + Sqrt[a^2 + b^2])*E^(-c - d*x))/a)))/(Sqrt[a^2 + b^2]*(b + Sqrt[a^2 + b^2])*d^3) + (6*b^2*(d^2*x^2*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) + 2*d*x*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2]]) - 2*PolyLog[3, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2])]))/(Sqrt[a^2 + b^2]*d^3) - (6*b^2*(d^2*x^2*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) + 2*d*x*PolyLog[2, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))] - 2*PolyLog[3, -((a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]))]))/(Sqrt[a^2 + b^2]*d^3) + (6*a*Cosh[d*x]*(-2*d*x*Cosh[c] + (2 + d^2*x^2)*Sinh[c]))/d^3 + (6*a*((2 + d^2*x^2)*Cosh[c] - 2*d*x*Sinh[c])*Sinh[d*x])/d^3*(b + a*Sinh[c + d*x]))/(12*a^2*(a + b*Csch[c + d*x])) - (e^2*Csch[c + d*x]*((b*Log[b + a*Sinh[c + d*x]])/a^2 - Sinh[c + d*x]/a)*(b + a*Sinh[c + d*x]))/(2*d*(a + b*Csch[c + d*x])) + (e*f*Csch[c + d*x]*(b + a*Sinh[c + d*x])*(-2*a*Cosh[c + d*x] - b*(2*c*(c + d*x) - (c + d*x)^2 + 2*(c + d*x)*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]) + 2*(c + d*x)*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]) - 2*c*Log[a - 2*...
```

3.27.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx \\
 & \quad \downarrow \text{6128} \\
 & \int \frac{(e+fx)^2 \sinh(c+dx) \cosh^3(c+dx)}{a \sinh(c+dx) + b} dx \\
 & \quad \downarrow \text{6113} \\
 & \frac{\int (e+fx)^2 \cosh^3(c+dx) dx}{a} - \frac{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx)^2 \sin\left(ic+idx+\frac{\pi}{2}\right)^3 dx}{a} \\
 & \quad \downarrow \text{3792}
 \end{aligned}$$

3.27. $\int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \frac{2f^2 \int \cosh^3(c+dx) dx}{9d^2} + \frac{2}{3} \int (e+fx)^2 \cosh(c+dx) dx - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh^2(c+dx)}{3d} \\
 & \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx} \\
 & \quad \downarrow \text{3042} \\
 & \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx} + \\
 & \quad \frac{2f^2 \int \sin(ic+idx+\frac{\pi}{2})^3 dx}{9d^2} + \frac{2}{3} \int (e+fx)^2 \sin(ic+idx+\frac{\pi}{2}) dx - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh^2(c+dx)}{3d} \\
 & \quad \quad \quad \frac{a}{\downarrow \text{3113}} \\
 & \quad \quad \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx} + \\
 & \quad \frac{2if^2 \int (\sinh^2(c+dx)+1)d(-i \sinh(c+dx))}{9d^3} + \frac{2}{3} \int (e+fx)^2 \sin(ic+idx+\frac{\pi}{2}) dx - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh^2(c+dx)}{3d} \\
 & \quad \quad \quad \frac{a}{\downarrow \text{2009}} \\
 & \quad \quad \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx} + \\
 & \quad \frac{2}{3} \int (e+fx)^2 \sin(ic+idx+\frac{\pi}{2}) dx + \frac{2if^2(-\frac{1}{3}i \sinh^3(c+dx)-i \sinh(c+dx))}{9d^3} - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh^2(c+dx)}{3d} \\
 & \quad \quad \quad \frac{a}{\downarrow \text{3777}} \\
 & \quad \quad \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx} + \\
 & \quad \frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2if \int -i(e+fx) \sinh(c+dx) dx}{d} \right) + \frac{2if^2(-\frac{1}{3}i \sinh^3(c+dx)-i \sinh(c+dx))}{9d^3} - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh^2(c+dx)}{3d} \\
 & \quad \quad \quad \frac{a}{\downarrow \text{26}} \\
 & \quad \quad \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx} + \\
 & \quad \frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2f \int (e+fx) \sinh(c+dx) dx}{d} \right) + \frac{2if^2(-\frac{1}{3}i \sinh^3(c+dx)-i \sinh(c+dx))}{9d^3} - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh^2(c+dx)}{3d} \\
 & \quad \quad \quad \frac{a}{\downarrow \text{3042}} \\
 & \quad \quad \quad \frac{a}{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx} + \\
 & \quad \frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} - \frac{2f \int -i(e+fx) \sin(ic+idx) dx}{d} \right) + \frac{2if^2(-\frac{1}{3}i \sinh^3(c+dx)-i \sinh(c+dx))}{9d^3} - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx) \cosh^2(c+dx)}{3d} \\
 & \quad \quad \quad \frac{a}{\downarrow}
 \end{aligned}$$

3.27. $\int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned} & \downarrow \mathbf{26} \\ & \frac{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{-} + \\ & \frac{\frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \int (e+fx) \sin(ic+idx) dx}{d} \right) + \frac{a}{2if^2} \left(-\frac{1}{3} i \sinh^3(c+dx) - i \sinh(c+dx) \right) - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{(e+fx)^2 \sinh(c+dx)}{9d^3}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow \mathbf{3777} \\ & \frac{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{-} + \\ & \frac{\frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \cosh(c+dx) dx}{d} \right)}{d} \right) + \frac{2if^2 \left(-\frac{1}{3} i \sinh^3(c+dx) - i \sinh(c+dx) \right) - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2}}{a}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow \mathbf{3042} \\ & \frac{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{-} + \\ & \frac{\frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \int \sin\left(ic+idx+\frac{\pi}{2}\right) dx}{d} \right)}{d} \right) + \frac{2if^2 \left(-\frac{1}{3} i \sinh^3(c+dx) - i \sinh(c+dx) \right) - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2}}{a}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow \mathbf{3117} \\ & \frac{b \int \frac{(e+fx)^2 \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{-} + \\ & \frac{\frac{2if^2 \left(-\frac{1}{3} i \sinh^3(c+dx) - i \sinh(c+dx) \right) - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{d} \right)}{a}}{a} \end{aligned}$$

$$\begin{aligned} & \downarrow \mathbf{6099} \\ & \frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx)^2 \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \int (e+fx)^2 \cosh(c+dx) dx}{a^2} + \frac{\int (e+fx)^2 \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{-} + \\ & \frac{\frac{2if^2 \left(-\frac{1}{3} i \sinh^3(c+dx) - i \sinh(c+dx) \right) - \frac{2f(e+fx) \cosh^3(c+dx)}{9d^2} + \frac{2}{3} \left(\frac{(e+fx)^2 \sinh(c+dx)}{d} + \frac{2if \left(\frac{i(e+fx) \cosh(c+dx)}{d} - \frac{if \sinh(c+dx)}{d^2} \right)}{d} \right)}{a}}{a} \end{aligned}$$

$$\downarrow \mathbf{3042}$$

3.27. $\int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) +$$

$$\frac{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\int(e+fx)^2\sin\left(ic+idx+\frac{\pi}{2}\right)dx}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a}\right)}{a}$$

a
↓ 3777

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) +$$

$$\frac{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} - \frac{2if\int-i(e+fx)\sinh(c+dx)dx}{d}\right)}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a}\right)}{a}$$

a
↓ 26

$$b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} - \frac{2f\int(e+fx)\sinh(c+dx)dx}{d}\right)}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a}\right) +$$

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) +$$

a
↓ 3042

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) +$$

$$b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} - \frac{2f\int-i(e+fx)\sin\left(ic+idx\right)dx}{d}\right)}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a}\right)$$

a
↓ 26

3.27. $\int \frac{(e+fx)^2\cosh^3(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \frac{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\int(e+fx)\sin(ic+idx)dx}{d}\right)}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a}\right)}{a}$$

a
↓ 3777

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \frac{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\int\cosh(c+dx)dx}{d}\right)}{a^2}\right)}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a}\right)}{a}$$

a
↓ 3042

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \frac{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\int\sin\left(ic+idx+\frac{\pi}{2}\right)dx}{d}\right)}{a^2}\right)}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a}\right)}{a}$$

a
↓ 3117

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \frac{b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} + \frac{\int(e+fx)^2\cosh(c+dx)\sinh(c+dx)dx}{a} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{a^2}\right)}{a^2}\right)}{a}$$

a
↓ 5969

3.27. $\int \frac{(e+fx)^2 \cosh^3(c+dx)}{a+b\cosh(c+dx)} dx$

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \dots$$

$$b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} + \frac{(e+fx)^2\sinh^2(c+dx)}{2d} - \frac{f\int(e+fx)\sinh^2(c+dx)dx}{a} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right)}{a^2}\right)$$

↓ 3042

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \dots$$

$$b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} + \frac{(e+fx)^2\sinh^2(c+dx)}{2d} - \frac{f\int(e+fx)\sin(ic+idx)^2dx}{a} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right)}{a^2}\right)$$

↓ 25

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \dots$$

$$b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} + \frac{(e+fx)^2\sinh^2(c+dx)}{2d} + \frac{f\int(e+fx)\sin(ic+idx)^2dx}{a} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right)}{a^2}\right)$$

↓ 3791

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \dots$$

$$b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} + \frac{f\left(\frac{1}{2}\int(e+fx)dx + \frac{f\sinh^2(c+dx)}{4d^2} - \frac{(e+fx)\sinh(c+dx)\cosh(c+dx)}{2d}\right)}{a} + \frac{(e+fx)^2\sinh^2(c+dx)}{2d} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right)}{a^2}\right)$$

↓ 17

3.27. $\int \frac{(e+fx)^2\cosh^3(c+dx)}{a+b\text{CSch}(c+dx)} dx$

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \dots$$

$$b\left(\frac{(a^2+b^2)\int\frac{(e+fx)^2\cosh(c+dx)}{b+a\sinh(c+dx)}dx}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right)}{a^2}\right) + \frac{f\left(\frac{f\sinh^2(c+dx)}{4d^2} - \frac{(e+fx)\sinh(c+dx)}{2d}\right)}{d}$$

a

↓ 6095

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \dots$$

$$b\left(\frac{(a^2+b^2)\left(\int\frac{e^{c+dx}(e+fx)^2}{e^{c+dx}a+b-\sqrt{a^2+b^2}}dx + \int\frac{e^{c+dx}(e+fx)^2}{e^{c+dx}a+b+\sqrt{a^2+b^2}}dx - \frac{(e+fx)^3}{3af}\right)}{a^2} - \frac{b\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right)}{a^2}\right)$$

a

↓ 2620

$$\frac{2if^2(-\frac{1}{3}i\sinh^3(c+dx)-i\sinh(c+dx))}{9d^3} - \frac{2f(e+fx)\cosh^3(c+dx)}{9d^2} + \frac{2}{3}\left(\frac{(e+fx)^2\sinh(c+dx)}{d} + \frac{2if\left(\frac{i(e+fx)\cosh(c+dx)}{d} - \frac{if\sinh(c+dx)}{d^2}\right)}{d}\right) + \dots$$

$$b\left(\frac{(a^2+b^2)\left(-\frac{2f\int(e+fx)\log\left(\frac{e^{c+dx}a}{b-\sqrt{a^2+b^2}}+1\right)dx}{ad} - \frac{2f\int(e+fx)\log\left(\frac{e^{c+dx}a}{b+\sqrt{a^2+b^2}}+1\right)dx}{ad} + \frac{(e+fx)^2\log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}+1\right)}{ad} + \frac{(e+fx)^2\log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b}+1\right)}{ad} - (e+fx)\right)}{a^2}\right)$$

input `Int[((e + f*x)^2*Cosh[c + d*x]^3)/(a + b*CsCh[c + d*x]),x]`

output `$Aborted`

3.27.3.1 Defintions of rubi rules used

- rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1)/(b*(m + 1))), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3791 $\text{Int}[(c_.) + (d_.)(x_.)]((b_.)\sin[(e_.) + (f_.)(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[d*(b*\sin[e + f*x])^n/(f^2*n^2), x] + (-\text{Simp}[b*(c + d*x)*\cos[e + f*x]*(b*\sin[e + f*x])^{(n - 1)}/(f*n), x] + \text{Simp}[b^2*((n - 1)/n) \text{Int}[(c + d*x)*(b*\sin[e + f*x])^{(n - 2)}, x], x]) /; \text{FreeQ}\{b, c, d, e, f, x\} \&\& \text{GtQ}[n, 1]$

rule 3792 $\text{Int}[(c_.) + (d_.)(x_.)]^{(m_.)}((b_.)\sin[(e_.) + (f_.)(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[d*m*(c + d*x)^{(m - 1)}*(b*\sin[e + f*x])^n/(f^2*n^2), x] + (-\text{Simp}[b*(c + d*x)^m*\cos[e + f*x]*(b*\sin[e + f*x])^{(n - 1)}/(f*n), x] + \text{Simp}[b^2*((n - 1)/n) \text{Int}[(c + d*x)^m*(b*\sin[e + f*x])^{(n - 2)}, x], x] - \text{Simp}[d^2*m*((m - 1)/(f^2*n^2)) \text{Int}[(c + d*x)^{(m - 2)}*(b*\sin[e + f*x])^n, x], x]) /; \text{FreeQ}\{b, c, d, e, f, x\} \&\& \text{GtQ}[n, 1] \&\& \text{GtQ}[m, 1]$

rule 5969 $\text{Int}[\cosh[(a_.) + (b_.)(x_.)]((c_.) + (d_.)(x_.)]^{(m_.)}\sinh[(a_.) + (b_.)(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(c + d*x)^m*(\sinh[a + b*x])^{(n + 1)}/(b*(n + 1))), x] - \text{Simp}[d*(m/(b*(n + 1))) \text{Int}[(c + d*x)^{(m - 1)}*\sinh[a + b*x]^{(n + 1)}, x], x] /; \text{FreeQ}\{a, b, c, d, n, x\} \&\& \text{IGtQ}[m, 0] \&\& \text{NeQ}[n, -1]$

rule 6095 $\text{Int}[(\cosh[(c_.) + (d_.)(x_.)]*((e_.) + (f_.)(x_.)]^{(m_.)})/((a_.) + (b_.)\sinh[(c_.) + (d_.)(x_.)]), x_Symbol] \rightarrow \text{Simp}[-(e + f*x)^{(m + 1)}/(b*f*(m + 1)), x] + (\text{Int}[(e + f*x)^m*(E^(c + d*x)/(a - \text{Rt}[a^2 + b^2, 2] + b*E^(c + d*x))), x] + \text{Int}[(e + f*x)^m*(E^(c + d*x)/(a + \text{Rt}[a^2 + b^2, 2] + b*E^(c + d*x))), x]) /; \text{FreeQ}\{a, b, c, d, e, f, x\} \&\& \text{IGtQ}[m, 0] \&\& \text{NeQ}[a^2 + b^2, 0]$

rule 6099 $\text{Int}[(\cosh[(c_.) + (d_.)(x_.)]^{(n_.)}*((e_.) + (f_.)(x_.)]^{(m_.)})/((a_.) + (b_.)\sinh[(c_.) + (d_.)(x_.)]), x_Symbol] \rightarrow \text{Simp}[-a/b^2 \text{Int}[(e + f*x)^m*\cosh[c + d*x]^{(n - 2)}, x], x] + (\text{Simp}[1/b \text{Int}[(e + f*x)^m*\cosh[c + d*x]^{(n - 2)}*\sinh[c + d*x], x], x] + \text{Simp}[(a^2 + b^2)/b^2 \text{Int}[(e + f*x)^m*(\cosh[c + d*x]^{(n - 2)}/(a + b*\sinh[c + d*x])), x], x]) /; \text{FreeQ}\{a, b, c, d, e, f, x\} \&\& \text{IGtQ}[n, 1] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{IGtQ}[m, 0]$

rule 6113 $\text{Int}[(\cosh[(c_.) + (d_.)(x_.)]^{(p_.)}*((e_.) + (f_.)(x_.)]^{(m_.)}\sinh[(c_.) + (d_.)(x_.)]^{(n_.)})/((a_.) + (b_.)\sinh[(c_.) + (d_.)(x_.)]), x_Symbol] \rightarrow \text{Simp}[1/b \text{Int}[(e + f*x)^m*\cosh[c + d*x]^p*\sinh[c + d*x]^{(n - 1)}, x], x] - \text{Simp}[a/b \text{Int}[(e + f*x)^m*\cosh[c + d*x]^p*(\sinh[c + d*x]^{(n - 1)}/(a + b*\sinh[c + d*x])), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, x\} \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[n, 0] \&\& \text{IGtQ}[p, 0]$

rule 6128 `Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.))/(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Int[(e + f*x)^m*Sinh[c + d*x]*(F[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && IntegersQ[m, n]`

3.27.4 Maple [F]

$$\int \frac{(fx + e)^2 \cosh(dx + c)^3}{a + b \operatorname{csch}(dx + c)} dx$$

input `int((f*x+e)^2*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x)`

output `int((f*x+e)^2*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x)`

3.27.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4887 vs. 2(592) = 1184.

Time = 0.34 (sec) , antiderivative size = 4887, normalized size of antiderivative = 7.68

$$\int \frac{(e + fx)^2 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

input `integrate((f*x+e)^2*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="fricas")`


```

output -1/432*(18*a^3*d^2*f^2*x^2 + 18*a^3*d^2*e^2 - 2*(9*a^3*d^2*f^2*x^2 + 9*a^3
*d^2*e^2 - 6*a^3*d*e*f + 2*a^3*f^2 + 6*(3*a^3*d^2*e*f - a^3*d*f^2)*x)*cosh
(dx + c)^6 - 2*(9*a^3*d^2*f^2*x^2 + 9*a^3*d^2*e^2 - 6*a^3*d*e*f + 2*a^3*f
^2 + 6*(3*a^3*d^2*e*f - a^3*d*f^2)*x)*sinh(dx + c)^6 + 12*a^3*d*e*f + 27*
(2*a^2*b*d^2*f^2*x^2 + 2*a^2*b*d^2*e^2 - 2*a^2*b*d*e*f + a^2*b*f^2 + 2*(2*
a^2*b*d^2*e*f - a^2*b*d*f^2)*x)*cosh(dx + c)^5 + 3*(18*a^2*b*d^2*f^2*x^2
+ 18*a^2*b*d^2*e^2 - 18*a^2*b*d*e*f + 9*a^2*b*f^2 + 18*(2*a^2*b*d^2*e*f -
a^2*b*d*f^2)*x - 4*(9*a^3*d^2*f^2*x^2 + 9*a^3*d^2*e^2 - 6*a^3*d*e*f + 2*a^
3*f^2 + 6*(3*a^3*d^2*e*f - a^3*d*f^2)*x)*cosh(dx + c))*sinh(dx + c)^5 +
4*a^3*f^2 - 54*((3*a^3 + 4*a*b^2)*d^2*f^2*x^2 + (3*a^3 + 4*a*b^2)*d^2*e^2
- 2*(3*a^3 + 4*a*b^2)*d*e*f + 2*(3*a^3 + 4*a*b^2)*f^2 + 2*((3*a^3 + 4*a*b^
2)*d^2*e*f - (3*a^3 + 4*a*b^2)*d*f^2)*x)*cosh(dx + c)^4 - 3*(18*(3*a^3 +
4*a*b^2)*d^2*f^2*x^2 + 18*(3*a^3 + 4*a*b^2)*d^2*e^2 - 36*(3*a^3 + 4*a*b^2)
*d*e*f + 36*(3*a^3 + 4*a*b^2)*f^2 + 10*(9*a^3*d^2*f^2*x^2 + 9*a^3*d^2*e^2
- 6*a^3*d*e*f + 2*a^3*f^2 + 6*(3*a^3*d^2*e*f - a^3*d*f^2)*x)*cosh(dx + c)
^2 + 36*((3*a^3 + 4*a*b^2)*d^2*e*f - (3*a^3 + 4*a*b^2)*d*f^2)*x - 45*(2*a^
2*b*d^2*f^2*x^2 + 2*a^2*b*d^2*e^2 - 2*a^2*b*d*e*f + a^2*b*f^2 + 2*(2*a^2*b
*d^2*e*f - a^2*b*d*f^2)*x)*cosh(dx + c))*sinh(dx + c)^4 - 144*((a^2*b +
b^3)*d^3*f^2*x^3 + 3*(a^2*b + b^3)*d^3*e*f*x^2 + 3*(a^2*b + b^3)*d^3*e^2*x
+ 6*(a^2*b + b^3)*c*d^2*e^2 - 6*(a^2*b + b^3)*c^2*d*e*f + 2*(a^2*b + b...

```

3.27.6 Sympy [F]

$$\int \frac{(e + fx)^2 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx)^2 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

```
input integrate((f*x+e)**2*cosh(d*x+c)**3/(a+b*csch(d*x+c)),x)
```

```
output Integral((e + f*x)**2*cosh(c + d*x)**3/(a + b*csch(c + d*x)), x)
```

3.27.7 Maxima [F]

$$\int \frac{(e + fx)^2 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^2 \cosh(dx + c)^3}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^2*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output

```
-1/24*e^2*((3*a*b*e^(-d*x - c) - a^2 - 3*(3*a^2 + 4*b^2)*e^(-2*d*x - 2*c))
*e^(3*d*x + 3*c)/(a^3*d) + 24*(a^2*b + b^3)*(d*x + c)/(a^4*d) + (3*a*b*e^(-
-2*d*x - 2*c) + a^2*e^(-3*d*x - 3*c) + 3*(3*a^2 + 4*b^2)*e^(-d*x - c))/(a^
3*d) + 24*(a^2*b + b^3)*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a
^4*d) - 1/432*(144*(a^2*b*d^3*f^2*e^(3*c) + b^3*d^3*f^2*e^(3*c))*x^3 + 43
2*(a^2*b*d^3*e*f*e^(3*c) + b^3*d^3*e*f*e^(3*c))*x^2 - 2*(9*a^3*d^2*f^2*x^2
*e^(6*c) + 6*(3*d^2*e*f - d*f^2)*a^3*x*e^(6*c) - 2*(3*d*e*f - f^2)*a^3*e^(
6*c))*e^(3*d*x) + 27*(2*a^2*b*d^2*f^2*x^2*e^(5*c) + 2*(2*d^2*e*f - d*f^2)*
a^2*b*x*e^(5*c) - (2*d*e*f - f^2)*a^2*b*e^(5*c))*e^(2*d*x) + 54*(6*(d*e*f
- f^2)*a^3*e^(4*c) + 8*(d*e*f - f^2)*a*b^2*e^(4*c) - (3*a^3*d^2*f^2*e^(4*c)
) + 4*a*b^2*d^2*f^2*e^(4*c))*x^2 - 2*(3*(d^2*e*f - d*f^2)*a^3*e^(4*c) + 4*
(d^2*e*f - d*f^2)*a*b^2*e^(4*c))*x)*e^(d*x) + 54*(6*(d*e*f + f^2)*a^3*e^(2
*c) + 8*(d*e*f + f^2)*a*b^2*e^(2*c) + (3*a^3*d^2*f^2*e^(2*c) + 4*a*b^2*d^2
*f^2*e^(2*c))*x^2 + 2*(3*(d^2*e*f + d*f^2)*a^3*e^(2*c) + 4*(d^2*e*f + d*f^
2)*a*b^2*e^(2*c))*x)*e^(-d*x) + 27*(2*a^2*b*d^2*f^2*x^2*e^c + 2*(2*d^2*e*f
+ d*f^2)*a^2*b*x*e^c + (2*d*e*f + f^2)*a^2*b*e^c)*e^(-2*d*x) + 2*(9*a^3*d
^2*f^2*x^2 + 6*(3*d^2*e*f + d*f^2)*a^3*x + 2*(3*d*e*f + f^2)*a^3)*e^(-3*d*
x))*e^(-3*c)/(a^4*d^3) + integrate(-2*((a^3*b*f^2 + a*b^3*f^2)*x^2 + 2*(a^
3*b*e*f + a*b^3*e*f)*x - ((a^2*b^2*f^2*e^c + b^4*f^2*e^c)*x^2 + 2*(a^2*b^2
*e*f*e^c + b^4*e*f*e^c)*x)*e^(d*x))/(a^5*e^(2*d*x + 2*c) + 2*a^4*b*e^(d...
```

3.27.8 Giac [F]

$$\int \frac{(e + fx)^2 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e)^2 \cosh(dx + c)^3}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)^2*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)^2*cosh(d*x + c)^3/(b*csch(d*x + c) + a), x)`

3.27.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx)^2 \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx)^3 (e + fx)^2}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)^3*(e + f*x)^2)/(a + b/sinh(c + d*x)),x)`output `int((cosh(c + d*x)^3*(e + f*x)^2)/(a + b/sinh(c + d*x)), x)`

3.28 $\int \frac{(e+fx) \cosh^3(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$

3.28.1	Optimal result	259
3.28.2	Mathematica [A] (verified)	260
3.28.3	Rubi [A] (verified)	261
3.28.4	Maple [B] (verified)	268
3.28.5	Fricas [B] (verification not implemented)	269
3.28.6	Sympy [F]	269
3.28.7	Maxima [F]	270
3.28.8	Giac [F]	270
3.28.9	Mupad [F(-1)]	271

3.28.1 Optimal result

Integrand size = 26, antiderivative size = 400

$$\int \frac{(e+fx) \cosh^3(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = -\frac{bfx}{4a^2d} + \frac{b(a^2+b^2)(e+fx)^2}{2a^4f} - \frac{2f \cosh(c+dx)}{3ad^2}$$

$$- \frac{b^2f \cosh(c+dx)}{a^3d^2} - \frac{f \cosh^3(c+dx)}{9ad^2}$$

$$- \frac{b(a^2+b^2)(e+fx) \log\left(1 + \frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d}$$

$$- \frac{b(a^2+b^2)(e+fx) \log\left(1 + \frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d}$$

$$- \frac{b(a^2+b^2)f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{a^4d^2}$$

$$- \frac{b(a^2+b^2)f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{a^4d^2}$$

$$+ \frac{2(e+fx) \sinh(c+dx)}{3ad} + \frac{b^2(e+fx) \sinh(c+dx)}{a^3d}$$

$$+ \frac{bf \cosh(c+dx) \sinh(c+dx)}{4a^2d^2}$$

$$+ \frac{(e+fx) \cosh^2(c+dx) \sinh(c+dx)}{3ad}$$

$$- \frac{b(e+fx) \sinh^2(c+dx)}{2a^2d}$$

output
$$-1/4*b*f*x/a^2/d+1/2*b*(a^2+b^2)*(f*x+e)^2/a^4/f-2/3*f*cosh(d*x+c)/a/d^2-b^2*f*cosh(d*x+c)/a^3/d^2-1/9*f*cosh(d*x+c)^3/a/d^2-b*(a^2+b^2)*(f*x+e)*ln(1+a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^4/d-b*(a^2+b^2)*(f*x+e)*ln(1+a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^4/d-b*(a^2+b^2)*f*polylog(2,-a*exp(d*x+c)/(b-(a^2+b^2)^(1/2)))/a^4/d^2-b*(a^2+b^2)*f*polylog(2,-a*exp(d*x+c)/(b+(a^2+b^2)^(1/2)))/a^4/d^2+2/3*(f*x+e)*sinh(d*x+c)/a/d+b^2*(f*x+e)*sinh(d*x+c)/a^3/d+1/4*b*f*cosh(d*x+c)*sinh(d*x+c)/a^2/d^2+1/3*(f*x+e)*cosh(d*x+c)^2*sinh(d*x+c)/a/d-1/2*b*(f*x+e)*sinh(d*x+c)^2/a^2/d$$

3.28.2 Mathematica [A] (verified)

Time = 1.18 (sec) , antiderivative size = 696, normalized size of antiderivative = 1.74

$$\int \frac{(e + fx) \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \frac{36a^2bc^2f + 36b^3c^2f - 36a^2bd^2fx^2 - 36b^3d^2fx^2 + 54a^3f \cosh(c + dx) + 72ab^2f \cosh(c + dx) + 18a^2bd^2f \cosh^2(c + dx) + 18a^2bd^2f \cosh(c + dx) + 18a^2bd^2f \cosh^2(c + dx)}{a^4d^2}$$

input `Integrate[((e + f*x)*Cosh[c + d*x]^3)/(a + b*Csch[c + d*x]),x]`

output
$$-1/72*(36*a^2*b*c^2*f + 36*b^3*c^2*f - 36*a^2*b*d^2*f*x^2 - 36*b^3*d^2*f*x^2 + 54*a^3*f*Cosh[c + d*x] + 72*a*b^2*f*Cosh[c + d*x] + 18*a^2*b*d*f*x*Cosh[2*(c + d*x)] + 2*a^3*f*Cosh[3*(c + d*x)] + 72*a^2*b*c*f*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2])] + 72*b^3*c*f*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2])] + 72*a^2*b*d*f*x*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2])] + 72*b^3*d*f*x*Log[1 + (a*E^(c + d*x))/(b - Sqrt[a^2 + b^2])] + 72*a^2*b*c*f*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])] + 72*b^3*c*f*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])] + 72*a^2*b*d*f*x*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])] + 72*b^3*d*f*x*Log[1 + (a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])] - 72*a^2*b*c*f*Log[a - 2*b*E^(c + d*x) - a*E^(2*(c + d*x))] - 72*b^3*c*f*Log[a - 2*b*E^(c + d*x) - a*E^(2*(c + d*x))] + 72*a^2*b*d*e*Log[b + a*Sinh[c + d*x]] + 72*b^3*d*e*Log[b + a*Sinh[c + d*x]] + 72*b*(a^2 + b^2)*f*PolyLog[2, (a*E^(c + d*x))/(-b + Sqrt[a^2 + b^2])] + 72*b*(a^2 + b^2)*f*PolyLog[2, -(a*E^(c + d*x))/(b + Sqrt[a^2 + b^2])] - 72*a^3*d*e*Sinh[c + d*x] - 72*a*b^2*d*e*Sinh[c + d*x] - 54*a^3*d*f*x*Sinh[c + d*x] - 72*a*b^2*d*f*x*Sinh[c + d*x] + 36*a^2*b*d*e*Sinh[c + d*x]^2 - 24*a^3*d*e*Sinh[c + d*x]^3 - 9*a^2*b*f*Sinh[2*(c + d*x)] - 6*a^3*d*f*x*Sinh[3*(c + d*x)]/(a^4*d^2)$$

3.28.3 Rubi [A] (verified)

Time = 2.30 (sec) , antiderivative size = 360, normalized size of antiderivative = 0.90, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6128, 6113, 3042, 3791, 3042, 3777, 26, 3042, 26, 3118, 6099, 3042, 3777, 26, 3042, 26, 3118, 5969, 3042, 25, 3115, 24, 6095, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(e+fx) \cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx \\
 & \quad \downarrow \text{6128} \\
 & \int \frac{(e+fx) \sinh(c+dx) \cosh^3(c+dx)}{a \sinh(c+dx) + b} dx \\
 & \quad \downarrow \text{6113} \\
 & \frac{\int (e+fx) \cosh^3(c+dx) dx}{a} - \frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \frac{\int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right)^3 dx}{a} \\
 & \quad \downarrow \text{3791} \\
 & \frac{\frac{2}{3} \int (e+fx) \cosh(c+dx) dx - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} - \frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
 & \frac{\frac{2}{3} \int (e+fx) \sin\left(ic+idx+\frac{\pi}{2}\right) dx - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & \quad \downarrow \text{3777} \\
 & -\frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
 & \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{if \int -i \sinh(c+dx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

3.28. $\int \frac{(e+fx) \cosh^3(c+dx)}{a+b\operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
& \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sinh(c+dx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
& \quad \frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
& \quad \downarrow \mathbf{3042} \\
& \quad \frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
& \quad \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int -i \sin(ic+idx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
& \quad \downarrow \mathbf{26} \\
& \quad \frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} + \\
& \quad \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} + \frac{if \int \sin(ic+idx) dx}{d} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
& \quad \downarrow \mathbf{3118} \\
& \quad \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
& \quad \frac{b \int \frac{(e+fx) \cosh^3(c+dx)}{b+a \sinh(c+dx)} dx}{a} \\
& \quad \downarrow \mathbf{6099} \\
& \quad \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
& \quad \frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \int (e+fx) \cosh(c+dx) dx}{a^2} + \frac{\int (e+fx) \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a} \\
& \quad \downarrow \mathbf{3042} \\
& \quad \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
& \quad \frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \int (e+fx) \sin(ic+idx + \frac{\pi}{2}) dx}{a^2} + \frac{\int (e+fx) \cosh(c+dx) \sinh(c+dx) dx}{a} \right)}{a} \\
& \quad \downarrow \mathbf{3777}
\end{aligned}$$

3.28. $\int \frac{(e+fx) \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{aligned}
 & \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{if \int -i \sinh(c+dx) dx}{d} \right)}{a^2} + \frac{\int (e+fx) \cosh(c+dx) \sinh(c+dx) dx}{a} \right) \\
 & \qquad \qquad \qquad \downarrow \text{26} \\
 & \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int \sinh(c+dx) dx}{d} \right)}{a^2} + \frac{\int (e+fx) \cosh(c+dx) \sinh(c+dx) dx}{a} \right) \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \int -i \sin(ic+idx) dx}{d} \right)}{a^2} + \frac{\int (e+fx) \cosh(c+dx) \sinh(c+dx) dx}{a} \right) \\
 & \qquad \qquad \qquad \downarrow \text{26} \\
 & \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} + \frac{if \int \sin(ic+idx) dx}{d} \right)}{a^2} + \frac{\int (e+fx) \cosh(c+dx) \sinh(c+dx) dx}{a} \right) \\
 & \qquad \qquad \qquad \downarrow \text{3118} \\
 & \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} + \frac{\int (e+fx) \cosh(c+dx) \sinh(c+dx) dx}{a} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{a^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{5969} \\
 & \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 & b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} + \frac{\frac{(e+fx) \sinh^2(c+dx)}{2d} - \frac{f \int \sinh^2(c+dx) dx}{2d}}{a} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{a^2} \right) \\
 & \qquad \qquad \qquad a
 \end{aligned}$$

3.28. $\int \frac{(e+fx) \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\begin{array}{c}
 \downarrow \text{3042} \\
 \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 \frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} + \frac{(e+fx) \sinh^2(c+dx)}{2d} - \frac{f \int \sin(ic+idx)^2 dx}{2d} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{a^2} \right)}{a} \\
 \downarrow \text{25} \\
 \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 \frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} + \frac{(e+fx) \sinh^2(c+dx)}{2d} + \frac{f \int \sin(ic+idx)^2 dx}{2d} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{a^2} \right)}{a} \\
 \downarrow \text{3115} \\
 \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 \frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} + \frac{f \left(\frac{\int 1 dx}{2} - \frac{\sinh(c+dx) \cosh(c+dx)}{2d} \right)}{2d} + \frac{(e+fx) \sinh^2(c+dx)}{2d} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{a^2} \right)}{a} \\
 \downarrow \text{24} \\
 \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 \frac{b \left(\frac{(a^2+b^2) \int \frac{(e+fx) \cosh(c+dx)}{b+a \sinh(c+dx)} dx}{a^2} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{a^2} + \frac{(e+fx) \sinh^2(c+dx)}{2d} + \frac{f \left(\frac{x}{2} - \frac{\sinh(c+dx) \cosh(c+dx)}{2d} \right)}{a} \right)}{a} \\
 \downarrow \text{6095} \\
 \frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{a} \\
 \frac{b \left(\frac{(a^2+b^2) \left(\int \frac{e^{c+dx} (e+fx)}{e^{c+dx} a + b - \sqrt{a^2+b^2}} dx + \int \frac{e^{c+dx} (e+fx)}{e^{c+dx} a + b + \sqrt{a^2+b^2}} dx - \frac{(e+fx)^2}{2af} \right)}{a^2} - \frac{b \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right)}{a^2} + \frac{(e+fx) \sinh^2(c+dx)}{2d} + \frac{f \left(\frac{x}{2} \right)}{a} \right)}{a} \\
 \downarrow \text{2620}
 \end{array}$$

3.28. $\int \frac{(e+fx) \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

$$\frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{b \left(\frac{(a^2+b^2) \left(-\frac{f \int \log\left(\frac{e^{c+dx} a}{b-\sqrt{a^2+b^2}}+1\right) dx}{ad} - \frac{f \int \log\left(\frac{e^{c+dx} a}{b+\sqrt{a^2+b^2}}+1\right) dx}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}+1\right)}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b}\right)}{ad} - \frac{(e+fx)^2}{2af} \right)}{a^2} \right)} - \frac{b \left(\frac{e^{c+dx}}{a} \right)}{a}$$

↓ 2715

$$\frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{b \left(\frac{(a^2+b^2) \left(-\frac{f \int e^{-c-dx} \log\left(\frac{e^{c+dx} a}{b-\sqrt{a^2+b^2}}+1\right) de^{c+dx}}{ad^2} - \frac{f \int e^{-c-dx} \log\left(\frac{e^{c+dx} a}{b+\sqrt{a^2+b^2}}+1\right) de^{c+dx}}{ad^2} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}+1\right)}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b}\right)}{ad} - \frac{(e+fx)^2}{2af} \right)}{a^2} \right)} - \frac{b \left(\frac{e^{c+dx}}{a} \right)}{a}$$

↓ 2838

$$\frac{\frac{2}{3} \left(\frac{(e+fx) \sinh(c+dx)}{d} - \frac{f \cosh(c+dx)}{d^2} \right) - \frac{f \cosh^3(c+dx)}{9d^2} + \frac{(e+fx) \sinh(c+dx) \cosh^2(c+dx)}{3d}}{b \left(\frac{(a^2+b^2) \left(\frac{f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}\right)}{ad^2} + \frac{f \operatorname{PolyLog}\left(2, -\frac{ae^{c+dx}}{b+\sqrt{a^2+b^2}}\right)}{ad^2} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{b-\sqrt{a^2+b^2}}+1\right)}{ad} + \frac{(e+fx) \log\left(\frac{ae^{c+dx}}{\sqrt{a^2+b^2}+b}\right)}{ad} - \frac{(e+fx)^2}{2af} \right)}{a^2} \right)} - \frac{b \left(\frac{e^{c+dx}}{a} \right)}{a}$$

input `Int[((e + f*x)*Cosh[c + d*x]^3)/(a + b*CSch[c + d*x]),x]`

output `(-1/9*(f*Cosh[c + d*x]^3)/d^2 + ((e + f*x)*Cosh[c + d*x]^2*Sinh[c + d*x])/(3*d) + (2*(-((f*Cosh[c + d*x])/d^2) + ((e + f*x)*Sinh[c + d*x])/d))/3/a - (b*(((a^2 + b^2)*(-1/2*(e + f*x)^2/(a*f) + ((e + f*x)*Log[1 + (a*E^(c + d*x))]/(b - Sqrt[a^2 + b^2])))/(a*d) + ((e + f*x)*Log[1 + (a*E^(c + d*x))]/(b + Sqrt[a^2 + b^2])))/(a*d) + (f*PolyLog[2, -(a*E^(c + d*x))/(b - Sqrt[a^2 + b^2]]))/(a*d^2) + (f*PolyLog[2, -(a*E^(c + d*x))/(b + Sqrt[a^2 + b^2]]))/(a*d^2))/a^2 - (b*(-((f*Cosh[c + d*x])/d^2) + ((e + f*x)*Sinh[c + d*x])/d))/a^2 + (((e + f*x)*Sinh[c + d*x]^2)/(2*d) + (f*(x/2 - (Cosh[c + d*x]*Sinh[c + d*x])/(2*d)))/(2*d))/a)/a`

3.28.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3118 `Int[sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-c + d*x)^m*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3791 `Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)*(b*Sine[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

rule 5969 `Int[Cosh[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[(c + d*x)^m*(Sinh[a + b*x]^(n + 1)/(b*(n + 1))), x] - Simp[d*(m/(b*(n + 1))) Int[(c + d*x)^(m - 1)*Sinh[a + b*x]^(n + 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 6095 `Int[(Cosh[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)), x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 + b^2, 2] + b*E^(c + d*x))), x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 + b^2, 2] + b*E^(c + d*x))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 + b^2, 0]`

rule 6099 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(n_))*((e_.) + (f_.)*(x_))^(m_.)/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-a/b^2 Int[(e + f*x)^m*Cosh[c + d*x]^(n - 2), x], x] + (Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^(n - 2)*Sinh[c + d*x], x], x] + Simp[(a^2 + b^2)/b^2 Int[(e + f*x)^m*(Cosh[c + d*x]^(n - 2)/(a + b*Sinh[c + d*x])), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[n, 1] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]`

rule 6113 `Int[(Cosh[(c_.) + (d_.)*(x_)]^(p_.))*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)]^(n_.)/((a_.) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[1/b Int[(e + f*x)^m*Cosh[c + d*x]^p*Sinh[c + d*x]^(n - 1), x], x] - Simp[a/b Int[(e + f*x)^m*Cosh[c + d*x]^p*(Sinh[c + d*x]^(n - 1)/(a + b*Sinh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6128 `Int[(((e_.) + (f_.)*(x_.))^(m_.)*(F_)[(c_.) + (d_.)*(x_.)]^(n_.))/(Csch[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Int[(e + f*x)^m*Sinh[c + d*x]*(F[c + d*x]^n/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && IntegersQ[m, n]`

3.28.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1101 vs. $2(372) = 744$.

Time = 17.62 (sec) , antiderivative size = 1102, normalized size of antiderivative = 2.76

method	result	size
risch	Expression too large to display	1102

input `int((f*x+e)*cosh(d*x+c)^3/(a+b*cosh(d*x+c)),x,method=_RETURNVERBOSE)`

output `-2/d^2*b^3/a^4*c*f*ln(exp(d*x+c))+1/d^2*b^3/a^4*c*f*ln(exp(2*d*x+2*c)*a+2*exp(d*x+c)*b-a)+1/d^2*b^3/a^4*f*c^2-1/d*b^3/a^4*e*ln(exp(2*d*x+2*c)*a+2*exp(d*x+c)*b-a)+2/d*b^3/a^4*e*ln(exp(d*x+c))-1/d^2*b^3/a^4*f*dilog((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))-1/d^2*b^3/a^4*f*dilog((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))+1/2/a^2*b*f*x^2-1/a^2*b*e*x-1/72*(3*d*f*x+3*d*e+f)/a/d^2*exp(-3*d*x-3*c)-1/16*b*(2*d*f*x+2*d*e-f)/a^2/d^2*exp(2*d*x+2*c)-1/8*(3*a^2+4*b^2)*(d*f*x+d*e+f)/a^3/d^2*exp(-d*x-c)-1/16*b*(2*d*f*x+2*d*e+f)/a^2/d^2*exp(-2*d*x-2*c)+2/d*b^3/a^4*f*c*x-1/d*b^3/a^4*f*ln((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))*x-1/d*b^3/a^4*f*ln((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))*x-1/d^2*b^3/a^4*f*ln((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))*c-1/d^2*b^3/a^4*f*ln((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))*c+1/72*(3*d*f*x+3*d*e-f)/a/d^2*exp(3*d*x+3*c)+1/2*b^3/a^4*f*x^2-b^3/a^4*e*x+1/8*(3*a^2*d*f*x+4*b^2*d*f*x+3*a^2*d*e+4*b^2*d*e-3*a^2*f-4*b^2*f)/a^3/d^2*exp(d*x+c)+2/d/a^2*b*f*c*x-1/d/a^2*b*f*ln((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))*x-1/d/a^2*b*f*ln((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))*x-1/d^2/a^2*b*f*ln((-a*exp(d*x+c)+(a^2+b^2)^(1/2)-b)/(-b+(a^2+b^2)^(1/2)))*c-1/d^2/a^2*b*f*ln((a*exp(d*x+c)+(a^2+b^2)^(1/2)+b)/(b+(a^2+b^2)^(1/2)))*c+1/d^2/a^2*b*c*f*ln(exp(2*d*x+2*c)*a+2*exp(d*x+c)*b-a)-2/d^2/a^2*b*c*f*ln(exp(d*x+c))+1/d^2/a^2*b*f*c^2-1/d^2/a^2*b*f*dilog(...`

3.28.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2465 vs. $2(370) = 740$.

Time = 0.31 (sec) , antiderivative size = 2465, normalized size of antiderivative = 6.16

$$\int \frac{(e + fx) \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((f*x+e)*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="fracas")
```

```
output 1/144*(2*(3*a^3*d*f*x + 3*a^3*d*e - a^3*f)*cosh(d*x + c)^6 + 2*(3*a^3*d*f*x
+ 3*a^3*d*e - a^3*f)*sinh(d*x + c)^6 - 6*a^3*d*f*x - 9*(2*a^2*b*d*f*x +
2*a^2*b*d*e - a^2*b*f)*cosh(d*x + c)^5 - 3*(6*a^2*b*d*f*x + 6*a^2*b*d*e -
3*a^2*b*f - 4*(3*a^3*d*f*x + 3*a^3*d*e - a^3*f)*cosh(d*x + c))*sinh(d*x +
c)^5 - 6*a^3*d*e + 18*((3*a^3 + 4*a*b^2)*d*f*x + (3*a^3 + 4*a*b^2)*d*e - (
3*a^3 + 4*a*b^2)*f)*cosh(d*x + c)^4 + 3*(6*(3*a^3 + 4*a*b^2)*d*f*x + 6*(3*
a^3 + 4*a*b^2)*d*e + 10*(3*a^3*d*f*x + 3*a^3*d*e - a^3*f)*cosh(d*x + c)^2
- 6*(3*a^3 + 4*a*b^2)*f - 15*(2*a^2*b*d*f*x + 2*a^2*b*d*e - a^2*b*f)*cosh(
d*x + c))*sinh(d*x + c)^4 - 2*a^3*f + 72*((a^2*b + b^3)*d^2*f*x^2 + 2*(a^2
*b + b^3)*d^2*e*x + 4*(a^2*b + b^3)*c*d*e - 2*(a^2*b + b^3)*c^2*f)*cosh(d*
x + c)^3 + 2*(36*(a^2*b + b^3)*d^2*f*x^2 + 72*(a^2*b + b^3)*d^2*e*x + 144*
(a^2*b + b^3)*c*d*e - 72*(a^2*b + b^3)*c^2*f + 20*(3*a^3*d*f*x + 3*a^3*d*e
- a^3*f)*cosh(d*x + c)^3 - 45*(2*a^2*b*d*f*x + 2*a^2*b*d*e - a^2*b*f)*cos
h(d*x + c)^2 + 36*((3*a^3 + 4*a*b^2)*d*f*x + (3*a^3 + 4*a*b^2)*d*e - (3*a^
3 + 4*a*b^2)*f)*cosh(d*x + c))*sinh(d*x + c)^3 - 18*((3*a^3 + 4*a*b^2)*d*f
*x + (3*a^3 + 4*a*b^2)*d*e + (3*a^3 + 4*a*b^2)*f)*cosh(d*x + c)^2 + 6*(5*(
3*a^3*d*f*x + 3*a^3*d*e - a^3*f)*cosh(d*x + c)^4 - 3*(3*a^3 + 4*a*b^2)*d*f
*x - 15*(2*a^2*b*d*f*x + 2*a^2*b*d*e - a^2*b*f)*cosh(d*x + c)^3 - 3*(3*a^3
+ 4*a*b^2)*d*e + 18*((3*a^3 + 4*a*b^2)*d*f*x + (3*a^3 + 4*a*b^2)*d*e - (3
*a^3 + 4*a*b^2)*f)*cosh(d*x + c)^2 - 3*(3*a^3 + 4*a*b^2)*f + 36*((a^2*b...
```

3.28.6 Sympy [F]

$$\int \frac{(e + fx) \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(e + fx) \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx$$

```
input integrate((f*x+e)*cosh(d*x+c)**3/(a+b*csch(d*x+c)),x)
```

```
output Integral((e + f*x)*cosh(c + d*x)**3/(a + b*csch(c + d*x)), x)
```

3.28. $\int \frac{(e+fx) \cosh^3(c+dx)}{a+b \operatorname{CSch}(c+dx)} dx$

3.28.7 Maxima [F]

$$\int \frac{(e + fx) \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e) \cosh(dx + c)^3}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/24*e*((3*a*b*e^(-d*x - c) - a^2 - 3*(3*a^2 + 4*b^2)*e^(-2*d*x - 2*c))*e^(3*d*x + 3*c)/(a^3*d) + 24*(a^2*b + b^3)*(d*x + c)/(a^4*d) + (3*a*b*e^(-2*d*x - 2*c) + a^2*e^(-3*d*x - 3*c) + 3*(3*a^2 + 4*b^2)*e^(-d*x - c))/(a^3*d) + 24*(a^2*b + b^3)*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a^4*d) - 1/144*f*((72*(a^2*b*d^2*e^(3*c) + b^3*d^2*e^(3*c))*x^2 - 2*(3*a^3*d*x*e^(6*c) - a^3*e^(6*c))*e^(3*d*x) + 9*(2*a^2*b*d*x*e^(5*c) - a^2*b*e^(5*c))*e^(2*d*x) + 18*(3*a^3*e^(4*c) + 4*a*b^2*e^(4*c) - (3*a^3*d*e^(4*c) + 4*a*b^2*d*e^(4*c))*x)*e^(d*x) + 18*(3*a^3*e^(2*c) + 4*a*b^2*e^(2*c) + (3*a^3*d*e^(2*c) + 4*a*b^2*d*e^(2*c))*x)*e^(-d*x) + 9*(2*a^2*b*d*x*e^c + a^2*b*e^c)*e^(-2*d*x) + 2*(3*a^3*d*x + a^3)*e^(-3*d*x))*e^(-3*c)/(a^4*d^2) - 18*integrate(16*((a^2*b^2*e^c + b^4*e^c)*x*e^(d*x) - (a^3*b + a*b^3)*x)/(a^5*e^(2*d*x + 2*c) + 2*a^4*b*e^(d*x + c) - a^5), x)`

3.28.8 Giac [F]

$$\int \frac{(e + fx) \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{(fx + e) \cosh(dx + c)^3}{b \operatorname{csch}(dx + c) + a} dx$$

input `integrate((f*x+e)*cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `integrate((f*x + e)*cosh(d*x + c)^3/(b*csch(d*x + c) + a), x)`

3.28.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(e + fx) \cosh^3(c + dx)}{a + b \operatorname{csch}(c + dx)} dx = \int \frac{\cosh(c + dx)^3 (e + fx)}{a + \frac{b}{\sinh(c + dx)}} dx$$

input `int((cosh(c + d*x)^3*(e + f*x))/(a + b/sinh(c + d*x)),x)`output `int((cosh(c + d*x)^3*(e + f*x))/(a + b/sinh(c + d*x)), x)`

3.29 $\int \frac{\cosh^3(c+dx)}{a+b\mathbf{csch}(c+dx)} dx$

3.29.1	Optimal result	272
3.29.2	Mathematica [A] (verified)	272
3.29.3	Rubi [A] (verified)	273
3.29.4	Maple [B] (verified)	275
3.29.5	Fricas [B] (verification not implemented)	276
3.29.6	Sympy [F]	276
3.29.7	Maxima [B] (verification not implemented)	277
3.29.8	Giac [A] (verification not implemented)	277
3.29.9	Mupad [B] (verification not implemented)	278

3.29.1 Optimal result

Integrand size = 21, antiderivative size = 85

$$\int \frac{\cosh^3(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = -\frac{b(a^2+b^2)\log(b+a\sinh(c+dx))}{a^4d} + \frac{(a^2+b^2)\sinh(c+dx)}{a^3d} - \frac{b\sinh^2(c+dx)}{2a^2d} + \frac{\sinh^3(c+dx)}{3ad}$$

output `-b*(a^2+b^2)*ln(b+a*sinh(d*x+c))/a^4/d+(a^2+b^2)*sinh(d*x+c)/a^3/d-1/2*b*sinh(d*x+c)^2/a^2/d+1/3*sinh(d*x+c)^3/a/d`

3.29.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.88

$$\int \frac{\cosh^3(c+dx)}{a+b\mathbf{csch}(c+dx)} dx = \frac{-6b(a^2+b^2)\log(b+a\sinh(c+dx)) + 6a(a^2+b^2)\sinh(c+dx) - 3a^2b\sinh^2(c+dx) + 2a^3\sinh^3(c+dx)}{6a^4d}$$

input `Integrate[Cosh[c + d*x]^3/(a + b*Csch[c + d*x]),x]`

output `(-6*b*(a^2 + b^2)*Log[b + a*Sinh[c + d*x]] + 6*a*(a^2 + b^2)*Sinh[c + d*x] - 3*a^2*b*Sinh[c + d*x]^2 + 2*a^3*Sinh[c + d*x]^3)/(6*a^4*d)`

3.29.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.88, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.524$, Rules used = {3042, 4360, 26, 26, 3042, 26, 3316, 26, 27, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(ic+idx)^3}{a+ib\operatorname{csc}(ic+idx)} dx \\
 & \quad \downarrow \text{4360} \\
 & \int \frac{i\sinh(c+dx)\cosh^3(c+dx)}{ia\sinh(c+dx)+ib} dx \\
 & \quad \downarrow \text{26} \\
 & i \int -\frac{i\cosh^3(c+dx)\sinh(c+dx)}{b+a\sinh(c+dx)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\sinh(c+dx)\cosh^3(c+dx)}{a\sinh(c+dx)+b} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i\sin(ic+idx)\cos(ic+idx)^3}{b-ia\sin(ic+idx)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\cos(ic+idx)^3\sin(ic+idx)}{b-ia\sin(ic+idx)} dx \\
 & \quad \downarrow \text{3316} \\
 & \frac{i \int -\frac{i\sinh(c+dx)(\sinh^2(c+dx)a^2+a^2)}{b+a\sinh(c+dx)} d(a\sinh(c+dx))}{a^3d} \\
 & \quad \downarrow \text{26} \\
 & \frac{\int \frac{\sinh(c+dx)(\sinh^2(c+dx)a^2+a^2)}{b+a\sinh(c+dx)} d(a\sinh(c+dx))}{a^3d}
 \end{aligned}$$

3.29. $\int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

$$\begin{array}{c}
\downarrow 27 \\
\int \frac{a \sinh(c+dx) (\sinh^2(c+dx)a^2+a^2)}{b+a \sinh(c+dx)} d(a \sinh(c+dx)) \\
a^4 d \\
\downarrow 522 \\
\int \left(\sinh^2(c+dx)a^2 + \left(\frac{b^2}{a^2} + 1\right) a^2 - b \sinh(c+dx)a - \frac{b(a^2+b^2)}{b+a \sinh(c+dx)} \right) d(a \sinh(c+dx)) \\
a^4 d \\
\downarrow 2009 \\
\frac{\frac{1}{3}a^3 \sinh^3(c+dx) + a(a^2+b^2) \sinh(c+dx) - b(a^2+b^2) \log(a \sinh(c+dx) + b) - \frac{1}{2}a^2 b \sinh^2(c+dx)}{a^4 d}
\end{array}$$

input `Int[Cosh[c + d*x]^3/(a + b*Csch[c + d*x]),x]`

output `(-(b*(a^2 + b^2)*Log[b + a*Sinh[c + d*x]]) + a*(a^2 + b^2)*Sinh[c + d*x] - (a^2*b*Sinh[c + d*x]^2)/2 + (a^3*Sinh[c + d*x]^3)/3)/(a^4*d)`

3.29.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 522 `Int[((e_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3316 Int[cos[(e_.) + (f_.)*(x_)]^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^m*(c + (d/b)*x)^n*(b^2 - x^2)^((p - 1)/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && IntegerQ[(p - 1)/2] && NeQ[a^2 - b^2, 0]
```

```
rule 4360 Int[(cos[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.), x_Symbol] := Int[(g*Cos[e + f*x])^p*((b + a*Sin[e + f*x])^m/Sin[e + f*x]^m), x] /; FreeQ[{a, b, e, f, g, p}, x] && IntegerQ[m]
```

3.29.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. 2(81) = 162.

Time = 17.86 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.87

method	result
risch	$\frac{bx}{a^2} + \frac{b^3x}{a^4} + \frac{e^{3dx+3c}}{24da} - \frac{be^{2dx+2c}}{8da^2} + \frac{3e^{dx+c}}{8ad} + \frac{e^{dx+c}b^2}{2a^3d} - \frac{3e^{-dx-c}}{8ad} - \frac{e^{-dx-c}b^2}{2a^3d} - \frac{be^{-2dx-2c}}{8da^2} - \frac{e^{-3dx-3c}}{24da}$
derivativedivides	$\frac{1}{3a(1+\tanh(\frac{dx}{2} + \frac{c}{2}))^3} - \frac{-a+b}{2a^2(1+\tanh(\frac{dx}{2} + \frac{c}{2}))^2} - \frac{2a^2-ab+2b^2}{2a^3(1+\tanh(\frac{dx}{2} + \frac{c}{2}))} + \frac{(a^2+b^2)b \ln(1+\tanh(\frac{dx}{2} + \frac{c}{2}))}{a^4} + \frac{2b(-\frac{a^2}{2} - \frac{b^2}{2}) \ln(\dots)}{a^4}$
default	$\frac{1}{3a(1+\tanh(\frac{dx}{2} + \frac{c}{2}))^3} - \frac{-a+b}{2a^2(1+\tanh(\frac{dx}{2} + \frac{c}{2}))^2} - \frac{2a^2-ab+2b^2}{2a^3(1+\tanh(\frac{dx}{2} + \frac{c}{2}))} + \frac{(a^2+b^2)b \ln(1+\tanh(\frac{dx}{2} + \frac{c}{2}))}{a^4} + \frac{2b(-\frac{a^2}{2} - \frac{b^2}{2}) \ln(\dots)}{a^4}$

```
input int(cosh(d*x+c)^3/(a+b*cSch(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output b*x/a^2+1/a^4*b^3*x+1/24/d/a*exp(3*d*x+3*c)-1/8*b/d/a^2*exp(2*d*x+2*c)+3/8/a/d*exp(d*x+c)+1/2/a^3/d*exp(d*x+c)*b^2-3/8/a/d*exp(-d*x-c)-1/2/a^3/d*exp(-d*x-c)*b^2-1/8*b/d/a^2*exp(-2*d*x-2*c)-1/24/d/a*exp(-3*d*x-3*c)+2*b/d/a^2*c+2*b^3/d/a^4*c-b/d/a^2*ln(exp(2*d*x+2*c))+2*b/a*exp(d*x+c)-1)-b^3/d/a^4*ln(exp(2*d*x+2*c)+2*b/a*exp(d*x+c)-1)
```

3.29.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 652 vs. 2(81) = 162.

Time = 0.27 (sec) , antiderivative size = 652, normalized size of antiderivative = 7.67

$$\int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$$

$$= \frac{a^3 \cosh(dx+c)^6 + a^3 \sinh(dx+c)^6 - 3a^2b \cosh(dx+c)^5 + 24(a^2b+b^3)dx \cosh(dx+c)^3 + 3(2a^3 \cosh(dx+c)^4 + 24a^2b \cosh(dx+c)^3 + 3(2a^3 \cosh(dx+c)^2 - 5a^2b \cosh(dx+c) + 3a^3 + 4a^2b^2) \sinh(dx+c)^4 - 3a^2b \cosh(dx+c) + 2(10a^3 \cosh(dx+c)^3 - 15a^2b \cosh(dx+c)^2 + 12(a^2b+b^3)dx + 6(3a^3 + 4a^2b^2) \cosh(dx+c)) \sinh(dx+c)^3 - a^3 - 3(3a^3 + 4a^2b^2) \cosh(dx+c)^2 + 3(5a^3 \cosh(dx+c)^4 - 10a^2b \cosh(dx+c)^3 + 24(a^2b+b^3)dx \cosh(dx+c) - 3a^3 - 4a^2b^2 + 6(3a^3 + 4a^2b^2) \cosh(dx+c)^2) \sinh(dx+c)^2 - 24((a^2b+b^3) \cosh(dx+c)^3 + 3(a^2b+b^3) \cosh(dx+c)^2 \sinh(dx+c) + 3(a^2b+b^3) \cosh(dx+c) \sinh(dx+c)^2 + (a^2b+b^3) \sinh(dx+c)^3) \log(2(a \sinh(dx+c) + b) / (\cosh(dx+c) - \sinh(dx+c))) + 3(2a^3 \cosh(dx+c)^5 - 5a^2b \cosh(dx+c)^4 + 24(a^2b+b^3)dx \cosh(dx+c)^2 + 4(3a^3 + 4a^2b^2) \cosh(dx+c)^3 - a^2b - 2(3a^3 + 4a^2b^2) \cosh(dx+c)) \sinh(dx+c) / (a^4d \cosh(dx+c)^3 + 3a^4d \cosh(dx+c)^2 \sinh(dx+c) + 3a^4d \cosh(dx+c) \sinh(dx+c)^2 + a^4d \sinh(dx+c)^3)}{1}$$

input `integrate(cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="fracas")`

output

```
1/24*(a^3*cosh(d*x + c)^6 + a^3*sinh(d*x + c)^6 - 3*a^2*b*cosh(d*x + c)^5
+ 24*(a^2*b + b^3)*d*x*cosh(d*x + c)^3 + 3*(2*a^3*cosh(d*x + c) - a^2*b)*s
inh(d*x + c)^5 + 3*(3*a^3 + 4*a*b^2)*cosh(d*x + c)^4 + 3*(5*a^3*cosh(d*x +
c)^2 - 5*a^2*b*cosh(d*x + c) + 3*a^3 + 4*a*b^2)*sinh(d*x + c)^4 - 3*a^2*b
*cosh(d*x + c) + 2*(10*a^3*cosh(d*x + c)^3 - 15*a^2*b*cosh(d*x + c)^2 + 12
*(a^2*b + b^3)*d*x + 6*(3*a^3 + 4*a*b^2)*cosh(d*x + c))*sinh(d*x + c)^3 -
a^3 - 3*(3*a^3 + 4*a*b^2)*cosh(d*x + c)^2 + 3*(5*a^3*cosh(d*x + c)^4 - 10*
a^2*b*cosh(d*x + c)^3 + 24*(a^2*b + b^3)*d*x*cosh(d*x + c) - 3*a^3 - 4*a*b
^2 + 6*(3*a^3 + 4*a*b^2)*cosh(d*x + c)^2)*sinh(d*x + c)^2 - 24*((a^2*b + b
^3)*cosh(d*x + c)^3 + 3*(a^2*b + b^3)*cosh(d*x + c)^2*sinh(d*x + c) + 3*(a
^2*b + b^3)*cosh(d*x + c)*sinh(d*x + c)^2 + (a^2*b + b^3)*sinh(d*x + c)^3)
*log(2*(a*sinh(d*x + c) + b)/(cosh(d*x + c) - sinh(d*x + c))) + 3*(2*a^3*c
osh(d*x + c)^5 - 5*a^2*b*cosh(d*x + c)^4 + 24*(a^2*b + b^3)*d*x*cosh(d*x +
c)^2 + 4*(3*a^3 + 4*a*b^2)*cosh(d*x + c)^3 - a^2*b - 2*(3*a^3 + 4*a*b^2)*
cosh(d*x + c))*sinh(d*x + c)/(a^4*d*cosh(d*x + c)^3 + 3*a^4*d*cosh(d*x +
c)^2*sinh(d*x + c) + 3*a^4*d*cosh(d*x + c)*sinh(d*x + c)^2 + a^4*d*sinh(d*
x + c)^3)
```

3.29.6 Sympy [F]

$$\int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = \int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$$

input `integrate(cosh(d*x+c)**3/(a+b*csch(d*x+c)),x)`

output `Integral(cosh(c + d*x)**3/(a + b*csch(c + d*x)), x)`

3.29. $\int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx$

3.29.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 183 vs. 2(81) = 162.

Time = 0.22 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.15

$$\int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = -\frac{(3abe^{(-dx-c)} - a^2 - 3(3a^2 + 4b^2)e^{(-2dx-2c)})e^{(3dx+3c)}}{24a^3d} - \frac{(a^2b + b^3)(dx + c)}{a^4d} - \frac{3abe^{(-2dx-2c)} + a^2e^{(-3dx-3c)} + 3(3a^2 + 4b^2)e^{(-dx-c)}}{24a^3d} - \frac{(a^2b + b^3)\log(-2be^{(-dx-c)} + ae^{(-2dx-2c)} - a)}{a^4d}$$

input `integrate(cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="maxima")`

output `-1/24*(3*a*b*e^(-d*x - c) - a^2 - 3*(3*a^2 + 4*b^2)*e^(-2*d*x - 2*c))*e^(3*d*x + 3*c)/(a^3*d) - (a^2*b + b^3)*(d*x + c)/(a^4*d) - 1/24*(3*a*b*e^(-2*d*x - 2*c) + a^2*e^(-3*d*x - 3*c) + 3*(3*a^2 + 4*b^2)*e^(-d*x - c))/(a^3*d) - (a^2*b + b^3)*log(-2*b*e^(-d*x - c) + a*e^(-2*d*x - 2*c) - a)/(a^4*d)`

3.29.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.71

$$\int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = \frac{a^2(e^{(dx+c)} - e^{(-dx-c)})^3 - 3ab(e^{(dx+c)} - e^{(-dx-c)})^2 + 12a^2(e^{(dx+c)} - e^{(-dx-c)}) + 12b^2(e^{(dx+c)} - e^{(-dx-c)})}{a^3} - \frac{24(a^2b + b^3)\log(|a(e^{(dx+c)} - e^{(-dx-c)})|)}{a^4} - \frac{24(a^2b + b^3)\log(|a(e^{(dx+c)} - e^{(-dx-c)})|)}{24d}$$

input `integrate(cosh(d*x+c)^3/(a+b*csch(d*x+c)),x, algorithm="giac")`

output `1/24*((a^2*(e^(d*x + c) - e^(-d*x - c)))^3 - 3*a*b*(e^(d*x + c) - e^(-d*x - c))^2 + 12*a^2*(e^(d*x + c) - e^(-d*x - c)) + 12*b^2*(e^(d*x + c) - e^(-d*x - c)))/a^3 - 24*(a^2*b + b^3)*log(abs(a*(e^(d*x + c) - e^(-d*x - c)) + 2*b))/a^4)/d`

3.29.9 Mupad [B] (verification not implemented)

Time = 2.47 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.12

$$\int \frac{\cosh^3(c+dx)}{a+b\operatorname{csch}(c+dx)} dx = \frac{x(a^2b+b^3)}{a^4} - \frac{e^{-3c-3dx}}{24ad} + \frac{e^{3c+3dx}}{24ad} - \frac{be^{-2c-2dx}}{8a^2d} - \frac{be^{2c+2dx}}{8a^2d} - \frac{e^{-c-dx}(3a^2+4b^2)}{8a^3d} - \frac{\ln(2be^{dx}e^c - a + ae^{2c}e^{2dx})(a^2b+b^3)}{a^4d} + \frac{e^{c+dx}(3a^2+4b^2)}{8a^3d}$$

input `int(cosh(c + d*x)^3/(a + b/sinh(c + d*x)),x)`

output `(x*(a^2*b + b^3))/a^4 - exp(- 3*c - 3*d*x)/(24*a*d) + exp(3*c + 3*d*x)/(24*a*d) - (b*exp(- 2*c - 2*d*x))/(8*a^2*d) - (b*exp(2*c + 2*d*x))/(8*a^2*d) - (exp(- c - d*x)*(3*a^2 + 4*b^2))/(8*a^3*d) - (log(2*b*exp(d*x)*exp(c) - a + a*exp(2*c)*exp(2*d*x))*(a^2*b + b^3))/(a^4*d) + (exp(c + d*x)*(3*a^2 + 4*b^2))/(8*a^3*d)`

APPENDIX

4.1 Listing of Grading functions	279
--	-----

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```



```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, CsCh,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```



```

if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):

```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #instance(expn,Pow)
    if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #instance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #instance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #instance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```



```
return grade, grade_annotation
```